



US 20030077747A1

(19) **United States**

(12) **Patent Application Publication**
Hillman et al.

(10) **Pub. No.: US 2003/0077747 A1**

(43) **Pub. Date: Apr. 24, 2003**

(54) **DELTA -6 DESATURASE HOMOLOGS**

Related U.S. Application Data

(75) Inventors: **Jennifer L. Hillman**, Santa Cruz, CA (US); **Karl J. Guegler**, Menlo Park, CA (US); **Neil C. Corley**, Castro Valley, CA (US); **Purvi Shah**, San Jose, CA (US)

(62) Division of application No. 09/048,888, filed on Mar. 26, 1998, now Pat. No. 6,492,108.

Publication Classification

Correspondence Address:

INCYTE GENOMICS, INC.
3160 PORTER DRIVE
PALO ALTO, CA 94304 (US)

(51) **Int. Cl.⁷** **C12N 9/04**; C07H 21/04;
C12P 21/02; C12N 1/21; C12N 5/06;
A01K 67/00

(52) **U.S. Cl.** **435/69.1**; 435/190; 435/252.3;
435/320.1; 435/325; 536/23.2;
800/8

(73) Assignee: **Incyte Genomics, Inc.**, Palo Alto, CA

(57) **ABSTRACT**

(21) Appl. No.: **10/262,617**

The invention provides a human delta-6 desaturase homolog (D6DH) and polynucleotides which identify and encode D6DH. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for treating or preventing disorders associated with expression of D6DH.

(22) Filed: **Sep. 30, 2002**

81	90	99	108
ATG GGC GGC GTC GGG GAG CCG GGA CCG CGG GAG			
M G G V G G E P G P R E			
117	126	135	144
GGA CCC GCG CAG CCG GGC GCA CCG CTG CCC ACC TTC TGC TGG GAG CAG ATC CGC			
G P A Q P G A P L P T F C W E Q I R			
171	180	189	198
GCG CAC GAC CAG CCC GGC GAC AAG TGG CTG GTC ATC GAG CGC CGC GTC TAC GAC			
A H D Q P G G D K W L V I E R R V Y D			
225	234	243	252
ATC AGC CGC TGG GCA CAG CCG CAC CCA GGG GGC AGC CGC CTC ATC GGC CAC CAC			
I S R W A Q R H P G G S R L I G H H			
279	288	297	306
GCG GCT GAG GAC GCC ACG GAT GCC TTC CGT GCC TTC CAT CAA GAT CTC AAT TTT			
G A E D A T D A F R A F H Q D L N F			
333	342	351	360
GTC CGC AAG TTC CTA CAG CCC CTG TTG ATT GGA GAG CTG GCT CCG GAA GAA CCC			
V R K F L Q CAG CCC CTG AAT GGC CAG CTG GTC GAT GGC GAA GAA CCC			
387	396	405	414
AGC CAG GAT GGA CCC CTG AAT GGC CAG CTG GTC GAG GAC TTC CGA GCC CTG CAC			
S Q D G P L N A Q L V E D F R A L H			

FIGURE 1A

441 450 459 468 477 486
 CAG GCA GCC GAG GAC ATG AAG CTG TTT GAT GCC AGT CCC ACC TTC TTT GCT TTC
 Q A A E D M K L F D A S P T F F A F

 495 504 513 522 531 540
 CTA CTG GGC CAC ATC CTG GCC ATG GAG GTG CTG GCC TGG CTC CTT ATC TAC CTC
 L L G H I L A M E V L A W L L I Y L

 549 558 567 576 585 594
 CTG GGT CCT GGC TGG GTG CCC AGT GCC CTG GCC GGC TTC ATC CTG GCC ATC TCT
 L G P G W V P S A L A A F I L A I S

 603 612 621 630 639 648
 CAG GCT CAG TCC TGG TGT CTG CAG CAT GAC CTG GGC CAT GCC TCC ATC TTC AAG
 Q A Q S W C L Q H D L G H A S I F K

 657 666 675 684 693 702
 AAG TCC TGG TGG AAC CAC GTG GCC CAG AAG TTC GTG ATG GGG CAG CTA AAG GGC
 K S W W N H V A Q K F V M G Q L K G

 711 720 729 738 747 756
 TTC TCC GCC CAC TGG TGG AAC TTC CGC CAC TTC CAG CAC CAC GCC AAG CCC AAC
 F S A H W W N F R H F Q H H A K P N

 765 774 783 792 801 810
 ATC TTC CAC AAA GAC CCA GAC GTG ACG GTG GCG CCC GTC TTC CTC CTG GGG GAG
 I F H K D P D V T V A P V F L L G E

FIGURE 1B

819 TCA TCC GTC GAG TAT GGC AAG AAG AAA CGC AGA TAC CTA CCC TAC AAC CAG CAG 864
 S S V E Y G TAT GGC AAG AAG AAA CGC AGA TAC CTA CCC TAC AAC CAG CAG
 828 GAG TAT GGC AAG AAG AAA CGC AGA TAC CTA CCC TAC AAC CAG CAG
 837 AAG AAG AAA CGC AGA TAC CTA CCC TAC AAC CAG CAG
 846 TAC CTA CCC TAC AAC CAG CAG
 855 TAC AAC CAG CAG
 864 CAG CAG CAG CAG
 873 CAC CTG TAC TTC TTC CTG ATC GGC CCG CTG CTC ACC CTG GTG AAC TTT GAA 918
 H L Y F F L I G P P L L L T L V N F E
 882 CTG ATC GGC CCG CTG CTC ACC CTG GTG AAC TTT GAA
 900 AAC TTT GAA
 891 CCG CTG CTC ACC CTG GTG AAC TTT GAA
 909 AAC TTT GAA
 927 GTG GAA AAT CTG GCG TAC ATG CTG GTG TGC ATG CAG TGG GCG GAT TTG CTC TGG 972
 V E N L A Y M L V C M Q W A D L L W
 936 TAC ATG CTG GTG TGC ATG CAG TGG GCG GAT TTG CTC TGG
 945 TGC ATG CAG TGG GCG GAT TTG CTC TGG
 954 TGC ATG CAG TGG GCG GAT TTG CTC TGG
 963 TGC ATG CAG TGG GCG GAT TTG CTC TGG
 972 TGC ATG CAG TGG GCG GAT TTG CTC TGG
 981 GCC GCC AGC TTC TAT GCC CGC TTC TTC TTA TCC TAC CTC CCC TTC TAC GGC GTC 1026
 A S F Y A R F F L S Y L P F Y G V
 990 TAT GCC CGC TTC TTC TTA TCC TAC CTC CCC TTC TAC GGC GTC
 1008 TCC TAC CTC CCC TTC TAC GGC GTC
 1017 TCC TTC TAC GGC GTC
 1026 TAC GGC GTC
 1035 CCT GGG GTG CTG CTC TTC TTT GTT GCT GTC AGG GTC CTG GAA AGC CAC TGG TTC 1080
 P G V L L F F V A V R V L E S H W F
 1044 CTG CTC TTC TTT GTT GCT GTC AGG GTC CTG GAA AGC CAC TGG TTC
 1053 GTC AGG GTC CTG GAA AGC CAC TGG TTC
 1062 GTC AGG GTC CTG GAA AGC CAC TGG TTC
 1071 GTC AGG GTC CTG GAA AGC CAC TGG TTC
 1080 GTC AGG GTC CTG GAA AGC CAC TGG TTC
 1089 GTG TGG ATC ACA CAG ATG AAC CAC ATC CCC AAG GAG ATC GGC CAC GAG AAG CAC 1134
 V W I T Q M N H I P K E I G H E K H
 1098 CAG ATG AAC CAC ATC CCC AAG GAG ATC GGC CAC GAG AAG CAC
 1107 CAC ATC CCC AAG GAG ATC GGC CAC GAG AAG CAC
 1116 CAC ATC CCC AAG GAG ATC GGC CAC GAG AAG CAC
 1125 CAC ATC CCC AAG GAG ATC GGC CAC GAG AAG CAC
 1134 CAC ATC CCC AAG GAG ATC GGC CAC GAG AAG CAC
 1143 CGG GAC TGG GTC AGC TCT CAG CTG GCA GCC ACC TGC AAC GTG GAG CCC TCA CTT 1188
 R D W V S S Q L A A T C N V E P S L
 1152 CAG CTG GCA GCC ACC TGC AAC GTG GAG CCC TCA CTT
 1161 CAG CTG GCA GCC ACC TGC AAC GTG GAG CCC TCA CTT
 1170 AAC GTG GAG CCC TCA CTT
 1179 AAC GTG GAG CCC TCA CTT
 1188 AAC GTG GAG CCC TCA CTT

FIGURE 1C

1197	1206	1215	1224	1233	1242
TTC ACC AAC TGG TTC AGC GGG CAC CTC AAC TTC CAG ATC GAG CAC CAC CTC TTC					
F T N W F S G H L N F Q I E H H L F					
1251	1260	1269	1278	1287	1296
CCC AGG ATG CCG AGA CAC AAC TAC AGC CCG GTG GCC CCG CTG GTC AAG TCG CTG					
P R M P R H N Y S R V A P L V K S L					
1305	1314	1323	1332	1341	1350
TGT GCC AAG CAC GGC CTC AGC TAC GAA GTG AAG CCC TTC CTC ACC GCG CTG GTG					
C A K H G L S Y E V K P F L T A L V					
1359	1368	1377	1386	1395	1404
GAC ATC GTC AGG TCC CTG AAG AAG TCT GGT GAC ATC TGG CTG GAC GCC TAC CTC					
D I V R S L K K S G D I W L D A Y L					

CAT CAG
H Q

FIGURE 1D

	90	99	108
	ATG GCC CCC GAC CCG GTG GCC GCC		
	M A P D P V A A		
	144	153	162
	GAG ACC GCG GCT CAG GGA CCT ACC CCG CGC TAC TTC ACC TGG GAC GAG GTG GCC		
	E T A A Q G G C C E E E E E E V A		
	180	189	207
	CAG CGC TCA GGG TGC GAG GAG GAG CCG TGG CTA GTG ATC GAC CGT AAG GTG TAC AAC		
	Q R S G C E E E E R W L V I D R K V Y N		
	225	234	261
	ATC AGC GAG TTC ACC CGC CGC CGG CAT CCA GGG GGC TCC CGG GTC ATC AGC CAC TAC		
	I S E F T R R R H P G G S R V I S H Y		
	279	288	315
	GCC GGG CAG GAT GCC ACG ACG GAT CCC TTT GTG GCC TTC CAC ATC AAC AAG GGC CTT		
	A G Q D A T A T D P F V A F H I N K G L		
	333	342	360
	GTG AAG AAG TAT ATG AAC TCT CTC CTG ATT GGA GAA CTG TCT CCA GAG CAG CCC		
	V K K Y M N S L L I G E L S P E Q P		
	387	396	414
	AGC TTT GAG CCC ACC AAG AAT AAA GAG CTG ACA GAT GAG TTC CGG GAG CTG CGG		
	S F E P T K N K E L T D E F R E L R		

FIGURE 2A

441 450 459 468 477 486
 GCC ACA GTG GAG CGG ATG GGG CTC ATG AAG GCC AAC CAT GTC TTC TTC CTG CTG CTG
 A T V E R M G L M K A N H V F F L L

 495 504 513 522 531 540
 TAC CTG CTG CAC ATC TTG TTG CTG CTG GGT GCA GCC TGG CTC ACC CTT TGG GTC
 Y L L H I L L L L D G A A W L T L W V

 549 558 567 576 585 594
 TTT GGG ACG TCC TTT TTG CCC TTC CTC CTC TGT GCG GTG CTG CTC AGT GCA GTT
 F G T S F L P F L L C A V L L S A V

 603 612 621 630 639 648
 CAG GCC CAG GCT GGC TGG CTG CAG CAT GAC TTT GGG CAC CTG TCG GTC TTC AGC
 Q A Q A G W L Q H D F G H L S V F S

 657 666 675 684 693 702
 ACC TCA AAG TGG AAC CAT CTG CTA CAT CAT TTT GTG ATT GGC CAC CTG AAG GGG
 T S K W N H L L L H H F V I G H L K G

 711 720 729 738 747 756
 GCC CCC GCC AGT TGG TGG AAC CAC ATG CAC TTC CAG CAC CAT GCC AAG CCC AAC
 A P A S W W N H M H F Q H H A K P N

 765 774 783 792 801 810
 TGC TTC CGC AAA GAC CCA GAC ATC AAC ATG CAT CCC TTC TCC TTT GCC TTG GGG
 C F R K D P D I N M H P F S F A L G

FIGURE 2B

819	AAG ATC CTC TCT GTG GAG CTT GGG AAA CAG AAG AAA AAA TAT ATG CCG TAC AAC	828	837	846	855	864
	K I L S V E L G K Q K K K Y M P Y N					
873	CAC CAG CAC AAA TAC TTC TTT CTA ATT GGG CCC CCA GCC TTG CTG CCT CTC TAC	882	891	900	909	918
	H Q H K Y F F L I G P P A L L P L Y					
927	TTC CAG TGG TAT ATT TTC TAT TTT GTT ATC CAG CGA AAG AAG TGG GTG GAC TTG	936	945	954	963	972
	F Q W Y I F Y Y F V I Q R K K W V D L					
981	GCC TGG ATG ATT ACC TTC TAC GTC CGC TTC TTC CTC ACT TAT GTG CCA CTA TTG	990	999	1008	1017	1026
	A W M I T F Y V R F F L T Y V P L L					
1035	GGG CTG AAA GCC TTC CTG GGC CTT TTC TTT ATA GTC AGG TTC CTG GAA AGC AAC	1044	1053	1062	1071	1080
	G L K A F L G L F F I V R F L E S N					
1089	TGG TTT GTG TGG GTG ACA CAG ATG AAC CAT ATT CCC ATG CAC ATT GAT CAT GAC	1098	1107	1116	1125	1134
	W F V W V T Q M N H I P M H I D H D					
1143	CGG AAC ATG GAC TGG GTT TCC ACC CAG CTC CAG GCC ACA TGC AAT GTC CAC AAG	1152	1161	1170	1179	1188
	R N M D W V S T Q L Q A T C N V H K					

FIGURE 2C

1	M	G	G	V	G	E	P	G	P	R	E	G	P	A	Q	P	G	A	P	L	2451043
1	M	A	-	-	-	-	P	D	P	V	A	A	E	T	A	A	Q	G	P	T	2056310
1	M	A	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	A	Q	I	2062403
1	M	V	-	-	-	-	S	P	S	I	E	V	L	N	S	I	A	A	D	G	1040729
21	P	T	-	F	C	W	E	Q	I	R	A	H	D	Q	P	G	D	K	W	L	2451043
17	P	R	Y	I	T	S	D	E	V	A	Q	R	S	G	C	E	E	R	W	L	2056310
6	K	K	Y	I	T	S	D	E	L	K	N	H	D	K	P	G	D	L	W	I	2062403
16	K	K	Y	I	T	S	K	E	L	K	H	N	N	P	N	D	L	L	W	I	1040729
40	V	I	E	R	R	V	Y	D	I	S	R	W	A	Q	R	H	P	G	G	S	2451043
37	V	I	D	R	K	V	Y	N	I	S	E	F	T	R	R	H	P	G	G	S	2056310
26	S	I	Q	G	K	A	Y	D	V	S	D	W	V	K	D	H	P	G	G	S	2062403
36	S	I	L	G	K	V	Y	N	V	T	E	W	A	K	E	H	P	G	G	D	1040729
60	R	L	I	G	H	H	G	A	E	D	A	T	D	A	F	R	A	F	H	Q	2451043
57	R	V	I	S	H	Y	A	G	Q	D	A	T	D	P	F	V	A	F	H	I	2056310
46	F	P	L	K	S	L	A	G	Q	E	V	T	D	A	F	V	A	F	H	P	2062403
56	A	P	L	I	N	L	A	G	Q	D	V	T	D	A	F	I	A	F	H	P	1040729
80	D	L	N	F	V	R	K	F	L	Q	P	L	L	I	G	E	L	A	P	E	2451043
77	N	K	G	L	V	K	K	Y	M	N	S	L	L	I	G	E	L	S	P	E	2056310
66	A	S	T	-	-	W	K	N	L	D	K	F	F	T	G	Y	L	K	D	2062403	
76	G	T	A	-	-	W	K	H	L	D	K	L	F	T	G	Y	H	L	K	D	1040729

FIGURE 3A

100	E	P	S	Q	D	G	P	L	N	A	Q	L	V	E	D	F	R	A	L	H	2451043
97	Q	P	S	F	E	P	T	K	N	K	E	L	T	D	E	F	R	E	L	R	2056310
84	Y	S	V	-	-	-	-	-	-	S	E	V	S	K	D	Y	R	K	L	V	2062403
94	Y	Q	V	-	-	-	-	-	-	S	D	I	S	R	D	Y	R	K	L	A	1040729
120	Q	A	A	E	D	M	K	L	F	-	-	D	A	S	P	T	F	F	A	F	2451043
117	A	T	V	E	R	M	G	L	M	-	-	K	A	N	H	V	F	F	L	L	2056310
98	F	E	F	S	K	M	G	L	Y	D	K	K	G	H	I	M	F	A	T	L	2062403
108	S	E	F	A	K	A	G	M	F	E	K	K	G	H	G	V	I	Y	S	L	1040729
138	L	L	G	H	I	L	A	M	E	V	L	A	W	L	L	I	Y	-	-	L	2451043
135	Y	L	L	H	I	L	L	D	G	A	A	W	L	T	L	W	-	-	V		2056310
118	C	F	I	A	M	L	L	F	A	-	-	-	M	S	V	Y	G	V	L	2062403	
128	C	F	V	S	L	L	L	S	-	-	-	-	A	C	V	Y	G	V	L	1040729	
156	L	G	P	G	W	V	P	S	A	L	A	A	F	I	L	A	I	S	Q	A	2451043
153	F	G	T	S	F	L	P	F	L	L	C	A	V	L	L	S	A	V	Q	A	2056310
133	F	C	E	G	V	L	V	H	L	L	F	S	G	C	L	M	G	F	L	I	2062403
143	Y	S	G	S	F	W	I	H	M	L	S	G	A	I	L	G	L	A	W	M	1040729
176	Q	S	-	W	C	L	Q	H	D	L	G	H	A	S	I	F	K	K	S	W	2451043
173	Q	A	G	W	-	L	Q	H	D	F	G	H	L	S	V	F	S	T	S	K	2056310
153	Q	S	G	W	-	I	G	H	D	A	G	H	Y	M	V	V	S	D	S	R	2062403
163	Q	I	A	Y	-	L	G	H	D	A	G	H	Y	Q	M	M	A	T	R	G	1040729

FIGURE 3B

195	W	N	H	V	A	Q	K	F	V	M	G	Q	L	K	G	F	S	A	H	W	2451043	
192	W	N	H	L	L	H	H	F	V	I	G	H	L	K	G	A	P	A	S	W	2056310	
172	L	N	K	F	M	G	I	F	A	A	N	C	L	S	G	T	S	I	G	W	2062403	
182	W	N	K	F	A	G	I	F	I	G	N	C	I	T	G	I	S	I	A	W	1040729	
215	W	N	F	R	H	F	Q	H	H	A	K	P	N	I	F	H	K	D	P	D	2451043	
212	W	N	H	M	H	F	Q	H	H	A	K	P	N	C	F	R	K	D	P	D	2056310	
192	W	K	W	N	H	N	A	H	H	I	A	C	N	S	L	E	Y	D	P	D	2062403	
202	W	K	W	T	H	N	A	H	H	I	A	C	N	S	L	D	Y	D	P	D	1040729	
235	V	T	V	A	P	V	-	F	L	L	G	E	S	-	S	-	-	-	-	-	2451043	
232	I	N	M	H	P	F	S	F	A	L	G	K	I	L	S	-	-	-	-	-	2056310	
212	L	Q	Y	I	P	F	L	V	V	S	S	K	F	F	G	S	L	T	S	H	2062403	
222	L	Q	H	L	P	M	L	L	A	V	S	S	K	L	F	N	S	I	T	S	V	1040729
248	-	-	-	-	-	-	-	V	E	Y	G	K	K	R	R	Y	L	P	Y	N	Q	2451043
247	-	-	-	-	-	-	-	V	E	L	G	K	Q	K	K	Y	M	P	Y	N	H	2056310
232	F	Y	E	K	R	L	T	F	F	D	S	L	S	R	F	F	V	S	Y	-	-	2062403
242	F	Y	G	R	Q	L	T	F	F	D	P	L	A	R	F	F	V	S	Y	-	-	1040729
263	Q	H	L	Y	F	F	L	I	G	P	P	L	L	T	L	V	N	F	E	V	2451043	
262	Q	H	K	Y	F	F	L	I	G	P	P	A	L	L	P	L	Y	F	Q	W	2056310	
250	Q	H	W	T	F	Y	P	I	M	C	A	A	R	L	N	M	Y	V	Q	S	2062403	
260	Q	H	Y	L	Y	P	P	I	M	C	V	A	R	V	N	L	Y	L	Q	T	1040729	

FIGURE 3C

283	E	N	L	A	Y	M	L	V	C	M	Q	W	-	-	A	D	L	L	W	A	2451043	
282	Y	I	F	Y	F	V	I	Q	R	K	K	W	-	-	V	D	L	A	W	M	2056310	
270	L	I	M	L	L	T	K	R	N	V	S	Y	R	A	H	E	L	L	G	C	2062403	
280	I	L	L	L	I	S	K	R	K	I	P	D	R	R	G	L	N	I	L	G	T	1040729
301	A	S	F	Y	A	R	F	-	-	F	L	S	Y	L	P	F	Y	G	V	P	2451043	
300	I	T	F	Y	V	R	F	-	-	F	L	T	Y	V	P	L	L	G	L	K	2056310	
290	L	V	F	S	I	W	Y	P	L	L	V	S	C	L	P	N	W	G	E	R	2062403	
300	L	I	F	W	T	W	F	P	L	L	V	S	R	L	P	N	W	P	E	R	1040729	
319	G	V	L	L	F	F	-	V	A	V	R	V	L	E	S	H	W	F	V	W	2451043	
318	A	F	L	G	L	F	-	F	I	V	R	F	L	E	S	N	W	F	V	W	2056310	
310	I	M	F	V	I	A	S	L	S	V	T	G	M	Q	Q	V	Q	F	S	-	2062403	
320	V	A	F	V	L	V	S	F	C	V	T	G	I	Q	H	I	Q	F	T	-	1040729	
338	I	T	Q	M	N	H	I	P	K	E	I	-	-	G	H	E	K	H	R	D	2451043	
337	V	T	Q	M	N	H	I	P	M	H	I	-	-	D	H	D	R	N	M	D	2056310	
329	-	-	-	L	N	H	F	S	S	V	Y	V	V	G	K	P	K	G	N	N	2062403	
339	-	-	-	L	N	H	F	S	G	D	V	V	V	G	P	P	K	G	D	N	1040729	
356	W	V	S	Q	L	A	A	T	C	N	V	E	P	S	L	F	T	N	W	W	2451043	
355	W	V	S	T	Q	L	Q	A	T	C	N	V	H	K	S	A	F	N	D	W	2056310	
346	W	F	E	K	Q	T	D	G	T	L	D	I	S	C	P	P	W	M	D	W	2062403	
356	W	F	E	K	Q	T	R	G	T	I	D	I	A	C	S	S	W	M	D	W	1040729	

FIGURE 3D

376	F	S	G	H	L	N	F	Q	I	E	H	H	L	F	P	R	M	P	R	H	2451043	
375	F	S	G	H	L	N	F	Q	I	E	H	H	L	F	P	T	M	P	R	H	2056310	
366	F	H	G	G	L	Q	F	Q	I	E	H	H	L	F	P	K	M	P	R	C	2062403	
376	F	F	G	G	L	Q	F	Q	L	E	H	H	L	F	P	R	L	P	R	C	1040729	
396	N	Y	S	R	V	A	P	L	V	K	S	L	C	A	K	H	G	L	S	Y	2451043	
395	N	Y	H	K	V	A	P	L	V	Q	S	L	C	A	K	H	G	I	E	Y	2056310	
386	N	L	R	K	I	S	P	Y	V	I	E	L	C	K	H	N	L	P	Y	2062403		
396	H	L	R	S	I	S	P	I	C	R	E	L	C	K	K	Y	N	L	P	Y	1040729	
416	E	V	K	P	F	L	T	A	L	V	D	I	V	R	S	L	K	K	S	G	2451043	
415	Q	S	K	P	L	L	S	A	F	A	D	I	I	H	S	L	K	E	S	G	2056310	
406	N	Y	A	S	F	S	K	A	N	E	M	T	L	R	T	L	R	N	T	A	2062403	
416	V	S	L	S	F	Y	D	A	N	V	T	T	L	K	T	L	R	T	A	A	1040729	
436	D	-	-	-	-	-	-	-	-	-	-	-	-	-	I	W	L	D	A	Y	2451043	
435	Q	-	-	-	-	-	-	-	-	-	-	-	-	-	L	W	L	D	A	Y	2056310	
426	L	Q	A	R	D	I	T	K	N	P	L	P	K	N	L	V	W	E	A	L	H	2062403
436	L	Q	A	R	D	L	T	N	P	P	A	P	Q	N	L	A	W	E	A	F	N	1040729
443	L	H	Q																			2451043
442	L	H	Q																			2056310
446	T	H	G																			2062403
456	T	H	G																			1040729

FIGURE 3E

DELTA -6 DESATURASE HOMOLOGS

[0001] This application is a divisional application of U.S. application Ser. No. 09/048,888, filed Mar. 26, 1998, entitled DELTA-6 DESATURASE HOMOLOGS, which is hereby expressly incorporated by reference herein.

FIELD OF THE INVENTION

[0002] This invention relates to nucleic acid and amino acid sequences of two delta-6 desaturase homologs and to the use of these sequences in the diagnosis, treatment, and prevention of cardiovascular diseases, disorders of aging, disorders of fatty acid metabolism, and cancer.

BACKGROUND OF THE INVENTION

[0003] Fatty acid desaturases catalyze the elongation and desaturation of the physiologically important hydrocarbon chain molecules known as fatty acids. Fatty acids are lipophilic modifiers of proteins, fuel molecules, insulating materials, and precursors to hormones including the prostaglandins, which are active in pyrogenesis, vasodilation, vasoconstriction, and pain reception. Phospholipids and glycolipids that compose cell membranes are made from fatty acids. Mammals derive unsaturated fatty acids from the unsaturated 16-carbon palmitate or diet-derived linoleate and linolenate, which are desaturated at two and three points, respectively. In eukaryotes, a variety of unsaturated fatty acids can be formed from oleate by a combination of elongation and desaturation reactions occurring on the cytosolic face of the endoplasmic reticulum membrane. Desaturation reactions require NADH and O₂ and are carried out by a complex consisting of a flavoprotein, a cytochrome, and a nonheme iron protein. The three components of the desaturase complex co-extract under pressure. It has been noted that each of the components of the delta-6 desaturase complex physically face a similar lipid environment in the microsomal membrane. (Leikin et al. (1994) *Biochim. Biophys. Acta* 1211:150-155.)

[0004] It is suggested that interaction between histidine-rich motifs in oleate desaturase could contribute to an iron binding site in the cytoplasmic domain of the protein that would facilitate interaction between the desaturase and cytochrome b5, the source of electrons for the desaturation reaction. (Okuley et al. (1994) *Plant Cell* 6:147-158.) Zinc also appears to play a role in delta-6 desaturation. (Eder, K. et al. (1996) *Biol. Trace Elem. Res.* 54:173-183.) A variety of delta-6 desaturase inhibitors are known, e.g., anti-oxidants and calcium channel blockers, and delta-6 desaturase activity in the body is regulated by several hormones. Inhibitors include, e.g., glucagon, epinephrine, ACTH, and glucocorticoids, while insulin and thyroxin are necessary to delta-6 desaturase activity, and growth hormone appears to enhance delta-6 desaturase activity. (Nakamura, M. T. et al. (1996) *Lipids* 31:139-143.) Expression of desaturase genes may also be influenced by diet. (Kaestner et al. (1989) *J. Biol. Chem.* 264:14755-14761.)

[0005] Several desaturases have been identified, including more than one delta-6 desaturase. (Marzo, I. et al. (1996) *Biochim. Biophys. Acta* 1301:263-272.) The presence of three histidine boxes and the HX(2-3)(XH)H amino acid motif are used to characterize membrane-bound fatty acid desaturases. (Schmidt et al. (1994) *Plant Mol. Biol.* 26:631-642; Mitchell et al. (1997) *J. Biol. Chem.* 272:28281-

28288.) A cytochrome b5 domain may also be used to characterize desaturases. A cDNA encoding the delta-6 desaturase from *Borago officinalis* that contained an N-terminal domain related to cytochrome b5 has been isolated. (Sayanova, O. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:4211-4216.)

[0006] Delta-6 desaturase is involved in a variety of diseases and disorders associated with fatty acid metabolism. A decrease in delta-6 desaturase activity is present in rats subjected to long-term exposure to ethanol. Diabetes mellitus involves many of the hormones that regulate delta-6 desaturase activity, such as insulin and glucagon. Obesity has been related to changes in delta-6 desaturase activity, and reduced delta-6 desaturase activity has also been implicated in coronary diseases. Delta-6 desaturation appears to decline with age, especially in women. Cancer cells show decreased delta-6 desaturase activity. Administration of delta-6 desaturation product, gamma linolenic acid, has been shown to ameliorate disorders of metabolism such as coronary disease, diabetes, osteoporosis, and ulcerative colitis, as well as disorders of cell proliferation such as tumorigenesis.

[0007] The discovery of a new delta-6 desaturase and the polynucleotides encoding it satisfies a need in the art by providing new compositions which are useful in the diagnosis, treatment, and prevention of cardiovascular diseases, disorders of aging, disorders of fatty acid metabolism, and cancer.

SUMMARY OF THE INVENTION

[0008] The invention features substantially purified polypeptides, delta-6 desaturase homologs, referred to collectively as "D6DH" and individually as "D6DH-1." and "D6DH-2." In one aspect, the invention provides a substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3.

[0009] The invention further provides a substantially purified variant having at least 90% amino acid identity to the amino acid sequences of SEQ ID NO:1 or SEQ ID NO:3, or to a fragment of either of these sequences. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3. The invention also includes an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3.

[0010] Additionally, the invention provides an isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3, as well as an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide encoding the polypeptide comprising the amino acid

sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3.

[0011] The invention also provides an isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, a fragment of SEQ ID NO:2, and a fragment of SEQ ID NO:4. The invention further provides an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide sequence comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, a fragment of SEQ ID NO:2, and a fragment of SEQ ID NO:4, as well as an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, a fragment of SEQ ID NO:2, and a fragment of SEQ ID NO:4.

[0012] The invention further provides an expression vector containing at least a fragment of the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3. In another aspect, the expression vector is contained within a host cell.

[0013] The invention also provides a method for producing a polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3, the method comprising the steps of: (a) culturing the host cell containing an expression vector containing at least a fragment of a polynucleotide encoding the polypeptide under conditions suitable for the expression of the polypeptide; and (b) recovering the polypeptide from the host cell culture.

[0014] The invention also provides a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3 in conjunction with a suitable pharmaceutical carrier.

[0015] The invention further includes a purified antibody which binds to a polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3, as well as a purified agonist and a purified antagonist to the polypeptide.

[0016] The invention also provides a method for treating or preventing a disorder of aging, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3.

[0017] The invention also provides a method for treating or preventing a cardiovascular disease, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially purified polypeptide having an amino

acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3.

[0018] The invention also provides a method for treating or preventing a disorder of fatty acid metabolism, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3.

[0019] The invention also provides a method for treating or preventing a cancer, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3.

[0020] The invention also provides a method for detecting a polynucleotide encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3 in a biological sample containing nucleic acids, the method comprising the steps of: (a) hybridizing the complement of the polynucleotide sequence encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, a fragment of SEQ ID NO:1, and a fragment of SEQ ID NO:3 to at least one of the nucleic acids of the biological sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of a polynucleotide encoding the polypeptide in the biological sample. In one aspect, the nucleic acids of the biological sample are amplified by the polymerase chain reaction prior to the hybridizing step.

BRIEF DESCRIPTION OF THE FIGURES

[0021] FIGS. 1A, 1B, 1C and 1D show the amino acid sequence (SEQ ID NO:1) and nucleic acid sequence (SEQ ID NO:2) of D6DH-1. The alignment was produced using MACDNASIS PRO software (Hitachi Software Engineering Co. Ltd., San Bruno, Calif.).

[0022] FIGS. 2A, 2B, 2C, and 2D show the amino acid sequence (SEQ ID NO:3) and nucleic acid sequence (SEQ ID NO:4) of D6DH-2. The alignment was produced using MACDNASIS PRO software.

[0023] FIGS. 3A, 3B, 3C, 3D, and 3E show the amino acid sequence alignments among D6DH-1 (2451043; SEQ ID NO:1), D6DH-2 (2056310; SEQ ID NO:3), delta-6 desaturase from *Borago officinalis* (GI 2062403; SEQ ID NO:5); and cytochrome b5 from *Helicanthus annus* (GI 1040729; SEQ ID NO:6), produced using the multisequence alignment program of LASERGENE software (DNASTAR Inc, Madison Wis.).

DESCRIPTION OF THE INVENTION

[0024] Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular methodology, protocols,

cell lines, vectors, and reagents described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

[0025] It must be noted that as used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to “a host cell” includes a plurality of such host cells, and a reference to “an antibody” is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

[0026] Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices, and materials are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, vectors, and methodologies which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

[0027] Definitions

[0028] “D6DH,” as used herein, refers to the amino acid sequences of substantially purified D6DH obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and preferably the human species, from any source, whether natural, synthetic, semi-synthetic, or recombinant.

[0029] The term “agonist,” as used herein, refers to a molecule which, when bound to D6DH, increases or prolongs the duration of the effect of D6DH. Agonists may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to and modulate the effect of D6DH.

[0030] An “allele” or an “allelic sequence,” as these terms are used herein, is an alternative form of the gene encoding D6DH. Alleles may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

[0031] “Altered” nucleic acid sequences encoding D6DH, as described herein, include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polynucleotide the same D6DH or a polypeptide with at least one functional characteristic of D6DH. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding D6DH, and improper or unexpected hybridization to alleles, with a locus other than the normal chromosomal locus for the polynucleotide sequence

encoding D6DH. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent D6DH. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of D6DH is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, positively charged amino acids may include lysine and arginine, and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine, isoleucine, and valine; glycine and alanine; asparagine and glutamine; serine and threonine; and phenylalanine and tyrosine.

[0032] The terms “amino acid” or “amino acid sequence,” as used herein, refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. In this context, “fragments,” “immunogenic fragments,” or “antigenic fragments” refer to fragments of D6DH which are preferably about 5 to about 15 amino acids in length and which retain some biological activity or immunological activity of D6DH. Where “amino acid sequence” is recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

[0033] “Amplification,” as used herein, relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art. (See, e.g., Dieffenbach, C. W. and G. S. Dveksler (1995) *PCR Primer; a Laboratory Manual*, Cold Spring Harbor Press, Plainview, N.Y., pp. 1-5.)

[0034] The term “antagonist,” as it is used herein, refers to a molecule which, when bound to D6DH, decreases the amount or the duration of the effect of the biological or immunological activity of D6DH. Antagonists may include proteins, nucleic acids, carbohydrates, antibodies, or any other molecules which decrease the effect of D6DH.

[0035] As used herein, the term “antibody” refers to intact molecules as well as to fragments thereof, such as Fa, F(ab)₂, and Fv fragments, which are capable of binding the epitopic determinant. Antibodies that bind D6DH polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

[0036] The term “antigenic determinant,” as used herein, refers to that fragment of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production

of antibodies which bind specifically to antigenic determinants (given regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

[0037] The term "antisense," as used herein, refers to any composition containing a nucleic acid sequence which is complementary to a specific nucleic acid sequence. The term "antisense strand" is used in reference to a nucleic acid strand that is complementary to the "sense" strand. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to block either transcription or translation. The designation "negative" can refer to the antisense strand, and the designation "positive" can refer to the sense strand.

[0038] As used herein, the term "biologically active," refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic D6DH, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

[0039] The terms "complementary" or "complementarity," as used herein, refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base pairing. For example, the sequence "A-G-T" binds to the complementary sequence "T-C-A." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands, and in the design and use of peptide nucleic acid (PNA) molecules.

[0040] A "composition comprising a given polynucleotide sequence" or a "composition comprising a given amino acid sequence," as these terms are used herein, refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation, an aqueous solution, or a sterile composition. Compositions comprising polynucleotide sequences encoding D6DH or fragments of D6DH may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

[0041] "Consensus sequence," as used herein, refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using XL-PCR (Perkin Elmer, Norwalk, Conn.) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of more than one Incyte Clone using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison, Wis.).

Some sequences have been both extended and assembled to produce the consensus sequence.

[0042] As used herein, the term "correlates with expression of a polynucleotide" indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding D6DH, by northern analysis is indicative of the presence of nucleic acids encoding D6DH in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding D6DH.

[0043] A "deletion," as the term is used herein, refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

[0044] The term "derivative," as used herein, refers to the chemical modification of D6DH, of a polynucleotide sequence encoding D6DH, or of a polynucleotide sequence complementary to a polynucleotide sequence encoding D6DH. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

[0045] The term "homology," as used herein, refers to a degree of complementarity. There may be partial homology or complete homology. The word "identity" may substitute for the word "homology." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially homologous." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially homologous sequence or hybridization probe will compete for and inhibit the binding of a completely homologous sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% homology or identity). In the absence of non-specific binding, the substantially homologous sequence or probe will not hybridize to the second non-complementary target sequence.

[0046] The phrases "percent identity" or "% identity" refer to the percentage of sequence similarity found in a comparison of two or more amino acid or nucleic acid sequences. Percent identity can be determined electronically, e.g., by using the MEGALIGN program (DNASTAR, Inc., Madison Wis.). The MEGALIGN program can create alignments between two or more sequences according to different methods, e.g., the clustal method. (See, e.g., Higgins, D. G. and P. M. Sharp (1988) *Gene* 73:237-244.) The clustal algorithm groups sequences into clusters by examining the distances between all pairs. The clusters are

aligned pairwise and then in groups. The percentage similarity between two amino acid sequences, e.g., sequence A and sequence B, is calculated by dividing the length of sequence A, minus the number of gap residues in sequence A, minus the number of gap residues in sequence B, into the sum of the residue matches between sequence A and sequence B, times one hundred. Gaps of low or of no homology between the two amino acid sequences are not included in determining percentage similarity. Percent identity between nucleic acid sequences can also be counted or calculated by other methods known in the art, e.g., the Jotun Hein method. (See, e.g., Hein, J. (1990) *Methods Enzymol.* 183:626-645.) Identity between sequences can also be determined by other methods known in the art, e.g., by varying hybridization conditions.

[0047] "Human artificial chromosomes" (HACs), as described herein, are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance. (See, e.g., Harrington, J. J. et al. (1997) *Nat. Genet.* 15:345-355.)

[0048] The term "humanized antibody," as used herein, refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

[0049] "Hybridization," as the term is used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

[0050] As used herein, the term "hybridization complex" as used herein, refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_{ot} or R_{ot} analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

[0051] The words "insertion" or "addition," as used herein, refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively, to the sequence found in the naturally occurring molecule.

[0052] "Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

[0053] The term "microarray," as used herein, refers to an arrangement of distinct polynucleotides arrayed on a substrate, e.g., paper, nylon or any other type of membrane, filter, chip, glass slide, or any other suitable solid support.

[0054] The terms "element" or "array element" as used herein in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

[0055] The term "modulate," as it appears herein, refers to a change in the activity of D6DH. For example, modulation may cause an increase or a decrease in protein activity,

binding characteristics, or any other biological, functional, or immunological properties of D6DH.

[0056] The phrases "nucleic acid" or "nucleic acid sequence," as used herein, refer to an oligonucleotide, nucleotide, polynucleotide, or any fragment thereof, to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material. In this context, "fragments" refers to those nucleic acid sequences which are greater than about 60 nucleotides in length, and most preferably are at least about 100 nucleotides, at least about 1000 nucleotides, or at least about 10,000 nucleotides in length.

[0057] The terms "operably associated" or "operably linked," as used herein, refer to functionally related nucleic acid sequences. A promoter is operably associated or operably linked with a coding sequence if the promoter controls the transcription of the encoded polypeptide. While operably associated or operably linked nucleic acid sequences can be contiguous and in the same reading frame, certain genetic elements, e.g., repressor genes, are not contiguously linked to the encoded polypeptide but still bind to operator sequences that control expression of the polypeptide.

[0058] The term "oligonucleotide," as used herein, refers to a nucleic acid sequence of at least about 6 nucleotides to 60 nucleotides, preferably about 15 to 30 nucleotides, and most preferably about 20 to 25 nucleotides, which can be used in PCR amplification or in a hybridization assay or microarray. As used herein, the term "oligonucleotide" is substantially equivalent to the terms "amplimer," "primer," "oligomer," and "probe," as these terms are commonly defined in the art.

[0059] "Peptide nucleic acid" (PNA), as used herein, refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA and RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell. (See, e.g., Nielsen, P. E. et al. (1993) *Anticancer Drug Des.* 8:53-63.)

[0060] The term "sample," as used herein, is used in its broadest sense. A biological sample suspected of containing nucleic acids encoding D6DH, or fragments thereof, or D6DH itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a solid support; a tissue; a tissue print; etc.

[0061] As used herein, the terms "specific binding" or "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

[0062] As used herein, the term "stringent conditions" refers to conditions which permit hybridization between

polynucleotide sequences and the claimed polynucleotide sequences. Suitably stringent conditions can be defined by, for example, the concentrations of salt or formamide in the prehybridization and hybridization solutions, or by the hybridization temperature, and are well known in the art. In particular, stringency can be increased by reducing the concentration of salt, increasing the concentration of formamide, or raising the hybridization temperature.

[0063] For example, hybridization under high stringency conditions could occur in about 50% formamide at about 37° C. to 42° C. Hybridization could occur under reduced stringency conditions in about 35% to 25% formamide at about 30° C. to 35° C. In particular, hybridization could occur under high stringency conditions at 42° C. in 50% formamide, 5×SSPE, 0.3% SDS, and 200 μ g/ml sheared and denatured salmon sperm DNA. Hybridization could occur under reduced stringency conditions as described above, but in 35% formamide at a reduced temperature of 35° C. The temperature range corresponding to a particular level of stringency can be further narrowed by calculating the purine to pyrimidine ratio of the nucleic acid of interest and adjusting the temperature accordingly. Variations on the above ranges and conditions are well known in the art.

[0064] The term “substantially purified,” as used herein, refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which they are naturally associated.

[0065] A “substitution,” as used herein, refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

[0066] “Transformation,” as defined herein, describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term “transformed” cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

[0067] A “variant” of D6DH, as used herein, refers to an amino acid sequence that is altered by one or more amino acids. The variant may have “conservative” changes, wherein a substituted amino acid has similar structural or chemical properties (e.g., replacement of leucine with isoleucine). More rarely, a variant may have “nonconservative” changes (e.g., replacement of glycine with tryptophan). Analogous minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, LASERGENE software.

[0068] The Invention

[0069] The invention is based on the discovery of new human delta-6 dehydrogenase homologs (D6DH), the polynucleotides encoding D6DH, and the use of these compositions for the diagnosis, treatment, or prevention of cardiovascular diseases, disorders of aging, disorders of fatty acid metabolism, and cancer.

[0070] Nucleic acids encoding D6DH-1 of the present invention were first identified in Incyte Clone 2451043 from an endothelial cell cDNA library (ENDANOT01) using a computer search for amino acid sequence alignments. A consensus sequence, SEQ ID NO:2, was derived from the following overlapping and/or extended nucleic acid sequences: Incyte Clones 2451043 (ENDANOT01), 2583121 (KIDNTUT13), 1986401 (LUNGAST01), 1432904 (BEPINON01), 1402462 (LATRUT02), 873359 (LUNGAST01), and 201775 (MPHGNOT02).

[0071] In one embodiment, the invention encompasses a polypeptide comprising the amino acid sequence of SEQ ID NO:1, as shown in FIGS. 1A-1D. D6DH-1 is 445 amino acids in length and has the three histidine boxes characteristic of delta-6 desaturases at residues H183-H187, H219-H223, and H351-H354. D6DH-1 has a potential N-glycosylation site at residue N396, a potential casein kinase II phosphorylation site at residue S246, and a potential protein kinase C phosphorylation site at S430. As shown in FIGS. 3A-3E, D6DH-1 has chemical and structural homology with delta-6 desaturase (GI 2062403; SEQ ID NO:5) and cytochrome b5 (GI 1040729; SEQ ID NO:6). In particular, D6DH-1 and delta-6 desaturase share 28% identity, including a histidine box at H219-H223, and D6DH-1 and cytochrome b5 share 27% homology. D6DH-1 and delta-6 desaturase have rather similar hydrophobicity plots and isoelectric points. The fragment of SEQ ID NO:2 from about nucleotide 104 to about nucleotide 128 is useful for designing oligonucleotides or to be used directly as a hybridization probe. Northern analysis shows the expression of this sequence in various libraries, 50% of which are cancerous, 20% are proliferating or immortalized, and 20% involve inflammation in the immune response. Of particular note is the expression of D6DH-1 in cardiovascular tissue (28%).

[0072] Nucleic acids encoding D6DH-2 of the present invention were first identified in Incyte Clone 2056310 from a bronchial epithelial cDNA library (BEPINOT01) using a computer search for amino acid sequence alignments. A consensus sequence, SEQ ID NO:4, was derived from the following overlapping and/or extended nucleic acid sequences: Incyte Clones 2056310 (BEPINOT01), 2290526 (BRAINON01), and 1362863 (LUNGNOT12).

[0073] In another embodiment, the invention encompasses a polypeptide comprising the amino acid sequence of SEQ ID NO:3, as shown in FIGS. 2A-2D. D6DH-2 is 444 amino acids in length and has the three histidine boxes characteristic of delta-6 desaturases at residues H179-H183, H217-H221, and H343-H347. D6DH-2 has a potential N-glycosylation site at residue N44, potential casein kinase II phosphorylation sites at residues T21, residues S29, and residues S429, and potential protein kinase C phosphorylation sites at residues T16, T49, T189, and S429. As shown in FIGS. 3A-3E, D6DH-2 has chemical and structural homology with delta-6 desaturase (GI 2062403; SEQ ID NO:5), and cytochrome b5 (GI 1040729; SEQ ID NO:6). In particular, D6DH-2 and delta-6 desaturase share 28% iden-

tity, including a histidine box at H217-H221, and D6DH-2 and cytochrome b5 share 27% identity. D6DH-2 and delta-6 desaturase have rather similar hydrophobicity plots and isoelectric points. The fragment of SEQ ID NO:2 from about nucleotide 102 to about nucleotide 134 is useful for designing oligonucleotides or to be used directly as a hybridization probe. Northern analysis shows the expression of this sequence in various libraries, 35% of which are cancerous, 32% of which involve the inflammatory response, and 22% of which are proliferating or immortalized. Of particular note is the expression of D6DH in cardiovascular tissue (29%).

[0074] The invention also encompasses D6DH variants. A preferred D6DH variant is one which has at least about 80%, more preferably at least about 90%, and most preferably at least about 95% amino acid sequence identity to the D6DH amino acid sequence, and which contains at least one functional or structural characteristic of D6DH.

[0075] The invention also encompasses polynucleotides which encode D6DH. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising the sequence of SEQ ID NO:2, which encodes D6DH-1. In a further embodiment, the invention encompasses a polynucleotide sequence comprising the sequence of SEQ ID NO:4, which encodes D6DH-2.

[0076] The invention also encompasses a variant of a polynucleotide sequence encoding D6DH. In particular, such a variant polynucleotide sequence will have at least about 80%, more preferably at least about 90%, and most preferably at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding D6DH. A particular aspect of the invention encompasses a variant of SEQ ID NO:2 which has at least about 80%, more preferably at least about 90%, and most preferably at least about 95% polynucleotide sequence identity to SEQ ID NO:2. The invention further encompasses a polynucleotide variant of SEQ ID NO:4 having at least about 80%, more preferably at least about 90%, and most preferably at least about 95% polynucleotide sequence identity to SEQ ID NO:4. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of D6DH.

[0077] It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding D6DH, some bearing minimal homology to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring D6DH, and all such variations are to be considered as being specifically disclosed.

[0078] Although nucleotide sequences which encode D6DH and its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring D6DH under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding D6DH or its derivatives possessing a substantially different codon usage. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular

prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding D6DH and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

[0079] The invention also encompasses production of DNA sequences which encode D6DH and D6DH derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents that are well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding D6DH or any fragment thereof.

[0080] Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:2, SEQ ID NO:4, a fragment of SEQ ID NO:2, or a fragment of SEQ ID NO:4, under various conditions of stringency. (See, e.g., Wahl, G. M. and S. L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A. R. (1987) *Methods Enzymol.* 152:507-511.)

[0081] Methods for DNA sequencing are well known and generally available in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (U.S. Biochemical Corp., Cleveland, Ohio), Taq polymerase (Perkin Elmer), thermostable T7 polymerase to (Amersham, Chicago, Ill.), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (GIBCO/BRL, Gaithersburg, Md.). Preferably, the process is automated with machines such as the Hamilton MICROLAB 2200 (Hamilton, Reno, N.Y.), Peltier PTC200 thermal cycler (MJ Research, Watertown, Mass.) and the ABI CATALYST and 373 and 377 DNA sequencers (Perkin Elmer).

[0082] The nucleic acid sequences encoding D6DH may be extended utilizing a partial nucleotide sequence and employing various methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal primers to retrieve unknown sequence adjacent to a known locus. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) In particular, genomic DNA is first amplified in the presence of a primer which is complementary to a linker sequence within the vector and a primer specific to a region of the nucleotide sequence. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

[0083] Inverse PCR may also be used to amplify or extend sequences using divergent primers based on a known region. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) The primers may be designed using commercially available software such as OLIGO 4.06 primer analysis software (National Biosciences Inc., Plymouth, Minn.) or

another appropriate program to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68° C. to 72° C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

[0084] Another method which may be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J. D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060.) Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries to walk genomic DNA (Clontech, Palo Alto, Calif.). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

[0085] When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Also, random-primed libraries are preferable in that they will include more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

[0086] Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

[0087] In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode D6DH may be used in recombinant DNA molecules to direct expression of D6DH, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced, and these sequences may be used to clone and express D6DH.

[0088] As will be understood by those of skill in the art, it may be advantageous to produce D6DH-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein

expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

[0089] The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter D6DH-encoding sequences for a variety of reasons including, but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

[0090] In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding D6DH may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of D6DH activity, it may be useful to encode a chimeric D6DH protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between the D6DH encoding sequence and the heterologous protein sequence, so that D6DH may be cleaved and purified away from the heterologous moiety.

[0091] In another embodiment, sequences encoding D6DH may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M. H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-232.) Alternatively, the protein itself may be produced using chemical methods to synthesize the amino acid sequence of D6DH, or a fragment thereof. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J. Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin Elmer). Additionally, the amino acid sequence of D6DH, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

[0092] The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R. M. and F. Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1983) *Proteins, Structures and Molecular Properties*, W H Freeman and Co., New York, N.Y.)

[0093] In order to express a biologically active D6DH, the nucleotide sequences encoding D6DH or derivatives thereof may be inserted into appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence.

[0094] Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding D6DH and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g.,

Sambrook, J. et al. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Plainview, N.Y., ch. 4, 8, and 16-17; and Ausubel, F. M. et al. (1995, and periodic supplements) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, N.Y., ch. 9, 13, and 16.)

[0095] A variety of expression vector/host systems may be utilized to contain and express sequences encoding D6DH. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus (CaMV) or tobacco mosaic virus (TMV)) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

[0096] The "control elements" or "regulatory sequences" are those non-translated regions, e.g., enhancers, promoters, and 5' and 3' untranslated regions, of the vector and polynucleotide sequences encoding D6DH which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters, e.g., hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, La Jolla, Calif.) or PSPORT1 plasmid (GIBCO/BRL), may be used. The baculovirus polyhedrin promoter may be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO, and storage protein genes) or from plant viruses (e.g., viral promoters or leader sequences) may be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding D6DH, vectors based on SV40 or EBV may be used with an appropriate selectable marker.

[0097] In bacterial systems, a number of expression vectors may be selected depending upon the use intended for D6DH. For example, when large quantities of D6DH are needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the sequence encoding D6DH may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced, and pIN vectors. (See, e.g., Van Heeke, G. and S. M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509.) pGEX vectors (Amersham Pharmacia Biotech, Uppsala, Sweden) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

[0098] In the yeast *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH, may be used. (See, e.g., Ausubel, supra; and Bitter, G. A. et al. (1987) *Methods Enzymol.* 153:516-544.)

[0099] In cases where plant expression vectors are used, the expression of sequences encoding D6DH may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV. (Takamatsu, N. (1987) *EMBO J.* 6:307-311.) Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews. (See, e.g., Hobbs, S. or Murry, L. E. in *McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York, N.Y.; pp. 191-196.)

[0100] An insect system may also be used to express D6DH. For example, in one such system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The sequences encoding D6DH may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of sequences encoding D6DH will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda* cells or *Trichoplusia* larvae in which D6DH may be expressed. (See, e.g., Engelhard, E. K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227.)

[0101] In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding D6DH may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing D6DH in infected host cells. (See, e.g., Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

[0102] Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes.

[0103] Specific initiation signals may also be used to achieve more efficient translation of sequences encoding D6DH. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding D6DH and its initiation codon and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be

needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

[0104] In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding, and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC, Bethesda, Md.) and may be chosen to ensure the correct modification and processing of the foreign protein.

[0105] For long term, high yield production of recombinant proteins, stable expression is preferred. For example, cell lines capable of stably expressing D6DH can be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

[0106] Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase genes and adenine phosphoribosyltransferase genes, which can be employed in tk⁻ and apr⁻ cells, respectively. (See, e.g., Wigler, M. et al. (1977) *Cell* 11:223-232; and Lowy, I. et al. (1980) *Cell* 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, dhfr confers resistance to methotrexate; npt confers resistance to the aminoglycosides neomycin and G-418; and als and pat confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci. USA* 77:3567-3570; Colbere-Garapin, F. et al (1981) *J. Mol. Biol.* 150:1-14; and Murry, supra.) Additional selectable genes have been described, e.g., trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine. (See, e.g., Hartman, S. C. and R. C. Mulligan (1988) *Proc. Natl. Acad. Sci. USA* 85:8047-8051.) Visible markers, e.g., anthocyanins, β glucuronidase and its substrate GUS, luciferase and its substrate luciferin may be used. Green fluorescent proteins (GFP) (Clontech, Palo Alto, Calif.) can also be used. These markers

can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (See, e.g., Rhodes, C. A. et al. (1995) *Methods Mol. Biol.* 55:121-131.)

[0107] Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding D6DH is inserted within a marker gene sequence, transformed cells containing sequences encoding D6DH can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding D6DH under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

[0108] Alternatively, host cells which contain the nucleic acid sequence encoding D6DH and express D6DH may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

[0109] The presence of polynucleotide sequences encoding D6DH can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding D6DH. Nucleic acid amplification based assays involve the use of oligonucleotides or oligomers based on the sequences encoding D6DH to detect transformants containing DNA or RNA encoding D6DH.

[0110] A variety of protocols for detecting and measuring the expression of D6DH, using either polyclonal or monoclonal antibodies specific for the protein, are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on D6DH is preferred, but a competitive binding assay may be employed. These and other assays are well described in the art. (See, e.g., Hampton, R. et al. (1990) *Serological Methods, a Laboratory Manual*, APS Press, St. Paul, Minn., Section IV; and Maddox, D. E. et al. (1983) *J. Exp. Med.* 158:1211-1216).

[0111] A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding D6DH include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding D6DH, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Pharmacia & Upjohn (Kalamazoo, Mich.), Promega (Madison, Wis.), and U.S. Biochemical Corp. (Cleveland, Ohio). Suitable reporter

molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

[0112] Host cells transformed with nucleotide sequences encoding D6DH may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode D6DH may be designed to contain signal sequences which direct secretion of D6DH through a prokaryotic or eukaryotic cell membrane. Other constructions may be used to join sequences encoding D6DH to nucleotide sequences encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). The inclusion of cleavable linker sequences, such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, Calif.), between the purification domain and the D6DH encoding sequence may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing D6DH and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on immobilized metal ion affinity chromatography (IMAC). (See, e.g., Porath, J. et al. (1992) *Prot. Exp. Purif.* 3: 263-281.) The enterokinase cleavage site provides a means for purifying D6DH from the fusion protein. (See, e.g., Kroll, D. J. et al. (1993) *DNA Cell Biol.* 12:441-453.)

[0113] Fragments of D6DH may be produced not only by recombinant production, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, T. E. (1984) *Protein: Structures and Molecular Properties*, pp. 55-60, W. H. Freeman and Co., New York, N.Y.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the Applied Biosystems 431A peptide synthesizer (Perkin Elmer). Various fragments of D6DH may be synthesized separately and then combined to produce the full length molecule.

[0114] Therapeutics

[0115] Chemical and structural homology exists among D6DH, delta-6 desaturase from *Borago officinalis* (GI2062403), and cytochrome b5 from *Helicanthus annus* (GI 1040729). In addition, D6DH-1 is expressed in cancer, proliferating or immortalized cells, the immune response, and cardiovascular tissue. Therefore, D6DH appears to play a role in cardiovascular disease, disorders of aging, disorders of fatty acid metabolism, and cancer.

[0116] Therefore, in one embodiment, D6DH or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder of aging. Such disorders can include, but are not limited to, Alzheimer's Disease, angio-immunoblastic lymphadenopathy, anorexia, atherosclerosis,

basal cell carcinoma, cardiac amyloidosis, cerebral amyloidosis, chronic lymphatic leukemia, decubitus ulcers, degenerative osteoarthritis, delirium, dementia, depression, diabetes, diabetic hyperosmolar nonketotic coma, dyskinesia, glaucoma, hypertension, hypercholesterolemia, hypothermia, metabolic bone disease including osteoporosis, nephrosclerosis, normal pressure hydrocephalus, osteoarthritis, Parkinson's disease, Reye's syndrome, polymyalgia rheumatica, prostatic carcinoma, renal amyloidosis, stroke, tuberculosis, and urinary incontinence.

[0117] In another embodiment, a vector capable of expressing D6DH or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder of aging including, but not limited to, those described above.

[0118] In a further embodiment, a pharmaceutical composition comprising a substantially purified D6DH in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder of aging including, but not limited to, those provided above.

[0119] In still another embodiment, an agonist which modulates the activity of D6DH may be administered to a subject to treat or prevent a disorder of aging including, but not limited to, those listed above.

[0120] In another embodiment, D6DH or a fragment or derivative thereof may be administered to a subject to treat or prevent a cardiovascular disease. Such disorders can include, but are not limited to, angina pectoris, atheroembolic disease, atherosclerosis, arteriosclerosis, cardiac ischemia, gangrene hypercholesterolemia, hyperlipidemia, hypertriglyceridemia, hypertension, mesenteric ischemia, renal artery stenosis, nephrosclerosis, and stroke.

[0121] In another embodiment, a vector capable of expressing D6DH or a fragment or derivative thereof may be administered to a subject to treat or prevent a cardiovascular disease including, but not limited to, those described above.

[0122] In a further embodiment, a pharmaceutical composition comprising a substantially purified D6DH in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a cardiovascular disease including, but not limited to, those provided above.

[0123] In still another embodiment, an agonist which modulates the activity of D6DH may be administered to a subject to treat or prevent a cardiovascular disease including, but not limited to, those listed above.

[0124] In another embodiment, D6DH or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder of fatty acid metabolism. Such disorders can include, but are not limited to, Addison's disease, cystic fibrosis, diabetes, fatty hepatocirrhosis, galactosemia, goiter, hyperadrenalism, hypoadrenalism, hyperparathyroidism, hypoparathyroidism, hypercholesterolemia, hyperthyroidism, hypothyroidism hyperlipidemia, hyperlipemia, lipid myopathies, obesity, lipodystrophies, and phenylketonuria.

[0125] In another embodiment, a vector capable of expressing D6DH or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder of fatty acid metabolism including, but not limited to, those described above.

[0126] In a further embodiment, a pharmaceutical composition comprising a substantially purified D6DH in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder of fatty acid metabolism including, but not limited to, those provided above.

[0127] In still another embodiment, an agonist which modulates the activity of D6DH may be administered to a subject to treat or prevent a disorder of fatty acid metabolism including, but not limited to, those listed above.

[0128] In another embodiment, D6DH or a fragment or derivative thereof may be administered to a subject to treat or prevent cancer. Such cancer can include, but are not limited to, adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

[0129] In another embodiment, a vector capable of expressing D6DH or a fragment or derivative thereof may be administered to a subject to treat or prevent a cancer including, but not limited to, those described above.

[0130] In a further embodiment, a pharmaceutical composition comprising a substantially purified D6DH in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a cancer including, but not limited to, those provided above.

[0131] In still another embodiment, an agonist which modulates the activity of D6DH may be administered to a subject to treat or prevent a cancer including, but not limited to, those listed above.

[0132] In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

[0133] An antagonist of D6DH may be produced using methods which are generally known in the art. In particular, purified D6DH may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind D6DH. Antibodies to D6DH may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

[0134] For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with D6DH or with any fragment

or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and *Corynebacterium parvum* are especially preferable.

[0135] It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to D6DH have an amino acid sequence consisting of at least about 5 amino acids, and, more preferably, of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of D6DH amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

[0136] Monoclonal antibodies to D6DH may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R. J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and Cole, S. P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

[0137] In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S. L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M. S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce D6DH-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D. R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

[0138] Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; and Winter, G. et al. (1991) *Nature* 349:293-299.)

[0139] Antibody fragments which contain specific binding sites for D6DH may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W. D. et al. (1989) *Science* 246:1275-1281.)

[0140] Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numer-

ous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between D6DH and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering D6DH epitopes is preferred, but a competitive binding assay may also be employed. (Maddox, supra.)

[0141] In another embodiment of the invention, the polynucleotides encoding D6DH, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding D6DH may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding D6DH. Thus, complementary molecules or fragments may be used to modulate D6DH activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding D6DH.

[0142] Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors which will express nucleic acid sequences complementary to the polynucleotides of the gene encoding D6DH. (See, e.g., Sambrook, supra; and Ausubel, supra.)

[0143] Genes encoding D6DH can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding D6DH. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system

[0144] As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding D6DH. Oligonucleotides derived from the transcription initiation site, e.g., between about positions 10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J. E. et al. (1994) in Huber, B. E. and B. I. Carr, *Molecular and Immunologic Approaches*, Futura Publishing Co., Mt. Kisco, N.Y., pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

[0145] Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The

mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding D6DH.

[0146] Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

[0147] Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding D6DH. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

[0148] RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

[0149] Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C. K. et al. (1997) *Nat. Biotechnol.* 15:462-466.)

[0150] Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

[0151] An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of D6DH,

antibodies to D6DH, and mimetics, agonists, antagonists, or inhibitors of D6DH. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

[0152] The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

[0153] In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of *Remington's Pharmaceutical Sciences* (Maack Publishing Co., Easton, Pa.).

[0154] Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

[0155] Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

[0156] Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

[0157] Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

[0158] Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

[0159] For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

[0160] The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

[0161] The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acid. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 MM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

[0162] After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of D6DH, such labeling would include amount, frequency, and method of administration.

[0163] Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

[0164] For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

[0165] A therapeutically effective dose refers to that amount of active ingredient, for example D6DH or fragments thereof, antibodies of D6DH, and agonists, antagonists or inhibitors of D6DH, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculat-

ing the ED₅₀ (the dose therapeutically effective in 50% of the population) or LD₅₀ (the dose lethal to 50% of the population) statistics. The dose ratio of therapeutic to toxic effects is the therapeutic index, and it can be expressed as the ED₅₀/LD50 ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED₅₀ with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

[0166] The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

[0167] Normal dosage amounts may vary from about 0.1 μ g to 100,000 μ g, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

[0168] Diagnostics

[0169] In another embodiment, antibodies which specifically bind D6DH may be used for the diagnosis of disorders characterized by expression of D6DH, or in assays to monitor patients being treated with D6DH or agonists, antagonists, or inhibitors of D6DH. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for D6DH include methods which utilize the antibody and a label to detect D6DH in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

[0170] A variety of protocols for measuring D6DH, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of D6DH expression. Normal or standard values for D6DH expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to D6DH under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, preferably by photometric means. Quantities of D6DH expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

[0171] In another embodiment of the invention, the polynucleotides encoding D6DH may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of D6DH may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of D6DH, and to monitor regulation of D6DH levels during therapeutic intervention.

[0172] In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding D6DH or closely related molecules may be used to identify nucleic acid sequences which encode D6DH. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low), will determine whether the probe identifies only naturally occurring sequences encoding D6DH, alleles, or related sequences.

[0173] Probes may also be used for the detection of related sequences, and should preferably have at least 50% sequence identity to any of the D6DH encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequences of SEQ ID NO:2, SEQ ID NO:4, or from genomic sequences including promoters, enhancers, and introns of the D6DH gene.

[0174] Means for producing specific hybridization probes for DNAs encoding D6DH include the cloning of polynucleotide sequences encoding D6DH or D6DH derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ³²P or ³⁵S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

[0175] Polynucleotide sequences encoding D6DH may be used for the diagnosis of a disorder associated with expression of D6DH. Examples of such a disorder include, but are not limited to, disorders of aging such as Alzheimer's Disease, angioimmunoblastic lymphadenopathy, anorexia, atherosclerosis, basal cell carcinoma, cardiac amyloidosis, cerebral amyloidosis, chronic lymphatic leukemia, decubitus ulcers, degenerative osteoarthritis, delirium, dementia, depression, diabetes, diabetic hyperosmolar nonketotic coma, dyskinesia, glaucoma, hypertension, hypercholesterolemia, hypothermia, metabolic bone disease including osteoporosis, nephrosclerosis, normal pressure hydrocephalus, osteoarthritis, Parkinson's disease, Reye's syndrome, polymyalgia rheumatica, prostatic carcinoma, renal amyloidosis, stroke, tuberculosis, and urinary incontinence; disorders of fatty acid metabolism such as Addison's disease, cystic fibrosis, diabetes, fatty hepatocirrhosis, galactosemia, goiter, hyperadrenalism, hypoadrenalism, hyperparathyroidism, hypoparathyroidism, hypercholesterolemia, hyperthyroidism, hypothyroidism, hyperlipidemia, hyperlipemia,

lipid myopathies, obesity, lipodystrophies, and phenylketonuria; cardiovascular disease such as angina pectoris, atheroembolic disease, atherosclerosis, arteriosclerosis, cardiac ischemia, gangrene hypercholesterolemia, hyperlipidemia, hypertriglyceridemia, hypertension, mesenteric ischemia, renal artery stenosis, nephrosclerosis, and stroke; and cancer such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

[0176] The polynucleotide sequences encoding D6DH may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; and in microarrays utilizing fluids or tissues from patients to detect altered D6DH expression. Such qualitative or quantitative methods are well known in the art.

[0177] In a particular aspect, the nucleotide sequences encoding D6DH may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding D6DH may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding D6DH in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

[0178] In order to provide a basis for the diagnosis of a disorder associated with expression of D6DH, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding D6DH, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

[0179] Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

[0180] With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of

the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

[0181] Additional diagnostic uses for oligonucleotides designed from the sequences encoding D6DH may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding D6DH, or a fragment of a polynucleotide complementary to the polynucleotide encoding D6DH, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantitation of closely related DNA or RNA sequences.

[0182] Methods which may also be used to quantitate the expression of D6DH include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P. C. et al. (1993) *J. Immunol. Methods* 159:235-244; and Duplaa, C. et al. (1993) *Anal. Biochem* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

[0183] In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

[0184] Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T. M. et al. (1995) U.S. Pat. No. 5,474,796; Schena, M. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R. A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-2155; and Heller, M. J. et al. (1997) U.S. Pat. No. 5,605,662.)

[0185] In another embodiment of the invention, nucleic acid sequences encoding D6DH may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Price, C. M. (1993) *Blood Rev.* 7:127-134; and Trask, B. J. (1991) *Trends Genet.* 7:149-154.)

[0186] Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al.

(1995) in Meyers, R. A. (ed.) *Molecular Biology and Biotechnology*, VCH Publishers New York, N.Y., pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) site. Correlation between the location of the gene encoding D6DH on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

[0187] In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., A1 to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R. A. et al. (1988) *Nature* 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

[0188] In another embodiment of the invention, D6DH, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between D6DH and the agent being tested may be measured.

[0189] Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with D6DH, or fragments thereof, and washed. Bound D6DH is then detected by methods well known in the art. Purified D6DH can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

[0190] In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding D6DH specifically compete with a test compound for binding D6DH. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with D6DH.

[0191] In additional embodiments, the nucleotide sequences which encode D6DH may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide

sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

[0192] The examples below are provided to illustrate the subject invention and are not included for the purpose of limiting the invention.

EXAMPLES

[0193] I. cDNA Library Construction

[0194] ENDANOT01

[0195] The ENDANOT01 cDNA library was constructed from an aortic endothelial cell line derived from explanted heart/aorta tissue obtained from a male (specimen # A062).

[0196] The frozen cells were homogenized and lysed using a Brinkmann POLYTRON PT-3000 homogenizer (Brinkmann Instruments, Westbury, N.Y.) in guanidinium isothiocyanate solution. The lysate was centrifuged over a 5.7 M CsCl cushion using a Beckman SW28 rotor in a Beckman L8-70M ultracentrifuge (Beckman Instruments) for 18 hours at 25,000 rpm at ambient temperature. The RNA was extracted with acid phenol pH 4.7, precipitated using 0.3 M sodium acetate and 2.5 volumes of ethanol, resuspended in RNase-free water, and DNase treated at 37° C. RNA extraction and precipitation were repeated twice as before. The mRNA was then isolated using the Qiagen OLIGOTEX kit (QIAGEN, Inc.; Chatsworth, Calif.) and used to construct the cDNA library.

[0197] The mRNA was handled according to the recommended protocols in the SUPERSRIPT plasmid system for cDNA synthesis and plasmid cloning (Cat. #18248-013; Gibco/BRL). The cDNAs were fractionated on a SEPHAROSE CL4B column (Cat. #275105-01; Pharmacia), and those cDNAs exceeding 400 bp were ligated into pINCY1. The plasmid pINCY1 was subsequently transformed into DH5a competent cells (Cat. #18258-012; Gibco/BRL).

[0198] BEPINOT01

[0199] The BEPINOT01 cDNA library was constructed from a bronchial epithelium primary cells line derived from a 54-year-old Caucasian male. The frozen tissue was homogenized and lysed using a Brinkmann POLYTRON PT-3000 homogenizer (Brinkmann Instruments, Westbury, N.Y.) in guanidinium isothiocyanate solution. The lysate was centrifuged over a 5.7 M CsCl cushion using a Beckman SW28 rotor in a Beckman L8-70M ultracentrifuge (Beckman Instruments) for 18 hours at 25,000 rpm at ambient temperature. The RNA was extracted with acid phenol pH 4.7, precipitated using 0.3 M sodium acetate and 2.5 volumes of ethanol, resuspended in RNase-free water, and DNase treated at 37° C. The RNA extraction and precipitation were repeated twice as before. The mRNA was then isolated using the Qiagen OLIGOTEX kit (QIAGEN, Inc., Chatsworth, Calif.) and used to construct the cDNA library.

[0200] The mRNA was handled according to the recommended protocols in the SUPERSRIPT plasmid system for cDNA synthesis and plasmid cloning (Catalog #18248-013, Gibco/BRL). The cDNAs were fractionated on a SEPHAROSE CL4B column (Catalog #275105-01, Pharmacia), and those cDNAs exceeding 400 bp were ligated

into PSPORT1. The plasmid PSPORT1 was subsequently transformed into DH5a competent cells (Catalog #18258-012, Gibco/BRL).

[0201] II. Isolation and Sequencing of cDNA Clones

[0202] Plasmid DNA was released from the cells and purified using the REAL Prep 96 plasmid kit (Catalog #26173; QIAGEN, Inc.). This kit enabled the simultaneous purification of 96 samples in a 96-well block using multi-channel reagent dispensers. The recommended protocol was employed except for the following changes: 1) the bacteria were cultured in 1 ml of sterile Terrific Broth (Catalog #22711, Gibco/BRL) with carbenicillin at 25 mg/L and glycerol at 0.4%; 2) after inoculation, the cultures were incubated for 19 hours and at the end of incubation, the cells were lysed with 0.3 ml of lysis buffer; and 3) following isopropanol precipitation, the plasmid DNA pellet was resuspended in 0.1 ml of distilled water. After the last step in the protocol, samples were transferred to a 96-well block for storage at 4° C.

[0203] The cDNAs were sequenced by the method of Sanger et al. (1975, *J. Mol. Biol.* 94:441f), using a Hamilton MICROLAB 2200 (Hamilton, Reno, Nev.) in combination with Peltier thermal cyclers (PTC200 from MJ Research, Watertown, Mass.) and Applied Biosystems 377 DNA sequencing systems.

[0204] III. Homology Searching of cDNA Clones and Their Deduced Proteins

[0205] The nucleotide sequences and/or amino acid sequences of the Sequence Listing were used to query sequences in the GenBank, SwissProt, BLOCKS, and Pima II databases. These databases, which contain previously identified and annotated sequences, were searched for regions of homology using BLAST (Basic Local Alignment Search Tool). (See, e.g., Altschul, S. F. (1993) *J. Mol. Evol.* 36:290-300; and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410.)

[0206] BLAST produced alignments of both nucleotide and amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST was especially useful in determining exact matches or in identifying homologs which may be of prokaryotic (bacterial) or eukaryotic (animal, fungal, or plant) origin. Other algorithms could have been used when dealing with primary sequence patterns and secondary structure gap penalties. (See, e.g., Smith, T. et al. (1992) *Protein Eng.* 5:35-51.) The sequences disclosed in this application have lengths of at least 49 nucleotides and have no more than 12% uncalled bases (where N is recorded rather than A, C, G, or T).

[0207] The BLAST approach searched for matches between a query sequence and a database sequence. BLAST evaluated the statistical significance of any matches found, and reported only those matches that satisfy the user-selected threshold of significance. In this application, threshold was set at 10^{-25} for nucleotides and 10^{-8} for peptides.

[0208] Incyte nucleotide sequences were searched against the GenBank databases for primate (pri), rodent (rod), and other mammalian sequences (mam), and deduced amino acid sequences from the same clones were then searched against GenBank functional protein databases, mammalian (mamp), vertebrate (vrtp), and eukaryote (eukp), for homology.

[0209] Additionally, sequences identified from cDNA libraries may be analyzed to identify those gene sequences encoding conserved protein motifs using an appropriate analysis program, e.g., the Block 2 Bioanalysis Program (Incyte, Palo Alto, Calif.). This motif analysis program, based on sequence information contained in the Swiss-Prot Database and PROSITE, is a method of determining the function of uncharacterized proteins translated from genomic or cDNA sequences. (See, e.g., Bairoch, A. et al. (1997) *Nucleic Acids Res.* 25:217-221; and Attwood, T. K. et al. (1997) *J. Chem. Inf. Comput. Sci.* 37:417-424.) PROSITE may be used to identify common functional or structural domains in divergent proteins. The method is based on weight matrices. Motifs identified by this method are then calibrated against the SWISS-PROT database in order to obtain a measure of the chance distribution of the matches.

[0210] In another alternative, Hidden Markov models (HMMs) may be used to find protein domains, each defined by a dataset of proteins known to have a common biological function. (See, e.g., Pearson, W. R. and D. J. Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85:2444-2448; and Smith, T. F. and M. S. Waterman (1981) *J. Mol. Biol.* 147:195-197.) HMMs were initially developed to examine speech recognition patterns, but are now being used in a biological context to analyze protein and nucleic acid sequences as well as to model protein structure. (See, e.g., Krogh, A. et al. (1994) *J. Mol. Biol.* 235:1501-1531; and Collin, M. et al. (1993) *Protein Sci.* 2:305-314.) HMMs have a formal probabilistic basis and use position-specific scores for amino acids or nucleotides. The algorithm continues to incorporate information from newly identified sequences to increase its motif analysis capabilities.

[0211] IV. Northern Analysis

[0212] Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; and Ausubel, supra, ch. 4 and 16.)

[0213] Analogous computer techniques applying BLAST are used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ database (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or homologous.

[0214] The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

[0215] The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Homolo-

gous molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

[0216] The results of northern analysis are reported as a list of libraries in which the transcript encoding D6DH occurs. Abundance and percent abundance are also reported. Abundance directly reflects the number of times a particular transcript is represented in a cDNA library, and percent abundance is abundance divided by the total number of sequences examined in the cDNA library.

[0217] V. Extension of D6DH Encoding Polynucleotides

[0218] The nucleic acid sequences of Incyte Clones 2451043 and 2056310 were used to design oligonucleotide primers for extending partial nucleotide sequences to full length. For each nucleic acid sequence, one primer was synthesized to initiate extension of an antisense polynucleotide, and the other was synthesized to initiate extension of a sense polynucleotide. Primers were used to facilitate the extension of the known sequence "outward" generating amplicons containing new unknown nucleotide sequence for the region of interest. The initial primers were designed from the cDNA using OLIGO 4.06 (National Biosciences, Plymouth, Minn.), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68° C. to about 72° C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

[0219] Selected human cDNA libraries (GIBCO/BRL) were used to extend the sequence. If more than one extension is necessary or desired, additional sets of primers are designed to further extend the known region.

[0220] High fidelity amplification was obtained by following the instructions for the XL-PCR kit (Perkin Elmer) and thoroughly mixing the enzyme and reaction mix. PCR was performed using the Peltier PTC200 thermal cycler (M.J. Research, Watertown, Mass.), beginning with 40 pmol of each primer and the recommended concentrations of all other components of the kit, with the following parameters:

Step 1	94° C. for 1 min (initial denaturation)
Step 2	65° C. for 1 min
Step 3	68° C. for 6 min
Step 4	94° C. for 15 sec
Step 5	65° C. for 1 min
Step 6	68° C. for 7 min
Step 7	Repeat steps 4 through 6 for an additional 15 cycles
Step 8	94° C. for 15 sec
Step 9	65° C. for 1 min
Step 10	68° C. for 7:15 min
Step 11	Repeat steps 8 through 10 for an additional 12 cycles
Step 12	72° C. for 8 min
Step 13	4° C. (and holding)

[0221] A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a low concentration (about 0.6% to 0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Bands thought to contain the largest products were excised from the gel, purified using QIAQUICK (QIAGEN Inc.), and trimmed of overhangs using Klenow enzyme to facilitate religation and cloning.

[0222] After ethanol precipitation, the products were redissolved in 13 μ l of ligation buffer, 1 μ l T4-DNA ligase (15 units) and 1 μ l T4 polynucleotide kinase were added, and the mixture was incubated at room temperature for 2 to 3 hours, or overnight at 16° C. Competent *E. coli* cells (in 40 μ l of appropriate media) were transformed with 3 μ l of ligation mixture and cultured in 80 μ l of SOC medium (See, e.g., Sambrook, supra, Appendix A, p. 2.) After incubation for one hour at 37° C., the *E. coli* mixture was plated on Luria Bertani (LB) agar (See, e.g., Sambrook, supra, Appendix A, p. 1) containing carbenicillin (2 \times carb). The following day, several colonies were randomly picked from each plate and cultured in 150 μ l of liquid LB/2 \times Carb medium placed in an individual well of an appropriate commercially-available sterile 96-well microtiter plate. The following day, 5 μ l of each overnight culture was transferred into a non-sterile 96-well plate and, after dilution 1:10 with water, 5 μ l from each sample was transferred into a PCR array.

[0223] For PCR amplification, 18 μ l of concentrated PCR reaction mix (3.3 \times) containing 4 units of rTth DNA polymerase, a vector primer, and one or both of the gene specific primers used for the extension reaction were added to each well. Amplification was performed using the following conditions:

Step 1	94° C. for 60 sec
Step 2	94° C. for 20 sec
Step 3	55° C. for 30 sec
Step 4	72° C. for 90 sec
Step 5	Repeat steps 2 through 4 for an additional 29 cycles
Step 6	72° C. for 180 sec
Step 7	4° C. (and holding)

[0224] Aliquots of the PCR reactions were run on agarose gels together with molecular weight markers. The sizes of the PCR products were compared to the original partial cDNAs, and appropriate clones were selected, ligated into plasmid, and sequenced.

[0225] In like manner, the nucleotide sequences of SEQ ID NO:2 and SEQ ID NO:4 are used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for 5' extension, and an appropriate genomic library.

[0226] VI. Labeling and Use of Individual Hybridization Probes

[0227] Hybridization probes derived from SEQ ID NO:2 and SEQ ID NO:4 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ -³²P] adenosine triphosphate (Amersham, Chicago, Ill.), and T4 polynucleotide kinase (DuPont NEN, Boston, Mass.). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine resin column (Pharmacia & Upjohn, Kalamazoo, Mich.). An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the

following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN, Boston, Mass.).

[0228] The DNA from each digest is fractionated on a 0.7 percent agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham, N.H.). Hybridization is carried out for 16 hours at 40° C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT AR film (Kodak, Rochester, N.Y.) is exposed to the blots for several hours, hybridization patterns are compared visually.

[0229] VII. Microarrays

[0230] A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, supra.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

[0231] Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE. Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; and Shalon, D. et al. (1996) *Genome Res.* 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

[0232] VIII. Complementary Polynucleotides

[0233] Sequences complementary to the D6DH-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring D6DH. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using Oligo 4.06 software and the coding sequence of D6DH. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the D6DH-encoding transcript.

[0234] IX. Expression of D6DH

[0235] Expression of D6DH is accomplished by subcloning the cDNA into an appropriate vector and transforming

the vector into host cells. This vector contains an appropriate promoter, e.g., β -galactosidase, upstream of the cloning site, operably associated with the cDNA of interest. (See, e.g., Sambrook, supra, pp. 404-433; and Rosenberg, M. et al. (1983) *Methods Enzymol.* 101:123-138.)

[0236] Induction of an isolated, transformed bacterial strain with isopropyl beta-D-thiogalactopyranoside (IPTG) using standard methods produces a fusion protein which consists of the first 8 residues of β -galactosidase, about 5 to 15 residues of linker, and the full length protein. The signal residues direct the secretion of D6DH into bacterial growth media which can be used directly in the following assay for activity.

[0237] X. Demonstration of D6DH Activity

[0238] D6DH activity is demonstrated by the method of Ivanetich, et al. (1996; *Biochim. Biophys. Acta* 1292:120-132), which measures the desaturation of linoleic acid to gamma linolenoyl CoA. The method uses HPLC for separation of the fatty acid substrate and product, exhibits a lower coefficient of variation (0.3%) than the reported TLC method (3.5%) and avoids the step of methylation of the saponified fatty acid substrate and product. The sample containing D6DH is incubated with 7.5 mM ATP, 1 mM CoA, 2.6 mM NADH, 40 mM KF, 0.33 mM nicotinamide, and 11.5 μ g/ μ g added fatty acid, at 35° C. with shaking at 60 cycles/min for times varying from 1 to 10 min. Standard concentrations of microsomal protein and substrate are 0.5 mg microsomal protein/ml and [1-¹⁴C]linoleic acid (0.45-10.9 nmol, 26-632 nCi). Product formation should be linear with time at the above protein concentrations. The fatty acid substrate and product of the delta-6 desaturation reaction are separated and quantified by HPLC separation of free fatty acids by a modification of the method of Aveldano et al. (1983; *J. Lipid Res.* 24:83-91). The desaturase reactions are terminated with 1 ml 10% KOH in methanol containing 0.005% butylated hydroxytoluene. Reaction mixtures, containing 1 mg of the relevant fatty acid carriers, are saponified for 30 min at 60° C. under Ar, acidified and extracted three times with 2 ml hexane. The hexane extract is dried under N at 45° C., fatty acids are dissolved in 0.5 ml ETOH and the solution passed through a 0.45 μ m filter prior to HPLC. The fatty acid substrate and product (50 μ l) are separated on an HPLC column. Fractions of 2 ml are collected and counted. After each run, the column is washed with acetonitrile. Delta-6 desaturase activity is calculated from the ratio: dpm product/(dpm product+dpm substrate).

[0239] XI. Production of D6DH Specific Antibodies

[0240] D6DH substantially purified using PAGE electrophoresis (see, e.g., Harrington, M. G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

[0241] Alternatively, the D6DH amino acid sequence is analyzed using LASERGENE software (DNASTAR Inc.) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel supra, ch. 11.)

[0242] Typically, oligopeptides 15 residues in length are synthesized using an Applied Biosystems peptide synthe-

-continued

Ala	Phe	Leu	Leu	Gly	His	Ile	Leu	Ala	Met	Glu	Val	Leu	Ala	Trp	140	145	150
Leu	Leu	Ile	Tyr	Leu	Leu	Gly	Pro	Gly	Trp	Val	Pro	Ser	Ala	Leu	155	160	165
Ala	Ala	Phe	Ile	Leu	Ala	Ile	Ser	Gln	Ala	Gln	Ser	Trp	Cys	Leu	170	175	180
Gln	His	Asp	Leu	Gly	His	Ala	Ser	Ile	Phe	Lys	Lys	Ser	Trp	Trp	185	190	195
Asn	His	Val	Ala	Gln	Lys	Phe	Val	Met	Gly	Gln	Leu	Lys	Gly	Phe	200	205	210
Ser	Ala	His	Trp	Trp	Asn	Phe	Arg	His	Phe	Gln	His	His	Ala	Lys	215	220	225
Pro	Asn	Ile	Phe	His	Lys	Asp	Pro	Asp	Val	Thr	Val	Ala	Pro	Val	230	235	240
Phe	Leu	Leu	Gly	Glu	Ser	Ser	Val	Glu	Tyr	Gly	Lys	Lys	Lys	Arg	245	250	255
Arg	Tyr	Leu	Pro	Tyr	Asn	Gln	Gln	His	Leu	Tyr	Phe	Phe	Leu	Ile	260	265	270
Gly	Pro	Pro	Leu	Leu	Thr	Leu	Val	Asn	Phe	Glu	Val	Glu	Asn	Leu	275	280	285
Ala	Tyr	Met	Leu	Val	Cys	Met	Gln	Trp	Ala	Asp	Leu	Leu	Trp	Ala	290	295	300
Ala	Ser	Phe	Tyr	Ala	Arg	Phe	Phe	Leu	Ser	Tyr	Leu	Pro	Phe	Tyr	305	310	315
Gly	Val	Pro	Gly	Val	Leu	Leu	Phe	Phe	Val	Ala	Val	Arg	Val	Leu	320	325	330
Glu	Ser	His	Trp	Phe	Val	Trp	Ile	Thr	Gln	Met	Asn	His	Ile	Pro	335	340	345
Lys	Glu	Ile	Gly	His	Glu	Lys	His	Arg	Asp	Trp	Val	Ser	Ser	Gln	350	355	360
Leu	Ala	Ala	Thr	Cys	Asn	Val	Glu	Pro	Ser	Leu	Phe	Thr	Asn	Trp	365	370	375
Phe	Ser	Gly	His	Leu	Asn	Phe	Gln	Ile	Glu	His	His	Leu	Phe	Pro	380	385	390
Arg	Met	Pro	Arg	His	Asn	Tyr	Ser	Arg	Val	Ala	Pro	Leu	Val	Lys	395	400	405
Ser	Leu	Cys	Ala	Lys	His	Gly	Leu	Ser	Tyr	Glu	Val	Lys	Pro	Phe	410	415	420
Leu	Thr	Ala	Leu	Val	Asp	Ile	Val	Arg	Ser	Leu	Lys	Lys	Ser	Gly	425	430	435
Asp	Ile	Trp	Leu	Asp	Ala	Tyr	Leu	His	Gln						440	445	

<210> SEQ ID NO 2

<211> LENGTH: 1717

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<223> OTHER INFORMATION: Incyte ID No: 2451043CB1

<400> SEQUENCE: 2

cgctcttcgc ttcctcggg gtattgctcg gacctcgcc accgcctggg atccccagga 60

-continued

```

ctcgtgcgtg cagcatgggc ggcgtcgggg agccgggacc gcgggagggg cccgcgcagc 120
cgggggcacc gctgcccacc ttctgctggg agcagatccg cgcgcacgac cagcccggcg 180
acaagtggct ggtcatcgag cgcgcgtct acgacatcag ccgctgggca cagcggcacc 240
cagggggcag ccgcctcacc gccaccacg gcgctgagga cgcacaggat gccttcctgt 300
ccttccatca agatctcaat ttgtgcgca agttcctaca gccctggtg attggagagc 360
tggctccgga agaaccacgc caggatggac cctggaatgc gcagctggtc gaggacttcc 420
gagccctgca ccaggcagcc gaggacatga agctgtttga tgccagtccc accttctttg 480
ctttcctact gggccacatc ctggccatgg aggtgctggc ctggctcctt atctacctcc 540
tgggtccctg ctgggtgccc agtgcctgg cgccttcat cctggccatc tctcaggctc 600
agtcctgggt tctgcagcat gaactgggcc atgcctccat ctcaagaag tcttggtgga 660
accacgtggc ccagaagtgc gtgatggggc agctaaaggg cttctccgcc cactggtgga 720
acttccgcca cttccagcac cagccaagc ccaacatctt ccacaaagac ccagacgtga 780
cggtgccgcc cgtcttcctc ctgggggagt catcgcgca gtatggcaag aagaaacgca 840
gatacctacc ctacaaccag cagcacctgt acttcttctt gatogggccg ccgctgctca 900
ccctggtgaa ctttgaagtg gaaaatctgg cgtacatgct ggtgtgcatg cagtgggagg 960
atgtgctctg ggccgccagc ttctatgccc gotttctt atcctaactc ccttctacg 1020
gcgtccctgg ggtgctgctc ttctttgttg ctgacagggt cctggaagc cactggttcg 1080
tgtggatcac acagatgaac cacatcccca aggagatcgg ccacgagaag caccgggact 1140
gggtcagctc tcagctggca gccacctgca acgtggagcc ctactttc accaactggt 1200
tcagcgggca cctcaacttc cagatcgagc accacctctt cccaggatg ccgagacaca 1260
actacagccg ggtggccccc ctggtcaagt cgctgtgtgc caagcacggc ctcagctacg 1320
aagtgaagcc cttcctcacc gcgctggtgg acatcgctcag gtcctgaag aagtctggtg 1380
acatctgctg ggacgcctac ctccatcagt gaaggcaaca cccaggcggg cagagaaggg 1440
ctcagggcac cagcaacca gcccagcccc ggccgggatc atacccccac cctccactg 1500
gccagcctgg ggtgcccctg cctgcctccc tggtagctgt gtcttccct cggccccctc 1560
acatgtgtat tcagcagccc tatggccttg gctctgggcc tgatgggaca ggggtagagg 1620
gaaggtgagc atagcacatt ttctagagc gagaattggg gaaaagctgt tattttata 1680
ttaaataca ttcagatgta aaaaaaaaa aaaaaaag 1717

```

```

<210> SEQ ID NO 3
<211> LENGTH: 444
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: Incyte ID No: 2056310CD1

```

<400> SEQUENCE: 3

```

Met Ala Pro Asp Pro Val Ala Ala Glu Thr Ala Ala Gln Gly Pro
 1             5             10             15
Thr Pro Arg Tyr Phe Thr Trp Asp Glu Val Ala Gln Arg Ser Gly
                20             25             30
Cys Glu Glu Arg Trp Leu Val Ile Asp Arg Lys Val Tyr Asn Ile
                35             40             45

```

-continued

Ser	Glu	Phe	Thr	Arg	Arg	His	Pro	Gly	Gly	Ser	Arg	Val	Ile	Ser	50	55	60
His	Tyr	Ala	Gly	Gln	Asp	Ala	Thr	Asp	Pro	Phe	Val	Ala	Phe	His	65	70	75
Ile	Asn	Lys	Gly	Leu	Val	Lys	Lys	Tyr	Met	Asn	Ser	Leu	Leu	Ile	80	85	90
Gly	Glu	Leu	Ser	Pro	Glu	Gln	Pro	Ser	Phe	Glu	Pro	Thr	Lys	Asn	95	100	105
Lys	Glu	Leu	Thr	Asp	Glu	Phe	Arg	Glu	Leu	Arg	Ala	Thr	Val	Glu	110	115	120
Arg	Met	Gly	Leu	Met	Lys	Ala	Asn	His	Val	Phe	Phe	Leu	Leu	Tyr	125	130	135
Leu	Leu	His	Ile	Leu	Leu	Leu	Asp	Gly	Ala	Ala	Trp	Leu	Thr	Leu	140	145	150
Trp	Val	Phe	Gly	Thr	Ser	Phe	Leu	Pro	Phe	Leu	Leu	Cys	Ala	Val	155	160	165
Leu	Leu	Ser	Ala	Val	Gln	Ala	Gln	Ala	Gly	Trp	Leu	Gln	His	Asp	170	175	180
Phe	Gly	His	Leu	Ser	Val	Phe	Ser	Thr	Ser	Lys	Trp	Asn	His	Leu	185	190	195
Leu	His	His	Phe	Val	Ile	Gly	His	Leu	Lys	Gly	Ala	Pro	Ala	Ser	200	205	210
Trp	Trp	Asn	His	Met	His	Phe	Gln	His	His	Ala	Lys	Pro	Asn	Cys	215	220	225
Phe	Arg	Lys	Asp	Pro	Asp	Ile	Asn	Met	His	Pro	Phe	Ser	Phe	Ala	230	235	240
Leu	Gly	Lys	Ile	Leu	Ser	Val	Glu	Leu	Gly	Lys	Gln	Lys	Lys	Lys	245	250	255
Tyr	Met	Pro	Tyr	Asn	His	Gln	His	Lys	Tyr	Phe	Phe	Leu	Ile	Gly	260	265	270
Pro	Pro	Ala	Leu	Leu	Pro	Leu	Tyr	Phe	Gln	Trp	Tyr	Ile	Phe	Tyr	275	280	285
Phe	Val	Ile	Gln	Arg	Lys	Lys	Trp	Val	Asp	Leu	Ala	Trp	Met	Ile	290	295	300
Thr	Phe	Tyr	Val	Arg	Phe	Phe	Leu	Thr	Tyr	Val	Pro	Leu	Leu	Gly	305	310	315
Leu	Lys	Ala	Phe	Leu	Gly	Leu	Phe	Phe	Ile	Val	Arg	Phe	Leu	Glu	320	325	330
Ser	Asn	Trp	Phe	Val	Trp	Val	Thr	Gln	Met	Asn	His	Ile	Pro	Met	335	340	345
His	Ile	Asp	His	Asp	Arg	Asn	Met	Asp	Trp	Val	Ser	Thr	Gln	Leu	350	355	360
Gln	Ala	Thr	Cys	Asn	Val	His	Lys	Ser	Ala	Phe	Asn	Asp	Trp	Phe	365	370	375
Ser	Gly	His	Leu	Asn	Phe	Gln	Ile	Glu	His	His	Leu	Phe	Pro	Thr	380	385	390
Met	Pro	Arg	His	Asn	Tyr	His	Lys	Val	Ala	Pro	Leu	Val	Gln	Ser	395	400	405
Leu	Cys	Ala	Lys	His	Gly	Ile	Glu	Tyr	Gln	Ser	Lys	Pro	Leu	Leu	410	415	420
Ser	Ala	Phe	Ala	Asp	Ile	Ile	His	Ser	Leu	Lys	Glu	Ser	Gly	Gln			

-continued

	425	430	435			
Leu Trp Leu Asp Ala Tyr Leu His Gln						
	440					
 <210> SEQ ID NO 4						
<211> LENGTH: 1928						
<212> TYPE: DNA						
<213> ORGANISM: Homo sapiens						
<220> FEATURE:						
<221> NAME/KEY: misc_feature						
<223> OTHER INFORMATION: Incyte ID No: 2056310CB1						
 <400> SEQUENCE: 4						
aaatccactc	ctggagcccg	cggaccccca	gcacgcgcct	gacagcccct	gctggcccgg	60
cgcgccggct	cgccaggcca	gctatggccc	cggacccggt	ggcgcgcgag	accgcggctc	120
agggacctac	cccgcgctac	ttcacctggg	acgaggtggc	ccagcgctca	gggtgcgagg	180
agcggtggtc	agtgatcgac	cgtaaggtgt	acaacatcag	cgagttcacc	cgccggcatc	240
caggggggct	ccgggtcadc	agccactacg	ccgggcagga	tgccacggat	ccctttgtgg	300
ccttccacat	caacaagggc	cttgtgaaga	agtatatgaa	ctctctcctg	attggagaac	360
tgctcccaga	gcagcccagc	tttgagccca	ccaagaataa	agagctgaca	gatgagttcc	420
gggagctgog	ggccacagtg	gagcggatgg	ggctcatgaa	ggccaacct	gtctttctcc	480
tgctgtacct	gctgcacatc	ttgctgctgg	atggtgcagc	ctggctcacc	ctttgggtct	540
ttgggacgtc	ctttttgccc	ttcctcctct	gtgcggtgct	gctcagtgca	gttcaggccc	600
aggctgggct	gctgcagcat	gactttgggc	acctgtcggg	cttcagcacc	tcaaagtggg	660
acctctgctc	acatcatttt	gtgattggcc	acctgaaggg	ggccccgcc	agttgggtgg	720
accacatgca	cttccagcac	catgccaagc	ccaactgctt	ccgcaaagac	ccagacatca	780
acatgcatcc	cttctccttt	gccttgggga	agatcctctc	tgtggagctt	gggaaacaga	840
agaaaaata	tatgccgtac	aaccaccagc	acaaatactt	cttctcaatt	gggccccag	900
ccttgctgoc	tctctacttc	cagtggtata	ttttctat	tgttatccag	cgaaagaagt	960
gggtggactt	ggcctggatg	attaccttct	acgtccgctt	cttctcact	tatgtgccac	1020
tattgggctc	gaaagccttc	ctgggccttt	tcttcatagt	caggttcctg	gaaagcaact	1080
ggtttgtgtg	ggtgacacag	atgaaccata	ttcccatgca	cattgatcat	gaccggaaca	1140
tggactgggt	ttccaccagc	ctccaggcca	catgcaatgt	ccacaagtct	gccttcaatg	1200
actggttacc	tggacacctc	aacttccaga	ttgagcacca	tctttttccc	acgatgcctc	1260
gacacaatta	ccacaaagtg	gctcccctgg	tgcagtcctt	gtgtgccaag	catggcatag	1320
agtaccagtc	caagcccctg	ctgtcagcct	tcgccgacat	catccactca	ctaaaggagt	1380
cagggcagct	ctggctagat	gcctatcttc	accaataaca	acagccacco	tgcccagtct	1440
ggaagaagag	gaggaagact	ctggagccaa	ggcagagggg	agcttgaggg	acaatgccac	1500
tatagtttaa	tactcagagg	gggttgggtt	tggggacata	aagcctctga	ctcaaacctc	1560
tcccttttat	cttctagcca	cagttctaag	acccaaagtg	gggggtggac	acagaagtcc	1620
ctaggagggg	aggagctggt	ggggcagggg	tgtaaattat	ttcctttttc	tagtttgcca	1680
catgcaggta	gttggtgaac	agagagaacc	aggagggtaa	cagaagagga	gggacctact	1740
gaaccacag	tcaggaagag	atttaacact	aaaattccac	tcatgccggg	cgtggtggca	1800

-continued

cgcgccctgta atcccagcta cccaggagggc tgaggcagga gaatcgcttg aaccggggag	1860
gtggaggttg cagtgaagctg agatcacgcc attgtactcg cagcgctggg cagacagagc	1920
acagctcc	1928

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:

- a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3,
- b) a polypeptide comprising a naturally occurring amino acid sequence at least 80% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3,
- c) a fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3, wherein the fragment has fatty acid desaturase activity, and
- d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3.

2. An isolated polypeptide of claim 1, comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3.

3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide encoding a polypeptide of claim 2.

5. An isolated polynucleotide of claim 4, comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:4.

6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

7. A cell transformed with a recombinant polynucleotide of claim 6.

8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

9. A method of producing a polypeptide of claim 1, the method comprising:

- a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

b) recovering the polypeptide so expressed.

10. A method of claim 9, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3.

11. An isolated antibody which specifically binds to a polypeptide of claim 1.

12. An isolated polynucleotide selected from the group consisting of:

- a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:4,

b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:4,

c) a polynucleotide complementary to a polynucleotide of a),

d) a polynucleotide complementary to a polynucleotide of b), and

e) an RNA equivalent of a)-d).

13. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 12.

14. A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 12, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and

b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

15. A method of claim 14, wherein the probe comprises at least 60 contiguous nucleotides.

16. A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 12, the method comprising:

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and

b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

17. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

18. A composition of claim 17, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3.

19. A method for treating a disease or condition associated with decreased expression of functional D6DH, comprising administering to a patient in need of such treatment the composition of claim 17.

20. A method of screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and

b) detecting agonist activity in the sample.

21. A composition comprising an agonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

22. A method for treating a disease or condition associated with decreased expression of functional D6DH, comprising administering to a patient in need of such treatment a composition of claim 21.

23. A method of screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

24. A composition comprising an antagonist compound identified by a method of claim 23 and a pharmaceutically acceptable excipient.

25. A method for treating a disease or condition associated with overexpression of functional D6DH, comprising administering to a patient in need of such treatment a composition of claim 24.

26. A method of screening for a compound that specifically binds to the polypeptide of claim 1, the method comprising:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and
- b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

27. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, the method comprising:

- a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

28. A method of screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
- b) detecting altered expression of the target polynucleotide, and
- c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

29. A method of assessing toxicity of a test compound, the method comprising:

- a) treating a biological sample containing nucleic acids with the test compound,

- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 12 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 12 or fragment thereof,

- c) quantifying the amount of hybridization complex, and

- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

30. A method for a diagnostic test for a condition or disease associated with the expression of D6DH in a biological sample, the method comprising:

- a) combining the biological sample with an antibody of claim 11, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex, and

- b) detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.

31. The antibody of claim 11, wherein the antibody is:

- a) a chimeric antibody,
- b) a single chain antibody,
- c) a Fab fragment,
- d) a F(ab')₂ fragment, or
- e) a humanized antibody.

32. A composition comprising an antibody of claim 11 and an acceptable excipient.

33. A method of diagnosing a condition or disease associated with the expression of D6DH in a subject, comprising administering to said subject an effective amount of the composition of claim 32.

34. A composition of claim 32, wherein the antibody is labeled.

35. A method of diagnosing a condition or disease associated with the expression of D6DH in a subject, comprising administering to said subject an effective amount of the composition of claim 34.

36. A method of preparing a polyclonal antibody which specifically binds to the polypeptide of claim 1, the method comprising:

- a) immunizing an animal with a polypeptide consisting of an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3, or an immunogenic fragment thereof, under conditions to elicit an antibody response,

- b) isolating antibodies from the animal, and

- c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which specifically binds to the polypeptide of claim 1.

37. A polyclonal antibody produced by a method of claim 36.

38. A composition comprising the polyclonal antibody of claim 37 and a suitable carrier.

39. A method of making a monoclonal antibody which specifically binds to the polypeptide of claim 1, the method comprising:

- a) immunizing an animal with an antigen polypeptide consisting of an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3, or an immunogenic fragment thereof, under conditions to elicit an antibody response,
- b) isolating antibody producing cells from the animal,
- c) fusing the antibody producing cells with immortalized cells to form monoclonal antibody-producing hybridoma cells,
- d) culturing the hybridoma cells, and
- e) isolating from the culture monoclonal antibody which specifically binds to the polypeptide of claim 1.

40. A monoclonal antibody produced by a method of claim 39.

41. A composition comprising the monoclonal antibody of claim 40 and a suitable carrier.

42. The antibody of claim 11, wherein the antibody is produced by screening a Fab expression library.

43. The antibody of claim 11, wherein the antibody is produced by screening a recombinant immunoglobulin library.

44. A method of detecting a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3 in a sample, the method comprising:

- a) incubating the antibody of claim 11 with the sample under conditions to allow specific binding of the antibody and the polypeptide, and
- b) detecting specific binding, wherein specific binding indicates the presence of a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3 in the sample.

45. A method of purifying a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3 from a sample, the method comprising:

- a) incubating the antibody of claim 11 with the sample under conditions to allow specific binding of the antibody and the polypeptide, and
- b) separating the antibody from the sample and obtaining the purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3.

46. A microarray wherein at least one element of the microarray is a polynucleotide of claim 13.

47. A method of generating an expression profile of a sample which contains polynucleotides, the method comprising:

- a) labeling the polynucleotides of the sample,
- b) contacting the elements of the microarray of claim 46 with the labeled polynucleotides of the sample under conditions suitable for the formation of a hybridization complex, and
- c) quantifying the expression of the polynucleotides in the sample.

48. An array comprising different nucleotide molecules affixed in distinct physical locations on a solid substrate, wherein at least one of said nucleotide molecules comprises a first oligonucleotide or polynucleotide sequence specifically hybridizable with at least 30 contiguous nucleotides of a target polynucleotide, and wherein said target polynucleotide is a polynucleotide of claim 12.

49. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to at least 30 contiguous nucleotides of said target polynucleotide.

50. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to at least 60 contiguous nucleotides of said target polynucleotide.

51. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to said target polynucleotide.

52. An array of claim 48, which is a microarray.

53. An array of claim 48, further comprising said target polynucleotide hybridized to a nucleotide molecule comprising said first oligonucleotide or polynucleotide sequence.

54. A method of claim 48, wherein a linker joins at least one of said nucleotide molecules to said solid substrate.

55. An array of claim 48, wherein each distinct physical location on the substrate contains multiple nucleotide molecules, and the multiple nucleotide molecules at any single distinct physical location have the same sequence, and each distinct physical location on the substrate contains nucleotide molecules having a sequence which differs from the sequence of nucleotide molecules at another distinct physical location on the substrate.

56. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

57. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.

58. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:2.

59. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:4.

60. A polypeptide of claim 1, comprising a naturally occurring amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:1.

61. A polypeptide of claim 1, comprising a naturally occurring amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO:3.

* * * * *