



US006716587B2

(12) **United States Patent**
Mosley et al.

(10) **Patent No.:** **US 6,716,587 B2**
(45) **Date of Patent:** **Apr. 6, 2004**

(54) **ANTIBODIES TO INTERLEUKIN-4
RECEPTORS AND USES THEREOF**

(75) Inventors: **Bruce A. Mosley**, Seattle, WA (US);
David J. Cosman, Bainbridge Island,
WA (US); **Linda S. Park**, Seattle, WA
(US); **M. Patricia Beckmann**,
Hansville, WA (US); **Carl J. March**,
Bainbridge Island, WA (US); **Rejean
Idzerda**, Seattle, WA (US)

(73) Assignee: **Immunex Corporation**, Seattle, WA
(US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/104,590**

(22) Filed: **Mar. 22, 2002**

(65) **Prior Publication Data**

US 2003/0185821 A1 Oct. 2, 2003

Related U.S. Application Data

(60) Division of application No. 09/094,917, filed on Jun. 15,
1998, now Pat. No. 6,391,581, which is a continuation of
application No. 07/480,694, filed on Feb. 14, 1990, now Pat.
No. 5,840,869, which is a continuation-in-part of application
No. 07/370,924, filed on Jun. 23, 1989, now abandoned,
which is a continuation-in-part of application No. 07/326,
156, filed on Mar. 20, 1989, now abandoned, which is a
continuation-in-part of application No. 07/319,438, filed on
Mar. 2, 1989, now abandoned, which is a continuation-in-
part of application No. 07/265,047, filed on Oct. 31, 1988,
now abandoned.

(51) **Int. Cl.⁷** **C01N 33/53**; C07K 76/00;
A61K 39/395

(52) **U.S. Cl.** **435/7.1**; 530/387.1; 530/388.1;
530/388.22; 424/143.1; 424/141.1

(58) **Field of Search** 435/7.1; 530/387.1,
530/388.1, 388.22; 424/143.1, 141.1

(56) **References Cited**

PUBLICATIONS

Larche M et al. A novel T-lymphocyte molecule that may
function in the induction of self-tolerance and MHC-re-
striction within th human thymic microenvironment. Immu-
nology. May 1988;64(1):101-5.*

Larche M et al. Functional evidence for a monoclonal
antibody that binds to the human IL-4 receptor. Immunol-
ogy. 1988 Dec; 65(4):617-22.*

* cited by examiner

Primary Examiner—Yvonne Eyler

Assistant Examiner—Joseph F. Murphy

(74) *Attorney, Agent, or Firm*—Kathryn A. Anderson

(57) **ABSTRACT**

Mammalian Interleukin-4 receptor proteins, DNAs and
expression vectors encoding mammalian IL-4 receptors, and
processes for producing mammalian IL-4 receptors as prod-
ucts of cell culture, are disclosed. A method for suppressing
an IL-4-dependent immune or inflammatory response in a
mammal, including a human, by administering an effective
amount of soluble IL-4 receptor (sIL-4R) and a suitable
diluent or carrier.

8 Claims, 22 Drawing Sheets

FIGURE 1




FIGURE 2A

ATG	GGG	CGG	CTT	TGC	ACC	AAG	TTC	CTG	ACC	TCT	GTG	GGC	TGT	CTG	-31
Met	Gly	Arg	Leu	Cys	Thr	Lys	Phe	Leu	Thr	Ser	Val	Gly	Cys	Leu	-11
ATT	TTG	CTG	TTG	GTG	ACT	GGA	TCT	GGG	AGC	ATC	AAG	GTC	CTG	GGT	15
Ile	Leu	Leu	Leu	Val	Thr	Gly	Ser	Gly	Ser	<u>Ile</u>	Lys	Val	Leu	Gly	5
GAG	CCC	ACC	TGC	TTC	TCT	GAC	TAC	ATC	CGC	ACT	TCC	ACG	TGT	GAG	60
Glu	Pro	Thr	Cys	Phe	Ser	Asp	Tyr	Ile	Arg	Thr	Ser	Thr	Cys	Glu	20
TGG	TTC	CTG	GAT	AGC	GCT	GTG	GAC	TGC	AGT	TCT	CAG	CTC	TGC	CTA	105
Trp	Phe	Leu	Asp	Ser	Ala	Val	Asp	Cys	Ser	Ser	Gln	Leu	Cys	Leu	35
CAC	TAC	AGG	CTG	ATG	TTC	TTC	GAG	TTC	TCT	GAA	AAC	CTC	ACA	TGC	150
His	Tyr	Arg	Leu	Met	Phe	Phe	Glu	Phe	Ser	Glu	Asn	Leu	Thr	Cys	50
ATC	CCG	AGG	AAC	AGT	GCC	AGC	ACT	GTG	TGT	GTG	TGC	CAC	ATG	GAA	195
Ile	Pro	Arg	Asn	Ser	Ala	Ser	Thr	Val	Cys	Val	Cys	His	Met	Glu	65
ATG	AAT	AGG	CCG	GTC	CAA	TCA	GAC	AGA	TAC	CAG	ATG	GAA	CTG	TGG	240
Met	Asn	Arg	Pro	Val	Gln	Ser	Asp	Arg	Tyr	Gln	Met	Glu	Leu	Trp	80
GCT	GAG	CAC	AGA	CAG	CTG	TGG	CAG	GGC	TCC	TTC	AGC	CCC	AGT	GGT	285
Ala	Glu	His	Arg	Gln	Leu	Trp	Gln	Gly	Ser	Phe	Ser	Pro	Ser	Gly	95
AAT	GTG	AAG	CCC	CTA	GCT	CCA	GAC	AAC	CTC	ACA	CTC	CAC	ACC	AAT	330
Asn	Val	Lys	Pro	Leu	Ala	Pro	Asp	Asn	Leu	Thr	Leu	His	Thr	Asn	110
GTG	TCC	GAC	GAA	TGG	CTG	CTG	ACC	TGG	AAT	AAC	CTG	TAC	CCA	TCG	375
Val	Ser	Asp	Glu	Trp	Leu	Leu	Thr	Trp	Asn	Asn	Leu	Tyr	Pro	Ser	125
AAC	AAC	TTA	CTG	TAC	AAA	GAC	CTC	ATC	TCC	ATG	GTC	AAC	ATC	TCC	420
Asn	Asn	Leu	Leu	Tyr	Lys	Asp	Leu	Ile	Ser	Met	Val	Asn	Ile	Ser	140
AGA	GAG	GAC	AAC	CCT	GCA	GAA	TTC	ATA	GTC	TAT	AAT	GTG	ACC	TAC	465
Arg	Glu	Asp	Asn	Pro	Ala	Glu	Phe	Ile	Val	Tyr	Asn	Val	Thr	Tyr	155
AAG	GAA	CCC	AGG	CTG	AGC	TTC	CCG	ATC	AAC	ATC	CTG	ATG	TCA	GGG	510
Lys	Glu	Pro	Arg	Leu	Ser	Phe	Pro	Ile	Asn	Ile	Leu	Met	Ser	Gly	170
GTC	TAC	TAT	ACG	GCG	CGT	GTG	AGG	GTC	AGA	TCC	CAG	ATA	CTC	ACT	555
Val	Tyr	Tyr	Thr	Ala	Arg	Val	Arg	Val	Arg	Ser	Gln	Ile	Leu	Thr	185
GGC	ACC	TGG	AGT	GAG	TGG	AGT	CCT	AGC	ATC	ACG	TGG	TAC	AAC	CAC	600
Gly	Thr	Trp	Ser	Glu	Trp	Ser	Pro	Ser	Ile	Thr	Trp	Tyr	Asn	His	200
TTC	CAG	CTG	CCC	CTG	ATA	CAG	CGC	CTT	CCA	CTG	GGG	GTC	ACC	ATC	645
Phe	Gln	Leu	Pro	Leu	Ile	Gln	Arg	<u>Leu</u>	<u>Pro</u>	<u>Leu</u>	<u>Gly</u>	<u>Val</u>	<u>Thr</u>	<u>Ile</u>	215
TCC	TGC	CTC	TGC	ATC	CCG	TTG	TTT	TGC	CTG	TTC	TGT	TAC	TTC	AGC	690
<u>Ser</u>	<u>Cys</u>	<u>Leu</u>	<u>Cys</u>	<u>Ile</u>	<u>Pro</u>	<u>Leu</u>	<u>Phe</u>	<u>Cys</u>	<u>Leu</u>	<u>Phe</u>	<u>Cys</u>	<u>Tyr</u>	<u>Phe</u>	<u>Ser</u>	230
ATT	ACC	AAG	ATT	AAG	AAG	ATA	TGG	TGG	GAC	CAG	ATT	CCC	ACC	CCA	735
<u>Ile</u>	<u>Thr</u>	Lys	Ile	Lys	Lys	Ile	Trp	Trp	Asp	Gln	Ile	Pro	Thr	Pro	245

FIGURE 2B

GCA	CGC	AGT	CCC	TTG	GTG	GCC	ATC	ATC	ATT	CAG	GAT	GCA	CAG	GTG	780
Ala	Arg	Ser	Pro	Leu	Val	Ala	Ile	Ile	Ile	Gln	Asp	Ala	Gln	Val	260
CCC	CTC	TGG	GAT	AAG	CAG	ACC	CGA	AGC	CAG	GAG	TCA	ACC	AAG	TAC	825
Pro	Leu	Trp	Asp	Lys	Gln	Thr	Arg	Ser	Gln	Glu	Ser	Thr	Lys	Tyr	275
CCG	CAC	TGG	AAA	ACT	TGT	CTA	GAC	AAG	CTG	CTG	CCT	TGC	TTG	CTG	870
Pro	His	Trp	Lys	Thr	Cys	Leu	Asp	Lys	Leu	Leu	Pro	Cys	Leu	Leu	290
AAG	CAC	AGA	GTA	AAG	AAG	AAG	ACA	GAC	TTC	CCG	AAG	GCT	GCC	CCA	915
Lys	His	Arg	Val	Lys	Lys	Lys	Thr	Asp	Phe	Pro	Lys	Ala	Ala	Pro	305
ACC	AAG	TCT	CTC	CAG	AGT	CCT	GGA	AAG	GCA	GGC	TGG	TGT	CCC	ATG	960
Thr	Lys	Ser	Leu	Gln	Ser	Pro	Gly	Lys	Ala	Gly	Trp	Cys	Pro	Met	320
GAG	GTC	AGC	AGG	ACC	GTC	CTC	TGG	CCA	GAG	AAT	GTT	AGT	GTC	AGT	1005
Glu	Val	Ser	Arg	Thr	Val	Leu	Trp	Pro	Glu	Asn	Val	Ser	Val	Ser	335
GTG	GTG	CGC	TGT	ATG	GAG	CTG	TTT	GAG	GCC	CCA	GTA	CAG	AAT	GTG	1050
Val	Val	Arg	Cys	Met	Glu	Leu	Phe	Glu	Ala	Pro	Val	Gln	Asn	Val	350
GAG	GAG	GAA	GAA	GAT	GAG	ATA	GTC	AAA	GAG	GAC	CTG	AGC	ATG	TCA	1095
Glu	Glu	Glu	Glu	Asp	Glu	Ile	Val	Lys	Glu	Asp	Leu	Ser	Met	Ser	365
CCT	GAG	AAC	AGC	GGA	GGC	TGC	GGC	TTC	CAG	GAG	aGC	CAG	GCA	GAC	1140
Pro	Glu	Asn	Ser	Gly	Gly	Cys	Gly	Phe	Gln	Glu	Ser	Gln	Ala	Asp	380
ATC	ATG	GCT	CGG	CTC	ACT	GAG	AAC	CTG	TTT	TCC	GAC	TTG	TTG	GAG	1185
Ile	Met	Ala	Arg	Leu	Thr	Glu	Asn	Leu	Phe	Ser	Asp	Leu	Leu	Glu	395
GCT	GAG	AAT	GGG	GGC	CTT	GGC	CAG	TCA	GCC	TTG	GCA	GAG	TCA	TGC	1230
Ala	Glu	Asn	Gly	Gly	Leu	Gly	Gln	Ser	Ala	Leu	Ala	Glu	Ser	Cys	410
TCC	CCT	CTG	CCT	TCA	GGA	AGT	GGG	CAG	GCT	TCT	GTA	TCC	TGG	GCC	1275
Ser	Pro	Leu	Pro	Ser	Gly	Ser	Gly	Gln	Ala	Ser	Val	Ser	Trp	Ala	425
TGC	CTC	CCC	ATG	GGG	CCC	AGT	GAG	GAG	GCC	ACA	TGC	CAG	GTC	ACA	1320
Cys	Leu	Pro	Met	Gly	Pro	Ser	Glu	Glu	Ala	Thr	Cys	Gln	Val	Thr	440
GAG	CAG	CCT	TCA	CAC	CCA	GGC	CCT	CTT	TCA	GGC	AGC	CCA	GCC	CAG	1365
Glu	Gln	Pro	Ser	His	Pro	Gly	Pro	Leu	Ser	Gly	Ser	Pro	Ala	Gln	455
AGT	GCA	CCT	ACT	CTG	GCT	TGC	ACG	CAG	GTC	CCA	CTT	GTC	CTT	GCA	1410
Ser	Ala	Pro	Thr	Leu	Ala	Cys	Thr	Gln	Val	Pro	Leu	Val	Leu	Ala	470
GAC	AAT	CCT	GCC	TAC	CGG	AGT	TTT	AGT	GAC	TGC	TGT	AGC	CCG	GCC	1455
Asp	Asn	Pro	Ala	Tyr	Arg	Ser	Phe	Ser	Asp	Cys	Cys	Ser	Pro	Ala	485
CCA	AAT	CCT	GGA	GAG	CTG	GCT	CCA	GAG	CAG	CAG	CAG	GCT	GAT	CAT	1500
Pro	Asn	Pro	Gly	Glu	Leu	Ala	Pro	Glu	Gln	Gln	Gln	Ala	Asp	His	500
CTG	GAA	GAA	GAG	GAG	CCT	CCA	AGC	CCG	GCT	GAC	CCC	CAT	TCT	TCA	1545
Leu	Glu	Glu	Glu	Glu	Pro	Pro	Ser	Pro	Ala	Asp	Pro	His	Ser	Ser	515

FIGURE 2C

GGG	CCA	CCA	ATG	CAG	CCA	GTG	GAG	AGC	TGG	GAG	CAG	ATC	CTT	CAC	1590
Gly	Pro	Pro	Met	Gln	Pro	Val	Glu	Ser	Trp	Glu	Gln	Ile	Leu	His	530
ATG	AGT	GTC	CTG	CAG	CAT	GGG	GCA	GCT	GCT	GGC	TCC	ACC	CCA	GCC	1635
Met	Ser	Val	Leu	Gln	His	Gly	Ala	Ala	Ala	Gly	Ser	Thr	Pro	Ala	545
CCT	GCC	GGT	GGC	TAC	CAG	GAG	TTT	GTG	CAG	GCA	GTG	AAG	CAG	GGT	1680
Pro	Ala	Gly	Gly	Tyr	Gln	Glu	Phe	Val	Gln	Ala	Val	Lys	Gln	Gly	560
GCC	GCC	CAG	GAT	CCT	GGG	GTG	CCT	GGT	GTC	AGG	CCT	TCT	GGA	GAC	1725
Ala	Ala	Gln	Asp	Pro	Gly	Val	Pro	Gly	Val	Arg	Pro	Ser	Gly	Asp	575
CCC	GGT	TAC	AAG	GCC	TTC	TCG	AGC	CTG	CTC	AGC	AGC	AAT	GGC	ATC	1770
Pro	Gly	Tyr	Lys	Ala	Phe	Ser	Ser	Leu	Leu	Ser	Ser	Asn	Gly	Ile	590
CGC	GGG	GAC	ACA	GCA	GCA	GCG	GGG	ACT	GAC	GAT	GGG	CAT	GGA	GGC	1815
Arg	Gly	Asp	Thr	Ala	Ala	Ala	Gly	Thr	Asp	Asp	Gly	His	Gly	Gly	605
TAC	AAG	CCC	TTC	CAG	AAT	CCT	GTT	CCT	AAC	CAG	TCC	CCT	AGC	TCC	1860
Tyr	Lys	Pro	Phe	Gln	Asn	Pro	Val	Pro	Asn	Gln	Ser	Pro	Ser	Ser	620
GTG	CCC	TTA	TTT	ACT	TTC	GGA	CTA	GAC	ACG	GAG	CTG	TCA	CCC	AGT	1905
Val	Pro	Leu	Phe	Thr	Phe	Gly	Leu	Asp	Thr	Glu	Leu	Ser	Pro	Ser	635
CCT	CTG	AAC	TCA	GAC	CCA	CCC	AAA	AGC	CCC	CCA	GAA	TGC	CTT	GGT	1950
Pro	Leu	Asn	Ser	Asp	Pro	Pro	Lys	Ser	Pro	Pro	Glu	Cys	Leu	Gly	650
CTG	GAG	CTG	GGG	CTC	AAA	GGA	GGT	GAC	TGG	GTG	AAG	GCC	CCT	CCT	1995
Leu	Glu	Leu	Gly	Leu	Lys	Gly	Gly	Asp	Trp	Val	Lys	Ala	Pro	Pro	665
CCT	GCA	GAT	GAG	GTG	CCC	AAG	CCC	TTT	GGG	GAT	GAC	CTG	GGC	TTT	2040
Pro	Ala	Asp	Glu	Val	Pro	Lys	Pro	Phe	Gly	Asp	Asp	Leu	Gly	Phe	680
GGT	ATT	GTG	TAC	TCG	TCC	CTC	ACT	TGC	CAC	TTG	TGT	GGC	CAC	CTG	2085
Gly	Ile	Val	Tyr	Ser	Ser	Leu	Thr	Cys	His	Leu	Cys	Gly	His	Leu	695
AAG	CAA	CAC	CAC	AGC	CAG	GAG	GAA	GGT	GGC	CAG	AGC	CCC	ATC	GTT	2130
Lys	Gln	His	His	Ser	Gln	Glu	Glu	Gly	Gly	Gln	Ser	Pro	Ile	Val	710
GCT	AGC	CCT	GGC	TGT	GGC	TGC	TGC	TAC	GAT	GAC	AGA	TCA	CCA	TCC	2175
Ala	Ser	Pro	Gly	Cys	Gly	Cys	Cys	Tyr	Asp	Asp	Arg	Ser	Pro	Ser	725
CTG	GGG	AGC	CTC	TCG	GGG	GCC	TTG	GAA	AGC	TGT	CCT	GAG	GGA	ATA	2220
Leu	Gly	Ser	Leu	Ser	Gly	Ala	Leu	Glu	Ser	Cys	Pro	Glu	Gly	Ile	740
CCA	CCA	GAA	GCC	AAC	CTC	ATG	TCA	GCA	CCC	AAG	ACA	CCC	TCA	AAC	2265
Pro	Pro	Glu	Ala	Asn	Leu	Met	Ser	Ala	Pro	Lys	Thr	Pro	Ser	Asn	755
TTG	TCA	GGG	GAG	GGC	AAG	GGC	CCT	GGT	CAC	TCT	CCT	GTT	CCC	AGC	2310
Leu	Ser	Gly	Glu	Gly	Lys	Gly	Pro	Gly	His	Ser	Pro	Val	Pro	Ser	770
CAG	ACG	ACC	GAG	GTG	CCT	GTG	GGC	GCC	CTG	GGC	ATT	GCT	GTT	TCT	2355
Gln	Thr	Thr	Glu	Val	Pro	Val	Gly	Ala	Leu	Gly	Ile	Ala	Val	Ser	785

Figure 3




FIGURE 4A

ATG	GGG	TGG	CTT	TGC	TCT	GGG	CTC	CTG	TTC	CCT	GTG	AGC	TGC	CTG	-31
Met	Gly	Trp	Leu	Cys	Ser	Gly	Leu	Leu	Phe	Pro	Val	Ser	Cys	Leu	-11
GTC	CTG	CTG	CAG	GTG	GCA	AGC	TCT	GGG	AAC	ATG	AAG	GTC	TTG	CAG	15
Val	Leu	Leu	Gln	Val	Ala	Ser	Ser	Gly	Asn	Met	Lys	Val	Leu	Gln	5
GAG	CCC	ACC	TGC	GTC	TCC	GAC	TAC	ATG	AGC	ATC	TCT	ACT	TGC	GAG	60
Glu	Pro	Thr	Cys	Val	Ser	Asp	Tyr	Met	Ser	Ile	Ser	Thr	Cys	Glu	20
TGG	AAG	ATG	AAT	GGT	CCC	ACC	AAT	TGC	AGC	ACC	GAG	CTC	CGC	CTG	105
Trp	Lys	Met	Asn	Gly	Pro	Thr	Asn	Cys	Ser	Thr	Glu	Leu	Arg	Leu	35
TTG	TAC	CAG	CTG	GTT	TTT	CTG	CTC	TCC	GAA	GCC	CAC	ACG	TGT	ATC	150
Leu	Tyr	Gln	Leu	Val	Phe	Leu	Leu	Ser	Glu	Ala	His	Thr	Cys	Ile	50
CCT	GAG	AAC	AAC	GGA	GGC	GCG	GGG	TGC	GTG	TGC	CAC	CTG	CTC	ATG	195
Pro	Glu	Asn	Asn	Gly	Gly	Ala	Gly	Cys	Val	Cys	His	Leu	Leu	Met	65
GAT	GAC	GTG	GTC	AGT	GCG	GAT	AAC	TAT	ACA	CTG	GAC	CTG	TGG	GCT	240
Asp	Asp	Val	Val	Ser	Ala	Asp	Asn	Tyr	Thr	Leu	Asp	Leu	Trp	Ala	80
GGG	CAG	CAG	CTG	CTG	TGG	AAG	GGC	TCC	TTC	AAG	CCC	AGC	GAG	CAT	285
Gly	Gln	Gln	Leu	Leu	Trp	Lys	Gly	Ser	Phe	Lys	Pro	Ser	Glu	His	95
GTG	AAA	CCC	AGG	GCC	CCA	GGA	AAC	CTG	ACA	GTT	CAC	ACC	AAT	GTC	330
Val	Lys	Pro	Arg	Ala	Pro	Gly	Asn	Leu	Thr	Val	His	Thr	Asn	Val	110
TCC	GAC	ACT	CTG	CTG	CTG	ACC	TGG	AGC	AAC	CCG	TAT	CCC	CCT	GAC	375
Ser	Asp	Thr	Leu	Leu	Leu	Thr	Trp	Ser	Asn	Pro	Tyr	Pro	Pro	Asp	125
AAT	TAC	CTG	TAT	AAT	CAT	CTC	ACC	TAT	GCA	GTC	AAC	ATT	TGG	AGT	420
Asn	Tyr	Leu	Tyr	Asn	His	Leu	Thr	Tyr	Ala	Val	Asn	Ile	Trp	Ser	140
GAA	AAC	GAC	CCG	GCA	GAT	TTC	AGA	ATC	TAT	AAC	GTG	ACC	TAC	CTA	465
Glu	Asn	Asp	Pro	Ala	Asp	Phe	Arg	Ile	Tyr	Asn	Val	Thr	Tyr	Leu	155
GAA	CCC	TCC	CTC	CGC	ATC	GCA	GCC	AGC	ACC	CTG	AAG	TCT	GGG	ATT	510
Glu	Pro	Ser	Leu	Arg	Ile	Ala	Ala	Ser	Thr	Leu	Lys	Ser	Gly	Ile	170
TCC	TAC	AGG	GCA	CGG	GTG	AGG	GCC	TGG	GCT	CAG	TGC	TAT	AAC	ACC	555
Ser	Tyr	Arg	Ala	Arg	Val	Arg	Ala	Trp	Ala	Gln	Cys	Tyr	Asn	Thr	185
ACC	TGG	AGT	GAG	TGG	AGC	CCC	AGC	ACC	AAG	TGG	CAC	AAC	TCC	TAC	600
Thr	Trp	Ser	Glu	Trp	Ser	Pro	Ser	Thr	Lys	Trp	His	Asn	Ser	Tyr	200
AGG	GAG	CCC	TTC	GAG	CAG	CAC	CTC	CTG	CTG	GGC	GTC	AGC	GTT	TCC	645
Arg	Glu	Pro	Phe	Glu	Gln	His	Leu	Leu	Leu	Gly	Val	Ser	Val	Ser	215
TGC	ATT	GTC	ATC	CTG	GCC	GTC	TGC	CTG	TTG	TGC	TAT	GTC	AGC	ATC	690
Cys	Ile	Val	Ile	Leu	Ala	Val	Cys	Leu	Leu	Cys	Tyr	Val	Ser	Ile	230
ACC	AAG	ATT	AAG	AAA	GAA	TGG	TGG	GAT	CAG	ATT	CCC	AAC	CCA	GCC	735
Thr	Lys	Ile	Lys	Lys	Glu	Trp	Trp	Asp	Gln	Ile	Pro	Asn	Pro	Ala	245

FIGURE 4B

CGC	AGC	CGC	CTC	GTG	GCT	ATA	ATA	ATC	CAG	GAT	GCT	CAG	GGG	TCA	780
Arg	Ser	Arg	Leu	Val	Ala	Ile	Ile	Ile	Gln	Asp	Ala	Gln	Gly	Ser	260
CAG	TGG	GAG	AAG	CGG	TCC	CGA	GGC	CAG	GAA	CCA	GCC	AAG	TGC	CCA	825
Gln	Trp	Glu	Lys	Arg	Ser	Arg	Gly	Gln	Glu	Pro	Ala	Lys	Cys	Pro	275
CAC	TGG	AAG	AAT	TGT	CTT	ACC	AAG	CTC	TTG	CCC	TGT	TTT	CTG	GAG	870
His	Trp	Lys	Asn	Cys	Leu	Thr	Lys	Leu	Leu	Pro	Cys	Phe	Leu	Glu	290
CAC	AAC	ATG	AAA	AGG	GAT	GAA	GAT	CCT	CAC	AAG	GCT	GCC	AAA	GAG	915
His	Asn	Met	Lys	Arg	Asp	Glu	Asp	Pro	His	Lys	Ala	Ala	Lys	Glu	305
ATG	CCT	TTC	CAG	GGC	TCT	GGA	AAA	TCA	GCA	TGG	TGC	CCA	GTG	GAG	960
Met	Pro	Phe	Gln	Gly	Ser	Gly	Lys	Ser	Ala	Trp	Cys	Pro	Val	Glu	320
ATC	AGC	AAG	ACA	GTC	CTC	TGG	CCA	GAG	AGC	ATC	AGC	GTG	GTG	CGA	1005
Ile	Ser	Lys	Thr	Val	Leu	Trp	Pro	Glu	Ser	Ile	Ser	Val	Val	Arg	335
TGT	GTG	GAG	TTG	TTT	GAG	GCC	CCG	GTG	GAG	TGT	GAG	GAG	GAG	GAG	1050
Cys	Val	Glu	Leu	Phe	Glu	Ala	Pro	Val	Glu	Cys	Glu	Glu	Glu	Glu	350
GAG	GTA	GAG	GAA	GAA	AAA	GGG	AGC	TTC	TGT	GCA	TCG	CCT	GAG	AGC	1095
Glu	Val	Glu	Glu	Glu	Lys	Gly	Ser	Phe	Cys	Ala	Ser	Pro	Glu	Ser	365
AGC	AGG	GAT	GAC	TTC	CAG	GAG	GGA	AGG	GAG	GGC	ATT	GTG	GCC	CGG	1140
Ser	Arg	Asp	Asp	Phe	Gln	Glu	Gly	Arg	Glu	Gly	Ile	Val	Ala	Arg	380
CTA	ACA	GAG	AGC	CTG	TTC	CTG	GAC	CTG	CTC	GGA	GAG	GAG	AAT	GGG	1185
Leu	Thr	Glu	Ser	Leu	Phe	Leu	Asp	Leu	Leu	Gly	Glu	Glu	Asn	Gly	395
GGC	TTT	TGC	CAG	CAG	GAC	ATG	GGG	GAG	TCA	TGC	CTT	CTT	CCA	CCT	1230
Gly	Phe	Cys	Gln	Gln	Asp	Met	Gly	Glu	Ser	Cys	Leu	Leu	Pro	Pro	410
TCG	GGA	AGT	ACG	AGT	GCT	CAC	ATG	CCC	TGG	GAT	GAG	TTC	CCA	AGT	1275
Ser	Gly	Ser	Thr	Ser	Ala	His	Met	Pro	Trp	Asp	Glu	Phe	Pro	Ser	425
GCA	GGG	CCC	AAG	GAG	GCA	CCT	CCC	TGG	GGC	AAG	GAG	CAG	CCT	CTC	1320
Ala	Gly	Pro	Lys	Glu	Ala	Pro	Pro	Trp	Gly	Lys	Glu	Gln	Pro	Leu	440
CAC	CTG	GAG	CCA	AGT	CCT	CCT	GCC	AGC	CCG	ACC	CAG	AGT	CCA	GAC	1365
His	Leu	Glu	Pro	Ser	Pro	Pro	Ala	Ser	Pro	Thr	Gln	Ser	Pro	Asp	455
AAC	CTG	ACT	TGC	ACA	GAG	ACG	CCC	CTC	GTC	ATC	GCA	GGC	AAC	CCT	1410
Asn	Leu	Thr	Cys	Thr	Glu	Thr	Pro	Leu	Val	Ile	Ala	Gly	Asn	Pro	470
GCT	TAC	CGC	AGC	TTC	AGC	AAC	TCC	CTG	AGC	CAG	TCA	CCG	TGT	CCC	1455
Ala	Tyr	Arg	Ser	Phe	Ser	Asn	Ser	Leu	Ser	Gln	Ser	Pro	Cys	Pro	485
AGA	GAG	CTG	GGT	CCA	GAC	CCA	CTG	CTG	GCC	AGA	CAC	CTG	GAG	GAA	1500
Arg	Glu	Leu	Gly	Pro	Asp	Pro	Leu	Leu	Ala	Arg	His	Leu	Glu	Glu	500
GTA	GAA	CCC	GAG	ATG	CCC	TGT	GTC	CCC	CAG	CTC	TCT	GAG	CCA	ACC	1545
Val	Glu	Pro	Glu	Met	Pro	Cys	Val	Pro	Gln	Leu	Ser	Glu	Pro	Thr	515

FIGURE 4C

ACT	GTG	CCC	CAA	CCT	GAG	CCA	GAA	ACC	TGG	GAG	CAG	ATC	CTC	CGC	1590
Thr	Val	Pro	Gln	Pro	Glu	Pro	Glu	Thr	Trp	Glu	Gln	Ile	Leu	Arg	530
CGA	AAT	GTC	CTC	CAG	CAT	GGG	GCA	GCT	GCA	GCC	CCC	GTC	TCG	GCC	1635
Arg	Asn	Val	Leu	Gln	His	Gly	Ala	Ala	Ala	Ala	Pro	Val	Ser	Ala	545
CCC	ACC	AGT	GGC	TAT	CAG	GAG	TTT	GTA	CAT	GCG	GTG	GAG	CAG	GGT	1680
Pro	Thr	Ser	Gly	Tyr	Gln	Glu	Phe	Val	His	Ala	Val	Glu	Gln	Gly	560
GGC	ACC	CAG	GCC	AGT	GCG	GTG	GTG	GGC	TTG	GGT	CCC	CCA	GGA	GAG	1725
Gly	Thr	Gln	Ala	Ser	Ala	Val	Val	Gly	Leu	Gly	Pro	Pro	Gly	Glu	575
GCT	GGT	TAC	AAG	GCC	TTC	TCA	AGC	CTG	CTT	GCC	AGC	AGT	GCT	GTG	1770
Ala	Gly	Tyr	Lys	Ala	Phe	Ser	Ser	Leu	Leu	Ala	Ser	Ser	Ala	Val	590
TCC	CCA	GAG	AAA	TGT	GGG	TTT	GGG	GCT	AGC	AGT	GGG	GAA	GAG	GGG	1815
Ser	Pro	Glu	Lys	Cys	Gly	Phe	Gly	Ala	Ser	Ser	Gly	Glu	Glu	Gly	605
TAT	AAG	CCT	TTC	CAA	GAC	CTC	ATT	CCT	GGC	TGC	CCT	GGG	GAC	CCT	1860
Tyr	Lys	Pro	Phe	Gln	Asp	Leu	Ile	Pro	Gly	Cys	Pro	Gly	Asp	Pro	620
GCC	CCA	GTC	CCT	GTC	CCC	TTG	TTC	ACC	TTT	GGA	CTG	GAC	AGG	GAG	1905
Ala	Pro	Val	Pro	Val	Pro	Leu	Phe	Thr	Phe	Gly	Leu	Asp	Arg	Glu	635
CCA	CCT	CGC	AGT	CCG	CAG	AGC	TCA	CAT	CTC	CCA	AGC	AGC	TCC	CCA	1950
Pro	Pro	Arg	Ser	Pro	Gln	Ser	Ser	His	Leu	Pro	Ser	Ser	Ser	Pro	650
GAG	CAC	CTG	GGT	CTG	GAG	CCG	GGG	GAA	AAG	GTA	GAG	GAC	ATG	CCA	1995
Glu	His	Leu	Gly	Leu	Glu	Pro	Gly	Glu	Lys	Val	Glu	Asp	Met	Pro	665
AAG	CCC	CCA	CTT	CCC	CAG	GAG	CAG	GCC	ACA	GAC	CCC	CTT	GTG	GAC	2040
Lys	Pro	Pro	Leu	Pro	Gln	Glu	Gln	Ala	Thr	Asp	Pro	Leu	Val	Asp	680
AGC	CTG	GGC	AGT	GGC	ATT	GTC	TAC	TCA	GCC	CTT	ACC	TGC	CAC	CTG	2085
Ser	Leu	Gly	Ser	Gly	Ile	Val	Tyr	Ser	Ala	Leu	Thr	Cys	His	Leu	695
TGC	GGC	CAC	CTG	AAA	CAG	TGT	CAT	GGC	CAG	GAG	GAT	GGT	GGC	CAG	2130
Cys	Gly	His	Leu	Lys	Gln	Cys	His	Gly	Gln	Glu	Asp	Gly	Gly	Gln	710
ACC	CCT	GTC	ATG	GCC	AGT	CCT	TGC	TGT	GGC	TGC	TGC	TGT	GGA	GAC	2175
Thr	Pro	Val	Met	Ala	Ser	Pro	Cys	Cys	Gly	Cys	Cys	Cys	Gly	Asp	725
AGG	TCC	TCG	CCC	CCT	ACA	ACC	CCC	CTG	AGG	GCC	CCA	GAC	CCC	TCT	2220
Arg	Ser	Ser	Pro	Pro	Thr	Thr	Pro	Leu	Arg	Ala	Pro	Asp	Pro	Ser	740
CCA	GGT	GGG	GTT	CCA	CTG	GAG	GCC	AGT	CTG	TGT	CCG	GCC	TCC	CTG	2265
Pro	Gly	Gly	Val	Pro	Leu	Glu	Ala	Ser	Leu	Cys	Pro	Ala	Ser	Leu	755
GCA	CCC	TCG	GGC	ATC	TCA	GAG	AAG	AGT	AAA	TCC	TCA	TCA	TCC	TTC	2310
Ala	Pro	Ser	Gly	Ile	Ser	Glu	Lys	Ser	Lys	Ser	Ser	Ser	Ser	Phe	770
CAT	CCT	GCC	CCT	GGC	AAT	GCT	CAG	AGC	TCA	AGC	CAG	ACC	CCC	AAA	2355
His	Pro	Ala	Pro	Gly	Asn	Ala	Gln	Ser	Ser	Ser	Gln	Thr	Pro	Lys	785
ATC	GTG	AAC	TTT	GTC	TCC	GTG	GGA	CCC	ACA	TAC	ATG	AGG	GTC	TCT	2400
Ile	Val	Asn	Phe	Val	Ser	Val	Gly	Pro	Thr	Tyr	Met	Arg	Val	Ser	800

FIGURE 5A

1 MGWLCSGLLFPVSCLVLLQVASSGNMKVLQEPTCVSDYMSISTCEWKMN 50
| | | | | | | | | | | | | | | | | |
1 MGR LCTKFLTSVGCLILLVLTGSGSIKVLGEPTCFSDYIRTSTCEWFLDS 50
51 PTNCSTELRLLYQLVFL.LSEAHTCIPENNGGAGCVCHLLMDDVVSADNY 99
| | | | | | | | | | | | | | | | | |
51 AVDCSSQLCLHYRLMFFEFSENLTICIPRNSASTVCVCHMEMNRPVQSDRY 100
100 TLDLWAGQQLLWKGSFKPSEHV KPRAPGNLTVHTNVSDTLLLTWSNPYP 149
| | | | | | | | | | | | | | | | | |
101 QMELWAEHRQLWQGSFSPSGNVKPLAPDNLT LHTNVSEWLLTWN NLYPS 150
150 DNYLYNHLTYAVNIWSENDPADFRIYNVTYLEPSLRIAASTLKSGISYRA 199
| | | | | | | | | | | | | | | | | |
151 NNLLYKDLISMVNISREDNPAEFIVYNVTYKEPRLSFPINILMSGVYYTA 200
200 RVRAWAQCYN TTWSEWSPSTKWHNSYREPFEQHLLLGVSVCIVILAVCL 249
| | | | | | | | | | | | | | | | | |
201 RVRVRSQILTGTWSEWSPSITWYNHFQLPLIQRLPLGVTISCLCIPLFCL 250
250 LCYVSITKIKKEWWDQIPNPARSRLVAII IQDAQGSQWEKRSRGQEPAC 299
| | | | | | | | | | | | | | | | | |
251 FCYFSITKIKKIWWDQIPTPARSPLVAII IQDAQVPLWDKQTRSQUESTKY 300
300 PHWKNCCLKLLPCFLEHNMKRDEDPHKA AKEMPFGSGKSAWCPVEISK 349
| | | | | | | | | | | | | | | | | |
301 PHWKTCLDKLLPCLLKHRVKKKTDFPKAAPT KSLQSPGKAGWCPMEVSRT 350
350 VLWPE..SISVVRCELFEAPVECEEEEEVEEEKGSFCASPSSRD.DFQ 396
| | | | | | | | | | | | | | | | | |
351 VLWPENVS SVVRCEMELFEAPVQNV EEEEEDEIVKEDLSMSPENSGGCGFQ 400
397 EGREGIVARLTESLFLDLLGEENGFCQQDMGESCLLPSPSGSTSAHMPWD 446
| | | | | | | | | | | | | | | | | |
401 ESQADIMARLTENLFSDLLEAENGGLGQSALAESCSPLPSGSGQASVSWA 450
447 EFPSAGPKEAPPWGKEQPLHLEPSPPASPTQSPDNLTCTETPLVIAGNPA 496
| | | | | | | | | | | | | | | | | |
451 CLPMGPSEEATCQVTEQPSHPGP.LSGSPAQSAPT LACTQVPLVLADNPA 499
497 YRSFSNSLSQSPCPRELGPDP LLARHLEEVEPEMPCVPQLSEPTTVPQPE 546
| | | | | | | | | | | | | | | | | |
500 YRSFSDCCSPAPNPGELAPEQQQADHLEEEEPSPADPHSSGP...PMQP 546
547 PETWEQILRRNVLQHGA AAAPVSAPTSGYQEFVHAVEQGGTQASAVVGLG 596
| | | | | | | | | | | | | | | | | |
547 VESWEQILHMSVLQHGA AAGSTPAPAGGYQEFVQAVKQGA AQDPGVPGVR 596
597 PPGEAGYKAFSSLLASSAVSPEKCGFGASSGEEGYKPFQDLIPGCPGDP 646
| | | | | | | | | | | | | | | | | |
597 PSGDPGYKAFSSLLSSNGIRGDTAAAGTDDGHGGYKPFQNPVP...NQS 642

FIGURE 5B

647 PVPVPLFTFGLDREPPRSPQSSHLPSSSPÉHLGLEPGEKVEDMPKPPLPQ 696
| | | | | | | | | | | | | | | | | | | | | |
643 PSSVPLFTFGLDTELSPLNSDPPKSPPECLGLELGLKGGDWVKAPPPA 692
697 EQATDPLVDSLGSIVYSALTCHLCGHLKQCHGQEDGGQTPVMASPCCGC 746
| | | | | | | | | | | | | | | | | | | | | |
693 DQVPKPFGDDLFGFIVYSSLTCHLCGHLKQHHSQEEGGQSPIVASPGCGC 742
747 CCGDRSSPPTTPLRAPDSPGGVPLEASLCPASLAPSGISEKSKSSSFH 796
| | | | | | | | | | | | | | | | | | | | | |
743 CYDDRSPSLGSLSGALESCPEGIPPEANLMSAPKTPSNLSGEGK..... 786
797 PAPGNAQSSSQTPKIVNFVSVGPTYMRVS 825
| | | | | | | | | | | | | | | | | | | | | |
787 .GPGHSPVPSQTTE....VPVGALGIAVS 810




FIGURE 6

Figure 7A




Figure 7D




Figure 7B




Figure 7E




Figure 7C





Figure 7F






FIGURE 8

Figure 9A




Figure 9B





Figure 9C






FIGURE 10





FIGURE 11

Figure 12






FIGURE 13


RAT anti-Ig D EFFECTS ON SERA Ig E LEVELS

FIGURE 14

**Effect of Treatment with sIL-1R or sIL-4R
on Contact Hypersensitivity Responses to DNFB**




FIGURE 15




FIGURE 16




FIGURE 17

ANTIBODIES TO INTERLEUKIN-4 RECEPTORS AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

This is a divisional of application Ser. No. 09/094,917, filed Jun. 15, 1998, now U.S. Pat. No. 6,391,581, now allowed, which is a continuation of application Ser. No. 07/480,694, filed Feb. 14, 1990, now U.S. Pat. No. 5,840,869, which is a continuation-in-part of application Ser. No. 07/370,924, filed Jun. 23, 1989, abandoned, which is a continuation-in-part of Ser. No. 07/326,156, filed Mar. 20, 1989, abandoned, which is a continuation-in-part of Ser. No. 07/319,438, filed Mar. 2, 1989, abandoned, which is a continuation-in-part of Ser. No. 07/265,047, filed Oct. 31, 1988, abandoned.

BACKGROUND OF THE INVENTION

The present invention relates generally to cytokine receptors and, more specifically, to Interleukin-4 receptors. Interleukin-4 (IL-4, also known as B cell stimulating factor, or BSF-1) was originally characterized by its ability to stimulate the proliferation of B cells in response to low concentrations of antibodies directed to surface immunoglobulin. More recently, IL-4 has been shown to possess a far broader spectrum of biological activities, including growth co-stimulation of T cells, mast cells, granulocytes, megakaryocytes, and erythrocytes. In addition, IL-4 stimulates the proliferation of several IL-2- and IL-3-dependent cell lines, induces the expression of class II major histocompatibility complex molecules on resting B cells, and enhances the secretion of IgE and IgG1 isotypes by stimulated B cells. Both murine and human IL-4 have been definitively characterized by recombinant DNA technology and by purification to homogeneity of the natural murine protein (Yokota et al., *Proc. Natl. Acad. Sci. USA* 83:5894, 1986; Noma et al., *Nature* 319:640, 1986; and Grabstein et al., *J. Exp. Med.* 163:1405, 1986).

The biological activities of IL-4 are mediated by specific cell surface receptors for IL-4 which are expressed on primary cells and in vitro cell lines of mammalian origin. IL-4 binds to the receptor, which then transduces a biological signal to various immune effector cells. Purified IL-4 receptor (IL-4R) compositions will therefore be useful in diagnostic assays for IL-4 or IL-4 receptor, and in raising antibodies to IL-4 receptor for use in diagnosis or therapy. In addition, purified IL-4 receptor compositions may be used directly in therapy to bind or scavenge IL-4, providing a means for regulating the biological activities of this cytokine.

Although IL-4 has been extensively characterized, little progress has been made in characterizing its receptor. Numerous studies documenting the existence of an IL-4 receptor on a wide range of cell types have been published; however, structural characterization has been limited to estimates of the molecular weight of the protein as determined by SDS-PAGE analysis of covalent complexes formed by chemical cross-linking between the receptor and radiolabeled IL-4 molecules. Ohara et al. (*Nature* 325:537, 1987) and Park et al. (*Proc. Natl. Acad. Sci. USA* 84:1669, 1987) first established the presence of an IL-4 receptor using radioiodinated recombinant murine IL-4 to bind a high affinity receptor expressed in low numbers on B and T lymphocytes and a wide range of cells of the hematopoietic lineage. By affinity cross-linking ¹²⁵I-IL-4 to IL-4R, Ohara et al. and Park et al. identified receptor proteins having

apparent molecular weights of 60,000 and 75,000 daltons, respectively. It is possible that the small receptor size observed on the murine cells represents a proteolytically cleaved fragment of the native receptor. Subsequent experiments by Park et al. (*J. Exp. Med.* 166:476, 1987) using yeast-derived recombinant human IL-4 radiolabeled with ¹²⁵I showed that human IL-4 receptor is present not only on cell lines of B, T, and hematopoietic cell lineages, but is also found on human fibroblasts and cells of epithelial and endothelial origin. IL-4 receptors have since been shown to be present on other cell lines, including CBA/N splenic B cells (Nakajima et al., *J. Immunol.* 139:774, 1987), Burkitt lymphoma Jijoye cells (Cabrillat et al., *Biochem. & Biophys. Res. Commun.* 149:995, 1987), a wide variety of hemopoietic and nonhemopoietic cells (Lowenthal et al., *J. Immunol.* 140:456, 1988), and murine Lyt-2⁺/L3T4⁺ thymocytes. More recently, Park et al. (UCLA Symposia, *J. Cell Biol.*, Suppl. 12A, 1988) reported that, in the presence of sufficient protease inhibitors, ¹²⁵I-IL-4-binding plasma membrane receptors of 138–145 kDa could be identified on several murine cell lines. Considerable controversy thus remains regarding the actual size and structure of IL-4 receptors.

Further study of the structure and biological characteristics of IL-4 receptors and the role played by IL-4 receptors in the responses of various cell populations to IL-4 or other cytokine stimulation, or of the methods of using IL-4 receptors effectively in therapy, diagnosis, or assay, has not been possible because of the difficulty in obtaining sufficient quantities of purified IL-4 receptor. No cell lines have previously been known to express high levels of IL-4 receptors constitutively and continuously, and in cell lines known to express detectable levels of IL-4 receptor, the level of expression is generally limited to less than about 2000 receptors per cell. Thus, efforts to purify the IL-4 receptor molecule for use in biochemical analysis or to clone and express mammalian genes encoding IL-4 receptor have been impeded by lack of purified receptor and a suitable source of receptor mRNA.

SUMMARY OF THE INVENTION

The present invention provides DNA sequences encoding mammalian Interleukin-4 receptors (IL-4R) or subunits thereof. Preferably, such DNA sequences are selected from the group consisting of: (a) cDNA clones having a nucleotide sequence derived from the coding region of a native IL-4R gene; (b) DNA sequences capable of hybridization to the cDNA clones of (a) under moderately stringent conditions and which encode biologically active IL-4R molecules; and (c) DNA sequences which are degenerate, as a result of the genetic code, to the DNA sequences defined in (a) and (b) and which encode biologically active IL-4R molecules. The present invention also provides recombinant expression vectors comprising the DNA sequences defined above, recombinant IL-4R molecules produced using the recombinant expression vectors, and processes for producing the recombinant IL-4R molecules using the expression vectors.

The present invention also provides substantially homogeneous protein compositions comprising mammalian IL-4R. The full length murine molecule is a glycoprotein having a molecular weight of about 130,000 to about 140,000 M_r by SDS-PAGE. The apparent binding affinity (K_d) for COS cells transfected with murine IL-4 receptor clones 16 and 18 from the CTLL 19.4 library is 1 to 8×10⁹ M⁻¹. The K_d for COS cells transfected with murine IL-4 receptor clones 7B9-2 and 7B9-4 from the murine 7B9 library is 2×10⁹ to 1×10¹⁰ M⁻¹. The mature murine IL-4 receptor molecule has an N-terminal amino acid sequence as follows: IKVLGEPTCFSDYIRTSTCEW.

The human IL-4R molecule is believed to have a molecular weight of between about 110,000 and 150,000 M_r and has an N-terminal amino acid sequence, predicted from the cDNA sequence and by analogy to the biochemically determined N-terminal sequence of the mature murine protein, as follows: MKVLQEPCTCVSDYMSISTCEW.

The present invention also provides compositions for use in therapy, diagnosis, assay of IL-4 receptor, or in raising antibodies to IL-4 receptors, comprising effective quantities of soluble receptor proteins prepared according to the foregoing processes. Such soluble recombinant receptor molecules include truncated proteins wherein regions of the receptor molecule not required for IL-4 binding have been deleted.

The present invention also provides a method for suppressing IL-4 mediated immune or inflammatory responses. This method comprises administering an effective quantity of soluble IL-4 receptor (sIL-4R), in association with a pharmaceutical carrier, to a mammal, including man. sIL-4R suppresses IL-4 dependent immune or inflammatory responses, including, for example, B cell mediated activities, such as B cell proliferation, immunoglobulin secretion, and expression of FcεR which are induced by IL-4. sIL-4R also suppresses cytotoxic T cell induction. Clinical applications of sIL-4R include, for example, use in allergy therapy to selectively suppress IgE synthesis and use in transplantation therapy to prevent allograft rejection. sIL-4R is also useful to suppress delayed-type hypersensitivity or contact hypersensitivity reactions. sIL-4R is highly specific in its immunosuppressive activity because it suppresses only IL-4 mediated immune responses.

These and other aspects of the present invention will become evident upon reference to the following detailed description and attached drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows restriction maps of cDNA clones containing the coding regions (denoted by a bar) of the murine and human IL-4R cDNAs. The restriction sites EcoRI, PvuII, Hinc II and Sst I are represented by the letters R, P, H and S, respectively.

FIGS. 2A–C depict the cDNA sequence and the derived amino acid sequence of the coding region of a murine IL-4 receptor, as derived from clone 7B9-2 of the 7B9 library. The N-terminal isoleucine of the mature protein is designated amino acid number 1. The coding region of the full-length membrane-bound protein from clone 7B9-2 is defined by amino acids 1–785. The ATC codon specifying the isoleucine residue constituting the mature N-terminus is underlined at position 1 of the protein sequence; the putative transmembrane region at amino acids 209–232 is also underlined. The sequences of the coding regions of clones 7B9-4 and clones CTLL-18 and CTLL-16 of the CTLL 19.4 library are identical to that of 7B9-2 except as follows. The coding region of CTLL-16 encodes a membrane-bound IL-4-binding receptor defined by amino acids –25 through 233 (including the putative 25 amino acid signal peptide sequence), but is followed by a TAG terminator codon (not shown) which ends the open reading frame. The nucleic acid sequence indicates the presence of a splice donor site at this position (indicated by an arrow in FIG. 1) and a splice acceptor site near the 3' end (indicated by a second arrow), suggesting that CTLL-16 was derived from an unspliced mRNA intermediate. Clones 7B94 and CTLL-18 encode amino acids 23 through 199 and –25 through 199, respec-

tively. After amino acid 199, a 114-base pair insert (identical in both clones and shown by an open box in FIG. 1) introduces six new amino acids, followed by a termination codon. This form of the receptor is soluble.

FIG. 3 is a schematic illustration of the mammalian high expression plasmid pCAV/NOT, which is described in greater detail in Example 8.

FIGS. 4A–C depict the coding sequence of a human IL-4 receptor cDNA from clone T22-8, which was obtained from a cDNA library derived from the T cell line T22. The predicted N-terminal methionine of the mature protein and the transmembrane region are underlined.

FIGS. 5A–B are a comparison of the predicted amino acid sequences of human (top line) and murine (bottom line) IL-4 receptor cDNA clones.

FIG. 6 shows the inhibition of B cell proliferation with IL-4 (panel A) or IL-1 (panel B) at various doses either alone (○) or in the presence of sIL-4R (□), sIL-1R (■) or anti-IL-4 antibody (●) as described in Example 15.

FIG. 7 shows the inhibition of B cell proliferation with fixed concentrations of 10 (●), 1 (□), 0.1 (○) or 0 (Δ) ng/ml of IL-4 (panels A–C) and IL-1 (panels D–F) at various doses of sIL-4R (panels A & D), anti-IL-4 antibody (panels B & E) and sIL-1R (panels C & F) as described in Example 15.

FIG. 8 shows the inhibition of immunoglobulin class switching with various doses of IL-4 and with sIL-4R (□), sIL-1R (■), or anti-IL-4 antibody (●) or medium control (Δ) as described in Example 15.

FIG. 9 shows the inhibition of IL-4-induced immunoglobulin class switching with fixed concentration of IL-4 and various doses of sIL-4R (□), sIL-1R (■), anti-IL-4 antibody (●) or medium control (Δ) as described in Example 16.

FIG. 10 shows the inhibition of MHC class II antigen expression with (dashed line) or without (solid line) IL-4 in the presence of medium control (panel A), sIL-4R (panel B), anti-IL-4 antibody (panel C) or sIL-1R (panel D) as described in Example 17.

FIG. 11 shows the inhibition of FcεR (CD23) expression with (dashed line) or without (solid line) IL-4 in the presence of medium control (panel A), sIL-4R (panel B), anti-IL-4 antibody (panel C) or sIL-1R (panel D) as described in Example 17.

FIG. 12 shows the inhibition of antigen specific polyclonal IgE levels by sIL-4R as described in Example 18.

FIG. 13 shows the inhibition of antigen specific anti-TNP-KLH IgE levels by sIL-4R as described in Example 18.

FIG. 14 shows that sIL-4R doses of 1, 5 or 25 ug on days –1, 0 and +1, as described in Example 19, do not significantly inhibit antigen specific anti-IgD IgE levels.

FIG. 15 shows the inhibition of contact hypersensitivity responses to DNFB with sIL-4R as described in Example 20.

FIG. 16 shows the inhibition of delayed-type hypersensitivity responses to SRBC with sIL-4R as described in Example 21.

FIG. 17 shows that sIL-4R, administered in Example 22, inhibits alloantigen induced proliferation (measured by weight gain) of lymph node cells. Mice are injected in the footpad with syngeneic cells (from the same species) and in the contralateral footpad with allogeneic cells (from another species). Large Δ values (e.g., for MSA or no reaction) indicate that the treatment did not inhibit alloantigen induced lymphocyte proliferation and inflammation. Small Δ values indicate that the treatment inhibited alloantigen induced lymphocyte proliferation and inflammation.

DETAILED DESCRIPTION OF THE
INVENTION

Definitions

"Interleukin-4" and "IL-4" (also referred to as B cell stimulating factor, or BSF-1) is a T cell-derived cytokine involved in the regulation of immune and inflammatory responses. The biological activities of IL-4 are mediated through binding to specific cell surface receptors, referred to as "Interleukin-4 receptors", "IL-4 receptors" or simply "IL-4R". "IL-4 mediated" immune or inflammatory responses include all biological responses which are caused by the binding of IL-4 to a native IL-4 receptor (bound to a cell surface) or which may be inhibited or suppressed by preventing IL-4 from binding to a native IL-4 receptor. IL-4 mediated biological responses include, for example, IL-4 induced proliferation of antigen-primed B lymphocytes, expression of class II major histocompatibility complex molecules on resting B cells, secretion and expression of antibodies of the IgE and IgG1 isotype, and regulation of the expression of the low affinity Fc receptor for IgE (CD23) on lymphocytes and monocytes. Outside the B lymphocyte compartment, IL-4 mediated biological responses include the proliferation of a variety of primary cells and in vitro cell lines, including factor-dependent T cell and mast cell lines, murine and human T lymphocytes, thymocytes, and connective tissue-type mast cells. IL-4 also induces both murine and human cytotoxic T cells. Under certain conditions, IL-4 inhibits the response of lymphoid cells to IL-2. IL-4 acts on both murine and human hematopoietic progenitor cells to either stimulate or suppress in vitro formation of colonies in combination with known colony stimulating factors. IL-4 also induces class I and class II MHC molecules on more mature cells of the monocytic lineage, enhances antigen presenting ability and promotes formation of giant multinucleated cells. Specific clinical conditions which may be mediated by IL-4 include, for example, graft rejection, graft versus host disease, allergy, asthma and delayed-type hypersensitivity responses.

As used herein, the terms "IL-4 receptor" or "IL-4R" refer to proteins which bind interleukin-4 (IL-4) molecules and, in their native configuration as intact human plasma membrane proteins, play a role in transducing the biological signal provided by IL-4 to a cell. Intact receptor proteins generally include an extracellular region which binds to a ligand, a hydrophobic transmembrane region which causes the protein to be immobilized within the plasma membrane lipid bilayer, and a cytoplasmic or intracellular region which interacts with cytoplasmic proteins and/or chemicals to deliver a biological signal to effector cells via a cascade of chemical reactions within the cytoplasm of the cell. The hydrophobic transmembrane region and a highly charged sequence of amino acids in the cytoplasmic region immediately following the transmembrane region cooperatively function to halt transport of the IL-4 receptor across the plasma membrane.

"IL-4 receptors" are proteins having amino acid sequences which are substantially similar to the native mammalian Interleukin-4 receptor amino acid sequences disclosed in FIGS. 2 and 4 or fragments thereof, and which are biologically active as defined below, in that they are capable of binding Interleukin-4 (IL-4) molecules or transducing a biological signal initiated by an IL-4 molecule binding to a cell, or cross-reacting with anti-IL-4R antibodies raised against IL-4R from natural (i.e., nonrecombinant) sources. The native human IL-4 receptor molecule has an apparent molecular weight by SDS-PAGE of about 140 kilodaltons (kDa). The native murine IL-4 receptor molecule

has an apparent molecular weight by SDS-PAGE of about 140 kilodaltons (kDa). The terms "IL-4 receptor" or "IL-4R" include, but are not limited to, soluble IL-4 receptors, as defined below. Specific IL-4 receptor polypeptides are designated herein by parenthetically indicating the amino acid sequence numbers, followed by any additional amino acid sequences. For Example, human IL-4R (1-207) refers to a human IL-4R protein having the sequence of amino acids 1-207 as shown in FIG. 4A. Human IL-4R (1-184) Pro Ser Asn Glu Asn refers to a human IL-4R protein having the sequence of amino acids 1-184 as shown in FIG. 4A, followed by the amino acid sequence Pro Ser Asn Glu Asn. As used throughout the specification, the term "mature" means a protein expressed in a form lacking a leader sequence as may be present in full-length transcripts of a native gene. Various bioequivalent protein and amino acid analogs are described in the detailed description of the invention.

"Substantially similar" IL-4 receptors include those whose amino acid or nucleic acid sequences vary from the native sequences by one or more substitutions, deletions, or additions, the net effect of which is to retain biological activity of the IL-4R protein. For example, nucleic acid subunits and analogs are "substantially similar" to the specific DNA sequences disclosed herein if: (a) the DNA sequence is derived from the coding region of a native mammalian IL-4R gene; (b) the DNA sequence is capable of hybridization to DNA sequences of (a) under moderately stringent conditions and which encode biologically active IL-4R molecules; or DNA sequences which are degenerate as a result of the genetic code to the DNA sequences defined in (a) or (b) and which encode biologically active IL-4R molecules. Substantially similar analog proteins will generally be greater than about 30 percent similar to the corresponding sequence of the native IL-4R. Sequences having lesser degrees of similarity but comparable biological activity are considered to be equivalents. More preferably, the analog proteins will be greater than about 80 percent similar to the corresponding sequence of the native IL-4R, in which case they are defined as being "substantially identical." In defining nucleic acid sequences, all subject nucleic acid sequences capable of encoding substantially similar amino acid sequences are considered substantially similar to a reference nucleic acid sequence. Percent similarity may be determined, for example, by comparing sequence information using the GAP computer program, version 6.0, available from the University of Wisconsin Genetics Computer Group (UWCGC). The GAP program utilizes the alignment method of Needleman and Wunsch (*J. Mol. Biol.* 48:443, 1970), as revised by Smith and Waterman (*Adv. Appl. Math.* 2:482, 1981). Briefly, the GAP program defines similarity as the number of aligned symbols (i.e., nucleotides or amino acids) which are similar, divided by the total number of symbols in the shorter of the two sequences. The preferred default parameters for the GAP program include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) for nucleotides, and the weighted comparison matrix of Gribskov and Burgess, *Nucl. Acids Res.* 14:6745, 1986, as described by Schwartz and Dayhoff, ed., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 353-358, 1979; (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps.

"Soluble IL-4 receptor" or "sIL-4-R" as used in the context of the present invention refers to a protein, or a substantially equivalent analog, having an amino acid sequence corresponding to the extracellular region of native

IL-4 receptors, for example, polypeptides having the amino acid sequences substantially equivalent to the sequences of amino acids 1–208 of FIG. 2, amino acids 1–207 of FIG. 4 or to the amino acid sequences discussed in Examples 8C and 9. Equivalent sIL-4Rs include polypeptides which vary from the sequences shown in FIGS. 2 or 4 by one or more substitutions, deletions, or additions, and which retain the ability to bind IL-4 and inhibit the ability of IL-4 to transduce a signal via cell surface bound IL-4 receptor proteins. Because sIL-4R proteins are devoid of a transmembrane region, they are secreted from the host cell in which they are produced. When administered in therapeutic formulations, sIL-4R proteins circulate in the body and bind to circulating IL-4 molecules, preventing interaction of IL-4 with natural IL-4 receptors and inhibiting transduction of IL-4 mediated biological signals, such as immune or inflammatory responses. The ability of a polypeptide to inhibit IL-4 signal transduction can be determined by transfecting cells with recombinant IL-4 receptor DNAs to obtain recombinant receptor expression. The cells are then contacted with IL-4 and the resulting metabolic effects examined. If an effect results which is attributable to the action of the ligand, then the recombinant receptor has signal transducing activity. Exemplary procedures for determining whether a polypeptide has signal transducing activity are disclosed by Idzerda et al., *J. Exp. Med.*, March 1990 in press, Curtis et al., *Proc. Natl. Acad. Sci. USA* 86:3045 (1989), Prywes et al., *EMBO J.* 5:2179 (1986) and Chou et al., *J. Biol. Chem.* 262:1842 (1987). Alternatively primary cells of cell lines which express an endogenous IL-4 receptor and have a detectable biological response to IL-4 could also be utilized. Such is the case with the CTLL-2 cell line which responds by short term proliferation in response to either IL-2 or IL-4; the IL-4 induced proliferation can be blocked specifically by the addition of exogenous soluble IL-4R (Mosley et al., *Cell* 59:335 (1989). In addition, any one of the in vivo or in vitro assays described in Examples 14–23 can be utilized to determine whether a soluble IL-4R inhibits transduction of a specific IL-4 mediated biological signal. The cloning, sequencing and expression of full-length and soluble forms of the receptor for murine IL-4 have recently been described by the applicants, Mosley et al., *Cell* 59:335, 1989, which publication is incorporated herein by reference.

“Recombinant,” as used herein, means that a protein is derived from recombinant (e.g., microbial or mammalian) expression systems. “Microbial” refers to recombinant proteins made in bacterial or fungal (e.g., yeast) expression systems. As a product, “recombinant microbial” defines a protein produced in a microbial expression system which is essentially free of native endogenous substances. Protein expressed in most bacterial cultures, e.g., *E. coli*, will be free of glycan. Protein expressed in yeast may have a glycosylation pattern different from that expressed in mammalian cells.

“Biologically active,” as used throughout the specification as a characteristic of IL-4 receptors, means that a particular molecule shares sufficient amino acid sequence similarity with the embodiments of the present invention disclosed herein to be capable of binding detectable quantities of IL-4, transducing an IL-4 signal to a cell, for example, as a component of a hybrid receptor construct, or cross-reacting with anti-IL-4R antibodies raised against IL-4R from natural (i.e., nonrecombinant) sources. Preferably, biologically active IL-4 receptors within the scope of the present invention are capable of binding greater than 0.1 nmoles IL-4 per nmole receptor, and most preferably, greater than 0.5 nmole IL-4 per nmole receptor in standard binding assays (see below).

“DNA sequence” refers to a DNA molecule, in the form of a separate fragment or as a component of a larger DNA construct, which has been derived from DNA isolated at least once in substantially pure form, i.e., free of contaminating endogenous materials and in a quantity or concentration enabling identification, manipulation, and recovery of the sequence and its component nucleotide sequences by standard biochemical methods, for example, using a cloning vector. Such sequences are preferably provided in the form of an open reading frame uninterrupted by internal nontranslated sequences, or introns, which are typically present in eucaryotic genes. Genomic DNA containing the relevant sequences could also be used. Sequences of non-translated DNA may be present 5' or 3' from the open reading frame, where the same do not interfere with manipulation or expression of the coding regions.

“Nucleotide sequence” refers to a heteropolymer of deoxyribonucleotides. DNA sequences encoding the proteins provided by this invention can be assembled from cDNA fragments and short oligonucleotide linkers, or from a series of oligonucleotides, to provide a synthetic gene which is capable of being expressed in a recombinant transcriptional unit.

“Recombinant expression vector” refers to a replicable DNA construct used either to amplify or to express DNA which encodes IL-4R and which includes a transcriptional unit comprising an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers, (2) a structural or coding sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription and translation initiation and termination sequences. Structural elements intended for use in yeast expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where recombinant protein is expressed without a leader or transport sequence, it may include an N-terminal methionine residue. This residue may optionally be subsequently cleaved from the expressed recombinant protein to provide a final product.

“Recombinant microbial expression system” means a substantially homogeneous monoculture of suitable host microorganisms, for example, bacteria such as *E. coli* or yeast such as *S. cerevisiae*, which have stably integrated a recombinant transcriptional unit into chromosomal DNA or carry the recombinant transcriptional unit as a component of a resident plasmid. Generally, cells constituting the system are the progeny of a single ancestral transformant. Recombinant expression systems as defined herein will express heterologous protein upon induction of the regulatory elements linked to the DNA sequence or synthetic gene to be expressed.

Proteins and Analogs

The present invention provides substantially homogeneous recombinant mammalian IL-4R polypeptides substantially free of contaminating endogenous materials and, optionally, without a associated native-pattern glycosylation. The native murine and human IL-4 receptor molecules are recovered from cell lysates as glycoproteins having an apparent molecular weight by SDS-PAGE of about 130–145 kilodaltons (kDa). Mammalian IL-4R of the present invention includes, by way of example, primate, human, murine, canine, feline, bovine, ovine, equine and porcine IL-4R. Derivatives of IL-4R within the scope of the invention also include various structural forms of the primary protein which retain biological activity. Due to the presence of ionizable amino and carboxyl groups, for example, an IL-4R protein may be in the form of acidic or basic salts, or in

neutral form. Individual amino acid residues may also be modified by oxidation or reduction.

The primary amino acid structure may be modified by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, lipids, phosphate, acetyl groups and the like, or by creating amino acid sequence mutants. Covalent derivatives are prepared by linking particular functional groups to IL-4R amino acid side chains or at the N- or C-termini. Other derivatives of IL-4R within the scope of this invention include covalent or aggregative conjugates of IL-4R or its fragments with other proteins or polypeptides, such as by synthesis in recombinant culture as N-terminal or C-terminal fusions. For example, the conjugated peptide may be a signal (or leader) polypeptide sequence at the N-terminal region of the protein which co-translationally or post-translationally directs transfer of the protein from its site of synthesis to its site of function inside or outside of the cell membrane or wall (e.g., the yeast α -factor leader). IL-4R protein fusions can comprise peptides added to facilitate purification or identification of IL-4R (e.g., poly-His). Specific examples of a poly-HIS fusion construct that is biologically active are soluble human IL-4R (1-207) His His and soluble human IL-4R (1-207) His His His His His His. The amino acid sequence of IL-4 receptor can also be linked to the peptide Asp-Tyr-Lys-Asp-Asp-Asp-Asp-Lys (DYKDDDDK) (Hopp et al., *Bio/Technology* 6:1204, 1988.) The latter sequence is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody, enabling rapid assay and facile purification of expressed recombinant protein. This sequence is also specifically cleaved by bovine mucosal enterokinase at the residue immediately following the Asp-Lys pairing. Fusion proteins capped with this peptide may also be resistant to intracellular degradation in *E. coli*. A specific example of such a peptide is soluble human IL-4R (1-207) Asp Tyr Lys Asp Asp Asp Asp Lys.

IL-4R derivatives may also be used as immunogens, reagents in receptor-based immunoassays, or as binding agents for affinity purification procedures of IL-4 or other binding ligands. IL-4R derivatives may also be obtained by cross-linking agents, such as M-maleimidobenzoyl succinimide ester and N-hydroxysuccinimide, at cysteine and lysine residues. IL-4R proteins may also be covalently bound through reactive side groups to various insoluble substrates, such as cyanogen bromide-activated, bisoxirane-activated, carbonyldiimidazole-activated or tosyl-activated agarose structures, or by adsorbing to polyolefin surfaces (with or without glutaraldehyde cross-linking). Once bound to a substrate, IL-4R may be used to selectively bind (for purposes of assay or purification) anti-IL-4R antibodies or IL-4.

The present invention also includes IL-4R with or without associated native-pattern glycosylation. IL-4R expressed in yeast or mammalian expression systems, e.g., COS-7 cells, may be similar or significantly different in molecular weight and glycosylation pattern than the native molecules, depending upon the expression system. Expression of IL-4R DNAs in bacteria such as *E. coli* provides non-glycosylated molecules. Functional mutant analogs of mammalian IL-4R having inactivated N-glycosylation sites can be produced by oligonucleotide synthesis and ligation or by site-specific mutagenesis techniques. These analog proteins can be produced in a homogeneous, reduced-carbohydrate form in good yield using yeast expression systems. N-glycosylation sites in eukaryotic proteins are characterized by the amino acid triplet Asn-A₁-Z, where A₁ is any amino acid except Pro, and Z is Ser or Thr. In this sequence, asparagine

provides a side chain amino group for covalent attachment of carbohydrate. Such a site can be eliminated by substituting another amino acid for Asn or for residue Z, deleting Asn or Z, or inserting a non-Z amino acid between A₁ and Z, or an amino acid other than Asn between Asn and A₁.

IL-4R derivatives may also be obtained by mutations of IL-4R or its subunits. An IL-4R mutant, as referred to herein, is a polypeptide homologous to IL-4R but which has an amino acid sequence different from native IL-4R because of a deletion, insertion or substitution. Like most mammalian genes, mammalian IL-4 receptors are presumably encoded by multi-exon genes. Alternative mRNA constructs which can be attributed to different mRNA splicing events following transcription, and which share large regions of identity or similarity with the cDNAs claimed herein, are considered to be within the scope of the present invention. Bioequivalent analogs of IL-4R proteins may be constructed by, for example, making various substitutions of residues or sequences or deleting terminal or internal residues or sequences not needed for biological activity. For example, cysteine residues can be deleted or replaced with other amino acids to prevent formation of incorrect intramolecular disulfide bridges upon renaturation. Other approaches to mutagenesis involve modification of adjacent dibasic amino acid residues to enhance expression in yeast systems in which KEX2 protease activity is present. Generally, substitutions should be made conservatively; i.e., the most preferred substitute amino acids are those having physicochemical characteristics resembling those of the residue to be replaced. Similarly, when a deletion or insertion strategy is adopted, the potential effect of the deletion or insertion on biological activity should be considered.

Subunits of IL-4R may be constructed by deleting terminal or internal residues or sequences. Particularly preferred subunits include those in which the transmembrane region and intracellular domain of IL-4R are deleted or substituted with hydrophilic residues to facilitate secretion of the receptor into the cell culture medium. The resulting protein is a soluble IL-4R molecule which may retain its ability to bind IL-4. Particular examples of soluble IL-4R include polypeptides having substantial identity to soluble murine IL-4R (1-208), soluble human IL-4R (10207) and soluble human IL-4R (1-198), all of which retain the biological activity of soluble human IL-4R (1-207). Chimeric polypeptides comprising fragments of human and murine IL-4R may also be constructed, for example, IL-4R (1-197) Pro Ser Asn Glu Asn Leu, which is comprised of the sequence of amino acids 1-197 of human IL-4R followed by the N-terminal six amino acids of soluble murine IL-4R clone 18. This polypeptide has been found to retain the biological activity of soluble IL-4R (1-207).

Mutations in nucleotide sequences constructed for expression of analog IL-4Rs must, of course, preserve the reading frame phase of the coding sequences and preferably will not create complementary regions that could hybridize to produce secondary mRNA structures, such as loops or hairpins, which would adversely affect translation of the receptor mRNA. Although a mutation site may be predetermined, it is not necessary that the nature of the mutation per se be predetermined. For example, in order to select for optimum characteristics of mutants at a given site, random mutagenesis may be conducted at the target codon and the expressed IL-4R mutants screened for the desired activity.

Not all mutations in the nucleotide sequence which encodes IL-4R will be expressed in the final product, for example, nucleotide substitutions may be made to enhance expression, primarily to avoid secondary structure loops in

the transcribed mRNA (see EPA 75,444A, incorporated herein by reference), or to provide codons that are more readily translated by the selected host, e.g., the well-known *E. coli* preference codons for *E. coli* expression.

Mutations can be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes an analog having the desired amino acid insertion, substitution, or deletion.

Alternatively, oligonucleotide-directed site-specific mutagenesis procedures can be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or insertion required. Exemplary methods of making the alterations set forth above are disclosed by Walder et al. (*Gene* 42:133, 1986); Bauer et al. (*Gene* 37:73, 1985); Craik (*Bio Techniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering: Principles and Methods*, Plenum Press, 1981); and U.S. Pat. Nos. 4,518,584 and 4,737,462, which are incorporated by reference herein. Expression of Recombinant IL-4R T

he present invention provides recombinant expression vectors which include synthetic or cDNA-derived DNA fragments encoding mammalian IL-4R or bioequivalent analogs operably linked to suitable transcriptional or translational regulatory elements derived from mammalian, microbial, viral or insect genes. Such regulatory elements include a transcriptional promoter, an optional operator sequence to control transcription, a sequence encoding suitable mRNA ribosomal binding sites, and sequences which control the termination of transcription and translation, as described in detail below. The ability to replicate in a host, usually conferred by an origin of replication, and a selection gene to facilitate recognition of transformants may additionally be incorporated. DNA regions are operably linked when they are functionally related to each other. For example, DNA for a signal peptide (secretory leader) is operably linked to DNA for a polypeptide if it is expressed as a precursor which participates in the secretion of the polypeptide; a promoter is operably linked to a coding sequence if it controls the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to permit translation. Generally, operably linked means contiguous and, in the case of secretory leaders, contiguous and in reading frame.

DNA sequences encoding mammalian IL-4 receptors which are to be expressed in a microorganism will preferably contain no introns that could prematurely terminate transcription of DNA into mRNA; however, premature termination of transcription may be desirable, for example, where it would result in mutants having advantageous C-terminal truncations, for example, deletion of a transmembrane region to yield a soluble receptor not bound to the cell membrane. Due to code degeneracy, there can be considerable variation in nucleotide sequences encoding the same amino acid sequence; exemplary DNA embodiments are those corresponding to the nucleotide sequences shown in the Figures. Other embodiments include sequences capable of hybridizing to the sequences of the Figures under moderately stringent conditions (50° C., 2xSSC) and other sequences hybridizing or degenerate to those described above, which encode biologically active IL-4 receptor polypeptides.

DNA which codes for soluble IL-4R proteins may be isolated using the cloning techniques described in the examples or may be made by constructing cDNAs which encode only the extracellular domain of IL-4 receptor

(devoid of a transmembrane region) using well-known methods of mutagenesis. Soluble forms of human IL-4 receptor are not yet known to exist and must therefore be constructed from isolated recombinant IL-4 receptor cDNAs. cDNAs which encode sIL-4R may be constructed, for example, by truncating a cDNA encoding the full length IL-4 receptor 5' of the transmembrane region, ligating synthetic oligonucleotides to regenerate truncated portions of the extracellular domain, if necessary, and providing a stop codon to terminate transcription. DNA sequences encoding the soluble IL-4 receptor proteins can be assembled from cDNA fragments and short oligonucleotide linkers, or from a series of oligonucleotides, to provide a synthetic gene which is capable of being expressed in a recombinant transcriptional unit. Such DNA sequences are preferably provided in the form of an open reading frame uninterrupted by internal nontranslated sequences, or introns, which are typically present in eukaryotic genes. Genomic DNA containing the relevant sequences could also be used. Sequences of non-translated DNA may be present 5' or 3' from the open reading frame, where the same do not interfere with manipulation or expression of the coding regions.

Transformed host cells are cells which have been transformed or transfected with IL-4R vectors constructed using recombinant DNA techniques. Transformed host cells ordinarily express IL-4R, but host cells transformed for purposes of cloning or amplifying IL-4R DNA do not need to express IL-4R. Expressed IL-4R will be deposited in the cell membrane or secreted into the culture supernatant, depending on the IL-4R DNA selected. Suitable host cells for expression of mammalian IL-4R include prokaryotes, yeast or higher eukaryotic cells under the control of appropriate promoters. Prokaryotes include gram negative or gram positive organisms, for example *E. coli* or bacilli. Higher eukaryotic cells include established cell lines of mammalian origin as described below. Cell-free translation systems could also be employed to produce mammalian IL-4R using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described by Pouwels et al. (*Cloning Vectors: A Laboratory Manual*, Elsevier, New York, 1985), the relevant disclosure of which is hereby incorporated by reference.

Prokaryotic expression hosts may be used for expression of IL-4Rs that do not require extensive proteolytic and disulfide processing. Prokaryotic expression vectors generally comprise one or more phenotypic selectable markers, for example a gene encoding proteins conferring antibiotic resistance or supplying an autotrophic requirement, and an origin of replication recognized by the host to ensure amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice.

Useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, Wis., USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. *E. coli* is typically transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species (Bolivar et al., *Gene* 2:95, 1977). pBR322 contains genes for

ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells.

Promoters commonly used in recombinant microbial expression vectors include the β -lactamase (penicillinase) and lactose promoter system (Chang et al., *Nature* 275:615, 1978; and Goeddel et al., *Nature* 281:544, 1979), the tryptophan (trp) promoter system (Goeddel et al., *Nucl. Acids Res.* 8:4057, 1980; and EPA 36,776) and tac promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, p. 412, 1982). A particularly useful bacterial expression system employs the phage λ P_L promoter and cl857ts thermolabile repressor. Plasmid vectors available from the American Type Culture Collection which incorporate derivatives of the λ P_L promoter include plasmid pHUB2, resident in *E. coli* strain JMB9 (ATCC 37092) and pPLc28, resident in *E. coli* RR1 (ATCC 53082).

Recombinant IL-4R proteins may also be expressed in yeast hosts, preferably from the *Saccharomyces* genus, such as *S. cerevisiae*. Yeast of other genera, such as *Pichia* or *Kluyveromyces* may also be employed. Yeast vectors will generally contain an origin of replication from the 2μ yeast plasmid or an autonomously replicating sequence (ARS), promoter, DNA encoding IL-4R, sequences for polyadenylation and transcription termination and a selection gene. Preferably, yeast vectors will include an origin of replication and selectable marker permitting transformation of both yeast and *E. coli*, e.g., the ampicillin resistance gene of *E. coli* and *S. cerevisiae* trp1 gene, which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, and a promoter derived from a highly expressed yeast gene to induce transcription of a structural sequence downstream. The presence of the trp1 lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan.

Suitable promoter sequences in yeast vectors include the promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* 255:2073, 1980) or other glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* 7:149, 1968; and Holland et al., *Biochem.* 17:4900, 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Suitable vectors and promoters for use in yeast expression are further described in Hitzeman, EPA 73,657.

Preferred yeast vectors can be assembled using DNA sequences from pBR322 for selection and replication in *E. coli* (Amp^r gene and origin of replication) and yeast DNA sequences including a glucose-repressible ADH2 promoter and α -factor secretion leader. The ADH2 promoter has been described by Russell et al. (*J. Biol. Chem.* 258:2674, 1982) and Beier et al. (*Nature* 300:724, 1982). The yeast α -factor leader, which directs secretion of heterologous proteins, can be inserted between the promoter and the structural gene to be expressed. See, e.g., Kurjan et al., *Cell* 30:933, 1982; and Bitter et al., *Proc. Natl. Acad. Sci. USA* 81:5330, 1984. The leader sequence may be modified to contain, near its 3' end, one or more useful restriction sites to facilitate fusion of the leader sequence to foreign genes.

Suitable yeast transformation protocols are known to those of skill in the art; an exemplary technique is described by Hinnen et al., *Proc. Natl. Acad. Sci. USA* 75:1929, 1978, selecting for Trp⁺ transformants in a selective medium consisting of 0.67% yeast nitrogen base, 0.5% casamino acids, 20% glucose, 10 μ g/ml adenine and 20 μ g/ml uracil.

Host strains transformed by vectors comprising the ADH2 promoter may be grown for expression in a rich medium consisting of 1% yeast extract, 2% peptone, and 1% glucose supplemented with 80 μ g/ml adenine and 80 μ g/ml uracil. Derepression of the ADH2 promoter occurs upon exhaustion of medium glucose. Crude yeast supernatants are harvested by filtration and held at 4° C. prior to further purification.

Various mammalian or insect cell culture systems can be employed to express recombinant protein. Baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow and Summers, *Bio/Technology* 6:47 (1988). Examples of suitable mammalian host cell lines include the COS-7 lines of monkey kidney cells, described by Gluzman (*Cell* 23:175, 1981), and other cell lines capable of expressing an appropriate vector including, for example, L cells, C127, 3T3, Chinese hamster ovary (CHO), HeLa and BHK cell lines. Mammalian expression vectors may comprise nontranscribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, and other 5' or 3' flanking nontranscribed sequences, and 5' or 3' untranslated sequences, such as necessary ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, and transcriptional termination sequences.

The transcriptional and translational control sequences in expression vectors to be used in transforming vertebrate cells may be provided by viral sources. For example, commonly used promoters and enhancers are derived from Polyoma, Adenovirus 2, Simian Virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide the other genetic elements required for expression of a heterologous DNA sequence. The early and late promoters are particularly useful because both are obtained easily from the virus as a fragment which also contains the SV40 viral origin of replication (Fiers et al., *Nature* 273:113, 1978). Smaller or larger SV40 fragments may also be used, provided the approximately 250 bp sequence extending from the Hind III site toward the Bgl I site located in the viral origin of replication is included. Further, mammalian genomic IL-4R promoter, control and/or signal sequences may be utilized, provided such control sequences are compatible with the host cell chosen. Additional details regarding the use of mammalian high expression vectors to produce a recombinant mammalian IL-4 receptor are provided in Example 8 below. Exemplary vectors can be constructed as disclosed by Okayama and Berg (*Mol. Cell. Biol.* 3:280, 1983).

A useful system for stable high level expression of mammalian receptor cDNAs in C127 murine mammary epithelial cells can be constructed substantially as described by Cosman et al. (*Mol. Immunol.* 23:935, 1986).

A particularly preferred eukaryotic vector for expression of IL-4R DNA is disclosed below in Example 2. This vector, referred to as pCAV/NOT, was derived from the mammalian high expression vector pDC201 and contains regulatory sequences from SV40, adenovirus-2, and human cytomegalovirus. pCAV/NOT containing a human IL-7 receptor insert has been deposited with the American Type Culture Collection (ATCC) under deposit accession number 68014.

Purification of IL-4 Receptors

Purified mammalian IL-4 receptors or analogs are prepared by culturing suitable host/vector systems to express the recombinant translation products of the DNAs of the present invention, and purifying IL-4 receptor from the culture media or cell extracts.

For example, supernatants from systems which secrete recombinant protein into culture media can be first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a suitable purification matrix. For example, a suitable affinity matrix can comprise an IL-4 or fectin or antibody molecule bound to a suitable support. Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups are preferred.

Finally, one or more reversed-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify an IL-4R composition. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a homogeneous recombinant protein.

Recombinant protein produced in bacterial culture is usually isolated by initial extraction from cell pellets, followed by one or more concentration, salting-out, aqueous ion exchange or size exclusion chromatography steps. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps. Microbial cells employed in expression of recombinant mammalian IL-4R can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Fermentation of yeast which express mammalian IL-4R as a secreted protein greatly simplifies purification. Secreted recombinant protein resulting from a large-scale fermentation can be purified by methods analogous to those disclosed by Urdal et al. (*J. Chromatog.* 296:171, 1984). This reference describes two sequential, reversed-phase HPLC steps for purification of recombinant human IL-2 on a preparative HPLC column.

Human IL-4R synthesized in recombinant culture is characterized by the presence of non-human cell components, including proteins, in amounts and of a character which depend upon the purification steps taken to recover human IL-4R from the culture. These components ordinarily will be of yeast, prokaryotic or non-human higher eukaryotic origin and preferably are present in innocuous contaminant quantities, on the order of less than about 1 percent by weight. Further, recombinant cell culture enables the production of IL-4R free of proteins which may be normally associated with IL-4R as it is found in nature in its species of origin, e.g. in cells, cell exudates or body fluids.

Administration of Soluble IL-4 Receptor Compositions

The present invention provides methods of using therapeutic compositions comprising an effective amount of soluble IL-4 receptor proteins and a suitable diluent and carrier, and methods for suppressing IL-4-dependent immune responses in humans comprising administering an effective amount of soluble IL-4 receptor protein.

For therapeutic use, purified soluble IL-4 receptor protein is administered to a patient, preferably a human, for treatment in a manner appropriate to the indication. Thus, for example, soluble IL-4 receptor protein compositions administered to suppress immune function can be given by bolus injection, continuous infusion, sustained release from implants, or other suitable technique. Typically, a soluble

IL-4 receptor therapeutic agent will be administered in the form of a composition comprising purified protein in conjunction with physiologically acceptable carriers, excipients or diluents. Such carriers will be nontoxic to recipients at the dosages and concentrations employed. Ordinarily, the preparation of such compositions entails combining the IL-4R with buffers, antioxidants such as ascorbic acid, low molecular weight (less than about 10 residues) polypeptides, proteins, amino acids, carbohydrates including glucose, sucrose or dextrans, chelating agents such as EDTA, glutathione and other stabilizers and excipients. Neutral buffered saline or saline mixed with conspecific serum albumin are exemplary appropriate diluents. Preferably, product is formulated as a lyophilizate using appropriate excipient solutions (e.g., sucrose) as diluents. Appropriate dosages can be determined in trials; generally, soluble IL-4 receptor dosages of from about 1 ng/kg/day to about 10 mg/kg/day, more preferably from about 500 ng/kg/day to about 5 mg/kg/day, and most preferably from about 5 ug/kg/day to about 2 mg/kg/day, are appropriate for inducing a biological effect. The amount and frequency of administration will depend, of course, on such factors as the nature and severity of the indication being treated, the desired response, the condition of the patient, and so forth.

Soluble IL-4R proteins are administered for the purpose of inhibiting IL-4 dependent responses, such as suppressing immune responses in a human. A variety of diseases or conditions are caused by IL-4 dependent immune responses as determined by the ability of sIL-4R to inhibit the response. Soluble IL-4R compositions may be used, for example, to regulate the function of B cells. Soluble IL-4R inhibits IL-4 dependent B cell proliferation and isotype specific (IgG1 and IgE) secretions. sIL-4R may therefore be used to suppress IgE antibody formation in the treatment of IgE-induced immediate hypersensitivity reactions, such as allergic rhinitis (common hay fever), bronchial asthma, atopic dermatitis and gastrointestinal food allergy.

sIL-4R compositions may also be used to regulate the function of T cells. Although T cell dependent functions were formerly thought to be mediated principally by IL-2, recent studies have shown that under some circumstances T cell growth and proliferation can be mediated by growth factors such as IL-4. Examples 20 through 23 below, for example, indicate that sIL-4R suppresses or inhibits T-cell dependent responses to alloantigen. A variety of diseases or conditions are caused by an immune response to alloantigen, including allograft rejection and graft-versus-host reaction. In alloantigen-induced immune responses, sIL-4R suppresses lymphoproliferation and inflammation which result upon activation of T cells. sIL-4R has therefore been shown to be potentially effective in the clinical treatment of, for example, rejection of allografts (such as skin, kidney, heart, lung liver and pancreas transplants), and graft-versus-host reactions in patients who have received bone marrow transplants.

sIL-4R may also be used in clinical treatment of autoimmune dysfunctions, such as rheumatoid arthritis, diabetes, which are dependent upon the activation of T cells against antigens not recognized as being indigenous to the host.

Because of the primary role IL-2 plays in the proliferation and differentiation of T cells, combination therapy using both IL-4 and IL-2 may be used in the treatment of T cell dependent dysfunctions. Use in conjunction with other soluble cytokine receptors, e.g., IL-1 receptor, is also contemplated.

The following examples are offered by way of illustration, and not by way of limitation.

EXAMPLES

Example 1

Binding Assays for IL-4 Receptor

A. Radiolabelling of IL-4. Recombinant murine and human IL-4 were expressed in yeast and purified to homogeneity as described by Park, et al., *Proc. Natl. Acad. Sci. USA* 84:5267 (1987) and Park et al., *J. Exp. Med.* 166:476 (1987), respectively. The purified protein was radiolabeled using a commercially available enzymobead radioiodination reagent (BioRad). In this procedure 2.5 μ g rIL-4 in 50 μ l 0.2 M sodium phosphate, pH 7.2 are combined with 50 μ l enzymobead reagent, 2 MCi of sodium iodide in 20 μ l of 0.05 M sodium phosphate pH 7.0 and 10 μ l of 2.5% b-D-glucose. After 10 min at 25° C., sodium azide (10 μ l of 50 mM) and sodium metabisulfite (10 μ l of 5 mg/ml) were added and incubation continued for 5 min. at 25° C. The reaction mixture was fractionated by gel filtration on a 2 ml bed volume of Sephadex® G-25 (Sigma) equilibrated in Roswell Park Memorial Institute (RPMI) 1640 medium containing 2.5% (w/v) bovine serum albumin (BSA), 0.20% (w/v) sodium azide and 20 mM Hepes pH 7.4 (binding medium). The final pool of 125 I-IL-4 was diluted to a working stock solution of 2×10^{-8} M in binding medium and stored for up to one month at 4° C. without detectable loss of receptor binding activity. The specific activity is routinely in the range of $1\text{--}2 \times 10^{16}$ cpm/mmol IL-4.

B. Binding to Adherent Cells. Binding assays done with cells grown in suspension culture (i.e., CTLL and CTLL-19.4) were performed by a phthalate oil separation method (Dower et al., *J. Immunol.* 132:751, 1984) essentially as described by Park et al., *J. Biol. Chem.* 261:4177, 1986 and Park et al., supra. Binding assays were also done on COS cells transfected with a mammalian expression vector containing cDNA encoding an IL-4 receptor molecule. For Scatchard analysis of binding to adherent cells, COS cells were transfected with plasmid DNA by the method of Luthman et al., *Nucl. Acids. Res.* 11:1295, 1983, and McCutchan et al., *J. Natl. Cancer Inst.* 41:351, 1968. Eight hours following transfection, cells were trypsinized, and reseeded in six well plates (Costar, Cambridge, Mass.) at a density of 1×10^4 COS-IL-4 receptor transfectants/well mixed with 5×10^5 COS control transfected cells as carriers. Two days later monolayers were assayed for 125 I-IL-4 binding at 4° C. essentially by the method described by Park et al., *J. Exp. Med.* 166:476, 1987. Nonspecific binding of 125 I-IL-4 was measured in the presence of a 200-fold or greater molar excess of unlabeled IL-4. Sodium azide (0.2%) was included in all binding assays to inhibit internalization of 125 I-IL-4 by cells at 37° C.

For analysis of inhibition of binding by soluble IL-4R, supernatants from COS cells transfected with recombinant IL-4R constructs were harvested three days after transfection. Serial two-fold dilutions of conditioned media were pre-incubated with 3×10^{-10} M 125 I-IL-4 (having a specific activity of about 1×10^{16} cpm/mmol) for one hour at 37° C. prior to the addition of 2×10^6 CTLL cells. Incubation was continued for 30 minutes at 37° C. prior to separation of free and cell-bound murine 125 I-IL-4.

C. Solid Phase Binding Assays. The ability of IL-4 receptor to be stably adsorbed to nitrocellulose from detergent extracts of CTLL 19.4 cells yet retain IL-4 binding activity provided a means of monitoring purification. One ml aliquots of cell extracts (see Example 3), IL-4 affinity column fractions (see Example 4) or other samples are placed on dry BA85/21 nitrocellulose membranes (Schleicher and Schuell, Keene, N.H.) and allowed to dry. The membranes are incubated in tissue culture dashes for 30

minutes in Tris (0.05 M) buffered saline (0.15 M) pH 7.5 containing 3% w/v BSA to block nonspecific binding sites. The membrane is then covered with 4×10^{-11} M 125 I-IL-4 in PBS+3% BSA with or without a 200 fold molar excess of unlabeled IL-4 and incubated for 2 hr at 4° C. with shaking. At the end of this time, the membranes are washed 3 times in PBS, dried and placed on Kodak X-Omat™ AR film for 18 hr at -70° C.

Example 2

Selection of CTLL Cells With High IL-4 Receptor Expression by Fluorescence Activated Cell Sorting (FACS)

The preferred cell line for obtaining high IL-4 receptor selection is CTLL, a murine IL-2 dependent cytotoxic T cell line (ATCC TIB 214) which typically exhibits 2,000 to 5,000 IL-4 receptors per cell and responds to IL-4 by short-term proliferation. To obtain higher levels of IL-4 receptor expression, CTLL cells (parent cells) were subjected to multiple rounds of fluorescence-activated cell sorting with labeled IL-4. A highly fluorescent derivative of IL-4 was derived by conjugating rmIL-4 fluorescein hydrazide to periodate oxidized sugar moieties of IL-4 which was produced in yeast as described by Park et al., *Proc. Natl. Acad. Sci. USA* 84:1669 (1987). The fluorescein-conjugated IL-4 was prepared by combining aliquots of hyperglycosylated rmIL-4 (300 μ g in 300 μ l of 0.1 M citrate-phosphate buffer, pH 5.5) with 30 μ l of 10 mM sodium m-periodate (Sigma), freshly prepared in 0.1 M citrate-phosphate, pH 5.5 and the mixture incubated at 4° C. for 30 minutes in the dark. The reaction was quenched with 30 μ l of 0.1 M glycerol and dialyzed for 18 hours at 4° C. against 0.1 M citrate-phosphate pH 5.5. Following dialysis, a 1/10 volume of 100 mM 5-(((2-(carbohydrazino)methyl)thio)acetyl)-aminofluorescein (Molecular Probes, Eugene Oreg.) dissolved in DMSO was added to the sample and incubated at 25° C. for 30 minutes. The IL-4-fluorescein was then exhaustively dialyzed at 4° C. against PBS, pH 7.4 and protein concentration determined by amino acid analysis. The final product was stored at 4° C. following the addition of 1% (w/v) BSA and sterile filtration.

CTLL cells (5×10^6) were then incubated for 30 min at 37° C. in 150 μ l PBS+1% BSA containing 1×10^{-9} M IL-4-fluorescein under sterile conditions. The mixture was then chilled to 4° C., washed once in a large volume of PBS+1% BSA and sorted using an EPICS® C flow cytometer (Coulter Instruments). The cells providing the highest level fluorescence signal (top 1.0%) were collected in bulk and the population expanded in liquid cell culture and subjected to additional rounds of sorting as described below. Alternatively, for single cell cloning, cells exhibiting a fluorescence signal in the top 1.0% were sorted into 96 well tissue culture microtiter plates at 1 cell per well.

Progress was monitored by doing binding assays with 125 I-IL-4 following each round of FACS selection. Unsorted CTLL cells (CTLL parent) typically exhibited 1000–2000 IL-4 receptors per cell. CTLL cells were subjected to 19 rounds of FACS selection. The final CTLL cells selected (CTLL-19) exhibited 5×10^5 to 1×10^6 IL-4 receptors per cell. At this point the CTLL-19 population was subjected to EPICS® C-assisted single cell cloning and individual clonal populations were expanded and tested for 125 I-IL-4 binding. A single clone, designated CTLL-19.4, exhibited 1×10^6 IL-4 receptors per cell and was selected for purification and cloning studies. While the calculated apparent K_a values are similar for the two lines, CTLL-19.4 expresses approximately 400-fold more receptors on its surface than does the CTLL parent.

Detergent Extraction of CTLL Cells

CTLL 19.4 cells were maintained in RPMI 1640 containing 10% fetal bovine serum, 50 U/ml penicillin, 50 µg/ml streptomycin and 10 ng/ml of recombinant human IL-2. Cells were grown to 5×10⁵ cells/ml in roller bottles, harvested by centrifugation, washed twice in serum free DMEM and sedimented at 2000×g for 10 minutes to form a packed pellet (about 2×10⁸ cells/ml). To the pellet was added an equal volume of PBS containing 1% Triton® X-100 and a cocktail of protease inhibitors (2 mM phenylmethylsulfonylfluoride, 10 µM pepstatin, 10 µM leupeptin, 2 mM o-phenanthroline and 2 mM EGTA). The cells were mixed with the extraction buffer by vigorous vortexing and the mixture incubated on ice for 20 minutes after which the mixture was centrifuged at 12,000×g for 20 minutes at 8° C. to remove nuclei and other debris. The supernatant was either used immediately or stored at -70° C. until use.

IL-4 Receptor Purification by IL-4 Affinity Chromatography

In order to obtain sufficient quantities of murine IL-4R to determine its N-terminal sequence or to further characterize human IL-4R, protein obtained from the detergent extraction of cells was further purified by affinity chromatography. Recombinant murine or human IL-4 was coupled to Affigel®-10 (BioRad) according to the manufacturer's suggestions. For example, to a solution of IL-4 (3.4 mg/ml in 0.4 ml of 0.1 M Hepes pH 7.4) was added 1.0 ml of washed Affigel®-10. The solution was rocked overnight at 4° C. and an aliquot of the supernatant tested for protein by a BioRad protein assay per the manufacturers instructions using BSA as a standard. Greater than 95% of the protein had coupled to the gel, suggesting that the column had a final load of 1.3 mg IL-4 per ml gel. Glycine ethyl ester was added to a final concentration of 0.05 M to block any unreacted sites on the gel. The gel was washed extensively with PBS-1% Triton® followed by 0.1 Glycine-HCl, pH 3.0. A 0.8×4.0 cm column was prepared with IL-4-coupled Affigel® prepared as described (4.0 ml bed volume) and washed with PBS containing 1% Triton® X-100 for purification of murine IL-4R. Alternatively, 50 µl aliquots of 20% suspension of IL-4-coupled Affigel® were incubated with ³⁵S-cysteine/methionine-labeled cell extracts for small-scale affinity purifications and gel electrophoresis.

Aliquots (25 ml) of detergent extracted IL-4 receptor bearing CTLL 19.4 cells were slowly applied to the murine IL-4 affinity column at 4° C. (flow rate of 3.0 ml/hr). The column was then washed sequentially with PBS containing 1% Triton® X-100, RIPA buffer (0.05 M Tris, 0.15 M NaCl, 1% NP-40, 1% deoxycholate and 0.1% SDS), PBS containing 0.1% Triton® X-100 and 10 mM ATP, and PBS with 1% Triton® X-100 to remove all contaminating material except the mIL-4R. The column was then eluted with pH 3.0 glycine HCl buffer containing 0.1% Triton® X-100 to remove the IL-4R and washed subsequently with PBS containing 0.1% Triton® X-100. One milliliter fractions were collected for the elution and 2 ml fractions collected during the wash. Immediately following elution, samples were neutralized with 80 µl of 1 M Hepes, pH 7.4. The presence of receptor in the fractions was detected by the solid phase binding assay as described above, using ¹²⁵I-

labeled IL-4. Aliquots were removed from each fraction for analysis by SDS-PAGE and the remainder frozen at -70° C. until use. For SDS-PAGE, 40 µl of each column fraction was added to 40 µl of 2×SDS sample buffer (0.125 M Tris HCl pH 6.8, 4% SDS, 20% glycerol, 10% 2-mercaptoethanol). The samples were placed in a boiling water bath for 3 minutes and 80 µl aliquots applied to sample wells of a 10% polyacrylamide gel which was set up and run according to the method of Laemmli (*Nature* 227:680, 1970). Following electrophoresis, gels were silver stained as previously described by Urdal et al. (*Proc. Natl. Acad. Sci. USA* 81:6481, 1984).

Purification by the foregoing process permitted identification by silver staining of polyacrylamide gels of two mIL-4R protein bands averaging 45-55 kDa and 30-40 kDa that were present in fractions exhibiting IL-4 binding activity. Experiments in which the cell surface proteins of CTLL-19.4 cells were radiolabeled and ¹²⁵I-labeled receptor was purified by affinity chromatography suggested that these two proteins were expressed on the cell surface. The ratio of the lower to higher molecular weight bands increased upon storage of fractions at 4° C., suggesting a precursor product relationship, possibly due to slow proteolytic degradation. The mIL-4 receptor protein purified by the foregoing process remains capable of binding IL-4, both in solution and when adsorbed to nitrocellulose.

Sequencing of IL-4 Receptor Protein

CTLL 19.4 mIL-4 receptor containing fractions from the mIL-4 affinity column purification were prepared for amino terminal protein sequence analysis by fractionating on an SDS-PAGE gel and then transferred to a PVDF membrane. Prior to running the protein fractions on polyacrylamide gels, it was first necessary to remove residual detergent from the affinity purification process. Fractions containing proteins bound to the mIL-4 affinity column from three preparations were thawed and concentrated individually in a speed vac under vacuum to a final volume of 1 ml. The concentrated fractions were then adjusted to pH 2 by the addition of 50% (v/v) TFA and injected onto a Brownlee RP-300 reversed-phase HPLC column (2.1×30 mm) equilibrated with 0.1% (v/v) TFA in H₂O at a flow rate of 200 µl/min running on a Hewlett Packard Model 1090M HPLC. The column was washed with 0.1% TFA in H₂O for 20 minutes post injection. The HPLC column containing the bound protein was then developed with a gradient as follows:

Time	% Acetonitrile in 0.1% TFA
0	0
5	30
15	30
25	70
30	70
35	100
40	0

1 ml fractions were collected every five minutes and analyzed for the presence of protein by SDS PAGE followed by silver staining.

Each fraction from the HPLC run was evaporated to dryness in a speed vac and then resuspended in Laemmli reducing sample buffer, prepared as described by Laemmli,

U.K. *Nature* 227:680, 1970. Samples were applied to a 5–20% gradient Laemmli SDS gel and run at 45 mA until the dye front reached the bottom of the gel. The gel was then transferred to PVDF paper and stained as described by Matsudaira, *J. Biol. Chem.* 262:10035, 1987. Staining bands were clearly identified in fractions from each of the three preparations at approximately 30,000 to 40,000 M_r.

The bands from the previous PVDF blotting were excised and subjected to automated Edman degradation on an Applied Biosystems Model 477A Protein Sequencer essentially as described by March et al. (*Nature* 315:641, 1985), except that PTH amino acids were automatically injected and analyzed on line with an Applied Biosystems Model 120A HPLC using a gradient and detection system supplied by the manufacturer. The following amino terminal sequence was determined from the results of sequencing: NH₂-Ile-Lys-Val-Leu-Gly-Glu-Pro-Thr-(Cys/Asn)-Phe-Ser-Asp-Tyr-Ile. Position 9 was assigned as a cysteine or glycosylated asparagine owing to the lack of an observable PTH-amino acid signal in the cycle. The bands from the second preparation used for amino terminal sequencing were treated with CNBr using the in situ technique described by March et al. (*Nature* 315: 641, 1985) to cleave the protein after internal methionine residues. Sequencing of the resulting cleavage products yielded the following data, indicating that the CNBr cleaved the protein after two internal methionine residues:

Cycle	Residues Observed
1	Val, Ser
2	Gly, Leu
3	Ile, Val
4	Tyr, Ser
5	Arg, Tyr
6	Glu, Thr
7	Asp, Ala
8	Asn, Leu
9	Pro, Val
10	Ala
11	Glu, Val
12	Phe, Gly
13	Ile, Asn
14	Val, Gln
15	Tyr, Ile
16	Lys, Asn
17	Val, Thr
18	Thr, Gly

When compared with the protein sequences derived from clones 16 and 18 (see FIG. 2), the sequences matched as follows:

151018

Sequence 1: (Met)-Val-Asn-Ile-Ser-Arg-Glu-Asp-Asn-Pro-Ala-Glu-Phe-Ile-Val-Tyr-Asn-Val-Thr

151018

Sequence 2: (Met)-Ser-Gly-Val-Tyr-Tyr-Thr-Ala-Arg-Val-Arg-Val-Arg-Ser-Gln-Ile-Leu-Thr-Gly

Identical matches were found for all positions of sequence 1 except Asn(2) and sequence 2, except Arg at positions 8, 10, and 12, Ser at position 13, and Leu at position 16. The above sequences correspond to amino acid residues 137–154 and 169–187 of FIG. 2A.

In addition, the amino terminal sequence matched a sequence derived from the clone with position 9 being defined as a Cys.

The above data support the conclusion that clones 16 and 18 are derived from the message for the IL-4 receptor.

Example 6

Synthesis of Hybrid-subtracted cDNA Probe

In order to screen a library for clones encoding a murine IL-4 receptor, a highly enriched IL-4 receptor cDNA probe was obtained using a subtractive hybridization strategy. Polyadenylated (polyA⁺) mRNA was isolated from two similar cell lines, the parent cell line CTLL (which expresses approximately 2,000 receptors per cell) and the sorted cell line CTLL 19.4 (which expresses 1×10⁶ receptors per cell). The mRNA content of these two cell lines is expected to be identical except for the relative level of IL-4 receptor mRNA. A radiolabeled single-stranded cDNA preparation was then made from the mRNA of the sorted cell line CTLL 19.4 by reverse transcription of polyadenylated mRNA from CTLL 19.4 cells by a procedure similar to that described by Maniatis et al., *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratory, New York, 1982). Briefly, polyA⁺ mRNA was purified as described by March et al. (*Nature* 315:641–647, 1985) and copied into cDNA by reverse transcriptase using oligo dT as a primer. To obtain a high level of ³²P-labeling of the cDNA, 100 μCi of ³²P-dCTP (s.a.=3000 Ci/mmol) was used in a 50 μl reaction with non-radioactive dCTP at 10 μM. After reverse transcription at 42° C. for 2 hours, EDTA was added to 20 mM and the RNA was hydrolyzed by adding NaOH to 0.2 M and incubating the cDNA mixture at 68° C. for 20 minutes. The single-stranded cDNA was extracted with a phenol/chloroform (50/50) mixture previously equilibrated with 10 mM Tris-Cl, 1 mM EDTA. The aqueous phase was removed to a clean tube and made alkaline again by the addition of NaOH to 0.5 M. The cDNA was then size-fractionated by chromatography on a 6 ml Sephadex® G50 column in 30 mM NaOH and 1 mM EDTA to remove small molecular weight contaminants.

The resulting size-fractionated cDNA generated from the sorted CTLL 19.4 cells was then hybridized with an excess of mRNA from the unsorted parental CTLL cells by ethanol-precipitating the cDNA from CTLL 19.4 cells with 30 μg of polyA⁺ mRNA isolated from unsorted CTLL cells, resuspending in 16 μl of 0.25 M NaPO₄, pH 6.8, 0.2% SDS, 2 mM EDTA and incubating for 20 hours at 68° C. The cDNAs from the sorted CTLL 19.4 cells that are complementary to mRNAs from the unsorted CTLL cells form double stranded cDNA/mRNA hybrids, which can then be separated from the single stranded cDNA based on their different binding affinities on hydroxyapatite. The mixture was diluted with 30 volumes of 0.02 M NaPO₄, pH 6.8, bound to hydroxya-

patite at room temperature, and single-stranded cDNA was then eluted from the resin with 0.12 M NaPO₄, pH 6.8, at 60° C., as described by Sims et al., *Nature* 312:541, 1984. Phosphate buffer was then removed by centrifugation over 2 ml Sephadex® G50 spin columns in water. This hybrid subtraction procedure removes a majority of common sequences between CTLL 19.4 and unsorted CTLL cells, and leaves a single-stranded cDNA pool enriched for radio-

labeled IL-4 receptor cDNA which can be used to probe a cDNA library (as described below).

Example 7

Synthesis of cDNA Library and Plaque Screening

A cDNA library was constructed from polyadenylated mRNA isolated from CTLL 19.4 cells using standard techniques (Gubler, et al., *Gene* 25:263, 1983; Ausubel et al., eds., *Current Protocols in Molecular Biology*, Vol. 1, 1987). After reverse transcription using oligo dT as primer, the single-stranded cDNA was rendered double-stranded with DNA polymerase I, blunt-ended with T4 DNA polymerase, methylated with EcoR I methylase to protect EcoR I cleavage sites within the cDNA, and ligated to EcoR I linkers. The resulting constructs were digested with EcoR I to remove all but one copy of the linkers at each end of the cDNA, and ligated to an equimolar concentration of EcoR I cut and dephosphorylated λ ZAP® arms and the resulting ligation mix was packaged in vitro (Gigapack®) according to the manufacturer's instructions. Other suitable methods and reagents for generating cDNA libraries in λ phage vectors are described by Huynh et al., *DNA Cloning Techniques: A Practical Approach*, IRL Press, Oxford (1984); Meissner et al., *Proc. Natl. Acad. Sci. USA* 84:4171 (1987), and Ausubel et al., supra. λ ZAP® is a phage λ cloning vector similar to λ gt11 (U.S. Pat. No. 4,788,135) containing plasmid sequences from pUC19 (Norlander et al., *Gene* 26:101, 1987), a polylinker site located in a lacZ gene fragment, and an f1 phage origin of replication permitting recovery of ssDNA when host bacteria are superinfected with f1 helper phage. DNA is excised in the form of a plasmid comprising the foregoing elements, designated Bluescript®. Gigapack® is a sonicated *E. coli* extract used to package λ phage DNA. λ ZAP®, Bluescript®, and Gigapack® are registered trademarks of Stratagene, San Diego, Calif., USA.

The radiolabeled hybrid-subtracted cDNA from Example 6 was then used as a probe to screen the cDNA library. The amplified library was plated on BB4 cells at a density of 25,000 plaques on each of 20 150 mm plates and incubated overnight at 37° C. All manipulations of λ ZAP® and excision of the Bluescript® plasmid were as described by Short et al., (*Nucl. Acids Res.* 16:7583, 1988) and Stratagene product literature. Duplicate plaque lift filters were incubated with hybrid-subtracted cDNA probes from Example 6 in hybridization buffer containing 50% formamide, 5xSSC, 5xDenhardt's reagent and 10% dextran sulfate at 42° C. for 48 hours as described by Wahl et al., *Proc. Natl. Acad. Sci. USA* 76:3683, 1979. Filters were then washed at 68° C. in 0.2xSSC. Sixteen positive plaques were purified for further analysis.

Bluescript® plasmids containing the cDNA inserts were excised from the phage as described by the manufacturer and transformed into *E. coli*. Plasmid DNA was isolated from individual colonies, digested with EcoR I to release the cDNA inserts and electrophoresed on standard 1% agarose gels. Four duplicate gels were blotted onto nylon filters to produce identical Southern blots for analysis with various probes which were (1) radiolabeled cDNA from unsorted CTLL cells, (2) radiolabeled cDNA from CTLL 19.4 sorted cells, (3) hybrid subtracted cDNA from CTLL 19.4 sorted cells, and (4) hybrid subtracted cDNA from CTLL 19.4 sorted cells after a second round of hybridization to poly A⁺ mRNA from an IL-4 receptor negative mouse cell line (LBRM 33 1A5B6). These probes were increasingly enriched for cDNA copies of mRNA specific for the sorted cell line CTLL 19.4. Of the 16 positive plaques isolated from

the library, four clones (11A, 14, 16 and 18) showed a parallel increase in signal strength with enrichment of the probe.

Restriction mapping (shown in FIG. 1) and DNA sequencing of the isolated CTLL clones indicated the existence of at least two distinct mRNA populations. Both mRNA types have homologous open reading frames over most of the coding region yet diverge at the 3' end, thus encoding homologous proteins with different COOH-terminal sequences. DNA sequence from inside the open reading frames of both clones code for protein sequence that is identical to protein sequence derived from sequencing of the purified IL-4 receptor described in more detail in Example 5. Clone 16 and clone 18 were used as the prototypes for these two distinct message types. Clone 16 contains an open reading frame that encodes a 258-amino acid polypeptide which includes amino acids -25 to 233 of FIG. 2A. Clone 18 encodes a 230-amino acid soluble receptor protein, the N-terminal 224 amino acids of which are identical to the N-terminus of clone 16 but diverge 9 amino acids upstream of the putative transmembrane region beginning with nucleotide number 598. This insertion adds the 3' nucleotide sequence CCAAGTAATGAAAATCTG which encodes the C-terminal 6 amino acids, Pro-Ser-Asn-Glu-Asn-Leu, followed by a termination codon TGA. Both clones were expressed in a mammalian expression system, as described in Example 8.

Example 8

Expression of IL-4R in Mammalian Cells

A. Expression in COS-7 Cells. A eukaryotic expression vector pCAV/NOT, shown in FIG. 3, was derived from the mammalian high expression vector pDC201, described by Sims et al., *Science* 241:585, 1988). pDC201 is a derivative of pMLSV, previously described by Cosman et al., *Nature* 312:768, 1984. pCAV/NOT is designed to express cDNA sequences inserted at its multiple cloning site (MCS) when transfected into mammalian cells and includes the following components: SV40 (hatched box) contains SV40 sequences from coordinates 5171-270 including the origin of replication, enhancer sequences and early and late promoters. The fragment is oriented so that the direction of transcription from the early promoter is as shown by the arrow. CMV contains the promoter and enhancer regions from human cytomegalovirus (nucleotides -671 to +7 from the sequence published by Boshart et al., *Cell* 41:521-530, 1985). The tripartite leader (stippled box) contains the first exon and part of the intron between the first and second exons of the adenovirus-2 tripartite leader, the second exon and part of the third exon of the tripartite leader and a multiple cloning site (MCS) containing sites for Xho I, Kpn I, Sma I, Not I and Bgl II. pA (hatched box) contains SV40 sequences from 4127-4100 and 2770-2533 that include the polyadenylation and termination signals for early transcription. Clockwise from pA are adenovirus-2 sequences 10532-11156 containing the VAI and VAI1 genes (designated by a black bar), followed by pBR322 sequences (solid line) from 4363-2486 and 1094-375 containing the ampicillin resistance gene and origin of replication. The resulting expression vector was designated pCAV/NOT.

Inserts in clone 16 and clone 18 were both released from Bluescript® plasmid by digestion with Asp 718 and Not I. The 3.5 kb insert from clone 16 was then ligated directly into the expression vector pCAV/NOT also cut at the Asp 718 and Not I sites in the polylinker region. The insert from clone 18 was blunt-ended with T4 polymerase followed by

ligation into the vector pCAV/NOT cut with Sma I and dephosphorylated.

Plasmid DNA from both IL-4 receptor expression plasmids were used to transfect a sub-confluent layer of monkey COS-7 cells using DEAE-dextran followed by chloroquine treatment, as described by Luthman et al. (*Nucl. Acids Res.* 11:1295, 1983) and McCutchan et al. (*J. Natl. Cancer Inst.* 41:351, 1968). The cells were then grown in culture for three days to permit transient expression of the inserted sequences. After three days, cell culture supernatants and the cell monolayers were assayed (as described in Example 1) and IL-4 binding was confirmed.

B. Expression in CHO Cells. IL-4R was also expressed in the mammalian CHO cell line by first ligating an Asp718/NotI restriction fragment of clone 18 into the pCAV/NOT vector as described in Example 8. The pCAV/NOT vector containing the insert from clone 18 was then co-transfected using a standard calcium phosphate method into CHO cells with the dihydrofolate reductase (DHFR) cDNA selectable marker under the control of the SV40 early promoter. The DHFR sequence enables methotrexate selection for mammalian cells harboring the plasmid. DHFR sequence amplification events in such cells were selected using elevated methotrexate concentrations. In this way, the contiguous DNA sequences are also amplified and thus enhanced expression is achieved. Mass cell cultures of the transfectants secreted active soluble IL-4R at approximately 100 ng/ml.

C. Expression in HeLa Cells. IL-4R was expressed in the human HeLa-EBNA cell line 653-6, which constitutively expresses EBV nuclear antigen-1 driven from the CMV immediate-early enhancer/promoter. The expression vector used was pHAV-EO-NEO, described by Dower et al., *J. Immunol.* 142:4314, 1989, a derivative of pDC201, which contains the EBV origin of replication and allows high level expression in the 653-6 cell line. pHAV-EO-NEO is derived from pDC201 by replacing the adenovirus major late promoter with synthetic sequences from HIV-1 extending from -148 to +78 relative to the cap site of the viral mRNA, and including the HIV-1 tat gene under the control of the SV-40 early promoter. It also contains a Bgl II-Sma I fragment containing the neomycin resistance gene of pSV2NEO (Southern & Berg, *J. Mol. Appl. Genet.* 1:332, 1982) inserted into the Bgl II and Hpa I sites and subcloning downstream of the Sal I cloning site. The resulting vector permits selection of transfected cells for neomycin resistance.

A 760 bp IL-4R fragment of clone C-18 from the CTLL 19.4 library was released from the Bluescript® plasmid of the λZAP® cloning system (Stratagene, San Diego, Calif., USA) by digesting with EcoN I and Sst I restriction enzymes. This fragment of clone C-18 corresponds to the nucleotide sequence set forth in FIG. 2, with the addition of a 5' terminal nucleotide sequence of TGCAGGCACCTTTTGTGTCCCA, a TGA stop codon which follows nucleotide 615 of FIG. 2A, and a 3' terminal nucleotide sequence of CTGAGTGACCTGGGGGCTGCGGTGGTGAGGAGAGCT. This fragment was then blunt-ended using T4 polymerase and subcloned into the Sal I site of pHAV-EO-NEO. The resulting plasmid was then transfected into the 653-6 cell line by a modified polybrene transfection method as described by Dower et al. (*J. Immunol.* 142:4314, 1989) or by electroporation with the exception that the cells were trypsinized at 2 days post-transfection and split at a ratio of 1:8 into media containing G418 (Gibco Co.) at a concentration of 1 mg/ml. Culture media were changed twice weekly until neomycin-resistant

colonies were established. Colonies were then either picked individually using cloning rings, or pooled together, to generate mass cultures. These cell lines were maintained under drug selection at a G418 concentration of 250 ug/ml.

In an effort to select cell colonies expressing high levels of soluble IL-4 receptor, a membrane filter assay was set up as follows. High expressing cell clones were isolated by seeding 450 HeLa IL-4R transfectant cells in a 20 cm plate and allowing the cells to grow for 10 days. Cell monolayers were then washed with a Tris-buffered saline (TBS) solution and overlaid with a nitrocellulose membrane. The overlay technique is essentially that of McCracken and Brown, *Biotechniques* 2:82, 1984, except that the nitrocellulose was overlaid with small glass beads to ensure that the membrane was kept flat. Cells were incubated for an additional 24 hours to allow secretion and adsorption of soluble IL-4 receptor to the nitrocellulose membrane. Finally the membrane was removed, washed gently in TBS containing 1% bovine serum albumin (BSA) for 30 minutes at room temperature, then incubated in TBS with 3% BSA containing ^{125}I -IL-4 ($4 \times 10^{-11}\text{M}$, specific activity $\sim 1 \times 10^{16}$ cpm/mmol) for hours at 4° C. Membranes were then washed 3 times with PBS, dried and exposed on Kodak X-omat™ film overnight at -70° C.

The developed film showed spots aligned with cells growing on the plates in culture. Cell colonies aligned with the darkest spots on the film (indicating the highest level of IL-4 receptor production by cells) were harvested, and grown up in culture. When the individual clones reached confluency, supernatants were tested for the presence of soluble IL-4 receptor in a binding inhibition assay as follows. Inhibition assays were performed by first incubating various concentrations of unlabeled IL-4 or soluble IL-4 receptor with 50 ul of ^{125}I -IL-4 ($1.65 \times 10^{-10}\text{M}$) for thirty minutes at 37° C. in binding medium (RPMI with 2.5% BSA, 0.2% sodium azide, 0.2% M Hepes, pH 7.4). Subsequently 2×10^6 CTLL-2 cells were added in 50 ul of binding medium and the incubation continued for an additional thirty minutes. Free and cell-bound ^{125}I -IL-4 were then separated by the pthalate oil separation method (Dower, S. K. et al., *J. Immunol.* 132:751, 1984). Percent specific inhibition was calculated using incubation of ^{125}I -IL-4 with excess unlabeled IL-4 ($4 \times 10^{-9}\text{M}$) as a positive control and 50 ul of binding medium as a negative control. The binding data were calculated and graphed using RS/1 (BBN Software Products, Cambridge, Mass.) as previously described (Dower, S. K. et al., *J. Exp. Med.* 162:501, 1985).

Initial cultures of cells produced ~ 100 – 600 ng/ml of soluble IL-4R protein, and several cell clones isolated with the membrane-trapping technique produced as much as 2–3 ug/ml of IL-4R protein. These cell lines are currently maintained under drug selection in G418 at a concentration of 250 ug/ml. The establishment of a stable cell clone, HeLa E3C3, producing soluble IL-4R enabled us to begin scaling up production and purification of the soluble recombinant IL-4 receptor. For soluble IL-4 receptor production, the HeLa E3C3 cells were seeded in expanded surface area roller bottles (1:20 split ratio), and were grown for four days with 250 ml of modified Dulbecco's Eagles medium, 5% fetal bovine serum and 1% penicillin, streptomycin and glutamine. Roller bottles were then switched to serum free media (300 ml/roller bottle) for three days. Soluble IL-4 receptor protein was purified from HeLa E3C3 culture supernatants by affinity chromatography on IL-4 linked to Affigel-10. Recombinant murine IL-K was coupled to Affigel®-10 (BioRad) according to the manufacturer's suggestions. Briefly, 1.0 ml of washed Affigel®-10 was added

to a solution of IL-4 (3.4 mg/ml in 0.4 ml of 0.1 M HEPES pH 7.4). The solution was rocked overnight at 4° C. and an aliquot of the supernatant tested for protein by a BioRad protein assay per the manufacturer's instructions using BSA as a standard. Greater than 95% of the protein had coupled to the gel, suggesting that the column had a final load of 1.3 mg IL-4 per ml gel. Glycine ethyl ester was added to a final concentration of 0.05 M to block any unreacted sites on the gel. The gel was washed extensively with PBS-1% Tritons followed by 0.1 Glycine-HCl, pH 3.0. A 0.8x4.0 cm column was prepared with IL-4-coupled Affigel® prepared as

α -factor (See Kujan et al., U.S. Pat. No. 4,546,082). An Asp 718 restriction site was introduced at position 237 in the α -factor signal peptide to facilitate fusion to heterologous genes. This was achieved by changing the thymidine residue at nucleotide 241 to a cytosine residue by oligonucleotide-directed in vitro mutagenesis as described by Craik, *BioTechniques*, January 1985, pp.12–19. A synthetic oligonucleotide containing multiple cloning sites and having the following sequence was inserted from the Asp718 site at amino acid 79 near the 3' end of the α -factor signal peptide to a SpeI site in the 2μ sequence:

[illegible]

described (4.0 ml bed volume) and washed with PBS containing 1% Triton® X-100 for purification of murine IL-4R. Alternatively, 50 μ l aliquots of 20% suspension of IL-4-coupled Affigel® were incubated with ³⁵S-cysteine/methionine-labeled cell extracts for small-scale affinity purifications and gel electrophoresis.

Aliquots (25 ml) of HeLa E3C3 culture supernatants (containing soluble IL-4 receptor) were slowly applied to the murine IL-4 affinity column at 4° C. (flow rate of 3.0 ml/hr). The column was then washed sequentially with PBS to remove all contaminating material except the bound mIL-4R. The column was then eluted with 0.01 M acetic acid, 0.15 M sodium chloride, pH 3.0 to remove the IL-4R and washed subsequently with PBS. One ml fractions were collected for the elution and 2 ml fractions collected during the wash. Immediately following elution, samples were neutralized with 80 ul of 1 M Hepes, pH 7.4. The presence of receptor in the fractions was detected by the inhibition binding assay described above.

From 100 ml of HeLa E3C3 culture supernatant approximately 600 ug of soluble IL-4 receptor protein was purified on a 4.0 ml affinity column. Purified receptor consisted of three major bands ranging from 30–39,000 daltons on SDS-PAGE. Heterogeneity in size of this preparation is due to variability in protein glycosylation, as treatment of the protein with N-glycanase to remove N-linked carbohydrates reduces the size of the protein to ~25,000 daltons on SDS-PAGE. In addition, amino acid sequencing confirmed that the bands have the same N-terminal sequence. Purity and protein concentrations were also confirmed by amino acid analysis.

Example 9

Expression of IL-4R in Yeast Cells

For expression of mIL-4R, a yeast expression vector derived from pIXY120 was constructed as follows. pIXY120 is identical to pYαHuGM (ATCC 53157), except that R contains no cDNA insert and includes a polylinker/multiple cloning site with an Nco I site. This vector includes DNA sequences from the following sources: (1) a large Sph I (nucleotide 562) to EcoR I (nucleotide 4361) fragment excised from plasmid pBR322 (ATCC 37017), including the origin of replication and the ampicillin resistance marker for selection in *E. coli*; (2) *S. cerevisiae* DNA including the TRP-1 marker, 2μ origin of replication, ADH2 promoter; and (3) DNA encoding an 85 amino acid signal peptide derived from the gene encoding the secreted peptide

pBC120 also varies from pY α HuGM by the presence of a 514 bp DNA fragment derived from the single-stranded phage f1 containing the origin of replication and intergenic region, which has been inserted at the Nru I site in the pBR322 sequence. The presence of an f1 origin of replication permits generation of single-stranded DNA copies of the vector when transformed into appropriate strains of *E. coli* and superinfected with bacteriophage f1, which facilitates DNA sequencing of the vector and provides a basis for in vitro mutagenesis. To insert a cDNA, pLXY120 is digested with Asp 718 which cleaves near the 3' end of the α -factor leader peptide (nucleotide 237) and, for example, BamH I which cleaves in the polylinker. The large vector fragment is then purified and ligated to a DNA fragment encoding the protein to be expressed.

To create a secretion vector for expressing mIL-4R, a cDNA fragment encoding mIL-4R was excised from the Bluescript® plasmid of Example 8 by digestion with Ppuu I and Bgl II to release an 831 bp fragment from the Ppuu I site (see FIGURE) to an Bgl II site located 3' to the open reading frame containing the mIL-4R sequence minus the first two 5' codons encoding Ile and Lys. pIXY120 was digested with Asp 718 near the 3' end of the α -factor leader and BamH I. The vector fragment was ligated to the Ppuu I/Bgl II hIL-4R cDNA fragment and the following fragment created by annealing a pair of synthetic oligonucleotides to recreate the last 6 amino acids of the α -factor leader and the first two amino acids of mature mIL-4R.

α-factor processing-->|
GTA CCT CTA GAT AAA AGA ATC AAG
GA GAT CTA TTT TCT TAG TTC CAG
Val Pro Leu Asp Lys Arg Ile Lys
|<--mIL-4R

The oligonucleotide also included a change from the nucleotide sequence TGG ATA to CTA GAT which introduces a Xba I restriction site, without altering the encoded amino acid sequence.

The foregoing expression vector was then purified and employed to transform a diploid yeast strain of *S. cerevisiae* (XV2181) by standard techniques, such as those disclosed in EPA 165,654, selecting for tryptophan prototrophs. The resulting transformants were cultured for expression of a secreted mIL-4R protein. Cultures to be assayed for biological activity were grown in 20–50 ml of YPD medium (1% yeast extract, 2% peptone, 1% glucose) at 37° C. to a cell density of 1–5×10⁸ cells/ml. To separate cells from medium, cells were removed by centrifugation and the medium filtered through a 0.45 μ cellulose acetate filter prior

to assay. Supernatants produced by the transformed yeast strain, or crude extracts prepared from disrupted yeast cells transformed the plasmid, were assayed to verify expression of a biologically active protein.

Example 10

Isolation of Full-length and Truncated Forms of Murine IL-4 Receptor cDNAs From Unsorted 7B9 Cells

Polyadenylated RNA was isolated from 7B9 cells, an antigen-dependent helper T cell clone derived from C57BL/6 mice, and used to construct a cDNA library in λ ZAP (Stratagene, San Diego), as described in Example 7. The λ ZAP library was amplified once and a total of 300,000 plaques were screened as described in Example 7, with the exception that the probe was a randomly primed 32 P-labeled 700 bp EcoR I fragment isolated from CTLL 19.4 clone 16. Thirteen clones were isolated and characterized by restriction analysis.

Nucleic acid sequence analysis of clone 7B9-2 revealed that it contains a polyadenylated tail, a putative polyadenylation signal, and an open reading frame of 810 amino acids (shown in FIG. 2), the first 258 of which are identical to those encoded by CTLL 19.4 clone 16, including the 25 amino acid putative signal peptide sequence. The 7B9-2 cDNA was subcloned into the eukaryotic expression vector, pCAV/NOT, and the resulting plasmid was transfected into COS-7 cells as described in Example 8. COS-7 transfectants were analyzed as set forth in Example 12.

A second cDNA form, similar to clone 18 in the CTLL 19.4 library, was isolated from the 7B9 library and subjected to sequence analysis. This cDNA, clone 7B9-4, is 376 bp shorter than clone 7B9-2 at the 5' end, and lacks the first 47 amino acids encoded by 7B9-2, but encodes the remaining N-terminal amino acids 23-199 (in FIG. 2). At position 200, clone 7B9-4 (like clone 18 from CTLL 19.4) has a 114 bp insert which changes the amino acid sequence to Pro Ser Asn Glu Asn Leu followed by a termination codon. The 114 bp inserts, found in both clone 7B9-4 and CTLL 19.4 clone 18 are identical in nucleic acid sequence. The fact that this cDNA form, which produces a secreted form of the IL-4 receptor when expressed in COS-7 cells, was isolated from these two different cell lines indicates that it is neither a cloning artifact nor a mutant form peculiar to the sorted CTLL cells.

Example 11

Isolation of Human IL-4 Receptor cDNAs From PBL and T22 Libraries by Cross-species Hybridization

Polyadenylated RNA was isolated from pooled human peripheral blood lymphocytes (PBL) that were obtained by standard Ficoll purification and were cultured in IL-2 for six days followed by stimulation with PMA and Con-A for eight hours. An oligo dT primed cDNA library was constructed in λ gt10 using techniques described in example 7. A probe was produced by synthesizing an unlabeled RNA transcript of the 7B9-4 cDNA insert using T7 RNA polymerase, followed by 32 P-labeled cDNA synthesis with reverse transcriptase using random primers (Boehringer-Mannheim). This murine single-stranded cDNA probe was used to screen 50,000 plaques from the human cDNA library in 50% formamide/0.4 M NaCl at 42° C., followed by washing in 2 \times SSC at 55° C. Three positive plaques were purified, and the EcoR I

inserts subcloned into the Bluescript® plasmid vector. Nucleic acid sequencing of a portion of clone PBL-1, a 3.4 kb cDNA, indicated the clone was approximately 67% homologous to the corresponding sequence of the murine IL-4 receptor. However, an insert of 68 bp, containing a termination codon and bearing no homology to the mouse IL-4 receptor clones, was found 45 amino acids downstream of the predicted N-terminus of the mature protein, suggesting that clone PBL-1 encodes a non-functional truncated form of the receptor. Nine additional human PBL clones were obtained by screening the same library (under stringent conditions) with a 32 P-labeled random-primed probe made from the clone PBL-1 (the 3.4 kb EcoR I cDNA insert). Two of these clones, PBL-11 and PBL-5, span the 5' region that contains the 68 bp insert in PBL-1, but lack the 68 bp insert and do not extend fully 3', as evidenced by their size, thus precluding functional analysis by mammalian expression. In order to obtain a construct expressible in COS-7 cells, the 5' Not I-Hinc II fragment of clones PBL-11 and PBL-5 were separately ligated to the 3' Hinc II-BamH I end of clone PBL-1, and subcloned into the pCAV/NOT expression vector cut with Not I and Bgl II described in Example 8. These chimeric human IL-4R cDNAs containing PBL-11/PBL-1 and PBL-5/PBL-1 DNA sequences have been termed clones A5 and B4, respectively, as further described in Example 12. These constructs were transfected into COS-7 cells, and assayed for IL-4 binding in a plate binding assay substantially as described in Sims et al. (*Science* 241:585, 1988). Both composite constructs encoded protein which exhibited IL-4 binding activity. The nucleotide sequence and predicted amino acid sequence of the composite A5 construct correspond to the sequence information set forth in FIGS. 4A-C, with the exception that a GTC codon encodes the amino acid Val at position 50, instead of Ile. No other clones that were sequenced contained this change. The consensus codon from clones PBL-1, PBL-5 and T22-8, however, is ATC and encodes Ile⁵⁰, as set forth in FIG. 4A. The nucleotide and predicted amino acid sequence of the composite B4 construct also shows that the 25 amino acid leader sequence of PBL-11 is replaced with the sequence Met-Gln-Lys-Asp-Ala-Arg-Arg-Glu-Gly-Asn.

Constructs expressing a soluble form of the human IL-4 receptor were made by excising a 5'-terminal 0.8 kb Sma I-Dra III fragment from PBL-5 and the corresponding 0.8 kb Asp718-Dra III fragment from PBL-11, of which the Dra III overhangs were blunt-ended with T4 polymerase. The PBL-5 and PBL-11 fragments were separately subcloned into CAV/NOT cut with Sma I or Asp 718 plus Sma I, respectively; these are called soluble hIL-4R-5 and soluble hIL-4R-11, respectively.

A second library made from a CD4+/CD8- human T cell clone, T22, (Acres et al., *J. Immunol.* 138:2132, 1987) was screened (using duplicate filters) with two different probes synthesized as described above. The first probe was obtained from a 220 bp Pvu II fragment from the 5' end of clone PBL-1 and the second probe was obtained from a 300 bp Pvu II-EcoR I fragment from the 3' end of clone PBL-1. Five additional cDNA clones were identified using these two probes. Two of these clones span the 5' region containing the 68 bp insert, but neither contain the insert. The third of these clones, T22-8, was approximately 3.6 kb in size and contained an open reading frame of 825 amino acids, including a 25 amino acid leader sequence, a 207 amino acid mature external domain, a 24 amino acid transmembrane region and a 569 amino acid cytoplasmic domain. The sequence of clone T22-8 is set forth in FIGS. 4A-4C. FIGS. 5A-5B compare the predicted human IL-4R amino acid sequence

with the predicted murine IL-4R sequence and show approximately 53% sequence identity between the two proteins.

Example 12

Analysis and Purification of IL-4 Receptor in COS Transfectants

Equilibrium binding studies were conducted for COS cells transfected with murine IL-4 receptor clones 16 and 18 from the CTLL 19.4 library. In all cases analysis of the data in the Scatchard coordinate system (Scatchard, *Ann. N.Y. Acad. Sci.* 51:660-672, 1949) yielded a straight line, indicating a single class of high-affinity receptors for murine IL-4. For COS pCAV-16 cells the calculated apparent K_a was $3.6 \times 10^9 \text{ M}^{-1}$ with 5.9×10^5 specific binding sites per cell. A similar apparent K_a was calculated for COS pCAV-18 cells at $1.5 \times 10^9 \text{ M}^{-1}$ but receptor number expressed at the cell surface was 4.2×10^4 . Equilibrium binding studies performed on COS cells transfected with IL-4R DNA clones isolated from the 7B9 cell library also showed high affinity binding of the receptor to IL-4. Specifically, studies using COS cells transfected with pCAV-7B9-2 demonstrated that the full length murine IL-4 receptor bound ^{125}I -IL-4 with an apparent K_a of about $1.4 \times 10^{10} \text{ M}^{-1}$ with 4.5×10^4 specific binding sites per cell. The apparent K_a of CAV-7B9-4 IL-4R was calculated to be about $1.7 \times 10^9 \text{ M}^{-1}$. Although absolute values for K_a and binding sites per cell varied between transfections, the binding affinities were generally similar (1×10^9 – $1 \times 10^{10} \text{ M}^{-1}$) and matched well with previously published affinity constants for IL-4 binding.

Inhibition of ^{125}I -mIL-4 binding to CTLL cells by conditioned media from COS cells transfected with plasmid pCAV, pCAV-18, or pCAV-7B9-4 was used to determine if these cDNAs encoded functional soluble receptor molecules. Approximately $1.5 \mu\text{l}$ of COS pCAV-18 conditioned media in a final assay volume of $150 \mu\text{l}$ gives approximately 50% inhibition of ^{125}I -IL-4 binding to the IL-4 receptor on CTLL cells. ^{125}I -IL-4 receptor competing activity is not detected in control pCAV transfected COS supernatants. From quantitative analysis of the dilution of pCAV-18 supernatant required to inhibit ^{125}I -IL-4 binding by 50%, it is estimated that approximately 60–100 ng/ml of soluble IL-4 receptor has been secreted by COS cells when harvested three days after transfection. Similar results were obtained utilizing supernatants from COS cells transfected with pCAV-7B9-4.

Conditioned medium from COS cells transfected with pCAV-18 or pCAV-7B9-4 (see Example 8) and grown in DMEM containing 3% FBS was harvested three days after transfection. Supernatants were centrifuged at 3,000 cpm for 10 minutes, and frozen until needed. Two hundred ml of conditioned media was loaded onto a column containing 4 ml of mIL-4 Affigel prepared as described above. The column was washed extensively with PBS and IL-4 receptor eluted with 0.1 M glycine, 0.15 M NaCl pH 3.0. Immediately following elution, samples were neutralized with 80 μl of 1 M HEPES pH 7.4. Samples were tested for their ability to inhibit binding of ^{125}I -mIL-4 to CTLL cells as set forth in Example 1B. Additionally samples were tested for purity by analysis on SDS-PAGE and silver staining as previously described. Alternative methods for testing functional soluble receptor activity or IL-4 binding inhibition include solid-phase binding assays, as described in Example 1C, or other similar cell free assays which may utilize either radioiodinated or colorimetrically developed IL-4 binding, such as RIA or ELISA. The protein analyzed by SDS-PAGE

under reducing conditions has a molecular weight of approximately 37,500, and appears approximately 90% pure by silver stain analysis of gels.

Purified recombinant soluble murine IL-4 receptor protein may also be tested for its ability to inhibit IL-4 induced ^3H -thymidine incorporation in CTLL cells. Pursuant to such methods, soluble IL-4 receptor has been found to block IL-4 stimulated proliferation, but does not affect IL-2 driven mitogenic response.

Molecular weight estimates were performed on mIL-4 receptor clones transfected into COS cells. Utilizing M2 monoclonal antibody prepared against murine CTLL 19.4 cells (see Example 13), IL-4 receptor is immunoprecipitated from COS cells transfected with CAV-16, CAV-7B9-2 and CAV-7B9-4 and labeled with ^{35}S -cysteine and ^{35}S -methionine. Cell associated receptor from CAV-7B9-4 shows molecular weight heterogeneity ranging from 32–39 kDa. Secreted CAV-7B9-4 receptor has molecular weight between 36 and 41 kDa. Cell associated receptor from CAV-16 transfected COS cells is about 40–41 kDa. This is significantly smaller than molecular weight estimations from crosslinking studies described by Park et al., *J. Exp. Med.* 166:476, 1987; *J. Cell. Biol.*, Suppl. 12A, 1988. Immunoprecipitation of COS CAV-7B9-2 cell-associated receptor showed a molecular weight of 130–140 kDa, similar to the estimates of Park et al., *J. Cell. Biol. Suppl.* 12A, 1988, estimated to be the full length IL-4 receptor. Similar molecular weight estimates of cell-associated CAV-16 and CAV-7B9-2 IL-4 receptor have also been made based on cross-linking ^{125}I -IL-4 to COS cells transfected with these cDNAs. Heterogeneity of molecular weight of the individual clones can be partially attributed to glycosylation. This data, together with DNA sequence analysis, suggests that the 7B9-2 cDNA encodes the full length cell-surface IL-4 receptor, whereas both 7B9-4 and clone 18 represent soluble forms of murine IL-4 receptor.

Receptor characterization studies were also done on COS cells transfected with hIL-4R containing expression plasmids. The two chimeric human IL-4R molecules A5 and B4 (defined in Example 11) were transfected into COS cells and equilibrium binding studies undertaken. The COS monkey cell itself has receptors capable of binding hIL-4; therefore the binding calculations performed on COS cells transfected with and overexpressing hIL-4R cDNAs represent background binding from endogenous monkey IL-4R molecules subtracted from the total binding. COS cells transfected with hIL-4R A5 had 5.3×10^4 hIL-4 binding sites with a calculated K_a of $3.48 \times 10^9 \text{ M}^{-1}$. Similarly, the hIL-4R B4 expressed in COS cells bound ^{125}I -hIL-4 with an affinity of $3.94 \times 10^9 \text{ M}^{-1}$ exhibiting 3.2×10^4 receptors per cell.

Molecular weight estimates of human IL-4R expressed in COS cells were also performed. COS cells transfected with clones A5 or B4 in pCAV/NOT were labeled with ^{35}S -cysteine/methionine and lysed. Human IL-4R was affinity purified from the resulting lysates with hIL-4-coupled Affigel® (as described in Example 4). The hIL-4R A5 and B4 eluted from this affinity support migrated at about 140,000 daltons on SDS-PAGE, agreeing well with previous estimates of hIL-4R molecular weight by cross-linking (Park et al., *J. Exp. Med.* 166:476, 1987), as well as with estimates of full-length mIL-4R presented here.

Because no soluble human IL-4R cDNA has thus far been found occurring naturally, as was the case for the murine receptor (clones 18 and 7B9-4), a truncated form was constructed as described in Example 11. Following expression in COS cells, supernatants were harvested three days

after transfection with soluble hIL-4R-11 and soluble hIL-4R-5 and tested for inhibition of ^{125}I -hIL-4 binding to the human B cell line Raji. Supernatants from two of the soluble hIL-4R-11 and one of the soluble hIL-4R-5 transfected plates contained 29–149 ng/ml of IL-4R competing activity into the medium. In addition, the truncated protein could be detected in ^{35}S -methionine/cysteine-labeled COS cell transfectants by affinity purification on hIL-4-coupled Affigel® as approximately a 44 kDa protein by SDS-PAGE.

Example 13

Preparation of Monoclonal Antibodies to IL-4R

Preparations of purified recombinant IL-4 receptor, for example, human or murine IL-4 receptor, transfected COS cells expressing high levels of IL-4 receptor or CTLL 19.4 cells are employed to generate monoclonal antibodies against IL-4 receptor using conventional techniques, such as those disclosed in U.S. Pat. No. 4,411,993. Such antibodies are likely to be useful in interfering with IL-4 binding to IL-4 receptors, for example, in ameliorating toxic or other undesired effects of IL-4.

To immunize rats, IL-4 receptor bearing CTLL 19.4 cells were used as immunogen emulsified in complete Freund's adjuvant and injected in amounts ranging from 10–100 μl subcutaneously into Lewis rats. Three weeks later, the immunized animals were boosted with additional immunogen emulsified in incomplete Freund's adjuvant and boosted every three weeks thereafter. Serum samples are periodically taken by retroorbital bleeding or tail-tip excision for testing by dot-blot assay, ELISA (enzyme-linked immunosorbent assay), or inhibition of binding of ^{125}I -IL-4 to extracts of CTLL cells (as described in Example 1). Other assay procedures are also suitable. Following detection of an appropriate antibody titer, positive animals were given a final intravenous injection of antigen in saline. Three to four days later, the animals were sacrificed, splenocytes harvested, and fused to the murine myeloma cell line AG8653. Hybridoma cell lines generated by this procedure were plated in multiple microtiter plates in a HAT selective medium (hypoxanthine, aminopterin, and thymidine) to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

Hybridoma clones thus generated were screened for reactivity with IL-4 receptor. Initial screening of hybridoma supernatants utilized an antibody capture and binding of partially purified ^{125}I -mIL-4 receptor. Two of over 400 hybridomas screened were positive by this method. These two monoclonal antibodies, M1 and M2, were tested by a modified antibody capture to detect blocking antibody. Only M1 was able to inhibit ^{125}I -rmIL-4 binding to intact CTLL cells. Both antibodies are capable of immunoprecipitating native mIL-4R protein from CTLL cells or COS-7 cells transfected with IL-4R clones labelled with ^{35}S -cysteine/methionine. M1 and M2 were then injected into the peritoneal cavities of nude mice to produce ascites containing high concentrations (>1 mg/ml) of anti-IL-4R monoclonal antibody. The resulting monoclonal antibody was purified by ammonium sulfate precipitation followed by gel exclusion chromatography, and/or affinity chromatography based on binding of antibody to Protein G.

A series of experiments (Examples 14–19) was conducted to show that sIL-4R inhibits IL-4 mediated B cell growth, differentiation and function in vitro. Each of these experiments utilized soluble IL-4R produced as described in Example 8C. Example 14 shows that sIL-4R inhibits the proliferation of stimulated B cells. Examples 15, 16 and 17

show, respectively, that sIL-4R inhibits IL-4 dependent B cell differentiation as measured by induction of IgG1 and IgE secretion by LPS activated B cells, down regulation of IgG3 secretion by LPS activated B cells, and increased Ia and Fc ϵ R (CD23) expression, respectively. In the following experiments, the activity of sIL-4R is compared with sIL-1R to show that the inhibitory effects of the soluble receptors are specific in that sIL-4R has no effect on IL-1 induced B cell activity and sIL-1R has no effect on IL-4 activity, thus demonstrating two independent pathways of B cell activation directed by IL-1 and IL-4.

Example 14

Inhibition of IL-4 Binding to B Cells in vitro by Soluble IL-4R

Untreated B cells express low, but detectable levels of IL-4 receptors. Upon stimulation with the B cell mitogen LPS, these cells show enhanced cell surface IL-4 receptor expression. The following experiments were conducted to show that sIL-4R inhibits radiolabeled IL-4 binding to LPS activated B lymphocytes.

B lymphocytes were first purified from spleens of 8 to 12 week-old C57BL/6 mice (Jackson Laboratory, Bar Harbor, Me. and Simonson, Gilroy, Calif.) as described by Grabstein, et al., *J. Exp. Med.* 163:1405, 1986. Briefly, murine splenocytes were depleted of T cells by incubation in a cocktail containing T24 rat anti-Thy 1 mAb (Dennert et al., *J. Immunol.* 131:2445 (1983), GK1.5 rat anti-mouse L3T4 mAb (Dalynas et al., *Cell. Immunol.* 53:350, (1980), rabbit anti-mouse thymocyte serum (absorbed with C57BL/6 liver and bone marrow), and rabbit complement (Pel-Freez Biologicals, Rogers, Ark.). Cells were then passed over Sephadex G-10 (Pharmacia Uppsala, Sweden) to remove adherent cells. B lymphocytes were positively selected by panning on petri dishes coated with affinity purified goat anti-mouse IgM (Organon Teknica Corp., West Chester, Pa.). The resultant preparations were >98% B cells as determined by flow cytometry.

The purified B cells were then cultured in RPMI 1640 supplemented with 5% fetal calf serum (Hazelton), sodium pyruvate (1 mM), nonessential amino acids (0.1 mM), penicillin (100 U/ml), streptomycin (100 ug/ml), L-glutamine (2 mM), and 2-mercaptoethanol (50 μM), as well as *Salmonella typhimurium* LPS (10 ug/ml; Difco Laboratories, Detroit, Mich.) to produce activated B cells.

Human rIL-1 β was produced in *Escherichia coli* and purified to homogeneity as described by Kronheim et al., *Bio/Technology* 4:1708, 1986. Recombinant murine IL-4 was produced in yeast, purified to homogeneity, and radiolabeled as described by Mosley et al., *Cell* 59:355, 1989, and Park et al., *Proc. Natl. Acad. Sci. USA* 84:1669, 1987.

Inhibition assays were performed by first incubating various concentrations of unlabeled cytokines (IL-1 or IL-4), soluble receptors (sIL-1R or sIL-4R), monoclonal antibody (11B11, a rat IgG1 anti-murine IL-4 antibody produced as described by Ohara et al., *Nature* 315:333, 1985) or medium control with 50 μl ^{125}I -labeled IL-4 (1.65×10^{-10} M) for thirty minutes in 10% CO_2 at 37° C. in binding medium (RPMI/2.5% BSA/0.20/sodium azide/0.2 M Hepes, pH 7.4). To these 2×10^6 murine B cells were added in 50 μl of binding medium for 30 min at 37° C. Cells were then separated by the phthalate oil method as described by Dower et al., *J. Immunol.* 132:751, 1984. Percent specific inhibition was calculated using incubation of ^{125}I -IL-4 with excess unlabeled IL-4 (4×10^{-9} M) as positive control an 50 μl of

binding medium as negative control. Each assay was performed with 3-fold dilutions in duplicate of each competitor compound through binding medium, and incubations carried out in 96-well round bottom plates (Unbro, Hamden, Conn.).

Results of the inhibition assays indicate that IL-4 binding was inhibited by unlabeled sIL-4R, unlabeled IL-4, and 11B11, an anti-L-4 specific mAb. The blocking effect was cytokine specific, and cross-competition between IL-4 and either 11B11 or sIL-4R generated similar inhibition constants as shown in Table A below. No competition of IL-4 binding was detected by IL-1 or sIL-1R.

TABLE A

Inhibition of Radiolabeled IL-4 Binding to LPS Blasts	
Inhibitor	Inhibition Constant (M ⁻¹)
IL-4	4.9 × 10 ¹⁰
sIL-4R	4.8 × 10 ⁹
11B11	7.9 × 10 ⁹
IL-1	No Inhibition
sIL-1R	No Inhibition

Example 15

Inhibition of Lymphokine Induced B Cell Proliferation in vitro by sIL-4R

Murine B cell proliferation is stimulated by treatment of anti-immunoglobulin and either IL-1 (Howard et al., *J. Exp. Med.* 157:1529, 1983; Booth et al., *J. Immunol.* 33:1346, 1984) or IL-4 (Grabstein et al., *J. Mol. Cell. Immunol.* 2:199,1986; Howard et al, *J. Exp. Med.* 155.914,1982). The ability of sIL-1R and sIL-4R to inhibit these B cell mitogenic responses was tested in a B cell proliferation assay as follows.

B cells were purified and cultured as described in Example 14 above. In order to determine the effect of various doses of inhibitors on B cell proliferation, the purified B cells were seeded at 1 x 10⁵ cells/well in 96-well flat-bottom tissue culture plates (Costar) in the presence of affinity purified goat anti-mouse IgM (2.5 ug/ml; Zymed Laboratories, Inc., So. San Francisco, Calif.) and various concentrations of IL-4 (panel A) or IL-1 (panel B), either alone (○) or in the presence of 1000 ng/ml sIL-4R (□), 1000 ng/ml sIL-1R (■), or 555 ng/ml 11B11 (●) as inhibitors. After 2 days, cultures received 2 uCi/well of [³H]thymidine (25 Ci/mmol; Amersham, Arlington Heights., Ill.) for 16 hours, and were then harvested onto glass fiber filters. Incorporation of radioactivity was measured by liquid scintillation spectrophotometry [cite reference]. Tritiated thymidine incorporation for triplicate wells was determined for the final 16 hours of a three day culture period. Results are presented as mean cpm±SEM.

In order to determine the effect of various doses of inhibitor on the inhibition of B cell proliferation by cytokines, purified B cells were seeded at 1x10⁵ cells/well in 96-well flat-bottom tissue culture plates (Costar) in the presence of affinity purified goat anti-mouse IgM (25 ug/ml; Zymed Laboratories, Inc., So. San Francisco, Calif.) and fixed concentrations of 10 (●),1 (□), 0.1 (○), or 0 (Δ) ng/ml of IL-4 (panels A–C) or IL-1 (panels D–F). Culture wells also included three-fold dilutions of sIL-4R (panels A, D), 11B11 (panels B,E), or sIL-1R (panels C,F). After 2 days, cultures received 2 uCi/well of [³H]thymidine (25 Ci/mmol; Amersham, Arlington Heights., Ill.) for 16 hours, and were then harvested onto glass fiber filters. Incorporation of

radioactivity was measured by liquid scintillation spectrophotometry as indicated above. Tritiated thymidine incorporation for triplicate wells was determined for the final 15 hours of a three day culture period. Results are presented as mean cpm±SEM.

FIGS. 6 and 7 show that sIL-4R and sIL-1R inhibitory activity was dose dependent and specific for the respective ligands. The inhibitory effects of the sIL-4R and 11B11 were virtually equivalent on a molar basis, with half-maximal inhibition of IL-4-induced proliferation requiring a 100–200 fold molar excess of either inhibitor. Half-maximal inhibition of IL-1 activity was achieved with a 300–400 fold molar excess of sIL-1R.

Example 16

Inhibition of IL-4 Induced Immunoglobulin Secretion in vitro by sIL-4R

IL-4 augments LPS-induced secretion of IgG1 and IgE and inhibits IgG3 production, possibly by a mechanism involving class switching from one isotype of an antibody to another isotype. The ability of sIL-4R to inhibit IL-4 induced class switching in LPS-stimulated B cells was tested in the following assay that measures immunoglobulin secretion from LPS treated B cells.

B cells were purified and cultured as described in Example 14 above. In order to determine the effect of various doses of IL-4 on IgG1, IgE and IgG3 secretion, the purified B cells (1x10⁵ cells/well) were grown in 96-well flat bottom plates in the presence of *Salmonella typhimurium* LPS (Difco Laboratories, Detroit, Mich.) and three-fold dilutions of IL-4 with sIL-4R (□), sIL-1R (■) or 11B11 (●), each at 555 ng/ml or medium control (Δ) (see FIG. 8). Six days after initiation of culture, cells were pelleted by centrifugation at 750xg and culture supernatant fluids were harvested.

Immunoglobulin (IgG1, IgG3, and IgE) levels were determined by an isotype specific sandwich ELISA technique as follows. 96-well flat-bottom Linbro plates (Flow Laboratories, Inc., McLean, Va.) were coated overnight with the appropriate (see below) first step isotype specific antibody (100 ul) and washed. This and all subsequent washing steps were done with phosphate buffered saline containing 0.05% Tween 20, 6 rinses per cycle. Nonspecific sites were blocked by incubation for one hour with 150 ul of 5% nonfat dry milk. Test material (100 ul), either culture supernatant or isotype standard curve solutions (all sample and antibody dilutions in PBS/3% BSA), was added to each well, incubated for 1 hour, then washed. 100 ul of the appropriate (see below) horseradish peroxidase-conjugated second step antibody was added and plates were incubated for 1 hour and washed. The presence of peroxidase-conjugated antibody was determined by using the TMB Microwell peroxidase substrate system (Kirkegaard & Perry Laboratories, Inc., Gaithersburg, Md.). Plates were read on a Dynatech ELISA reader. Immunoglobulin concentrations in test samples were determined by comparing triplicate test values with isotype control standard curves, using the DeltaSoft 1.8 ELISA analysis program for the Macintosh (Biometallics, Inc., Princeton, N.J.).

For the IgG1 and IgG3 assays, unconjugated and horseradish peroxidase-conjugated affinity purified goat anti-mouse isotype specific reagents (Southern Biotechnology Associates, Inc., Birmingham, Ala.) were used as plate coating and second step reagents, respectively. Standard curves for IgG1 and IgG3 were run with isotype matched

murine myeloma proteins (Southern). For the IgE assay, the EM95 IgG2a anti-mouse IgE mAb (Baniyash et al., *Eur. J. Immunol.* 14:797, 1984) (provided by Dr. Fred Finkelman, Uniformed Services, Bethesda, Md.) was used as plate coating step reagent and biotinylated rat anti-mouse IgE (Bioproducts for Science, Inc., Indianapolis, Ind.) was used as second step reagent, and horse radish peroxidase-conjugated streptavidin (Zymed) was used in the third step. Standard curves were established with a murine anti-dinitrophenol specific IgE myeloma antibody (ATCC No. TIB 141). All three ELISA assays were determined to be specific based upon cross-reactivity experiments using all individual murine antibody isotypes as controls.

The effect of various doses of inhibitor on the inhibition of IgG1, IgG3 and IgE secretion is shown in FIG. 9. In this experiment, purified B cells (1×10^5 cells/well) were grown in 96well flat bottom plates in the presence of *Salmonella typhimurium* LPS (Difco Laboratories, Detroit, Mich.) and IL-4 (30 ng/ml for IgE; 3 ng/ml for IgG1 and IgG3) in the presence of three-fold dilutions of sIL-4R (\square), sIL-1R (\blacksquare) or 11B11 (\bullet). Six days after initiation of culture, cells were pelleted by centrifugation at $750 \times g$ and culture supernatant fluids were harvested. The supernatants were analyzed for IgG1, IgG3 and IgE secretion using the isotype specific sandwich ELISA technique described above.

FIG. 8 (panels A and B) shows that IgG1 and IgE secretion from LPS treated B cells was induced by IL-4 and that these activities were inhibited by both the sIL-4R as well as 11B11. In contrast, panel C shows that IgG3 secretion was induced by LPS directly in the absence of exogenous cytokines. When IL-4 was present at concentrations of 10 ng/ml or less, LPS induced IgG3 secretion was ablated. sIL-4R blocked this inhibitory effect of IL-4, shifting the IL-4 dose response curve and effectively permitting induction of IgG3 secretion in the presence of otherwise inhibitory doses of IL-4.

FIG. 9 shows that the inhibition of IL-4 induced class switching by sIL-4R was dose dependent: with increasing concentrations of inhibitors, progressively lower levels of IgG1 and IgE and progressively higher levels of IgG3 were secreted. sIL-1R had no such effect, even at the higher concentrations (>1 ug/ml).

Example 17

Inhibition of IL-4 Induced Cell Surface Antigen Expression on B Cells in vitro by sIL-4R

IL-4 Induces increased expression of various cell surface antigens on resting murine B cells. In order to determine the effect of sIL-4R on the inhibition of cell surface antigen expression, fluorescent-labeled antibodies to two specific cell surface antigens, MHC class II (Ia) antigens and Fc ϵ R (CD23), were used to measure the level of antigen expressed on B cells as follows.

Inhibition of MHC class II (Ia) Antigens. Purified B cells (5×10^5 cells/ml) were cultured for 16 h in 24-well plates (Costar) with or without IL-4 (0.1 ng/ml) in the presence of sIL-4R, 11B11 or sIL-1R each at 500 ng/ml or medium control with or without inhibitors. Cells were washed and preincubated for 20 min on ice with the rat IgG2b anti-murine Fc γ R mAb 2.4G2 (Unkeless, et al., *J. Exp. Med.* 150:580,1979) to block IgG Fc receptors. Fluorescinated mAbs (25-9-17, a murine IgG2a anti-murine I-Ab antibody, described by Ozato et al., *J. Immunol.* 126:317, 1981; or control murine IgG1) were added directly and cells were incubated for 30 min at 4° C. and washed. The antibody

diluent and wash solution was PBS/1% fetal calf serum/0.01% NaN_3 . Stained cells were analyzed on a FACScan flow cytometer (Beckton-Dickinson, San Jose, Calif.) using a logarithmic fluorescence intensity scale.

FIG. 10 shows that in medium control (panel A) Ia expression in B cells is significantly greater in the presence of IL-4 (dashed line) than without IL-4 (solid line). Addition of sIL-4R (panel B) or 11B11 (panel C) at the onset of culture returned Ia expression to constitutive levels, whereas addition of sIL-1R (panel D) had no effect. The results shown in FIG. 10 are representative of 3 separate experiments.

Inhibition of Fc ϵ R (CD23). IL-4 also induces expression of CD23 on murine and human B cells. B cells were cultured without or without IL-4 in the presence of sIL-4R, 11B11, sIL-1R or medium control as described above. Cells were stained with FITC-labeled anti-CD23 antibody (B3B4, a rat IgG2a anti-murine Fc ϵ R [CD23], described by Rao et al., *J. Immunol.* 138:1845, 1987) as described above.

FIG. 11 shows that in medium control (panel A) CD23 expression in B cells is significantly greater in the presence of IL-4 (dashed line) than without IL-4 (solid line). Addition of sIL-4R (panel B) or 11B11 (panel C) returned cell surface CD23 expression to constitutive levels, whereas addition of sIL-1R (panel D) had no effect. As with Ia expression, the sIL-4R and 11B11 inhibitors did not diminish the constitutive expression of CD23, indicating that the inhibition was limited to the IL-4 dependent increase. The results shown in FIG. 10 are also representative of 3 separate experiments.

Examples 18 and 19 are experiments which show the effect of sIL-4R on the inhibition of IgE responses in vivo. Example 18 shows that sIL-4R inhibits an IgE response to a specific antigen. Example 19 shows that administration of sIL-4R in doses ranging from 1–25 ug twice daily on days –1, 0 and +1 do not inhibit an IgE response to a cocktail of anti-IgD antibodies.

Example 18

Inhibition of IL-4 Dependent Antigen-specific IgE Response of B Cells by sIL-4R in vivo

Animals immunized with the hapten-carrier conjugate TNP-KLH (trinitrophenol-keyhole limpet hemocyanin) in alum generate a strong anti-TNP IgE antibody response. In order to determine the effect of sIL-4R administration on the IgE response, the following experiment was conducted.

Balb/c mice (3 mice/group, 7 groups) were immunized i.p. with 1 ug of TNP-KLH in alum on day 0. On day 21, the mice were boosted with the same amount of TNP-KLH, then bled 5 days later. Serum was then assayed by immunoglobulin isotype-specific ELISA for levels of both polyclonal and antigen-specific (TNP-specific) immunoglobulin. This secondary antibody response is characterized by, although not restricted to, the generation of a strong IgE response, most of which is anti-TNP specific.

On days –1, 0, and +1 of the secondary immunization, mice were given twice-daily injections of sIL-4R in total daily doses of 25, 5, and 1 ug/mouse. Thus, each mouse received a total of 75, 15, or 3 ug of sIL-4R over the three day treatment period. Serum was prepared from each animal, and analyzed for polyclonal and anti-TNP IgE concentrations.

The results of these experiments, shown in FIG. 12, indicate that untreated mice (saline control), primary-immunized mice (1-no boost), and primed and boosted mice (1–2) displayed polyclonal IgE levels of approximately 1

ug/ml, 5 ug/ml, and 25 ug/ml, respectively. Treatment of boosted mice with 11B11 anti-IL-4 antibody lowered IgE levels to approximately 2 ug/ml. Treatment of boosted mice with sIL-4R lowered IgE levels significantly, with the highest concentration of sIL-4R (25 ug/hit) resulting in greater than 80% reduction in IgE. Lower doses of sIL-4R inhibited the polyclonal IgE response, although less dramatically. Thus, the sIL-4R acts as an inhibitor of an antigen-induced polyclonal IgE response.

FIG. 13 shows that priming of mice with TNP-KLH (1-no boost) resulted in a detectable anti-TNP response of the IgE isotype. Boosting with TNP-KLH caused a significant increase in the anti-TNP IgE titre. Treatment of primed and boosted mice with 11B11 at the time of secondary immunization diminished the antigen-specific IgE levels to less than levels seen in primed-only mice. The highest concentration of sIL-4R also dramatically decreased anti-TNP specific levels. Lower concentrations of sIL-4R had no discernible effect upon the secondary IgE response to TNP-KLH.

Example 19

In vivo Inhibition of IL-4-Dependent Polyclonal IgE Response of B Cells by sIL-4R

Animals immunized with a cocktail of monoclonal IgD antibodies specific for murine IgD (allotype specific) generate a strong polyclonal IgE antibody response. One likely mechanism of anti-IgD action involves crosslinking of surface immunoglobulin on B cells, internalization and processing of the anti-IgD, and presentation to helper T cells. The Ig-allotype specific T cells are thus triggered to provide signals (presumably cytokines) which induce immunoglobulin class switching and secretion by B cells.

In order to determine the effect of sIL-4R administration on the IgD-induced polyclonal IgE response, the following experiment was conducted. BALB/c mice (3 mice/group) immunized i.v. with 800 ug of anti-IgD were treated twice daily with three different doses of sIL-4R (12.5, 2.5, or 0.5 ug/injection) on days -1, 0, and +1. Mice were bled on day 9 and serum IgE levels were determined. FIG. 14 shows that anti-IgD treatment (MSA control) caused large increases in levels of secreted IgE when compared with unimmunized controls (saline control). This effect was blocked by anti-IL-4 (11B11) administration, but not by any of the doses of sIL-4R. Anti-IgD treatment also caused large increases in IgG1, IgG2a, and IgG3. Whereas sIL-4R administration had no effect upon these isotypes, 11B11 administration resulted in increased IgG2a and IgG3 secretion.

The failure of sIL-4R to inhibit the IgE stimulatory effect of anti-IgD may be due to the fact that the inhibitor must be present for longer than one day after anti-IgD treatment, that the 11B11 antibody has a longer serum half-life than the sIL-4R, or that higher doses of sIL-4R are required.

Example 20

Use of Soluble IL-4 Receptor to Inhibit Contact Hypersensitivity Responses to DNFB

The effect of soluble IL-4 receptor (sIL-4R) and soluble IL-1 receptor (sIL-1R) on contact hypersensitivity (CHS) responses were evaluated in a murine system using 2,4-dinitrofluorobenzene (DNFB) as the contact sensitizer. Groups of female BALB/c mice (5 mice per group) were treated with either sIL-4R or sIL-1R on either days -1 through +1, days 4 and 5 or days -1 through 5 with 500 ng,

b.i.d. via intraperitoneal injection. Control mice were treated with equivalent doses of the carrier solution containing mouse serum albumin (MSA). Epicutaneous sensitization with DNFB was performed by application of 25 ul of a solution of 0.5% DNFB in 4:1 mixture of acetone:olive oil to the shaved backs of mice on day 0. Negative control mice were not sensitized to DNFB. CHS responses were elicited by challenging all groups of mice on day 5 by application of 10 ul of the DNFB solution to the right rear footpads of the mice in each of the treatment groups. The extent of CHS induction was determined by measuring the difference in thickness (in units of 10^{-2} mm) between the challenged right and unchallenged left rear footpads (as measured with a dial micrometer) 24 hours later. FIG. 15 shows the results of these experiments. The data are presented as mean footpad swelling \pm SEM.

As shown in FIG. 15, mice treated with sIL-1R developed CHS responses that were not significantly different from control mice treated with MSA regardless of the treatment regimen used. Mice treated with sIL-4R on days -1 through 1 developed CHS responses that were not significantly different from the MSA control group. Although there appeared to be a slight increase in the CHS responses to DNFB induced in mice treated with sIL-4R on days 4 and 5, inspection of the data indicated that this was primarily due to the enhanced response of only one out of five of the mice in the group. No significant effect of treatment (either enhancement or inhibition) is observed if the response of this single mouse is considered to be an outlier and is not considered in the evaluation of the data (FIG. 15). However, mice treated with sIL-4R during the entire period (days -1 through 5) were significantly ($p<0.001$) inhibited relative to MSA-treated control mice, and not significantly different from mice which had not been sensitized, but only challenged, with DNFB. This data thus indicates that sIL-4R is effective in inhibiting contact hypersensitivity responses to DNFB.

Example 21

Use of Soluble IL-4 Receptor to Inhibit Delayed-Type Hypersensitivity Responses to SRBC

The effect of soluble IL-4 receptor (sIL-4R) and soluble IL-1 receptor (sIL-1R) on delayed-type hypersensitivity (DTH) responses to sheep red blood cells (SRBC) were evaluated in a murine system as described by Kitamura, *J. Immunol. Meth.* 39:277, 1980. Three groups of female BALB/c mice (4 mice per group) were sensitized on day 0 by i.v. injection of 2×10^5 SRBC. A fourth group of negative control mice were not sensitized to SRBC. DTH responses were elicited in the sensitized mice by challenging the mice on day 4 with 1×10^8 SRBC via intracutaneous injection in the right rear footpads of the mice and 100 ul normal saline in the contralateral footpad as a control. The mice were treated with either 0.1 ug or 1.0 ug of sIL-4R in 100 ul MSA via intraperitoneal injection, on the day of challenge and the day of immunization. Control mice were treated with equivalent doses of the carrier solution (100 ul) containing mouse serum albumin (MSA). The fourth group of negative control mice not sensitized to SRBC were treated with 100 ul of MSA. The extent of the DTH response induced was determined by excising the footpads at the tarsus and measuring the difference in weight between SRBC challenged and saline challenged footpads. FIG. 16 shows the results of these experiments. The data are presented as mean footpad swelling \pm SEM.

As shown in FIG. 16, the DTH response was almost totally blocked by intraperitoneal injection with 1 ug sIL-4R

on the day of challenge and the day of immunization. Treatment with 0.1 ug sIL-4R was less effective. Treatment with 0.2 ug or 2 ug of sIL-1R inhibited the response (not shown). These data thus indicate that sIL-4R is effective in inhibiting delayed-type hypersensitivity responses.

Example 22

Use of Soluble IL-4R to Suppress Immune Response to Alloantigen in vivo

Experiments were conducted to show that systemic administration of sIL-4R suppresses a localized, T cell-dependent, immune response to alloantigen presented by allogeneic cells. The response to allogeneic cells in vivo was quantified using the popliteal lymph node enlargement assay described by Twist et al., *Transplantation* 15: 182, 1973, which is used as a measure of allograft transplant immunity (see Grebe et al., *Adv. Immunol.* 22:119, 1976). In this assay mice are injected in the footpad with irradiated, allogeneic spleen cells. The mice are then injected in the contralateral footpad with irradiated, syngeneic spleen cells. An alloreactive response (marked by proliferation of lymphocytes and inflammation) occurs in the footpad receiving the allogeneic cells, which can be measured by determining the increase in size and weight of the popliteal lymph node draining the site of antigen deposition relative to controls or by an increase in cellularity.

Specific pathogen free 8–12 week old BALB/c (H-2^d) and C57BL/6 (H-2^b) mice (Jackson Laboratory, Bar Harbor, Me.) were used in this experiment. 9 BALB/c mice were divided into 3 groups, each having 3 mice. Each group of mice received a different mode of treatment as indicated below in Tables B. On day 0 the left footpads of all mice were injected intracutaneously with 10⁷ irradiated (2500R), allogeneic spleen cells from C57BL/6 mice in 50 ul of RPMI-1640 (Gibco) as antigen and the right contralateral footpads of the same mice were injected with 10⁷ irradiated (2500R), syngeneic spleen cells from BALB/c mice. All doses of soluble murine IL-4 receptor (sIL-4R) were diluted in phosphate buffered saline (PBS). On days -1, 0 and +1 three mice were injected (intravenously on days -1 and 0, and subcutaneously on day +1) with 100 ng of purified smuIL-4R, three mice were injected intravenously with 1 ug of smuIL-4R, three mice were injected with 2 ug of smuIL-4R and three mice were injected with MSA (control).

Seven days after antigen administration, the mice were sacrificed and the popliteal lymph nodes (PLN) were removed from the right and left popliteal fossa by surgical dissection. Lymph nodes were weighed and the results expressed as the difference (Δ) in weight (mg) of the lymph node draining the site of allogeneic cell injection and the weight of the node draining the syngeneic cell injection site (Table B; FIG. 17). The mean difference in weight of the lymph nodes from the sites of allogeneic and syngeneic spleen cells was approximately 2.5 mg for the mice treated with MSA, 1 mg for the mice treated with 100 ng of sIL-4R, and 0.5 mg for mice treated with 1 ug sIL-4R. No detectable difference in weight of lymph nodes was ascertainable for the mice treated with 2 ug sIL-4R. Lymph nodes draining the syngeneic cell injection site weighed approximately 1 mg, regardless of whether they were obtained from mice treated with MSA or smuIL-4R, and did not differ significantly in weight from nodes obtained from mice given no cell injection. Values for statistical significance were calculated using the two-tailed Student's t-test. Thus, IL-4R significantly (p<0.1 in all groups, using a two-tailed T test) suppressed the in vivo lymphoproliferative response in a dose dependent fashion relative to control mice.

TABLE B

Effect of smuIL-4R Administration on Proliferation of Lymph Node Cells			
Treatment	Weight (mg) of Lymph Node		
Group	Allogeneic	Syngeneic	Δ
MSA	3.9 ± 0.06	1.27 ± 0.07	2.63 ± 0.1
100 ng smuIL-1R	2.3 ± 0.03	1.3 ± 0.03	1.0 ± 0.06
1 ug smuIL-1R	2.1 ± 0.9	1.9 ± 0.3	0.23 ± 0.6
2 ug smuIL-1R	1.6 ± 0.3	1.5 ± 0.1	0.0 ± 0.4

Table B (illustrated graphically in FIG. 17) shows that systemic administration of sIL-4R for 3 days beginning on day -1 relative to alloantigenic challenge resulted in a dramatic decrease in the size of lymph nodes, indicating that the lymphoproliferative response is inhibited. The effect was dose dependent and, in some cases, the response was virtually eliminated.

Example 23

Use of Soluble IL-4 Receptor to Suppress Allograft Rejection

Soluble murine IL-4 receptor also suppresses rejection of organ grafts in vivo. In order to demonstrate this, neonatal C57BL/6 (H-2^b) hearts were transplanted into the ear pinnae of adult BALB/c (H-2^d) recipients utilizing the method of Fulmer et al., *Am. J. Anat.* 113:273, 1963, modified as described by Trager et al., *Transplantation* 47:587,1989, and Van Buren et al., *Transplant. Proc.* 15:2967, 1983. Survival of the transplanted hearts was assessed by visually inspecting the grafts for pulsatile activity. Pulsatile activity was determined by examining the ear-heart grafts of anesthetized recipients under a dissecting microscope with soft reflected light beginning on day 5 or 6 post transplant. The time of graft rejection was defined as the day after transplantation on which contractile activity ceases.

Recipient mice were divided into two groups, a primary treatment group and a secondary treatment group. The primary treatment group were not exposed to antigen from C57BL/6 mice previous to being treated, while the secondary treatment group had been exposed to antigen. All mice were transplanted on day 0 and injected with either smuIL-4R (1000 ng/day) plus MSA (mouse serum albumin, 100 ng) or with MSA alone on days 0 through 2, i.p. The results of this experiment are reported below in Table C. The probability (p value) that the survival time for the group treated with smuIL-4R differs by chance alone from the group treated with MSA is less than 0.04 when analyzed by the Student's t-test for the primary treatment group. The corresponding p values for secondary treatment group are not significant.

TABLE C

Effects of smuIL-1R Treatment on Nonvascularized Heterotopic Cardiac Allograft Survival		
Treatment Group	Survival Time (days)	Median Survival Time ±S.D.
Primary Treatment		
MSA(100 ng)	9, 10, 12, 14	11.3 ± 1.1
smuIL-1R (100 ng)	10, 10, 10, 12	10.5 ± 0.5
smuIL-4R (1000 ng)	12, 14, 14, 16, 17, 19	15.3 ± 1.0

TABLE C-continued

Effects of smuIL-1R Treatment on Nonvascularized Heterotopic Cardiac Allograft Survival		
Treatment Group	Survival Time (days)	Median Survival Time \pm S.D.
Secondary Treatment		
MSA(100 ng)	8, 8, 8	8 \pm 0.0
smuIL-1R (1000 ng)	8, 10, 12	10 \pm 1.2

Table C shows that heart allografts survived 10–14 days in individual control mice treated with MSA. When primary allograft recipients were given 3 daily injections of 1000 ng smuIL-4R, graft survival was prolonged. The median graft survival time in smuIL-4R treated mice (14–19 days) was approximately four days longer than the median graft survival time of identical grafts in control mice. A subtle increase in graft survival following secondary transplantation suggests that acute rejection episodes are influenced by smuIL-4R administration as well. This data is evidence of the therapeutic potential of soluble human IL-4 receptor in humans for the suppression of heart allograft rejection.

We claim:

1. A method of inhibiting binding of interleukin-4 (IL-4) to an IL-4 Receptor (IL-4R), said method comprising contacting an IL-4R with an antibody to IL-4R, wherein said antibody inhibits binding of IL-4 to the IL-4R,

wherein the antibody is reactive with an IL-4R selected from the group consisting of:

- a) a human IL-4R comprising amino acids 1 to 800 of FIGS. 4A–4C;
- b) a soluble human IL-4R comprising amino acids 1 to 207 of FIG. 4A;
- c) a murine IL-4R comprising amino acids 1 to 785 of FIGS. 2A–2C; and
- d) a soluble murine IL-4R comprising amino acids 1 to 208 of FIG. 2A.

2. A method of claim 1, wherein the antibody is a monoclonal antibody.

3. A method of claim 1, wherein the antibody is reactive with a soluble human IL-4R, wherein the amino acid sequence of said soluble human IL-4R consists of amino acids 1 to 207 of FIG. 4A.

4. A method of claim 3, wherein the antibody is a monoclonal antibody.

5. A method of claim 1, wherein the antibody inhibits binding of IL-4 to IL-4R on a cell.

6. A method of claim 2, wherein the monoclonal antibody inhibits binding of IL-4 to IL-4R on a cell.

7. A method of claim 3, wherein the antibody inhibits binding of IL-4 to IL-4R on a cell.

8. A method of claim 4, wherein the monoclonal antibody inhibits binding of IL-4 to IL-4R on a cell.

* * * * *