




US 20050147604A1

(19) **United States**(12) **Patent Application Publication** (10) **Pub. No.: US 2005/0147604 A1****Barden et al.**(43) **Pub. Date:****Jul. 7, 2005**(54) **MEANS AND METHODS FOR DIAGNOSING  
AND TREATING AFFECTIVE DISORDERS**(75) Inventors: **Nicholas Barden**, Sainte-Foy (CA);  
**Inge Sillaber**, Munich (DE); **Marcelo  
Paez-Pereda**, Munich (DE)Correspondence Address:  
**FOLEY AND LARDNER**  
**SUITE 500**  
**3000 K STREET NW**  
**WASHINGTON, DC 20007 (US)**(73) Assignee: **NeuroNova AG**(21) Appl. No.: **10/825,593**(22) Filed: **Apr. 16, 2004****Related U.S. Application Data**(60) Provisional application No. 60/474,232, filed on May  
30, 2003. Provisional application No. 60/501,011,  
filed on Sep. 9, 2003.(30) **Foreign Application Priority Data**Apr. 17, 2003 (EP) ..... 03 00 8753.0  
Sep. 9, 2003 (EP) ..... 03 01 9626.5**Publication Classification**(51) **Int. Cl.<sup>7</sup>** ..... **C12Q 1/68**; C07H 21/04;  
A61K 39/395; C07K 14/705;  
C07K 16/28  
(52) **U.S. Cl.** ..... **424/143.1**; 435/6; 435/69.1;  
435/320.1; 435/325; 530/350;  
536/23.5; 530/388.22(57) **ABSTRACT**

The present invention relates to nucleic acid molecules, preferably genomic sequences, encoding an ATP-gated ion channel P2X7R which contain a mutation in the 5'UTR or 3'UTR regions, a mutation in exon 3, 5, 6, 8 or 13 or in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 or a deletion in exon 13, which allow to diagnose affective disorders. The invention further relates to polypeptides encoded by said nucleic acid molecules vectors and host cells comprising said nucleic acid molecules as well as to methods for producing polypeptides encoded by said nucleic acid molecules. The present invention also provides antibodies specifically directed to polypeptides encoded by said nucleic acid molecules and aptamers specifically binding said nucleic acid molecules.

Additionally, primers for selectively amplifying said nucleic acid molecules are provided in the present invention as well as kits, compositions, particularly pharmaceutical and diagnostic compositions comprising said nucleic acid molecules, vectors, polypeptides, aptamers, antibodies and/or primers. Moreover, the present invention relates to methods for diagnosing affective disorders associated with a non-functional P2X7R protein, an altered ATP-gating of the P2X7R protein, an over- or underexpression of the P2X7R protein or associated with the presence of any one of the aforementioned nucleic acid molecules or polypeptides encoded thereby. Additionally, the present invention relates to uses and methods for treating affective disorders employing a functional or non-functional ATP-gated ion-channel P2X7R. The present invention also relates to uses of modulators of P2X7R activity for treating affective diseases.

Furthermore, the present invention also relates to methods for identifying and characterizing compounds which are capable of specifically interacting with or altering the characteristics of the polypeptides of the present invention as well as to methods for the production of pharmaceutical compositions.



**Figure 1a**




Figure 1b




Figure 1c






Figure 1d




Figure 1e




Figure 1f




Figure 1g




Figure 1h




Figure 1i




Figure 1j




Figure 1k






Figure 11




Figure 2





Figure 3



**Figure 4**




Figure 5




Figure 6




Figure 7

paroxetine-treated mouse  
(stressful handling)



vehicle-treated mouse  
(stressful handling)



stress-free mouse



Figure 8






Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15

P2X7v01 MPACCSCDVFQYETNKNVTRIQSMNYGTIKWFFHVIIFSVCFALVSDKLYQRKEPVISS  
 P2X7v04 MPPVD-----AFPCLPFS---FALVSDKLYQRKEPVISS  
 P2X7v02 MPACCSCDVFQYETNKNVTRIQSMNYGTIKWFFHVIIFSVCFALVSDKLYQRKEPVISS  
 P2X7v03 MPACCSCDVFQYETNKNVTRIQSMNYGTIKWFFHVIIFSVCFALVSDKLYQRKEPVISS  
 1.....10.....20.....30.....40.....50  
  
 P2X7v01 VHTKVKGIAEVKEEIVENGVKKLVHVSVDADYTFPLQGNSSFFVMTNFLKTEGQEQR LCP  
 P2X7v04 VHTKVKGIAEVKEEIVENGVKKLVHVSVDADYTFPLQGNSSFFVMTNFLKTEGQEQR LCP  
 P2X7v02 VHTKVKGIAEVKEEIVENGVKKLVHVSVDADYTFPLQGNSSFFVMTNFLKTEGQEQR LCP  
 P2X7v03 VHTKVKGIAEVKEEIVENGVKKLVHVSVDADYTFPLQGNSSFFVMTNFLKTEGQEQR LCP  
 61.....70.....80.....90.....100.....110

Figure 16a



P2X7v01 EYPTRRTLCSSTRGCKKGWMDPQSKGIQTGRVCVHEGNQKTCEVSAWCPIEAVEEAPRPA  
 P2X7v04 EYPTRRTLCSSTRGCKKGWMDPQSKGIQTGRVCVHEGNQKTCEVSAWCPIEAVEEAPRPA  
 P2X7v02 EYPTRRTLCSSTRGCKKGWMDPQSKGLLS-----  
 P2X7v03 EYPTRRTLCSSTRGCKKGWMDPQSKGIQTGRVCVHEGNQKTCEVSAWCPIEAVEEAPRPA  
 121.....130.....140.....150.....160.....170  
  
 P2X7v01 LLNSAENFTVLKNNIDFPGHNYTTRNLPGLNITCTFHKTQNPQCPIFRLGDIFRETGD  
 P2X7v04 LLNSAENFTVLKNNIDFPGHNYTTRNLPGLNITCTFHKTQNPQCPIFRLGDIFRETGD  
 P2X7v02 -----  
 P2X7v03 LLNSAENFTVLKNNIDFPGHNYTTRNLPGLNITCTFHKTQNPQCPIFRLGDIFRETGD  
 181.....190.....200.....210.....220.....230

Figure 16b

P2X7v01 NFSDVAIQGGIMGIEIYWDCNLD RWFHHCHPKYSFRRLLDDKTTNVSLYPGYNFRYAKYYK  
 P2X7v04 NFSDVAIQGGIMGIEIYWDCNLD RWFHHCHPKYSFRRLLDDKTTNVSLYPGYNFRYAKYYK  
 P2X7v02 -----  
 P2X7v03 NFSDVAIQGGIMGIEIYWDCNLD RWFHHCHPKYSFRRLLDDKTTNVSLYPGYNFRYAKYYK  
 241 .....250.....260.....270.....280.....290  
  
 P2X7v01 ENNVEKRTLKVFGRFDILVFGTGGKFDIIQLVYIGSTLSYFGLAAVFIDFLIDTYSS  
 P2X7v04 ENNVEKRTLKVFGRFDILVFGTGGKFDIIQLVYIGSTLSYFGLAAVFIDFLIDTYSS  
 P2X7v02 -----  
 P2X7v03 ENNVEKRTLKVFGRFDILVFGTGGKFDIIQLVYIGSTLSYFGLVRDSL FHALGKWFG  
 301 .....310.....320.....330.....340.....350


Figure 16c

P2X7v01	NCCRS	HI	P	W	C	K	C	C	Q	P	C	V	N	E	Y	Y	R	K	K	C	E	S	I	V	E	P	K	P	T	L	K	Y	V	S	F	V	D	E	S	H	I	R	M	V	N	Q	Q	L	L	G	R	S					
P2X7v04	NCCRS	HI	P	W	C	K	C	C	Q	P	C	V	N	E	Y	Y	R	K	K	C	E	S	I	V	E	P	K	P	T	L	K	Y	V	S	F	V	D	E	S	H	I	R	M	V	N	Q	Q	L	L	G	R	S					
P2X7v02	-----																																																								
P2X7v03	EGSD	-----																																																							
	361	.....	370	.....	380	.....	390	.....	400	.....	410																																														
P2X7v01	LQD	V	K	G	Q	E	V	P	R	P	A	M	D	F	T	D	L	S	R	L	P	L	A	L	H	D	T	P	P	I	P	G	Q	P	E	E	I	Q	L	L	R	K	E	A	T	P	R	S	R	D	S	P	W	C	Q	C	G
P2X7v04	LQD	V	K	G	Q	E	V	P	R	P	A	M	D	F	T	D	L	S	R	L	P	L	A	L	H	D	T	P	P	I	P	G	Q	P	E	E	I	Q	L	L	R	K	E	A	T	P	R	S	R	D	S	P	W	C	Q	C	G
P2X7v02	-----																																																								
P2X7v03	-----																																																								
	421	.....	430	.....	440	.....	450	.....	460	.....	470																																														

Figure 16d

P2X7v01	SCLPSQLPESHRCLEELCCRKPGACITTS	ELFRKLVLSRHLVQLFLLYQEPLLDVDS
P2X7v04	SCLPSQLPESHRCLEELCCRKPGACITTS	ELFRKLVLSRHLVQLFLLYQEPLLDVDS
P2X7v02	-----	
P2X7v03	-----	
	481.....490.....500.....510.....520.....530	
P2X7v01	TNSRLRHCA YRCYATWRFSGQDMADFA	ILPSCCRWRIRKEFPKSEGQYSGFKSPY
P2X7v04	TNSRLRHCA YRCYATWRFSGQDMADFA	ILPSCCRWRIRKEFPKSEGQYSGFKSPY
P2X7v02	-----	
P2X7v03	-----	
	541.....550.....560.....570.....580.....590	

Figure 16e



**Figure 17**




Figure 18




Figure 19a




Figure 19b






Figure 19c




Figure 20




Figure 21




Figure 22




Figure 23




Figure 24

## MEANS AND METHODS FOR DIAGNOSING AND TREATING AFFECTIVE DISORDERS

[0001] The present invention relates to nucleic acid molecules, preferably genomic sequences, encoding an ATP-gated ion channel P2X7R which contain a mutation in the 5'UTR or 3'UTR regions, a mutation in exon 3, 5, 6, 8 or 13 or in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 or a deletion in exon 13, which allow to diagnose affective disorders. The invention further relates to polypeptides encoded by said nucleic acid molecules vectors and host cells comprising said nucleic acid molecules as well as to methods for producing polypeptides encoded by said nucleic acid molecules. The present invention also provides antibodies specifically directed to polypeptides encoded by said nucleic acid molecules and aptamers specifically binding said nucleic acid molecules.

[0002] Additionally, primers for selectively amplifying said nucleic acid molecules are provided in the present invention as well as kits, compositions, particularly pharmaceutical and diagnostic compositions comprising said nucleic acid molecules, vectors, polypeptides, aptamers, antibodies and/or primers. Moreover, the present invention relates to methods for diagnosing affective disorders associated with a non-functional P2X7R protein, an altered ATP-gating of the P2X7R protein, an over- or underexpression of the P2X7R protein or associated with the presence of any one of the aforementioned nucleic acid molecules or polypeptides encoded thereby. Additionally, the present invention relates to uses and methods for treating affective disorders employing a functional or non-functional ATP-gated ion-channel P2X7R. The present invention also relates to uses of modulators of P2X7R activity for treating affective diseases.

[0003] Furthermore, the present invention also relates to methods for identifying and characterizing compounds which are capable of specifically interacting with or altering the characteristics of the polypeptides of the present invention as well as to methods for the production of pharmaceutical compositions.

[0004] Up to 10% of persons visiting a physician are afflicted with an affective disorder (also known as behavioural disorder, mood disorder). Nonetheless, most cases remain undiagnosed or inadequately treated. Affective disorders include among others, depression, anxiety, and bipolar disorder. These diseases are well described in the literature; see, for example, Diagnostic and Statistical Manual of Mental Disorders—4th Edition Text Revision (DSM-IV-TR), American Psychiatric Press, 2000.

[0005] Depression, also known as unipolar affective disorder, is characterized by a combination of symptoms such as lowered mood, loss of energy, loss of interest, feeling of physical illness, poor concentration, altered appetite, altered sleep and a slowing down of physical and mental functions resulting in a relentless feeling of hopelessness, helplessness, guilt, and anxiety. The primary subtypes of this disease are major depression, dysthymia (milder depression), and atypical depression. Other important forms of depression are premenstrual dysphoric disorder and seasonal affective disorder. Present treatment of depression consists of psychotherapy, antidepressant drugs, or a combination of both. Most antidepressive drugs target the transport of the neurotransmitters serotonin and/or norepinephrine, or the activ-

ity of the enzyme monoamine oxidase. They include: Selective serotonin-reuptake inhibitors (e.g., fluoxetine, paroxetine, sertraline, fluvoxamine), tricyclic antidepressants (e.g., amitriptyline, imipramine, desipramine, nortriptyline), monoamine oxidase inhibitors (e.g., phenelzine, isocarboxazid, tranylcypromine), and designer antidepressants such as mirtazapine, reboxetine, nefazodone. However, all existing antidepressive drugs possess shortcomings such as long latency until response, high degree of non-responders and undesirable side effects (Holsboer, *Biol. Psychol.* 57 (2001), 47-65). Therefore, a need exists in the medical community for new antidepressive drugs with improved pharmacological profile (Baldwin, *Hum. Psychopharmacol. Clin. Exp.* 16 (2001), S93-S99).

[0006] Anxiety disorders are defined by an excessive or inappropriate aroused state characterized by feelings of apprehension, uncertainty, or fear. They are classified according to the severity and duration of their symptoms and specific affective characteristics. Categories include: (1) Generalized anxiety disorder, (2) panic disorder, (3) phobias, (4) obsessive-compulsive disorder, (5) post-traumatic stress disorder, and (6) separation anxiety disorder. The standard treatment for most anxiety disorders is a combination of cognitive-behavioural therapy with antidepressant medication. Additional medications include benzodiazepines and buspirone.

[0007] Bipolar disorder, also known as manic-depression, is characterized by mood swings between periods of mania (i.e. mood elevation including exaggerated euphoria, irritability) and periods of depression. Bipolar disorder is classified according to the severity of the symptoms. Patients diagnosed with bipolar disorder type I suffer from manic or mixed episodes with or without major depression. In Bipolar Disorder type II, patients have episodes of hypomania and episodes of major depression. With hypomania the symptoms of mania (euphoria or irritability) appear in milder forms and are of shorter duration. The current drugs used to treat bipolar disorders are lithium, valproate and lamotrigine, which stimulates the release of the neurotransmitter glutamate. As with antidepressive drugs, they take weeks to become effective and can result in undesirable side effects, for example, high levels of lithium in the blood can be fatal.

[0008] Compelling evidence suggest that affective disorders are biological diseases. However, there are no laboratory tests or other procedures that a common physician can use to make a definitive diagnosis. Instead, a specially trained physician or psychiatrist must diagnose the illness based on a group of symptoms that occur together. This process is often time consuming and laborious requiring several visits for the physician to perform a careful history of the symptoms that the patient is currently experiencing as well as any symptoms he or she has had in the past. Therefore, an easy and effective method for the accurate diagnosis of affective disorders is of high interest to the medical community (Wittchen et al., *J. Clin. Psychiatry* 62, suppl. 26 (2001), 23-28).

[0009] Most patients afflicted with affective disorders have family antecedents and identical twins studies suggest a strong genetic component. For example, genetic mapping on an isolated population of the central valley of Costa Rica suggests a locus for severe bipolar disorder at chromosome 18q22-q23 (Freimer et al., *Nature Genetics* 12 (1996),

436-441). Moreover, genetic studies performed on the Old Order Amish population suggest that genes on chromosomes 6, 13, and 15 may contribute to the susceptibility of bipolar affective disorder (Ginns et al., *Nature Genetics* 12 (1996), 431-435). Recently, a genome-wide search in a homogenous population found in the Saguenay/Lac-St-Jean region of Quebec suggests the presence of a major locus for bipolar disorder on chromosome 12q23-q24 (Morissette et al., *Am. J. Med. Genet. (Neuropsychiatr. Genet.)* 88 (1999), 567-587). Susceptibility loci on chromosomes 5 and 21 were also found in this study. Other groups report minimal evidence for linkage in the region of 12q23 (Kelsoe et al., *Proc. Natl. Acad. Sci. USA* 98 (2001), 585-590; Sklar, *Annu. Rev. Genomics Hum. Genet.* 3 (2002), 371-413). Given the various loci mentioned in the above studies (e.g., links to chromosomes 5, 6, 12, 13, 15, 18, 21), a definite genetic link for affective diseases remains to be found.

[0010] Thus, although several genes have been assumed to be linked with affective disorders as mentioned hereinabove, however, no clear correlation has so far been shown. Since no well-suited medication nor diagnosis on a molecular level for affective disorders is available, there is a need for identifying a gene whose mutations cause the whole spectrum of affective disorders as well as for providing medicaments and methods for diagnosis and treatment of affective disorders.

[0011] Thus, the technical problem underlying the present invention is to provide means and methods for diagnosis and treating affective disorders.

[0012] The solution to said technical problem is achieved by providing the embodiments characterized in the claims.

[0013] Accordingly, the present invention relates to a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of:

[0014] (a) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 5'UTR region corresponding to positions 362, 532, 1100, 1122, 1171 or 1702 of the genomic sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;

[0015] (b) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of the ATP-gated ion channel P2X7R, wherein in the exon as indicated in column "Exon" of the following Table A the amino acid residue as indicated in column "Amino acid residue" of Table A corresponding to the position as indicated in column "Position in wild-type" of Table A of the wild-type ATP-gated ion channel P2X7R amino acid sequence as depicted in SEQ ID NO: 3 or 4 is replaced by another amino acid residue

TABLE A

Exon	Amino acid residue	Position in wild-type
exon 3	R (Arg)	117
exon 5	G (Gly)	150
exon 6	E (Glu)	186
exon 6	L (Leu)	191

TABLE A-continued

Exon	Amino acid residue	Position in wild-type
exon 8	R (Arg)	270
exon 13	I (Ile)	568
exon 13	R (Arg)	578

[0016] (c) a nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in exon 5 or 8 corresponding to position 32548 or position 37633 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide

[0017] (d) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted;

[0018] (e) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R, wherein in the intron as indicated in column "Intron" of the following Table B the nucleotide as indicated in column "Replaced nucleotide" of Table B corresponding to the position as indicated in column "Position in wild-type" of Table B of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1 is replaced by another nucleotide

TABLE B

Intron	REPLACED NUCLEOTIDE	Position in wild-type
intron 1	G	3166
intron 1	C	24778
intron 1	C	24830
intron 3	A	26308
intron 3	G	26422
intron 4	G	32394
intron 4	T	32434
intron 5	A	32783
intron 6	G	35641
intron 6	A	35725
intron 6	T	36001
intron 7	G	36378
intron 7	T	36387
intron 7	G	36398
intron 9	C	47214
intron 11	T	47563
intron 12	C	54307
intron 12	G	54308

[0019] (f) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 3'UTR region corresponding to position 55169, 55170, 55171, 55917 or 54925 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;

[0020] (g) a nucleotide sequence comprising at least 20 or 21 nucleotides and comprising the mutations or deletions as defined in any one of (a) to (f);



- [0021] (h) a nucleic acid sequence comprising a nucleotide sequence as shown in any one of SEQ ID NOs: 13 to 51;
- [0022] (i) a nucleic acid sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NOs: 5 to 12;
- [0023] (j) a nucleotide sequence which hybridizes to a nucleotide sequence defined in any one of (a) to (g) or to the nucleotide sequence of (h) and having a mutation as defined in any one of (a) to (f); and
- [0024] (k) a nucleic acid sequence being degenerate as a result of the genetic code to the nucleic acid sequence as defined in (j).

[0025] It has surprisingly been found that mutations in the P2X7R gene which encodes the ATP-gated ion channel P2X7R can cause the whole spectrum of affective disorders. Six different mutations in the 5'UTR of the P2X7R gene, seven different mutations in exons 3, 5, 6, 8 and 13 of the P2X7R gene leading to an amino acid replacement of the corresponding amino acid in the wild-type sequence of P2X7R depicted in SEQ ID NO: 3 or 4 and two mutations in exons 5 and 8 of said gene, respectively, leading to a replacement of a nucleotide by another nucleotide, a deletion of nucleotides in exon 13 of said gene, 18 mutations in introns 1, 3, 4, 5, 6, 7, 9, 11 and 12 and 5 mutations in the 3'UTR of the P2X7R gene have been identified to co-segregate with the affection status in 41 unrelated families affected with affective disorders. The term "affective disorder" when used in the context of the present invention means to include, but is not limited to, depression, anxiety, unipolar disorder, bipolar disorder type I, bipolar disorder type II, mania, attention deficit hyperactive disorder, substance abuse, and any other disorders affecting the normal behaviour, or mood of an individual.

[0026] Each mutation causes alterations that can explain affective disorders as shown in the Examples hereinbelow.

[0027] P2X7R is an ATP-gated ion channel belonging to the P2X ionotropic channel family. The gene was first isolated from rat brain (Surprenant et al., (1996), 272, 735-738; Genbank accession number NM\_019256) and subsequently from a human monocyte library (Rassendren et al., J. Biol. Chem. 272 (1997), 5482-5486; Genbank accession numbers NM\_002562, Y09561) by virtue of its sequence homology with the other members of the P2X family. It was later found that P2X7R corresponded to the unidentified P2Z receptor which mediates the permeabilising action of ATP on mast cells and macrophages (Dahlqvist and Diamant, Acta Physiol. Scand. 34 (1974), 368-384; Steinberg and Silverstein, J. Biol. Chem. 262 (1987), 3118-3122; Gordon, Biochem. J. 233 (1986), 309-319). The P2X7R has two hydrophobic membrane-spanning domains, an extracellular loop, and forms transmembrane ion channels. P2X7 receptors seem to function only in homooligomeric form and bear a pharmacological profile markedly different from other P2X homo- or heteromers (North and Surprenant, Annual Rev. Pharmacology Toxicology 40 (2000), 563-580). P2X7R requires levels of ATP in excess of 1 mM to achieve activation, whereas other P2X receptors activate at ATP concentrations of  $\leq 100 \mu\text{M}$  (Steinberg et al., J. Biol. Chem. 262 (1987), 8884-8888; Greenberg et al., J. Biol. Chem. 263 (1988), 10337-10343) 32). While all P2X

receptors demonstrate non-selective channel-like properties following ligation, the channels formed by the P2X7R can rapidly transform into pores that can allow the passage of molecules of up to 900 Dalton (Virginio et al., J. Physiol. 519 (1999), 335-346).

[0028] P2X7R is expressed in hematopoietic cells, mast cells and macrophages (Surprenant et al., Science 272 (1996), 3118-3122), where it is organized in tetrameric or hexameric form (Kim et al., J. Biol. Chem. 276 (2001), 23262-23267). P2X7R is inter alia involved in the regulation of the immune function and inflammatory response. Activation of P2X7R by ATP in macrophages is associated with mitogenic stimulation of T cells (Baricordi et al., Blood 87 (1996), 682-690), the release of cytokines such as interleukin-1 $\beta$  (Griffiths et al., J. Immunol. 154 (1995), 2821-2828), and formation of macrophage polykaryons (Falzoni et al., J. Clin. Invest. 95 (1995), 1207-1216). Stimulation of the P2X7R with ATP can also result in cell death by triggering massive transmembrane ion fluxes (particularly influx of Ca<sup>2+</sup> and Na<sup>+</sup>, and efflux of K<sup>+</sup>) and the formation of non-selective plasma membrane pores (Di Virgilio et al., Cell Death Differ. 5 (1998), 191-199).

[0029] In the brain, P2X7R was originally thought to be restricted to microglia (resident macrophage of the brain) and ependymal cells rather than neurons (Collo et al., Neuropharmacology 36 (1997), 1277-1283) suggesting a role of P2X7R in neurodegeneration. However, P2X7R has since been found in neurons of the rat retina (Brandle et al., Brain Research Molecular Brain Res. 62 (1998), 106-109), cochlear ganglion cells (Brandle et al., Neuroscience Letters 273 (1999), 105-108), and presynaptic terminals of neurons throughout the brainstem and spinal cord (Deuchards et al., J. Neurosci. 21 (2001), 7143-7152). Subsequent studies also suggest that P2X7R regulates the release of neurotransmitters such as glutamate and GABA in neurons of the hippocampus (Armstrong et al., J. Neuroscience 22 (2002), 5938-5945, Sperl gh et al., J. Neurochem. 81 (2002), 1196-1211). Organisation of P2X7R in glial cells and astrocytes of the brain appears monomeric (Kim et al., J. Biol. Chem. 276 (2001), 23262-23267).

[0030] Several agonists and antagonists of P2X7R have been identified. Brilliant Blue (Jiang et al., Mol. Pharmacol. 58 (2000), 82-88), the isoquinolines 1-[N,O-Bis(5-isoquinolinesulfonyl)-N-methyl-L-tyrosyl]-4-phenylpiperazine and N-[1-[N-methyl-p-(5-isoquinolinesulfonyl)benzyl]-2-(4-phenylpiperazine)ethyl]-5-isoquinolinesulfonamide (Humphreys et al., Mol. Pharmacol., 54 (1998), 22-32), adamantane derivatives (WO 99/29660, WO 99/29661, WO 00/61569, WO 01/42194, WO 01/44170, WO 01/44213), substituted phenyl compounds (WO 00/71529), piperidine and piperazine derivatives (WO 01/46200) are antagonists of P2X7R while Oxidized ATP (oATP) acts as an irreversible inhibitor of the receptor (Chen et al., J. Biol. Chem., 268 (1993), 8199-8203). Some of these antagonists are presently being evaluated for the treatment of inflammatory, immune, and cardiovascular diseases. BzATP (2'-3'-O-(4-Benzoylbenzoyl)adenosine 5'-triphosphate (C<sub>24</sub>H<sub>24</sub>N<sub>5</sub>O<sub>15</sub>P<sub>3</sub>)) acts as agonist of P2X7R (North and Surprenant, Annu. Rev. Pharmacol. Toxicol. 40 (2000), 563-580). WO 99/55901 describes a method for identifying compounds that modulate the activity of a mammalian purinoreceptor selected from the group consisting of P2X2, P2X3, P2X4, P2X5, P2X6 and P2X7 and suggests a role of said purinoreceptors in

therapy of behavioural disorders such as epilepsy, depression and aging-associated degenerative diseases.

[0031] Mutant mice lacking P2X7R are healthy, fertile and demonstrate no overt phenotype. However, in contrast to their wild-type counterparts, LPS-activated peritoneal macrophages from P2X7R<sup>-/-</sup> animals fail to generate mature interleukin-1 $\beta$  (IL-1 $\beta$ ) when challenged with ATP suggesting an inability of peritoneal macrophages to release IL-1 in response to ATP (Solle et al, J. Biol. Chem. 276 (2001), 125-132). A detailed behavioural study of the P2X7R<sup>-/-</sup> mice was not performed. In humans, a Glu-496 to Ala polymorphism leads to the loss of P2X7 function (Gu et al., J. Biol. Chem. 276 (2001), 11135-11142) and is associated with B-cell chronic lymphocytic leukaemia (Thunberg, et al, The Lancet 360 (2002), 1935-1939). Additional polymorphisms in the putative P2X7R promoter region, and coding region have been reported (Li et al., FEBS Lett. 531 (2002), 127-131; EP 1199372).

[0032] Despite the abundant literature concerning P2X7R, a role in affective disorders has never been suggested or alluded to in the prior art.

[0033] Before the present invention is described in detail, it is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors, and reagents described herein as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art.

[0034] Preferably, the terms used herein are defined as described in "A multilingual glossary of biotechnological terms: (IUPAC Recommendations)", Leuenberger, H. G. W, Nagel, B. and Kolbl, H. eds. (1995), Helvetica Chimica Acta, CH-4010 Basel, Switzerland).

[0035] Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integer or step.

[0036] Several documents are cited throughout the text of this specification. Each of the documents cited herein (including all patents, patent applications, scientific publications, manufacturer's specifications, instructions, etc.), whether supra or infra, are hereby incorporated by reference in their entirety. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

[0037] It must be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the", include plural referents unless the context clearly indicates otherwise. Thus, for example, reference to "a reagent" includes one or more of such different reagents, and reference to "the method" includes reference to equivalent steps and methods known to those of ordinary skill in the art that could be modified or substituted for the methods described herein.

[0038] In accordance with the present invention, the term "nucleic acid sequence" means the sequence of bases comprising purine- and pyrimidine bases which are comprised by nucleic acid molecules, whereby said bases represent the primary structure of a nucleic acid molecule. Nucleic acid sequences include DNA, cDNA, genomic DNA, RNA, synthetic forms and mixed polymers, both sense and antisense strands, or may contain non-natural or derivatized nucleotide bases, as will be readily appreciated by those skilled in the art.

[0039] When used herein, the term "polypeptide" means a peptide, a protein, or a polypeptide which encompasses amino acid chains of a given length, wherein the amino acid residues are linked by covalent peptide bonds. However, peptidomimetics of such proteins/polypeptides wherein amino acid(s) and/or peptide bond(s) have been replaced by functional analogs are also encompassed by the invention as well as other than the 20 gene-encoded amino acids, such as selenocysteine. Peptides, oligopeptides and proteins may be termed polypeptides. The terms polypeptide and protein are often used interchangeably herein. The term polypeptide also refers to, and does not exclude, modifications of the polypeptide, e.g., glycosylation, acetylation, phosphorylation and the like. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature.

[0040] The term "position" used in accordance with the present invention means the position of either an amino acid within an amino acid sequence depicted herein or the position of a nucleotide within a nucleic acid sequence depicted herein.

[0041] The term "ATP-gated ion channel P2X7R", in accordance with this invention, denotes a polypeptide which can be classified as a member of the P2X ionotropic receptor family. They are also known as purinergic receptors. P2X receptors are ligand-gated ion channels. The ligand for these receptors may be ATP and/or another natural nucleotide such as ADP, UTP and UDP, or a synthetic nucleotide such as 2-methylthioATP. The criteria for the classification are: (1) a sequence homology that is higher than 39% across the family or different species; (2) signal transduction mechanism involving ion conductance (Khakh et al., Pharmacol Rev. 253 (2001), 107-18). Accordingly, the term "ATP-gated ion channel P2X7R" is interchangeable with the terms "ionotropic receptor" or "purinergic receptor". Preferably, the term "ATP-gated ion channel P2X7R" denotes a polypeptide which can be classified as an ATP-gated ion channel P2X7R on the basis of one or more structural and/or functional characteristics, preferably those described above. Structural characteristics refer to certain structural features which allow to classify a polypeptide as being a P2X7R protein. One such feature is the amino acid sequence. In the context of the present invention a polypeptide is classified as an ATP-gated ion channel P2X7R if it shows a certain degree of sequence identity over its own length to the amino acid sequence of the human P2X7R protein depicted in SEQ ID NO: 3 or 4. This degree of sequence identity is at least 40%, more preferably at least 50%, even more preferably at least 60%, at least 70%, at least 80%, at least 90% or at least 95%. It is particularly preferred that the degree of sequence identity is at least 65%.

[0042] Moreover, structural characteristics of P2X7R proteins are two hydrophobic membrane-spanning domains, an

extra cellular loop which could be analyzed by using the program TMPRED (Hofmann *Biol. Chem.* 347 (1993), 166) or TMHMM (Krogh *J. Mol. Bio.* 305 (2001), 567-580). Additionally, P2X7R may exist as a single polypeptide, as dimer, tetramer or the like.

**[0043]** Thus, in the context of the present invention a protein is preferably classified as a P2X7R protein if it displays at least one of the above-mentioned structural characteristics. Functional characteristics refer to properties related to the biological activity of the P2X7R protein. In particular, P2X7R is an ATP-gated ion channel which allows calcium and sodium ions to pass from extracellular solution to intracellular solution, and allows potassium ions to pass from intracellular to extracellular solution. Moreover, the ATP-gated ion channel P2X7R forms naturally a homooligomeric form. The characteristics of P2X7R receptor proteins can be determined as mentioned hereinbelow. The term "ATP-gated ion channels P2X7R" comprises functional and non-functional forms of the ATP-gated ion channels P2X7R. A functional ATP-gated ion channel P2X7R is understood to be a P2X7R protein which has at least one of the above-mentioned functional characteristics which can be measured by methods known in the art. A non-functional ATP-gated ion channel P2X7R is a protein which can be classified as a P2X7R protein due to structural characteristics as described above but which has lost at least one, preferably all, functional characteristics of a P2X7R protein as described above. Non-functionality of the P2X7R protein can, e.g., be determined by measuring whether calcium and sodium ions can flow into cells or whether potassium ions can exit from cells. Thus, it is possible to determine the occurrence of a mutation in the ATP-gated ion channel P2X7R by measuring either calcium and/or sodium influx or efflux of cells. Cells harbouring a mutation in the P2X7R gene show an altered ion influx and/or efflux in comparison to cells harbouring a wild-type P2X7R protein. Additionally, there are different methods that could be used to determine whether the P2X7R is functional or non-functional, for example, altered. One method consists of measuring the rate of ATP-induced incorporation of ethidium into cells, e.g. cells isolated from an individual. Ethidium is incorporated into the cells through P2X7R pores, when the pore formation is activated by ATP. Cells are then incubated with or without ATP in the presence of ethidium, then they are analyzed by flow cytometry. Ethidium fluorescence is measured and compared in the presence or absence of ATP. If the P2X7R has lower activity, the ethidium fluorescence induced by ATP will be lower than in control cells. Such a method was used to verify P2X7R activity in isolated B-lymphocytes and T-lymphocytes from leukaemia patients (Wiley et al., *Lancet* 359 (2002), 1114-1119). Briefly, isolated cells are incubated in 1 ml of Hepes buffered potassium chloride at 37° C. with continuous stirring. Ethidium is then added at a concentration of 25 mol/l, followed 40 seconds later by the addition of 10  $\mu$ l of 100 mmol/l ATP stock. Cells are analyzed at 1,000 events/s by flow cytometry using a Coulter Elite flow cytometer (Coulter, Hialeah, Fla.) with argon laser excitation at 488 nm. Fluorescent emission was collected using a 590-nm long-pass filter. The linear mean channel fluorescence intensity for each gated subpopulation over successive 5-s intervals was analyzed with the use of Win-MDI software (Joseph Trotter, version 2.7) and plotted against time.

**[0044]** Another method of determining P2X7R activity is to measure calcium entry into isolated cells incubated with

fluorescent dyes that emit only upon binding to calcium. The cells have to be loaded with the dye and then the calcium entry has to be stimulated. Examples of such dyes include Fura-2, Calcium green, calcium orange, calcium crimson (all available from Molecular Probes). Methods of measuring calcium transport are well known in the art; see for example, Takahashi et al., *Physiol. Rev.* 79 (1999), 1089-1125. Furthermore, calcium entry into the cells produces changes in the membrane electric potential. This changes can be measured by electrophysiology (patch clamp) or by using dyes which are sensitive to voltage change. Such methods are also well known in the art, see for example, Gonzalez et al., *DDT* 4 (1999), 431-439; Gonzalez and Tsien, *Chemistry & Biology* 4 (1997), 269-277; Gonzalez and Tsien, *Biophysical Journal* 69 (1995), 1272-1280.

**[0045]** Yet another method is to measure uptake of  $^{133}\text{Ba}^{21}$ . Ba21 is a good surrogate for Ca21 and once inside the cell is neither pumped nor sequestered by transport ATPases. Ba21 uptake can be measured over 60 s using  $^{133}\text{BaCl}_2$  (final concentration, 0.2 mM). At time 0, a prewarmed stock solution of  $^{133}\text{Ba}^{21}$  (0.4 mM and 1  $\mu\text{Ci/ml}$ ) is added in equal volumes to prewarmed isolated cells in 150 mM KCl with HEPES (pH 7.4) at 37° C. ATP (1 mM) is added either 10 minutes before or simultaneously with the  $^{133}\text{Ba}^{21}$  isotope. Aliquots of 0.8 ml are taken at time points between 0 and 60 s and are immediately mixed with 0.2 ml of ice-cold 50 mM  $\text{MgCl}_2$  (in KCl-HEPES medium) that had been previously layered over 250  $\mu\text{l}$  of oil mixture (di-n-butyl phthalate and di-iso-octyl phthalate, 7:3 vol/vol) and then centrifuged at 8,000 g for 30 s. The supernatants and the oil are aspirated, and the cell pellets are counted in a Wallac Wizard 3 automatic gamma-counter or in any other suitable gamma measuring unit.

**[0046]** The present invention is based on the finding that mutations of different kinds in the P2X7R gene are linked to the occurrence of affective disorders. The first type of mutations are mutations in the 5'UTR. Examples of such mutations are single nucleotide replacements at positions corresponding to positions 362, 532, 1100, 1122, 1171 or 1702 of the genomic sequence of the wild-type ATP-gated ion channel P2X7R as described in SEQ ID NO: 1.

**[0047]** The position with respect to nucleotide sequences mentioned herein refer to the sequence shown in SEQ ID NO: 1. This sequence represents the nucleic acid sequence of the P2X7R gene encoding the ATP-gated ion channel P2X7R. It is possible for the skilled person to identify the position in the genomic sequence corresponding to a position in SEQ ID NO: 1 by aligning the sequences. Moreover, the exact locations of the exons and introns are indicated in SEQ ID NO: 1 hereinbelow. Additionally, the person skilled in the art is able to identify exons and introns of the P2X7R gene by comparing SEQ ID NO: 1 with SEQ ID NO: 2 which shows the cDNA sequence of the P2X7R gene.

**[0048]** Preferably, at position 362 in the 5'UTR of the genomic sequence of the P2X7R gene depicted in SEQ ID NO: 1 a thymine (T) is replaced by another nucleotide, preferably a purine base. More preferably, at said position said thymidine is replaced by a pyrimidine base. Particularly preferred, said thymine is replaced by a cytosine (C).

**[0049]** At position 532 in the 5'UTR of the genomic sequence of the P2X7R gene depicted in SEQ ID NO: 1 a thymine (T) is preferably replaced by another nucleotide,

preferably a pyrimidine base. More preferably, at said position said thymine is replaced by a purine base. Particularly preferred, said thymidine is replaced by a guanine (G).

[0050] The adenine (A) residues at positions 1100 and 1122, respectively, in the 5'UTR of the genomic sequence of the P2X7R gene is preferably replaced by a pyrimidine base. More preferably, said adenine is replaced by a purine base and particularly preferred said adenine is replaced by a guanine (G).

[0051] At position 1171 in the 5'UTR of the genomic sequence of the P2X7R gene depicted in SEQ ID NO: 1 a cytidine (C) is replaced by another nucleotide, preferably by a pyrimidine base. More preferably, said cytidine is replaced by a purine base and even more preferred, said cytidine is replaced by a guanine (G).

[0052] The guanine at position 1702 in the 5'UTR of the genomic sequence of the gene P2X7R depicted in SEQ ID NO: 1 is replaced by another nucleotide, preferably by a pyrimidine base. More preferably, said guanine is replaced by a purine base and particularly preferred it is replaced by an adenine (A).

[0053] A second type of mutation found in the P2X7R gene are mutations in exons which lead to amino acid substitutions in the corresponding amino acid sequence. These are the mutations listed under item (b), supra. In this context, the term "an amino acid residue as indicated in column 'Amino acid residue' of Table A corresponding to position X of the wild-type ATP-gated ion channel P2X7R as depicted in column 'Position in wild-type'" has the following meaning: The amino acid residue in question would be located at position X in the sequence of SEQ ID NO: 3 or 4 if the sequence in which said amino acid residue occurs is compared and aligned with the amino acid sequence of SEQ ID NO: 3 or 4. The amino acid sequence shown in SEQ ID NO: 3 or 4 is the amino acid sequence of the human P2X7R gene and is used as a reference sequence in the present invention.

[0054] In order to determine whether an amino acid residue or nucleotide residue in a given P2X7R sequence corresponds to a certain position in the amino acid sequence or nucleotide sequence of SEQ ID NO: 1, 3 or 4, the skilled person can use means and methods well-known in the art, e.g., alignments, either manually or by using computer programs such as those mentioned further down below in connection with the definition of the term "hybridization" and degrees of homology.

[0055] For example, BLAST2.0, which stands for Basic Local Alignment Search Tool (Altschul, Nucl. Acids Res. 25 (1997), 3389-3402; Altschul, J. Mol. Evol. 36 (1993), 290-300; Altschul, J. Mol. Biol. 215 (1990), 403410), can be used to search for local sequence alignments. BLAST produces alignments of both nucleotide and amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST is especially useful in determining exact matches or in identifying similar sequences. The fundamental unit of BLAST algorithm output is the High-scoring Segment Pair (HSP). An HSP consists of two sequence fragments of arbitrary but equal lengths whose alignment is locally maximal and for which the alignment score meets or exceeds a threshold or cutoff score set by the user. The BLAST approach is to look for

HSPs between a query sequence and a database sequence, to evaluate the statistical significance of any matches found, and to report only those matches which satisfy the user-selected threshold of significance. The parameter E establishes the statistically significant threshold for reporting database sequence matches. E is interpreted as the upper bound of the expected frequency of chance occurrence of an HSP (or set of HSPs) within the context of the entire database search. Any database sequence whose match satisfies E is reported in the program output.

[0056] Analogous computer techniques using BLAST (Altschul (1997), loc. cit.; Altschul (1993), loc. cit.; Altschul (1990), loc. cit.) are used to search for identical or related molecules in nucleotide databases such as GenBank or EMBL. This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

[0057] and it takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1-2% error; and at 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

[0058] As mentioned above, the second group of mutations identified in the P2X7R gene are mutations in the exons of the P2X7R gene which lead to amino acid substitutions. In this respect SEQ ID NO 2 shows the cDNA sequence of the P2X7R gene. In exon 3 at position 117 of the corresponding wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 an arginine (R) residue is replaced by another amino acid residue, preferably by an aliphatic, acidic or basic amino acid residue. More preferably, by an aromatic amino acid residue which is particularly preferred to be a tryptophane (W). The resulting polypeptide is shown in SEQ ID NO: 5.

[0059] In exon 5 at position 150 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 a glycine (G) residue is replaced by another amino acid residue, preferably by an aliphatic, aromatic or acidic amino acid residue. More preferably, by a basic amino acid residue and particularly preferred by an arginine (R). The resulting polypeptide is shown in SEQ ID NO: 6.

[0060] At position 186 in exon 6 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 a glutamate residue (E) is replaced by another amino acid residue, preferably by an aliphatic, aromatic or acidic amino acid residue. More preferably, said glutamate is replaced by a basic amino acid residue which is particularly preferred a lysine (K). The resulting polypeptide is shown in SEQ ID NO: 7.

[0061] In exon 6 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 at position 191 a leucine residue (L) is replaced by another amino acid

residue. Said amino acid residue is preferably an aliphatic, acidic or basic amino acid residue. More preferably, said amino acid residue is an aromatic amino acid residue which is particularly preferred to be a proline (P). The resulting polypeptide is shown in SEQ ID NO: 8.

**[0062]** In exon 8 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 at position 270 an arginine residue (R) is replaced by another amino acid residue. Said amino acid residue is preferably an aromatic, acidic or basic amino acid residue. More preferably, said amino acid residue is an aliphatic amino acid residue which is particularly preferred to be a cysteine (C). The resulting polypeptide is shown in SEQ ID NO: 9.

**[0063]** At position 568 in exon 13 of the wild-type amino acid residue of P2X7R depicted in SEQ ID NO: 3 or 4 an isoleucine (I) residue is replaced by another amino acid residue. More preferably, said isoleucine is replaced by an aromatic, basic or acidic amino acid residue. Even more preferred, said isoleucine is replaced by an aliphatic amino acid residue which is particularly preferred to be an asparagine (N). The resulting polypeptide is shown in SEQ ID NO: 10.

**[0064]** In exon 13 at position 578 in the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 an arginine residue (R) is replaced by another amino acid residue. Said amino acid residue is preferably an aromatic, acidic or basic amino acid residue. More preferably, it is an aliphatic amino acid residue and particularly preferred it is a glutamine (Q) residue. The resulting polypeptide is shown in SEQ ID NO: 12.

**[0065]** It is envisaged that the above-mentioned mutations in the exons of the P2X7R gene occur due to point mutations caused by, e.g. chemical and/or physical means or inaccuracy of the replication complex followed by a failure of the reparation machinery of a cell, can result in a change of a single codon. Possible types of point mutations are transitions, i.e. change of a purine or pyrimidine base for another purine or pyrimidine base, e.g. adenine to guanine or thymidine to cytosine or transversions, i.e. change of a purine or pyrimidine base for another pyrimidine or purine base, e.g., adenine to thymidine or guanine to cytosine. Additionally a point mutation can also be caused by insertion or deletion of one or more nucleotides.

**[0066]** The mutations leading to the replacement of the amino acids as mentioned hereinabove and hereinbelow are indicated in Table 1 hereinbelow.

**[0067]** The third group of mutations in the P2X7R gene has been identified to be in exons 5 and 8 of the P2X7R gene depicted in SEQ ID NO: 1 and to be silent, i.e. they do not lead to amino acid changes. In particular, at position 32548 in exon 5 of the wild-type genomic sequence P2X7R gene depicted in SEQ ID NO: 1 a cytidine residue is replaced by another nucleotide. Said nucleotide is preferably a pyrimidine base and particularly preferred a thymine. The exchange of the cytidine residue at position 32548 in exon 5 of the P2X7R gene by another nucleotide preferably does not lead to the replacement of the amino acid cysteine by another amino acid residue.

**[0068]** In exon 8 of the wild-type P2X7R gene depicted in SEQ ID NO: 1 at position 37633 a cytidine residue is replaced by another nucleotide residue. Said nucleotide

residue is preferably a pyrimidine base and particularly preferred thymine. Due to this replacement the amino acid aspartate (D) encoded by the respective codon in which at position 37633 a replacement has taken place is preferably not replaced by another amino acid residue.

**[0069]** The above-mentioned mutations in exons 5 and 8 at positions 32548 and 37633, respectively, of the wild-type P2X7R gene depicted in SEQ ID NO: 1 are mutations at the third position of a triplet codon, i.e. at the wobble base, which lead to so-called silent mutations. Silent mutations do normally not lead to a change of the amino acid due to the degeneracy of the genetic code, i.e. 64 triplets encode at all 20 naturally occurring amino acids. However, said silent mutations lead to a change in the codon encoding its respective amino acids insofar that the newly generated codon may not fit so well into the codon usage of an organism. Namely, the newly generated codon is not translated by the ribosome with the same efficiency as the "old" codon. This may lead to insufficient amounts of the corresponding polypeptide causing a distinct phenotype.

**[0070]** The fourth group of mutations in the P2X7R gene described hereinabove in item (d) is a deletion of 7 amino acids corresponding to positions 488 to 494 of the wild-type P2X7R amino acid sequence as depicted in SEQ ID NO: 3 or 4. Thus, the present invention also relates to nucleic acid sequences encoding a P2X7R protein in which amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted. This means, according to the present invention, that a fragment encompassing amino acid positions 488 to 494 of the corresponding wild-type amino acid sequence depicted in SEQ ID NO: 3 or 4 is deleted which results in a shortened polypeptide. An example for such a shortened polypeptide is depicted in SEQ ID NO: 11. This type of mutation as described herein preferably encodes a non-functional ATP-gated ion channel P2X7R. In the present invention the deletion of a fragment encompassing amino acids 488 to 494 of the wild-type amino acid sequence depicted in SEQ ID NO: 3 or 4 is the result of a deletion in exon 13. The resulting protein depicted in SEQ ID NO: 11 lacks amino acids 488 to 494 of the corresponding wild type amino acid sequence depicted in SEQ ID NO: 3 or 4 such that amino acid position 494 of the deleted polypeptide depicted in SEQ ID NO: 11 corresponds to amino acid position 502 of the wild-type amino acid sequence depicted in SEQ ID NO: 3 or 4. Preferably, the nucleic acid sequence of the invention encodes a P2X7R polypeptide in which exactly amino acids corresponding to positions 488 to 494 of SEQ ID NO: 3 or 4 are deleted. However, also mutants are comprised in which either more or less amino acids within the P2X7R amino acid sequence set forth in SEQ ID NO: 3 or 4 may be deleted due to, for example, atypical splicing or deletion of nucleotides of the nucleic acid molecule encoding P2X7R or wrong posttranslational processes, as long as the P2X7R ATP-gated ion channel is non-functional. For example, it is also possible that further amino acids preceding amino acid position 488 or amino acids succeeding amino acid position 494 may be deleted or that less amino acids are deleted. Preferably at least one, more preferably at least two, even more preferably at least three and most preferably at least 5 amino acid residues are further deleted upstream from the position

corresponding to amino acid residue 488 and/or downstream of the position corresponding to amino acid residue 494 of SEQ ID NO: 3 or 4.

[0071] However, it is preferred that not more than 20, preferably not more than 15, even more preferably not more than 10 and most preferably not more than 7 amino acid residues are further deleted upstream of the position corresponding to amino acid residue 488 of SEQ ID NO: 3 or 4 or downstream of the position corresponding to amino acid residue 494 of SEQ ID NO: 3 or 4.

[0072] Another group of mutation (mentioned in item (c), supra) resides in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 of the wild-type genomic sequence of P2X7R depicted in SEQ ID NO: 1. Said mutations in said introns are point mutations as shown in Table B hereinabove and in Table 1, hereinbelow.

[0073] At the respective position indicated in the column "Position in wild-type" in Table B or indicated in the column "Polymorphism" in Table 1 the position of the nucleotide residue in the respective intron which is replaced by another nucleotide residue is shown. Accordingly, the term "a nucleotide as indicated in column "Intron" of the Table B corresponding to the position as indicated in column "Replaced nucleotide" of Table B corresponding to the position as indicated in column "Position in wild-type" of Table B is replaced by another nucleotide means that a nucleotide residue in a P2X7R encoding sequence would be located at position Y in SEQ ID NO: 1 when the P2X7R sequence is compared and aligned with the sequence of SEQ ID NO: 1.

[0074] If the nucleotide at the respective position is a purine base such as adenine or guanine it is preferred that due to a transition it is replaced by another purine base. For example, an adenine is replaced by a guanine or a guanine is replaced by an adenine. If the nucleotide at the respective position is a pyrimidine base it is preferred that due to a transition it is replaced by another pyrimidine base. For example, thymine is replaced by a cytidine and a cytidine is replaced by a thymine.

[0075] It is also preferred that due to a transversion a purine base is replaced by a pyrimidine base or vice versa. For example, an adenine is replaced by a thymine and a guanine is replaced by a cytidine. Particularly preferred, said nucleotide in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 of the P2X7R gene depicted in SEQ ID NO: 1 is replaced by the nucleotide depicted in column "Polymorphism" of Table 1, hereinbelow.

[0076] A last group of mutations that has been identified relates to mutations which reside in the 3'UTR of the wild-type P2X7R gene depicted in SEQ ID NO: 1. The mutations were found at positions 54925, 55169, 55170, 55171 or 55917 respectively, of the wild-type P2X7R gene depicted in SEQ ID NO: 1.

[0077] At position 54925 a guanine residue was found to be replaced by another nucleotide. Preferably, said guanine residue is replaced by a pyrimidine base, more preferably by a purine base and particularly preferred by an adenine.

[0078] At position 55169 a cytidine residue is replaced by another nucleotide, preferably by a pyrimidine base. More preferably, it is replaced by a purine base and particularly preferred, it is replaced by an adenine.

[0079] At positions 55170 and 55171 an adenine residue is replaced by another nucleotide residue, preferably by a purine base. More preferably, said adenine residue is replaced by a pyrimidine base and particularly preferred said adenine residue is replaced by a cytidine residue. It was also found that at position 55917 a cytidine residue is replaced by another nucleotide. Preferably, said nucleotide residue is a purine base, more preferably a pyrimidine base and particularly preferred a thymine.

[0080] As is evident from the above, not all identified mutations are located in exons or lead to a change in the amino acid sequence. Some of the mutations are located in the 5'UTR, the 3'UTR or in introns.

[0081] It is known that polymorphisms in promoter and enhancer regions can affect gene function by modulating transcription, particularly if they are situated at recognition sites for DNA binding proteins (Fishman et al., J. Clin. Invest. 102 (1998), 1369-1376). The term "polymorphism" which is used in the present invention means single nucleotide substitution, nucleotide insertion and nucleotide deletion which in the case of insertion and deletion includes insertion or deletion of one or more nucleotides at a position of a gene and corresponding alterations in expressed proteins. Polymorphisms in the 5' untranslated region (5'UTR) of genes can affect the efficiency with which proteins are translated. A representative example of this is in the c-myc gene where a C-G SNP that creates an internal ribosome entry site is associated with increased efficiency of c-myc translation and myeloma (Chappell et al., Oncogene 19 (2000), 4437-4440). Polymorphisms in the 3'UTR can affect gene function by altering the secondary structure of RNA and efficiency of translation or by affecting motifs in the RNA that bind proteins which regulate RNA degradation. Polymorphisms within introns can affect gene function by affecting RNA splicing resulting in aberrant polypeptides. Another way in which intronic polymorphisms can affect gene function is when they affect regulatory motifs within introns. Examples are the Sp1 binding site polymorphism within intron 1 of the COL1A1 gene (Mann et al., J. Clin. Invest. 107 (2001), 899-907) and a repeat polymorphisms within the IL-1Ra gene (Keen et al., Bone 23 (1998), 367-371). Further examples between intronic SNPs and gene function are described in Caceres and Kornblith, Trends Genet. 4 (2002), 186-93. Example 4 on page 52, line 30 to page 53, line 51 of the text describes potential alternative splicing events and aberrant protein production associated with three SNPs disclosed in the application.

[0082] The nucleic acid sequences described hereinabove may comprise at least 56580 nucleotides, preferably at least 10000 nucleotides, at least 5000 nucleotides, at least 1000 nucleotides, at least 500 nucleotides, at least 100 nucleotides. More preferably, said nucleic acid sequences comprise at least 50 nucleotides and particularly preferred they comprise at least 20 or 21 nucleotides comprising the mutations or deletions as described hereinabove. Most preferably such a nucleic acid sequence has a sequence as depicted in any one of SEQ ID NOs: 13 to 51.

[0083] The nucleic acid sequences described hereinabove which comprise mutations in exons leading to a replacement of the corresponding amino acid sequence of the P2X7R wild-type polypeptide depicted in SEQ ID NO: 3 or 4 encode polypeptides shown in SEQ ID NOs: 5 to 10 and 12

[0084] Additionally, the nucleic acid sequences described hereinabove which comprise a deletion leading to a truncated polypeptide in comparison to the full-length polypeptide of the wild-type P2X7R polypeptide shown in SEQ ID NO: 3 or 4 is shown in SEQ ID NO: 11.

[0085] The present invention also relates to nucleic acid molecules which hybridize to one of the above described nucleic acid molecules and which shows a mutation as described hereinabove.

[0086] The term “hybridizes” as used in accordance with the present invention may relate to hybridizations under stringent or non-stringent conditions. If not further specified, the conditions are preferably non-stringent. Said hybridization conditions may be established according to conventional protocols described, for example, in Sambrook, Russell “Molecular Cloning, A Laboratory Manual”, Cold Spring Harbor Laboratory, N.Y. (2001); Ausubel, “Current Protocols in Molecular Biology”, Green Publishing Associates and Wiley Interscience, N.Y. (1989), or Higgins and Hames (Eds.) “Nucleic acid hybridization, a practical approach” IRL Press Oxford, Washington DC, (1985). The setting of conditions is well within the skill of the artisan and can be determined according to protocols described in the art. Thus, the detection of only specifically hybridizing sequences will usually require stringent hybridization and washing conditions such as 0.1×SSC, 0.1% SDS at 65° C. Non-stringent hybridization conditions for the detection of homologous or not exactly complementary sequences may be set at 6×SSC, 1% SDS at 65° C. As is well known, the length of the probe and the composition of the nucleic acid to be determined constitute further parameters of the hybridization conditions. Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt’s reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility. Hybridizing nucleic acid molecules also comprise fragments of the above described molecules. Such fragments may represent nucleic acid sequences which code for a non-functional ATP-gated ion channel P2X7R or a non-functional fragment thereof, and which have a length of at least 12 nucleotides, preferably at least 15, more preferably at least 18, more preferably of at least 21 nucleotides, more preferably at least 30 nucleotides, even more preferably at least 40 nucleotides and most preferably at least 60 nucleotides. Furthermore, nucleic acid molecules which hybridize with any of the aforementioned nucleic acid molecules also include complementary fragments, derivatives and allelic variants of these molecules. Additionally, a hybridization complex refers to a complex between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary G and C bases and between complementary A and T bases; these hydrogen bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization complex may be formed in solution (e.g., Cot or Rot analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., membranes, filters, chips, pins or glass slides to which, e.g.,

cells have been fixed). The terms complementary or complementarity refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, the sequence “A-G-T” binds to the complementary sequence “T-C-A”. Complementarity between two single-stranded molecules may be “partial”, in which only some of the nucleic acids bind, or it may be complete when total complementarity exists between single-stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands.

[0087] The term “hybridizing sequences” preferably refers to sequences which display a sequence identity of at least 40%, preferably at least 50%, more preferably at least 60%, even more preferably at least 70%, particularly preferred at least 80%, more particularly preferred at least 90%, even more particularly preferred at least 95% and most preferably at least 97% identity with a nucleic acid sequence as described above encoding a P2X7R protein having a described mutation. Moreover, the term “hybridizing sequences” preferably refers to sequences encoding a P2X7R protein having a sequence identity of at least 40%, preferably at least 50%, more preferably at least 60%, even more preferably at least 70%, particularly preferred at least 80%, more particularly preferred at least 90%, even more particularly preferred at least 95% and most preferably at least 97% identity with an amino acid sequence of a P2X7R mutant as described herein above.

[0088] In accordance with the present invention, the term “identical” or “percent identity” in the context of two or more nucleic acid or amino acid sequences, refers to two or more sequences or subsequences that are the same, or that have a specified percentage of amino acid residues or nucleotides that are the same (e.g., 60% or 65% identity, preferably, 70-95% identity, more preferably at least 95% identity), when compared and aligned for maximum correspondence over a window of comparison, or over a designated region as measured using a sequence comparison algorithm as known in the art, or by manual alignment and visual inspection. Sequences having, for example, 60% to 95% or greater sequence identity are considered to be substantially identical. Such a definition also applies to the complement of a test sequence. Preferably the described identity exists over a region that is at least about 15 to 25 amino acids or nucleotides in length, more preferably, over a region that is about 50 to 100 amino acids or nucleotides in length. Those having skill in the art will know how to determine percent identity between/among sequences using, for example, algorithms such as those based on CLUSTALW computer program (Thompson Nucl. Acids Res. 2 (1994), 4673-4680) or FASTDB (Brutlag Comp. App. Biosci. 6 (1990), 237-245), as known in the art.

[0089] Although the FASTDB algorithm typically does not consider internal non-matching deletions or additions in sequences, i.e., gaps, in its calculation, this can be corrected manually to avoid an overestimation of the % identity. CLUSTALW, however, does take sequence gaps into account in its identity calculations. Also available to those having skill in this art are the BLAST and BLAST 2.0 algorithms (Altschul Nucl. Acids Res. 25 (1977), 3389-

3402). The BLASTN program for nucleic acid sequences uses as defaults a word length (W) of 11, an expectation (E) of 10, M=5, N=4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, and an expectation (E) of 10. The BLOSUM62 scoring matrix (Henikoff Proc. Natl. Acad. Sci., USA, 89, (1989), 10915) uses alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

**[0090]** Moreover, the present invention also relates to nucleic acid molecules the sequence of which is degenerate in comparison with the sequence of an above-described hybridizing molecule. When used in accordance with the present invention the term "being degenerate as a result of the genetic code" means that due to the redundancy of the genetic code different nucleotide sequences code for the same amino acid.

**[0091]** The present invention also related to nucleic acid molecules which comprise one or more of the above-described mutations or deletions.

**[0092]** The nucleic acid molecules according to the invention may be derived from any organism encoding corresponding P2X7R ATP-gated ion channels. For example, P2X7R ATP-gated ion channels have been reported in various organisms, for example, rat (see, Suprenant (1996), loc. cit.), mouse (Genbank Accession No. AJ 489297), xenopus (Genbank Accession No. AJ 345114), chicken (Genbank Accession No. BM 491404) or Bos Taurus (Genbank Accession No. AF 083073). In a preferred embodiment the nucleic acid molecule of the invention is derived from a vertebrate, preferably from a mammal, even more preferably the nucleic acid molecule is derived from rabbit or guinea pig, and most preferably the nucleic acid is derived from mouse, rat or human.

**[0093]** The nucleic acid molecule according to the invention may be any type of nucleic acid, e.g. DNA, RNA or PNA (peptide nucleic acid).

**[0094]** For the purposes of the present invention, a peptide nucleic acid (PNA) is a polyamide type of DNA analog and the monomeric units for adenine, guanine, thymine and cytosine are available commercially (Perceptive Biosystems). Certain components of DNA, such as phosphorus, phosphorus oxides, or deoxyribose derivatives, are not present in PNAs. As disclosed by Nielsen et al., Science 254:1497 (1991); and Egholm et al., Nature 365:666 (1993), PNAs bind specifically and tightly to complementary DNA strands and are not degraded by nucleases. In fact, PNA binds more strongly to DNA than DNA itself does. This is probably because there is no electrostatic repulsion between the two strands, and also the polyamide backbone is more flexible. Because of this, PNA/DNA duplexes bind under a wider range of stringency conditions than DNA/DNA duplexes, making it easier to perform multiplex hybridization. Smaller probes can be used than with DNA due to the strong binding. In addition, it is more likely that single base mismatches can be determined with PNA/DNA hybridization because a single mismatch in a PNA/DNA 15-mer lowers the melting point (T<sub>sub.m</sub>) by 8°-20° C., vs. 4°-16° C. for the DNA/DNA 15-mer duplex. Also, the absence of charge groups in PNA means that hybridization can be done at low ionic strengths and reduce possible interference by salt during the analysis.

**[0095]** The DNA may, for example, be cDNA. In a preferred embodiment it is a genomic DNA. The RNA may be, e.g., mRNA. The nucleic acid molecule may be natural, synthetic or semisynthetic or it may be a derivative, such as peptide nucleic acid (Nielsen, Science 254 (1991), 1497-1500) or phosphorothioates. Furthermore, the nucleic acid molecule may be a recombinantly produced chimeric nucleic acid molecule comprising any of the aforementioned nucleic acid molecules either alone or in combination.

**[0096]** Preferably, the nucleic acid molecule of the present invention is part of a vector. Therefore, the present invention relates in another embodiment to a vector comprising the nucleic acid molecule of this invention. Such a vector may be, e.g., a plasmid, cosmid, virus, bacteriophage or another vector used e.g. conventionally in genetic engineering, and may comprise further genes such as marker genes which allow for the selection of said vector in a suitable host cell and under suitable conditions.

**[0097]** The nucleic acid molecules of the present invention may be inserted into several commercially available vectors. Nonlimiting examples include plasmid vectors compatible with mammalian cells, such as pUC, pBluescript (Stratagene), pET (Novagen), pREP (Invitrogen), pCRTopo (Invitrogen), pcDNA3 (Invitrogen), pCEP4 (Invitrogen), pMC1 neo (Stratagene), pXT1 (Stratagene), pSG5 (Stratagene), EBO-pSV2neo, pBPV-1, pDBPVM Tneo, pRS-Vgpt, pRSVneo, pSV2-dhfr, pUCtag, pLZD35, PLXIN and PSIR (Clontech) and pIRES-EGFP (Clontech). Baculovirus vectors such as pBlueBac, BacPacZ Baculovirus Expression System (CLONTECH), and MaxBac™ Baculovirus Expression System, insect cells and protocols (Invitrogen) are available commercially and may also be used to produce high yields of biologically active protein. (see also, Miller (1993), Curr. Op. Genet. Dev., 3, 9; O'Reilly, Baculovirus Expression Vectors: A Laboratory Manual, p. 127). In addition, prokaryotic vectors such as pcDNA2; and yeast vectors such as pYes2 are nonlimiting examples of other vectors suitable for use with the present invention. For vector modification techniques, see Sambrook and Russel (2001), loc. cit. Vectors can contain one or more replication and inheritance systems for cloning or expression, one or more markers for selection in the host, e. g., antibiotic resistance, and one or more expression cassettes.

**[0098]** The coding sequences inserted in the vector can be synthesized by standard methods, isolated from natural sources, or prepared as hybrids. Ligation of the coding sequences to transcriptional regulatory elements (e. g., promoters, enhancers, and/or insulators) and/or to other amino acid encoding sequences can be carried out using established methods.

**[0099]** Furthermore, the vectors may, in addition to the nucleic acid sequences of the invention, comprise expression control elements, allowing proper expression of the coding regions in suitable hosts. Such control elements are known to the artisan and may include a promoter, translation initiation codon, translation and insertion site or internal ribosomal entry sites (IRES) (Owens, Proc. Natl. Acad. Sci. USA 98 (2001), 1471-1476) for introducing an insert into the vector. Preferably, the nucleic acid molecule of the invention is operatively linked to said expression control sequences allowing expression in eukaryotic or prokaryotic cells. Particularly preferred are in this context control



sequences which allow for correct expression in neuronal cells and/or cells derived from nervous tissue.

**[0100]** Control elements ensuring expression in eukaryotic and prokaryotic cells are well known to those skilled in the art. As mentioned above, they usually comprise regulatory sequences ensuring initiation of transcription and optionally poly-A signals ensuring termination of transcription and stabilization of the transcript. Additional regulatory elements may include transcriptional as well as translational enhancers, and/or naturally-associated or heterologous promoter regions. Possible regulatory elements permitting expression in for example mammalian host cells comprise the CMV-HSV thymidine kinase promoter, SV40, RSV-promoter (Rous sarcoma virus), human elongation factor 1 $\alpha$ -promoter, CMV enhancer, CaM-kinase promoter or SV40-enhancer.

**[0101]** For the expression for example in nervous tissue and/or cells derived therefrom, several regulatory sequences are well known in the art, like the minimal promoter sequence of human neurofilament L (Charron, J. Biol. Chem. 270 (1995), 25739-25745). For the expression in prokaryotic cells, a multitude of promoters including, for example, the tac-lac-promoter, the lacUV5 or the trp promoter, has been described. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pRc/CMV, pcDNA1, pcDNA3 (In-VitroGene, as used, inter alia in the appended examples), pSPORT1 (GIBCO BRL) or pGEMHE (Promega), or prokaryotic expression vectors, such as lambda gt11.

**[0102]** An expression vector according to this invention is at least capable of directing the replication, and preferably the expression, of the nucleic acids and protein of this invention. Suitable origins of replication include, for example, the Col E1, the SV40 viral and the M 13 origins of replication. Suitable promoters include, for example, the cytomegalovirus (CMV) promoter, the iacZ promoter, the gai10 promoter and the Autographa californica multiple nuclear polyhedrosis virus (AcMNPV) polyhedral promoter. Suitable termination sequences include, for example, the bovine growth hormone, SV40, iacZ and AcMNPV polyhedral polyadenylation signals. Examples of selectable markers include neomycin, ampicillin, and hygromycin resistance and the like. Specifically-designed vectors allow the shuttling of DNA between different host cells, such as bacteria-yeast, or bacteria-animal cells, or bacteria-fungal cells, or bacteria invertebrate cells.

**[0103]** Beside the nucleic acid molecules of the present invention, the vector may further comprise nucleic acid sequences encoding for secretion signals. Such sequences are well known to the person skilled in the art. Furthermore, depending on the expression system used leader sequences capable of directing the expressed polypeptide to a cellular compartment may be added to the coding sequence of the nucleic acid molecules of the invention and are well known in the art. The leader sequence(s) is (are) assembled in appropriate phase with translation, initiation and termination sequences, and preferably, a leader sequence capable of

directing secretion of translated protein, or a part thereof, into, inter alia, the extracellular membrane. Optionally, the heterologous sequence can encode a fusion protein including an C— or N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product. Once the vector has been incorporated into the appropriate host, the host is maintained under conditions suitable for high level expression of the nucleotide sequences, and, as desired, the collection and purification of the proteins, antigenic fragments or fusion proteins of the invention may follow. Of course, the vector can also comprise regulatory regions from pathogenic organisms.

**[0104]** Furthermore, said vector may also be, besides an expression vector, a gene transfer and/or gene targeting vector. Gene therapy, which is based on introducing therapeutic genes (for example for vaccination) into cells by ex-vivo or in-vivo techniques is one of the most important applications of gene transfer. Suitable vectors, vector systems and methods for in-vitro or in-vivo gene therapy are described in the literature and are known to the person skilled in the art; see, e.g., Giordano, Nature Medicine 2 (1996), 534-539; Schaper, Circ. Res. 79 (1996), 911-919; Anderson, Science 256 (1992), 808-813; Isner, Lancet 348 (1996), 370-374; Muhlhauser, Circ. Res. 77 (1995), 1077-1086; Wang, Nature Medicine 2 (1996), 714-716; WO 94/29469; WO 97/00957; Schaper, Current Opinion in Biotechnology 7 (1996), 635-640 or Verma, Nature 389 (1997), 239-242 and references cited therein. The nucleic acid molecules of the invention and vectors as described herein above may be designed for direct introduction or for introduction via liposomes, or viral vectors (e.g. adenoviral, retroviral) into the cell. Additionally, baculoviral systems or systems based on vaccinia virus or Semliki Forest Virus can be used as eukaryotic expression system for the nucleic acid molecules of the invention. In addition to recombinant production, fragments of the protein, the fusion protein or antigenic fragments of the invention may be produced by direct peptide synthesis using solid-phase techniques (cf Stewart et al. (1969) Solid Phase Peptide Synthesis; Freeman Co, San Francisco; Merrifield, J. Am. Chem. Soc. 85 (1963), 2149-2154). In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer, Foster City Calif.) in accordance with the instructions provided by the manufacturer. Various fragments may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

**[0105]** The present invention in addition relates to a host transformed with a vector of the present invention or to a host comprising the nucleic acid molecule of the invention. Said host may be produced by introducing said vector or nucleotide sequence into a host cell which upon its presence in the cell mediates the expression of a protein encoded by the nucleotide sequence of the invention or comprising a nucleotide sequence or a vector according to the invention wherein the nucleotide sequence and/or the encoded polypeptide is foreign to the host cell.

**[0106]** By "foreign" it is meant that the nucleotide sequence and/or the encoded polypeptide is either heterologous with respect to the host, this means derived from a cell or organism with a different genomic background, or is

homologous with respect to the host but located in a different genomic environment than the naturally occurring counterpart of said nucleotide sequence. This means that, if the nucleotide sequence is homologous with respect to the host, it is not located in its natural location in the genome of said host, in particular it is surrounded by different genes. In this case the nucleotide sequence may be either under the control of its own promoter or under the control of a heterologous promoter. The location of the introduced nucleic acid molecule or the vector can be determined by the skilled person by using methods well-known to the person skilled in the art, e.g., Southern Blotting. The vector or nucleotide sequence according to the invention which is present in the host may either be integrated into the genome of the host or it may be maintained in some form extrachromosomally. In this respect, it is also to be understood that the nucleotide sequence of the invention can be used to restore or create a mutant gene via homologous recombination.

**[0107]** Said host may be any prokaryotic or eukaryotic cell. Suitable prokaryotic/bacterial cells are those generally used for cloning like *E. coli*, *Salmonella typhimurium*, *Serratia marcescens* or *Bacillus subtilis*. Said eukaryotic host may be a mammalian cell, an amphibian cell, a fish cell, an insect cell, a fungal cell, a plant cell or a bacterial cell (e.g., *E. coli* strains HB101, DH5a, XLI Blue, Y1090 and JM101). Eukaryotic recombinant host cells are preferred. Examples of eukaryotic host cells include, but are not limited to, yeast, e.g., *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis* or *Pichia pastoris* cells, cell lines of human, bovine, porcine, monkey, and rodent origin, as well as insect cells, including but not limited to, *Spodoptera frugiperda* insect cells and *Drosophila*-derived insect cells as well as zebra fish cells. Mammalian species-derived cell lines suitable for use and commercially available include, but are not limited to, L cells, CV-1 cells, COS-1 cells (ATCC CRL 1650), COS-7 cells (ATCC CRL 1651), HeLa cells (ATCC CCL 2), C1271 (ATCC CRL 1616), BS-C-1 (ATCC CCL 26) and MRC-5 (ATCC CCL 171).

**[0108]** In a particularly preferred embodiment said mammalian cell is a neuronal cell and/or a cultured cell like, inter alia, a HEK 293 (human embryonic kidney) cell, a CHO, HeLa, NIH3T3, BHK, PC12 cell or a neuronal stem cell preferably derived from a mammal and more preferably from a human.

**[0109]** In another more preferred embodiment said amphibian cell is an oocyte. In an even more preferred embodiment said oocyte is a frog oocyte, particularly preferred a *Xenopus laevis* oocyte.

**[0110]** In a more preferred embodiment, the host according to the invention is a non-human transgenic organism. Said non-human organism may be a mammal, amphibian, a fish, an insect, a fungus or a plant. Particularly preferred non-human transgenic animals are *Drosophila* species, *Caenorhabditis elegans*, *Xenopus* species, zebra fish, *Spodoptera frugiperda*, *Autographa californica*, mice and rats. Transgenic plants comprise, but are not limited to, wheat, tobacco, parsley and *Arabidopsis*. Transgenic fungi are also well known in the art and comprise, inter alia, yeasts, like *S. pombe* or *S. cerevisiae*, or *Aspergillus*, *Neurospora* or *Ustilago* species or *Pichia* species.

**[0111]** In another embodiment, the present invention relates to a method for producing the polypeptide encoded

by a nucleic acid molecule of the invention comprising culturing/raising the host of the invention and isolating the produced polypeptide. A large number of suitable methods exist in the art to produce polypeptides in appropriate hosts. If the host is a unicellular organism or a mammalian or insect cell, the person skilled in the art can revert to a variety of culture conditions that can be further optimized without an undue burden of work. Conveniently, the produced protein is harvested from the culture medium or from isolated (biological) membranes by established techniques. Furthermore, the produced polypeptide may be directly isolated from the host cell. Said host cell may be part of or derived from a part of a host organism, for example said host cell may be part of the CNS of an animal or the harvestable part of a plant. Additionally, the produced polypeptide may be isolated from fluids derived from said host, like blood, milk or cerebrospinal fluid.

**[0112]** Additionally the present invention relates to polypeptides depicted in SEQ ID NOs: 5 to 12 which are encoded by the nucleic acid molecules of the invention or produced by the method of the invention. The polypeptide of the invention may accordingly be produced by microbiological methods or by transgenic mammals. It is also envisaged that the polypeptide of the invention is recovered from transgenic plants. Alternatively, the polypeptide of the invention may be produced synthetically or semi-synthetically.

**[0113]** For example, chemical synthesis, such as the solid phase procedure described by Houghton Proc. Natl. Acad. Sci. USA (82) (1985), 5131-5135, can be used. Another method is in vitro translation of mRNA. A preferred method involves the recombinant production of protein in host cells as described above. For example, nucleotide acid sequences comprising all or a portion of any one of the nucleotide sequences according to the invention can be synthesized by PCR, inserted into an expression vector, and a host cell transformed with the expression vector. Thereafter, the host cell is cultured to produce the desired polypeptide, which is isolated and purified. Protein isolation and purification can be achieved by any one of several known techniques; for example and without limitation, ion exchange chromatography, gel filtration chromatography and affinity chromatography, high pressure liquid chromatography (HPLC), reversed phase HPLC, preparative disc gel electrophoresis. In addition, cell-free translation systems can be used to produce the polypeptides of the present invention. Suitable cell-free expression systems for use in accordance with the present invention include rabbit reticulocyte lysate, wheat germ extract, canine pancreatic microsomal membranes, *E. coli* S30 extract, and coupled transcription/translation systems such as the TNT-system (Promega). These systems allow the expression of recombinant polypeptides or peptides upon the addition of cloning vectors, DNA fragments, or RNA sequences containing coding regions and appropriate promoter elements. As mentioned supra, protein isolation/purification techniques may require modification of the proteins of the present invention using conventional methods. For example, a histidine tag can be added to the protein to allow purification on a nickel column. Other modifications may cause higher or lower activity, permit higher levels of protein production, or simplify purification of the protein.

[0114] In a further embodiment, the present invention relates to an antibody specifically directed to a polypeptide of the invention, wherein said antibody specifically reacts with an epitope generated and/or formed by the mutation in the ATP-gated ion channel P2X7R selected from the group consisting of:

[0115] (i) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein the R (Arg), G (Gly), E (Glu), L (Leu), R (Arg), I (Ile) or R (Arg) residue corresponding to position 117, 150, 186, 191, 270, 568 or 578 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 is replaced by another amino acid residue; and

[0116] (ii) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted. With respect to preferred embodiments of (i) and (ii) the same applies as described above in connection with the nucleic acid molecules. The term “specifically” in this context means that the antibody reacts with the mutant P2X7R protein but not with a wild-type P2X7R protein. Preferably this term also means that such an antibody does not bind to other mutant forms of the P2X7R protein, in particular those described herein. Whether the antibody specifically reacts as defined herein above can easily be tested, *inter alia*, by comparing the reaction of said antibody with a wild-type ATP-gated ion channel P2X7R (or a sub-unit or a fragment thereof) with the reaction of said antibody with a mutant P2X7R polypeptide of the invention.

[0117] The antibody of the present invention can be, for example, polyclonal or monoclonal. The term “antibody” also comprises derivatives or fragments thereof which still retain the binding specificity. Techniques for the production of antibodies are well known in the art and described, e.g. in Harlow and Lane “Antibodies, A Laboratory Manual”, CSH Press, Cold Spring Harbor, 1988. These antibodies can be used, for example, for the immunoprecipitation and immunolocalization of the polypeptides of the invention as well as for the monitoring of the presence of such polypeptides, for example, in recombinant organisms or in diagnosis. They can also be used for the identification of compounds interacting with the proteins according to the invention (as mentioned herein below). For example, surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of the polypeptide of the invention (Schier, *Human Antibodies Hybridomas* 7 (1996), 97-105; Malmberg, *J. Immunol. Methods* 183 (1995), 7-13).

[0118] The present invention furthermore includes chimeric, single chain and humanized antibodies, as well as antibody fragments, like, *inter alia*, Fab fragments. Antibody fragments or derivatives further comprise F(ab')<sub>2</sub>, Fv or scFv fragments; see, for example, Harlow and Lane, *loc. cit.* Various procedures are known in the art and may be used for the production of such antibodies and/or fragments. Thus, the (antibody) derivatives can be produced by peptidomimetics.

Further, techniques described for the production of single chain antibodies (see, *inter alia*, U.S. Pat. No. 4,946,778) can be adapted to produce single chain antibodies to polypeptide(s) of this invention. Also, transgenic animals may be used to express humanized antibodies to polypeptides of this invention. Most preferably, the antibody of this invention is a monoclonal antibody. For the preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples for such techniques include the hybridoma technique (Köhler and Milstein *Nature* 256 (1975), 495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor, *Immunology Today* 4 (1983), 72) and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. (1985), 77-96). Techniques describing the production of single chain antibodies (e.g., U.S. Pat. No. 4,946,778) can be adapted to produce single chain antibodies to immunogenic polypeptides as described above. Furthermore, transgenic mice may be used to express humanized antibodies directed against said immunogenic polypeptides. It is in particular preferred that the antibodies/antibody constructs as well as antibody fragments or derivatives to be employed in accordance with this invention or capable to be expressed in a cell. This may, *inter alia*, be achieved by direct injection of the corresponding proteinaceous molecules or by injection of nucleic acid molecules encoding the same. Furthermore, gene therapy approaches are envisaged. Accordingly, in context of the present invention, the term “antibody molecule” relates to full immunoglobulin molecules as well as to parts of such immunoglobulin molecules. Furthermore, the term relates, as discussed above, to modified and/or altered antibody molecules, like chimeric and humanized antibodies. The term also relates to monoclonal or polyclonal antibodies as well as to recombinantly or synthetically generated/synthesized antibodies. The term also relates to intact antibodies as well as to antibody fragments thereof, like, separated light and heavy chains, Fab, Fab/c, Fv, Fab', F(ab')<sub>2</sub>. The term “antibody molecule” also comprises bifunctional antibodies and antibody constructs, like single chain Fvs (scFv) or antibody-fusion proteins. It is also envisaged in context of this invention that the term “antibody” comprises antibody constructs which may be expressed in cells, e.g. antibody constructs which may be transfected and/or transduced via, *inter alia*, viruses or vectors. It is in particular envisaged that such antibody constructs specifically recognize the polypeptides of the present invention. It is, furthermore, envisaged that said antibody construct is employed in gene therapy approaches.

[0119] The present invention relates also to an aptamer specifically binding to a polypeptide according to the invention wherein said aptamer reacts with an epitope of a polypeptide of the present invention. The present invention furthermore relates to an aptamer specifically directed to a corresponding nucleic acid molecule according to the invention.

[0120] In accordance with the present invention, the term “aptamer” means nucleic acid molecules that can bind to target molecules. Aptamers commonly comprise RNA, single stranded DNA, modified RNA or modified DNA molecules. The preparation of aptamers is well known in the

art and may involve, inter alia, the use of combinatorial RNA libraries to identify binding sides (Gold, *Ann. Rev. Biochem.* 64 (1995), 763-797).

[0121] Furthermore, the present invention relates to a primer or pair of primers capable of specifically amplifying the nucleic acid molecules of the present invention. The term "primer" when used in the present invention means a single-stranded nucleic acid molecule capable of annealing the nucleic acid molecule of the present application and thereby being capable of serving as a starting point for amplification. Said term also comprises oligoribo- or desoxyribonucleotides which are complementary to a region of one of the strands of a nucleic acid molecule of the present invention. According to the present invention the term "pair of primers" means a pair of primers that are with respect to a complementary region of a nucleic acid molecule directed in the opposite direction towards each other to enable, for example, amplification by polymerase chain reaction (PCR).

[0122] The term "amplifying" refers to repeated copying of a specified sequence of nucleotides resulting in an increase in the amount of said specified sequence of nucleotides and allows the generation of a multitude of identical or essentially identical (i.e. at least 95% more preferred at least 98%, even more preferred at least 99% and most preferred at least 99.5% such as 99.9% identical) nucleic acid molecules or parts thereof. Such methods are well established in the art; see Sambrook et al. "Molecular Cloning, A Laboratory Manual", 2<sup>nd</sup> edition 1989, CSH Press, Cold Spring Harbor. They include polymerase chain reaction (PCR) and modifications thereof, ligase chain reaction (LCR) to name some preferred amplification methods.

[0123] When used in the context of primers the term "specifically" means that only the nucleic acid molecules as described herein above are amplified and nucleic acid molecules encoding the wild-type P2X7R ATP-gated receptor as depicted in SEQ ID NO: 1 are not amplified. Thus, a primer according to the invention is preferably a primer which binds to a region of a nucleic acid molecule of the invention which is unique for this molecule and which is not present in the wild-type P2X7R encoding sequence, i.e. the primer binds in a region in which one of the above described mutations occur. In connection with a pair of primers according to the invention it is possible that one of the primers of the pair is specific in the above described meaning or both of the primers of the pair are specific. In both cases, the use of such a pair of primers would allow to specifically amplify a mutant of the invention as described herein-above but not the wild-type P2X7R encoding sequence.

[0124] The 3'-OH end of a primer is used by a polymerase to be extended by successive incorporation of nucleotides. The primer or pair of primers of the present invention can be used, for example, in primer extension experiments on template RNA according to methods known by the person skilled in the art. Preferably, the primer or pair of primers of the present invention are used for amplification reactions on template RNA or template DNA, preferably cDNA or genomic DNA. The terms "template DNA" or "template RNA" refers to DNA or RNA molecules or fragments thereof of any source or nucleotide composition, that comprise a target nucleotide sequence as defined above. The primer or pair of primers can also be used for hybridization

experiments as known in the art. Preferably, the primer or pair of primers are used in polymerase chain reactions to amplify sequences corresponding to a sequence of the nucleic acid molecule of the present invention. It is known that the length of a primer results from different parameters (Gillam, *Gene* 8 (1979), 81-97; Innis, *PCR Protocols: A guide to methods and applications*, Academic Press, San Diego, USA (1990)). Preferably, the primer should only hybridize or bind to a specific region of a target nucleotide sequence. The length of a primer that statistically hybridizes only to one region of a target nucleotide sequence can be calculated by the following formula:  $(1/4)^x$  (whereby x is the length of the primer). For example a hepta- or octanucleotide would be sufficient to bind statistically only once on a sequence of 37 kb. However, it is known that a primer exactly matching to a complementary template strand must be at least 9 base pairs in length, otherwise no stable-double strand can be generated (Goulian, *Biochemistry* 12 (1973), 2893-2901). It is also envisaged that computer-based algorithms can be used to design primers capable of amplifying the nucleic acid molecules of the invention. Preferably, the primers of the invention are at least 10 nucleotides in length, more preferred at least 12 nucleotides in length, even more preferred at least 15 nucleotides in length, particularly preferred at least 18 nucleotides in length, even more particularly preferred at least 20 nucleotides in length and most preferably at least 25 nucleotides in length. The invention, however, can also be carried out with primers which are shorter or longer.

[0125] It is also envisaged that the primer or pair of primers is labeled. The label may, for example, be a radioactive label, such as  $^{32}\text{P}$ ,  $^{33}\text{P}$  or  $^{35}\text{S}$ . In a preferred embodiment of the invention, the label is a non-radioactive label, for example, digoxigenin, biotin and fluorescence dye or a dye.

[0126] In another preferred embodiment said primers are selected from the group consisting of SEQ ID NOs: 52 to 111.

[0127] In yet another embodiment, the present invention relates to a composition comprising a nucleic acid molecule, a vector, a polypeptide, an antibody, an aptamer and/or a primer or pair of primers of the invention.

[0128] The term "composition", as used in accordance with the present invention, relates to compositions which comprise at least one nucleic acid molecule, vector, polypeptide, an antibody and/or primer or pair of primers of this invention. It may, optionally, comprise further molecules capable of altering the characteristics of the component of the invention thereby, for example, suppressing, blocking, modulating and/or activating their function which have neuroprotective, nootropic, antidepressive and/or cell-protective properties as will also be described herein below. The composition may be in solid, liquid or gaseous form and may be, inter alia, in the form of (a) powder(s), (a) tablet(s), (a) solution(s) or (an) aerosol(s).

[0129] In a preferred embodiment the composition according to the invention is a diagnostic composition, optionally further comprising suitable means for detection. As described above, the present invention is based on the surprising finding that mutations in the P2X7R protein are connected with affective disorders. Thus, this knowledge now allows to diagnose affective disorders in an easy way. The diagnostic composition comprises at least one of the

aforementioned compounds of the invention. The diagnostic composition may be used, *inter alia*, for methods for determining the presence and/or expression of the nucleic acids and/or polypeptides of the invention. This may be effected by detecting, e.g., the presence of a corresponding gene in the genetic material of an individual or the presence of the corresponding mRNA which comprises isolation of DNA or RNA from a cell derived from said individual, contacting the DNA or RNA so obtained with a nucleic acid probe as described above under hybridizing conditions, and detecting the presence of mRNAs hybridized to the probe. Alternatively, the diagnostic composition may also be used for detecting the presence of a nucleic acid molecule of the invention by PCR. Furthermore, polypeptides of the invention can be detected with methods known in the art, which comprise, *inter alia*, immunological methods, like, RIA, FIA, ELISA, FACS or Western blotting.

[0130] Furthermore, the diagnostic composition of the invention may be useful, *inter alia*, in detecting the prevalence, the onset or the progress of a disease related to the expression of a polypeptide of the invention. Accordingly, the diagnostic composition of the invention may be used, *inter alia*, for assessing the prevalence, the onset and/or the disease status of affective disorders, as defined herein above. It is also contemplated that the diagnostic composition of the invention may be useful in discriminating (the) stage(s) of a disease.

[0131] The diagnostic composition optionally comprises suitable means for detection. The nucleic acid molecule(s), vector(s), host(s), antibody(ies), aptamer(s), polypeptide(s) described above are, for example, suitable for use in immunoassays in which they can be utilized in liquid phase or bound to a solid phase carrier. Examples of well-known carriers include glass, polystyrene, polyvinyl ion, polypropylene, polyethylene, polycarbonate, dextran, nylon, amyloses, natural and modified celluloses, polyacrylamides, agaroses, and magnetite. The nature of the carrier can be either soluble or insoluble for the purposes of the invention.

[0132] Solid phase carriers are known to those in the art and may comprise polystyrene beads, latex beads, magnetic beads, colloid metal particles, glass and/or silicon chips and surfaces, nitrocellulose strips, membranes, sheets, duracytes and the walls of wells of a reaction tray, plastic tubes or other test tubes. Suitable methods of immobilizing nucleic acid molecule(s), vector(s), host(s), antibody(ies), aptamer(s), polypeptide(s), etc. on solid phases include but are not limited to ionic, hydrophobic, covalent interactions or (chemical) crosslinking and the like. Examples of immunoassays which can utilize said compounds of the invention are competitive and non-competitive immunoassays in either a direct or indirect format. Commonly used detection assays can comprise radioisotopic or non-radioisotopic methods. Examples of such immunoassays are the radioimmunoassay (RIA), the sandwich (immunometric assay) and the Northern or Southern blot assay. Furthermore, these detection methods comprise, *inter alia*, IRMA (Immune Radioimmunometric Assay), EIA (Enzyme Immuno Assay), ELISA (Enzyme Linked Immuno Assay), FIA (Fluorescent Immuno Assay), and CLIA (Chemiluminescent Immune Assay). Furthermore, the diagnostic compounds of the present invention may be employed in techniques like FRET (Fluorescence Resonance Energy Transfer) assays.

[0133] Appropriate labels and methods for labeling are known to those of ordinary skill in the art. Examples of the types of labels which can be used in the present invention include *inter alia*, fluorochromes (like fluorescein, rhodamine, Texas Red, etc.), enzymes (like horse radish peroxidase,  $\beta$ -galactosidase, alkaline phosphatase), radioactive isotopes (like  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^{35}\text{S}$  or  $^{125}\text{I}$ ), biotin, digoxigenin, colloidal metals, chemi- or bioluminescent compounds (like dioxetanes, luminol or acridiniums).

[0134] A variety of techniques are available for labeling biomolecules, are well known to the person skilled in the art and are considered to be within the scope of the present invention and comprise, *inter alia*, covalent coupling of enzymes or biotinyl groups, phosphorylations, biotinylations, random priming, nick-translations, tailing (using terminal transferases). Such techniques are, e.g., described in Tiossen, "Practice and theory of enzyme immunoassays", Burden and von Knippenburg (Eds), Volume 15 (1985), "Basic methods in molecular biology", Davis L G, Dabmer M D, Battey Elsevier (1990); Mayer, (Eds) "Immunochemical methods in cell and molecular biology" Academic Press, London (1987); or in the series "Methods in Enzymology", Academic Press, Inc.

[0135] Detection methods comprise, but are not limited to, autoradiography, fluorescence microscopy, direct and indirect enzymatic reactions, etc.

[0136] Said diagnostic composition may be used for methods for detecting the presence and/or abundance of a nucleic acid molecule of the invention in a biological and/or medical sample and/or for detecting expression of such a nucleic acid molecule (e.g. by determining the mRNA or the expressed polypeptide). Furthermore, said diagnostic composition may also be used in methods of the present invention, *inter alia*, for the detection of specific antagonists or agonists for P2X7R ATP-gated ion channels (see herein below).

[0137] In a further embodiment the present invention provides a method of diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2X7R protein expressed in the cells of said individual is non-functional, shows an altered ATP-gating in comparison to the wild-type P2X7R protein or is over- or under-expressed in comparison to the P2X7R protein level of an unaffected individual.

[0138] The term "over- or under-expressed in comparison to the P2X7R protein level" in the context of the present invention means that the P2X7R protein level is higher or lower than the P2X7R level of a healthy individual, i.e. an individual not affected with an affective disorder. The over-expression may result, e.g. from an increased amount of P2X7R mRNA caused by enhanced transcription rates due to increased activity of the RNA-polymerase II. The amount of mRNA may accordingly lead to an increased translation and, thus, to a higher protein level of P2X7R. It may also be possible that a higher amount of P2X7R protein is caused by increased stability of the protein. An under-expression of P2X7R protein may be caused by low transcription rates of the P2X7R gene and, thus, insufficient amounts of P2X7R mRNA give only rise to a low P2X7R protein amount. Another reason may be that the P2X7R protein is unstable and, thus, is not present in amounts comparable to the wild-type protein level.

[0139] The under- or over-expression of P2X7R protein may be determined by methods well-known to the person skilled in the art. These include, but are not limited, to methods for determining the amount of mRNA or the amount and/or activity of the protein. Examples are Northern Blot analysis or immuno based techniques, such as Western Blotting.

[0140] “Non-functional” means that the P2X7R protein has lost at least one functional property displayed by the wild-type P2X7R protein as described herein above. Preferably, “non-functional” means that the P2X7R protein does no longer function as a channel. Non-functionality may, e.g., be caused by the fact that one allele occurring in an individual codes for a P2X7R protein which leads to non-functional dimers (dominant negative mutation). Whether a P2X7R protein in an individual is functional or non-functional can be determined by the methods described herein above and in the examples.

[0141] The term “altered ATP-gating” means that the respective P2X7R protein reacts in a different way to ATP than the wild-type P2X7R protein. This can be determined as described in the appended examples or as described hereinabove.

[0142] In the context of diagnosis, not only the activity of the P2X7R could be of diagnostic value but also the amount of expression. For example, if a polymorphism affects RNA stability or translation efficiency, this could lead to lower expression of the P2X7 protein not only in the hippocampus but also in the blood. Therefore, one could speculate that a lower amount of P2X7 detected by western blot in blood cells could be related to depression.

[0143] Another aspect of the present invention is a method for diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2X7R gene sequence or encoded protein thereof comprises a mutation in comparison to the wild-type P2X7R sequence.

[0144] A preferred embodiment of the present invention is a method, wherein a mutation is a mutation in a P2X7R sequence as defined hereinabove and/or a nucleotide replacement or deletion selected from the following Table C indicating in column “Region of P2X7R” the region of the P2X7R genomic nucleotide sequence in which the replacement or deletion occurs, in column “Nucleotide” of Table C the nucleotide which is replaced by another nucleotide or the nucleotides which are deleted and in column “Position in wild-type” of Table C the corresponding position in the nucleotide sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1

TABLE C

Region of P2X7R	Nucleotide	Position in wild-type
5'UTR	T	362
5'UTR	T	532
5'UTR	A	1100
5'UTR	A	1122
5'UTR	C	1171
5'UTR	T	1351
5'UTR	G	1702
5'UTR	T	1731

TABLE C-continued

Region of P2X7R	Nucleotide	Position in wild-type
5'UTR	C	1860
5'UTR	C	2162
5'UTR	C	2238
5'UTR	A	2373
5'UTR	G	2569
5'UTR	G	2702
intron 1	G	3166
intron 1	C	24778
intron 1	C	24830
exon 2	T	24942
exon 3	C	26188
exon 3	A	26308
exon 3	G	26422
intron 4	G	32394
intron 4	T	32434
exon 5	G	32493
exon 5	G	32506
exon 5	C	32507
exon 5	C	32548
intron 5	A	32783
intron 5	T	35309
intron 5	C	35374
intron 5	A	35378
exon 6	G	35438
exon 6	T	35454
intron 6	T	35549
intron 6	G	35641
intron 6	A	35725
intron 6	T	36001
intron 6	A	36064
intron 6	deletion of GTTT	36091 to 36094
intron 6	C	36108
intron 7	C	36374
intron 7	G	36378
intron 7	T	36387
intron 7	G	36398
intron 7	C	37439
intron 7	T	37513
exon 8	C	37604
exon 8	G	37605
exon 8	G	37623
exon 8	C	37633
intron 9	C	47214
exon 11	G	47383
exon 11	C	47411
intron 11	T	47563
intron 12	C	54307
intron 12	G	54308
exon 13	C	54399
exon 13	A	54480
exon 13	C	54523
exon 13	deletion of CCCTGAGAGCCACAGGTGCCT	54562 to 54582
exon 13	A	54588
exon 13	C	54664
exon 13	G	54703
exon 13	A	54804
exon 13	G	54834
exon 13	G	54847
3'UTR	G	54925
3'UTR	C	55169
3'UTR	A	55170
3'UTR	A	55171
3'UTR	C	55917

[0145] As indicated hereinabove, if the respective nucleotide which is replaced by another nucleotide is a purine base, it is preferred to be replaced by another purine base. If it is a pyrimidine base, it is preferred to be replaced by another pyrimidine base. It is also preferred that a purine base is replaced by a pyrimidine base and that a pyrimidine

base is replaced by a purine base. Most preferably, the nucleotides indicated in Table C are replaced by the nucleotides indicated at the respective position in Table 12 hereinbelow (see Example 3).

**[0146]** In a preferred embodiment the present invention relates to diagnostic composition designed for use in a method in which the occurrence of the mutation in the ATP-gated ion channel P2X7R gene is determined by PCR, immunological methods and/or electrophysiological methods as described herein below and in the appended Examples. Additionally, it is possible to determine the occurrence of a mutation in the ATP-gated ion channel P2X7R as described hereinabove.

**[0147]** In yet another aspect the present invention relates to the use of a nucleic acid molecule, a vector, a polypeptide, an antibody, aptamer and/or a primer or pair of primers of the present invention for the preparation of a diagnostic composition for the detection of an affective disorder.

**[0148]** It is also envisaged that the present invention relates to methods of diagnosing an affective disorder of an individual comprising:

**[0149]** (a) isolating DNA from cells obtained from an individual;

**[0150]** (b) determining all or part of the nucleotide composition of the P2X7R gene; and

**[0151]** (c) analyzing said nucleotide composition of P2X7R for the presence of one or more polymorphism, mutation or allelic variation.

**[0152]** The term "gene" means a nucleotide sequence associated with the production of a protein, including promoter sequences, enhancer sequences, intron sequences, exon sequences, coding regions, 5' untranslated region (5'UTR), 3' untranslated region (3'UTR), and splice variants.

**[0153]** In a preferred embodiment of the described method the individual is a mammal and more preferably human. Moreover, the cells are preferably derived from skin, blood, urine or cerebral spinal fluid.

**[0154]** The method of the present invention allows for the diagnosis of an affective disorder according to the composition of a genetic marker corresponding to the P2X7R gene. As is demonstrated by the appended examples, polymorphisms in the P2X7R are genetically linked to patients suffering from an affective disorder.

**[0155]** In accordance with this embodiment of the present invention, the diagnosis of an affective disorder can, e.g., be effected by isolating cells from an individual, and isolating the genomic DNA of said cells. Such cells can be collected from body fluids, skin, hair, biopsies and other sources. Collection and analysis of cells from bodily fluids such as blood, urine and cerebrospinal fluid is well known to the art; see for example, Rodak, "Haematology: Clinical Principles & Applications" second ed., WB Saunders Co, 2002; Brunzel, "Fundamentals of Urine and Body Fluids Analysis", WB Saunders Co, 1994; Herndon and Brumback (Ed.), "Cerebrospinal Fluid", Kluwer Academic Pub., 1989. In addition, methods for DNA isolation are well described in the art; see, for example, Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3<sup>rd</sup> edition, Cold Spring Harbor Laboratory, 2001.

**[0156]** Once DNA has been isolated, various oligonucleotide primers spanning the P2X7R locus may be designed in order to amplify the genetic material by Polymerase Chain Reaction (PCR). Conventional methods for designing, synthesizing, producing said oligonucleotide primers and performing PCR amplification may be found in standard textbooks, see, for example Agrawal (Ed.), "Protocols for Oligonucleotides and Analogs: Synthesis and Properties (Methods in Molecular Biology, 20)", Humana Press, 1993; Innis et al. (Ed.), "PCR Applications: Protocols for Functional Genomics", Academic Press, 1999; Chen and Janes (Ed.), "PCR Cloning Protocols: From Molecular Cloning to Genetic", 2<sup>nd</sup> edition, Humana Press, 2002. Primers for the detection of P2X7R polymorphisms are also given in, but not limited to, SEQ ID NO: 52 to SEQ ID NO: 111. Once DNA has been amplified, nucleotide structure can be analysed by sequencing methods and compared to normal P2X7R DNA.

**[0157]** Sequencing may be performed manually by any molecular biologist of ordinary skills or by an automated sequencing apparatus. These procedures are common in the art, see, for example, Adams et al. (Ed.), "Automated DNA Sequencing and Analysis", Academic Press, 1994; Alihey, "DNA Sequencing: From Experimental Methods to Bioinformatics", Springer Verlag Publishing, 1997.

**[0158]** Detection and analysis of polymorphisms in P2X7R can also be performed using amplification refractory mutation system (ARMS<sup>TM</sup>), amplification refractory mutation system linear extension (ALEX<sup>TM</sup>), single-strand conformation polymorphism (SSCP), heteroduplex analysis, PCR-SSCP, fluorescent SSCP in an automated DNA sequencer, denaturing gradient gel electrophoresis, RNase protection assays, detection of mutations by sequence specific oligonucleotide hybridization, chemical cleavage methods, enzyme mismatch cleavage methods, cleavage fragment length methods, allele-specific oligonucleotide hybridization on DNA chips, and other such methods known in the art, see, for example Nollau et al, Clin. Chem. 43 (1997), 1114-1128; Burczak and Mardis (Ed.), "Polymorphism Detection & Analysis Techniques", Eaton Pub Co, 2000; Cotton et al. (Ed.), "Mutation Detection: A Practical Approach", Irl Press, 1998; Taylor (Ed.), "Laboratory Methods for the Detection of Mutations and Polymorphisms in DNA", CRC Press, 1997.

**[0159]** The present invention also relates to a method of diagnosing an affective disorder in an individual comprising:

**[0160]** (a) isolating RNA from cells obtained from an individual;

**[0161]** (b) converting the RNA into cDNA;

**[0162]** (c) determining all or part of the nucleotide composition of the cDNA so obtained; and

**[0163]** (c) analyzing said nucleotide composition for the presence of one or more polymorphism(s) or allelic variation.

**[0164]** With respect to the preferred embodiments the same applies as already described above.

**[0165]** Detection and analysis of polymorphisms in the P2X7R RNA can be performed according to the methods described above.

[0166] The present invention also relates to a method for diagnosing an affective disorder in an individual comprising:

[0167] (a) isolating RNA or proteins from cells obtained from an individual;

[0168] (b) determining the levels of P2X7R RNA or protein; and

[0169] (c) comparing the levels of P2X7R RNA or protein with the corresponding levels from a normal individual not afflicted with an affective disorder.

[0170] With respect to the preferred embodiments the same applies as already described above.

[0171] As is demonstrated by the appended examples, a relationship exists between the expression, or protein level of P2X7R and an affective disorder. This and other embodiments of the present invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

[0172] According to another aspect on the invention, there is provided a polynucleotide comprising at least 20 bases of the human P2X7R gene and comprising a mutation or polymorphism selected from any of the following:

TABLE 1

Novel polymorphisms in the human P2X7R		
Region in P2X7	Polymorphism	Protein Modification
5'UTR	362 T-C	Arg117Trp
5'UTR	532 T-G	
5'UTR	1100 A-G	
5'UTR	1122 A-G	
5'UTR	1171 C-G	
5'UTR	1702 G-A	
Intron01	3166 G-C	
Intron01	24778 C-T	
Intron01	24830 6C-T	
Exon03	26188 C-T	
Intron03	26308 A-G	
Intron03	26422 G-A	
Intron04	32394 G-A	
Intron04	32434 T-C	

TABLE 1-continued

Novel polymorphisms in the human P2X7R		
Region in P2X7	Polymorphism	Protein Modification
Exon05	32493 G-A	Gly150Arg
Exon05	32548 C-T	Silent
Intron05	32783 A-C	Cys168
Exon06	35438 G-A	Glu186Lys Leu191Pro
Exon06	35454 T-C	
Intron06	35641 G-C	
Intron06	35725 A-C	Arg270Cys Silent Asp279
Intron06	36001 T-G	
Intron07	36378 G-A	
Intron07	36387 T-A	
Intron07	36398 G-C	
Exon08	37604 C-T	
Exon08	37633 C-T	
Intron09	47214 C-T	
Intron11	47563 T-C	
Intron12	54307 C-T	
Intron12	54308 G-A	deletion of 7aa 488 to 494 (PESHRL)
Exon13	54562-54582 deletion of CCCTGAGA GCCACAGGTGCCT	
Exon13	54804 A-T	
Exon13	54834 G-A	Ile568Asn
3'UTR	55169 C-A	Arg578Gln
3'UTR	55170 A-C	
3'UTR	55171 A-C	
3'UTR	55917 C-T	
3'UTR	54925 G-A	

[0173] The polymorphism describes the position and the variation observed. The position and numbering of the polymorphism corresponds to the human P2X7R gene as defined in SEQ ID No 1. Primers used for SNP amplification and sequencing are shown in Table 1a and listed in SEQ ID NO: 52 to SEQ ID NO: 111.

TABLE 1a

Primer sequences for SNP amplification and sequencing				
Primer Name	Orientation	Sequence	Begin	End
P2RX7_01.for	Sense	cgtaggacttggcgcttct	2785	2803
P2RX7_01.rev	Anti sense	gagcacgtctcagattcgaaa	3224	3244
P2RX7_02.for	Sense	ccatgaggcaggtatgactattc	24665	24687
P2RX7_02.rev	Antisense	ctcctggatctcacccagtt	25168	25187
P2RX7_03.for	Sense	ctcgtccagctttgatattaagc	25966	25988
P2RX7_03.rev	Antisense	ggtccctagtgtctagaaccaga	26426	26447
P2RX7_04.for	Sense	attcatccgtcagtgcc	30794	30811
P2RX7_04.rev	Antisense	gccatgtgaattttctaccgat	31277	31298
P2RX7_05.for	Sense	ttcgttggttaggatggg	32314	32333



TABLE 1a-continued

<u>Primer sequences for SNP amplification and sequencing</u>				
Primer Name	Orientation	Sequence	Begin	End
P2RX7_05.rev	Antisense	caaggatgctcagggtagtagc	32805	32826
P2RX7_06.for	Sense	cactaggtttgctgtatccatttct	35277	35301
P2RX7_06.rev	Antisense	gcaactgtgtgagagcttg	35731	35750
P2RX7_07.for	Sense	tcaacctggtccagtgtg	35950	35968
P2RX7_07.rev	Antisense	caccaagtagctctcactcataagg	36424	36448
P2RX7_08.for	Sense	caataacacttgtgcgagttaggt	37380	37403
P2RX7_08.rev	Antisense	catcttggtgccttgaaacc	37750	37770
P2RX7_09.for	Sense	gtgagtggtaatcctgctactgc	45321	45343
P2RX7_09.rev	Antisense	aggccactcctgtactcg	45743	45761
P2RX7_10_11.for	Sense	ccaagtcacagcatgaggc	47119	47137
P2RX7_10_11.rev	Antisense	accagcgacgtatccac	47632	47649
P2RX7_12.for	Sense	aagcatggggttcatttc	50252	50268
P2RX7_12.rev	Antisense	gcataaaagggaactcctgctagta	50691	50714
P2RX7_13a.for	Sense	gcttacagaacacatgcatgg	54232	54252
P2RX7_13a.rev	Antisense	gcacctgtaggcacagtgc	54739	54757
P2RX7_13b.for	Sense	atcaccacctcagagctgttc	54620	54640
P2RX7_13b.rev	Antisense	gttaacatggctactgcagcc	55203	55223
P2XR7_13d.for	Sense	gcttagaaaggaggcgactcc	54484	54504
P2XR7_Pro13.for	Sense	ttgtgacatttgcaaggctgcc	2617	2638
P2XR7_Pro7.rev	Antisense	tctgaagctctgctcctgag	1955	1974
P2XR7_Pro8.rev	Antisense	ctcaccttctggcttccagt	1611	1630
P2XR7_Pro9.for	Sense	cttaccactcccaggactaa	1496	1515
P2XR7_Pro10.for	Sense	gtctgcctgttcaactgccat	1149	1168
P2XR7_Pro1.for	Sense	cagagaccttcagaaacttcg	1841	1861
P2XR7_Pro2.rev	Antisense	agatcaccaggacacagtg	2261	2280
P2XR7_Pro3.for	Sense	ctcaactccactttcctcgg	2133	2152
P2XR7_Pro4.rev	Antisense	cctttcacttttttggtctcatg	2655	2677
P2XR7_Pro5.for	Sense	gggagaattctgaaaatgcc	2691	2711
P2XR7_Pro6.rev	Antisense	ggaccagagctctactcttc	2951	2970
P2XR7_Pro11.for	Sense	aggatcatagatcgacctgcc	2296	2315
P2XR7_Pro12.rev	Antisense	aagaagcgccaagtcctacg	2785	2804
P2XR7_Pro14.for	Sense	gcaatccagactgaagttgac	2051	2071
P2XR7_Pro15.rev	Antisense	actctggtctgcagttggtg	2428	2447
P2XR7_Pro21.for	Sense	cctttaaaatcagagaccttcaga	1831	1854
P2XR7_Pro22.for	Sense	gcccacctctgaacacat	2708	2727
P2XR7_3UTR10.for	Sense	cccttggaactcttgctatcg	55804	55824

TABLE 1a-continued

<u>Primer sequences for SNP amplification and sequencing</u>				
Primer Name	Orientation	Sequence	Begin	End
P2XR7_3UTR1.for	Sense	ggcagtacagtggcttcaaga	54858	54878
P2XR7_3UTR2.rev	Antisense	gtgggacagtttgctgtgcct	55150	55170
P2XR7_3UTR3.for	Sense	gagtccttaccaatagcagg	55183	55202
P2XR7_3UTR4.rev	Antisense	gtcaaagaatttggtgccacc	55643	55663
P2XR7_3UTR5.for	Sense	catgaactgtcttttaatgtgtaaa	55515	55540
		g		
P2XR7_3UTR6.rev	Antisense	gagatacggtttcacatgttg	55955	55976
P2XR7_3UTR7.for	Sense	aattagctgggcatggtgcg	55992	56011
P2XR7_3UTR8.rev	Antisense	ttgagatggagtctcgtctg	56122	56140
P2XR7_3UTR9.rev	Antisense	cactgtccacgtgactgctt	56208	56227
P2RX7_11.For	Sense	tcctacttcggtctgtaagagatt	47281	47305
P2RX7_11.Rev	Antisense	gggcctaattttcgtgcat	47591	47609
P2RX7_13G.For	Sense	aagaacctagaacctgagggctt	54333	54355
P2RX7_13G.Rev	Antisense	ttgagatgggaggcagctt	54541	54559
P2RX7_13H.For	Sense	ttcggctcccaggacat	54773	54789
P2RX7_13H.Rev	Antisense	cacagagctttgcaggtgaa	55248	55267

[0174] Another aspect of the present invention is in the form of a diagnostic kit for affective disorders comprising a specific oligonucleotide probe, or primer corresponding to P2X7R polymorphisms. The diagnostic kit may comprise appropriate packaging and instructions for the use in the method of the invention. Said kit may further comprise appropriate buffer, and enzymes such as reverse transcriptase, and thermostable polymerases.

[0175] In a preferred embodiment of the invention, diagnosis can be performed on a mouse, rat or human. The invention is generally applied in vitro, e.g. using cells or other material obtained from an individual. However, it can also be applied on a living individual, or post mortem.

[0176] In accordance with the embodiments of the present invention, diagnosis of an affective disorder may be followed by prescription, or administration of an antidepressant drug. Administration and dosage of antidepressive drugs can vary between patients and are well known in the medical art, see, for example Benkert and Hippus, "Kompendium der Psychiatrischen Pharmakotherapie", Springer Verlag Publishing, 2000; Albers, "Handbook of Psychiatric Drugs: 2001-2002 Edition", Current Clinical Strategies Publishing, 2000. Preferred examples include between 5 mg and 80 mg per day, preferably 20 mg, fluoxetine; between 5 mg and 50 mg per day, preferably 20 mg, paroxetine; between 5 mg and 200 mg per day, preferably 50 mg, sertraline; between 5 mg and 300 mg per day, preferably 100 mg, fluvoxamine; between 5 mg and 100 mg per day, preferably 30 mg, mirtazapine; between 4 mg and 50 mg, preferably 8 mg, reboxetine; between 5 mg and 600 mg per day, preferably

200 mg, nefazodone; between 450 mg and 1800 mg per day, preferably 900 mg, lithium carbonate.

[0177] The P2X7R protein is also useful for monitoring the efficacy and/or dosing of a drug or the likelihood of a patient to respond to a drug. Thus, in yet another embodiment the invention relates to a method for, monitoring the efficacy and/or dosing of a drug, e.g. an antidepressive drug, and/or the likelihood of a patient to respond to said drug which comprises determining the level of expression and/or activity of the P2X7R protein in a patient before and after administration of the respective drug. As presented in the examples below, treatment with an antidepressive drug results in an upregulation in P2X7R activity. In humans, P2X7R activity can be monitored by Positron Emission Tomography (PET) or Single Photon Emission Computerised Tomography (SPECT) using a radiolabelled ligand tracer for P2X7R. Examples of P2X7R ligands can be, but are not limited to, ATP, an antagonist binding P2X7R, an agonist binding P2X7R, or a small polynucleotide comprising at least 20 bases of the human P2X7R gene. A modulation of P2X7R activity, membrane distribution or expression levels would reflect the activity and potency of the antidepressive drug. Methods and techniques required for PET analysis are well known in the art, see, for example Paans and Vaalburg, Curr. Pharmac. Design 6 (2000), 1583-1591; van Waarde, Curr. Pharmac. Design. 6 (2000), 1593-1610; Paans et al, Methods 27 (2002), 195-207; Passchier et al., Methods 27 (2002), 278-286; Laruelle et al., Methods 27 (2002), 287-299.

[0178] In accordance with the present invention by the term "sample" is intended any biological sample obtained

from an individual, cell line, tissue culture, or other source containing polynucleotides or polypeptides or portions thereof. As indicated, biological samples include body fluids (such as blood, sera, plasma, urine, synovial fluid and spinal fluid) and tissue sources found to express the polynucleotides of the present invention. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. A biological sample which includes genomic DNA, mRNA or proteins is preferred as a source.

**[0179]** As described herein above, mutations of the P2X7R encoding gene can occur on DNA level or on mRNA level and may result in an altered expression of P2X7R or in the expression of P2X7R ATP-gated ion channels which show either an altered function or no function when compared to the wild-type P2X7R ATP-gated ion channel as described herein. Thus, various methods on DNA level, RNA level or protein level exist for determining whether the ATP-gated ion channel P2X7R gene shows a mutation as described herein above. Consequently, mRNA, cDNA, DNA and genomic DNA are the preferred nucleic acid molecules to be used in the below mentioned methods. Also polypeptides or fragments thereof are preferred if a mutation in the P2X7R ATP-gated ion channel protein as described herein is to be determined.

**[0180]** Preferably, a point mutation leading to the replacement of an amino acid residue at the positions as indicated in Table 1 of the corresponding wild-type P2X7R amino acid sequence depicted in SEQ ID NO: 3 or 4 by another amino acid can be determined by PCR. Said PCR is followed by a restriction fragment length polymorphism (RFLP) analysis if due to the point mutation a recognition site for a restriction endonuclease is generated which is not present in the wild-type nucleotide sequence or a recognition site for a restriction enzyme is created which does not occur in the wild-type P2X7R. More preferably said mutation can be determined by PCR using primers and conditions that allow only an amplification of the wild-type nucleotide sequence encoding the corresponding wild-type amino acid at the respective position, but not of the nucleotide sequence of a nucleic acid molecule encoding a different amino acid residue at the corresponding position. It is even more preferred that PCR is performed to determine a mutation using primers and conditions that allow no amplification if the wild-type nucleotide sequence is present, but only if another amino acid residue is encoded at the respective position. Particularly preferred is a method using PCR and primers under conditions that allow amplification of a fragment comprising at least the nucleotide residues encoding the amino acid residue corresponding to positions of SEQ ID NO: 1.

**[0181]** Said PCR is followed by e.g., sequencing and/or single strand conformation analysis (SSCA). Said fragment is preferably of at least 25 nucleotides in length, more preferred of at least 50 nucleotide in length, even more preferred of at least 75 nucleotides in length, particularly preferred of at least 100 nucleotides in length, more particularly preferred of at least 200 nucleotides in length, also more particularly preferred at least 250 nucleotides in length, even more particularly preferred at least 300 nucleotides in length and most particularly preferred at least 600 nucleotides in length. Said primers are preferably of at least 12 nucleotides in length, more preferred of at least 15 nucleotides in length, even more preferred of at least 18

nucleotides in length and most preferred of at least 21 nucleotides in length as depicted in SEQ ID NOs: 52 to 111. The temperature for annealing said primers is preferably at least 50° C., more preferred at least 55° C. and most preferred at least 58° C. The temperature for denaturation is preferably at least 95° C. for preferably at least 10 sec, more preferably at least 20 sec, even more preferred at least 30 sec and most preferred at least 60 sec. However, depending on the length and the G-C content of the nucleic acid sequence to be amplified the temperature for denaturation may be shorter or longer. The temperature for extension of the annealed primers is preferably at least 10 sec, more preferably at least 20 sec, even more preferred at least 30 sec and most preferred at least 60 sec. A PCR reaction comprising the aforementioned conditions is exemplified in the Examples herein below. The subsequent sequencing and/or SSCA is carried out as known in the art. Preferably, the PCR fragments are separated on a 10% polyacrylamide gel at 4° C. or also preferred at room temperature. PCR fragments showing a SSCA band shift are amplified with the primers under conditions as mentioned above and are subsequently sequenced. Alternatively, it is also possible to directly sequence genomic DNA in order to determine whether a mutation in the CLCN2 gene has occurred. A direct genomic sequencing approach is, for example, demonstrated for baker's yeast in Horecka, Yeast 16 (2000), 967-970.

**[0182]** Preferably, a deletion is determined by using hybridization techniques as known in the art. In particular, a primer is designed as mentioned herein above that is capable to only hybridize to wild-type genomic DNA as depicted in SEQ ID NO: 1 but not to a nucleotide sequence comprising a deletion of a fragment between nucleotides 54562 and 54582 of SEQ ID NO: 1. Also preferred is the method of fluorescent in situ hybridization (FISH) for determining on whole chromosomes, in particular on chromosome 12q23-q24 that said chromosome has the above mentioned deletion. Even more preferred is that a deletion of nucleotide residues as described herein may be determined by using PCR, wherein one primer of a pair of primers is located within the region of genomic DNA comprising said deletion. Preferably, said deletion is between nucleotide positions 54562 and 54582 as depicted in SEQ ID NO: 1. Thus, under the appropriate conditions no PCR fragment will result if the genomic DNA comprises said deletion. It is particularly preferred that PCR using primers which are located upstream or downstream of the deletion is performed to determine said deletion. Under appropriate conditions as mentioned herein above, both a fragment of genomic DNA of the wild-type nucleotide sequence as set forth in SEQ ID NO: 1 and a fragment of the nucleotide sequence comprising a deletion of preferably the nucleotides between positions 54562 and 54582 as depicted in SEQ ID NO: 1 will be amplified.

**[0183]** It is also possible to determine the above-described P2X7R mutations on the protein level. Some of the mutations described above lead to shortened versions of the P2X7R protein. Thus, it is conceivable to determine the occurrence of these mutations by determining the length or molecular weight of the P2X7R protein expressed in an individual, e.g. by SDS PAGE.

**[0184]** It is also possible to determine the mutations of the P2X7R ATP-gated channel as described herein by using the antibodies of the present invention. Said antibodies specific

for said mutations of P2X7R proteins will be determined by assay techniques such as radioimmunoassays, competitive-binding assays, Western blot analysis and ELISA assay. Also preferred are classical immunohistological methods.

**[0185]** The finding, described in the present invention, that certain mutations in the P2X7R encoding gene and/or the corresponding protein are connected with affective disorder is indicative that the non- or dysfunction of the P2X7R protein is responsible for various forms of affective disorders. Thus, the finding of these mutations not only allows the diagnosis of affective disorders by determining whether the above-described mutations occur in an individual. It also allows to develop a treatment of affective disorders which has been diagnosed to be the result of a mutation in the P2X7R encoding gene. Such a treatment can, e.g., comprise the introduction of a nucleic acid molecule encoding a non-functional or functional wild-type P2X7R protein thereby restoring in said individual the P2X7R activity or the activation or repression of (a) P2X7R gene(s) in vivo. The term “activation or repression” in this context means that the expression of the gene is either enhanced (activation) or reduced (repression). An enhancement of expression can, e.g., be achieved by increasing the efficiency of transcription initiation, for example, by using suitable compounds which have an activating effect on transcription. Alternatively, an enhancement can be achieved by replacing the naturally occurring promoter by a more efficient promoter.

**[0186]** A repression may be achieved by suppressing expression of the gene, e.g., by specifically suppressing transcription from the respective promoter by suitable compounds or by rendering the promoter less efficient or non-functional.

**[0187]** In another embodiment the present invention also relates to a pharmaceutical composition. In accordance with the present invention the term “pharmaceutical composition” relates to a composition comprising a nucleic acid molecule comprising a nucleotide sequence which encodes an ATP-gated ion channel P2X7R and which is selected from the group consisting of:

**[0188]** (a) a nucleotide sequence encoding a polypeptide comprising the amino acid sequence as depicted in SEQ ID NO: 3 or 4;

**[0189]** (b) a nucleotide sequence comprising the nucleotide sequence as depicted in SEQ ID NO: 1 or SEQ ID NO: 2;

**[0190]** (c) a nucleotide sequence which hybridizes to the nucleotide sequence of (a) or (b); and

**[0191]** (d) a nucleotide sequence which is degenerated as a result of the genetic code to the nucleotide sequence of (c).

**[0192]** Such pharmaceutical compositions comprise a therapeutically effective amount of a nucleic acid molecule encoding a functional P2X7R protein and, optionally, a pharmaceutically acceptable carrier. The pharmaceutical composition may be administered with a physiologically acceptable carrier to a patient, as described herein. In a specific embodiment, the term “pharmaceutically acceptable” means approved by a regulatory agency or other generally recognized pharmacopoeia for use in animals, and

more particularly in humans. The term “carrier” refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered.

**[0193]** Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium ion, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in “Remington’s Pharmaceutical Sciences” by E. W. Martin. Such compositions will contain a therapeutically effective amount of the aforementioned compounds, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

**[0194]** In another preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

**[0195]** The pharmaceutical composition of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

**[0196]** In vitro assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of

administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems. Preferably, the pharmaceutical composition is administered directly or in combination with an adjuvant.

**[0197]** The pharmaceutical composition is preferably designed for the application in gene therapy. The technique of gene therapy has already been described above in connection with the nucleic acid molecules of the invention and all what has been said there also applies in connection with the pharmaceutical composition. For example, the nucleic acid molecule in the pharmaceutical composition is preferably in a form which allows its introduction, expression and/or stable integration into cells of an individual to be treated.

**[0198]** For gene therapy, various viral vectors which can be utilized, for example, adenovirus, herpes virus, vaccinia, or, preferably, an RNA virus such as a retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can also incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. By inserting a P2X7R sequence of interest encoding a functional P2X7R protein into the viral vector, along with another gene which encodes, for example, the ligand for a receptor on a specific target cell, for example, the vector is now target specific.

**[0199]** Retroviral vectors can be made target specific by inserting, for example, a polynucleotide encoding a sugar, a glycolipid, or a protein. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome to allow target specific delivery of the retroviral vector containing the inserted polynucleotide sequence.

**[0200]** Since recombinant retroviruses are preferably defective, they require assistance in order to produce infectious vector particles. This assistance can be provided, for example, by using helper cell lines that contain plasmids encoding all of the structural genes of the retrovirus under the control of regulatory sequences within the LTR. These plasmids are missing a nucleotide sequence which enables the packaging mechanism to recognize an RNA transcript for encapsidation. Helper cell lines which have deletions of the packaging signal include, but are not limited to w2, PA317 and PA12, for example. These cell lines produce empty virions, since no genome is packaged. If a retroviral vector is introduced into such cells in which the packaging signal is intact, but the structural genes are replaced by other genes of interest, the vector can be packaged and vector virion produced. Alternatively, NIH 3T3 or other tissue culture cells can be directly transfected with plasmids encoding the retroviral structural genes gag, pol and env, by conventional calcium phosphate transfection. These cells are then transfected with the vector plasmid containing the genes of interest. The resulting cells release the retroviral

vector into the culture medium. Another targeted delivery system for P2X7R polynucleotides is a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The preferred colloidal system of this invention is a liposome. Liposomes are artificial membrane vesicles which are useful as delivery vehicles in vitro and in vivo. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0  $\mu\text{m}$  can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley, et al., *Trends Biochem. Sci.*, 6:77, 1981). In addition to mammalian cells, liposomes have been used for delivery of polynucleotides in plant, yeast and bacterial cells. In order for a liposome to be an efficient gene transfer vehicle, the following characteristics should be present: (1) encapsulation of the genes of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information (Mannino, et al., *Biotechniques*, 6:682, 1988). The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations. Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebrosides, and gangliosides. Particularly useful are diacylphosphatidylglycerols, where the lipid moiety contains from 14-18 carbon atoms, particularly from 16-18 carbon atoms, and is saturated. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine. The targeting of liposomes can be classified based on anatomical and mechanistic factors. Anatomical classification is based on the level of selectivity, for example, organ-specific, cell-specific, and organelle-specific. Mechanistic targeting can be distinguished based upon whether it is passive or active. Passive targeting utilizes the natural tendency of liposomes to distribute to cells of the reticulo-endothelial system (RES) in organs which contain sinusoidal capillaries.

**[0201]** In another aspect the present invention relates to a method of treating an affective disorder comprising administering a therapeutically effective amount of the pharmaceutical composition comprising a nucleotide sequence encoding a functional ATP-gated ion channel as described herein above to a subject suffering from said disorder.

**[0202]** In the context of the present invention the term "subject" means an individual in need of a treatment of an affective disorder. Preferably, the subject is a vertebrate, even more preferred a mammal, particularly preferred a human.

**[0203]** The term "administered" means administration of a therapeutically effective dose of the aforementioned nucleic

acid molecule encoding a functional P2X7R protein to an individual. By "therapeutically effective amount" is meant a dose that produces the effects for which it is administered. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques. As is known in the art and described above, adjustments for systemic versus localized delivery, age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by those skilled in the art.

[0204] The methods are applicable to both human therapy and veterinary applications. The compounds described herein having the desired therapeutic activity may be administered in a physiologically acceptable carrier to a patient, as described herein. Depending upon the manner of introduction, the compounds may be formulated in a variety of ways as discussed below. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt %. The agents may be administered alone or in combination with other treatments.

[0205] The administration of the pharmaceutical composition can be done in a variety of ways as discussed above, including, but not limited to, orally, subcutaneously, intravenously, intra-arterial, intranodal, intramedullary, intrathecal, intraventricular, intranasally, intrabronchial, transdermally, intranodally, intrarectally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, or intraocularly. In some instances, for example, in the treatment of wounds and inflammation, the candidate agents may be directly applied as a solution dry spray.

[0206] The attending physician and clinical factors will determine the dosage regimen. As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. A typical dose can be, for example, in the range of 0.001 to 1000  $\mu$ g; however, doses below or above this exemplary range are envisioned, especially considering the aforementioned factors.

[0207] The dosages are preferably given once a week, however, during progression of the treatment the dosages can be given in much longer time intervals and in need can be given in much shorter time intervals, e.g., daily. In a preferred case the immune response is monitored using herein described methods and further methods known to those skilled in the art and dosages are optimized, e.g., in time, amount and/or composition. Dosages will vary but a preferred dosage for intravenous administration of DNA is from approximately  $10^6$  to  $10^{12}$  copies of the DNA molecule. If the regimen is a continuous infusion, it should also be in the range of 1  $\mu$ g to 10 mg units per kilogram of body weight per minute, respectively. Progress can be monitored by periodic assessment. The pharmaceutical composition of the invention may be administered locally or systemically. Administration will preferably be parenterally, e.g., intravenously. Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous

carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium ion solution, Ringer's dextrose, dextrose and sodium ion, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

[0208] It is also envisaged that the pharmaceutical compositions are employed in co-therapy approaches, i.e. in co-administration with other medicaments or drugs, for example other drugs for preventing, treating or ameliorating affective disorders.

[0209] Another aspect of the present invention is a pharmaceutical composition comprising a compound, the administration of which to cells leads to a reduction or increase of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells or comprising a nucleic acid molecule the expression of which in cells or the administration of which to cells leads to a reduction or increase of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells. Said pharmaceutical composition may be useful for treating individuals having an increased or reduced amount of the P2X7R protein or expression level as described hereinabove. Preferably, said pharmaceutical composition leads to an increase of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells.

[0210] It is envisaged that the above-mentioned pharmaceutical composition, the administration of which to cells leads to a reduction of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R is an antisense nucleic acid, a ribozyme, a co-suppressive nucleic acid, iRNA or siRNA.

[0211] An siRNA approach is, for example, disclosed in Elbashir ((2001), Nature 411, 494-498)). It is also envisaged in accordance with this invention that for example short hairpin RNAs (shRNAs) are employed in accordance with this invention as pharmaceutical composition. The shRNA approach for gene silencing is well known in the art and may comprise the use of st (small temporal) RNAs; see, inter alia, Paddison (2002) Genes Dev. 16, 948-958.

[0212] As mentioned above, approaches for gene silencing are known in the art and comprise "RNA"-approaches like RNAi or siRNA. Successful use of such approaches has been shown in Paddison (2002) loc. cit., Elbashir (2002) Methods 26, 199-213; Novina (2002) Mat. Med. Jun. 3, 2002; Donze (2002) Nucl. Acids Res. 30, e46; Paul (2002) Nat. Biotech 20, 505-508; Lee (2002) Nat. Biotech. 20, 500-505; Mivagashi (2002) Nat. Biotech. 20, 497-500; Yu (2002) PNAS 99, 6047-6052 or Brummelkamp (2002), Science 296, 550-553. These approaches may be vector-based, e.g. the pSUPER vector, or RNA polIII vectors may be employed as illustrated, inter alia, in Yu (2002) loc. cit.; Mivagashi (2002) loc. cit. or Brummelkamp (2002) loc. cit.

[0213] A compound which leads to a reduction of the expression of the P2X7R gene may, e.g., be a compound which acts on the regulatory region of the gene and thereby reduces the level of transcription. Such compounds can be identified by methods as described herein below.

[0214] The invention also relates to the use of a nucleic acid molecule encoding a functional P2X7R protein as described herein above in connection with the pharmaceutical composition for the preparation of a pharmaceutical composition for treating an affective disorder.

[0215] Furthermore, the present invention relates to a method of treating an affective disorder comprising administering a therapeutically effective amount of the nucleic acid molecule according to the invention or a therapeutically effective amount of the corresponding encoded polypeptide to a subject suffering from said disorder.

[0216] In another preferred embodiment the present invention relates to a pharmaceutical composition comprising, inter alia, the polynucleotides according to the present invention, i.e. polynucleotides having mutations and/or deletions as described hereinabove. Such pharmaceutical compositions may, e.g., be useful for treating individuals having an increased or decreased amount of the P2X7R protein or having a P2X7R protein showing an increased or decreased activity which can be determined as described hereinabove. Preferably, such a pharmaceutical composition may be useful for treating individuals having a decreased amount of the P2X7R protein or having a P2X7R protein showing a decreased activity which can be determined as described hereinabove. It is envisaged that, e.g. a non-functional or preferably a hyperfunctional P2X7R protein comprised by said pharmaceutical composition is incorporated in a P2X7R complex which naturally exists in cells as described hereinabove. It is also envisaged that the above-described techniques for gene therapy can be used for treating an individual with the nucleic acid molecules of the present invention, *mutatis mutandis*.

[0217] With respect to the possible modes of administration and preferred embodiments the same applies as has been set forth above.

[0218] Additionally, the present invention also envisages the use of the nucleic acid molecules, the vectors, the polypeptides, the antibody and/or the aptamer according to the invention for the preparation of a pharmaceutical composition for the treatment of an affective disorder.


[0219] A further aspect of the present invention is the use of a modulator of P2X7R activity or expression for the preparation of a pharmaceutical composition for treating an affective disorder. In the context of the present invention the term "modulator" means (a) compound(s), a complex of compounds, (a) substance(s) or complex of substances which can modify, i.e. modulate the activity of P2X7R or the expression of P2X7R either directly or indirectly. The modulation can, for example, occur at the protein level. Particularly, the P2X7R protein may interfere with the modulator such that it is either more active or less active. The modulation can also occur on nucleic acid level. Namely, the gene is transcribed more frequently or less frequently giving rise to more or less protein. Modulation can also influence RNA or protein stability. Since it was surprisingly found that agonists of P2X7R improve the symptoms of mice selected for anxiety and depressive behaviour, the modulator of P2X7R activity used for the preparation of a pharmaceutical composition for treating an affective disorder is preferably an agonist. The term "agonist" means an agent or a compound that can interact with a receptor and initiate a physiological or a pharmacological response characteristic of that

receptor. Examples of P2X7R agonist include but are not restricted to ATP, ATP-4, BZATP (2'-3'-O-(4-Benzoylbenzoyl)adenosine 5'-triphosphate ( $C_{24}H_{24}N_5O_{15}P_3$ )) and tenidap (5-chloro-2,3-dihydro-2-oxo-3-(2-thienylcarbonyl)-indole-1-carboxamide, i.e.  $C_{15}H_{11}ClN_2O_2S$ ) or a derivative thereof. Particularly preferred, said agonist used to treat depression or anxiety is BZATP as demonstrated in Example 9 hereinbelow.

[0220] Although it was reported that activation of P2X7R could induce apoptosis and cell death in vitro (Di Virgilio et al., *Cell Death Differ.* 5 (1988), 191-199; Virginio et al., *J. Physiol.* 519 (1999), 335-346), the present application demonstrates in Example 9 hereinbelow that treatment of the brain of mice selected for anxiety and depressive behaviour with BZATP revealed no significant difference in the numbers of apoptotic cells between control mice and mice treated with BZATP. This result indicates that activation of P2X7R did not result in cerebral cell death in vivo which, thus, renders BZATP to be a candidate drug for treatment of affective disorders.

[0221] It is furthermore envisaged that said modulator is selected from the group consisting of piperidine and piperazine derivatives, adamantane derivatives, substituted phenyl compounds, oxidized ATP, 2-O-(4-benzoylbenzoyl)adenosine-5-triphosphate and 3-O-(4-benzoylbenzoyl)adenosine-5-triphosphate as, for example, described in WO 99/29660, WO 99/29661; WO 99/296896; WO 00/61569; WO 01/42194; WO 01/44170; WO 01/44213; WO 00/71529; WO 01/46200. The following compounds illustrate compounds which are also preferred to be used as modulators of P2X7R activity.


[0222] A compound of general formula:



[0223] where A is phenyl or a 5- or 6-membered heterocyclic ring containing one or two heteroatoms selected from O, N or S; and optionally substituted by  $C_{1-6}$ alkyl, halogen, nitro, amino, alkylamino,  $CF_3$ ,  $SO_2Me$ ,  $NHSO_2Me$  or cyano; B is  $C=O$ ,  $NH$  or  $SO_2$ ; X is  $C=O$ ,  $CH(Me)$ , O or  $(CH_2)_p$  where p is 0 or 1; Y is O,  $CH_2$ ,  $NH$  or S; Z is  $C=O$  or  $SO_2$ , provided that when Z is  $C=O$ , then Y is O,  $CH_2$  or S; R is hydrogen or  $C_{1-6}$ alkyl;  $R^1$  is hydrogen, halogen;  $R^2$  is phenyl optionally substituted by  $CO_2H$ ,  $CO_2$ alkyl,  $CONH$ , or  $R^2$  is  $OH$ ,  $NHR^3$ ,  $NHCH(R^4)(CHR^5)_nR^6$ ,  $NH-R^7-R^8$ ,  $SO_2NH$ alkyl,  $NHCO$ alkyl,  $NHSO_2$ alkyl, morpholine,  $NR^9R^{10}$ , piperazine substituted by phenyl, alkoxyphenyl, pyridyl or fluorophenyl; n is 0, 1 or 2;  $R^3$  is hydrogen, a bi- or tricyclic saturated ring system optionally containing a nitrogen atom, piperidinyl, alkylpyrrolidine, ethinylcyclohexyl, a 5-membered aromatic ring containing 2 or 3 heteroatoms,  $C_{4-6}$  cycloalkyl optionally substituted by alkyl, cyano or hydroxy, or  $C_{1-8}$  alkyl optionally containing an oxygen atom in the alkyl chain and being optionally substituted by one or more substituents selected from ethy-

nyl, cyano, fluoro, dialkylamino, hydroxy, thioalkyl,  $\text{CO}_2\text{R}^{11}$  or  $\text{CONH}_2$ ;  $\text{R}^4$  is hydrogen or alkyl optionally substituted by hydroxy or alkoxy;  $\text{R}^5$  is hydrogen or hydroxy;  $\text{R}^6$  is  $\text{CO}_2\text{R}^{12}$ ,  $\text{NHCO}_2\text{R}^{12}$ ,  $\text{CONH}_2$  or a 5 or 6-membered saturated ring containing an oxygen atom, a 5-membered heterocyclic ring containing one or two heteroatoms selected from O, N or S, or phenyl optionally substituted by one or more groups selected from alkyl, hydroxy, amino, alkoxy, or nitro;  $\text{R}^6$  is alkyl;  $\text{R}^7$  is a cyclopentane ring;  $\text{R}^8$  is phenyl;  $\text{R}^9$  and  $\text{R}^{10}$  are independently hydrogen, benzyl, alkenyl, cycloalkyl, alkyl optionally substituted by hydroxy, alkoxy, cyano, dialkylamino, phenyl, pyridyl or  $\text{CO}_2\text{R}^{11}$  or  $\text{R}^9$  and  $\text{R}^{10}$  together form a 5- to 7-membered saturated or partially saturated ring optionally containing a further heteroatom and optionally substituted by one or more groups selected from alkyl (optionally containing an oxygen atom in the chain and optionally substituted 14 by hydroxy),  $\text{COalkyl}$ ,  $\text{CO}_2\text{R}^{11}$ ,  $\text{COR}^{13}\text{R}^{14}$ ,  $\text{CHO}$  or piperidine,  $\text{R}^{11}$  is hydrogen or alkyl;  $\text{R}^{12}$  is alkyl; and  $\text{R}^{13}$  and  $\text{R}^{14}$  are independently hydrogen or alkyl, is or a pharmaceutically acceptable salt or solvate thereof.


[0224] A compound of general formula:




[0225] wherein A represents a group  $\text{CH}_2$  or an oxygen atom; B represents a hydrogen or halogen atom; D represents a group  $\text{CH}_2$ ,  $\text{OCH}_2$ ,  $\text{NHCH}_2$  or  $\text{CH}_2\text{CH}_2$ ; R represents a phenyl, benzothiazolyl, indolyl, indazolyl, purinyl, pyridyl, pyrimidinyl or thiophenyl group, each of which may be optionally substituted by one or more substituents independently selected from a halogen atom or a cyano, carboxyl, hydroxyl, nitro, halo- $\text{C}_1\text{-C}_6\text{-alkyl}$ ,  $-\text{N}(\text{R}^1)-\text{C}(=\text{O})-\text{R}^2$ ,  $-\text{C}(\text{O})-\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{R}^6$ ,  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$ , 3- to 8-membered heterocyclyl,  $\text{C}_3\text{-C}_8\text{-cycloalkyloxy}$ ,  $\text{C}_1\text{-C}_6\text{-alkylcarbonyl}$ , phenoxy, benzyl,  $\text{C}_1\text{-C}_6\text{-alkylthio}$ , phenylthio,  $\text{C}_1\text{-C}_6\text{-alkoxycarbonyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkylsulphinyl}$  or  $\text{C}_1\text{-C}_6\text{-alkylsulphonyl}$  group, or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_1\text{-C}_6\text{-alkoxy}$  group optionally substituted by one or more substituents independently selected from a halogen atom or an amino, carboxyl, hydroxyl,  $\text{C}_1\text{-C}_6\text{-alkoxy}$ , (di) $\text{C}_1\text{-C}_6\text{-alkylamino}$ ,  $\text{C}_1\text{-C}_6\text{-alkoxycarbonyl}$ , imidazolyl, morpholinyl, piperidinyl or pyrrolidinyl group;  $\text{R}^1$  represents a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^2$  represents a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group; and  $\text{R}^3$ ,  $\text{R}^4$ ,  $\text{R}^5$  and  $\text{R}^6$  each independently represent a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group; with the provisos that when A is  $\text{CH}_2$ , B is H and D is  $\text{CH}_2$ , then R does not represent a phenyl, ortho-carboxyphenyl, methylphenyl or para-phenoxyphenyl group, and that when A is  $\text{CH}_2$ , D is  $\text{CH}_2$  or  $\text{CH}_2\text{CH}_2$  and R represents a substituted phenyl group, the substituent or substituents present do not comprise, in an ortho position, a  $\text{C}_1\text{-C}_6\text{-alkoxy}$  group substituted by an amino, (di) $\text{C}_1\text{-C}_6\text{-alkylamino}$ , imidazolyl, mor-

pholinyl, piperidinyl or pyrrolidinyl group; or a pharmaceutically acceptable salt or solvate thereof.

[0226] A compound of general formula:




[0227] wherein x represents 1 or 2; A represents a group  $\text{CH}_2$  or an oxygen atom; B represents a hydrogen or halogen atom; R represents a phenyl, pyridyl, indolyl, indazolyl, pyrimidinyl or thiophenyl group, each of which may be optionally substituted by one or more substituents independently selected from a halogen atom or an amino, cyano, carboxyl, hydroxyl, nitro,  $\text{C}_1\text{-C}_6\text{-alkyl}$ , halo- $\text{C}_1\text{-C}_6\text{-alkyl}$ ,  $-\text{N}(\text{R}^1)-\text{C}(=\text{O})-\text{R}^2$ ,  $-\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{R}^6$ ,  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$ , 3- to 8-membered heterocyclyl,  $\text{C}_3\text{-C}_8\text{-cycloalkyloxy}$ ,  $\text{C}_1\text{-C}_6\text{-alkylcarbonyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkoxycarbonyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkylsulphinyl}$  or  $\text{C}_1\text{-C}_6\text{-alkylsulphonyl}$  group, or a  $\text{C}_1\text{-C}_6\text{-alkoxy}$ ,  $\text{C}_1\text{-C}_6\text{-alkylamino}$ , phenoxy, benzyl,  $\text{C}_1\text{-C}_6\text{-alkylthio}$  or phenylthio group optionally substituted by one or more substituents independently selected from a 15 halogen atom or an amino, cyano, carboxyl, hydroxyl, nitro, 1-pyrrolidinyl, 1-piperidinyl,  $\text{C}_1\text{-C}_6\text{-alkyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkoxy}$ , (di) $\text{C}_1\text{-C}_6\text{-alkylamino}$ , halo- $\text{C}_1\text{-C}_6\text{-alkyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkoxycarbonyl}$  or one of the following groups:




[0228]  $\text{R}^1$  represents a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^2$  represents a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^3$  and  $\text{R}^4$  each independently represent a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^5$  represents a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^6$  represents a  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group and, additionally, a  $\text{C}_1\text{-C}_6\text{-alkyl}$  group when  $\text{R}^5$  is not a hydrogen atom;  $\text{R}^7$  represents a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^8$  represents a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^9$  represents a hydrogen atom or a hydroxyl group; and  $\text{R}^{10}$  represents a hydrogen atom or a phenyl or imidazolyl group; with the provisos that R does not represent an unsubstituted pyridyl group when A represents a group  $\text{CH}_2$  and B represents a hydrogen atom, and that when R represents a substituted phenyl, indolyl or indazolyl group, the substituent or substituents present do not comprise an amido, carboxyl, (di) $\text{C}_1\text{-C}_6\text{-alkylamido}$  or  $\text{C}_1\text{-C}_6\text{-alkoxycarbonyl}$  group in an ortho position; or a pharmaceutically acceptable salt or solvate thereof.



[0229] A compound of general formula:




[0230] wherein D represents  $\text{CH}_2$  or  $\text{CH}_2\text{CH}_2$ ; E represents  $\text{C}(\text{O})\text{NH}$  or  $\text{NHC}(\text{O})$ ;  $\text{R}^1$  and  $\text{R}^2$  each independently represent a hydrogen or halogen atom, or an amino, nitro,  $\text{C}_1$ - $\text{C}_6$ -alkyl or trifluoromethyl group;  $\text{R}^3$  represents a group of formula:




[0231] X represents an oxygen or sulphur atom or a group  $\text{NH}$ ,  $\text{SO}$  or  $\text{SO}_2$ ; Y represents an oxygen or sulphur atom or a group  $\text{NR}^{11}$ ,  $\text{SO}$  or  $\text{SO}_2$ ; Z represents a group  $-\text{OH}$ ,  $-\text{SH}$ ,  $-\text{CO}_2\text{H}$ ,  $\text{C}_1$ - $\text{C}_6$ -alkoxy,  $\text{C}_1$ - $\text{C}_6$ -alkylthio,  $\text{C}_1$ - $\text{C}_6$ -alkylsulphinyl,  $\text{C}_1$ - $\text{C}_6$ -alkylsulphonyl,  $-\text{NR}^6\text{R}^7$ ,  $-\text{C}(\text{O})\text{NR}^8\text{R}^9$ , imidazolyl, 1-methylimidazolyl,  $-\text{N}(\text{R}^{10})\text{C}(\text{O})-\text{C}_1$ - $\text{C}_6$ -alkyl,  $\text{C}_1$ - $\text{C}_6$ -alkylcarbonyloxy,  $\text{C}_1$ - $\text{C}_6$ -alkoxycarbonyloxy,  $-\text{OC}(\text{O})\text{NR}^{12}\text{R}^{13}$ ,  $-\text{OCH}_2\text{OC}(\text{O})\text{R}^{14}$ ,  $-\text{OCHOC}(\text{O})\text{OR}^{15}$  or  $-\text{OC}(\text{O})\text{OCH}_2\text{OR}^{16}$ ,  $\text{R}^4$  represents a  $\text{C}_2$ - $\text{C}_6$ -alkyl group;  $\text{R}^5$  represents a  $\text{C}_1$ - $\text{C}_6$ -alkyl group;  $\text{R}^6$ ,  $\text{R}^7$ ,  $\text{R}^8$ ,  $\text{R}^9$ ,  $\text{R}^{10}$ ,  $\text{R}^{12}$ , and  $\text{R}^{13}$  each independently represent a hydrogen atom, or a  $\text{C}_1$ - $\text{C}_6$ -alkyl group optionally substituted by at least one hydroxyl group;  $\text{R}^{11}$  represents a hydrogen atom, or a  $\text{C}_1$ - $\text{C}_6$ -alkyl group optionally substituted by at least one substituent independently selected from hydroxyl and  $\text{C}_1$ - $\text{C}_6$ -alkoxy; and  $\text{R}^{14}$ ,  $\text{R}^{15}$  and  $\text{R}^{16}$  each independently represent a  $\text{C}_1$ - $\text{C}_6$ -alkyl group; with the provisos that (i) when E represents  $\text{NEC}(\text{O})$ , X represents O, S or NH and Y represents O, then Z represents  $-\text{NR}^6\text{R}^7$  where  $\text{R}^6$  represents a hydrogen atom and  $\text{R}^7$  represents either a hydrogen atom or a  $\text{C}_1$ - $\text{C}_6$ -alkyl group substituted by at least one hydroxyl group, and (ii) when E represents  $\text{NHC}(\text{O})$ , X represents O, S or NH, Y represents NH, and  $\text{R}^5$  represents  $\text{CH}_2\text{CH}_2$ , then Z is not  $-\text{OH}$  or imidazolyl; or a pharmaceutically acceptable salt or solvate thereof.

[0232] A compound of general formula:



[0233] wherein D represents  $\text{CH}_2$  or  $\text{CH}_2\text{CH}_2$ ; E represents  $\text{C}(\text{O})\text{NH}$  or  $\text{NHC}(\text{O})$ ;  $\text{R}^1$  and  $\text{R}^2$  each independently

represent hydrogen, halogen, amino, nitro,  $\text{C}_1$ - $\text{C}_6$ -alkyl or trifluoromethyl, but  $\text{R}^1$  and  $\text{R}^2$  may not both simultaneously represent hydrogen;  $\text{R}^3$  represents a group of formula




[0234]  $\text{R}^4$  represents a  $\text{C}_1$ - $\text{C}_6$ -alkyl group; X represents an oxygen or sulphur atom or a group  $\text{NR}^{13}$ ,  $\text{SO}$  or  $\text{SO}_2$ ;  $\text{R}^5$  represents hydrogen, or  $\text{R}^5$  represents  $\text{C}_1$ - $\text{C}_6$ -alkyl or  $\text{C}_2$ - $\text{C}_6$ -alkenyl, each of which may be optionally substituted by at least one substituent selected from halogen, hydroxyl, (di) $\text{C}_1$ - $\text{C}_6$ -alkylamino,  $-\text{Y}-\text{R}^6$ ,




[0235] and a 5- or 6-membered heteroaromatic ring comprising from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulphur which heteroaromatic ring may itself be optionally substituted by at least one substituent selected from halogen, hydroxyl and  $\text{C}_1$ - $\text{C}_6$ -alkyl; Y represents an oxygen or sulphur atom or a group  $\text{NH}$ ,  $\text{SO}$  or  $\text{SO}_2$ ;  $\text{R}^6$  represents a group  $-\text{R}^7\text{Z}$  where  $\text{R}^7$  represents a  $\text{C}_2$ - $\text{C}_6$ -alkyl group and Z represents an  $-\text{OH}$ ,  $-\text{CO}_2\text{H}$ ,  $-\text{NR}^8\text{R}^9$ ,  $-\text{C}(\text{O})\text{NR}^{10}\text{R}^{11}$  or  $-\text{N}(\text{R}^{12})\text{C}(\text{O})-\text{C}_1$ - $\text{C}_6$ -alkyl group, and, in the case where Y represents an oxygen or sulphur atom or a group  $\text{NH}$ ,  $\text{R}^6$  additionally represents hydrogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl,  $\text{C}_1$ - $\text{C}_6$ -alkylcarbonyl,  $\text{C}_1$ - $\text{C}_6$ -alkoxycarbonyl,  $-\text{C}(\text{O})\text{NR}^{14}\text{R}^{15}$ ,  $-\text{CH}_2\text{OC}(\text{O})\text{R}^{16}$ ,  $-\text{CH}_2\text{OC}(\text{O})\text{OR}^{17}$  or  $-\text{C}(\text{O})\text{OCH}_2\text{OR}^{18}$ ;  $\text{R}^8$ ,  $\text{R}^9$ ,  $\text{R}^{10}$ ,  $\text{R}^{11}$  and  $\text{R}^{12}$  each independently represent a hydrogen atom or a  $\text{C}_1$ - $\text{C}_6$ -alkyl group;  $\text{R}^{13}$  represents hydrogen,  $\text{C}_3$ - $\text{C}_8$ -cycloalkyl,  $\text{C}_3$ - $\text{C}_8$ -cycloalkylmethyl, or  $\text{R}^{13}$  represents a  $\text{C}_1$ - $\text{C}_6$ -alkyl group optionally substituted by at least one substituent selected from hydroxyl to and  $\text{C}_1$ - $\text{C}_6$ -alkoxy; and  $\text{R}^{14}$ ,  $\text{R}^{15}$ ,  $\text{R}^{16}$ ,  $\text{R}^{17}$  and  $\text{R}^{18}$  each independently represent a  $\text{C}_1$ - $\text{C}_6$ -alkyl group; with the proviso that when E is  $\text{C}(\text{O})\text{NH}$ , X is O, NH or  $\text{N}(\text{C}_1\text{-C}_6\text{-alkyl})$ , then  $\text{R}^5$  is other than a hydrogen atom or an unsubstituted  $\text{C}_1$ - $\text{C}_6$ -alkyl group; or a pharmaceutically acceptable salt or solvate thereof.

[0236] A compound of general formula:




[0237] wherein m represents 1, 2 or 3; each  $\text{R}^1$  independently represents a hydrogen or halogen atom; A represents  $\text{C}(\text{O})\text{NH}$  or  $\text{NHC}(\text{O})$ ; Ar represents a group



[0238] X represents a bond, an oxygen atom or a group CO, (CH<sub>2</sub>)<sub>1-6</sub>, CH=, (CH<sub>2</sub>)<sub>1-6</sub>O, O(CH<sub>2</sub>)<sub>1-6</sub>, O(CH<sub>2</sub>)<sub>2-6</sub>O, O(CH<sub>2</sub>)<sub>2-3</sub>O(CH<sub>2</sub>)<sub>1-3</sub>, CR'(OH), (CH<sub>2</sub>)<sub>1-3</sub>O(CH<sub>2</sub>)<sub>1-3</sub>, (CH<sub>2</sub>)<sub>1-3</sub>O(CH<sub>2</sub>)<sub>2-3</sub>O, NR<sup>5</sup>, (CH<sub>2</sub>)<sub>1-6</sub>NR<sup>5</sup>, NR<sup>5</sup>(CH<sub>2</sub>)<sub>1-6</sub>, (CH<sub>2</sub>)<sub>1-3</sub>NR<sup>5</sup>(CH<sub>2</sub>)<sub>1-3</sub>, O(CH<sub>2</sub>)<sub>2-6</sub>NR<sup>5</sup>, O(CH<sub>2</sub>)<sub>2-3</sub>NR<sup>5</sup>(CH<sub>2</sub>)<sub>1-3</sub>, (CH<sub>2</sub>)<sub>1-3</sub>NR<sup>5</sup>(CH<sub>2</sub>)<sub>2-3</sub>O, NR<sup>5</sup>(CH<sub>2</sub>)<sub>1-3</sub>, CONR<sup>5</sup>, NR<sup>5</sup>CO, S(O)<sub>n</sub>, S(O)<sub>n</sub>CH<sub>2</sub>, CH<sub>2</sub>S(O)<sub>n</sub>, SO<sub>2</sub>NR<sup>5</sup> or NR<sup>5</sup>SO<sub>2</sub>; n is 0, 1 or 2; R<sup>1</sup> represents a hydrogen atom or a C<sub>1</sub>-C<sub>6</sub>-alkyl group; one of R<sup>2</sup> and R<sup>3</sup> represents a halogen, cyano, nitro, amino, hydroxyl, or a group selected from (i) C<sub>1</sub>-C<sub>6</sub>-alkyl optionally substituted by at least one C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, (ii) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, (iii) C<sub>1</sub>-C<sub>6</sub>-alkyloxy optionally substituted by at least one C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, and (iv) C<sub>3</sub>-C<sub>8</sub>-cycloalkyloxy, each of these groups being optionally substituted by one or more fluorine atoms, and the other of R<sup>2</sup> and R<sup>3</sup> represents a hydrogen or halogen atom; either R<sup>4</sup> represents a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom, the heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl, carboxyl, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, —NR<sup>6</sup>R<sup>7</sup>, —(CH<sub>2</sub>)<sub>n</sub>NR<sup>6</sup>R<sup>7</sup> and CONR<sup>6</sup>R<sup>7</sup>, r is 1, 2, 3, 4, 5 or 6; R<sup>5</sup> represents a hydrogen atom or a C<sub>1</sub>-C<sub>6</sub>-alkyl or C<sub>3</sub>-C<sub>8</sub>-cycloalkyl group; R<sup>6</sup> and R<sup>7</sup> each independently represent a hydrogen atom or a C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-hydroxyalkyl or C<sub>3</sub>-C<sub>8</sub>-cycloalkyl group, or R<sup>6</sup> and R<sup>7</sup> together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring; with the provisos that, (a) when A represents C(O)NH and R<sup>4</sup> represents an unsubstituted 3- to 8-membered saturated aliphatic heterocyclic ring system containing one nitrogen atom, then X is other than a bond, and (b) when A represents C(O)NH and X represents a group (CH<sub>2</sub>)<sub>1-6</sub> or O(CH<sub>2</sub>)<sub>1-6</sub>, then R<sup>4</sup> does not represent an unsubstituted imidazolyl, unsubstituted morpholinyl, unsubstituted piperidinyl or unsubstituted pyrrolidinyl group, and (c) when A represents NHC(O) and R<sup>4</sup> represents an unsubstituted 3- to 8-membered saturated aliphatic heterocyclic ring system containing one nitrogen atom, then X is other than a bond, and (d) when A represents NHC(O) and X represents O(CH<sub>2</sub>)<sub>1-6</sub>, NH(CH<sub>2</sub>)<sub>1-6</sub> or SCH<sub>2</sub>, then R<sup>4</sup> does not represent an unsubstituted 1-piperidinyl or unsubstituted 1-pyrrolidinyl group, and (e) when A represents NHC(O) and X represents O(CH<sub>2</sub>)<sub>2-3</sub>NH(CH<sub>2</sub>)<sub>2</sub>, then R<sup>4</sup> does not represent an imidazolyl group; or a pharmaceutically acceptable salt or solvate thereof.


[0239] A compound of general formula:




[0240] X represents a nitrogen atom or a group C(R<sup>5</sup>), Y represents an oxygen or sulphur atom or a group NR<sup>6</sup>; either R<sup>1</sup> and R<sup>2</sup> each independently represent a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub>-alkyl group but do not both simultaneously represent a hydrogen atom, or R<sup>1</sup> and R<sup>2</sup> together represent a group —CH<sub>2</sub>ZCH<sub>2</sub>—; Z represents a bond, an oxygen or sulphur atom or a group CH<sub>2</sub> or NR<sup>7</sup>; m is 0 or 1; R<sup>3</sup> represents a 5- to 10-membered unsaturated ring system which may comprise from 1 to 4 ring heteroatoms independently selected from nitrogen, oxygen and sulphur, the ring system being optionally substituted by one or more substituents independently selected from halogen, nitro, cyano, NR<sup>8</sup>R<sup>9</sup>, C<sub>1</sub>-C<sub>4</sub>-alkyl-C(O)NH—, NHR<sup>12</sup>C(O)—, C<sub>1</sub>-C<sub>4</sub>-alkyl-SO<sub>2</sub>—, C<sub>1</sub>-C<sub>4</sub>-alkyl-SO<sub>2</sub>NH—, C<sub>1</sub>-C<sub>4</sub>-alkyl-NHSO<sub>2</sub>—, C<sub>1</sub>-C<sub>4</sub>-alkoxy, and C<sub>1</sub>-C<sub>4</sub>-alkyl optionally substituted by one or more fluorine atoms; R<sup>4</sup> represents a phenyl or pyridinyl group, each of which is substituted in an ortho position with a substituent selected from halogen, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, and C<sub>1</sub>-C<sub>4</sub>-alkyl optionally substituted by one or more fluorine atoms, the phenyl or pyridinyl group being optionally further substituted by one or more substituents independently selected from halogen, cyano, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkyl-NH—, NHR<sup>13</sup>-C<sub>1</sub>-C<sub>4</sub>-alkyl-, C<sub>1</sub>-C<sub>4</sub>-alkyl-SO<sub>2</sub>—, C<sub>1</sub>-C<sub>4</sub>-alkyl-SO<sub>2</sub>NH—, C<sub>1</sub>-C<sub>4</sub>-alkyl-NHSO<sub>2</sub>—, C<sub>1</sub>-C<sub>4</sub>-alkyl-C(O)NH—, C<sub>1</sub>-C<sub>4</sub>-alkyl-NHC(O)—, -D-G, C<sub>1</sub>-C<sub>4</sub>-alkoxy optionally substituted by —NR<sup>14</sup>R<sup>15</sup> or by R<sup>16</sup>, and C<sub>1</sub>-C<sub>4</sub>-alkyl optionally substituted by one or more fluorine atoms or by one or more hydroxyl groups, or R<sup>4</sup> represents a 9- or 10-membered unsaturated bicyclic ring system which may comprise from 1 to 4 ring heteroatoms independently selected from nitrogen, oxygen and sulphur, the bicyclic ring system being optionally substituted by one or more substituents independently selected from halogen, oxo, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio and —NR<sup>10</sup>R<sup>11</sup>; D represents an oxygen atom or a group (CH<sub>2</sub>)<sub>n</sub> or CH<sub>2</sub>NH; n is 1, 2 or 3; G represents a piperazinyl, morpholinyl or 2,5-diazabicyclo[2.2.1]heptyl group, or G represents a piperidinyl group optionally substituted by amino; R<sup>5</sup> represents a hydrogen atom, or a hydroxyl or C<sub>1</sub>-C<sub>4</sub>-alkoxy group; R<sup>6</sup> represents a hydrogen atom, or a cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-alkoxy group; R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> each independently represent a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub>-alkyl group; R<sup>10</sup> and R<sup>11</sup> each independently represent a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub>-alkyl group, or R<sup>10</sup> and R<sup>11</sup> together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated heterocyclic ring comprising one or two ring nitrogen atoms; R<sup>12</sup> represents a hydrogen atom, or a C<sub>1</sub>-C<sub>4</sub>-alkyl group optionally substituted by amino; R<sup>13</sup> represents a hydrogen atom, or a C<sub>1</sub>-C<sub>4</sub>-alkyl group optionally substituted by hydroxyl; R<sup>14</sup> and R<sup>15</sup> each independently represent a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub>-alkyl group optionally substituted by hydroxyl, or R<sup>14</sup> and R<sup>15</sup> together

with the nitrogen atom to which they are attached form a 5- or 6-membered saturated heterocyclic ring comprising one or two ring nitrogen atoms; and  $R^{16}$  represents a 1-( $C_1$ - $C_4$ -alkyl)-piperidinyl group; with the proviso that when  $m$  is 0,  $X$  is N and  $Y$  is O, then  $R^4$  does not represent 2-benzothiazolyl; or a pharmaceutically acceptable salt or solvate thereof.

[0241] A compound of general formula:




[0242] wherein: each  $R^1$  independently represents a hydrogen or halogen atom, or a trifluoromethyl, cyano, nitro,  $C_1$ - $C_6$ -alkyl or  $C_1$ - $C_6$ -alkoxy group;  $T$  represents an oxygen atom or a group  $NH$ ;  $U$  represents an oxygen or sulphur atom or a group  $NH$ ;  $Ar$  represents a group:



[0243]  $X$  represents a bond, an oxygen atom or a group  $CO$ ,  $CH_2$ ,  $CH_2O$ ,  $O(CH_2)_m$ ,  $CH_2OCH_2$ ,  $NR^5$ ,  $CH_2NR^5$ ,  $NR^5CH_2$ ,  $CH_2NR^5CH_2$ ,  $CONR^5$ ,  $S(O)_n$  or  $SO_2NR^5$ ,  $m$  is 1, 2 or 3;  $n$  is 0, 1 or 2; one of  $R^2$  and  $R^3$  represents a halogen, cyano, nitro, amino, hydroxyl, or a group selected from  $C_1$ - $C_6$ -alkyl optionally substituted by at least one  $C_3$ - $C_6$ -cycloalkyl,  $C_3$ - $C_8$ -cycloalkyl,  $C_1$ - $C_6$ -alkyloxy optionally substituted by at least one  $C_3$ - $C_6$ -cycloalkyl,  $C_3$ - $C_8$ -cycloalkyloxy,  $S(O)_p$ - $C_1$ - $C_6$ -alkyl or  $S(O)_q$ - $C_3$ - $C_8$ -cycloalkyl, each of these groups being optionally substituted by one or more fluorine atoms, and the other of  $R^2$  and  $R^3$  represents a hydrogen or halogen atom or a methyl group;  $p$  is 0, 1 or 2;  $q$  is 0, 1 or 2;  $R^4$  represents  $di(C_{1-2}alkyl)N(CH_2)_t$ , where  $t$  is 0, 1 or 2 or imidazolyl, or  $R^4$  represents a 3- to 9-membered saturated heterocyclic ring system containing one or two nitrogen atoms, the heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl,  $C_1$ - $C_6$ -alkyl, acetyl, hydroxy- $C_1$ - $C_6$ -alkyl,  $-NR^6R^7$ ,  $-(CH_2)_rNR^6R^7$ ,  $-CONR^6R^7$  and pyrimidinyl, or  $R^4$  represents a 3- to 8-membered saturated carbocyclic ring system substituted by one or more substituents independently selected from  $-NR^6R^7$ ,  $-(CH_2)_rNR^6R^7$ ,  $-CONR^6R^7$  the ring system being optionally further substituted by one or more substituents independently selected from fluorine atoms, hydroxyl and  $C_1$ - $C_6$ -alkyl;  $r$  is 1, 2, 3, 4, 5 or 6;  $R^5$  represents a hydrogen atom or a  $C_1$ - $C_6$ -alkyl or  $C_3$ - $C_8$ -

cycloalkyl group; and  $R^6$  and  $R^7$  each independently represent a hydrogen atom or a  $C_1$ - $C_6$ -alkyl or  $C_3$ - $C_8$ -cycloalkyl group, or  $R^6$  and  $R^7$  together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring, provided that when  $R^3$  represents a cyano group, then  $X$  is other than a bond; or a pharmaceutically acceptable salt or solvate thereof.

[0244] A compound of general formula:




[0245] wherein  $X$  represents an oxygen or sulphur atom or a group  $NH$ ,  $CH_2$ ,  $CH_2CH_2$  or  $OCH_2$ ;  $Y$  represents a group  $CH_2$  or  $C=O$ ;  $R^1$  represents a pyridyl or pyrimidinyl group;  $R^2$  represents a phenyl, pyridyl or pyrimidinyl group, each of which may be optionally substituted by one or more substituents independently selected from a halogen atom or an amino, cyano, hydroxyl, nitro,  $C_1$ - $C_6$ -alkyl, halo- $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -alkylthio, (di) $C_1$ - $C_6$ -alkylamino,  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkoxycarbonyl,  $C_1$ - $C_6$ -alkylsulphinyl,  $C_1$ - $C_6$ -alkylsulphonyl,  $-NR^3SO_2R^4$  or  $-SO_2NR^3R^6$  group, or a group  $-Z-(CH_2)_p-Z-(CH_2)_q-H$  wherein each  $Z$  independently represents a nitrogen or oxygen atom,  $p$  is an integer from 2 to 5 and  $q$  is 0 or an integer from 1 to 5;  $R^3$  and  $R^4$  each independently represent a hydrogen atom or a  $C_1$ - $C_6$ -alkyl group; and  $R^5$  and  $R^6$  each independently represent a hydrogen atom or a  $C_1$ - $C_6$ -alkyl group, or together with the nitrogen atom to which they are attached form a pyrrolidinyl or piperidinyl group; or a pharmaceutically acceptable salt or solvate thereof.

[0246] An additional embodiment of the invention provides a method for treating affective disorders by administering an agent modulating the activity of P2X7R, such as an antagonist of the P2X7R. The term "antagonist" means an agent or drug or a compound that opposes the physiological effects of another. Examples of P2X7R antagonists include, but are not restricted to, adamantane derivatives, isoquinolines and their derivatives, substituted phenyl compounds, piperidine derivatives, piperazine derivatives. P2X7R antagonists are described in the art and include the compounds found in Chen et al., Bioconjugate Chem. 13 (2002), 1100-1111; WO 99/29660; WO 99/29661; WO 99/296896; WO 00/61569; WO 01/42194; WO 01/44170; WO 01/44213; WO 00/71529; WO 01/46200. P2X7R activity can also be modulated by RNA-based interference mechanisms and methods such as, but not limited to, small interference RNA (siRNA) molecules, and long double-stranded RNA (dsRNA).

[0247] Since it was unexpectedly found that agonists of P2X7R improve the symptoms of mice selected for anxiety and depressive behaviour, the present invention relates in a further embodiment to a method for treating affective disorders such as anxiety and depressive behaviour by administering an agent modulating the activity of P2X7R such as an agonist of the P2X7R. What is even more striking is the

finding of the present application shown in Example 10 that antagonists of P2X7R have no antidepressive effects although this is being taught by the prior art, for example, WO 03/042190, WO 03/042191, WO 03/049353 or US 2004/0029841. WO 03/042190, WO 03/042191, WO 03/059353 or US 2004/0029841 describe compositions of P2X7R antagonists and methods of treating P2X7 mediated diseases by administering these compounds. However, while the prior art may make a link between P2X7R and, e.g., treating depression, it is generally implied that antagonists of P2X7R have to be used in the treatment of, e.g., depression. Thus, the finding of the present invention that antagonists have no antidepressive effects, but rather agonists of P2X7R have an antidepressive effect could not have been expected and is, thus, even more surprising. The present application, therefore, provides, inter alia, the basis for the development of effective medicaments having therapeutic benefits against, e.g., depression. In particular, said medicaments comprising an agonist of P2X7R are effective in treating affective disorders, in particular for treating those disorders mentioned herein, and in particular for treating depression. Examples of P2X7R agonists include but are not restricted to ATP, ATP-4, and BZATP (2'-3'-O-(4-Benzoyl-benzoyl)adenosine 5'-triphosphate ( $C_{24}H_{24}N_5O_{15}P_3$ )). Preferably, the P2X7R agonist to be used for treating affective disorders is BZATP. More preferably, BZATP is used to treat depression or anxiety as demonstrated in Example 9 hereinbelow.

[0248] The present application provides another unexpected finding in that the chemical compound called tenidap or a derivative thereof also functions as a modulator of P2X7R which can thus be used for treating affective disorders. So far, various medical applications are described for tenidap or a derivative thereof. However, the use of tenidap or a derivative thereof for treating affective disorders is neither known nor suggested in the prior art. Accordingly, the present application relates to the use of tenidap or a derivative thereof or 3-substituted-2-oxindole-1-carboxamides for the preparation of a pharmaceutical composition for treating an affective disorder. Of course, also a method of treatment of an affective disorder comprising administering a therapeutically effective amount of tenidap or a derivative thereof or 3-substituted-2-oxindole-1-carboxamides to a subject suffering from said disorder is envisaged. The composition of tenidap (5-chloro-2,3-dihydro-2-oxo-3-(2-thienylcarbonyl)-indole-1-carboxamide, i.e.  $C_{15}H_{11}ClN_2O_2S$ ) having the following structural formula



[0249] and other 3-substituted-2-oxindole-1-carboxamides and their use as anti-inflammatory and analgesic agents,

and as inhibitors of both the cyclooxygenase (Cox) and lipoxygenase (5-LPO) enzymes was first disclosed in U.S. Pat. No. 4,556,672. Of course, various modifications of, e.g., side groups or atoms which are well known in the art can be made to the composition of tenidap.

[0250] Derivatives of tenidap or 3-substituted-2-oxindole derivatives are described, for example, in U.S. Pat. No. 4,556,672; U.S. Pat. No. 4,658,037; U.S. Pat. No. 4,721,712; US 5,290,802; U.S. Pat. No. 5,118,703; U.S. Pat. No. 5,270,331; U.S. Pat. No. 5,298,522; U.S. Pat. No. 5,086,186; U.S. Pat. No. 5,449,788 and U.S. Pat. No. 5,795,902. Various processes for the synthesis of tenidap and other 3-substituted-2-oxindole-1-carboxamides are well known in the art, see for example U.S. Pat. No. 4,652,658; U.S. Pat. No. 4,665,194; U.S. Pat. No. 4,952,703; EP-B1 155 828, WO 90/04393, WO 94/07488, WO 94/17061, WO 95/20574; WO 97/36895; van Deurzen et al., J. Mol. Catal. B-Enzym., 2 (1996), 33-42; Porcs-Makkav and Simici, Org. Process. Res. Dev., 4 (2000), 10-16; Kumar et al., Org. Process. Res. Dev., 5 (2001), 61-64. The anhydrous crystalline form of the sodium salt of tenidap is described in U.S. Pat. No. 5,036,099 and WO 88/05656. Injectable composition and pharmaceutical composition for rectal administration of tenidap are mentioned in EP-B1 508 311 and EP-B1 508 310, respectively. It is also described in the prior art that tenidap or a derivative thereof can be administered in combination with tetracycline (U.S. Pat. No. 5,308,839) or methotrexate (WO 96/35419) for the treatment of rheumatoid arthritis. Inhibition of the photodecomposition of tenidap and other 3-substituted-2-oxindole-1-carboxamides is disclosed in WO 96/33701.

[0251] Further applications of tenidap or a derivative thereof and other 3-substituted-2-oxindole-1-carboxamides have been described for the inhibition of interleukin-1 biosynthesis in mammals and for the treatment interleukin-1 mediated disorders (U.S. Pat. No. 4,861,794); for the inhibition of elastase release from neutrophils (U.S. Pat. No. 5,006,547); for the suppression of T-cell function in mammals and to treat T-cell mediated autoimmune disorders of the systemic or organ specific type (U.S. Pat. No. 4,853,409; Dolhain et al., Scand. J. Immunol. 42(1995), 686-693). Tenidap is also used for the treatment of Alzheimer's disease (WO 96/31209; U.S. Pat. No. 5,545,656).

[0252] Tenidap or a derivative thereof or its pharmaceutically base salts have also been shown to inhibit activation of collagenase, treat collagenase mediated disorders and diseases, and inhibit the activity of myeloperoxidase in mammals (U.S. Pat. No. 5,008,283). Tenidap can reduce total serum cholesterol, LDL cholesterol and triglycerides (U.S. Pat. No. 5,122,534), and can be used for the treatment of ischemia induced myocardial injury and cytokine mediated myocardial injury (EP-B1 679 396). However, none of the aforementioned documents discloses a use of tenidap or derivatives thereof or 3-substituted-2-oxindole-1-carboxamides thereof for treating, for example, affective disorders such as depression.

[0253] Sanz et al., Eur. J. Pharmacol. 355 (1998), 235-244 suggest that tenidap can enhance the activity of the P2X7 receptor. It is suggested that tenidap may act by increasing ATP levels or improving the effect of ATP on P2X7. ATP is the natural ligand of P2X7R. Accordingly, tenidap or a derivative thereof is a modulator of P2X7R as is described

herein since a modulator is defined as either directly or indirectly modulating the activity or expression of P2X7R. By making use of the teaching of the present invention that modulators of P2X7R are useful for treating affective disorders, it is envisaged that tenidap or a derivative thereof is used as a modulator of P2X7R activity for the preparation of a pharmaceutical composition for the treatment of an affective disorder, examples of which are described herein. The preparation of pharmaceutical compositions, the modes of administration etc. are described supra and infra and apply to the use of tenidap or a derivative thereof for the preparation of a pharmaceutical composition, *mutatis mutandis*. Moreover, also the embodiments relating to the uses of and methods for treating affective disorders described herein apply to the use of tenidap or a derivative thereof for treating affective disorders or the corresponding method of treatment, *mutatis mutandis*.

**[0254]** The present application moreover envisages that modulators of P2X7R activity can be used in any combinations thereof for treating an affective disorder. For example, BZATP and tenidap or a derivative thereof or 3-substituted-2-oxindole-1-carboxamides may be used together, e.g., simultaneously or by successive administration for treating an affective disorder.

**[0255]** In a preferred embodiment the pharmaceutical composition described herein optionally comprises further molecules which have cell protective properties capable of altering the characteristics of the components of the invention thereby, for example, modulating, preferably blocking possible undesired, adverse or negative side effects of these components. One such possible undesired, adverse or negative side effect is the formation of pores in the cell membrane of treated cells which ultimately leads to apoptosis. Accordingly, said further molecules belong to the class of beta-adrenergic receptor modulators including agonists or antagonists having membrane-stabilizing properties. Beta-adrenergic receptor modulators including agonists and antagonists are compounds which decrease or increase the positive chronotropic, positive inotropic, bronchodilator and vasodilator responses caused by beta-adrenergic receptor agonists or antagonists. The magnitude of this decreased or increased response is proportional to the existing sympathetic tone and the concentration of beta-adrenergic receptor blocking agent which reaches the receptor sites. A beta-adrenergic receptor modulator in the context of the present invention is thus an antagonist or agonist. The activity of a beta-adrenergic receptor antagonist or agonist can be determined as is well known in the art. The activity of beta-adrenergic receptors can be determined by measuring the accumulation of cyclic adenosine mono-phosphate (cAMP) in Chinese hamster ovary (CHO) cells. CHO cells can be uniquely transfected with the cDNA coding for the human beta1-, beta2-, or beta3-adrenergic receptor under the control of the CMV promoter or any other suitable promoter element. Transfection of the cells is performed using standard cell transfection methods, see for example, Joyner, "Gene Targeting: A Practical Approach", Oxford University Press, New York, 1993. Cells overexpressing one of the beta-adrenergic gene are then grown to confluence in Ham's F12 media (Gibco BRL) containing 10% fetal bovine serum, 500 mg/ml Geneticin, 100 U/ml penicillin, 100 mg/ml streptomycin and 250 ng/ml fungizone according to the procedure described in American Type Culture Collection Catalogue of Cell Lines and Hybridomas, Seventh Edition,

1992, p. 36, ATCC CCL 61 CHO-K1. Beta-adrenergic modulator compounds can be prepared as 10 mM stock solutions in DMSO (0.1% DMSO, final concentration), diluted in Ham's F12 media and added to the cells at  $10^{-10}$  to  $10^{-5}$  M along with  $10^{-3}$  M isobutylmethylxanthine to inhibit phosphodiesterase activity. The media and cells are then incubated for 5 minutes at 37° C. At the end of this period, the media is aspirated and the cells lysed in 0.01N HCl. The cellular content of cAMP can then be determined by radioimmunoassay (RIA) using a kit from New England Nuclear (Burlington, Mass.). There is a direct correlation between the cellular content of cAMP and the activation/inhibition of the beta-adrenergic receptor. Other methods for determining the activity of a beta-adrenergic receptors are well described in the art, see for example, Vansal and Feller, J. Recept. Signal. Transduct. Res. 19 (1999) 853-863; Durocher et al., Anal. Biochem. 284 (2000) 316-326.

**[0256]** Examples of compounds which fit the definition of a beta-adrenergic receptor modulating agent include but are not limited to known beta-adrenergic receptor antagonists such as timolol, sotalol, esmolol, catechol, propranolol, betaxolol, penbutolol, metoprolol, acebutolol, atenolol, metoprolol, pindolol, and bisoprolol, and their salts, hydrates, solvates and any crystal forms in which they may occur. Further examples of beta-adrenergic receptor blocking agents are described in U.S. Pat. No. 5,776,930. Preferred examples of beta-adrenergic receptor antagonists are DL-propranolol, D-propranolol and labetalol. DL-propranolol and labetalol are beta-adrenergic receptor antagonists with membrane-stabilizing properties, while D-propranolol is an optical isomer with poor beta-adrenergic blocking activity. The optional addition of beta-adrenergic receptor antagonists or agonists to the pharmaceutical composition of the present invention for treating an affective disorder may be useful in the context of administering P2X7R agonists. This is because the prior art suggests that the activation of P2X7R by agonists may result in cell death by triggering the formation of pores within the cell membrane. However, as is demonstrated in Example 9 of the present application the pore-forming activity effect described for P2X7R agonists by the prior art was only observed for very few cells. Recently, Alzola et al., Cell Signal. 13 (2001), 465-473 have shown that concentrations of 10 to 300  $\mu$ M DL-propranolol, D-propranolol or labetalol can inhibit the pore-forming activity of P2X7R without affecting the opening of the cation channel activity of P2X7R. From FR 2768626 it is also known that beta-adrenergic modulators, e.g. agonists are useful as apoptosis inhibiting agents. Accordingly, in a preferred embodiment of the present invention, beta-adrenergic receptor modulators including antagonists or agonists are administered in combination with P2X7R agonists for the treatment of affective disorders. Said beta-adrenergic receptor antagonists or agonists are preferably administered in a concentration of 10 to 300  $\mu$ M.

**[0257]** Dosage, pharmaceutical preparation and delivery of P2X7R modulating agent for use in accordance with the present invention may be formulated in conventional manner according to methods found in the art, using one or more physiological carriers or excipient, see, for example Ansel et al., "Pharmaceutical Dosage Forms and Drug Delivery Systems", 7th edition, Lippincott Williams & Wilkins Publishers, 1999. Thus, the P2X7R modulating agent and its physiologically acceptable salts and solvates may be formulated

for administration by inhalation, insufflation (either through the mouth, or nose), oral, buccal, parenteral, or rectal administration.

[0258] For oral administration, the pharmaceutical composition of the P2X7 modulating agent may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutical acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone, hydroxypropyl methylcellulose), fillers (e.g., lactose, microcrystalline cellulose, calcium hydrogen phosphate), lubricants (e.g., magnesium stearate, talc, silica), disintegrants (e.g., potato starch, sodium starch glycolate), or wetting agents (e.g., sodium lauryl sulphate). Liquid preparations for oral administration may take the form of, for example, solutions, syrups, or suspensions, or may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparation may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol, syrup, cellulose derivatives, hydrogenated edible fats), emulsifying agents (e.g., lecithin, acacia), non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol, fractionated vegetable oils), preservatives (e.g., methyl or propyl-p-hydroxycarbonates, sorbic acids). The preparations may also contain buffer salts, flavouring, coloring and sweetening agents as deemed appropriate. Preparations for oral administration may be suitably formulated to give controlled release of the agent modulating P2X7R activity.

[0259] For administration by inhalation, the agent modulating P2X7R activity for use according to the present invention is conveniently delivered in the form of an aerosol spray presentation from a pressurised pack or a nebulizer, with the use of a suitable propellant (e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas). In the case of a pressurised aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, for example, gelatine, for use in an inhaler or insufflator may be formulated containing a powder mix of the P2X7R activity modulating agent and a suitable powder base such as lactose or starch.

[0260] An agent modulating P2X7R activity may be formulated for parenteral administration by injection, for example, by bolus injection or continuous infusion. Site of injections include intravenous, intra-peritoneal or sub-cutaneous. Formulations for injection may be presented in units dosage form (e.g., in phial, in multi-dose container), and with an added preservative. The agent modulating P2X7R activity may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing, or dispersing agents. Alternatively, the agent may be in powder form for constitution with a suitable vehicle (e.g., sterile pyrogen-free water) before use.

[0261] An agent modulating P2X7R activity may, if desired, be presented in a pack, or dispenser device which may contain one or more unit dosage forms containing the said agent. The pack may for example comprise metal or plastic foil, such as blister pack. The pack or dispenser device may be accompanied with instruction for administration.

[0262] In a more preferred embodiment the aforementioned methods or uses are envisaged to treat affective disorders selected from the group consisting of major depression, generalized anxiety disorder and bipolar disorder.

[0263] In a particularly preferred embodiment said major depression is selected from the group consisting of major depression, dysthymia, atypical depression, premenstrual dysphoric disorder and seasonal affective disorder.

[0264] In another particularly preferred embodiment said generalized anxiety disorder is selected from the group consisting of panic disorder, phobias, agoraphobia, social phobia, specific phobia, obsessive-compulsive disorder, post-traumatic stress disorder, separation anxiety disorder, mania, hypomania and cyclothymic disorder.

[0265] A still also particularly preferred embodiment is that said bipolar disorder is bipolar disorder type I or bipolar disorder type II.

[0266] Additionally, the present invention relates to a kit comprising the nucleic acid molecule, the vector, the host, the polypeptide, the antibody or the aptamer, the primer or pair of primers of the invention or the molecule as identified or characterized in a method herein below of the present invention.

[0267] Advantageously, the kit of the present invention further comprises, optionally (a) reaction buffer(s), storage solutions and/or remaining reagents or materials required for the conduct of scientific or diagnostic assays or the like. Furthermore, parts of the kit of the invention can be packaged individually in vials or bottles or in combination in containers or multicontainer units.

[0268] The kit of the present invention may be advantageously used, inter alia, for carrying out the method of producing a polypeptide of the invention, the method(s) of identification and/or characterization of molecules specifically interacting with P2X7R ATP-gated ion channels as described herein below and/or it could be employed in a variety of applications referred herein, e.g., as diagnostic kits, as research tools or therapeutic tools. Additionally, the kit of the invention may contain means for detection suitable for scientific, medical and/or diagnostic purposes. The manufacture of the kits follows preferably standard procedures which are known to the person skilled in the art.

[0269] Furthermore, the present invention relates to a method for identifying compounds or mixtures of compounds which are capable of specifically interacting with a polypeptide of the present invention, comprising the steps of (a) contacting a polypeptide of the present invention with a candidate compound or mixture of compounds to be tested; and (b) determining whether said is capable of specifically interacting with said polypeptide. The polypeptide may be provided directly or by expression of a corresponding nucleic acid molecule or vector of the invention, e.g., in vitro or in a suitable host cell.

[0270] Additionally, the present invention relates to a method for the characterization of compounds which are capable of altering characteristics of the polypeptides of the present invention, comprising the steps of (a) contacting a polypeptide of the invention with said compound; and (b) determining whether the compound alters a characteristic of said polypeptide.

[0271] The term "altering characteristic of the polypeptide of the present invention" means that the functional characteristics to the polypeptides of the present invention in comparison to functional characteristics which they had before being contacted with the compounds identified by the above-described method: as described hereinabove are altered; i.e. changed.

[0272] Said identification and/or characterization of which are capable of interacting with or altering characteristics of the polypeptide of this invention, may be, inter alia, achieved by transfecting an appropriate host with a nucleic acid molecule of invention. Said hosts comprise, but are not limited to, HEK 293 cells or are injected into frog oocytes, preferably a *Xenopus* oocyte for functional expression (Goldin, *Methods Enzymol.* 207 (1992), 266). Expressed P2X7R ATP-gated channels can be examined using standard two-electrode voltage clamp techniques (Stuhmer, *Methods Enzymol.* 207 (1992), 319; Kohler, *Science* 273 (1996), 1709). After expression of a P2X7R ATP-gated ion channel as defined herein, membrane currents may be deduced in the absence and/or presence of the molecule to be identified and/or characterized. Methods for the deduction of membrane currents are well known in the art and comprise, e.g., patch clamp methods as described in Hamill, *Pflügers Arch.* 391 (1981), 85-100 or two-electrode voltage clamp in oocytes, as described in Methfessel, *Pflügers Archive* 407 (1986), 577-588. In accordance with the present invention the term "interacting with the polypeptides of the present invention" means that the polypeptides of the present invention interact directly and/or indirectly with compounds identified by the method described above. Furthermore, the present invention relates to a method of screening for molecules which are capable of interacting with the polypeptide of this invention, comprising the steps of (a) contacting a polypeptide of the invention with a molecule; and (b) measuring and/or detecting a response; and (c) comparing said response to a standard response as measured in the absence of said candidate molecule.

[0273] The present invention also relates to a method for identifying a compound which is capable of enhancing or reducing the expression of the P2X7R gene comprising the steps of contacting a cell which expresses the P2X7R gene from its natural promoter or a reporter gene driven by the P2X7R promoter and determining whether the expression of the gene is increased or reduced when compared to conditions in which the compound is not present.

[0274] Potential candidate molecules or candidate mixtures of molecules may be, inter alia, substances, compounds or compositions which are of chemical or biological origin, which are naturally occurring and/or which are synthetically, recombinantly and/or chemically produced or compounds or compositions described hereinabove. Thus, candidate molecules may be proteins, protein-fragments, peptides, amino acids and/or derivatives thereof or other compounds, such as ions, which bind to and/or interact with wild-type P2X7R ATP-gated ion channels. Such binding and/or interacting candidate compounds may be found employing, inter alia, yeast two-hybrid systems or modified yeast two-hybrid systems as described, for example in Fields, *Nature* 340 (1989), 245-246; Gyuris, *Cell* 75 (1993), 791-801; or Zervos, *Cell* 72 (1993), 223-232.

[0275] Furthermore, potential candidate molecules may be contacted with a cell, such as an oocyte or a HEK 293 cell,

which expresses a polypeptide of the invention or with a membrane patch comprising a polypeptide of the invention and a corresponding response (inter alia, a dose-response response, a current-response, or single current channel response) may be measured in order to elucidate any effect said candidate molecule causes.

[0276] Within the scope of the present invention are also methods for identifying, characterizing and for screening of molecules which are capable of interacting with the P2X7R ATP-gated ion channels according to the invention which comprise so-called high-throughput screening methods and similar approaches which are known in the art (Spencer, *Biotechnol. Bioeng.* 61 (1998), 61-67; Oldenburg, *Annu. Rev. Med. Chem.* 33 (1998), 301-311) carried out using 96-well, 384-well, 1536-well (and other) commercially available plates. Further methods to be employed in accordance with the present invention comprise, but are not limited to, homogenous fluorescence readouts in high-throughput screenings (as described, inter alia, in Pope, *Drug Discovery Today* 4 (1999), 350-362). The method of the present invention for identification, characterization and/or screening of molecules capable of interacting with P2X7R ATP-gated ion channels can, inter alia, employ hosts as defined herein which express the polypeptide of the present invention. Cell-based assays, instrumentation for said assays and/or measurements are well-known in the art and described, inter alia, in Gonzalez, *Drug Discovery Today* 4 (1999), 431-439 or Ramm, *Drug Discovery Today* 4 (1999), 401-410. It is also envisaged that the high throughput screens described herein are conducted by using, for example cRNA, i.e. synthetic RNA from a cDNA construct) that can be introduced in host cells, such as *Xenopus* oocytes using routine methods in the art. As an example, direct nucleic acid injection can be employed, such as the Eppendorf microinjection system (Micromanipulator 5171 and Transjector 5242). The injected/transformed cells can be analyzed for ion currents about 4 hours later using patch-clamp techniques which are commonly practiced in the art.

[0277] Additionally, the present invention relates to a method for the production of a pharmaceutical composition comprising the steps of a method of the invention for identifying, characterizing and/or screening of molecules which are capable of interacting with and/or altering the characteristics of a P2X7R ATP-gated ion channel of the invention and further comprising a step, wherein a derivative of said identified, characterized and/or screened molecule is generated. Such a derivative may be generated by, inter alia, peptidomimetics.

[0278] The invention furthermore relates to a method for the production of a pharmaceutical composition comprising the steps of a method of the invention for identifying, characterizing, screening and/or derivatizing of molecules which are capable of interacting with and/or altering the characteristics of a P2X7R ATP-gated ion channel and formulating the molecules identified, characterized, screened and/or derivatized in pharmaceutically acceptable form.

[0279] In a more preferred embodiment the present invention relates to a method wherein said molecule(s) comprise(s) (a) neuroprotective, (a) nootropic and/or (a) anti-epileptic molecule(s).

[0280] Yet another embodiment of the invention is the use of a P2X7R polypeptide, in particular those according to the

present invention, to identify biological, chemical, or pharmacological agents that can have an antidepressive effect. The term 'agent' refers to a chemical compound or composition capable of inducing a desired therapeutic or prophylactic effect when properly administered to a subject or cell. For example, the present invention allows the generation of cells expressing P2X7R for the identification and characterization of agents which modulate ionic influx and efflux. For example, HEK293 cells, or other cell lines (e.g., HCN-1A, HCN-2, HIT-T15, RIN-m5F, betaTC3, PC12, HT22, SH-SY5Y, Neuro2A or CA77), can be stably transfected with cDNA encoding the human P2X7R and plated in 12, 96 and 384 well plates. Said cells are cultured in appropriate medium. Examples of such medium are well known in the art, see, for example Freshney, "Culture of Animal Cells: A Manual of Basic Technique, 4th edition, Wiley-Liss Publishing, 2000.

[0281] Said cells can then be pre-incubated with said agents for 15 min prior to stimulation with 3 mM ATP for 10 minutes. Reactions are then terminated by rapid aspiration of the extracellular medium in each well. The cells in each well are subsequently extracted overnight with 1 ml 10% HNO<sub>3</sub>. Potassium (K<sup>+</sup>) content in the extracts can be determined by atomic absorbance spectrophotometry. Agent function is then measured by the percent inhibition or stimulation of the K<sup>+</sup> release triggered by 3 mM ATP and compared to K<sup>+</sup> release in the absence of the agents. P2X7R activity can also be monitored according to the movement of calcium (Ca<sup>2+</sup>; see Denver et al., *Drug Discov. Today* 7 (1998), 323-332; González et al., *Drug Discov. Today* 9 (1999), 431-439; Helmchen and Waters, *Eur. J. Pharmacol.* 447 (2002), 119-129). Agents can also be verified in the absence of ATP.

[0282] P2X7R activity can also be monitored according to secretion of neurotransmitters such as glutamate and GABA. Neurotransmitter levels in treated cells can be quantified by suitable methods, e.g., Enzyme Linked Immunoabsorbent Assay (ELISA), Radio Immuno Assay (RIA), High Performance Liquid Chromatography (HPLC). Using these methods, a large number of compounds can be screened for increase in neurotransmitter (for example, glutamate) secretion. The release of glutamate can be measured for example by Fluorometric glutamate release assays (e.g., Amplex Red Glutamic Acid/Glutamate Oxidase Assay Kit, Molecular Probes) or High-Throughput ElectroPhysiology.

[0283] In a further aspect the present invention uses the P2X7R polypeptides disclosed herein or polypeptides of the present invention in a method for identifying compounds or agents having agonist activity to said P2X7R polypeptides or to the polypeptides of the present invention. Agents and compounds are defined and described hereinabove and hereinbelow. In particular, cells that express the P2X7 gene are contacted with candidate agents, molecules or compounds as described hereinabove and either calcium influx or ethidium bromide entry is measured by methods known in the art, described hereinabove and in particular described in Example 8 hereinbelow. The cells used in the method for identifying agonists to P2X7R are preferably cells of a hippocampal cell line. Hippocampal cell lines are prepared by methods known in the art, for example, described in EP 0 773 287 or EP 0 773 292. Non limiting examples of hippocampal cell lines are rat H19-7 hippocampal cells (ATCC-2526) described in Eves et al. *Proc. Natl. Acad. Sci.*

USA 89 (1992), 4373-4377, mouse HN9.10 hippocampal cells described in Lee et al. *J. Neurosci.* 10 (1992), 1779-1787 or rat Hi5B hippocampal cells described in Renfranz et al., *Cell* 66 (1991), 713-729. Preferably, the hippocampal cells used in accordance with the aforementioned method are cells of the HT-series (see Davis and Maher (1994), *Brain Res.* 652, 169-173), Morimoto and Koshland (1990), *Neuron* 5, 875-880). It is also preferred that the hippocampal cells express the endogenous P2X7R gene. However, it is also envisaged that such cells may be genetically modified by introducing an exogenous P2X7R gene using methods commonly known in the art. More preferably, HT-22 cells are used for identifying agonists to the P2X7R polypeptides described herein or to the polypeptides of the present invention and HT-39 cells are used as a negative control as described in Example 8 hereinbelow.

[0284] In another embodiment, cells are transfected with nucleic acid constructs encoding a reporter gene regulated by the P2X7R promoter (see above), an increase or decrease in the expression of the reporter gene in response to biological or pharmaceutical agents can be analyzed using methods that detect levels or status of protein or mRNA present in the corresponding cell or detect biological activities of the reporter gene. Suitable reporter molecules or labels, which may be used, include radionucleotides, enzymes, fluorescent, chemiluminescent or chromogenic agents as well as substrates, co-factors, inhibitors, magnetic particles, and the like. Designing such drug screening assays are well known in the art; see Harvey ed., 'Advances in drug discovery techniques', John Wiley and Sons, 1998; Vogel and Vogel eds., 'Drug discovery and evaluation: Pharmaceutical assays', Springer-Verlag Berlin, 1997). For example, drug screening in animal models, in vitro tests using animal cells, or in vivo tests involving toxicology tests in animals. An in vitro model can be used for screening libraries of compounds in any of a variety of drug screening techniques.

[0285] Candidate agents encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 Daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise carbocyclic or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups.

[0286] Candidate agents are also found among biomolecules including peptides, amino acids, saccharides, fatty acids, steroids, purines, pyrimidines, nucleic acids and derivatives, structural analogs or combinations thereof. Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical,



physical and biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, etc. to produce structural analogs.

[0287] Another technique for drug screening, which may be used, provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO 84/03564. In this method, as applied to the proteins of the invention large numbers of different small test compounds, e.g. aptamers, peptides, low-molecular weight compounds etc., are provided or synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with the proteins or fragments thereof, and washed. Bound proteins are then detected by methods well known in the art. Purified proteins can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support. In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding the protein specifically compete with a test compound for binding the protein. In this manner, the antibodies can be used to detect the presence of any peptide, which shares one or more antigenic determinants with the protein.

[0288] The present invention further particularly provides a method, wherein the pharmaceutical composition to be produced further comprises neuroprotective substances, nootropic substances, brilliant blue, piperidine or derivatives thereof, adamantane derivatives, substituted phenyl compounds, oxidized ATP, 2-O-(4-benzoylbenzoyl)adenosine-5-triphosphate or 3-O-(4-benzoylbenzoyl)adenosine-5-triphosphate. It is also envisaged that the pharmaceutical compositions to be produced further comprise antidepressants such as fluoxetine, paroxetine, sertraline, fluvoxamine, mirtazapine, reorepine, nefazodone or lithium carbonate.

[0289] In a preferred embodiment of the present invention, the compounds of the aforementioned methods comprise antagonist(s), partial antagonist(s), partial agonist(s) and/or agonist(s) for an altered ATP-gated ion channel P2X7R.

[0290] In accordance with the present invention, the term "antagonist" denotes molecules/substances, which are capable of inhibiting and/or reducing an agonistic effect. The term "antagonist" comprises competitive, non-competitive, functional and chemical antagonists as described, inter alia, in Mutschler, "Arzneimittelwirkungen" (1986), Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, Germany. The term "partial antagonist" in accordance with the present invention means a molecule/substance that is capable of incompletely blocking the action of agonists through, inter alia, a non-competitive mechanism.

[0291] In the context of the present invention, an antagonist is preferably a drug that does not provoke a response itself, but blocks agonist-mediated responses. It is a chemical entity that opposes the receptor-associated responses normally induced by another bioactive agent. For the P2X7R, the antagonists have an IC50 between 10 nanomolar and 300 micromolar.

[0292] As "agonist", in accordance with this invention, molecules/substances are denoted which have an affinity as

well as an intrinsic activity. Mostly, said intrinsic activity ( $\alpha$ ) is defined as being proportional to the quotient of the effect, triggered by said agonist (EA) and the effect which can be maximally obtained in a given biological system (Emax): therefore, the intrinsic activity can be defined as

$$\alpha \sim \frac{E_A}{E_{max}}$$

[0293] The highest relative intrinsic activity results from EA/Emax=1. Agonists with an intrinsic activity of 1 are full agonists, whereas substances/molecules with an intrinsic activity of >0 and <1 are partial agonists. Partial agonists show a dualistic effect, i.e. they comprise agonistic as well as antagonistic effects.

[0294] Preferably, in the context of the present invention, an agonist (or full agonist) is an endogenous substance or a drug that can interact with a receptor and initiate a maximal or complete physiological or a pharmacological response characteristic of that receptor. ATP, the natural ligand for the P2X7R, is an agonist with an EC50 of 300 micromolar while the synthetic P2X7R agonist Bz-ATP has an EC50 of 8 micromolar. Thus, agonists of P2X7R have an EC50 equal or below 300 micromolar. The EC50 is defined as the concentration of agonist that provokes a response half way between the baseline response and maximum response on a dose response curve where the X-axis plots concentration of an agonist and the Y-axis plots ion current. An inverse agonist (also called negative antagonist) is a drug which acts at the same receptor as that of an agonist, yet produces an opposite effect. A partial agonist is an endogenous substance or a drug that also provokes physiological or a pharmacological response but, the maximum response is less than the maximum response to a full agonist, regardless of the amount of drug applied. In the case of P2X7R, partial agonists have EC50s higher than 300 micromolar.

[0295] The person skilled in the art can, therefore, easily employ the compounds and the methods of this invention in order to elucidate the agonistic and/or antagonistic effects and/or characteristics of a compound/molecule/substance to be identified and/or characterized in accordance with any of the above described methods. Preferably, an identified antagonist of the ATP-gated ion channel P2X7R comprising the mutation(s) and/or deletion(s) described hereinabove may be useful to reestablish the properties normally shown by wild-type P2X7R ATP-gated ion channels. An identified agonist of the ATP-gated ion channel P2X7R comprising the mutation(s) and/or deletion(s) described hereinabove may be useful to reestablish the lost functionality of the P2X7R ATP-gated ion channel.

The Figures show:

[0296] FIG. 1a. Genomic map of the region on the human chromosome 12 associated to bipolar affective disorder. Genes found between markers NBG 11 and NBG2 are depicted.

[0297] FIG. 1b. Graphic illustrating the multipoint analysis using ASPEX on independent sib-pairs.

[0298] FIG. 1c. Graphic illustrating the multipoint analysis using ASPEX on all sib-pairs

[0299] **FIG. 1d.** Graphic illustrating the ASPEX sib\_phase by considering only independent sib-pairs

[0300] **FIG. 1e.** Graphic illustrating the ASPEX sib\_phase by considering all sib-pairs

[0301] **FIG. 1f.** Effect of the P2XR7v13A polymorphism on basal cortisol levels before and after administration of dexamethasone (DST test). Individuals were subjected to the test within the first ten days of admission. Individuals with the AG and GG genotypes have significantly lower cortisol levels pre- and post-dexamethasone administration.

[0302] **FIG. 1g.** Effect of the P2XR7v13A polymorphism on cortisol response during the Dex/CRH test. Individuals were subjected to the test within the first ten days of admission (i.e. at admission) and at the last ten days before discharge (i.e. at discharge). Individuals with the GG genotype have lower cortisol levels in response to the Dex/CRH test at admission and at discharge. These results are indicative of an abnormal HPA axis.

[0303] **FIG. 1h.** Effect of the P2XR7v13A polymorphism on ACTH response during the Dex/CRH test. Individuals were subjected to the test within the first ten days of admission (i.e. at admission) and at the last ten days before discharge (i.e. at discharge). Individuals with the GG genotype have lower ACTH levels in response to the Dex/CRH test, at admission and at discharge. These results are indicative of an abnormal HPA axis.

[0304] **FIG. 1i.** Duration of antidepressant treatment until remission. Depression is diagnosed according to the Hamilton Depression Rating Scale (HAM-D; Hamilton, Br. J. Soc. Clin. Psychol. 6 (1967) 278-296). A HAM-D score of 10 or below is regarded as remission of the depressive symptoms.

[0305] **FIG. 1j.** Effect of the P2XR7v13C polymorphism on basal cortisol levels before and after administration of dexamethasone (DST test). Individuals were subjected to the test within the first ten days of admission. Individuals with the CC genotypes have elevated cortisol levels post-dexamethasone administration.

[0306] **FIG. 1k.** Effect of the P2XR7v13C polymorphism on cortisol response during the Dex/CRH test. Individuals were subjected to the test within the first ten days of admission (i.e. at admission) and at the last ten days before discharge (i.e. at discharge). Individuals with the AC or CC genotype have elevated cortisol levels in response to the Dex/CRH test at admission, indicating an abnormal HPA axis.

[0307] **FIG. 1l.** Effect of the P2XR7v13C polymorphism on ACTH response during the Dex/CRH test. Individuals were subjected to the test within the first ten days of admission (i.e. at admission) and at the last ten days before discharge (i.e. at discharge). Individuals with the CC genotype have lower ACTH levels in response to the Dex/CRH test, at admission and at discharge. These results are indicative of an abnormal HPA axis.

[0308] **FIG. 2.** RT-PCR analysis of the complete coding sequence of P2X7R in different tissues

[0309] **FIG. 3.** P2X7R expression in the olfactory bulb, hypothalamus and ependymal cells in the brain of a stress-free mouse. Magnification 100x.

[0310] **FIG. 4.** P2X7R expression in the hippocampus/dentate gyrus and subcommissural organ in the brain of a stress-free mouse. Magnification 100x.

[0311] **FIG. 5.** Floating behaviour in the forced swim test. Passive stress coping behaviour decreased after long-term treatment with the antidepressant paroxetine (Par28: treated with paroxetine for 28 days, per os). Basal n=8; vehicle n=8; Par28 n=8.

[0312] **FIG. 6.** Comparative analysis of P2X7R expression in the olfactory bulb of stress-free, vehicle-treated and antidepressant-treated mice. Magnification 100x.

[0313] **FIG. 7.** Comparative analysis of P2X7R expression in the hypothalamus of stress-free, treated-treated and antidepressant-treated mice. Magnification 100x.

[0314] **FIG. 8.** Comparative analysis of P2X7R expression in ependymal cells of stress-free, vehicle-treated and antidepressant-treated mice. Magnification 100x.

[0315] **FIG. 9.** Comparative analysis of P2X7R expression in the hippocampus of stress-free, vehicle-treated and antidepressant-treated mice. Magnification 25x.

[0316] **FIG. 10.** P2X7R expression in the hippocampus of a vehicle treated mouse. Magnification 25x.

[0317] **FIG. 11.** P2X7R expression in the hippocampus of a mouse treated with the antidepressant paroxetine. Magnification 25x.

[0318] **FIG. 12.** Detailed expression of P2X7R in the dentate gyrus of a mouse treated with the antidepressant paroxetine. Magnification 400x.

[0319] **FIG. 13.** Comparative analysis of P2X7R expression and apoptotic cells in the hippocampus of a mouse treated with the antidepressant paroxetine. Magnification 100x.

[0320] **FIG. 14.** Floating behaviour in the forced swim test. Passive stress coping behaviour increased after acute intrahippocampal (bilateral, dentate gyrus) of siRNA targeting P2X7R. Vehicle n=7; control RNA n=10; P2X7R siRNA n=9.

[0321] **FIG. 15.** Comparative analysis of P2X7R expression in the hippocampus of mice treated with vehicle, control RNA and of siRNA targeting P2X7R. Magnification 100x upper row, 25x lower row.

[0322] **FIG. 16a, b, c, d, e.** Three splicing variants caused by polymorphisms in the introns of P2X7R.

[0323] **FIG. 17.** Expression of P2X7R in immortalized hippocampal cell lines.

[0324] **FIG. 18.** Increase calcium influx in hippocampal cells treated with a P2X7R agonist compound (BzATP).

[0325] **FIG. 19a, b.** Entry of ethidium bromide dye into hippocampal cells (a) treated with P2X7R agonist compound (BzATP) or (b) pre-treated with a P2X7R antagonist compound.

[0326] **FIG. 19c.** Agonist action of BzATP and tenidap on P2X7R activity. The calcium channel activity of human P2X7R was measured under basal conditions for four seconds to 10 seconds. A. Negative control consisting of cells loaded with 10  $\mu$ M Fluo-4-AM without further treatment. B.

Cells treated with 20  $\mu$ M BzATP after four seconds of basal measurement. C. Cells treated with 50  $\mu$ M tenidap after four seconds of basal measurement.

[0327] **FIG. 20.** Effect of intrahippocampal injection of a P2X7R agonist compound (BzATP) on behaviour in the forced swim test.

[0328] **FIG. 21.** Open field test measuring locomotor activity of mice treated with a P2X7R agonist compound (BzATP).

[0329] **FIG. 22.** Comparative analysis of apoptotic cells in the hippocampus of a mouse treated with control vehicle solution or a P2X7R agonist compound (BzATP).

[0330] **FIG. 23.** Effect of intrahippocampal injection of the P2X7R antagonist KN-62 and oATP on behaviour during the forced swim test

[0331] **FIG. 24.** Open field test measuring locomotor activity of mice treated with the P2X7R antagonist KN-62 and oATP.

[0332] A better understanding of the present invention and of its many advantages will be had from the following examples, offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

#### EXAMPLE 1

##### Linkage Analysis of Bipolar Affective Disorder in a Homogeneous Human Population

[0333] 41 families of different sizes containing a total of 485 sampled individuals from the region of Saguenay/Lac St-Jean were used in the linkage analysis. Individuals were distributed according to their diagnoses as follows: 105 individuals afflicted with Bipolar Disorder type I (BPI) or schizoaffective disorder bipolar type; 42 individuals diagnosed with Bipolar Disorder type II (BPII); 54 individuals with recurrent major depression; and 57 individuals with single episode major depression. The remaining 227 individuals were unaffected and normal. For the purpose of the calculation, the following classification was used: individuals diagnosed with either BPI, schizoaffective disorder, bipolar type, BPII and recurrent major depression were considered as affected (n=201); individuals with a single major depression episode were scored as unknown phenotype (n=57); and all others diagnoses as unaffected (n=227).

[0334] Blood samples from each individual were collected in 10-ml K3 EDTA Vacutainer tube (Becton-Dickinson) and genomic DNA was isolated by Puregene DNA Isolation kit (Gentra Systems). Blood was poured into 50 ml conical tube and diluted with four volume of Red Blood Cell Lysis

Solution. After an incubation of 10 minutes at room temperature, the tube was centrifuged for 10 minutes at 2,000 g and supernatant was removed leaving behind cell pellet and 200-400  $\mu$ l of the residual liquid. Cells were resuspended by vortexing the tube and 9 ml of Cell Lysis Solution were added with up and down pipetting. 40  $\mu$ l of RNase A Solution (20 mg/ml) were added and the sample was mixed by inverting the tube several times. Sample was incubated at 37° C. for 15 minutes and cooled to room temperature. 3 ml of Protein Precipitation Solution were added to cell lysate. Tube was vigorously vortexed for 30 seconds and centrifuged at 2,000 g for 10 minutes. Supernatant was poured into a new tube containing 9 ml of 100% isopropanol. Sample was mixed by inverting gently several times. Tube was centrifuged at 2,000 g for 5 minutes. The DNA white pellet was washed with 10 ml 70% ethanol and the tube was centrifuged at 2,000 g for 3 minutes. Ethanol was poured off and pellet allowed to partially air dry. DNA was solubilized in 500  $\mu$ l of DNA Hydration Solution. Final concentration was adjusted to 300-400  $\mu$ g/ml.

[0335] A fluorescent-based method was used for the genotyping of microsatellite markers. Briefly, the region encompassing each repeated sequence was amplified by PCR using an unlabeled primer and a fluorescent-labeled primer (Applied biosystems inc, CA, USA). The marker-associated dyes and the corresponding PCR product length are listed in table 2. The PCR reaction was performed using 10 ng of DNA sample, 0.2 unit of Taq platinum DNA polymerase (Invitrogene, CA, USA), 20 mM Tris-Cl (pH 8.4), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 100  $\mu$ M of dNTP, and 1.5  $\mu$ M of each primer in a final volume of 7  $\mu$ l. The samples were incubated at 95° C. for 3 minutes to activate the Taq platinum DNA polymerase, then 10 cycles of PCR amplification were performed as follows: 95° C. for 15 seconds; 58° C. for 15 seconds; 72° C. for 30 seconds; after that 15 cycles were performed as follow: 89° C. for 15 seconds; 58° C for 15 seconds; 72° C. for 30 seconds. Finally, the samples were incubated at 72° C. for 30 min. Following the PCR amplification samples were pooled according to their dye-labeled primer and their PCR product length (pool of four samples). Pooled sample were separated on an ABI 3100 DNA analyzer (Applied Biosystems inc, CA, USA). The resulting data were analysed using Genemapper2 (Applied Biosystems inc, CA, USA), and compiled in a 4D database (ACIUS) designed in a Macintosh environment as previously described (Morissette et al., Am. J. Med. Genet. (Neuropsychiatr. Genet.) 88 (1999), 567-587)

[0336] Markers used in the following linkage analysis are shown in table 2. Recombination fraction (q) between successive markers was computed according to the analyzed families.

TABLE 2

Genomic markers used for the linkage analysis					
Locus	Associated dye	Allele length (bp)	Distance (q)	Cumulative distance (cM)	Heterozygosity (%)
D12S1619	VIC	170-210	0.0135	0.00	74.5
NBG11	VIC	204-218	0.006	1.37	65.5
D12S1666	FAM	241-281	0.001	1.97	66.9
NBG5	VIC	253-261	0.001	2.07	38.3

TABLE 2-continued

Genomic markers used for the linkage analysis					
Locus	Associated dye	Allele length (bp)	Distance (q)	Cumulative distance (cM)	Heterozygosity (%)
D12S1721	VIC	263–299	0.005	2.17	72.1
NBG8	VIC	166–188	0.011	2.67	73.3
NBG6	NED	182–218	0.0115	3.79	73.9
NBG9	VIC	156–180	0.0035	4.95	68.9
NBG10	FAM	174–186	0.001	5.30	49.7
NBG12	NED	165–207	0.009	5.40	64.2
NBG4	NED	171–199	0.001	6.31	66.4
NBG3	VIC	182–206	0.006	6.41	64.8
NBG2	VIC	171–199		7.01	54.2

[0337] Haldane's map function was used for cumulative distance in cMorgans.

[0338] For bipoint parametric analysis, MOD score analysis were used where parametric LOD score were maximized over genetic models.

[0339] The following results were obtained under MOD score analysis for recessive models.

TABLE 3

MOD score analysis for recessive models			
Locus	Distance (q)	Cumulative distance (cM)	LOD score ( $q_{max}$ )
D12S1619	0.0135	0.00	3.46 (0.10)
NBG11	0.006	1.37	4.06 (0.04)
D12S1666	0.001	1.97	1.22 (0.14)
NBG5	0.001	2.07	0.66 (0.16)
D12S1721	0.005	2.17	2.82 (0.10)
NBG8	0.011	2.67	1.51 (0.00)
NBG6	0.0115	3.79	4.77 (0.06)
NBG9	0.0035	4.95	0.75 (0.22)
NBG10	0.001	5.30	0.74 (0.00)
NBG12	0.009	5.40	1.41 (0.16)
NBG4	0.001	6.31	3.56 (0.08)
NBG3	0.006	6.41	3.96 (0.08)
NBG2		7.01	2.59 (0.10)

[0340] Model-free LOD score studies using ANALYZE, sib\_phase from the ASPEX V1.85 package (David Hinds and Neil Risch 1999; [ftp://lahmed.stanford.edu/pub/aspex](http://lahmed.stanford.edu/pub/aspex), see also <http://watson.hgen.pitt.edu/docs/usage.html>) and SIMWALK2 (Sobel and Lange, Am J Hum Genet 58 (1996), 1323-1337) were performed to analyze the allele sharing among affected sib-pairs. The ANALYZE program weights sibships according to their size. The ASPEX sib\_phase program uses allele frequencies to reconstruct missing information, and is tailored for data sets where parents are missing, but additional typed children may be used to reconstruct and phase the parents. SimWalk2 is a statistical genetics computer application for haplotype, parametric linkage, non-parametric linkage (NPL), identity by descent (IBD) and mistyping analyses on any size of pedigree. SimWalk2 uses Markov chain Monte Carlo (MCMC) and simulated annealing algorithms to perform these multipoint analyses.

[0341] ASPEX sib\_phase was used with two computational strategies: First, by using strictly independent sib pairs; secondly, by using all affected sib-pair combinations. ASPEX was performed for bi-point and multipoint calculations.

[0342] The bi-point results observed with ANALYZE and ASPEX are shown in Table 4.

TABLE 4

Bi-point results observed with ANALYZE and ASPEX					
Locus	Distance (q)	Cumulative distance (cM)	Sib-pair from ANALYZE LOD score	sib_phase LOD score indep. sib-pairs	sib_phase LOD score all sib-pairs
D12S1619	0.0135	0.00	2.31	2.55	3.14
NBG11	0.006	1.37	2.83	2.72	3.27
D12S1666	0.001	1.97	1.01	2.52	3.14
NBG5	0.001	2.07	0.50	2.52	3.13
D12S1721	0.005	2.17	1.57	2.51	3.12
NBG8	0.011	2.67	0.51	2.24	2.75
NBG6	0.0115	3.79	2.55	2.11	2.64
NBG9	0.0035	4.95	0.49	1.65	1.97
NBG10	0.001	5.30	0.77	1.45	2.10
NBG12	0.009	5.40	0.47	1.44	2.17
NBG4	0.001	6.31	1.21	1.29	3.07
NBG3	0.006	6.41	1.84	1.29	3.07
NBG2		7.01	1.24	1.22	3.00

[0343] SIMWALK2 computed four different statistics based on descent trees. These statistics measure the degree of clustering among the marker alleles descending from the founders.

[0344] Statistic A is the number of different founder-alleles contributing alleles to the affected it is most powerful at detecting linkage to a recessive trait. Statistic B is the maximum number of alleles among the affected descended from any one founder-allele and most powerful at detecting linkage to a dominant trait. Statistic C is the 'entropy' of the marker alleles among the affected. Statistic D is the extent of allele sharing among all affected pairs as measured by their IBD kinship coefficient. Statistics C and D are more general statistics indicating whether a few founder-alleles are overly represented among the affected.

[0345] Table 5 shows the results observed with SIMWALK2. The authors signal that p-values should be generally conservative. They are expressed as  $-\text{Log}(\text{p-values})$ . For correspondence purpose,  $-\text{Log}(0.05)=1.30$ ,  $-\text{Log}(0.01)=2$ ,  $-\text{Log}(0.001)=3$  etc.

within the selected regions, the proportion of alleles shared IBD by affected sib-pairs was estimated with ASPEX (sib phase). For each region retained, the proportion of shared alleles was used as variable for a Principal Component Analysis and the first principal component as an index of linkage. Correlation analysis was done on these indexes to detect heterogeneity (correlation<0) or epistasis (correlation>0). Fisher algorithm was used to classify into two groups of families as linked or unlinked to a particular locus. A negative correlation was observed between the chromosome 12 region and the chromosome 15 area ( $r=-0.51$ ;  $p=0.023$ ). Cluster analysis suggested that 11 families out of 20 were linked to chromosome 12. This sub-sample was called the strata.

[0350] This strata included 11 families (266 sampled individuals) that include 52 BPI or schizoaffective disorder, bipolar type, 20 BP11 and 28 recurrent major depression

[0351] The following MOD score values illustrated in Table 6 were obtained under recessive models.

TABLE 5

SIMWALK2 analysis						
Locus	Distance (q)	Cumul. Distance (cM)	STAT(A)-Log (p-value)	STAT(B)-Log (p-value)	STAT(C)-Log (p-value)	STAT(D)-Log (p-value)
D12S1619	0.0135	0.00	1.4550	0.4103	1.1306	1.1310
NBG11	0.006	1.37	2.0157	1.4375	1.5955	1.9845
D12S1666	0.001	1.97	2.0236	0.9765	1.4727	1.4614
NBG5	0.001	2.07	1.7596	0.8558	1.3866	1.3602
D12S1721	0.005	2.17	1.6628	1.1692	1.4235	1.6384
NBG8	0.011	2.67	1.5374	0.6940	1.0623	1.1552
NBG6	0.0115	3.79	1.5896	0.4452	1.0935	1.1786
NBG9	0.0035	4.95	1.2677	0.3815	0.8412	0.9133
NBG10	0.001	5.30	1.1117	0.3642	0.6987	0.7554
NBG12	0.009	5.40	1.0809	0.3485	0.6694	0.7179
NBG4	0.001	6.31	1.1024	0.4148	0.6368	0.8544
NBG3	0.006	6.41	1.1040	0.4146	0.6373	0.8559
NBG2		7.01	1.0963	0.5380	0.6587	0.9356

[0346] Multipoint result observed with ASPEX when only independent sib-pairs were used (FIG. 1b). The maximum LOD score value was observed at NBG11.

[0347] Multipoint result observed with ASPEX when all sib-pairs were considered (FIG. 1c). The maximum LOD score value was observed at NBG11 but a second peak appeared at NBG4 and NBG3.

[0348] Multipoint and bi-point LOD score values computed by ASPEX were similar. The second peak, observed when all sib-pairs are used, may be explained by the presence of a recombinant affected individual, with many affected sibs, sharing the chromosomal region telomeric to NBG12. This kind of individuals has a large impact on LOD score values when all sib-pairs are used instead of one sib-pair. This situation was observed in two sibships.

[0349] Strata analysis was subsequently performed. Although HOMOG did not detect evidence for heterogeneity, a homogeneity test was constructed based on allele sharing found in selected chromosomal regions. Only 20 of the 41 families were used for this analysis since the others were not genotyped in all these regions. For each marker

TABLE 6

MOD scores under recessive models			
Locus	Distance (q)	Cumulative distance (cM)	LOD score ( $q_{\max}$ )
D12S1619	0.0135	0.00	4.03 (0.08)
NBG11	0.006	1.37	4.98 (0.00)
D12S1666	0.001	1.97	1.49 (0.12)
NBG5	0.001	2.07	0.79 (0.14)
D12S1721	0.005	2.17	4.23 (0.06)
NBG8	0.011	2.67	2.79 (0.00)
NBG6	0.0115	3.79	5.06 (0.06)
NBG9	0.0035	4.95	1.57 (0.14)
NBG10	0.001	5.30	1.73 (0.00)
NBG12	0.009	5.40	1.65 (0.12)
NBG4	0.001	6.31	4.60 (0.08)
NBG3	0.006	6.41	4.84 (0.06)
NBG2		7.01	2.80 (0.06)

[0352] Model-free LOD score results obtained with ANALYZE and ASPEX applied to the strata are shown in Table 7.

TABLE 7

Model-free LOD score obtained with ANALYZE and ASPEX					
Locus	Distance (q)	Cumulative distance (cM)	ANALYZE LOD score	sib_phase LOD score independent sib-pairs	sib_phase LOD score all sib-pairs
D12S1619	0.0135	0.00	4.54	5.29	7.65
NBG11	0.006	1.37	4.29	5.34	7.70
D12S1666	0.001	1.97	2.77	5.36	7.74
NBG5	0.001	2.07	0.67	5.36	7.74
D12S1721	0.005	2.17	4.48	5.35	7.74
NBG8	0.011	2.67	2.97	4.87	7.00
NBG6	0.0115	3.79	4.05	4.59	6.76
NBG9	0.0035	4.95	2.03	3.72	5.41
NBG10	0.001	5.30	2.00	3.42	5.89
NBG12	0.009	5.40	0.89	3.44	6.11
NBG4	0.001	6.31	2.84	3.71	9.00
NBG3	0.006	6.41	3.89	3.71	9.01
NBG2		7.01	1.91	3.52	8.73

[0353] Model-free results observed with SIMWALK2 are illustrated in Table 8.

TABLE 8

Model-free LOD score obtained with SIMWALK2						
Locus	Distance (q)	Cumulative Distance (cM)	STAT(A)-Log(p-value)	STAT(B)-Log(p-value)	STAT(C)-Log(p-value)	STAT(D)-Log(p-value)
D12S1619	0.0135	0.00	2.5963	0.9565	3.2156	2.3584
NBG11	0.006	1.37	3.0698	1.7400	3.7747	3.0103
D12S1666	0.001	1.97	2.9340	1.6546	3.5812	2.7223
NBG5	0.001	2.07	2.9781	1.2722	3.6505	2.7846
D12S1721	0.005	2.17	2.9680	1.2630	3.6844	2.7752
NBG8	0.011	2.67	3.0954	1.0804	3.4399	2.5654
NBG6	0.0115	3.79	3.1632	1.0672	3.2670	2.5956
NBG9	0.0035	4.95	2.2106	1.0137	2.7765	2.4456
NBG10	0.001	5.30	2.5513	1.0251	2.7625	2.1914
NBG12	0.009	5.40	2.4893	0.9868	2.6841	2.0920
NBG4	0.001	6.31	2.9028	1.1312	3.4063	2.8156
NBG3	0.006	6.41	2.9070	1.1326	3.4637	2.8300
NBG2		7.01	2.8430	1.1108	3.3135	2.7978

[0354] Multipoint results on the strata with ASPEX sib\_phase by considering only independent sib-pairs (FIG. 1b) or all sib-pairs (FIG. 1c) are shown in FIGS. 1d and 1e. As previously reported a second peak appeared when all sib-pairs were observed.

[0355] A confidence interval was calculated. GENEFINDER (Liang et al., Am. J. Hum. Genet. 66 (2000), 1631-1641) was used to estimate the location of the susceptibility gene (say t). The method is based on the IBD (Identity by Descent) sharing of affected sib-pairs for multiple markers. For the purpose of our analysis, pedigrees were divided into sibship. 56 nuclear families and 183 sib-pairs were used. Liang KY, Huang CY, Beaty TH (2000) A unified sampling approach for multipoint analysis of qualitative and quantitative traits in sib pairs. Am J Hum Genet 66:1631-1641

[0356] The GENEFINDER results points to localization of a susceptibility gene for affective disorders at 3.19±0.446 cM telomeric to the marker D12S1721 (D12S1721 is

approximately located at 136.82 cM on the sex-averaged Marshfield chromosome 12 map).

[0357] 95% C.I.: [2.32, 4.06];

[0358] 99% C.I.: [2.03, 4.35]

[0359] 99.9% C.I.: [1.71, 4.67]

[0360] From the strata, 24 nuclei, and 107 sib-pairs were obtained, and the location of the susceptibility gene was estimated at 3.07±0.57 (see map above). The following confidence interval (C.I.) was obtained:

[0361] 95% C.I.: [1.95, 4.19];

[0362] 99% C.I.: [1.59, 4.55]

[0363] 99.9% C.I.: [1.18, 4.96]

[0364] An association study using the NBG microsatellite markers was done with CLUMP (Sham & Curtis, Ann. Hum. Genet. 59 (1995), 97-105). Samples were distributed as

follow: 83 male/case; 124 female/case; 95 male/control; and 101 female/control. One thousand simulations were used to estimate p-values. The observed results are summarized in Table 9.

TABLE 9

Association study using the NBG microsatellite						
Locus	Sample		T1 statistic	T2 statistic	T3 statistic	T4 statistic
	Case	Control	(p-value)	(p-value)	(p-value)	(p-value)
NBG11	204	129	0.226	0.562	0.410	0.421
NBG5	206	194	0.972	0.980	0.948	0.971
NBG8	206	194	0.983	1.000	0.994	0.978
NBG6	206	194	0.147	0.074	0.759	0.485
NBG9	206	190	0.512	0.940	0.786	0.583
NBG10	206	190	0.594	0.480	0.403	0.709
NBG12	206	190	0.002	0.019	0.003	0.117

T1 statistic is the usual chi-squared statistic on the raw contingency table  
T2 statistic is the usual chi-squared statistic apply on he contingency table obtained after collapsing columns with small expected values together  
T3 statistic is the largest chi-squared statistic got by comparing one column of the original table against the total of the others columns  
T4 statistic is the largest chi-squared statistic got by comparing any combination of alleles against the rest.

[0365] Only the NBG12 marker gave significant association at the 1% level. For the others markers, there was no single alleles that seems to be associated with bipolar disorder. It seems that no founder-alleles was overly represented among the affected. There is no significant result for association of genotypes with the NBG markers.

[0366] Further microsatellite marker based association studies using CLUMP was performed on samples containing additional control and case individuals. One thousand simulations were used to estimate p-values.

[0367] HWE hypothesis was satisfied at the 5% level for each microsatellite marker after application of the conservative Bonferroni corrections for multiple testing (Bland & Altman, Brit. J. Med. 310 (1995) 170). Table 9a lists empirical p-values observed with CLUMP for allele and genotype association analyses. Empirical p-values less than 0.005 were observed at marker NBG12 for T1 and T3 statistics under allelic association analysis. T1 statistic suggested allelic association between bipolar affective disorders and NBG6 (empirical p-value=0.008). Moreover, a barely significant empirical p-value of 0.023 was observed at the most distal marker D12S2075.

[0368] In conclusion, the parametric and model-free multipoint results suggest to investigate genes located between D12S1619 and D12S1666. Moreover, according to GENEFINDER results, genes situated centromeric to NBG9 should be considered for association and linkage disequilibrium analysis. Moreover, positive association was seen with the NBG6 marker, which is located in intron 9 of the P2X7R gene.

## EXAMPLE 2

#### Physical Mapping and Mutation Analysis of Chromosome 12 Associating the P2X7R to Bipolar Affective Disorders

[0369] The most conservative prediction for the disease-associated region is included between markers NBG11 and NBG2 (see FIG. 1a). This region was delimited according to linkage and association analysis described in Example 1, using genethon markers and NBG markers. The approximate length of this region is 5.2 Mb. Two major gaps (between FLJ10701 and FLJ32372, and between FLJ1466 and MONDOA) were included in this region. At least 73

TABLE 9a

Empirical p-values observed with CLUMP for statistics T1 and T3 for allelic and genotypic analyses of microsatellite markers						
Name	Effective		Alleles		Genotypes	
	case	controls	T1 (p-value)	T3 (p-value)	T1 (p-value)	T3 (p-value)
NBG11	204	98	0.250	0.421	0.680	0.553
D12S1666	208	175	0.366	0.543	0.393	0.476
NBG5	213	179	0.969	0.934	0.997	1.000
D12S1721	210	176	0.693	0.463	0.805	0.838
NBG8	213	179	0.754	0.921	0.973	0.929
NBG6	213	179	0.008	0.356	0.172	0.449
NBG9	213	175	0.759	0.768	0.690	0.606
NBG10	213	175	0.521	0.178	0.122	0.173
D12S1349	212	180	0.887	0.864	0.782	0.816
NBG12	213	175	0.002	<10 <sup>-3</sup>	0.018	0.552
NBG4	207	178	0.418	0.506	0.813	0.545
NBG3	209	175	0.171	0.829	0.601	0.897
D12S378	211	180	0.171	0.405	0.540	0.560
NBG2	210	170	0.896	0.749	0.210	0.613
D12S1614	210	179	0.803	0.692	0.710	0.831
D12S342	211	180	0.394	0.740	0.445	0.622
D12S340	209	179	0.890	0.869	0.895	0.838
D12S1639	209	180	0.087	0.170	0.652	0.295
D12S1634	211	181	0.361	0.248	0.505	0.590
D12S2075	203	181	0.023	0.157	0.085	0.451

genes were listed in this area, where 48 are known genes and 25 are unknown but associated to mRNA and/or EST clusters based on the last genome assembly available at UCSC (November 2002). Predicted genes were not listed. However, the estimation of CI 99% (confidence interval) using GENEFINDER has limited the most interesting region between markers D12S1666 and NBG9. This genomic region covers 1.6 Mb and includes at least 28 genes, and has no major gap. Thus, the term fBAD (familial Bipolar Affective Disorders) region was used to describe the genomic segment between D12S1666 and NBG9. Genes found within this region include CaMKK2, CABP, P2X7, P2X4, PIN, PLA2, G1B, CIT, PXN, Rab35, and APC5. However, given the present art, it would not have been obvious to an ordinary person skilled in the art to select P2X7R as the gene associated with affective diseases. Other genes from the ones listed above would be obvious.

**[0370]** For example, the CaMKK2 gene (also known as Ca<sup>2+</sup>/Calmodulin-dependent protein kinase kinase beta, or CaMKKb) is a serine/threonine protein kinase involved in Ca<sup>2+</sup> dependent signalling pathways. CaMKK2 can activate in vitro the downstream kinases CaMKIV and CaMKI, which modulate gene transcription through phosphorylation of transcription factors (e.g., CREB, SRF, MEF2; Corcoran and Means, *J. Biol. Chem.* 276 (2001), 2975-2978; Soderling, *Trends Biochem. Sci.* 24 (1999), 232-236). Its role in the Ca<sup>2+</sup> cascade is not critical. Some studies suggest that CaMKs could be activated without the CaMKKs phosphorylation (Matsushita and Nairn, *J. Biol. Chem.* 274 (1998), 10086-10093). However, CaMKK phosphorylation step would contribute to amplification of the Ca<sup>2+</sup> signal since CaMKK is more sensitive to activation by Ca<sup>2+</sup>/Calmodulin, therefore CaMKK would be an important mediator when the levels of intracellular Ca<sup>2+</sup> are low (Anderson et al., *J. Biol. Chem.* 273 (1998), 31880-31889).

**[0371]** CaMKK2 is an obvious target for depression since prior art suggest that cAMP-dependent signaling pathways (mediated by PKA activation) is affected in brain from patients with Bipolar Affective Disorders (Field et al., *J. Neurochem.* 73 (1997), 1704-1710; Rahman et al., *J. Neurochem.* 68 (1997), 297-304; Takahashi et al., *J. Neurosci.* 19 (1999), 610-618). According to a study using lymphoblastic cell lines, Bipolar disorder could be related to a elevated intracellular calcium levels (Yoon et al., *Mol. Psychiatry* 6 (2001), 678-683). Moreover, some groups found relations between antidepressant drugs and CaMK activation (Budziszewska et al., *Br. J. Pharmacol.* 130 (2000), 1385-1393; Consogno et al., *Neuropsychopharmacology* 24 (2001), 21-30; Mori et al., *Neuropharmacology* 40 (2001), 448-456; Zanotti et al., *Neuropharmacology* 37 (1998), 1081-1089). Furthermore, inhibition of CaMKK by PKA-mediated phosphorylation suggest a close relationship between both pathways (Matsushita et al., *J. Biol. Chem.* 273 (1999), 21473-21481). These observations would suggest to a person skilled in the art that CaMKK2 is the gene responsible for bipolar affective disease.

**[0372]** Another obvious candidate for affective disorders would have been the CABP1 gene which generates four neuronal Ca<sup>2+</sup>-binding protein by alternative usage of the 9 coding exons, which are L-CABP, S-CABP, calbrain, and caldendrin (Haeseleer et al., *J. Biol. Chem.* 275 (2000), 1247-1260). Their expression is almost totally restricted to brain tissues. A functional study on calbrain reveals its negative effect on Ca<sup>2+</sup>/Calmodulin-dependent CaMKII activity by competitively interacts with the CaM-binding

domain of CaMKII (Yamagushi et al., *J. Biol. Chem.* 274 (1999), 3610-3616). One would expect similar roles in Ca<sup>2+</sup> signaling for other CABP1 alternative products. Participation of CABP1 gene in Ca<sup>2+</sup>-dependent signaling pathways would make it obvious to one skilled in the art to select this gene as a candidate for bipolar affective disorder. However, all CABP1 exons were analyzed for the presence of mutations, and surprisingly only two mutations were detected in noncoding regions.

**[0373]** The PIN gene (Protein inhibitor of NOS (Nitric oxide synthase)) is another obvious candidate responsible for bipolar affective disorder. Nitric oxide (NO) in the brain, may be involved in apoptosis, synaptogenesis, and neuronal development. Because NO cannot be stored in vesicles like other neurotransmitters, its release is regulated by the activity of NOS (Nitric oxide synthase). PIN is a direct inhibitor of NOS by binding and destabilizing the active homodimer complex of NOS (Jaffrey et al., *Science* 274 (1996), 774-777). PIN is highly conserved throughout the evolution and is expressed in many cell types. A recent clinical study evaluating plasma nitrate levels in depressive states suggests that NO production is increased in depression (Suzuki et al., *J. Affect. Disord.* 63 (2001), 221-224) and may result from a deficiency in NOS inhibition. Moreover in a mouse model, NO synthase antagonists have been linked to antidepressant properties (Harkin et al., 1999; Karolewicz et al., *Eur. J. Pharmacol.* 372 (1999), 215-220). Thus, PIN would be an obvious However, due to the pleiotrophic action of NO, a deficiency in PIN function would generate many unrelated disorders throughout the body. Thus, without the information presented in the disclosure herein, a person of ordinary skills in the art would have predicted PIN and not P2X7R as the gene associated with affective disorders.

**[0374]** The human phospholipase A2 group IB (PLA2G1B) catalyses the release of fatty acids from glycerol-3-phosphocholines. Phospholipase A2 genes (PLA2) are expressed in many tissues. Some studies have demonstrated associations between excessive PLA2 activity in brain and affective disorders (Chang et al., *Neurochem. Res.* 23 (1998), 887-892; Hibbeln et al., *Biol. Psychiatry* 25 (1989), 945-961). Moreover, other genetic studies have found associations between PLA2G1B gene and bipolar affective disorder (Dawson et al., *Psychiatr. Genet.* 5 (1995), 177-180). Thus, PLAG1B represent a likely candidate for affective disorders. However in the present example, only a single silent mutation was found within exon 3 of the PLAG1B gene.

**[0375]** The human citron kinase gene, Rho-associated protein (CIT) is a 183 kDa protein which associates to the GTPase Rho. CIT shares strong similarity with ROCK and ROK proteins which are other Rho-associated kinases (Madaule et al., *Nature* 394 (1998), 491-494). Rho GTPases are involved in many processes such as cytoskeletal organization, membrane trafficking, cell growth, and transcriptional activation (Van Aelst and D'Souza-Schorey, *Genes Dev.* 11 (1997), 2295-2322). Studies on brain variants of Citron-K (without the kinase domain) reveal the association with postsynaptic density proteins (PSD-95), suggesting a role in either synapse organization or function (Zhang et al., *J. Neurosci.* 19 (1999), 96-108; Furuvashiki et al., *J. Neurosci.* 19 (1999), 109-118).

**[0376]** The human paxillin (PXN) gene encodes for a 68 kDa protein found in focal adhesions. It is within focal adhesions where adhesion molecules dynamically interact with the cytoskeleton (Salgia et al., *J. Biol. Chem.* 270



(1995), 5039-5047). The signaling pathways that regulate these dynamic interactions begin to be elucidated. Many observations suggest that paxillin is involved in transducing signals from growth factor receptors to focal adhesions. The paxillin is expressed in many tissues including brain.

[0377] However as set forth below, the gene causative for affective diseases is identified as being the P2X7 receptor (P2X7R).

[0378] Mutations were searched in coding sequences and exon-intron boundaries of the above mentioned genes since such mutations are more likely to give a functionally significant Single Nucleotide Polymorphisms (SNP). The starting sample was composed of 16 unrelated affected individu-

als from the Saguenay/Lac St-Jean region, which gives an 80% power to detect polymorphisms with a frequency of 0.05. To identify polymorphisms, targeted sequences were first amplified by PCR. Then, PCR products are purified on Whatman GF/C membranes (VWR, Montreal, Canada), and quantified using the PicoGreen dsDNA quantitation assay (Molecular probes, Oregon, USA). 4 ng of purified PCR products are sequenced using the DYEnamic ET terminator cycle sequencing kit (Amersham Biosciences, Baie D'Urfé, Canada). The sequencing products are resolved on an ABI PRISM 3730XL DNA analyzer, and an ABI PRISM 3700 DNA analyzer. The PCR products are sequenced in both directions. The SNPs identified in studied genes are listed in Table 10.

TABLE 10

Mutation analysis between markers D12S1666 and NBG9				
Genes	Positions	Variations	Alleles	Modifications
Rab35	Exon06	RABE06A	486G-A	Silent Asn162
Rab35	Intron04	RABI04A	51C-T	unknown
Rab35	Intron03	RABI03A	33G-A	unknown
Rab35	Intron02	RABI02B	85G-A	unknown
Rab35	Intron02	RABI02A	76C-G	unknown
PXN	Exon11	PXNE11A	1527C-T	Silent Thr509
PXN	Exon06	PXNE06A	750C-T	Silent Ser250
PXN	Exon02	PXNE02A	217G-A	Gly73Ser
PLA2G1B	Exon03	PLA2G1BE03A	294C-T	Silent Ser98
PIN	5'UTR01	PINUTR01A	-49T-G	unknown
PIN	5'UTR01	PINUTR01B	-80T-C	unknown
PIN	Intron02	PINI02A	26C-T	unknown
PIN	Intron02	PINI02B	50C-T	unknown
CaBP	Intron04	CaBPI04A	35C-T	unknown
CaBP	exon01	CaBPE01A	-23A-G	unknown
OASL	Exon02	OASLE02A	213G-T	Silent Gly72
OASL	Exon02	OASLE02B	408C-T	Silent Leu136
OASL	Exon05	OASLE05A	1042G-A	Val348Met
OASL	Exon06	OASLE06A	1509G-A	Silent Ser503
P2X7R	5'UTR	P2XR7UTR5L	362T-C	unknown
P2X7R	5'UTR	P2XR7UTR5M	532T-G	unknown
P2X7R	5'UTR	P2XR7UTR5K	1100A-G	unknown
P2X7R	5'UTR	P2XR7UTR5J	1122A-G	unknown
P2X7R	5'UTR	P2XR7UTR5I	1171C-G	unknown
P2X7R	5'UTR	P2XR7UTR5F	1351T-C	unknown
P2X7R	5'UTR	P2XR7UTR5N	1702G-A	unknown
P2X7R	5'UTR	P2XR7UTR5G	1731T-G	unknown
P2X7R	5'UTR	P2XR7UTR5H	1860C-T	unknown
P2X7R	5'UTR	P2XR7UTR5A	2162C-A	unknown
P2X7R	5'UTR	P2XR7UTR5B	2238C-T	unknown
P2X7R	5'UTR	P2XR7UTR5D	2373A-G	unknown
P2X7R	5'UTR	P2XR7UTR5E	2569G-A	unknown
P2X7R	5'UTR	P2XR7UTR5C	2702G-A	unknown
P2X7R	Intron01	P2XR7I01C	3166G-C	unknown
P2X7R	Intron01	P2XR7I01A	24778C-T	unknown
P2X7R	Intron01	P2XR7I01B	24830C-T	unknown
P2X7R	Exon02	P2XR7v02A	24942T-C	Val76Ala
P2X7R	Exon03	P2XR7E03A	26188C-T	Arg117Trp
P2X7R	Intron03	P2XR7I03A	26308A-G	unknown
P2X7R	Intron03	P2XR7I03B	26422G-A	unknown
P2X7R	Intron04	P2XR7I04A	32394G-A	unknown
P2X7R	Intron04	P2XR7v05B	32434T-C	unknown
P2X7R	Exon05	P2XR7E05D	32493G-A	Gly150Arg
P2X7R	Exon05	P2XR7v05A	32507C-T	Tyr155His
P2X7R	Exon05	P2XR7E05C	32783C-T	Silent Cys168
P2X7R	Intron05	P2XR7I05C	32783A-C	unknown
P2X7R	Intron05	P2XR7I05D	35309T-C	unknown
P2X7R	Intron05	P2XR7I05B	35374C-T	unknown
P2X7R	Intron05	P2XR7I05A	35378A-C	unknown
P2X7R	Exon06	P2XR7E06A	35438G-A	Glu186Lys
P2X7R	Exon06	P2XR7E06B	35454T-C	Leu191Pro
P2X7R	Intron06	P2XR7I06C	35549T-C	unknown
P2X7R	Intron06	P2XR7I06G	35641G-C	unknown
P2X7R	Intron06	P2XR7I06D	35725A-C	unknown

TABLE 10-continued

Mutation analysis between markers D12S1666 and NBG9				
Genes	Positions	Variations	Alleles	Modifications
P2X7R	Intron06	P2XR7I06F	36001T-G	unknown
P2X7R	Intron06	P2XR7I06E	36064A-T	unknown
P2X7R	Intron06	P2XR7I06A	36091DelGTTT	unknown
P2X7R	Intron06	P2XR7I06B	36108C-G	unknown
P2X7R	Intron07	P2XR7I07A	36374C-T	unknown
P2X7R	Intron07	P2XR7I07B	36378G-A	unknown
P2X7R	Intron07	P2XR7I07C	36387T-A	unknown
P2X7R	Intron07	P2XR7I07D	36398G-C	unknown
P2X7R	Intron07	P2XR7I07E	37439C-T	unknown
P2X7R	Intron07	P2XR7I07F	37513T-C	unknown
P2X7R	Exon08	P2XR7E08C	37604C-T	Arg270Cys
P2X7R	Exon08	P2XR7v08A	37605G-A	Arg270His
P2X7R	Exon08	P2XR7v08B	37623G-A	Arg276His
P2X7R	Exon08	P2XR7E08D	37633C-T	Silent Asp279
P2X7R	Intron09	P2XR7v11A	47214C-T	unknown
P2X7R	Exon11	P2XR7v11B	47383G-A	Ala348Thr
P2X7R	Exon11	P2XR7v11C	47411C-G	Thr357Ser
P2X7R	Intron11	P2XR7I11D	47563T-C	unknown
P2X7R	Intron12	P2XR7I12A	54307C-T	unknown
P2X7R	Intron12	P2XR7I12B	54308G-A	unknown
P2X7R	Exon13	P2XR7v13F	54399C-T	Ala433Val
P2X7R	Exon13	P2XR7v13A	54480A-G	Gln460Arg
P2X7R	Exon13	P2XR7v13B	54523C-T	Silent Pro474
P2X7R	Exon13	P2XR7v13G	54562DelCCCTGAGAG CCACAGGTGCCT	Del of 7aa 488 to 494 PESHRL
P2X7R	Exon13	P2XR7v13C	54588A-C	Glu496Ala
P2X7R	Exon13	P2XR7v13H	54664C-G	Silent His521
P2X7R	Exon13	P2XR7E13D	54703G-T	Silent Leu534
P2X7R	Exon13	P2XR7E13J	54804A-T	Ile568Asn
P2X7R	Exon13	P2XR7v13I	54834G-A	Arg578Gln
P2X7R	Exon13	P2XR7v13E	54847G-A	Silent Pro582
P2X7R	3'UTR	P2XR7UTR3A	55169C-A	unknown
P2X7R	3'UTR	P2XR7UTR3B	55170A-C	unknown
P2X7R	3'UTR	P2XR7UTR3C	55171A-C	unknown
P2X7R	3'UTR	P2XR7UTR3D	55917C-T	unknown
P2X7R	3'UTR	P2XR7UTR3E	54925G-A	unknown
P2X4R	5'UTR	P2XR4UTR5I	-1956G-A	unknown
P2X4R	5'UTR	P2XR4UTR5H	-1649G-A	unknown
P2X4R	5'UTR	P2XR4UTR5G	-800G-A	unknown
P2X4R	5'UTR	P2XR4UTR5A	-648C-A	unknown
P2X4R	5'UTR	P2XR4UTR5B	-537A-G	unknown
P2X4R	5'UTR	P2XR4UTR5C	-437A-G	unknown
P2X4R	5'UTR	P2XR4UTR5J	-206VNRG	unknown
P2X4R	5'UTR	P2XR4UTR5D	-211C-G	unknown
P2X4R	5'UTR	P2XR4UTR5F	-150VNRGGGCCCC	unknown
P2X4R	5'UTR	P2XR4UTR5E	-98G-T	unknown
P2X4R	Intron01	P2XR4I01A	31G-T	unknown
P2X4R	Exon02	P2XR4E02A	262G-A	Silent mutation Ala87
P2X4R	Intron02	P2XR4I02A	4600C-T	unknown
P2X4R	Intron03	P2XR4I03A	15G-A	unknown
P2X4R	Intron03	P2XR4I03B	72G-A	unknown
P2X4R	Exon04	P2XR4E04A	355G-A	Ile119Val
P2X4R	Exon04	P2XR4E04A	375G-A	Silent Val125
P2X4R	Intron04	P2XR4I04B	17T-C	unknown
P2X4R	Intron04	P2XR4I04A	32G-A	unknown
P2X4R	Exon05	P2XR4E05A	465T-C	Silent Ser155
P2X4R	Exon07	P2XR4E07A	724A-G	Ser242Gly
P2X4R	Intron08	P2XR4I08A	DelT	unknown
P2X4R	Exon09	P2XR4E09A	944A-G	Tyr315Cys
P2X4R	Intron10	P2XR4I10A	11G-T	unknown
P2X4R	Intron10	P2XR4I10B	G-C	unknown
P2X4R	Intron10	P2XR4I10C	A-G	unknown
P2X4R	Intron11	P2XR4I11B	C-G	unknown
P2X4R	Intron11	P2XR4I11C	T-A	unknown
P2X4R	Intron11	P2XR4I11A	374C-T	unknown
CaMKK2	3'UTR	CaMKK2UTR3bA	733C-T	unknown
CaMKK2	3'UTR	CaMKK2UTR3aB	390G-A	unknown
CaMKK2	3'UTR	CaMKK2UTR3aA	239G-A	unknown
CaMKK2	Intron15	CaMKK2I15B	325T-C	unknown
CaMKK2	Intron15	CaMKK2I15A	169G-A	unknown
CaMKK2	Intron14	CaMKK2I14A	224A-G	unknown
CaMKK2	Intron10	CaMKK2I10A	156DelGTGATCCGCCTG	unknown

TABLE 10-continued

Mutation analysis between markers D12S1666 and NBG9				
Genes	Positions	Variations	Alleles	Modifications
CaMKK2	intron09	CaMKK2I09B	528A-G	unknown
CaMKK2	intron09	CaMKK2I09A	521A-G	unknown
CaMKK2	Exon09	SNP6f18v5	1095C-A	Silent Ile365
CaMKK2	Exon09	SNP6f18v4	1087C-T	Arg363Cys
CaMKK2	Exon05	CaMKKE05A	687C-T	Silent Pro229
CaMKK2	Intron03	CaMKK2I03A	10C-T	unknown
CaMKK2	Intron02	CaMKK2I02A	39C-T	unknown
CaMKK2	Intron01	CaMKK2I01B	2911G-C	unknown
CaMKK2	Intron01	CaMKK2I01A	89C-A	unknown
CaMKK2	Exon01	SNP6f18v2	253A-T	Thr85Ser
CaMKK2	Exon01	SNP6f18v1	29G-A	Ser10Asn
CaMKK2	5'UTR01	CaMKK2UTR01B	253T-C	unknown
CaMKK2	5'UTR01	CaMKK2UTR01A	63C-A	unknown
APC5	Intron01	APC5I01A	10G-T	unknown
APC5	Intron01	APC5I01B	50A-T	unknown
APC5	Intron05	APC5I05A	73T-C	unknown
APC5	Intron06	APC5I06A	73T-G	unknown
APC5	Exon11	APC5E11A	1416C-T	Silent His472

[0379] Each SNP in genes Rab35, PXN, PLA2G1B, PIN, CaBP, OASL, P2X4R, CaMKK2 and APC5 was designated according to the gene where it was found, and its location in that gene (intronic or exonic regions). Each SNP in the P2X7R gene was designated according to their position on SEQ ID NO: 1. The allele describes the position and the variation observed. In coding regions, the position is relative to the start codon, whereas the intronic SNPs are positioned relative to the beginning of the corresponding intron (when known). Primers used for identifying the SNPs in the P2X7R and the location of each SNPs included in tables 2 and 12 are defined in table 1a and SEQ ID NOs 52 to 111.

[0380] Association studies using missense SNPs were performed. Missense SNPs or SNPs that could be close to the splice sites were used, because it is more likely that diseases would be associated to an improper function in proteins. Case group was composed by bipolar I individuals, schizoaffective bipolar type (182 subjects) and bipolar II diagnosed persons (31 subjects). Many controls from the Saguenay/Lac-St-Jean region, were sampled from Steinert, Glaucoma and Paget DNA banks. The control individuals were not diagnosed for affective disorders. According to the

lifetime risks of bipolar disorders (1%), there is no need to screen controls for psychiatric disorders.

[0381] Direct sequencing of PCR products is by far the most accurate method of analysis and is the method of choice in view of our sequencing platform capacity. PCR products were analyzed by direct sequencing as described above. After sequencing analysis, individuals are automatically typed for the corresponding SNP using a home-developed program, GENO.pl. The results of SNP genotyping are compiled in a 4D database.

[0382] The association hypothesis was tested with CLUMP (Sham & Curtis 1995, Ann. Hum. Genet. 59:97-105). One thousand simulations were used to estimate p-values. Results are illustrated in table 11. The T1 statistic, which is the usual chi-squared statistic on the raw contingency table, was used to test for allelic association. Moreover, the largest chi-squared statistic got by comparing one column of the original table against the total of the other columns, called T3 statistic, was added to the previous one to test for potential genotype association since T1 statistic results may be biased when the contingency table contains cells with low values.

TABLE 11

Association hypothesis using CLUMP						
gene	SNPs	Effective		Allele Analysis	Genotype Analysis	
		Cases	Controls	p-value	p-value	p-value
				(T1)	(T2)	(T3)
P2X7R	P2XR7v11B	208	211	0.795	0.036	0.028
	P2XR7v13A	212	214	0.344	0.250	0.186
	P2XR7v13E	212	211	0.780	0.017	0.017
CAMKK2	SNP6f18v5	206	135	1.00	1.00	1.00
	SNP6f18v4	206	135	0.816	0.962	0.841
	SNP6f18v2	205	135	0.057	0.110	0.095
	SNP6f18v1	206	135	0.512	0.532	0.385

[0383] The association studies using SNPs in P2X7, P2X4, and CaMKK2 reveal associations significant at level of about 5% or less. Three genotype associations in P2X7 were observed. However, SNPs P2XR7v11B and P2XR7v13E are closely linked together based on a contingency table. There is also an allele association at level of

5.7% for SNP6f18v2 in CaMKK2. The information associated to each relevant SNP can be found in Tables 10 and 12.

[0384] Further association studies using CLUMP were performed on samples that contain more case and control individuals. One thousand simulations were used to estimate p-values.

TABLE 11a

Empirical p-values and odds ratio (OR) with 95% confidence interval observed with CLUMP for alleles and genotypes analysis of SNPs									
Gene	Marker (marker rank)	Allele Frequencies	Alleles						
			Effective		OR			Genotypes	
			case	controls	p-value	OR	CI	p-value	p-value
P2XR7	P2XR7UTR5F (1)	C (0.18); T (0.82)	212	208	0.280	1.21	0.86–1.71	0.067	0.069
	P2XR7UTR5G (2)	G (0.09); T (0.91)	211	204	0.481	1.19	0.76–1.87	0.261	0.231
	P2XR7UTR5H (3)	C (0.95); T (0.05)	210	202	0.549	1.19	0.67–2.13	0.768	0.582
	P2XR7UTR5A (4)	A (0.05); C (0.95)	210	207	0.526	1.26	0.68–2.34	0.754	0.517
	P2XR7UTR5B (5)	C (0.78); T (0.22)	211	207	0.629	1.09	0.79–1.50	0.104	0.128
	P2XR7UTR5D (6)	A (0.96); G (0.04)	211	205	0.268	1.43	0.77–2.65	0.598	0.240
	P2XR7UTR5E (7)	A (0.04); G (0.96)	211	210	0.658	1.23	0.65–2.33	0.139	0.234
	P2XR7UTR5C (8)	A (0.22); G (0.78)	208	210	0.889	1.04	0.75–1.44	0.168	0.293
	P2XR7I01B (9)	C (0.98); T (0.02)	210	207	0.352	1.71	0.67–4.39	0.348	0.348
	P2XR7v02A (10)	C (0.05); T (0.95)	211	208	0.189	1.49	0.84–2.64	0.397	0.167
	P2XR7I04A (11)	A (0.01); G (0.99)	211	211	0.344	0.25	0.03–2.23	0.356	0.356
	P2XR7v05B (12)	C (0.75); T (0.25)	212	211	0.854	1.03	0.76–1.41	0.234	0.335
	P2XR7E05D (13)	A (0.01); G (0.99)	211	211	0.726	1.51	0.42–5.38	0.735	0.735
	P2XR7v05A (14)	C (0.48); T (0.52)	211	209	0.638	1.07	0.82–1.40	0.895	0.895
	P2XR7E05C (15)	C (0.97); T (0.03)	210	211	0.195	0.45	0.16–1.31	0.349	0.276
	P2XR7I07E (16)	C (0.64); T (0.36)	208	214	0.394	0.87	0.66–1.16	0.057	0.064
	P2XR7v08A (17)	A (0.24); G (0.76)	210	212	0.221	1.22	0.90–1.67	0.433	0.496
	P2XR7v08B (18)	A (0.05); G (0.95)	210	213	0.386	0.71	0.36–1.41	0.520	0.662
	P2XR7V11A (19)	C (0.88); T (0.12)	213	149	0.394	0.80	0.50–1.29	0.387	0.463
	P2XR7v11B (20)	A (0.36); G (0.64)	208	211	0.795	1.04	0.79–1.38	0.036	0.028
	P2XR7v11C (21)	C (0.89); G (0.11)	211	212	0.409	0.82	0.52–1.28	0.303	0.661
	P2XR7v13F (22)	C (0.99); T (0.01)	196	207	0.030	3.24	1.04–10.12	0.039	0.039
	P2XR7v13A (23)	A (0.84); G (0.16)	212	214	0.344	1.21	0.85–1.72	0.250	0.186
	P2XR7v13B (24)	C (0.89); T (0.11)	207	212	0.494	0.83	0.53–1.31	0.315	0.699
	P2XR7v13C (25)	A (0.77); C (0.23)	211	213	0.731	0.95	0.68–1.31	0.557	0.616
	P2XR7V13H (26)	C (0.98); G (0.02)	211	213	0.238	1.75	0.68–4.49	0.236	0.236
	P2XR7E13D (27)	G (0.89); T (0.11)	211	213	0.435	0.82	0.53–1.28	0.268	0.680

TABLE 11a-continued

Empirical p-values and odds ratio (OR) with 95% confidence interval observed with CLUMP for alleles and genotypes analysis of SNPs									
Gene	Marker (marker rank)	Allele Frequencies	Alleles						
			Effective		T1 p-value	OR		Genotypes	
			case	controls		95%	CI	T1 p-value	T3 p-value
P2XR4	P2XR7E13J (28)	A (0.03); T (0.97)	204	199	0.179	0.48	0.16–1.42	0.329	0.329
	P2XR7v13E (29)	A (0.36); G (0.64)	212	213	0.841	1.04	0.79–1.37	0.026	0.025
	P2XR7UTR3E (30)	A (0.04); G (0.96)	205	197	1.000	0.96	0.45–2.04	1.000	1.000
	P2XR7UTR3A (31)	A (0.47); C (0.53)	208	209	0.932	0.99	0.75–1.30	0.264	0.239
	P2XR7UTR3B (32)	A (0.92); C (0.08)	208	210	0.174	0.65	0.38–1.14	0.151	0.303
	P2XR7UTR3C (33)	A (0.95); C (0.05)	208	210	0.395	0.71	0.36–1.40	0.508	0.667
	UTR5A	A (0.18); C (0.82)	212	210	0.285	0.82	0.57–1.18	0.514	0.484
	UTR5B	A (0.69); G (0.31)	212	210	0.670	0.93	0.70–1.25	0.833	0.833
	I06A	C (0.84); T (0.16)	207	192	0.212	0.78	0.53–1.16	0.398	0.217
	E07A	A (0.84); G (0.16)	212	208	0.294	0.81	0.55–1.19	0.536	0.479
	UTR3A	C (0.74); G (0.26)	211	203	0.015	1.50	1.11–2.02	0.021	0.014
	UTR3B	A (0.97); T (0.03)	211	209	0.653	0.81	0.33–1.97	0.649	0.649
	UTR3C	C (0.03); G (0.97)	211	209	0.653	0.81	0.33–1.97	0.672	0.672
CAMKK2	E09B	A (0.03); C (0.97)	208	214	0.830	0.85	0.36–2.00	0.829	0.829
	E09A	C (0.83); T (0.17)	208	214	0.202	0.78	0.54–1.14	0.446	0.473
	E01B	A (0.35); T (0.65)	207	214	0.048	1.33	1.01–1.76	0.126	0.218
	E01A	C (0.93); T (0.07)	208	214	0.189	1.44	0.86–2.39	0.439	0.237

[0385] Thirty-three SNPs in P2X7R, seven SNPs in P2X4R, and four SNPs in CAMKK2, with minor allele frequency higher or equal to 1 % were genotyped (Table 11a). The genotype distributions of these SNPs did not deviate significantly from HWE. At the 5% level, statistically significant increases of minor allele frequency were observed in the bipolar affective disorder group at p2XR7v13F (p-value=0.030, OR=3.24, 95% CI=1.04-10.12), P2XR4UTR3A (p-value=0.015, OR=1.50, 95% CI=1.11-2.02) and CAMKK2E01B (p-value=0.048, OR=1.33, 95% CI=1.01-1.76). The distribution of genotypes at SNPs P2XR7v13F and P2XR4UTR3A also differed significantly at this level for T1 and T3 statistics, with an increase of heterozygotes in the case sample. One SNP from exon 11 of P2X7R, P2XR7v11B, and another from exon 13, P2XR7v13E, displayed difference in genotype distributions with minimum p-value of 0.028 and 0.025 observed both

with T3 statistic. Again, increase in heterozygote frequency of 12% and 13% were respectively observed in the bipolar sample at these polymorphisms.

[0386] Significant haplotypic association tests led to p-values less than 0.5% for different SNP groups overlapping the P2X7R gene (Table 11b). Considering the SNPs collection ranging from SNP32507 to SNP54847 (table 11c) as an example for haplotype distribution, we observed the largest difference of frequencies between cases and controls with the haplotype no 1 (table 11d). The haplotype no 2 is another example of haplotype that is more frequently observed in cases group. On the other hand, the frequency for haplotype no 3 is slightly increased in control sample (difference of frequencies=0.91). Table 11e presents the peptidic products derived from the nucleotidic haplotypes shown in table 11d.

TABLE 11b

Haplotypes showing allelic association significant at the 0.5% level for T1 or T3 statistics.							
Haplotype (marker ranks <sup>1</sup> )	#SNPs	Distance <sup>2</sup> (bp)	Haplotype effective		T1 statistic	T3 statistic	#haplotype <sup>3</sup>
			case	control	(p-value)	(p-value)	
P2XR7I01B- P2XR7v13A (9–23)	15	29618	361	257	0.0003	0.0252	20
P2XR7v02A- P2XR7v13B (10–24)	15	29550	361	260	0.00008	0.0294	20
P2XR7I04A- P2XR7v13C (11–25)	15	22164	360	264	0.0003	0.0323	19
P2XR7v05B- P2XR7v13H (12–26)	15	22200	361	265	0.0004	0.0065	18
P2XR7E05D- P2XR7E13D (13–27)	15	22180	365	268	0.0035	0.0287	16
P2XR7v05A- P2XR7E13J (14–28)	15	22267	352	246	0.0007	0.0163	15
P2XR7E05C- P2XR7v13E (15–29)	15	22269	353	250	0.0012	0.0200	10
P2XR7I07E- P2XR7UTR3E (16–30)	15	17452	355	247	0.0020	0.0192	11

<sup>1</sup>The marker ranks of SNPs in the haplotype indicated in table 11b refer to those genotyped in table 11a

<sup>2</sup>Distance between the two most distal SNPs of the haplotype

<sup>3</sup>Number of haplotypes with frequencies >1% in case or control groups.

[0387]

TABLE 11c

Position and Allele for haplotype-forming SNPs. Haplotypes are described in table 11d.			
SEQ ID NO	Polymorphism	Position	
1	C-T	32507	
1	C-T	32548	
1	C-T	37439	
1	G-A	37605	
1	G-A	37623	
1	C-T	47214	
1	G-A	47383	
1	C-G	47411	
1	C-T	54399	
1	A-G	54480	
1	C-T	54523	
1	A-C	54588	
1	C-T	54664	
1	G-T	54703	
1	T-A	54804	
1	G-A	54847	

[0388]

TABLE 11d

haplotypes with significant difference of frequencies between affected and control individuals.									
Haplotype	32507	32548	37439	37605	37623	47214	47383	47411	54399
No 1	C	C	C	A	G	C	G	C	C
No 2	C	C	C	G	G	C	G	C	C
No 3	C	C	T	G	G	C	A	C	C
Haplotype	54480	54523	54588	54664	54703	54804	54847	F <sub>affected</sub>	F <sub>controls</sub>
No 1	A	C	A	C	G	T	G	0.20	0.13
No 2	A	C	C	C	G	T	G	0.05	0.01
No 3	A	C	A	C	G	T	A	0.11	0.20

[0389]

TABLE 11e

Corresponding amino acids for cSNPs described in table 11c. They are positioned according to SEQ ID NO 3.							
Position in SEQ							
ID NO 3	155	168	270	276	348	357	433
Haplotype 1	Y	C	H	R	A	T	A
Haplotype 2	Y	C	R	R	A	T	A
Haplotype 3	Y	C	R	R	T	T	A
Position in SEQ							
ID NO 3	460	474	496	521	534	568	582
Haplotype 1	Q	P	E	H	L	I	P
Haplotype 2	Q	P	A	H	L	I	P
Haplotype 3	Q	P	E	H	L	I	P

## EXAMPLE 3

## Polymorphisms Found in the P2X7R in Individuals Suffering From Depression

[0390] Association studies using SNPs in the P2X7R gene was performed in a case/control sample (535 individuals) from a German population. The case group was composed of 36 individuals diagnosed with bipolar type I or type II, and 279 individuals diagnosed with unipolar disorders (i.e. depression) representing 133 affected males and 182 affected females. Among controls, we count The remaining 220 control individuals were normal (i.e. diagnosed as non depressive), and comprising 81 males, 182 females and 14

of unknown gender. The same sexual distribution was noted in both groups.

[0391] SNPs were identified in this sample by using a subgroup of 24 affected individuals. SNPs in the P2X7R gene detected in the German population were similar if not identical to the SNPs seen in the Saguenay/Lac-St-Jean population (see table 12). Other rare missense SNPs were also noted in the German population, such as Arg117Trp (P2XR7E03A), Glu186Lys (P2XR7E06A), Leu191Pro (P2XR7E06B), Ile568Asn (P2XR7E13J). These amino acids are quite conserved between ortholog P2X7 genes. It is possible that the Ile568Asn (P2XR7E13J) mutation may be involved in the surface expression of P2X7.

TABLE 12

Comparison between polymorphisms in the Saguenay/Lac-St-Jean population and the German population in the human P2X7R gene						
Associated exons or introns	Variation (SNP or others)	Allele	Position*	Modification	Frequency (Canada)	Frequency (Germany)
5'UTR	P2XR7UTR5L	T-C	362	unknown	0.13	0.08
5'UTR	P2XR7UTR5M	T-G	532	unknown	0.16	0.1
5'UTR	P2XR7UTR5K	A-G	1100	unknown	0.13	0.13
5'UTR	P2XR7UTR5J	A-G	1122	unknown	0.13	0.13
5'UTR	P2XR7UTR5I	C-G	1171	unknown	0.06	0.02
5'UTR	P2XR7UTR5F	T-C	1351	unknown	0.3	0.12
5'UTR	P2XR7UTR5N	G-A	1702	unknown	—	0.02
5'UTR	P2XR7UTR5G	T-G	1731	unknown	0.17	0.15
5'UTR	P2XR7UTR5H	C-T	1860	unknown	0.07	0.15
5'UTR	P2XR7UTR5A	C-A	2162	unknown	0.07	0.12
5'UTR	P2XR7UTR5B	C-T	2238	unknown	0.3	0.27
5'UTR	P2XR7UTR5D	A-G	2373	unknown	0.07	0.12
5'UTR	P2XR7UTR5E	G-A	2569	unknown	0.1	0.02
5'UTR	P2XR7UTR5C	G-A	2702	unknown	0.31	0.27
Intron01	P2XR7I01C	G-C	3166	unknown	0.03	—
Intron01	P2XR7I01A	C-T	24778	unknown	0.03	—
Intron01	P2XR7I01B	C-T	24830	unknown	0.03	RARE
Exon02	P2XR7v02A	T-C	24942	Val76Ala	0.06	0.08
Exon03	P2XR7E03A	C-T	26188	Arg117Trp	—	RARE
Intron03	P2XR7I03A	A-G	26308	unknown	0.7	0.44
Intron03	P2XR7I03B	G-A	26422	unknown	0.18	0.12
Intron04	P2XR7I04A	G-A	32394	unknown	0.03	0.01
Intron04	P2XR7v05B	T-C	32434	unknown	0.33	0.29
Exon05	P2XR7E05D	G-A	32493	Gly150Arg	RARE	0.02
Exon05	P2XR7E05E	G-A	32506	Silent	—	RARE
Exon05	P2XR7v05A	C-T	32507	Tyr155His	0.33	0.44
Exon05	P2XR7E05C	C-T	32548	Silent	RARE	0.02
Intron05	P2XR7I05C	A-C	32783	Cys168	0.25	—
Intron05	P2XR7I05D	T-C	35309	unknown	ND	0.35
Intron05	P2XR7I05B	C-T	35374	unknown	0.7	0.67
Intron05	P2XR7I05A	A-C	35378	unknown	0.7	0.65
Exon06	P2XR7E06A	G-A	35438	Glu186Lys	—	0.02
Exon06	P2XR7E06B	T-C	35454	Leu191Pro	—	0.02
Intron06	P2XR7I06C	T-C	35549	unknown	0.04	0.08
Intron06	P2XR7I06G	G-C	35641	unknown	—	0.02
Intron06	P2XR7I06D	A-C	35725	unknown	0.21	0.27
Intron06	P2XR7I06F	T-G	36001	unknown	0.17	0.3
Intron06	P2XR7I06E	A-T	36064	unknown	0.11	0.1
Intron06	P2XR7I06A	DelGTTT	36091–36094	unknown	0.14	0.3
Intron06	P2XR7I06B	C-G	36108	unknown	0.14	0.29
Intron07	P2XR7I07A	C-T	36374	unknown	0.07	—
Intron07	P2XR7I07B	G-A	36378	unknown	0.21	0.28
Intron07	P2XR7I07C	T-A	36387	unknown	0.21	0.28
Intron07	P2XR7I07D	G-C	36398	unknown	0.42	0.4
Intron07	P2XR7I07E	C-T	37439	unknown	0.41	—
Intron07	P2XR7I07F	T-C	37513	unknown	—	RARE
Exon08	P2XR7E08C	C-T	37604	Arg270Cys	RARE	—

TABLE 12-continued

Comparison between polymorphisms in the Saguenay/Lac-St-Jean population and the German population in the human P2X7R gene						
Associated exons or introns	Variation (SNP or others)	Allele	Position*	Modification	Frequency (Canada)	Frequency (Germany)
Exon08	P2XR7v08A	G-A	37605	Arg270His	0.46	0.24
Exon08	P2XR7v08B	G-A	37623	Arg276His	0.03	0.02
Exon08	P2XR7E08D	C-T	37633	Silent Asp279	RARE	—
Intron09	P2XR7v11A	C-T	47214	unknown	0.08	0.03
Exon11	P2XR7v11B	G-A	47383	Ala348Thr	0.5	0.44
Exon11	P2XR7v11C	C-G	47411	Thr357Ser	0.08	0.07
Intron11	P2XR7I11D	T-C	47563	unknown	0.43	0.44
Intron12	P2XR7I12A	C-T	54307	unknown	0.32	—
Intron12	P2XR7I12B	G-A	54308	unknown	0.03	—
Exon13	P2XR7v13F	C-T	54399	Ala433Val	0.13	—
Exon13	P2XR7v13A	A-G	54480	Gln460Arg	0.13	0.17
Exon13	P2XR7v13B	C-T	54523	Silent Pro474	0.1	0.07
Exon13	P2XR7v13G	DelCCCTGAGA GCCACAGG TGCCT	54562-54582	Del of 7aa 488 to 494 (PESHRL)	RARE	—
Exon13	P2XR7v13C	A-C	54588	Glu496Ala	0.13	0.06
Exon13	P2XR7v13H	C-G	54664	His521Gln	0.03	—
Exon13	P2XR7E13D	G-T	54703	Silent Leu534	0.1	0.02
Exon13	P2XR7E13J	A-T	54804	Ile568Asn	—	0.01
Exon13	P2XR7v13I	G-A	54834	Arg578Gln	—	RARE
Exon13	P2XR7v13E	G-A	54847	Silent Pro582	0.4	0.45
3'UTR	P2XR7UTR3A	C-A	55169	unknown	0.48	0.37
3'UTR	P2XR7UTR3B	A-C	55170	unknown	0.09	0.1
3'UTR	P2XR7UTR3C	A-C	55171	unknown	0.05	0.06
3'UTR	P2XR7UTR3D	C-T	55917	unknown	0.001	—
3'UTR	P2XR7UTR3E	G-A	54925	unknown	—	0.01

[0392] The position and numbering of the polymorphism corresponds to the human P2X7R gene as defined in SEQ ID NO: 1. To identify the genomic organization of the P2X7R gene, BAC clones were firstly organized using known polymorphic markers, sequence tag sites (STSs), BAC-end sequences and expressed sequence tags (ESTs). Unorientated and unordered DNA regions were reassembled into a sequences using Phrap and reordered the pieces using P2X7R exons as scaffolds. No complete gene organization for P2X7R has been done. There is only a partial gene

structure from exon6 to 13, NT\_037809. Therefore, this genomic sequence encompassing the P2X7R gene as depicted in SEQ ID NO: 1 could contain some sequence errors, specifically in intronic regions. Primers used for SNP amplification and sequencing are shown in Table 1a and depicted in SEQ ID NOs: 52 to 111.

[0393] Statistical analysis was performed according to the CLUMP method (Sham & Curtis 1995, Ann. Hum. Genet. 59:97-105). Table 13 resumes the allelic and genotypic association studies for SNPs in P2X7 gene.

TABLE 13

Allelic and genotypic association studies using CLUMP						
Locus	Allele Frequencies*	Effective		Allele Analysis	Genotype Analysis	
		Cases	Controls	p-value (T1)	p-value (T1)	p-value (T3)
P2XR7UTR5F	2(0.23); 4(0.77)	311	217	0.109	0.319	0.339
P2XR7UTR5N	1(0.001); 3(0.999)**	314	218	0.038	0.048	0.048
P2XR7UTR5G	2(0.001); 3(0.105); 4(0.894)	314	218	0.993	0.714	0.761
P2XR7UTR5H	2(0.92); 4(0.08)	312	215	0.743	0.884	0.754
P2XR7UTR5A	1(0.08); 2(0.92)	312	219	0.557	0.786	0.678
P2XR7UTR5B	2(0.73); 4(0.27)	310	218	0.485	0.761	0.814
P2XR7UTR5D	1(0.92); 3(0.08)	311	217	0.555	0.787	0.691
P2XR7v02A	2(0.09); 4(0.91)	313	218	0.501	0.729	0.591
P2XR7I04A	1(0.04); 3(0.96)	314	220	0.604	0.433	0.348
P2XR7v05B	2(0.69); 4(0.31)	314	220	0.133	0.270	0.325
P2XR7E05D	1(0.03); 3(0.97)	314	220	0.842	0.827	0.827



TABLE 13-continued

Allelic and genotypic association studies using CLUMP						
Locus	Allele Frequencies*	Effective		Allele	Genotype Analysis	
		Cases	Controls	Analysis p-value (T1)	p-value (T1)	p-value (T3)
P2XR7E05E	1(0.006); 3(0.994)**	314	220	0.048	0.045	0.045
P2XR7v05A	2(0.60); 4(0.40)	314	220	0.038	0.144	0.219
P2XR7E05C	2(0.98); 4(0.02)	314	220	1.000	1.000	1.000
P2XR7I07F	2(0.002); 4(0.98)	315	219	1.000	1.000	1.000
P2XR7v08A	1(0.23); 3(0.77)	315	219	0.454	0.673	0.634
P2XR7v08B	1(0.02); 3(0.98)	315	219	0.636	0.638	0.638
P2XR7v11A	2(0.95); 4(0.05)	311	218	0.348	0.391	0.436
P2XR7v11B	1(0.45); 3(0.55)	312	218	0.605	0.803	0.790
P2XR7v11C	2(0.93); 3(0.07)	312	218	0.793	0.256	0.924
P2XR7I11D	2(0.45); 4(0.55)	312	219	0.665	0.735	0.740
P2XR7v13A	1(0.87); 3(0.13)	305	215	0.017	<0.001	<0.001
P2XR7v13B	2(0.93); 4(0.07)	305	216	1.000	0.228	0.677
P2XR7V13C	1(0.91); 2(0.09)	305	216	0.151	0.006	0.008
P2XR7E13D	3(0.94); 4(0.06)	315	219	0.402	0.429	0.474
P2XR7E13J	1(0.01); 4(0.99)	315	219	0.618	0.603	0.603
P2XR7E13I	1(0.004); 3(0.996)	315	219	0.999	1.000	1.000
P2XR7v13E	1(0.46); 3(0.54)	314	219	0.699	0.866	0.845
P2XR7UTR3A	1(0.518); 2(0.482)	314	219	0.617	0.850	0.875
P2XR7UTR3B	1(0.966); 2(0.034)	313	219	0.522	0.850	0.643
P2XR7UTR3C	1(0.979); 2(0.021)	313	219	0.636	0.505	0.382
P2XR7UTR3E	1(0.02); 3(0.98)	315	219	0.147	0.161	0.161

\*The column Allele Frequencies presents the allele for each SNP (A = 1, C = 2, G = 3, T = 4) and their respective frequency.

\*\*For this SNP we observed a zero cell in both (allele and genotype) 2 × 2 contingency tables. p-value <0.045 was observed exact Fisher test.

[0394] For the SNP analysis, the Hardy-Weinberg (HW) equilibrium was controlled in the control samples. The Hardy-Weinberg principle (HWP) may be stated as follow: In a large, randomly mating population, in which there is no migration, or selection against a particular genotype and the mutation rate remains constant, the proportions of the various genotypes will remain unchanged from one generation to another. Take a two allele system with alleles A and a. If the proportion of A in the population is represented as p and the proportion of a as q, then p plus q represent the sum total of alleles at this locus, that is p+q=1. The HWP is useful to evaluate some population problems like marital assortment, Inbreeding, population stratification, admixture, decreased viability of a particular genotype. The SNP P2XR7v13A did not respect the Hardy-Weinberg equilibrium.

[0395] The association hypothesis was also tested using an allele positivity table known to be suitable for the detection of susceptibility alleles showing a dominant mode of inheritance (Ohashi and Tokunaga, J. Hum. Genet. 44 (1999), 246-248; Ohashi et al., Ann. Hum. Genet. 65 (2001), 197-206). Similar results were obtained using this method as those obtained using the allele frequency tables, with the exception of P2XR7v05A where the p-values were 0.253. Thus, P2XR7v05A presented a less significant association in this analysis. This difference can be attributed to the mode of inheritance.

[0396] The proportion of unipolar individuals in analysis of the German population is quite important since the American Psychiatric Association (Diagnostic and Statistical Manual of Mental Disorders-4th Edition Text Revision (DMS-IV-TR), American Psychiatric Press, 2000) has reported an increase in susceptibility for unipolar disorders in female groups. To determine whether the sexual variable

could influence the association analysis, additional association studies were performed by controlling the sexual parameter. Normal individuals in the German population without gender information were omitted from the study. Then, a logistic regression model was derived by including the sex as factor. In order to obtain a model that is as stable as possible, the regression model was minimised by using the difference between log-likelihood's for models with or without interaction (Hosmer, and Lemeshow, "Applied logistic regression", John Wiley and Sons, 1989). The strategy used for handling the zero cells from contingency tables was to eliminate associated category completely. Calculations were done with SAS v8.0 SAS is a statistical software package that allows the user to manipulate and analyze data in many different ways. Because of its capabilities, this software package is used in many disciplines, including medical sciences, biological sciences, and social sciences.

[0397] The introduction of a sexual parameter did not perturb the association already observed in previous analysis. Moreover, this analysis model revealed additional results: a potential allele association with P2XR7v05B (p=0.064), and a genotypic association for P2XR7v08A (p=0.042) was observed.

[0398] Association studies using pooled samples was performed by merging individuals from the samples of the Saguenay/Lac St-Jean with those of the German population. Results are illustrated in table 14. The aim of this analysis is to highlight common features between both populations. However, according to differences between both samples (mainly the phenotype of affected individuals i.e. bipolar

disorder in the Saguenay/Lac St-Jean samples, versus mostly unipolar disorder in the German population) some parameters were controlled, including sex and ethnicity. The modelling strategy for logistic regressions was described above.

TABLE 14

Association studies using pooled samples from both populations				
Locus	Allele analysis		Genotype analysis	
	p-value for SNP	p-value for sex	p-value for SNP	p-value for sex
P2XR7v02A	0.8254	0.0085	0.8650	0.4531
P2XR7v05B	0.1751	0.3714	0.2034	0.5110
P2XR7v05A	0.3808	0.0266	0.0885	0.1392
P2XR7v08A	0.0452	0.0041	0.1021	0.3452
P2XR7v08B	0.3471	0.0040	0.3413	0.3617
P2XR7v11A	0.3559	0.0136	0.5888	0.4404
P2XR7v11B	0.5902	0.0093	0.3897	0.4302
P2XR7v11C	0.3731	0.0094	0.7648	0.4615
P2XR7v13A	0.0047	0.0209	<0.0001	0.4814
P2XR7v13B	0.5129	0.2352	0.9584	0.4092
P2XR7v13C	0.2466	0.0284	0.2225	0.4228
P2XR7v13E	0.8168	0.0159	0.3713	0.4990

[0399] An allelic and genotypic association was observed for the P2XR7v13A locus ( $p=0.0047$ ) which was stronger than in the separate analyses. A significant allelic association was also noted for the P2XR7v08A locus ( $p=0.0452$ ). In addition, the present analysis also demonstrate the potential relationship between SNP P2XR7v05A and the origin with a  $p\text{-value}=0.0515$  (not shown in the table) which is in agreement with previous association analysis done in both samples separately (see Table 13).

[0400] The haplotype analysis was performed using the German population. The PHASE program (Stephens et al., Am. J. Hum. Genet. 68 (2001), 978-989) was used to estimate SNPs haplotypes within exons of the P2XR7R gene. Haplotypes were created for each exon having more than one associated SNP (see Table 15 for exon-associated

SNPs). Case groups varied from 218-220 individuals, whereas control groups varied between 312-316 individuals. Association hypothesis was tested with the CLUMP method since many haplotypes were created for each exon. T1 and T3 statistic tests performed as described above. T2 and T4 statistics were also calculated owing to the presence of small effective cells in the contingency tables. T2 statistic is the usual chi-squared statistic applied on the contingency table obtained after collapsing columns with small expected values. T4 statistic is the largest chi-squared statistic obtained by comparing one column of the original table against the total of the other columns. One thousand simulations were used to estimate p-values. The resulting data was analyzed with the logistic regression model (describe above) using SDAS V8.0 in order the sexual parameter (for these tests the sample was reduced by 14 normal individuals). However, this analysis method is limited by the reliability of reconstructed haplotypes.

TABLE 15

Exon-associated SNPs	
Exons	Associated SNPs
5	P2XR7E05D P2XR7E05E P2XR7v05A P2XR7E05C
8	P2XR7v08A P2XR7v08B
11	P2XRv11B P2XRv11C
13	P2XR7v13A P2XR7v13B P2XR7v13C P2XR7E13D P2XR7E13J P2XR7v13I P2XR7v13E

[0401]

TABLE 16

Genotypic association with haplotypes in exon 13 of P2XR7R						
Exon (haplotype)	Allele analysis			Genotype analysis		
	Clump*	p-value(sex)	p-value(haplo)	Clump	p-value(sex)	p-value(haplo)
5(5)	T1: 0.032	0.3133	0.1947	T1: 0.193	0.460	0.5355
	T2: 0.068			T2: 0.159		
	T3: 0.054			T3: 0.099		
	T4: 0.059			T4: 0.304		
	T1: 0.551			T1: 0.812		
8(3)	72: 0.585	0.3813	0.3064	T2: 0.689	0.5428	0.6652
	T3: 0.646			T3: 0.644		
	T4: 0.646			T4: 0.756		
	T1: 0.750			T1: 0.625		
	T2: 0.786			T2: 0.919		
11(3)	T3: 0.726	0.0886	0.7396	T3: 0.929	0.2305	0.9494
	T4: 0.726			T4: 0.921		
	T1: 0.088			T1: 0.001		
	T2: 0.079			T2: 0.002		
	T3: 0.147			T3: 0.057		
13(15**)	T4: 0.072	0.1871	0.1264	T4: <0.001	0.4610	0.019

\*T1 test should not be considered because of contingency tables with zero cells.

\*\*Among these 15 haplotypes, we observed 8 haplotypes where case cells have less than 3 individuals.

[0402] Table 16 illustrates a genotypic association with haplotypes in exon 13 of the P2X7R genes. Interestingly, many haplotypes for the exon 13 were observed. The differences between statistics in exon 13 (T3 less significant) can be explained by the involvement of more than one genotype of haplotypes in the disease. A potential allelic association was also noted with haplotypes in exon 5 of the P2X7R gene.

[0403] The following are clinical results illustrating the functional consequences of polymorphisms in P2X7R.

[0404] The development and course of depression is causally linked to impairments in the central regulation of the hypothalamic-pituitary-adrenocortical (HPA) axis. Abnormalities in the HPA axis can be measured using the dexamethasone-suppression test (DST) or the combined dexamethasone/corticotropin-releasing hormone (Dex/CRH) test. Changes in cortisol and/or adrenocorticotrophic hormone (ACTH) measurements during the DST or Dex/CHR test are indicative of HPA dysfunction in depressed patients (Heuser et al, J. Psychiat. Res. 28 (1994) 341-356; Rybakowski and Twardowska, J. Psychiat. Res. 33 (1999) 363-370; Zobel et al, J. Psychiat. Res. 35 (2001) 83-94; Künzel et al, Neuropsychopharmacology 28 (2003) 2169-2178). In order to demonstrate that P2X7R SNPs associated with affective disorders also correlate with changes in the HPA axis, cortisol and ATCH levels in response to the DST and Dex/CRH test were measured for the P2XR7v13A and P2XR7v13C SNPs. P2XR7v13A consist of an A to G nucleotide change resulting in a Gln460Arg modification in the P2X7R protein. The P2XR7v13C SNP corresponds to an A to C nucleotide change resulting in a Glu496Ala modification that has been shown to drastically reduce protein activity (Wiley et al, Drug Dev. Res. 53 (2001) 72-76).

[0405] Methods and conditions for performing the DST and Dex/CRH test are well known in the art, see for example Heuser et al, J. Psychiat. Res. 28 (1994) 341-356; Künzel et al, Neuropsychopharmacology 28 (2003) 2169-2178. Briefly, individuals were pre-treated at 23:00 with an oral administration of 1.5 mg dexamethasone. For the DST test, a blood sample was drawn at 8:00 prior to dexamethasone administration (i.e. pre-dexamethasone) and at 8:00 the morning following dexamethasone administration (i.e. post-dexamethasone). For the Dex/CRH test, a venous catheter was inserted at 14:30 the day following dexamethasone administration and blood was collected at 15:00, 15:30, 15:45, 16:00, and 16:15 into tubes containing EDTA and trasylol (Bayer Inc., Germany). At 15:02, 100 mg of human CRH (Ferring Inc., Germany) was administered intravenously. Measurement of plasma cortisol concentrations was done using a commercial radioimmunoassay kit (ICN Biomedicals, USA) while plasma ACTH concentrations was measured using a commercial immunometric assay (Nichols Institute, USA). Both assays were performed according to the manufacturer specifications.

[0406] For the P2XR7v13A SNP, a decrease in basal cortisol levels was seen at admission in individuals with an

AG or GG allele when compared to individuals with the AA allele (FIG. 1f). During the Dex/CRH test, a reduction in cortisol and ATCH response was measured in individuals with the GG allele when compared to individuals with an AA or AG allele (FIGS. 1g and 1h). Furthermore, response to antidepressant treatment was delayed in GG individuals (FIG. 1i).

[0407] For the P2XR7v13C SNP, an increase in basal cortisol levels was measured post-dexamethasone administration (FIG. 1j). During the Dex/CRH test, individuals with the CC allele displayed elevated cortisol response (FIG. 1k), but reduced ATCH response (FIG. 1l) when compared to AA and AC individuals. These results are indicative of dysregulation of the HPA axis.

[0408] Thus, SNPs in P2X7R correlate with dysfunction in the HPA axis and demonstrate the functional and clinical consequences of polymorphisms in P2X7R.

#### EXAMPLE 4

##### P2X7R Gene Structure and mRNA Expression and Transcript Sequence

[0409] A 1700 bp nucleotide sequence corresponding to the human P2X7R promoter was analyzed by using MatInspector V2.2 and Transfac 4.0 algorithms. This analysis showed that the P2X7R gene does not contain a standard TATA box, but has SP1 sites that can make up for transcriptional initiation. Besides the SP1 sequences, there are binding sites for the transcription factors GATA, Oct and Ikari. These sites are thought to provide tissue specificity. Interestingly, the P2X7R promoter has binding sites that suggest responsiveness to different cytokines such as AP-1, NFAT and CEBPB.

[0410] P2X7R possesses 13 exons and 12 introns (Buell et al., Receptors Channels 5 (1998), 347), providing a basis for alternative splicing that would yield in theory different transcripts and produce different isoforms with possible different functions. No alternatively spliced variant was clearly identified. However, experiments of EST clustering allowed the description of three splicing variants. One is defined by the lack of the exon 5. This P2X7v02 variant corresponds to the clone IMAGE: 3628076 isolated from brain-derived cell lines. The P2x7v02 lacking the exon 5 produces a frame shift, thus generating a shorter polypeptide. The second splicing variant, P2X7v03, is characterized by the presence of the short intron 10 into the mRNA. This variant is supported by two high quality sequences, the cDNA clone BRAMY2008977 (AC number: AK090866) from human amygdala and the EST clone dbEST:7339877 derived from an unknown human tumor. The last variant, P2X7v04, is defined by the lack of the first exon that suggests an alternative promoter usage closed to the exon 2. A high quality EST clone dbEST:4782844 derived from a head and neck tumor supports this variant. These variants are shown in FIGS. 16a to 16e.

#### P2X7 variants.

P2X7v01	MPACCSCSDVDFQYETNKVTRIQSMNYGTIKWFFHVIIFSVCALVSDKLYQRKEPVISS
P2X7v04	MPPVD-----AFPCLPFS---FALVSDKLYQRKEPVISS

## -continued

P2X7v02 MPACCCSDVDFQYETNKVTRIQSMNYGTIKWFFHVIIFSIVCFALVSDKLYQRKEPVISS  
P2X7v03 MPACCCSDVDFQYETNKVTRIQSMNYGTIKWFFHVIIFSIVCFALVSDKLYQRKEPVISS  
1.....10.....20.....30.....40.....50  
P2X7v01 VHTKVKGIAEVKEEIVENGVKLVHSVFDTADYTFPLQGNSFFVMTNFLKTEGQEQR LCP  
P2X7v04 VHTKVKGIAEVKEEIVENGVKLVHSVFDTADYTFPLQGNSFFVMTNFLKTEGQEQR LCP  
P2X7v02 VHTKVKGIAEVKEEIVENGVKLVHSVFDTADYTFPLQGNSFFVMTNFLKTEGQEQR LCP  
P2X7v03 VHTKVKGIAEVKEEIVENGVKLVHSVFDTADYTFPLQGNSFFVMTNFLKTEGQEQR LCP  
61.....70.....80.....90.....100.....110  
P2X7v01 EYPTRRRLCSDRGCKKGWMDPQSKGIQTGR CVVHEGNQKTCEVSAWCPIEAVEEAPRPA  
P2X7v04 EYPTRRRLCSDRGCKKGWMDPQSKGIQTGR CVVHEGNQKTCEVSAWCPIEAVEEAPRPA  
P2X7v02 EYPTRRRLCSDRGCKKGWMDPQSKGLLS-----  
P2X7v03 EYPTRRRLCSDRGCKKGWMDPQSKGIQTGR CVVHEGNQKTCEVSAWCPIEAVEEAPRPA  
121.....130.....140.....150.....160.....170  
P2X7v01 LLNSAENFTVLIKNNIDFPGHNYTTRN ILPGLNITCTFHKTQNPQCP IFR LGDIFRETGD  
P2X7v04 LLNSAENFTVLIKNNIDFPGHNYTTRN ILPGLNITCTFHKTQNPQCP IFR LGDIFRETGD  
P2X7v02 -----  
P2X7v03 LLNSAENFTVLIKNNIDFPGHNYTTRN ILPGLNITCTFHKTQNPQCP IFR LGDIFRETGD  
181.....190.....200.....210.....220.....230  
P2X7v01 NFSDVAIQGGIMGIEIYWCNLD RWFHCHPKYSFRRLDDKTTNVS LYPGYNFRYAKYYK  
P2X7v04 NFSDVAIQGGIMGIEIYWCNLD RWFHCHPKYSFRRLDDKTTNVS LYPGYNFRYAKYYK  
P2X7v02 -----  
P2X7v03 NFSDVAIQGGIMGIEIYWCNLD RWFHCHPKYSFRRLDDKTTNVS LYPGYNFRYAKYYK  
241.....250.....260.....270.....280.....290  
P2X7v01 ENNVEKRTLKIVFGIRFDILVFGTGGKFDI IQLVVYIGSTLSYFGLAAV FIDFLIDTYSS  
P2X7v04 ENNVEKRTLKIVFGIRFDILVFGTGGKFDI IQLVVYIGSTLSYFGLAAV FIDFLIDTYSS  
P2X7v02 -----  
P2X7v03 ENNVEKRTLKIVFGIRFDILVFGTGGKFDI IQLVVYIGSTLSYFGLVRDSL FHALGKWFG  
301.....310.....320.....330.....340.....350  
P2X7v01 NCCRS HIYPWCKCCQPCVVNEY YRKKCESIVEPKPTLK YVSFVDESHIR MVNQQLLGRS  
P2X7v04 NCCRS HIYPWCKCCQPCVVNEY YRKKCESIVEPKPTLK YVSFVDESHIR MVNQQLLGRS  
P2X7v02 -----  
P2X7v03 EGSD-----  
361.....370.....380.....390.....400.....410  
P2X7v01 LQDVKGQEVPRPAMDFTDLSRLPLALHD TPPIPGQP EEIQLLRKEATPRSRDSPVWCQCG  
P2X7v04 LQDVKGQEVPRPAMDFTDLSRLPLALHD TPPIPGQP EEIQLLRKEATPRSRDSPVWCQCG  
P2X7v02 -----  
P2X7v03 -----  
421.....430.....440.....450.....460.....470

-continued

```

P2X7v01      SCLPSQLPESHRCLEELCCRKKPGACITTTSELFRKLVLSRHVLQFLLLYQEPLALDVDS
P2X7v04      SCLPSQLPESHRCLEELCCRKKPGACITTTSELFRKLVLSRHVLQFLLLYQEPLALDVDS
P2X7v02      -----
P2X7v03      -----

481.....490.....500.....510.....520.....530

P2X7v01      TNSRLRHCAIRCYATWRFGSQDMADFAILPSCCRWRIRKEFPKSEGQYSGFKSPY
P2X7v04      TNSRLRHCAIRCYATWRFGSQDMADFAILPSCCRWRIRKEFPKSEGQYSGFKSPY
P2X7v02      -----
P2X7v03      -----

541.....550.....560.....570.....580.....590

```

[0411] Therefore the transcriptional and translational start sequences of the human P2X7R were analyzed using Blast, Genescan and HMMgene computer software. This analysis indicated that P2X7R possesses with high probability only one translation start site. Most P2X7R expression sequence tags (ESTs; Unique cluster Hs. 193470) having a reliable 5' end showed identical transcriptional start site. None of the ESTs showed any indication of alternative splicing. Therefore, in silico analysis suggests that there is a low probability to find different transcripts produced by alternative splicing or alternative promoter usage.

[0412] The above mentioned in silico data were confirmed by RT-PCR analysis spanning the whole predicted human P2X7R coding sequence using 14 and 19 bases (5'-ATGC-CGGCTTGCTG-3'; 5'-GTAGGGATACTTGAAGCCA-3') oligonucleotides corresponding to the beginning and end of the coding sequence, respectively. Total RNA from whole brain, different dissected brain areas, thymus, spleen and kidney were isolated and analyzed for P2X7R expression. RT-PCR reactions were performed using the C. Therm One Step polymerase system (Roche Applied Science) and a protocol for touch down PCR with hot start. Briefly, Reverse Transcription was performed at 52° C. according to the manufacturer's conditions. PCR reactions were executed with an annealing temperatures of 64° C. for the first five cycles and of 54° C. for the next 30 cycles.

[0413] A single specific band of the size of 1785 bp corresponding to the complete coding sequence of P2X7R was detected. P2X7R mRNA was detected in the whole brain, hippocampus, cerebellum, leukocytes and thymus but not in cerebral cortex, hypothalamus, spleen and kidney (FIG. 2). All PCR products were cloned using the pGEM-T-Easy plasmid (Promega), selected in Top-10 bacteria (Invitrogen) by blue-white selection and tested by EcoRI digestion. Clones having fragments of the expected size were amplified and purified for sequencing. The sequence confirmed the identity of the 1785 bp clones as the complete coding sequence of wild-type P2X7R. Therefore, in all the tissues tested, wild-type P2X7R is expressed as a single transcript which includes the complete coding sequence. The presence of tissue specific isoforms is unlikely. These studies provide useful information about the P2X7R mRNA expression and transcript processing. This information can be used to synthesize riboprobes for in situ hybridization,

Northern and Southern blot as well as engineering cells for the overexpression of P2X7R.

#### EXAMPLE 5

##### P2X7R Expression in the Mouse Brain

[0414] The expression of P2X7R was further studied by immunohistochemistry of serial sections of complete mouse brains using a polyclonal antibody directed against an internal peptide of P2X7R (Santa Cruz Biotechnology). The brains from stress-free mice were shock frozen, cut into 16  $\mu$ m slices and fixed with paraformaldehyde for 5 minutes. The sections were blocked for 30 minutes at room temperature with 1:10 horse serum. All antibodies were diluted in TBST buffer (Tris-buffered saline with 0.05% Tween-20). The first antibody was used in a dilution 1:200 and incubated overnight. All washes were performed with TBST buffer. As a secondary antibody, an anti-goat IgG biotinylated (Vector Laboratories) was used and detection was performed using the streptavidin-biotin-horse-radish peroxidase complex system (Vector Laboratories) in combination with diaminobenzidine. Slides were counterstained with toluidine blue using standard procedures. The same procedure in the absence of the primary antibody was performed as a negative control. As a positive control to test the Preservation of the tissue was verified with an antibody specific for the protein Patched1 (Santa Cruz Biotechnology). Patched1 was used as positive control since it stains all relevant brain structures and is not affected by stress or antidepressants. Very specific staining pattern was detected, consistent with the specific subcellular localization of P2X7R in brain cells. Negative controls were completely devoid of signal. Positive control with Patched1 showed identical signal intensity and distribution in all samples, indicating that all tissues were equally well preserved and processed.

[0415] Proceeding from frontal to caudal, P2X7R protein was observed in the glomerular layer of the olfactory bulb at low levels (FIG. 3). P2X7R was also present at very low levels in a restricted area of the periventricular hypothalamic nucleus (FIG. 3). Ependymal cells surrounding the lateral ventricles also showed a faint staining (FIG. 3). A stronger signal was detected in restricted areas of the hippocampus, where the signal was present in single cells of the polymorph layer, the lacunosum moleculare and the oriens layer (FIG. 4). In more posterior areas of the hippocampus,

the signal was present in the molecular layer, stratum radiatum and near the CA3. In a further caudal position, P2X7R was expressed in the subcomissural organ (**FIG. 4**). Therefore, the basal P2X7R expression in the brain of stress-free mice is restricted to areas that had been previously associated with depression, stress, learning and memory.

#### EXAMPLE 6

##### P2X7R is Modulated in Mice Treated With an Antidepressant

[0416] Further validation of role of P2X7R in affective disorders was performed by examining its expression pattern in response to stress and treatment with antidepressant drugs. A treatment schedule which has been proven to produce antidepressant effects on the behavioural level was administered to mice which were characterized as antidepressant-responsive by using a variety of behavioural paradigms suitable to detect anxiolytic and antidepressant effects of classical antidepressants like the selective serotonin reuptake inhibitor paroxetine. Paroxetine was delivered by gavage to naive male mice over a time period of 28 days at a dosage of 10 mg/kg bodyweight twice per day. In parallel, a control group of mice was given vehicle solution (i.e. without paroxetine) using the same treatment regimen while a second control group of mice was left undisturbed and stress-free (i.e. untreated) during the same period of the experiments. At the end of the long-term treatment, part of the mice of each experimental group were tested in the dark/light box (test of anxiety behaviour) and in the Porsolt's forced swim test (test of depressive-like behaviour) to confirm the effectiveness of the treatment (**FIG. 5**). Passive stress coping behaviour decreased after long-term treatment with the antidepressant paroxetine. The other part of the experimental groups (i.e. mice without test experience) were decapitated, brains rapidly removed and frozen at  $-80^{\circ}\text{C}$ . until usage.

[0417] The expression of P2X7R in the brains of mice under stress-free conditions, and mice under mild stress produced by the vehicle application, and mice under paroxetine treatment was evaluated using three different brains from each group. Serial slides from each group of animals were analyzed in parallel by immunohistochemistry using the same materials in order to produce completely comparable results. No significant change in P2X7R expression in the olfactory bulb was seen in response to stress or to paroxetine treatment (**FIG. 6**). However, in the periventricular nucleus of the hypothalamus, paroxetine produced a slight inhibition of P2X7R expression (**FIG. 7**). No significant change was observed in the ependymal cells from different brain areas (**FIG. 8**). The most dramatic changes were observed in the hippocampus, where P2X7R was strongly inhibited by stressful handling whereas paroxetine treatment produced a marked stimulation above basal levels (**FIGS. 9, 10 and 11**). This effect was observed all along the hippocampus but was more evident in the polymorph layer near the dentate gyrus. In the subcommissural organ, P2X7R expression remained unchanged by the different treatments. Therefore, P2X7R expression is strongly regulated in two specific brain areas involved in depression and stress. Other brain areas, which showed low levels of P2X7R and are not directly involved in depression, did not show changes.

[0418] In the samples from mice treated with paroxetine and showing a strong P2X7R expression, it was possible to analyze the distribution of P2X7R in more detail (**FIGS. 10 and 11**). The P2X7R protein was not only present in cell bodies but also was clearly detected in projections innervating the granular layer of the dentate gyrus (**FIG. 12**). This subcellular localization of P2X7R is consistent with a role in neurotransmitter release and long term potentiation.

[0419] Since some reports (Muria et al., *Biochem. J.* 288 (1992) 897-901; Ferrari et al., *FEBS Lett.* 447 (1999), 71-75) suggest that chronic and high dose stimulation of P2X7R may cause apoptosis in some cell types, the hippocampus of the above described animals were analyzed for the co-localization of apoptotic cells and P2X7R expressing cells, in consecutive sections, using TUNNEL staining and immunohistochemistry. In correlative sections, only few apoptotic cells were detected and they were present along the granular layers of the hippocampus where no P2X7R expression was observed (**FIG. 13**). No significant differences in the numbers of apoptotic cells were observed between the different treatment conditions. Therefore, the location and number of apoptotic cells did not correlate with the location and number of cells expressing P2X7R and rules out an involvement of P2X7R in the induction of apoptosis in the hippocampus.

[0420] Thus, P2X7R expression is considerably restricted to specific brain areas involved in depression. Moreover, P2X7R expression is inhibited by stress and strongly stimulated by antidepressant treatment in these specific areas. Therefore, P2X7R fulfils all criteria required for the actions of antidepressants according to the highest standards in the field of depression research. In addition, these results suggest that modulation of function of P2X7R is associated with chronic stress, which serves as a model for several aspects of affective disorders.

#### EXAMPLE 7

##### The Behavioural Effect of P2X7R Inhibition in Mice

[0421] To demonstrate that P2X7R inhibition acts as a causative agent for affective disorders, P2X7R function was specifically inhibited in distinct regions of the brain without affecting any other brain function. This was achieved by delivering double stranded small interference RNA molecules (siRNA) into restricted areas of the brain.

[0422] According to the observed expression pattern of P2X7R in the hippocampus (**FIGS. 9, 10, and 11**) and the known involvement of the hippocampus in depression, the dentate gyrus (hippocampus) was selected as target region for siRNA application. Male, naive mice were bilaterally implanted with a guide cannulae (23 gauge, length 8 mm) by means of a stereotactic instrument. The coordinates, in relation to bregma, were  $-2.0\text{ mm}$  posterior,  $\pm 1.0\text{ mm}$  lateral, and  $-1.0\text{ mm}$  ventral. Following a recovery period of 5 days, the mice were divided into three experimental groups: vehicle (veh), control double stranded RNA (control), and P2X7R specific double stranded siRNA (siRNA). Sequences used for P2X7R siRNA are 5'-GUGGGUCUUGCAUGAUCTT-3' and 5'-GAUCAUGUGCAAGACCACTT-3'. Both sequences were annealed and injected together as a double stranded RNA. On day 6 after

surgery, mice were slightly anaesthetized with Isofluran and injections of siRNA were carried out. The concentration of the control and siRNA was 0.1 nmol/ $\mu$ l, and a volume of 1  $\mu$ l per side was infused using specifically adapted injection systems (30 gauge, length 9 mm). The anaesthesia for the infusion was of short duration and the mice were awake immediately or few seconds after the manipulation.

**[0423]** Once delivered into the brain the siRNA molecules specific for P2X7R were taken up by brain cells and specifically induce the degradation of the complementary P2X7R mRNA with high efficiency. As a result, P2X7R function was specifically inhibited for a short period without affecting any other brain function. In this regard, injection of vehicle or control siRNA did not result in any obvious changes in normal behaviour, i.e., food and water intake, or motor behaviour in the home cage.

**[0424]** The effects of P2X7R inhibition on depressive-like behaviour was assessed 24 hours and 48 hours after infusion of siRNA, control or vehicle according to the standard test paradigm, the Porsolt's forced swim test (Porsolt et al., Arch. Int. Pharmacodyn. 229 (1977), 327-336; Porsolt, Rev. Neurosci. 11 (2000), 53-58). The parameter used to evaluate depressive-like behaviour is the time the animal is floating in the water, a behaviour which is associated with behavioural despair as the animal does not make any effort to actively cope with the stressful situation. Compared to vehicle application, no influence of control double stranded RNA (5'-CAACUUAUCUUCUACGCGTT-3') on floating behaviour (passive stress coping) was detected. In contrast, compared to controls, mice infused with P2X7R specific siRNA showed a significant increase in passive behaviour, which is construed as depressive-like behaviour (**FIG. 14**). This interpretation becomes moreover evident when the effects of antidepressants on passive stress coping behaviour in the forced swim test are visualized (**FIG. 5**). Passive stress coping behaviour increased after acute intrahippocampal injection (bilateral, dentate gyrus) of siRNA targeting P2X7R. The Porsolt's forced swim test is a standard test used to assess the effectiveness of antidepressants and it has been proven by many studies that the test is selectively sensitive for these effects, given that the right animal model is used. The paradigm has been widely used to test pharmaceutical compounds and to validate animal models of depression, which show an increase in passive behaviour as do the mice where P2X7R has been inhibited (siRNA).

**[0425]** At the end of the experiment, the mice were sacrificed and the brains were examined to confirm the location and efficiency of the siRNA injections. For this purpose the brains were cut into sections and the slides were stained by immunohistochemistry using the above mentioned protocols. Brains from mice injected with the specific double stranded siRNA, with control double stranded RNA and with vehicle were examined in parallel. Under these conditions, the specific siRNA directed against P2X7R injected near the dentate gyrus induced on average an 80% inhibition of P2X7R protein expression as compared to the samples from mice injected with vehicle or with control double stranded RNA. Both the number of cells expressing P2X7R as well as the intensity of the expression were strongly reduced (**FIG. 15**). The injections with siRNA did not produce any sign of local inflammation or infiltration at the hippocampus. Thus, P2X7R expression is specifically and locally inhibited by siRNA application in vivo. This

inhibition produced behavioural changes indicating a causative role for P2X7R in affective disorders. These results in combination with those mentioned above support and confirm the observation of mutations in P2X7R being associated with affective diseases in humans and that modulation of P2X7R activity has antidepressive effects.

## EXAMPLE 8

### Drug Screening Assay

**[0426]** Methods for identifying P2X7R agonists were established using an immortalised mouse hippocampal cell line expressing the endogenous P2X7 gene. Briefly, the expression of P2X7 was confirmed by culturing the cells at 37° C./5% CO<sub>2</sub> in DMEM with 10% foetal calf serum (Gibco). Upon reaching 80% confluence, cells were collected in PBS and homogenized by repeated passage through a syringe (18G needle). The amount of total protein was measured by the Bradford assay (Sigma; diluted 1:5, O.D. measured at 595 nm) according to the manufacturer's recommendation. Protein homogenates were then mixed with an equal volume of loading buffer (50 mM Tris-Cl pH 6.8; 25% glycerol; 7.2 mM bromophenol blue; 2% SDS; 200 nM  $\beta$ -mercaptoethanol) and subsequently denatured in boiling water for 10 minutes. 20 mg of each sample were loaded onto a 10% polyacrylamide gel containing 0.4 % SDS. Electrophoresis and Western blot transfer were performed according to conventional protocols described, for example, in Sambrook, Russell "Molecular Cloning, A Laboratory Manual", Cold Spring Harbor Laboratory, N.Y. (2001). Membranes were then blocked with 5% dry milk and incubated with an antibody against P2X7R (1:1000 dilution; Santa Cruz Biotech) followed by incubation with a horse anti-goat peroxidase-coupled secondary antibody (1: 10000 dilution; Santa Cruz Biotech). Membranes were then incubated for 1 hour at 37° C. in Lumi-Light Western Blotting Substrate (Roche Applied Science) followed by a 10 minute exposure on a BioMax MR Film (Kodak).

**[0427]** A 70 kD band corresponding to the expected size of the P2X7R protein was detected in HT-22 cells demonstrating expression of the endogenous mouse P2X7 gene (**FIG. 17**). A second mouse hippocampal cell line (HT-39) did not express P2X7.

**[0428]** Since P2X7R is an ATP-gated ion channel which allows the entry of calcium and sodium ions into cells, a method for identifying P2X7R agonist was established by monitoring calcium influx into HT-22 cells. Cells were first loaded with the fluorescent dye Oregon green AM ester (Molecular Probes) for 30 minutes at room temperature, washed 2 times with DMEM/10% foetal calf serum to remove excess dye and cultured for 15 minutes in the presence of 100  $\mu$ M 2- and 2'-3'-O-(4-Benzoylbenzoyl)adenosine 5'-triphosphate (BzATP: C<sub>24</sub>H<sub>24</sub>N<sub>5</sub>O<sub>15</sub>P<sub>3</sub>). BzATP is a known agonist of P2X7R (North and Surprenant, Annu. Rev. Pharmacol. Toxicol. 40 (2000), 563-580). Calcium movement into the cells was visualised under a fluorescent microscope with a fluorescein filter (wavelength 492/517 nm). Oregon green AM ester is a fluorescent dye that binds to intracellular calcium. Accordingly, an increase in green fluorescence was observed in cells treated with BzATP (**FIG. 18**) signalling an activation of P2X7R which results in an influx of calcium into the cells and an increase binding of Oregon green dye to intracellular calcium. Alternatively,

Oregon green AM ester can be replaced by Fluo-3, fluo-4, fluo-5F, fluo-5N, fluo-4FF, Fluo-4 dextran, Fluo-3 AM, Fluo-4 AM, Fluo-5F AM, Fluo-5N AM and Fluo-4FF AM (Molecular Probes). Calcium influx in HT-22 cells can also be measured in 96-well and 384-well microplate using the Calcium Plus Assay Kit (Molecular Device) or FLIPR® Calcium Assay Kit for Fluorometric Imaging Plate Reader Systems (Molecular Device). HT-22 cells can be replaced by any cells expressing P2X7R, including cells that have been genetically modified by introducing an exogenous P2X7 gene.

[0429] The specificity of the P2X7R agonist on calcium influx was confirmed by pre-treatment of HT-22 cells with 100 mM Oxidized ATP (oATP; Sigma) for 1 hour before the addition of BzATP. oATP is an irreversible inhibitor of the receptor (Chen et al., *J. Biol. Chem.*, 268 (1993), 8199-8203). Activation of P2X7R by the agonist was inhibited by oATP (FIG. 18) as illustrated by the absence of green fluorescence in the cells.

[0430] Yet another method of measuring P2X7R activity involves the entry of ethidium bromide into P2X7R expressing cells. Activation of P2X7R by an agonist allows the entry of ethidium bromide which binds nuclear DNA and emits a fluorescence signal. Alternatively, the propidium dye YOPRO-1 can be substituted for ethidium bromide. An increase in fluorescence can be used as a measure of P2X7 receptor activation. Therefore, the assay can be used to test and quantify the effect of an agent or compound with agonist properties on P2X7R. In the present example,  $10^3$  HT-22 cells were seeded per well in a 96-well flat bottom microtitre plates and incubated at 37° C./5% CO<sub>2</sub> in DMEM medium containing 10% FCS until the cells attached to the culture surface. Once attached, cells were incubated for 60 minutes in DMEM medium containing 10% FCS,  $10^4$ -M ethidium bromide and increasing concentrations of BzATP (1  $\mu$ M, 10  $\mu$ M, 100  $\mu$ M, 500  $\mu$ M, 1 mM). The number of fluorescent cells which have integrated the ethidium bromide to the DNA can then be counted using a fluorescent microscope (Zeiss, Germany). Concentrations above 100  $\mu$ M BzATP increased the number of fluorescent nuclei signalling activation of P2X7R (FIG. 19a). Alternatively, ethidium bromide fluorescence can be measured using a Perkin-Elmer fluorescent plate reader (excitation 520 nm, emission 595 nm, slit widths: Ex 15 nm, Em 20 nm). From the readings obtained, a pIC<sub>50</sub> figure can be calculated for each candidate agent or compound. Accordingly, a P2X7R agonist is defined as an agent or a compound with an EC<sub>50</sub> equal or below 300 micromolar, whereas the term EC<sub>50</sub> is defined as the concentration eliciting 50% of maximal response to an agonist (North and Surprenant, *Annu. Rev. Pharmacol. Toxicol.* 40 (2000), 563-580). The specificity of an agonist for P2X7R can be evaluated by pre-incubation of the cells for 60 minutes with 100  $\mu$ M o-ATP before adding the agonist and ethidium bromide dye. Under these conditions, activation of P2X7R by the agonist is inhibited by oATP resulting in a reduction in the number of fluorescent cells (FIG. 19b).

[0431] Yet another method for identifying P2X7R agonists was devised by generating a immortalised mouse cell line that overexpresses the human P2X7R gene under the control of the human cytomegalovirus (CMV) early promoter/enhancer region. The human P2X7R cDNA was inserted into the pcDNA3.1 vector (Invitrogen) and transfected into the mouse hippocampal cell line HT-22 using Lipofectamine

(Invitrogen) according to the manufacturer's specifications. One day after transfection, culture medium containing 500  $\mu$ g/ml G418 was added to the cells. Resistant clones were separately isolated and cultured 14 days after applying the selection medium.

[0432] The agonistic activity of a compound was evaluated by measuring calcium entry in the cells that overexpress the human P2X7R. Cells were cultured in 96 well plates and incubated at 37° C. with 5% CO<sub>2</sub> DMEM with 10% foetal calf serum (Gibco) until they reached confluence. Cells were then loaded for one hour with 10  $\mu$ M of Fluo-4 AM (Molecular Probes). Fluo-4 AM is a fluorescent dye that binds to intracellular calcium. After loading, cells were washed once with a buffer containing 0.5 mM CaCl<sub>2</sub> and 20 mM Hepes and were treated with 20  $\mu$ M BzATP or 50  $\mu$ M tenidap. Agonist activity was detected by measuring an increase in calcium influx which results in increased binding to Fluo-4 AM and increased fluorescence. Changes in fluorescence signal are measured using a Fluostar Optima plate reader (BMG biotech). Both BzATP and tenidap produced a rapid increase in fluorescence intensity which declined slowly over time (FIG. 19c). Thus, both compound stimulated the activity of P2X7R which results in an influx of ions into the cells.

#### EXAMPLE 9

##### Activation of P2X7R With Agonists Has Antidepressive Effects

[0433] To demonstrate that activation of P2X7R has therapeutic effects on affective disorders, the P2X7R agonist BzATP (2'-3'-O-(4-Benzoylbenzoyl)adenosine 5'-triphosphate (C<sub>24</sub>H<sub>24</sub>N<sub>5</sub>O<sub>15</sub>P<sub>3</sub>)) was administered to a selected DBA/20la mouse strain that displays characteristics of being highly anxious, responding to antidepressants, and showing anxiolysis after subchronic antidepressant treatment (Lucki et al., *Psychopharmacology* 155 (2001), 315-322). BzATP is a compound with strong specificity to P2X7R (North and Surprenant, *Annu. Rev. Pharmacol. Toxicol.* 40 (2000), 563-580). In the present example, the P2X7R agonist was directly injected into the hippocampus of mice. However, a P2X7R agonist agent or compound could also be delivered orally, subcutaneously, intravenously, intra-arterial, intranodal, intramedullary, intrathecal, intraventricular, intranasally, intrabronchial, transdermally, intrarectally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, or intraocularly.

[0434] Four months old male mice were bilaterally implanted with guide cannulae (23 gauge, length 8 mm) by means of a stereotactic instrument (David Kopf Instruments). The coordinates, in relation to bregma, were -2.0 mm posterior,  $\pm$ 1.0 mm lateral, and -1.0 mm ventral. After surgery, the mice were allowed to recover for 10 to 112 days. Following this recovery period, mice were injected with 1  $\mu$ l vehicle solution (0.5% DMSO, Sigma) or 50  $\mu$ M BzATP (Sigma, prepared in 0.5% DMSO) in each side of the brain over a period of 60 seconds. Injections were performed using a 9 mm-30 gauge needle inserted into the guide cannulae and connected via tubing to a 10  $\mu$ l Hamilton syringe.

[0435] The behaviour of individual mice was assessed using the Porsolt's forced swim test 24 hours after injection



of vehicle solution or BzATP. A pre-exposure of 5 minutes to the test was done 10-15 minutes after vehicle or BzATP injection. The forced swim test is a standard test that measures primary stress-induced reductions in avoidance or escape, termed behavioural despair. The test is used to determine the effectiveness of antidepressants, test new pharmaceutical compounds and validate animal models of depression (Porsolt et al., *Arch. Int. Pharmacodyn.* 229 (1977), 327-336; Porsolt, *Rev. Neurosci.* 11 (2000), 53-58; Rénier et al., *Behav. Brain Res.* 136 (2002), 521-532; Page et al., *Psychopharmacology* 165 (2003), 194-201; Kelliher et al., *Psychoneuroendocrinology* 28 (2003), 332-347). The test consists of placing a mouse for a period of 5 minutes into a glass cylinder containing water. Under such circumstances, the mouse cannot touch the bottom of the cylinder and is thus forced to swim. Time, latency and frequency of struggling versus floating are scored as behavioural parameters. Floating (i.e. movements made only for keeping balance and breath) is a passive behaviour associated with despair and represents a depressive-like symptom since the animal does not make any effort to actively cope with the stressful situation. Increased struggling (i.e. active attempts to escape) indicates active coping behaviour that can be interpreted as an improvement of depression-like symptoms. For example, treatment with serotonergic antidepressants reduce the total time spent floating (Borsini, *Neurosci. Biobehav. Rev.* 19 (1995), 377-395; Redrobe and Bourin, *Psychopharmacology* 138 (1998), 198-206, and in parallel increases the time of active behaviour (i.e. swimming or struggling; Lucki et al., *Psychopharmacology* 155 (2001), 315-322).

[0436] The P2X7R agonist BzATP was found to increase active escape attempts (i.e. increase in time and frequency of struggling, decrease in latency of struggling) while a decrease in passive behaviour (i.e. decrease in time and frequency of floating, increase in latency of floating) was measured when compared to control mice injected with vehicle solution (FIG. 20). Observed results were verified statistically using Mann-Whitney U and one-way MANOVA tests. The differences in time struggling, latency of floating and frequency of floating were found to be statistically significant. While latency and frequency of struggling and time floating results were not supported statistically, they still represented a tendency towards improvement in stress coping behaviour. These results demonstrate that a P2X7R agonist can lead to improvements in depressive-like symptoms.

[0437] Since conclusions drawn from the forced swim test can be influenced by unspecific effects of an agent or compound on animal activity (i.e. increase in struggling behaviour can be the result of hyperactivity instead of increased active coping behaviour), the potential effect of BzATP on locomotor activity was assessed by the open field test (Crawley "What's wrong with my mouse: Behavioral phenotyping of transgenic and knockout mice", Wiley-Liss (2000)). Locomotor activity in mice treated with control vehicle solution or 50  $\mu$ M BzATP was assessed 24 hours after injection by placing individual animal in a dark-grey wooden box (30x30x40 cm). Locomotor activity was monitored for a period of 30 minutes using a video camera. Overall distance travelled by the animals during the testing period was then analysed by means of VideoMot2 computer software (TSE GmbH, Bad Homburg). No difference in locomotor activity was measured between mice treated with control vehicle solution and BzATP (FIG. 21). Therefore,

the application of BzATP did not induce hyperactivity. These results confirm that activation of P2X7R by an agonist agent or compound leads to improvements in depressive-like symptoms and is not the result of an unspecific effect on animal activity per se.

[0438] Several reports suggest that activation of P2X7R can induce apoptosis and cell death in vitro (Di Virgilio et al., *Cell Death Differ.* 5 (1998), 191-199, Virginio et al., *J. Physiol.* 519 (1999), 335-346). To test whether P2X7R activation in the hippocampus resulted in cell death, apoptosis levels were quantified in the brain of the mice treated with BzATP. Mice were sacrificed at the end of the behavioural experiments, the brains were removed, shock frozen and sectioned into 16  $\mu$ m slices. Brain sections were then studied for apoptosis using the DeadEnd fluorometric TUNEL system according to the manufacturer's recommendation (Promega Corporation). The TUNEL system measures the fragmented DNA of apoptotic cells. Positive control for the assay are made by pre-treating brain sections for 10 minutes with 1 unit/ml of DNase I.

[0439] Very few apoptotic cells (i.e. less than one cell per brain section) were observed in brains of mice treated with control vehicle or the P2X7R agonist (FIG. 22) when compared to positive control sections pre-treated with DNase. Moreover, no significant differences in the numbers of apoptotic cells was observed between the control animals and mice treated with BzATP, indicating that activation of P2X7R did not result in cerebral cell death in vivo.

#### EXAMPLE 10

##### b 2X7R Antagonists Have No Antidepressive Effects

[0440] The P2X7R antagonists KN-62 (1-(N,O-bis[5-isoquinolinesulphonyl]-N-methyl-L-tyrosyl)-4-phenylpiperazine) and oxidized ATP (oATP) were administered to DBA/201a mice (Harlan Winkelmann, Germany) that exhibit the behavioural characteristic of being highly anxious. KN-62 has been shown to be a non competitive antagonist of P2X7R (Chessel et al., *Brit. J. Pharmacol.*, 124 (1998), 1314-1320) while oATP acts as an irreversible inhibitor of P2X7R (Chen et al., *J. Biol. Chem.*, 268 (1993), 8199-8203).

[0441] In the present example, the P2X7R antagonists were directly injected into the dentate gyrus region of the hippocampus. Briefly, three months old male mice were bilaterally implanted with guide cannulae (23 gauge, length 8 mm) by means of a stereotactic instrument (David Kopf Instruments). The coordinates, in relation to bregma, were -1.5 mm posterior,  $\pm$ 1.0 mm lateral, and -0.8 mm ventral. Mice were allowed to recover for 10 to 13 days after surgery. Following this recovery period, mice were injected with 1  $\mu$ l vehicle solution (0.01% DMSO, Sigma), or 100 nM KN-62 (Sigma, prepared in 0.01% DMSO), or 10  $\mu$ M oATP (Sigma, prepared in PBS) in each side of the brain over a period of 60 seconds. All injections were performed using a 9 mm-31 gauge needle inserted into the guide cannulae and connected via tubing to a 10  $\mu$ l Hamilton syringe.

[0442] The behaviour of individual mice was assessed using the Porsolt's forced swim test 24 hours after injection of vehicle solution, KN-62, or oATP. A pre-exposure of 5

minutes to the test was performed 15-17 minutes after administration of vehicle, KN-62, or oATP. A description of the Porsolt's forced swim test is given in example 9. In the present example, no changes in active escape attempts (i.e. time, frequency, latency of struggling) or in passive behaviour (i.e. time, frequency, latency of floating) was measured between vehicle, KN-62 or oATP treated mice (**FIG. 23**). Observed results were verified statistically using one-way MANOVA test. The differences seen in the different parameters between vehicle, KN-62 or oATP treated mice were not supported statistically. These results demonstrate that P2X7R antagonists do not improve depressive-like symptoms and have no antidepressive action.

[0443] Since conclusions drawn from the forced swim test can be influenced by unspecific effects of an agent or compound on animal activity (i.e. increase in struggling

behaviour can be the result of hyperactivity instead of increased active coping behaviour), the potential effect of the P2X7R antagonist oATP on locomotor activity was assessed by performing the open field test. Locomotor activity in mice treated with control vehicle solution, 10  $\mu$ M oATP, or 50  $\mu$ M oATP was assessed 15 minutes after injection by placing individual animal in a dark-grey wooden box (30×30×40 cm). Locomotor activity was monitored for a period of 30 minutes using a video camera. Overall distance travelled by the animals during the testing period was then analysed by means of VideoMot2 computer software (TSE GmbH, Bad Homburg). No difference in locomotor activity was measured between mice treated with control vehicle solution and oATP (**FIG. 24**). Therefore, the application of a P2X7R antagonist did not induce hypo- or hyperactivity in the animals.

---

#### SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 111

<210> SEQ ID NO 1  
 <211> LENGTH: 56580  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: exon1  
 <222> LOCATION: (3000)..(3124)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:  
 <221> NAME/KEY: exon2  
 <222> LOCATION: (24841)..(25009)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:  
 <221> NAME/KEY: exon3  
 <222> LOCATION: (26134)..(26202)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:  
 <221> NAME/KEY: exon4  
 <222> LOCATION: (30958)..(31030)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:  
 <221> NAME/KEY: exon5  
 <222> LOCATION: (32481)..(32577)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:  
 <221> NAME/KEY: exon6  
 <222> LOCATION: (35416)..(35496)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:  
 <221> NAME/KEY: exon7  
 <222> LOCATION: (36113)..(36242)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:  
 <221> NAME/KEY: exon8  
 <222> LOCATION: (37541)..(37677)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:  
 <221> NAME/KEY: exon9  
 <222> LOCATION: (45470)..(45560)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:  
 <221> NAME/KEY: exon10  
 <222> LOCATION: (47229)..(47295)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:  
 <221> NAME/KEY: exon11  
 <222> LOCATION: (47380)..(47529)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:

-continued

<221> NAME/KEY: exon12  
 <222> LOCATION: (50438)..(50539)  
 <223> OTHER INFORMATION:  
 <220> FEATURE:  
 <221> NAME/KEY: exon13  
 <222> LOCATION: (54392)..(54889)  
 <223> OTHER INFORMATION:

<400> SEQUENCE: 1

tgtggtccca gctactcagg aggctgtggt agggggatca ctggagccca gaagtttgaa	60
gctgcagtga gctatgattg cgccactgca ctccagcctg gacgacagag caagacccag	120
tctctaaaaat aaattaatta aattaatttt ttaaaaagaa agataagata aagagtcata	180
gaagtacaaa tgagaaacca ggagaacatg ggacccagga aagtcatgga aaccaaggga	240
aaatactggt ccaagatgca gaaagtagcc agtgatgttg aatgcaccca agagatcaag	300
taacataagg aataaaatat atccaccag tctggcaact gagaagtctt tggtaacctc	360
attgagaaca gaatccatgg agtggaggag gtagcagtga tgaatatagcg gacaagggct	420
gaaggagtgg agaggggaga gcatgaggga ccagtgcaag agtgcagcag aaagacaaca	480
aacaggccgg gtgtagtggc tcacgcctgt aatcccaaca ctttggtagg ctaaggcggg	540
cagatcgctt gtgaccagga gttcgagacc aacctagcca acatggtgaa accccatctc	600
tactaaaaac acaaaaatca gctgagcgtg cgcatggctg taattgcaga tactcatgag	660
gctgaggcac aagaatcgct tgaaccacag aggcaggggt tgcagtgacc caaagtcttg	720
ccactgcact acagcctggg caacaggggg agactccatc tcaaaaaaac aaaaaaaaaa	780
aagaaagaaa gaaaagacaa agaaagaaa agaaagaaa agaaagaaa aaagaaagaa	840
agaaagaaa aaagaaagaa agaaagaaa agaaagaaa aaagaaagac aagaaaatag	900
atgtctttga gtggtagcaa gagcaacct gaagtgattt gcctattctg gacatttcct	960
ataaatggag tcgtaaaaga tgtatccttt tgtgtctggc ctctttcact taagcgtagt	1020
gttttcaagg ttcattccatg ttggagcatg ggtcagtggt tcattccttt ttatggctga	1080
ataatattcc ctgtatgga ctttcgtgct tttgtgtatc catccatcag ctgatggaca	1140
tttgacttgt ctgcctgttc actgccatat ctactattcc tggtgctctt gggaatgagt	1200
gaatgaatag cataaacaga gctgtccaag gtcacagagc tggtaagatg tgaagccagg	1260
atcggaagcc aggccattag tcccctagag cctatgttct aagcatcagg ctttacctgt	1320
gaatctcctc tttttacaga tgaagatgac tgtatcactc agattcccgg caggaaagca	1380
atggcatact caagtggggt aactaatgat ggaaccattt acaaggtgt ggacagagtt	1440
aagaaaaagc aataggagat agtgagcttc ctggggctgg taagagtggg gagcccttac	1500
cactcccagg actaaaggag ggagtgggct ccagaagccc tgcctatatg caactgagaa	1560
gggcagggcc agggagtac gtccatcctc actgctctcc agtctcctga actggaagcc	1620
agaaggtgag gggaaccctg atgcagtttg tatgtgtgag aaagtacaat tagtttagac	1680
tgaaaaactg aaaatctacc cggccactta gcaggctgga ataacagaaa tggatcaagc	1740
cagctgtaaa gataacaggg aacaataatt ctctgtagct gtaaagtgat aatacaaccc	1800
tgcactcttg agtgactgct gaaacattgt cctttaaaat cagagacctt cagaaaactc	1860
gctgtttgaa attacatgac taagactgaa atattccaat tttgcctgga agatttaagt	1920
catcttgaca cagagaagca gcctcaattt acaactcagg agcagagctt cagataaaga	1980

## -continued

ttttctggac acatttgaca tgtatcttag ctatgttgct tcctaggaaa cagggccctg	2040
ggctcctctt gcaatccaga ctgaagtga ctgctttgta caaacctgtt ttgctttgag	2100
tccatcaaaa catgacttca tttagatttt atctcaactc cactttcctc ggaatcctat	2160
actaaattgc tgttttcctt tgtttggtga tgtgcgtagc tcttctgggtg ggtggtgtcc	2220
ctcactgaat aggtcaacaa acctaacctt gttggactgc cactgtgtcc ctggtgatct	2280
ttggctgatt ggtctaggtc atagatcgac ctgccggggt gcagaggagg gtggagagta	2340
actcagaggg tcaagcatga aagatctggc agaaaaataa agcccctcca cccccaccac	2400
ccctaccctt gcaaactcga tttccccac caactgcaga ccagagtatt ataagggcg	2460
gtggaagaag agggggagat cttcatttac ccagagctcc tatacatcag gggctgaata	2520
aagggttgta gaaatgaatg aatcaatctc tgagtggggc ttcaggcagt ggaagatct	2580
cagtcctttt ctgaggcata atggaagctc ccagtcttgt gacatttgca aggctgcccc	2640
tttctcccaa gagacatgag accaaaaaag tgaaaggaaa ggggggaaa gggagaattc	2700
tgaaaatgcc catcctctga acaccatctt tgtgtaggca tctgggggag gccagctggg	2760
gtgaggtcat ctgccagcca ggcccgtagg acttggcgct tcttgtttat cacagccaca	2820
tgtggggcca ctgccagggc cgcgcccaac tctgcagtca ttggaggagc ttgaagttaa	2880
agactcctgc taaaaaccag tacgtttcat tttgcagtta ctgggagggg gcttgcgtgtg	2940
gccctgtcag gaagagtaga gctctggctc agctccgcgc agggagggag gctgtcacca	3000
tgccggcctg ctgcagctgc agtgatgtt tccagtatga gacgaacaaa gtcactcgga	3060
tccagagcat gaattatggc accattaagt ggttcttcca cgtgatcatc ttttcctacg	3120
tttggttaagt gggatctggg gaggaccag atctctgcag tggccgacag cacagaaagc	3180
cccagcgggc agcttcagggt gcacattctg aatctcacat ggttttcgaa tctgagacgt	3240
gctctcacag ccagctgggc gggaggagg aagcagcagc aggcagagg aaacggtgcc	3300
aggctgcagc agagagaagc cacaggacaa gcgggattcc tttctgctct acttcaggcc	3360
cgccagggcg cgcaggcgag ggcgtgcctg gggaggttag gaaagcgag ggcaacaccc	3420
tggatcccca gggaggagc gaggatctca gggcagcctt ggtgatcatg ctggcatctg	3480
agtcaccatg cttgggagga ataggaccag gcttgaaaat gtgttataac tttaggtcct	3540
caccaacgtc aggaaggccc tgcttttttg tttttgttct tctaaaaga aacttactga	3600
gatataatth atacaccata caattgaccc atttaaaggg taccatttaa tgattttcag	3660
attattccca gaggttgtga accatcccca aaatcaattt tagaatattt taatcgactc	3720
aaaaggaaac ccacactcct tcaccatcat ttccaacact gttcctcctc cttcccaccc	3780
atcaatttcc tttctgcctc tatggatttg cagattcttg acatttcata taaatggaat	3840
cacataatat gtggctcctt gtggcactta gcgtgttttc aatgctcatc catgttgtag	3900
catgtgttga tacttcattc aatttttttt tttaaagaga cggggtctca ctattttgtc	3960
caggctggtc tcaaactcct ggactcaagt gatctgcctg cctcgccctc ccaaagtgtc	4020
aggattacag gcgtgagcca ttgcacccgg ctgatacttc attccttttt acggctgagt	4080
agtactccat tgcatggata gaccactttt ttctatccat tcatccattg atggacattg	4140
gggtgtgttc tctttttttg ctatcatgaa taatgccaca tgaacatttg tgtacaaggt	4200
tttatgtgga tatatatctt cctttctctc gaatatgtac ctaagagtaa aaattgctag	4260

## -continued

gtcatatggt aactatgttt cacctttggg gggaaatgtg agctgaattt cacagcagct	4320
gcagtttttt acattcctat cagcagagta tgagggatcc aatttctcca catcctcacc	4380
aacgcttggt atcgtctgtc tttttgggtt ttgttggtgt ctttttggtt ttgtctttga	4440
gatgaagtct tgctctgttg cccaggtggt agtgcagtgg cgcaatcttg gctcactgca	4500
acctccacct ccccggttca agcgattctc ctgcctcagc ctcatgagta gctgggatta	4560
cagggtgtgc tcaccactcc tactaattt ttgtattttt agtagagatg aggtttcgcc	4620
atgtaggcca ggctggtctc aaactcctga cctcaagtga tccgccacc ttggcctccc	4680
aaagtgtggt gattacaggc atgagccgcc acgcccggct gattgtctgt cttttttatt	4740
atagccatgc tagtgggtgt gaagtggtag ttcattgttg ttgtgatttg aatttccctg	4800
atggtgagtg cctcttattc tctgtgctga ttgataatga tgatgaaggc aatttgtatc	4860
tatagagtgg cagtgtagtt tactaagagt tagggtaact tatttcatag tactggctat	4920
gtcttctggg ccaagtcatt aacttctctg agcctcagtt tctgcactctg ttcataggg	4980
tggtgcaatt aacccaaaaa aaaaggcatg aacagccctt atcatgatga ctgacatagg	5040
ataagagctc cataactagt atctattttt aaaaataatc tttttaagtc tgggagtgg	5100
ggctcacacc tgtaatccca acactttggg aggccgaggc gggtggtatca cgaagtcagg	5160
agtttgagac cagcctggcc aatatggtga aaccccatct ctactaaaaa tacaaaaatt	5220
agtggggagt ggtggtgcac acctgtaatc ccagctacta gggaggctga ggcaggagaa	5280
tcgcttgaa cgggaggcgg aggttgagct gagccgagat caagccactg cactccagcc	5340
tgggtgacag agcaagactc catctcaaaa taataataat agtaataatt tttttgatta	5400
tataatagta tatatgtata taaaatacat gtatgtattt ttatctatat cctctgctct	5460
gacctcaaa gtaaccacgt ccaagttcag gatttgaaat ctggaaacgt ggattcaaaa	5520
atccttcacc tctttgagcc ttggtttcat catctgtaaa atggggagaa ttgttgatag	5580
gaatattaaa tgaactaata aatgcaaagc tgtttgagaa atatatggca tatagtaatc	5640
cctgattaag tgtagttctt tattattaat aatgctatta ttaggattat tattattcga	5700
ttcatatggt tactgttcaa caaatattga atgataaaca tatatgctgg gtcgggcatg	5760
gtggcccatg cctgtaatto cagcactttg ggaggccaag gcgggcaggt cacttgaggt	5820
caagagtttg agaccagcct ggccaacgtg gtggaaactc cgtctgtgct aaaaatacaa	5880
aaattagccg ggcattggtg tgggtgctg taatcccagc tactcgggag gctgagacag	5940
gagaatcact tgaaccagc aggtggaggt tgcagtgagc caagattgca ccactgcact	6000
ccagcctgag ccacagagca agactctgtc tcaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	6060
tatatatata tatatatata tatatatata tagtattttt agtagagatg gggttttgcc	6120
atctcttata tttttttata tttttatatt ttttatatat ttatatatat ttatatgtat	6180
tttatatatt tatatatatg tgctgaaatc taaaataaga catatcactt aaaccagcta	6240
tcacaggacc aatggtaaa agcataatta gtaaatgccc ccttctgcat taatacgtag	6300
attgtatgtc acaatttcat gtagcaggtg atggggcttc ctgacattct tgggcccagc	6360
atacaatctt tttatttctt ttttagagac aggttctcat tgtgttgccc taggctggag	6420
tgacgtgtca ttatcacagc tcgccacagc ctcaacttct caggctcaag tgatcctccc	6480
gcctcatcct cccaagtagc tgggaccata ggcacacgcc atgatgtctg actaattttt	6540

## -continued

tat	tttttttt	tgtggagatg	ggggtctccc	tatgttgetc	aggctagtct	caaactcctg	6600	
gact	caagtg	atcctcctgc	tttgccctcc	caaagtgctg	ggattatagg	catgagccac	6660	
tttgt	ctggc	ctacaattcc	ttttctttct	ttctttcttt	tttttttttt	tttttgagtt	6720	
gagt	ctcact	ctgtcaccta	ggctgtagt	cagtgggtg	atctcagctc	actgcaacct	6780	
atac	ctccca	ggcccaagca	atcctccac	ctcaacctcc	cgagtagctg	ggaccacagg	6840	
tgtac	accac	cacgcccagc	taattttttg	tatttttggt	agagacagg	tttcaccata	6900	
gtgt	gccagg	ctggctctga	actcctgagc	tcaagcaatc	cacccgcctt	ggcctcccaa	6960	
agt	gctggga	ttacagggtg	gggtcaccat	gccagcctt	caatttcttc	agattattgt	7020	
ataa	taataa	aatttctaaa	aacagattac	tatgagaaca	ccttgtaatg	ataaatattt	7080	
ttct	tttcaca	tggtgaaatc	ataacttaca	tgtatttaac	tccctcaa	atcagttatac	7140	
atgt	tccatg	gggacatgga	gagagagaga	acaatataaa	taatgcaatg	attcagcaac	7200	
cggg	ttcgc	gtgagtaggtg	tgagatggga	tactggcatc	tgggtgtccc	tccgtcaatc	7260	
tctg	ttccca	cgtgcttttt	agcgagcatc	ttgctgctct	acttattaat	ctaggtacat	7320	
at	ttttcagc	cacatgaccc	ctaaaataag	tcaactcctt	caccttcacc	tggagcaaaa	7380	
ctg	taaaaa	agcaacttgt	tctagatctg	gaggcaaaat	tggttccagc	agaccactg	7440	
gga	agtgga	ggtgtttgct	atgctttgtg	gacaatttaa	gtgtttaaaa	ggcaattttg	7500	
atg	agccatg	at	tttcatct	taagtgccaa	ctttagaaga	taaccttc	atataa	7560
ct	tacagtt	tgtattttaa	aataataaa	at	ggaagaaaa	actatcacca	gctttgcttt	7620
tt	ctctctg	gttaattggac	tatctattaa	cgtctcacta	aatggccaaa	tggagtctca	7680	
cag	tacacag	tttgggaaat	gctgccttaa	gttgcctctg	gaagccttaa	gttcgtgtaa	7740	
cc	atgactta	gccatctgtg	aaatgggtag	acccatgggg	cataggaatt	ggggaatgta	7800	
tt	ttgcatca	agtaggactg	gtgtggacca	caggtggaaa	tccagacagg	ctcgagcaat	7860	
gt	tcaggata	cagcagcaag	tgacagggtc	tggccagctg	tccctactgg	ggctgcaagg	7920	
act	aagagcc	aggataagga	gacagggcct	cggttattga	aaatgggata	caaagcagg	7980	
act	ggttatg	gcggaaatta	accaagaggc	tggttcagctt	cttctattct	cacaaccaat	8040	
tc	agatctg	attcaaaggc	agggatgcag	ctacaagagg	agtttgtgtt	ggaggctctg	8100	
g	tagatagga	tgctaagcag	gtctgtgtca	gccaggatcg	tttgggcagt	aagtgcaga	8160	
a	acccaactt	acgttaacca	caaagcctca	aaggggaatt	tattggatca	tgtaactggc	8220	
c	agttgaaga	gatgggtgctg	tcctcggacg	tggatgaaga	taagagttca	aactgtgtca	8280	
tc	aggacgga	tccgtctccc	tacatcctgg	ctccatttgc	ttctatttgg	cttccttctt	8340	
ga	acaactct	gtatgtacaa	gatatcctca	cattctcttc	ttttaacaac	ctcagaggag	8400	
caa	acaagtc	tttcctgata	gctctgctaa	gaaagtccca	ggaagactc	tgattgatca	8460	
gat	gtgagtt	acagatccat	ccctgaacca	atcactgtgg	ccagggccaa	ggtgtcatcc	8520	
cat	tgccag	cctgggtcat	gtgctggctc	cogtagcagg	aagtagggag	aagacgagg	8580	
ag	cgtgtga	gccctacgtg	aaccacatcc	agtggatttt	ccttaggaag	gaagattctg	8640	
tg	accagacc	aaatggaaaa	ggaagtgggg	agggtaggca	gacaaaaatc	ggtgatctat	8700	
t	acaaggtct	ttccaaaagc	cagaaattag	ggatttctaa	gtgtacctct	ctagtcctgg	8760	
ct	tgtctcag	ccctgcagct	gggctgtgga	gactccatca	gcaattgctg	accccctatt	8820	

## -continued

cttacctcgg ccccttcctg agtcccacca ggagccctgc tcctcatctg ggacatgctt	8880
cttggcccat ggaggttgcc agctcacett ggtactttac tactatttgg tttctgtgac	8940
caccaagctg ttgcttaaac caagccgcct atggtggcac cactttggag aattcttttt	9000
tattttttga gacaggttct cactctggtt accgagactg gagtgcagtg gtatgatctt	9060
ggctcactgc agcctcgacc tcccgggttc aggtgatttt cccacctcag cctccccagt	9120
agctgggact attggtgcat gccactacac ccggtcatt tttgtatttt tagtagagac	9180
gggggtttgc catgttggtc gggctggtct tgaactctg acctcaggtg atctgcccc	9240
ctcgccctgt caaagtgtg aaattacagg cgtgagccac cgcacccggc tggataattc	9300
tttagatttt attccaagt gggtgagaag ccattgggaag attgtaagca aaggaataaa	9360
agttggttg attctctttg gaggatttcc ctcttccctg gagaccagtc ctacagaaag	9420
cttttccaat ggccacagaa gactaaggaa gattgagaga ggcacaggaa acaattccac	9480
tcagagctgc tgcttcgcca ctgccagtc agcctccact ggcgccaaca cgagcctatg	9540
ctgtcattcc tggggccagg ctggctgtcg ctgtggagat ggtgctgctg tgctagtttg	9600
tggctgttaa aatggagctc ctggctggac acctgtttcc cagtgggtgg ccacaaacct	9660
gtgagtctg gcctcctcca gctgagtaga agtgagccca gcacagaggc cggggaggcc	9720
ccagagccca ggaggagcaa atccaaatcc aacccaagca tggccctct gacctgacc	9780
atcttttttt ttttttttt ttttgagaca gagtctcct ctgtcaccga gactggagt	9840
cagtggcacg atcttggctc attgcaacct ccacctcctg ggttcatgctg attctcctgc	9900
ctcagcctcc ctagtaggga ctacaggcat gcaccactac acccagataa tgtctgtatt	9960
tttagtagag atgggggttct gctatgttgg ctaggctggt ctggaactcc tggcctcaag	10020
cgatccatcc gtcttggctt cccaaaatgc tgagattaca ggtgtgagcc actgcgccca	10080
gcctgcccc gactgccttc agaaaagtag aagcaggagg tctgccacat tctcatccca	10140
tccttgacac ctgggtgatg tgctagagct tgaggggaag gtcagaggta acccagttcg	10200
ggaagtgacc ccgtaggagg agaagaatcc agccacgtga cacacagtac gaagagaaca	10260
gtaaagtcaa agacctgaa acgaggaaca gcaaggacag catgtcccag aacctgggga	10320
gagaggagg agcagccgga gctgagggca gctcagggga agagttgcta tttgattttc	10380
agtgcattgg aagccactga aaagtattat gcagagaagt aaaaccatct cactcatggt	10440
tctagaagcc ttatgggcca ctgtgcagag catagagtgt ggtgggacaa gaatgacggt	10500
gaggaaccag tgagaagggt cctggagtca ccagatgcga ggagatagag gcttgggcta	10560
gaatgggctg gcggagctgg agagagggtg acaagctga gatgtgtttt ggaggcagaa	10620
atgatagatt tgccaggagg tgaaagatgg ctctagattc ctgaggaacg atgatgcctt	10680
ttaccaacct gaagaagacc aaggagaggc ttgaggacta acacggaact gtgaaggagg	10740
ccaggcaaa gtggagactg ccgtgagctg tgatcacgcc actgcacct cgcctgggtg	10800
acagactgag acctgtctt aaaatatata tgtaataaaa aataataaat aaagggccag	10860
gcgtggtggc tcaacgcctg taatcccaac actgggaggc tgaagcaggt ggatcacctg	10920
aggtcagaag ttcaagacca gcctggtcaa catggcgaaa tcccatctat aataaaaaa	10980
caaaaattag ccagacatgg tggcacatgc ctgtaatccc agctacttgg gaggtgagg	11040
caggagaatc acttgaacct gggaggcaga ggttgcatg agcctaaatc acaccactgc	11100

## -continued

actccagcct	gggcaacaga	gtgagacccc	ataaataaat	aaataaatac	ataaataaag	11160
gatatcaggc	aagcaattaa	aacacctgcc	tactaccatc	tatccgcac	ctttttttca	11220
tgtctaacac	caaaacacca	aaatagctgg	aaacatatca	tgtcaaaaga	taagcatgat	11280
tctttcaaat	gaatggctgt	ggacttatga	aaagctgcct	ccgcgtctga	ttttcttggt	11340
tcaaggggta	atgatggagg	tttctctcct	gcacctgcag	gcctccgggg	atgctgcccc	11400
agtgcagcct	cccaccttg	gctggagcct	cccagccccg	ctgcatgggt	agcacagtgt	11460
tgggaagagg	aaggagcagt	ttgtctggct	ttttcacagg	cccacagctg	gccgggtacag	11520
tggcagagaa	ggcgaggctg	aggtggggct	ggggaagcca	gcaggggcct	gatcatggag	11580
cgccttgagc	tgcatccac	aggtactgag	aaagcatgga	gagctgtccc	ccagccccca	11640
gcactcttatt	agaaaatatt	gaaacataac	agaaaacatg	aagggtttta	tgttcatatt	11700
ttcatcttag	agagatcatt	ctgcttgaaa	tataaacct	actgagtact	gctgaggact	11760
tacctgtgc	cgggttctgt	tcttagcatg	tgacatgtat	tggcctatct	gattctccta	11820
cataatcctg	gaagattggt	tgtgggagtg	ggtgatcatg	acctgaacag	cctaaccatg	11880
tttagggaac	actctgtgtt	atgcgttcca	ccacacaatt	catgtcacag	ctgaatgtcc	11940
cagtttccct	tcagctagg	acacaacggg	agacccaagc	tttccaaaca	tatgccccag	12000
gcaacacttt	gatacacaag	gaataaact	ggatgctggg	tgtgcacaga	attcaaattc	12060
aggaatggag	ggaagtactg	ccctttgacc	catacaatag	gggatccct	tccttgagcc	12120
ccatggctta	gagggcccca	ctctgtccct	ctgcagactg	tgagctgccc	tatggaatga	12180
gggtgtctg	aggtgttatt	aatttctatt	tgttacacag	cagtggaaac	taatacagtc	12240
ccacaggcct	aatcaaggc	gtcagcaggc	tgagttcctg	cctggagact	ctggggaaaa	12300
ttcacttcca	aactcatttt	tggtgttgcc	aaaattcaat	cccttgaggc	tcagggactg	12360
ggctctgctc	ctcactggct	gtcagcctgg	ggccactctc	agattctgaa	gtcctccccc	12420
attccctgcc	acatggctct	ctccatctcc	aagcctgcta	tgacacaacg	cattgttctc	12480
atgcttcaaa	tctctgactt	ctctcccggg	accagaagaa	gaatactctc	ttcttttaag	12540
gggtcatgt	gattgggtca	ggtcaactct	atcatatcat	ggagtctcac	cgtcttgccc	12600
aggtgggtct	tgaacacctg	ggctcaagt	atcctcctgc	gttggcctcc	caaagtgtg	12660
ggattacagg	cgggaactgc	tgtgcccagc	caaggccagc	atcttcagac	ctctctgccc	12720
tgttttcaca	tggctctctc	ctctttgtgt	gtgtgtcaaa	tctcctctctg	cctctcttat	12780
aaggacgtgt	gtaatagcac	tcagggccca	cctggatgac	acagggtcat	ctcgccatct	12840
caaaatcggt	aactttggcc	aggtgcagtg	gctcatgcct	gtaatctcag	cactttggga	12900
ggctgagaca	ggtggatcac	ttgaggtcag	gagttcaaga	ccagcctggc	cagcatgggtg	12960
aaacctgtgc	tctactaaaa	atacaaaagt	tagctgggca	cgcacctgta	gtcccagcta	13020
ctcaggagac	tgaggcaaaa	gaatctctag	aacctgggag	gcagagactg	cagtgaagcca	13080
agattatgcc	actgcacttc	cagcctgggg	gacagagtga	gacttcatct	caaaaaaaaa	13140
aaaaaaaaaga	tgactgggca	cgggtggctca	catctgtaat	cccagcattt	gggaggccca	13200
ggtgggcaga	tcacctgggt	tcaggagtgc	attcaagacc	agcctggcca	acatgggtgaa	13260
agtctctctc	tactaaaaaa	aaaaaaaaata	caaaaattac	ccagatgtgg	tggcaggtgc	13320
ctgtaatccc	agctacttgg	caggccgagg	caggagaatc	acttaaacct	gggaggcaga	13380



## -continued

gggtgcagtg agctgagatc actctgctgc actccagcct ggacgacaga gcgaaactcc	13440
atctcagaaa aacaaattaa ccaggcatgg tgctgcgtgc ctgtagtccc agctactggg	13500
aaggctgagg tgggaggatt gcttgatcct gggaggttga ggctgcagtg agctgtgttt	13560
gtgccactgc actccagcct gggcgacaga gtgagaccct atctgaaaaa aaaaaaatcg	13620
gccaggcgcg gtggctcacg tctgtaatcc cagaactctg ggagactgag gtgggtggat	13680
cacctgaggt caggagtctg agaccagcct gaccaacagg gtgaaacccc gtctctacta	13740
aaaatacaaa aattagctgg gcatgggtggc aggcacctat ataatcccag ctactcgga	13800
ggctgaggca ggagaatcgc ttgaactcag gaggcagagg ttgcagtgag ccgaggtcac	13860
ggcattgcac tccagcctgg tggcagagtg agactccgtc tcaaaaaaa gaaagtctct	13920
ttttacaact tttttgaggt ataacttaca tatcagaaaa tcaccggttt taaatataca	13980
tttcaacgac ttttagtaaa tttccaagt tgtgcaacca tcatcacaat ccagtttcag	14040
aacatttcat cagcccagta agagccctca caccatttaa cagtactccc cactccccc	14100
ttctctctgg tggttgtcga acccaggtaa ccatgaatct attctctgcc tctggagatt	14160
tgtcttttct ggacaatctt tcttcttaaa ctacgcagag tgagttctgt tgtttagaac	14220
tagaaacctt gcccaataga atactaatat cctctgtttt tgcaaggagg ttagagtctc	14280
gctctgtcgc ccaggctgga gtgcagtggg atgatctcgc ctacttcga cctctacctt	14340
ccggattcaa gcgattctcc tacctcagct tccctgagtag ctgggattac agacgtgcac	14400
caacacacct ggctaatttt tgtactttta gtagagatgg ggtctcacca tgttgccaa	14460
gctggtcttg aactcctgac ttcaaatgag ccccccgcct tggcctccca aagttctggg	14520
attacaggca tgagccactg ccccagccc taatatcctc ttattacaaa agaagaaaca	14580
atgactcaga aagattaagt cacttaccca aggtcatgta gtcagtaagc aatagagctg	14640
agactgaacc tactccaaa gtctatgctc tactaaagt attgcagtgt agagagtaga	14700
ggggacaggg tgagactgga ggaaggagg gcatttagga gactattggg atgtccagat	14760
ctgagagctg ttggcgacct ggactgggga ggtgatatag gacagaggaga gaagtagatg	14820
aatttgagca tattaggaag tgaacagta ggatttgga actgaacggg tgagtggaca	14880
agaaggagag gatgagtcaa agagtgcacc atgcttctgg cttgggcaac tgatgggtac	14940
tggataaact tgctgggata ggaaatgcag gagaacagc caatttgggg tggagatga	15000
ggtcaggact aaaggcattg agtttgaggt gtctatggaa tgtacaaatg gagatgttga	15060
atggatgggt ggatatagca atctgcagct gagatgtagg tttctatatc accagcatat	15120
agacggcaac cagaacctg gaagaagggt caaccgacca gcaagagaag atgtagatga	15180
gaaagcaaaa gggcctagaa cgcgcccga aggagtgcata acctttaagg cgtgggagcg	15240
cagaagtccc caaaagattc tgaggagtag ccacggcgat ccagtgaag tcaggcacia	15300
gtggactcct gcaagctaag ggaaggaggc gagggaagtg gggcatggc acagtacgtg	15360
tcaactgctg ctgagagttc cagggaagtc aggtccatgg tgtatctatt ggatttagtg	15420
acaacgacat tgttgtttac ttggcaagag aggcctttaga gaagggatgg ggtgggagc	15480
cagactagag tgggtagagg agtgagtga tcaagatgcg gaagtggatg tgatgactcg	15540
ccctttcttt cttctttctt tctttttttt ttttttgaga tgcagtcttc ctctgtcgcc	15600
caggctggag tgcgatgggt tgatctccac tcaccacaac ctctgcctcc tgggttcaag	15660

## -continued

tgattctcct	gcctcagcct	tccaagtagc	tgggattaca	ggcatgtgcc	accacacccg	15720
cctaattttt	tgtattttta	gtagagacgg	ggtttcacca	tgttgccag	gctggtctca	15780
aactcctgac	ctcttgatcc	acctgcctca	gcctcccaaa	gtgctgggat	tacaggcgtg	15840
agccaccgta	cccggcctgg	tgactcactc	tttcaagggg	ctcagctata	aagaagagag	15900
gcatggagtg	gtggtaggat	ggattaagct	ggggagggga	ggcattttta	agatgagaaa	15960
tgatgtttta	atgcagatgg	aaatgggaca	ggagagaaag	ggaagtgtgat	gggataaatg	16020
aaaattatca	tcatcatctt	cctcaccatc	accagcagca	accacaagga	aagctcgttc	16080
tgctctggga	ctgccctgtt	agacatttct	tccttatact	gagccacaat	ctatctccct	16140
gtaactttcg	cctttgggta	gagttctgcc	ttgtgtccct	cttctactct	tcaaattggct	16200
gcacgctgct	actatgtctg	tcctcttctt	caaactcaca	gcccccatcc	tacggggcct	16260
gtgtcccag	cctggtgacc	cgaccctcta	ctgtcccagc	caagttcaca	tgaccatcct	16320
cttatttgtc	aagcctctct	tttttttttt	tttttttttt	tttttgggag	gagtcttact	16380
ctgtcccagg	ctggagggca	tggcacaatc	tcagctcact	gcaacctctg	cctcctgggt	16440
tcaagtgatt	ctcctgcctc	agcctcctga	atgctgggat	tacaggcata	catcgccatg	16500
cctggttaat	ttttgtatct	ttagtagaga	cgggggttca	ccatgttggc	caggctggtc	16560
ttgaactcct	gacctccagt	gatccccctg	cctcagcttc	ccaaagtgt	gggattacag	16620
cgtgagccac	tgagccgggc	caagcccctc	tgaagaggca	gtgtgcacac	tcttgctttg	16680
ctgatagaag	catgaaatgg	ttcaagtttt	tgtgatttgg	gaaaatgaac	tcagctataa	16740
aatgtgcaca	cccagaagtt	gtcctatgga	cttaatgtct	taggggtgag	aagaagcttg	16800
tgcagggatt	cctggcaaca	tgaatgcggg	atggacagac	agtgagcaca	gtgttgagga	16860
gcacagcctg	agtgcactc	cctgagatca	ggaacaggac	aaggatgccc	gctttcaccg	16920
ctgctgttcg	acattgtact	ggaagttcta	gccagagcaa	ttaggcaaga	aaaagcaata	16980
aaaggcatcc	agattgtaaa	ggatgaagta	aaactatctt	tagtcacagg	tggcacaatc	17040
ccatataggg	agaatcccaa	agaattcaca	agaaaactgt	tagagcaaata	aaactaattc	17100
agtcaagttg	caaggaacaa	gatcaaccac	aaaaattaat	tgtgtggggt	ttttttacac	17160
cagcaatgaa	catttgaaaa	ggaaattcga	aaaagtattc	cattcactgt	agcatttaag	17220
aaagctgagg	ctgtgaagct	acatggctgt	gtttggagct	ggtgctgcta	tttaccatgt	17280
gacctgggac	aagcttctca	cttcattcat	gacaccgttt	attcagctat	taagtggggg	17340
taataatagt	accttcctta	tagagttggg	tttttgtttt	tgtttttggt	tttgggtggg	17400
gagagtaagg	attgcaagag	ttaatatatg	ttagcaaaat	ggacttagaa	gagtgggttg	17460
cacatggtaa	ggactatatt	agggttacca	aaaaatgttt	ataattggga	aaattgaaca	17520
aaggttccag	taaatatcca	tcccagaggc	tgggtgacaa	attatcaca	tgtacctttt	17580
agcttgaacc	atatggaact	gcctatatcc	tatagctatg	gcctacaaaa	tgcaaaatgg	17640
gcagtttttt	ttgttggtgt	tggttggttg	ttgttggtgt	tgtcttgaga	cagtctcagt	17700
ctgtcaccag	gctgaagtgc	agtggcacga	tcttggtcca	ctgcaacctc	tgccctccag	17760
gttcaagcga	ttctcctgcc	tcagcctccc	aagtagctgg	gattacaggt	gtgcaccacc	17820
acgccagct	aatttttgta	tttttagtag	agatgggggt	tccaccatgt	tggccaggcg	17880
ggtcttgaac	tcctggcctc	aattaatctg	cctgcctcgg	cctcccagag	ttctgggatt	17940

## -continued

---

acatgagtga gccaccgagc ccagccaaaa tgggccgttt catatgttca atttaaacac	18000
taagatactc tttagccact gaaaatgact gtgtatcata gatctatctt tatcacagat	18060
ctatctttac tgtcatagaa agaactaata acatattatt gaacaaaaat aacaagttac	18120
aaaaacagctt gcagccacgt ttgtaacatc agtatatcta tttgtgtgtc taggtgtgga	18180
cataaagaga tatctagaag acactcccat tgcctatggg aatgacttgg gctgggatct	18240
tttttattta tttatttatt tatttattta tttatttatt tattttgaga cagagtctca	18300
ctctgtcacc caggctggag tgcaatggca tgatcttggc tcaactgcaac ctccgcctcc	18360
caggttcaaa cgattctccc acctcagcct cccaagtagc tggaattata ggcacccatc	18420
accgcacctc gctaattttt gccttttttag tagaaatggg gtttcgccat gttgcccagg	18480
ctggtctcaa actcttgatc tcaaattatc cacctgcctc agcctcccaa agtgctggaa	18540
ttacaggcgt gagtccaccg acccagctgt gatcttaatt tataattttt ctttttaacc	18600
cttgattttt accctgtttg actgacttta ggagtctgac ccccaaaact gcaaaagaat	18660
aaattagtgt tgttgaaagc cactaagttt gtggttaattt gttagagcaa caaccagaaa	18720
ctaatacaat cgggattaaa cttgccttgg ggggtgctggc tctccccac tttcttctca	18780
gagcatctct tgggtcctcg ccccttgctg agccccacc atggccgggt ggaccagcga	18840
tggacctgat cggattctcc tcttgggatt ttaaataaga gatgcagagg ctgcacgcca	18900
agcttctctg tgccgactg caggcacagt cctggcgcta ctgttcttta ggggtccatg	18960
aagatgtttt ttggcctggc ttagggtgta gctaagagat acaagggata cactttggag	19020
gctcagccct ttgatgttac ctttgcgctg gaagcccagg ggctgtaaga cggagaggaa	19080
ctttgctgaa agggagggtcc tccctgcaga gcttgaagag caatctaacc atgctggcat	19140
ggggcccatc ccacctgtga tgccagtgtg agtctgtgac agtcactcat attcagagca	19200
tgtggattga gactctgaag aacgacttta tgggtggcac cttcatctcc atctggtggg	19260
tccagagcaa gcagcctcag ggtcccatgg gcccaacctc accctccatc cttatttccc	19320
tgccactcca gggctcctcc tcagccagac ttgcctgtga gtgccccttg atggctgtat	19380
tagtctgttc tcacgtctgt aataaagaca tacccaagac caggttaattt ataaaggaaa	19440
gagggttaat gggcccacag ttccacatgg ctgggaaggc ctcacaatca tggcagaggg	19500
cgaaggcat gtcttacatg gtggcaggca aaaagtgaat gacaaccaag caaaagaaga	19560
aacctcttat aaaacctca gatcccatga gacttactca ctaccacgag aacagtatgg	19620
gggaaactgc ccccatgatt caattatctc ccaccaggtc cctcccacaa cacatggaaa	19680
ttatgggagc tataattcaa gatgaaattt ggatggggac acagccaaac catatcagtg	19740
gccctggctt ctctcccat agctctgttc tacttcccaa aatctttcca gctttgttct	19800
tttttagaga ctggagtctc actgtgttgc ccaggcttgt cttgaactcc tgggctcaag	19860
ggatcctcct acctcggtct aacaagtagc tgccaatata ggtgcacacc accacacccc	19920
cacaacttac ccctcctccc cggtcagtg aaatccctag gtccttcaca ctcccgtag	19980
ctccaggtac cagcggcttt gcctcttaca gaatgcataa cattatagcc tcttagagca	20040
aaaaggtcat ggggatggg aaggctggga gttctctctt cattcaggga ttaaacctac	20100
cacctggagg ggtgacatgg ctcaccttag tcaccaagca aggtcgggag agtatttctg	20160
ggatggtcat tagcactttc tttccaaagg gaggatacta caaactcact ttcagatca	20220

## -continued

---

cagtatcccc	cccgaacaggt	aaagtgtgca	aactcgcttt	cattcccttt	ttgtctttga	20280
cgcagctctg	aagcatttta	attcttaaaa	tttacctgtt	catgatgaca	cagagacacg	20340
ttttctgttt	aggagagggc	ctgattccat	taaggctgat	cgatcatatg	ttgcacattc	20400
tgggataaat	atgagagatc	agaattgagc	ttcccaaatg	gtgggtcaca	tgaccacgtg	20460
ttgttccagg	agaacgttgc	aacatccagc	atgagaaaagc	gctgcatgca	ccatgtatgt	20520
gttgggatgc	tgggtgaaa	tgtattaatg	agcaagtgtt	atcatatttg	tctatggaat	20580
tctatttaat	agctatgtct	cgtatgtgct	gattgggtct	caaactctggg	tatcttacct	20640
caacatcttt	cttgaaaatc	cttcttttta	aataaatggt	ggccgtgtgc	actggcttat	20700
gcctgtaatc	ccaacacttt	gggaggccaa	ggcgggtgga	tcacctgagg	tcaggagttc	20760
aagaccagcc	tgaccaacat	ggagaaaccc	catctctact	aaaaatacaa	aaattagctg	20820
ggcacgatgg	tggatgcctg	taatcccagc	tacttgggag	gctgaggcag	gagaatcact	20880
tgaacctggg	aggcggagggt	tgacagtaagc	tgagatcgcg	ccattgcact	ccagccttgg	20940
caacagagtg	agactccgct	tcaaataagt	aataaataaa	tgttatagaa	atactaaagt	21000
gtattgtgca	taataataca	cccacgtacc	accatccaac	ttaagaaata	aagcattact	21060
aatccagttt	tgctccctgc	aaactcatcc	ttcatcccat	tctctttcct	tttgctttat	21120
aaacatcctg	cagccaggca	cagtggctgc	acctgcagcc	tgagctactt	gggaggctga	21180
gatgggagga	tcacttgagc	ccaggagttt	gtggctgtcg	tgtgctgatt	gcacctgtga	21240
aatagccact	gtgttccagc	atgggcaaca	tagcaagact	ccatctctaa	atgaaatcct	21300
tttttttttt	tttctttttg	agacagagtc	tcactctatc	accaggetg	gagtgcagtg	21360
gcgccatctc	ggctcactgc	aacctccgct	tccagggttc	ctgcctcagc	ctccggagta	21420
gctgggacta	caggcacatg	ccaccacaac	tggctaattt	ttgtattttt	agtagagacg	21480
gggtttggcc	aggttggttt	cgaacccagg	acctcagatg	atccacctgc	ctcggcctcc	21540
caaagattac	aggcatgagc	caccatgccc	agcctaaaag	aaatcctgaa	tttgacattt	21600
ttaattccct	tgtttgtttt	aatatttcta	cagcacatat	atgtctacat	atataaaaaa	21660
tataattttt	tattggggtg	aaattcacat	aacgtaaaat	taaccactgt	aaacatgaac	21720
aattcagtgg	catttaggac	attcatagta	ctgtgcaacc	accacctcta	tctagctcca	21780
gaacattttc	atcaccccaa	aaggacactc	catgcccccc	aggagtcact	ccccatttct	21840
tcccactctg	tcccctgcaa	cgaccaatct	actctctgtc	tctgtggatt	cacctattct	21900
ggataattca	tataaatgga	atcatacaat	atgtggcctt	ttgtgactgg	tctctttgac	21960
ttagctgggt	gtggtggcac	gagctaccca	cctacttggg	aggctgaggc	aggagaaatc	22020
cttgaactcc	agccataaat	tatatgtttt	tcaaggttca	tcaatattac	tcgccagggc	22080
tggagtgcag	taatgtgacc	ttggccccct	gcaacctcca	cctccccggt	tcaaaggatt	22140
ctcctgcctc	accctcccga	gtagttggga	ttacaggcgc	cccccccccc	caccgggcta	22200
attttttgta	tttttagtag	agagagggtt	tcaccatggt	ggatcatgctg	gtcctgaact	22260
cctaattctca	agtatccac	ccaccttggc	ctcccaaaat	gctgagatta	caggtgtgag	22320
acactatgtc	aggcctagtc	aattcatttt	actgctgca	aagcatctca	ttttaaaaaa	22380
ttattttatt	attatttttt	tgagacaggg	tctcactctg	ttgccagggc	tggagtacag	22440
gggtgcaatc	acagcttact	atagccttga	cctcccgggc	ccaagcaatc	ctcccactgt	22500

## -continued

---

ggcctccaga gtagctggga ctacaggcat gtgccaccac acacagctaa tttttgttct	22560
ttgttttttg ttttgttttg ttttgttttt tgagatggag ctttgctctt gttgcccagg	22620
ctggagtgcg gtggcgcgat ctacagctcac cacaacctcc atctcccagg ttcaagcgat	22680
tctcctgcct cagcctcccg aataggtggg attacaggct cagccaccg taccagttta	22740
atttttatth ttagtagaga tgaggtttca ccatattggc caggatggcc tcaaactcct	22800
gacctcaggt gatccatccg ccttggcatc ccaaagttct gggattacag gcgtgagcca	22860
ctgtgcctgg ctcaaagcat ctcatgtgat gaatccaaca atttatttat ccattcgttg	22920
tttgagtaca ttaggtttat ttcctgcttt ccctcttaca agcaatgctg cagaaacatt	22980
cttgccata gtcctcccg cacatgtaca agagcttctc tagggcaca atgtaagagt	23040
tttattgctg ggccatggg aatgctcttc ttcaaagta cttgctattg tcaaactctga	23100
gctatctttt caggagccta ctactgggtc tcagccctgg ctatgcatta gaatcacttg	23160
ggaactttaa aaaaatatgt gggtatcttg ataactccc cgaccctgg aatcctagtc	23220
tccagggatg ggactgggga cttgaaagct ccctgggtga ttacacagcc aaggtggaga	23280
acagctggga ctgagctctg gccactctg tctcagcttc ccaaggtcag ctcatagagg	23340
gcagccatca tcttagactt gctaatatca ctacagcagct gagcagatag aagggactaa	23400
ataaataaag tctgaatgag taaataaagc atttaaaaca aaccttttag ggctgggtgt	23460
ggtggtcac acctgtaatc ccagcacttt gggaggccaa ggtgggtgga tcccttgagg	23520
ttgggcattc aagaccagcc tgaccaacat ggcgaacct tgtctctact aaaaatacaa	23580
aaattagcaa agtggtgtg tacgcgctg taatcccagc tactcggaag gctgaggcag	23640
gagaatcact tgaaccgcgc aggcagaagt tgcagtgagc tgagatggag tcaactgact	23700
ccagcccggt ctacagagca agactccatc tcaaaaaaa cataataaaa caaacctttt	23760
aggctgggtg cgggtggcga tgccggtaat cctggtaact atctcctact gttgtgacga	23820
ttaaaggaga tgggtgatgc agagtacttg gcacatatga agtgctcaat aaagagtaac	23880
tctgcagcc tgggtgtgtg gccacacct gtaaccacaa catgttgga ggtcaacat	23940
gttggcagga ggattgcttg aggccaaagg tttgagacca gcctgggcaa cataaagaga	24000
ccccctctc tataaaaaat aataaaaact agctgagcat ggtgatacat gcctgtagtt	24060
ccagctactc aggaggtgta ggtgggagga tcaacttgag ctcgggagtt ggaggctgca	24120
gtgagctacg atcacgccac tgccctccag cctgggcaac agagcaaggc cctttctctg	24180
gaataaaaaa aaaagagagg ctgggcgcag tggctcacac ctgtaatccc agctacttg	24240
gaggctgagg caggagaatc acttgaacct gggaggtgga agttgcagtg agccaagatc	24300
gcaccattgc actccagtct gggcaacaag agcaaaactc tgtttcaaaa aaaaaaaag	24360
agagagagag agagtactg ccattttcaa tgttattgat aaagcaggtg ggtgtttctg	24420
aaaatgcacc agatgcagcc ttagactgga aggtgctgat gtgttactga gcctccagac	24480
aaagctggcg aaccaccggt ggcgagactt tatgacttga gaagctgtcc tatctatgag	24540
ggtccatggt agggaatatt aaagaatagg attgggggca gtagctttcc ttagcaagt	24600
gctagctatg tgccagatag tgggcaagg agtctgcaga cgtctctcat ttcacctca	24660
caatccatga ggcaggtatg actattcttc ccattatcca gagaggga acaagtcaca	24720
cgaagcaag tcacgcagca gagctaggat tggaacagaa gtgcctgcat cctccaacgc	24780

## -continued

ctgcatccca	acccgctgtg	ctatgctccc	cggtgatgct	ttcccattgc	tgccatttag	24840
ctttgctctg	gtgagtgaca	agctgtacca	gcggaagag	cctgtcatca	gttctgtgca	24900
caccaagggtg	aaggggatag	cagaggtgaa	agaggagatc	gtggagaatg	gagtgaagaa	24960
gttgggtgcac	agtgctcttg	acaccgcaga	ctacaccttc	cctttgcagg	tgagcacctc	25020
gtagcattct	cccaggctcg	tcgctgggta	ccgtcgccag	ggcctagctc	ccttccccta	25080
ggatctacag	cctcactcca	gaaaaaacgc	tggtcctatt	ttaaaagctc	tgtgaacatc	25140
ccaactgaga	aaacctaaaa	attgcaaaac	tgggtgagat	ccaggagatg	attccttgct	25200
ttatcaagtc	ctacagggtt	tctcaaaata	gcctcatgtc	ctggtgcatt	gaaaacagga	25260
tttgattcac	tcgcgcttga	tttaccgcga	gctttttaga	cgcacagcca	ccaagccagc	25320
aaggctctgcc	gagattcaga	aggacagaga	actttctacc	aaagagcacc	agcccttcta	25380
gattgtgtag	catttcatca	acaaaaatga	ctccaggaga	ccattcttgg	gtacattttt	25440
ccacaaagaa	aggatgtaaa	tgacaaaaag	aagaaaggca	tcggtcactg	agagaaggcg	25500
tgtgacttct	agatttgcaa	gcagggaaaa	tgaagcaga	attgccaaaa	aagagaaatt	25560
aaagtagggt	tgaggagtcc	aaaaaacaga	taagggggcg	agagggacaa	aggggcacgt	25620
ttgcggcttg	gggggctaaa	atgctttgca	aagacctatg	aagcccaagc	tgctgtttgt	25680
ttctaacagt	gggtcattaa	aatcctgtga	ttgctcaggt	gggtggatgc	cctagtgcct	25740
tttcattaaa	gtctgggaaa	atctgaacag	tgttgtgatg	aaggctgctc	ccctaccctc	25800
gcctccccag	gtctcctgag	tttcaattta	atgagatttt	tactgcgtaa	aaaaaaaaaa	25860
aaacaaaaac	aaaaacatga	aaaggctgct	tttgagactg	catttggtaaa	tgactcttca	25920
acccattcaa	acgctccttc	actctccctc	agctcagcag	ggctgctcgt	ccagctttga	25980
tattaagccc	ttggcatatt	ccaagttgcc	cacagatcct	gatttctaga	agcttagaaa	26040
agtgagagag	ttcgcccagc	aagctggatt	attataatta	agtagttctc	ttttcaaagg	26100
ccttgcatth	tcttagcctc	tccttctcca	cagggggaact	ccttcttcgt	gatgacaaac	26160
tttctcaaaa	cagaaggcca	agagcagcgg	ttgtgtcccg	aggtaaggag	gggacctgga	26220
gtggtgggtc	aggtcttaag	agttcctggg	ggagggtgca	gtcggaaaga	gcagaaatgc	26280
ggacctgggg	gtgtatttga	gccacaaaat	atctactgag	cacctgcctc	ttttgtgggg	26340
tagggctggg	ctgagtggag	gaaggggagag	aacacatggc	agtgctctct	catggccacc	26400
agatctttgt	ctgcctgcct	cggtctctgg	ttctagcact	agggaccctg	gcagacggca	26460
acctgtctct	ttctatctct	ggctctcttc	gcagcttatg	acaatgggtg	ttcttgccat	26520
ttgcacaaca	tttcacattt	tacttgaggc	tcacaatgac	tttaggagaa	aaatggcatt	26580
ttacaaaacca	gaacacagat	tcagagatgt	tgtgacttcc	cccagggtca	cacagcttgc	26640
tgacggcaga	gccagggtta	ggacctgagt	gtcctcactc	acagcttctc	ctaccatggt	26700
tcacagtga	ctgtctgcag	ggctggctct	ggaaaaccct	ggcacacagc	tctcttgctc	26760
attcatccac	ttcttagagt	ctcagcctct	ccatcacgtg	catgtgcccc	catgactccc	26820
tgccaccacc	ccagacaatc	tttcccaact	tggctttgct	ctctccaaag	agccgcctct	26880
ttctggaaga	gcttcttccc	tatcaaatca	tatttggttt	attctgggct	aggcacgggt	26940
gctcacacct	ataatctcag	cactttggga	ggccaaggca	ggaggatcac	tggagcccag	27000
gagttcaaga	ccagcctggg	caacatagtg	agaccttgct	tctaaaaaca	aaacaaaaca	27060

## -continued

aaaaactgtat ttgtcctatt ctttaacctca ccatacctgct ctctcctctt aaaccagctt	27120
ttcttctaag ttatctata ccagcgtgtt actgctaggc attacctgta actcccaacc	27180
tcccagcctc ccctctcagg taacgtaact gtgctgtacc agggaccatt tccaacactg	27240
accgttgtag cagcgtgtgt gatacccagc cagatcctcc actgatgagc aggaccagcc	27300
ctgccggtgg agttttgctc acgggcccac tctggcaaca aggtctagct tcctgagtgt	27360
ccaggacctc tgggcaaaag tgctgtcttt gactacagtc tttgccatca tttgtacca	27420
aattcagaac caagataaca ttcccttggt gtctatcttg ataactctcc gtctaaaagg	27480
agagctttct acatttgttt tgcctcatc accccttaac tgcctctgct gctcagatga	27540
aatctctgac agccaggcca gccaaagcct cccttctgct tcccttccca gcagtgttcc	27600
gtctccaatc cagtttgggg gcctgtttcc cttttgtccc atcctttggt catctctaca	27660
ccacactcta cctccatgta cacataaaga cacttttggg cttattgaga cttggaagaa	27720
aaagagttag gatagcccag agattaagag cctgggctca gaagctagac cgctaaaggt	27780
tcaaaactta aacttcttaa ctctgtgacc tcaggcaggt gactcttcac ctctctgtgc	27840
caagttgtgt gcatctgtaa aatggggctg agaacaaaac ctgctgttc ggattattga	27900
aacaaatgag tgaatacctg tgacacgctt aaaacaaggc atggtacatt ataagcactc	27960
aatgaatgtc agttgttatg attcaaatga catgttactc ctcagatagg atgtcactca	28020
gagtgactgg aaccacaacag gttccatcga agcacttcca gccagagag cacatgtgac	28080
tctttgggat ttaacttat gctccaggct gggttgggct ggtctaattg ggcatttgcc	28140
tcctctctgc ccatcaacac ccctctatct cgggttgtct cggtaaatat gaccaagttc	28200
cagttacaca ccaacatgaa tagatatttg ctccatttat ctgtggctgc aaaacaaact	28260
accctgatca tgggtgggctt aacacaagca ttttattaac tctcacagtt cagtggttga	28320
ctgagttcag ctggttgggt cttctgctgg tctcactagg gatctctcaa ggagctgcag	28380
tcagatggca gctggggttg gcatcccttg gagcctcaat tggcacccag ggatagctag	28440
acatctcttc ctctctggga acccctaaga gttaccctct cttcaaggtc tgtccgcatt	28500
gtctctccag cagggtagtc aaactttcta atagtggctc aggttcctca aaacaaaaca	28560
aaaaaataga agcttccagg cttctttaag gcttaggcct ggaactggca acagcattcc	28620
ttctgctgct tctattggct ttgcagtcac agcacaggct cagcttcaga aagtagggga	28680
aacatccac ctctcaatgg agaaagtgac caagaatttg tggccatctt taatccatgc	28740
aacaatcaca aaagagaaga ttaaaacaaa actaatgtgc tgttttttaa aaaaaaatt	28800
agaattcctt ttggagagca aaataaaact gatttctctt ctttctgag attcatctcc	28860
agttgataat gaaaatgtaa ctttccatga gccacaatga atgatcacac attcattcct	28920
tccatgacta tcaagacctc ctatttgcca ggcacttggg ctgcaaagat atgaccttg	28980
ctctgtattc aaagggttca taatctaac tggggagtca caaggggagg agaccaacgg	29040
catgccatgt gacaaggcag cctgggaaat aaagagatca tgagagagtt tgaaaactac	29100
aagaaacatt cattgagcat gtactatgtg ctaaaccctt tcttttcata actcacttaa	29160
tcttcacaac aaccaggcaa gacaatttgt tggctgggca tgatggctca cgcctataat	29220
cccagcatt tgggaggtcg aggcaggcag atcatgaggt caggagtca agaccagcct	29280
ggccaagatg gggaacacc gtctctacta aaaatacaaa aattagccag gcatggtggt	29340

## -continued

---

gcgtgcctgt aatcccagct actcaggagg ctgaggcagg agaattgcct gaactcagga	29400
ggtggagggt acagtgagcc aagatcacac cattgcactc tagcctggtc gacagagcaa	29460
gactctgtct ccaaaaaaaaa aaaaaaagaca aatctgttca cctgtctttt	29520
acagatgaac agtgagctgg ccaaggctctc tcagctagaa aatgggtgcag ctactgctga	29580
accagatata cctgggtcca agacctaggt ccttgacctg aatcactgta gaagccacaa	29640
attgcacaga tgacaagag cagaggccat tttcacacct atgtcatgtt ctggcttccc	29700
caagactggc catgaagcct gctccctaga aaacaggaag agaaatatgc ataaaccagg	29760
gcagctcatt tctctcttcc gtgttctgtc cttcagccct gagaccagc tggctctcatt	29820
ttcttgggcc aggcgagtgt cctggccaat gtctctcagt tccatgtgtc ctgctgagtc	29880
tccgcacctg ttgtccctgc tcagtccttt cagccacagg gccacttga ccaagccacc	29940
tgcccccat cccagccagc atccctgaaa gtgatcccta gcttcccaga cctctgcttt	30000
tcccattttt atttatttgt attgggatat aattcacata ccataaacc caccatttta	30060
aagtgtacat acggttcagt ggtttttagt ataatcacag agttgtgcaa ccatcaccac	30120
cgtctaattc cagaatattt tcttccttct ttctttcttt ttttttctt tctttctttc	30180
tttttctttc ttttctctt ccttcctttc tttcttttt taggtggagt ttgctcttg	30240
tcaccaggc tggagtgcag tgacgcaatc tcagctcact gcaacctcca cctcccagg	30300
tcaagtgatt ctctgcctc agcccccta gtagctggga ttacaggcgc acaccaccac	30360
atctgataat ttttgtattt ttagtagaga cggggtttca ccatgttgag caggctggtc	30420
tcagactcct gacctcaggt gatctgccg ccttggcctc ccaaagtgtc gggattacag	30480
gcgtgataag ccacggcgcc cagccccag aacattttca tcacctaca aggaaacccc	30540
aaatccagta gcagtcactc cccattctcc ccttcccctg tccctggcca cagtctactt	30600
tctgtctcta tagatgccta ttctggacat ttcctataaa tagaattgta tatggtgtgg	30660
ccttttgtgt ctgtcttctt tcaactcagca tcatgttctc caggtccatc catgttgtag	30720
cctgtgtcat tgcttcatcc ttcttatggc taaataagat tctgtgtatg aatgtaccac	30780
attttatttg tccattcctc cgtcagtgcc cacttgcag gtttccactt ttttggcgat	30840
tctgagtagt gctgctataa gcattcgtgt gcacattctg gtggatatcg aatcacttct	30900
ccacatctta gtaacacacg tcaacttactc cccactctgt catccttcta tctgcagtat	30960
cccaccgcga ggacgtctgt ttcctctgac cgagggtgta aaaagggatg gatggaccgc	31020
cagagcaaaag gtaccttctg tttcttttcc cgagacccta ggggtggatg gtctggcatc	31080
ttggtgacat ttgtgatgcc caggtcaggt cttcagcctc tgctctcagc tgcctcttct	31140
caccatcacc aagccatagg cgagtctgcc catgcttcgg ctctgtcccc agcagaccag	31200
ctgctgactg taaacatgac tccagttttc cagtgcagga agaagctcct aaaaacctag	31260
caggttcagg attctaactg gtagaaaatt cacatggcct atagcatcat ctgagtattc	31320
taaaactttc ccctgaattt cctcaaaggt tgaggacat gaacttttac cccaggga	31380
cctggcagca ataccatata taacctgcag aatttttttt gttttttatt ttattttatt	31440
ttttaaacat tttttgact gttttatttt gattttgatt ttgattttat ttatatctaa	31500
gtgcagtgtc attgcgatac ctgcagaatt tctttatctc acattttaac ttaaaaaggc	31560
acagggcagc gagcgagag gctgggtgcct gtaatcccag cactttggga gggtgaggca	31620



## -continued

gatggatgct	tgaggtcagg	ggttcgagaa	cagcctggaa	aacatggtga	aaccccgctct	31680
ctactaaaa	tacaaaaatc	agccagacat	ggtggcacac	gcttataatc	ccagctactt	31740
gggaggctga	gacgtgagaa	tcacttgaa	ctggaaggca	gaggttgag	tgagccaaga	31800
tcattgccact	gcactccagc	atgggtgaca	gagcgagacc	cctttaaaaa	aaaaaaaaaa	31860
aggcacagg	caatttttaa	aatactgcaa	atagtaaaa	aaaaaaaaatc	agtgggtata	31920
atgcaaacac	acacaaaaag	gcataatgcc	attactgcat	tctactccat	actgtatgtg	31980
tatttgagtt	agtataaaag	ttattttaac	attgctcact	atttaattaa	ttctcccttg	32040
gaaactgatt	aatcatcctg	gcactccagg	aagatgtgcc	atgctgattt	catggctttg	32100
cacatcctgg	gcaggctgtg	tacccttgga	gggacttggtg	ccccttgag	aggccatgtt	32160
ctagtccatt	tatactaagt	gagagcatac	acctgttccg	ctccctcat	gggcaccttt	32220
tcttataaag	aaacaaaaga	gccagcagaa	tccacagtct	ttctgtgttc	tctctgatct	32280
ttattatgtt	ttgcttgttt	gccttgccct	gtgttcgttg	tggttaggat	gggcttgatg	32340
gaagctgaag	ctgcgtgggt	tggaagcct	ggtcaaagcc	tagtctctcg	ccggggtga	32400
gttaatgatg	tccctcctgg	agaacgtcct	ctctgcagtt	ctttcacatc	tgtggttcta	32460
cgatgctttg	acccctatag	gaattcagac	cgggaagggtg	gtagtgcag	aagggaacca	32520
gaagacctgt	gaagtctctg	cctgtgtccc	catcgaggca	gtggaagagg	ccccccggtg	32580
agtcgcatgg	ggagacagac	acagtgggcc	tcagcggcga	ccagatgagg	ccttgccgag	32640
gtgcttggtg	ccttcccctc	tcagcacagc	cctgcaaagt	cctgggtcct	accggcttgg	32700
ggacccctgc	gctctggatg	cactgcttgg	cacaaactag	tatctctggg	agggccatgg	32760
tggttggttaa	actgttgtaa	caatcctgta	ccaactggta	aatagctact	accctgagca	32820
tccttggtgtg	tccctggccc	cttccttccc	ccagatcttc	cagggtaccc	ccagaccccc	32880
tcctgtagtgt	ccacagcagg	atcccttctg	acttgtcagt	gtccatactg	agtgatcaag	32940
gatagaaagg	aaggaggagg	atggaaggga	aggacgaagc	gaggaaagag	aagggaagg	33000
ggaggaaaaa	gcaaaagggg	tgagggtaaa	agggggggga	aaggaaagttt	tctcagatta	33060
aatgcttaca	atgacataca	gatttggtgg	tcccttgat	tgatgcttcg	cttcaataca	33120
caaagtcaca	atgttaaatc	tcagaagcca	caagggtgta	tgtatttcag	cagagaatag	33180
ttagaaagac	ctggattcaa	ttcctaactc	taacaccatt	ttgctgtgtg	tccttgggaa	33240
aatggcttaa	cctctctgag	tttcagtgtc	ctcacctgta	aaagcagaat	aataatttca	33300
ccaacttcat	agggctgttg	taaggattaa	atgagatgat	actgttacag	ttattgtaag	33360
gtaagcccca	tgcattgctg	gcttacacac	acacacacac	acacacacac	acacgcacac	33420
acgcacacac	acacacacac	acacaatcta	cccctagaag	tgtggtggtt	ctagaccagc	33480
actgtccaat	tgaacttgat	gcagtgatgg	aaatttctgt	atctgtgctg	tccaataggg	33540
cagctactag	gtacatgtgg	ctattgagta	catgaaatgc	gactactgaa	tttttgaaag	33600
agatgataga	tgatagatag	aaagatagat	agatagataa	atagataata	gatagataga	33660
caggtagata	gatagataga	tagatagata	gatagataga	tagatagata	gatagagttt	33720
tgctatgttg	cccaggctgg	ttttgaaact	ctgggctcaa	gcgatcctcc	tgcttggcc	33780
tcccaaagt	ctggggttac	aggtttgagc	cattgctccc	agcctgaatt	tttaattaaa	33840
tttaaattta	aatagccaca	catgtctagt	ggctaccata	ttggacagcg	cagttctaga	33900

## -continued

ccgatgtgat	tcaggatcat	tccctcagca	tcgtggggca	aagagaaaac	tgccccaaagc	33960
tggcctgtag	aaggctcagg	cgaaggttct	ccaatgccgg	gatgggggggt	gcgctcagca	34020
gcataccccc	ttatgattct	caatcgctaa	tagctccact	caggttcatt	tctcggtcag	34080
gggcattttct	ttgggaatca	cccagctctg	ggagatacag	cagcctccac	tcaggtagtc	34140
cttgttcaag	acaagcggcc	cttgactgac	tgcaagttca	gttccagctc	tgctatcaac	34200
tcactcatta	aataaactgc	atctccagtg	tgctgcctc	tgggctggat	tttgacgtga	34260
cctgggcaag	caactccctg	aacttcagtt	tctcatatat	tatatgaatt	agctaagatg	34320
gttcgtttaa	tcattcattc	aacacatcca	tcaccacgta	gtagggtgta	gatattttatt	34380
tcatacgtaa	ctacgcataa	gagactttgc	taagtttttag	gtaaaataca	agtcccagat	34440
acggagcaag	tctcaaccac	tgtacatacc	tgaatgtgta	attacatcac	tgtaagaggt	34500
gccacagtaa	atgccactgg	gtcttgtgtt	agtccattct	cacacaaaga	actacctagc	34560
cagggtcggt	ggctcacgcc	tgtaatccca	acactttggc	aggctgaggc	aggcggatca	34620
cttgaggtea	ggagtttgat	accagcctgg	caaacatggt	gaaaccccat	ctctattaaa	34680
aaatgcataa	attagccagg	tgtggtggca	cacgcctgta	atcccggtca	ctcgggagggc	34740
tgaggcagga	gagtcgcttg	aacccgggag	gtggaggttg	cagtgaccgg	agatcgcgcc	34800
actgcactcc	agcctgggtg	acagagtggg	actccatctc	agaaaaaat	aaaaataaaa	34860
ataaagaact	acctgagacc	aaatacttta	cgaaaaaaa	gaggtttaat	tgactcacag	34920
ctccacaggc	ttaacaggaa	gcctcaggag	acttacaatc	atggcagaag	gcgaagggga	34980
agcaaacaca	tcttaccatg	atggagcagg	aggcgagttt	cgggggatgt	gccgcacact	35040
tttaaatgat	cagatctcgt	gagaactcac	tcactatcac	gagaacagca	aggaggaagt	35100
ccgcccccat	gattcagtca	cctcccacca	ggcccctcct	ctggcacatg	gggattacaa	35160
ttcaagatga	gatttggttg	gggacacaga	gccaaccat	atcagatctc	aagaagggag	35220
aaattcttct	tgaggagct	ggaggggctt	tgtggagagt	ttcagaatgc	tttgcccact	35280
aggtttgctg	tatccatttc	tcttcatgta	tcccaaagac	caagccaaga	aaccagaagc	35340
ctctggtccc	actggcccat	gggtccctc	ggtccccacc	gtcactaatg	gccattttgc	35400
atgtctctct	cccaggcctg	ctctcttgaa	cagtgcggaa	aacttcactg	tgctcatcaa	35460
gaacaatatc	gacttccccg	gccacaacta	caccacgtaa	gtgccaggc	tgcttggtg	35520
tcttagttat	ctactgctga	gtaataaatt	atcccaaacc	tcagaagcct	gaaacaacaa	35580
acgcctattg	tctcccacgg	tttctgtggg	tcaggaaatc	gggaatgact	ttgctgcgtg	35640
gttctggctc	aaggctctgc	aggttgtagc	caagctgtca	accagggtg	cagtcatttc	35700
taggcttgac	tggggttgga	gaacactttt	ccaagctctc	acacagttgc	tcgtggggaga	35760
gctcagttcc	tcaccacgtg	aacctcgccc	tagaccactt	gagtatcctt	ggtatatggt	35820
ggctggcttc	tcccagagca	agtgaaccac	gagagacaga	gcaagcaacc	aagagtataa	35880
ccaagatgga	agccacagtc	tttgggggga	gaccccaaca	cttctgcoat	atgccattgg	35940
tcacacagat	caacctgggt	ccagtgtgag	aggccactgc	ccagggttcc	caggaggcag	36000
tgatcatttg	gggctttcat	ggaacctctc	caccacactg	gctcactcct	gggaaagaga	36060
cagatctgtt	ttcaatcgag	atgtttgttt	gtttgctttt	aattatgcac	aggagaaaca	36120
tcctgccagg	tttaaacatc	acttgtaact	tcacaaagac	tcagaatcca	cagtggtcca	36180

## -continued

ttttccgact	aggagacatc	ttccgagaaa	caggcgataa	tttttcagat	gtggcaattc	36240
aggttggtgg	tgctttgtac	actgggatgt	ggggctgtgt	gtctagggat	ggaggatgtc	36300
aaacagccaa	gaggccgggc	cactgggtct	tcataatgtg	gctcacattt	actgagcatt	36360
tagtaaatcc	acccgctgcg	ctaagggtctt	tacctacgct	acctcgtaa	atcccaaac	36420
aatccttatg	agtgagagct	acttggtgta	ttcctttcct	gtggctgtg	tagcaagtta	36480
tcaaagctta	gtggcttcaa	acaacacata	tttgcttatg	ttgccagaga	tcagaagtgt	36540
gagatgattt	tccctgagcc	agggcggtgc	tccctccggg	actttaaggg	agaatccagt	36600
tcctcagctt	ttccaccttc	tggagctgca	ttccttgcat	ttcttcaaag	ccagcagcat	36660
aacatcttgc	ctcagtggcc	actttcactc	cctatcctgt	gtccaatctc	cctttgcctc	36720
tgtcttacaa	agagagagag	catttacaag	agggggcatt	taaggaccaa	ctggataatc	36780
caggataatc	tcccatctca	agatccttca	tttaggctgg	gcacgggtggc	tcatgcctgt	36840
aatcccagca	ctttgggagg	ctgaggtggg	tggatcacct	gaggtcagga	gttcaacacc	36900
agcctggcca	acatggtgaa	agcccatctt	tactaaaaat	acaaaaaaaa	aaaaaaaaatag	36960
ccgggcatga	ttgcaggctc	ctgtaatccc	agctactcgg	gaggctgaga	caggagaatc	37020
gcttgaacct	gggaggcaga	ggttgcatgt	agccgagatc	gcaccactgc	actccagcct	37080
aggtgacaag	agcgaactc	catctcaaaa	aaaaaaaaaa	aatccttcat	gtattcgcct	37140
ctgcaaagag	ctttccctag	gggagtacta	ggaggtaag	cagaaaagat	atttgataga	37200
gtgccctgaa	ttccagtcta	ataagtttgg	acttgatctt	taatgggggc	gtggggggca	37260
ttaaagggtg	ttgggtacag	gagtggtctg	ttgaaagttg	tatttttagga	caatgagttt	37320
aacagtgatg	tgtcccagac	gggggtaggg	agagtgagga	gatgcgattg	tggctgccac	37380
aataacactt	gtgcgagtta	ggtggggctg	tacatatggt	tcttcaatca	gcatttttcc	37440
tctaaaaacc	ttaagcaatc	ctgggtatgc	agggagatgt	ctggcggttg	cgtaactcac	37500
acccagcagc	catagagact	gtcccttggt	gatccttcag	ggcggataa	tgggcattga	37560
gatctactgg	gactgcaacc	tagaccgttg	gttccatcac	tgcctgcca	aatacagttt	37620
ccgtgcctt	gacgacaaga	ccaccaacgt	gtccttgtag	cctggctaca	acttcaggta	37680
actccaaggc	ccagggtcaa	ctcaccacgt	ggctgaatcg	cattcccagg	aactggtgag	37740
actaattttg	gtttccaagg	caacaagatg	aatgaaaaaa	gactttctct	aagaactagg	37800
tgataactga	attttttcca	taatttttta	aaattctcaa	aagagatgca	cactctttat	37860
ttttttactt	attttttttt	ttttgaaatg	gagtctcact	ctgtcaccga	ggctgaagtg	37920
cagtggcgcc	atctcagtca	ctgcaaacct	ccgcctccca	ggttcaagcg	actctcctgc	37980
ctcagcctcc	caagtagctg	agattacagg	cggatgcaca	ctgtttataa	aacaaaacta	38040
ttgggaaaca	gaaaagcata	gagggggatc	aaaatcacc	ataattcccc	taccctgaaa	38100
taatcaataa	caaccctcgg	gggaattttc	ctcatctgta	ccaattatct	catacagctc	38160
ctatgagata	atagcatata	tatatatatc	ttgtggtatt	ctgcagggtt	tttcatacca	38220
cagccactca	aaattctttg	taaccatcac	attaatgata	ataacattcc	attttgtagg	38280
tgaacaaata	acaactgcta	caattcagga	agtgttttct	tttcttttct	tttcttttct	38340
tttttttttt	tagatggagt	cacactctgc	ttgccaggc	tggagtgcag	tggcatgata	38400
tcagctcact	gcaacctctg	cctcctaggt	ccaagcgatc	ctcccacctc	ccaagtttct	38460

## -continued

---

gggaccacag	gcatgtgcca	ccacaccacag	ctaatttttg	tatattcagt	agagatgggg	38520
tttcaactgtg	ttggccagtc	tggctctgaa	ctcttgacct	caagtgatct	tcccaccttg	38580
gcttcccaaa	gtgctaggat	tacaggcatg	agccactgtg	cctggcccaa	ggagggtttt	38640
ccatatacca	agcactcccc	atcgccatcc	ctaaatctcc	caacaaccct	ggaaggaaga	38700
tattgtttct	ggaagatgat	ttgccaaga	cccacagctg	atagtacatg	ttgcataatt	38760
ctaaccacag	tcactctgac	cccacactca	cactccatcc	cttccttccc	atctcatgat	38820
tttctcacct	acgcctccat	gattgaatat	ttgagttgct	tccagttttt	ctattacaag	38880
taaccacagt	gtgcatcttt	gcacataaac	ttctctttga	attccagggt	acttccttag	38940
gataaaattc	tagacttatt	gaatcaaagg	ttgtgaacat	tttatcatat	gctttttatt	39000
tttaaaaaata	tctatagtta	taatgtttca	tttttttttc	tgagacagag	tctcaactctg	39060
tcacccaggt	tgagtggtgac	cgggtgcaat	tatagctcac	tgcaacctct	gcctcccagg	39120
cccaagtgat	cctcccacct	cagcctcctg	agtagctagg	actacagggtg	cagccacca	39180
tgcccagcta	attttttaaaa	tttttttgta	gagtcgggggt	ctcactatat	tgcccaggct	39240
ggtctcaaac	tcctggctca	agcaatccgc	ctgccttggc	ctcctaaagc	gttgggattg	39300
cagggtgaag	ccactgcacc	tggcctataa	ttttcatttc	taggattttt	atttggtgct	39360
ttttcaaagt	catctattat	tgttccagtg	agtccatttc	ttaccttaag	gacccacttc	39420
cttctgtcca	ttctactgta	tcattccttt	cataccgacc	tattatctga	agtaacttgg	39480
gtgggagttc	tcctcgtggg	ctttgaaata	ctgtcttcag	tagaaaagat	cttatgcaaa	39540
gttctgcatg	tgctgtgtgt	tatggagtat	tccttatata	ttactcaaag	cagctccata	39600
aactggcagg	cagccctcta	aatgtgttct	tgtttctggt	tttttgtttt	ttggtctgca	39660
tgatgtttta	aaacttgaga	caggccaggt	gcggtggcct	atgcctgtaa	tcccagcact	39720
ttgggaggcc	aaggtgggag	gatcacttga	acccgggggt	ttgaggctag	cctgggcaac	39780
atagggagac	cccatctcta	caaataataa	tttttaaaac	atcagccagg	tgtgggggca	39840
tgcaacctctg	gtcccagcta	ctcaggaggc	tgaggcagga	ggattgcttg	aacctgggag	39900
gtggaggctg	cagtgaaccg	tggttgtgcc	actgcactcc	agcctgggtg	acagagttag	39960
acccttacat	ggtggccact	ggctggagct	gagtatcagt	ggcctatatt	agaaaggggc	40020
tgggctttct	ggttcacac	tgccccgcc	actccttagt	gcttatacct	ggcccacatc	40080
actcattttct	gtcatctgcc	tggccctgtg	gtagacattt	gagtttgaaa	cccttgactc	40140
aaaggcaggc	tgatgtcttt	tgcttcctct	ggatcaatgg	aatttcattcc	aaggcaatgg	40200
aaccagctcc	taattgtgat	aactttttga	tcaccccttg	ccactgtgct	gaataggctt	40260
atggccatta	agaagagagg	acaggatgaa	tggccccctg	actgactgcc	cttcagggtt	40320
gttttttttag	ctgtcagcag	aagcatgctg	ggcagtgtac	caatcagggtg	tcagcaagtg	40380
tccttcacgc	gactgagttg	agcacaaaaa	catctgctcc	tggagagacc	tgagccctct	40440
gaaggcctca	gccagttatg	atgttaatgt	tcttttagaa	caaagtggta	gagcatttgt	40500
tgtctggaat	gagcaccaga	agaaatctct	ggccaaaaat	aaattgttga	gggccgggag	40560
cagtggctta	tacctgtaat	cccaacactt	tgggaggccg	aagcaggagg	atcacctgag	40620
gtcaggagtt	cgagaccagc	ctggccaaca	tggtgaaacc	ctgtctctcc	taaaaataca	40680
aaaattagct	ggacgtgggtg	acagggtgcct	gtaatacagc	tactgggagg	ctgaggcagg	40740

## -continued

---

gagaactgct tgaacccgga aggtggaggt tgcagtgagc cgggggtcatg ccacttcact	40800
ccagcctggg caacagagca aagctctgtc tttaaaaaca acttaaataa ataaataaat	40860
tgttgaggtc tgatgagtaa gtggacaagt tattttccag cagacacaca aaagagaagg	40920
aaattacagg ttatacgagg tatttcagaa aatataactt tctaaaacat aggaagttga	40980
agaagttgat cacattacag aattctgttg tttagaaaat gacctgtggc gaaatgtcct	41040
tattcagtgat ataggtgatt ccgcttatgc acgacctgtg tgaagtggat caggccaccc	41100
agaatgcacg acgcgttctt caggcccagc aggagtatgt gtctgtgtta atttcctgtg	41160
gctattatga ctaattgccca caaatgtggg ggcttaaaac aacagaaatt aatcttctta	41220
tagttctgga agccagaagt ttggaatcaa gatgtcagca gggccacact cgctctgatg	41280
ctctacggga gggctctctc ttgcctcttc cagcgtcttg tggctccagg cattccataa	41340
ctttatagca gcgtcccaca aatctctgcc tccatcctca catggccttc tccactgtgt	41400
ctctatgtct tcaatctctt tttttttttt tttttttttt gaggcagggt ttcactccag	41460
tctcctagac tgaagtgcaa tggcgtaatt tcgggtcact gcaacctctg cctcccgggc	41520
tcaagcgatt tgatctctcc tttatcttat aaagatacta gtcattggat ttggggctta	41580
ccctaaatcc aggataatct catcttgaga tgtttaactt aattatatct gcaaacactg	41640
tatttccaaa taaggtcata tcacagccac tagggattag atacttgaac atatcttatt	41700
tgggggctca acccattcca gtgtacgaaa aacactcttg ttcaaggccc gatgtttctc	41760
agggcatagc cactgacta cctgcatcag aataatcact tggtagctgt actggaaata	41820
cagactccta gaaacatctc agagcttctg caaccactct ttgagtgagg ggctcaggag	41880
tctgcctctg aacacactca cccaagtga ttctttcttt ctttcttttt tttttttttt	41940
tttgagatgg agtcttgctc tgtcccccag gctggagtgc agtggcgtga tctcggtca	42000
ctgcaagctc cgctcccgg gtacacagca ttctcttgcc tcagcctccc gagtagctgg	42060
gactacaggt gcccgcacc acaccggct aattttttgc atttttagta cagacggggt	42120
ttcacatgt tagccaggat ggtctcgatc tctgacctc gtgatctgcc tgcctcgcc	42180
tcccaaagt ctgggattac aggcagagc caccgcgcc agccatcacc ccaagtgatt	42240
cttgccttca gtttaagagc cactgtaaca agactatgga agcagaaatt cacgtgctta	42300
ctacacaatg ttaaccttc caggcaaacc aactcacata gggagataat gccaatccca	42360
gggcaggcag tggcaatgca tgcttgcttg cgaattaaaa aacaaatcac tgcctgggca	42420
cgttggctca tgcctataat cccaacaatt tgggaggctg aggtgggtgc atcacttgag	42480
ctcaggagtt caagccagc tgggcaacac agtgagacca tcatctctat aaaaaattt	42540
taaaaattag tcgggtgtaa tggtttgac ctgtagtccc agctgccggg aggtgaagt	42600
gggaggatga cttgagcccg ggaggcggag gccacagtga gctgtgttca agccactgca	42660
ctccagcctg gatgacagag ccagatcctg tctcaaaaa aacaaaaaca aacaacaaca	42720
acaaagataa atcactcaat acatcagcaa gagaaaaagc tctcttgaaa tagtcacatg	42780
caaaagaaat gaattccctc cagtacagaa gagccactaa agtgcctgag aatatctgat	42840
cgatttcaat gtcaggtttt gagaggtttt ttaaaaacag tttcagatgt ttcttactat	42900
tttttggcag aacatctgcc atctgcttcc ttctctccct acatcttgta actagacggg	42960
gaattaataa ctcaaaaaa ataaacatgt atatgtaact ttatctaaaa agaatacatca	43020

## -continued

aagtgtggtg	agaagaaggg	gcagatttaa	aagttttatg	aagcgttcat	tttaagcctc	43080
cttaattatt	cttgaaaaac	aaaacacacc	acttttcctg	actgcagcac	tggtgagggg	43140
tgcatatcac	gggtgactgt	gatgatttgt	gcctggcgct	taatttttaa	agttagtact	43200
gagtgatgac	agagagaggt	caatgccact	caaagaatat	tttgtttggt	tggttggttg	43260
tttgtttggt	tgagaggcag	tcttactctc	tcaactcaggc	cggagtgcag	tggtgtgata	43320
tcggctcact	gcaacctctg	cctcctgggt	tcaagtgatt	ctcctgcctc	agcttcctga	43380
gtagctggga	ttacaggcgt	ctgccaccac	atccagctaa	tttttgattt	tttagtagag	43440
atggggtttc	atcatgttgg	ccaggctggg	ctcaaactcc	tgacctgagg	tgatctgcct	43500
accttgccct	cccaaagtgc	tgggattaca	ggcgtgagcc	attgcacctg	gccaaagaaga	43560
atttattaca	tacttttccc	aagagatggg	gccacaccac	accataccag	gccacaccag	43620
gccatgccat	gctagacagg	gccacaggag	gaagtaccag	atccgctcaa	gaggcagaat	43680
aaagggtaaa	gaatggtcca	gagctttatt	gtgttactca	gtggaagggc	aaggcaggac	43740
tagggaaaca	gcttaggggt	aactactttg	aataatgccca	gtaggttctg	agttacagga	43800
atggtctcta	aatgtctggc	cctcacccta	cctgtcccta	aggagaaata	ctggagagtt	43860
agaaaaggag	gtggttgagg	gattggttgg	agggtttgta	atatgatttt	cacaccctca	43920
caaaagctgg	attgcagagg	agatgtaaac	aacttcagcc	ttgggaggcc	aagatgaaag	43980
gatggcttga	agccaggagt	tcaaggctgc	aacaagctat	gattacacca	ctgcaactcca	44040
gcctggttga	aggaatgaga	tcctgattct	acaaaacatt	tttgaaaaaa	acttttttat	44100
tttttctttt	cctttctttt	tttttttttt	tttttgagac	agtcttcttc	tgctacccag	44160
gctagagcgc	agtgatacga	tctcggctca	ctgcgacctc	catctcctgg	gttcaagtga	44220
ttctcgtgcc	tcagcgtcct	gagtagctgg	gattacaggt	gcctgccacc	atgcacggct	44280
aatttttgta	tttttagtag	agatgggggt	tcaccatggt	gtccgggctg	gtcttgaact	44340
cctgacctca	agtgatccac	ccgcctcggc	ctcccaaagt	gctgggatta	caggtgtgag	44400
ccacggtgcc	cgaccaaaaa	atttaaaaat	aaaaattagc	cacctgtagt	ggtgcacgct	44460
tgtagtccca	cccagctact	tgggaggctg	aggtgggagt	atcacttggg	cccaggaggt	44520
ggaggctgca	gtgagctctg	atcatgccag	tgctctccaa	cctgggtgac	agaacaagac	44580
ccccactcaa	aacaaaaaaa	aacaaaaaac	acaacttcag	cctttagttt	ggccctgtga	44640
ttaatgattg	ccaaataggc	atacacagaa	tctaagaaaa	tacagtttgc	tgagggtgtgc	44700
ctgtctccag	tccagtaatt	agtatgcaag	atttaccacc	gcccactccc	actttgttct	44760
agatgccaaa	cctcttcttc	ccccttaagg	aatagtcata	ttgcttgaag	tttttttttt	44820
ttaattttct	tgcttgttgt	taatcctgtg	ttggtttaaa	atgtgcattt	taatcttaag	44880
cgacaagctg	attttccgtc	actctgagat	gatgcaggta	caggtggtga	catggggagg	44940
ggggactgct	ctttggttcc	aggtggttgg	agagagaccc	agggctttgg	attgtgttcc	45000
ttccctgcc	acctgtcaca	gagccagggg	acagaccagg	aggactagac	aggccactgt	45060
tttggttttt	ccttccataa	aataccagca	ttttttgcca	gatgcagtgg	ctcatgcctg	45120
taaccccaaa	actttgggag	cctgaggcag	gcaaatcact	tgagctcagg	agttcgagac	45180
cagcctgggc	aacatggcaa	aagccogtct	ctacaaaaaa	tacaaaaaat	tagccgagca	45240
cgggtgtgca	tgctcttagt	cccagctacc	tgggagactg	aggtggggaga	atcatctgag	45300

## -continued

---

cctagcaagt caaggctgca gtgagtggta atcctgctac tgcactccag cctgggtgac	45360
agcgtgagac cctgtctcaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa ccccaaaacc	45420
cagcactttc aaaggatctt tacaaatata gatccttttc ttcctacaga tacgccaagt	45480
actacaagga aaacaatgtt gagaaacgga ctctgataaa agtcttcggg atccgttttg	45540
acatcctggt ttttggcacc gtaagtctcg tttcccagct ccgggcaccg gcatcctatg	45600
actgtgtcct aattactgct gtggggcctc catggaggga agggtttttg tctcagcctt	45660
cagctagcac tgggcatttc gcacatggga taaaagagga agaagatgtt ctcgggctgc	45720
tgccctgaa tgagtcatct gacgagtaca ggagtgggcc tggaaagcaa ttctaacatt	45780
cggctttaaa aatactctga tacttaacaa gagagagaaa gaaatctctc tgtaaaatag	45840
tttgtaagta acagaatcca attcaaattg gcttaagggg aaaaaaaaaa gttttgagt	45900
ggcttttttg ggttcacgta actgaaaaaa tccagggggt atttttaagc tacaagcatg	45960
gttgatcca ggtgctcaaa caacaacatc atcatcaaga acttgccacc ttcctgtctt	46020
tatatagctc ctcttggctc ttggcagcca tctctctatg aaggatcag gcagcacctt	46080
cgatctgacc agtttagcaa cctcagtcaa aagaaaacac ctctttgcca gaaggttcag	46140
caaaggtctc agacaaatgt cattggctct gattggccca ctatggatga catgcccact	46200
gtgaaactaa tcaactgtac cagggtcatg ccatactccc agtggttagt gtcgaaccag	46260
ccctaacctc gcccggggag cagcagagag gccaacacc cagccccttc aacacacaca	46320
cacacacaca cacacacaca cacacacaca cactacatgg actgaaaaga aaattggagt	46380
gttgcttctt gaagatggaa ggaaaagatg ccagctatgc aaaaatcaac atatccacg	46440
acatcctgtc caagagcatg ttatctatgt aacacagtga taagcaggac tagaagcaat	46500
aagatatagc tgagagaatg caaagaccgc tgggaataaa aaccacaagg ctgggccggg	46560
cacgggtggc catgcctgta atcccagctc tttgggaggc tgaggcagggt ggatcacaag	46620
gtcaggagct caagaccagc ctggccaaca tggtgaaacc ctgtctctac taaaaatgca	46680
aaaaattagc tgggcgtggc ggtgggtact tgtaatccca gctacttggg aggctgaggc	46740
aggagtatct cttgaaccca ggaggcggag cttgcagtga gccgagattg cgccactgca	46800
ctccagcctg ggcaaaagtg cgagactctg totcaaaaaa aaaaaaaaaa aaaaaccac	46860
gaggccacc agacattcct tccctctgct tctgaagtaa tgtcaaacct tttggaggag	46920
aagcttgtct gaaaccctca ccttgtatgg agaaatgata gcccttcagt ggtctctccc	46980
ttaatgtgtg ggcccagtg cttgtctatg taaaaccttt atgggtccaa gcacaccctg	47040
cacggctgaa gcaggatgcc tgagagtcat tttcagctc cgtagtctct gcctcagcca	47100
gcacttgaac gcatctatcc aagtcacagc atgaggctcc gctccctgat agaaccaaca	47160
attgcacgtt gaagcaaaag agcgttgctc tgaatttcac ctgagtaaac tctccactc	47220
tgtttttagg gaggaatttg tgacattatc cagctggttg tgtacatcgg ctcaaccctc	47280
tcctacttcg gtcgtgtaag agattctctt ttccatgctt taggaaaatg gtttgagaa	47340
ggaagtgact aacgcagcgc ttgtctgcat tctccccagg ccgctgtgtt catcgacttc	47400
ctcatcgaca ctactccag taactgctgt cgctcccata tttatccctg gtgcaagtgc	47460
tgtcagccct gtgtggtcaa cgaatactac tacaggaaga agtgcgagtc cattgtggag	47520
ccaaagccgg tgaggccgct gtgttcacag gacaccaaga catggagaga ttccatgaaa	47580

## -continued

---

tcactcagaa atgcacgaaa attagggcca aatcacaggc ttcacctgt agtggatacg	47640
tcgctgggtt ctaccccgat caaccaactc tcagataaat tttttgtct tagagaagaa	47700
tggaaacaaa aatggagggg caagatagag ggaaggcaaa tttttatgtc taggacttgc	47760
caattttgtc atttatttat ttattttttt atttatttat ttattttttt atttattttc	47820
agatggagtc ttgctctatc acccaggctg gagtgcagag gcatgatctc ggctcactgc	47880
agcctctgcc tcccaggttt aagtgtattc cctgcctcaa cctccaagt agctgggact	47940
acagggggcac accaccacac ctggctaatt tttgtatttt tagtagagat ggtgttttagc	48000
tatgttggcc aggctggtct caaactcctg acttcaaag atccaccac tcggcctccc	48060
acagtgtcgg gattataagt gtgagccact gagccagcc tgttttgtca tttattaaat	48120
tgttatagcc aaaaaagaaa aaagaaaaga aaaataacaa ctttgagaaa caatttggca	48180
gtgactaatg tttaaatggg acatacttta cagcctggca ttttcagttc tccatacctg	48240
ccctagagaa acactcacat gagtaccgcg agtacctgag tacaagatgt tcaaagcagg	48300
attgtttatt aaattatcaa taatatatga ttattaacag tgaagaaata gcatctaacc	48360
aaatatccaa cagggtgaatg gtgaaactat ggtaaatcca tagaaaggaa taccaggcac	48420
cagcttaaaa aaatgagata gataataata atggcaaaca tttacacagc acttctaaat	48480
gctttatcta ttaactcaat cttcacaaca acctgatgta gacagatgct actattatct	48540
ctttctatat atgaggaaat tgagccacag aaaggttaaa taattggccc aaggctgggc	48600
acagtggctc acgcctataa tcccacactc ttgggagggc gaggcaggca gatcacttga	48660
ggtaagagtc tcgagaccag cctggccaac atgggtgaaac cccatctcta ctaaaaaaac	48720
aaaaattagc caggcgtggt ggtgcatgcc tgaagtcca gctacttggg aggctgaggc	48780
aggagaattg cttgaacctg gtaggcagag gttgcagtga gctgagatca tgccattgca	48840
ctccagcctg agtgacagag ggaaactctg tctcaaaaaa aaaaaaaattat	48900
aggacttgac aagagcagat ctccaagaca ttttgttaaa agaaaaaaag caaactgtag	48960
aacaatatat gttatatgat atataggta aaaaaaaatc actagacaca gaagctaatac	49020
tgtatatatt ctgcatgtat attctatata tacacaagtg tacacacaca cacacataca	49080
tatacatatg tgtgggtgta tatatgtaca caaaattgta ccagtggctg cttctgga	49140
ggagcttagg gtgggggaggt aggagtagtc aaagagattt tagcgtcatc tgtattgttt	49200
tgatttgatt aattcagact ttatcaagca ggtcctctgc gttcaactcc atgattttcc	49260
ccaaagataa atctctggta cctaaaaaca aagacgattg gctagacgtg gtggctcacg	49320
cctgtagtcc cagtactttg ggaggccaag gaggtggat cacctgaggt caggagtcca	49380
aaaccagcct ggccaacatg gagaaacccc gtctctacta aaaatacaaa aaattagcca	49440
ggcatggtgg cgcctgcctg taatccagc ttctctagag gctgaggcag gagaatcgct	49500
tgaaccagg aggtggaggt tgtgtgagc caggattgca ccattgcact ccagcctggg	49560
caacaagagc gaaactccgt ctcaaaaaa aaaaaaaaac aaagacgatt tctttgtctt	49620
tccctcatcc aagaacatga ttgtcctgtt ccagcagctg atgcacaatt cactgtccat	49680
tgtatatgca ttcacaattt gaaataaaag ttcactcttg cagctaaaac taatcaccac	49740
ttcatggccc aagatgagat gaaatttaac aaacatgtaa ataatttaag ttgcaatagt	49800
acaaatttct ggagatactg aatctagagt tactgaaatt gacagaatac aacaaagaaa	49860



## -continued

---

ttttatgcag caactggggg gtccaatgta aaaacattaa gcagtaagct gtggctgtgt	49920
tgaatttaca agttaagatg catgggggtc cgcctggcgt ggtggctcac tcctgtaatc	49980
ccagcacttt ggtagcccaa ggcgggcgga tcaactgagg tcaggagtgc aagaccagcc	50040
tgaccaacat ggagaaaccc cgtctctact aaaaatacaa aattagctgg gcatgatggt	50100
gcatgcctgt aatcccagct actcaggagg ctgaggcagg agaatcattt gaacctggga	50160
ggcggagggt gcaagtggcc aagatcatgc cattgcactc cagcctggca acaagagtga	50220
actccaactc aaaaaaaaaa aaaaaaaaaa agcatggggt tccatttctg atttatcttt	50280
agactcagaa atcattaatt cttgggtaat gagagttttg agccagcttg ttcaatagtc	50340
tatcatttgg caaataggaa ttacagttgc ctttagatag gcaattcttg ataattctgt	50400
acaaaaatgg gtaaaacttc aaaccatctt ttcctagaca ttaaagtatg tgtcctttgt	50460
ggatgaatcc cacattagga tgggtaacca gcagctacta gggagaagtc tgcaagatgt	50520
caagggccaa gaagtcacag taagttaaatt cttttgtct tttttttttt ttttaagaaa	50580
tttactgtta aatataaaca catctagaaa cttgtacaaa tcaaaactga tggattttta	50640
caaagtaaac atactcatat aaccggcact cagattaaac aattgaaaat tactagcagg	50700
agtccctttt atgccccctt ccaatcacta ccctctcttt ctcccttttt aattttttaa	50760
atttggtctg gcacggtggc tcacacctgc aatcccagca ctttgggagg ccaagggtgg	50820
tggatcactt gaggtcagga gttcgagacc aacctagcca acatggtgaa acctcgtctc	50880
tactaaaaat acaaaaatta gccgggtggt gtggtggcac acatgtaatc ccagctactc	50940
agaagactga ggcaggagaa ctgcttgaac ccaggagggt gaggttgagc tgagccgaga	51000
ttgtgccatt gcactccagc ctgggtgaca gagcaagact ccactcaca aaaaaaaag	51060
aaaaaaatgt atattcttaa attacaaacg aggtctcact atgttgcca ggcgtgctc	51120
aagcagtcct cccacctcag ctgtaccaag cccaccaact gccctctctt aaaagtagtc	51180
attatcctgt ttccaagat taattttact ttggctagaa ttttctaaaa actgaatcac	51240
gtagtatgta agcggatat acgcggttga gtgtctggct tcctttactc aacattatct	51300
ttgtgagagt tgttcgatg gccatgtata gttcattctc attgtataat tctgttttat	51360
aaatatccaa ctatttcagc catcctactg ttgatggaca tttgggtagt gtccagtctg	51420
gggctaatac caataacgct gctatgtcga acatattgga ctctactgga cagttaccta	51480
agagtggagt tgctgagtca taaggcagac atatgttcgg ttttaggaga tactaacaaa	51540
cgggtgctgaa aaatgggtgt tcacatttgc actctcccca gcagtctctg ttgctgggca	51600
tcttcaatct cctagggctg aattaccaca aactaagtgg cttaaaacaa cagaaatggc	51660
caggcatggt ggctcatgcc tgtaattcca gcactttggg aggctgaggc aggcagatca	51720
cctgagggtc ggagtttgag accagcctga ccaacatgga gaaaccccat ctctactaaa	51780
aatacaaaaa aatagccagg catggtggca ggcgactgta gtcccagcca ctccaggaggc	51840
tgaggcagga gaatggcatg aacctgggag gcagagcttg cagttagcca agattgcacc	51900
actttgagag gccaaaggtg gcggatcacc tgaggtcagg agttcaggaa cagcctggcc	51960
aacatggcga aacctgtct ctattaaaaa tacaaaaatt agccaggcgt ggtggtgcac	52020
acctgtaatc ccagctactc aggacgtga ggtgggagaa ttgcttgaac ctgggaggca	52080
gaggctgcgg ttagctgaga tcacacctct gcactctagc ctgggcaaca gaacaagctc	52140

## -continued

---

catctaaaaa aaaaaaaaaa aaaaagtctt ttctgctcct tttgaaatat cagtgaagttt	52200
tcttcttttt ttctgttagt tgaattgtac tgattgattt tcaaatatta agccaaactt	52260
gcattcctga agtaactca atttgaatgt gttgtactat tctttgtatt tattgctgaa	52320
ttccattcac taatatattag gatttttaca tctcttcttg agaagactg accaaagtgt	52380
ttccattcct gtaatgttct tgttggaattt gtgtatgaag tgaactacag tcatgcatca	52440
cttaatgatg gggatatgtt ctgaaaaacg catcagtagc tgattctgtg gttgtctgaa	52500
tatcatggac tctatttaca caaacctaaa tagaatagcc tattatactt aggttatatg	52560
gtgtagtcta ttgctcctag gctgcaaacc tgtacagcat gttactgtac tgaatacggg	52620
aagcaactgt aacagaatgg taagatttgt atatctaaat atagaaaagg tacagtga	52680
atatgatata aaagcttaaa aatggtacat ctgcataggg cacttaccat gaatggagct	52740
tgtaggactg aaagttgctc tgggtgagtc agtgagtagt gagtgaatgt gaaagcctag	52800
gtgtttaccg tgcactacag tagacttcat aaacactgta cacttaggct acactaaatt	52860
tacttcaaaa tatttatctt tcttcaataa taaattaatc ttagcttact gtgactgttt	52920
tactttataa atttttaaat tttttaaact ctgtacagtg gtataaaaat attttctttc	52980
tcaccgggag tgggtgctca tgcctgtaat cccatcactt tgggaggccg aggcaggcgg	53040
atcacaagat caggagattg agaccatcct ggccaacatg gtgaaacccc atctctacta	53100
aaaatacaaa aaaatagcca ggcattggtg caggcgcctg tagtcccagc tactcgggag	53160
gctgaggcag gagaatggcg tgaacccagg aggcagagct tgcagtgagc caagatcaca	53220
ccactgcact ccagcctggg cgacagagca agactccatc tcaaaaaaaaa aaaaattctt	53280
tctctatata cttattctat atactttttt ctattttttaa cattttttat ttttattttt	53340
acttttttaa tatttttgtt aaaaactaag tcatggcccg gtgcagtggc tcacgcttgt	53400
aatctcagta ctttgggagg ctgaggtggg tggatcactc gaggtcagga gttcaacacc	53460
agcctggcca atatggtgaa actctgtctc tactaaaaat ataaaaatta gccgggtgtg	53520
gtggtgcacg cctgtagtcc cagctactca ggaggctgag gcagaagaat cgcttgaacc	53580
caggaggcgg aggttgagc aagccaagat catgccactg cactccagcc tgggagacag	53640
agcaagactc catctcaaaa aaaaaaaaaa acaacaaaca cacacacaca cacacacaca	53700
aaaaccattt ccaatagtgc taagtcttat gaagaatgtg gaataccaca gtgtgataag	53760
ggaatcatgt gggagaaaag ctgctagata ggggtgtcag gacaagaggt gacatctcaa	53820
cagaggcctg gcttttctga atttcatttt ccaaaatctg taaaaataggc caggtgcagt	53880
ggctcatgcc tgtaatccca gcactttgag aggctgaggg aggtggatca cctgaggtca	53940
ggagttccaa acaagcctgg ccaacatgtt gaaatcctgt ttctactaga aatacgaaag	54000
aattagctgg gcatggtggc atgcacctgt aataccagct acttaggagg ttaaggcatg	54060
aaaattgtct gaacctggga ggtgaaagt gcagtgaacc aagatcacac cactgtgctc	54120
cagcctgtgc gacagagtga gaccctctct caaaaaaaaa aaaaaaatct gtaaaataaa	54180
gacaaggata cattatctca caagcgtctt caaaggcctg aatgaggcaa tgcttacaga	54240
acacatgcat ggtcctgata tctacaccta ataaatgacg gctactataa atcatgtaat	54300
attaacgta actttataag ttaataaaat taaagaacct agaacctgag ggcttgtcat	54360
ggctaataag ttgtgaaact tgctttttca gagacctgag atggacttca cagatttgtc	54420

## -continued

caggctgccc	ctggccctcc	atgacacacc	cccgattcct	ggacaaccag	aggagataca	54480
gctgcttaga	aaggaggcga	ctcctagatc	cagggatagc	cccgtctggt	gccagtgtgg	54540
aagctgcctc	ccatctcaac	tccctgagag	ccacaggtgc	ctggaggagc	tgtgctgccg	54600
gaaaagccg	ggggcctgca	tcaccacctc	agagctgttc	aggaaagctgg	tcctgtccag	54660
acacgtcctg	cagttcctcc	tgctctacca	ggagcccttg	ctggcgctgg	atgtggattc	54720
caccaacagc	cggctgcggc	actgtgccta	caggtgctac	gccacctggc	gcttcggctc	54780
ccaggacatg	gctgactttg	ccaacctgcc	cagctgctgc	cgctggagga	tccggaaaga	54840
gtttccgaag	agtgaagggc	agtacagtgg	cttcaagagt	ccttactgaa	gccaggcacc	54900
gtggctcacg	tctgtaatcc	cagcgctttg	ggaggccgag	gcaggcagat	cacctgagat	54960
cgggagttgg	agaccgcctc	ggctaacaag	gcgaaatcct	gtctgtacta	aaaatacaaa	55020
aatcagccag	acatggtggc	atgcacctgc	aatcccagct	actcgggagg	ctgaggcaca	55080
agaatcactt	gaaccgggga	ggcagagggt	gtagtgaacc	cagattgtgc	cactgtcttc	55140
cagcctggga	ggcacagcaa	actgtcccca	aaaaaaaaaa	aagagtcctt	accaatagca	55200
ggggctgcag	tagccatggt	aacatgacat	ttaccagcaa	cttgaacttc	acctgcaaag	55260
ctctgtggcc	acatttttcag	ccaaagggaa	atatgctttc	atcttctgtt	gctctctgtg	55320
tctgagagca	aagtgacctg	gttaaacaaa	ccagaatccc	tctacatgga	ctcagagaaa	55380
agagattgag	atgtaagtct	caactctgtc	cccagggaagt	tgtgtgaccc	taggcctctc	55440
acctctgtgc	ctctgtctcc	ttgttgccca	actactatct	cagagatatt	gtgaggacaa	55500
attgagacag	tgacatgaa	ctgtctttta	atgtgtaaag	atctacatga	atgcaaaaca	55560
tttcattatg	aggtcagact	aggataatgt	ccaactaaaa	acaaaccctt	ttcatcctgg	55620
ctggagaatg	tggagaacta	aaggtggcca	caaattcttt	gacactcaag	tcccccaaga	55680
cctaagggtt	ttatctctcc	cccttgaata	tgggtggctc	tgattgcttt	atccaaaagt	55740
ggaagtgaca	ttgtgtcagt	ttcagatcct	gatcttaaga	ggctgacagc	ttctacttgc	55800
tgcccttg	aactctgtct	atcggggaag	ccagacgcca	tttaaaagtc	tgccctatcct	55860
ggccaggtgt	ggtggctcac	acctgtaatc	ccagcacttt	gggagaccaa	ggcgggcgga	55920
tcacttaaa	tcaggagtcc	aagaccagac	tcgccaacat	ggtgaaaccg	tatctctaata	55980
aaaaatacaa	aaattagctg	ggcatggtgc	gggcacctgt	agtcctagct	atcaagaggc	56040
tgagacagga	gaaacacttg	aacctgggag	gtggagggtg	cattgagctg	agatcggtgc	56100
actgcactcc	aggctgggtg	acagagcgag	actccatctc	aaaaaaaaaa	aaaaaagaaa	56160
aaaaaaatgt	ctgcctatcc	tgagactgcc	ctgctgtgag	gaagcccaag	cagtcacgtg	56220
gacagtcctc	gaccagcccc	agctttcaag	ccatccaagc	ccagtcacca	aacatgagag	56280
agaagaagcc	ttcagggtgat	tctggactcc	actaacatat	gactgatacc	gcatgataca	56340
tcccaagtga	gaactgcccc	ataaatccag	aaaaccacat	tgctatctta	agtccctaag	56400
tttggggctt	atttgttcca	cagcaacagg	taactggaac	agagggcaag	cctgatgaat	56460
gggcacacag	actcagccca	taccttcctc	ggttctaata	ttctcaggga	gcccgaccca	56520
acctggggag	cctcaggaac	ttaggtttcc	actggacagt	tctagaaggg	ctatagacca	56580

<210> SEQ ID NO 2  
<211> LENGTH: 2164  
<212> TYPE: DNA

-continued

&lt;213&gt; ORGANISM: Homo sapiens

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: CDS

&lt;222&gt; LOCATION: (79)..(1863)

&lt;223&gt; OTHER INFORMATION:

&lt;400&gt; SEQUENCE: 2

```

ggcacgaggg cttgctgtgg ccctgtcagg aagagtagag ctctggtcca gctccgcgca      60
gggagggagg ctgtcacc  atg ccg gcc tgc tgc agc tgc agt gat gtt ttc      111
                        Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe
                        1          5          10
cag tat gag acg aac aaa gtc act cgg atc cag agc atg aat tat ggc      159
Gln Tyr Glu Thr Asn Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly
                        15          20          25
acc att aag tgg ttc ttc cac gtg atc atc ttt tcc tac gtt tgc ttt      207
Thr Ile Lys Trp Phe Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe
                        30          35          40
gct ctg gtg agt gac aag ctg tac cag cgg aaa gag cct gtc atc agt      255
Ala Leu Val Ser Asp Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser
                        45          50          55
tct gtg cac acc aag gtg aag ggg ata gca gag gtg aaa gag gag atc      303
Ser Val His Thr Lys Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile
                        60          65          70          75
gtg gag aat gga gtg aag aag ttg gtg cac agt gtc ttt gac acc gca      351
Val Glu Asn Gly Val Lys Lys Leu Val His Ser Val Phe Asp Thr Ala
                        80          85          90
gac tac acc ttc cct ttg cag ggg aac tct ttc ttc gtg atg aca aac      399
Asp Tyr Thr Phe Pro Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn
                        95          100          105
ttt ctc aaa aca gaa ggc caa gag cag cgg ttg tgt ccc gag tat ccc      447
Phe Leu Lys Thr Glu Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro
                        110          115          120
acc cgc agg acg ctc tgt tcc tct gac cga ggt tgt aaa aag gga tgg      495
Thr Arg Arg Thr Leu Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp
                        125          130          135
atg gac ccg cag agc aaa gga att cag acc gga agg tgt gta gtg cat      543
Met Asp Pro Gln Ser Lys Gly Ile Gln Thr Gly Arg Cys Val Val His
                        140          145          150          155
gaa ggg aac cag aag acc tgt gaa gtc tct gcc tgg tgc ccc atc gag      591
Glu Gly Asn Gln Lys Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu
                        160          165          170
gca gtg gaa gag gcc ccc cgg cct gct ctc ttg aac agt gcc gaa aac      639
Ala Val Glu Glu Ala Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn
                        175          180          185
ttc act gtg ctc atc aag aac aat atc gac ttc ccc ggc cac aac tac      687
Phe Thr Val Leu Ile Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr
                        190          195          200
acc acg aga aac atc ctg cca ggt tta aac atc act tgt acc ttc cac      735
Thr Thr Arg Asn Ile Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His
                        205          210          215
aag act cag aat cca cag tgt ccc att ttc cga cta gga gac atc ttc      783
Lys Thr Gln Asn Pro Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe
                        220          225          230          235
cga gaa aca ggc gat aat ttt tca gat gtg gca att cag ggc gga ata      831
Arg Glu Thr Gly Asp Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile
                        240          245          250
atg ggc att gag atc tac tgg gac tgc aac cta gac cgt tgg ttc cat      879
Met Gly Ile Glu Ile Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His

```

## -continued

255	260	265	
cac tgc cat ccc aaa tac agt ttc cgt cgc ctt gac gac aag acc acc His Cys His Pro Lys Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr 270 275 280			927
aac gtg tcc ttg tac cct ggc tac aac ttc aga tac gcc aag tac tac Asn Val Ser Leu Tyr Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr 285 290 295			975
aag gaa aac aat gtt gag aaa cgg act ctg ata aaa gtc ttc ggg atc Lys Glu Asn Asn Val Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile 300 305 310 315			1023
cgt ttt gac atc ctg gtt ttt ggc acc gga gga aaa ttt gac att atc Arg Phe Asp Ile Leu Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile 320 325 330			1071
cag ctg gtt gtg tac atc ggc tca acc ctc tcc tac ttc ggt ctg gcc Gln Leu Val Val Tyr Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala 335 340 345			1119
gct gtg ttc atc gac ttc ctc atc gac act tac tcc agt aac tgc tgt Ala Val Phe Ile Asp Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys 350 355 360			1167
cgc tcc cat att tat ccc tgg tgc aag tgc tgt cag ccc tgt gtg gtc Arg Ser His Ile Tyr Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val 365 370 375			1215
aac gaa tac tac tac agg aag aag tgc gag tcc att gtg gag cca aag Asn Glu Tyr Tyr Tyr Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys 380 385 390 395			1263
ccg aca tta aag tat gtg tcc ttt gtg gat gaa tcc cac att agg atg Pro Thr Leu Lys Tyr Val Ser Phe Val Asp Glu Ser His Ile Arg Met 400 405 410			1311
gtg aac cag cag cta cta ggg aga agt ctg caa gat gtc aag ggc caa Val Asn Gln Gln Leu Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln 415 420 425			1359
gaa gtc cca aga cct gcg atg gac ttc aca gat ttg tcc agg ctg ccc Glu Val Pro Arg Pro Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro 430 435 440			1407
ctg gcc ctc cat gac aca ccc ccg att cct gga caa cca gag gag ata Leu Ala Leu His Asp Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile 445 450 455			1455
cag ctg ctt aga aag gag gcg act cct aga tcc agg gat agc ccc gtc Gln Leu Leu Arg Lys Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val 460 465 470 475			1503
tgg tgc cag tgt gga agc tgc ctc cca tct caa ctc cct gag agc cac Trp Cys Gln Cys Gly Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His 480 485 490			1551
agg tgc ctg gag gag ctg tgc tgc ccg aaa aag ccg ggg gcc tgc atc Arg Cys Leu Glu Glu Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile 495 500 505			1599
acc acc tca gag ctg ttc agg aag ctg gtc ctg tcc aga cac gtc ctg Thr Thr Ser Glu Leu Phe Arg Lys Leu Val Leu Ser Arg His Val Leu 510 515 520			1647
cag ttc ctc ctg ctc tac cag gag ccc ttg ctg gcg ctg gat gtg gat Gln Phe Leu Leu Leu Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp 525 530 535			1695
tcc acc aac agc ccg ctg ccg cac tgt gcc tac agg tgc tac gcc acc Ser Thr Asn Ser Arg Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr 540 545 550 555			1743
tgg cgc ttc ggc tcc cag gac atg gct gac ttt gcc atc ctg ccc agc Trp Arg Phe Gly Ser Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser 560 565 570 575			1791

## -continued

560	565	570	
tgc tgc cgc tgg agg atc cgg aaa gag ttt ccg aag agt gaa ggg cag			1839
Cys Cys Arg Trp Arg Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln			
575	580	585	
tac agt ggc ttc aag agt cct tac tgaagccagg caccgtggct cacgtctgta			1893
Tyr Ser Gly Phe Lys Ser Pro Tyr			
590	595		
atcccagcgc ttggggaggc cgaggcaggc agatcacctg aggtcgggag ttggagaccc			1953
gcctgggctaa caaggcgaaa tcctgtctgt actaaaaata caaaaatcag ccagacatgg			2013
tggcattgcac ctgcaatccc agctactcgg gaggctgagg cacaagaatc acttgaaccc			2073
gggaggcaga ggttgtagt agcccagatt gtgccactgc tctccagcct gggaggcaca			2133
gcaaactgtc ccaaaaaaaaa aaaaaaaaaa a			2164

&lt;210&gt; SEQ ID NO 3

&lt;211&gt; LENGTH: 595

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 3

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn		
1	5	10
Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe		
20	25	30
Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp		
35	40	45
Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys		
50	55	60
Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val		
65	70	75
Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro		
85	90	95
Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu		
100	105	110
Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu		
115	120	125
Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser		
130	135	140
Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys		
145	150	155
Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala		
165	170	175
Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile		
180	185	190
Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile		
195	200	205
Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro		
210	215	220
Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp		
225	230	235
Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile		
245	250	255

-continued

---

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys His Pro Lys  
                   260                                  265                                  270  
 Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
                   275                                  280                                  285  
 Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
                   290                                  295                                  300  
 Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
   305                                  310                                  315                                  320  
 Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
                   325                                  330                                  335  
 Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp  
                   340                                  345                                  350  
 Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr  
                   355                                  360                                  365  
 Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr  
                   370                                  375                                  380  
 Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
   385                                  390                                  395                                  400  
 Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
                   405                                  410                                  415  
 Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
                   420                                  425                                  430  
 Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
                   435                                  440                                  445  
 Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
                   450                                  455                                  460  
 Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
   465                                  470                                  475                                  480  
 Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
                   485                                  490                                  495  
 Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
                   500                                  505                                  510  
 Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
                   515                                  520                                  525  
 Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
                   530                                  535                                  540  
 Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
   545                                  550                                  555                                  560  
 Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg  
                   565                                  570                                  575  
 Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
                   580                                  585                                  590  
 Ser Pro Tyr  
                   595

<210> SEQ ID NO 4  
 <211> LENGTH: 595  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: DOMAIN  
 <222> LOCATION: (1)..(20)  
 <223> OTHER INFORMATION: Intracellular domain  
 <220> FEATURE:

---

-continued

---

<221> NAME/KEY: TRANSMEM  
<222> LOCATION: (21)..(46)  
<223> OTHER INFORMATION:  
<220> FEATURE:  
<221> NAME/KEY: DOMAIN  
<222> LOCATION: (47)..(320)  
<223> OTHER INFORMATION: Extracellular domain  
<220> FEATURE:  
<221> NAME/KEY: TRANSMEM  
<222> LOCATION: (321)..(356)  
<223> OTHER INFORMATION:  
<220> FEATURE:  
<221> NAME/KEY: DOMAIN  
<222> LOCATION: (357)..(595)  
<223> OTHER INFORMATION: Intracellular domain

<400> SEQUENCE: 4

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile  
180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys  
260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu



-continued

305		310		315		320
Val Phe Gly Thr Gly	Gly Lys Phe Asp Ile	Ile Gln Leu Val Val Tyr				
	325	330			335	
Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp		345			350	
	340					
Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr		360			365	
	355					
Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr		375			380	
	370					
Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr		390			395	400
Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu		405		410		415
Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro		420		425		430
Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp		435		440		445
Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys		450		455		460
Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly		465		470		475
						480
Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu		485		490		495
Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu		500		505		510
Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu		515		520		525
Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg		530		535		540
Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser		545		550		555
						560
Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg		565		570		575
Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys		580		585		590
Ser Pro Tyr						
	595					

&lt;210&gt; SEQ ID NO 5

&lt;211&gt; LENGTH: 595

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 5

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn									
1		5			10			15	
Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe									
	20			25			30		
Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp									
	35		40			45			
Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys									
	50		55			60			

## -continued

Val	Lys	Gly	Ile	Ala	Glu	Val	Lys	Glu	Glu	Ile	Val	Glu	Asn	Gly	Val	65	70	75	80
Lys	Lys	Leu	Val	His	Ser	Val	Phe	Asp	Thr	Ala	Asp	Tyr	Thr	Phe	Pro	85	90	95	
Leu	Gln	Gly	Asn	Ser	Phe	Phe	Val	Met	Thr	Asn	Phe	Leu	Lys	Thr	Glu	100	105	110	
Gly	Gln	Glu	Gln	Trp	Leu	Cys	Pro	Glu	Tyr	Pro	Thr	Arg	Arg	Thr	Leu	115	120	125	
Cys	Ser	Ser	Asp	Arg	Gly	Cys	Lys	Lys	Gly	Trp	Met	Asp	Pro	Gln	Ser	130	135	140	
Lys	Gly	Ile	Gln	Thr	Gly	Arg	Cys	Val	Val	His	Glu	Gly	Asn	Gln	Lys	145	150	155	160
Thr	Cys	Glu	Val	Ser	Ala	Trp	Cys	Pro	Ile	Glu	Ala	Val	Glu	Glu	Ala	165	170	175	
Pro	Arg	Pro	Ala	Leu	Leu	Asn	Ser	Ala	Glu	Asn	Phe	Thr	Val	Leu	Ile	180	185	190	
Lys	Asn	Asn	Ile	Asp	Phe	Pro	Gly	His	Asn	Tyr	Thr	Thr	Arg	Asn	Ile	195	200	205	
Leu	Pro	Gly	Leu	Asn	Ile	Thr	Cys	Thr	Phe	His	Lys	Thr	Gln	Asn	Pro	210	215	220	
Gln	Cys	Pro	Ile	Phe	Arg	Leu	Gly	Asp	Ile	Phe	Arg	Glu	Thr	Gly	Asp	225	230	235	240
Asn	Phe	Ser	Asp	Val	Ala	Ile	Gln	Gly	Gly	Ile	Met	Gly	Ile	Glu	Ile	245	250	255	
Tyr	Trp	Asp	Cys	Asn	Leu	Asp	Arg	Trp	Phe	His	His	Cys	Arg	Pro	Lys	260	265	270	
Tyr	Ser	Phe	Arg	Arg	Leu	Asp	Asp	Lys	Thr	Thr	Asn	Val	Ser	Leu	Tyr	275	280	285	
Pro	Gly	Tyr	Asn	Phe	Arg	Tyr	Ala	Lys	Tyr	Tyr	Lys	Glu	Asn	Asn	Val	290	295	300	
Glu	Lys	Arg	Thr	Leu	Ile	Lys	Val	Phe	Gly	Ile	Arg	Phe	Asp	Ile	Leu	305	310	315	320
Val	Phe	Gly	Thr	Gly	Gly	Lys	Phe	Asp	Ile	Ile	Gln	Leu	Val	Val	Tyr	325	330	335	
Ile	Gly	Ser	Thr	Leu	Ser	Tyr	Phe	Gly	Leu	Ala	Ala	Val	Phe	Ile	Asp	340	345	350	
Phe	Leu	Ile	Asp	Thr	Tyr	Ser	Ser	Asn	Cys	Cys	Arg	Ser	His	Ile	Tyr	355	360	365	
Pro	Trp	Cys	Lys	Cys	Cys	Gln	Pro	Cys	Val	Val	Asn	Glu	Tyr	Tyr	Tyr	370	375	380	
Arg	Lys	Lys	Cys	Glu	Ser	Ile	Val	Glu	Pro	Lys	Pro	Thr	Leu	Lys	Tyr	385	390	395	400
Val	Ser	Phe	Val	Asp	Glu	Ser	His	Ile	Arg	Met	Val	Asn	Gln	Gln	Leu	405	410	415	
Leu	Gly	Arg	Ser	Leu	Gln	Asp	Val	Lys	Gly	Gln	Glu	Val	Pro	Arg	Pro	420	425	430	
Ala	Met	Asp	Phe	Thr	Asp	Leu	Ser	Arg	Leu	Pro	Leu	Ala	Leu	His	Asp	435	440	445	
Thr	Pro	Pro	Ile	Pro	Gly	Gln	Pro	Glu	Glu	Ile	Gln	Leu	Leu	Arg	Lys	450	455	460	
Glu	Ala	Thr	Pro	Arg	Ser	Arg	Asp	Ser	Pro	Val	Trp	Cys	Gln	Cys	Gly				

## -continued

---

465		470		475		480
Ser Cys Leu Pro	Ser Gln Leu Pro Glu	Ser His Arg Cys Leu Glu Glu				
	485	490			495	
Leu Cys Cys Arg	Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu					
	500	505			510	
Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu						
	515	520			525	
Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg						
	530	535			540	
Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser						
	545	550			555	560
Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg						
	565	570			575	
Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys						
	580	585			590	
Ser Pro Tyr						
	595					

<210> SEQ ID NO 6  
 <211> LENGTH: 595  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn		
1	5	10 15
Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe		
	20	25 30
Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp		
	35	40 45
Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys		
	50	55 60
Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val		
	65	70 75 80
Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro		
	85	90 95
Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu		
	100	105 110
Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu		
	115	120 125
Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser		
	130	135 140
Lys Gly Ile Gln Thr Arg Arg Cys Val Val His Glu Gly Asn Gln Lys		
	145	150 155 160
Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala		
	165	170 175
Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile		
	180	185 190
Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile		
	195	200 205
Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro		
	210	215 220

-continued

Gln	Cys	Pro	Ile	Phe	Arg	Leu	Gly	Asp	Ile	Phe	Arg	Glu	Thr	Gly	Asp	
225					230					235					240	
Asn	Phe	Ser	Asp	Val	Ala	Ile	Gln	Gly	Gly	Ile	Met	Gly	Ile	Glu	Ile	
				245					250					255		
Tyr	Trp	Asp	Cys	Asn	Leu	Asp	Arg	Trp	Phe	His	His	Cys	Arg	Pro	Lys	
			260					265					270			
Tyr	Ser	Phe	Arg	Arg	Leu	Asp	Asp	Lys	Thr	Thr	Asn	Val	Ser	Leu	Tyr	
		275					280					285				
Pro	Gly	Tyr	Asn	Phe	Arg	Tyr	Ala	Lys	Tyr	Tyr	Lys	Glu	Asn	Asn	Val	
	290					295					300					
Glu	Lys	Arg	Thr	Leu	Ile	Lys	Val	Phe	Gly	Ile	Arg	Phe	Asp	Ile	Leu	
305					310					315					320	
Val	Phe	Gly	Thr	Gly	Gly	Lys	Phe	Asp	Ile	Ile	Gln	Leu	Val	Val	Tyr	
				325					330					335		
Ile	Gly	Ser	Thr	Leu	Ser	Tyr	Phe	Gly	Leu	Ala	Ala	Val	Phe	Ile	Asp	
			340					345					350			
Phe	Leu	Ile	Asp	Thr	Tyr	Ser	Ser	Asn	Cys	Cys	Arg	Ser	His	Ile	Tyr	
		355					360					365				
Pro	Trp	Cys	Lys	Cys	Cys	Gln	Pro	Cys	Val	Val	Asn	Glu	Tyr	Tyr	Tyr	
	370					375					380					
Arg	Lys	Lys	Cys	Glu	Ser	Ile	Val	Glu	Pro	Lys	Pro	Thr	Leu	Lys	Tyr	
385					390					395					400	
Val	Ser	Phe	Val	Asp	Glu	Ser	His	Ile	Arg	Met	Val	Asn	Gln	Gln	Leu	
				405					410					415		
Leu	Gly	Arg	Ser	Leu	Gln	Asp	Val	Lys	Gly	Gln	Glu	Val	Pro	Arg	Pro	
			420					425					430			
Ala	Met	Asp	Phe	Thr	Asp	Leu	Ser	Arg	Leu	Pro	Leu	Ala	Leu	His	Asp	
		435				440						445				
Thr	Pro	Pro	Ile	Pro	Gly	Gln	Pro	Glu	Glu	Ile	Gln	Leu	Leu	Arg	Lys	
	450					455					460					
Glu	Ala	Thr	Pro	Arg	Ser	Arg	Asp	Ser	Pro	Val	Trp	Cys	Gln	Cys	Gly	
465					470					475					480	
Ser	Cys	Leu	Pro	Ser	Gln	Leu	Pro	Glu	Ser	His	Arg	Cys	Leu	Glu	Glu	
				485					490					495		
Leu	Cys	Cys	Arg	Lys	Lys	Pro	Gly	Ala	Cys	Ile	Thr	Thr	Ser	Glu	Leu	
			500					505					510			
Phe	Arg	Lys	Leu	Val	Leu	Ser	Arg	His	Val	Leu	Gln	Phe	Leu	Leu	Leu	
		515					520					525				
Tyr	Gln	Glu	Pro	Leu	Leu	Ala	Leu	Asp	Val	Asp	Ser	Thr	Asn	Ser	Arg	
	530					535					540					
Leu	Arg	His	Cys	Ala	Tyr	Arg	Cys	Tyr	Ala	Thr	Trp	Arg	Phe	Gly	Ser	
545					550					555					560	
Gln	Asp	Met	Ala	Asp	Phe	Ala	Ile	Leu	Pro	Ser	Cys	Cys	Arg	Trp	Arg	
				565					570					575		
Ile	Arg	Lys	Glu	Phe	Pro	Lys	Ser	Glu	Gly	Gln	Tyr	Ser	Gly	Phe	Lys	
			580					585					590			
Ser	Pro	Tyr														
		595														

&lt;210&gt; SEQ ID NO 7

&lt;211&gt; LENGTH: 595

&lt;212&gt; TYPE: PRT

-continued

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 7

```

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
 1          5          10          15
Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
 20          25          30
Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
 35          40          45
Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
 50          55          60
Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
 65          70          75          80
Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
 85          90          95
Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
100          105          110
Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
115          120          125
Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
130          135          140
Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
145          150          155          160
Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
165          170          175
Pro Arg Pro Ala Leu Leu Asn Ser Ala Lys Asn Phe Thr Val Leu Ile
180          185          190
Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
195          200          205
Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
210          215          220
Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
225          230          235          240
Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
245          250          255
Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys
260          265          270
Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
275          280          285
Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
290          295          300
Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
305          310          315          320
Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
325          330          335
Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
340          345          350
Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
355          360          365
Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
370          375          380

```

## -continued

---

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
 385 390 395 400  
 Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
 405 410 415  
 Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
 420 425 430  
 Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
 435 440 445  
 Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
 450 455 460  
 Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
 465 470 475 480  
 Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
 485 490 495  
 Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
 500 505 510  
 Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
 515 520 525  
 Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
 530 535 540  
 Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
 545 550 555 560  
 Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg  
 565 570 575  
 Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
 580 585 590  
 Ser Pro Tyr  
 595

&lt;210&gt; SEQ ID NO 8

&lt;211&gt; LENGTH: 595

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 8

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
 1 5 10 15  
 Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
 20 25 30  
 Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
 35 40 45  
 Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
 50 55 60  
 Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
 65 70 75 80  
 Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
 85 90 95  
 Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
 100 105 110  
 Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
 115 120 125  
 Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
 130 135 140

-continued

---

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
 145 150 155 160  
 Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
 165 170 175  
 Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Pro Ile  
 180 185 190  
 Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205  
 Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
 210 215 220  
 Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
 225 230 235 240  
 Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
 245 250 255  
 Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys  
 260 265 270  
 Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
 275 280 285  
 Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
 290 295 300  
 Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
 305 310 315 320  
 Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
 325 330 335  
 Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp  
 340 345 350  
 Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr  
 355 360 365  
 Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr  
 370 375 380  
 Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
 385 390 395 400  
 Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
 405 410 415  
 Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
 420 425 430  
 Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
 435 440 445  
 Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
 450 455 460  
 Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
 465 470 475 480  
 Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
 485 490 495  
 Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
 500 505 510  
 Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
 515 520 525  
 Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
 530 535 540

-continued

---

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
 545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg  
 565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
 580 585 590

Ser Pro Tyr  
 595

<210> SEQ ID NO 9  
 <211> LENGTH: 595  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
 165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile  
 180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
 210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
 225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
 245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Cys Pro Lys  
 260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
 275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
 290 295 300



-continued

---

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
 305 310 315 320  
 Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
 325 330 335  
 Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp  
 340 345 350  
 Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr  
 355 360 365  
 Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr  
 370 375 380  
 Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
 385 390 395 400  
 Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
 405 410 415  
 Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
 420 425 430  
 Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
 435 440 445  
 Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
 450 455 460  
 Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
 465 470 475 480  
 Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
 485 490 495  
 Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
 500 505 510  
 Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
 515 520 525  
 Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
 530 535 540  
 Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
 545 550 555 560  
 Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg  
 565 570 575  
 Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
 580 585 590  
 Ser Pro Tyr  
 595

<210> SEQ ID NO 10

<211> LENGTH: 595

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
 1 5 10 15  
 Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
 20 25 30  
 Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
 35 40 45  
 Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys

-continued

50					55					60					
Val 65	Lys	Gly	Ile	Ala	Glu 70	Val	Lys	Glu	Glu	Ile 75	Val	Glu	Asn	Gly	Val 80
Lys	Lys	Leu	Val	His 85	Ser	Val	Phe	Asp	Thr 90	Ala	Asp	Tyr	Thr	Phe 95	Pro
Leu	Gln	Gly	Asn 100	Ser	Phe	Phe	Val	Met 105	Thr	Asn	Phe	Leu	Lys 110	Thr	Glu
Gly	Gln	Glu	Gln 115	Arg	Leu	Cys	Pro 120	Glu	Tyr	Pro	Thr	Arg 125	Arg	Thr	Leu
Cys	Ser 130	Ser	Asp	Arg	Gly	Cys 135	Lys	Lys	Gly	Trp	Met 140	Asp	Pro	Gln	Ser
Lys 145	Gly	Ile	Gln	Thr	Gly 150	Arg	Cys	Val	Val	His 155	Glu	Gly	Asn	Gln	Lys 160
Thr	Cys	Glu	Val 165	Ser	Ala	Trp	Cys	Pro	Ile 170	Glu	Ala	Val	Glu	Glu 175	Ala
Pro	Arg	Pro	Ala 180	Leu	Leu	Asn	Ser	Ala 185	Glu	Asn	Phe	Thr	Val 190	Leu	Ile
Lys	Asn 195	Asn	Ile	Asp	Phe	Pro	Gly 200	His	Asn	Tyr	Thr	Thr 205	Arg	Asn	Ile
Leu 210	Pro	Gly	Leu	Asn	Ile	Thr 215	Cys	Thr	Phe	His 220	Lys	Thr	Gln	Asn	Pro
Gln 225	Cys	Pro	Ile	Phe	Arg 230	Leu	Gly	Asp	Ile	Phe 235	Arg	Glu	Thr	Gly	Asp 240
Asn	Phe	Ser	Asp 245	Val	Ala	Ile	Gln	Gly	Gly 250	Ile	Met	Gly	Ile	Glu 255	Ile
Tyr	Trp	Asp	Cys 260	Asn	Leu	Asp	Arg	Trp 265	Phe	His	His	Cys	Arg 270	Pro	Lys
Tyr	Ser	Phe 275	Arg	Arg	Leu	Asp	Asp 280	Lys	Thr	Thr	Asn	Val 285	Ser	Leu	Tyr
Pro	Gly 290	Tyr	Asn	Phe	Arg	Tyr 295	Ala	Lys	Tyr	Tyr	Lys 300	Glu	Asn	Asn	Val
Glu 305	Lys	Arg	Thr	Leu	Ile 310	Lys	Val	Phe	Gly	Ile 315	Arg	Phe	Asp	Ile	Leu 320
Val	Phe	Gly	Thr 325	Gly	Lys	Phe	Asp	Ile 330	Ile	Gln	Leu	Val	Val 335	Tyr	
Ile	Gly	Ser	Thr 340	Leu	Ser	Tyr	Phe	Gly 345	Leu	Ala	Ala	Val 350	Phe	Ile	Asp
Phe	Leu	Ile 355	Asp	Thr	Tyr	Ser	Ser 360	Asn	Cys	Cys	Arg	Ser 365	His	Ile	Tyr
Pro	Trp 370	Cys	Lys	Cys	Cys	Gln 375	Pro	Cys	Val	Val	Asn 380	Glu	Tyr	Tyr	Tyr
Arg 385	Lys	Lys	Cys	Glu	Ser 390	Ile	Val	Glu	Pro	Lys 395	Pro	Thr	Leu	Lys	Tyr 400
Val	Ser	Phe	Val 405	Asp	Glu	Ser	His	Ile 410	Arg	Met	Val	Asn	Gln	Gln	Leu
Leu	Gly	Arg	Ser 420	Leu	Gln	Asp	Val	Lys 425	Gly	Gln	Glu	Val 430	Pro	Arg	Pro
Ala	Met	Asp 435	Phe	Thr	Asp	Leu	Ser 440	Arg	Leu	Pro	Leu	Ala 445	Leu	His	Asp
Thr	Pro 450	Pro	Ile	Pro	Gly 455	Gln	Pro	Glu	Glu	Ile 460	Gln	Leu	Leu	Arg	Lys

-continued

---

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
 465 470 475 480  
 Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
 485 490 495  
 Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
 500 505 510  
 Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
 515 520 525  
 Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
 530 535 540  
 Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
 545 550 555 560  
 Gln Asp Met Ala Asp Phe Ala Asn Leu Pro Ser Cys Cys Arg Trp Arg  
 565 570 575  
 Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
 580 585 590  
 Ser Pro Tyr  
 595

&lt;210&gt; SEQ ID NO 11

&lt;211&gt; LENGTH: 588

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 11

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
 1 5 10 15  
 Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
 20 25 30  
 Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
 35 40 45  
 Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
 50 55 60  
 Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
 65 70 75 80  
 Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
 85 90 95  
 Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
 100 105 110  
 Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
 115 120 125  
 Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
 130 135 140  
 Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
 145 150 155 160  
 Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
 165 170 175  
 Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile  
 180 185 190  
 Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205  
 Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro

-continued

210	215	220
Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp 225 230 235 240		
Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile 245 250 255		
Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys 260 265 270		
Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr 275 280 285		
Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val 290 295 300		
Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu 305 310 315 320		
Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr 325 330 335		
Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp 340 345 350		
Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr 355 360 365		
Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr 370 375 380		
Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr 385 390 395 400		
Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu 405 410 415		
Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro 420 425 430		
Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp 435 440 445		
Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys 450 455 460		
Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly 465 470 475 480		
Ser Cys Leu Pro Ser Gln Leu Glu Glu Leu Cys Cys Arg Lys Lys Pro 485 490 495		
Gly Ala Cys Ile Thr Thr Ser Glu Leu Phe Arg Lys Leu Val Leu Ser 500 505 510		
Arg His Val Leu Gln Phe Leu Leu Leu Tyr Gln Glu Pro Leu Leu Ala 515 520 525		
Leu Asp Val Asp Ser Thr Asn Ser Arg Leu Arg His Cys Ala Tyr Arg 530 535 540		
Cys Tyr Ala Thr Trp Arg Phe Gly Ser Gln Asp Met Ala Asp Phe Ala 545 550 555 560		
Ile Leu Pro Ser Cys Cys Arg Trp Arg Ile Arg Lys Glu Phe Pro Lys 565 570 575		
Ser Glu Gly Gln Tyr Ser Gly Phe Lys Ser Pro Tyr 580 585		

&lt;210&gt; SEQ ID NO 12

&lt;211&gt; LENGTH: 595

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

-continued

---

<400> SEQUENCE: 12

```

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
1          5          10          15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
          20          25          30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
          35          40          45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
50          55          60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
65          70          75          80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
          85          90          95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
100          105          110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
115          120          125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
130          135          140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
145          150          155          160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
165          170          175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile
180          185          190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
195          200          205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
210          215          220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
225          230          235          240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
245          250          255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys
260          265          270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
275          280          285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
290          295          300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
305          310          315          320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
325          330          335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
340          345          350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
355          360          365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
370          375          380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr

```

-continued

385	390	395	400
Val Ser Phe Val Asp	Glu Ser His Ile Arg	Met Val Asn Gln Gln Leu	
	405	410	415
Leu Gly Arg Ser Leu	Gln Asp Val Lys Gly	Gln Glu Val Pro Arg Pro	
	420	425	430
Ala Met Asp Phe Thr Asp	Leu Ser Arg Leu Pro	Leu Ala Leu His Asp	
	435	440	445
Thr Pro Pro Ile Pro Gly	Gln Pro Glu Glu Ile	Gln Leu Leu Arg Lys	
	450	455	460
Glu Ala Thr Pro Arg Ser	Arg Asp Ser Pro Val	Trp Cys Gln Cys Gly	
	465	470	475
Ser Cys Leu Pro Ser Gln	Leu Pro Glu Ser His	Arg Cys Leu Glu Glu	
	485	490	495
Leu Cys Cys Arg Lys Lys	Pro Gly Ala Cys Ile	Thr Thr Ser Glu Leu	
	500	505	510
Phe Arg Lys Leu Val Leu	Ser Arg His Val Leu	Gln Phe Leu Leu Leu	
	515	520	525
Tyr Gln Glu Pro Leu Leu	Ala Leu Asp Val Asp	Ser Thr Asn Ser Arg	
	530	535	540
Leu Arg His Cys Ala Tyr	Arg Cys Tyr Ala Thr	Trp Arg Phe Gly Ser	
	545	550	555
Gln Asp Met Ala Asp Phe	Ala Ile Leu Pro Ser	Cys Cys Arg Trp Arg	
	565	570	575
Ile Gln Lys Glu Phe Pro	Lys Ser Glu Gly Gln	Tyr Ser Gly Phe Lys	
	580	585	590
Ser Pro Tyr			
	595		

<210> SEQ ID NO 13  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

gtaacctcac tgagaacaga

20

<210> SEQ ID NO 14  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

tttggttaggc gaaggcgggc

20

<210> SEQ ID NO 15  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

ccttgtaggc gctttcgtgc t

21

<210> SEQ ID NO 16  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

---

-continued

---

&lt;400&gt; SEQUENCE: 16

ttgtgtatcc gtccatcagc t

21

&lt;210&gt; SEQ ID NO 17

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 17

actgccatat gtactattcc

20

&lt;210&gt; SEQ ID NO 18

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 18

aaatctaccc agccacttag

20

&lt;210&gt; SEQ ID NO 19

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 19

tgcagtggcc cacagcacag

20

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 20

catcctccaa tgctctgcatc

20

&lt;210&gt; SEQ ID NO 21

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 21

tttcccatgt ttgccattta

20

&lt;210&gt; SEQ ID NO 22

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 22

caagagcagt ggttgtgtcc

20

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 23

gagcccacag atatctactg

20

---

-continued

---

<210> SEQ ID NO 24  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

tgccctgcctc agtctctggt 20

<210> SEQ ID NO 25  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

tctctcgccc aggttgagtt 20

<210> SEQ ID NO 26  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

acgtcctctc cgcagttctt 20

<210> SEQ ID NO 27  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

attcagaccg aaagggtgtg 20

<210> SEQ ID NO 28  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

tgccctggtgt cccatcgagg 20

<210> SEQ ID NO 29  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

gttgtaacac tcctgtacca 20

<210> SEQ ID NO 30  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

aacagtgcc aaaaacttcac 20

<210> SEQ ID NO 31  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31



---

-continued

---

tcactgtgcc catcaagaac 20

<210> SEQ ID NO 32  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32

tgctgcgtgc ttctggctca 20

<210> SEQ ID NO 33  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33

ctggagaacc cttttccaag 20

<210> SEQ ID NO 34  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 34

caggaggcag ggatcatttg 20

<210> SEQ ID NO 35  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 35

tccaccgct acgctaaggt 20

<210> SEQ ID NO 36  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 36

tgcgctaagg actttaccta 20

<210> SEQ ID NO 37  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 37

tttacctacc ctacctcgtc 20

<210> SEQ ID NO 38  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 38

ccatcactgc tgtcccaaat 20

<210> SEQ ID NO 39  
<211> LENGTH: 20

---

-continued

---

<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 39

tcgccttgat gacaagacca 20

<210> SEQ ID NO 40  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 40

gtaaactctt ccactctgtt 20

<210> SEQ ID NO 41  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

accaagacac ggagagattc 20

<210> SEQ ID NO 42  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 42

taatattaaa tgtaacttta 20

<210> SEQ ID NO 43  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 43

aatatataaac ataactttat 20

<210> SEQ ID NO 44  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 44

ccatctcaac tggaggagct 20

<210> SEQ ID NO 45  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 45

gactttgccca tcctgcccag 20

<210> SEQ ID NO 46  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 46

tggaggatcc agaaagagtt 20

---

-continued

---

<210> SEQ ID NO 47  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 47

aatcccagca ctttgggagg

20

<210> SEQ ID NO 48  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 48

aaactgtccc aaaaaaaaaa

20

<210> SEQ ID NO 49  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 49

aactgtcccc caaaaaaaaaa

20

<210> SEQ ID NO 50  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 50

ctgtccccac aaaaaaaaaa

20

<210> SEQ ID NO 51  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 51

caaggcgggt ggatcactta

20

<210> SEQ ID NO 52  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 52

cgtaggactt ggcgcttct

19

<210> SEQ ID NO 53  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 53

gagcagctct cagattcgaa a

21

<210> SEQ ID NO 54  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

---

-continued

---

&lt;400&gt; SEQUENCE: 54

ccatgaggca ggtatgacta ttc 23

&lt;210&gt; SEQ ID NO 55

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 55

ctcctggatc tcacccagtt 20

&lt;210&gt; SEQ ID NO 56

&lt;211&gt; LENGTH: 23

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 56

ctcgtccagc ttgatatta agc 23

&lt;210&gt; SEQ ID NO 57

&lt;211&gt; LENGTH: 22

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 57

ggtccctagt gctagaacca ga 22

&lt;210&gt; SEQ ID NO 58

&lt;211&gt; LENGTH: 18

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 58

attcatccgt cagtggcc 18

&lt;210&gt; SEQ ID NO 59

&lt;211&gt; LENGTH: 22

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 59

gccatgtgaa ttttctaccg at 22

&lt;210&gt; SEQ ID NO 60

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 60

ttcgttgtgg ttaggatggg 20

&lt;210&gt; SEQ ID NO 61

&lt;211&gt; LENGTH: 22

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 61

caaggatgct cagggtagta gc 22

---

-continued

---

<210> SEQ ID NO 62  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 62

cactaggttt gctgtatcca tttct 25

<210> SEQ ID NO 63  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 63

gcaactgtgt gagagcttgg 20

<210> SEQ ID NO 64  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 64

tcaaccctgg tccagtgtg 19

<210> SEQ ID NO 65  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 65

caccaagtag ctctcactca taagg 25

<210> SEQ ID NO 66  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 66

caataacact tgtgcgagtt aggt 24

<210> SEQ ID NO 67  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 67

catcttggtg ccttggaac c 21

<210> SEQ ID NO 68  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 68

gtgagtggta atcctgctac tgc 23

<210> SEQ ID NO 69  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 69

---

-continued

---

aggcccactc ctgtactcg 19

<210> SEQ ID NO 70  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 70

ccaagtcaca gcatgaggc 19

<210> SEQ ID NO 71  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 71

acccagcgac gtatccac 18

<210> SEQ ID NO 72  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 72

aagcatgggg ttccatttc 19

<210> SEQ ID NO 73  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 73

gcataaaagg gactcctgct agta 24

<210> SEQ ID NO 74  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 74

gcttacagaa cacatgcatg g 21

<210> SEQ ID NO 75  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 75

gcacctgtag gcacagtgc 19

<210> SEQ ID NO 76  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 76

atcaccacct cagagctggt c 21

<210> SEQ ID NO 77  
<211> LENGTH: 21

---

-continued

---

<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 77

gttaacatgg ctactgcagc c 21

<210> SEQ ID NO 78  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 78

gcttagaaaag gaggcgactc c 21

<210> SEQ ID NO 79  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 79

ttgtgacatt tgcaaggctg cc 22

<210> SEQ ID NO 80  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 80

tctgaagctc tgctcctgag 20

<210> SEQ ID NO 81  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 81

ctcaccttct ggcttccagt 20

<210> SEQ ID NO 82  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 82

cttaccactc ccaggactaa 20

<210> SEQ ID NO 83  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 83

gtctgcctgt tcactgccat 20

<210> SEQ ID NO 84  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 84

cagagacctt cagaaacttc g 21

---

-continued

---

<210> SEQ ID NO 85  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 85

agatcaccag ggacacagtg

20

<210> SEQ ID NO 86  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 86

ctcaactcca ctttcctcgg

20

<210> SEQ ID NO 87  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 87

cctttcactt ttttggtctc atg

23

<210> SEQ ID NO 88  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 88

gggagaattc tgaaaatgcc c

21

<210> SEQ ID NO 89  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 89

ggaccagagc tctactcttc

20

<210> SEQ ID NO 90  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 90

aggtcataga tcgacctgcc

20

<210> SEQ ID NO 91  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 91

aagaagcgcc aagtcctacg

20

<210> SEQ ID NO 92  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens



---

-continued

---

&lt;400&gt; SEQUENCE: 92

gcaatccaga ctgaagttga c

21

&lt;210&gt; SEQ ID NO 93

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 93

actctggtct gcagttggtg

20

&lt;210&gt; SEQ ID NO 94

&lt;211&gt; LENGTH: 24

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 94

cctttaaaat cagagacctt caga

24

&lt;210&gt; SEQ ID NO 95

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 95

gccccatcctc tgaacacccat

20

&lt;210&gt; SEQ ID NO 96

&lt;211&gt; LENGTH: 21

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 96

cccttggaac tcttgctatc g

21

&lt;210&gt; SEQ ID NO 97

&lt;211&gt; LENGTH: 21

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 97

ggcagtacag tggcttcaag a

21

&lt;210&gt; SEQ ID NO 98

&lt;211&gt; LENGTH: 21

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 98

gtgggacagt ttgctgtgcc t

21

&lt;210&gt; SEQ ID NO 99

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 99

gagtccttac caatagcagg

20

---

-continued

---

<210> SEQ ID NO 100  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 100

gtcaaagaat ttgtggccac c 21

<210> SEQ ID NO 101  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 101

catgaactgt cttttaatgt gtaaag 26

<210> SEQ ID NO 102  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 102

gagatacggg ttcacccatgt tg 22

<210> SEQ ID NO 103  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 103

aattagctgg gcatggtgcg 20

<210> SEQ ID NO 104  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 104

ttgagatgga gtctcgctct g 21

<210> SEQ ID NO 105  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 105

cactgtccac gtgactgctt 20

<210> SEQ ID NO 106  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 106

tcctacttcg gtctggtaag agatt 25

<210> SEQ ID NO 107  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 107

-continued

---

```

gggcctaatt ttcgtgcat                                19

<210> SEQ ID NO 108
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 108
aagaacctag aacctgaggg ctt                            23

<210> SEQ ID NO 109
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 109
ttgagatggg aggcagctt                                19

<210> SEQ ID NO 110
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 110
ttcggtctcc aggacat                                  17

<210> SEQ ID NO 111
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 111
cacagagctt tgcaggtgaa                                20

```

---

1. An isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of:

(a) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 5'UTR region corresponding to positions 362, 532, 1100, 1122, 1171 or 1702 of the genomic sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;

(b) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of the ATP-gated ion channel P2X7R, wherein in the exon as indicated in column "Exon" of the following Table A the amino acid residue as indicated in column "Amino acid residue" of Table A corresponding to the position as indicated in column "Position in wild-type" of Table A of the wild-type ATP-gated ion channel P2X7R amino acid sequence as depicted in SEQ ID NO: 3 or 4 is-replaced by another amino acid residue

TABLE A

Exon	Amino acid residue	Position in wild-type
exon 3	R (Arg)	117
exon 5	G (Gly)	150
exon 6	E (Glu)	186
exon 6	L (Leu)	191
exon 8	R (Arg)	270
exon 13	I (Ile)	568
exon 13	R (Arg)	578

(c) a nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in exon 5 or 8 corresponding to position 32548 or position 37633 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide

(d) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to

positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted;

- (e) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R, wherein in the intron as indicated in column "Intron" of the following Table B the nucleotide as indicated in column "Replaced nucleotide" of Table B corresponding to the position as indicated in column "Position in wild-type" of Table B of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1 is replaced by another nucleotide

TABLE B

Intron	REPLACED NUCLEOTIDE	Position in wild-type
intron 1	G	3166
intron 1	C	24778
intron 1	C	24830
intron 3	A	26308
intron 3	G	26422
intron 4	G	32394
intron 4	T	32434
intron 5	A	32783
intron 6	G	35641
intron 6	A	35725
intron 6	T	36001
intron 7	G	36378
intron 7	T	36387
intron 7	G	36398
intron 9	C	47214
intron 11	T	47563
intron 12	C	54307
intron 12	G	54308

- (f) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 3'UTR region corresponding to position 54925, 55169, 55170, 55171 or 55917 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;
- (g) a nucleotide sequence comprising at least 20 or 21 nucleotides and comprising the mutations or deletions as defined in any one of (a) to (f);
- (h) a nucleic acid sequence comprising a nucleotide sequence as shown in any one of SEQ ID NOS: 13 to 51;
- (i) a nucleic acid sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NOS: 5 to 12;
- (j) a nucleotide sequence which hybridizes to a nucleotide sequence defined in any one of (a) to (g) or to the nucleotide sequence of (h) and having a mutation as defined in any one of (a) to (f); and
- (k) a nucleic acid sequence being degenerate as a result of the genetic code to the nucleic acid sequence as defined in (j).

2. The nucleic acid molecule of claim 1 derived from mouse, rat or human.

3. The nucleic acid molecule of claim 1 which is DNA, RNA, PNA or phosphorothioates.

4. A vector comprising the nucleic acid molecule of claim 1.

5. The vector of claim 4 which is an expression vector, a gene targeting vector and/or a gene transfer vector.

6. A host transformed with the nucleic acid molecule of claim 1 or a vector comprising said nucleic acid molecule.

7. The host of claim 6 which is a mammalian cell, an amphibian cell, a fish, an insect cell, a fungal cell, a plant cell or a bacterial cell.

8. The host of claim 7, wherein said mammalian cell is selected from the group consisting of CHO cells, HEK293 cells, COS-7 cells or PC12 cells.

9. The host of claim 7, wherein said amphibian cell is an oocyte, preferably a *Xenopus* oocyte.

10. The host of claim 9, wherein said oocyte is a frog oocyte.

11. The host of claim 6 which is a non-human transgenic organism.

12. The host of claim 11, wherein said non-human organism is a mammal, amphibian, a fish, an insect, a fungus or a plant.

13. A method for producing the polypeptide encoded by a nucleic acid molecule of claim 1(b) or 1(d) comprising culturing/raising a host transformed with the nucleic acid molecule of claim 1 or a vector comprising said nucleic acid molecule and isolating the produced polypeptide.

14. A polypeptide encoded by the nucleic acid molecule of claim 1(b) or 1(d) or produced by a method comprising culturing/raising a host transformed with the nucleic acid molecule of claim 1 or a vector comprising said nucleic acid molecule.

15. An antibody specifically directed to the polypeptide of claim 14, wherein said antibody specifically reacts with an epitope generated and/or formed by the mutation in the ATP-gated ion channel P2X7R selected from the group consisting of:

- (i) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein the R (Arg), G (Gly), E (Glu), L (Leu), R (Arg), I (Ile) or R (Arg) residue corresponding to position 117, 150, 186, 191, 270, 568 or 578 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 is replaced by another amino acid residue; and
- (ii) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted.

16. The antibody of claim 15 which is a monoclonal antibody.

17. An aptamer specifically binding to a nucleic acid molecule of claim 1 or to the polypeptide encoded by the nucleic acid molecule of claim 1.

18. A primer or pair of primers capable of specifically amplifying a nucleic acid molecule as defined in claim 1.

19. The primer or pair of primers of claim 18, which is selected from the group consisting of SEQ ID NOS: 52 to 111.

20. A composition comprising the nucleic acid molecule of claim 1, a vector comprising said nucleic acid molecule, a polypeptide encoded by said nucleic acid molecule, an

antibody specifically directed to said polypeptide, an aptamer specifically binding to said nucleic acid molecule and/or a primer or pair of primers capable of specifically amplifying said nucleic acid molecule.

21. The composition of claim 20 which is a diagnostic composition.

22. The composition of claim 21, optionally further comprising suitable means for detection.

23. (canceled)

24. A method of diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2X7R protein expressed in the cells of said individual is non-functional, shows an altered ATP-gating in comparison to the wild-type P2X7R protein or is over- or under-expressed in comparison to the P2X7R protein level an unaffected individual.

25. A method for diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2X7R gene sequence or encoded protein thereof comprises a mutation in comparison to the wild-type P2X7R sequence.

26. A method for diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2X7R gene sequence or encoded protein thereof comprises a mutation in comparison to the wild-type P2X7R sequence, wherein said mutation is a mutation as defined in claim 1 and/or a nucleotide replacement or deletion selected from the following Table C indicating in column "Region of P2X7R" the region of the P2X7R genomic nucleotide sequence in which the replacement or deletion occurs, in column "Nucleotide" of Table C the nucleotide which is replaced by another nucleotide or the nucleotides which are deleted and in column "Position in wild-type" of Table C the corresponding position in the nucleotide sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1

TABLE C

Region of P2X7R	NUCLEOTIDE	Position in wild-type
5'UTR	T	362
5'UTR	T	532
5'UTR	A	1100
5'UTR	A	1122
5'UTR	C	1171
5'UTR	T	1351
5'UTR	G	1702
5'UTR	T	1731
5'UTR	C	1860
5'UTR	C	2162
5'UTR	C	2238
5'UTR	A	2373
5'UTR	G	2569
5'UTR	G	2702
intron 1	G	3166
intron 1	C	24778
intron 1	C	24830
exon 2	T	24942
exon 3	C	26188
exon 3	A	26308
exon 3	G	26422
intron 4	G	32394
intron 4	T	32434
exon 5	G	32493

TABLE C-continued

Region of P2X7R	NUCLEOTIDE	Position in wild-type
exon 5	G	32506
exon 5	C	32507
exon 5	C	32548
intron 5	A	32783
intron 5	T	35309
intron 5	C	35374
intron 5	A	35378
exon 6	G	35438
exon 6	T	35454
intron 6	T	35549
intron 6	G	35641
intron 6	A	35725
intron 6	T	36001
intron 6	A	36064
intron 6	deletion of GTTT	36091 to 36094
intron 6	C	36108
intron 7	C	36374
intron 7	G	36378
intron 7	T	36387
intron 7	G	36398
intron 7	C	37439
intron 7	T	37513
exon 8	C	37604
exon 8	G	37605
exon 8	G	37623
exon 8	C	37633
intron 9	C	47214
exon 11	G	47383
exon 11	C	47411
intron 11	T	47563
intron 12	C	54307
intron 12	G	54308
exon 13	C	54399
exon 13	A	54480
exon 13	C	54523
exon 13	deletion of CCCTGAGAGCCACAGGTGCCT	54562 to 54582
exon 13	A	54588
exon 13	C	54664
exon 13	G	54703
exon 13	A	54804
exon 13	G	54834
exon 13	G	54847
3'UTR	G	54925
3'UTR	C	55169
3'UTR	A	55170
3'UTR	A	55171
3'UTR	C	55917

27. The method of claim 26, wherein the occurrence of the mutation in the ATP-gated ion channel P2X7R gene is determined by PCR or immunological methods.

28. The composition of claim 20 which is a pharmaceutical composition.

29. The pharmaceutical composition of claim 28, optionally further comprising a pharmaceutically acceptable carrier.

30. A method of treating an affective disorder comprising administering a therapeutically effective amount of the nucleic acid molecule as defined in claim 1 or a therapeutically effective amount of a polypeptide encoded by said nucleic acid molecule to a subject suffering from said disorder.

31. (canceled)

32. A pharmaceutical composition comprising a nucleic acid molecule comprising a nucleotide sequence which encodes a functional ATP-gated ion channel P2X7R and which is selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide comprising the amino acid sequence as depicted in SEQ ID NO: 3 or 4;
- (b) a nucleotide sequence comprising the nucleotide sequence as depicted in SEQ ID NO: 1 or 2;
- (c) a nucleotide sequence which hybridizes to the nucleotide sequence of (a) or (b); and
- (d) a nucleotide sequence which is degenerated as a result of the genetic code to the nucleotide sequence of (c).

33. A pharmaceutical composition comprising a compound the administration of which to cells leads to an increase of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells or comprising a nucleic acid molecule the expression of which in cells or the administration of which to cells leads to an increase of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells.

34. A method of treating an affective disorder comprising administering a therapeutically effective amount of the pharmaceutical composition as defined in claim 32, a pharmaceutical composition comprising a compound the administration of which to cells leads to an increase of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells, a pharmaceutical composition comprising a nucleic acid molecule the expression of which in cells or the administration of which to cells leads to an increase of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells or a pharmaceutical composition comprising a polypeptide encoded by said nucleic acid molecule to a subject suffering from said disorder.

35. (canceled)

36. A method of treating an affective disorder comprising administering a therapeutically effective amount of a pharmaceutical composition comprising a modulator of P2X7R activity to a subject suffering from said affective disorder.

37. The method of claim 36, wherein said modulator is an agonist.

38. The method of claim 37, wherein said agonist is selected from the group consisting of ATP, ATP-4 and BzATP (2'-3'-O-(4-Benzoylbenzoyl)adenosine 5'-triphosphate ( $C_{24}H_{24}N_5O_{15}P_3$ )).

39. The method of claim 37, wherein said agonist is tenidap ( $C_{15}H_{11}ClN_2O_2S$ ) or a derivative thereof or 3-substituted-2-oxindole-1-carboxamides.

40. The method of claim 36 wherein said pharmaceutical composition optionally further comprises a  $\beta$ -adrenergic receptor modulator.

41. The method of claim 40, wherein said  $\beta$ -adrenergic receptor modulator is a  $\beta$ -adrenergic receptor antagonist selected from the group consisting of DL-propanolol, D-propanolol and labetolol.

42. The method of claim 36, wherein said affective disorder is selected from the group consisting of major depression, generalized anxiety disorder and bipolar disorder.

43. The method of claim 42, wherein said major depression is selected from the group consisting of major depression, dysthymia, atypical depression, premenstrual dysphoric disorder and seasonal affective disorder.

44. The method of claim 42, wherein said generalized anxiety disorder is selected from the group consisting of panic disorder, phobias, agoraphobia, social phobia, specific

phobia, obsessive-compulsive disorder, post-traumatic stress disorder, separation anxiety disorder, mania, hypomania and cyclothymic disorder.

45. The method of claim 42, wherein said bipolar disorder is bipolar disorder type I or bipolar disorder type II.

46. A kit comprising the nucleic acid molecule of claim 1, a vector comprising said nucleic acid molecule, a host comprising said nucleic acid molecule, a polypeptide encoded by said nucleic acid molecule, an antibody specifically directed to said polypeptide, an aptamer specifically binding to said nucleic acid molecule and/or a primer or pair of primers capable of specifically amplifying said nucleic acid molecule.

47. A method for identifying compounds which are capable of specifically interacting with the polypeptide of claim 14, comprising the steps of

contacting a polypeptide of claim 14 with a compound or a candidate mixture of compounds to be tested; and

determining whether said compound or a candidate mixture of compounds is capable of specifically interacting with said polypeptide.

48. A method for the characterization of compounds which are capable of altering characteristics of the polypeptide of claim 14, comprising the steps of

contacting a polypeptide of claim 14 with a compound or a candidate mixture; and

determining whether the compound or a candidate mixture alters a characteristic of the polypeptide of claim 14.

49. A method of screening for compounds which are capable of interacting with the polypeptide of claim 14, comprising the steps of

(a) contacting a polypeptide of claim 14 with a compound or a candidate mixture of compounds;

(b) measuring and/or detecting a response; and

(c) comparing said response to a standard response as measured in the absence of said candidate molecule.

50. A method for the production of a pharmaceutical composition comprising the steps of the method of claim 47 and comprising a further step, wherein a derivative of said identified, characterized and/or screened molecule is generated.

51. A method for the production of a pharmaceutical composition comprising the steps of the method of claim 47 and formulating the molecules identified, characterized, screened and/or derivatized in pharmaceutically acceptable form.

52. The method of claim 51, wherein the pharmaceutical composition to be produced further comprises neuroprotective substances, nootropic substances, brilliant blue, piperidine or piperazine derivatives thereof, adamantane derivatives, substituted phenyl compounds, oxidized ATP, 2-O-(4-benzoylbenzoyl)adenosine-5-triphosphate, 3-O-(4-benzoylbenzoyl)adenosine-5-triphosphate or  $\beta$ -adrenergic receptor modulators.

53. The method of claim 47, wherein said compound(s) or candidate mixture(s) of compounds comprise(s) antagonist(s), partial antagonist(s), partial agonist(s) and/or agonist(s) for an altered ATP-gated ion channel P2X7R.

54. A method for diagnosing an affective disorder of an individual comprising:

- (a) isolating DNA from cells obtained from an individual;
- (b) determining all or part of the nucleotide composition of the P2X7R gene; and
- (c) analyzing said nucleotide composition of P2X7R for the presence of one or more polymorphism(s) mutation or allelic variation.

**55.** A method for diagnosing an affective disorder of an individual comprising:

- (a) isolating RNA from cells obtained from an individual;
- (b) converting said RNA into cDNA;
- (c) determining all or part of the nucleotide composition of the P2X7R gene; and

- (d) analyzing said nucleotide composition of P2X7R for the presence of one or more polymorphism(s), mutation or allelic variation.

**56.** A method for diagnosing an affective disorder of an individual comprising:

- (a) isolating RNA or protein from cells obtained from an individual;
- (b) determining the levels of P2X7R RNA or protein; and
- (c) comparing the levels of P2X7R RNA or protein with the corresponding levels from a normal individual not afflicted with an affective disorder.

\* \* \* \* \*