



- (51) **International Patent Classification:**
A01N 65/00 (2009.01)
- (21) **International Application Number:**
PCT/US2013/033393
- (22) **International Filing Date:**
22 March 2013 (22.03.2013)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
13/443,572 10 April 2012 (10.04.2012) US
- (72) **Inventors; and**
- (71) **Applicants :** VAN ALLER, Robert, Thomas [US/US]; 3004 Raphael Drive, Hattiesburg, MS 39402 (US). VAN ALLER, Robert, Merrick [US/US]; 4004 Samoa Road, Pearlinton, MS 39572 (US). VAN ALLER, Geoffrey, Thomas [US/US]; 3912 Cabildo Place, Ocean Springs, MS 39564 (US).
- (74) **Agent:** KEATY, Thomas, S.; Keaty Law Firm, LLC, 365 Canal Street, Suite 2410, New Orleans, LA 70130 (US).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with amended claims (Art. 19(1))



(54) **Title:** HERBAL COMPOSITION AND A METHOD OF MAKING THEREOF

(57) **Abstract:** A topical composition is prepared using macerated leaves of St. Andrew's Cross (*hypericum hypericoides*). An aqueous extract of the leaves is mixed with ethanol to form a spray or to saturate a wet wipe and administer to a mammal skin for treatment a variety of skin conditions, including eczema, psoriasis, rosacea, cellulitis, contact dermatitis, seborrheic dermatitis, nummular dermatitis, dandruff, stasis dermatitis, perioral dermatitis, dermatitis herpetiformis, ecthyma, impetigo, shingles, urticaria, tinea pedis, tinea manuum, tinea cruris, sunburn, insect bites and stings, jellyfish stings, granuloma annulare, bullous pemphigoid, lichen planus, bed sore, inflammations caused by waterborne vectors, diaper rash, heat rash, pityriasis rosea, jungle rot, scleroderma, skin rashes and photo-aging.

HERBAL COMPOSITION AND A METHOD OF MAKING THEREOF

BACKGROUND OF THE INVENTION

[0001] This invention relates to an herbal composition containing *Hypericum hypericoides* (St. Andrew's Cross) and a method of making thereof. More particularly, the present invention relates to an herbal composition designed for treatment of skin conditions.

[0002] The skin is the largest organ of the human body. Skin infections can be either bacterial, viral or fungal in nature. A significant portion of Americans as well as the global population are afflicted with skin conditions including but not limited to eczema, psoriasis, rosacea, cellulitis, contact dermatitis (both irritant and allergic) including poison ivy, poison oak and poison sumac from contact with urushiol (organic allergen), seborrhoeic dermatitis, nummular dermatitis, dandruff, stasis dermatitis, perioral dermatitis, dermatitis herpetiformis, ecthyma, impetigo, shingles, urticaria, tinea pedis (athlete's foot), Tinea Manuum, tinea cruris, sunburn, inflammation from insect bites and stings, inflammation from jellyfish stings, granuloma annulare, bullous pemphigoid, lichen planus, bed sore, inflammation from puncture wounds from catfish and other waterborne creatures, diaper rash, heat rash,

pityriasis rosea, jungle rot (trench foot) and scleroderma. Skin rashes, which often go misdiagnosed or undiagnosed, are very common and fall into at least one of the above-mentioned categories of conditions.

[0003] Some of these cases can be severe and can result in a rash, itching, bleeding sores, hair loss, permanent scarring and subsequent breakdown of one's immune system and internal organs. Particular types of systemic cases can often cause people to become isolated from the general population due to embarrassment and shame and can have profound psychological and physiological effects on their entire lives. Death can occur as a result of some of these skin problems and is well documented in medical literature all over the world.

[0004] Systemic steroids, also called corticosteroids, glucocorticoids or cortisones, are commonly prescribed for the treatment of the above-mentioned skin conditions. These medications are administered orally or by injections. They are synthetic derivatives of the natural steroid, Cortisol, which is produced by the adrenal glands. Topical corticosteroids are also used to treat the skin conditions; they are applied directly to the skin. Inhaled steroids are breathed in.

[0005] High dose steroids can trigger depression, anxiety, panic, mood swings, fluid retention, increased blood pressure, sleep disturbances, and psychiatric side effects. Weight gain, redistribution of body fat, skin and hair changes, eye changes, musculoskeletal, gastrointestinal, cardiovascular, and immune changes are among other possible side effects.

[0006] High doses of systemic corticosteroids such as prednisone are sometimes prescribed for inflammatory diseases, which manifest themselves in the skin. These high doses can cause a condition known as Cushing Syndrome. Symptoms of Cushing Syndrome include easy bruising and purple striae (stretch

marks) of the skin over the abdomen, buttocks and thighs, telangiectatic cheeks (broken capillaries), fragile skin and poor wound healing, acne and excessive hair growth in women on their faces, necks, chests, abdomens and thighs. Women may also show clitoral hypertrophy and male-pattern baldness.

[0007] Doctors often prescribe topical steroids to patients with the above-mentioned skin conditions. While topical steroids are effective in reducing inflammation, they also have significant side effects. Most of these side effects are seen with long-term use, but some may be noticed within days of starting therapy.

[0008] Tachyphylaxis is the tolerance the skin develops to the vasoconstrictive action of topical steroids. After repeated use of topical steroids, the capillaries in the skin do not constrict as well, requiring higher doses or more frequent application of the steroid. With high-potency steroids this effect has been documented after four days of applying a topical steroid three times a day. The ability of the blood vessels to constrict returns about four days after stopping therapy.

[0009] Steroid rosacea is a side effect commonly observed in fair-skinned people who already have rosacea. A typical example occurs when a person uses a very mild steroid on the face to counteract the facial flushing. This gives pleasing results, but tolerance develops, causing the person to use a higher strength steroid. At this point any attempt to cut down on the steroid application or stop altogether causes intense facial redness and pustules.

[0010] Repeated use of topical steroids in the same area can cause thinning of the epidermis and changes in the connective tissue of the dermis or skin atrophy. The skin becomes lax, wrinkled, and shiny. Affected areas can be depressed below the level of normal skin with visible telangiectasias, hypopigmentation, and prominence

of underlying veins. In most cases the atrophy is reversible once topical steroid use is stopped, but it may take months for the skin to "thicken" again.

[0011] Repeated use of topical steroids in areas where skin touches skin such as the groin and armpits can result in striae, or stretch marks. Stretch marks from topical steroids are permanent and irreversible. These stretch marks can be very itchy and may require a lower potency steroid to relieve the itching. It is recommended to progressively decrease the steroid potency until topical steroids therapy in these areas can be terminated.

[0012] Because topical steroids change the way the immune system functions, they can inhibit the skin's ability to fight off bacterial or fungal infections. A typical example of this is seen when someone applies a topical steroid to an itchy groin rash. If this is a fungal infection, the rash gets redder, itchier, and spreads more extensively than a typical fungal infection. The resulting rash is a bizarre pattern of widespread inflammation with pustules called tinea incognito.

[0013] Some people may be allergic to a component of the topical steroid base, or vehicle. Patch-testing of a group of patients with dermatitis revealed that about 4-5% of them were allergic to topical steroids. People who have chronic skin conditions and use multiple prescription or over-the-counter topical steroids are at higher risk of developing allergies to topical steroids.

[0014] Glaucoma is a disease in which the pressure inside the eye increases to the point of damaging the optic nerve. There are isolated cases reports of people developing glaucoma after long-term use of topical steroids around the eyes. How topical steroids applied around the eyes cause glaucoma is not completely understood, but it is believed that enough of the steroid can be absorbed and get into the eye.

[0015] Topical immunomodulators (TIMs) including brand name products Protopic (Tacrolimus) and Elidel (Pimecrolimus) work by altering the body's immune response to allergens, preventing flare ups of eczema and other skin rashes. In 2005, the FDA warned doctors to prescribe Elidel and Protopic with caution due to concerns over a possible cancer risk associated with their use. The two creams carry the FDA's strongest "black box" warning on their packaging to alert doctors and patients to these potential risks. The warning advises doctors to prescribe short-term use of Elidel and Protopic only after other available eczema treatments have failed in adults and children over the age of two.

[0016] Biologies such as Enbrel (Etanercept) and Humira (adalimumab) can be prescribed for severe skin conditions and their prolonged use can lead to pneumonia, cellulitis, septic arthritis, bronchitis, gastroenteritis, pyelonephritis, sepsis, abscess and osteomyelitis, tuberculosis, cancer, congestive heart failure, hepatitis, and a number of other fungal, viral and bacterial infections.

[0017] The cause of most skin disorders is unknown. According to some statistics, one percent of the U. S. population dies every year from the overuse of medication or mistreatment of skin disorders using conventional medicine.

[0018] There exists therefore a need for a safer treatment of skin disorders, particularly a treatment that avoids the use of harsh medications and the associated serious side effects.

SUMMARY OF THE INVENTION

[0019] It is, therefore, an object of the present invention to provide a natural composition that can relieve a variety of skin conditions.

[0020] It is another object of the invention to provide a plant-based composition that can relieve serious skin disorders without introducing life-endangering side effects.

[0021] It is a further object of the present invention to provide an herbal composition that can be administered topically or aurally to a person suffering from skin disorders.

[0022] It is still a further object of the invention to provide a method of making an herbal composition that can be used for treatment of a variety of skin conditions.

[0023] It is still a further object of the invention to provide a method of administering an herbal composition in a form of a spray or wet wipes for treatment of skin conditions.

[0024] These and other objects of the invention are achieved through a provision of an herbal topical composition, which is prepared using macerated leaves of St. Andrew's Cross (*hypericum hypericoides*). An aqueous extract of the leaves is mixed with a hydroxyl to form a spray or to saturate a wet wipe and topically administer to a mammal skin for treatment a variety of skin conditions. The skin conditions include eczema, psoriasis, rosacea, cellulitis, contact dermatitis, seborrhoeic dermatitis, nummular dermatitis, dandruff, stasis dermatitis, perioral dermatitis, dermatitis herpetiformis, ecthyma, impetigo, shingles, urticaria, tinea pedis, tinea manuum, tinea cruris, sunburn, insect bites and stings, jellyfish stings, granuloma annulare, bullous pemphigoid, lichen planus, bed sore, inflammations caused by waterborne vectors, diaper rash, heat rash, pityriasis rosea, jungle rot, scleroderma, skin rashes and photo-aging.

[0025] The aqueous extract is prepared using boiling distilled water, to which the plant material is added, mixture stirred to increase contact with water and then vacuum filtered. The filtered material is then mixed with alcohol and used as a topical agent to treat a variety of skin conditions.

[0026] In the preferred embodiment, ethanol is used as a preservative of the extract. Optionally, glycerin can be added.

DETAIL DESCRIPTION OF THE INVENTION

[0027] The herbal composition of the present invention is completely natural and does not contain steroids or heavy metals (mercury, lead, etc.), which can be found in conventional creams and ointments used for the treatment of skin disorders. Unlike synthetic drugs and invasive method of treatment, the application of the composition according to the instant invention has no side effects such as irritation of the digestive tract and skin, allergic reactions and pain caused by medications, and unlike other herbal preparations, has a wider range and stronger action.

[0028] An aqueous solution of *Hypericum hypericoides*, commonly known as St. Andrew's Cross, has been found to have a remarkably positive effect on the above-mentioned skin conditions, including eczema, psoriasis, rosacea, cellulitis, contact dermatitis (both irritant and allergic) including poison ivy, poison oak and poison sumac from contact with urushiol (organic allergen), seborrheic dermatitis, nummular dermatitis, dandruff, stasis dermatitis, perioral dermatitis, dermatitis herpetiformis, ecthyma, impetigo, shingles, urticaria, tinea pedis (athlete's foot), Tinea Manuum, tinea cruris, sunburn, inflammation from insect bites and stings, inflammation from jellyfish stings, granuloma annulare, bullous pemphigoid, lichen planus, bed sores, inflammation from puncture wounds from catfish and other

waterborne vectors, diaper rash, heat rash, pityriasis rosea, jungle rot (trench foot), scleroderma, skin rashes and photo-aging.

[0029] *Hypericum hypericoides* (L.) Crantz ssp. *Hypericoides* is a perennial flowering shrub native to Southern and Southeastern regions of the United States. It is classified in Dicot group, Clusiaceae family, genus *Hypericum*. It grows 1 - 3 feet in size, blooming from June to September. Its native habitat is dry woods, pine barrens, sand hills and ridges and floodplains.

[0030] Some evidence exists of the root of *Hypericum hypericoides* having been chewed as an antidote to rattlesnake bites. A tea made from the root has been used to relieve colic, fevers, pain and diarrhea. It has been applied externally to ulcerated breasts. A tea made from the leaves has been used for kidney and bladder ailments and skin problems. A milky substance from the plant has been rubbed on sores. This evidence is almost anecdotal. However, the tests conducted by the inventors support the claims that when topically applied, an aqueous extract of the leaves of *hypericum hypericoides* relieves a variety of skin conditions, as will be described in more details hereinafter.

[0031] According to the method of the present invention, preparation of the herbal composition consists of several phases:

[0032] 1. Maceration - maceration is the first step of preparation, where the leaves of St. Andrew's Cross (can be used fresh and dried) are chopped in small pieces. Plants can be chopped and ground in a blender, mixer and chopper grinder.

[0033] 2. Extraction - Extraction is the second step of preparation where the plant material (macerated leaves) is mixed with boiling distilled water at a rate of 1:25 to 1:30, preferably at a rate of 1:27. This step can be carried out in a glass or Pyrex vessel.

[0034] After mixing, the mixture is removed from heat and stirred with a non-metal stirrer every two minutes for about 12 - 18 minutes. Between the stirring cycles, the vessel is covered with a lid. Stirring increases the contact area of water with the plant material and speeds up the process of forming the final composition. In addition, the dissolution or breakdown of the ingredients that make the basic structure of the leaves creates a series of aromatic alcohols, aldehydes, pentoses, hexoses and tannins in the herbal composition.

[0035] 3. Filtering - Filtering is the third step of preparation, where the mixture is filtered using a fine vacuum filter. Suspended solids and water are removed at this stage.

[0036] 4. Preservation - The filtered extract is immediately preserved by mixing it with a stabilizing/preservative agent selected from the hydroxyl group, such as for instance pure alcohol in the form of ethanol or methanol. The stabilizing preserving agent is added at a ratio of between 4 - 6% by volume of the filtered composition, preferably at a ratio of 3% by volume. The composition of the aqueous extract of hypericum hypericoides and alcohol is refrigerated and kept in the refrigerated state at a temperature of about 3 - 5 degrees Celsius (37.4 - 41 degrees Fahrenheit).

[0037] Example: According to one example, an extract of hypericum hypericoides was made to isolate the active components of hypericum hypericoides in an aqueous solution. According to the method, about 6 liters (about 1.6 gallons) of distilled water are deposited in a heat-proof container and the water is brought to a boil. Then, about 225 grams (about 0.5 pounds) of macerated plant leaves were deposited in the container of boiling distilled water. The container is then removed from heat. The contents of the container were stirred with a non-metallic rod every

two minutes. A lid was placed over the container top to keep it covered while not stirring. The stirring cycle continues for about 16 minutes.

[0038] Next, the mixture was forced through a 10 micron vacuum filter to remove suspended solids and as much water from the plant material as possible. The resultant mixture was immediately preserved with 3% of ethyl alcohol by volume and refrigerated to 3.3 Celsius (38 degrees Fahrenheit).

[0039] It is noted that the active compounds in the extract are sensitive to the following and should not be contaminated at any point with any of these compounds: acids, surfactants, sodium benzoate, methyl paraben and amphoteric metals. The tests demonstrated that the extract preserved with ethyl alcohol and refrigerated has the durability over one year.

[0040] The aqueous extract contains two active compounds and demonstrates the following properties:

[0041] 1. The extract as tested had a concentration of active compounds of 0.311% - 0.362% by weight. The pH was from 6.30 to 6.62.

[0042] 2. Over time there is some slight settling of very fine particulate matter. This is harmless and can be filtered out if desired.

[0043] It was noted that when the plant is extracted in the presence of aluminum, severe precipitation and darkening occurs. As with other flavonoids care should be taken to prevent oxidation. It is preferred that the extract be preserved quickly and remained refrigerated when not used. The extract may be frozen if it is not to be used for an extended period of time. The air space in the storage container(s) should be minimized if not eliminated. Oxygen was noted to shorten the useful life of the product.

[0044] The tests showed that glycerin will prevent oxidation of the active compounds at room temperature but does not necessarily prevent bacterial and/or fungal growth. Ethanol acts as a preservative and stabilizer of the active compounds and prevents bacterial and fungal growth if the extraction is refrigerated and any oxygen is kept out of the container.

[0045] The following concentrations were also obtained from extractions with the percentages of solids by weight of the two active compounds in the extract:

[0046] 4.0 grams plant material/100 ml distilled water yielded 0.339% solids
by weight

[0047] 2.0 grams plant material/100 ml distilled water yielded 0.238% solids
by weight

[0048] 1.0 grams plant material/100 ml distilled water yielded 0.138% solids
by weight

[0049] 0.5 grams plant material/100 ml distilled water yielded 0.081% solids
by weight

[0050] The tests demonstrated that these different concentrations are effective for varying degrees of severity of the skin ailments described above. It should also be noted this may vary depending on the time of the year the plant is harvested (warmer months will yield higher concentrations given an equal amount of plant material in a given amount of distilled water).

The pH of the extract is slightly acidic due to the phenolic groups of the two active compounds.

[0051] Use of Extract as a Topical Agent

[0052] The product in the form of a spray has been tested in the following manner to treat the above-mentioned skin condition: it was sprayed on the affected

area(s) 3-4 times per day and allowed to air dry on the skin. The tests showed that a small amount may be poured in the hands and rubbed on the affected area(s). A wipe can be saturated with the herbal composition and used to wipe the affected area. Itching or burning in all tested cases ceased immediately or in less than one to two minutes. Rash in the form of bumps or redness, scabs, dryness, sores, etc. cleared up within two-three days.

[0053] One particular case tested on a patient suffering from cellulitis took approximately one week to relieve. After the cellulitis returned in a 14 days it was treated again. This happened twice and the herbal composition was topically applied again. In that case, the bacterial infection did not return.

[0054] The tests showed that severe cases of eczema and psoriasis may take considerably longer to relieve (1 to 2 weeks depending on the severity of the case). Some cases of eczema and psoriasis did return but often several weeks later and the returning cases were less severe (quite possibly due to the discontinuation of steroid use). These return cases were easily relieved within two or three days of treatment using the herbal composition. It must be noted that some of the tested eczema and psoriasis cases were individuals with a history of long term steroid treatment use, whose immune systems seem took longer to respond to the topical treatment using the extract of this invention.

[0055] In another test, the extract in the form of a spray was used to treat sunburn. The relief was instantaneous and there was no subsequent peeling or itching. The redness associated with sunburn disappeared within 24-36 hours.

[0056] The extract in the form of a spray was also tested to relieve insect bites and stings including yellow jackets, wasps, gnats, deer flies, etc. The relief was instantaneous.

[0057] During tests it was noted that extract helps in the treatment of photo-aging. Topical application of the herbal composition resulted in diminished appearance of wrinkles on the face and other parts of the body.

[0058] The herbal composition was also successfully tested to relieve razor burn and an astringent. The extract was topically applied and demonstrated beneficial results almost instantaneously.

[0059] Some eczema manifests itself on the lining of the inner ear and is a dermatological concern for professionals. At present, the tinnitus problem has no known cure and all treatments are judged by the user of the treatment. Treatments considered to be the most effective are steroidal and considered marginally effective and then only at the onset of the problem. Some tests were conducted by applying the herbal composition to the subject's ears for treatment of tinnitus (aural administration). While the studies of these problems using the instant herbal composition are ongoing, preliminary results demonstrate subject's positive response to the treatment when the spray was administered directly into the ear cavity.

[0060] Preliminary toxicology studies indicate that the extract is not toxic to humans or other mammals. One of the inventors personally consumed four ounces per day of the extract for two months with no noticeable side effects.

[0061] The extract prepared according to this invention has been tested on dogs for itching and works remarkably well. It has been tested as a nasal mist to stop the production of mucous on surgical sores in the nasal passages. It has been tested on one case of post operative contact dermatitis which could not be cured with conventional medicine over a period of 8 months. The case was relieved in approximately one week after using this extraction of this invention.

[0062] The tests showed that the extract has a potential for treatment of fungal, bacterial and viral infections (athlete's foot, cellulitis and shingles respectively) to a great extent.

[0063] The compounds in the extract are flavonoids. It is noted that the flavonoids may modify allergens, viruses and carcinogens acting as biological "response modifiers." The inventors believe that the mode of action of the extract is the interruption of the immune system response which is caused by the complicated reactions of T-Cells and subsequent histamine production by the mast cells. The flavonoids in the extract are believed to bind dermal proteins and other foreign proteins like wasp venom.

[0064] Although not empirically confirmed, it is also believed that the mechanism of action of the herbal composition of this invention on mammal skin is associated with interruption of the body immune response for a period long enough for the skin to heal. It was noted that most skin disorders return, except for certain types of contact dermatitis, but they return at a lower frequency and severity and can be kept under control with relative ease.

[0065] It is envisioned that the composition provided by this invention can be used as medical cosmetics for treating various diseases and changes of the skin, such as erythema (caused by shaving and make up). Using of composition after shaving, leads to a mild burning and calming the skin. After frequent use of make-up, or due to lack of safety measures, tension and irritation of the skin can occur. During preliminary tests, the composition was applied directly to the face, and the first noticeable signs of the disappearance of itching and rough skin were reported within several hours of application as a spray or saturated wipe.

[0066] The inventors were unable to identify any side effects when using this extract in the manner described.

[0067] The herbal composition can be used in the form of spray, wet wipes, and directly to the affected areas. Other forms suitable for topical application are envisioned. This composition is formulated to influence several diseases and disorders with active ingredients that have antioxidant, antibacterial, antiviral, antifungal, anti-inflammatory, antiseptic, regenerative, stimulatory, cosmetic and other effects.

[0068] The combination of ingredients shows synergistic effect in contact with skin. Unlike other herbal-based compositions, the composition provided by this invention has a greater effect due to the unique extract of hypericum hypericoides and ways of applying this composition by spraying with spray or applying directly to skin or using saturated wet wipes which increases the action of active agents because of their uniform application and facilitates penetration into the skin pores.

[0069] This invention is a natural herbal composition which is based on an aqueous extract of hypericum hypericoides leaves. It contains iron, various tannins, (mostly catechins), essential oils, resins, anthocyanins, hypericin - red color, carotene, choline, vitamin C, and some traces of alkaloids. It has a broad antibacterial, antiviral and anti-inflammatory effect. St. Andrew's Cross extract treats inflammatory processes on the skin, superficial wounds and burns by reducing inflammation of the skin.

[0070] Many changes and modifications can be made to the composition and method of the present invention without departing from the spirit thereof. We, therefore, pray that our rights to the present invention be limited only by the scope of the appended claims.

We claim:

1. An herbal composition based on an aqueous extract of hypericum hypericoides stabilized/preserved with a topically acceptable stabilizing/preservative agent in proportions suitable to form a topical agent for treatment of skin conditions.
2. The herbal composition of claim 1 in the form of spray and wet wipes.
3. The herbal composition of claim 1, wherein the skin disorder is selected from the group consisting of eczema, psoriasis, rosacea, cellulitis, contact dermatitis, seborrhoeic dermatitis, nummular dermatitis, dandruff, stasis dermatitis, perioral dermatitis, dermatitis herpetiformis, ecthyma, impetigo, shingles, urticaria, tinea pedis, tinea manuum, tinea cruris, sunburn, insect bites and stings, jellyfish stings, granuloma annulare, bullous pemphigoid, lichen planus, bed sore, inflammations caused by waterborne vectors, diaper rash, heat rash, pityriasis rosea, jungle rot, scleroderma, skin rashes and photo-aging.
4. The herbal composition of claim 1, wherein the acceptable stabilizing/preservative agent is a compound selected from a hydroxyl group.
5. The herbal composition of claim 4, wherein the stabilizing/preservative agent selected from the hydroxyl group is ethanol.
6. The herbal composition of claim 1, wherein the plant material is mixed with distilled boiling water at a rate of 1:25 to 1:30.
7. The herbal composition of claim 4, wherein the extract of hypericum hypericoides is mixed with a stabilizing/preservative agent at a ratio of 3% by volume of the filtered mixture.
8. The herbal composition of claim 1, wherein the concentration of active compounds is between 0.311% and 0.362% by weight of the total composition.

9. A process of obtaining an herbal composition on the basis of hypericum hypericoides comprising the steps of macerating plant material of hypericum hypericoides and mixing macerated plant material with boiling water, then stirring for a predetermined period of time, forcing the composition through a vacuum filter to obtain an herbal extract and then mixing the filtered herbal extract with a stabilizing/preservative agent to form a topical composition.

10. The process of claim 9, wherein the plant material comprises leaves of hypericum hypericoides.

11. The process of claim 9, wherein the plant material is mixed with distilled boiling water at a rate of 1:25 to 1:30.

12. The process of claim 9, wherein the stabilizing/preservative agent is alcohol.

13. The process of claim 9, wherein the filtered extract is mixed with the stabilizing/preservative agent at a ratio of 4 - 6% by volume of the stabilizing/preservative agent.

14. The process of claim 9, wherein the filtered extract is mixed with the stabilizing/preservative agent at a ratio of 3% by volume of the stabilizing/preservative agent.

15. The process of claim 9, wherein the mixture stabilizing/preservative agent is ethanol.

16. The process of claim 15, comprising a step of refrigerating the vacuum-filtered mixture.

17. The process of claim 9, wherein the concentration of active compounds is between 0.311% and 0.362% by weight of the total composition.

18. A method for treating a skin condition comprising topically administering a composition comprising an effective skin-penetrating amount of an aqueous extract of hypericum hypericoides.
19. The method of claim 18, wherein the composition additionally comprises a topically acceptable stabilizing/preservative agent.
20. The method of claim 19, wherein the topically acceptable stabilizing preservative agent is selected from a hydroxyl group.
21. The method of claim 20, wherein topically acceptable stabilizing preservative agent is ethanol.
22. The method of claim 20, wherein the topically acceptable stabilizing preservative agent is present at a ratio of 4 - 6% by volume of the composition.
23. The method of claim 18, wherein the skin condition is selected from the group consisting of eczema, psoriasis, rosacea, cellulitis, contact dermatitis, seborrhoeic dermatitis, nummular dermatitis, dandruff, stasis dermatitis, perioral dermatitis, dermatitis herpetiformis, ecthyma, impetigo, shingles, urticaria, tinea pedis, tinea manuum, tinea cruris, sunburn, insect bites and stings, jellyfish stings, granuloma annulare, bullous pemphigoid, lichen planus, bed sore, inflammations caused by waterborne vectors, diaper rash, heat rash, pityriasis rosea, jungle rot, scleroderma, skin rashes and photo-aging.
24. The method of claim 18, wherein the concentration of active compounds is between 0.311% and 0.362% by weight of the total composition.
25. The method of claim 18, wherein macerated leaves of hypericum hypericoides are mixed with distilled boiling water at a rate of 1:25 to 1:30.
26. The method of claim 18, wherein the composition is administered directly to skin in the form of a spray.

27. The method of claim 18, wherein the composition is administered directly to skin in the form of a wet wipe.
28. The method of claim 18, wherein the composition is aurally administered.
29. A method for treating a skin condition comprising topically administering a composition comprising 0.311% and 0.362% by weight of active compounds of an aqueous extract of hypericum hypericoides and between 4 and 6% by volume of the composition being alcohol.
30. The method of claim 29, wherein the skin condition is selected from the group consisting of eczema, psoriasis, rosacea, cellulitis, contact dermatitis, seborrhoeic dermatitis, nummular dermatitis, dandruff, stasis dermatitis, perioral dermatitis, dermatitis herpetiformis, ecthyma, impetigo, shingles, urticaria, tinea pedis, tinea manuum, tinea cruris, sunburn, insect bites and stings, jellyfish stings, granuloma annulare, bullous pemphigoid, lichen planus, bed sore, inflammations caused by waterborne vectors, diaper rash, heat rash, pityriasis rosea, jungle rot, scleroderma, skin rashes and photo-aging.
31. The method of claim 29, wherein alcohol is ethanol.
32. The method of claim 29, wherein alcohol is present at the ratio of 3% by volume.
33. The method of claim 29, wherein the composition is administered in the form of a spray or wet wipe directly to affected skin.
34. The method of claim 33, wherein the composition is administered aurally.

AMENDED CLAIMS

received by the International Bureau on 08 August 2013 (08.08.2013)

1. An herbal composition based on an aqueous extract of hypericum hypericoides stabilized/preserved with a topically acceptable stabilizing/preservative agent in proportions suitable to form a topical agent for treatment of skin conditions, said aqueous extract being prepared by adding distilled water heated to 100°C to hypericum hypericoides, stirring and then vacuum filtering the mixture to remove solids and water before combining filtered extract with the stabilizing/preservative agent
2. The herbal composition of claim 1 in the form of spray and wet wipes.
3. The herbal composition of claim 1, wherein the skin disorder is selected from the group consisting of eczema, psoriasis, rosacea, cellulitis, contact dermatitis, scborrheic dermatitis, nummular dermatitis, dandruff, stasis dermatitis, perioral dermatitis, dermatitis herpetiformis, ecthyma, impetigo, shingles, urticaria, tinea pedis, tinea manuum, tinea cruris, sunburn, insect bites and stings, jellyfish stings, granuloma annulare, bullous pemphigoid, lichen planus, bed sore, inflammations caused by waterborne vectors, diaper rash, heat rash, pityriasis rosea, jungle rot, scleroderma, and skin rashes.
4. The herbal composition of claim 1, wherein the acceptable stabilizing/preservative agent is a compound selected from a hydroxyl group.
5. The herbal composition of claim 4, wherein the stabilizing/preservative agent selected from the hydroxyl group is ethanol.
6. The herbal composition of claim 1, wherein the plant material is mixed with distilled boiling water at a rate of 1:25 to 1:30.
7. The herbal composition of claim 4, wherein the extract of hypericum hypericoides is mixed with a stabilizing/preservative agent at a ratio of 3% by volume of the filtered mixture.
8. The herbal composition of claim 1, wherein the concentration of active compounds is between 0.311% and 0.362% by weight of the total composition.

9. A process of obtaining an herbal composition on the basis of hypericum hypericoides comprising the steps of macerating plant material of hypericum hypericoides and mixing macerated plant material with boiling water, then stirring for a predetermined period of time, forcing the composition through a vacuum filter to remove solids and water to obtain an herbal extract and then mixing the filtered herbal extract with a hydroxyl stabilizing/preservative agent to form a topical composition having about 3% by volume of filtered extract.
10. The process of claim 9, wherein the plant material comprises leaves of hypericum hypericoides.
11. The process of claim 9, wherein the plant material is mixed with distilled boiling water at a rate of 1:25 to 1:30.
12. The process of claim 9, wherein the stabilizing/preservative agent is alcohol.
13. The process of claim 9, wherein the filtered extract is mixed with the stabilizing/preservative agent at a ratio of 4 - 6% by volume of the stabilizing/preservative agent.
14. The process of claim 9, wherein the filtered extract is mixed with the stabilizing/preservative agent at a ratio of 3% by volume of the stabilizing/preservative agent.
15. The process of claim 9, wherein the mixture stabilizing/preservative agent is ethanol.
16. The process of claim 15, comprising a step of refrigerating the vacuum-filtered mixture.
17. The process of claim 9, wherein the concentration of active compounds is between 0.311% and 0.362% by weight of the total composition.
18. A method for treating a skin condition comprising topically administering a composition comprising an effective skin-penetrating amount of an aqueous extract of hypericum hypericoides prepared by adding distilled water heated to 100°C to hypericum hypericoides, stirring and then vacuum filtering the mixture to remove solids and water before combining filtered extract with the stabilizing/preservative agent
19. The method of claim 18, wherein the composition additionally comprises a topically acceptable stabilizing/preservative agent.
20. The method of claim 19, wherein the topically acceptable stabilizing preservative agent is selected from a hydroxyl group.

21. The method of claim 20, wherein topically acceptable stabilizing preservative agent is cLhanoI.
22. The method of claim 20, wherein the topically acceptable stabilizing preservative agent is present at a ratio of 4 - 6% by volume of the composition.
23. The method of claim 18, wherein the skin condition is selected from the group consisting of eczema, psoriasis, rosacea, cellulitis, contact dermatitis, seborrhoeic dermatitis, nummular dermatitis, dandruff, stasis dermatitis, perioral dermatitis, dermatitis herpetiformis, ecthyma, impetigo, shingles, urticaria, tinea pedis, tinea manuum, tinea cruris, sunburn, insect bites and stings, jellyfish stings, granuloma annulare, bullous pemphigoid, lichen planus, bed sore, inflammations caused by waterborne vectors, diaper rash, heat rash, pityriasis rosea, jungle rot, scleroderma, and skin rashes.
24. The method of claim 18, wherein the concentration of active compounds is between 0.311% and 0.362% by weight of the total composition.
25. The method of claim 18, wherein macerated leaves of hypericum hypericoides are mixed with distilled boiling water at a rate of 1:25 to 1:30.
26. The method of claim 18, wherein the composition is administered directly to skin in the form of a spray.
27. The method of claim 18, wherein the composition is administered directly to skin in the form of a wet wipe.
28. The method of claim 18, wherein the composition is orally administered.
29. A method for treating a skin condition comprising topically administering a composition comprising 0.311% and 0.362% by weight of active compounds of an aqueous extract of hypericum hypericoides and between 4 and 6% by volume of the composition being alcohol prepared by adding distilled water heated to 100°C to hypericum hypericoides, stirring and then vacuum filtering the mixture to remove solids and water before combining filtered extract with the stabilizing/preservative agent.

30. The method of claim 29, wherein the skin condition is selected from the group consisting of eczema, psoriasis, rosacea, cellulitis, contact dermatitis, seborrhoeic dermatitis, nummular dermatitis, dandruff, stasis dermatitis, perioral dermatitis, dermatitis herpetiformis, ecthyma, impetigo, shingles, urticaria, tinea pedis, tinea manuum, tinea cruris, sunburn, insect bites and stings, jellyfish stings, granuloma annulare, bullous pemphigoid, lichen planus, bed sore, inflammations caused by waterborne vectors, diaper rash, heat rash, pityriasis rosea, jungle rot, scleroderma, and skin rashes.
31. The method of claim 29, wherein alcohol is ethanol.
32. The method of claim 29, wherein alcohol is present at the ratio of 3% by volume.
33. The method of claim 29, wherein the composition is administered in the form of a spray or wet wipe directly to affected skin.
34. The method of claim 33, wherein the composition is administered orally.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 13/33393

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A01N 65/00 (2013.01)**USPC - 424/725**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) - A01N 65/00 (2013.01)

USPC - 424/725

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
See search terms below:

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PatBase, PubWest, Freepatentsonline, DialogWeb, Google Scholar, Google

Search Terms: hypericum hypericoid, hypericum, skin, water, ethanol, St. Andrew's cross, tutsan, extract, ethanol, alcohol, hypericin, aqueous, leaf, boil, flavonoid, bioflavonoid

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2007/0122492 A1 (Behr et al.) 31 May 2007 (31.05.2007) para [0026], [0068], [0071]-[0073], [0075], [0079]-[0083], [0126], [0159], [0182], [0197], [0270]	1-34
Y	Foster et al. A Field Guide to Medicinal Plants and Herbs of Eastern and Central North America. Houghton Mifflin Company. 2000, pg105, In5 from bottom; pg106, In20 from bottom	1-34
Y	US 2008/0241 101 A1 (Amano et al.) 02 October 2008 (02.10.2008), para [0050], [0052]	1, 9, 18
Y	US 6,699,512 B2 (Quintanilla Almagro et al.) 02 March 2004 (02.03.2004) col2, lin40-46, 55-57	1, 9, 18
A	US 6,291,241 B1 (Castor et al.) 18 September 2001 (18.09.2001) col 2, ln 54-57	9, 18,
A, P	KR 2012/0058830 A (Kim et al.) 08 June 2012 (08.06.2012), abstract	1, 9, 18

 Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 May 2013 (14.05.2013)

Date of mailing of the international search report

10 JUN 2013

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774