



(12)

Oversættelse af
europæisk patentskrift

Patent- og
Varemærkestyrelsen

(51) Int.Cl.: **C 07 D 215/227 (2006.01)** **A 23 L 13/00 (2016.01)** **A 23 L 27/40 (2016.01)**
C 07 D 407/02 (2006.01)

(45) Oversættelsen bekendtgjort den: **2019-01-21**

(80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2018-10-24**

(86) Europæisk ansøgning nr.: **14705458.9**

(86) Europæisk indleveringsdag: **2014-02-07**

(87) Den europæiske ansøgnings publiceringsdag: **2016-02-17**

(86) International ansøgning nr.: **US2014015192**

(87) Internationalt publikationsnr.: **WO2014124191**

(30) Prioritet: **2013-02-08 US 201361762781 P** **2013-02-08 US 201361762792 P**
2013-02-08 US 201361762798 P **2013-02-08 US 201361762804 P**
2013-02-11 US 201361763244 P **2013-02-11 US 201361763274 P**
2013-02-11 US 201361763300 P

(84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV**
MC MK MT NL NO PL PT RO RS SE SI SK SM TR

(73) Patenthaver: **General Mills, Inc., PO Box 1113, Number One General Mills Boulevard, Minneapolis, Minnesota 55440, USA**

(72) Opfinder: **VAN LENGERICH, Bernhard H., 1855 Troy Lane North, Plymouth, Minnesota 55447, USA**
GRUESS, Olaf, 12725 91st Avenue North, Maple Grove, Minnesota 55369, USA
HANS, Joachim, Bergblick 7a, 37603 Holzminden, Tyskland
HAUSTEDT, Lars Ole, Clara-Zetkin Str. 27, 14471 Potsdam, Tyskland
HOCHHEIMER, Andreas, Reichsstrasse 4a, 93055 Regensburg, Tyskland
KROHN, Micheal, Alexanderstrasse, 64653 Lorsch, Tyskland
MULLER, Jens-Peter, Varziner Str. 10, 12161 Berlin, Tyskland
NOWAKOWSKI, Christine M., 2745 Norwood Lane North, Plymouth, Minnesota 55441, USA
PECORE, Suzanne Denise, 331 Fillmore Street, Pasadena, California 91106, USA
RATHJEN-NOWAK, Candace Michelle, 4562 McAllister Avenue NE, St. Michael, Minnesota 55376, USA
SCARABOTTOLO, Lia, Via Unbria 60, I-20068 Peschiera Borromeo, Italien
SIEMS, Karsten, Feuerbachstr. 7, 14552 Michendorf, Tyskland

(74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**

(54) Benævnelse: **FØDEVAREPRODUKT MED REDUCERET NATRIUM**

(56) Fremdragne publikationer:
WO-A1-2010/124905
ANJAN REDDY K ET AL: "REDUCING THE SODIUM CONTENT OF FOODS: A REVIEW", JOURNAL OF FOOD PROTECTION, INTERNATIONAL ASSOCIATION FOR FOOD PROTECTION, US, vol. 54, no. 2, 1 January 1991

(1991-01-01), pages 138-150, XP008078157, ISSN: 0362-028X

BEST D: "Compensating for sodium: the low-salt solution", PREPARED FOODS, GORMAN PUB., CHICAGO, IL, US, vol. 158, no. 22, 1 February 1989 (1989-02-01), pages 97-98, XP008098474, ISSN: 0747-2536

GYU-HEE LEE: "A salt substitute with low sodium content from plant aqueous extracts", FOOD RESEARCH INTERNATIONAL, vol. 44, no. 2, 1 March 2011 (2011-03-01), pages 537-543, XP055113729, ISSN: 0963-9969, DOI: 10.1016/j.foodres.2010.11.018

DESMOND ET AL: "Reducing salt: A challenge for the meat industry", MEAT SCIENCE, ELSEVIER SCIENCE, GB, vol. 74, no. 1, 1 September 2006 (2006-09-01), pages 188-196, XP027989851, ISSN: 0309-1740 [retrieved on 2006-09-01]

DESCRIPTION

FIELD

[0001] This disclosure generally relates to, among other things, food products having a compound that modifies or enhances the taste of the food product, for example, the saltiness of the food product.

BACKGROUND

[0002] Sodium chloride, ordinary table salt, is the prototypical compound for eliciting the perception of salty taste. However, attempts to reduce sodium consumption have led investigators to find suitable substitutes for sodium chloride or to reduce sodium chloride amounts, without sacrificing salty taste.

[0003] Salts can elicit complex tastes, including mixtures of sweet, bitter, sour, umami, and salty perceptual components. It is believed that the cations of salts impart the perceptual taste component, while the anions, in addition to contributing to tastes of their own, modify the perception of the taste of the cations. By way of example, sodium and lithium are believed to impart only salty tastes, while potassium and other alkaline earth cations produce both salty and bitter tastes. Among the anions commonly found in foods, the chloride ion is considered to be the least inhibitory to the salty taste, while the citrate anion is more inhibitory.

[0004] Many attempts have been made to provide salty tasting compositions as a substitute for table salt which will give the same or a similar seasoning effect and which are comprised of substantially reduced quantities of sodium chloride. To this end, potassium chloride, ammonium chloride, and similar compounds have been suggested. The use of such salts, and combinations of such salts, leaves much to be desired as to taste. None of them individually or in combination positively affects other taste modalities and tastes like sodium chloride. Each alone has a disagreeable taste, as do mixtures of such salts. For example, potassium chloride has a strong aftertaste that is characterized as "bitter" by most people. Ammonium chloride also has a bitter aftertaste.

[0005] WO 2010/124905 A1 discloses a process for preparing a low-sodium salt product. A review about reducing the sodium content of foods is provided by K.A. Reddy et al. in Journal of Food Protection, International Association for Food Protection, vol. 54, no. 2 (1991), 138-150. D. Best deals with "Compensation for sodium: The low-salt solution" in Prepared Foods, Gorman Pub., US, vol. 158, no. 22 (1989), 97-98. G.-H. Lee describes "A salt substitute with low sodium content from plant aqueous extracts" in Food Research International, vol. 44, no. 2 (2011), 537-543. E. Desmond deals with "Reducing salt: A challenge for the meat industry" in Meat Science, Elsevier Science, GB, vol. 74, no. 1 (2006), 188-196.

SUMMARY

[0006] The present invention is directed to food products as defined in the claims.

[0007] This disclosure describes, among other things, compounds that elicit or enhance the perception of salty taste, or another taste associated with consumption of sodium chloride or other salts, or that interact with a receptor or ion channel associated with the perception of salty taste or another complex taste associated with consumption of sodium chloride or other salts. In embodiments, the compounds are naturally derived taste modulating compounds used as ingredients in food products to elicit or enhance perception of salty taste. In embodiments, the food products contain lower amounts of sodium than normal.

[0008] As described herein, a number of derived compounds were screened for their ability to modulate activity of a sodium channel *in vitro*. Many of the identified compounds were found to enhance the saltiness of a composition containing sodium chloride.

[0009] One or more of the compounds, compositions, food products or methods described herein provide one or more advantages over prior compounds, compositions, food products or methods. For example, food products that include one or more taste modulating or salty taste modulating compounds described herein may have lower sodium content relative to food products that do not include such taste modulating or salty compounds while imparting a similar level of saltiness. This and other advantages will be readily understood from the following detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010]

FIG. 1 is a table that provides results of DAP score testing regarding the perception of saltiness of various combinations of compounds in sodium chloride solution.

FIG. 2 is a table that provides results of DAP score testing regarding the perception of saltiness of various combinations of compounds in combination in broth solution.

DETAILED DESCRIPTION

[0011] The present invention is directed to food products as defined in the claims.

[0012] This disclosure describes, among other things, compounds that elicit or enhance the perception of salty taste or another taste associated with consumption of sodium chloride. In embodiments, the compounds are taste modulating compounds used as ingredients in food products to elicit or enhance perception of salty taste. In embodiments, the food products are food products that contain reduced amounts of sodium, while imparting a salty taste typically associated with higher amounts of sodium.

[0013] The food product includes at least one ingredient, at least one salt that imports a salty taste and at least one compound of Formula B3 described below, wherein the compound of Formula B3 is either derived from a natural product or synthesized, and if derived from a natural product, it is separated from its naturally occurring environment.

[0014] As used herein, a "food product" is a food in a form that does not exist in nature. In embodiments, a food product includes at least two edible ingredients that do not exist together in nature. A "food" is a nutritious substance that animals, including humans, pets and livestock, eat or drink. A "nutritious substance" is a macronutrient such as a fat, carbohydrate or protein, or a micronutrient such as an essential or non-essential vitamin or mineral.

[0015] One or more taste modulating or salty taste modulating compounds described herein or derivatives thereof, alone or in combination, may be incorporated into a food product. The one or more compounds may elicit a perception of saltiness when the food product is consumed. In embodiments, the one or more compounds are included in a food product that contains a salt that imparts a salty taste. Preferably, at least one of the one or more compounds is a taste modulating compound or salty taste modulating compound.

[0016] The food product includes an ingredient, a salt that imparts a salty taste, and at least one compound of formula B3. The ingredient may be a nutritious ingredient; that is, an ingredient that is a nutritious substance. The taste modulating or salty taste modulating compound of formula B3 may be present in the food product in an amount sufficient to enhance the salty taste of the food product. In embodiments, the ingredient, the salt and the taste modulating or salty taste modulating compound are present in the food product in amounts or concentrations not found in naturally existing food products, such as bananas, peppers, avocados, wheat, or the like.

[0017] In embodiments, at least one of the one or more compounds is a salty taste modulating compound and is present in the food product in an amount or concentration sufficient to elicit or enhance the perception of saltiness. In embodiments, the one or more salty taste modulating compounds are present in the food product in an amount or concentration sufficient to elicit or enhance the perception of salty taste such that less salt may be included in the food product to elicit a similar perception of saltiness as a substantially similar food product that does not include the one or more salty taste modulating compounds. Preferably, the reduced salt food product elicits the same or a similar perception of saltiness as a substantially similar food product that does not include the one or more salty taste modulating compounds.

[0018] In embodiments, the one or more taste modulating or salty taste modulating compounds are present in a food product in an amount or concentration sufficient to elicit or enhance the perception of salty taste such that the amount of sodium may be reduced by about 10 mg or more per serving relative to a substantially similar food product that does not have the one or more taste modulating or salty taste modulating compounds while having a similar salty taste. In embodiments, the one or more taste modulating or salty taste modulating compounds are present in a food product in an amount or concentration sufficient to elicit or enhance the perception of salty taste such that the amount of sodium in a serving of a food product may be reduced to about 150 mg or less, more particularly to about 100 mg or less, more particularly to about 75 mg or less, more particularly to about 25 mg or less, more particularly to about 10 mg or less. By way of example, it may be desirable to reduce sodium by about 10 mg or more in cereals or snacks per serving relative to a substantially similar food product that does not have the one or more taste modulating or salty taste modulating compounds while having a similar salty taste. It may be desirable to reduce sodium to about 150 mg or less, more particularly to about 100 mg or less, more particularly to about 75 mg or less, more particularly to about 25 mg or less, more particularly to about 10 mg or less in cereals or snacks per serving. For cereal, a typical serving size is 50 grams. Of course, cereals may have other serving sizes.

[0019] In embodiments, the one or more taste modulating or salty taste modulating compounds are present in a food product in an amount or concentration sufficient to elicit or enhance the perception of salty taste such that the amount of sodium may be reduced by about 20 mg or more per serving relative to a substantially similar food product that does not have the one or more taste modulating or salty taste modulating compounds while having a similar salty taste. In embodiments, the one or more taste modulating or salty taste modulating compounds are present in a serving of a food product in an amount or concentration sufficient to elicit or enhance the perception of salty taste such that the amount of sodium may be reduced to about 800 mg or less, more particularly to about 500 mg or less, more particularly to about 300 mg or less, more particularly to about 100 mg or less, more particularly to about 20 mg or less. By way of example, it may be desirable to reduce sodium by about 20 mg or more in meals per serving. It may be desirable to reduce sodium to about 800 mg or less, more particularly to about 500 mg or less, more particularly to about 300 mg or less, more particularly to about 100 mg or less, more particularly to about 20 mg or less in meals per serving.

[0020] In embodiments, the one or more taste modulating or salty taste modulating compounds are present in a food product in an amount or concentration sufficient to elicit or enhance the perception of salty taste such that the amount of sodium may be reduced by about 100 mg or more per serving relative to a substantially similar food product that does not have the one or more taste modulating or salty taste modulating compounds while having a similar salty taste. In embodiments, the one or more taste modulating or salty taste modulating compounds are present in a serving of a food product in an amount or concentration sufficient to elicit or enhance the perception of salty taste such that the amount of sodium may be reduced to about 800 mg or less, more particularly to about 500 mg or less, more particularly

to about 300 mg or less, more particularly to about 200 mg or less, more particularly to about 100 mg or less relative to a substantially similar food product that does not have the one or more taste modulating or salty taste modulating compounds while having a similar salty taste. By way of example, it may be desirable to reduce sodium by about 100 mg or more in soups per serving. It may be desirable to reduce sodium to about 800 mg or less, more particularly to about 500 mg or less, more particularly to about 300 mg or less, more particularly to about 200 mg or less, more particularly to about 100 mg or less in soups per serving. For soup, a typical serving size is 250 grams. Of course, soups may have other serving sizes.

[0021] Any suitable combination of compounds described herein, or derivatives thereof, may be included in a food product. In embodiments, a food product includes a combination of compounds such that the combination includes at least two structurally diverse taste modulating or salty taste modulating compounds.

[0022] A food product or composition may include one or more compounds described herein, or derivatives thereof, in any suitable concentration. By way of example, a compound described herein, or a derivate thereof, such as a taste modulating or salty taste modulating compound may be present in a food product at a concentration of about 0.01% by weight or greater, about 2% by weight or less, or from about 0.01 % by weight to about 2% by weight. It will be understood that the concentration of the salt or salts in the food product may affect the desired concentration of a taste modulating or salty taste modulating compound. For example, if more salt is present, less taste modulating or salty taste modulating compound may be desired. In addition, it will be understood that the presence of more than one taste modulating or salty taste modulating compound may affect the desired concentration of other taste modulating or salty taste modulating compounds, particularly if the effects of the taste modulating or salty taste modulating compounds are additive or synergistic.

[0023] Any salt that imparts a salty taste may be present or incorporated into a food product that contains a bioactive, taste modulating, or salty taste modulating compound. The most commonly used salt for food applications is sodium chloride (typically referred to as common table salt). Other illustrative sources of sodium salts that may be present or incorporated into a food product include sodium phosphates, mono sodium glutamate, sodium nitrite, sodium nitrate, sodium bicarbonate, sodium lactate, sodium citrate, and sodium stearoyl lactylate. Similar lithium, potassium, ammonium or other alkali earth salts may be present or included in addition or as an alternative to one or more sodium salts.

[0024] In embodiments, a food product includes sodium chloride as a salt that imparts a salty taste. Sodium chloride may be present in the food product at any suitable amount or concentration. In embodiments, sodium chloride is present in the food product in an amount up to about 10.0 weight percent, more particularly, up to about 5.0 weight percent, even more particularly up to about 1.2 weight percent, or in the range of about 0.017 to about 1.2 weight percent, or about 0.1 to about 1, or about 0.4 to about 0.6 weight percent. In embodiments, a food product that includes one or more bioactive, taste modulating, or salty taste modulating compounds comprises no more than 0.04 weight percent, no more than 0.1 weight percent

sodium, no more than 0.2 weight percent, no more than 0.25 weight percent sodium, no more than 0.3 weight percent, no more than 0.4 weight percent, no more than 0.5 weight percent sodium, no more than 0.75 weight percent sodium, no more than 1 weight percent sodium, no more than 5 weight percent sodium, or no more than 10 weight percent sodium. It will be understood that a desired weight percent of sodium may vary depending on the type of food product. For example, it may be desirable for a seasoning to have a higher weight percent sodium than a soup or a breakfast cereal. In embodiments, a food product that includes one or more taste modulating or salty taste modulating compounds comprises no more than 100 mg sodium per serving, no more than 250 mg sodium per serving, no more than 500 mg sodium per serving.

[0025] One or more taste modulating or salty taste modulating compounds may be utilized in connection with virtually any food product for which it is desired to elicit or enhance the perception of a salty taste or other taste associated with consumption of a salt. The taste modulating or salty taste modulating compounds can find application for imparting saltiness to beverages or food dishes or as an ingredient in snack foods or other food products in which saltiness is desired.

[0026] Examples of food products that may incorporate one or more taste modulating or salty taste modulating compound include a confectionary, a gum, a bakery product, an ice cream, a dairy product, a fruit snack, a chip or crisp, an extruded snack, a tortilla chip or corn chip, a popcorn, a pretzel, a nut, a snack bar, a meal replacement, a ready meal, a soup, a pasta, a canned food, a frozen processed food, a dried processed food, an instant noodle, a chilled processed food, an oil or fat, a sauce dressing or condiment, a dip, a pickled product, a seasoning, a baby food, a spread, a chip or a crisp such as chips or crisps comprising potato, corn, rice, vegetable (including raw, pickled, cooked and dried vegetables), a fruit, a grain, a soup, a seasoning, a baked product such as a ready-to-eat breakfast cereal, hot cereal or dough, an ice cream such as a frozen yogurt, a dairy products such as a yogurt or cheese, ready meal, a soup, a pasta, a canned food, a frozen processed food, a dried processed food, an instant noodle, or a chilled processed food, a beverage including beverages that include fiber or protein a meat or a meat substitute, a pet food, an animal product, a medical food, a nutritional supplement, a vitamin supplement, and an infant formula product.

[0027] In embodiments, one or more bioactive, taste modulating, or salty taste modulating compounds are incorporated into a medicinal or pharmaceutical product, or the like.

[0028] In embodiments, a food product is a processed food product. Food processing includes the transformation of raw ingredients into food or transforming forms of food into other forms of food. Food processing often includes using harvested crops or animal products to produce marketable products sold to consumers at stores, restaurants and the like. Processed food products include products for which additional processing by a consumer occurs after purchase but prior to consumption (e.g., heating, cooking, baking, or the like).

[0029] Particularly suitable food products including soup, meal kits, grain products such as

ready-to-eat cereals, snacks, bars and baked dough, and dairy products such as ice cream, yogurt and cheese. In some aspects, a bioactive, taste modulating, or salty taste modulating compound is used to reduce the amount of sodium salt that is typically included in soups, including (but not limited to) chicken or poultry broth, chicken- or poultry-based soups (such as chicken noodle soup), tomato-based soups, and the like. In some aspects, a taste modulating or salty taste modulating compound is used to reduce sodium salt in meal kits, such as kits that include ingredients to be combined with meat to prepare a meal. Such meal kits can include dried components (such as noodles, rice, dried potatoes, or the like) and seasoning packages. In some aspects, a taste modulating or salty taste modulating compound is used to reduce the sodium chloride that is typically added to a snack food to enhance its flavor. Exemplary snack foods include potato chips, corn chips, pretzels, fruit-type snacks, and snack mixes including any mixes of any of these foods with other ingredients (such as cereals).

[0030] In some aspects, a taste modulating or salty taste modulating compound is used to reduce the amount of sodium salt that is typically included in a ready-to-eat cereal or other grain-based food products, such as dough, baked goods, grain snacks, grain bars, or the like. In some aspects, a taste modulating or salty taste modulating compound is used to reduce the amount of sodium salt that is typically included in dairy-based food products, such as fresh or frozen dairy products, which may include yogurt, ice cream, or the like. In some aspects, a taste modulating or salty taste modulating compound is used to reduce the amount of sodium salt that is typically included in packaged meal food products, such as packaged meals that contain rice, potatoes, or vegetables, dry packaged meals, frozen packaged meals, or the like.

[0031] For the purposes of the present disclosure "grain" includes grain and pseudograin. Examples of food grains include corn; sorghum; fonio; millet such as pearl millet, proso millet, finger millet, foxtail millet, Japanese millet, kodo millet and the like; Job's tears; wheat; rice; rye; barley; oat; triticale; wild rice; teff; amaranth; quinoa; buckwheat; and the like.

[0032] A taste modulating or salty taste modulating compound can also be used in connection with soup, broth, sauce (such as basting sauce), various seasoning sauces, ketchup, dressings, and other like foods.

[0033] In embodiments, a food product into which a taste modulating or salty taste modulating compound or composition is included has a water content of about 30% or more by weight. For example, the food product may have a water content of about 35% or more, or about 40% or more by weight. Non-limiting examples of food products that typically have water contents of about 30% or more by weight include soups, beverages, batters and dough.

[0034] In embodiments, a food product into which a taste modulating or salty taste modulating compound or composition is included has a water content of about 50% or more by weight. For example, the food product may have a water content of about 60% or more, or about 70% or more by weight. Non-limiting examples of food products that typically have water contents of about 50% or more by weight include soups and beverages. For example, a soup containing a taste modulating or salty taste modulating compound or composition may contain from about

50% water by weight to about 90% water by weight.

[0035] In embodiments, a food product into which a taste modulating or salty taste modulating composition is included has a water content of about 20% or less by weight. For example, the food product may be incorporated into dry food products that having low water contents. In embodiments, a taste modulating or salty taste modulating food product is included in a dried for as a seasoning. In embodiments, the seasoning comprises, consists essentially of, or consists of one or more taste modulating or salty taste modulating compounds, one or more carriers, and one or more salts.

[0036] A taste modulating or salty taste modulating compound can be employed to elicit the perception of salty taste or enhance the perceived salt taste of any salts used in food or beverage products. The preferred salt taste to be elicited or enhanced by the salty compounds is that of sodium chloride.

[0037] Moreover, a taste modulating or salty taste modulating compound described herein can be used to elicit or enhance the perceived salt taste of known salty tasting compounds that may be used as salt substitutes. Such compounds include amino acids such as cationic amino acids and low molecular weight peptides such as dipeptides and tripeptides. Specific examples of these compounds include arginine hydrochloride, lysine hydrochloride, and lysine-ornithine hydrochloride. These compounds exhibit a salty taste but are typically useful only at low concentrations since they exhibit a bitter flavor at higher concentrations. Ordinarily, these salt-tasting compounds will be used in concentrations in the range of about 1 to about 40 mM, or about 10 to about 30 mM. Thus, it is feasible to reduce the sodium chloride content of a food or beverage product by first formulating a food or beverage with less sodium chloride than is necessary to achieve a desired salt taste and then adding to the food or beverage a taste modulating or salty taste modulating compound described herein in an amount sufficient to enhance the salt taste of the salted food or beverage to reach the desired taste. In addition, sodium chloride content may be further reduced by substituting a salt-tasting cationic amino acid, a low molecular weight dipeptide, or mixtures thereof, for at least a portion of the salt.

[0038] In embodiments, a method includes setting a target salty taste of a food product, including an amount of a salt that imparts a salty taste in the food product, where the amount of the salt does not achieve the target level of salty taste, and including an amount of a salty taste enhancing compound (or more than one salty taste enhancing compounds) to achieve the desired salty taste. In embodiments, a method includes setting a target salty taste of a food product, including an amount of a salty-taste imparting sodium salt in the food product that does not achieve the target level of salty taste, including an amount of a non-sodium salt that imparts a salty taste and an amount of a salty taste enhancing compound (or more than one salty taste enhancing compounds) to achieve the desired salty taste.

PROCESSING

[0039] A taste modulating or salty taste modulating compound, or derivative thereof, described herein can be added to food products in dry or liquid form. For example, a taste modulating or salty taste modulating compound that is in the liquid form can be prepared by simply dissolving or suspending the compound in an appropriate relative amount in an aqueous liquid. Useful aqueous liquids include water, alcohol-water mixtures, triacetin, propylene glycol, and triglycerides and other known organic solvents. Depending upon the concentration of the taste modulating or salty taste modulating compound, it can be desirable to heat the mixture to dissolve the compound.

[0040] Taste modulating or salty taste modulating compounds that exist in a dry state, such as powders or granules, can be prepared by either mixing or blending the compounds with other components in the dry state. The dry blending or mixing can be carried out in any conventional suitable apparatus. In some aspects, the taste modulating or salty taste modulating compounds described herein can be prepared into dry compositions by commonly used methods of granulation from mixtures of the several ingredients, preferably initially conveniently smaller than forty mesh. Such starting mixtures can be wetted in known manner, granulated, and their granulations dried as usual and screened to give a product approximately the typical size of common table salt, for example, by taking the fraction passing through the thirty mesh screen and retained on the forty mesh screen.

[0041] Taste modulating or salty taste modulating compounds that exist in a dry composition state can be alternatively prepared by first forming a solution, emulsion or suspension of the compounds and other individual components, and then extruding or drying the solution or suspension. The preparation of the solution or suspension of the components can be carried out as described above in the context of preparing the liquid flavoring agents. The thus-prepared solution, emulsion or suspension can then be dried using any conventional suitable apparatus, such as a rotary drier, a drum drier, or a fluidized bed drier or spray drier.

[0042] Taste modulating or salty taste modulating compounds described herein can be prepared by thoroughly mixing the compounds with other components in the indicated proportions until a suitably mixed (for example, homogeneous) product is attained.

[0043] Compositions or formulations containing the taste modulating or salty taste modulating compounds can then be combined with a food product.

PERCEPTION OF SALTINESS

[0044] In embodiments, a composition that includes a salty taste modulating compound is perceived as imparting a quantity of saltiness equal to a substantially similar composition that does not include the salty taste modulating compound but that has a higher concentration of the salt. Preferably, the composition that includes the salty taste modulating compound imparts a perception of saltiness equal to the substantially similar composition that does not have the salty taste modulating compound when the composition has less salt than the substantially

similar composition (e.g., salt reduced by about 1% or more). For example, the composition that includes the salty taste modulating compound may impart a perception of saltiness equal to the substantially similar composition that does not have the salty compound when the composition that includes the salty taste modulating compound has a salt concentration reduced by about 2% or more, about 5% or more, about 7% or more, about 8% or more, about 9% or more, about 10% or more, about 11% or more, about 15% or more, about 20% or more, about 30% or more, about 35% or more, about 40% or more, or about 50% or more, relative to the substantially similar composition. In embodiments, one or more salty taste modulating compounds may be present in a food product in an amount sufficient to reduce the amount of a salt, such as sodium chloride, by about 1% or more, about 2% or more, about 5% or more, about 7% or more, about 8% or more, about 10% or more, about 11% or more, about 12% or more, about 15% or more, about 20% or more, about 22% or more, about 25% or more, about 30% or more, about 35% or more, about 40% or more, about 45% or more, about 50% or more, about 55% or more, about 60% or more, about 65% or more, about 70% or more, about 75% or more, about 80% or more, about 85% or more, about 90% or more, about 95% or more, or the like. Preferably, the reduced salt food product elicits the same or similar perception of saltiness as a substantially similar food product that does not include the one or more salty taste modulating compounds.

[0045] Perception of saltiness may be evaluated in any suitable manner. In embodiments, saltiness is determined by a trained analytical sensory panel. In embodiments, the trained sensory panel determines the saltiness of a composition having a salty taste modulating compound relative to a substantially similar composition having increased sodium chloride content.

[0046] Sensory panelists may be trained in any suitable manner. Preferably, the panelists are trained to discern salty taste or other attributes without reference to liking or acceptability. The panelists are also preferably trained to accurately quantify salty taste or other attributes according to an intensity scale. General information that may be helpful in understanding beneficial training protocols can be found in, for example, *Sensory Evaluation Techniques*, 4th Ed by Meilgaard M., Civille G.V. and Carr B.T (2007), CRC Press, pages 147-152. Prescreening, selection, and training of panelists may occur as described in one or more standards, such as *Hootman RC, Manual 13 MNL13 Manual on Descriptive Analysis Testing for Sensory Evaluation*, ASTM (1992); *STP758 Guidelines for the Selection and Training of Sensory Panel Members*, ASTM (1981); and *Munoz A.M and Civille, G.V., MLN13: The Spectrum Descriptive Analysis Method*, ASTM (1992). Preferably panelists are trained according to the *Spectrum Method* (Munoz A.M and Civille, G.V., MLN13: The Spectrum Descriptive Analysis Method, ASTM 1992).

[0047] Preferably, average scores regarding saltiness from more than one panelist trained to discern salty taste or other attributes using the same training are considered in determining whether a reduced salt food product elicits the same or similar perception of saltiness as a substantially similar food product that does not include the one or more salty taste modulating compounds. For example, a panel may contain three or more trained panelists, 5 or more

trained panelists, 7 or more trained panelists, 10 or more trained panelists, or the like.

[0048] A taste modulating or a salty taste modulating compound may be a compound that directly acts to elicit or enhance the perception of salty taste of a salt or may be a compound that is converted, when ingested, into a compound that directly acts to elicit enhance the perception of salty taste of the salt.

TASTE MODULATING, AND SALTY TASTE MODULATING COMPOUNDS

[0049] A variety of naturally-derived compounds were tested *in vitro* for their ability to activate or facilitate activation of a TrpML3 channel, a TrpV1 channel or an ENaC channel.

[0050] The TrpML3 (transient receptor potential cation channel, mucolipin subfamily, member 3) channel, also known as Mucolipin-3 is a protein that, in humans, is encoded by the MCOLN3 gene. The TrpV1 (transient receptor potential cation channel subfamily V member 1) channel, also known as the capsaicin receptor and the vanilloid receptor 1, is a protein that, in humans, is encoded by the TrpV1 gene. The ENaC (epithelial sodium channel), also known as sodium channel non-neuronal 1 (SCNN1) or amiloride sensitive sodium channel (ASSC) is a membrane-bound ion-channel that is permeable for Li⁺-ions, protons and especially Na⁺-ions.

[0051] Any compound that interacts with one or more of the TrpML3 channel, the TrpV1 channel and the ENaC channel may be useful for modulating taste or saltiness of a food product into which the compound is incorporated.

[0052] It is estimated that natural products, extracts, and isolated compounds that collectively contained about 2,000,000 potential taste modulating or salty taste modulating compounds were tested for sodium channel activity. About 600 of the 2,000,000 compounds had some level of sodium channel activity. About 300 of the 600 compounds had an increased threshold level of activity. Further analysis, including structure based toxicological analysis, resulted in 99 initial compounds being selected as candidates for taste modulating or salty taste modulating compounds. Presented herein are naturally-derived compounds and compound classes that have been identified as acting at one or more of these channels, or that otherwise may function as bioactive, taste modulating, or salty taste modulating compounds.

[0053] A listing of the 99 initially selected compounds (Cmpd), as well as the common names (where known), Chemical Abstract Service (CAS) Registry numbers where known (CAS-RN), Sources/Taxons (where known) from which the compounds were isolated (Source/Taxon) and common name of the sources (Common Name), is presented in **Table 1** below. The structures of the compounds are also presented herein. To the extent the structures conflict with other information provided, the structures of the 99 initially selected compounds should be considered determinative.

Table 1: Information regarding initially selected compounds

(according to claim 1, the food product comprises a compound of formula B3)

Cmpd	CAS-RN	Common name	Source/Taxon	Common Name
1 *	104264-55-3	12-Gingerol	Zingiber officinalis	Ginger
2 *	36752-54-2	[10]-Shogaol	Zingiber officinalis	Ginger
3 *	143615-75-2	[6]-Gingerdiacetate	Zingiber officinalis	Ginger
4 *	555-66-8	6-Shogaol	Aframomum meleguata, Zingiber officinalis	Grains of paradise; Ginger
5 *	39886-76-5	[6]-Gingerol	Zingiber officinalis	Ginger
6 *	53172-04-6	[7]-Paradol	Aframomum meleguata, Zingiber officinalis	Grains of paradise; Ginger
7 *	27113-22-0	[6]-Paradol or [6]-Gingerone	Aframomum meleguata, Zingiber officinalis	Grains of paradise; Ginger
8	626200-64-4	5-methoxy-[6]-Gingerol	Aframomum meleguata, Zingiber officinalis	Grains of paradise; Ginger
9 *	23513-08-8	8-Gingerol	Zingiber officinalis	Ginger
10	1083195-12-3	a pentadecatrienyl-1,3-benzenediol	Embelia ribes	False Black Pepper, white-flowered Embelia
11	79559-60-7	a 1-Ph-4-hepten-3-one	Kaempferia galanga	Kencur, aromatic ginger, sand ginger, cutcherry, resurrection lily
12	79559-61-8	a 1-Ph-5-OH-3-heptanone	Kaempferia galanga	Kencur, aromatic ginger, sand ginger, cutcherry, resurrection lily
13 *	205687-01-0	Capsiate	Capsiate	
14 *	147030-09-9	Pipersintenamide	Piper longum	Long pepper, Indian long pepper
15 *	55038-30-	Guineensine	Piper longum	Long pepper, Indian

(according to claim 1, the food product comprises a compound of formula B3)

Cmpd	CAS-RN	Common name	Source/Taxon	Common Name
	7			long pepper
16 *	182056-19-5		Evodia rutaecarpa	Evodia fruit
17 *	15266-38-3	Evocarpine	Evodia rutaecarpa	Evodia fruit
18 *	94-62-2	Piperine	Piper longum	Long pepper, Indian long pepper
19	52483-20-2	Irisresorcinol	Ardisia silvestris	
20	79559-61-8	a 1-Ph-5-OH-3-heptanone	Alpinia officinarum	Lesser galangal (ginger family)
21*	19408-84-5	Dihydrocapsaicin	Capsicum annuum	Serrano pepper
22 *	30511-77-4	Isochavicine	Piper longum, Piper nigrum	Long pepper, Indian long pepper; black pepper
23 *	517-73-7	Melicopicine	Teclea trichocarpa	
24 *			Zanthoxylum esquirolii	
25 *	41303-25-7	O-Methylglycosolone	Zanthoxylum esquirolii	
26 *	5307-59-5	Robustic acid	Derris robusta	
27 *	84-99-1	Xanthoxyletin	Toddalia asiatica; Millettia pulchra	Orange climber
28 *	4335-12-0	Toddaculin	Toddalia asiatica	Orange climber
29 *	351427-18-4	Vitetrifolin D	Vitex agnus	Vitex, Chaste Tree, Chasteberry, Abraham's Balm, Monk's Pepper
30 *	61263-52-3		Vitex agnus	Vitex, Chaste Tree, Chasteberry, Abraham's Balm, Monk's Pepper
31 *	465-92-9	Marrubin	Marrubium vulgare	White Horehound, Common Horehound
32 *	238088-78-3	Orthosiphon I	Orthosiphon stamineus	Cat whiskers

(according to claim 1, the food product comprises a compound of formula B3)

Cmpd	CAS-RN	Common name	Source/Taxon	Common Name
33 *	345905-36-4	Orthosiphon M	Orthosiphon stamineus	Cat whiskers
34 *	254896-53-2	Acsculioside A	Acsculus hippocastaneum	Horse-chestnut, conker tree
35 *			Gleditschia australis	Locust tree
36 *	1383715-41-0		Pithecoctenium echinatum	Monkey comb
37 *			Yucca gloriosa	Spanish dagger
38 *			Nephelium cuspidatum	Bayong
39 *	20874-52-6	Saikosaponin D	Bupleurum falcatum	Chinese thoroughwax; Sickle hare's car
40 *	1217879-76-9	a 3-benzofurancarboxylic ester	Salvia miltiorrhiza	Red sage, Chinese sage, tan shen, or danshen
41 *	27994-11-2	Cimigoside; b-D-Xylopyranoside	Cimicifuga racemosa	Black cohosh, black bugbane, black snakeroot, fairy candle
42 *	4373-41-5	Maslinic acid, Crategolic acid	Alchemilla xanthochlora	Lady's Mantle
43 *	77-52-1	Ursolic acid	Lavandula officinalis	Lavender, English lavender, common lavender, true lavender, narrow-leaved lavender
44 *	58546-54-6	Schizandrol B; Gomisin A	Schisandra chinensis	Five flavor berry
45 *	61281-38-7	Schisandrin A, Deoxyschizandrin	Schisandra chinensis	Five flavor berry
46 *	61281-37-6	Schizandrin B	Schisandra chinensis	Five flavor berry
47 *	102036-29-3	Protosappanin B		
48 *	129102-89-2	Secoisolariciresinol; Secoisolariciresin-4-yl b-D-glucopyranoside	Angelica archangelica	Garden Angelica, Holy Ghost, Wild Celery, Norwegian angelica
49 *	66322-34-7	Dihydrogualaretic acid	Schisandra chinensis	Five flavor berry

(according to claim 1, the food product comprises a compound of formula B3)

Cmpd	CAS-RN	Common name	Source/Taxon	Common Name
50 *	193816-85-2	Epicalyxin C	Alpinia katsumadai	Greater galangal
51 *	181490-70-0	Icariin	Angelica sinensis	dong quai or female ginseng
52 *	446030-43-9	Myrrhanone B	Commiphora mukul	Guggul, Indian Bdellium
53 *	936499-55-7	Brevifolin	Zanthoxylum piperitum	Japanese pepper, Japanese pricklyash
54 *	1083202-45-2		Fungus	Strain code: 01469fxxx000005
55 *			Erythrina variegata	Tiger's claw or Indian coral tree
56 *	155485-76-0	Senecrassidiol	Psidium guajava	Apple guava or common guava
57 *	84-26-4	Rutaecarpine	Evodia rutaecarpa	Evodia fruit
58 *	992-20-1	Salannin	Azadirachta indica	Neem, Nimtree and Indian Lilac
59 *	69222-20-4	Isoanthricin	Podophyllum peltatum	May Apple, hogapple, Indian apple, mayflower, umbrella plant, wild mandrake, American mandrake
60 *	850494-43-8	Mammea A/AD cyclo F	Mesua ferrea	Ceylon ironwood, Indian rose chestnut, or Cobra's saffron
61 *	7282-19-1	Atanine	Zanthoxylum piperitum	Japanese pepper, Japanese pricklyash
62 *	484-20-8	Bergapten	Petroselinum sativum	Parsley
63 *	133164-11-1	Prangol	Petroselinum sativum	Parsley
64 *	482-44-0	Imperatorin	Petroselinum sativum	Parsley
65 *	2543-94-4	Phelopterin	Petroselinum sativum	Parsley
66 *	497226-80-9	terpenoid coumarins	Ferula assa-foetida	Giant fennel, asant, food of the gods, jowani badian, stinking gum, Devil's dung
67 *	484-33-3	Pongamol	Millettia pulchra	

(according to claim 1, the food product comprises a compound of formula B3)

Cmpd	CAS-RN	Common name	Source/Taxon	Common Name
68 *			artificial	
69 *	1176891-50-1	Sakisacaulon A	Lichen	
70 *	1268481-32-8	Chalepin		
71 *	1092383-76-0	Rutamarin		
72 *	13164-03-9	Halepensin		
73 *	143-62-4	Digitoxigenin	Xysmalobium undulatum	Uzara
74 *	26241-51-0	Azadiradione	Azadirachta indica	Neem, Nimtree, Indian Lilac
75 *	95975-55-6	(Z)-Guggulsterone	Commiphora mukul	Guggul, Indian Bdellium
76 *	1941-73-7	Apobioside	Apocynum cannabinum	Dogbane, Amy Root, Hemp Dogbane, Indian Hemp, Rheumatism Root, Wild Cotton
77 *	3751-87-9	Apocannoside	Apocynum cannabinum	Dogbane, Amy Root, Hemp Dogbane, Indian Hemp, Rheumatism Root, Wild Cotton
78 *	86894-26-0	Hebelomic acid A methyl ester	Hebeloma senescens	mushroom
79 *	2221-82-1	β-Cyclocostunolide	Critonia morifolia	
80a *	546-43-0	Alantolactone	Inula helenium	Elecampane, horse-heal, marchalan
80b *	470-17-7	Isoalantolactone	Inula helenium	Elecampane, horse-heal, marchalan
81 *		a 2-octenyl-3-hydroxy-1,5-pentanedioic acid	Fungus	Strain code 02295fxxx000001
82 *	83797-45-9	16-Heptadecene-1,2,4-triol; Avocadene	Persea gratissima	Avocado
83 *	1356361-43-7	16-Heptadecene-1,2,4-triol, 1,4-	Persea gratissima	Avocado

(according to claim 1, the food product comprises a compound of formula B3)

Cmpd	CAS-RN	Common name	Source/Taxon	Common Name
		diacetate		
84 *	16423-52-2	N-Decyl acetamide	Bacteria	Strain code 0172axxx000002
85 *	21402-68-6	9-hydroxy-10,12,15-Octadecatrienoic acid	Marrubium vulgare	White Horehound, Common Horehound
86 *	167936-49-4	12-hydroxy-9,13,15-Octadecatrienoic acid	Petroselinum sativum	Parsley
87 *			Ricinus communis	Castor oil plant
88 *	15514-85-9	Dimorphecolic acid	Podophyllum peltatum	May Apple, hogapple, Indian apple, mayflower, umbrella plant, wild mandrake, American mandrake
89 *	463-40-1	Linolenic acid	Mesua ferrea	Ceylon ironwood, Indian rose chestnut, Cobra's saffron
90 *	35949-86-1	Gingerglycolipid C	Zingiber officinalis	Ginger
91 *	187218-23-1	Capsianoside E	Capsicum annuum	Serrano pepper
92 *	131580-15-9	Capsianoside D	Capsicum annuum	Serrano pepper
93 *	22338-69-8	Grandifloric acid	Aralia cordata, Espeletia spp.	Spikenard, udo
94 *	6619-97-2	Xylopic acid	Xylopia aethiopica	Bitterwood
95 *	32381-03-6	Angeloylegrandifloric acid	Sideritis hirsuta	Hairy ironwort
96 *	482-00-8	Lanceolatin B		
97 *	101140-06-1	Biapigenin	Fagopyrum esculentum, Hypericum perforatum	
98 *	64125-32-2		Millettia pulchra	
99 *	36640-12-7		Lichen	

* not within formula B3



[0054] Compounds 12 and 20 are the same compound isolated from different sources.

[0055] The CAS registry numbers presented in **Table 1** above reflect a compound or an isomer thereof. It will be understood that other isomers may have other CAS registry numbers. Further, the structures presented herein, to the extent that they show stereochemistry may not match the particular isomer of the CAS registry number presented in **Table 1**.

[0056] Those compounds for which no CAS registry numbers are provided in **Table 1**, as well as those for which registry numbers are provided, may be isolated or purified in any suitable manner. For example, the natural source of the compound, which is presented in **Table 1**, may be fractionated and the fractions subjected to chromatography, such as gas chromatography or HPLC, or other suitable separation process to isolate or purify the compound. The selection of, for example, a chromatography column and parameters can be readily identified based on the chemical structure of the compound. To facilitate isolation or purification or for verification, obtained fractions, subfractions, or individual compounds may be tested for ability to activate a sodium channel, for example, expressed in cells in culture, cell membrane, or the like and employing an appropriate assay, such as an electrophysiological assay, a colorimetric assay, or the like.


[0057] Alternatively or in addition, the compounds listed in **Table 1** may be synthesized. Alternatively or in addition, companies that have access to the appropriate natural sources or the ability to test for sodium channel activity may be contracted to isolate the compounds. Companies that have access to natural products or natural product libraries that may include sources presented in **Table 1** or that have expertise in development of assays for identification of compounds or fractions containing compounds capable of activating a sodium channel include Biotechnology Research And Information Network AG (Zwingenberg, Germany); AnalytiCon Discovery, GmbH (Potsdam, Germany); Albany Molecular Research, Inc. (Albany, New York, USA); Axxam SpA (Milan, Italy); Boulder BioPharmaceuticals, LLC, Boulder, CO; ChromaDcx (Irvine, California, USA); Enzo Life Sciences, Inc. (Farmingdale, New York, USA); IMD Natural Solutions GmbH (Dortmund, Germany); TimTec LLC (Newark, Delaware, USA); and The Natural Products Discovery Institute (Doylestown, Pennsylvania, USA).

[0058] The structures of the initially selected compounds are as follows:







(1), not within formula B3




(3), not within formula B3




(4), not within formula B3





(5), not within formula B3




(6), not within formula B3




(7), not within formula B3




(9), not within formula B3




(10),




(11),




(12),




(13), not within formula B3




(14), not within formula B3




(15), not within formula B3




(16), not within formula B3




(17), not within formula B3




(18), not within formula B3




(19),




(20),




(21), not within formula B3




(22), not within formula B3




(23), not within formula B3




(24), not within formula B3




(25), not within formula B3




(26), not within formula B3




(27), not within formula B3




(28), not within formula B3




(29), not within formula B3




(30), not within formula B3




(31), not within formula B3







(32), not within formula B3




(33), not within formula B3




(34), not within formula B3







(35), not within formula B3




(36), not within formula B3




(37), not within formula B3







(38), not within formula B3




(39), not within formula B3




(40), not within formula B3




(41), not within formula B3




(42), not within formula B3




(43), not within formula B3




(44), not within formula B3




(45), not within formula B3




(46), not within formula B3




(47), not within formula B3







(48), not within formula B3




(49), not within formula B3



(50), not within formula B3




(51), not within formula B3







(52), not within formula B3




(53), not within formula B3




(54), not within formula B3




(55), not within formula B3




(56), not within formula B3




(57), not within formula B3




(58), not within formula B3








(59), not within formula B3




(60), not within formula B3




(61), not within formula B3







(66), not within formula B3




(67), not within formula B3




(68), not within formula B3




(69), not within formula B3




(70), not within formula B3




(71), not within formula B3




(72), not within formula B3




(73), not within formula B3




(74), not within formula B3




(75), not within formula B3




(76), not within formula B3




(77), not within formula B3




(78), not within formula B3




(79), not within formula B3




(80a), not within formula B3




(80b), not within formula B3



(81), not within formula B3




(82), not within formula B3




ö


(83), not within formula B3




(84),




(85), not within formula B3




(86), not within formula B3




(87), not within formula B3




(88), not within formula B3




(89), not within formula B3



(90), not within formula B3




(91), not within formula B3




(92), not within formula B3







(93), not within formula B3




(94), not within formula B3




(95), not within formula B3




(96), not within formula B3






(97), not within formula B3



(98) not within formula B3,

and



(99). not within formula B3

[0059] Presented below are chemical compound structures based on groupings of structural relationships of the 99 initially selected compounds. The compounds are grouped into 21 categories (A through U) containing one or more of the 99 initially selected compounds. Within some categories, sub-categories are described.

According to claim 1, the food product comprises a compound of formula B3,


[0060] In some instances, a compound may be in more than one category due to its structural similarity to compounds in more than one category. It will be understood that structural similarities of the various compounds other than those presented herein exist and that groupings into categories other than those presented herein are possible and contemplated.

[0061] Each of the 22 categories of compounds presented herein is discussed independently. That is, discussion of substituents with regard to one category should not be construed to limit discussion of substituents with regard to another category. For example, R¹ for the group A compounds is independently defined relative to R¹ for the group B compounds. In addition,

discussion of substituents with regard to subgroups is independently defined. For example, R¹ for the group J1 compounds is independently defined relative to R¹ for the group J2 compounds, unless otherwise stated.

GROUP A COMPOUNDS

[0062] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:




where:

R¹ is H or C₁-C₁₀ alkyl;

R² is H or C₁-C₃ alkyl;

X is CHOR³ or C=O;


R³ is H, C₁-C₃ alkyl, or



R⁴ is H or C₁-C₃ alkyl;

Y is CR⁵=CH or CHR⁵-CH₂;

R⁵ is H, OH, -OCH₃, -OCH₂CH₃, -O-OCH₂CH₂CH₃, or




and

R⁶ is H or C₁-C₃ alkyl.

[0063] In embodiments, R¹ is C₂-C₈ alkyl. In embodiments, R² is H. In embodiments, X is C=O or






where R^4 is CH_3 . In embodiments, when Y is $\text{CR}^5=\text{CH}$, R^5 is H . In embodiments, when Y is CHR^5-CH_2 , R^5 is OH or $-\text{OCH}_3$. In embodiments, R^6 is CH_3 .

GROUP B COMPOUNDS (in accordance with claim 1, the food product comprises a compound of formula B3)


Group B1 Compounds

[0064] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:





where:








R^1 and R^2 are each independently OH or $\text{C}_1\text{-C}_3$ alkoxy or where R^1 and R^2 together with the carbons to which they are bound form a five-membered ring having two oxygen heteroatoms to form a compound having the following structure




and

R^3 is






O




or



Group B2 Compounds


[0065] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where R^1 is $C_{10}-C_{15}$ alkyl or alkenyl.

Group B3 Compounds (these compounds are used in the food product of claim 1)

[0066] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:




where:

R^1 and R^2 are each independently OH or C_1-C_3 alkoxy, and


R^3 is selected from the group consisting of

1. (i) $C_{10}-C_{20}$ unsubstituted straight or branched chain alkenyl with one or more double bonds; and

2. (ii)




where X , Y and Z are independently CH , CH_2 , CO or $CHOR^5$ where R^5 is H or C_1-C_3 alkyl, provided that if one of X or Y are CH then Z is also CH , and where R^4 is C_1-C_8 straight or branched chain unsubstituted alkyl or




where n is 1-5, provided that if R^4 is C_1-C_8 straight or branched chain alkyl, then Y is $CHOR^5$ where R^5 is C_1-C_3 alkyl.

[0067] In embodiments, R^1 and R^2 are OH . In such embodiments, R^1 may be substituted at the 5 position, R^2 may be substituted at the 3 position, and R^3 may be substituted at the 1 position.

[0068] In embodiments, R^3 is $C_{10}-C_{20}$ unsubstituted straight or branched chain alkenyl with one or more double bonds. In such embodiments, R^3 may be $C_{12}-C_{18}$ unsubstituted straight or branched chain alkenyl. For example, R^3 may be $C_{13}-C_{17}$ unsubstituted straight or branched chain alkenyl, such as C_{15} unsubstituted straight or branched chain alkenyl. In embodiments, R^3 has 1-5 double bonds. For example, R^3 may have 1-4 double bonds, such as 1-3 double bonds. In embodiments, R^3 is a straight chain alkenyl. By way of example, R^3 may be




or




[0069] In embodiments, R^1 is C_1-C_3 alkoxy, such as methoxy, and R^2 is OH . In such embodiments, R^1 may be substituted at the 3 position, R^2 may be substituted at the 4 position,

and R³ may substituted at the 1 position.



[0070] In embodiments, R³ is




In such embodiments, R⁴ may be



In embodiments, n is 2. In embodiments, Z is CH or CH₂, X and Z are each independently CH or CH₂, depending on whether Z is CH or CH₂. In embodiments, X is CHOR⁵. R⁵ may be H. In embodiments, Y is CO. In embodiments, X is CO. By way of example, R³ may be




or








[0071] In embodiments, Y is CHOR⁵. R⁵ may be C₁-C₃ alkyl. For example, R⁵ may be methyl. In some embodiments, where Y is CHOR⁵, R⁴ is C₁-C₈ straight or branched chain alkyl. For example, R⁴ may be C₄-C₆ straight or branched chain alkyl, such as C₅ straight or branched chain alkyl. IN embodiments, R⁴ is a straight chain alkyl.

[0072] In embodiments, R³ is






[0073] In embodiments, R^3 is selected from the group consisting of:




and



Group B4 Compounds

[0074] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where:

A and B are each independently NCH₃ or C(O), with the proviso that one of A and B is NCH₃ and the other of A and B is C(O); and

R₁ and R₂ are independently selected from H or C₈-C₁₆ unsaturated alkyl, with the proviso that if one of R₁ and R₂ is H, then the other of R₁ and R₂ is C₈-C₁₆ unsaturated alkyl.


[0075] In embodiments, R₁ and R₂ are independently selected from H or C₈-C₁₆ unsaturated alkyl, wherein the C₈-C₁₆ unsaturated alkyl contains 1 to 3 double bonds. In embodiments, R₁ and R₂ are independently selected from H or C₈-C₁₆ unsaturated alkyl, wherein the C₈-C₁₆ unsaturated alkyl contains only 1 double bond.

[0076] In embodiments, B is NCH₃ and R₂ is C₈-C₁₆ unsaturated alkyl.

[0077] In embodiments, R₁ and R₂ are independently selected from H or C₁₁-C₁₅ unsaturated alkyl, such as C₁₃ unsaturated alkyl.

GROUP C COMPOUNDS

[0078] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where:

X is C or N;

R¹ is H, OH, C₁-C₃ alkoxy, or C₁-C₆ alkyl;

R² and R³ are each independently selected from H; OH; C₁-C₃ alkoxy; straight or branched chain, saturated or unsaturated C₁-C₆ alkyl or alkenyl; or R¹ and R² together with the carbons to which they are bound form a part of a five or six membered ring structure;

R⁴ is H or C₁-C₃ alkyl;

R⁵ is H, OH, C₁-C₃ alkoxy or C₁-C₃ alkyl;


A and B are each independently selected from CH, C=O, C-benzyl methoxy, C-CH₂-R⁶ or C-C(O)R⁶ where R⁶ is straight or branched chain, saturated or unsaturated C₁-C₆ alkyl, or A and B together are part of an aromatic six-membered ring structure sharing a side with the remainder of the structure of Formula (C); and


Y is O, CH, C=O, or C-O-R⁷, where R⁷ is H or C₁-C₃ alkyl.

[0079] In embodiments, when X is N, R⁵ is C₁-C₃ alkyl, such as methyl. In embodiments, when X is N, Y is C=O or C-O-R⁷, such as C-O-Me. In embodiments when X is C, R⁵ is H or OH. In embodiments when X is C, Y is O. In embodiments, R¹ is H or methoxy. In embodiments, one of A or B is C=O and the other is H, C-benzyl methoxy, C-CH₂CHC(CH₃)₂ or C-C(O)CHC(CH₃)₂.

[0080] In embodiments, R² and R³ together with the carbons to which they are bound form a part of a six membered ring structure. In embodiments, the six membered ring structure includes an oxygen or nitrogen heteroatom. In embodiments, the six membered ring structure contains one or more carbon atoms substituted with one or more C₁-C₆ alkyl, such as methyl. In embodiments, one carbon atom of the ring structure is substituted with two methyl groups. In embodiments, the ring structure is an unsubstituted six carbon aromatic ring structure.


[0081] In embodiments, a compound according to Formula (C) has the following structure:





where A, B, X, Y, R¹, R⁴ and R⁵ are as described above for Formula (C).

[0082] In embodiments, a compound according to Formula (C) has the following structure:



where:

R¹, R² and R³ are each independently H, OH or C₁-C₃ alkoxy;

R⁴ and R⁵ are each independently H or C₁-C₃ alkyl;

Y is C=O or C-O-R⁷, where R⁷ is H or C₁-C₃ alkyl; and

A is C-CH₂-R⁶ or C-C(O)R⁶ where R⁶ is straight or branched chain, saturated or partially unsaturated C₁-C₆ alkyl, B is C=O, or A and B together are part of an aromatic six-carbon membered ring structure sharing a side with the remainder of the structure of Formula (C'').


[0083] In embodiments of a compound according to Formula (C''), R¹, R² and R³ are H.


[0084] In embodiments, A is C-CH₂-R⁶ or C-C(O)R⁶ where R⁶ is straight or branched chain, saturated or partially unsaturated C₁-C₆ alkyl and B is C=O.

[0085] In embodiments, R¹, R² and R³ are each independently OH or C₁-C₃ alkoxy. In some embodiments, R¹, R² and R³ are the same.

[0086] In embodiments, A and B together are part of an aromatic six-carbon membered unsubstituted ring structure.

[0087] In embodiments, a compound according to Formula (C) has the following structure:





where:

R¹, R² and R³ are each independently H, OH or C₁-C₃ alkoxy;


R⁴ and R⁵ are each independently H or C₁-C₃ alkyl;

Y is C=O or C-O-R⁷, where R⁷ is H or C₁-C₃ alkyl; and

A is C-CH₂-R⁶ or C-C(O)R⁶ where R⁶ is straight or branched chain, saturated or partially unsaturated C₁-C₆ alkyl, B is C=O, or A and B together are part of an aromatic six-carbon membered ring structure sharing a side with the remainder of the structure of Formula (C'').

[0088] In embodiments of a compound according to Formula (C''), R¹, R² and R³ are H. In embodiments, A is C-CH₂-R⁶ or C-C(O)R⁶ where R⁶ is straight or branched chain, saturated or partially unsaturated C₁-C₆ alkyl and B is C=O. In embodiments, R¹, R² and R³ are each independently OH or C₁-C₃ alkoxy. In embodiments, R¹, R² and R³ are the same. In embodiments, A and B together are part of an aromatic six-carbon membered unsubstituted ring structure.

[0089] In embodiments, a compound according to Formula (C) has the following structure:



where:

X is N;

R¹ is H or C₁-C₃ alkyl;

R^2 and R^3 are each independently selected from H and straight or branched chain, saturated or unsaturated C_1 - C_3 alkyl or alkenyl;

R^4 is H or C_1 - C_3 alkyl;

R^5 is H or C_1 - C_3 alkyl;


A and B are each independently selected from $C=O$ and $C-C(O)R^6$ where R^6 is straight or branched chain, saturated or partially unsaturated C_1 - C_6 alkyl; and

Y is $C-O-R^7$, where R^7 is H or C_1 - C_3 alkyl.

[0090] In embodiments of a compound according to Formula (C''), one or more of R^1 , R^2 and R^3 are H. In embodiments, each of R^1 , R^2 and R^3 are H. In embodiments, R^4 and R^5 are independently C_1 - C_3 alkyl. In embodiments, R^4 and R^5 are methyl. In embodiments, wherein A is $C-C(O)R^6$. In embodiments, B is $C=O$.

GROUP D COMPOUNDS

[0091] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where:


R^1 is H, methyl, $OCOCH_3$ or forms together with R^6 a five membered ring structure in which R^1 and R^6 together are $C=O$ or CH_2 ;

R^6 is H, $C=OCH_3$, or together forms a five membered ring structure in which R^1 and R^6 together are $C=O$ or CH_2 ;

R^7 is $OCOCH_3$;


A and B are C, CH or CCH_3 , wherein when A and B are both C a double bond is formed between A and B; and

R^4 and R^5 are independently selected from OH, methyl,




or

R^4 and R^5 together with the carbon to which they are bound form




to form a compound of the following formula



where R^1 , R^6 and R^7 are as described above.

GROUP E COMPOUNDS


[0092] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where R^1 , R^2 and R^3 are independently selected from the group consisting of H and $COCH_3$.

GROUP F COMPOUNDS

[0093] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where:

R^{42} is H or OH;

R^{43} are R^{44} each independently H or CH_3 ;

R^1 and R^2 are each independently OH, C_1-C_3 alkyl or C_1-C_3 hydroxyl;

R^3 and R^4 are each independently H, OH or C_1-C_3 hydroxyl;

R^5 is H or $-OC(O)R^8$, where R^8 is C_1-C_8 straight or branched chain, saturated or unsaturated

alkyl;

W is CH;


Y is CH, X is CH or CH₂, and Z is C or CR⁹, provided that Z is C when X is CH₂, where R⁹ is H or together with R⁶ and the carbons to which they are bound and the intervening carbon of W form a five membered ring with an oxygen heteroatom;

R⁶ is C₁-C₃ hydroxyl, C(O)R¹⁰, (CH₂)_pR¹⁰ where p' is zero or 1, or together with R⁹ and the carbons to which they are bound and the intervening carbon of W form a five membered ring with an oxygen heteroatom,

where R¹⁰ is H, OH, or saccharidyl; and


R⁷ is H, OH, C₁-C₃ hydroxyl, or (CH₂)_pR⁴⁴, where p is zero or 1 and R⁴⁴ is saccharidyl.

[0094] In embodiments, R¹⁰ is saccharidyl and is



where a and b are each independently zero or 1;

where R¹¹ is H or




where d is zero or 1,

R¹⁴ is H, OH, CH₃ or C₁-C₃ hydroxyl, and

R¹⁵ is H, C(O)R¹⁶ where R¹⁶ is C₁-C₁₅ straight or branched chain, saturated or unsaturated alkyl unsubstituted or substituted with hydroxyl,

where R^{12} and R^{13} are each independently H, $C(O)R^{41}$ where R^{41} is C_1-C_{15} straight or branched chain, saturated or unsaturated alkyl unsubstituted or substituted with hydroxyl, or




where e is zero or 1,

R^{17} is H, OH or CH_3 , and

R^{18} is H, $C(O)R^{42}$ where R^{42} is C_1-C_{15} straight or branched chain, saturated or unsaturated alkyl unsubstituted

or substituted with hydroxyl, or




where f is zero or 1,

R^{19} is H, OH or CH_3 , and

R^{20} and R^{21} are each independently H, $C(O)R^{43}$

where R^{43} is C_1-C_{15} straight or branched chain, saturated or unsaturated alkyl unsubstituted or substituted with hydroxyl,

or




where

g is zero or 1, and

R^{22} , R^{23} and R^{24} are each independently H, OH or CH_3 .


[0095] In embodiments, R^{44} is saccharidyl and is



where

i is zero or 1,


R^{25} , R^{26} , R^{27} and R^{28} are each independently H, OH, CH_3 , C_1 - C_3 hydroxyl, COOH, $(CH_2)_nOC(O)R^{29}$ where n is zero, 1, 2 or 3 and R^{29} is H or C_1 - C_3 alkyl, or $(CH_2)_mOR^{30}$ where m is zero or 1 and R^{30} is



where

j is zero or 1,

R^{31} , R^{32} , R^{33} and R^{34} are each independently H, OH, CH_3 , C_1 - C_3 hydroxyl, COOH, $(CH_2)_qOC(O)R^{35}$ where q is zero, 1, 2 or 3 and R^{35} is H or C_1 - C_3 alkyl, or $(CH_2)_tOR^{36}$ where t is zero or 1 and R^{36} is




where


k is zero or 1, and

R^{37} , R^{38} , R^{39} and R^{40} are each independently H, OH, CH_3 , C_1-C_3 hydroxyl, $COOH$, $(CH_2)_uOC(O)R^{41}$ where u is zero, 1, 2 or 3 and R^{41} is H or C_1-C_3 alkyl.

[0096] In embodiments, Z is C. In embodiments, Z is CR^9 and wherein R^9 together with R^6 and the carbons to which they are bound and the intervening carbon of W for a five membered ring with an oxygen heteroatom. In embodiments, R^6 is $C(O)R^{10}$. In embodiments, R^{10} is H or OH. In embodiments, R^7 is OH. In embodiments, R^1 and R^2 are each independently C_1-C_3 alkyl or C_1-C_3 hydroxyl. In embodiments, R^3 is H or OH. In embodiments, R^4 is H. In embodiments, R^5 is H. In embodiments, one or more of R^{16} , if present, R^{42} and R^{43} are



[0097] In embodiments, a compound according to Formula (F) has the following structure:



where:

R^1 and R^2 are each independently OH, C_1-C_3 alkyl or C_1-C_3 hydroxyl;

R^3 and R^4 are each independently H, OH or C_1-C_3 hydroxyl;

R^5 is H;

W is CH;

Y is CH, X is CH or CH_2 , and Z is C or CR^9 , provided that Z is C when X is CH_2 , where R^9 is H


or together with R⁶ and the carbons to which they are bound and the intervening carbon of W form a five membered ring with an oxygen heteroatom;

R⁶ is C₁-C₃ hydroxyl, C(O)R¹⁰, (CH₂)_pR¹⁰ where p' is zero or 1, or together with R⁹ and the carbons to which they are bound and the intervening carbon of W form a five membered ring with an oxygen heteroatom,

where R¹⁰ is H or saccharidyl; and


R⁷ is H, OH, C₁-C₃ hydroxyl, or (CH₂)_pOR⁴⁴, where p is zero or 1 and R⁴⁴ is saccharidyl.

[0098] In embodiments of a compound according to **Formula F'**, R¹⁰ is saccharidyl and is



where a and b are each independently zero or 1;

R¹¹ is H or





where d is zero or 1,

R¹⁴ is H, OH, CH₃ or C₁-C₃ hydroxyl, and

R¹⁵ is H, C(O)R¹⁶ where R¹⁶ is C₁-C₁₅ straight or branched chain, saturated or unsaturated alkyl unsubstituted or substituted with hydroxyl,

where R¹² and R¹³ are each independently H, C(O)R⁴¹ where R⁴¹ is C₁-C₁₅ straight or branched chain, saturated or unsaturated alkyl unsubstituted or substituted with hydroxyl, or






where e is zero or 1,

R¹⁷ is H, OH or CH₃ and

R¹⁸ is H, C(O)R⁴² where R⁴² is C₁-C₁₅ straight or branched chain, saturated or unsaturated alkyl unsubstituted


or substituted with hydroxyl, or



where f is zero or 1,

R¹⁹ is H, OH or CH₃, and

R²⁰ and R²¹ are each independently H, C(O)R⁴³ where R⁴³ is C₁-C₁₅ straight or branched chain, saturated or unsaturated alkyl unsubstituted or substituted with hydroxyl, or




where

g is zero or 1, and

R²², R²³ and R²⁴ are each independently H, OH or CH₃.

[0099] In embodiments of a compound according to **Formula F'**, R⁴⁴ is saccharidyl and is






where

i is zero or 1, and


R²⁵, R²⁶, R²⁷ and R²⁸ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_nOC(O)R²⁹ where n is zero, 1, 2 or 3 and R²⁹ is H or C₁-C₃ alkyl, or (CH₂)_mOR³⁰ where m is zero or 1 and R³⁰ is



where

j is zero or 1, and

R³¹, R³², R³³ and R³⁴ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_qOC(O)R³⁵ where q is zero, 1, 2 or 3 and R³⁵ is H or C₁-C₃ alkyl, or (CH₂)_tOR³⁶ where t is zero or 1 and R³⁶ is




where


k is zero or 1, and

R³⁷, R³⁸, R³⁹ and R⁴⁰ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_uOC(O)R⁴¹ where u is zero, 1, 2 or 3 and R⁴¹ is H or C₁-C₃ alkyl.

[0100] In embodiments of a compound according to **Formula F'**, Z is C. In embodiments, Z is CR⁹ and wherein R⁹ together with R⁶ and the carbons to which they are bound and the intervening carbon of W for a five membered ring with an oxygen heteroatom. In embodiments, R⁶ is C(O)R¹⁰. In embodiments, R¹ and R² are each independently C₁-C₃ alkyl or C₁-C₃ hydroxyl. In embodiments, R³ is H or OH. In embodiments, R⁴ is H. In embodiments, R¹⁶, if present, is



[0101] In embodiments, a compound according to Formula (F) has the following structure:



where:

R¹ and R² are each independently OH, C₁-C₃ alkyl or C₁-C₃ hydroxyl;

\mathbb{R}^3 is H ;

R^4 is H, OH or C₁-C₃ hydroxyl;


\mathbb{R}^5 is H :

W is CH;

Y is CH, X is CH or CH_2 , and Z is C or CR^9 , provided that Z is C when X is CH_2 , where R^9 is H or together with R^6 and the carbons to which they are bound and the intervening carbon of W form a five membered ring with an oxygen heteroatom;


R^6 is, together with R^9 and the carbons to which they are bound and the intervening carbon of W form a five membered ring with an oxygen heteroatom, or $C(O)R^{10}$

where R^{10} is H or



where a and b are each independently zero or 1;

where R¹¹ is H, OH, or




where d is zero or 1,

R¹⁴ is H, OH, CH₃ or C₁-C₃ hydroxyl, and

R¹⁵ is H, C(O)R¹⁶ where R¹⁶ is C₁-C₁₅ straight or branched chain, saturated or unsaturated alkyl unsubstituted or substituted with hydroxyl,

where R¹² and R¹³ are each independently H, C(O)R⁴¹ where R⁴¹ is C₁-C₁₅ straight or branched chain, saturated or unsaturated alkyl unsubstituted or substituted with hydroxyl, or




where e is zero or 1,

R¹⁷ is H, OH or CH₃ and

R¹⁸ is H, C(O)R⁴² where R⁴² is C₁-C₁₅ straight or branched chain, saturated or unsaturated alkyl unsubstituted or substituted with hydroxyl, or






where f is zero or 1,

R¹⁹ is H, OH or CH₃, and

R²⁰ and R²¹ are each independently H, C(O)R⁴³ where R⁴³ is C₁-C₁₅ straight or branched chain, saturated or unsaturated alkyl unsubstituted or substituted with hydroxyl, or




where

g is zero or 1, and

R²², R²³ and R²⁴ are each independently H, OH or CH₃; and

R⁷ is H, OH, C₁-C₃ hydroxyl, or (CH₂)_pR⁴⁴, where p is zero or 1 and R⁴⁴ is




where

i is zero or 1, and

R²⁵, R²⁶, R²⁷ and R²⁸ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_nOC(O)R²⁹ where n is zero, 1, 2 or 3 and R²⁹ is H or C₁-C₃ alkyl, or (CH₂)_mOR³⁰ where


m is zero or 1 and R³⁰ is



where

j is zero or 1, and

R³¹, R³², R³³ and R³⁴ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_qOC(O)R³⁵ where q is zero, 1, 2 or 3 and R³⁵ is H or C₁-C₃ alkyl, or (CH₂)_tOR³⁶ where t is zero or 1 and R³⁶ is



where


k is zero or 1, and


R³⁷, R³⁸, R³⁹ and R⁴⁰ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_uOC(O)R⁴¹ where u is zero, 1, 2 or 3 and R⁴¹ is H or C₁-C₃ alkyl.

[0102] In embodiments of a compound according to Formula (F''), R⁴ is H. In embodiments, R¹ is CH₃. In embodiments, R² is CH₃.

GROUP G COMPOUND


[0103] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:





GROUP H COMPOUNDS

[0104] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where:

R^1 is H or saccharidyl;


R^2 is H or OH;


R^3 and R^4 are independently selected from H or methyl or together form CH_2 ;

R^5 is CH_2 or CH;


R^6 is CH or C, provided that when R^5 is CH, R^6 is C;

R^7 and R^8 together with the carbons to which they are bound form








or




to form a compound having the following structure

 (H') , (H'') ,

or






and


R⁹, R¹⁰, and R¹¹ are independently H or methyl.

[0105] In embodiments, R¹ is saccharidyl and is



GROUP I COMPOUNDS

[0106] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where:

X is O or CH₂;

R¹ and R⁹ are independently selected from H, OH, C₁-C₃ alkoxy, C₁-C₃ alkyl, and (CH₂)_nOH, where n is an integer from 1 to 3;

R² and R¹⁰ are independently selected from H, OH and C₁-C₃ alkyl;

R³ and R⁴ are independently selected from H and C₁-C₃ alkyl or together form CH₂;


R^5 , R^6 and R^7 are independently selected from H and C₁-C₃ alkoxy; and

R^8 is H, OH or C₁-C₃ alkoxy.

[0107] In embodiments, R^5 , R^6 , R^7 and R^8 are independently C₁-C₃ alkoxy. In embodiments, R^5 , R^6 , R^7 and R^8 are methoxy. In embodiments, R^3 and R^4 together form CH₂. In embodiments, R^3 and R^4 are independently C₁-C₃ alkyl. In embodiments, R^3 and R^4 are methyl. In embodiments, at least two of R^1 , R^2 , R^9 and R^{10} are independently C₁-C₃ alkyl. In embodiments, at least two of R^1 , R^2 , R^9 and R^{10} are methyl. In embodiments, X is CH₂.

[0108] In embodiments, X is O. In some embodiments where X is O, R^2 and R^{10} are H. In some embodiments where X is O, two or more of R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^{10} are H. In some embodiments where X is O, each of R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^{10} are H. In some embodiments where X is O, R^8 is OH or C₁-C₃ alkoxy. In some embodiments where X is O, R^1 and R^9 are independently selected from OH, C₁-C₃ alkoxy, and (CH₂)_nOH. In some embodiments where X is O, one of R^1 and R^9 is selected from OH and C₁-C₃ alkoxy, and the other of R^1 and R^9 is (CH₂)_nOH.

[0109] In embodiments, a compound according to Formula (I) has the following structure:



where:

X is O;

R^1 and R^9 are independently selected from H, OH, C₁-C₃ alkoxy, C₁-C₃ alkyl, and (CH₂)_nOH, where n is an integer from 1 to 3;

R^2 and R^{10} are independently selected from H, OH and C₁-C₃ alkyl;

R^3 and R^4 are independently selected from H and C₁-C₃ alkyl;

R^5 , R^6 and R^7 are H; and


R^8 is H, OH or C₁-C₃ alkoxy.

[0110] In embodiments of a compound according to Formula (I'), R^2 and R^{10} are H. In embodiments, each of R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^{10} are H. In embodiments, R^8 is OH or C₁-C₃ alkoxy. In embodiments, R^1 and R^9 are independently selected from OH, C₁-C₃ alkoxy, and (CH₂)_nOH. In embodiments, one of R^1 and R^9 is selected from OH and C₁-C₃ alkoxy, and the other of R^1 and R^9 is (CH₂)_nOH.

GROUP J COMPOUNDS

Group J1 Compounds

[0111] In embodiments, a bioactive, taste modulating, or salty taste modulating is a compound having the following structure:



where:


R^1 and R^6 are each independently OH, C₁-C₃ hydroxyl, C₁-C₃ alkoxy or saccharidyl;

R^2 , R^3 , R^7 and R^8 are each independently H, OH, C₁-C₃ hydroxyl, C₁-C₃ alkoxy;

R^4 and R^5 are each independently H, OH, C₁-C₃ alkyl, C₁-C₃ alkoxy, or C₁-C₃ hydroxyl;

Y is CHCH and n is zero or 1; and


X is CHR⁹ and m is zero or 1, where R⁹ is




where R¹⁰, R¹¹ and R¹² are each independently OH, C₁-C₃ alkoxy, or C₁-C₃ hydroxyl,

provided that if n is 1, then m is 1.

[0112] In embodiments of a compound according to **Formula J1**, R¹, R⁶, or R¹ and R⁶ are saccharidyl and are




[0113] In embodiments of a compound according to **Formula J1**, R¹ is OH. In embodiments, R², R³, R⁷ and R⁸ are each independently H or C₁-C₃ alkoxy. In embodiments, where one of wherein R² and R³ is H and the other is C₁-C₃ alkoxy, one of R⁷ and R⁸ is H and the other is C₁-C₃ alkoxy. In embodiments, R², R³, R⁷ and R⁸ are each H. In embodiments, R¹⁰ is present and is OH. In embodiments, R¹¹ is present and is C₁-C₃ alkoxy. In embodiments, R¹² is present and is OH. In embodiments, R⁹ is present and is



where R¹⁰,


R^{11} , and R^{12} are as defined above.

[0114] In embodiments of a compound according to **Formula J1**, the compound has the following structure:




where R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , and R^9 are as defined above.

[0115] In embodiments of a compound according to **Formula J1**, the compound has the following structure:



where R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 are as defined above.

[0116] In embodiments of a compound according to **Formula J1**, the compound has the following structure:



where:


R^1 and R^6 are each independently OH, C₁-C₃ hydroxyl, C₁-C₃ alkoxy or saccharidyl;

R^2 , R^3 , R^7 and R^8 are each independently H, OH, C₁-C₃ hydroxyl, C₁-C₃ alkoxy;

R^4 and R^5 are each independently H, OH, C₁-C₃ alkoxy, or C₁-C₃ hydroxyl;

Y is CHCH and n is zero or 1; and


X is CHR^9 and m is zero or 1, where R^9 is




where R^{10} , R^{11} and R^{12} are each independently OH, C₁-C₃ alkoxy, or C₁-C₃ hydroxyl,

provided that if n is 1, then m is 1.

[0117] In embodiments, R^1 , R^6 or R^1 and R^6 are saccharidyl and are




In embodiments, R^1 is OH. In embodiments, R^2 , R^3 , R^7 and R^8 are each independently H or C₁-C₃ alkoxy. In embodiments, R^2 and R^3 is H and the other is C₁-C₃ alkoxy, and one of R^7 and R^8 is H and the other is C₁-C₃ alkoxy. In embodiments, R^2 , R^3 , R^7 and R^8 are each H. In embodiments, R^{10} is present and is OH. In embodiments, R^{11} is present and is C₁-C₃ alkoxy. In embodiments, R^{12} is present and is OH. In embodiments, R^9 is present and is



where R^{10} , R^{11} , and R^{12} are as defined above.

Group J2 Compounds

[0118] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:




where

R^1 is H, OH, or saccharidyl;


R² and R³ are each independently H or C₁-C₃ alkyl;

R^4 and R^5 are each independently H ,



or $\text{CHC}(\text{R}^{10})\text{R}^{11}$, where

R^{10} and R^{11} are each independently H or C₁-C₅ alkyl unsubstituted or substituted with one or more of OH and



X is $C \equiv \Omega$ or Ω .


Y is C≡O or O, provided that when X is O, Y is C≡O, or when X is C≡O, Y is O.

A is CHR^{12} or CR^{12} , where

R¹² is H, OH, or saccharidyl:

B is C or CH, provided that if B is C then A is CR¹².

[0119] In embodiments of a compound according to **Formula J2**, R¹ is saccharidyl and is




where

a is zero or 1, and

R⁶, R⁷, R⁸, and R⁹ are each independently H, OH, CH₃, or CH₂OH.

[0120] In embodiments of a compound according to **Formula J2**, R¹² is saccharidyl and is




where


b is zero or 1, and

R¹³, R¹⁴, R¹⁵, and R¹⁶ are each independently H, OH, CH₃, or CH₂OH.

[0121] In embodiments of a compound according to **Formula J2**, R⁴ is CHC(CH₃)₂ or




[0122] In embodiments of a compound according to **Formula J2**, R⁵ is H or




GROUP K COMPOUND

[0123] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



GROUP L COMPOUNDS


[0124] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:




where:

R¹, R³, R⁴, R⁵ and R⁶ are each independently H, OH, C₁-C₃ hydroxyl or C₁-C₃ alkoxy; and


R² is H, C₁-C₃ alkyl, or





where R⁷, R⁸ and R⁹ are each independently OH, C₁-C₃ hydroxyl or C₁-C₃ alkoxy.

[0125] In embodiments, R² is C₁-C₃ alkyl, such as methyl. In embodiments, R⁵ is OH or C₁-C₃ hydroxyl. In embodiments, R³ is H. In embodiments, R⁴ is C₁-C₃ alkoxy, such as methoxy. In embodiments, R⁶ is C₁-C₃ alkyl, such as methyl or propyl. In embodiments, R¹ is H. In embodiments, R¹ is C₁-C₃ hydroxyl, such as C₂ hydroxyl. In embodiments, R² is



In embodiments, R⁷, R⁸ and R⁹ are each independently OH or C₁-C₃ alkoxy. In embodiments, R⁷ is OH and R⁸ and R⁹ are each independently C₁-C₃ alkoxy, such as methoxy.

[0126] In embodiments, R³ is OH, C₁-C₃ hydroxyl, or C₁-C₃ alkoxy.


[0127] In embodiments, R⁵ is H.


[0128] In embodiments, R⁶ is C₁-C₃ alkoxy, such as methoxy.

[0129] In embodiments, R¹ is C₁-C₃ alkoxy, such as methoxy.


[0130] In embodiments, R⁴ is H.

[0131] In embodiments of a compound according to **Formula L**, (i) R¹ is OH, C₁-C₃ hydroxyl or C₁-C₃ alkoxy; (ii) R² is H, C₁-C₃ alkyl, or







where R⁷, R⁸ and R⁹ are each independently OH, C₁-C₃ hydroxyl or C₁-C₃ alkoxy; and (iii) R³, R⁴, R⁵ and R⁶ are each independently H, OH, C₁-C₃ hydroxyl or C₁-C₃ alkoxy. In embodiments, R² is C₁-C₃ alkyl, such as methyl. In embodiments, R⁵ is OH or C₁-C₃ hydroxyl. In embodiments, R³ is H. In embodiments, R⁴ is C₁-C₃ alkoxy, such as methoxy. In embodiments, R⁶ is C₁-C₃ alkyl, such as butyl. In embodiments, R¹ is C₁-C₃ hydroxyl. In some embodiments where R² is




R⁷, R⁸ and R⁹ may each independently be OH or C₁-C₃ alkoxy. For example, R⁷ may be OH and R⁸ and R⁹ may each independently be C₁-C₃ alkoxy, such as methoxy. In some embodiments where R² is



R³ is OH, C₁-C₃ hydroxyl, or C₁-C₃ alkoxy. In some embodiments where R² is




R² is H. In some embodiments where R² is




I^7 ,

R^6 is $\text{C}_1\text{-}\text{C}_3$ alkoxy, such as methoxy. In some embodiments where R^2 is




R^1 is $\text{C}_1\text{-}\text{C}_3$ alkoxy, such as methoxy. In some embodiments where R^2 is



R^4 is H.

[0132] In embodiments, a compound according to **Formula L** is a compound having the following structure:



where:

R^1 , R^3 , R^4 , R^5 and R^6 are each independently H, OH, $\text{C}_1\text{-}\text{C}_3$ hydroxyl or $\text{C}_1\text{-}\text{C}_3$ alkoxy; and


R^7 , R^8 and R^9 are each independently OH, $\text{C}_1\text{-}\text{C}_3$ hydroxyl or $\text{C}_1\text{-}\text{C}_3$ alkoxy.

[0133] In embodiments of a compounds according to **Formula L'**, R^7 , R^8 and R^9 are each independently OH or $\text{C}_1\text{-}\text{C}_3$ alkoxy. In embodiments, R^7 is OH and R^8 and R^9 are each independently $\text{C}_1\text{-}\text{C}_3$ alkoxy, such as methoxy. In embodiments, R^3 is OH, $\text{C}_1\text{-}\text{C}_3$ hydroxyl, or $\text{C}_1\text{-}\text{C}_3$ alkoxy. In embodiments, R^5 is H. In embodiments, R^6 is $\text{C}_1\text{-}\text{C}_3$ alkoxy, such as methoxy.

In embodiments, R¹ is C₁-C₃ alkoxy, such as methoxy. In embodiments, R⁴ is H.


GROUP M COMPOUND

[0134] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:




GROUP N COMPOUND


[0135] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



GROUP O COMPOUND

[0136] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:






GROUP P COMPOUNDS

Group P1 Compounds

[0137] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:




where:

X is CH or CH_2 ;

Y is CH or C, with the caveat that if Y is C, then X is CH;

R^1 is H, or



where R^5 is OH, $\text{C}(\text{O})\text{OH}$, or $\text{OC}(\text{O})\text{R}^6$, where R^6 is a $\text{C}_1\text{-C}_3$ alkyl;

R^2 and R^3 are independently H, or substituted with hydroxyl or unsubstituted saturated or unsaturated branched or straight $\text{C}_1\text{-C}_6$ alkoxy; and

R^4 is H, or a saturated or unsaturated branched or straight $\text{C}_1\text{-C}_6$ alkyl.


[0138] In embodiments, at least one of R^2 or R^3 is H. In embodiments, both R^2 and R^3 are H.

In embodiments, at least one of R² and R³ is a substituted saturated or unsaturated branched or straight C₁-C₆ alkoxy substituted with at least one hydroxyl group. In embodiments, both R² and R³ are independently selected from a substituted saturated or unsaturated branched or straight C₁-C₆ alkoxy substituted with at least one hydroxyl group. For example, at least one of R² and R³ can be methoxy. In embodiments, at least one of R² and R³ is unsaturated. In embodiments, at least one of R² and R³ is branched. In embodiments, R² is a straight unsubstituted saturated C₁-C₃ alkoxy; and R³ is a branched unsubstituted, unsaturated C₃-C₆ alkoxy.

[0139] In embodiments, R⁴ is saturated or unsaturated branched or straight C₁-C₆ alkyl. For example, R⁴ can be C₅ alkyl. In embodiments, R⁴ is unsaturated. In embodiments, R⁴ is CH(CH₃)₂CHCH₂.

Group P2 compounds


[0140] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:




where

X is selected from NH, O, C(O), CR⁵ or CR⁵R⁶, where R⁵ and R⁶ are independently selected from H, OH, a straight or branched C₁ to C₆ hydroxyl, or a C₁ to C₆ alkyl; and

Y, and Z are independently selected from O, CR⁵, CR⁵R⁶, where R⁵ and R⁶ are independently selected from H, OH, a straight or branched C₁ to C₆ hydroxyl, or a C₁ to C₆ alkyl, or






,


with the proviso that (i) no more than two of X, Y and Z are O, and (ii) if Y is CR⁵ then X or Z is also CR⁵;

R¹ is H, C₁-C₃ alkoxy, C(O)R⁷, where R⁷ is a C₁ to C₆ straight or branched alkyl, or




where R⁸, R⁹ and R¹⁰ are independently selected from: H, OH, C₁ to C₃ hydroxyl, C₁ to C₃ alkoxy, or C₃ to C₆ alkyl;

R² is H, OH, or



where R¹¹ is H, OH, or C₁ to C₃ hydroxyl;

R³ is H, C₁-C₃ alkoxy, or together with R⁴ form a six membered ring having an oxygen heteroatom to form a compound having a structure of:




R⁴ is H, C₁ to C₃ alkoxy, or together with R³ forms the structure above.

[0141] In embodiments, X is CR^5R^6 . In embodiments, X is $C(OH)(CH_2CH_2CH_2CH_3)$. In embodiments, X is O. In embodiments, X is CH_2 .

[0142] In embodiments, Y is O. In embodiments, Y is CR^5R^6 , where R^5 and R^6 are H and a branched C₁ to C₆ hydroxyl, such as $C(CH_3)(CH_3)OH$. In embodiments, Y is CH.

[0143] In embodiments, Z is C(O). In embodiments, Z is CH_2 . In embodiments, Z is O.


[0144] In embodiments, R^1 is




In embodiments, R^1 is a C₁-C₃ alkoxy. In embodiments, R is $C(O)R$, where R^7 is a C₁ to C₆ straight or branched alkyl.

[0145] In embodiments, R^7 is a branched C₃ alkyl.

[0146] In embodiments, R^2 is H. In embodiments, R^2 is OH. In embodiments, R^2 is



[0147] In embodiments, R^3 together with R^4 form a six membered ring having an oxygen heteroatom to form a compound having the structure:




$\begin{array}{c} | \\ R^1 \end{array}$
(P2').

[0148] In embodiments, R^3 is H. In embodiments, R^3 is a C_1 - C_3 alkoxy. In embodiments, R^4 is H.


[0149] In embodiments, X is NH.

[0150] In embodiments, Z is



Group P3 Compounds

[0151] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:




(P3), where

Y is independently selected from CR^5 , CR^5R^6 , where R^5 and R^6 are independently selected


from H, OH, a straight or branched C₁ to C₆ hydroxyl, or a C₁ to C₆ alkyl;

R¹ is H, C₁-C₃ alkoxy, C(O)R⁷, where R⁷ is a C₁ to C₆ straight or branched alkyl, or




where R⁸, R⁹ and R¹⁰ are independently selected from: H, OH, C₁ to C₃ hydroxyl, C₁ to C₃ alkoxy, or C₃ to C₆ alkyl;

R² is H or



where R¹¹ is H, OH, or C₁ to C₃ hydroxyl;

R³ is H, C₁-C₃ alkoxy, or together with R⁴ form a six membered ring having an oxygen heteroatom to form a compound having a structure of:




R⁴ is H, C₁ to C₃ alkoxy, or together with R³ forms the structure above.

[0152] In embodiments, R¹ is C₁ to C₃ alkoxy. In embodiments, R¹ is H. In embodiments, R² is H. In embodiments, R³ is H. In embodiments, R⁴ is H.

Group P4 Compounds

[0153] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:




where

R¹ is H or a saturated or unsaturated C₁ to C₆ alkyl;

R² is H or a C₁ to C₃ alkoxy; and


R³ is H or



where R⁴, R⁵, R⁶, R⁷, and R⁸ are independently selected from: H, OH, C₁ to C₃ hydroxyl, C₁ to C₃ alkoxy, and C₁ to C₃ alkyl; and


X is NH or O.

[0154] In embodiments, R¹ is CH₂CHC=(CH₃)₂. In embodiments, R¹ is H. In embodiments, R² is methoxy. In embodiments, R² is H. In embodiments, R³ is H. In embodiments, R³ is



Group P5 Compound


[0155] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



(P5, 59).

GROUP Q COMPOUNDS**Group Q1 Compounds**

[0156] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



(Q1)


where A is:

1. a. CR^9 where R^9 is H, saturated or unsaturated C_1 to C_6 alkyl, C_1 to C_3 hydroxyl or CHR^{10} , where R^{10} is C_1 to C_6 alkyl;

2. b. $CCR^{64}R^{65}R^{66}$, where R^{64} , R^{65} , and R^{66} are independently selected from: C_1 to C_{10} saturated or unsaturated, branched or straight chain alkyl, or $(CH_2)_aR^{11}$ where a is zero or 1 and R^{11} is H, OH, or saccharidyl,


3. c. CR^{23} , where R^{23} is saturated or unsaturated, substituted with an oxygen or unsubstituted five membered ring including at least one oxygen heteroatom,

4. d. CR^{24} , where R^{24} is



where R^{25} and R^{26} are independently selected from H, OH, or C_1 to C_6 branched or straight hydroxyl,

5. e. or along with Q and the atoms attached thereto form a ring structure of formula (Q1A) below,




where R^{27} and R^{28} are H, OH, C_1 to C_3 alkyl, C_1 to C_3 alkoxy, or C_1 to C_3 hydroxyl;

B is CH_2 , or CH, with the caveat that if B is CH then D is C;

D, E, F, and G are independently selected from: C, CR^{29} , where R^{29} is H, OH, C_1 to C_3 alkyl, or C_1 to C_3 hydroxyl, with the caveat that if D is C, then B is CH; if E is C, then G is C; and if F is C, then u is zero;

J is $C(R^{30})_vR^{31}$, where v is zero or 1 and R^{30} and R^{31} are independently selected from H, OH, C_1 to C_3 alkyl, C_1 to C_3 hydroxyl, or



where T and X are CH_2 and where f and g are independently selected from zero (0), 1, or 2, and Z is selected from $CR^{32}R^{33}$, where R^{32} and R^{33} are independently selected from H, OH, C_1 to C_3 hydroxyl, or C_1 to C_3 alkyl, with the caveat that if v is zero then k is also zero;

K is $CR^5_uR^6$, where u is zero or 1 and R^5 and R^6 are independently selected from H, OH, C_1 to C_3 alkyl, C_1 to C_3 hydroxyl, C_1 to C_3 alkoxy, with the caveat that if u is zero then F is C;

L is $CR^{34}kR^{35}$ where k is zero or 1 and R^{34} and R^{35} are independently selected from H, OH, C_1 to C_3 alkyl, C_1 to C_3 hydroxyl, with the caveat that if v is zero then k is zero;

M is $C(O)$, $COC(O)R^{36}$ where R^{36} is C_1 to C_3 alkyl, $CH(OH)$, $CH(CH_2)_jR^{37}$ where j is zero, 1, or 2 and R^{37} is H, OH, or saccharidyl;

Q is $C(O)$, or $CR^{61}R^{62}$, where R^{61} and R^{62} are independently selected from H, OH, C_1 to C_3 alkyl, C_1 to C_3 hydroxyl or along with A form a five membered ring containing an oxygen heteroatom, or structure Q1A;


R^1 , R^3 , and R^7 are independently selected from H, OH or C_1 to C_3 alkyl;

R^2 , is selected from H, O, OH or C_1 to C_3 alkyl; and

R^4 and R^8 are independently selected from H, $C(O)$, OH, C_1 to C_3 alkyl, $C(O)R^{63}$ where R^{63} is H or C_1 to C_3 alkyl, or $OC(O)R^{64}$ where R^{64} is C_1 to C_3 alkyl, or H.

[0157] In embodiments, A is CR^{23} , wherein R^{23} is saturated or unsaturated, substituted with an oxygen or unsubstituted five membered ring including at least one oxygen heteroatom.

[0158] In embodiments, R^{23} is




[0159] In embodiments, A is CR^9 , where R^9 is H, saturated or unsaturated C_1 to C_6 alkyl, C_1 to C_3 hydroxyl or CHR^{10} , where R^{10} is C_1 to C_6 alkyl.

[0160] In embodiments, R^9 is CHR^{10} . In embodiments, R^{10} is CH_3 .


[0161] In embodiments, A along with Q and the atoms attached thereto form a ring structure of

formula (Q1A) below,




where R²⁷ and R²⁸ are H, OH, C₁ to C₃ alkyl, C₁ to C₃ alkoxy, or C₁ to C₃ hydroxyl. In embodiments, R²⁷ and R²⁸ are each CH₃.

[0162] In embodiments, A is CCR⁶⁴R⁶⁵R⁶⁶, where R⁶⁴, R⁶⁵, and R⁶⁶ are independently selected from: C₁ to C₁₀ saturated or unsaturated, branched or straight chain alkyl, or (CH₂)_aR¹¹ where a is zero or 1 and R¹¹ is H, OH, or





where a' is zero or 1, R¹², R¹³, R¹⁴, and R¹⁵ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_bOC(O)R¹⁶ where b is zero, 1,2 or 3 and R¹⁶ is H or C₁-C₃ alkyl, or (CH₂)_dO(CH₂)_{d'}R¹⁷, where d and d' are independently zero (0) or 1 and R¹⁷ is H or




where R¹⁸, R¹⁹, R²⁰ and R²¹ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_eOC(O)R²² where e is zero, 1,2 or 3 and R²² is H or C₁-C₃ alkyl.

[0163] In embodiments, R⁶⁴ is CH₃, R⁶⁵ is a C₃ to C₈ unsaturated, branched or chain alkyl, and R⁶⁶ is (CH₂)_aO(CH₂)_{a'}R¹¹ where a and a' are both zero and R¹¹ is H or





where R^{12} , R^{13} , and R^{15} are each independently H, OH, CH_3 , C_1 - C_3 hydroxyl, and R^{14} is $(CH_2)_dO(CH_2)_{d'}R^{17}$, where d and d' are independently zero (0) or 1 and R^{17} is H or




where R^{18} , R^{19} , R^{20} and R^{21} are each independently H, OH, CH_3 , C_1 - C_3 hydroxyl, $COOH$, $OC(O)R^{22}$ where R^{22} is H or C_1 - C_3 alkyl.

[0164] In embodiments, R^{12} , R^{13} , and R^{15} are each independently OH or CH_3 .

[0165] In embodiments, R^{18} , R^{19} , and R^{21} are each independently OH or H and R^{20} is $OC(O)CH_3$.

[0166] In embodiments, A is CR^{24} , where R^{24} is



where R^{25} and R^{26} are independently selected from H, OH, or C_1 to C_6 branched or straight hydroxyl. In embodiments, R^{26} is OH and R^{25} is $C(CH_3)_2OH$.

[0167] In embodiments, Q is $CR^{61}R^{62}$, where R^{61} and R^{62} are independently selected from H, OH, C_1 to C_3 alkyl, C_1 to C_3 hydroxyl. In embodiments, Q is CH_2 . In embodiments, Q is $CHOH$. In embodiments, Q is $C(O)$.

[0168] In embodiments, B is CH_2 . In embodiments, B is CH and D is C.

[0169] In embodiments, D is CR^{29} , where R^{29} is H, OH, C_1 to C_3 alkyl, or C_1 to C_3 hydroxyl.

[0170] In embodiments, R²⁹ is OH. In embodiments, R²⁹ is H. In embodiments, R²⁹ is CH₃.


[0171] In embodiments, E is CR²⁹, where R²⁹ is H, OH, C₁ to C₃ alkyl, or C₁ to C₃ hydroxyl. In embodiments, R²⁹ is H. In embodiments, R²⁹ is CH₃. In embodiments, E is C and G is C.

[0172] In embodiments, G is CH.

[0173] In embodiments, F is CH. In embodiments, K is CH and F is C.

[0174] In embodiments, K is CR⁵_uR⁶, where u is 1 and R⁵ and R⁶ are independently selected from C₁ to C₃ alkyl. In embodiments, R⁵ and R⁶ are both CH₃. In embodiments, R⁵ and R⁶ are both H.


[0175] In embodiments, both J and L are CH. In embodiments, J is CH₂. In embodiments, J is C(R³⁰)_vR³¹, where v is 1, R³⁰ and R³¹ is




where T and X are CH₂ and where f and g are independently selected from zero, 1, or 2, and Z is CR³²R³³, where R³² and R³³ are independently selected from H, OH, C₁ to C₃ hydroxyl, or C₁ to C₃ alkyl. In embodiments, f and g are both zero and R³² and R³³ are OH and C₁ to C₃ alkyl.

[0176] In embodiments, L is CH₂.


[0177] In embodiments, M is CHOH. In embodiments, M is CH(CH₂)_jR³⁷ where j is zero, 1, or 2 and R³⁷ is H or




where j' is zero or 1, R³⁸, R³⁹, R⁴⁰, and R⁴¹ are each independently selected from H, OH, CH₃, C₁-C₃ hydroxyl, C₁-C₃ alkoxy, COOH, (CH₂)_mOC(O)R⁴² where m is zero, 1, 2 or 3 and R⁴² is H or C₁-C₃ alkyl, or (CH₂)_nO(CH₂)_{n'}R⁴³, where n and n' are independently zero or 1 and R⁴³ is H or



where R⁴⁴, R⁴⁵, R⁴⁶ and R⁴⁷ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_pOC(O)R⁴⁸ where p is zero, 1, 2 or 3 and R⁴⁸ is H or C₁-C₃ alkyl, or (CH₂)_qO(CH₂)_{q'}R⁴⁹ where q and q' are independently zero or 1 and R⁴⁹ is H or





where R⁵⁰, R⁵¹, R⁵² and R⁵³ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_rOC(O)R⁵⁴ where r is zero, 1, 2 or 3 and R⁵⁴ is H or C₁-C₃ alkyl, or (CH₂)_sO(CH₂)_{s'}R⁵⁵ where s and s' are independently zero or 1 and R⁵⁵ is H or



where R⁵⁶, R⁵⁷, R⁵⁸ and R⁵⁹ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_tOC(O)R⁶⁰ where t is zero, 1, 2 or 3 and R⁶⁰ is H or C₁-C₃ alkyl.

[0178] In embodiments, j and j' are both zero and R³⁸ is CH₃; R³⁹ is OH; R⁴⁰ is OCH₃; and R⁴¹ is H. In embodiments, j and j' are both zero and R³⁹ is (CH₂)_nO(CH₂)_{n'}R⁴³, where n and n' are independently zero or 1 and R⁴³ is H or






where R^{44} , R^{45} , R^{46} and R^{47} are each independently H, OH, CH_3 , C_1 - C_3 hydroxyl, COOH, $(CH_2)_pOC(O)R^{48}$ where p is zero, 1,2 or 3 and R^{48} is H or C_1 - C_3 alkyl.


[0179] In embodiments, R^{44} is CH_2OH ; and R^{45} , R^{46} , and R^{47} are all OH.

[0180] In embodiments, M is $CH(CH_2)_jR^{37}$ where R^{37} is H, OH, or




where j is zero or 1, R^{38} and R^{41} are each independently selected from H, OH, CH_3 , C_1 - C_3 hydroxyl, C_1 - C_3 alkoxy, COOH, $(CH_2)_mOC(O)R^{42}$ where m is zero, 1,2 or 3 and R^{42} is H or C_1 - C_3 alkyl;

where R^{39} is OR^{43} , where R^{43} is H or




where R^{44} , R^{45} and R^{47} are each independently H, OH, CH_3 , C_1 - C_3 hydroxyl, COOH, $(CH_2)_pOC(O)R^{48}$ where p is zero, 1,2 or 3 and R^{48} is H or C_1 - C_3 alkyl; and R^{46} is $(CH_2)_qO(CH_2)_{q'}R^{49}$ where q and q' are independently zero or 1 and R^{49} is H or




where R^{50} , R^{51} and R^{53} are each independently H, OH, CH_3 , C_1 - C_3 hydroxyl, COOH,

$(\text{CH}_2)_r\text{OC(O)R}^{54}$ where r is zero, 1, 2 or 3 and R^{54} is H or $\text{C}_1\text{-C}_3$ alkyl; and R^{52} is OR^{55} where R^{55} is H or



where R^{56} , R^{57} , R^{58} and R^{59} are each independently H, OH, CH_3 or $\text{C}_1\text{-C}_3$ hydroxyl; and R^{40} is OR^{43} , and R^{43} is H or



where

R^{44} , R^{45} , R^{46} and R^{47} are each independently OH, or $\text{C}_1\text{-C}_3$ hydroxyl.

[0181] In embodiments, M is CO.

[0182] In embodiments, R^1 is H or $\text{C}_1\text{-C}_3$ alkyl. In embodiments R^1 is CH_3 .

[0183] In embodiments, R^2 is H, OH, O, or $\text{C}_1\text{-C}_3$ alkyl.

[0184] In embodiments, R^3 is H or $\text{C}_1\text{-C}_3$ alkyl.

[0185] In embodiments, R^4 is CH_3 . In embodiments, R^4 is CHO.

[0186] In embodiments, R^7 and R^8 are both H. In embodiments, R^8 is OCOCH_3 .

[0187] In embodiments, F is CR^{29} , where R^{29} is H, OH, $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ hydroxyl.


[0188] In embodiments, K is CR^5R^6 .

[0189] In embodiments, M is $\text{C}(\text{O})$, COC(O)R^{36} , where R^{36} is $\text{C}_1\text{-C}_3$ alkyl, $\text{CH}(\text{CH}_2)_j\text{R}^{37}$ where j

is zero or 1.


Group Q1B Compounds

[0190] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where A is:

1. a. $CCR^{64}R^{65}R^{66}$, where R^{64} , R^{65} , and R^{66} are independently selected from: C_1 to C_{10} saturated or unsaturated, branched or straight chain alkyl, or $(CH_2)_aR^{11}$ where a is zero or 1 and R^{11} is H, OH, or saccharidyl,
2. b. or along with Q and the atoms attached thereto form a ring structure of formula (Q1B') below,



where R^{27} and R^{28} are H, OH, C_1 to C_3 alkyl, C_1 to C_3 alkoxy, or C_1 to C_3 hydroxyl;

D, E, F, and G are independently selected from: CR^{29} , where R^{29} is H, OH, C_1 to C_3 alkyl, or C_1 to C_3 hydroxyl;

K is CR^5R^6 , where R^5 and R^6 are independently selected from H, or C_1 to C_3 alkyl;


M is $CH(CH_2)_jR^{37}$ where j is zero, 1, or 2 and R^{37} is H, OH, or saccharidyl;

Q is $CHOH$, or along with A form a five membered ring containing an oxygen

heteroatom, or structure II;


R^1 , R^2 R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are independently selected from H, OH, O, or C_1 to C_3 alkyl.

[0191] In embodiments, A along with Q and the atoms attached thereto form a ring structure of formula (Q1B') below,




where R^{27} and R^{28} are H, OH, C_1 to C_3 alkyl, C_1 to C_3 alkoxy, or C_1 to C_3 hydroxyl. In embodiments R^{27} and R^{28} are each CH_3 .

[0192] In embodiments, A is $CCR^{64}R^{65}R^{66}$, where R^{64} , R^{65} , and R^{66} are independently selected from: C_1 to C_{10} saturated or unsaturated, branched or straight chain alkyl, or $(CH_2)_aO(CH_2)_{a'}R^{11}$ where a and a' are independently zero or 1 and R^{11} is saccharidyl and is:




where a' is zero or 1, R^{12} , R^{13} , R^{14} , and R^{15} are each independently H, OH, CH_3 , C_1 - C_3 hydroxyl, COOH, $(CH_2)_bOC(O)R^{16}$ where b is zero, 1, 2 or 3 and R^{16} is H or C_1 - C_3 alkyl, or $(CH_2)_dO(CH_2)_{d'}R^{17}$, where d and d' are independently zero (0) or 1 and R^{17} is H or




where R^{18} , R^{19} , R^{20} and R^{21} are each independently H, OH, CH_3 , C_1 - C_3 hydroxyl, COOH,

$(\text{CH}_2)_e\text{OC(O)R}^{22}$ where e is zero, 1, 2 or 3 and R^{22} is H or $\text{C}_1\text{-C}_3$ alkyl.

[0193] In embodiments, R^{64} is CH_3 , R^{65} is a C_3 to C_8 unsaturated, branched or chain alkyl, and R^{66} is $(\text{CH}_2)_a\text{O}(\text{CH}_2)_{a'}\text{R}^{11}$ where a and a' are both zero and R^{11} is saccharidyl and is:



where R^{12} , R^{13} , and R^{15} are each independently H, OH, CH_3 , $\text{C}_1\text{-C}_3$ hydroxyl, and R^{14} is $(\text{CH}_2)_d\text{O}(\text{CH}_2)_{d'}\text{R}^{17}$, where d and d' are independently zero (0) or 1 and R^{17} is H or




where R^{18} , R^{19} , R^{20} and R^{21} are each independently H, OH, CH_3 , $\text{C}_1\text{-C}_3$ hydroxyl, COOH, OC(O)R^{22} where R^{22} is H or $\text{C}_1\text{-C}_3$ alkyl.

[0194] In embodiments, R^{12} , R^{13} , and R^{15} are each independently OH or CH_3 .


[0195] In embodiments, R^{18} , R^{19} , and R^{21} are each independently OH or H and R^{20} is OC(O)CH_3 .

[0196] In embodiments, M is $\text{CH}(\text{CH}_2)_j\text{R}^{37}$ where j is zero, 1, or 2 and R^{37} is saccharidyl and is:




where j is zero or 1, R^{38} , R^{39} , R^{40} , and R^{41} are each independently selected from H, OH, CH_3 , $\text{C}_1\text{-C}_3$ hydroxyl, $\text{C}_1\text{-C}_3$ alkoxy, COOH, $(\text{CH}_2)_m\text{OC(O)R}^{42}$ where m is zero, 1, 2 or 3 and R^{42} is


H or C₁-C₃ alkyl, or (CH₂)_nO(CH₂)_{n'}R⁴³, where n and n' are independently zero (0) or 1 and R⁴³ is H or



where R⁴⁴, R⁴⁵, R⁴⁶ and R⁴⁷ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_pOC(O)R⁴⁸ where p is zero, 1, 2 or 3 and R⁴⁸ is H or C₁-C₃ alkyl, or (CH₂)_qO(CH₂)_{q'}R⁴⁹ where q and q' are independently zero or 1 and R⁴⁹ is H or



where R⁵⁰, R⁵¹, R⁵² and R⁵³ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_rOC(O)R¹⁴ where r is zero, 1, 2 or 3 and R⁵⁴ is H or C₁-C₃ alkyl, or (CH₂)_sO(CH₂)_{s'}R⁵⁵ where s and s' are independently zero or 1 and R⁵⁵ is H or




where R⁵⁶, R⁵⁷, R⁵⁸ and R⁵⁹ are each independently H, OH, CH₃, C₁-C₃ hydroxyl, COOH, (CH₂)_tOC(O)R⁶⁰ where t is zero, 1, 2 or 3 and R⁶⁰ is H or C₁-C₃ alkyl.

[0197] In embodiments, j and j' are both zero and R³⁸ is CH₃; R³⁹ is OH; R⁴⁰ is OCH₃; and R⁴¹ is H.

[0198] In embodiments, j and j' are both zero and R³⁹ is (CH₂)_nO(CH₂)_{n'}R⁴³, where n and n' are independently zero or 1 and R⁴³ is saccharidyl and is:





where R^{44} , R^{45} , R^{46} and R^{47} are each independently H, OH, CH_3 , C_1 - C_3 hydroxyl, $COOH$, $(CH_2)_pOC(O)R^{48}$ where p is zero, 1,2 or 3 and R^{48} is H or C_1 - C_3 alkyl.

[0199] In embodiments, R^{44} is CH_2OH ; and R^{45} , R^{46} , and R^{47} are all OH.

[0200] In embodiments, D is CR^{29} , where R^{29} is C_1 to C_3 alkyl.

[0201] In embodiments, E is CR^{29} , where R^{29} is C_1 to C_3 alkyl.

[0202] In embodiments, G is CH.

[0203] In embodiments, F is CH.

[0204] In embodiments, K is CH and F is C.

[0205] In embodiments, K is $CR^{5u}R^6$, where u is 1 and R^5 and R^6 are independently selected from C_1 to C_3 alkyl. In embodiments, R^5 and R^6 are both CH_3 .


[0206] In embodiments, R^2 , R^4 , R^8 , and R^9 are all H.


[0207] In embodiments, R^1 is C_1 - C_3 alkyl. In embodiments, R^1 is H.

[0208] In embodiments, R^3 is O. In embodiments, R^3 is H.

Group Q2 Compounds

[0209] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:






where A is C or CR⁴, where R₄ is H, OH, C₁ to C₃ alkyl, with the caveat that if A is C, then B is CR⁶;


B is CR⁵_aR⁶, where a is zero (0) or 1, and R⁵ and R⁶ are independently selected from H, OH, C₁ to C₃ alkyl, with the caveat that if A is C, then B is CR⁶;

X is H, or along with Y and the atoms attached thereto form a ring structure having formula (Q2A) below,



where R⁹ and R¹⁰ are independently selected from O and CH₂;

Z is H or along with Y and the atoms attached thereto form a ring structure having formula (Q2B) below,




where R¹³ and R¹⁴ are independently selected from O and CH₂;

Y together with either X or Z forms a ring structure of formula Q2A or Q2B above; and

R¹, R² and R³ are independently selected from H, CH₂, C₁ to C₃ alkyl.


[0210] In embodiments, A is CH_3 . In embodiments, R^1 is CH_2 . In embodiments, R^2 is H. In embodiments, R^3 is H. In embodiments, A is CR^4 , where R^4 is C_1 to C_3 alkyl. In embodiments, B is CH_2 .

[0211] In embodiments, X and Y along with the atoms attached thereto form a ring structure having formula (Q2A) below,



In embodiments, R^9 is O and R^{10} is CH_2 .


[0212] In embodiments, Y and Z along with the atoms attached thereto form a ring structure having formula (Q2B) below,




where R^{13} and R^{14} are independently selected from O and CH_2 . In embodiments, R^3 is CH_3 . In embodiments, R^1 is H. In embodiments, R^2 is CH_2 . In embodiments, R^2 is CH_3 .

Group Q3 Compounds

[0213] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:





where A is C or CR⁴, where R₄ is H, OH, C₁ to C₃ alkyl, with the caveat that if A is C, then B is CR⁶;

B is CR⁵_aR⁶, where a is zero or 1, and R⁵ and R⁶ are independently selected from H, OH, C₁ to C₃ alkyl;

Z is H or CR¹¹R¹² where R¹¹ and R¹² are independently selected from H, OH, C₁ to C₃ alkyl;

R¹, R² and R³ are independently selected from H, CH₂, or a saturated or unsaturated C₁ to C₆ alkyl; and


R⁹ and R¹⁰ are independently selected from O or CH₂.

[0214] In embodiments, A is CCH₃. In embodiments, B is CH₂. In embodiments, Z is H. In embodiments, R¹ is CH₂. In embodiments, R₂ is H. In embodiments, R₃ is H. In embodiments, R⁹ is O. In embodiments, R¹⁰ is CH₂.

GROUP R COMPOUNDS

Group R1 Compounds

[0215] In embodiments a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where R¹ is OH or saccharidyl;

R^2 and R^3 are independently H, OH, and COOH; and


R^4 is C₃-C₁₄ saturated or unsaturated alkyl optionally substituted with one or more hydroxyls.

[0216] In embodiments R^1 is OH. In embodiments R^2 is H. In embodiments, R^2 is OH. In embodiments, R^3 is H. In embodiments, R^3 is COOH. In embodiments, R^4 is C₉ to C₁₃ unsaturated alkyl. In embodiments, R^4 is C₉ to C₁₃ unsaturated alkyl with at least one hydroxyl. In embodiments, R^4 is C₁₀ to C₁₂ alkyl with at least two double bonds, such as C₁₁ with at least two double bonds. In embodiments, R^4 is C₁₁ with three double bonds. In embodiments, R^4 is C₁₁ with three double bonds and one hydroxyl. In embodiments, R^4 is C₃ to C₇ with at least one double bond.

[0217] In embodiments of a compound according to **Formula R1** is a compound where R^2 and R^3 are independently H, OH, and COOH; and R^4 is C₃-C₁₀ saturated or unsaturated alkyl optionally substituted with one or more hydroxyls. In embodiments, R^2 is OH. In embodiments, R^3 is COOH. In embodiments, R^4 is C₃ to C₇ with at least one double bond.

Group R2 Compounds

[0218] In embodiments a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where


A is (CH₂)_x, where x is 0 or 1;

R^5 is saccharidyl;

R^2 and R^3 are independently H, OH, and COOH; and

R^4 is C₈-C₁₆ saturated or unsaturated alkyl optionally substituted with one or more hydroxyls.

[0219] In embodiments, a compound of formula R2 the following structure:



(R2A),

where

A, D, E and G are independently $(CH_2)_x$, where x is 0 or 1;

R^1 is H or saccharidyl;


R^2 and R^3 are independently H, OH, and COOH; and

R^4 is C_8 - C_{16} saturated or unsaturated alkyl optionally substituted with one or more hydroxyls.

[0220] In embodiments, R^1 is H. In embodiments, R^2 is H. In embodiments, R^3 is H. In embodiments, R^4 is C_8 to C_{14} unsaturated alkyl optionally substituted with one or more hydroxyls, such as C_{10} to C_{13} unsaturated alkyl optionally substituted with one or more hydroxyls. In embodiments, R^4 is C_{11} unsaturated alkyl optionally substituted with one or more hydroxyls. In embodiments, R^4 is C_{11} unsaturated alkyl with one double bond optionally substituted with one or more hydroxyls. In embodiments, x is one in A and E; and x is zero in D and E.

Group R3 Compounds

[0221] In embodiments a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where

R^1 is H, or OH;

R^2 and R^3 are independently H, OH, or CH_2OH ;

X is O or $CHOH$;

Y is $C=O$ or CH_2 ; and

R^4 is C_3-C_{12} unsaturated alkyl.

[0222] In embodiments, R^1 is OH. In embodiments, R^2 is H. In embodiments, X is $CHOH$. In embodiments, Y is CH_2 . In embodiments, R^3 is H.


[0223] In embodiments, R^4 is a C_3 to C_7 unsaturated alkyl. In embodiments, R^4 is a C_5 unsaturated alkyl. In embodiments, R^4 is a C_5 alkyl with one double bond.

[0224] In embodiments, R^3 is OH.

[0225] In embodiments, R^4 is a C_9 to C_{12} unsaturated alkyl. In embodiments, R^4 is a C_{10} unsaturated alkyl. In embodiments, R^4 is a C_{10} alkyl with one double bond.

Group R4 Compounds

[0226] In embodiments a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where

X is O or NH;

R^1 is H, OH, or CH_2OH ;

R^2 is H or $OCOR^5$, wherein R^5 is a C_1-C_3 alkyl; and


R^3 is H or C_1-C_{10} saturated or unsaturated alkyl.

[0227] In embodiments, X is O. In embodiments, R¹ is OH. In embodiments, R² is OCOR⁵. In some embodiments where R² is OCOR⁵, R⁵ is CH₃. In embodiments, R³ is C₅ to C₉ saturated or unsaturated alkyl, such as C₇ unsaturated alkyl. In embodiments, R³ is C₇ alkyl with one double bond.

[0228] In embodiments, X is NH. In some embodiments where X is NH, R¹ and R² are both H. In some embodiments where X is NH, R³ is H.

GROUP S COMPOUNDS

[0229] In embodiments a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:




where:


X and Y are each independently C₇-C₂₀ straight or branched chain unsaturated alkyl unsubstituted or substituted with one or more OH;

A is saccharidyl; and

R¹⁰ and R¹¹ are each independently H, OH or saccharidyl.

[0230] In embodiments a compound according to **Formula S** has the following structure:





where:


a, b, d, e, f and g are each independently zero or 1;

X and Y are each independently C₇-C₂₀ straight or branched chain unsaturated alkyl unsubstituted or substituted with one or more OH;

R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, and R⁹ are each independently H, OH, CH₃, CH₂CH₃, and CH₂OH; and

R¹⁰ and R¹¹ are each independently H, OH or saccharidyl.

[0231] In embodiments, one or both of R¹⁰ and R¹¹ are saccharidyl and are independently




where

i, j and k are each independently zero or 1, and

R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷ and R¹⁸ are each independently H, OH, CH₃, CH₂CH₃ and CH₂OH.

[0232] In embodiments, X and Y are each independently C₈-C₁₆ straight or branched chain unsaturated alkyl unsubstituted or substituted with one or more OH and having 2 to 6 double bounds. In embodiments, X and Y are each independently C₁₀-C₁₅ straight or branched chain unsaturated alkyl unsubstituted or substituted with one or more OH and having 2 to 4 double bounds.

[0233] In embodiments a compound according to **Formula S** has the following structure:




(S’’),

where a, b, d, e, f and g, and R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰ and R¹¹ are as defined above; and wherein

R¹⁹, R²⁰, R²¹, R²² and R²³ are each independently H, CH₃ or OH.

[0234] In embodiments a compound according to **Formula S** has the following structure:




(S'''),

where R¹⁹, R²⁰, R²¹, R²³ and R²⁴ are each independently H, CH₃ or OH.

GROUP T COMPOUNDS

[0235] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:



where:

R¹ is H, or COR⁴ where R⁴ is H, or saturated or unsaturated C₁-C₆ straight or branched alkyl;


and

R^2 , R^3 , R^a , R^b , R^c , R^d , R^e , R^f , R^g , R^h , R^i , and R^j are independently selected from H, or saturated or unsaturated C₁-C₃ alkyl.

[0236] In embodiments, R^2 , R^3 , R^a , R^b , R^c , R^d , R^e , R^f , R^g , R^h , R^i , and R^j are independently selected from H, or saturated or unsaturated C₁-C₃ alkyl with the caveat that saturated or unsaturated C₁-C₃ alkyl substituents are not directly adjacent each other.

[0237] In embodiments, R^2 is a saturated or unsaturated C₁-C₃ alkyl, such as methyl. In embodiments, R^3 is a saturated or unsaturated C₁-C₃ alkyl, such as methyl. In embodiments, R^2 is CH₃ and R^3 is CH₃.

[0238] In embodiments, R^1 is H. In embodiments, R^1 is COCH₃. In embodiments, R^1 is COR⁴ and R⁴ is




where R⁵ and R⁶ are independently selected from H, or saturated or unsaturated C₁-C₄.

[0239] In embodiments, R⁵ and R⁶ are both CH₃.

[0240] In embodiments a compound according to **Formula T** is a compound where (i) R^2 , R^3 , R^a , R^b , R^c , R^d , R^e , R^f , R^g , R^h , R^i , and R^j are independently selected from H, or saturated or unsaturated C₁-C₃ alkyl; and (ii) R^7 is H or saturated or unsaturated C₁-C₃ alkyl. In such embodiments, R^7 may be CH₃. R^2 , R^3 , R^a , R^b , R^c , R^d , R^e , R^f , R^g , R^h , R^i , and R^j may independently be selected from H, or saturated or unsaturated C₁-C₃ alkyl with the caveat that saturated or unsaturated C₁-C₃ alkyl substituents are not directly adjacent each other. R^2 may be a saturated or unsaturated C₁-C₃ alkyl, such as methyl. R^3 may be a saturated or unsaturated C₁-C₃ alkyl, such as methyl. In some of such embodiment, R^2 is CH₃ and R^3 is CH₃.

[0241] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:






where

R^2 , R^3 , R^a , R^b , R^c , R^d , R^e , R^f , R^g , R^h , R^i , and R^j are independently selected from H, or saturated or unsaturated C₁-C₃ alkyl; and

R^7 is H or saturated or unsaturated C₁-C₃ alkyl.

GROUP U COMPOUNDS

[0242] In embodiments, a bioactive, taste modulating, or salty taste modulating compound is a compound having the following structure:




where:


R^1 is H, OH, or C₁-C₃ alkoxy;

R^a , R^b , R^c , R^d , and R^e are each independently selected from: H, OH, or C₁-C₃ alkoxy;


R^2 is H, OH, C₁-C₃ alkoxy, or R^2 and R^3 together form




or



to form a compound of the following structure



or




(U'')

R^3 is H, OH, C₁-C₃ alkoxy, or R^2 and R^3 together form a ring structure as indicated above;

R^4 is H, OH, or C₁-C₃ alkoxy;

X is O or CH; and

Y is O or CR⁵, where R⁵ is H or



where

R⁶, R⁷, and R⁸ are each independently selected from: H, OH, or C₁-C₃ alkoxy;

R^f, R^g, R^h, R¹, R^j, and R^k are each independently selected from: H, OH, or C₁-C₃ alkoxy,


wherein the at least one compound of formula U is present in the food product in an amount sufficient to enhance a perception of saltiness of the food product.

[0243] In embodiments, R¹ is H. In embodiments, R¹ is -OCH₃. In embodiments, R² is H. In embodiments, R³ is H. In embodiments, both R² and R³ are H.

[0244] In embodiments where R² and R³ together form



to form a compound of the following structure




(U'),


R⁴ is H.

[0245] In embodiments where R² and R³ form

/




to form a compound of the following structure



R⁴ is H.


[0246] In embodiments, R^a, R^b, R^c, R^d, and R^e are independently H or OH.


[0247] In embodiments, Y is O. In embodiments, Y is CR⁵, where CR⁵ is



R⁶ may be OH. R⁷ may be OH. R⁸ may be OH. R^f, R^g, R^h, Rⁱ, R^j, or R^k may independently be H or OH.

[0248] In embodiments, a compound according to Formula U is a compound having the following structure:





where:

$R^1, R^2, R^3, R^4, R^6, R^7$, and R^8 are each independently H, OH, or C_1-C_3 alkoxy;

X and Z are each independently O or CH; and

$R^a, R^b, R^c, R^d, R^e, R^f, R^g, R^h, R^i, R^j$, and R^k are each independently H, OH, or C_1-C_3 alkoxy.

[0249] In embodiments, a compound of **Formula UA** is a compound where R^1 is OH or C_1-C_3 alkoxy. In embodiments, R^2 is H. In embodiments, R^3 is OH or C_1-C_3 alkoxy. In embodiments, R^4 is OH or C_1-C_3 alkoxy. In embodiments, R^6 is OH or C_1-C_3 alkoxy. In embodiments, R^7 is OH or C_1-C_3 alkoxy. In embodiments, $R^a, R^b, R^c, R^d, R^e, R^f, R^g, R^h, R^i, R^j$, and R^k are each independently H or OH. In embodiments, $R^a, R^b, R^c, R^d, R^e, R^f, R^g, R^h, R^i, R^j$, and R^k are each H. In embodiments, X is O. In embodiments, Z is O.

[0250] For purpose of convenience, **Table 2A** lists the initially selected 99 compounds (with compounds 12 and 20 being the same compound isolated from different sources) and indicates which categories and subcategories, if appropriate, into which each of the compounds falls. **Table 2B** lists the categories and the compound that fall within each category.

Table 2A: Listing of compounds according to category (according to claim 1, the food product comprises a compound of formula B3)

Compound	Category,	Sub-Categories
1	A	-
2	A	-
3	A	-
4	A	-
5	A	-
6	A	-
7	A	-
8	A and B	B1,B3

Compound	Category,	Sub-Categories
9	A	-
10	B	B1,B3
11		B1,B3
12		B1,B3
13		B1
14		B1'
15		B1'
16		B2, B4
17		B2, B4
18		B1'
19		B1,B3
20		B1, B3
21		B1
22		B1'
23	C	C" ,C'''
24		C",C"',C'''
25		C", C'''
26		C',C'''
27		C'
28		-
29	D	-
30		-
31		-
32	E	-
33		-
34	F	-
35		F', F''
36		F'
37	Q	Q1, Q1A, Q1B
38		Q1, Q1B
39	F	F'
40	G	-
41	H	-
42	F	-
43		-

Compound	Category,	Sub-Categories
44	I	-
45		-
46		-
47		I'
48	J	J1,J1"
49		J1,J1"
50		J1, J1', J2
51		J2
52	K	-
53		-
54		-
55		L'
56	M	-
57		-
58		-
59		P5
60	P	P2, P2'
61		P4
62		P1
63		P1
64		P1
65		P1
66		P4
67		P2
68		P2, P3
69		P2
70		P1
71		P1
72		P1
73	Q	Q1
74		Q1
75		Q1
76		Q1
77		Q1

Compound	Category,	Sub-Categories
78		Q1
79		Q2, Q2A, Q3
80a		Q2, Q2B
80b		Q2, Q2B
81	R	R1
82		R3
83		R4
84		R4
85		R1
86		R1
87		R3
88		R1
89		R1
90		R2
91	S	-
92		-
93		-
94	T	T2
95		-
96		U'
97		UA
98	U	U"
99		-

Table 2B: Listing of compounds according to category

Category	Compounds
A	1-9
B *	8, 10-22
C	23-28
D	29-31
E	32, 33
F	34-36, 39, 42, 43
G	40
H	41
I	44-47
J	48-51

Category	Compounds
K	52
L	53-55
M	56
N	57
O	58
P	59-72
Q	37, 38, 73-79, 80a, 80b
R	81-90
S	91, 92
T	93-95
U	96-99

* Compounds 8, 10-12, 19, and 20 fall within formula B3; according to claim 1, a compound of formula B3 is present in the food product.

[0251] As can be seen from the structures of the compounds provided above, many of the compounds have structural similarities. Accordingly, it is believed that structural derivatives of the specific compounds presented above would also have the ability to elicit the perception of saltiness or enhance saltiness. Combinations of the compounds could also serve to elicit the perception of saltiness or enhance saltiness. In addition or alternatively, one or more of the compounds may elicit the perception of other simple or complex flavors, other than or in addition to saltiness.

[0252] Many of the structural similarities between the compounds are reflected in the compounds of Formulas **A** through **U**, as well as subclasses thereof, presented above. It will be further understood, based on the compounds identified herein, that one or more of gingerols, alkyl substituted phenols, acridone alkaloids, labdanes, primaranes, saponines, neolignans, pentacyclic triterpenes, 2,2'-cyclohexanes, dibenzylbutane lignans, bicyclic triterpenes, phloroglucinones, caryophyllenes, beta-carbolines, limnoids, cumarines, cardanolide steroids, fatty acids, and derivatives may be candidates for taste modulating compounds. It will be further understood that other structural similarities of the compounds presented herein may be exploited to develop taste modulating compounds.

[0253] By way of example, many of the compounds presented herein have unsaturated carbon chains of at least 11 carbons with attached hydroxyl groups and may be amphiphilic, with hydrophobic head portions and hydrophobic tails. Other compounds having, for example C₅-C₂₀ alkane or alkene tails may elicit or enhance saltiness. Similarly, other compounds with differently substituted carboxyl or hydroxyl groups may elicit or enhance the perception of saltiness.

[0254] Many compounds presented herein have large numbers of cyclic groups having a central portion that may be hydrophobic and peripheral regions that may be hydrophilic. More specifically, some compounds include pentacyclohexane with hydroxyl groups, attached sugars and at least one ester linkage. Substitution of the central hydrophobic ring structure with, for example, C₅-C₂₀ hydrophobic alkyl or alkene groups, may result in compounds that may elicit or enhance saltiness. Alternative substitution of hydroxyl groups at the peripheral regions or substitution with carboxylic groups may also result in compounds that elicit or enhance the perception of saltiness.

[0255] Many of the compounds presented above have one or more aromatic ring structures, with some being substituted and some being unsubstituted. Similar substitution or unsubstitution of such compounds may result in compounds that enhance or elicit saltiness.

[0256] A plurality of compounds presented herein include saturated carbon chains of at least 9 carbons and one oxygen containing group such as hydroxyl, carbonyl, carboxyl, or ester. Other compounds having unsaturated carbon chains of, for example, 5 to 15 carbons and an oxygen group may have similar effects with regard to salty taste.

[0257] Many compounds presented herein have at least one phenol group with an ether group and a carbon side chain comprising at least seven carbons. Other similar compounds may have similar effects with regard to salty taste.

[0258] A number of compounds presented herein have a benzyl heterocyclic furan with various attached groups containing unsaturated carbon linkages and at least one carbonyl group. Other similar compounds may have similar activity with regard to salty taste.

[0259] A plurality of compounds presented above contain a cyclopentaphenthrene group. Other compounds having a cyclopentaphenthrene and similar substituents may have similar activity with regard to salty taste.

[0260] A number of the compounds presented herein include a benzopyranone group. Other compounds having a benzopyranone and similar substituents may have similar activity with regard to salty taste.

[0261] Some compounds presented above have unsubstituted carbon chain with a minimum of 13 carbons and at least one carbonyl group. Other similar compounds may have similar activity with regard to salty taste.

[0262] A plurality of compounds presented herein have a methoxymethyltetrahydrobenzo-cyclooctabenzo-dioxole or -annulene group. Other compounds having such groups may have similar activity with regard to salty taste.

[0263] A number of compounds presented above have tetracyclohexane with an attached ester or carbonyl moiety. Other similar compounds may have similar activity with regard to salty

taste.

[0264] Many of the compounds can be classified as lactones, lignol-like compounds, oxylipins, polyisoprene glycosides, triterpenoid glycosides, alkylamides, or gamma pyrenes. Other similar compounds may also elicit or enhance salty taste perception.

[0265] It will be understood that derivations of the compounds discussed above are provided for purposes of example and that other derivatives or derivations may be made based on structural similarities between the various compounds, resulting in compounds that elicit or enhance perception of saltiness.

EVALUATION OF SALTY TASTE OR SALTY TASTE ENHANCEMENT

[0266] Many of the identified compounds were tested by tasters and rated for perception of saltiness in combination with reduced amounts of sodium chloride and were assigned a rating (DAP score) for saltiness. Briefly, each individual tested compound was placed in water and in sodium solution to test saltiness and saltiness enhancement potential. Tests in water were executed in a compound concentration of 10 ppm. Tests in sodium solution were executed in compound concentrations of 0.1, 1 and 10 ppm. Two control sodium solutions with known organoleptic salt intensities were provided as references for each test. The test for individual compounds was also conducted using simple broth instead of sodium solution. A number of compound combinations identified from the Na-solution DAP test were used for the broth DAP test. Tests were executed with a trained panel of 9-12 assessors. For Na-solution DAP tests, a DAP score of greater than 3.1 indicates saltiness or salt enhancement. The DAP score can be correlated with a sodium reduction potential by subtracting 3.1 from the DAP score. For example, a DAP score of 4.0 would result in a 9% sodium reduction potential $((4.0 - 3.1) * 10 = 9\%)$, which means that 9% less sodium may be present in a food product having the salty compound relative to a substantially similar food product that does not have the salty compound while producing a similar saltiness. For broth DAP tests, a DAP score of greater than 7.6 indicates saltiness or salt enhancement. The DAP score can be correlated with a sodium reduction potential by subtracting 7.6 from the DAP score. For example, a DAP score of 8.5 would result in a 9% sodium reduction potential $((8.5 - 7.6) * 10 = 9\%)$, which means that 9% less sodium may be present in a food product having the salty compound relative to a substantially similar food product that does not have the salty compound while producing a similar saltiness.

[0267] A summary of the compounds, the channels for which *in vitro* thresholds were identified, and the DAP scores are provided below in **Table 3**.

Table 3. Activity of identified compounds in Na-Solution and in simple broth (according to claim 1, the food product comprises a compound of formula B3)

Compound	DAP in Na-Sol.	DAP in Broth
1 *	3.6	7.8

Compound	DAP in Na-Sol.	DAP in Broth
2 *	3.5	7.8
3 *	4.1	8.8
4 *	3.5	N.T.
5 *	3.6	N.T.
6 *	3.8	N.T.
7 *	3.7	N.T.
8	4	N.T.
9 *	3.8	N.T.
10	3.9	7.9
11	2.6	8.1
12	4	7.6
13 *	3.8	7.9
14 *	3.6	N.T.
15 *	3.5	8.1
16 *	3.3	8.0
17 *	3.6	8.2
18 *	3.8	8.1
19	N.T.	N.T.
20	4	N.T.
21 *	4	N.T.
22 *	N.T.	N.T.
23 *	2.9	8.3
24 *	N.T.	N.T.
25 *	3.9	N.T.
26 *	3.4	N.T.
27 *	3.4	N.T.
28 *	3.6	N.T.
29 *	3.5	8.0
30 *	3.1	N.T.
31 *	3.2	N.T.
32 *	3.5	8.1
33 *	3.4	7.7
34 *	3.9	7.7
35 *	3.7	8.1
36 *	3.9	8.2

Compound	DAP in Na-Sol.	DAP in Broth
37 *	3.7	8.1
38 *	3.9	8.1
39 *	N.T.	N.T.
40 *	3.5	7.8
41 *	3.9	7.5
42 *	3.1	8.1
43 *	3.7	8.0
44 *	3.8	8.6
45 *	3.5	8.2
46 *	3.6	7.8
47 *	N.T.	N.T.
48 *	3.7	7.8
49 *	3.6	8.2
50 *	4	N.T.
51 *	N.T.	N.T.
52 *	3.6	N.T.
53 *	3.7	8.5
54 *	3.4	8.1
55 *	N.T.	N.T.
56 *	4	7.7
57 *	3.1	N.T.
58 *	3.3	8.0
59 *	3.3	7.9
60 *	3.8	8.1
61 *	3.6	N.T.
62 *	3.9	8.0
63 *	3.5	8.0
64 *	3.7	7.8
65 *	3.3	7.9
66 *	3.2	7.8
67 *	N.T.	N.T.
68 *	N.T.	N.T.
69 *	N.T.	N.T.
70 *	N.T.	N.T.

Compound	DAP in Na-Sol.	DAP in Broth
71 *	N.T.	N.T.
72 *	N.T.	N.T.
73 *	3.5	8.0
74 *	3.2	7.7
75 *	3.6	N.T.
76 *	3.2	8.0
77 *	3.4	8.1
78 *	N.T.	N.T.
79 *	N.T.	N.T.
80a *	3.9	N.T.
80b *	3.9	N.T.
81 *	3.2	7.8
82 *	3.8	8.3
83 *	4.2	8.1
84 *	3.8	8.1
85 *	2.9	N.T.
86 *	3.9	8.0
87 *	3.9	8.0
88 *	3.2	8.0
89 *	3.7	7.6
90 *	N.T.	N.T.
91 *	3.7	N.T.
92 *	4	N.T.
93 *	N.T.	N.T.
94 *	4.2	N.T.
95 *	N.T.	N.T.
96 *	N.T.	N.T.
97 *	N.T.	N.T.
98 *	N.T.	N.T.
99 *	4.1	N.T.
* not within formula B3		

[0268] In Table 3 above, "N.T." with regard to a DAP score, means the compound was not taste tested.

[0269] The data in **Table 3** above reflect the best DAP score for the tested concentrations of 0.1 parts per million (ppm), 1 ppm and 10 ppm. The highest concentrations did not always result in the highest DAP scores.

[0270] Compounds in water with no sodium were also tested. The compounds in water with no sodium elicited no appreciably discernable salty taste, even at the 10 ppm concentration (data not shown).

[0271] DAP score testing results for various pairs of compounds are presented in **FIG. 1** (sodium solution) and **FIG. 2** (broth). Certain combinations were tested twice. For these combinations, two DAP scores are shown in the tables presented in **FIGS 1-2**. As shown in the results presented in **FIGS. 1-2**, certain combinations of compounds can enhance the perception of saltiness. Some combinations resulted in DAP scores as high as 4.5 in some tests. See, for example, the combination of compound **83** and **13** in **FIG. 1** (sodium solution) and the combination of compound **12** and **18** in **FIG. 2** (broth). Such DAP scores may result in a sodium reduction potential of about 14%. The combinations tested in **FIGS. 1-2** are representative of the combinations that may be used in a food product to enhance the perception of saltiness or reduce sodium content. It will be understood that any other suitable combination of compounds may be employed.

[0272] Additional testing of combinations of pairs of compounds was performed in sodium solution. The DAP scores from this additional testing is shown below in **Table 4**.

Table 4: Activity of a combination of bioactive compounds in sodium solution
(according to claim 1, the food product comprises a compound of formula B3)

	Compound 66: (0.1ppm)	Compound 29: (10ppm)	Compound 16: (1ppm)	Compound 33: (1ppm)	Compound 73: (1ppm)
Compound 83: (1 ppm)	3.7	4.0	3.8	3.6	4.0
Compound 10: (1 ppm) *	3.3	3.7	3.8	2.9	3.6
Compound 45: (10 ppm)	3.2	3.3	3.8	3.7	4.2

* compound of formula B3

[0273] In addition, more than two bioactive, taste modulating or salty taste modulating compounds described herein may be included in a food product. By way of example, **Table 5** below shows DAP scores obtained from testing sodium solutions and chicken broth containing a combination of compounds **12**, **13** and **83**. As shown in **Table 5**, such a combination resulted

in a DAP score of 5.3 when tested in a sodium solution. Accordingly, such a combination may result in a sodium reduction potential of about 22%. Of course, other suitable combinations of three or more compounds may be used or included in a food product to enhance the perception of saltiness or to reduce sodium content.

Table 5: Activity of a combination of bioactive compounds in sodium solution and broth

Compound	Conc.	In Na Solution	In Broth
12 *	1 ppm		
13	10 ppm	5.3	8.1
83	1 ppm		

* compound of formula B3

[0274] Some illustrative examples of combinations of compounds that may produce desired or beneficial effect, for example when incorporated in a food product, include combinations that include at least one compound selected from the group consisting of compounds **3, 10, 12, 13, 16, 18, 29, 33, 36, 37, 41, 43, 44, 45, 48, 53, 56, 62, 66, 73, 82, 83, and 84**. Another illustrative example is a combination that includes at least one compound selected from the group consisting of compounds **10, 12, 13, 18, 36, 45, 56, 82, and 83**. Yet another illustrative example is a combination that includes compounds **12, 13 and 83**. Of course, any other suitable or desirable combination may be used.

[0275] DAP scores for combinations of three different compounds in broth are shown in **Table 6** below.

Table 6: Activity of a combination of bioactive compounds in broth

(according to claim 1, the food product comprises a compound of formula B3)	
Compound (concentration)	DAP score
3 (0.1 ppm)	
36 (0.1 ppm)	7.8
44 (10 ppm)	
3 (0.1 ppm)	
36 (0.1 ppm)	7.7
53 (1 ppm)	
3 (0.1 ppm)	
36 (0.1 ppm)	7.9
18 (10 ppm)	
13 (10 ppm)	
84 (1ppm)	8.1

Compound (concentration)	DAP score
44 (10 ppm)	
13 (10 ppm)	
84 (1ppm)	7.9
53 (1 ppm)	
13 (10 ppm)	
84 (1ppm)	8.0
3 (0.1 ppm)	
18 (10 ppm)	
12 (1 ppm) *	8.1
44 (10 ppm)	
18 (10 ppm)	
12 (1 ppm) *	8.2
53 (1 ppm)	
18 (10 ppm)	
12 (1 ppm) *	8.3
3 (0.1 ppm)	
* compound of formula B3	

SOURCING

[0276] Natural sources of the mentioned taste modulating or salty taste modulating compounds can be extracted by a variety of methods such as, but not exclusive to, water, solvent extractions (ethanol/water combinations), or supercritical carbon dioxide or other volatilization methods. These concentrated extracts or isolates could be stabilized physically by encapsulation, for example, or chemical reaction to non-reactive compounds such as simple sugars or small chain fatty acids. Compounds may be altered for their solubility in aqueous solutions by hybridization to larger sized molecules and additionally processed or reacted to create an impacting ingredient in either a dry or aqueous form.

[0277] In embodiments, a composition comprises a bioactive, taste modulating or salty taste

modulating compound described herein. The composition may be included in a food product. In embodiments, the composition comprises one or more natural extracts. In another embodiment, the extract is selected from a plant or microbial (e.g., fungi or bacterial) source. Examples of suitable natural extracts include extracts derived from *Aesculus hippocastaneum*; *Alchemilla xanthochlora*; *Angelica archangelica*; *Apocynum cannabinum*; *Azadirachta indica*; *Actinomycete* bacteria (Strain code: 01702axxx000002); *Capsicum annuum*; *Cimicifuga racemosa*; *Commiphora mukul*; *Embelia ribes*; *Evodia rutaecarpa*; *Ferula assa-foetida*; *Fungi* (Strain code: 02295fxxx000001; Strain code: 01469fxxx000005); *Gleditschia australis*; *Kaempferia galanga*; *Lavandula officinalis*; *Marrubium vulgare*; *Mesua ferrea*; *Nephelium cuspidatum*; *Orthosiphon stamineus*; *Persea gratissima*; *Petroselinum sativum*; *Piper longum*; *Pithecoctenium echinatum*; *Podophyllum peltatum*; *Psidium guajava*; *Ricinus communis*; *Salvia miltiorrhiza*; *Schisandea chinensis*; *Teclea trichocarpa*; *Vitex agnus*; *Xysmalobium undulatum*; *Yucca gloriosa*; *Zanthoxylum piperitum*; *Zingiber officinalis* and others. The composition may be in a dry or liquid form. The liquid composition may be a solution, suspension, colloidal suspension, microencapsulated suspension, emulsion, or the like, or combinations thereof. The dry composition may be a microencapsulation solid, agglomeration, or the like or combinations thereof.

[0278] In embodiments, a bioactive, taste modulating or salty taste modulating compound described herein is included in a composition comprising a carrier. The composition comprising the carrier may be incorporated into a food product. Any suitable carrier may be used. Examples of suitable carriers include propylene glycol, ethanol, water, or oil. In embodiments, the carrier is a starch, such as a starch comprising carbohydrate, a maltodextrin, a cyclodextrin or another dextrin, or a liposome. In embodiments, the carrier is an encapsulant or the carrier may comprise an embedded bioactive, taste modulating or salty taste modulating compound.

DEFINITIONS

[0279] All scientific and technical terms used herein have meanings commonly used in the art unless otherwise specified. The definitions provided herein are to facilitate understanding of certain terms used frequently herein and are not meant to limit the scope of the present disclosure.

[0280] As used in this specification and the appended claims, the singular forms "a", "an", and "the" encompass embodiments having plural referents, unless the content clearly dictates otherwise.

[0281] As used in this specification and the appended claims, the term "or" is generally employed in its sense including "and/or" unless the content clearly dictates otherwise. The term "and/or" means one or all of the listed elements or a combination of any two or more of the listed elements.

[0282] As used herein, "have", "having", "include", "including", "comprise", "comprising" or the

like are used in their open ended sense, and generally mean "including, but not limited to". It will be understood that "consisting essentially of," "consisting of," and the like are subsumed in "comprising" and the like. As used herein, "consisting essentially of," as it relates to an composition, product, method or the like, means that the components of the composition, product, method or the like are limited to the enumerated components and any other components that do not materially affect the basic and novel characteristic(s) of the composition, product, method or the like.

[0283] The words "preferred" and "preferably" refer to embodiments of the invention that may afford certain benefits, under certain circumstances. However, other embodiments may also be preferred, under the same or other circumstances. Furthermore, the recitation of one or more preferred embodiments does not imply that other embodiments are not useful, and is not intended to exclude other embodiments from the scope of the disclosure, including the claims.

[0284] Also herein, the recitations of numerical ranges by endpoints include all numbers subsumed within that range (e.g., 1 to 5 includes 1, 1.5, 2, 2.75, 3, 3.80, 4, 5, etc. or 10 or less includes 10, 9.4, 7.6, 5, 4.3, 2.9, 1.62, 0.3, etc.). Where a range of values is "up to" a particular value, that value is included within the range.

[0285] As used herein, a "bioactive compound" is a compound that alters the flow of ions through one or more channels associated with the perception of salty taste or another taste associated with consumption of sodium chloride.

[0286] As used herein, a "taste modulating compound" is a compound that modifies the taste of a food product. By way of example, a taste modulating compound may modify the taste of a food product due to a particular taste imparted by the taste modulating compound, due to a modification of the perceived taste of the food product, or a component thereof, or the like. In embodiments, a taste modulating compound is a salty taste modulating compound.

[0287] As used herein a "salty taste modulating compound" is a compound that, when ingested, elicits or enhances a perception of salty taste alone or in the presence of a salt, such as sodium chloride.

[0288] As used herein, a composition that is "substantially similar" to another composition contains substantially the same concentration of components (e.g., within about 5%) except for the specifically enumerated components that make the compositions different. For example, a composition that includes a salty compound may be substantially similar to a composition that does not have the salty compound, if the components of the compositions, other than the salt and the salty compound, are present in a substantially similar concentration.

[0289] As used herein, a compound "derived" from a natural product is a compound that exists in a natural product, whose identity is verified. The compound derived from the natural product may be extracted from, for example, a plant or microbial source as opposed to being produced synthetically. Extraction or isolation of the naturally-derived compound may be facilitated by


simple chemical reactions such as acidification, basification, ion exchange, hydrolysis, and salt formation as well as microbial fermentation, and the like. In embodiments, a taste modulating or salty taste modulating compound is derived from natural sources such as natural plant, fungi, and bacterial sources. Examples of such natural sources include, but are not limited to *Aesculus hippocastaneum*; *Alchemilla xanthochlora*; *Angelica archangelica*; *Apocynum cannabinum*; *Azadirachta indica*; *Actinomycete* bacteria (Strain code: 01702axxx000002); *Capsicum annuum*; *Cimicifuga racemosa*; *Commiphora mukul*; *Embelia ribes*; *Evodia rutaecarpa*; *Ferula assa-foetida*; *Fungi* (Strain code: 02295fxxx000001; Strain code: 01469fxxx000005); *Gleditschia australis*; *Kaempferia galanga*; *Lavandula officinalis*; *Marrubium vulgare*; *Mesua ferrea*; *Nephelium cuspidatum*; *Orthosiphon stamineus*; *Persea gratissima*; *Petroselinum sativum*; *Piper longum*; *Pithecoctenium echinatum*; *Podophyllum peltatum*; *Psidium guajava*; *Ricinus communis*; *Salvia miltiorrhiza*; *Schisandea chinensis*; *Teclea trichocarpa*; *Vitex agnus*; *Xysmalobium undulatum*; *Yucca gloriosa*; *Zanthoxylum piperitum*; *Zingiber officinalis*; and others. In embodiments, one or more compounds derived from *Persea gratissima* are combined with one or more compounds derived from *Kaempferia galanga* or one or more compounds derived from *Capsicum annuum*; and others.

[0290] As used herein, an "isolated" or "purified" compound is a compound that is substantially separated from other components of the source of the compound. For example, if the source of the compound is a natural product, an isolated or purified compound may be a compound that is separated from its naturally occurring environment. If the compound is synthesized, the compound may be separated from unreacted reagents, reaction byproducts, solvents, or the like.

[0291] As used herein a "synthetic compound" is a compound that is synthesized via chemical reaction *in vitro*. A compound that is "synthesized" is a synthetic compound. A synthesized compound may be identical to a compound derived from a natural product.

[0292] For the purposes of this disclosure, reference to a compound includes reference to salts of the compound, hydrates of the compound, polymorphs of the compound, isomers of the compound (including constitutional isomers and stereoisomers such as enantiomers and diasteriomers), and the like.

[0293] For the purposes of the present disclosure, it will be understood that a ring structure having a structure of




or the like, will be considered to be aromatic.

[0294] It will be understood that compounds disclosed herein may be glycosylated or may be substituted with one or more saccharides. In various embodiments, specific or generic


compounds substituted with one or more saccharides are disclosed. However, it will be understood that other saccharide substitutions are possible and are contemplated.

[0295] As used herein, a "saccharide" is a monosaccharide or an oligosaccharide. A monosaccharide may be a diose, a triose, a tetrose, a pentose, a hexose, a heptose, and so one. Monosaccharides include aldoses and ketoses. Examples of monosaccharides include glyceraldehyde, dihydroxyacetone, erythrose, threose, erythrulose, arabinose, lyxose, ribose, xylose, ribulose, xylulose, allose, altrose, galactose, glucose, gulose, idose, nannose, talose, fructose, psicose, sorbose, tagatose, mannoheptulose, sedoheptulose, 2-keto-3-deoxy-manno-actonate, and sialose. Monosaccharides may be acyclic or cyclic. Cyclic isomers include furanoses and pyranoses.

[0296] As used herein, "monosaccharide" includes deoxygenated variants of monosaccharides that are deoxygenated at one or more positions. As used herein, "monosaccharide" includes monosaccharides having carbon atoms of a monosaccharide chain or ring that are substituted with one or more of the following: H(H), CH₂OH, OH, COOH, OCOR¹⁰⁰, CH₃, OCH₃, C(CH₃)₂OH, and




where R¹⁰⁰ is selected from the group consisting of




[0297] An "oligosaccharide" is a chain of two or more monosaccharides where each monosaccharide is bound by a glycosidic bond.


[0298] As used herein, "saccharidyl" means a monosaccharide or oligosaccharide substituent. The monosaccharide or oligosaccharide substituent may be a terminal substituent or an internal substituent. That is, a saccharidyl group may be bound to one or more parent compounds (e.g., parent structure 1-saccharidyl or parent structure 1-saccharidyl-parent structure 2).

[0299] For purposes of example, a generic structure representing a monosaccharide is presented below:







where one of A', B', D', E', F' and G' is O and where each of the rest of A', B', D', E', F' and G' is independently selected from the group consisting of CH₂, CHO, CHCH₂OH, CHCH₃, CHCOOH, CHOCOR¹⁰¹, CHOCH₃, CHC(CH₃)₂OH, and




where R¹⁰¹ is selected from the group consisting of




[0300] For purposes of Example, a generic structure representing an embodiment of an oligosaccharide that is a disaccharide is presented below:



(i) where (a) one of A', B', D', E', and F' is O, (b) one of A'', B'', D'', E'', and F'' is O, and (c) each of the rest of A', B', D', E', F', A'', B'', D'', E'', and F'' is independently selected from the group consisting of CH₂, CHO, CHCH₂OH, CHCH₃, CHCOOH, CHOCOR¹⁰², CHOCH₃, CHC(CH₃)₂OH, and




where R¹⁰² is selected from the group consisting of



and (ii) where x' and y' are independently selected from zero or 1. Typically, the oxygen heteroatom of a monosaccharide ring will be in an ortho or meta position relative to a


saccharide substitution.

[0301] For purposes of example, a generic structure representing an embodiment of an oligosaccharide that is a trisaccharide is presented below:




1. (i) where (a) one of A'', B'', D'', E'', and F'' is O, (b) one of A'', B'', D'', E'', and F'' is O, and (c) each of the rest of A', B', D', A'', B'', D'', E'', F'', A''', B''', D''', E''', and F''' is independently selected from the group consisting of CH_2 , CHOH , CHCH_2OH , CHCH_3 ,


CHCOOH , CHOCOR^{103} , CHOCH_3 , $\text{CHC}(\text{CH}_3)_2\text{OH}$, and



where R^{103} is selected from the group consisting of





and




2. and (ii) where f', g', x' and y' are independently selected from zero or 1. In the structure depicted above, the oxygen heteroatom of a monosaccharide ring is in an ortho position relative to the other saccharide substitutions. Of course, trisaccharides (or tetra-, penta-, etc-saccharides) where the where substitutions of the other saccharides are ortho/meta, ortho/para, meta/para or meta/meta relative to the oxygen of the central ring are also possible. Typically, trisaccharides (or tetra-, penta-, etc-saccharides) will be ortho/ortho, meta/meta, or ortho/meta regarding the saccharide substitutions relative to the oxygen heteroatom of a central ring.

[0302] For purposes of example, a generic structure representing an embodiment of an oligosaccharide that is a tetrasaccharide is presented below:






1. (i) where (a) one of A'', B'', D'', E'', and F''' is O, (b) one of A'', B'', D'', E'', and F''' is O, (c) one of A'', B'', D'', E'', and F''' is O, and (d) each of the rest of A', B', A'', B'', D'', E'', F'', A'', B'', D'', E'', F'', A''', B''', D''', E''', and F'''' is independently selected from the group consisting of CH_2 , CHOH , CHCH_2OH , CHCH_3 , CHCOOH , CHOCOR^{104} , CHOCH_3 , $\text{CHC}(\text{CH}_3)_2\text{OH}$, and



where R^{104} is selected from the group consisting of



2. and (ii) where f, g', p', q', x' and y' are independently selected from zero or 1. In the structure depicted above, the oxygen heteroatom of a monosaccharide ring is in an ortho position relative to the other saccharide substitutions. The example of a tetrasaccharide depicted below is a branched chain saccharide. However, tetrasaccharides (or penta-, hexa, etc-saccharides) may have linear chains.

[0303] It will be understood that the structures of the saccharides presented above are examples of saccharides and that other saccharides are contemplated herein. Any compound described herein may be optionally saccharidyl substituted at any suitable position. Examples of some saccharidyl substituents are presented herein below (e.g., with regard to compounds 34-39, 41, 48, 51, 76, 77, and 90-92). However, it will be understood that similar compounds with different saccharide substitutions are contemplated herein.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO2010124905A1 [0005]

Non-patent literature cited in the description


- **K.A. REDDY et al.** Journal of Food Protection, International Association for Food Protection, 1991, vol. 54, 2138-150 [0005]
- Compensation for sodium: The low-salt solution Prepared Foods Gorman Pub. 19890000 vol. 158, 97-98 [0005]
- **G.-H. LEEA** salt substitute with low sodium content from plant aqueous extracts Food Research International, 2011, vol. 44, 2537-543 [0005]
- Reducing salt: A challenge for the meat industry **E. DESMOND** Meat Science Elsevier Science 20060000 vol. 74, 188-196 [0005]
- **MEILGAARD M.CIVILLE G.V.CARR B.T** Sensory Evaluation Techniques CRC Press 20070000147-152 [0046]
- **HOOTMAN RC** Manual 13 MNL13 Manual on Descriptive Analysis Testing for Sensory Evaluation ASTM 19920000 [0046]
- STP758 Guidelines for the Selection and Training of Sensory Panel Members ASTM 19810000 [0046]
- **MUNOZ A.MCIVILLE, G.V.MLN13:** The Spectrum Descriptive Analysis Method ASTM 19920000 [0046] [0046]

Patentkrav**1. Fødevareprodukt omfattende:**

mindst én bestanddel;

mindst ét salt der bibringer et indtryk af en saltagtig smag; og

5 mindst én forbindelse ifølge formel **B3**:




hvor:

R¹ og R² hver uafhængigt er OH eller C₁-C₃ alkoxy, og


R³ er valgt fra gruppen bestående af

10 (i) C₁₀-C₂₀ usubstitueret lige eller forgrenet kæde alkenyl med én eller flere dobbeltbindinger; og

(ii)



15 hvor X, Y og Z uafhængigt er CH, CH₂, CO eller CHOR⁵ hvor R⁵ er H eller C₁-C₃ alkyl, forudsat at hvis en af X eller Y er CH så er Z også CH, og hvor R⁴ er C₁-C₈ lige eller forgrenet usubstitueret alkyl eller



20 hvor n er 1-5, forudsat at hvis R⁴ er C₁-C₈ lige eller forgrenet kæde alkyl, så er Y CHOR⁵ hvor R⁵ er C₁-C₃ alkyl;
hvor fødevareproduktet ikke eksisterer i naturen;
og hvor forbindelsen med formel B3 enten er syntetiseret eller afledt

af et naturprodukt og hvis afledt fra et naturprodukt er det separeret fra dets naturligt forekommende miljø.

2. Fødevareprodukt ifølge krav 1, hvor R^1 er C_1 - C_3 alkoxy og R^2 er OH.


5

3. Fødevareprodukt ifølge krav 2, hvor R^1 er methoxy.

4. Fødevareprodukt ifølge krav 2 eller krav 3, hvor R^1 er substitueret ved 3-positionen, R^2 er substitueret ved 4-positionen, og R^3 er substitueret ved 1-


10 positionen.

5. Fødevareprodukt ifølge et hvilket som helst af kravene 1-4, hvor R^3 er



hvor X, Y og Z uafhængigt er CH, CH_2 , CO eller $CHOR^5$ hvor R^5 er H eller C_1 - C_3


15 alkyl, forudsat at hvis en af X eller Y er CH så er Z også CH, og hvor R^4 er C_1 - C_8 lige eller forgrenet kæde usubstitueret alkyl eller



hvor n er 1-5, forudsat at hvis R^4 er C_1 - C_8 lige eller forgrenet kæde alkyl, så er Y $CHOR^5$ hvor R^5 er C_1 - C_3 alkyl.

20

6. Fødevareprodukt ifølge krav 5, hvor R^4 er



25 **7.** Fødevareprodukt ifølge krav 6, hvor Z er CH eller CH_2 .


8. Fødevareprodukt ifølge krav 6 eller krav 7, hvor X er $CHOR^5$.

9. Fødevareprodukt ifølge krav 8, hvor R^5 er H.

10. Fødevareprodukt ifølge et hvilket som helst af kravene 6-9, hvor Y er CO.


5 11. Fødevareprodukt ifølge et hvilket som helst af kravene 6-10, hvor n er 2.

12. Fødevareprodukt ifølge krav 5, hvor R^3 er





10

13. Fødevareprodukt ifølge krav 1, hvor forbindelsen med formel B3 er valgt fra gruppen bestående af:




15


(8),



(11),



og



5

14. Fødevareprodukt ifølge et hvilket som helst af kravene 1 til 13, hvor den mindst ene forbindelse med formel B3 er til stede i fødevareproduktet i en mængde tilstrækkelig til at øge et indtryk af saltagtig smag af fødevareproduktet.

10 **15.** Fødevareprodukt ifølge et hvilket som helst af kravene 1 til 14, hvor fødevareproduktet har et vandindhold på mindst omkring 30 vægtprocent.

DRAWINGS

Conc.	12	13	83	10	37	36	45	18	56	82	3	86	43	48	53	44	62	43
12	1 ppm																	
13	10 ppm	3.4																
83	1 ppm	3.2, 4.2	3.7, 4.5															
10	1 ppm	3.9	3.1	3.9														
37	10 ppm	3.0, 4.8	3.5	4.1	4.1													
36	0.1 ppm	3.9	3.8	3.5	3.3, 3.4	3.1												
45	10 ppm	3.6, 3.9	3.6, 4.1	2.4, 3.4	3.4	3.3	4											
18	10 ppm	2.6, 4.0	3.1	3.2, 3.9	3.3	3.8	3.4	2.7, 3.3										
56	10 ppm	2.6	3.4	3.8	4.1	3.4, 4.4	3.1, 3.7	3.1	3.4									
22	1 ppm					3.2				3.4								
3	0.1 ppm						3.5											
83	2 ppm			3.8		3.2				3.5								
43	10 ppm				3.3					3								
48	10 ppm					3.4	3.3				3.3							
53	1 ppm							2.3				3.0						
44	10 ppm		3.2				3.3	3.1							3	2.8		
52	10 ppm				3.0	3.1					3.0	2.3						
43	10 ppm		3.1				3.2					3.1	3.2		3.1			

FIG. 1

26