

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 October 2007 (18.10.2007)

PCT

(10) International Publication Number
WO 2007/115821 A3

(51) International Patent Classification:

C07D 211/26 (2006.01) **A61K 31/439** (2006.01)
C07D 211/96 (2006.01) **A61P 3/10** (2006.01)
C07D 471/08 (2006.01)

(21) International Application Number:

PCT/EP2007/003185

(22) International Filing Date: 10 April 2007 (10.04.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

0607309.2 11 April 2006 (11.04.2006) GB
06122445.7 17 October 2006 (17.10.2006) EP

(71) Applicant (for all designated States except US): **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BAESCHLIN, Daniel, Kaspar** [CH/CH]; Austr. 1A, CH-4144 Arlesheim (CH). **FENTON, Garry** [GB/GB]; Argenta Discovery Ltd., Units 7/9, Spire Green Centre, Flexmeadow, Harlow Essex CM19 5TR (GB). **NAMOTO, Kenji** [JP/CH]; Kandererstrasse 29, CH-4057 Basel (CH). **OSTERMANN, Nils** [DE/DE]; Mühlenstrasse 43, 79589 Binzen (DE). **SEDRANI, Richard** [LU/CH]; Herrenggrabenweg 15, CH-4054 Basel (CH). **SIROCKIN, Finton** [FR/FR]; 17 Rue Des Asperges, F-68730 St. Louis (FR).

(74) Agent: **LIGIBEL, Jean-Marc**; Novartis AG, Corporate Intellectual Property, CH-4002 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))


Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

10 July 2008

(54) Title: ORGANIC COMPOUNDS



(57) Abstract: The present invention relates to compounds of the formula; and their use in therapy.

- 1 -

Organic Compounds

Field of the Invention

The present invention relates to compounds and their use in therapy.

Background to the Invention

Dipeptidylpeptidase-IV (DPP-IV) is a serine protease which cleaves N-terminal dipeptides from a peptide chain containing, in general, a proline residue in the penultimate position. DPP-IV is widely expressed in mammalian tissue as a type II integral membrane protein. The protease is expressed on the surface of differentiated epithelial cells of the intestine, liver, kidney proximal tubules, prostate, corpus luteum, and on leukocyte subsets such as lymphocytes and macrophages. A soluble form of the enzyme is found in serum that has structure and function identical to the membrane-bound form of the enzyme but lacks the hydrophobic transmembrane domain.

DPP-IV has many physiologically relevant substrates including chemokines (e.g. eotaxin and macrophage-derived chemokine), neuropeptides (e.g. neuropeptide Y and substance P), vasoactive peptides, and incretins (e.g. GLP-1 and GIP). GLP-1 (glucagon-like peptide-1) is a hormone produced in the L cells of the distal small intestine in response to ingested nutrients. GLP-1 receptor binding on various tissues stimulates insulin gene expression, biosynthesis and glucose-dependent insulin secretion, inhibits glucagon secretion, promotes satiety, slows gastric emptying and promotes growth of pancreatic beta cells.

Although the biological role of DPP-IV in mammalian systems has not been completely established, it is believed to play an important role in neuropeptide metabolism, T-cell activation, attachment of cancer cells to the endothelium and the entry of HIV into lymphoid cells. It has also been discovered that DPP-IV is responsible for inactivating glucagon-like peptide-1 (GLP-1). Since GLP-1 is a major stimulator of pancreatic insulin secretion and has direct beneficial effects on glucose disposal, DPP-IV inhibition appears to represent an attractive approach for treating, for example, non-insulin-dependent diabetes mellitus (NIDDM).


- 2 -

DPP-IV has also been shown to play a part in the immune response. Expressed by T-CD4+ lymphocytes, where it is synonymous with the antigen CD26, DPP-IV plays an important part in the mechanism of transplant rejection (Transplantation 1997, 63 (10), 1495-500). By allowing more selective suppression of the immune response, inhibition of DPP-IV accordingly represents an extremely promising approach in the prevention of transplant rejection in transplant patients.

Inhibitors of DPP-IV are described *inter alia* in WO-A-03/000180, WO-A-000181, WO-A-004498, WO-A-03/082817, WO-A-04/032836, WO-A-04/007468, EP1679069 and WO-A-05/121089.

Summary of the Invention

A first aspect of the invention is a compound of the Formula (I):



(I)

wherein

the asterisk * designates a chiral centre of (*R*) or (*S*) configuration;

V is absent or is ethylene;

W is -C(O)- or -S(O)-;

X is a linker having 1 to 12 (e.g. 1 to 6) in-chain atoms and comprising one or more linkages selected from -O-, -C(O)-, -S(O)-, -N(R⁹)- and hydrocarbylene optionally substituted with 1, 2, 3, 4 or 5 R¹¹;

Y is a linker selected from -O-, -N(R⁹)-, -C(O)-, -C(O)O-, -C(O)N(R⁹)-, -S(O)- and -S(O)N(R⁹)-;

- 3 -

R^1 is selected from hydrogen; $-N(R^9)(R^{10})$; hydrocarbyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} ; hydrocarbyloxy optionally substituted with 1, 2, 3, 4 or 5 R^{11} ; and $-(CH_2)_k$ -heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

or, when Y is $-N(R^9)-$, R^1 and R^9 taken together with the nitrogen atom to which they are attached may form a heterocycle, wherein said heterocycle is bound to X via said nitrogen atom and is optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

R^2 and R^3 are each independently selected from R^8 , $-OR^8$, $-C(O)R^8$, $-C(O)OR^8$ and $-S(O)_2R^8$;

R^4 and R^5 are each independently selected from hydrogen, hydroxy, halogen and C_{1-6} alkyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

R^6 is aryl or heteroaryl, either of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

R^8 is selected from hydrogen; hydrocarbyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} ; and $-(CH_2)_k$ -heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

R^9 and R^{10} are each independently selected from R^8 , $-OR^8$, $-C(O)R^8$, $-C(O)OR^8$ and $-S(O)_2R^8$; or R^9 and R^{10} taken together with a nitrogen atom to which they are attached form heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

each R^{11} is independently selected from R^{12} ; hydrocarbyl optionally substituted with 1, 2, 3, 4 or 5 R^{12} ; and $-(CH_2)_k$ -heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{12} ;

R^{12} is independently selected from halogen, trifluoromethyl, cyano, nitro, oxo, $=NR^{13}$, $-OR^{13}$, $-C(O)R^{13}$, $-C(O)N(R^{13})R^{14}$, $-C(O)OR^{13}$, $-OC(O)R^{13}$, $-S(O)_2R^{13}$, $-S(O)_2N(R^{13})R^{14}$, $-N(R^{13})R^{14}$, $-N(R^{13})N(R^{13})R^{14}$, $-N(R^{13})C(O)R^{14}$ and $-N(R^{13})S(O)_2R^{13}$;

R^{13} and R^{14} are each independently hydrogen or selected from hydrocarbyl and $-(CH_2)_k$ -heterocyclyl, either of which is optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from halogen, cyano, amino, hydroxy, C_{1-6} alkyl and C_{1-6} alkoxy; or

- 4 -

R^{13} and R^{14} are each independently hydrogen or selected from hydrocarbyl and $-(CH_2)_k$ -heterocyclyl, either of which is optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from oxo, halogen, cyano, amino, hydroxy, C_{1-6} alkyl and C_{1-6} alkoxy;

k is 0, 1, 2, 3, 4, 5 or 6; and

l is 0, 1, or 2;

or a pharmaceutically acceptable salt or prodrug thereof.

In some embodiments, when X is C_{1-6} alkylene, $-O-C_{1-6}$ alkylene- or $-N(R^9)-C_{1-6}$ alkylene- and Y is $-O-$, $-S-$ or $-N(R^9)-$, said C_{1-6} alkylene linkage of X is substituted with 1, 2, 3, 4 or 5 R^{11} , wherein at least one of said R^{11} is other than halogen or C_{1-6} alkyl.

A second aspect of the invention is a compound of the invention for therapeutic use.

Another aspect of the invention is a pharmaceutical formulation comprising a compound of the invention and, optionally, a pharmaceutically acceptable diluent or carrier.

A further aspect of the invention is a product comprising a compound of the invention and a therapeutic agent; as a combined preparation for simultaneous, separate or sequential use in therapy.

Another aspect of the invention is the use of a compound of the invention for the manufacture of a medicament for the treatment or prevention of a disease or condition selected from non-insulin-dependent diabetes mellitus, arthritis, obesity, allograft transplantation, calcitonin-osteoporosis, heart failure, impaired glucose metabolism or impaired glucose tolerance, neurodegenerative diseases, cardiovascular or renal diseases, and neurodegenerative or cognitive disorders.

Another aspect of the invention is the use of a compound of the invention for the manufacture of a medicament for producing a sedative or anxiolytic effect, attenuating post-

- 5 -

surgical catabolic changes or hormonal responses to stress, reducing mortality and morbidity after myocardial infarction, modulating hyperlipidemia or associated conditions, or lowering VLDL, LDL or Lp(a) levels.

Another aspect of the invention is a method of treating or preventing a disease or condition in a patient, which comprises administering a therapeutically effective amount of a compound of the invention.

The compounds of the invention can exist in different forms, such as free acids, free bases, esters and other prodrugs, salts and tautomers, for example, and the disclosure includes all variant forms of the compounds.

The extent of protection includes counterfeit or fraudulent products which contain or purport to contain a compound of the invention irrespective of whether they do in fact contain such a compound and irrespective of whether any such compound is contained in a therapeutically effective amount.

Included in the scope of protection are packages which include a description or instructions which indicate that the package contains a species or pharmaceutical formulation of the invention and a product which is or comprises, or purports to be or comprise, such a formulation or species. Such packages may be, but are not necessarily, counterfeit or fraudulent.

Features, integers, characteristics, compounds, chemical moieties or groups described in conjunction with a particular aspect, embodiment or example of the invention are to be understood to be applicable to any other aspect, embodiment or example described herein unless incompatible therewith.

Description of Various Embodiments

Hydrocarbyl

The term "hydrocarbyl" as used herein includes reference to a moiety consisting exclusively of hydrogen and carbon atoms; such a moiety may comprise an aliphatic and/or an aromatic moiety. The moiety may comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18,

- 6 -

19 or 20 carbon atoms. Examples of hydrocarbonyl groups include C₁₋₆ alkyl (e.g. C₁, C₂, C₃ or C₄ alkyl, for example methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl or tert-butyl); C₁₋₆ alkyl substituted by aryl (e.g. benzyl) or by cycloalkyl (e.g. cyclopropylmethyl); cycloalkyl (e.g. cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl); aryl (e.g. phenyl, naphthyl or fluorenyl) and the like. When hydrocarbonyl is a cycloalkyl it can be attached to the chemical moiety in the form of a spiro substitution.

Alkyl

The terms "alkyl" and "C₁₋₆ alkyl" as used herein include reference to a straight or branched chain alkyl moiety having 1, 2, 3, 4, 5 or 6 carbon atoms. This term includes reference to groups such as methyl, ethyl, propyl (n-propyl or isopropyl), butyl (n-butyl, sec-butyl or tert-butyl), pentyl, hexyl and the like. In particular, alkyl may have 1, 2, 3 or 4 carbon atoms.

Alkenyl

The terms "alkenyl" and "C₂₋₆ alkenyl" as used herein include reference to a straight or branched chain alkyl moiety having 2, 3, 4, 5 or 6 carbon atoms and having, in addition, at least one double bond, of either E or Z stereochemistry where applicable. This term includes reference to groups such as ethenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 1-hexenyl, 2-hexenyl and 3-hexenyl and the like.

Alkynyl

The terms "alkynyl" and "C₂₋₆ alkynyl" as used herein include reference to a straight or branched chain alkyl moiety having 2, 3, 4, 5 or 6 carbon atoms and having, in addition, at least one triple bond. This term includes reference to groups such as ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-pentylnyl, 2-pentylnyl, 3-pentylnyl, 1-hexynyl, 2-hexynyl and 3-hexynyl and the like.

Alkoxy

The terms "alkoxy" and "C₁₋₆ alkoxy" as used herein include reference to -O-alkyl, wherein alkyl is straight or branched chain and comprises 1, 2, 3, 4, 5 or 6 carbon atoms. In one class of embodiments, alkoxy has 1, 2, 3 or 4 carbon atoms. This term includes reference to

- 7 -

groups such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, tert-butoxy, pentoxy, hexoxy and the like.

Cycloalkyl

The term "cycloalkyl" as used herein includes reference to an alicyclic moiety having 3, 4, 5, 6, 7 or 8 carbon atoms. The group may be a bridged or polycyclic ring system. More often cycloalkyl groups are monocyclic. This term includes reference to groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, norbornyl, bicyclo[2.2.2]octyl and the like.

Aryl

The term "aryl" as used herein includes reference to an aromatic ring system comprising 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16 ring carbon atoms. Aryl is often phenyl but may be a polycyclic ring system, having two or more rings, at least one of which is aromatic. This term includes reference to groups such as phenyl, naphthyl, fluorenyl, azulenyl, indenyl, anthryl and the like.

Carbocyclyl

The term "carbocyclyl" as used herein includes reference to a saturated (e.g. cycloalkyl) or unsaturated (e.g. aryl) ring moiety having 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16 carbon ring atoms. In particular, carbocyclyl includes a 3- to 10-membered ring or ring system and, in particular, a 5- or 6-membered ring, which may be saturated or unsaturated. A carbocyclic moiety is, for example, selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, norbornyl, bicyclo[2.2.2]octyl, phenyl, naphthyl, fluorenyl, azulenyl, indenyl, anthryl and the like.

Heterocyclyl

The term "heterocyclyl" as used herein includes reference to a saturated (e.g. heterocycloalkyl) or unsaturated (e.g. heteroaryl) heterocyclic ring moiety having from 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16 ring atoms, at least one of which is selected from nitrogen, oxygen, phosphorus, silicon and sulphur. In particular, heterocyclyl includes a 3- to 10-membered ring or ring system and more particularly a 5- or 6-membered ring, which may

be saturated or unsaturated. When the heterocyclyl is a heterocycloalkyl it can be attached to the chemical moiety in the form of a spiro substitution.

A heterocyclic moiety is, for example, selected from oxiranyl, aziranyl, 1,2-oxathiolanyl, imidazolyl, thienyl, furyl, tetrahydrofuryl, pyranal, thiopyranal, thianthrenyl, isobenzofuranyl, benzofuranyl, chromenyl, 2*H*-pyrrolyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolyl, imidazolidinyl, benzimidazolyl, pyrazolyl, pyrazinyl, pyrazolidinyl, thiazolyl, isothiazolyl, dithiazolyl, oxazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, piperidyl, piperazinyl, pyridazinyl, morpholinyl, thiomorpholinyl, especially thiomorpholino, indoliziny, 1,3-Dioxo-1,3-dihydro-isoindolyl, 3*H*-indolyl, indolyl, benzimidazolyl, cumaryl, indazolyl, triazolyl, tetrazolyl, purinyl, 4*H*-quinoliziny, isoquinolyl, quinolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, decahydroquinolyl, octahydroisoquinolyl, benzofuranyl, dibenzofuranyl, benzothiophenyl, dibenzothiophenyl, phthalazinyl, naphthyridinyl, quinoxalyl, quinazolinyl, quinazolinyl, cinnolyl, pteridinyl, carbazolyl, β -carbolinyl, phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, furazanyl, phenazinyl, phenothiazinyl, phenoxazinyl, chromenyl, isochromanyl, chromanyl, 3,4-dihydro-2*H*-isoquinolin-1-one, 3,4-dihydro-2*H*-isoquinolinyl, and the like.

Heterocycloalkyl

The term "heterocycloalkyl" as used herein includes reference to a saturated heterocyclic moiety having 3, 4, 5, 6 or 7 ring carbon atoms and 1, 2, 3, 4 or 5 ring heteroatoms selected from nitrogen, oxygen, phosphorus and sulphur. The group may be a polycyclic ring system but more often is monocyclic. This term includes reference to groups such as azetidiny, pyrrolidinyl, tetrahydrofuranyl, piperidinyl, oxiranyl, pyrazolidinyl, imidazolyl, indolizidinyl, piperazinyl, thiazolidinyl, morpholinyl, thiomorpholinyl, quinolizidinyl, tetrahydropyranal, tetrahydrofuranyl, 1,3-dihydro-isoindolyl, 1,3-Dioxo-1,3-dihydro-isoindolyl, 3,4-dihydro-2*H*-isoquinolinyl, 3,4-dihydro-2*H*-isoquinolinyl-1-one and the like.

Heteroaryl

The term "heteroaryl" as used herein includes reference to an aromatic heterocyclic ring system having 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16 ring atoms, at least one of which is selected from nitrogen, oxygen and sulphur. The group may be a polycyclic ring system, having two or more rings, at least one of which is aromatic, but is more often monocyclic.


This term includes reference to groups such as pyridazinyl, pyrimidinyl, furanyl, benzo[b]thiophenyl, thiophenyl, pyrrolyl, imidazolyl, pyrrolidinyl, pyridinyl, benzo[b]furanyl, pyrazinyl, purinyl, indolyl, benzimidazolyl, quinolinyl, phenothiazinyl, triazinyl, phthalazinyl, 2H-chromenyl, oxazolyl, isoxazolyl, thiazolyl, isoindolyl, indazolyl, purinyl, isoquinolinyl, quinazolinyl, pteridinyl and the like.

Halogen

The term "halogen" as used herein includes reference to F, Cl, Br or I. In a particular, halogen may be F or Cl, of which F is more common.

Spiro

The term "spiro" as used herein includes 3- to 6-cycloalkyl or 5- to 6-heterocycloalkyl groups which can optionally be substituted by 1, 2, 3 or 4, R^{13} . Non limitative examples of spiro groups are;



Substituted

The term "substituted" as used herein in reference to a moiety means that one or more, especially up to 5, more especially 1, 2 or 3, of the hydrogen atoms in said moiety are replaced independently of each other by the corresponding number of the described substituents. The term "optionally substituted" as used herein means substituted or unsubstituted.

- 10 -

It will, of course, be understood that substituents are only at positions where they are chemically possible, the person skilled in the art being able to decide (either experimentally or theoretically) without inappropriate effort whether a particular substitution is possible. For example, amino or hydroxy groups with free hydrogen may be unstable if bound to carbon atoms with unsaturated (e.g. olefinic) bonds. Additionally, it will of course be understood that the substituents described herein may themselves be substituted by any substituent, subject to the aforementioned restriction to appropriate substitutions as recognised by the skilled man.

Pharmaceutically acceptable


The term "pharmaceutically acceptable" as used herein includes reference to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings or animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. This term includes acceptability for both human and veterinary purposes.

Independently

Where two or more moieties are described as being "each independently" selected from a list of atoms or groups, this means that the moieties may be the same or different. The identity of each moiety is therefore independent of the identities of the one or more other moieties.

Compounds

The invention provides compounds of the Formula (I):



(I)


wherein

the asterisk * designates a chiral centre of (*R*) or (*S*) configuration; and

V, W, X, Y, R¹, R², R³, R⁴, R⁵ and R⁶ are as defined herein;

or a pharmaceutically acceptable salt or prodrug thereof.

As indicated above, the stereochemical configuration of the chiral centre indicated by * (i.e. the carbon atom to which the group -N(R²)(R³) is attached) may be (*R*) or (*S*). Of particular mention are compounds of the invention in which the stereochemical configuration at said carbon atom is (*R*), i.e. compounds of the following Formula:



(I, (*R*))

or a pharmaceutically acceptable salt or prodrug thereof.

A compound of the invention may be in the form of a racemate or in a substantially pure form (e.g. a form having a purity of greater than 80% purity, in particular greater than 90%, 95% or


- 12 -

99%) of a single enantiomer or diastereomer. Thus, one embodiment of the invention is a substantially pure form of a compound of Formula (I, (*R*)).


Embodiments of the invention are described below. It will be appreciated that the features specified in each embodiment may be combined with other specified features, to provide further embodiments.

V

V is absent or is ethylene. The invention therefore includes compounds of the following Formulae:



(II)




(III)

or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

Where V is ethylene (i.e. compounds of Formula (III)), the relative orientation of the ethylene bridge and the hydrogen atom located *p*- to the nitrogen atom of the piperidine ring is often *exo*, as illustrated below:

- 13 -



The term “exo” in this context means that the ethylene bridge and the hydrogen atom shown in the Formula above are on the same side of the piperidine ring.

The present invention also concerns the herein described compounds and claims wherein the bridged piperidine ring is in the configuration *exo* as described in the above paragraph.

-W-X-Y-

In Formula (I), W is -C(O)- or -S(O)_n-; X is a linker having 1 to 12 (e.g. 1 to 6) in-chain atoms and comprising one or more linkages selected from -O-, -C(O)-, -S(O)_n-, -N(R⁹)- and hydrocarbylene optionally substituted with 1, 2, 3, 4 or 5 R¹¹; and Y is a linker selected from -O-, -N(R⁹)-, -C(O)-, -C(O)O-, -C(O)N(R⁹)-, -S(O)_n- and -S(O)_nN(R⁹)-.

W is usually -C(O)- or -S(O)₂-.

Where X comprises one or more hydrocarbylene linkages, the or each linkage may be aliphatic and/or carbocyclic (e.g. cycloalkylene). Of particular mention are aliphatic, e.g. alkylenic, hydrocarbylene linkages. Aliphatic linkages are usually C₁₋₆ aliphatic linkages, examples including C₁₋₆ alkylene linkages. Carbocyclylene is usually C₃₋₇ carbocyclylene, including cycloalkylene (e.g. C₃₋₆ cycloalkylene, especially cyclopropylene). In the case of carbocyclylene-containing linkages, at least one (usually 1, 2, 3 or 4, especially 1 or 2 but in other cases 3 or 4) of the in-ring atoms forms or is included in the linkage. A hydrocarbylene linkage is often aliphatic, in particular C₁₋₆ alkylene. In one class of compounds, X comprises a hydrocarbylene linkage which is directly bonded to W. In another class of compounds, X comprises an arylene (e.g. phenylene) linkage which is optionally substituted with 1, 2, 3, 4 or 5 R¹¹. In certain compounds, said arylene linkage is directly bonded in W. Of particular

- 14 -

mention are compounds comprising a phenylene linkage optionally substituted with 1, 2, 3, 4 or 5 R¹¹, wherein said phenylene linkage is directly bonded to W.

X may comprise at least one linkage selected from -N(R⁹)-, and aliphatic or cyclic hydrocarbylene (e.g. C₁₋₆ alkylene or cycloalkylene), either of which is optionally substituted with 1, 2, 3, 4 or 5 R¹¹. In particular, X may comprise at least one linker selected from -N(R⁹)- and C₁₋₆ alkylene (e.g. C₁, C₂ or C₃ alkylene) optionally substituted with 1, 2, 3, 4 or 5 R¹¹. When X comprises -N(R⁹)-, R⁹ is usually hydrogen or selected from C₁₋₆ alkyl (e.g. C₁, C₂, C₃ or C₄ alkyl), -(CH₂)_k-carbocyclyl and -(CH₂)_k-heterocyclyl, any of which is optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

Of mention are compounds in which X is a linker having 1 to 12 (e.g. 1 to 6) in-chain atoms and comprising one or more linkages selected from -O-, -C(O)-, -S(O)-, -N(R⁹)- and C₁₋₆ aliphatic (e.g. C₁₋₆ alkylene) optionally substituted with 1, 2, 3, 4 or 5 R¹¹;

In one embodiment, X is selected from one of the following linkers:

-X¹-;
 -X¹-X²-;
 -X¹-X²-X³-;
 -X¹-X²-X³-X⁴-;
 -X¹-X²-X³-X⁴-X⁵-;
 -X¹-X²-X³-X⁴-X⁵-X⁶-;
 -X¹-X²-X³-X⁴-X⁵-X⁶-X⁷-; and
 -X¹-X²-X³-X⁴-X⁵-X⁶-X⁷-X⁸-;

wherein X¹, X², X³, X⁴, X⁵, X⁶, X⁷ and X⁸ are each independently selected from -O-, -C(O)-, -S(O)-, -N(R⁹)- and hydrocarbylene (e.g. C₁₋₆ alkylene) optionally substituted with 1, 2, 3, 4 or 5 R¹¹. More usually, X is -X¹- or -X¹-X²-.


Of mention are compounds in which X¹ is hydrocarbylene (e.g. C₁₋₆ alkylene, C₂₋₆ alkenylene or carbocyclylene) optionally substituted with 1, 2, 3, 4 or 5 R¹¹. For example, X¹ may be C₁₋₆ alkylene optionally substituted with 1, 2, 3, 4 or 5 R¹¹. In particular, X¹ may be methylene or ethylene, either of which is optionally substituted with 1, 2, 3 or 4 substituents independently selected from halogen (e.g. fluorine or chlorine), amino and hydroxy. Alternatively, X¹ may be arylene, e.g. phenylene, optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

- 15 -

Also of mention is a class of compounds in which X^2 is $-N(R^9)-$, wherein R^9 is usually hydrogen or selected from C_{1-6} alkyl (e.g. C_1 , C_2 , C_3 or C_4 alkyl), $-(CH_2)_k$ -carbocyclyl and $-(CH_2)_k$ -heterocyclyl, any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . In particular, $-N(R^9)-$ may be $-NH-$ or $-N(CH_3)-$. In compounds where X^1 is arylene (e.g. phenylene), X^2 is often C_{1-6} alkylene (e.g. C_1 or C_2 alkylene) optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

Also included are compounds in which X^1 is C_{1-6} alkylene substituted, e.g. at the 1- or 2-position relative to W (which is typically carbonyl or sulphonyl), by at least one R^{11} . In this case, the at least one R^{11} is often selected from halogen, trifluoromethyl, cyano, nitro, oxo, $=NR^{13}$, $-OR^{13}$, $-C(O)R^{13}$, $-C(O)N(R^{13})R^{14}$, $-C(O)OR^{13}$, $-OC(O)R^{13}$, $-S(O)R^{13}$, $-S(O)N(R^{13})R^{14}$, $-N(R^{13})R^{14}$, $-N(R^{13})N(R^{13})R^{14}$, $-N(R^{13})C(O)R^{14}$, $-N(R^{13})S(O)R^{13}$, hydrocarbyl (which is other than C_{1-6} alkyl, and is often $-(CH_2)_k$ -carbocyclyl) optionally substituted with 1, 2, 3, 4 or 5 R^{12} ; and $-(CH_2)_k$ -heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{12} . In this case, R^{13} and R^{14} are often each independently selected from hydrogen, C_{1-6} alkyl, $-(CH_2)_k$ -carbocyclyl (e.g. phenyl, cyclopropyl or benzyl) and $-(CH_2)_k$ -heterocyclyl. Exemplary R^{11} moieties include carbamate, phenyl, benzyl, $-NH-C(O)-(C_{1-6} \text{ alkyl})$, oxo, sulphonamido, urea, thiourea and acyl groups.

Also included are compounds in which X^1 is an arylene preferably selected from;



In certain compounds, Y is $-C(O)-$, $-C(O)O-$, $-C(O)N(R^9)-$, $-S(O)-$, $-S(O)_2-$ or $-S(O)N(R^9)-$. In particular, Y is often $-C(O)-$, $-C(O)O-$, $-S(O)-$ or $-S(O)_2-$.


In some embodiments, when X is C_{1-6} alkylene, $-O-C_{1-6}$ alkylene- or $-N(R^9)-C_{1-6}$ alkylene-, Y is $-O-$, $-S-$ or $-N(R^9)-$ and R^7 is hydrogen, said C_{1-6} alkylene linkage of X is substituted with 1, 2, 3, 4 or 5 R^{11} , wherein at least one of said R^{11} is other than halogen or C_{1-6} alkyl. In this case, at least one R^{11} , e.g. the or each R^{11} is often independently selected from trifluoromethyl,

- 16 -

cyano, nitro, oxo, $=NR^{13}$, $-OR^{13}$, $-C(O)R^{13}$, $-C(O)N(R^{13})R^{14}$, $-C(O)OR^{13}$, $-OC(O)R^{13}$, $-S(O)_hR^{13}$, $-S(O)_hN(R^{13})R^{14}$, $-N(R^{13})R^{14}$, $-N(R^{13})N(R^{13})R^{14}$, $-N(R^{13})C(O)R^{14}$, $-N(R^{13})S(O)_hR^{13}$, hydrocarbonyl (which is other than C_{1-6} alkyl, and is often $-(CH_2)_k$ -carbocyclyl) optionally substituted with 1, 2, 3, 4 or 5 R^{12} ; and $-(CH_2)_k$ -heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{12} . In this case, R^{13} and R^{14} are often each independently selected from hydrogen, C_{1-6} alkyl, $-(CH_2)_k$ -carbocyclyl (e.g. phenyl, cyclopropyl or benzyl) and $-(CH_2)_k$ -heterocyclyl. Exemplary R^{11} moieties include carbamate, sulphonamido, urea, thiourea and acyl groups. Carbocyclyl and heterocyclyl may for example be 5, 6 or 7-membered saturated or unsaturated rings, e.g. phenyl.

The linker $-W-X-Y-$ often comprises 3, 4 or 5, especially 4 or 5, in-chain atoms.

Of particular mention are compounds of the following Formula:



(IV)

wherein X^1 is $-N(R^9)-$ or hydrocarbylene (e.g. C_{1-6} alkylene) optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

or a pharmaceutically acceptable salt or prodrug thereof.

With regard to Formula (IV), Y is often $-C(O)-$, $-C(O)O-$, $-C(O)N(R^9)-$, $-S(O)-$, $-S(O)_2-$ or $-S(O)_hN(R^9)-$. In particular, Y is often $-C(O)-$, $-C(O)-$, $-S(O)-$ or $-S(O)_2-$.

Examples of the linker $-W-X^1-Y-$ are described in the table below:

No.	W	X^1	Y
1	$-C(O)-$	$-N(R^9)-$	$-C(O)-$
2	$-C(O)-$	C_{1-6} alkylene	$-C(O)-$





- 17 -

3	-C(O)-	-N(R ⁹)-	-S(O) ₂ -
4	-C(O)-	C ₁₋₆ alkylene	-S(O) ₂ -
5	-S(O) ₁ -	-N(R ⁹)-	-C(O)-
6	-S(O) ₁ -	C ₁₋₆ alkylene	-C(O)-
7	-S(O) ₁ -	C ₁₋₆ alkylene	-S(O) ₂ -

Further examples of the linker -W-X¹-Y- are described in the table below:

No.	W	X ¹	Y
1	-C(O)-	C ₁₋₆ alkylene	-S(O) ₂ -
2	-S(O) ₁ -	C ₁₋₆ alkylene	-C(O)-

Further examples of the linker -W-X¹-Y- are described in the table below:

No.	W	X ¹	Y
1	-C(O)-	C ₁₋₆ alkylene	-C(O)-N(R ⁹)-
2	-C(O)-	-CH ₂ -	-C(O)-NH-
3	-C(O)-		-C(O)-
4	-C(O)-		-C(O)-
5	-C(O)-		-N(R ⁹)-
6	-C(O)-		-N(R ⁹)-
7	-C(O)-	arylene	-C(O)-
8	-C(O)-	C ₁₋₆ alkylene	-C(O)-

- 18 -

9	-C(O)-	arylene	-N(R ⁹)-
10	-C(O)-	-CH ₂ -	-C(O)-
11	-C(O)-	-CH(CH ₃)-	-C(O)-
12	-C(O)-	-C(CH ₃) ₂ -	-C(O)-
13	-C(O)-	-CH ₂ CH ₂ -	-C(O)-

In one embodiment in the above compounds wherein Y is -N(R⁹)-, R¹ and R⁹ taken together form a 5- or 6-membered heterocyclyl optionally substituted by 1, 2, 3, 4 or 5 R¹¹. The formed 5- or 6-membered heterocyclyl is for instance 1-methyl-imidazolidine-2,4-dione, imidazolidine-2,4-dione, thiazolidine-2,4-dione, pyrrolidine-2,5-dione, isoindole-1,3-dione, or pyrrolidiny-2-oxo.

Further examples of the linker -W-X¹-Y- are described in the table below:

No.	W	X ¹	Y
1	-C(O)-	-CH ₂ -	-S(O) ₂ -
2	-C(O)-	-CH(CH ₃)-	-S(O) ₂ -
3	-C(O)-	-C(CH ₃) ₂ -	-S(O) ₂ -
4	-C(O)-	-CH ₂ CH ₂ -	-S(O) ₂ -
5	-S(O) ₂ -	-CH ₂ -	-C(O)-
6	-S(O) ₂ -	-CH(CH ₃)-	-C(O)-
7	-S(O) ₂ -	-C(CH ₃) ₂ -	-C(O)-
8	-S(O) ₂ -	-CH ₂ CH ₂ -	-C(O)-

Further examples of the linker -W-X¹-Y- are described in the table below:

No.	W	X ¹	Y
1	-C(O)-	C ₁₋₆ alkylene	-C(O)-
2	-S(O) ₂ -	C ₁₋₆ alkylene	-S(O) ₂ -

Further examples of the linker -W-X¹-Y- are described in the table below:

No.	W	X ¹	Y
1	-C(O)-	-CH ₂ -	-C(O)-

- 19 -

2	-C(O)-	-CH(CH ₃)-	-C(O)-
3	-C(O)-	-C(CH ₃) ₂ -	-C(O)-
4	-C(O)-	-CH ₂ CH ₂ -	-C(O)-

Also of particular mention are compounds of the above Formula wherein X¹ is carbocyclylene (e.g. cycloalkylene or arylene) optionally substituted with 1, 2, 3, 4 or 5 R¹¹; or a pharmaceutically acceptable salt or prodrug thereof.

Further examples of the linker -W-X¹-Y- are described in the table below:

No.	W	X ¹	Y
1	-C(O)-	cycloalkylene	-S(O) ₂ -
2	-S(O) ₂ -	cycloalkylene	-C(O)-
3	-C(O)-	cycloalkylene	-C(O)-
4	-S(O) ₂ -	cycloalkylene	-S(O) ₂ -

Further examples of the linker -W-X¹-Y- are described in the table below:

No.	W	X ¹	Y
1	-C(O)-	cyclopropylene	-S(O) ₂ -
2	-S(O) ₂ -	cyclopropylene	-C(O)-
3	-C(O)-	cyclopropylene	-C(O)-
4	-S(O) ₂ -	cyclopropylene	-S(O) ₂ -

Further examples of the linker -W-X¹-Y- are described in the table below:


No.	W	X ¹	Y
1	-C(O)-	arylene	-S(O) ₂ -
2	-S(O) ₂ -	arylene	-C(O)-
3	-C(O)-	arylene	-C(O)-
4	-S(O) ₂ -	arylene	-S(O) ₂ -

Further examples of the linker -W-X¹-Y- are described in the table below:

- 20 -

No.	W	X ¹	Y
1	-C(O)-	phenylene	-S(O) ₂ -
2	-S(O) _r -	phenylene	-C(O)-
3	-C(O)-	phenylene	-C(O)-
4	-S(O) _r -	phenylene	-S(O) ₂ -

Also of mention are compounds of the following Formula:



(V)

wherein one of X¹ and X² is -N(R⁹)-; and the other is C₁₋₆ alkylene optionally substituted with 1, 2, 3, 4 or 5 R¹¹;

or a pharmaceutically acceptable salt or prodrug thereof.

With regard to Formula (V), Y is often -C(O)-, -C(O)O-, -C(O)N(R⁹)-, -S(O)-, -S(O)₂- or -S(O)N(R⁹)-. In particular, Y is often -C(O)-, -C(O)-, -S(O)- or -S(O)₂-.





Examples of the linker -W-X¹-X²-Y- are described in the table below:

No.	W	X ¹	X ²	Y
1	-C(O)-	C ₁₋₆ alkylene	-O-	-C(O)-
2	-C(O)-	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-
3	-C(O)-	-O-	C ₁₋₆ alkylene	-C(O)-
4	-C(O)-	-N(R ⁹)-	C ₁₋₆ alkylene	-C(O)-

- 21 -

No.	W	X ¹	X ²	Y
5	-C(O)-	C ₁₋₆ alkylene	-N(R ⁹)-	-S(O) ₂ -
6	-C(O)-	-O-	C ₁₋₆ alkylene	-S(O) ₂ -
7	-C(O)-	-N(R ⁹)-	C ₁₋₆ alkylene	-S(O) ₂ -
8	-S(O)-	C ₁₋₆ alkylene	-O-	-C(O)-
9	-S(O)-	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-
10	-S(O)-	-N(R ⁹)-	C ₁₋₆ alkylene	-C(O)-
11	-S(O)-	C ₁₋₆ alkylene	-N(R ⁹)-	-S(O) ₂ -
12	-S(O)-	-N(R ⁹)-	C ₁₋₆ alkylene	-S(O) ₂ -

Examples of the linker -W-X¹-X²-Y- are described in the table below:

1	-S(O)-	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-O-
2	-S(O)-	-CH ₂ CH ₂ -	-N(R ⁹)-	-C(O)-O-
3	-C(O)-	arylene	-N(R ⁹)-	-C(O)-
4	-C(O)-		-N(R ⁹)-	-C(O)-
5	-C(O)-		-N(R ⁹)-	-C(O)-
6	-C(O)-		C ₁₋₆ alkylene	-N(R ⁹)-
7	-C(O)-		-CH ₂ CH ₂ -	-N(R ⁹)-

Further examples of the linker -W-X¹-X²-Y- are described in the table below:

No.	W	X ¹	X ²	Y
1	-C(O)-	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-

- 22 -

2	-C(O)-	C ₁₋₆ alkylene	-N(R ⁹)-	-S(O) ₂ -
3	-S(O) ₂ -	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-
4	-S(O) ₂ -	C ₁₋₆ alkylene	-N(R ⁹)-	-S(O) ₂ -

Further examples of the linker -W-X¹-X²-Y- are described in the table below:

No.	W	X ¹	X ²	Y
1	-C(O)-	-CH ₂ -	-N(R ⁹)-	-C(O)-
2	-C(O)-	-CH ₂ -	-N(R ⁹)-	-S(O) ₂ -
3	-C(O)-	-CH(CH ₃)-	-N(R ⁹)-	-C(O)-
4	-C(O)-	-CH(CH ₃)-	-N(R ⁹)-	-S(O) ₂ -
5	-C(O)-	-C(CH ₃) ₂ -	-N(R ⁹)-	-C(O)-
6	-C(O)-	-C(CH ₃) ₂ -	-N(R ⁹)-	-S(O) ₂ -
7	-C(O)-	-CH ₂ CH ₂ -	-N(R ⁹)-	-C(O)-
8	-C(O)-	-CH ₂ CH ₂ -	-N(R ⁹)-	-S(O) ₂ -
9	-S(O) ₂ -	-CH ₂ -	-N(R ⁹)-	-C(O)-
10	-S(O) ₂ -	-CH ₂ -	-N(R ⁹)-	-S(O) ₂ -
11	-S(O) ₂ -	-CH(CH ₃)-	-N(R ⁹)-	-C(O)-
12	-S(O) ₂ -	-CH(CH ₃)-	-N(R ⁹)-	-S(O) ₂ -
13	-S(O) ₂ -	-C(CH ₃) ₂ -	-N(R ⁹)-	-C(O)-
14	-S(O) ₂ -	-C(CH ₃) ₂ -	-N(R ⁹)-	-S(O) ₂ -
15	-S(O) ₂ -	-CH ₂ CH ₂ -	-N(R ⁹)-	-C(O)-
16	-S(O) ₂ -	-CH ₂ CH ₂ -	-N(R ⁹)-	-S(O) ₂ -

A further example of the linker -W-X¹-X²-Y- is described in the table below:

No.	W	X ¹	X ²	Y
1	-C(O)-	-CH(CH ₃)-	-NH-	-C(O)-

In each of the various tables above, R⁹ is usually hydrogen or selected from C₁₋₆ alkyl (e.g. C₁, C₂, C₃ or C₄ alkyl), -(CH₂)_k-carbocyclyl and -(CH₂)_k-heterocyclyl, any of which is optionally substituted with 1, 2, 3, 4 or 5 R¹¹. In particular, -N(R⁹)- may be -NH- or -N(CH₃)₂-.

Where C₁₋₆ alkylene (e.g. -CH₂-, -CH(CH₃)-, -C(CH₃)₂- or -CH₂CH₂-), cycloalkylene (e.g. cyclopropylene) or arylene (e.g. phenylene) are mentioned in the various tables, they may be

- 23 -

substituted with 1, 2, 3, 4 or 5 R^{11} , more usually being unsubstituted or substituted by 1 or 2 substituents selected from hydroxy, amino, halogen (e.g. fluorine or chlorine), C_{1-6} alkyl (e.g. C_1 , C_2 , C_3 or C_4 alkyl) optionally substituted with 1, 2, 3, 4 or 5 R^{12} , and arylene (e.g. phenylene) optionally substituted with 1, 2, 3, 4 or 5 R^{12} , where R^{12} is, for example, $-C(O)NH_2$. Also, where C_{1-6} alkylene is mentioned, it may be exchanged for C_{3-6} carbocyclylene (e.g. cyclopropylene). Where cyclopropylene is mentioned, it may be regarded as having 1 or 2 in-chain atoms, typically 1 in-chain atom.

R^1

R^1 is selected from hydrogen; $-N(R^9)(R^{10})$; hydrocarbyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} ; hydrocarboxy optionally substituted with 1, 2, 3, 4 or 5 R^{11} ; and $-(CH_2)_k$ -heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Alternatively, when Y is $-N(R^9)-$, R^1 and R^9 taken together with the nitrogen atom to which they are attached form a heterocycle, wherein said heterocycle is bound to X via said nitrogen atom and is optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

In one embodiment of the invention, R^1 is $-N(R^9)(R^{10})$. In this case, R^9 and R^{10} are usually each independently hydrogen or selected from C_{1-6} alkyl (e.g. C_1 , C_2 , C_3 or C_4 alkyl), $-(CH_2)_k$ -carbocyclyl (e.g. $-(CH_2)_k$ -cycloalkyl or $-(CH_2)_k$ -aryl) and $(CH_2)_k$ -heterocyclyl, any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . For example, R^9 and R^{10} may be each independently hydrogen or C_{1-6} alkyl (e.g. methyl or ethyl) optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Of mention are compounds in which at least one of R^9 and R^{10} is C_{1-6} alkyl group substituted with C_{1-6} alkoxy. In particular, $-N(R^9)(R^{10})$ may be amino, methylamino, dimethylamino or (methoxymethyl)methylamino.

In another embodiment of the invention, R^1 is hydrocarbyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} . In this case, R^1 is often selected from C_{1-6} alkyl (e.g. C_1 , C_2 , C_3 or C_4 alkyl) or $-(CH_2)_k$ -carbocyclyl (e.g. $-(CH_2)_k$ -cycloalkyl or $-(CH_2)_k$ -aryl), either of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . In particular, R^1 may be C_{1-6} alkyl (e.g. C_1 , C_2 , C_3 or C_4 alkyl), $-(CH_2)_k$ -cycloalkyl (e.g. cyclopropyl, cyclobutyl or cyclopropylmethyl) or $-(CH_2)_k$ -aryl (e.g. phenyl or benzyl), any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Of particular mention are methyl; methoxymethyl; cyclopropyl optionally substituted with 1 or 2 R^{11} ; and phenyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

- 24 -

In another embodiment of the invention, R^1 is hydrocarbyloxy optionally substituted with 1, 2, 3, 4 or 5 R^{11} . In this case, R^1 is often selected from C_{1-6} alkyl (e.g. C_1 , C_2 , C_3 or C_4 alkyl) or $-O-(CH_2)_k$ -carbocyclyl (e.g. $-O-(CH_2)_k$ -cycloalkyl or $-O-(CH_2)_k$ -aryl), either of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . In particular, R^1 may be C_{1-6} alkyl (e.g. C_1 , C_2 , C_3 or C_4 alkyloxy), $-O-(CH_2)_k$ -cycloalkyl (e.g. cyclopropyloxy, cyclobutyloxy or cyclopropylmethyloxy) or $-O-(CH_2)_k$ -aryl (e.g. $-O$ -phenyl or $-O$ -benzyl), any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Of particular mention are methyl; methoxymethyl; cyclopropyl optionally substituted with 1 or 2 R^{11} ; and phenyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

In a further embodiment, R^1 is $-(CH_2)_k$ -heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Typically, k is 0, 1 or 2, more usually 0. The heterocyclyl group may be heterocycloalkyl or heteroaryl, either of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Exemplary heterocyclyl groups include oxiranyl, aziriny, 1,2-oxathiolanyl, imidazolyl, thienyl, furyl, tetrahydrofuryl, pyranly, imidazolyl, thiopyranly, thianthrenyl, isobenzofuranyl, benzofuranyl, chromenyl, 2*H*-pyrrolyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolyl, imidazolidinyl, benzimidazolyl, pyrazolyl, pyrazinyl, pyrazolidinyl, thiazolyl, isothiazolyl, dithiazolyl, oxazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pirdinyl, piperidyl, piperazinyl, pyridazinyl, morpholinyl, thiomorpholinyl, especially thiomorpholino, indolizinyl, isoindolyl, 3*H*-indolyl, indolyl, benzimidazolyl, cumaryl, indazolyl, triazolyl, tetrazolyl, purinyl, 4*H*-quinolizinyl, isoquinolyl, quinolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, 3,4-dihydro-2*H*-isoquinolin-1-one, 3,4-dihydro-2*H*-isoquinolinyl, tetrahydropyranly, 1,3-dihydro-isoindolyl, 1,3-Dioxo-1,3-dihydro-isoindolyl, 3,4-dihydro-2*H*-isoquinolinyl, 3,4-dihydro-2*H*-isoquinolinyl-1-one, decahydroquinolyl, octahydroisoquinolyl, benzofuranyl, dibenzofuranyl, benzothiophenyl, dibenzothiophenyl, phthalazinyl, naphthyridinyl, quinoxalyl, quinazolinyl, quinazolinyl, cinnolinyl, pteridinyl, carbazolyl, β -carbolinyl, phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, furazanyl, phenazinyl, phenothiazinyl, phenoxazinyl, chromenyl, isochromanyl and chromanyl, any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Of particular mention are imidazolyl, oxazolyl, morpholinyl, 1,4-benzodioxanyl, pyrimidyl, and pyrazinyl, any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Also of mention are 1,3-dioxo-isoindolyl, 2-oxo-pyrrolidinyl and 2,4-dioxo-thiazolidin-3-yl, any of which is optionally substituted with 1, 2 or 3 R^{11} .

In compounds in which Y is $-C(O)-$, R^1 is often C_{1-6} alkyl (e.g. C_1 , C_2 , C_3 or C_4 alkyl),

- 25 -

cycloalkyl (e.g. cyclopropyl), aryl (e.g. phenyl) or heterocyclyl (e.g. imidazolyl, oxazolyl, morpholinyl, 1,4-benzodioxanyl or pyrazinyl), any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

In compounds in which Y is $-S(O)_2-$ or $-S(O)N(R^9)-$, R^1 is often hydrogen, C_{1-6} alkyl, cycloalkyl (e.g. cyclopropyl or cyclopropylmethyl), aryl (e.g. phenyl) or heterocyclyl (e.g. imidazolyl, oxazolyl, morpholinyl, 1,4-benzodioxanyl or pyrazinyl), any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Alternatively, R^1 may often be $-N(R^9)(R^{10})$, e.g. amino, C_{1-6} alkylamino or di(C_{1-6} alkyl)amino.

Of particular mention are compounds in which R^1 is morpholin-4-yl or cyclopropyl, either of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

In another embodiment, Y is $-N(R^9)-$, and $-N(R^9)R^1$ taken together form a nitrogen-containing heterocycle optionally substituted with 1, 2, 3, 4 or 5 R^{11} . The heterocyclyl group may be heterocycloalkyl or heteroaryl, either of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Exemplary nitrogen-containing heterocyclyl groups include aziriny, imidazolyl, pyranly, 2*H*-pyrrolyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolidinyl, benzimidazolyl, pyrazolyl, pyrazinyl, pyrazolidinyl, thiazolyl, isothiazolyl, dithiazolyl, oxazolyl, isoxazolyl, pyridyl, pyrimidinyl, piperidyl, piperazinyl, pyridazinyl, morpholinyl, thiomorpholinyl, piridinyl, indolizinyl, isoindolyl, 3*H*-indolyl, indolyl, indazolyl, triazolyl, tetrazolyl, purinyl, 4*H*-quinolizinyl, isoquinolyl, quinolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, decahydroquinolyl, octahydroisoquinolyl, phthalazinyl, naphthyridinyl, quinoxalyl, quinazolinyl, quinazolinyl, cinnolinyl, pteridinyl, carbazolyl, β -carbolinyl, phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, furazanyl, phenazinyl, phenothiazinyl, phenoxazinyl, 2,3-Dihydro-benzo[1,4]dioxine, Dihydro-benzodioxine, tetrahydropyranyl, tetrahydrofuranyl, 1,3-Dioxo-1,3-dihydro-isoindolyl any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Of mention are indolyl, isoindolyl, pyrrolidinyl and thiazolidin-3-yl, any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Of particular mention are 1,3-dioxo-isoindolyl, 2-oxo-pyrrolidinyl and 2,4-dioxo-thiazolidin-3-yl, any of which is optionally substituted with 1, 2 or 3 R^{11} .

Where R^1 is substituted with R^{11} , the or each R^{11} is often independently selected from halogen (e.g. fluorine, chlorine or bromine), hydroxy, cyano, amino, $-C(O)OH$, C_{1-6} alkyl, C_{1-6} alkoxy (e.g. C_1 , C_2 , C_3 or C_4 alkoxy), $-C(O)-C_{1-6}$ alkyl, $-C(O)O-C_{1-6}$ alkyl, $-S(O)_r-C_{1-6}$ alkyl, $-NH(C_{1-6}$ alkyl), and $-N(C_{1-6}$ alkyl)₂, wherein any C_{1-6} alkyl group present is optionally

- 26 -

substituted with 1, 2, 3, 4 or 5 substituents independently selected from halogen, cyano, amino, hydroxy and C₁₋₆ alkoxy.

Where R¹ is substituted with R¹¹, the or each R¹¹ is often independently selected from halogen (e.g. fluorine, chlorine or bromine), hydroxy, cyano, amino, oxo, -C(O)OH, -C(O)-NH₂, -C(O)-NH-C₁₋₆ alkyl, C₁₋₆ alkyl, C₁₋₆ alkoxy (e.g. C₁, C₂, C₃ or C₄ alkoxy), -O-aryl (e.g. -O-phenyl), -(CH₂)_k-phenyl, -(CH₂)_k-heterocycyl, -(CH₂)_k-cycloalkyl, -C(O)-heterocycyl, -C(O)-C₁₋₆ alkyl, -C(O)O-C₁₋₆ alkyl, -S(O)_r-C₁₋₆ alkyl, -NH(C₁₋₆ alkyl), -NH-C(O)-(C₁₋₆ alkyl), 6- or 5-membered cycloaryl, a spiro group (e.g. via a cyclopropyl) and -N(C₁₋₆ alkyl)₂, wherein any C₁₋₆ alkyl group present is optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from halogen, cyano, amino, hydroxy and C₁₋₆ alkoxy.

R² & R³

R² and R³ are each independently selected from R⁸, -OR⁸, -C(O)R⁸, -C(O)OR⁸ and -S(O)_rR⁹.

In one embodiment of the invention, R² and R³ are each independently hydrogen; hydroxy; or selected from C₁₋₆ alkyl, C₁₋₆ alkoxy, -(CH₂)_k-cycloalkyl, -(CH₂)_k-heterocycloalkyl, -(CH₂)_k-aryl and -(CH₂)_k-heteroaryl, any of which is optionally substituted with 1, 2, 3, 4, 5 or 6 R¹¹.

In another embodiment, R² is hydrogen; and R³ is hydrogen, hydroxy, C₁₋₆ alkyl, C₁₋₆ alkoxy, -(CH₂)_k-cycloalkyl or -(CH₂)_k-aryl, any of which alkyl, alkoxy, cycloalkyl and aryl groups is optionally substituted with 1, 2, 3, 4 or 5 R¹¹, wherein the or each R¹¹ is, for example, hydroxy, halogen (for example, chlorine or fluorine); C₁, C₂, C₃ or C₄ alkyl, for example methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl or tert-butyl, any of which is optionally substituted with 1, 2, 3 or 4 halogen (e.g. fluorine or chlorine) atoms, an example being trifluoromethyl; C₁, C₂, C₃ or C₄ alkoxy, for example methoxy, ethoxy, propoxy, isopropoxy, butoxy or tert-butoxy, any of which is optionally substituted with 1, 2, 3 or 4 halogen (e.g. fluorine or chlorine) atoms, and in particular R³ is hydrogen, methyl, cyclopropylmethyl or benzyl.

In a further embodiment, R² and R³ are each hydrogen.

R⁴ & R⁵

- 27 -

R^4 and R^5 are each independently selected from hydrogen, hydroxy, halogen and C_{1-6} alkyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

In one embodiment, R^4 and R^5 are each independently hydrogen, hydroxy, halogen (for example, chlorine or fluorine); or C_1 , C_2 , C_3 or C_4 alkyl, for example methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl or tert-butyl, any of which is optionally substituted with 1, 2, 3 or 4 halogen (e.g. fluorine or chlorine) atoms, an example being trifluoromethyl.

In another embodiment, R^4 is hydrogen, hydroxy, halogen (for example, chlorine or fluorine); or C_1 , C_2 , C_3 or C_4 alkyl, for example methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl or tert-butyl, any of which is optionally substituted with 1, 2, 3 or 4 halogen (e.g. fluorine or chlorine) atoms, an example being trifluoromethyl; and R^5 is typically hydrogen.

In a further embodiment, R^4 is hydrogen, hydroxy, fluorine, chlorine or C_1 , C_2 , C_3 or C_4 alkyl; and R^5 is hydrogen.

In a further embodiment, R^4 is hydrogen, hydroxy, fluorine, chlorine or methyl; and R^5 is hydrogen.

In a further embodiment, R^4 and R^5 are each hydrogen.


R^6

R^6 is aryl or heteroaryl, either of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

In one embodiment of the invention, R^6 is aryl, in particular phenyl or naphthyl, either of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} . In embodiments, R^6 is phenyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} , wherein the or each R^{11} is, for example, hydroxy, halogen (for example, chlorine or fluorine); C_1 , C_2 , C_3 or C_4 alkyl, for example methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl or tert-butyl, any of which is optionally substituted with 1, 2, 3 or 4 halogen (e.g. fluorine or chlorine) atoms, an example being trifluoromethyl; or C_1 , C_2 , C_3 or C_4 alkoxy, for example methoxy, ethoxy, propoxy, isopropoxy, butoxy, tert-butoxy, any of which is optionally substituted with 1, 2, 3 or 4 halogen (e.g. fluorine or chlorine) atoms. For example, R^6 may be phenyl optionally substituted with 1, 2, 3, 4 or 5 halogen (e.g. fluorine) atoms.

- 28 -

In a further embodiment, R^6 is a group selected from:



In a further embodiment, R^6 is 2,4,5-trifluorophenyl.

In another embodiment, R^6 is heteroaryl (often monocyclic), for example, thienyl or benzothiophenyl, and is optionally substituted with 1, 2, 3, 4 or 5 R^{11} , wherein the or each R^{11} is, for example, hydroxy, halogen (for example, chlorine or fluorine); C_1 , C_2 , C_3 or C_4 alkyl, for example methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl or tert-butyl, any of which is optionally substituted with 1, 2, 3 or 4 halogen (e.g. fluorine or chlorine) atoms, an example being trifluoromethyl; or C_1 , C_2 , C_3 or C_4 alkoxy, for example methoxy, ethoxy, propoxy, isopropoxy, butoxy, tert-butoxy, any of which is optionally substituted with 1, 2, 3 or 4 halogen (e.g. fluorine or chlorine) atoms.

R^{11}

Each R^{11} is independently selected from R^{12} ; hydrocarbyl (e.g. C_{1-6} alkyl or $-(CH_2)_k$ -aryl) optionally substituted with 1, 2, 3, 4 or 5 R^{12} ; and $-(CH_2)_k$ -heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{12} ; wherein R^{12} is independently selected from halogen, trifluoromethyl, cyano, nitro, oxo, $=NR^{13}$, $-OR^{13}$, $-C(O)R^{13}$, $-C(O)N(R^{13})R^{14}$, $-C(O)OR^{13}$, $-OC(O)R^{13}$, $-S(O)_2R^{13}$, $-S(O)_2N(R^{13})R^{14}$, $-N(R^{13})R^{14}$, $-N(R^{13})N(R^{13})R^{14}$, $-N(R^{13})C(O)R^{14}$ and $-N(R^{13})S(O)_2R^{13}$; and R^{13} and R^{14} are each independently hydrogen or selected from hydrocarbyl (e.g. C_{1-6} alkyl or $-(CH_2)_k$ -aryl, or $-(CH_2)_k$ -cycloalkyl) and $-(CH_2)_k$ -heterocyclyl, either of which is optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from halogen, cyano, amino, hydroxy, C_{1-6} alkyl and C_{1-6} alkoxy, or selected from oxo, halogen, cyano, amino, hydroxy, C_{1-6} alkyl and C_{1-6} alkoxy.

- 29 -


Typically, each R^{11} is independently selected from halogen (e.g. fluorine, chlorine or bromine), hydroxy, cyano, amino, $-C(O)OH$, C_{1-6} alkyl, C_{1-6} alkoxy (e.g. C_1 , C_2 , C_3 or C_4 alkoxy), $-C(O)-C_{1-6}$ alkyl, $-C(O)O-C_{1-6}$ alkyl, $-S(O)_r-C_{1-6}$ alkyl- $NH(C_{1-6}$ alkyl) and $-N(C_{1-6}$ alkyl) $_2$, wherein any C_{1-6} alkyl group present is optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from halogen, cyano, amino, hydroxy and C_{1-6} alkoxy.

Typically, each R^{11} is independently selected from halogen (e.g. fluorine, chlorine or bromine), hydroxy, cyano, amino, $-C(O)OH$, C_{1-6} alkyl, C_{1-6} alkoxy (e.g. C_1 , C_2 , C_3 or C_4 alkoxy), $-C(O)-C_{1-6}$ alkyl, $-C(O)O-C_{1-6}$ alkyl, $-S(O)_r-C_{1-6}$ alkyl, $-(CH_2)_k$ -cycloalkyl, $-(CH_2)_k$ -aryl, $-(CH_2)_k$ -heterocyclyl, $-NH-(CH_2)_k$ -aryl, $-NH-(CH_2)_k$ -cycloalkyl, $-NH-C(O)-(CH_2)_k$ -aryl, $-NH-C(O)-(CH_2)_k$ -cycloalkyl, $-N(C_{1-6}$ alkyl)- $(CH_2)_k$ -aryl, $-N(C_{1-6}$ alkyl)- $(CH_2)_k$ -cycloalkyl, $-NH(C_{1-6}$ alkyl) and $-N(C_{1-6}$ alkyl) $_2$, wherein any C_{1-6} alkyl group or aryl or present is optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from halogen, cyano, amino, hydroxy and C_{1-6} alkoxy.

For the avoidance of doubt, where a group is substituted with more than one R^{11} , each R^{11} is independently selected from the range of substituents specified. The same applies to compounds of the invention comprising more than one R^{11} substituent; each R^{11} is selected independently of any other R^{11} substituent present in the compound. As previously indicated, where R^{11} is halo, particularly fluoro, any number of hydrogens may in principle be replaced.

- 30 -


A particular embodiment of the invention is a compound of the following Formula:




(VI)

or a pharmaceutically acceptable salt or prodrug thereof.

The invention therefore includes compounds of the following Formulae:




(VII)



(VIII)

or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

Another embodiment of the invention is a compound of the following Formula:




(IX)

- 31 -


wherein X^1 is selected from $-O-$, $-C(O)-$, $-S(O)_2-$, $-N(R^9)-$ and hydrocarbylene (e.g. C_{1-6} alkylene) optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

or a pharmaceutically acceptable salt or prodrug thereof.

The invention therefore includes compounds of the following Formulae:




(X)



(XI)

or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

A further embodiment of the invention is a compound of the following Formula:




(XII)


wherein p is 0, 1, 2, 3, 4 or 5 or p is 0, 1, 2 or 3;

or a pharmaceutically acceptable salt or prodrug thereof.


The invention therefore includes compounds of the following Formulae:



(XIII)



(XIV)



(XIVb)

or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

With regard to Formulae (IX) to (XIV) and (XIVb), X^1 may be defined as in Formula (IV), i.e. X^1 is $-N(R^9)-$ or hydrocarbylene (e.g. C_{1-6} alkylene) optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Y is often $-C(O)-$, $-C(O)O-$, $-C(O)N(R^9)-$, $-S(O)-$, $-S(O)_2-$ or $-S(O)N(R^9)-$. In particular, Y is often $-C(O)-$, $-C(O)-$, $-S(O)-$ or $-S(O)_2-$.

Of particular mention are compounds of any of Formulae (IX) to (XIV) and (XIVb) in which $-W-X^1-Y-$ is one of the following linkers:

No.	W	X^1	Y
1	$-C(O)-$	$-N(R^9)-$	$-C(O)-$
2	$-C(O)-$	C_{1-6} alkylene	$-C(O)-$
3	$-C(O)-$	$-N(R^9)-$	$-S(O)_2-$





- 33 -

4	-C(O)-	C ₁₋₆ alkylene	-S(O) ₂ -
5	-S(O) _r -	-N(R ⁹)-	-C(O)-
6	-S(O) _r -	C ₁₋₆ alkylene	-C(O)-
7	-S(O) _r -	C ₁₋₆ alkylene	-S(O) ₂ -

Also of mention are compounds of any of said Formulae in which -W-X¹-Y- is one of the following linkers:

No.	W	X ¹	Y
1	-C(O)-	C ₁₋₆ alkylene	-S(O) ₂ -
2	-S(O) _r -	C ₁₋₆ alkylene	-C(O)-

Also of mention are compounds of any of said Formulae in which -W-X¹-Y- is one of the following linkers:

No.	W	X ¹	Y
1	-C(O)-	C ₁₋₆ alkylene	-C(O)-N(R ⁹)-
2	-C(O)-	-CH ₂ -	-C(O)-NH-
3	-C(O)-		-C(O)-
4	-C(O)-		-C(O)-
5	-C(O)-		-N(R ⁹)-
6	-C(O)-		-N(R ⁹)-
7	-C(O)-	arylene	-C(O)-

- 34 -

8	-C(O)-	C ₁₋₆ alkylene	-C(O)-
9	-C(O)-	arylene	-N(R ⁹)-
10	-C(O)-	-CH ₂ -	-C(O)-
11	-C(O)-	-CH(CH ₃)-	-C(O)-
12	-C(O)-	-C(CH ₃) ₂ -	-C(O)-
13	-C(O)-	-CH ₂ CH ₂ -	-C(O)-

In one embodiment in the above compounds wherein Y is -N(R⁹)-, R¹ and R⁹ taken together form a 5- or 6-membered heterocyclyl optionally substituted by 1, 2, 3, 4 or 5 R¹¹. The formed 5- or 6-membered heterocyclyl is for instance 1-methyl-imidazolidine-2,4-dione, imidazolidine-2,4-dione, thiazolidine-2,4-dione, pyrrolidine-2,5-dione, isoindole-1,3-dione, or pyrrolidiny-2-oxo.

Also of mention are compounds of any of said Formulae in which -W-X¹-Y- is one of the following linkers:

No.	W	X ¹	Y
1	-C(O)-	-CH ₂ -	-S(O) ₂ -
2	-C(O)-	-CH(CH ₃)-	-S(O) ₂ -
3	-C(O)-	-C(CH ₃) ₂ -	-S(O) ₂ -
4	-C(O)-	-CH ₂ CH ₂ -	-S(O) ₂ -
5	-S(O) ₂ -	-CH ₂ -	-C(O)-
6	-S(O) ₂ -	-CH(CH ₃)-	-C(O)-
7	-S(O) ₂ -	-C(CH ₃) ₂ -	-C(O)-
8	-S(O) ₂ -	-CH ₂ CH ₂ -	-C(O)-

In the case of the table directly above, R¹ is often a 6-membered heterocyclyl which may be unsubstituted or substituted with 1, 2, 3, 4 or 5 R¹¹. In the case of the table directly above, in particular the first row thereof, R¹ is often morpholiny (e.g. morpholin-4-yl) which may be unsubstituted or substituted with 1, 2, 3, 4 or 5 R¹¹.

Also of mention are compounds of any of said Formulae in which -W-X¹-Y- is one of the following linkers:

- 35 -

No.	W	X ¹	Y
1	-C(O)-	C ₁₋₆ alkylene	-C(O)-
2	-S(O) _r -	C ₁₋₆ alkylene	-S(O) ₂ -

Also of mention are compounds of any of said Formulae in which -W-X¹-Y- is one of the following linkers:

No.	W	X ¹	Y
1	-C(O)-	-CH ₂ -	-C(O)-
2	-C(O)-	-CH(CH ₃)-	-C(O)-
3	-C(O)-	-C(CH ₃) ₂ -	-C(O)-
4	-C(O)-	-CH ₂ CH ₂ -	-C(O)-

Also of particular mention are compounds of the above Formula wherein X¹ is carbocyclylene (e.g. cycloalkylene or arylene) optionally substituted with 1, 2, 3, 4 or 5 R¹¹; or a pharmaceutically acceptable salt or prodrug thereof.

Also of mention are compounds of any of said Formulae in which -W-X¹-Y- is one of the following linkers:

No.	W	X ¹	Y
1	-C(O)-	cycloalkylene	-S(O) ₂ -
2	-S(O) _r -	cycloalkylene	-C(O)-
3	-C(O)-	cycloalkylene	-C(O)-
4	-S(O) _r -	cycloalkylene	-S(O) ₂ -

Also of mention are compounds of any of said Formulae in which -W-X¹-Y- is one of the following linkers:

No.	W	X ¹	Y
1	-C(O)-	cyclopropylene	-S(O) ₂ -
2	-S(O) _r -	cyclopropylene	-C(O)-
3	-C(O)-	cyclopropylene	-C(O)-
4	-S(O) _r -	cyclopropylene	-S(O) ₂ -

- 36 -


Also of mention are compounds of any of said Formulae in which $-W-X^1-Y-$ is one of the following linkers:

No.	W	X ¹	Y
1	-C(O)-	arylene	-S(O) ₂ -
2	-S(O) _r -	arylene	-C(O)-
3	-C(O)-	arylene	-C(O)-
4	-S(O) _r -	arylene	-S(O) ₂ -

Also of mention are compounds of any of said Formulae in which $-W-X^1-Y-$ is one of the following linkers:

No.	W	X ¹	Y
1	-C(O)-	phenylene	-S(O) ₂ -
2	-S(O) _r -	phenylene	-C(O)-
3	-C(O)-	phenylene	-C(O)-
4	-S(O) _r -	phenylene	-S(O) ₂ -

A further embodiment of the invention is a compound of the following Formula:




(XV)

wherein X^1 and X^2 are each independently selected from $-O-$, $-C(O)-$, $-S(O)_r-$, $-N(R^9)-$ and C_{1-6} alkylene optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;


or a pharmaceutically acceptable salt or prodrug thereof.

The invention therefore includes compounds of the following Formulae:

- 37 -




(XVI)



(XVII)

or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

A further embodiment of the invention is a compound of the following Formula:




(XVIII)

wherein p is 0, 1, 2, 3, 4 or 5;


or a pharmaceutically acceptable salt or prodrug thereof.

The invention therefore includes compounds of the following Formulae:



(XIX)

- 38 -



or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

With regard to Formulae (XV) to (XX) or (XXb), X^1 and X^2 may be defined as in Formula (V), i.e. one of X^1 and X^2 is $-N(R^9)-$; and the other is C_{1-6} alkylene optionally substituted with 1, 2, 3, 4 or 5 R^{11} . Y is often $-C(O)-$, $-C(O)O-$, $-C(O)N(R^9)-$, $-S(O)-$, $-S(O)_2-$ or $-S(O)N(R^9)-$. In particular, Y is often $-C(O)-$, $-C(O)-$, $-S(O)-$ or $-S(O)_2-$.





Of particular mention are compounds of any of Formulae (XV) to (XX) in which $-W-X^1-X^2-Y-$ is one of the following linkers:

No.	W	X^1	X^2	Y
1	$-C(O)-$	C_{1-6} alkylene	$-O-$	$-C(O)-$
2	$-C(O)-$	$-O-$	$-N(R^9)-$	$-C(O)-$
3	$-C(O)-$	C_{1-6} alkylene	$-N(R^9)-$	$-C(O)-$
4	$-C(O)-$	$-O-$	C_{1-6} alkylene	$-C(O)-$
5	$-C(O)-$	$-N(R^9)-$	C_{1-6} alkylene	$-C(O)-$
6	$-C(O)-$	C_{1-6} alkylene	$-N(R^9)-$	$-S(O)_2-$
7	$-C(O)-$	$-O-$	C_{1-6} alkylene	$-S(O)_2-$
8	$-C(O)-$	$-N(R^9)-$	C_{1-6} alkylene	$-S(O)_2-$
9	$-S(O)-$	C_{1-6} alkylene	$-O-$	$-C(O)-$

- 39 -

No.	W	X ¹	X ²	Y
10	-S(O) ₁ -	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-
11	-S(O) ₁ -	-N(R ⁹)-	C ₁₋₆ alkylene	-C(O)-
12	-S(O) ₁ -	C ₁₋₆ alkylene	-N(R ⁹)-	-S(O) ₂ -
13	-S(O) ₁ -	-N(R ⁹)-	C ₁₋₆ alkylene	-S(O) ₂ -

Also of mention are compounds of any of said Formulae in which -W-X¹-X²-Y- is one of the following linkers:

1	-S(O) ₁ -	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-O-
2	-S(O) ₁ -	-CH ₂ CH ₂ -	-N(R ⁹)-	-C(O)-O-
3	-C(O)-	arylene	-N(R ⁹)-	-C(O)-
4	-C(O)-		-N(R ⁹)-	-C(O)-
5	-C(O)-		-N(R ⁹)-	-C(O)-
6	-C(O)-		C ₁₋₆ alkylene	-N(R ⁹)-
7	-C(O)-		-CH ₂ CH ₂ -	-N(R ⁹)-

Also of mention are compounds of any of said Formulae in which -W-X¹-X²-Y- is one of the following linkers:

No.	W	X ¹	X ²	Y
1	-C(O)-	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-
2	-C(O)-	C ₁₋₆ alkylene	-N(R ⁹)-	-S(O) ₂ -
3	-S(O) ₂ -	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-

- 40 -

4	-S(O) ₂ -	C ₁₋₆ alkylene	-N(R ⁹)-	-S(O) ₂ -
---	----------------------	---------------------------	----------------------	----------------------

Also of mention are compounds of any of said Formulae in which -W-X¹-X²-Y- is one of the following linkers:

No.	W	X ¹	X ²	Y
1	-C(O)-	-CH ₂ -	-N(R ⁹)-	-C(O)-
2	-C(O)-	-CH ₂ -	-N(R ⁹)-	-S(O) ₂ -
3	-C(O)-	-CH(CH ₃)-	-N(R ⁹)-	-C(O)-
4	-C(O)-	-CH(CH ₃)-	-N(R ⁹)-	-S(O) ₂ -
5	-C(O)-	-C(CH ₃) ₂ -	-N(R ⁹)-	-C(O)-
6	-C(O)-	-C(CH ₃) ₂ -	-N(R ⁹)-	-S(O) ₂ -
7	-C(O)-	-CH ₂ CH ₂ -	-N(R ⁹)-	-C(O)-
8	-C(O)-	-CH ₂ CH ₂ -	-N(R ⁹)-	-S(O) ₂ -
9	-S(O) ₂ -	-CH ₂ -	-N(R ⁹)-	-C(O)-
10	-S(O) ₂ -	-CH ₂ -	-N(R ⁹)-	-S(O) ₂ -
11	-S(O) ₂ -	-CH(CH ₃)-	-N(R ⁹)-	-C(O)-
12	-S(O) ₂ -	-CH(CH ₃)-	-N(R ⁹)-	-S(O) ₂ -
13	-S(O) ₂ -	-C(CH ₃) ₂ -	-N(R ⁹)-	-C(O)-
14	-S(O) ₂ -	-C(CH ₃) ₂ -	-N(R ⁹)-	-S(O) ₂ -
15	-S(O) ₂ -	-CH ₂ CH ₂ -	-N(R ⁹)-	-C(O)-
16	-S(O) ₂ -	-CH ₂ CH ₂ -	-N(R ⁹)-	-S(O) ₂ -

A further example of the linker -W-X¹-X²-Y- is described in the table below:

No.	W	X ¹	X ²	Y
1	-C(O)-	-CH(CH ₃)-	-NH-	-C(O)-

In the case of the linker described in the table directly above, R¹ is especially cycloalkyl (e.g. cyclopropyl) and may be unsubstituted or substituted with 1, 2, 3, 4 or 5 R¹¹.

In each of the various tables above, R⁹ is usually hydrogen or selected from C₁₋₆ alkyl (e.g. C₁, C₂, C₃ or C₄ alkyl), -(CH₂)_k-carbocyclyl and -(CH₂)_k-heterocyclyl, any of which is optionally substituted with 1, 2, 3, 4 or 5 R¹¹. In particular, -N(R⁹)- may be -NH- or -N(CH₃)-.

- 41 -


Where C₁₋₆ alkylene (e.g. -CH₂-, -CH(CH₃)-, -C(CH₃)₂- or -CH₂CH₂-), cycloalkylene (e.g. cyclopropylene) or arylene (e.g. phenylene) are mentioned in the various tables, they may be substituted with 1, 2, 3, 4 or 5 R¹¹, more usually being unsubstituted or substituted by 1 or 2 substituents selected from hydroxy, amino, halogen (e.g. fluorine or chlorine), C₁₋₆ alkyl (e.g. C₁, C₂, C₃ or C₄ alkyl) optionally substituted with 1, 2, 3, 4 or 5 R¹², and arylene (e.g. phenylene) optionally substituted with 1, 2, 3, 4 or 5 R¹², where R¹² is, for example, -C(O)NH₂. Also, where C₁₋₆ alkylene is mentioned, it may be exchanged for C₃₋₆ carbocyclylene (e.g. cyclopropylene). Where cyclopropylene is mentioned, it may be regarded as having 1 or 2 in-chain atoms, typically 1 in-chain atom.

In one embodiment are the hereinabove described compounds wherein X¹ or X² is an alkylene or arylene and wherein X¹ or X² is substituted by 1, 2, 3, 4 or 5 R¹¹, preferably 1 or 2 R¹¹. Preferably the R¹¹ substituents are independently selected from a spiro group, C₁₋₆ alkyl, -O-C₁₋₆ alkyl, -NH-C(O)-(C₁₋₆ alkyl), phenyl, benzyl, hydroxy, halogen, amino, wherein the alkyl group is optionally substituted by amino, hydroxy, -C(O)-N(C₁₋₆ alkyl)(C₁₋₆ alkyl), -C(O)-NH(C₁₋₆ alkyl), -C(O)-NH₂, (C₁₋₆ alkyl), or halogen.


In one embodiment in the above compounds wherein Y is -N(R⁹)-, R¹ and R⁹ taken together form a 5- or 6-membered heterocyclyl optionally substituted by 1, 2, 3, 4 or 5 R¹¹. The formed 5- or 6-membered heterocyclyl is for instance 1-methyl-imidazolidine-2,4-dione, imidazolidine-2,4-dione, thiazolidine-2,4-dione, pyrrolidine-2,5-dione or pyrrolidinyl-2-oxo.

Examples of compounds of the invention include those shown below. It will of course be appreciated that, where appropriate, each compound may be in the form of the free compound, an acid or base addition salt, or a prodrug. Where a nitrogen atom forming only two bonds is shown, this generally represents NH.


- 42 -




B1




C1




C2




D1




D2




D3



D4




D5




D6


- 43 -




D7




D8




D9




D10




D11




D12



D13




D14




D15


- 44 -




D16




D17




D18




D19



E1




E2




E3


- 45 -




E4




E5




E6




E7




E8




E9



E10




E11




E12


- 46 -




E13




E14




E15




E16




E17




E18



E19





E20




E21



- 47 -




E22




E23




F1



F2



F3



F4




F5




F6




- 48 -




F7




F8




F9




G1




G2




G3



G4




G5




G6


- 49 -




G7




G8




G9




G10




G11




G12



G13




G14




G15


- 50 -




G16




G17




G18




G19




G20




G21



G22




G23




G24


- 51 -




G25




G26




G27




G28




G29




G30



G31




G32




G33


- 52 -




G34




H1




H2




H3



H4




H5




H6


- 53 -




H7




H8




H9




H10



H11




H12




H13


- 54 -




11




12




13




14




15




16



17




18




19


- 55 -




I10




K1




K4




K2



K5




K3




K6


- 56 -




K7




K8




K9




K10




K11




K12



K13




K14




K15


- 57 -




K16




K17




K18



K19




L1




L2


Further compounds of the invention include:




X1




X2




X3




X4




X5




X6




X7




X8




X9




X10




X11




X12




X13




X14




X15



X16




X17




X18


- 60 -



X19



X20



X21

Compounds of the invention may be in the form of pharmaceutically acceptable salts. The pharmaceutically acceptable salts of the present disclosure can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, Pa., US, 1985, p. 1418, the disclosure of which is hereby incorporated by reference; see also Stahl et al, Eds, "*Handbook of Pharmaceutical Salts Properties Selection and Use*", Verlag Helvetica Chimica Acta and Wiley-VCH, 2002.

The disclosure thus includes pharmaceutically-acceptable salts of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. for example the conventional non-toxic salts or the quaternary ammonium salts which are formed, e.g. from inorganic or organic acids or bases. Examples of such acid addition salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, and undecanoate. Base salts include ammonium salts, alkali metal salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups may be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl; and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides and others.

The invention includes prodrugs for the active pharmaceutical species of the invention, for example in which one or more functional groups are protected or derivatised but can be converted *in vivo* to the functional group, as in the case of esters of carboxylic acids

convertible *in vivo* to the free acid, or in the case of protected amines, to the free amino group. The term "prodrug," as used herein, represents in particular compounds which are rapidly transformed *in vivo* to the parent compound, for example, by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987; H Bundgaard, ed, Design of Prodrugs, Elsevier, 1985; and Judkins, et al. Synthetic Communications, 26(23), 4351-4367 (1996), each of which is incorporated herein by reference.

Prodrugs therefore include drugs having a functional group which has been transformed into a reversible derivative thereof. Typically, such prodrugs are transformed to the active drug by hydrolysis. As examples may be mentioned the following:

Functional Group	Reversible derivative
<i>Carboxylic acid</i>	<i>Esters, including e.g. acyloxyalkyl esters, amides</i>
<i>Alcohol</i>	<i>Esters, including e.g. sulfates and phosphates as well as carboxylic acid esters</i>
<i>Amine</i>	<i>Amides, carbamates, imines, enamines,</i>
<i>Carbonyl (aldehyde, ketone)</i>	<i>Imines, oximes, acetals/ketals, enol esters, oxazolidines and thiazoxolidines</i>

Prodrugs also include compounds convertible to the active drug by an oxidative or reductive reaction. As examples may be mentioned:

Oxidative activation

- N- and O- dealkylation
- Oxidative deamination
- N-oxidation
- Epoxidation

Reductive activation

- Azo reduction
- Sulfoxide reduction

- 63 -

- Disulfide reduction
- Bioreductive alkylation
- Nitro reduction.

Also to be mentioned as metabolic activations of prodrugs are nucleotide activation, phosphorylation activation and decarboxylation activation. For additional information, see "The Organic Chemistry of Drug Design and Drug Action", R B Silverman (particularly Chapter 8, pages 497 to 546), incorporated herein by reference.

The use of protecting groups is fully described in 'Protective Groups in Organic Chemistry', edited by J W F McOmie, Plenum Press (1973), and 'Protective Groups in Organic Synthesis', 2nd edition, T W Greene & P G M Wutz, Wiley-Interscience (1991).

Thus, it will be appreciated by those skilled in the art that, although protected derivatives of compounds of the disclosure may not possess pharmacological activity as such, they may be administered, for example parenterally or orally, and thereafter metabolised in the body to form compounds of the invention which are pharmacologically active. Such derivatives are therefore examples of "prodrugs". All prodrugs of the described compounds are included within the scope of the disclosure.

Some groups mentioned herein (especially those containing heteroatoms and conjugated bonds) may exist in tautomeric forms and all these tautomers are included in the scope of the disclosure. More generally, many species may exist in equilibrium, as for example in the case of organic acids and their counterpart anions; a reference herein to a species accordingly includes reference to all equilibrium forms thereof.

The compounds of the disclosure may also contain one or more asymmetric carbon atoms and may therefore exhibit optical and/or diastereoisomerism. All diastereoisomers may be separated using conventional techniques, e.g. chromatography or fractional crystallisation. The various stereoisomers may be isolated by separation of a racemic or other mixture of the compounds using conventional, e.g. fractional crystallisation or HPLC, techniques. Alternatively the desired optical isomers may be made by reaction of the appropriate optically active starting materials under conditions which will not cause racemisation or epimerisation, or by derivatisation, for example with a homochiral acid followed by separation of the

diastereomeric derivatives by conventional means (e.g. HPLC, chromatography over silica). All stereoisomers are included within the scope of the disclosure. Where a single enantiomer or diastereomer is disclosed, the disclosure also covers the other enantiomers or diastereomers, and also racemates; in this regard, particular reference is made to the specific compounds listed herein. As mentioned above, the stereochemical configuration of the carbon atom to which the group $-N(R^2)(R^3)$ is attached may be (*R*) or (*S*), especially (*R*).


Geometric isomers may also exist in the compounds of the present disclosure. The present disclosure contemplates the various geometric isomers and mixtures thereof resulting from the arrangement of substituents around a carbon-carbon double bond and designates such isomers as of the *Z* or *E* configuration, wherein the term "*Z*" represents substituents on the same side of the carbon-carbon double bond and the term "*E*" represents substituents on opposite sides of the carbon-carbon double bond.

The disclosure therefore includes all variant forms of the defined compounds, for example any tautomer or any pharmaceutically acceptable salt, ester, acid or other variant of the defined compounds and their tautomers as well as substances which, upon administration, are capable of providing directly or indirectly a compound as defined above or providing a species which is capable of existing in equilibrium with such a compound.

Synthesis


By way of illustration, a compound of the invention may be prepared according to the following Schemes, in which Scheme 1 shows the synthesis of an intermediate in the form of a racemate; Schemes 2 to 4 show synthesis of compounds of the invention in (*S*) form, and Schemes 5 and 6 show synthesis of compounds in (*R*) form. R^x , R^y and R^z may each be any suitable group. For example, R^x and R^y may be methyl and benzyl respectively, while $-N(R^z)_2$ may form a phthalimido group.

- 65 -




Scheme 1

- 66 -




Scheme 2

- 67 -




Scheme 3

- 68 -




Scheme 4

- 69 -



Scheme 5

- 70 -



Scheme 6

- 71 -

It will be understood that the processes detailed above are solely for the purpose of illustrating the invention and should not be construed as limiting. A process utilising similar or analogous reagents and/or conditions known to one skilled in the art may also be used to obtain a compound of the invention.

Any mixtures of final products or intermediates obtained can be separated on the basis of the physico-chemical differences of the constituents, in a known manner, into the pure final products or intermediates, for example by chromatography, distillation, fractional crystallisation, or by the formation of a salt if appropriate or possible under the circumstances.

Administration & Pharmaceutical Formulations

The compounds of the invention will normally be administered orally, intravenously, subcutaneously, buccally, rectally, dermally, nasally, tracheally, bronchially, by any other parenteral route, as an oral or nasal spray or via inhalation. The compounds may be administered in the form of pharmaceutical preparations comprising prodrug or active compound either as a free compound or, for example, a pharmaceutically acceptable non-toxic organic or inorganic acid or base addition salt, in a pharmaceutically acceptable dosage form. Depending upon the disorder and patient to be treated and the route of administration, the compositions may be administered at varying doses.

Typically, therefore, the pharmaceutical compounds of the invention may be administered orally or parenterally ("parenterally" as used herein, refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion) to a host to obtain an protease-inhibitory effect. In the case of larger animals, such as humans, the compounds may be administered alone or as compositions in combination with pharmaceutically acceptable diluents, excipients or carriers.

Actual dosage levels of active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active compound(s) that is effective to achieve the desired therapeutic response for a particular patient, compositions, and mode of administration. The selected dosage level will depend upon the activity of the particular compound, the route of administration, the severity of the condition being treated and the

- 72 -

condition and prior medical history of the patient being treated. However, it is within the skill of the art to start doses of the compound at levels lower than required for to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

In the treatment, prevention, control, amelioration, or reduction of risk of conditions which require inhibition of DPP-IV enzyme activity, an appropriate dosage level will generally be about 0.01 to 500 mg per kg patient body weight per day which can be administered in single or multiple doses. Preferably, the dosage level will be about 0.1 to about 250 mg/kg per day; more preferably about 0.5 to about 100 mg/kg per day. A suitable dosage level may be about 0.01 to 250 mg/kg per day, about 0.05 to 100 mg/kg per day, or about 0.1 to 50 mg/kg per day. Within this range the dosage may be 0.05 to 0.5, 0.5 to 5 or 5 to 50 mg/kg per day. For oral administration, the compositions are preferably provided in the form of tablets containing 1.0 to 1000 milligrams of the active ingredient, particularly 1.0, 5.0, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 100.0, 150.0, 200.0, 250.0, 300.0, 400.0, 500.0, 600.0, 750.0, 800.0, 900.0 and 1000.0 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day. The dosage regimen may be adjusted to provide the optimal therapeutic response.

According to a further aspect of the invention there is thus provided a pharmaceutical composition including a compound of the invention, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier.

Pharmaceutical compositions of this invention for parenteral injection suitably comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol and the like), and suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

These compositions may also contain adjuvants such as preservative, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol or phenol sorbic acid. It may also be desirable to include isotonic agents such as sugars or sodium chloride, for example. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents (for example aluminum monostearate and gelatin) which delay absorption.

In some cases, in order to prolong the effect of the drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are suitably made by forming microencapsule matrices of the drug in biodegradable polymers, for example polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations may also be prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues. The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable media just prior to use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound is typically mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or one or more: a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol and silicic acid; b) binders such as carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose and acacia; c) humectants such as glycerol; d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates and sodium carbonate; e) solution retarding agents such as paraffin; f) absorption accelerators such as quaternary ammonium compounds; g) wetting

agents such as cetyl alcohol and glycerol monostearate; h) absorbents such as kaolin and bentonite clay and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycol, for example.

Suitably, oral formulations contain a dissolution aid. The dissolution aid is not limited as to its identity so long as it is pharmaceutically acceptable. Examples include nonionic surface active agents, such as sucrose fatty acid esters, glycerol fatty acid esters, sorbitan fatty acid esters (e.g. sorbitan trioleate), polyethylene glycol, polyoxyethylene hydrogenated castor oil, polyoxyethylene sorbitan fatty acid esters, polyoxyethylene alkyl ethers, methoxypolyoxyethylene alkyl ethers, polyoxyethylene alkylphenyl ethers, polyethylene glycol fatty acid esters, polyoxyethylene alkylamines, polyoxyethylene alkyl thioethers, polyoxyethylene polyoxypropylene copolymers, polyoxyethylene glycerol fatty acid esters, pentaerythritol fatty acid esters, propylene glycol monofatty acid esters, polyoxyethylene propylene glycol monofatty acid esters, polyoxyethylene sorbitol fatty acid esters, fatty acid alkylolamides, and alkylamine oxides; bile acid and salts thereof (e.g. chenodeoxycholic acid, cholic acid, deoxycholic acid, dehydrocholic acid and salts thereof, and glycine or taurine conjugate thereof); ionic surface active agents, such as sodium laurylsulfate, fatty acid soaps, alkylsulfonates, alkylphosphates, ether phosphates, fatty acid salts of basic amino acids; triethanolamine soap, and alkyl quaternary ammonium salts; and amphoteric surface active agents, such as betaines and aminocarboxylic acid salts.

The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and may also be of a composition such that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, and/or in delayed fashion. Examples of embedding compositions include polymeric substances and waxes.

The active compounds may also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

The active compounds may be in finely divided form, for example it may be micronised.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan and mixtures thereof. Besides inert diluents, the oral compositions may also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring and perfuming agents. Suspensions, in addition to the active compounds, may contain suspending agents such as ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth and mixtures thereof.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

Compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals which are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolisable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilisers, preservatives, excipients and the like. The preferred lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art, for example, Prescott, Ed., *Methods in Cell Biology*, Volume XIV, Academic Press, New York, N.Y. (1976), p 33 et seq.

Dosage forms for topical administration of a compound of this invention include powders, sprays, ointments and inhalants. The active compound is mixed under sterile conditions with

- 76 -

a pharmaceutically acceptable carrier and any needed preservatives, buffers or propellants which may be required. Ophthalmic formulations, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

Advantageously, the compounds of the invention may be orally active, have rapid onset of activity and low toxicity.

The compounds of the invention may have the advantage that they are more efficacious, less toxic, longer acting, have a broader range of activity, more potent, produce fewer side effects, more easily absorbed than, or have other useful pharmacological properties over, compounds known in the prior art.

Combination therapies

Compounds of the invention may be administered in combination with one or more additional therapeutic agents. Accordingly, the invention provides a pharmaceutical composition comprising an additional agent. The invention also provides a product comprising a compound of the invention and an agent; as a combined preparation for simultaneous, separate or sequential use in therapy.

In particular, a composition or product of the invention may further comprise a therapeutic agent selected from anti-diabetic agents, hypolipidemic agents, anti-obesity or appetite-regulating agents, anti-hypertensive agents, HDL-increasing agents, cholesterol absorption modulators, Apo-A1 analogues and mimetics, thrombin inhibitors, aldosterone inhibitors, inhibitors of platelet aggregation, estrogen, testosterone, selective estrogen receptor modulators, selective androgen receptor modulators, chemotherapeutic agents, and 5-HT₃ or 5-HT₄ receptor modulators; or pharmaceutically acceptable salts or prodrugs thereof.

Examples of anti-diabetic agents include insulin, insulin derivatives and mimetics; insulin secretagogues, for example sulfonylureas (e.g. glipizide, glyburide or amaryl); insulinotropic sulfonylurea receptor ligands, for example meglitinides (e.g. nateglinide or repaglinide); insulin sensitisers, for example protein tyrosine phosphatase-1B (PTP-1B) inhibitors (e.g. PTP-112); GSK3 (glycogen synthase kinase-3) inhibitors, for example SB-517955, SB-4195052, SB-216763, NN-57-05441 or NN-57-05445; RXR ligands, for example GW-0791 or AGN-194204; sodium-dependent glucose cotransporter inhibitors, for example T-1095;

- 77 -

glycogen phosphorylase A inhibitors, for example BAY R3401; biguanides, for example metformin; alpha-glucosidase inhibitors, for example acarbose; GLP-1 (glucagon like peptide-1), GLP-1 analogues and mimetics, for example exendin-4; DPPIV (dipeptidyl peptidase IV) inhibitors, for example DPP728, LAF237 (vildagliptin), MK-0431, saxagliptin or GSK23A; AGE breakers; and thiazolidone derivatives, for example glitazone, pioglitazone, rosiglitazone or (R)-1-{4-[5-methyl-2-(4-trifluoromethyl-phenyl)-oxazol-4-ylmethoxy]-benzenesulfonyl}-2,3-dihydro-1H-indole-2-carboxylic acid (compound 4 of Example 19 of WO 03/043985) or a non-glitazone type PPAR- agonist (e.g. GI-262570); or pharmaceutically acceptable salts or prodrugs thereof.

Examples of hypolipidemic agents include 3-hydroxy-3-methyl-glutaryl coenzyme A (HMG-CoA) reductase inhibitors, for example lovastatin, pitavastatin, simvastatin, pravastatin, cerivastatin, mevastatin, velostatin, fluvastatin, dalvastatin, atorvastatin, rosuvastatin or rivastatin; squalene synthase inhibitors; FXR (farnesoid X receptor) ligands; LXR (liver X receptor) ligands; cholestyramine; fibrates; nicotinic acid; and aspirin; or pharmaceutically acceptable salts or prodrugs thereof.

Examples of anti-obesity/appetite-regulating agents include phentermine, leptin, bromocriptine, dexamphetamine, amphetamine, fenfluramine, dexfenfluramine, sibutramine, orlistat, dexfenfluramine, mazindol, phentermine, phendimetrazine, diethylpropion, fluoxetine, bupropion, topiramate, diethylpropion, benzphetamine, phenylpropanolamine or ecopipam, ephedrine, pseudoephedrine and cannabinoid receptor antagonists; or pharmaceutically acceptable salts or prodrugs thereof.

Examples of anti-hypertensive agents include loop diuretics, for example ethacrynic acid, furosemide or torsemide; diuretics, for example thiazide derivatives, chlorithiazide, hydrochlorothiazide or amiloride; angiotensin converting enzyme (ACE) inhibitors, for example benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perinodopril, quinapril, ramipril ortrandolapril; Na-K-ATPase membrane pump inhibitors, for example digoxin; neutralendopeptidase (NEP) inhibitors, for example thiorphan, tertio-thiorphan or SQ29072; ECE inhibitors, for example SLV306; dual ACE/NEP inhibitors, for example omapatrilat, sampatrilat or fasidotril; angiotensin II antagonists, for example candesartan, eprosartan, irbesartan, losartan, telmisartan or valsartan; renin inhibitors, for example aliskiren, telakiren, ditekiren, RO-66-1132 or RO-66-1168; b-adrenergic receptor blockers, for example acebutolol, atenolol, betaxolol, bisoprolol, metoprolol, nadolol, propranolol, sotalol

- 78 -

or timolol; inotropic agents, for example digoxin, dobutamine or milrinone; calcium channel blockers, for example amlodipine, bepridil, diltiazem, felodipine, nicardipine, nimodipine, nifedipine, nisoldipine or verapamil; aldosterone receptor antagonists; and aldosterone synthase inhibitors; or pharmaceutically acceptable salts or prodrugs thereof.

Examples of cholesterol absorption modulators include Zetia® and KT6-971, or pharmaceutically acceptable salts or prodrugs thereof.

Examples of aldosterone inhibitors include anastrozole, fadrazole and eplerenone, or pharmaceutically acceptable salts or prodrugs thereof.

Examples of inhibitors of platelet aggregation include aspirin or clopidogrel bisulfate, or pharmaceutically acceptable salts or prodrugs thereof.

Examples of chemotherapeutic agents include compounds decreasing the protein kinase activity, for example PDGF receptor tyrosine kinase inhibitors (e.g. imatinib or 4-methyl-N-[3-(4-methyl-imidazol-1-yl)-5-trifluoromethyl-phenyl]-3-(4-pyridin-3-yl-pyrimidin-2-ylamino)-benzamide), or pharmaceutically acceptable salts or prodrugs thereof.

Examples of 5-HT₃ or 5-HT₄ receptor modulators include tegaserod, tegaserod hydrogen maleate, cisapride or cilansetron, or pharmaceutically acceptable salts or prodrugs thereof.

The weight ratio of the compound of the present invention to the further active ingredient(s) may be varied and will depend upon the effective dose of each ingredient. Generally, an effective dose of each will be used. Thus, for example, when a compound of the present invention is combined with another agent, the weight ratio of the compound of the present invention to the other agent will generally range from about 1000: 1 to about 1: 1000, preferably about 200: 1 to about 1: 200.

Combinations of a compound of the present invention and other active ingredients will generally also be within the aforementioned range, but in each case, an effective dose of each active ingredient should be used.

- 79 -

In such combinations the compound of the present invention and other active agents may be administered separately or in conjunction. In addition, the administration of one element may be prior to, concurrent to, or subsequent to the administration of other agent(s).

Use

Compounds of the invention may be useful in the therapy of a variety of diseases and conditions.

In particular, compounds of the invention may be useful in the treatment or prevention of a disease or condition selected from non-insulin-dependent diabetes mellitus, arthritis, obesity, allograft transplantation, osteoporosis, heart failure, impaired glucose metabolism or impaired glucose tolerance, neurodegenerative diseases (for example Alzheimer's disease or Parkinson disease), cardiovascular or renal diseases (for example diabetic cardiomyopathy, left or right ventricular hypertrophy, hypertrophic medial thickening in arteries and/or in large vessels, mesenteric vasculature hypertrophy or mesangial hypertrophy), neurodegenerative or cognitive disorders, hyperglycemia, insulin resistance, lipid disorders, dyslipidemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, low HDL levels, high LDL levels, atherosclerosis, vascular restenosis, irritable bowel syndrome, inflammatory bowel disease (e.g. Crohn's disease or ulcerative colitis), pancreatitis, retinopathy, nephropathy, neuropathy, syndrome X, ovarian hyperandrogenism (polycystic ovarian syndrome), type 2 diabetes, growth hormone deficiency, neutropenia, neuronal disorders, tumor metastasis, benign prostatic hypertrophy, gingivitis, hypertension and osteoporosis.

The compounds may also be useful in producing a sedative or anxiolytic effect, attenuating post-surgical catabolic changes or hormonal responses to stress, reducing mortality and morbidity after myocardial infarction, modulating hyperlipidemia or associated conditions; and lowering VLDL, LDL or Lp(a) levels.

Examples

The following Examples illustrate the invention.

Terms used in the Examples:


ACN: acetonitrile

HPLC: high performance liquid chromatography

Cbz: carbobenzyloxy

Intermediates A1 & A2

Intermediates A1 and A2 are prepared according to Scheme A:



Scheme A

A) 3-Oxo-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester

To tropinone (20g, 142 mmol) in toluene (600 mL) are added benzyl chloroformate (42.2 mL, 284 mmol) and K_2CO_3 (118 mg, 0.853 mmol) and the resulting solution is stirred at reflux during 16h. After evaporation of the solvent, the residue is treated with dichloromethane / aqueous saturated $NaHCO_3$, the organic phase is dried, filtered and evaporated to give a yellow oil.

MS: 260 $[M + H]^+$

TLC, Rf (cyclohexane/ethyl acetate 1/1) = 0.7.

B) 3-[1-Methoxy-methylidene]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester

To methoxymethyltriphenyl phosphonium chloride (46.6 g, 133 mmol) in tetrahydrofuran (900 mL) is added dropwise at $-40^\circ C$ sodium bis(trimethylsilyl) amide in tetrahydrofuran (2M, 67 mL, 130 mmol) then the resulting red solution is stirred at $-40^\circ C$ during 1h and at $0^\circ C$ during 30 min. After cooling to $-40^\circ C$, 3-oxo-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester (27 g, 83 mmol) in tetrahydrofuran (100 mL) is added. The resulting mixture is warmed to $0^\circ C$ and stirred during 30 min. It is quenched with aqueous saturated NH_4Cl , extracted with dichloromethane and washed with aqueous saturated $NaHCO_3$. The organic phase is dried, filtered and evaporated to give a residue, which is purified by flash chromatography on silica gel (eluent: cyclohexane/ethyl acetate 100/0 to 7/3) to yield a yellow oil.

MS: 288 $[M + H]^+$

TLC, Rf (cyclohexane/ethyl acetate 1/1) = 0.75.

C) 3-exo-Formyl-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester

To 3-[1-methoxy-methylidene]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester (13g, 40.7 mmol) in acetone (450 mL) and water (50 mL) is added 37% conc HCl in water (0.920 mL, 9.4 mmol). The resulting mixture is stirred at $50^\circ C$ during 24h. After evaporation of the solvent, the residue is treated with dichloromethane / aqueous saturated $NaHCO_3$, the organic phase is dried, filtered and evaporated to give a residue, which is purified by flash chromatography on silica gel (eluent: cyclohexane/ethyl acetate 100/0 to 7/3) to yield a yellow oil.

MS: 274 $[M + H]^+$

- 82 -

TLC, Rf (cyclohexane/ethyl acetate 1/1) = 0.42.

D) 3-exo-[[[(S)-2-Methyl-propane-2-sulfinylimino]-methyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester

(S)-2-methyl-2-propanesulfinamide (543 mg, 4.39 mmol), pyridinium toluene-4-sulfonate (46 mg, 0.183 mmol) and MgSO₄ (2.21 g, 18.3 mmol) are stirred in dichloroethane (15 mL) for 1h, before the addition of 3-exo-formyl-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester (1g, 3.66 mmol) in dichloroethane (5 mL). The resulting mixture is stirred at 50°C during 16h, it is filtered and evaporated to give a residue, which is purified by flash chromatography on silica gel (eluent: cyclohexane/ethyl acetate 100/0 to 50/50) to yield a yellow oil.

MS: 377 [M + H]⁺

TLC, Rf (cyclohexane/ethyl acetate 1/1) = 0.37.

E) 3-exo-[(R)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester (Intermediate A1); and 3-exo-[(S)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester (Intermediate A2)

To stirred magnesium turnings (18 mg, 0.75 mmol) in diethyl ether (0.5 mL) is added 2,4,5-trifluorobenzylbromide (112 mg, 0.499 mmol) in diethyl ether (0.5mL). After stirring during 30 min at rt, the resulting suspension is added at 0°C to a solution of 3-exo-[[[(S)-2-methyl-propane-2-sulfinylimino]-methyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester (179 mg, 0.333 mmol) in dichloromethane (1 mL). The resulting mixture is stirred for 4h at rt, before it is quenched at 0°C with saturated aqueous NH₄Cl and extracted with dichloromethane. The organic phase is dried, filtered and evaporated to give a residue, which is purified by flash chromatography on silica gel (eluent: cyclohexane/ethyl acetate).

Intermediate A1: MS: 523 [M + H]⁺


TLC, Rf (cyclohexane/ethyl acetate 1/1) = 0.17

Intermediate A2: MS: 523 [M + H]⁺

TLC, Rf (cyclohexane/ethyl acetate 1/1) = 0.12

Intermediates A'1 & A'2

Intermediates A'1 and A'2 are prepared according to Scheme A' :



Scheme A'

4-[(R)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidine-1-carboxylic acid benzyl ester (Intermediate A1); and 4-[(S)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidine-1-carboxylic acid benzyl ester (Intermediate A2)

The title compounds are prepared analogously as described in Scheme A using 4-Formyl-piperidine-1-carboxylic acid benzyl ester instead of 3-exo-formyl-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester

Intermediate A'1: MS: 497.1 [M + H]⁺

TLC, R_f (cyclohexane/ethyl acetate 1/1) = 0.23


Intermediate A'2: MS: 497.1 [M + H]⁺

TLC, R_f (cyclohexane/ethyl acetate 1/1) = 0.10

Example B1: N-(2-{3-exo-[(S)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide

This compound is prepared according to Scheme B:

- 84 -



Scheme B

A) 3-*exo*-[(S)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane

To 3-*exo*-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester (Intermediate A2; 300mg, 0.574 mmol) in ethylacetate (10 mL) is added 10% Pd/C (610mg, 0.57mmol). The resulting mixture is stirred and put under H₂-atmosphere. After 1h of stirring, the suspension is filtered through celite

and evaporated to give a residue, which is purified by preparative HPLC (Column Interchrom C18 ODB 10µm 28x250, Gradient: 0-2.5 min 5% ACN, 2.5-25.5 min 5-100% ACN, 25.5-30 min 100% CAN) to yield the title compound.

MS: 389 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.06 min.

B) N-(2-{3-exo-[(S)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-phthalimide

To 3-exo-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane (540 mg, 1.2 mmol) in dichloromethane/1M NaOH (1/1, 5 mL) is added 2-phthalimidoethane sulfonyl chloride (1.101 g, 3.59 mmol) then the resulting mixture is stirred at 50°C during 16h. It is extracted with dichloromethane, the organic phase is dried and evaporated to give a white solid.

MS: 626 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.44 min.

C) 2-{3-exo-[(S)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine

To N-(2-{3-exo-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-phthalimide (100 mg, 0.160 mmol) in ethanol (2 mL) is added hydrazine monohydrate (0.396 mL, 8 mmol) and the resulting solution is stirred at rt during 12h. After extraction with dichloromethane and aqueous saturated NaHCO₃, the organic phase is dried and evaporated to give a residue, which is purified by preparative HPLC (Column Waters C18 ODB 5µm 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield a white solid.

MS 496 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.05 min.

D) N-(2-{3-exo-[(S)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide

- 86 -

To 2-{3-exo-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine (27 mg, 0.054 mmol) in acetic acid (1 mL) is added acetic anhydride (7.86 μ L, 0.081 mmol) and the resulting solution is stirred at rt during 3h. After extraction with dichloromethane and aqueous saturated NaHCO₃, the organic phase is dried and evaporated to give a colorless gum.

MS 538 [M+H]⁺

E) N-(2-{3-exo-[(S)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide

To N-(2-{3-exo-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide (29.3 mg, 0.545 mmol) in dioxane (3 mL) is added 4N HCl in dioxane (2 mL). The resulting mixture is stirred at rt during 1h. The solution is frozen and lyophilised to give a white solid.


MS: 435 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.95 min.

Example C1: N-(2-{3-exo-[(S)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-acetamide

This compound is prepared according to Scheme C:

- 87 -



Scheme C

A) N-(2-{3-exo-[(S)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-phthalimide

To a solution of N-phthaloylglycine (291 mg, 1.41 mmol) in acetonitrile (7 mL) is added benzotriazol-1-yloxytripyrrolidinophosphonium hexafluorophosphate (805 mg, 1.54 mmol) and the resulting mixture is stirred at rt during 1h before addition of 3-exo-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane

- 88 -

(500 mg, 1.29 mmol) and triethylamine (720 μ L, 5.16 mmol) in acetonitrile (5 mL). The resulting solution is stirred at rt during 16h and is evaporated before purification by preparative HPLC (Column Interchrom C18 ODB 10 μ m 28x250, Gradient: 0-2.5 min 5% ACN, 2.5-25.5 min 5-100% ACN, 25.5-30 min 100% ACN) to yield a yellow solid.

MS: 576 [M+H]⁺

B) N-(2-{3-exo-[(S)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-acetamide

The title compound is prepared analogously as described in example B1 using N-(2-{3-exo-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-phthalimide instead of N-(2-{3-exo-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-phthalimide.

MS: 384 [M+H]⁺

HPLC (Nucleosil 100-5 C18, 10min method (0-1min 10%ACN, 1-6min 10-100%ACN, 6-8.5min 100%ACN, 8.5-9min 100-10%ACN, 9-10min 10%ACN): 3.28 min.

Example C2: Cyclopropanesulfonic acid (2-{3-exo-[(S)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

This compound is also prepared according to Scheme C.

A) Cyclopropanesulfonic acid (2-{3-exo-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

To a solution of 2-{3-exo-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethylamine (50 mg, 0.112 mmol) in dichloromethane (1 mL) are added triethylamine (32 μ L, 0.224 mmol) and cyclopropanesulfonyl chloride (14 μ L, 0.134 mmol). The resulting solution is stirred at rt during 16h before evaporation and purification by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield a colorless gum.

MS 550 [M+H]⁺

- 89 -

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.17 min.

B) Cyclopropanesulfonic acid (2-{3-exo-[(S)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

To cyclopropanesulfonic acid (2-{3-exo-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide (27 mg, 0.049 mmol) in dioxane (0.5 mL) is added 4N HCl in dioxane (0.5 mL). The resulting mixture is stirred at rt during 1h. The solution is frozen and lyophilised to give a white solid before purification by preparative HPLC (Column Waters C18 ODB 5µm 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield a colorless gum.


MS 446 [M+H]⁺

HPLC (Nucleosil 100-5 C18, 10min method (0-1min 10%ACN, 1-6min 10-100%ACN, 6-8.5min 100%ACN, 8.5-9min 100-10%ACN, 9-10min 10%ACN): 3.5 min.

Example D1: 5-Methyl-pyrazine-2-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

This compound is prepared according to Scheme D:

- 90 -



Scheme D

- 91 -

A) 3-*exo*-[(*R*)-1-((*S*)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane

To 3-*exo*-[(*R*)-1-((*S*)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester (Intermediate A1; 7.74g, 14.8 mmol) in ethanol (200 mL) is added 10% Pd/C (16g, 15mmol), the resulting mixture is stirred and put under H₂-atmosphere. After 1h of stirring, the suspension is filtered through celite and evaporated to yield the title compound.

MS: 389 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.91 min.

B) N-(2-{3-*exo*-[(*R*)-1-((*S*)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-phthalimide

To 3-*exo*-[(*R*)-1-((*S*)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane (265 mg, 0.68 mmol) in dichloromethane/1M NaOH (1/1, 5 mL) is added 2-phthalimidoethane sulfonyl chloride (577 mg, 2.05 mmol) then the resulting mixture is stirred at 50°C during 16h. It is extracted with dichloromethane, the organic phase is dried and evaporated to give a white solid.

MS: 626 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.25 min.

C) 2-{3-*exo*-[(*R*)-1-((*S*)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine

To N-(2-{3-*exo*-[(*R*)-1-((*S*)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-phthalimide (256mg, 0.409mmol) in ethanol (3 mL) is added hydrazine monohydrate (1.01 mL, 20.5 mmol) and the resulting solution is stirred at rt during 12h. After extraction with dichloromethane and aqueous saturated NaHCO₃, the organic phase is dried and evaporated to yield a colorless oil.

MS: 496 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.95min.

- 92 -

D) 5-Methyl-pyrazine-2-carboxylic acid (2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

To 5-methylpyrazinecarboxylic acid (18.8 mg, 0.133 mmol) in acetonitrile (2 mL) is added benzotriazol-1-yloxytripyrrolidinophosphonium hexafluorophosphate (75.6 mg, 0.145 mmol) and the resulting mixture is stirred at rt during 1h. After this time, 2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine (60 mg, 0.121) and triethylamine (50.5 μ L, 0.363 mmol) in acetonitrile are added and the solution is shaken at rt during 16h. The solution is purified by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% CAN) to yield a white solid.

MS: 616 [M +H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.18min.

E) 5-Methyl-pyrazine-2-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

To 5-methyl-pyrazine-2-carboxylic acid (2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide (48.5 mg, 0.0788 mmol) in dioxane (3 mL) is added 4N HCl in dioxane (1 mL). The resulting mixture is stirred at rt during 1h before it is purified by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% CAN) to yield a white solid .

MS 513 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.95 min.

Example D2: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

This compound is prepared according to Scheme D.

- 93 -

A) Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

To a solution of 2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine (40 mg, 0.078 mmol) in dichloroethane (2 mL) are added triethylamine (32.7 μ L, 0.235 mmol), 4-dimethylaminopyridine (0.95 mg, 0.007 mmol) and cyclopropanesulfonyl chloride (16.8 μ L, 0.157 mmol). The resulting solution is stirred at rt during 2h before evaporation to yield the title compound.

MS: 600 [M+H]⁺

HPLC (Waters Symmetry C18 3.5 μ m 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 3.28 min.

B) Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

To Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide (63 mg, 0.102 mmol) in dioxane (2 mL) is added 4N HCl in dioxane (3 mL). The resulting mixture is stirred at rt during 1h. The solution is frozen and lyophilised to give a white solid before purification by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield title compound.


MS: 496 [M+H]⁺

HPLC (Waters Symmetry C18 3.5 μ m 2.1 x 50mm, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 5%ACN): 2.56 min.

Example D2a: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide maleate

This compound is prepared according to Scheme Da:

- 94 -




Scheme Da

A solution of maleic acid in ethyl acetate (1 molequivalent; 0.1 M) is added to a solution of Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide freebase (20 mg) in ethyl acetate (1 mL) under stirring. The obtained mixture is left to precipitate for one hour before it is stirred overnight. The resulting white suspension is filtered and the obtained solid is dried to yield the title compound.

Example D2b: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide toluene-4-sulfonate

This compound is prepared according to Scheme Db:

- 95 -



Scheme Db

A solution of toluene-4-sulfonic acid in ethyl acetate (1 molequivalent; 0.1 M) is added to a solution of Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide freebase (50 mg) in ethyl acetate (1 mL) under stirring. The obtained mixture is left to precipitate for 2 hours before it is stirred overnight. The resulting white suspension is filtered and the obtained solid is dried to yield the title compound.

Example D3: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide

This compound is prepared according to Scheme D:

A) N-(2-{3-exo-[(R)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide

To 2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine (100 mg, 0.202 mmol) in acetic acid (2 mL) is added acetic anhydride (29.1 μ L, 0.303 mmol) and the resulting solution is stirred at rt during 3h. After extraction with dichloromethane and aqueous saturated NaHCO_3 , the organic phase is dried and evaporated to give the title compound.

MS: 538 $[\text{M}+\text{H}]^+$

B) N-(2-{3-*exo*-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide

To N-(2-{3-*exo*-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide (108 mg, 0.199 mmol) in dioxane (2 mL) is added 4N HCl in dioxane (3 mL). The resulting mixture is stirred at rt during 1h. The solution is frozen and lyophilised to give a white solid before purification by preparative HPLC (Column Waters C18 ODB 5µm 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield the title compound.

MS: 434 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.84 min.

Example D3a: N-(2-{3-*exo*-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide maleate

The title compound is prepared analogously as described in example D2a using N-(2-{3-*exo*-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide instead of cyclopropanesulfonic acid (2-{3-*exo*-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example D3b: N-(2-{3-*exo*-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-(2-{3-*exo*-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-acetamide instead of cyclopropanesulfonic acid (2-{3-*exo*-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example D4: Dimethylsulfamic acid (2-{3-*exo*-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

The title compound is prepared analogously as described in example D2 using dimethylsulfamoyl chloride instead of cyclopropanesulfonyl chloride.

MS: 499 [M+H]

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 0.94min.

Example D4a: Dimethylsulfamic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide maleate

The title compound is prepared analogously as described in example D2a using Dimethylsulfamic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example D4b: Dimethylsulfamic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using Dimethylsulfamic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example D5: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-4-methoxy-benzenesulfonamide

The title compound is prepared analogously as described in example D2 using 4-methoxyphenylsulfonyl chloride instead of cyclopropanesulfonyl chloride.

MS: 562 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.0 min.

Example D5a: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-4-methoxy-benzenesulfonamide maleate

The title compound is prepared analogously as described in example D2a using N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-4-methoxy-benzenesulfonamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example D5b: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-4-methoxy-benzenesulfonamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-4-methoxy-benzenesulfonamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example D6: Tetrahydropyran-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

The title compound is prepared analogously as described in example D1 using tetrahydro-2H-pyran-4-carboxylic acid instead of 5-methylpyrazinecarboxylic acid.

MS: 505 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.92 min.

Example D7: Morpholine-4-sulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

The title compound is prepared analogously as described in example D2 using 4-morpholinesulfonyl chloride instead of cyclopropanesulfonyl chloride.

MS: 541 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.13 min.

Example D8: 2-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

A) 2-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonic acid (2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

To a solution of 2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine (100 mg, 0.202 mmol) in dichloromethane / 2N NaOH (2 mL, 1/1) are 2-phtalimidoethanesulfonyl chloride (167 mg, 0.605 mmol). The resulting solution is stirred at 50°C during 2h before extraction and evaporation of organic phase to yield the crude compound. It is purified by preparative HPLC (Column Waters C18 ODB 5µm 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield title compound.

MS: 734 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.43 min.

B) 2-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

To 2-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonic acid (2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide (49 mg, 0.057 mmol) in dioxane (3 mL) is added 4N HCl in dioxane (3 mL). The resulting mixture is stirred at rt during 1h. The solution is frozen and lyophilised to give a white solid before purification by preparative HPLC (Column Waters C18 ODB 5µm 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield title compound.

MS: 629 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.22 min.

- 100 -

Example D9: **N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionamide**

The title compound is prepared analogously as described in example D1 using 2-phtamididoethanecarboxylic acid instead of 5-methylpyrazinecarboxylic acid.

MS: 593 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.8 min.

Example D10: **(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-carbamic acid 2-methoxy-ethyl ester**

The title compound is prepared analogously as described in example D8 using chloroformic acid 2-methoxyethylester instead of 2-phtalimidoethanesulfonyl chloride.

MS: 494 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.14 min.

Example D10a: **(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-carbamic acid 2-methoxy-ethyl ester maleate**

The title compound is prepared analogously as described in example D2a using (2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-carbamic acid 2-methoxy-ethyl ester instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example D10b: **(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-carbamic acid 2-methoxy-ethyl ester toluene-4-sulfonate**

The title compound is prepared analogously as described in example D2b using (2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-carbamic acid 2-methoxy-ethyl ester instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example D11: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-4-methoxy-benzamide

The title compound is prepared analogously as described in example D1 using p-anisic acid instead of 5-methylpyrazinecarboxylic acid.

MS: 526 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.18 min.

Example D12: (2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-carbamic acid 1,1-dioxo-benzothiophen-2-yl-methyl ester

The title compound is prepared analogously as described in example D8 using 1,1-dioxobenzothiophen-2-yl-methylchloroformate instead of 2-phtalimidoethanesulfonyl chloride.

MS: 614 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.21min.

Example D13: (2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-carbamic acid ethyl ester

The title compound is prepared analogously as described in example D2 using ethylchloroformate instead of cyclopropanesulfonyl chloride.

MS: 464 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.96 min.

Example D14: Pyrrolidine-1-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

The title compound is prepared analogously as described in example D2 using 1-pyrrolidinecarbonyl chloride instead of cyclopropanesulfonyl chloride.

MS: 489 [M+H]

- 102 -

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.97 min.

Example D15: Morpholine-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

The title compound is prepared analogously as described in example D2 using 4-morpholinecarbonyl chloride instead of cyclopropanesulfonyl chloride.

MS: 505 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.92 min.

Example D15a: Morpholine-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide maleate

The title compound is prepared analogously as described in example D2a using morpholine-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example D15b: Morpholine-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using morpholine-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example D16: 1-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-3-(2-methoxy-ethyl)-urea

A) (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-(3-exo-8-{2-[3-(2-methoxy-ethyl)-ureido]-ethanesulfonyl]-8-aza-bicyclo[3.2.1]oct-3-yl)-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide

- 103 -

To a solution of 2-methoxyethylamine (29.8 μ L, 0.344 mmol) in dimethyl formamide (2 mL) is added 1,1'-carbonyldiimidazole (58.6 mg). After stirring at rt, 2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine (180 mg, 0.312 mmol) is added. The resulting solution is stirred at 80°C during 3h before purification by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield title compound.

MS: 597 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.32 min.

B) 1-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-3-(2-methoxy-ethyl)-urea

To (S)-2-methyl-propane-2-sulfinic acid [(R)-1-(3-exo-8-{2-[3-(2-methoxy-ethyl)-ureido]-ethanesulfonyl}-8-aza-bicyclo[3.2.1]oct-3-yl)-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide (143 mg, 0.206 mmol) in dioxane (3 mL) is added 4N HCl in dioxane (3 mL). The resulting mixture is stirred at rt during 1h. The solution is frozen and lyophilised to give a white solid before purification by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield title compound.

MS: 493 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.12 min.

Example D17: 2-{1,3-Dioxo-1,3-dihydro-isoindol-2-yl}-ethanesulfonic acid (2-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidine-1-sulfonyl}-ethyl)-amide

The title compound is prepared analogously as described in example D8 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane.

MS: 603 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.17 min.

Example D18: N-[2-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylsulfamoyl)-ethyl]-benzamide

A) 2-Amino-ethanesulfonic acid (2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide

The title compound is prepared analogously as described in example D1 using 2-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonic acid (2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide instead of N-(2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-phthalimide.

MS: 603 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.22 min.

B) N-[2-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylsulfamoyl)-ethyl]-benzamide

The title compound is prepared analogously as described in example D1 using 2-amino-ethanesulfonic acid (2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide instead of 2-{3-exo-[(R)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine and benzoic acid instead of 5-methyl-pyrazine-2-carboxylic acid.

MS: 601 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.16 min.

Example D19: Oxazole-5-carboxylic acid [2-(2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylsulfamoyl)-ethyl]-amide

The title compound is prepared analogously as described in example D1 using oxazole-5-carboxylic acid instead of 5-methylpyrazinecarboxylic acid.


MS: 488 [M+H]

- 105 -

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.17 min.

Example E1: Cyclopropanecarboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluorophenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

This compound is prepared according to Scheme E:



Scheme E

A) (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-{3-exo-8-[2-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-acetyl]-8-aza-bicyclo[3.2.1]oct-3-yl}-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide

To a solution of N-phthaloylglycine (291 mg, 1.41 mmol) in acetonitrile (7 mL) is added benzotriazol-1-yloxytripyrrolidinophosphonium hexafluorophosphate (805 mg, 1.54 mmol) and the resulting mixture is stirred at rt during 1h before addition of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane (500 mg, 1.29 mmol) and triethylamine (720 μ L, 5.16 mmol) in acetonitrile (5 mL). The resulting solution is stirred at rt during 16h and is evaporated before purification by preparative HPLC (Column Interchrom C18 ODB 10 μ m 28x250, Gradient: 0-2.5 min 5% ACN, 2.5-25.5 min 5-100% ACN, 25.5-30 min 100% ACN) to yield a yellow solid.

MS: 576 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.22 min.

B) (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-{3-exo-8-(2-amino-acetyl)-8-aza-bicyclo[3.2.1]oct-3-yl}-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide

To (S)-2-methyl-propane-2-sulfinic acid [(R)-1-{3-exo-8-[2-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-acetyl]-8-aza-bicyclo[3.2.1]oct-3-yl}-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide (300mg, 0.521mmol) in ethanol (3.5 mL) is added hydrazine monohydrate (1.3 mL, 26.05 mmol) and the resulting solution is stirred at rt during 16h. After extraction with dichloromethane and aqueous saturated NaHCO₃, the organic phase is dried and evaporated to yield a white solid.

MS: 446 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.96min.

C) Cyclopropanecarboxylic acid (2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

To a solution of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-{3-exo-8-(2-amino-acetyl)-8-aza-bicyclo[3.2.1]oct-3-yl}-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide (50 mg, 0.112 mmol) in dichloromethane (1 mL) are added triethylamine (32 μ L, 0.224 mmol) and cyclopropanecarbonyl chloride (14 μ L, 0.134 mmol). The resulting solution is stirred at rt

- 107 -

during 2h before evaporation and purification by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield a white solid.

MS: 514 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 2.87 min.

D) Cyclopropanecarboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

To cyclopropanecarboxylic acid (2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide (26.4 mg, 0.052 mmol) in dioxane (0.5 mL) is added 4N HCl in dioxane (0.5 mL). The resulting mixture is stirred at rt during 1h. The solution is frozen and lyophilised to give a white solid.

MS: 410 [M+H]⁺

HPLC (Nucleosil 100-5 C18, 10min method (0-1min 10%ACN, 1-6min 10-100%ACN, 6-8.5min 100%ACN, 8.5-9min 100-10%ACN, 9-10min 10%ACN): 3.55 min.

Example E1a: Cyclopropanecarboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide maleate

The title compound is prepared analogously as described in example D2a using Cyclopropanecarboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example E1b: Cyclopropanecarboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using Cyclopropanecarboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example E2: 5-Methyl-pyrazine-2-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

This compound is also prepared according to Scheme E and analogously to example D1 using 2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethylamine instead of 2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine.

MS: 462 [M+H]

HPLC (Nucleosil 100-5 C18, 10min method (0-1min 10%ACN, 1-6min 10-100%ACN, 6-8.5min 100%ACN, 8.5-9min 100-10%ACN, 9-10min 10%ACN): 3.59 min.

Example E3: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

This compound is prepared analogously to example E1 using cyclopropanesulfonyl chloride instead of cyclopropanecarbonyl chloride.

MS: 446 [M+H]

HPLC (Nucleosil 100-5 C18, 10min method (0-1min 10%ACN, 1-6min 10-100%ACN, 6-8.5min 100%ACN, 8.5-9min 100-10%ACN, 9-10min 10%ACN): 3.56 min.

Example E3a: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide maleate

The title compound is prepared analogously as described in example D2a using cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example E3b: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-

- 109 -

bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example E4: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-acetamide

A) N-(2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-acetamide

To 2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethylamine (50 mg, 0.112 mmol) in acetic acid (2 mL) is added acetic anhydride (16 μ L, 0.168 mmol) and the resulting solution is stirred at rt during 3h. After extraction with dichloromethane and aqueous saturated NaHCO₃, the organic phase is dried and evaporated to give a residue which is purified by preparative HPLC (Column Interchrom C18 ODB 10 μ m 28x250, Gradient: 0-2.5 min 5% ACN, 2.5-25.5 min 5-100% ACN, 25.5-30 min 100% ACN) to yield a colorless gum.

MS: 488 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.09 min.

B) N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-acetamide

To N-(2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-acetamide (38 mg, 0.078 mmol) in dioxane (0.5 mL) is added 4N HCl in dioxane (0.5 mL). The resulting mixture is stirred at rt during 1h. The solution is frozen and lyophilised to give a white solid before purification by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield a light yellow solid.

MS 384 [M+H]⁺

HPLC (Nucleosil 100-5 C18, 10min method (0-1min 10%ACN, 1-6min 10-100%ACN, 6-8.5min 100%ACN, 8.5-9min 100-10%ACN, 9-10min 10%ACN): 3.28 min.

Example E5: Morpholine-4-sulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-amide

- 110 -

The title compound is prepared analogously as described in example E1 using morpholinesulfonyl chloride instead of cyclopropanecarbonyl chloride.

MS: 491 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.92 min.

Example E6: Morpholine-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example E1 using morpholinecarbonyl chloride instead of cyclopropanecarbonyl chloride.

MS: 455 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.87 min.

Example E6a: Morpholine-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide maleate

The title compound is prepared analogously as described in example D2a using morpholine-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example E6b: Morpholine-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using morpholine-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example E7: 1-Hydroxy-cyclopropanecarboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

- 111 -

The title compound is prepared analogously as described in example E2 using 1-hydroxycyclopropane-1-carboxylic acid instead of 4-methylpyrazinecarboxylic acid.

MS: 426 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.87 min.

Example E8: Tetrahydropyran-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example E2 using tetrahydropyran-4-carboxylic acid instead of 4-methylpyrazinecarboxylic acid.

MS: 454 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.18 min.

Example E9: Cyclobutanecarboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example E2 using cyclobutanecarboxylic acid instead of 4-methylpyrazinecarboxylic acid.

MS: 424 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.22 min.

Example E10: 3-Methyl-3H-imidazole-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example E2 using 3-methyl-3H-imidazole-4-carboxylic acid instead of 4-methylpyrazinecarboxylic acid.

MS: 449 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.76 min.

Example E11: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-4-fluoro-benzamide

- 112 -

This compound is prepared analogously to example E1 using 4-fluorobenzoyl chloride instead of cyclopropanecarbonyl chloride.

MS: 464 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.13 min.

Example E11a: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-4-fluoro-benzamide maleate

The title compound is prepared analogously as described in example D2a using N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-4-fluoro-benzamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example E11b: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-4-fluoro-benzamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-4-fluoro-benzamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example E12: 3H-Imidazole-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

This compound is prepared analogously to example E2 using N-boc-3H-imidazole-4-carboxylic acid instead of 4-methylpyrazinecarboxylic acid.

MS: 436 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.81 min.

Example E13: Pyrazine-2-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

- 113 -

This compound is prepared analogously to example E2 using pyrazine-2-carboxylic acid instead of 4-methylpyrazinecarboxylic acid.

MS: 448 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.2 min.

Example E14: 4-Methyl-oxazole-5-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

This compound is prepared analogously to example E2 using 4-methyl-oxazole-5-carboxylic acid instead of 4-methylpyrazinecarboxylic acid.

MS: 451 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.9 min.

Example E15: 2-Amino-pyrimidine-5-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

This compound is prepared analogously to example E2 using 2-amino-pyrimidine-5-carboxylic acid instead of 4-methylpyrazinecarboxylic acid.

MS: 463 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.11 min.

Example E16: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-2-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-acetamide

This compound is prepared analogously to example E2 using N-phthaloylglycine instead of 4-methylpyrazinecarboxylic acid.

MS: 530 [M-H]⁺

HPLC (waters symmetry C18, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 5%ACN): 2.87 min.

Example E17: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-4-fluoro-benzenesulfonamide

- 114 -

This compound is prepared analogously to example E1 using 4-fluorophenylsulfonyl chloride instead of cyclopropanecarbonyl chloride.

MS: 500 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.14 min.

Example E18: N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionamide

A) (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-[3-exo-8-(3-amino-propionyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide

This compound is prepared analogously to example E1 using N-phthaloyl-beta-glycine instead of N-phthaloylglycine.

MS: 460 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.17 min.

B) N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionamide

This compound is prepared analogously to example E2 using N-phthaloyl-beta-glycine instead of 4-methylpyrazinecarboxylic acid.

MS: 557 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.17 min.

Example E19: Cyclopropanesulfonic acid (3-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-amide

This compound is prepared analogously to example E3 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-[3-exo-8-(3-amino-propionyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-[3-exo-8-(2-amino-acetyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

- 115 -

MS: 460 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.11 min.

Example E20: N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-acetamide

This compound is prepared analogously to example E4 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-[3-exo-8-(3-amino-propionyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-[3-exo-8-(2-amino-acetyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 398 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.07 min.

Example E20a: N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-acetamide maleate

The title compound is prepared analogously as described in example D2a using N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-acetamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example E20b: N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-acetamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-acetamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example E21: N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-benzamide

This compound is prepared analogously to example E19 using benzoylchloride instead of cyclopropanesulfonyl chloride.

MS: 460 [M-H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.12 min.

Example E22: Cyclopropanecarboxylic acid (2-{4-[(R)-1-amino-2-(2,4,5-trifluorophenyl)-ethyl]-piperidin-1-yl}-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example E1 (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane.

MS: 384 [M+H]

HPLC (YMC Pack ODS-AQ 3 µm 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.6 min.

Example E23: N-(2-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-oxo-ethyl)-benzamide

This compound is prepared analogously to example E22 using benzoylchloride instead of cyclopropanecarbonyl chloride.

MS: 420 [M+H]

HPLC (YMC Pack ODS-AQ 3 µm 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.83 min.

Example E24: Cyclopropanecarboxylic acid (2-[(1S,3S,5R)-3-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-2-oxo-ethyl)-isopropyl-amide

A) (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-[(1S,5R)-8-(2-isopropylamino-acetyl)-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide

To (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-[(1S,5R)-8-(2-amino-acetyl)-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide (150 mg, 0.33 mmol) in methanol (3 mL) is added acetic acid until to have pH 5-5.5. After addition of acetone (124 µL, 1.68 mmol), the reaction

- 117 -

is stirred at rt during 1h and sodium cyanoborohydride (43 mg, 0.67 mmol) is added before stirring at rt during 16h. The reaction is quenched with ethylacetate and an aqueous saturated NaHCO₃ solution, the organic phase is dried and evaporated to afford the crude compound before purification onto a SCX-2 cartridge (5g, DCM/MeOH 1:1, then 2N NH₃ in MeOH) to yield the title compound.

MS: 488 [M+H]

HPLC (Luna 3 microns C18(2) 30 x 4.6mm, 6 min method, 0-0.5 min 5% ACN, 0.5-5.5min 5-95% ACN, 5.5-6 min 5% ACN): 2.17 min.

B) Cyclopropanecarboxylic acid (2-((1S,3S,5R)-3-((R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-isopropyl-amide

This compound is prepared analogously to example E1 using (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-((1S,5R)-8-(2-isopropylamino-acetyl)-bicyclo[3.2.1]oct-3-yl)-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-[3-exo-8-(2-amino-acetyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 452 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 6.39 min.

Example E25: N-(2-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-3-phenyl-propionamide

The title compound is prepared analogously as described in example E2 using 3-Phenylpropionic acid instead of 4-methylpyrazinecarboxylic acid.

MS:474 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 6.74 min.

Example E25a: N-(2-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-3-phenyl-propionamide maleate

The title compound is prepared analogously as described in example D2a using N-(2-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-yl)-2-

oxo-ethyl)-3-phenyl-propionamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example E25b: N-(2-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-2-oxo-ethyl)-3-phenyl-propionamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-(2-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-2-oxo-ethyl)-3-phenyl-propionamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example E26: N-(2-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-2-oxo-ethyl)-2-phenoxy-acetamide

The title compound is prepared analogously as described in example E2 using 2-Phenoxyacetic acid instead of 4-methylpyrazinecarboxylic acid.

MS:476 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 6.67 min.

Example E27: N-(2-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-2-oxo-ethyl)-2-(3-methyl-isoxazol-5-yl)-acetamide

The title compound is prepared analogously as described in example E2 using (3-Methyl-isoxazol-5-yl)-acetic acid instead of 4-methylpyrazinecarboxylic acid.

MS:465 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 5.76 min.

Example E28: N-(2-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-2-oxo-ethyl)-2-morpholin-4-yl-acetamide

The title compound is prepared analogously as described in example E2 using Morpholin-4-yl-acetic acid instead of 4-methylpyrazinecarboxylic acid.

MS:469 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 4.09 min.

Example E29: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-2-phenyl-acetamide

The title compound is prepared analogously as described in example E2 using Phenylacetic acid instead of 4-methylpyrazinecarboxylic acid.

MS:460 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 6.38 min.

Example E30: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-C-phenyl-methanesulfonamide

The title compound is prepared analogously as described in example E1 using Phenylmethanesulfonyl chloride instead of cyclopropanecarbonyl chloride.

MS:496 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 6.76 min.

Example E30a: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-C-phenyl-methanesulfonamide maleate

The title compound is prepared analogously as described in example D2a using N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-C-phenyl-methanesulfonamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example E30b: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-C-phenyl-methanesulfonamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-

oxo-ethyl)-C-phenyl-methanesulfonamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example E31: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-2-[1,2,4]triazol-1-yl-isobutyramide

The title compound is prepared analogously as described in example E2 using 2-Methyl-2-[1,2,4]triazol-1-yl-propionic acid instead of 4-methylpyrazinecarboxylic acid.

MS:479 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 5.53 min.

Example E32: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-2-(tetrahydro-pyran-4-yl)-acetamide

The title compound is prepared analogously as described in example E2 using (Tetrahydro-pyran-4-yl)-acetic acid instead of 4-methylpyrazinecarboxylic acid.

MS:468 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 5.56 min.

Example E33: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-2-methoxy-benzamide

The title compound is prepared analogously as described in example E2 using 2-methoxybenzoic acid instead of 4-methylpyrazinecarboxylic acid.

MS:476 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 6.62 min.

Example E34: 2,3-Dihydro-benzo[1,4]dioxine-5-carboxylic acid (2-((1S,3S,5R)-3-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-amide

- 121 -

The title compound is prepared analogously as described in example E2 using 2,3-Dihydrobenzo[1,4]dioxine-5-carboxylic acid instead of 4-methylpyrazinecarboxylic acid.

MS:504 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 6.43 min.

Example E35: Pyridazine-3-carboxylic acid (2-((1S,3S,5R)-3-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example E2 using Pyridazine-3-carboxylic acid instead of 4-methylpyrazinecarboxylic acid.

MS:448 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 5.56 min.

Example E36: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-2-(3H-imidazol-4-yl)-acetamide

The title compound is prepared analogously as described in example E2 using (3H-Imidazol-4-yl)-acetic acid instead of 4-methylpyrazinecarboxylic acid.

MS:450 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 0.84-3.14 min.

Example E37: 2-Phenyl-ethanesulfonic acid (2-((1S,3S,5R)-3-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example E1 using Phenylethanesulfonyl chloride instead of cyclopropanecarbonyl chloride.

MS:510 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 6.35 min.

Example E38: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-4-methyl-benzenesulfonamide

The title compound is prepared analogously as described in example E1 using toluenesulfonyl chloride instead of cyclopropanecarbonyl chloride.

MS:496 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 6.11 min.

Example E39: 2,3-Dihydro-benzo[1,4]dioxine-6-sulfonic acid (2-((1S,3S,5R)-3-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example E1 using 2,3-Dihydro-benzo[1,4]dioxine-6-sulfonyl chloride instead of cyclopropanecarbonyl chloride.

MS:540 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 5.87 min.

Example E40: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-2-methoxy-benzenesulfonamide

The title compound is prepared analogously as described in example E1 using 2-Methoxybenzenesulfonyl chloride instead of cyclopropanecarbonyl chloride.

MS:512 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 5.74 min.

Example E40a: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-2-methoxy-benzenesulfonamide maleate

The title compound is prepared analogously as described in example D2a using N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-2-methoxy-benzenesulfonamide instead of cyclopropanesulfonic acid (2-(3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example E40b: N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-2-methoxy-benzenesulfonamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-(2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-2-methoxy-benzenesulfonamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example E41: 3,5-Dimethyl-isoxazole-4-sulfonic acid (2-((1S,3S,5R)-3-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example E1 using 3,5-Dimethyl-isoxazole-4-sulfonyl chloride instead of cyclopropanecarbonyl chloride.

MS:501 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 5.55 min.

Example E42: 1,3,5-Trimethyl-1H-pyrazole-4-sulfonic acid (2-((1S,3S,5R)-3-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example E1 using 1,3,5-Trimethyl-1H-pyrazole-4-sulfonyl chloride instead of cyclopropanecarbonyl chloride.


MS:514 [M+H]

HPLC (Higgins Cliepus 5 microns C18(2) 100 x 3mm, 25 min method, 0-1 min 5% ACN, 1-15min 5-95% ACN, 15-20 95% CAN, 20-22 min 95-5% CAN, 22-25min 5% ACN): 5.15 min.

Example F1: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-2-oxo-ethyl)-methyl-amide

This compound is prepared according to Scheme F, in which "Cbz" is carbobenzoxy:

- 124 -



Scheme F

A) Methyl-(2-{3-*exo*-[(*R*)-1-((*S*)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-carbamic acid benzyl ester

The title compound is prepared analogously as described in example E1 using N-methyl-N-Cbz-glycine instead of N-phthaloylglycine.

MS: 594 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.25 min.

- 125 -

B) Methyl-2-{3-*exo*-[(*R*)-1-((*S*)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethylamine

To methyl-(2-{3-*exo*-[(*R*)-1-((*S*)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-carbamic acid benzyl ester (340 mg, 0.573 mmol) in ethanol (5 mL) is added palladium on charcoal (122 mg, 0.114mmol) and the resulting mixture is stirred at rt during 4h under H₂-atmosphere. The suspension is filtered through celite and the filtrate is evaporated to yield a grey solid.

MS: 460 [M+H]

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.97 min.

C) Cyclopropanesulfonic acid methyl-(2-{3-*exo*-[(*R*)-1-((*S*)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide

To a solution of methyl-2-{3-*exo*-[(*R*)-1-((*S*)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethylamine (100 mg, 0.217 mmol) in dichloromethane (2 mL) are added triethylamine (91 µL, 0.651 mmol) and cyclopropanesulfonyl chloride (33 µL, 0.326 mmol). The resulting solution is stirred at rt during 3h before evaporation and purification by preparative HPLC (Column Waters C18 ODB 5µm 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield a white solid.

MS: 564 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.18 min.

D) Cyclopropanesulfonic acid (2-{3-*exo*-[(*R*)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-methyl-amide

To cyclopropanesulfonic acid methyl-(2-{3-*exo*-[(*R*)-1-((*S*)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-amide (62 mg, 0.11 mmol) in dioxane (1 mL) is added 4N HCl in dioxane (1 mL). The resulting mixture is stirred at rt during 1h. The solution is frozen and lyophilised to give a white solid.

MS 460 [M+H]⁺

- 126 -

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.95 min.

Example F1a: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-methyl-amide maleate

The title compound is prepared analogously as described in example D2a using cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-methyl-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example F1b: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-methyl-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-methyl-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example F2: Cyclopropanesulfonic acid ((S)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-hydroxymethyl-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example F1 using N-Cbz-serine(tBu)-OH instead of N-methyl-N-Cbz-glycine.

MS: 476 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.88 min.

Example F3: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-N-methyl-acetamide

The title compound is prepared analogously as described in example E4 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-[3-exo-8-(2-methylamino-acetyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-[3-exo-8-(2-amino-acetyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 398 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.86 min.

Example F3a: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-N-methyl-acetamide maleate

The title compound is prepared analogously as described in example D2a using N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-N-methyl-acetamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example F3b: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-N-methyl-acetamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-N-methyl-acetamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example F4: Cyclopropanecarboxylic acid ((S)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-hydroxymethyl-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example F1 using cyclopropane carbonyl chloride instead of cyclopropane sulfonyl chloride.

MS: 424 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 2.74 min.

Example F5: Cyclopropanecarboxylic acid ((S)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example F4 using N-Cbz-(S)-alanine instead of N-methyl-N-Cbz-glycine.

MS: 460 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 2.80 min.

Example F5a: Cyclopropanecarboxylic acid ((S)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-amide maleate

The title compound is prepared analogously as described in example D2a using cyclopropanecarboxylic acid ((S)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example F5b: Cyclopropanecarboxylic acid ((S)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using cyclopropanecarboxylic acid ((S)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example F6: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-4-fluoro-N-methyl-benzenesulfonamide

The title compound is prepared analogously as described in example F1 using 4-fluorophenylsulfonyl chloride instead of cyclopropane sulfonyl chloride.

MS: 514 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 3.13 min.

Example F7: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1,1-dimethyl-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example F4 using N-Cbz-methyl-alanine instead of N-methyl-N-Cbz-glycine.

MS: 474 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.93 min.

Example F7a: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1,1-dimethyl-2-oxo-ethyl)-amide maleate

The title compound is prepared analogously as described in example D2a using cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1,1-dimethyl-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example F7b: Cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1,1-dimethyl-2-oxo-ethyl)-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1,1-dimethyl-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example F8: N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-N-methyl-acetamide

The title compound is prepared analogously as described in example F3 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-[3-exo-8-(3-methylamino-propionyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-

- 130 -

1-[3-exo-8-(2-methylamino-acetyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 412 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.09 min.

Example F9: Cyclopropanecarboxylic acid ((R)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-hydroxymethyl-2-oxo-ethyl)-amide

A) (R)-2-Benzyloxycarbonylamino-3-(tert-butyl-dimethyl-silanyloxy)-propionic acid methyl ester

To (R)-2-Benzyloxycarbonylamino-3-hydroxy-propionic acid methyl ester (1g, 4 mmol) in dimethyl formamide are added triethylamine (1.2 mL, 8 mmol), tert-butyldimethylsilyl chloride (895 mg, 6 mmol) and 4-dimethylaminopyridine (50 mg, 0.4 mmol). After stirring at rt during 16h, the reaction is quenched with water and aqueous 1N HCl and extracted with ethyl acetate. The organic phase is dried and evaporated to give a residue, which is purified by flash chromatography (Silica gel, ethyl acetate / cyclohexane 1/9 to 1/4) to yield a light yellow gum.

MS: 368 [M+H]⁺

HPLC (waters symmetry C18, 6min method (0-3.5min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 4.63 min.

B) (R)-2-Benzyloxycarbonylamino-3-(tert-butyl-dimethyl-silanyloxy)-propionic acid

To (R)-2-Benzyloxycarbonylamino-3-(tert-butyl-dimethyl-silanyloxy)-propionic acid methyl ester (1.450 g, 3.95 mmol) in tetrahydrofuran / water (2/1) is added lithium hydroxide (250 mg, 5.92 mmol). After stirred at rt during 16h, the solution is treated with ethyl acetate and the pH is decrease to 2 with aqueous 1N HCl. The organic phase is dried and evaporated to yield the title compound.

MS: 354 [M+H]⁺

HPLC (waters symmetry C18, 6min method (0-3.5min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 4.25 min.

C) Cyclopropanecarboxylic acid ((R)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-hydroxymethyl-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example F2 using (R)-2-benzyloxycarbonylamino-3-(tert-butyl-dimethyl-silanyloxy)-propionic acid using instead of N-Cbz-serine(tBu)-OH and cyclopropylcarboxylic acid instead of cyclopropylsulfonyl chloride.

MS: 440 [M+H]⁺

HPLC (waters symmetry C18, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 5%ACN): 2.65 min.

Example F10: N-((S)-2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-1-methyl-2-oxo-ethyl)-isobutyramide

The title compound is prepared analogously as described in example F5 using isobutyroyl chloride instead of cyclopropane carbonyl chloride.

MS: 426 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.239 min.

Example F11: N-((R)-2-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-1-methyl-2-oxo-ethyl)-isobutyramide

The title compound is prepared analogously as described in example F10 using N-Cbz-(R)-alanine instead of N-Cbz-(S)-alanine.

MS: 426 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.236 min.

Example F12: Cyclopropanecarboxylic acid ((R)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-amide

The title compound is prepared analogously as described in example F5 using N-Cbz-(R)-alanine instead of N-Cbz-(S)-alanine.

MS: 424 [M+H]⁺

- 132 -

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.224 min.

Example F12a: Cyclopropanecarboxylic acid ((R)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-amide maleate

The title compound is prepared analogously as described in example D2a using cyclopropanecarboxylic acid ((R)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.


Example F12b: Cyclopropanecarboxylic acid ((R)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using cyclopropanecarboxylic acid ((R)-2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example G1: N-[(S)-2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-acetamide

This compound is prepared according to Scheme G:

- 133 -



Scheme G

A) N-((S)-1-Methyl-2-{3-*exo*-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-acetamide

The title compound is prepared analogously as described in example E1 using N-acetyl-L-alanine instead of N-phtaloylglycine.

MS: 502 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.11 min.

B) N-((S)-2-{3-*exo*-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-acetamide

To N-((S)-1-Methyl-2-{3-*exo*-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-acetamide (60 mg, 0.120 mmol) in dioxane (1 mL) is added 4N HCl in dioxane (1 mL). The resulting mixture is stirred at rt during 1h. The solution is frozen and lyophilised to give a white solid before purification by preparative HPLC (Column Waters C18 ODB 5µm 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield the title compound.

MS: 398 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.91 min.

Example G1a: N-((S)-2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-acetamide maleate

The title compound is prepared analogously as described in example D2a using N-((S)-2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-acetamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G1b: N-((S)-2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-acetamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-((S)-2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-acetamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G2: N-((R)-2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-methyl-2-oxo-ethyl)-acetamide

The title compound is prepared analogously as described in example G1 using N-acetyl-D-alanine instead of N-acetyl-L-alanine.

MS: 398 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.88 min.

Example G3: N-(2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-benzamide

The title compound is prepared analogously as described in example G1 using N-benzoylglycine instead of N-acetyl-L-alanine.

MS: 446 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.94 min.

Example G4: (R)-1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-hydroxy-2-phenyl-propan-1-one

The title compound is prepared analogously as described in example G1 using (R)-2-hydroxy-2-phenyl-propionic acid instead of N-acetyl-L-alanine.

MS: 433 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.00 min.

Example G5: N-((S)-5-Acetylamino-1-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl)-pentyl)-acetamide

The title compound is prepared analogously as described in example G1 using (S)-2,6-bis-acetylamino-hexanoic acid instead of N-acetyl-L-alanine.

MS: 497 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.10 min.

Example G6: ((S)-2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-1-phenyl-ethyl)-carbamic acid ethyl ester

The title compound is prepared analogously as described in example G1 using (S)-ethoxycarbonylamino-phenyl-acetic acid instead of N-acetyl-L-alanine.

MS: 490 [M+H]⁺

HPLC (waters symmetry C18, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 5%ACN): 3.14 min.

Example G7: ((R)-2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-1-phenyl-ethyl)-carbamic acid ethyl ester

The title compound is prepared analogously as described in example G1 using (R)-ethoxycarbonylamino-phenyl-acetic acid instead of N-acetyl-L-alanine.

MS: 490 [M+H]⁺

HPLC (waters symmetry C18, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 5%ACN): 3.14 min.

Example G8: (S)-1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-methoxy-2-phenyl-ethanone

The title compound is prepared analogously as described in example G1 using (S)-methoxyphenyl-acetic acid instead of N-acetyl-L-alanine.

MS: 433 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 2.49 min.

Example G9: (R)-1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-methoxy-2-phenyl-ethanone

The title compound is prepared analogously as described in example G1 using (R)-methoxyphenyl-acetic acid instead of N-acetyl-L-alanine.

MS: 433 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 2.47 min.

Example G10: N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-benzenesulfonamide

The title compound is prepared analogously as described in example G1 using 3-Benzenesulfonylamino-propionic acid instead of N-acetyl-L-alanine.

MS: 496 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.17 min.

Example G10a: N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-benzenesulfonamide maleate

The title compound is prepared analogously as described in example D2a using N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-benzenesulfonamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example G10b: N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-benzenesulfonamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-benzenesulfonamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example G11: 2,3-Dihydro-benzo[1,4]dioxine-6-sulfonic acid (3-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-amide

The title compound is prepared analogously as described in example G1 using 3-(2,3-dihydro-benzo[1,4]dioxine-6-sulfonylamino)-propionic acid instead of N-acetyl-L-alanine.

MS: 554 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.18 min.

Example G12: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-(2,3-dihydro-benzo[1,4]dioxine-6-sulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G1 using 3-(2,3-dihydro-benzo[1,4]dioxine-6-sulfonyl)-propionic acid instead of N-acetyl-L-alanine.

MS: 539 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.97 min.

Example G13: N-[4-{3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propane-1-sulfonyl}-phenyl]-acetamide

The title compound is prepared analogously as described in example G1 using 3-(4-acetyl-amino-benzenesulfonyl)-propionic acid instead of N-acetyl-L-alanine.

MS: 574 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.12 min.

Example G14: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-(4-fluoro-benzenesulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G1 using 3-(4-fluoro-benzenesulfonyl)-propionic acid instead of N-acetyl-L-alanine.

MS: 499 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.28 min.

Example G15: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-(4-thiophen-2-yl-6-trifluoromethyl-pyrimidine-2-sulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G1 using 3-(4-thiophen-2-yl-6-trifluoromethyl-pyrimidine-2-sulfonyl)-propionic acid instead of N-acetyl-L-alanine.

MS: 663 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.21 min.

Example G16: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-phenylmethanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example G1 using 3-phenylmethanesulfonyl-propionic acid instead of N-acetyl-L-alanine.

MS: 495 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.18 min.

Example G17: 2-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-benzyl)-isoindole-1,3-dione

The title compound is prepared analogously as described in example G1 using 3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl-methyl)-benzoic acid instead of N-acetyl-L-alanine.

- 139 -

MS: 548 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.30 min.

Example G18: N-{3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-benzyl)-benzamide

The title compound is prepared analogously as described in example E21 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-{3-exo-8-[3-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-benzoyl]-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-{3-exo-8-[3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionyl]-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 522 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.28 min.

Example G19: 3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-N-benzyl-3-oxo-propionamide

The title compound is prepared analogously as described in example G1 using N-benzyl-malonamic acid instead of N-acetyl-L-alanine.

MS: 462 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.25 min.

Example G20: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-morpholin-4-yl-propane-1,3-dione

The title compound is prepared analogously as described in example G1 using N-morpholin-malonamic acid instead of N-acetyl-L-alanine.

MS: 440 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.16 min.

- 140 -

Example G21: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-phenylmethanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example G16 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane.

MS: 469 [M+H]⁺

HPLC (YMC Pack ODS-AQ 3 µm 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.98 min.

Example G21a: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-phenylmethanesulfonyl-propan-1-one maleate

The title compound is prepared analogously as described in example D2a using 1-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-phenylmethanesulfonyl-propan-1-one instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G21b: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-phenylmethanesulfonyl-propan-1-one toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 1-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-phenylmethanesulfonyl-propan-1-one instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G22: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(4-fluoro-benzenesulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G14 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane.

MS: 473 [M+H]⁺

- 141 -

HPLC (YMC Pack ODS-AQ 3 μ m 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.78 min.

Example G23: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-benzenesulfonyl-propan-1-one

The title compound is prepared analogously as described in example G1 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane and 3-benzenesulfonyl-propionic acid instead of N-acetyl-L-alanine.

MS: 455 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.13 min.

Example G24: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(2,3-dihydro-benzo[1,4]dioxine-6-sulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G12 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane.

MS: 513 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.15 min.

Example G25: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-methanesulfonyl-ethanone

The title compound is prepared analogously as described in example G22 using methanesulfonyl-acetic acid instead of 3-(4-fluoro-benzenesulfonyl)-propionic acid.

MS: 401 [M+H]⁺

HPLC (YMC Pack ODS-AQ 3 μ m 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.48 min.

- 142 -

Example G26: 3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-N-methyl-benzamide

The title compound is prepared analogously as described in example G1 using N-methyl-isophthalamic acid instead of N-acetyl-L-alanine.

MS: 446 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.17 min.

Example G26a: 3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-N-methyl-benzamide maleate

The title compound is prepared analogously as described in example D2a using 3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-N-methyl-benzamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G26b: 3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-N-methyl-benzamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-N-methyl-benzamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G27: {3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-[3-(morpholine-4-carbonyl)-phenyl]-methanone

The title compound is prepared analogously as described in example G1 using 3-(morpholine-4-carbonyl)-benzoic acid instead of N-acetyl-L-alanine.

MS: 502 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.19 min.

- 143 -

Example G28: 1-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-benzyl)-pyrrolidin-2-one

The title compound is prepared analogously as described in example G1 using 3-(2-oxo-pyrrolidin-1-ylmethyl)-benzoic acid instead of N-acetyl-L-alanine.

MS: 486 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.23 min.

Example G29: 3-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-benzyl)-thiazolidine-2,4-dione

The title compound is prepared analogously as described in example G1 using 3-(2,4-dioxo-thiazolidin-3-ylmethyl)-benzoic acid instead of N-acetyl-L-alanine.

MS: 518 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.25 min.

Example G30: 1-(4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl)-3-(propane-2-sulfonyl)-propan-1-one

A) 3-Isopropylsulfonyl-propionic acid benzyl ester

To benzylacrylate (100 mg, 0.617 mmol) in ethanol (2 mL) are added triethylamine (95 μ L, 0.679 mmol) and 2-propanethiol (58 μ L, 0.617 mmol). After stirring at rt during 2h, the solvent is evaporated to give a residue which is treated with dichloromethane and water. The organic phase is dried and evaporated to afford a colorless oil before purification by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield the title compound.

MS: 239 [M+H]⁺

HPLC (Waters Symmetry C18 3.5 μ m 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 3.97 min.

B) 3-(Propane-2-sulfonyl)-propionic acid benzyl ester

- 144 -

To 3-isopropylsulfonyl-propionic acid benzyl ester (46 mg, 0.193 mmol) in acetic acid (500 μ L) is added an aqueous 30% H₂O₂ solution (82 μ L). After stirring at 80°C during 2h, the mixture is quenched with saturated NaHCO₃ solution and extracted with dichloromethane. The organic phase is dried and evaporated to give the title compound.

MS: 271 [M+H]⁺

HPLC (Waters Symmetry C18 3.5 μ m 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 2.92 min.

C) 3-(Propane-2-sulfonyl)-propionic acid

To 3-(propane-2-sulfonyl)-propionic acid benzyl ester (49.3 mg, 0.182 mmol) in methanol (1 mL) is added Pd/C (5 mg) and the mixture is stirred at rt during 16h under H₂ atmosphere. The resulting suspension is filtered through celite and the filtrate is evaporated to yield the title compound.

D) 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(propane-2-sulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G24 using 3-(propane-2-sulfonyl)-propionic acid instead of 3-(2,3-dihydro-benzo[1,4]dioxine-6-sulfonylamino)-propionic acid.

MS: 421 [M+H]⁺

HPLC (YMC Pack ODS-AQ 3 μ m 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.6 min.

Example G31: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-(4-trifluoromethyl-pyrimidine-2-sulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G30 using 4-trifluoromethyl-pyrimidine-2-thiol instead of 2-propanethiol and 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 551 [M+H]⁺

- 145 -

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.25 min.

Example G32: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-benzenesulfonyl-propan-1-one

The title compound is prepared analogously as described in example G1 using 3-benzenesulfonyl-propionic acid instead of N-acetyl-L-alanine.

MS: 481 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.2 min.

Example G33: 1-{4-[1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-(propane-2-sulfonyl)-ethanone

A) Isopropylsulfanyl-acetic acid benzyl ester

To benzyl bromoacetate (100 mg, 0.437 mmol) in dimethyl formamide (2 mL) is added triethylamine (67 μ L, 0.481 mmol) and 2-propanethiol (41 μ L, 0.437 mmol). After stirring at rt during 2 h, the solvent is evaporated, the residue is treated with dichloromethane and water. The organic phase is dried and evaporated to give a crude compound before purification by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield the title compound.

MS: 225 [M+H]⁺

HPLC (Waters Symmetry C18 3.5 μ m 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 3.84 min.

B) 1-{4-[1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-(propane-2-sulfonyl)-ethanone

The title compound is prepared analogously as described in example G30 using isopropylsulfanyl-acetic acid benzyl ester instead of 3-isopropylsulfanyl-propionic acid benzyl ester.

MS: 551 [M+H]⁺

- 146 -

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.25 min.

Example G34: Cyclopropanecarboxylic acid ((S)-1-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-3-carbamoyl-propyl)-amide

A) (S)-4-Carbamoyl-2-(cyclopropanecarbonyl-amino)-butyric acid

To (S)-2-Amino-4-carbamoyl-butyric acid (100 mg, 0.684 mmol) in water (16 mL) are added Na₂CO₃ (218 mg, 2.052 mmol) and a solution of cyclopropylcarboxylic acid (62.1 μ L, 0.684 mmol) in tetrahydrofuran (8 mL). After stirring at rt during 2h, ethyl acetate is added and the pH is decreased to 3 by addition of aqueous 1N HCl. The aqueous phase is evaporated, the residue is mixed with methanol and filtrated. The filtrate is evaporated to yield the title compound.

MS: 237 [M+H]⁺

HPLC (Waters Symmetry C18 3.5 μ m 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 0.504 min.

B) Cyclopropanecarboxylic acid ((S)-1-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-3-carbamoyl-propyl)-amide

The title compound is prepared analogously as described in example G1 using (S)-4-carbamoyl-2-(cyclopropanecarbonyl-amino)-butyric acid instead of N-acetyl-L-alanine.

MS: 482 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.13 min.

Example G35: N-[4-{3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propane-1-sulfonyl}-phenyl]-acetamide

The title compound is prepared analogously as described in example G21 using 3-(4-acetylaminobenzenesulfonyl)-propionic acid instead of 3-phenylmethanesulfonyl-propionic acid.

MS: 512 [M+H]⁺

- 147 -

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.19 min.

Example G35a: N-[4-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propane-1-sulfonyl)-phenyl]-acetamide maleate

The title compound is prepared analogously as described in example D2a using N-[4-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propane-1-sulfonyl)-phenyl]-acetamide instead of cyclopropanesulfonic acid (2-{3-*exo*-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G35b: N-[4-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propane-1-sulfonyl)-phenyl]-acetamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-[4-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propane-1-sulfonyl)-phenyl]-acetamide instead of cyclopropanesulfonic acid (2-{3-*exo*-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G36: 1-[4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl]-3-(tetrahydro-furan-2-ylmethanesulfonyl)-propan-1-one

A) 3-(2-Benzyloxycarbonyl-ethyl)disulfanyl-propionic acid benzyl ester

To 3,3'-dithiodipropionic acid (1 g, 4.75 mmol) in DCM (6 mL) are added DIPEA (2.86 mL, 16.64 mmol), DMAP (76 mg, 0.618 mmol) and benzyl bromide (1.42 mL, 11.89 mmol). After stirring at rt during 16h, the mixture is washed with water, brine, an aqueous 1N HCl solution and an aqueous 10% NaHCO₃ solution. The organic phase is dried and evaporated to afford a orange oil before purification by flash chromatography on silica (cyclohexane / ethyl acetate 1/0 to 8/2) to yield a yellow oil.

MS: 391 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 4.206 min.

- 148 -

B) 3-Mercapto-propionic acid benzyl ester

A stirred solution of 3-(2-Benzoyloxycarbonyl-ethyl-disulfanyl)-propionic acid benzyl ester (1.73 g, 4.353 mmol) in THF (15 mL) and water (1.5 mL) is deoxygenated during 15 min using a stream of N₂. After addition of tributylphosphine (2.15 mL, 8.706 mmol), the mixture is stirred at rt during 6h and the solvent is evaporated. The aqueous phase is treated with DCM and an aqueous 1N HCl solution, the aqueous phase is extracted with DCM and the combined organic phases are dried and evaporated to afford a yellow oil before purification by flash chromatography on silica (cyclohexane / ethyl acetate 1/0 to 8/2) to yield a yellow oil.

MS: 219 [M+Na]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 3.365 min.

C) 3-(Tetrahydro-furan-2-ylmethylsulfanyl)-propionic acid benzyl ester

To 3-Mercapto-propionic acid benzyl ester (200 mg, 1.019 mmol) are added tetrahydrofurfuryl bromide (505 mg, 3.057 mmol) and sodium methoxide (55 mg, 1.019 mmol). After stirring at rt during 2h, the mixture is quenched with ethyl acetate and water, the organic phase is washed with water and brine, dried and evaporated to afford a yellow oil before purification by preparative HPLC (Column interchim C18 ODB 5µm 19x50, Gradient: 0-5 min 20% ACN, 5-15 min 20-100% ACN, 15-20 min 100% ACN) to yield a colorless oil.

MS: 281 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 3.522 min.

D) 3-(Tetrahydro-furan-2-ylmethanesulfonyl)-propionic acid

The title compound is prepared analogously as described in example G30 using 3-(Tetrahydro-furan-2-ylmethylsulfonyl)-propionic acid benzyl ester instead of 3-isopropylsulfonyl-propionic acid benzyl ester.

MS: 245 [M+Na]⁺

E) 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(tetrahydro-furan-2-ylmethanesulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G24 using 3-(Tetrahydro-furan-2-ylmethanesulfonyl)-propionic acid instead of 3-(2,3-dihydrobenzo[1,4]dioxine-6-sulfonylamino)-propionic acid.

MS: 463 [M+H]⁺

HPLC (YMC Pack ODS-AQ 3 µm 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.469 min.

Example G37: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-cyclopentanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example G30 using cyclopentyl mercaptan instead of 2-propanethiol.

MS: 447 [M+H]⁺

HPLC (YMC Pack ODS-AQ 3 µm 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.641 min.

Example G38: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-ethanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example G30 using ethanthiol instead of 2-propanethiol.

MS: 407 [M+H]⁺

HPLC (YMC Pack ODS-AQ 3 µm 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.43 min.

Example G39: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(2-methyl-propane-2-sulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G30 using tertbutylthiol instead of 2-propanethiol.

MS: 435 [M+H]⁺

HPLC (YMC Pack ODS-AQ 3 µm 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.63 min.

Example G40: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(tetrahydro-pyran-2-ylmethanesulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G36 using 2-(Bromomethyl)tetrahydro-2H-pyran instead of tetrahydrofurfuryl bromide.

MS: 477 [M+H]⁺

HPLC (YMC Pack ODS-AQ 3 µm 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.585 min.

Example G41: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(2-methoxy-ethanesulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G36 using 2-bromoethyl-methyl ether instead of tetrahydrofurfuryl bromide.

MS: 437 [M+H]⁺

HPLC (YMC Pack ODS-AQ 3 µm 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.295 min.

Example G41a: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(2-methoxy-ethanesulfonyl)-propan-1-one maleate

The title compound is prepared analogously as described in example D2a using 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(2-methoxy-ethanesulfonyl)-propan-1-one instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G41b: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(2-methoxy-ethanesulfonyl)-propan-1-one toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(2-methoxy-ethanesulfonyl)-propan-1-one instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G42: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-(2-methyl-propane-2-sulfonyl)-ethanone

The title compound is prepared analogously as described in example G33 using tertbutylthiol instead of 2-propanethiol.

MS: 421 [M+H]⁺

HPLC (YMC Pack ODS-AQ 3 µm 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.56 min.

Example G43: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-cyclopentanesulfonyl-ethanone

The title compound is prepared analogously as described in example G33 using cyclopentyl mercaptan instead of 2-propanethiol.

MS: 433 [M+H]⁺

HPLC (YMC Pack ODS-AQ 3 µm 2.1 x 50mm, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 5%ACN): 2.68 min.

Example G44: 1-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]oct-8-yl]-2-(2-methyl-propane-2-sulfonyl)-ethanone

The title compound is prepared analogously as described in example G42 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 447 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.659 min.

Example G45: 1-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]oct-8-yl]-2-cyclopentanesulfonyl-ethanone

The title compound is prepared analogously as described in example G43 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

- 152 -

MS: 459 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.755 min.

Example G46: 1-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-yl)-2-(propane-2-sulfonyl)-ethanone

The title compound is prepared analogously as described in example G33 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 433 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.601 min.

Example G47: N-[4-(3-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-yl)-3-oxo-propane-1-sulfonyl)-phenyl]-acetamide

The title compound is prepared analogously as described in example G35 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 574 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.12 min.

Example G48: 1-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-yl)-3-(3-methoxy-benzenesulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G44 using 3-methoxythiophenol instead of tertbutylthiol.

MS: 511 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.26 min.

Example G49: 1-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-azabicyclo[3.2.1]oct-8-yl)-3-(toluene-3-sulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G44 using 3-thiocresol instead of tertbutylthiol.

MS: 495 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.27 min.

Example G50: 1-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-azabicyclo[3.2.1]oct-8-yl)-3-(pyrimidine-2-sulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G44 using Pyrimidine-2-thiol instead of tertbutylthiol.

MS: 483 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.17 min.

Example G51: 1-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-azabicyclo[3.2.1]oct-8-yl)-3-(6-methyl-pyridine-2-sulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G44 using 6-Methyl-pyridine-2-thiol instead of tertbutylthiol.

MS: 498 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.24 min.

Example G52: 7-(3-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-azabicyclo[3.2.1]oct-8-yl)-3-oxo-propane-1-sulfonyl)-3,4-dihydro-2H-isoquinolin-1-one

The title compound is prepared analogously as described in example G44 using 7-Mercapto-3,4-dihydro-2H-isoquinolin-1-one instead of tertbutylthiol.

MS: 550 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.20 min.

Example G53: 3-(3-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-oxo-propane-1-sulfonyl)-benzamide

The title compound is prepared analogously as described in example G44 using 3-Mercapto-benzamide instead of tertbutylthiol.

MS: 525 [M+H]⁺

Example G53a: 3-(3-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-oxo-propane-1-sulfonyl)-benzamide maleate

The title compound is prepared analogously as described in example D2a using 3-(3-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-oxo-propane-1-sulfonyl)-benzamide instead of cyclopropanesulfonic acid (2-(3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example G53b: 3-(3-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-oxo-propane-1-sulfonyl)-benzamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 3-(3-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-oxo-propane-1-sulfonyl)-benzamide instead of cyclopropanesulfonic acid (2-(3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example G54: 1-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-cyclopentanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example G37 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 473 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.731 min.

Example G55: 1-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-yl)-3-(tetrahydro-furan-2-ylmethanesulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G36 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 489 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.565 min.

Example G56: 1-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-yl)-3-(2-methyl-propane-2-sulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G39 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 461 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.700 min.

Example G57: 1-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-yl)-3-ethanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example G38 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 433 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.572 min.

Example G58: 1-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-yl)-3-(tetrahydro-pyran-2-ylmethanesulfonyl)-propan-1-one

- 156 -

The title compound is prepared analogously as described in example G40 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 503 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.666 min.

Example G59: 1-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]oct-8-yl)-3-(2-methoxy-ethanesulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G41 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 463 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.435 min.

Example G60: 3-(3-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]octane-8-carbonyl)-benzyl)-1-methyl-imidazolidine-2,4-dione

A) 3-(3-Methyl-2,5-dioxo-imidazolidin-1-ylmethyl)-benzoic acid methyl ester

To methylhydantoine (50 mg, 0.438 mmol) in DMF (2 mL) is added at 0°C sodium hydride (23 mg, 0.525 mmol). After stirring at 0°C during 1h, methyl 2-bromomethylphenylcarboxylate (121 mg, 0.525 mmol) is added and the mixture is stirred at rt during 2h before quenching with an aqueous saturated NaHCO₃ solution and DCM. The organic phase is dried and evaporated to give a crude compound before purification by preparative HPLC (Column Waters C18 ODB 5µm 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield the title compound.

MS: 285 [M+Na]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.216 min.

- 157 -

B) 3-(3-Methyl-2,5-dioxo-imidazolidin-1-ylmethyl)-benzoic acid

To 3-(3-Methyl-2,5-dioxo-imidazolidin-1-ylmethyl)-benzoic acid methyl ester (100 mg, 0.381 mmol) in THF/water (2/1, 2 mL) is added lithium hydroxyde monohydrate (24 mg, 0.572 mmol) and the reaction is stirred at rt during 16h before quenching with water and ethylacetate. The separated aqueous phase is acidified to pH 2 with an aqueous 1N HCl solution, extracted with ethylacetate and the organic phase is dried and evaporated to give the title compound.

MS: 267 [M+H₂O+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.25 min.

C) 3-(3-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]octane-8-carbonyl)-benzyl)-1-methyl-imidazolidine-2,4-dione

The title compound is prepared analogously as described in example G17 using 3-(3-Methyl-2,5-dioxo-imidazolidin-1-ylmethyl)-benzoic acid instead of 3-(1,3-dioxo-1,3-dihydro-isindol-2-yl-methyl)-benzoic acid.

MS: 515 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 5%ACN): 2.91 min.

Example G61: 3-(4-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]octane-8-carbonyl)-benzyl)-thiazolidine-2,4-dione

The title compound is prepared analogously as described in example G1 using 4-(2,4-Dioxo-thiazolidin-3-ylmethyl)-benzoic acid instead of N-acetyl-L-alanine.

MS: 518 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.245 min.

Example G62: 1-(4-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]octane-8-carbonyl)-benzyl)-pyrrolidine-2,5-dione

- 158 -

A) 2-Methyl-propane-2-sulfinic acid [(R)-1-[(1S,3S,5R)-8-[3-(2,5-dioxo-pyrrolidin-1-ylmethyl)-benzoyl]-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide

To 2-Methyl-propane-2-sulfinic acid [(R)-1-[(1S,3S,5R)-8-(3-aminomethyl-benzoyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide (228 mg, 0.437 mmol) in toluene (10 ml) are added succinic anhydride (53 mg, 0.524 mmol) and molecular sieves (200 mg). After stirring at 110°C during 6h, CDI (107 mg, 0.655 mmol) and triethylamine (183 μ L, 1.311 mmol) are added and the mixture is stirred at 110°C during 24h. After filtration and evaporation of the solvent, the residue is treated with DCM and an aqueous saturated NaHCO₃ solution, the organic phase is dried and evaporated to give a crude compound before purification by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield the title compound.

MS: 604 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.370 min.

B) 1-{4-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl]-benzyl}-pyrrolidine-2,5-dione

The title compound is prepared analogously as described in example G1 using (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-[(1S,3S,5R)-8-[3-(2,5-dioxo-pyrrolidin-1-ylmethyl)-benzoyl]-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of N-[(S)-1-Methyl-2-{3-exo-[(R)-1-[(S)-2-methyl-propane-2-sulfinylamino]-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethyl)-acetamide.

MS: 500 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.212 min.

Example G63: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-[2-(1,1-dioxo-1 λ 6*-thiomorpholin-4-yl)-2-oxo-ethanesulfonyl]-propan-1-one

A) 3-Carboxymethylsulfonyl-propionic acid benzyl ester

- 159 -

The title compound is prepared analogously as described in example G36 using Bromoacetic acid instead of tetrahydrofurfuryl bromide.

MS: 255 [M+H]⁺

HPLC (Waters Symmetry C18 3.5μm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 2.841 min.

B) 3-(2-Oxo-2-thiomorpholin-4-yl-ethylsulfanyl)-propionic acid benzyl ester

To 3-Carboxymethylsulfanyl-propionic acid benzyl ester (763 mg, 3 mmol), HBTU (1.71 g, 4.5 mmol) and DIPEA (2.05 mL, 12 mmol) in DCM (10 mL) is added thiomorpholine (283 μL, 3 mmol). After stirring at rt during 2h and evaporation of the solvent, the residue is dissolved with ethylacetate and washed with an aqueous 1N HCl solution and an aqueous saturated NaHCO₃ solution. The organic phase is dried and evaporated to afford the title compound.

MS: 340 [M+H]⁺

HPLC (Waters Symmetry C18 3.5μm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 3.529 min.

C) 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-[2-(1,1-dioxo-1λ⁶-thiomorpholin-4-yl)-2-oxo-ethanesulfonyl]-propan-1-one

The title compound is prepared analogously as described in example G36 using 3-(2-Oxo-2-thiomorpholin-4-yl-ethylsulfanyl)-propionic acid benzyl ester instead of 3-(Tetrahydro-furan-2-ylmethylsulfonyl)-propionic acid benzyl ester.

MS: 554 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.214 min.

Example G63a: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-[2-(1,1-dioxo-1λ⁶-thiomorpholin-4-yl)-2-oxo-ethanesulfonyl]-propan-1-one maleate

The title compound is prepared analogously as described in example D2a using 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-[2-(1,1-dioxo-1λ⁶-thiomorpholin-4-yl)-2-oxo-ethanesulfonyl]-propan-1-one instead of cyclopropanesulfonic acid

(2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G63b: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-[2-(1,1-dioxo-1 λ 6*-thiomorpholin-4-yl)-2-oxo-ethanesulfonyl]-propan-1-one toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-[2-(1,1-dioxo-1 λ 6*-thiomorpholin-4-yl)-2-oxo-ethanesulfonyl]-propan-1-one instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example G64: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(2-morpholin-4-yl-2-oxo-ethanesulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G63 using morpholine instead of thiomorpholine.

MS: 506 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.296 min.

Example G65: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-(tetrahydro-pyran-2-ylmethanesulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G58 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane.

MS: 477 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.372 min.

Example G66: 1-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]oct-8-yl)-3-[2-(1,1-dioxo-1 λ 6*-thiomorpholin-4-yl)-2-oxo-ethanesulfonyl]-propan-1-one

The title compound is prepared analogously as described in example G63 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 580 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.352 min.

Example G67: 1-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]oct-8-yl)-3-(2-morpholin-4-yl-2-oxo-ethanesulfonyl)-propan-1-one

The title compound is prepared analogously as described in example G64 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.


MS: 532 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.379 min.

Example H1: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]oct-8-yl}-2-(morpholine-4-sulfonyl)-ethanone

This compound is prepared according to Scheme H:

- 162 -



Scheme H

A) Chlorosulfonyl-acetic acid methyl ester

To chlorosulfonyl chloride (3.34 g, 17.9 mmol) in diethyl ether (30 mL) is added at 0°C methanol (800 μ L, 19.7 mmol). The resulting mixture is stirred at 0°C during 1h and the solvent is evaporated to give the title compound.

B) (Morpholine-4-sulfonyl)-acetic acid methyl ester

To chlorosulfonyl-acetic acid methyl ester (3.39 g, 19.6 mmol) in dichloromethane (50 mL) is added morpholine (8.6 mL, 98 mmol). The resulting mixture is stirred at rt during 2h and the solvent is evaporated to yield the title compound.

MS: 223 [M-H]⁺

C) (Morpholine-4-sulfonyl)-acetic acid

(Morpholine-4-sulfonyl)-acetic acid methyl ester (0.5 g, 2.24 mmol) is dissolved in 0.66N KOH aqueous ethanol solution (1/1, 150 mL) and the mixture is heated at reflux during 16h. The solvent is evaporated and residue is extracted with dichloromethane/methanol 4:1 to yield the title compound.

MS: 232 [M+Na]

D) 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-(morpholine-4-sulfonyl)-ethanone

The title compound is prepared analogously as described in example G1 using (morpholine-4-sulfonyl)-acetic acid instead of N-acetyl-L-alanine.

MS: 476 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.94 min.

Example H1a: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-(morpholine-4-sulfonyl)-ethanone maleate

The title compound is prepared analogously as described in example D2a using 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-(morpholine-4-sulfonyl)-ethanone instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example H1b: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-(morpholine-4-sulfonyl)-ethanone toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-(morpholine-4-sulfonyl)-ethanone instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example H2: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-benzenesulfonyl-ethanone

The title compound is prepared analogously as described in example H1 using phenylsulfonylacetic acid instead of (Morpholine-4-sulfonyl)-acetic acid.

MS: 466 [M+H]⁺

HPLC (Nucleosil 100-5 C18, 10min method (0-1min 10%ACN, 1-6min 10-100%ACN, 6-8.5min 100%ACN, 8.5-9min 100-10%ACN, 9-10min 10%ACN): 3.88 min.

Example H3: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-methanesulfonyl-ethanone

The title compound is prepared analogously as described in example H1 using methanesulfonylacetic acid instead of (Morpholine-4-sulfonyl)-acetic acid.

MS: 405 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 0.87 min.

Example H4: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-(piperidine-1-sulfonyl)-ethanone

The title compound is prepared analogously as described in example H1 using piperidine instead of morpholine.

MS: 474 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.28 min.

Example H5: 2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethanesulfonic acid benzylamide

The title compound is prepared analogously as described in example H1 using benzylamine instead of morpholine.

MS: 496 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.28 min.

Example H6: 2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethanesulfonic acid benzyl-methyl-amide

The title compound is prepared analogously as described in example H1 using N-methylbenzylamine instead of morpholine.

MS: 510 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.30 min.

Example H7: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-methyl-2-(morpholine-4-sulfonyl)-propan-1-one

A) 2-Methyl-2-(morpholine-4-sulfonyl)-propionic acid methyl ester

To a solution of (Morpholine-4-sulfonyl)-acetic acid methyl ester (500 mg, 2.24 mmol) in tetrahydrofuran (10 mL) is added potassium bis(trimethylsilyl) amide (1.18g, 5.6 mmol). After stirring at rt during 1h, methyl iodide (349 μ L, 5.6 mmol) is added and the resulting mixture is stirred at rt during 1h and at 50°C during the weekend. The solution is purified by flash chromatography on silica gel (eluent: cyclohexane/ethyl acetate 100/0 to 0/100) to yield a yellow solid.

MS: 252 [M + H]⁺

B) 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-methyl-2-(morpholine-4-sulfonyl)-propan-1-one

The title compound is prepared analogously as described in example H1 using 2-methyl-2-(morpholine-4-sulfonyl)-propionic acid methyl ester instead of (Morpholine-4-sulfonyl)-acetic acid methyl ester.

MS: 504 [M + H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.26 min.

Example H8: 2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethanesulfonic acid (2-methoxy-ethyl)-methyl-amide

The title compound is prepared analogously as described in example H1 using N-methyl-2-methoxyethylamine instead of morpholine.

MS: 478 [M + H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.22 min.

Example H8a: 2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethanesulfonic acid (2-methoxy-ethyl)-methyl-amide maleate

The title compound is prepared analogously as described in example D2a using 2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethanesulfonic acid (2-methoxy-ethyl)-methyl-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example H8b: 2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethanesulfonic acid (2-methoxy-ethyl)-methyl-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-oxo-ethanesulfonic acid (2-methoxy-ethyl)-methyl-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example H9: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-(morpholine-4-sulfonyl)-ethanone

The title compound is prepared analogously as described in example H1 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane.

MS: 450 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 2.70 min.

Example H10: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-methyl-1-oxo-propane-2-sulfonic acid (2-methoxy-ethyl)-methyl-amide

- 167 -

The title compound is prepared analogously as described in example H7 using N-methyl-2-methoxyethylamine instead of morpholine.

MS: 506 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.29 min.

Example H11: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-methyl-1-oxo-propane-2-sulfonic acid benzyl-methyl-amide

The title compound is prepared analogously as described in example H7 using N-methyl-benzylamine instead of morpholine.

MS: 538 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.36 min.

Example H12: {3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-[1-(morpholine-4-sulfonyl)-cyclopropyl]-methanone

The title compound is prepared analogously as described in example H7 using 1,2-dibromoethane instead of methyl iodide.

MS: 502 [M+H]⁺

Example H13: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-(1,3-dihydro-isoindole-2-sulfonyl)-ethanone

The title compound is prepared analogously as described in example H1 using isoindoline instead of morpholine.

MS: 508 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.26 min.

Example H14: 1-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-2-(4,4-difluoro-piperidine-1-sulfonyl)-ethanone

- 168 -

The title compound is prepared analogously as described in example H1 using 4,4-Difluoropiperidine instead of morpholine.


MS: 510 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.23 min.

Example I1: N-(3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]oct-8-yl}-3-oxo-propyl)-3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionamide

This compound is prepared according to Scheme I :

- 169 -



Scheme I

A) (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide

The title compound is prepared analogously as described in example B1 using 4-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidine-1-carboxylic acid benzyl ester instead of 3-*exo*-[(S)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester.

MS: 363 [M+H]⁺

- 170 -

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 2.45 min.

B) (3-{4-[(R)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-carbamic acid benzyl ester

To a solution of Cbz-beta-alanine (339 mg, 1.52 mmol) in dichloromethane (10 mL) are added O-benzotriazol-1-yl-tetramethyluronium hexafluorophosphate (785 mg, 2.07 mmol) and diisopropylethylamine (960 µL, 5.52 mmol) before addition of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide (500 mg, 1.379 mmol). The resulting solution is stirred at rt during 2h and washed with water and brine. The organic phase is dried and evaporated before purification by preparative HPLC (Column Interchrom C18 ODB 10µm 28x250, Gradient: 0-2.5 min 5% ACN, 2.5-25.5 min 5-100% ACN, 25.5-30 min 100% ACN) to yield a yellow solid.

MS: 568 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.41 min.

C) (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-[1-(3-amino-propionyl)-piperidin-4-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide

To (3-{4-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-carbamic acid benzyl ester (684 mg, 1.205 mmol) in methanol (10 mL) are added ammonium formate (379 mg, 6.025 mmol) and Pd/C (171 mg). After stirring at rt during 72h, the solution is filtered through celite and evaporated to yield a yellow solid.

MS: 434 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.17 min.

D) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-(3-{4-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-propionamide

To 2-methyl-propane-2-sulfinic acid [(R)-1-[1-(3-amino-propionyl)-piperidin-4-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide (104 mg, 0.241 mmol) in dichloromethane (3 mL) are added

- 171 -

triethylamine (101 μ L, 0.723), N,N-dimethylamidodisulfonyl chloride (28.5 μ L, 0.265 mmol) and 4-dimethylaminopyridine (6 mg, 0.05 mmol). After stirring during 4h at rt, the mixture is washed with saturated NaHCO₃ solution, the organic phase is dried and evaporated to give a residue which is purified by preparative HPLC (Column Interchrom C18 ODB 5 μ m 19x50, Gradient: 0-5 min 10% ACN, 5-15 min 10-90% ACN, 15-20 min 90% ACN) to yield a yellow oil.

MS: [M+H]⁺ 541

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.309 min.

E) N-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionamide

To 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-(3-{4-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-propionamide (22.3 mg, 0.041 mmol) is added 4N HCl in dioxane (2 mL). The resulting mixture is stirred at rt during 1h before it is frozen, lyophilized and purified by preparative HPLC (Column YMC ODS-AQ 20x50 5 μ m, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% CAN) to yield a white solid.

MS 437 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.09 min.

Example I1a: N-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionamide maleate

The title compound is prepared analogously as described in example D2a using N-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example I1b: N-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using N-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-3-(1,3-dioxo-1,3-dihydro-isindol-2-yl)-propionamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example 12: Cyclopropanesulfonic acid (3-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-amide

The title compound is prepared analogously as described in example 11 using cyclopropanesulfonyl chloride instead of N,N-dimethylamidosulfonyl chloride.

MS: 434 [M+H]⁺

HPLC (Nucleosil C18 HD CC70, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 5%ACN): 4.008 min.

Example 13: Ethylsulfonic acid (3-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-amide

The title compound is prepared analogously as described in example 11 using ethylsulfonyl chloride instead of N,N-dimethylamidosulfonyl chloride.

MS: 422 [M+H]⁺

HPLC (Nucleosil C18 HD CC70, 6min method (0-3.5min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 5%ACN): 3.957 min.

Example 14: methylsulfonic acid (3-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-amide

The title compound is prepared analogously as described in example 11 using methylsulfonyl chloride instead of N,N-dimethylamidosulfonyl chloride.

MS: 408 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.58 min.

Example 15: N-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-formamide

- 173 -

The title compound is prepared analogously as described in example I1 using 3-formylamino-propionic acid instead of Cbz-beta-alanine.

MS: 358 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.45 min.

Example I6:

A) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-(3-{4-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-propionamide

To a solution of 3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionic acid (83 mg, 0.381 mmol) in dichloromethane (2 mL) are added O-benzotriazol-1-yl-tetramethyluronium hexafluorophosphate (196 mg, 0.519 mmol) and diisopropylethylamine (237 μ L, 1.384 mmol) before addition of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-[1-(3-amino-propionyl)-piperidin-4-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide (150 mg, 0.346 mmol). The resulting solution is stirred at rt during 16h and washed with aqueous 1N HCl and saturated aqueous NaHCO₃. The organic phase is dried and evaporated before purification by preparative HPLC (Column Interchrom C18 ODB 10 μ m 50x28, Gradient: 0-10 min 5% ACN, 10-20 min 5-90% ACN, 20-25 min 90% ACN) to yield an orange solid.

MS: 635 [M+H]⁺

HPLC (Waters Symmetry C18 3.5 μ m 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 3.14 min.

B) N-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionamide

To 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-(3-{4-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-propionamide (137 mg, 0.216 mmol) is added 4N HCl in dioxane (2 mL). The resulting mixture is stirred at rt during 1h before it is frozen, lyophilized and purified by preparative HPLC (Column Interchrom C18 ODS-AQ 10 μ m 50x20, Gradient: 0-2.5 min 2% ACN, 2.5-12.5 min 2-90% ACN, 12.5-15 min 90% ACN) to yield a light yellow solid.

MS: 553 [M+H]⁺

- 174 -

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.87 min.

Example 17: N-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-isobutyramide

The title compound is prepared analogously as described in example 16 using isobutyric acid instead of 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-propionic acid.

MS: 400 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.096 min.

Example 18: Cyclopropanecarboxylic acid (3-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-amide

The title compound is prepared analogously as described in example 16 using cyclopropanecarboxylic acid instead of 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-propionic acid.

MS: 398 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 1.75 min.

Example 19: 5-Oxo-pyrrolidine-2-carboxylic acid (3-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-amide

The title compound is prepared analogously as described in example 16 using 5-oxo-pyrrolidine-2-carboxylic acid instead of 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-propionic acid.

MS: 441 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.40 min.

Example 19a: 5-Oxo-pyrrolidine-2-carboxylic acid (3-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-amide maleate

The title compound is prepared analogously as described in example D2a using 5-Oxo-pyrrolidine-2-carboxylic acid (3-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-

- 175 -

3-oxo-propyl)-amide instead of cyclopropanesulfonic acid (2-{3-*exo*-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example I9b: 5-Oxo-pyrrolidine-2-carboxylic acid (3-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 5-Oxo-pyrrolidine-2-carboxylic acid (3-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-amide instead of cyclopropanesulfonic acid (2-{3-*exo*-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example I10: Pyridazine-4-carboxylic acid (3-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-amide

The title compound is prepared analogously as described in example I6 using pyridazine-4-carboxylic acid instead of 3-(1,3-Dioxo-1,3-dihydro-isindol-2-yl)-propionic acid.


MS: 436 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.49 min.

Example K1: 3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N-cyclopropyl-3-oxo-propionamide

This compound is prepared according to Scheme I :

- 176 -



Scheme K

A) 3-{4-[(R)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propionic acid benzyl ester

The title compound is prepared analogously as described in example I1 using malonic acid monobenzyl ester instead of Cbz-beta-alanine.

MS: 539 [M+H]⁺

HPLC (Waters Symmetry C18 3.5μm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 3.44 min.

- 177 -

B) 3-{4-[(R)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propionic acid

The title compound is prepared analogously as described in example I1 using 3-{4-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propionic acid benzyl ester instead of 3-{4-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propyl)-carbamic acid benzyl ester.

MS: 449 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.24 min.

C) N-Cyclopropyl-3-{4-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propionamide

To a solution of 3-{4-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propionic acid (45 mg, 0.0.1 mmol) in dichloromethane (2 mL) are added O-benzotriazol-1-yl-tetramethyluronium hexafluorophosphate (57 mg, 0.15 mmol) and diisopropylethylamine (69 uL, 0.4 mmol) before addition of cyclopropylamine (8 uL, 0.11 mmol). The resulting solution is stirred at rt during 3h and washed with water and brine. The organic phase is dried and evaporated before purification by preparative HPLC (Column Interchrom C18 ODB 10µm 50x28, Gradient: 0-2 min 10% ACN, 2-12 min 10-100% ACN, 12-15 min 100% ACN) to yield a colorless oil.

MS: 488 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.27 min.

D) 3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N-cyclopropyl-3-oxo-propionamide

To N-cyclopropyl-3-{4-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propionamide (40 mg, 0.082 mmol) is added 4N HCl in dioxane (2 mL). The resulting mixture is stirred at rt during 1h before it is frozen, lyophilized and purified by preparative HPLC (Column nucleosil C18HD 5 um 50x21, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield a white solid.

- 178 -

MS: 384 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.57 min.

Example K1a: 3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N-cyclopropyl-3-oxo-propionamide maleate

The title compound is prepared analogously as described in example D2a using 3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N-cyclopropyl-3-oxo-propionamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example K1b: 3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N-cyclopropyl-3-oxo-propionamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N-cyclopropyl-3-oxo-propionamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example K2: 3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N,N-diethyl-3-oxo-propionamide

The title compound is prepared analogously as described in example K1 using diethylamine instead of cyclopropylamine.

MS: 400 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.75 min.

Example K3: 3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N-((R)-2-hydroxy-propyl)-3-oxo-propionamide

The title compound is prepared analogously as described in example K1 using (R)-1-Amino-propan-2-ol instead of cyclopropylamine.

MS: 402 [M+H]⁺

- 179 -

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.39 min.

Example K4: 4-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N,N-dimethyl-4-oxo-butyramide

The title compound is prepared analogously as described in example K1 using succinic acid monobenzyl ester instead malonic acid monobenzyl ester and dimethylamine instead of cyclopropylamine.

MS: 386 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.260 min.

Example K5: 4-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N-(4-hydroxy-cyclohexyl)-4-oxo-butyramide

The title compound is prepared analogously as described in example K4 using 4-amino-cyclohexanol instead of dimethylamine.

MS: 456 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.46 min.

Example K6: 4-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N-cyclopropyl-4-oxo-butyramide

The title compound is prepared analogously as described in example K1 using succinic acid monobenzyl ester instead malonic acid monobenzyl ester.

MS: 398 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.56 min.

Example K7: 3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-N-cyclopropyl-3-oxo-propionamide

- 180 -

The title compound is prepared analogously as described in example K1 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 412 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.56 min.

Example K8 3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-N-(4-tert-butyl-cyclohexyl)-3-oxo-propionamide

The title compound is prepared analogously as described in example K7 using 4-terbutylcyclohexylamine instead of cyclopropylamine.

MS: 508 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 3.38 min.

Example K9: {3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-[1-(morpholine-4-carbonyl)-cyclopropyl]-methanone

A) 1-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid benzyl ester

The title compound is prepared analogously as described in example K7 using cyclopropane-1,1-dicarboxylic acid benzyl ester instead of malonic acid monobenzyl ester.

MS: 591 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.5 min.

B) 1-{3-exo-[(R)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid

To 1-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid benzyl ester (50 mg, 0.085 mmol) in MeOH (0.425 mL) is added an aqueous 1N LiOH solution (93.5 μ L, 0.093 mmol).

- 181 -

After stirring at rt during 16h, the pH is adjusted to 3 with aqueous 1N HCl, the mixture is extracted with chloroform and the organic phase is dried and evaporated to give a crude compound which is purified by preparative HPLC (Column Waters ODB 19x50 5 μ M, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield a white solid.

MS 501 [M+H]⁺

LCMS (Waters Symmetry C18 3.5 μ m 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 3.3 min.

C) 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-[1-(morpholine-4-carbonyl)-cyclopropyl]-methanone

The title compound is prepared analogously as described in example K7 using 1-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid instead of 3-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-3-oxo-propionic acid and morpholine instead of cyclopropylamine.

MS: 466 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.23 min.

Example K10: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid benzylamide

The title compound is prepared analogously as described in example K9 using benzylamine instead of morpholine.

MS: 487 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.17 min.

Example K10a: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid benzylamide maleate

The title compound is prepared analogously as described in example D2a using 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-

- 182 -

cyclopropanecarboxylic acid benzylamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example K10b: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid benzylamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid benzylamide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example K11: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid benzyl-methyl-amide

The title compound is prepared analogously as described in example K9 using N-methylbenzylamine instead of morpholine.

MS: 500 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.27 min.

Example K12: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid (2-methoxy-ethyl)-methyl-amide

The title compound is prepared analogously as described in example K9 using N-methyl-2-methoxyethylamine instead of morpholine.

MS: 468 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.22 min.

Example K13: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid dimethylamide

- 183 -

The title compound is prepared analogously as described in example K9 using dimethylamine instead of morpholine.

MS: 424 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.22 min.

Example K14: 1-{3-exo-[(R)-1-((S)-2-Methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-carbonyl}-cyclopropanecarboxylic acid amide

The title compound is prepared analogously as described in example K9 using ammonium carbonate instead of morpholine.

MS: 396 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.11 min.

Example K15: 4-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-N-cyclopropyl-4-oxo-butyramide

The title compound is prepared analogously as described in example K6 using 4-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-4-oxo-butyric acid instead of 4-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-4-oxo-butyric acid.

MS: 424 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.61 min.

Example K16: 4-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-N-(4-tert-butyl-cyclohexyl)-4-oxo-butyramide

The title compound is prepared analogously as described in example K15 using 4-tert-butylcyclohexylamine instead of cyclopropylamine.

MS: 522 [M+H]⁺

- 184 -

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 3.36 min.

Example K17: 4-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-N-benzyl-4-oxo-butylamide

The title compound is prepared analogously as described in example K15 using benzylamine instead of cyclopropylamine.

MS: 473 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.89 min.

Example K18: 3-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-N-benzyl-2,2-dimethyl-3-oxo-propionamide

The title compound is prepared analogously as described in example K7 using using dimethyl-1,1-dicarboxylic acid benzyl ester instead of cyclopropane-1,1-dicarboxylic acid benzyl ester and benzyl amine instead of morpholine.

MS: 489 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.29 min.

Example K19: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2,2-dimethyl-3-morpholin-4-yl-propane-1,3-dione

The title compound is prepared analogously as described in example K18 using morpholine instead of benzylamine.

MS: 468 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.22 min.

Example K20: {4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-[1-(morpholine-4-carbonyl)-cyclopropyl]-methanone

- 185 -

The title compound is prepared analogously as described in example K9 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane.

MS: 440 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.594 min.

Example K20a: {4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-[1-(morpholine-4-carbonyl)-cyclopropyl]-methanone maleate

The title compound is prepared analogously as described in example D2a using {4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-[1-(morpholine-4-carbonyl)-cyclopropyl]-methanone instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example K20b: {4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-[1-(morpholine-4-carbonyl)-cyclopropyl]-methanone toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using {4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-[1-(morpholine-4-carbonyl)-cyclopropyl]-methanone instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example K21: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidine-1-carbonyl}-cyclopropanecarboxylic acid benzyl-methyl-amide

The title compound is prepared analogously as described in example K11 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane.

MS: 474 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 3.00 min.

Example K22: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidine-1-carbonyl}-cyclopropanecarboxylic acid benzylamide

The title compound is prepared analogously as described in example K10 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane.

MS: 457 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.881 min.

Example K23 3-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-N-(4-hydroxy-cyclohexyl)-3-oxo-propionamide

The title compound is prepared analogously as described in example K7 using 4-Aminocyclohexanol instead of cyclopropylamine.

MS: 468 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.500 min.

Example K24 3-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-N,N-dimethyl-3-oxo-propionamide

The title compound is prepared analogously as described in example K7 using dimethylamine instead of cyclopropylamine.

MS: 398 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.413 min.

Example K25: {(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-(1-[(1S,3S,5R)-3-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-bicyclo[3.2.1]octane-8-carbonyl]-cyclopropyl)-methanone

The title compound is prepared analogously as described in example K9 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of morpholine.

MS: 663 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 2.974 min.

Example K26: 4-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-N,N-dimethyl-4-oxo-butamide

The title compound is prepared analogously as described in example K4 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 412 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.643 min.

Example K27: 4-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-N-(4-hydroxy-cyclohexyl)-4-oxo-butamide

The title compound is prepared analogously as described in example K5 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.


MS: 482 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.559 min.

Example L1: Tetrahydropyran-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-methyl-amide

This compound is prepared according to Scheme L:

- 188 -



Scheme L

A) Methyl-(2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-carbamic acid benzyl ester

The title compound is prepared analogously as described in example D1 using (2-chlorosulfonyl-ethyl)-methyl-carbamic acid benzyl ester instead of 2-phthalimidoethane sulfonyl chloride.

MS: 644 [M+H]⁺

- 189 -

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.53 min.

B) (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-[3-exo-8-(2-methylamino-ethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide

To methyl-(2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-carbamic acid benzyl ester (530 mg, 0.799 mmol) in ethanol (20 mL) is added Pd/C (85 mg, 0.799 mmol). After stirring at rt during 2h under H₂ atmosphere, the solution is filtered through celite and evaporated to yield a yellow solid.

MS: 510 [M+H]⁺

C) Tetrahydropyran-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-methyl-amide

The title compound is prepared analogously as described in example D6 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-[3-exo-8-(2-methylamino-ethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine.

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.13 min.

Example L1a: Tetrahydropyran-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-methyl-amide maleate

The title compound is prepared analogously as described in example D2a using tetrahydropyran-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-methyl-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example L2b: Tetrahydropyran-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-methyl-amide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using tetrahydropyran-4-carboxylic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-methyl-amide instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example L2: (2-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-methyl-carbamic acid ethyl ester

The title compound is prepared analogously as described in example D13 using (S)-2-methyl-propane-2-sulfinic acid [(R)-1-{3-exo-8-(2-methylamino-ethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-yl}-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of 2-{3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethylamine.


MS: 478 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.21 min.

Example M1: 1-{3-exo-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl}-2-(morpholine-4-sulfonyl)-ethanone

This compound is prepared according to Scheme M:

- 191 -



Scheme M

A) Cyclohexylsulfanyl-acetic acid

To cyclohexylmercaptane (147 μ L, 1.2 mmol) in DMF (1 mL) are added bromoacetic acid (167 mg, 1.2 mmol) and DIPEA (616 μ L, 3.6 mmol). The resulting mixture is stirred at rt during 2h and used such as in the next step.

B) (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-[1-(2-cyclohexylsulfanyl-acetyl)-piperidin-4-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide

To (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide (200 mg, 0.552 mmol) in dichloromethane (1 mL) are added EDC (127mg, 0.662 mmol), HOBt (97 mg, 0.718 mmol), DIPEA (283 μ L, 1.656 mmol) and cyclohexylsulfanyl-acetic acid in DMF (2 mL, 1.2 mmol). After stirring at rt during 3 days, the mixture is washed with an aqueous 1N HCl solution, the organic phase is dried and evaporated to give the title compound.

MS 519 [M+H]⁺

LCMS (Waters Symmetry C18 3.5 μ m 2.1 x 50mm, 6min method (0-3min 20-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-20%ACN, 5.55-6min 20%ACN): 4.236 min.

- 192 -

C) 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-cyclohexylsulfanyl-ethanone

To (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-[1-(2-cyclohexylsulfanyl-acetyl)-piperidin-4-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide (367 mg, 0.552 mmol) is added 4N HCl in dioxane (2 mL). The resulting mixture is stirred at rt during 20 min. The solution is frozen and lyophilised to give a yellow oil before purification by preparative HPLC HPLC (Column YMC ODS-AQ 20x50 5 uM, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield a colorless oil .

MS 415 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.287 min.

D) 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-cyclohexanesulfonyl-ethanone

To 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-cyclohexylsulfanyl-ethanone (84 mg, 0.203 mmol) in acetic acid (465 uL) is added an aqueous 30% H₂O₂ solution (83 uL, 0.812 mmol). The resulting mixture is stirred at 80°C during 30 min. The solution is frozen and lyophilised to give a colorless oil

MS: 447 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.219 min.

Example M1a: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-cyclohexanesulfonyl-ethanone maleate

The title compound is prepared analogously as described in example D2a using 1-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-cyclohexanesulfonyl-ethanone instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example M1b: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-cyclohexanesulfonyl-ethanone toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 1-{4-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-2-cyclohexanesulfonyl-ethanone instead of cyclopropanesulfonic acid (2-{3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-azabicyclo[3.2.1]octane-8-sulfonyl}-ethyl)-amide.

Example M2: 2-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propane-1-sulfonyl)-N,N-diethyl-acetamide

A) 3-Diethylcarbamoylmethylsulfonyl-propionic acid

To 3-mercaptopropionic acid (200 mg, 1.884 mmol) in methanol (3 mL) are added sodium methoxide (204 mg, 3.768 mmol) and 2-Bromo-N,N-diethyl-acetamide (402 mg, 2.070 mmol). After stirring at 80°C during 4h and evaporation of solvent, the residue is treated with ethyl acetate and water, the pH is adjusted to 10 with an aqueous NaHCO₃ solution and the aqueous phase is extracted with ethyl acetate. After acidification of the aqueous phase to pH 2 with an aqueous 1N HCl solution, it is extracted with ethyl acetate, then the organic phase is dried and evaporated to give the title compound.

MS: 220 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 0.9 min.

B) 2-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propane-1-sulfonyl)-N,N-diethyl-acetamide

The title compound is prepared analogously as described in example M1 using 3-Diethylcarbamoylmethylsulfonyl-propionic acid instead of Cyclohexylsulfonyl-acetic acid.

MS: 492 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.548 min.

Example M3: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-cyclopropylmethanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example M2 using Bromomethylcyclopropane instead of 2-Bromo-N,N-diethyl-acetamide.

- 194 -

MS: 433 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.48 min.

Example M4: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-methanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example M1 using 3-Methylthiopropionic acid instead of Cyclohexylsulfanyl-acetic acid.

MS: 392 [M+H]⁺

Example M5: 1-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-cyclohexanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example M1 using 3-bromopropionic acid instead of bromoacetic acid.

MS: 461 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.221 min.

Example M6: 7-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propane-1-sulfonyl)-3,4-dihydro-2H-isoquinolin-1-one

A) 1-Oxo-1,2,3,4-tetrahydro-isoquinoline-7-sulfonyl chloride

To Chlorosulfonic acid (30 mL) is added 3,4-Dihydro-2H-isoquinolin-1-one (5 g, 34 mmol) at 0°C. After stirring at rt during 1h, the mixture is heated at 50°C during 16h, then it is poured carefully into an ice bath and stirred at 0°C during 30min. The precipitate is filtered and dried in oven at 60°C to give the title compound.

MS: 246 [M+H]⁺

B) 7-Mercapto-3,4-dihydro-2H-isoquinolin-1-one

To a zinc powder (2.79 g, 42.7 mmol) and Dichlorodimethylsilane (5.15 mL, 42.7 mmol) in DCE (45 mL) is added a solution of 1-Oxo-1,2,3,4-tetrahydro-isoquinoline-7-sulfonyl chloride

- 195 -

(3 g, 12.2 mmol) and 1,3-dimethylimidazolidin-2-one (3.96 mL, 36.6 mmol) in DCE (5 mL).

After stirring at rt during 2h, the mixture is concentrated.

MS: 180 [M+H]⁺

C) 7-(3-{4-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-piperidin-1-yl}-3-oxo-propane-1-sulfonyl)-3,4-dihydro-2H-isoquinolin-1-one

The title compound is prepared analogously as described in example M1 using 7-Mercapto-3,4-dihydro-2H-isoquinolin-1-one instead of cyclohexylmercaptane.

MS: 510 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 5%ACN): 2.776 min.

Example M7: 1-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-2-cyclohexanesulfonyl-ethanone

The title compound is prepared analogously as described in example M1 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 473 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.239 min.

Example M8: 2-(3-[(1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl]-3-oxo-propane-1-sulfonyl)-N,N-diethyl-acetamide

The title compound is prepared analogously as described in example M2 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 518 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.648 min.

Example M8a: 2-(3-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-oxo-propane-1-sulfonyl)-N,N-diethyl-acetamide maleate

The title compound is prepared analogously as described in example D2a using 2-(3-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-oxo-propane-1-sulfonyl)-N,N-diethyl-acetamide instead of cyclopropanesulfonic acid (2-(3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example M8b: 2-(3-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-oxo-propane-1-sulfonyl)-N,N-diethyl-acetamide toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 2-(3-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-oxo-propane-1-sulfonyl)-N,N-diethyl-acetamide instead of cyclopropanesulfonic acid (2-(3-exo-[(R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example M9: 1-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-cyclopropylmethanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example M3 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 459 [M+H]⁺

HPLC (YMC, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 20%ACN): 2.573 min.

Example M10: 1-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-cyclohexanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example M5 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-

- 197 -

bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 487 [M+H]⁺


HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.241 min.

Example M11: 1-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-yl)-3-methanesulfonyl-propan-1-one

The title compound is prepared analogously as described in example M4 using 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-piperidin-4-yl-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

Example N1: 2-(3-((1S,3S,5R)-3-[(R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]oct-8-ylmethyl)-benzyl)-isoindole-1,3-dione

This compound is prepared according to Scheme N:



Scheme N

A) 2-(3-Hydroxymethyl-benzyl)-isoindole-1,3-dione

- 198 -

To 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-benzoic acid (300 mg, 1.066 mmol) in DCM (1 mL) are added at 0°C triethylamine (164 μ L, 1.173 mmol) and ethylchloroformate (112 μ L, 1.173 mmol). After stirring at 0°C during 15 min and filtration, the obtained solution is added at 0°C to NaBH₄ (61 mg, 1.599 mmol) in water (400 μ L) before stirring at 0°C during 30 min and at rt during 2h. The solution is acidified to pH 3 with an aqueous 1N HCl solution, extracted with ethylacetate, the organic phase is dried and evaporated to give a crude compound before purification by flash chromatography on silica gel (eluent: cyclohexane/ethyl acetate 9/1 to 1/1) to yield a colorless gum.

TLC, R_f (cyclohexane/ethyl acetate 1/1) = 0.15.

B) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-benzaldehyde

To oxalylchloride (43 μ L, 0.449 mmol) in DCM (400 μ L) is added at -78°C DMSO (43 μ L, 0.598 mmol). After stirring at -78°C during 15 min, 2-(3-Hydroxymethyl-benzyl)-isoindole-1,3-dione (80 mg, 0.299 mmol) in DCM (600 μ L) is added and the mixture is stirred at -78°C during 45 min before addition of triethylamine (210 μ L, 1.495 mmol). After stirring at rt during 4h, the reaction is quenched with water, extracted with DCM, the organic phase is washed with an aqueous 10% NaHSO₄ solution and an aqueous saturated NaHCO₃ solution. The organic phase is dried and evaporated to give a crude compound before purification by flash chromatography on silica gel (eluent: cyclohexane/ethyl acetate 9/1 to 2/1) to yield a colorless gum.

MS: 266 [M+H]⁺

TLC, R_f (cyclohexane/ethyl acetate 1/1) = 0.7.

C) (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-((1S,3S,5R)-8-[3-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-benzyl]-8-aza-bicyclo[3.2.1]oct-3-yl)-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide

3-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-benzaldehyde (44 mg, 0.166 mmol) and 3-exo-[(R)-1-((S)-2-methyl-propane-2-sulfinylamino)-2-(2,4,5-trifluoro-phenyl)-ethyl]-8-aza-bicyclo[3.2.1]octane (54 mg, 0.138 mmol) in DCE (500 μ L) are stirred at rt during 1h before added of sodium triacetoxymethylborohydride (74 mg, 0.345 mmol). The mixture is stirred at rt during 16h and the solvent is evaporated to give a crude compound before purification by preparative HPLC (Column Waters C18 ODB 5 μ m 19x50, Gradient: 0-2.5 min 5% ACN, 2.5-12.5 min 5-100% ACN, 12.5-15 min 100% ACN) to yield the title compound.

- 199 -

MS: 638 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.361 min.

D) 2-(3-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-ylmethyl)-benzyl)-isoindole-1,3-dione

The title compound is prepared analogously as described in example G17 using S)-2-Methyl-propane-2-sulfinic acid [(R)-1-((1S,3S,5R)-8-[3-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-benzyl]-8-aza-bicyclo[3.2.1]oct-3-yl)-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-((1S,3S,5R)-8-[3-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-benzoyl]-8-aza-bicyclo[3.2.1]oct-3-yl)-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 534 [M+H]⁺

HPLC (Zorbax SB C18, 2min method (0-0.8min 10-95%ACN, 0.8-1.5min 95%ACN, 1.5-1.6min 95-10%ACN, 1.6-2min 10%ACN): 1.228 min.

Example N1a: 2-(3-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-ylmethyl)-benzyl)-isoindole-1,3-dione maleate

The title compound is prepared analogously as described in example D2a using 2-(3-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-ylmethyl)-benzyl)-isoindole-1,3-dione instead of cyclopropanesulfonic acid (2-(3-exo-((R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example N1b: 2-(3-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-ylmethyl)-benzyl)-isoindole-1,3-dione toluene-4-sulfonate

The title compound is prepared analogously as described in example D2b using 2-(3-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-ylmethyl)-benzyl)-isoindole-1,3-dione instead of cyclopropanesulfonic acid (2-(3-exo-((R)-1-amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]octane-8-sulfonyl)-ethyl)-amide.

Example N2: N-(3-((1S,3S,5R)-3-((R)-1-Amino-2-(2,4,5-trifluoro-phenyl)-ethyl)-8-aza-bicyclo[3.2.1]oct-8-ylmethyl)-benzyl)-benzamide

- 200 -

The title compound is prepared analogously as described in example G18 using (S)-2-Methyl-propane-2-sulfinic acid [(R)-1-[(1S,3S,5R)-8-[3-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-benzyl]-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide instead of (S)-2-methyl-propane-2-sulfinic acid [(R)-1-[3-*exo*-8-[3-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-benzoyl]-8-aza-bicyclo[3.2.1]oct-3-yl]-2-(2,4,5-trifluoro-phenyl)-ethyl]-amide.

MS: 508 [M+H]⁺

HPLC (Waters Symmetry C18 3.5µm 2.1 x 50mm, 6min method (0-3min 5-95%ACN, 3.5-5.5min 95%ACN, 5.5-5.55min 95-5%ACN, 5.55-6min 5%ACN): 2.863 min.

Example P: Activity Assay

Various compounds of the invention were tested for their inhibitory activity to human DPP-IV.

Materials

Human DPP-IV consisting of amino acids 39 to 766 followed by a C-terminal Streptavidin-tag was expressed using the baculovirus system and purified to >80% purity. The enzyme was stored in 25 mM Tris buffer, pH 9.0, containing 300 mM NaCl at -80° C.

The fluorogenic substrates H-Gly-Pro-AMC was purchased from Bachem AG (Bubendorf, Switzerland). The substrate was kept as a 5 mM stock solution in DMSO at -20° C. All other chemicals were purchased from Sigma (Buchs, Switzerland).

The assay buffer for the DPP-IV reaction was 25 mM Tris/HCl, pH 7.5, containing 140 mM NaCl, 10 mM KCl and 0.05% (w/v) CHAPS.

Compound and liquid handling

The test compounds were dissolved in 90% DMSO/10% H₂O (v/v). Serial dilutions of the compounds from 3 mM to 0.03 µM in 90% DMSO/10% H₂O (v/v) followed by a 1:33.3 dilution in assay buffer was done in 96-well polypropylene plates using a CyBio Dilus 8-channel pipettor (CyBio AG, Jena, Germany) with tip change after each pipetting step. The compound solutions as well as the substrate and the enzyme solutions were transferred to the assay plates (384-well black Cliniplate; cat. no. 95040020 Labsystems Oy, Finland) by means of a CyBi-Well 96-channel pipettor (CyBio AG, Jena, Germany).

Kinetic measurements

- 201 -

Enzyme kinetics were measured by mixing 10 μ l of a 3-fold concentrated substrate solution in assay buffer (final substrate concentration was 10 μ M) with 10 μ l of the corresponding compound solution. The reactions were initiated by addition of 10 μ l of a 3-fold concentrated solution of the enzyme in assay buffer. Final enzyme (active site) concentrations in the assay was 10 pM for DPP-IV. Fluorescence product (AMC) formation was monitored for 1 hour at room temperature at 35 second intervals by measuring the fluorescence emission at 500 nm using an excitation wavelength of 350 nm in a TECAN Ultra fluorescence reader (TECAN, Maennedorf, Switzerland). The fluorescence in each well was excited by one flash per measurement. The Origin software package (Origin 7.5 Mircocal, Northampton, MA, USA) was used to generate all graphs and to perform the IC₅₀ calculations.

Results

The inhibitory activities (IC₅₀ values) of the compounds to human DPP-IV were found to be 4.7 μ M or less and in many cases 0.01 μ M or less. In the case of exemplary compounds, their IC₅₀ values were found to be between 4.7 and 0.0001 μ M or between 4.7 μ M and 0.0053 μ M.


Representative examples.

Examples	hDPP-IV IC ₅₀ (μ M)
D1	0.0077
E7	0.0006
G4	0.01
H13	0.007

- 202 -

Claims

1. A compound of Formula (I):



(I)

wherein

the asterisk * designates a chiral centre of (*R*) or (*S*) configuration;

V is absent or is ethylene;

W is -C(O)- or -S(O)_{*l*};

X is a linker having 1 to 12 (e.g. 1 to 6) in-chain atoms and comprising one or more linkages selected from -O-, -C(O)-, -S(O)_{*l*}-, -N(R⁹)- and hydrocarbylene optionally substituted with 1, 2, 3, 4 or 5 R¹¹;

Y is a linker selected from -O-, -N(R⁹)-, -C(O)-, -C(O)O-, -C(O)N(R⁹)-, -S(O)_{*l*}- and -S(O)_{*l*}N(R⁹)-;

R¹ is selected from hydrogen; -N(R⁹)(R¹⁰); hydrocarbyl optionally substituted with 1, 2, 3, 4 or 5 R¹¹; hydrocarbyloxy optionally substituted with 1, 2, 3, 4 or 5 R¹¹; and -(CH₂)_{*k*}-heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R¹¹;

or, when Y is -N(R⁹)-, R¹ and R⁹ taken together with the nitrogen atom to which they are attached may form a heterocycle, wherein said heterocycle is bound to X via said nitrogen atom and is optionally substituted with 1, 2, 3, 4 or 5 R¹¹;

- 203 -

R^2 and R^3 are each independently selected from R^8 , $-OR^8$, $-C(O)R^8$, $-C(O)OR^8$ and $-S(O)_lR^9$;

R^4 and R^5 are each independently selected from hydrogen, hydroxy, halogen and C_{1-6} alkyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

R^6 is aryl or heteroaryl, either of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

R^8 is selected from hydrogen; hydrocarbyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} ; and $-(CH_2)_k$ -heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

R^9 and R^{10} are each independently selected from R^8 , $-OR^8$, $-C(O)R^8$, $-C(O)OR^8$ and $-S(O)_lR^9$; or R^9 and R^{10} taken together with a nitrogen atom to which they are attached form heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

each R^{11} is independently selected from R^{12} ; hydrocarbyl optionally substituted with 1, 2, 3, 4 or 5 R^{12} ; and $-(CH_2)_k$ -heterocyclyl optionally substituted with 1, 2, 3, 4 or 5 R^{12} ;

R^{12} is independently selected from halogen, trifluoromethyl, cyano, nitro, oxo, $=NR^{13}$, $-OR^{13}$, $-C(O)R^{13}$, $-C(O)N(R^{13})R^{14}$, $-C(O)OR^{13}$, $-OC(O)R^{13}$, $-S(O)_lR^{13}$, $-S(O)_lN(R^{13})R^{14}$, $-N(R^{13})R^{14}$, $-N(R^{13})N(R^{13})R^{14}$, $-N(R^{13})C(O)R^{14}$ and $-N(R^{13})S(O)_lR^{13}$;

R^{13} and R^{14} are each independently hydrogen or selected from hydrocarbyl and $-(CH_2)_k$ -heterocyclyl, either of which is optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from oxo, halogen, cyano, amino, hydroxy, C_{1-6} alkyl and C_{1-6} alkoxy;


k is 0, 1, 2, 3, 4, 5 or 6; and

l is 0, 1, or 2;

or a pharmaceutically acceptable salt or prodrug thereof.


- 204 -

2. A compound according to claim 1, which is of the Formula (II):



or a pharmaceutically acceptable salt or prodrug thereof.


3. A compound according to claim 1, which is of the Formula (III):



or a pharmaceutically acceptable salt or prodrug thereof.

4. A compound according to any preceding claim, wherein X is $-X^1-$ and wherein X^1 is selected from $-O-$, $-C(O)-$, $-S(O)-$, $-N(R^9)-$ and hydrocarbylene (e.g. C_{1-6} alkylene) optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

5. A compound according to claim 4, which is of the Formula (IV):



- 205 -

wherein X^1 is $-N(R^9)-$ or hydrocarbylene (e.g. C_{1-6} alkylene) optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

or a pharmaceutically acceptable salt or prodrug thereof.

6. A compound according to claim 5, wherein X^1 is C_{1-6} alkylene optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

7. A compound according to claim 6, wherein $-W-X^1-Y-$ is one of the following linkers:

No.	W	X^1	Y
1	$-C(O)-$	C_{1-6} alkylene	$-S(O)_2-$
2	$-S(O)_2-$	C_{1-6} alkylene	$-C(O)-$

wherein X^1 is optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

8. A compound according to claim 7, wherein $-W-X^1-Y-$ is one of the following linkers:

No.	W	X^1	Y
1	$-C(O)-$	$-CH_2-$	$-S(O)_2-$
2	$-C(O)-$	$-CH(CH_3)-$	$-S(O)_2-$
3	$-C(O)-$	$-C(CH_3)_2-$	$-S(O)_2-$
4	$-C(O)-$	$-CH_2CH_2-$	$-S(O)_2-$
5	$-S(O)_2-$	$-CH_2-$	$-C(O)-$
6	$-S(O)_2-$	$-CH(CH_3)-$	$-C(O)-$
7	$-S(O)_2-$	$-C(CH_3)_2-$	$-C(O)-$
8	$-S(O)_2-$	$-CH_2CH_2-$	$-C(O)-$

wherein X^1 is optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

9. A compound according to claim 4, wherein $-W-X^1-Y-$ is one of the following linkers:

- 206 -

No.	W	X ¹	Y
1	-C(O)-	cycloalkylene	-S(O) ₂ -
2	-S(O) _r -	cycloalkylene	-C(O)-
3	-C(O)-	cycloalkylene	-C(O)-
4	-S(O) _r -	cycloalkylene	-S(O) ₂ -

wherein X¹ is optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

10. A compound according to claim 9, wherein -W-X¹-Y- is one of the following linkers:


No.	W	X ¹	Y
1	-C(O)-	cyclopropylene	-S(O) ₂ -
2	-S(O) _r -	cyclopropylene	-C(O)-
3	-C(O)-	cyclopropylene	-C(O)-
4	-S(O) _r -	cyclopropylene	-S(O) ₂ -

wherein X¹ is optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

11. A compound according to any of claims 1 to 3, wherein X is -X¹-X²-, wherein X¹ and X² are each independently selected from -O-, -C(O)-, -S(O)_r-, -N(R⁹)-, and C₁₋₆ alkylene optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

- 207 -

12. A compound according to claim 11, which is of the Formula (V):



(V)

wherein

one of X¹ and X² is -N(R⁹)-; and the other is C₁₋₆ alkylene optionally substituted with 1, 2, 3, 4 or 5 R¹¹;

or a pharmaceutically acceptable salt or prodrug thereof.

13. A compound according to claim 11 or claim 12, wherein -W-X¹-X²-Y- is one of the following linkers:

No.	W	X ¹	X ²	Y
1	-C(O)-	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-
2	-C(O)-	C ₁₋₆ alkylene	-N(R ⁹)-	-S(O) ₂ -
3	-S(O) ₂ -	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-
4	-S(O) ₂ -	C ₁₋₆ alkylene	-N(R ⁹)-	-S(O) ₂ -

wherein X¹ is optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

14. A compound according to claim 13, wherein -W-X¹-X²-Y- is one of the following linkers:

No.	W	X ¹	X ²	Y
1	-C(O)-	-CH ₂ -	-N(R ⁹)-	-C(O)-
2	-C(O)-	-CH ₂ -	-N(R ⁹)-	-S(O) ₂ -
3	-C(O)-	-CH(CH ₃)-	-N(R ⁹)-	-C(O)-
4	-C(O)-	-CH(CH ₃)-	-N(R ⁹)-	-S(O) ₂ -
5	-C(O)-	-C(CH ₃) ₂ -	-N(R ⁹)-	-C(O)-
6	-C(O)-	-C(CH ₃) ₂ -	-N(R ⁹)-	-S(O) ₂ -

- 208 -

No.	W	X ¹	X ²	Y
7	-C(O)-	-CH ₂ CH ₂ -	-N(R ⁹)-	-C(O)-
8	-C(O)-	-CH ₂ CH ₂ -	-N(R ⁹)-	-S(O) ₂ -
9	-S(O) ₂ -	-CH ₂ -	-N(R ⁹)-	-C(O)-
10	-S(O) ₂ -	-CH ₂ -	-N(R ⁹)-	-S(O) ₂ -
11	-S(O) ₂ -	-CH(CH ₃)-	-N(R ⁹)-	-C(O)-
12	-S(O) ₂ -	-CH(CH ₃)-	-N(R ⁹)-	-S(O) ₂ -
13	-S(O) ₂ -	-C(CH ₃) ₂ -	-N(R ⁹)-	-C(O)-
14	-S(O) ₂ -	-C(CH ₃) ₂ -	-N(R ⁹)-	-S(O) ₂ -
15	-S(O) ₂ -	-CH ₂ CH ₂ -	-N(R ⁹)-	-C(O)-
16	-S(O) ₂ -	-CH ₂ CH ₂ -	-N(R ⁹)-	-S(O) ₂ -

wherein X¹ is optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

15. A compound according to any preceding claim, wherein R¹ is C₁₋₆ alkyl, -(CH₂)_k-cycloalkyl or -(CH₂)_k-aryl, any of which is optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

16. A compound according to claim 15, wherein R¹ is selected from methyl; cyclopropyl optionally substituted with 1 or 2 R¹¹; and phenyl optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

17. A compound according to any of claims 1 to 14, wherein R¹ is heterocyclyl (e.g. morpholinyl) optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

18. A compound according to any preceding claim, wherein R² and R³ are each hydrogen.

19. A compound according to any preceding claim, wherein R⁴ and R⁵ are each hydrogen.


20. A compound according to any preceding claim, wherein R⁶ is aryl optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

21. A compound according to claim 20, wherein R⁶ is phenyl optionally substituted with 1, 2, 3, 4 or 5 R¹¹.


- 209 -

22. A compound according to claim 19, wherein the or each R^{11} is independently halogen.

23. A compound according to any of claims 5 to 10, or any of claims 15 to 17, of the following Formulae:



(XIII)




(XIV)

wherein;

- p is 1, 2, 3, 4 or 5, preferably 0, 1, 2 or 3,
- R^{11} is independently halogen.

or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

24. A compound of the formula (XIV) according to claim 23, which is in the exo configuration as represented by the following Formulae (XIVb);




(XIVb)


or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

25. A compound according to any of claims 11 to 17, of the following Formulae:

- 210 -



(XIX)




(XX)

wherein;

- p is 1, 2, 3, 4 or 5, preferably 0, 1, 2 or 3,
- R¹¹ is independently halogen.

or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

26. A compound of the formula (XX) according to claim 25, which is in the exo configuration as represented by the following Formulae (XXb);



(XXb)

or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

27. A compound according to any of claims 22 to 26, wherein R⁶ is 2,4,5-trifluorophenyl, or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

28. A compound according to any of claims 1 to 21, wherein V is ethylene, and which is in an "exo" configuration, or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

- 211 -

29. A compound according to any of claims 1 to 28, wherein;


i) Y is $-\text{C}(\text{O})-$; R^1 is C_{1-6} alkyl, cycloalkyl, aryl or heterocyclyl, any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} , wherein R^{11} is independently selected from halogen, hydroxy, cyano, amino, $-\text{C}(\text{O})\text{OH}$, C_{1-6} alkyl, C_{1-6} alkoxy, $-\text{C}(\text{O})-\text{C}_{1-6}$ alkyl, $-\text{C}(\text{O})\text{O}-\text{C}_{1-6}$ alkyl, $-\text{S}(\text{O})-\text{C}_{1-6}$ alkyl, $-\text{NH}(\text{C}_{1-6}$ alkyl) and $-\text{N}(\text{C}_{1-6}$ alkyl) $_2$, wherein any C_{1-6} alkyl group present is optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from halogen, cyano, amino, hydroxy and C_{1-6} alkoxy,

ii) Y is $-\text{S}(\text{O})_2-$ or $-\text{S}(\text{O})\text{N}(\text{R}^9)-$; R^1 is hydrogen, C_{1-6} alkyl, cycloalkyl, aryl, or heterocyclyl, any of which is optionally substituted with 1, 2, 3, 4 or 5 R^{11} wherein R^{11} is independently selected from halogen, hydroxy, cyano, amino, $-\text{C}(\text{O})\text{OH}$, C_{1-6} alkyl, C_{1-6} alkoxy, $-\text{C}(\text{O})-\text{C}_{1-6}$ alkyl, $-\text{C}(\text{O})\text{O}-\text{C}_{1-6}$ alkyl, $-\text{S}(\text{O})-\text{C}_{1-6}$ alkyl, $-\text{NH}(\text{C}_{1-6}$ alkyl) and $-\text{N}(\text{C}_{1-6}$ alkyl) $_2$, wherein any C_{1-6} alkyl group present is optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from halogen, cyano, amino, hydroxy and C_{1-6} alkoxy, or

iii) R^1 is $-\text{N}(\text{R}^9)(\text{R}^{10})$,

or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

30. A compound according to claim 1, which is of the Formula (VI):




(VI)

or a pharmaceutically acceptable salt or prodrug thereof.

31. A compound according to claim 20, which is of the Formula (IX):

- 212 -




(IX)

wherein X^1 is selected from $-O-$, $-C(O)-$, $-S(O)_2-$, $-N(R^9)-$ and hydrocarbylene (e.g. C_{1-6} alkylene) optionally substituted with 1, 2, 3, 4 or 5 R^{11} ;

or a pharmaceutically acceptable salt or prodrug thereof.

32. A compound according to claim 31, which is of the Formula (XII):



(XII)


wherein p is 0, 1, 2, 3, 4 or 5;

or a pharmaceutically acceptable salt or prodrug thereof.

33. A compound according to claim 31 or claim 32, wherein X^1 is;

- i) $-N(R^9)-$ or hydrocarbylene (e.g. C_{1-6} alkylene) optionally substituted with 1, 2, 3, 4 or 5 R^{11} ,
or
- ii) $-N(R^9)-$ or C_{1-6} alkylene optionally substituted with 1, 2, 3, 4 or 5 R^{11} .

34. A compound according to claim 31, which is of the Formula (XV):




- 213 -

(XV)

wherein X^1 and X^2 are each independently selected from -O-, -C(O)-, -S(O)-, -N(R⁹)- and C₁₋₆ alkylene optionally substituted with 1, 2, 3, 4 or 5 R¹¹;

or a pharmaceutically acceptable salt or prodrug thereof.

35. A compound according to claim 34, which is of the Formula (XVIII):




(XVIII)

wherein p is 0, 1, 2, 3, 4 or 5;




or a pharmaceutically acceptable salt or prodrug thereof.

36. A compound according to claim 34 or claim 35, wherein one of X^1 and X^2 is -N(R⁹)-; and the other is C₁₋₆ alkylene optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

37. A compound according to claim 11 or claim 12, or any of claims 15 to 22, wherein -W-X¹-Y- is one of the following linkers:



No.	W	X ¹	Y
1	-C(O)-	C ₁₋₆ alkylene	-C(O)-N(R ⁹)-
2	-C(O)-	-CH ₂ -	-C(O)-NH-
3	-C(O)-		-C(O)-

- 214 -



4	-C(O)-		-C(O)-
5	-C(O)-		-N(R ⁹)-
6	-C(O)-		-N(R ⁹)-
7	-C(O)-	arylene	-C(O)-
8	-C(O)-	C ₁₋₆ alkylene	-C(O)-
9	-C(O)-	arylene	-N(R ⁹)-
10	-C(O)-	-CH ₂ -	-C(O)-
11	-C(O)-	-CH(CH ₃)-	-C(O)-
12	-C(O)-	-C(CH ₃) ₂ -	-C(O)-
13	-C(O)-	-CH ₂ CH ₂ -	-C(O)-

wherein X¹ is optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

38. A compound according to claim 13, or any of claims 15 to 22, wherein -W-X¹-Y- is one of the following linkers:

1	-S(O) _r -	C ₁₋₆ alkylene	-N(R ⁹)-	-C(O)-O-
2	-S(O) _r -	-CH ₂ CH ₂ -	-N(R ⁹)-	-C(O)-O-
3	-C(O)-	arylene	-N(R ⁹)-	-C(O)-
4	-C(O)-		-N(R ⁹)-	-C(O)-
5	-C(O)-		-N(R ⁹)-	-C(O)-

- 215 -

6	-C(O)-		C ₁₋₆ alkylene	-N(R ⁹)-
7	-C(O)-		-CH ₂ CH ₂ -	-N(R ⁹)-

wherein X¹ is optionally substituted with 1, 2, 3, 4 or 5 R¹¹ or X² is optionally substituted with 1, 2, 3, 4 or 5 R¹¹ when X² is alkylene.

39. A compound according to any of the previous claims wherein X¹ or X² is an alkylene or arylene substituted by 1, 2, 3, 4 or 5 R¹¹, preferably 1 or 2 R¹¹.


40. A compound according to claim 39 wherein R¹¹ is selected from a spiro group, C₁₋₆ alkyl, -O-C₁₋₆ alkyl, -NH-C(O)-(C₁₋₆ alkyl), phenyl, benzyl, hydroxy, halogen, amino, wherein the alkyl group is optionally substituted by amino, hydroxy, -C(O)-N(C₁₋₆ alkyl)(C₁₋₆ alkyl), -C(O)-NH(C₁₋₆ alkyl), -C(O)-NH₂, (C₁₋₆ alkyl), or halogen.

41. A compound according to any of the previous claims wherein R¹ is


- i) hydrocarbyloxy, optionally substituted with 1, 2, 3, 4 or 5 R¹¹, or
- ii) C₁₋₆ alkoxy, optionally substituted with 1, 2, 3, 4 or 5 R¹¹.

42. A compound according to claim 1, selected from:


- 216 -




B1




C1




C2




D1




D2




D3



D4




E1




E2


- 217 -




E3




E4




E5




E6




F1




F2



G1




G2




G3


- 218 -




G4




H1




H2




H3




X1




X2




X3




X4




X5



X6




X7




X8


- 219 -




X9




X10




X11




X12




X13




X14




X15




X16




X17




X18



X19




X20




X21

or, in each case, a pharmaceutically acceptable salt or prodrug thereof.


43. A compound according to claim 1, selected from:




D5




D6




D7




D8




D9




D10



D11




D12




D13


- 221 -




D14




D15




D16




D17




D18




D19



E7




E8




E9


- 222 -




E10




E11




E12




E13




E14




E15



E16




E17




E18


- 223 -




E19




E20




E21




E22




E23




E24



E25




E26




E27


- 224 -




E28




E29




E30




E31




E32




E33



E34




E35




E36


- 225 -




E37




E38




E39




E40




E41




E42



F3




F4




F5


- 226 -




F6




F7




F8




F9



F10




F11




F12


- 227 -




G5




G6




G7




G8




G9




G10



G11




G12




G13


- 228 -




G14




G15




G16




G17




G18




G19



G20




G21




G22


- 229 -




G23




G24




G25




G26




G27




G28



G29




G30




G31


- 230 -




G32




G33




G34




G35




G36




G37



G38




G39




G40


- 231 -




G41




G42




G43




G44




G45




G46



G47




G48




G49


- 232 -




G50




G51




G52




G53




G54




G55



G56




G57




G58


- 233 -




G59




G60




G61




G62




G63




G64



G65




G66




G67


- 234 -




H4




H5




H6




H7




H8




H9



H10




H11




H12


- 235 -




H13




H14




I1




I2




I3



I4




I5




I6


- 236 -




17




18




19




110



K1






K2




K3


- 237 -




K4




K5




K6




K7



K8



K9




K10


K11

K12


- 238 -




K13




K14




K15




K16




K17




K18



K19




K20




K21


- 239 -




K22




K23




K24




K25



K26




L1




L2

K27


- 240 -




M1




M2




M3




M4




M5




M6



M7




M8




M9


- 241 -




M10



M11



N1



N2

or, in each case, a pharmaceutically acceptable salt or prodrug thereof.

44. A compound according to any preceding claim, in a substantially pure form of the (*R*) enantiomer, wherein (*R*) refers to the stereochemical configuration at C^{*}.

45. A compound according to any preceding claim, in a substantially pure form of the *exo* configuration of the bridged piperidine ring.

46. A compound according to claim 45, wherein the purity is greater than 95 %.

47. A compound according to any preceding claim for use in therapy.

48. A pharmaceutical formulation comprising a compound of any of claims 1 to 43.

- 242 -

49. A formulation according to claim 48, which further comprises a pharmaceutically acceptable excipient or carrier.

50. A formulation according to claim 48 or claim 49, which further comprises a therapeutic agent selected from anti-diabetic agents, hypolipidemic agents, anti-obesity or appetite-regulating agents, anti-hypertensive agents, HDL-increasing agents, cholesterol absorption modulators, Apo-A1 analogues and mimetics, thrombin inhibitors, aldosterone inhibitors, inhibitors of platelet aggregation, estrogen, testosterone, selective estrogen receptor modulators, selective androgen receptor modulators, chemotherapeutic agents, and 5-HT₃ or 5-HT₄ receptor modulators; or pharmaceutically acceptable salts or prodrugs thereof.

51. A formulation according to claim 50, wherein the agent is tegaserod, imatinib, vildagliptin, metformin, a thiazolidone derivative, a sulfonylurea receptor ligand, aliskiren, valsartan, orlistat or a statin, or pharmaceutically acceptable salts or prodrugs.

52. A product comprising a compound of any of claims 1 to 43 and an agent as defined in claim 50; as a combined preparation for simultaneous, separate or sequential use in therapy.

53. A product according to claim 52, wherein the agent is as defined in claim 51.

54. Use of a compound of any of claims 1 to 43 for the manufacture of a medicament for the treatment or prevention of a disease or condition selected from non-insulin-dependent diabetes mellitus, arthritis, obesity, allograft transplantation, calcitonin-osteoporosis, heart failure, impaired glucose metabolism or impaired glucose tolerance, neurodegenerative diseases, cardiovascular or renal diseases, and neurodegenerative or cognitive disorders, hyperglycemia, insulin resistance, lipid disorders, dyslipidemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, low HDL levels, high LDL levels, atherosclerosis, vascular restenosis, irritable bowel syndrome, inflammatory bowel disease, pancreatitis, retinopathy, nephropathy, neuropathy, syndrome X, ovarian hyperandrogenism (polycystic ovarian syndrome), type 2 diabetes, growth hormone deficiency, neutropenia, neuronal disorders, tumor metastasis, benign prostatic hypertrophy, gingivitis, hypertension and osteoporosis.

55. Use according to claim 54, wherein the disease or condition is Alzheimer's disease, Parkinson's disease, Crohn's disease or ulcerative colitis.

- 243 -

56. Use according to claim 54, wherein the disease is diabetic cardiomyopathy, left or right ventricular hypertrophy, hypertrophic medial thickening in arteries and/or in large vessels, mesenteric vasculature hypertrophy or mesangial hypertrophy.

57. Use of a compound of any of claims 1 to 43 for the manufacture of a medicament for producing a sedative or anxiolytic effect, attenuating post-surgical catabolic changes or hormonal responses to stress, reducing mortality and morbidity after myocardial infarction, modulating hyperlipidemia or associated conditions, or lowering VLDL, LDL or Lp(a) levels.

58. A method of treating or preventing a disease or condition in a patient, which comprises administering a therapeutically effective amount of a compound of any of claims 1 to 43.

59. A method according to claim 58, wherein the disease or condition is as defined in any of claims 54 to 56.