



(86) Date de dépôt PCT/PCT Filing Date: 2009/08/07
(87) Date publication PCT/PCT Publication Date: 2010/02/11
(85) Entrée phase nationale/National Entry: 2011/02/03
(86) N° demande PCT/PCT Application No.: US 2009/004543
(87) N° publication PCT/PCT Publication No.: 2010/016935
(30) Priorités/Priorities: 2008/08/07 (US61/188,192);
2008/12/02 (US61/200,628)

(51) Cl.Int./Int.Cl. *A61K 38/16* (2006.01),
A61K 38/22 (2006.01), *A61K 47/48* (2006.01),
A61P 3/04 (2006.01), *A61P 3/10* (2006.01),
A61P 5/48 (2006.01), *C07K 14/575* (2006.01),
C07K 17/08 (2006.01)

(71) Demandeur/Applicant:
IPSEN PHARMA S.A.S., FR

(72) Inventeurs/Inventors:
DONG, ZHENG XIN, US;
SHEN, YEELANA, US;
DEOLIVEIRA, DANIEL B., US

(74) Agent: MOFFAT & CO.

(54) Titre : ANALOGUES TRONQUES DE POLYPEPTIDE INSULINOTROPE GLUCOSE-DEPENDANT
(54) Title: TRUNCATED ANALOGUES OF GLUCOSE-DEPENDENT INSULINOTROPIC POLYPEPTIDE

(57) **Abrégé/Abstract:**

There is provided a novel series of analogues of glucose-dependent insulintropic polypeptide, pharmaceutical compositions containing said compounds, and the use of said compounds as GIP- receptor agonists or antagonists for treatment of GIP-receptor mediated conditions, such as non- insulin dependent diabetes mellitus and obesity.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 February 2010 (11.02.2010)

PCT

(10) International Publication Number
WO 2010/016935 A3

(51) International Patent Classification:
A61K 38/00 (2006.01)

(21) International Application Number:
PCT/US2009/004543

(22) International Filing Date:
7 August 2009 (07.08.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/188,192 7 August 2008 (07.08.2008) US
61/200,628 2 December 2008 (02.12.2008) US

(71) Applicant (for all designated States except US): **IPSEN PHARMA S.A.S.** [FR/FR]; 65, Quai Georges Gorse, F-92100 Boulogne-billancourt (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DONG, Zheng, Xin** [US/US]; 66 Fairview Street, Holliston, MA 01746 (US). **SHEN, Yeelana** [US/US]; 5 Crystal Pond Lane, Franklin, MA 02038 (US). **DEOLIVEIRA, Daniel, B.** [US/US]; 279 Center St., Bellingham, MA 02019 (US).

(74) Agents: **FEENEY, Alan, F.** et al.; Biomeasure, Incorporated, 27 Maple Street, Milford, MA 01757-3650 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,

HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the identity of the inventor (Rule 4.17(i))
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(88) Date of publication of the international search report:
8 April 2010

(54) Title: TRUNCATED ANALOGUES OF GLUCOSE-DEPENDENT INSULINOTROPIC POLYPEPTIDE

(57) Abstract: There is provided a novel series of analogues of glucose-dependent insulinotropic polypeptide, pharmaceutical compositions containing said compounds, and the use of said compounds as GIP- receptor agonists or antagonists for treatment of GIP-receptor mediated conditions, such as non- insulin dependent diabetes mellitus and obesity.



WO 2010/016935 A3

TRUNCATED ANALOGUES OF GLUCOSE-DEPENDENT INSULINOTROPIC POLYPEPTIDE

FIELD OF THE INVENTION

5 The present invention relates to the area of novel analogues of glucose-dependent insulintropic polypeptide, pharmaceutical compositions containing said compounds, and the use of said compounds as GIP-receptor agonists or antagonists for treatment of GIP-receptor mediated conditions, such as non-insulin dependent diabetes mellitus and obesity.

BACKGROUND ART

10 Glucose-dependent insulintropic polypeptide ("GIP", also known as "gastric inhibitory polypeptide") is a 42-residue peptide secreted by enteroendocrine K-cells of the small intestine into the bloodstream in response to oral nutrient ingestion. GIP inhibits the secretion of gastric acid, and it has been shown to be a potent stimulant for the secretion of insulin from pancreatic beta cells after oral
15 glucose ingestion (the "incretin effect") (Creutzfeldt, W., *et al.*, 1979, *Diabetologia*, 16:75-85).

 Insulin release induced by the ingestion of glucose and other nutrients is due to both hormonal and neural factors (Creutzfeldt, W., *et al.*, 1985, *Diabetologia*, 28:565-573). Several gastrointestinal regulatory peptides have been proposed as incretins, and among these candidates, only GIP and glucagon-like peptide 1 ("GLP-1") appear to fulfill the requirements to be considered physiological
20 stimulants of postprandial insulin release (Nauck, *et al.*, 1989, *J. Clin. Endocrinol. Metab.*, 69:654-662). It has been shown that the combined effects of GIP and GLP-1 are sufficient to explain the full incretin effect of the enteroinsular axis (Fehmann, H. C., *et al.*, 1989, *FEBS Lett.*, 252:109-112).

 As is well known to those skilled in the art, the known and potential uses of GIP are varied and multitudinous. Thus, the administration of the compounds of this invention for purposes of
25 eliciting an agonist effect can have the same effects and uses as GIP itself. These varied uses of GIP may be summarized as follows: treating a disease selected from the group consisting of type 1 diabetes, type 2 diabetes (Visboll, T., 2004, *Dan. Med. Bull.*, 51:364-70), insulin resistance (WO 2005/082928), obesity (Green, B. D., *et al.*, 2004, *Current Pharmaceutical Design*, 10:3651-3662), metabolic disorder (Gault, V. A., *et al.*, 2003, *Biochem. Biophys. Res. Commun.*, 308:207-213),
30 central nervous system disease, neurodegenerative disease, congestive heart failure, hypoglycemia, and disorders wherein the reduction of food intake and weight loss are desired. In pancreatic islets, GIP not only enhances insulin secretion acutely, but it also stimulates insulin production through enhancement of proinsulin transcription and translation (Wang, *et al.*, 1996, *Mol. Cell. Endocrinol.*, 116:81-87) and enhances the growth and survival of pancreatic beta cells (Trumper, *et al.*, 2003, *Diabetes*, 52:741-750). In addition to effects on the pancreas to enhance insulin secretion, GIP also
35 has effects on insulin target tissues directly to lower plasma glucose: enhancement of glucose uptake

in adipose (Eckel, *et al.*, 1979, *Diabetes*, 28:1141-1142) and muscle (O'Harte, *et al.*, 1998, *J. Endocrinol.*, 156:237-243), and inhibition of hepatic glucose production (Elahi, D., *et al.*, 1986, *Can. J. Physiol. Pharmacol.*, 65:A18).

In addition, a GIP receptor antagonist in accordance with the present invention inhibits, blocks or reduces glucose absorption from the intestine of an animal. In accordance with this observation, therapeutic compositions containing GIP antagonists may be used in patients with non-insulin dependent diabetes mellitus to improve tolerance to oral glucose in mammals, such as humans, to prevent, inhibit or reduce obesity by inhibiting, blocking or reducing glucose absorption from the intestine of the mammal.

The use of unmodified GIP as a therapeutic, however, is limited by the short *in vivo* half-life of about 2 minutes (Said and Mutt, 1970, *Science*, 169:1217-1218). In serum, both incretins, GIP and GLP-1, are degraded by dipeptidyl peptidase IV ("DPPIV"). Improving the stability of GIP to proteolysis not only maintains the activity of GIP at its receptor but, more importantly, prevents the production of GIP fragments, some of which act as GIP receptor antagonists (Gault, *et al.*, 2002, *J. Endocrinol.*, 175:525-533). Reported modifications have included protection of the N-terminus of GIP from proteolysis by DPPIV through modification of the N-terminal tyrosine (O'Harte, *et al.*, 2002, *Diabetologia*, 45:1281-1291), mutation of the alanine at position 2 (Hinke, *et al.*, 2002, *Diabetes*, 51:656-661), mutation of glutamic acid at position 3 (Gault, *et al.*, 2003, *Biochem. Biophys. Res. Commun.*, 308:207-213), and mutation of alanine at position 13 (Gault, *et al.*, 2003, *Cell Biol. International*, 27:41-46),

The following patent applications have been filed related to the effects of GIP analogues on the function of various target organs and their potential use as therapeutic agents:

PCT publication WO 00/58360 discloses peptidyl analogues of GIP which stimulate the release of insulin. In particular, this application discloses specific peptidyl analogues comprising at least 15 amino acid residues from the N-terminal end of GIP(1-42), *e.g.*, an analogue of GIP containing exactly one amino acid substitution or modification at positions 1, 2 and 3, such as [Pro³]GIP(1-42).

PCT publication WO 98/24464 discloses an antagonist of GIP consisting essentially of a 24-amino acid polypeptide corresponding to positions 7-30 of the sequence of GIP, a method of treating non-insulin dependent diabetes mellitus and a method of improving glucose tolerance in a non-insulin dependent diabetes mellitus patient.


PCT publication WO 03/082898 discloses C-terminal truncated fragments and N-terminal modified analogues of GIP, as well as various GIP analogues with a reduced peptide bond or alterations of the amino acids close to the DPPIV-specific cleavage site. This application further discloses analogues with different linkers between potential receptor binding sites of GIP. The

compounds of this application are alleged to be useful in treating GIP-receptor mediated conditions, such as non-insulin dependent diabetes mellitus and obesity.

There exists a need for improved analogues of GIP, which are stable in formulation and have long plasma half-life *in vivo* resulting from decreased susceptibility to proteolysis and decreased clearance while maintaining binding affinity to a GIP receptor to elicit respective agonistic or antagonistic effects. Moreover, among other therapeutic effects of the compounds of the present invention as illustrated herein, tighter control of plasma glucose levels may prevent long-term diabetic complications, thereby providing an improved quality of life for patients.

SUMMARY OF THE INVENTION

In one aspect, the invention relates to peptide variants of GIP of the following formula (I):



wherein:

A^1 is deleted;

A^2 is Ala, Abu, D-Abu, Acc, Aib, β -Ala, D-Ala, Gaba, Gly, 4Hppa, Ser, D-Ser, Thr, D-Thr, Val, D-Val, or deleted;

A^3 is Glu, Aib, Asp, NMe-Asp, Dhp, Dmt, NMe-Glu, 3Hyp, 4Hyp, 4Ktp, Pro, hPro, Thz, Tic, or deleted;

A^4 is Gly, Acc, Aib, β -Ala, or deleted;

A^5 is Thr, Acc, Aib, Ser, or deleted;

A^6 is Phe, Acc, Aib, Aic, Cha, 1Nal, 2Nal, 2-Pal, 3-Pal, 4-Pal, (X^4, X^5, X^6, X^7, X^8)Phe, Trp, or deleted;

A^7 is Ile, Abu, Acc, Aib, Ala, Cha, Leu, Nle, Phe, Tle, Val, or deleted;

A^8 is Ser, Aib, Chc-Ser, Thr, or deleted;

A^9 is Asp, Aib, Glu, or deleted;

A^{10} is Tyr, Acc, Cha, 1Nal, 2Nal, 2-Pal, 3-Pal, 4-Pal, Phe, (X^4, X^5, X^6, X^7, X^8)Phe, or deleted;

A^{11} is Ser, Acc, Aib, Thr, or deleted;

A^{12} is Ile, Abu, Acc, Aib, Ala, Cha, Leu, Nle, Phe, Tle, Val, or deleted;

A^{13} is Ala, β -Ala, D-Ala, Acc, Aib, Gly, Ser, or deleted;

A^{14} is Met, Abu, Acc, Aib, Ala, Cha, Ile, Leu, Nle, Phe, Tle, or Val;

A^{15} is Asp, Aib, or Glu;

A^{16} is Lys, Amp, Apc, Arg, hArg, Orn, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-

NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃);

A¹⁷ is Ile, Abu, Acc, Aib, Ala, Cha, Leu, Nle, Phe, Tle, or Val;

A¹⁸ is His, Amp, Arg, 2-Pal, 3-Pal, 4-Pal, Phe, or Tyr;

5 A¹⁹ is Gln, Aib, or Asn;

A²⁰ is Gln, Aib, or Asn;

A²¹ is Asp, Aib, or Glu;

A²² is Phe, Acc, Aib, Aic, Cha, 1Nal, 2Nal, 2-Pal, 3-Pal, 4-Pal, (X⁴,X⁵,X⁶,X⁷,X⁸)Phe or Trp;

A²³ is Val, Abu, Acc, Aib, Ala, Cha, Ile, Leu, Nle, or Tle;

10 A²⁴ is Asn, Aib, or Gln;

A²⁵ is Trp, Acc, Aib, 1Nal, 2Nal, 2-Pal, 3-Pal, 4-Pal, Phe, or (X⁴,X⁵,X⁶,X⁷,X⁸)Phe;

A²⁶ is Leu, Acc, Aib, Cha, Ile, Nle, Phe, (X⁴,X⁵,X⁶,X⁷,X⁸)Phe, or Tle;

A²⁷ is Leu, Acc, Aib, Cha, Ile, Nle, Phe, (X⁴,X⁵,X⁶,X⁷,X⁸)Phe or Tle;

A²⁸ is Ala, Acc, or Aib;

15 A²⁹ is Gln, Aib, Asn, or deleted;

A³⁰ is Lys, Amp, Apc, Arg, hArg, Orn, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

20 A³¹ is Gly, Aib, Acc, β-Ala, 2Nal, D-2Nal, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

25 A³² is Lys, Amp, Apc, Arg, hArg, Orn, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

30 A³³ is Lys, Amp, Apc, Arg, hArg, Orn, Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

35 A³⁴ is Asn, Aib, Gln, Ser, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-

CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A³⁵ is Asp, Aib, Glu, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl),

5 hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A³⁶ is Trp, Acc, Aib, 1Nal, 2Nal, 2-Pal, 3-Pal, 4-Pal, Phe, (X⁴,X⁵,X⁶,X⁷,X⁸)Phe, HN-

10 CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

15 A³⁷ is Lys, Amp, Apc, Arg, hArg, Orn, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

20 A³⁸ is His, Amp, Phe, 2-Pal, 3-Pal, 4-Pal, Tyr, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

25 CH₃), or deleted;

A³⁹ is Asn, Aib, Gln, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl),

hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

30 C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A⁴⁰ is Ile, Acc, Aib, Ser, Thr, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl),

hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

35 C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A⁴¹ is Thr, Acc, Aib, Asn, Gln, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl),

hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-

CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A⁴² is Gln, Aib, Acc, Asn, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl),

5 hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A⁴³ is Acc, Aib, Ala, Asp, Gln, Phe, Thr, Trp, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O),

10 Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

15 R¹ is OH, NH₂, (C₁-C₃₀)alkoxy, or NH-X²-CH₂-Z⁰, wherein X² is a (C₀-C₃₀)hydrocarbon moiety, and Z⁰ is H, OH, CO₂H or CONH₂;

each of R², R³, R⁴ and R⁵ is independently selected from the group consisting of H, (C₁-C₃₀)alkyl, (C₁-C₃₀)heteroalkyl, (C₁-C₃₀)acyl, (C₂-C₃₀)alkenyl, (C₂-C₃₀)alkynyl, aryl(C₁-C₃₀)alkyl, aryl(C₁-C₃₀)acyl, substituted (C₁-C₃₀)alkyl, substituted (C₁-C₃₀)heteroalkyl, substituted (C₁-C₃₀)acyl, substituted (C₂-C₃₀)alkenyl, substituted (C₂-C₃₀)alkynyl, substituted aryl(C₁-C₃₀)alkyl, and substituted aryl(C₁-C₃₀)acyl; provided that when R² is (C₁-C₃₀)acyl, aryl(C₁-C₃₀)acyl, substituted (C₁-C₃₀)acyl, or substituted aryl(C₁-C₃₀)acyl, then R³ is H, (C₁-C₃₀)alkyl, (C₁-C₃₀)heteroalkyl, (C₂-C₃₀)alkenyl, (C₂-C₃₀)alkynyl, aryl(C₁-C₃₀)alkyl, substituted (C₁-C₃₀)alkyl, substituted (C₁-C₃₀)heteroalkyl, substituted (C₂-C₃₀)alkenyl, substituted (C₂-C₃₀)alkynyl, or substituted aryl(C₁-C₃₀)alkyl; further provided that

25 when R⁴ is (C₁-C₃₀)acyl, aryl(C₁-C₃₀)acyl, substituted (C₁-C₃₀)acyl, or substituted aryl(C₁-C₃₀)acyl, then R⁵ is H, (C₁-C₃₀)alkyl, (C₁-C₃₀)heteroalkyl, (C₂-C₃₀)alkenyl, (C₂-C₃₀)alkynyl, aryl(C₁-C₃₀)alkyl, substituted (C₁-C₃₀)alkyl, substituted (C₁-C₃₀)heteroalkyl, substituted (C₂-C₃₀)alkenyl, substituted (C₂-C₃₀)alkynyl, or substituted aryl(C₁-C₃₀)alkyl;

n is, independently for each occurrence, an integer from 1 to 5 inclusive;

30 s, t, x and y each is, independently for each occurrence, an integer from 1 to 30 inclusive;

X⁴, X⁵, X⁶, X⁷ and X⁸ each is, independently for each occurrence, H, F, Cl, Br, I, (C₁₋₁₀)alkyl, substituted (C₁₋₁₀)alkyl, aryl, substituted aryl, OH, NH₂, NO₂, or CN; and

provided that when A² is 4Hppa, then R² and R³ are deleted.

A subset (A) of the compounds covered by the above formula (I) are those in which:

35 A² is Ala, 4Hppa, or deleted;

A³ is Glu, 4Hyp, Pro, or deleted;

- A⁴ is Gly or deleted;
 A⁵ is Thr or deleted;
 A⁶ is Phe or deleted;
 A⁷ is Ile, A5c, A6c, or deleted;
 5 A⁸ is Ser, Chc-Ser, or deleted;
 A⁹ is Asp or deleted;
 A¹⁰ is Tyr or deleted;
 A¹¹ is Ser, A6c, Aib, or deleted;
 A¹² is Ile or deleted;
 10 A¹³ is Ala, Aib, or deleted;
 A¹⁴ is Met, A6c, or Nle;
 A¹⁵ is Asp;
 A¹⁶ is Lys;
 A¹⁷ is Ile;
 15 A¹⁸ is His;
 A¹⁹ is Gln;
 A²⁰ is Gln;
 A²¹ is Asp;
 A²² is Phe;
 20 A²³ is Val;
 A²⁴ is Asn;
 A²⁵ is Trp;
 A²⁶ is Leu;
 A²⁷ is Leu;
 25 A²⁸ is Ala;
 A²⁹ is Gln;
 A³⁰ is Lys;
 A³¹ is Gly, Cys(Hsu), Cys(Psu), 2Nal, D-2Nal, Orn(N-C(O)-(CH₂)₄-CH₃), Orn(N-C(O)-
 (CH₂)₈-CH₃), Orn(N-C(O)-(CH₂)₁₂-CH₃), or deleted;
 30 A³² is Lys, Cys(Psu), or deleted;
 A³³ is Lys, Cys(Psu), or deleted;
 A³⁴ is Asn, Cys(Psu), Orn(N-C(O)-(CH₂)₈-CH₃), or deleted;
 A³⁵ is Asp, Cys(Psu), or deleted;
 A³⁶ is Trp, Cys(Psu), or deleted;
 35 A³⁷ is Lys, Cys(Psu), or deleted;
 A³⁸ is His, Cys(Psu), or deleted;
 A³⁹ is Asn, Cys(Psu), or deleted;

A⁴⁰ is Ile, Cys(Psu), or deleted;

A⁴¹ is Thr or deleted;

A⁴² is Gln or deleted;

A⁴³ is Gln or deleted; and

5 provided that at least one of A², A³, A⁴, A⁵, A⁶, A⁷, A⁸, A⁹, A¹¹, A¹³, A¹⁴, A³¹, A³², A³³, A³⁴, A³⁵, A³⁶, A³⁷, A³⁸, A³⁹ and A⁴⁰ is not the amino acid residue of the corresponding position of the native GIP.

A subset of the compounds of the preceding subset (A) are those in which:

A² is deleted;

10 A³ is deleted;

A⁴ is deleted;

A⁵ is deleted;

A⁶ is deleted;

A⁷ is A5c or A6c;

15 A⁸ is Ser;

A⁹ is Asp;

A¹⁰ is Tyr;

A¹¹ is Ser;

A¹² is Ile;

20 A¹³ is Ala;

A¹⁴ is Met;

A³¹ is Gly, Cys(Hsu), Cys(Psu), 2Nal, D-2Nal, Orn(N-C(O)-(CH₂)₄-CH₃), Orn(N-C(O)-(CH₂)₈-CH₃), or Orn(N-C(O)-(CH₂)₁₂-CH₃);

A³² is Lys or Cys(Psu);

25 A³³ is Lys or Cys(Psu);

A³⁴ is Asn, Cys(Psu), or Orn(N-C(O)-(CH₂)₈-CH₃);

A³⁵ is Asp or Cys(Psu);

A³⁶ is Trp or Cys(Psu);

A³⁷ is Lys or Cys(Psu);

30 A³⁸ is His or Cys(Psu);

A³⁹ is Asn or Cys(Psu);

A⁴⁰ is Ile or Cys(Psu);

A⁴¹ is Thr;

A⁴² is Gln; and

35 A⁴³ is deleted.

A subset of the compounds of the preceding subset (A) are those in which A⁴³ is deleted, A² is 4Hppa, and at least one of A³, A⁷, A¹¹, A¹³ and A¹⁴ is not the amino acid residue of the corresponding position of the native GIP.

Another subset of the compounds of the preceding subset (A) are those in which A² to A⁵ and
5 A³¹ to A⁴³ are deleted, and at least one of A⁶, A⁷, A¹¹ and A¹⁴ is not the amino acid residue of the corresponding position of the native GIP.

Another subset of the compounds of the preceding subset (A) are those in which A² to A⁷ and A⁴³ are deleted, and at least one of A⁸ and A³¹ is not the amino acid residue of the corresponding position of the native GIP.

10 Another subset of the compounds of the preceding subset (A) are those in which A² to A⁵ and A⁴³ are deleted, and at least one of A⁶, A⁷ and A³¹ is not the amino acid residue of the corresponding position of the native GIP.

Another subset of the compounds of the preceding subset (A) are those in which A² to A⁵ and A³² to A⁴³ are deleted, and at least one of A⁶, A⁷ and A³¹ is not the amino acid residue of the
15 corresponding position of the native GIP.

Preferred compounds of formula (I) are:

- Example 1: (Ac-A6c⁷)hGIP(7-42)-OH (SEQ ID NO:4);
 Example 2: [Ac-A6c⁷, Cys(Psu)⁴⁰]hGIP(7-42)-OH (SEQ ID NO:5);
 Example 3: [Ac-A6c⁷, Cys(Psu)³⁹]hGIP(7-42)-OH (SEQ ID NO:6);
 20 Example 4: [Ac-A6c⁷, Cys(Psu)³⁸]hGIP(7-42)-OH (SEQ ID NO:7);
 Example 5: [Ac-A6c⁷, Cys(Psu)³⁶]hGIP(7-42)-OH (SEQ ID NO:8);
 Example 6: [Ac-A6c⁷, Cys(Psu)³⁵]hGIP(7-42)-OH (SEQ ID NO:9);
 Example 7: [Ac-A6c⁷, Cys(Psu)³⁴]hGIP(7-42)-OH (SEQ ID NO:10);
 Example 8: [Ac-A6c⁷, Cys(Psu)³³]hGIP(7-42)-OH (SEQ ID NO:11);
 25 Example 9: [Ac-A6c⁷, Cys(Psu)³²]hGIP(7-42)-OH (SEQ ID NO:12);
 Example 10: [Ac-A6c⁷, Cys(Psu)³¹]hGIP(7-42)-OH (SEQ ID NO:13);
 Example 11: [Ac-A6c⁷, Cys(Psu)³⁷]hGIP(7-42)-OH (SEQ ID NO:14);
 Example 12: [Ac-A6c⁷, Orn³¹(N-C(O)-(CH₂)₁₂-CH₃)]hGIP(7-42)-OH (SEQ ID NO:15);
 Example 13: [Ac-A6c⁷, Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:16);
 30 Example 14: [A6c⁷, Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:17);
 Example 15: [CH₃-(CH₂)₈-C(O)-A6c⁷, Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH
 (SEQ ID NO:18);
 Example 16: [Ac-A6c⁷, Orn³¹(N-C(O)-(CH₂)₄-CH₃)]hGIP(7-42)-OH (SEQ ID NO:19);
 Example 17: [A6c⁷, Orn³¹(N-C(O)-(CH₂)₄-CH₃)]hGIP(7-42)-OH (SEQ ID NO:20);
 35 Example 18: [CH₃-(CH₂)₄-C(O)-A6c⁷, Orn³¹(N-C(O)-(CH₂)₄-CH₃)]hGIP(7-42)-OH

- (SEQ ID NO:21);
- Example 19: [Ac-A6c⁷, Orn³⁴(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:22);
- Example 20: [A6c⁷, Orn³⁴(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:23);
- Example 21: [CH₃-(CH₂)₈-C(O)-A6c⁷, Orn³⁴(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH
 5 (SEQ ID NO:24);
- Example 22: [Ac-A6c⁷, Cys(Hsu)³¹]hGIP(7-42)-OH (SEQ ID NO:25);
- Example 23: [A6c⁷, Cys(Hsu)³¹]hGIP(7-42)-OH (SEQ ID NO:26);
- Example 24: (Ac-A6c⁷, 2Nal³¹)hGIP(7-42)-OH (SEQ ID NO:27);
- Example 25: (Ac-A6c⁷, D-2Nal³¹)hGIP(7-42)-OH;
- 10 Example 26: (Ac-4Hyp³, A6c⁷)hGIP(3-42)-OH (SEQ ID NO:28);
- Example 27: (Ac-A6c⁷, Gln⁴³)hGIP(7-43)-OH (SEQ ID NO:29);
- Example 28: [Ac-A6c⁷, Cys(Psu)³¹]hGIP(7-34)-NH₂ (SEQ ID NO:30);
- Example 29: [Ac-A6c⁷, Cys(Psu)³¹]hGIP(7-31)-NH₂ (SEQ ID NO:31);
- Example 30: [Ac-Phe⁶, A6c⁷, Cys(Psu)³¹]hGIP(6-42)-OH (SEQ ID NO:32);
- 15 Example 31: [A6c⁷, Cys(Psu)³¹]hGIP(6-42)-OH (SEQ ID NO:33);
- Example 32: (Ac-Phe⁶, A6c⁷)hGIP(6-30)-NH₂ (SEQ ID NO:34);
- Example 33: [Ac-Phe⁶, A6c⁷, Cys(Psu)³¹]hGIP(6-31)-NH₂ (SEQ ID NO:35);
- Example 34: [A6c⁷, Cys(Psu)³¹]hGIP(6-31)-NH₂ (SEQ ID NO:36);
- Example 35: (A5c⁷, Nle¹⁴)hGIP(6-30)-NH₂ (SEQ ID NO:37);
- 20 Example 36: (A6c⁷, Nle¹⁴)hGIP(6-30)-NH₂ (SEQ ID NO:38);
- Example 37: (Aib¹¹, Nle¹⁴)hGIP(6-30)-NH₂ (SEQ ID NO:39);
- Example 38: [Ac-Asp⁹, Cys(Psu)³³]hGIP(9-42)-OH (SEQ ID NO:40);
- Example 39: [Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(8-42)-OH (SEQ ID NO:41);
- Example 40: [Chc-Ser⁸, Cys(Psu)³¹]hGIP(8-42)-OH (SEQ ID NO:42);
- 25 Example 41: [CH₃-(CH₂)₄-C(O)-Ser⁸, Cys(Psu)³¹]hGIP(8-42)-OH (SEQ ID NO:43);
- Example 42: (4Hppa², 4Hyp³, A6c⁷)hGIP(2-42)-OH (SEQ ID NO:44);
- Example 43: (4Hppa², Pro³, Nle¹⁴)hGIP(2-42)-OH (SEQ ID NO:45);
- Example 44: (4Hppa², Aib¹³)hGIP(2-42)-OH (SEQ ID NO:46);
- Example 45: (4Hppa², A6c¹⁴)hGIP(2-42)-OH (SEQ ID NO:47);
- 30 Example 46: (4Hppa², A6c¹¹)hGIP(2-42)-OH (SEQ ID NO:48); and
- Example 47: [Aib², A5c¹¹, Nle¹⁴, Lys⁴³(N-C(O)-(CH₂)₁₀-CH₃)]hGIP(2-43)-OH (SEQ ID NO:49).

According to another aspect of the present invention, a compound according to the present invention as summarized hereinabove and claimed in the appended claims may further comprise a covalently linked PEG moiety, in which said PEG moiety is covalently linked to the compound via a
 35 Cys(maleimide), hCys(maleimide), or Pen(maleimide) linker, to form Cys(succinimide-N-PEG), hCys(succinimide-N-PEG), or Pen(succinimide-N-PEG), wherein "succinimide-N-PEG" is either

linear or branched as defined hereinbelow. Such PEG moiety has average molecular weight of from about 2,000 to about 80,000, and preferably such PEG moiety is selected from the group consisting of 5K PEG, 10K PEG, 20K PEG, 30K PEG, 40K PEG, 50K PEG, and 60K PEG, to form

Cys(succinimide-N-5K PEG), Cys(succinimide-N-10K PEG), Cys(succinimide-N-20K PEG),
 5 Cys(succinimide-N-30K PEG), Cys(succinimide-N-40K PEG), Cys(succinimide-N-50K PEG),
 Cys(succinimide-N-60K PEG), hCys(succinimide-N-5K PEG), hCys(succinimide-N-10K PEG),
 hCys(succinimide-N-20K PEG), hCys(succinimide-N-30K PEG), hCys(succinimide-N-40K PEG),
 hCys(succinimide-N-50K PEG), hCys(succinimide-N-60K PEG), Pen(succinimide-N-5K PEG),
 Pen(succinimide-N-10K PEG), Pen(succinimide-N-20K PEG), Pen(succinimide-N-30K PEG),
 10 Pen(succinimide-N-40K PEG), Pen(succinimide-N-50K PEG), or Pen(succinimide-N-60K PEG).

PEGylation occurs at any one of amino acid residue positions 16, 30, and 31-43, and preferably at any one of amino acid residue positions 32, 33 and 43, whereby Cys(succinimide-N-PEG), hCys(succinimide-N-PEG), or Pen(succinimide-N-PEG) is placed in any one of such amino acid residue positions.

15 Further, the above formula (I) may be expanded to provide PEGylation sites at positions A⁴⁴-A⁴⁷. The C-terminus of such PEGylated compounds of the present invention may be amidated, *e.g.*, (Ac-A6c⁷)hGIP(7-42)-NH₂ (SEQ ID NO:50), or it may remain as free acid, *e.g.*, (Ac-A6c⁷)hGIP(7-42)-OH (SEQ ID NO:4).

20 DETAILED DESCRIPTION OF THE INVENTION

The application employs the following commonly understood abbreviations:

	Abu:	α -aminobutyric acid
	Acc:	1-amino-1-cyclo(C ₃ -C ₉)alkyl carboxylic acid
25	A3c:	1-amino-1-cyclopropanecarboxylic acid
	A4c:	1-amino-1-cyclobutanecarboxylic acid
	A5c:	1-amino-1-cyclopentanecarboxylic acid
	A6c:	1-amino-1-cyclohexanecarboxylic acid
	Act:	4-amino-4-carboxytetrahydropyran
30	Ado:	12-aminododecanoic acid
	Aib:	α -aminoisobutyric acid
	Aic:	2-aminoindan-2-carboxylic acid
	Ala or A:	alanine
	β -Ala:	beta-alanine
35	Amp:	4-amino-phenylalanine;
	Apc:	4-amino-4-carboxypiperidine:

	Arg or R:	arginine
	hArg:	homoarginine
	Asn or N:	asparagine
	Asp or D:	aspartic acid
5	Aun:	11-aminoundecanoic acid
	Ava:	5-aminovaleric acid
	Cha:	β -cyclohexylalanine
	Chc:	cyclohexyl carboxylic acid
	Cys or C:	cysteine
10	D-Ala:	D-alanine
	Dhp:	3,4-dehydroproline
	Dmt:	5,5-dimethylthiazolidine-4-carboxylic acid
	Gaba:	γ -aminobutyric acid
	Gln or Q:	glutamine
15	Glu or E:	glutamic acid
	Gly or G:	glycine
	His or H:	histidine
	4Hppa:	3-(4-hydroxyphenyl)propionic acid
	Hsu:	N-hexylsuccinimide
20	3Hyp:	3-hydroxyproline
	4Hyp:	4-hydroxyproline
	hPro:	homoproline
	Ile or I:	isoleucine
	4Ktp:	4-ketoproline
25	Leu or L:	leucine
	Lys or K:	lysine
	Met or M:	methionine
	Nle:	norleucine
	NMe-Tyr:	N-methyl-tyrosine
30	1Nal or 1-Nal:	β -(1-naphthyl)alanine
	2Nal or 2-Nal:	β -(2-naphthyl)alanine
	Nle:	norleucine
	Nva:	norvaline
	Orn:	ornithine
35	2Pal or 2-Pal:	β -(2-pyridinyl)alanine
	3Pal or 3-Pal:	β -(3-pyridinyl)alanine
	4Pal or 4-Pal:	β -(4-pyridinyl)alanine

	Pen:	penicillamine
	Phe or F:	phenylalanine
	(3,4,5F)Phe:	3,4,5-trifluorophenylalanine
	(2,3,4,5,6)Phe:	2,3,4,5,6-pentafluorophenylalanine
5	Pro or P:	proline
	Psu:	N-propylsuccinimide
	Ser or S:	serine
	Taz:	β -(4-thiazolyl)alanine
	3Thi:	β -(3-thienyl)alanine
10	Thr or T:	threonine
	Thz:	thioprolin
	Tic:	tetrahydroisoquinoline-3-carboxylic acid
	Tle:	tert-leucine
	Trp or W:	tryptophan
15	Tyr or Y:	tyrosine
	Val or V:	valine

Certain other abbreviations used herein are defined as follows:

	Act:	acetonitrile
20	Boc:	<i>tert</i> -butoxycarbonyl
	BSA:	bovine serum albumin
	DCM:	dichloromethane
	DIPEA:	diisopropylethyl amine
	DMF:	dimethylformamide
25	DTT:	dithiothreitol
	ESI:	electrospray ionization
	Fmoc:	9-fluorenylmethoxycarbonyl
	HBTU:	2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
	HOBT:	1-hydroxybenzotriazole
30	HPLC:	high performance liquid chromatography
	IBMX:	isobutylmethylxanthine
	LC-MS:	liquid chromatography-mass spectrometry
	Mtt:	methyltrityl
	NMP:	N-methylpyrrolidone
35	5K PEG:	polyethylene glycol, which may include other functional groups or moieties

such as a linker, and which is either linear or branched as defined hereinbelow, with an average total molecular weight of about 5,000

10K PEG: polyethylene glycol, which may include other functional groups or moieties such as a linker, and which is either linear or branched as defined hereinbelow, with an average total molecular weight of about 10,000

20K PEG: polyethylene glycol, which may include other functional groups or moieties such as a linker, and which is either linear or branched as defined hereinbelow, with an average total molecular weight of about 20,000

30K PEG: polyethylene glycol, and which is either linear or branched as defined hereinbelow, which may include other functional groups or moieties such as a linker, with an average total molecular weight of about 30,000

40K PEG: polyethylene glycol, which may include other functional groups or moieties such as a linker, and which is either linear or branched as defined hereinbelow, with an average total molecular weight of about 40,000

50K PEG: polyethylene glycol, which may include other functional groups or moieties such as a linker, and which is either linear or branched as defined hereinbelow, with an average total molecular weight of about 50,000

60K PEG: polyethylene glycol, which may include other functional groups or moieties such as a linker, and which is either linear or branched as defined hereinbelow, with an average total molecular weight of about 60,000


tBu: *tert*-butyl

TIS: triisopropylsilane


Trt: trityl

TFA: trifluoro acetic acid


Z: benzyloxycarbonyl




“Cys(succinimide-N-alkyl)” has the structure of:



“Cys(Hsu)” has the structure of:




“Cys(Psu)” has the structure of:



“Orn(N-C(O)-(CH₂)₁₂-CH₃)” has the structure of:


“Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃)” has the structure of:

5




wherein, $x = 1-30$, and $y = 1-30$.

“hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃)” has the structure of:




10 wherein, $x = 1-30$, and $y = 1-30$.

“Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃)” has the structure of:




wherein, $x = 1-30$, and $y = 1-30$.

“Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃)” has the structure of:




wherein, $s = 1-30$, and $t = 1-30$.

“hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃)” has the structure of:




wherein $s = 1-30$, and $t = 1-30$.


“Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃)” has the structure of:




wherein $s = 1-30$, and $t = 1-30$.



“Cys(succinimide-N-PEG)” has the structure of:




“hCys(succinimide-N-PEG)” has the structure of:




“Pen(succinimide-N-PEG)” has the structure of:

“Cys(succinimide-N-(CH₂)₂-C(O)NH-(CH₂)₃-PEG)” has the structure of:



“Cys(succinimide-N-(CH₂)₂-C(O)NH-(CH₂)₃-O-CH₂-CH(PEG)-CH₂-PEG)” has the structure of:



With the exception of the N-terminal amino acid, all abbreviations (*e.g.*, Ala) of amino acids in this disclosure stand for the structure of -NH-C(R)(R')-CO-, wherein R and R' each is, independently, hydrogen or the side chain of an amino acid (*e.g.*, R = CH₃ and R' = H for Ala), or R and R' may be joined to form a ring system. For the N-terminal amino acid, the abbreviation stands for the structure of (R²R³)N-C(R)(R')-CO-, wherein R² and R³ are as defined in the above formula (I).

The term “(C₁-C₃₀)hydrocarbon moiety” encompasses alkyl, alkenyl and alkynyl, and in the case of alkenyl and alkynyl there are C₂-C₃₀.

A peptide of this invention is also denoted herein by another format, *e.g.*, (A5c²)hGIP(1-42)-OH (SEQ ID NO:3), with the substituted amino acids from the natural sequence placed between the brackets (*e.g.*, A5c² for Ala² in hGIP). The numbers between the parentheses refer to the number of amino acids present in the peptide (*e.g.*, hGIP(1-42)-OH (SEQ ID NO:1) is amino acids 1 through 42 of the peptide sequence for hGIP). The designation “NH₂” in hGIP(1-30)-NH₂ (SEQ ID NO:2) indicates that the C-terminus of the peptide is amidated; hGIP(1-42) (SEQ ID NO:1) or hGIP(1-42)-OH (SEQ ID NO:1) means that the C-terminus is the free acid.

Human GIP (“hGIP”) has the amino acid sequence of:

Tyr	Ala	Glu	Gly	Thr	Phe	Ile	Ser	Asp	Tyr	Ser	Ile	Ala	Met	Asp	Lys	Ile	His	Gln	Gln	Asp	Phe	Val
1				5					10					15					20			
Asn	Trp	Leu	Leu	Ala	Gln	Lys	Gly	Lys	Lys	Asn	Asp	Trp	Lys	His	Asn	Ile	Thr	Gln				
		25			30				35					40								

(SEQ ID NO:1)

“Acyl” refers to R"-C(O)-, where R" is H, alkyl, substituted alkyl, heteroalkyl, substituted heteroalkyl, alkenyl, substituted alkenyl, aryl, alkylaryl, or substituted alkylaryl.

“Alkyl” refers to a hydrocarbon group containing one or more carbon atoms, where multiple carbon atoms if present are joined by single bonds. The alkyl hydrocarbon group may be straight-

chain or contain one or more branches or cyclic groups.

“Substituted alkyl” refers to an alkyl wherein one or more hydrogen atoms of the hydrocarbon group are replaced with one or more substituents selected from the group consisting of halogen, (*i.e.*, fluorine, chlorine, bromine, and iodine), -OH, -CN, -SH, -NH₂, -NHCH₃, -NO₂, -C₁₋₂₀ alkyl substituted with halogens, -CF₃, -OCH₃, -OCF₃, and -(CH₂)₀₋₂₀-COOH. In different embodiments 1, 2, 3 or 4 substituents are present. The presence of -(CH₂)₀₋₂₀-COOH results in the production of an alkyl acid. Examples of alkyl acids containing, or consisting of, -(CH₂)₀₋₂₀-COOH include 2-norbornane acetic acid, *tert*-butyric acid and 3-cyclopentyl propionic acid.

“Heteroalkyl” refers to an alkyl wherein one or more of the carbon atoms in the hydrocarbon group are replaced with one or more of the following groups: amino, amido, -O-, -S- or carbonyl. In different embodiments 1 or 2 heteroatoms are present.

“Substituted heteroalkyl” refers to a heteroalkyl wherein one or more hydrogen atoms of the hydrocarbon group are replaced with one or more substituents selected from the group consisting of halogen, -OH, -CN, -SH, -NH₂, -NHCH₃, -NO₂, -C₁₋₂₀ alkyl substituted with halogens, -CF₃, -OCH₃, -OCF₃, and -(CH₂)₀₋₂₀-COOH. In different embodiments 1, 2, 3 or 4 substituents are present.

“Alkenyl” refers to a hydrocarbon group made up of two or more carbons wherein one or more carbon-carbon double bonds are present. The alkenyl hydrocarbon group may be straight-chain or contain one or more branches or cyclic groups.

“Substituted alkenyl” refers to an alkenyl wherein one or more hydrogens are replaced with one or more substituents selected from the group consisting of halogen, -OH, -CN, -SH, -NH₂, -NHCH₃, -NO₂, -C₁₋₂₀ alkyl substituted with halogens, -CF₃, -OCH₃, -OCF₃, and -(CH₂)₀₋₂₀-COOH. In different embodiments 1, 2, 3 or 4 substituents are present.

“Aryl” refers to an optionally substituted aromatic group with at least one ring having a conjugated pi-electron system, containing up to three conjugated or fused ring systems. Aryl includes carbocyclic aryl, heterocyclic aryl and biaryl groups. Preferably, the aryl is a 5 or 6 membered ring. Preferred atoms for a heterocyclic aryl are one or more sulfur, oxygen, and/or nitrogen. Examples of aryl include phenyl, 1-naphthyl, 2-naphthyl, indole, quinoline, 2-imidazole, and 9-anthracene. Aryl substituents are selected from the group consisting of -C₁₋₂₀ alkyl, -C₁₋₂₀ alkoxy, halogen, -OH, -CN, -SH, -NH₂, -NO₂, -C₁₋₂₀ alkyl substituted with halogens, -CF₃, -OCF₃, and -(CH₂)₀₋₂₀-COOH. In different embodiments the aryl contains 0, 1, 2, 3, or 4 substituents.

“Alkylaryl” refers to an “alkyl” joined to an “aryl”.

Synthesis

The peptides of this invention can be prepared by standard solid phase peptide synthesis. See, *e.g.*, Stewart, J. M., *et al.*, 1984, *Solid Phase Synthesis*, Pierce Chemical Co., 2d ed. If R^1 is $NH-X^2-CH_2-CONH_2$, *i.e.*, $Z^0 = CONH_2$, the synthesis of the peptide starts with Fmoc- $NH-X^2-CH_2-CONH_2$ which is coupled to Rink amide MBHA resin. If R^1 is $NH-X^2-CH_2-COOH$, *i.e.*, $Z^0 = COOH$, the synthesis of the peptide starts with Fmoc- $NH-X^2-CH_2-COOH$ which is coupled to Wang resin. For this particular step, 2 molar equivalents of Fmoc- $NH-X^2-COOH$, HBTU and HOBt and 10 molar equivalents of DIPEA are used. The coupling time is about 8 hours.

In the synthesis of a GIP analogue of this invention containing A5c, A6c, and/or Aib, the coupling time is 2 hrs for these residues and the residue immediately following them.

The substituents R^2 and R^3 of the above generic formula can be attached to the free amine of the N-terminal amino acid A^1 by standard methods known in the art. For example, alkyl groups, *e.g.*, (C_1-C_{30}) alkyl, can be attached using reductive alkylation. Hydroxyalkyl groups, *e.g.*, (C_1-C_{30}) hydroxyalkyl, can also be attached using reductive alkylation wherein the free hydroxy group is protected with a *tert*-butyl ester. Acyl groups, *e.g.*, $-C(O)X^3$, can be attached by coupling the free acid, *e.g.*, $-X^3COOH$, to the free amine of the N-terminal amino acid by mixing the completed resin with 3 molar equivalents of both the free acid and diisopropylcarbodiimide in methylene chloride for about one hour. If the free acid contains a free hydroxy group, *e.g.*, 3-fluoro-4-hydroxyphenylacetic acid, then the coupling should be performed with an additional 3 molar equivalents of HOBt.

The following examples describe synthetic methods for making a peptide of this invention, which methods are well-known to those skilled in the art. Other methods are also known to those skilled in the art. The examples are provided for the purpose of illustration and are not meant to limit the scope of the present invention in any manner.

Example 2: [Ac-A6c⁷, Cys(Psu)⁴⁰]hGIP(7-42)-OH

Solid-phase peptide synthesis was used to assemble the peptide using microwave-assisted Fmoc Chemistry on a Liberty Peptide Synthesizer (CEM; Matthews, NC, USA) at the 0.1 mmole scale. Pre-loaded Fmoc-Gln(Trt)-Wang resin (0.59 mmole/g; Novabiochem, San Diego, CA, USA) was used to generate the C-terminal acid peptide. The resin (0.17 g) was placed in a 50 ml conical tube along with 15 ml of dimethylformamide (DMF) and loaded onto a resin position on the synthesizer. The resin was then quantitatively transferred to the reaction vessel via the automated process. The standard Liberty synthesis protocol for 0.1 mmole scale synthesis was used. This protocol involves deprotecting the N-terminal Fmoc moiety via an initial treatment with 7 ml of 20% piperidine, containing 0.1M N-hydroxybenzotriazole (HOBt), in DMF. The initial deprotection step was for 30 seconds with microwave power (45 watts, maximum temperature of 75 °C), and nitrogen bubbling (3 seconds on / 7 seconds off). The reaction vessel was then drained and a second piperidine

treatment, identical to the first treatment, except that it was for a 3-minute duration. The resin was then drained and thoroughly washed with DMF several times. The protected amino acid, Fmoc-Thr(tBu)-OH, prepared as 0.2M stock solution in DMF, was then added (2.5 ml, 5 equivalents), followed by 1.0 ml of 0.45M (4.5 eq.) HBTU [2-(1H-benzo-triazole-1-yl)-1,1,3,3-tetramethyluronium
 5 hexafluorophosphate] in DMF. This was followed by the addition of 0.5 ml of 2M (10 eq.) DIPEA (diisopropylethylamine) in NMP (N-methylpyrrolidinone). The coupling step was performed for 5 minutes using 20 watts of microwave power, a max temperature of 75 °C, and the same rate of nitrogen bubbling.

Following the initial coupling step, the reaction vessel was drained to waste and the coupling
 10 step repeated. Cycle 2 was then initiated similar to cycle 1. All amino acids were introduced similarly and a double-coupling strategy was employed throughout the entire sequence. Cycles 1-3, 19-20, 25-26, and 30-34 contained a capping procedure immediately following the coupling step. Capping was performed by adding 7 ml of 0.5M acetic anhydride, containing 0.015M HOBT in NMP, along with 2 ml of the 2M DIPEA solution using a multi-step microwave protocol: 50 watts of power
 15 for 30 seconds (65 °C max temperature), followed by 30 seconds of microwave power off, followed by a second round of 30 seconds of microwave power on (50 watts), and then again 30 seconds of no microwave power. The resin was then drained and thoroughly washed with DMF. The following amino acids (Advanced Chemtech; Louisville, KY, USA) were used: Cycle 1: Fmoc-Thr(OtBu)-OH; Cycle 2: Fmoc-Cys(Trt)-OH; Cycle 3: Fmoc-Asn(Trt)-OH; Cycle 4: Fmoc-His(Trt)-OH; Cycle 5:
 20 Fmoc-Lys(Boc)-OH; Cycle 6: Fmoc-Trp(Boc)-OH; Cycle 7: Fmoc-Asp(OtBu)-OH; Cycle 8: Fmoc-Asn(Trt)-OH; Cycle 9: Fmoc-Lys(Boc)-OH; Cycle 10: Fmoc-Lys(Boc)-OH; Cycle 11: Fmoc-Gly-OH; Cycle 12: Fmoc-Lys(Boc)-OH; Cycle 13: Fmoc-Gln(Trt)-OH; Cycle 14: Fmoc-Ala-OH; Cycle 15: Fmoc-Leu-OH; Cycle 16: Fmoc-Leu-OH; Cycle 17: Fmoc-Trp(Boc)-OH; Cycle 18: Fmoc-Asn(Trt)-OH; Cycle 19: Fmoc-Val-OH; Cycle 20: Fmoc-Phe-OH; Cycle 21: Fmoc-Asp(OtBu)-OH; Cycle 22:
 25 Fmoc-Gln(Trt)-OH; Cycle 23: Fmoc-Gln(Trt)-OH; Cycle 24: Fmoc-His(Trt)-OH; Cycle 25: Fmoc-Ile-OH; Cycle 26: Fmoc-Lys(Boc)-OH; Cycle 27: Fmoc-Asp(OtBu)-OH; Cycle 28: Fmoc-Met-OH; Cycle 29: Fmoc-Ala-OH; Cycle 30: Fmoc-Ile-OH; Cycle 31: Fmoc-Tyr(tBu)-Ser(psiMe,Me,Pro)-OH; Cycle 32: Fmoc-Asp(OtBu)-OH; Cycle 33: Fmoc-Ser(tBu)-OH; and Cycle 34: Fmoc-A6c-OH. Once the peptide backbone was complete, the resin was treated with piperidine solution to remove the N-
 30 terminal Fmoc group, followed by treatment with the standard capping procedure in order to acetylate the N-terminus. The resin was then thoroughly washed with DMF and then transferred back to the 50 ml conical tube using DMF as the transfer solvent.

The resin was deprotected and cleaved from the resin via treatment with 5 ml of the following reagent; 5% TIS, 2% water, 5% (w/v) dithiothreitol (DTT), 88% TFA, and allowed to mix for 3.5
 35 hours. The filtrate was collected into 45 ml of cold anhydrous ethyl ether. The precipitate was pelleted for 10 minutes at 3500 RPM in a refrigerated centrifuge. The ether was decanted, and the

peptide re-suspended in fresh ether. The ether workup was performed a total of 2 times. Following the last ether wash the peptide was allowed to air dry to remove residual ether. The peptide pellet was resuspended in 8 ml of acetonitrile (Acn) followed by 8ml of de-ionized water, and allowed to fully dissolve. The peptide solution was then analyzed by mass spectrometry. Mass analysis employing electro-spray ionization identified a main product containing a mass of 4358.0 Daltons; corresponding to the acetylated, linear product. The crude product (approximately 500 mg) was analysed by HPLC, employing a 250 x 4.6 mm C18 column (Phenomenex; Torrance, CA, USA) using a gradient of 2-80% acetonitrile (0.1% TFA) over 30 minutes. Analytical HPLC identified a product with 38% purity. The crude peptide was then purified on a preparative HPLC equipped with a C18 reverse phase column using a 10-60% acetonitrile (0.1% TFA) over 50 minutes at a 10 ml/min flowrate. The purified peptide was then lyophilized yielding 15 mg of peptide. The linear peptide was then derivatized with N-propylmaleimide (Pma) to generate the propylsuccinimide (Psu) derivative on the Cysteine side chain. The purified linear peptide was brought up in water, adjusted to pH 6.5 with ammonium carbonate, at 5mg/ml. Five equivalents of Pma was added with constant stirring for 30 seconds. The derivatized peptide solution was then analyzed by mass spectrometry. Mass analysis identified a main product containing a mass of 4498.6 Daltons; corresponding to the desired Psu derivatized product. The product was then re-purified via preparative HPLC using a similar gradient as before. The purified product was analyzed by HPLC for purity (95.2%) and mass spectrometry (4498.6 Daltons) and subsequently lyophilized. Following lyophilization, 4.3 mg of purified product was obtained representing a 1% yield.

Example 12: [Ac-A6c⁷, Orn(N-C(O)-(CH₂)₁₂-CH₃)³¹]hGIP(7-42)-OH

Solid-phase peptide synthesis was used to assemble the peptide using microwave-assisted Fmoc Chemistry on a Liberty Peptide Synthesizer (CEM; Matthews, NC, USA) at the 0.1 mmole scale. Pre-loaded Fmoc-Gln(Trt)-Wang resin (0.59 mmole/g; Novabiochem, San Diego, CA, USA) was used to generate the C-terminal acid peptide. The resin (0.17 g) was placed in a 50 ml conical tube along with 15 ml of dimethylformamide (DMF) and loaded onto a resin position on the synthesizer. The resin was then quantitatively transferred to the reaction vessel via the automated process. The standard Liberty synthesis protocol for 0.1 mmole scale synthesis was used. This protocol involves deprotecting the N-terminal Fmoc moiety via an initial treatment with 7 ml of 20% piperidine, containing 0.1M N-hydroxybenzotriazole (HOBt), in DMF. The initial deprotection step was for 30 seconds with microwave power (45 watts, maximum temperature of 75 °C), and nitrogen bubbling (3 seconds on / 7 seconds off). The reaction vessel was then drained and a second piperidine treatment, identical to the first treatment, except that it was for a 3-minute duration. The resin was then drained and thoroughly washed with DMF several times. The protected amino acid, Fmoc-Thr(tBu)-OH, prepared as 0.2M stock solution in DMF, was then added (2.5 ml, 5 eq.), followed by 1.0 ml of 0.45M (4.5 eq.) HBTU [2-(1H-benzo-triazole-1-yl)-1,1,3,3-tetramethyluronium

hexafluorophosphate] in DMF. This was followed by the addition of 0.5 ml of 2M (10 eq.) DIPEA (diisopropylethylamine) in NMP (N-methylpyrrolidinone). The coupling step was performed for 5 minutes using 20 watts of microwave power, a max temperature of 75 °C, and the same rate of nitrogen bubbling.

5 Following the initial coupling step the reaction vessel was drained to waste and the coupling step repeated. Cycle 2 was then initiated similar to cycle 1. All amino acids were introduced similarly and a double-coupling strategy was employed throughout the entire sequence. Cycles 1-3, 19-20, 25-26, and 30-34 contained a capping procedure immediately following the coupling step. Capping was performed by adding 7ml of 0.5M acetic anhydride, containing 0.015M HOBt in NMP, 10 along with 2 ml of the 2M DIPEA solution using a multi-step microwave protocol: 50 watts of power for 30 seconds (65 °C max temperature), followed by 30 seconds of microwave power off, followed by a second round of 30 seconds of microwave power on (50 watts), and then again 30 seconds of no microwave power. The resin was then drained and thoroughly washed with DMF. The following amino acids (Advanced Chemtech; Louisville, KY, USA) were used; Cycle 1: Fmoc-Thr(tBu)-OH; 15 Cycle 2: Fmoc-Ile-OH; Cycle 3: Fmoc-Asn(Trt)-OH; Cycle 4: Fmoc-His(Trt)-OH; Cycle 5: Fmoc-Lys(Boc)-OH; Cycle 6: Fmoc-Trp(Boc)-OH; Cycle 7: Fmoc-Asp(OtBu)-OH; Cycle 8: Fmoc-Asn(Trt)-OH; Cycle 9: Fmoc-Lys(Boc)-OH; Cycle 10: Fmoc-Lys(Boc)-OH; Cycle 11: Fmoc-Orn(Mtt)-OH; Cycle 12: Fmoc-Lys(Boc)-OH; Cycle 13: Fmoc-Gln(Trt)-OH; Cycle 14: Fmoc-Ala-OH; Cycle 15: Fmoc-Leu-OH; Cycle 16: Fmoc-Leu-OH; Cycle 17: Fmoc-Trp(Boc)-OH; Cycle 18: 20 Fmoc-Asn(Trt)-OH; Cycle 19: Fmoc-Val-OH; Cycle 20: Fmoc-Phe-OH; Cycle 21: Fmoc-Asp(OtBu)-OH; Cycle 22: Fmoc-Gln(Trt)-OH; Cycle 23: Fmoc-Gln(Trt)-OH; Cycle 24: Fmoc-His(Trt)-OH; Cycle 25: Fmoc-Ile-OH; Cycle 26: Fmoc-Lys(Boc)-OH; Cycle 27: Fmoc-Asp(OtBu)-OH; Cycle 28: Fmoc-Met-OH; Cycle 29: Fmoc-Ala-OH; Cycle 30: Fmoc-Ile-OH; Cycle 31: Fmoc-Tyr(tBu)-Ser(psiMe,Me,Pro)-OH; Cycle 32: Fmoc-Asp(OtBu)-OH; Cycle 33: Fmoc-Ser(tBu)-OH; and Cycle 25 34: Fmoc-A6c-OH. The coupling protocol for Fmoc-His(Trt)-OH was a slightly modified version of the standard protocol. The microwave power was off for the first 2 minutes, followed by 4 minutes with microwave power on (20 watts; max temperature of 50 °C). Once the peptide backbone was complete, the resin was treated with piperidine solution to remove the N-terminal Fmoc group, followed by treatment with the standard capping procedure in order to acetylate the N-terminus. The 30 resin was then treated with 12 ml of 1% trifluoroacetic acid (TFA) / 5% triisopropylsilane (TIS) in dichloromethane (DCM) for 5 minutes and a N₂ sparge rate of 5 seconds on and 10 seconds off. The resin was then drained and again treated with the 1% TFA / 5% TIS in DCM solution for 5 minutes. This was performed a total of 7 times to effectively remove the Mtt moiety from the Ornithine side chain. The resin was thoroughly washed with DCM several times, and then treated with the standard 35 piperidine treatment in order to neutralize residual TFA salt on the δN of ornithine. Myristic acid, (CH₃-(CH₂)₁₂-COOH; Aldrich, St. Louis, MO, USA) prepared as a 0.2M solution in DMF, was

coupled to the ornithine side chain using the standard amino acid coupling protocol. The resin was then thoroughly washed with DMF and then transferred back to the 50 ml conical tube using DMF as the transfer solvent.

The resin was deprotected and cleaved from the resin via treatment with 5 ml of the following reagent; 5% TIS, 2% water, 5% (w/v) dithiothreitol (DTT), 88% TFA, and allowed to mix for 3.5 hours. The filtrate was collected into 45 ml of cold anhydrous ethyl ether. The precipitate was pelleted for 10 minutes at 3500 RPM in a refrigerated centrifuge. The ether was decanted, and the peptide re-suspended in fresh ether. The ether workup was performed a total of 2 times. Following the last ether wash the peptide was allowed to air dry to remove residual ether. The peptide pellet was resuspended in 8 ml of acetonitrile (Acn) followed by 8ml of de-ionized water, and allowed to fully dissolve. The peptide solution was then analyzed by mass spectrometry. Mass analysis employing electrospray ionization identified a main product containing a mass of 4636.5 Daltons; corresponding to the desired product. The crude product was analysed by HPLC, employing a 250 x 4.6 mm C18 column (Phenomenex; Torrance, CA, USA) using a gradient of 2-80% acetonitrile (0.1% TFA) over 30 minutes. Analytical HPLC identified a product with 37% purity. The peptide was then purified on a preparative HPLC equipped with a C18 column using a similar elution gradient. The purified product was re-analyzed by HPLC for purity (95.20%) and mass spectrometry (4636.6 Daltons) and subsequently lyophilized. Following lyophilization, 3 mg of purified product was obtained representing a 0.6% yield.

Example 22: [Ac-A6c⁷, Cys(Hsu)³¹]hGIP(7-42)-OH

Solid-phase peptide synthesis was used to assemble the peptide using microwave-assisted Fmoc Chemistry on a Liberty Peptide Synthesizer (CEM; Matthews, NC, USA) at the 0.1 mmole scale. Pre-loaded Fmoc-Gln(Trt)-Wang resin (0.59 mmole/g; Novabiochem, San Diego, CA, USA) was used to generate the C-terminal acid peptide. The resin (0.17 g) was placed in a 50 ml conical tube along with 15 ml of dimethylformamide (DMF) and loaded onto a resin position on the synthesizer. The resin was then quantitatively transferred to the reaction vessel via the automated process. The standard Liberty synthesis protocol for 0.1 mmole scale synthesis was used. This protocol involves deprotecting the N-terminal Fmoc moiety via an initial treatment with 7 ml of 20% piperidine, containing 0.1M N-hydroxybenzotriazole (HOBt), in DMF. The initial deprotection step was for 30 seconds with microwave power (45 watts, maximum temperature of 75°C), and nitrogen bubbling (3 seconds on / 7 seconds off). The reaction vessel was then drained and a second piperidine treatment, identical to the first treatment, except that it was for a 3-minute duration. The resin was then drained and thoroughly washed with DMF several times. The protected amino acid, Fmoc-Thr(tBu)-OH, prepared as 0.2M stock solution in DMF, was then added (2.5 ml, 5 eq.), followed by 1.0 ml of 0.45M (4.5 eq.) HBTU [2-(1H-benzo-triazole-1-yl)-1,1,3,3-tetramethyluronium

hexafluorophosphate] in DMF. This was followed by the addition of 0.5 ml of 2M (10 eq.) DIPEA (diisopropylethylamine) in NMP (N-methylpyrrolidinone). The coupling step was performed for 5 minutes using 20 watts of microwave power, a max temperature of 75°C, and the same rate of nitrogen bubbling.

5 Following the initial coupling step the reaction vessel was drained to waste and the coupling step repeated. Cycle 2 was then initiated similar to cycle 1. All amino acids were introduced similarly and a double-coupling strategy was employed throughout the entire sequence. Cycles 1-3, 19-20, 25-26, and 30-34 contained a capping procedure immediately following the coupling step. Capping was performed by adding 7 ml of 0.5M acetic anhydride, containing 0.015M HOBT in NMP, 10 along with 2 ml of the 2M DIPEA solution using a multi-step microwave protocol: 50 watts of power for 30 seconds (65 °C max temperature), followed by 30 seconds of microwave power off, followed by a second round of 30 seconds of microwave power on (50 watts), and then again 30 seconds of no microwave power. The resin was then drained and thoroughly washed with DMF. The following amino acids (Advanced Chemtech, Louisville, KY, USA) were used: Cycle 1: Fmoc-Thr(OtBu)-OH; 15 Cycle 2: Fmoc-Ile-OH; Cycle 3: Fmoc-Asn(Trt)-OH; Cycle 4: Fmoc-His(Trt)-OH; Cycle 5: Fmoc-Lys(Boc)-OH; Cycle 6: Fmoc-Trp(Boc)-OH; Cycle 7: Fmoc-Asp(OtBu)-OH; Cycle 8: Fmoc-Asn(Trt)-OH; Cycle 9: Fmoc-Lys(Boc)-OH; Cycle 10: Fmoc-Lys(Boc)-OH; Cycle 11: Fmoc-Cys(Trt)-OH; Cycle 12: Fmoc-Lys(Boc)-OH; Cycle 13: Fmoc-Gln(Trt)-OH; Cycle 14: Fmoc-Ala-OH; Cycle 15: Fmoc-Leu-OH; Cycle 16: Fmoc-Leu-OH; Cycle 17: Fmoc-Trp(Boc)-OH; Cycle 18: Fmoc-Asn(Trt)-OH; Cycle 19: Fmoc-Val-OH; Cycle 20: Fmoc-Phe-OH; Cycle 21: Fmoc-Asp(OtBu)-OH; 20 Cycle 22: Fmoc-Gln(Trt)-OH; Cycle 23: Fmoc-Gln(Trt)-OH; Cycle 24: Fmoc-His(Trt)-OH; Cycle 25: Fmoc-Ile-OH; Cycle 26: Fmoc-Lys(Boc)-OH; Cycle 27: Fmoc-Asp(OtBu)-OH; Cycle 28: Fmoc-Met-OH; Cycle 29: Fmoc-Ala-OH; Cycle 30: Fmoc-Ile-OH; Cycle 31: Fmoc-Tyr(tBu)-Ser(psiMe,Me,Pro)-OH; Cycle 32: Fmoc-Asp(OtBu)-OH; Cycle 33: Fmoc-Ser(tBu)-OH; and Cycle 25 34: Fmoc-A6c-OH. Once the peptide backbone was complete, the resin was treated with piperidine solution to remove the N-terminal Fmoc group, followed by treatment with the standard capping procedure in order to acetylate the N-terminus. The resin was then thoroughly washed with DMF and then transferred back to the 50 ml conical tube using DMF as the transfer solvent.

30 The resin was deprotected and cleaved from the resin via treatment with 5 ml of the following reagent; 5% TIS, 2% water, 5% (w/v) dithiothreitol (DTT), 88% TFA, and allowed to mix for 3.5 hours. The filtrate was collected into 45 ml of cold anhydrous ethyl ether. The precipitate was pelleted for 10 minutes at 3500 RPM in a refrigerated centrifuge. The ether was decanted, and the peptide re-suspended in fresh ether. The ether workup was performed a total of 2 times. Following the last ether wash the peptide was allowed to air dry to remove residual ether. The peptide pellet was 35 resuspended in 8 ml of acetonitrile (Acn) followed by 8 ml of de-ionized water, and allowed to fully dissolve. The peptide solution was then analyzed by mass spectrometry. Mass analysis employing

electrospray ionization identified a main product containing a mass of 4414.9 Daltons; corresponding to the linear product. The crude product (approximately 500 mg) was analysed by HPLC, employing a 250 x 4.6 mm C18 column (Phenomenex; Torrance, CA, USA) using a gradient of 2-80% acetonitrile (0.1% TFA) over 30 minutes. Analytical HPLC identified a product with 58% purity.

- 5 The crude peptide was then derivatized with N-hexylmaleimide (Hma) to generate the hexylsuccinimide (Hsu) derivative on the Cysteine side chain. The crude linear peptide was brought up in water, adjusted to pH 6.5 with ammonium carbonate, at 5 mg/ml. Five equivalents of Hma was added with constant stirring for 30 seconds. Excess Hma was quenched using 5 eq. of dithiothreitol (DTT). The derivatized peptide solution was then analyzed by mass spectrometry. Mass analysis
- 10 identified a main product containing a mass of 4596.1 Daltons; corresponding to the desired Hsu derivatized product. The product was then re-purified via preparative HPLC using a similar gradient as before. The purified product was analyzed by HPLC for purity (95.4%) and mass spectrometry (4596.4 Daltons) and subsequently lyophilized. Following lyophilization, 28.1 mg of purified product was obtained representing a 6.1% yield.

- 15 The PEGylated GIP compounds disclosed herein can be synthesized substantially according to the procedure described for the synthesis of the compound of Example 2, by using PEG-maleimide as the starting material instead of N-propylmaleimide used in Example 2.

- Other peptides of the invention can be prepared by a person of ordinary skill in the art using synthetic procedures analogous to those disclosed in the foregoing examples. Physical data for the
- 20 compounds exemplified herein are given in Table 1.

TABLE 1

Example Number	Mol. Wt. (Expected)	Mol. Wt. (ESI-MS)	% Purity (HPLC)
1	4368.92	4368.8	96.70
2	4498.06	4498.6	95.20
3	4497.12	4497.5	99.90
4	4474.08	4473.9	99.90
5	4425.01	4425.0	99.90
6	4496.13	4496.5	99.90
7	4497.12	4496.7	99.90
8	4483.05	4482.4	99.90
9	4483.05	4482.7	99.90
10	4554.17	4554.0	99.90
11	4483.05	4482.7	98.40
12	4636.38	4636.6	95.20
13	4580.27	4580.7	96.70
14	4538.23	4538.8	99.90
15	4692.48	4693.1	95.00
16	4524.16	4524.9	96.20
17	4482.13	4482.6	99.90
18	4580.27	4580.9	95.70
19	4523.22	4523.7	95.50

20	4481.18	4481.4	99.90
21	4635.43	4636.0	95.30
22	4596.25	4596.4	95.40
23	4554.21	4554.6	95.80
24	4509.11	4509.8	99.10
25	4509.11	4509.9	99.90
26	4787.37	4788.4	99.90
27	4497.05	4496.8	99.90
28	3530.07	3530.0	99.90
29	3159.62	3159.6	96.40
30	4701.35	4701.7	96.40
31	4659.31	4660.0	95.10
32	3064.50	3064.7	99.90
33	3306.80	3306.7	96.30
34	3264.76	3264.6	98.20
35	2990.40	2990.8	96.83
36	3004.42	3004.7	99.90
37	2990.44	2990.8	97.20
38	4270.80	4270.6	99.90
39	4413.06	4413.5	99.90
40	4497.12	4497.6	96.90
41	4485.11	4485.7	96.40
42	4893.49	4894.4	99.90
43	4847.45	4848.1	99.90
44	4911.51	4911.4	99.90
45	4891.45	4891.0	99.90
46	4935.58	4935.8	99.90
47	5150.9	5151.4	99.9

Functional Assays

A. In Vitro hGIP Receptor Binding Assay

5

10

15

Membranes for *in vitro* receptor binding assays were prepared by homogenizing the CHO-K1 clonal cells expressing the human recombinant GIP receptor, with a Brinkman Polytron (setting 6, 15 sec), in ice-cold 50 mM Tris-HCl and then subjected to two centrifugations at 39,000 g for 10 minutes, with a resuspension in fresh buffer in between. For the assay, aliquots of the washed membrane preparations were incubated (100 minutes at 25°C with 0.05 nM [¹²⁵I]GIP (approximately 2200 Ci/mmol) in 50mM Tris-HCl, 0.1mg/ml bacitracin, and 0.1% BSA. The final assay volume was 0.5 ml. The incubations were terminated by rapid filtration through GF/C filters (pre-soaked in 0.5% polyethylenimine) using a Brandel filtration manifold. Each tube and filter were then washed three times with 5-ml aliquots of ice-cold buffer. Specific binding was defined as the total radioligand bound minus that bound in the presence of 1000 nM GIP. *In vitro* hGIP receptor binding data for the compounds exemplified herein are given in Table 2.

B. Human and Rat Plasma Half-Life Assay

GIP peptide (50 μ L 1 mg/ml) was added to 450 μ L plasma (human or rat), vortexed briefly and incubated at 37°C. 50 μ L was removed at various times, like at 0, 1, 2, 3, 4, 8, 24, 32, 48, 56, 72
 5 hours, mixed with 5 μ L formic acid and 150 μ L acetonitrile in a microcentrifuge tube, vortexed, and centrifuged for 10 minutes at 10K rpm. The supernatant was transferred to an injection vial and analyzed by LC-MS. The LC-MS system consisted of an API4000 mass spectrometer with an ESI probe. Positive ion mode and full scan detection were used. HPLC separation was carried out on a Luna 3 μ C8 (2), 2 x 30 mm column with a gradient from 90% A to 90% B in 10 minutes at a flow rate
 10 of 0.3 ml/min. Buffer A was 1% formic acid in water and buffer B was 1% formic acid acetonitrile. Human and rat plasma half-life data for the compounds exemplified herein are given in Table 2.

TABLE 2

Example Number	Ki (nM)	Human Plasma T $\frac{1}{2}$ (hr)	Rat Plasma T $\frac{1}{2}$ (hr)
1	N/A	>72	11.0
2	532.19	13.7	7.2
3	75.73	16.4	5.6
4	332.97	10.0	6.0
5	442.49	7.8	8.5
6	486.41	8.0	3.2
7	735.40	7.9	1.6
8	416.57	N/A	N/A
9	686.96	N/A	N/A
10	963.06	8.0	2.6
11	127.00	N/A	N/A
12	178.00	7.3	>72
13	N/A	17.8	19.0
14	N/A	4.1	15.3
15	N/A	5.2	>72
16	N/A	>50	30.0
17	N/A	>50	9.4
18	N/A	13.8	6.3
19	N/A	18.9	10.7
20	274.00	9.4	13.9
21	163.50	8.0	>72
22	N/A	7.7	5.0
23	772.00	11.7	5.3
24	194.33	26.3	12.2
25	159.39	>50	13.7
26	546.10	30.1	17.5
27	7.92	N/A	N/A
28	114.78	8.8	4.1
29	48.32	12.2	7.0
30	574.00	7.2	7.6
31	277.01	4.4	5.1
32	68.54	24.1	60.3
33	77.48	13.6	10.2

34	101.42	11.3	7.4
35	734.33	16.5	23.0
36	212.33	21.9	21.6
37	170.00	13.5	28.5
38	472.00	N/A	N/A
39	N/A	31.1	13.7
40	403.33	4.1	6.9
41	205.48	5.9	3.7
42	293.90	>51	10.3
43	800.01	>53	2.3
44	12.89	48.8	9.9
45	50.43	64.2	4.3
46	91.78	30.8	11.9
47	N/A	N/A	N/A

C. Determination of cyclic AMP stimulation

1 x 10⁵ CHO-K1 cells expressing the human recombinant GIP receptor or RIN-5F insulinoma cells were seeded overnight into 24-well cell culture plates (Corning Incorporate, Corning, NY, USA). For the assay, the cells were preincubated in 500 µl of Hanks balanced salt solution (Sigma, St. Louis, MO, USA) with 0.55 mM IBMX (Sigma, St. Louis, MO, USA) adjusted to pH 7.3 for 10 minutes. GIP or its analogs was then added at a concentration of 100 nM. Following a 30-minute incubation at 37 °C, the plates were placed on ice and 500 µl of ice-cold absolute ethanol was added to stop the reaction. The contents of the wells were collected, spun at 2,700 g for 20 minutes at 4 °C to remove cellular debris. The cAMP levels in the supernatants were determined by radioimmunoassay (New England Nuclear, Boston, MA, USA).

Administration

The peptides of this invention can be provided in the form of pharmaceutically acceptable salts. Examples of such salts include, but are not limited to, those formed with organic acids (*e.g.*, acetic, lactic, maleic, citric, malic, ascorbic, succinic, benzoic, methanesulfonic, toluenesulfonic, or pamoic acid), inorganic acids (*e.g.*, hydrochloric acid, sulfuric acid, or phosphoric acid), and polymeric acids (*e.g.*, tannic acid, carboxymethyl cellulose, polylactic, polyglycolic, or copolymers of polylactic-glycolic acids). A typical method of making a salt of a peptide of the present invention is well known in the art and can be accomplished by standard methods of salt exchange. Accordingly, the TFA salt of a peptide of the present invention (the TFA salt results from the purification of the peptide by using preparative HPLC, eluting with TFA containing buffer solutions) can be converted into another salt, such as an acetate salt by dissolving the peptide in a small amount of 0.25 N acetic acid aqueous solution. The resulting solution is applied to a semi-prep HPLC column (Zorbax, 300 SB, C-8). The column is eluted with (1) 0.1N ammonium acetate aqueous solution for 0.5 hrs, (2) 0.25N acetic acid aqueous solution for 0.5 hrs, and (3) a linear gradient (20% to 100% of solution B over 30 minutes) at a flow rate of 4 ml/min (solution A is 0.25N acetic acid aqueous solution; solution

B is 0.25N acetic acid in acetonitrile/water, 80:20). The fractions containing the peptide are collected and lyophilized to dryness.

The dosage of active ingredient in the compositions of this invention may be varied; however, it is necessary that the amount of the active ingredient be such that a suitable dosage form is obtained.

5 The selected dosage depends upon the desired therapeutic effect, on the route of administration, and on the duration of the treatment. In general, an effective dosage for the activities of this invention is in the range of 1×10^{-7} to 200 mg/kg/day, preferably 1×10^{-4} to 100 mg/kg/day, which can be administered as a single dose or divided into multiple doses.

10 The compounds of this invention can be administered by oral, parenteral (*e.g.*, intramuscular, intraperitoneal, intravenous or subcutaneous injection, or implant), nasal, vaginal, rectal, sublingual, or topical routes of administration, and can be formulated with pharmaceutically acceptable carriers to provide dosage forms appropriate for each route of administration.

15 Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound is admixed with at least one inert pharmaceutically acceptable carrier such as sucrose, lactose, or starch. Such dosage forms can also comprise, as is normal practice, additional substances other than such inert diluents, *e.g.*, lubricating agents such as magnesium stearate. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

20 Liquid dosage forms for oral administration include, without limitation, pharmaceutically acceptable emulsions, solutions, suspensions, syrups, elixirs, and the like, containing inert diluents commonly used in the art, such as water. Besides such inert diluents, compositions can also include adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring and perfuming agents.

25 Preparations according to this invention for parenteral administration include, without limitation, sterile aqueous or non-aqueous solutions, suspensions, emulsions, and the like. Examples of non-aqueous solvents or vehicles include propylene glycol, polyethylene glycol, vegetable oils, such as olive oil and corn oil, gelatin, and injectable organic esters such as ethyl oleate. Such dosage forms may also contain adjuvants such as preserving, wetting, emulsifying, and dispersing agents. They may be sterilized by, for example, filtration through a bacteria-retaining filter, by incorporating
30 sterilizing agents into the compositions, by irradiating the compositions, or by heating the compositions. They can also be manufactured in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use.

Compositions for rectal or vaginal administration are preferably suppositories which may contain, in addition to the active substance, excipients such as coca butter or a suppository wax.

Compositions for nasal or sublingual administration are also prepared with standard excipients well known in the art.


Further, a compound of this invention can be administered in a sustained release composition such as those described in the following patents and patent applications. U.S. Patent No. 5,672,659 teaches sustained release compositions comprising a bioactive agent and a polyester. U.S. Patent No. 5,595,760 teaches sustained release compositions comprising a bioactive agent in a gelable form. U.S. Patent No. 5,821,221 teaches polymeric sustained release compositions comprising a bioactive agent and chitosan. U.S. Patent No. 5,916,883 teaches sustained release compositions comprising a bioactive agent and cyclodextrin. PCT publication WO99/38536 teaches absorbable sustained release compositions of a bioactive agent. PCT publication WO00/04916 teaches a process for making microparticles comprising a therapeutic agent such as a peptide in an oil-in-water process.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Also, all publications, patent applications, patents and other references mentioned herein are hereby incorporated by reference, each in its entirety.

CLAIMS

What is claimed is:

5 1. A compound of formula (I),



wherein:

10 A^1 is deleted;

A^2 is Ala, Abu, D-Abu, Acc, Aib, β -Ala, D-Ala, Gaba, Gly, 4Hppa, Ser, D-Ser, Thr, D-Thr, Val, D-Val, or deleted;

A^3 is Glu, Aib, Asp, NMe-Asp, Dhp, Dmt, NMe-Glu, 3Hyp, 4Hyp, 4Ktp, Pro, hPro, Thz, Tic, or deleted;

15 A^4 is Gly, Acc, Aib, β -Ala, or deleted;

A^5 is Thr, Acc, Aib, Ser, or deleted;

A^6 is Phe, Acc, Aib, Aic, Cha, 1Nal, 2Nal, 2-Pal, 3-Pal, 4-Pal, (X^4, X^5, X^6, X^7, X^8)Phe, Trp, or deleted;

A^7 is Ile, Abu, Acc, Aib, Ala, Cha, Leu, Nle, Phe, Tle, Val, or deleted;

20 A^8 is Ser, Aib, Chc-Ser, Thr, or deleted;

A^9 is Asp, Aib, Glu, or deleted;

A^{10} is Tyr, Acc, Cha, 1Nal, 2Nal, 2-Pal, 3-Pal, 4-Pal, Phe, (X^4, X^5, X^6, X^7, X^8)Phe, or deleted;

A^{11} is Ser, Acc, Aib, Thr, or deleted;

A^{12} is Ile, Abu, Acc, Aib, Ala, Cha, Leu, Nle, Phe, Tle, Val, or deleted;

25 A^{13} is Ala, β -Ala, D-Ala, Acc, Aib, Gly, Ser, or deleted;

A^{14} is Met, Abu, Acc, Aib, Ala, Cha, Ile, Leu, Nle, Phe, Tle, or Val;

A^{15} is Asp, Aib, or Glu;

A^{16} is Lys, Amp, Apc, Arg, hArg, Orn, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃);

A^{17} is Ile, Abu, Acc, Aib, Ala, Cha, Leu, Nle, Phe, Tle, or Val;

A^{18} is His, Amp, Arg, 2-Pal, 3-Pal, 4-Pal, Phe, or Tyr;

35 A^{19} is Gln, Aib, or Asn;

A^{20} is Gln, Aib, or Asn;

A^{21} is Asp, Aib, or Glu;

A²² is Phe, Acc, Aib, Aic, Cha, 1Nal, 2Nal, 2-Pal, 3-Pal, 4-Pal, (X⁴,X⁵,X⁶,X⁷,X⁸)Phe or Trp;

A²³ is Val, Abu, Acc, Aib, Ala, Cha, Ile, Leu, Nle, or Tle;

A²⁴ is Asn, Aib, or Gln;

A²⁵ is Trp, Acc, Aib, 1Nal, 2Nal, 2-Pal, 3-Pal, 4-Pal, Phe, or (X⁴,X⁵,X⁶,X⁷,X⁸)Phe;

5 A²⁶ is Leu, Acc, Aib, Cha, Ile, Nle, Phe, (X⁴,X⁵,X⁶,X⁷,X⁸)Phe, or Tle;

A²⁷ is Leu, Acc, Aib, Cha, Ile, Nle, Phe, (X⁴,X⁵,X⁶,X⁷,X⁸)Phe or Tle;

A²⁸ is Ala, Acc, or Aib;

A²⁹ is Gln, Aib, Asn, or deleted;

10 A³⁰ is Lys, Amp, Apc, Arg, hArg, Orn, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

15 A³¹ is Gly, Aib, Acc, β-Ala, 2Nal, D-2Nal, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

20 A³² is Lys, Amp, Apc, Arg, hArg, Orn, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

25 A³³ is Lys, Amp, Apc, Arg, hArg, Orn, Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

30 A³⁴ is Asn, Aib, Gln, Ser, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

35 A³⁵ is Asp, Aib, Glu, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-

(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A³⁶ is Trp, Acc, Aib, 1Nal, 2Nal, 2-Pal, 3-Pal, 4-Pal, Phe, (X⁴,X⁵,X⁶,X⁷,X⁸)Phe, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A³⁷ is Lys, Amp, Apc, Arg, hArg, Orn, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A³⁸ is His, Amp, Phe, 2-Pal, 3-Pal, 4-Pal, Tyr, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A³⁹ is Asn, Aib, Gln, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A⁴⁰ is Ile, Acc, Aib, Ser, Thr, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A⁴¹ is Thr, Acc, Aib, Asn, Gln, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A⁴² is Gln, Aib, Acc, Asn, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O), Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Pen(succinimide-N-(CH₂)_x-C(O)-NH-

(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), or deleted;

A⁴³ is Acc, Aib, Ala, Asp, Gln, Phe, Thr, Trp, HN-CH((CH₂)_n-N(R⁴R⁵))-C(O),

Cys(succinimide-N-alkyl), hCys(succinimide-N-alkyl), Pen(succinimide-N-alkyl), Cys(succinimide-

5 N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), hCys(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃),

Pen(succinimide-N-(CH₂)_x-C(O)-NH-(CH₂)_y-CH₃), Cys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃),

hCys(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-CH₃), Pen(succinimide-N-(CH₂)_s-NH-C(O)-(CH₂)_t-

CH₃), or deleted;

10 R¹ is OH, NH₂, (C₁-C₃₀)alkoxy, or NH-X²-CH₂-Z⁰, wherein X² is a (C₀-C₃₀)hydrocarbon moiety, and Z⁰ is H, OH, CO₂H or CONH₂;

each of R², R³, R⁴ and R⁵ is independently selected from the group consisting of H, (C₁-C₃₀)alkyl, (C₁-C₃₀)heteroalkyl, (C₁-C₃₀)acyl, (C₂-C₃₀)alkenyl, (C₂-C₃₀)alkynyl, aryl(C₁-C₃₀)alkyl, aryl(C₁-C₃₀)acyl, substituted (C₁-C₃₀)alkyl, substituted (C₁-C₃₀)heteroalkyl, substituted (C₁-C₃₀)acyl, substituted (C₂-C₃₀)alkenyl, substituted (C₂-C₃₀)alkynyl, substituted aryl(C₁-C₃₀)alkyl, and substituted

15 aryl(C₁-C₃₀)acyl; provided that when R² is (C₁-C₃₀)acyl, aryl(C₁-C₃₀)acyl, substituted (C₁-C₃₀)acyl, or substituted aryl(C₁-C₃₀)acyl, then R³ is H, (C₁-C₃₀)alkyl, (C₁-C₃₀)heteroalkyl, (C₂-C₃₀)alkenyl, (C₂-C₃₀)alkynyl, aryl(C₁-C₃₀)alkyl, substituted (C₁-C₃₀)alkyl, substituted (C₁-C₃₀)heteroalkyl, substituted (C₂-C₃₀)alkenyl, substituted (C₂-C₃₀)alkynyl, or substituted aryl(C₁-C₃₀)alkyl; further provided that

20 when R⁴ is (C₁-C₃₀)acyl, aryl(C₁-C₃₀)acyl, substituted (C₁-C₃₀)acyl, or substituted aryl(C₁-C₃₀)acyl, then R⁵ is H, (C₁-C₃₀)alkyl, (C₁-C₃₀)heteroalkyl, (C₂-C₃₀)alkenyl, (C₂-C₃₀)alkynyl, aryl(C₁-C₃₀)alkyl, substituted (C₁-C₃₀)alkyl, substituted (C₁-C₃₀)heteroalkyl, substituted (C₂-C₃₀)alkenyl, substituted (C₂-C₃₀)alkynyl, or substituted aryl(C₁-C₃₀)alkyl;

n is, independently for each occurrence, an integer from 1 to 5 inclusive;

s, t, x and y each is, independently for each occurrence, an integer from 1 to 30 inclusive;

25 X⁴, X⁵, X⁶, X⁷ and X⁸ each is, independently for each occurrence, H, F, Cl, Br, I, (C₁₋₁₀)alkyl, substituted (C₁₋₁₀)alkyl, aryl, substituted aryl, OH, NH₂, NO₂, or CN; and

provided that when A² is 4Hppa, then R² and R³ are deleted;

or a pharmaceutically acceptable salt thereof.

30 2. A compound according to claim 1, wherein:

A² is Ala, 4Hppa, or deleted;

A³ is Glu, 4Hyp, Pro, or deleted;

A⁴ is Gly or deleted;

A⁵ is Thr or deleted;

35 A⁶ is Phe or deleted;

A⁷ is Ile, A5c, A6c, or deleted;

A⁸ is Ser, Chc-Ser, or deleted;

A⁹ is Asp or deleted;
 A¹⁰ is Tyr or deleted;
 A¹¹ is Ser, A6c, Aib, or deleted;
 A¹² is Ile or deleted;
 5 A¹³ is Ala, Aib, or deleted;
 A¹⁴ is Met, A6c, or Nle;
 A¹⁵ is Asp;
 A¹⁶ is Lys;
 A¹⁷ is Ile;
 10 A¹⁸ is His;
 A¹⁹ is Gln;
 A²⁰ is Gln;
 A²¹ is Asp;
 A²² is Phe;
 15 A²³ is Val;
 A²⁴ is Asn;
 A²⁵ is Trp;
 A²⁶ is Leu;
 A²⁷ is Leu;
 20 A²⁸ is Ala;
 A²⁹ is Gln;
 A³⁰ is Lys;
 A³¹ is Gly, Cys(Hsu), Cys(Psu), 2Nal, D-2Nal, Orn(N-C(O)-(CH₂)₄-CH₃), Orn(N-C(O)-
 (CH₂)₈-CH₃), Orn(N-C(O)-(CH₂)₁₂-CH₃), or deleted;
 25 A³² is Lys, Cys(Psu), or deleted;
 A³³ is Lys, Cys(Psu), or deleted;
 A³⁴ is Asn, Cys(Psu), Orn(N-C(O)-(CH₂)₈-CH₃), or deleted;
 A³⁵ is Asp, Cys(Psu), or deleted;
 A³⁶ is Trp, Cys(Psu), or deleted;
 30 A³⁷ is Lys, Cys(Psu), or deleted;
 A³⁸ is His, Cys(Psu), or deleted;
 A³⁹ is Asn, Cys(Psu), or deleted;
 A⁴⁰ is Ile, Cys(Psu), or deleted;
 A⁴¹ is Thr or deleted;
 35 A⁴² is Gln or deleted;
 A⁴³ is Gln or deleted; and

provided that at least one of A², A³, A⁴, A⁵, A⁶, A⁷, A⁸, A⁹, A¹¹, A¹³, A¹⁴, A³¹, A³², A³³, A³⁴, A³⁵, A³⁶, A³⁷, A³⁸, A³⁹ and A⁴⁰ is not the amino acid residue of the corresponding position of the native GIP;

or a pharmaceutical salt thereof.

5

3. A compound according to claim 2, wherein said compound is:

(Ac-A6c⁷)hGIP(7-42)-OH (SEQ ID NO:4);

[Ac-A6c⁷, Cys(Psu)⁴⁰]hGIP(7-42)-OH (SEQ ID NO:5);

[Ac-A6c⁷, Cys(Psu)³⁹]hGIP(7-42)-OH (SEQ ID NO:6);

10

[Ac-A6c⁷, Cys(Psu)³⁸]hGIP(7-42)-OH (SEQ ID NO:7);

[Ac-A6c⁷, Cys(Psu)³⁶]hGIP(7-42)-OH (SEQ ID NO:8);

[Ac-A6c⁷, Cys(Psu)³⁵]hGIP(7-42)-OH (SEQ ID NO:9);

[Ac-A6c⁷, Cys(Psu)³⁴]hGIP(7-42)-OH (SEQ ID NO:10);

[Ac-A6c⁷, Cys(Psu)³³]hGIP(7-42)-OH (SEQ ID NO:11);

15

[Ac-A6c⁷, Cys(Psu)³²]hGIP(7-42)-OH (SEQ ID NO:12);

[Ac-A6c⁷, Cys(Psu)³¹]hGIP(7-42)-OH (SEQ ID NO:13);

[Ac-A6c⁷, Cys(Psu)³⁷]hGIP(7-42)-OH (SEQ ID NO:14);

[Ac-A6c⁷, Orn³¹(N-C(O)-(CH₂)₁₂-CH₃)]hGIP(7-42)-OH (SEQ ID NO:15);

[Ac-A6c⁷, Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:16);

20

[A6c⁷, Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:17);

[CH₃-(CH₂)₈-C(O)-A6c⁷, Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:18);

[Ac-A6c⁷, Orn³¹(N-C(O)-(CH₂)₄-CH₃)]hGIP(7-42)-OH (SEQ ID NO:19);

[A6c⁷, Orn³¹(N-C(O)-(CH₂)₄-CH₃)]hGIP(7-42)-OH (SEQ ID NO:20);

[CH₃-(CH₂)₄-C(O)-A6c⁷, Orn³¹(N-C(O)-(CH₂)₄-CH₃)]hGIP(7-42)-OH (SEQ ID NO:21);

25

[Ac-A6c⁷, Orn³⁴(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:22);

[A6c⁷, Orn³⁴(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:23);

[CH₃-(CH₂)₈-C(O)-A6c⁷, Orn³⁴(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:24);

[Ac-A6c⁷, Cys(Hsu)³¹]hGIP(7-42)-OH (SEQ ID NO:25);

[A6c⁷, Cys(Hsu)³¹]hGIP(7-42)-OH (SEQ ID NO:26);

30

(Ac-A6c⁷, 2NaI³¹)hGIP(7-42)-OH (SEQ ID NO:27);

(Ac-A6c⁷, D-2NaI³¹)hGIP(7-42)-OH;

(Ac-4Hyp³, A6c⁷)hGIP(3-42)-OH (SEQ ID NO:28);

(Ac-A6c⁷, Gln⁴³)hGIP(7-43)-OH (SEQ ID NO:29);

[Ac-A6c⁷, Cys(Psu)³¹]hGIP(7-34)-NH₂ (SEQ ID NO:30);

35

[Ac-A6c⁷, Cys(Psu)³¹]hGIP(7-31)-NH₂ (SEQ ID NO:31);

[Ac-Phe⁶, A6c⁷, Cys(Psu)³¹]hGIP(6-42)-OH (SEQ ID NO:32);

[A6c⁷, Cys(Psu)³¹]hGIP(6-42)-OH (SEQ ID NO:33);

- (Ac-Phe⁶, A6c⁷)hGIP(6-30)-NH₂ (SEQ ID NO:34);
 [Ac-Phe⁶, A6c⁷, Cys(Psu)³¹]hGIP(6-31)-NH₂ (SEQ ID NO:35);
 [A6c⁷, Cys(Psu)³¹]hGIP(6-31)-NH₂ (SEQ ID NO:36);
 (A5c⁷, Nle¹⁴)hGIP(6-30)-NH₂ (SEQ ID NO:37);
 5 (A6c⁷, Nle¹⁴)hGIP(6-30)-NH₂ (SEQ ID NO:38);
 (Aib¹¹, Nle¹⁴)hGIP(6-30)-NH₂ (SEQ ID NO:39);
 [Ac-Asp⁹, Cys(Psu)³³]hGIP(9-42)-OH (SEQ ID NO:40);
 [Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(8-42)-OH (SEQ ID NO:41);
 [Chc-Ser⁸, Cys(Psu)³¹]hGIP(8-42)-OH (SEQ ID NO:42);
 10 [CH₃-(CH₂)₄-C(O)-Ser⁸, Cys(Psu)³¹]hGIP(8-42)-OH (SEQ ID NO:43);
 (4Hppa², 4Hyp³, A6c⁷)hGIP(2-42)-OH (SEQ ID NO:44);
 (4Hppa², Pro³, Nle¹⁴)hGIP(2-42)-OH (SEQ ID NO:45);
 (4Hppa², Aib¹³)hGIP(2-42)-OH (SEQ ID NO:46);
 (4Hppa², A6c¹⁴)hGIP(2-42)-OH (SEQ ID NO:47);
 15 (4Hppa², A6c¹¹)hGIP(2-42)-OH (SEQ ID NO:48); or
 [Aib², A5c¹¹, Nle¹⁴, Lys⁴³(N-C(O)-(CH₂)₁₀-CH₃)]hGIP(2-43)-OH (SEQ ID NO:49);
 or a pharmaceutically acceptable salt thereof.

4. A compound according to claim 2, wherein:

- 20 A² is deleted;
 A³ is deleted;
 A⁴ is deleted;
 A⁵ is deleted;
 A⁶ is deleted;
 25 A⁷ is A5c or A6c;
 A⁸ is Ser;
 A⁹ is Asp;
 A¹⁰ is Tyr;
 A¹¹ is Ser;
 30 A¹² is Ile;
 A¹³ is Ala;
 A¹⁴ is Met;
 A³¹ is Gly, Cys(Hsu), Cys(Psu), 2Nal, D-2Nal, Orn(N-C(O)-(CH₂)₄-CH₃), Orn(N-C(O)-
 (CH₂)₈-CH₃), or Orn(N-C(O)-(CH₂)₁₂-CH₃);
 35 A³² is Lys or Cys(Psu);
 A³³ is Lys or Cys(Psu);
 A³⁴ is Asn, Cys(Psu), or Orn(N-C(O)-(CH₂)₈-CH₃);

A³⁵ is Asp or Cys(Psu);

A³⁶ is Trp or Cys(Psu);

A³⁷ is Lys or Cys(Psu);

A³⁸ is His or Cys(Psu);

5 A³⁹ is Asn or Cys(Psu);

A⁴⁰ is Ile or Cys(Psu);

A⁴¹ is Thr;

A⁴² is Gln; and

A⁴³ is deleted;

10 or a pharmaceutical salt thereof.

5. A compound according to claim 4, wherein said compound is:

(Ac-A6c⁷)hGIP(7-42)-OH (SEQ ID NO:4);

[Ac-A6c⁷, Cys(Psu)⁴⁰]hGIP(7-42)-OH (SEQ ID NO:5);

15 [Ac-A6c⁷, Cys(Psu)³⁹]hGIP(7-42)-OH (SEQ ID NO:6);

[Ac-A6c⁷, Cys(Psu)³⁸]hGIP(7-42)-OH (SEQ ID NO:7);

[Ac-A6c⁷, Cys(Psu)³⁶]hGIP(7-42)-OH (SEQ ID NO:8);

[Ac-A6c⁷, Cys(Psu)³⁵]hGIP(7-42)-OH (SEQ ID NO:9);

[Ac-A6c⁷, Cys(Psu)³⁴]hGIP(7-42)-OH (SEQ ID NO:10);

20 [Ac-A6c⁷, Cys(Psu)³³]hGIP(7-42)-OH (SEQ ID NO:11);

[Ac-A6c⁷, Cys(Psu)³²]hGIP(7-42)-OH (SEQ ID NO:12);

[Ac-A6c⁷, Cys(Psu)³¹]hGIP(7-42)-OH (SEQ ID NO:13);

[Ac-A6c⁷, Cys(Psu)³⁷]hGIP(7-42)-OH (SEQ ID NO:14);

[Ac-A6c⁷, Orn³¹(N-C(O)-(CH₂)₁₂-CH₃)]hGIP(7-42)-OH (SEQ ID NO:15);

25 [Ac-A6c⁷, Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:16);

[A6c⁷, Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:17);

[CH₃-(CH₂)₈-C(O)-A6c⁷, Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:18);

[Ac-A6c⁷, Orn³¹(N-C(O)-(CH₂)₄-CH₃)]hGIP(7-42)-OH (SEQ ID NO:19);

[A6c⁷, Orn³¹(N-C(O)-(CH₂)₄-CH₃)]hGIP(7-42)-OH (SEQ ID NO:20);

30 [CH₃-(CH₂)₄-C(O)-A6c⁷, Orn³¹(N-C(O)-(CH₂)₄-CH₃)]hGIP(7-42)-OH (SEQ ID NO:21);

[Ac-A6c⁷, Orn³⁴(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:22);

[A6c⁷, Orn³⁴(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:23);

[CH₃-(CH₂)₈-C(O)-A6c⁷, Orn³⁴(N-C(O)-(CH₂)₈-CH₃)]hGIP(7-42)-OH (SEQ ID NO:24);

[Ac-A6c⁷, Cys(Hsu)³¹]hGIP(7-42)-OH (SEQ ID NO:25);

35 [A6c⁷, Cys(Hsu)³¹]hGIP(7-42)-OH (SEQ ID NO:26);

(Ac-A6c⁷, 2NaI³¹)hGIP(7-42)-OH (SEQ ID NO:27); or

(Ac-A6c⁷, D-2NaI³¹)hGIP(7-42)-OH;

or a pharmaceutically acceptable salt thereof.

6. A compound according to claim 2, wherein: A⁴³ is deleted; A² is 4Hppa; and at least one of A³, A⁷, A¹¹, A¹³ and A¹⁴ is not the amino acid residue of the corresponding position of the native GIP; or a pharmaceutical salt thereof.

7. A compound according to claim 6, wherein said compound is:

(4Hppa², 4Hyp³, A6c⁷)hGIP(2-42)-OH (SEQ ID NO:44);

(4Hppa², Pro³, Nle¹⁴)hGIP(2-42)-OH (SEQ ID NO:45);

10. (4Hppa², Aib¹³)hGIP(2-42)-OH (SEQ ID NO:46);

(4Hppa², A6c¹⁴)hGIP(2-42)-OH (SEQ ID NO:47); or

(4Hppa², A6c¹¹)hGIP(2-42)-OH (SEQ ID NO:48);

or a pharmaceutically acceptable salt thereof.

15. 8. A compound according to claim 2, wherein: A² to A⁵ and A³¹ to A⁴³ are deleted; and at least one of A⁶, A⁷, A¹¹ and A¹⁴ is not the amino acid residue of the corresponding position of the native GIP; or a pharmaceutical salt thereof.

9. A compound according to claim 8, wherein said compound is:

20. (Ac-Phe⁶, A6c⁷)hGIP(6-30)-NH₂ (SEQ ID NO:34);

(A5c⁷, Nle¹⁴)hGIP(6-30)-NH₂ (SEQ ID NO:37);

(A6c⁷, Nle¹⁴)hGIP(6-30)-NH₂ (SEQ ID NO:38); or

(Aib¹¹, Nle¹⁴)hGIP(6-30)-NH₂ (SEQ ID NO:39);

or a pharmaceutically acceptable salt thereof.

25. 10. A compound according to claim 2, wherein: A² to A⁷ and A⁴³ are deleted; and at least one of A⁸ and A³¹ is not the amino acid residue of the corresponding position of the native GIP; or a pharmaceutical salt thereof.

30. 11. A compound according to claim 10, wherein said compound is:

[Orn³¹(N-C(O)-(CH₂)₈-CH₃)]hGIP(8-42)-OH (SEQ ID NO:41);

[Chc-Ser⁸, Cys(Psu)³¹]hGIP(8-42)-OH (SEQ ID NO:42); or

[CH₃-(CH₂)₄-C(O)-Ser⁸, Cys(Psu)³¹]hGIP(8-42)-OH (SEQ ID NO:43);

or a pharmaceutically acceptable salt thereof.

35. 12. A compound according to claim 2, wherein: A² to A⁵ and A⁴³ are deleted; and at least one of A⁶, A⁷ and A³¹ is not the amino acid residue of the corresponding position of the native GIP; or a pharmaceutical salt thereof.

13. A compound according to claim 12, wherein said compound is:

[Ac-Phe⁶, A6c⁷, Cys(Psu)³¹]hGIP(6-42)-OH (SEQ ID NO:32); or

[A6c⁷, Cys(Psu)³¹]hGIP(6-42)-OH (SEQ ID NO:33);

5 or a pharmaceutically acceptable salt thereof.

14. A compound according to claim 2, wherein: A² to A⁵ and A³² to A⁴³ are deleted; and at least one of A⁶, A⁷ and A³¹ is not the amino acid residue of the corresponding position of the native GIP; or a pharmaceutical salt thereof.

10

15. A compound according to claim 14, wherein said compound is:

[Ac-Phe⁶, A6c⁷, Cys(Psu)³¹]hGIP(6-31)-NH₂ (SEQ ID NO:35); or

[A6c⁷, Cys(Psu)³¹]hGIP(6-31)-NH₂ (SEQ ID NO:36);

or a pharmaceutically acceptable salt thereof.

15

16. A compound according to any one of claims 1-15, further comprising a covalently linked PEG moiety, or a pharmaceutically acceptable salt thereof.

17. A compound according to claim 16, wherein said PEG moiety is linked to the
20 compound via a Cys(maleimide), hCys(maleimide), or Pen(maleimide) linker, to form Cys(succinimide-N-PEG), hCys(succinimide-N-PEG), or Pen(succinimide-N-PEG), or a pharmaceutically acceptable salt thereof.

18. A compound according to claim 17, wherein PEGylation occurs at any one of amino
25 acid residue positions 16, 30, and 31-43, whereby Cys(succinimide-N-PEG), hCys(succinimide-N-PEG), or Pen(succinimide-N-PEG) is placed in any one of amino acid residue positions 16, 30, and 31-43, or a pharmaceutically acceptable salt thereof.

19. A compound according to claim 18, wherein PEGylation occurs at any one of amino
30 acid residue positions 32, 33 and 43, whereby Cys(succinimide-N-PEG), hCys(succinimide-N-PEG), or Pen(succinimide-N-PEG) is placed in any one of amino acid residue positions 32, 33 and 43, or a pharmaceutically acceptable salt thereof.

20. A compound according to claim 19, wherein said PEG moiety has average molecular
35 weight of from about 2,000 to about 80,000, or a pharmaceutically acceptable salt thereof.

21. A compound according to claim 20, wherein said PEG moiety is selected from the group consisting of 5K PEG, 10K PEG, 20K PEG, 30K PEG, 40K PEG, 50K PEG, and 60K PEG, to form Cys(succinimide-N-5K PEG), Cys(succinimide-N-10K PEG), Cys(succinimide-N-20K PEG),

Cys(succinimide-N-30K PEG), Cys(succinimide-N-40K PEG), Cys(succinimide-N-50K PEG), Cys(succinimide-N-60K PEG), hCys(succinimide-N-5K PEG), hCys(succinimide-N-10K PEG), hCys(succinimide-N-20K PEG), hCys(succinimide-N-30K PEG), hCys(succinimide-N-40K PEG), hCys(succinimide-N-50K PEG), hCys(succinimide-N-60K PEG), Pen(succinimide-N-5K PEG),
5 Pen(succinimide-N-10K PEG), Pen(succinimide-N-20K PEG), Pen(succinimide-N-30K PEG), Pen(succinimide-N-40K PEG), Pen(succinimide-N-50K PEG), or Pen(succinimide-N-60K PEG), or a pharmaceutically acceptable salt thereof.

10 22. A compound according to any one of claims 17-21, wherein said succinimide-N-PEG is linear, or a pharmaceutically acceptable salt thereof.

23. A compound according to claim 22, wherein said linear succinimide-N-PEG is succinimide-N-(CH₂)₂-C(O)NH-(CH₂)₃-PEG, or a pharmaceutically acceptable salt thereof.

15 24. A compound according to any one of claims 17-21, wherein said succinimide-N-PEG is branched, or a pharmaceutically acceptable salt thereof.

20 25. A compound according to claim 24, wherein said branched succinimide-N-PEG is succinimide-N-(CH₂)₂-C(O)NH-(CH₂)₃-O-CH₂-CH(PEG)-CH₂-PEG, or a pharmaceutically acceptable salt thereof.

26. A pharmaceutical composition comprising an effective amount of a peptide analogue of any one of claims 1-25.

25 27. A pharmaceutical composition of claim 26, further comprising a pharmaceutically acceptable carrier.

30 28. A method of eliciting an agonist effect from a GIP receptor in a subject in need thereof which comprises administering to said subject a therapeutically effective amount of a peptide analogue of any one of claims 1-25 or a pharmaceutical composition of claim 26 or claim 27.

29. A method of eliciting an antagonist effect from a GIP receptor in a subject in need thereof which comprises administering to said subject a therapeutically effective amount of a peptide analogue of any one of claims 1-25 or a pharmaceutical composition of claim 26 or claim 27.

35 30. A method for treating conditions or diseases mediated by GIP-receptor binding, comprising the step of administering to a subject in need thereof a therapeutically effective amount of a peptide analogue of any one of claims 1-25 or a pharmaceutical composition of claim 26 or claim 27.

31. The method of claim 30, wherein said condition or disease mediated by GIP-receptor binding is selected from the group consisting of type 1 diabetes, type 2 diabetes, obesity, insulin resistance, glucose intolerance, fatty liver, glucagonomas, secretory disorders of the airway, metabolic disorders, arthritis, osteoporosis, central nervous system disease, restenosis, neurodegenerative disease, renal failure, congestive heart failure, nephrotic syndrome, cirrhosis, pulmonary edema, hypertension, and disorders wherein the reduction of food intake and/or losing body weight is desired.

32. A method for treating diabetes, comprising the step of administering to a subject in need thereof a therapeutically effective amount of a peptide analogue of any one of claims 1-25 or a pharmaceutical composition of claim 26 or claim 27.

33. The method of claim 32, wherein said diabetes is type 2 diabetes.

34. A method of treating diabetes-related disorders, comprising the step of administering to a subject in need thereof a therapeutically effective amount of a peptide analogue of any one of claims 1-25 or a pharmaceutical composition of claim 26 or claim 27.

35. The method of claim 34, wherein said diabetes-related disorder is selected from the group consisting of hyperglycemia, hyperinsulinemia, impaired glucose tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, and insulin resistance.

36. A method of treating or preventing secondary causes of diabetes, comprising the step of administering to a subject in need thereof a therapeutically effective amount of a peptide analogue of any one of claims 1-25 or a pharmaceutical composition of claim 26 or claim 27.

37. The method of claim 36, wherein said secondary cause is selected from the group consisting of glucocorticoid excess, growth hormone excess, pheochromocytoma, and drug-induced diabetes.

38. A method of treating obesity, comprising the step of administering to a subject in need thereof a therapeutically effective amount of a peptide analogue of any one of claims 1-25 or a pharmaceutical composition of claim 26 or claim 27.

39. A method of stimulating insulin secretion in a subject in need thereof by administering to said subject a therapeutically effective amount of a peptide analogue of any claims 1-25 or a pharmaceutical composition of claim 26 or claim 27.

40. Use of a peptide analogue of any one of claims 1-25 for the manufacture of a medicament for GIP-receptor binding for the prevention or treatment of diseases or conditions related to impaired binding of GIP-receptor analogues.

41. Use according to claim 40 for the manufacture of a medicament for the prevention or treatment of pancreatic beta cell apoptosis.

5 42. Use according to claim 40 for the manufacture of a medicament for the potentiation of glucose dependent proliferation of pancreatic beta cells.