

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
28 April 2005 (28.04.2005)

PCT

(10) International Publication Number
WO 2005/037305 A1

- (51) International Patent Classification⁷: **A61K 38/19**, A61P 29/00, 31/00, 37/00
- (21) International Application Number: PCT/EP2004/052572
- (22) International Filing Date: 18 October 2004 (18.10.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 03078308.8 16 October 2003 (16.10.2003) EP
- (71) Applicant (for all designated States except US): **APPLIED RESEARCH SYSTEMS ARS HOLDING N.V.** [NL/—]; Pietermaai 15, Curacao (AN).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **PROUDFOOT, Amanda** [CH/FR]; Sous Chens, F-74140 Chens Sur Lemans (FR). **SHAW, Jeffrey** [IT/CH]; Chemin Thury 10, CH-1206 Geneva (CH). **JOHNSON, Zoe** [GB/CH]; Rue du 1er Juin 6, CH-1207 Geneva (CH).
- (74) Agent: **SERONO INTERNATIONAL S.A. INTELLECTUAL PROPERTY**; 12, Chemin des Aulx, CH-1228 Plan-les-Ouates (CH).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- with international search report
 - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
 - with sequence listing part of description published separately in electronic form and available upon request from the International Bureau
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2005/037305 A1

(54) Title: THERAPEUTIC USES OF CHEMOKINE VARIANTS

(57) Abstract: Variants of homodimer-forming chemokines, such as human CCL2, having a single amino acid substitution in the dimerization interface that alters the pattern of hydrogen bonds and acting as an obligate monomer, can antagonize natural chemokines and have anti-inflammatory activity *in vivo*. These variants can be used as active ingredient in pharmaceutical compositions for the treatment of inflammatory, autoimmune, or infectious diseases.

THERAPEUTIC USES OF CHEMOKINE VARIANTS

FIELD OF THE INVENTION

The invention relates to novel therapeutic uses of chemokine variants, and in particular of human CCL2 variants.

BACKGROUND OF THE INVENTION

Chemokines are small, secreted pro-inflammatory proteins, which mediate directional migration of leukocytes from the blood to the site of injury. Depending on the position of the conserved cysteines characterizing this family of proteins, the chemokine family can be divided structurally in C, C-C, C-X-C and C-X₃-C chemokines which bind to a series of membrane receptors (Baggiolini M et al., 1997; Fernandez EJ and Lolis E, 2002).

These membrane receptors, all heptahelical G-protein coupled receptors, allow chemokines to exert their biological activity on the target cells, which may present specific combinations of receptors according to their state and/or type. The physiological effects of chemokines result from a complex and integrated system of concurrent interactions: the receptors often have overlapping ligand specificity, so that a single receptor can bind different chemokines, as well a single chemokine can bind to different receptors.

Usually chemokines are produced at the site of injury and cause leukocyte migration and activation, playing a fundamental role in inflammatory, immune, homeostatic, hematopoietic, and angiogenic processes. Even though there are potential drawbacks in using chemokines as therapeutic agents (tendency to aggregate and promiscuous binding, in particular), these molecules are considered good target

candidates for therapeutic intervention in diseases associated to such processes, by inhibiting specific chemokines and their receptors at the scope to preventing the excessive recruitment and activation of leukocytes (Baggiolini M, 2001; Loetscher P and Clark-Lewis I, 2001; Godessart N and Kunkel SL, 2001).

5 Studies on structure-activity relationships indicate that chemokines have two main sites of interaction with their receptors, the flexible amino-terminal region and the conformationally rigid loop that follows the second cysteine. Chemokines are thought to dock onto receptors by means of the loop region, and this contact is believed to facilitate the binding of the amino-terminal region that results in receptor activation.

10 This importance of the amino-terminal region has been also demonstrated by testing natural and synthetic chemokines in which this domain is modified or shortened. This processing, following proteolytic digestion, mutagenesis, or chemical modification of amino acids, can either activate or render these molecules inactive, generating compounds with agonistic and/or antagonistic activity. Thus, chemokines with specific

15 modifications in the amino-terminal region have therapeutic potential for inflammatory and autoimmune diseases (Schwarz and Wells, 1999).

 CCL2, also known as Monocyte Chemoattractant Protein 1 (MCP-1) or Monocyte Chemotactic And Activating Factor (MCAF), has been identified as having a central role in inflammation, being capable of promoting the recruitment of monocytes and

20 lymphocytes in response to injury and infection signals in various inflammatory diseases, different types of tumors, cardiac allograft, AIDS, and tuberculosis (Yoshimura T et al., 1989; Gu L et al., 1999). The physiological activities associated with CCL2 have been extensively studied by means of transgenic animals, which allowed the demonstration that this chemokine controls not only monocyte recruitment in

25 inflammatory models, but also the expression of cytokines related to T helper

responses and the initiation of atherosclerosis (Gu L et al., 2000; Gosling J et al, 1999; Lu B et al, 1998).

Structurally, CCL2 consists of a N-terminal loop and a beta sheet overlaid by an alpha helix at the C-terminal end, and forms homodimers, even though has been
5 detected as a monomer in specific experimental conditions (Handel T et al., 1996; Kim KS et al., 1996; Lubkowski J, et al., 1997). As for many other chemokines, the literature provides many examples of structure-activity studies in which CCL2 mutants have been generated by expressing N-terminal truncated or single site substituted variants
10 activities and other properties (Gong J and Clark-Lewis I, 1995; Zhang Y et al., 1996; Steitz SA et al., 1998; Gu L et al., 1999; Hemmerich S et al., 1999; Seet BT et al., 2001).

In particular, the role of dimerization in CCL2 receptor binding and activation was studied showing that different mutations in the terminal region hindering dimerization
15 may alter some CCL2 activities such as receptor binding affinity, stimulation of chemotaxis, inhibition of adenylate cyclase, and stimulation of calcium influx (Paavola C et al, 1998). While one mutant described by Paavola, herein called P8A*, does not dimerize, but maintains original potency and efficacy, another obligate monomeric mutant described by Paavola, herein called Y13A*, was shown to have a 100-fold
20 weaker binding affinity *in vitro*, to be a much less potent inhibitor of adenylate cyclase and stimulator of calcium influx *in vitro*, and unable to stimulate chemotaxis in cell culture. Similarly to Y13A*, a mutant, [1+9-76]MCP-1 (a CCL2 variant lacking residues 2-8), antagonizes CCL2 activities *in vitro*.

The binding properties of the CCL2 mutant containing the P8A substitution were
25 also studied in an experimental model based upon the recognition of the viral

chemokine binding protein M3, demonstrating the efficient binding of this viral protein to this CCL2 mutant (Alexander JM et al., 2002). Moreover it has been shown that monomeric variants of chemokines, such as CCL2-P8A, are devoid of activity *in vivo*, although fully active and indistinguishable from the dimeric form *in vitro* (Proudfoot A et al., 2003).

Examples of CCL2 mutants involving residues not affecting the dimerization profile of the resulting protein have been already described in the literature as leading to molecules having antagonistic properties towards CCL2 (US 5739103, WO 03/84993). However, there is not indication in the prior art that a specific chemokine mutant, and in particular a CCL2 mutant, being an obligate monomer due to a single site substitution (for example, involving a Proline), may act as a chemokine antagonist.

SUMMARY OF THE INVENTION

It has been surprisingly found that variants of homodimer-forming chemokines, such as CCL2, having a single amino acid substitution in the dimerization interface that alters the pattern of hydrogen bonds, so as to result in an obligate monomer that binds to the receptor and has agonistic properties *in vitro*, can antagonize natural chemokines and have anti-inflammatory activity *in vivo*.

The variants described herein are useful as a medicament. Polypeptides comprising SEQ ID NO: 2 and those comprising SEQ ID NO: 2 with a further mutation to isoleucine at position 64 of SEQ ID NO: 2 (SEQ ID NO: 4) are useful as medicaments.

The variants described herein are useful as medicaments for the treatment of autoimmune, inflammatory or infectious diseases. Polypeptides comprising SEQ ID NO: 2 and those comprising SEQ ID NO: 2 with a further mutation to isoleucine at

position 64 of SEQ ID NO: 2 (SEQ ID NO: 4) are useful as medicaments for the treatment of autoimmune, inflammatory and infectious diseases. In one example, polypeptides comprising SEQ ID NO: 2 and those comprising SEQ ID NO: 2 with a further mutation to isoleucine at position 64 of SEQ ID NO: 2 (SEQ ID NO: 4) are useful
5 as medicaments for the treatment of multiple sclerosis.

The variants described herein are further useful in methods for treatment of autoimmune, inflammatory and infectious diseases. Such methods comprise the administration of an effective amount of a monomeric variant of the invention, wherein said variant results from at least an amino acid substitution that alters the pattern of
10 hydrogen bonds at the dimerization interface of said chemokine. Examples of such monomeric variants that can be used in such methods are: a) a polypeptide comprising SEQ ID NO: 2; b) a polypeptide comprising SEQ ID NO: 4; c) an active mutant of a) or b); d) a polypeptide comprising a), b) or c) and an amino acid sequence belonging to a protein other than said chemokine; as well as the corresponding
15 molecules in the form of their active fractions, precursors, salts, derivatives, complexes or conjugates.

The invention further includes pharmaceutical compositions containing a polypeptide comprising the obligate monomeric antagonistic chemokine variants of the Invention.

20 The Invention further includes fusion proteins comprising an obligate monomeric antagonistic chemokine variant described herein fused to a non-chemokine protein sequence, for example the amino acid sequence of SEQ ID NO: 2 fused to the constant region of a human immunoglobulin heavy chain (SEQ IN NO: 5). The Invention further includes methods for producing these fusion proteins.

The Invention further includes methods for screening for obligate monomeric antagonist chemokine variants described herein comprising:

- a) making single point mutations in CCL2 that block its ability to dimerize;
- b) identifying said variants that bind to the receptor and show agonistic properties *in vitro*;
- c) identifying said variants from the group identified in (b) above that are further characterized as inhibiting peritoneal cell recruitment.

Other features and advantages of the invention will be apparent from the following detailed description.

10

DESCRIPTION OF THE FIGURES

Figure 1: amino acid sequences of human mature CCL2 (residues 24-99 of SWISSPROT Acc. N° P13500; SEQ ID NO: 1), mature CCL2-P8A (SEQ ID NO: 2), mature CCL2* (SEQ ID NO: 3), mature CCL2*-P8A (SEQ ID NO: 4). The mutation in position 64 to Isoleucin (M64I) is boxed in CCL2* and CCL2*-P8A; the relevant mutation P8A is bold and underlined.

15

Figure 2: graph comparing the cell recruiting activity of recombinant human CCL2 and CCL2-P8A with the baseline in the peritoneal cell recruiting assay .

Figure 3: graph showing the dose-response inhibiting activity of CCL2-P8A in peritoneal cell recruitment assays using two different inducers: CCL2 (A) or Thioglycollate (B); the results are compared with the baseline level, a negative control (vehicle only, saline) and a positive control, Dexamethasone (Dex; only in B).

20

- Figure 4: graph showing the inhibiting activity of CCL2-P8A in ovalbumin-induced lung inflammation assay; the result is compared with the baseline level, and a negative control (saline).
- Figure 5: Graph showing the efficacy of CCL2-P8A in reducing clinical scores in an animal model for multiple sclerosis called EAE (Experimental Autoimmune Encephalomyelitis), either when the animals present a mild disease with low clinical score (not more than 2; A) or a severe disease with high clinical score (at least 3; B); the data are compared to those observed using the vehicle only (negative control) or positive control for the EAE model (mouse Interferon beta); the number of asterisks on the bottom of each graph indicate the level of statistical significance of the effect observed when compared to the negative control at each time point (* means $p < 0.05$, ** means $p < 0.01$, and *** means $p < 0.001$, as calculated by one-way ANOVA followed by Fisher test).
- Figure 6: graph showing the inhibiting activity of CCL2-P8A when administered at different dosages in an animal model for skin inflammation (DNFB-induced inflammation on the ear of sensitized mice); the effect of CCL2-P8A on the volume of the ear swelling volume after the treatment, and it is compared with the following controls: animals not sensitized but challenged at day 5 with DNFB (A), animal sensitized and challenged with DNFB but only vehicle treated (B), and animals treated with Dexamethasone (Dex; administered in the same manner of CCL2-P8A). The values were measured at day 6 (the day after the treatment and the challenge).
- Figure 7: Alternative forms of CCL2-P8A that can be generated by single/multiple amino acid addition and/or substitution (A) or by fusion to residues 243 –

474 of human immunoglobulin lambda heavy chain IgG1 (B; CCL2 signal sequence is double underlined, mature CCL2-P8A is underlined).

DETAILED DESCRIPTION OF THE INVENTION

5 In view of the above mentioned evidences in the prior art, there is no indication that a CCL2 variant, resulting from at least single amino acid substitution in the dimerization interface that alters the pattern of hydrogen bonds so as to result in an obligate monomer binding to the receptor and having agonistic properties *in vitro*, can antagonize CCL2 *in vivo*, and in general can inhibit cell recruitment and/or
10 inflammatory reactions.

The single amino acid substitution is preferably at position 8 in human mature CCL2, and in particular it consists of replacing Proline in position 8 with an Alanine. These variants do not contain an additional mutation in the position corresponding to 9, 10, or 13 of human mature CCL2.

15 An example of the above described substitutions is monomeric variant of the mature form of CCL2 called CCL2-P8A which can be expressed either as including the single substitution of the Proline in position 8 to Alanine (SEQ ID NO: 2) or as including a further substitution (improving expression of the protein) of Methionine 64 to Isoleucine (SEQ ID NO: 4). The substitution of Proline 8 to Alanine (an amino acid
20 having a different orientation of the chemical group possibly forming hydrogen bonds) generate a CCL2 variant acting as an obligate monomer. CCL2-P8A and CCL2*-P8A can be considered as molecule having equivalent activity.

Pharmaceutical uses, methods, and compositions that can be consequently envisaged for these specific obligate monomeric variants of CCL2, called CCL2-P8A
25 and CCL2*-P8A, can be also envisaged for any other obligate monomeric variant of

CCL2, or of other chemokines that are naturally active as dimers, that are generated in the same manner. These mutants should present antagonistic and inhibitory properties similar to those of CCL2-P8A and CCL2*-P8A against the natural chemokine, and therefore they should have a medical applicability.

5 One aspect of the present invention is the use of an obligate monomeric variant of a homodimer-forming chemokine, for example CCL2 which is known to be a therapeutic target for various diseases, such as autoimmune, inflammatory, or infectious diseases. Such variants result from at least an amino acid substitution that alters the pattern of hydrogen bonds at the dimerization interface of said chemokine, so
10 as to result in an obligate monomer that binds to the receptor and has agonistic properties *in vitro* and antagonistic properties *in vivo*, as active ingredient in a pharmaceutical composition, in particular for the treatment or prevention of autoimmune, inflammatory, or infectious diseases. These *in vivo* antagonistic properties may become evident in assays allowing the evaluation of properties such as
15 the effect cell recruitment in the peritoneum following the induction with an inflammatory molecule (a chemokine such CCL2 itself, thioglycollate, or ovalbumin).

The obligate monomeric variant may result from a single amino acid substitution in the dimerization interface that alters the pattern of hydrogen bonds. More preferably, being proline an amino acid well known to be particularly relevant for establishing
20 stereospecific hydrogen bonds involved in the formation of protein complexes such as homodimers, the single amino acid substitution should be the substitution of a proline with non-proline amino acid. Alternatively, the single amino acid substitution should be the substitution of non-proline amino acid with a proline, so that the stereospecificity of hydrogen bonds results altered.

Alternative forms of the monomeric variants of the homodimer-forming chemokines above defined that can be used as active ingredients in pharmaceutical compositions include:

- a) an active mutant; or
- 5 b) a polypeptide comprising said variant, or said active mutant, and an amino acid sequence belonging to a protein sequence other than to said chemokine.

The term "active" means that such alternative compounds should maintain the functional features of the CCL2 mutants of the present invention, i.e. should antagonize CCL2 *in vivo* and inhibit cell recruitment and/or inflammatory reactions.

10 An obligate monomeric chemokine antagonist as defined herein, as an active compound according to the Invention, can be in the form of their active fractions, precursors, salts, derivatives, complexes or conjugates.

These alternative compounds are intended to comprehend molecules with changes to the sequence of the monomeric variants of the homodimer-forming chemokines that do not affect the basic characteristics disclosed herein. For example,
15 CCL2*-P8A has been generated on the basis of a mutant of CCL2 (CCL2-M64I or CCL2*) that maintains normal CCL2 activities but has improved properties regarding the recombinant expression.

The antagonistic properties of the monomeric variants of homodimer-forming chemokines defined above, and exemplified herein using CCL2-P8A as CCL2
20 antagonist, can be maintained, or even potentiated, in the active mutants. This category of molecules includes natural or synthetic analogs of said sequence, wherein one or more amino acid residues have been added, deleted, or substituted, provided they display the same biological activity characterized in the present invention at

comparable or higher levels, as determined by means known in the art and disclosed in the Examples below.

Natural analogs are intended to be the corresponding sequences of human chemokines proteins identified in other organisms, like mouse CCL2 (SWISSPROT 5 Acc. N° P10148). Artificial analogs of the monomeric variants of homodimer-forming chemokines are intended to be polypeptides prepared by known chemical synthesis and/or by site-directed mutagenesis techniques, combinatorial technologies at the level of encoding DNA sequence (such as DNA shuffling, phage display/selection), by computer-aided design studies, or by any other known technique suitable thereof, 10 which provide a finite set of substantially corresponding mutated or shortened peptides or polypeptides which can be routinely obtained and tested by one of ordinary skill in the art using the teachings presented in the prior art and in the Examples below.

For example, specific artificial mutants may have one or more amino acids being substituted in other positions of CCL2 and found relevant for generating CCL2 15 antagonists (WO 03/84993).

In accordance with the present invention, preferred changes in the active mutants are commonly known as "conservative" or "safe" substitutions, and involve non-basic residues. Conservative amino acid substitutions are those with amino acids having sufficiently similar chemical properties, in order to preserve the structure and the 20 biological function of the molecule. It is clear that insertions and deletions of amino acids may also be made in the above defined sequences without altering their function, particularly if the insertions or deletions only involve a few amino acids, e.g., under ten, and preferably under three, and do not remove or displace amino acids which are critical to the functional conformation of a protein or a peptide.

The literature provide many models on which the selection of conservative amino acids substitutions can be performed on the basis of statistical and physico-chemical studies on the sequence and/or the structure of natural protein (Rogov SI and Nekrasov AN, 2001). Protein design experiments have shown that the use of specific subsets of amino acids can produce foldable and active proteins, helping in the classification of amino acid "synonymous" substitutions which can be more easily accommodated in protein structure, and which can be used to detect functional and structural homologs and paralogs (Murphy LR et al., 2000). The synonymous amino acid groups and more preferred synonymous groups are those defined in Table I. For example, the substitution of Methionine 64 with Isoleucine common to CCL2* and CCL2*-P8A have been chosen using a similar criteria.

A further group of active mutants of the monomeric variants of the homodimer-forming chemokines defined above are peptide mimetics (also called peptidomimetics), in which the nature of peptide or polypeptide has been chemically modified at the level of amino acid side chains, of amino acid chirality, and/or of the peptide backbone. These alterations are intended to provide monomeric variants of the homodimer-forming chemokines having similar or improved properties in terms of preparation, potency and/or pharmacokinetics features.

For example, when the peptide is susceptible to cleavage by peptidases following injection into the subject is a problem, replacement of a particularly sensitive peptide bond with a non-cleavable peptide mimetic can provide a peptide more stable and thus more useful as a therapeutic. Similarly, the replacement of an L-amino acid residue is a standard way of rendering the peptide less sensitive to proteolysis, and finally more similar to organic compounds other than peptides. Also useful are amino-terminal blocking groups such as t-butyloxycarbonyl, acetyl, theyl, succinyl, methoxysuccinyl,

suberyl, adipyl, azelayl, dansyl, benzyloxycarbonyl, fluorenyl methoxycarbonyl, methoxyazelayl, methoxyadipyl, methoxysuberyl, and 2,4-dinitrophenyl. Many other modifications providing increased potency, prolonged activity, easiness of purification, and/or increased half-life are known in the art (WO 02/10195; Villain M et al., 2001).

5 Preferred alternative, "synonymous" groups for amino acids included in peptide mimetics are those defined in Table II. . A non-exhaustive list of amino acid derivatives also include aminoisobutyric acid (Aib), hydroxyproline (Hyp), 1,2,3,4-tetrahydro-isoquinoline-3-COOH, indoline-2carboxylic acid, 4-difluoro-proline, L- thiazolidine -4-carboxylic acid, L-homoproline, 3,4-dehydro-proline, 3,4-dihydroxy-phenylalanine,
10 cyclohexyl-glycine, and phenylglycine.

By "amino acid derivative" is intended an amino acid or amino acid-like chemical entity other than one of the 20 genetically encoded naturally occurring amino acids. In particular, the amino acid derivative may contain substituted or non-substituted alkyl moieties that can be linear, branched, or cyclic, and may include one or more
15 heteroatoms. The amino acid derivatives can be made de novo or obtained from commercial sources (Calbiochem-Novabiochem AG, Switzerland; Bachem, USA).

The techniques for the synthesis and the development of peptide mimetics, as well as non-peptide mimetics, are well known in the art (Hruby VJ and Balse PM, 2000; Golebiowski A et al., 2001). Various methodology for incorporating unnatural amino
20 acids into proteins, using both *in vitro* and *in vivo* translation systems, to probe and/or improve protein structure and function are also disclosed in the literature (Dougherty DA, 2000).

The present Invention discloses the use of monomeric variants of the homodimer-forming chemokines, and their active mutants, as active ingredients in
25 pharmaceutical compositions, as well as of proteins comprising their amino acid

sequence and an amino acid sequence belonging to a protein sequence other than said chemokine. This heterologous latter sequence should provide additional properties without impairing the pharmaceutical applicability. Examples of such additional properties are an easier purification procedure, a longer lasting half-life in body fluids, or extracellular localization. This latter feature is of particular importance for defining a specific group of fusion or chimeric proteins included in the above definition since it allows these monomeric variants to be localized in the space where not only where the isolation and purification of these peptides is facilitated, but also where CCL2 naturally interacts with other molecules.

Design of the moieties, ligands, and linkers, as well methods and strategies for the construction, purification, detection and use of fusion proteins are widely discussed in the literature (Nilsson J et al., 1997; "Applications of chimeric genes and hybrid proteins" Methods Enzymol. Vol. 326-328, Academic Press, 2000; WO 01/77137). Additional protein sequences which can be used to generate alternative forms of these obligate monomeric variants of homodimer-forming chemokines as defined above are the ones of extracellular domains of membrane-bound protein, immunoglobulin constant region (Fc region), multimerization domains, extracellular proteins, signal peptide-containing proteins, export signal-containing proteins. The choice of one or more of these sequences to be fused to the monomeric variant is functional to specific use of said agent. When the additional protein sequence, as in the case of the sequence of extracellular, export signal, or signal-peptide containing proteins, allows the monomeric variant to be secreted in the extracellular space, the agent can be more easily collected and purified from cultured cells in view of further processing or, alternatively, the cells can be directly used or administered.

The obligate monomeric variants of homodimer-forming chemokines defined above can be also used in other preferred forms, for example as active fractions, precursors, salts, derivatives, conjugates or complexes.

The term "fraction" refers to any fragment of the polypeptidic chain of the compound itself, alone or in combination with related molecules or residues bound to it, for example residues of sugars or phosphates, or aggregates of the original polypeptide or peptide. Such molecules can result also from other modifications which do not normally alter primary sequence, for example *in vivo* or *in vitro* chemical derivativization of peptides (acetylation or carboxylation), those made by modifying the pattern of phosphorylation (introduction of phosphotyrosine, phosphoserine, or phosphothreonine residues) or glycosylation (by exposing the peptide to enzymes which affect glycosylation e.g., mammalian glycosylating or deglycosylating enzymes) of a peptide during its synthesis and processing or in further processing steps.

The "precursors" are compounds which can be converted into the compounds of present invention by metabolic and enzymatic processing prior or after the administration to the cells or to the body.

The term "salts" herein refers to both salts of carboxyl groups and to acid addition salts of amino groups of the peptides, polypeptides, or analogs thereof, of the present invention. Salts of a carboxyl group may be formed by means known in the art and include inorganic salts, for example, sodium, calcium, ammonium, ferric or zinc salts, and the like, and salts with organic bases as those formed, for example, with amines, such as triethanolamine, arginine or lysine, piperidine, procaine and the like. Acid addition salts include, for example, salts with mineral acids such as, for example, hydrochloric acid or sulfuric acid, and salts with organic acids such as, for example,

acetic acid or oxalic acid. Any of such salts should have substantially similar activity to the peptides and polypeptides of the invention or their analogs.

The term "derivatives" as herein used refers to derivatives which can be prepared from the functional groups present on the lateral chains of the amino acid moieties or on the N- or C-terminal groups according to known methods. Such derivatives include
5 for example esters or aliphatic amides of the carboxyl-groups and N-acyl derivatives of free amino groups or O-acyl derivatives of free hydroxyl-groups and are formed with acyl-groups as for example alcanoyl- or aroyl-groups.

Useful conjugates or complexes of obligate monomeric variants of homodimer-
10 forming chemokines defined above can be generated, using molecules and methods known in the art of the interaction with receptor or other proteins (radioactive or fluorescent labels, biotin), therapeutic efficacy (cytotoxic agents), or improving the agents in terms of drug delivery efficacy, such as polyethylene glycol and other natural or synthetic polymers (Harris JM and Chess RB, 2003; Greenwald RB et al., 2003;
15 Pillai O and Panchagnula R, 2001). Residues can be used for attachment, provided they have a side-chain amenable for polymer attachment (i.e., the side chain of an amino acid bearing a functional group, e.g., lysine, aspartic acid, glutamic acid, cysteine, histidine, etc.). Alternatively, a residue at these sites can be replaced with a different amino acid having a side chain amenable for polymer attachment. Also, the
20 side chains of the genetically encoded amino acids can be chemically modified for polymer attachment, or unnatural amino acids with appropriate side chain functional groups can be employed. Polymer attachment may be not only to the side chain of the amino acid naturally occurring in a specific position of the antagonist or to the side chain of a natural or unnatural amino acid that replaces the amino acid naturally

occurring in a specific position of the antagonist, but also to a carbohydrate or other moiety that is attached to the side chain of the amino acid at the target position.

Polymers suitable for these purposes are biocompatible, namely, they are non-toxic to biological systems, and many such polymers are known. Such polymers may be hydrophobic or hydrophilic in nature, biodegradable, non-biodegradable, or a combination thereof. These polymers include natural polymers (such as collagen, gelatin, cellulose, hyaluronic acid), as well as synthetic polymers (such as polyesters, polyorthoesters, polyanhydrides). Examples of hydrophobic non-degradable polymers include polydimethyl siloxanes, polyurethanes, polytetrafluoroethylenes, polyethylenes, polyvinyl chlorides, and polymethyl methacrylates. Examples of hydrophilic non-degradable polymers include poly(2-hydroxyethyl methacrylate), polyvinyl alcohol, poly(N-vinyl pyrrolidone), polyalkylenes, polyacrylamide, and copolymers thereof. Preferred polymers comprise as a sequential repeat unit ethylene oxide, such as polyethylene glycol (PEG).

The preferred method of attachment employs a combination of peptide synthesis and chemical ligation. Advantageously, the attachment of a water-soluble polymer will be through a biodegradable linker, especially at the amino-terminal region of a protein. Such modification acts to provide the protein in a precursor (or "pro-drug") form, that, upon degradation of the linker releases the protein without polymer modification.

As a general procedure, the obligate monomeric variants of homodimer-forming chemokines defined above can be produced may be prepared by any procedure known in the art, including recombinant DNA-related technologies and chemical synthesis technologies.

Many books and reviews provides teachings on how to clone and produce recombinant proteins using vectors and prokaryotic (e.g. E. coli) or eukaryotic host

cells, such as some titles in the series "A Practical Approach" published by Oxford University Press ("DNA Cloning 2: Expression Systems", 1995; "DNA Cloning 4: Mammalian Systems", 1996; "Protein Expression", 1999; "Protein Purification Techniques", 2001).

5 Another embodiment according to the Invention is the nucleic acid sequence encoding for the obligate monomeric chemokine variant antagonist described herein.

The DNA sequence coding for the obligate monomeric variants of homodimer-forming chemokines can be inserted and ligated into a suitable episomal or non-homologously integrating vectors, which can be introduced in the appropriate host
10 cells by any suitable means (transformation, transfection, conjugation, protoplast fusion, electroporation, calcium phosphate-precipitation, direct microinjection, etc.) to transform them. Factors of importance in selecting a particular plasmid or viral vector include: the ease with which recipient cells that contain the vector, may be recognized and selected from those recipient cells which do not contain the vector; the number of
15 copies of the vector which are desired in a particular host; and whether it is desirable to be able to "shuttle" the vector between host cells of different species.

The vectors should allow the expression of the isolated or fusion protein including the antagonist of the invention in the prokaryotic or eukaryotic host cell under the control of transcriptional initiation / termination regulatory sequences, which are
20 chosen to be constitutively active or inducible in said cell. A cell line substantially enriched in such cells can be then isolated to provide a stable cell line.

For eukaryotic hosts (e.g. yeasts, insect or mammalian cells), different transcriptional and translational regulatory sequences may be employed, depending on the nature of the host. They may be derived from viral sources, such as adenovirus,
25 bovine papilloma virus, Simian virus or the like, where the regulatory signals are

associated with a particular gene which has a high level of expression. Examples are the TK promoter of the Herpes virus, the SV40 early promoter, the yeast gal4 gene promoter, etc. Transcriptional initiation regulatory signals may be selected which allow for repression and activation, so that expression of the genes can be modulated. The cells which have been stably transformed by the introduced DNA can be selected by also introducing one or more markers which allow for selection of host cells which contain the expression vector. The marker may also provide for phototrophy to an auxotrophic host, biocide resistance, e.g. antibiotics, or heavy metals such as copper, or the like. The selectable marker gene can either be directly linked to the DNA gene sequences to be expressed, or introduced into the same cell by co-transfection. Additional elements may also be needed for optimal synthesis of the proteins.

Host cells may be either prokaryotic or eukaryotic. Preferred are eukaryotic hosts, e.g. mammalian cells, such as human, monkey, mouse, and Chinese Hamster Ovary (CHO) cells, because they provide post-translational modifications to protein molecules, including correct folding or glycosylation at correct sites. Also yeast cells can carry out post-translational peptide modifications including glycosylation. A number of recombinant DNA strategies exist which utilize strong promoter sequences and high copy number of plasmids which can be utilized for production of the desired proteins in yeast. Yeast recognizes leader sequences in cloned mammalian gene products and secretes peptides bearing leader sequences (i.e., pre-peptides).

For long-term, high-yield production of a recombinant polypeptide, stable expression is preferred. For example, cell lines which stably express the polypeptide of interest may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may

be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells that successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type. A cell line substantially enriched in such cells can be then isolated to provide a stable cell line.

Mammalian cell lines available as hosts for expression are known in the art and include many immortalised cell lines available from the American Type Culture Collection (ATCC) including, but not limited to, Chinese hamster ovary (CHO), HeLa, baby hamster kidney (BHK), monkey kidney (COS), C127, 3T3, BHK, HEK 293, Bowes melanoma and human hepatocellular carcinoma (for example Hep G2) cells and a number of other cell lines. In the baculovirus system, the materials for baculovirus/insect cell expression systems are commercially available in kit form from, inter alia, the "MaxBac" kit (Invitrogen).

Examples of chemical synthesis technologies are solid phase synthesis and liquid phase synthesis. As a solid phase synthesis, for example, the amino acid corresponding to the C-terminus of the peptide to be synthesized is bound to a support which is insoluble in organic solvents, and by alternate repetition of reactions, one wherein amino acids with their amino groups and side chain functional groups protected with appropriate protective groups are condensed one by one in order from the C-terminus to the N-terminus, and one where the amino acids bound to the resin or the protective group of the amino groups of the peptides are released, the peptide chain is thus extended in this manner. Solid phase synthesis methods are largely classified by the tBoc method and the Fmoc method, depending on the type of protective group used. Typically used protective groups include tBoc (t-

butoxycarbonyl), Cl-Z (2-chlorobenzyloxycarbonyl), Br-Z (2-bromobenzyloxycarbonyl), Bzl (benzyl), Fmoc (9-fluorenylmethoxycarbonyl), Mbh (4,4'-dimethoxydibenzhydryl), Mtr (4-methoxy-2,3,6-trimethylbenzenesulphonyl), Trt (trityl), Tos (tosyl), Z (benzyloxycarbonyl) and Cl₂-Bzl (2,6-dichlorobenzyl) for the amino groups; NO₂ (nitro) and Pmc (2,2,5,7,8-pentamethylchromane-6-sulphonyl) for the guanidino groups; and tBu (t-butyl) for the hydroxyl groups). After synthesis of the desired peptide, it is subjected to the de-protection reaction and cut out from the solid support. Such peptide cutting reaction may be carried with hydrogen fluoride or trifluoromethane sulfonic acid for the Boc method, and with TFA for the Fmoc method.

10 Totally synthetic CCL2 proteins are disclosed in the literature (Brown A et al., 1996).

Purification of synthetic or recombinant monomeric variants of homodimer-forming chemokines defined above can be carried out by any one of the methods known for this purpose, i.e. any conventional procedure involving extraction, precipitation, chromatography, electrophoresis, or the like. A further purification procedure that may be used in preference for purifying the protein of the invention is affinity chromatography using monoclonal antibodies or affinity groups, which bind the target protein and are produced and immobilized on a gel matrix contained within a column. Impure preparations containing the proteins are passed through the column. The protein will be bound to the column by heparin or by the specific antibody while the impurities will pass through. After washing, the protein is eluted from the gel by a change in pH or ionic strength. Alternatively, HPLC (High Performance Liquid Chromatography) can be used. The elution can be carried using a water-acetonitrile-based solvent commonly employed for protein purification.

15

20

The monomeric variant of a homodimer-forming chemokine can be used in the pharmaceutical composition for the treatment or prevention of autoimmune,

25

inflammatory, or infectious diseases. In particular, the results provided in the examples regarding an animal model for multiple sclerosis shows that these monomeric variants can be used in the pharmaceutical composition for the treatment or prevention of multiple sclerosis.

5 Another aspect of the invention is a monomeric variant of a homodimer-forming chemokine, wherein said variant results from at least an amino acid substitution that alters the pattern of hydrogen bonds at the dimerization interface of said chemokine, used as a medicament. Examples of such variants are disclosed herein as CCL2-P8A and CCL2*-P8A. However, the teaching of the Invention allows the identification,
10 production and testing of similar molecules on the basis of the sequence and the activities of other chemokines.

In particular, these monomeric variants can be chosen from:

- a) CCL2-P8A (SEQ ID NO: 2);
 - b) CCL2*-P8A (SEQ ID NO: 4);
 - 15 c) An active mutant of (a) or (b); or
 - d) A polypeptide comprising (a), (b), or (c), and an amino acid sequence belonging to a protein sequence other than said chemokine;
- as well as the corresponding monomeric variants in the form of their active fractions, precursors, salts, derivatives, complexes or conjugates.

20 Another aspect of the invention is a pharmaceutical composition containing a monomeric variant of a homodimer-forming chemokine as active ingredient, wherein said variant result from at least an amino acid substitution that alters the pattern of hydrogen bonds at the dimerization interface of said chemokine, such as CCL2-P8A or CCL2*-P8A, optionally in the forms defined above (such as active mutants,

polypeptides comprising them, or conjugates) as well as DNA coding or cells expressing them.

The pharmaceutical compositions of the invention may contain suitable pharmaceutically acceptable carriers, biologically compatible vehicles and additives
5 that are suitable for administration to an animal (for example, physiological saline) and eventually comprising auxiliaries (like excipients, stabilizers or diluents) that facilitate the processing of the active compounds into preparations that can be used pharmaceutically. The pharmaceutical compositions may be formulated in any acceptable way to meet the needs of the mode of administration. For example, the use
10 of biomaterials and other polymers for drug delivery, as well the different techniques and models to validate a specific mode of administration are disclosed in literature (Luo B and Prestwich GD, 2001; Cleland JL et al., 2001).

An "effective amount" refers to an amount of the active ingredients that is sufficient to affect the course and the severity of the disease, leading to the reduction
15 or remission of such pathology. The effective amount will depend on the route of administration and the condition of the patient.

"Pharmaceutically acceptable" is meant to encompass any carrier, which does not interfere with the effectiveness of the biological activity of the active ingredient and that is not toxic to the host to which is administered. For example, for parenteral
20 administration, the above active ingredients may be formulated in unit dosage form for injection in vehicles such as saline, dextrose solution, serum albumin and Ringer's solution. Carriers can be selected also from starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk, glycerol, propylene

glycol, water, ethanol, and the various oils, including those of petroleum, animal, vegetable or synthetic origin (peanut oil, soybean oil, mineral oil, sesame oil).

Any accepted mode of administration can be used and determined by those skilled in the art to establish the desired blood levels of the active ingredients. For example, administration may be by various parenteral routes such as subcutaneous, intravenous, intradermal, intramuscular, intraperitoneal, intranasal, transdermal, oral, or buccal routes. Parenteral administration can be by bolus injection, by gradual perfusion over time or controlled release dosage forms, including depot injections, osmotic pumps, and the like, for the prolonged administration of the polypeptide at a predetermined rate, preferably in unit dosage forms suitable for single administration of precise dosages. Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions, which may contain auxiliary agents or excipients known in the art, and can be prepared according to routine methods. In addition, suspension of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides. Aqueous injection suspensions that may contain substances increasing the viscosity of the suspension include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers. Pharmaceutical compositions include suitable solutions for administration by injection, and contain from about 0.01 to 99 percent, preferably from about 20 to 75 percent of active compound together with the excipient. Compositions that can be administered rectally include suppositories.

It is understood that the dosage administered will be dependent upon the age, sex, health, and weight of the recipient, kind of concurrent treatment, if any, frequency

of treatment, and the nature of the effect desired. The dosage will be tailored to the individual subject, as is understood and determinable by one of skill in the art. The total dose required for each treatment may be administered by multiple doses or in a single dose. The pharmaceutical composition of the present invention may be administered
5 alone or in conjunction with other therapeutics directed to the condition, or directed to other symptoms of the condition. Usually a daily dosage of active ingredient is comprised between 0.01 to 100 milligrams per kilogram of body weight. Ordinarily 1 to 40 milligrams per kilogram per day given in divided doses or in sustained release form is effective to obtain the desired results. Second or subsequent administrations can be
10 performed at a dosage, which is the same, less than, or greater than the initial or previous dose administered to the individual.

Another object of the present invention is a method for treating or preventing autoimmune or inflammatory (such as multiple sclerosis), or infectious diseases comprising the administration of an effective amount of a monomeric variants of
15 homodimer-forming chemokines, wherein said variant result from at least an amino acid substitution that alters the pattern of hydrogen bonds at the dimerization interface of said chemokine. Examples of such monomeric variants that can be used in such methods are:

- a) CCL2-P8A (SEQ ID NO: 2);
- 20 b) CCL2*-P8A (SEQ ID NO: 4);
- c) An active mutant of (a) or (b);
- d) A polypeptide comprising (a), (b), or (c), and an amino acid sequence belonging to a protein sequence other than said chemokine;
as well as the corresponding monomeric variants in the form of their active
25 fractions, precursors, salts, derivatives, complexes or conjugates.

A non-limitative list of examples for autoimmune, inflammatory, or infectious diseases mentioned above regarding the uses, the variants, and the methods of the invention are the following: arthritis, rheumatoid arthritis (RA), psoriatic arthritis, osteoarthritis, systemic lupus erythematosus (SLE), systemic sclerosis, scleroderma, polymyositis, glomerulonephritis, fibrosis, fibrosis, allergic or hypersensitivity diseases, dermatitis, asthma, chronic obstructive pulmonary disease (COPD), inflammatory bowel disease (IBD), Crohn's diseases, ulcerative colitis, multiple sclerosis, cancer, septic shock, viral or HIV infections, transplantation, , airways inflammation, graft-versus-host disease (GVHD) and atherosclerosis.

10 The therapeutic applications of the polypeptides of the invention and of the related reagents can be evaluated (in terms of safety, pharmacokinetics and efficacy) by the means of the *in vivo* or *in vitro* assays making use of animal cell, tissues and models (Coleman RA et al., 2001; Li AP, 2001; Methods Mol. Biol. vol. 138, "Chemokines Protocols", edited by Proudfoot A et al., Humana Press Inc., 2000; 15 Methods Enzymol, vol. 287 and 288, Academic Press, 1997).

Another aspect of the Invention are methods for screening for obligate monomeric antagonist chemokine variants described herein comprising:

- a) making single point mutations in CCL2 that block its ability to dimerize;
- b) identifying said variants that bind to the receptor and show agonistic 20 properties *in vitro*;
- c) identifying said variants from the group identified in (b) above that are further characterized as inhibiting peritoneal cell recruitment.

This evaluation of these properties can be made using techniques known in the art, and shown in the examples, applying a molecule known to induce inflammation and

peritoneal cell recruitment, for example a chemokine such as CCL2 itself, thioglycollate, or ovalbumin.

The present invention has been described with reference to the specific embodiments, but the content of the description comprises all modifications and substitutions, which can be brought by a person skilled in the art without extending
5 beyond the meaning and purpose of the claims.

The invention will now be described by means of the following Examples, which should not be construed as in any way limiting the present invention.

10

EXAMPLES

Example 1: cloning, expression, and purification of the recombinant proteins

Mature CCL2 (fig. 1; SEQ ID NO: 1) and the CCL2-P8A mutant proteins (fig. 1; SEQ ID NO: 2), mature CCL2* (fig. 1; SEQ ID NO: 3) and the CCL2*-P8A mutant
15 proteins (fig. 1; SEQ ID NO: 4) were generated and expressed as recombinant proteins in *E.coli* as described in the literature (Paavola CD et al, 1998) on the basis of the sequence of the mature form of human CCL2/MCP-1, corresponding to the segment 24-99 of the precursor molecule. For CCL2* and CCL2*-P8A, the substitution of a Methionine with an Isoleucine in position 64 improves the purity and homogeneity of
20 the mutants by eliminating the formation of species containing methionine-sulfoxide at position 64.

Briefly, the genes for CCL2 for CCL2* was constructed by standard gene synthesis techniques with optimal codon usage for expression in *E. coli* and a codon for methionine added at the 5' end of the sequence encoding mature human CCL2 or
25 CCL2*. Mutant constructs including Alanine in position 8 were made by polymerase

chain reaction mutagenesis of the CCL2 or CCL2* template and cloned into a pET3a vector (Novagen) between the Xho I and Nde I sites.

Plasmids encoding CCL2*-based proteins were used to transform TAP302 cells, which are BL21 pLys S cells engineered with a thioredoxin reductase knockout to make the intracellular redox potential more conducive to disulfide bond formation. Using this
5 strain, disulfide bonds appear to be formed in the cell, eliminating the need for a refolding step.

All constructs were obtained and controlled by standard molecular biology technologies (PCR mutagenesis and amplification, DNA sequencing, restriction
10 digestion). One of the clones containing the correct sequence of MCP-1 (P8A) was subsequently used to produce the protein in *E. coli*. The plasmids were also used to transform BL21 Star™ (DE3) (Invitrogen Cat n° C6010-03) or BL21 DE3 (Novagen Cat n° 69387-3).

CCL2, CCL2* and CCL2*-P8A are expressed and purified as described in the
15 original article, using a sonication step, a lysis step and a chromatographic step (SP-Sepharose column; elution with a gradient of NaCl in 10 mM K₂PO₄, pH 7.5). Peak fractions were pooled and further purified by reversed-phase HPLC (C18 column with a 5- μ m particle size and 300-Å pore size). Proteins were eluted using a gradient of increasing acetonitrile containing 0.1% trifluoroacetic acid; typically, proteins eluted at
20 34 \pm 5% acetonitrile. They were then lyophilized, dissolved at 1 mg/ml in 35 mM Tris, pH 8, reacted with 15 μ g of aminopeptidase (Peptotech, Rock Hill, NJ)/1 mg of protein for 36 h at room temperature, and repurified by reversed-phase HPLC. Aminopeptidase treatment removes only the N-terminal methionine, generating either a N-terminal Glutamin or N-terminal pyro-Glutamic acid (there is no effect on biological activity due
25 to this difference), as observed by N-terminal sequencing of the recombinant protein.

The protein was then lyophilized, redissolved in water at 1-5 mg/ml, and stored in small aliquots at 80 °C.

Larger amounts of the recombinant proteins, and in particular of CCL2-P8A, were obtained also by using an alternative protocol designed for purifications starting from the cell pellet obtained from large fermentation cultures of E. coli strains producing these proteins. Generally a 5 L fermentor produces approximately 200 g wet weight cell pellet, and a fermentor of 50 L produces 1.8 kg wet weight cell pellet. The purification procedure described here treats 200 g wet weight cell pellet.

The cell pellet was thawed and 3 ml of breakage buffer per gram wet weight (50 mM Tris/HCl buffer, pH 8.0 (Cat. 20092391, Biosolve) containing 10 mM MgCl₂ (Cat. 63065, Fluka), 5 mM Benzamidine/HCl (Cat. 12073, Fluka), 1 mM 1,4 DL dithiothreitol (DTT) (Cat. 43819, Fluka), 1 mM phenylmethylsulfonyl fluoride (PMSF) (Cat. 78830, Fluka)* 20 mg/L DNase (Fluka) (Cat. DN-25, Sigma) was added. The suspension was homogenized with a Polytron to obtain a good homogenate devoid of fragments or clumps. All manipulations were carried out at 4 °C. The homogenized bacterial suspension was transferred to a French Press cell mechanical disrupter (differential pressure). The number of passages was typically 2-4 under 1500 bar. The cell break was monitored by SDS-PAGE stained with Coomassie blue.

The lysate was dispensed into GSA centrifuge tubes and centrifuged at 10'000 rpm with the Sorval RC5C (16'300 x g) for 90 minutes at 4°C. After centrifugation, the supernatant was discarded after confirmation by SDS-PAGE analysis that no soluble protein of interest could be detected in the supernatant. The pellet was removed with a spatula from the centrifuge tube and transferred it to a pre-weighed beaker to weigh the pellet. The pellet was washed with de-ionized water by adding 5 ml of water per gram of pellet in a beaker and stirred for 30 min at 4°C with a magnetic stirrer. The

suspension was centrifuged in GSA centrifuge tubes at 10'000 rpm with the Sorval RC5C centrifuge (16'300 x g) for 60 min at 4°C. The wash step was repeated 3 times. After each centrifugation, the supernatant was discarded after confirmation by SDS-PAGE analysis that no soluble protein of interest could be detected in the supernatant.

5 The pellet was solubilised in inclusion body freshly prepared extraction buffer (100 mM Tris/HCl buffer, pH 8.0 (Cat. 20092391, Biosolve) containing 1 mM 1,4 DL dithiotreitol (DTT) (Cat. 43819, Fluka) and 6 M Guanidium/HCl (Cat. 50950, Fluka)) in a ratio of 100 ml of buffer for 25 g cell pellet using a Polytron. The solution was heated for 30 min at 60°C and stirred to ensure monomerisation, then cooled to room
10 temperature. The homogenate was dispensed into Ti45 centrifuge tubes and ultracentrifuged at 35'000 rpm with the Beckman L-60 (100'000 x g). The supernatant was filtered with a 0.8-0.2 mm filter (SpiralCap (Cat. 12069, PALL), analyzed by SDS-PAGE and quantified using the Coomassie protein assay reagent (Pierce) following the protocol supplied with the kit.

15 The recombinant protein of interest is captured on a FineLine 35 Pilot column containing Source 30 RPC resin packed following the supplier's instructions (Amersham Pharmacia). After use, the column is regenerated following the cleaning procedure supplied by the manufacturer (Amersham Pharmacia). For 100 grams of cell pellet, a column of 3.5 cm diameter x 23 cm height giving a total of 220 ml (equivalent
20 to a 1 Column Volume) of Source 30 RPC is packed. The column was installed on an AKTA FPLC (Amersham Pharmacia). The flow rate was 10 ml/min, and the maximum pressure was 1 MPa. Before loading the sample the column was washed with deionised water for 2 Column Volume (440 ml). After washing, the column was equilibrated with 5 CVs (Column Volumes) of equilibration buffer (100 mM Tris/HCl

buffer, (Cat. 20092391, Biosolve) adjusted to pH 7.5 with fuming 37% HCl (Cat. 84426, Fluka).

The dissolved Ibs (Inclusion Bodies) in Guanidium/HCl were loaded onto the column at a flow rate of 5 ml/min. The column was then washed with 5 Column Volume of equilibration buffer followed by 5 CVs of buffer A (0.1 % TFA Trifluoroacetic acid (Cat. 28904, Pierce) and 99.9 % Distilled water). The protein is eluted using a linear gradient from 0% to 100% of buffer B (0.1% TFA Trifluoroacetic acid (Cat. 28904, Pierce) 9.9 % distilled water, 90% Acetonitrile (UN1648, Baker), over 10 CV with a flow rate of 10 ml/min, and a 1 MPA pressure limit. 10 ml fractions were collected. All peaks detected were analysed by SDS-PAGE, HPLC and quantified by UV-spectroscopy. The fractions containing the protein of interest were pooled and the amount measured by UV-spectroscopy.

The protein was renatured by a 10-fold dilution into renaturation buffer (100 mM Tris/HCl buffer, pH 8.0 (Cat. 20092391, Biosolve) containing 0.1 mM Reduced Glutathione (Cat.G-4251, Sigma) and 0.01 mM Oxidised Glutathione (Cat.120 000 250, Acros Organic) to obtain a final concentration of approximately 0.1 mg/ml. The pool of Source 30 RPC was added dropwise into the renaturation buffer. If the volume is large, this can be carried out using a peristaltic pump. The solution was stirred overnight at 4°C. The solution often appears cloudy due to the precipitation of protein that has not renatured. Final concentrations ranging from 0.1 to 0.4 mg/ml in the renaturation buffer yielded equivalent amounts of renatured protein (40 to 50%). The pH and acetonitrile in the starting material does not affect the renaturation step. Renaturation can be followed by HPLC to follow the refolding.

The renaturation solution was filtered using a High Flow peristaltic pump with a double filter, consisting of a prefilter of 0.8 mm followed by a 0.22 mm. The clarified

solution was then concentrated by cation exchange on Hiload SP Sepharose HP after quantification by a UV-spectrum. The size of the ion exchange column depends of the amount of protein. For < 500 milligrams, a 16/10 column (1 CV = 20 ml) was used; 500 - 1000 mg, a 26/10 column (1 CV = 50 ml) and for 1 – 2 grams, a 50/5 column (1 CV= 5 100 ml).

The column was packed according to the supplier's instructions (Amersham Pharmacia). The column was washed with 2 CVs of deionised water, and then equilibrated with 4 CVs of cation exchange buffer A (50 mM Acetic acid (Fluka), adjusted to pH to 4.5 with NaOH (Cat.71690 Fluka). The solution was adjusted to pH 10 4.5 with acetic acid and the conductivity adjusted to <10 mS. After loading the protein solution at the flow rate recommended by the supplier for the column chosen, the column was washed with 5 CVs of buffer A. The protein was eluted with a linear gradient from 0% to 100% of buffer B (buffer A, containing 2 M NaCl (Cat. 71380, Fluka) over 20 CVs. The fraction size was determined by the column size. All peaks 15 detected were analysed by SDS-PAGE, HPLC and quantified by UV-spectroscopy. After analysis, the fractions containing AS900652 were pooled, quantified by UV-spectrum and analysed by HPLC.

The removal the N-terminal Methionine was performed enzymatically using Methionine Aminopeptidase (MAP), followed by a purification step. Briefly, the sample 20 was first dialysed using membrane tubing with a cut-off of 3.5 kD into cleavage buffer (35 mM Tris/HCl buffer, pH 7.5 (Cat. 20092391, Biosolve). The dialysis buffer was changed three times over 24 hours. Methionine Aminopeptidase (MAP) was added to the protein solution at a ratio of 1:10000 (w:w, enzyme:protein). The digestion was performed at room temperature for 48 hours. The digested protein was then purified by

cation exchange is carried out as described above. The protein was > 98 % pure as estimated by SDS-PAGE.

The AKTA purifier system (Pharmacia) was used to further purify the desired protein. The system was cleaned for 1 hour with 1 M NaOH, washed with sterile water, and equilibrated in buffer filtered with a 0.22 mm filter (0.1% TFA (Trifluoroacetic acid) (Cat. 28904, Pierce) 99.9 % Distilled water). The protein was desalted using G-25 fine Sepharose XK50/30 column. The column was washed with 1 CV of 1 M NaOH followed by 4 CVs of sterile water and then equilibrated with 5 CVs 0.1% TFA. For optimal desalting conditions, 50-to100 ml samples are desalted on the 450 ml G-25 fine Sepharose. For volumes larger than 100 ml, the desalting step is repeated. The sample was filtered with 0.22 mm filter before loading. The column is eluted with 1.5 CV 0.1% TFA with a flow rate of 10 ml/min and a maximum pressure of 1 MPa. Fractions of 10 ml are collected into sterile tubes. After analysis, the fractions containing the protein are pooled, quantified by UV-spectrum under sterile conditions and analysed by HPLC, SDS-PAGE and mass spectrometry.

The remaining contaminants are removed by using a preparative reverse phase chromatography (RPC) on DeltaPrep HPLC (WATERS). The sample was acidified to 0.1 % TFA and loaded onto a Vydac C8 RPC (Cat. 208TB101522, Vydac) equilibrated in buffer A (0.1 % TFA Trifluoroacetic acid (Cat. 28904, Pierce) and 99.9 % distilled water). The protein was eluted using a linear gradient from 0% to 100% of buffer B (0.1% TFA Trifluoroacetic acid (Cat. 28904, Pierce), 99% Acetonitrile (UN1648, Baker), over 10 CV with a flow rate of 25 ml/min, and a 700 bar pressure limit. All peaks detected were analysed by SDS-PAGE, HPLC and quantified by UV-spectroscopy. After analysis, the fractions containing AS900652 were pooled, quantified by UV-

spectrum, aliquoted as required and lyophilised. The protein is stored at -20°C or -80°C .

The recombinant protein is quantified by UV spectroscopy using a UV-VIS spectrophotometer (Uvikon system, KONTRON). A quartz cuvette (QS 1.000, HELLMA) was used for the buffer reference and the other cuvette contains the sample. A scan from 350 nm to 240 nm was measured and the absorptions at 280 nm was used to determine the quantification according the extinction coefficient obtained from the amino acid composition using ProtParam, Expasy. The value used was 1.1 for a solution of 1 mg/ml for the oxidised protein containing disulfide bonds.

SDS-PAGE analyses were carried out using NuPAGE 10% gel (Cat.NP0301, Invitrogen). The sample was diluted 2 fold in sample buffer (cat. LC2676, Invitrogen) and heated for 5 minutes at 95°C . The Benchmark protein ladder was used as molecular weight standards. 10 μl of molecular weight standard solution and 20 μl of protein sample were loaded in the appropriate wells. The electrophoresis running buffer was MES (Cat.NP0002, Invitrogen). The migration was carried out according to the supplier's instructions of 200 V, 12 mA and 25 W for 35 minutes (PowerEase500, Invitrogen).

NuPAGE gels were stained with 0.1% R250 Coomassie blue in 10% acetic acid, 30% methanol in distilled water for 30 minutes, and de-stained in 10% acetic acid, 30% methanol in distilled water under slow rocking motion until background level is not coloured. The gel was then washed several times in water prior to embedding in drying solution (Invitrogen) sandwiched between two cellophane paper sheets (Invitrogen) for 10 minutes and then mounted in the miniature press allowing the gel to be stored as a fine sheet. Alternatively the gel was stained with the SimplyBlue Safe stain protocol (Cat.LC6065, Invitrogen).

The Alliance HPLC system supplied by WATERS was used with an analytical C8 Aquapore RP-300 7m (0.2 cm diam. x 22 cm) equilibrated in 0.1 % TFA. 10-50 mg previously acidified 0.1% final TFA were injected. Proteins were eluted with a gradient of 25 to 50 % acetonitrile over 20 CVs.

5 The identity of the recombinant protein was confirmed by mass spectral analyses and N-terminal sequence analysis. The correct N-terminal sequence QPDAINAAVT was obtained for the purified material.

A mass of 8655 Da was obtained for the main species corresponding to the theoretical mass of the protein chain with 2 disulfide bonds. A second species with a
10 mass lower by 17 Da was also observed corresponding to the modification of the N-terminal Glutamine residue into a pyroglutamic acid. The presence of this modification has no influence on the activity of the protein.

Example 2: Cell Based Assays

15 Materials and methods

Assays for chemokine-induced peritoneal cellular recruitment

Female Balb/C mice (Janvier, France) of 8 to 12 weeks were housed under normal animal holding conditions with a standard 12-h light/dark cycle and free access to food and water. Groups composed of 3-6 mice were injected intraperitoneally with
20 200 µl of saline (sterile LPS-free NaCl 0.9% (w/v) or of this solution containing CCL2 or CCL2-P8A at 10 µg per injection. For studies investigating the inhibitory effects of CCL2-P8A on CCL2-induced peritoneal cell recruitment, these molecules were administered intraperitoneally 30 minutes before the intraperitoneal injection of CCL2. All the molecules were administered at the concentration and in buffer above indicated

(saline). The mice were sacrificed on at 4 hours after the CCL2 or CCL2-P8A final injection.

The assay for thioglycollate-induced peritoneal cell recruitment has been published (Mishell B, 1980). Briefly, thioglycollate medium was prepared by suspending
5 30 g of dehydrated thioglycollate medium (Becton Dickinson) in 1 liter of cold distilled water, then heated until boiling to dissolve the powder completely. The medium was then aliquoted into 100 ml bottles and autoclaved. After cooling, the medium was stored in the dark at room temperature for at least one month. Cellular recruitment was induced by intraperitoneal injection of mice in groups of 3 with 200 μ l of a 3% solution
10 of thioglycollate on Day 1, 30 minutes after CCL2*-P8A administration. CCL2* was administered daily thereafter for 3 days (Days 2, 3 and 4). Dexamethasone (Sigma) was used as a positive control and administered at 10 mg/kg intraperitoneally. The mice were sacrificed on Day 5.

Peritoneal lavages to assess cell recruitment in the previous assays were
15 performed as follows. Mice were sacrificed by asphyxiation with rising concentrations of CO₂ in a plexiglass box. Skin was cleaned with 70% ethanol. The outer layer of skin was removed, exposing the peritoneal membrane. The peritoneal cavity was lavaged 3 times with 5 ml ice cold PBS (phosphate buffered saline) and fluid was pooled in a 15 ml polystyrene Falcon tube (Becton Dickinson) on ice. Each lavage was accompanied
20 with a light massage of the peritoneal cavity. Lavage fluid was centrifuged at 425xg, the supernatant discarded and the resultant cell pellet was resuspended by gentle multiple pipetting in 1 ml PBS. 10 μ l cell suspension was stained with 90 μ l trypan blue and total cell counts were enumerated with a Neubauer haemocytometer by counting 4 areas each of 1 mm². The mean of the 4 counts was taken, multiplied by the dilution factor of
25 10, and multiplied again by 10 to give the number of cells per μ l, according to the

directions for use accompanying the haemocytometer. Finally the total value was multiplied by 1000 (to equal 1 ml) to arrive at the total cell number recovered.

Results

Recombinant mature human CCL2/MCP-1 and the mutants called CCL2* and the corresponding obligate monomer mutants called CCL2-P8A and CCL2*-P8A (fig. 1) were expressed in *E coli*.

The literature clearly shows that P8A mutation in CCL2 blocks the formation of CCL2 dimers, without affecting the binding to cells expressing the receptor or to a viral receptor-like protein, but also without showing the activities of a known CCL2 antagonist in relevant assays (see (1+9-76)MCP-1 in table I of Paaavola CD et al., 1998; Alexander JM et al., 2002)

The obligate monomeric form of CCL2 presents specific and unexpected properties in assays performed in cell-based assays. In the peritoneal cell recruitment assay, CCL2*-P8A and CCL2-P8A are unable to recruit cells compared with natural CCL2 (Figure 2). Moreover, these molecules are able, in a dose dependent manner, to inhibit CCL2-induced (Figure 3A) and thioglycollate-induced macrophage recruitment (Figure 3B). In the latter assay, CCL2-P8A appears as effective as the positive control (dexamethasone, a known anti-inflammatory compound).

20 **Example 3: CCL2-P8A properties in animal models for diseases**

Materials and methods

Ovalbumin-induced lung inflammation model

The ovalbumin-induced lung inflammation model was performed as published (Blyth DI et al., 1996). Groups of 6 mice were sensitised by an intraperitoneal injection of 10 µg chicken egg albumin precipitated in 2 mg aluminium hydroxide 2% (Serva) in a

total volume of 200 μ l, which were prepared by mixing 25 μ l ovalbumin (2mg/ml), 250 μ l aluminium hydroxide in 725 μ l LPS-free 0.9% NaCl (saline) and precipitated 3-4 hours at 4°C. Fifteen days after sensitisation, mice were treated and challenged in groups of 6 mice with the intranasal administration of 15 μ g ovalbumin in 50 μ l saline, under inhaled anaesthesia (Isoflurane) daily from day 15 to 19. CCL2-P8A (200 μ l, 10 μ g per intraperitoneal injection) was administered 30 minutes before each challenge. Peritoneal lavages to assess cell recruitment and cell counts were performed as described above in Example 2.

EAE (Experimental Autoimmune Encephalomyelitis) model

10 C57Bl/6 mice from Charles River Italy (the selected strain has documented susceptibility to EAE; Sahrbacher UC et al., 1998.) are immunized (day=0) by injecting s.c. in the left flank 0.2 mL of an emulsion composed of 200 μ g MOG₃₅₋₅₅ peptide (Neosystem, Strasbourg, France) in Complete Freund's Adjuvant (CFA, Difco, Detroit, U.S.A.) containing 0.5 mg of *Mycobacterium tuberculosis*. Immediately after, they receive an i.p. injection of 500 ng pertussis toxin (List Biological Lab., Campbell, CA, U.S.A.) dissolved in 400 μ L of buffer (0.5 M NaCl, 0.017% Triton X-100, 0.015 M Tris, pH=7.5). On day 2, the animals are given a second i.p. injection of 500 ng pertussis toxin. On day 7, the mice receive a second dose of 200 μ g of MOG₃₅₋₅₅ peptide in CFA injected s.c. in the right flank. Starting approximately from day 8-10, this procedure results in a gradually progressing paralysis, arising from the tail and ascending up to the forelimbs. Starting from day 7 the animals are individually examined for the presence of paralysis by means of a clinical score as follows:

0 = no sign of disease

0.5 = partial tail paralysis

25 1 = tail paralysis

1.5 = tail paralysis + partial unilateral hindlimb paralysis

2 = tail paralysis + hindlimb weakness or partial hindlimb paralysis

2.5 = tail paralysis + partial hindlimb paralysis (lowered pelvi)

3 = tail paralysis + complete hindlimb paralysis

5 3.5 = tail paralysis + complete hindlimb paralysis + incontinence

4 = tail paralysis + hindlimb paralysis + weakness or partial paralysis of forelimbs

5 = moribund or dead

The treatment with compounds or vehicle starts for each animal at day 7 post immunization and is continued for 21 consecutive days (10-12 animals per treatment
10 group). Interferon beta and CCL2-P8A were administered s.c. or i.p., respectively, once a day solubilized in 10 ml/kg PBS at the doses indicated in the figure.

Delayed Contact Hypersensitivity Model

The mouse ear-swelling test to measure contact hypersensitivity was performed as previously described (Garrigue JL et al., 1994). Briefly, mice were pre-sensitized
15 topically by applying 25 μ l of 0.5% 2,4-dinitrofluorobenzene (DNFB; Sigma Chemical Co.) solution in acetone/ olive oil (4:1) to the shaved abdomen. Five days later, 20 μ l of 0.2% DNFB in the same vehicle was applied to the right ears, and vehicle alone to the left ears. Mice (n=6 per group) were treated daily on Day 5 with an intraperitoneal administration of 0.05, 0.5 or 5 mg/kg (1,10 or 100 micrograms/mouse, respectively) of
20 CCL2-P8A, Dexamethasone (1 mg/kg), or PBS only in the control group. The treatment was administered 30 minutes prior to the DNFB challenge. Ear thickness was measured with a dial thickness gauge (Mitutoyo Corp.), Ear swelling was estimated by subtracting the pre-challenge from the post-challenge value, and by further subtracting any swelling detected in the vehicle-challenged contralateral ear.

25 Results

The potential therapeutic activities of CCL2-P8A as chemokine antagonist have been tested in animal models for inflammatory and autoimmune diseases.

CCL2*-P8A was tested in a disease model, the ovalbumin-induced lung inflammation. In this classic model for allergic lung inflammation, the mice are sensitised with ovalbumin, with an adjuvant of aluminium hydroxide during sensitisation phase to boost the immune response, and then challenged by intranasal administration of ovalbumin over a period of 5 consecutive days, wherein CCL2-P8A was administered intraperitoneally throughout this phase. Also in this case, CCL2-P8A was capable to inhibit cell recruitment (figure 4).

In a second model, CCL2-P8A was tested in the EAE (Experimental Autoimmune Encephalomyelitis) model, a well known model for multiple sclerosis that has been used to validate antagonists of chemokines (including CCL2) for the treatment of this autoimmune, inflammatory demyelinating disease of the human central nervous system (Mahad DJ and Ransohoff RM, 2003; Izikson L et al., 2002). CCL-P8A was tested in animals showing either mild or severe level of the disease, as evaluated by clinical score, following the treatment of the EAE-inducing compounds. Each of the two groups of animals were divided in five subgroups: three of them were treated with different amounts of CCL2-P8A, and the two others were used as either negative control (treated with vehicle only) or as positive control (treated with Interferon -beta, a common therapeutic product for the treatment of multiple sclerosis). The evolution of the state of the animals was compared on the basis of the clinical score measured during the treatment period (21 days). In both disease models, the administration of CCL2 -P8A (at a dosage down to 0.15 mg/kg) improves the state of the animals in a statistically significant manner. The observed decrease of the clinical score using CCL-P8A is at least comparable to that observed when interferon beta is used as treatment (figure 5).

Another disease model, the contact hypersensitivity model was used to evaluate the potential therapeutic efficacy of CCL2-P8A on skin inflammation mediated by T cells. Contact hypersensitivity (CHS) is a Langerhans cell (LC)-dependent, T cell-mediated cutaneous immune response, reflecting a culmination of LC activities *in vivo* (uptake of epicutaneous antigens, migration into lymph nodes, and presentation of antigens to naive T cells). The model is well established for characterization of compound for dermatological indications like psoriasis and allergic contact dermatitis (Xu H et al. 1996). It involves a sensitisation phase and a subsequent challenge with an antigen, resulting in a skin inflammation with formation of edema and cellular infiltrates in the skin. The edema can be measured by caliper at the challenged site (ear of the mice). The involvement of chemokines, and of CCL2 in particular, in the development of this excessive response disease have been demonstrated (Mitsui G et al., 2003; Mizumoto N et al., 2001). Intraperitoneal administration of CCL2-P8A 30 minutes before a challenge with the antigen (DNFB, in this case) results in a decrease of the swelling comparable to that observed using a known anti-inflammatory compound (Dexamethasone) one day after the treatment. Control mice were obtained by challenging them with the antigen, but with or without previous sensitisation, so that T cell dependent inflammation and edema is formed or not (figure 6).

Therefore, a monomeric variant of a homodimer-forming CC-chemokine, wherein said variant result from at least an amino acid substitution that alters the pattern of hydrogen bonds at the dimerization interface, are inhibitors of chemokine-mediated cell recruitment in *in vivo* cell recruitment assays as well as in animal models for human diseases, implying that this is a novel strategy for generating chemokine variants which can be used for preparing pharmaceutical compositions and in therapeutic methods.

Example 4: Alternative forms of CCL2-P8A

Alternative forms of the chemokine variants disclosed above can be generated by introducing mutations known in the art as improving specific features.

One or more single amino acid substitutions and/or additions can be introduced in different position of CCL2-P8A (figure 7A). CCL2-P8A can be expressed as a mature protein missing the natural Glutamine N-terminal residue, or by adding an additional small residue (such as Alanine or Glycine) at the N-terminus before Glutamine, so that this residue does not remain exposed and does not get converted spontaneously into the pyroglutamate form (Gong J and Clark-Lewis I, 1995). CCL-P8A can also be mutated in way that a fifth Cysteine is available to allow specific PEGylation reactions. These PEGylation sites can be integrated at the level of either an internal amino acid (for example at Asparagin 14 or 17, and even at position 8, so that a single modification can allow both monomerization and PEGylation) or of the C-terminus (by directly adding a Cysteine after the natural C-terminal Threonine).

A further variant of CCL2-P8A can be obtained by fusing this sequence to an immunoglobulin domain constant region, a protein domain known to improve the stability and the efficacy of recombinant proteins in the circulation. The resulting fusion protein can be expressed directly by mammalian cells (such as CHO or HEK293 cells) using the appropriate expression vectors so that the fusion protein is secreted in the culture medium. In a preferred arrangement, the nucleic acid sequence encoding the mature CCL2-P8A can be cloned in an expression vector fused to a nucleic acid sequence encoding the human CCL2 signal sequence at its 5' end, and the nucleic acid sequence encoding the constant region (segment 246-467) of human immunoglobulin lambda heavy chain IgG1 (NCBI Acc. No. CAA75302) at its 3' end. The resulting vector can be used to transform a CHO or HEK293 cell line and the

clones stably expressing and secreting the recombinant fusion protein having CCL2 -
P8A at the N-terminus and the IgG1 sequence at the C-terminus (figure 7B) can be
selected. This clone then can be used for scaling up the production and for purifying
the recombinant fusion protein from the culture medium. Alternatively, the position of
5 the nucleic acid encoding the constant region (segment 243-474) of human
immunoglobulin lambda heavy chain IgG1 and CCL2-P8A can be inverted, and the
resulting protein can be expressed and secreted using still the human CCL2 signal
sequence, or any other signal sequence.

TABLE I

Amino Acid	Synonymous Groups	More Preferred Synonymous Groups
Ser	Gly, Ala, Ser, Thr	Thr, Ser
Arg	Asn, Lys, Gln, Arg, His	Arg, Lys, His
Leu	Phe, Ile, Val, Leu, Met	Ile, Val, Leu, Met
Pro	Pro, Ala, Ser, Thr	Pro
Thr	Gly, Ala, Ser, Thr	Thr, Ser
Ala	Gly, Thr, Ser	Gly, Ala
Val	Met, Phe, Ile, Leu, Val	Met, Ile, Val, Leu
Gly	Ala, Thr, Ser, Gly	Gly, Ala
Ile	Phe, Ile, Val, Leu, Met	Ile, Val, Leu, Met
Phe	Trp, Phe, Tyr	Tyr, Phe
Tyr	Trp, Phe, Tyr	Phe, Tyr
Cys	Ser, Thr, Cys	Cys
His	Asn, Lys, Gln, Arg, His	Arg, Lys, His
Gln	Glu, Asn, Asp, Gln	Asn, Gln
Asn	Glu, Asn, Asp, Gln	Asn, Gln
Lys	Asn, Lys, Gln, Arg, His	Arg, Lys, His
Asp	Glu, Asn, Asp, Gln	Asp, Glu
Glu	Glu, Asn, Asp, Gln	Asp, Glu
Met	Phe, Ile, Val, Leu, Met	Ile, Val, Leu, Met
Trp	Trp, Phe, Tyr	Trp

TABLE II

Amino Acid	Synonymous Group
Ser	D-Ser, Thr, D-Thr, allo-Thr, Met, D-Met, Met(O), D-Met(O), L-Cys, D-Cys
Arg	D-Arg, Lys, D-Lys, homo-Arg, D-homo-Arg, Met, Ile, D-.Met, D-Ile, Orn, D-Orn
Leu	D-Leu, Val, D-Val, AdaA, AdaG, Leu, D-Leu, Met, D-Met
Pro	D-Pro, L-l-thioazolidine-4-carboxylic acid, D-or L-1-oxazolidine-4-carboxylic acid
Thr	D-Thr, Ser, D-Ser, allo-Thr, Met, D-Met, Met(O), D-Met(O), Val, D-Val
Ala	D-Ala, Gly, Aib, B-Ala, Acp, L-Cys, D-Cys
Val	D-Val, Leu, D-Leu, Ile, D-Ile, Met, D-Met, AdaA, AdaG
Gly	Ala, D-Ala, Pro, D-Pro, Aib, .beta.-Ala, Acp
Ile	D-Ile, Val, D-Val, AdaA, AdaG, Leu, D-Leu, Met, D-Met
Phe	D-Phe, Tyr, D-Thr, L-Dopa, His, D-His, Trp, D-Trp, Trans-3,4, or 5-phenylproline, AdaA, AdaG, cis-3,4, or 5-phenylproline, Bpa, D-Bpa
Tyr	D-Tyr, Phe, D-Phe, L-Dopa, His, D-His
Cys	D-Cys, S--Me--Cys, Met, D-Met, Thr, D-Thr
Gln	D-Gln, Asn, D-Asn, Glu, D-Glu, Asp, D-Asp
Asn	D-Asn, Asp, D-Asp, Glu, D-Glu, Gln, D-Gln
Lys	D-Lys, Arg, D-Arg, homo-Arg, D-homo-Arg, Met, D-Met, Ile, D-Ile, Orn, D-Orn
Asp	D-Asp, D-Asn, Asn, Glu, D-Glu, Gln, D-Gln
Glu	D-Glu, D-Asp, Asp, Asn, D-Asn, Gln, D-Gln
Met	D-Met, S--Me--Cys, Ile, D-Ile, Leu, D-Leu, Val, D-Val

REFERENCES

- Alexander JM et al., *Cell*, 111; 343-356, 2002.
- Baggiolini M et al., *Annu Rev Immunol*, 15: 675-705, 1997.
- Baggiolini M, *J Intern Med*, 250: 91-104, 2001.
- 5 Blyth DI et al., *Am J Respir Cell Mol Biol.*, 14: 425-438, 1996.
- Brown A et al., *J Pept Sci* 2 :40-46, 1996.
- Cleland JL et al., *Curr Opin Biotechnol*,12: 212-219, 2001.
- Coleman RA et al., *Drug Disc Toay*, 6:1116-1126, 2001.
- Dougherty DA, *Curr Opin Chem Bio*, 4: 645-652, 2000.
- 10 Fernandez EJ and Lolis E, *Annu Rev Pharmacol Toxicol*, 42:469-499, 2002.
- Garrigue JL et al., *Contact Dermatitis*, 30: 231-237, 1994.
- Godessart N and Kunkel SL, *Curr Opin Immunol*, 13: 670-675, 2001.
- Golebiowski A et al., *Curr Opin Drug Discov Devel*, 4: 428-434, 2001.
- Gong J and Clark-Lewis I, *J Exp Med* 181: 631-640, 1995.
- 15 Gosling J et al., *J Clin Invest*, 103: 773-778, 1999.
- Greenwald RB et al., *Adv Drug Deliv Rev*, 55: 217-50, 2003.
- Gu L et al., *Chem Immunol*, 72: 7-29, 1999.
- Gu L et al., *Nature*, 404: 407-411, 2000.
- Harris JM and Chess RB, *Nat Rev Drug Discov*, 2: 214-21, 2003.
- 20 Handel T et al., *Biochemistry*,35 : 6569-6584, 1996.
- Hemmerich S et al., *Biochemistry*, 38: 13013-13025, 1999.
- Hruby VJ and Balse PM, *Curr Med Chem*, 7:945-970, 2000.
- Izikson L et al., *Clin Immunol*, 103: 125-31 2002.
- Kim KS et al.,, *FEBS Lett*, 395:277-282, 1996.
- 25 Li AP, *Drug Disc Today*, 6: 357-366, 2001

- Loetscher P and Clark-Lewis I, *J Leukoc Biol*, 69: 881-884, 2001.
- Lu B et al., *J Exp Med*, 187: 601-608, 1998.
- Lubkowski J et al., *Nat Struct Biol*, 4: 64-69, 1997.
- Luo B and Prestwich GD, *Exp Opin Ther Patents*, 11: 1395-1410, 2001.
- 5 Mishell B, *Immunopharm*, 2: 233-245. 1980.
- Mahad DJ and Ransohoff RM, *Semin Immunol*, 15: 23-32, 2003.
- Mitsui G et al., *Immunol Lett*, 86: 191-7, 2003.
- Mizumoto N et al., *Immunobiology*, 204:477-93, 2001.
- Murphy LR et al., *Protein Eng*, 13:149-152, 2000.
- 10 Nilsson J et al., *Protein Expr Purif*, 11: 1-16, 1997.
- Paavola C et al., *J Biol Chem*, 273: 33157-33165, 1998.
- Pillai O and Panchagnula R, *Curr Opin Chem Biol*, 5: 447-451, 2001.
- Proudfoot A et al., *Proc Natl Acad Sci U S A*, 100: 1885-1890, 2003.
- Rogov SI and Nekrasov AN, *Protein Eng*, 14: 459-463, 2001.
- 15 Sahrbacher UC et al., *Eur J Immunol*, 28: 1332-8, 1998.
- Schwarz MK and Wells TN, *Curr Opin Chem Biol*, 3: 407-417, 1999.
- Seet BT et al., *Proc Natl Acad Sci U S A*, 98: 9008-9013, 2001.
- Steitz SA et al., *FEBS Lett*, 430: 158-164, 1998.
- Villain M et al., *Chem Biol*, 8: 673-679, 2001.
- 20 Yoshimura T et al., *FEBS Lett*, 244: 487-493, 1989.
- Xu H et al., *J Exp Med*, 183: 1001-12, 1996.
- Zhang Y et al., *Methods*, 10: 93-103, 1996.

CLAIMS

1. The use of a polypeptide comprising SEQ ID NO: 2 as a medicament.
- 5 2. The use according to claim 1 wherein the polypeptide further comprises an isoleucine at position 64 of SEQ ID NO: 2.
3. The use according to claim 1 or 2 further characterized in that said polypeptide does not contain a mutation in position 9, 10, or 13 in the corresponding
10 sequence of SEQ ID NO: 2 and SEQ ID NO: 4.
4. The use according to claim 1 or 2 further characterized in that said polypeptide contains, in the corresponding sequence of SEQ ID NO: 2 and SEQ ID NO: 4:
 - a) a Cysteine in position 8, 14, 17, or 77; or
 - 15 b) an Alanine or a Glycine in position 1.
5. The use of any of the claims from 1 to 4, wherein said polypeptide comprises the constant region of a human immunoglobulin heavy chain.
- 20 6. The use of a polypeptide comprising SEQ ID NO: 2 for the manufacture of a medicament for the treatment of autoimmune, inflammatory or infectious diseases.
7. The use according to claim 65 wherein the polypeptide further comprises an
25 isoleucine at position 64 of SEQ ID NO: 2.

8. The use according to claim 6 or 76 further characterized in that said polypeptide does not contain a mutation in position 9, 10, or 13 in the corresponding sequence of SEQ ID NO: 2 and SEQ ID NO: 4.

5

9. The use according to claim 6 or 7 further characterized in that said polypeptide contains, in the corresponding sequence of SEQ ID NO: 2 and SEQ ID NO: 4:

a) a Cysteine in position 8, 14, 17, or 77; or

b) an Alanine or a Glycine in position 1.

10

10. The use of any of the claims from 6 to 9, wherein said polypeptide comprises the constant region of a human immunoglobulin heavy chain.

11. The use according to claim 6 wherein said disease is selected from the group consisting of: arthritis, rheumatoid arthritis (RA), psoriatic arthritis, osteoarthritis, systemic lupus erythematosus (SLE), systemic sclerosis, scleroderma, polymyositis, glomerulonephritis, fibrosis, fibrosis, allergic or hypersensitivity diseases, dermatitis, asthma, chronic obstructive pulmonary disease (COPD), inflammatory bowel disease (IBD), Crohn's diseases, ulcerative colitis, multiple sclerosis, cancer, septic shock, viral or HIV infections, transplantation, , airways inflammation, graft-versus-host disease (GVHD) and atherosclerosis.

20

12. The use according to claim 7 wherein said disease is selected from the group consisting of: arthritis, rheumatoid arthritis (RA), psoriatic arthritis, osteoarthritis, systemic lupus erythematosus (SLE), systemic sclerosis, scleroderma,

25

polymyositis, glomerulonephritis, fibrosis, fibrosis, allergic or hypersensitivity diseases, dermatitis, asthma, chronic obstructive pulmonary disease (COPD), inflammatory bowel disease (IBD), Crohn's diseases, ulcerative colitis, multiple sclerosis, cancer, septic shock, viral or HIV infections, transplantation, , airways inflammation, graft-versus-host disease (GVHD) and atherosclerosis.

5

13. The use according to claim 11 wherein the disease is multiple sclerosis.

14. The use according to claim 12 wherein the disease is multiple sclerosis.

10

15. The fusion polypeptide amino acid sequence of SEQ ID NO: 2 fused to the constant region of a human immunoglobulin heavy chain of SEQ ID NO: 5.

16. The nucleic acid sequence encoding for the fusion polypeptide of SEQ ID NO:

15

5.

17. Method for producing the fusion polypeptide of claim 15 comprising:

a) cloning of the nucleic acid sequence encoding the mature CCL2-P8A in an expression vector fused to a nucleic acid sequence encoding the human CCL2 signal sequence at its 5' end, and the nucleic acid sequence encoding the constant region (segment 243-474) of human immunoglobulin lambda heavy chain IgG1 at its 3' end;

20

b) transforming a CHO or HEK293 cell line with the resulting vector;

- c) selecting the clones stably expressing and secreting the recombinant fusion protein having CCL2-P8A at the N-terminus and the IgG1 sequence at the C-terminus;
- d) purifying the fusion protein from the culture medium.

5

18. Methods for screening for obligate monomeric antagonist chemokine variants described herein comprising:

- a) making single point mutations in CCL2 that block its ability to dimerize;
- b) identifying said variants that bind to the receptor and show agonistic properties in vitro;
- c) identifying said variants from the group identified in (b) above that are further characterized as inhibiting peritoneal cell recruitment.

10

19. The pharmaceutical composition comprising a monomeric variant of a homodimer-forming chemokine as active ingredient, wherein said variant result from at least an amino acid substitution that alters the pattern of hydrogen bonds at the dimerization interface of said chemokine.

15

20. The pharmaceutical composition of claim 19 wherein the monomeric variant is chosen from:

20

- a) CCL2-P8A (SEQ ID NO: 2);
- b) CCL2*-P8A (SEQ ID NO: 4);
- c) An active mutant of (a) or (b); or
- d) A polypeptide comprising (a), (b), or (c), and an amino acid sequence belonging to a protein sequence other than said chemokine.

25

21. The pharmaceutical composition of claim 18 or 19, wherein said monomeric variants is in the form of an active fraction, precursors, salt, derivative, complex or conjugate.
- 5
22. A method for treating or preventing autoimmune, inflammatory, or infectious diseases comprising the administration of an effective amount of a monomeric variants of a homodimer-forming chemokine, wherein said variant result from at least an amino acid substitution that alters the pattern of hydrogen bonds at the
- 10 dimerization interface of said chemokine.
23. The method of claim 21 wherein the monomeric variant is chosen from:
- a) CCL2-P8A (SEQ ID NO: 2);
 - b) CCL2*-P8A (SEQ ID NO: 4);
 - 15 c) An active mutant of (a) or (b); or
 - d) A polypeptide comprising (a), (b), or (c), and an amino acid sequence belonging to a protein sequence other than said chemokine.

1/7

Figure 1

CCL2	1	QPDAINAPVT	CCYNFTNRKI	SVQRLASYRR	ITSSKCPKEA	VIFKTIVAKE	50
CCL2-P8A	1	QPDAINA <u>A</u> VT	CCYNFTNRKI	SVQRLASYRR	ITSSKCPKEA	VIFKTIVAKE	50
CCL2*	1	QPDAINAPVT	CCYNFTNRKI	SVQRLASYRR	ITSSKCPKEA	VIFKTIVAKE	50
CCL2*-P8A	1	QPDAINA <u>A</u> VT	CCYNFTNRKI	SVQRLASYRR	ITSSKCPKEA	VIFKTIVAKE	50
CCL2	51	ICADPKQKWV	QDSMDHLDKQ	TQTPKT	76		
CCL2-P8A	51	ICADPKQKWV	QDSMDHLDKQ	TQTPKT	76		
CCL2*	51	ICADPKQKWV	QDSI DHLDKQ	TQTPKT	76		
CCL2*-P8A	51	ICADPKQKWV	QDSI DHLDKQ	TQTPKT	76		

2/7

Figure 2






Figure 3

A)



B)



4/7

Figure 4






Figure 5



B)



6/7

Figure 6





Figure 7

(A)



(B)

1 MKVSAAALCL LLIAATFIPO GLAQPDAINA AVTCCYNFTN RKISVQRLAS 50
 51 YRRITSSKCP KEAVIFKTIV AKEICADPKQ KVVQDSMDHL DKQTQTPKT E 100
 101 PKSCDKTHTC PPCAPELGG GPSVFLFPPK PKDTLMISRT PEVTCVVVDV 150
 151 SHEDPEVKFN WYVDGVEVHN AKTKPREEQY NSTYRVVSVL TVLHQDWLNG 200
 201 KEYKCKVSNK ALPAPIEKTI SKAKGQPREP QVYTLPPSRE EMTKNQVSLT 250
 251 CLVKGFYPSD IAVEWESNGQ PENNYKTTPP VLDSGSEFFL YSKLTVDKSR 300
 301 WQQGNVFSCS VMHEALHNHY TQKSLSLSPG K

SEQUENCE LISTING

<110> Applied Research Systems ARS holding

5 <120> THERAPEUTIC USES OF CHEMOKINE VARIANTS

<130> WO933

<160> 5

10 <170> PatentIn version 3.0

<210> 1

15 <211> 76

<212> PRT

<213> Homo sapiens

<220>

<223> Human CCL2

20 <400> 1

Gln Pro Asp Ala Ile Asn Ala Pro Val Thr Cys Cys Tyr Asn Phe Thr
 1 5 10 15

25 Asn Arg Lys Ile Ser Val Gln Arg Leu Ala Ser Tyr Arg Arg Ile Thr
 20 25 30

30 Ser Ser Lys Cys Pro Lys Glu Ala Val Ile Phe Lys Thr Ile Val Ala
 35 40 45

Lys Glu Ile Cys Ala Asp Pro Lys Gln Lys Trp Val Gln Asp Ser Met
 50 55 60

35 Asp His Leu Asp Lys Gln Thr Gln Thr Pro Lys Thr
 65 70 75

<210> 2

40 <211> 76

<212> PRT

<213> synthetic construct

<220>

<223> Human CCL2-P8A

45 <400> 2

Gln Pro Asp Ala Ile Asn Ala Ala Val Thr Cys Cys Tyr Asn Phe Thr
 1 5 10 15

50 Asn Arg Lys Ile Ser Val Gln Arg Leu Ala Ser Tyr Arg Arg Ile Thr
 20 25 30

55 Ser Ser Lys Cys Pro Lys Glu Ala Val Ile Phe Lys Thr Ile Val Ala
 35 40 45

Lys Glu Ile Cys Ala Asp Pro Lys Gln Lys Trp Val Gln Asp Ser Met
 50 55 60

<213> synthetic construct

<220>

<223> Human CCL2-P8A_IgG1 fusion protein

5 <400> 5

Met Lys Val Ser Ala Ala Leu Leu Cys Leu Leu Leu Ile Ala Ala Thr
1 5 10 15

10 Phe Ile Pro Gln Gly Leu Ala Gln Pro Asp Ala Ile Asn Ala Ala Val
20 25 30

Thr Cys Cys Tyr Asn Phe Thr Asn Arg Lys Ile Ser Val Gln Arg Leu
35 40 45

15 Ala Ser Tyr Arg Arg Ile Thr Ser Ser Lys Cys Pro Lys Glu Ala Val
50 55 60

20 Ile Phe Lys Thr Ile Val Ala Lys Glu Ile Cys Ala Asp Pro Lys Gln
65 70 75 80

Lys Trp Val Gln Asp Ser Met Asp His Leu Asp Lys Gln Thr Gln Thr
85 90 95

25 Pro Lys Thr Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro
100 105 110

Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro
115 120 125

30 Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr
130 135 140

35 Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn
145 150 155 160

Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg
165 170 175

40 Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val
180 185 190

Leu His Asn Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser
195 200 205

45 Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys
210 215 220

50 Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu
225 230 235 240

Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe
245 250 255

55 Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Gln Gly Gln Pro Glu
260 265 270

Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe

		275					280									285
	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly
5		290					295					300				
	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr
	305					310					315					320
10	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys					
					325						330					

INTERNATIONAL SEARCH REPORT

PCT/EP2004/052572

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K38/19 A61P29/00 A61P31/00 A61P37/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>PAAVOLA CHAD D ET AL: "Monomeric monocyte chemoattractant protein-1 (MCP-1) binds and activates the MCP-1 receptor CCR2B" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 50, 11 December 1998 (1998-12-11), pages 33157-33165, XP002283331 ISSN: 0021-9258 cited in the application</p>	2, 3, 19-22
Y	<p>the whole document</p> <p style="text-align: center;">----- -/--</p>	4, 5, 15-17



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

25 February 2005

Date of mailing of the international search report

10/03/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Bayrak, S

INTERNATIONAL SEARCH REPORT

PCT/EP2004/052572

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HEMMERICH S ET AL: "Identification of residues in the monocyte chemotactic protein-1 that contact the MCP-1 receptor, CCR2." BIOCHEMISTRY. 5 OCT 1999, vol. 38, no. 40, 5 October 1999 (1999-10-05), pages 13013-13025, XP002319264 ISSN: 0006-2960	1,3, 19-22
Y		4,5, 15-17
X	US 5 739 103 A (ZHANG YUJUN ET AL) 14 April 1998 (1998-04-14) column 4, line 43 - line 50 column 6, line 55 - line 59 column 10, line 46 - line 50	19-22
X	WO 03/083059 A (CENTOCOR, INC) 9 October 2003 (2003-10-09) the whole document	19-22
P,X	WO 03/084993 A (KOSCO-VILBOIS MARIE ; PROUDFOOT AMANDA (FR); APPLIED RESEARCH SYSTEMS) 16 October 2003 (2003-10-16) page 5 page 8; examples	19-22
A	KIM KEY-SUN ET AL: "Structural characterization of a monomeric chemokine: Monocyte chemoattractant protein-3" FEBS LETTERS, vol. 395, no. 2-3, 1996, pages 277-282, XP002283332 ISSN: 0014-5793 page 281, column 2, line 9 - line 11	1-23

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 22,23 are directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.
2. Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

PCT/EP2004/052572

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5739103	A	14-04-1998	US 5459128 A	17-10-1995
			EP 0828833 A1	18-03-1998
			JP 11506005 T	02-06-1999
			WO 9638559 A1	05-12-1996
			US 5705360 A	06-01-1998
			US 5854412 A	29-12-1998
			AT 204583 T	15-09-2001
			CA 2176217 A1	18-05-1995
			DE 69428049 D1	27-09-2001
			DE 69428049 T2	18-04-2002
			DK 725794 T3	10-12-2001
			EP 0725794 A1	14-08-1996
			ES 2162908 T3	16-01-2002
			JP 9505053 T	20-05-1997
			PT 725794 T	28-02-2002
WO 9513295 A1	18-05-1995			
WO 03083059	A	09-10-2003	AU 2003224761 A1	13-10-2003
			EP 1497423 A2	19-01-2005
			WO 03083059 A2	09-10-2003
WO 03084993	A	16-10-2003	AU 2003240765 A1	20-10-2003
			WO 03084993 A1	16-10-2003
			EP 1495050 A1	12-01-2005