



US00PP14379P29

(12) **United States Plant Patent**
Noodeljik

(10) **Patent No.:** **US PP14,379 P2**
(45) **Date of Patent:** **Dec. 16, 2003**

(54) **CHRYSANTHEMUM PLANT NAMED**
'MITHRA'

(50) Latin Name: *Chrysanthemum morifolium*
Varietal Denomination: **Mithra**

(75) Inventor: **Robert Noodeljik**, Woubrugge (NL)

(73) Assignee: **Chrysanthemum Breeders Association**
N.V. (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(22) Filed: **Dec. 12, 2002**

(51) **Int. Cl.**⁷ **A01H 5/00**

(52) **U.S. Cl.** **Plt./293**

(58) **Field of Search** **Plt./293**

Primary Examiner—Anne Marie Grunberg

(74) *Attorney, Agent, or Firm*—Parkhurst & Wendel, L.L.P.

(57) **ABSTRACT**

A chrysanthemum plant named 'Mithra' characterized by its small sized blooms with dark red ray florets and prolific branching; natural season flower date August 26–31; blooming for a period of 5 weeks.

(21) Appl. No.: **10/317,043**

3 Drawing Sheets

1

BACKGROUND OF THE INVENTION

'Mithra' is a product of a breeding and selection program for outdoor pot mums (garden mums) which had the objective of creating new chrysanthemum cultivars with a decorative (pompom) type flower, a natural season flower date around August 26–31; blooming for a period of 5 weeks. The new plant of the present invention comprises a new and distinct cultivar of Chrysanthemum plant 'Mithra' is a seedling resulting from the open pollination among groups of chrysanthemum cultivars maintained under the control of the inventor for breeding purposes. The new and distinct cultivar was discovered and selected as one flowering plant by Rob Noodeljik on a cultivated field in Rijsenhout Holland in August 2000. The plant has been asexually reproduced by cuttings in greenhouses at Rijsenhout Holland. The new cultivar has been found to retain its distinctive characteristics through successive propagations.

BRIEF DESCRIPTION OF THE DRAWINGS

The present invention of a new and distinct variety of chrysanthemum is shown in the accompanying drawings, the color being as nearly true as possible with color photographs of this type.

FIG. 1 shows a plant of the cultivar in full bloom.

FIG. 2 shows the various stages of bloom of the new cultivar.

FIG. 3 shows the petiole and foliage of the new cultivar.

DESCRIPTION OF THE INVENTION

This new variety of chrysanthemum is of the botanical classification *Chrysanthemum morifolium*. The observations and measurements were gathered from plants grown out door in Rijsenhout, Holland under natural day length and temperature and planted week 22 in 2000 and 2001. The natural blooming date of this crop was August 26–31 (week 35). The average height of the plants was 26 cms. No growth retardants were used. No tests were done on disease or insect resistance or susceptibility. No tests were done on cold or drought tolerance. This new variety produces small sized blooms with dark ray florets blooming for a period of 5 weeks.

2

From the cultivars known to inventor the most similar existing cultivar in comparison to 'Mithra' is 'Glenda'. When 'Glenda' and 'Mithra' are being compared the following differences are noticed: The differences of 'Glenda' and 'Mithra' are (1) Flower color. 'Mithra' is more dark red than 'Glenda'. (2) Natural flowering date. 'Mithra' flowers earlier than 'Glenda'.

The following is a description of the plant and characteristics that distinguish 'Mithra' as a new and distinct variety. The color designations are taken from the plant itself. Accordingly, any discrepancies between the color designations and the colors depicted in the photographs are due to photographic tolerances. The color chart used in this description is: The Royal Horticultural Society Colour Chart, edition 1995.

Table 1. Botanical Description of cultivar 'Mithra'

Bud:

Size.—Medium; cross-section 1.3 cm, height 0.7 cm.

Outside color.—Red 53 C.

Involucral bracts.—2 rows, length 7 mm, width 3 mm.

Involucral bracts among disc-florets.—Not present.

Involucral bracts color.—Green 138 C.

Bloom:

Type.—Decorative (pompom).

Height.—Medium. 1.5 cm.

Size.—Small.

Fully expanded.—3.5 cm.

Number of blooms per branch.—Approx. 7–9 blooms per branch.

Performance on the plant.—5 weeks.

Seeds.—Produced in small quantities, ovate grey-brown 199A, 1½ mm in length.

Fragrance.—Typical chrysanthemum, slightly.

Color:

Center of the flower.—Immature red 46 A. Mature red 46 A.

Color of upper surface of the ray-florets.—Red 46 A.

Color of the lower surface of the ray-florets.—Red 53 C.

Tonality from distance.—A garden mum with dark red flowers.

Color of the upper surface of the plant after aging of the plant.—Red 46 A.

Ray florets:

Texture.—Upper and under side smooth.
Number.—130–140.
Cross-section.—Concave.
Longitudinal axis of majority.—Incurving.
Length of corolla tube.—Short 0.2 cm.
Ray-floret margin.—Entire.
Ray-floret length.—0.9–1.0 cm.
Ray-floret width.—0.4 cm.
Ratio length / width.—Medium.
Shape of tip.—Dentate.
Disc florets.—None.
Receptacle shape.—Domed raised.

Reproductive organs:

Stamen.—Absent.
Pollen.—Absent.
Styles.—Thin.
Style color.—Yellow-green 144 A.
Style length.—4 mm.
Stigma color.—Yellow-green 144 A.
Stigma width.—1 mm.
Ovaries.—Enclosed in calyx.

Plant:

Shape.—Grown as a spray-type pot mum, outdoor mounded and round.
Growth habit.—Spreading.
Growth rate.—Moderate.
Height.—26 cm.
Width.—31–33 cm.
Stem color.—Yellow-green 147 B.
Stem strength.—Strong.
Stem brittleness.—Present.
Stem anthocyanin coloration.—Absent.
Length of lateral branch.—From top to bottom 12–13 cm.
Lateral branch color.—Yellow-green 147 B.
Lateral branch, attachment.—Brittle.

Branching (average number of lateral branches).—

Good with 7–8 breaks after pinching.

Peduncle length.—4.0–4.5 cm.

Peduncle color.—Yellow-green 147 B.

Natural season blooming date.—August 26–31.

Foliage:

Color.—Upper side yellow-green 147 A. Under side yellow-green 147 B.

Size.—Small; length 5.0 cm, width 3.0 cm.

Quantity (number per lateral branch).—10.

Shape.—Ovate.

Texture upper side.—Glabrous.

Texture under side.—Pubescent.

Venation arrangement.—Palmate.

Shape of the margin.—Serrated.

Shape of base of sinus between lateral lobes.—Acute.

Margin of sinus between lateral lobes.—Diverging.

Shape of base.—Acute-Rounded.

Apex.—Mucronate.

TABLE 2

Differences with the comparison varieties		
	'MITHRA'	'GLENDA'
Color of the upper surface of the ray-florets	Red 46 A	Red 45 A
Natural blooming date	August 26–31	October 1–8

I claim:

1. A new and distinct variety of chrysanthemum plant as described and illustrated.

* * * * *




FIG. 1




FIG. 2



FIG. 3