



(86) Date de dépôt PCT/PCT Filing Date: 2012/04/26
(87) Date publication PCT/PCT Publication Date: 2012/11/01
(85) Entrée phase nationale/National Entry: 2014/01/09
(86) N° demande PCT/PCT Application No.: US 2012/035251
(87) N° publication PCT/PCT Publication No.: 2012/149186
(30) Priorité/Priority: 2011/04/29 (US61/480,641)

(51) Cl.Int./Int.Cl. *C07D 401/12* (2006.01),
A61K 31/4725 (2006.01), *A61P 29/00* (2006.01),
C07D 401/14 (2006.01)

(71) Demandeur/Applicant:
THE TRUSTEES OF THE UNIVERSITY OF
PENNSYLVANIA, US

(72) Inventeurs/Inventors:
AMARAVADI, RAVI K., US;
WINKLER, JEFFREY, US

(74) Agent: DIMOCK STRATTON LLP

(54) Titre : NOUVEAUX COMPOSES BISAMINOQUINOLEINES, COMPOSITIONS PHARMACEUTIQUES PREPAREES
A PARTIR DE CEUX-CI ET LEUR UTILISATION

(54) Title: NOVEL BISAMINOQUINOLINE COMPOUNDS, PHARMACEUTICAL COMPOSITIONS PREPARED
THEREFROM AND THEIR USE

(57) **Abrégé/Abstract:**

The present invention relates to novel bisaminoquinoline compounds, pharmaceutical compositions comprising these novel compounds and methods for inhibiting autophagy in biological systems. Methods of treating cancer in patients in need using compounds and/or compositions according to the present invention alone or in combination with at least one additional anticancer agent represent additional aspects of the invention. Methods of treating disease states and/or conditions in which inhibition of autophagy plays a favorable treatment role including rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria and Sjogren's disease, with compounds according to the present invention represent additional aspects of the invention.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property
Organization
International Bureau



WIPO | PCT



(10) International Publication Number
WO 2012/149186 A9

(43) International Publication Date
1 November 2012 (01.11.2012)

(51) International Patent Classification:

C07D 401/12 (2006.01) *A61K 31/4725* (2006.01)
C07D 401/14 (2006.01) *A61P 29/00* (2006.01)

(21) International Application Number:

PCT/US2012/035251

(22) International Filing Date:

26 April 2012 (26.04.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/480,641 29 April 2011 (29.04.2011) US

(71) Applicant (*for all designated States except US*): **THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA** [US/US]; 3160 Chestnut Street, Suite 200, Philadelphia, PA 19104 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **AMARAVADI, Ravi, K.** [US/US]; 706 Iris Lane, Media, PA 19063 (US). **WINKLER, Jeffrey** [US/US]; 541 Ballytore Road, Wybbwood, PA 19096 (US).

(74) Agent: **COLEMAN, Henry, D.**; Coleman Sudol Sapone, P.C., 714 Colorado Avenue, Bridgeport, CT 06605-1601 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,

DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with information concerning authorization of rectification of an obvious mistake under Rule 91.3 (b) (Rule 48.2(i))

(88) Date of publication of the international search report:

3 January 2013

(48) Date of publication of this corrected version:

21 February 2013

(15) Information about Correction:

see Notice of 21 February 2013

(54) Title: NOVEL BISAMINOQUINOLINE COMPOUNDS, PHARMACEUTICAL COMPOSITIONS PREPARED THEREFROM AND THEIR USE

(57) Abstract: The present invention relates to novel bisaminoquinoline compounds, pharmaceutical compositions comprising these novel compounds and methods for inhibiting autophagy in biological systems. Methods of treating cancer in patients in need using compounds and/or compositions according to the present invention alone or in combination with at least one additional anticancer agent represent additional aspects of the invention. Methods of treating disease states and/or conditions in which inhibition of autophagy plays a favorable treatment role including rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria and Sjogren's disease, with compounds according to the present invention represent additional aspects of the invention.



WO 2012/149186 A9

Novel Bisaminoquinoline Compounds, Pharmaceutical Compositions Prepared Therefrom and Their Use

Related Applications and Grant Support

The present application claims the benefit of priority of United States provisional application no.US61/480,641, filed April 29, 2011, of identical title to the present application, the entire contents of which are incorporated by reference herein.

This invention was made with government support under the Abramson Cancer Center Pilot grant awarded by the National Cancer Institute. The government has certain rights in the invention.

Field of the Invention

The present invention relates to novel bisaminoquinoline compounds, pharmaceutical compositions comprising these novel compounds and methods for inhibiting autophagy in biological systems. Methods of treating cancer in patients in need using compounds and/or compositions according to the present invention alone or in combination with at least one additional anticancer agent represent additional aspects of the invention. The use of radiation therapy in combination with the present compounds, alone or in combination with an additional anticancer agent as otherwise disclosed herein, represents an additional aspect of the invention. Methods of treating disease states and/or conditions in which inhibition of autophagy plays a favorable treatment role including rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria and Sjogren's disease, with compounds according to the present invention represent additional aspects of the invention.

Background of the Invention

Autophagy consists of the sequestration of organelles and proteins in autophagic vesicles (AV) and degradation of this cargo through lysosomal fusion (1). Autophagy allows tumor cells to survive metabolic and therapeutic stresses (2-5). Multiple publications indicate therapy-induced autophagy is a key resistance mechanism to many anti-cancer agents.

Chloroquine (CQ) (Compound 1, Figure 1) derivatives block autophagy by inhibiting the lysosome (3, 6, 7). A randomized phase III trial of CQ versus placebo with carmustine and radiation in patients with glioma reported a trend towards a doubling in duration of survival in the patients treated with CQ (8). Based on these findings, clinical trials combining cancer therapies with hydroxychloroquine (HCQ; Figure 1 Compound 2), (which is safer than CQ to dose escalate) have been launched. Preliminary results indicate these combinations have activity (9), but it is still unclear if this activity is consistently due to the addition of HCQ. High micromolar concentrations of HCQ are required to inhibit autophagy. While there is some pharmacodynamic evidence of autophagy inhibition with HCQ in cancer patients, it is inconsistent because adequate concentrations are not achieved in all patients (10). There is an unmet need to develop more potent inhibitors of autophagy. The design and synthesis of dimeric analogs of CQ, that exploit the thermodynamic advantages imparted by polyvalency (11, 12), has been a subject of intensive study for over 10 years (13-15). An early report by Vennerstrom(14) described the synthesis of heteroalkane-bridged bisquinolines as potential antimalarials, but none of the compounds had sufficient antimalarial activity to warrant further investigation. Subsequently, Sergheraert (13) reported that tetraquinolines, i.e., dimers of bisquinolines, afforded potent antimalarials, confirming the possibility that the application of the polyvalency strategy could afford increased potency, at least with respect to antimalarial activity.

More recently, Lee(16) has described the potentiation of AKT inhibitors by fluorinated quinoline analogs. Solomon(17) has reported the preparation of "repositioned" chloroquine dimers, based on the use of a piperazine connector. These results suggest that these chloroquine analogs could serve as bases for the development of a new group of effective cancer chemotherapeutics.

Objects of the Invention

It is an object of the invention to provide novel compounds for inhibiting autophagy in biological systems, especially including patients or subjects in need.

It is another object of the invention to treat disease states and/or conditions in which inhibition of autophagy is beneficial to the disease states and/or conditions a patient or subject.

It is yet a further object of the invention to provide pharmaceutical compositions which may be used to inhibit autophagy, especially autophagy associated with disease states and/or conditions including cancer and its metastasis.


It is still a further object of the invention to inhibit, treat or prevent cancer, including the metastasis of cancer in patients or subjects in need utilizing compounds, compositions and/or methods which are presented herein.

It is still another object of the invention to inhibit, treat or prevent diseases in which the inhibition of autophagy provides a favorable effect, including rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria and Sjogren's disease, among others.

Any one or more of these and/or other objects of the invention may be readily gleaned from a description of the invention which follows.

Brief Description of the Invention

The present invention relates to compounds according to the chemical structure I



Wherein R^1 and $R^{1'}$ are each independently H, halo (F, Cl, Br or I), CN, NO₂, optionally substituted C₁-C₆ alkyl (when substituted, preferably substituted with 1 or 2 hydroxyl groups or 3-5 fluoro groups), optionally substituted O-C₁-C₆ alkyl (preferably, OCH₃), optionally substituted C₂-C₇ acyl (preferably acetyl) or optionally substituted C₂-C₇ ester (oxycarbonyl ester or carboxyester, preferably carboxyester);

R and R' are each independently H, a C₁-C₆ optionally substituted alkyl group, a C₁-C₇ (preferably C₂-C₇) optionally substituted acyl group, a C₂-C₇ optionally substituted carboxy ester group (which forms a urethane group with the nitrogen atom to which R or R' is bonded);

L is a $-(CH_2Y)_n-X-(Y'CH_2)_n-$ group or a $A-(CH_2CH_2-Z)_n-A'$ group (either A

or A' may be bonded to either of the two amine groups in compound I) wherein at least one of the CH₂ groups in L is optionally substituted with a C₁-C₃ alkyl group which itself is optionally substituted with one or two hydroxyl groups;

X is absent, (CH₂)_j, O, S or N-R'';

Y is absent, CH₂, O, CH₂O or N-R'' and Y' is absent, CH₂, O, OCH₂ or N-R'', with the proviso that when one or more of X, Y and Y' is present, each of X and Y, X and Y' or Y and Y', when present, forms a stable bond;

R'' is H or an optionally substituted C₁-C₆ (preferably C₁-C₃) alkyl group;

j is 1, 2 or 3 (preferably 1 or 2);

n is 0, 1, 2, 3 or 4, with the proviso that when n is 0, X is (CH₂)_j where j is at least 1 and at least one CH₂ group is optionally substituted with a C₁-C₃ alkyl group which itself is optionally substituted with one or two hydroxyl groups;

A is absent or (CH₂)_j and A' is (CH₂)_j wherein at least one CH₂ group in A or A' is optionally substituted with a C₁-C₃ alkyl group which is itself optionally substituted with one or two hydroxyl groups;

Z is O or N-R^Z;

R^Z is H or an optionally substituted C₁-C₃ alkyl group,

or a pharmaceutically acceptable salt, enantiomer, diastereomer, solvent or polymorph thereof.

In preferred aspects of the invention, R¹ and R^{1'} are each independently H, a halo group, a nitro group or a trifluoromethyl group, preferably a chloro group. R and R' are preferably each independently H, a C₁-C₃ optionally substituted alkyl group itself preferably substituted with at least one hydroxyl group, an alkoxy group, an amine, monoalkyl amine or dialkyl amine group, wherein said amine group or said monoalkyl amine group is optionally substituted on the amine position with one or two 7-substituted-4-quinolinyl group(s) wherein the amine binds to the 4-position of the quinolinyl group and the 7-position of each quinolinyl group is optionally substituted, preferably with a R¹ and/or R^{1'} group as broadly

described for generic structure I above, or one or both alkyl groups of said monoalkyl amine or dialkyl amine is itself further optionally substituted with at least one hydroxyl group, an alkoxy group, an amine, a monoalkyl amine or a dialkyl amine wherein the amine or monoalkyl amine is optionally substituted on the amine position with one or two 7-substituted-quinolinyl group(s) wherein the amine binds to the 4-position of the quinolinyl group and the 7-position of each quinolinyl group is optionally substituted, preferably with R^1 and/or $R^{1'}$ as broadly described for generic structure I above, and each of said alkoxy groups (e.g. methoxy or ethoxy) is optionally further substituted with an alkoxy group, preferably a methoxy group, thus forming a diether substituent.

In certain preferred aspects of the invention L is a $-(CH_2Y)_n-X-(Y'CH_2)_n-$ group, where X is N- R'' , Y and Y' are each independently absent or CH_2 , and R'' is H or a C_1 - C_3 alkyl group which is optionally substituted with at least one hydroxyl group, an alkoxy group, an amine, monoalkyl amine or dialkyl amine group, wherein said amine group or said monoalkyl amine group is optionally substituted on the amine position with one or two 7-substituted-4-quinolinyl group wherein the amine binds to the 4-position of the quinolinyl group and the 7-position of each quinolinyl group is optionally substituted, preferably with R^1 and/or $R^{1'}$ as broadly described for generic structure I above, or one or both alkyl groups of said monoalkyl amine or dialkyl amine is itself further optionally substituted with at least one hydroxyl group, an alkoxy group, an amine, a monoalkyl amine or a dialkyl amine wherein the amine or monoalkyl amine is optionally substituted on the amine position with one or two 7-substituted-quinolinyl group(s) wherein the amine binds to the 4-position of the quinolinyl group and the 7-position of each quinolinyl group is optionally substituted, preferably with R^1 and/or $R^{1'}$ as broadly described for generic structure I above, and each of said alkoxy groups (e.g. methoxy or ethoxy) is optionally further substituted with an alkoxy group, preferably a methoxy group, thus forming a diether substituent.

Further preferred compounds according to the present invention include those which are presented in the various schemes which are presented in Scheme 1 and Schemes 3-10 and figures 14, 15 and 15 A as presented herein.

In another aspect of the invention, a pharmaceutical composition comprises a compound according to formula I above or as otherwise described herein in combination with

a pharmaceutically acceptable carrier, additive or excipient, optionally in combination with at least one additional anticancer agent.

Methods of inhibiting autophagy in a biological system, in particular a patient or subject is a further aspects of this invention. In this aspect of the invention, a bisaminoquinoline compound as otherwise described herein is presented to the biological system, including administration to a patient or subject in need, in order to inhibit autophagy. The resulting inhibition may be monitored or applied in the biological system to effect a favorable result, including the inhibition, treatment and/or prevention of cancer, including metastasis of cancer, or the inhibition, treatment and/or prevention of one or more disease states or conditions in which the inhibition of autophagy provides a favorable result including rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria and Sjogren's disease, among others.

Methods of inhibiting, treating and/or reducing the likelihood of cancer, including metastasis of cancer and drug resistant cancer, comprises administering to a patient in need at least one compound according to the present invention, optionally in combination with at least one additional anticancer agent as otherwise described herein.

The present invention also relates to treating, inhibiting and/or preventing diseases, diseases states and/or conditions in a patient in need in which the inhibition of autophagy provides a favorable outcome, including rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria and Sjogren's disease, the method comprising administering to said patient at least one compound according to the present invention.

Brief Description of the Figures

Figure 1. Chemical structure of mono and bisaminoquinolines.

Figure 2. Synthesis scheme for bisaminoquinolines

Figure 3. Effects of Lys01-Lys04 on LC3 immunoblotting Immunoblotting and quantification of LC3II/LC3I ratio in lysates from LN229 cells treated for 4 hours. The

graphs show (mean $\pm$ SD) LC3II/LC3I ratios of each treatment normalized to the LC3II/LC3I ratio of control treated cells for each experiment.

Figure 4. Autophagy inhibition and cytotoxicity of Lys01 compared to HCQ. (A) Representative images of LN229 GFP-LC3 cells treated as indicated for 4 hours. White arrows: small puncta; red arrows: dense puncta. Graphs show mean $\pm$ SEM puncta/cell. (B) Representative electron micrographs of LN229-GFP-LC3 cells treated (4 hours) with DMSO, HCQ 10 μ M, or Lys01 10 μ M. Arrows: Autophagic vesicles. (C) LC3 immunoblotting of LN229 cells treated for 24 hours as indicated; calculated ratio of LC3II/LC3I ratios for bafilomycin versus control co-treatment. Above the dashed line indicates an autophagy inducer or control, below the dashed line indicates an autophagy inhibitor. (D) MTT assay (72 hours) for 4 cell lines. Red: Lys01, Blue: Lys02, Purple: Lys03, Green: Lys04, Orange: HCQ. Values presented are means $\pm$ SEM with 5 replicates per treatment.

Figure 5. Autophagy inhibition and cytotoxicity of Lys05, the water soluble salt of Lys01. (A) Immunoblotting against LC3 and p62 in c8161 cells treated as indicated. (B) MTT assay in c8161 cells at 72 hours. HCQ: Hydroxychloroquine. Values presented are mean $\pm$ SD with 5 replicates per treatment condition. * No remaining cells for analysis.

Figure 6. In vivo autophagy inhibition and antitumor activity of Lys05. (A) Representative electron micrographs (12,000x) of c8161 xenograft tumors harvested after 2 days of daily i.p. treatment with PBS, HCQ 60 mg/kg, or Lys05 76 mg/kg. Arrows: autophagic vesicles; scale bar 2 μ m. (B) Quantification of mean $\pm$ SEM number of autophagic vesicles/cell from two representative tumors from each treatment group. (C-D) 1205Lu xenografts were treated with PBS (blue), HCQ 60 mg/kg ip (green), or Lys05 76 mg/kg (red) i.p. every 3/5 days. (C) Tumor volumes over 14 days. (D) Daily tumor growth rate. (E-G) HT29 xenografts were generated in the flanks of nude mice and treated with PBS, Lys05 10 mg/kg ip daily, Lys05 40 mg/kg ip daily, or Lys05 80 mg/kg ip every 3/5 days. (E) Average daily tumor growth rate. (F) Tumor volumes over 14 days. (G) Weight of excised tumors, * $p < 0.05$.

Figure 7. Autophagy inhibition and tumor necrosis in melanoma and colon cancer xenografts treated with Lys05 or HCQ. (A) Immunoblotting against LC3 in lysates from individual c8161 tumors treated as indicated with daily i.p. injections for 48 hours. Quantification of LC3II/LC3I ratio (mean $\pm$ SEM). (B) Tumor necrosis (arrows) in H&E stained sections of

1205Lu tumor xenografts harvested after 14 days of treatment; Electron micrographs (7000-12000X) of melanoma tumor cells. Arrows: Autophagic vesicle (white); apoptotic cell (orange) (C) Immunoblotting against LC3 in HT29 xenografts treated with daily dosing (10, 40 mg/kg) or 3/5 days (80 mg/kg) for 14 days.

Figure 8. Toxicity associated with Lys05 76 mg.kg ip 3/5 days. (A) Mice were lethargic with arched backs after 3 days of dosing. (B) 3/10 mice developed bowel obstruction. (C) Dysmorphic paneth cells (arrows) in the terminal ileum of one mouse.

Figure 9. Lys05 treatment at the highest dose reproduces the intestinal phenotype of a genetic autophagy deficiency. (A-F) Weight and intestines were analyzed for mice bearing HT29 xenografts treated with PBS, Lys05 10-80 mg/kg. (A) Daily weight (B) Representative excised gastrointestinal tracts after 14 days of treatment (C) Representative images (40X) of hemotoxylin and eosin stained ileal crypts from mice bearing HT29 xenografts (14 days), (40X) arrows: paneth cells. (D) Paneth cell number per crypt (E) Paneth cell dysfunction score, * $p < 0.05$ (F) Scoring of lysozyme positive cells, * $p = 0.001$. Representative images of lysosome immunofluorescence (green) of ileum in mice treated with PBS and Lys05 80 mg/kg ip 3/5 days.

Figure 10. Paneth cell dysfunction scale. Under 40X power the size and number of eosinophilic granules per Paneth cell was scored for 10 Paneth cells per sample: A0= normal size and number A1: Decreased size, normal number; A2: Normal size, decreased number A3: decreased size and number.

Figure 11. Lys05 inhibits autophagy by accumulating in and deacidifying the lysosome. (A) 1205Lu cells (treated with PBS, HCQ 10 μ M, or Lys05 10 μ M for 24 hours) and harvested 1205 Lu xenograft tumors (treated with PBS, HCQ 60 mg/kg i.p. 3/5 days, or Lys05 76 mg/kg i.p. 3/5 days for 14 days) were homogenized and fractionated into whole cell (WC) and Lysosomal (L) fractions. LAMP2 immunoblotting confirmed isolation of concentrated lysosomes for analysis. (B) Concentrations of HCQ or Lys05 in cells and tumor whole cell and lysosomal homogenates. (C) Fluorescence imaging of 1205 Lu cells treated as indicated for 30 minutes and stained with LysoTracker Red. LysoTracker puncta (red) per cell was scored for three high powered fields. Blue: nuclear DAPI staining. Data presented is mean $\pm$

SEM. (D) Fluorescence imaging of c8161 cells treated as indicated for 24 hours and stained with acridine orange (AO): orange: aggregated AO, green: diffuse AO.

Figure 12. High performance liquid chromatography tandem mass spectrometry assay for HCQ and Lys05. 1205Lu cells (24 hours) and 1205Lu tumors (14 days). WC: Whole cell homogenate L: Lysosomal subfraction HCQ: hydroxychloroquine

Figure 13. Impairment of lysosomal enzymes and extralysosomal leakage associated with Lys05 treatment. (A) Acid phosphatase activity and (B) Cathepsin D immunoblotting in whole cell (white, WC) and lysosomal (Black; L) fractions of 1205 Lu cells treated with PBS, HCQ 10 μ M, Lys05 10 μ M for 24 hours. Graphs show the mean $\pm$ SEM for three independent experiments. (C) Acid phosphatase activity and (D) Cathepsin D immunoblotting in whole cell (white, WC) and lysosomal (Black; L) fractions of 1205 Lu xenografts treated with PBS, HCQ 60 mg/kg, Lys05 76 mg/kg i.p. 3/5 days (tumors). Whole cell homogenates (white) and lysosomal homogenates (black) were prepared from three separate tumors were pooled together. * $p < 0.05$.

Figure 14. Chemical structures of synthesized compounds Lys06-Lys12; The chemical structures of Lys06-Lys12 are shown.

Figure 15. Chemical structures of synthesized compounds Lys13-Lys18; The chemical structures of Lys13-Lys18 are shown.

Figure 15A shows a number of additional bisaminoquinoline autophagy inhibitors under investigation.

Figure 16, Table 1 provides MTT IC₅₀ values in LN229 for select compounds of the present invention as presented.

Figure 17, Table 12 provides IC₅₀ (M) values of a number of compounds according to the present invention in *P. falciparum*.

Detailed Description of the Invention

The following terms shall be used throughout the specification to describe the present invention. Where a term is not specifically defined herein, that term shall be understood to be used in a manner consistent with its use by those of ordinary skill in the art.

Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges is also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either both of those included limits are also included in the invention. In instances where a substituent is a possibility in one or more Markush groups, it is understood that only those substituents which form stable bonds are to be used.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described.

It must be noted that as used herein and in the appended claims, the singular forms "a," "and" and "the" include plural references unless the context clearly dictates otherwise.

Furthermore, the following terms shall have the definitions set out below.

The term "patient" or "subject" is used throughout the specification within context to describe an animal, generally a mammal, especially including a domesticated animal and preferably a human, to whom treatment, including prophylactic treatment (prophylaxis), with the compounds or compositions according to the present invention is provided. For treatment of those infections, conditions or disease states which are specific for a specific animal such

as a human patient, the term patient refers to that specific animal. In most instances, the patient or subject of the present invention is a human patient of either or both genders.

The term “effective” is used herein, unless otherwise indicated, to describe an amount of a compound or component which, when used within the context of its use, produces or effects an intended result, whether that result relates to the prophylaxis and/or therapy of an infection and/or disease state or as otherwise described herein. The term effective subsumes all other effective amount or effective concentration terms (including the term “therapeutically effective”) which are otherwise described or used in the present application.

The term “compound” is used herein to describe any specific compound or bioactive agent disclosed herein, including any and all stereoisomers (including diastereomers), individual optical isomers (enantiomers) or racemic mixtures, pharmaceutically acceptable salts and prodrug forms. The term compound herein refers to stable compounds. Within its use in context, the term compound may refer to a single compound or a mixture of compounds as otherwise described herein. It is understood that the choice of substituents or bonds within a Markush or other group of substituents or bonds is provided to form a stable compound from those choices within that Markush or other group.

The term “bioactive agent” refers to any biologically active compound or drug which may be formulated for use in the present invention. Exemplary bioactive agents include the compounds according to the present invention which are used to inhibit autophagy and to treat cancer as well as other compounds or agents which are otherwise described herein.

The terms “treat”, “treating”, and “treatment”, are used synonymously to refer to any action providing a benefit to a patient at risk for or afflicted with a disease, including improvement in the condition through lessening or suppression of at least one symptom, delay in progression of the disease, prevention or delay in the onset of the disease, etc.

Treatment, as used herein, encompasses both prophylactic and therapeutic treatment, principally of cancer. Compounds according to the present invention can, for example, be administered prophylactically to a mammal in advance of the occurrence of disease to reduce the likelihood of that disease. Prophylactic administration is effective to reduce or decrease

the likelihood of the subsequent occurrence of disease in the mammal, or decrease the severity of disease that subsequently occurs, especially including metastasis of cancer. Alternatively, compounds according to the present invention can, for example, be administered therapeutically to a mammal that is already afflicted by disease. In one embodiment of therapeutic administration, administration of the present compounds is effective to eliminate the disease and produce a remission or substantially eliminate the likelihood of metastasis of a cancer. Administration of the compounds according to the present invention is effective to decrease the severity of the disease or lengthen the lifespan of the mammal so afflicted, in the case of cancer.

The term “pharmaceutically acceptable” as used herein means that the compound or composition is suitable for administration to a subject to achieve the treatments described herein, without unduly deleterious side effects in light of the severity of the disease and necessity of the treatment.

The term “inhibit” as used herein refers to the partial or complete elimination of a potential effect, while inhibitors are compounds that have the ability to inhibit.

The term “prevention” when used in context shall mean “reducing the likelihood” or preventing a disease, condition or disease state from occurring as a consequence of administration or concurrent administration of one or more compounds or compositions according to the present invention, alone or in combination with another agent. It is noted that prophylaxis will rarely be 100% effective; consequently the terms prevention and reducing the likelihood are used to denote the fact that within a given population of patients or subjects, administration with compounds according to the present invention will reduce the likelihood or inhibit a particular condition or disease state (in particular, the worsening of a disease state such as the growth or metastasis of cancer) or other accepted indicators of disease progression from occurring.

The term “autophagy” or “autophagocytosis” is used to describe a catabolic process in cells which involves the degradation of a cell’s own components through lysosomes. Autophagy is a highly regulated process of biological systems that plays a normal part in cell growth development and homeostasis, helping to maintain a balance between the synthesis,

degradation, and subsequent recycling of cellular products. It is a major mechanism by which a cell allocates nutrients from unnecessary processes to more-essential processes.

A number of autophagic processes occur in nature, all of which have the degradation of intracellular components via the lysosome as a common feature. A well-known mechanism of autophagy involves the formation of a membrane around a targeted region of a cell, separating the contents from the rest of the cytoplasm. The resultant vesicle then fuses with a lysosome which subsequently degrades the contents.

Autophagy consists of the sequestration of organelles and proteins in autophagic vesicles (AV) and degradation of this cargo through lysosomal fusion (1). Autophagy allows tumor cells to survive metabolic and therapeutic stresses (2-5). Multiple publications indicate therapy-induced autophagy is a key resistance mechanism to many anti-cancer agents.

Diseases, disease states and/or conditions which benefit from the inhibition of autophagy include cancer (including the metastasis of cancer), rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria and Sjogren's disease.

The term "cancer" shall refer to a proliferation of tumor cells having the unique trait of loss of normal controls, resulting in unregulated growth, lack of differentiation, local tissue invasion, and/or metastasis. As used herein, neoplasms include, without limitation, morphological irregularities in cells in tissue of a subject or host, as well as pathologic proliferation of cells in tissue of a subject, as compared with normal proliferation in the same type of tissue. Additionally, neoplasms include benign tumors and malignant tumors (e.g., colon tumors) that are either invasive or noninvasive. Malignant neoplasms are distinguished from benign neoplasms in that the former show a greater degree of dysplasia, or loss of differentiation and orientation of cells, and have the properties of invasion and metastasis. The term cancer also within context, includes drug resistant cancers, including multiple drug resistant cancers. Examples of neoplasms or neoplasias from which the target cell of the present invention may be derived include, without limitation, carcinomas (e.g., squamous-cell carcinomas, adenocarcinomas, hepatocellular carcinomas, and renal cell carcinomas), particularly those of the bladder, bone, bowel, breast, cervix, colon (colorectal), esophagus, head, kidney, liver, lung, nasopharyngeal, neck, ovary, pancreas, prostate, and stomach; leukemias, such as acute myelogenous leukemia, acute lymphocytic leukemia, acute

promyelocytic leukemia (APL), acute T-cell lymphoblastic leukemia, adult T-cell leukemia, basophilic leukemia, eosinophilic leukemia, granulocytic leukemia, hairy cell leukemia, leukopenic leukemia, lymphatic leukemia, lymphoblastic leukemia, lymphocytic leukemia, megakaryocytic leukemia, micromyeloblastic leukemia, monocytic leukemia, neutrophilic leukemia and stem cell leukemia; benign and malignant lymphomas, particularly Burkitt's lymphoma, Non-Hodgkin's lymphoma and B-cell lymphoma; benign and malignant melanomas; myeloproliferative diseases; sarcomas, particularly Ewing's sarcoma, hemangiosarcoma, Kaposi's sarcoma, liposarcoma, myosarcomas, peripheral neuroepithelioma, and synovial sarcoma; tumors of the central nervous system (e.g., gliomas, astrocytomas, oligodendrogliomas, ependymomas, glioblastomas, neuroblastomas, ganglioneuromas, gangliogliomas, medulloblastomas, pineal cell tumors, meningiomas, meningeal sarcomas, neurofibromas, and Schwannomas); germ-line tumors (e.g., bowel cancer, breast cancer, prostate cancer, cervical cancer, uterine cancer, lung cancer (e.g., small cell lung cancer, mixed small cell and non-small cell cancer, pleural mesothelioma, including metastatic pleural mesothelioma small cell lung cancer and non-small cell lung cancer), ovarian cancer, testicular cancer, thyroid cancer, astrocytoma, esophageal cancer, pancreatic cancer, stomach cancer, liver cancer, colon cancer, and melanoma; mixed types of neoplasias, particularly carcinosarcoma and Hodgkin's disease; and tumors of mixed origin, such as Wilms' tumor and teratocarcinomas, among others. It is noted that certain epithelial tumors including ovarian, breast, colon, head and neck, medulloblastoma and B-cell lymphoma, among others are shown to exhibit increased autophagy and are principal target cancers for compounds and therapies according to the present invention.

The term "additional anti-cancer agent" is used to describe an additional compound which may be coadministered with one or more compounds of the present invention in the treatment of cancer. Such agents include, for example, everolimus, trabectedin, abraxane, TLK 286, AV-299, DN-101, pazopanib, GSK690693, RTA 744, ON 0910.Na, AZD 6244 (ARRY-142886), AMN-107, TKI-258, GSK461364, AZD 1152, enzastaurin, vandetanib, ARQ-197, MK-0457, MLN8054, PHA-739358, R-763, AT-9263, a FLT-3 inhibitor, a VEGFR inhibitor, an EGFR TK inhibitor, an aurora kinase inhibitor, a PIK-1 modulator, a Bcl-2 inhibitor, an HDAC inhibitor, a c-MET inhibitor, a PARP inhibitor, a Cdk inhibitor, an EGFR TK inhibitor, an IGFR-TK inhibitor, an anti-HGF antibody, a PI3 kinase inhibitors, an AKT inhibitor, a JAK/STAT inhibitor, a checkpoint-1 or 2 inhibitor, a focal adhesion kinase inhibitor, a Map kinase kinase (mek) inhibitor, a VEGF trap antibody, pemetrexed, erlotinib,

dasatanib, nilotinib, decatanib, panitumumab, amrubicin, oregovomab, Lep-etu, nolatrexed, azd2171, batabulin, ofatumumab, zanolimumab, edotecarin, tetrandrine, rubitecan, tesmilifene, oblimersen, ticilimumab, ipilimumab, gossypol, Bio 111, 131-I-TM-601, ALT-110, BIO 140, CC 8490, cilengitide, gimatecan, IL13-PE38QQR, INO 1001, IPdR₁ KRX-0402, lucanthone, LY 317615, neuradiab, vitespan, Rta 744, Sdx 102, talampanel, atrasentan, Xr 311, romidepsin, ADS- 100380, sunitinib, 5-fluorouracil, vorinostat, etoposide, gemcitabine, doxorubicin, irinotecan, liposomal doxorubicin, 5'-deoxy-5-fluorouridine, vincristine, temozolomide, ZK-304709, seliciclib; PD0325901, AZD-6244, capecitabine, L-Glutamic acid, N-[4-[2-(2-amino-4,7-dihydro-4-oxo-1 H - pyrrolo[2,3- d]pyrimidin-5-yl)ethyl]benzoyl]-, disodium salt, heptahydrate, camptothecin, PEG-labeled irinotecan, tamoxifen, toremifene citrate, anastrozole, exemestane, letrozole, DES(diethylstilbestrol), estradiol, estrogen, conjugated estrogen, bevacizumab, IMC-1C11, CHIR-258,); 3-[5-(methylsulfonylpiperadinemethyl)- indolyl]-quinolone, vatalanib, AG-013736, AVE-0005, the acetate salt of [D- Ser(Bu t) 6 ,Azgly 10] (pyro-Glu-His-Trp-Ser-Tyr-D-Ser(Bu t)-Leu-Arg-Pro- Azgly-NH₂ acetate [C₅₉H₈₄N₁₈O₁₄ -(C₂H₄O₂)_x where x = 1 to 2.4], goserelin acetate, leuprolide acetate, triptorelin pamoate, medroxyprogesterone acetate, hydroxyprogesterone caproate, megestrol acetate, raloxifene, bicalutamide, flutamide, nilutamide, megestrol acetate, CP-724714; TAK-165, HKI-272, erlotinib, lapatanib, canertinib, ABX-EGF antibody, erbitux, EKB-569, PKI-166, GW-572016, Ionafarnib, BMS-214662, tipifarnib; amifostine, NVP-LAQ824, suberoyl analide hydroxamic acid, valproic acid, trichostatin A, FK-228, SU11248, sorafenib, KRN951, aminoglutethimide, arnsacrine, anagrelide, L-asparaginase, Bacillus Calmette-Guerin (BCG) vaccine, bleomycin, buserelin, busulfan, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, diethylstilbestrol, epirubicin, fludarabine, fludrocortisone, fluoxymesterone, flutamide, gemcitabine, gleevac, hydroxyurea, idarubicin, ifosfamide, imatinib, leuprolide, levamisole, lomustine, mechlorethamine, melphalan, 6-mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, octreotide, oxaliplatin, pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, streptozocin, teniposide, testosterone, thalidomide, thioguanine, thiotepa, tretinoin, vindesine, 13-cis-retinoic acid, phenylalanine mustard, uracil mustard, estramustine, altretamine, floxuridine, 5-deoxyuridine, cytosine arabinoside, 6-mecaptopurine, deoxycoformycin, calcitriol, valrubicin, mithramycin, vinblastine, vinorelbine, topotecan, razoxin, marimastat, COL-3, neovastat, BMS-275291, squalamine, endostatin, SU5416, SU6668, EMD121974, interleukin-12, IM862, angiostatin,

vitaxin, droloxifene, idoxifene, spironolactone, finasteride, cimitidine, trastuzumab, denileukin diftitox, gefitinib, bortezomib, paclitaxel, irinotecan, topotecan, doxorubicin, docetaxel, vinorelbine, bevacizumab (monoclonal antibody) and erbitux, cremophor-free paclitaxel, epithilone B, BMS- 247550, BMS-310705, droloxifene, 4-hydroxytamoxifen, pipendoxifene, ERA- 923, arzoxifene, fulvestrant, acolbifene, lasofoxifene, idoxifene, TSE- 424, HMR- 3339, ZK186619, PTK787/ZK 222584, VX-745, PD 184352, rapamycin, 40-O-(2-hydroxyethyl)-rapamycin, temsirolimus, AP-23573, RAD001 , ABT-578, BC-210, LY294002, LY292223, LY292696, LY293684, LY293646, wortmannin, ZM336372, L- 779,450, PEG-filgrastim, darbepoetin, erythropoietin, granulocyte colony-stimulating factor, zolendronate, prednisone, cetuximab, granulocyte macrophage colony-stimulating factor, histrelin, pegylated interferon alfa-2a, interferon alfa-2a, pegylated interferon alfa-2b, interferon alfa-2b, azacitidine, PEG-L-asparaginase, lenalidomide, gemtuzumab, hydrocortisone, interleukin-11 , dexrazoxane, alemtuzumab, all-transretinoic acid, ketoconazole, interleukin-2, megestrol, immune globulin, nitrogen mustard, methylprednisolone, ibritumomab tiuxetan, androgens, decitabine, hexamethylmelamine, bexarotene, tositumomab, arsenic trioxide, cortisone, editronate, mitotane, cyclosporine, liposomal daunorubicin, Edwina-asparaginase, strontium 89, casopitant, netupitant, an NK-1 receptor antagonists, palonosetron, aprepitant, , diphenhydramine, hydroxyzine, metoclopramide, lorazepam, alprazolam, haloperidol, droperidol, dronabinol, dexamethasone, methylprednisolone, prochlorperazine, granisetron, ondansetron, dolasetron, tropisetron, sspegfilgrastim, erythropoietin, epoetin alfa and darbepoetin alfa, ipilimumab, vemurafenib among others.

The term "alkyl" is used herein to refer to a fully saturated monovalent radical containing carbon and hydrogen (up to 10 carbon atoms or as otherwise indicated), and which may be a straight chain, branched or cyclic. Examples of alkyl groups are methyl, ethyl, n-butyl, n-heptyl, isopropyl, 2-methyl propyl, tert-butyl, neopentyl, hexyl, heptyl, octyl, nonyl, decyl, etc.

The term "substituted" as that term relates to alkyl groups which are described above include one or more functional groups such as lower alkyl groups containing 1-6 carbon atoms which are optionally substituted with 1 or 2 hydroxyl groups or between 1 and 5 (preferably 3-5) fluoro groups, acyl (C₁-C₆), halogen (F, Cl, Br, I, e.g., alkyl halos, e.g., CF₃), amido, hydroxyl, carboxy/carboxylic acid, thioamido, cyano, nitro, alkenyl (C₂-C₆) alkynyl

(C₂-C₆), azido, alkoxy (C₁-C₆), (including alkoxy groups which are further substituted with a C₁-C₆ alkoxy group thus producing a diether group), amino, C₁-C₆ alkylamino and dialkylamino, where the alkyl groups may be optionally substituted with 1 or 2 hydroxyl groups or an amine, aminoalkyl or dialkyl group which itself is substituted one or two alkyl groups or a 7-substituted-4-quinolinyl group, C₂-C₆ acylamino, C₂-C₆ oxyacylester or carboxyester, aryloxy, aryloxy(C₁-C₆)alkyl, carboxamido, thio, C₂-C₆ ether or thioether, a 7-substituted-4-aminoquinolinyl group (or a substitution on an amine group which forms a 7-substituted-4-aminoquinolinyl group) and the like. Preferred substituents on alkyl groups (within context, especially on the amino group of the 7-substituted-4-aminoquinoline) or a linker which contains at least one amine group, include, for example, at least one hydroxyl group, an amine, monoalkyl amine or dialkyl amine (where one or both alkyl groups is itself further optionally substituted with a dialkyl amine or an amine substituted with one or two (preferably one) 7-substituted-4-quinolinyl group(s) where the amine group is bonded to the 4-position of the quinolinyl group) or an alkoxy group (e.g. methoxy or ethoxy) which may be further substituted with an alkoxy group, preferably a methoxy group, thus forming a diether substituent.

The term "aryl" refers to a substituted or unsubstituted monovalent aromatic radical having a single ring (e.g., phenyl) or multiple condensed rings (e.g., naphthyl). Other examples include heterocyclic aromatic (heteroaromatic or heteroaryl) ring groups having one or more nitrogen, oxygen, or sulfur atoms in the ring, in particular, quinoline groups, in particular, 7-substituted-amino quinoline groups, as well as other groups.

The term "substituted" as used in the term "substituted aryl, substituted aromatic, substituted heteroaryl, or substituted heteroaromatic" herein signifies that a substitution on the 7-position of 4-aminoquinoline may be present, said substituents being selected from atoms and groups, which when present enhance the activity of the compound as an inhibitor of autophagy. Examples of substituents that may be present in a substituted aromatic or heteroaromatic group include, but are not limited to, groups such as H, halo (F, Cl, Br or I), CN, NO₂, optionally substituted C₁-C₆ alkyl (when substituted, preferably substituted with 1 or 2 hydroxyl groups or 3-5 fluoro groups), optionally substituted O-C₁-C₆ alkyl (preferably, OCH₃), optionally substituted C₂-C₇ acyl (preferably acetyl) or optionally substituted C₂-C₇ ester (oxycarbonyl ester or carboxyester, preferably carboxyester);. It is noted that each of the substituents disclosed herein may themselves be substituted.

The term “co-administration” or “adjunct therapy” shall mean that at least two compounds or compositions are administered to the patient at the same time, such that effective amounts or concentrations of each of the two or more compounds may be found in the patient at a given point in time. Although compounds according to the present invention may be co-administered to a patient at the same time, the term embraces both administration of two or more agents at the same time or at different times, including sequential administration. Preferably, effective concentrations of all co-administered compounds or compositions are found in the subject at a given time. The term co-administration or adjunct therapy also contemplates other bioactive agents being coadministered with pharmaceutical compositions according to the present invention, especially where a cancer has metastasized or is at risk for metastasis.

The term “radiotherapy” or “radiation therapy” is used to describe therapy for cancer which may be used in conjunction with the present compounds. Radiation therapy uses high doses of radiation, such as X-rays, or other energy sources such as radioisotopes (gamma, beta or alpha emitters), to destroy cancer cells. The radiation damages the genetic material of the cells so that they can't grow. Although radiation damages normal cells as well as cancer cells, the normal cells can repair themselves and function, while the cancer cells cannot.

Radiation therapy may be used in combination with the presently claimed compounds, alone or in combination with additional anticancer compounds as otherwise disclosed herein, depending on the cancer to be treated. Radiotherapy therapy is most effective in treating cancers that have not spread outside the area of the original cancer, but it also may be used if the cancer has spread to nearby tissue. Radiotherapy is sometimes used after surgery to destroy any remaining cancer cells and to relieve pain from metastatic cancer.

Pharmaceutical Compositions

Compounds according to the present invention may be readily formulated into pharmaceutical compositions, useful in the inhibition of autophagy in a biological system and/or the inhibition, treatment or prevention of diseases states and/or conditions which benefit from the inhibition of autophagy including cancer (and its metastasis), rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus (systemic lupus erythematosus), chronic urticaria and Sjogren's disease. Pharmaceutical compositions comprise an effective

amount of one or more compounds according to the present invention in combination with a pharmaceutically acceptable carrier, additive or excipient, optionally in combination with at least one additional agent, in the case of cancer, preferably an anticancer agent as otherwise described herein. .

As noted above, the compounds and method of the invention may be used to inhibit autophagy as otherwise described herein, and are useful for the inhibition (including prophylaxis) and/or treatment of cancer and its metastasis, rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus (systemic lupus erythematosus), chronic urticaria and Sjogren's disease. The treatment of cancer or malaria are important aspects of the present invention.

In methods according to the present invention, subjects or patients in need are treated with the present compounds, pharmaceutical compositions in order to inhibit, reduce the likelihood or treat a disease state, condition and/or infection as otherwise described herein. The disease states, conditions and infections treated by the present compounds and compositions are readily recognized and diagnosed by those of ordinary skill in the art and treated by administering to the patient an effective amount of one or more compounds according to the present invention.

Generally, dosages and routes of administration of the compound are determined according to the size and condition of the subject, according to standard pharmaceutical practices. Dose levels employed can vary widely, and can readily be determined by those of skill in the art. Typically, amounts in the milligram up to gram quantities are employed. The composition may be administered to a subject by various routes, e.g. orally, transdermally, perineurally or parenterally, that is, by intravenous, subcutaneous, intraperitoneal, or intramuscular injection, among others, including buccal, rectal and transdermal administration. Subjects contemplated for treatment according to the method of the invention include humans, companion animals, laboratory animals, and the like.

Formulations containing the compounds according to the present invention may take the form of solid, semi-solid, lyophilized powder, or liquid dosage forms, such as, for example, tablets, capsules, powders, sustained-release formulations, solutions, suspensions,

emulsions, suppositories, creams, ointments, lotions, aerosols, patches or the like, preferably in unit dosage forms suitable for simple administration of precise dosages.

Pharmaceutical compositions according to the present invention typically include a conventional pharmaceutical carrier or excipient and may additionally include other medicinal agents, carriers, adjuvants, additives and the like. Preferably, the composition is about 0.1% to about 85%, about 0.5% to about 75% by weight of a compound or compounds of the invention, with the remainder consisting essentially of suitable pharmaceutical excipients. For oral administration, such excipients include pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, gelatin, sucrose, magnesium carbonate, and the like. If desired, the composition may also contain minor amounts of non-toxic auxiliary substances such as wetting agents, emulsifying agents, or buffers.

Liquid compositions can be prepared by dissolving or dispersing the compounds (about 0.5% to about 20% by weight or more), and optional pharmaceutical adjuvants, in a carrier, such as, for example, aqueous saline, aqueous dextrose, glycerol, or ethanol, to form a solution or suspension. For use in oral liquid preparation, the composition may be prepared as a solution, suspension, emulsion, or syrup, being supplied either in liquid form or a dried form suitable for hydration in water or normal saline.

When the composition is employed in the form of solid preparations for oral administration, the preparations may be tablets, granules, powders, capsules or the like. In a tablet formulation, the composition is typically formulated with additives, e.g. an excipient such as a saccharide or cellulose preparation, a binder such as starch paste or methyl cellulose, a filler, a disintegrator, and other additives typically used in the manufacture of medical preparations.

An injectable composition for parenteral administration will typically contain the compound in a suitable i.v. solution, such as sterile physiological salt solution. The composition may also be formulated as a suspension in a lipid or phospholipid, in a liposomal suspension, or in an aqueous emulsion.

Methods for preparing such dosage forms are known or is apparent to those skilled in the art; for example, see Remington's Pharmaceutical Sciences (17th Ed., Mack Pub. Co., 1985). The composition to be administered will contain a quantity of the selected compound in a pharmaceutically effective amount for inhibiting autophagy in a biological system, including a patient or subject according to the present invention.

Synthesis of Compounds According to the Present Invention

Strategy for synthesis of bivalent aminoquinoline autophagy inhibitors

The inventors have examined the application of the strategy of polyvalency (11, 12) to the synthesis of novel autophagy inhibitors by preparing a dimeric chloroquine (Figure 1, compound **3**: **Lys01**), from commercially available materials. Based on literature precedent(14), we envisioned the preparation of compound **3** from one equivalent of compound **5** and two equivalents of compound **6** (14), as outlined retrosynthetically in Figure 2.

While compound **4** (R=Cl) is a known compound (14), the bisquinoline compound **3** (R₂=Me) has not been described in the literature. Due its putative lysosomotropism, we refer to compound **3** as Lys01. Reaction of compound **5** with two equivalents of compound **4** led to the formation of a mixture of the desired product compound **3** along with some of the monoquinoline compound **7** (Figure 1, Lys02, Figure 2), the synthesis of which was previously reported by Higuchi (18). To examine the role of the C-7 chlorine substituents in compound **3**, we prepared compound **9** (Figure 1, compound **9**: Lys 03), the dimethoxy analog of compound **3**.

To determine the importance of the polyamine connector of compound **3**, we prepared the polyether analog compound **11** (Figure 1, compound **11**: Lys04) of compound **3** from the commercially available 2,2'-(ethylenedioxy)bis(ethylamine) **10** (see Figure 2).

In an effort to obtain SAR data on the lead compound **3** (**Lys01**), the inventors examined the systematic modification of the structure of compound **3** (i.e., **12** R=Cl), as outlined in Scheme A of Figure 15A. Initial efforts focused on changes in three different areas of the structure of compound **12** as examined: 1) modification of the C-7 chlorine substituent present in compound **3** (X in compound **12** of figure 15A) ; 2) modification of the


C-4 nitrogen substituent, i.e., N-alkylation or acylation, as well as the adjacent carbon atom (that contains a stereocenter in CQ and HCQ (Figure 1); and 3) modification of the N-methyl group in compound **12** (Figure 15A).

Each of the requisite starting compounds outlined in Figure 15A (compounds **13-16**) are known or commercially available, facilitating the synthesis of a family of analogs that differ from compound **3** (**12** R=Cl) by incorporating different electron-withdrawing groups, based on the work of Lee (16).

The inventors also examine the biological activity of N-alkylated and N-acylated analogs of compound **12**, compound **17** and compound **18** (Figure 15A). The preparation of these novel compounds would proceed directly from **12** (R=Cl), either by direct alkylation, reductive alkylation, or acylation.

The inventors also examine the introduction into compound **12** of the chirality that is present in CQ and HCQ (Figure 1) as outlined in compound **19** (Scheme C of Figure 15A). The requisite linker compound **20** could be obtained using the method of Kokotos (J. Chem. Res, Synopses 1992, 12, 391).


Finally, the modification of the structure of compound **12** via replacement of the N-methyl group (Scheme A of Figure 15A) with other functionalities is examined. Two intriguing possibilities are shown in Scheme D of Figure 15A, i.e., compound **21** and compound **22**, in which the N-methyl group is replaced with the hydroxyethyl group present in HCQ, and in which the N-methyl group is replaced by another quinoline moiety to produce tris-quinoline compound **22**. The synthetic routes for the formation of **21** and **22** are based closely on the work of Lee (19) and Solomon(17), respectively.




While characterization of the second generation compounds described continue, there are additional compounds which are believed to represent more potent autophagy inhibitors. Taking the lead compound as compound **12** (scheme 1; Lys01), the inventors describe more extensive systematic modification of three parts of the structure of **12**, as outlined in the generic structure **23** (see above) as the next logical step in SAR analysis. Each of the three parts of the structure (R1, R2 and R3) is modified, as outlined below.

The Role of R1

The effect of substituting the chlorine moiety in chloroquine **1** (scheme 2) has been examined




by Egan and coworkers (20), who established that electron-withdrawing groups are important for the antiparasmodial activity of these 7-substituted quinolines. The inventors therefore examined these same substitutions in the case of the generic structure **23**, substituting both R¹ and R^{1'} in **23**, and also examining the effect of monosubstitution, in which the R^{1'} substituent is then either Cl as in **12** or hydrogen (X=H). The Egan study indicates that all of the requisite 4-chloro-7-substituted-quinolines are known, thereby facilitating the preparation of each of the compounds shown in Scheme 3. None of the compounds outlined in scheme 3 are known, although the synthesis of linked bis-quinolines (R1=H) has been described (21).




The Role of R2


The role of $R^2=H$ in lead structure **3** is examined by either acetylation ($R^2=MeCO-$; mono- or di-) or methylation ($R^2=Me$; mono- or di-) to give the structures **41-44** shown in Scheme 4. While the homologous analogs **45** and **46** (containing one or two propylene chains between the nitrogen atoms) are known (FR 1345573; CAN 60:68181), each of the analogs **41-44** represents a novel structure.



Well-tolerated substitution at R^2 further directs substitution of R^2 with a hydroxyethyl moiety, as found in hydroxychloroquine **2** (scheme 5). The reaction of anilines with ethylene oxide to give the corresponding hydroxyethyl compounds is well-precedented (22), so that conversion of the lead structure **3** to either mono- or di-hydroxyethylated analogs **47** and **48** is readily achieved.




The inventors also examine the preparation of tri- and tetra-quinoline containing structures by oxidation of the primary alcohols in **47** and **48** to give the corresponding aldehydes, which on reductive alkylation with 7-chloro-4-aminoquinoline gives the tri- and tetraquinolines **49** and **50**, based on the work of Bailey and coworkers (23), as shown in scheme 6.




The inventors also examine the effect of incorporation of both lipophilic groups, i.e., long chain alkyl, as well as more polar substituents in the place of R^2 in the lead structure in Scheme 7, via alkylation of the secondary amines ($R^2=H$) with the commercially available alkylating agents **59-62**, leading to the preparation of **51-58** via mono- and dialkylation of the


two secondary amine functionalities, based on the work of Drefahl and König (Chem. Ber. 1954, 87, 1632-4).



The inventors examine the effect of changing R^3 in the lead structure **23** (see scheme 1, above). Certain substitutions for R^3 are already known, such as shown in **63-65** in Scheme 8, below. The inventors prepare a series of new analogs, based on the alkylation of the known secondary amine **63** (13) with the alkylating agents shown in scheme 7 (**59-62**) to generate the novel structures (**66-69**). The substrate **70** in which $R^3 = CH_2CH_2OH$, i.e., the analog of **63** that corresponds most closely to hydroxychloroquine is examined. This compound is available by the same sequence used to prepare the analogs shown in Scheme 5, or by demethylation of **66**.




Another important difference between the lead structure **3** and chloroquine **1** (scheme 9) is the presence of the stereocenter next to the nitrogen atom in **1**. The inventors also prepare hybrid structures, in which either one or both of the sides of **23** more closely resemble **1**, such as **71/72** and **73/74**. The requisite diamines in each case are prepared starting from (L) glutamic acid, following the procedure employed by Craig in the stereoselective synthesis of chloroquine (J. Org. Chem. 1988, 53, 1167-1170).



The inventors prepare the analog of the lead structure **3** containing the stereocenter present in chloroquine **1**, i.e., **75/76** and **77/78**, which are available from alanine and serine, respectively, using the method of Charlton and coworkers (24) via reductive alkylation.

Finally, the inventors examine a series of compounds that contain four nitrogen atoms in the tether connecting the quinoline rings, instead of the three nitrogen atoms that are present in the connector chain in **3** (scheme 9), as illustrated in scheme 10, below. Denny and coworkers (24) have described the synthesis of the requisite tetramines **81** and **82**.

Attachment of the additional chloroquine moieties present in **80** takes place via the same methodology employed in Scheme 8 for the synthesis of **65**.



Method of Treatment


According to one aspect of the invention, a method is provided for treating a mammalian patient or subject to inhibit autophagy in that patient or subject. Compounds according to the present invention described herein may be used to inhibit autophagy in a manner consistent with inhibiting, treating and/or preventing disease states and/or conditions including cancer (including metastasis of cancer), rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria and Sjogren's disease.

According to the present invention, in patients or subjects in need thereof, are treated by administering to the patient or subject an effective amount of one or more compounds according to the present invention, optionally in combination with at least one additional bioactive agent useful for treating the same disease state or condition. Compounds according to the present invention may be used to inhibit, reduce the likelihood or treat cancer, including the metastasis of cancer in a patient or subject in need of such treatment. The treatment is useful for any cancer for which inhibition of autophagy represents a favorable result or for which metastasis is a risk element. Therapy with at least one additional

anticancer agent as otherwise described herein is also contemplated in the present methods. The numerous cancers which may be treated pursuant to the present method are described hereinabove.

In another aspect the present invention is directed to a method for treating a disease state and/or condition which benefits from the inhibition of autophagy, including rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria and Sjorgen's disease. In this method, a patient or subject in need of treatment is administered an effective amount of a compound as otherwise described herein optionally in combination with a pharmaceutically acceptable carrier, additive or excipient in order to inhibit, treat and/or prevent the above disease states of conditions.

In the present invention, the method of treatment comprises administering to the subject in need of treatment, in a pharmaceutically acceptable carrier, an effective amount of a compound according to I below:



Wherein R^1 and $R^{1'}$ are each independently H, halo (F, Cl, Br or I), CN, NO₂, optionally substituted C₁-C₆ alkyl (when substituted, preferably substituted with 1 or 2 hydroxyl groups or 3-5 fluoro groups), optionally substituted O-C₁-C₆ alkyl (preferably, OCH₃), optionally substituted C₂-C₇ acyl (preferably acetyl) or optionally substituted C₂-C₇ ester (oxycarbonyl ester or carboxyester, preferably carboxyester);

R and R' are each independently H, a C₁-C₆ optionally substituted alkyl group, a C₁-C₇ (preferably C₂-C₇) optionally substituted acyl group, a C₂-C₇ optionally substituted carboxy ester group (which forms a urethane group with the nitrogen atom to which R or R' is bonded);

L is a $\text{---}(\text{CH}_2\text{Y})_n\text{---X---}(\text{Y}'\text{CH}_2)_n\text{---}$ group or a $\text{A---}(\text{CH}_2\text{---CH}_2\text{---Z})_n\text{---A}'$ group (either A or A' may be bonded to either of the two amine groups in compound I) wherein at least one of the CH_2 groups in L is optionally substituted with a $\text{C}_1\text{--C}_3$ alkyl group which itself is optionally substituted with one or two hydroxyl groups;

X is absent, $(\text{CH}_2)_j$, O, S or $\text{N--R}''$;

Y is absent, CH_2 , O, CH_2O or $\text{N--R}''$ and Y' is absent CH_2 , O, OCH_2 or $\text{N--R}''$, with the proviso that when one or more of X, Y and Y' is present, each of X and Y, X and Y' or Y and Y', when present, forms a stable bond;

R'' is H or an optionally substituted $\text{C}_1\text{--C}_6$ (preferably $\text{C}_1\text{--C}_3$) alkyl group;

j is 1, 2 or 3 (preferably 1 or 2);

n is 0, 1, 2, 3 or 4, with the proviso that when n is 0, X is $(\text{CH}_2)_j$ where j is at least 1 and at least one CH_2 group is optionally substituted with a $\text{C}_1\text{--C}_3$ alkyl group which itself is optionally substituted with one or two hydroxyl groups;

A is absent or $(\text{CH}_2)_j$ and A' is $(\text{CH}_2)_j$ wherein at least one CH_2 group in A or A' is optionally substituted with a $\text{C}_1\text{--C}_3$ alkyl group which is itself optionally substituted with one or two hydroxyl groups;

Z is O or N--R^Z ;

R^Z is H or an optionally substituted $\text{C}_1\text{--C}_3$ alkyl group,

or a pharmaceutically acceptable salt, enantiomer, diastereomer, solvent or polymorph thereof.

In certain preferred methods of the invention, R^1 and $\text{R}^{1'}$ are each independently H, a halo group, a nitro group or a trifluoromethyl group, preferably a chloro group. R and R' are preferably each independently H, a $\text{C}_1\text{--C}_3$ optionally substituted alkyl group itself preferably substituted with at least one hydroxyl group, an amine, monoalkyl amine or dialkyl amine group, wherein said amine group or said monoalkyl amine group is optionally substituted on the amine position with a 7-substituted-4-quinolinyl group wherein the amine binds to the 4-position of the quinolinyl group, or one or both alkyl groups of said monoalkyl amine or dialkyl amine is itself further optionally substituted with at least one hydroxyl group, an amine, a monoalkyl amine or a dialkyl amine wherein the amine or monoalkyl amine is optionally substituted on the amine position with one or two 7-substituted-quinolinyl group(s) (the 7-position of each quinolinyl group may be substituted with R^1 and/or $\text{R}^{1'}$ as broadly described for generic structure I above), or an alkoxy group (e.g. methoxy or ethoxy) which

alkoxy group may be further substituted with an alkoxy group, preferably a methoxy group (thus forming a diether substituent).

In other preferred methods of the invention L is a $\text{---}(\text{CH}_2\text{Y})_n\text{---X---}(\text{Y}'\text{CH}_2)_n\text{---}$ group, where X is N-R'', Y and Y' are each independently absent or CH₂, and R'' is H or a C₁-C₃ alkyl group which is optionally substituted with at least one hydroxyl group, an amine, monoalkyl amine or dialkyl amine group, wherein said amine group or said monoalkyl amine group is optionally substituted on the amine position with a 7-substituted-4-quinolinyl group wherein the amine binds to the 4-position of the quinolinyl group, or one or both alkyl groups of said monoalkyl amine or dialkyl amine is itself further optionally substituted with at least one hydroxyl group, an amine, a monoalkyl amine or a dialkyl amine wherein the amine or monoalkyl amine is optionally substituted on the amine position with one or two 7-substituted-quinolinyl group(s) (the 7-position of each quinolinyl group may be substituted with R¹ and/or R^{1'} as broadly described for generic structure I above) for the), or an alkoxy group (e.g. methoxy or ethoxy) which alkoxy group may be further substituted with an alkoxy group, preferably a methoxy group (thus forming a diether substituent).

Further preferred methods relate to the use/administration of the compounds according to the present invention which are presented in the various schemes which are presented in Scheme 1 and Schemes 3-10 and figures 14, 15 and 15A as presented herein.

In the methods treating or inhibiting cancer or the metastasis of cancer, the compounds described above may be coadministered with at least one additional anticancer agent including, for example, everolimus, trabectedin, abraxane, TLK 286, AV-299, DN-101, pazopanib, GSK690693, RTA 744, ON 0910.Na, AZD 6244 (ARRY-142886), AMN-107, TKI-258, GSK461364, AZD 1152, enzastaurin, vandetanib, ARQ-197, MK-0457, MLN8054, PHA-739358, R-763, AT-9263, a FLT-3 inhibitor, a VEGFR inhibitor, an EGFR TK inhibitor, an aurora kinase inhibitor, a PIK-1 modulator, a Bcl-2 inhibitor, an HDAC inhibitor, a c-MET inhibitor, a PARP inhibitor, a Cdk inhibitor, an EGFR TK inhibitor, an IGFR-TK inhibitor, an anti-HGF antibody, a PI3 kinase inhibitors, an AKT inhibitor, a JAK/STAT inhibitor, a checkpoint-1 or 2 inhibitor, a focal adhesion kinase inhibitor, a Map kinase kinase (mek) inhibitor, a VEGF trap antibody, pemetrexed, erlotinib, dasatanib, nilotinib, decatanib, panitumumab, amrubicin, oregovomab, Lep-etu, nolatrexed, azd2171, batabulin,

ofatumumab, zanolimumab, edotecarin, tetrandrine, rubitecan, tesmilifene, oblimersen, ticilimumab, ipilimumab, gossypol, Bio 111, 131-I-TM-601, ALT-110, BIO 140, CC 8490, cilengitide, gimatecan, IL13-PE38QQR, INO 1001, IPdR₁ KRX-0402, lucanthone, LY 317615, neuradiab, vitespan, Rta 744, Sdx 102, talampanel, atrasentan, Xr 311, romidepsin, ADS- 100380, sunitinib, 5-fluorouracil, vorinostat, etoposide, gemcitabine, doxorubicin, irinotecan, liposomal doxorubicin, 5'-deoxy-5-fluorouridine, vincristine, temozolomide, ZK-304709, seliciclib; PD0325901, AZD-6244, capecitabine, L-Glutamic acid, N-[4-[2-(2-amino-4,7-dihydro-4-oxo-1 H - pyrrolo[2,3- d]pyrimidin-5-yl)ethyl]benzoyl]-, disodium salt, heptahydrate, camptothecin, PEG-labeled irinotecan, tamoxifen, toremifene citrate, anastrozole, exemestane, letrozole, DES(diethylstilbestrol), estradiol, estrogen, conjugated estrogen, bevacizumab, IMC-1C11, CHIR-258,); 3-[5-(methylsulfonylpiperadinemethyl)-indolyl]-quinolone, vatalanib, AG-013736, AVE-0005, the acetate salt of [D- Ser(Bu t) 6 ,Azgly 10] (pyro-Glu-His-Trp-Ser-Tyr-D-Ser(Bu t)-Leu-Arg-Pro- Azgly-NH₂ acetate [C₅₉H₈₄N₁₈Oi₄ -(C₂H₄O₂)_x where x = 1 to 2.4], goserelin acetate, leuprolide acetate, triptorelin pamoate, medroxyprogesterone acetate, hydroxyprogesterone caproate, megestrol acetate, raloxifene, bicalutamide, flutamide, nilutamide, megestrol acetate, CP-724714; TAK-165, HKI-272, erlotinib, lapatanib, canertinib, ABX-EGF antibody, erbitux, EKB-569, PKI-166, GW-572016, Ionafarnib, BMS-214662, tipifarnib; amifostine, NVP-LAQ824, suberoyl analide hydroxamic acid, valproic acid, trichostatin A, FK-228, SU11248, sorafenib, KRN951, aminoglutethimide, arnsacrine, anagrelide, L-asparaginase, Bacillus Calmette-Guerin (BCG) vaccine, bleomycin, buserelin, busulfan, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, diethylstilbestrol, epirubicin, fludarabine, fludrocortisone, fluoxymesterone, flutamide, gemcitabine, gleevac, hydroxyurea, idarubicin, ifosfamide, imatinib, leuprolide, levamisole, lomustine, mechlorethamine, melphalan, 6-mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, octreotide, oxaliplatin, pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, streptozocin, teniposide, testosterone, thalidomide, thioguanine, thiotepa, tretinoin, vindesine, 13-cis-retinoic acid, phenylalanine mustard, uracil mustard, estramustine, altretamine, floxuridine, 5-deoxyuridine, cytosine arabinoside, 6-mecaptopurine, deoxycorformycin, calcitriol, valrubicin, mithramycin, vinblastine, vinorelbine, topotecan, razoxin, marimastat, COL-3, neovastat, BMS-275291, squalamine, endostatin, SU5416, SU6668, EMD121974, interleukin-12, IM862, angiostatin, vitaxin, droloxifene, idoxifene, spironolactone, finasteride, cimitidine, trastuzumab, denileukin diftitox, gefitinib, bortezomib, paclitaxel,

irinotecan, topotecan, doxorubicin, docetaxel, vinorelbine, bevacizumab (monoclonal antibody) and erbitux, cremophor-free paclitaxel, epithilone B, BMS- 247550, BMS-310705, droloxifene, 4-hydroxytamoxifen, pipendoxifene, ERA- 923, arzoxifene, fulvestrant, acolbifene, lasofoxifene, idoxifene, TSE-424, HMR- 3339, ZK186619, PTK787/ZK 222584, VX-745, PD 184352, rapamycin, 40-O-(2-hydroxyethyl)-rapamycin, temsirolimus, AP-23573, RAD001 , ABT-578, BC-210, LY294002, LY292223, LY292696, LY293684, LY293646, wortmannin, ZM336372, L-779,450, PEG-filgrastim, darbepoetin, erythropoietin, granulocyte colony-stimulating factor, zoledronate, prednisone, cetuximab, granulocyte macrophage colony-stimulating factor, histrelin, pegylated interferon alfa-2a, interferon alfa-2a, pegylated interferon alfa-2b, interferon alfa-2b, azacitidine, PEG-L-asparaginase, lenalidomide, gemtuzumab, hydrocortisone, interleukin-11 , dexrazoxane, alemtuzumab, all-transretinoic acid, ketoconazole, interleukin-2, megestrol, immune globulin, nitrogen mustard, methylprednisolone, ibritumomab tiuxetan, androgens, decitabine, hexamethylmelamine, bexarotene, tositumomab, arsenic trioxide, cortisone, editronate, mitotane, cyclosporine, liposomal daunorubicin, Edwina-asparaginase, strontium 89, casopitant, netupitant, an NK-1 receptor antagonists, palonosetron, aprepitant, , diphenhydramine, hydroxyzine, metoclopramide, lorazepam, alprazolam, haloperidol, droperidol, dronabinol, dexamethasone, methylprednisolone, prochlorperazine, granisetron, ondansetron, dolasetron, tropisetron, pegfilgrastim, erythropoietin, epoetin alfa and darbepoetin alfa, among others, and mixtures thereof.

In methods involving infections, disease states and/or conditions caused by rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria and Sjogren's disease, the compounds according to the present invention may be coadministered with additional agents which are traditionally used in therapy for these disease states and/or conditions.

Examples

The following examples illustrate and describe the present invention but are not intended to limit the invention in any way.

Synthesis of Compound 3 (Lys01). A round-bottom flask was charged with the 4-bromo-7-chloroquinoline (compound 5) (734 mg, 3.0 mmol), Pd(OAc)₂ (23 mg, 0.1 mmol),

BINAP (125 mg, 0.2 mmol), K_3PO_4 (1.06g, 5.0 mmol), and triamine (compound **6**) (117 mg, 1.0 mmol). Dioxane (10 mL) was introduced through the septum. The resulting suspension was stirred under argon at 90 °C for 18 h and cooled. The mixture was adsorbed onto silica gel and purified by flash chromatography ($CH_2Cl_2/MeOH$: 90/9/1) to afford compound **3** (387 mg, 88%) as a yellow solid. mp 199-200 °C; R_f = 0.28 (silica gel, $CH_2Cl_2/MeOH/NH_4OH$: 90/9/1); ^1H NMR (500 MHz, $CDCl_3$): δ 8.53 (d, J = 5.5 Hz, 2H), 7.94 (d, J = 2.0 Hz, 2H), 7.41 (d, J = 9.0 Hz, 2H), 6.98 (dd, J = 9.0, 2.0 Hz, 2H), 6.39 (d, J = 5.0 Hz, 2H), 5.44 (s, 2H), 3.42 (q, J = 5.0 Hz, 4H), 2.90 (t, J = 6.0 Hz, 2H), 2.46 (s, 9H). ^{13}C NMR (125 MHz, $CDCl_3$): δ 152.1, 149.5, 149.1, 135.1, 128.9, 125.5, 120.6, 117.1, 99.3, 55.5, 42.4, 40.3 FTIR (thin film): 3215, 2917, 1609, 1579, 1449. HRMS-ESI (m/z): calcd for $C_{23}H_{24}N_5Cl_2$ $[M+H]^+$: 440.1409, found: 440.1406.

Synthesis of Compound 11 (Lys 05). To generate a water soluble salt of compound **3**, a suspension of compound **3** (896 mg, 2.04 mmol) in MeOH (40 mL) was bubbled with HCl gas for 10 min at 0 °C. The mixture was stirred for another 12 h at room temperature. The solvent was removed by rotary evaporation and the residue was dried under vacuum at 50 °C overnight to afford the salt **3** (1.13g, 100%) as a yellow solid. mp 270 °C (decomp.); ^1H NMR (500 MHz, D_2O): δ 8.12 (d, J = 7.0 Hz, 2H), 7.73 (d, J = 9.0 Hz, 2H), 7.58 (d, J = 2.0 Hz, 2H), 7.26 (dd, J = 9.0, 2.0 Hz, 2H), 6.62 (d, J = 2.0 Hz, 2H), 3.89 (br, 4H), 3.68 (br, 4H), 3.12 (s, 3H). ^{13}C NMR (125 MHz, D_2O): δ 155.8, 142.8, 140.2, 137.2, 128.1, 123.8, 119.1, 114.8, 98.7, 52.9, 42.7, 38.2. FTIR (thin film): 3376, 3019, 2914, 1631, 1612, 1215 cm^{-1} . HRMS-ESI (m/z): calcd for $C_{23}H_{24}N_5Cl_2$ $[M-3HCl+H]^+$: 440.1409, found: 440.1408.

Biological Testing

Lys01 is a more potent autophagy inhibitor than HCQ or CQ.

LN229 (human glioblastoma) were treated with Lys01 and derivatives Lys02, Lys03, Lys04, HCQ and CQ. Near complete cell death of cultured cells was observed in cells treated with Lys01 at concentrations of 10 μM or higher between 4- 24 hours. LC3 is a ubiquitin-like protein which exists as an unconjugated form (LC3I) or conjugated to AV membranes (LC3II)(25). The ratio of LC3II/LC3I reflects the accumulation of AV in cells, and therefore effective autophagy inhibition. LC3 immunoblotting (Figure 3) demonstrated that Lys01 is a >10-fold more potent autophagy inhibitor than HCQ or CQ at a concentration of 10 μM .

Lys02 and Lys03 produced dose-response relationships for LC3 immunoblotting similar to HCQ or CQ, whereas Lys04, which retains the two chloroquinoline rings present in Lys01, demonstrated intermediate potency in the LC3 autophagy assay

To further characterize the effects of Lys01 on autophagy, LN229 GFP-LC3 cells were treated with Lys01 or HCQ (Figure 4A). Within 4 hours of treatment, in cells treated with HCQ 1 μ M, punctate fluorescence, indicating an accumulation of ineffective autophagic vesicles, was observed in a minority of cells. HCQ 10 μ M produced numerous small puncta, and HCQ 100 μ M resulted in larger dense puncta that represent fusion of accumulated autophagic vesicles. Lys01 1 μ M produced numerous small puncta, whereas in cells treated with Lys01 10 μ M dense puncta similar in appearance to those observed in cells treated with HCQ 100 μ M were apparent. All cells treated with Lys01 100 μ M were dead by 4 hours. Quantification of the GFP-LC3 puncta per cell demonstrated a significant 5-fold increase in GFP-LC3 puncta between 10 μ M Lys01 compared to 10 μ M HCQ treatments. The average number of vesicles per cell in cells treated with Lys01 10 μ M was higher than in cells treated with 100 μ M HCQ (Figure 4A). Electron micrographs of LN229 GFP-LC3 cells treated with DMSO, HCQ, or Lys01 further characterized the significant morphological difference in the size and number of vesicles produced by blockade of autophagy with these agents (Figure 4B). Thus, Lys01 produces morphological changes more pronounced than HCQ, a known lysosomal inhibitor, at 10-fold lower concentrations. To determine if Lys01 treatment was inducing production of new autophagic vesicles (an autophagy inducer) or blocking the clearance of autophagy vesicles (an autophagy inhibitor), a bafilomycin clamp experiment was performed (Figure 4C). LN229 GFP-LC3 cells were treated with DMSO, rapamycin, HCQ 10 μ M, and Lys01 10 μ M, in the absence or presence of bafilomycin. At 24 hours, rapamycin treatment resulted in a further increase in the LC3II/LC3 ratio in bafilomycin treated cells compared to control cells whereas HCQ- or Lys05-treated cells did not demonstrate an increase in LC3II/LC3I ratio in bafilomycin treated cells compared to control, providing further evidence that Lys01 is an autophagy inhibitor (Figure 4C).

To determine the implications of more potent autophagy inhibition on cytotoxicity, LN229 (glioma), 1205Lu (melanoma), HT-29 (colon) and c8161 (melanoma) cells were treated with Lys01, Lys02, Lys03 Lys04, and HCQ at concentrations between 0.01-100 μ M (Figure 4D). The MTT assay was used to assess viable cells at 72 hours. In the 4 cell lines tested, the IC50 of Lys01 was 4-8 μ M (Supplemental Table 2). Near complete cell death after

24 hours was observed in 1205Lu and HCC827 (cell lines which are highly resistant to HCQ) cells treated with 10 μ M Lys01. In contrast the IC₅₀ for Lys02, which is a monofunctional CQ derivative (35-91 μ M), Lys03, the bisaminoquinoline with methoxy groups replacing chlorine (24-53 μ M), or HCQ (15-42 μ M) were collectively 9-30-fold less potent than Lys01. Lys04, which retains the bivalent aminoquinoline rings but has an altered linker, had intermediate activity, with IC₅₀ of 10-17 μ M. These studies demonstrate that Lys01 is consistently more cytotoxic than other aminoquinolines tested or HCQ. Together with the LC3 western blot data, these results indicate that the most potent cytotoxic autophagy inhibitors contain two aminoquinoline rings, the triamine linker present in Lys01 and a chlorine substituent at the C-7 position of the aminoquinoline ring.

In vivo autophagy inhibition and antitumor efficacy Lys05.

Lys05, the trihydrochloride salt of Lys01 was synthesized to enhance aqueous solubility and to enable in vivo studies. Lys01 and Lys05 produced equivalent dose-dependent increases in the LC3II/LC3I ratio, and accumulation of the autophagy cargo protein p62 (26), and identical IC₅₀ values in the MTT assay, (Figure 5A, B). To investigate the safety of Lys05 and its in vivo effects on autophagy, c8161 xenografts matched for tumor size were treated with intraperitoneal (i.p.) daily PBS, or equimolar doses of HCQ or Lys05 (HCQ 60 mg/kg (138 nmoles/g), Lys05 76 mg/kg (138 nmoles/g)) for 48 hours. With this high dose, short term treatment no mice died, but after 2 days of dosing, mice treated with Lys05 76 mg/kg i.p. were observed to have arched backs and lethargy. After 48 hours of treatment mice were euthanized, and tumors were processed for electron microscopy (EM). Morphologically, EM demonstrated that cells with intact nuclear and cytoplasmic membranes contained large AV in Lys05- treated tumors (Figure 6A). Quantification of the mean number of AV/cell in two representative tumor from each treatment group found a significant >2-fold increase in the mean number of AV/cell in Lys05 treated tumors compared to control- or HCQ-treated tumors (Figure 6B). Significantly higher LC3II/LC3I levels were observed in Lys05-treated tumors compared to control- or HCQ –treated tumors providing further evidence of in vivo autophagy inhibition (Figure 7A). After 48 hours of treatment, cleaved caspase 3 levels indicative of apoptosis were elevated in Lys05 treated tumors compared to HCQ- or PBS-treated tumors.

Except for certain models of pancreatic cancer (27), in many animal tumor models, where high levels of autophagy are likely present in untreated tumors(4, 28), treatment with single agent HCQ does not impair tumor growth(29, 30). To determine if a more potent autophagy inhibitor such as Lys05 could significantly impair tumor growth as a single agent 1205Lu xenografts were generated in the flanks of nude mice. For chronic treatment experiments the 1205Lu melanoma model was chosen over the c8161 xenograft model because c8161 xenografts tend to spontaneously ulcerate confounding tumor measurements and safety analysis. Ten mice bearing 1205Lu xenografts were matched for tumor volume per cohort were assigned to either PBS, HCQ 60 mg/kg i.p. or Lys05 76 mg/kg i.p. (equimolar dosing) dosed for 3 days of daily treatment with 2 days off treatment (3/5 days) for all 3 treatment groups, to allow for symptom recovery and to avoid excess toxicity. This schedule was tolerated well for a 14 day period. Tumor growth curves for each of the 3 groups indicated tumor growth was significantly impaired in Lys05 treated tumors compared to controls (Figure 6C). Lys05 treatment resulted in a 53% reduction in the average daily tumor growth rate compared to vehicle treated controls (31.2 v. 14.6 mm³/day; p=0.002; Fig 6D). A significant accumulation of AV was observed at the end of 14 days of treatment in both HCQ-and Lys05-treated tumors, but Lys05 treated tumors had a 6-fold increase in AV/cell whereas HCQ-treated tumors had a 3-fold increase in AV/cell compared to control treated tumors (Figure 7B). Extensive tumor necrosis was observed in the center of Lys05 treated tumors (Figure 7B).

To determine if lower doses of Lys05 could produce antitumor activity, mice bearing HT-29 colon cancer xenografts were treated with PBS, or Lys05 at 10mg/kg i.p. daily, 40 mg/kg i.p. daily, or 80 mg/kg i.p. 3/5days off. Clinical toxicity was observed only in the 80 mg/kg cohort, with 2/8 mice euthanized early for bowel obstruction. Daily dosing for the 10 mg/kg and 40 mg/kg cohorts was well tolerated. The average daily tumor growth rate was significantly impaired in a dose-dependent fashion with Lys05 treatment (Figure 6E). Tumor growth curves demonstrated that all 3 doses of Lys05 produced significant tumor growth impairment compared to control (Figure 6F). At the end of the experiment excised tumor weights demonstrated that significant antitumor activity was observed with 10 mg/kg daily dosing (Figure 6G). Immunoblotting against LC3 in tumor lysates harvested after 14 days of treatment revealed a significant increase in LC3II/LC3I ratio in all Lys05 treated tumors including the 10 mg/kg dosed tumors compared to control (Figure 7C).

Intestinal toxicity at the maximal administered dose of Lys05 resembles a genetic autophagy deficiency.

In the 1205lu xenograft experiment, individual animals treated with Lys05 76 mg/kg i.p. 3/5 days appeared lethargic with arched backs (Figure 8A). Three out of ten mice treated with Lys05 developed signs of bowel obstruction (Figure 8B). Inspection of the bowel found dilated proximal small intestine with a pseudostricture of the terminal ileum. Histological examination of the ileum revealed no evidence of excess inflammation, fibrosis, or mechanical obstruction, indicating that the obstructive signs observed in the mice were due to pseudo-obstruction or functional ileus. While intestinal villi and crypt architecture were intact, dysmorphic Paneth cells (Figure 8C) were observed. Paneth cell dysfunction, including reduced size and number of eosinophilic lysozyme-containing granules, has previously been described as the pathognomonic sign of autophagy deficiency in mice and a subset of Crohn's disease patients that have a genetic deficiency in the essential autophagy gene *ATG16L1* (31).

In the HT29 dose-finding xenograft experiment there was no significant weight loss observed in any dose cohort (Figure 9A). Resection of the entire gastrointestinal tract from mice bearing HT-29 tumors after 14 days of treatment demonstrated bowel thickening and obstruction was limited to 80 mg/kg dose cohort (Figure 9B). Histological examination of the terminal ileum resected from mice bearing HT-29 xenografts treated with PBS ip daily, or Lys05 10 mg/kg ip daily, 40 mg/kg ip daily, and 80 mg/kg ip daily every 3/5 days for 14 days demonstrated dose dependent effects on Paneth cell morphology (Figure 9C). While the number of Paneth cells/crypt did not change with treatment (Figure 9D), the size and number of granules decreased in a dose dependent manner. Scoring on a Paneth cell dysfunction scale (Figure 10) indicated that Paneth cell dysfunction was observed at all doses tested of Lys05, despite signs and symptoms of toxicity being restricted to the 80 mg/kg dose (Figure 9E). In mice treated with Lys05 40 mg/kg or 80 mg/kg, but not 10 mg/kg, lysozyme was significantly reduced or absent in Paneth cells (Figure 9F). Taken together these findings indicate that Lys05-associated Paneth cell dysfunction mimics *ATG16L1* deficiency, and lower doses of Lys05 produce significant antitumor activity without dose-limiting toxicity.

Lys05 inhibits autophagy by deacidifying the lysosome.

To compare the relative lysosomal accumulation of Lys05 compared to HCQ, lysosomes were subfractionated from 1205Lu cells treated with PBS, HCQ 10 μ M, or Lys05 10 μ M, and 1205Lu tumors harvested after 14 days of treatment with PBS, HCQ 60 mg/kg i.p., or Lys05 76 mg./kg i.p. every 3/5 days. Immunoblotting against the lysosomal marker LAMP2 confirmed adequate separation of the lysosomal and whole cell population in both cells and tumor samples (Figure 12A). HPLC tandem mass spectrometry (MS/MS) measurements (Figure 11) determined that the concentrations of Lys05 and HCQ in the whole cell homogenate treated with Lys05 10 μ M or HCQ 10 μ M for 24 hours were 57 μ M and 8 μ M respectively, indicating an 6-fold higher concentration of Lys05 within the cell compared to HCQ. The concentration of Lys05 and HCQ in the lysosomal fraction of cells treated with Lys05 10 μ M or HCQ 10 μ M were 105 μ M and 13 μ M respectively, indicating an 8-fold higher concentration of Lys05 in the lysosome compared to HCQ. This difference in cellular and lysosomal accumulation of Lys05 and HCQ was more marked in tumor tissue. There was an 11-fold higher concentration and a 34-fold higher concentration of Lys05 compared to HCQ in whole cell homogenates and lysosomes, respectively, within tumors (Figure 12B).

Having established that Lys05 more effectively accumulates in the lysosome than HCQ, the functional effects of this accumulation were investigated. 1205Lu cells were treated with vehicle, Lys05 and HCQ, and stained the LysoTracker Red (Figure 12C). Within 30 minutes of treatment, fewer LysoTracker-positive puncta were observed in Lys05 treated cells at both 10 μ M and 100 μ M concentrations. In contrast, a significant decrease in LysoTracker positive puncta was observed in cells treated with HCQ 100 μ M, but not observed in cells treated with HCQ 10 μ M. To understand the implications of this more potent and complete lysosomal inhibition, 1205Lu cells were treated with vehicle, Lys05, or HCQ and stained with acridine orange (AO; a dye which aggregates in all endovesicular acidic compartments) at 24 hours (Figure 12D). HCQ produced a dose-dependent accumulation of acidic vesicles. In contrast, Lys05 caused an accumulation of acidic vesicles at lower doses (10 μ M), but at higher doses (50 μ M), no acidic vesicles were observed, indicating the complete deacidification of the endovesicular system.

Finally, the functional consequences of lysosomal deacidification were investigated by measuring enzymatic activity of acid phosphatase. In 1205Lu cells treated with PBS, HCQ 10 μ M or Lys05 10 μ M, within 24 hours there is a 43% reduction in acid phosphatase activity in the lysosomal fraction of Lys05 treated versus PBS treated cells (Figure 13A). Leakage of certain lysosomal enzymes such as activated cathepsins, could lead to an autophagy-independent cell death. Within 24 hours of treatment of 1205Lu cells with PBS, HCQ or Lys05, there is decreased acid-dependent processing of immature cathepsin D to the mature activated form within the lysosome in Lys05-treated compared to HCQ- or PBS-treated cells (Figure 13B). In 1205Lu xenograft tumors, after 14 days of treatment there was a 1.75-fold increase in extralysosomal acid phosphatase activity in the Lys05-treated tumors, suggesting that chronic treatment can lead to extralysosomal leakage of enzymes (Figure 13C). but increased acid-dependent processing of cathepsin D within the whole cell homogenate was not observed in Lys05 treated tumors (Figure 13D). These results indicate that high doses of Lys05 cause lysosomal dysfunction by deacidifying the lysosome, leading to impairment of lysosomal enzymes, and effective autophagy inhibition, whereas high doses of HCQ incompletely deacidify the lysosome, leading to incomplete autophagy inhibition associated with less cell death.

Additional compounds Lys06-Lys18 Additional Lys01 derivatives have been synthesized and tested (Figure 14, Figure 15). In 72 hour MTT assays the IC₅₀'s of compounds Lys01-Lys13 demonstrate increased or decreased activity compared to HCQ and CQ (Table 1, figure 16). In most cases the Lys01 derivatives are more active than CQ or HCQ. These findings further refine the starting point for further drug development of Lys01 derivatives.

Activity of Lys01 derviatives in malaria. Table 2, Figure 17 shows the IC₅₀ values for Lys01 derivative- induced cell death in the human cancer cell LN229 and a number of strains of *P. Falciparum* grown in invitro in human RBC. There is a similar activity profile for anticancer activity of Lys01 dereivatives and malaria cytotoxicity. Lys01 was more active than artesunate in some CQ-resistant cell lines.

Conclusions

Potential Commercial Uses and Applications: Lys01 and Lys05 are lead compounds with great potential to be optimized further for potency as a novel autophagy inhibitor. Autophagy inhibition is a new therapeutic strategy in cancer that is applicable to every cancer. There are currently over 30 HCQ trials in cancer patients involving nearly every tumor type. Due to its low potency and poor pharmacology, in humans HCQ will likely not yield the promising augmentation of anticancer therapy observed in laboratory models. An optimized derivative of Lys01 could be developed as a second generation autophagy inhibitor. The GI toxicity associated with Paneth cell dysfunction observed at LD30 doses of Lys05, support the mechanism of action of the drug, and also suggests that colon cancers, which often share features with Paneth cells, may be a tumor type that may be especially sensitive to Lys05 and its optimized derivatives. Additional cancers worth investigating include melanoma, and non small cell lung cancer, since melanoma cell lines demonstrated the highest difference in sensitivity to Lys01 compared to HCQ, and an EGFR mutated lung cancer cell line demonstrated sensitivity to both HCQ and Lys05. The synthesis of Lys01 was designed such that there is no overlap with other patented and/or published aminoquinoline compounds. Further mechanistic studies are planned that are to identify pharmacodynamics assays that guide drug development. Pharmacokinetic studies planned in mice establish initial in vivo profile.

Other similar technologies and competing products: Novel chloroquine derivatives for use as anticancer agents is an active area of investigation (16). Autophagy has been identified as one of the top ten areas of research in which the NIH will invest in the next few years. No studies to date have leveraged the potential of bivalency as the inventors provide here. In addition, most studies lack the in vivo studies and the mechanistic studies herein reported that can guide further development of optimize lead compounds for drug development.

Advantages over other similar technologies and products: Thus, the present application has shows that the disclosed series of bisaminoquinolines are potent autophagy inhibitors that have single agent antitumor activity in an in vivo tumor model.

References


1. Lum JJ, DeBerardinis RJ, Thompson CB. Autophagy in metazoans: cell survival in the land of plenty. *Nat Rev Mol Cell Biol* 2005;6: 439-48.
2. Amaravadi RK, Thompson CB. The roles of therapy-induced autophagy and necrosis in cancer treatment. *Clin Cancer Res* 2007;13: 7271-9.
3. Amaravadi RK, Yu D, Lum JJ, *et al.* Autophagy inhibition enhances therapy-induced apoptosis in a Myc-induced model of lymphoma. *J Clin Invest* 2007;117: 326-36.
4. Degenhardt K, Mathew R, Beaudoin B, *et al.* Autophagy promotes tumor cell survival and restricts necrosis, inflammation, and tumorigenesis. *Cancer Cell* 2006;10: 51-64.
5. Amaravadi RK. Autophagy-induced tumor dormancy in ovarian cancer. *J Clin Invest* 2008.
6. Carew JS, Nawrocki ST, Kahue CN, *et al.* Targeting autophagy augments the anticancer activity of the histone deacetylase inhibitor SAHA to overcome Bcr- Abl-mediated drug resistance. *Blood* 2007.
7. Degtyarev M, De Maziere A, Orr C, *et al.* Akt inhibition promotes autophagy and sensitizes PTEN-null tumors to lysosomotropic agents. *J Cell Biol* 2008;183: 101-16.
8. Sotelo J, Briceno E, Lopez-Gonzalez MA. Adding chloroquine to conventional treatment for glioblastoma multiforme: a randomized, double-blind, placebo-controlled trial. *Ann Intern Med* 2006;144: 337-43.
9. Amaravadi RK, Lippincott-Schwartz J, Yin XM, *et al.* Principles and Current Strategies for Targeting Autophagy for Cancer Treatment. *Clin Cancer Res* 2011;17: 654-66.
10. Rosenfeld MR GS, Brem S, Mikkelsen T, Wang D, Piao S, Davis L, O'Dwyer PJ, Amaravadi RK Pharmacokinetic analysis and pharmacodynamic evidence of autophagy inhibition in patients with newly diagnosed glioblastoma treated on a phase I trial of hydroxychloroquine in combination with adjuvant temozolomide and radiation (ABTC 0603). *J Clin Oncol* 2010;28: Abstract # 3086.
11. Vance D, Shah M, Joshi A, Kane RS. Polyvalency: a promising strategy for drug design. *Biotechnol Bioeng* 2008;101: 429-34.
12. Shrivastava A, Nunn AD, Tweedle MF. Designer peptides: learning from nature. *Curr Pharm Des* 2009;15: 675-81.

13. Girault S, Grellier P, Berecibar A, *et al.* Antiplasmodial activity and cytotoxicity of bis-, tris-, and tetraquinolines with linear or cyclic amino linkers. *J Med Chem* 2001;44: 1658-65.
14. Vennerstrom JL, Ager AL, Jr., Dorn A, *et al.* Bisquinolines. 2. Antimalarial N,N-bis(7-chloroquinolin-4-yl)heteroalkanediamines. *J Med Chem* 1998;41: 4360-4.
15. Burnett JC, Schmidt JJ, Stafford RG, *et al.* Novel small molecule inhibitors of botulinum neurotoxin A metalloprotease activity. *Biochem Biophys Res Commun* 2003;310: 84-93.
16. Hu C, Raja Solomon V, Cano P, Lee H. A 4-aminoquinoline derivative that markedly sensitizes tumor cell killing by Akt inhibitors with a minimum cytotoxicity to non-cancer cells. *Eur J Med Chem* 2010;45: 705-9.
17. Solomon VR, Hu C, Lee H. Design and synthesis of chloroquine analogs with anti-breast cancer property. *Eur J Med Chem* 2010;45: 3916-23.
18. T. Higuchi HO, M. Umezawa, H. Kim, Y. Wataya,. Compound with Antimalarial Activity and Antimalarial Drug Containing the Same as Active Ingredient. WO/2007/097450.
19. Lee WW, Berridge BJ, Jr., Ross LO, Goodman L. Synthesis of Mustards from Putrescine, Cadaverine, and 1,3-Diaminopropane. *J Med Chem* 1963;6: 567-9.
20. Kaschula CH, Egan TJ, Hunter R, *et al.* Structure-activity relationships in 4-aminoquinoline antiplasmodials. The role of the group at the 7-position. *J Med Chem* 2002;45: 3531-9.
21. Adams A, Jarrott B, Elmes BC, Denny WA, Wakelin LP. Interaction of DNA-intercalating antitumor agents with adrenoceptors. *Mol Pharmacol* 1985;27: 480-91.
22. Gourdie TA, Valu KK, Gravatt GL, *et al.* DNA-directed alkylating agents. 1. Structure-activity relationships for acridine-linked aniline mustards: consequences of varying the reactivity of the mustard. *J Med Chem* 1990;33: 1177-86.
23. Bailey DM, DeGrazia CG, Hoff SJ, *et al.* Bispyridinamines: a new class of topical antimicrobial agents as inhibitors of dental plaque. *J Med Chem* 1984;27: 1457-64.
24. Vicker N, Burgess L, Chuckowree IS, *et al.* Novel angular benzophenazines: dual topoisomerase I and topoisomerase II inhibitors as potential anticancer agents. *J Med Chem* 2002;45: 721-39.
25. Tanida I, Ueno T, Kominami E. LC3 conjugation system in mammalian autophagy. *Int J Biochem Cell Biol* 2004;36: 2503-18.

26. Pankiv S, Clausen TH, Lamark T, *et al.* p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of ubiquitinated protein aggregates by autophagy. *J Biol Chem* 2007;282: 24131-45.
27. Yang S, Wang X, Contino G, *et al.* Pancreatic cancers require autophagy for tumor growth. *Genes Dev* 2011;25: 717-29.
28. Ma X, Piao S, Wang DW, *et al.* Measurements of tumor cell autophagy predict invasiveness, resistance to chemotherapy, and survival in melanoma. *Clin Cancer Res* 2011.
29. Fan QW, Cheng C, Hackett C, *et al.* Akt and autophagy cooperate to promote survival of drug-resistant glioma. *Sci Signal* 2010;3: ra81.
30. Saleem A, Dvorzhinski D, Santanam U, *et al.* Effect of dual inhibition of apoptosis and autophagy in prostate cancer. *Prostate* 2012.
31. Cadwell K, Liu JY, Brown SL, *et al.* A key role for autophagy and the autophagy gene Atg16l1 in mouse and human intestinal Paneth cells. *Nature* 2008;456: 259-63.

Claims:

1. A compound according to the chemical structure I



Wherein R^1 and $R^{1'}$ are each independently H, halo (F, Cl, Br or I), CN, NO₂, optionally substituted C₁-C₆ alkyl (when substituted, preferably substituted with 1 or 2 hydroxyl groups or 3-5 fluoro groups), optionally substituted O-C₁-C₆ alkyl (preferably, OCH₃), optionally substituted C₂-C₇ acyl (preferably acetyl) or optionally substituted C₂-C₇ ester (oxycarbonyl ester or carboxyester, preferably carboxyester);

R and R' are each independently H, a C₁-C₆ optionally substituted alkyl group, a C₁-C₇ (preferably C₂-C₇) optionally substituted acyl group, a C₂-C₇ optionally substituted carboxy ester group (which forms a urethane group with the nitrogen atom to which R or R' is bonded);

L is a $-(CH_2Y)_n-X-(Y'CH_2)_n-$ group or a $A-(CH_2-CH_2-Z)_n-A'$ group (either A or A' may be bonded to either of the two amine groups in compound I) wherein at least one of the CH₂ groups in L is optionally substituted with a C₁-C₃ alkyl group which itself is optionally substituted with one or two hydroxyl groups;

X is absent, (CH₂)_j, O, S or N-R'';

Y is absent, CH₂, O, CH₂O or N-R'' and Y' is absent, CH₂, O, OCH₂ or N-R'', with the proviso that when one or more of X, Y and Y' is present, each of X and Y, X and Y' or Y and Y', when present, forms a stable bond;

R'' is H or an optionally substituted C₁-C₆ (preferably C₁-C₃) alkyl group;

j is 1, 2 or 3 (preferably 1 or 2);

n is 0, 1, 2, 3 or 4, with the proviso that when n is 0, X is (CH₂)_j where j is at least 1 and at least one CH₂ group is optionally substituted with a C₁-C₃ alkyl group which itself is optionally substituted with one or two hydroxyl groups;

A is a bond (absent) or (CH₂)_j and A' is a bond (absent) or (CH₂)_j wherein at least one CH₂ group in A or A' is optionally substituted with a C₁-C₃ alkyl group which is itself optionally substituted with one or two hydroxyl groups;

Z is O or N-R^Z;

R^Z is H or an optionally substituted C₁-C₃ alkyl group,

or a pharmaceutically acceptable salt, enantiomer, diastereomer, solvate or polymorph thereof.

2. The compound according to claim 1, wherein R¹ and R^{1'} are each independently H, a halo group, a nitro group or a trifluoromethyl group.

3. The compound according to claim 1 or 2 wherein R and R' are each independently H, a C₁-C₃ alkyl group substituted with at least one hydroxyl group, alkoxy group, an amine, monoalkyl amine or dialkyl amine group, wherein said amine group or said monoalkyl amine group is optionally substituted on the amine position with a 7-substituted-4-quinolinyl group wherein the amine binds to the 4-position of the quinolinyl group and the 7 position of said quinolinyl group is substituted by R¹ or R^{1'} of claim 1, or one or both alkyl groups of said monoalkyl amine or dialkyl amine is itself further optionally substituted with at least one hydroxyl group, alkoxy group, an amine, a monoalkyl amine, a dialkyl amine wherein the amine or monoalkyl amine is optionally substituted on the amine position with one or two 7-substituted-quinolinyl group(s) wherein the amine binds to the 4-position of the quinolinyl group and the 7 position of said quinolinyl group is substituted by R¹ or R^{1'} of claim 1, and each of said alkoxy groups may be further substituted with an alkoxy group, preferably a methoxy group (thus forming a diether substituent).

4. The compound according to any of claims 1-3 wherein L is a

—(CH₂Y)_n—X—(Y'CH₂)_n— group, where X is N-R^{''}; n is 1, 2, or 3; Y and Y' are each independently absent or CH₂; and R^{''} is H or a C₁-C₃ alkyl group which is optionally substituted with at least one hydroxyl group, an alkoxy group, an amine, monoalkyl amine, dialkyl amine group, wherein said amine group or said monoalkyl amine group is optionally

substituted on the amine position with a 7-substituted-4-quinolinyl group wherein the amine binds to the 4-position of the quinolinyl group and the 7 position of said quinolinyl group is substituted by R¹ or R1' of claim 1, or one or both alkyl groups of said monoalkyl amine or dialkyl amine is itself further optionally substituted with at least one hydroxyl group, alkoxy group, an amine, a monoalkyl amine or a dialkyl amine wherein the amine or monoalkyl amine is optionally substituted on the amine position with one or two 7-substituted-quinolinyl group(s) wherein the amine binds to the 4-position of the quinolinyl group and the 7 position of said quinolinyl group is substituted by R¹ or R1' of claim 1, and each of said alkoxy group may be further substituted with a second alkoxy group, thus forming a diether substituent.

5. A compound according to claim 3 or 4 wherein said C₁-C₃ alkyl group is substituted with an amine group which itself is substituted with one 7-substituted-4-quinolinyl group.
6. The compound according to claim 5 wherein said alkyl group is a C₂ alkyl group.
7. The compound according to claim 3 or 4 wherein said C₁-C₃ alkyl group is substituted with an alkoxy group which itself is substituted with a second alkoxy group thus forming a diether group.
8. The compound according to claim 7 wherein said second alkoxy group is a methoxy group or an ethoxy group.
9. A compound as depicted in Scheme 1 or 3-10 hereof.
10. A compound of claim 9 which is any one of compounds 28-40 of scheme 3 hereof.
11. A compound of claim 9 which is any one of compounds 41-46 of scheme 4 hereof.
12. A compound of claim 9 which is either of compounds 47-48 of scheme 5 hereof.
13. A compound of claim 9 which is either of compounds 49-50 of scheme 6 hereof.
14. A compound of claim 9 which is any one of compounds 51-58 of scheme 7 hereof.

15. A compound of claim 9 which is any one of compounds 63-70 of scheme 8 hereof.
16. A compound of claim 9 which is any one of compounds 71-78 of scheme 9 hereof.
17. A compound of claim 9 which is any one of compounds 79-82 of scheme 10 hereof.
18. A compound according to claim 9 which is compound 3 of scheme 1 hereof.
19. A compound as depicted in Figures 14, 15 or 15A hereof.

20. The compound:

N^1 -(7-chloroquinolin-4-yl)- N^2 -(2-((7-chloroquinolin-4-yl)amino)ethyl)- N^2 -methylethane-1,2-diamine;

N^1 -(7-chloroquinolin-4-yl)- N^2 -(2-((7-chloroquinolin-4-yl)amino)ethyl)ethane-1,2-diamine;

N,N' -((ethane-1,2-diylbis(oxy))bis(ethane-2,1-diyl))bis(7-chloroquinolin-4-amine);

N^1 -(7-methoxyquinolin-4-yl)- N^2 -(2-((7-methoxyquinolin-4-yl)amino)ethyl)- N^2 -methylethane-1,2-diamine;

N,N' -((ethane-1,2-diylbis(oxy))bis(ethane-2,1-diyl))bis(7-chloroquinolin-4-amine);

N^1 -(7-chloroquinolin-4-yl)- N^2 -(2-((7-chloroquinolin-4-yl)amino)ethyl)- N^2 -methylethane-1,2-diamine trihydrochloride;

N^1 -(7-chloroquinolin-4-yl)- N^2 -(2-((7-chloroquinolin-4-yl)(methyl)amino)ethyl)- N^1,N^2 -dimethylethane-1,2-diamine;

N,N' -((methylazanediyl)bis(ethane-2,1-diyl))bis(N -(7-chloroquinolin-4-yl)acetamide);

(*S*)-*N*²-(7-chloroquinolin-4-yl)-*N*¹-((*S*)-2-((7-chloroquinolin-4-yl)amino)propyl)-*N*¹-methylpropane-1,2-diamine;

2-(bis(2-((7-chloroquinolin-4-yl)amino)ethyl)amino)ethanol;

*N*¹-(7-chloroquinolin-4-yl)-*N*²,*N*²-bis(2-((7-chloroquinolin-4-yl)amino)ethyl)ethane-1,2-diamine, or

a pharmaceutically acceptable salt thereof.

21. The compound according to claim 1 which is *N*¹-(7-chloroquinolin-4-yl)-*N*²-(2-((7-chloroquinolin-4-yl)amino)ethyl)-*N*²-methylethane-1,2-diamine or a pharmaceutically acceptable salt thereof.

22. The compound according to claim 1 which is *N*¹-(7-chloroquinolin-4-yl)-*N*²-(2-((7-chloroquinolin-4-yl)amino)ethyl)-*N*²-methylethane-1,2-diamine trihydrochloride.

23. A pharmaceutical composition comprising an effective amount of at least one compound according to any of claims 1-22 in combination with a pharmaceutically acceptable carrier, additive or excipient and optionally in combination with at least one additional anticancer agent.

24. A method of inhibiting autophagy in a biological system in which inhibition of autophagy is desired, said method comprising exposing said biological system to an effective amount of at least one compound according to any of claims 1-22 hereof.

25. A method of inhibiting or treating cancer in a patient in need comprising administering to said patient an effective amount of at least one compound according to any of claims 1-22, optionally in combination with at least one additional anticancer agent.

26. The method according to claim 25 wherein said cancer is metastatic.

27. The method according to claim 26 wherein said cancer is a drug-resistant cancer.

28. A method of reducing the likelihood that cancer will occur in a patient or that a cancer will metastasize in a patient comprising administering at least one compound according to any of claims 1-22, optionally in combination with at least one additional anticancer agent.

29. The method according to claim 25 wherein said cancer is a carcinoma, cancer of the esophagus, head, kidney, liver, lung, nasopharyngeal, neck, ovary, pancreas, prostate, and stomach; a leukemia, a malignant lymphoma, a malignant melanoma; myeloproliferative diseases; a sarcoma, a tumor of the central nervous system, a germ-line tumor, lung cancer, ovarian cancer, testicular cancer, thyroid cancer, astrocytoma, esophageal cancer, pancreatic cancer, stomach cancer, liver cancer, colon cancer, melanoma, a mixed type of neoplasia,

30. The method according to claim 28 wherein said leukemia is acute myelogenous leukemia, acute lymphocytic leukemia, acute promyelocytic leukemia (APL), acute T-cell lymphoblastic leukemia, adult T-cell leukemia, basophilic leukemia, eosinophilic leukemia, granulocytic leukemia, hairy cell leukemia, leukopenic leukemia, lymphatic leukemia, lymphoblastic leukemia, lymphocytic leukemia, megakaryocytic leukemia, micromyeloblastic leukemia, monocytic leukemia, neutrophilic leukemia and stem cell leukemia

31. The method according to claim 28 wherein said lymphoma is Burkitt's lymphoma, Non-Hodgkin's lymphoma or B-cell lymphoma.

32. The method according to claim 28 wherein said sarcoma is Ewing's sarcoma, hemangiosarcoma, Kaposi's sarcoma, liposarcoma, myosarcomas, peripheral neuroepithelioma or synovial sarcoma.

33. The method according to claim 28 wherein said tumor of the central nervous system is a glioma, astrocytoma, oligodendroglioma, ependymoma, glioblastoma, neuroblastoma, ganglioneuroma, ganglioglioma, medulloblastoma, pineal cell tumor, meningioma, meningeal sarcoma, neurofibroma, or Schwannoma.

34. The method according to claim 28 wherein said germ-line tumor is bowel cancer, breast cancer, prostate cancer, cervical cancer or uterine cancer.

35. The method according to claim 28 wherein said lung cancer is small cell lung cancer, mixed small cell and non-small cell cancer, pleural mesothelioma, metastatic pleural mesothelioma, small cell lung cancer or non-small cell lung cancer.
36. The method according to claim 28 wherein said mixed neoplasia is carcinosarcoma and Hodgkin's disease and said tumors of mixed origin is Wilms' tumor and teratocarcinomas.
37. The method according to claim 28 wherein said cancer is ovarian, breast, colon, head and neck, medulloblastoma or B-cell lymphoma,
38. The method according to claim 28 wherein said cancer is melanoma or non-small cell lung cancer.
39. The method according to any of claims 25-38 wherein said additional anticancer agent is selected from the group consisting of everolimus, trabectedin, abraxane, TLK 286, AV-299, DN-101, pazopanib, GSK690693, RTA 744, ON 0910.Na, AZD 6244 (ARRY-142886), AMN-107, TKI-258, GSK461364, AZD 1152, enzastaurin, vandetanib, ARQ-197, MK-0457, MLN8054, PHA-739358, R-763, AT-9263, pemetrexed, erlotinib, dasatinib, nilotinib, decatanib, panitumumab, amrubicin, oregovomab, Lep-etu, nolatrexed, azd2171, batabulin, ofatumumab, zanolimumab, edotecarin, tetrandrine, rubitecan, tesmilifene, oblimersen, ticilimumab, ipilimumab, gossypol, Bio 111, 131-I-TM-601, ALT-110, BIO 140, CC 8490, cilengitide, gimatecan, IL13-PE38QQR, INO 1001, IPdR₁ KRX-0402, lucanthone, LY 317615, neuradiab, vitespan, Rta 744, Sdx 102, talampanel, atrasentan, Xr 311, romidepsin, ADS- 100380, sunitinib, 5-fluorouracil, vorinostat, etoposide, gemcitabine, doxorubicin, irinotecan, liposomal doxorubicin, 5'-deoxy-5-fluorouridine, vincristine, temozolomide, ZK-304709, seliciclib; PD0325901, AZD-6244, capecitabine, L-Glutamic acid, N-[4-[2-(2-amino-4,7-dihydro-4-oxo-1 H - pyrrolo[2,3- d]pyrimidin-5-yl)ethyl]benzoyl]-, disodium salt, heptahydrate, camptothecin, PEG-labeled irinotecan, tamoxifen, toremifene citrate, anastrozole, exemestane, letrozole, DES(diethylstilbestrol), estradiol, estrogen, conjugated estrogen, bevacizumab, IMC-1C11, CHIR-258,); 3-[5-(methylsulfonylpiperadinemethyl)-indolyl]-quinolone, vatalanib, AG-013736, AVE-0005, the acetate salt of [D- Ser(Bu t) 6 , Azgly 10] (pyro-Glu-His-Trp-Ser-Tyr-D-Ser(Bu t)-Leu-Arg-Pro- Azgly-NH₂ acetate [C₅₉H₈₄N₁₈O₁₄ -(C₂H₄O₂)_x where x = 1 to 2.4], goserelin acetate, leuprolide acetate, triptorelin pamoate, medroxyprogesterone acetate, hydroxyprogesterone caproate, megestrol

acetate, raloxifene, bicalutamide, flutamide, nilutamide, megestrol acetate, CP-724714; TAK-165, HKI-272, erlotinib, lapatanib, canertinib, ABX-EGF antibody, erbitux, EKB-569, PKI-166, GW-572016, Ionafernib, BMS-214662, tipifarnib; amifostine, NVP-LAQ824, suberoyl analide hydroxamic acid, valproic acid, trichostatin A, FK-228, SU11248, sorafenib, KRN951, aminoglutethimide, arnsacrine, anagrelide, L-asparaginase, Bacillus Calmette-Guerin (BCG) vaccine, bleomycin, buserelin, busulfan, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, diethylstilbestrol, epirubicin, fludarabine, fludrocortisone, fluoxymesterone, flutamide, gemcitabine, gleevac, hydroxyurea, idarubicin, ifosfamide, imatinib, leuprolide, levamisole, lomustine, mechlorethamine, melphalan, 6-mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, octreotide, oxaliplatin, pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, streptozocin, teniposide, testosterone, thalidomide, thioguanine, thiotepa, tretinoin, vindesine, 13-cis-retinoic acid, phenylalanine mustard, uracil mustard, estramustine, altretamine, floxuridine, 5-deoxyuridine, cytosine arabinoside, 6-mecaptopurine, deoxycoformycin, calcitriol, valrubicin, mithramycin, vinblastine, vinorelbine, topotecan, razoxin, marimastat, COL-3, neovastat, BMS-275291, squalamine, endostatin, SU5416, SU6668, EMD121974, interleukin-12, IM862, angiostatin, vitaxin, droloxifene, idoxyfene, spironolactone, finasteride, cimitidine, trastuzumab, denileukin diftitox, gefitinib, bortezomib, paclitaxel, irinotecan, topotecan, doxorubicin, docetaxel, vinorelbine, bevacizumab (monoclonal antibody) and erbitux, cremophor-free paclitaxel, epithilone B, BMS-247550, BMS-310705, droloxifene, 4-hydroxytamoxifen, pipendoxifene, ERA-923, arzoxifene, fulvestrant, acolbifene, lasofoxifene, idoxifene, TSE-424, HMR-3339, ZK186619, PTK787/ZK222584, VX-745, PD184352, rapamycin, 40-O-(2-hydroxyethyl)-rapamycin, temsirolimus, AP-23573, RAD001, ABT-578, BC-210, LY294002, LY292223, LY292696, LY293684, LY293646, wortmannin, ZM336372, L-779,450, PEG-filgrastim, darbepoetin, erythropoietin, granulocyte colony-stimulating factor, zolendronate, prednisone, cetuximab, granulocyte macrophage colony-stimulating factor, histrelin, pegylated interferon alfa-2a, interferon alfa-2a, pegylated interferon alfa-2b, interferon alfa-2b, azacitidine, PEG-L-asparaginase, lenalidomide, gemtuzumab, hydrocortisone, interleukin-11, dexrazoxane, alemtuzumab, all-transretinoic acid, ketoconazole, interleukin-2, megestrol, immune globulin, nitrogen mustard, methylprednisolone, ibritumomab tiuxetan, androgens, decitabine, hexamethylmelamine, bexarotene, tositumomab, arsenic trioxide, cortisone, editronate, mitotane, cyclosporine, liposomal daunorubicin, Edwina-asparaginase, strontium 89,

casopitant, netupitant, an NK-1 receptor antagonists, palonosetron, aprepitant, ,
diphenhydramine, hydroxyzine, metoclopramide, lorazepam, alprazolam, haloperidol,
droperidol, dronabinol, dexamethasone, methylprednisolone, prochlorperazine, granisetron,
ondansetron, dolasetron, tropisetron, sspegfilgrastim, erythropoietin, epoetin alfa and
darbepoetin alfa, ipilimumab, vemurafenib and mixtures thereof.

40. The method according to any of claims 25-38 wherein said additional anticancer agent is
a FLT-3 inhibitor, a VEGFR inhibitor, an EGFR TK inhibitor, an aurora kinase inhibitor, a
PIK-1 modulator, a Bcl-2 inhibitor, an HDAC inhibitor, a c-MET inhibitor, a PARP inhibitor,
a Cdk inhibitor, an EGFR TK inhibitor, an IGFR-TK inhibitor, an anti-HGF antibody, a PI3
kinase inhibitors, an AKT inhibitor, a JAK/STAT inhibitor, a checkpoint-1 or 2 inhibitor, a
focal adhesion kinase inhibitor, a Map kinase kinase (mek) inhibitor, a VEGF trap antibody
or a mixture thereof.

41. A method of treating a disease state or condition in a patient in need wherein said disease
state or condition responds favorably to inhibition of autophagy comprising administering to
said patient an effective amount of a compound according to any of claims 1-22 to said
patient.

42. The method according to claim 41 wherein said disease state or condition is rheumatoid
arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria or Sjogren's
disease.

43. The method according to claim 42 wherein said disease state is malaria.

44. Use of a compound according to any of claims 1-22 in the manufacture of a medicament
for the inhibition of autophagy in a biological system in which inhibition of autophagy is
desired.

45. Use of a compound according to any of claims 1-22 in the manufacture of a medicament
for use in inhibiting or treating cancer in a patient in need, optionally in combination with at
least one additional anticancer agent.

46. The use according to claim 45 wherein said cancer is metastatic.

47. The use according to claim 45 wherein said cancer is drug resistant.
48. Use of a compound according to any of claims 1-22 in the manufacture of a medicament for reducing the likelihood that cancer will occur in a patient or that a cancer will metastasize in a patient, optionally in combination with at least one additional anticancer agent.
49. Use according to any of claims 45-48 wherein said cancer is a carcinoma, cancer of the esophagus, head, kidney, liver, lung, nasopharyngeal, neck, ovary, pancreas, prostate, and stomach; a leukemia, a malignant lymphoma, a malignant melanoma; myeloproliferative diseases; a sarcoma, a tumor of the central nervous system, a germ-line tumor, lung cancer, ovarian cancer, testicular cancer, thyroid cancer, astrocytoma, esophageal cancer, pancreatic cancer, stomach cancer, liver cancer, colon cancer, melanoma, a mixed type of neoplasia,
50. Use according to claim 49 wherein said leukemia is acute myelogenous leukemia, acute lymphocytic leukemia, acute promyelocytic leukemia (APL), acute T-cell lymphoblastic leukemia, adult T-cell leukemia, basophilic leukemia, eosinophilic leukemia, granulocytic leukemia, hairy cell leukemia, leukopenic leukemia, lymphatic leukemia, lymphoblastic leukemia, lymphocytic leukemia, megakaryocytic leukemia, micromyeloblastic leukemia, monocytic leukemia, neutrophilic leukemia and stem cell leukemia
51. Use according to claim 49 wherein said lymphoma is Burkitt's lymphoma, Non-Hodgkin's lymphoma or B-cell lymphoma.
52. Use according to claim 49 wherein said sarcoma is Ewing's sarcoma, hemangiosarcoma, Kaposi's sarcoma, liposarcoma, myosarcomas, peripheral neuroepithelioma or synovial sarcoma.
53. Use according to claim 49 wherein said tumor of the central nervous system is a glioma, astrocytoma, oligodendroglioma, ependymoma, glioblastoma, neuroblastoma, ganglioneuroma, ganglioglioma, medulloblastoma, pineal cell tumor, meningioma, meningeal sarcoma, neurofibroma, or Schwannoma.

54. Use according to claim 49 wherein said germ-line tumor is bowel cancer, breast cancer, prostate cancer, cervical cancer or uterine cancer.

55. Use according to claim 49 wherein said lung cancer is small cell lung cancer, mixed small cell and non-small cell cancer, pleural mesothelioma, metastatic pleural mesothelioma, small cell lung cancer or non-small cell lung cancer.

56. Use according to claim 49 wherein said mixed neoplasia is carcinosarcoma and Hodgkin's disease and said tumors of mixed origin is Wilms' tumor and teratocarcinomas.

57. Use according to claim 49 wherein said cancer is ovarian, breast, colon, head and neck, medulloblastoma or B-cell lymphoma,

58. Use according to claim 49 wherein said cancer is melanoma or non-small cell lung cancer.

59. Use according to any of claims 45-59 wherein said additional anticancer agent is selected from the group consisting of everolimus, trabectedin, abraxane, TLK 286, AV-299, DN-101, pazopanib, GSK690693, RTA 744, ON 0910.Na, AZD 6244 (ARRY-142886), AMN-107, TKI-258, GSK461364, AZD 1152, enzastaurin, vandetanib, ARQ-197, MK-0457, MLN8054, PHA-739358, R-763, AT-9263, pemetrexed, erlotinib, dasatanib, nilotinib, decatanib, panitumumab, amrubicin, oregovomab, Lep-etu, nolatrexed, azd2171, batabulin, ofatumumab, zanolimumab, edotecarin, tetrandrine, rubitecan, tesmilifene, oblimersen, ticilimumab, ipilimumab, gossypol, Bio 111, 131-I-TM-601, ALT-110, BIO 140, CC 8490, cilengitide, gimatecan, IL13-PE38QQR, INO 1001, IPdR₁ KRX-0402, lucanthone, LY 317615, neuradiab, vitespan, Rta 744, Sdx 102, talampanel, atrasentan, Xr 311, romidepsin, ADS- 100380, sunitinib, 5-fluorouracil, vorinostat, etoposide, gemcitabine, doxorubicin, irinotecan, liposomal doxorubicin, 5'-deoxy-5-fluorouridine, vincristine, temozolomide, ZK-304709, seliciclib; PD0325901, AZD-6244, capecitabine, L-Glutamic acid, N-[4-[2-(2-amino-4,7-dihydro-4-oxo-1 H - pyrrolo[2,3- d]pyrimidin-5-yl)ethyl]benzoyl]-, disodium salt, heptahydrate, camptothecin, PEG-labeled irinotecan, tamoxifen, toremifene citrate, anastrozole, exemestane, letrozole, DES(diethylstilbestrol), estradiol, estrogen, conjugated estrogen, bevacizumab, IMC-1C11, CHIR-258,); 3-[5-(methylsulfonylpiperadinemethyl)-indolyl]-quinolone, vatalanib, AG-013736, AVE-0005, the acetate salt of [D- Ser(Bu t) 6

, Azgly 10] (pyro-Glu-His-Trp-Ser-Tyr-D-Ser(Bu t)-Leu-Arg-Pro- Azgly-NH₂ acetate [C₅₉H₈₄N₁₈O₁₄-(C₂H₄O₂)_x where x = 1 to 2.4], goserelin acetate, leuprolide acetate, triptorelin pamoate, medroxyprogesterone acetate, hydroxyprogesterone caproate, megestrol acetate, raloxifene, bicalutamide, flutamide, nilutamide, megestrol acetate, CP-724714; TAK-165, HKI-272, erlotinib, lapatanib, canertinib, ABX-EGF antibody, erbitux, EKB-569, PKI-166, GW-572016, Ionafernib, BMS-214662, tipifarnib; amifostine, NVP-LAQ824, suberoyl analide hydroxamic acid, valproic acid, trichostatin A, FK-228, SU11248, sorafenib, KRN951, aminoglutethimide, arnsacrine, anagrelide, L-asparaginase, Bacillus Calmette-Guerin (BCG) vaccine, bleomycin, buserelin, busulfan, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, diethylstilbestrol, epirubicin, fludarabine, fludrocortisone, fluoxymesterone, flutamide, gemcitabine, gleevac, hydroxyurea, idarubicin, ifosfamide, imatinib, leuprolide, levamisole, lomustine, mechlorethamine, melphalan, 6-mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, octreotide, oxaliplatin, pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, streptozocin, teniposide, testosterone, thalidomide, thioguanine, thiotepa, tretinoin, vindesine, 13-cis-retinoic acid, phenylalanine mustard, uracil mustard, estramustine, altretamine, floxuridine, 5-deoxyuridine, cytosine arabinoside, 6-mecaptopurine, deoxycorformycin, calcitriol, valrubicin, mithramycin, vinblastine, vinorelbine, topotecan, razoxin, marimastat, COL-3, neovastat, BMS-275291, squalamine, endostatin, SU5416, SU6668, EMD121974, interleukin-12, IM862, angiostatin, vitaxin, droloxifene, idoxyfene, spironolactone, finasteride, cimitidine, trastuzumab, denileukin diftitox, gefitinib, bortezimib, paclitaxel, irinotecan, topotecan, doxorubicin, docetaxel, vinorelbine, bevacizumab (monoclonal antibody) and erbitux, cremophor-free paclitaxel, epithilone B, BMS- 247550, BMS-310705, droloxifene, 4-hydroxytamoxifen, pibendoxifene, ERA- 923, arzoxifene, fulvestrant, acolbifene, lasofoxifene, idoxifene, TSE-424, HMR- 3339, ZK186619, PTK787/ZK 222584, VX-745, PD 184352, rapamycin, 40-O-(2-hydroxyethyl)-rapamycin, temsirolimus, AP-23573, RAD001, ABT-578, BC-210, LY294002, LY292223, LY292696, LY293684, LY293646, wortmannin, ZM336372, L-779,450, PEG-filgrastim, darbepoetin, erythropoietin, granulocyte colony-stimulating factor, zolendronate, prednisone, cetuximab, granulocyte macrophage colony-stimulating factor, histrelin, pegylated interferon alfa-2a, interferon alfa-2a, pegylated interferon alfa-2b, interferon alfa-2b, azacitidine, PEG-L-asparaginase, lenalidomide, gemtuzumab, hydrocortisone, interleukin-11, dexrazoxane, alemtuzumab, all-transretinoic acid, ketoconazole, interleukin-2, megestrol, immune globulin, nitrogen

mustard, methylprednisolone, ibritumomab tiuxetan, androgens, decitabine, hexamethylmelamine, bexarotene, tositumomab, arsenic trioxide, cortisone, editronate, mitotane, cyclosporine, liposomal daunorubicin, Edwina-asparaginase, strontium 89, casopitant, netupitant, an NK-1 receptor antagonists, palonosetron, aprepitant, , diphenhydramine, hydroxyzine, metoclopramide, lorazepam, alprazolam, haloperidol, droperidol, dronabinol, dexamethasone, methylprednisolone, prochlorperazine, granisetron, ondansetron, dolasetron, tropisetron, sspegfilgrastim, erythropoietin, epoetin alfa and darbepoetin alfa, ipilimumab, vemurafenib and mixtures thereof.

60. Use according to any of claims 45-58 wherein said additional anticancer agent is a FLT-3 inhibitor, a VEGFR inhibitor, an EGFR TK inhibitor, an aurora kinase inhibitor, a PIK-1 modulator, a Bcl-2 inhibitor, an HDAC inhibitor, a c-MET inhibitor, a PARP inhibitor, a Cdk inhibitor, an EGFR-TK inhibitor, an IGFR-TK inhibitor, an anti-HGF antibody, a PI3 kinase inhibitors, an AKT inhibitor, a JAK/STAT inhibitor, a checkpoint-1 or 2 inhibitor, a focal adhesion kinase inhibitor, a Map kinase kinase (mek) inhibitor, a VEGF trap antibody or a mixture thereof.

61. Use of a compound according to any of claims 1-22 in the manufacture of a medicament for treating a disease state or condition in a patient in need wherein said disease state or condition responds favorably to inhibition of autophagy..

62. Use according to claim 61 wherein said disease state or condition is rheumatoid arthritis, malaria, antiphospholipid antibody syndrome, lupus, chronic urticaria or Sjogren's disease.

63. Use according to claim 62 wherein said disease state or condition is malaria.

Application number / Numéro de demande: 2841452


Figures: _____

Pages: 4, 5, 6, 9 — 16, 19-21,


Unscannable items received with this application
(Request original documents in File Prep. Section on the 10th floor)

Documents reçu avec cette demande ne pouvant être balayés
(Commander les documents originaux dans la section de la préparation
des dossiers au 10ième étage)

1/26

**Figure 1**

2/26

Figure 2

3/26




Figure 3

7/26




Figure 4D




Figure 5

17/26






Figure 11

18/26

A**B****Figure 12A-B**




FIGURE 14

23/26




FIGURE 15




FIGURE 15A

Table 1. 72 hr MTT IC50 in LN229 cells	
Compound	IC50 (M)
Lyso12: Trifunctional	3.19E-06
Lyso1: Bifunctional	4.60E-06
Lyso11: Short ether	5.29E-06
Lyso5: Bifunctional HCL	5.57E-06
Lyso8: Trifluoro	5.60E-06
Lyso13: Central NH	6.02E-06
Lyso9: Nitro	7.08E-06
Lyso4: Long ether	8.56E-06
Lyso7: Fluoro	1.12E-05
Lyso6: Dechloro	1.19E-05
Lyso3: Methoxy	1.67E-05
CQ	2.25E-05
Lyso2: Mono + linker	2.97E-05
HCC	3.08E-05
Lyso10: Methylketone	1.08E-04

Figure 16

Table 2. IC50 (M) of Lys01 derivatives in P. Falciparum						
	CQ- sensitive	CQ-sensitive	CQ-resistant	CQ-resistant	CQ-resistant	CQ-resistant
Name	3D7	HB3W	Dd2	7G8	K1	
Lys012: Trifunctional	1.09E-07	1.04E-07	4.19E-08	3.87E-08	4.65E-08	
Lys01: Bifunctional	3.74E-09		1.57E-08	8.10E-09	6.02E-09	
Lys011: Short ether	6.32E-09	6.44E-09	1.14E-08	3.74E-08	2.12E-08	
Lys05: Bifunctional HCL	4.00E-09		1.11E-08	5.38E-09	4.38E-09	
Lys08: Trifluoro	2.35E-08	2.42E-08	1.93E-08	1.51E-08	2.53E-08	
Lys013: Central NH						
Lys09: Nitro	4.10E-08	3.05E-08	4.70E-08	5.26E-08	4.30E-08	
Lys04: Long ether	4.39E-09		1.07E-08	8.12E-09	8.66E-09	
Lys07: Fluoro	1.26E-08	1.24E-08	1.33E-08	3.97E-08	1.42E-08	
Lys06: Dechloro	8.58E-08	8.48E-08	3.60E-07	8.63E-07	1.22E-06	
Lys03: Methoxy	2.45E-07		4.10E-07	5.18E-07	6.11E-07	
CQ	4.60E-09	4.50E-09	2.05E-08	3.23E-08	4.09E-08	
Lys02: Mono + linker	2.84E-08		1.28E-07	3.07E-07	1.48E-07	
Artesunate	2.21E-08	1.73E-08	1.26E-08	7.19E-08	9.35E-08	

Figure 17