



(86) Date de dépôt PCT/PCT Filing Date: 2004/08/24
(87) Date publication PCT/PCT Publication Date: 2005/03/17
(45) Date de délivrance/Issue Date: 2009/10/20
(85) Entrée phase nationale/National Entry: 2006/02/28
(86) N° demande PCT/PCT Application No.: IB 2004/002744
(87) N° publication PCT/PCT Publication No.: 2005/023780
(30) Priorité/Priority: 2003/09/05 (US60/500,733)

(51) Cl.Int./Int.Cl. *C07D 239/42* (2006.01)

(72) Inventeurs/Inventors:
KATH, JOHN CHARLES, US;
RICHTER, DANIEL TYLER, US;
LUZZIO, MICHAEL JOSEPH, US


(73) Propriétaire/Owner:
PFIZER PRODUCTS INC., US

(74) Agent: SMART & BIGGAR

(54) Titre : SYNTHESE SELECTIVE DE PYRIMIDINES SUBSTITUEES EN CF3
(54) Title: SELECTIVE SYNTHESIS OF CF3-SUBSTITUTED PYRIMIDINES

(57) **Abrégé/Abstract:**

The present invention relates to a method of making a compound of the formula (I), wherein X^1 , X^2 and R^3 - R^4 are as defined herein. The method includes reacting a compound of the formula (II) with an amine of formula (III) (HNR^3R^4) in the presence of a Lewis Acid and a non-nucleophilic base. The 2,4-diamino pyrimidine moiety is a common component in a variety of biologically active drug-like molecules and pyrimidine derivatives have been found to be useful in the treatment of abnormal cell growth, such as cancer, in mammals.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
17 March 2005 (17.03.2005)

PCT

(10) International Publication Number
WO 2005/023780 A1

(51) International Patent Classification⁷: **C07D 239/42**

(74) Agent: **FULLER, Grover, F., Jr.**; Pfizer Inc., 201 Tabor Road, Morris Plains, NJ 07950 (US).

(21) International Application Number:
PCT/IB2004/002744

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 24 August 2004 (24.08.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/500,733 5 September 2003 (05.09.2003) US

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **PFIZER PRODUCTS INC.** [US/US]; Eastern Point Road, Groton, CT 06340 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **KATH, John, Charles** [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **RICHTER, Daniel, Tyler** [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **LUZZIO, Michael, Joseph** [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US).

Published:


— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.


(54) Title: SELECTIVE SYNTHESIS OF CF₃-SUBSTITUTED PYRIMIDINES

(57) Abstract: The present invention relates to a method of making a compound of the formula (I), wherein X¹, X² and R³-R⁴ are as defined herein. The method includes reacting a compound of the formula (II) *with an amine of formula (III) (HNR³R⁴) in the presence of a Lewis Acid and a non-nucleophilic base*. The 2,4-diamino pyrimidine moiety is a common component in a variety of biologically active drug-like molecules and pyrimidine derivatives have been found to be useful in the 10 treatment of abnormal cell growth, such as cancer, in mammals.

WO 2005/023780 A1

SELECTIVE SYNTHESIS OF CF₃-SUBSTITUTED PYRIMIDINESBackground of the Invention


The 2,4-diamino pyrimidine moiety (**1**) is a common component in a variety of biologically active drug-like molecules. Pyrimidine derivatives have been found to be useful in the treatment of abnormal cell growth, such as cancer, in mammals. These moieties are commonly synthesized starting with pyrimidine intermediate **2** (where "X" is a leaving group; most commonly a halogen) and an equivalent of amine **3**, HNR³R⁴ (see Scheme 1 below). For the vast majority of reactions involving pyrimidines of formula **2** and amines of formula **3**, it is well known that this first amine addition occurs preferentially (or exclusively) at the more reactive pyrimidine 4-position (*Chemistry of Heterocyclic Compounds, The Pyrimidines*, Volume 52, Wiley, New York 1994, p.371.) to provide intermediate **4**. The primary factors that influence the selectivity of this initial amine addition are the stereoelectronic effects associated with substituents present in both pyrimidine **2** and amine **3** and to a lesser extent the reaction solvent. Subsequent heating of **4** with a second amine (**5**) provides the desired 2,4-diaminopyrimidine **1**.

SCHEME 1

A representative example of the aforementioned chemistry can be found in WO00391901 and is highlighted in Scheme 2. Other examples utilizing this general synthetic scheme include Montebugnoli et. al. *Tetrahedron* 2002, (58), p. 2147. *Chemistry of Heterocyclic Compounds, The Pyrimidines*, Volume 52, Wiley, New York 1994, pp. 371-417. Selective amine addition to 2,4-dichloro-5-carboxamidopyrimidines is described in WO 02/04429. Selective amine additions to 2,4-dichloro-5-halopyrimidines are described in WO 01/65655.


-2-

SCHEME 2



While there are a number of general examples where specific pyrimidines (2), amines (3) or reaction conditions provide non-selective mixtures of the 2-chloro-4-amino-pyrimidine (4) and the isomeric 2-amino-4-chloro-pyrimidine (6) (Scheme 3), these reactions are of limited utility not only due to their lack of selectivity (and its impact on overall yield) but also because separation of the resulting isomers is generally extremely difficult. Preparative HPLC is generally required as a means to individually isolate the pure isomers (4 and 6), which can then be transformed further into compounds such as 1 or its isomer 7 respectively.

SCHEME 3




An example of such a reaction that provides mixtures of isomers is the addition of 4-methyl-aniline to 2,4-dichloro-5-trifluoromethylpyrimidine (Scheme 4). This electron deficient pyrimidine has a slight preference for amine addition to the pyrimidine 2-position. HPLC analysis of the crude reaction mixture shows a 1.4 to 1 mixture of (4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-p-tolyl-amine (8) and (2-Chloro-5-trifluoromethyl-pyrimidin-4-yl)-p-tolyl-amine (9). Other examples of non-selective amine addition to 2,4 di-halogenopyrimidines are described in *Chemistry of Heterocyclic Compounds, The Pyrimidines*, Volume 52, Wiley, New York 1994, pp. 371-417. Luo et. al. *Tetrahedron Lett.* 2002, (43) p. 5739. Yoshida et. al. *J. Chem. Soc, Perkin Trans. I: Organic and Bioorganic*

-3-


Chemistry, 1992 (7) p. 919. EP 647639 describes additions of piperidines to 2,4-dichloropyrimidine.

SCHEME 4




5 In contrast to the reactions described above, there are only a few very specific examples where an amine (3) is added to a pyrimidine of formula 2 in a selective manner to provide preferentially the 2-amino-4-chloro-pyrimidine 6. The most notable example of this type of reaction is the addition of N-methyl piperidine to 2,4-dichloro-5-methyl pyrimidine to provide 4-chloro-5-methyl-2-piperidinopyrimidine (Scheme 5) found in Yoshida et. al. *J.*
 10 *Chem. Soc, Perkin Trans. I: Organic and Bioorganic Chemistry*, 1992 (7) p.919. In this case, the steric effect of the 5-methyl substituent on the pyrimidine coupled with the fact that the amine nucleophile is a tertiary (rather than a primary or secondary) amine provides for selective addition of piperidine to the pyrimidine 2-position.

SCHEME 5

Summary of the Invention

15 It has been surprisingly found that by adding a Lewis acid to the reaction medium, one can selectively add an amine functionality to the C-2 position of a CF₃ substituted-pyrimidine ring. Thus, the present invention relates to a process of making a compound of the
 20 formula 11




wherein X² is a leaving group such as a halide, arylsulfonate, alkylsulfonate, perfluoroalkylsulfonate, arylsulfinates or alkylsulfinates; and

-4-

R^3 and R^4 are substituents independently selected from the group consisting of hydrogen, an aromatic group and an aliphatic group; or taken together $-NR^3R^4$ can form a 4-11 membered aromatic or aliphatic ring;

wherein the process comprises reacting a compound of formula **10**



5

with an amine of formula **3** (HNR^3R^4) in the presence of a Lewis Acid and a non-nucleophilic base to form a compound of formula **11**, wherein X^1 is a leaving group such as a halide, arylsulfonate, alkylsulfonate, perfluoroalkylsulfonate, arylsulfinate or alkylsulfinate.

10 In a preferred embodiment, amine **3** is an aromatic amine and 0.25-10 equivalents of Lewis Acid are used relative to the amount of pyrimidine **10**, and preferably 0.5-3.0 equivalents of Lewis Acid are used relative to pyrimidine **10**.

In an alternative preferred embodiment, amine **3** is an aliphatic amine and 0.5-10 equivalents of Lewis Acid are used relative to the amount of pyrimidine **10**, and preferably 1-4 equivalents of Lewis Acid are used relative to pyrimidine **10**.

15 In a further preferred embodiment, X^1 and X^2 are the same or different and are each independently halides and the Lewis Acid is a salt of zinc or magnesium.

In the most preferred embodiment, X^1 and X^2 are chloride and the Lewis Acid is zinc chloride

20 The compounds prepared by the method of the present invention include all stereoisomers (e.g., cis and trans isomers) and all optical isomers of compounds of the formula **11** (e.g., R and S enantiomers), as well as racemic, diastereomeric and other mixtures of such isomers.

25 The compounds and salts prepared by the method of the present invention can exist in several tautomeric forms, including the enol and imine form, and the keto and enamine form and geometric isomers and mixtures thereof. The preparation of all such tautomeric forms is included within the scope of the present invention. Tautomers exist as mixtures of a tautomeric set in solution. In solid form, usually one tautomer predominates. Even though the preparation of one tautomer may be described, the present invention encompasses the preparation of all tautomers of the present compounds.

30 The present invention also includes the preparation of atropisomers of the present invention. Atropisomers refer to compounds of formula **11** that can be separated into rotationally restricted isomers.

35 The compounds prepared by the method of the invention may contain olefin-like double bonds. When such bonds are present, the compounds exist as cis and trans configurations and as mixtures thereof and the present invention contemplates the preparation of such compounds.

As used herein the term "aromatic", and specifically, an "aromatic group" refers to an aryl or heteroaryl radical as defined herein.

Further, an "aromatic amine" or "aromatic amine radical" refers to any amine or amine radical bound to at least one sp^2 carbon atom that is part of an aryl or heteroaryl group. An
5 amine or amine radical will be referred to as an aromatic amine or radical even if the amine nitrogen is bound to a hydrogen or an sp^3 carbon atom, in addition to the one sp^2 carbon atom. Thus, for example, $-HN(C_6-C_{10})\text{aryl}$ and $-N((C_1-C_6)\text{alkyl})((C_6-C_{10})\text{aryl})$ each refer to aromatic amine radicals as defined herein, despite the fact that each amine nitrogen is attached to non-aromatic substituents.

10 The term "aryl" refers to aromatic radicals such as phenyl, naphthyl, tetrahydronaphthyl, indanyl and the like. An "aryl" group may be optionally substituted with 1-3 suitable substituents, as defined herein. "Aryl" also refers to a phenyl radical fused to a non-aromatic heterocycle. Examples of such groups include but are not limited to 2-oxo-indoliny, chromanyl, indoliny and 2-oxo-3,4-dihydroquinoliny optionally substituted by 1 to 3
15 suitable substituents.

As used herein, the term "heteroaryl" refers to an aromatic heterocyclic group usually with one heteroatom selected from O, S and N in the ring, wherein the aromatic heterocyclic group may be substituted by up to three suitable substituents as defined herein. In addition to
20 in the ring. Examples of heteroaryl groups include but are not limited to pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, thienyl, furyl, imidazolyl, pyrrolyl, oxazolyl (e.g., 1,3-oxazolyl, 1,2-oxazolyl), thiazolyl (e.g., 1,2-thiazolyl, 1,3-thiazolyl), pyrazolyl, tetrazolyl, triazolyl (e.g., 1,2,3-triazolyl, 1,2,4-triazolyl), oxadiazolyl (e.g., 1,2,3-oxadiazolyl), thiadiazolyl (e.g., 1,3,4-thiadiazolyl), quinolyl, isoquinolyl, benzothienyl, benzofuryl, indolyl, and the like; optionally
25 substituted by 1 to 3 suitable substituents. Alternatively, any ring carbon, $-CH-$, of the aforementioned heteroaryl group, may be replaced by a group selected from $-C=O$ or $-SO_2$. "Heteroaryl" also refers to one of the aforementioned heteroaryl groups fused to a non-aromatic heterocycle. Examples of such groups include but are not limited to 1,3-Dihydro-pyrrolo[2,3-b]pyridin-2-one, 3,4-Dihydro-1H-[1,8]naphthyridin-2-one, 1,3-Dihydro-pyrrolo[2,3-b]pyridine and
30 3,4-Dihydro-2H-pyrano[2,3-b]pyridine.

"Aliphatic group" refers to an alkyl, cycloalkyl, or heterocycloalkyl radical, as defined herein. Aliphatic groups may be substituted with up to three suitable substituents as defined herein.

As used herein, the term "aliphatic amine" or "aliphatic amino radical" refers to any
35 amine or amine radical in which the amine or radical nitrogen atom is bound to an sp^3 carbon that is part of an alkyl, cycloalkyl, or heterocycloalkyl group. Aliphatic amine groups may be substituted with up to three suitable substituents as defined herein.

The term "alkyl" refers to C₁-C₁₀ linear or branched alkyl groups (such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl etc.) optionally substituted by 1 to 3 suitable substituents as defined herein.

The term "cycloalkyl" or "cyclyl" refers to C₃-C₁₂ mono, bicyclic or tricyclic carbocyclic ring (e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclopentenyl, cyclohexenyl, bicyclo[2.2.1]heptanyl, bicyclo[3.2.1]octanyl and bicyclo[5.2.0]nonanyl, etc.) that is optionally substituted by 1 to 3 suitable substituents as defined herein. Bicyclic or tricyclic species may be fused, bridged or spirocyclic. Thus, examples of "cycloalkyl" or "cyclyl" groups, as defined herein, include, but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclohexenyl, bicyclo[2.2.1]heptanyl, bicyclo[3.2.1]octanyl, bicyclo[3.1.0]hexyl and spiro[2.4]heptyl.

The term "heterocycloalkyl" or "heterocyclyl" or "heterocycle" refers to a mono, bicyclic or tricyclic group containing 3 to 9 carbon atoms and 1 to 4 heteroatoms selected from -N-, -NR-, -O-, -S-, -SO or -SO₂, wherein the cyclic radical is optionally substituted by 1 to 3 suitable substituents as defined herein. Bicyclic or tricyclic species may be fused, bridged or spirocyclic. Examples of such groups include but are not limited to azetidiny, pyrrolidiny, piperidiny, morpholinyl, piperazinyl, tetrahydrofuranyl, oxetanyl, thiomorpholinyl, quinuclidiny, 5-aza-spiro[2.4]heptyl and 3-aza-bicyclo[3.1.0]hexyl.

As used herein, the term "halogen" includes fluoro, chloro, bromo or iodo or fluoride, chloride, bromide or iodide.

As used herein, the term "carbonyl" or "(C=O)" (as used in phrases such as alkylcarbonyl, alkyl-(C=O)- or alkoxy carbonyl) refers to the joinder of the >C=O moiety to a second moiety such as an alkyl or amino group (i.e. an amido group).

When -NR³R⁴ or -NR⁵R⁶ taken together form a cyclic amine, said amine can be a mono, bicyclic or tricyclic ring comprising 3 to 9 carbon atoms and 0 to 3 heteroatoms selected from -N-, -O-, -S-, -SO or -SO₂ (excluding the nitrogen atom in -NR³R⁴ or -NR⁵R⁶). The cyclic amine may be optionally substituted with 1 to 3 suitable substituents as defined herein. Bicyclic or tricyclic species may be fused bridged or spirocyclic. Examples of such cyclic amines include but are not limited to morpholine, azetidine, piperazine, piperidine, pyrrolidine, indoline, thiomorpholine.


A "suitable substituent" is intended to mean a chemically and pharmaceutically acceptable functional group. Such suitable substituents for the aforementioned aryl, heteroaryl, alkyl, cycloalkyl, heterocycloalkyl groups may be routinely described by those skilled in the art. Illustrative examples of said suitable substituents include, but are not limited to hydrogen, halo groups, perfluoroalkyl groups, perfluoroalkoxy groups, alkyl groups, alkenyl groups, alkynyl groups, hydroxy groups, oxo groups, alkylthio groups, arylthio groups, alkylsulfonyl groups, arylsulfonyl groups, heteroarylsulfonyl groups, alkylsulfonate groups,

-7-

arylsulfonate groups, perfluoroalkylsulfonate groups, alkoxy groups, aryl or heteroaryl groups, cycloalkyl or heterocycloalkyl groups, aryloxy or heteroaryloxy groups, aralkyl or heteroaralkyl groups, aralkoxy or heteroaralkoxy groups, HO-(C=O)- groups, amino groups, alkyl- and dialkylamino groups, carbamoyl groups, alkylcarbonyl groups, alkoxy carbonyl groups, alkylaminocarbonyl groups, dialkylamino carbonyl groups, sulfonamido groups, alkylsulfonamido groups, dialkylsulfonamido groups, amido groups, N-acyl groups, arylcarbonyl groups, aryloxy carbonyl groups and the like. Methylene groups may also be substituted for a carbonyl (C=O) group. Those skilled in the art will appreciate that many substituents can be substituted by additional substituents.

“Embodiment” as used herein refers to specific groupings of compounds or uses into discrete subgenera. Such subgenera may be cognizable according to one particular substituent such as a specific R³ or R⁴ group. Other subgenera are cognizable according to combinations of various substituents, such as all compounds wherein R³ is hydrogen and R⁴ is (C₁-C₆)alkyl, optionally substituted by -(C₃-C₁₀)cyclyl.


Detailed Description of the Invention



X=leaving group

Compounds of the formula **11** can be prepared by reaction of pyrimidine **10** and a primary or secondary amine nucleophile (**3**) in the presence of a Lewis acid and a non-nucleophilic base in an organic solvent or mixture of solvents. Leaving groups (X¹, X², which may be the same or different) on pyrimidine **10** suitable for displacement, include but are not limited to halides, sulfonates and sulfinates. Preferentially, each leaving group is a halide. In a further preferred embodiment, the halide is chloride. Suitable primary and secondary amine nucleophiles include aromatic, aliphatic or cyclic amines as described above. Lewis acids include but are not limited to the salts of Zn, Mg, Sn, Ti, Al, B, Li, Ag, Na, K, Ca, V, Cr, Mn, Fe, Co, Ni, In, Zr, Sm and Cu. In a preferred embodiment, salts of Zn, Mg, Sn, Ti, Al, B, Li, Ag, and Cu are preferred. Most preferably, a salt of Zn or Mg is used as the Lewis acid. Suitable equivalents of Lewis Acid relative to pyrimidine **10** necessary to effect selective amine addition range from 0.25-10 equivalents, and preferably 0.5-3 equivalents when the amine nucleophile is aromatic or 0.5-10 equivalents, and preferably 1-4 equivalents when the amine is aliphatic (or an aromatic amine capable of forming a multidentate ligand with the Lewis Acid). Suitable organic solvents include but are not limited tetrahydrofuran, 1,2-dichloroethane, *t*-butanol, ether, methylene chloride, acetonitrile, methanol, ethanol, 2-propanol, dioxane, 1,2-dimethoxyethane, toluene, chloroform, ethyl acetate or mixtures

Compounds of the formula **11**, when combined with a primary or secondary amine HNR^5R^6 (**5**) are useful in the preparation of compounds of formula **12**, where R^5 and R^6 independently represent hydrogen, an aromatic group or an aliphatic group, or NR^5R^6 taken together can form a 4-11 membered aromatic or aliphatic ring. Alternatively, compounds of formula **11** are useful in the preparation of compounds of formula **13a** or **13b** by addition of an oxygen or sulfur nucleophile, respectively. Compounds of formula **11** (where X is a halide or perfluoroalkylsulfonate) are also useful in preparing compounds of formula **14** or **15a/b**, wherein R^7 is an aromatic or an aliphatic group, via palladium catalyzed carbon-carbon bond formation.



EXAMPLES

25 The following examples illustrate the preparation of the compounds of the present invention. NMR data are reported in parts per million and are referenced to the deuterium lock signal from the sample solvent. Commercial reagents were utilized without further purification. THF refers to tetrahydrofuran and DMF refers to N,N-dimethylformamide. Chromatography refers to column chromatography performed using .040 mm silica gel and
30 executed under flash chromatography conditions. Low Resolution Mass Spectra (LRMS) were recorded on a Fisons Atmospheric Pressure Chemical Ionization platform, which uses a 50/50 mixture of acetonitrile/water with 0.1% formic acid as the ionizing agent. All non-

50054-130

-9-

aqueous reactions were run under a nitrogen atmosphere for convenience and to maximize yields. Concentration under reduced pressure means that a rotary evaporator was used. Aromatic or aliphatic amine nucleophiles were either purchased and used without further purification or prepared by standard methods of amine synthesis known to those skilled in the art.

Where analytical HPLC chromatography is referred to in the preparations and examples below, the general conditions used, unless otherwise indicated, are as follows. The column used is a ZORBAX^{*} RXC18 column (manufactured by Hewlett Packard) of 150 mm distance and 4.6 mm interior diameter. The samples are run on a Hewlett Packard-1100 system. A gradient solvent method is used running 100 percent ammonium acetate / acetic acid buffer (0.2 M) to 100 percent acetonitrile over 10 minutes. The system then proceeds on a wash cycle with 100 percent acetonitrile for 1.5 minutes and then 100 percent buffer solution for 3 minutes. The flow rate over this period is a constant 3 mL/ minute.

Example 1

Preparation of 2,4-dichloro-5-trifluoromethylpyrimidine: 5-Trifluoromethyluracil

(250g, 1.39 mol) and phosphorous oxychloride (655 mL, 6.94 mol, 5 equiv) were charged to a 3L 4-neck flask equipped with overhead stirrer, a reflux condenser, an addition funnel and an internal thermocouple. The contents were maintained under a nitrogen atmosphere as concentrated phosphoric acid (85 wt%, 9.5 mL, 0.1 equiv) was added in one portion to the slurry, resulting in a moderate exotherm. Diisopropylethylamine (245 mL, 1.39 mol, 1 equiv) was then added dropwise over 15 min at such a rate that the internal temperature of the reaction reached 85-90 °C by the end of the addition. By the end of the amine addition the reaction mixture was a homogenous light-orange solution. Heating was initiated and the orange solution was maintained at 100 °C for 20h, at which time HPLC analysis of the reaction mixture indicated that the starting material was consumed. External heating was removed and the contents of the flask were cooled to 40 °C and then added dropwise to a cooled mixture of 3N HCl (5 L, 10 equiv) and diethyl ether (2L) keeping the temperature of the quench pot between 10 and 15 °C. The layers were separated, and the aqueous layer was extracted once with ether (1L). The combined organic layers were combined, washed with water until the washes were neutral (5 x 1.5L washes), dried with MgSO₄ and concentrated to provide 288g (95% yield) of a light yellow-orange oil of 96% purity (HPLC). This material can be further purified by distillation (bp 109 °C at 79 mmHg).

Example 2:

General procedure for non-selective amine addition:

Method A: A mixture of (2-Chloro-5-trifluoromethyl-pyrimidin-4-yl)-p-tolyl-amine (9) and (4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-p-tolyl-amine (8) To a solution of 5-trifluoromethyl-2,4-dichloropyrimidine (500 mg; 2.3 mmol) in DCE/t-butanol (20 mL) was

*Trade-mark

-10-

added 4-methylaniline (247 mg; 1 eq) followed by dropwise addition triethylamine (1.1 eq). After stirring overnight the reaction was concentrated and taken up in ethyl acetate, washed with sat. NaHCO_3 , dried over Na_2SO_4 , and the solvent removed. Analytical HPLC analysis revealed that the crude reaction contained a 1.4:1 mixture of 8:9. The resulting mixture of isomers was separated on a Shimadzu^{*} preparative HPLC system using a standard gradient (Waters XTerra^{*} Prep MS C_{18} column 5m, 50 x 50 mm; 0.1% NH_4OH in 40-90% $\text{ACN}/\text{H}_2\text{O}$ 75 mL/min, 15 min gradient elution) to provide; (2-Chloro-5-trifluoromethyl-pyrimidin-4-yl)-p-tolyl-amine (9) (122 mg): ^1H NMR (CDCl_3 , 400 MHz) δ 2.35 (s, 3H), 6.99 (br s, 1H), 7.19 (d, J = 8.3 Hz, 2H), 7.38 (d, J = 8.3 Hz, 2H), 8.38 (s, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 163.9, 157.6, 156.0 (q, J = 5 Hz), 136.1, 133.7, 130.0, 123.8 (q, J = 270 Hz), 122.8, 106.6 (q, J = 32 Hz), 21.2; HPLC ret. Time: 7.236 min. LRMS (M^+): 288.1, 290.1. and ((4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-p-tolyl-amine (8) (205 mg): ^1H NMR (CDCl_3 , 400 MHz) 2.33 (s, 3H), 7.17 (d, J = 8.3 Hz, 2H), 7.42 (d, J = 8.3 Hz, 2H), 7.46 (br s, 1H), 8.52 (s, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) 160.9, 157.6 (br), 134.9, 134.8, 129.9, 122.0 (q, J = 269 Hz), 121.1, 113.7 (q, J = 34 Hz), 21.1; HPLC ret. Time: 8.137 min. LRMS (M^+): 288.1, 290.1. Confirmation of the structures of the two isomers was obtained by single crystal X-ray analysis.

General procedures for selective addition of amines using a Lewis Acid:

Method B: (4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-p-tolyl-amine (8). To a solution of 5-trifluoromethyl-2,4-dichloropyrimidine (2 g; 9.2 mmol) in 1:1 DCE/*t*-BuOH (80 mL) was added Zinc chloride (11 mL of a 1M solution in ether; 1.2 eq) at 0° C. After 1 hour, 4-methylaniline (988 mg; 1 eq) was added followed by dropwise addition of a solution of triethylamine (1.03 g; 1.1 eq) in 10mL of DCE/*t*-BuOH. After stirring for 1.5 hours the reaction was concentrated. Analytical HPLC analysis revealed that the crude reaction contained <5% of isomer 9. The desired product 8 was obtained as a white solid (2.25g; 85%) following crystallization from methanol. HPLC ret. Time: 8.169 min. LRMS (M^+): 288.2, 290.1.

The application of Method B with a variety of other amines allows for the preparation of a variety of other 2-amino-4-chloro-5-trifluoromethylpyrimidines including, but not limited to those outlined below.

5-(4-Chloro-5-trifluoromethyl-pyrimidin-2-ylamino)-1,3-dihydro-indol-2-one: ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ 3.29 (s, 2H), 6.76 (d, J = 7.9 Hz, 2H), 7.39 (d, J = 8.3 Hz), 7.51 (br s, 1H), 8.71 (s, 1H), 10.33 (s, 1H), 10.49 (s, 1H). ^{13}C NMR ($\text{DMSO}-d_6$, 100 MHz) δ 177.0, 161.3, 158.7 (br), 140.7, 132.8, 126.9, 123.7 (q, J = 268 Hz), 121.0, 118.7, 111.2 (q, J = 32 Hz), 109.6, 36.7; HPLC ret. time: 5.759 min. LRMS (M^+) 329.1, 331.1.

(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-(4-methoxy-phenyl)-amine : ^1H NMR (CDCl_3 , 400 MHz) δ 3.80 (s, 3H), 6.91 (d, J = 9.1 Hz, 2H), 7.38 (br s, 1H), 7.43 (d, J = 8.7 Hz, 2H), 8.50 (s, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 161.1, 157.6 (br), 157.2, 130.3, 123.2, 122.9

*Trade-mark

(q, J = 269 Hz), 114.5, 113.5 (q, J = 34 Hz), 112.5, 55.7; HPLC ret. time: 7.550 min. LRMS (M+) 304.2, 306.1.

(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-(4-nitro-phenyl)-amine: ¹H NMR (CDCl₃, 400 MHz) δ 7.80 (br s, 1H), 7.82 (d, J = 24 Hz, 2H), 8.26 (d, J = 23 Hz, 2H), 8.67 (s, 1H); ¹³C NMR (DMSO-d₆, 100 MHz) δ 160.7, 158.9 (q, 4.5 Hz), 158.6, 145.7, 142.7, 125.6, 123.3 (q, J = 269 Hz), 120.0, 113.8 (q, J = 34 Hz); HPLC ret. time: 7.720 min. LRMS (M+) 318.3, 320.3.

(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-(3,4-dichloro-phenyl)-amine: ¹H NMR (CDCl₃, 400 MHz) δ 7.39 (m, 3H), 7.86 (s, 1H), 8.60 (s, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ 160.2, 159.8, 157.6 (q, J = 5 Hz), 137.2, 133.2, 130.9, 127.9, 112.6 (q, J = 270 Hz), 121.9, 119.5, 115.0 (q, J = 34 Hz); HPLC ret. time: 8.837 min. LRMS (M+) 342.1, 344.1.

(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-o-tolyl-amine: ¹H NMR (CDCl₃, 400 MHz) δ 2.30 (s, 3H), 7.15 (m, 2H), 7.26 (m, 3H), 7.35 (d, J = 7.5 Hz, 1H), 8.51 (s, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ 161.5, 159.7, 157.8 (q, J = 4.5 Hz), 135.4, 131.1, 127.0, 126.3, 124.0, 122.8 (q, J = 270 Hz), 113.8 (q, J = 34 Hz), 18.3; HPLC ret. time: 7.663 min. LRMS (M+) 288.1, 290.1.

(3-Chloro-phenyl)-(4-chloro-5-trifluoromethyl-pyrimidin-2-yl)-amine: ¹H NMR (CDCl₃, 400 MHz) δ 7.11 (d, J = 9.1 Hz, 1H), 7.28 (t, J = 8.1 Hz, 1H), 7.38 (br s, 1H), 7.41 (d, J = 9.1 Hz, 1H), 7.75 (s, 1H), 8.59 (s, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ 60.4, 159.7, 157.6 (q, J = 4.5 Hz), 138.8, 135.1, 124.6, 122.6 (q, J = 269 Hz), 120.4, 118.3, 114.7 (q, J = 34 Hz); HPLC ret. time: 8.301 min. LRMS (M+) 308.1, 310.1.

(4-Chloro-phenyl)-(4-chloro-5-trifluoromethyl-pyrimidin-2-yl)-amine: ¹H NMR (CDCl₃, 400 MHz) δ 7.33 (d, J = 9.1 Hz, 2H), 7.42 (s, 1H), 7.53 (d, J = 8.7 Hz, 2H), 8.56 (s, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ 160.5, 159.7, 157.6 (q, J = 5 Hz), 136.2, 129.8, 129.4, 122.7 (q, J = 270 Hz), 118.6, 114.4 (q, J = 34 Hz); HPLC ret. time: 8.316 min. LRMS (M+) 308.1, 310.0.

(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-methyl-p-tolyl-amine: ¹H NMR (CDCl₃, 400 MHz) δ 2.36 (s, 3H), 3.52 (s, 3H), 7.13 (d, J = 8.3 Hz, 2H), 7.23 (d, J = 9.1 Hz, 2H), 8.39 (br s, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ 162.4, 159.1, 157.2, 141.3, 137.3, 130.4, 126.4, 123.2 (q, J = 269 Hz), 111.7 (q, J = 34 Hz), 39.6, 21.4; HPLC ret. time: 8.708 min. LRMS (M+) 302.2, 304.1.

(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-(3-oxazol-5-yl-phenyl)-amine: ¹H NMR (DMSO-d₆, 400 MHz) δ 7.44 (m, 2H), 7.63 (s, 1H), 7.65 (m, 1H), 8.07 (s, 1H), 8.44 (s, 1H), 8.82 (s, 1H), 10.78 (s, 1H); ¹³C NMR (DMSO-d₆, 100 MHz) δ 161.2, 158.9 (br), 158.4, 152.6, 151.0, 139.8, 130.2, 128.5, 123.6 (q, J = 269 Hz), 122.9, 121.2, 120.1, 116.4, 112.3 (q, J = 34 Hz); HPLC ret. time: 7.374 min. LRMS (M+) 341.2, 343.1.

4-(4-Chloro-5-trifluoromethyl-pyrimidin-2-ylamino)-benzamide: ¹H NMR (DMSO-d₆, 400 MHz) δ 7.26 (s, 1H), 7.74 (d, J = 8.7 Hz, 2H), 7.83 (d, J = 9.1 Hz, 2H), 8.44 (br s, 1H),

-12-

8.84 (s, 1H), 10.87 (s, 1H); ^{13}C NMR (DMSO- d_6 , 100 MHz) δ 168.0, 161.0, 158.9 (br), 158.4, 141.8, 129.6, 129.0, 123.5 (q, J = 268 Hz), 119.9, 112.6 (q, J = 34 Hz); HPLC ret. time: 5.605 min. LRMS (M $^{+}$) 317.1, 319.3.

(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-(4-methanesulfonyl-phenyl)-amine: ^1H NMR (DMSO- d_6 , 400 MHz) δ 3.15 (s, 3H), 7.87 (d, J = 8.7 Hz, 2H), 7.93 (d, J = 8.7 Hz, 2H), 8.89 (s, 1H), 11.10 (s, 1H); ^{13}C NMR (DMSO- d_6 , 100 MHz) δ 60.9, 158.9 (br), 158.5, 143.9, 135.3, 128.8, 123.4 (q, J = 269 Hz), 120.4, 113.3 (q, J = 34 Hz), 44.5; HPLC ret. time: 6.542 min. LRMS (M $^{+}$) 352.1, 354.1.

4-(4-Chloro-5-trifluoromethyl-pyrimidin-2-ylamino)-benzenesulfonamide: ^1H NMR (DMSO- d_6 , 400 MHz) δ 7.25 (s, 2H), 7.76 (d, J = 9.1 Hz, 2H), 7.83 (d, J = 9.1, 2H), 8.85 (s, 1H), 10.98 (s, 1H); ^{13}C NMR (DMSO- d_6 , 100 MHz) δ 161.0, 158.9 (br), 158.5, 142.2, 139.2, 127.3, 123.5 (q, J = 269 Hz), 120.3, 113.0 (q, J = 33 Hz); HPLC ret. time: 5.983 min. LRMS (M $^{+}$) 353.2, 355.2.

For those cases in which the amine nucleophile is aliphatic or is an aromatic amine capable of serving as a multidentate ligand for the Lewis Acid, additional equivalents of Lewis Acid are necessary for the reaction to proceed selectively.

Method C: N-(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-N',N'-dimethyl-benzene-1,4-diamine: To a solution of 5-trifluoromethyl-2,4-dichloropyrimidine (500 mg; 2.3 mmol) in 1:1 DCE/*t*-BuOH (20 mL) was added Zinc chloride (5.1 mL of a 1M solution in ether; 2.2 eq) at 0° C. After 1 hour, N,N-dimethyl-1,4-phenylenediamine (313 mg; 1 eq) was added followed by dropwise addition of a solution of triethylamine (279 mg; 1.1 eq) in 5 mL of DCE/*t*-BuOH. After stirring for 24 hours the reaction was concentrated. The product was obtained as a pale green solid (531 mg; 73%) following crystallization from 25% H₂O/methanol. ^1H NMR (DMSO- d_6 , 400 MHz) δ 2.83 (s, 6H), 6.69 (m, 2H), 7.39 (m, 2H), 8.65 (s, 1H), 10.32 (s, 1H); ^{13}C NMR (CDCl₃, 100 MHz) δ 161.3, 158.8 (br), 158.5 (br), 148.3, 128.2, 123.8 (q, J = 268 Hz), 123.0, 113.1, 110.8 (br), 41.0; HPLC ret. time: 7.901 min. LRMS (M $^{+}$) 317.3, 319.3.

The application of Method C with a variety of other amines allows for the preparation of a variety of other 2-amino-4-chloro-5-trifluoromethylpyrimidines including, but not limited to those outlined below.

(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-(2-methoxy-phenyl)-amine: ^1H NMR (CDCl₃, 400 MHz) δ 3.90 (s, 3H), 6.91 (d, J = 8.1 Hz, 1H), 7.01 (m, 1H), 7.07 (m, 1H), 8.09 (s, 1H), 8.38 (d, J = 7.9 Hz, 1H), 8.57 (s, 1H); ^{13}C NMR (CDCl₃, 100 MHz) δ 160.4, 159.3, 157.5 (q, J = 4.5 Hz), 148.6, 127.4, 124.1, 122.9 (q, J = 269 Hz), 121.1, 119.8, 113.7 (q, J = 34 Hz), 110.4, 56.0; HPLC ret. time: 8.151 min. LRMS (M $^{+}$) 304.2, 306.2.

(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-(4-methyl-benzyl)-amine: ^1H NMR (DMSO- d_6 , 400 MHz) δ 2.23 (s, 3H), 4.46 (m, 2H), 7.09 (m, 2H), 7.15 (m, 2H), 8.57 (2s, 1H), 8.98 (m, 1H); HPLC ret. time: 8.238 min. LRMS (M $^{+}$) 302.0, 304.0.

50054-130

-13-

(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-cyclohexyl-amine: ^1H NMR (DMSO- d_6 , 400 MHz) δ 1.10 (m, 1H), 1.23 (m, 4H), 1.55 (m, 1H), 1.65 (m, 2H), 1.81 (m, 2H), 3.69 (m, 2H), 8.47 (m, 1H), 8.55 (2 br s, 1H); HPLC ret. time: 8.548 min. LRMS (M+) 280.1, 282.1.

(4-Chloro-5-trifluoromethyl-pyrimidin-2-yl)-cyclohexylmethyl-amine: ^1H NMR (DMSO- d_6 , 400 MHz) δ 0.86 (m, 2H), 1.11 (m, 3H), 1.50 (m, 1H), 1.58 (m, 1H), 1.64 (m, 4H), 3.12 (m, 2H), 8.55 (m, 2H); HPLC ret. time: 9.039 min. LRMS (M+) 294.1, 296.1.


4-Chloro-2-piperidin-1-yl-5-trifluoromethyl-pyrimidine ^1H NMR (CDCl_3 - d_6 , 400 MHz) δ 1.60 (m, 4H), 1.68 (m, 2H), 3.82 (m, 4H), 8.36 (s, 1H); ^{13}C NMR (DMSO- d_6 , 100 MHz) δ 161.5, 159.1, 157.2 (q, J = 4.5 Hz), 123.4 (q, J = 268 Hz), 109.8 (q, J = 34 Hz), 45.4, 25.9, 24.7; HPLC ret. time: 8.915 min. LRMS (M+) 266.1, 268.2.

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and the accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

-14-

CLAIMS


1. A method of making a compound of the formula



wherein X^2 is a leaving group, and

- 5 R^3 and R^4 are substituents independently selected from the group consisting of hydrogen, an aromatic group and an aliphatic group; or taken together $-NR^3R^4$ can form a 4-11 membered aromatic or aliphatic ring;

wherein the method comprises reacting a compound of formula 10



- 10 with an amine of formula 3 (HNR^3R^4) in the presence of a Lewis Acid and a non-nucleophilic base to form a compound of formula 11, wherein X^1 is a leaving group.

2. A method of claim 1 wherein X^1 and X^2 are the same or different leaving groups independently selected from the group consisting of halide, arylsulfonate, alkylsulfonate, perfluoroalkylsulfonate, arylsulfinate or alkylsulfinate.

- 15 3. A method of claim 2 wherein X^1 and X^2 are the same or different and are each independently halides.

4. A method of claim 3 wherein X^1 and X^2 are chloride.

5. A method of claim 1 wherein said amine, HNR^3R^4 , is an aromatic amine and 0.25-10 equivalents of Lewis Acid are used relative to the amount of pyrimidine 10.

- 20 6. A method of claim 1 wherein said amine, HNR^3R^4 , is an aromatic amine and 0.5-3 equivalents of Lewis Acid are used relative to the amount of pyrimidine 10.

7. A method of claim 5 wherein the Lewis Acid is a salt of a metal ion selected from the group consisting of Zn, Mg, Sn, Ti, Al, B, Li, Ag, and Cu.

8. A method of claim 7 wherein said Lewis Acid is a salt of Zn or Mg.

- 25 9. A method of claim 7 wherein said Lewis Acid is $ZnCl_2$.

10. A method of claim 1 wherein said amine, HNR^3R^4 , is an aliphatic amine or is an aromatic amine capable of serving as a multidentate ligand for the Lewis Acid and 0.5-10 equivalents of Lewis Acid are used relative to the amount of pyrimidine 10.

- 30 11. A method of claim 1 wherein said amine, HNR^3R^4 , is an aliphatic amine or is an aromatic amine capable of serving as a multidentate ligand for the Lewis Acid and 1-4 equivalents of Lewis Acid are used relative to the amount of pyrimidine 10.

-15-

12. A method of claim 10 wherein the Lewis Acid is a salt of a metal ion selected from the group consisting of Zn, Mg, Sn, Ti, Al, B, Li, Ag, and Cu.
13. A method of claim 12 wherein said Lewis acid is a salt of Zn or Mg.
14. A method of claim 12 wherein said Lewis Acid is ZnCl_2 .