



(86) Date de dépôt PCT/PCT Filing Date: 2002/10/24  
(87) Date publication PCT/PCT Publication Date: 2003/05/01  
(45) Date de délivrance/Issue Date: 2009/10/13  
(85) Entrée phase nationale/National Entry: 2003/06/23  
(86) N° demande PCT/PCT Application No.: JP 2002/011040  
(87) N° publication PCT/PCT Publication No.: 2003/035015  
(30) Priorités/Priorities: 2001/10/26 (JP2001-329600);  
2001/11/30 (JP2001-365423);  
2002/04/05 (JP2002-103976)

(51) Cl.Int./Int.Cl. *A61K 8/92* (2006.01),  
*A61K 8/04* (2006.01), *A61K 8/37* (2006.01),  
*A61K 8/39* (2006.01), *A61K 8/55* (2006.01),  
*A61Q 19/02* (2006.01)  
(72) Inventeurs/Inventors:  
TAKASE, YOSHIHIKO, JP;  
UCHIDA, KAZUHITO, JP;  
HONDA, NAOTERU, JP;  
KATO, TOMOHARU, JP;  
YAMAZAKI, NAGAHIRO, JP  
(73) Propriétaire/Owner:  
TAIYO KAGAKU CO., LTD., JP  
(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : COMPOSITION D'UN AEROSOL HUILEUX EXPANSIBLE  
(54) Title: COMPOSITION FOR OILY FOAMABLE AEROSOL

(57) **Abrégé/Abstract:**

The present invention provides a composition for an oily foamable aerosol, characterized in that the composition comprises a fat or oil, and a polyglycerol fatty acid ester and/or an organic and fatty acid esters of glycerol; and an aerosol comprising the composition as defined in above and a propellant.

## ABSTRACT

The present invention provides a composition for an oily foamable aerosol, characterized in that the composition comprises a fat or oil, and a polyglycerol fatty acid ester and/or an organic and fatty acid esters of glycerol; and an aerosol comprising the composition as defined in above and a propellant.

DESCRIPTIONCOMPOSITION FOR OILY FOAMABLE AEROSOL5 TECHNICAL FIELD

The present invention relates to a composition for oily foamable aerosol which is useful as a cleansing agent or the like for oily cosmetics.

BACKGROUND ART

10 Conventional compositions for aerosol include those which form stable-type foams, fast-disintegrating-type foams, spray-type foams and the like according to the combinations of a stock solution and a propellant. The stable-type foams are constituted by an O/W type emulsion, comprising two phases of an inner phase comprising an oily phase in which a propellant is dissolved, and  
15 an aqueous phase having poor compatibility with the propellant. The composition for the foamable aerosol of an O/W emulsion includes cosmetics for cleansing. However, since the cosmetics have low detergency, they have been inappropriate for cleaning oily make-ups. In order to remove oily make-ups, an oily component is most effective, and it is most preferable to obtain a stable-type  
20 foam from stock solutions of an oily component and a propellant. However, since the oily component itself is not foamable, it has been difficult to obtain a stable-type foam.

DISCLOSURE OF INVENTION

25 An object of the present invention is to provide a composition for an oily

foamable aerosol which is excellent in detergency, applicability, and favorable wash-up property and stability of the foam.

As a result of intensive studies in order to solve the above-mentioned problems, the present inventors have found that a composition comprising a fat or oil, and a polyglycerol fatty acid ester and/or an organic and fatty acid esters of glycerol is excellent in detergency, applicability, wash-up property and stability of the foam (excellent form-keeping property or form-sustaining property or the like of the foam formed by spraying an aerosol obtained by combining the composition and a propellant, or blowing air into the composition to mix the composition with air. The present invention has been accomplished thereby.

Specifically, the present invention relates to:

- [1] a composition for an oily foamable aerosol, characterized in that the composition comprises a fat or oil, and a polyglycerol fatty acid ester and/or an organic and fatty acid esters of glycerol;
- [2] the composition according to the above [1], further comprising a phospholipids;
- [3] the composition according to the above [1], wherein a constituting fatty acid of the polyglycerol fatty acid ester has 14 or less carbon atoms;
- [4] the composition according to any one of the above [1] to [3], wherein the polyglycerol fatty acid ester is a diglycerol fatty acid ester and/or a triglycerol fatty acid ester;
- [5] the composition according to any one of the above [1] to [4], wherein the organic and fatty acid esters of glycerol is a succinic and fatty acid esters of glycerol and/or a citric and fatty acid esters of glycerol;
- [6] The composition according to any one of the above [1] to [5], wherein a

monoester is contained in the polyglycerol fatty acid ester in an amount of 50% by weight or more; and

[7] an aerosol comprising the composition as defined in any one of the above [1] to [6] and a propellant.

5

#### BEST MODE FOR CARRYING OUT THE INVENTION

The composition for an oily foamable aerosol (hereinafter referred to as “composition for aerosol”) of the present invention is a composition comprising a fat or oil and a specified surfactant. The composition is very excellent in  
10 detergency, applicability, wash-up property and stability of the foam, as compared to those of a similar conventional composition for aerosol.

The composition for aerosol of the present invention comprises as essential components a fat or oil, and a polyglycerol fatty acid ester and/or an organic and fatty acid esters of glycerol. There are roughly the following three  
15 embodiments.

(Embodiment A)

A composition for an oily foamable aerosol, characterized in that the composition comprises at least one member of each of a fat or oil, a polyglycerol fatty acid ester and/or an organic and fatty acid esters of glycerol, and a  
20 phospholipids.

(Embodiment B)

A composition for an oily foamable aerosol, characterized in that the composition comprises at least two members selected from the group consisting of a polyglycerol fatty acid ester, an organic and fatty acid esters of glycerol, and  
25 a phospholipid, and a fat or oil.

(Embodiment C)

A composition for an oily foamable aerosol, characterized in that the composition comprises a polyglycerol fatty acid ester of which constituting fatty acid is composed of 14 or less carbon atoms, and a fat or oil.

5

1. Composition for Aerosol of Embodiment A

The fat or oil is not particularly limited, as long as the fat or oil is in a liquid state during foaming. The fat or oil may be a natural product or a chemically synthesized product. The fat or oil includes oily components such as hydrocarbons, fatty acids, higher alcohols, silicones and esters, besides general liquid fats and oils. Each of the fat or oil can be used alone or in admixture of at least two kinds.

The term “(being) in a liquid state during foaming” as referred to herein means that the fat or oil shows a liquid state at a temperature at which the composition for aerosol is generally used as an aerosol, for instance, 15°C.

The liquid fats and oils include, for instance, avocado oil, camellia oil, turtle oil, macadamia nut oil, corn oil, mink oil, olive oil, rapeseed oil, egg yolk oil, sesame oil, persic oil, wheat germ oil, *sasanqua* oil, castor oil, linseed oil, safflower oil, cottonseed oil, nettle tree oil, soybean oil, peanut oil, tea oil, *kaya* oil, rice bran oil, Chinese wood oil, wood oil, jojoba oil, germ oil and the like, and glycerol trioctanoate, glycerol triisopalmitate, and the like. The hydrocarbon includes, for instance, liquid paraffin, squalene, squalane, pristan, and the like. The fatty acid includes, for instance, oleic acid, isostearic acid and the like. The higher alcohol includes, for instance, lauryl alcohol, oleyl alcohol, isostearyl alcohol, octyl dodecanol and the like. The silicone includes, for instance, methyl

polysiloxane, methyl phenyl polysiloxane, methyl hydrogen polysiloxane, decamethyl polysiloxane, and the like. The ester includes, for instance, isopropyl myristate, isopropyl palmitate, cetyl octanoate, hexyl laurate, oleyl oleate, decyl oleate, octyl dodecyl myristate, hexyl decyl dimethyloctanoate, diethyl phthalate, dibutyl phthalate and the like. In addition, those obtained by 5  
subjecting the above-mentioned fats and oils to such treatments as hydrogen-addition (referred to as hydrogenation in some cases) or separation may be included in the fats and oils. The fat or oil is preferably a combination of one or more kinds of the hydrocarbons and the esters, from the viewpoint of improving 10  
detergency for the oily make-ups. The fat or oil may contain an unsaturated fatty acid, side-chain fatty acid, a diglyceride, a monoglyceride or the like as the glyceride component as long as the amount of the glyceride component does not inhibit the exhibition of the desired effects of the present invention.

The content of the fat or oil in the composition for aerosol of the present 15  
invention is not particularly limited. The content of the fat or oil is preferably from 10 to 99% by weight, more preferably from 30 to 95% by weight, still more preferably from 60 to 95% by weight.

The polyglycerol of the polyglycerol fatty acid ester includes, but not particularly limited thereto, for instance, glycerol polymers such as diglycerol, 20  
triglycerol, tetraglycerol, pentaglycerol, hexaglycerol, heptaglycerol, octaglycerol, nonaglycerol and decaglycerol. The polyglycerol is preferably diglycerol and triglycerol, from the viewpoint of improving stability of the foam. In other words, the polyglycerol fatty acid esters are preferably a diglycerol fatty acid ester and a triglycerol fatty acid ester. Also, the fatty acid of the 25  
polyglycerol fatty acid ester is not particularly limited. As the fatty acid, a

saturated or unsaturated, linear or branched fatty acid having 8 to 22 carbon atoms and a hydroxy-compound thereof are preferred, and a fatty acid having 14 to 18 carbon atoms is more preferred, from the viewpoint of improving stability of the foam.

5           As the polyglycerol fatty acid ester, there can be preferably used, for instance, diglycerol myristate, diglycerol palmitate, diglycerol stearate, triglycerol myristate, triglycerol palmitate, triglycerol stearate, and the like.

          Each of the polyglycerol fatty acid esters can be used alone or in admixture of at least two kinds. Also, the polyglycerol fatty acid ester may be a  
10           mixture of the polyglycerol fatty acid esters having a given esterification degree. The degree of esterification is not particularly limited. When expressed as the monoester content of the mixture, the monoester content is preferably 50% by weight or more, more preferably 70% by weight or more from the viewpoint of improving wash-up property and stability of the foam.

15           The polyglycerol fatty acid ester is obtained by, for instance, carrying out esterification reaction of a polyglycerol with a fatty acid with heating in the presence of an inert gas such as nitrogen gas in accordance with a known method, and further subjecting the resulting product to distillation and purification as desired, without being limited to this process.

20           The content of the polyglycerol fatty acid ester in the composition for aerosol of the present invention is not particularly limited. The content of the polyglycerol fatty acid ester is preferably from 0.01 to 50 parts by weight, more preferably from 0.05 to 30 parts by weight, still more preferably from 0.1 to 20 parts by weight, especially preferably from 1 to 20 parts by weight, based on  
25           100 parts by weight of the fat or oil, which is the constituent of the present

invention.

Incidentally, when the content of the constituents other than the fat or oil in the composition for aerosol of the present invention is calculated on the basis of the content of the fat or oil, if other oily components besides the constituents  
5 of the composition are contained in addition to the fat or oil, the amount of the oily components is included in the amount of the fat or oil.

The organic and fatty acid esters of glycerol refers to a compound in which one or two organic acids are bonded to a glycerol fatty acid ester. When the organic acids are bonded in a plural number, those organic acids may be  
10 identical to or different from each other.

The organic acid is not particularly limited. Acetic acid, lactic acid, citric acid, succinic acid and diacetylated tartaric acid are preferred, and citric acid and/or succinic acid is more preferred, from the viewpoint of improving stability of the foam. Especially, an organic and fatty acid esters of glycerol in which one  
15 or two succinic acids are bonded to glycerol is preferably used. In other words, as the organic and fatty acid esters of glycerol, a succinic and fatty acid esters of glycerol and a citric and fatty acid esters of glycerol are preferable, and the succinic and fatty acid esters of glycerol is more preferable.

The fatty acid of the organic and fatty acid esters of glycerol is not  
20 particularly limited. As the fatty acid, a saturated or unsaturated, linear or branched fatty acid having 8 to 22 carbon atoms, or a hydroxy-compound thereof is preferred, and a fatty acid having 12 to 18 carbon atoms is more preferred, from the viewpoint of improving wash-up property and stability of the foam.

As the organic and fatty acid esters of glycerol, for instance, there can be  
25 suitably used succinic and lauric acid esters of glycerol, succinic and myristic

acid esters of glycerol, succinic and oleic acid esters of glycerol, succinic and palmitic acid esters of glycerol, succinic and stearic acid esters of glycerol, citric and lauric acid esters of glycerol, citric and myristic acid esters of glycerol, citric and oleic acid esters of glycerol, citric and palmitic acid esters of glycerol, citric and stearic acid esters of glycerol, and the like.

Each of the organic and fatty acid esters of glycerol can be used alone or in admixture of at least two kinds.

The organic and fatty acid esters of glycerol is obtained by, for instance, carrying out esterification reaction of a glycerol fatty acid ester with an organic acid with heating in the presence of an inert gas in accordance with a known method, and further subjecting the resulting product to distillation and purification as desired, without being limited to this process.

The content of the organic and fatty acid esters of glycerol in the composition for aerosol of the present invention is not particularly limited. The content of the organic and fatty acid esters of glycerol is preferably from 0.01 to 50 parts by weight, more preferably from 0.05 to 20 parts by weight, still more preferably from 1 to 10 parts by weight, based on 100 parts by weight of the fat or oil, which is the constituent of the present invention.

At least one of the polyglycerol fatty acid ester or the organic and fatty acid esters of glycerol may be contained in the composition for aerosol of this embodiment. When both the compounds are contained, the mixing ratio of the both compounds (polyglycerol fatty acid ester/organic and fatty acid esters of glycerol) is preferably from 9/1 to 1/9, more preferably from 7/3 to 3/7, on a weight basis. Also, the total content of the polyglycerol fatty acid ester and the organic and fatty acid esters of glycerol is preferably from 0.01 to 50 parts by

weight, more preferably from 0.1 to 30 parts by weight, still more preferably from 1 to 20 parts by weight, based on 100 parts by weight of the fat or oil, which is the constituent of the present invention.

As the phospholipid usable in the present invention, natural phospholipids such as egg yolk lecithin, soybean lecithin, rapeseed lecithin, cottonseed lecithin corn lecithin, which are extracted from animals or plants, or hydrogenated products thereof, enzymatically degraded products thereof, acetylated products thereof, hydroxylated products thereof, halogenated products thereof and sulfonated products thereof are preferred. The egg yolk lecithin, the soybean lecithin, or hydrogenated products thereof are more preferable from the viewpoint of improving wash-up property and stability of the foam. Each of those phospholipids can be used alone or in admixture of at least two kinds.

The content of the phospholipid in the composition for aerosol of the present invention is not particularly limited. The content of the phospholipid is preferably from 0.01 to 30 parts by weight, more preferably from 0.05 to 20 parts by weight, still preferably from 1 to 10 parts by weight, based on 100 parts by weight of the fat or oil, which is the constituent of the present invention.

In the composition for aerosol of the present invention, other components can be properly formulated in addition to the above-mentioned components, within the range in which the object of the present invention would be achieved. The other components include, for instance, a known surfactant, a powder of titanium oxide, talc, bentonite, cation and mica, preservatives, coloring agents, antioxidants, perfumes, chemicals and solid oil components other than the above-mentioned fat or oil, semi-solid oily component, alcohols, moisturizing agents, and the like.

As the composition for aerosol of this embodiment, it is especially preferable that the content of the fat or oil is from 60 to 95% by weight, the content of the polyglycerol fatty acid ester and/or the organic and fatty acid esters of glycerol is from 1 to 20% by weight, the content of the phospholipid is from 1 to 10% by weight, and the content of the other components is from 0 to 10% by weight.

## 2. Composition for Aerosol of Embodiment B

The polyglycerol fatty acid ester, the organic and fatty acid esters of glycerol, the phospholipids, the fat or oil, and optionally formulated other components, which are constituents of the composition for aerosol of this embodiment, and their preferred embodiments and contents are the same as mentioned above.

The composition for aerosol of this embodiment may contain at least two kinds of any one of a polyglycerol fatty acid ester, an organic and fatty acid esters of glycerol and a phospholipid.

For instance, when the polyglycerol fatty acid ester and the organic and fatty acid esters of glycerol are contained, it is preferable that their mixing ratio (polyglycerol fatty acid ester/organic and fatty acid esters of glycerol) is the same as that of the above-mentioned Embodiment A. When the polyglycerol fatty acid ester or the organic and fatty acid esters of glycerol and the phospholipids are contained, their mixing ratio (polyglycerol fatty acid ester or organic and fatty acid esters of glycerol/phospholipids), is preferably from 1/1 to 3/1, more preferably from 1/1 to 2/1, on a weight basis. Also, the total content of any of the two kinds of these components is preferably from 0.02 to 50 parts by

weight, more preferably from 0.1 to 30 parts by weight, still more preferably from 1 to 20 parts by weight, based on 100 parts by weight of the fat or oil, which is the constituent of the present invention.

As the composition for aerosol of this embodiment, it is especially  
5 preferable that the content of the fat or oil is from 60 to 95% by weight, the content of at least two kinds selected from the group consisting of the polyglycerol fatty acid esters, the organic and fatty acid esters of glycerol and the phospholipids is from 1 to 30% by weight, and the content of the other  
10 components is from 0 to 10% by weight.

### 3. Composition for Aerosol of Embodiment C

The fat or oil, which is the constituent of the composition for aerosol of this embodiment, and its preferred embodiment and content are the same as described above. On the other hand, as the polyglycerol fatty acid ester, those of  
15 which constituting fatty acid has 14 or less carbon atoms are used, and its content is 0.1 to 50 parts by weight, more preferably from 1 to 30 parts by weight based on 100 parts by weight of the fat or oil, which is the constituent of the present invention.

The polyglycerol fatty acid ester of which constituting fatty acid has 14 or  
20 less carbon atoms refers more specifically to an ester of a saturated or unsaturated, linear or branched fatty acid having 14 or less carbon atoms and a polyglycerol; an ester of a hydroxy-compound of the fatty acid and a polyglycerol; and the like. There can be cited an ester of a fatty acid having preferably 8 to 14 carbon atoms, more preferably 12 or 14, or its hydroxy-  
25 compound and a polyglycerol, from the viewpoint of improving detergency and

wash-up property of the oily make-up, and stability of the foam. In addition, it is preferable that the polyglycerol, which is a constituting unit, is a diglycerol or a triglycerol. As to the esterification degree of the polyglycerol fatty acid ester, those within the preferred ranges are preferable. The polyglycerol fatty acid ester used in this embodiment includes, for instance, various polyglycerol fatty acid esters exemplified above, of which constituting fatty acid has 14 or less carbon atoms.

As other components, for instance, those exemplified above are used within the range so that the exhibition of the desired effects of the present invention would not be hindered.

As the composition for aerosol of this embodiment, it is especially preferable that the content of the fat or oil is from 60 to 95% by weight, the content of the polyglycerol fatty acid esters of which constituting fatty acid has 14 or less carbon atoms is from 1 to 30% by weight, and the content of the other components is from 0 to 15% by weight.

The composition for aerosol of the present invention as described above can be prepared in accordance with a known method. For instance, the composition can be prepared by appropriately mixing various components mentioned above, heating the components to dissolve them at, for instance, 40° to 80°C with stirring, and cooling the resulting solution to an ambient temperature.

The application for the composition for aerosol of the present invention includes, but not particularly limited to, for instance, such applications as a detergent for oil stains, a cleansing agent for oily cosmetics, a suntan oil, a baby

oil, a hair oil and foamy massage oil and the like, and the application as a cleansing agent is preferable.

The composition for aerosol of the present invention is preferably used as aerosol described below in which the composition is combined with a propellant.

5 On the other hand, since the composition can be foamed by blowing air into the composition to mix the composition with air without formulating a propellant, which is a so-called non-gas type, the composition can be used by packing into a container in which mixing by air-blowing during spraying can be carried out.

The aerosol of the present invention comprises the composition for  
10 aerosol of the present invention and an arbitrary propellant, which can be prepared by a known process. For instance, the aerosol can be prepared by introducing the composition for aerosol of the present invention into a given container, and filling the container with a propellant. As the container, for instance, there can be preferably used an aerosol can and the like. Also, as the  
15 propellant, there can be preferably used, for instance, chlorofluorocarbons such as trichlorofluoromethane, dichlorodifluoromethane, dichlorofluoromethane, trichlorotrifluoromethane and dichlorotetrafluoromethane; and liquefied gases such as propane, isobutane, isopentane, normal butane and mixtures thereof, which are liquefied petroleum gases (which may be hereinafter referred to as  
20 "LPG"). In addition to the above-mentioned propellant, dimethyl ether, carbon dioxide gas, and nitrogen gas may be used as a propellant. Each of the propellants may be used alone or in admixture of at least two kinds. The content of the propellant in the aerosol of the present invention is preferably from 2 to 90% by weight, more preferably from 3 to 90% by weight, still more preferably  
25 from 5 to 70% by weight, from the viewpoint of obtaining an excellent foam

formation. The content of the composition for aerosol is preferably from 10 to 98% by weight, more preferably from 10 to 97% by weight, still more preferably from 30 to 95% by weight.

5 Next, the present invention will be described in further detail by means of Examples, without intending to limit the present invention thereto.

Example 1-1 Oily Wash-Away type Cleansing Oil Cosmetics

10 Each of oily wash-away type cleansing oil cosmetics was produced by the following process on the basis of the composition of the raw materials shown in Table 1.

Table 1

Raw Materials	Inventive Product A (g)	Inventive Product B (g)	Inventive Product C (g)	Inventive Product D (g)
1 Liquid Paraffin	34.8	34.8	34.8	34.8
2 Olive Oil	20	20	20	20
3 Isopropyl Myristate	20	25	25	25
4 Dibutylhydroxytoluene	0.2	0.2	0.2	0.2
5 Diglycerol Monopalmitate	5	5	5	—
6 Soybean Lecithin	5	5	—	5
7 Palmitic and Succinic Acid Esters of Glycerol	5	—	5	5

15 Ingredients 1 to 7 were heated and dissolved at 80°C, and the resulting solution was cooled to ambient temperature, to give a composition for aerosol of the above-mentioned cosmetics. Thereafter, the composition was packed into an

aerosol can, a valve was set thereto, and 10 g of a liquefied petroleum gas was packed therein as a propellant to 90 g of the composition, to give an aerosol of the above-mentioned cosmetics. The resulting cosmetics were referred to as the inventive products A to D, respectively.

5

Comparative Example 1-1 Oily Wash-Away type Cleansing Oil Cosmetics

Each of oily wash-away type cleansing oil cosmetics was produced in the same manner as in Example 1-1 on the basis of the composition of the raw materials shown in Table 2.

10

Table 2

Raw Materials	Comparative Product E (g)	Comparative Product F (g)	Comparative Product G (g)	Comparative Product H (g)
1 Liquid Paraffin	44.8	39.8	39.8	39.8
2 Olive Oil	20	20	20	20
3 Isopropyl Myristate	25	25	25	25
4 Dibutylhydroxytoluene	0.2	0.2	0.2	0.2
5 Diglycerol Monopalmitate	—	5	—	—
6 Soybean Lecithin	—	—	—	5
7 Palmitic and Succinic Acid Esters of Glycerol	—	—	5	—

15

Specifically, aerosols of the above-mentioned cosmetics as comparative products to the inventive products A to D were prepared in the same manner as in the inventive products A to D. The same type oily component-formulated composition as the inventive product but not containing any of polyglycerol fatty acid ester, phospholipid, and organic and fatty acid esters of glycerol is referred

to as a comparative product E; the same type oily component-formulated composition containing only polyglycerol fatty acid ester is referred to as a comparative product F; the same type oily component-formulated composition containing only organic and fatty acid esters of glycerol is referred to as a comparative product G; and the same type oily component-formulated composition containing only phospholipid is referred to as a comparative product H.

#### Test Example 1-1

10 The inventive products A to D were compared with the comparative products E to H.

As a result, the inventive products A to D formed fine and stable foams. By contrast, the comparative product E did not form foams, and the comparative examples F to H formed rough foams which disappeared immediately.

15 The results clearly show the followings: The compositions of the present invention had excellent foam stability after spraying, whereas the composition for aerosol not containing polyglycerol fatty acid ester, phospholipid, and organic and fatty acid esters of glycerol (comparative product E) did not form foams. In addition, the compositions containing only one kind of polyglycerol fatty acid ester, phospholipid, and organic and fatty acid esters of glycerol (comparative products F, G, H) formed rough foams and had poor foam stability.

#### Test Example 1-2

25 Detergency for the inventive products A to D of Example 1-1 was compared to that of conventional O/W emulsion type foamable aerosol cleansing

cosmetics having the raw material composition shown in Table 3.

Ten female panelists were asked to apply an oily foundation. The results of the test for the degree of removal of the foundation are shown in Table 4. Each of the numerical figures in Table 4 shows the number of female panelists which were judged that the foundation was “completely removed,” “slightly removed,” or “not removed” by the use of each of the cosmetics. Here, “slightly removed” includes a case where the degree of removal of the foundation is insufficient.

10

Table 3

	Raw Materials	Weight (g)
1	Polyoxyethylene Lauryl Ether	10
2	Liquid Paraffin	40
3	Purified Water	Proper Amount
4	Liquefied Petroleum Gas	10
5	Diethyl Ether	5
	Total	100

The O/W emulsion type foamable aerosol cleansing cosmetics were produced by the following process on the basis of the composition of the raw materials shown in Table 3.

15

Ingredients 1 to 3 were dissolved with stirring. Thereafter, the resulting solution was packed into an aerosol can, a valve was set, and Ingredients 4 and 5 were packed therein, to give an O/W emulsion type aerosol of the above-mentioned cosmetics.

Table 4

	Completely Removed	Slightly Removed	Not Removed
Inventive Product A	9	1	0
Inventive Product B	9	1	0
Example 1-1 Inventive Product C	9	1	0
Inventive Product D	9	1	0
O/W Emulsion type Aerosol	0	2	8

Table 4 clearly shows the followings: The inventive products A to D had very high detergency as compared to the conventional O/W emulsion type foamable aerosol cleansing cosmetics. Also, the inventive products had the advantages of not dripping when taking onto a palm, easily applying to face, etc. as compared that of conventional pump-type cleansing cosmetics, so that their applicability was excellent. Further, the inventive products had substantially no greasy texture upon wash up, thereby showing excellent wash-up property.

10

Example 1-2 Suntan Oil

	Raw Materials	Weight (g)
1	Octamethyl Cyclotetrasiloxane	15.0
2	Decamethyl Cyclopentanesiloxane	15.0
3	C11-C15 Alkyl Benzoate	5.0
4	Octyl Paradimethylaminobenzoate	7.0
5	Oxybenzone	0.5
6	Olive Oil	11.0

7	Diglycerol Dimyristate	5.0
8	Hydrogenated Soybean Lecithin	5.0
9	Stearyl Glycyrrhetinate	0.2
10	Cetyl Isooctanoate	21.3
11	Liquefied Petroleum Gas	10.0
12	Diethyl Ether	5.0

Ingredients 1 to 10 were dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of a suntan oil.

Thereafter, the composition was packed into an aerosol can, a valve was set, and

5 Ingredients 11 and 12 were packed therein, to give an aerosol for the suntan oil.

The resulting suntan oil was excellent in foam stability and applicability.

#### Example 1-3 Wipe-off type Cleansing Cosmetics

	Raw Materials	Weight (g)
1	Isopropyl Myristate	42.0
2	Oleyl Alcohol	40.0
3	2-Hydroxy-4-methoxybenzophenone	3.0
4	White Vaseline	2.0
5	Diglycerol Monomyristate	7.0
6	Egg Yolk Lecithin	6.0

10 Ingredients 1 to 6 were dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of the above-mentioned cosmetics. Thereafter, air was blown into and mixed with the resulting

cosmetics to form a foam. As a result, its foam stability was excellent. Also, the cosmetics were excellent in applicability.

Example 1-4 Wash-Away type Cleansing Cosmetics

	Raw Materials	Weight (g)
1	Castor Oil	19.88
2	Oleyl Oleate	15.0
3	Diethyl Phthalate	15.0
4	Tocopherol Acetate	0.01
5	Perfume	0.1
6	Triglycerol Monopalmitate	9.0
7	Enzyme-Degradable Soybean Lecithin	5.0
8	Trichlorocarbanilide	1.0
9	Stearyl Glycyrrhetinate	0.01
10	LPG	35.0

5

Ingredients 1 to 9 were dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of the above-mentioned cosmetics. Thereafter, the composition was packed into an aerosol can, a valve was set, and Ingredient 10 was packed therein, to give an aerosol for the above-mentioned cosmetics. The resulting cosmetics were excellent in foam stability and applicability.

10

Example 1-5 Wash-Away type Cleansing Cosmetics

	Raw Materials	Weight (g)
1	Castor Oil	19.88
2	Oleyl Oleate	15.0
3	Diethyl Phthalate	10.0
4	Tocopherol Acetate	0.01
5	Perfume	0.1
6	Triglycerol Monopalmitate	9.0
7	Enzyme-Degradable Soybean Lecithin	5.0
8	Palmitic and Succinic Acid Esters of Glycerol	5.0
9	Trichlorocarbanilide	1.0
10	Stearyl Glycyrrhetinate	0.01
11	LPG	35.0

Ingredients 1 to 10 were dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of the above-mentioned cosmetics. Thereafter, the composition was packed into an aerosol can, a valve was set, and Ingredient 11 was packed therein, to give an aerosol for the above-mentioned cosmetics. The resulting cosmetics were excellent in foam stability and applicability.

Example 1-6 Baby Oil

	Raw Materials	Weight (g)
1	Squalane	34.44
2	Oleyl Alcohol	20.0
3	Polymethylsiloxane	5.0
4	Stearyl Glycyrhetinate	0.01
5	Beeswax	0.5
6	Tocopherol Acetate	0.05
7	Monooleic and Disuccinic Acid Esters of Glycerol	5.0
8	Soybean Lecithin	5.0
9	Trichlorotrifluoromethane	20.0
10	Dimethyl Ether	10.0

Ingredients 1 to 8 were dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of a baby oil. Thereafter, the composition was packed into an aerosol can, a valve was set, and Ingredients 9 and 10 were packed therein, to give an aerosol for the baby oil. The resulting baby oil was excellent in foam stability and applicability.

Example 1-7 Hair Oil

	Raw Materials	Weight (g)
1	Camellia Oil	30.0
2	Hexyl Laurate	22.0
3	Tocopherol Acetate	0.01

4	Perfume	0.5
5	Monopalmitic and Disuccinic Acid Esters of Glycerol	4.0
6	Soybean Lecithin	4.0
7	Trichlorofluoromethane	40.0
8	Nitrogen gas	0.1

Ingredients 1 to 6 were dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of a hair oil. Thereafter, the composition was packed into an aerosol can, a valve was set, and Ingredients 5 7 and 8 were packed therein, to give an aerosol for the hair oil. The resulting hair oil was excellent in foam stability and applicability.

#### Example 1-8 Hair Oil

	Raw Materials	Weight (g)
1	Camellia Oil	30.0
2	Hexyl Laurate	22.0
3	Tocopherol Acetate	0.01
4	Perfume	0.5
5	Monopalmitic and Disuccinic Acid Esters of Glycerol	4.0
6	Diglycerol Monolaurate	4.0
7	Trichlorofluoromethane	40.0
8	Nitrogen gas	0.1

10 Ingredients 1 to 6 were dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of a hair oil. Thereafter,

the composition was packed into an aerosol can, a valve was set, and Ingredients 7 and 8 were packed therein, to give an aerosol for the hair oil. The resulting hair oil was excellent in foam stability and applicability.

5 Example 1-9 Foamy Massage Oil Cosmetics

	Raw Materials	Weight (g)
1	Liquid Paraffin	52.0
2	Polyoxyethylene Glyceryl Triisostearate	4.0
3	Squalane	10.0
4	Perfume	0.8
5	Macadamia Nut Oil	5.0
6	Isostearyl Glycyrrhetinate	0.1
7	Monopalmitic and Disuccinic Acid Esters of Glycerol	4.0
8	Diglycerol Monolaurate	4.0
9	Trichlorofluoromethane	20.0
10	Nitrogen gas	0.1

Ingredients 1 to 8 were dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of the above-mentioned cosmetics. Thereafter, the composition was packed into an aerosol can, a valve was set, and Ingredients 9 and 10 were packed therein, to give an aerosol for the above-mentioned cosmetics. The resulting cosmetics were excellent in foam stability and applicability.

Example 2-1 Oily Wash-Away type Cleansing Oil Cosmetics

Each of oily wash-away type cleansing oil cosmetics was produced by the following process on the basis of the composition of the raw materials shown in Table 5.

5

Table 5

	Raw Materials	Inventive Products (g)						
		A	B	C	D	E	F	G
1	Liquid Paraffin	34.8	34.8	34.8	34.8	34.8	34.8	34.8
2	Olive Oil	25	20	20	20	20	20	20
3	Isopropyl Myristate	25	25	25	25	20	20	20
4	Dibutylhydroxytoluene	0.2	0.2	0.2	0.2	0.2	0.2	0.2
5	Diglycerol Monomyristate	5	—	5	5	—	5	5
6	Diglycerol Monolaurate	—	10	—	—	—	—	—
7	Diglycerol Monocaprata	—	—	5	—	5	—	—
8	Triglycerol Monolaurate	—	—	—	5	10	—	5
9	Triglycerol Monomyristate	—	—	—	—	—	10	5

Ingredients 1 to 9 were mixed, and heated and dissolved at 80°C, and the resulting solution was cooled to ambient temperature, to give a composition for aerosol of the above-mentioned cosmetics. Thereafter, the composition was packed into an aerosol can, a valve was set thereto, and 10 g of a liquefied petroleum gas was packed therein as a propellant to 90 g of the composition, to give an aerosol of the above-mentioned cosmetics. The resulting cosmetics were referred to as the inventive products A to G, respectively.

15

#### Comparative Example 2-1 Oily Wash-Away type Cleansing Oil Cosmetics

Each of oily wash-away type cleansing oil cosmetics was produced by the

following process on the basis of the composition of the raw materials shown in Table 6.

Table 6

Raw Materials		Comparative Products (g)				
		H	I	J	K	L
1	Liquid Paraffin	44.8	39.8	39.8	39.8	39.8
2	Olive Oil	20	20	20	20	20
3	Isopropyl Myristate	25	25	25	25	25
4	Dibutylhydroxytoluene	0.2	0.2	0.2	0.2	0.2
5	Polyoxyethylene(6) Oleyl Ether	—	10	—	—	—
6	Triglycerol Monopalmitate	—	—	10	—	—
7	Diglycerol Monostearate	—	—	—	10	—
8	Diglycerol Monopalmitate	—	—	—	—	10

5 \*: The number inside the parenthesis ( ) in Raw Material 5 shows the number of moles of ethylene oxide added, hereinafter the same.

Compositions for aerosols and aerosols were obtained in accordance with the method described in Example 2-1 using Raw Materials 1 to 8. Here, the product not containing a surfactant is referred to as a comparative product H; a product containing polyoxyethylene(6) oleyl ether, Raw Material 5, is referred to as a comparative product I; a product containing triglycerol monopalmitate, Raw Material 6, is referred to as a comparative product J; a product containing diglycerol monostearate, Raw Material 7, is referred to as a comparative product K; and a product containing diglycerol monopalmitate, Raw Material 8, is referred to as a comparative product L.

Test Example 2-1

The inventive products A to G were compared with the comparative products H to L. The evaluation methods and the evaluation criteria are as follows.

## 5 (State of Foam Formation)

The state of foam formation was observed by naked eyes. The evaluations were: those which are excellent were noted with ○, and those which are poor were noted with ×. The case where fine foams were formed when each of the aerosol was sprayed on the acrylic plate (length: 10 cm, width 10 cm) was  
10 judged “excellent,” and other cases were judged “poor.”

## (Foam Stability)

The maintenance of the formed foams was evaluated. The evaluations were: those which are excellent were noted with ○, and those which are poor  
15 were noted with ×. The case where the foams substantially did not disappear for at least one minute after spraying each aerosol was judged “excellent,” and other cases were judged “poor.”

## (Wash-Up Property)

20 The wash-up property during water rinsing was evaluated. The evaluations were: those which are excellent were noted with ○, and those which are poor were noted with ×. One female panelist was asked to apply an oily foundation, and the foundation was cleansed with each of the cosmetics, and the cosmetics were further washed away with water. The case where the foundation  
25 can be completely washed away without giving at least a greasy wash-away state

was judged “excellent,” and other cases were judged “poor.”

The results are shown in Table 7.

		State of Foam Formation	Foam Stability	Wash-Up Property
Example 2-1	Inventive Product A	○	○	○
	Inventive Product B	○	○	○
	Inventive Product C	○	○	○
	Inventive Product D	○	○	○
	Inventive Product E	○	○	○
	Inventive Product F	○	○	○
	Inventive Product G	○	○	○
Comparative Example 2-1	Comparative Product H	×	×	×
	Comparative Product I	×	×	×
	Comparative Product J	×	×	×
	Comparative Product K	×	×	×
	Comparative Product L	×	×	×

5

It is clear from Table 7 that the inventive products A to G formed stable fine foams, and gave excellent wash-up, whereas the comparative products H to L formed rough foams and gave much greasy texture during wash-up.

10

Test Example 2-2

Detergency for the inventive products A to G of Example 2-1 was compared to that of the above-mentioned conventional O/W emulsion type foamable aerosol cleansing cosmetics.

5 Ten female panelists were asked to apply an oily foundation. The results for the test for the degree of removal of the foundation are shown in Table 8.

Table 8

	Completely Removed	Slightly Removed	Not Removed
Inventive Product A	9	1	0
Inventive Product B	9	1	0
Inventive Product C	9	1	0
Example 2-1 Inventive Product D	9	1	0
Inventive Product E	9	1	0
Inventive Product F	9	1	0
Inventive Product G	9	1	0
O/W Emulsion type Aerosol	0	2	8

10 Table 8 clearly shows the followings: The inventive products A to G had very high detergency as compared to the conventional O/W emulsion type foamable aerosol cleansing cosmetics. Also, the inventive products had the advantages of not dripping when taking onto a palm, and easily applying to face, etc. as compared those of conventional pump-type cleansing cosmetics, so that their applicability was excellent. Further, the inventive products had  
15 substantially no greasy texture upon wash up, thereby showing excellent wash-up property.

Example 2-2 Suntan Oil

	Raw Materials	Weight (g)
1	Octamethyl Cyclotetrasiloxane	15.0
2	Decamethyl Cyclopentanesiloxane	15.0
3	C11-C15 Alkyl Benzoate	5.0
4	Octyl Paradimethylaminobenzoate	7.0
5	Oxybenzone	0.5
6	Olive Oil	11.0
7	Diglycerol Monomyristate	10.0
8	Stearyl Glycyrrhettinate	0.2
9	Cetyl Isooctanoate	21.3
10	Liquefied Petroleum Gas	10.0
11	Diethyl Ether	5.0

Raw Materials 1 to 9 were mixed and dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of a suntan oil.

5 Thereafter, the composition was packed into an aerosol can, a valve was set, and Raw Materials 10 and 11 were packed therein, to give an aerosol for the suntan oil. The resulting suntan oil was excellent in foam stability and applicability.

Example 2-3 Wipe-off type Cleansing Cosmetics

	Raw Materials	Weight (g)
1	Isopropyl Myristate	42.0
2	Oleyl Alcohol	40.0

3	2-Hydroxy-4-methoxybenzophenone	3.0
4	White Vaseline	2.0
5	Diglycerol Monolaurate	3.0
6	Liquefied Petroleum Gas	10.0

Raw Materials 1 to 5 were mixed and dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of the above-mentioned cosmetics. Thereafter, the composition was packed into an aerosol can, a valve was set, and Raw Material 6 was packed therein, to give an aerosol for the above-mentioned cosmetics. The resulting cosmetics were excellent in foam stability and applicability.

#### Example 2-4 Wash-Away type Cleansing Cosmetics

	Raw Materials	Weight (g)
1	Castor Oil	19.88
2	Oleyl Oleate	15.0
3	Diethyl Phthalate	10.0
4	Tocopherol Acetate	0.01
5	Perfume	0.1
6	Triglycerol Monomyristate	5.0
7	Diglycerol Monolaurate	9.0
8	Diglycerol Monocaprylate	5.0
9	Trichlorocarbanilide	1.0
10	Stearyl Glycyrrhetinate	0.01
11	Liquefied Petroleum Gas	35.0

Raw Materials 1 to 10 were mixed and dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of the above-mentioned cosmetics. Thereafter, the composition was packed into an aerosol can, a valve was set, and Raw Material 11 was packed therein, to give an aerosol for the above-mentioned cosmetics. The resulting cosmetics were excellent in foam stability and applicability.

Example 2-5 Baby Oil

	Raw Materials	Weight (g)
1	Squalane	34.44
2	Oleyl Alcohol	20.0
3	Polymethylsiloxane	5.0
4	Stearyl Glycyrrhettinate	0.01
5	Beeswax	0.5
6	Tocopherol Acetate	0.05
7	Polyoxyethylene(1) Lauryl Ether Phosphate	5.0
8	Diglycerol Monolaurate	5.0
9	Liquefied Petroleum Gas	20.0
10	Dimethyl Ether	10.0

10

Raw Materials 1 to 8 were mixed and dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of a baby oil. Thereafter, the composition was packed into an aerosol can, a valve was set, and Raw Materials 9 and 10 were packed therein, to give an aerosol for the baby oil.

The resulting baby oil was excellent in foam stability and applicability.

Example 2-6 Hair Oil

	Raw Materials	Weight (g)
1	Camellia Oil	30.0
2	Hexyl Laurate	22.0
3	Tocopherol Acetate	0.01
4	Perfume	0.5
5	Diglycerol Monocaprylate	2.0
6	Diglycerol Monomyristate	6.0
7	Liquefied Petroleum Gas	39.39
8	Nitrogen gas	0.1

5 Raw Materials 1 to 6 were mixed and dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of a hair oil. Thereafter, the composition was packed into an aerosol can, a valve was set, and Raw Materials 7 and 8 were packed therein, to give an aerosol for the hair oil. The resulting hair oil was excellent in foam stability and applicability.

10

Example 2-7 Foamy Massage Oil Cosmetics

	Raw Materials	Weight (g)
1	Liquid Paraffin	52.0
2	Polyoxyethylene Glyceryl Triisostearate	4.0
3	Squalane	10.0

4	Perfume	0.8
5	Macadamia Nut Oil	5.0
6	Isostearyl Glycyrrhetinate	0.1
7	Diglycerol Monocaprato	2.0
8	Diglycerol Monolaurate	6.0
9	Liquefied Petroleum Gas	20.0
10	Nitrogen gas	0.1

Raw Materials 1 to 8 were mixed and dissolved at 80°C with stirring, and cooled to ambient temperature, to give a composition for aerosol of the above-mentioned cosmetics. Thereafter, the composition was packed into an aerosol can, a valve was set, and Raw Materials 9 and 10 were packed therein, to give an aerosol for the above-mentioned cosmetics. The resulting cosmetics were excellent in foam stability and applicability.

#### INDUSTRIAL APPLICABILITY

The composition for aerosol provided by the present invention is useful as a cleansing agent for oily cosmetics, and the composition is excellent in detergency, applicability, wash-up property and foam stability.

CLAIMS

1. A composition for an oily foamable aerosol comprising:
  - a) a fat or oil that is in a liquid state during foaming; and
  - 5 b) a diglycerol or triglycerol fatty acid ester having a constituting fatty acid of 14 or less carbon atoms and a monoester content of 50% by weight or more.
- 10 2. The composition according to claim 1, further comprising a phospholipid.
3. An aerosol comprising the composition as defined in any one of claims 1 to 2 and a propellant.