



US00PP26751P2

(12) **United States Plant Patent**  
**Kientzler**

(10) **Patent No.:** **US PP26,751 P2**

(45) **Date of Patent:** **May 24, 2016**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED ‘KIRORUF113’**

(50) Latin Name: *Impatiens hawkeri*  
Varietal Denomination: **Kiroruf113**

(71) Applicant: **Ludwig Kientzler**, Gensingen (DE)

(72) Inventor: **Ludwig Kientzler**, Gensingen (DE)

(73) Assignee: **Visioplant**, Alajuela (CR)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/121,723**

(22) Filed: **Oct. 11, 2014**

(51) **Int. Cl.**  
**A01H 5/02** (2006.01)

(52) **U.S. Cl.**  
USPC ..... **Plt./318.3**

(58) **Field of Classification Search**  
USPC ..... Plt./318.3  
CPC ..... A01H 5/0261  
See application file for complete search history.

*Primary Examiner* — Kent L Bell

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of New Guinea *Impatiens* plant named ‘Kiroruf113’, characterized by its compact, upright and mounded plant habit; freely branching habit; dense and bushy growth habit; dark green-colored leaves; and numerous salmon peach-colored flowers with undulate margins and positioned above and beyond the foliar plane.

**1 Drawing Sheet**

**1**

Botanical designation: *Impatiens hawkeri*.  
Cultivar denomination: ‘KIRORUF113’.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of New Guinea *Impatiens* plant, botanically known as *Impatiens hawkeri*, and hereinafter referred to by the name ‘Kiroruf113’.

The new *Impatiens* plant is a product of a planned breeding program conducted by the Inventor in Gensingen, Germany. The objective of the breeding program was to develop new compact and freely branching *Impatiens* plants with numerous flowers and attractive leaf and flower colors.

The new *Impatiens* plant originated from a cross-pollination made by the Inventor in November, 2010, of a proprietary selection of *Impatiens hawkeri* identified as code number 09-307, not patented, as the female, or seed, parent with a proprietary selection of the *Impatiens hawkeri* identified as code number 09-029, not patented, as the male, or pollen, parent. The new *Impatiens* plant was discovered and selected by the Inventor as a flowering plant within the progeny of the stated cross-pollination in a controlled greenhouse environment in Gensingen, Germany in April, 2011.

Asexual reproduction of the new *Impatiens* plant by terminal cuttings propagated in a controlled greenhouse environment in Gensingen, Germany since May, 2011 has shown that the unique features of this new *Impatiens* plant are stable and reproduced true to type in successive generations of asexual reproduction.

**SUMMARY OF THE INVENTION**

Plants of the new *Impatiens* have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity, without, however, any variance in genotype.

**2**

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Kiroruf113’. These characteristics in combination distinguish ‘Kiroruf113’ as a new and distinct *Impatiens* plant:

1. Compact, upright and mounded plant habit.
2. Freely branching habit; dense and bushy growth habit.
3. Dark green-colored leaves.
4. Numerous salmon peach-colored flowers with undulate margins and positioned above and beyond the foliar plane.

Plants of the new *Impatiens* differ primarily from plants of the female parent selection in flower color. In addition, plants of the new *Impatiens* are more freely branching than plants of the female parent selection.

Plants of the new *Impatiens* differ primarily from plants of the male parent selection in growth habit as plants of the new *Impatiens* are more freely branching than plants of the male parent selection.

Plants of the new *Impatiens* can be compared to plants of New Guinea *Impatiens* ‘Visinfsalimp’, disclosed in U.S. Plant Pat. No. 20,971. In side-by-side comparisons conducted in Gensingen, Germany, plants of the new *Impatiens* differed primarily from plants of ‘Visinfsalimp’ in the following characteristics:

1. Flower petals of plants of the new *Impatiens* had undulate margins whereas flower petals of plants of ‘Visinfsalimp’ did not have undulate margins.
2. Plants of the new *Impatiens* and ‘Visinfsalimp’ differed in flower color as plants of ‘Visinfsalimp’ had salmon-colored flowers.

Plants of the new *Impatiens* can be compared to plants of New Guinea *Impatiens* ‘Visinfdsg’, disclosed in U.S. Plant Pat. No. 16,194. In side-by-side comparisons conducted in Gensingen, Germany, plants of the new *Impatiens* differed primarily from plants of ‘Visinfdsg’ in the following characteristics:

1. Flower petals of plants of the new *Impatiens* had undulate margins whereas flower petals of plants of ‘Visinfdsg’ did not have undulate margins.

2. Flowers of plants of the new *Impatiens* were lighter in color than flowers of plants of 'Visinfdsq'.

#### BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs illustrate the overall appearance of the new *Impatiens* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Impatiens* plant.

The photograph at the bottom of the sheet comprises a side perspective view of a typical flowering plant of 'Kioruf113' grown in a container.

The photograph at the top of the sheet is a close-up view of a typical flowering plant of 'Kioruf113'.

#### DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations and measurements describe plants grown in Bonsall, Calif. during the summer in 11.5-cm containers in a polypropylene-covered shadehouse and under cultural practices typical of commercial New Guinea *Impatiens* production. During the production of the plants, day temperatures ranged from 24° C. to 27° C., night temperatures averaged 18° C. and light levels ranged from 2,500 to 4,000 foot-candles. Measurements and numerical values represent averages for typical flowering plants which were pinched one time and had been growing for four weeks after planting rooted young plants when the photographs and detailed description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2007 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens hawkeri* 'Kioruf113'.

Parentage:

*Female, or seed, parent.*—Proprietary selection of *Impatiens hawkeri* identified as code number 09-307, not patented.

*Male, or pollen parent.*—Proprietary selection of *Impatiens hawkeri* identified as code number 09-029, not patented.

Propagation:

*Type.*—Vegetative cuttings.

*Time to initiate roots, summer.*—About 14 days at temperatures about 20° C. to 22° C.

*Time to initiate roots, winter.*—About 18 days at temperatures about 19° C. to 21° C.

*Time to produce a rooted young plant, summer.*—About 18 days at temperatures about 20° C. to 22° C.

*Time to produce a rooted young plant, winter.*—About 21 days at temperatures about 19° C. to 21° C.

*Root description.*—Fibrous, medium in thickness; white in color.

*Rooting habit.*—Freely branching; medium density.

Plant description:

*Plant habit.*—Compact, upright and mounded plant habit.

*Growth and branching habit.*—Rapid growth rate; vigorous and freely branching habit; about ten lateral branches developing at the base; dense and bushy growth habit; pinching, that is, removal of the terminal apices, is typically not required, but will enhance branching.

*Plant height.*—About 18.8 cm.

*Plant diameter or spread.*—About 30.5 cm.

*Lateral branches.*—Length: About 15.5 cm. Diameter: About 6 mm. Internode length: About 4.8 cm. Texture: Smooth, glabrous. Color: Close to 187B.

Leaf description:

*Arrangement.*—Opposite or in whorls, simple.

*Length.*—About 9.6 cm.

*Width.*—About 3.6 cm.

*Shape.*—Elliptical.

*Apex.*—Acuminate.

*Base.*—Attenuate.

*Margin.*—Serrulate with ciliation.

*Texture, upper and lower surfaces.*—Smooth, glabrous.

*Venation pattern.*—Pinnate.

*Color.*—Developing leaves, upper surface: Close to N137B. Developing leaves, lower surface: Close to 147B. Fully expanded leaves, upper surface: Close to N137A; venation, close to 182B. Fully expanded leaves, lower surface: Close to 147B; venation, close to 185A.

*Petioles.*—Length: About 2 cm. Diameter: About 4 mm.

Texture, upper and lower surfaces: Smooth, glabrous.

Color, upper and lower surfaces: Close to 185A.

Flower description:

*Flower type and flowering habit.*—Single axillary flowers; freely flowering habit with usually about eight flowers and flower buds per lateral branch; flowers positioned above and beyond the foliar plane and typically face upright and outwardly.

*Flower longevity.*—Flowers last about one week under greenhouse conditions; petals self-cleaning, gynoecium persistent.

*Fragrance.*—None detected.

*Natural flowering season.*—Year-round under greenhouse conditions; in the garden, flowering from spring until fall in California; plants begin flowering about four weeks after planting young plants.

*Flower size.*—Diameter: About 4.8 cm. Depth: About 3.4 cm.

*Flower buds.*—Length: About 2.3 cm. Diameter: About 1.3 cm. Shape: Ovoid. Color: Close to 54C to 54D.

*Petals.*—Quantity and arrangement: Five petals arranged in a single whorl. Length, banner petal: About 2.8 cm. Length, lateral petals: About 2.5 cm. Length, lower petals: About 2.5 cm. Width, banner petal: About 3.5 cm. Width, lateral petals: About 2.9 cm. Width, lower petals: About 3.1 cm. Shape: Cordate. Apex: Emarginate. Base: Attenuate. Margin: Entire with irregular sinuate edges; undulate. Texture, upper and lower surfaces: Smooth, glabrous; velvety. Color: When opening, upper surface: Close to 54C; towards the margins, close to 52C. When opening, lower surface: Close to 55B. Fully opened, upper surface: Center, close to 54C; towards the margins, close to 52B to 52C; towards the base, close to 60B; colors do not fade with development. Fully opened, lower surface: Close to 55B.

*Sepals.*—Quantity and arrangement: Three arranged in a single whorl; one modified into an elongated spur. Length: About 7 mm. Width: About 5 mm. Shape: Elliptical. Apex: Acuminate. Base: Obtuse. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: Close to 59B. Spur length: About 3.4 cm. Spur diameter: At

flower, about 2 mm; at apex, less than 1 mm. Spur texture: Smooth, glabrous. Spur color: Close to 59B.

*Peduncles*.—Length: About 4.2 cm. Diameter: About 2 mm. Angle: About 25° to 35° from vertical. Strength: Strong. Texture: Smooth, glabrous. Color: Close to 59B.

*Reproductive organs*.—Stamens: Quantity: Five fused at anthers; filaments free. Anther size: About 3 mm by 5 mm. Anther color: Close to 160D. Pollen amount: Moderate. Pollen color: Close to 158A. Pistils: Quantity per flower: One. Pistil length: About 5 mm. Stigma shape: Round. Stigma color: Close to 145D. Style color: Close to 151B. Ovary color: Close to 151B.

*Seeds and fruit*.—Seed and fruit development have not been observed on plants of the new *Impatiens*.

Disease & pest resistance: Plants of the new *Impatiens* have not been observed to be resistant to pathogens and pests common to New Guinea *Impatiens* plants.

Temperature tolerance: Plants of the new *Impatiens* have been observed to tolerate temperatures from about 16° C. to about 35° C.

It is claimed:

1. A new and distinct New Guinea *Impatiens* plant named 'Kiroruf113' as illustrated and described.

\* \* \* \* \*

