

- (51) International Patent Classification:

C07K 7/06 (2006.01)

A61P 17/00 (2006.01)

A61K 38/08 (2019.01)

A61P 11/02 (2006.01)

A61K 8/64 (2006.01)

A61P 11/00 (2006.01)

A61P 29/00 (2006.01)

A61P 1/04 (2006.01)

A61P 17/06 (2006.01)

A61P 33/12 (2006.01)

A61P 17/10 (2006.01)

A61Q 19/00 (2006.01)

A61P 17/02 (2006.01)
- (21) International Application Number:

PCT/CN2020/133438
- (22) International Filing Date:

02 December 2020 (02.12.2020)
- (25) Filing Language:

English
- (26) Publication Language:


English
- (30) Priority Data:

PCT/CN2019/122391

02 December 2019 (02.12.2019) CN
- (71) Applicant: JIANGYIN USUN BIOCHEMICAL TECHNOLOGY CO., LTD. [CN/CN]; Bldg. C3, No. 6 West Dongsheng Rd, Jiangyin, Jiangsu 214437 (CN).
- (72) Inventors: JANSON, Jan-Christer; Husargatan 3, S-751 23 Uppsala (SE). GU, Ming; B13, No. 2, Dongsheng West Road, Jiangyin, Jiangsu 214437 (CN). SONG, Mao-qian; B13, No. 2, Dongsheng West Road, Jiangyin, Jiangsu 214437 (CN). SAMUELSSON, Bengt Ingemar; Bio-medicum, 9A, Floor 9, Solnavagen 9, S-171 65 Solna (SE).
- (74) Agent: WU, FENG & ZHANG; 3 FL, Building 2, Yard 3, FengXiuZhongLu Road, Haidian District, Beijing 100094 (CN).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Published:

— with international search report (Art. 21(3))

(54) Title: NEW CONJUGATES OF PEPTIDES AND POLYSACCHARIDE



(57) Abstract: Provided is a conjugate formed between one or more linear polysaccharide chains and a peptide selected from one or more of the peptide components (a), (b), (c) or (d) as defined below: (a) a peptide component of formula (I), A-Q-B I wherein Q represents a structural fragment of formula (II), wherein: the squiggly lines and m have meanings given in the description and A and B have meanings given in the description, but may represent a peptide component of the amino acid sequence:[W-Lys-X1-Ser-U-X2-Y] n-W-Lys-X1-Ser-U-X2-Y--- (SEQ ID No: 3), wherein the dashed line, n, W, X1, U, X2 and Y have meanings given in the description; (b) a peptide component of the amino acid sequence: [Ala-Lys-X1-Ser-U-X2-Y] p-Ala-Lys-X1-Ser-U-X1-Y-G (SEQ ID No: 4) wherein p, G, X1, U, X2 and Y have meanings given in the description; (c) a peptide component of the amino acid sequence: W-Lys-X1-Ser-U-X2-Y-G (SEQ ID No: 5), wherein W, X1, U, X2, Y and G have meanings given in the description; or (d) a peptide component of the amino acid sequence: K-W1-Lys-X1-Ser-U1-X2-Y1-I-J (SEQ ID No: 6), wherein K, W1, U1, Y1, I, J, X1 and X2 have meanings as well as regioisomers, stereoisomers, and pharmaceutically-or cosmetically-acceptable salts of said conjugates, which conjugates are particularly useful in the treatment of conditions characterised by inflammation, including wounds, burns, and disorders of the mucosa, such as anorectal diseases, inflammatory bowel diseases, gynaecological diseases and dental diseases. Preferred linear polysaccharides include hyaluronic acid.

NEW CONJUGATES OF PEPTIDES AND POLYSACCHARIDE

Field of the Invention

- 5 This invention relates to new peptide conjugates, the use of such conjugates in human medicine, and to pharmaceutical compositions comprising them. In particular, the invention relates to the use of those conjugates and compositions in the treatment of e.g. inflammation.

10 Background and Prior Art

Inflammation is typically characterized as a localised tissue response to e.g. invasion of microorganisms, certain antigens, damaged cells or physical and/or chemical factors. The inflammatory response is normally a protective mechanism which serves to
15 destroy, dilute or sequester both the injurious agent and the injured tissue, as well as to initiate tissue healing.

Inflammation may result from physical trauma, infection, some chronic diseases (e.g. psoriasis and autoimmune diseases, such as rheumatoid arthritis) and/or chemical
20 and/or physiological reactions to external stimuli (e.g. as part of an allergic response). A complex series of events may be involved, in which inflammatory mediators increase blood flow and dilation of local blood vessels, resulting in redness and heat, the exudation of fluids, often resulting in localised swelling, leukocytic migration into the inflamed area, and pain.

25 Many conditions/disorders are characterized by, and/or are caused by, abnormal, tissue-damaging inflammation. Such conditions are typically characterized by activation of immune defence mechanisms, resulting in an effect that is more harmful than beneficial to the host, and are generally associated with varying degrees of tissue
30 redness or hyperemia, swelling, hyperthermia, pain, itching, cell death, tissue destruction, cell proliferation and/or loss of function. Examples include inflammatory bowel diseases, rheumatoid arthritis, multiple sclerosis, psoriasis, glomerulonephritis and transplant rejection.

35 Typically, a complex series of events results in inflammatory changes such as increased blood flow through dilation of local blood vessels, resulting in redness and heat, the extravasation of leukocytes and plasma, often resulting in localised swelling, activation of sensory nerves (resulting in pain in some tissues) and loss of function.

These inflammatory changes are triggered by a cascade of cellular and biochemical events involving cells like neutrophils, monocytes, macrophages and lymphocytes together with inflammatory mediators such as vasoactive amines, cytokines, complement factors and reactive oxygen species.

5

Amongst other things, inflammation plays a key role in the wound healing process. Wounds and burns can therefore be classified as conditions with which inflammation is associated. Traditional thinking in the art is that anti-inflammatory drugs should not be applied directly to open wounds, as this would be detrimental to the progress of wound healing.

10

Fibrosis is defined by the excessive accumulation of fibrous connective tissue (components of the extracellular matrix (ECM) such as collagen and fibronectin) in and around inflamed or damaged tissue. Although collagen deposition is typically a reversible part of wound healing, it can often evolve into a progressively irreversible fibrotic response if tissue injury is severe, or if the wound-healing response itself becomes dysregulated. Furthermore, fibrogenesis is known to be a major cause of morbidity and mortality in many chronic inflammatory diseases, as well as end-stage liver disease, kidney disease, idiopathic pulmonary fibrosis (IPF) and heart failure. It is also a pathological feature of many chronic autoimmune diseases, such as scleroderma, rheumatoid arthritis, Crohn's disease, ulcerative colitis, myelofibrosis and systemic lupus erythematosus. Fibrosis may also influence the pathogenesis of many progressive myopathies, metastasis and graft rejection.

15

20

25

Mussel adhesive protein (MAP), also known as *Mytilus edulis* foot protein (mefp), is a protein that is secreted by marine shellfish species, such as *Mytilus edulis*, *Mytilus coruscus* and *Perna viridis*. Eleven identified separate adhesive protein subtypes have been derived from mussels, including the collagens pre-COL-P, pre-COL-D and pre-COL-NG; the mussel feet matrix proteins PTMP (proximal thread matrix protein) and DTMP (distal thread matrix protein); and mfp proteins mfp-2 (sometimes referred to as "mefp-2", hereinafter used interchangeably), mfp-3/mefp-3, mfp-4/mefp-4, mfp-5/mefp-5, mfp-6/mefp-6 and, most preferably mfp-1/mefp-1 (see, for example, Zhu *et al.*, *Advances in Marine Science*, **2014**, 32, 560-568 and Gao *et al.*, *Journal of Anhui Agr. Sci.*, **2011**, 39, 19860-19862).

30

35

A significant portion of mefp-1 consists of 70 to 90 tandem repeats of the decapeptide: Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 1; see Waite, *Int. J. Adhesion and Adhesives*, **1987**, 7, 9-14). This decapeptide sequence may be isolated

as a low molecular weight derivative of naturally-occurring MAPs, or may be synthesized, for example as described by Yamamoto in *J. Chem. Soc., Perkin Trans.*, **1987**, 1, 613-618. See also Dalsin *et al.*, *J. Am. Chem. Soc.*, **2003**, 125, 4253-4258.

- 5 Analogues of the decapeptide, notably Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 2) have also been disclosed. See, for example, US 5,616,311 and WO 96/39128.

The use of lysine amino acid residues to prepare multi antigen peptides has been disclosed in, for example, Tam, *Proc. Natl. Acad., Sci. USA*, **1988**, 85, 5409-5413, Rao *et al.*, *J. Am. Chem. Soc.*, **1994**, 116, 6975-6976, US 5,229,490 and WO 2010/038220.

The use of peptide-based scaffolds as drug delivery vehicles has been disclosed. See, for example, Brokx *et al.*, *J. Control. Release*, **2002**, 78, 115-123.

15 Hyaluronic acid (HA) and sodium hyaluronate are widely used to prepare biomaterials for tissue engineering (see, for example, Khunmanee *et al.*, *J. Tissue Eng.*, **2017**, 8, 1-16). HA is a naturally occurring polysaccharide that is distributed widely throughout the human body in connective, epithelial, and neural tissues. It is a major component of the synovial fluid and is one of the fluid's main lubricating components. It is also an important component of articular cartilage and plays a part in the resistance of cartilage to compression. As a major component of skin, it is involved in tissue repair and also contributes to tissue hydrodynamics, movement and proliferation of cells. As a main component of the extracellular matrix, it has a key role in tissue regeneration, the inflammatory response and angiogenesis (phases of skin wound repair).

HA can form a gel with water. This makes it useful in osteoarthritis as an intra-articular injection, as well as in skin treatments as a dermal filler for treating facial wrinkles in cosmetic surgery. HA is approved by the US Food and Drug Administration (FDA) for this purpose, where treatment involves repeated injection at 6 to 12-month intervals using either a hypodermic needle or a micro-cannula.

The cross-linking of HA is required to prevent biodegradation from free radicals and enzymes and to increase the HA filler duration. Presently, chemical cross-linkers such as 1,4-butanediol diglycidyl ether (BDDE) are employed, but the presence of these cross-linkers in the crosslinked HA product is known to cause problems. For example, BDDE has been found to be mutagenic in the *Drosophila* model organism (Foureman

et al., *Environ. Mol. Mutagen.*, **1994**, 23, 51-63). Moreover, BDDE is a suspected carcinogen (WO 2017/076495).

There is a clear need for new and/or improved medicines that may be used in the treatment of inflammation and conditions characterised thereby.

Disclosure of the Invention

According to a first aspect of the invention, there is provided a conjugate formed between one or more linear polysaccharide chains and a peptide component, preferably selected from one or more of the peptide components (a), (b), (c) or (d) as defined below:


(a) a peptide component of formula I,

A-Q-B I

wherein:

A and B independently represent Z or A¹-Q¹-B¹;

Q represents a structural fragment of formula II,




wherein:

the squiggly lines represent points of attachment of Q to A and/or B; and m represents an integer 1 to 4;

A¹ and B¹ independently represent Z or A²-Q²-B²;

A² and B² independently represent Z or Z-Q³-Z;


Q¹, Q² and Q³ independently represent structural fragments of formula III,



wherein:

the squiggly lines adjacent to the NH groups represent the points of attachment of Q¹, Q² and Q³ to A¹ and/or B¹, A² and/or B², and Z, respectively; and the squiggly line adjacent to the O atom represents the point of attachment of Q¹, Q² and Q³ to Q, Q¹ and Q², respectively; and m is as defined above;

on each occasion that it is employed, Z represents a peptide component of the amino acid sequence:



wherein:

the dashed line represents the point of attachment of Z to the rest of the molecule;

n represents 0 or an integer 1 to 4; and,

on each occasion that they are employed:

W represents a 1 or 2 amino acid sequence, in which the amino acids are selected from one or more of the group Lys, Ala, DOPA and a 3,4-dihydrocinnamic acid (HCA) residue, provided that, when present, the HCA residue is located at the N-terminus of the peptide sequence Z;


X¹ represents Pro, Hyp or diHyp;

U represents Tyr or, DOPA;

X² represents Ser, Pro, Hyp or diHyp; and

Y represents a 1 to 5 (e.g. a 1 to 4) amino acid sequence, in which the amino acids are selected from one or more of the group Lys, Ala, Pro, Hyp, diHyp, Thr, DOPA and Tyr; or

(b) a peptide component of the amino acid sequence:



wherein


p represents an integer 1 to 4;

G may be absent (in which case Y is the C-terminal amino acid) or G may represent DOPA or dopamine (or, more properly, 'a dopamine fragment'); and

X¹, U, X² and Y are as defined above;

or

(c) a peptide component of the amino acid sequence:



wherein W, X¹, U, X², Y and G are as defined above; or

(d) a peptide component of the amino acid sequence:

K-W¹-Lys-X¹-Ser-U¹-X²-Y¹-I-J (SEQ ID No: 6)

5

wherein K represents an optional N-terminal HCA group;

W¹ may be absent (in which case Lys is the N-terminal amino acid) or W¹ may represent a 1 or 2 amino acid sequence, in which the amino acids are selected from one or more of the group Ser, Lys, Ala and DOPA;

10 U¹ represents Tyr, DOPA or a single bond (i.e. is absent);

Y¹ represents a single bond (i.e. is absent) or represents Y;

I represents Pro, Hyp, diHyp, Thr, DOPA or Tyr; and

J represents Lys or is absent (in which case I represents the C-terminal amino acid); and

15 X¹, X² and Y are as defined above, as well as regioisomers, stereoisomers, and pharmaceutically- or cosmetically-acceptable salts of said conjugates, which conjugates, regioisomers, stereoisomers and salts are referred to together hereinafter as 'the conjugates of the invention'.

20 It is preferred that the one or more linear polysaccharide chains comprises hyaluronic acid (HA).

Conjugates of the invention that may be mentioned include those in which:

the peptide components are selected from one or more of (a), (b) and/or (c);

25 W represents a 1 or 2 amino acid sequence, in which the amino acids are selected from one or more of the group Lys, Ala and DOPA;

X¹ represents Pro;

X² represents Ser, Pro or Hyp;

Y represents a 1 to 5 (e.g. a 1 to 4) amino acid sequence, in which the amino acids

30 are selected from one or more of the group Lys, Ala, Pro, Hyp, Thr, DOPA and Tyr.

Preferred conjugates of the invention include those in which:

X¹ represents Hyp or, more preferably, Pro;

X² represents Pro or, more preferably, Hyp;

35 W represents HCA, HCA-Ala-, preferably Ala or Lys-Ala or, more preferably DOPA or DOPA-Ala-; and/or

Y represents a 5, preferably a 3 or, more preferably, a 4 amino acid sequence, in which the amino acids are selected from one or more of the group Lys, Ala, Hyp, Thr, DOPA and Tyr.

- 5 More preferably, conjugates of the invention also include those in which Y represents a 4 amino acid sequence selected from the group -Pro-Y¹-Y²-Lys- or, more preferably, -Hyp-Y¹-Y²-Lys- and -Thr-Y¹-Y²-Lys-, wherein Y¹ and Y² are each independently selected from the group Pro or, more preferably, Ala, Hyp, Thr, DOPA and Tyr.
- 10 Wherein Y represents a 4 amino acid sequence, preferred conjugates of the invention include those in which the amino acid sequence defined by Y is selected from the group:
 -Pro-Thr-DOPA-Lys-;
 -Pro-Thr-Tyr-Lys-;
 -Thr-Tyr-Pro-Lys-; and
 15 -Thr-DOPA-Pro-Lys-; and, more preferably,
 -Hyp-Thr-Tyr-Lys-;
 -Hyp-Thr-DOPA-Lys-;
 -Hyp-Thr-Ala-Lys-;
 -Thr-Tyr-Hyp-Lys-;
 20 -Thr-DOPA-Hyp-Lys-; and
 -Thr-Ala-Hyp-Lys-.

When Y represents a 2 amino acid sequence, preferred conjugates of the invention include those in which the amino acid sequence defined by Y is selected from the group
 25 -Hyp-Thr-, -Thr-Tyr-, -Pro-Thr- and -Thr-DOPA-.

Other preferred conjugates of the invention that may be mentioned include those in which the amino acid sequence defined by Y is selected from -Thr-Tyr-Lys-, -Tyr-Pro-Lys-, -DOPA-Pro-Lys-, -Hyp-Thr-Tyr-, -Hyp-Thr-Tyr-Hyp-Lys- and, more preferably,
 30 the groups -Thr-Tyr-Hyp-Lys-DOPA- and -Hyp-Thr-DOPA-.

When conjugates of the invention comprise peptide components of formula I (as defined under (a) above), those that may be mentioned are those wherein m represents 1, 3 or, more preferably 4, such that one or more of Q, Q¹, Q² and Q³
 35 represent Lys or, more properly, 'a Lys fragment', in accordance with what are defined above as 'the structural fragments of formulae II and III' (as appropriate).

On each occasion that they are employed, Q, Q¹, Q² and Q³ may each be attached to zero, one or two Z groups.

In this respect, preferred conjugates of the invention include those in which, in the peptide component of formula I:

one of A or B represents Z and the other represents A¹-Q¹-B¹; or, more preferably, A and B both represent Z, or both represent A¹-Q¹-B¹, in which, in each case, Q¹ preferably represents a Lys fragment and Z is as hereinbefore defined.

10

Further preferred conjugates of the invention also include those in which:

one of A¹ and B¹ represents Z and the other represents A²-Q²-B²; or, more preferably, A¹ and B¹ both represent Z, or both represent A²-Q²-B²,

in which, in each case, Q² preferably represents a Lys fragment, and Z is as hereinbefore defined.

15

Further preferred conjugates of the invention also include those in which:

one of A² and B² represents Z and the other represents Z-Q³-Z; or, more preferably, A² and B² both represent Z, or both represent Z-Q³-Z,

in which, in each case, Q³ preferably represents a Lys fragment, and Z is as hereinbefore defined.

20


More preferred conjugates of the invention include those in which A² and B² both represent Z.

25

Peptide components of conjugates of the invention that may be mentioned include those in which n is 0, 1 or 4, or, more preferably, n is 0.

When conjugates of the invention comprises peptide components as defined under (b) or (c) above, the terms 'dopamine' and 'dopamine fragment' that may be defined by the substituent G refer to a structural fragment of formula IV,

30



wherein the squiggly line represents the point of attachment to Y.

35

Preferred values of p in peptide components as defined under (b) above are, in ascending order of preference 2, 3, 1 and 4.

Particular conjugates of the invention comprising peptide components as defined under (b) above that may be mentioned are those where G is absent and, in this respect, preferred peptide components include those of the amino acid sequence:

Ala-Lys-Pro-Ser-Tyr-Ser-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Ser-Hyp-Thr-Tyr-Lys (SEQ ID No: 7);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 8);

Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 9);

Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 10);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 11);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 12);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 13); and

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 14).

Conjugates of the invention that may be mentioned include those in which:

U represents Tyr; and/or

W represents Ala, and, in this respect, conjugates of the invention comprising: peptide components of formula I as defined under (a) above that may be mentioned include those wherein Z is selected from the group:

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 2);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 1);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 15);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 16);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys-DOPA--- (SEQ ID No: 17);

Ala-Lys-Pro-Ser-Tyr-Ser-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Ser-Hyp-Thr-Tyr-Lys--
(SEQ ID No: 18); and

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-
Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-

5 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 19); and

peptide components as defined under (c) above that may be mentioned include those
of the amino acid sequence:

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys-DOPA (SEQ ID No: 20);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys-Dopamine (SEQ ID No: 21);

10 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-Dopamine (SEQ ID No: 22);

Ala-Lys-Pro-Ser-Tyr-Pro-Pro-Thr-DOPA-Lys (SEQ ID No: 23);

Ala-Lys-Pro-Ser-Tyr-Pro-Thr-Tyr-Pro-Lys (SEQ ID No: 24);

Ala-Lys-Pro-Ser-Tyr-Pro-Thr-DOPA-Pro-Lys (SEQ ID No: 25);

Ala-Lys-Pro-Ser-Tyr-Pro-Hyp-Thr-Tyr-Lys (SEQ ID No: 26);

15 Ala-Lys-Pro-Ser-Tyr-Pro-Hyp-Thr-DOPA-Lys (SEQ ID No: 27);

Ala-Lys-Pro-Ser-Tyr-Hyp-Pro-Thr-Tyr-Lys (SEQ ID No: 28);

Ala-Lys-Pro-Ser-Tyr-Hyp-Pro-Thr-DOPA-Lys (SEQ ID No: 29);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 30);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 31);

20 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-DOPA (SEQ ID No: 32);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-DOPA (SEQ ID No: 33);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Dopamine (SEQ ID No: 34);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys-DOPA (SEQ ID No: 35); and

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys-Dopamine (SEQ ID No: 36).

25

Conjugates of the invention that may be mentioned include those in which:

U represents Tyr; and/or

W represents Lys-Ala-, and, in this respect, conjugates of the invention comprising:

peptide components of formula I as defined under (a) above that may be mentioned

30 include those wherein Z is selected from the group:

Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr--- (SEQ ID No: 37);

Lys-Ala-Lys-Hyp-Ser-Tyr-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 38); and, more preferably,

Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 39); and

peptide components as defined under (c) above that may be mentioned include those

35 of the amino acid sequence:

Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA (SEQ ID No: 40);

Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Dopamine (SEQ ID No: 41);

Lys-Ala-Lys-Hyp-Ser-Tyr-Hyp-Hyp-Thr-DOPA (SEQ ID No: 42);

Lys-Ala-Lys-Hyp-Ser-Tyr-Hyp-Hyp-Thr-Tyr (SEQ ID No: 43); and
Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr (SEQ ID No: 44).

Further conjugates of the invention that may be mentioned include those in which:

5 U represents Tyr; and/or

W represents HCA, HCA-Ala- or, more preferably, DOPA or DOPA-Ala-, and, in this respect, conjugates of the invention comprising:

peptide components of formula I as defined under (a) above that may be mentioned include those wherein Z is selected from the group:

10 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 45);

HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 46);

HCA-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Ala-Lys--- (SEQ ID No: 47);

HCA-Lys-Pro-Ser-Tyr-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 48);

and, more preferably, wherein Z is selected from the group:

15 DOPA-Lys-Pro-Ser-Tyr-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 49);

DOPA-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Ala-Lys--- (SEQ ID No: 50);

DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 51); and

DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 52); and

peptide components as defined under (c) above that may be mentioned include those

20 of the amino acid sequence:

DOPA-Lys-Pro-Ser-Tyr-Hyp-Thr-Ala-Hyp-Lys (SEQ ID No: 53);

DOPA-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Ala-Lys (SEQ ID No: 54);

DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 55);

DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 56);

25 DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 57);

DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 58);

HCA-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Ala-Lys (SEQ ID No: 59);

HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 60);

HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 61);

30 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 62);

HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 63); and

HCA-Lys-Pro-Ser-Tyr-Hyp-Thr-Ala-Hyp-Lys (SEQ ID No: 64).

Other conjugates of the invention that may be mentioned include those in which:

35 U represents DOPA; and/or

W represents Ala or Lys-Ala-, and, in this respect, conjugates of the invention comprising peptide components of formula I as defined under (a) above that may be mentioned include those wherein Z is selected from the group:

U represents DOPA; and/or

W represents HCA, HCA-Ala- or, more preferably, DOPA or DOPA-Ala-, and, in this respect, particular conjugates of the invention comprising peptide components of formula I as defined under (a) above that may be mentioned include those wherein Z

5 is selected from the group:

HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 94);

HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 95);

HCA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 96);

HCA-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Ala-Lys--- (SEQ ID No: 97);

10 and, more preferably, wherein Z is selected from the group:

DOPA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 98);

DOPA-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Ala-Lys--- (SEQ ID No: 99);

DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 100);

DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 101); and

15 DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 102); and

peptide components as defined under (c) above that may be mentioned include those of the amino acid sequence:

DOPA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys (SEQ ID No: 103);

DOPA-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Ala-Lys (SEQ ID No: 104);

20 DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 105);

DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 106);

DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 107).

HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 108);

HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 109);

25 HCA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys (SEQ ID No: 110);

HCA-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Ala-Lys (SEQ ID No: 111);

DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 112);

HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 113); and

HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 114).

30

When conjugates of the invention comprise a peptide component of formula I as defined under (a) above, those that may be mentioned include those in which, in the peptide component of formula I:

A and B both represent Z;

35 one, or preferably both, Z groups represent:

HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 45),

HCA-Lys-Pro-Ser-Tyr-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 48),

DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 51),

- Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 65),
 Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 67)
 HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 94),
 DOPA-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Ala-Lys--- (SEQ ID No: 99),
 5 DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 100),
 or, more preferably, one, or preferably both, Z groups represent:
 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 15), or
 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 16);
 or, even more preferably, one, or preferably both, Z groups represent:
 10 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 2), or
 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 1),
 and
 Q represents a Lys fragment.
- 15 Further conjugates of the invention that comprise a peptide component of formula I as
 defined under (a) above that may be mentioned include those in which, in the peptide
 component of formula I:
 A and B both represent $A^1-Q^1-B^1$;
 A^1 and B^1 both represent Z;
 20 one, or preferably both, Z groups represent:
 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 15),
 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 16),
 Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 39),
 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 45),
 25 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 46),
 DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 51),
 Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 65),
 Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 66),
 Lys-Ala-Lys-Hyp-Ser-Tyr-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 38),
 30 HCA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 96),
 DOPA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 98),
 DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 100),
 or, more preferably, one, or preferably both, Z groups represent:
 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 2) or
 35 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 1); and
 Q^1 represents a Lys fragment.

Further conjugates of the invention that comprise a peptide component of formula I as defined under (a) above that may be mentioned include those in which, in the peptide component of formula I:

A and B both represent $A^1-Q^1-B^1$;

5 A^1 and B^1 both represent $A^2-Q^2-B^2$;

A^2 and B^2 both represent Z;

one, or preferably both, Z groups represent:

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 15),

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 16),

10 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 45),

DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 51),

Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 66),

DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 100),

or, more preferably, one, or preferably both, Z groups represent:

15 DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 52),

or, even more preferably, one, or preferably both, Z groups represent:

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 2), or

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 1); and

Q^1 and Q^2 both represent Lys fragments.

20

Further conjugates of the invention that comprise a peptide component of formula I as defined under (a) above that may be mentioned include those in which, in the peptide component of formula I:

A and B both represent $A^1-Q^1-B^1$;

25 A^1 and B^1 both represent $A^2-Q^2-B^2$;

A^2 and B^2 both represent $Z-Q^3-Z$;

one, or preferably both, Z groups represent:

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 2), or

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 1); and

30 Q^1 , Q^2 and Q^3 all represent Lys fragments.

When conjugates of the invention comprise a peptide component as defined under (d) above, those that may be mentioned include those in which:

W^1 represents Ala or Ser, or is absent (in which case, Lys is the N-terminal amino acid);

35 X^2 represents Pro, Hyp or diHyp; and/or

when K is not present, W^1 represents Ala or is absent and J represents Lys, then I represents Pro, Hyp, diHyp or Thr (i.e. I does not represent DOPA or Tyr).

Preferred conjugates of the invention comprising a peptide component as defined under (d) above include those in which:

U¹ represents DOPA or, more preferably Tyr;

X¹ represents Hyp or, more preferably, Pro;

5 X² represents diHyp or, more preferably, Hyp; and/or

Y¹ represents a 1 to 4, such as a 3, 1 or, preferably, 2 amino acid sequence, in which the amino acids are selected from the group Pro, Hyp, Thr, DOPA and Tyr.

10 Peptide components as defined under (d) above that may be mentioned include those in which W¹ represents Ser.

However, more preferred peptide components as defined under (d) above include those in which W¹ is absent or, more preferably, W¹ represents Ala.

15 Preferred peptide components as defined under (d) above also include those in which J represents Lys.

More preferably, peptide components as defined under (d) above also include those in which I represents DOPA or Tyr, more preferably Pro or, especially, Hyp.

20 Preferred peptide components as defined under (d) above also include those in which, when J represents Lys, Z represents DOPA or Tyr, more preferably Pro or, especially, Hyp.

25 Further preferred peptide components as defined under (d) above include those in which the amino acids in the sequence defined by Y¹ are selected from Pro, preferably DOPA, more preferably Hyp, Thr and Tyr.

30 Especially preferred peptide components as defined under (d) above include those in which, in the sequence defined by Y¹:
the amino acid DOPA, preferably Thr or, more preferably, Tyr is linked to I; and/or
the amino acid Pro, or more preferably Hyp or Thr is linked to X².

35 Preferred values of Y¹ in peptide components as defined under (d) above include, when it is a 3-membered amino acid sequence, -Hyp-Thr-Tyr or, more preferably -Hyp-Thr-DOPA-, and, when it is a 2-membered amino acid sequence, -Thr-Tyr- or, more preferably, -Thr-DOPA-, or -Pro-Thr- or, more preferably, -Hyp-Thr-.

Particular conjugates of the invention comprising peptide components as defined under (d) above that may be mentioned include those in which K is absent.

In this respect, peptide components as defined under (d) above include those comprising the amino acid sequence:

Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 115);

Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 116);

Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 117);

Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 118);

Ser-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 119);

Ser-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 120);

Ser-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 121);

Ser-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 122);

More preferred conjugates of the invention comprising peptide components as defined under (d) above include those comprising the amino acid sequence:

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 123);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 124); more preferably those comprising the amino acid sequence:

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 125); and particularly those comprising the amino acid sequence:

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 126).

Further conjugates of the invention comprising peptide components as defined under (d) above that may be mentioned include those in which J is absent, such as those comprising the amino acid sequence:

Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Hyp (SEQ ID No: 127);

Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Hyp (SEQ ID No: 128);

Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp (SEQ ID No: 129);

Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp (SEQ ID No: 130);

Ser-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr (SEQ ID No: 131);

Ser-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp (SEQ ID No: 132);

Ser-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp (SEQ ID No: 133);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Hyp (SEQ ID No: 134);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Hyp (SEQ ID No: 135);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp (SEQ ID No: 136);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp (SEQ ID No: 137); and particularly, those comprising the amino acid sequence:

Ser-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA (SEQ ID No: 138).

Further conjugates of the invention comprising peptide components as defined under (d) above include those in which K is an N-terminal HCA group, include those comprising the amino acid sequence:

HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 139); and, more preferably, that defined by the amino acid sequence:

HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 140).

Further preferred conjugates of the invention comprising peptide components as defined under (d) above that may be mentioned include those in which W¹ is Ala and J is Lys, such as those comprising the amino acid sequence:

Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 141);

Ala-Lys-Pro-Ser-Pro-Thr-Tyr-Pro-Lys (SEQ ID No: 142);

Ala-Lys-Hyp-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 143);

Ala-Lys-Hyp-Ser-DOPA-Hyp -Thr-DOPA-Hyp-Lys (SEQ ID No: 144); and

particularly, those defined by the amino acid sequence:

Ala-Lys-Hyp-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 145).

Further preferred conjugates of the invention comprising peptide components as defined under (d) above that may be mentioned include those in which J is absent, such as those comprising the amino acid sequence:

HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr (SEQ ID No: 146);

HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp (SEQ ID No: 147);

Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp (SEQ ID No: 148);

Ala-Lys-Pro-Ser-Pro-Thr-Tyr-Pro (SEQ ID No: 149);

Ala-Lys-Hyp-Ser-Tyr-Hyp-Thr-Tyr-Hyp (SEQ ID No: 150);

Ala-Lys-Hyp-Ser-DOPA-Hyp -Thr-DOPA-Hyp (SEQ ID No: 151); and

Ala-Lys-Hyp-Ser-Tyr-Hyp-Thr-DOPA-Hyp (SEQ ID No: 152).

30

The skilled person will understand that a conjugate is a compound formed by electrostatically linking and/or covalently linking a chemical compound to a different chemical compound.

The term 'electrostatic cross-linking' will be understood by the skilled person to include the association of disordered molecules into an ordered state by virtue of its nature or by electrostatic interactions (also referred to as 'self-assembly', which is a primary

mechanism of gelation observed in amphiphilic peptide molecules (Hauser et al., *Biomed. Mat.* **2015**, *11*, 014103).

5 In this case, conjugates of the invention are preferably formed by covalently linking one or more linear polysaccharide chains to one or more of the peptide components as defined under one or more of formulae (a), (b), (c) and/or (d) above.

10 In this respect, conjugates of the invention may comprise one or more (and preferably, at least two) linear polysaccharide, such as HA, chains. In other words, such linear polysaccharide chains (e.g. HA) can be cross-linked through linking to one or more of the peptide components as defined under one or more of formulae (a), (b), (c) and/or (d) above.

15 For example, conjugates of the invention feature at least one covalent bond (e.g. an amide bond) formed by a reaction between an amine (i.e. -NH₂) group present in a peptide component as defined under (a), (b), (c) and/or (d) above and a carboxylic acid (i.e. -CO₂H) moiety present in the one or more HA chains. For example, an amide bond may be formed between an amine group of one of the Lys residues in a peptide component of formula I and a carboxylic acid group of one of the glucuronic acid
20 residues of HA.

25 Preferably, in conjugates of the invention, at least about 0.1 (e.g. at least about 1%) of the carboxylic acid groups in the HA chains form an amide bond with the amine groups of a such a peptide component.

Other conjugates of the invention that may be mentioned are those there up to about 25% (e.g. up to about 5%) of the carboxylic acid groups in the HA chains form an amide bond with the amine groups of such a peptide component.

30 Particular conjugates of the invention that may be mentioned are those where from about 0.1% to about 25% (e.g. from about 1% to about 5%) of the carboxylic acid groups in the HA chains form an amide bond with the amine groups of such a peptide component.

35 It will be appreciated by those skilled in the art that a cross-link is a link from one polymer chain (e.g. a polysaccharide chain, such as HA) to another. In the context of the present invention, one or more of the peptide components as defined under (a), (b), (c) and/or (d) above may be chemically bonded to each of the two or more e.g.

HA chains such that the conjugate of the invention is formed through a cross link between the HA chains.

Particular conjugates of the invention are those formed between any of the specific peptide components defined under one or more (a), (b), (c) and/or (d) above, and in particular peptide components of formula I as defined under (a) above, in which:

A and B both represent Z or A¹-Q¹-B¹;

A¹ and B¹ both represent Z or A²-Q²-B²;

A² and B² both represent Z-Q³-Z;

Q¹, Q² and Q³ all represent 'a Lys fragment', in accordance with what is defined above; and

W represents DOPA or DOPA-Ala-; and/or

U represents DOPA,

and the conjugate comprises to or more and two or more hyaluronic acid molecules.

15

Further particular conjugates of the invention are those formed between peptide components of formula I as defined under (a) above, in which:

A and B both represent Z; and

W represents DOPA or DOPA-Ala-; and/or

20 U represents DOPA,

and the conjugate comprises to or more and two or more hyaluronic acid molecules.

As used herein, Pro represents proline, Ala represents alanine, Ser represents serine, Tyr represents tyrosine, Hyp represents hydroxyproline (including 3-hydroxyproline (3Hyp) and 4-hydroxyproline (4Hyp)), diHyp represents dihydroxyproline (including 3,4-dihydroxyproline (3,4diHyp), 3,5-dihydroxyproline (3,5diHyp) and 4,5-dihydroxyproline (4,5diHyp)), Thr represents threonine, Lys represents lysine, Ala represents alanine and DOPA represents 3,4-dihydroxyphenylalanine. 3,4-Dihydrocinnamic acid (HCA) residues are essentially DOPA residues but without the -NH₂ group in the 2- or α-carbon position relative to the carboxylic acid that is attached to the N-terminal amino acid (whether Lys or Ala).

Conjugates of the invention, whether in the form of salts or otherwise, include regioisomers within amino acids of the peptides (for example diHyp, Hyp and Tyr moieties), as well as mixtures of such regioisomers. For example, included within the definition of Tyr are, not only tyrosine (4-hydroxyphenylalanine), but also 2- and 3-hydroxyphenylalanine. Included within the definition of Hyp are 4-hydroxyproline (4Hyp), 3-hydroxyproline (3Hyp) and 5-hydroxyproline (5Hyp). It is more preferred

that Hyp residues are 4-hydroxyproline. Similarly, included within the definition of diHyp are 3,4-dihydroxyproline (3,4diHyp), 3,5-dihydroxyproline (3,5diHyp) and 4,5-dihydroxyproline (4,5diHyp). It is more preferred that diHyp residues are 3,4-dihydroxyproline (3,4diHyp).

5

Also, in addition to the standard central carbon atom of the amino acids in the conjugates of the invention (which are normally but not exclusively in the L-configuration), certain amino acids in the sequence comprise further chiral carbon atoms. All such stereoisomers and mixtures (including racemic mixtures) thereof are included within the scope of the invention. In respect, included within the definition of Hyp are *trans*-4-hydroxy-L-proline, *cis*-4-hydroxy-L-proline, *trans*-3-hydroxy-L-proline, *cis*-3-hydroxy-L-proline, *trans*-5-hydroxy-L-proline and *cis*-5-hydroxy-L-proline, however we prefer that the Hyp that is employed in conjugates of the invention is 4-hydroxy-L-proline. Similarly, corresponding definitions may be applied to diHyp, in which the two hydroxy groups can also be *cis* or *trans* relative to each other. In any event, individual enantiomers of peptide components as defined under formulae (a), (b), (c) or (d) above that may form part of a conjugate of the invention are included within the scope of the invention.

Conjugates of the invention may be in the form of salts. Salts that may be mentioned include pharmaceutically-acceptable and/or cosmetically-acceptable salts, such as pharmaceutically- and/or cosmetically-acceptable acid addition salts and base addition salts. Such salts may be formed by conventional means, for example by reaction of a conjugate of the invention with one or more equivalents of an appropriate acid or base, optionally in a solvent, or in a medium in which the salt is insoluble, followed by removal of said solvent, or said medium, using standard techniques (e.g. *in vacuo*, by freeze-drying or by filtration). Salts may also be prepared by exchanging a counter-ion of the conjugate of the invention in the form of a salt with another counter-ion, for example using a suitable ion exchange resin.

30

Preferred salts include, for example, acetate, hydrochloride, bisulfate, maleate, mesylate, tosylate, alkaline earth metal salts, such as calcium and magnesium, or alkali metal salts, such as sodium and potassium salts. Most preferably, conjugates of the invention may be in the form of acetate salts.

35

Conjugates of the invention may be prepared by way of conventional techniques, for example by way of standard amino acid coupling techniques, using standard coupling reagents and solvents, for example as described hereinafter. Conjugates of the

- invention may be synthesised from available starting materials using appropriate reagents and reaction conditions. In this respect, the skilled person may refer to *inter alia* "Comprehensive Organic Synthesis" by B. M. Trost and I. Fleming, Pergamon Press, 1991. Further references that may be employed include "Heterocyclic Chemistry" by J. A. Joule, K. Mills and G. F. Smith, 3rd edition, published by Chapman & Hall, "Comprehensive Heterocyclic Chemistry II" by A. R. Katritzky, C. W. Rees and E. F. V. Scriven, Pergamon Press, 1996 and "Science of Synthesis", Volumes 9-17 (Heteroarenes and Related Ring Systems), Georg Thieme Verlag, 2006.
- Conjugates of the invention may be isolated from their reaction mixtures and, if necessary, purified using conventional techniques as known to those skilled in the art. Thus, processes for preparation of conjugates of the invention as described herein may include, as a final step, isolation and optionally purification of the conjugate of the invention.
- It will be appreciated by those skilled in the art that, in the processes described above and hereinafter, the functional groups of intermediate compounds may need to be protected by protecting groups. The protection and deprotection of functional groups may take place before or after a reaction.
- Protecting groups may be applied and removed in accordance with techniques that are well-known to those skilled in the art and as described hereinafter. For example, protected compounds/intermediates described herein may be converted chemically to unprotected compounds using standard deprotection techniques. The type of chemistry involved will dictate the need, and type, of protecting groups as well as the sequence for accomplishing the synthesis. The use of protecting groups is fully described in 'Protective Groups in Organic Synthesis', 5th edition, T.W. Greene & P.G.M. Wutz, Wiley-Interscience (2014), the contents of which are incorporated herein by reference.
- Conjugates of the invention are useful as human and animal medicine. They are therefore indicated as pharmaceuticals (and/or in veterinary science), although they may also be used as cosmetics and/or as part of a medical device.
- Conjugates of the invention may also possess pharmacological activity as such, certain pharmaceutically-acceptable (e.g. 'protected') derivatives of conjugates of the invention may exist or may be prepared which may not possess such activity, but which may be administered and thereafter be metabolised or chemically transformed to form

conjugates of the invention. Such compounds (which may possess some pharmacological activity, provided that such activity is appreciably lower than that of the active conjugates to which they are metabolised/transformed) may therefore be described as 'prodrugs' of conjugates of the invention.

5

As used herein, references to prodrugs will include conjugates that form a conjugate of the invention, in an experimentally-detectable amount, within a predetermined time, following administration. All prodrugs of the conjugates of the invention are included within the scope of the invention.

10

When conjugates of the invention possess pharmacological activity, they are particularly useful in the treatment of inflammation.

The term 'treatment of inflammation' includes the treatment of inflammation in any organ of the body (including soft tissue, joints, nerves, the vascular system, internal organs, especially mucosal surfaces, and particularly the skin), irrespective of the cause, and also includes all such inflammatory disorders or conditions, and/or disorders or conditions characterized by inflammation (e.g. as a symptom).

Inflammatory disorders and/or conditions may be (and are typically) characterized by activation of immune defence mechanisms, resulting in an effect that is more harmful than beneficial to the host. Such conditions are generally associated with varying degrees of tissue redness or hyperemia, swelling, edema, hyperthermia, pain (including aching), exudation of body fluids, itching (pruritis), cell death and tissue destruction, cell proliferation, and/or loss of function.

Inflammatory conditions that may be mentioned include arteritis, diabetes mellitus, metabolic syndrome, rosacea, asthma and allergy, ankylosing spondylitis, chronic obstructive pulmonary disease, gouty arthritis, inflammatory bowel disease (such as Crohn's disease and ulcerative colitis), multiple sclerosis, osteoarthritis, pancreatitis, prostatitis, psoriatic arthritis, rheumatoid arthritis, tendinitis, bursitis, Sjögren's syndrome, systemic lupus erythematosus, uveitis, urticaria, vasculitis, mastocytosis, diabetic vascular complications, migraine, atherosclerosis and associated cardiovascular disorders. A disease state characterised by inflammation that may be mentioned is chronic obstructive pulmonary disease (COPD). A further disease state characterised by inflammation that may be mentioned is inflammatory bowel diseases including Crohn's disease and, especially, ulcerative colitis. Other disease states characterized by inflammation that may be mentioned are gynaecological diseases,

such as cervicitis, vaginitis (e.g. radiation vaginitis) and colpitis. Diseases that affect the gastrointestinal tract, such as gastrohelcosis (e.g. gastritis, gastric ulcer, gastric cancer and other stomach mucosa diseases) as well as gastroesophageal reflux disease (GERD), constipation, and gastritis, inflammation associated with cancers and
5 infections (e.g. viral infections, such as the common cold or influenza).

Inflammatory conditions that may be more especially mentioned include inflammations of the skin or mucosa (including the oral, nasal, ocular, vaginal, cervical and/or anorectal mucosae, more particularly the oral or nasal mucosae), such as inflammation
10 resulting from infections (such as viral and/or bacterial infections), or allergic/atopic conditions (such as rhinitis (e.g. allergic rhinitis), pharyngitis, periodontitis, gingivitis, xerophthalmia, conjunctivitis (e.g. allergic conjunctivitis), dermatitis, urticaria (hives) and food allergy); and other inflammatory conditions, such as herpes, drug eruptions, polymorphous light eruptions, sunburn, early manifestations of skin cancers
15 (erythema-like skin lesions), pathological hair loss (including following skin grafting), chemo rash, psoriasis, erythema multiforme, folliculitis, eczema and external otitis. A disease state that may be mentioned is polymorphous light eruptions.

More particularly, conjugates of the invention may be used to treat certain conditions
20 characterized by inflammation, and/or with which inflammation is associated. Such conditions may include wounds (including abrasions (scratches), incisions (including operative incisions), lacerations, punctures, avulsions, bruising and scarring), and burns (including inflammation resulting from surgery following burns, such as skin grafting) and other conditions, such as hemorrhoids. Wounds may be acute or chronic,
25 and/or may result from one or more inflammatory disorders as defined herein.

Wounds of the skin or mucosa may arise from internal or external physical injury to the membrane surface, or may be caused by (i.e. be a symptom of) an underlying physiological disorder.
30

Physical (e.g. 'open') wounds may be caused by sharp objects (cuts, incisions, punctures) or blunt objects/mechanical forces (lacerations, abrasions, avulsions), physical blows (bruises), heat or chemicals (burns and blisters), UV light (sunburn), cold (chilblains or frostbite). Wounds may be superficial (damage only to the
35 epidermis and/or dermis) or may be full thickness wounds (damage below the epidermis and/or dermis). In serious cases, subcutaneous and/or submucosal tissues, such as muscles, bones, joints, and even internal organs, may be damaged.

Conjugates of the invention may be used to relieve the pain (including aching) associated with inflammation and/or wounding. In particular, conjugates of the invention may be used to relieve procedural pain and/or non-procedural pain. The skilled person will understand that the term 'procedural pain' (i.e. operation pain) refers to acute pain that is associated with medical investigations and treatments conducted for the purpose of healthcare. The term 'non-procedural' refers to general pain that is associated with inflammation and/or wounding (e.g. pain associated with dental ulcers, burns and/or scars), and is not a consequence of a particular medical intervention.

10

Conjugates of the invention may be used to treat not only the inflammation, pain (including aching) and/or pruritis (itching) associated with the wound itself and the healing process, but also to prevent the exudation of body fluids from wounds, the risk of infection, and the prevention of physiological reactions that result from inflammation and/or wound healing processes, such as scarring and melanin pigmentation.

15

Scarring is a consequence of inflammation and/or wound healing and is a general term for the formation of fibrotic tissue that is a consequence of such inflammation/healing.

Conjugates of the invention may also be useful in the suppression of the production of melanin pigmentation, which may or may not result from inflammation and/or wound healing. Conjugates of the invention may also be useful in the suppression of disorders associated with melanin pigmentation, such as chloasma, freckles, melanosis, malar rash and other chromatosis, skin cancers with melanoma, and chromatosis that is caused by exposure to the sun or skin diseases like acne.

20

Wounds may also arise as a consequence of (e.g. inflammatory) diseases or disorders. Such wounds may include blistering and/or ulcers of the skin and mucosa. These are common conditions that are often long-lasting and difficult to treat. Skin tissues can often be damaged, removed, liquefied, infected and/or necrotic. Ulcers can lead to secondary consequences to health particularly if they become infected, are hard to heal and are costly to treat. They can also cause significant psychological stress and economic loss to patients, affecting both general well-being and quality of life.

30

In the alternative, inflammatory skin conditions or diseases in which conjugates of the invention find particular utility include psoriasis, acne, eczema and dermatitis, especially allergic/atopic dermatitis, as well as in the treatment of mucosal

35

inflammation as characterized by rhinitis, especially allergic rhinitis, hemorrhoids, chronic obstructive pulmonary disease and ulcerative colitis, for example.

5 Psoriasis is a chronic, inflammatory skin disease with a tendency to recur (some patients never heal during their entire life). Clinical manifestations of psoriasis mainly include erythema and scales. It can occur over the whole body, but is more commonly observed on the scalp and limbs.

10 Acne is a follicular (pilosebaceous unit) chronic, inflammatory skin disease, the occurrence of which is closely related to main factors like hypersebum production, blocked pilosebaceous ducts (including closed and open comedones), bacterial infection and inflammatory reactions, that tends to occur during youth, characterized by multiform skin lesions on the face. The term acne thus includes regular acne and acne rosacea (i.e. copper nose).

15 Eczema is a skin inflammatory reaction with strong itching caused by a variety of internal and external factors. It has three phases, acute, sub-acute, and chronic. In the acute phase, there is a tendency for the production of exudates, while the chronic phase includes infiltration and hypertrophy. Skin lesions are often itchy and recur
20 easily.

Dermatitis is a common skin disease characterized by coarseness, redness, itching, eczema, and dryness. Small lumps, refractory ulcers, and pigmented spots caused by dermatitis may, if not treated promptly, develop to basal cell carcinoma, squamous cell carcinoma, and malignant melanoma. Dermatitis may be caused by various internal
25 and external infectious or non-infectious factors, including substances (contact dermatitis) or allergy (allergic/atopic dermatitis). Also included is seborrheic dermatitis (seborrheic eczema) and all forms of steroid-dependent dermatitis (including light-sensitive seborrheic, perioral dermatitis, rosacea-like dermatitis, steroid-rosacea, steroid-induced rosacea, rosacea, steroid dermatitis resembling
30 rosacea, topical corticosteroid-induced rosacea-like dermatitis and, more particularly, facial corticosteroid addictive dermatitis (FCAD) or facial corticosteroid-dependent dermatitis (FCDD), as characterized by flushing, erythema, telangiectasia, atrophy, papules and/or pustules in the facial area after long-term treatment with (including
35 uncontrolled use, abuse or misuse of) topical corticosteroids; see, for example, Xiao *et al.*, *J. Dermatol.*, **2015**, 42, 697-702 and Lu *et al.*, *Clin. Exp. Dermatol.*, **2009**, 35, 618-621).

Rhinitis is irritation and inflammation of the mucous membrane inside the nose. Common symptoms of rhinitis include a stuffy nose, runny nose, sneezing and post-nasal drip. The most common kind of rhinitis is allergic rhinitis, caused by an allergen, such as pollen, dust, mould, or flakes of skin from certain animals. It has been
5 surprisingly found that patients with allergic rhinitis who were treated with conjugates of the invention experienced relief of eye itchiness, even when conjugates of the invention were administered nasally (i.e. to the nasal mucosa).

Hemorrhoids are swellings caused by inflammation of the hemorrhoidal blood vessels
10 found inside or around the rectum and the anus. Symptoms include bleeding (i.e. wounding) after the passage of a stool, prolapse of the hemorrhoid, mucus discharge and itchiness, soreness, redness and swelling in the area of the anus. Hemorrhoids are believed to be a consequence of an increase of pressure in the abdomen, for example, as a result of constipation or diarrhea.

15 Chronic obstructive pulmonary disease (COPD) is the name for a group of lung conditions that cause breathing difficulties, including emphysema (damage to the alveoli) and chronic bronchitis (long-term inflammation of the airways). COPD occurs when the lungs become inflamed, damaged and narrowed. The damage to the lungs
20 is usually irreversible and results in an impairment of the flow of air into and out of the lungs. Symptoms of COPD include breathlessness, productive cough, frequent chest infections and persistent wheezing. The most common cause of the disease is smoking, although other risk factors include high levels of air pollution and occupational exposure to dust, chemicals and fumes.

25 Conjugates of the invention may have positive effects in mitigating erythema, redness and swelling, edema, blisters, and bullous pemphigoid caused by various conditions including those mentioned generally and specifically herein, and may inhibit exudation of subcutaneous tissue fluid, and suppressing itching and pain caused by such
30 inflammatory conditions.

Other inflammatory conditions that may be mentioned include:

(a) Mucosal inflammation, such as oral mucositis, aphthous ulcers, otitis media,
35 laryngitis, tracheitis, esophagitis, gastritis, enteritis and enterocolitis (including bacillary dysentery, chronic amoebic dysentery, schistosomiasis, nonspecific ulcerative colitis and regional enteritis), cervicitis and endocervicitis, endometritis, inflammation caused by inhalation injury and the like, as well as mucosal inflammation associated

with cancers, and infections (e.g. viral infections, such as the common cold or influenza), that affect mucosal surfaces, such as those in the oral cavity, the nasopharynx, the ear, the throat, the trachea, the gastrointestinal tract, the cervix, etc.

5

(b) Orthopedic inflammation associated with, for example bone fractures, pyogenic infection of bones and joints, inflammation caused by rheumatic bone diseases, as well as pyogenic osteomyelitis (acute, chronic, localized, sclerotic, post-traumatic), pyogenic arthritis; bone tumors (osteoma, osteoid osteoma, chondroma), bone cysts, osteoclastoma, primary bone sarcoma (osteosarcoma, chondrosarcoma, osteofibrosarcoma, Ewing's sarcoma, non-Hodgkin's lymphoma, myeloma, chordoma), metastatic bone tumors, tumor-like lesions of bone (bone cyst, aneurysmal bone cyst, eosinophilic granuloma, fibrous dysplasia); and rheumatic arthritis.

15 (c) Nerve inflammation, such as peripheral polyneuritis, facial neuritis, peripheral neuritis, subcutaneous neuritis, ulnar neuritis, intercostal neuritis, etc.

(d) Subcutaneous and submucosal soft tissue inflammation, such as myositis, ligamentitis, tendonitis, panniculitis capsulitis, lymphadenitis, bubonadenitis, tonsillitis, synovitis, fasciitis, and soft tissue inflammation caused by injuries, contusion or laceration of muscles, ligaments, fascia, tendons, membrana synovialis, fat, articular capsules, and lymphoid tissue.

(e) Vascular inflammation, such as allergic leukocytoclastic vasculitis, allergic cutaneous vasculitis, polyarteritis nodosa, thrombotic vasculitis, granulomatous vasculitis, lymphocytic vasculitis, vasculitis with abnormalities in blood composition, and rheumatic vasculitis, as well as vascular inflammation associated with vascular cancers caused by allergic leukocytoclastic vasculitis, polyarteritis nodosa, thrombotic vasculitis, granulomatous vasculitis, lymphocytic vasculitis, vasculitis with abnormalities in blood composition, and rheumatic vasculitis.

(f) Inflammation of the internal organs, such as the heart, stomach, intestine, lung, liver, spleen, kidney, pancreas, bladder, ovary, and prostate, including but not limited to pericarditis, myocarditis, endocarditis, pneumonia, hepatitis, splenitis, nephritis, pancreatitis, cystitis, oophoritis, prostatitis and treatment of gastric ulcer.

(g) Inflammation of the eye and surrounding area, such as conjunctivitis, keratitis (e.g. acute epithelial keratitis, nummular keratitis, interstitial keratitis, disciform keratitis,

neurotrophic keratitis, mucous plaque keratitis, herpes simplex keratitis, herpes zoster keratitis, bacterial keratitis, fungal keratitis, acanthamoebic keratitis, onchocercal keratitis, superficial punctate keratitis, ulcerative keratitis, exposure keratitis, photokeratitis and contact lens acute red eye), optic neuritis, etc.

5

(h) Inflammation of the gums and the oral cavity, such as periodontitis, gingivitis, dental ulcers, etc.

(i) Inflammation associated with rheumatism, such as rheumatic vasculitis, rheumatoid arthritis, rheumatic bone diseases, ankylosing spondylitis, bursitis, Crohn's disease, gout, infectious arthritis, juvenile idiopathic arthritis, osteoarthritis, osteoporosis, polymyalgia rheumatica, polymyositis, psoriatic arthritis, scleroderma, Sjögren's syndrome, spondyloarthropathies, systemic lupus erythematosus, tendinitis, etc.

15 Conjugates of the invention may also be used in the treatment of certain specific diseases of the digestive system, such as gastroesophageal reflux disease (GERD), which may be characterized by an acidic taste in the mouth, regurgitation, heartburn, pain with swallowing and/or sore throat, increased salivation (water
brash), nausea, chest pain, and coughing. GERD may cause injury of the esophagus,
20 including reflux esophagitis (i.e. inflammation of the esophageal epithelium which may cause ulceration at or around the junction of the stomach and esophagus), esophageal strictures (i.e. the persistent narrowing of the esophagus caused by reflux-induced inflammation), Barrett's esophagus (i.e. intestinal metaplasia (i.e. changes of
epithelial cells from squamous to intestinal columnar epithelium of the distal esophagus)
25 and/or esophageal adenocarcinoma (a form of cancer)).

Conjugates of the invention may also be used in the treatment of certain specific diseases of the respiratory system, such as pulmonary cystic fibrosis, usual interstitial pneumonia, allergic pneumonia, asbestosis, emphysema, pulmonary heart disease,
30 pulmonary embolism, etc. A specific disease state that may be mentioned in idiopathic pulmonary fibrosis (IPF).

IPF is a diffuse and fatal pulmonary interstitial disease with pathological features including alveolar epithelial damage, massive proliferation of lung fibroblasts,
35 excessive deposition of extracellular matrix, ultimately leading to irreversible lung tissue damage. In the latter stages of the disease, subjects with IPF experience respiratory failure and death. It has been found that conjugates of the invention may

find utility in the treatment of IPF and/or alleviation of the symptoms associated with the disease.

Conjugates of the invention are particularly useful in the treatment of the following lung and/or fibrotic conditions (whether otherwise mentioned herein or not): lung fibrosis, renal fibrosis, liver fibrosis, silicosis, acute bronchitis, chronic bronchitis, tracheobronchitis, bronchial asthma, status asthmatics, bronchiectasis, upper respiratory tract infections (including the common cold and influenza), allergic airway inflammation, bacterial pneumonia, viral pneumonia, mycoplasma pneumonia, reckettsia, radiation pneumonia, pneumococcal (including staphylococcal, streptococcal and gram-negative bacillus) pneumonia, pulmonary candidiasis (including aspergillosis, mucormycosis, histoplasmosis, actinomycosis and nocardiosis), pulmonary mycosis, cryptococcosis, lung abscesses, anaphylactic pneumonia, extrinsic allergic alveolitis, pulmonary eosinophilia (including Loeffler's syndrome and eosinophilosis), obstructive pulmonary emphysema, pulmonary edema, pulmonary tuberculosis, respiratory alkalosis/acidosis, acute lung injury, interstitial lung disease, empyema, lung fibroma and cor pulmonale.

Particular mucosal disorders and disease in which conjugates of the invention find utility include anorectal diseases, such as diarrhea, hemorrhoids, abscesses, fistula, fissures, anal itching, anal sinusitis, warts and rectal prolapse; inflammatory bowel disease, including Crohn's disease and, particularly, ulcerative colitis; gynaecological diseases, such as cervicitis, vaginitis, pelvic pain and disorders; and dental diseases, such as paradentitis, for example.

25

Conjugates of the invention may further possess an antioxidation effect, by increasing SOD (superoxide dismutase) production and reducing lipid oxidation. Conjugates of the invention may therefore be considered to have antioxidant properties.

Conjugates of the invention may also possess antipyretic properties that allow for the treatment of a fever and/or alleviate the symptoms thereof; for example, by reducing a subject's body temperature, which results in a reduction of fever. Conjugates of the invention and formulations including them may therefore be considered to be antipyretics.

35

According to a further aspect of the invention there is provided a method of treatment of inflammation, of an inflammatory disorder, and/or of a disorder/condition characterised by inflammation (for example as a symptom), which method comprises

the administration of a conjugate of the invention or a salt thereof to a patient in need of such treatment.

For the avoidance of doubt, in the context of the present invention, the terms
5 'treatment', 'therapy' and 'therapy method' include the therapeutic, or palliative, treatment of patients in need of, as well as the prophylactic treatment and/or diagnosis of patients which are susceptible to, inflammation and/or inflammatory disorders.

Conjugates of the invention may further possess antiviral properties that may allow for
10 the treatment of a viral infection *per se*, that is treatment of a viral infection, or a viral disease, by interfering with the replication of the virus within a host, as opposed to the treatment of any symptoms of any viral infection or disease, such as pain and/or inflammation. Such antiviral properties may also allow for the prevention of the onset of such an infection or disease, the protection of cells in a host from (e.g. further) viral
15 infection, prevention or arrest of the spread of viral infection or disease (within a single host, or from one host to a new host), or for the prevention of reactivation of a virus after latency in a host.

According to a further aspect of the invention there is provided a method of treatment
20 of a viral infection, which method comprises the administration of a conjugate of the invention or a salt thereof to a patient in need of such treatment.

Viral infections that may be mentioned include those caused by viruses in the following families: adenoviridae (e.g. adenovirus), papillomaviridae (e.g. human papillomavirus),
25 polyomaviridae (e.g. BK virus; JC virus), herpesviridae (e.g. herpes simplex, type 1; herpes simplex, type 2; varicella-zoster virus; Epstein-Barr virus; human cytomegalovirus; human herpes virus, type 8), poxviridae (e.g. smallpox), hepadnaviridae (e.g. hepatitis B virus), parvoviridae (e.g. parvovirus B19), astroviridae (e.g. human astrovirus), caliciviridae (e.g. norovirus; Norwalk virus), picornaviridae
30 (e.g. coxsackievirus, hepatitis A virus; poliovirus; rhinovirus), coronaviridae (e.g. severe acute respiratory syndrome virus), flaviviridae (e.g. hepatitis C virus; yellow fever virus; dengue virus; West Nile virus; tick-borne encephalitis virus), retroviridae (e.g. human immunodeficiency virus; HIV), togaviridae (e.g. rubella virus), arenaviridae (e.g. Lassa virus), bunyaviridae (e.g. hantavirus; Crimean-Congo hemorrhagic fever virus; Hantaan virus), filoviridae (e.g. Ebola virus; Marburg virus; Ravn virus), orthomyxoviridae (e.g. influenza viruses, including influenza A virus (e.g. H1N1 and H3N2 viruses), influenza B virus or influenza C virus), paramyxoviridae (e.g. measles virus; mumps virus; parainfluenza virus, respiratory syncytial virus),
35

rhabdoviridae (e.g. rabies virus), hepeviridae (e.g. hepatitis E virus), reoviridae (e.g. rotavirus; orbivirus; coltivirus; Banna virus), as well as viruses not assigned to families, such as hepatitis D virus.

- 5 Viruses that may be more specifically mentioned include herpes simplex, type 1 and herpes simplex, type 2 viruses, human papillomavirus, influenza virus and parainfluenza virus.

10 Conjugates of the invention may further possess antibacterial and/or bacteriostatic properties that may allow for the treatment of a bacterial infection *per se*, that is treatment of a bacterial infection, or a bacterial disease, by interfering with bacterial growth or proliferation in a host, as opposed to the treatment of any symptoms of any bacterial infection or disease, such as pain and/or inflammation. Conjugates of the invention may therefore be considered to be bacteriocides and/or, preferably,
15 bacteriostatic agents.

Such antibacterial properties may also allow for the prevention of the onset of such an infection or disease, the protection of cells in a host from (e.g. further) bacterial infection, prevention or arrest of the spread of bacterial infection or disease (within a
20 single host, or from one host to a new host), or for the prevention of reactivation of a bacterium after latency in a host.

According to a further aspect of the invention there is provided a method of treatment of a bacterial infection, which method comprises the administration of a conjugate of
25 the invention or a salt thereof to a patient in need of such treatment.

As disclosed herein, conjugates of the invention may further possess anticancer properties that may allow for the treatment of a cancer *per se*, that is treatment of a cancer by interfering with the cancer as opposed to the treatment of any symptoms of
30 the cancer, such as pain and/or inflammation. Such anticancer properties may also include the prevention of the onset of such a disease e.g. by treating inflammation and thereby preventing such onset.

According to another aspect of the invention, there is provided a method of treatment
35 of cancer, which method comprises the administration of a conjugate of the invention or a salt thereof to a patient in need of such treatment.

Particular cancers that may be mentioned include oral cancer, a nasopharynx cancer, a middle ear cancer, a conjunctival cancer, a throat cancer, a tracheal cancer, an esophageal cancer, a gastric cancer, an intestinal cancer, a cervical cancer, an endometrial cancer, skin cancer and the like caused by oral mucositis, rhinitis, otitis media, conjunctivitis, pharyngitis, laryngitis, tracheitis, esophagitis, gastritis, enterocolitis, cervicitis, endometritis, erythema-like skin lesions and the like. A particular skin cancer that may be mentioned is basal cell carcinoma.

'Patients' include reptilian, avian and, preferably, mammalian (particularly human) patients.

In accordance with the invention, conjugates of the invention are preferably administered locally or systemically, for example orally, intravenously or intraarterially (including by intravascular and other perivascular devices/dosage forms (e.g. stents)), intramuscularly, cutaneously, subcutaneously, transmucosally (e.g. sublingually or buccally), rectally, intravaginally, intradermally, transdermally, nasally, pulmonarily (e.g. tracheally or bronchially), preferably topically, or by any other parenteral route, in the form of a pharmaceutical preparation comprising the conjugate(s) in pharmaceutically acceptable dosage form(s).

Administration by inhalation (e.g. nasally) is particularly useful when the condition to be treated is rhinitis or inflammation resulting from viral infections of the airways (e.g. upper respiratory tract infections, such as the common cold and influenza).

Pulmonary administration is particularly useful when the condition to be treated is COPD or IPF. Topical forms of administration may be enhanced by creating a spray comprising the conjugates of the invention, e.g. by using a powder aerosol or by way of an aqueous mist using an appropriate atomisation technique or apparatus, such as a nebulizer.

Anorectal administration is particularly useful when the condition to be treated is hemorrhoids or ulcerative colitis, using an appropriate delivery means, such as a solution of foam to be injected or a suppository.

Administration to the lower gastrointestinal tract may also be achieved by parenteral, and particularly by peroral, delivery, by means of standard delayed- or extended-release coating techniques known to those skilled in the art. In particular, distinct parts of the upper or lower intestine may be targeted. For example, colonic

administration can also be achieved by way of colon-targeted drug delivery means that are initially administered perorally or parenterally.

5 Conjugates of the invention may in the alternative be administered by direct systemic parenteral administration. Such administration may be useful in methods of treatment of an inflammatory and/or fibrotic disorder or condition of one or more internal organs of a patient.

10 Internal organs that may be mentioned include the stomach, the intestines, the pancreas, the liver, the spleen, the bladder, the vascular system, the ovaries, the prostate, preferably the heart and the kidneys and more preferably the lungs.

Fibrotic conditions of internal organs that may be mentioned include acute and/or severe internal fibrotic conditions characterised by the excessive accumulation of
15 fibrous connective tissues (as described above) in and around inflamed or damaged tissues. Formulations of the invention may thus be useful in the treatment or prevention of fibrogenesis (as described above) and the morbidity and mortality that may be associated therewith. Thus, (e.g. acute and/or severe) fibrotic conditions of the internal organs that may be treated with formulations of the invention include
20 fibrosis of the liver, the kidneys, the lungs, the cardiovascular system, including the heart and the vascular system, the pancreas, the spleen, the central nervous system (nerve fibrosis), bone marrow fibrosis, the eyes, the vagina, the cervix, etc.

Inflammatory conditions of internal organs include any condition that is, or may
25 develop into a condition that is, severe (i.e. one that requires intensive medical treatment), and in which some sort of inflammatory component is apparent, as may be characterised by detectable inflammation, and further in which morbidity is manifested (or is expected) and/or is life-threatening.

30 Inflammatory conditions that may be mentioned include one or more acute disorders or conditions of internal organs (i.e. one or more conditions that require, or may develop into a condition that requires, immediate medical interventions) that are characterized by inflammation (e.g. as a symptom), such as acute internal injuries, in one or more internal organs (including any of the organs mentioned hereinbefore). By
35 treating such acute inflammatory disorders, formulations of the invention may prevent or arrest the development of symptoms (acute or chronic) that are associated with such conditions, and also may arrest the progress of morbidity and/or mortality that is associated with such conditions.

Acute inflammatory conditions that may be mentioned thus include conditions such as peritonitis, pancreatitis, colitis, proctitis, gastritis, duodenitis, pharyngitis, GERD, parodontitis and stomatitis. Particular acute inflammatory conditions that may be mentioned include acute injury to one or more internal organs (including any of those mentioned hereinbefore), such as acute lung injury, inhalation injury (such as burns), acute respiratory distress syndrome (ARDS), severe acute respiratory syndrome (SARS), and multiple-organ inflammation, injury and/or failure.

Such conditions may be caused by internal or external trauma (e.g. injury or a burn), or by an infection by e.g. viruses, bacteria or fungi.

For example, proctitis (which includes eosinophilic, gonorrheal and/or ulcerative proctitis) may be caused by inflammatory bowel disease, infections, radiation (e.g. for cancer), drugs such as antibiotics, surgery or allergic conditions, such as food intolerances.

For example, multiple-organ inflammation, injury and/or failure may result from extensive and/or traumatic external injuries, including traumatic and/or extensive external burns. Traumatic external burns will be understood to include second-degree, and more particularly third-degree burns and fourth-degree, burns. Extensive external burns will be understood to include burns that affect at least about 10%, such as at least about 15%, including at least about 20% of a patient's body area. External (and internal) burns may result from exposure to heat, chemicals and the like.

Acute inflammatory and/or fibrotic conditions may also result from sepsis or septic shock, which can be caused by viral, bacterial or fungal infection. Furthermore, acute lung injury, ARDS and, particularly, SARS may be caused by viruses, such as coronaviruses, include the novel SARS coronavirus 2 (SARS-CoV-2).

Thus, in addition, one or more of the aforementioned (e.g. acute) inflammatory conditions may (indeed in some cases will likely) result in some form of internal tissue damage and/or dysfunction of relevant internal tissues. Relevant tissues thus include (e.g. mucosal) tissues, such as the respiratory epithelium. Such tissue damage may also give rise to one or more of the fibrotic conditions mentioned hereinbefore. For example, the SARS disease caused by the novel coronavirus SARS-CoV-2 (coronavirus disease 2019 or COVID-19) is known in many cases to result in fibrosis, which arise from one or more of a number of factors, including inflammation.

In this respect, conjugates of the invention and salts thereof find particular utility in the treatment of relevant inflammatory and/or fibrotic conditions on the basis that such conditions are often characterized by one or more comorbidities. By conditions that
5 are 'characterized by comorbidities', we include that the main condition in question results in (or from) one more further medical conditions, including (and indeed preferably) those mentioned hereinbefore, at the same time, which conditions may interact and/or overlap with each other in some way.

10 Thus, there are provided:

- methods of treatment of at least one inflammatory and/or fibrotic disorder or condition of one or more internal organs of a patient, which method comprises direct systemic parenteral administration of a conjugate of the invention, or a pharmaceutically-acceptable salt thereof, to a patient in need of such
15 treatment;
- a method of treatment of two or more inflammatory and/or fibrotic disorders or conditions of one or more internal organs of a patient, which method comprises direct systemic parenteral administration of a conjugate of the invention, or a pharmaceutically-acceptable salt thereof, to a patient in need
20 of such treatment; and
- a method of reduction in the incidence of morbidity and/or mortality that is or may be associated with one or more inflammatory and/or fibrotic disorders or conditions of one or more internal organs of a patient, which method comprises direct systemic parenteral administration of a conjugate of the invention, or a
25 pharmaceutically-acceptable salt thereof, to a patient in need of such treatment.

When conjugates of the invention/salts thereof are administered directly and parenterally, they may be administered intravenously, intraarterially, intravascularly,
30 perivascularly, intramuscularly, cutaneously, and/or subcutaneously, for example by way of direct injection, or by way of any other parenteral route, in the form of a conjugate of the invention or salt thereof in the form of a pharmaceutically-acceptable dosage form.

35 Pharmaceutically-acceptable formulations for use in such administration may thus comprise conjugates of the invention in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier, which may be selected with due regard to the intended route of direct parenteral administration and standard pharmaceutical practice. Such

pharmaceutically-acceptable carriers may be chemically inert to the active compounds and may have no detrimental side effects or toxicity under the conditions of use. Such pharmaceutically-acceptable carriers may also impart an immediate, or a modified, release of the conjugate of the invention.

5

Formulations for injection may thus be in the form of an aqueous formulation such as an a suspension and/or, more preferably a solution (e.g. an (optionally) buffered aqueous formulation (e.g. solution), such as a physiological saline-containing formulation (e.g. solution), a phosphate-containing formulation (e.g. solution), an acetate-containing formulation (e.g. solution) or a borate-containing formulation (e.g. solution), or a freeze-dried powder that may be reconstituted with a vehicle, such as an aqueous vehicle prior to use (e.g. injection)).

Formulations for injection may include other suitable excipients known to those skilled in the art, such as solvents (e.g. water), co-solvents, solubilizing agents (e.g. cyclodextrins), wetting agents, suspending agents, emulsifying agents, thickening agents, chelating agents, antioxidants, reducing agents, antimicrobial preservatives, bulking agents and/or protectants.

Formulations for injection are preferably buffered by standard techniques to physiologically-acceptable pH values (e.g. pHs of between about 4.5 and about 9.5, e.g. about 6 and about 9, such as between about 6.5 and about 8.5) using buffers and/or pH modifiers as described herein, and/or may further comprise tonicity-modifying agents (such as sodium chloride).

25

The above notwithstanding, preferred modes of delivery of conjugates of the invention include topically to the site of inflammation (e.g. the mucosa, including the oral and/or nasal mucosa, the lung, the anorectal area and/or the colon or, more preferably, the skin) in an appropriate (for example pharmaceutically- and topically-acceptable) vehicle suitable for application to the skin and/or the appropriate mucosal surface, and/or a commercially-available formulation, but may also include oral, intravenous, cutaneous or subcutaneous, nasal, intramuscular, intraperitoneal, or pulmonary delivery.

Administration by injection is particularly useful for administering the conjugates of the invention, in the form of a solution of suspension into e.g. the dermis (e.g. intradermal injection), joint cavity or the eyes.

Administration by intradermal injection (e.g. intradermally) is particularly useful for administering the conjugates of the invention, in the form of a solution or suspension (e.g. a dermal filler), into the dermis. This is particularly useful as a means of administration for melanin pigmentation therapy as described hereinbefore or for the use of the conjugates of the invention in the treatment of, e.g. wrinkles.

Administration by injection is particularly useful to fill, e.g. the surgical site of the nasal cavity, the anal fistula, the space between the gingival and the root or the sinus. This is particularly useful for shaping support and/or lubrication.

Conjugates of the invention will generally be administered in the form of one or more for example pharmaceutical formulations in admixture with a (e.g. pharmaceutically acceptable) adjuvant, diluent or carrier, which may be selected with due regard to the intended route of administration (e.g. topical to the relevant mucosa (including the lung) or, preferably, the skin) and standard pharmaceutical or other (e.g. cosmetic) practice. Such pharmaceutically acceptable carriers may be chemically inert to the active compounds and may have no detrimental side effects or toxicity under the conditions of use. Such pharmaceutically acceptable carriers may also impart an immediate, or a modified, release of the conjugate of the invention.

Suitable pharmaceutical formulations may be commercially available or otherwise prepared according to techniques that are described in the literature, for example, Remington *The Science and Practice of Pharmacy*, 22nd edition, Pharmaceutical Press (2012) and *Martindale – The Complete Drug Reference*, 38th Edition, Pharmaceutical Press (2014) and the documents referred to therein, the relevant disclosures in all of which documents are hereby incorporated by reference. Otherwise, the preparation of suitable formulations including conjugates of the invention may be achieved non-inventively by the skilled person using routine techniques.

Conjugates of the invention may be in the form of an aqueous formulation such as an emulsion, a suspension and/or a solution (e.g. an (optionally) buffered aqueous formulation (e.g. solution), such as a physiological saline-containing formulation (e.g. solution), a phosphate-containing formulation (e.g. solution), an acetate-containing formulation (e.g. solution) or a borate-containing formulation (e.g. solution)), or a freeze-dried powder.

Conjugates of the invention may further and/or in the alternative be combined with appropriate excipients to prepare:

- gel formulations (for which suitable gel matrix materials include cellulose derivatives, carbomer and alginates, gummi tragacanthae, gelatin, pectin, carrageenan, gellan gum, starch, Xanthan gum, cationic guar gum, agar, noncellulosic polysaccharides, saccharides such as glucose, glycerin, propanediol, vinyl polymers, acrylic resins, polyvinyl alcohol, carboxyvinyl polymer and, particularly, hyaluronic acid);
- lotions (for which suitable matrix materials include cellulose derivatives, glycerin, noncellulosic polysaccharides, polyethylene glycols of different molecular weights and propanediol);
- pastes or ointments (for which suitable paste matrix materials include glycerin, vaseline, paraffin, polyethylene glycols of different molecular weights, etc.);
- creams or foams (for which suitable excipients (e.g. foaming agents) include hydroxypropyl methyl cellulose, gelatin, polyethylene glycols of different molecular weights, sodium dodecyl sulfate, sodium fatty alcohol polyoxyethylene ether sulfonate, corn gluten powder and acrylamide);
- powder aerosols (for which suitable excipients include mannitol, glycine, dextrin, dextrose, sucrose, lactose, sorbitol and polysorbates, e.g. a dry powder inhalant);
- liquid, for example, water (aerosol) sprays for oral use or for inhalation (for which suitable excipients include viscosity modifiers, such as hyaluronic acid, sugars, such as glucose and lactose, emulsifiers, buffering agents, alcohols, water, preservatives, sweeteners, flavours, etc.); and/or
- injectable solutions or suspensions (which may be aqueous or otherwise and for which suitable excipients include solvents and co-solvents, solubilizing agents, wetting agents, suspending agents, emulsifying agents, thickening agents, chelating agents, antioxidants, reducing agents, antimicrobial preservatives, buffers and/or pH modifiers, bulking agents, protectants and tonicity-modifying agents), particular injectable solutions or suspensions that may be mentioned include dermal fillers (i.e. injectable fillers or soft-tissue fillers), particularly when the conjugate of the invention is combined with hyaluronic acid.

Moisturizing agents, such as glycerol, glycerin, polyethylene glycol, trehalose, glycerol, petrolatum, paraffin oil, silicone oil, hyaluronic acid and salts (e.g. sodium and potassium salts) thereof, octanoic/caprylic triglyceride, and the like; and/or antioxidants, such as vitamins and glutathione; and/or pH modifiers, such as acids, bases and pH buffers, may also be included in such formulations, as appropriate. Furthermore, surfactants/emulsifiers, such as hexadecanol (cetyl alcohol), fatty acids (e.g. stearic acid), sodium dodecyl sulfate (sodium lauryl sulfate), sorbitan esters (e.g. sorbitan stearate, sorbitan oleate, etc.), monoacyl glycerides (such as glyceryl

monostearate), polyethoxylated alcohols, polyvinyl alcohols, polyol esters, polyoxyethylene alkyl ethers (e.g. polyoxyethylene sorbitan monooleate), polyoxyethylene castor oil derivatives, ethoxylated fatty acid esters, polyoxylglycerides, lauryl dimethyl amine oxide, bile salts (e.g. sodium deoxycholate, sodium cholate),
5 lipids (e.g. fatty acids, glycerolipids, glycerophospholipids, sphingolipids, sterols, prenols, saccharolipids, polyketides), phospholipids, N,N-dimethyldodecylamine-N-oxide, hexadecyltrimethyl-ammonium bromide, poloxamers, lecithin, sterols (e.g. cholesterol), sugar esters, polysorbates, and the like; preservatives, such as phenoxyethanol, ethylhexyl glycerin, and the like; and thickeners, such as
10 acryloyldimethyltaurate/VP copolymer, may be included. In particular, stearic acid, glyceryl monostearate, hexadecanol, sorbitan stearate, cetyl alcohol, octanoic/capric glyceride etc. may be included, particularly in cream formulations.

Conjugates of the invention, and (e.g. pharmaceutical) formulations (e.g. aqueous
15 solutions, gels, creams, ointments, lotions, foams, pastes and/or dry powders as described above) including them, may further be combined with an appropriate matrix material to prepare a dressing or a therapeutic patch for application on a biological surface, such as the skin or a mucosal surface. Such formulations may thus be employed to impregnate a matrix material, such as gauze, non-woven cloth or silk
20 paper. The therapeutic patch may alternatively be, for example, a band-aid, a facial mask, an eye mask, a hand mask, a foot mask, etc.

Vaseline may be employed for use in applying such dressings to wounds, but we have also found that ointments based on PEGs (e.g. PEG 400) may be combined with matrix
25 materials to prepare dressings without the need to use Vaseline.

Conjugates of the invention may also be used in combination with solid supports (such as nasal dressings (for example, to stop nasal bleeding), dermal scaffolds (for example, in wound healing) or artificial bones (for example, in the case of bone
30 grafting/implantation).

Conjugates of the invention may be administered for inhalation by way of suspension, a dry powder or a solution. Suitable inhalation devices include pressurized metered-dose inhalers (pMDIs), which may be hand-or breath-actuated and employed with or
35 without a standard spacer device, dry powder inhalers (DPIs), which may be single-dose, multi-dose, and power-assisted, and soft mist inhalers (SMIs) or nebulizers, in which aerosol drug in a fine mist is delivered with slower velocity than a spray delivered using, for example, a pMDI.

In pMDIs, conjugates of the invention may be administered as a pressurized suspension of micronized particles distributed in a propellant (e.g. HFA, along with excipients, such as mannitol, lactose, sorbitol, etc.), or as an ethanolic solutions, to deliver one or more metered dose of between about 20 and about 100 μ L with each actuation. Actuation may be effected by hand (e.g. pressing) or by inhalation (breath-actuation), involving a flow-triggered system driven by a spring.

In DPIs, conjugates of the invention may be administered in the form of micronized drug particles (of a size between about 1 and about 5 μ m), either alone or blended with inactive excipient of larger particle size (e.g. mannitol), inside a capsule, which may be pre-loaded or manually loaded into the device. Inhalation from a DPI may de-aggregate the medication particles and disperse them within the airways.

In SMIs, conjugates of the invention may be stored as a solution inside a cartridge, which is loaded into the device. A spring may release the dose into a micropump, such that the dose is released when a button is pressed, releasing jet streams of drug solution.

Various nebulizers may also be used to administer conjugates of the invention in the form of a fine mist of aerosolized solution. Nebulizers may include breath-enhanced jet nebulizer (in which, with the assistance of a compressor, an air stream moves through jet causing drug solution to be aerosolized); breath-actuated jet nebulizers (in which, after a patient inhales, with the assistance of a compressor, an air stream moves through a tube causing the drug solution to be aerosolized); ultrasonic nebulizers (in which piezoelectric crystals vibrate causing aerosolization by heating causing nebulization); vibrating mesh nebulizers (in which piezoelectric crystals vibrate a mesh plate causing aerosolization to give very fine droplets without a significant change in temperature of the solution during nebulization).

According to a further aspect of the invention there is provided a process for the preparation of a pharmaceutical composition/formulation, as defined herein, which process comprises bringing into association a conjugate of the invention, as hereinbefore defined, with one or more pharmaceutically-acceptable excipient, as hereinbefore defined.

Conjugates of the invention may also be combined in treatment with one or more growth factors selected from platelet-type growth factors (including platelet-derived

growth factors, PDGFs); osteosarcoma-derived growth factors (ODGF), epidermal growth factors (EGFs), transforming growth factors (TGF α and TGF β), fibroblast growth factors (α FGF, β FGF), insulin-like growth factors (IGF-I, IGF-II), nerve growth factors (NGF), interleukin-type growth factors (IL-1, IL-1, IL-3), erythropoietin (EPO), and
5 colony stimulating factor (CSF).

According to a further aspect of the invention there is provided a (e.g. pharmaceutical) composition comprising a conjugate of the invention and one or more pharmaceutically-acceptable excipient, such as an adjuvant, diluent or carrier.
10 Preferred formulations are suitable for application locally to e.g. the mucosa (including the oral and/or nasal mucosa, the lung, the anorectal area and/or the colon) or, more preferably, the skin and therefore comprise a topically-acceptable adjuvant, diluent or carrier.

15 There is, thus, further provided pharmaceutical compositions comprising conjugates of the invention that are suitable for, adapted for, and/or packaged and presented for topical administration (e.g. to the mucosa, including the oral and/or nasal mucosa, the lung, the anorectal area and/or the colon, or, preferably, to the skin), as well as the use of such a formulation in the treatment of a disorder including inflammation, an
20 inflammatory disorder and/or a condition characterized by inflammation (e.g. as a symptom) by way of direct topical administration of that formulation (e.g. to the mucosa, including the oral and/or nasal mucosa, the lung, the anorectal area and/or the colon, or, preferably, to the skin).

25 In relation to this aspect of the invention, for the avoidance of doubt, topical formulations comprising conjugates of the invention may be used in any and all conditions described herein, including treatments of inflammation, in the treatment of any and all inflammatory disorder(s), and/or in the treatment of any and all condition(s) characterized by inflammation, as hereinbefore mentioned, defined or described.
30 Similarly, topical formulations comprising conjugates of the invention that may be mentioned include any and all of those mentioned, defined or described herein. Any and all of the relevant disclosures herein are hereby incorporated by reference in conjunction with this aspect of the invention.

35 Topical (e.g. liquid- or (e.g. aqueous) solution-based) formulations comprising conjugates of the invention may be particularly useful in wound recovery, and may alleviate pain (including aching) and, particularly, pruritis/itching that is associated with the wound itself and the wound healing process. Such topical formulations

comprising conjugates of the invention may be particularly useful in the prevention and/or suppression of the exudation of body fluids from wounds, particularly during the acute inflammation stage, for example during the first 48 hours, after a burn or wound has been inflicted. This prevents the risk of infection, and other physiological reactions. Such topical formulations comprising conjugates of the invention may also be particularly useful in the prevention and/or suppression of scarring and melanin pigmentation (*vide supra*), whether associated with wounds or otherwise.

Administration of the conjugates of the invention may be continuous or intermittent. The mode of administration may also be determined by the timing and frequency of administration, but is also dependent, in the case of the therapeutic treatment of inflammation, on the severity of the condition.

Depending on the disorder, and the patient, to be treated, as well as the route of administration, conjugates of the invention may be administered at varying therapeutically effective doses to a patient in need thereof.

Similarly, the amount of the conjugate of the invention in a formulation will depend on the severity of the condition, and on the patient, to be treated, but may be determined by the skilled person.

In any event, the medical practitioner, or other skilled person, will be able to determine routinely the actual dosage, which will be most suitable for an individual patient, depending on the severity of the condition and route of administration. The dosages mentioned herein are exemplary of the average case; there can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

Doses may be administered between once and four (e.g. three) times daily.

Appropriate concentrations of conjugates of the invention in an aqueous solution product may be about 0.01 (e.g. about 0.1) to about 15.0 mg/mL, in all cases calculated as the free (non-salt) conjugate.

Appropriate topical doses of conjugates of the invention are in the range of about 0.05 to about 50 $\mu\text{g}/\text{cm}^2$ of treated area, such as about 0.1 (e.g. about 0.5) to about 20 $\mu\text{g}/\text{cm}^2$ of treated area, including about 1 to about 10 $\mu\text{g}/\text{cm}^2$ of treated area, such as about 5 $\mu\text{g}/\text{cm}^2$ of treated area, in all cases calculated as the free (non-salt) conjugate.

Appropriate doses of conjugates of the invention for nasal administration (e.g. by inhalation) are in the range of about 0.01 µg to about 2000 mg, for example between about 0.1 µg to about 500 mg, or between 1 µg to about 100 mg. Particular doses
5 for nasal administration that may be mentioned include between about 10 µg to about 1 mg, particularly a dose of about 0.1 mg (i.e. about 100 µg). Nasal administration of about 0.1 mg per day of conjugates of the invention has been found to be particularly effective in the treatment of conditions associated with inflammation of the nasal passages and mucosae, such as rhinitis (e.g. allergic rhinitis) and/or conditions
10 associated with nasosinusitis surgery.

Appropriate doses of conjugates of the invention for pulmonary administration (e.g. by inhalation) are in the range of about 0.01 µg to about 2000 mg, for example between about 0.1 µg to about 500 mg, or between 1 µg to about 100 mg. Particular doses
15 for pulmonary administration that may be mentioned include between about 10 µg to about 10 mg, particularly a dose of about 0.6 mg (i.e. 60 µg) to 6 mg (e.g. for use in treating COPD or IPF).

We prefer that pH values of formulations comprising conjugates of the invention are in
20 the range of about 1.0 to about 9.0 (for example about 3.0 to about 8.0).

In any event, the dose administered to a mammal, particularly a human, in the context of the present invention should be sufficient to effect a therapeutic response in the mammal over a reasonable timeframe (as described hereinbefore). One skilled in the
25 art will recognize that the selection of the exact dose and composition and the most appropriate delivery regimen will also be influenced by *inter alia* the pharmacological properties of the formulation, the nature and severity of the condition being treated, and the physical condition and mental acuity of the recipient, as well as the age, condition, body weight, sex and response of the patient to be treated, and the
30 stage/severity of the disease, as well as genetic differences between patients.

Conjugates of the invention are useful in human and animal medicine. In this respect, and as described above, conjugates of the invention that possess an appropriate degree of relevant pharmacological (or biological) activity *per se* may be used as
35 human, and/or animal, medicines.

Certain conjugates of the invention comprising peptide components as defined under (a), (b), (c) or (d) above, preferably those in which W/W¹ represents HCA, HCA-Ala or,

more preferably, DOPA or DOPA-Ala, and/or U/U¹ represents DOPA, may in addition and/or instead of possessing the aforementioned biological activity, possess adhesive properties.

- 5 These adhesive properties stem from the fact that the relevant W/W¹ and/or U/U¹ groups are capable of cross-linking with each other in order to form three-dimension networks.

Such conjugates of the invention may adhere to a number of substrates including
10 inorganic substrates, such as glass, metal and the like, as well as organic substrates, such as biological tissue.

In respect, such conjugates of the invention may also be used as wound surface repair products, wound surface protecting products, medical biological adhesive products,
15 medical coating products, industrial coating products (e.g. in corrosion prevention in ships, electronic apparatuses, pipelines and the like), biochemical reagents, medical products, sterilization products, culture vessels for cell culture and the like.

Such conjugates of the invention may form a film over various skin and mucous wound
20 surfaces such as burns, scalds, ulcers, chilblains, and bedsores to aid in recovery. Such conjugates of the invention may also be used in surgery, e.g. in the closure of surgical incisions, adhesion of fractured bones, adhesion of mucous membranes, coatings of human body implants such as artificial bones, cartilage brackets, periosteal, artificial joints, dental implants, plugging stents, spinal fusion devices, spinal spacers
25 and organ patches.

According to a further aspect of the invention, there is provided a conjugate of the invention comprising one or more peptide components as defined under any of (a), (b), (c) and/or (d) above, preferably in which W/W¹ represents HCA, HCA-Ala- or, more
30 preferably, DOPA or DOPA-Ala-, and U/U¹ represents DOPA, as an adhesive or a film-forming material.

As discussed hereinbefore, naturally occurring MAP is known for its adhesive properties, but it should be remembered that such adhesives properties may arise from the fact
35 that that is a high molecular weight, linear peptide that can exist in multiple conformations, enabling inter- and intramolecular reactions/cross-linking of DOPA residues in molecules, and thereby adhesion. Conjugates of the invention (made with both linear polypeptides and/or proteins or multiply-branched lower molecular weight

residues) have proved to possess similar properties (whether adhesive or biological) to naturally-occurring MAP.

In addition to this property, such conjugates of the invention that may be employed as pharmaceutical excipients may be mixed with relevant active ingredients either before or after crosslinking and/or at least partial crosslinking, in order to form a stable pharmaceutical composition in which a conjugate of the invention is an excipient, such as a carrier.

Such crosslinking may be carried out by a variety of chemical (e.g. iodine vapour, glutaraldehyde, N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride and N-hydroxysuccinimide (EDC/NHS), 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM), or other water soluble condensation agents) or enzymatic means (e.g. tyrosinase, or as described hereinafter). This notwithstanding, irrespective of the level of pharmacological activity that conjugates of the invention may possess, they may in any event be (and/or may be further) combined with active pharmaceutical ingredients, either in combination therapy (as described hereinafter), or by performing a function either as, or as part of, a pharmaceutically-acceptable excipient (e.g. an adjuvant, diluent or carrier), as part of a medical device, and/or as part of a drug-medical device combination.

In this respect, certain conjugates of the invention may thus be described as novel multifunctional excipients, which may be used for a variety of applications in the pharmaceutical field. In this respect, conjugates of the invention may be used as adhesives or as film-forming agents (for example as described hereinafter) and/or may be used as release retarding polymers, as binders, as suspending agents, as gelling agents, as coating agents, as diluents or as carriers for drugs of varying solubilities.

In this respect, conjugates of the invention may be combined with a multitude of known pharmaceutically-active ingredients, including any agent, or drug, capable of producing some sort of physiological effect (whether in a therapeutic or prophylactic capacity against a particular disease state or condition) in a living subject, including, in particular, mammalian and especially human subjects (patients). This is the case irrespective of whether the conjugate of the invention is employed:

- as a separate pharmaceutically-active ingredient *per se* in combination therapy;
- as, or as part of, a medical device;
- as, or as the medical device part of, a drug-medical device combination; or
- as a pharmaceutically-acceptable excipient.

Such patients may also (and/or may already) be receiving therapy based upon administration of one or more of such other, known pharmaceutically-active ingredients, by which we mean receiving a prescribed dose of one or more of the active ingredients mentioned herein, prior to, in addition to, and/or following, treatment with a conjugate of the invention.

Pharmaceutically-active agents that may be co-administered with a conjugate of the invention include any agent, or drug, that is capable of producing some sort of physiological effect (whether in a therapeutic or prophylactic capacity against a particular disease state or condition) in a living subject, including, in particular, mammalian and especially human subjects (patients).

Pharmaceutically-active agents may, for example, be selected from anti-inflammatory agents, pro-inflammatory agents, antibiotics, anti-bacterial and/or antiprotozoal agents, antiviral agents (e.g. protease inhibitors), anaesthetics and wound recovery drugs (e.g. growth factors).

In this respect conjugates of the invention may be combined with a multitude of known pharmaceutically-active ingredients, including any agent, or drug, capable of producing some sort of physiological effect (whether in a therapeutic or prophylactic capacity against a particular disease state or condition) in a living subject, including, in particular, mammalian and especially human subjects (patients).

Biologically-active agents may, for example, be selected from anti-inflammatory agents, pro-inflammatory agents, antibiotics, anti-bacterial and/or antiprotozoal agents, antiviral agents (e.g. protease inhibitors), anaesthetics and wound recovery drugs (e.g. growth factors).

Non-limiting examples of anti-inflammatory drugs which may be used also include those used in the treatment of rheumatic diseases and/or arthritis (such as cataflam, betamethasone, naproxen, cyclosporin, chondroitin, celecoxib, etodolac, meclofenamate, salsalate, methylprednisolone, and piroxicam); osteoarthritis (such as sulindac, meloxicam, fenoprofen, etoricoxib, and nabumetone); inflammation and its symptoms, e.g. fever, pain, itchiness and/or swelling (such as mefenamic acid, indomethacin, aspirin, ketorolac, fluorometholone, loteprednol, hydrocortisone, fluorometholone, bromfenac, prednisolone acetate, indomethacin, and ibuprofen); allergies and their symptoms (such as pheniramine, diphenhydramine, naphazoline,

antazoline, prednisolone, lodoxamide, pemirolast, oxymetazoline, ketotifen, naphazoline, emestine fumarate, olopatadine, azelastine, tranilast, levocabastine, cortisone, ephedrine, cetirizine, levocetirizine, pseudophedrine, fexofenadine, terfenadine, loratadine, and alexis); respiratory diseases, including asthma and/or
 5 COPD (such as budesonide, ciclesonide, nedocromil, dexamethasone, ambroxol, and pranlukast); skin diseases (such as mometasone, triamcinolone, desonide, sulfacetamide, tacrolimus, allantoin, and triamcinolone); mastocytosis (such as cromolyn); gout (such as diclofenac, and febuxostat); conjunctivitis (such as hydrobenzole, pranoprofen, and zinc sulfate); eye diseases (such as dextran 70,
 10 thyroxine/liothyronine, and ocular extractives), known or commercially-available pharmaceutically acceptable salts of any of the foregoing, and combinations of any of the foregoing compounds and/or salts.

Antiinflammatory drugs that may be mentioned include endogenous (and/or
 15 exogenous) lipid-based pro-resolving, antiinflammatory molecules or mediators, such as lipoxins, resolvins, and protectins. Pro-inflammatory agents that may be mentioned include prostaglandins (e.g. latanoprost, prostaglandin E1, and prostaglandin E2), and leukotrienes (e.g. Leukotriene B4).

20 Non-limiting examples of anti-bacterial drugs which may be used also include chloramphenicol, ofloxacin, levofloxacin, tobramycin, norfloxacin, ciprofloxacin, lomefloxacin, lincomycin, fluconazole, enoxacin, furazolidone, nitrofurazone, rifampicin, micronomicin, gentamicin, cetylpyridinium, neomycin, roxithromycin, sulfadiazine silver, clarithromycin, clindamycin, metronidazole, azithromycin, mafenide,
 25 sulfamethoxazole, paracetamol, chloramphenicol, pseudoephedrine, mupirocin, amoxicillin, amoxicillin/clavulanic acid, trimethoprim/sulfamethoxazole, cefalexin, moxifloxacin, known or commercially-available pharmaceutically acceptable salts of any of the foregoing, and combinations of any of the foregoing compounds and/or salts.

30 Non-limiting examples of antiviral drugs which may be used also include tobramycin ribavirin, acyclovir, moroxydine, foscarnet, ganciclovir, idoxuridine, trifluridine, brivudine, vidarabine, entecavir, telbivudine, foscarnet, zidovudine, didanosine, zalcitabine, stavudine, lamivudine, abacavir, emtricitabine, nevirapine, delavirdine, efavirenz, etravirine, rilpivirine, saquinavir, ritonavir, indinavir, nelfinavir, amprenavir,
 35 lopinavir, ritonavir, atazanavir, fosamprenavir, tipranavir, darunavir, telaprevir, boceprevir, simeprevir, asunaprevir, raltegravir, elvitegravir, dolutegravir, rsv-igiv, palivizumab, docosanol, enfuvirtide, maraviroc, vzig, varizig, acyclovir, ganciclovir, famciclovir, valacyclovir, penciclovir, valganciclovir, cidofovir, tenofovir disoproxil

fumarate, adefovir dipivoxil, fomivirsen, podofilox, imiquimod, sinecatechins, interferon- α 2b (recombinant, human), known or commercially-available pharmaceutically acceptable salts of any of the foregoing, and combinations of any of the foregoing compounds and/or salts.

5

Non-limiting examples of anaesthetics which may be used also include articaine, dextropropoxyphene, sevoflurane, cophenylcaine, lidocaine, prilocaine, pramoxine, benzocaine, dibucaine, diclonine, tetracaine, bupivacaine and known or commercially-available pharmaceutically acceptable salts of any of the foregoing, and combinations

10 of any of the foregoing compounds and/or salts.

Non-limiting examples of wound recovery drugs which may be used also include basic fibroblast growth factor (recombinant, human; recombinant, bovine), epidermal growth factor (recombinant, human; yeast), rhEGF (I), acidic fibroblast growth factor

15 (recombinant, human), granulocyte macrophage stimulating factor (recombinant, human), sulfadiazine silver, sulfadiazine zinc, fusidic Acid, bacitracin, chlorhexidine, silver nitrate, triethanolamine, ethacridine, retinoids, calf blood deproteinized extract, carraghenates, amiotide and known or commercially-available pharmaceutically acceptable salts of any of the foregoing, and combinations of any of the foregoing

20 compounds and/or salts.

Such pharmaceutically-active ingredients include those that may be administered topically, e.g. to the skin or to a mucosal surface along with a conjugate of the invention. In this respect, preferred active ingredients from the above list include cyclosporin,

25 chondroitin, loteprednol, fluorometholone, bromfenac, prednisolone acetate, indomethacin, oxymetazoline, ketotifen, naphazoline, emestine fumarate, olopatadine, azelastine, tranilast, levocabastine, cortisone, ephedrine, cetirizine, pseudoephedrine, levocetirizine, fexofenadine, terfenadine, loratadine, alexis, dexamethasone, ambroxol), sulfacetamide, tacrolimus, allantoin, triamcinolone, cromolyn, nedocromil,

30 diclofenac, hydrobenzole, pranoprofen, zinc sulfate, dextran 70, thyroxine/liothyronine, ocular extractives, chloramphenicol, ofloxacin, levofloxacin, tobramycin, norfloxacin, ciprofloxacin, lomefloxacin, lincomycin, fluconazole, enoxacin, furazolidone, nitrofurazone, rifampicin, micronomicin, gentamicin, cetylpyridinium, neomycin, roxithromycin, sulfadiazine silver, clarithromycin, sulfamethoxazole, chloramphenicol,

35 tobramycin ribavirin, acyclovir, moroxydine, foscarnet, ganciclovir, interferon- α 2b (recombinant, human), articaine, dextropropoxyphene, sevoflurane, cophenylcaine, lidocaine, prilocaine, pramoxine, benzocaine, dibucaine, diclonine, tetracaine, bupivacaine, basic fibroblast growth factor (recombinant, human; recombinant,

bovine), epidermal growth factor (recombinant, human; yeast), rhEFG (I), acidic fibroblast growth factor (recombinant, human), granulocyte macrophage stimulating factor (recombinant, human), sulfadiazine silver, sulfadiazine zinc, fusidic acid, bacitracin, chlorhexidine, silver nitrate, triethanolamine, ethacridine, retinoids, calf
5 blood deproteinized extract, carraghenates, amiotide, and known or commercially-available pharmaceutically acceptable salts of any of the foregoing, and combinations of any of the foregoing compounds and/or salts.

Other pharmaceutically-active ingredients that may be co-administered with a
10 conjugate of the invention include those that may be administered to treat one or of the gastrointestinal disorders mentioned hereinbefore.

Non-limiting examples of gastrointestinal drugs include oxalazine (olsalazine), sulfasalazine, domperidone, erythromycin, berberine, dexamethasone, cefuroxime
15 axetil, levofloxacin, mesalazine, belladonna, sulfobenzidine, azathioprine, sulfasalazine, live bacillus (such as clostridium butyricum, licheniformis, cereus), probiotics (such as bifidobacterium) tegafur, nifuratel, amoxicillin, ampicillin, nystatin, allicin, cefadroxil, dyclonine, carmofur, fluorouracil, mosapride, sodium carbosulfan, thrombin, pantoprazole, cimetidine, cisapride, ethylenediamine diacetamine, nimustine,
20 famotidine, barium sulfate, aminocaproic acid, roxatidine acetate, vincristine, azasetron, lentinan, bismuth salts (e.g. aluminate, potassium citrate) in combination with e.g. magnesium salts, magnesium trisilicate, bicarbonate, vitamin U, aluminium hydroxide, belladonna extract, famotidine and calcium carbonate, magnesium hydroxide, hydrotalcite, proton pump inhibitors (such as omeprazole, lansoprazole,
25 rabeprazole, pantoprazole, dexlansoprazole or esomeprazole), glycine, trypsin, allantoin aluminium hydroxide, sodium L-glutamine gualenate, rebampette, rotundine, quixipite, lafutidine, thymus protein, hericium erinaceus, irsogladine maleate, nizatidine, L-glutamine and sodium azulene sulfonate (sodium gualenate), ranitidine, bismuth citrate, lactobacillin, bisacordine, dimethylsiloxane, live clostridium butyricum,
30 loperamide hydrochloride, dibazol, secnidazole, zinc acephate, montmorillonite, tegafur/gimeracil/oteracil, famotidine, oteracil, doxifluridine, capecitabine and known or commercially-available pharmaceutically acceptable salts of any of the foregoing.

Pharmaceutically-active ingredients that may be mentioned for use in combination with
35 conjugates of the invention include active ingredients that are useful in the treatment of inflammation and/or inflammatory disorders (other anti-inflammatory agents).

Anti-inflammatory agents that may be used in combination with conjugates of the invention in the treatment of inflammation include therapeutic agents that are useful in the treatment of inflammation and/or of diseases characterized by inflammation as one of its symptoms, including those described hereinbefore. Depending on the condition to be treated, such anti-inflammatory agents may include NSAIDs (e.g. aspirin), aminosalysates (e.g. 5-aminosalicylic acid (mesalazine)), leukotriene receptor antagonists (e.g. montelukast, pranlukast, and zafirlukast), corticosteroids, analgesics and certain enzymes, such as trypsin, for example as described hereinafter. Conjugates of the invention may also be combined with leukotrienes (e.g. cysteinyl leukotrienes, and leukotriene B₄).

Other preferred agents that may be combined with conjugates of the invention include LTB₄ (to treat wounds and burns), NSAIDs (e.g. aspirin) or montelukast (to treat inflammation generally) and trypsin (to treat inflammation of the mucosa associated with e.g. viral infections).

Conjugates of the invention may also be combined with other therapeutic agents which, when administered, are known to give rise to inflammation as a side-effect.

Conjugates of the invention may also be combined with stem cells (e.g. totipotent (omnipotent), pluripotent (such as embryonic or induced pluripotent stem cells), multipotent (such as mesenchymal stem cells), oligopotent (such as hematopoietic stem cells), or unipotent (such as muscle stem cells)).

Other known pharmaceutically-active ingredients may also be administered in combination with conjugates of the invention in numerous ways.

For example, conjugates of the invention may be 'combined' with the (or with the other) pharmaceutically-active ingredients (or 'therapeutic agents') for administration together in the same (e.g. pharmaceutical) formulation, or administration separately (simultaneously or sequentially) in different (e.g. pharmaceutical) formulations.

Thus, such combination products provide for the administration of conjugates of the invention in conjunction with the (or with the other) therapeutic agent, and may thus be presented either as separate formulations, wherein at least one of those formulations comprises a conjugate of the invention, and at least one comprises the (or the other) therapeutic agent, or may be presented (i.e. formulated) as a combined

preparation (i.e. presented as a single formulation including a conjugate of the invention and the (or the other) therapeutic agent).

Thus, there is further provided:

5

(1) a (e.g. pharmaceutical) formulation including a conjugate of the invention; another pharmaceutically-active ingredient; and, optionally, a pharmaceutically-acceptable inactive excipient (e.g. adjuvant, diluent or carrier), which formulation is hereinafter referred to as a 'combined preparation'; and

10

(2) a kit of parts comprising components:

(A) a conjugate of the invention, optionally in the form of an (e.g. pharmaceutical) formulation in admixture with a pharmaceutically-acceptable inactive excipient (e.g. adjuvant, diluent or carrier); and

15

(B) another pharmaceutically-active ingredient, optionally in the form of a (e.g. pharmaceutical) formulation in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier,

which components (A) and (B) are each provided in a form that is suitable for administration in conjunction with the other.

20

In a further aspect of the invention, there is provided a process for the preparation of a combined preparation (1) as hereinbefore defined, which process comprises bringing into association a conjugate of the invention, the other pharmaceutically-active ingredient, and at least one (e.g. pharmaceutically-acceptable) excipient.

25

In a further aspect of the invention, there is provided a process for the preparation of a kit-of-parts (2) as hereinbefore defined, which process comprises bringing into association components (A) and (B). As used herein, references to bringing into association will mean that the two components are rendered suitable for administration in conjunction with each other.

30

Thus, in relation to the process for the preparation of a kit-of-parts as hereinbefore defined, by bringing the two components 'into association with' each other, we include that the two components of the kit-of-parts may be:

35

(i) provided as separate formulations (i.e. independently of one another), which are subsequently brought together for use in conjunction with each other in combination therapy; or

(ii) packaged and presented together as separate components of a 'combination pack' for use in conjunction with each other in combination therapy.

Thus, there is further provided a kit of parts comprising:

- 5 (I) one of components (A) and (B) as defined herein; together with
(II) instructions to use that component in conjunction with the other of the two components.

In relation to kits of parts described above, although the conjugate of the invention
10 may be provided in the form of a (e.g. pharmaceutical) formulation, in admixture with one or more additional pharmaceutically-acceptable excipients (e.g. adjuvants, diluents or carriers), when the compound of the invention is provided with a view to it primarily performing its function as a medical device or as an excipient, it may not be provided along with such additional pharmaceutically-acceptable excipients. In any
15 event, it is preferred that the (other) pharmaceutically-active ingredient of the kit of parts is provided in the form of a pharmaceutical formulation in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.

The kits of parts described herein may comprise more than one (e.g. formulation
20 including an) appropriate quantity/dose of a conjugate of the invention, and/or more than one (e.g. formulation including an) appropriate quantity/dose of the other pharmaceutically-active ingredient, in order to provide for repeat dosing. If more than one formulation comprising or quantity/dose of either of the foregoing is present, such may be the same, or may be different in terms of the dose of either compound,
25 chemical composition(s) and/or physical form(s).

With respect to the kits of parts as described herein, by 'administration in conjunction with', we include that respective components are administered, sequentially, separately and/or simultaneously, over the course of treatment of the relevant
30 condition.

Thus, in respect of the combination product according to the invention, the term 'administration in conjunction with' includes that the two components of the combination product (conjugate of the invention and other pharmaceutically-active
35 ingredient) are administered (optionally repeatedly), either together, or sufficiently closely in time, to enable a beneficial effect for the patient, that is greater, over the course of the treatment of the relevant condition, than if either the conjugate of the invention, or (e.g. a formulation comprising) the other agent, are administered

(optionally repeatedly) alone, in the absence of the other component, over the same course of treatment. Determination of whether a combination provides a greater beneficial effect in respect of, and over the course of treatment of, a particular condition will depend upon the condition to be treated or prevented, but may be
5 achieved routinely by the skilled person.

Further, in the context of a kit of parts according to the invention, the term 'in conjunction with' includes that one or other of the two components may be administered (optionally repeatedly) prior to, after, and/or at the same time as,
10 administration of the other component. When used in this context, the terms 'administered simultaneously' and 'administered at the same time as' include that individual quantities/doses of the relevant conjugate of the invention and other active pharmaceutical ingredient are administered within 48 hours (e.g. 24 hours) of each other.

15 Depending on the disorder, and the patient, to be treated, as well as the route of administration, conjugates of the invention may be administered at varying therapeutically effective doses to a patient in need thereof.

20 In relation to combined preparations and kits of parts described above, it is preferred that the other pharmaceutically-active ingredient is an anti-inflammatory agent, or agent known to give rise to inflammation as a side-effect, as hereinbefore described.

Wherever the word 'about' is employed herein, for example in the context of amounts,
25 such as concentrations and/or doses of the conjugates of the invention and/or the pharmaceutically-active ingredients, molecular weights or pHs, it will be appreciated that such variables are approximate and as such may vary by $\pm 10\%$, for example $\pm 5\%$ and preferably $\pm 2\%$ (e.g. $\pm 1\%$) from the numbers specified herein. In this respect, the term 'about 10%' means e.g. $\pm 10\%$ about the number 10, i.e. between
30 9% and 11%.

Conjugates of the invention have the advantage that they have a wide variety of uses including:

- as biologically-active agents in variety of conditions characterised by
35 inflammation, whether that condition is an organic inflammatory disease *per se* or is associated with, or is characterised by, inflammation (e.g. a wound or a burn), and/or in surgical and/or cosmetic applications as described hereinbefore

- in combination with active pharmaceutical ingredients, either in combination therapy, or by performing a more inert function either as, or as part of:
 - a pharmaceutically-acceptable excipient (e.g. an adjuvant, diluent or carrier),
 - 5 ○ a medical device, and/or
 - the medical device part of a drug-medical device combination.

The conjugates, uses and methods described herein may also have the advantage that, in the treatment of the conditions mentioned hereinbefore, they may be more
10 convenient for the physician and/or patient than, be more efficacious than, be less toxic than, have a broader range of activity than, be more potent than, produce fewer side effects than, or that it/they may have other useful pharmacological properties over, similar compounds or methods (treatments) known in the prior art, whether for use in the treatment of inflammation, inflammatory disorders, or disorders
15 characterised by inflammation as a symptom (including wounds), or otherwise.

The invention is illustrated, but in no way limited, by the following examples

Examples

20

Illustrative Example 1

(Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys)₂ (SEQ ID No: 11)

Fmoc-Lys(Boc)-Wang resin (9.15 g, GLS180322-41301, GL Biochem, Shanghai, China)
25 was loaded into a glass reaction column.

Methylene chloride (DCM, 200 mL; Shandong Jinling Chemical Industry Co. Ltd., Shandong, China) was added to the column and allowed to soak the resin for about half an hour. The DCM was then removed by vacuum filtration.

30

The resin was washed 3 times with *N,N*-dimethylformamide (DMF, 200 mL; Shandong Shitaifeng Fertilizer Industry Co. Ltd., Shandong, China).

A 20% piperidine solution in DMF (200 mL; Shandong Shitaifeng Fertilizer Industry Co. Ltd., Shandong, China) and was added as deprotection solution and reacted for 20
35 minutes. The solution was then removed by vacuum filtration and the column was washed with DMF six times.

- Fmoc-DOPA(Acetonide)-OH (4.14 g; GLS190219-21003, GL Biochem, Shanghai, China) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium tetrafluoroborate (TBTU, 2.89 g; GLS170805-00705, GL Biochem, Shanghai, China) were added to the resin. DMF (150 mL) was added to the reaction column, followed by *N,N*-diisopropylethylamine (DIPEA, 2.33 g; Suzhou Highfine Biotech Co. Ltd., Jiangsu, China). A Kiaser test was carried out with few of the resin after 30 min reaction. colour reaction was detected in the resin after 30 minutes, indicating the reaction was complete. The solvent was removed by vacuum filtration.
- 10 The above coupling steps were repeated to couple the remaining amino acids in the same amounts (by mols): Fmoc-Thr(tBu)-OH, Fmoc-4-Hyp(tBu)-OH, Fmoc-4-Hyp(tBu)-OH, Fmoc-Tyr(tBu)-OH, Fmoc-Ser(tBu)-OH, Fmoc-Pro-OH, Fmoc-Lys(Boc)-OH and Fmoc-Ala-OH.
- 15 After the Fmoc-Ala-OH was coupled to the resin, the above coupling steps were repeated starting with Fmoc-Lys(Boc)-OH and followed by Fmoc-DOPA(Acetonide)-OH, Fmoc-Thr(tBu)-OH, Fmoc-4-Hyp(tBu)-OH, Fmoc-4-Hyp(tBu)-OH, Fmoc-Tyr(tBu)-OH, Fmoc-Ser(tBu)-OH, Fmoc-Pro-OH, Fmoc-Lys(Boc)-OH and Fmoc-Ala-OH.
- 20 In a separate procedure, after Fmoc-Ala-OH was coupled on the resin, a deprotection step was carried out to remove the Fmoc protection on Dopa. The resin was washed 3 times with DMF (200 mL each time). A 20% piperidine solution in DMF (200 mL) was added as a deprotection solution and reacted for 20 minutes. Then, the resin was washed three times each with the following solvents, DMF (200 mL each time), DCM
- 25 (200 mL each time) and methanol (200 mL each time; Xilong Scientific Co. Ltd., Guangdong, China). The resin was dried under vacuum for about 2 hours.
- 160.0 mL (i.e. 10 mL per gram of the dried resin) of lysate, which comprised of 95% trifluoroacetic acid (TFA), 2.5% water and 2.5% triisopropylsilane (Tis), were added to
- 30 immerse the resin-bounded peptide-containing compound. After cleavage for about 2 hours, the solid support was removed by filtration and the filtrate was collected under reduced pressure. The filtrate was precipitated with 1600 mL (i.e. 10 mL per mL of the filtrate) of diethyl ether (Xilong Scientific Co., Ltd., Guangdong, China) and the sediment was collected by filtration. The sediment was dried by vacuum for about 2
- 35 hours, yielding 7.53 g of the crude title compound.

The crude product was firstly analyzed as a 1 mg/mL sample in pure water and detected using a Shimadzu LCMS-8050 system. The analysis column was an Agilent

ZORBAX Eclipse SB-C18 (4.6 × 250 mm, 5 μm column; detection: UV at 220 nm; solvent A: 0.1% TFA in MeCN, solvent B: 0.1% TFA in water, with a linear gradient from 5%-90% solvent A concentration in 50 minutes; flow rate 1.0 mL/min; sample volume: 10 μL).

5

The target peak was eluted at 11.926 minutes and had the expected molecular weight, with a purity of 60.345%.

MS: m/z 2380.6

10

7.5 g of the crude product was then dissolved in 80 mL of pure water and purified using LC3000 semi-preparation equipment. The preparation column model was a Dubhe-C18 model (Hanbon Sci. & Tech. Co., Ltd., Jiangsu, China) (50*250 mm, 100Å column; detection: UV at 220 nm). The appropriate gradient for elution was calculated from LCMS detection step (Solvent A: 0.1% TFA in MeCN, solvent B: 0.1% TFA in water, with a linear gradient from 5%-20% solvent A concentration in 30 minutes; flow rate 60.0 mL/min;). The fractions were collected and analyzed using a Shimadzu LC-20 HPLC system (column as above, except with a linear gradient from 5%-30% solvent A concentration in 25 minutes).

20

Fractions with a purity of 98% were then mixed for an anion exchange step. This was achieved using a LC3000 semi-preparation equipment (preparation column model: Dubhe-C18 model (as above). The fractions were diluted one time with pure water and loaded to the column directly, after that the column was washed with 0.37% of ammonium acetate in pure water for about 20 minutes followed by pure water for another 20 minutes at the flow rate of 60 mL/min, then eluted with the following gradient (Solvent A: 0.1% HAc in MeCN, solvent B: 0.1% HAc in water, with a linear gradient from 5%-20% solvent A concentration in 30 minutes; flow rate 60.0 mL/min). The fractions were collected and analyzed using Shimadzu LC-20 HPLC system (column and conditions as above). Fractions with a purity of 98% were mixed and freeze-dried to give 3.06 g of the purified title compound.

35

Illustrative Example 2

40

(Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys)₂-Lys (SEQ ID No: 153)

Fmoc-Lys(fmoc)-Wang resin (9.9 g, GLS191010-41303, GL Biochem, Shanghai, China) was loaded into a glass reaction column.

The method was the same as described in Illustrative Example 1, except that Fmoc-Lys(Boc)-OH was coupled to the resin first followed by Fmoc-Dopa(Acetonide)-OH, Fmoc-Thr(tBu)-OH, Fmoc-4-Hyp(tBu)-OH, Fmoc-4-Hyp(tBu)-OH, Fmoc-Tyr(tBu)-OH, Fmoc-Ser(tBu)-OH, Fmoc-Pro-OH, Fmoc-Lys(Boc)-OH and Fmoc-Ala-OH, and the amounts of the amino acids, TBTU and DIPEA were doubled (by mols) compared to Illustrative Example 1.

10

MS: m/z 2508.8

Repeating essentially the same procedure gave a further batch of crude title compound (yield 7.89 g). Analysis showed a target peak that was eluted at 11.376 minutes with the expected molecular weight (*MS: m/z 2508.8*). The purity was 68.985%.

15

7.8 g of the crude product was then purified as described in Illustrative Example 1 above to give 2.57 g of pure title compound after freeze-drying.

20

MS: m/z 2508.8

Illustrative Example 3

{[(Dopa-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys)₂-Lys]₂-Lys}₂-Lys (SEQ ID No: 154)

25

Fmoc-Lys(fmoc)-Wang resin (9.9 g, GLS191010-41303, GL Biochem, Shanghai, China) was loaded into a glass reaction column.

The method was the same as described in Illustrative Example 2, except that Fmoc-Lys(Fmoc)-OH was coupled to the resin first followed by Fmoc-Lys(Fmoc)-OH, Fmoc-Lys(Boc)-OH, Fmoc-Tyr(tBu)-OH, Fmoc-Thr(tBu)-OH, Fmoc-4-Hyp(tBu)-OH, Fmoc-4-Hyp(tBu)-OH, Fmoc-Tyr(tBu)-OH, Fmoc-Ser(tBu)-OH, Fmoc-Pro-OH, Fmoc-Lys(Boc)-OH, Fmoc-Ala-OH and Fmoc-Dopa(Acetonide)-OH, and the amounts of the amino acids, TBTU and DIPEA were quadrupled (by mols) compared to Illustrative Example 2.

35

MS: m/z 11671.1

Repeating essentially the same procedure gave a further batch of crude title compound (yield 28.89 g). Analysis showed a target peak that was eluted at 11.896 minutes with the expected molecular weight (*MS: m/z 11671.1*). The purity was 29.985%.

- 5 28.8 g of the crude product was then purified as described in Illustrative Example 1 above to give 5.57 g of pure title compound after freeze-drying.

Illustrative Example 4

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 2)

10

The title peptide was synthesised using essentially the same procedure as that described in Illustrative Example 1 above, except that the appropriate amino acids were used in the appropriate peptide coupling sequences.

15 *MS: m/z 1183.3*

Example 1

(Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys)₂-Lys (SEQ ID No: 153) HA Conjugate

- 20 A 1% HA solution (50 mM of carboxyl groups) was made by dissolving HA-EP2 (1 g, 19072911, Furida Biotech., Shandong, China) in 100 mL of pure water for 8 hours.

DMTMM (275 mg, Sigma) and the product of Example 2 (22 mg) were dissolved in 20 mL of pure water to make a DMTMM (50mM) and peptide (2.5mM of amino groups) solution. 20 mL of the 1% HA solution was added to the 20 mL DMTMM and peptide solution and stirred at room temperature for one day.

After one day, 0.1 g of the reacted gel was taken and mixed well with 0.4 mL 96% ethanol. The sample was then centrifuged at 15000 rpm for 15 minutes. The supernatant was taken and detected using a Shimadzu LC-20 HPLC system equipped with LC-20AT binary pump, a DGU-20A5 degasser, a SIL-20AC autosampler, a CTO-20AC column oven, and a SPD-M20A photodiode array detector (Shimadzu, Japan).

The sample was analysed on an Agilent SB-C18 column (5 µm, 4.6* 250 mm; Agilent, USA) under the following conditions:

- 35
- the mobile phase: 0.1% TFA in acetonitrile (solvent A) and 0.1% TFA in water (solvent B);
 - linear gradient program: 0-25 min, 10%-35% solvent A;
 - flow rate: 1.0 mL/min;

- column temperature: 30 °C;
- the PDA detector recorded UV spectra in the range of from 190 nm to 400 nm;
- the HPLC chromatogram was monitored at 220 nm.

5 The concentration of the product of Example 2 in the supernatant (un-coupled free peptide) was calculated according its standard curve to be 0.052 mg/mL. Thus, about 0.29 mg/mL ($=22/40-0.052*5$) of peptide was coupled to HA-EP2, which is about 11.6 mg ($=0.29*40$) peptide per 200 mg ($0.2g=20*1\%$) HA-EP2.

10 Products are isolated by precipitation by dropwise addition of 96% ethanol (100 mL) to the reaction mixture. The white powder is obtained and thoroughly washed with water:ethanol 1:4, ethanol 96% and finally with absolute ethanol. The product is dried under vacuum for 3 days at 38°C.

15 Example 2

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 2) HA Conjugate

HA (200 mg, HA-EP2, Bloomage Biotechnology Co. Ltd., Jinan, China) was taken in a test tube, 2 mL of 0.2M NaOH was added to dissolve the HA powder and mixed well.

20

The peptide (SEQ ID No: 2) (220 mg, P200921, USUN Pharmaceutical Co. Ltd., Jiangyin, China) was taken into another test tube. 1mL of 0.4M HCl was used to dissolve the peptide powder, DMTMM (275 mg, Aladdin, Shanghai, China) was then added to dissolve and mixed well.

25

The mixture was then transferred to the HA solution's tube and mixed well, the test tube was then covered, heated at 35°C, and let to react for about 16 hours.

30 After 16 hours, the solution became a colorless rigid gel. The gel was removed and cut with 20 mesh screen once. The cut gel was then immersed into 100 mL HA buffer (which contained 9.0 g NaCl, 30 mg KH_2PO_4 and 140 mg $\text{Na}_2\text{HPO}_4 \cdot 12 \text{H}_2\text{O}$ in a total volume of 1 L pure water) for one hour. The gel swelled after soaking. The HA buffer was removed by filtration and the gel was cut with 200 mesh screen three times to make small and even particles.

35

The small particles were then immersed into 100 mL HA buffer for 1 hour to remove the condensation agent (DMTMM) (dialysis-like step), the HA buffer was then removed

by filtration. This step was repeated for additional 7 times to exhaustively remove DMTMM. The smaller particles were obtained at a concentration of about 30 mg/g.

HA (10 mg, HA-EP2, Bloomage Biotechnology Co. Ltd., Jinan, China) was dissolved
5 into 1 mL HA buffer, 1 g of the prepared small particles was then added and mixed well. The final product was obtained at 20 mg/g.

The obtained product was then filled in the final packing material and underwent moist heat sterilization.

10

The obtained product can be used for filling out wrinkles and folds.

Example 3

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 2) HA Conjugate Combined
15 with Celecoxib

The HA conjugate is obtained as described in Example 2.

The HA conjugate and celecoxib are mixed according to the mass ratio of 10:1 and
20 stirred to uniform. The obtained product can be used for intra-articular injection to relieve the symptoms of osteoarthritis and relieve pain.

Example 4

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 2) HA Conjugate Combined
25 with Diclofenac Sodium

The HA conjugate is obtained as described in Example 2.

The HA conjugate and diclofenac sodium are mixed according to the mass ratio of 5:1
30 and stirred to uniform. The obtained product can be used for intra-articular injection as an analgesic and/or anti-inflammatory drug.

35

Example 5

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 2) HA Conjugate Combined with Rifampicin

The HA conjugate is obtained as described in Example 2.

5

The HA conjugate and Rifampicin are mixed according to the mass ratio of 20:1 and stirred to uniform. The obtained product can be used in ophthalmic surgery as an anti-inflammatory, detumescence and/or analgesic.

10 Example 6

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 2) HA Conjugate Against Bleeding

15 A 49-years old women had allergic rhinitis for about 5 years. She was subject to seasonal allergy every spring and autumn. She was prescribed Fluticasone Propionate Nasal Spray to control the symptoms but started to have dry nose since the previous winter. She had nose bleeds after constant usage of fluticasone propionate nasal spray. When she had another allergy attack in Spring, she used Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 2) HA conjugate spray (comprising small particles of the conjugate prepared according to Example 2 above (0.5 mg) dissolved in 1 mL HA buffer in addition to the prescribed drug. She used the conjugate spray 4-5 times during the day. The bleeding stopped and the dry nose felt much better. This indicated that the title compound can be used to relief dry nose symptoms.

25 Example 7

{[(Dopa-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys)₂-Lys]₂-Lys}₂-Lys (SEQ ID No. 154) HA gel

30 A gel comprising {[(Dopa-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys)₂-Lys]₂-Lys}₂-Lys (SEQ ID No. 154) was made by mixing an appropriate amount of the peptide component with methyl cellulose (2.2 g; Shandong Guangda Technology Development Co., Ltd., ShanDong, China) and purified water (75.3 g), and stirring until a homogeneous colloidal suspension was formed. Then, 0.5 g of HA powder (HA-EP2, Bloomage Biotechnology Co. Ltd., Jinan, China), glycerin (11 g) and propanediol 11 g (both from Sinopharm Chemical Reagent Co. Ltd.) were added to the methyl cellulose/water mixture, and the resultant mixture was quickly stirred for 5 minutes to obtain the title product.

The product of Example 7 is referred to hereinafter as 'HA gel'.

Example 8

Mouse Ear Swelling Model

5

20 healthy male BALB/c mice of 6-8 weeks of age and average body weight of 18-25 g were supplied by Changzhou Cvens Experimental Animal Co., Ltd. and housed and cared for about for 1 week prior to the experiment. The housing temperature was 25-27°C with 74% humidity, alternating 12-hour periods of light and darkness, and free access to food and water. The mice were randomly divided into 4 groups as described in Table 1 below, with 5 mice in each group.

10

The left ear of each mouse was used as autologous control. The right ear of each mouse was treated by various different treatments, as summarized in Table 1 below. 20 µL of xylene (Shanghai Aladdin Bio-Chem Technology Co., Ltd., Shanghai, China) was applied to the right ear of each mouse, both inside and outside. The ear started to swell in about 4 minutes. Then, 40 µL of treatments or vehicle were applied to the right ears in each group. The mice were put back into their cages.

15

Dexamethasone Acetate Cream (DEX, Tianjin Jinyao Pharmaceutical Co., Ltd.) was used as positive control. The HA gel was synthesized as described in Example 7 above.

20

Table 1

Group	Drug concentration	Drug administration on right ear	Total amount of drugs (µg/mouse)
Model	/	Xylene	/
Vehicle	/	Xylene + gel containing HA (without peptide)	/
DEX	10 µg/µL	Xylene + Dex cream	400
HA gel	0.5 mg/g	Xylene + 'HA gel'	40

25

The mice were sacrificed by cervical dislocation after 40 minutes. The left and right ears were cut off. A skin pouch (Electron Microscopy Sciences, Hatfield, PA, USA) with a diameter of 8 mm was used to take a piece of the ear from the same site of both ears. The weights were recorded, and the swelling rates were calculated as follows:

30

Swelling rate = (right ear weight - left ear weight) / left ear weight × 100%

and the results showed in Table 2 below.

Table 2

	Model	Vehicle	DEX	HA gel
Mean	97.4	97%	45%	50%
SD	0.03	0.35	0.19	0.25

5

The above results show that HA gel could reduce the xylene induced swelling.

Claims

1. A conjugate formed between one or more linear polysaccharide chains and a peptide, which peptide component is selected from one or more of the peptide components (a), (b), (c) or (d) as defined below:


(a) a peptide component of formula I,



wherein:

A and B independently represent Z or $A^1-Q^1-B^1$;

- 10 Q represents a structural fragment of formula II,



wherein:


the squiggly lines represent points of attachment of Q to A and/or B; and

m represents an integer 1 to 4;

- 15 A^1 and B^1 independently represent Z or $A^2-Q^2-B^2$;

A^2 and B^2 independently represent Z or $Z-Q^3-Z$;

Q^1 , Q^2 and Q^3 independently represent structural fragments of formula III,



wherein:

- 20 the squiggly lines adjacent to the NH groups represent the points of attachment of Q^1 , Q^2 and Q^3 to A^1 and/or B^1 , A^2 and/or B^2 , and Z, respectively; and the squiggly line adjacent to the O atom represents the point of attachment of Q^1 , Q^2 and Q^3 to Q, Q^1 and Q^2 , respectively; and m is as defined above;

on each occasion that it is employed, Z represents a peptide component of the amino acid sequence:

- 25 $[W-Lys-X^1-Ser-U-X^2-Y]_n-W-Lys-X^1-Ser-U-X^2-Y---$ (SEQ ID No: 3)

wherein:

the dashed line represents the point of attachment of Z to the rest of the molecule;

n represents 0 or an integer 1 to 4; and,

on each occasion that they are employed:

W is absent or represents a 1 or 2 amino acid sequence, in which the amino acids are
 5 selected from one or more of the group Lys, Ala, DOPA and a 3,4-dihydrocinnamic acid
 (HCA) residue, provided that, when present, the HCA residue is located at the N-
 terminus of the peptide sequence Z;

X¹ represents Pro, Hyp or diHyp;

U represents Tyr, DOPA or a single bond;

10 X² represents Ser, Pro, Hyp or diHyp;

Y represents a single bond or represents a 1 to 5 amino acid sequence, in which the
 amino acids are selected from one or more of the group Lys, Ala, Pro, Hyp, diHyp, Thr,
 DOPA and Tyr; or

(b) a peptide component of the amino acid sequence:

15 [Ala-Lys-X¹-Ser-U-X²-Y]_p-Ala-Lys-X¹-Ser-U-X²-Y-G (SEQ ID No: 4)

wherein:

p represents an integer 1 to 4;

G is absent or represents DOPA or dopamine; and

X¹, U, X² and Y are as defined above; or

20 (c) a peptide component of the amino acid sequence:

W-Lys-X¹-Ser-U-X²-Y-G (SEQ ID No: 5)

wherein W, X¹, U, X², Y and G are as defined above; or

(d) a peptide component of the amino acid sequence:

K-W¹-Lys-X¹-Ser-U¹-X²-Y¹-I-J (SEQ ID No: 6)

25 wherein K represents an optional N-terminal HCA group;

W¹ may be absent or represents a 1 or 2 amino acid sequence, in which the amino
 acids are selected from one or more of the group Ser, Lys, Ala and DOPA;

U¹ represents Tyr, DOPA or a single bond;

Y¹ represents a single bond or represents Y;

30 I represents Pro, Hyp, diHyp, Thr, DOPA or Tyr; and

J represents Lys or is absent; and

X¹, X² and Y are as defined above,

as well as regioisomers, stereoisomers, and pharmaceutically- or cosmetically-
 acceptable salts of said conjugate.

35

2. A conjugate as claimed in Claim 1, wherein the polysaccharide chain is hyaluronic
 acid.

3. A conjugate as claimed in in Claim 1 or 2, wherein X^1 represents Pro.
4. A conjugate as claimed in in any one of the preceding claims, X^2 represents Pro or Hyp.
5. A conjugate as claimed in any one of the preceding claims, wherein W represents HCA, HCA-Ala-, Ala, DOPA, Lys-Ala or DOPA-Ala-.
6. A conjugate as claimed in Claim 5, wherein W represents HCA, HCA-Ala-, DOPA, or DOPA-Ala-.
7. A conjugate as claimed in any one of the preceding claims, wherein Y represents a 4 amino acid sequence, in which the amino acids are selected from one or more of the group Lys, Ala, Hyp, Thr, DOPA and Tyr.
8. A conjugate as claimed in Claim 7, wherein Y represents an amino acid sequence selected from the group -Hyp- Y^1 - Y^2 -Lys- and -Thr- Y^1 - Y^2 -Lys-, wherein Y^1 and Y^2 are each independently selected from the group Ala, Hyp, Thr, DOPA and Tyr.
9. A conjugate as claimed in Claim 7, wherein the amino acid sequence defined by Y is selected from the group -Pro-Thr-DOPA-Lys-, -Pro-Thr-Tyr-Lys-, -Thr-Tyr-Pro-Lys-, -Thr-DOPA-Pro-Lys-, -Hyp-Thr-Tyr-Lys-, -Hyp-Thr-Tyr-Lys-, -Hyp-Thr-DOPA-Lys-, -Hyp-Thr-Ala-Lys-, -Thr-Tyr-Hyp-Lys-, -Thr-DOPA-Hyp-Lys-, -Thr-Ala-Hyp-Lys-, -Hyp-Thr-, -Thr-Tyr-, -Pro-Thr-, -Thr-DOPA-, -Thr-Tyr-Lys-, -Tyr-Pro-Lys-, -DOPA-Pro-Lys, -Hyp-Thr-Tyr-, -Thr-Tyr-Hyp-Lys-DOPA-, -Hyp-Thr-Tyr-Hyp-Lys and -Hyp-Thr-DOPA.
10. A conjugate as claimed in any one of the preceding claims, wherein m represents 4.
11. A conjugate as claimed in any one of the preceding claims, wherein A and B both represent Z, or both represent A^1 - Q^1 - B^1 .
12. A conjugate as claimed in any one of the preceding claims, wherein A^1 and B^1 both represent Z, or both represent A^2 - Q^2 - B^2 .
13. A conjugate as claimed in any one of the preceding claims, wherein A^2 and B^2 both represent Z, or both represent Z- Q^3 -Z.
14. A conjugate as claimed in any one of the preceding claims, wherein n is 0.
15. A conjugate as claimed in any one of the preceding claims, wherein G is absent.

16. A conjugate as claimed in Claim 15, wherein the peptide component as defined under (b) comprises the amino acid sequence:

Ala-Lys-Pro-Ser-Tyr-Ser-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Ser-Hyp-Thr-Tyr-Lys (SEQ ID No: 7);

5 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 8);

Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 9);

10 Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 10);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-

15 DOPA-Lys (SEQ ID No: 11);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 12);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-

20 Hyp-Thr-DOPA-Lys (SEQ ID No: 13); or

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 14).

25 17. A conjugate as claimed in any one of Claims 1 to 15, wherein U represents Tyr and/or W represents Ala-.

18. A conjugate as claimed in Claim 17, wherein Z in the peptide component defined under (a) is selected from the group:

30 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 2);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 1);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 15);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 16);

Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys-DOPA--- (SEQ ID No: 17);

35 Ala-Lys-Pro-Ser-Tyr-Ser-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Ser-Hyp-Thr-Tyr-Lys-- (SEQ ID No: 18); and

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 19).

19. A conjugate as claimed in any one of Claims 1 to 15, wherein U represents Tyr and/or W represents Lys-Ala-.

5

20. A conjugate as claimed in Claim 19, wherein Z in the peptide component defined under (a) is selected from the group:

Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr--- (SEQ ID No: 37);

Lys-Ala-Lys-Hyp-Ser-Tyr-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 38); and

10 Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 39).

21. A conjugate as claimed in any one of Claims 1 to 15, wherein U represents Tyr and/or W represents HCA-, HCA-Ala-, DOPA- or DOPA-Ala-.

15 22. A conjugate as claimed in Claim 21, wherein Z in the peptide component defined under (a) is selected from the group:

HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 45);

HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 46);

HCA-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Ala-Lys--- (SEQ ID No: 47);

20 HCA-Lys-Pro-Ser-Tyr-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 48);

DOPA-Lys-Pro-Ser-Tyr-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 49);

DOPA-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Ala-Lys--- (SEQ ID No: 50);

DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 51); and

DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 52).

25

23. A conjugate as claimed in any one of Claims 1 to 15, wherein U represents DOPA and/or W represents Ala or Lys-Ala-.

24. A conjugate as claimed in Claim 23, wherein Z in the peptide component defined under (a) is selected from the group:

30

Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 65);

Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 66);

Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 67);

Lys-Ala-Lys-Hyp-Ser-DOPA-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 68);

35 Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 69); and

Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-

DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-

Hyp-Hyp-Thr-DOPA-Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 70).

25. A conjugate as claimed in any one of Claims 1 to 15, wherein U represents
5 DOPA and/or W represents HCA-, HCA-Ala-, DOPA- or DOPA-Ala-.

26. A conjugate as claimed in Claim 25, wherein Z is selected from the group:
HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 94);
HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 95);
10 HCA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 96);
HCA-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Ala-Lys--- (SEQ ID No: 96);
DOPA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 98);
DOPA-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Ala-Lys--- (SEQ ID No: 99);
DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 100);
15 DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 101); and
DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 102).

27. A conjugate as claimed in any one of the preceding claims, wherein, in the
peptide component defined under (a), A and B both represent Z, one or both Z groups
20 represent:

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 1);
Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 2);
Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 15);
Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 16);
25 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 45);
HCA-Lys-Pro-Ser-Tyr-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 48);
DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 51);
Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 65);
Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 67);
30 HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 94);
DOPA-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Ala-Lys--- (SEQ ID No: 99); or
DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 100),
and
Q represents a Lys fragment.

35

28. A conjugate as claimed in any one of Claims 1 to 26, wherein, in the peptide
component defined under (a), A and B both represent A¹-Q¹-B¹, A¹ and B¹ both
represent Z, one or both Z groups represent:

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 1);
 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 2);
 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 15);
 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 16);
 5 ;Lys-Ala-Lys-Hyp-Ser-Tyr-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 38);
 Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA--- (SEQ ID No: 39);
 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 45);
 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 46);
 DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 51);
 10 Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 65);
 Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 66);
 HCA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 96);
 DOPA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys--- (SEQ ID No: 98); or
 DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 100,
 15 and
 Q¹ represents a Lys fragment.

29. A conjugate as claimed in any one of Claims 1 to 26, wherein, in the peptide component defined under (a), A and B both represent A¹-Q¹-B¹, A¹ and B¹ both
 20 represent A²-Q²-B², A² and B² both represent Z, one or both Z groups represent:
 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 1);
 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 2);
 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 15);
 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 16);
 25 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 45);
 DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys--- (SEQ ID No: 51);
 DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 52);
 Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 66); or
 DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys--- (SEQ ID No: 100),
 30 and
 Q¹ and Q² both represent Lys fragments.

30. A conjugate as claimed in any one of Claims 1 to 26, wherein, in the peptide component defined under (a), A and B both represent A¹-Q¹-B¹, A¹ and B¹ both
 35 represent A²-Q²-B², A² and B² both represent Z-Q³-Z, one or both Z groups represent:
 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys--- (SEQ ID No: 1); or
 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys--- (SEQ ID No: 2), and
 Q¹, Q² and Q³ all represent Lys fragments.

31. A conjugate as claimed in Claim 17, wherein the peptide component defined under (c) comprises the amino acid sequence:

- Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys-DOPA (SEQ ID No: 20);
- 5 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys-Dopamine (SEQ ID No: 21);
- Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-Dopamine (SEQ ID No: 22);
- Ala-Lys-Pro-Ser-Tyr-Pro-Pro-Thr-DOPA-Lys (SEQ ID No: 23);
- Ala-Lys-Pro-Ser-Tyr-Pro-Thr-Tyr-Pro-Lys (SEQ ID No: 24);
- Ala-Lys-Pro-Ser-Tyr-Pro-Thr-DOPA-Pro-Lys (SEQ ID No: 25);
- 10 Ala-Lys-Pro-Ser-Tyr-Pro-Hyp-Thr-Tyr-Lys (SEQ ID No: 26);
- Ala-Lys-Pro-Ser-Tyr-Pro-Hyp-Thr-DOPA-Lys (SEQ ID No: 27);
- Ala-Lys-Pro-Ser-Tyr-Hyp-Pro-Thr-Tyr-Lys (SEQ ID No: 28);
- Ala-Lys-Pro-Ser-Tyr-Hyp-Pro-Thr-DOPA-Lys (SEQ ID No: 29);
- Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 30);
- 15 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 31);
- Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys-DOPA (SEQ ID No: 32);
- Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-DOPA (SEQ ID No: 33);
- Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys-Dopamine (SEQ ID No: 34);
- Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys-DOPA (SEQ ID No: 35); or
- 20 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys-Dopamine (SEQ ID No: 36).

32. A conjugate as claimed in Claim 19, wherein the peptide component defined under (c) comprises the amino acid sequence:

- Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA (SEQ ID No: 40);
- 25 Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Dopamine (SEQ ID No: 41); and
- Lys-Ala-Lys-Hyp-Ser-Tyr-Hyp-Hyp-Thr-DOPA (SEQ ID No: 42);
- Lys-Ala-Lys-Hyp-Ser-Tyr-Hyp-Hyp-Thr-Tyr (SEQ ID No: 43); or
- Lys-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr (SEQ ID No: 44).

30 33. A conjugate as claimed in Claim 21, wherein the peptide component defined under (c) comprises the amino acid sequence:

- DOPA-Lys-Pro-Ser-Tyr-Hyp-Thr-Ala-Hyp-Lys (SEQ ID No: 53);
- DOPA-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Ala-Lys (SEQ ID No: 54);
- DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 55);
- 35 DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 56);
- DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 57);
- DOPA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 58);
- HCA-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Ala-Lys (SEQ ID No: 59);

- HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 60);
 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 61);
 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 62);
 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 63); or
 5 HCA-Lys-Pro-Ser-Tyr-Hyp-Thr-Ala-Hyp-Lys (SEQ ID No: 64).

34. A conjugate as claimed in Claim 23, wherein the peptide component defined under (c) comprises the amino acid sequence:

- Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Dopamine (SEQ ID No: 71);
 10 Lys-Ala-Lys-Hyp-Ser-DOPA-Hyp-Hyp-Thr-DOPA (SEQ ID No: 72);
 Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA (SEQ ID No: 73);
 Ala-Lys-Pro-Ser-DOPA-Pro-Pro-Thr-Tyr-Lys (SEQ ID No: 74);
 Ala-Lys-Pro-Ser-DOPA-Pro-Pro-Thr-DOPA-Lys (SEQ ID No: 75);
 Ala-Lys-Pro-Ser-DOPA-Pro-Thr-Tyr-Pro-Lys (SEQ ID No: 76);
 15 Ala-Lys-Pro-Ser-DOPA-Pro-Thr-DOPA-Pro-Lys (SEQ ID No: 77);
 Ala-Lys-Pro-Ser-DOPA-Pro-Hyp-Thr-Tyr-Lys (SEQ ID No: 78);
 Ala-Lys-Pro-Ser-DOPA-Pro-Hyp-Thr-DOPA-Lys (SEQ ID No: 79);
 Ala-Lys-Pro-Ser-DOPA-Hyp-Pro-Thr-Tyr-Lys (SEQ ID No: 80);
 Ala-Lys-Pro-Ser-DOPA-Hyp-Pro-Thr-DOPA-Lys (SEQ ID No: 81);
 20 Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 82);
 Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 83);
 Lys-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Tyr (SEQ ID No: 84);
 Lys-Ala-Lys-Hyp-Ser-DOPA-Hyp-Hyp-Thr-Tyr (SEQ ID No: 85);
 Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Tyr-Lys-DOPA (SEQ ID No: 86);
 25 Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-DOPA (SEQ ID No: 87);
 Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Tyr-Lys-Dopamine (SEQ ID No: 88);
 Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys-Dopamine (SEQ ID No: 89);
 Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-Tyr-Hyp-Lys-DOPA (SEQ ID No: 90);
 Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys-DOPA (SEQ ID No: 91);
 30 Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-Tyr-Hyp-Lys-Dopamine (SEQ ID No: 92); or
 Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys-Dopamine (SEQ ID No: 93).

35. A conjugate as claimed in Claim 25, wherein the peptide component defined under (c) comprises the amino acid sequence:

- 35 DOPA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys (SEQ ID No: 103);
 DOPA-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Ala-Lys (SEQ ID No: 104);
 DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 105);
 DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 106);

- DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 107);
HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 108);
HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 109);
HCA-Lys-Pro-Ser-DOPA-Hyp-Thr-Ala-Hyp-Lys (SEQ ID No: 110);
5 HCA-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Ala-Lys (SEQ ID No: 111);
DOPA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 112);
HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 113); or
HCA-Ala-Lys-Pro-Ser-DOPA-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 114).
- 10 36. A conjugate as claimed in any one of Claims 1 to 4 or 7 to 9, wherein, in the peptide component defined under (d), W^1 represents Ala or Ser, or is absent.
37. A conjugate as claimed in any one of Claims 1 to 4 or 7 to 9 or 36, wherein, in the peptide component defined under (d), X^2 represents Pro, Hyp or diHyp.
- 15 38. A conjugate as claimed in any one of Claims 1 to 4 or 7 to 9, 36 or 37, wherein, in the peptide component defined under (d), when K is not present, W^1 represents Ala or is absent and J represents Lys, then I represents Pro, Hyp, diHyp or Thr.
- 20 39. A conjugate as claimed in any one of Claims 1 to 4 or 7 to 9, 36 to 38, wherein, in the peptide component defined under (d), in the sequence defined by Y^1 :
the amino acid DOPA, Thr or Tyr is linked to I; and/or
the amino acid Pro, Hyp or Thr is linked to X^2 .
- 25 40. A conjugate as claimed in any one of Claims 1 to 4, 7 to 9 or 36 to 39, wherein, in the peptide component defined under (d), K is absent.
41. A conjugate as claimed in Claim 40, wherein the peptide component defined under (d) comprises the amino acid sequence:
- 30 Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 115);
Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 116);
Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 117);
Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 118);
Ser-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Lys (SEQ ID No: 119);
35 Ser-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 120);
Ser-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 121);
Ser-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 122);
Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 123);

Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 124);
 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 125); or
 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 126).

- 5 42. A conjugate as claimed in any one of Claims 1 to 4, 7 to 9, 36, 37, 39 or 40, wherein, in the peptide component defined under (d), J is absent.

43. A conjugate as claimed in Claim 42, wherein the peptide component defined under (d) comprises the amino acid sequence:

- 10 Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Hyp (SEQ ID No: 127);
 Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Hyp (SEQ ID No: 128);
 Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp (SEQ ID No: 129);
 Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp (SEQ ID No: 130);
 Ser-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr (SEQ ID No: 131);
 15 Ser-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp (SEQ ID No: 132);
 Ser-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp (SEQ ID No: 133);
 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA-Hyp (SEQ ID No: 134);
 Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Hyp (SEQ ID No: 135);
 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-DOPA-Hyp (SEQ ID No: 136);
 20 Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp (SEQ ID No: 137);
 Ser-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-DOPA (SEQ ID No: 138);
 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr (SEQ ID No: 146);
 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp (SEQ ID No: 147);
 Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp (SEQ ID No: 148);
 25 Ala-Lys-Pro-Ser-Pro-Thr-Tyr-Pro (SEQ ID No: 149);
 Ala-Lys-Hyp-Ser-Tyr-Hyp-Thr-Tyr-Hyp (SEQ ID No: 150);
 Ala-Lys-Hyp-Ser-DOPA-Hyp -Thr-DOPA-Hyp (SEQ ID No: 151); or
 Ala-Lys-Hyp-Ser-Tyr-Hyp-Thr-DOPA-Hyp (SEQ ID No: 152).

- 30 44. A conjugate as claimed in any one of Claims 1 to 4, 7 to 9, 36 to 39 or 42, wherein, in the peptide component defined under (d), K is an N-terminal HCA group.

45. A conjugate as claimed in any one of Claim 44, wherein the peptide component defined under (d) comprises the amino acid sequence:

- 35 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Hyp-Thr-Tyr-Lys (SEQ ID No: 139); or
 HCA-Ala-Lys-Pro-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 140).

46. A conjugate as claimed in any one of Claims 1 to 4, 7 to 9, 36 to 40 or 44, wherein,

in the peptide component defined under (d), W¹ is Ala and J is Lys.

47. A conjugate as claimed in Claim 46, wherein the peptide component defined under (d) comprises the amino acid sequence:

- 5 Ala-Lys-Pro-Ser-DOPA-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 141);
Ala-Lys-Pro-Ser-Pro-Thr-Tyr-Pro-Lys (SEQ ID No: 142);
Ala-Lys-Hyp-Ser-Tyr-Hyp-Thr-Tyr-Hyp-Lys (SEQ ID No: 143);
Ala-Lys-Hyp-Ser-DOPA-Hyp -Thr-DOPA-Hyp-Lys (SEQ ID No: 144); or
Ala-Lys-Hyp-Ser-Tyr-Hyp-Thr-DOPA-Hyp-Lys (SEQ ID No: 145).

10

48. A conjugate as defined in any one of the preceding claims, for use in human or animal medicine.

49. A conjugate as defined in any one of Claims 1 to 47, for use as a pharmaceutical.

15

50. A pharmaceutical formulation comprising a conjugate as defined in any one of Claims 1 to 47 and a pharmaceutically- or cosmetically-acceptable, adjuvant, diluent or carrier.

20 51. A pharmaceutical formulation as claimed in Claim 50 that is suitable for, adapted for, and/or packaged and presented for, topical administration, wherein the pharmaceutically- or cosmetically-acceptable adjuvant, diluent or carrier is a topical adjuvant, diluent or carrier.

25 52. A pharmaceutical formulation as claimed in Claim 50 or Claim 51, which is in the form of a gel, a spray, a cream, an ointment or a dry powder.

53. A pharmaceutical formulation as claimed in Claim 50 that is suitable for, adapted for, and/or packaged and presented for, administration by injection.

30

54. A pharmaceutical formulation as claimed in Claim 51 or Claim 53 in the use of the treatment of melanin pigmentation, wrinkles and/or arthritis.

35 55. A pharmaceutical formulation as claimed in any one of Claims 50 to 54, which further includes a, or a further, pharmaceutically-active ingredient.

56. A kit of parts comprising components:

(A) a conjugate as defined in any one of Claims 1 to 47, or a pharmaceutical formulation as defined in any one of Claims 50 to 54; and

(B) a pharmaceutical formulation including a, or a further, pharmaceutically-active ingredient in admixture with a pharmaceutically-acceptable adjuvant, diluent or

5 carrier,

which components (A) and (B) are each provided in a form that is suitable for administration in conjunction with the other.

57. A pharmaceutical formulation as claimed in Claim 55, or a kit of parts as claimed
10 in Claim 56 wherein the pharmaceutically-active ingredient is an anti-inflammatory agent, a pro-inflammatory agent, an antibiotic, an anti-bacterial and/or antiprotozoal agent, an antiviral agent, an anaesthetic and/or a wound recovery drug.

58. A pharmaceutical formulation or a kit of parts as claimed in Claim 57, wherein
15 the pharmaceutically-active ingredient is an antiinflammatory agent.

59. A conjugate as defined in any one of Claims 1 to 47, a formulation as claimed
in any one of Claims 50 to 55, 57 or 58, or a kit of parts as claimed in Claim 56, for
use in the treatment of inflammation, an inflammatory disorder, and/or of a disorder
20 characterised by inflammation.

60. The use of a conjugate as defined in any one of Claims 1 to 47, a formulation
as claimed in any one of Claims 50 to 55, 57 or 58, or a kit of parts as claimed in Claim
56, for the manufacture a medicament for the treatment of inflammation, an
25 inflammatory disorder, and/or of a disorder characterised by inflammation.

61. A method of treatment of inflammation, an inflammatory disorder, and/or of a
disorder characterised by inflammation, which method comprises the administration of
a conjugate as defined in any one of Claims 1 to 47, a formulation as claimed in any
30 one of Claims 50 to 55, 57 or 58, or a kit of parts as claimed in Claim 56, to a patient
in need of such treatment.

62. A conjugate, formulation or a kit of parts for use as claimed in Claim 59, a use
as claimed in Claim 60, or a method as claimed in Claim 61, wherein the inflammation,
35 inflammatory disorder, and/or disorder characterised by inflammation is selected from
the group of radiation vaginitis, fibrosis of the vagina and/or nasosinusitis.

63. A conjugate, formulation or kit of parts for use as claimed in Claim 59, a use as claimed in Claim 60, or a method as claimed in Claim 61, wherein the disorder characterised by inflammation is, or results in, a wound or a burn.

5 64. A conjugate, formulation or kit of parts for use, use or method as claimed in Claim 63, wherein the disorder resulting in a wound is haemorrhoids or ulcerative colitis.

65. A conjugate, formulation or kit of parts for use, use or method as claimed in any one of Claims 48 to 64 (as appropriate), wherein the conjugate(s) or salt thereof
10 is administered topically in the form of a topical formulation.

66. A conjugate, formulation or kit of parts for use, use or method as claimed in Claim 65 wherein the relevant condition is treated by way of direct topical administration to the skin.

15

67. A conjugate, formulation or kit of parts for use, use or method as claimed in Claim 65, wherein the relevant condition is treated by way of direct topical administration to a mucosal surface.

20 68. A conjugate, a formulation, or a kit of parts, for use, a use or a method as claimed in any one of Claims 48 to 68 (as appropriate), wherein the conjugate(s) is/are administered by oral, intravenous, cutaneous or subcutaneous, nasal, intramuscular, intraperitoneal, pulmonary or anorectal delivery.

25 69. A conjugate, a formulation, or a kit of parts, for use, a use or a method as claimed in any one of Claims 48 to 69 (as appropriate), wherein the conjugate as defined in any of Claim 1 to 47 acts as an excipient, as a medical device, or as a medical device component of a drug-medical device combination.

30 70. A conjugate, a formulation, or a kit of parts, for use, a use or a method as claimed in any one of Claims 48 to 67 (as appropriate), wherein the conjugate as defined in any of Claim 1 to 47 is crosslinked, before or after administration to a subject.

71. A conjugate as defined in Claim 70 for use as an adhesive or as a film-forming
35 material.

72. A conjugate for use as claimed in Claim 71, wherein the use is as a wound surface repair product, a wound surface protecting product, a medical biological

adhesive product, a medical coating product, an industrial coating product, a biochemical reagent, a medical product, a sterilization product or as a culture vessel for cell culture.

- 5 73. A conjugate for use as claimed in Claim 71 or Claim 72, wherein a film is formed over a skin or a mucosal wound surfaces to aid in recovery.

74. A conjugate for use as claimed in any one of Claims 71 to 73, wherein the use is in closure of a surgical incision, adhesion of a fractured bone, adhesion of a mucous
10 membrane or coating of a human body implant.

75. A pharmaceutical formulation comprising a conjugate as defined in Claim 71 and a pharmaceutically- or cosmetically-acceptable, adjuvant, diluent or carrier

- 15 76. A pharmaceutical formulation as claimed in Claim 75 that is suitable for, adapted for, and/or packaged and presented for, topical administration, wherein the pharmaceutically- or cosmetically-acceptable adjuvant, diluent or carrier is a topical adjuvant, diluent or carrier.

- 20 77. A pharmaceutical formulation as claimed in Claim 75 or Claim 76, which is in the form of a gel, a spray, a cream, an ointment or a dry powder.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN2020/133438

A. CLASSIFICATION OF SUBJECT MATTER

C07K 7/06(2006.01)i; A61K 38/08(2019.01)i; A61K 8/64(2006.01)i; A61P 29/00(2006.01)i; A61P 17/06(2006.01)i; A61P 17/10(2006.01)i; A61P 17/02(2006.01)i; A61P 17/00(2006.01)i; A61P 11/02(2006.01)i; A61P 11/00(2006.01)i; A61P 1/04(2006.01)i; A61P 33/12(2006.01)i; A61Q 19/00(2006.01)i

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07K; A61K; A61P; A61Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CNTEXT,CNABS,VEN,WOTXT,EPTXT,USTXT,PUBMED,BAIDU,ISI Web of Knowledge,CNKI,EBI,GENBANK: conjugate,inflammation,mfp,mefp,MAP,DOPA,3, 4-dihydrocinnamic acid,polysaccharide,hyaluronic acid,HCA, Ala-Lys-Pro-Ser-Tyr-Hyp, W-Lys-X1-Ser-U-X2-Y,SEQ ID No:1-152.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0359996 A2 (BIO-POLYMERS, INC.) 28 March 1990 (1990-03-28) Claims 1-26, Description page 3 line 45-page 6 line 50	1-50
Y	EP 0359996 A2 (BIO-POLYMERS, INC.) 28 March 1990 (1990-03-28) Claims 1-26, Description page 3 line 45-page 6 line 50	51-77
Y	CN 110312513 A (JIANGYIN USUN PHARMACEUTICAL CO., LTD.) 08 October 2019 (2019-10-08) Claims 1-26, Description paragraphs 8-109	51-77
Y	CN 110139662 A (JIANGYIN USUN PHARMACEUTICAL CO., LTD.) 16 August 2019 (2019-08-16) abstract, claims 1-15, description paragraphs 8-90	51-77
A	EP 2990412 A1 (National Cerebral and Cardiovascular Center, et al) 02 March 2016 (2016-03-02) the whole document	1-77
A	WO 2005/118831 A2 (NORTHWESTERN UNIVERSITY) 15 December 2005 (2005-12-15) the whole document	1-77



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
“A” document defining the general state of the art which is not considered to be of particular relevance	“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
“E” earlier application or patent but published on or after the international filing date	“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
“L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	“&” document member of the same patent family
“O” document referring to an oral disclosure, use, exhibition or other means	
“P” document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

18 February 2021

Date of mailing of the international search report

08 March 2021

Name and mailing address of the ISA/CN

National Intellectual Property Administration, PRC
6, Xitucheng Rd., Jimen Bridge, Haidian District, Beijing
100088
China

Authorized officer

LIU,Miao

Facsimile No. (86-10)62019451

Telephone No. 86-(10)-53961924

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN2020/133438

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2012/067837 A1 (AVON PRODUCTS, INC.) 24 May 2012 (2012-05-24) the whole document	1-77
A	LEE, M.S.et al. "Target-specific delivery of siRNA by stabilized calcium phosphate nanoparticles using dopa–hyaluronic acid conjugate" <i>Journal of Controlled Release</i> , Vol. 192, 01 July 2014 (2014-07-01), Pages 122-130	1-77
A	DALSIN, J.L.et al. "Mussel Adhesive Protein Mimetic Polymers for the Preparation of Nonfouling Surfaces" <i>J. AM. CHEM. SOC.</i> , Vol. 125, 31 December 2003 (2003-12-31), Pages 4253-4258	1-77

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
- a. ☐ forming part of the international application as filed:
- ☐ in the form of an Annex C/ST.25 text file.
- ☐ on paper or in the form of an image file.
- b. ☐ furnished together with the international application under PCT Rule 13*ter*.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
- c. ☒ furnished subsequent to the international filing date for the purposes of international search only:
- ☒ in the form of an Annex C/ST.25 text file (Rule 13*ter*.1(a)).
- ☐ on paper or in the form of an image file (Rule 13*ter*.1(b) and Administrative Instructions, Section 713).
2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **61,62-70(partial)**
because they relate to subject matter not required to be searched by this Authority, namely:

[1] The subject-matter of claim 61,62-70(partial) relates to a method of treatment of inflammation, an inflammatory disorder, and/or of a disorder characterised by inflammation, therefore does not warrant an international search according to the criteria set out in Rule 39.1(iv). However, the search has also been carried out based on the manufacture of a medicament for treating inflammation, an inflammatory disorder, and/or of a disorder characterized by inflammation.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/CN2020/133438

Patent document cited in search report			Publication date (day/month/year)	Patent family member(s)			Publication date (day/month/year)
EP	0359996	A2	28 March 1990	AT	104318	T	15 April 1994
				FI	893854	A	23 February 1990
				NO	175006	C	17 August 1994
				NO	175006	B	09 May 1994
				CN	1042162	A	16 May 1990
				NO	893350	D0	21 August 1989
				EP	0359996	A3	07 August 1991
				DE	68914559	T2	27 October 1994
				NO	893350	L	23 February 1990
				US	4908404	A	13 March 1990
				AU	618834	B2	09 January 1992
				DK	410889	A	23 February 1990
				NO	893350	A	23 February 1990
				JP	H02191629	A	27 July 1990
				FI	893854	A0	16 August 1989
				DE	68914559	D1	19 May 1994
				AU	4001489	A	22 February 1990
				DK	410889	D0	21 August 1989
				EP	0359996	B1	13 April 1994
CN	110312513	A	08 October 2019	AU	2018295944	A1	20 February 2020
				TW	201919633	A	01 June 2019
				KR	20200039677	A	16 April 2020
				WO	2019007356	A1	10 January 2019
				SG	11201911841P	A	30 January 2020
				JP	2020527134	A	03 September 2020
				CA	3068818	A1	10 January 2019
				EP	3648767	A1	13 May 2020
CN	110139662	A	16 August 2019	US	2020148722	A1	14 May 2020
				TW	201920232	A	01 June 2019
				JP	2020527135	A	03 September 2020
				KR	20200027493	A	12 March 2020
				WO	2019007355	A1	10 January 2019
				EP	3648785	A1	13 May 2020
EP	2990412	A1	02 March 2016	US	10351600	B2	16 July 2019
				JP	2014214140	A	17 November 2014
				US	2017327539	A1	16 November 2017
				CN	105263951	A	20 January 2016
				EP	2990412	B1	08 November 2017
				CN	105263951	B	10 December 2019
				JP	6202669	B2	27 September 2017
				EP	2990412	A4	28 December 2016
				WO	2014175445	A1	30 October 2014
				US	2016137694	A1	19 May 2016
WO	2005/118831	A2	15 December 2005	EP	1735456	A4	16 November 2011
				AU	2005250314	A1	15 December 2005
				WO	2005118831	A3	28 June 2007
				JP	2007527871	A	04 October 2007
				JP	5133048	B2	30 January 2013
				EP	1735456	A2	27 December 2006

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CN2020/133438

Patent document cited in search report			Publication date (day/month/year)	Patent family member(s)			Publication date (day/month/year)
				CA	2557330	A1	15 December 2005
WO	2012/067837	A1	24 May 2012	TW	201223557	A	16 June 2012
				MX	2013005346	A	03 July 2013
				EP	2640470	A1	25 September 2013
				EP	2640470	A4	17 February 2016
				JP	6010541	B2	19 October 2016
				CN	103140262	A	05 June 2013
				CA	2811274	A1	24 May 2012
				US	2013224269	A1	29 August 2013
				JP	2013543865	A	09 December 2013
				AR	083881	A1	27 March 2013
				BR	112013005304	A2	02 May 2017