

(19) **DANMARK**

(10) **DK/EP 3823970 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 D 487/04 (2006.01)** **A 61 P 37/00 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2024-02-05**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2024-01-03**
- (86) Europæisk ansøgning nr.: **19742204.1**
- (86) Europæisk indleveringsdag: **2019-07-22**
- (87) Den europæiske ansøgnings publiceringsdag: **2021-05-26**
- (86) International ansøgning nr.: **EP2019069610**
- (87) Internationalt publikationsnr.: **WO2020016452**
- (30) Prioritet: **2018-07-20 EP 18184607**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Grünenthal GmbH, Zieglerstrasse 6, 52078 Aachen, Tyskland**
- (72) Opfinder: **JAKOB, Florian, Bachstrasse 50, 52066 Aachen, Tyskland**
ALEN, Jo, Prins de Merodelaan 24, 3271 Averbode, Belgien
LUCAS, Simon, Berggartenstraße 36, 64732 Bad König, Tyskland
CRAAN, Tobias, Marienbader Straße 5, 63843 Niedernberg, Tyskland
KONETZKI, Ingo, Auf der Haag 23, 52156 Monschau, Tyskland
KLESS, Achim, Berensberger Winkel 14, 52072 Aachen, Tyskland
SCHUNK, Stefan, Maargasse 20, 52080 Aachen, Tyskland
RATCLIFFE, Paul, Im Vennbahnbogen 20, 52076 Aachen, Tyskland
WACHTEN, Sebastian, Aussemstraße 24, 52066 Aachen, Tyskland
CRUWYS, Simon, Nizzaallee 26, 52072 Aachen, Tyskland
- (74) Fuldmægtig i Danmark: **Budde Schou A/S, Dronningens Tværgade 30, 1302 København K, Danmark**
- (54) Benævnelse: **YDERLIGERE SUBSTITUERED E TRIAZOLOQUINOXALINDERIVATER**
- (56) Fremdragne publikationer:
US-A1- 2013 303 537

Description

[0001] The present invention relates to compounds according to general formulae (II) or (III), which act as modulators of the glucocorticoid receptor and can be used in the treatment and/or prophylaxis of disorders which are at least partially mediated by the glucocorticoid receptor.

[0002] Glucocorticoids (GC) exert strong anti-inflammatory, immunosuppressive and disease-modifying therapeutic effects mediated by the glucocorticoid receptor (GR). They have been widely used to treat inflammatory and immune diseases for decades and still represent the most effective therapy in those conditions. However, chronic GC treatment of inflammatory diseases is hampered by GC-associated adverse effects. These undesired side effects include insulin resistance, diabetes, hypertension, glaucoma, depression, osteoporosis, adrenal suppression and muscle wasting with osteoporosis and diabetes being the most severe ones from the physician's point of view (Hapgood JP. et al., Pharmacol Ther. 2016 Sep; 165: 93-113; Buttgereit F. et al, Clin Exp Rheumatol. 2015 Jul-Aug;33(4 Suppl 92):S29-33; Hartmann K. et al, Physiol Rev. 2016 Apr;96(2):409-47).

[0003] One example of an oral glucocorticoid is prednisone which is frequently prescribed for the treatment of several inflammatory disorders (De Bosscher Ket al., Trends Pharmacol Sci. 2016 Jan;37(1):4-16; Buttgereit F. et al., JAMA. 2016;315(22):2442-2458). As GC cause adrenal suppression, prednisolone withdrawal symptoms can be severe if the drug is discontinued abruptly when all the signs of the disease have disappeared. Thus gradual GC tapering to physiological doses is frequently part of treatment protocols to reduce the risk of relapse and other withdrawal symptoms (Liu D. et al., Allergy Asthma Clin Immunol. 2013 Aug 15;9(1):30). Therefore, there is high medical need for novel potent anti-inflammatory drugs with less adverse effects.

[0004] Recent research has focused on the development of partial agonists or selective glucocorticoid receptor modulators which activate the pathways for the inhibition of inflammation but avoid targeting the pathways that lead to the GC-associated adverse effects. Most of these effects have been demonstrated to be mediated by different GR-dependent genomic mechanisms termed transactivation and transrepression. The anti-inflammatory actions of GC are mainly attributable to the transrepression of inflammatory genes while certain side effects are predominantly mediated via transactivation of several genes. According to the nature of a ligand the GR can be selectively modulated in a specific conformation which favors transrepression over transactivation resulting in an improved therapeutic benefit (De Bosscher Ket al., Trends Pharmacol Sci. 2016 Jan;37(1):4-16). The concept of such dissociating ligands was already defined about two decades ago and several compounds have been identified and were evaluated in preclinical and clinical testing but none of them has as yet been approved for clinical use.


[0005] Compounds which are active as modulators of the glucocorticoid receptor are also known from WO 2009/035067 and WO 2017/034006.

[0006] It was an object of the present invention to provide novel compounds which are modulators of the glucocorticoid receptor and which preferably have advantages over the compounds of the prior art. The novel compounds should in particular be suitable for use in the treatment and/or prophylaxis of disorders or diseases which are at least partially mediated by the glucocorticoid receptor.

[0007] This object has been achieved by the subject-matter of the patent claims.

[0008] It was surprisingly found that the compounds according to the present invention are highly potent modulators of the glucocorticoid receptor.

[0009] The present invention relates to a compound according to general formulae (II) or (III):



wherein

R¹ represents H; C₁₋₁₀-alkyl; C₃₋₁₀-cycloalkyl; 3 to 7 membered heterocycloalkyl; aryl; or 5 or 6-membered heteroaryl; wherein C₃₋₁₀-cycloalkyl, 3 to 7 membered heterocycloalkyl, aryl and 5 or 6-membered heteroaryl can optionally be bridged via C₁₋₆-alkylene;

R² represents H; F; Cl; Br; I; CN; C₁₋₁₀-alkyl; C₃₋₁₀-cycloalkyl; O-C₁₋₁₀-alkyl; N(H)(C₁₋₁₀-alkyl), N(C₁₋₁₀-alkyl)₂; C(O)-C₁₋₁₀-alkyl; C(O)-O-C₁₋₁₀-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₁₀-alkyl); C(O)-N(C₁₋₁₀-alkyl)₂; O-C₃₋₁₀-cy-

cloalkyl; N(H)(C₃₋₁₀-cycloalkyl), N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); C(O)-C₃₋₁₀-cycloalkyl; C(O)-O-C₃₋₁₀-cycloalkyl; C(O)-N(H)(C₃₋₁₀-cycloalkyl) or C(O)-N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); wherein C₃₋₁₀-cycloalkyl can optionally be bridged via C₁₋₆-alkylene;

5 R³ represents H; F; Cl; Br; I; CN; C₁₋₁₀-alkyl; C₃₋₁₀-cycloalkyl; O-C₁₋₁₀-alkyl; N(H)(C₁₋₁₀-alkyl); N(C₁₋₁₀-alkyl)₂; C(O)-C₁₋₁₀-alkyl; C(O)-O-C₁₋₁₀-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₁₀-alkyl); C(O)-N(C₁₋₁₀-alkyl)₂; O-C₃₋₁₀-cycloalkyl; N(H)(C₃₋₁₀-cycloalkyl), N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); C(O)-C₃₋₁₀-cycloalkyl; C(O)-O-C₃₋₁₀-cycloalkyl; C(O)-N(H)(C₃₋₁₀-cycloalkyl) or C(O)-N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); wherein C₃₋₁₀-cycloalkyl can optionally be bridged via C₁₋₆-alkylene;

10 R⁴ represents F or Cl;

R⁵ and R⁶ represent independently from one another H or unsubstituted C₁₋₄-alkyl;

15 X represents N or NR⁷;

Z represents N, NR⁷ or CR⁹; with the proviso that when X represents NR⁷, Z represents Nor CR⁹; when X represents N, Z represents NR⁷;

20 R⁷ represents H or L-R⁸; wherein

L represents bond; S(O); S(O)₂; C₁₋₆-alkylene; C(O); C₁₋₆-alkylene-C(O); C(O)-O; C₁₋₆-alkylene-C(O)-O; C₁₋₆-alkylene-N(H)-C(O); C₁₋₆-alkylene-N(C₁₋₁₀-alkyl)-C(O); C₁₋₆-alkylene-N(H)-C(O)-O; C₁₋₆-alkylene-N(C₁₋₁₀-alkyl)-C(O)-O; O; NH or N(C₁₋₁₀-alkyl);

25 R⁸ represents C₁₋₁₀-alkyl; C₃₋₁₀-cycloalkyl or 3 to 7 membered heterocycloalkyl; wherein C₃₋₁₀-cycloalkyl and 3 to 7 membered heterocycloalkyl can optionally be bridged via C₁₋₆-alkylene;

R⁹ and R¹⁰ represent independently from one another H; F; Cl; Br; I; CN; C₁₋₁₀-alkyl; C₃₋₁₀-cycloalkyl; 3 to 7 membered heterocycloalkyl; S(O)-(C₁₋₁₀-alkyl); S(O)-(C₃₋₁₀-cycloalkyl); S(O)-(3 to 7-membered heterocycloalkyl); S(O)₂-(C₁₋₁₀-alkyl); S(O)₂-(C₃₋₁₀-cycloalkyl); S(O)₂-(3 to 7-membered heterocycloalkyl); P(O)-(C₁₋₁₀-alkyl)₂; P(O)(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); P(O)(C₁₋₁₀-alkyl)(3 to 7-membered heterocycloalkyl); P(O)-(O-C₁₋₁₀-alkyl)₂; P(O)(O-C₁₋₁₀-alkyl)(O-C₃₋₁₀-cycloalkyl); P(O)(O-C₁₋₁₀-alkyl)(O-(3 to 7-membered heterocycloalkyl)); O-C₁₋₁₀-alkyl; S-C₁₋₁₀-alkyl; N(H)(C₁₋₁₀-alkyl), N(C₁₋₁₀-alkyl)₂; C(O)-C₁₋₁₀-alkyl; C(O)-O-C₁₋₁₀-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₁₀-alkyl); C(O)-N(C₁₋₁₀-alkyl)₂; O-C₃₋₁₀-cycloalkyl; N(H)(C₃₋₁₀-cycloalkyl), N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); C(O)-C₃₋₁₀-cycloalkyl; C(O)-O-C₃₋₁₀-cycloalkyl; C(O)-N(H)(C₃₋₁₀-cycloalkyl); C(O)-N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); O-3 to 7-membered heterocycloalkyl; N(H)(3 to 7-membered heterocycloalkyl), N(C₁₋₁₀-alkyl)(3 to 7-membered heterocycloalkyl); C(O)-3 to 7-membered heterocycloalkyl; C(O)-O-(3 to 7-membered heterocycloalkyl); C(O)-N(H)(3 to 7-membered heterocycloalkyl) or C(O)-N(C₁₋₁₀-alkyl)(3 to 7-membered heterocycloalkyl); wherein C₃₋₁₀-cycloalkyl and 3 to 7 membered heterocycloalkyl can optionally be bridged via C₁₋₆-alkylene;

40 R¹¹ represents F; Cl; Br; I; CN; C₁₋₁₀-alkyl; O-C₁₋₁₀-alkyl; NO₂; OH, NH₂; C₃₋₁₀-cycloalkyl; 3 to 7-membered heterocycloalkyl; S(O)-(C₁₋₁₀-alkyl); S(O)-(C₃₋₁₀-cycloalkyl); S(O)-(3 to 7-membered heterocycloalkyl); S(O)₂-(C₁₋₁₀-alkyl); S(O)₂-(C₃₋₁₀-cycloalkyl); S(O)₂-(3 to 7-membered heterocycloalkyl); P(O)-(C₁₋₁₀-alkyl)₂; P(O)(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); P(O)(C₁₋₁₀-alkyl)(3 to 7-membered heterocycloalkyl); P(O)-(O-C₁₋₁₀-alkyl)₂; P(O)(O-C₁₋₁₀-alkyl)(O-C₃₋₁₀-cycloalkyl); P(O)(O-C₁₋₁₀-alkyl)(O-(3 to 7-membered heterocycloalkyl)); O-Ci-io-alkyl; N(H)(Ci-io-alkyl), N(C₁₋₁₀-alkyl)₂; C(O)-C₁₋₁₀-alkyl; C(O)-O-C₁₋₁₀-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₁₀-alkyl); C(O)-N(C₁₋₁₀-alkyl)₂; O-C₃₋₁₀-cycloalkyl; N(H)(C₃₋₁₀-cycloalkyl), N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); C(O)-C₃₋₁₀-cycloalkyl; C(O)-O-C₃₋₁₀-cycloalkyl; C(O)-N(H)(C₃₋₁₀-cycloalkyl); C(O)-N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); O-3 to 7-membered heterocycloalkyl; N(H)(3 to 7-membered heterocycloalkyl), N(C₁₋₁₀-alkyl)(3 to 7-membered heterocycloalkyl); C(O)-3 to 7-membered heterocycloalkyl; C(O)-O-(3 to 7-membered heterocycloalkyl); C(O)-N(H)(3 to 7-membered heterocycloalkyl) or C(O)-N(C₁₋₁₀-alkyl)(3 to 7-membered heterocycloalkyl); wherein C₃₋₁₀-cycloalkyl and 3 to 7 membered heterocycloalkyl can optionally be bridged via C₁₋₆-alkylene;

55 n represents 0, 1, 2 or 3;

wherein C₁₋₁₀-alkyl, C₁₋₄-alkyl and C₁₋₆-alkylene in each case independently from one another is linear or branched, saturated or unsaturated;

wherein C₁₋₁₀-alkyl, C₁₋₄-alkyl, C₁₋₆-alkylene, C₃₋₁₀-cycloalkyl and 3 to 7 membered heterocycloalkyl in each case independently from one another are unsubstituted or mono- or polysubstituted with one or more substituents selected from F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; CF₂Cl; CFCF₂; C(O)-C₁₋₆-alkyl; C(O)-OH; C(O)-OC₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; =O; OCF₃; OCF₂H; OCFH₂; OCF₂Cl; OCFCl₂; O-C₁₋₆-alkyl; O-C(O)-C₁₋₆-alkyl; O-C(O)-O-C₁₋₆-alkyl; O-(CO)-N(H)(C₁₋₆-alkyl); O-C(O)-N(C₁₋₆-alkyl)₂; O-S(O)₂-NH₂; O-S(O)₂-N(H)(C₁₋₆-alkyl); O-S(O)₂-N(C₁₋₆-alkyl)₂; NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; N(H)-C(O)-C₁₋₆-alkyl; N(H)-C(O)-O-C₁₋₆-alkyl; N(H)-C(O)-NH₂; N(H)-C(O)-N(H)(C₁₋₆-alkyl); N(H)-C(O)-N(C₁₋₆-alkyl)₂; N(C₁₋₆-alkyl)-C(O)-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-C(O)-O-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-C(O)-NH₂; N(C₁₋₆-alkyl)-C(O)-N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)-C(O)-N(C₁₋₆-alkyl)₂; N(H)-S(O)₂OH; N(H)-S(O)₂-C₁₋₆-alkyl; N(H)-S(O)₂-O-C₁₋₆-alkyl; N(H)-S(O)₂-NH₂; N(H)-S(O)₂-N(H)(C₁₋₆-alkyl); N(H)-S(O)₂-N(C₁₋₆-alkyl)₂; N(C₁₋₆-alkyl)-S(O)₂-OH; N(C₁₋₆-alkyl)-S(O)₂-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-S(O)₂-O-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-S(O)₂-NH₂; N(C₁₋₆-alkyl)-S(O)₂-N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)-S(O)₂-N(C₁₋₆-alkyl)₂; SCF₃; SCF₂H; SCFH₂; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; S(O)₂-OH; S(O)₂-O-C₁₋₆-alkyl; S(O)₂-NH₂; S(O)₂-N(H)(C₁₋₆-alkyl); S(O)₂-N(C₁₋₆-alkyl)₂; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; phenyl; 5 or 6-membered heteroaryl; O-C₃₋₆-cycloalkyl; O-(3 to 6-membered heterocycloalkyl); O-phenyl; O-(5 or 6-membered heteroaryl); C(O)-C₃₋₆-cycloalkyl; C(O)-(3 to 6-membered heterocycloalkyl); C(O)-phenyl; C(O)-(5 or 6-membered heteroaryl); S(O)₂-(C₃₋₆-cycloalkyl); S(O)₂-(3 to 6-membered heterocycloalkyl); S(O)₂-phenyl or S(O)₂-(5 or 6-membered heteroaryl);

wherein aryl and 5 or 6-membered heteroaryl in each case independently from one another are unsubstituted or mono- or polysubstituted with one or more substituents selected from F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; CF₂Cl; CFCF₂; C₁₋₄-alkylene-CF₃; C₁₋₄-alkylene-CF₂H; C₁₋₄-alkylene-CFH₂; C(O)-C₁₋₆-alkyl; C(O)-OH; C(O)-OC₁₋₆-alkyl; C(O)-N(H)(OH); C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; OCF₃; OCF₂H; OCFH₂; OCF₂Cl; OCFCl₂; O-Ci-e-alkyl; O-C₃₋₆-cycloalkyl; O-(3 to 6-membered heterocycloalkyl); NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; N(H)-C(O)-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-C(O)-C₁₋₆-alkyl; N(H)-C(O)-NH₂; N(H)-C(O)-N(H)(C₁₋₆-alkyl); N(H)-C(O)-N(C₁₋₆-alkyl)₂; N(C₁₋₆-alkyl)-C(O)-N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)-C(O)-N(C₁₋₆-alkyl)₂; N(H)-S(O)₂-C₁₋₆-alkyl; SCF₃; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; S(O)₂-NH₂; S(O)₂-N(H)(C₁₋₆-alkyl); S(O)₂-N(C₁₋₆-alkyl)₂; C₃₋₆-cycloalkyl; C₁₋₄-alkylene-C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; C₁₋₄-alkylene-(3 to 6-membered heterocycloalkyl); phenyl or 5 or 6-membered heteroaryl;

[0010] in the form of the free compound or a physiologically acceptable salt thereof.

[0011] In a preferred embodiment, the compound according to the present invention is present in form of the free compound. For the purpose of specification, "free compound" preferably means that the compound according to the present invention is not present in form of a salt. Methods to determine whether a chemical substance is present as the free compound or as a salt are known to the skilled artisan such as ¹⁴N or ¹⁵N solid state NMR, x-ray diffraction, x-ray powder diffraction, IR, Raman, XPS. ¹H-NMR recorded in solution may also be used to consider the presence of protonation.

[0012] In another preferred embodiment, the compound according to the present invention is present in form of a physiologically acceptable salt. For the purposes of this specification, the term "physiologically acceptable salt" preferably refers to a salt obtained from a compound according to the present invention and a physiologically acceptable acid or base.

[0013] According to the present invention, the compound according to the present invention may be present in any possible form including solvates, cocrystals and polymorphs. For the purposes of this specification, the term "solvate" preferably refers to an adduct of (i) a compound according to the present invention and/or a physiologically acceptable salt thereof with (ii) distinct molecular equivalents of one or more solvents.

[0014] Further, the compound according to the present invention may be present in form of the racemate, enantiomers, diastereomers, tautomers or any mixtures thereof.

[0015] The present invention also includes isotopic isomers of a compound of the invention, wherein at least one atom of the compound is replaced by an isotope of the respective atom which is different from the naturally predominantly occurring isotope, as well as any mixtures of isotopic isomers of such a compound. Preferred isotopes are ²H (deuterium), ³H (tritium), ¹³C and ¹⁴C. Isotopic isomers of a compound of the invention can generally be prepared by conventional procedures known to a person skilled in the art.

[0016] According to the present invention, the terms "C₁₋₁₀-alkyl", "C₁₋₈-alkyl", "C₁₋₆-alkyl" and "C₁₋₄-alkyl" preferably mean acyclic saturated or unsaturated aliphatic (i.e. non-aromatic) hydrocarbon residues, which can be linear (i.e. unbranched) or branched and which can be unsubstituted or mono- or polysubstituted (e.g. di- or trisubstituted), and which contain 1 to 10 (i.e. 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10), 1 to 8 (i.e. 1, 2, 3, 4, 5, 6, 7 or 8), 1 to 6 (i.e. 1, 2, 3, 4, 5 or 6) and 1 to 4 (i.e. 1, 2, 3 or 4) carbon atoms, respectively. In a preferred embodiment, C₁₋₁₀-alkyl, C₁₋₈-alkyl, C₁₋₆-alkyl and C₁₋₄-alkyl are saturated. In another preferred embodiment, C₁₋₁₀-alkyl, C₁₋₈-alkyl, C₁₋₆-alkyl and C₁₋₄-alkyl are not saturated. According to this embodiment, C₁₋₁₀-alkyl, C₁₋₈-alkyl, C₁₋₆-alkyl and C₁₋₄-alkyl comprise at least one C-C double bond (a C=C-bond) or at least one C-C triple bond (a C≡C-bond). In still another preferred embodiment, C₁₋₁₀-alkyl,

C₁₋₈-alkyl, C₁₋₆-alkyl and C₁₋₄-alkyl are (i) saturated or (ii) not saturated, wherein C₁₋₁₀-alkyl, C₁₋₈-alkyl, C₁₋₆-alkyl and C₁₋₄-alkyl comprise at least one, preferably one, C-C triple bond (a C≡C-bond). Preferred C₁₋₁₀-alkyl groups are selected from methyl, ethyl, ethenyl (vinyl), n-propyl, 2-propyl, 1-propynyl, 2-propynyl, propenyl (-CH₂CH=CH₂, -CH=CH-CH₃, -C(=CH₂)-CH₃), n-butyl, 1-butynyl, 2-butynyl, 1-butenyl, 2-butenyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, 2-pentyl, 3-pentyl, 1-pentenyl, 2-pentenyl, 1-pentynyl, 2-pentynyl, 2-methylbutyl, 3-methylbutyl, 3-methylbut-2-yl, 2-methylbut-2-yl, 3-methylbut-1-ynyl, 2,2-dimethylpropyl, n-hexyl, 2-hexyl, 3-hexyl, 2-methylpentyl, 4-methylpentyl, 4-methylpent-2-yl, 2-methylpent-2-yl, 3,3-dimethylbutyl, 3,3-dimethylbut-2-yl, 3-methylpentyl, 3-methylpent-2-yl and 3-methylpent-3-yl; more preferably methyl, ethyl, n-propyl, 2-propyl, 1-propynyl, 2-propynyl, propenyl (-CH₂CH=CH₂, -CH=CH-CH₃, -C(=CH₂)-CH₃), n-butyl, 1-butynyl, 2-butynyl, 1-butenyl, 2-butenyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, 2-pentyl, 3-pentyl, 1-pentenyl, 2-pentenyl, 1-pentynyl, 2-pentynyl, 2-methylbutyl, 3-methylbutyl, 3-methylbut-2-yl, 2-methylbut-2-yl, 3-methylbut-1-ynyl, 2,2-dimethylpropyl, n-hexyl, n-heptyl, n-octyl, n-nonyl and n-decyl. Preferred C₁₋₈-alkyl groups are selected from methyl, ethyl, ethenyl (vinyl), n-propyl, 2-propyl, 1-propynyl, 2-propynyl, propenyl (-CH₂CH=CH₂, -CH=CH-CH₃, -C(=CH₂)-CH₃), n-butyl, 1-butynyl, 2-butynyl, 1-butenyl, 2-butenyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, 2-pentyl, 3-pentyl, 1-pentenyl, 2-pentenyl, 1-pentynyl, 2-pentynyl, 2-methylbutyl, 3-methylbutyl, 3-methylbut-2-yl, 2-methylbut-2-yl, 3-methylbut-1-ynyl, 2,2-dimethylpropyl, n-hexyl, 2-hexyl, 3-hexyl, 2-methylpentyl, 4-methylpentyl, 4-methylpent-2-yl, 2-methylpent-2-yl, 3,3-dimethylbutyl, 3,3-dimethylbut-2-yl, 3-methylpentyl, 3-methylpent-2-yl and 3-methylpent-3-yl; more preferably methyl, ethyl, n-propyl, 2-propyl, 1-propynyl, 2-propynyl, propenyl (-CH₂CH=CH₂, -CH=CH-CH₃, -C(=CH₂)-CH₃), n-butyl, 1-butynyl, 2-butynyl, 1-butenyl, 2-butenyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, 2-pentyl, 3-pentyl, 1-pentenyl, 2-pentenyl, 1-pentynyl, 2-pentynyl, 2-methylbutyl, 3-methylbutyl, 3-methylbut-2-yl, 2-methylbut-2-yl, 3-methylbut-1-ynyl, 2,2-dimethylpropyl, n-hexyl, n-heptyl and n-octyl. Preferred C₁₋₆-alkyl groups are selected from methyl, ethyl, ethenyl (vinyl), n-propyl, 2-propyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, 2-pentyl, 3-pentyl, 2-methylbutyl, 3-methylbutyl, 3-methylbut-2-yl, 2-methylbut-2-yl, 2,2-dimethylpropyl, n-hexyl, 2-hexyl, 3-hexyl, 2-methylpentyl, 4-methylpentyl, 4-methylpent-2-yl, 2-methylpent-2-yl, 3,3-dimethylbutyl, 3,3-dimethylbut-2-yl, 3-methylpentyl, 3-methylpent-2-yl and 3-methylpent-3-yl; more preferably methyl, ethyl, n-propyl, 2-propyl, 1-propynyl, 2-propynyl, propenyl (-CH₂CH=CH₂, -CH=CH-CH₃, -C(=CH₂)-CH₃), n-butyl, 1-butynyl, 2-butynyl, 1-butenyl, 2-butenyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, 2-pentyl, 3-pentyl, 1-pentenyl, 2-pentenyl, 1-pentynyl, 2-pentynyl, 2-methylbutyl, 3-methylbutyl, 3-methylbut-2-yl, 2-methylbut-2-yl, 3-methylbut-1-ynyl, 2,2-dimethylpropyl, n-hexyl. Particularly preferred C₁₋₆-alkyl groups are selected from C₁₋₄-alkyl groups. Preferred C₁₋₄-alkyl groups are selected from methyl, ethyl, ethenyl (vinyl), n-propyl, 2-propyl, 1-propynyl, 2-propynyl, propenyl (-CH₂CH=CH₂, -CH=CH-CH₃, -C(=CH₂)-CH₃), n-butyl, 1-butynyl, 2-butynyl, 1-butenyl, 2-butenyl, isobutyl, sec-butyl, tert-butyl and 3-methylbut-1-ynyl.

[0017] Further according to the present invention, the terms "C₁₋₆-alkylene", "C₁₋₄-alkylene" and "C₁₋₂-alkylene" relate to a linear or branched, preferably linear, and preferably saturated aliphatic residues which are preferably selected from the group consisting of methylene (-CH₂-), ethylene (-CH₂CH₂-), propylene (-CH₂CH₂CH₂- or -C(CH₃)₂-), butylene (-CH₂CH₂CH₂CH₂-), pentylene (-CH₂CH₂CH₂CH₂CH₂-) and hexylene (-CH₂CH₂CH₂CH₂CH₂CH₂-); more preferably methylene (-CH₂-) and ethylene (-CH₂CH₂-) and most preferably methylene (-CH₂-). Preferably, C₁₋₆-alkylene is selected from C₁₋₄-alkylene, more preferably from C₁₋₂-alkylene.

[0018] Still further according to the present invention, the terms "C₃₋₁₀-cycloalkyl" and "C₃₋₆-cycloalkyl" preferably mean cyclic aliphatic hydrocarbons containing 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms and 3, 4, 5 or 6 carbon atoms, respectively, wherein the hydrocarbons in each case can be saturated or unsaturated (but not aromatic), unsubstituted or mono- or polysubstituted. Preferably, C₃₋₁₀-cycloalkyl and C₃₋₆-cycloalkyl are saturated. The C₃₋₁₀-cycloalkyl and C₃₋₆-cycloalkyl can be bound to the respective superordinate general structure via any desired and possible ring member of the cycloalkyl group. The C₃₋₁₀-cycloalkyl and C₃₋₆-cycloalkyl groups can also be condensed with further saturated, (partially) unsaturated, (hetero)cyclic, aromatic or heteroaromatic ring systems, i.e. with cycloalkyl, heterocyclyl, aryl or heteroaryl residues, which in each case can in turn be unsubstituted or mono- or polysubstituted. Further, C₃₋₁₀-cycloalkyl and C₃₋₆-cycloalkyl can be singly or multiply bridged such as, for example, in the case of adamantyl, bicyclo[2.2.1]heptyl or bicyclo[2.2.2]octyl. However, preferably, C₃₋₁₀-cycloalkyl and C₃₋₆-cycloalkyl are neither condensed with further ring systems nor bridged. More preferably, C₃₋₁₀-cycloalkyl and C₃₋₆-cycloalkyl are neither condensed with further ring systems nor bridged and are saturated. Preferred C₃₋₁₀-cycloalkyl groups are selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopentenyl, cyclohexenyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, adamantyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, bicyclo[2.2.1]heptyl and bicyclo[2.2.2]octyl. Particularly preferred C₃₋₁₀-cycloalkyl groups are selected from C₃₋₆-cycloalkyl groups. Preferred C₃₋₆-cycloalkyl groups are selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopentenyl and cyclohexenyl. Particularly preferred C₃₋₆-cycloalkyl groups are selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl, most preferably cyclopropyl.

[0019] According to the present invention, the terms "3 to 7-membered heterocycloalkyl" and "3 to 6-membered heterocycloalkyl" preferably mean heterocycloaliphatic saturated or unsaturated (but not aromatic) residues having 3 to 7, i.e. 3, 4, 5, 6 or 7 ring members and 3 to 6, i.e. 3, 4, 5 or 6 ring members, respectively, wherein in each case at least one, if appropriate also two or three carbon atoms are replaced by a heteroatom or a heteroatom group each selected

independently of one another from the group consisting of O, S, S(=O), S(=O)₂, N, NH and N(C₁₋₄-alkyl) such as N(CH₃), wherein the carbon atoms of the ring can be unsubstituted or mono- or polysubstituted. Preferably, 3 to 7-membered heterocycloalkyl and 3 to 6-membered heterocycloalkyl are saturated. The 3 to 7-membered heterocycloalkyl and the 3 to 6-membered heterocycloalkyl groups can also be condensed with further saturated or (partially) unsaturated cycloalkyl or heterocyclyl, aromatic or heteroaromatic ring systems. However, more preferably, 3 to 7-membered heterocycloalkyl and 3 to 6-membered heterocycloalkyl are not condensed with further ring systems. Still more preferably, 3 to 7-membered heterocycloalkyl and 3 to 6-membered heterocycloalkyl are not condensed with further ring systems and are saturated. The 3 to 7-membered heterocycloalkyl and the 3 to 6-membered heterocycloalkyl group can be bound to the superordinate general structure via any desired and possible ring member of the heterocycloaliphatic residue if not indicated otherwise. In a preferred embodiment, 3 to 7-membered heterocycloalkyl and 3 to 6-membered heterocycloalkyl are bound to the superordinate general structure via a carbon atom.

[0020] Preferred 3 to 7-membered heterocycloalkyl groups are selected from the group consisting of azepanyl, dioxepanyl, oxazepanyl, diazepanyl, thiazolidinyl, tetrahydrothiophenyl, tetrahydropyridinyl, thiomorpholinyl, tetrahydropyranyl, oxetanyl, oxiranyl, tetrahydrofuranlyl, morpholinyl, pyrrolidinyl, 4-methylpiperazinyl, morpholinonyl, azetidiny, aziridinyl, dithiolanyl, dihydropyrrolyl, dioxanyl, dioxolanyl, dihydropyridinyl, dihydrofuranlyl, dihydroisoxazolyl, dihydrooxazolyl, imidazolidinyl, isoxazolidinyl, oxazolidinyl, piperazinyl, piperidinyl, pyrazolidinyl, pyranlyl; tetrahydropyrrolyl, dihydroquinolinyl, dihydroisoquinolinyl, dihydroindolinyl, dihydroisindolyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl and tetrahydroindolinyl. Particularly preferred 3 to 7-membered heterocycloalkyl groups are selected from 3 to 6-membered heterocycloalkyl groups. Preferred 3 to 6-membered heterocycloalkyl groups are selected from the group consisting of tetrahydropyranyl, oxetanyl, oxiranyl, tetrahydrofuranlyl, thiazolidinyl, tetrahydrothiophenyl, tetrahydropyridinyl, thiomorpholinyl, morpholinyl, pyrrolidinyl, 4-methylpiperazinyl, morpholinonyl, azetidiny, aziridinyl, dithiolanyl, dihydropyrrolyl, dioxanyl, dioxolanyl, dihydropyridinyl, dihydrofuranlyl, dihydroisoxazolyl, dihydrooxazolyl, imidazolidinyl, isoxazolidinyl, oxazolidinyl, piperazinyl, piperidinyl, pyrazolidinyl, pyranlyl, tetrahydropyrrolyl, dihydroindolinyl, dihydroisindolyl and tetrahydroindolinyl. Particularly preferred 3 to 6-membered heterocycloalkyl groups are selected from the group consisting of tetrahydropyranyl, oxetanyl, oxiranyl, and tetrahydrofuranlyl.

[0021] According to the present invention, the term "aryl" preferably means aromatic hydrocarbons having 6 to 14, i.e. 6, 7, 8, 9, 10, 11, 12, 13 or 14 ring members, preferably having 6 to 10, i.e. 6, 7, 8, 9 or 10 ring members, including phenyls and naphthyls. Each aryl residue can be unsubstituted or mono- or polysubstituted. The aryl can be bound to the superordinate general structure via any desired and possible ring member of the aryl residue. The aryl residues can also be condensed with further saturated or (partially) unsaturated cycloalkyl or heterocycloalkyl, aromatic or heteroaromatic ring systems, which can in turn be unsubstituted or mono- or polysubstituted. In a preferred embodiment, aryl is condensed with a further ring system. Examples of condensed aryl residues are 2H-benzo[b][1,4]oxazin-3(4H)-onyl, 1H-benzo[d]imidazolyl, 2,3-dihydro-1H-indenyl, tetrahydronaphthalenyl, isochroman, 1,3-dihydroisobenzofuranlyl, benzodioxolanyl and benzodioxanyl. Preferably, aryl is selected from the group consisting of phenyl, 1H-benzo[d]imidazolyl, 2H-benzo[b][1,4]oxazin-3(4H)-onyl, 2,3-dihydro-1H-indenyl, tetrahydronaphthalenyl, isochroman, 1,3-dihydroisobenzofuranlyl, 1-naphthyl, 2-naphthyl, fluorenyl and anthracenyl, each of which can be respectively unsubstituted or mono- or polysubstituted. In another preferred embodiment, aryl is not condensed with any further ring system. A particularly preferred aryl is phenyl, unsubstituted or mono- or polysubstituted.

[0022] According to the present invention, the term "5- to 6-membered heteroaryl" preferably means a 5 or 6-membered cyclic aromatic residue containing at least 1, if appropriate also 2, 3, 4 or 5 heteroatoms, wherein the heteroatoms are each selected independently of one another from the group S, N and O and the heteroaryl residue can be unsubstituted or mono- or polysubstituted, if not indicated otherwise. In the case of substitution on the heteroaryl, the substituents can be the same or different and be in any desired and possible position of the heteroaryl. The binding to the superordinate general structure can be carried out via any desired and possible ring member of the heteroaryl residue if not indicated otherwise. Preferably, the 5- to 6-membered heteroaryl is bound to the superordinate general structure via a carbon atom of the heterocycle. The heteroaryl can also be part of a bi- or polycyclic system having up to 14 ring members, wherein the ring system can be formed with further saturated or (partially) unsaturated cycloalkyl or heterocycloalkyl, aromatic or heteroaromatic ring systems, which can in turn be unsubstituted or mono- or polysubstituted, if not indicated otherwise. In a preferred embodiment, the 5- to 6-membered heteroaryl is part of a bi- or polycyclic, preferably bicyclic, system. In another preferred embodiment, the 5- to 6-membered heteroaryl is not part of a bi- or polycyclic system. Preferably, the 5- to 6-membered heteroaryl is selected from the group consisting of pyridyl (i.e. 2-pyridyl, 3-pyridyl, 4-pyridyl), pyrimidinyl, pyridazinyl, pyrazinyl, pyrrolyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, furanyl, thienyl (thiophenyl), triazolyl, thiadiazolyl, 4,5,6,7-tetrahydro-2H-indazolyl, 2,4,5,6-tetrahydrocyclopenta[c]pyrazolyl, benzofuranlyl, benzoimidazolyl, benzothienyl, benzothiadiazolyl, benzothiazolyl, benzotriazolyl, benzooxazolyl, benzooxadiazolyl, quinoxalinyl, quinoxalinyl, carbazolyl, quinolinyl, dibenzofuranlyl, dibenzothienyl, imidazothiazolyl, indazolyl, indoliziny, indolyl, isoquinolinyl, naphthyridinyl, oxazolyl, oxadiazolyl, phenazinyl, phenothiazinyl, phthalazinyl, purinyl, phenazinyl, tetrazolyl and triazinyl. Particularly preferred 5- to 6-membered heteroaryl are selected from the group consisting of pyridyl (i.e. 2-pyridyl, 3-pyridyl, 4-pyridyl).

[0023] The compounds according to the present invention are defined by substituents, for example by R¹, R² and R³ (1st generation substituents) which may optionally be for their part themselves be substituted (2nd generation substituents). Depending on the definition, these substituents of the substituents can optionally be for their part resubstituted (3rd generation substituents). If, for example, R¹ = a C₁₋₁₀-alkyl (1st generation substituent), then the C₁₋₁₀-alkyl can for its part be substituted, for example with a N(H)(C₁₋₆-alkyl) (2nd generation substituent). This produces the functional group R¹ = (C₁₋₁₀-alkyl-NH-C₁₋₆-alkyl). The NH-C₁₋₆-alkyl can then for its part be resubstituted, for example with Cl (3rd generation substituent). Overall, this produces the functional group R¹ = C₁₋₁₀-alkyl-NH-C₁₋₆-alkyl, wherein the C₁₋₆-alkyl of the NH-C₁₋₆-alkyl is substituted by Cl. However, in a preferred embodiment, the 3rd generation substituents may not be resubstituted, i.e. there are then no 4th generation substituents. More preferably, the 2nd generation substituents may not be resubstituted, i.e. there are no 3rd generation substituents.

[0024] If a residue occurs multiply within a molecule, then this residue can have respectively different meanings for various substituents: if, for example, both R² and R³ denote C₁₋₆-alkyl, then C₁₋₆-alkyl can e.g. represent ethyl for R² and can represent methyl for R³.

[0025] In connection with the terms "C₁₋₁₀-alkyl", "C₁₋₆-alkyl", "C₁₋₄-alkyl", "C₃₋₁₀-cycloalkyl", "C₃₋₆-cycloalkyl", "3 to 7 membered heterocycloalkyl", "3 to 6-membered heterocycloalkyl", "C₁₋₆-alkylene", "C₁₋₄-alkylene" and "C₁₋₂-alkylene", the term "substituted" refers in the sense of the present invention, with respect to the corresponding residues or groups, to the single substitution (monosubstitution) or multiple substitution (polysubstitution), e.g. disubstitution or trisubstitution; more preferably to monosubstitution or disubstitution; of one or more hydrogen atoms each independently of one another by at least one substituent. In case of a multiple substitution, i.e. in case of polysubstituted residues, such as di- or trisubstituted residues, these residues may be polysubstituted either on different or on the same atoms, for example trisubstituted on the same carbon atom, as in the case of CF₃, CH₂CF₃ or disubstituted as in the case of 1,1-difluorocyclohexyl, or at various points, as in the case of CH(OH)-CH=CH-CHCl₂ or 1-chloro-3-fluorocyclohexyl. The multiple substitution can be carried out using the same or using different substituents.

[0026] In relation to the terms "aryl", "phenyl", "heteroaryl" and "5- to 6-membered heteroaryl", the term "substituted" refers in the sense of this invention to the single substitution (monosubstitution) or multiple substitution (polysubstitution), e.g. disubstitution or trisubstitution, of one or more hydrogen atoms each independently of one another by at least one substituent. The multiple substitution can be carried out using the same or using different substituents.

[0027] According to the present invention, preferably C₁₋₁₀-alkyl, C₁₋₆-alkyl, C₁₋₄-alkyl, C₃₋₁₀-cycloalkyl, C₃₋₆-cycloalkyl, 3 to 7 membered heterocycloalkyl, 3 to 6-membered heterocycloalkyl, C₁₋₆-alkylene, C₁₋₄-alkylene and C₁₋₂-alkylene in each case independently from one another are unsubstituted or mono- or polysubstituted with one or more substituents selected from F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; CF₂Cl; CFC₂Cl; C(O)-C₁₋₆-alkyl; C(O)-OH; C(O)-OC₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; =O; OCF₃; OCF₂H; OCFH₂; OCF₂Cl; OCFCl₂; O-C₁₋₆-alkyl; O-C(O)-C₁₋₆-alkyl; O-C(O)-O-C₁₋₆-alkyl; O-(CO)-N(H)(C₁₋₆-alkyl); O-C(O)-N(C₁₋₆-alkyl)₂; O-S(O)₂-NH₂; O-S(O)₂-N(H)(C₁₋₆-alkyl); O-S(O)₂-N(C₁₋₆-alkyl)₂; NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; N(H)-C(O)-C₁₋₆-alkyl; N(H)-C(O)-O-C₁₋₆-alkyl; N(H)-C(O)-NH₂; N(H)-C(O)-N(H)(C₁₋₆-alkyl); N(H)-C(O)-N(C₁₋₆-alkyl)₂; N(C₁₋₆-alkyl)-C(O)-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-C(O)-O-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-C(O)-NH₂; N(C₁₋₆-alkyl)-C(O)-N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)-C(O)-N(C₁₋₆-alkyl)₂; N(H)-S(O)₂OH; N(H)-S(O)₂-C₁₋₆-alkyl; N(H)-S(O)₂-O-C₁₋₆-alkyl; N(H)-S(O)₂-NH₂; N(H)-S(O)₂-N(H)(C₁₋₆-alkyl); N(H)-S(O)₂-N(C₁₋₆-alkyl)₂; N(C₁₋₆-alkyl)-S(O)₂-OH; N(C₁₋₆-alkyl)-S(O)₂-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-S(O)₂-O-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-S(O)₂-NH₂; N(C₁₋₆-alkyl)-S(O)₂-N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)-S(O)₂-N(C₁₋₆-alkyl)₂; SCF₃; SCF₂H; SCFH₂; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; S(O)₂-OH; S(O)₂-O-C₁₋₆-alkyl; S(O)₂-NH₂; S(O)₂-N(H)(C₁₋₆-alkyl); S(O)₂-N(C₁₋₆-alkyl)₂; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; phenyl; 5 or 6-membered heteroaryl; O-C₃₋₆-cycloalkyl; O-(3 to 6-membered heterocycloalkyl); O-phenyl; O-(5 or 6-membered heteroaryl); C(O)-C₃₋₆-cycloalkyl; C(O)-(3 to 6-membered heterocycloalkyl); C(O)-phenyl; C(O)-(5 or 6-membered heteroaryl); S(O)₂-C₃₋₆-cycloalkyl; S(O)₂-(3 to 6-membered heterocycloalkyl); S(O)₂-phenyl and S(O)₂-(5 or 6-membered heteroaryl).

[0028] Preferred substituents of C₁₋₁₀-alkyl, C₁₋₆-alkyl, C₁₋₄-alkyl, C₃₋₁₀-cycloalkyl, C₃₋₆-cycloalkyl, 3 to 7 membered heterocycloalkyl, 3 to 6-membered heterocycloalkyl, C₁₋₆-alkylene and C₁₋₄-alkylene are selected from the group consisting of F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; OCF₃; OCF₂H; OCFH₂; O-C₁₋₆-alkyl; NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; SCF₃; SCF₂H; SCFH₂; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; phenyl and 5 or 6-membered heteroaryl; and particularly preferably F, CN, CH₃, CH₂CH₃, CF₃; CF₂H; CFH₂; C(O)-NH₂; C(O)-N(H)(CH₃); C(O)-N(CH₃)₂; OH, NH₂, OCH₃, SCH₃, S(O)₂(CH₃), S(O)(CH₃), N(CH₃)₂, cyclopropyl and oxetanyl. According to this embodiment, C₁₋₁₀-alkyl, C₁₋₆-alkyl, C₁₋₄-alkyl, C₃₋₁₀-cycloalkyl, C₃₋₆-cycloalkyl, 3 to 7 membered heterocycloalkyl, 3 to 6-membered heterocycloalkyl are preferably each independently from one another unsubstituted, mono- di- or trisubstituted, more preferably unsubstituted or monosubstituted or disubstituted with a substituent selected from the group consisting of F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; OCF₃; OCF₂H; OCFH₂; O-C₁₋₆-alkyl; NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; SCF₃; SCF₂H; SCFH₂; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; phenyl and 5 or 6-membered heteroaryl. Preferably, C₁₋₆-alkylene groups and

C₁₋₄-alkylene groups are unsubstituted.

[0029] According to the present invention, preferably aryl, phenyl and 5 or 6-membered heteroaryl in each case independently from one another are unsubstituted or mono- or polysubstituted with one or more substituents selected from F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; CF₂Cl; CFC₂Cl; C₁₋₄-alkylene-CF₃; C₁₋₄-alkylene-CF₂H; C₁₋₄-alkylene-CFH₂; C(O)-C₁₋₆-alkyl; C(O)-OH; C(O)-OC₁₋₆-alkyl; C(O)-N(H)(OH); C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; OCF₃; OCF₂H; OCFH₂; OCF₂Cl; OCFCl₂; O-C₁₋₆-alkyl; O-C₃₋₆-cycloalkyl; O-(3 to 6-membered heterocycloalkyl); NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; N(H)-C(O)-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-C(O)-C₁₋₆-alkyl; N(H)-C(O)-NH₂; N(H)-C(O)-N(H)(C₁₋₆-alkyl); N(H)-C(O)-N(C₁₋₆-alkyl)₂; N(C₁₋₆-alkyl)-C(O)-N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)-C(O)-N(C₁₋₆-alkyl)₂; N(H)-S(O)₂-C₁₋₆-alkyl; SCF₃; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; S(O)₂-NH₂; S(O)₂-N(H)(C₁₋₆-alkyl); S(O)₂-N(C₁₋₆-alkyl)₂; C₃₋₆-cycloalkyl; C₁₋₄-alkylene-C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; C₁₋₄-alkylene-(3 to 6-membered heterocycloalkyl); phenyl or 5 or 6-membered heteroaryl.

[0030] Preferred substituents of aryl, phenyl and 5 or 6-membered heteroaryl are selected from the group consisting of F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; C₁₋₄-alkylene-CF₃; C₁₋₄-alkylene-CF₂H; C₁₋₄-alkylene-CFH₂; OH; OCF₃; OCF₂H; OCFH₂; O-C₁₋₆-alkyl; O-C₃₋₆-cycloalkyl and C₃₋₆-cycloalkyl; and particularly preferably of F; Cl; Br; CN; CH₃; CH₂CH₃; CF₃; CF₂H; CFH₂; CH₂-CF₃; OH; OCF₃; OCF₂H; OCFH₂; O-CH₃; O-cyclopropyl and cyclopropyl. According to this embodiment, aryl, phenyl and 5 or 6-membered heteroaryl are preferably each independently from one another unsubstituted, mono- di- or trisubstituted, more preferably unsubstituted or monosubstituted or disubstituted with a substituent selected from the group consisting of F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; C₁₋₄-alkylene-CF₃; C₁₋₄-alkylene-CF₂H; C₁₋₄-alkylene-CFH₂; OH; OCF₃; OCF₂H; OCFH₂; O-Ci-e-alkyl; O-C₃₋₆-cycloalkyl and C₃₋₆-cycloalkyl.

[0031] In a preferred embodiment, X represents NR⁷ and Z represents N or CR⁹. More preferably, X represents NR⁷ and Z represents CR⁹. In another preferred embodiment, X represents N and Z represents NR⁷.

[0032] More preferably, the compound according to the present invention is according to general formula (II) or (III), wherein X represents NR⁷ and Z represents N or CR⁹, more preferably CR⁹.

[0033] In a preferred embodiment, R¹ represents H; C₁₋₆-alkyl; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; phenyl; or 5 or 6-membered heteroaryl; wherein C₃₋₆-cycloalkyl, 3 to 6-membered heterocycloalkyl, phenyl and 5 or 6-membered heteroaryl can optionally be bridged via C₁₋₄-alkylene. According to this embodiment, preferably C₁₋₆-alkyl; C₃₋₆-cycloalkyl; C₁₋₄-alkylene and 3 to 6-membered heterocycloalkyl in each case independently from one another are unsubstituted or mono-, di- or trisubstituted with one or more substituents selected from F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; CF₂Cl; CFC₂Cl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; ; OCF₃; OCF₂H; OCFH₂; ; O-C₁₋₆-alkyl; O-C(O)-C₁₋₆-alkyl; NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; N(H)-C(O)-C₁₋₆-alkyl; N(H)-C(O)-O-C₁₋₆-alkyl; SCF₃; SCF₂H; SCFH₂; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; ;S(O)₂-O-C₁₋₆-alkyl; S(O)₂-NH₂; S(O)₂-N(H)(C₁₋₆-alkyl); S(O)₂-N(C₁₋₆-alkyl)₂; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; phenyl; 5 or 6-membered heteroaryl; O-C₃₋₆-cycloalkyl; O-(3 to 6-membered heterocycloalkyl); O-phenyl; O-(5 or 6-membered heteroaryl); C(O)-C₃₋₆-cycloalkyl; C(O)-(3 to 6-membered heterocycloalkyl); C(O)-phenyl; C(O)-(5 or 6-membered heteroaryl); S(O)₂-(C₃₋₆-cycloalkyl); S(O)₂-(3 to 6-membered heterocycloalkyl); S(O)₂-phenyl or S(O)₂-(5 or 6-membered heteroaryl); more preferably F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; OCF₃; OCF₂H; OCFH₂; O-C₁₋₆-alkyl; NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; SCF₃; SCF₂H; SCFH₂; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; phenyl and 5 or 6-membered heteroaryl; and particularly preferably F, CN, CH₃, CH₂CH₃, CF₃; CF₂H; CFH₂; C(O)-NH₂; C(O)-N(H)(CH₃); C(O)-N(CH₃)₂; OH, NH₂, OCH₃, SCH₃, S(O)₂(CH₃), S(O)(CH₃), N(CH₃)₂, cyclopropyl and oxetanyl;

and preferably phenyl and 5 or 6-membered heteroaryl in each case independently from one another are unsubstituted or mono-, di- or trisubstituted with one or more substituents selected from F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; C₁₋₄-alkylene-CF₃; C₁₋₄-alkylene-CF₂H; C₁₋₄-alkylene-CFH₂; C(O)-C₁₋₆-alkyl; C(O)-OC₁₋₆-alkyl; C(O)-; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; OCF₃; OCF₂H; OCFH₂; O-C₁₋₆-alkyl; O-C₃₋₆-cycloalkyl; O-(3 to 6-membered heterocycloalkyl); NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; SCF₃; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; C₃₋₆-cycloalkyl; C₁₋₄-alkylene-C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; C₁₋₄-alkylene-(3 to 6-membered heterocycloalkyl); phenyl or 5 or 6-membered heteroaryl; more preferably F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; C₁₋₄-alkylene-CF₃; C₁₋₄-alkylene-CF₂H; C₁₋₄-alkylene-CFH₂; OH; OCF₃; OCF₂H; OCFH₂; O-C₁₋₆-alkyl; O-C₃₋₆-cycloalkyl and C₃₋₆-cycloalkyl; and particularly preferably F; Cl; Br; CN; CH₃; CH₂CH₃; CF₃; CF₂H; CFH₂; CH₂-CF₃; OH; OCF₃; OCF₂H; OCFH₂; O-CH₃; O-cyclopropyl and cyclopropyl.

[0034] In another preferred embodiment, R¹ represents H; C₁₋₄-alkyl; C₃₋₆-cycloalkyl, optionally bridged via C₁₋₂-alkylene; 3 to 6-membered heterocycloalkyl, optionally bridged via C₁₋₂-alkylene; phenyl, optionally bridged via C₁₋₂-alkylene; or 5 or 6-membered heteroaryl, optionally bridged via C₁₋₂-alkylene; preferably wherein

C₁₋₄-alkyl, C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl in each case independently from one another is unsubstituted, mono-, di- or trisubstituted with one or more substituents selected from the group consisting of F,

CN, CH₃, CH₂CH₃, CF₃, CF₂H; CFH₂; C(O)-NH₂; C(O)-N(H)(CH₃); C(O)-N(CH₃)₂; OH, NH₂, OCH₃, SCH₃, S(O)₂(CH₃), S(O)(CH₃), N(CH₃)₂, cyclopropyl and oxetanyl; and
 C₁₋₂-alkylene is unsubstituted; and
 phenyl and 5 or 6-membered heteroaryl in each case independently from one another is unsubstituted, mono-, di-
 or trisubstituted with one or more substituents selected from the group consisting of F; Cl; Br; CN; CH₃; CH₂CH₃;
 CF₃; CF₂H; CFH₂; CH₂-CF₃; OH; OCF₃; OCF₂H; OCFH₂; O-CH₃; O-cyclopropyl and cyclopropyl.

[0035] In still another preferred embodiment, R¹ represents H; CH₃, CF₃, CF₂H; CFH₂; ethyl, n-propyl, 2-propyl, 1-propynyl, 2-propynyl, propenyl (-CH₂CH=CH₂, -CH=CH-CH₃, -C(=CH₂)-CH₃), n-butyl, 1-butylnyl, 2-butylnyl, 1-butenyl, 2-butenyl, isobutyl, sec-butyl; tert-butyl; (CH₂)₁₋₂OCH₃; (CH₂)₁₋₂OH; (CH₂)₀₋₂C(H)(OH)-(CH₂)₀₋₂CH₃; (CH₂)₁₋₂SCH₃; (CH₂)₁₋₂N(CH₃)₂; (CH₂)₁₋₂S(O)CH₃; (CH₂)₁₋₂S(O)₂CH₃; (CH₂)₁₋₂CN; (CH₂)₀₋₂C(H)(CN)-(CH₂)₀₋₂CH₃; (CH₂)₀₋₂-cyclopropyl, (CH₂)₀₋₂-cyclobutyl, (CH₂)₀₋₂-cyclopentyl and (CH₂)₀₋₂-cyclohexyl; (CH₂)₀₋₂-tetrahydropyranyl, (CH₂)₀₋₂-oxetanyl, (CH₂)₀₋₂-oxiranyl, (CH₂)₀₋₂-tetrahydrofuranyl; (CH₂)₀₋₂-phenyl; (CH₂)₀₋₂-pyridyl, (CH₂)₀₋₂-pyrimidinyl, (CH₂)₀₋₂-pyridazinyl, (CH₂)₀₋₂-thienyl, (CH₂)₀₋₂-oxazolyl or (CH₂)₀₋₂-thiazolyl.

[0036] Preferably, R² represents H; F; Cl; Br; CN; C₁₋₆-alkyl; C₃₋₆-cycloalkyl; O-C₁₋₆-alkyl; N(H)(C₁₋₆-alkyl), N(C₁₋₆-alkyl)₂; C(O)-C₁₋₆-alkyl; C(O)-O-C₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; O-C₃₋₆-cycloalkyl; N(H)(C₃₋₆-cycloalkyl), N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); C(O)-C₃₋₆-cycloalkyl; C(O)-O-C₃₋₆-cycloalkyl; C(O)-N(H)(C₃₋₆-cycloalkyl) or C(O)-N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); wherein C₃₋₆-cycloalkyl can optionally be bridged via C₁₋₄-alkylene.

[0037] In a preferred embodiment, R² represents H; F; Cl; Br; CN; C₁₋₆-alkyl; C₃₋₆-cycloalkyl; O-C₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; C(O)-N(H)(C₃₋₆-cycloalkyl) or C(O)-N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); wherein C₃₋₆-cycloalkyl can optionally be bridged via C₁₋₄-alkylene; preferably wherein

C₁₋₆-alkyl and C₃₋₆-cycloalkyl in each case independently from one another is unsubstituted, mono-, di- or trisubstituted with one or more substituents selected from the group consisting of F, CN, CH₃, CH₂CH₃, CF₃; CF₂H; CFH₂; C(O)-NH₂; C(O)-N(H)(CH₃); C(O)-N(CH₃)₂; OH, NH₂, OCH₃, SCH₃, S(O)₂(CH₃), S(O)(CH₃), N(CH₃)₂, cyclopropyl and oxetanyl; and
 C₁₋₄-alkylene is unsubstituted.

[0038] In a particularly preferred embodiment, R² represents H; F; Cl; Br; CN; methyl; ethyl; ethenyl (vinyl); n-propyl; 2-propyl; 1-propynyl; 2-propynyl; propenyl (-CH₂CH=CH₂; -CH=CH-CH₃; -C(=CH₂)-CH₃); n-butyl; isobutyl; sec-butyl; tert-butyl; CF₃; CH₂CF₃; CHF₂; CH₂CHF₂; CH₂F; CH₂CH₂F; OCH₃; OCH₂CH₃; OC(H)(CH₃)₂; OCH₂CH₂CH₃; O-C(CH₃)₃; OCF₃; OCH₂CF₃; OCHF₂; OCH₂CHF₂; OCH₂F; OCH₂CH₂F; CH₂OH; CH₂CH₂OH; CH₂C(H)(OH)CH₃; CH₂CH₂CH₂OH; CH₂CH₂CH₂CH₂OH; C(CH₃)₂CH₂OH; C(O)-CH₃; C(O)-CH₂CH₃; C(O)-C(H)(CH₃)₂; C(O)-CH₂CH₂CH₃; C(O)-C(CH₃)₃; cyclopropyl; cyclobutyl; cyclopentyl; cyclohexyl; CH₂-cyclopropyl; CH₂-cyclobutyl; CH₂-cyclopentyl; CH₂-cyclohexyl; O-cyclopropyl; O-cyclobutyl; O-cyclopentyl; O-cyclohexyl; C(O)-cyclopropyl; C(O)-cyclobutyl; C(O)-cyclopentyl; C(O)-cyclohexyl or C(O)-NH₂.

[0039] Preferably, R³ represents H; F; Cl; Br; CN; C₁₋₆-alkyl; C₃₋₆-cycloalkyl; O-C₁₋₆-alkyl; N(H)(C₁₋₆-alkyl), N(C₁₋₆-alkyl)₂; C(O)-C₁₋₆-alkyl; C(O)-O-C₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; O-C₃₋₆-cycloalkyl; N(H)(C₃₋₆-cycloalkyl), N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); C(O)-C₃₋₆-cycloalkyl; C(O)-O-C₃₋₆-cycloalkyl; C(O)-N(H)(C₃₋₆-cycloalkyl) or C(O)-N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); wherein C₃₋₆-cycloalkyl can optionally be bridged via C₁₋₄-alkylene.

[0040] In a preferred embodiment, R³ represents H; F; Cl; Br; CN; C₁₋₆-alkyl; C₃₋₆-cycloalkyl; O-C₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; C(O)-N(H)(C₃₋₆-cycloalkyl) or C(O)-N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); wherein C₃₋₆-cycloalkyl can optionally be bridged via C₁₋₄-alkylene; preferably wherein

C₁₋₆-alkyl and C₃₋₆-cycloalkyl in each case independently from one another is unsubstituted, mono-, di- or trisubstituted with one or more substituents selected from the group consisting of F, CN, CH₃, CH₂CH₃, CF₃; CF₂H; CFH₂; C(O)-NH₂; C(O)-N(H)(CH₃); C(O)-N(CH₃)₂; OH, NH₂, OCH₃, SCH₃, S(O)₂(CH₃), S(O)(CH₃), N(CH₃)₂, cyclopropyl and oxetanyl; and
 C₁₋₄-alkylene is unsubstituted.

[0041] In a particularly preferred embodiment, R³ represents H; F; Cl; Br; CN; methyl; ethyl; ethenyl (vinyl); n-propyl; 2-propyl; 1-propynyl; 2-propynyl; propenyl (-CH₂CH=CH₂; -CH=CH-CH₃; -C(=CH₂)-CH₃); n-butyl; isobutyl; sec-butyl; tert-butyl; CF₃; CH₂CF₃; CHF₂; CH₂CHF₂; CH₂F; CH₂CH₂F; OCH₃; OCH₂CH₃; OC(H)(CH₃)₂; OCH₂CH₂CH₃; O-C(CH₃)₃; OCF₃; OCH₂CF₃; OCHF₂; OCH₂CHF₂; OCH₂F; OCH₂CH₂F; CH₂OH; CH₂CH₂OH; CH₂C(H)(OH)CH₃; CH₂CH₂CH₂OH;

CH₂CH₂CH₂CH₂OH; C(CH₃)₂CH₂OH; C(O)-CH₃; C(O)-CH₂CH₃; C(O)-C(H)(CH₃)₂ C(O)-CH₂CH₂CH₃; C(O)-C(CH₃)₃; cyclopropyl; cyclobutyl; cyclopentyl; cyclohexyl; CH₂-cyclopropyl; CH₂-cyclobutyl; CH₂-cyclopentyl; CH₂-cyclohexyl; O-cyclopropyl; O-cyclobutyl; O-cyclopentyl; O-cyclohexyl; C(O)-cyclopropyl; C(O)-cyclobutyl; C(O)-cyclopentyl; C(O)-cyclohexyl; C(O)-NH₂.

5 **[0042]** In a preferred embodiment,

R² represents H; F; Cl; Br; CN; C₁₋₆-alkyl; C₃₋₆-cycloalkyl; O-C₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; C(O)-N(H)(C₃₋₆-cycloalkyl) or C(O)-N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); wherein C₃₋₆-cycloalkyl can optionally be bridged via C₁₋₄-alkylene; and/or

10 R³ represents H; F; Cl; Br; CN; C₁₋₆-alkyl; C₃₋₆-cycloalkyl; O-C₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; C(O)-N(H)(C₃₋₆-cycloalkyl) or C(O)-N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); wherein C₃₋₆-cycloalkyl can optionally be bridged via C₁₋₄-alkylene.

[0043] According to the present invention, R⁴ represents F or Cl.

15 **[0044]** According to the present invention, R⁵ and R⁶ represent independently from one another H or unsubstituted C₁₋₄-alkyl. Preferably, R⁵ and R⁶ represent independently from one another H, CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂ or CH₂CH₂CH₂CH₃, more preferably H, CH₃, CH₂CH₃ or CH₂CH₂CH₃; still more preferably H, CH₃ or CH₂CH₃; most preferably H or CH₃. In a particularly preferred embodiment, R⁵ and R⁶ both represent CH₃.

[0045] In a particularly preferred embodiment, the compound according to the present invention is according to general formula (II) or (III), wherein X represents NR⁷ and Z represents N or CR⁹, more preferably CR⁹, and wherein R⁵ and R⁶ both represent CH₃.

[0046] In a preferred embodiment, at least one of R¹, R², R³, R⁵ and R⁶ is not H. More preferably, both R⁵ and R⁶ are not H and at least one of R¹, R², and R³ is not H. In a preferred embodiment, both R⁵ and R⁶ are not H and one of R¹, R², and R³ is not H. In another preferred embodiment, both R⁵ and R⁶ are not H and two of R¹, R², and R³ are not H.

25 In yet another preferred embodiment, R¹, R², R³, R⁵ and R⁶ are not H.

[0047] In a particularly preferred embodiment,

- the compound according to the present invention is according to general formula (II) or (III); and/or
- X represents NR⁷ and Z represents Nor CR⁹; and/or
- R⁵ and R⁶ both represent methyl; and/or
- at least one of R¹, R², and R³ is not H.

[0048] According to the present invention, R⁷ represents H or L-R⁸.

[0049] In a preferred embodiment, R⁷ represents H. According to this embodiment, preferably X represents NH and Z represents N or CR⁹; or X represents N and Z represents NH. More preferably, X represents NH and Z represents CR⁹.

[0050] Still further according to this embodiment, preferably the compound according to the present invention is according to general formula (II) or (III), wherein X represents NH and Z represents N or CR⁹, more preferably CR⁹.

[0051] In another preferred embodiment, R⁷ is not H. According to this preferred embodiment, R⁷ represents L-R⁸.

40 **[0052]** In a preferred embodiment, L represents bond; S(O); S(O)₂; C₁₋₄-alkylene; C(O); C₁₋₄-alkylene-C(O); C(O)-O; C₁₋₄-alkylene-C(O)-O; C₁₋₄-alkylene-N(H)-C(O); C₁₋₄-alkylene-N(C₁₋₆-alkyl)-C(O); C₁₋₄-alkylene-N(H)-C(O)-O or C₁₋₄-alkylene-N(C₁₋₆-alkyl)-C(O)-O. According to this embodiment, C₁₋₄-alkylene and C₁₋₆-alkyl are preferably unsubstituted.

[0053] More preferably, L represents bond; S(O); S(O)₂; C₁₋₄-alkylene; C(O); C₁₋₄-alkylene-C(O); C(O)-O; C₁₋₄-alkylene-C(O)-O; C₁₋₄-alkylene-N(H)-C(O) or C₁₋₄-alkylene-N(H)-C(O)-O; still more preferably bond; S(O); S(O)₂; CH₂; CH₂CH₂; C(CH₃)₂; CH₂CH₂CH₂; C(O); CH₂-C(O); CH₂CH₂-C(O); CH₂CH₂CH₂-C(O); C(CH₃)₂-C(O); C(O)-O; CH₂-C(O)-O; CH₂CH₂-C(O)-O; CH₂CH₂CH₂-C(O)-O; C(CH₃)₂-C(O)-O; CH₂-N(H)-C(O); CH₂CH₂-N(H)-C(O); C(CH₃)₂-N(H)-C(O); CH₂CH₂CH₂-N(H)-C(O); CH₂-N(H)-C(O)-O; CH₂CH₂-N(H)-C(O)-O; C(CH₃)₂-N(H)-C(O)-O or CH₂CH₂CH₂-N(H)-C(O)-O; most preferably bond; S(O)₂; CH₂; C(O); C(O)-O; CH₂-C(O)-O; CH₂CH₂-C(O)-O; CH₂CH₂-N(H)-C(O) or CH₂CH₂-N(H)-C(O)-O.

50 **[0054]** In a preferred embodiment, R⁸ represents C₁₋₆-alkyl; C₃₋₆-cycloalkyl or 3 to 6-membered heterocycloalkyl; wherein C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl can optionally be bridged via C₁₋₄-alkylene; and preferably wherein C₁₋₆-alkyl; C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl in each case independently from one another are unsubstituted or mono- or polysubstituted with one or more substituents selected from the group consisting of F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; OCF₃; OCF₂H; OCFH₂; O-C₁₋₆-alkyl; NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; SCF₃; SCF₂H; SCFH₂; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; phenyl and 5 or 6-membered heteroaryl.

[0055] More preferably R⁸ represents

C₁₋₆-alkyl, which is selected from the group consisting of methyl, ethyl, ethenyl (vinyl), n-propyl, 2-propyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, 2-pentyl, 3-pentyl, 2-methylbutyl, 3-methylbutyl, 3-methylbut-2-yl, 2-methylbut-2-yl, 2,2-dimethylpropyl, n-hexyl, 2-hexyl, 3-hexyl, 2-methylpentyl, 4-methylpentyl, 4-methylpent-2-yl, 2-methylpent-2-yl, 3,3-dimethylbutyl, 3,3-dimethylbut-2-yl, 3-methylpentyl, 3-methylpent-2-yl and 3-methylpent-3-yl; preferably methyl, ethyl, ethenyl (vinyl), n-propyl, 2-propyl, 1-propynyl, 2-propynyl, propenyl (-CH₂CH=CH₂, -CH=CH-CH₃, -C(=CH₂)-CH₃), n-butyl, 1-butylnyl, 2-butylnyl, 1-butenyl, 2-butenyl, isobutyl, sec-butyl and tert-butyl;

C₃₋₆-cycloalkyl, which is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopentenyl and cyclohexenyl; preferably cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl;
or

3 to 6-membered heterocycloalkyl, which is selected from the group consisting of tetrahydropyranyl, oxetanyl, oxiranyl, tetrahydrofuranyl, thiazolidinyl, tetrahydrothiophenyl, tetrahydropyridinyl, thiomorpholinyl, morpholinyl, pyrrolidinyl, 4-methylpiperazinyl, morpholinonyl, azetidiny, aziridinyl, dithiolanyl, dihydropyrrolyl, dioxanyl, dioxolanyl, dihydropyridinyl, dihydrofuranyl, dihydroisoxazolyl, dihydrooxazolyl, imidazolidinyl, isoxazolidinyl, oxazolidinyl, piperazinyl, piperidinyl, pyrazolidinyl, pyranyl, tetrahydropyrrolyl, dihydroindolyl, dihydroisoindolyl and tetrahydroindolyl; preferably tetrahydropyranyl, oxetanyl, oxiranyl and tetrahydrofuranyl;

wherein C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl can optionally be bridged via C₁₋₄-alkylene; and

wherein C₁₋₆-alkyl, C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl in each case independently from one another are unsubstituted or mono- or polysubstituted with one or more substituents selected from the group consisting of F, CN, CH₃, CH₂CH₃, CF₃, CF₂H, CFH₂, C(O)-NH₂, C(O)-N(H)(CH₃), C(O)-N(CH₃)₂, OH, NH₂, OCH₃, SCH₃, S(O)₂(CH₃), S(O)(CH₃), N(CH₃)₂, cyclopropyl and oxetanyl.

[0056] Most preferably R⁸ represents methyl, ethyl, n-propyl, 2-propyl, n-butyl, isobutyl, sec-butyl, tert-butyl, CF₃, CHF₂, CH₂F, CH₂CF₃, CH₂CHF₂, CH₂CH₂F, CHFCH₃, CHFCH₂F, CHFCHF₂, CHFCHF₃, CF₂CH₃, CF₂CH₂F, CF₂CHF₂, CF₂CF₃, CH₂CH₂CF₃, CH₂CH₂CHF₂, CH₂CH₂CH₂F, CH₂CHFCH₃, CH₂CHFCH₂F, CH₂CHFCHF₂, CH₂CHFCHF₃, CH₂CF₂CH₃, CH₂CF₂CH₂F, CH₂CF₂CHF₂, CH₂CF₂CF₃, CH₂OH, CH₂CH₂OH, C(H)(OH)CH₃, CH₂CH₂CH₂OH, C(CH₃)₂OH, C(H)(OH)CH₂CH₃, C(H)(CH₃)-CH₂OH, CH₂C(H)(OH)-CH₃, CH₂CH₂CH₂CH₂OH, CH₂CH₂C(H)(OH)CH₃, CH₂C(H)(OH)CH₂CH₃, C(H)(OH)CH₂CH₂CH₃, CH₂-C(CH₃)₂-OH, C(CH₃)₂CH₂OH, C(H)(OH)CH₂CF₃, C(H)(OH)CH₂CHF₂, C(H)(OH)CH₂CH₂F, CH₂C(H)(OH)-CF₃, CH₂C(H)(OH)-CHF₂, CH₂C(H)(OH)-CH₂F, CH₂OCH₃, CH₂CH₂OCH₃, C(H)(CH₃)-OCH₃, CH₂CH₂CH₂OCH₃, C(CH₃)₂OCH₃, C(H)(OCH₃)CH₂CH₃, C(H)(CH₃)-CH₂OCH₃, CH₂C(H)(OCH₃)-CH₃, CH₂CH₂CH₂CH₂OCH₃, CH₂CH₂C(H)(OCH₃)CH₃, CH₂C(H)(OCH₃)CH₂CH₃, C(H)(OCH₃)CH₂CH₂CH₃, CH₂-C(CH₃)₂-OCH₃, C(CH₃)₂CH₂OCH₃, CH₂NH₂, CH₂CH₂NH₂, CH₂CH₂CH₂NH₂, C(CH₃)₂NH₂, C(H)(NH₂)CH₂CH₃, C(H)(CH₃)-CH₂NH₂, CH₂C(H)(NH₂)-CH₃, CH₂CH₂CH₂CH₂NH₂, CH₂CH₂C(H)(NH₂)CH₃, CH₂C(H)(NH₂)CH₂CH₃, C(H)(NH₂)CH₂CH₂CH₃, CH₂-C(CH₃)₂-NH₂, C(CH₃)₂CH₂NH₂, CH₂N(CH₃)₂, CH₂CH₂N(CH₃)₂, CH₂CH₂CH₂N(CH₃)₂, C(CH₃)₂N(CH₃)₂, C(H)(N(CH₃)₂)CH₂CH₃, C(H)(CH₃)-CH₂N(CH₃)₂, CH₂C(H)(N(CH₃)₂)-CH₃, CH₂CH₂CH₂CH₂N(CH₃)₂, CH₂CH₂C(H)(N(CH₃)₂)CH₃, CH₂C(H)(N(CH₃)₂)CH₂CH₃, CH₂-C(CH₃)₂-N(CH₃)₂, C(CH₃)₂CH₂N(CH₃)₂, CH₂-C(O)N(CH₃)₂, CH₂CH₂-C(O)N(CH₃)₂, CH₂CH₂CH₂-C(O)N(CH₃)₂, C(CH₃)₂-C(O)N(CH₃)₂, C(H)(C(O)N(CH₃)₂)CH₂CH₃, C(H)(CH₃)-CH₂-C(O)N(CH₃)₂, CH₂C(H)(C(O)N(CH₃)₂)-CH₃, CH₂CH₂CH₂CH₂-C(O)N(CH₃)₂, CH₂CH₂C(H)(C(O)N(CH₃)₂)CH₃, CH₂C(H)(C(O)N(CH₃)₂)CH₂CH₃, C(H)(C(O)N(CH₃)₂)CH₂CH₂CH₃, CH₂-C(CH₃)₂-C(O)N(CH₃)₂, C(CH₃)₂CH₂-C(O)N(CH₃)₂, (CH₂)₀₋₂-cyclopropyl, (CH₂)₀₋₂-cyclobutyl, (CH₂)₀₋₂-cyclopentyl, (CH₂)₀₋₂-cyclohexyl, (CH₂)₀₋₂-tetrahydropyranyl, (CH₂)₀₋₂-oxetanyl, (CH₂)₀₋₂-oxiranyl or (CH₂)₀₋₂-tetrahydrofuranyl.

[0057] Preferred embodiments (E1 to E45) encompass those wherein L and R⁸ have the meaning as given in the table below:

	L	R ⁸		L	R ⁸
E1	bond	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl	E24	C ₁₋₆ -alkylene-C(O)-O	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl
E2	bond	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl	E25	C ₁₋₆ -alkylene-N(H)-C(O)	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl
E3	bond	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl	E26	C ₁₋₆ -alkylene-N(H)-C(O)	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl

(continued)

	L	R ⁸		L	R ⁸
E4	S(O)	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl	E27	C ₁₋₆ -alkylene-N (H)-C(O)	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl
E5	S(O)	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl	E28	C ₁₋₆ -alkylene-N (C ₁₋₁₀ -alkyl)-C(O)	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl
E6	S(O)	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl	E29	C ₁₋₆ -alkylene-N (C ₁₋₁₀ -alkyl)-C(O)	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl
E7	S(O) ₂	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl	E30	C ₁₋₆ -alkylene-N (C ₁₋₁₀ -alkyl)-C(O)	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl
E8	S(O) ₂	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl	E31	C ₁₋₆ -alkylene-N (H)-C(O)-O	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl
E9	S(O) ₂	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl	E32	C ₁₋₆ -alkylene-N (H)-C(O)-O	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl
E10	C ₁₋₆ -alkylene	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl	E33	C ₁₋₆ -alkylene-N (H)-C(O)-O	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl
E11	C ₁₋₆ -alkylene	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl	E34	C ₁₋₆ -alkylene-N (C ₁₋₁₀ -alkyl)-C (O)-O	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl
E12	C ₁₋₆ -alkylene	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl	E35	C ₁₋₆ -alkylene-N (C ₁₋₁₀ -alkyl)-C (O)-O	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl
E13	C(O)	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl	E36	C ₁₋₆ -alkylene-N (C ₁₋₁₀ -alkyl)-C (O)-O	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl
E14	C(O)	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl	E37	O	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl
E15	C(O)	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl	E38	O	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl
E16	C ₁₋₆ -alkylene- C(O)	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl	E39	O	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl
E17	C ₁₋₆ -alkylene- C(O)	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl	E40	NH	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl
E18	C ₁₋₆ -alkylene- C(O)	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl	E41	NH	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl
E19	C(O)-O	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl	E42	NH	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl
E20	C(O)-O	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl	E43	N(C ₁₋₁₀ -alkyl)	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl
E21	C(O)-O	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl	E44	N(C ₁₋₁₀ -alkyl)	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl
E22	C ₁₋₆ -alkylene- C(O)-O	(CH ₂) ₀₋₆ -C ₁₋₁₀ -alkyl	E45	N(C ₁₋₁₀ -alkyl)	(CH ₂) ₀₋₆ -3 to 7 membered heterocycloalkyl
E23	C ₁₋₆ -alkylene- C(O)-O	(CH ₂) ₀₋₆ -C ₃₋₁₀ -cycloalkyl			

[0058] Preferably,

L represents bond; S(O); S(O)₂; C₁₋₄-alkylene; C(O); C₁₋₄-alkylene-C(O); C(O)-O; C₁₋₄-alkylene-C(O)-O;

C₁₋₄-alkylene-N(H)-C(O); C₁₋₄-alkylene-N(C₁₋₆-alkyl)-C(O); C₁₋₄-alkylene-N(H)-C(O)-O or C₁₋₄-alkylene-N(C₁₋₆-alkyl)-C(O)-O; and

R⁸ represents C₁₋₆-alkyl; C₃₋₆-cycloalkyl or 3 to 6-membered heterocycloalkyl; wherein C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl can optionally be bridged via C₁₋₄-alkylene.

[0059] In a particularly preferred embodiment,

L represents bond; S(O); S(O)₂; C₁₋₄-alkylene; C(O); C₁₋₄-alkylene-C(O); C(O)-O; C₁₋₄-alkylene-C(O)-O; C₁₋₄-alkylene-N(H)-C(O) or C₁₋₄-alkylene-N(H)-C(O)-O;

R⁸ represents C₁₋₆-alkyl; C₃₋₆-cycloalkyl or 3 to 6-membered heterocycloalkyl; wherein C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl can optionally be bridged via C₁₋₄-alkylene.

[0060] More preferably,

L represents bond; S(O); S(O)₂; C₁₋₄-alkylene; C(O); C₁₋₄-alkylene-C(O); C(O)-O; C₁₋₄-alkylene-C(O)-O; C₁₋₄-alkylene-N(H)-C(O) or C₁₋₄-alkylene-N(H)-C(O)-O; still more preferably bond; S(O); S(O)₂; CH₂; CH₂CH₂; C(CH₃)₂; CH₂CH₂CH₂; C(O); CH₂-C(O); CH₂CH₂-C(O); CH₂CH₂CH₂-C(O); C(CH₃)₂-C(O); C(O)-O; CH₂-C(O)-O; CH₂CH₂-C(O)-O; CH₂CH₂CH₂-C(O)-O; C(CH₃)₂-C(O)-O; CH₂-N(H)-C(O); CH₂CH₂-N(H)-C(O); C(CH₃)₂-N(H)-C(O); CH₂CH₂CH₂-N(H)-C(O); CH₂-N(H)-C(O)-O; CH₂CH₂-N(H)-C(O)-O; C(CH₃)₂-N(H)-C(O)-O or CH₂CH₂CH₂-N(H)-C(O)-O; most preferably bond; S(O)₂; CH₂; C(O); C(O)-O; CH₂-C(O)-O; CH₂CH₂-C(O)-O; CH₂CH₂-N(H)-C(O) or CH₂CH₂-N(H)-C(O)-O; and

R⁸ represents C₁₋₆-alkyl; C₃₋₆-cycloalkyl or 3 to 6-membered heterocycloalkyl; wherein C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl can optionally be bridged via C₁₋₄-alkylene; and preferably wherein C₁₋₆-alkyl; C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl in each case independently from one another are unsubstituted or mono- or polysubstituted with one or more substituents selected from the group consisting of F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; OCF₃; OCF₂H; OCFH₂; O-C₁₋₆-alkyl; NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; SCF₃; SCF₂H; SCFH₂; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; phenyl and 5 or 6-membered heteroaryl.

[0061] Preferably, **R⁹** represents H; F; Cl; Br; I; CN; C₁₋₆-alkyl; C₃₋₆-cycloalkyl, 3 to 6-membered heterocycloalkyl; S(O)-(C₁₋₆-alkyl); S(O)-(C₃₋₆-cycloalkyl); S(O)-(3 to 6-membered heterocycloalkyl); S(O)₂-(C₁₋₆-alkyl); S(O)₂-(C₃₋₆-cycloalkyl); S(O)₂-(3 to 6-membered heterocycloalkyl); P(O)-(C₁₋₆-alkyl)₂; P(O)(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); P(O)(C₁₋₆-alkyl)(3 to 6-membered heterocycloalkyl); P(O)-(O-C₁₋₆-alkyl)₂; P(O)(O-C₁₋₆-alkyl)(O-C₃₋₆-cycloalkyl); P(O)(O-C₁₋₆-alkyl)(O-(3 to 6-membered heterocycloalkyl)); O-C₁₋₆-alkyl; S-C₁₋₆-alkyl; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; C(O)-C₁₋₆-alkyl; C(O)-O-C₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; O-C₃₋₆-cycloalkyl; N(H)(C₃₋₆-cycloalkyl); N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); C(O)-C₃₋₆-cycloalkyl; C(O)-O-C₃₋₆-cycloalkyl; C(O)-N(H)(C₃₋₆-cycloalkyl); C(O)-N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); O-(3 to 6-membered heterocycloalkyl); N(H)(3 to 6-membered heterocycloalkyl); N(C₁₋₆-alkyl)(3 to 6-membered heterocycloalkyl); C(O)-3 to 6-membered heterocycloalkyl; C(O)-O-(3 to 6-membered heterocycloalkyl); C(O)-N(H)(3 to 6-membered heterocycloalkyl) or C(O)-N(C₁₋₆-alkyl)(3 to 6-membered heterocycloalkyl);

wherein C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl can optionally be bridged via C₁₋₄-alkylene; preferably wherein C₁₋₆-alkyl, C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl is in each case unsubstituted.

[0062] More preferably, **R⁹** represents H; F; Cl; Br; I; CN; methyl, ethyl, n-propyl, 2-propyl, n-butyl, isobutyl, sec-butyl, tert-butyl, CF₃, CHF₂, CH₂F, CH₂CF₃, CH₂CHF₂, CH₂CH₂F, cyclopropyl, cyclobutyl, tetrahydropyranyl, oxetanyl, oxiranyl, tetrahydrofuranyl, S(O)-(CH₃); S(O)-(CH₂CH₃); S(O)-((CH₂)₀₋₂-cyclopropyl); S(O)-((CH₂)₀₋₂-cyclobutyl); S(O)-((CH₂)₀₋₂-cyclopentyl); S(O)-((CH₂)₀₋₂-cyclohexyl); S(O)-((CH₂)₀₋₂-tetrahydropyranyl); S(O)-((CH₂)₀₋₂-oxetanyl); S(O)-((CH₂)₀₋₂-oxiranyl); S(O)-((CH₂)₀₋₂-tetrahydrofuranyl); S(O)₂-(CH₃); S(O)₂-(CH₂CH₃); S(O)₂-(CH₂)₀₋₂-cyclopropyl); S(O)₂-(CH₂)₀₋₂-cyclobutyl); S(O)₂-(CH₂)₀₋₂-cyclopentyl); S(O)₂-(CH₂)₀₋₂-cyclohexyl); S(O)₂-(CH₂)₀₋₂-tetrahydropyranyl); S(O)₂-(CH₂)₀₋₂-oxetanyl); S(O)₂-(CH₂)₀₋₂-oxiranyl); S(O)₂-(CH₂)₀₋₂-tetrahydrofuranyl); O-CH₃, O-CH₂CH₃, O-CH₂CH₂CH₃, O-C(H)(CH₃)₂, N(H)(CH₃), N(H)(CH₂CH₃), N(CH₃)₂ or N(CH₃)(CH₂CH₃).

[0063] In a particularly preferred embodiment, **R⁹** represents H; F; CN; methyl; ethyl; n-propyl; 2-propyl; CF₃; CH₂CF₃; CHF₂; CH₂CHF₂; CH₂F; CH₂CH₂F; S(O)-CH₃; S(O)-CH₂CH₃; S(O)-CH₂CH₂CH₃; S(O)-CH(CH₃)₂; S(O)₂-CH₃; S(O)₂-CH₂CH₃; S(O)₂-CH₂CH₂CH₃ or S(O)₂-CH(CH₃)₂.

[0064] Preferably, **R¹⁰** represents H; F; Cl; Br; I; CN; C₁₋₆-alkyl; C₃₋₆-cycloalkyl, 3 to 6-membered heterocycloalkyl; S(O)-(C₁₋₆-alkyl); S(O)-(C₃₋₆-cycloalkyl); S(O)-(3 to 6-membered heterocycloalkyl); S(O)₂-(C₁₋₆-alkyl); S(O)₂-(C₃₋₆-cy-

cloalkyl); S(O)₂-(3 to 6-membered heterocycloalkyl); P(O)-(C₁₋₆-alkyl)₂; P(O)(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); P(O)(C₁₋₆-alkyl)(3 to 6-membered heterocycloalkyl); P(O)-(O-C₁₋₆-alkyl)₂; P(O)(O-C₁₋₆-alkyl)(O-C₃₋₆-cycloalkyl); P(O)(O-C₁₋₆-alkyl)(O-(3 to 6-membered heterocycloalkyl)); O-C₁₋₆-alkyl; S-C₁₋₆-alkyl; N(H)(C₁₋₆-alkyl), N(C₁₋₆-alkyl)₂; C(O)-C₁₋₆-alkyl; C(O)-O-C₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; O-C₃₋₆-cycloalkyl; N(H)(C₃₋₆-cycloalkyl), N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); C(O)-C₃₋₆-cycloalkyl; C(O)-O-C₃₋₆-cycloalkyl; C(O)-N(H)(C₃₋₆-cycloalkyl); C(O)-N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); O-(3 to 6-membered heterocycloalkyl); N(H)(3 to 6-membered heterocycloalkyl), N(C₁₋₆-alkyl)(3 to 6-membered heterocycloalkyl); C(O)-3 to 6-membered heterocycloalkyl; C(O)-O-(3 to 6-membered heterocycloalkyl); C(O)-N(H)(3 to 6-membered heterocycloalkyl) or C(O)-N(C₁₋₆-alkyl)(3 to 6-membered heterocycloalkyl);

wherein C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl can optionally be bridged via C₁₋₄-alkylene; and preferably wherein C₁₋₆-alkyl; C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl in each case independently from one another are unsubstituted or mono- or polysubstituted with one or more substituents selected from the group consisting of F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; OCF₃; OCF₂H; OCFH₂; O-C₁₋₆-alkyl; NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; SCF₃; SCF₂H; SCFH₂; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; phenyl and 5 or 6-membered heteroaryl.

[0065] In another preferred embodiment, R¹⁰ represents H; F; Cl; Br; CN; C₁₋₆-alkyl; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; P(O)-(C₁₋₆-alkyl)₂; P(O)(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); P(O)(C₁₋₆-alkyl)(3 to 6-membered heterocycloalkyl); P(O)-(O-C₁₋₆-alkyl)₂; P(O)(O-C₁₋₆-alkyl)(O-C₃₋₆-cycloalkyl); P(O)(O-C₁₋₆-alkyl)(O-(3 to 6-membered heterocycloalkyl));

preferably wherein C₁₋₆-alkyl; C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl in each case independently from one another are unsubstituted or mono- or polysubstituted with one or more substituents selected from the group consisting of F; Cl; Br; I; CN; C₁₋₄-alkyl; CF₃; CF₂H; CFH₂; OH; OCF₃; OCF₂H; OCFH₂; O-C₁₋₄-alkyl; NH₂; N(H)(C₁₋₄-alkyl); N(C₁₋₄-alkyl)₂; S-C₁₋₄-alkyl; S(O)-C₁₋₄-alkyl and S(O)₂-C₁₋₄-alkyl; more preferably F, Cl, Br, CF₃, OCH₃, OCF₃, OCHF₂, OCH₂F, OH and NH₂.

[0066] More preferably, R¹⁰ represents H; F; Cl; Br; I; CN; methyl, ethyl, n-propyl, 2-propyl, 1-propynyl, 2-propynyl, propenyl (-CH₂CH=CH₂, -CH=CH-CH₃, -C(=CH₂)-CH₃), n-butyl, 1-butyryl, 2-butyryl, 1-butenyl, 2-butenyl, 3-methyl-1-butyryl, n-butyl, isobutyl, sec-butyl, tert-butyl, CF₃, CHF₂, CH₂F, CH₂CF₃, CH₂CHF₂, CH₂CH₂F, CH₂CHFCH₃, CH₂CHFCH₂F, CH₂CHFCHF₂, CH₂CHFCH₂F, CH₂CF₂CH₃, CH₂CF₂CH₂F, CH₂CF₂CHF₂, CH₂CF₂CF₃, CH₂OH, CH₂CH₂OH, C(H)(OH)CH₃, CH₂CH₂CH₂OH, C(CH₃)₂OH, C(H)(OH)CH₂CH₃, C(H)(CH₃)-CH₂OH, CH₂C(H)(OH)-CH₃, CH₂CH₂CH₂CH₂OH, CH₂CH₂C(H)(OH)CH₃, CH₂C(H)(OH)CH₂CH₃, C(H)(OH)CH₂CH₂CH₃, CH₂-C(CH₃)₂-OH, C(CH₃)₂CH₂OH, C=C-C(H)(OH)CH₃, C(H)(OH)-C=C-CH₃, C=C-C(CH₃)(OH)CH₃, C(CH₃)(OH)-C=C-CH₃, C(H)(OH)CH₂CF₃, C(H)(OH)CH₂CHF₂, C(H)(OH)CH₂CH₂F, CH₂C(H)(OH)-CF₃, CH₂C(H)(OH)-CHF₂, CH₂C(H)(OH)-CH₂F, CH₂OCH₃, CH₂CH₂OCH₃, C(H)(CH₃)-OCH₃, CH₂CH₂CH₂OCH₃, C(CH₃)₂OCH₃, C(H)(OCH₃)CH₂CH₃, C(H)(CH₃)-CH₂OCH₃, CH₂C(H)(OCH₃)-CH₃, CH₂CH₂CH₂CH₂OCH₃, CH₂CH₂C(H)(OCH₃)CH₃, CH₂C(H)(OCH₃)CH₂CH₃, C(H)(OCH₃)CH₂CH₂CH₃, CH₂-C(CH₃)₂-OCH₃, C(CH₃)₂CH₂OCH₃, CH₂NH₂, CH₂CH₂NH₂, CH₂CH₂CH₂NH₂, C(CH₃)₂NH₂, C(H)(NH₂)CH₂CH₃, C(H)(CH₃)-CH₂NH₂, CH₂C(H)(NH₂)-CH₃, CH₂CH₂CH₂CH₂NH₂, CH₂CH₂C(H)(NH₂)CH₃, CH₂C(H)(NH₂)CH₂CH₃, C(H)(NH₂)CH₂CH₂CH₃, CH₂-C(CH₃)₂-NH₂, C(CH₃)₂CH₂NH₂, C=C-C(H)(NH₂)CH₃, C(H)(NH₂)-C=C-CH₃, C≡C-C(CH₃)(NH₂)CH₃, C(CH₃)(NH₂)-C≡C-CH₃, CH₂N(CH₃)₂, CH₂CH₂N(CH₃)₂, CH₂CH₂CH₂N(CH₃)₂, C(CH₃)₂N(CH₃)₂, C(H)(N(CH₃)₂)CH₂CH₃, C(H)(CH₃)-CH₂N(CH₃)₂, CH₂C(H)(N(CH₃)₂)-CH₃, CH₂CH₂CH₂CH₂N(CH₃)₂, CH₂CH₂C(H)(N(CH₃)₂)CH₃, CH₂C(H)(N(CH₃)₂)CH₂CH₃, C(H)(N(CH₃)₂)CH₂CH₂CH₃, CH₂-C(CH₃)₂-N(CH₃)₂, C(CH₃)₂CH₂N(CH₃)₂, S(O)-CH₃, S(O)-CH₂CH₃, S(O)-((CH₂)₀₋₂-cyclopropyl), S(O)-((CH₂)₀₋₂-cyclobutyl), S(O)-((CH₂)₀₋₂-cyclopentyl), S(O)-((CH₂)₀₋₂-cyclohexyl), S(O)₂-CH₃, S(O)₂-CH₂CH₃, S(O)₂-((CH₂)₀₋₂-cyclopropyl), S(O)₂-((CH₂)₀₋₂-cyclobutyl), S(O)₂-((CH₂)₀₋₂-cyclopentyl), S(O)₂-((CH₂)₀₋₂-cyclohexyl), P(O)-(CH₃)₂, P(O)-(CH₃)(CH₂CH₃), P(O)-(CH₃)((CH₂)₀₋₂-cyclopropyl), P(O)-(CH₃)((CH₂)₀₋₂-cyclobutyl), P(O)-(CH₃)((CH₂)₀₋₂-cyclopentyl), P(O)-(CH₃)((CH₂)₀₋₂-cyclohexyl), (CH₂)₀₋₂-cyclopropyl, (CH₂)₀₋₂-cyclobutyl, (CH₂)₀₋₂-tetrahydropyranyl, (CH₂)₀₋₂-oxetanyl, (CH₂)₀₋₂-oxiranyl, (CH₂)₀₋₂-tetrahydrofuranyl, O-CH₃, O-CH₂CH₃, O-CH₂CH₂CH₃, O-C(H)(CH₃)₂, N(H)(CH₃), N(H)(CH₂CH₃), N(CH₃)₂ or N(CH₃)(CH₂CH₃).

[0067] In a preferred embodiment, at least one of R⁹ and R¹⁰ is H.

[0068] According to the present invention, n represents 0, 1, 2 or 3. In a preferred embodiment, n represents 0. In another preferred embodiment, n represents 1. In yet another preferred embodiment, n represents 2. In still another preferred embodiment, n represents 3. More preferably, n represents 1 or 2, most preferably 1.

[0069] Preferably, R¹¹ represents F; Cl; Br; I; CN; C₁₋₆-alkyl; O-C₁₋₆-alkyl; NO₂; OH; NH₂; C₃₋₆-cycloalkyl; 3 to 6-membered heterocycloalkyl; S(O)-(C₁₋₆-alkyl); S(O)₂-(C₁₋₆-alkyl); P(O)-(C₁₋₆-alkyl)₂; O-C₁₋₆-alkyl; N(H)(C₁₋₆-alkyl), N(C₁₋₆-alkyl)₂;

preferably wherein C₁₋₆-alkyl; C₃₋₆-cycloalkyl and 3 to 6-membered heterocycloalkyl in each case independently from

one another are unsubstituted or mono- or polysubstituted with one or more substituents selected from the group consisting of F; Cl; Br; I; CN; C₁₋₄-alkyl; CF₃; CF₂H; CFH₂; OH; OCF₃; OCF₂H; OCFH₂; O-C₁₋₄-alkyl; NH₂; N(H)(C₁₋₄-alkyl); N(C₁₋₄-alkyl)₂; S-C₁₋₄-alkyl; S(O)-C₁₋₄-alkyl and S(O)₂-C₁₋₄-alkyl; more preferably F, Cl, Br, CF₃, OCH₃, OCF₃, OCHF₂, OCH₂F, OH and NH₂.

[0070] In a particularly preferred embodiment, R¹¹ represents F; Cl; Br; I; CN; C₁₋₆-alkyl or O-C₁₋₆-alkyl; preferably wherein C₁₋₆-alkyl in each case independently from one another is unsubstituted or mono- or polysubstituted with one or more substituents selected from the group consisting of F and CF₃.

[0071] More preferably, R¹¹ represents F; Cl; Br; I; CN; CH₃, CH₂CH₃, O-CH₃ or O-CH₂CH₃.

[0072] In a particularly preferred embodiment, R¹¹ represents F; Cl; Br; I; CN; C₁₋₆-alkyl or O-C₁₋₆-alkyl; and/or n represents 0, 1 or 2.

[0073] In a preferred embodiment, the compound according to the present invention is selected from the group consisting of

38 8-(3-Cyclopropyl-5-fluoro-1H-indol-7-yl)-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

48 9-Ethyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

49 9-Ethyl-6-fluoro-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline

53 9-Ethyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

54 6-Fluoro-8-[6-fluoro-1-(2-methoxy-ethyl)-1H-indol-4-yl]-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

55 2-[6-Fluoro-4-(6-fluoro-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-1-yl]-EtOH

56 6-Fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

57 6-Fluoro-9-methoxy-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline

58 8-[1-(2,2-Difluoro-ethyl)-6-fluoro-1H-indol-4-yl]-6-fluoro-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

59 1-Cyclopropyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-9-methoxy-4,4-dimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

60 1-Cyclopropyl-6-fluoro-9-methoxy-4,4-dimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline

61 2-[4-(1-Cyclopropyl-6-fluoro-9-methoxy-4,4-dimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluoro-1H-indol-1-yl]-EtOH

62 1-Cyclopropyl-6-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-9-methoxy-4,4-dimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

63 1-Cyclopropyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

64 1-Cyclopropyl-6-fluoro-4,4,9-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline

65 2-[4-(1-Cyclopropyl-6-fluoro-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluoro-1H-indol-1-yl]-EtOH

66 1-Cyclopropyl-6-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

300 6-Fluoro-8-[1-(2-methoxy-ethyl)-1H-indol-4-yl]-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

301 2-[4-(6-Fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-1-yl]-EtOH

304 6-Fluoro-1,4,4,9-tetramethyl-8-(1-methyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline

305 8-(1-Cyclopropyl-1H-indol-4-yl)-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

308 1-Ethyl-6-fluoro-4,4,9-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline

316 1-Ethyl-6-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

322 1-Ethyl-6-fluoro-8-[1-(2-methoxy-ethyl)-1H-indol-4-yl]-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

323 1-Ethyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

326 8-[1-(2,2-Difluoro-ethyl)-1H-indol-4-yl]-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

- 5 **340** 6-Fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 349** 6-Fluoro-8-[1-(fluoro-methylsulfonyl)-1H-indol-4-yl]-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 350** 8-[1-(Ethylsulfonyl)-1H-indol-4-yl]-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 352** 6-Fluoro-1,4,4,9-tetramethyl-8-(3-methyl-1H-indazol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 355** 1-Ethyl-6-fluoro-4,4,9-trimethyl-8-(3-methyl-1H-indol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 10 **366** 6-Fluoro-8-[1-(isopropylsulfonyl)-1H-indol-4-yl]-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 374** 1-Ethyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 375** [2-[4-(6-Fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-1-yl]-ethyl]-dimethylamine
- 15 **378** 6-Fluoro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 389** 1-Ethyl-6-fluoro-4,4,9-trimethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 391** 8-[1-(Cyclopropylsulfonyl)-1H-indol-4-yl]-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 446** 8-[1-(Ethylsulfonyl)-6-fluoro-1H-indol-4-yl]-6,7-difluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 20 **447** 6,7-Difluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 448** 6,7-Difluoro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 449** 2-[4-(6,7-Difluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluoro-1H-indol-1-yl]-EtOH
- 25 **450** 8-(1-Cyclopropyl-1H-indol-4-yl)-6-fluoro-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline-9-carbonitrile
- 451** 6,7-Difluoro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 452** 6,7-Difluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 453** [6-Fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-MeOH
- 30 **454** 6-Chloro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 455** 9-(Difluoro-methyl)-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 456** 6-Chloro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 35 **457** 6-Chloro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 458** 6-Chloro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 459** 9-(Difluoro-methyl)-6-fluoro-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 40 **460** 9-(Difluoro-methyl)-8-[1-(ethylsulfonyl)-6-fluoro-1H-indol-4-yl]-6-fluoro-1,4,4-trimethyl-SH-[1,2,4]triazolo[4,3-a]quinoxaline
- 461** 9-(Difluoro-methyl)-8-[1-(ethylsulfonyl)-1H-indol-4-yl]-6-fluoro-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 462** 9-(Difluoro-methyl)-6-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4-trimethyl-SH-[1,2,4]triazolo[4,3-a]quinoxaline
- 45 **463** 2-[4-(6-Chloro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluoro-1H-indol-1-yl]-EtOH
- 464** 2-[4-(9-Ethyl-6-fluoro-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluoro-1H-indol-1-yl]-EtOH
- 465** 6-Fluoro-8-(6-fluoro-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 466** 9-(Difluoro-methyl)-6-fluoro-1,4,4-trimethyl-8-(3-methyl-1H-indazol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 50 **467** 9-Cyclopropyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 468** 6-Fluoro-8-(1H-indol-4-yl)-1,4,4,9-tetramethyl-SH-[1,2,4]triazolo[4,3-a]quinoxaline
- 469** 6-Fluoro-8-(1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-SH-[1,2,4]triazolo[4,3-a]quinoxaline
- 55 **470** 6-Fluoro-8-(6-fluoro-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 471** 9-Cyclopropyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 472** 9-Cyclopropyl-6-fluoro-1,4,4-trimethyl-8-(3-methyl-1H-indol-7-yl)-SH-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

	473	6-Fluoro-9-methoxy-1,4,4-trimethyl-8-[6-(trifluoromethyl)-1H-indol-4-yl]-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	474	9-Cyclopropyl-6-fluoro-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
5	475	6-Fluoro-1,4,4,9-tetramethyl-8-(6-methyl-1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	476	6-Fluoro-9-methoxy-1,4,4-trimethyl-8-(6-methyl-1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
10	477	4-(6-Fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indole-6-carbonitrile
	478	8-(3-Cyclopropyl-5-fluoro-1H-indol-7-yl)-6-fluoro-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	479	[8-[1-(Ethylsulfonyl)-6-fluoro-1H-indol-4-yl]-6-fluoro-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-methyl-amine
15	480	2-[6-Fluoro-4-(6-fluoro-1,4,4-trimethyl-9-methylamino-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-1-yl]-EtOH
	481	[8-(1-Cyclopropyl-1H-indol-4-yl)-6-fluoro-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-methyl-amine
20	482	6-Chloro-9-methoxy-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	483	6-Chloro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	484	6-Fluoro-1,4,4,9-tetramethyl-8-(1-methyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	485	6-Fluoro-9-methoxy-1,4,4-trimethyl-8-(1-methyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
25	486	[6-Fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-methyl-amine
	487	[6-Fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-methyl-amine
30	488	6-Chloro-8-(6-fluoro-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	489	6-Fluoro-8-(7-fluoro-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	490	6-Fluoro-8-(7-fluoro-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	491	8-[1-(Cyclopropyl-methylsulfonyl)-1H-indol-4-yl]-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
35	492	6-Fluoro-9-methoxy-1,4,4-trimethyl-8-[1-methylsulfonyl-6-(trifluoromethyl)-1H-indol-4-yl]-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	493	6-Chloro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	494	6-Fluoro-8-(7-fluoro-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
40	495	6-Fluoro-8-(6-fluoro-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	496	6-Fluoro-8-(6-fluoro-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	497	8-[1-(Cyclopropyl-methylsulfonyl)-6-fluoro-1H-indol-4-yl]-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
45	498	6,7-Difluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	499	6,7-Difluoro-9-methoxy-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	500	6,7-Difluoro-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-9-(trifluoromethyl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
50	501	6,7-Difluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	502	6-Chloro-7-fluoro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	503	6-Chloro-7-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
55	504	6-Fluoro-8-(1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
	505	8-[1-(Cyclopropylsulfonyl)-1H-indol-4-yl]-6-fluoro-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline


(continued)

- 506 8-(6-Chloro-1H-indol-4-yl)-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 507 6-Fluoro-1,4,4,9-tetramethyl-8-[6-(trifluoromethyl)-1H-indazol-4-yl]-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 5 508 6-Chloro-7-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 509 6-Chloro-7-fluoro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 10 510 6-Chloro-7-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 511 8-(1-Cyclopropyl-1H-indol-4-yl)-6-fluoro-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 512 6,7-Difluoro-8-(1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 513 6,7-Difluoro-1,4,4,9-tetramethyl-8-(3-methyl-1H-indol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 15 514 6,7-Difluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 515 6-Fluoro-8-(5-fluoro-1H-indol-7-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 516 6,7-Difluoro-8-(6-fluoro-1H-indol-4-yl)-1,4,4-trimethyl-9-(trifluoromethyl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 517 6-Fluoro-8-(7-fluoro-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 518 6,7-Difluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-9-(trifluoromethyl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 20 519 6,7-Difluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4-trimethyl-9-(trifluoromethyl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 520 4-(6,7-Difluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indole-7-carbonitrile
- 521 6,7-Difluoro-8-(6-fluoro-1-methyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 25 522 6,7-Difluoro-8-(6-methoxy-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 523 6-Fluoro-8-(6-methoxy-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 30 524 6-Fluoro-9-methoxy-8-(6-methoxy-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 525 7-Chloro-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 526 7-Chloro-6-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
- 35 527 7-Chloro-6-fluoro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline

[0074] in the form of the free compound or a physiologically acceptable salt thereof.

[0075] The compounds according to the present invention can be synthesized by standard reactions in the field of organic chemistry known to the person skilled in the art or in a manner as described herein (cf. Reaction Scheme 1 below) or analogously. The reaction conditions in the synthesis routes described herein are known to the skilled person and are for some cases also exemplified in the Examples described herein.

Reaction scheme 1:



[0076] Substituted indole/indazole moiety in compounds of formula (VII) can be introduced by subjecting a compound of formula (IV) in a metal catalyzed C-C coupling reaction. Metal catalyzed C-C coupling reactions are known in the art (cf. Metal Catalyzed Cross-Coupling Reactions and More, 3 Volume Set Wiley, 2014; Angew. Chem. Int. Ed., 2012, 51, 5062 - 5085). Favorable C-C coupling reactions are palladium catalyzed cross coupling reactions (cf. Angew. Chem., 2005, 117, 4516 - 4563). Triazole cyclization of compound (II) gives access to compounds of general formula (III). Triazole formation on quinoxalines is known in the art (cf. Heterocycles, 1992, 34, 771 - 780; Biological and Pharmaceutical Bulletin, 2005, 28, 1216 - 1220). Electrophilic aromatic bromination of compound (III) gives compound (IV). Bromination reactions of aromatic compounds are generally known (cf. Science of Synthesis, Compounds with One Saturated Carbon-Heteroatom Bond, Volume 35, Houben-Weyl, 2007). If desired, compound of formula (II) can be brominated to compound (V) which can undergo a metal catalyzed cross-coupling reaction to give compound (VI). Triazole formation leads to an alternate route to compounds of general formula (VII). Copper mediated quinoxaline cyclization of compound (I) to compound (II) is known in the art (cf. Adv. Synth. Catal., 2010, 352, 2531 - 2537). Compounds of formula (I) are commercially available or can be prepared according to methods known in the art.

[0077] The compounds according to the present invention can be produced in the manner described here or in an analogous manner.

[0078] In a preferred embodiment, the compounds according to the present invention are modulators of the glucocorticoid receptor. In the sense of the present invention, the term "selective modulator of the glucocorticoid receptor (glucocorticoid receptor modulator)" preferably means that the respective compound exhibits in a cellular target engagement assay for agonistic or antagonistic potency on the glucocorticoid receptor an EC₅₀ or IC₅₀ value on the glucocorticoid receptor of at most 15 μ M ($10 \cdot 10^{-6}$ mol/L) or at most 10 μ M; more preferably at most 1 μ M; still more preferably at most 500 nM (10^{-9} mol/L); yet more preferably at most 300 nM; even more preferably at most 100 nM; most preferably at most 10 nM; and in particular at most 1 nM.

[0079] The person skilled in the art knows how to test compounds for modulation (agonistic or antagonistic) of the activity of the glucocorticoid receptor. Preferred target engagement assays for testing compounds for their agonistic or antagonistic potency (EC₅₀, IC₅₀) on the glucocorticoid receptor are described herein below:

Glucocorticoid receptor cell-based assays

[0080] Potential selective glucocorticoid receptor modulators of this intervention can be tested for modulation of the

activity of the glucocorticoid receptor using cell-based assays. These assays involve a Chinese hamster ovary (CHO) cell line which contains fragments of the glucocorticoid receptor as well as fusion proteins. The glucocorticoid receptor fragments used are capable of binding the ligand (e.g. beclomethasone) to identify molecules that compete for binding with glucocorticoid receptor ligands. In more detail, the glucocorticoid receptor ligand binding domain is fused to the DNA binding domain (DBD) of the transcriptionfactor GAL4 (GAL4 DBD-GR) and is stably integrated into a CHO cell line containing a GAL4-UAS-Luciferase reporter construct. To identify selective glucocorticoid receptor modulators, the reporter cell line is incubated with the molecules using an 8-point half-log compound dilution curve for several hours. After cell lysis the luminescence that is produced by luciferase after addition of the substrate is detected and EC₅₀ or IC₅₀ values can be calculated. Engagement of molecules which induce gene expression via glucocorticoid receptor binding to the DNA leads to expression of the luciferase gene under the control of the fusion protein GAL4 DBD-GR and therefore to a dose-dependent increase of the luminescence signal. Binding of molecules which repress beclomethasone-induced gene expression of the luciferase gene under the control of the fusion protein GAL4 DBD-GR leads to a dose-dependent reduction of the luminescence signal.

[0081] In a preferred embodiment, the compound according to the present invention exhibits in a cellular target engagement assay for agonistic or antagonistic potency on the glucocorticoid receptor an EC₅₀ or IC₅₀ value on the glucocorticoid receptor of at most 1 μ M (10^{-6} mol/L); still more preferably at most 500 nM (10^{-9} mol/L); yet more preferably at most 300 nM; even more preferably at most 100 nM; most preferably at most 50 nM; and in particular at most 10 nM or at most 1 nM.

[0082] In a preferred embodiment, the compound according to the present invention exhibits in a cellular target engagement assay for agonistic or antagonistic potency on the glucocorticoid receptor an EC₅₀ or IC₅₀ value on the glucocorticoid receptor in the range of from 0.1 nM (10^{-9} mol/L) to 1000 nM; still more preferably 1 nM to 800 nM; yet more preferably 1 nM to 500 nM; even more preferably 1 nM to 300 nM; most preferably 1 nM to 100 nM; and in particular 1 nM to 80 nM.

[0083] Preferably, the compounds according to the present invention are useful as selective modulators of the glucocorticoid receptor.

[0084] Therefore, the compounds according to the present invention are preferably useful for the in vivo treatment or prevention of diseases in which participation of the glucocorticoid receptor is implicated.

[0085] The present invention therefore further relates to a compound according to the present invention for use in the modulation of glucocorticoid receptor activity.

[0086] Therefore, another aspect of the present invention relates to a compound according to the present invention for use in the treatment and/or prophylaxis of a disorder which is mediated at least in part by the glucocorticoid receptor. Still another aspect of the present invention relates to a method of treatment of a disorder which is mediated at least in part by the glucocorticoid receptor comprising the administration of a therapeutically effective amount of a compound according to the present invention to a subject in need thereof, preferably a human.

[0087] A further aspect of the invention relates to the use of a compound according to the present invention as medicament.

[0088] Another aspect of the present invention relates to a pharmaceutical dosage form comprising a compound according to the present invention. Preferably, the pharmaceutical dosage form comprises a compound according to the present invention and one or more pharmaceutical excipients such as physiologically acceptable carriers, additives and/or auxiliary substances; and optionally one or more further pharmacologically active ingredient. Examples of suitable physiologically acceptable carriers, additives and/or auxiliary substances are fillers, solvents, diluents, colorings and/or binders. These substances are known to the person skilled in the art (see H. P. Fiedler, Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzende Gebiete, Editio Cantor Aulendorf).

[0089] The pharmaceutical dosage form according to the present invention is preferably for systemic, topical or local administration, preferably for oral administration. Therefore, the pharmaceutical dosage form can be in form of a liquid, semisolid or solid, e.g. in the form of injection solutions, drops, juices, syrups, sprays, suspensions, tablets, patches, films, capsules, plasters, suppositories, ointments, creams, lotions, gels, emulsions, aerosols or in multiparticulate form, for example in the form of pellets or granules, if appropriate pressed into tablets, decanted in capsules or suspended in a liquid, and can also be administered as such.

[0090] The pharmaceutical dosage form according to the present invention is preferably prepared with the aid of conventional means, devices, methods and processes known in the art. The amount of the compound according to the present invention to be administered to the patient may vary and is e.g. dependent on the patient's weight or age and also on the type of administration, the indication and the severity of the disorder. Preferably 0.001 to 100 mg/kg, more preferably 0.05 to 75 mg/kg, most preferably 0.05 to 50 mg of a compound according to the present invention are administered per kg of the patient's body weight.

[0091] The glucocorticoid receptor is believed to have potential to modify a variety of diseases or disorders in mammals such as humans. These include in particular inflammatory diseases.

[0092] Another aspect of the present invention relates to a compound according to the present invention for use in

the treatment and/or prophylaxis of pain and/or inflammation; more preferably inflammatory pain.

[0093] A further aspect of the present invention relates to a method of treatment of pain and/or inflammation; more preferably inflammatory pain.


Examples

[0094] The following abbreviations are used in the descriptions of the experiments:

AcOH = acetic acid; Ac = acetyl group; Attaphos = bis(di-tert-butyl(4 dimethylaminophenyl)phosphine)dichloropalladium(II); Ar = argon; BISPIN (or Bis-Pin) = bis(pinacolato)diborane; dba = dibenzylideneacetone; DAST = (diethylamino)sulfur trifluoride; DCM = DCM; DIPEA = N,N-diisopropylethylamine; DIBAL-H = diisobutylaluminium hydride; DMA = dimethylacetamide; DMADMF = N,N-dimethylformamide dimethylacetal; DMAP = 4-(dimethylamino)-pyridine; DMF = N,N-dimethylformamide; DMSO = dimethylsulfoxide; dppe = 1,1'-bis(diphenylphosphanyl)ferrocene; EtOAc = EtOAc; EtOH = EtOH; h = hour; LAH = lithium aluminium hydride; LDA = lithiumdiisopropylamide; LiHMDS = lithium bis(trimethylsilyl)amide; m_c = multiplet centered; MeOH = MeOH; min = minute; MTBE = methyl tert-butyl ether; n-BuLi = n-butyllithium; RT = room temperature; Rt = retention time; tert = tertiary; TEA = triethylamine; THF = tetrahydrofuran; p-TSA = para-toluene sulfonic acid; TMSCl = trimethylsilyl chloride; X-Phos = 2-Dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl; XANTphos = 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene.

Synthesis of 6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (Intermediate A-1)

[0095]




[0096] Step1: To a stirring solution of 2-bromo-4-fluoro-6-nitrotoluene (4.69 g, 20 mmol, 1eq) in 1,4-dioxane (25 ml) was slowly added N,N-dimethylformamide dimethylacetal (13.3 mL, 100 mmol, 5eq) and pyrrolidine (1.47 mL, 20 mmol, 1eq). The reaction mixture was then stirred for 18 h at 100°C. The reaction mixture was concentrated to a dark residue. To this residue were added AcOH (30 mL) and iron powder (11 g, 200mmol, 10 eq) and then the reaction mixture was refluxed for 1 h. The reaction mixture was then cooled to RT and then filtered through a celite bed. The filtrate was neutralised by 50% sodium hydroxide solution and then extracted with EtOAc (2 × 100 mL). Combined organic layers was washed with water (100 mL), brine (100 mL), dried over anhydrous Na₂SO₄ and evaporated to get the crude which was purified by column chromatography to afford 4-bromo-6-fluoro-1H-indole (1.3 g, 30%) as brown liquid.

[0097] Step2: To a stirring suspension of 4-bromo-6-fluoro-1H-indole (1.1 g, 5.1 mmol, 1 eq), bis(pinacolato)diborane (2.6 g, 10.2 mmol, 2eq) and potassium acetate (2.0 g, 20.4 mmol, 4 eq) in 1,4-dioxane (20 mL) was deoxygenated by Ar for 10 min. Pd₂(dba)₃ (0.07 g, 0.07 mmol, 0.015 eq) and tricyclohexylphosphine (0.102 g, 0.36 mmol, 0.07 eq) was then added to the reaction mixture and again deoxygenated by Ar for 10 min. The reaction mixture was then stirred for 14 h at 110°C. The reaction mixture then cooled to RT and then filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude material which was purified by column chromatography to afford 6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (1.1 g, 82%) as light yellow solid.

Synthesis of 6-fluoro-1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (Intermediate A-2)

[0098]




[0099] Step 1: To a stirring solution of 4-bromo-6-fluoro-1H-indole (0.18 g, 0.841 mmol, 1 eq) in DMF (5 mL) was portion wise added sodium hydride (60%, 0.07 g, 1.68 mmol, 2 eq) at 0°C. The reaction mixture was then stirred for 30 min at RT. Methanesulfonylchloride (0.114 mL, 1.26 mmol, 1.5 eq) then added to the reaction mixture at 0°C. The reaction mixture then stirred for 2 h at RT. Reaction mixture was diluted with EtOAc (50 mL). Combined organic layers was washed with water (5 × 10 mL), brine (10 mL), dried over anhydrous Na₂SO₄ and evaporated under reduced pressure. Crude product was purified by column chromatography to afford 4-bromo-6-fluoro-1-(methylsulfonyl)-1H-indole (0.1 g, 41%) as off-white solid.

[0100] Step2: To a stirring suspension of 4-bromo-6-fluoro-1-(methylsulfonyl)-1H-indole (1.2 g, 3.53 mmol, 1 eq), bis-pinacolatodiborane (1.79 g, 7.06 mmol, 2 eq) and potassium acetate (1.39 g, 10.62 mmol, 4eq) in 1,4-dioxan (20 mL) was deoxygenated by Ar for 10 min. Pd₂(dba)₃ (0.048 g, 0.052 mmol, 0.015 eq) and tricyclohexylphosphine (0.071 g, 0.25 mmol, 0.07 eq) was then added to the reaction mixture and again deoxygenated by Ar for 10 min. The reaction mixture then stirred for 14 h at 110°C. The reaction mixture then cooled to RT and then filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude product which was purified by column chromatography to afford 6-fluoro-1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (1.0 g, 80%) as light yellow solid.

Synthesis of 1-(ethylsulfonyl)-6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**Intermediate A-8**)

[0101]




[0102] Step1: To a stirring solution of 4-bromo-6-fluoro-1H-indole (1.0 g, 4.67 mmol, 1 eq) in DMF (24 mL) was portion wise added sodium hydride (60%, 0.224 g, 9.34 mmol, 2 eq) at 0°C. The reaction mixture was then stirred for 30 min at RT. Ethanesulfonylchloride (0.604 mL, 7 mmol, 1.5 eq) then added to the reaction mixture at 0°C. The reaction mixture then stirred for 2 h at RT. Reaction mixture was diluted with EtOAc (200 mL). Combined organic layers was washed with water (5 × 30 mL), brine (30 mL), dried over anhydrous Na₂SO₄ and evaporated under reduced pressure. Crude product was purified by column chromatography to afford 4-bromo-1-(ethylsulfonyl)-6-fluoro-1H-indole (0.513 g, 36%) as off white solid.

[0103] Step2: To a stirring suspension of 4-bromo-1-(ethylsulfonyl)-6-fluoro-1H-indole (0.51 g, 1.6 mmol, 1 eq), bis(pinacolato)diborane (0.843 g, 3.2 mmol, 2 eq) and potassium acetate (0.653 g, 6.4 mmol, 4 eq) in 1,4-dioxan (15 mL) was deoxygenated by Ar for 10 min. Pd₂(dba)₃ (0.023 g, 0.025 mmol, 0.015 eq) and tricyclohexylphosphine (0.036 g, 0.12 mmol, 0.072 eq) was then added to the reaction mixture and again deoxygenated by Ar for 10 min. The reaction mixture was then stirred for 14 h at 110°C. The reaction mixture cooled to RT and then filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude material which was purified by column chromatography to afford 1-(ethylsulfonyl)-6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (0.212 g, 38%) as off white solid.

Synthesis of 1-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indol-1-yl)ethanone (**Intermediate A-10**)


[0104]



[0105] Step1: To a stirred solution of 4-bromo-1H-indole (0.5 g, 2.55 mmol, 1 eq) in THF (25 mL) was added sodium hydride (60%) (0.122 g, 3.06 mmol, 1.2eq) at 0°C and continued stirred at RT for 30 min. Acetyl chloride (0.02 mL, 3.06 mmol, 1.2 eq) was then added to the reaction mixture and again stirred for another 2 h. The reaction mixture was quenched with water and extracted with EtOAc (2 x 100 mL). Combined organic layers were washed with water (100 mL), brine (100 mL), dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product which was purified by column chromatography to afford 1-(4-bromo-1H-indol-1-yl)ethanone (0.55 g, 91%) as brown liquid. Step2: To a stirred solution of 1-(4-bromo-1H-indol-1-yl)ethanone (0.55 g, 2.31 mmol, 1 eq), bis(pinacolato)diborane (0.707 g, 4.62 mmol, 2 eq) and potassium acetate (0.680 g, 6.93 mmol, 3 eq) in 1,4-dioxan (20 mL) was deoxygenated by Ar for 10 min. Pd₂(dba)₃ (0.106g, 0.1155 mmol, 0.08 eq) and Cy₃P (0.052 g, 0.1848 mmol, 0.08 eq) was then added to the reaction mixture and reflux at 90°C for another 16 h. The reaction mixture was cooled to RT and filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude material which was purified by column chromatography to afford 1-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indol-1-yl)ethanone (0.600 g, 92%) as brown liquid.

Synthesis of 6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-(2,2,2-trifluoroethyl)-1H-indole (**Intermediate A-11**)

[0106]




[0107] Step1: To a solution of 4-bromo-6-fluoro-1H-indole (2.0 g, 9.345 mmol, 1 eq.) in DMF (25 mL) was added Cs₂CO₃ (15.18 g, 46.72 mmol, 5 eq.) and 1,1,1-trifluoro-2-iodoethane (5.8 g, 28.037 mmol, 3.0 eq) in a sealed tube. The reaction mixture was refluxed at 50°C for 24 h. The reaction mixture was filtered through sintered and the filtrate was diluted with EtOAc (100 mL). Organic layer was washed with cold water (3 x 50 mL), brine (25 mL), dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product, which was purified by flash column chromatography to afford mixture which further purified by Prep HPLC to afford 4-bromo-6-fluoro-1-(2,2,2-trifluoroethyl)-1H-indole (0.400 g, 14%) as off white solid.

[0108] Step2: To a stirred solution of 4-bromo-6-fluoro-1-(2,2,2-trifluoroethyl)-1H-indole (0.450 g, 1.52 mmol, 1eq), bis(pinacolato)diborane (0.461 g, 1.824 mmol, 1.2eq) and potassium acetate (0.446 g, 4.56 mmol, 3 eq) in 1,4-dioxan (20 mL) was deoxygenated by Ar for 10 min. PdCl₂(dppf)-DCM (0.124 g, 0.152 mmol, 0.1 eq) was then added to the reaction mixture and stirred at 90°C for another 16 h. The reaction mixture was filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude material which was used in next step without further purification.

Synthesis of 1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**Intermediate A-12**)

[0109]




[0110] Step1: To a stirring solution of 4-bromo-1H-indole (1.0 g, 5.1mmol, 1 eq) in DMF (20ml) was portion wise added sodium hydride (60%, 0.245 g, 10.2 mmol, 2 eq) at 0°C. The reaction mixture was then stirred for 30 min at RT. Methanesulfonylchloride (0.584 ml, 7.6 mmol, 1.5 eq) then added to the reaction mixture at 0°C. The reaction mixture was stirred for 2 h at RT. Reaction mixture was diluted with EtOAc (100 mL). Combined organic layers was washed with water (5 x 20 mL), brine (20 mL), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure. The crude product was purified by column chromatography to afford 4-bromo-1-(methylsulfonyl)-1H-indole (0.532 g, 38%) as off white solid.

[0111] Step2: To a stirring suspension of 1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (0.36g, 1.31mmol, 1eq), bis(pinacolato)diborane (0.66 g, 2.62 mmol, 2eq) and potassium acetate (0.57 g, 5.25 mmol, 4 eq) in 1,4-dioxan (10 LI) was deoxygenated by Ar for 10 min. Pd₂(dba)₃ (0.018g, 0.019mmol, 0.015 eq) and tricyclohexylphosphine (0.027 g, 0.094 mmol, 0.072 eq) was then added to the reaction mixture and again deoxygenated by Ar for 10 min. The reaction mixture was then stirred for 14 h at 110°C. The reaction mixture then cooled to RT and then filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude material which was purified by column chromatography to afford 1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (0.31 g, 73%) as off white solid.

Synthesis of 5-fluoro-3-methyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**Intermediate A-15**)


[0112]



[0113] Step1: To a solution of 2-bromo-4-fluoro-1-nitrobenzene (0.5 g, 2.27 mmol, 1 eq.) in THF (20 mL) was added (E)-prop-1-en-1-ylmagnesium bromide (0.5 M in THF) (13.6 mL, 6.818 mmol, 3 eq) at -60°C under nitrogen atmosphere. Then the reaction mixture was stirred at the same temperature for 4 h. The reaction was quenched with saturated ammonium chloride solution at -60°C. Then the resulting mixture was extracted with EtOAc (2 x 100 mL), washed with brine solution and concentrated under reduced pressure to give the crude product which was purified by flash column chromatography to afford 7-bromo-5-fluoro-3-methyl-1H-indole (0.3 g, 58%) as dense yellow liquid. **Step2:** To a solution of 7-bromo-5-fluoro-3-methyl-1H-indole (0.8 g, 3.669 mmol, 1 eq) in 1,4-dioxane (15.0 mL) were added KOAc (1.43 g, 14.67 mmol, 4 eq) and bispinacolatodiborane (1.12 g, 7.33 mmol, 2 eq). The solution was degassed with Ar for 20 min followed by addition of Pd₂(dba)₃ (0.16 g, 0.183 mmol, 0.05 eq) and Cy₃P (0.082 g, 0.293 mmol, 0.08 eq). The reaction mixture was refluxed for 16 h. After completion of reaction (monitored by TLC), solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography to afford 5-fluoro-3-methyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (0.7 g, 70%), as brown solid.

Synthesis of 3-cyclopropyl-5-fluoro-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (Intermediate A-16)

[0114]



[0115] Step1: To a stirring solution of 7-bromo-5-fluoroindole (7.0 g, 32.7 mmol, 1 eq) in DMF (175 mL) was added powdered potassium hydroxide (4.56 g, 81.77 mmol, 2.5 eq). The reaction mixture was then stirred for 30 min at RT. Iodine (12.46 g, 49.06 mmol, 1.5 eq) was then added to the reaction mixture and finally stirred for 2 h at RT. The reaction mixture was diluted with EtOAc (1000 mL) and washed with water (5 x 100 mL) followed brine (100 mL). The organic layer was dried over anhydrous Na_2SO_4 and the solvent was evaporated to get the crude product, which was purified by column chromatography to afford 7-bromo-5-fluoro-3-iodo-1H-indole (6.2, 56 %) as brown solid. **Step2:** To a stirring solution of 7-bromo-5-fluoro-3-iodo-1H-indole (6.2 g, 18.23 mmol, 1 eq) in THF (109 mL) was added drop wise LiHMDS (1M) (91.15 mL, 91.15 mmol, 5 eq) at -78°C under inert atmosphere. The reaction mixture was stirred for 30min at same condition. MOMCl (5.83 g, 72.94 mmol, 4 eq) was then added to the reaction mixture at -78°C . The reaction mixture was allowed to warm up to RT and then stirred for 16 h. The reaction mixture was quenched by addition of saturated solution of ammonium chloride (100 mL). Organic layer was separated and the aqueous layer was extracted with EtOAc (100 mL). The combined organic layers were washed with brine (100 mL). The organic layer was dried over anhydrous Na_2SO_4 and the solvent was evaporated to get the crude product, which was purified by column chromatography to afford 7-bromo-5-fluoro-3-iodo-1-(methoxymethyl)-1H-indole (5.4 g, 57%) as off white solid.

[0116] Step3: To a stirred suspension of 7-bromo-5-fluoro-3-iodo-1-(methoxymethyl)-1H-indole (2.7 g, 7.03 mmol, 1 eq), cyclopropylboronic acid (1.84 g, 2.03 mmol, 3 eq) and K_3PO_4 (4.5 g, 21.05 mmol, 3 eq) in 1,4-dioxan (45 mL) was deoxygenated by Ar for 10 min. $\text{Pd}(\text{OAc})_2$ (0.08 g, 0.3525 mmol, 0.05 eq) and xantphos (0.407 g, 0.713 mmol, 0.1 eq) were then added to the reaction mixture and again deoxygenated for 10 min. Finally the reaction mixture was stirred at 100°C for 16 h. The reaction mixture was cooled to RT and then filtered through celit bed. The filtrate was concentrated under reduced pressure to get the crude material which was purified by column chromatography to afford 7-bromo-3-cyclopropyl-5-fluoro-1-(methoxymethyl)-1H-indole (0.65 g, 31%) as off white solid.


[0117] Step4: To a stirring solution of 7-bromo-3-cyclopropyl-5-fluoro-1-(methoxymethyl)-1H-indole (1.25 g, 4.19 mmol, 1 eq) in mixture of MeOH and water (3:1) (66 mL) was added oxalic acid (1.13 g, 12.58 mmol, 3 eq). The reaction mixture was then stirred at 90°C for 18 h. The reaction mixture was cooled to RT and concentrated under reduced pressure to get the residue. The residue was diluted with EtOAc (100 mL) and washed with water (2 x 40mL) and brine (40 mL). The organic layer was dried over anhydrous Na_2SO_4 and the solvent was evaporated to get the crude product, which was purified by column chromatography to afford 7-bromo-3-cyclopropyl-5-fluoro-1H-indole (0.57g, 54%) as color less liquid.

[0118] Step5: To a stirring suspension of 7-bromo-3-cyclopropyl-5-fluoro-1H-indole (0.57 g, 2.24 mmol, 1 eq), bis-pinacolatodiboron (1.7 g, 6.73 mmol, 3 eq) and potassium acetate (0.66 g, 6.73 mmol, 3 eq) in 1,4-dioxan (20 mL) was deoxygenated by Ar for 10 min. $\text{Pd}_2(\text{dba})_3$ (0.031g, 0.033 mmol, 0.015 eq) and tricyclohexylphosphine (0.047 g, 0.168 mmol, 0.075 eq) was then added to the reaction mixture and again deoxygenated by Ar for 10 min. The reaction mixture was then stirred for 14 h at 110°C . The reaction mixture was then cooled to RT and then filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude material which was purified by column chromatography

to afford 3-cyclopropyl-5-fluoro-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (0.35 g, 52%) as off white solid.

Synthesis of 2-(6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indol-1-yl)ethan-1-ol (**Intermediate A-20**)

[0119]




[0120] Step1: To a solution of 4-bromo-6-fluoro-1H-indole (0.5 g, 2.34 mmol, 1 eq.) in DMF (5 mL) was added sodium hydride (0.130 g, 2.80 mmol, 1.2 eq) at 0°C. The solution was stirred at RT for 30 min followed by addition of (2-bromoethoxy)(tert-butyl)dimethylsilane (1.17g, 4.67 mmol, 2.0 eq) and reaction mixture was stirred at RT for 2 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (20 mL) and organic layer was washed with cold water (5 x 10 mL), brine (10 mL), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure. Crude product was purified by column chromatography to afford 4-bromo-1-(2-((tert-butyldimethylsilyl)oxy)ethyl)-6-fluoro-1H-indole (0.85g, 98%) as brown liquid having (2-bromoethoxy)(tert-butyl)dimethylsilane as impurity.

[0121] Step2: To a stirred solution of 4-bromo-1-(2-((tert-butyldimethylsilyl)oxy)ethyl)-6-fluoro-1H-indole (1.3 g, 3.49 mmol, 1 eq.) in THF (15 mL) was added TBAF (3.49 mL) (1M) at RT and the mixture was stirred for 16 h. After completion of reaction (monitored by LCMS & TLC), reaction mixture was diluted with EtOAc (20 mL) and organic layer was washed with cold water (5 x 10 mL), brine (10 mL), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure. Crude product was purified by column chromatography to afford 2-(4-bromo-6-fluoro-1H-indol-1-yl)ethan-1-ol (0.55 g, 61%) as brown liquid.

[0122] Step3: To a stirred solution of 2-(4-bromo-6-fluoro-1H-indol-1-yl)ethan-1-ol (0.55 g, 2.13 mmol, 1 eq), bis(pinacolato)diborane (0.647 g, 2.55 mmol, 1.2 eq) and potassium acetate (0.626 g, 6.393 mmol, 3 eq) in 1,4-dioxane (20 mL) was deoxygenated by Ar for 10 min. PdCl₂(dppf)-DCM (0.173 g, 0.213 mmol, 0.1 eq) was then added to the reaction mixture and the mixture was stirred at 90°C for 16 h. After completion of reaction (monitored by TLC), reaction mixture was filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude 2-(6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indol-1-yl)ethan-1-ol which was used in next step without further purification.

Synthesis of 2-(6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazol-1-yl)ethan-1-ol (**Intermediate A-21**)

[0123]




[0124] Step1: To a stirred solution of 4-bromo-6-fluoro-1H-indazole (0.2 g, 0.93 mmol, 1 eq) in DMF (5 mL) was added K₂CO₃ (0.38 g, 2.79 mmol, 3.0 eq) at RT and the mixture was stirred for 20 min. Then bromoethan-1-ol (0.07 mL, 0.93 mmol, 1 eq) was added and the mixture was stirred for 16h at 50°C. After completion of the reaction (monitored by TLC),

the reaction mass quenched with ice cold water and extracted with EtOAc (3 x 20 mL), washed with H₂O (3 x 20 mL), brine (25 mL), dried over Na₂SO₄ and concentrated. The crude product was purified by column chromatography to afford 2-(4-bromo-6-fluoro-1H-indazol-1-yl)ethan-1-ol (0.12 g, 50%) as white solid.

[0125] Step2: To a solution of 2-(4-bromo-6-fluoro-1H-indazol-1-yl)ethan-1-ol (0.9 g, 3.473 mmol, 1 eq) in 1,4-dioxane (60.0 mL) were added KOAc (1.02 g, 10.419 mmol, 3 eq) and bispincolatediborane (1.76 g, 6.947 mmol, 2.0 eq). The solution was degassed with Ar for 20 min followed by addition of Pd₂(dba)₃ (0.17 g, 0.173mmol, 0.05 eq) and Cy₃P (0.077 g, 0.277 mmol, 0.08 eq). The reaction mixture was refluxed for 16 h. After completion of reaction (monitored by TLC), solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography to afford 2-(6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazol-1-yl)ethan-1-ol (0.95 g, 89%) as brown solid.

Synthesis of 1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole (**Intermediate A-22**)

[0126]




[0127] Step 1: To a stirring solution of 4-bromo-1H-indazole (1.0 g, 5.07 mmol, 1 eq) in DMF (25ml) was portion wise added sodium hydride (60%, 0.406 g, 10.152mmol, 2 eq) at 0°C. The reaction mixture was stirred for 30 min at RT. Methanesulfonylchloride (0.59 mL, 7.6 mmol, 1.5 eq) was added to the reaction mixture at 0°C. The reaction mixture was stirred for 2 h at RT. Reaction mixture was diluted with EtOAc (150 mL). Combined organic layers were washed with water (5 x 30 mL), brine (30 mL), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure. Crude product was purified by column chromatography (230-400 mesh silica gel 10% EtOAc/hexane; R_F-value-0.5) to afford 4-bromo-1-(methylsulfonyl)-1H-indazole (0.95 g, 69%) as light yellow solid.

[0128] Step 2: To a stirring suspension of 4-bromo-1-(methylsulfonyl)-1H-indazole (0.95, 3.45 mmol, 1 eq), bis(pinacolato)diborane (1.75 g, 6.91 mmol, 2eq) and potassium acetate (1.01 g, 10.36 mmol, 3 eq) in 1,4-dioxane (35 mL) was deoxygenated by Ar for 10 min. Pd(dppf)Cl₂·DCM (0.141g, 0.1727mmol, 0.05 eq) was added to the reaction mixture and again deoxygenated by Ar for 10 min. The reaction mixture was stirred for 14 h at 110°C. The reaction mixture was cooled to RT and then filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude material which was purified by column chromatography (230-400 mesh silica gel, 10% EtOAc/hexane; R_F-value-0.45) to afford 1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole (0.9 g, 85.4%) as off white solid.

Synthesis of 6-fluoro-1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole (**Intermediate A-23**)

[0129]




[0130] Step 1: To a stirring solution of 4-bromo-6-fluoro-1H-indazole (1.2 g, 5.58 mmol, 1 eq) in DMF (30 mL) was

portion wise added sodium hydride (60%, 0.446 g, 11.16mmol, 2 eq) at 0°C. The reaction mixture was then stirred for 30 min at RT. Methanesulfonylchloride (0.65 ml, 8.37 mmol, 1.5 eq) was added to the reaction mixture at 0°C. The reaction mixture was stirred for 2 h at RT. Reaction mixture was diluted with EtOAc (150 mL). Combined organic layers were washed with water (5 x 30 mL), brine (30 mL), dried over anhydrous Na₂SO₄ and evaporated under reduced pressure. Crude product was purified by column chromatography (230-400 mesh silica gel 10% EtOAc/hexane; R_F-value-0.5) to afford 4-bromo-6-fluoro-1-(methylsulfonyl)-1H-indazole (1.3 g, 80%) as light yellow solid.

[0131] Step 2: To a stirring suspension of 4-bromo-6-fluoro-1-(methylsulfonyl)-1H-indazole (1.3, 4.43 mmol, 1eq), bis(pinacolato)diborane (2.25 g, 8.87 mmol, 2 eq) and potassium acetate (1.3 g, 13.3 mmol, 3 eq) in 1,4-dioxane (45 mL) was deoxygenated by Ar for 10 min. Pd(dppf)Cl₂-DCM (0.18 g, 0.22 mmol, 0.05 eq) and was then added to the reaction mixture and again deoxygenated by Ar for 10 min. The reaction mixture was stirred for 14 h at 110°C. The reaction mixture was cooled to RT and then filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude material which was purified by column chromatography (230-400 mesh silica gel, 10% EtOAc/hexane; R_F-value-0.45) to afford 6-fluoro-1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole (1.1 g, 73%) as off white solid.

Synthesis of 6-fluoro-1-(2-methoxyethyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**Intermediate A-24**)


[0132]



[0133] Step1: To a stirring solution of 4-bromo-6-fluoro-1H-indole (0.5 g, 2.34 mmol, 1 eq) in DMF (5 mL) was portion wise added sodium hydride (0.112 g, 2.8 mmol, 1.2 eq.) at 0°C. The reaction mixture was then stirred for 30 min at RT. 1-Bromo-2-methoxyethane (0.812 mL, 5.84 mmol, 2.5 eq) was then added to the reaction mixture at 0°C. The reaction mixture was stirred for 2 h at RT. Reaction mixture was diluted with EtOAc (50 mL). Combined organic layers were washed with water (5x10 mL), brine (10 mL), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure. Crude product was purified by column chromatography (230-400 mesh silica gel 20% EtOAc/hexane; R_F-value-0.6) to afford 4-bromo-6-fluoro-1-(2-methoxyethyl)-1H-indole (0.63 g, 99%) as brown gum. **Step2:** To a stirring suspension of 4-bromo-6-fluoro-1-(2-methoxyethyl)-1H-indole (0.8 g, 2.94 mmol, 1eq), bis-pinacolatodiborane(1.2 g, 4.4 mmol, 1.5eq.) and potassium acetate (0.865 g, 8.823 mmol, 3eq.) in 1,4-dioxan (20 mL) was deoxygenated by Ar for 10 min. PdCl₂(dppf)-DCM (0.239 g, 0.29 mmol, 0.01 eq.) was then added and the reaction mixture was stirred for 14 h at 90°C. The reaction mixture was cooled to RT and then filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel 5% EtOAc/hexane; R_F-value-0.6) to afford 6-fluoro-1-(2-methoxyethyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (0.93 g, 99%) as light brown gummy solid.

Synthesis of 1-(2,2-difluoroethyl)-6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**Intermediate A-25**)

[0134]




[0135] **Step1:** To a stirred solution of 4-bromo-6-fluoro-1H-indole (0.1 g, 0.469 mmol, 1 eq) in DMF (25 ml) was added Cs_2CO_3 (0.457 g, 1.407 mmol, 3 eq) followed by LiBr (86.84 g, 0.469 mmol, 1 eq) and the mixture was stirred at RT for 10 minutes. Then 2,2-difluoroethyl 4-methylbenzenesulfonate (0.133 g, 0.563 mmol, 1.2 eq) was added to the reaction mixture and heated at 80°C for 3 h (TLC). Reaction mixture was then diluted with water (10 mL) and EtOAc (15 mL).

The organic layer was washed with cold water (3x10 mL) brine (10 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to get crude product which was purified by column chromatography (100-200 mesh silica gel; TLC system: EtOAc/Hexane (3:7); R_f -value-0.5) to afford 4-bromo-1-(2,2-difluoroethyl)-6-fluoro-1H-indole (0.75 g, 58%).

[0136] **Step2:** A suspension of 4-bromo-1-(2,2-difluoroethyl)-6-fluoro-1H-indole (0.2 g, 0.722 mmol, 1 eq), bis(pinacolato)diboron (0.275 g, 1.083 mmol, 1.5 eq) and potassium acetate (0.212 g, 2.166 mmol, 3 eq) in 1,4-dioxane (10 mL) was deoxygenated well by Ar for 10 min. 1,1'-Bis(diphenylphosphino)ferrocene palladium(II)dichloride DCM complex (0.03 g, 0.0361 mmol, 0.05 eq) was then added to the reaction mixture and the reaction mixture heated at 100°C for 16 h (LCMS). The reaction mixture was then cooled to RT, filtered through celite pad and the filtrate was concentrated under reduced pressure to get the crude 1-(2,2-difluoroethyl)-6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole which was used in the next step without further purification (Yield ~ 49% in LCMS).

Synthesis of 1-cyclopropyl-6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**Intermediate A-27**)

[0137]




[0138] **Step1:** To a stirring solution of 4-bromo-6-fluoro-1H-indole (5.4g, 25.23mmol, 1 eq) in toluene (45 mL) were added cyclopropylboronic acid (4.33 g, 50.46 mmol, 2 eq), $\text{Cu}(\text{OAc})_2$ (0.46 g, 2.52 mmol, 0.1 eq), 2-(4,5-dihydro-1H-imidazol-2-yl)phenol (0.41 g, 2.52 mmol, 0.1 eq) and pyridine (6.0 g, 75.7 mmol, 3 eq). The reaction mixture was stirred for 24 h at 65°C. Solvent was removed under reduced pressure and azeotrope by MeOH twice. The residue was purified by column chromatography (230-400 mesh silica gel; 10% EtOAc/hexane; R_f -value-0.6) to afford 4-bromo-1-cyclopropyl-6-fluoro-1H-indole (0.85 g, 13%) as brown liquid.

[0139] **Step2:** To a stirring suspension of 4-bromo-1-cyclopropyl-6-fluoro-1H-indole (0.85 g, 3.35 mmol, 1 eq), bis(pinacolato)diborane (1.7 g, 6.7 mmol, 2 eq) and potassium acetate (1.31 g, 13.38 mmol, 4 eq) in 1,4-dioxane (20 mL) was deoxygenated by Ar for 10 min. $\text{Pd}_2(\text{dba})_3$ (0.046 g, 0.05 mmol, 0.015 eq) and ommtetricyclohexylphosphine (0.067 g, 0.24 mmol, 0.072 eq) was then added to the reaction mixture and again deoxygenated by Ar for 10 min. The reaction mixture was stirred for 14 h at 110°C. The reaction mixture was cooled to RT and then filtered through celite bed. Filtrate was concentrated under reduced pressure to get the crude material which was purified by column chromatography (230-400 mesh silica gel, 20% EtOAc/hexane; R_f -value-0.6) to afford 1-cyclopropyl-6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (0.61 g, 61%) as light yellow solid.

[0140] $^1\text{H-NMR}$ (400 MHz; $\text{DMSO}-d_6$, 20°C): δ 7.45 (dd, 1H), 7.35 (d, 1H), 7.15 (dd, 1H), 6.67 (d, 1H), 3.41 (m, 1H), 1.32 (12H), 1.03-1.08 (2H), 0.82-0.92 (2H).

Synthesis of 5-fluoro-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**Intermediate A-71**)

[0141]











[0142] A mixture of 7-bromo-5-fluoro-1H-indole (1 g, 4.7 mmol, 1 eq), bis(pinacolato)diborane (2.02 g, 7.9 mmol, 1.7 eq), potassium acetate (917 mg, 9.4 mmol, 2 eq), 1,1'-Bis(diphenylphosphino)ferrocene palladium(II)dichloride dichloromethane complex (382 mg, 0.467 mmol, 0.1 eq) in 1,4-dioxane (13 mL) was degassed with nitrogen, and the reaction mixture was stirred at 60°C. After completion of the reaction (monitored by LCMS), a sat. sodiumbicarbonate solution was added to the reaction mixture, which was then extracted with EtOAc (2 x). The combined organic layers were washed with brine, dried over anhydrous MgSO_4 and evaporated under reduced pressure. The crude product was purified by column chromatography (silica gel; EtOAc/cyclo-Hexane as eluent) to afford 5-fluoro-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (940 mg, 77%).

[0143] The intermediates in Table 1 were synthesized in analogy to Intermediate A-1 to Intermediate A-27.

Intermediate	Synthesized in analogy to	Structure
Int-A-30	Int-A-2	
Int-A-42	Int-A-11	
Int-A-43	Int-A-11	
Int-A-44	Int-A-27	
Int-A-45	Int-A-2	
Int-A-46	Int-A-11	

(continued)

Intermediate	Synthesized in analogy to	Structure
Int-A-47	Int-A-2	
Int-A-48	Int-A-2	
Int-A-54	Int-A-22	
Int-A-63	Int-A-22	
Int-A-59	Int-A-21	
Int-A-66	Int-A-22	
Int-A-68	Int-A-22	
Int-A-72	Int-A-22	









(continued)

Intermediate	Synthesized in analogy to	Structure
Int-A-73	Int-A-21	
Int-A-75	Int-A-71	
Int-A-76	Int-A-22	
Int-A-77	Int-A-22	










[0144] The Intermediates in Table 2 are commercially available:

Name	Structure
4-(tetramethyl-1,3,2-dioxaborolan-2-yl)-6-(trifluoromethyl)-1H-indole	
5-fluoro-1H-indole-4-boronic acid pinacol ester	
2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2H-indazole	









(continued)

Name	Structure
1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	
4-(tetramethyl-1,3,2-dioxaborolan-2-yl)-2-(trifluoromethyl)-1H-indole	
4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	
tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole-1-carboxylate	
indole-4-boronic acid	
1,1-dimethylethyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole-1-carboxylate	
3-methyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole	
(7-chloro-1H-indazol-4-yl)boronic acid	
7-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	




(continued)

Name	Structure
7-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	
4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7-(trifluoromethyl)-1H-indazole	
(6-(trifluoromethyl)-1H-indazol-4-yl)boronic acid	
(6-fluoro-1H-indazol-4-yl)boronic acid	
7-chloro-6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	
7-fluoro-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-indole	
7-methyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-indole	
6-methoxy-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-indole	
(1H-indol-7-yl)boronic acid	

(continued)


Name	Structure
3-Methyl-1H-indazole-7-boronic acid	
4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole-6-carbonitrile	
7-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	
7-fluoro-4-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole	
6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	
6-chloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole	
7-chloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole	
7-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole	

(continued)

Name	Structure
6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	
6-chloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole	
4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-6-(trifluoromethyl)-1H-indazole	

Synthesis of 8-bromo-6-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**Intermediate B-7**):

[0145]



[0146] Step1: To a stirred solution of 5-fluoro-2-methyl-phenylamine (100 g, 0.8 mol) in DCM (1500 mL) was added pyridine (129 mL, 1.6 mol) followed by DMAP (1 g, 0.008 mol) at RT. To this reaction mixture was added drop wise pivoyl chloride (109 mL, 0.88 mol) at 0°C. Resulting reaction mixture was stirred at 0°C for 2 h. After completion of starting material reaction mixture was poured into ice cooled 1(N) HCl solution (1500 mL) and stirred for 30 min. The two layers were separated and the organic layer was washed with 1(N) HCl solution (1000 mL) followed by saturated NaHCO₃ solution (1000 mL) followed by brine (1000 mL) and then dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude product. Crude product was triturated using hexane to afford N-(5-fluoro-2-methyl-phenyl)-2,2-dimethyl-propionamide (158 g, 95%) as white solid.

[0147] Step2: To a stirring solution N-(5-fluoro-2-methyl-phenyl)-2,2-dimethyl-propionamide (90 g, 0.43 mol) in toluene (2000 mL) was added para-toluene sulphonic acid monohydrate (82.0 g, 0.43 mol), Pd(OAc)₂ (9.66 g, 0.043 mol)

followed by N-bromo succinimide (84.5 g, 0.473 mol) at RT. Resulting reaction mixture was stirred at RT for 48 h. Reaction monitoring TLC in 20% EA-Hex, showed formation of desired product ($R_f=0.4$) along with the -60% unreacted SM ($R_f=0.5$). Reaction mixture was then concentrated, obtained residue was diluted with EtOAc and washed with water (twice). Total organic layer was dried over anhydrous Na_2SO_4 and concentrated to afford crude product. Two combined

batches of crude product (obtained from 90g and 100g reaction of N-(5-fluoro-2-methyl-phenyl)-2,2-dimethyl-propionamide) were purified by flash chromatography to afford N-(2-bromo-3-fluoro-6-methyl-phenyl)-2,2-dimethyl-propionamide (60 g, 23%) as white solid and recovered un-reacted N-(5-fluoro-2-methyl-phenyl)-2,2-dimethyl-propionamide (125 g). **[0148]** Step3: To a pre cooled solution of 70% H_2SO_4 in water (430 mL) was added N-(2-bromo-3-fluoro-6-methyl-phenyl)-2,2-dimethyl-propionamide (60 g, 0.208 mol) portion wise at 0°C . Resulting reaction mixture was stirred at RT for 10 min and then heated to 100°C for 2 h. After completion of starting material (monitored by TLC in 10% EA-Hex, $R_f=0.8$) reaction mixture was cooled to 0°C and pH adjusted to ~ 10 with 10% NaOH solution. Resulting basic aqueous was extracted with EtOAc (3 x 500 mL). Total organic layers were washed with water followed by brine, dried over anhydrous Na_2SO_4 and concentrated to afford crude product. Crude product was purified by column chromatography to afford 2-bromo-3-fluoro-6-methyl-phenylamine (36 g, 85%) as pale yellow liquid.

[0149] Step4: To the stirred suspension of 2-bromo-3-fluoro-6-methyl-phenylamine (36 g, 0.1764 mol) in dry DMSO (540 mL) was added 2-amino-2-methyl-propionic acid (16.9 g, 0.164 mol) followed by K_3PO_4 (75 g, 0.353 mol) at RT. Resulting reaction mixture was degassed with nitrogen for 30 min, then CuCl (1.75 g, 0.0176 mol) was added and reaction mixture was heated at 140°C for 5 h. After completion of the starting material (monitored by TLC, 20% EA-Hexane, $R_f 0.4$) reaction mixture was cooled to RT and filtered through celite. Celite bed was washed with EtOAc (500 mL). Resulting filtrate was poured into ice cold water (2500 mL). Resulting aqueous layer was extracted with EtOAc (2 x 750 mL). Total organic part was washed with water (2 x 750 mL), followed by brine (500 mL), dried over anhydrous Na_2SO_4 and evaporated under reduced pressure to afford crude product. Obtained crude product was triturated with hexane and filtered and dried to afford 5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxalin-2-one (19 g, 52%) as brown solid.

[0150] Step5: To a solution of 5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxalin-2-one (19 g, 91.2 mmol) in toluene (380 mL) was added Lawesson's reagent (55.5 g, 137 mmol) at RT and the reaction mixture was refluxed at 120°C for 1 h. After completion of starting material (monitored by TLC in 20% EA-Hexane, $R_f 0.7$), the reaction mass was cooled to RT and quenched with sat. NaHCO_3 solution (250 mL) and resulting aqueous layer was extracted with EtOAc (3 x 250 mL). Combined organic layers were washed with water (250 mL), followed by brine (250 mL), dried over anhydrous Na_2SO_4 and the solvent was evaporated to afford crude product. Obtained crude product was purified by column chromatography to afford 5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxaline-2-thione (19 g, 93%) as yellow solid.


[0151] Step6: To a stirred solution of 5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxaline-2-thione (20.5 g, 91.5 mmol) in THF (512 mL) was added drop wise hydrazine hydrate (13.5 mL, 274.5 mmol) at 0°C . The reaction mixture was stirred at RT for 16 h. TEA (64 mL, 457.5 mmol) followed acetyl chloride (21.5 mL, 274.5 mmol) were added to the reaction mixture drop-wise at 0°C and stirred for 2 h at RT. After completion of starting material (monitored by LCMS) reaction mixture diluted with water (500 mL) and extracted by 10% MeOH-DCM (5 x 500 mL). The total organic part was washed by brine (250 mL), dried over Na_2SO_4 and concentrated under reduced pressure to afford acetic acid (5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxalin-2-ylidene)-hydrazide (21 g, 87%, crude) as pale yellow solid.

[0152] Step7: Acetic acid (5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxalin-2-ylidene)-hydrazide (10.5 g, 0.04 mol) was taken in round bottom flask (100 mL) and then cooled to -10°C . Then phosphorus oxychloride (18.5 mL, 0.2 mol) was added drop wise followed by drop wise addition of TEA (5.6 mL, 0.04 mol). After that the reaction mixture was stirred at -10°C for 10 min and then 10 min at RT and finally at reflux condition for 1 h. After completion of starting material (monitored by LCMS) reaction mixture cooled to 0°C and quenched with crushed ice water (100 mL). The aqueous part was then basified using cold aqueous ammonia solution (100 mL) drop-wise. Resulting basic aqueous was then extracted with EtOAc (3 x 150 mL). Total organic part was washed with brine (150 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford crude compound. Crude product co-distillation with MTBE twice, then trituration with hexane and dried to afford 6-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (6 g, 61%) as pale yellow solid.

[0153] Step8: To a solution of 6-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (12 g, 0.048 mol) in DMF (360 mL) was added NBS (9.39 g, 0.0528 mol) portion wise at -10°C . Resulting reaction mixture was stirred at RT for 4 h. After completion of starting material (monitored by LCMS), reaction mixture was diluted with water (500 mL) and extracted with EtOAc (2 x 500 mL). Combined organic layers were washed with water (750 mL) followed by brine (400 mL), dried over anhydrous Na_2SO_4 and concentrated to afford crude compound. Obtained crude product was purified by column chromatography followed by trituration using MTBE to afford 8-bromo-6-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (9.1 g, 57%) as off white solid.

Synthesis of 8-bromo-9-ethyl-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (Intermediate B-9):

[0154]



[0155] Step1: A stirring suspension of $\text{Pd}(\text{dppf})\text{Cl}_2 \cdot \text{DCM}$ (1.03 g, 1.27 mmol, 0.1 eq) and Cs_2CO_3 (16.5g, 50.63 mmol, 4 eq) in DMF (78 mL) was deoxygenated by Ar for 5 min. 5-Fluoro-2-iodoaniline (3.0 g, 12.65 mmol, 1 eq) and triethylborane (1M) (16.45 mL, 16.45 mmol, 1.5 eq) was added to the reaction mixture and again deoxygenated by Ar for 10 min. Finally the reaction mixture was stirred for 16 h at 70°C. The reaction mixture was cooled to RT and then diluted with EtOAc (100 mL). The organic layer was washed with water (5 x 20 mL) and brine (20 mL), dried over Na_2SO_4 , concentrated under reduced pressure to get the crude product. Crude product was purified by column chromatography (230-400 mesh silica gel; 10% EtOAc/hexane; R_f -value-0.5) to afford 2-ethyl-5-fluoroaniline (1.20 g, 68%) as off white solid.

[0156] Step2: To a stirring solution of 2-ethyl-5-fluoroaniline (2.0 g, 14.38 mmol 1 eq), pyridine (2.31 mL, 28.77 mmol, 2 eq), and DMAP (0.018 g, 0.144 mmol, 0.01 eq) in DCM (50 mL) was drop wise added pivaloyl chloride (1.91 g, 15.82 mmol, 1.1 eq) at 0°C. The reaction mixture was then stirred for another 1 h at 0°C and then poured into ice cooled HCl (20 ml) (1M) solution. The organic layer was separated and the aqueous layer was extracted with Et_2O (2 x 20 mL). The combined organic layer was washed with water (2 x 20 mL) and brine (20 mL). The organic layer was dried over anhydrous Na_2SO_4 , concentrated under reduced pressure to get the crude material. Crude product was purified by column chromatography (230-400 mesh silica gel; 10%EtOAc/hexane; R_f -value-0.5) to afford N-(2-ethyl-5-fluorophenyl)pivalamide (2.3 g, 72%) as off white solid.

[0157] Step3: To a stirring solution of N-(2-ethyl-5-fluorophenyl)pivalamide (25.0 g, 112.1 mmol, 1 eq), in toluene (500 mL) was added p-TSA (19.3 g, 112.1 mmol, 1 eq), $\text{Pd}(\text{OAc})_2$ (2.52 g, 11.21 mmol, 0.1 eq) and NBS (21.5 g, 123.3 mmol, 1.1 eq) respectively at RT. The reaction mixture was stirred for 48 h at RT under air. Solvent was evaporated and the residue was dissolved in EtOAc (1000 mL) and washed with water (2 x 200 mL) followed by brine (200 mL). The organic layer was dried over anhydrous Na_2SO_4 , concentrated under reduced pressure to get the crude material. Crude product was purified by column chromatography (230-400mesh silica gel; 10%EtOAc/hexane; R_f -value-0.45) to afford N-(2-bromo-6-ethyl-3-fluorophenyl)pivalamide (5.0 g, 15%) as off white solid.

[0158] Step4: 70% H_2SO_4 (70 mL) was added to N-(2-bromo-6-ethyl-3-fluorophenyl)pivalamide (9.2 g, 30.46 mmol, 1 eq) in a round bottom flask at 0°C. The reaction mixture was stirred for 20 min at 0°C and for 2 h 110°C. The reaction mixture then cooled to 0°C and basified by 20%-NaOH solution up to pH-14. The aqueous part was extracted with EtOAc (3 x 150 mL). The combined organic layers were washed with water (2 x 100 mL) and brine (100 mL). The organic layer was dried over anhydrous Na_2SO_4 , concentrated under reduced pressure to get the crude material. Crude product was purified by column chromatography (230-400mesh silica gel; 5% EtOAc/hexane; R_f -value-0.5) to afford 2-bromo-6-ethyl-3-fluoroaniline (5.9 g, 89%) as brown liquid.

[0159] Step5: A suspension of 2-bromo-6-ethyl-3-fluoroaniline (4.9 g, 22.47 mmol, 1 eq), 2-amino-2-methylpropanoic acid (4.63 g, 44.94 mmol, 2 eq), K_3PO_4 (9.54 g, 44.94 mmol, 2 eq) and cuprous chloride (0.22 g, 2.247 mmol, 0.1 eq) in dry DMSO (75 mL) in a sealed tube was deoxygenated with Ar for 20 min. Reaction mixture was then stirred at 140°C for 2 h. After completion of the reaction, the reaction mixture cooled to RT and filtered through celite bed and washed with EtOAc (100 mL). The filtrate was diluted with EtOAc (500 mL) and washed with water (3 x 150 mL), brine (200 mL), dried over anhydrous Na_2SO_4 and the solvent was evaporated under reduced pressure. Crude product was purified by column chromatography (230-400 mesh silica gel; 20% EtOAc/hexane; R_f -value-0.4) to afford 8-ethyl-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (2.83 g, 57%) light brown solid.

[0160] Step6: To a solution of 8-ethyl-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (3.4 g, 15.3 mmol, 1 eq.) in toluene (110 mL) was added Lawesson's reagent (8.03 g, 19.89 mmol, 1.3 eq.) at RT and the reaction mixture was

refluxed at 120°C for 40 min. After completion of reaction (monitored by TLC), the reaction mixture was quenched with sat. NaHCO₃ solution (80 mL) followed by extraction with EtOAc (2 x 70 mL). Combined organic layers were washed with water (80 mL), brine (80 mL), dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product which was purified by column chromatography (230-400 mesh silica gel; 20% EtOAc/hexane; R_F-value-0.6) to afford 8-ethyl-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxaline-2(1H)-thione (3.0 g, 82%) as yellow solid. **Step7:** To a stirring solution of 8-ethyl-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxaline-2(1H)-thione (3.0 g, 12.58 mmol, 1 eq) in THF (50 mL) was drop wise added hydrazine hydrate (3.03 mL, 62.93 mmol, 5 eq) at 0°C. The reaction mixture was stirred for 16 h at RT. TEA (8.56 mL, 62.93 mmol, 5 eq) was added to the reaction mixture and stirred for another 10 min. Acetyl chloride (2.7 mL, 37.74 mmol, 3 eq) was added to the reaction mixture very slowly at 0°C and then stirred for 2 h at RT. The reaction mixture was diluted with water (100 mL) and extracted with DCM (3 x 100 mL). The combined organic layer was washed with brine (100 mL). The organic layer was dried over Na₂SO₄, concentrated under reduced pressure to get the crude material which purified by washing with diethyl ether to afford the N'-(8-ethyl-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-ylidene)acetohydrazide (3.4 g, 97%) as off white solid.


[0161] Step8: N'-(8-ethyl-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-ylidene)acetohydrazide (3.4 g, 12.23 mmol, 1 eq) was taken in round bottom flask (100 mL) and cooled to -10°C. Phosphorus oxychloride (11.7 mL, 122.3 mmol, 10 eq) was then added drop wise to the compound followed by drop wise addition of TEA (1.66 mL, 12.23 mmol, 1 eq). The reaction mixture was stirred at -10°C for 10 min and then for 10 min at RT and finally heated to reflux for 2 h. The reaction mixture was cooled to 0°C and then drop wise added into crushed ice with constant stirring. To this aqueous part was slowly added cold ammonium solution up to pH~12. The aqueous part was extracted with DCM (3 x 100 mL). The combined organic layer was washed with brine (100 mL). The organic layer was dried over anhydrous Na₂SO₄, concentrated under reduced pressure to get the crude material. Crude product was purified by column chromatography (230-400mesh silica gel; 5% MeOH/DCM; R_F-value-0.4) to afford 9-ethyl-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (1.4 g, 44%) as light yellow solid.

[0162] Step9: A stirred solution of 9-ethyl-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (1.4 g, 5.38 mmol, 1 eq) in DMF (30 mL) at -10°C was treated portion wise over 10 min with solid N-bromosuccinimide (1.01 g, 5.65 mmol, 1.05 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (300 mL) and organic layers was washed with water (5 x 50 mL), brine (50 mL), dried over anhydrous Na₂SO₄ and evaporated under reduced pressure. Crude product was purified by silica gel (230-400) column chromatography (5% MeOH/DCM; R_F-value-0.4) to afford 8-bromo-9-ethyl-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (1.45 g, 80%) as off white solid.

[0163] 1H-NMR (400 MHz; DMSO-D₆, 20°C): δ 7.54-7.56 (1H), 6.69 (s, 1H), 2.82-2.85 (2H), 2.38 (s, 3H), 1.46 (bs, 3H), 0.94-0.97 (3H)

Synthesis of 8-bromo-6-fluoro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**Intermediate B-10**):

[0164]



[0165] Step1: To an ice cold solution of 3-bromo-4-fluorophenol (100 g, 523.56 mmol, 1 eq) and ammonium nickel(II) sulfate hexahydrate (103.4 g, 261.77 mmol, 0.5 eq) in DCM (1000 mL) fuming nitric acid was added drop wise and the reaction mixture was stirred at same temperature for 3 h (TLC). Reaction mixture was poured into crushed ice and diluted with DCM. The extracted organic layer was washed with brine; the organic layer was dried over Na₂SO₄ and concentrated. The crude product was purified by column chromatography (230-400 mesh silica gel, TLC system: EtOAc / hexane (2:8);

$R_f = 0.2$) to give 3-bromo-4-fluoro-2-nitrophenol (35 g, 28%).

[0166] Step2: To a stirred solution of 3-bromo-4-fluoro-2-nitrophenol (35 g, 148.30 mmol, 1 eq) in ACN (40 mL) potassium carbonate (61.48 g, 444.91 mmol, 3 eq) and iodo methane (63.15 g, 444.91 mmol, 3 eq) were added and the reaction mixture was heated at 80°C for 4 h. After completion (monitored by TLC) the reaction mixture was diluted with EtOAc and washed with water. The extracted organic layer was washed with brine, dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude product was purified by column chromatography (230-400 mesh silica gel, TLC system: EtOAc / hexane (2:8); $R_f = 0.6$) to give 2-bromo-1-fluoro-4-methoxy-3-nitrobenzene (35 g, 94%).

[0167] Step3: To a stirred solution of 2-bromo-1-fluoro-4-methoxy-3-nitrobenzene (35 g, 140 mmol, 1 eq) in EtOH and water (300 mL, 1:1), iron powder (78.17 g, 1400 mmol, 10 eq) and ammonium chloride (74.886 g, 1400 mmol, 10 eq) were added. The reaction mixture was heated to reflux for 12 h. After completion (monitored by TLC) the reaction mixture was filtered through a pad of celite. The filtrate was concentrated under reduced pressure to get crude product. The crude product was purified by column chromatography (230-400 mesh silica gel, TLC system: EtOAc / hexane (2:8); $R_f = 0.6$) to give 2-bromo-3-fluoro-6-methoxyaniline (30 g, 97%).

[0168] Step4: A suspension of 2-bromo-3-fluoro-6-methoxyaniline (30 g, 136.36 mmol, 1 eq), 2-aminoisobutamic acid (44.65 g, 340.83 mmol, 2.5 eq), DBU (49.70 g, 272.21 mmol, 2 eq) and copper iodide (2.59 g, 13.59 mmol, 0.1 eq.) in dry DMA (300 mL) in a round bottom flask was deoxygenated with Ar for 20 min. Reaction mixture was then stirred at 120°C for 16 h. After completion of the reaction, (monitored by TLC), it was filtered through a pad of celite and washed with EtOAc (500 mL), washed with water (3x500 mL), brine (500 mL), dried over anhydrous Na_2SO_4 and evaporated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; TLC system: EtOAc / hexane (2:8); $R_f = 0.3$) to give 5-fluoro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (14 g, 46%) as brown solid.

[0169] Step5: To a solution of 5-fluoro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (14 g, 62.22 mmol, 1 eq) in toluene (150 mL) was added Lawesson's reagent (37.75 g, 93.33 mmol, 1.5 eq) at RT and the reaction mixture was refluxed at 120°C for 1 h. After completion of reaction (monitored by TLC), the reaction mixture was quenched with sat. NaHCO_3 solution (200 mL) followed by extraction with EtOAc (2x200 mL). Combined organic layers were washed with water (300 mL), brine (100 mL), dried over anhydrous Na_2SO_4 and the solvent was evaporated to get the crude product which was purified by column chromatography using 230-400 mesh silica gel and 10% EtOAc in hexane as an eluting solvent to afford 5-fluoro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxaline-2(1H)-thione (10 g, 67%) as a yellow solid. TLC system: EtOAc / hexane (2:8); $R_f = 0.4$.


[0170] Step6: To a solution of 5-fluoro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxaline-2(1H)-thione (10 g, 41.61 mmol, 1 eq) in n-BuOH (100 mL) was added acetic hydrazide (10.80 g, 145.93 mmol, 3.5 eq) followed by addition of acetic acid (10 mL) and then the reaction mixture was stirred at 140°C for 16 h. After completion of reaction (monitored by TLC) reaction mixture was diluted with water (100 mL) and extracted with EtOAc (2x50 mL). Combined organic layers were washed with water (100 mL), brine (100 mL), dried over anhydrous Na_2SO_4 and evaporated to get the crude product which was purified by column chromatography using 5% MeOH in DCM as an eluting solvent and 230-400 silica gel to afford 6-fluoro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (3.5 g, 32%) as off white solid (TLC system, 5% MeOH in DCM, $R_f=0.2$).

[0171] Step7: A stirred solution of 6-fluoro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (3.5 g, 13.34 mmol, 1 eq) in DMF (30 mL) at 0°C was treated portion wise over 10 min with solid N-bromosuccinamide (2.48 g, 13.93 mmol, 1.05 eq). Reaction mixture was allowed to warm to RT and stirred for 30 min. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (50 mL) and organic layers were washed with water (3x100 mL), brine (100 mL), dried over anhydrous Na_2SO_4 and the solvent was evaporated under reduced pressure. Crude product was purified by column chromatography using 1.5% MeOH in DCM as an eluting solvent and 230-400 silica gel to afford 8-bromo-6-fluoro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (1.9 g, 42%) as off white solid (TLC system, 5% MeOH in DCM, $R_f=0.4$).

[0172] $^1\text{H-NMR}$ (400 MHz; $\text{DMSO}-d_6$, 20°C): δ 7.58 (d, 1H), 6.73 (s, 1H), 3.56 (s, 3H), 2.44 (s, 3H), 1.49 (s, 3H), 1.46 (s, 3H).

Synthesis of 8-bromo-1-cyclopropyl-6-fluoro-9-methoxy-4,4-dimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (Intermediate B-11):

[0173]




[0174] Step1: To a solution of 5-fluoro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxaline-2(1H)-thione (10 g, 41.61 mmol, 1 eq) in *n*-BuOH (100 mL) was added cyclopropanecarbohydrazide (14.58 g, 145.62 mmol, 3.5 eq) followed by addition of acetic acid (10 mL) and then the reaction mixture was stirred at 140°C for 16 h. After completion of reaction (monitored by TLC) reaction mixture was diluted with water (100 mL) and extracted with EtOAc (2 x 50 mL). Combined organic layers washed with water (100 mL), brine (100 mL), dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product which was purified by column chromatography using 5% MeOH in DCM as an eluting solvent and 230-400 silica gel to afford 1-cyclopropyl-6-fluoro-9-methoxy-4,4-dimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (2.6 g, 32%) as off white solid.

[0175] Step2: A stirred solution of 1-cyclopropyl-6-fluoro-9-methoxy-4,4-dimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (2.6 g, 9.02 mmol, 1 eq) in DMF (30 mL) at 0°C was treated portion wise over 10 min with solid N-bromosuccinamide (1.68 g, 9.43 mmol, 1.05 eq). Reaction mixture was allowed to warm up to RT and stirred for 30 min. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (50 mL) and organic layers were washed with water (3x100 mL), brine (100 mL), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure. Crude product was purified by column chromatography using 1.5% MeOH in DCM as an eluting solvent and 230-400 silica gel to afford 8-bromo-1-cyclopropyl-6-fluoro-9-methoxy-4,4-dimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (1.8 g, 55%) as off white solid.

Synthesis of 8-bromo-1-cyclopropyl-6-fluoro-4,4,9-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**Intermediate B-12**):

[0176]



[0177] Step1: To a stirring solution of 5-fluoro-3,3,8-trimethyl-3,4-dihydroquinoxaline-2(1H)-thione (1.7 g, 7.58 mmol, 1 eq) in THF (40 mL) was drop wise added hydrazine hydrate (1.72 mL, 37.94 mmol, 5 eq) at 0°C. The reaction mixture was stirred for 16 h at RT. TEA (5.13 mL, 37.94 mmol, 5 eq) was added to the reaction mixture and stirred for another 10 min. Cyclopropanecarbonyl chloride (2.39 g, 22.76 mmol, 3 eq) was added to the reaction mixture very slowly at 0°C and then stirred for 2 h at RT. The reaction mixture was diluted with water (100 mL) and extracted with DCM (3x100 mL). The combined organic layers were washed with brine (100 mL). The organic layer was dried over Na₂SO₄, concentrated under reduced pressure to get the crude product which was purified by washing with diethyl ether to afford N'-(5-fluoro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-ylidene)cyclopropanecarbohydrazide (2.1 g, 95%) as off white solid.

[0178] Step2: N'-(5-fluoro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-ylidene)cyclopropanecarbohydrazide (1.44 g, 4.96 mmol, 1 eq) was taken in a round bottom flask (50 mL) and then cooled to -10°C. Phosphorus oxychloride (4.7 mL, 49.65 mmol, 10 eq) was then added drop wise to the compound followed by drop wise addition of TEA (10.67 mL, 4.96 mmol, 1 eq). After that the reaction mixture was stirred at -10°C for 10 min and then 10 min at RT and finally at reflux condition for 2 h. The reaction mixture was cooled to 0°C and then it was drop wise added into crushed ice with constant stirring. To this aqueous part was slowly added cold ammonium solution up to pH-12. The aqueous part was extracted with DCM (3x50 mL). The combined organic layer were washed with brine (50 mL). The organic layer was dried over anhydrous Na₂SO₄, concentrated under reduced pressure to get the crude product. Crude product was purified by column chromatography (230-400 mesh silica gel; 5% MeOH/DCM; R_f-value-0.4) to afford 1-cyclopropyl-6-fluoro-4,4,9-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (1.8 g, 55%) as off white solid.


trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.24 g, 18%) as light yellow solid.

[0179] Step3: A stirred solution of 1-cyclopropyl-6-fluoro-4,4,9-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.55 g, 2.02 mmol, 1 eq) in DMF (15 mL) at -10°C was treated portion wise over 10 min with solid N-bromosuccinimide (0.36 g, 2.02 mmol, 1 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (80 mL) and organic layers were washed with water (5x20 mL), brine (20 mL), dried over anhydrous Na₂SO₄ and solvent was evaporated under reduced pressure. Crude product was purified by silica gel (230-400) column chromatography (5% MeOH/DCM; R_f-value-0.4) to afford 8-bromo-1-cyclopropyl-6-fluoro-4,4,9-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.55 g, 78%) as off white solid.

[0180] ¹H-NMR (400 MHz; DMSO-D₆, 20°C): δ 7.58 (d, 1H), 6.58 (s, 1H), 2.45 (s, 3H), 1.71-1.77 (1H), 1.44 (bs, 6H), 1.02-1.27 (4H).

Synthesis of 8-bromo-1-ethyl-6-fluoro-4,4,9-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**Intermediate B-30**):

[0181]



[0182] Step1: To a stirred solution of 5-fluoro-2-methyl-phenylamine (100 g, 0.8 mol, 1 eq) in DCM (1.5 L) was added pyridine (129 mL, 1.6 mol, 2 eq), followed by DMAP (1 g, 0.008 mol, 0.01 eq) at room temperature. To this reaction mixture was added dropwise pivoyl chloride (109 mL, 0.88 mol, 1.1 eq) at 0°C and the resulting reaction mixture was stirred at 0°C for 2 h. After complete consumption of starting material (monitored by TLC in 10% EtOAc-Hex, R_f 0.7), the reaction mixture was poured into an ice-cooled 1N HCl solution (1.5 L) and stirred for 30 min. The two layers were separated and the organic layer was washed with an 1N HCl solution (1 L), followed by saturated NaHCO₃ solution (1 L) and brine (1 L), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. This crude residue was triturated using hexane to afford N-(5-fluoro-2-methyl-phenyl)-2,2-dimethyl-propionamide (158 g, 95%) as a white solid.

[0183] Step2: To a stirring solution of N-(5-fluoro-2-methyl-phenyl)-2,2-dimethyl-propionamide (90 g, 0.43 mol, 1 eq) in toluene (2 L) were added *para*-toluene sulfonic acid monohydrate (82.0 g, 0.43 mol, 1 eq) and Pd(OAc)₂ (9.66 g, 0.043 mol, 0.1 eq) followed by N-bromo succinimide (84.5 g, 0.473 mol, 1.1 eq) at room temperature and the resulting reaction mixture was stirred at room temperature for 48 h. Reaction monitoring by TLC (20% EtOAc in Hexane) showed formation of desired product (R_f 0.4) along with ~60% unreacted starting material (R_f 0.5). A second batch was then run starting from 100 g of N-(5-fluoro-2-methyl-phenyl)-2,2-dimethyl-propionamide, following the same procedure as described above.

[0184] The combined reaction mixtures of the two batches (190 g of starting material in total) were then concentrated and the obtained residue was diluted in EtOAc and washed with water (twice). The combined organic layers were then dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude residue was purified by column chromatography (100-200 mesh silica gel; 1-5% EtOAc/hexane as eluent) to afford N-(2-bromo-3-fluoro-6-methylphenyl)-2,2-dimethyl-propionamide (60 g, 22.9%) as a white solid along with recovered unreacted starting material (125 g).

[0185] Step3: To a pre-cooled solution of 70% H₂SO₄ in water (430 mL) was added N-(2-bromo-3-fluoro-6-methylphenyl)-2,2-dimethyl-propionamide (60 g, 0.208 mol, 1 eq) portion wise at 0°C and the resulting reaction mixture was stirred at room temperature for 10 min and then heated at 100°C for 2 h. After complete consumption of starting material (monitored by TLC in 10% EtOAc-Hexane, R_f 0.8), the reaction mixture was cooled to 0°C and the pH adjusted to ~10 with 10% NaOH solution. The resulting basic aqueous fraction was extracted with EtOAc (3 x 500 mL), the combined organic layers were washed with water and brine subsequently, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford a crude residue. This was purified by column chromatography (silica 100-200 mesh; 1-3% EtOAc/hexane as eluent) to afford 2-bromo-3-fluoro-6-methyl-phenylamine (36 g, 85%) as a pale yellow liquid.

[0186] Step4: To a stirred suspension of 2-bromo-3-fluoro-6-methyl-phenylamine (36 g, 0.1764 mol, 1 eq) in dry DMSO

(540 ml) was added 2-amino-2-methyl-propionic acid (16.9 g, 0.1636 mol, 0.93 eq), followed by K_3PO_4 (75 g, 0.3528 mol, 2 eq) at room temperature. The resulting reaction mixture was degassed with nitrogen for 30 minutes, then $CuCl$ (1.75 g, 0.0176 mol, 0.1 eq) was added and the reaction mixture was heated at $140^\circ C$ for 5 h. After complete consumption of the starting material (monitored by TLC, 20% EtOAc-Hexane, Rf 0.4), the reaction mixture was cooled to room temperature and filtered over a bed of celite. After washing with EtOAc (500 mL), the resulting filtrate was poured into ice-cold water (2.5 L) and the resulting aqueous fraction was extracted with EtOAc (2 x 750 mL). The combined organic layers were washed with water (2 x 750 mL) and brine (500 ml), dried over anhydrous Na_2SO_4 and evaporated under reduced pressure to afford a crude residue. This was triturated with hexane, filtered and dried to afford 5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxalin-2-one (19 g, 52%) as a brown solid.

[0187] Step5: To a solution of 5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxalin-2-one (19, 91.2 mmol, 1 eq) in toluene (380 mL) was added Lawesson's reagent (55.5 g, 137 mmol, 1.5 eq) at RT and the reaction mixture was refluxed at $120^\circ C$ for 1 h. After complete consumption of starting material (monitored by TLC in 20% EtOAc-Hexane, Rf 0.7), the reaction mass was cooled to room temperature and quenched with sat. $NaHCO_3$ solution (250 mL) and the resulting aqueous fraction was extracted with EtOAc (3 x 250 mL). The combined organic layers were washed with water (250 mL) and brine (250 mL), dried over anhydrous Na_2SO_4 and evaporated under reduced pressure. The obtained crude residue was purified by column chromatography (silica 100-200 mesh; 3-5% EtOAc/hexane) to afford 5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxaline-2-thione (19 g, 93%) as a yellow solid.


[0188] Step6: To a stirring solution of 5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxaline-2-thione (9 g, 0.0401 mol, 1 eq) in tetrahydrofuran (225 mL) was added dropwise hydrazine hydrate (6.02 g, 0.1203 mol, 3 eq) at $0^\circ C$ and the reaction mixture was stirred at room temperature for 16 h. Triethyl amine (27.8 mL, 0.2006 mol, 5 eq) and propionyl chloride (10.5 mL, 0.1203 mol, 3 eq) were added to the reaction mixture dropwise at $0^\circ C$ and this was stirred for 2 h at room temperature. After complete consumption of starting material (monitored by LCMS), the reaction mixture was diluted with water (250 mL) and extracted with 10% MeOH-DCM (5 x 250 mL). The combined organic layers were washed with brine (250 mL), dried over Na_2SO_4 and concentrated under reduced pressure to afford propionic acid (5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxalin-2-ylidene)-hydrazide (12.5 g crude material) as an off-white solid.

[0189] Step7: Crude propionic acid (5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxalin-2-ylidene)-hydrazide (12.5 g, 0.0449 mol, 1 eq) was taken up in a round-bottom flask, that was then cooled to $-10^\circ C$. Phosphorus oxydichloride (20.9 mL, 0.2243 mol, 5 eq) was then added dropwise to the compound, followed by dropwise addition of triethyl amine (6.25 mL, 0.0449 mol, 1 eq). After that, the reaction mixture was stirred at $-10^\circ C$ for 10 min followed by stirring at room temperature for 10 min and finally at reflux conditions for 4 h. After complete consumption of starting material (monitored by LCMS), the reaction mixture was cooled to $0^\circ C$ and quenched with crushed ice in water (250 mL). The aqueous part was then basified by dropwise addition of a cold ammonia solution (250 mL). The resulting basic aqueous fraction was extracted with EtOAc (3 x 500 mL) and the combined organic layers were washed with brine (250 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained crude residue was purified by trituration using MTBE to afford 1-ethyl-6-fluoro-4,4,9-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (6.5 g, 56%) as an off-white solid.

[0190] Step8: To a solution of 1-ethyl-6-fluoro-4,4,9-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (10 g, 0.0384 mol, 1 eq) in DMF (250 mL) was added NBS (7.52 g, 0.0422 mol, 1.1 eq) portion wise at $-10^\circ C$ and the resulting reaction mixture was stirred at room temperature for 4 h. After complete consumption of starting material (monitored by LCMS), the reaction mixture was diluted with water (500 mL) and extracted with EtOAc (2 x 250 mL). The combined organic layers were washed with water (500 mL) and brine (400 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained crude residue was purified by column chromatography (230-400 mesh silica gel; 5% MeOH/DCM as eluent) to afford 8-bromo-1-ethyl-6-fluoro-4,4,9-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (7.5 g, 58%) as an off-white solid.

Synthesis of 8-bromo-6,7-difluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**Intermediate B-38**):

[0191]



[0192] Step1: To a suspension of 4,5-difluoro-2-methylaniline (0.5 g, 3.49 mmol, 1 eq) in DCM (30 ml), Br₂ (0.55 g, 0.17 ml, 3.49 mmol, 1 eq) in DCM (20 ml) was added at 0°C. Then the reaction was stirred at RT for 2 h. After completion of starting material, the reaction mass was quenched with saturated NaHCO₃ solution (50 ml). The organic layer was separated and aqueous layer was extracted with DCM (50 ml). Combined organic layers were washed with water (150 ml), followed by brine (150 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated to afford crude product which was purified by column chromatography (230-400 mesh silica gel; 10% EtOAc/hexane; R_F-value-0.6) to afford 2-bromo-3,4-difluoro-6-methylaniline (0.4 g, 52%).

[0193] Step2: A suspension of 2-bromo-3,4-difluoro-6-methylaniline (0.25 g, 1.13 mmol, 1 eq), 2-aminobutyric acid (0.23 g, 2.26 mmol, 2 eq) in DMA (10 ml) in a sealed tube was deoxygenated with Ar for 20 min. DBU (0.35 ml, 2.26 mmol, 2 eq) and CuI (0.02 g, 0.113 mmol, 0.1 eq) were added and reaction mixture was stirred at 140°C for 16 h. After completion of the reaction, it was filtered through celite bed and washed by EtOAc (100 ml). The filtrate was diluted with EtOAc (100 ml) and washed with water (3x150 ml), brine (200 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography (100-200 mesh silica gel; 30% EtOAc/hexane; R_F-value-0.4) to afford 5,6-difluoro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-one (0.2 g, 75%).

[0194] Step3: To a solution of 5,6-difluoro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-one (3.66 g, 17.6 mmol, 1 eq) in toluene (75 ml) was added Lawesson's reagent (10.67 g, 26.2 mmol, 1.5 eq) at RT and the reaction mixture was heated at 120°C for 40 min. After completion of reaction (monitored by TLC), the reaction mixture was quenched with sat. NaHCO₃ solution (100 ml) followed by extraction with EtOAc (2x100 ml). Combined organic layers were washed with water (100 ml), brine (100 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product which was purified by column chromatography (230-400 mesh silica gel; 20% EtOAc/hexane; R_F-value-0.4) to afford 5,6-difluoro-3,3,8-trimethyl-3,4-dihydroquinoxaline-2(1H)-thione (2.8 g, 71%) as yellow solid.

[0195] Step4: To a stirring solution of 5,6-difluoro-3,3,8-trimethyl-3,4-dihydroquinoxaline-2(1H)-thione (5.50 g, 24.55 mmol, 1 eq) in THF (30 ml) was drop wise added hydrazine hydrate (5.17 ml, 122.76 mmol, 5 eq) at 0°C. The reaction mixture then stirred for 16 h at RT. TEA (16.7 ml, 122.76 mmol, 5 eq) was added to the reaction mixture and stirred for another 10 min. Acetyl chloride (5.78 g, 73.65 mmol, 3 eq) was added very slowly at 0°C and then the mixture was stirred for 2 h at RT. The reaction mixture was diluted with water (50 ml) and extracted by DCM (5x100 ml). The combined organic layers were washed by brine (100 ml). The organic layer was dried over Na₂SO₄, concentrated under reduced pressure to get the crude material which purified by washing with diethyl ether to afford N'-(5,6-difluoro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-ylidene)acetohydrazide (5.5 g, 85%) as a white solid.


[0196] Step5: N'-(5,6-difluoro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-ylidene)acetohydrazide (5.5 g, 20.8 mmol, 1 eq) was taken in round bottom flask (50 ml) and then cooled to -10°C. Phosphorus oxychloride (18.4 ml, 197.6 mmol, 9.5 eq) was then added drop wise to the compound followed by drop wise addition of TEA (2.9 ml, 20.8 mmol, 1 eq). After that the reaction mixture was stirred at -10°C for 10 min and then 10 min at RT and finally at reflux condition for 4 h. The reaction mixture was cooled to 0°C and then drop wise added into crushed ice with constant stirring. To this aqueous part was slowly added cold ammonium solution (100 ml). The aqueous part was extracted by DCM (2x100 ml). The combined organic layer was washed by brine (100 ml). The organic layer was dried over anhydrous Na₂SO₄, concentrated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; 5% MeOH/DCM; R_F-value-0.4) to afford 6,7-difluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (3.0 g, 59%) as yellow solid.

[0197] Step6: A stirred solution of 6,7-difluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (4.0 g,

16.2 mmol, 1 eq) in DMF (40 ml) at -10°C was treated portion wise over 10 min with solid N-bromosuccinimide (3.1 g, 17.1 mmol, 1.05 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (300 ml) and organic layers were washed with water (5x50 ml), brine (50 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure to get the crude product which was purified by silica gel (230-400) column chromatography (5% MeOH/DCM; R_f-value-0.3) to afford 8-bromo-6,7-difluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (3.3 g, 63%) as off white solid.

Synthesis of 8-bromo-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline-9-carbonitrile (**intermediate B-39**):


[0198]



[0199] A stirred solution of 6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline-9-carbonitrile (for synthesis see **intermediate B-41**) (2.0 g, 7.78 mmol, 1 eq) in DMF (20 ml) at 0°C was treated portion wise over 10 min with solid N-bromosuccinimide (1.38 g, 7.78 mmol, 1 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (100 ml) and organic layers were washed with water (5x100 ml), brine (100 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure to get the crude product which was purified by silica gel (230-400) column chromatography (5% MeOH/DCM; R_f-value-0.3) to afford 8-bromo-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline-9-carbonitrile (1.5 g, 58%) as off white solid.

Synthesis of (8-bromo-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl)MeOH (**intermediate B-41**):

[0200]



[0201] Step1: A solution of 6-fluoro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (for synthesis see **intermediate B-10**) (12 g, 45.80 mmol, 1 eq) in aqueous HBr (350 ml) was heated up to 80°C and then acetic acid (100 ml) was added and heating was continued at 120°C for 48 h. After completion of reaction (monitored by TLC) reaction mixture is diluted with ice cold water (1000 ml) and basified with sodium bicarbonate (pH>8). The reaction

mixture was extracted with EtOAc (3x500 ml). Combined organic layers were washed with water (1000 ml), brine (1000 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product which was purified by column chromatography using 3% MeOH in DCM as an eluting solvent and 230-400 silica gel to afford 6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-ol (8 g, 70%) as off white solid (TLC system, 5% MeOH in DCM, R_f-0.2).

[0202] Step2: To an ice cold solution of 6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-ol (8 g, 32.25 mmol, 1 eq) in DCM, TEA (4.89g, 48.32 mmol, 1.5 eq) and DMAP (5.91 g, 48.37 mmol, 1.5 eq) were added and the mixture was stirred at same temperature for 10 min. Then triflic anhydride (10.92 g, 38.70 mmol, 1.2 eq) was added drop wise and the mixture was stirred for 16h. After completion of reaction (monitored by TLC) reaction mixture was diluted with ice cold water (1000 ml) and the reaction mixture was extracted with DCM (2 x 500 ml). Combined organic layers were washed with water (1000 ml), brine (1000 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product which was purified by column chromatography using 2% MeOH in DCM as an eluting solvent and 230-400 silica gel to afford 6-fluoro-1,4-dimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl trifluoromethanesulfonate (4 g, 33%) as liquid. (TLC system, 5% MeOH in DCM, R_f-0.5).

[0203] Step3: To an argon purged solution of 6-fluoro-1,4-dimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl trifluoromethanesulfonate (4 g, 10.52 mmol, 1 eq) in 1,4-dioxan, N,N'-dimethylethylenediamine (0.611 g, 37.67 mmol, 0.5 eq) and zinc cyanide (0.962 g, 10.52 mmol, 1 eq) were added and the mixture was stirred for 10 min. Then Pd₂(dba)₃ (1.089g, 1.052mmol, 0.1 eq) and xantphos (0.611 g, 1.05 mmol, 0.1 eq) were added and the reaction mixture was purged for another 5 min and the reaction mixture was heated to 100°C for 12 h. After completion of reaction (monitored by TLC) reaction mixture was filtered through a pad of celite. The filtrate was concentrated under vacuo to get the crude product which was purified by column chromatography using 4% MeOH in DCM as an eluting solvent and 230-400 silica gel to afford 6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline-9-carbonitrile (3 g, 37%) as liquid (TLC system, 5% MeOH in DCM, R_f-0.4).


[0204] Step4: To a solution of 6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline-9-carbonitrile (0.1 g, 0.389 mmol, 1 eq) in toluene (5 ml), DIBAL-H (0.2 ml, 0.389 mmol, 1 eq) was added at 0°C. The reaction was stirred at same temperature for 3 h. After completion of reaction (monitored by TLC), reaction mixture was quenched with NH₄Cl and extracted with EA. The organic layer was washed with water, brine, dried over Na₂SO₄, filtered and the solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography to afford 6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline-9-carbaldehyde (0.04 g, 34%) as brown solid.

[0205] Step5: To a solution of 6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline-9-carbaldehyde (0.04 g, 0.15 mmol, 1 eq) in THF (5 ml), LAH (0.005 g, 0.15 mmol, 1 eq) was added at 0°C. The reaction mixture was stirred at same temperature for 3 h. After completion of reaction (monitored by TLC), reaction mixture was quenched with saturated Na₂SO₄ solution and extracted with EA. The organic layer was washed with water, brine, dried over Na₂SO₄, filtered and the solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography to afford (6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl)MeOH (0.01 g, 25%) as brown gum.

[0206] Step6: A stirred solution of (6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl)MeOH (0.18 g, 0.692 mmol, 1 eq) in DMF (15 ml) at 0°C was treated portion wise over 10 min with solid N-bromosuccinimide (0.123 g, 0.692 mmol, 1 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (30 ml) and organic layers were washed with water (5x50 ml), brine (50 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure to get the crude product which was purified by silica gel (230-400) column chromatography (5% MeOH/DCM; R_f-value-0.3) to afford (8-bromo-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl)MeOH (0.12 g, 50%) as off white solid.

Synthesis of 8-bromo-6-chloro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**intermediate B-42**):

[0207]



[0208] Step1: A stirred solution of 5-chloro-2-methylaniline (25.0 g, 176.55 mmol, 1 eq) in DMF (530 ml) at -10°C was treated portion wise over 10 min with solid N-bromosuccinimide (69.13 g, 388.4 mmol, 2.2 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction, reaction mixture was diluted with EtOAc (3000 ml) and organic layers was washed with water (5x500 ml), brine (500 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure to get the crude product which was purified by silica gel (230-400) column chromatography (10% EtOAc/hexane; R_F-value-0.6) to afford 2,4-dibromo-3-chloro-6-methylaniline (35 g, 66%) as brown solid.

[0209] Step2: A solution of 2,4-dibromo-3-chloro-6-methylaniline (7.5 g, 25.08 mmol, 1 eq), 2-amino-2-methylpropanoic acid (5.16 g, 50.16 mmol, 2 eq) and DBU (9.54 g, 44.94 mmol, 2 eq) in dry DMA (112 ml) in a sealed tube were deoxygenated by Ar for 10 min. Cuprous iodide (0.479 g, 2.5 mmol, 0.1 eq) was added to the reaction mixture and again deoxygenated by Ar for 10 min. Reaction mixture was then stirred at 140°C for 16 h. After completion of the reaction, the reaction mixture was cooled to RT and diluted with EtOAc (1000 ml) and washed with water (4 x 150 ml), brine (200 ml), dried over anhydrous Na₂SO₄ and evaporated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; 20% EtOAc/hexane; R_F-value-0.4) to afford 6-bromo-5-chloro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-one (2.2 g, 29%) light brown solid.

[0210] Step3: To a stirring solution of 6-bromo-5-chloro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-one (3.9 g, 12.87 mmol, 1 eq) in a mixture of 1,4-dioxan and water (1:1) (100 ml), was added ammonium chloride (10.32 g, 193.06 mmol, 15 eq) and zinc dust (12.26 g, 193.06 mmol, 15 eq). The reaction mixture was stirred for 16 h at reflux condition. The reaction mixture was cooled to RT and then filtered through sintered funnel. The filtrate was concentrated under reduced pressure to get the residue, which was dissolved in EtOAc (100 ml) and washed by water (2x30 ml) followed by brine (30 ml). The organic layer was dried over Na₂SO₄, concentrated under reduced pressure to get 5-chloro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-one (2.3 g, 80%) as off white solid.

[0211] Step4: To a solution of 5-chloro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-one (2.3 g, 10.23 mmol, 1 eq) in toluene (75 ml) was added Lawesson's reagent (5.37 g, 13.3 mmol, 1.3 eq) at RT and the reaction mixture was refluxed at 120°C for 40 min. After completion of reaction (monitored by TLC), the reaction mixture was quenched with sat. NaHCO₃ solution (80 ml) followed by extraction with EtOAc (2x70 ml). Combined organic layers were washed with water (80 ml), brine (80 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product which was purified by column chromatography (230-400 mesh silica gel; 20% EtOAc/hexane; R_F-value-0.6) to afford 5-chloro-3,3,8-trimethyl-3,4-dihydroquinoxaline-2(1H)-thione (2.0 g, 81%) as yellow solid.

[0212] Step5: To a stirring solution of 5-chloro-3,3,8-trimethyl-3,4-dihydroquinoxaline-2(1H)-thione (2.0 g, 8.33 mmol, 1 eq) in THF (50 ml) was drop wise added hydrazine hydrate (2.08 ml, 41.64 mmol, 5 eq) at 0°C. The reaction mixture was stirred for 16 h at RT. TEA (5.8 ml, 41.65 mmol, 5 eq) was added and the reaction mixture was stirred for another 10 min. Acetyl chloride (1.78 ml, 24.99 mmol, 3 eq) was added to the reaction mixture very slowly at 0°C and then stirred for 2 h at RT. The reaction mixture was diluted with water (50 ml) and extracted by DCM (3x100 ml). The combined organic layer was washed by brine (100 ml). The organic layer was dried over Na₂SO₄, concentrated under reduced pressure to get the crude product which was purified by washing with diethyl ether to afford N'-(5-chloro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-ylidene)acetohydrazide (2.2 g, 94%) as off white solid.


[0213] Step6: N'-(5-chloro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-ylidene)acetohydrazide (5.2 g, 18.57 mmol, 1 eq) was taken in round bottom flask (100 ml) and then cooled to -10°C. Phosphorus oxychloride (17.8 ml, 185.7 mmol, 10 eq) was then added drop wise to the compound followed by drop wise addition of TEA (2.52 ml, 18.57 mmol, 1 eq). After that the reaction mixture was stirred at -10°C for 10min and then 10 min at RT and finally at reflux condition for 2 h. The reaction mixture was cooled to 0°C and drop wise added into crushed ice with constant stirring. To this aqueous

part was slowly added cold ammonium solution up to pH~12. The aqueous part was extract by DCM (3x100 ml). The combined organic layer was washed by brine (100 ml). The organic layer was dried over anhydrous Na_2SO_4 , concentrated under reduced pressure to get the crude product which was purified by column chromatography (230-400mesh silica gel; 5% MeOH/DCM; R_f -value-0.4) to afford 6-chloro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (3.25 g, 67%) as light yellow solid.

[0214] Step7: A stirred solution of 6-chloro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (3.25 g, 12.4 mmol, 1 eq) in DMF (60 ml) at -10°C was treated portion wise over 10 min with solid N-bromosuccinimide (2.32 g, 13.02 mmol, 1.05 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (500 ml) and organic layers was washed with water (5x100 ml), brine (100 ml), dried over anhydrous Na_2SO_4 and the solvent was evaporated under reduced pressure to get the crude product which was purified by silica gel (230-400) column chromatography (5% MeOH/DCM; R_f -value-0.4) to afford 8-bromo-6-chloro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (3.0 g, 71%) as off white solid.

Synthesis of 8-bromo-9-(difluoromethyl)-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**intermediate B-43**):

[0215]



[0216] Step1: To a stirred solution of 4-fluoro-2-nitrobenzaldehyde (20 g, 118.27 mmol, 1 eq) in DCM (600 mL) was added DAST (23.26 ml, 177.51 mmol, 1.5 eq) at 0°C . The reaction mixture was stirred at RT for 4 h. After completion of starting material reaction mixture was quenched with sat. NaHCO_3 solution and extracted with DCM (2x500 mL). Combined organic layers was washed with water (500 mL) followed by brine (500 mL), dried over anhydrous Na_2SO_4 and concentrated. Crude product was purified by column chromatography using 2-3% EtOAc/hexane to afford 1-(difluoromethyl)-4-fluoro-2-nitrobenzene (22 g, 88%) as yellow liquid.

[0217] Step2: To a stirred solution of 1-(difluoromethyl)-4-fluoro-2-nitrobenzene (10 g, 52.16 mmol, 1 eq) in EtOH (372 mL) was added $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (47 g, 209.46 mmol, 4 eq) followed by con. HCl (35 mL) at 0°C . The reaction mixture was stirred at RT for 2 h. After completion of starting material reaction mixture was concentrated, residue was basified with 5N NaOH solution and extracted with MTBE (2 x 500 mL). Combined organic layers were washed with water (500 mL) followed by brine (500 mL), dried over anhydrous Na_2SO_4 and concentrated. Crude product was purified by column chromatography using 2-3% EtOAc/hexane to afford 2-(difluoromethyl)-5-fluoroaniline (7.0 g, 83%) as yellow gummy liquid.

[0218] Step3: To a solution of 2-(difluoromethyl)-5-fluoroaniline (4 g, 24 mmol, 1.0 eq) in DMF (40 mL) was added NBS (13 g, 74 mmol, 3.0 eq) portion wise at -10°C . Resulting reaction mixture was stirred at 0°C for 1 h. After completion of starting material (monitored by LCMS), reaction mixture was diluted with water (100 mL) and extracted with EtOAc (2 x 100 mL). Combined organic layers were washed with water (2 x 100 mL) and brine, dried over anhydrous Na_2SO_4 and concentrated to get crude 2,4-dibromo-6-(difluoromethyl)-3-fluoroaniline (7 g, 85%) was used for the next step without further purification.

[0219] Step4: A suspension of crude 2,4-dibromo-6-(difluoromethyl)-3-fluoroaniline (8 g, 25.2 mmol, 1 eq) in DMSO (80 ml), 2-aminobutyric acid (5.2 g, 50 mmol, 2 eq) was added at RT. The reaction mixture was then deoxygenated with Ar for 20 min. K_3PO_4 (10.6 g, 50 mmol, 2 eq) and CuCl (0.024 g, 2.5 mmol, 0.1 eq) were added and reaction mixture was then stirred at 140°C for 16 h. After completion of the reaction, it was filtered through celite bed and washed by EtOAc (100 ml). The filtrate was diluted with EtOAc (100 ml) and washed with water (3x150 ml), brine (200 ml), dried

over anhydrous Na_2SO_4 and the solvent was evaporated under reduced pressure to get crude product which was purified by column chromatography (100-200 mesh silica gel; 30% EtOAc/hexane; R_f -value-0.4) to afford 6-bromo-8-(difluoromethyl)-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (1.5 g, 20%).

[0220] Step5: To a stirring solution of 6-bromo-8-(difluoromethyl)-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (3.0 g, 9.3 mmol, 1 eq) in a mixture of 1,4-dioxan and water (1:1) (100 ml), was added NH_4Cl (7.47 g, 139.0 mmol, 15 eq) and zinc dust (9.08 g, 139.0 mmol, 15 eq). The reaction mixture was stirred for 16 h at reflux. The reaction mixture was cooled to RT and then filtered through sintered funnel. The filtrate was concentrated under reduced pressure to get the product, which was dissolved in EtOAc (100 ml) and washed by water (2x30 ml) followed by brine (30 ml). The organic layer was dried over Na_2SO_4 , concentrated under reduced pressure to get 8-(difluoromethyl)-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (2.0 g, 90%) as off white solid.


[0221] Step6: To a solution of 8-(difluoromethyl)-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (1 g, 4.09 mmol, 1 eq) in toluene (10 ml) was added Lawesson's reagent (2.48 g, 6.14 mmol, 1.5 eq) at RT and the reaction mixture was refluxed at 120°C for 40 min. After completion of reaction (monitored by TLC), the reaction mixture was quenched with sat. NaHCO_3 solution (100 ml) followed by extraction with EtOAc (2 x 100 ml). Combined organic layers were washed with water (100 ml), brine (100 ml), dried over anhydrous Na_2SO_4 and the solvent was evaporated to get the crude product which was purified by column chromatography (230-400 mesh silica gel; 20% EtOAc/hexane; R_f -value-0.4) to afford 8-(difluoromethyl)-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxaline-2(1H)-thione (0.6 g, 65%) as yellow solid.

[0222] Step7: To a stirring solution of 8-(difluoromethyl)-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxaline-2(1H)-thione (2.5 g, 9.6 mmol, 1 eq) in THF (30 ml) was drop wise added hydrazine hydrate (2.4 ml, 48 mmol, 5 eq) at 0°C . The reaction mixture then stirred for 16 h at RT. Then the reaction mixture was concentrated under reduced pressure to get the crude material. Then triethyl ortho acetate (20 ml) was added to the reaction mixture. The reaction mixture was stirred at 140°C for 16 h. After completion of reaction (monitored by TLC), reaction mixture was diluted with water (50 ml) and extracted by DCM (5x100 ml). The combined organic layer was washed by brine (100 ml). The organic layer was dried over Na_2SO_4 , concentrated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; 5% MeOH/DCM; R_f -value-0.4) to afford 9-(difluoromethyl)-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (2.0 g, 74%) as a off white solid.

[0223] Step8: A stirred solution of 9-(difluoromethyl)-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (2.0 g, 7.08 mmol, 1 eq) in DMF (30 ml) at -10°C was treated portion wise over 10 min with solid N-bromosuccinimide (1.3 g, 7.79 mmol, 1.05 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (300 ml) and organic layers were washed with water (5x50 ml), brine (50 ml), dried over anhydrous Na_2SO_4 and the solvents were evaporated under reduced pressure to get the crude product which was purified by silica gel (230-400) column chromatography (5% MeOH/DCM; R_f -value-0.3) to afford 8-bromo-9-(difluoromethyl)-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (1.5 g, 75%) as off white solid.

Synthesis of 8-bromo-9-cyclopropyl-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**intermediate B-45**):

[0224]



[0225] Step1: 2-Bromo-5-fluoro-phenylamine (50 g, 0.262 mol, 1 eq) was dissolved in acetic anhydride (25 mL) at 10°C and stirred at room temperature for 3 h. After completion of the reaction (monitored by TLC, 10% EtOAc/hexane), the thick reaction mass was diluted with n-hexane and filtered. The solid material was washed with n-hexane and dried under reduced pressure to afford N-(2-bromo-5-fluoro-phenyl)-acetamide (55 g, 90%) as an off-white solid.

[0226] Step2: To a stirred solution of N-(2-bromo-5-fluoro-phenyl)-acetamide (50 g, 0.259 mol, 1 eq) in a toluene:water

(1:1, 1 L) mixture was added tricyclohexyl phosphine (7.26 g, 0.0259 mol, 0.1 eq), followed by K_3PO_4 (192 g, 0.906 mol, 3.5 eq) at room temperature and the mixture was degassed with argon for 30 minutes. Cyclopropyl boronic acid (28.98 g, 0.336 mol, 1.3 eq) and subsequently $Pd(OAc)_2$ (2.9 g, 0.0129, 0.05 eq) were added and the reaction mixture was heated at 100°C for 16 h. After completion of the reaction (monitored by TLC, 20% EtOAc-Hexane, Rf 0.4), the reaction mixture was cooled to room temperature, diluted with EtOAc (1 L), washed with water (2 x 500 mL) and brine (500 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained crude residue was purified by column chromatography (100-200 mesh silica gel; 15-20% EtOAc/hexane as eluent) to afford N-(2-cyclopropyl-4-fluoro-phenyl)-acetamide (36 g, 86%) as a brownish solid.

[0227] Step3: A stirred suspension of N-(2-cyclopropyl-4-fluoro-phenyl)-acetamide (54 g, 0.279 mol, 1 eq) in aqueous HCl (2.1 L, 2M) was heated at 90°C for 16 h. After completion of the reaction (monitored by TLC, 20% EtOAc-Hexane, Rf 0.6), the reaction mixture was cooled to room temperature and basified to pH~13-14 with a NaOH solution (2M). This mixture was extracted with EtOAc (1 L), washed with water (2 x 500 mL) and brine (500 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford 2-cyclopropyl-5-fluoro-phenylamine (48 g, crude) as a dark brown liquid.

[0228] Step4: To a stirred solution of 2-cyclopropyl-5-fluoro-phenylamine (27 g, 0.18 mol, 1 eq) in DMF (480 mL) was added NBS (79.5 g, 0.447 mol, 2.5 eq) portion wise at -10°C and the resulting reaction mixture was stirred at 0°C for 1 h. After completion of the reaction (monitored by TLC, 20% EtOAc/hexane), the reaction mixture was diluted with water (1 L) and extracted with MTBE (2 x 750 mL). The combined organic layers were washed with cold brine (3 x 500 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained crude residue was purified by column chromatography (100-200 mesh silica gel; 5-10% EtOAc/hexane as eluent) to afford 2,4-dibromo-6-cyclopropyl-3-fluoro-phenylamine (22 g, 45% over two steps) as a brownish liquid.

[0229] Step5: To a stirred suspension of 2,4-dibromo-6-cyclopropyl-3-fluoro-phenylamine (20 g, 0.065 mol, 1 eq) in dry DMA (300 mL) was added 2-amino-2-methyl-propionic acid (13.35 g, 0.129 mol, 2 eq) followed by DBU (19.2 mL, 0.129 mol, 2 eq) at room temperature. The resulting reaction mixture was degassed with argon for 30 minutes, CuI (1.2 g, 0.006 mol, 0.1 eq) was added and the reaction mixture was heated at 140°C for 16 h. After complete consumption of the starting material (monitored by TLC, 30% EtOAc-Hexane, Rf 0.4), the reaction mixture was cooled to room temperature and filtered over a bed of celite, that was then washed with EtOAc (500 mL). The organic fraction was washed with water (2 x 750 mL) and brine (500 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained crude residue was purified by column chromatography (15-16% EtOAc-hexane) to afford 6-bromo-8-cyclopropyl-5-fluoro-3,3-dimethyl-3,4-dihydro-1H-quinoxalin-2-one (13 g, 64%) as a brown solid.

[0230] Step6: To a suspension of 6-bromo-8-cyclopropyl-5-fluoro-3,3-dimethyl-3,4-dihydro-1H-quinoxalin-2-one (10 g, 0.031 mol, 1 eq) in a dioxane:water (2:1, 200 mL) mixture was added Zn-powder (12.52 g, 0.191 mol, 6 eq), followed by ammonium chloride (10.25 g, 0.191 mol, 6 eq) at room temperature and the reaction was stirred at 100°C for 16 h. After completion of the reaction (monitored by TLC, 30% EtOAc-Hexane, Rf 0.4), the reaction mixture was cooled to room temperature and filtered through a celite bed, that was then washed with EtOAc (500 mL). The organic fraction was washed with water (2 x 750 mL) and brine (500 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained crude residue was purified by column chromatography using 15-20% EtOAc-hexane to afford 8-cyclopropyl-5-fluoro-3,3-dimethyl-3,4-dihydro-1H-quinoxalin-2-one (4.3 g, 59%) as a brown solid.


[0231] Step7: To a solution of 8-cyclopropyl-6-fluoro-3,3-dimethyl-3,4-dihydro-1H-quinoxalin-2-one (4.3 g, 18.376 mmol, 1 eq) in toluene (86 mL) was added Lawesson's reagent (11.16 g, 27.564 mmol, 1.5 eq) at room temperature and the reaction mixture was refluxed at 120°C for 1 h. After completion of the reaction (monitored by TLC in 20% EtOAc-Hexane, Rf 0.6), the reaction was cooled to room temperature and quenched with a sat. $NaHCO_3$ solution. The resulting aqueous fraction was extracted with EtOAc (3 x 200 mL) and the combined organic layers were washed with water (200 mL) and brine (200 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained crude residue was purified by column chromatography (100-200 mesh silica; 5-10% EtOAc/hexane) to afford 8-cyclopropyl-5-fluoro-3,3-dimethyl-3,4-dihydro-1H-quinoxaline-2-thione (4.2 g, 92%) as a yellow solid.

[0232] Step8: To a stirred solution of 8-cyclopropyl-5-fluoro-3,3-dimethyl-3,4-dihydro-1H-quinoxaline-2-thione (4 g, 16 mmol, 1 eq) in tetrahydrofuran (80 mL) was added hydrazine hydrate (2.35 mL, 48 mmol, 3 eq) drop wise at 0°C and the reaction mixture was stirred at room temperature for 16 h. The solvent was then evaporated and the residue was taken up in triethyl orthoacetate and heated at 140°C for 48 h. The reaction mixture was concentrated under reduced pressure and purified by column chromatography (100-200 mesh silica gel; 2% MeOH/ DCM as eluent) to afford 9-cyclopropyl-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (2.7 g, 59%) as a white solid. Step9: To a stirred solution of 9-cyclopropyl-6-fluoro-4,4-dimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (2.27 g, 8.345 mmol, 1 eq) in DMF (40 mL) was added NBS (1.33 g, 7.511 mmol, 0.9 eq) portion wise at -10°C and the resulting reaction mixture was stirred at 0°C for 1 h. After complete consumption of starting material (monitored by LCMS), the reaction mixture was diluted with water (20 mL) and extracted with EtOAc (2 x 30 mL). The combined organic layers were washed with water (2 x 50 mL) and brine, dried over anhydrous Na_2SO_4 and concentrated. The crude residue was combined with another batch (starting from 800 mg of 9-cyclopropyl-6-fluoro-4,4-dimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline

and following the same procedure as described above) and the total crude amount was then purified by column chromatography (100-200 mesh silica gel; 5% MeOH/DCM as eluent) to afford 8-bromo-9-cyclopropyl-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (2.2 g, 56 %) as an off-white solid.

Synthesis of 8-bromo-6-fluoro-N,1,4,4-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-amine (Intermediate B-46)

[0233]



[0234] Step1: To a stirred solution of 1,4-difluoro-2-nitrobenzene (120 g, 754.7 mmol, 1.0 eq) in DMF was added sodium bicarbonate (82.41 g, 981.07 mmol, 1.3 eq) and the mixture was stirred for 10 min. Then N-methyl benzyl amine (109.58 g, 905.6 mmol, 1.2 eq) was added and the mixture was stirred at RT for 16 h. After completion (monitored by TLC) the reaction mixture was diluted with EtOAc and washed with water. The extracted organic layer was washed with brine, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography (230-400 mesh silica gel, TLC system: EtOAc / hexane (2:8); R_f = 0.3) to give N-benzyl-4-fluoro-N-methyl-2-nitroaniline (110 g, 56%).

[0235] Step2: To an ice cold solution of N-benzyl-4-fluoro-N-methyl-2-nitroaniline (110 g, 423.07 mmol, 1.0 eq) in EtOH (800 mL) was added Tin (II) chloride (381.8 g, 1692.3 mmol, 4.0 eq). The reaction mixture was stirred for 10 min at RT. Concentrated HCl (250 ml) was drop wise added at 0°C. The reaction mixture was stirred at RT for 2 h. After completion (monitored by TLC) the reaction mixture was concentrated under vacuo. The residue was dissolved in cold water and basified with NaOH pellets upto (pH~12) and the aqueous part was diluted with EtOAc. The extracted organic layer was washed with brine, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel), (TLC system: EtOAc / hexane (2:8); R_f = 0.4) to give N¹-benzyl-4-fluoro-N¹-methylbenzene-1,2-diamine (100 g, 100%).

[0236] Step3: A stirred solution of N¹-benzyl-4-fluoro-N¹-methylbenzene-1,2-diamine (100 g, 434.7 mmol, 1 eq) in DMF at -10°C was treated portion wise over 10 min with solid N-bromosuccinimide (81.26 g, 456.51 mmol, 1.05 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (300 ml) and organic layers were washed with water (5x50 ml), brine (50 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure. Crude product was purified by silica gel (230-400) column chromatography (TLC system: EtOAc / hexane (2:8); R_f = 0.3) to afford N¹-benzyl-5-bromo-4-fluoro-N¹-methylbenzene-1,2-diamine (60 g, 45%) as color less liquid.

[0237] Step4: A stirred solution of N¹-benzyl-5-bromo-4-fluoro-N¹-methylbenzene-1,2-diamine (60 g, 194.17 mmol, 1 eq) in DMF at -10°C was treated portion wise over 10 min with solid N-bromosuccinimide (34.56 g, 194.17 mmol, 1.05 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (300 ml) and organic layers were washed with water (5x50 ml), brine (50 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure to get the crude product which

was purified by silica gel (230-400) column chromatography (TLC system: EtOAc / hexane (2:8); $R_f = 0.3$) to afford N¹-benzyl-3,5-dibromo-4-fluoro-N¹-methylbenzene-1,2-diamine (30 g, 40%) as off white solid.

[0238] Step5: A suspension of N¹-benzyl-3,5-dibromo-4-fluoro-N¹-methylbenzene-1,2-diamine (30 g, 77.31 mmol, 1 eq), 2-aminoisobutaric acid (15.94 g, 154.63 mmol, 2 eq), DBU (29.42 g, 193.29 mmol, 2.5 eq) and cuprous iodide (1.47 g, 7.73 mmol, 0.1 eq) in DMA (300 ml) in a round bottom flask were deoxygenated with Ar for 20 min. Reaction mixture was then stirred at 130°C for 16 h. After completion (monitored by TLC) the reaction mixture was diluted with EtOAc (500 ml) and water (5 x 100 ml), brine (100 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; (TLC system: EtOAc / hexane (2:8); $R_f = 0.2$) to afford 8-(benzyl(methyl)amino)-6-bromo-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (8.0 g, 26%) as brown solid.

[0239] Step6: To a solution of 8-(benzyl(methyl)amino)-6-bromo-5-fluoro-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (8 g, 20.40 mmol, 1 eq) in EtOH 10% Pd/C (2 g) was added. The reaction was performed in par autoclave under hydrogen atmosphere (200 psi) for 8 h. After completion of reaction the reaction mixture was filtered through a pad of celite it was washed with EtOAc. The filtrate was evaporated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; (TLC system: EtOAc / hexane (4:6); $R_f = 0.3$) to afford 5-fluoro-3,3-dimethyl-8-(methylamino)-3,4-dihydroquinoxalin-2(1H)-one (2.0 g, 22%) as white solid.


[0240] Step7: To a solution of 5-fluoro-3,3-dimethyl-8-(methylamino)-3,4-dihydroquinoxalin-2(1H)-one (2 g, 8.95 mmol, 1 eq) in toluene (30 ml) was added Lawesson's reagent (5.43 g, 13.43 mmol, 1.5 eq) at RT and the reaction mixture was heated to 120°C for 40 min. After completion of reaction (monitored by TLC), the reaction mixture was quenched with sat. NaHCO₃ solution (100 ml) followed by extraction with EtOAc (2 x 200 ml). Combined organic layers were washed with water (300 ml), brine (100 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product which was purified by column chromatography (230-400 mesh silica gel; (TLC system: EtOAc / hexane (4:6); $R_f = 0.3$) to afford 5-fluoro-3,3-dimethyl-8-(methylamino)-3,4-dihydroquinoxaline-2(1H)-thione (1.85 g, 86%) as yellow solid.

[0241] Step8: To a stirred solution of 5-fluoro-3,3-dimethyl-8-(methylamino)-3,4-dihydroquinoxaline-2(1H)-thione (1.85 g, 7.73 mmol, 1 eq) and acetic hydrazide (2 g, 27.05 mmol, 1 eq) in n-butanol (20 ml) catalytic amount of acetic acid was added and the mixture was heated to reflux for 16 h. After completion of reaction the solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; (TLC system: MeOH / DCM (1:9); $R_f = 0.2$) to afford 6-fluoro-N,1,4,4-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-amine (1.0 g, 50%) as liquid.

[0242] Step9: A stirred solution of 6-fluoro-N,1,4,4-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-amine (1.0 g, 3.82 mmol, 1 eq) in DMF (20 ml) at -10°C was treated portion wise over 10 min with solid N-bromosuccinimide (0.681 g, 3.82 mmol, 1.05 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (300 ml) and organic layers was washed with water (5x50 ml), brine (50 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; (TLC system: MeOH / DCM (1:9); $R_f = 0.4$) to afford 8-bromo-6-fluoro-N,1,4,4-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-amine (1.0 g, 77%) as liquid.

Synthesis of 8-bromo-6-chloro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**Intermediate B-47**)

[0243]



[0244] Step1: To an ice cold solution of 5-chloro-2-methoxyaniline (25 g, 158.62 mmol, 1.0 eq) in DMF a solution of N-bromosuccinimide (56.47 g, 317.25 mmol, 1.0 eq) was added in portion wise and the mixture was stirred at same temperature for 1 h. After completion (monitored by TLC) the reaction mixture was diluted with EtOAc and washed with water. The extracted organic layer was washed with brine, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel, TLC system: EtOAc / hexane (1:9); R_f = 0.5) to give 2,4-dibromo-3-chloro-6-methoxyaniline (30 g, 60%).

[0245] Step2: To a suspension of 2,4-dibromo-3-chloro-6-methoxyaniline (30 g, 95.12 mmol, 1 eq), 2-aminoisobutamic acid (24.92 g, 190.24 mmol, 2 eq), DBU (43.44 g, 285.36 mmol, 3 eq) and CuI (1.81 g, 9.51 mmol, 0.1 eq.) in DMA (300 ml) in a round bottom flask were deoxygenated with Ar for 20 min. Reaction mixture was then stirred at 130°C for 16 h. After completion (monitored by TLC) the reaction mixture was diluted with EtOAc (500 ml) and water (5 x 100 ml), brine (400 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; (TLC system: EtOAc / hexane (2:8); R_f = 0.2) to afford 6-bromo-5-chloro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (3.0 g, 10%) as brown solid.

[0246] Step3: A stirred solution of 6-bromo-5-chloro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (3 g, 9.38 mmol, 1 eq) in a mixture of water and 1,4-dioxane (1:1) zinc (8.94 g, 140.78 mmol, 15 eq) and ammonium chloride (7.53 g, 140.74 mmol, 15 eq) were added and the mixture was heated to reflux for 16 h. After completion (monitored by TLC) the reaction mixture was filtered through a pad of celite. The filtrate was diluted with EtOAc (500 ml) and water (5 x 100 ml), brine (100 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; (TLC system: EtOAc / hexane (2:8); R_f = 0.4) to afford 5-chloro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (2 g, 88%) as white solid.


[0247] Step4: To a solution of 5-chloro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (2.5 g, 10.38 mmol, 1 eq) in toluene (30 ml) was added Lawesson's reagent (6.30 g, 15.58 mmol, 1.5 eq.) at RT and the reaction mixture was heated to 120°C for 40 min. After completion of reaction (monitored by TLC), the reaction mixture was quenched with sat. NaHCO₃ solution (100 ml) followed by extraction with EtOAc (2 x 200 ml). Combined organic layers were washed with water (300 ml), brine (100 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product which was purified by column chromatography (230-400 mesh silica gel; (TLC system: EtOAc / hexane (3:7); R_f = 0.3) to afford 5-chloro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxaline-2(1H)-thione (2.1 g, 79%) as yellow solid.

[0248] Step5: To a stirred solution of 5-chloro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxaline-2(1H)-thione (2.1 g, 8.17 mmol, 1 eq) and acetic hydrazide (2.13 g, 28.75 mmol, 3.5 eq) were taken in n-butanol (20 ml) and catalytic amount of acetic acid was added and the mixture was heated to reflux for 16 h. After completion of reaction the solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; (TLC system: MeOH / DCM (1:9); R_f = 0.2) to afford 6-chloro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (1.2 g, 53%) as liquid.

[0249] Step6: A stirred solution of 6-chloro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (1.2 g, 4.30 mmol, 1 eq) in DMF (20 ml) at -10°C was treated portion wise over 10 min with solid N-bromosuccinimide (0.804 g, 4.52 mmol, 1.05 eq). Reaction mixture was allowed to warm to RT and stirred for 1.5 h. After completion of reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (300 ml) and organic layers was washed with water (5x50 ml), brine (50 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; (TLC system: MeOH / DCM (1:9); R_f = 0.4) to afford 8-bromo-6-chloro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.9 g, 59%) as white solid.

Synthesis of 8-bromo-6,7-difluoro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**Intermediate B-49**)

[0250]



[0251] Step1: To a stirred solution of 4,5-difluoro-2-nitrophenol (20 g, 114.226 mmol, 1 eq) in DMF (100 ml), potassium carbonate (47.2 g, 342.68 mmol, 3 eq) and iodo methane (21.34 ml, 342.68 mmol, 3 eq) were added into the reaction mixture at RT for 4 h. After completion (monitored by TLC) the reaction mixture was diluted with EtOAc and washed with water. The extracted organic layer was washed with brine, dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel, TLC system: EtOAc/Hexane (2:8); R_f = 0.6) to give 1,2-difluoro-4-methoxy-5-nitrobenzene (20 g, 93%).

[0252] Step2: To a stirred solution of 1,2-difluoro-4-methoxy-5-nitrobenzene (20.0 g, 105.75 mmol, 1 eq) in MeOH (200 ml), Pd/C (10.0 g, 10% moist) was added and the reaction mixture was stirred with hydrogen balloon for 2 h at RT. After completion (monitored by TLC, TLC system 20% EtOAc in Hexane, R_f -0.4), the reaction mixture was filtered through celite bed and washed 2-3 times with MeOH. The filtrate was concentrated to get the desired 4,5-difluoro-2-methoxyaniline (16.0 g, 95%).

[0253] Step3: To a stirred solution of 4,5-difluoro-2-methoxyaniline (8.0 g, 50.273 mmol, 1 eq) in MeOH:DCM (50 ml:100 ml) was added benzyltrimethylammonium tribromide (8.82 g, 60.328 mmol, 1.2 eq) portion wise at RT. Resulting reaction mixture was stirred at RT for 12 h. After completion of starting material reaction mixture was concentrated to afford crude product which was purified by column chromatography (230-400 mesh silica gel; 15% EtOAc/Hexane; R_f -value-0.4) to afford 2-bromo-3,4-difluoro-6-methoxyaniline (0.5 g, 4%) as yellow solid.

[0254] Step4: To a stirred solution of 2-bromo-3,4-difluoro-6-methoxyaniline (0.413 g, 1.735 mmol, 1 eq) in dry DMSO (10 ml) was added 2-amino-2-methyl-propionic acid (0.536 g, 5.205 mmol, 3.0 eq) followed by K_3PO_4 (0.919 g, 4.337 mmol, 2.5 eq) at RT. Resulting reaction mixture was degassed with Ar for 10 min, then CuI (0.033 g, 0.1735 mmol, 0.1 eq) was added and reaction mixture was heated at 140°C for 12 h. After completion of the starting material reaction mixture was cooled to RT and filtered through celite. Celite bed was washed with EtOAc (500 ml). Resulting filtrate was washed with water (2 x 50 ml), followed by brine (50 ml), dried over anhydrous Na_2SO_4 and the solvent was evaporated under reduced pressure to afford crude product which was purified by column chromatography (100-200 mesh silica gel; 20% EtOAc/hexane; R_f -value-0.2) to afford 5,6-difluoro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (0.3 g, 71%) as brown solid.

[0255] Step5: To a solution of 5,6-difluoro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxalin-2(1H)-one (0.3 g, 1.238 mmol, 1 eq) in toluene (50 ml) was added Lawesson's reagent (0.65 g, 1.61 mmol, 1.5 eq) at RT and the reaction mixture was heated to 120°C for 1 h. After completion of starting material, the reaction mass was cooled to RT and quenched with saturated NaHCO_3 solution (50 ml). The organic layer was separated and aqueous layer was extracted with EtOAc (50 ml). Combined organic layers were washed with water (50 ml), followed by brine (50 ml), dried over anhydrous Na_2SO_4 and the solvent was evaporated to afford crude product which was purified by column chromatography (230-400 mesh silica gel; 20% EtOAc/hexane; R_f -value-0.6) to afford 5,6-difluoro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxaline-2(1H)-thione (0.3 g, 94%) as yellow solid.


[0256] Step6: To a solution of 5,6-difluoro-8-methoxy-3,3-dimethyl-3,4-dihydroquinoxaline-2(1H)-thione (0.1 g, 0.3871 mmol, 1 eq) in n-BuOH (10 ml) was added acetic hydrazide (0.094 g, 1.277 mmol, 3.3 eq) followed by addition of acetic acid (1 ml) and then the reaction mixture was stirred at 140°C for 12 h. After completion of reaction (monitored by TLC) reaction mixture was diluted with water (10 ml) and extracted with EtOAc (2x20 ml). Combined organic layer was washed with water (20 ml), brine (20 ml), dried over anhydrous Na_2SO_4 and the solvent was evaporated to get the crude product which was purified by column chromatography using 5% MeOH in DCM as an eluting solvent and 230-400 silica gel to afford 6,7-difluoro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.05 g, 46%) as off white solid.

[0257] Step7: To a stirred solution of 6,7-difluoro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.8 g, 2.854 mmol, 1 eq) in DMF (20 ml) at 0°C was treated portion wise over 10 min with solid N-bromosuccinimide (0.6 g, 3.425 mmol, 1.2 eq). Reaction mixture was allowed to warm to RT and stirred for 30 min. After completion of

reaction (monitored by LCMS), reaction mixture was diluted with EtOAc (50 ml) and organic layers were washed with water (3x50 ml), brine (100 ml), dried over anhydrous Na_2SO_4 and the solvents were evaporated under reduced pressure to get crude product which was purified by column chromatography using 1.5% MeOH in DCM as an eluting solvent and 230-400 silica gel to afford 8-bromo-6,7-difluoro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.25 g, 42%) as off white solid.

Synthesis of 8-bromo-6,7-difluoro-1,4,4-trimethyl-9-(trifluoromethyl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**Intermediate B-50**)

[0258]



[0259] Step1: A solution of 1-bromo-4,5-difluoro-2-nitrobenzene (1.0 g, 4.2 mmol, 1 eq) in DMF (15 ml) was degassed with Ar for 20 min followed by addition of CuBr (0.06 g, 0.42 mmol, 0.1 eq) and methyl 2,2-difluoro-2-(fluorosulfonyl)acetate (0.88 ml, 6.93 mmol, 1.65 eq). The reaction mixture was stirred at 80°C for 16 h. After completion of reaction (monitored by TLC), reaction mixture was diluted with diethyl ether and washed with ice cold water (2-3 times). Combined organic layer was evaporated to get the crude product, which was purified by column chromatography using 100-200 silica gel and hexane as eluting solvent to afford 1,2-difluoro-4-nitro-5-(trifluoromethyl)benzene (0.6 g, 45%) as light yellow liquid.

[0260] Step2: A solution of 1,2-difluoro-4-nitro-5-(trifluoromethyl)benzene (0.25 g, 1.1 mmol, 1 eq.) in EtOH (10 ml) was cooled to 0°C. $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ was added portionwise (0.994 g, 4.4 mmol, 4.0 eq.) followed by drop wise addition of conc. HCl (2.5 ml). The reaction mixture was stirred at RT for 2 h. After completion of reaction (monitored by TLC), reaction mixture was basified with 6N NaOH solution and diluted with DCM. The organic layer was washed with water (2-3 times). Combined organic layer was evaporated to get the desired 4,5-difluoro-2-(trifluoromethyl)aniline (0.21 g, 93%) as light yellow liquid compound.

[0261] Step3: A solution of 4,5-difluoro-2-(trifluoromethyl)aniline (0.20 g, 1.01 mmol, 1 eq.) in DMF (10 ml) was cooled to 0°C. NBS was added portionwise (0.217 g, 1.21 mmol, 1.2 eq.). The reaction mixture was gradually allowed to RT for 5 h. After completion of reaction (monitored by TLC), reaction mixture was diluted with diethylether. The organic layer was washed with ice cold water (2-3 times). Combined organic layer was evaporated to get the crude product which was purified by column chromatography using 100-200 silica gel and hexane as eluting solvent to afford 2-bromo-3,4-difluoro-6-(trifluoromethyl)aniline (0.19 g, 63%) as light yellow liquid compound.

[0262] Step4: A solution of 2-bromo-3,4-difluoro-6-(trifluoromethyl)aniline (5.0 g, 18.11 mmol, 1 eq.), 2-aminoisobutamic acid (3.76 g, 36.23 mmol, 2 eq.), K_3PO_4 (9.6 g, 45.28 mmol, 2.5 eq.) in dry DMSO (86 ml) taken in a sealed tube were deoxygenated with Ar. CuI (0.34 g, 1.8 mmol, 0.1 eq.) was added and again deoxygenated for 5 min. Reaction mixture was then stirred at 120°C for 6 h. After completion of the reaction, (monitored by TLC, 20% EtOAc in hexane, R_f -0.2), it was filtered through celite bed and washed by EtOAc (100 ml), washed with water (3x100 ml), brine (100 ml), dried over anhydrous Na_2SO_4 and evaporated under reduced pressure to get the crude product which was purified by column chromatography (230-400 mesh silica gel; 20% EtOAc/hexane; R_f -value-0.2) to afford 5,6-difluoro-3,3-dimethyl-8-(trifluoromethyl)-3,4-dihydroquinoxalin-2(1H)-one (0.7g, 14%) as brown solid.

[0263] Step5: To a solution of 5,6-difluoro-3,3-dimethyl-8-(trifluoromethyl)-3,4-dihydroquinoxalin-2(1H)-one (0.7 g, 2.5 mmol, 1 eq.) in toluene (15 ml) was added Lawesson's reagent (1.5 g, 3.75 mmol, 1.5 eq.) at RT and the reaction mixture was refluxed at 120°C for 1.5 h. After completion of reaction (monitored by TLC, 20% EtOAc in hexane, R_f -0.5), the reaction mixture was quenched with sat. NaHCO_3 solution (100 ml) followed by extraction with EtOAc (2 x 100 ml). Combined organic layers were washed with water (100 ml), brine (100 ml), dried over anhydrous Na_2SO_4 and the solvent was evaporated to get the crude product which was purified by column chromatography using 230-400 mesh silica gel and 10% EtOAc in hexane as an eluting solvent to afford 5,6-difluoro-3,3-dimethyl-8-(trifluoromethyl)-3,4-dihydroquinoxaline-2(1H)-thione (0.7g, 94%) as yellow solid.


[0264] Step6: To a solution of 5,6-difluoro-3,3-dimethyl-8-(trifluoromethyl)-3,4-dihydroquinoxaline-2(1H)-thione (0.7 g, 2.36 mmol, 1 eq.) in THF (30 ml) was added hydrazine hydrate (0.59 g, 11.82 mmol, 5.0 eq.) at RT and the reaction mixture was stirred at RT for 1 h. After completion of reaction (monitored by TLC, 5% MeOH-DCM, Rf-0.5), the reaction mixture was concentrated. Reaction mixture was dissolved in THF and cooled to 0°C. TEA (1.6 ml, 11.82 mmol, 5.0 eq.) and acetyl chloride (0.5 ml, 7.09 mmol, 3.0 eq.) was added and stirred at RT for 1 h. After completion of reaction (monitored by TLC, 5% MeOH-DCM, Rf-0.5), reaction mixture was diluted with EtOAc. Combined organic layers were washed with water, brine, dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude N'-(5,6-difluoro-3,3-dimethyl-8-(trifluoromethyl)-3,4-dihydroquinoxalin-2(1H)-ylidene)acetohydrazide which was taken to the next step without further purification.

[0265] Step7: A solution of N'-(5,6-difluoro-3,3-dimethyl-8-(trifluoromethyl)-3,4-dihydroquinoxalin-2(1H)-ylidene)acetohydrazide (2.36 mmol, 1 eq) in POCl₃ (15 ml) was cooled to 0°C. TEA (0.33 ml, 2.36 mmol, 1.0 eq) was added and the mixture was stirred at the same temperature for 10 min and then at reflux for 6 h. After completion of reaction (monitored by TLC, 50% Acetone-Hexane, Rf-0.5), the reaction mixture was concentrated. Reaction mixture was cooled to 0°C and basified using aqueous ammonia solution and diluted with EtOAc. Combined organic layers were washed with water, brine, dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product which was purified by column chromatography using 230-400 mesh silica gel and 50% acetone in hexane as an eluting solvent to afford 6,7-difluoro-1,4,4-trimethyl-9-(trifluoromethyl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.5 g, 66%) as brown solid.

[0266] Step8: A solution of 6,7-difluoro-1,4,4-trimethyl-9-(trifluoromethyl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.5 g, 1.57 mmol, 1 eq) in DMF (15 ml) was cooled to 0°C. NBS (0.559 g, 6.28 mmol, 2.0 eq.) was added and the mixture was stirred at RT for 10 min and then at reflux for 2 h. After completion of reaction (monitored by TLC, 50% Acetone-Hexane, Rf-0.5), the reaction mixture was diluted with EtOAc. Combined organic layers were washed with ice cold water, brine, dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product which was purified by column chromatography using 230-400 mesh silica gel and 50% acetone in hexane as an eluting solvent to afford 8-bromo-6,7-difluoro-1,4,4-trimethyl-9-(trifluoromethyl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.45 g, 72%) as brown solid.

Synthesis of 8-bromo-6-chloro-7-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**Intermediate B-51**)

[0267]



[0268] Step1: To a stirred solution of NiCl₂·6H₂O (12.54 g, 0.0527 mol, 1 eq) in MeOH (526 mL) was added NaBH₄ (5.97 g, 0.158 mol, 3 eq) portion wise at 0°C and the resulting suspension was stirred at 0°C for 1 h. Then, 1-chloro-2-fluoro-4-methyl-5-nitro-benzene (20 g, 0.1055 mol, 2 eq) was added, followed by further addition of NaBH₄ (13.95 g, 0.369 mol, 7 eq) portion wise and the reaction mixture was stirred at 0°C for 2h. After complete consumption of starting material (monitored by TLC), the reaction mixture was concentrated and the obtained residue was diluted with 1N HCl (150 mL) followed by aqueous ammonia (150 mL) and EtOAc (200 mL). This mixture was stirred at room temperature for 15 min, filtered over a celite bed, and the obtained aqueous filtrate was extracted with EtOAc (2 x 250 mL). The combined organic fractions were washed with water (350 mL) and brine (250 mL), dried over anhydrous Na₂SO₄ and concentrated to afford 5-chloro-4-fluoro-2-methyl-phenylamine (14 g, 83%) as an off-white solid.

[0269] Step2: To a solution of 5-chloro-4-fluoro-2-methyl-phenylamine (14 g, 0.087 mol, 1 eq) in DCM (280 mL) was added a solution of Br₂ (4.97 mL, 0.098 mol, 1.1 eq) in DCM (70 mL) drop-wise at 0°C and the resulting reaction mixture was stirred at 0°C for 2 h. After complete consumption of starting material (monitored by TLC in 10% EtOAc-Hexane),

the reaction mixture was diluted with water (300 mL) and extracted with DCM (2 x 150 mL). The combined organic layers were washed with water (2 x 250 mL), a saturated sodium thiosulfate solution (2 x 250 mL) and brine (250 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained crude residue was purified by column chromatography (100-200 mesh silica gel; 5% EtOAc/hexane as eluent) to yield 2-bromo-3-chloro-4-fluoro-6-methyl-phenylamine (11.2 g, 53%) as a brown solid.

[0270] Step3: To a stirred suspension of 2-bromo-3-chloro-4-fluoro-6-methyl-phenylamine (18 g, 0.075 mol, 1 eq) in dry DMAc (270 mL) was added 2-amino-2-methyl-propionic acid (15.56 g, 0.15 mol, 2 eq), followed by DBU (22.57 mL, 0.15 mol, 2 eq) at room temperature and the resulting reaction mixture was degassed with nitrogen for 30 minutes. Then, CuI (1.43 g, 0.0075 mol, 0.1 eq) was added and the reaction mixture was heated at 140°C for 16 h. After complete consumption of the starting material (monitored by TLC, 20% EtOAc-Hexane, Rf 0.4), the reaction mixture was cooled to room temperature and poured into ice-cold water (1 L). The resulting aqueous mixture was extracted with EtOAc (2 x 400 mL) and the combined organic fractions were washed with water (2 x 500 mL) and brine (400 mL), dried over anhydrous Na_2SO_4 and evaporated under reduced pressure. The obtained crude residue was triturated with hexane to afford 5-chloro-6-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxalin-2-one (8 g, 44%) as a brown liquid.


[0271] Step4: To a solution of 5-chloro-6-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxalin-2-one (12 g, 0.049 mol, 1 eq) in toluene (150 mL) was added Lawesson's reagent (29.97 g, 0.074 mol, 1.5 eq) at room temperature and the reaction mixture was refluxed at 120°C for 2 h. After complete consumption of starting material (monitored by TLC in 20% EtOAc-Hexane, Rf 0.6), the reaction mixture was quenched with a sat. NaHCO_3 solution (200 mL) and the resulting aqueous mixture was extracted with EtOAc (2 x 350 mL). The combined organic layers were washed with water (200 mL) and brine (200 mL), dried over anhydrous Na_2SO_4 and evaporated under reduced pressure. The obtained crude residue was purified by column chromatography (100-200 mesh silica gel; 10% EtOAc/hexane as eluent) to afford 5-chloro-6-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxaline-2-thione (128 g, 93%) as a yellow solid.

[0272] Step5: To a stirring solution of 5-chloro-6-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxaline-2-thione (11 g, 0.042 mol, 1 eq) in tetrahydrofuran (275 mL) was added hydrazine hydrate (6.38 g, 0.127 mol, 3 eq) drop wise at 0°C and the resulting reaction mixture was stirred at room temperature for 16 h. After formation of imine intermediate, the reaction mixture was concentrated and azeotroped with toluene twice. The obtained residue was dissolved in triethyl-orthoacetate (120 mL) and heated to reflux for 24 h. After complete consumption of starting material (monitored by LCMS), the reaction mixture was concentrated and the obtained crude residue was purified by column chromatography (230-400 mesh silica gel; 5% MeOH/DCM as eluent) to afford 6-chloro-7-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (8.5 g, 71%) as an off-white solid.

[0273] Step6: To a solution of 6-chloro-7-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (8 g, 0.028 mol, 1 eq) in DMF (160 mL) was added NBS (5.07 g, 0.028 mol, 1 eq) portion wise at 0°C and the resulting reaction mixture was stirred at room temperature for 3 h. After complete consumption of starting material (monitored by LCMS), the reaction mixture was diluted with water (500 mL) and extracted with EtOAc (2 x 250 mL). The combined organic layers were washed with water (2 x 250 mL) and brine (250 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained crude residue was purified by column chromatography (230-400 mesh silica gel; 5% MeOH/DCM as eluent) to afford 8-bromo-6-chloro-7-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (6.2 g, 60%) as an off-white solid.

Synthesis of 8-bromo-7-chloro-6-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**Intermediate B-53**)

[0274]



[0275] Step1: To a stirred solution of 1-chloro-2-fluoro-5-methyl-4-nitro-benzene (10.0 g, 52.91 mmol, 1 eq) in an EtOH-water (1:1, 120 mL) mixture was added a conc. aqueous HCl solution (6 mL), followed by addition of iron powder

(10.3 g, 185.18 mmol, 3.5 eq) at room temperature and the reaction mixture was stirred for 16 h at 110°C. After completion of the reaction (monitored by TLC, 20% EtOAc/hexane, Rf 0.3), the reaction mixture was filtered through a celite bed and washed with EtOAc (200 mL x 3). The combined organic fractions were concentrated under reduced pressure, diluted with EtOAc (500 mL), washed with water (100 mL x 2) and brine (200 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The obtained crude residue was purified by column chromatography (100-200 mesh silica gel; 20% EtOAc/hexane as eluent) to afford 4-chloro-5-fluoro-2-methyl-phenylamine (7.5 g, 89%) as a light brown solid.

[0276] Step2: To a stirred solution of 4-chloro-5-fluoro-2-methyl-phenylamine (5.5 g, 34.81 mmol, 1 eq) in DCM (150 mL) was added a solution of Br₂ (1.79 mL, 34.81 mmol, 1 eq) in DCM (60 mL) drop wise at 0°C and the resulting reaction mixture was stirred at 0°C for 2 h. After complete consumption of starting material (monitored by TLC in 20% EtOAc-Hexane, Rf 0.6), the reaction mixture was quenched with a saturated NaHCO₃ solution (400 mL) and the organic layer was separated. The aqueous fraction was extracted and the combined organic layers were washed with water (2 x 300 mL) and brine (200 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The obtained crude residue was purified by column chromatography (100-200 mesh silica gel; 10% EtOAc/hexane as eluent) to afford 2-bromo-4-chloro-3-fluoro-6-methylaniline (4.5 g, 54%) as a brown solid.

[0277] Step3: To a stirred solution of 2-bromo-4-chloro-3-fluoro-6-methylaniline (5 g, 21.008 mmol, 1 eq) in dry DMAc (100 mL) were added 2-amino-2-methyl-propionic acid (4.4 g, 42.016 mmol, 2 eq) and DBU (5.46 mL, 42.016 mmol, 2 eq) at room temperature and the resulting reaction mixture was degassed with nitrogen for 30 minutes. CuI (400 mg, 2.1006 mmol, 0.1 eq) was added to this reaction mixture, which was heated at 140°C for 16 h. After complete consumption of the starting material (monitored by TLC, 20% EtOAc-Hexane, Rf 0.3), the reaction mixture was cooled to room temperature and filtered over a celite bed, that was then washed with EtOAc (2 x 300 mL). The resulting filtrate was poured into ice-cold water (200 mL) and extracted with EtOAc (2 x 400 mL). The combined organic fractions were washed with cold water (3 x 100 mL) and brine (100 mL), dried over anhydrous Na₂SO₄ and evaporated under reduced pressure. The crude residue was purified by chromatography (100-200 mesh silica gel; 20% EtOAc/hexane as eluent) to afford 6-chloro-5-fluoro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-one (28 g, 69%) as an off-white solid.


[0278] Step4: To a stirred solution of 6-chloro-5-fluoro-3,3,8-trimethyl-3,4-dihydroquinoxalin-2(1H)-one (12 g, 49.58 mmol, 1 eq) in toluene (240 mL) was added Lawesson's reagent (30.08 g, 74.38 mmol, 1.5 eq) at room temperature and the reaction mixture was refluxed at 120°C for 2 h. After consumption of starting material (monitored by TLC in 20% EtOAc-Hexane, Rf 0.6), the reaction mixture was concentrated under reduced pressure. The obtained solid residue was quenched with sat. NaHCO₃ solution (500 mL) and the resulting aqueous phase was extracted with EtOAc (3 x 300 mL), after which the combined organic layers were washed with water (400 mL) and brine (200 mL), dried over anhydrous Na₂SO₄ and evaporated under reduced pressure. The obtained crude residue was purified by column chromatography (100-200 mesh silica gel; 10% EtOAc/hexane as eluent) to afford 6-chloro-5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxaline-2-thione (11.5 g, 90%) as a yellow solid.

[0279] Step5: To a stirred solution of 6-chloro-5-fluoro-3,3,8-trimethyl-3,4-dihydro-1H-quinoxaline-2-thione (8 g, 31.007 mmol, 1 eq) in tetrahydrofuran (200 mL) was drop wise added hydrazine hydrate (6.209 g, 124 mmol, 4 eq) at 0°C and the reaction mixture was stirred at room temperature for 16 h. After complete consumption of starting material (monitored by TLC in 5% MeOH-DCM, Rf 0.4), the reaction mixture was concentrated to afford crude 7-chloro-8-fluoro-3-hydrazineylidene-2,2,5-trimethyl-1,2,3,4-tetrahydroquinoxaline (8 g), which was used in the next step without purification.

[0280] Step6: To crude 7-chloro-8-fluoro-3-hydrazineylidene-2,2,5-trimethyl-1,2,3,4-tetrahydroquinoxaline (8 g, 34.06 mmol, 1 eq), triethyl orthoacetate (240 mL) was added at room temperature, and the solution was stirred at 140°C for 48 h. After consumption of starting material (monitored by LCMS), the reaction mixture was concentrated and purified by column chromatography (100-200 mesh silica gel; 2 % MeOH/ DCM as eluent) to afford 7-chloro-6-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (6 g, 69% over two steps) as a white solid.


[0281] Step7: To a stirred solution of 7-chloro-6-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (3 g, 10.686 mmol, 1 eq) in DMF (90 mL) was added NBS (1.99 g, 11.22 mmol, 1.05 eq) portion wise at 0°C and the resulting reaction mixture was stirred at 0°C for 1 h. After consumption of starting material (monitored by LCMS), the reaction mixture was diluted with water (60 mL) and extracted with EtOAc (2 x 100 mL). The combined organic layers were washed with water (2 x 60 mL) and brine, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. Three batches (3 x 3 g) were done in parallel and the combined crude material was purified by column chromatography (100-200 mesh silica gel; 5% MeOH/DCM as eluent) to afford 8-bromo-7-chloro-6-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (7.6 g, 66%) as a yellow solid.

[0282] Any of intermediates-A can be coupled to any of intermediates-B in standard chemical reactions which are known to the person skilled in the art, e.g. those as described herein below.

Example 38: 8-(3-cyclopropyl-5-fluoro-1H-indol-7-yl)-6-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline**[0283]**

[0284] To a solution of **intermediate B-7** (0.30 g, 0.923 mmol, 1 eq) in toluene:EtOH (2:1) (9 ml) were added 10 % Na_2CO_3 (1.0 ml) solution and 3-cyclopropyl-5-fluoro-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**intermediate A-16**) (0.361 g, 1.2 mmol, 1.3 eq) in a sealed tube. The solution was degassed with Ar for 20 min followed by addition of $\text{Pd}(\text{PPh}_3)_4$ (0.053 g, 0.0461 mmol, 0.05 eq). The reaction mixture was refluxed at 110°C for 16 h. After completion of reaction, the reaction mixture was evaporated to dryness and the residue was diluted with EtOAc (50 ml). The organic layer was washed with water (2 x 30 ml), brine (30 ml), dried over anhydrous Na_2SO_4 and the solvents were evaporated to get the crude product which was purified by silica gel (230-400) column chromatography (5% MeOH/DCM; R_f -value-0.4) to afford 8-(3-cyclopropyl-5-fluoro-1H-indol-7-yl)-6-fluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.140 g, 36%) as off white solid.


[0285] ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 10.64 (s, 1H), 7.36 (d, 1H), 7.17 (d, 1H), 7.08 (s, 1H), 6.9 (d, 1H), 6.49 (s, 1H), 2.5 (s, 3H), 1.98 (s, 3H), 1.93-1.9 (m, 1H), 1.53-1.48 (m, 6H), 0.85 (d, 2H), 0.60 (d, 2H).

Example 49: 9-ethyl-6-fluoro-1,4,4-trimethyl-8-(1-(methylsulfonyl)-1H-indol-4-yl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline**[0286]**

[0287] To a solution of **intermediate B-9** (0.25 g, 0.737 mmol, 1 eq) in 1,4-dioxan and water (15 ml and 1.5 ml) were added CsF (0.336 g, 2.21 mmol, 3 eq) and **intermediate A-2** (0.344 g, 1.1 mmol, 1.5 eq) in a sealed tube. The solution was degassed with Ar for 20 min followed by addition of $\text{Pd}(\text{PPh}_3)_4$ (0.043 g, 0.0368 mmol, 0.05 eq). The reaction mixture was stirred at 90°C for 16 h. After completion of reaction (monitored by LCMS), the solvents were evaporated to dryness and the residue was diluted with EtOAc (50 ml). The organic layer was washed with water (2 x 20 ml), brine (20 ml), dried over anhydrous Na_2SO_4 and the solvent was evaporated to get the crude product which was purified by silica gel (230-400) column chromatography (5% MeOH/DCM; R_f -value-0.4) to afford 9-ethyl-6-fluoro-1,4,4-trimethyl-8-(1-(methylsulfonyl)-1H-indol-4-yl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.08 g, 24%) as off white solid.

[0288] ^1H NMR (400 MHz, $\text{DMSO}-d_6$, at 100°C): 7.92 (d, 1H), 7.62 (s, 1H), 7.48 (t, 1H), 7.34 (d, 1H), 7.08 (d, 1H), 6.58 (s, 1H), 6.21 (s, 1H), 3.43 (s, 3H), 2.68-2.66(m, 2H), 2.53 (s, 3H), 1.57 (s, 6H), 0.51 (t).

Example 58: 8-(1-(2,2-difluoroethyl)-6-fluoro-1H-indol-4-yl)-6-fluoro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline**[0289]**




[0290] To a solution of **intermediate B-10** (0.1 g, 0.294 mmol, 1 eq) and **intermediate A-25** (0.115 g, 0.353 mmol, 1.2 eq) in t-amyl alcohol (3 ml) / dioxane (3 ml) / water (0.3 ml) mixture was added K_2CO_3 (0.122 g, 0.882 mmol, 3 eq). The solution was then degassed (N_2) for 10 min followed by addition of Attaphos catalyst (0.011 g, 0.0147 mmol, 0.05 eq). The reaction mixture was then heated at 100°C for 16 h. After completion (LCMS), reaction mixture was filtered through celite pad. The filtrate was concentrated under reduced pressure to get crude product which was purified by preparative HPLC to afford 8-(1-(2,2-difluoroethyl)-6-fluoro-1H-indol-4-yl)-6-fluoro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.071 g, 50%).

[0291] 1H -NMR (400 MHz; $DMSO-d_6$): δ 7.51 (dd, 1H), 7.41 (d, 1H), 7.25 (d, 1H), 7.05 (dd, 1H), 6.67 (s, 1H), 6.54-6.27 (m, 2H), 4.71 (td, 1H), 3.16 (s, 3H), 2.55 (d, 3H), 1.55 (s, 3H).

Example 446: 8-(1-(ethylsulfonyl)-6-fluoro-1H-indol-4-yl)-6,7-difluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline

[0292]




[0293] To a solution of 8-bromo-6,7-difluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**intermediate B-38**) (0.6 g, 1.75 mmol, 1.0 eq) in t-amyl alcohol:1,4-dioxane: H_2O (3:2:1) (100 ml), K_2CO_3 (0.725 g, 5.25 mmol, 3 eq) was added at RT. The solution was degassed with Ar for 20 min followed by addition of 1-(ethylsulfonyl)-6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**intermediate A-8**) (0.930 g, 2.63 mmol, 1.2 eq) and Attaphos (0.062 g, 0.0875 mmol, 0.05 eq). The reaction mixture was then placed to pre-heated oil bath at 90°C for 2 h. After completion of reaction (monitored by TLC & LCMS), reaction mixture was diluted with water (100 mL) and extracted with EtOAc (2x250 mL). The organic layer was washed with brine (75 mL), dried over Na_2SO_4 , filtered and the solvent was evaporated under reduced pressure to get the crude product which was initially purified by column chromatography using MeOH-DCM as eluent. After that it was again repurified in combiflash column chromatography using acetone-hexane as eluent. The product was then washed with ether to afford pure 8-(1-(ethylsulfonyl)-6-fluoro-1H-indol-4-yl)-6,7-difluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.32 g, 38%) as white solid.

[0294] 1H NMR (400 MHz, $DMSO-d_6$): δ 7.71 (d, 1H), 7.66 (d, 1H), 7.37 (d, 1H), 6.93 (s, 1H), 6.59 (d, 1H), 3.75 (q, 2H), 2.45 (s, 3H), 1.97 (s, 3H), 1.57 (s, 3H), 1.51 (s, 3H), 1.11 (t, 3H).

Example 448: 6,7-difluoro-1,4,4,9-tetramethyl-8-(1-(methylsulfonyl)-1H-indol-4-yl)-4,5-dihydro-[1,2]triazolo[4,3-a]quinoxaline

[0295]




[0296] To a solution of 8-bromo-6,7-difluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**inter-**

mediate B-38) (0.05 g, 0.146 mmol, 1 eq) in 1,4-dioxan (9 ml) were added CsF (0.067 g, 0.438 mmol, 3 eq) and 1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**Intermediate A-12**) (0.07 g, 0.219 mmol, 1.5 eq) in a sealed tube. The solution was degassed with Ar for 20 min followed by addition of Pd(PPh₃)₄ (0.013 g, 0.0073 mmol, 0.05 eq). The reaction mixture was refluxed at 110°C for 16 h. After completion of reaction (monitored by LCMS), reaction mixture was evaporated to dryness and the residue was diluted with EtOAc (50 ml). The organic layer was washed with water (2x20 ml), brine (20 ml), dried over anhydrous Na₂SO₄ and the solvent was evaporated to get the crude product, which was purified by preparative HPLC (5% MeOH/DCM; R_f-value-0.4) to afford 6,7-difluoro-1,4,4,9-tetramethyl-8-(1-(methylsulfonyl)-1H-indol-4-yl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.04 g, 61%) as a white solid.

[0297] ¹H NMR (DMSO-d₆) δ= 7.94 (d, 1H), 7.65 (d, 1H), 7.51 (t, 1H), 7.35 (d, 1H), 6.88 (s, 1H), 6.57 (d, 1H), 3.53 (s, 3H), 2.44 (s, 3H), 1.95 (s, 3H), 1.58 (s, 3H), 1.49 (s, 3H).

Example 450: 8-(1-cyclopropyl-1H-indol-4-yl)-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline-9-carbonitrile

[0298]




[0299] To a solution of 8-bromo-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline-9-carbonitrile (**Intermediate B-39**) (0.130 g, 0.388 mmol, 1.00 eq) in dioxane:water (10:1, 10 ml) were added CsF (0.176 g, 1.164 mmol, 3.00 eq) and 1-cyclopropyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**Intermediate A-44**) (0.219 g, 0.776 mmol, 2.00 eq). The solution was degassed with Ar for 20 min followed by the addition of Pd(PPh₃)₄ (0.022 g, 0.019 mmol, 0.05 eq). The reaction mixture was then heated to 90°C for 16 h. After completion of the reaction (monitored by TLC) the reaction mixture was filtered through a celite pad and the celite pad was washed with EtOAc. The combined organic layers were evaporated under reduced pressure to get the crude compound, which was purified by prep. HPLC to afford 8-(1-cyclopropyl-1H-indol-4-yl)-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline-9-carbonitrile (0.045 g, 28.3%) as a white solid.

[0300] ¹H NMR (DMSO-d₆) δ= 7.71 (d, 1H), 7.52-7.50 (m, 1H), 7.47 (d, 1H), 7.34-7.30 (m, 1H), 7.26-7.25 (m, 1H), 7.13 (s, 1H), 6.42-6.41 (m, 1H), 3.52-3.49 (m, 1H), 2.75 (s, 3H), 1.57 (s, 6H), 1.11-1.08 (m, 2H), 1.02-1.00 (m, 2H).

Example 455: 9-(difluoromethyl)-6-fluoro-8-(6-fluoro-1-(methylsulfonyl)-1H-indol-4-yl)-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline

[0301]




[0302] To a solution of 8-bromo-9-(difluoromethyl)-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**Intermediate B-43**) (0.470 g, 1.388 mmol, 1.50 eq) in t-amyl alcohol:dioxane:water (3:2:1, 30 ml) was added K₂CO₃ (0.287 g, 2.082 mmol, 3.00 eq) at ambient temperature. The solution was degassed with Ar for 20 minutes followed by the addition of 6-fluoro-1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**Intermediate A-2**) (0.250 g, 0.694 mmol, 1.00 eq) and Attaphos (0.025 g, 0.035 mmol, 0.05 eq). The reaction mixture was then heated to 90°C for 2 h. After completion of the reaction (monitored by TLC & LCMS) the reaction mixture was diluted with water and was extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄ and evaporated under reduced pressure to get the crude compound. This crude material was then purified via column chromatography using MeOH-DCM as eluent and again purified via column chromatography using acetone-hexane as eluent. The resulting

material was then washed with ether to afford 9-(difluoromethyl)-6-fluoro-8-(6-fluoro-1-(methylsulfonyl)-1H-indol-4-yl)-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.150 g, 42%) as a white solid.

[0303] ^1H NMR ($\text{DMSO}-d_6$) δ : 7.68-7.65 (m, 1H), 7.61 (d, 1H), 7.29-7.26 (m, 2H), 7.18-6.92 (m, 2H), 6.56-6.55 (m, 1H), 3.56 (s, 3H), 1.54 (bs, 6H); (CH_3 - omitted by DMSO).

Example 465: 6-fluoro-8-(6-fluoro-1H-indol-4-yl)-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline

[0304]




[0305] Into a microwave vial were weighed out 6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**intermediate A-1**) (144.5 mg, 0.554 mmol, 2.0 eq), 8-bromo-6-fluoro-1,4,4,9-tetramethyl-SH-[1,2,4]triazolo[4,3-a]quinoxaline (**intermediate B-7**) (90.0 mg, 0.277 mmol, 1.0 eq) and bis(tri-*tert*-butylphosphine)palladium(0) (14.2 mg, 0.028 mmol, 0.1 eq). A stir bar was added, the vial was sealed and the vial was purged with nitrogen for 5 minutes under stirring. Then, THF (1.8 mL) and 2 M Na_2CO_3 (0.6 mL) solution were added and the mixture was heated to 60 °C for 16 hours. The mixture was then cooled back to ambient temperature, followed by the addition of DCM and water. The mixture was filtered through a hydrophobic frit, and the organic layer was evaporated under reduced pressure. The crude remains were then purified using silica gel column chromatography to yield 84.0 mg (80%) of 6-fluoro-8-(6-fluoro-1H-indol-4-yl)-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**example 465**) as a white solid.

[0306] ^1H NMR ($\text{DMSO}-d_6$) δ : 11.30 (s, 1H), 7.40 (t, 1H), 7.22 (dd, 1H), 7.18 (d, 1H), 6.91 (dd, 1H), 6.46 (d, 1H), 6.19 (t, 1H), 2.50 (s, 3H), 2.05 (s, 3H), 1.53 (s, 6H).

Example 466: 9-(difluoromethyl)-6-fluoro-1,4,4-trimethyl-8-(3-methyl-1H-indazol-7-yl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline

[0307]



[0308] Step1: To a solution of 8-bromo-9-(difluoromethyl)-6-fluoro-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**intermediate B-43**) (0.05 g, 0.138 mmol, 1 eq) in dioxane (10.0 ml) were added KOAc (0.041 g, 0.414 mmol, 3 eq) and bispinacolatodiborane (0.42 g, 0.166 mmol, 1.2 eq). The solution was degassed with Ar for 20 min followed by addition of $\text{PdCl}_2(\text{dppf}) \cdot \text{DCM}$ (0.011 g, 0.0138 mmol, 0.01 eq). The reaction mixture was refluxed for 16 h. After completion of reaction (monitored by TLC), solvent was evaporated under reduced pressure to get the crude product which was purified by column chromatography to afford 9-(difluoromethyl)-6-fluoro-1,4,4-trimethyl-8-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.05 g, 89%), as brown solid.


[0309] Step2: To a solution of 7-bromo-3-methyl-1H-indazole (0.10 g, 0.473 mmol, 1.0 eq) in t-amyl alcohol:dioxane: H_2O (3:2:1) (30 ml), K_2CO_3 (0.196 g, 1.419 mmol, 3 eq) was added at RT. The solution was degassed with Ar for 20 min followed by addition of 9-(difluoromethyl)-6-fluoro-1,4,4-trimethyl-8-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.231 g, 0.568 mmol, 1.2 eq) and Attaphos (0.017 g, 0.023 mmol, 0.05 eq). The reaction mixture was then placed to pre-heated oil bath at 90 °C for 2 h. After completion of reaction (monitored by TLC & LCMS), reaction mixture was diluted with water and extracted with EA. The organic layer was washed with brine, dried over Na_2SO_4 , filtered and the solvent was evaporated under reduced pressure to get the crude product

which was initially purified by column chromatography using MeOH-DCM as eluent. After that it was again repurified in combiflash column chromatography using acetone-hexane as eluent. The product was then washed with ether to afford pure 9-(difluoromethyl)-6-fluoro-1,4,4-trimethyl-8-(3-methyl-1H-indazol-7-yl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.08 g, 40%) as white solid.

[0310] ¹H NMR (400 MHz, dms_o-d₆): δ 12.55 (s, 1H), 7.72 (d, 1H), 7.26-7.22 (m, 2H), 7.14-7.12 (m, 1H), 7.11-6.82 (m, 2H), 2.51 (s, 3H), 1.54-1.51 (m, 6H).

Example 481: 8-(1-cyclopropyl-1H-indol-4-yl)-6-fluoro-N,1,4,4-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-amine

[0311]




[0312] To a solution of 8-bromo-6-fluoro-N,1,4,4-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-amine (**intermediate B-46**) (0.15 g, 0.44 mmol, 1 eq.) in t-amyl alcohol:dioxane (2:1) (5.0-2.5 ml) were added 2 M K₂CO₃ (1.0 ml) solution and 1-cyclopropyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**intermediate A-44**) (0.15 g, 0.53 mmol, 1.2 eq). The solution was degassed with Ar for 20 min followed by addition of Attaphos (0.016 g, 0.02 mmol, 0.05 eq.). The reaction mixture was heated to 90°C for 16 h. After completion of reaction (monitored by TLC), reaction mixture was filtered through celite pad and washed with EtOAc. Combined organic layer was evaporated to get the crude product which was purified by prep-HPLC (R_F-value-0.3: 5% MeOH/DCM) to afford 8-(1-cyclopropyl-1H-indol-4-yl)-6-fluoro-N,1,4,4-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-9-amine (0.09 g, 49%) as white solid.

[0313] ¹H NMR (400 MHz, dms_o-d₆): δ 7.63-7.61 (m, 1H), 7.40 (d, 1H), 7.31-7.27 (m, 1H), 7.08 (d, 1H), 7.00-6.98 (m, 1H), 6.46 (s, 1H), 6.20 (d, 1H), 4.07-4.05 (m, 1H), 3.48-3.47 (m, 1H), 2.53 (s, 3H), 2.16-2.14 (m, 3H), 1.58-1.49 (m, 6H), 1.10-1.08 (m, 2H), 0.99-0.98 (m, 2H).

Example 483: 6-chloro-8-(6-fluoro-1-(methylsulfonyl)-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline

[0314]




[0315] To a solution of 8-bromo-6-chloro-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**intermediate B-47**) (0.150 g, 0.42 mmol, 1 eq) in t-amyl alcohol:dioxane:H₂O (3:2:1) (10 ml) were added K₂CO₃ (0.174 g, 1.26 mmol, 3 eq) and 6-fluoro-1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**intermediate A-2**) (0.171 g, 0.50 mmol, 1.2 eq). The solution was degassed with Ar for 20 min followed by addition of Attaphos (0.015 g, 0.021 mmol, 0.05 eq.). The reaction mixture was heated to 90°C for 30 min. After completion of reaction (monitored by TLC), reaction mixture is filtered through celite pad and celite pad was washed with EtOAc. Combined organic layer was evaporated to get the crude product which was purified by prep-HPLC (R_F-value-0.3: 50% acetone/hexane) to afford 6-chloro-8-(6-fluoro-1-(methylsulfonyl)-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.05 g, 24%) as white solid.

[0316] ¹H NMR (400 MHz, dms_o-d₆): δ 7.68-7.65 (m, 1H), 7.64-7.63 (m, 1H), 7.40 (s, 1H), 7.33-7.30 (m, 1H), 6.68-6.67 (m, 1H), 6.33 (s, 1H), 3.56 (s, 3H), 3.21 (s, 3H), 2.55 (s, 3H), 1.57 (s, 6H).

Example 490: 6-fluoro-8-(7-fluoro-1H-indol-4-yl)-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline

[0317]




[0318] Into a microwave vial were weighed out 7-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (128.5 mg, 0.492 mmol, 2.0 eq), 8-bromo-6-fluoro-1,4,4,9-tetramethyl-SH-[1,2,4]triazolo[4,3-a]quinoxaline (**intermediate B-7**) (80.0 mg, 0.246 mmol, 1.0 eq) and Pd(PPh₃)₄. A stir bar was added, the vial was sealed and the vial was purged with nitrogen under stirring for 5 minutes. Then, toluene (2.1 mL), EtOH (0.6 mL) and 2M Na₂CO₃ solution (0.5 mL) were added successively. The reaction mixture was then heated to 90 °C for 16 hours. The mixture was allowed to cool to ambient temperature, diluted with water and DCM and was filtered through a hydrophobic frit. The organic layer was evaporated under reduced pressure to obtain the crude material, which was then purified via silica gel column chromatography and HPLC to obtain 6-fluoro-8-(7-fluoro-1H-indol-4-yl)-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**example 490**) in 71% (66.0 mg) yield as a white solid.

[0319] ¹H NMR (DMSO-d₆) δ: 11.76 (t, 1H), 7.47 (t, 1H), 7.13 (d, 1H), 7.01 (dd, 1H), 6.96 (dd, 1H), 6.42 (d, 1H), 6.27 (m, 1H), 2.49 (s, 3H), 2.02 (s, 3H), 1.52 (s, 6H).

Example 500: 6,7-difluoro-1,4,4-trimethyl-8-(1-(methylsulfonyl)-1H-indol-4-yl)-9-(trifluoromethyl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline

[0320]




[0321] To a solution of 8-bromo-6,7-difluoro-1,4,4-trimethyl-9-(trifluoromethyl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**intermediate B-50**) (0.07 g, 0.17 mmol, 1 eq) in t-amyl alcohol:dioxane:H₂O (3:2:1) (3.0 ml:2.0 ml: 1ml) were added K₂CO₃ (0.073 g, 0.53 mmol, 3 eq) and 1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**intermediate A-12**) (0.085 g, 0.26 mmol, 1.5 eq). The solution was degassed with Ar for 20 min followed by addition of Attaphos (0.006 g, 0.008 mmol, 0.05 eq). The reaction mixture was heated to 90°C for 16 h. After completion of reaction (monitored by TLC), reaction mixture was filtered through celite pad and washed with EtOAc. Combined organic layer was evaporated to get the crude product which was purified by prep-HPLC (R_f-value-0.3: 50% acetone/hexane) to afford 6,7-difluoro-1,4,4-trimethyl-8-(1-(methylsulfonyl)-1H-indol-4-yl)-9-(trifluoromethyl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.037 g, 39%) as white solid.

[0322] ¹H NMR (400 MHz, dmsO-d₆): δ 8.00-7.98 (m, 1H), 7.61 (s, 1H), 7.51-7.47 (m, 1H), 7.38 (s, 1H), 7.25 (s, 1H), 6.63 (s, 1H), 3.46 (s, 3H), 2.49 (s, 3H), 1.85 (s, 3H), 1.40 (s, 3H).

Example 518: 6,7-difluoro-8-(6-fluoro-1-(methylsulfonyl)-1H-indol-4-yl)-1,4,4-trimethyl-9-(trifluoromethyl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline

[0323]




[0324] To a solution of 8-bromo-6,7-difluoro-1,4,4-trimethyl-9-(trifluoromethyl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (**intermediate B-50**) (0.07 g, 0.17 mmol, 1 eq) in t-amyl alcohol:dioxane (1:1) (2.5 ml-2.5 ml) were added K_2CO_3 (0.073 g, 0.53 mmol, 3 eq) and 6-fluoro-1-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (**intermediate A-2**) (0.09 g, 0.26 mmol, 1.5 eq). The solution was degassed with Ar for 20 min followed by addition of X-phos (0.017 g, 0.035 mmol, 0.2 eq.) and $Pd_2(dba)_3$ (0.016 g, 0.017 mmol, 0.1 eq.). The reaction mixture was refluxed at 90°C for 16 h. After completion of reaction (monitored by TLC), reaction mixture was filtered through celite pad and washed with EtOAc. Combined organic layer was evaporated to get the crude product which was purified by prep-HPLC (R_f -value-0.3: 50% acetone/hexane) to afford 6,7-difluoro-8-(6-fluoro-1-(methylsulfonyl)-1H-indol-4-yl)-1,4,4-trimethyl-9-(trifluoromethyl)-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxaline (0.037 g, 37%) as white solid.

[0325] 1H NMR (400 MHz, $DMSO-d_6$): δ 7.77 (d, 1H), 7.63 (d, 1H), 7.28-7.34 (m, 2H), 6.64 (s, 1H), 3.51 (s, 3H), 2.47 (s, 3H), 1.77 (s, 3H), 1.38 (s, 3H).

Example 520: 4-(6,7-difluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indole-7-carbonitrile

[0326]



[0327] Into a microwave vial were weighed out 8-bromo-6,7-difluoro-1,4,4,9-tetramethyl-SH-[1,2,4]triazolo[4,3-a]quinoxaline (**intermediate B-38**) (60.0 mg, 0.175 mmol, 1.0 eq), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole-7-carbonitrile (**intermediate A-75**) (93.8 mg, 0.350 mmol, 2.0 eq), Pd_2dba_3 (16.0 mg, 0.018 mmol, 0.1 eq) and X-Phos (16.9 mg, 0.036 mmol, 0.2 eq). A stir bar was then added and the vial was sealed. The vial was then evacuated and backfilled with nitrogen three times, followed by the addition of 1,4-dioxane (1.3 mL), *tert*-amyl alcohol (1.3 mL) and 2M K_2CO_3 (0.4 mL). The reaction mixture was then sparged with nitrogen under ultra sonication, followed by heating to 60 °C for four hours. The mixture was allowed to cool to ambient temperature, diluted with water and DCM and was filtered through a hydrophobic frit. The organic layer was evaporated under reduced pressure to obtain the crude material, which was then purified via silica gel column chromatography and HPLC to obtain 4-(6,7-difluoro-1,4,4,9-tetramethyl-4,5-dihydro-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indole-7-carbonitrile (**example 520**) in 76% (54.0 mg) yield as a white solid.

[0328] 1H NMR ($DMSO-d_6$) δ : 12.20 (s, 1H), 7.72 (d, 1H), 7.57 (t, 1H), 7.22 (d, 1H), 6.90 (d, 1H), 6.36 (dd, 1H), 2.45 (s, 3H), 1.95 (s, 3H), 1.59 (s, 3H), 1.50 (s, 3H)

[0329] In the following Tables 3, 4 and 5 it is summarized how the remaining examples have been obtained.

Table 3 :

Ex. #	Intermediates	Synthesis in analogy to	yield (mol-%)	1H -NMR
48	Int-A-2 + Int-B-9	Ex. 49	31%	1H NMR (400 MHz, $DMSO-d_6$, at 100°C): 7.68 (d, 1H), 7.63 (s, 1H), 7.24 (d, 1H, $J=10.16$ Hz), 7.11 (d, 1H), 6.57 (s, 1H), 6.26 (s, 1H), 3.48 (s, 3H), 2.69-2.67(m, 2H), 2.53 (s, 3H), 1.57 (s, 6H), 0.54 (t, 3H).

(continued)

Ex. #	Intermediates	Synthesis in analogy to	yield (mol-%)	¹ H-NMR
53	Int-A-23 + Int-B-9	Ex. 49	15%	¹ H NMR (400 MHz, DMSO-d ₆ , at 100°C): 8.33 (s, 1H), 7.73 (d, 1H, J=9.0 Hz), 7.4 (d, 1H), 7.2 (d, 1H), 6.36 (s, 1H), 3.53 (s, 3H), 2.74-2.66 (m, 2H), 2.56 (s, 3H), 1.58 (s, 6H), 0.56 (t, 3H).
54	Int-A-24 + Int-B-10	Ex. 49	40%	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.44-7.38 (m, 2H), 7.24 (d, 1H), 7.0 (dd, , 1H), 6.67 (s, 1H), 6.33 (d, 1H), 4.33 (t, 2H), 3.67 (t, , 2H), 3.24 (s, 3H), 3.16 (s, 3H), 2.55 (s, 3H), 1.55 (s, 6H).
55	Int-A-20 + Int-B-10	Ex. 49	63%	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.40-7.39 (m, 2H), 7.23 (d, 1H), 7.0 (dd, , 1H), 6.67 (s, 1H), 6.33 (d, 1H), 4.9 (t, 1H), 4.22 (t, 2H), 3.74 (q, 2H), 3.17 (s, 3H), 2.56 (s, 3H), 1.55 (s, 6H).
56	Int-A-2 + Int-B-10	Ex. 49	34%	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.68-7.62 (m, 2H), 7.33-7.26 (m, 2H), 6.66 (s, 1H), 6.69 (d, 1H), 3.56 (s, 3H), 3.18 (s, 3H), 2.57 (s, 3H), 1.55 (s, 6H).
57	Int-A-12 + Int-B-10	Ex. 49	33%	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.89 (d, 1H), 7.62 (d, 1H), 7.48 (d, 1H), 7.38 (d, 1H), 7.23 (d, 1H), 6.69 (d, 1H), 3.5 (s, 3H), 3.14 (s, 3H), 2.57 (s, 3H), 1.55 (s, 6H).
59	Int-A-2 + Int-B-11	Ex. 49	20%	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.66 (dd, 1H), 7.63 (d, 1H), 7.31-7.36 (m, 1H), 7.19 (d, 1H), 6.55-6.56 (m, 2H), 3.56 (s, 3H), 2.18 (s, 3H), 1.95 (m, 1H), 1.50 (bs, 6H), 1.08 (bs, 4H).
60	Int-A-12 + Int-B11	Ex. 49	26%	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.88 (d, 1H), 7.62 (d, 1H), 7.46 (t, 1H), 7.33-7.35 (m, 1H), 7.16 (d, J = 10.4 Hz, 1H), 6.55 (d, 1H), 6.50 (s, 1H), 3.50 (s, 3H), 2.15 (s, 3H), 1.90 (m, 1H), 1.50 (bs, 6H), 1.07-1.11 (m, 4H).
61	Int-A-20 + Int-B-11	Ex. 49	27%	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.38-7.40 (m, 2H), 7.15 (d, 1H), 6.96 (d, 1H), 6.46 (s, 1H), 6.16 (d, 1H), 4.90 (t, J = 5.2 Hz, 1H), 4.21 (t, 2H), 3.73 (q, 2H), 2.19 (s, 3H), 1.92 (m, 1H), 1.50 (bs, 6H), 1.07-1.12 (m, 4H).
62	Int-A-15 + Int-B-11	Ex. 49	29%	¹ H NMR (400 MHz, DMSO-d ₆): δ 10.61 (s, 1H), 7.26 (dd, 1H), 7.15-7.18 (m, 2H), 6.93 (d, 1H), 6.46 (s, 1H), 2.25 (s, 3H), 2.07-2.08 (m, 4H), 1.51 (bs, 6H), 1.09 (bs, 4H).
63	Int-A-2 + Int-B-12	Ex. 49	20%	¹ H NMR (400 MHz, DMSO-d ₆): δ 10.61 (s, 1H), 7.26 (dd, 1H), 7.15-7.18 (m, 2H), 6.93 (d, 1H), 6.46 (s, 1H), 2.25 (s, 3H), 2.07-2.08 (m, 4H), 1.51 (bs, 6H), 1.09 (bs, 4H).
64	Int-A-22 + Int-B-12	Ex. 49	26%	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.88 (d, 1H), 7.62 (d, 1H), 7.46 (t, J = 8.0 Hz, 1H), 7.33-7.35 (m, 1H), 7.16 (d, J = 10.4 Hz, 1H), 6.55 (d, 1H), 6.50 (s, 1H), 3.50 (s, 3H), 2.15 (s, 3H), 1.90 (m, 1H), 1.50 (bs, 6H), 1.07-1.11 (m, 4H).
65	Int-A-20 + Int-B-12	Ex. 49	27%	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.38-7.40 (m, 2H), 7.15 (d, 1H), 6.96 (d, J = 10.0 Hz, 1H), 6.46 (s, 1H), 6.16 (d, 1H), 4.90 (t, 1H), 4.21 (t, 2H), 3.73 (q, 2H), 2.19 (s, 3H), 1.92 (m, 1H), 1.50 (bs, 6H), 1.07-1.12 (m, 4H).
66	Int-A-15 + Int-B-12	Ex. 49	29%	¹ H NMR (400 MHz, DMSO-d ₆): δ 10.61 (s, 1H), 7.26 (dd, 1H), 7.15-7.18 (m, 2H), 6.93 (d, 1H), 6.46 (s, 1H), 2.25 (s, 3H), 2.07-2.08 (m, 4H), 1.51 (bs, 6H), 1.09 (bs, 4H).

Table 4:

Ex. #	Intermediates	Synthesis in analogy to	yield (mol-%)	m/z [M+1]	¹ H-NMR
300	Int-B-7 + Int-A-43	Exp. 465	22%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.54 (m _c , 1H), 7.41 (d, 1H), 7.23 (dd, 1H), 7.13 (d, 1H), 7.02 (d, 1H), 6.44 - 6.40 (m, 1H), 6.21 - 6.17 (m, 1H), 4.38 (t, 2H), 3.70 (t, 2H), 3.25 (d, 3H), 2.50 (s, 3H), 2.03 (s, 3H), 1.55 - 1.51 (m, 6H)
304	Int-B-7 + 1-methyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-indazole	Exp. 490	57%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.86 (s, 1H), 7.68 (d, 1H), 7.49 (dd, 1H), 7.20 (d, 1H), 7.16 (d, 1H), 6.51 (s, 1H), 4.10 (s, 3H), 2.52 (s, 3H), 2.04 (s, 3H), 1.53 (s, 6H)
305	Int-B-7 + Int-A-44	Exp. 465	88%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.61 (dd, 1H), 7.38 (d, 1H), 7.27 (dd, 1H), 7.11 (d, 1H), 7.08 - 7.03 (m, 1H), 6.42 (d, 1H), 6.15 (dd, 1H), 2.49 (s, 3H), 2.02 (s, 3H), 1.52 (s, 6H), 1.34 - 1.25 (m, 1H), 1.13 - 1.05 (m, 2H), 1.02 - 0.96 (m, 2H)
308	Int-B-30 + Int-A-12	Exp. 465	70%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.91 (d, 1H), 7.66 (d, 1H), 7.48 (dd, 1H), 7.31 (d, 1H), 7.15 (d, 1H), 6.55 (dd, 1H), 6.50 (d, 1H), 3.51 (s, 3H), 2.84 (s, 2H), 2.01 (s, 3H), 1.75 - 1.34 (m, 6H), 1.29 (t, 3H)
316	Int-B-30 + Int-A-15	Exp. 465	80%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 10.59 (d, 1H), 7.27 (dd, 1H), 7.21 - 7.12 (m, 2H), 6.90 (d, 1H), 6.45 (d, 1H), 2.86 (s, 2H), 2.27 (d, 3H), 1.98 (s, 3H), 1.69 - 1.39 (m, 6H), 1.29 (t, 3H)
322	Int-B-30 + Int-A-43	Exp. 465	66%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.53 (dd, 1H), 7.41 (d, 1H), 7.22 (dd, 1H), 7.12 (d, 1H), 7.01 (d, 1H), 6.41 (d, 1H), 6.16 (m _c , 1H), 4.37 (t, 2H), 3.70 (t, 2H), 3.25 (d, 3H), 2.84 (s, 2H), 2.02 (s, 3H), 1.52 (s, 6H), 1.28 (t, 3H)
323	Int-B-30 + Int-A-2	Exp. 465	82%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.69 - 7.67 (m, 1H), 7.66 (d, 1H), 7.27 (d, 1H), 7.19 (d, 1H), 6.54 (d, 1H), 6.53 (dd, 1H), 3.56 (s, 3H), 2.83 (s, 2H), 2.03 (s, 3H), 1.74 - 1.33 (m, 6H), 1.28 (t, 3H)
326	Int-B-7 + Int-A-42	Exp. 490	62%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.60 (d, 1H), 7.44 (d, 1H), 7.27 (dd, 1H), 7.14 (d, 1H), 7.08 - 7.04 (m, 1H), 6.51 - 6.30 (m, 2H), 6.27 (dd, 1H), 4.73 (td, 2H), 2.50 (s, 3H), 2.02 (s, 3H), 1.52 (s, 6H)
340	Int-B-7 + Int-A-23	Exp. 465	77%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.47 - 8.43 (m, 1H), 7.72 (dd, 1H), 7.47 (dd, 1H), 7.29 (dd, 1H), 6.63 (d, 1H), 3.57 (d, 3H), 2.54 (d, 3H), 2.08 (s, 3H), 1.53 (s, 6H)
349	Int-B-7 + Int-A-45	Exp. 465	57%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.89 (d, 1H), 7.65 (d, 1H), 7.54 - 7.46 (m, 1H), 7.35 (d, 1H), 7.17 (d, 1H), 6.66 (d, 1H), 6.51 (d, 1H), 6.07 (d, 2H), 2.50 (s, 3H), 2.01 (s, 3H), 1.58 (s, 3H), 1.48 (s, 3H)

(continued)

Ex. #	Intermediates	Synthesis in analogy to	yield (mol-%)	m/z [M+1]	¹ H-NMR
350	Int-B-7 + Int-A-47	Exp. 465	56%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.90 (m _c , 1H), 7.65 (m _c , 1H), 7.47 (m _c , 1H), 7.31 (m _c , 1H), 7.17 (m _c , 1H), 6.58 (m _c , 1H), 6.50 (t, 1H), 3.66 (m _c , 2H), 2.50 - 2.49 (m, 3H), 2.00 (s, 3H), 1.45 (s, 6H), 1.10 (m _c , 3H)
352	Int-B-7 + 3-Methyl-1H-indazole-7-boronic acid	Exp. 465	20%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 12.56 (s, 1H), 7.74 (dd, 1H), 7.26 (dd, 1H), 7.20 - 7.12 (m, 2H), 6.47 (d, 1H), 2.55 (s, 3H), 2.53 (s, 3H), 1.97 (s, 3H), 1.53 (s, 6H)
355	Int-B-30 + 3-methyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole	Exp. 465	81%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 10.47 (d, 1H), 7.51 (d, 1H), 7.14 - 7.06 (m, 3H), 6.99 (d, 1H), 6.40 (d, 1H), 2.93 - 2.78 (m, 2H), 2.29 (d, 3H), 1.96 (s, 3H), 1.67 - 1.43 (m, 6H), 1.28 (t, 3H)
366	Int-B-7 + Int-A-30	Exp. 465	97%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.90 (d, 1H), 7.64 (d, 1H), 7.47 (dd, 1H), 7.31 (d, 1H), 7.18 (d, 1H), 6.58 (d, 1H), 6.50 (d, 1H), 3.84 (m _c , 1H), 2.50 (s, 3H), 2.00 (s, 3H), 1.64 - 1.42 (m, 6H), 1.23 (s, 3H), 1.22 (s, 3H)
374	Int-B-30 + Int-A-23	Exp. 465	85%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.42 (d, 1H), 7.75 - 7.70 (m, 1H), 7.49 - 7.44 (m, 1H), 7.29 (d, 1H), 6.62 (d, 1H), 3.57 (s, 3H), 2.94 - 2.80 (m, 2H), 2.10 - 2.06 (m, 3H), 1.67 - 1.42 (m, 6H), 1.30 (t, 3H)
375	Int-B-7 + Int-A-46	Exp. 465	67%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.52 (d, 1H), 7.45 (d, 1H), 7.23 (dd, 1H), 7.13 (d, 1H), 7.02 (d, 1H), 6.42 (d, 1H), 6.20 - 6.13 (m, 1H), 4.30 (t, 2H), 2.65 (t, 2H), 2.49 (d, 3H), 2.21 (d, 6H), 2.03 (s, 3H), 1.57 - 1.48 (m, 6H)
378	Int-B-7 + Int-A-22	Exp. 465	99%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.45 (s, 1H), 8.03 (d, 1H), 7.73 (dd, 1H), 7.46 (d, 1H), 7.25 (d, 1H), 6.58 (d, 1H), 3.53 (s, 3H), 2.52 (d, 3H), 2.05 (s, 3H), 1.54 (s, 6H)
389	Int-B-30 + Int-A-22	Exp. 465	96%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.41 (d, 1H), 8.02 (d, 1H), 7.73 (ddd, 1H), 7.46 (d, 1H), 7.24 (d, 1H), 6.57 (d, 1H), 3.53 (d, 3H), 2.86 (s, 2H), 2.04 (s, 3H), 1.72 - 1.38 (m, 6H), 1.30 (t, 3H)
391	Int-B-7 + Int-A-48	Exp. 465	86%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.95 (dd, 1H), 7.66 (dd, 1H), 7.50 - 7.44 (m, 1H), 7.31 (d, 1H), 7.18 (dd, 1H), 6.58 (dd, 1H), 6.50 (d, 1H), 3.15 (m _c , 1H), 2.53 - 2.47 (m, 3H), 2.01 (s, 3H), 1.53 (m, 6H), 1.33 - 1.25 (m, 2H), 1.15 - 1.07 (m, 2H)
464	Int-B-9 + Int-A-20	Exp. 465	43%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.43 (d, 1H), 7.42 - 7.38 (m, 1H), 7.13 (d, 1H), 6.55 (d, 1H), 6.28 - 6.18 (m, 1H), 4.23 (t, 2H), 3.76 (m _c , 2H), 2.67 (s, 2H), 2.54 (s, 3H), 1.78 - 1.35 (m, 6H), 0.49 (t, 3H)

(continued)

Ex. #	Intermediates	Synthesis in analogy to	yield (mol-%)	m/z [M+1]	¹ H-NMR
467	Int-B-45 + Int-A-23	Exp. 465	50%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.52 (s, 1H), 7.72 (dd, 1H), 7.52 (s, 1H), 7.29 (d, 1H), 6.64 (d, 1H), 3.57 (s, 3H), 2.65 (s, 3H), 2.02 (s, 1H), 1.80 (s, 3H), 1.19 (s, 3H), 0.65 (s, 1H), 0.25 (s, 1H), -0.20 (d, 2H)
468	Int-B-7 + indole-4-boronic acid	Exp. 465	99%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 11.24 (s, 1H), 7.44 (d, 1H), 7.40 (t, 1H), 7.18 (t, 1H), 7.14 (d, 1H), 6.99 (d, 1H), 6.40 (d, 1H), 6.19 (dd, 1H), 2.50 (d, 3H), 2.03 (s, 3H), 1.53 (s, 6H)
469	Int-B-10 + indole-4-boronic acid	Exp. 465	71%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 11.19 (s, 1H), 7.44 (d, 1H), 7.37 (t, 1H), 7.23 - 7.15 (m, 2H), 7.10 (d, 1H), 6.57 (d, 1H), 6.35 (t, 1H), 3.13 (s, 3H), 2.57 (s, 3H), 1.56 (s, 6H)
470	Int-B-10 + Int-A-1	Exp. 465	75%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 11.26 (s, 1H), 7.38 (t, 1H), 7.25 (d, 1H), 7.22 (dd, 1H), 6.99 (dd, 1H), 6.64 (d, 1H), 6.35 (t, 1H), 3.16 (s, 3H), 2.57 (s, 3H), 1.56 (s, 6H)
471	Int-B-45 + Int-A-2	Exp. 465	67%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.69 - 7.63 (m, 2H), 7.47 - 7.23 (m, 1H), 7.19 (d, 1H), 6.69 - 6.64 (m, 1H), 6.56 (d, 1H), 3.54 (s, 3H), 2.61 (s, 3H), 2.08-1.82 (m, 1H), 1.79 (s, 3H), 1.18 (s, 3H), 0.62 (s, 1H), 0.21 (s, 1H), -0.19 (d, 2H)
472	Int-B-45 + 3-methyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole	Exp. 465	63%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 10.49 (s, 1H), 7.50 (d, 1H), 7.27 - 6.70 (m, 4H), 6.42 (d, 1H), 2.64 (s, 3H), 2.30 (s, 3H), 2.04-1.86 (m, 1H), 1.79 (s, 3H), 1.19 (s, 3H), 0.55 (s, 1H), 0.06 (s, 1H), -0.14 - -0.34 (m, 2H)
473	Int-B-10 + 4-(tetramethyl-1,3,2-dioxaborolan-2-yl)-6-(trifluoromethyl)-1H-indole	Exp. 465	68%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 11.64 (s, 1H), 7.78 (d, 1H), 7.64 (t, 1H), 7.36 (d, 1H), 7.28 (d, 1H), 6.67 (d, 1H), 6.47 (t, 1H), 3.15 (s, 3H), 2.57 (s, 3H), 1.57 (s, 6H)
474	Int-B-45 + Int-A-12	Exp. 465	45%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.89 (d, 1H), 7.64 (d, 1H), 7.48 (t, 1H), 7.36 (s, 1H), 7.15 (d, 1H), 6.67 (d, 1H), 6.51 (d, 1H), 3.49 (s, 3H), 3.35 (s, 1H), 2.61 (s, 3H), 1.79 (s, 3H), 1.18 (s, 3H), 0.60 (s, 1H), 0.13 (s, 1H), -0.20 (s, 2H)
475	Int-B-7 + Int-A-54	Exp. 465	91%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.16 (d, 1H), 7.15 (d, 1H), 6.48 (d, 1H), 7.72 (d, 1H), 7.55 (d, 1H), 6.52 (dd, 1H), 3.48 (s, 3H), 2.52 (s, 3H), 2.50 (s, 3H), 2.01 (s, 3H), 1.73 - 1.17 (m, 6H)
476	Int-B-10 + Int-A-54	Exp. 465	63%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.72 (m _c , 1H), 7.53 (d, 1H), 7.23 (t, 1H), 7.21 (d, 1H), 6.66 (d, 1H), 6.63 (dd, 1H), 3.47 (s, 3H), 3.16 (s, 3H), 2.58 (s, 3H), 2.53 (s, 3H), 1.56 (s, 6H)

(continued)

Ex. #	Intermediates	Synthesis in analogy to	yield (mol-%)	m/z [M+1]	¹ H-NMR
477	Int-B-7 + 4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole-6-carbonitrile	Exp. 465	67%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 11.84 - 11.80 (m, 1H), 7.95 (d, 1H), 7.71 (t, 1H), 7.38 (d, 1H), 7.20 (d, 1H), 6.50 (d, 1H), 6.34 (t, 1H), 2.51 (s, 3H), 2.02 (s, 3H), 1.55 - 1.49 (m, 6H)
478	Int-B-10 + Int-A-16	Exp. 465	48%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 10.56 (d, 1H), 7.36 (dd, 1H), 7.22 (d, 1H), 7.08 (d, 1H), 6.98 (dd, 1H), 6.64 (d, 1H), 3.18 (d, 3H), 2.57 (s, 3H), 1.92 (m _C , 1H), 1.56 (s, 6H), 0.86 (m _C , 2H), 0.66 - 0.55 (m, 2H)
484	Int-B-7 + Int-A-59	Exp. 465	24%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.48 (dd, 1H), 7.38 (d, 1H), 7.25 (dd, 1H), 7.12 (d, 1H), 7.06 - 7.00 (m, 1H), 6.42 (d, 1H), 6.18 (dd, 1H), 3.84 (s, 3H), 2.49 (s, 3H), 2.02 (s, 3H), 1.52 (s, 6H)
485	Int-B-10 + Int-A-59	Exp. 465	52%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.47 (d, 1H), 7.35 (d, 1H), 7.25 (dd, 1H), 7.19 (d, 1H), 7.14 (dd, 1H), 6.58 (d, 1H), 6.33 (dd, 1H), 3.83 (s, 3H), 3.12 (s, 3H), 2.57 (s, 3H), 1.56 (s, 6H)
489	Int-B-10 + 7-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	Exp. 465	36%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.06 (dd, 1H), 7.30 (d, 1H), 7.28 (d, 1H), 7.19 (dd, 1H), 6.69 (s, 1H), 3.13 (s, 3H), 2.61 (s, 3H), 1.56 (s, 6H)
491	Int-B-25 + Int-A-53	Exp. 465	57%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.90 (d, 1H), 7.62 (d, 1H), 7.46 (t, 1H), 7.30 (d, 1H), 7.17 (d, 1H), 6.58 (d, 1H), 6.50 (d, 1H), 3.61 (d, 2H), 2.50 (s, 3H), 2.01 (s, 3H), 1.77 - 1.33 (m, 6H), 0.90 - 0.76 (m, 1H), 0.42 - 0.27 (m, 2H), -0.00 - -0.11 (m, 2H)
492	Int-B-10 + Int-A-66	Exp. 465	59%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.22 - 8.18 (m, 1H), 7.88 (d, 1H), 7.69 (d, 1H), 7.32 (d, 1H), 6.82 (dd, 1H), 6.77 (d, 1H), 3.64 (s, 3H), 3.18 (s, 3H), 2.58 (s, 3H), 1.57 (s, 6H)
494	Int-B-10 7-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole	Exp. 490	50%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 11.69 (t, 1H), 7.44 (t, 1H), 7.20 (d, 1H), 7.09 - 6.98 (m, 2H), 6.59 (d, 1H), 6.41 (m, 1H), 3.12 (s, 3H), 2.57 (s, 3H), 1.56 (s, 6H)
495	Int-B-10 + 6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	Exp. 465	22%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 13.21 (s, 1H), 7.99 (s, 1H), 7.39 - 7.32 (m, 2H), 7.14 (dd, 1H), 6.78 - 6.73 (m, 1H), 3.17 (s, 3H), 2.61 (s, 3H), 1.56 (s, 6H)
496	Int-B-7 + 6-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	Exp. 465	51%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 13.28 (s, 1H), 7.89 (d, 1H), 7.41 - 7.34 (m, 1H), 7.26 (d, 1H), 7.09 (dd, 1H), 6.55 (d, 1H), 2.53 (s, 3H), 2.07 (s, 3H), 1.53 (s, 6H)

(continued)

Ex. #	Intermediates	Synthesis in analogy to	yield (mol-%)	m/z [M+1]	¹ H-NMR
497	Int-B-7 + Int-A-68	Exp. 465	20%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.66 (dd, 1H), 7.63 (dd, 1H), 7.29 - 7.23 (m, 1H), 7.21 (d, 1H), 6.58 (d, 1H), 6.55 (d, 1H), 3.68 (d, 2H), 2.51 - 2.50 (m, 3H), 2.03 (s, 3H), 1.53 (s, 6H), 0.86 (tt, 1H), 0.36 (d, 2H), -0.01 - -0.06 (m, 2H)
502	Int-B-51 + Int-A-12	Exp. 465	37%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.95 (m, 1H), 7.65 (d, 1H), 7.51 (dd, 1H), 7.34 (d, 1H), 6.55 (d, 1H), 6.45 (s, 1H), 3.52 (s, 3H), 2.44 (s, 3H), 1.98 (s, 3H), 1.63 (s, 3H), 1.52 (s, 3H)
503	Int-B-51 + Int-A-2	Exp. 465	70%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.73 (m, 1H), 7.66 (d, 1H), 7.35 (dd, 1H), 6.56 (d, 1H), 6.50 (s, 1H), 3.59 (s, 3H), 2.45 (s, 3H), 2.01 (s, 3H), 1.64 - 1.60 (m, 3H), 1.53 (s, 3H),
504	Int-B-7 + 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	Exp. 465	13%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.88 (s, 1H), 7.58 (d, 1H), 7.45 (dd, 1H), 7.21 (d, 1H), 7.12 (d, 1H), 6.50 (d, 1H), 2.52 (s, 3H), 2.05 (s, 3H), 1.53 (s, 6H)
505	Int-B-10 + Int-A-48	Exp. 465	62%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.94 (dt, 1H), 7.64 (d, 1H), 7.47 (dd, 1H), 7.38 (dd, 1H), 7.24 (d, 1H), 6.72 - 6.64 (m, 2H), 3.17 - 3.09 (m, 4H), 2.58 (s, 3H), 1.56 (s, 6H), 1.28 (dd, 2H), 1.11 (dd, 2H)
506	Int-B-7 + Int-A-69	Exp. 465	90%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 11.37 (t, 1H), 7.49 (dd, 1H), 7.45 (t, 1H), 7.17 (d, 1H), 7.03 (d, 1H), 6.47 (d, 1H), 6.20 (ddd, 1H), 2.49 (s, 3H), 2.03 (s, 3H), 1.52 (s, 6H)
507	Int-B-7 + 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-6-(trifluoromethyl)-1H-indazole	Exp. 465	35%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.06 (s, 1H), 7.98 (s, 1H), 7.41 (s, 1H), 7.29 (d, 1H), 6.57 (d, 1H), 2.54 (s, 3H), 2.04 (s, 3H), 1.53 (s, 6H)
508	Int-B-51 + Int-A-15	Exp. 465	67%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 10.70 (d, 1H), 7.33 (dd, 1H), 7.18 (dd, 1H), 6.97 (dd, 1H), 6.41 (s, 1H), 2.45 (s, 3H), 2.27 (s, 3H), 1.96 (s, 3H), 1.61 (s, 3H), 1.55 (s, 3H)
509	Int-B-51 + Int-A-22	Exp. 465	39%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.46 (s, 1H), 8.08 (d, 1H), 7.77 (dd, 1H), 7.49 (d, 1H), 6.52 (s, 1H), 3.56 (s, 3H), 2.47 (s, 3H), 2.01 (s, 3H), 1.64 (s, 3H), 1.53 (s, 3H)
510	Int-B-51 + Int-A-23	Exp. 465	4%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.47 (s, 1H), 7.81 - 7.75 (m, 1H), 7.54 (dd, 1H), 6.58 (s, 1H), 3.60 (s, 3H), 2.48 (s, 3H), 2.04 (s, 3H), 1.58 (d, 6H)
511	Int-B-10 + Int-A-44	Exp. 465	20%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.60 (dd, 1H), 7.35 (d, 1H), 7.27 (dd, 1H), 7.21 - 7.13 (m, 2H), 6.59 (d, 1H), 6.29 (dd, 1H), 3.48 (tt, 1H), 3.13 (s, 3H), 2.56 (s, 3H), 1.55 (s, 6H), 1.09 (td, 2H), 1.02 - 0.96 (m, 2H)

(continued)

Ex. #	Intermediates	Synthesis in analogy to	yield (mol-%)	m/z [M+1]	¹ H-NMR
512	Int-B-38 + indole-4-boronic acid	Exp. 465	52%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 11.27 (s, 1H), 7.49 (d, 1H), 7.40 (t, 1H), 7.21 (t, 1H), 7.01 (d, 1H), 6.77 (s, 1H), 6.14 (t, 1H), 2.45 (s, 3H), 1.97 (s, 3H), 1.60 (s, 3H), 1.49 (s, 3H)
513	Int-B-38 + Int-3-methyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole	Exp. 465	73%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 10.59 (d, 1H), 7.57 (d, 1H), 7.17 - 7.08 (m, 2H), 7.05 (dd, 1H), 6.78 (d, 1H), 2.47 (s, 3H), 2.30 (d, 3H), 1.93 (s, 3H), 1.56 (d, 6H)
514	Int-B-38 + Int-A-15	Exp. 465	55%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.33 (dd, 1H), 7.19 (s, 1H), 6.99 (dd, 1H), 6.84 (d, 1H), 2.47 (s, 3H), 2.27 (d, 3H), 1.95 (s, 3H), 1.57 (s, 3H), 1.55 (s, 3H)
515	Int-B-10 + Int-A-71	Exp. 465	4%	396.1	
517	Int-B-7 7-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole	Exp. 465	46%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.99 (d, 1H), 7.28 (dd, 1H), 7.20 (d, 1H), 7.09 (dd, 1H), 6.50 (d, 1H), 2.52 (s, 3H), 2.03 (s, 3H), 1.53 (s, 6H)
521	Int-B-38 + Int-A-73	Exp. 465	59%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.42 (dt, 1H), 7.38 (d, 1H), 6.99 (dd, 1H), 6.84 (d, 1H), 6.14 (d, 1H), 3.82 (d, 3H), 2.44 (s, 3H), 1.97 (s, 3H), 1.59 (s, 3H), 1.50 (s, 3H),
522	Int-B-38 + Int-A-77	Exp. 520	55%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.49 (d, 1H), 7.46 - 7.42 (m, 1H), 7.01 (d, 1H), 6.87 (d, 1H), 6.46 (d, 1H), 3.88 (s, 3H), 2.45 (s, 3H), 1.97 (s, 3H), 1.58 (s, 3H), 1.49 (s, 3H) (MeO- omitted by water)
523	Int-B-7 + Int-A-77	Exp. 520	87%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.49 (d, 1H), 7.42 - 7.38 (m, 1H), 7.16 (d, 1H), 6.96 (d, 1H), 6.51 (d, 1H), 6.50 - 6.45 (m, 1H), 3.89 (s, 3H), 3.48 (s, 3H), 2.50 (s, 3H), 2.02 (s, 3H), 1.52 (s, 6H)
524	Int-B-10 + Int-A-77	Exp. 465	86%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.47 (d, 1H), 7.42 - 7.38 (m, 1H), 7.24 (d, 1H), 7.03 (d, 1H), 6.69 (d, 1H), 6.60 (dd, 1H), 3.89 (s, 3H), 3.46 (s, 3H), 3.18 (s, 3H), 2.57 (s, 3H), 1.55 (s, 6H)
525	Int-B-53 + Int-A-23	Exp. 465	41%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.48 (s, 1H), 7.78 (d, 1H), 7.54 (d, 1H), 6.58 (s, 1H), 3.60 (s, 3H), 2.48 (d, 3H), 2.05 (s, 3H), 1.62 (s, 3H), 1.55 (s, 3H)
526	Int-B-53 + Int-A-15	Exp. 465	61%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 10.72 - 10.68 (m, 1H), 7.33 (dd, 1H), 7.21 - 7.12 (m, 1H), 6.97 (dd, 1H), 6.41 (s, 1H), 2.45 (s, 3H), 2.27 (d, 3H), 1.96 (s, 3H), 1.61 (s, 3H), 1.55 (s, 3H)
527	Int-B-53 + Int-A-22	Exp. 465	73%		¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.46 (s, 1H), 8.08 (d, 1H), 7.81 - 7.75 (m, 1H), 7.49 (d, 1H), 6.53 (s, 1H), 3.56 (d, 3H), 2.47 (s, 3H), 2.01 (s, 3H), 1.65 (s, 3H), 1.53 (s, 3H)

Table 5 :

Ex. #	Intermediates	Synthesis in analogy to	yield (mol-%)	¹ H-NMR
447	Int-A2 + Int-B-38	Ex. 446	56%	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.72 (dd, 1H), 7.66 (d, 1H), 7.36 (dd, 1H), 6.93 (s, 1H), 6.58 (d, 1H), 3.59 (s, 3H), 2.45 (s, 3H), 1.97 (s, 3H), 1.57 (s, 3H), 1.50 (s, 3H).
449	Int-A-20 + Int-B-38	Ex. 448	59%	¹ H NMR (DMSO-d ₆) δ= 7.46 (d, 1H), 7.41 (d, 1H), 6.98 (d, 1H), 6.85 (s, 1H), 6.13 (s, 1H), 4.92 (s, 1H), 4.22 (s, 2H), 3.74 (s, 2H), 2.44 (s, 3H), 1.97 (s, 3H), 1.58 (s, 3H), 1.48 (s, 3H).
451	Int-A-22 + Int-B-38	Ex. 448	54%	¹ H NMR (DMSO-d ₆) δ= 8.49 (s, 1H), 8.07 (d, 1H), 7.77 (t, 1H), 7.51 (d, 1H), 6.95 (s, 1H), 3.55 (s, 3H), 2.48 (s, 3H), 1.98 (s, 3H), 1.52-1.58 (m, 6H).
452	Int-A-23 + Int-B-38	Ex. 448	58%	¹ H NMR (DMSO-d ₆) δ= 8.51 (s, 1H), 7.77 (d, 1H), 7.57 (d, 1H), 7.02 (s, 1H), 3.59 (s, 3H), 2.02 (s, 3H), 1.53-1.56 (m, 6H). (CH ₃ -omitted by DMSO)
453	Int-A-2 + Int-B-41	Ex. 450	28%	¹ H NMR (DMSO-d ₆) δ= 7.66-7.64 (m, 2H), 7.29-7.26 (m, 1H), 7.18-7.16 (m, 1H), 6.58-6.56 (m, 2H), 4.84 (s, 1H), 4.55 (s, 2H), 3.56 (s, 3H), 1.52 (s, 6H). (CH ₃ -omitted by DMSO)
454	Int-A-15 + Int-B-42	Ex. 450	54%	¹ H NMR (DMSO-d ₆) δ= 10.62 (s, 1H), 7.28-7.25 (m, 2H), 7.17 (s, 1H), 6.90-6.88 (m, 1H), 6.06 (s, 1H), 2.25 (s, 3H), 1.97 (s, 3H), 1.62-1.48 (m, 6H). (CH ₃ -omitted by DMSO)
456	Int-A-12 + Int-B-42	Ex. 450	55%	¹ H NMR (DMSO-d ₆) δ= 7.91 (d, 1H), 7.65-7.64 (m, 1H), 7.49 (t, 1H), 7.31-7.29 (m, 1H), 7.27 (s, 1H), 6.57-6.56 (m, 1H), 6.10 (s, 1H), 3.50 (s, 3H), 2.47 (s, 3H), 2.01 (s, 3H), 1.62-1.47 (m, 6H).
457	Int-A-2 + Int-B-42	Ex. 450	52%	¹ H NMR (DMSO-d ₆) δ= 7.68-7.65 (m, 2H), 7.31-7.27 (m, 2H), 6.56-6.55 (m, 1H), 6.15 (s, 1H), 3.56 (s, 3H), 2.03 (s, 3H), 1.59-1.48 (m, 6H). (CH ₃ -omitted by DMSO)
458	Int-A-22 + Int-B-42	Ex. 450	57%	¹ H NMR (DMSO-d ₆) δ= 8.43 (s, 1H), 8.03 (d, 1H), 7.74 (t, 1H), 7.45 (d, 1H), 7.35 (s, 1H), 6.17 (s, 1H), 3.53 (s, 3H), 2.04 (s, 3H), 1.55 (s, 6H). (CH ₃ -omitted by DMSO)
459	Int-A-12 + Int-B-43	Ex. 455	40%	¹ H NMR (400 MHz, dmso-d ₆): δ 7.90-7.88 (m, 1H), 7.59 (d, 1H), 7.45-7.41 (m, 1H), 7.33-7.31 (m, 1H), 7.24-7.21 (m, 1H), 7.12-6.86 (m, 2H), 6.56-6.55 (m, 1H), 3.50 (s, 3H), 1.54 (bs, 6H). (CH ₃ -omitted by DMSO)
460	Int-A-8 + Int-B-43	Ex. 455	40%	¹ H NMR (400 MHz, dmso-d ₆): δ 7.67-7.65 (m, 1H), 7.61 (d, 1H), 7.31-7.25 (m, 2H), 7.17-6.91 (m, 2H), 6.56-6.55 (m, 1H), 3.75-3.69 (m, 2H), 1.61-1.54 (m, 6H), 1.14-1.07 (m, 3H). (CH ₃ -omitted by DMSO)
461	Int-A-47 + Int-B-43	Ex. 455	40%	¹ H NMR (400 MHz, dmso-d ₆): δ 7.89 (d, 1H), 7.60 (d, 1H), 7.44-7.40 (m, 1H), 7.33-7.31 (m, 1H), 7.26-7.23 (m, 1H), 7.11-6.85 (m, 2H), 6.56-6.55 (m, 1H), 3.69-3.63 (m, 2H), 1.54-1.50 (m, 6H), 1.14-1.05 (m, 3H). (CH ₃ -omitted by DMSO)
462	Int-A-15 + Int-B-43	Ex. 455	40%	¹ H NMR (400 MHz, dmso-d ₆): δ 10.57 (s, 1H), 7.28-7.22 (m, 2H), 7.14 (s, 1H), 7.08-6.82 (m, 3H), 2.25 (s, 3H), 1.55-1.48 (m, 6H). (CH ₃ -omitted by DMSO)
463	Int-A-20 + Int-B-42	Ex. 450	43%	¹ H NMR (400 MHz, dmso-d ₆): δ 7.42-7.39 (m, 2H), 7.28 (s, 1H), 6.93-6.91 (m, 1H), 6.16-6.15 (m, 1H), 6.07 (m, 1H), 4.91 (t, 1H), 4.23 (t, 2H), 3.75-3.71 (m, 2H), 2.47 (s, 3H), 2.05 (s, 3H), 1.54 (s, 6H).

(continued)

Ex. #	Intermediates	Synthesis in analogy to	yield (mol-%)	¹ H-NMR
479	Int-A-8 + Int-B-46	Ex. 481	37%	¹ H NMR (400 MHz, dmso-d ₆): δ 7.66-7.64 (m, 2H), 7.24-7.21 (m, 1H), 7.06-7.03 (m, 1H), 6.57-6.55 (m, 2H), 4.45-4.44 (m, 1H), 3.73-3.68 (m, 2H), 2.54 (s, 3H), 2.17-2.15 (m, 3H), 1.60-1.47 (m, 6H), 1.14 (t, 3H). (CH ₃ -omitted by DMSO)
480	Int-A-20 + Int-B-46	Ex. 481	52%	¹ H NMR (400 MHz, dmso-d ₆): δ 7.43-7.41 (m, 2H), 7.04-7.02 (m, 1H), 6.92-6.89 (m, 1H), 6.51 (s, 1H), 6.23 (d, 1H), 4.89-4.86 (m, 1H), 4.23-4.21 (m, 2H), 4.17-4.16 (m, 1H), 3.75-3.71 (m, 2H), 2.56 (s, 3H), 2.17-2.16 (m, 3H), 1.54 (bs, 6H).
482	Int-A-12 + Int-B-47	Ex. 455	20%	¹ H NMR (400 MHz, dmso-d ₆): δ 7.90-7.88 (m, 1H), 7.62 (d, 1H), 7.47-7.45 (m, 1H), 7.39-7.35 (m, 2H), 6.68 (d, 1H), 6.27 (s, 1H), 3.50 (s, 3H), 3.17 (s, 3H), 2.55 (s, 3H), 1.57 (s, 6H).
486	Int-A-2 + Int-B-46	Ex. 481	43%	¹ H NMR (400 MHz, dmso-d ₆): δ 7.67-7.63 (m, 2H), 7.24-7.21 (m, 1H), 7.05-7.02 (m, 1H), 6.56-6.55 (m, 2H), 4.41-4.39 (m, 1H), 3.53 (s, 3H), 2.54 (s, 3H), 2.17-2.16 (m, 3H), 1.61-1.47 (m, 6H).
487	Int-A-15 + Int-B-46	Ex. 481	50%	¹ H NMR (400 MHz, dmso-d ₆ , 100°C): δ 10.29 (s, 1H), 7.27-7.25 (m, 1H), 7.14 (s, 1H), 7.00-6.97 (m, 1H), 6.86-6.84 (m, 1H), 6.16 (s, 1H), 3.86-3.85 (m, 1H), 2.56 (s, 3H), 2.28-2.23 (m, 6H), 1.60 (s, 6H).
488	Int-A-1 + Int-B-47	Ex. 483	29%	¹ H NMR (400 MHz, dmso-d ₆): δ 11.28 (s, 1H), 7.38-7.37 (m, 2H), 7.22-7.19 (m, 1H), 6.99-6.96 (m, 1H), 6.31 (s, 1H), 6.23 (s, 1H), 3.19 (s, 3H), 2.54 (s, 3H), 1.57 (s, 6H).
493	Int-A-43 + Int-B-47	Ex. 483	29%	¹ H NMR (400 MHz, dmso-d ₆): δ 8.46 (s, 1H), 7.72-7.70 (m, 1H), 7.53 (s, 1H), 7.49-7.46 (m, 1H), 6.43 (s, 1H), 3.57 (s, 3H), 3.22 (s, 3H), 2.60 (s, 3H), 1.58 (s, 6H).
498	Int-A-2 + Int-B-49	Ex. 455	29%	¹ H NMR (400 MHz, dmso-d ₆): δ 7.73-7.71 (m, 1H), 7.65-7.64 (m, 1H), 7.38-7.35 (m, 1H), 7.12 (s, 1H), 6.67-6.66 (m, 1H), 3.59 (s, 3H), 3.22 (s, 3H), 1.65 (s, 3H), 1.49 (s, 3H). (CH ₃ -omitted by DMSO)
499	Int-A-12 + Int-B-49	Ex. 455	36%	¹ H NMR (400 MHz, dmso-d ₆): δ 7.94-7.92 (m, 1H), 7.63 (d, 1H), 7.52-7.48 (m, 1H), 7.40-7.38 (m, 1H), 7.05 (s, 1H), 6.66-6.65 (m, 1H), 3.53 (s, 3H), 3.18 (s, 3H), 1.65 (s, 3H), 1.49 (s, 3H).
501	Int-A-23 + Int-B-49	Ex. 500	39%	¹ H NMR (400 MHz, dmso-d ₆): δ 8.53 (s, 1H), 7.78 (d, 1H), 7.55 (d, 1H), 7.23 (s, 1H), 3.60 (s, 3H), 3.22 (s, 3H), 2.54 (s, 3H), 1.68 (s, 3H), 1.48 (s, 3H).
516	Int-A-1 + Int-B-50	Ex. 500	44%	¹ H NMR (400 MHz, DMSO-d ₆): δ 11.33 (d, 1H), 7.51 (d, 1H), 7.36-7.39 (m, 1H), 7.28-32 (m, 1H), 7.02 (dd, 1H), 6.23 (d, 1H), 2.41 (s, 2H), 1.82 (s, 3H), 1.35 (s, 3H). (CH ₃ -omitted by DMSO)
519	Int-A-23 + Int-B-50	Ex. 518	17%	¹ H NMR (400 MHz, DMSO-d ₆): δ 8.65 (d, 1H), 7.83 (d, 1H), 7.59-7.67 (m, 2H), 3.62 (d, 3H), 2.49 (s, 3H), 1.83 (s, 3H), 1.37 (d, , 3H).

[0330] The example in Table 6 (below) was synthesized in standard chemical reactions which are known to the person skilled in the art:

Ex. #	Intermediates	Yield (%)	¹ H NMR
301	Demethylation of Ex. 300 (DL-Methionin)	19%	¹ H NMR (DMSO- <i>d</i> ₆) δ: 7.53 (d, 1H), 7.42 (d, 1H), 7.22 (dd, 1H), 7.12 (d, 1H), 7.01 (d, 1H), 6.42 (d, 1H), 6.20 - 6.16 (m, 1H), 4.91 (t, 1H), 4.27 (t, 2H), 3.77 (q, 2H), 2.50 (s, 3H), 2.04 (s, 3H), 1.55 - 1.51 (m, 6H)

[0331] The molecular structures and chemical names of the Examples summarized in Tables 3, 4, 5 and 6 are given in Table 7 below:

Ex. #	Structure	Name
48		9-Ethyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
53		9-Ethyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
54		6-Fluoro-8-[6-fluoro-1-(2-methoxy-ethyl)-1H-indol-4-yl]-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
55		2-[6-Fluoro-4-(6-fluoro-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-1-yl]-EtOH

(continued)

Ex. #	Structure	Name
56		6-Fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
57		6-Fluoro-9-methoxy-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
59		1-Cyclopropyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-9-methoxy-4,4-dimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
60		1-Cyclopropyl-6-fluoro-9-methoxy-4,4-dimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
61		2-[4-(1-Cyclopropyl-6-fluoro-9-methoxy-4,4-dimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluoro-1H-indol-1-yl]-EtOH
62		1-Cyclopropyl-6-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-9-methoxy-4,4-dimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
63		1-Cyclopropyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
64		1-Cyclopropyl-6-fluoro-4,4,9-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
65		2-[4-(1-Cyclopropyl-6-fluoro-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluoro-1H-indol-1-yl]-EtOH
66		1-Cyclopropyl-6-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
300		6-Fluoro-8-[1-(2-methoxy-ethyl)-1H-indol-4-yl]-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
301		2-[4-(6-Fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-1-yl]-EtOH

(continued)

Ex. #	Structure	Name
304		6-Fluoro-1,4,4,9-tetramethyl-8-(1-methyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
305		8-(1-Cyclopropyl-1H-indol-4-yl)-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
308		1-Ethyl-6-fluoro-4,4,9-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
316		1-Ethyl-6-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
322		1-Ethyl-6-fluoro-8-[1-(2-methoxy-ethyl)-1H-indol-4-yl]-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
323		1-Ethyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
326		8-[1-(2,2-Difluoro-ethyl)-1H-indol-4-yl]-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
340		6-Fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
349		6-Fluoro-8-[1-(fluoro-methylsulfonyl)-1H-indol-4-yl]-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
350		8-[1-(Ethylsulfonyl)-1H-indol-4-yl]-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
352		6-Fluoro-1,4,4,9-tetramethyl-8-(3-methyl-1H-indazol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
355		1-Ethyl-6-fluoro-4,4,9-trimethyl-8-(3-methyl-1H-indol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
366		6-Fluoro-8-[1-(isopropylsulfonyl)-1H-indol-4-yl]-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
374		1-Ethyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
375		[2-[4-(6-Fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-1-yl]-ethyl]-dimethyl-amine
378		6-Fluoro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
389		1-Ethyl-6-fluoro-4,4,9-trimethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
391		8-[1-(Cyclopropylsulfonyl)-1H-indol-4-yl]-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
447		6,7-Difluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
449		2-[4-(6,7-Difluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluoro-1H-indol-1-yl]-EtOH
451		6,7-Difluoro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
452		6,7-Difluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
453		[6-Fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-MeOH
454		6-Chloro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
456		6-Chloro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
457		6-Chloro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
458		6-Chloro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
459		9-(Difluoro-methyl)-6-fluoro-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
460		9-(Difluoro-methyl)-8-[1-(ethylsulfonyl)-6-fluoro-1H-indol-4-yl]-6-fluoro-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
461		9-(Difluoro-methyl)-8-[1-(ethylsulfonyl)-1H-indol-4-yl]-6-fluoro-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
462		9-(Difluoro-methyl)-6-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
463		2-[4-(6-Chloro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluoro-1H-indol-1-yl]-EtOH
464		2-[4-(9-Ethyl-6-fluoro-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluoro-1H-indol-1-yl]-EtOH
467		9-Cyclopropyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
468		6-Fluoro-8-(1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
469		6-Fluoro-8-(1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
470		6-Fluoro-8-(6-fluoro-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
471		9-Cyclopropyl-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
472		9-Cyclopropyl-6-fluoro-1,4,4-trimethyl-8-(3-methyl-1H-indol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
473		6-Fluoro-9-methoxy-1,4,4-trimethyl-8-[6-(trifluoromethyl)-1H-indol-4-yl]-5H-[1,2,4]triazolo[4,3-a]quinoxaline
474		9-Cyclopropyl-6-fluoro-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
475		6-Fluoro-1,4,4,9-tetramethyl-8-(6-methyl-1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
476		6-Fluoro-9-methoxy-1,4,4-trimethyl-8-(6-methyl-1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
477		4-(6-Fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indole-6-carbonitrile
478		8-(3-Cyclopropyl-5-fluoro-1H-indol-7-yl)-6-fluoro-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
479		[8-[1-(Ethylsulfonyl)-6-fluoro-1H-indol-4-yl]-6-fluoro-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-methyl-amine

(continued)

Ex. #	Structure	Name
480		2-[6-Fluoro-4-(6-fluoro-1,4,4-trimethyl-9-methylamino-5H-[1,2,4]triazolo[4,3-a] quinoxalin-8-yl)-1H-indol-1-yl]-EtOH
482		6-Chloro-9-methoxy-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
484		6-Fluoro-1,4,4,9-tetramethyl-8-(1-methyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
485		6-Fluoro-9-methoxy-1,4,4-trimethyl-8-(1-methyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
486		[6-Fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-methyl-amine
487		[6-Fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-methyl-amine

(continued)

Ex. #	Structure	Name
488		6-Chloro-8-(6-fluoro-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
489		6-Fluoro-8-(7-fluoro-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
491		8-[1-(Cyclopropyl-methylsulfonyl)-1H-indol-4-yl]-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
492		6-Fluoro-9-methoxy-1,4,4-trimethyl-8-[1-methylsulfonyl-6-(trifluoromethyl)-1H-indol-4-yl]-5H-[1,2,4]triazolo[4,3-a]quinoxaline
493		6-Chloro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
494		6-Fluoro-8-(7-fluoro-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
495		6-Fluoro-8-(6-fluoro-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
496		6-Fluoro-8-(6-fluoro-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
497		8-[1-(Cyclopropyl-methylsulfonyl)-6-fluoro-1H-indol-4-yl]-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
498		6,7-Difluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
499		6,7-Difluoro-9-methoxy-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
501		6,7-Difluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
502		6-Chloro-7-fluoro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
503		6-Chloro-7-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
504		6-Fluoro-8-(1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
505		8-[1-(Cyclopropylsulfonyl)-1H-indol-4-yl]-6-fluoro-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
506		8-(6-Chloro-1H-indol-4-yl)-6-fluoro-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
507		6-Fluoro-1,4,4,9-tetramethyl-8-[6-(trifluoromethyl)-1H-indazol-4-yl]-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
508		6-Chloro-7-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
509		6-Chloro-7-fluoro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
510		6-Chloro-7-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
511		8-(1-Cyclopropyl-1H-indol-4-yl)-6-fluoro-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
512		6,7-Difluoro-8-(1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
513		6,7-Difluoro-1,4,4,9-tetramethyl-8-(3-methyl-1H-indol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
514		6,7-Difluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
515		6-Fluoro-8-(5-fluoro-1H-indol-7-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
516		6,7-Difluoro-8-(6-fluoro-1H-indol-4-yl)-1,4,4-trimethyl-9-(trifluoromethyl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
517		6-Fluoro-8-(7-fluoro-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
519		6,7-Difluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4-trimethyl-9-(trifluoromethyl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline
521		6,7-Difluoro-8-(6-fluoro-1-methyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
522		6,7-Difluoro-8-(6-methoxy-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
523		6-Fluoro-8-(6-methoxy-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
524		6-Fluoro-9-methoxy-8-(6-methoxy-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
525		7-Chloro-6-fluoro-8-(6-fluoro-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline
526		7-Chloro-6-fluoro-8-(5-fluoro-3-methyl-1H-indol-7-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxaline

(continued)

Ex. #	Structure	Name
527		7-Chloro-6-fluoro-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxaline

Biological Assays

Agonistic mode of action on the glucocorticoid receptor

[0332] The reporter cell line CHO-Gal4/GR consisted of a chinese hamster ovary (CHO) cell line (*Leibniz Institute DSMZ* - German Collection of Microorganisms and Cell Cultures GmbH: ACC-110) containing a firefly luciferase gene under the control of the GR ligand binding domain fused to the DNA binding domain (DBD) of GAL4 (GAL4 DBD-GR) stably integrated into CHO cells. This cell line was established by stable transfection of CHO cells with a GAL4-UAS-Luciferase reporter construct. In a subsequent step the ligand binding domain of the GR cloned into pIRES2-EGFP-GAL4 containing the DNA binding domain of GAL4 from pFA-AT2 was transfected. This fusion construct activated firefly luciferase expression under the control of a multimerized GAL4 upstream activation sequence (UAS). The signal of the emitted luminescence was recorded by the FLIPR^{TETRA}. This allowed for specific detection of ligand-induced activation of the GR and therefore for the identification of compounds with agonistic properties. The GAL4/LTAS reporter was premixed with a vector that constitutively expressed Renilla luciferase, which served as an internal positive control for transfection efficiency.

[0333] The complete culture medium for the assay was:

- DMEM F-12 (1:1) MIXTURE (LONZA cat. N°: BE04-687F/U1) 500mL
- 5 mL of 100 mM Sodium Pyruvate (LONZA cat. N°: BE12-115E)
- 25 mL of 7.5% Sodium Bicarbonate (LONZA cat. N° BE17-613E)
- 6.5 mL of 1 M Hepes (LONZA cat. N°: BE17-737E)
- 5 mL of 100X Penicillin/Streptomycin (LONZA cat. N° DE17-602E)
- 50 mL of Fetal Bovine Serum (Euroclone cat. N° ECS 0180L)
- 0.25 mL of 10mg/mL Puromycin (InvivoGen cat. : ant-pr-1)
- 0.5 mL of 100 mg/mL Zeocin (InvivoGen cat. : ant-zn-1)

[0334] Cryo-preserved CHO-Gal4/GR cells were suspended in complete medium and 5000 cells/25 μ L/well were seeded into the wells of 384-well polystyrene assay plates (Thermo Scientific, cat.# 4332) and cultured at 37°C, 5% CO₂ and 95% humidity. After 24 hours growth medium was carefully removed and replaced by 30 μ L Opti-MEM (GIBCO, cat.# 31985062) as assay buffer. To test the compounds an 8-point half log compound dilution curve was generated in 100% DMSO starting from a 2mM stock and compounds were then diluted 1:50 in Opti-MEM. 10 μ L of compounds were then added to the wells containing 30 μ L Opti-MEM resulting in a final assay concentration range from 10 μ M to 0.003 μ M in 0.5% DMSO. Compounds were tested at 8 concentrations in quadruplicate data points. Cells were incubated for 6 hour with compounds and beclometasone (Sigma, cat.# Y0000351) as control compound at 37°C, 5% CO₂ and 95% humidity in a total volume of 40 μ L. Finally, cells were lysed with 20 μ L of Triton/Luciferin solution and the signal of the emitted luminescence was recorded at the FLIPR^{TETRA} for 2 minutes.

[0335] The relative efficacy of a compound (% effect) was calculated based on the full effect of the agonist beclom-

etasone:

$$\% \text{ effect} = ((\text{compound} - \text{min}) / (\text{max} - \text{min})) \times 100$$

[min=Opti-MEM only, max=beclometasone]

[0336] To calculate EC₅₀, max, min and slope factor for each compound a concentration response curve was fitted by plotting %effect versus compound concentration using a 4 parameter logistic equation:

$$y = A + (B-A) / (1 + ((1/C)^x)^D)$$

[A=min y, B=max y, C=logEC₅₀, D=slope]

Antagonistic mode of action on the glucocorticoid receptor

[0337] The reporter cell line CHO-Gal4/GR consisted of a chinese hamster ovary (CHO) cell line (*Leibniz Institute DSMZ* - German Collection of Microorganisms and Cell Cultures GmbH: ACC-110) containing a firefly luciferase gene under the control of the GR ligand binding domain fused to the DNA binding domain (DBD) of GAL4 (GAL4 DBD-GR) stably integrated into CHO cells. This cell line was established by stable transfection of CHO cells with a GAL4-UAS-Luciferase reporter construct. In a subsequent step the ligand binding domain of the GR cloned into pIRES2-EGFP-GAL4 containing the DNA binding domain of GAL4 from pFA-AT2 was transfected. This fusion construct activated firefly luciferase expression under the control of a multimerized GAL4 upstream activation sequence (UAS). The signal of the emitted luminescence was recorded by the FLIPR^{TETRA}. This allowed for specific detection of antagonistic properties of compounds by measuring the ligand-induced inhibition of beclometasone-activated GR. The GAL4/LTAS reporter was premixed with a vector that constitutively expressed Renilla luciferase, which served as an internal positive control for transfection efficiency.

[0338] The complete culture medium for the assay was:

- DMEM F-12 (1:1) MIXTURE (LONZA cat. N°: BE04-687F/U1) 500mL
- 5 mL of 100 mM Sodium Pyruvate (LONZA cat. N°: BE12-115E)
- 25 mL of 7.5% Sodium Bicarbonate (LONZA cat. N° BE17-613E)
- 6.5 mL of 1 M Hepes (LONZA cat. N°: BE17-737E)
- 5 mL of 100X Penicillin/Streptomycin (LONZA cat. N° DE17-602E)
- 50 mL of Fetal Bovine Serum (Euroclone cat. N° ECS 0180L)
- 0.25 mL of 10mg/mL Puromycin (InvivoGen cat. : ant-pr-1)
- 0.5 mL of 100 mg/mL Zeocin (InvivoGen cat. : ant-zn-1)

[0339] Cryo-preserved CHO-Gal4/GR cells were suspended in complete medium and 5000 cells/25µl/well were seeded into the wells of 384-well polystyrene assay plates (Thermo Scientific, cat.# 4332) and cultured at 37°C, 5% CO₂ and 95% humidity. After 24 hours growth medium was carefully removed and replaced by 20µl Opti-MEM (GIBCO, cat.# 31985062) as assay buffer. For testing compounds an 8-point half log compound dilution curve was generated in 100% DMSO starting from a 2mM stock and compounds were then diluted 1:50 in Opti-MEM. To test the compounds in the antagonist mode 10µl of compounds were then added to the wells containing 20µl Opti-MEM and incubated for 10 min. After this pre-incubation 10µl of the reference agonist beclometasone (Sigma, cat.# Y0000351) at an EC₅₀ of 2.5 nM were added resulting in a final assay concentration range from 10 µM to 0.003 µM in 0.5% DMSO in a total volume of 40 µl. Compounds were tested at 8 concentrations in quadruplicate data points. Cells were incubated for 6 hour with compounds and mifepristone as control compound (Sigma, cat.# M8046) at 37°C, 5% CO₂ and 95% humidity. Finally, cells were lysed with 20µl of Triton/Luciferin solution and the signal of the emitted luminescence was recorded at the FLIPR^{TETRA} for 2 minutes.

[0340] The relative efficacy of a compound (% effect) was calculated based on the full effect of the antagonist mifepristone:

$$\% \text{ effect} = ((\text{compound} - \text{min}) / (\text{max} - \text{min})) \times 100$$

[min=Opti-MEM only, max=mifepristone]

[0341] To calculate IC₅₀, max, min and slope factor for each compound a concentration response curve was fitted by

plotting %effect versus compound concentration using a 4 parameter logistic equation:

$$y = A + (B-A)/(1 + ((10C)^x)^D)$$

[A=min y, B=max y, C=logIC₅₀, D=slope]

[0342] In Table 7 below, the IC₅₀ or EC₅₀ ranges of the Examples are summarized which were observed in the agonistic assay or the antagonistic assay described above.

Table 7 (A < 100nM; B = 100nM-1μM; C = 1μM-15μM; n.d. = not determined):



38	A	449	B	489	B
48	A	450	A	490	B
49	B	451	B	491	A
53	B	452	A	492	B
54	B	453	A	493	B
55	B	454	A	494	B
56	B	455	A	495	B
57	B	456	A	496	B
58	B	457	A	497	A
59	C	458	B	498	A
60	n.d.	459	A	499	B
61	C	460	A	500	B
62	B	461	B	501	B
63	B	462	A	502	A
64	C	463	A	503	C
65	C	464	B	504	B
66	B	465	A	505	B
300	B	466	B	506	B
301	B	467	B	507	B
304	B	468	A	508	B
305	A	469	B	509	B
308	A	470	B	510	A
316	B	471	A	511	B
322	n.d.	472	B	512	A
323	B	473	B	513	B
326	B	474	B	514	A
340	B	475	B	515	B
349	B	476	B	516	A
350	A	477	B	517	B
352	B	478	A	518	B
355	B	479	B	519	B
366	C	480	B	520	B
374	n.d.	481	B	521	A

(continued)

375	B	482	B	522	B
378	C	483	A	523	B
389	n.d.	484	A	524	B
391	A	485	B	525	A
446	A	486	B	526	A
447	A	487	A	527	B
448	A	488	A		

5

10

Patentkrav**1. Forbindelse ifølge almen formel (II) eller (III)**

hvor

R¹ betegner H; C₁₋₁₀-alkyl; C₃₋₁₀-cycloalkyl; 3- til 7-leddet heterocycloalkyl; aryl; eller 5- eller 6-leddet heteroaryl;

hvor C₃₋₁₀-cycloalkyl, 3- til 7-leddet heterocycloalkyl, aryl og 5- eller 6-leddet heteroaryl eventuelt kan broforbindes via C₁₋₆-alkylen;

R² betegner H; F; Cl; Br; I; CN; C₁₋₁₀-alkyl; C₃₋₁₀-cycloalkyl; O-C₁₋₁₀-alkyl; N(H)(C₁₋₁₀-alkyl); N(C₁₋₁₀-alkyl)₂; C(O)-C₁₋₁₀-alkyl; C(O)-O-C₁₋₁₀-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₁₀-alkyl); C(O)-N(C₁₋₁₀-alkyl)₂; O-C₃₋₁₀-cycloalkyl; N(H)(C₃₋₁₀-cycloalkyl); N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); C(O)-C₃₋₁₀-cycloalkyl; C(O)-O-C₃₋₁₀-cycloalkyl; C(O)-N(H)(C₃₋₁₀-cycloalkyl) eller C(O)-N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl);

hvor C₃₋₁₀-cycloalkyl eventuelt kan broforbindes via C₁₋₆-alkylen;

R³ betegner H; F; Cl; Br; I; CN; C₁₋₁₀-alkyl; C₃₋₁₀-cycloalkyl; O-C₁₋₁₀-alkyl; N(H)(C₁₋₁₀-alkyl); N(C₁₋₁₀-alkyl)₂; C(O)-C₁₋₁₀-alkyl; C(O)-O-C₁₋₁₀-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₁₀-alkyl); C(O)-N(C₁₋₁₀-alkyl)₂; O-C₃₋₁₀-cycloalkyl; N(H)(C₃₋₁₀-cycloalkyl); N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); C(O)-C₃₋₁₀-cycloalkyl; C(O)-O-C₃₋₁₀-cycloalkyl; C(O)-N(H)(C₃₋₁₀-cycloalkyl) eller C(O)-N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl);

hvor C₃₋₁₀-cycloalkyl eventuelt kan broforbindes via C₁₋₆-alkylen;

R⁴ betegner F eller Cl;

R⁵ og R⁶ uafhængigt af hinanden betegner H eller usubstitueret C₁₋₄-alkyl;

X betegner N eller NR⁷;

Z betegner N, NR⁷ eller CR⁹;

med det forbehold at

Z, når X betegner NR⁷, betegner N eller CR⁹;

Z, når X betegner N, betegner NR⁷;

R⁷ betegner H eller L-R⁸; hvor

L betegner binding; S(O); S(O)₂; C₁₋₆-alkylen; C(O); C₁₋₆-alkylen-C(O); C(O)-O; C₁₋₆-alkylen-C(O)-O; C₁₋₆-alkylen-N(H)-C(O); C₁₋₆-alkylen-N(C₁₋₁₀-alkyl)-C(O); C₁₋₆-alkylen-N(H)-C(O)-O; C₁₋₆-alkylen-N(C₁₋₁₀-alkyl)-C(O)-O; O; NH eller N(C₁₋₁₀-alkyl);

R⁸ betegner C₁₋₁₀-alkyl; C₃₋₁₀-cycloalkyl eller 3- til 7-leddet heterocycloalkyl; hvor C₃₋₁₀-cycloalkyl og 3- til 7-leddet heterocycloalkyl eventuelt kan broforbindes via C₁₋₆-alkylen;

R⁹ og R¹⁰ uafhængigt af hinanden betegner H; F; Cl; Br; I; CN; C₁₋₁₀-alkyl; C₃₋₁₀-cycloalkyl; 3- til 7-leddet heterocycloalkyl; S(O)-(C₁₋₁₀-alkyl); S(O)-(C₃₋₁₀-cycloalkyl); S(O)-(3- til 7-leddet heterocycloalkyl); S(O)₂-(C₁₋₁₀-alkyl); S(O)₂-(C₃₋₁₀-cycloalkyl); S(O)₂-(3- til 7-leddet heterocycloalkyl); P(O)-(C₁₋₁₀-alkyl)₂; P(O)(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); P(O)(C₁₋₁₀-alkyl)(3- til 7-leddet heterocycloalkyl); P(O)-(O-C₁₋₁₀-alkyl)₂; P(O)(O-C₁₋₁₀-alkyl)(O-C₃₋₁₀-cycloalkyl); P(O)(O-C₁₋₁₀-alkyl)(O-(3- til 7-leddet heterocycloalkyl)); O-C₁₋₁₀-alkyl; S-C₁₋₁₀-alkyl; N(H)(C₁₋₁₀-alkyl), N(C₁₋₁₀-alkyl)₂; C(O)-C₁₋₁₀-alkyl; C(O)-O-C₁₋₁₀-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₁₀-alkyl); C(O)-N(C₁₋₁₀-alkyl)₂; O-C₃₋₁₀-cycloalkyl; N(H)(C₃₋₁₀-cycloalkyl), N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); C(O)-C₃₋₁₀-cycloalkyl; C(O)-O-C₃₋₁₀-cycloalkyl; C(O)-N(H)(C₃₋₁₀-cycloalkyl); C(O)-N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); O-3- til 7-leddet heterocycloalkyl; N(H)(3- til 7-leddet heterocycloalkyl), N(C₁₋₁₀-alkyl)(3- til 7-leddet heterocycloalkyl); C(O)-3- til 7-leddet heterocycloalkyl; C(O)-O-(3- til 7-leddet heterocycloalkyl); C(O)-N(H)(3- til 7-leddet heterocycloalkyl) eller C(O)-N(C₁₋₁₀-alkyl)(3- til 7-leddet heterocycloalkyl); hvor C₃₋₁₀-cycloalkyl og 3- til 7-leddet heterocycloalkyl eventuelt kan broforbindes via C₁₋₆-alkylen;

R¹¹ betegner F; Cl; Br; I; CN; C₁₋₁₀-alkyl; O-C₁₋₁₀-alkyl; NO₂; OH, NH₂; C₃₋₁₀-cycloalkyl; 3- til 7-leddet heterocycloalkyl; S(O)-(C₁₋₁₀-alkyl); S(O)-(C₃₋₁₀-cycloalkyl); S(O)-(3- til 7-leddet heterocycloalkyl); S(O)₂-(C₁₋₁₀-alkyl); S(O)₂-(C₃₋₁₀-cycloalkyl); S(O)₂-(3- til 7-leddet heterocycloalkyl); P(O)-(C₁₋₁₀-alkyl)₂; P(O)(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); P(O)(C₁₋₁₀-alkyl)(3- til 7-leddet heterocycloalkyl); P(O)-(O-C₁₋₁₀-alkyl)₂; P(O)(O-C₁₋₁₀-alkyl)(O-C₃₋₁₀-cycloalkyl); P(O)(O-C₁₋₁₀-alkyl)(O-(3- til 7-leddet heterocycloalkyl)); O-C₁₋₁₀-alkyl; N(H)(C₁₋₁₀-alkyl), N(C₁₋₁₀-alkyl)₂; C(O)-C₁₋₁₀-alkyl; C(O)-O-C₁₋₁₀-alkyl; C(O)-NH₂; C(O)-

N(H)(C₁₋₁₀-alkyl); C(O)-N(C₁₋₁₀-alkyl)₂; O-C₃₋₁₀-cycloalkyl; N(H)(C₃₋₁₀-cycloalkyl), N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); C(O)-C₃₋₁₀-cycloalkyl; C(O)-O-C₃₋₁₀-cycloalkyl; C(O)-N(H)(C₃₋₁₀-cycloalkyl); C(O)-N(C₁₋₁₀-alkyl)(C₃₋₁₀-cycloalkyl); O-3- til 7-leddet heterocycloalkyl; N(H)(3- til 7-leddet heterocycloalkyl), N(C₁₋₁₀-alkyl)(3- til 7-leddet heterocycloalkyl); C(O)-3- til 7-leddet heterocycloalkyl; C(O)-O-(3- til 7-leddet heterocycloalkyl); C(O)-N(H)(3- til 7-leddet heterocycloalkyl) eller C(O)-N(C₁₋₁₀-alkyl)(3- til 7-leddet heterocycloalkyl); hvor C₃₋₁₀-cycloalkyl og 3- til 7-leddet heterocycloalkyl eventuelt kan broforbindes via C₁₋₆-alkylen;

n betegner 0, 1, 2 eller 3;

hvor C₁₋₁₀-alkyl, C₁₋₄-alkyl og C₁₋₆-alkylen i hvert tilfælde uafhængigt af hinanden er ligekædet eller forgrenet, mættet eller umættet;

hvor C₁₋₁₀-alkyl, C₁₋₄-alkyl, C₁₋₆-alkylen, C₃₋₁₀-cycloalkyl og 3- til 7-leddet heterocycloalkyl i hvert tilfælde uafhængigt af hinanden er usubstitueret eller mono- eller polysubstitueret med én eller flere substituent, der er valgt blandt F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; CF₂Cl; CFC₂Cl₂; C(O)-C₁₋₆-alkyl; C(O)-OH; C(O)-OC₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; =O; OCF₃; OCF₂H; OCFH₂; OCF₂Cl; OCFCl₂; O-C₁₋₆-alkyl; O-C(O)-C₁₋₆-alkyl; O-C(O)-O-C₁₋₆-alkyl; O-(CO)-N(H)(C₁₋₆-alkyl); O-C(O)-N(C₁₋₆-alkyl)₂; O-S(O)₂-NH₂; O-S(O)₂-N(H)(C₁₋₆-alkyl); O-S(O)₂-N(C₁₋₆-alkyl)₂; NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; N(H)-C(O)-C₁₋₆-alkyl; N(H)-C(O)-O-C₁₋₆-alkyl; N(H)-C(O)-NH₂; N(H)-C(O)-N(H)(C₁₋₆-alkyl); N(H)-C(O)-N(C₁₋₆-alkyl)₂; N(C₁₋₆-alkyl)-C(O)-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-C(O)-O-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-C(O)-NH₂; N(C₁₋₆-alkyl)-C(O)-N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)-C(O)-N(C₁₋₆-alkyl)₂; N(H)-S(O)₂OH; N(H)-S(O)₂-C₁₋₆-alkyl; N(H)-S(O)₂-O-C₁₋₆-alkyl; N(H)-S(O)₂-NH₂; N(H)-S(O)₂-N(H)(C₁₋₆-alkyl); N(H)-S(O)₂-N(C₁₋₆-alkyl)₂; N(C₁₋₆-alkyl)-S(O)₂OH; N(C₁₋₆-alkyl)-S(O)₂-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-S(O)₂-O-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-S(O)₂-NH₂; N(C₁₋₆-alkyl)-S(O)₂-N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)-S(O)₂-N(C₁₋₆-alkyl)₂; SCF₃; SCF₂H; SCFH₂; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; S(O)₂-OH; S(O)₂-O-C₁₋₆-alkyl; S(O)₂-NH₂; S(O)₂-N(H)(C₁₋₆-alkyl); S(O)₂-N(C₁₋₆-alkyl)₂; C₃₋₆-cycloalkyl; 3- til 6-leddet heterocycloalkyl; phenyl; 5- eller 6-leddet heteroaryl; O-C₃₋₆-cycloalkyl; O-(3- til 6-leddet heterocycloalkyl); O-phenyl; O-(5- eller 6-leddet heteroaryl); C(O)-C₃₋₆-cycloalkyl; C(O)-(3- til 6-leddet heterocycloalkyl); C(O)-phenyl; C(O)-(5- eller 6-leddet heteroaryl); S(O)₂-(C₃₋₆-cycloalkyl); S(O)₂-

(3- til 6-leddet heterocycloalkyl); S(O)₂-phenyl eller S(O)₂-(5- eller 6-leddet heteroaryl);

hvor aryl og 5- eller 6-leddet heteroaryl i hvert tilfælde uafhængigt af hinanden er usubstitueret eller mono- eller polysubstitueret med én eller flere substituent, der er valgt blandt F; Cl; Br; I; CN; C₁₋₆-alkyl; CF₃; CF₂H; CFH₂; CF₂Cl; CFC₂Cl; C₁₋₄-alkylen-CF₃; C₁₋₄-alkylen-CF₂H; C₁₋₄-alkylen-CFH₂; C(O)-C₁₋₆-alkyl; C(O)-OH; C(O)-OC₁₋₆-alkyl; C(O)-N(H)(OH); C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; OH; OCF₃; OCF₂H; OCFH₂; OCF₂Cl; OCFC₂Cl; O-C₁₋₆-alkyl; O-C₃₋₆-cycloalkyl; O-(3- til 6-leddet heterocycloalkyl); NH₂; N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)₂; N(H)-C(O)-C₁₋₆-alkyl; N(C₁₋₆-alkyl)-C(O)-C₁₋₆-alkyl; N(H)-C(O)-NH₂; N(H)-C(O)-N(H)(C₁₋₆-alkyl); N(H)-C(O)-N(C₁₋₆-alkyl)₂; N(C₁₋₆-alkyl)-C(O)-N(H)(C₁₋₆-alkyl); N(C₁₋₆-alkyl)-C(O)-N(C₁₋₆-alkyl)₂; N(H)-S(O)₂-C₁₋₆-alkyl; SCF₃; S-C₁₋₆-alkyl; S(O)-C₁₋₆-alkyl; S(O)₂-C₁₋₆-alkyl; S(O)₂-NH₂; S(O)₂-N(H)(C₁₋₆-alkyl); S(O)₂-N(C₁₋₆-alkyl)₂; C₃₋₆-cycloalkyl; C₁₋₄-alkylen-C₃₋₆-cycloalkyl; 3- til 6-leddet heterocycloalkyl; C₁₋₄-alkylen-(3- til 6-leddet heterocycloalkyl); phenyl eller 5- eller 6-leddet heteroaryl;

i form af den frie forbindelse eller et fysiologisk acceptabelt salt deraf.

2. Forbindelse ifølge krav 1, hvor R⁵ og R⁶ begge betegner CH₃.

3. Forbindelse ifølge krav 1 eller 2, hvor X betegner NR⁷, og Z betegner N eller CR⁹.

4. Forbindelse ifølge et hvilket som helst af de foregående krav, hvor

R¹ betegner H; C₁₋₆-alkyl; C₃₋₆-cycloalkyl; 3- til 6-leddet heterocycloalkyl; phenyl; eller 5- eller 6-leddet heteroaryl;

hvor C₃₋₆-cycloalkyl, 3- til 6-leddet heterocycloalkyl, phenyl og 5- eller 6-leddet heteroaryl eventuelt kan broforbindes via C₁₋₄-alkylen.

5. Forbindelse ifølge et hvilket som helst af de foregående krav, hvor

R² betegner H; F; Cl; Br; CN; C₁₋₆-alkyl; C₃₋₆-cycloalkyl; O-C₁₋₆-alkyl; C(O)-NH₂; C(O)-N(H)(C₁₋₆-alkyl); C(O)-N(C₁₋₆-alkyl)₂; C(O)-N(H)(C₃₋₆-cycloalkyl) eller C(O)-N(C₁₋₆-alkyl)(C₃₋₆-cycloalkyl); hvor C₃₋₆-cycloalkyl eventuelt kan broforbindes via C₁₋₄-alkylen; og/eller

R^3 betegner H; F; Cl; Br; CN; C_{1-6} -alkyl; C_{3-6} -cycloalkyl; O- C_{1-6} -alkyl; C(O)-NH₂; C(O)-N(H)(C_{1-6} -alkyl); C(O)-N(C_{1-6} -alkyl)₂; C(O)-N(H)(C_{3-6} -cycloalkyl) eller C(O)-N(C_{1-6} -alkyl)(C_{3-6} -cycloalkyl); hvor C_{3-6} -cycloalkyl eventuelt kan broforbindes via C_{1-4} -alkylen.

6. Forbindelse ifølge et hvilket som helst af de foregående krav, hvor R^7 betegner H.
7. Forbindelse ifølge et hvilket som helst af kravene 1 til 5, hvor R^7 betegner L- R^8 ; hvor L betegner binding; S(O); S(O)₂; C_{1-4} -alkylen; C(O); C_{1-4} -alkylen-C(O); C(O)-O; C_{1-4} -alkylen-C(O)-O; C_{1-4} -alkylen-N(H)-C(O) eller C_{1-4} -alkylen-N(H)-C(O)-O; R^8 betegner C_{1-6} -alkyl; C_{3-6} -cycloalkyl eller 3- til 6-leddet heterocycloalkyl; hvor C_{3-6} -cycloalkyl og 3- til 6-leddet heterocycloalkyl eventuelt kan broforbindes via C_{1-4} -alkylen.
8. Forbindelse ifølge et hvilket som helst af de foregående krav, hvor R^9 betegner H; F; CN; methyl; ethyl; n-propyl; 2-propyl; CF₃; CH₂CF₃; CHF₂; CH₂CHF₂; CH₂F; CH₂CH₂F; S(O)-CH₃; S(O)-CH₂CH₃; S(O)-CH₂CH₂CH₃; S(O)-CH(CH₃)₂; S(O)₂-CH₃; S(O)₂-CH₂CH₃; S(O)₂-CH₂CH₂CH₃ eller S(O)₂-CH(CH₃)₂.
9. Forbindelse ifølge et hvilket som helst af de foregående krav, hvor R^{10} betegner H; F; Cl; Br; CN; C_{1-6} -alkyl; C_{3-6} -cycloalkyl; 3- til 6-leddet heterocycloalkyl; P(O)-(C₁₋₆-alkyl)₂; P(O)(C₁₋₆-alkyl)(C_{3-6} -cycloalkyl); P(O)(C₁₋₆-alkyl)(3- til 6-leddet heterocycloalkyl); P(O)-(O- C_{1-6} -alkyl)₂; P(O)(O- C_{1-6} -alkyl)(O- C_{3-6} -cycloalkyl); P(O)(O- C_{1-6} -alkyl)(O-(3- til 6-leddet heterocycloalkyl)).
10. Forbindelse ifølge et hvilket som helst af de foregående krav, hvor R^{11} betegner F; Cl; Br; I; CN; C_{1-6} -alkyl eller O- C_{1-6} -alkyl; og/eller n betegner 0, 1 eller 2.
11. Forbindelse ifølge et hvilket som helst af de foregående krav, der er valgt fra gruppen bestående af

38	8-(3-cyclopropyl-5-fluor-1H-indol-7-yl)-6-fluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
-----------	---

48	9-ethyl-6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
49	9-ethyl-6-fluor-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
53	9-ethyl-6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
54	6-fluor-8-[6-fluor-1-(2-methoxy-ethyl)-1H-indol-4-yl]-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
55	2-[6-fluor-4-(6-fluor-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-1-yl]-EtOH
56	6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
57	6-fluor-9-methoxy-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
58	8-[1-(2,2-difluor-ethyl)-6-fluor-1H-indol-4-yl]-6-fluor-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
59	1-cyclopropyl-6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-9-methoxy-4,4-dimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
60	1-cyclopropyl-6-fluor-9-methoxy-4,4-dimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
61	2-[4-(1-cyclopropyl-6-fluor-9-methoxy-4,4-dimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluor-1H-indol-1-yl]-EtOH
62	1-cyclopropyl-6-fluor-8-(5-fluor-3-methyl-1H-indol-7-yl)-9-methoxy-4,4-dimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
63	1-cyclopropyl-6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
64	1-cyclopropyl-6-fluor-4,4,9-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin

65	2-[4-(1-cyclopropyl-6-fluor-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluor-1H-indol-1-yl]-EtOH
66	1-cyclopropyl-6-fluor-8-(5-fluor-3-methyl-1H-indol-7-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
300	6-fluor-8-[1-(2-methoxy-ethyl)-1H-indol-4-yl]-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
301	2-[4-(6-fluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-1-yl]-EtOH
304	6-fluor-1,4,4,9-tetramethyl-8-(1-methyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
305	8-(1-cyclopropyl-1H-indol-4-yl)-6-fluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
308	1-ethyl-6-fluor-4,4,9-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
316	1-ethyl-6-fluor-8-(5-fluor-3-methyl-1H-indol-7-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
322	1-ethyl-6-fluor-8-[1-(2-methoxy-ethyl)-1H-indol-4-yl]-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
323	1-ethyl-6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
326	8-[1-(2,2-difluor-ethyl)-1H-indol-4-yl]-6-fluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
340	6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
349	6-fluor-8-[1-(fluor-methylsulfonyl)-1H-indol-4-yl]-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
350	8-[1-(ethylsulfonyl)-1H-indol-4-yl]-6-fluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin

352	6-fluor-1,4,4,9-tetramethyl-8-(3-methyl-1H-indazol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
355	1-ethyl-6-fluor-4,4,9-trimethyl-8-(3-methyl-1H-indol-7-yl)-5H-[1,2,4]triazolo [4,3-a]quinoxalin
366	6-fluor-8-[1-(isopropylsulfonyl)-1H-indol-4-yl]-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
374	1-ethyl-6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indazol-4-yl)-4,4,9-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
375	[2-[4-(6-fluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-1-yl]-ethyl]-dimethyl-amin
378	6-fluor-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
389	1-ethyl-6-fluor-4,4,9-trimethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo [4,3-a]quinoxalin
391	8-[1-(cyclopropylsulfonyl)-1H-indol-4-yl]-6-fluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
446	8-[1-(ethylsulfonyl)-6-fluor-1H-indol-4-yl]-6,7-difluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
447	6,7-difluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
448	6,7-difluor-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
449	2-[4-(6,7-difluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluor-1H-indol-1-yl]-EtOH
450	8-(1-cyclopropyl-1H-indol-4-yl)-6-fluor-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-carbonitril
451	6,7-difluor-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo [4,3-a]quinoxalin

452	6,7-difluor-8-(6-fluor-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
453	[6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-MeOH
454	6-chlor-8-(5-fluor-3-methyl-1H-indol-7yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
455	9-(difluor-methyl)-6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
456	6-chlor-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
457	6-chlor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
458	6-chlor-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
459	9-(difluor-methyl)-6-fluor-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
460	9-(difluor-methyl)-8-[1-(ethylsulfonyl)-6-fluor-1H-indol-4-yl]-6-fluor-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
461	9-(difluor-methyl)-8-[1-(ethylsulfonyl)-1H-indol-4-yl]-6-fluor-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
462	9-(difluor-methyl)-6-fluor-8-(5-fluor-3-methyl-1H-indol-7-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
463	2-[4-(6-chlor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluor-1H-indol-1-yl]-EtOH
464	2-[4-(9-ethyl-6-fluor-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-6-fluor-1H-indol-1-yl]-EtOH
465	6-fluor-8-(6-fluor-H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin

466	9-(difluor-methyl)-6-fluor-1,4,4-trimethyl-8-(3-methyl-1H-indazol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
467	9-cyclopropyl-6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
468	6-fluor-8-(1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
469	6-fluor-8-(1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
470	6-fluor-8-(6-fluor-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
471	9-cyclopropyl-6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
472	9-cyclopropyl-6-fluor-1,4,4-trimethyl-8-(3-methyl-1H-indol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
473	6-fluor-9-methoxy-1,4,4-trimethyl-8-[6-(trifluormethyl)-1H-indol-4-yl]-5H-[1,2,4]triazolo[4,3-a]quinoxalin
474	9-cyclopropyl-6-fluor-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
475	6-fluor-1,4,4,9-tetramethyl-8-(6-methyl-1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
476	6-fluor-9-methoxy-1,4,4-trimethyl-8-(6-methyl-1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
477	4-(6-fluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-6-carbonitril
478	8-(3-cyclopropyl-5-fluor-1H-indol-7-yl)-6-fluor-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
479	[8-[1-(ethylsulfonyl)-6-fluor-1H-indol-4-yl]-6-fluor-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-methyl-amin

480	2-[6-fluor-4-(6-fluor-1,4,4-trimethyl-9-methylamino-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-1-yl]-EtOH
481	[8-(1-cyclopropyl-1H-indol-4-yl)-6-fluor-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-methyl-amin
482	6-chlor-9-methoxy-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
483	6-chlor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
484	6-fluor-1,4,4,9-tetramethyl-8-(1-methyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
485	6-fluor-9-methoxy-1,4,4-trimethyl-8-(1-methyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
486	[6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-methyl-amin
487	[6-fluor-8-(5-fluor-3-methyl-1H-indol-7-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-9-yl]-methyl-amin
488	6-chlor-8-(6-fluor-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
489	6-fluor-8-(7-fluor-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
490	6-fluor-8-(7-fluor-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
491	8-[1-(cyclopropyl-methylsulfonyl)-1H-indol-4-yl]-6-fluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
492	6-fluor-9-methoxy-1,4,4-trimethyl-8-[1-methylsulfonyl-6-(trifluormethyl)-1H-indol-4-yl]-5H-[1,2,4]triazolo[4,3-a]quinoxalin
493	6-chlor-8-(6-fluor-1-methylsulfonyl-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin

494	6-fluor-8-(7-fluor-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
495	6-fluor-8-(6-fluor-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
496	6-fluor-8-(6-fluor-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
497	8-[1-(cyclopropyl-methylsulfonyl)-6-fluor-1H-indol-4-yl]-6-fluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
498	6,7-difluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
499	6,7-difluor-9-methoxy-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
500	6,7-difluor-1,4,4-trimethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-9-(trifluormethyl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
501	6,7-difluor-8-(6-fluor-1-methylsulfonyl-1H-indazol-4-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
502	6-chlor-7-fluor-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
503	6-chlor-7-fluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
504	6-fluor-8-(1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
505	8-[1-(cyclopropylsulfonyl)-1H-indol-4-yl]-6-fluor-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
506	8-(6-chlor-1H-indol-4-yl)-6-fluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
507	6-fluor-1,4,4,9-tetramethyl-8-[6-(trifluormethyl)-1H-indazol-4-yl]-5H-[1,2,4]triazolo[4,3-a]quinoxalin

508	6-chlor-7-fluor-8-(5-fluor-3-methyl-1H-indol-7-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
509	6-chlor-7-fluor-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
510	6-chlor-7-fluor-8-(6-fluor-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
511	8-(1-cyclopropyl-1H-indol-4-yl)-6-fluor-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
512	6,7-difluor-8-(1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
513	6,7-difluor-1,4,4,9-tetramethyl-8-(3-methyl-1H-indol-7-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
514	6,7-difluor-8-(5-fluor-3-methyl-1H-indol-7-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
515	6-fluor-8-(5-fluor-1H-indol-7-yl)-9-methoxy-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
516	6,7-difluor-8-(6-fluor-1H-indol-4-yl)-1,4,4-trimethyl-9-(trifluormethyl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
517	6-fluor-8-(7-fluor-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
518	6,7-difluor-8-(6-fluor-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-9-(trifluormethyl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
519	6,7-difluor-8-(6-fluor-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4-trimethyl-9-(trifluormethyl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin
520	4-(6,7-difluor-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin-8-yl)-1H-indol-7-carbonitril
521	6,7-difluor-8-(6-fluor-1-methyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin

522	6,7-difluor-8-(6-methoxy-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
523	6-fluor-8-(6-methoxy-1-methylsulfonyl-1H-indol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
524	6-fluor-9-methoxy-8-(6-methoxy-1-methylsulfonyl-1H-indol-4-yl)-1,4,4-trimethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
525	7-chlor-6-fluor-8-(6-fluor-1-methylsulfonyl-1H-indazol-4-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
526	7-chlor-6-fluor-8-(5-fluor-3-methyl-1H-indol-7-yl)-1,4,4,9-tetramethyl-5H-[1,2,4]triazolo[4,3-a]quinoxalin
527	7-chlor-6-fluor-1,4,4,9-tetramethyl-8-(1-methylsulfonyl-1H-indazol-4-yl)-5H-[1,2,4]triazolo[4,3-a]quinoxalin

i form af den frie forbindelse eller et fysiologisk acceptabelt salt deraf.

12. Farmaceutisk doseringsform, der omfatter en forbindelse ifølge et hvilket som helst af kravene 1 til 11.

13. Forbindelse ifølge et hvilket som helst af kravene 1 til 11 til anvendelse ved behandling og/eller profylakse af smerter og/eller inflammation.

14. Forbindelse til anvendelse ifølge krav 13, hvor anvendelsen er ved behandling og/eller profylakse af inflammatoriske smerter.