





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/00	A2	(11) International Publication Number: WO 00/09103 (43) International Publication Date: 24 February 2000 (24.02.00)
(21) International Application Number: PCT/US99/18231 (22) International Filing Date: 12 August 1999 (12.08.99) (30) Priority Data: 09/134,471 14 August 1998 (14.08.98) US (71) Applicant: GUILFORD PHARMACEUTICALS INC. [US/US]; 6611 Tributary Street, Baltimore, MD 21224 (US). (72) Inventors: ROSS, Douglas, T.; 316 South Main Street, North Wales, PA 19454 (US). SAUER, Hansjorg; 10617 Lorrain Avenue, Silver Spring, MD 20901 (US). HAMILTON, Gregory, S.; 6501 Frederick Road, Catonsville, MD 21228 (US). STEINER, Joseph, P.; 4150 Louisville Road, Finksburg, MD 21048 (US). (74) Agent: NATH, Gary, M.; Nath & Associates, 6th Floor, 1030 15th Street, N.W., Washington, DC 20005-1503 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: N-LINKED SULFONAMIDES OF N-HETEROCYCLIC CARBOXYLIC ACIDS OR ISOSTERES FOR VISION AND MEMORY DISORDERS (57) Abstract This invention relates to novel compositions and uses of an N-linked sulfonamide of an N-heterocyclic carboxylic acid or isostere thereof for treating a vision disorder or improving vision or treating memory impairment or enhancing memory performance in an animal. GPI 1046 protects ganglion cells against degeneration due to 1 hour of retinal ischemia Fluorogold labelled retinal ganglion cells in wholemount, 28 days after ischemic episode  A. Labeled retinal ganglion cells in the Normal central retina  B. 1 hour of retinal ischemia produces extensive loss of ganglion cells  B. Administration of GPI 1046 1 hour before retinal ischemia and for 4 days subsequently produces significant protection of vulnerable retinal ganglion cells		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

N-LINKED SULFONAMIDES OF N-HETEROCYCLIC CARBOXYLIC ACIDS OR ISOSTERES
FOR VISION AND MEMORY DISORDERS

BACKGROUND OF THE INVENTION

5

1. Field of Invention

This invention relates to pharmaceutical compositions and methods for treating vision loss, preventing vision degeneration, and promoting vision regeneration ("neopsis") using low molecular weight, small molecule derivatives.

10

2. Description of Related Art

The visual system is composed of the eyes, ocular adnexa and the visual pathways. Dysfunction of the visual system may lead to permanent or temporary visual impairment, i.e. a deviation from normal in one or more functions of the eye. Visual impairment manifests itself in various ways and includes a broad range of visual dysfunctions and disturbances. Without limitation, these dysfunctions and disturbances include partial or total loss of vision, the need for correction of visual acuity for objects near and far, loss of visual field, impaired ocular motility without diplopia (double vision), impaired or skewed color perception, limited adaptation to light and dark, diminished accommodation, metamorphopsic

15

20

25

distortion, impaired binocular vision, paresis of accommodation, iridoplegia, entropion, ectropion, epiphora, lagophthalmos, and scarring. See Physicians' Desk Reference (PDR) for Ophthalmology, 16th Edition, 6:47 (1988). The visual system may be adversely affected by various ophthalmologic disorders, diseases, injuries, and complications, including, without limitation, genetic disorders; [non-genetic disorders;] disorders associated with aging or degenerative diseases; disorders correlating to physical injury to the eye, head, or other parts of the body resulting from external forces; disorders resulting from environmental factors; disorders resulting from a broad range of diseases; and combinations of any of the above.

The visual system is a complex system composed of numerous components. Visual impairment can involve the entire visual system, any one component, or any combination of components, depending upon the precise nature of the circumstances. The eye is composed of a lens, which is suspended in the zonules of Zinn and is focused by the ciliary body. The ciliary body also secretes aqueous humor, which fills the posterior chamber, passes through the pupil into the anterior chamber, then drains primarily via the canal of Schlemm. The iris regulates the quantity of light

entering the eye by adjusting the size of its central opening, the pupil. A visual image is focused onto the retina, the fovea centralis being the retinal area of sharpest visual acuity. The conjunctiva is the
5 mucus membrane which lines the eyelids and the eyeball, and ends abruptly at the limbus conjunctivae, the edge of the conjunctiva overlapping the cornea. The cornea is the clear, transparent anterior portion of the fibrous coat of the eye; it is important in
10 light refraction and is covered with an epithelium that differs in many respects from the conjunctival epithelium.

The retina is the innermost, light sensitive portion of the eye, containing two types of
15 photoreceptors, cones, which are responsible for color vision in brighter light, and rods, which are essential for vision in dim light but do not perceive colors. After light passes through the cornea, lens system, and the vitreous humor, it enters the retina
20 from the inside; that is, it passes through the ganglion cells and nerve fibers, the inner and outer plexiform layers, the inner and outer nuclear layers, and the internal and external limiting membranes before it finally reaches the layer of photoreceptors
25 located near the outside of the retina, just inside the outermost pigment epithelium layer. The cells of

the pigment epithelium layer act as an anatomical barrier to liquids and substances located outside of the eye, forming the "blood-retina" barrier, and provide nourishment, oxygen, a source of functionally useful substances like vitamin A, and phagocytosis of decomposition products to photoreceptor cells. There is no anatomical connection between the pigment epithelium and the photoreceptor layer, permitting separation of the layers in some pathological situations.

When rods or cones are excited by light, signals are transmitted through successive neurons in the retina itself, into the optic nerve fibers, and ultimately to the cerebral cortex. Both rods and cones contain molecules that decompose on exposure to light and, in the process, excite the nerve fibers leading from the eye. The molecule in rods is rhodopsin. The three light-sensitive molecules in cones, collectively called iodopsin, have compositions only slightly different from that of rhodopsin and are maximally excited by red, blue, or green light, respectively.

Neither rods nor cones generate action potentials. Rather, the light-induced membrane hyperpolarization generated in the outer, photosensitive segment of a rod or cone cell is

transmitted from the outer segment through the inner segment to the synaptic body by direct conduction of the electrical voltage itself, a process called electrotonic conduction. At the synaptic body, the membrane potential controls the release of an unknown transmitter molecule. In low light, rod and cone cell membranes are depolarized and the rate of transmitter release is greatest. Light-induced hyperpolarization causes a marked decrease in the release of transmitter molecules.

The transmitters released by rod and cone cells induce signals in the bipolar neurons and horizontal cells. The signals in both these cells are also transmitted by electrotonic conduction and not by action potential.

The rod bipolar neurons connect with as many as 50 rod cells, while the dwarf and diffuse bipolar cells connect with one or several cone cells. A depolarizing bipolar cell is stimulated when its connecting rods or cones are exposed to light. The release of transmitter molecules inhibits the depolarizing bipolar cell. Therefore, in the dark, when the rods and cones are secreting large quantities of transmitter molecules, the depolarizing bipolar cells are inhibited. In the light, the decrease in release of transmitter molecules from the rods and

cones reduces the inhibition of the bipolar cell, allowing it to become excited. In this manner, both positive and negative signals can be transmitted through different bipolar cells from the rods and
5 cones to the amacrine and ganglion cells.

As their name suggests, horizontal cells project horizontally in the retina, where they may synapse with rods, cones, other horizontal cells, or a combination of cells types. The function of
10 horizontal cells is unclear, although some mechanism in the convergence of photoreceptor signaling has been postulated.

All types of bipolar cells connect with ganglion cells, which are of two primary types. A-type
15 ganglion cells predominately connect with rod bipolar cells, while B-type ganglion cells predominately connect with dwarf and diffuse bipolar cells. It appears that A-type ganglion cells are sensitive to contrast, light intensity, and perception of movement,
20 while B-type ganglion cells appear more concerned with color vision and visual acuity.

Like horizontal cells, the Amacrine cells horizontally synapse with several to many other cells, in this case bipolar cells, ganglion cells, and other
25 Amacrine cells. The function of Amacrine cells is also unclear.

The axons of ganglion cells carry signals into the nerve fiber layer of the eye, where the axons converge into fibers which further converge at the optic disc, where they exit the eye as the optic nerve. The ganglion cells transmit their signals through the optic nerve fibers to the brain in the form of action potentials. These cells, even when unstimulated, transmit continuous nerve impulses at an average, baseline rate of about 5 per second. The visual signal is superimposed onto this baseline level of ganglion cell stimulation. It can be either an excitatory signal, with the number of impulses increasing above the baseline rate, or an inhibitory signal, with the number of nerve impulses decreasing below the baseline rate.

As part of the central nervous system, the eye is in some ways an extension of the brain; as such, it has a limited capacity for regeneration. This limited regeneration capacity further complicates the challenging task of improving vision, resolving dysfunction of the visual system, and/or treating or preventing ophthalmologic disorders. Many disorders of the eye, such as retinal photic injury, retinal ischemia-induced eye injury, age-related macular degeneration, free radical-induced eye diseases, as well as numerous other disorders, are considered to be

entirely untreatable. Other ophthalmologic disorders, e.g., disorders causing permanent visual impairment, are corrected only by the use of ophthalmic devices and/or surgery, with varying degrees of success.

5 The immunosuppressant drugs FK506, rapamycin, and cyclosporin are well known as potent T-cell specific immunosuppressants, and are effective against autoimmunity, transplant or graft rejection, inflammation, allergic responses, other autoimmune or
10 immune-mediated diseases, and infectious diseases. It has been disclosed that application of Cyclosporin, FK-506, Rapamycin, Buspirone, Spiperone, and/or their derivatives are effective in treating some ophthalmologic disorders of these types. Several
15 ophthalmologic disorders or vision problems are known to be associated with autoimmune and immunologically-mediated activities; hence, immunomodulatory compounds are expected to demonstrate efficacy for treating those types of ophthalmologic disorders or vision
20 problems.

 The effects of FK506, Rapamycin, and related agents in the treatment of ophthalmologic diseases are disclosed in several U.S. patents (Goulet et al., U.S. Patent No. 5,532,248; Mochizuki et al., U.S. Patent
25 No. 5,514,686; Luly et al., U.S. Patent No. 5,457,111; Russo et al., U.S. Patent No. 5,441,937; Kulkarni,

U.S. Patent No. 5,387,589; Asakura et al., U.S. Patent No. 5,368,865; Goulet et al., U.S. Patent No. 5,258,389; Armistead et al., U.S. Patent No. 5,192,773; Goulet et al., U.S. Patent No. 5,189,042; and Fehr, U.S. Patent No. 5,011,844). These patents claim FK506 or Rapamycin related compounds and disclose the known use of FK506 or Rapamycin related compounds in the treatment of ophthalmologic disorders in association with the known immunosuppressive effects of FK506 and Rapamycin. The compounds disclosed in these patents are relatively large. Further, the cited patents relate to immunomodulatory compounds limited to treating autoimmunity or related diseases, or immunologically-mediated diseases, for which the efficacy of FK506 and Rapamycin is well known.

Other U.S. patents disclose the use of cyclosporin, Spiperone, Buspirone, their derivatives, and other immunosuppressive compounds for use in the treatment of ophthalmologic diseases (Sharpe et al., U.S. Patent No. 5,703,088; Sharpe et al., U.S. Patent No. 5,693,645; Sullivan, U.S. Patent No. 5,688,765; Sullivan, U.S. Patent No. 5,620,921; Sharpe et al., U.S. Patent No. 5,574,041; Eberle, U.S. Patent No. 5,284,826; Sharpe et al., U.S. Patent No. 5,244,902; Chiou et al., U.S. Patent Nos. 5,198,454 and

5,194,434; and Kaswan, U.S. Patent No. 4,839,342).

These patents also relate to compounds useful for treating autoimmune diseases and cite the known use of cyclosporin, Spiperone, Buspirone, their derivatives, and other immunosuppressive compounds in treating ocular inflammation and other immunologically-mediated ophthalmologic diseases.

The immunosuppressive compounds disclosed in the prior art suppress the immune system, by definition, and also exhibit other toxic side effects. Accordingly, there is a need for non-immunosuppressant, small molecule compounds, and compositions and methods for use of such compounds, that are useful in improving vision; preventing, treating, and/or repairing visual impairment or dysfunction of the visual system; and preventing, treating, and/or resolving ophthalmologic disorders.

There are also a number of patents on non-immunosuppressive compounds disclosing methods of use for permitting or promoting wound healing (whether from injury or surgery); controlling intraocular pressure (often resulting from glaucoma); controlling neurodegenerative eye disorders, including damage or injury to retinal neurons, damage or injury to retinal ganglion cells, and macular degeneration; stimulating neurite outgrowth; preventing or reducing oxidative

damage caused by free radicals; and treating impaired oxygen and nutrient supply, as well as impaired waste product removal, resulting from low blood flow. These non-immunosuppressive substances fall into one of two
5 general categories: naturally occurring molecules, such as proteins, glycoproteins, peptides, hormones, and growth factors; and synthetic molecules.

Within the group of naturally occurring non-immunosuppressive molecules, several hormones, growth
10 factors, and signaling molecules have been patented for use as supplements to naturally occurring quantities of such molecules, as well as for targeting of specific cells where the particular molecule does not naturally occur in a mature individual. These
15 patents generally claim methods of use for reducing or preventing the symptoms of ocular disease, or arresting or reversing vision loss.

Specifically, Louis et al., U.S. Patent Nos. 5,736,516 and 5,641,749, disclose the use of a glial
20 cell line derived neurotrophic factor (GDNF) to stop or reverse the degeneration of retinal neurons (i.e. photoreceptors) and retinal ganglion cells caused by glaucoma, or other degenerative or traumatic retinal diseases or injuries. O'Brien, et al., U.S. Patent
25 Nos. 5,714,459 and 5,700,909, disclose the use of a glycoprotein, Saposin, and its derivatives for

stimulating neurite outgrowth and increasing myelination. To stop or reverse degeneration of retinal neurons, LaVail et al., U.S. Patent No. 5,667,968, discloses the use of a variety of neurotrophic proteins, including brain-derived neurotrophic factor, ciliary neurotrophic factor, neurotrophin-3 or neurotrophin-4, acidic or basic fibroblast growth factors, interleukin, tumor necrosis factor- α , insulin-like growth factor-2 and other growth factors. Wong et al., U.S. Patent No. 5,632,984, discloses the use of interferons, especially interferon α -2a, for treating the symptoms of macular degeneration by reducing hemorrhage and limiting neovascularization. Finally, Wallace et al., U.S. Patent No. 5,441,937, discloses the use of a lung-derived neurotrophic factor (NTF) to maintain the functionality of ciliary ganglion and parasympathetic neuron cells.

A key characteristic of factors derived from specific cell lines is their localization to specific cell lines or tissues; systemic treatment with these molecules would run a substantial risk of unintended, and potentially dangerous, effects in cell lines where the genes encoding these molecules are inactive. Similarly, hormones and growth factors often activate a large number of genes in many cell lines; again,

non-localized application of these molecules would run a substantial risk of provoking an inappropriate, and potentially dangerous, response.

5 Within the category of synthetic molecules, most of the patented compounds are immunosuppressive and disclose uses in treating inflammatory, autoimmune, and allergic responses, as discussed above. A few others are non-immunosuppressive and claim the ability to treat cellular degeneration, and in some cases
10 promote cellular regeneration, most often in the context of their antioxidant properties.

Specifically, Tso et al., U.S. Patent No. 5,527,533, discloses the use of astaxanthin, a carotenoid antioxidant, for preventing or reducing
15 photoreceptor damage resulting from the presence of free radicals. Similarly, Babcock et al., U.S. Patent No. 5,252,319, discloses the use of antioxidant aminosteroids for treating eye disease and injury, by increasing resistance to oxidative damage. Freeman,
20 U.S. Patent No. 5,468,752, discloses the use of the antiviral phosphonylmethoxyalkylcytosines to reduce abnormally increased intraocular pressure.

Hamilton and Steiner disclose in U.S. Patent No. 5,614,547 novel pyrrolidine carboxylate compounds
25 which bind to the immunophilin FKBP12 and stimulate nerve growth, but which lack immunosuppressive


effects. Unexpectedly, it has been discovered that these non-immunosuppressant compounds promote improvements in vision and resolve ophthalmologic disorders. Yet their novel small molecule structure and non-immunosuppressive properties differentiate them from FK506 and related immunosuppressive compounds found in the prior art.

Further, these compounds may be differentiated from the non-immunosuppressive compounds used to treat vision disorders by their novel small molecule structure and their lack of general, systemic effects. Naturally occurring hormones, growth factors, cytokines, and signaling molecules are generally multifunctional and activate many genes in diverse cell lines. The present compounds do not, thus avoiding the unexpected, and potentially dangerous, side effects of systemic use. Similarly, the present compounds also avoid the potential unexpected side effects of introducing cell line-specific molecules into other cell lines where they do not naturally occur.

SUMMARY OF THE INVENTION

The present invention relates to the surprising discovery that a N-linked sulfonamide of an N-heterocyclic carboxylic acid or isostere may be useful for treating a vision disorder or improving vision or treating memory impairment or enhancing memory performance in an animal. Accordingly, novel compositions and methods of using a N-linked sulfonamide of an N-heterocyclic carboxylic acid or isostere are provided. A preferred feature of the compounds of the present invention is that they do not exert any significant immunosuppressive activity.

Preferred embodiments of this invention include methods and compositions containing a compound having the formula (I):



I

where

n is 1-3;

R_1 is selected from the group consisting of hydrogen, C_1-C_9 straight or branched chain alkyl, C_2-C_9 straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

5 D is a bond, or a C_1-C_{10} straight or branched chain alkyl, C_2-C_{10} alkenyl or C_2-C_{10} alkynyl;


R_2 is a carboxylic acid or a carboxylic acid isostere; wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or carboxylic acid isostere is

10 optionally substituted with one or more substituents selected from R^3 , where

R^3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C_1-C_6 straight or branched chain alkyl, C_2-C_6 straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO_2R^4 where R^4 is hydrogen or C_1-C_9 straight or branched chain alkyl or alkenyl;

20 or a pharmaceutically acceptable salt, ester or solvate thereof.

Especially preferred embodiments of this invention are where R_2 is selected from the group below:



where the atoms of said ring structure may be optionally substituted at one or more positions with R^3 ,

where

5 R^3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl or
10 alkynyl, aryl, heteroaryl, carbocycle, heterocycle, and CO_2R^4 where R^4 is hydrogen or C_1 - C_9 straight or branched chain alkyl or alkenyl.

Another preferred embodiment of this invention is where R_2 is selected from the group consisting of -COOH,
15 - SO_3H , - SO_2HNR^3 , - $PO_2(R^3)_2$, -CN, - $PO_3(R^3)_2$, - OR^3 , - SR^3 , - $NHCOR^3$, - $N(R^3)_2$, - $CON(R^3)_2$, - $CONH(O)R^3$, - $CONHNHSO_2R^3$, - $COHNSO_2R^3$, and - $CONR^3CN$.

Brief Description of the Drawings

5 Figure 1 A, B and C show that GPI 1046 protects retinal ganglion cells against degeneration following retinal ischemia.

10 Figure 2 shows that GPI 1046 prevents degeneration of optic nerve axons and myelin following retinal ischemia.

15 Figure 3 shows that GPI 1046 provides moderate protection against retinal ganglion cell death after optic nerve transection.

20 Figure 4 shows that GPI 1046 treatment duration significantly affects the process of optic nerve axonal degeneration after transection.

25 Figure 5 shows that GPI 1046 treatment produces a greater effect on optic nerve axons than ganglion cell bodies.

30 Figure 6 shows that GPI 1046 treatment for 28 days after optic nerve transection prevents myelin degeneration in the proximal stump.

Figure 7 shows that FKBP-12 immunohistochemistry labels oligodendroglia (large dark cells with fibrous processes), the cells which produce myelin, located
5 between the fascicles of optic nerve fibers, and also some optic nerve axons.

Figure 8 shows GPI 1046 treatment for 28 days after optic nerve transection prevents myelin degeneration in
10 the distal stump.

Figure 9 shows that 28 day treatment with GPI 1046 treatment beginning 8 weeks after onset of streptozotocin induced diabetes decreases the extent of
15 neovascularization in the inner and outer retina and protects neurons in the inner nuclear layer (INL) and ganglion cell layer (GCL) from degeneration.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

"Eye" refers to the anatomical structure responsible for vision in humans and other animals, and encompasses the following anatomical structures, without limitation: lens, vitreous body, ciliary body, posterior chamber, anterior chamber, pupil, cornea, iris, canal of Schlemm, zonules of Zinn, limbus, conjunctiva, choroid, retina, central vessels of the retina, optic nerve, fovea centralis, macula lutea, and sclera.

"Alkyl" means a branched or unbranched saturated hydrocarbon chain comprising a designated number of carbon atoms. For example, C₁-C₆ straight or branched alkyl hydrocarbon chain contains 1 to 6 carbon atoms, and includes but is not limited to substituents such as methyl, ethyl, propyl, iso-propyl, butyl, iso-butyl, tert-butyl, n-pentyl, n-hexyl, and the like. It is also contemplated as within the scope of the present invention that "alkyl" may also refer to a hydrocarbon chain wherein any of the carbon atoms of said alkyl are optionally replaced with O, NH, S, or SO₂. For example, carbon 2 of n-pentyl can be replaced with O to form propyloxymethyl.

"Alkenyl" means a branched or unbranched unsaturated hydrocarbon chain comprising a designated number of carbon atoms. For example, C₂-C₆ straight or branched alkenyl hydrocarbon chain contains 2 to 6 carbon atoms having at least one double bond, and includes but is not limited to substituents such as ethenyl, propenyl, iso-propenyl, butenyl, iso-butenyl, tert-butenyl, n-pentenyl, n-hexenyl, and the like. It is also contemplated as

within the scope of the present invention that "alkenyl" may also refer to an unsaturated hydrocarbon chain wherein any of the carbon atoms of said alkenyl are optionally replaced with O, NH, S, or SO₂. For example, carbon 2 of 4-pentene can be replaced with O to form (2-propene)oxymethyl.

"Alkoxy" means the group -OR wherein R is alkyl as herein defined. Preferably, R is a branched or unbranched saturated hydrocarbon chain containing 1 to 6 carbon atoms.

Aryl, heteroaryl, carbocycle, or heterocycle means a cyclic or fused cyclic ring and includes a mono-, bi- or tricyclic, carbo- or heterocyclic ring, wherein the ring is either unsubstituted or substituted in one or more position(s) with hydrogen, hydroxy, carbonyl, amino, amido, cyano, isocyano, nitro, nitroso, nitrilo, isonitrilo, imino, azo, diazo, sulfonyl, sulfhydryl, sulfoxy, thio, thiocarbonyl, thiocyano, formanilido, thioformamido, sulfhydryl, halo, haloalkyl, trifluoromethyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, alkylamino, aminoalkyl, thioalkyl, alkylthio, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R' where R' is hydrogen or C₁-C₆ straight or branched chain alkyl and carbocyclic and heterocyclic moieties. Carbocyclic

moieties include alicyclic and aromatic structures; wherein the individual ring sizes are 5-8 members; wherein the heterocyclic ring contains 1-4 heteroatom(s) selected from the group consisting of O, N, or S; wherein aromatic or tertiary alkyl amines are optionally oxidized to a corresponding N-oxide. Examples of useful alkyl groups include, without limitation, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, n-pentyl, 2-methyl pentyl and the like. Examples of useful carbocyclic and heterocyclic moieties include, without limitation, phenyl, benzyl, naphthyl, indenyl, azulenyl, fluorenyl, anthracenyl, indolyl, isoindolyl, indolinyl, cyclohexyl, benzofuranyl, benzothiophenyl, indazolyl, benzimidazolyl, benzthiazolyl, tetrahydrofuranyl, tetrahydropyranyl, pyridyl, pyrrolyl, pyrrolidinyl, pyridinyl, pyrimidinyl, purinyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, quinolizinyll, furyl, thiophenyl, imidazolyl, oxazolyl, benzoxazolyl, thiazolyl, isoxazolyl, isotriazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, trithianyl, indolizinyll, pyrazolyl, pyrazolinyl, pyrazolidinyl, thienyl, tetrahydroisoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, and adamantyl.

"Halo" means at least one fluoro, chloro, bromo, or

iodo moiety.

The term "pharmaceutically acceptable salt, ester, or solvate" refers to salt, ester, or solvates of the subject compounds which possess the desired pharmacological activity and which are neither biologically nor otherwise undesirable. The salt, ester, or solvates can be formed with inorganic or organic acids such as acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, gluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, naphthylate, 2-naphthalenesulfonate, nicotinate, oxalate, sulfate, thiocyanate, tosylate and undecanoate. Base salt, ester, or solvates include ammonium salts, alkali metal salts such as lithium, sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salt with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups can be quarternized with such agents as: 1) lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; 2)

dialkyl sulfates like dimethyl²⁶, diethyl, dibutyl and diamyl sulfates; 3) long chain alkyls such as decyl, lauryl, myristyl and stearyl substituted with one or more halide such as chloride, bromide and iodide; and 4) aryl or arylalkyl halides like benzyl and phenethyl bromide and others.

The compounds of this invention may possess at least one asymmetric center and thus can be produced as mixtures of stereoisomers or as individual enantiomers or diastereomers. The individual stereoisomers may be obtained by using an optically active starting material, by resolving a racemic or non-racemic mixture of an intermediate at some appropriate stage of the synthesis, or by resolution of the compound of formula (I). It is understood that the individual stereoisomers as well as mixtures (racemic and non-racemic) of stereoisomers are encompassed by the scope of the present invention. The S-stereoisomer at atom 1 of formula I is a most preferred embodiment of the invention.

"Stereoisomers" are isomers that differ only in the way the atoms are arranged in space.

"Isomers" are different compounds that have the same molecular formula and includes cyclic isomers such as (iso)indole and other isomeric forms of cyclic moieties.

"Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of each other.




















"Diastereoisomers" are stereoisomers which are not mirror images of each other.

"Racemic mixture" means a mixture containing equal parts of individual enantiomers. "Non-racemic mixture" is a mixture containing unequal parts of individual enantiomers or stereoisomers.

"Isosteres" are different compounds that have different molecular formulae but exhibit the same or similar properties. For example, tetrazole is an isostere of carboxylic acid because it mimics the properties of carboxylic acid even though they both have very different molecular formulae. Tetrazole is one of many possible isosteric replacements for carboxylic acid. Other carboxylic acid isosteres contemplated by the present invention include

$-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{HNR}^3$, $-\text{PO}_2(\text{R}^3)_2$, $-\text{CN}$, $-\text{PO}_3(\text{R}^3)_2$, $-\text{OR}^3$, $-\text{SR}^3$, $-\text{NHCOR}^3$, $-\text{N}(\text{R}^3)_2$, $-\text{CON}(\text{R}^3)_2$, $-\text{CONH}(\text{O})\text{R}^3$, $-\text{CONHNHSO}_2\text{R}^3$, $-\text{COHNSO}_2\text{R}^3$, and $-\text{CONR}^3\text{CN}$.

In addition, carboxylic acid isosteres can include 5-7 membered carbocycles or heterocycles containing any combination of CH_2 , O , S , or N in any chemically stable oxidation state, where any of the atoms of said ring structure are optionally substituted in one or more positions. The following structures are non-limiting examples of preferred carbocyclic and heterocyclic isosteres contemplated by this invention.



where the atoms of said ring structure may be optionally substituted at one or more positions with R^3 . The present invention contemplates that when chemical substituents are added to a carboxylic isostere then the inventive compound retains the properties of a carboxylic isostere. The present invention contemplates that when a carboxylic isostere is optionally substituted with one or more moieties selected from R^3 , then the substitution can not eliminate the carboxylic acid isosteric properties of the inventive compound. The present invention contemplates that the placement of one or more R^3 substituents upon a carbocyclic or heterocyclic carboxylic acid isostere shall not be at an atom(s) which maintains or is integral to the carboxylic acid isosteric properties of the inventive compound if such a substituent(s) would destroy the carboxylic acid isosteric properties of the inventive compound.

Other carboxylic acid isosteres not specifically exemplified or described in this specification are also contemplated by the present invention.

The term "treatment" as used herein covers any treatment of a disease and/or condition in an animal, particularly a human, and includes:

- (i) preventing a disease and/or condition from occurring in a subject which may be predisposed to the disease and/or condition but has not yet been diagnosed as having it;
- (ii) inhibiting the disease and/or condition, i.e., arresting its development; or
- (iii) relieving the disease and/or condition, i.e., causing regression of the disease and/or condition.

The system used in naming the compounds of the present invention is shown below, using a compound of formula I as an example.

A compound of the present invention, especially formula I, wherein n is 1, D is a bond, R₁ is phenylmethyl, and R₂ is -CN, is named (2S)-1-(phenylmethyl) sulfonyl-2-pyrrolidine carbonitrile.

15 "Enhancing memory performance" refers to
improving or increasing the mental faculty by which to
register, retain or recall past experiences,
knowledge, ideas, sensations, thoughts or impressions.

20 "Memory impairment" refers to a diminished mental
registration, retention or recall of past experiences,
knowledge, ideas, sensations, thoughts or impressions.
Memory impairment may affect short and long-term
information retention, facility with spatial
relationships, memory (rehearsal) strategies, and
verbal retrieval and production. Common causes of

memory impairment are age, severe head trauma, brain anoxia or ischemia, alcoholic-nutritional diseases, and drug intoxications. Examples of memory impairment include, without limitation, benign forgetfulness, amnesia and any disorder in which memory deficiency is present, such as Korsakoff's amnesic psychosis, dementia and learning disorders.

"Neopsic factors" or "neopsics" refers to compounds useful in treating vision loss, preventing vision degeneration, or promoting vision regeneration.

"Neopsis" refers to the process of treating vision loss, preventing vision degeneration, or promoting vision regeneration.

"Ophthalmological" refers to anything about or concerning the eye, without limitation, and is used interchangeably with "ocular," "ophthalmic," "ophthalmologic," and other such terms, without limitation.

"Preventing vision degeneration" refers to the ability to prevent degeneration of vision in patients newly diagnosed as having a degenerative disease

affecting vision, or at risk of developing a new degenerative disease affecting vision, and for preventing further degeneration of vision in patients who are already suffering from or have symptoms of a degenerative disease affecting vision.

"Promoting vision regeneration" refers to maintaining, improving, stimulating or accelerating recovery of, or revitalizing one or more components of the visual system in a manner which improves or enhances vision, either in the presence or absence of any ophthalmologic disorder, disease, or injury.

"Treating" refers to:

(i) preventing a disease and/or condition from occurring in a subject which may be predisposed to the disease and/or condition but has not yet been diagnosed as having it;

(ii) inhibiting the disease and/or condition, i.e., arresting its development; or

(iii) relieving the disease and/or condition, i.e., causing regression of the disease and/or condition.

"Vision" refers to the ability of humans and other animals to process images, and is used interchangeably with "sight", "seeing", and other such terms, without limitation.

"Vision disorder" refers to any disorder that affects or involves vision, including without

limitation visual impairment, orbital disorders, disorders of the lacrimal apparatus, disorders of the eyelids, disorders of the conjunctiva, disorders of the cornea, cataracts, disorders of the uveal tract, disorders of the retina, disorders of the optic nerve or visual pathways, free radical induced eye disorders and diseases, immunologically-mediated eye disorders and diseases, eye injuries, and symptoms and complications of eye disease, eye disorder, or eye injury.

"Visual impairment" refers to any dysfunction in vision including without limitation disturbances or diminution in vision (e.g., binocular, central, peripheral, scotopic), visual acuity for objects near and far, visual field, ocular motility, color perception, adaptation to light and dark, accommodation, refraction, and lacrimation. See Physician's Desk Reference (PDR) for Ophthalmology, 16th Edition, 6:47 (1988).

Methods of the Present Invention.

The present invention relates to a method of treating a vision disorder, improving vision, treating memory impairment, or enhancing memory performance in an animal, which comprises administering to said animal an effective amount of a derivative.

The inventive methods are particularly useful for treating various eye disorders including but not limited to visual disorders, diseases, injuries, and complications, genetic disorders; disorders associated with aging or degenerative vision diseases; vision disorders correlating to physical injury to the eye, head, or other parts of the body resulting from external forces; vision disorders resulting from environmental factors; vision disorders resulting from a broad range of diseases; and combinations of any of the above.

In particular, the compositions and methods of the present invention are useful for improving vision, or correcting, treating, or preventing visual (ocular) impairment or dysfunction of the visual system, including permanent and temporary visual impairment, without limitation. The present invention is also useful in preventing and treating ophthalmologic diseases and disorders, treating damaged and injured eyes, and preventing and treating diseases, disorders, and injuries which result in vision deficiency, vision loss, or reduced capacity to see or process images, and the symptoms and complications resulting from same. The eye diseases and disorders which may be treated or prevented by the compositions and methods of the present invention are not limited with regard to the cause of said diseases or disorders.

Accordingly, said compositions and methods are applicable whether the disease or disorder is caused by genetic or environmental factors, as well as any other influences. The compositions and methods of the present invention are particularly useful for eye problems or vision loss or deficiency associated with all of the following, without limitation: aging, cellular or physiological degeneration, central nervous system or neurological disorder, vascular defects, muscular defects, and exposure to adverse environmental conditions or substances.

The compositions and methods of the present invention are particularly useful in correcting, treating, or improving visual impairment, without limitation. Visual impairment in varying degrees occurs in the presence of a deviation from normal in one or more functions of the eye, including (1) visual acuity for objects at distance and near; (2) visual fields; and (3) ocular motility without diplopia. See *Physicians' Desk Reference (PDR) for Ophthalmology*, 16th Edition, 6:47 (1988). Vision is imperfect without the coordinated function of all three. *Id.*

Said compositions and methods of use are also useful in correcting, treating, or improving other ocular functions including, without limitation, color perception, adaptation to light and dark, accommodation, metamorphopsia, and binocular vision.

The compositions and methods of use are particularly useful in treating, correcting, or preventing ocular disturbances including, without limitation, paresis of accommodation, iridoplegia, entropion, ectropion, 5 epiphora, lagophthalmos, scarring, vitreous opacities, non-reactive pupil, light scattering disturbances of the cornea or other media, and permanent deformities of the orbit.

The compositions and methods of use of the present invention are also highly useful in improving 10 vision and treating vision loss. Vision loss ranging from slight loss to absolute loss may be treated or prevented using said compositions and methods of use. Vision may be improved by the treatment of eye 15 disorders, diseases, and injuries using the compositions and methods of the invention. However, improvements in vision using the compositions and methods of use are not so limited, and may occur in the absence of any such disorder, disease, or injury.

20 The compositions and methods of the present invention are also useful in the treatment or prevention of the following non-limiting exemplary diseases and disorders, and symptoms and complications resulting therefrom.

25 Vision disorders include but are not limited to the following:

visual impairment, such as diminished visual

acuity for objects near and far, visual fields, and ocular motility;

orbital disorders, such as orbital cellulitis, periorbital cellulitis, cavernous sinus thrombosis, and exophthalmos (proptosis);

disorders of the lacrimal apparatus, such as dacryostenosis, congenital dacryostenosis, and dacryocystitis (acute or chronic);

disorders of the eyelids, such as lid edema, blepharitis, ptosis, Bell's palsy, blepharospasm, hordeolum (stye), external hordeolum, internal hordeolum (meibomian stye), chalazion, entropion (inversion of the eyelid), ectropion (eversion of the eyelid), tumors (benign and malignant), xanthelasma, basal cell carcinoma, squamous cell carcinoma, meibomian gland carcinoma, and melanoma;

disorders of the conjunctiva, such as pinguecula, pterygium, and other neoplasms, acute conjunctivitis, chronic conjunctivitis, adult gonococcal conjunctivitis, neonatal conjunctivitis, trachoma (granular conjunctivitis or Egyptian ophthalmia), inclusion conjunctivitis (inclusion blenorrhea or swimming pool conjunctivitis), neonatal inclusion conjunctivitis, adult inclusion conjunctivitis, vernal keratoconjunctivitis, keratoconjunctivitis sicca (keratitis sicca or dry eye syndrome), episcleritis, scleritis, cicatricial pemphigoid (ocular cicatricial

pemphigoid or benign mucous membrane pemphigoid), and subconjunctival hemorrhage;

disorders of the cornea, such as superficial punctate keratitis, corneal ulcer, indolent ulcer, recurrent corneal erosion, corneal epithelial basement membrane dystrophy, corneal endothelial cell dystrophy, herpes simplex keratitis (herpes simplex keratoconjunctivitis), dendritic keratitis, disciform keratitis, ophthalmic herpes zoster, phlyctenular keratoconjunctivitis (phlyctenular or eczematous conjunctivitis), interstitial keratitis (parenchymatous keratitis), peripheral ulcerative keratitis (marginal keratolysis or peripheral rheumatoid ulceration), keratomalacia (xerotic keratitis), xerophthalmia, keratoconus, bullous keratopathy;

cataracts, including developmental or congenital cataracts, juvenile or adult cataracts, nuclear cataract, posterior subcapsular cataracts;

disorders of the uveal tract, such as uveitis (inflammation of the uveal tract or retina), anterior uveitis, intermediate uveitis, posterior uveitis, iritis, cyclitis, choroiditis, ankylosing spondylitis, Reiter's syndrome, pars planitis, toxoplasmosis, cytomegalovirus (CMV), acute retinal necrosis, toxocariasis, birdshot choroidopathy, histoplasmosis (presumed ocular histoplasmosis syndrome), Behcet's

syndrome, sympathetic ophthalmia, Vogt-Koyanagi-Harada syndrome, sarcoidosis, reticulum cell sarcoma, large cell lymphoma, syphilis, tuberculosis, juvenile rheumatoid arthritis, endophthalmitis, and malignant melanoma of the choroid;

disorders of the retina, such as vascular retinopathies (e.g., arteriosclerotic retinopathy and hypertensive retinopathy), central and branch retinal artery occlusion, central and branch retinal vein occlusion, diabetic retinopathy (e.g., proliferative retinopathy and non-proliferative retinopathy), macular degeneration of the aged (age-related macular degeneration or senile macular degeneration), neovascular macular degeneration, retinal detachment, retinitis pigmentosa, retinal photic injury, retinal ischemia-induced eye injury, and glaucoma (e.g., primary glaucoma, chronic open-angle glaucoma, acute or chronic angle-closure, congenital (infantile) glaucoma, secondary glaucoma, and absolute glaucoma);

disorders of the optic nerve or visual pathways, such as papilledema (choked disk), papillitis (optic neuritis), retrobulbar neuritis, ischemic optic neuropathy, toxic amblyopia, optic atrophy, higher visual pathway lesions, disorders of ocular motility (e.g., third cranial nerve palsies, fourth cranial nerve palsies, sixth cranial nerve palsies, internuclear ophthalmoplegia, and gaze palsies);

free radical induced eye disorders and diseases;
and

immunologically-mediated eye disorders and
diseases, such as Graves' ophthalmopathy, conical
5 cornea, dystrophia epithelialis corneae, corneal
leukoma, ocular pemphigus, Mooren's ulcer, scleritis,
and sarcoidosis (See *The Merck Manual*, Sixteenth
Edition, 217:2365-2397 (1992) and *The Eye Book*,
Cassel, Billig, and Randall, The Johns Hopkins
10 University Press (1998)).

The compositions and methods of the present
invention are also useful in the treatment of the
following non-limiting eye injuries, and symptoms and
complications resulting therefrom: conjunctival and
15 corneal foreign body injuries, corneal abrasion,
intraocular foreign body injuries, lacerations, lid
lacerations, contusions, lid contusions (black eye),
trauma to the globe, laceration of the iris, cataract,
dislocated lens, glaucoma, vitreous hemorrhage,
20 orbital-floor fractures, retinal hemorrhage or
detachment, and rupture of the eyeball, anterior
chamber hemorrhage (traumatic hyphema), burns, eyelid
burns, chemical burns, chemical burns of the cornea
and conjunctiva, and ultraviolet light burns
25 (sunburn). See *The Merck Manual*, Sixteenth Edition,
217:2364-2365 (1992).

The compositions and methods of the present

invention are also useful in treating and/or preventing the following non-limiting exemplary symptoms and complications of eye disease, eye disorder or eye injury: subconjunctival hemorrhages, vitreous hemorrhages, retinal hemorrhages, floaters, retinal detachments, photophobia, ocular pain, scotomas (negative and positive), errors of refraction, emmetropia, ametropia, hyperopia (farsightedness), myopia (nearsightedness), astigmatism, anisometropia, aniseikonia, presbyopia, bleeding, recurrent bleeding, sympathetic ophthalmia, inflammation, swelling, redness of the eye, irritation of the eye, corneal ulceration and scarring, iridocyclitis, perforation of the globe, lid deformities, exophthalmos, impaired mobility of the eye, lid swelling, chemosis, loss of vision, including partial or total blindness, optic neuritis, fever, malaise, thrombophlebitis, cavernous sinus thrombosis, panophthalmitis, infection of the meninges and brain, papilledema, severe cerebral symptoms (headache, decreased level of consciousness, and convulsions), cranial nerve palsies, epiphora (chronic or persistent tearing), copious reflux of mucus or pus, follicular subconjunctival hyperplasia, corneal vascularization, cicatrization of the conjunctiva, cornea, and lids, pannus, hypopyon, lagophthalmos, phlyctenules, rubeosis iridis, bitemporal hemianopia, and homonymous

hemianopia. See *The Merck Manual, Sixteenth Edition*, 217:2362-2363 (1992).

5 The derivative may be administered in combination with an effective amount of one or more factor(s) useful in treating vision disorder, improving vision, treating memory impairment, or enhancing memory performance.

10 In a preferred embodiment, the factor(s) to be combined with the derivative is/are selected from the group consisting of immunosuppressants for treating autoimmune, inflammatory, and immunologically-mediated disorders; wound healing agents for treating wounds resulting from injury or surgery; antiglaucomatous medications
15 for treating abnormally elevated intraocular pressure; neurotrophic factors and growth factors for treating neurodegenerative disorders or stimulating neurite outgrowth; compounds effective in limiting or preventing hemorrhage or neovascularization for
20 treating macular degeneration; and antioxidants for treating oxidative damage to eye tissues.

Pharmaceutical Compositions of the Present Invention


25 The present invention also relates to a pharmaceutical composition comprising:

- (i) an effective amount of a derivative for treating a vision disorder, improving vision, treating memory impairment, or enhancing memory performance in an animal; and

- (ii) a pharmaceutically acceptable carrier.

5 The derivative may be administered in combination with an effective amount of one or more factor(s) useful in treating vision disorders, improving vision, treating memory impairment, or enhancing memory performance.









TABLE A











No.	n	D	R2	R1
1	1	bond	COOH	Benzyl
2	1	bond	COOH	α -MethylBenzyl
3	1	bond	COOH	4-MethylBenzyl
4	1	bond	Tetrazole	Benzyl
5	1	bond	SO ₃ H	α -MethylBenzyl
6	1	CH ₂	COOH	4-MethylBenzyl
7	1	bond	SO ₂ HNMe	Benzyl
8	1	bond	CN	α -MethylBenzyl
9	1	bond	PO ₃ H ₂	4-MethylBenzyl
10	2	bond	COOH	Benzyl
11	2	bond	COOH	α -MethylBenzyl
12	2	bond	COOH	4-MethylBenzyl
13	2	bond	COOH	3,4,5-trimethoxy phenyl
14	2	bond	COOH	Cyclohexyl
15	2	bond	PO ₃ HEt	i-propyl
16	2	bond	PO ₃ HPropyl	ethyl
17	2	bond	PO ₃ (Et) ₂	Methyl
18	2	bond	OMe	tert-butyl
19	2	bond	OEt	n-pentyl
20	2	bond	OPropyl	n-hexyl
21	1	bond	OButyl	Cyclohexyl
22	1	bond	OPentyl	cyclopentyl
23	1	bond	OHexyl	n-heptyl
24	1	bond	SMe	n-octyl
25	1	bond	SEt	n-nonyl
26	2	bond	SPropyl	2-indolyl
27	2	bond	SButyl	2-furyl
28	2	bond	NHCOMe	2-thiazolyl
29	2	bond	NHCOEt	2-thienyl
30	1	CH ₂	N(Me) ₂	2-pyridyl
31	1	(CH ₂) ₂	N(Me)Et	benzyl
32	1	(CH ₂) ₃	CON(Me) ₂	benzyl
33	1	(CH ₂) ₄	CONHMe	benzyl
34	1	(CH ₂) ₅	CONHEt	benzyl
35	1	(CH ₂) ₆	CONHPropyl	1,1-dimethylpropl
No.	n	D	R2	R1
36	1	bond	CONH(O)Me	Benzyl

37	1	bond	CONH(O)Et	α -Methylphenyl
38	1	bond	CONH(O)Propyl	4-Methylphenyl
39	2	bond	COOH	Benzyl
40	2	bond	COOH	α -Methylphenyl
41	2	bond	COOH	4-Methylphenyl
42	1	CH ₂	COOH	benzyl
43	1	(CH ₂) ₂	COOH	benzyl
44	1	(CH ₂) ₃	COOH	benzyl
45	1	(CH ₂) ₄	COOH	benzyl
46	1	(CH ₂) ₅	COOH	benzyl
47	1	(CH ₂) ₆	COOH	benzyl
48	1	(CH ₂) ₇	COOH	benzyl
49	1	(CH ₂) ₈	COOH	benzyl
50	1	(CH ₂) ₉	COOH	benzyl
51	1	(CH ₂) ₁₀	COOH	benzyl
52	1	C ₂ H ₂	COOH	benzyl
53	1	2-OH, Et	COOH	benzyl
54	1	2butylene	COOH	benzyl
55	1	i-Pro	COOH	benzyl
56	1	tert-Bu	COOH	benzyl
57	1	2-nitro	COOH	benzyl
		Hexyl		
58	3	(CH ₂) ₂	CN	benzyl
59	1	(CH ₂) ₃	CN	benzyl
60	3	bond	CONHNHSO ₂ Me	Benzyl
61	3	bond	CONHNHSO ₂ Et	α -Methylphenyl
62	3	bond	CONHSO ₂ Me	4-Methylphenyl
63	2	bond	CONHNHSO ₂ Et	Phenyl
64	2	bond	CON(Me)CN	α -Methylphenyl
65	2	bond	CON(Et)CN	4-Methylphenyl
66	1	(CH ₂) ₂	COOH	methyl
67	1	(CH ₂) ₃	COOH	ethyl
68	1	(CH ₂) ₄	COOH	n-propyl
69	1	(CH ₂) ₅	COOH	t-butyl
70	1	(CH ₂) ₆	COOH	Pentyl
71	1	(CH ₂) ₇	COOH	Hexyl
72	1	(CH ₂) ₈	COOH	Septyl
73	1	(CH ₂) ₉	COOH	Octyl
74	1	(CH ₂) ₁₀	COOH	Nonyl
75	1	C ₂ H ₂	COOH	Cyclohexyl




47


No.	n	D	R2	R1
76	1	bond		benzyl
77	1	bond		benzyl
78	1	bond		benzyl
79	1	bond		benzyl
80	1	bond		benzyl
81	1	bond		benzyl
82	1	bond		benzyl
83	1	bond		benzyl

48

No.	n	D	R2	R1
84	1	bond		benzyl
85	1	bond		benzyl
86	1	bond		benzyl
87	1	bond		benzyl
88	1	bond		benzyl
89	1	bond		benzyl
90	1	bond		benzyl
91	1	bond		benzyl

49

No.	n	D	R2	R1
92	1	bond		benzyl
93	1	bond		benzyl
94	1	bond		benzyl
95	1	bond	CH ₂ OH	benzyl
96	1	bond	CONH ₂	benzyl
97	1	bond	CN	benzyl



No.	n	D	R ₂	L	R ₁
101	1	CH ₂	OH	1,2-dioxoethyl	benzyl
102	1	bond	-CN	1,2-dioxoethyl	1,1-dimethylpropyl
103	1	bond	tetrazole	1,2-dioxoethyl	1,1-dimethylpropyl
104	2	bond	CONH ₂	1,2-dioxoethyl	1,1-dimethylpropyl
105	1	bond	COOH	1,2-dioxoethyl	1,1-dimethylpropyl
106	2	bond	COOH ⁺	1,2-dioxoethyl	1,1-dimethylpropyl

Affinity for FKBP12

10 The compounds used in the inventive methods and
pharmaceutical compositions have an affinity for the
FK506 binding protein, particularly FKBP12. The
inhibition of the prolyl peptidyl *cis-trans* isomerase
activity of FKBP may be measured as an indicator of
15 this affinity.

K_i Test Procedure

 Inhibition of the peptidyl-prolyl isomerase
(rotamase) activity of the compounds used in the
20 inventive methods and pharmaceutical compositions can
be evaluated by known methods described in the
literature (Harding et al., *Nature*, 1989, 341:758-760;
Holt et al. *J. Am. Chem. Soc.*, 115:9923-9938). These

values are obtained as apparent K_i 's.

The *cis-trans* isomerization of an alanine-proline bond in a model substrate, N-succinyl-Ala-Ala-Pro-Phe-
5 p-nitroanilide, is monitored spectrophotometrically in a chymotrypsin-coupled assay, which releases para-nitroanilide from the *trans* form of the substrate. The inhibition of this reaction caused by the addition of different concentrations of inhibitor is
10 determined, and the data is analyzed as a change in first-order rate constant as a function of inhibitor concentration to yield the apparent K_i values.

In a plastic cuvette are added 950 μ l of ice cold assay buffer (25 mM HEPES, pH 7.8, 100 mM NaCl), 10 μ l
15 of FKBP (2.5 mM in 10 mM Tris-Cl pH 7.5, 100 mM NaCl, 1 mM dithiothreitol), 25 μ l of chymotrypsin (50 mg/ml in 1 mM HCl) and 10 μ l of test compound at various concentrations in dimethyl sulfoxide. The reaction is initiated by the addition of 5 μ l of substrate
20 (succinyl-Ala-Phe-Pro-Phe-para-nitroanilide, 5 mg/ml in 2.35 mM LiCl in trifluoroethanol).

The absorbance at 390 nm versus time is monitored for 90 seconds using a spectrophotometer and the rate constants are determined from the absorbance versus
25 time data files.

Route of Administration

45 To effectively treat vision loss or promote
vision regeneration, the compounds used in the
inventive methods and pharmaceutical compositions must

readily affect the targeted areas. For these purposes, the compounds are preferably administered [topically to the skin.]

[For topical application to the skin, the compounds can be formulated into suitable ointments containing the compounds suspended or dissolved in, for example, mixtures with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated into suitable lotions or creams containing the active compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl ester wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.]

Other routes of administration known in the pharmaceutical art are also contemplated by this invention.

Dosage

Dosage levels on the order of about 0.1 mg to about 10,000 mg of the active ingredient compound are useful in the treatment of the above conditions, with preferred levels of about 0.1 mg to about 1,000 mg. The specific dose level for any particular patient

will vary depending upon a variety of factors, including the activity of the specific compound employed; the age, body weight, general health, sex and diet of the patient; the time of administration; 5 the rate of excretion; drug combination; the severity of the particular disease being treated; and the form of administration. Typically, *in vitro* dosage-effect results provide useful guidance on the proper doses for patient administration. Studies in animal models 10 are also helpful. The considerations for determining the proper dose levels are well known in the art.



The compounds can be administered with other agents for treating vision loss, preventing vision degeneration, or promoting vision regeneration. 15 Specific dose levels for such other agents will depend upon the factors previously stated and the effectiveness of the drug combination.

The following examples are illustrative of preferred embodiments of the invention and are not to be construed as limiting the invention thereto. All polymer molecular weights are mean average molecular weights. All percentages are based on the percent by weight of the final delivery system or formulation prepared unless otherwise indicated and all totals equal 100% by weight.


EXAMPLES

The inventive compounds may be prepared by a variety of synthetic sequences that utilize established chemical transformations. An exemplary general pathway to the present compounds is described in Scheme I, Scheme II, and Scheme III.

57

SCHEME ISCHEME II

58

SCHEME IIIEXAMPLE 1

Synthesis of (2S)-N-(benzylsulfonyl)-2-
pyrrolidinecarboxylic acid (Compound 1) (A)

To a cooled (0°C) solution of proline methyl ester hydrochloride salt (5.0 g; 30.19 mmol) in 200 mL of methylene chloride was added triethylamine (35mL) and benzenesulfonyl chloride (5.75 g; 30.19 mmol). The mixture was stirred for one hour at 0°C and then washed with 2 x 100 mL of water. The organic phase was dried and concentrated. Chromatography eluting with 50% EtOAc/hexane delivered 8.14 g (5%) of the N-sulfonamide methyl ester, which was dissolved in 120 mL of methanol, cooled to 0°C , and treated with 40 mL of 1 N lithium hydroxide. The mixture was stirred for 1 hour at 0°C and then overnight at room temperature. After making the reaction mixture acidic (pH 1) with 1 N HCl, the product was extracted into methylene chloride and dried and

concentrated to yield 4.25 g of (2S)-N-(benzylsulfonyl)-2-pyrrolidinecarboxylic acid (A) as a white solid, ¹H NMR (CDCl₃, 400 MHz): δ 1.85-1.90 (m, 2H); 2.08 (m, 1H); 2.18 (m, 1H); 3.04 (m, 1H); 3.27 (m, 1H); 4.32-4.35 (m, 2H); 4.45 (m, 1H); 4.45 (m, 2H); 7.36 (m, 3H); 7.48 (m, 2H); 10.98 (br, 1H).

EXAMPLE 2

Synthesis of (2S)-1-(phenylmethylsulfonyl)-2-hydroxymethyl pyrrolidine (Compound 95) (E).

To a solution of (S)-(+)-2-pyrrolidinemethanol (1.01 g, 10 mmol) and triethylamine (1.5 ml, 11 mmol) in 30 ml methylene chloride was added 1.9 g (10 mmol) α-toluenesulfonyl chloride at 0°C with stirring. The reaction was gradually warmed up to room temperature and stirred overnight. The mixture was diluted with water, and extracted into 200 ml methylene chloride. The organic extract was concentrated and further purified by silica gel to give 1.5 g product as a white solid (58.9% yield). ¹H NMR (CDCl₃): δ 0.71-1.88 (m, 4H); 2.05 (br, 1H, OH); 3.22 (m, 2H); 3.47 (m, 2H); 3.67 (m, 1H); 4.35 (s, 2H); 7.26-7.44 (m, 5H, aromatic).

EXAMPLE 3

Synthesis of (2S)-1-(phenylmethylsulfonyl)-2-

pyrrolidinecarboxamide (Compound 96) (B).

To a solution of L-prolinamide (2.28 g, 20 mmol) and triethylamine (5.76 ml, 42 mmol) in 40 ml methylene chloride was added 3.92 g (20 mmol) α -toluenesulfonyl chloride at 0°C with stirring. The reaction was gradually warmed up to room temperature and stirred overnight. The mixture was diluted with water, and extracted into 200 ml methylene chloride. The organic extract was concentrated and further purified by silica gel to give 3.0 g product as a white solid (55.7% yield). $^1\text{H NMR}$ (CDCl_3): δ 01.89 (m, 3H); 2.25 (m, 1H); 3.40 (m, 1H); 3.50 (m, 1H); 3.96 (m, 1H); 4.35 (s, 2H); 7.39-7.45 (m, 5H, aromatic).

EXAMPLE 4Synthesis of (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinecarbonitrile (Compound 97) (C).

To a solution of 0.67 ml DMF (8.7 mmol) in 10 ml acetonitrile at 0°C was added 0.70 ml (8.0 mmol) oxalyl chloride. A white precipitate was formed immediately and was accompanied by gas evolution. When complete, a solution of 2.0 g (7.5 mmol) of (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinecarboxamide in 5.0 ml acetonitrile was added. When the mixture became homogeneous, 1.35 ml

(16.5 mmol) pyridine was added. After 5 min., the mixture was diluted with water, and extracted by 200 ml ethyl acetate. The organic layer was concentrated and further purified by silica gel to give 1.5 g product as a white solid (80% yield). ¹H NMR (CDCl₃): δ 01.92 (m, 2H); 2.01 (m, 1H); 2.11 (m, 1H); 3.45 (m, 2H); 4.35 (s, 2H); 4.65 (m, 1H); 7.26-7.45 (m, 5H, aromatic).

EXAMPLE 5

Synthesis of (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinecarboxamide (Compound 4) (D).

A mixture of (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinecarboxamide (250 mg, 1 mmol), NaN₃ (81 mg, 1.3 mmol) and NH₄Cl (70 mg, 1.3 mmol) in 3 ml DMF was stirred at 130°C for 16 hours. The mixture was concentrated and purified by silica gel to give 120 mg product as a white solid (41.1% yield). ¹H NMR (CDCl₃): δ 01.95 (m, 2H); 2.21 (m, 1H); 2.90 (m, 1H); 3.40 (m, 2H); 4.27 (s, 2H); 5.04 (m, 1H); 7.36-7.41 (m, 5H, aromatic); 8.05 (s, 1H, NH).

Example 6Synthesis of 3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate (1)Methyl (2S)-1-(1,2-dioxo-2-methoxyethyl)-2-pyrrolidinecarboxylate

A solution of L-proline methyl ester hydrochloride (3.08 g; 18.60 mmol) in dry methylene chloride was cooled to 0°C and treated with triethylamine (3.92 g; 38.74 mmol; 2.1 eq). After stirring the formed slurry under a nitrogen atmosphere for 15 min, a solution of methyl oxalyl chloride (3.20 g; 26.12 mmol) in methylene chloride (45 ml) was added dropwise. The resulting mixture was stirred at 0°C for 1.5 hour. After filtering to remove solids, the organic phase was washed with water, dried over MgSO₄, and concentrated. The crude residue was purified on a silica gel column, eluting with 50% ethyl acetate in hexane, to obtain 3.52 g (88%) of the product as a reddish oil. Mixture of cis-trans amide rotamers; data for trans rotamer given. ¹H NMR (CDCl₃): δ 1.93 (dm, 2H); 2.17 (m, 2H); 3.62 (m, 2H); 3.71 (s, 3H); 3.79, 3.84 (s, 3H total); 4.86 (dd, 1H, J = 8.4, 3.3).

Methyl (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylate

A solution of methyl (2S)-1-(1,2-dioxo-2-methoxyethyl)-2-pyrrolidinecarboxylate (2.35 g; 10.90 mmol) in 30 ml of tetrahydrofuran (THF) was cooled to -78°C and treated with 14.2 ml of a 1.0 M solution of 1,1-dimethylpropylmagnesium chloride in THF. After

stirring the resulting homogeneous mixture at -78°C for three hours, the mixture was poured into saturated ammonium chloride (100 ml) and extracted into ethyl acetate. The organic phase was washed with water, 5 dried, and concentrated, and the crude material obtained upon removal of the solvent was purified on a silica gel column, eluting with 25% ethyl acetate in hexane, to obtain 2.10 g (75%) of the oxamate as a colorless oil. ^1H NMR (CDCl_3): d 0.88 (t, 3H); 1.22, 10 1.26 (s, 3H each); 1.75 (dm, 2H); 1.87-2.10 (m, 3H); 2.23 (m, 1H); 3.54 (m, 2H); 3.76 (s, 3H); 4.52 (dm, 1H, $J = 8.4, 3.4$).

Synthesis of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylic acid

15 A mixture of methyl (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylate (2.10 g; 8.23 mmol), 1 N LiOH (15 ml), and methanol (50 ml) was stirred at 0°C for 30 minutes and at room temperature overnight. The mixture was acidified to pH 1 with 1 N 20 HCl, diluted with water, and extracted into 100 ml of methylene chloride. The organic extract was washed with brine and concentrated to deliver 1.73 g (87%) of snow-white solid which did not require further purification. ^1H NMR (CDCl_3): d 0.87 (t, 3H); 1.22, 25 1.25 (s, 3H each); 1.77 (dm, 2H); 2.02 (m, 2H); 2.17 (m, 1H); 2.25 (m, 1H); 3.53 (dd, 2H, $J = 10.4, 7.3$); 4.55 (dd, 1H, $J = 8.6, 4.1$).

3-Phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate (1)

A mixture of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidine-carboxylic acid (600 mg; 2.49 mmol), 3-phenyl-1-propanol (508 mg; 3.73 mmol),
5 dicyclohexylcarbodiimide (822 mg; 3.98 mmol), camphorsulfonic acid (190 mg; 0.8 mmol) and 4-dimethylaminopyridine (100 mg; 0.8 mmol) in methylene chloride (20 ml) was stirred overnight under a
10 nitrogen atmosphere. The reaction mixture was filtered through Celite to remove solids and concentrated in vacuo, and the crude material was purified on a flash column (25% ethyl acetate in hexane) to obtain 720 mg (80%) of Example 1 as a
15 colorless oil. ¹H NMR (CDCl₃): δ 0.84 (t, 3H); 1.19 (s, 3H); 1.23 (s, 3H); 1.70 (dm, 2H); 1.98 (m, 5H); 2.22 (m, 1H); 2.64 (m, 2H); 3.47 (m, 2H); 4.14 (m, 2H); 4.51 (d, 1H); 7.16 (m, 3H); 7.26 (m, 2H).

Figure 1. GPI 1046 protects retinal ganglion cells against degeneration following retinal ischemia.

Retinal ganglion cells were retrogradely labeled in adult rats by bilateral injection of fluorogold in their lateral geniculate nuclei. Labeled ganglion cells in the normal rat retina appear as white profiles against the dark background (Figure 1A). Complete retinal ischemia was produced by infusing normal saline solution into the retinal vitreous cavity of each eye until the intraocular pressure exceeded arterial blood pressure. 28 days after the ischemic episode extensive degeneration of retinal ganglion cell was evidenced by massive reduction in the density of fluorogold labeled cells (Figure 1B). Administration of GPI 1046 (10mg/kg, s.c.) 1 hour prior to the ischemic episode and at 10mg/kg/day for the next four days produced noticeable protection of a large proportion of the vulnerable ganglion cell population (Figure 1C).

Figure 2. GPI 1046 prevents degeneration of optic nerve axons and myelin following retinal ischemia

Examination of the optic nerves from the same retinal ischemia cases reveals that GPI 1046 produces dramatic protection of optic nerve element from ischemic degeneration. Toluidine blue staining of epon embedded optic nerve cross sections revealed the detail of myelin sheaths (white circles) and optic nerve axons

(black centers) in the normal rat optic nerve. Optic nerves from vehicle treated cases examined 28 days after a 1 hour retinal ischemic episode are characterized by a decreased density of optic nerve axons and the appearance of numerous degenerating myelin figures (bright white filled circles).

Treatment with GPI 1046 protected the majority of optic nerve axons from degeneration and also dramatically decreased the density of degenerating myelin figures.

Figure 3. GPI 1046 provides moderate protection against retinal ganglion cell death after optic nerve transection

Complete transection of the optic nerve 5 mm from the eyeball produces massive degeneration of retinal ganglion cells, representing loss of >87% of the normal ganglion cell population 90 days after the injury (Table 1). Few spared fluorogold pre labeled ganglion cells are present in vehicle treated cases (large white figures) among a population of small microglia that digest the debris of the degenerating cells and take up the fluorogold label (Figure 3A). Treatment with GPI 1046 for 14 days resulted in a small but not significant increase in the density of retinal ganglion cells that survived 90 days after transection (Table 1) but treatment with GPI 1046 for the first 28 days after transection produced moderate but significant

protection of 12.6% of the vulnerable ganglion cell population (Table 1, Figure 3B).

5 **Figure 4. GPI 1046 treatment duration significantly affects the process of optic nerve axonal degeneration after transection.**

Examination of optic nerve axon density in the proximal stump of the optic nerve from the same cases revealed a more dramatic protection afforded by GPI 1046
10 treatment. 90 days after transection few ganglion cell axons remain within the optic nerve (Figure 4B), representing only 5.6% of the normal population. The loss of axons reflects both the death of retinal ganglion cells and the regression or "dying back" of
15 the axons of ~ 70% of the small surviving ganglion cell population into the retina itself (Table 1). Treatment with GPI 1046 for the first 14 days after optic nerve transection produced a small but significant 5.3% protection of optic nerve axons (Figure 4D, Table 1),
20 but treatment with the same dose of GPI 1046 for 28 days resulted in the protection of optic nerve axons for the vast majority (81.4%) of spared retinal ganglion cells (Figure 4C, Table 1).

25 **Figure 5. GPI 1046 treatment produces a greater effect on optic nerve axons than ganglion cell bodies**
This summary figure shows data from Figure 3 ganglion cell protection and higher power photomicrographs of

optic nerve axon protection (Figure 5A&B, upper panels). 28 day treatment with GPI 1046 produced a significant increase in the density of large, and particularly medium and small caliber optic nerve axons (Figure 5C&D, lower panels).

Figure 6. GPI 1046 treatment for 28 days after optic nerve transection prevents myelin degeneration in the proximal stump

Myelin basic protein immunohistochemistry labels fascicles (darker labeled 'islands') of myelinated axons in the normal optic nerve (Figure 6A, upper left). 90 days after transection extensive degeneration of myelin is evident in vehicle treated cases, characterized by the loss of fascicular organization and the appearance of numerous large dense degenerating myelin figures (Figure 6B, upper right). Treatment with GPI 1046 for the first 14 days after optic nerve transection did not alter the pattern of myelin degeneration (Figure 6C, lower left panel), and yielded an insignificant 1.6% quantitative recovery in myelin density (Table 1). Extending the GPI 1046 treatment course through the first 28 days after optic nerve transection produced a dramatic preservation of the fascicular staining pattern for myelin basic protein in the proximal stump of the optic nerve and decreased the density of degenerating myelin figures

(Figure 6D, lower right panel), representing a '70% recovery of myelin density (Table 1).

Figure 7. FKBP-12 immunohistochemistry labels
5 oligodendroglia (large dark cells with fibrous processes), the cells which produce myelin, located between the fascicles of optic nerve fibers, and also some optic nerve axons.

10 Figure 8. GPI 1046 treatment for 28 days after optic nerve transection prevents myelin degeneration in the distal stump.
Complete transection of the optic nerve leads to degeneration of the distal segments (axon fragments
15 disconnected from the ganglion cell bodies), and the degeneration of their myelin sheaths. 90 days after transection (Figure 8B) myelin basic protein immunohistochemistry reveals the near total loss of fascicular organization (present in the normal optic
20 nerve, Figure 8A) and the presence of numerous dense degenerating myelin figures. Quantitation reveals that the cross sectional area of the transected distal stump shrinks by 31% and loses approximately 1/2 of its myelin (Table 1). Treatment with GPI 1046 for the
25 first 14 days after transection did not protect against shrinkage of the distal stump but did slightly increase the density of myelin, though the density of

degenerating myelin figures remained high (Figure 8C, Table 1). GPI 1046 treatment through the first 28 days produced dramatic protection of the fascicular pattern of myelin labeling, decreased the density of
5 degenerating myelin figures, prevented cross sectional shrinkage of the distal stump of the transected nerve and maintained the myelin levels at ~99% of normal levels (Figure 8D, Table 1).

10 Figure 9. 28 day treatment with GPI 1046 treatment beginning 8 weeks after onset of streptozotocin induced diabetes decreases the extent of neovascularization in the inner and outer retina and protects neurons in the inner nuclear layer (INL) and ganglion cell layer (GCL)
15 from degeneration.

Negative images of cresyl violet stained tangential retinal sections reveals perikarya in the three cellular layers (Figure 9A). The retinae of streptozotocin treated animals administered only
20 vehicle (Figure 9B) exhibited loss of cells from the ONL and INL, decreased thickness of the Outer plexiform layer (the dark area between ONL and INL) and a dramatic increase in the size and density of retinal blood vessels (large black circular outlines) in the
25 INL, OPL, ONL and the photoreceptor layer (PR, the gray fuzzy area above the ONL). GPI 1046 treatment reduced neovascularization (i.e. prevented the proliferation of

blood vessels) in the PR, ONL, OPL and INL. Although
GPI 1046 did not appear to protect against neuronal
loss in the ONL, it appeared to decrease the loss of
neurons in both the INL and GCL compared to
5 streptozotocin/vehicle treated controls.

Example 7

10 In Vivo Retinal Ganglion Cell
 and Optic Nerve Axon Tests

 The extent of degeneration reduction or prevention in retinal ganglion cells and optic nerve axons was determined in a vision loss model utilizing surgical optic nerve transection to simulate
15 mechanical damage to the optic nerve. The effects of several neuroimmunophilin FKBP ligands on retinal ganglion cells neuroprotection and optic nerve axon density was determined experimentally, comparing 14
20 day and 28 day neuroimmunophilin FKBP ligand treatments. The effects of treatment with neuroimmunophilin FKBP ligands on retinal ganglion cells and optic nerve axons was correlated.

Surgical Procedures

25 Adult male Sprague Dawley rats (3 months old, 225-250 grams) were anesthetized with a ketamine

(87mg/kg) and xylazine (13mg/kg) mixture. Retinal ganglion cells were pre-labeled by bilateral stereotaxic injection of the fluorescent retrogradely transported marker fluoro-gold (FG, 0.5 microliters of 2.5% solution in saline) at the coordinates of the LGNd (4.5 millimeters post β , 3.5 millimeters lateral, 4.6 millimeters below dura). Four days later, FG labeled rats underwent a second surgery for microsurgical bilateral intraorbital optic nerve transection 4-5 millimeters behind the orbit.

Experimental animals were divided into six experimental groups of six rats (12 eyes) per group. One group received a neuroimmunophilin FKBP ligand (10 milligrams per kg per day sc in PEG vehicle (20 percent propylene glycol, 20 percent ethanol, and 60 percent saline)) for 14 days. A second group received the same neuroimmunophilin FKBP ligand dose for 28 days. Each treated group had a corresponding sham/surgery and transection control group which received corresponding 14 or 28 day dosing with the vehicle only.

All animals were sacrificed 90 days after optic nerve transection and perfused pericardially with formalin. All eyes and optic nerves stumps were removed. Cases were excluded from the study if the optic nerve vasculature was damaged or if FG labeling

was absent in the retina.

Retinal Ganglion Cell Counts

Retinas were removed from eyes and prepared for wholemount analysis. For each group, five eyes with
5 dense and intense FG labeling were selected for quantitative analysis using a 20 power objective. Digital images were obtained from five fields in the central retina (3-4 millimeters radial to optic nerve head). FG labeled Large ($>18\text{ }\mu\text{m}$), medium ($12\text{-}16\text{ }\mu\text{m}$),
10 and small ($<10\text{ }\mu\text{m}$) ganglion cells and microglia were counted in five $400\text{ }\mu\text{m}$ by $400\text{ }\mu\text{m}$ fields per case, 5 cases per group.

Examination of Optic Nerves

Proximal and distal optic nerve stumps were
15 identified, measured, and transferred to 30% sucrose saline. The proximal stumps of five nerves were blocked and affixed to a chuck, and 10 micron cross sections were cut on a cryostat; one in ten sections were saved per set. Sections including the region 1-2
20 mm behind the orbit were reacted for RT97 neurofilament immunohistochemistry. Analysis of optic nerve axon density was performed using a 63 power oil immersion lens, a Dage 81 camera, and the Simple Image Analysis program. RT97 positive optic nerve axons
25 were counted in three $200\text{ }\mu\text{m}$ by $200\text{ }\mu\text{m}$ fields per nerve. The area of the nerve was also determined for

each case at 10 power.

As depicted graphically in Table I&II, the 14 day course of treatment with a neuroimmunophilin FKBP ligand provided moderate neuroprotection of retinal ganglion cells observed 28 days after optic nerve transection. However, by 90 days after transection, only 5% of the ganglion cell population remained viable.

90 days after optic nerve transection the number of axons persisting in the proximal stump of the optic nerve represented approximately one half of the number of surviving ganglion cells in groups of animals that received vehicle alone or the 14 day course of treatment with a neuroimmunophilin FKBP ligand. These results indicate that over half of the transected ganglion cell axons retract beyond the optic nerve head, and that treatment with a neuroimmunophilin FKBP ligand during the first 14 days after optic nerve transection is not sufficient to arrest this retraction.

As depicted graphically in Table I&II, more prolonged treatment with a neuroimmunophilin FKBP ligand during the 28 day course of treatment produced a moderate increase in retinal ganglion cell neuroprotection. Approximately 12% of the vulnerable retinal ganglion cell population was protected. A

Table 1.
Effect of prolonged GPI 1046 treatment on retinal ganglion cell survival,
optic nerve axon preservation, and myelination 90 days after optic nerve transection

GROUP	RGC Counts ¹	ON Axon density ²	ON head area (%sham)	% RGC's Rescued	increased ON axon density ³	Spared RGC population	ON axon Count ⁴	% surviving RGC's with ON axons	Proximal optic nerve myelin basic protein Density ⁵	Distal optic nerve myelin basic protein Density ⁵
Sham	290 ± 14.8	7600*	100%			120,000*	120,000	100%	normal	
ONT/Vehicle	35.9 ± 2.8	428 ± 34	68%	(87% loss)		14,855 *	4593	30.9%	52 ± 5.2 SEM % loss	31% shrinkage 52.1% loss
ONT/ 14 days GPI 1046	49 ± 5.3	569 ± 23	76%	5.3%	1.5X	20,275	6820	33.6%	1.6 ± 3.0 SEM % recovery	31% shrinkage 47% loss
ONT/ 28 days GPI 1046	67.9 ± 5.8*	1526 ± 120*	95%*	12.6%*	5.0X	28,096*	22,861*	81.4%	70 ± 6.3 SEM % recovery*	56% less shrinkage* 99% myelin preservation*

*Significance p < 0.01

¹ Mean density + SEM of Fluoro-gold labeled retinal ganglion cells (RGC) in 400 µm x 400 µm sample gridfields.

² mean density + SEM of RT97 neurofilament antibody labeled optic nerve (ON) axons in 200 µm x 200 µm region of interest

³ estimate for 200 µm x 200 µm region in normal optic nerve assuming 120,000 RGC axons in normal rat optic nerve, measured to be 0.630 mm² mean cross sectional area


⁴ adjusted for optic nerve diameter

⁵ calculated by multiplying axonal density by ON area

⁶ determined from 20X analysis of % areal coverage of optic nerve cross section

⁷ shrinkage determined by comparing cross sectional area to sham control, myelin levels determined by multiplying cross sectional area by myelin density

TABLE II



similar proportion (~50%) of optic nerve axon density sparing was also observed. These results demonstrate the startling result that extending the duration of treatment with a neuroimmunophilin FKBP ligands to 28 days after transection completely arrests the regression of damaged axons for essentially the entire surviving population of retinal ganglion cells.

Additional results are set forth in Tables III & IV.

TABLE III

Correlation between Retinal Ganglion Cell and Optic Nerve Axon Sparing at 90 days following optic nerve transection and 14 or 28 day GPI 1046 treatment





TABLE IV

GPI 1046 preserves optic nerve axons
in the proximal stump following transection



Row Numbers

Example 8

A patient is suffering from macular degeneration.
A derivative as identified above, alone or
in combination with one or more other neoplastic factors,
5 or a pharmaceutical composition comprising the same,
may be administered to the patient. A reduction in
vision loss, prevention of vision degeneration, and/or
promotion of vision regeneration are/is expected to
occur following treatment.

10

Example 9

A patient is suffering from glaucoma, resulting
in cupping of the optic nerve disc and damage to nerve
fibers. A derivative as identified above,
15 alone or in combination with one or more other neoplastic
factors, or a pharmaceutical composition comprising
the same, may be administered to the patient. A
reduction in vision loss, prevention of vision
degeneration, and/or promotion of vision regeneration
20 are/is expected to occur following treatment.

Example 10

A patient is suffering from cataracts requiring
surgery. Following surgery, a derivative
25 as identified above, alone or in combination with one
or more other neoplastic factors, or a pharmaceutical

composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 11

A patient is suffering from an impairment or blockage of retinal blood supply relating to diabetic retinopathy, ischemic optic neuropathy, or retinal artery or vein blockage. A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 12

A patient is suffering from a detached retina. A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or

promotion of vision regeneration are/is expected to occur following treatment.

Example 13

5 A patient is suffering from tissue damage caused by inflammation associated with uveitis or conjunctivitis. A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 14

15 A patient is suffering from photoreceptor damage caused by chronic or acute exposure to ultraviolet light. A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 15

A patient is suffering from optic neuritis. A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 16

A patient is suffering from tissue damage associated with a "dry eye" disorder. A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 17

Efficacy of representative compounds from different immunophilin ligand series in protecting retinal ganglion cell axons from degeneration following optic nerve transection is set forth in Table V.

**Efficacy of representative compounds from
different immunophilin ligand series
in protecting retinal ganglion cell axons from
degeneration following optic nerve transection**







Compound	Structure	Comments	RT-PCR GC analysis of retinal ganglion cell axons following optic nerve transection
B		Adamantyl Thioester of urea Ki rotomase = 149 nM Clearance = 7 µl/min	100.0 % ±5.2 % SEM
A GPI 1046		Ester Ki rotomase = 7.5 nM Clearance = 63.8 µl/min	60.5 % ±3.9 SEM
C		Sulfonamide Ki rotomase = 107 nM Clearance = 31.1 µl/min	60.4 % ±3.1 % SEM
D		Pipercolic sulfonamide Ki rotomase = nM Clearance = µl/min	58.4 % ±6.4 % SEM
E		Ester of pipercolic acid Ki rotomase = 20 nM Clearance = 41.8 µl/min	56.6 % ±9.4 % SEM
F		Proline heterocycle Analog of GPI 1046 Ki rotomase = 272 nM Clearance = ? µl/min	55.1 % ±5.9 % SEM

TABLE V









G		Pipericolic acid dimethyl ketone Ki rotomase >10,000 nM Clearance=? μ l/min	34.0% $\pm$ 4.8% SEM
H		Ki rotomase = nM Clearance=? μ l/min	30.3% $\pm$ 8.0% SEM
I		Ester of Thiourea Ki rotomase= 131 nM Clearance=8.0 μ l/min	23.8% $\pm$ 5.3 SEM
J		Ketone - analog of GPI 1046 Ki rotomase= 210nM Clearance=1.5 μ l/min	15.8% $\pm$ 4.5% SEM
K		Pipericolic acid Thioester Ki rotomase= 86nM Clearance= 4.5 μ l/min	13.0% $\pm$ 4.2% SEM
L		Prolyl acid Ki rotomase= >7743nM Clearance=5.2 μ l/min	7.8% $\pm$ 3.0% SEM
M		Thioester Ki rotomase =7nM Clearance=12.5 μ l/min	-6.3% $\pm$ 3.9% SEM
N		Ki rotomase = 722 nM Clearance= 21.9 μ l/ml	

TABLE V continued

Example 18

5 THE FKBP NEUROIMMUNOPHILIN LIGAND GPI-1046
ENHANCES RETINAL GANGLION CELL SURVIVAL
AND ARRESTS AXONAL DYING BACK
FOLLOWING OPTIC NERVE TRANSECTION

10

Transection of the mammalian optic nerve results in a brief period of abortive regeneration, but the majority of axotomized neurons die and the axons from many persisting ganglion cells die back beyond the optic nerve head. The present Example was designed to examine the neuroprotective effects of GPI-1046 following optic nerve transection.

15

Retinal ganglion cells in adult male Sprague
20 Dawley rats were retrogradely labeled by fluorogold injection in the LGNd and four days later the optic nerves were transected 5 mm behind the globe. Groups of animals received either GPI-1046 10mg/kg/day s.c. or vehicle for 28 days. All experimental animals and
25 controls were sacrificed 90 days after transection.

By 90 days only - 10% of the FG labeled ganglion cell population survived but less than half of these neurons maintained axons that extended past the optic nerve head, as detected with RT97 neurofilament immunohistochemistry. GPI-1046 treatment produced a moderate degree of perikaryal neuroprotection, sparing

30

25% of the ganglion cell population, and preserved the axons of virtually all protected neurons in the proximal stump of the transected nerve. These results indicate that treatment with the FKBP neuroimmunophilin ligand GPI-1046 produces a fundamental alteration in the pathological process following injury to CNS tracts.

These results also demonstrate that the small molecule FKBP neuroimmunophilin ligand GPI 1046 enhances neurite outgrowth in culture, enhance peripheral nerve regeneration, and stimulate sprouting within the CNS following partial deafferentation.

Example 19

5 NEUROIMMUNOPHILIN LIGANDS PROMOTE RECOVERY
FROM THE PERIPHERAL SENSORY NEUROPATHY ASSOCIATED
WITH STREPTOZOTOCIN-INDUCED DIABETES

10 Peripheral neuropathy is a common debilitating
complication of Type 2 diabetes in some 30-40% of
diabetic patients. Neurotrophic factors such as nerve
growth factor (NGF) are known to promote survival of
developing and adult neurons of the peripheral nervous
15 system (PNS), and have also been evaluated as
treatments for diabetic peripheral neuropathy. Some of
the selective ligands of the neuroimmunophilin FKBP-12
such as the small molecule GPI-1046, have also been
shown to promote repair and regeneration in the central
20 and peripheral nervous systems (Proc. Nat'l. Acad. Sci.
USA 94, 2019-2024, 1997).

 In this Example the potential therapeutic effects
of GPI-1046 were evaluated for its ability to improve
25 sensory function in the streptozotocin-induced diabetic
rat. The procedure involved using Male Wistar rats
which were given a single injection of streptozotocin
(65 mg/kg i.v.). Blood glucose levels were determined
weekly for the first three weeks and on the last week
30 of the experiment. Animals were evaluated weekly for
signs of sensory neuropathy using the conventional hot
plate and tail flick apparatus test procedures. After

six weeks, treatment either with GPI-1046 or vehicle was initiated.

5 The results demonstrated that behavioral testing
using the hot plate and the tail flick apparatus
indicated improvement in latency in lesioned animals
treated for 6 weeks with GPI-1046 at 10 mg/kg s.c. The
results also showed that GPI-1046 ameliorates the
behavioral sequelae of diabetic sensory neuropathy and
10 may offer some relief for patients suffering from
diabetic peripheral neuropathy.

Morris Watermaze/Aging and Memory Test Procedure

5

Aged rodents exhibit marked individual differences in performance on a variety of behavioral tasks, including two-choice spatial discrimination in a modified T-maze, spatial discrimination in a circular platform task,
10 passive avoidance, radial maze tasks, and spatial navigation in a water pool.

15

In all of these tasks, a proportion of aged rats or mice perform as well as the vast majority of young control animals, while other animals display severe impairments in memory function compared to young animals. For example, Fischer and colleagues showed that the proportion of rats displaying significant impairments in spatial navigation increases with age, (Fischer et al.
20 1991b) with 8% of all 12 month old, 45% of 18 month old, 53% of 24 month old, and 90% of all 30 month old rats displaying impairments in spatial acquisition of the Morris watermaze task relative to young controls.

25

Specifically, rodent spatial learning and memory decline during aging has been accepted by many investigators as an intriguing correlative animal model of human senile dementia. Cholinergic function in the hippocampus has

been extensively studied as a component of spatial learning in rodents, and declining hippocampal cholinergic function has been noted in parallel with the development of learning and memory impairments. In addition, other neurotransmitter systems have been shown to contribute to spatial learning, and to decline with age, such as the dopaminergic and noradrenergic, serotonergic, and glutamatergic systems.

Also, reports on age-related deficits of hippocampal long-term potentiation (LTP)-induction, a reduction in theta rhythm frequency, a loss of experience-dependent plasticity of hippocampal place-units, and reductions in hippocampal protein kinase C are in keeping with the concept that no single underlying pathology can be identified as the cause of age-related behavioral impairment in rodents. However, the various experimental therapeutic approaches that have been undertaken to improve memory function in aged rodents have been somewhat slanted towards the cholinergic hypothesis.

The Morris watermaze is widely used for assessing spatial memory formation and retention in experimental animals. The test depends on the animal's ability to utilize spatial visual information in order to locate a submerged escape platform in a water tank. It is important that the tank itself be as devoid of specific visual features

as possible - thus, it is always circular in shape, the sides are kept smooth and in uniform dull colors, and the water is rendered opaque with nontoxic watercolour pigment or powdered milk. This is to ensure that the animal navigates only by the use of more distant visual cues, or by the use of intra-maze cues specifically provided by the experimenter.

The tank is filled to a level which forces the animal to swim actively. Normal mice and rats react aversively to the swimming part of the test and will climb onto, and remain on, an escape platform from which they are removed to a heated resting cage.

If the platform is visible (i.e. above the surface), animals placed in the tank will quickly learn to home in on the platform and climb out onto it. Testing with a visible platform will also ensure that the experimental animals are not blind and show sufficient motivation and stamina to perform the task, which can be important in experiments involving aged rodents. If the platform is invisible (i.e. submerged just below the surface), normal animals learn to use distant visual cues in the test room for orientation in the test tank, and, when placed in the tank, will quickly home in on the approximate location of the platform and circle in that area until the platform is found.

The animals' path, speed, and swim time are tracked with a ceiling camera for later computerized analysis. Over the course of several successive trials, spatial learning can therefore be defined as a drop of distance swum, or time elapsed, from placement in the tank until escape onto the invisible platform.

The test can be adapted to assess several aspects of spatial memory: a) acquisition of a cued task, where the animal's ability to link one visual cue directly with the escape platform depends on cortical function (i.e. a ball is suspended over the escape platform and the animal learns to follow this cue to find the platform); b) acquisition of a spatial task, where the animal's ability to learn the location of a submerged escape platform based on a combination of distant visual cues is dependent upon hippocampal function (i.e. the animal learns to triangulate its position in the tank by visually aligning the paper-tower dispenser with the door and ceiling lamp); c) retention of a successfully acquired spatial task, which is predominantly dependant on cortical function (i.e. the animal must remember the spatial location of the platform over several weeks); d) a hippocampus-dependant reversal task where the animals must reacquire a new spatial platform location (i.e. the platform is moved to a new location between swim trials

and the animal must abandon its previous search strategy and acquire a new one).

5 These different modifications of the Morris watermaze procedure can be applied in sequence to the same set of experimental animals and allow for a thorough characterization of their spatial memory performance and its decline with normal ageing. Moreover, such a series of sequential memory tests sheds some light on the functional integrity of the specific brain systems involved in the acquisition and retention of spatial memory (e.g. rats with cholinergic lesions of the hippocampus may remember a platform location acquired weeks before, but persevere over the old platform location after the platform is moved).

10

15

Example 20

EFFECTS OF CHRONIC GPI-1046 ADMINISTRATION 20 ON SPATIAL LEARNING AND MEMORY IN AGED RODENTS

This Example shows the effects of chronic treatment with the systemically available FKBP-ligand GPI-1046 on spatial learning and memory in aged rodents.

25

The procedure involved using three-month old (young) and 18-19 month old male C57BL/6N-Nia (aged) mice which

habituated to the well known and conventional Morris watermaze during a 4 trials/day, 3-4 day visible platform training phase. Subsequent spatial acquisition testing was conducting as follows: All mice were given 4 trials/day (block), for 5 days. Maximum swim time was 90 seconds. Aged mice were allocated to an "aged impaired" group if their performance during blocks 4 or 5 of the acquisition phase was >1 S.D. above the mean of "young" mice, and to an "aged non-impaired" group if their performance was < 0.5 S.D. above the mean of "young" mice. Aged groups were then split into statistically similar "GPI-1046" and "vehicle" groups.

Daily treatment with 10mg/kg GPI-1046 was initiated 3 days after the end of acquisition training, and continued through retention testing. Retention testing began after 3 weeks of dosing using the same methods as the acquisition phase. Swim Distances (cm) were analyzed in a 7 X 5 ANOVA including Groups and Blocks (1-5) as factors in the analysis, treating Blocks as a repeated measure.

The results showed that planned contrasts revealed that there were significant differences between the "young", and "aged impaired-vehicle and GPI-1046" treated groups at the end of the acquisition phase, $F_{1,58} = 26.75$, $P=0.0001$, and $F_{1,58} = 17.70$, $P=0.0001$ respectively. While

there were no significant differences between the two
"aged impaired" groups, $F_{1,58} = 0.67$, $P = 0.42$. During
retention testing, however, "aged impaired-vehicle"
treated animals performed significantly poorer than "aged
5 impaired - GPI-1046", and "young" animals, $F_{1,59} = 8.11$,
 $P = 0.006$, and $F_{1,59} = 25.45$, $P = 0.0001$ respectively.
There was no longer any statistically significant
difference between the "young" and "aged impaired" - GPI-
1046" treated groups during the retention phase, $F_{1,59} =$
10 3.09, $P = 0.08$. In summary, systemic treatment with GPI-
1046 significantly enhanced spatial memory performance of
mice with age-related spatial memory impairments.

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such
5 modification are intended to be included within the scope of the following claims.

100

What is claimed is:

1. A method for treating a vision disorder,
5 improving vision, treating memory impairment or enhancing
memory performance in an animal, which comprises
administering to said animal an effective amount of a N-
linked sulfonamide of an N-heterocyclic carboxylic acid
or isostere thereof.

10 2. The method of claim 1, wherein the N-linked
sulfonamide of an N-heterocyclic carboxylic acid or
isostere thereof is immunosuppressive or non-
immunosuppressive.

15 3. The method of claim 1, wherein the N-linked
sulfonamide of an N-heterocyclic carboxylic acid or
isostere thereof has an affinity for an FKBP-type
immunophilin.

20 4. The method of claim 3, wherein the FKBP-type
immunophilin is FKBP-12.


25 5. The method of claim 1, wherein the vision
disorder is selected from the group consisting of: visual
impairments; orbital disorders; disorders of the lacrimal

101

appartus; disorders of the eyelids; disorders of the
conjunctiva; disorders of the cornea; cataract; disorders
of the uveal tract; disorders of the retina; disorders of
the optic nerve or visual pathways; free radical induced
5 eye disorders and diseases; immunologically-mediated eye
disorders and disorders; eye injuries; and symtoms and
complications of eye disease, eye disorder, or eye
injury.

10 6. The method of claim 1, which is for improving
naturally-occurring vision in an animal, in the absence
of any ophthalmologic disorder, disease, or injury.

15 7. The method of claim 1, wherein the N-linked
sulfonamide of an N-heterocyclic carboxylic acid or
isostere thereof is a compound having the formula (I):



where

20 n is 1-3;

102

R_1 is selected from the group consisting of hydrogen, C_1-C_9 straight or branched chain alkyl, C_2-C_9 straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

5 D is a bond, or a C_1-C_{10} straight or branched chain alkyl, C_2-C_{10} alkenyl or C_2-C_{10} alkynyl;

R_2 is a carboxylic acid or a carboxylic acid isostere; wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or carboxylic acid isostere is optionally substituted with one or more substituents selected from R^3 , where

R^3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C_1-C_6 straight or branched chain alkyl, C_2-C_6 straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO_2R^4 where R^4 is hydrogen or C_1-C_9 straight or branched chain alkyl or alkenyl;

20 or a pharmaceutically acceptable salt, ester or solvate thereof.



















8. The method of claim 7, wherein R_2 is a carbocycle or heterocycle containing any combination of CH_2 , O, S, or N in any chemically stable oxidation state, where any of the atoms of said ring structure are optionally

substituted in one or more positions with R^3 , wherein
 R^3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl,
alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy,
cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl,
5 thioalkyl, alkylthio, sulfonyl, C_1 - C_6 straight or branched
chain alkyl, C_2 - C_6 straight or branched chain alkenyl or
alkynyl, aryl, heteroaryl, carbocycle, heterocycle, and
 CO_2R^4 where R^4 is hydrogen or C_1 - C_6 straight or branched
chain alkyl or alkenyl.

10

9. The method of claim 7, wherein R_2 is selected
from the group below:

104



where the atoms of said ring structure R_2 may be optionally substituted at one or more positions with R^3 , wherein

R^3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, and CO_2R^4 where R^4 is hydrogen or C_1 - C_6 straight or branched chain alkyl or alkenyl.

10. The method of claim 7, wherein R_2 is selected from the group consisting of $-COOH$, $-SO_3H$, $-SO_2HNR^3$, $-PO_2(R^3)_2$, $-CN$, $-PO_3(R^3)_2$, $-OR^3$, $-SR^3$, $-NHCOR^3$, $-N(R^3)_2$, $-CON(R^3)_2$, $-CONH(O)R^3$, $-CONHNHSO_2R^3$, $-COHNSO_2R^3$, and $-CONR^3CN$.

11. The method of claim 7, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or isostere thereof is selected from the group consisting of:

(2S)-1-(phenylmethyl)sulfonyl-2-hydroxymethyl pyrrolidine;

(2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinetetrazole; and compounds 1-97 disclosed herein.

12. A pharmaceutical composition for treating a vision disorder, improving vision, treating memory impairment or enhancing memory performance in an animal, comprising:

- 5 a) an effective amount for treating a vision disorder, improving vision, treating memory impairment or enhancing memory performance in an animal of a N-linked sulfonamide of an N-heterocyclic carboxylic acid or isostere thereof; and
- 10 b) a pharmaceutically acceptable carrier.

13. The pharmaceutical composition of claim 12, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or isostere thereof is immunosuppressive

15 or non-immunosuppressive.

14. The pharmaceutical composition of claim 12, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or isostere thereof has an affinity for

20 an FKBP-type immunophilin.

15. The pharmaceutical composition of claim 14, wherein the FKBP-type immunophilin is FKBP-12.


25 16. The pharmaceutical composition of claim 12, wherein the vision disorder is selected from the group

107

consisting of: visual impairments; orbital disorders;
disorders of the lacrimal apparatus; disorders of the
eyelids; disorders of the conjunctiva; disorders of the
cornea; cataract; disorders of the uveal tract; disorders
of the retina; disorders of the optic nerve or visual
pathways; free radical induced eye disorders and
diseases; immunologically-mediated eye disorders and
disorders; eye injuries; and symptoms and complications of
eye disease, eye disorder, or eye injury.

17. The pharmaceutical composition of claim 12,
which is for improving naturally-occurring vision in an
animal, in the absence of any ophthalmologic disorder,
disease, or injury.

18. The pharmaceutical composition of claim 12,
wherein the N-linked sulfonamide of an N-heterocyclic
carboxylic acid or isostere thereof comprises a compound
of formula (I):



I

where

n is 1-3;

R₁ is selected from the group consisting of hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl;

R₂ is a carboxylic acid or a carboxylic acid isostere;

wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or carboxylic acid isostere is optionally substituted with one or more substituents selected from R³, where

R³ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R⁴ where R⁴ is hydrogen or C₁-C₉ straight or branched chain alkyl or alkenyl;


or a pharmaceutically acceptable salt, ester or solvate thereof.

19. The pharmaceutical composition of claim 18, wherein R₂ is a carbocycle or heterocycle containing any

combination of CH₂, O, S, or N in any chemically stable oxidation state, wherein any of the atoms of said ring structure are optionally substituted in one or more positions with R³.

5

20. The pharmaceutical composition of claim 18,
wherein R₂ is selected from the following group:




where the atoms of said ring structure may be optionally substituted at one or more positions with R^3 .

21. The pharmaceutical composition of claim 18,
5. wherein R_2 is selected from the group consisting of:
-COOH; -SO₃H; -SO₂HNR³; -PO₂(R³)₂; -CN; -PO₃(R³)₂; -OR³; -
SR³; -NHCOR³; -N(R³)₂; -CON(R³)₂; -CONH(O)R³; -CONHNHSO₂R³;
-COHNSO₂R³; and -CONR³CN.

- 10 22. The pharmaceutical composition of claim 18,
wherein the N-linked sulfonamide of an N-heterocyclic
carboxylic acid or isostere thereof is selected from the
group consisting of:
(2S)-1-(phenylmethyl)sulfonyl-2-hydroxymethyl
15 pyrrolidine;
(2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinetetrazole; and
compounds 1-97 disclosed herein.

Figure 1


GPI 1046 protects ganglion cells against degeneration due to 1 hour of retinal ischemia
Fluorogold labelled retinal ganglion cells in wholemount, 28 days after ischemic episode



A. Labelled retinal ganglion cells in the
Normal central retina




B. 1 hour of retinal ischemia produces
extensive loss of ganglion cells



B. Administration of GPI 1046 1 hour before retinal ischemia
and for 4 days subsequently produces significant protection of vulnerable retinal ganglion cells


GPI 1046 Protects retinal ganglion cell axons and prevents myelin degeneration
in the optic nerve induced by 1 hour of complete retinal ischemia,
toluidine blue stained optic nerve cross sections, 630X



A. Normal optic nerve



B. Vehicle treated optic nerve 28 days
after 1 hour complete retinal ischemia



C. GPI 1046 treated optic nerve 28 days
after 1 hour complete retinal ischemia

Figure 3

**GPI 1046 administration for 28 days
provides only moderate protection of
axotomized retinal ganglion cells**



Fluorogold labelled RGCs 90 days following transection.
Treatment with vehicle alone for 1st 28 days

Fluorogold labelled RGCs 90 days following transection.
Treatment with GPI 1046 for 1st 28 days
Treatment with vehicle alone for 1st 28 days






Figure 4

GPI 1046 prevents axonal degeneration in the proximal stump of the optic nerve
RT97 neurofilament immunohistochemistry,
optic nerve cross sections, 90 days after complete transection



A. Sham



B. Optic nerve transection (ONT) 90 days survival




C. optic nerve 90 days after transection,
GPI 1046 treatment days 1-28D. optic nerve 90 days after transection,
GPI 1046 treatment days 1-14


Figure 5

**GPI 1046 administration for 28 days provides only moderate protection
of axotomized retinal ganglion cells**

Fluorogold labelled retinal ganglion cells 90 days following transection



vehicle administered for 1st 28 days




GPI 1046 administered for 1st 28 days

**GPI 1046 administration for 28 days preserves optic nerve axons
of surviving retinal ganglion cells**

RT 97 neurofilament immunohistochemistry 90 days after transection




vehicle administered for 1st 28 days




GPI 1046 administered for 1st 28 days

Figure 6


Preservation of myelin in the proximal stump of the optic nerve 90 days after transection
14 vs 28 days treatment with GPI 1046 10mg/kg s.c.




Normal(sham) Optic nerve



90 days after optic nerve transection, vehicle treated



90 days after optic nerve transection, 14 days GPI 1046



90 days after optic nerve transection, 28 days GPI 1046

myelin basic protein immunohistochemistry (SMI-94), 20X






Figure 7
FKBP-12 immunohistochemistry labels oligodendroglia and axons in the normal optic nerve

Figure 8


GPI 1046 treatment prevents myelin degeneration
in the distal stump of the optic nerve
Myelin basic protein immunohistochemistry 90 days after transection




A. Normal optic nerve



B. Distal optic nerve stump
90 days after complete transection



C. Distal optic nerve stump
90 days after complete transection
GPI 1046 administered 1-14 days
after transection



D. Distal optic nerve stump
90 days after complete transection
GPI 1046 administered 1-28 days
after transection

Figure 9


GPI 1046 decreases neovascularization and prevents neuronal loss in the inner retinal in the Streptozotocin model of diabetic retinopathy

**A. Normal
retina
Cross section
Cresyl violet**

Outer Nuclear
layer (ONL)

Inner Nuclear
layer (INL)

Ganglion cell
layer (GCL)




**B. retina from
Streptozotocin
/vehicle case**

ONL

INL

GCL



**C. Retina from
Streptozotocin
/GPI 1046 case**

ONL

INL

GCL

