




(86) Date de dépôt PCT/PCT Filing Date: 1998/02/09
(87) Date publication PCT/PCT Publication Date: 1998/08/13
(45) Date de délivrance/Issue Date: 2007/04/10
(85) Entrée phase nationale/National Entry: 1999/07/08
(86) N° demande PCT/PCT Application No.: JP 1998/000522
(87) N° publication PCT/PCT Publication No.: 1998/034484
(30) Priorité/Priority: 1997/02/10 (JP9/26953)

(51) Cl.Int./Int.Cl. *A01N 51/00* (2006.01),
A01N 25/04 (2006.01), *A01N 25/10* (2006.01),
A01N 25/30 (2006.01)
(72) Inventeurs/Inventors:
KOIKE, MASAHIKO, JP;
SAKAI, TOMOYO, JP
(73) Propriétaire/Owner:
SUMITOMO CHEMICAL TAKEDA AGRO COMPANY,
LIMITED, JP
(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : SUSPENSION AQUEUSE AGROCHIMIQUE
(54) Title: AQUEOUS SUSPENSION OF AGROCHEMICAL



(57) **Abrégé/Abstract:**

The present invention relates to an aqueous suspension comprising (i) a compound of formula (I) or a salt thereof, (ii) a condensate of formaldehyde with an aromatic sulfonic acid or a salt thereof or a polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer. The aqueous suspension of the invention can be used with advantage as a stable aqueous suspension of low viscosity providing for excellent delivery from a container, with excellent dispersibility in diluent water and excellent long-term fluidity free from caking due to precipitation of the suspended particles.



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶:

A01N 51/00, 25/04, 25/30, 25/10

A1

(11) International Publication Number:

WO 98/34484

(43) International Publication Date:

13 August 1998 (13.08.98)

(21) International Application Number: PCT/JP98/00522

(22) International Filing Date: 9 February 1998 (09.02.98)

(30) Priority Data:

9/26953

10 February 1997 (10.02.97)

JP

(71) Applicant (for all designated States except US): TAKEDA
CHEMICAL INDUSTRIES, LTD. [JP/JP]; 1-1,
Doshomachi 4-chome, Chuo-ku, Osaka-shi, Osaka
541-0045 (JP).


(72) Inventors; and

(75) Inventors/Applicants (for US only): KOIKE, Masahiko
[JP/JP]; 7-9-1403, Kasuga 1-chome, Tsukuba-shi, Ibaraki
305-0821 (JP). SAKAI, Tomoyo [JP/JP]; 11-16-707,
Sengen 2-chome, Tsukuba-shi, Ibaraki 305-0047 (JP).(74) Agents: ASAHINA, Tadao et al.; Osaka Plant of Takeda
Chemical Industries, Ltd., 17-85, Jusohonmachi 2-chome,
Yodogawa-ku, Osaka-shi, Osaka 532-0024 (JP).(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY,
CA, CN, CU, CZ, EE, GE, GW, HU, ID, IL, IS, KG, KR,
KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO,
NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA,
US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW,
SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ,
MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI
patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE,
SN, TD, TG).**Published***With international search report.**Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*

(54) Title: AQUEOUS SUSPENSION OF AGROCHEMICAL

(57) Abstract

The present invention relates to an aqueous suspension comprising (i) a compound of formula (I) or a salt thereof, (ii) a condensate of formaldehyde with an aromatic sulfonic acid or a salt thereof or a polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer. The aqueous suspension of the invention can be used with advantage as a stable aqueous suspension of low viscosity providing for excellent delivery from a container, with excellent dispersibility in diluent water and excellent long-term fluidity free from caking due to precipitation of the suspended particles.



DESCRIPTION

Aqueous Suspension of Agrochemical

5 [Technical Field]

The present invention relates to an agricultural aqueous suspension characterized by excellent pesticidal activity, low viscosity, and excellent suspension stability.

10

[Background Art]

Heretofore, as dosage forms for agricultural pesticides, dusts and granules have been primarily used. Those dosage forms are prepared by formulating a small amount of a pesticidally active substance with a large amount of a carrier such as clay, bentonite, calcium carbonate, etc. and, therefore, such formulations must be applied in quantities as large as 2 to 6 kilograms/10 ares in order to insure a uniform distribution of the active substance. This imposes a great burden on the farmers and, moreover, there are many disadvantages in transportation and storage. For this reason, so-called concentrates, which are supplied in small packages to provide for a greater ease of handling, are being developed.

25


Those concentrates are rich in the active substances and can be applied after dispersion in a suitable diluent, such as water, in the field. Examples include emulsifiable concentrates and wettable powders. However, emulsifiable concentrates require large quantities of an organic solvent for dilution, while it is unavoidable with wettable powders that preparation of sprays in the field involves contaminations with flying solid particles. For those reasons, the recent trend is toward refraining from using those formulations. Wettable granules, which are

30

35

granular versions of wettable powders have almost overcome the drawbacks of wettable powders but because the size of particles dispersed in water is as large as about 10 and odd μm , application of a dispersion of high concentration causes clogging of the spray nozzle, among other disadvantages.

The compound of the formula (I):



15 or a salt thereof, is an agricultural pesticide having potent pest-controlling activity.

The demand exists today for development of a stable aqueous suspension of the compound (I) or a salt thereof which should have a low viscosity enough to permit smooth delivery from a container as well as satisfactory dispersibility in diluent water and long-term fluidity free from caking due to precipitation of the suspended particles.


25 [Disclosure of Invention]

The inventors of the present invention have made much research to overcome the above-mentioned disadvantages of the prior art and found that an aqueous suspension comprising (i) the compound (I) or a salt thereof, (ii) a condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or a polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer in combination may have a low viscosity of not more than 700 mPa·s with excellent dispersibility in diluent water and, in addition, have a long-term stability. The inventors

have made further research and have completed the present invention.

Namely, the present invention relates to:

(1) an aqueous suspension which comprises (i) a compound of the formula (I):



or a salt thereof, (ii) a condensate of formaldehyde with an aromatic sulfonic acid or a salt thereof or a polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer,


(2) the aqueous suspension as described in (1) above, wherein the aqueous suspension contains about 1 to 10 weight % of the condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate in the total aqueous suspension,

(3) the aqueous suspension as described in (1) above, wherein the absorptive water-soluble polymer is one or more (preferably one to three) polymers selected from the group consisting of polyvinylpyrrolidone, poly(alkylated vinylpyrrolidone), vinylpyrrolidone-styrene block copolymer and methyl vinyl ether-maleic anhydride copolymer,

(4) the aqueous suspension as described in (1) above, which further comprises an unabsorptive water-soluble polymer,

(5) the aqueous suspension as described in (4) above, wherein the unabsorptive water-soluble polymer is one or more (preferably one to three) polymers selected from the group consisting of a polymer in the polyol

- series, a polysaccharides and polyethylene glycol,
 (6) the aqueous suspension as described in (4) above,
 wherein the unabsorptive water-soluble polymer is one
 or more (preferably one to three) polymers selected
 5 from the group consisting of xanthan gum, starch and
 polyethylene glycol,
 (7) the aqueous suspension as described in (1) above,
 which further comprises an auxiliary suspending agent,
 (8) the aqueous suspension as described in (7) above,
 10 wherein the auxiliary suspending agent is one or more
 (preferably one to three) substances selected from the
 group consisting of montmorillonite, colloidal silicon
 oxide, silicon oxide-aluminum oxide colloidal mixture
 and sepiolite,
 15 (9) a method for producing the aqueous suspension as
 described in (1) above, which comprises suspending (i)
 the compound of the formula (I):



- 25 or a salt thereof, (ii) the condensate of formaldehyde
 with an aromatic sulfonic acid or a salt thereof or the
 polyoxyalkylene allyl phenyl ether sulfate, and (iii)
 the absorptive water-soluble polymer in water, and
 (10) a method for controlling a pest which comprises
 30 scattering the aqueous suspension as described in (1)
 above into a field.

[Best Mode for Carrying Out the Invention]

35 The compound (I) which is guanidine derivative for
 use in the aqueous suspension of the present invention
 occurs as trans (E) and cis (Z) isomers in respect of

the geometrical orientation of NO₂. Preferred, however, is (E)-1-(2-chlorothiazol-5-ylmethyl)-3-methyl-2-nitroguanidine (hereinafter referred to briefly as compound (Ia)).

5 The salt of compound (I) includes agrochemically acceptable salts with inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, phosphoric acid, sulfuric acid, perchloric acid, etc. and salts with organic acids such as formic acid,
10 acetic acid, tartaric acid, malic acid, citric acid, oxalic acid, succinic acid, benzoic acid, picric acid, p-toluenesulfonic acid, etc.

 The production method for the compound (I) or a salt thereof is described in, inter alia, JP-A
15 157308/1991.

 The proportion of compound (I) or a salt thereof in the aqueous suspension of the present invention is generally about 5-60 weight %, preferably about 5-50 weight %, and more preferably about 10-50 weight %,
20 further more preferably 15-45 weight %.

 The condensate of formaldehyde with aromatic sulfonic acid or a salt thereof which can be used in preparing the aqueous suspension of the present invention includes a condensate of formaldehyde with a
25 sulfonic acid such as a phenylsulfonic acid, an aryl-substituted phenylsulfonic acid, a phenolsulfonic acid, an aryl-substituted phenolsulfonic acid, p-toluenesulfonic acid, a naphthalene sulfonic acid, an alkyl-substituted naphthalene sulfonic acid, an
30 aralkyl-substituted naphthalene sulfonic acid, an anthracene sulfonic acid, or an alkyl-substituted anthracene sulfonic acid.

 The aryl group of the aryl-substituted phenylsulfonic acid or aryl-substituted phenolsulfonic
35 acid as mentioned above includes C₆₋₁₄ aryl groups such as phenyl and naphthyl. The number of the substituents

26456-117

6

is 1 to 3.

The alkyl group of the alkyl-substituted naphthalene sulfonic acid as mentioned above includes C₁₋₆ alkyl groups such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl and tert-butyl. The number of the substituents is 1 to 7.

The aralkyl group of the aralkyl-substituted naphthalene sulfonic acid as mentioned above includes phenyl-C₁₋₄ alkyl groups such as benzyl, phenethyl and phenylpropyl; benzhydryl; and trityl. The number of the substituents is 1 to 3.

The alkyl group of the alkyl-substituted anthracene sulfonic acid as mentioned above includes a C₁₋₆ alkyl group such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl and tert-butyl. The number of the substituents is 1 to 9.

The condensed degree between an aromatic sulfonic acid and formaldehyde is 1.2 to 30, preferably 1.2 to 10.

Among them as mentioned above, a naphthalene sulfonate-formaldehyde condensate is preferable.

The salt of the aromatic sulfonic acid includes alkaline metals such as sodium.

Specific examples of the condensate of formaldehyde with aromatic sulfonic acid or a salt thereof include New Kalgen^{*} FS-4 (sodium naphthalenesulfonate-formaldehyde condensate; Takemoto oil & fat co., Ltd.), New Kalgen^{*} 9131 (phenylphenolsulfonic acid-formaldehyde condensate; Takemoto oil & fat co., Ltd.), Lavelin^{*} FM-L (sodium β -naphthalenesulfonate-formaldehyde condensate; Daiichi Kogyo Seiyaku), New Kalgen^{*} WG-2 (sodium β -naphthalenesulfonate-formaldehyde condensate; Takemoto oil & fat co., Ltd.), Demol^{*} N (sodium β -naphthalenesulfonate-formaldehyde condensate; Kao Corporation), and Demol^{*} C (sodium aromaticsulfonate-

*Trade-mark

26456-117

7

formaldehyde condensate; Kao Corporation).

Among them, especially New Kalgen* FS-4 is preferable.

5 The polyoxyalkylene allyl phenyl ether sulfate includes those having a molecular weight of 500-5000.

The polyoxyalkylene in the polyoxyalkylene allyl phenyl ether sulfate includes polyoxyethylene, polyoxypropylene or polyoxybutylene. Especially, New Kalgen* FS-7 (Takemoto oil & fat co., Ltd.) which
10 contains mainly polyoxyethylene allyl phenyl ether sulfate is preferable.

The proportion of the condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate in the total
15 aqueous suspension of the present invention is generally about 1-10 weight %, preferably about 1-7 weight %.

One or more (preferable one to three) other
20 surfactants such as nonionic or anionic surfactants besides the condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate, may be used in combination for preparing the aqueous suspension of the present
25 invention.

The nonionic surfactant include a nonionic surfactant in the ether series such as polyoxyalkylene allyl phenyl ether, polyoxyethylene allyl phenyl ether-formaldehyde condensate, polyoxyethylene alkyl phenyl
30 ethers, polyoxyethylene nonyl phenyl ether, polyoxyethylene distyrenated phenyl ether, polyoxyethylene alkyl allyl ethers, polyoxyethylene alkyl ethers, and polyoxyethylene-polyoxypropylene block copolymer. The nonionic surfactant that can be
35 used in preparing the aqueous suspension of the invention is preferably one having an HLB number in the

*Trade-mark

26456-117

8

range of 10-20. The more preferred HLB range is 12-18. Among such surfactants, which are commercially available, are New Kalgen* FS-1 (Takemoto oil & fat co., Ltd., polyoxyalkylene allyl phenyl ether), New Kalgen* FS-11 (Takemoto oil & fat co., Ltd., polyoxyethylene allyl phenyl ether-formaldehyde condensate), Nonipol* 100 (Sanyo Chemical Industries, LTD., polyoxyethylene nonyl phenyl ether), Noigen* EA-177 (Daiichi Kogyo Seiyaku, polyoxyethylene distyrenated phenyl ether), Noigen* EA-142 (Daiichi Kogyo Seiyaku, polyoxyethylene alkyl allyl ether), and Noigen* EA-157 (Daiichi Kogyo Seiyaku, polyoxyethylene alkyl ether).

The anionic surfactant includes metal ligninsulfonates, polyoxyethylene allyl phenyl ether phosphate amine salts, polyoxyethylene alkyl phenyl ether phosphate amine salts, and sodium dialkylsulfosuccinates. To mention commercial products, there can be mentioned New Kalgen* WG-4 (Takemoto oil & fat co., Ltd., metal ligninsulfonate), New Kalgen* FS-3 (Takemoto oil & fat co., Ltd., polyoxyethylene allyl phenyl ether phosphate amine salt), and New Kalgen* FS-31 and FS-32 (Takemoto oil & fat co., Ltd., polyoxyethylene alkyl phenyl ether phosphate amine salt).

The proportion of such a surfactant which may be used in combination in the total aqueous suspension of the present invention is generally about 1-10 weight % and preferably about 1-5 weight %.

To broaden the spectrum of utility of the aqueous suspension of the present invention, the suspension may be supplemented with one or more (preferably one to three) other agrochemicals. As such supplemental agrochemicals, insecticides, bactericides, etc. can be mentioned, which if such are solid at atmospheric temperature. The following are examples of such agrochemicals.

[Insecticides]

*Trade-mark

26456-117

9

Pyridafenthion, dimethoate, PMP, CVMP, dimethyl-
vinphos, acephate, salithion, DEP, NAC, MTMC, MIPC,
PHC, MPMC, XMC, bendiocarb, pirimicarb, mesomile,
oxamyl, thiodicarb, cypermethrin, caltap hydrochloride,
5 thiocyclam, bensultap, diflubenzuron, teflubenzuron,
chlorfluazron, buprofezin, hexythiazox, phenbutatin
oxide, pyridaben, clofentezine, nitenpyram, etc.

[Bactericides]

Thiuram, captan, TPN, phthalide, trichlorophos
10 methyl, phosethyl, methyl thiophanate, benomyl,
carbendazole, thiabendazole, diethofencarb, iprodione,
vinclozolin, procymidone, fluorimide, oxycarboxin,
mepronil, flutolanil, bencyclane, metalaxyl, oxadixyl,
triadimefon, hexaconazole, trifolin, blastidicid S,
15 kasugamycin, polyoxin, validamycin A, mildéomycin,
PCNB, hydroxyisoxazole, dazomet, dimethirimol,
diclomezine, triazine, ferimzone, probenazole,
isoprothiolane, tricyclazole, pyroquilon, oxolinic
acid, etc.

20 The agrochemicals are not limited to the above-
mentioned ones but other suitable agrochemicals which
are solid at atmospheric temperature can be added.

The absorptive water-soluble polymer is protective
colloids which can absorb other hydrophobic substances
25 by a hydrophobic group in the molecular, and which
includes a polyvinylpyrrolidone, a poly(alkylated
vinylpyrrolidone), a vinylpyrrolidone-styrene block
copolymer and a methylvinyl ether-maleic anhydride
copolymer.

30 The polyvinylpyrrolidone includes those having a
mean molecular weight of 5,000-2,000,000 such as
Agrimer* 15 (mean molecular weight: about 8000), Agrimer*
30 (mean molecular weight: about 57500), Agrimer* 60
(mean molecular weight: about 406000) and Agrimer* 90
35 (mean molecular weight: about 1270000), any of which
are manufactured by ISP Ltd.

*Trade-mark

The alkyl of poly(alkylated vinylpyrrolidone) includes C₁₋₃₀ alkyl groups such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, pentyl, isopentyl, tert-pentyl, neopentyl, 5 hexyl, isohexyl, 2,2-dimethylbutyl, heptyl, 2-methylhexyl, octyl, 2,2,4-trimethylpentyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, icosyl, henicosyl, dococyl, tricosyl and 10 tetracosyl. Among them, C₁₋₂₄ alkyl groups are preferable. Especially, C₄, C₁₆ and C₂₀ alkyl groups are preferable. Moreover, degree of alkylation of the poly(alkylated vinylpyrrolidone) is preferably 5-90%, more preferably 5-15% and 40-60%. The mean molecular weight 15 of the poly(alkylated vinylpyrrolidone) is preferably about 1,000 to about 100,000. Examples of the poly(alkylated vinylpyrrolidone) include Agrimer*AL-10 (number of carbon atom in the alkyl group: 4, degree of alkylation: about 10%, mean molecular weight: about 70000), 20 Agrimer*AL-10LC (number of carbon atom in the alkyl group: 4, degree of alkylation: about 10%, mean molecular weight: about 70000), Agrimer*AL-25 (number of carbon atom in the alkyl group: 16, degree of alkylation: about 50%, mean molecular weight: about 9500), 25 and Agrimer*AL-904 (number of carbon atom in the alkyl group: 4, degree of alkylation: about 10%, mean molecular weight: about 16,000), any of which are manufactured by ISP Ltd.

The vinylpyrrolidone-styrene block copolymer 30 includes those having a mean molecular weight of about 40000 such as Agrimer*ST (ISP Ltd.).

The methylvinylether-maleic anhydride copolymer includes free acid form having a mean molecular weight of about 100,000 to 300,000 such as Agrimer*VEMA H-240 35 (ISP Ltd.), anhydride form having a mean molecular weight of about 100,000 to about 300,000 such as

*Trade-mark

26456-117

11

Agrimer^{*} VEMA AN216 (ISP Ltd.), or half esters form having a mean molecular weight of about 20,000 to about 60,000 such as Agrimer^{*} ES-22 (ISP Ltd.).

5 As other absorptive water-soluble polymer which may be used in combination for preparing the aqueous suspension, poly(sodium acrylate) and carboxymethylcellulose are also mentioned.

As the absorptive water-soluble polymer, poly(alkylated vinylpyrrolidone) is preferable.

10 The proportion of the absorptive water-soluble polymer in the total aqueous suspension of the present invention is generally about 0.05-5 weight %, preferably about 0.1-3 weight %, more preferably about 0.1-2 weight %.

15 For the purpose of stabilizing and adjusting the viscosity of the suspension, the aqueous suspension of the present invention may be further supplemented with an unabsorptive water-soluble polymer and/or an auxiliary suspending agent.

20 The unabsorptive water-soluble polymer includes polymers in the polyol series (e.g. Agrisol^{*} FL-104FA, Kao Corporation); polysaccharides such as gum arabic, gelatin, guar gum, xanthan gum (e.g. Rhodopol^{*} 23; Rhone-Poulenc), starch (e.g. Stabirose^{*} MKK 100; Matsutani Chemical Industry co., LTD.), starch derivatives such as hydrolyzed starch (e.g. Pine-Dex^{*}; Matsutani Chemical Industry co., LTD.), soluble-starch (e.g. Stabirose^{*} S-10; Matsutani Chemical Industry co., LTD.), enzymatically denatured starch (e.g. Amicol^{*} 6L; Nippon Starch Refining co., LTD.) and pregelatinized starch (e.g. Cornalpha^{*}; Nippon Starch Refining co., LTD.) and dextrans (e.g. dextrin, cyclodextrin); and polyethylene glycol (e.g. PEG 6000; Sanyo Chemical Industries, LTD.).

35 Among them, the polysaccharides such as xanthan gum and starch, and polyethylene glycol are preferable.

*Trade-mark

26456-117

12

When the unabsorptive water-soluble polymer is used, it is used in a proportion of generally about 0.05-10 weight %, preferably about 0.05-3 weight % based on the total suspension.

5 The auxiliary suspending agent is water-insoluble additive and includes montmorillonite (e.g. Kunipia* F; Kunimine Industries co., LTD., Laponite*; Nisshin Chemical Industries, LTD.), colloidal silicon oxide (e.g. Aerosil* series such as Aerosil* 200; Aerosil* 10 NIPPON), silicon oxide-aluminum oxide colloidal mixture (e.g. Aerosil* COK84: $\text{SiO}_2/\text{Al}_2\text{O}_3=82$ to 86 (%) / 14 to 18 (%); Aerosil* NIPPON) and sepiolite (e.g. Aidplus* ML70DS-NV; Mizusawa Industrial Chemicals Ltd.).

Among them, montmorillonite is preferable.

15 When the auxiliary suspending agent is used, it is used in proportion of generally about 0.05-10 weight %, preferably about 0.1-3 weight % based on the total suspension.

20 Those additives such as unabsorptive water-soluble polymers and auxiliary suspending agent can be used each alone or in combination.

 In addition, other additives for routine use in aqueous suspensions in general can also be incorporated in the aqueous suspension of the present invention.
25 For example, a pH control agent, an antifreeze agent, an antifoam agent, and/or an antiseptic agent can be added liberally but selectively according to the amount of the agrochemical used. The pH control agent may for example be citric acid monohydrate. The antifreeze
30 agent that can be used includes, but is not limited to, ethylene glycol, diethylene glycol, and glycerin. The antifoam agent may for example be a silicone series defoaming agent (e.g. Anti-foam E-20; KAO Corporation). The antiseptic agent includes, but is not limited to,
35 n-butyl p-hydroxybenzoate, potassium sorbate, and sorbic acid.

*Trade-mark

The pH control agent, when used, is added in a proportion of generally about 0.01-5 weight % or preferably about 0.05-3 weight % based on the total suspension. The antifreeze agent, when used, is used in a proportion of generally about 1-20 weight % or preferably about 3-10 weight % based on the total suspension. The antifoam agent, when used, is added in a proportion of generally about 0.05-0.5 weight % or preferably about 0.05-0.3 weight % based on the total suspension. The antiseptic agent, when used, is added in an amount of generally about 0.01-3 weight % or preferably about 0.01-1.5 weight % of the total suspension.

The aqueous suspension of the present invention can be produced by suspending (i) the compound (I) or a salt thereof, (ii) a condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or a polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer in water in accordance with the per se known method or any other technique analogous thereto. For example, there can be mentioned a process (1) which comprises mixing all the components together thoroughly in a high-speed mixer for about 20-60 minutes and pulverizing and dispersing the mixture by means of a wet mill such as Dynomill or a microfluidizer and a process (2) which comprises grinding bulk powders of the active substance, e.g. compound (I) or a salt thereof, in a dry mill such as Jetmizer and blending and dispersing the finely divided powders with the other components in a high-speed mixer for about 30-90 minutes.

As it may be understood from the above description of the production method, the effective ingredient for the aqueous suspension of the present invention is dispersed in finely divided form in water. In the presence of other powder components, too, the effective

ingredient is dispersed as efficiently in water. The mean diameter of the dispersed particles is not greater than about 10 μm , preferably 0.1-5 μm .

The crop plant to which the aqueous suspension of the present invention can be applied includes rice, wheat, barley, sugar beet, maize (Indian corn), cotton, vegetables (e.g. cabbage, Chinese cabbage, radish, cucumber, eggplant, potato, etc.), fruit-bearing trees (e.g. mandarin orange, peach, pear, etc.), tea, and tobacco.

The pest which can be controlled with the aqueous suspension of the present invention includes insects of the order Hemiptera {e.g. cabbage stink bug [Eurydema rugosum Motschulsky], black rice stink bug [Scotinophara lurida Burmeister], bean bug [Riptortus clavatus Thunberg], pear lace-bug [Stephanitis nashi Esaki et Takeya], brown planthopper [Nilaparvata lugens Stal], white-back planthopper [Sogota furcifera Horvath], smaller brown planthopper [Delphacodes striatella Fallén], green rice leafhopper [Nephotettix bipunctatus cincticeps Uhler], tea green leafhopper [Chlorita onukii Matsuda], greenhouse whitefly [Trialeurodes vaporariorum], arrow-head scale [Prontaspis yanonensis Kuwana], mulberry mealy bug [Pseudococcus comstocki Kuwana], soybean aphid [Aphis glycines Matsumura], turnip aphid [Rhopalosiphum pseudobrassicae Davis], cabbage aphid [Brevicoryne brassicae Linné], greenhouse aphid [Myzus persicae Sulzer], cotton aphid [Aphis gossypii Glover], apple aphid [Aphis pomi DeGeer], etc.}, the order Lepidoptera {e.g. tobacco cutworm [Prodenia litura Fabricius], diamond-back moth [Plutella maculipennis Curtis], cabbage butterfly [Pieris rapae Linné], Asiatic rice borer [Chilo suppressalis Walker], beet worm [Plusia nigrisigna Walker], oriental tobacco budworm [Helicoverpa assulta assulta Guenée], armyworm

[Leucania separata Walker], cabbage armyworm [Mamestra brassicae Linné], smaller tea tortrix [Adoxophyes orana Fischer von Röslerstamm] (apple), cotton leaf roller [Sylepta derogata Fabricius], grass leaf roller
 5 [Cnaphalocrocis medinalis Guénéée], potato tuberworm [Phthorimaea operculella Zeller], tea leaf roller [Caloptilia theivora Walsingham], smaller tea tortrix [Adoxophyes orana Fischer von Röslerstamm] (tea),
 etc.), the order Coleoptera {e.g. potato lady beetle
 10 [Epilachna sparsa orientalis Dieke], cucurbit leaf beetle [Rhaphidopalpa femoralis Motschulsky], striped cabbage flea-beetle [Phyllotreta vittata Fabricius], Colorado potato bug [Leptinotarsa decemlineata], beet
 tortoise beetle [Cassida nebulosa Linné], rice leaf
 15 beetle [Oulema oryzae Kuwayama], rice plant weevil [Echinocnemus squameus Billberg], vegetable weevil [Listroderes obliquus Klug], etc.), the order Diptera {e.g. oriental house fly [Musca domestica vicina Macquart], pale house mosquito [Culex pipiens pallens
 20 Coquillett], common horse fly [Tabanus trigonus Coquillett], onion maggot [Hylemya antiqua Meigen], seed-corn maggot [Hylemya platura Meigen], etc.), the order Orthoptera {e.g. Asiatic locust [Locusta migratoria Linné], African mole cricket [Gryllotalpa africana Palisot de Beauvois], etc.), reticulopterous
 25 insects {e.g. German cockroach [Blattella germanica Linné], smoky-brown cockroach [Periplaneta fuliginosa Serville], etc.), spider mites {e.g. two-spotted spider mite [Tetranychus urticae Koch], citrus red mite
 30 [Panonychus citri McGregor], Kanzawa spider mite [Tetranychus kanzawai Kishida], carmine mite [Tetranychus telarius Linné], European red mite [Panonychus ulmi Koch], Japanese citrus rust mite [Aculus pelekassi Keifer], etc.), nematodes {e.g. rice
 35 white-tip nematode [Aphelenchoides besseyi Christie] etc.) and so on.

The aqueous suspension of the present invention is a safe and useful agrochemical composition with low toxicity against plants and mammals. This agrochemical composition can be used either as it is or as dilution in a suitable medium, e.g. water, when used. The recommended method for application depends on the crop plant applied, but is generally scattering by means of spraying or dropping, or dipping to field, i.e. paddy field, plow-land, orchard, grass-land or non-crop land in accordance with the per se known method. The specific method for application includes aerial broadcast, soil treatment, folial application, nursery box treatment, hand application along rows, seed dressing, bed soil treatment, directed application to infesting pests, application to paddy flood water, etc.

The recommended dose of the aqueous suspension of the present invention depends on the concentration of the agrochemical active ingredient but is generally about 1-500 g, preferably about 1-200 g, as active ingredient, per 10 ares for paddy field and plow-land (growing tea, wheat, sugar beet, maize, potato, cotton, etc.) pests and orchard pests.

[Examples]

The following examples and test examples are intended to illustrate the present invention in further detail. The compound (Ia) used here was prepared in accordance with the procedure described in JP-A 157308/1991 (Table 4; Compound No. 19). It should be understood that all parts and percents (%) are by weight unless otherwise indicated.

[Example 1]

	Compound (Ia)	20 parts
	New Kalgen FS-4	5 parts
35	Agrimer 30	1 part
	Polyethylene glycol (PEG6000)	2 parts

	Rhodopol 23	0.1 part
	Anti-foam E-20	0.2 part
	Ethylene glycol	4 parts
	Kunipia F	1 part
5	n-Butyl p-hydroxybenzoate	0.1 part
	Water	add 100 parts

The above components were thoroughly mixed in a high-speed mixer and wet-pulverized (one pass) by means of Dynomill (Shinmaru Enterprises, 1.0 mm glass beads, packing rate 80%, peripheral speed 15 m/s) to provide an aqueous suspension containing 20% of compound (Ia).

The term of "add 100 parts" means to make 100 parts of aqueous suspension by adding water besides other components.

15 [Example 2]

	Compound (Ia)	20 parts
	New Kalgen FS-4	2 parts
	Agrimer AL-10	1 part
	Stavirose MKK 100	0.5 part
20	Anti-foam E-20	0.2 part
	Ethylene glycol	4 parts
	Aerosil COK84	1 part
	n-Butyl p-hydroxybenzoate	0.1 part
	Water	add 100 parts

25 Using the above components, the procedure of Example 1 was repeated to provide an aqueous suspension containing 20% of compound (Ia).

[Example 3]

	Compound (Ia)	20 parts
30	New Kalgen FS-7	3 parts
	Agrimer ST	1 part
	Anti-foam E-20	0.2 part
	Ethylene glycol	4 parts
	Aidplus ML70DS-NV	1 part
35	n-Butyl p-hydroxybenzoate	0.1 part
	Water	add 100 parts

Using the above components, the procedure of Example 1 was repeated to provide an aqueous suspension containing 20% of compound (Ia).

[Example 4]

5	Compound (Ia)	20 parts
	New Kalgen FS-7	3 parts
	Agrimer VEMA H-240	0.5 part
	Anti-foam E-20	0.2 part
	Ethylene glycol	4 parts
10	Kunipia F	1.5 part
	Citric acid monohydrate	0.1 part
	n-Butyl p-hydroxybenzoate	0.1 part
	Water	add 100 parts

Using the above components, the procedure of Example 1 was repeated to provide an aqueous suspension containing 20% of compound (Ia).

[Example 5]

	Compound (Ia)	40 parts
	New Kalgen FS-7	5 parts
20	Agrimer AL-30	1 part
	Anti-foam E-20	0.2 part
	Ethylene glycol	7 parts
	Aerosil COK84	1.5 part
	Rhodopol 23	0.2 part
25	n-Butyl p-Hydroxybenzoate	0.1 part
	Water	add 100 parts

Using the above components, the procedure of Example 1 was repeated to provide an aqueous suspension containing 40% of compound (Ia).

30 [Example 6]

	Compound (Ia)	21 parts
	New Kalgen FS-4	2 parts
	Agrimer AL-10	0.5 part
	Stabirose MKK 100	0.3 part
35	Rhodopol 23	0.1 part
	Kunipia F	0.8 part

n-Butyl p-hydroxybenzoate	0.1 part
Anti-foam E-20	0.2 part
Ethylene glycol	4 parts
Water	add 100 parts

5 Using the above components, the procedure of Example 1 was repeated to provide an aqueous suspension containing 21% of compound (Ia).

[Example 7]

Compound (Ia)	21 parts
10 New Kalgen FS-4	2 parts
Agrimer AL-10	0.5 part
Polyethylene glycol (PEG6000)	2.0 part
Rhodopol 23	0.05 part
Kunipia F	0.65 part
15 n-Butyl p-hydroxybenzoate	0.1 part
Anti-foam E-20	0.2 part
Ethylene glycol	4 parts
Water	add 100 parts

20 Using the above components, the procedure of Example 1 was repeated to provide an aqueous suspension containing 21% of compound (Ia).

[Example 8]

Compound (Ia)	21 parts
New Kalgen FS-4	2 parts
25 Agrimer AL-10	0.5 part
Polyethylene glycol (PEG6000)	2 parts
Kunipia F	0.8 part
n-Butyl p-hydroxybenzoate	0.1 part
Anti-foam E-20	0.2 part
30 Ethylene glycol	4 parts
Water	add 100 parts

 Using the above components, the procedure of Example 1 was repeated to provide an aqueous suspension containing 21% of compound (Ia).

35 [Test Example 1]

 Using the aqueous suspensions prepared in Examples

1-8, the viscosity, water dispersibility, stability of 10-fold diluted suspension and tendency toward caking of each aqueous suspension were evaluated by the following methods.

5 (1) Viscosity

Viscosity was measured with a Type B viscosimeter at 25°C.

(2) Water dispersibility

10 A hollow cylinder of 500 ml capacity was filled in with 500 ml of water. Then, 0.5 ml of the test suspension was added dropwise and the dispersion was evaluated according to the criteria: (A) excellent, (B) good, (C) poor, and (D) not dispersed.

(3) Stability test of 10-fold diluted suspension

15 A hollow cylinder of 500 ml capacity was filled in with 450 ml of water and 50 ml of diluted suspension (flowable), and then turned upside down 10 times.

20 The test suspensions were respectively allowed to stand for 15 hours and the degree of the separation of water-phase was investigated.

(4) Tendency toward caking

25 The test suspensions were respectively allowed to stand at room temperature for 12 months following preparation and the degree of caking on the container bottom was investigated.

The results are shown in Table 1.

[Table 1]

30	Sample	Viscosity (mPa·s)	Water disper- sibility	Degree of separation (%)	Caking
	Ex. 1	230	A	2.1	non
	Ex. 2	150	A	1.8	non
	Ex. 3	65	A	3.2	non
35	Ex. 4	370	A	3.0	non
	Ex. 5	280	A	1.1	non
	Ex. 6	290	A	2.1	non
	Ex. 7	390	A	2.4	non
40	Ex. 8	250	A	2.1	non

It is clear from Table 1 that the aqueous suspension of the present invention is of low viscosity, i.e. not higher than 700 mPa·s, with good water dispersibility and has a good shelf-life (stability of the stock suspension).

[Industrial Applicability]

The aqueous suspension comprising (i) the compound (I) or a salt thereof, (ii) the condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer can be used with advantage as a highly stable agrochemical preparation with a low viscosity, i.e. not higher than 700 mPa·s, thus providing for excellent delivery from a container with excellent dispersibility in diluent water and excellent long-term fluidity free from caking due to precipitation of the suspended particles.


26456-117

22

CLAIMS:

1. An aqueous suspension which comprises the following ingredients:

(i) a compound of the formula (I):



or an agrochemically acceptable salt thereof,

10 (ii) a condensate of formaldehyde with an aromatic sulfonic acid or a salt thereof or a polyoxyalkylene allyl phenyl ether sulfate, and

(iii) an absorptive water-soluble polymer,

wherein the aqueous suspension has a viscosity of not more than 700 mPa·s; and

15 the absorptive water-soluble polymer (iii) is at least one polymer selected from the group consisting of polyvinylpyrrolidone, poly(alkylated vinylpyrrolidone), vinylpyrrolidone-styrene block copolymer and methyl vinyl ether-maleic anhydride copolymer.

20 2. The aqueous suspension as claimed in claim 1, wherein the compound of the formula (I) is (E)-1-(2-chlorothiazol-5-ylmethyl)-3-methyl-2-nitroguanidine.

3. The aqueous suspension as claimed in claim 1 or 2,
25 which contains 1 to 10 weight % of the ingredient (ii) in the aqueous suspension.

26456-117

23

4. The aqueous suspension as claimed in any one of claims 1 to 3, wherein the absorptive water-soluble polymer is poly(alkylated vinylpyrrolidone).

5. The aqueous suspension as claimed in any one of
5 claims 1 to 3, wherein the absorptive water-soluble polymer is polyvinylpyrrolidone.

6. The aqueous suspension as claimed in any one of claims 1 to 5, wherein the ingredient (iii) is contained in an amount of 0.05-5 weight % based on the aqueous
10 suspension.

7. The aqueous suspension as claimed in any one of claims 1 to 6, which further comprises an unabsorptive water-soluble polymer.

8. The aqueous suspension as claimed in claim 7,
15 wherein the unabsorptive water-soluble polymer is at least one polymer selected from the group consisting of a polysaccharide, polyethylene glycol, and another polyol.

9. The aqueous suspension as claimed in claim 7,
wherein the unabsorptive water-soluble polymer is at least
20 one polymer selected from the group consisting of xanthan gum, starch and polyethylene glycol.

10. The aqueous suspension as claimed in any one of claims 7 to 9, wherein the unabsorptive water-soluble polymer is contained in an amount of 0.05-10 weight % based
25 on the aqueous solution.

11. The aqueous suspension as claimed in any one of claims 1 to 10, which further comprises an auxiliary suspending agent.


26456-117

24

12. The aqueous suspension as claimed in claim 11,
wherein the auxiliary suspending agent is at least one
substance selected from the group consisting of
montmorillonite, colloidal silicon oxide, silicon oxide-
5 aluminum oxide colloidal mixture and sepiolite.

13. The aqueous suspension as claimed in claim 11 or 12, wherein the auxiliary suspending agent is contained in an amount of 0.05-10 weight % based on the aqueous solution.

10 14. A method for producing the aqueous suspension as
defined in claim 1 which comprises suspending (i) the
compound of the formula (I):



or an agrochemically acceptable salt thereof, (ii) the condensate of formaldehyde with an aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate, and (iii) the absorptive water-soluble polymer in

20 water.

15. A method for controlling a pest which comprises scattering the aqueous suspension as defined in any one of claims 1 to 13 into a field.

16. The aqueous suspension as claimed in any one of
25 claims 1 to 13, wherein the ingredient (i) is contained in
an amount of 5-60 weight % based on the aqueous suspension.

17. The aqueous suspension as claimed in any one of claims 1 to 13 or claim 16, wherein the ingredient (ii) is

26456-117

25

the condensate of formaldehyde with aromatic sulfonic acid or a salt thereof.

18. The aqueous suspension as claimed in claim 17, wherein the ingredient (ii) is an alkali metal salt of naphthalene-sulfonate formaldehyde condensate.


19. The aqueous suspension as claimed in any one of claims 1 to 13 or claim 16, wherein the ingredient (ii) is the polyoxyalkylene allyl phenyl ether sulfate.

20. The aqueous suspension as claimed in claim 19, wherein the polyoxyalkylene allyl phenyl ether sulfate is polyoxyethylene allyl phenyl ether sulfate having a molecular weight of 500 - 5,000.

FETHERSTONHAUGH & CO.

OTTAWA, CANADA

PATENT AGENTS



(I)