

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 April 2007 (12.04.2007)

PCT

(10) International Publication Number
WO 2007/040650 A2

(51) International Patent Classification:

C07D 295/14 (2006.01) C07D 491/08 (2006.01)
C07D 487/08 (2006.01) A61K 31/496 (2006.01)

(21) International Application Number:

PCT/US2006/018799

(22) International Filing Date: 12 May 2006 (12.05.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/680,107 12 May 2005 (12.05.2005) US
60/718,618 20 September 2005 (20.09.2005) US
11/432,937 12 May 2006 (12.05.2006) US

(71) Applicant (for all designated States except US): **ABBOTT LABORATORIES** [US/US]; Dept. 377 Bldg Ap6a-1, 100 Abbott Park Road, Abbott Park, Illinois 60064-6008 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BRUNCKO, Milan** [US/US]; 13317 W. Heiden Circle, Lake Bluff, Illinois 60044 (US). **DING, Hong** [CN/US]; 7465 Clem, Gurnee, Illinois 60031 (US). **ELMORE, Steven, W.** [US/US]; 6405 Lone Tree Court, Gurnee, Illinois 60031 (US). **KUNZER, Aaron, R.** [US/US]; 322 Woodbury Court #2, Schaumburg, Illinois 60193 (US). **LYNCH, Christopher,**

L. [US/US]; 23530 112th Street United States Of America, Trevor, Wisconsin 53179 (US). **MCCELLAN, William, J.** [US/US]; 1212 N. Sheridan, Waukegan, Illinois 60085 (US). **PARK, Cheol Min** [KR/US]; 1407 Almaden Lane, Gurnee, Illinois 60031 (US). **SONG, Xiahong** [CN/US]; 4107 Greenleaf Court, # 203, Park City, Illinois 60085 (US). **WANG, Xilu** [CN/US]; 340656 Sulky Drive, Grayslake, IL 60030 (US).


(74) Agents: **DONNER, B., Gregory** et al.; Dept. 377 Bldg Ap6a-1, 100 Abbott Park Road, Abbott Park, Illinois 60064-6008 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,

[Continued on next page]

(54) Title: APOPTOSIS PROMOTERS



(57) Abstract: Compounds which inhibit the activity of anti-apoptotic family protein members, compositions containing the compounds and methods of treating diseases during which occur expression one or more than one of an anti-apoptotic family protein member are disclosed.

WO 2007/040650 A2



ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

- *without international search report and to be republished upon receipt of that report*

APOPTOSIS PROMOTERS

FIELD OF THE INVENTION

This invention comprises compounds which inhibit the activity of anti-apoptotic Bcl-2 family protein members, compositions containing the compounds and methods of treating diseases during which are expressed one or more than one of an anti-apoptotic family protein member.

5

BACKGROUND OF THE INVENTION

Anti-apoptotic family protein members are associated with a number of diseases and thus are under investigation as potential therapeutic drug targets. Important targets for interventional therapy are the Bcl-2 family of proteins which include, for example, Bcl-2, Bcl-Xl and Bcl-w. Recently inhibitors of Bcl-2 family members have been reported in the literature, see, for example, WO 2005/049594, US 6,720,338 and US 7,030,115. While this art teaches inhibitors having high binding to the target enzyme, this is only one of many parameters that must be considered as a compound is investigated for further or continued drug development. This invention is directed to a series of compounds that promote apoptosis and that demonstrate enhanced and unexpected properties with respect to cellular potency, oral bioavailability, pharmacodynamic activity, and/or efficacy.

10
15

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows comparative antitumorigenesis of EXAMPLE 1, etoposide and combinations thereof on B-cell lymphoma.

20

FIG. 2 shows comparative antitumorigenesis of EXAMPLE 1, vincristine and combinations thereof on B-cell lymphoma.

FIG. 3 shows comparative antitumorigenesis of EXAMPLE 1, CHOP and combinations thereof on B-cell lymphoma.

25

FIG. 4 shows comparative antitumorigenesis of EXAMPLE 1, rituximab and combinations thereof on B-cell lymphoma.

FIG. 5 shows comparative antitumorigenesis of EXAMPLE 1, rapamycin and combinations thereof on B-cell lymphoma.


30

FIG. 6 shows comparative antitumorigenesis of EXAMPLE 1, R-CHOP and combinations thereof on mantle cell lymphoma.

FIG. 7 shows comparative antitumorigenesis of EXAMPLE 1, bortezomib and combinations thereof on mantle cell lymphoma.

SUMMARY OF THE INVENTION


5 One embodiment of this invention comprises compounds having formula (II)



(II),

and therapeutically acceptable salts, prodrugs, salts of prodrugs and metabolites thereof, wherein X^3 is Cl or F;

10 X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $N(CH_3)_2$, $N(CH_3)(CH(CH_3)_2)$, 7-azabicyclo[2.2.1]heptan-1-yl or 2-oxa-5-azabicyclo[2.2.1]hept-5-yl, and R^0 is




, wherein

X^5 is CH_2 , $C(CH_3)_2$ or CH_2CH_2 ;

X^6 and X^7 are both hydrogen or are both methyl; and


15 X^8 is F, Cl, Br or I; or

X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $N(CH_3)(CH(CH_3)_2)$ or 7-azabicyclo[2.2.1]heptan-1-yl, and R^0 is




; or

X^4 is $N(CH_3)_2$ or morpholin-1-yl, and R^0 is



Another embodiment comprises compounds having formula (II), and therapeutically acceptable salts, prodrugs, salts of prodrugs and metabolites thereof, wherein X^3 is Cl or F;

X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $N(CH_3)_2$, $N(CH_3)(CH(CH_3)_2)$, 7-azabicyclo[2.2.1]heptan-1-yl or 2-oxa-5-azabicyclo[2.2.1]hept-5-yl, and R^0 is




, wherein

X^5 is CH_2 , $C(CH_3)_2$ or CH_2CH_2 ;

X^6 and X^7 are both hydrogen or are both methyl; and


X^8 is F, Cl, Br or I; or

X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $N(CH_3)(CH(CH_3)_2)$ or 7-azabicyclo[2.2.1]heptan-1-yl, and R^0 is




; or

X^4 is $N(CH_3)_2$ or morpholin-1-yl, and R^0 is



Still another embodiment comprises compounds having formula (II), and therapeutically acceptable salts, prodrugs, salts of prodrugs and metabolites thereof, wherein X^3 is Cl or F;


X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $N(CH_3)_2$, $N(CH_3)(CH(CH_3)_2)$, 7-azabicyclo[2.2.1]heptan-1-yl or 2-oxa-5-azabicyclo[2.2.1]hept-5-yl, and R^0 is



, wherein X^5 is CH_2 , $\text{C}(\text{CH}_3)_2$ or CH_2CH_2 , and X^6 and X^7 are both hydrogen or are both methyl; and X^8 is F, Cl, Br or I.

Still another embodiment comprises compounds having formula (II), and therapeutically acceptable salts, prodrugs, salts of prodrugs and metabolites thereof, wherein X^3 is Cl or F;


X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $\text{N}(\text{CH}_3)(\text{CH}(\text{CH}_3)_2)$ or 7-azabicyclo[2.2.1]heptan-1-yl;
 R^0 is



10 , wherein X^6 and X^7 are both hydrogen or are both methyl; and X^8 is F, Cl, Br or I.


Still another embodiment comprises compounds having formula (II), and therapeutically acceptable salts, prodrugs, salts of prodrugs and metabolites thereof, wherein X^3 is Cl or F;

15 X^4 is $\text{N}(\text{CH}_3)_2$ or morpholin-1-yl;
 R^0 is



; and X^8 is F, Cl, Br or I.

20 Still another embodiment comprises a compound having formula (II), and therapeutically acceptable salts, prodrugs, salts of prodrugs and metabolites thereof, wherein X^3 is F; X^4 is morpholin-1-yl;
 R^0 is



Still another embodiment comprises

- 5 N-(4-(4-((2-(4-chlorophenyl)-5,5-dimethyl-1-cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(morpholin-4-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
- 3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(morpholin-4-yl)-1-((phenylsulfanyl)methyl)propyl)amino)benzenesulfonamide,
- 10 3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)-1-cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(morpholin-4-yl)-1-((phenylsulfanyl)methyl)propyl)amino)benzenesulfonamide,
- N-(4-(4-((2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(isopropyl(methyl)amino)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
- 15 N-(4-(4-((2-(4-chlorophenyl)-1-cyclohepten-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(isopropyl(methyl)amino)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
- 3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-((1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl)-1-((phenylsulfanyl)methyl)propyl)amino)benzenesulfonamide,
- 20 N-(4-(4-((2-(4-chlorophenyl)cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(morpholin-4-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
- 25 4-(((1R)-3-(7-azabicyclo[2.2.1]hept-7-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-N-(4-(4-((2-(4-chlorophenyl)cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
- N-(4-(4-((2-(4-chlorophenyl)cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
- 30

- N-(4-(4-((2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)-5,5-
5 dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)-1-((phenylsulfanyl)methyl)propyl)amino)benzenesulfonamide,
N-(4-(4-((2-(4-chlorophenyl)cyclohept-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
10 3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)cyclohept-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)-1-((phenylsulfanyl)methyl)propyl)amino)benzenesulfonamide,
N-(4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(isopropyl(methyl)amino)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
15 N-(4-(4-((2-(4-chlorophenyl)cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(1,4-oxazepan-4-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
4-(((1R)-3-(azepan-1-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-N-(4-(4-((2-(4-chlorophenyl)-1-cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
20 N-(4-(4-((2-(4-chlorophenyl)cyclohept-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(dimethylamino)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,
25 3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(dimethylamino)-1-((phenylsulfanyl)methyl)propyl)amino)benzenesulfonamide,
3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(morpholin-4-yl)-1-((phenylsulfanyl)methyl)propyl)amino)benzenesulfonamide,
30 N-(4-(4-((2-(4-chlorophenyl)-5,5-dimethyl-1-cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(dimethylamino)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,

N-(4-(4-((4-(4-chlorophenyl)-5,6-dihydro-2H-pyran-3-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(morpholin-4-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,

3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(isopropyl(methyl)amino)-1-((phenylsulfanyl)methyl)propyl)amino)benzenesulfonamide,

3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(isopropyl(methyl)amino)-1-((phenylsulfanyl)methyl)propyl)amino)benzenesulfonamide,

N-(4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(morpholin-4-yl)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,

3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-((1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl)-1-((phenylsulfanyl)methyl)propyl)amino)benzenesulfonamide,

N-(4-(4-((2-(4-chlorophenyl)cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(dimethylamino)-1-((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,

N-(4-(4-((2-(4-chlorophenyl)cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-1-((phenylsulfanyl)methyl)-3-(pyrrolidin-1-yl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,

3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(dimethylamino)-1-((phenylsulfanyl)methyl)propyl)amino)benzenesulfonamide,

N-(4-(4-((2-(4-chlorophenyl)cyclohept-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-1-((phenylsulfanyl)methyl)-3-(pyrrolidin-1-yl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide,

3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)-5,5-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-1-((phenylsulfanyl)methyl)-3-(pyrrolidin-1-yl)propyl)amino)benzenesulfonamide,

3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-1-((phenylsulfanyl)methyl)-3-(pyrrolidin-1-yl)propyl)amino)benzenesulfonamide,

3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)cyclohept-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-1-((phenylsulfanyl)methyl)-3-(pyrrolidin-1-yl)propyl)amino)benzenesulfonamide,

N-(4-(4-((2-(4-chlorophenyl)cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-
5 (((1R)-3-(isopropyl(methyl)amino)-1-((phenylsulfanyl)methyl)propyl)amino)-3-
((trifluoromethyl)sulfonyl)benzenesulfonamide,

N-(4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-1-((phenylsulfanyl)methyl)-3-(pyrrolidin-1-yl)propyl)amino)-3-
((trifluoromethyl)sulfonyl)benzenesulfonamide,

10 3-((chloro(difluoro)methyl)sulfonyl)-N-(4-(4-((2-(4-chlorophenyl)cyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-1-((phenylsulfanyl)methyl)-3-(pyrrolidin-1-yl)propyl)amino)benzenesulfonamide,

N-(4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-(dimethylamino)-1-((phenylsulfanyl)methyl)propyl)amino)-3-
15 ((trifluoromethyl)sulfonyl)benzenesulfonamide,

N-(4-(4-((2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl)methyl)piperazin-1-yl)benzoyl)-4-(((1R)-3-((1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl)-1-
((phenylsulfanyl)methyl)propyl)amino)-3-((trifluoromethyl)sulfonyl)benzenesulfonamide

N-(4-(4-((4'-chloro(1,1'-biphenyl)-2-yl)methyl)-1-piperazinyl)benzoyl)-4-(((1R)-3-
20 (dimethylamino)-1-((phenylsulfanyl)methyl)propyl)amino)-3-
((trifluoromethyl)sulfonyl)benzenesulfonamide and

N-(4-(4-((4'-chloro(1,1'-biphenyl)-2-yl)methyl)-1-piperazinyl)benzoyl)-4-(((1R)-3-(4-
morpholinyl)-1-((phenylsulfanyl)methyl)propyl)amino)-3-
((trifluoromethyl)sulfonyl)benzenesulfonamide,

25 and therapeutically acceptable salts, prodrugs, salts of prodrugs and metabolites thereof.

Still another embodiment comprises compositions for treating diseases during which are expressed one or more than one of antiapoptotic Bcl-X_L protein, antiapoptotic Bcl-2 protein or antiapoptotic Bcl-w protein, said compositions comprising an excipient and a therapeutically effective amount of the compound having formula (II).

30 Still another embodiment comprises methods of treating diseases in a patient during which are expressed one or more than one of antiapoptotic Bcl-X_L protein, antiapoptotic Bcl-2 protein or antiapoptotic Bcl-w protein, said methods comprising administering to the patient a therapeutically effective amount of a compound having formula (II).

Still another embodiment comprises compositions comprising an excipient and a
35 therapeutically effective amount of the compound having formula (II) for treating diseases of

abnormal cell growth and/or dysregulated apoptosis, such as cancer, mesothelioma, bladder cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, ovarian cancer, breast cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, bone cancer, ovarian cancer, cervical cancer, colon cancer, rectal cancer, cancer of the anal region, stomach cancer, gastrointestinal (gastric, colorectal, and duodenal), chronic lymphocytic leukemia, esophageal cancer, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, testicular cancer, hepatocellular cancer (hepatic and biliary duct), primary or secondary central nervous system tumor, primary or secondary brain tumor, Hodgkin's disease, chronic or acute leukemia, chronic myeloid leukemia, lymphocytic lymphomas, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, multiple myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, prostate cancer, small cell lung cancer, cancer of the kidney and ureter, renal cell carcinoma, carcinoma of the renal pelvis, neoplasms of the central nervous system, primary central nervous system lymphoma, non Hodgkin's lymphoma, spinal axis tumors, brain stem glioma, pituitary adenoma, adrenocortical cancer, gall bladder cancer, cancer of the spleen, cholangiocarcinoma, fibrosarcoma, neuroblastoma, retinoblastoma, or a combination thereof.

Still another embodiment comprises methods of treating mesothelioma, bladder cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, ovarian cancer, breast cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, bone cancer, ovarian cancer, cervical cancer, colon cancer, rectal cancer, cancer of the anal region, stomach cancer, gastrointestinal (gastric, colorectal, and duodenal), chronic lymphocytic leukemia, esophageal cancer, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, testicular cancer, hepatocellular cancer (hepatic and biliary duct), primary or secondary central nervous system tumor, primary or secondary brain tumor, Hodgkin's disease, chronic or acute leukemia, chronic myeloid leukemia, lymphocytic lymphomas, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, multiple myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, prostate cancer, small cell lung cancer, cancer of the kidney and ureter, renal cell carcinoma, carcinoma of the renal pelvis, neoplasms of the central nervous system, primary central nervous system lymphoma,

non Hodgkin's lymphoma, spinal axis tumors, brains stem glioma, pituitary adenoma, adrenocortical cancer, gall bladder cancer, cancer of the spleen, cholangiocarcinoma, fibrosarcoma, neuroblastoma, retinoblastoma, or a combination of one or more of the above cancers in a patient, said methods comprising administering thereto a therapeutically effective amount of a compound having formula (II).

Still another embodiment comprises compositions for treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, prostate cancer, small cell lung cancer and spleen cancer, said compositions comprising an excipient and a therapeutically effective amount of the compound having formula (II).

Still another embodiment comprises methods of treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer, hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, prostate cancer, small cell lung cancer and spleen cancer in a patient, said methods comprising administering to the patient a therapeutically effective amount of a compound having formula (II).

Still another embodiment comprises compositions for treating diseases in a patient during which are expressed one or more than one of antiapoptotic Bcl-X_L protein, antiapoptotic Bcl-2 protein or antiapoptotic Bcl-w protein, said compositions comprising an excipient and a therapeutically effective amount of the compound having formula (II) and a therapeutically effective amount of one additional therapeutic agent or more than one additional therapeutic agent.

Still another embodiment comprises methods of treating diseases in a patient during which is expressed one or more than one of antiapoptotic Bcl-X_L protein, antiapoptotic Bcl-2 protein or antiapoptotic Bcl-w protein, said methods comprising administering to the patient a therapeutically effective amount of a compound having formula (II) and a therapeutically effective amount of one additional therapeutic agent or more than one additional therapeutic agent.

Still another embodiment comprises compositions for treating mesothelioma, bladder cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, ovarian cancer, breast cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma

of the vulva, bone cancer, ovarian cancer, cervical cancer, colon cancer, rectal cancer, cancer of the anal region, stomach cancer, gastrointestinal (gastric, colorectal, and duodenal), chronic lymphocytic leukemia, esophageal cancer, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, testicular cancer, hepatocellular cancer (hepatic and biliary duct), primary or secondary central nervous system tumor, primary or secondary brain tumor, Hodgkin's disease, chronic or acute leukemia, chronic myeloid leukemia, lymphocytic lymphomas, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, multiple myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, prostate cancer, small cell lung cancer, cancer of the kidney and ureter, renal cell carcinoma, carcinoma of the renal pelvis, neoplasms of the central nervous system, primary central nervous system lymphoma, non Hodgkin's lymphoma, spinal axis tumors, brains stem glioma, pituitary adenoma, adrenocortical cancer, gall bladder cancer, cancer of the spleen, cholangiocarcinoma, fibrosarcoma, neuroblastoma, retinoblastoma, or a combination of one or more of the above cancers, said compositions comprising an excipient and therapeutically effective amount of a compound having formula (II) and one additional therapeutic agent or more than one additional therapeutic agent.

Still another embodiment comprises methods of treating mesothelioma, bladder cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, ovarian cancer, breast cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, bone cancer, ovarian cancer, cervical cancer, colon cancer, rectal cancer, cancer of the anal region, stomach cancer, gastrointestinal (gastric, colorectal, and duodenal), chronic lymphocytic leukemia, esophageal cancer, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, testicular cancer, hepatocellular cancer (hepatic and biliary duct), primary or secondary central nervous system tumor, primary or secondary brain tumor, Hodgkin's disease, chronic or acute leukemia, chronic myeloid leukemia, lymphocytic lymphomas, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, multiple myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, prostate cancer, small cell lung cancer, cancer of the kidney and ureter, renal cell carcinoma, carcinoma of the renal pelvis, neoplasms of the central nervous system, primary central nervous system lymphoma, non Hodgkin's lymphoma, spinal axis tumors, brains stem glioma, pituitary adenoma,

adrenocortical cancer, gall bladder cancer, cancer of the spleen, cholangiocarcinoma, fibrosarcoma, neuroblastoma, retinoblastoma, or a combination of one or more of the above cancers in a patient, said methods comprising administering thereto therapeutically effective amounts of a compound having formula (II) and one additional therapeutic agent or more than
5 one additional therapeutic agent.

Still another embodiment comprises methods of treating mesothelioma, bladder cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, ovarian cancer, breast cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma
10 of the vulva, bone cancer, ovarian cancer, cervical cancer, colon cancer, rectal cancer, cancer of the anal region, stomach cancer, gastrointestinal (gastric, colorectal, and duodenal), chronic lymphocytic leukemia, esophageal cancer, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, testicular
15 cancer, hepatocellular cancer (hepatic and biliary duct), primary or secondary central nervous system tumor, primary or secondary brain tumor, Hodgkin's disease, chronic or acute leukemia, chronic myeloid leukemia, lymphocytic lymphomas, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies of T-cell or B-cell origin, melanoma, multiple myeloma, oral cancer, ovarian cancer, non-small cell lung cancer, prostate cancer, small cell
20 lung cancer, cancer of the kidney and ureter, renal cell carcinoma, carcinoma of the renal pelvis, neoplasms of the central nervous system, primary central nervous system lymphoma, non Hodgkin's lymphoma, spinal axis tumors, brain stem glioma, pituitary adenoma, adrenocortical cancer, gall bladder cancer, cancer of the spleen, cholangiocarcinoma, fibrosarcoma, neuroblastoma, retinoblastoma, or a combination of one or more of the above
25 cancers in a patient, said methods comprising administering thereto therapeutically effective amounts of a compound having formula (II) and one or more than one of etoposide vincristine CHOP, rituximab, rapamycin, R-CHOP or bortezomib.

Still another embodiment comprises methods of treating B-cell lymphoma in a patient comprising administering thereto a therapeutically acceptable amounts of a compound having
30 formula (II) and etoposide.

Still another embodiment comprises methods of treating B-cell lymphoma in a patient comprising administering thereto therapeutically acceptable amounts of a compound having formula (II) and vincristine.

Still another embodiment comprises methods of treating B-cell lymphoma in a patient comprising administering thereto therapeutically acceptable amounts of a compound having formula (II) and CHOP.

5 Still another embodiment comprises methods of treating B-cell lymphoma in a patient comprising administering thereto therapeutically acceptable amounts of a compound having formula (II) and rituximab.

Still another embodiment comprises methods of treating B-cell lymphoma in a patient comprising administering thereto therapeutically acceptable amounts of a compound having formula (II) and rapamycin.

10 Still another embodiment comprises methods of treating mantle cell lymphoma in a patient comprising administering thereto therapeutically acceptable amounts of a compound having formula (II) and R-CHOP.

15 Still another embodiment comprises methods of treating mantle cell lymphoma in a patient comprising administering thereto therapeutically acceptable amounts of a compound having formula (II) and bortezomib.

DETAILED DESCRIPTION OF THE INVENTION

Variable moieties of compounds herein are represented by identifiers (capital letters with numerical and/or alphabetical superscripts) and may be specifically embodied.

20 It is meant to be understood that proper valences are maintained for all moieties and combinations thereof and that monovalent moieties having more than one atom are attached through their left ends.

It is also meant to be understood that a specific embodiment of a variable moiety may be the same or different as another specific embodiment having the same identifier.

25 The term "antitumorogenesis," as used herein, means reduction of tumor growth.

Compounds of this invention may contain asymmetrically substituted carbon atoms in the R or S configuration, wherein the terms "R" and "S" are as defined in Pure Appl. Chem. (1976) 45, 13-10. Compounds having asymmetrically substituted carbon atoms with equal amounts of R and S configurations are racemic at those atoms. Atoms having excess of one configuration over the other are assigned the configuration in excess, preferably an
30 excess of about 85%-90%, more preferably an excess of about 95%-99%, and still more preferably an excess greater than about 99%. Accordingly, this invention is meant to embrace racemic mixtures and relative and absolute diastereoisomers of the compounds thereof.

Compounds of this invention may also contain carbon-carbon double bonds or carbon-nitrogen double bonds in the Z or E configuration, in which the term "Z" represents the larger two substituents on the same side of a carbon-carbon or carbon-nitrogen double bond and the term "E" represents the larger two substituents on opposite sides of a carbon-carbon or carbon-nitrogen double bond. The compounds of this invention may also exist as a mixture of "Z" and "E" isomers.

Compounds of this invention may also exist as tautomers or equilibrium mixtures thereof wherein a proton of a compound shifts from one atom to another. Examples of tautomers include, but are not limited to, keto-enol, phenol-keto, oxime-nitroso, nitro-aci, imine-enamine and the like.

Compounds having formula (II) having NH, C(O)OH, OH or SH moieties may have attached thereto prodrug-forming moieties. The prodrug-forming moieties are removed by metabolic processes and release the compounds having the freed NH, C(O)OH, OH or SH in vivo. Prodrugs are useful for adjusting such pharmacokinetic properties of the compounds as solubility and/or hydrophobicity, absorption in the gastrointestinal tract, bioavailability, tissue penetration, and rate of clearance.

Metabolites of compounds having formula (II), produced by in vitro or in vivo metabolic processes, may also have utility for treating diseases associated with expression of an anti-apoptotic family protein member such as of BCL-X_L protein, Bcl-2 protein or Bcl-w protein.

Compounds having formula (II) may also be radiolabeled with a radioactive isotope such as a radioactive isotope of carbon (i.e. ¹³C), hydrogen (i.e. ³H), nitrogen (i.e. ¹⁵N), phosphorus (i.e. ³²P), sulfur (i.e. ³⁵S) or iodide (i.e. ¹²⁵I). Radioactive isotopes may be incorporated into the compounds having formula (II) by reacting the same and a radioactive derivitizing agent or by incorporating a radiolabeled intermediate into their syntheses. The radiolabeled compounds of formula (II) are useful for both prognostic and diagnostic applications as well as for in vivo and in vitro imaging.

Certain precursor compounds which may be metabolized in vitro or in vivo to form compounds having formula (II) may also have utility for treating diseases associated with expression of an anti-apoptotic family protein member such as of BCL-X_L protein, Bcl-2 protein or Bcl-w protein.

Compounds having formula (II) may exist as acid addition salts, basic addition salts or zwitterions. Salts of compounds having formula (II) are prepared during their isolation or following their purification. Acid addition salts are those derived from the reaction of a compound having formula (II) with acid. Accordingly, salts including the acetate, adipate,

alginate, bicarbonate, citrate, aspartate, benzoate, benzenesulfonate (besylate), bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, formate, fumarate, glycerophosphate, glutamate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, lactobionate, lactate, maleate, mesitylenesulfonate, methanesulfonate, naphthylenesulfonate, 5 nicotinate, oxalate, pamoate, pectinate, persulfate, phosphate, picrate, propionate, succinate, tartrate, thiocyanate, trichloroacetic, trifluoroacetic, para-toluenesulfonate and undecanoate salts of the compounds having formula (II) are meant to be embraced by this invention. Basic addition salts of compounds are those derived from the reaction of the compounds having formula (II) with the bicarbonate, carbonate, hydroxide or phosphate of cations such as lithium, sodium, potassium, calcium and magnesium. 10

Compounds having formula (II) may be administered, for example, buccally, ophthalmically, orally, osmotically, parenterally (intramuscularly, intraperitoneally intrasternally, intravenously, subcutaneously), rectally, topically, transdermally, vaginally and intraarterially as well as by intraarticular injection, infusion, and placement in the body, 15 such as, for example, the vasculature by means of, for example, a stent.

Therapeutically effective amounts of a compound having formula (II) depend on recipient of treatment, disease treated and severity thereof, composition comprising it, time of administration, route of administration, duration of treatment, potency, rate of clearance and whether or not another drug is co-administered. The amount of a compound having 20 formula (II) used to make a composition to be administered daily to a patient in a single dose or in divided doses is from about 0.03 to about 200 mg/kg body weight. Single dose compositions contain these amounts or a combination of submultiples thereof.

Compounds having formula (II) may be administered with or without an excipient. Excipients include, but are not limited to, encapsulators and additives such as absorption 25 accelerators, antioxidants, binders, buffers, coating agents, coloring agents, diluents, disintegrating agents, emulsifiers, extenders, fillers, flavoring agents, humectants, lubricants, perfumes, preservatives, propellants, releasing agents, sterilizing agents, sweeteners, solubilizers, wetting agents, mixtures thereof and the like.

Excipients for preparation of compositions comprising a compound having 30 formula (II) to be administered orally include, but are not limited to, agar, alginic acid, aluminum hydroxide, benzyl alcohol, benzyl benzoate, 1,3-butylene glycol, carbomers, castor oil, cellulose, cellulose acetate, cocoa butter, corn starch, corn oil, cottonseed oil, cross-povidone, diglycerides, ethanol, ethyl cellulose, ethyl laureate, ethyl oleate, fatty acid esters, gelatin, germ oil, glucose, glycerol, groundnut oil, hydroxypropylmethyl cellulose, 35 isopropanol, isotonic saline, lactose, magnesium hydroxide, magnesium stearate, malt,

mannitol, monoglycerides, olive oil, peanut oil, potassium phosphate salts, potato starch, povidone, propylene glycol, Ringer's solution, safflower oil, sesame oil, sodium carboxymethyl cellulose, sodium phosphate salts, sodium lauryl sulfate, sodium sorbitol, soybean oil, stearic acids, stearyl fumarate, sucrose, surfactants, talc, tragacanth, 5 tetrahydrofurfuryl alcohol, triglycerides, water, mixtures thereof and the like. Excipients for preparation of compositions comprising a compound having formula (II) to be administered ophthalmically or orally include, but are not limited to, 1,3-butylene glycol, castor oil, corn oil, cottonseed oil, ethanol, fatty acid esters of sorbitan, germ oil, groundnut oil, glycerol, isopropanol, olive oil, polyethylene glycols, propylene glycol, sesame oil, water, mixtures 10 thereof and the like. Excipients for preparation of compositions comprising a compound having formula (II) to be administered osmotically include, but are not limited to, chlorofluorohydrocarbons, ethanol, water, mixtures thereof and the like. Excipients for preparation of compositions comprising a compound having formula (II) to be administered parenterally include, but are not limited to, 1,3-butanediol, castor oil, corn oil, cottonseed oil, 15 dextrose, germ oil, groundnut oil, liposomes, oleic acid, olive oil, peanut oil, Ringer's solution, safflower oil, sesame oil, soybean oil, U.S.P. or isotonic sodium chloride solution, water, mixtures thereof and the like. Excipients for preparation of compositions comprising a compound having formula (II) to be administered rectally or vaginally include, but are not limited to, cocoa butter, polyethylene glycol, wax, mixtures thereof and the like.

20 This invention also comprises combination therapeutic methods of treating disease conditions involving abnormal cell growth and/or dysregulated apoptosis, such as cancer, in a patient comprising administering thereto a therapeutically effective amount of a pharmaceutical composition comprising a compound having formula (II) and a therapeutically effective amount of one or more than one additional therapeutic agents or ionizing radiation.

25 The combination therapeutic methods include administering compositions of a compound having formula (II) and one or more than one additional therapeutic agents or ionizing radiation to a patient using any desired dosing and/or scheduling regimen.

30 Compounds having formula (II) may be administered with one or more than one additional therapeutic agents, wherein the additional therapeutic agents include ionizing radiation or chemotherapeutic agents, wherein chemotherapeutic agents include, but are not limited to, carboplatin, cisplatin, cyclophosphamide, dacarbazine, dexamethasone, docetaxel, doxorubicin, etoposide, fludarabine, irinotecan, CHOP (C: Cytosan® (cyclophosphamide); H: Adriamycin® (hydroxydoxorubicin); O: Vincristine (Oncovin®); P: prednisone), paclitaxel, rapamycin, Rituxin® (rituximab), vincristine and the like.

Compounds having formula (II) are also expected to be useful as chemotherapeutic agents in combination with therapeutic agents that include, but are not limited to, angiogenesis inhibitors, antiproliferative agents, kinase inhibitors, receptor tyrosine kinase inhibitors, aurora kinase inhibitors, polo-like kinase inhibitors, bcr-abl kinase inhibitors, growth factor inhibitors, COX-2 inhibitors, non-steroidal anti-inflammatory drugs (NSAIDS), antimitotic agents, alkylating agents, antimetabolites, intercalating antibiotics, platinum containing agents, growth factor inhibitors, ionizing radiation, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biologic response modifiers, immunologicals, antibodies, hormonal therapies, retinoids/deltoids plant alkaloids, proteasome inhibitors, HSP-90 inhibitors, histone deacetylase inhibitors (HDAC) inhibitors, purine analogs, pyrimidine analogs, MEK inhibitors, CDK inhibitors, ErbB2 receptor inhibitors, mTOR inhibitors and combinations thereof as well as other antitumor agents.

Angiogenesis inhibitors include, but are not limited to, EGFR inhibitors, PDGFR inhibitors, VEGFR inhibitors, TIE2 inhibitors, IGF1R inhibitors, matrix metalloproteinase 2 (MMP-2) inhibitors, matrix metalloproteinase 9 (MMP-9) inhibitors, thrombospondin analogs such as thrombospondin-1 and N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-NHCH₂CH₃ or a salt thereof and analogues of N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-NHCH₂CH₃ such as N-Ac-GlyVal-D-alle-Ser-Gln-Ile-Arg-ProNHCH₂CH₃ or a salt thereof.

Examples of EGFR inhibitors include, but are not limited to, Iressa (gefitinib), Tarceva (erlotinib or OSI-774), Erbitux (cetuximab), EMD-7200, ABX-EGF, HR3, IgA antibodies, TP-38 (IVAX), EGFR fusion protein, EGF-vaccine, anti-EGFr immunoliposomes and Tykerb (lapatinib).

Examples of PDGFR inhibitors include, but are not limited to, CP-673,451 and CP-868596.

Examples of VEGFR inhibitors include, but are not limited to, Avastin (bevacizumab), Sutent (sunitinib, SU11248), Nexavar (sorafenib, BAY43-9006), CP-547,632, axitinib (AG13736), Zactima (vandetanib, ZD-6474), AEE788, AZD-2171, VEGF trap, Vatalanib (PTK-787, ZK-222584), Macugen, IM862, Pazopanib (GW786034), ABT-869 and angiozyme.

Examples of thrombospondin analogs include, but are not limited to, TSP-1 and ABT-510.

Examples of aurora kinase inhibitors include, but are not limited to, VX-680, AZD-1152 and MLN-8054.

Example of polo-like kinase inhibitors include, but are not limited to, BI-2536.

Examples of bcr-abl kinase inhibitors include, but are not limited to, Gleevec (imatinib) and Dasatinib (BMS354825).

Examples of platinum containing agents includes, but are not limited to, cisplatin, Paraplatin (carboplatin), eptaplatin, lobaplatin, nedaplatin, Eloxatin (oxaliplatin) or
5 satraplatin.

Examples of mTOR inhibitors includes, but are not limited to, CCI-779, rapamycin, temsirolimus, everolimus, RAD001, and AP-23573.

Examples of HSP-90 inhibitors includes, but are not limited to, geldanamycin, radicicol, 17-AAG, KOS-953, 17-DMAG, CNF-101, CNF-1010, 17-AAG-nab, NCS-683664,
10 Mycograb, CNF-2024, PU3, PU24FC1, VER49009, IPI-504, SNX-2112 and STA-9090.

Examples of histone deacetylase inhibitors (HDAC) includes, but are not limited to, Suberoylanilide hydroxamic acid (SAHA), MS-275, valproic acid, TSA, LAQ-824, Trapoxin, and Depsipeptide.

Examples of MEK inhibitors include, but are not limited to, PD325901,
15 ARRY-142886, ARRY-438162 and PD98059.

Examples of CDK inhibitors include, but are not limited to, flavopyridol, MCS-5A, CVT-2584, seliciclib (CYC-202, R-roscovitine), ZK-304709, PHA-690509, BMI-1040, GPC-286199, BMS-387,032, PD0332991 and AZD-5438.

Examples of COX-2 inhibitors include, but are not limited to, CELEBREX™
20 (celecoxib), parecoxib, deracoxib, ABT-963, MK-663 (etoricoxib), COX-189 Lumiracoxib), BMS347070, RS 57067, NS-398, Bextra (valdecoxib), paracoxib, Vioxx (rofecoxib), SD-8381, 4-Methyl-2-(3,4-dimethylphenyl)-1-(4-sulfamoyl-phenyl)-1H-pyrrole, T-614, JTE-522, S-2474, SVT-2016, CT-3, SC-58125 and Arcoxia (etoricoxib).

Examples of non-steroidal anti-inflammatory drugs (NSAIDs) include, but are not
25 limited to, Salsalate (Amigesic), Diflunisal (Dolobid), Ibuprofen (Motrin), Ketoprofen (Orudis), Nabumetone (Relafen), Piroxicam (Feldene), Naproxen (Aleve, Naprosyn), Diclofenac (Voltaren), Indomethacin (Indocin), Sulindac (Clinoril), Tolmetin (Tolectin), Etodolac (Lodine), Ketorolac (Toradol) and Oxaprozin (Daypro).

Examples of ErbB2 receptor inhibitors include, but are not limited to, CP-724-714,
30 CI-1033, (canertinib), Herceptin (trastuzumab), Omitarg (2C4, petuzumab), TAK-165, GW-572016 (Ionafarnib), GW-282974, EKB-569, PI-166, dHER2 (HER2 Vaccine), APC8024 (HER2 Vaccine), anti-HER2/neu bispecific antibody, B7.her2IgG3, AS HER2 trifunctional bispecific antibodies, mAB AR-209 and mAB 2B-1.

Examples of alkylating agents include, but are not limited to, nitrogen mustard N-
35 oxide, cyclophosphamide, ifosfamide, trofosfamide, Chlorambucil, melphalan, busulfan,

mitobronitol, carboquone, thiotepa, ranimustine, nimustine, temozolomide, AMD-473, altretamine, AP-5280, apaziquone, brostallicin, bendamustine, carmustine, estramustine, fotemustine, glufosfamide, KW-2170, mafosfamide, and mitolactol, carmustine (BCNU), lomustine (CCNU), Busulfan, Treosulfan, Decarbazine and Temozolomide.

5 Examples of antimetabolites include but are not limited to, methotrexate, 6-mercaptapurine riboside, mercaptopurine, uracil analogues such as 5-fluorouracil (5-FU) alone or in combination with leucovorin, tegafur, UFT, doxifluridine, carmofur, cytarabine, cytarabine ocfosfate, enocitabine, S-1, Alimta (premetrexed disodium, LY231514, MTA), Gemzar (gemcitabine), fludarabine, 5-azacitidine, capecitabine, cladribine, clofarabine, 10 decitabine, eflornithine, ethnylcytidine, cytosine arabinoside, hydroxyurea, TS-1, melphalan, nelarabine, nolatrexed, ocfosfate, disodium premetrexed, pentostatin, pelitrexol, raltitrexed, triapine, trimetrexate, vidarabine, vincristine, vinorelbine, mycophenolic acid, tiazofurin, Ribavirin, EICAR, hydroxyurea and deferoxamine.

 Examples of antibiotics include intercalating antibiotics but are not limited to, 15 aclerubicin, actinomycins such as actinomycin D, amrubicin, annamycin, adriamycin, bleomycin a, bleomycin b, daunorubicin, doxorubicin, elsamitruicin, epirbucin, glarbuicin, idarubicin, mitomycin C, nemorubicin, neocarzinostatin, peplomycin, pirarubicin, rebeccamycin, stimalamer, streptozocin, valrubicin, zinostatin and combinations thereof.

 Examples of topoisomerase inhibiting agents include, but are not limited to, one or 20 more agents selected from the group consisting of aclerubicin, amonafide, belotecan, camptothecin, 10-hydroxycamptothecin, 9-aminocamptothecin, diflomotecan, irinotecan HCL (Camptosar), edotecarin, epirubicin (Ellence), etoposide, exatecan, gimatecan, lurtotecan, orathecin (Supergen), BN-80915, mitoxantrone, pirarubicin, pixantrone, rubitecan, sobuzoxane, SN-38, tafluposide and topotecan.

25 Examples of antibodies include, but are not limited to, Rituximab, Cetuximab, Bevacizumab, Trastuzimab, specific CD40 antibodies and specific IGF1R antibodies,

 Examples of hormonal therapies include, but are not limited to, exemestane (Aromasin), leuprolide acetate, anastrozole (Arimidex), fosreltin (Zoladex), goserelin, doxercalciferol, fadrozole, formestane, tamoxifen citrate (tamoxifen), Casodex, Abarelix, 30 Trelstar, finasteride, fulvestrant, toremifene, raloxifene, lasofoxifene, letrozole, flutamide, bicalutamide, megestrol, mifepristone, nilutamide, dexamethasone, prednisone and other glucocorticoids.

 Examples of retinoids/deltoids include, but are not limited to, seocalcitol (EB 1089, CB 1093), lexacalcitrol (KH 1060), fenretinide, Aliretinoin, Bexarotene and LGD-1550.

Examples of plant alkaloids include, but are not limited to, vincristine, vinblastine, vindesine and vinorelbine.

Examples of proteasome inhibitors include, but are not limited to, bortezomib (Velcade), MG132, NPI-0052 and PR-171.

5 Examples of immunologicals include, but are not limited to, interferons and numerous other immune enhancing agents. Interferons include interferon alpha, interferon alpha-2a, interferon, alpha-2b, interferon beta, interferon gamma-1a, interferon gamma-1b (Actimmune), or interferon gamma-n1 and combinations thereof. Other agents include filgrastim, lentinan, sizofilan, TheraCys, ubenimex, WF-10, aldesleukin, alemtuzumab, 10 BAM-002, decarbazine, daclizumab, denileukin, gemtuzumab ozogamicin, ibritumomab, imiquimod, lenograstim, lentinan, melanoma vaccine (Corixa), molgramostim, OncoVAC-CL, sargamostim, tasonermin, teceleukin, thymalasin, tositumomab, Virulizin, Z-100, epratuzumab, mitumomab, oregovomab, pentumomab (Y-muHMFG1), Proveng (Dendreon), CTLA4 (cytotoxic lymphocyte antigen 4) antibodies and agents capable of 15 blocking CTLA4 such as MDX-010.

Examples of biological response modifiers are agents that modify defense mechanisms of living organisms or biological responses, such as survival, growth, or differentiation of tissue cells to direct them to have anti-tumor activity. Such agents include krestin, lentinan, sizofiran, picibanil and ubenimex.

20 Examples of pyrimidine analogs include, but are not limited to, 5-Fluorouracil, Floxuridine, Doxifluridine, Ratitrexed, cytarabine (ara C), Cytosine arabinoside, Fludarabine, and Gemcitabine.

Examples of purine analogs include but are not limited to, Mercaptopurine and thioguanine.

25 Examples of antimetabolic agents include, but are not limited to, ABT-751, paclitaxel, docetaxel, epothilone D (KOS-862) and ZK-EPO.

Compounds of the present invention are also intended to be used as a radiosensitizer that enhances the efficacy of radiotherapy. Examples of radiotherapy include but are not limited to, external beam radiotherapy (XBRT), or teletherapy, brachtherapy or sealed source 30 radiotherapy, unsealed source radiotherapy.

Additionally, compounds having formula (II) may be combined with other antitumor agents selected from the following agents: Genasense, Panitumumab, Zevalin, Bexar (Corixa), Abarelix, Alimta, EPO906, discodermolide, Neovastat, enzastaurin, Combrestatin A4P, ZD-6126, AVE-8062, DMXAA, Thymitaq, Temodar, Revlimid, Cypat, Histerelin, 35 Plenaizis, Atrasentan, Satraplatin, thalomid (Thalidomide), theratope, Temilifene, ABI-007,

Evista, Atamestane, Xyotax, Targretin, Triazone, Aposyn, Nevastat, Ceplene, Orathecin, Virulizin, Gastrimmune, DX-8951f, Onconase, BEC2, Xcytrin, CeaVac, NewTrexin, OvaRex, Osidem, Advexin, RSR13 (efaproxiral, Cotara, NBI-3001 (IL-4), Canvaxin, GMK vaccine, PEG Interferon A, Taxoprexin, gene therapy agents such as TNFerade (GeneVac),
5 Interferon-alpha, Interferon-gamma, Tumor necrosis factor, Lovastatin, staurosporine, dactinomycin, zorubicin, Bosentan, ampligen, ibandronic acid, miltefosine, L-asparaginase, procarbazine, hydroxycarbamide, pegaspargase, pentostatin, tazarotne, Telcyta, tretinoin, acitretin, zolendronic acid, halofuginone, rebimastat, removab, squalamine, ukrain, paditaxel, Zinecard, Vitaxin, anthracyclines, antifolates, antiestrogen agents, antimicrotubule agents,
10 anti-androgens, aromatase inhibitors, Ca²⁺ adenosine triphosphate (ATP)ase inhibitors, cytosine analogs, deldihydrofolate reductase inhibitors, deoxyribonucleic acid (DNA) topoisomerase inhibitors, (HSP)-90 inhibitors, immunotherapeutic agents, inosine monophosphate (IMP) dehydrogenase inhibitors, isoprenylation inhibitors, luteinizing hormone-releasing hormone agonists, mammalian target of rapomycin (mtor) inhibitors,
15 multi-drug resistance (MDR) inhibitors, mytomyins, photodynamic therapies, ribonucleotide reductase inhibitors, thrombospondin mimetics, vinca alkaloids, vitamin D3 analogs, 17-allylamino-17-demethoxygeldanamycin, N-(4-(3-amino-1H-indazol-4-yl)phenyl)-N'-(2-fluoro-5-methylphenyl)urea or a salt thereof, N-(4-(4-aminothieno[2,3-d]pyrimidin-5-yl)phenyl)-N'-(2-fluoro-5-(trifluoromethyl)phenyl)urea or a salt thereof, campathecins,
20 CB1093, CHIR258, CNF-101, CNF-1001, CP547632, demethoxyhypocrellin A, 17-dimethylaminoethylamino-17-demethoxygeldanamycin, EB1089, eothilone D, epirubicin, 5-ethynyl-1-β-D-ribofuranosylimidazole-4-carboxamide (EICAR), erlotinib, N-(2-(4-hydroxyanilino)-3-pyridinyl)-4-methoxybenzenesulfonamide or a salt thereof, imatinab, IPI-504, KH 1060, LAQ824, lomustine, 1-methyl-4-phenylpyridinium, MLN-518,
25 nitrosoureas, photosensitizer Pc4, phtalocyanine, plicamycin, retinoids such as pheuretinide, sunitinib, taxol, teniposide, thapsigargin, trichostatin A, verapamil, vertoporfin, ZK-EP, Polo-like kinase inhibitors, proteasome inhibitors or combinations thereof.

BAX and BAD peptides are reported in Zhang, H. C., Nimmer, P., Rosenberg, S. H., Ng, S. C., and Joseph, M. (2002). Development of a High-Throughput Fluorescence
30 Polarization Assay for Bcl-x(L). Analytical Biochemistry 307, 70-75.

Binding affinity of compounds having formula (II) to Bcl-X_L protein is indicia of their inhibition of the activity of this protein. To determine the binding affinity of compounds having formula (II) to Bcl-X_L protein, representative examples were diluted in DMSO to concentrations between 100 μM and 1 pM and added to each well of a 96-well microtiter plate. A mixture
35 comprising 125 μL per well of assay buffer (20 mM phosphate buffer (pH 7.4), 1 mM EDTA, 50

mM NaCl, 0.05% PF-68), 6 nM of Bcl-X_L protein (prepared as described in Science 1997, 275, 983-986), 1 nM fluorescein-labeled BAD peptide (prepared in-house) and the DMSO solution of the compound was shaken for 2 minutes and placed in a LJL Analyst (LJL Bio Systems, CA). A negative control (DMSO, 15 nM BAD peptide, assay buffer) and a positive control (DMSO, 1 nM BAD peptide, 6 nM Bcl-X_L, assay buffer) were used to determine the range of the assay. Polarization was measured at room temperature using a continuous Fluorescein lamp (excitation 485 nm, emission 530 nm). Percentage of inhibition was determined by $(1 - ((\text{mP value of well-negative control}) / \text{range})) \times 100\%$. The results are shown in TABLE 1.

Binding affinity of compounds having formula (II) to Bcl-2 protein is indicia of their inhibition of the activity of this protein. To determine the binding affinity of compounds having formula (II) to Bcl-2, representative examples were diluted in DMSO to concentrations between 10 μM and 10 pM and added to each well of a 96-well microtiter plate. A mixture comprising 125 μL per well of assay buffer (20 mM phosphate buffer (pH 7.4), 1 mM EDTA, 50 mM NaCl, 0.05% PF-68), 10 nM of Bcl-2 protein (prepared according to the procedure described in PNAS 2001, 98, 3012 - 3017), 1 nM fluorescein-labeled BAX peptide (prepared in-house) and the DMSO solution of the representative EXAMPLE was shaken for 2 minutes and placed in a LJL Analyst (LJL Bio Systems, CA). Polarization was measured at room temperature using a continuous Fluorescein lamp (excitation 485 nm, emission 530 nm). The results are also shown in TABLE 1.

These data demonstrate the utility of compounds having formula (II) as binders to and inhibitors of anti-apoptotic Bcl-X_L protein and anti-apoptotic Bcl-2.

It is expected that, because compounds having formula (II) bind to and inhibit the activity of Bcl-X_L and Bcl-2, they would also have utility as inhibitors of anti-apoptotic family protein members having close structural homology to Bcl-X_L and Bcl-2 such as, for example, anti-apoptotic Bcl-w protein.

Accordingly, compounds having formula (II) are expected to have utility in treatment of diseases during which anti-apoptotic Bcl-X_L protein, anti-apoptotic Bcl-2 protein, anti-apoptotic Bcl-w protein or a combination thereof, are expressed.

Determination of Cellular Efficacy in Human Tumor Cell Line

NCI-H146 (ATCC, Manassas, VA.) human small cell lung carcinoma cells were plated 50,000 cells per well in 96-well tissue culture plates in a total volume of 100 μL tissue culture medium supplemented with 10% human serum (Invitrogen, Carlsbad, CA.) instead of fetal bovine serum and treated with a 2-fold serial dilution of the compounds of interest from 10 μM to 0.020 μM . Each concentration was tested in duplicate at least 3 separate times. The number of viable cells following 48 hours of compound treatment was determined using the

CellTiter 96® AQueous non-radioactive cell proliferation MTS assay according to manufacturer's recommendations (Promega Corp., Madison, WI). The results are also shown in TABLE 1.

Pharmacokinetic Evaluation of Selected Compounds in Rat

5 The pharmacokinetic behavior of compounds of this invention was determined following a single 2 mg/kg intravenous or 5 mg/kg oral dose in male Sprague-Dawley derived rats (n=3 per group). The compounds were prepared as 2 mg/mL solution in a 10% DMSO in PEG-400 formulation for both oral and intravenous administration. The 1 mL/kg intravenous dose was administered as a slow bolus (about 1-2 minutes) in the jugular vein of a rat under light ether anesthetic. The oral dose was administered by gavage. Serial blood samples were obtained from a tail vein of each rat prior 0.1 (IV only), 0.25, 0.5, 1, 1.5, 2, 3, 4, 6, 8 and 24 hours after dosing. The heparinized samples were thoroughly mixed and placed in an ice bath. Plasma was separated by centrifugation and stored frozen prior to analysis. The results are also shown in TABLE 1.

15 The compounds of interest were separated from the plasma using protein precipitation with acetonitrile. A plasma (100-200 µL, sample or spiked standard) aliquot was combined with 50 µL of internal standard (structurally related analog prepared in acetonitrile) and 1 ml acetonitrile in a 96-well polypropylene deep well plate. The plates were vortexed for 30 seconds followed by centrifugation (3500 rpm x 15 minutes, 4°C). In an automated manner, the supernatant was transferred to a clean 96-well plate. The samples were evaporated to near dryness on a Micro-Vap™ under a stream of dry nitrogen over low heat (~37°C). The samples were reconstituted vortexing with 0.2 mL 5% DMSO in acetonitrile. A 0.1-0.2 ml aliquot of acetonitrile: 0.1% trifluoroacetic acid (20:80, by volume) was added to each well, followed by an additional 30 second vortexing. The plates were centrifuged (3500 rpm x 15 minutes, 4°C) prior to HPLC-MS/MS analysis. Samples were analyzed simultaneously with spiked plasma standards. All samples from each study were analyzed as a single batch on the LC-MS/MS.

20 The compounds of interest and the internal standard were separated from each other and co-extracted contaminants on a 50 x 3 mm Keystone Betasil CN 5 µm column with an acetonitrile: 0.1% trifluoroacetic acid mobile phase (50:50, by volume) at a flow rate of 0.7 ml/min. Analysis was performed on a Sciex API 300™ Biomolecular Mass Analyzer using a heated nebulizer interface. Peak areas of the title compounds and internal standards were determined using the Sciex MacQuan™ software. Calibration curves were derived from peak area ratio (parent drug/internal standard) of the spiked rat plasma standards using least squares linear regression of the ratio versus the theoretical concentration. The methods were

generally linear over the range of the standard curve (correlation coefficients > 0.99) with an estimated quantitation limit of 0.01 $\mu\text{g/mL}$. The plasma concentration data for each animal were submitted to multi-exponential curve fitting using WinNonlin. The area under the plasma concentration-time curve from 0 to t hours (time of the last measurable plasma concentration) after dosing (AUC_{0-t}) was calculated using the linear trapezoidal rule for the plasma concentration-time profiles. The residual area extrapolated to infinity, determined as the final measured plasma concentration (C_t) divided by the terminal elimination rate constant (β), was added to AUC_{0-t} to produce the total area under the curve. The results are also shown in TABLE 1.

10

TABLE 1

Ex.	Ki (Bcl-2)	Ki (Bcl-X _L)	EC ₅₀	AUC _{0-∞}	AUC/EC ₅₀
1	≤ 0.001 μM	≤ 0.001 μM	0.0891 μM	4.88 μM	56.2
2	≤ 0.001 μM	≤ 0.001 μM	0.2907 μM	8.54 μM	241
3	≤ 0.001 μM	≤ 0.001 μM	0.02876 μM	4.92 μM	143.6
4	≤ 0.001 μM	≤ 0.001 μM	0.05868 μM	6.09 μM	108
5	≤ 0.001 μM	≤ 0.001 μM	0.03884 μM	2.11 μM	57.4
6	≤ 0.001 μM	≤ 0.001 μM	0.00916 μM	0.89 μM	91.6
7	≤ 0.001 μM	≤ 0.001 μM	0.0589 μM	3.66 μM	65.6
8	≤ 0.001 μM	≤ 0.001 μM	0.02123 μM	1.05 μM	51.7
9	≤ 0.001 μM	≤ 0.001 μM	0.1366 μM	1.80 μM	137.5
10	≤ 0.001 μM	≤ 0.001 μM	0.03421 μM	2.45 μM	72.6
11	≤ 0.002 μM	≤ 0.003 μM	0.02055 μM	2.41 μM	116.9
12	≤ 0.001 μM	≤ 0.001 μM	0.02707 μM	2.01 μM	76.3
13	≤ 0.001 μM	≤ 0.001 μM	0.01900 μM	2.04 μM	108.6
14	≤ 0.001 μM	≤ 0.001 μM	0.03089 μM	4.06 μM	110.6
15	≤ 0.001 μM	≤ 0.001 μM	0.00985 μM	1.49 μM	127.3
16	≤ 0.002 μM	≤ 0.002 μM	0.37390 μM	2.81 μM	63.3
17	≤ 0.001 μM	≤ 0.001 μM	0.02872 μM	1.01 μM	30.7
18	≤ 0.001 μM	≤ 0.001 μM	0.01582 μM	0.7 μM	39.5
19	≤ 0.001 μM	≤ 0.001 μM	0.1582 μM	7.06 μM	450.3
20	≤ 0.001 μM	≤ 0.001 μM	0.02766 μM	2.9 μM	112.4
21	≤ 0.001 μM	≤ 0.001 μM	0.06425 μM	2.13 μM	28.2
22	≤ 0.001 μM	≤ 0.001 μM	0.11922 μM	4.92 μM	32.4
23	≤ 0.001 μM	≤ 0.001 μM	0.05281 μM	4.27 μM	67.1

24	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.04429 μM	9.67 μM	181.5
25	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.01640 μM	2.06 μM	102
26	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.02434 μM	0.91 μM	33
27	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.01854 μM	1.94 μM	112.4
28	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.02420 μM	7.42 μM	260.5
29	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.02982 μM	1.64 μM	58.2
30	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.03165 μM	3.31 μM	107.3
31	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.13040 μM	4.97 μM	390.9
32	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.01872 μM	1.33 μM	73.9
33	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.03778 μM	2.81 μM	79.8
34	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.20030 μM	10.6 μM	552.6
35	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.00764 μM	1.19 μM	164.5
36	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.07094 μM	8.47 μM	102.9
37	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.2707 μM	5.12 μM	155.7
38	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.03900 μM	0.67 μM	17.05
39	$\leq 0.001 \mu\text{M}$	$\leq 0.001 \mu\text{M}$	0.083 μM	1.66 μM	19.96

The compounds of the present invention were also tested against compounds disclosed in WO 2005/049594, identified herein as EXAMPLES A-N, by determining the ratio of potency to exposure. This measure, sometimes reported as EC_{50}/AUC , is well known to those skilled in the art of pharmaceutical drug discovery and drug development as a useful measurement of pharmacodynamic activity.

The examples of the present invention and compounds disclosed in WO 2005/049594 were tested in both H146 cell assay and for pharmacokinetic evaluation in rat, both as previously described herein. The results are shown in TABLES 2 and 3. As can be seen with reference to the data, the compounds of the present invention have a more preferred pharmacodynamic profile as compared to the compounds known in the art. From these results a number of observations can be drawn. It can be observed that the compounds having a NO_2 moiety at position W^1 tend to have good to excellent cellular potency. However, when the oral bioavailability of these same compounds is determined, it can be seen that the exposure is poor, resulting in EC_{50}/AUC ratios of from 0.5 to 19.7. On the other hand, when compounds having a CF_3 moiety at position W^1 are tested in the cellular assay, the results demonstrate that these derivatives have relatively poor potency while at the same time having suitable oral exposure. Again, this combination provides overall ratios from about 2.8 to about < 7.4 . Surprisingly, compounds of the present invention demonstrate cellular potency on par with compounds having an NO_2 moiety while maintaining the oral bioavailability of

compounds having the CF₃ moiety. The resulting ratios for the compounds of the invention are from about 20 to about 550.







TABLE 2

EXAMPLE	W ¹	W ²	W ³	H146, EC ₅₀ , (μM)	C _{max} , (μM)	AUC, (μM)	AUC/EC ₅₀
38	SO ₂ CF ₃	N(CH ₃) ₂	H	0.039	0.072	0.665	16.9
A	CF ₃	N(CH ₃) ₂	H	1.599	0.371	4.412	2.8
B	NO ₂	N(CH ₃) ₂	H	0.063	0.039	0.283	4.5
C	CO ₂ CF ₃	N(CH ₃) ₂	H	2.011	0.094	0.989	0.5
D	CN	N(CH ₃) ₂	H	1.807	0.315	1.917	1.1
E	CF ₃	N(CH ₃) ₂	F	7.329	0.386	3.827	0.5
39	SO ₂ CF ₃		H	0.083	0.290	1.657	20.0
F	NO ₂		H	0.974	0.195	1.157	1.2
G	CF ₃		H	> 1.00	0.592	7.365	<7.4












TABLE 3

EXAMPLE	W ¹	W ²	H146 EC ₅₀ , (μM)	C _{max} , (μM)	AUC, (μM)	AUC/EC ₅₀
18	SO ₂ CF ₂ Cl	N(CH ₃) ₂	0.015	0.059	0.609	39.5
26	SO ₂ CF ₃	N(CH ₃) ₂	0.024	0.097	0.803	33.0
H	NO ₂	N(CH ₃) ₂	0.026	0.057	0.507	19.7
I	CF ₃	N(CH ₃) ₂	0.410	0.215	1.973	4.8
3	SO ₂ CF ₂ Cl		0.029	0.385	4.131	143.6
7	SO ₂ CF ₃		0.059	0.518	3.867	65.6
K	NO ₂		0.094	0.267	1.977	21.0
6	SO ₂ CF ₂ Cl		0.010	0.113	0.913	91.6
9	SO ₂ CF ₃		0.014	0.156	1.878	137.5
L	NO ₂		0.028	0.047	0.402	14.3
8	SO ₂ CF ₃		0.021	0.071	1.098	51.7
M	NO ₂		0.049	0.077	0.368	7.5
33	SO ₂ CF ₃	N(<i>i</i> -Pr)CH ₃	0.038	0.129	3.013	79.8
N	NO ₂	N(<i>i</i> -Pr)CH ₃	0.034	0.0367	0.615	18.2

i-Pr means iso-propyl

As shown in Figures 1-7, studies pertaining to the efficacy of EXAMPLE 1 in combination with etoposide, vincristine, CHOP, rituximab, rituximab with CHOP, rapamycin, and velcade demonstrated that EXAMPLE 1 synergistically enhanced efficacy of these cytotoxic agents during combination therapy.

Further, combinations comprising EXAMPLE 1 and vincristine resulted in 10% complete tumor regression.

Still further, combinations comprising EXAMPLE 1 and rituximab resulted in 70% complete tumor regression whereas no tumor regressions were observed for rituximab alone.

Still further, combinations comprising EXAMPLE 1 and rapamycin resulted in 70% complete tumor regression whereas 10% tumor regressions were observed for rapamycin alone.

Still further, combinations comprising EXAMPLE 1 and rituximab with CHOP resulted in 90% complete tumor regression whereas 10% tumor regressions were observed for rituximab with CHOP only.

Still further, combinations comprising EXAMPLE 1 and bortezomib resulted in 10% complete tumor regression whereas no tumor regressions were observed for bortezomib alone.

Diseases during which anti-apoptotic Bcl-X_L protein, anti-apoptotic Bcl-2 protein, anti-apoptotic Bcl-w protein or a combination thereof, are expressed include, but are not limited to, cancer and autoimmune disorders, wherein cancer includes, but is not limited to, acoustic neuroma, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia (monocytic, myeloblastic, adenocarcinoma, angiosarcoma, astrocytoma, myelomonocytic and promyelocytic), acute t-cell leukemia, basal cell carcinoma, bile duct carcinoma, bladder cancer, brain cancer, breast cancer, bronchogenic carcinoma, cervical cancer, chondrosarcoma, chordoma, choriocarcinoma, chronic leukemia, chronic lymphocytic leukemia, chronic myelocytic (granulocytic) leukemia, colorectal cancer, craniopharyngioma, cystadenocarcinoma, diffuse large B-cell lymphoma, dysproliferative changes (dysplasias and metaplasias), embryonal carcinoma, endometrial cancer, endotheliosarcoma, ependymoma, epithelial carcinoma, erythroleukemia, esophageal cancer, estrogen-receptor positive breast cancer, essential thrombocythemia, Ewing's tumor, fibrosarcoma, follicular lymphoma, germ cell testicular cancer, glioma, heavy chain disease, hemangioblastoma, hepatoma, hepatocellular cancer, hormone insensitive prostate cancer, leiomyosarcoma, liposarcoma, lung carcinoma, lymphagioendotheliosarcoma, lymphangiosarcoma, lymphoblastic leukemia, lymphoma (Hodgkin's and non-Hodgkin's), malignancies and hyperproliferative disorders of the bladder, breast, colon, lung, ovaries, pancreas, prostate, skin and uterus, lymphoid malignancies of T-cell or B-cell origin, leukemia, lymphoma, medullary carcinoma, medulloblastoma, melanoma, meningioma, mesothelioma, multiple myeloma, myelogenous leukemia, myeloma, myxosarcoma, neuroblastoma, non-small cell lung cancer, oligodendroglioma, oral cancer, osteogenic sarcoma, ovarian cancer, pancreatic cancer, papillary adenocarcinomas, papillary carcinoma, pinealoma, polycythemia vera, prostate cancer, renal cell carcinoma, retinoblastoma, rhabdomyosarcoma, sarcoma, sebaceous gland carcinoma, seminoma, small cell lung carcinoma, solid tumors (carcinomas and sarcomas), small cell lung cancer squamous cell carcinoma, synovioma, sweat gland carcinoma, Waldenström's macroglobulinemia, testicular tumors, uterine cancer and Wilms' tumor, (Cancer Res., 2000, 60, 6101-10 and Medicine, 2d Ed., J.B. Lippincott Co., Philadelphia (1985)); autoimmune disorders include, but are not limited to, acquired immunodeficiency

disease syndrome, autoimmune lymphoproliferative syndrome, hemolytic anemia, inflammatory diseases, and thrombocytopenia (Current Allergy and Asthma Reports 2003, 3:378-384; Br. J. Haematol. 2000 Sep; 110(3): 584-90; Blood 2000 Feb 15;95(4):1283-92; and New England Journal of Medicine 2004 Sep; 351(14): 1409-1418).

5 It is also expected that compounds having formula (II) would inhibit the growth of cells derived from a cancer or neoplasm such as breast cancer (including estrogen-receptor positive breast cancer), colorectal cancer, endometrial cancer, lung cancer (including small cell lung cancer), lymphoma (including follicular or Diffuse Large B-cell), lymphoma (including non-Hodgkin's lymphoma), neuroblastoma, ovarian cancer, prostate cancer
10 (including hormone-insensitive prostate cancer), testicular cancer (including germ cell testicular cancer).

It is also expected that compounds having formula (II) would inhibit the growth of cells derived from a pediatric cancer or neoplasm such as embryonal rhabdomyosarcoma, pediatric acute lymphoblastic leukemia, pediatric acute myelogenous leukemia, pediatric
15 alveolar rhabdomyosarcoma, pediatric anaplastic ependymoma, pediatric anaplastic large cell lymphoma, pediatric anaplastic medulloblastoma, pediatric atypical teratoid/rhabdoid tumor of the central nervous system, pediatric biphenotypic acute leukemia, pediatric Burkitts lymphoma, pediatric cancers of Ewing's family of tumors such as primitive neuroectodermal tumors, pediatric diffuse anaplastic Wilm's tumor, pediatric favorable histology Wilm's
20 tumor, pediatric glioblastoma, pediatric medulloblastoma, pediatric neuroblastoma, pediatric neuroblastoma-derived myelocytomatosis, pediatric pre-B-cell cancers (such as leukemia), pediatric osteosarcoma, pediatric rhabdoid kidney tumor, pediatric rhabdomyosarcoma, and pediatric T-cell cancers such as lymphoma and skin cancer (commonly-owned United States Application Ser No. 10/988,338), Cancer Res., 2000, 60, 6101-10); autoimmune disorders
25 include, but are not limited to, acquired immunodeficiency disease syndrome, autoimmune lymphoproliferative syndrome, hemolytic anemia, inflammatory diseases, and thrombocytopenia (Current Allergy and Asthma Reports 2003, 3:378-384; Br. J. Haematol. 2000 Sep; 110(3): 584-90; Blood 2000 Feb 15;95(4):1283-92; and New England Journal of Medicine 2004 Sep; 351(14): 1409-1418).

30 Compounds having formula (II) may be made by synthetic chemical processes, examples of which are shown hereinbelow. It is meant to be understood that the order of the steps in the processes may be varied, that reagents, solvents and reaction conditions may be substituted for those specifically mentioned, and that vulnerable moieties may be protected and deprotected, as necessary.

Protecting groups for C(O)OH moieties include, but are not limited to, acetoxymethyl, allyl, benzoylmethyl, benzyl, benzyloxymethyl, tert-butyl, tert-butyldiphenylsilyl, diphenylmethyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclopropyl, diphenylmethylsilyl, ethyl, para-methoxybenzyl, methoxymethyl, methoxyethoxymethyl, methyl, methylthiomethyl, naphthyl, para-nitrobenzyl, phenyl, n-propyl, 2,2,2-trichloroethyl, triethylsilyl, 2-(trimethylsilyl)ethyl, 2-(trimethylsilyl)ethoxymethyl, triphenylmethyl and the like.

Protecting groups for C(O) and C(O)H moieties include, but are not limited to, 1,3-dioxylketal, diethylketal, dimethylketal, 1,3-dithianylketal, O-methyloxime, O-phenyloxime and the like.

Protecting groups for NH moieties include, but are not limited to, acetyl, alanyl, benzoyl, benzyl (phenylmethyl), benzylidene, benzyloxycarbonyl (Cbz), tert-butoxycarbonyl (Boc), 3,4-dimethoxybenzyloxycarbonyl, diphenylmethyl, diphenylphosphoryl, formyl, methanesulfonyl, para-methoxybenzyloxycarbonyl, phenylacetyl, phthaloyl, succinyl, trichloroethoxycarbonyl, triethylsilyl, trifluoroacetyl, trimethylsilyl, triphenylmethyl, triphenylsilyl, para-toluenesulfonyl and the like.

Protecting groups for OH and SH moieties include, but are not limited to, acetyl, allyl, allyloxycarbonyl, benzyloxycarbonyl (Cbz), benzoyl, benzyl, tert-butyl, tert-butyldimethylsilyl, tert-butyldiphenylsilyl, 3,4-dimethoxybenzyl, 3,4-dimethoxybenzyloxycarbonyl, 1,1-dimethyl-2-propenyl, diphenylmethyl, formyl, methanesulfonyl, methoxyacetyl, 4-methoxybenzyloxycarbonyl, para-methoxybenzyl, methoxycarbonyl, methyl, para-toluenesulfonyl, 2,2,2-trichloroethoxycarbonyl, 2,2,2-trichloroethyl, triethylsilyl, trifluoroacetyl, 2-(trimethylsilyl)ethoxycarbonyl, 2-trimethylsilylethyl, triphenylmethyl, 2-(triphenylphosphonio)ethoxycarbonyl and the like.


The following abbreviations have the meanings indicated.

ADDP means 1,1'-(azodicarbonyl)dipiperidine; AD-mix- β means a mixture of (DHQD)₂PHAL, K₃Fe(CN)₆, K₂CO₃ and K₂SO₄; AIBN means 2,2'-azobis(2-methylpropionitrile); 9-BBN means 9-borabicyclo[3.3.1]nonane; (DHQD)₂PHAL means hydroquinidine 1,4-phthalazinediyl diethyl ether; DBU means 1,8-diazabicyclo[5.4.0]undec-7-ene; DIBAL means diisobutylaluminum hydride; DIEA means diisopropylethylamine; DMAP means N,N-dimethylaminopyridine; DME means 1,2-dimethoxyethane; DMF means N,N-dimethylformamide; dmpe means 1,2-bis(dimethylphosphino)ethane; DMSO means dimethylsulfoxide; dppb means 1,4-bis(diphenylphosphino)butane; dppe means 1,2-bis(diphenylphosphino)ethane; dppf means 1,1'-bis(diphenylphosphino)ferrocene; dppm means 1,1-bis(diphenylphosphino)methane; EDAC means 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide; Fmoc means fluorenylmethoxycarbonyl; HATU means O-(7-

azabenzotriazol-1-yl)-N,N'N'N'-tetramethyluronium hexafluorophosphate; HMPA means hexamethylphosphoramide; IPA means isopropyl alcohol; LDA means lithium diisopropylamide; LHMDS means lithium bis(hexamethyldisilylamide); MP-BH₃ means macroporus triethylammonium methylpolystyrene cyanoborohydride; LAH means lithium aluminum hydride; NCS means N-chlorosuccinimide; PyBOP means benzotriazol-1-yloxytripyrrolidinophosphonium hexafluorophosphate; TDA-1 means tris(2-(2-methoxyethoxy)ethyl)amine;


TEA means triethylamine; TFA means trifluoroacetic acid; THF means tetrahydrofuran; NCS means N-chlorosuccinimide; NMM means N-methylmorpholine; NMP means N-methylpyrrolidine; PPh₃ means triphenylphosphine.

SCHEME 1



As shown in SCHEME 1, compounds having formula (1) may be converted to compounds having formula (2) by reacting the former, chlorosulfonic acid, and ammonia.

SCHEME 2



Compounds having formula (2) may be converted to compounds having formula (II) by reacting the former and compounds having formula (3) and a coupling agent, with or without a base. Examples of coupling agents include EDCI, CDI, and PyBop. Examples of bases include TEA, DIEA, DMAP, and mixtures thereof.

Compounds having formula (2) may be converted to compounds having formula (II) by reacting the former and compounds having formula Z^1 -COCl and the base.

The following examples are presented to provide what is believed to be the most useful and readily understood description of procedures and conceptual aspects of this invention.

EXAMPLE 1A

3-(R)-((Carbobenzyloxy)amino)- γ -butyrolactone, prepared as described in J. Am. Chem. Soc. 1986, 108, 4943-4952, (62 g) and morpholine (46 mL) in dioxane (700 mL) at 65°C was stirred for 24 hours, cooled and concentrated. The concentrate was chromatographed on silica gel with 10% methanol/ethyl acetate.

EXAMPLE 1B

EXAMPLE 1A (16.5 g), diphenyl disulfide (14.5 g) and tributylphosphine (16.6 mL) in toluene (250 mL) at 80°C was stirred for 24 hours, cooled and concentrated. The concentrate was chromatographed on silica gel with 1:1 ethyl acetate/hexanes.

EXAMPLE 1C

EXAMPLE 1B (18 g) in 30% HBr in acetic acid (250 mL) at 25°C was stirred for 24 hours, concentrated, poured into 1M HCl and extracted with diethyl ether. The extract was extracted with 1M HCl, and this extract was cooled to 0°C, adjusted to pH 12 with KOH and extracted with dichloromethane. The extract was washed with brine and dried (Na_2SO_4), filtered and concentrated.

EXAMPLE 1D

EXAMPLE 1C (45.4 g) in THF (500 mL) at 55°C was treated with 1M $\text{BH}_3 \cdot \text{THF}$ (650 mL) over 2 hours, stirred for 24 hours, cooled to 0°C, treated with methanol (80 mL), poured into methanol (500 mL) and concentrated. A mixture of the concentrate in methanol (400 mL) was treated with a HCl-saturated methanol (800 mL), refluxed for 24 hours, cooled, concentrated, poured into 2M NaOH and extracted with ethyl acetate. The extract was washed with 1M NaOH and brine and dried (Na_2SO_4), filtered and concentrated. The concentrate was chromatographed on silica gel with ethyl acetate 10% methanol/ethyl acetate and 10% methanol/10% acetonitrile/5% TEA/75% ethyl acetate.

EXAMPLE 1E

Methyl viologen hydrochloride (1.17 g) in DMF (80 mL) at 25°C was saturated with trifluoromethyl iodide, treated with 2-fluorobenzenethiol (9.7 mL) and TEA (20 mL), stirred for 24 hours, diluted with water (240 mL) and extracted with diethyl ether. The extract was washed with 1M NaOH, saturated ammonium chloride and brine and concentrated.

5

EXAMPLE 1F

EXAMPLE 1E (17.346 g) in 1:1:2 carbon tetrachloride/acetonitrile/water (800 mL) at 25°C was treated with sodium periodate (56.8 g) and ruthenium(III) chloride hydrate (183 mg), stirred for 18 hours, diluted with dichloromethane (100 mL) and filtered through diatomaceous earth (Celite®). The filtrate was washed with saturated sodium bicarbonate and extracted with dichloromethane. The extract was washed with brine and dried (MgSO₄), filtered and concentrated. The concentrate was filtered through silica gel.

10

EXAMPLE 1G

EXAMPLE 1F (37.3 g) in chlorosulfonic acid (32.8 mL) at 120°C was stirred for 18 hours, cooled to 25°C and pipetted onto crushed ice. The mixture was extracted with ethyl acetate, and the extract was washed with water and brine and dried (MgSO₄), filtered and concentrated.

15

EXAMPLE 1H

EXAMPLE 1G (23 g) in isopropanol (706 mL) at -78°C was treated with ammonium hydroxide (98 mL) over 1 hour, stirred for 1 hour, quenched with 6M HCl (353 mL), warmed to 25°C and concentrated. The concentrate was mixed with water and extracted with ethyl acetate. The extract was dried (MgSO₄), filtered and concentrated. The concentrate was recrystallized from ethyl acetate/hexane.

20

EXAMPLE 1I

EXAMPLE 1H (13.48 g) and EXAMPLE 1D (11.56 g) in THF (218 mL) was treated with DIEA (15.1 mL), stirred at 50°C for 4 hours, cooled, treated with saturated sodium bicarbonate and extracted with ethyl acetate. The extract was dried (MgSO₄), filtered and concentrated. The concentrate was recrystallized from hexanes/ethyl acetate.

30

EXAMPLE 1J

DMF (10 mL) and chloroform (80 mL) at 3°C was treated with PBr₃ (12 mL), stirred for 20 minutes at 25°C, treated with 4,4-dimethylcyclohexanone (7.15 g) in chloroform

35

(50 mL), stirred for 18 hours, poured onto ice, neutralized with solid sodium bicarbonate and extracted with diethyl ether. The extract was washed with brine and dried (MgSO_4), filtered and concentrated. The concentrate was chromatographed on silica gel with 0-10% ethyl acetate/hexanes.

5

EXAMPLE 1K

EXAMPLE 1J (1.7 g) and 4-piperazin-1-ylbenzoic acid ethyl ester (1.9 g) in methanol (30 mL) was treated with sodium cyanoborohydride (0.6 g), adjusted to pH 5 with acetic acid, stirred for 18 hours and filtered through diatomaceous earth (Celite®). The filtrate was concentrated, and the concentrate was chromatographed on silica gel on silica gel with 10-30% ethyl acetate/hexanes.

10

EXAMPLE 1L

EXAMPLE 1K (1.1 g), 4-chlorophenylboronic acid (0.6 g), 2M Na_2CO_3 (2 mL) and $\text{PdCl}_2(\text{PPh}_3)_2$ (0.1 g) in 7:3:2 DME/water/ethanol (20 mL) was stirred at 85°C for 18 hours, filtered through diatomaceous earth (Celite®) and concentrated. The concentrate was chromatographed on silica gel with 10-30% ethyl acetate/hexanes.

15

EXAMPLE 1M

EXAMPLE 1L (4.59 g) and LiOH (1.25 g) in dioxane (75 mL) and water (10 mL) was stirred at 100°C for 18 hours, cooled to 25°C and concentrated. The concentrate was dissolved in water, heated to reflux, neutralized with 1M HCl (28.5 mL), cooled to 25°C, filtered and concentrated.

20

EXAMPLE 1N

EXAMPLE 1M (31.5 g), EXAMPLE 1I (39.93 g), EDAC·HCl (20.60 g) and DMAP (13.15 g) in dichloromethane (500 mL) at 25°C was stirred for 18 hours, diluted with dichloromethane, washed with saturated ammonium chloride and brine and dried (MgSO_4), filtered and concentrated. The concentrate was chromatographed on silica gel with 0-10% methanol/ammonia-saturated dichloromethane. ^1H NMR (300MHz, DMSO-d_6) δ 8.12 (d, 1H), 7.94 (dd, 1H), 7.71 (d, 2H), 7.38 (d, 2H), 7.30 (m, 4H), 7.18 (m, 1H), 7.12 (d, 2H), 6.98 (d, 1H), 6.85 (d, 3H), 4.07 (m, 1H), 3.53 (br, 4H), 3.28 (m, 12H), 2.44 (m, 8H), 1.99 (m, 3H), 1.80 (m, 1H), 1.44 (t, 2H), 0.97 (s, 6H).

25

30

35

EXAMPLE 2A

Powdered NaOH (31.2 g), TDA-1 (5 mL) and 2-fluorobenzene thiol (33.6 mL) in benzene (400 mL) was saturated with chlorodifluoromethane, stirred at 80°C for 30 minutes and filtered through diatomaceous earth (Celite®). The filtrate was washed with saturated NaHCO₃ and the water layer was extracted with diethyl ether. The extracts were combined and dried (MgSO₄), filtered and concentrated.

EXAMPLE 2B

EXAMPLE 2A (46 g) in 1:1:2 CCl₄/CH₃CN/water (1.2L) at 25°C was treated with NaIO₄ (165.6 g) and RuCl₃·xH₂O (534 mg), stirred for 18 hours, diluted with dichloromethane and filtered through diatomaceous earth (Celite®). The filtrate was washed with saturated NaHCO₃ and dried (Na₂SO₄), filtered and concentrated. The concentrate was filtered through silica gel.

EXAMPLE 2C

EXAMPLE 2B (25 g) and NCS (17.55 g) in THF (700 mL) at -78 °C was treated with LHMDS (178.5 mL) over 1 hour, stirred for 1 hour and quenched with ammonium chloride. The mixture was extracted with ethyl acetate, and the extract was washed with brine and dried (MgSO₄), filtered and concentrated. The concentrate was chromatographed on silica gel with 0-5% ethyl acetate/hexanes.

EXAMPLE 2D

EXAMPLE 2C (44 g) in chlorosulfonic acid (36.7 mL) at 120°C was stirred for 18 hours, cooled to 25°C, pipetted onto crushed ice and extracted with ethyl acetate. The extract was washed with water and brine and dried (MgSO₄), filtered and concentrated.

EXAMPLE 2E

EXAMPLE 2D (22 g) in isopropanol (700 mL) at -78 °C was treated with aqueous ammonia (90 mL) over 1 hour, stirred for another hour, quenched with 6M HCl (300 mL), warmed to 25°C and concentrated. The concentrate was mixed with water and extracted with ethyl acetate. The extract was dried (MgSO₄), filtered and concentrated. The concentrate was recrystallized from hexanes/ethyl acetate.

EXAMPLE 2F

EXAMPLE 2E (2.89 g) and EXAMPLE 1D (2.39 g) in THF (20 mL) was treated with diisopropylethylamine (3.2 mL), stirred at 60°C for 18 hours, cooled, treated with saturated

sodium bicarbonate and extracted with ethyl acetate. The extract was dried (MgSO_4), filtered and concentrated. The concentrate was chromatographed on silica gel with 1.5-5% 7M ammonia in methanol/dichloromethane.

5 EXAMPLE 2G

Hexane-washed 60% oily NaH (17 g) in dichloromethane (300 mL) at -5°C was treated with 4,4-dimethyl-2-oxo-cyclohexanecarboxylic acid methyl ester, prepared as described in Tetrahedron (1992), 48 (21), 4459-64, (53.89 g), stirred for 30 minutes, cooled to -78°C , treated with trifluoromethanesulfonic anhydride, warmed to 25°C , stirred for 18 hours, washed with brine and dried (MgSO_4), filtered and concentrated.

EXAMPLE 2H

EXAMPLE 2G (86 g), 4-chlorophenylboronic acid (50 g), CsF (104 g) and tetrakis(triphenylphosphine)palladium(0) (2.5 g) in 2:1 DME/methanol (600 mL) at 70°C was stirred for 18 hours and concentrated. The concentrate was dissolved in diethyl ether, and the solution was dried (MgSO_4), filtered and concentrated. The concentrate was filtered through silica gel with 20% ethyl acetate/hexanes.

EXAMPLE 2I

Lithium borohydride (18 g) was treated with EXAMPLE 2H (76 g) in diethyl ether (400 mL) and methanol (23 mL), stirred at reflux for 4 hours, cooled, quenched with 1M HCl, diluted with water and extracted with diethyl ether. The extract was dried (MgSO_4), filtered and concentrated. The concentrate was chromatographed on silica gel with 0-30% ethyl acetate/hexanes.

EXAMPLE 2J

EXAMPLE 2I (17.5 g) in dichloromethane (100 mL) at 0°C was treated simultaneously with methanesulfonyl chloride (5.6 mL) and TEA (21 mL), stirred for 5 minutes, treated with 4-piperazin-1-ylbenzoic acid ethyl ester (17 g), stirred at 25°C for 18 hours, washed with ammonium chloride and dried (Na_2SO_4), filtered and concentrated. The concentrate was chromatographed on silica gel with 10% ethyl acetate/hexanes.

EXAMPLE 2K

This example was prepared by substituting EXAMPLE 2J for EXAMPLE 1L in EXAMPLE 1M.

EXAMPLE 2L

EXAMPLE 2K (16.9 g) and EXAMPLE 2F (22 g) in dichloromethane (200 mL) at 25°C was treated with EDAC·HCl (11.06 g) and DMAP (7.06 g), stirred for 18 hours, diluted
5 with dichloromethane (400 mL), washed with saturated ammonium chloride and brine and dried (MgSO₄), filtered and concentrated. The concentrate was chromatographed on silica gel with 0-10% methanol/ammonia-saturated dichloromethane. ¹H NMR (400MHz, DMSO-d₆) δ 8.07 (d, 1H), 7.90 (dd, 1H), 7.71 (d, 2H), 7.36 (m, 4H), 7.29 (m, 2H), 7.20 (m, 1H), 7.09 (d, 2H), 6.86 (d, 1H), 6.80 (d, 2H), 6.76 (d, 1H), 4.02 (m, 1H), 3.50 (m, 4H),
10 3.33 (m, 2H), 3.16 (m, 4H), 2.81 (s, 2H), 2.29 (m, 12H), 1.99 (s, 2H), 1.94 (m, 1H), 1.71 (m, 1H), 1.42 (t, 2H), 0.96 (s, 6H).

EXAMPLE 3A

This example was prepared by substituting 2-bromo-cyclohex-1-enecarbaldehyde,
15 prepared as described in Collect. Czech. Chem. Commun., 1961, 26, 3059.) for EXAMPLE 1J in EXAMPLE 1K.

EXAMPLE 3B

This example was prepared by substituting EXAMPLE 3A for EXAMPLE 1K in
20 EXAMPLE 1L.

EXAMPLE 3C

This example was prepared by substituting EXAMPLE 3B for EXAMPLE 1L in
25 EXAMPLE 1M.

EXAMPLE 3D

This example was prepared by substituting EXAMPLE 3C and EXAMPLE 2F for
EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (400MHz, DMSO-d₆) δ 8.11 (d, 1H), 7.92 (dd, 1H), 7.71 (d, 2H), 7.37 (d, 2H), 7.34 (m, 2H), 7.27 (t, 2H), 7.18 (t, 1H), 7.12 (d, 2H), 6.94 (d, 1H), 6.84 (m, 3H), 4.04 (m, 1H), 3.51 (br, 4H),
30 3.27 (br, 10H), 2.84 (br, 2H), 2.33 (br, 6H), 2.18 (br, 4H), 1.97 (m, 1H), 1.76 (m, 1H), 1.66 (s, 4H).

EXAMPLE 4A

A solution of 3-(R)-((carbobenzyloxy)amino)- γ -butyrolactone (prepared according to the procedure described in J. Am. Chem. Soc. 1986, 108, 4943-4952, 7.72 g, 32.8 mmol) in THF (100mL) was saturated with gaseous dimethylamine, stirred at room temperature for 16 hours, and concentrated. The residue was filtered through a plug of silica gel eluting with
5 50% acetone in hexanes to give the desired product.

EXAMPLE 4B

A solution of EXAMPLE 4A (8.45 g, 30.14 mmol) in toluene (15 mL) was treated with tributylphosphine (9.76 mL, 39.20 mmol) and diphenyldisulfide (7.30 g, 39.20 mmol)
10 and heated to 80 °C for 16 hours. The reaction mixture was concentrated and purified by column chromatography on silica gel eluting with a gradient of 0–50% ethyl acetate in hexanes to give the desired product.

EXAMPLE 4C

EXAMPLE 4B (7.5g) and bis(cyclopentadienyl)zirconium(IV) chloride hydride (10.31 g) in THF (100 mL) at 25°C was stirred for 20 minutes and concentrated. The concentrate was chromatographed on silica gel with with 50% ethyl acetate in hexane.
15

EXAMPLE 4D

EXAMPLE 4C (2.87 g) and N-isopropylmethylamine (1.92 g) in 1,2-dichloroethane (50 mL) at 25°C was treated with sodium triacetoxyborohydride (3 g), stirred for 2 hours, diluted with ethyl acetate, washed with 2M NaOH, water and brine and dried (Na₂SO₄), filtered and concentrated. The concentrate was chromatographed on silica gel with with
20 1% methanol/dichloromethane.

EXAMPLE 4E

This example was prepared by substituting EXAMPLE 4D for EXAMPLE 1B in EXAMPLE 1C.
25

EXAMPLE 4F

This example was prepared by substituting EXAMPLE 4E for EXAMPLE 1D in EXAMPLE 1I.
30

EXAMPLE 4G

This example was prepared by substituting EXAMPLE 4F for EXAMPLE 1I in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ 8.08 (s, 1H), 7.98 (d, 1H), 7.71 (d, 2H), 7.37 (m, 4H), 7.28 (t, 2H), 7.20 (t, 1H), 7.12 (d, 2H), 6.89 (d, 1H), 6.78 (d, 2H), 6.70 (d, 1H), 4.01 (m, 1H), 3.13 (m, 6H), 2.75 (m, 2H), 2.28 (m, 6H), 2.04 (m, 4H), 1.99 (m, 2H), 1.43 (m, 2H), 1.12 (m, 10H), 0.97 (s, 6H).

EXAMPLE 5

This example was prepared by substituting 4-(4-(2-(4-chlorophenyl)cyclo-1-enylmethyl)piperazin-1-yl)benzoic acid, prepared as described in commonly-owned U.S. Patent Application Ser. No. 10/988,338, and EXAMPLE 4F for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ 9.00 (m, 1H), 8.09 (s, 1H), 7.98 (d, 1H), 7.71 (d, 2H), 7.36 (m, 4H), 7.30 (t, 2H), 7.20 (t, 1H), 7.09 (d, 2H), 6.90 (d, 1H), 6.78 (d, 2H), 6.65 (d, 1H), 4.00 (m, 2H), 3.13 (m, 4H), 2.78 (m, 2H), 2.55 (m, 2H), 2.40 (m, 4H), 2.31 (m, 4H), 2.00 (m, 3H), 1.79 (m, 4H), 1.58 (m, 4H), 1.51 (m, 2H), 1.12 (m, 6H).

EXAMPLE 6A

This example was prepared by substituting EXAMPLE 4B for EXAMPLE 1B in EXAMPLE 1C.

EXAMPLE 6B

EXAMPLE 6A (6.13 g) in THF (200 mL) at 25°C was treated with di-tert-butylidicarbonate (7 g), stirred for 4 hours and concentrated. The concentrate was dissolved into ethyl acetate (500 mL), washed with 1M NaOH, water and brine and dried (Na₂SO₄), filtered and concentrated. The concentrate in THF (200 mL) at 25°C to was treated with 1M NaOH (200 mL), stirred for 5 hours and isolated. The water layer was extracted with ethyl acetate, and the THF and ethyl acetate extracts were combined, washed with water and brine and dried (Na₂SO₄), filtered and concentrated.

EXAMPLE 6C

This example was prepared by substituting EXAMPLE 6B for EXAMPLE 5B in EXAMPLE 4C.

EXAMPLE 6D

This example was prepared by substituting EXAMPLE 6C and 2-oxa-5-aza-bicyclo[2.2.1]heptane, prepared as described in commonly-owned U.S. Patent Application Ser. No. 10/988,338, for EXAMPLE 4C and N-isopropylmethyl amine in EXAMPLE 4D.

5
EXAMPLE 6E

EXAMPLE 6D (7.86 g) in dichloromethane (200 mL) at 25°C was treated with 2M HCl in diethyl ether (200 mL), stirred for 18 hours and concentrated.

EXAMPLE 6F

10

This example was prepared by substituting EXAMPLE 6E for EXAMPLE 1D in EXAMPLE 2F.

EXAMPLE 6G

15

This example was prepared by substituting EXAMPLE 6F and EXAMPLE 3C for EXAMPLE 1I and EXAMPLE 1M in EXAMPLE 1N. ¹H NMR (300 MHz, DMSO-d₆) δ 8.07 (s, 1H), 7.92 (d, 1H), 7.70 (d, 2H), 7.37 (m, 4H), 7.30 (t, 2H), 7.21 (t, 1H), 7.12 (d, 2H), 6.84 (d, 1H), 6.79 (d, 2H), 4.21 (m, 1H), 4.09 (m, 1H), 4.01 (m, 2H), 3.82 (m, 2H), 3.46 (m, 1H), 3.18 (m, 6H), 2.86 (m, 4H), 2.75 (m, 4H), 2.28 (m, 2H), 2.18 (m, 4H), 1.88 (m, 4H), 1.66 (m, 4H).

20
EXAMPLE 725

This example was prepared by substituting EXAMPLE 3C for EXAMPLE 1M in EXAMPLE 1N. ¹H NMR (300MHz, DMSO-d₆) δ 8.09 (d, 1H), 7.92 (dd, 1H), 7.71 (d, 2H), 7.31 (m 7H), 7.18 (tt, 1H), 7.12 (dt, 2H), 6.92 (d, 1H), 6.82 (m, 3H), 4.04 (m, 1H), 3.51 (m, 4H), 3.26 (m, 10H), 2.82 (m, 2H), 2.30 (m, 10H), 1.94 (m, 1H), 1.72 (m, 5H).

EXAMPLE 8A

30

A solution of 3-(9H-fluoren-9-ylmethoxycarbonylamino)-4-phenylsulfanylbutyric acid, prepared as described in commonly-owned U.S. Patent Application Ser. No. 10/988,338 and HATU in DMF was treated with 7-aza-bicyclo[2.2.1]heptane (prepared as described in Org. Lett., 2001, 3, 1371-1374; and N-methylmorpholine, stirred at ambient temperature for 30min, diluted with ethyl acetate, washed with 1.5% HCl, NaHCO₃(aq), H₂O and brine, dried (Na₂SO₄), filtered and concentrated to give the desired product.

35
EXAMPLE 8B

A solution of EXAMPLE 8A in THF was treated with diethyl amine, stirred at ambient temperature for 2 hours and concentrated. The residue was purified by silica gel chromatography eluting with CH₂Cl₂ (saturated with NH₃), followed by ethyl acetate to give the desired product.

5
EXAMPLE 8C

This example was prepared by substituting EXAMPLE 8B for EXAMPLE 1C in EXAMPLE 1D.

10
EXAMPLE 8D

This example was prepared by substituting EXAMPLE 8C for EXAMPLE 1D in EXAMPLE 1I.

15
EXAMPLE 8E

This example was prepared by substituting EXAMPLE 8D and EXAMPLE 3C for EXAMPLE 1I and EXAMPLE 1M, respectively, in EXAMPLE 1N. ¹H NMR (300MHz, DMSO-d₆) δ 9.19 (m, 1H), 8.07 (d, 1H), 7.97 (d, 1H), 7.71 (d, 2H), 7.31 (m 6H), 7.20 (tt, 1H), 7.12 (dt, 2H), 6.89 (d, 1H), 6.76 (d, 2H), 6.65 (d, 1H), 4.03 (m, 2H), 3.31 (m, 4H), 3.12 (m, 4H), 2.90 (br, 2H), 2.76 (m 2H), 1.96 (m, 21H).

20
EXAMPLE 9A

This example was prepared by substituting EXAMPLE 6E for EXAMPLE 1D in EXAMPLE 1I.

25
EXAMPLE 9B

This example was prepared by substituting EXAMPLE 9A and EXAMPLE 3C for EXAMPLE 1I and EXAMPLE 1M, respectively, in EXAMPLE 1N. ¹H NMR (300MHz, DMSO-d₆) δ 8.08 (d, 1H), 7.94 (d, 1H), 7.71 (d, 2H), 7.32 (m 7H), 7.20 (tt, 1H), 7.12 (dt, 2H), 6.87 (d, 1H), 6.78 (d, 3H), 4.40 (m, 1H), 4.03 (m, 2H), 3.83 (m, 2H), 3.54 (m, 2H), 3.26 (m, 2H), 3.14 (m, 4H), 2.80 (br, 2H), 2.78 (m, 4H), 1.97 (m, 14H).

30
EXAMPLE 10

This example was prepared by substituting EXAMPLE 9A for EXAMPLE 1I in EXAMPLE 1N. ¹H NMR (300MHz, DMSO-d₆) δ 8.09 (d, 1H), 7.95 (d, 1H), 7.71 (d, 2H), 7.33 (m 7H), 7.20 (tt, 1H), 7.12 (dt, 2H), 6.90 (d, 1H), 6.79 (d, 3H), 4.44 (m, 1H), 4.03

(m, 1H), 3.84 (m, 1H), 3.57 (m, 1H), 3.02 (m, 13H), 2.25 (m, 6H), 1.99 (m, 6H), 1.43 (t, 2H), 0.97 (s, 6H).

EXAMPLE 11

5 This example was prepared by substituting EXAMPLE 6F for EXAMPLE 1I in EXAMPLE 1N. ¹H NMR (300MHz, DMSO-d₆) δ 8.07 (d, 1H), 7.93 (d, 1H), 7.70 (d, 2H), 7.33 (m 7H), 7.20 (tt, 1H), 7.12 (dt, 2H), 6.81 (m, 4H), 4.41 (m, 1H), 4.06 (m, 1H), 3.83 (m, 1H), 3.47 (m, 1H), 3.02 (m, 13H), 2.25 (m, 6H), 1.99 (m, 6H), 1.43 (t, 2H), 0.97 (s, 6H).

EXAMPLE 12

10 This example was prepared by substituting 4-(4-(2-(4-chlorophenyl)cyclohept-1-enylmethyl)piperazin-1-yl)benzoic acid, prepared as described in commonly-owned U.S. Patent Application Ser. No. 10/988,338, and EXAMPLE 9A for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (300MHz, DMSO-d₆) δ 8.09 (d, 1H), 7.95 (dd, 1H), 7.71 (d, 2H), 7.32 (m 7H), 7.19 (tt, 1H), 7.09 (dt, 2H), 6.90 (d, 1H), 6.79 (d, 3H), 4.45 (m, 1H), 4.03 (m, 2H), 3.85 (m, 2H), 3.55 (m, 2H), 3.04 (m, 8H), 2.34 (m, 8H), 1.85 (m, 7H), 1.54 (m, 5H).

EXAMPLE 13

20 This example was prepared by substituting 4-(4-(2-(4-chlorophenyl)cyclohept-1-enylmethyl)piperazin-1-yl)benzoic acid, prepared as described in commonly-owned U.S. Patent Application Ser. No. 10/988,338, and EXAMPLE 6F for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (300MHz, DMSO-d₆) δ 8.07 (d, 1H), 7.93 (dd, 1H), 7.71 (d, 2H), 7.32 (m 7H), 7.20 (tt, 1H), 7.09 (dt, 2H), 6.87 (d, 1H), 6.79 (d, 3H), 4.45 (m, 1H), 4.02 (m, 2H), 3.84 (m, 2H), 3.56 (m, 2H), 3.07 (m, 8H), 2.33 (m, 8H), 1.85 (m, 7H), 1.54 (m, 5H).

EXAMPLE 14

30 This example was prepared by substituting EXAMPLE 2K and EXAMPLE 4F for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (300MHz, DMSO-d₆) δ 8.07 (d, 1H), 7.96 (dd, 1H), 7.71 (d, 2H), 7.33 (m 7H), 7.20 (tt, 1H), 7.09 (dt, 2H), 6.87 (d, 1H), 6.77 (d, 2H), 6.72 (d, 1H), 4.00 (m, 1H), 3.28 (m, 4H), 3.12 (m, 4H), 2.79 (m, 2H), 2.48 (m, 2H), 2.23 (m, 8H), 2.02 (m, 4H), 1.42 (t, 2H), 1.08 (m, 6H), 0.96 (s, 6H).

EXAMPLE 15A

This example was prepared by substituting EXAMPLE 6C and 1,4-oxazepane for EXAMPLE 4C and N-isopropyl-N-methylamine in EXAMPLE 4D.

EXAMPLE 15B

5 This example was prepared by substituting EXAMPLE 15A for EXAMPLE 6D in EXAMPLE 6E.

EXAMPLE 15C

10 This example was prepared by substituting EXAMPLE 15B for EXAMPLE 1D in EXAMPLE 1I.

EXAMPLE 15D

15 This example was prepared by substituting EXAMPLE 15C and EXAMPLE 3C for EXAMPLE 1I and EXAMPLE 1M, respectively, in EXAMPLE 1N. ¹H NMR (400MHz, CDCl₃) δ 8.32 (s, 1H), 8.01 (br, 1H), 7.67 (d, 2H), 7.34 (t, 4H), 7.24 (m, 3H), 6.99 (m, 3H), 6.67 (br, 3H), 3.97 (br, 1H), 3.88 (s, 2H), 3.79 (s, 2H), 3.73-3.23 (br m, 12H), 3.14 (m, 6H), 2.29 (s, 6H), 2.08 (m, 2H), 1.74 (s, 4H).

EXAMPLE 16A

20 This example was prepared by substituting azepane for N-isopropyl-N-methylamine in EXAMPLE 4D.

EXAMPLE 16B

25 This example was prepared by substituting EXAMPLE 16A for EXAMPLE 4D in EXAMPLE 4E.

EXAMPLE 16C

30 This example was prepared by substituting EXAMPLE 16A for EXAMPLE 1D in EXAMPLE 1I.

EXAMPLE 16D

This example was prepared by substituting EXAMPLE 16C and EXAMPLE 3C for EXAMPLE 1I and EXAMPLE 1M, respectively, in EXAMPLE 1N. ¹H NMR (400MHz, CDCl₃) δ 8.33 (s, 1H), 8.01 (d, 1H), 7.67 (d, 2H), 7.34 (t, 4H), 7.23 (m, 3H), 6.98 (m, 3H),

6.67 (m, 3H), 3.99 (m, 1H), 3.82-3.19 (br m, 10H), 3.12 (s, 4H), 2.86 (m, 2H), 2.55 (br, 2H), 2.29 (s, 4H), 2.06 (m, 1H), 1.93 (m, 3H), 1.74 (s, 8H), 1.60 (m, 2H).

EXAMPLE 17A

5 This example was prepared by substituting EXAMPLE 6A for EXAMPLE 1C in EXAMPLE 1D.

EXAMPLE 17B

10 This example was prepared by substituting EXAMPLE 17B for EXAMPLE 1D in EXAMPLE 1I.

EXAMPLE 17C

This example was prepared by substituting 4-(4-(2-(4-chlorophenyl)cyclohept-1-enylmethyl)piperazin-1-yl)benzoic acid, prepared as described in commonly-owned U.S. Patent Application Ser. No. 10/988,338, and EXAMPLE 17B for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (400MHz, DMSO-d₆) δ 9.58 (brs, 1H), 9.46 (brs, 1H), 8.18 (d, 1H), 8.00 (dd, 1H), 7.77 (d, 2H), 7.41 (d, 2H), 7.29 (d, 2H), 7.24 (m, 2H), 7.13 (m, 4H), 6.96 (m, 3H), 4.12 (m, 1H), 3.87 (m, 1H), 3.63 (m, 1H), 3.38 (m, 4H), 3.15 (m, 4H), 3.02 (m, 2H), 2.74 (s, 6H), 2.46 (m, 4H), 2.09 (m, 2H), 1.81 (m, 2H), 1.57 (m, 4H).

15
20

EXAMPLE 18A

This example was prepared by substituting EXAMPLE 2E and EXAMPLE 17B for EXAMPLE 1H and EXAMPLE 1D in EXAMPLE 1I.

25

EXAMPLE 18B

This example was prepared by substituting EXAMPLE 18A and EXAMPLE 3C for EXAMPLE 1I and EXAMPLE 1M, respectively, in EXAMPLE 1N. ¹H NMR (400MHz, DMSO-d₆) δ 9.60 (brs, 1H), 9.47 (brs, 1H), 8.18 (d, 1H), 7.99 (dd, 1H), 7.77 (d, 2H), 7.41 (d, 2H), 7.30 (d, 2H), 7.24 (m, 2H), 7.15 (m, 3H), 7.12 (d, 1H), 6.96 (m, 3H), 6.92 (d, 1H), 4.10 (m, 1H), 3.91 (m, 2H), 3.60 (m, 2H), 3.37 (m, 4H), 3.15 (m, 2H), 3.02 (m, 1H), 2.74 (s, 6H), 2.25 (d, 4H), 2.08 (m, 2H), 1.71 (m, 4H).

30

EXAMPLE 19

This example was prepared by substituting EXAMPLE 2F for EXAMPLE 1I in EXAMPLE 1N. ¹H NMR (400MHz, DMSO-d₆) δ 8.11 (d, 1H), 7.93 (dd, 1H), 7.71 (d, 2H), 7.36 (m, 4H), 7.27 (m, 2H), 7.18 (m, 1H), 7.12 (d, 2H), 6.97 (d, 1H), 6.85 (m, 3H), 4.05 (m, 1H), 3.53 (m, 4H), 3.23 (m, 1H), 2.83 (m, 1H), 2.34 (m, 8H), 2.22 (m, 2H), 1.99 (m, 2H),
5 1.96 (m, 1H), 1.77 (m, 1H), 1.44 (t, 2H), 0.97 (s, 6H).

EXAMPLE 20

This example was prepared by substituting EXAMPLE 17B for EXAMPLE 1I in EXAMPLE 1N. ¹H NMR (500 MHz, DMSO-d₆) δ 9.46 (brs, 1H), 8.18 (d, 1H), 7.99 (dd, 1H), 7.76 (d, 2H), 7.40 (d, 2H), 7.29 (d, 2H), 7.23 (t, 2H), 7.14 (s, 4H), 6.95 (m, 3H), 4.11 (m, 1H), 3.88 (m, 2H), 3.58 (m, 4H), 3.08 (m, 4H), 2.73 (s, 6H), 2.27 (m, 2H), 2.08 (m, 2H),
10 2.02 (s, 2H), 1.47 (t, 2H), 1.00 (s, 6H).

EXAMPLE 21

This example was prepared by substituting 4-(4-(4-(4-chlorophenyl)-5,6-dihydro-2H-pyran-3-ylmethyl)piperazin-1-yl)benzoic acid, prepared as described in commonly-owned U.S. Patent Application Ser. No. 10/988,338, for EXAMPLE 1M in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ 8.27 (d, 1H), 8.11 (d, 1H), 7.89 (d, 2H), 7.59 (d, 2H), 7.48 (m, 4H), 7.37 (m, 3H), 7.13 (m, 1H), 7.01 (m, 3H), 4.35 (s, 2H), 4.24 (m, 1H), 3.97 (m, 2H),
15 3.68 (m, 4H), 3.36 (m, 6H), 3.07 (m, 3H), 2.68 (s, 2H), 2.59 (m, 4H), 2.14 (m, 2H), 1.93 (m, 2H).

EXAMPLE 22A

This example was prepared by substituting EXAMPLE 4E for EXAMPLE 2E in EXAMPLE 2F.
25

EXAMPLE 22

This example was prepared by substituting EXAMPLE 2K and EXAMPLE 22A for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ 9.21 (brs, 1H), 8.17 (m, 1H), 8.00 (dd, 1H), 7.77 (d, 2H), 7.42 (m, 2H), 7.31 (m, 2H), 7.24 (m, 2H), 7.14 (m, 4H), 6.97 (m, 3H), 4.11 (m, 1H), 3.90 (m, 1H), 3.12 (m, 6H), 2.84 (m, 3H), 2.63 (m, 3H), 2.25 (m, 2H), 2.07 (m, 4H), 1.49 (t, 2H), 1.16 (m, 6H), 0.97 (s, 6H).
30

EXAMPLE 23

35

This example was prepared by substituting EXAMPLE 22A for EXAMPLE 1I in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ 9.21 (brs, 1H), 8.18 (m, 1H), 7.99 (dd, 1H), 7.78 (d, 2H), 7.40 (d, 2H), 7.30 (d, 2H), 7.24 (m, 2H), 7.15 (m, 4H), 6.97 (m, 3H), 4.11 (m, 1H), 3.89 (m, 1H), 3.13 (m, 6H), 2.84 (m, 2H), 2.63 (m, 3H), 2.28 (m, 2H), 2.07 (m, 4H), 1.48 (t, 2H), 1.17 (m, 6H), 1.00 (s, 6H).

EXAMPLE 24

This example was prepared by substituting EXAMPLE 2K for EXAMPLE 1M in EXAMPLE 1N. ¹H NMR (500 MHz, DMSO-d₆) δ 12.14 (brs, 1 H), 9.89 (brs, 1H), 9.52 (s, 1H), 8.18 (d, 1H), 8.00 (dd, 1H), 7.77 (d, 2H), 7.41 (d, 2H), 7.30 (d, 2H), 7.24 (t, 2H), 7.14 (m, 4H), 6.96 (m, 3H), 4.12 (m, 1H), 3.93 (m, 3H), 3.63 (m, 4H), 2.93 (m, 10H), 2.24 (m, 2H), 2.09 (m, 4H), 1.48 (t, 2H), 0.97 (s, 6H).

EXAMPLE 25

This example was prepared by substituting EXAMPLE 2K and EXAMPLE 6F for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (500 MHz, DMSO-d₆) δ 9.58 (brs, 1H), 9.39 (brs, 1H), 8.17 (s, 1H), 7.99 (dd, 1H), 7.77 (d, 2H), 7.41 (d, 2H), 7.30 (d, 2H), 7.24 (t, 2H), 7.14 (m, 4H), 6.97 (m, 3H), 4.63 (d, 1H), 4.43 (d, 1H), 4.13 (m, 1H), 3.92 (m, 2H), 3.69 (m, 2H), 3.52 (m, 2H), 3.01 (m, 6H), 2.25 (m, 2H), 2.04 (m, 6H), 1.49 (m, 2H), 0.98 (s, 6H).

EXAMPLE 26

This example was prepared by substituting EXAMPLE 17B and EXAMPLE 3C for EXAMPLE 1I and EXAMPLE 1M, respectively, in EXAMPLE 1N. ¹H NMR (500 MHz, DMSO-d₆) δ 9.49 (brs, 1H), 8.08 (d, 1H), 7.95 (dd, 1H), 7.71 (d, 2H), 7.36 (m, 4H), 7.30 (t, 2H), 7.20 (t, 1H), 7.12 (d, 2H), 6.84 (m, 2H), 6.78 (d, 2H), 3.98 (m, 1H), 3.28 (m, 2H), 3.12 (brs, 4H), 2.81 (brs, 1H), 2.77 (s, 1H), 2.46 (s, 6H), 2.28 (s, 4H), 2.19 (m, 4H), 2.00 (m, 1H), 1.90 (m, 1H), 1.65 (m, 4H).

EXAMPLE 27A

This example was prepared by substituting pyrrolidine for N-isopropylethylamine in EXAMPLE 4D.

EXAMPLE 27B

This example was prepared by substituting EXAMPLE 27A for EXAMPLE 4D in EXAMPLE 4E.

EXAMPLE 27C

5 This example was prepared by substituting EXAMPLE 27B for EXAMPLE 1D in EXAMPLE 1I.

EXAMPLE 27D

10 This example was prepared by substituting EXAMPLE 27C and EXAMPLE 3C for EXAMPLE 1I and EXAMPLE 1M, respectively, in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ 8.08 (d, 1H), 7.96 (dd, 1H), 7.71 (d, 2H), 7.36 (m, 4H), 7.30 (t, 2H), 7.20 (t, 1H), 7.12 (d, 2H), 6.87 (m, 1H), 6.77 (d, 2H), 6.72 (d, 1H), 4.00 (m, 1H), 3.26 (m, 2H), 3.12 (brs, 4H), 2.97 (m, 6H), 2.76 (s, 1H), 2.28 (brs, 4H), 2.19, (m, 4H), 2.05 (m, 1H), 1.95 (m, 1H), 1.82 (brs, 4H), 1.65 (m, 4H).

15

EXAMPLE 28

This example was prepared by substituting EXAMPLE 2K and EXAMPLE 17B for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (500 MHz, DMSO-d₆) δ 9.59 (brs, 1H), 8.08 (d, 1H), 7.94 (dd, 1H), 7.70 (d, 2H), 7.36 (m, 4H), 7.30 (t, 2H), 7.21 (tt, 1H), 7.09 (d, 2H), 6.83 (d, 1H), 6.78 (d, 3H), 3.97 (m, 1H), 3.28 (m, 2H), 3.13 (brs, 4H), 2.90 (brs, 2H), 2.79 (s, 2H), 2.55 (s, 6H), 2.28 (brs, 4H), 2.20 (m, 2H), 1.99 (s, 2H), 1.90 (m, 2H), 1.42 (t, 2H), 0.96 (s, 6H).

20

EXAMPLE 29

25 This example was prepared by substituting 4-(4-(2-(4-chlorophenyl)cyclo-1-enylmethyl)piperazin-1-yl)benzoic acid, prepared as described in commonly-owned U.S. Patent Application Ser. No. 10/988,338, and EXAMPLE 27C for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ 9.60 (brs, 1H), 8.08 (d, 1H), 7.94 (dd, 1H), 7.71 (d, 2H), 7.36 (m, 4H), 7.30 (t, 2H), 7.20 (t, 1H), 7.09 (d, 2H), 6.83 (d, 1H), 6.77 (d, 3H), 3.99 (m, 1H), 3.26 (m, 2H), 3.12 (brs, 4H), 2.80 (m, 5H), 2.76 (s, 2H), 2.40 (m, 4H), 2.31 (brs, 4H), 1.99 (m, 1H), 1.89 (m, 1H), 1.77 (brs, 6H), 1.58 (m, 2H), 1.51 (m, 2H).

30

EXAMPLE 30A

This example was prepared by substituting EXAMPLE 2E and EXAMPLE 27B for EXAMPLE 1H and EXAMPLE 1D in EXAMPLE 1I.

EXAMPLE 30B

5 This example was prepared by substituting EXAMPLE 30A for EXAMPLE 1I and EXAMPLE 1D in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ 9.57 (brs, 1H), 8.07 (d, 1H), 7.93 (dd, 1H), 7.70 (d, 2H), 7.36 (m, 4H), 7.30 (t, 2H), 7.21 (t, 1H), 7.12 (d, 2H), 6.81 (d, 1H), 6.77 (d, 3H), 3.99 (m, 1H), 3.26 (m, 2H), 3.12 (brs, 4H), 2.80 (m, 5H), 2.76 (s, 2H), 2.27 (m, 4H), 2.22 (m, 2H), 1.99 (m, 3H), 1.88 (m, 1H), 1.77 (brs, 4H), 1.43 (t, 2H),
10 0.97 (s, 6H).

EXAMPLE 31

This example was prepared by substituting EXAMPLE 2K and EXAMPLE 30A for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (400 MHz,
15 DMSO-d₆) δ 9.52 (brs, 1H), 8.06 (d, 1H), 7.92 (dd, 1H), 7.70 (d, 2H), 7.36 (m, 4H), 7.30 (t, 2H), 7.21 (t, 1H), 7.08 (d, 2H), 6.81 (d, 1H), 6.77 (d, 3H), 3.99 (m, 1H), 3.26 (m, 2H), 3.12 (brs, 4H), 2.76 (s, 2H), 2.75 (m, 5H), 2.26 (m, 4H), 2.20 (m, 2H), 1.99 (m, 3H), 1.86 (m, 1H), 1.76 (brs, 4H), 1.42 (t, 2H), 0.96 (s, 6H).

EXAMPLE 32

20 This example was prepared by substituting 4-(4-(2-(4-chlorophenyl)cyclohept-1-enylmethyl)piperazin-1-yl)benzoic acid, prepared as described in commonly-owned U.S. Patent Application Ser. No. 10/988,338, and EXAMPLE 30A for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ 9.50 (brs, 1H), 8.08 (d, 1H), 7.94 (dd, 1H), 7.71 (d, 2H), 7.36 (m, 4H), 7.30 (t, 2H), 7.20 (t, 1H), 7.09 (d, 2H), 6.83 (d, 1H), 6.77 (d, 3H), 3.99 (m, 1H), 3.26 (m, 2H), 3.12 (brs, 4H), 2.80 (m, 5H), 2.76 (s, 2H), 2.40 (m, 4H), 2.31 (brs, 4H), 1.98 (m, 1H), 1.87 (m, 1H), 1.76 (brs, 6H), 1.58 (m, 2H), 1.51 (m, 2H).

EXAMPLE 33

30 This example was prepared by substituting EXAMPLE 4F and EXAMPLE 3C for EXAMPLE 1I and EXAMPLE 1M, respectively, in EXAMPLE 1N. ¹H NMR (300MHz, DMSO-d₆) δ 9.03 (s, 1H), 8.08 (d, 1H), 7.97 (d, 1H), 7.71 (d, 2H), 7.39-7.34 (m, 4H), 7.30 (t, 2H), 7.20 (tt, 1H), 7.13 (dt, 2H), 6.88 (m, 1H), 6.78 (d, 2H), 6.70 (m, 1H), 3.99 (m, 1H),

3.37-3.26 (m, 4H), 3.12 (s, 4H), 2.76 (s, 2H), 2.68-2.53 (m, 2H), 2.34-2.13 (m, 10H), 2.10-1.95 (m, 2H), 1.66 (s, 4H), 1.13 (m, 6H).

EXAMPLE 34

5 This example was prepared by substituting EXAMPLE 2K and EXAMPLE 27C for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (300MHz, DMSO-d₆) δ 9.53 (s, 1H), 8.08 (d, 1H), 7.97 (dd, 1H), 7.71 (d, 2H), 7.40-7.34 (m, 4H), 7.30 (t, 2H), 7.20 (tt, 1H), 7.09 (d, 2H), 6.89 (d, 1H), 6.78 (d, 2H), 6.71 (d, 1H), 4.01 (m, 1H), 3.38-3.27 (m, 4H), 3.20-2.84 (m, 10H), 2.79 (s, 2H), 2.27 (s, 4H), 2.20 (t, 2H), 2.03 (m, 2H),
10 1.85 (m, 4H), 1.42 (t, 2H), 0.96 (s, 6H).

EXAMPLE 35

This example was prepared by substituting EXAMPLE 3C and EXAMPLE 30A for EXAMPLE 1M and EXAMPLE 1I, respectively in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ ppm 8.06 (d, 1H), 7.93 (dd, 1H), 7.71 (d, 2H), 7.36 (m, 4H), 7.30 (m, 2H), 7.21 (m, 1H), 7.12 (d, 2H), 6.81 (d, 1H), 6.77 (d, 3H), 3.97 (m, 1H), 3.26 (m, 4H), 3.12 (s, 4H), 2.78 (m, 6H), 2.27 (s, 4H), 2.18 (m, 4H), 1.99 (m, 1H), 1.87 (m, 1H), 1.76 (s, 4H), 1.66 (s, 4H).

20

EXAMPLE 36

This example was prepared by substituting EXAMPLE 2K and EXAMPLE 9A for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ 8.09 (d, 1H), 7.97 (dd, 1H), 7.71 (d, 2H), 7.33 (m, 6H), 7.21 (m, 1H), 7.08 (d, 2H), 6.87 (m, 1H), 6.78 (m, 3H), 3.99 (m, 1H), 3.14 (m, 4H), 2.95 (m, 1H), 2.80 (m, 3H),
25 2.58 (s, 6H), 2.28 (m, 4H), 2.20 (m, 2H), 1.99 (m, 4H), 1.42 (t, 2H), 0.96 (s, 6H).

EXAMPLE 37

This example was prepared by substituting EXAMPLE 2K and EXAMPLE 17B for EXAMPLE 1M and EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (400 MHz, DMSO-d₆) δ 8.09 (d, 1H), 7.97 (dd, 1H), 7.71 (d, 2H), 7.33 (m, 6H), 7.21 (m, 1H), 7.08 (d, 2H), 6.87 (m, 1H), 6.78 (m, 3H), 3.99 (m, 1H), 3.14 (m, 4H), 2.95 (m, 1H), 2.80 (m, 3H), 2.58 (s, 6H), 2.28 (m, 4H), 2.20 (m, 2H), 1.99 (m, 4H), 1.42 (t, 2H), 0.96 (s, 6H).

EXAMPLE 38


This example was prepared by substituting 4-(4-(1,1'-biphenyl-2-ylmethyl)-1-piperazinyl)benzoic acid, prepared as described in commonly-owned U.S. Patent Application Ser. No. 10/988,338, for EXAMPLE 1M and EXAMPLE 17B for EXAMPLE 1I, respectively, in EXAMPLE 1N. ¹H NMR (400MHz, DMSO-d₆) δ 8.19 (d, 1H), 7.99 (dd, 1H), 7.76 (d, 3H), 7.52 (d, 4H), 7.40 (d, 2H), 7.35 (m, 1H), 7.30 (d, 2H), 7.24 (t, 2H), 7.16 (t, 2H), 6.96 (m, 3H), 4.25 (br, 2H), 4.12 (m, 1H), 3.37 (m, 2H), 3.14 (m, 1H), 3.10 (br, 8H), 2.74 (s, 6H), 2.10 (m, 2H).

EXAMPLE 39

This example was prepared by substituting 4-(4-(1,1'-biphenyl-2-ylmethyl)-1-piperazinyl)benzoic acid, prepared as described in commonly-owned U.S. Patent Application Ser. No. 10/988,338, for EXAMPLE 1M in EXAMPLE 1N. ¹H NMR (400MHz, DMSO-d₆) δ 8.19 (d, 1H), 8.00 (dd, 1H), 7.76(d, 2H), 7.52 (m, 5H), 7.14 (m, 8H), 6.96 (m, 3H), 4.29 (m, 2H), 4.14 (m, 2H), 4.02 (m, 1H), 3.10 (m, 8H), 2.13 (m, 2H).

The foregoing is meant to illustrate the invention but not to limit it. Variations and changes obvious to one skilled in the art are intended to be within the scope of the invention as defined in the claims.

1. A compound having formula (II)




(II),

or a therapeutically acceptable salt thereof, wherein X^3 is Cl or F;

5

X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $N(CH_3)_2$, $N(CH_3)(CH(CH_3)_2)$, 7-azabicyclo[2.2.1]heptan-1-yl or 2-oxa-5-azabicyclo[2.2.1]hept-5-yl, and R^0 is




, wherein

X^5 is CH_2 , $C(CH_3)_2$ or CH_2CH_2 ;

10 X^6 and X^7 are both hydrogen or are both methyl; and


X^8 is F, Cl, Br or I; or

X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $N(CH_3)(CH(CH_3)_2)$ or 7-azabicyclo[2.2.1]heptan-1-yl, and R^0 is




; or

15 X^4 is $N(CH_3)_2$ or morpholin-1-yl, and R^0 is



2. A compound of claim 1, or a therapeutically acceptable salt thereof, wherein X^3 is Cl or F;

X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $N(CH_3)_2$, $N(CH_3)(CH(CH_3)_2)$, 7-azabicyclo[2.2.1]heptan-1-yl or 2-oxa-5-azabicyclo[2.2.1]hept-5-yl, and R^0 is




, wherein

X^5 is CH_2 , $C(CH_3)_2$ or CH_2CH_2 ;

X^6 and X^7 are both hydrogen or are both methyl; and


X^8 is F, Cl, Br or I; or

10 X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $N(CH_3)(CH(CH_3)_2)$ or 7-azabicyclo[2.2.1]heptan-1-yl, and R^0 is




; or

X^4 is $N(CH_3)_2$ or morpholin-1-yl, and R^0 is



3. A compound of claim 1, or a therapeutically acceptable salt thereof, wherein X^3 is Cl or F;


X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $N(CH_3)_2$, $N(CH_3)(CH(CH_3)_2)$, 7-azabicyclo[2.2.1]heptan-1-yl or 2-oxa-5-azabicyclo[2.2.1]hept-5-yl, and R^0 is



, wherein X^5 is CH_2 , $\text{C}(\text{CH}_3)_2$ or CH_2CH_2 , and X^6 and X^7 are both hydrogen or are both methyl; and X^8 is F, Cl, Br or I.

4. A compound of claim 1, or a therapeutically acceptable salt thereof, wherein X^3 is Cl or F;


X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $\text{N}(\text{CH}_3)(\text{CH}(\text{CH}_3)_2)$ or 7-azabicyclo[2.2.1]heptan-1-yl; R^0 is



, wherein X^6 and X^7 are both hydrogen or are both methyl; and X^8 is F, Cl, Br or I.


5. A compound of claim 1, or a therapeutically acceptable salt thereof, wherein X^3 is Cl or F;

X^4 is $\text{N}(\text{CH}_3)_2$ or morpholin-1-yl; R^0 is



; and X^8 is F, Cl, Br or I.

6. A compound of claim 1, or a therapeutically acceptable salt thereof, wherein X^3 is F; X^4 is morpholin-1-yl; R^0 is




, wherein X^5 is $\text{C}(\text{CH}_3)_2$; X^6 and X^7 are both methyl; and

X^8 is Cl.

5

7. A compound of claim 1, or a therapeutically acceptable salt thereof, wherein X^3 is Cl or F;


X^4 is azepan-1-yl, morpholin-1-yl, pyrrolidin-1-yl, $N(CH_3)(CH(CH_3)_2)$ or 7-azabicyclo[2.2.1]heptan-1-yl; R^0 is



5 X^6 and X^7 are both hydrogen or are both methyl; and X^8 is F, Cl, Br or I.


8. A compound of claim 1, or a therapeutically acceptable salt thereof, wherein X^3 is Cl or F;

X^4 is $N(CH_3)_2$ or morpholin-1-yl; R^0 is



5 X^8 is F, Cl, Br or I.

9. A compound of claim 1, or a therapeutically acceptable salt thereof, wherein X^3 is Cl or F; X^4 is morpholin-1-yl; R^0 is



5 X^5 is $C(CH_3)_2$; X^6 and X^7 are both methyl; and X^8 is Cl.

10. A composition comprising an excipient and a therapeutically effective amount of a compound of claim 1.

11. A method of treating bladder cancer, brain cancer, breast cancer, bone marrow cancer, cervical cancer, chronic lymphocytic leukemia, colorectal cancer, esophageal cancer,

hepatocellular cancer, lymphoblastic leukemia, follicular lymphoma, lymphoid malignancies
of T-cell or B-cell origin, melanoma, myelogenous leukemia, myeloma, oral cancer, ovarian
5 cancer, non-small cell lung cancer, prostate cancer, small cell lung cancer or spleen cancer in
a patient, said method comprising administering to the patient a therapeutically effective
amount of a compound of claim 1.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

