

(19) **DANMARK**

(10) **DK/EP 2436700 T5**



Patent- og
Varemærkestyrelsen

(12) **Rettet oversættelse af
europæisk patentskrift**

-
- (51) Int.Cl.: **C 08 B 37/00 (2006.01)** **A 61 K 47/64 (2017.01)** **C 12 P 19/04 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-09-03**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-06-20**
- (86) Europæisk ansøgning nr.: **11194173.8**
- (86) Europæisk indleveringsdag: **2008-03-20**
- (87) Den europæiske ansøgnings publiceringsdag: **2012-04-04**
- (30) Prioritet: **2007-03-23 US 896616 P**
- (62) Stamansøgningsnr: **08732592.4**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR**
- (73) Patenthaver: **Wyeth LLC, 235 East 42nd Street, New York, NY 10017, USA**
- (72) Opfinder: **Yuan, Yonghui, 118 Newport Avenue, Tappan, NY New York 10983, USA**
Ruppen, Mark, 6 Lea Avenue, Garnerville, NY New York 10923, USA
Sun, Wei-Qiang, 30 Windmill Drive, Morristown, NJ New Jersey 07960, USA
Chu, Ling, 5 Potter Lane, Suffern, NY New York 10901, USA
Simpson, John, 417 Maple Avenue, Upper Nyack, NY New York 10960, USA
Patch, James, 11 River Avenue, Cornwall On Hudson, NY New York 12520, USA
Fink Charbonneau, Pamela, 8 Eldor Avenue, New City, NY New York 10956, USA
Moran, Justin K., 682 Sierra Vista Lane, Valley Cottage, NY New York 10989, USA
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **AFKORTET OPRENSNINGSFREMANGSMÅDE TIL FREMSTILLING AF STREPTOCOCCUS PNEUMONIAE-KAPSELPOLYSACCHARIDER**
- (56) Fremdragne publikationer:
EP-A- 0 002 404
EP-A- 1 762 245
WO-A-82/01995
US-A- 5 623 057
US-A1- 2006 228 381

DESCRIPTION

FIELD OF THE INVENTION

[0001] This invention relates to methods for removing excess soluble protein and other impurities from cellular lysates of *Streptococcus pneumoniae* (*S. pneumoniae*) serotypes used in the production of purified pneumococcal polysaccharides.

BACKGROUND OF THE INVENTION

[0002] *Streptococcus pneumoniae* are Gram-positive, lancet shaped cocci that are usually seen in pairs (diplococci), but also in short chains or as single cells. They grow readily on blood agar plates with glistening colonies and display alpha hemolysis unless grown anaerobically where they show beta hemolysis. They are sensitive to bile salts that can break down the cell wall with the presence of the cells' own enzyme, autolysin. The organism is an aerotolerant anaerobe and is fastidious in that it has complex nutritional requirements.

[0003] The cells of most pneumococcal serotypes have a capsule which is a polysaccharide coating surrounding each cell. This capsule is a determinant of virulence in humans because it interferes with phagocytosis by preventing antibodies from attaching to the bacterial cells. There are currently 90 capsular serotypes identified, with 23 serotypes responsible for about 90% of invasive disease. As a vaccine the polysaccharide can confer a reasonable degree of immunity to *S. pneumoniae* in individuals with developed or unimpaired immune systems. However, when the polysaccharide is conjugated with a high molecular weight protein such as CRM₁₉₇ and formulated into a vaccine containing conjugates of multiple serotypes, such conjugate vaccines allow for an immune response in infants and elderly who are also most at risk for pneumococcal infections.

[0004] The capsular polysaccharide for each *S. pneumoniae* serotype utilized for vaccine products is produced by growing the organism in liquid medium. The population of the organism is often scaled up from a seed vial to seed bottles and passed through one or more seed fermentors of increasing volume until production scale fermentation volumes are reached. The end of the growth cycle can be determined by one of several means, at which point the cells are lysed through the addition of a detergent or other reagent which aids in the cell wall breakdown and release of autolysin which causes cellular lysis when the cells reach stationary phase. The broth is then harvested for downstream (purification) processing. The major contaminants are cellular proteins, nucleic acids, C-polysaccharide and medium components.

[0005] For most of the serotypes for the currently marketed 7-valent pneumococcal conjugate (7vPnC) vaccine (Prevnar®), as well as the newly developed 13-valent pneumococcal

conjugate (13vPnC) vaccine, the current purification process requires sixteen steps involving many expensive, labor intensive and technologically demanding operations, such as chromatography and multiple membrane separations. Previous attempts at improving purification processes for *S. Pneumoniae* polysaccharides have included, for example, pH manipulation during fermentation and recovery (see U.S. Patent App. Pub. No. 2006/0228381) and solvent and detergent precipitation. However, the removal of impurities in these processes is still spread over many labor intensive and costly steps. Protein level is the most problematic specification to meet due to the physical and chemical properties of the soluble proteins.

[0006] Thus, there is a need for a simplified purification process to reduce the soluble protein levels in *S. pneumoniae* lysates and eliminate inefficiencies of the current purification process to produce substantially purified capsular polysaccharides suitable for incorporation into pneumococcal conjugate vaccines.

SUMMARY OF THE INVENTION

[0007] The present invention relates to a process for producing substantially purified capsular polysaccharides from a *Streptococcus pneumoniae* cell lysate. This process comprises the steps of:

1. (a) providing a fermentation broth comprising bacterial cells that produce a selected *Streptococcus pneumoniae* serotype;
2. (b) lysing the bacterial cells in step (a) with a lytic agent, thereby producing a cell lysate comprising cell debris, soluble proteins, nucleic acids, and polysaccharides;
3. (c) clarifying the cell lysate of step (b) using centrifugation or filtration to remove cell debris, thereby producing a clarified cell lysate;
4. (d) ultrafiltering and diafiltering the clarified cell lysate of step (c) to remove low molecular weight impurities and increase polysaccharide concentration, thereby producing a retentate;
5. (e) lowering the pH of the retentate of step (d) to less than 4.5 using an organic or a mineral acid to precipitate protein and nucleic acids, thereby forming an acidified retentate solution;
6. (f) holding the acidified retentate solution formed in step (e) for a time sufficient to allow settling of the precipitate, followed by filtration or centrifugation of the acidified retentate solution, thereby producing a clarified polysaccharide solution;
7. (g) filtering the clarified polysaccharide solution of step (f) through an activated carbon filter;
8. (h) ultrafiltering and diafiltering the filtered solution produced by step (g), thereby producing a concentrated purified polysaccharide solution; and
9. (i) filtering the concentrated purified polysaccharide solution produced by step (h) using a sterile filter;

whereby substantially purified capsular polysaccharides in the form of a solution are produced.

Exemplary, non-limiting *S. pneumoniae* serotypes selected for this embodiment of the invention are 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F. In a particular embodiment, the pH of step (e) is lowered to about 3.5. In another embodiment, the diafiltration of step (h) comprises a pH adjustment to between about 5.5 to about 7.5. In another embodiment, the diafiltration of step (h) comprises a pH adjustment to between about 7.0 to about 7.5. In another embodiment, the diafiltration of step (h) comprises a pH adjustment to about 7.4. In still another embodiment, step (e) removes at least 98% of protein from the retentate of step (d). In another embodiment, step (g) removes at least 90% of the protein from the clarified polysaccharide solution of step (f). In another embodiment, the activated carbon filter of step (g) comprises wood-based phosphoric acid-activated carbon. In another embodiment, step (f) comprises holding the acidified retentate solution formed in step (e) for at least 2 hours. In still another embodiment, the lytic agent of step (b) is deoxycholate sodium (DOC). In another embodiment, the lytic agent of step (b) is a non-animal derived lytic agent. In still another embodiment, the lytic agent of step (b) is the non-animal derived lytic agent N-lauryl sarcosine sodium (NLS).

[0008] The substantially purified capsular polysaccharide produced by a process of the invention from a *Streptococcus pneumoniae* cell lysate may be used in the manufacture of a pneumococcal vaccine, preferably a pneumococcal vaccine containing polysaccharide conjugated with a protein carrier.

[0009] The present invention also relates to a process for producing substantially purified capsular polysaccharides from a *Streptococcus pneumoniae* cell lysate comprising serotype 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F, or 23F. This process comprises the steps of:

1. (a) providing a fermentation broth comprising bacterial cells that produce *Streptococcus pneumoniae* serotype 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F, or 23F;
2. (b) lysing the bacterial cells in step (a) with a lytic agent, thereby producing a cell lysate comprising cell debris, soluble proteins, nucleic acids, and polysaccharides;
3. (c) clarifying the cell lysate of step (b) using centrifugation or filtration to remove cell debris, thereby producing a clarified cell lysate;
4. (d) ultrafiltering and diafiltering the clarified cell lysate of step (c) at room temperature at neutral pH in salt free media to remove low molecular weight impurities and increase polysaccharide concentration, thereby producing a salt free retentate;
5. (e) lowering the pH of the salt free retentate of step (d) to less than 4.5 using an organic or a mineral acid to precipitate protein and nucleic acids, thereby forming an acidified retentate solution;
6. (f) holding the acidified retentate solution formed in step (e) for at least 2 hours at room temperature to allow settling of the precipitate, followed by filtration or centrifugation of the acidified retentate solution, thereby producing a clarified polysaccharide solution;
7. (g) filtering the clarified polysaccharide solution of step (f) through an activated carbon filter;
8. (h) ultrafiltering and diafiltering the filtered solution produced by step (g), thereby producing a concentrated purified polysaccharide solution; and

9. (i) filtering the concentrated purified polysaccharide solution produced by step (h) using a sterile filter;

whereby substantially purified capsular polysaccharides comprising serotype 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F, or 23F in the form of a solution are produced. In a particular embodiment, the pH of step (e) is lowered to about 3.5. In another embodiment, the diafiltration of step (h) comprises a pH adjustment to between about 5.5 to about 7.5. In another embodiment, the diafiltration of step (h) comprises a pH adjustment to between about 7.0 to about 7.5. In another embodiment, the diafiltration of step (h) comprises a pH adjustment to about 7.4. In still another embodiment, step (e) removes at least 98% of protein from the salt free retentate of step (d). In another embodiment, step (g) removes at least 90% of the protein from the clarified polysaccharide solution of step (f). In another embodiment, the activated carbon filter of step (g) comprises wood-based phosphoric acid-activated carbon. In still another embodiment, the lytic agent of step (b) is deoxycholate sodium (DOC). In another embodiment, the lytic agent of step (b) is a non-animal derived lytic agent. In still another embodiment, the lytic agent of step (b) is the non-animal derived lytic agent N-lauryl sarcosine sodium (NLS).

[0010] The present invention also relates to a process for producing substantially purified capsular polysaccharides from a *Streptococcus pneumoniae* cell lysate comprising serotype 19A. This process comprises the steps of:

1. (a) providing a fermentation broth comprising bacterial cells that produce *Streptococcus pneumoniae* serotype 19A;
2. (b) lysing the bacterial cells in step (a) with a lytic agent, thereby producing a cell lysate comprising cell debris, soluble proteins, nucleic acids, and polysaccharides;
3. (c) clarifying the cell lysate of step (b) using centrifugation or filtration to remove cell debris, thereby producing a clarified cell lysate;
4. (d) ultrafiltering and diafiltering the clarified cell lysate of step (c) at about 4 °C at a pH of about 6 in sodium phosphate buffer to remove low molecular weight impurities and increase polysaccharide concentration, thereby producing a retentate;
5. (e) lowering the pH of the retentate of step (d) to less than 4.5 using an organic or a mineral acid to precipitate protein and nucleic acids, thereby forming an acidified retentate solution;
6. (f) holding the acidified retentate solution formed in step (e) for at least 2 hours at about 4 °C to allow settling of the precipitate, followed by filtration or centrifugation of the acidified retentate solution, thereby producing a clarified polysaccharide solution;
7. (g) adjusting the pH of the clarified polysaccharide solution of step (f) to about 6, thereby producing a pH-adjusted clarified polysaccharide solution;
8. (h) filtering the pH-adjusted clarified polysaccharide solution of step (g) through an activated carbon filter;
9. (i) ultrafiltering and diafiltering the filtered solution produced by step (h), thereby producing a concentrated purified polysaccharide solution; and
10. (j) filtering the concentrated purified polysaccharide solution produced by step (i) using a sterile filter;

whereby substantially purified capsular polysaccharides comprising serotype 19A in the form of a solution are produced. In a particular embodiment, the pH of step (e) is lowered to about 3.5. In another embodiment, the diafiltration of step (i) comprises a pH adjustment to between about 5.5 to about 7.5. In another embodiment, the diafiltration of step (i) comprises a pH adjustment to between about 7.0 to about 7.5. In another embodiment, the diafiltration of step (i) comprises a pH adjustment to about 7.4. In still another embodiment, step (e) removes at least 98% of protein from the retentate of step (d). In another embodiment, step (h) removes at least 90% of the protein from the pH-adjusted clarified polysaccharide solution of step (g). In another embodiment, the activated carbon filter of step (h) comprises wood-based phosphoric acid-activated carbon. In another embodiment, the sodium phosphate buffer of step (d) is 25 mM sodium phosphate. In still another embodiment, the lytic agent of step (b) is deoxycholate sodium (DOC). In another embodiment, the lytic agent of step (b) is a non-animal derived lytic agent. In still another embodiment, the lytic agent of step (b) is the non-animal derived lytic agent N-lauryl sarcosine sodium (NLS).

BRIEF DESCRIPTION OF THE DRAWINGS

[0011]

Figure 1 shows average in-process polysaccharide (PS) yield, protein/PS ratio, and nucleic acid (NA)/PS ratio for serotype 5 using the shortened purification process of the invention. Results are shown for each purification step.

Figure 2 shows NMR spectra for serotype 4 PS from the current purification process (A) compared to serotype 4 PS from the shortened purification process (B). No significant differences between the two spectra were observed. The second peak from the right in both spectra was pyruvate, and the pyruvate group peak height was comparable in both spectra.

Figure 3 shows average in-process PS yield, protein/PS ratio, and NA/PS ratio for serotype 4 using the shortened purification process of the invention. Results are shown for each purification step.

Figure 4 shows average in-process PS yield, protein/PS ratio, and NA/PS ratio for serotype 19A using the shortened purification process of the invention. Results are shown for each purification step.

Figure 5 shows average in-process PS yield, protein/PS ratio, and NA/PS ratio for serotype 7F using the shortened purification process of the invention. Results are shown for each purification step.

Figure 6 shows average in-process PS yield, protein/PS ratio, and NA/PS ratio for serotype 6B using the shortened purification process of the invention. Results are shown for each purification step.

Figure 7 shows average in-process PS yield, protein/PS ratio, and NA/PS ratio for serotype 6A using the shortened purification process of the invention. Results are shown for each purification step.

Figure 8 shows average in-process PS yield, protein/PS ratio, and NA/PS ratio for serotype 1 using the shortened purification process of the invention. Results are shown for each purification step.

Figure 9 shows average in-process PS yield, protein/PS ratio, and NA/PS ratio for serotype 14 using the shortened purification process of the invention. Results are shown for each purification step.

Figure 10 shows a comparison of PS in-process yields for serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, and 19F purified using the shortened purification process of the invention.

Figure 11 shows a comparison of protein/PS ratios for serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, and 19F purified using the shortened purification process of the invention. Results are compared for each purification step.

Figure 12 shows a comparison of NA/PS ratios for serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, and 19F purified using the shortened purification process of the invention. Results are compared for each purification step.

Figure 13 shows protein removal efficiency attributable to the acidification step of the shortened purification process of the invention. The difference in protein concentration (SDS-PAGE) before and after acidification is plotted against initial protein concentration before acidification for batches of serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, and 19F. The protein concentration difference divided by the initial protein concentration reflected the protein removal rate by the acidification step.

Figure 14 shows protein removal efficiency attributable to the carbon adsorption step of the shortened purification process of the invention. The amount of protein removed (adsorbed on carbon) was plotted against initial protein loading amounts for batches of serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, and 19F. The amount of protein removed divided by the initial protein loading amounts reflected the protein removal rate by the carbon adsorption step.

DETAILED DESCRIPTION OF THE INVENTION

[0012] The present invention relates to a shortened purification process to reduce the soluble protein levels in cellular *Streptococcus pneumoniae* lysates to produce substantially purified capsular polysaccharides suitable for incorporation into pneumococcal conjugate vaccines. For most of the serotypes for the currently marketed 7-valent pneumococcal conjugate (7vPnC) vaccine (Pneumovax®), as well as the newly developed 13-valent pneumococcal conjugate

(13vPnC) vaccine, the current polysaccharide purification process requires up to sixteen steps. These steps involve many expensive, labor-intensive, and technologically demanding operations, such as chromatography and multiple membrane separations. The process of the present invention eliminates up to eight of these steps while accomplishing the same purification and eliminates the need for chromatography. Thus, the present invention relates to a more efficient purification process that is less expensive, takes less time, and involves fewer steps.

[0013] The shortened purification process of the present invention relates to the discovery that ultrafiltering and diafiltering a clarified cellular *S. pneumoniae* lysate broth, followed by acidification of the concentrated lysate broth with an organic or a mineral acid to a pH of less than 4.5, preferably about 3.5, precipitates at least 98% of the protein in the solution without seriously affecting polysaccharide yield. By ultrafiltering and diafiltering the lysed and clarified fermentation broth before acidification to a pH of less than 4.5, preferably around 3.5, "salting in" effects of proteins are eliminated and the fraction of protein that is "salted out" is increased. "Salting in" refers to increased solubility of proteins while "salting out" refers to precipitation of proteins in solution as they reach their isoelectric points. The ultrafiltration and diafiltration step also prevents the foaming observed when sodium-carbonate treated broth undergoes acidification even to a pH of 5.0 (see U.S. Patent App. Pub. No. 2006/0228381). Thus, ultrafiltration and diafiltration of the clarified lysate broth makes it possible to use any low molecular weight pH titrant such as sodium carbonate during *S. pneumoniae* serotype fermentation and prevents foaming of the clarified lysate broth when acidified to a pH of less than 4.5.

[0014] "Clarified lysate broth" refers to a lysate broth that has undergone centrifugation or filtration to remove cell debris.

[0015] "Diafiltering," "diafiltration," "DF," and like terms refer to, for example, using semi-permeable membranes with appropriate physical and chemical properties to remove small molecules from a solution.

[0016] "Ultrafiltering," "ultrafiltration," "UF," and like terms refer to, for example, using semi-permeable membranes with appropriate physical and chemical properties to discriminate between molecules in a solution and concentrate like molecules into a smaller volume of solution.

[0017] Within the methods of the present invention, ultrafiltration and diafiltration typically comprise "cross-flow" or "tangential-flow" filtration in order to avoid clogging of the filter membranes. In "cross-flow" filtration, the solution to be filtered is passed across the surface of the membrane. Materials which pass through the membrane are referred to as the permeate. The materials which do not pass through the membrane are referred to as the retentate. The retentate is recycled to a feed reservoir to be refiltered.

[0018] As used herein, an organic or a mineral acid is used to lower the pH of the ultrafiltered

and diafiltered lysate broth so long as a pH of less than 4.5, particularly about 3.5, is achieved. Accordingly, both organic and mineral acids may be used within the methods of the invention. As used herein, the term "mineral acid" refers to an acid derived from inorganic mineral by chemical reaction as opposed to organic acids. Exemplary, non-limiting examples of mineral acids that may be used within the methods of the present invention include hydrochloric acid, nitric acid, phosphoric acid, and sulphuric acid. In particular embodiments, the pH of the concentrated lysate broth is lowered to less than 4.4, 4.3, 4.2, 4.1, 4.0, 3.9, 3.8, 3.7, 3.6, 3.5, 3.4, 3.3, 3.2, 3.1, or 3.0. In other embodiments, the pH of the concentrated lysate broth is lowered to about 4.4, about 4.3, about 4.2, about 4.1, about 4.0, about 3.9, about 3.8, about 3.7, about 3.6, about 3.5, about 3.4, about 3.3, about 3.2, about 3.1, about 3.0, about 2.9, about 2.8, about 2.7, about 2.6, about 2.5, about 2.4, about 2.3, about 2.2, about 2.1, or about 2.0.

[0019] The shortened purification process of the present invention also relates to the discovery that, in combination with the concentration and low pH steps described above, filtration using activated carbon precipitates at least 90% of remaining protein without seriously affecting polysaccharide yield. In particular embodiments, carbon filtration using carbon derived from sawdust or other wood products and activated with phosphoric acid was found to be more effective at reducing or removing protein impurities than carbons used within current carbon filtration methods.

[0020] Accordingly, the present invention relates to a process for producing substantially purified capsular polysaccharides from a *Streptococcus pneumoniae* cell lysate comprising the steps of:

1. (a) providing a fermentation broth comprising bacterial cells that produce a selected *Streptococcus pneumoniae* serotype;
2. (b) lysing the bacterial cells in step (a) with a lytic agent, thereby producing a cell lysate comprising cell debris, soluble proteins, nucleic acids, and polysaccharides;
3. (c) clarifying the cell lysate of step (b) using centrifugation or filtration to remove cell debris, thereby producing a clarified cell lysate;
4. (d) ultrafiltering and diafiltering the clarified cell lysate of step (c) to remove low molecular weight impurities and increase polysaccharide concentration, thereby producing a retentate;
5. (e) lowering the pH of the retentate of step (d) to less than 4.5, particularly about 3.5, using an organic or a mineral acid to precipitate protein and nucleic acids, thereby forming an acidified retentate solution;
6. (f) holding the acidified retentate solution formed in step (e) for a time sufficient to allow settling of the precipitate, particularly for at least 2 hours with or without agitation, followed by filtration or centrifugation of the acidified retentate solution, thereby producing a clarified polysaccharide solution;
7. (g) filtering the clarified polysaccharide solution of step (f) through an activated carbon filter, particularly an activated carbon filter comprising wood-based phosphoric acid-activated carbon;

8. (h) ultrafiltering and diafiltering the filtered solution produced by step (g), thereby producing a concentrated purified polysaccharide solution; and
9. (i) filtering the concentrated purified polysaccharide solution produced by step (h) using a sterile filter;

whereby substantially purified capsular polysaccharides in the form of a solution are produced. The sterile filtration of step (i) is useful to remove bacteria and particles from the concentrated purified polysaccharide solution. Exemplary, non-limiting *S. pneumoniae* serotypes selected for this embodiment of the invention are 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F. In a particular embodiment, step (e) removes at least 98% of protein from the retentate of step (d). In another embodiment, step (g) removes at least 90% of the protein from the clarified polysaccharide solution of step (f). In another embodiment, the diafiltration of step (h) comprises a pH adjustment to between about 5.5 to about 7.5. For improved stability of the substantially purified capsular polysaccharide during long-term storage, however, the diafiltration of step (h) comprises a pH adjustment to between about 7.0 to about 7.5, and more particularly to about 7.4.

[0021] As used herein, the term "substantially purified capsular polysaccharide-containing lysate" or "solution containing substantially purified capsular polysaccharides" refers to a cellular *Streptococcus pneumoniae* lysate or solution from which protein has been removed such that the percent ratio of protein to polysaccharide (protein/PS) is less than 10%, less than 9%, less than 8%, less than 7%, less than 6%, less than 5%, less than 4%, less than 3%, less than 2%, or less than 1% and the percent ratio of nucleic acid to polysaccharide (NA/PS) is less than 5%, less than 4%, less than 3%, less than 2%, or less than 1%. In particular embodiments, the percent ratios of protein/PS and NA/PS for substantially purified capsular polysaccharide-containing lysates or solutions comprising specific serotypes are as follows: for serotype 1 the ratio of protein/PS is less than 2% and the ratio of NA/PS is less than 2%, for serotype 4 the ratio of protein/PS is less than 3% and the ratio of NA/PS is less than 2%, for serotype 5 the ratio of protein/PS is less than or equal to 7.5% and the ratio of NA/PS is less than or equal to 2%, for serotype 6A the ratio of protein/PS is less than 2% and the ratio of NA/PS is less than 2%, for serotype 6B the ratio of protein/PS is less than 4% and the ratio of NA/PS is less than 1%, for serotype 7F the ratio of protein/PS is less than 5% and the ratio of NA/PS is less than 2%, for serotype 9V the ratio of protein/PS is less than 2% and the ratio of NA/PS is less than 1%, for serotype 14 the ratio of protein/PS is less than 3% and the ratio of NA/PS is less than 2%, for serotype 18C the ratio of protein/PS is less than 2% and the ratio of NA/PS is less than 2%, for serotype 19A the ratio of protein/PS is less than 2% and the ratio of NA/PS is less than 2%, for serotype 19F the ratio of protein/PS is less than 3% and the ratio of NA/PS is less than 2%, and for serotype 23F the ratio of protein/PS is less than 2% and the ratio of NA/PS is less than 2%. Methods for the quantification of protein, polysaccharide, and nucleic acid concentrations in a cellular lysate or solution are well known in the art and include, for example, SDS-PAGE (Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis) analysis, HPLC (High Performance Liquid Chromatography) and SEC (Size Exclusion Chromatography), modified Lowry assays, spectrophotometry, SEC-MALLS (Size-Exclusion Chromatography/Multi-Angle Laser Light Scattering), and NMR (Nuclear Magnetic

Resonance).

[0022] Within the methods of the present invention, the bacterial cells may be lysed using any lytic agent. A "lytic agent" is any agent that aids in cell wall breakdown and release of autolysin which causes cellular lysis including, for example, detergents. As used herein, the term "detergent" refers to any anionic or cationic detergent capable of inducing lysis of bacterial cells. Representative examples of such detergents for use within the methods of the present invention include deoxycholate sodium (DOC), N-lauryl sarcosine (NLS), chenodeoxycholic acid sodium, and saponins.

[0023] In one embodiment of the present invention, the lytic agent used for lysing bacterial cells is DOC. DOC is the sodium salt of the bile acid deoxycholic acid, which is commonly derived from biological sources such as cows or oxen. DOC activates the LytA protein, which is an autolysin that is involved in cell wall growth and division in *Streptococcus pneumoniae*. The LytA protein has choline binding domains in its C-terminal portion, and mutations of the *lytA* gene are known to produce LytA mutants that are resistant to lysis with DOC.

[0024] Although there is no evidence that the use of DOC during *Streptococcus pneumoniae* polysaccharide purification poses a health risk, the use of such biologically derived reagents could raise potential regulatory concerns. Accordingly, in one embodiment of the present invention, the lytic agent used for lysing bacterial cells is a non-animal derived lytic agent. Non-animal derived lytic agents for use within the methods of the present invention include agents from non-animal sources with modes of action similar to that of DOC (*i.e.*, that affect LytA function and result in lysis of *Streptococcus pneumoniae* cells). Such non-animal derived lytic agents include, but are not limited to, analogs of DOC, surfactants, detergents, and structural analogs of choline, and may be determined using procedures as described in the Experimental section herein below. In one embodiment, the non-animal derived lytic agent is selected from the group consisting of decanesulfonic acid, tert-octylphenoxy poly(oxyethylene)ethanols (*e.g.* Igepal® CA-630, CAS #: 9002-93-1, available from Sigma Aldrich, St. Louis, MO), octylphenol ethylene oxide condensates (*e.g.* Triton® X-100, available from Sigma Aldrich, St. Louis, MO), N-lauryl sarcosine sodium (NLS), lauryl iminodipropionate, sodium dodecyl sulfate, chenodeoxycholate, hyodeoxycholate, glycodeoxycholate, taurodeoxycholate, taurochenodeoxycholate, and cholate. In another embodiment, the non-animal derived lytic agent is NLS.

[0025] The present invention also relates to serotype-specific modifications to the process described above. For example, because the serotype 19A polysaccharide is unstable and its molecular weight changes during purification, it was discovered that modifications to the process described were useful in stabilizing the 19A polysaccharide. These modifications included carrying out the ultrafiltration and diafiltration step prior to acidification at about 4 °C at a pH of about 6 in sodium phosphate buffer, holding the acidified retentate solution for at least 2 hours at about 4 °C to allow settling of the precipitate, and adjusting the pH of the clarified polysaccharide solution to 6 prior to the activated carbon filtration step. By contrast, it was discovered that for serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F, and 23F, less

polysaccharide loss and more protein removal was achieved when the ultrafiltration and diafiltration step prior to acidification was carried out in salt free media such as water, and this step could be carried out at room temperature at neutral pH.

[0026] Accordingly, the present invention also relates to a process for producing substantially purified capsular polysaccharides from a *Streptococcus pneumoniae* cell lysate comprising serotype 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F, or 23F comprising the steps of:

1. (a) providing a fermentation broth comprising bacterial cells that produce *Streptococcus pneumoniae* serotype 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F, or 23F;
2. (b) lysing the bacterial cells in step (a) with a detergent, thereby producing a cell lysate comprising cell debris, soluble proteins, nucleic acids, and polysaccharides;
3. (c) clarifying the cell lysate of step (b) using centrifugation or filtration to remove cell debris, thereby producing a clarified cell lysate;
4. (d) ultrafiltering and diafiltering the clarified cell lysate of step (c) at room temperature at neutral pH in salt free media to remove low molecular weight impurities and increase polysaccharide concentration, thereby producing a salt free retentate;
5. (e) lowering the pH of the salt free retentate of step (d) to less than 4.5, particularly about 3.5, using an organic or a mineral acid to precipitate protein and nucleic acids, thereby forming an acidified retentate solution;
6. (f) holding the acidified retentate solution formed in step (e) for at least 2 hours at room temperature with or without agitation to allow settling of the precipitate, followed by filtration or centrifugation of the acidified retentate solution, thereby producing a clarified polysaccharide solution;
7. (g) filtering the clarified polysaccharide solution of step (f) through an activated carbon filter, particularly an activated carbon filter comprising wood-based phosphoric acid-activated carbon;
8. (h) ultrafiltering and diafiltering the filtered solution produced by step (g), thereby producing a concentrated purified polysaccharide solution; and
9. (i) filtering the concentrated purified polysaccharide solution produced by step (h) using a sterile filter;

whereby substantially purified capsular polysaccharides comprising serotype 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F, or 23F in the form of a solution are produced. In a particular embodiment, step (e) removes at least 98% of protein from the salt free retentate of step (d). In another embodiment, step (g) removes at least 90% of the protein from the clarified polysaccharide solution of step (f). In another embodiment, the diafiltration of step (h) comprises a pH adjustment to between about 5.5 to about 7.5. For improved stability of the substantially purified capsular polysaccharide during long-term storage, however, the diafiltration of step (h) comprises a pH adjustment to between about 7.0 to about 7.5, and more particularly to about 7.4.

[0027] The present invention also relates to a process for producing substantially purified capsular polysaccharides from a *Streptococcus pneumoniae* cell lysate comprising serotype 19A comprising the steps of

1. (a) providing a fermentation broth comprising bacterial cells that produce *Streptococcus pneumoniae* serotype 19A;
2. (b) lysing the bacterial cells in step (a) with a detergent, thereby producing a cell lysate comprising cell debris, soluble proteins, nucleic acids, and polysaccharides;
3. (c) clarifying the cell lysate of step (b) using centrifugation or filtration to remove cell debris, thereby producing a clarified cell lysate;
4. (d) ultrafiltering and diafiltering the clarified cell lysate of step (c) at about 4 °C at a pH of about 6 in sodium phosphate buffer, 25 mM sodium phosphate, to remove low molecular weight impurities and increase polysaccharide concentration, thereby producing a retentate;
5. (e) lowering the pH of the retentate of step (d) to less than 4.5, particularly about 3.5, using an organic or a mineral acid to precipitate protein and nucleic acids, thereby forming an acidified retentate solution;
6. (f) holding the acidified retentate solution formed in step (e) for at least 2 hours at about 4 °C with or without agitation to allow settling of the precipitate, followed by filtration or centrifugation of the acidified retentate solution, thereby producing a clarified polysaccharide solution;
7. (g) adjusting the pH of the clarified polysaccharide solution of step (f) to about 6, thereby producing a pH-adjusted clarified polysaccharide solution;
8. (h) filtering the pH-adjusted clarified polysaccharide solution of step (g) through an activated carbon filter, particularly an activated carbon filter comprising wood-based phosphoric acid-activated carbon;
9. (i) ultrafiltering and diafiltering the filtered solution produced by step (h), thereby producing a concentrated purified polysaccharide solution; and
10. (j) filtering the concentrated purified polysaccharide solution produced by step (i) using a sterile filter;

whereby substantially purified capsular polysaccharides comprising serotype 19A in the form of a solution are produced. In a particular embodiment, step (e) removes at least 98% of protein from the retentate of step (d). In another embodiment, step (h) removes at least 90% of the protein from the pH-adjusted clarified polysaccharide solution of step (g). In another embodiment, the diafiltration of step (i) comprises a pH adjustment to between about 5.5 to about 7.5. For improved stability of the substantially purified 19A polysaccharide during long-term storage, however, the diafiltration of step (i) comprises a pH adjustment to between about 7.0 to about 7.5, and more particularly to about 7.4.

[0028] The following examples are offered by way of illustration and not by way of limitation.

EXPERIMENTAL

[0029] The following Examples present results for *S. pneumoniae* polysaccharide serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, and 19F purified at 10L scale using the improved process of

the present invention, and results are compared with those of the current purification process.

Example 1. Shortened Purification Process for *S. pneumoniae* Polysaccharide Serotypes 1, 4, 5, 6A, 6B, 7F, 14 and 19A.

[0030] *S. pneumoniae* fermentation broths lysed with deoxycholate sodium (DOC) were obtained either within two days following their harvest or stored in 4°C and processed within the following week. As described below, the present purification process included the following changes compared to the existing purification process: 1) the acidification step was moved from the beginning to after the first ultrafiltration/diafiltration (UF/DF) step, and the pH was adjusted to 3.5 instead of 5; 2) the diafiltration buffer was changed from 0.025 M phosphate to de-ionized (DI) water; 3) carbon adsorption was changed to 2 CUNO R32SP carbon disks (CUNO Inc., Wayne, NJ) using wood-based phosphoric acid-activated carbon, and the adsorption time was extended from 4 to 12 turnovers (each turnover was 22 minutes); and 4) pH was adjusted to 7.4 during the last 30K diafiltration step when diafiltered about 5 times. The same purification procedure was applied to serotypes 1, 4, 5, 6A, 6B, or 7F. For serotype 19A, the steps were modified further and the purification was conducted in a chill room, as described below.

Purification Steps

[0031] All steps were conducted at room temperature except for type 19A, in which case the process was conducted at 4°C in a chill room.

[0032] Clarification of the lysate: The purpose of this step was to remove cell debris and clarify the broth. This was accomplished either by centrifugation or filtration. The broth was centrifuged at 10,000 g for 30 min or until the broth was clear at 20°C (4°C for type 19A), or filtered with a Millipore Prefilter (Millipore Corp., Billerica, MA) with the addition of Celpure® filter aids (Advanced Minerals, Santa Barbara, CA). The clarified lysate was collected for further processing and the pellets were discarded.

[0033] First UF/DF (Ultrafiltration/Diafiltration): This step provided volume reduction and buffer exchange and also removed low molecular weight impurities. The clarified lysate was concentrated to about 1/8th original volume. Diafiltration was performed using about 10 volumes DI water (pH 6, 25 mM phosphate for 19A).

[0034] Acidification: More than 98% of the proteins were removed in this step. While stirring, concentrated phosphoric acid was added carefully to the retentate. The pH of the retentate was adjusted to a target value of pH 3.5. The acidified retentate was stirred for half an hour and aged in room temperature to age overnight (2 hours at 4°C for 19A), resulting in the precipitation of protein and nucleic acids.

[0035] Clarification of Acidified Retentate: This was the clarification step to remove the precipitates after acidification. The slurry of acidified solution was centrifuged in a rotor at 10,000 rpm (17,000 Relative Centrifugal Force or RCF) for one hour at 20°C (except for 6B, in which case the centrifugation was 6 hours at 37°C). The supernatant was collected and the pellet was discarded. Depth filtration with a Millipore Prefilter (Millipore Corp., Billerica, MA) with the addition of Celpure® filter aids (Advanced Minerals, Santa Barbara, CA) can also be used for this step.

[0036] Carbon Adsorption: In most cases, there was a slight yellow color after centrifugation of the acidified 100K retentate. The color removal was achieved by the carbon adsorption using wood-based phosphoric acid-activated carbon. This step also removed residual protein that remained after acidification. The clarified polysaccharide solution was recirculated through the carbon filter for 5-6 hours or overnight (for 19A, the pH was adjusted to 6 before carbon adsorption).

[0037] Final 30K UF/DF: This was another concentration and buffer exchange step to concentrate the solution to a final polysaccharide (PS) concentration of >2 g/L, which was diafiltered into deionized (DI) water. The carbon-filtered PS solution was concentrated. Then the concentrate was dialfiltered 10X with DI water. The pH was adjusted to 7.4 during the diafiltration.

[0038] Final 0.2 µm Sterile Filtration: The final PS solution was sterile filtered with a 0.22 µm filter or sterile disposable filter unit and stored in a 4°C refrigerator.

Analytical Methods

[0039] Quantification of protein, PS, and nucleic acid concentrations were carried out using methods that are well known in the art, including SDS-PAGE (Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis) analysis, HPLC (High Performance Liquid Chromatography) and SEC (Size Exclusion Chromatography), modified Lowry assays, spectrophotometry, SEC-MALLS (Size-Exclusion Chromatography/Multi-Angle Laser Light Scattering), and NMR (Nuclear Magnetic Resonance).

Results and Discussion

[0040] Type 5 Shortened Purification Batches: The comparison of final PS yield, PS molecular weight and major impurity levels of three purification batches using the shortened process and those using the current process for Type 5 are shown in Table 1. All the starting broths were DOC lysed high cell density fermentation batches, which had a higher polysaccharide concentration (~0.5 g/L) than the standard SOP fermentation broth (~0.3 g/L). A 30K or 50K membrane was used for the first UF/DF step. The impurity levels were calculated using

impurity/PS ratios. The same approach was used for all other serotypes described herein.

Table 1. Testing Data for Type 5 Shortened Purification Process Batches.

BATCH #	PS YIELD (%)	PROTEIN RATIO (%)	NA RATIO (%)	PS MOLECULAR (KG/MOL)	C-PS RATIO (%)
L29276-27 (30K UF/DF)	56.7	1.2	0.12	299	20.7
L29276-32 (30K UF/DF)	54.6	1.8	0.12	282	24.7
L29276-52 (50K UF/DF)	60.2	1.4	0.03	275	22.8
Current process	57	3.6	0.18	320	15.9
Specification or expected value	50-60	≤7.5	≤2.0	NA	≤35

[0041] As the data in Table 1 show, the protein ratios of all three batches using the shortened purification process met the specification of ≤7.5%, and were also comparable to that of the current process. The nucleic acid (NA) as well as C-polysaccharide (C-PS) ratios were also well below the specifications of ≤2.0% and ≤35%, respectively and were comparable to that of the current process. The results of the final PS yield and impurity levels of the three batches shown in Table 1 demonstrate the reproducibility and robustness of the shortened process.

[0042] There was some concern about whether polysaccharides could be hydrolyzed at a lower pH of 3.5. PS retention time change during the purification process was therefore monitored, and the molecular weight of the final purified PS was measured. No significant changes in serotype 5 PS retention time from the HPLC chromatogram were observed. There were also no significant differences in molecular weights for the purified PS by the shortened process and that of the current process (284kg/mol) based on the MALLS measurement. Thus it was concluded that the molecular weight of purified PS was not adversely affected by the shortened purification process.

[0043] The in-process polysaccharide yield, protein/polysaccharide ratio and nucleic acid/polysaccharide ratio at each of the processing steps for the three shortened process batches are summarized in Table 2.

Table 2. Type 5 In-Process PS yield, Protein (SDS-PAGE) and Nucleic Acid Ratios.

STEP	PS YIELD (%)			PROTEIN/PS (%)		
	L29276-27	L29276-32	L29276-52	L29276-27	L29276-32	L29276-52
Broth	100.0	100.0	100.0	384.40	1026.60	193.26
Centrifugation	99.9	95.6	94.6	274.60	642.00	209.78
30/50K UF/DF	71.1	82.8	93.9	152.10	561.00	204.76
Acidification	56.1	65.9	58.6	9.00	7.50	0.20
Carbon	57.7	55.7	58.7	0.20	1.00	0.50

Carbon	27.7	22.7	20.7	0.20	1.20	0.09
30K UF/DF	54.5	54.6	60.3	0.10	1.00	0.09

Table 2, Continued.			
STEP	NUCLEIC ACID/PS (%)		
	L29276-27	L29276-32	L29276-52
Broth	825.80	579.83	344.70
Centrifugation	505.10	484.30	311.90
30/50K UF/DF	107.30	139.90	137.60
Acidification	9.44	16.40	7.38
Carbon	0.22	0.30	0.74
30K UF/DF	0.11	0.12	0.19

[0044] There is always a loss of product during each step of any purification process. For these three shortened process batches, PS loss occurred mostly in the first UF/DF step and the acidification step. The loss of PS at the first UF/DF step was due to adsorption of PS or PS-protein complex to the surface of the membrane. This loss was minimized by rinsing the retentate side of the membrane with DI water after the diafiltration and combining the rinse with the original retentate. The loss of PS at the acidification step could be due to two reasons: physical adsorption of PS to the precipitation solids, and polysaccharide binding to the protein with co-precipitation during acidification. This second possibility was further investigated and results showed evidence of PS/protein binding.

[0045] Figure 1 shows the reduction of protein/PS ratio at each purification step. Although the centrifugation step removed a small percentage of proteins, the majority of protein was removed at the acidification step. Only a trace amount of protein was detected after pH 3.5 treatment even for the highest protein potency batch.

[0046] Similar to protein/PS ratio, the nucleic acid/PS ratio showed variability of impurity levels among batches. The 30/50K UF/DF step removed a significant amount of nucleic acid as compared to protein removal in the same processing step. Without being bound by theory, this may have been due to the molecular size of nucleic acids being smaller than that of proteins, making them relatively easier to be removed via the 30/50K UF/DF step. The first centrifugation and acidification step also removed a considerable amount of nucleic acid.

[0047] Table 2 shows that the activated carbon adsorption step also reduces protein/PS and NA/PS level. The percentage reduction was not as significant as the first two steps, but this step was important for removing the color of the solution and ensured that the impurity level met the specification.

[0048] Type 4 Shortened Purification Batches: A summary of three shortened purification batches for Type 4 is shown in Table 3. The feed broths for all three batches were DOC-lysed.

Table 3. Type 4 Shortened Purification Batches Summary.

BATCH	PS YIELD (%)	PROTEIN/PS (%)	NUCLEIC ACID/PS (%)	PS MOLECULAR WEIGHT (KG/MOL)	C-PS RATIO (%)
L29276-47	56.3	0.7	0.04	289	15.4
L29276-55	53.0	1.3	0.03	354	14.7
L29276-148	57.3	1.16	0.02	293	10.0
Current process	71.1	0.05	0.02	285	
Specification	NA	<3.0	<2.0	>350	<25

[0049] The type 4 PS yield was also between 50-60%. Protein/PS ratio and nucleic acid/PS ratio were well within their specifications. C-PS ratio was also well within the specification. The molecular weight of all of the three batches were close to ~300 kg/mol. Serotype 4 PS purified by the current purification process using the similar fermentation broth also gave a lower molecular weight 285 kg/mol. Comparison of the HPLC chromatographs of the PS from fermentation broth and the final purified solution showed that there were no differences in the PS retention time. This suggested that the difference in molecular weight difference was not caused by the process change, but rather due to intrinsic nature of the fermentation process.

[0050] The Type 4 PS contains a pyruvate group in the purified molecule, and this pyruvate was important for conjugation for use in a pneumococcal conjugate vaccine. To ensure that the pyruvate amount was not adversely affected by acid treatment, NMR analysis was conducted. Figure 2 shows the NMR spectra for standard type 4 PS and that from batch L29276-47. No significant differences between the two spectra were observed. The second peak from the right in both spectra was pyruvate, and the pyruvate group peak height was comparable in both spectra. The pyruvate ratios for all three shortened process batches were 0.8 mol/mol, and met the specification of >0.7 mol/mol.

[0051] The in-process PS yield, protein/PS and NA/PS ratio for the three type 4 batches are summarized in Table 4. PS loss occurred mostly at the first centrifugation, acidification and activated carbon adsorption steps, with an average of 10%, 8% and 20%, respectively. The overall PS yield was close to that of type 5, around 55%.

Table 4. Type 4 In-Process PS Yield, Protein (SDS-PAGE)/PS Ratio, and NA/PS Ratio.

STEP	PS YIELD (%)			PROTEIN/PS (%)		
	L29276-47	L29276-55	L29276-148	L29276-47	L29276-55	L29276-148
Broth	100.0	100.0	100.0	231.22	297.33	499.75
Centrifugation	85.8	97.8	88.0	198.69	247.40	534.61
50/100K UF/DF	85.1	86.4	87.7	235.90	249.24	364.98
Acidification	84.2	72.0	78.6	0.42	1.81	0.41
Carbon	65.3	48.9	58.6	0.09	0.39	0.34

30K UF/DF	57.9	50.6	57.4	0.08	0.09	1.31
-----------	------	------	------	------	------	------

Table 4, Continued.			
STEP	NA/PS (%)		
	L29276-47	L29276-55	L29276-148
Broth	312.84	282.50	302.71
Centrifugation	301.40	261.82	243.28
50/100K UF/DF	80.99	67.67	92.00
Acidification	4.24	5.01	1.17
Carbon	0.08	0.92	0.38
30K UF/DF	0.05	0.06	0.07

[0052] Figure 3 shows average PS yield, protein/PS and NA/PS ratio change at each of the purification steps for the three batches in Table 4. The protein removal occurred mostly at the acidification step as expected. The first centrifugation and UF/DF steps also collectively removed a certain amount of protein, but the protein reduction was less than the acidification step.

[0053] Similar to type 5, the most nucleic acid/PS ratio reduction took place at the 50/100K UF/DF, first centrifugation, and the acidification steps, and the NA reduction at the first UF/DF step was more significant than that for proteins. Again, as shown in Figure 3, the activated carbon step removed a certain amount of protein and NA and brought the impurity levels below the specifications. It also removed the color of the solution as well.

[0054] Type 19A Shortened Purification Batches: Type 19A polysaccharide is unstable and the molecular weight changes during purification. The shortened purification process was modified slightly in order to stabilize the 19A polysaccharide. These modifications are summarized as follows: 1) the purification steps were mostly conducted at 4°C in the chill room; 2) the first 100K diafiltration was chilled using 25 mM phosphate buffer (4°C) with a pH of 6 instead of using room temperature water; 3) the acidification holding time was reduced from overnight to 2 hours; and 4) after clarifying the acidified 100K retentate, the pH was adjusted to 6 and activated carbon adsorption was conducted at pH 6 instead of 3.5.

[0055] The results of two 19A batches purified by the shortened purification process are shown in Table 5.

Table 5. Type 19A Shortened Purification Batches Summary.					
BATCH	PS YIELD (%)	PROTEIN/PS (%)	NA/PS (%)	MOLECULAR WEIGHT (KG/MOL)	C-PS (%)
L29276-116	61.58	0.14	0.01	525	3.8
L29276-143	76.15	1.43	0.01	488	2.8
Specification	NA	<2	<2	NA	<10

[0056] The PS yields of the two shortened purification batches were 62 and 76% respectively. Final protein/PS ratio, nucleic acid ratio, and C-PS ratio all met their respective specifications. The final molecular weights of polysaccharides from the two batches were 525 kg/mol and 488 kg/mol, respectively, and were close to that of the PS of 19A used in phase III clinical trials (486kg/mol).

[0057] The protein/PS ratio for the two batches both met the specification of <2%. Both NA and C-PS ratio of the two batches were well within their specifications.

[0058] The PS yield, protein and NA reduction at each of the purification steps are shown in Table 6 and Figure 4. The results showed PS loss at each of the purification steps except the first centrifugation step. Like serotypes 5 and 4, protein and NA removal mostly took place at the first three steps, and there was hardly any detectable protein and NA left after acidification. Although the activated carbon adsorption step did not remove a significant amount of protein and NA, possibly due to very low protein and NA concentration after the acidification, the step was still needed for color removal.

Table 6. Type 19A In-Process PS Yield, Protein/PS and NA/PS Ratio.						
STEP	PS YIELD (%)		PROTEIN/PS (%)		NA/PS (%)	
	L29276-116	L29276-143	L29276-116	L29276-143	L29276-116	L29276-143
Broth	100.0	100.0	224.41	144.08	277.55	134.73
Centrate	100.8	101.2	121.60	119.11	189.64	100.64
100K UF/DF	87.7	90.3	103.83	105.16	56.87	26.36
Acidification	90.4	72.1	0.28	0.84	0.03	0.03
Carbon	77.8	71.7	0.96	0.03	0.04	0.02
30K UF/DF	61.0	76.1	0.54	0.00	0.02	0.01

[0059] Type 7F Shortened Purification Batches: Type 7F is a non-ionic polysaccharide, which typically requires change in steps during purification using the existing process compared to the serotypes described above). However, the shortened purification process of the present invention was successfully applied to serotype 7F without the need for process deviation. Two batches of type 7F broth were purified using the shortened process. One was a standard fermentation broth (L29276-107) and one was a high cell density broth (L29276-157). The results of the two batches are summarized in Table 7.

Table 7. Type 7F Shortened Purification Batches Summary.					
BATCH #	PS YIELD (%)	PROTEIN/PS (%)	NA/PS (%)	MOLECULAR WEIGHT (KG/MOL)	C-PS (%)
L29276-107	65.27	0.49	0.02	968	3.3
L29276-157	79.06	0.26	0.04	881	2.9

Table 7. Type 7F Shortened Purification Batches Summary.

BATCH #	PS YIELD (%)	PROTEIN/PS (%)	NA/PS (%)	MOLECULAR WEIGHT (KG/MOL)	C-PS (%)
Specification	NA	<5	<2	NA	<17

[0060] The PS yield of type 7F was actually higher than the other serotypes, possibly due to less binding of the non-ionic polymer to the charged protein molecules. Final protein, NA and C-PS ratios were all well within their specifications. Molecular weight of type 7F was comparable to that of standard batches and even though the 7F molecular weight was quite high, the PS solution was not very viscous due to smaller excluded volume of the nonionic polymer.

[0061] Type 7F PS yield, protein and NA ratio at each of the purification steps are shown in Table 8 and the average of the two batches was plotted in Figure 5.

Table 8. Type 7F In-Process PS Yield, Protein/PS, and NA/PS Ratio.

STEP	PS YIELD (%)		PROTEIN/ PS (%)		NUCLEIC ACID/PS (%)	
	L29276-107	L29276-157	L29276-107	L29276-157	L29276-107	L29276-157
Broth	100.00	100.00	274.10	189.24	258.55	134.64
Centrifugation	100.31	95.57	110.60	107.20	163.38	83.14
100K UF/DF	84.19	95.39	139.38	27.53	54.02	32.74
Acidification	70.89	84.08	0.44	0.10	1.80	0.94
Carbon	63.96	79.40	0.70	0.09	2.29	0.23
30K UF/DF	57.24	77.70	0.30	0.05	0.06	0.00

[0062] There was some Type 7F PS loss at each purification step. Overall, PS loss was less than that of serotypes 5 and 4. Protein and NA ratio reduction was mostly by the first centrifugation, 100K UF/DF, and acidification steps. Similar to other serotypes, the 100K UF/DF was more efficient in removing NA than proteins. Although the activated carbon adsorption step did not remove a significant amount of protein and NA due to very low impurity levels after acidification, the step was still needed for color removal.

[0063] Type 6B Shortened Purification Batches: Two batches of 6B were purified using the shortened purification process. It was found the clarification of the acidified 100K retentate took a longer time than the other serotypes (6 hours instead of 1 hour). Except for this difference, the purification process was similar to that of the serotypes. The results of the two batches are summarized in Table 9.

Table 9. Type 6B Shortened Purification Summary.

BATCH	PS YIELD (%)	PROTEIN/PS (%)	NA/PS (%)	MOLECULAR WEIGHT (KG/MOL)	C-PS (%)
L29276-161	69.50	1.73	0.28	857	3.6
L29276-164	74.50	1.41	0.17	1142	3.9
Specification	NA	<4	<1	>800	<10

[0064] All impurity levels (protein, NA, and C-PS) were well within their specifications. The PS yield was relatively high, and so were the protein and NA ratio compared to the other serotypes.

[0065] The in-process PS yield, protein and NA ratios at each of the purification steps are shown in Table 10 and Figure 6.

Table 10. Type 6B Average In-Process PS Yield, Protein, and NA Ratio.

STEP	PS YIELD (%)		PROTEIN/PS (%)		NUCLEIC ACID/PS (%)	
	L29276-161	L29276-164	L29276-161	L29276-164	L29276-161	L29276-164
Broth	100.00	100.00	238.20	288.80	215.12	237.67
Centrifugation	101.43	116.54	95.88	123.59	104.08	104.47
100K UF/DF	104.11	105.47	68.15	88.14	22.13	47.81
Acidification	91.98	101.45	0.71	0.43	1.06	2.24
Carbon	80.71	91.42	0.39	0.23	0.45	0.74
30K UF/DF	72.67	77.28	1.23	0.46	0.37	0.25

[0066] Similar to 19A, PS loss occurred at each of the purification steps except the first centrifugation step, in which there was a slight increase of PS. Removal of protein and NA was achieved mostly by the first centrifugation, the first 100K UF/DF, and acidification steps. However, there was also some reduction of protein and NA ratio at the activated carbon adsorption step.

[0067] Type 6A Shortened Purification Batches: Two batches of type 6A were purified using the shortened purification process. The PS yield, impurity levels and molecular weight of the final solutions are summarized in Table 11.

Table 11. Type 6A Shortened Purification Batches Summary.

BATCH	BROTH	PS YIELD (%)	PROTEIN/PS (%)	NA/PS (%)	MOLECULAR WEIGHT (KG/MOL)	C-PS (%)
L29276-138	Standard	75.75	1.23	0.04	640	7.3
L29276-183	High cell density	72.56	0.20	0.01	670	5.1
Specification		NA	<2.00	<2.00	NA	<15

[0068] The final PS yields of the two 6A batches were both >70%, which were the highest among the serotypes processed using the shortened process. Protein, NA, and C-PS ratios were all within specifications.

[0069] Table 12 and Figure 7 show in-process PS yield, protein and NA ratio change at each of the purification steps. There was hardly any loss of PS at the first centrifugation and 100K UF/DF steps. About 10-15% PS loss occurred at the acidification and activated carbon adsorption steps, which was close to that of the other serotypes. The most protein and NA ratio reduction was by the first centrifugation, 100K UF/DF, and acidification. After acidification, both protein and NA ratios were already below their specifications. For protein, the acidification was the most efficient step, while for NA, 100K UF/DF reduced the most NA/PS ratio. Although the activated carbon adsorption step did not remove a significant amount of protein and NA due to very low impurity levels after acidification, the step was still needed for color removal.

Table 12. Type 6A In-Process PS Yield, Protein and NA Ratio.

STEP	PS YIELD (%)		PROTEIN/PS (%)		NUCLEIC ACID/PS (%)	
	L29276-138	L29276-183	L29276-138	L29276-183	L29276-138	L29276-183
Broth	100.00	100.00	214.35	138.71	211.45	163.88
Centrifugation	99.35	97.80	183.43	149.71	177.03	150.41
100K UF/DF	93.19	100.40	114.72	103.00	43.38	40.43
Acidification	89.87	87.90	0.16	0.87	1.09	0.97
Carbon	76.02	75.70	0.62	0.85	0.09	0.07
30K UF/DF	75.57	72.90	0.00	0.07	0.04	0.02

[0070] Type 1 Shortened Purification Batches: Two batches of type 1 purified by the shortened process are summarized in Table 13. Batch L29276-170 was purified from high cell density fermentation broth and L29276-173 was from standard fermentation broth.

Table 13. Type 1 Shortened Purification Batch Summary.

BATCH	BROTH	PS YIELD (%)	PROTEIN/PS (%)	NA/PS (%)	MOLECULAR WEIGHT (KG/MOL)	C-PS (%)
L29276-170	High cell density	50.96	0.44	0.08	501	5.3
L29276-173	SOP	53.84	1.23	0.01	458	11.9
Specification		NA	<2	<2	NA	<15

[0071] The PS yields of both batches were about 50%, and the impurity levels for protein, NA and C-PS were all within their specifications.

[0072] The in-process PS yield, protein and NA ratios after each of the purification steps are shown in Table 14 and Figure 8. The trends were similar to other serotypes purified. PS loss occurred mostly at the acidification and activated carbon adsorption steps, and most of protein and NA removal occurred at the first three steps. More NA was removed at the 100K UF/DF step than proteins. Although the activated carbon adsorption step did not remove a significant amount of protein and NA due to very low impurity levels after acidification, the step was still needed for color removal.

Table 14. Type 1 Average In-Process PS Yield, Protein and NA Ratio.

STEP	PS YIELD (%)		PROTEIN/PS (%)		NA/PS (%)	
	L29276-170	L29276-173	L29276-170	L29276-173	L29276-170	L29276-173
Broth	100.00	100.00	453.30	595.20	594.41	1178.98
Centrifugation	93.99	105.63	301.86	396.97	340.55	1018.47
50/100K UF/DF	88.33	105.62	232.34	235.08	17.49	182.56
Acidification	66.54	87.67	0.54	4.87	1.09	5.85
Carbon	55.85	59.69	1.91	1.86	0.49	0.01
30K UF/DF	51.24	55.59	0.36	0.92	1.65	0.01

[0073] Type 14 Shortened Purification Batches: Two batches of serotype 14 were purified using the shortened purification process with no process deviations. The final PS yield, and impurity levels are summarized in Table 15.

Table 15. Type 14 Shortened Purification Batches Summary.

BATCH	BROTH	PS YIELD (%)	PROTEIN/PS (%)	NA/PS (%)	MOLECULAR WEIGHT (KG/MOL)	C-PS (%)
L32874-155	High cell density	51.3	2.06	0.03	520	4.2
L32874-163	SOP	56.7	1.39	0.03	662	2.9
Specification		NA	<3	<2	>400	<15

[0074] Similar to serotype 7F, serotype 14 is a non-ionic polysaccharide, and its current purification process is slightly different from the other serotypes. However, the shortened purification process of the present invention was successfully applied to serotype 14 without the need for such additional steps. The PS yields of the two shortened purification process batches were 50-60%, and protein and NA ratios were within their specifications. The molecular weights of the purified PS also met the specification.

[0075] In-process PS yield, protein and NA ratios are summarized in Table 16 and Figure 9. Similar trends of PS loss, protein and NA removal were observed as for the other tested serotypes described above.

Table 16. Type 14 Average In-Process PS Yield, Protein, and NA Ratio.

STEP	PS YIELD (%)		PROTEIN/PS (%)	
	L32874-155	L232874-163	L32874-155	L232874-163
Broth	100.00	100.00	109.73	286.56
Centrifugation	91.40	90.80	110.88	201.48
50/100K UF/DF	82.40	68.70	45.52	165.69
Acidification	65.40	93.90	0.65	0.47
Carbon	59.40	74.60	0.60	0.10
30K UF/DF	53.10	57.60	0.57	0.41

Table 16, Continued.

STEP	NA/PS (%)	
	L32874-155	L232874-163
Broth	192.80	272.50
Centrifugation	135.50	201.82
50/100K UF/DF	47.10	64.54
Acidification	0.89	1.08
Carbon	0.17	0.21
30K UF/DF	0.13	0.25

Example 2. Comparison of Different Serotypes (Serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, and 19F).

[0076] To compare purification of the different serotypes using the shortened purification process of the present invention, PS removal for all the serotypes described in Example 1 plus serotypes 9V, 18C, and 19F were plotted in one graph in Figure 10. Most of the serotypes followed a similar trend with PS loss at each of the purification steps. There was an increase of PS yield for type 6B at the first centrifugation step and type 14 at the acidification step. PS percentage loss varied from serotype to serotype. Type 5 seemed to lose most PS at the acidification step and type 4 at the activated carbon adsorption step.

[0077] Protein/PS ratios for each purification step for each of the serotypes are shown in Figure 11. The figure shows that there was a difference in initial protein/PS ratios for different serotypes, with types 1, 4, 5, 9V, 19F, and 18C having the highest initial protein/PS ratios. Even though the protein/PS ratios were much higher for types 1, 4, 5, 9V, 19F, and 18C compared to the other serotypes even after the first UF/DF step, the acidification step greatly reduced the protein/PS ratio and not much protein was left after this step.

[0078] As shown in Figure 12, NA/PS ratios also varied from serotype to serotype. Types 1 and 5 had the highest NA/PS ratio, followed by types 18C and 4. The first centrifugation and UF/DF step removed a significant amount of NA for these serotypes and type 1 seemed to be the most efficient. There was very little NA left after the acidification step.

Example 3. Acidification and Activated Carbon Adsorption Step Efficiency Analysis (Serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, and 19F).

[0079] For purification of PnC polysaccharides, the most significant and difficult impurity to remove is proteins. To better understand the impurity removal efficiency of the shortened process, two of the major purification steps, acidification and activated carbon adsorption, were analyzed for protein removal. For the acidification step, the difference in protein concentration (SDS-PAGE) before and after acidification was plotted against the initial protein concentration before acidification for serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, and 19F using the shortened purification process of the invention (Figure 13).

[0080] Due to relatively small changes in solution volume before and after the acidification step, the protein concentration difference divided by the initial protein concentration reflected the protein removal rate by the acidification step. Figure 13 shows the slope of the linear fit of protein concentration difference with respect to initial protein concentration (by SDS-PAGE assay). A very good linear relationship was observed between the protein concentration change and the initial concentration, with the R^2 close to 1. The slope was 0.9876, which corresponds to a 98.76% protein removal (assuming negligible solution volume change). Therefore, for all serotypes studied, the acidification step was very efficient and on average it removed more than 98% protein.

[0081] Similar to the acidification step, the efficiency of the activated carbon adsorption step

was also evaluated by plotting the amount of protein removed (adsorbed on carbon) with respect to initial protein loading (Figure 14). A good linear relationship between the removed protein and the initial protein loading was observed, although the R^2 (0.9723) was not as high as for the acidification step. Because the slope of this linear fit corresponds to the protein removal rate, the activated carbon adsorption step produced 93.87% protein removal.

[0082] Based on an analysis of the acidification and activated carbon adsorption steps, after the two steps only about 0.1% protein was left in the solution.

Conclusions for Examples 1-3

[0083] A shortened purification process was developed to replace the current purification process for capsular polysaccharides of *S. pneumoniae*. The shortened purification process was directly applied to serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, and 19F, and to 19A with slight deviation. The process was conducted at 10L scale using DOC-lysed fermentation broths. The impurity levels, including protein/PS, NA/PS and C-PS/PS ratios, all met their respective specifications. The PS yields were over 50% and comparable to that of the current purification process. In-process PS yield, protein/PS and NA/PS ratios were plotted to compare the behavior of different serotypes. Serotypes 1, 5, 9V, 19F, and 18C were found to be the most difficult to purify based on protein/PS ratios before and after the 100K UF/DF step.

[0084] Step analysis showed that the acidification and activated carbon adsorption steps removed more than 98% and 90% protein, respectively (as measured by SDS-PAGE).

Example 4. Substitution of Non-Animal Derived Lytic Agents for Deoxycholate Sodium in the Production of Polysaccharides

[0085] The present example investigated whether non-animal derived lytic agents could be used as a substitute for deoxycholate sodium (DOC) within the process described above for the production of substantially purified capsular *Streptococcus pneumoniae* polysaccharides. As described above, DOC activates the LytA protein, which is an autolysin that is involved in cell wall growth and division in *Streptococcus pneumoniae*. The LytA protein has choline binding domains in its C-terminal portion, and mutations of the *lytA* gene are known to produce LytA mutants that are resistant to lysis with DOC.

[0086] A rapid microtiter plate assay was developed to identify compounds that cause cell lysis by a mechanism similar to that of DOC. Several non-animal derived alternatives to DOC were identified which were equally effective as DOC at killing *Streptococcus pneumoniae* cells and releasing polysaccharide. Following processing using standard conditions, the size and purity of the polysaccharides produced with the non-animal derived compounds were identical to those produced with DOC.

Methods

[0087] Cell Lysis: Compounds to be tested were added to fermentation broth at a final concentration of 0.1 to 0.01% (v/v) and the solution was allowed to incubate at 37°C for one hour. The solution was then stored at 2-8°C overnight. The following morning the solution was centrifuged to remove all cell debris. The cell-free broth was analyzed by SEC-HPLC to determine the concentration of released polysaccharide. The lysis could be performed at any temperature between 2-37°C, preferably at a pH range of 6.0-7.5. The concentration of the detergent was typically between 0.05-0.2% depending on the particular detergent.

[0088] Mitrotiter Assay: A rapid microtiter plate assay was devised for examining *lytA*-dependent lysis of pneumococci by different detergents or surfactants. Two pairs of isogenic strains of *S. pneumoniae* were used in the assay; one member of each pair was wild type for the *lytA* gene while the other strain carried a deletion in the *lytA* gene. Thus, if lysis were dependent on an active *lytA* function, that detergent would not lyse the mutant strains. The four strains, R6X, R6X $\Delta lytA$, D39 and D39 $\Delta lytA$, were cultivated in HySoy medium to approximately mid-log phase ($OD_{600} \sim 0.2-0.8$; OD_{600} = Optical Density at 600 nm). Cells were then centrifuged and cell pellets were resuspended in HySoy medium. To each well of the microtiter plate, 100 μ L of cell suspension was added, along with 10 μ L of detergent stocks or water as a control. After about 15 minutes at 36°C, the OD_{600} of the samples was measured with a Spectramax® spectrophotometer (Molecular Devices, Sunnyvale, CA). The following results were observed for freshly prepared cells or frozen and thawed cells: Wild- type cells exposed to DOC, sodium dodecyl sulfate, Triton® X-100 and N-laurylsarcosine all had OD values comparable to the medium blank, which indicated that lysis had occurred. On the other hand, the $\Delta lytA$ cells did not lyse in the presence of these detergents, which indicates that *lytA* function is required for these detergents to lyse the cells.

[0089] Isolation of PS Used For Comparative Analytical Studies: Cultures were grown in Hy-Soy medium in 10 L bioreactors. The pH was controlled at around 7.0 using either NaOH or Na_2CO_3 . At the end of growth (as indicated by no further increase in optical density), the cultures were treated with either 0.12% DOC or 0.1% NLS. The cultures were incubated for 12-16 hours. The effectiveness of cell killing was confirmed by plating a sample of treated broth on TSA-Blood Agar plates. Polysaccharide was purified from the clarified lysate using standard procedures (as described above). Purified polysaccharide was examined using a variety of standard analytical techniques appropriate to determine the purity and identity of the material.

[0090] The PS content of the lysate was determined using SEC-HPLC coupled to a refractive index (RI) detector.

Screening for Non-Animal Derived Lytic Agents Using Serotypes 1 and 6B

[0091] *S. pneumoniae* serotypes 1 and 6B were separately grown in Hy-Soy based media. The cultures were separately harvested and separately dispensed into tubes. The non-animal derived compounds to be screened for lytic activity comparable to DOC were prepared as stock solutions (in suitable solvents) and added to the cultures. After overnight incubation, the tubes were centrifuged and the PS content of the lysates for each serotype were determined by SEC-HPLC and compared to DOC.

Screening for Non-Animal Derived Lytic Agents Using *lytA* Mutants

[0092] Isogenic pairs of strains containing the *lytA* mutation were grown in a Hy-Soy based medium. The cells were harvested and dispensed into wells in a microtiter plate. The test compound was added and the culture was incubated. After 15 min at 36°C, the optical density (OD₆₀₀) of each well was determined using a SpectraMax® plate reader (Molecular Devices, Sunnyvale, CA)(see Tables 17 and 18, which summarize results from two separate tests for exemplary compounds).

Table 17. Change in Optical Density for <i>lytA</i> Mutant Strains (Test 1)				
	Change in OD ₆₀₀			
Compound	R6X	R6X Δ <i>lytA</i>	D39	D39 Δ <i>lytA</i>
Blank				
No addn				
Deoxycholic acid, 0.1%	100%	34%	99%	47%
Lithocholic acid, 0.1%	20%	16%	8%	-2%
Tauroglycocholic acid, 0.1 %	51 %	-16%	47%	-20%
Heptanoic acid, 0.1 %	25%	-57%	7%	-30%
SDS, 0.1%	99%	28%	95%	36%
Octanesulfonic acid, 0.1%	43%	6%	34%	16%
Triton® X-100, 0.1%	91%	22%	97%	30%
Tween 80, 0.1%	36%	14%	20%	19%
Tween 20, 0.1%	44%	22%	28%	19%
N-lauryl sarcosine, 0.1 %	100%	-34%	89%	-44%
Pluronic L31, 0.1%	39%	0%	17%	-38%
Pluronic L-61, 0.1%	23%	-3%	-19%	14%
Pluronic L81, 0.1%	21%	-3%	-14%	10%
Antarox 17-R-2, 0.1 %	37%	24%	7%	16%

Table 18. Change in Optical Density for *lytA* Mutant Strains (Test 2)

Compound	Change in OD ₆₀₀			
	R6X	R6X Δ <i>lytA</i>	D39	D39 Δ <i>lytA</i>
Blank				
No addn				
Water	-8%	-8%	3%	-8%
Deoxycholic acid, 0.1 %	103%	20%	101%	41%
SDS, 0.1%	102%	-13%	100%	16%
N-lauryl sarcosine, 0.1 %	102%	-40%	101%	14%
Triton® X-100, 0.1%	101%	-19%	100%	4%
Tween 20, 0.1%	6%	-17%	-5%	-3%
Octanesulfonic acid, 0.1%	13%	3%	18%	-3%

[0093] Based on the screening studies described above, the following non-animal derived lytic agent alternatives to DOC were identified: decanesulfonic acid, Igepal® CA-630 (tert-octylphenoxy poly(oxyethylene)ethanol; CAS #: 9002-93-1; available from Sigma Aldrich, St. Louis, MO), N-lauryl sarcosine sodium (NLS), lauryl iminodipropionate, sodium dodecyl sulfate, Triton® X-100, chenodeoxycholate, hyodeoxycholate, glycodeoxycholate, taurodeoxycholate, taurochenodeoxycholate, and cholate.

Comparison of DOC-Lysed PS to NLS-Lysed PS

[0094] *S. pneumoniae* polysaccharide serotypes 1, 4, 5, 6A, 6B, and 7F were purified at the 10L scale as described above in Examples 1 and 2 using the improved process of the present invention. However, in one group NLS (0.1%) was used as the lytic agent while in another group DOC (0.12%) was used as the lytic agent.

[0095] PS yield, protein/PS ratios, NA/PS ratios, and PS molecular weight were measured as described above, and results are summarized in Table 19. These results showed that the use of NLS as a lytic agent within the purification methods of the invention produced relatively high PS yields with relatively low protein and nucleic acid levels. In fact, for the majority of serotypes tested, use of NLS as a lytic agent produced higher PS yields compared to the use of DOC.

Table 19. PS Characterization For Different Serotypes Using DOC vs. NLS

Serotype	Lytic Agent	PS Yield (%)	Protein/PS (%)	Nucleic Acid/PS (%)	SEC MALLS MW (g/mol, 10 ⁶)
1	0.12% DOC	20%	4.2%	0.04%	0.615
	0.1%				

Table 19. PS Characterization For Different Serotypes Using DOC vs. NLS					
Serotype	Lytic Agent	PS Yield (%)	Protein/PS (%)	Nucleic Acid/PS (%)	SEC MALLS MW (g/mol, 10 ⁶)
1	NLS	41%	2%	0.04%	0.65
4	0.12% DOC	72%	0.05%	0.02%	0.38
4	0.1% NLS	57%	0.67%	0.25%	0.34
5	0.12% DOC	57%	3.6%	0.18%	0.32
5	0.1% NLS	63%	1.8%	0.01%	0.35
6A	0.12% DOC	56%	0.7%	0.00%	0.55
6A	0.1% NLS	66%	1.1%	0.05%	0.43
6B	0.12% DOC	38%	0.6%	0.05%	0.969
6B	0.1% NLS	55%	0.1%	0.01%	1.0
7F	0.12% DOC	73%	0.7%	0.06%	0.814
7F	0.1% NLS	76%	0.5%	0.02%	0.912

[0096] All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains.

[0097] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

[0098] Particular embodiments of the invention are set forth in the following numbered paragraphs:

1. 1. A process for producing substantially purified capsular polysaccharides from a *Streptococcus pneumoniae* cell lysate, the process comprising the steps of:
 1. (a) providing a fermentation broth comprising bacterial cells that produce a selected *Streptococcus pneumoniae* serotype;
 2. (b) lysing the bacterial cells in step (a) with a lytic agent, thereby producing a cell lysate comprising cell debris, soluble proteins, nucleic acids, and polysaccharides;
 3. (c) clarifying the cell lysate of step (b) using centrifugation or filtration to remove cell debris, thereby producing a clarified cell lysate;

4. (d) ultrafiltering and diafiltering the clarified cell lysate of step (c) to remove low molecular weight impurities and increase polysaccharide concentration, thereby producing a retentate;
5. (e) lowering the pH of the retentate of step (d) to less than 4.5 using an organic or a mineral acid to precipitate protein and nucleic acids, thereby forming an acidified retentate solution;
6. (f) holding the acidified retentate solution formed in step (e) for a time sufficient to allow settling of the precipitate, followed by filtration or centrifugation of the acidified retentate solution, thereby producing a clarified polysaccharide solution;
7. (g) filtering the clarified polysaccharide solution of step (f) through an activated carbon filter;
8. (h) ultrafiltering and diafiltering the filtered solution produced by step (g), thereby producing a concentrated purified polysaccharide solution; and
9. (i) filtering the concentrated purified polysaccharide solution produced by step (h) using a sterile filter;

whereby substantially purified capsular polysaccharides in the form of a solution are produced.

2. 2. The process of paragraph 1, wherein the selected *Streptococcus pneumoniae* serotype is selected from the group consisting of 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F.
3. 3. The process of paragraph 1 or 2, wherein the pH of step (e) is lowered to about 3.5.
4. 4. The process of paragraph 1, 2 or 3, wherein the diafiltration of step (h) comprises a pH adjustment to between about 5.5 to about 7.5.
5. 5. The process of paragraph 1, 2 or 3, wherein the diafiltration of step (h) comprises a pH adjustment to between about 7.0 to about 7.5.
6. 6. The process of paragraph 1, 2 or 3, wherein the diafiltration of step (h) comprises a pH adjustment to about 7.4.
7. 7. The process of any one of paragraphs 1 to 6, wherein step (e) removes at least 98% of protein from the retentate of step (d).
8. 8. The process of any one of paragraphs 1 to 7, wherein step (g) removes at least 90% of the protein from the clarified polysaccharide solution of step (f).
9. 9. The process of any one of paragraphs 1 to 8, wherein the activated carbon filter of step (g) comprises wood-based phosphoric acid-activated carbon.
10. 10. The process of any one of paragraphs 1 to 9, wherein step (f) comprises holding the acidified retentate solution formed in step (e) for at least 2 hours.
11. 11. The process of any one of paragraphs 1 to 10, wherein the lytic agent of step (b) is deoxycholate sodium.
12. 12. The process of any one of paragraphs 1 to 10, wherein the lytic agent of step (b) is a non-animal derived lytic agent.
13. 13. The process of paragraph 12, wherein said non-animal derived lytic agent is selected from the group consisting of: decanesulfonic acid, tert-octylphenoxy poly(oxy ethyl ene)ethanols, octylphenol ethylene oxide condensates, N-lauryl sarcosine sodium (NLS), lauryl iminodipropionate, sodium dodecyl sulfate, chenodeoxycholate, hyodeoxycholate, glycodeoxycholate, taurodeoxycholate, taurochenodeoxycholate, and

cholate.

14. 14. The process of paragraph 12, wherein said non-animal derived lytic agent is N-lauryl sarcosine sodium.
15. 15. A process for producing substantially purified capsular polysaccharides from a *Streptococcus pneumoniae* cell lysate comprising serotype 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F, or 23F, the process comprising the steps of:
 1. (a) providing a fermentation broth comprising bacterial cells that produce *Streptococcus pneumoniae* serotype 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F, or 23F;
 2. (b) lysing the bacterial cells in step (a) with a lytic agent, thereby producing a cell lysate comprising cell debris, soluble proteins, nucleic acids, and polysaccharides;
 3. (c) clarifying the cell lysate of step (b) using centrifugation or filtration to remove cell debris, thereby producing a clarified cell lysate;
 4. (d) ultrafiltering and diafiltering the clarified cell lysate of step (c) at room temperature at neutral pH in salt free media to remove low molecular weight impurities and increase polysaccharide concentration, thereby producing a salt free retentate;
 5. (e) lowering the pH of the salt free retentate of step (d) to less than 4.5 using an organic or a mineral acid to precipitate protein and nucleic acids, thereby forming an acidified retentate solution;
 6. (f) holding the acidified retentate solution formed in step (e) for at least 2 hours at room temperature to allow settling of the precipitate, followed by filtration or centrifugation of the acidified retentate solution, thereby producing a clarified polysaccharide solution;
 7. (g) filtering the clarified polysaccharide solution of step (f) through an activated carbon filter;
 8. (h) ultrafiltering and diafiltering the filtered solution produced by step (g), thereby producing a concentrated purified polysaccharide solution; and
 9. (i) filtering the concentrated purified polysaccharide solution produced by step (h) using a sterile filter;

whereby substantially purified capsular polysaccharides comprising serotype 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F, or 23F in the form of a solution are produced.
16. 16. The process of paragraph 15, wherein the pH of step (e) is lowered to about 3.5.
17. 17. The process of paragraph 15 or 16, wherein the diafiltration of step (h) comprises a pH adjustment to between about 5.5 to about 7.5.
18. 18. The process of paragraph 15 or 16, wherein the diafiltration of step (h) comprises a pH adjustment to between about 7.0 to about 7.5.
19. 19. The process of paragraph 15 or 16, wherein the diafiltration of step (h) comprises a pH adjustment to about 7.4.
20. 20. The process of any one of paragraphs 15 to 19, wherein step (e) removes at least 98% of protein from the salt free retentate of step (d).
21. 21. The process of any one of paragraphs 15 to 20, wherein step (g) removes at least 90% of the protein from the clarified polysaccharide solution of step (f).
22. 22. The process of any one of paragraphs 15 to 21, wherein the activated carbon filter of step (g) comprises wood-based phosphoric acid-activated carbon.

23. 23. The process of any one of paragraphs 15 to 22, wherein the lytic agent of step (b) is deoxycholate sodium.
24. 24. The process of any one of paragraphs 15 to 22, wherein the lytic agent of step (b) is a non-animal derived lytic agent.
25. 25. The process of paragraph 24, wherein said non-animal derived lytic agent is selected from the group consisting of: decanesulfonic acid, tert-octylphenoxy poly(oxyethylene)ethanols, octylphenol ethylene oxide condensates, N-lauryl sarcosine sodium (NLS), lauryl iminodipropionate, sodium dodecyl sulfate, chenodeoxycholate, hyodeoxycholate, glycodeoxycholate, taurodeoxycholate, taurochenodeoxycholate, and cholate.
26. 26. The process of paragraph 24, wherein said non-animal derived lytic agent is N-lauryl sarcosine sodium.
27. 27. A process for producing substantially purified capsular polysaccharides from a *Streptococcus pneumoniae* cell lysate comprising serotype 19A, the process comprising the steps of:
1. (a) providing a fermentation broth comprising bacterial cells that produce *Streptococcus pneumoniae* serotype 19A;
 2. (b) lysing the bacterial cells in step (a) with a lytic agent, thereby producing a cell lysate comprising cell debris, soluble proteins, nucleic acids, and polysaccharides;
 3. (c) clarifying the cell lysate of step (b) using centrifugation or filtration to remove cell debris, thereby producing a clarified cell lysate;
 4. (d) ultrafiltering and diafiltering the clarified cell lysate of step (c) at about 4 °C at a pH of about 6 in sodium phosphate buffer to remove low molecular weight impurities and increase polysaccharide concentration, thereby producing a retentate;
 5. (e) lowering the pH of the retentate of step (d) to less than 4.5 using an organic or a mineral acid to precipitate protein and nucleic acids, thereby forming an acidified retentate solution;
 6. (f) holding the acidified retentate solution formed in step (e) for at least 2 hours at about 4°C to allow settling of the precipitate, followed by filtration or centrifugation of the acidified retentate solution, thereby producing a clarified polysaccharide solution;
 7. (g) adjusting the pH of the clarified polysaccharide solution of step (f) to about 6, thereby producing a pH-adjusted clarified polysaccharide solution;
 8. (h) filtering the pH-adjusted clarified polysaccharide solution of step (g) through an activated carbon filter;
 9. (i) ultrafiltering and diafiltering the filtered solution produced by step (h), thereby producing a concentrated purified polysaccharide solution; and
 10. (j) filtering the concentrated purified polysaccharide solution produced by step (i) using a sterile filter;
- whereby substantially purified capsular polysaccharides comprising serotype 19A in the form of a solution are produced.
28. 28. The process of paragraph 27, wherein the pH of step (e) is lowered to about 3.5.
29. 29. The process of paragraph 27 or 28, wherein the diafiltration of step (i) comprises a

- pH adjustment to between about 5.5 to about 7.5.
30. 30. The process of paragraph 27 or 28, wherein the diafiltration of step (i) comprises a pH adjustment to between about 7.0 to about 7.5.
 31. 31. The process of paragraph 27 or 28, wherein the diafiltration of step (i) comprises a pH adjustment to about 7.4.
 32. 32. The process of any one of paragraphs 27 to 31, wherein step (e) removes at least 98% of protein from the retentate of step (d).
 33. 33. The process of any one of paragraphs 27 to 32, wherein step (h) removes at least 90% of the protein from the pH-adjusted clarified polysaccharide solution of step (g).
 34. 34. The process of any one of paragraphs 27 to 33, wherein the activated carbon filter of step (h) comprises wood-based phosphoric acid-activated carbon.
 35. 35. The process of any one of paragraphs 27 to 34, wherein the sodium phosphate buffer of step (d) is 25 mM sodium phosphate.
 36. 36. The process of any one of paragraphs 27 to 35, wherein the lytic agent of step (b) is deoxycholate sodium.
 37. 37. The process of any one of paragraphs 27 to 35, wherein the lytic agent of step (b) is a non-animal derived lytic agent.
 38. 38. The process of paragraph 37, wherein said non-animal derived lytic agent is selected from the group consisting of: decanesulfonic acid, tert-octylphenoxy poly(oxyethylene)ethanols, octylphenol ethylene oxide condensates, N-lauryl sarcosine sodium (NLS), lauryl iminodipropionate, sodium dodecyl sulfate, chenodeoxycholate, hyodeoxycholate, glycodeoxycholate, taurodeoxycholate, taurochenodeoxycholate, and cholate.
 39. 39. The process of paragraph 37, wherein said non-animal derived lytic agent is N-lauryl sarcosine sodium.
 40. 40. A process for the manufacture of a pneumococcal vaccine, which comprises producing substantially purified capsular polysaccharides from a *Streptococcus pneumoniae* cell lysate by a process as described in any one of paragraphs 1 to 39 and using said substantially purified capsular polysaccharides in the manufacture of a pneumococcal vaccine.
 41. 41. A process as described in paragraph 40 wherein the pneumococcal vaccine containing polysaccharide conjugated with a protein carrier.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US20060228381A [0005] [0013]

Non-patent literature cited in the description

- CHEMICAL ABSTRACTS, 9002-93-1 [0024] [0093]

P a t e n t k r a v

1. Fremgangsmåde til fremstilling af oprensede kapselpolysaccharider ud fra et *Streptococcus pneumoniae*-cellelysat, hvilken fremgangsmåde omfatter trinnene:

(a) at tilvejebringe en fermenteringssuppe omfattende bakterieceller, der producerer en udvalgt *Streptococcus pneumoniae*-serotype;

(b) at lysere bakteriecellerne fra trin (a) med et lytisk middel, hvorved der fremstilles et cellelysat omfattende celledbris, opløselige proteiner, nukleinsyrer og polysaccharider;

(c) at klare cellelysatet fra trin (b) under anvendelse af centrifugering eller filtrering til fjernelse af celledbris, hvorved der fremstilles et klaret cellelysat;

(d) at ultrafiltrere og diafiltrere det klarede cellelysat fra trin (c) til fjernelse af urenheder med lav molekylvægt og øgning af polysaccharidkoncentrationen, hvorved der fremstilles et retentat;

(e) at sænke pH-værdien af retentatet fra trin (d) til mindre end 4,5 under anvendelse af en organisk syre eller en mineralsyre til udfældning af protein og nukleinsyrer, hvorved der dannes en forsuret retentatopløsning;

(f) at fastholde den forsurede retentatopløsning dannet i trin (e) i et tidsrum, der er tilstrækkeligt til at muliggøre, at udfældningsproduktet bundfælder sig, efterfulgt af filtrering eller centrifugering af den forsurede retentatopløsning, hvorved der fremstilles en polysaccharidopløsning;

(g) at filtrere den klarede polysaccharidopløsning fra trin (f) gennem et aktive-ret kulfilter;

(h) at ultrafiltrere og diafiltrere den filtrerede opløsning fremstillet ved trin (g), hvorved der fremstilles en koncentreret oprenset polysaccharidopløsning; og

(i) at filtrere den koncentrede oprensede polysaccharidopløsning fremstillet ved trin (h) under anvendelse af et sterilt filter;

hvorved der fremstilles oprensede kapselpolysaccharider i form af en opløsning.

2. Fremgangsmåde ifølge krav 1, hvor den valgte *Streptococcus pneumoniae*-serotype udvælges fra gruppen bestående af 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F og 23F.

3. Fremgangsmåde ifølge krav 1, omfattende trinnene:

(a) at tilvejebringe en fermenteringssuppe omfattende bakterieceller, der producerer *Streptococcus pneumoniae*-serotype 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F eller 23F;

5 (b) at lysere bakteriecellerne fra trin (a) med et lytisk middel, hvorved der fremstilles et cellelysat omfattende celledbris, opløselige proteiner, nukleinsyrer og polysaccharider;

(c) at klare cellelysatet fra trin (b) under anvendelse af centrifugering eller filtrering til fjernelse af celledbris, hvorved der fremstilles et klaret cellelysat;

10 (d) at ultrafiltrere og diafiltrere det klarede cellelysat fra trin (c) ved rumtemperatur ved neutral pH i saltfri medier til fjernelse af urenheder med lav molekylvægt og øgning af polysaccharidkoncentrationen, hvorved der fremstilles et saltfrit retentat;

15 (e) at sænke pH-værdien af det saltfrie retentat fra trin (d) til mindre end 4,5 under anvendelse af en organisk syre eller en mineralsyre til udfældning af protein og nukleinsyrer, hvorved der dannes en forsuret retentatopløsning;

(f) at fastholde den forsurede retentatopløsning dannet i trin (e) i mindst 2 timer ved rumtemperatur til muliggørelse af, at udfældningsproduktet bundfælder sig, efterfulgt af filtrering eller centrifugering af den forsurede retentatopløsning, hvorved der fremstilles en polysaccharidopløsning;

20 (g) at filtrere den klarede polysaccharidopløsning fra trin (f) gennem et aktivret kulfilter;

(h) at ultrafiltrere og diafiltrere den filtrerede opløsning fremstillet ved trin (g), hvorved der fremstilles en koncentreret oprenset polysaccharidopløsning; og

25 (i) at filtrere den koncentrerede oprensede polysaccharidopløsning fremstillet ved trin (h) under anvendelse af et sterilt filter;

hvorved der fremstilles oprensede kapselpolysaccharider omfattende serotype 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19F eller 23F i form af en opløsning.

30 **4.** Fremgangsmåde ifølge krav 1, 2 eller 3, hvor pH-værdien fra trin (e) sænkes til ca. 3,5.

5. Fremgangsmåde ifølge krav 1, 2, 3 eller 4, hvor diafiltreringen i trin (h) omfatter en pH-justering til mellem ca. 5,5 til ca. 7,5, fortrinsvis hvor diafiltreringen i trin (h) omfatter en pH-justering til mellem ca. 7,0 til ca. 7,5, endnu mere fortrinsvis hvor diafiltreringen i trin (h) omfatter en pH-justering til ca. 7,4.

35

6. Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 5, hvor trin (e) fjerner mindst 98 % af protein fra retentatet fra trin (d).

5 7. Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 6, hvor trin (g) fjerner mindst 90 % af proteinet fra den klarede polysaccharidopløsning fra trin (f).

10 8. Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 7, hvor det aktiverede kulfilter i trin (g) omfatter træbaseret phosphorsyreaktiveret kul.

9. Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 8, hvor trin (f) omfatter at fastholde den forsuredede retentatopløsning dannet i trin (e) i mindst 2 timer.

15 10. Fremgangsmåde ifølge krav 1, omfattende trinnene:

(a) at tilvejebringe en fermenteringssuppe omfattende bakterieceller, der producerer *Streptococcus pneumoniae*-serotype 19A;

20 (b) at lysere bakteriecellerne fra trin (a) med et lytisk middel, hvorved der fremstilles et cellelysats omfattende celledetrit, opløselige proteiner, nukleinsyrer og polysaccharider;

(c) at klare cellelysats fra trin (b) under anvendelse af centrifugering eller filtrering til fjernelse af celledetrit, hvorved der fremstilles et klaret cellelysats;

25 (d) at ultrafiltrere og diafiltrere det klarede cellelysats fra trin (c) ved ca. 4 °C ved en pH-værdi på ca. 6 i natriumphosphatbuffer til fjernelse af urenheder med lav molekylvægt og øgning af polysaccharidkoncentrationen, hvorved der fremstilles et retentat;

(e) at sænke pH-værdien af retentatet fra trin (d) til mindre end 4,5 under anvendelse af en organisk syre eller en mineralsyre til udfældning af protein og nukleinsyrer, hvorved der dannes en forsuret retentatopløsning;

30 (f) at fastholde den forsuredede retentatopløsning dannet i trin (e) i mindst 2 timer ved ca. 4 °C til muliggørelse af, at udfældningsproduktet bundfælder sig, efterfulgt af filtrering eller centrifugering af den forsuredede retentatopløsning, hvorved der fremstilles en polysaccharidopløsning;

35 (g) at justere pH-værdien af den klarede polysaccharidopløsning fra trin (f) til ca. 6, hvorved der fremstilles en pH-justeret klaret polysaccharidopløsning;

(h) at filtrere den pH-justerede klarede polysaccharidopløsning fra trin (g) gennem et aktiveret kulfilter;

(i) at ultrafiltrere og diafiltrere den filtrerede opløsning fremstillet ved trin (h), hvorved der fremstilles en koncentreret oprenset polysaccharidopløsning; og

5 (j) at filtrere den koncentrerede oprensede polysaccharidopløsning fremstillet ved trin (i) under anvendelse af et sterilt filter; hvorved der fremstilles oprensede kapselpolysaccharider omfattende serotype 19A i form af en opløsning.

10 **11.** Fremgangsmåde ifølge krav 10, hvor pH-værdien fra trin (e) sænkes til ca. 3,5.

15 **12.** Fremgangsmåde ifølge krav 10 eller 11, hvor diafiltreringen i trin (i) omfatter en pH-justering til mellem ca. 5,5 til ca. 7,5, fortrinsvis hvor diafiltreringen i trin (i) omfatter en pH-justering til mellem ca. 7,0 til ca. 7,5, endnu mere foretrukket hvor diafiltreringen i trin (i) omfatter en pH-justering til ca. 7,4.

20 **13.** fremgangsmåde ifølge et hvilket som helst af kravene 10 til 12, hvor trin (e) fjerner mindst 98 % af protein fra retentatet i trin (d).

14. Fremgangsmåde ifølge et hvilket som helst af kravene 10 til 13, hvor trin (h) fjerner mindst 90 % af proteinet fra den pH-justerede klarede polysaccharidopløsning fra trin (g).

25 **15.** Fremgangsmåde ifølge et hvilket som helst af kravene 10 til 13, hvor det aktiverede kulfilter i trin (h) omfatter træbaseret phosphorsyreaktiveret kul.

16. Fremgangsmåde ifølge et hvilket som helst af kravene 10 til 15, hvor natriumphosphatbufferen i trin (d) er 25 mM natriumphosphat.

30

17. Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 16, hvor det lytiske middel i trin (b) er deoxycholatnatrium.

35 **18.** Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 16, hvor det lytiske middel i trin (b) er et lytisk middel af ikke-animalsk oprindelse, fortrinsvis

5 hvor det lytiske middel af ikke-animalsk oprindelse er udvalgt fra gruppen bestående af: decansulfonsyre, tert-octylphenoxy-poly(oxyethylen)ethanoler, octylphenoxyethylenoxidkondensater, N-lauryl-sarcosinnatrium (NLS), lauryliminodipropionat, natriumdodecylsulfat, chenodeoxycholat, hyodeoxycholat, glycodeoxycholat, taurodeoxycholat, taurochenodeoxycholat og cholat, endnu mere foretrukket hvor det lytiske middel af ikke-animalsk oprindelse er N-lauryl-sarcosinnatrium.

10 **19.** Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 18, hvor mineralsyren er udvalgt fra gruppen bestående af saltsyre, salpetersyre, phosphorsyre og svovlsyre.

15 **20.** Fremgangsmåde til fremstilling af en pneumokokvaccine, som omfatter at fremstille oprensede kapselpolysaccharider fra et *Streptococcus pneumoniae*-cellelysat ved en fremgangsmåde ifølge et hvilket som helst af kravene 1 til 19 og at anvende de oprensede kapselpolysaccharider til fremstilling af en pneumokokvaccine.

20 **21.** Fremgangsmåde ifølge krav 20, hvor pneumokokvaccinen indeholder polysaccharid konjugeret med en proteinbærer.

DRAWINGS

Figure 1





Figure 2

A.



B.




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14

