

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2016322617 B2**

(54) Title
Novel lactobacillus having various functions, and use thereof

(51) International Patent Classification(s)
C12N 1/20 (2006.01) **A61K 35/747** (2015.01)
A23L 33/135 (2016.01) **C12R 1/225** (2006.01)
A61K 35/745 (2015.01) **C12R 1/24** (2006.01)

(21) Application No: **2016322617** (22) Date of Filing: **2016.09.07**

(87) WIPO No: **WO17/047968**

(30) Priority Data

(31) Number	(32) Date	(33) Country
10-2016-0005018	2016.01.15	KR
10-2015-0130124	2015.09.15	KR

(43) Publication Date: **2017.03.23**

(44) Accepted Journal Date: **2020.01.16**

(71) Applicant(s)
Navipharm Co, Ltd;University-Industry Cooperation Group of Kyung Hee University

(72) Inventor(s)
Kim, Dong Hyun;Han, Myung Joo

(74) Agent / Attorney
FPA Patent Attorneys Pty Ltd, Level 43 101 Collins Street, Melbourne, VIC, 3000, AU

(56) Related Art
WO 2013190068 A1

(12) 특허협력조약에 의하여 공개된 국제출원

(19) 세계지식재산권기구
국제사무국

(43) 국제공개일
2017년 3월 23일 (23.03.2017)



(10) 국제공개번호
WO 2017/047968 A1

- (51) 국제특허분류:
C12N 1/20 (2006.01) *A61K 35/745* (2014.01)
A23L 33/135 (2016.01) *C12R 1/225* (2006.01)
A61K 35/747 (2014.01) *C12R 1/24* (2006.01)
- (21) 국제출원번호: PCT/KR2016/009994
- (22) 국제출원일: 2016년 9월 7일 (07.09.2016)
- (25) 출원언어: 한국어
- (26) 공개언어: 한국어
- (30) 우선권정보:
10-2015-0130124 2015년 9월 15일 (15.09.2015) KR
10-2016-0005018 2016년 1월 15일 (15.01.2016) KR
- (71) 출원인: 경희대학교 산학협력단 (UNIVERSITY-INDUSTRY COOPERATION GROUP OF KYUNG HEE UNIVERSITY) [KR/KR]; 17104 경기도 용인시 기흥구 덕영대로 1732 (서천동, 경희대학교 국제캠퍼스내), Gyeonggi-do (KR).
- (72) 발명자: 김동현 (KIM, Dong Hyun); 02823 서울시 성북구 선잠로 92-24 (성북동), Seoul (KR). 한명주 (HAN, Myung Joo); 02823 서울시 성북구 선잠로 92-24 (성북동), Seoul (KR).
- (74) 대리인: 김영갑 (KO, Young Kap); 13636 경기도 성남시 분당구 성남대로 69, 410 호 (구미동,로드랜드 EZ 타워), Gyeonggi-do (KR).


- (81) 지정국 (별도의 표시가 없는 한, 가능한 모든 종류의 국내 권리의 보호를 위하여): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) 지정국 (별도의 표시가 없는 한, 가능한 모든 종류의 역내 권리의 보호를 위하여): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), 유라시아 (AM, AZ, BY, KG, KZ, RU, TJ, TM), 유럽 (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

공개:

- 국제조사보고서와 함께 (조약 제 21 조(3))
- 청구범위 보정 기한 만료 전의 공개이며, 보정서를 접수하는 경우 그에 관하여 별도 공개함 (규칙 48.2(h))
- 명세서의 서열목록 부분과 함께 (규칙 5.2(a))

(54) Title: NOVEL LACTOBACILLUS HAVING VARIOUS FUNCTIONS, AND USE THEREOF

(54) 발명의 명칭: 다양한 기능성을 가진 신규 유산균 및 이의 용도



(57) Abstract: The present invention provides a novel *Lactobacillus* sp. strain, a novel *Bifidobacterium* sp. strain or a lactobacillus mixture thereof, which is isolated from kimchi or human feces. According to the present invention, the specific *Lactobacillus* sp. strain or the specific *Bifidobacterium* sp. strain is isolated from kimchi or human feces, thereby having high safety and various physiological activities such as an antioxidant activity, a β -glucuronidase inhibitory activity, a lipopolysaccharide (LPS) generation inhibitory activity or a tight junction protein expression-inducing activity. Therefore, according to the present invention, the specific *Lactobacillus* sp. strain, the specific *Bifidobacterium* sp. strain or a lactobacillus mixture thereof can be used as a functional food and drug material useful for preventing, alleviating or treating a bowel injury, a liver injury, allergies, inflammatory diseases or obesity.

(57) 요약서: 본 발명은

[다음 쪽 계속]



WO 2017/047968 A1



김치 또는 사람의 분변에서 분리된 신규 락토바실러스속 균주, 신규 비피도박테리움속 균주 또는 이들의 혼합 유산균을 제공한다. 본 발명에 따른 특정 락토바실러스속 균주 또는 특정 비피도박테리움속 균주는 김치 또는 사람의 분변에서 분리되어 안전성이 높고 항산화 활성, 베타-글루쿠로니다제(β -glucuronidase) 저해 활성, 지질다당류(lipopolysaccharide, LPS) 생성 억제 활성 또는 밀착연접단백질(tight junction protein) 발현 유도 활성 등과 같은 다양한 생리활성을 갖는다. 따라서, 본 발명에 따른 특정 락토바실러스속 균주, 특정 비피도박테리움속 균주 또는 이들의 혼합 유산균은 장 손상, 간 손상, 알러지 질환, 염증 질환, 또는 비만 등을 예방, 개선 또는 치료하는데에 유용한 기능성 식의약 소재로 사용될 수 있다.

【DESCRIPTION】

【Invention Title】

NOVEL LACTOBACILLUS HAVING VARIOUS FUNCTIONS, AND USE
THEREOF

5 **【Technical Field】**

The present invention relates to novel lactic acid
bacteria and the like, and more particularly to novel lactic
acid bacteria or novel lactic acid bacteria mixtures, which are
isolated from kimchi or human feces and have various
10 physiological activities such as antioxidant activity, β -
glucuronidase-inhibitory activity, lipopolysaccharide (LPS)
production-inhibitory activity or tight junction protein
expression-inducing activity. Moreover, the present invention
relates to various food and medicinal uses of novel lactic acid
15 bacteria or novel lactic acid bacteria mixtures.

【Background Art】

As humanity has developed into a prosperous society, the
fast food has been rapidly become popular and the pattern of
diseases has also changed dramatically. In particular, in
20 modern people, intestinal flora disturbance, intestinal
permeability syndrome, colitis, liver diseases, allergic
diseases, obesity and the like are increasing due to fast food
eating habits based on meat and fat, irregular meal, excessive
drinking, lack of exercise, excessive stress, exposure to
25 harmful environments, and the like.

Intestinal Flora Disturbance

There are many bacteria living in the gastrointestinal tract of the human body. The human body has about 10 trillion normal cells, but has about 100 trillion bacteria which are about 10-fold larger than the normal cells. These bacteria can be divided into beneficial bacteria that help human intestinal health and harmful bacteria that are harmful to human health. The health of human body can be maintained when beneficial bacterial such as *Lactobacillus*, *Bifidobacterium*, *Streptococcus*, *Leuconostoc*, *Pediococcus*, *Sporolactobacillus* and the like are more dominant in the gastrointestinal tract than harmful bacteria. Otherwise, diseases can be caused, such as obesity, intestinal permeability syndrome, liver diseases, accelerated aging, enteritis and the like.

Intestinal Permeability Syndrome

The gastrointestinal tract of the human body is composed of mucus and villi, which efficiently absorb nutrient components, but prevent the absorption of pathogenic microorganisms having a high molecular weight or toxins produced by these microorganisms. In addition, the human body has an immune system capable of protecting the body from invasion of external antigens having a high molecular weight. However, due to infection with many pathogenic microorganisms or toxins, excessive stress, intake of foods such as high-fat diets capable of proliferating harmful bacterial living in the

gastrointestinal tract, excessive alcohol intake, the abuse of drugs (e.g., antibiotics) and the like, intestinal flora is disturbed, abnormalities in the gastrointestinal tract's immune system occur, and expression of tight junction proteins is inhibited. If expression of tight junction proteins is inhibited, tight junction of intestinal mucosa becomes loosened, and the invasion into the body of large molecules due to the loosened gap and abnormalities in the immune system. Intestinal permeability syndrome is also known as leaky gut syndrome, and refers to a condition in which external such as less digested foods, pathogenic microorganisms, toxins or the like are continuously introduced into blood, because the tight junction barrier system of epithelial cells forming the gastrointestinal tract is not smoothly operated. When intestinal permeability syndrome occurs, external antigens that are generally not absorbed into the body enter the body, thus causing ulcerative colitis, Crohn's disease, liver injury, liver dysfunction, allergic diseases (including asthma), atopy, autoimmune diseases, steatorrhea, digestive absorption disorder, acne, accelerated aging, endotoxemia, intestinal infection, eczema, irritable bowel syndrome, chronic fatigue syndrome, psoriasis, rheumatoid arthritis, pancreatic insufficiency, inflammatory joint diseases or the like.

Colitis

Although it was previously known that the incidence of

ulcerative colitis and Crohn's disease is high in Europeans, the number of patients with ulcerative colitis and Crohn's disease in oriental countries including Korea has recently increased rapidly due to changes in lifestyles such as eating habits. However, the cause is unclear, and thus a fundamental treatment method for these diseases has not yet been established. For this reason, drugs are used which do not aim to completely treat, but aim to relieve symptoms and maintain this relieved state over the longest possible period. As drugs for this symptomatic therapy, aminosalicylic acid agents, adrenocorticosteroid agents, immunosuppressants and the like are mainly used, but have been reported to cause various side effects. For example, sulfasalazine which is frequently used as an aminosalicylic acid agent was reported to cause side effects, including nausea, vomiting, anorexia, rash, headache, liver injury, leukocytopenia, abnormal red blood cells, proteinuria, diarrhea and the like. In addition, adrenocorticosteroid agents are generally used by prednisolone oral administration, infusion, suppository, intravenous injection or the like, but cause strong side effects such as gastric ulcer or femoral necrosis upon long-term use. However, discontinuation of medication can cause symptoms to recur, and thus these drugs must be continuously used. Accordingly, there is a need to develop agents for treating intestinal bowel diseases, such as ulcerative colitis, Crohn's diseases and the

like, which have excellent effects, are safe and cause no side effects. Irritable bowel syndrome (IBS) is also a chronic abdominal disease whose cause is unclear. Currently, there is no fundamental therapeutic agent for IBS, and symptomatic
5 therapy is performed for the purpose of relieving symptoms of each type of IBS. For example, for diarrhea-IBS, an anticholinergic agent having spasmolytic action that suppresses the contraction of smooth muscles is used, and for constipation-IBS, salt laxatives are used. For alternating-IBS
10 difficult to control with drugs, an agent for improving gastrointestinal motor function is fundamentally used.

Liver Diseases

The liver in the human body plays roles such as energy metabolism (nutrient treatment and storage, and waste
15 excretion), detoxification of toxins, synthesis of serum proteins, and smooth absorption of fat in the bowel by bile juice secretion, and is also important in immunity maintenance (body defense) and vitamin metabolism. However, infection with hepatitis viruses or excessive intake of alcohol or high-fat
20 meals may cause liver diseases such as hepatitis, fatty liver or liver cirrhosis. In addition, liver diseases may also be caused by drugs (tuberculosis therapeutic drugs, aspirin, antibiotics, anesthetics, antihypertensive drugs, oral contraceptives, etc.), congenital metabolic disorders, heart
25 failure, shock, or the like. When liver disease occurs, it can

develop into chronic hepatitis, starting with acute hepatitis with fatigue, vomiting, diarrhea, anorexia, jaundice, right upper quadrant pain, fever or muscle pain.

Allergic Diseases

5 As society has become more complicated and the industry and civilization has developed, environmental pollution and stress have increased, and as eating habits have changed, patients with allergic diseases have increased every year. Patients with allergic diseases such as atopy, anaxylolosis,
10 asthma and the like were less than 1% in 1980, but increased rapidly to 5% or more in 2000s, and are estimated to be more than 10%, including potential patients. Allergic diseases are caused by excessive immune responses of a body, which result from antigen-antibody reactions, and allergic diseases are
15 classified into types 1 to 4 hypersensitivity reactions based on response time and whether complements are involved. Type 1 hypersensitivity reactions include atopy, anaphylactic shock, bronchial asthma, urticaria, pollinosis and the like; type 2 hypersensitivity reactions include inadequate transfusion,
20 autoimmune hemolytic anemia, hemolytic anemia caused by drugs, granulocytopenia, thrombocytopenic purpura and the like; type 3 hypersensitivity reactions include erythema, lymphatic swelling, arthralgia, arthritis, nephritis, acute glomerulonephritis following streptococcal infection, and the like; and type 4
25 hypersensitivity reactions include chronic inflammation and the

like. To improve allergic diseases, it is preferable to remove allergens (house dust, mites, etc.) from the skin by showering or bathing and avoid allergen intake. However, when allergic diseases are not improved, drugs such as steroids, antihistamines, immunosuppressants or the like are used, which easily cause side effects such as skin atrophy, vasodilation, discoloration, purpura (steroids), drowsiness (antihistamines), kidney failure (immunosuppressants) and the like. Among the drugs developed so far, there is no drug that can completely cure allergies, and these drugs are expected to improve symptoms, but have the problem of causing significant side effects.

Obesity

Obesity is a metabolic disorder caused by the imbalance of calorie intake and consumption, and is caused by the increased size (hypertrophy) or increased number (hyperplasia) of *in vivo* adipocytes in morphological terms. Obesity is not only the most common malnutrition disorder in western society, and the prevalence of obesity in Korea is also rapidly increasing due to the improvement of eating habits and westernization of lifestyles. Therefore, the importance of treatment and prevention of obesity has been greatly emphasized. Obesity is an important factor that disturbs the individual in psychological terms and also increases the risk of various adult diseases in social terms. Obesity is known to be

directly related to the increased prevalence of various adult diseases such as type 2 diabetes, hypertension, hyperlipidemia, heart disease and the like (Cell 87:377, 1999), and diseases related to obesity are collectively referred to as metabolic syndrome or insulin resistance syndrome, and these diseases have been reported to cause arteriosclerosis and cardiovascular diseases. Obesity therapeutic agents known so far Xenical (Roche Pharmaceuticals, Switzerland), Reductil (Abbott, USA), Exolise (Arkopharma, France) and the like, and are largely classified into appetite suppressants, energy expenditure promoters, and fat absorption inhibitors. Most obesity therapeutic agents are appetite suppressants that suppress appetite by controlling the neurotransmitters associated with the hypothalamus. However, conventional therapeutic agents cause side effects such as heart diseases, respiratory diseases, neurological diseases and the like, and the persistence of their effects is also low. Thus, the development of improved obesity therapeutic agents is required. In addition, among currently developed products, there are little or no therapeutic agents that have satisfactory therapeutic effect without causing side effects, and thus the development of a new therapeutic agent for obesity is required.

Probiotics are collectively referred to as live microorganisms that improve the host's microbial environment in the gastrointestinal tract of animals, including humans, and

have beneficial effects on the host's health. In order to be effective as probiotics, it is necessary to have excellent acid resistance, bile resistance and adherence to epithelial cells, because most of these probiotics should reach the small intestine upon oral administration and must be adhered to the intestinal surface. Lactic acid bacteria are used as probiotics because they play a role in decomposing fibrous and complex proteins to make important nutrients while living in the digestive system of the human body. Lactic acid bacteria have been reported to exhibit effects such as maintenance of intestinal normal flora, improvement of intestinal flora, anti-diabetic and anti-hyperlipidemic effects, inhibition of carcinogenesis, inhibition of colitis, and nonspecific activity of the host's immune system. Among these lactic acid bacteria, *Lactobacillus* sp. strains are major members of normal microbial communities living in the bowel of the human body and have long been known to be important in maintaining a healthy digestive tract and vaginal environment. Currently, according to the U.S. Public Health Service guidelines, all the *Lactobacillus* strains deposited with the American Type Culture Collection (ATCC) are classified as 'Bio-Safety Level 1', which is recognized as having no known potential risk of causing disease in humans or animals. Meanwhile, lactic acid bacteria of kimchi that are involved in kimchi fermentation have been reported to have immune enhancement effects, antimicrobial effects, antioxidant

effects, anti-cancer effects, anti-obesity effects, hypertension preventive effects or constipation preventive effects [Hivak P, Odrska J, Ferencik M, Ebringer L, Jahnova E, Mikes Z. : One-year application of Probiotic strain
5 *Enterococcus faecium* M-74 decreases Serum cholesterol levels. : Bratisl lek Listy 2005; 106(2); 67-72; Agerholm-Larsen L. Bell ML. Grunwald GK. Astrup A. : The effect of a probiotic milk product on plasma cholesterol : a metaanalysis of short-term intervention studies ; Eur J Clin Nutr. 2000; 54(11) 856-860;
10 Renato Sousa, Jaroslava Helper, Jian Zhang, Strephen J Lewis and Wani O Li ; Effect of *Lactobacillus acidophilus* supernants on body weight and leptin expression in rats; BMC complementary and alternative medicine. 2008; 8(5)1-8].

Since various bioactive activities of lactic acid
15 bacteria were known, studies have recently been conducted to develop lactic acid bacterial strains that have excellent functions while being safe for the human and to apply these strains to medicines or functional foods. For example, Korean Patent Application Publication No. 10-2009-0116051 discloses
20 *Lactobacillus brevis* HY7401 having the effects of treating and preventing colitis. Furthermore, Korean Patent Application Publication No. 10-2006-0119045 discloses lactic acid bacteria for preventing or treating atopic dermatitis, which is selected from the group consisting of *Leuconostoc citreum* KACC91035,
25 *Leuconostoc mesenteroides* subsp. *mesenteroides* KCTC 3100 and

2016322617 03 Dec 2019

Lactobacillus brevis KCTC 3498. Furthermore, Korean Patent Application Publication No. 10-2013-0092182 discloses a functional health food for preventing alcoholic liver disease or relieving hangovers, which comprises *Lactobacillus brevis*

5 HD-01 (accession number: KACC91701P) having an excellent ability to decompose alcohol. In addition, Korean Patent Application Publication No. 10-2010-0010015 discloses a *Lactobacillus johnsonii* HFI 108 strain (KCTC 11356BP) having blood cholesterol lowering and anti-obesity activities. In
10 addition, Korean Patent Application Publication No. 10-2014-0006509 discloses a composition for preventing or treating obesity comprising a *Bifidobacterium longum* CGB-C11 strain (accession number: KCTC 11979BP) that produces conjugated linoleic acid as an active ingredient.

15 However, there has been no report of lactic acid bacteria-related technology capable of alleviating or treating all of intestinal flora disturbance, intestinal permeability syndrome, colitis, liver diseases, allergic diseases, obesity and the like, which are increasing in modern humans. Therefore,
20 there is a need to screen a novel strain having various functionalities and to develop medicines, functional foods and the like by use of this strain.

Reference to any prior art in the specification is not an acknowledgement or suggestion that this prior art forms part of
25 the common general knowledge in any jurisdiction or that this prior art could reasonably be expected to be combined with any other piece of prior art by a skilled person in the art.

【Disclosure】**【Technical Problem】**

The present invention has been made under the Background Art as described above, and one aspect the present invention
5 seeks to provide novel lactic acid bacteria having various physiological activities or functionalities required for probiotics, and the food and medicinal uses thereof.

In another aspect of the present invention, the invention
10 seeks to provide a novel lactic acid bacteria mixture capable of exhibiting various maximized physiological activities or functionalities, and the food and medicinal uses thereof.

【Technical Solution】

The present inventors have screened numerous lactic acid bacteria from kimchi or human feces, and have found that, among
15 these screened lactic acid bacteria, a certain *Lactobacillus* sp. strain, a certain *Bifidobacterium* sp. strain or a mixture thereof has an excellent effect on the improvement of intestinal damage such as intestinal permeability syndrome, liver injury such as fatty liver, allergic diseases such as
20 atopic dermatitis, inflammatory diseases such colitis, or obesity, thereby completing the present invention.

To achieve the above aspects, an embodiment of the present invention provides a lactic acid bacteria selected from *Lactobacillus brevis* comprising a 16S rDNA nucleotide sequence
25 represented by SEQ ID NO: 1, *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 3,

2016322617 03 Dec 2019

Lactobacillus plantarum comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 4, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 5 or *Bifidobacterium longum* comprising a 16S rDNA
5 nucleotide sequence represented by SEQ ID NO: 7. The *Lactobacillus brevis*, *Lactobacillus plantarum* or *Bifidobacterium longum* has antioxidant activity, β -glucuronidase-inhibitory activity, lipopolysaccharide (LPS) production-inhibitory activity or tight junction protein
10 expression-inducing activity. Another embodiment of the present invention provides a pharmaceutical composition for preventing or treating intestinal damage, liver injury, allergic disease, inflammatory disease or obesity comprising lactic acid bacteria selected from *Lactobacillus brevis* comprising a 16S rDNA
15 nucleotide sequence represented by SEQ ID NO: 1, *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 3, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 4, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide
20 sequence represented by SEQ ID NO: 5 or *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 7, a culture of the lactic acid bacteria, a lysate of the lactic acid bacteria or an extract of the lactic acid bacteria as an active ingredient. Still another embodiment of the
25 present invention provides a food composition for preventing or alleviating intestinal damage, liver injury, allergic disease, inflammatory disease or obesity comprising *Lactobacillus brevis* comprising a 16S rDNA nucleotide sequence represented by SEQ ID

NO: 1, *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 3, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 4, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 5 or *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 7, a culture of the lactic acid bacteria, a lysate of the lactic acid bacteria or an extract of the lactic acid bacteria as an active ingredient.

10 To achieve other aspects of the present invention, an embodiment of the present invention provides a mixture of two or more lactic acid bacteria selected from the group consisting of *Lactobacillus brevis* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 1, *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 3, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 4, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 5, and *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 7. The mixture of lactic acid bacteria has antioxidant activity, β -glucuronidase-inhibitory activity, lipopolysaccharide (LPS) production-inhibitory activity or tight junction protein expression-inducing activity. Another embodiment of the

present invention provides a composition for preventing or treating intestinal damage, liver injury, allergic disease, inflammatory disease or obesity comprising a mixture of two or more lactic acid bacteria selected from the group consisting of

5 *Lactobacillus brevis* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 1, *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 3, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 4, *Lactobacillus plantarum*

10 comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 5 and *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 7, a culture of the mixture of lactic acid bacteria, a lysate of the mixture of lactic acid bacteria or an extract of the mixture of lactic

15 acid bacteria as an active ingredient. Still another embodiment of the present invention provides a food composition for preventing or alleviating intestinal damage, liver injury, allergic disease, inflammatory disease or obesity comprising a mixture of two or more lactic acid bacteria selected from the

20 group consisting of *Lactobacillus brevis* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 1, *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 3, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID

25 NO: 4, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide

sequence represented by SEQ ID NO: 5 and *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 7, a culture of the mixture of lactic acid bacteria, a lysate of the mixture of lactic acid bacteria, or an extract of the mixture of lactic acid bacteria as an active ingredient.

【Advantageous Effects】

A certain *Lactobacillus* sp. strain or certain *Bifidobacterium* sp. strain according to the present invention is isolated from kimchi or human feces, and thus is highly safe, and has various physiological activities such as antioxidant activity, β -glucuronidase-inhibitory activity, lipopolysaccharide (LPS) production-inhibitory activity or tight junction protein expression-inducing activity. Accordingly, a certain *Lactobacillus* sp. strain, certain *Bifidobacterium* sp. strain or mixture thereof according to the present invention may be used as a functional food or medicinal material useful for preventing, alleviating or treating of intestinal damage, liver injury, allergic disease, inflammatory disease or obesity.

20 【Description of Drawings】

FIG. 1 is a graph showing the change in GOT value when lactic acid bacteria were administered to model animals having liver injury induced by D-galactosamine, in a first experiment of the present invention.

25 FIG. 2 is a graph showing the change in GPT value when

lactic acid bacteria were administered to model animals having liver injury induced by D-galactosamine, in a first experiment of the present invention.

FIG. 3 is a graph showing the change in MDA value when
5 lactic acid bacteria were administered to model animals having liver injury induced by D-galactosamine, in a first experiment of the present invention.

FIG. 4 is a graph showing the effect of lactic acid bacteria, screened in a first experiment of the present
10 invention, on the lipopolysaccharide (LPS)-induced inflammatory response of dendritic cells. The left graph in FIG. 4 shows the effect of lactic acid bacteria on cells not treated with LPS (lipopolysaccharide), and the right graph shows the effect of lactic acid bacteria on cells treated with LPS
15 (lipopolysaccharide).

FIG. 5 is a graph showing the effect of *Bifidobacterium longum* CH57 on the LPS (lipopolysaccharide)-induced inflammatory response of macrophages in a first experiment of the present invention.

20 FIG. 6 shows the results of analyzing the effect of *Lactobacillus brevis* CH23 on the differentiation of T cells (isolated from spleen) into Th17 cells or Treg cells by a fluorescence-activated cell sorting system in a first experiment of the present invention.

25 FIG. 7 shows the results of analyzing the effect of

Lactobacillus brevis CH23, *Bifidobacterium longum* CH57 or a mixture thereof on ZO-1 protein expression in CaCO2 cells in a first experiment of the present invention.

FIG. 8 shows the colon appearance or myeloperoxidase (MPO) activity indicating the effect of *Bifidobacterium longum* CH57 on model animals having acute colitis induced by TNBS, in a first experiment of the present invention.

FIG. 9 depicts histological images showing the effect of *Bifidobacterium longum* CH57 on model animals having acute colitis induced by TNBS, in a first experiment of the present invention.

FIG. 10 shows inflammation-related cytokine levels indicating the effect of *Bifidobacterium longum* CH57 on model animals having acute colitis induced by TNBS, in a first experiment of the present invention.

FIG. 11 shows the colon appearance or myeloperoxidase (MPO) activity indicating the effect of *Lactobacillus brevis* CH23 on model animals having acute colitis induced by TNBS, in a first experiment of the present invention.

FIG. 12 depicts histological images of colon, which show the effect of *Lactobacillus brevis* CH23 on model animals having acute colitis induced by TNBS, in a first experiment of the present invention.

FIG. 13 shows T-cell differentiation patterns indicating the effect of *Lactobacillus brevis* CH23 on model animals having

acute colitis induced by TNBS, in a first experiment of the present invention.

FIG. 14 shows inflammation-related cytokine levels indicating the effect of *Lactobacillus brevis* CH23 on model
5 animals having acute colitis induced by TNBS, in a first experiment of the present invention.

FIG. 15 shows the colon appearance or myeloperoxidase (MPO) activity indicating the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on
10 model animals having acute colitis induced by TNBS, in a first experiment of the present invention.

FIG. 16 depicts histological images showing the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on model animals having acute colitis induced by
15 TNBS, in a first experiment of the present invention.

FIG. 17 shows inflammation-related cytokine levels indicating the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on model animals having acute colitis induced by TNBS, in a first experiment of the
20 present invention.

FIG. 18 shows weight changes indicating the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on obesity-induced model animals, in a first experiment of the present invention.

25 FIG. 19 shows the appearance of colon, myeloperoxidase

(MPO) activity, histological images of colon, and the like indicating the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on obesity-induced model animals, in a first experiment of the present invention.

5 FIG. 20 shows inflammation-related cytokine levels indicating the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on obesity-induced model animals, in a first experiment of the present invention.

10 FIG. 21 shows inflammatory response markers indicating the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on obesity-induced model animals, in a first experiment of the present invention.

15 FIG. 22 shows the differentiation patterns of T cells into Th17 cells indicating the effect of lactic acid bacteria on model animals having acute colitis induced by TNBS, in a second experiment of the present invention.

20 FIG. 23 shows the differentiation patterns of T cells into Treg cells indicating the effect of lactic acid bacteria on model animals having acute colitis induced by TNBS, in a second experiment of the present invention.

FIG. 24 shows inflammatory response markers indicating the effect of lactic acid bacteria on model animals having acute colitis induced by TNBS, in a second experiment of the present invention.

25 FIG. 25 depicts images showing the effect of lactic acid

bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in a second experiment of the present invention.

FIG. 26 shows the gross gastric lesion score indicating the effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in a second experiment of the present invention.

FIG. 27 shows the ulcer index indicating the effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in a second experiment of the present invention.

FIG. 28 shows the histological activity index indicating the effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in a second experiment of the present invention.

FIG. 29 shows the myeloperoxidase (MPO) activity indicating the effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in a second experiment of the present invention.

FIG. 30 shows CXCL4 expression levels indicating the effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in a second experiment of the present invention.

FIG. 31 shows TNF- α expression levels indicating the effect of lactic acid bacteria on the stomach mucosa of mice

having gastric ulcer induced by ethanol, in a second experiment of the present invention.

【Mode for Invention】

As used herein, terms used in the present invention will
5 be defined.

As used herein, the term "culture" means a product obtained by culturing a microorganism in a known liquid medium or solid medium, and thus is intended to include a microorganism.

10 As used herein, the terms "pharmaceutically acceptable" and "sitologically acceptable" means neither significantly stimulating an organism nor inhibiting the biological activity and characteristics of an active material administered.

As used herein, the term "preventing" refers to all
15 actions that inhibit symptoms or delay the progression of a particular disease by administering the composition of the present invention.

As used herein, the term "treating" refers to all actions that alleviate or beneficially change the symptoms of a
20 particular disease by administering the composition of the present invention.

As used herein, the term "alleviating" refers to all actions that at least reduce a parameter related to the condition to be treated, for example, the degree of symptom.

As used herein, the term "administering" means providing the composition of the present invention to a subject by any suitable method. As used herein, the term "subject" means all animals, including humans, monkeys, dogs, goats, pigs or rats, 5 which have a particular disease whose symptoms can be alleviated by administering the composition of the present invention.

As used herein, the term "pharmaceutically effective amount" refers to an amount sufficient to treat diseases, at a 10 reasonable benefit/risk ratio applicable to any medical treatment. The pharmaceutically effective amount may be determined depending on factors including the kind of subject's disease, the severity of the disease, the activity of the drug, sensitivity to the drug, the time of administration, the route 15 of administration, excretion rate, the duration of treatment and drugs used in combination with the composition, and other factors known in the medical field.

Hereinafter, the present invention will be described in detail.

20 One aspect of the present invention is related to novel lactic acid bacteria having various physiological activities or to novel lactic acid bacteria mixture which may have increased physiological activities.

A novel lactic acid bacteria according to one embodiment 25 of the present invention is selected from *Lactobacillus brevis*

comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 1, *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 3, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 4, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 5 or *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 7, and has antioxidant activity, β -glucuronidase inhibitory activity, lipopolysaccharide (LPS) production inhibitory activity or tight junction protein expression-inducing activity.

The *Lactobacillus brevis* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 1 is an anaerobic bacillus isolated from kimchi, is positive to gram staining, can survive in a wide temperature range, low pHs and high salt concentrations, and produces glucosidase. Furthermore, the *Lactobacillus brevis* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 1 utilizes D-ribose, D-xylose, D-glucose, D-fructose, esculin, salicin, maltose, melibiose, 5-keto-gluconate and the like as carbon sources. In addition, the *Lactobacillus brevis* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 1 is preferably *Lactobacillus brevis* CH23 (accession number: KCCM 11762P). The *Bifidobacterium longum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 3 is an anaerobic bacillus

isolated from human feces, is positive to gram staining, and produces glucosidase. Furthermore, the *Bifidobacterium longum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 3 utilizes D-galactose, D-glucose, D-fructose and the like as carbon sources. In addition, the *Bifidobacterium longum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 3 is preferably *Bifidobacterium longum* CH57 (accession number: KCCM 11764P). The *Lactobacillus plantarum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 4 is an anaerobic bacillus isolated from kimchi and is positive to gram staining. Furthermore, the *Lactobacillus plantarum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 4 utilizes D-ribose, D-galactose, D-glucose, D-fructose, D-mannose, mannitol, sorbitol, N-acetylglucosamine, amygdalin, arbutin, esculin, salicin, cellobiose, maltose, melibiose, sucrose, trehalose, melezitose and the like as carbon sources. In addition, the *Lactobacillus plantarum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 4 is preferably *Lactobacillus plantarum* LC5 (accession number: KCCM 11800P). The *Lactobacillus plantarum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 5 is an anaerobic bacillus isolated from kimchi, and is positive to gram staining. Furthermore, the *Lactobacillus plantarum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 5 utilizes L-arabinose, D-ribose, D-glucose, D-fructose,

D-mannose, mannitol, sorbitol, N-acetyl-glucosamine, amygdalin, arbutin, esculin, salicin, cellobiose, maltose, lactose, melibiose, sucrose, trehalose, melezitose and the like as carbon sources. In addition, the *Lactobacillus plantarum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 5 is preferably *Lactobacillus plantarum* LC27 (accession number: KCCM 11801P). The *Bifidobacterium longum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 7 is an anaerobic bacillus isolated from human feces, and is positive to gram staining. Furthermore, the *Bifidobacterium longum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 7 utilizes L-arabinose, D-xylose, D-glucose, D-fructose, esculin, maltose, lactose, melibiose, sucrose and the like as carbon sources. In addition, the *Bifidobacterium longum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 7 is preferably *Bifidobacterium longum* LC67 (accession number: KCCM 11802P).

A mixture of lactic acid bacteria according to an embodiment of the present invention is a mixture of two or more lactic acid bacteria selected from *Lactobacillus brevis* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 1, *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 3, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 4, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide

sequence represented by SEQ ID NO: 5 and *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 7. In view of the synergistic effect of lactic acid bacteria, the mixture of lactic acid bacteria according to the embodiment of the present invention is preferably a combination of *Lactobacillus brevis* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 1 and *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 3. In addition, in view of the synergistic effect of lactic acid bacteria, the mixture of lactic acid bacteria according to the embodiment of the present invention is preferably a combination of one or more *Lactobacillus* sp. selected from *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 4 or *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 5; and *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 7. The mixture of lactic acid bacteria has higher antioxidant activity, β -glucuronidase inhibitory activity, lipopolysaccharide (LPS) production inhibitory activity or tight junction protein expression-inducing activity than a single lactic acid bacteria due to the synergistic effect of a specific *Lactobacillus* sp. strain and a specific *Bifidobacterium* sp. strain, and is more advantageous in terms of functional food and medicinal materials. In the mixture of

lactic acid bacteria according to the embodiment of the present invention, the *Lactobacillus brevis* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 1 is preferably *Lactobacillus brevis* CH23 (accession number: KCCM 11762P); the
5 *Bifidobacterium longum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 3 is preferably *Bifidobacterium longum* CH57 (accession number: KCCM 11764P); the *Lactobacillus plantarum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 4 is preferably
10 *Lactobacillus plantarum* LC5 (accession number: KCCM 11800P); the *Lactobacillus plantarum* comprising the 16S rDNA nucleotide sequence represented by SEQ ID NO: 5 is preferably *Lactobacillus plantarum* LC27 (accession number: KCCM 11801P); and the *Bifidobacterium longum* comprising the 16S rDNA
15 nucleotide sequence represented by SEQ ID NO: 7 is preferably *Bifidobacterium longum* LC67 (accession number: KCCM 11802P).

Another aspect of the present invention is related to various uses of the novel lactic acid bacteria or the novel lactic acid bacteria mixture. As the use of the novel lactic
20 acid bacteria, the present invention provides a composition for preventing, alleviating or treating intestinal damage, liver injury, allergic disease, inflammatory disease or obesity comprising a lactic acid bacteria from *Lactobacillus brevis* comprising a 16S rDNA nucleotide sequence represented by SEQ ID
25 NO: 1, *Bifidobacterium longum* comprising a 16S rDNA nucleotide

sequence represented by SEQ ID NO: 3, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 4, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 5 or *Bifidobacterium longum* 5 comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 7, a culture thereof, a lysate thereof or an extract thereof as an active ingredient. Furthermore, as the use of the novel lactic acid bacteria mixture, the present invention provides a composition for preventing, alleviating or treating 10 intestinal damage, liver injury, allergic disease, inflammatory disease or obesity comprising a mixture of two or more lactic acid bacteria selected from *Lactobacillus brevis* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 1, *Bifidobacterium longum* comprising a 16S rDNA nucleotide 15 sequence represented by SEQ ID NO: 3, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 4, *Lactobacillus plantarum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 5 and *Bifidobacterium longum* comprising a 16S rDNA nucleotide sequence represented by SEQ ID 20 NO: 7, a culture thereof, a lysate thereof or an extract thereof as an active ingredient. In the composition of the present invention, the technical characteristics of the *Lactobacillus brevis*, *Lactobacillus plantarum* and *Bifidobacterium longum* are as described above, and thus the 25 description thereof is omitted. The intestinal damage refers

to a condition in which the function of the intestines (particularly small intestine or large intestine) is abnormal due to intestinal flora disturbance or the like. Preferably, the intestinal damage is intestinal permeability syndrome.

5 Furthermore, the liver injury refers to a condition in which the function of the liver is abnormal due to external factors or internal factors. Preferably, the liver injury is selected from hepatitis, fatty liver or liver cirrhosis. Furthermore, the hepatitis includes all non-alcoholic hepatitis and

10 alcoholic hepatitis. Moreover, the fatty liver includes all non-alcoholic fatty liver and alcoholic fatty liver. Furthermore, the allergic disease is not limited in its kind if it caused by excessive immune responses of a living body, and is preferably selected from atopic dermatitis, asthma,

15 pharyngitis or chronic dermatitis. Furthermore, the inflammatory disease is not limited in its kind if it caused by inflammatory responses, and is preferably selected from gastritis, gastric ulcer, arthritis or colitis. Moreover, the arthritis includes rheumatoid arthritis. The colitis refers to

20 a condition in which inflammation occurred in the large intestine due to bacterial infection or pathological fermentation of intestinal contents. The colitis includes infectious colitis and non-infectious colitis. Specific examples of the colitis include inflammatory bowel diseases,

25 irritable bowel syndrome and the like. Furthermore, the

inflammatory bowel diseases include ulcerative colitis, Crohn's disease and the like.

In the present invention, a culture of the lactic acid bacteria or a culture of the lactic acid bacteria mixture is a
5 produced by culturing a certain strain or a mixture of strains in a medium. The medium may be selected from known liquid media or solid media, and may be, for example, MRS liquid medium, MRS agar medium or BL agar medium.

In the present invention, the composition may be embodied
10 as a pharmaceutical composition, a food additive, a food composition (particularly, a functional food composition), a feed additive or the like depending on the intended use or aspect. In addition, the content of the lactic acid bacteria or the lactic acid bacteria mixture as an active ingredient may
15 also be adjusted within a wide range depending on the specific type, intended use or aspect of the composition.

The content of the novel lactic acid bacteria, the novel lactic acid bacteria mixture, a culture thereof, a lysate thereof or an extract thereof as an active ingredient in the
20 pharmaceutical composition according to the present invention is not particularly limited. For example, the content may be 0.01 to 99 wt%, preferably 0.5 to 50 wt%, more preferably 1 to 30 wt%, based on the total weight of the composition. In addition, the pharmaceutical composition according to the
25 present invention may further contain, in addition to the

active ingredient, additives such as pharmaceutically acceptable carriers, excipients or diluents. Carriers, excipients and diluents, which may be contained in the pharmaceutical composition according to the present invention, 5 include lactose, dextrose, sucrose, sorbitol, mannitol, xylitol, erythritol, maltitol, starch, acacia gum, alginate, gelatin, calcium phosphate, calcium silicate, cellulose, methyl cellulose, microcrystalline cellulose, polyvinyl pyrrolidone, water, methyl hydroxybenzoate, propyl hydroxylbenzoate, talc, 10 magnesium stearate and mineral oil. In addition, the pharmaceutical composition according to the present invention may further contain, in addition to the novel lactic acid bacteria, the novel lactic acid bacteria mixture, a culture thereof, a lysate thereof or an extract thereof, one or more 15 active ingredients having the effect of preventing or treating intestinal damage, liver injury, allergic disease, inflammatory disease or obesity. The pharmaceutical composition according to the present invention may be prepared as formulations for oral administration or formulations for parenteral 20 administration, and the formulations may be prepared using diluents or excipients, such as fillers, extenders, binders, wetting agents, disintegrants, surfactants and the like, which are commonly used. Solid formulations for oral administration include tablets, pellets, powders, granules, capsules and the 25 like, and such solid formulations may be prepared by mixing the

active ingredient with at least one excipient, for example, starch, calcium carbonate, sucrose, lactose or gelatin. In addition to simple excipients, lubricants such as magnesium stearate or talc may also be used. Liquid formulations for oral administration include suspensions, solutions, emulsions and syrup, and may contain various excipients, for example, wetting agents, flavoring agents, aromatics, preservatives and the like, in addition to water and liquid paraffin which are frequently used simple diluents. Formulations for parenteral administration include sterilized aqueous solutions, non-aqueous solutions, suspensions, emulsions, freeze-dried preparations and suppositories. Propylene glycol, polyethylene glycol, plant oils such as olive oil, injectable esters such as ethyl oleate and the like may be used as non-aqueous solvents or suspending agents. As the base of the suppositories, witepsol, Macrogol, Tween 61, cacao butter, laurin fat, glycerogelatin and the like may be used. Furthermore, the composition may preferably be formulated depending on each disease or component by a suitable method known in the art or the method disclosed in Remington's Pharmaceutical Science (the latest edition), Mack Publishing Company, Easton PA. The pharmaceutical composition of the present invention may be administered orally or parenterally to mammals, including humans, according to a desired method. Routes for parenteral administration include skin external application,

intraperitoneal injection, intrarectal injection, subcutaneous injection, intravenous injection, intramuscular injection, intrathoracic injection or the like. The dose of the pharmaceutical composition of the present invention is not particularly limited as long as it is a pharmaceutically effective amount. The dose may vary depending on the patient's weight, age, sex, health condition, diet, administration time, administration mode, excretion rate and the severity of the disease. The daily dose of the pharmaceutical composition of the present invention is not particularly limited, but is preferably 0.1 to 3000 mg/kg based on an active ingredient, more preferably 1 to 2000 mg/kg based on an active ingredient and may be administered once or several times a day.

Furthermore, the content of the novel lactic acid bacteria, the novel lactic acid bacteria mixture, a culture thereof, a lysate thereof or an extract thereof as an active ingredient in the food composition according to the present invention is 0.01 to 99 wt%, preferably 0.1 to 50 wt%, more preferably 0.5 to 25 wt%, based on the total weight of the composition, but is not limited thereto. The food composition of the present invention may be in the form of pellets, powders, granules, infusions, tablets, capsules, liquid or the like, and specific examples of the food may include meats, sausages, breads, chocolates, candies, snacks, confectionaries, pizzas, ramens, other noodles, gums, dairy products including

ice creams, various kinds of soups, beverages, teas, functional water, drinks, alcoholic beverages, vitamin complexes and the like, and may include all health foods in a general sense. The food composition of the present invention may further contain
5 sitologically acceptable carriers, various flavoring agents or natural carbohydrates as additional ingredients, in addition to the active ingredient. Additionally, the food composition of the present invention may contain various nutrients, vitamins, electrolytes, flavoring agents, coloring agents, pectic acid
10 and its salt, alginic acid and its salt, an organic acid, a protective colloidal thickener, a pH adjusting agent, a stabilizer, a preservative, glycerin, alcohol, a carbonating agent used for carbonated drinks and the like. Additionally, the food composition of the present invention may contain fruit
15 flesh for preparing natural fruit juices, fruit juice drinks and vegetable drinks. These ingredients may be used independently or as a mixture. The above-described natural carbohydrates may include monosaccharides such as glucose and fructose, disaccharides such as maltose and sucrose,
20 polysaccharides such as dextrin and cyclodextrin and sugar alcohols such as xylitol, sorbitol, and erythritol. As a flavoring agent, a natural flavoring agent such as thaumatin or a stevia extract, or a synthetic flavoring agent such as saccharin or aspartame may be used.

Hereinafter, the present invention will be described in further detail with reference to examples. It is to be understood, however, that these examples are merely intended to clearly illustrate the technical characteristics of the present invention and do not limit the scope of the present invention.

I. First Experiment for Screening of Lactic Acid Bacteria and Evaluation of the Effects Thereof

1. Isolation and Identification of Lactic Acid Bacteria

(1) Isolation of Lactic Acid Bacteria from Kimchi

Each of Chinese cabbage kimchi, radish kimchi and green onion kimchi was crushed, and the crushed liquid was suspended in MRS liquid medium (MRS Broth; Difco, USA). Next, the supernatant was collected, transferred to MRS agar medium (Difco, USA) and cultured anaerobically at 37°C for about 48 hours, and then strains that formed colonies were isolated.

(2) Isolation of Lactic Acid Bacteria from Human Feces

Human feces were suspended in GAM liquid medium (GAM broth; Nissui Pharmaceutical, Japan). Next, the supernatant was collected, transferred to BL agar medium (Nissui Pharmaceutical, Japan) and cultured anaerobically at 37°C for about 48 hours, and then *Bifidobacterium* sp. strains that formed colonies were isolated.

(3) Identification of Screened Lactic Acid Bacteria

The physiological characteristics and 16S rDNA sequences of the strains isolated from kimchi or human feces were

analyzed to identify the species of the strains, and names were given to the strains. Table 1 below the control numbers and strain names of the lactic acid bacteria isolated from Chinese cabbage kimchi, radish kimchi, green onion kimchi and human feces.

Table 1

Control No.	Strain name	Control No.	Strain name
1	<i>Lactobacillus acidophilus</i> CH1	31	<i>Lactobacillus sakei</i> CH31
2	<i>Lactobacillus acidophilus</i> CH2	32	<i>Lactobacillus johnsonii</i> CH32
3	<i>Lactobacillus acidophilus</i> CH3	33	<i>Lactobacillus sakei</i> CH33
4	<i>Lactobacillus brevis</i> CH4	34	<i>Lactobacillus sakei</i> CH34
5	<i>Lactobacillus curvatus</i> CH5	35	<i>Lactobacillus plantarum</i> CH35
6	<i>Lactobacillus brevis</i> CH6	36	<i>Lactobacillus sanfranciscensis</i> CH36
7	<i>Lactobacillus casei</i> CH7	37	<i>Bifidobacterium pseudocatenulatum</i> CH37
8	<i>Lactobacillus planantrum</i> CH8	38	<i>Bifidobacterium pseudocatenulatum</i> CH38
9	<i>Lactobacillus sakei</i> CH9	39	<i>Bifidobacterium adolescentis</i> CH39
10	<i>Lactobacillus curvatus</i> CH10	40	<i>Bifidobacterium adolescentis</i> CH40
11	<i>Lactobacillus sakei</i> CH11	41	<i>Bifidobacterium adolescentis</i> CH41
12	<i>Lactobacillus curvatus</i> CH12	42	<i>Bifidobacterium animalis</i> CH42
13	<i>Lactobacillus plantarum</i> CH13	43	<i>Bifidobacterium animalis</i> CH43
14	<i>Lactobacillus fermentum</i> CH14	44	<i>Bifidobacterium bifidum</i> CH44
15	<i>Lactobacillus fermentum</i> CH15	45	<i>Bifidobacterium bifidum</i> CH45
16	<i>Lactobacillus gasseri</i> CH16	46	<i>Bifidobacterium breve</i> CH46
17	<i>Lactobacillus paracasei</i> CH17	47	<i>Bifidobacterium breve</i> CH47
18	<i>Lactobacillus helveticus</i> CH18	48	<i>Bifidobacterium breve</i> CH48
19	<i>Lactobacillus helveticus</i> CH19	49	<i>Bifidobacterium catenulatum</i> CH49
20	<i>Lactobacillus johnsonii</i> CH20	50	<i>Bifidobacterium catenulatum</i> CH50
21	<i>Lactobacillus johnsonii</i> CH21	51	<i>Bifidobacterium dentium</i> CH51
22	<i>Lactobacillus johnsonii</i> CH22	52	<i>Bifidobacterium infantis</i> CH52
23	<i>Lactobacillus brevis</i> CH23	53	<i>Bifidobacterium infantis</i> CH53
24	<i>Lactobacillus paracasei</i> CH24	54	<i>Bifidobacterium infantis</i> CH54
25	<i>Lactobacillus kimchi</i> CH25	55	<i>Bifidobacterium longum</i> CH55

26	<i>Lactobacillus gasseri</i> CH26	56	<i>Bifidobacterium longum</i> CH56
27	<i>Lactobacillus paracasei</i> CH27	57	<i>Bifidobacterium longum</i> CH57
28	<i>Lactobacillus pentosus</i> CH28	58	<i>Bifidobacterium longum</i> CH58
29	<i>Lactobacillus pentosus</i> CH29	59	<i>Bifidobacterium longum</i> CH59
30	<i>Lactobacillus reuteri</i> CH30	60	<i>Bifidobacterium longum</i> CH60

Among the strains shown in Table 1 above, *Lactobacillus brevis* CH23 was a gram-positive anaerobic bacillus, did not form spores, and could survive even under aerobic conditions. Furthermore, *Lactobacillus brevis* CH23 survived at 10 to 42°C and was an acid-resistant strain stable at pH 2 for 2 hours. Furthermore, *Lactobacillus brevis* CH23 survived even in 2% sodium chloride solution and actively produced glucosidase. In addition, to chemically classify *Lactobacillus brevis* CH23, the 16S rDNA thereof was analyzed, and as a result, it was shown that *Lactobacillus brevis* CH23 had a nucleotide sequence of SEQ ID NO: 1. The 16S rDNA nucleotide sequence of *Lactobacillus brevis* CH23 was identified by BLAST in the Genbank (<http://www.ncbi.nlm.nih.gov/>), and as a result, a *Lactobacillus brevis* strain having the same 16S rDNA nucleotide sequence as that of *Lactobacillus brevis* CH23 was not found, and *Lactobacillus brevis* CH23 showed a homology of 99% with the 16S rDNA sequence of *Lactobacillus brevis* strain FJ004.

Among the strains shown in Table 1 above, *Lactobacillus johnsonii* CH32 was a gram-positive anaerobic bacillus, did not form spores, and could survive under aerobic conditions.

Furthermore, *Lactobacillus johnsonii* CH32 survived stably at a temperature of up to 45°C, and was an acid-resistant strain stable in pH 2 for 2 hours. Moreover, *Lactobacillus johnsonii* CH32 actively produced β -glucosidase, but did not produce β -glucuronidase. In addition, to chemically classify *Lactobacillus johnsonii* CH32, the 16S rDNA thereof was analyzed, and as a result, it was shown that *Lactobacillus johnsonii* CH32 had a nucleotide sequence of SEQ ID NO: 2. The 16S rDNA nucleotide sequence of *Lactobacillus johnsonii* CH32 was identified by BLAST in Genebank (<http://www.ncbi.nlm.nih.gov/>), and as a result, a *Lactobacillus johnsonii* strain having the same 16S rDNA nucleotide sequence as that of *Lactobacillus johnsonii* CH32 was not found, and *Lactobacillus johnsonii* CH32 showed a homology of 99% with the 16S rDNA sequence of *Lactobacillus johnsonii* strain JCM 2012.

Among the strains shown in Table 1 above, *Bifidobacterium longum* CH57 was a gram-positive anaerobic bacillus, did not form spores, and showed very low viability under aerobic conditions. Furthermore, *Bifidobacterium longum* CH57 was thermally unstable. Furthermore, *Bifidobacterium longum* CH57 actively produced glucosidase, but did not produce β -glucuronidase. In addition, to chemically classify *Bifidobacterium longum* CH57, the 16S rDNA thereof was analyzed, and as a result, it was shown that *Bifidobacterium longum* CH57 had a nucleotide sequence of SEQ ID NO: 3. The 16S rDNA

nucleotide sequence of *Bifidobacterium longum* CH57 was identified by BLAST in the Genebank (<http://www.ncbi.nlm.nih.gov/>), and as a result, a *Bifidobacterium longum* strain having the same 16S rDNA
5 nucleotide sequence as that of *Bifidobacterium longum* CH57 was not found, and *Bifidobacterium longum* CH57 showed a homology of 99% with the 16S rDNA sequence of *Bifidobacterium longum* strain CBT-6.

In addition, among the physiological characteristics of
10 *Lactobacillus brevis* CH23, *Lactobacillus johnsonii* CH32 and *Bifidobacterium longum* CH57, the carbon source utilization was analyzed using a sugar fermentation by an API kit (model: API 50 CHL; manufactured by BioMerieux's, USA). Table 2 below shows the results of analyzing the carbon source utilization of
15 *Lactobacillus brevis* CH23; Table 3 below shows the results of analyzing the carbon source utilization of *Lactobacillus johnsonii* CH32; and Table 4 below shows the results of analyzing the carbon source utilization of *Bifidobacterium longum* CH57. In Tables 2, 3 and 4, "+" indicates the case in
20 which carbon source utilization is positive; "-" indicates the case in which carbon source utilization is negative; and "±" indicates the case in which carbon source utilization is ambiguous. As shown in Tables 2, 3 and 4 below, *Lactobacillus brevis* CH23, *Lactobacillus johnsonii* CH32 and *Bifidobacterium longum* CH57 showed carbon source utilization different from
25

that of other strains of the same species with respect to some carbon sources.

Table 2

Carbon source	Strain name		Carbon source	Strain name	
	<i>L. brevis</i> ¹⁾	<i>L. brevis</i> CH23		<i>L. brevis</i> ¹⁾	<i>L. brevis</i> CH23
glycerol	-	-	salicin	+	+
erythritol	-	-	cellobiose	+	-
D-arabinose	-	-	maltose	+	+
L-arabinose	+	-	lactose	+	-
D-ribose	+	+	melibiose	-	+
D-xylose	+	+	sucrose	+	-
L-xylose	-	-	trehalose	+	-
D-adonitol	-	-	inulin	+	-
methyl- β -D-xylopyranoside	-	-	melezitose	+	-
D-galactose	+	-	raffinose	-	-
D-glucose	+	+	starch	-	-
D-fructose	+	+	glycogen	-	-
D-mannose	+	-	xylitol	-	-
L-sorbose	-	-	gentiobiose	+	-
L-rhamnose	-	-	D-turanose	+	-
dulcitol	+	-	D-lyxose	-	-
inositol	-	-	D-tagatose	+	-
mannitol	+	-	D-fucose	-	-
sorbitol	+	-	L-fucose	-	-
α -methyl-D-mannoside	-	-	D-arabitol	-	-
α -methlyl-D-glucoside	-	-	L-arabitol	-	-
N-acetyl-glucosamine	+	±	gluconate	+	±
amygdalin	+	-	2-keto-gluconate	-	-
arbutin	+	-	5-keto-gluconate	-	+
esculin	+	+			

¹⁾ Suriasih K., Aryanta WR, MahardikaG, Astawa NM.

5 Microbiological and Chemical Properties of Kefir Made of Bali Cattle Milk. Food Science and Quality Management 2012;6:112-22.

Table 3

Carbon source	Strain name		Carbon source	Strain name	
	<i>L. johnsonii</i> ²⁾	<i>L. johnsonii</i> CH32		<i>L. johnsonii</i> ²⁾	<i>L. johnsonii</i> CH32
glycerol	-	-	salicin	-	-
erythritol	-	-	cellobiose	+	-
D-arabinose	-	-	maltose	-	+
L-arabinose	-	-	lactose	-	+
D-ribose	-	-	melibiose	+	-
D-xylose	-	-	sucrose	+	+
L-xylose	-	-	trehalose	+	-
D-adonitol	-	-	inulin	-	-
methyl- β -D-xylopyranoside	-	-	melezitose	-	-
D-galactose	-	-	raffinose	+	-
D-glucose	-	+	starch	-	-
D-fructose	-	+	glycogen	-	-
D-mannose	+	+	xylitol	-	-
L-sorbose	-	-	gentiobiose	-	+
L-rhamnose	-	-	D-turanose	-	-
dulcitol	-	-	D-lyxose	-	-
inositol	-	-	D-tagatose	-	-
mannitol	-	-	D-fucose	-	-
sorbitol	-	-	L-fucose	-	-
α -methyl-D-mannoside	-	-	D-arabitol	-	-
α -methyl-D-glucoside	-	-	L-arabitol	-	-
N-acetyl-glucosamine	+	+	gluconate	-	-
amygdalin	-	-	2-keto-gluconate	-	-
arbutin	-	-	5-keto-gluconate	-	-
esculin	-	-			

²⁾ Pridmore RD, Berger B, Desiere F, Vilanova D, Barretto C, Pittet AC, Zwahlen MC, Rouvet M, Altermann E, Barrangou R, Mollet B, Mercenier A, Klaenhammer T, Arigoni F, Schell MA. The

genome sequence of the probiotic intestinal bacterium *Lactobacillus johnsonii* NCC 533. Proc Natl Acad Sci U S A. 2004 Feb 24;101(8):2512-7.

Table 4

Carbon source	Strain name		Carbon source	Strain name	
	<i>B. longum</i> ³⁾	<i>B. longum</i> CH57		<i>B. longum</i> ³⁾	<i>B. longum</i> CH57
glycerol	±	-	salicin	±	-
erythritol	-	-	cellobiose	±	±
D-arabinose	-	-	maltose	-	-
L-arabinose	-	-	lactose	-	-
D-ribose	±	-	melibiose	-	-
D-xylose	-	-	sucrose	+	±
L-xylose	-	-	trehalose	±	-
D-adonitol	-	-	inulin	-	-
methyl-β-D-xylopyranoside	-	-	melezitose	-	-
D-galactose	+	+	raffinose	-	-
D-glucose	+	+	starch	-	-
D-fructose	+	+	glycogen	-	-
D-mannose	-	-	xylitol	-	-
L-sorbose	-	-	gentiobiose	-	-
L-rhamnose	-	-	D-turanose	-	-
dulcitol	-	-	D-lyxose	-	-
inositol	-	-	D-tagatose	-	-
mannitol	+	-	D-fucose	-	-
sorbitol	-	-	L-fucose	-	-
α-methyl-D-mannoside	-	-	D-arabitol	-	-
α-methyl-D-glucoside	-	-	L-arabitol	-	-
N-acetyl-glucosamine	±	-	gluconate	±	-
amygdalin	-	-	2-keto-gluconate	-	-
arbutin	±	-	5-keto-gluconate	-	-
esculin	-	-			

3) Lukacova D, Karovucova J, Greifova M, Greif G, Sovcikova A, Kohhajdova Z. In vitro testing of selected probiotic

characteristics of *Lactobacillus plantarum* and *Bifidobacterium longum*. Journal of Food and Nutrition Research 2006; 45: 77-83.

(4) Information on Deposition of Lactic Acid Bacteria

The present inventors deposited *Lactobacillus brevis* CH23
5 with the Korean Culture Center of Microorganisms (address: Yurim Building, 45, Hongjenae 2ga-gil, Seodaemun-gu, Seoul, Korea), an international depositary authority, on September 1, 2015 under accession number KCCM 11762P. Furthermore, the present inventors deposited *Lactobacillus johnsonii* CH32 with
10 the Korean Culture Center of Microorganisms (address: Yurim Building, 45, Hongjenae 2ga-gil, Seodaemun-gu, Seoul, Korea), an international depositary authority, on September 1, 2015, under accession number KCCM 11763P. Furthermore, the present inventors deposited *Bifidobacterium longum* CH57 with the Korean
15 Culture Center of Microorganisms (address: Yurim Building, 45, Hongjenae 2ga-gil, Seodaemun-gu, Seoul, Korea), an international depositary authority, on September 1, 2015 under accession number KCCM 11764P.

2. Evaluation of the Effect of Lactic Acid Bacteria on 20 **Alleviation of Intestinal Damage or Intestinal Permeability**

In order to evaluate the effect of the lactic acid bacteria isolated from kimchi or human feces, on the alleviation of intestinal damage or internal permeability, the antioxidant activity, lipopolysaccharide (LPS) production
25 inhibitory activity, β -glucuronidase (harmful intestinal

enzyme) inhibitory activity and tight junction protein expression-inducing activity of the lactic acid bacteria were measured.

(1) Experimental Methods

5 * Antioxidant activity

DPPH (2,2-diphenyl-1-picrylhydrazyl) was dissolved in ethanol to a concentration of 0.2 mM to prepare a DPPH solution. A lactic acid bacteria suspension (1×10^8 CFU/ml) or a vitamin C solution (1 g/ml) was added to 0.1 ml of the DPPH solution and
10 cultured at 37°C for 20 minutes. The culture was centrifuged at 3000 rpm for 5 minutes, and the supernatant was collected. Next the absorbance of the supernatant at 517 nm was measured, and the antioxidant activity of the lactic acid bacteria was calculated.

15 * Lipopolysaccharide (LPS) production-inhibitory activity

0.1 g of human fresh feces was suspended in 0.9 ml of sterile physiological saline and diluted 100-fold with general anaerobic medium to prepare a fecal suspension. 0.1 ml of the fecal suspension and 0.1 ml of lactic acid bacteria (1×10^4 or
20 1×10^5 CFU) were added to 9.8 ml of sterile anaerobic medium (Nissui Pharmaceuticals, Japan) and cultured anaerobically for 24 hours. Next, the culture was sonicated for about 1 hour to disrupt the outer cell membrane of the bacteria, and centrifuged at 5000×g, and the supernatant was collected. Next,
25 the content of LPS (lipopolysaccharide) (which is a typical

endotoxin) in the supernatant was measured by a LAL (Limulus Amoebocyte Lysate) assay kit (manufactured by Cape Cod Inc., USA). In addition, in order to evaluate the *E. coli* proliferation inhibitory activity of the lactic acid bacteria, the culture obtained through the same experiment as described above was diluted 1000-fold and 100000-fold and cultured in DHL medium, and then the number of *E. coli* cells was counted.

* β -glucuronidase inhibitory activity

0.1 ml of 0.1 mM p-nitrophenyl- β -D-glucuronide solution, 0.2 ml of 50 mM phosphate buffered saline and 0.1 ml of a lactic acid bacteria suspension (prepared by suspending of a lactic acid bacteria culture in 5 ml of physiological saline) were placed in a reactor and subjected to an β -glucuronidase enzymatic reaction, and 0.5 ml of 0.1 mM NaOH solution was added to stop the reaction. Next, the reaction solution was centrifuged at 3000 rpm for 5 minutes, and the supernatant was collected. Then, the absorbance of the supernatant at 405 nm was measured.

* Tight junction protein expression-inducing activity

Caco2 cells obtained from the Korean Cell Line Bank were cultured in RPMI 1640 medium for 48 hours, and then the cultured Caco2 cells were dispensed to each well of a 12-well plate at a density of 2×10^6 cells/well. Next, each well was treated with 1 μ g of LPS (lipopolysaccharide) or a combination of 1 μ g of LPS (lipopolysaccharide) and 1×10^3 CFU of lactic acid

bacteria and incubated for 24 hours. Next, the cultured cells were collected from each well, and the expression level of tight junction protein ZO-1 in the cells was measured by an immunoblotting method.

5 (2) Experimental Results

 The antioxidant activity, lipopolysaccharide (LPS) production inhibitory activity, β -glucuronidase inhibitory activity and tight junction protein expression-inducing activity of the lactic acid bacteria isolated from kimchi or
10 human feces were measured, and the results of the measurement are shown in Tables 5 and 6 below. As shown in Tables 5 and 6 below, *Lactobacillus curvatus* CH5, *Lactobacillus sakei* CH11, *Lactobacillus brevis* CH23, *Lactobacillus johnsonii* CH32, *Bifidobacterium pseudocatenulatum* CH38 and *Bifidobacterium*
15 *longum* CH57 had excellent antioxidant activity, strongly inhibited lipopolysaccharide (LPS) production and β -glucuronidase activity, and strongly induced the expression of tight junction protein. These lactic acid bacteria have an excellent antioxidant effect, have an excellent effect of
20 inhibiting the enzymatic activity of intestinal flora's harmful bacteria associated with inflammation and carcinogenesis, inhibit the production of endotoxin LPS (lipopolysaccharide) produced by intestinal flora's harmful bacteria, and induce the expression of tight junction protein. Thus, these lactic acid
25 bacteria can improve intestinal permeability syndrome.

Table 5

Control No.	Strain name	Antioxidant activity	Beta-glucuronidase inhibitory activity	LPS production inhibitory activity	Tight junction protein expression inducing activity
1	<i>Lactobacillus acidophilus</i> CH1	+	+	-	-
2	<i>Lactobacillus acidophilus</i> CH2	+	+	+	-
3	<i>Lactobacillus acidophilus</i> CH3	+	+	+	-
4	<i>Lactobacillus brevis</i> CH4	+	+	-	-
5	<i>Lactobacillus curvatus</i> CH5	+++	+	+++	++
6	<i>Lactobacillus brevis</i> CH6	+	+	-	-
7	<i>Lactobacillus casei</i> CH7	+	+	-	-
8	<i>Lactobacillus planantrum</i> CH8	+	+	-	-
9	<i>Lactobacillus sakei</i> CH9	-	+	-	-
10	<i>Lactobacillus curvatus</i> CH10	-	+	-	-
11	<i>Lactobacillus sakei</i> CH11	+++	+	+++	++
12	<i>Lactobacillus curvatus</i> CH12	-	+	-	+
13	<i>Lactobacillus plantarum</i> CH13	-	+	-	-
14	<i>Lactobacillus fermentum</i> CH14	-	+	-	-
15	<i>Lactobacillus fermentum</i> CH15	+++	+	++	-
16	<i>Lactobacillus gasseri</i> CH16	+	+	-	-
17	<i>Lactobacillus paracasei</i> CH17	+	+	-	-
18	<i>Lactobacillus helveticus</i> CH18	+	+	-	-
19	<i>Lactobacillus helveticus</i> CH19	+	+	-	-
20	<i>Lactobacillus johnsonii</i> CH20	+	+	-	+
21	<i>Lactobacillus johnsonii</i> CH21	+	+	-	+
22	<i>Lactobacillus johnsonii</i> CH22	+	+	-	+
23	<i>Lactobacillus brevis</i> CH23	+++	+	++	++
24	<i>Lactobacillus paracasei</i> CH24	+	+	-	-
25	<i>Lactobacillus kimchi</i> CH25	+	+	-	-
26	<i>Lactobacillus gasseri</i> CH26	+	+	-	-
27	<i>Lactobacillus paracasei</i> CH27	+	+	-	+
28	<i>Lactobacillus pentosus</i> CH28	+	+	-	-
29	<i>Lactobacillus pentosus</i> CH29	+	+	-	-
30	<i>Lactobacillus reuteri</i> CH30	+	-	-	-

Table 6

Control No.	Strain name	Antioxidant activity	Beta-glucuronidase inhibitory activity	LPS production inhibitory activity	Tight junction protein expression inducing activity
31	<i>Lactobacillus sakei</i> CH31	-	+	-	+
32	<i>Lactobacillus johnsonii</i> CH32	+++	+	++	++
33	<i>Lactobacillus sakei</i> CH33	+	+	-	+
34	<i>Lactobacillus sakei</i> CH34	+	+	-	+
35	<i>Lactobacillus plantarum</i> CH35	+	+	+	+
36	<i>Lactobacillus sanfranciscensis</i> CH36	+	+	+	+
37	<i>Bifidobacterium pseudocatenulatum</i> CH37	-	+	-	+
38	<i>Bifidobacterium pseudocatenulatum</i> CH38	+++	+	++	++
39	<i>Bifidobacterium adolescentis</i> CH39	-	+	-	+
40	<i>Bifidobacterium adolescentis</i> CH40	-	+	+++	+
41	<i>Bifidobacterium adolescentis</i> CH41	+	+	-	+
42	<i>Bifidobacterium animalis</i> CH42	+	+	-	-
43	<i>Bifidobacterium animalis</i> CH43	+	+	-	-
44	<i>Bifidobacterium bifidum</i> CH44	+	+	-	-
45	<i>Bifidobacterium bifidum</i> CH45	+	+	-	-
46	<i>Bifidobacterium breve</i> CH46	+	-	-	-
47	<i>Bifidobacterium breve</i> CH47	+	+	-	+
48	<i>Bifidobacterium breve</i> CH48	+	+	-	+
49	<i>Bifidobacterium catenulatum</i> CH49	+	+	-	++
50	<i>Bifidobacterium catenulatum</i> CH50	-	+	-	-
51	<i>Bifidobacterium dentium</i> CH51	+	-	-	-
52	<i>Bifidobacterium infantis</i> CH52	-	+	-	-
53	<i>Bifidobacterium infantis</i> CH53	-	+	-	-

54	<i>Bifidobacterium infantis</i> CH54	+	+	-	-
55	<i>Bifidobacterium longum</i> CH55	+	+	-	+
56	<i>Bifidobacterium longum</i> CH56	+++	+	++	+
57	<i>Bifidobacterium longum</i> CH57	+++	+	+++	++
58	<i>Bifidobacterium longum</i> CH58	+	+	+	+
59	<i>Bifidobacterium longum</i> CH59	+	+	+	+
60	<i>Bifidobacterium longum</i> CH60	+	-	+	+

* The final concentration of lactic acid bacteria in measurement of antioxidant activity: 1×10^4 CFU/ml; the concentration of lactic acid bacteria added for measurement of beta-glucuronidase inhibitory activity and lipopolysaccharide (LPS) production inhibitory activity: 1×10^4 CFU/ml; the concentration of lactic acid bacteria in measurement of tight junction protein expression-inducing activity: 1×10^4 CFU/ml.

* Criteria for measurement of various activities of lactic acid bacteria: very strongly (+++; >90%); strongly (++; >60-90%); weakly (+; >20-60%); not or less than 20% (-; <20%).

3. Evaluation of the Effect of Lactic Acid Bacteria on Alleviation of Liver injury

Based on evaluation of the effect of the lactic acid bacteria on the alleviation of intestinal damage or intestinal permeability syndrome, the following seven strains were selected: *Lactobacillus curvatus* CH5, *Lactobacillus sakei* CH11, *Lactobacillus fermentum* CH15, *Lactobacillus brevis* CH23, *Lactobacillus johnsonii* CH32, *Bifidobacterium pseudocatenulatum* CH38 and *Bifidobacterium longum* CH57. The effect of each of these selected strains or a mixture of these strains on the

alleviation of liver injury was evaluated using various liver injury model animals.

(1) Measurement of the liver injury-alleviating effect of lactic acid bacteria by an experiment using model animals

5 having liver injury induced by D-galactosamine

1) Experimental Method

Mice (C57BL/6, male) were divided into several groups, each consisting of 6 animals. D-Galactosamine was administered intraperitoneally to the test animals of groups other than a
10 normal group at a dose of 800 mg/kg to induce liver injury. From 2 hours after intraperitoneal administration of D-galactosamine, 1×10^9 CFU of lactic acid bacteria were administered orally to the test animals of groups other than the normal group and the negative control group, once a day for
15 3 days. In addition, silymarin in place of lactic acid bacteria was administered orally to the test animals of the positive control group at a dose of 100 mg/kg, once a day for 3 days. At 6 hours after the last administration of the drug, blood was taken from the heart. The taken blood was allowed to
20 stand at room temperature for 60 minutes, and centrifuged at 3,000 rpm for 15 minutes to separate serum. The GPT (glutamic pyruvate transaminase) and GOT (glutamic oxalacetic transaminase) levels in the separated serum were measured using a blood assay kit (ALT & AST measurement kit; Asan Pharm. Co.,
25 Korea).

In addition, liver tissue was dissected from the test animals, and the amount of malondialdehyde (MDA) present in the liver tissue was measured. Malondialdehyde is a marker of lipid peroxidation. Specifically, 0.5 g of the dissected liver tissue was added to a 16-fold volume of RIPA solution (0.21M mannitol, 0.1M EDTA-2Na, 0.07M sucrose, 0.01M Trizma base), and then homogenized using a homogenizer. The homogenized solution was centrifuged at 3,000 rpm for 10 minutes, and the liver homogenate was collected. 0.5 ml of the liver homogenate was added to 0.4 ml of 10% SDS, incubated at 37°C for 30 minutes, and cooled, and then 3 ml of 1% phosphate buffer and 1 ml of 0.6% TBA were added thereto and heated on a water bath at 100°C for 45 minutes to develop the sample solution. The developed sample solution was added to and mixed with 4 ml of n-butanol, and then centrifuged at 3000 rpm for 10 minutes, and the supernatant was collected. The absorbance of the collected supernatant at 535 nm was measured to quantify MDA. In addition, a calibration curve for MDA measurement was plotted using 1,1,3,3-tetraethoxypropane.

20 2) Experimental Results

FIG. 1 is a graph showing the change in GOT value when lactic acid bacteria were administered to model animals having liver injury induced by D-galactosamine; FIG. 2 is a graph showing the change in GPT value when lactic acid bacteria were administered to model animals having liver injury induced by D-

galactosamine; and FIG. 3 is a graph showing the change in MDA value when lactic acid bacteria were administered to model animals having liver injury induced by D-galactosamine.

On the x-axis of FIGS. 1 to 3, "Nor" indicates a normal group; "Con" indicates a negative control group in which any drugs were not administered to model animals having liver injury induced by D-galactosamine; "ch11" indicates a group administered with *Lactobacillus sakei* CH11; "ch15" indicates a group administered with *Lactobacillus fermentum* CH15; "ch23" indicates a group administered with *Lactobacillus brevis* CH23; "ch32" indicates a group administered with *Lactobacillus johnsonii* CH32; "ch38" indicates a group administered with *Bifidobacterium pseudocatenulatum* CH38; "ch57" indicates a group administered with *Bifidobacterium longum* CH57; "ch57+ch11" indicates a group administered with a lactic acid bacteria mixture prepared by mixing *Bifidobacterium longum* CH57 and *Lactobacillus sakei* CH11 in the same amount; "ch57+ch23" indicates a group administered with a lactic acid bacteria mixture prepared by mixing *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 in the same amount; "ch57+ch32" indicates a group administered with a lactic acid bacteria mixture prepared by mixing *Bifidobacterium longum* CH57 and *Lactobacillus johnsonii* CH32 in the same amount; and "SM" indicates a positive control group administered with silymarin instead of lactic acid bacteria.

As shown in FIGS. 1 to 3, when each of *Lactobacillus brevis* CH23, *Lactobacillus johnsonii* CH32 and *Bifidobacterium longum* CH57 was administered to the model animals in which GOT, GPT and MAD values increased due to liver injury, the liver injury was alleviated. Particularly, when a lactic acid bacteria mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 or a lactic acid bacteria mixture of *Bifidobacterium longum* CH57 and *Lactobacillus johnsonii* CH32 was administered, the liver injury was greatly alleviated. In addition, specific lactic acid bacteria or a mixture of lactic acid bacteria selected therefrom showed a better effect on the alleviation of liver injury than silymarin which is used as a drug for treating liver injury. These results suggest that specific lactic acid bacteria or a mixture of lactic acid bacteria selected therefrom is effective in alleviating fatty liver induced by alcohol and high-fat diets, or in alleviating liver diseases resulting from oxidative stress.

(2) Measurement of the liver injury-alleviating effect of lactic acid bacteria by an experiment using model animals having liver injury induced by tert-butylperoxide

1) Experimental Method

Mice (C57BL/6, male) were divided into several groups, each consisting of 6 animals. Tert-butylperoxide was administered intraperitoneally to the test animals of groups other than a normal group in an amount of 2.5 mmol/kg to induce

liver injury. From 2 hours after intraperitoneal administration of tert-butylperoxide, 2×10^9 CFU of lactic acid bacteria were administered orally to the test animals of groups other than the normal group and the negative control group, once a day for 3 days. In addition, silymarin in place of lactic acid bacteria was administered orally to the test animals of the positive control group at a dose of 100 mg/kg, once a day for 3 days. At 6 hours after the last administration of the drug, blood was taken from the heart. The taken blood was allowed to stand at room temperature for 60 minutes, and centrifuged at 3,000 rpm for 15 minutes to separate serum. The GPT (glutamic pyruvate transaminase) and GOT (glutamic oxalacetic transaminase) levels in the separated serum were measured using a blood assay kit ((ALT & AST measurement kit; Asan Pharm. Co., Korea).

2) Experimental Results

Table 7 below shows the changes in GOT and GPT values when lactic acid bacteria were administered to the model animals having liver injury induced by tert-butylperoxide. As shown in Table 7 below, *Lactobacillus brevis* CH23, *Lactobacillus johnsonii* CH32 and *Bifidobacterium longum* CH57 showed excellent effects on the alleviation of liver injury compared to silymarin, and a lactic acid bacteria mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 or a lactic acid bacteria mixture of *Bifidobacterium longum* CH57

and *Lactobacillus johnsonii* CH32 showed a better effect on the alleviation of liver injury.

Table 7

Test groups	GOT(IU/L)	GPT(IU/L)
Normal group	36.1	26.3
Negative control group	84.1	96.1
Group administered with CH23	58.0	74.2
Group administered with CH32	53.0	70.5
Group administered with CH57	57.6	71.2
Group administered with CH57+CH23	48.6	64.3
Group administered with CH57+CH32	51.2	68.4
Group administered with silymarin	61.7	69.1

5 In Table 7 above, "CH23" indicates *Lactobacillus brevis* CH23; "CH32" indicates *Lactobacillus johnsonii* CH32; "CH57" indicates *Bifidobacterium longum* CH57; "CH57+CH23" indicates a lactic acid bacteria mixture prepared by mixing *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 in the same amount; 10 and "CH57+CH32" indicates a lactic acid bacteria mixture prepared by mixing *Bifidobacterium longum* CH57 and *Lactobacillus johnsonii* CH32 in the same amount.

4. Evaluation of the Effect of Lactic Acid Bacteria on Alleviation of Allergy

15 (1) Measurement of the Inhibition of Degranulation by Lactic Acid Bacteria

The RBL-2H3 cell line (rat mast cell line, the Korean Cell Line Bank, Cat. No.22256) was cultured with DMEM (Dulbeccos' modified Eagle's medium, Sigma, 22256) containing 10% FBS

(fetal bovine serum) and L-glutamine in a humidified 5% CO₂ incubator at 37°C. The cells contained in the culture medium were floated using trypsin-EDTA solution, and the floated cells were isolated, collected and used in the experiment. The
5 collected RBL-2H3 cells were dispensed into a 24-well plate at a density of 5×10⁵ cells/well and sensitized by incubation with 0.5 µg/ml of mouse monoclonal IgE for 12 hours. The sensitized cells were washed with 0.5 ml of siraganian buffer (119 mM NaCl, 5 mM KCl, 0.4 mM MgCl₂, 25 mM PIPES, 40 mM NaOH, pH 7.2), and
10 then incubated with 0.16 ml of siraganian buffer (supplemented with 5.6 mM glucose, 1 mM CaCl₂, 0.1% BSA) at 37°C for 10 minutes. Next, lactic acid bacteria as a test drug were added to the cell culture to a concentration of 1×10⁴ CFU/ml, or 0.04 ml of DSCG (disodium cromoglycate) as a control drug was added
15 to the cell culture, and after 20 minutes, the cells were activated with 0.02 ml of antigen (1 µg/ml DNP-BSA) at 37°C for 10 minutes. Next, the cell culture was centrifuged at 2000 rpm for 10 minutes, and the supernatant was collected. 0.025 ml of the collected supernatant was transferred to a 96-well plate,
20 and then 0.025 ml of 1mM p-NAG (a solution of p-nitrophenyl-N-acetyl-β-D-glucosamide in 0.1M citrate buffer, pH 4.5) was added thereto, and then the mixture was allowed to react at 37°C for 60 minutes. Next, the reaction was stopped by addition of 0.2 ml of 0.1M Na₂CO₃/NaHCO₃, and the absorbance at
25 405 nm was measured by an ELISA analyzer.

(2) Measurement of the Inhibition of Itching by Lactic Acid Bacteria

BALB/c mice were divided into several groups, each consisting of 5 animals. 1×10^9 CFU of lactic acid bacteria as a test drug were administered orally to test groups other than a normal group and a control group, once a day for 3 days, or DSCG (disodium cromoglycate) or Azelastine as a control drug was administered orally in an amount of 0.2 mg/mouse, once a day for 3 days. At 1 hour after the last administration of the drug, the mice were allowed to stand in an observation cage (24 cm $\times$ 22 cm $\times$ 24 cm) for 10 minutes so as to be acclimated to the environment, and then the back portion of the head was shaved. Next, the mice of the normal group were injected with physiological saline, and the mice of the other test groups were injected with an itching inducer (50 μ g of compound 48/80; Sigma, USA) by a 29-gauge needle. Next, each mouse was immediately confined in an observation cage, and the itching behavior was observed under the unattended condition by recording with an 8-mm video camera (SV-K80, Samsung) for 1 hour. Scratching the injection area with the back foot was regarded as the itching behavior, and scratching other portions was not regarded.

(3) Measurement of the Inhibition of Vascular Permeability by Lactic Acid Bacteria

It is known that itching-induced areas have increased vascular permeability. This experiment was performed in order to examine whether lactic acid bacteria could efficiently inhibit vascular permeability caused by various compounds.

5 According to the same method as the above-described experiment for measurement of itching inhibitory activity, the drug was administered to the same mice. Next, physiological saline was injected subcutaneously into the back portion of the head of the mice of the normal group, and an itching inducer (50 µg of

10 compound 48/80; Sigma, USA) was injected subcutaneously into the back portion of the head of the mice of the other test group. Next, 0.2 ml of 1% Evans blue solution (Sigma, USA) was injected into the tail vein, and after 1 hour, the mice were euthanized. The skin of the subcutaneously injected area was

15 dissected, and incubated in 1 ml of 1N KOH at 37°C overnight. On the next day, the incubated skin tissue was mixed with 4 ml of 0.6N phosphoric acid-acetone (5:13) mixture and centrifuged at 3000 rpm for 15 minutes, and the supernatant was collected and measured for absorbance at 620 nm. Inhibition (%) of

20 vascular permeability was calculated using the following equation:

Inhibition (%) = {1- [absorbance of area treated with drug and itching inducer - absorbance of area not treated with itching inducer]/[absorbance of area treated with itching

inducer - absorbance of area not treated with itching inducer]]
x 100

(4) Experimental Results

Table 8 below shows the results of measuring the
5 degranulation inhibition rate, itching inhibition rate and
capillary permeability inhibition rate of the lactic acid
bacteria. In Table 8 below, "CH5" indicates *Lactobacillus*
curvatus CH5; "CH11" indicates *Lactobacillus sakei* CH11; "CH15"
indicates *Lactobacillus fermentum* CH15; "CH23" indicates
10 *Lactobacillus brevis* CH23; "CH32" indicates *Lactobacillus*
johnsonii CH32; "CH38" indicates *Bifidobacterium*
pseudocatenulatum CH38; "CH57" indicates *Bifidobacterium longum*
CH57; "CH57+CH11" indicates a lactic acid bacteria mixture
prepared by mixing *Bifidobacterium longum* CH57 and
15 *Lactobacillus sakei* CH11 in the same amount; "CH57+CH23"
indicates a lactic acid bacteria mixture prepared by mixing
Bifidobacterium longum CH57 and *Lactobacillus brevis* CH23 in
the same amount; and "CH57+CH32" indicates a lactic acid
bacteria mixture prepared by mixing *Bifidobacterium longum* CH57
20 and *Lactobacillus johnsonii* CH32 in the same amount.

As shown in Table 8 below, *Lactobacillus curvatus* CH5,
Lactobacillus brevis CH23, *Lactobacillus johnsonii* CH32 and
Bifidobacterium longum CH57 effectively inhibited the
degranulation of basophils, and *Bifidobacterium longum* CH57
25 very strongly inhibited itching and vascular permeability. In

addition, in comparison with these lactic acid bacteria alone, a mixture of these lactic acid bacteria, particularly a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 or a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus johnsonii* CH32 showed higher degranulation inhibition rate, itching inhibition rate and vascular permeability inhibition rate. Thus, it can be seen that these lactic acid bacteria or mixtures thereof can very effectively alleviate allergic atopy, asthma, pharyngitis, chronic dermatitis or the like.

10 Table 8

Drug	Inhibition (%)		
	Degranulation	Itching	vascular permeability
None	0	2	1
CH5	53	46	45
CH11	47	46	45
CH15	48	42	42
CH23	54	47	47
CH32	52	45	46
CH38	44	45	42
CH57	55	55	52
CH57+CH11	59	56	54
CH57+CH23	63	62	61
CH57+CH32	61	58	56
DSCG (disodium cromoglycate)	62	25	37
Azelastine	-	65	68

5. In Vitro Evaluation of the Anti-inflammatory Effect and Intestinal Permeability Inhibitory Effect of Lactic Acid Bacteria

(1) Isolation of Dendritic Cells and Measurement of Inflammatory Marker

Immune cells were isolated from the bone marrow of C57BL/6 mice (male, 20-23 g) by use of RPMI 1640 (containing 10% FBS, 1% antibiotics, 1% glutamax, 0.1% mercaptoethanol). The isolated cells were treated with RBC lysis buffer, washed, dispensed into each well of a 24-well plate, treated with GM-CSF and IL-4 at a ratio of 1:1000, and cultured. On 5 days of the culturing, the medium was replaced with fresh medium, and on 8 days, the cells were collected and used as dendritic cells. Next, the dendritic cells were seeded on a 24-well plate at a density of 0.5×10^6 cells/well and treated with lactic acid bacteria (test substance) and the inflammation inducer LPS (lipopolysaccharide) for 2 hours or 24 hours, and then the supernatant and the cells were collected. Using the collected supernatant, the expression levels of IL-10 and IL-12 were measured by an immunoblotting method.

FIG. 4 is a graph showing the effect of lactic acid bacteria screened in the present invention, on the lipopolysaccharide (LPS)-induced inflammatory response of dendritic cells. The left graph in FIG. 4 shows the effect of lactic acid bacteria on cells not treated with LPS (lipopolysaccharide), and the right graph shows the effect of lactic acid bacteria on cells treated with LPS (lipopolysaccharide). In addition, on the x-axis of FIG. 4,

"Nor" indicates a group not treated with the test lactic acid bacteria and the inflammation inducer LPS (lipopolysaccharide); "LPS" indicates a group treated with the inflammation inducer LPS (lipopolysaccharide); "ch11" indicates a group treated with
5 *Lactobacillus sakei* CH11; "ch15" indicates a group treated with *Lactobacillus fermentum* CH15; "ch23" indicates a group treated with *Lactobacillus brevis* CH23; "ch32" indicates a group treated with *Lactobacillus johnsonii* CH32; "ch38" indicates a group treated with *Bifidobacterium pseudocatenulatum* CH38;
10 "ch57" indicates a group treated with *Bifidobacterium longum* CH57; "ch57+ch11" indicates a group treated with a lactic acid bacteria mixture prepared by mixing *Bifidobacterium longum* CH57 and *Lactobacillus sakei* CH11 in the same amount; "ch57+ch23" indicates a group treated with a lactic acid bacteria mixture
15 prepared by mixing *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 in the same amount; and "ch57+ch32" indicates a group treated with a lactic acid bacteria mixture prepared by mixing *Bifidobacterium longum* CH57 and *Lactobacillus johnsonii* CH32 in the same amount.

20 As shown in FIG. 4, *Lactobacillus sakei* CH11, *Lactobacillus brevis* CH23 and *Lactobacillus johnsonii* CH32 induced IL-10 production of the dendritic cells obtained by differentiation after isolation from the bone marrow, effectively inhibited LPS (lipopolysaccharide)-induced
25 production of IL-12, and also the effects were increased when

used in combination with *Bifidobacterium longum* CH57. In particular, a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 exhibited the best effect on the inhibition of inflammation. When dendritic cells are controlled, Treg cells (regulatory T cells) can be efficiently controlled. For this reason, the lactic acid bacteria screened in the present invention can effectively alleviate chronic inflammatory diseases such as colitis, autoimmune diseases such as rheumatoid arthritis and the like.

(2) Isolation of Macrophages and Measurement of Inflammatory Marker

6-week-old C57BL/6J male mice (20-23g) were purchased from RaonBio Co., Ltd. 2 ml of 4% sterile thioglycolate was administered into the abdominal cavity of each mouse, and after 96 hours, the mice were anesthetized, and 8 ml of RPMI 1640 medium was administered into the abdominal cavity of each mouse. After 5 to 10 minutes, the RPMI medium (including macrophages) in the abdominal cavity of the mice was taken out, centrifuged at 1000 rpm for 10 minutes, and then washed twice with RPMI 1640 medium. The macrophages were seeded on a 24-well plate at a density of 0.5×10^6 cells/well and treated with the test substance lactic acid bacteria and the inflammation inducer LPS (lipopolysaccharide) for 2 hours or 24 hours, and then the supernatant and the cells were collected. The collected cells were homogenized in buffer (Gibco). Using the collected

supernatant, the expression levels of cytokines such as TNF- α and IL-1 β were measured by an ELISA kit. In addition, using the collected cells, the expression levels of p65 (NF-kappa B), p-p65 (phosphor-NF-kappa B) and β -actin were measured by an immunoblotting method. Specifically, 50 μ g of the supernatant was taken and electrophoresed on SDS 10% (w/v) polyacrylamide gel for 1 hour and 30 minutes. The electrophoresed sample was transferred to a nitrocellulose membrane under the conditions of 100 V and 400 mA for 1 hour and 10 minutes. The sample-transferred nitrocellulose membrane was blocked with 5% skim milk for 30 minutes, and then washed three times with PBS-Tween for 5 minutes each time, and incubated with a 1:100 dilution of primary antibody (Santa Cruz Biotechnology, USA) overnight. Next, the membrane was washed three times for 10 minutes each time, and incubated with a 1:1000 dilution of secondary antibody (Santa Cruz Biotechnology, USA) for 1 hour and 20 minutes. Next, the membrane was washed three times for 15 minutes each time, and it was developed by fluorescence and visualized.

FIG. 5 is a graph showing the effect of *Bifidobacterium longum* CH57 on the LPS (lipopolysaccharide)-induced inflammatory response of macrophages. As shown in FIG. 5, *Bifidobacterium longum* CH57 effectively inhibited the LPS (lipopolysaccharide)-induced inflammatory response.

(3) Isolation of T cells from Spleen and Measurement of Differentiation into Th17 Cells or Treg Cells

Spleen was separated from C56BL/6J mice, crushed suitably, and suspended in 10% FCS-containing RPMI 1640 medium, and CD4 T cells were isolated therefrom using a CD4 T cell isolation kit (MiltenyiBiotec, Bergisch Gladbach, Germany). The isolated CD4 T cells were seeded in a 12-well plate at a density of 5×10^5 cells/well, and anti-CD3 (1 $\mu\text{g/ml}$, MiltenyiBiotec, Bergisch Gladbach, Germany) and anti-CD28 (1 $\mu\text{g/ml}$, MiltenyiBiotec, Bergisch Gladbach, Germany) were added thereto, or anti-CD3 (1 $\mu\text{g/ml}$, MiltenyiBiotec, Bergisch Gladbach, Germany), anti-CD28 (1 $\mu\text{g/ml}$, MiltenyiBiotec, Bergisch Gladbach, Germany), recombinant IL-6 (20 ng/ml, MiltenyiBiotec, Bergisch Gladbach, Germany) and recombinant transforming growth factor beta (1 ng/ml, MiltenyiBiotec, Bergisch Gladbach, Germany) were added. While the cells were cultured, 1×10^3 or 1×10^5 CFU of the lactic acid bacteria were added thereto, and the cells were cultured for 4 days. Next, the cells of the culture were stained with anti-FoxP3 or anti-IL-17A antibody, and the distribution of Th17 and Treg cells was analyzed using a FACS (fluorescence-activated cell sorting) system (C6 Flow Cytometer® System, San Jose, CA, USA).

FIG. 6 shows the results of analyzing the effect of *Lactobacillus brevis* CH23 on the differentiation of T cells (isolated from spleen) into Th17 cells or Treg cells by a

fluorescence-activated cell sorting system. As shown in FIG. 6, *Lactobacillus brevis* CH23 inhibited the differentiation of T cells into Th17 cells (T helper 17 cells) and promoted the differentiation of T cells into Treg cells. These results suggest that *Lactobacillus brevis* CH23 can effectively alleviate inflammatory diseases such as colitis and arthritis.

(4) Measurement of the Effect of Lactic Acid Bacteria on ZO-1 Protein Expression of CaCO₂ Cells

Caco2 cells obtained from the Korean Cell Line Bank were cultured in RPMI 1640 medium for 48 hours, and then the cultured Caco2 cells were dispensed into a 12-well plate at a density of 2×10^6 cells/well. Next, each well was treated with 1 μ g of LPS (lipopolysaccharide) alone or a combination of 1 μ g of LPS (lipopolysaccharide) and 1×10^3 CFU or 1×10^5 CFU of lactic acid bacteria, and then incubated for 24 hours. Next, the cultured cells were collected from each well, and the expression level of tight junction protein ZO-1 was measured by an immunoblotting method.

FIG. 7 shows the results of analyzing the effect of *Lactobacillus brevis* CH23, *Bifidobacterium longum* CH57 or a mixture thereof on ZO-1 protein expression of CaCO₂ cells. In FIG. 7, "CH23" indicates *Lactobacillus brevis* CH23; "CH57" indicates *Bifidobacterium longum* CH57; "mix" indicates a lactic acid bacteria mixture prepared by mixing *Bifidobacterium longum* CH57 and *Lactobacillus johnsonii* CH32 in the same amount. As

shown in FIG. 7, treatment with *Lactobacillus brevis* CH23 and *Bifidobacterium longum* CH57 increased the expression of tight junction protein ZO-1, and treatment with a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus johnsonii* CH32 synergistically increased the expression of tight junction protein ZO-1. When the expression of tight junction protein increases, *in vivo* penetration of toxic substances can be blocked, thereby prevents the worsening of colitis, arthritis and liver injury.

10 **6. In Vivo Evaluation of the Anti-inflammatory and Colitis-Alleviating Effects of Lactic Acid Bacteria**

(1) Test Animals

5-Week-old C57BL/6 male mice (24-27g) were purchased from OrientBio, and housed under controlled environmental conditions (humidity: 50±10%, temperature: 25±2°C, 12-hr light/12-hr dark cycle), and then used in the experiment. As feed, standard experimental feed (Samyang, Korea) was used, and the animals had access to drinking water *ad libitum*. In all the experiments, one group consisted of 6 animals.

20 (2) Colitis Induction by TNBS and Sample Administration

One group of the test animals was used as a normal group, and the test animals of the other groups were treated with 2,4,6-trinitrobenzenesulfonic acid (TNBS) to induce acute colitis. Specifically, the test animals were lightly anesthetized with ether, and then a mixture solution of 2.5 g

of TNBS (2,4,6-trinitrobenzene sulfonic acid) and 100 ml of 50% ethanol was administered into the colon through the anal in an amount of 0.1 ml each time by use of a 1-ml round-tip syringe, lifted vertically and maintained for 30 seconds, thereby inducing inflammation. On the other hand, the normal group was administered orally with 0.1 ml of saline. On the next day, the lactic acid bacteria or the lactic acid bacteria mixture as a test sample was suspended in saline and administered orally to each mouse in an amount of 2.0×10^9 CFU, once a day for three days. On the next day following the end of sample administration, the animals were killed with carbon dioxide, and a colon portion ranging from the cecum to the site just before the anus was dissected and used. Meanwhile, the test animals of the normal group were orally administered with saline alone instead of the lactic acid bacteria. In addition, the test animals of the negative control group were orally administered with saline alone instead of the lactic acid bacteria after the induction of colitis by TNBS. Furthermore, the test animals of the positive control group were orally administered with 50 mg/kg of sulfasalazine, which is a drug for treating colitis, instead of the lactic acid bacteria.

(3) Macroscopic Analysis of Colon

The length and appearance of the dissected colon were observed, and the appearance was analyzed by scoring according to the criteria (Hollenbach et al., 2005, Criteria for Degree

of Colitis) shown in Table 9 below. After complete removal of colon contents, the colon tissue was washed with saline. A portion of the washed colon tissue was fixed with 4% formaldehyde solution in order to use it as a pathological tissue sample, and the remainder was freeze-stored at -80°C for molecular biological analysis.

Table 9

Macroscopic score	Criteria
0	Any ulcer and inflammation are not found.
1	Edema without bleeding is found.
2	Ulcer with edema is found.
3	Ulcer and inflammation are found at only one site.
4	Ulcer and inflammation are found at two or more sites.
5	Ulcer has an increased size of 2 cm or more.

(4) Measurement of Myeloperoxidase (MPO) Activity

100 mg of colon tissue was homogenized in 200 µl of 10 mM potassium phosphate buffer (pH 7.0) containing 0.5% hexadecyl trimethyl ammonium bromide. The homogenized tissue was centrifuged at 10,000×g and 4°C for 10 minutes, and the supernatant was collected. 50 µl of the supernatant was added to 0.95 ml of a reaction solution (containing 1.6mM tetramethyl benzidine and 0.1mM H₂O₂) and allowed to react at 37°C, and the absorbance at 650 nm was measured at various time points during the reaction. To calculate myeloperoxidase (MPO) activity, 1 µmol/ml of peroxide produced by the reaction was used as 1 unit.

(5) Measurement of Inflammatory Marker

Using a Western blotting method, inflammatory markers such as p-p65, p65, iNOS, COX-2 and β -actin were measured. Specifically, according to the same method as the experiment for measurement of myeloperoxidase (MPO) activity, a
5 supernatant was obtained. 50 μ g of the supernatant was taken and electrophoresed on SDS 10% (w/v) polyacrylamide gel for 1 hour and 30 minutes. The electrophoresed sample was transferred to a nitrocellulose membrane under the conditions of 100 V and 400 mA for 1 hour and 10 minutes. The sample-
10 transferred nitrocellulose membrane was blocked with 5% skim milk for 30 minutes, and then washed three times with PBS-Tween for 5 minutes each time, and incubated with a 1:100 dilution of primary antibody (Santa Cruz Biotechnology, USA) overnight. Next, the membrane was washed three times for 10 minutes each
15 time, and incubated with a 1:1000 dilution of secondary antibody (Santa Cruz Biotechnology, USA) for 1 hour and 20 minutes. Next, the membrane was washed three times for 15 minutes each time, and it was developed by fluorescence and visualized.

20 In addition, inflammation-related cytokines such as TNF- α , IL-1 β and the like were measured using an ELISA kit.

(6) Experimental Results

FIG. 8 shows the colon appearance or myeloperoxidase (MPO) activity indicating the effect of *Bifidobacterium longum* CH57
25 on model animals having acute colitis induced by TNBS; FIG. 9

depicts histological images of colon, which show the effect of *Bifidobacterium longum* CH57 on model animals having acute colitis induced by TNBS; and FIG. 10 shows inflammation-related cytokine levels indicating the effect of *Bifidobacterium longum* CH57 on model animals having acute colitis induced by TNBS. In FIGS. 8 to 10, "NOR" indicates a normal group; "TNBS" indicates a negative control group; "CH57" indicates a group administered with *Bifidobacterium longum* CH57; and "SS50" indicates a group administered with sulfasalazine. As shown in FIGS. 8 to 10, *Bifidobacterium longum* CH57 effectively alleviated colitis in view of the weight of the model animals having TNBS-induced acute colitis, the colitis markers, the colon length, myeloperoxidase (MPO) activity, and the like, and showed a better effect on the alleviation of colitis than sulfasalazine. In addition, *Bifidobacterium longum* CH57 inhibited inflammatory cytokine production and increased the production of the anti-inflammatory cytokine IL-10 in the model animals having TNBS-induced acute colitis.

FIG. 11 shows the colon appearance or myeloperoxidase (MPO) activity indicating the effect of *Lactobacillus brevis* CH23 on model animals having acute colitis induced by TNBS; FIG. 12 depicts histological images of colon, which show the effect of *Lactobacillus brevis* CH23 on model animals having acute colitis induced by TNBS; FIG. 13 shows T-cell differentiation patterns indicating the effect of *Lactobacillus brevis* CH23 on

model animals having acute colitis induced by TNBS; and FIG. 14 shows inflammation-related cytokine levels indicating the effect of *Lactobacillus brevis* CH23 on model animals having acute colitis induced by TNBS. In FIGS. 11 to 14, "N" indicates a normal group; "TNBS" indicates a negative control group; "CH23" indicates a group administered with *Lactobacillus brevis* CH23; and "SS" indicates a group administered with sulfasalazine. As shown in FIGS. 11 to 14, *Lactobacillus brevis* CH23 effectively alleviated colitis in view of the weight of the model animals having TNBS-induced acute colitis, the colitis markers, the colon length, myeloperoxidase (MPO) activity and the like, and showed a better effect on the alleviation of colitis than sulfasalazine. In addition, *Lactobacillus brevis* CH23 inhibited the differentiation of T cells into Th17 cells and induced the differentiation of T cells into Treg cells in the model animals having TNBS-induced acute colitis. Furthermore, *Lactobacillus brevis* CH23 inhibited inflammatory cytokine production and increased the production of the anti-inflammatory cytokine IL-10 in the model animals having TNBS-induced acute colitis.

FIG. 15 shows the colon appearance or myeloperoxidase (MPO) activity indicating the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on model animals having acute colitis induced by TNBS; FIG. 16 depicts histological images showing the effect of a mixture of

Bifidobacterium longum CH57 and *Lactobacillus brevis* CH23 on model animals having acute colitis induced by TNBS; and FIG. 17 shows inflammation-related cytokine levels indicating the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on model animals having acute colitis induced by TNBS. In FIGS. 15 to 17, "NOR" indicates a normal group; "TNBS" indicates a negative control group; "BL" indicates a group administered with a lactic acid bacteria mixture prepared by mixing *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 in the same amount; and "SS50" indicates a group administered with sulfasalazine. As shown in FIGS. 15 to 17, a lactic acid bacteria mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 significantly improved effects against the reduced weight of the model animals having TNBS-induced acute colitis, increased colitis marker levels, shortened colon lengths and increased myeloperoxidase (MPO) activity, and the effect thereof on the alleviation of colitis was significantly better than that of sulfasalazine. In addition, the lactic acid bacteria mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 significantly inhibited inflammatory cytokine production and dramatically increased the production of anti-inflammatory cytokine IL-10 in the model animals having TNBS-induced acute colitis.

7. In Vivo Evaluation of the Obesity-Reducing and Anti-

Inflammatory Effects of Lactic Acid Bacteria

(1) Experimental Method

A total of 24 C57BL6/J mice were purchased from RaonBio Co., Ltd., and acclimated with chow diet (Purina) under the conditions of temperature of $20\pm 2^{\circ}\text{C}$, humidity of $50\pm 10\%$ and 12-hr light/12-hr dark cycle for 1 week. Next, the test animals were divided into three groups (LFD, HFD, and HFD+BL), each consisting of 8 animals, the LFD group was fed with a normal diet (LFD, 10% of calories from fat; Research, NJ, USA) for 4 weeks, and the HFD group and the HFD+BL group were fed with a high-fat diet (HFD, 60% of calories from fat; Research, NJ, USA) for 4 weeks. Next, the LFD group was orally administered with PBS while fed with the normal diet for 4 weeks. Furthermore, the HFD group was orally administered with PBS while fed with the high-fat diet for 4 weeks. In addition, the HFD+BL group was orally administered with a PBS suspension of 2×10^9 CFU of a lactic acid bacteria mixture while fed with the high-fat diet for 4 weeks. The lactic acid bacteria mixture was prepared by mixing *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 in the same amount.

(2) Analysis of the Anti-obesity Effect and Anti-inflammatory Effect of Lactic Acid Bacteria Mixture

The anti-obesity effect of the lactic acid bacteria mixture was analyzed through weight change. In addition, the anti-inflammatory effect of the lactic acid bacteria mixture

was analyzed using the same method as that used in the experiment on the model animals having TNBS-induced acute colitis.

(3) Experimental Results

5 FIG. 18 shows weight changes indicating the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on obesity-induced model animals; FIG. 19 shows the appearance of colon, myeloperoxidase (MPO) activity, histological images of colon and the like, which indicate the effect of a mixture of *Bifidobacterium longum* CH57 and
10 *Lactobacillus brevis* CH23 on obesity-induced model animals; FIG. 20 shows inflammation-related cytokine levels indicating the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on obesity-induced model animals; and
15 FIG. 21 shows inflammatory response markers indicating the effect of a mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23 on obesity-induced model animals. As shown in FIGS. 18 to 21, the lactic acid bacteria mixture of *Bifidobacterium longum* CH57 and *Lactobacillus brevis* CH23
20 greatly reduced the increased weight, increased colitis marker levels and increased myeloperoxidase (MPO) activity of the model animals having obesity induced by the high-fat diet, and inhibited the development of colitis. In addition, the lactic acid bacteria mixture of *Bifidobacterium longum* CH57 and
25 *Lactobacillus brevis* CH23 greatly inhibited inflammatory

cytokine production and increased the production of anti-inflammatory cytokine IL-10 in the model animals having obesity induced by the high-fat diet.

II. Second Experiment for Screening of Lactic Acid

5 Bacteria and Evaluation of the Effects Thereof

1. Isolation and Identification of Lactic Acid Bacteria

(1) Isolation of Lactic Acid Bacteria from Kimchi

Each of Chinese cabbage kimchi, radish kimchi and green onion kimchi was crushed, and the crushed liquid was suspended
10 in MRS liquid medium (MRS Broth; Difco, USA). Next, the supernatant was collected, transferred to MRS agar medium (Difco, USA), and cultured anaerobically at 37°C for about 48 hours, and then *Bifidobacterium longum* strains that formed colonies were separated according to shape.

15 (2) Isolation of Lactic Acid Bacteria from Human Feces

Human feces were suspended in GAM liquid medium (GAM broth; Nissui Pharmaceutical, Japan). Next, the supernatant was collected, transferred to BL agar medium (Nissui Pharmaceutical, Japan) and cultured anaerobically at 37°C for
20 about 48 hours, and then *Bifidobacterium* sp. strains that formed colonies were isolated.

(3) Identification of Screened Lactic Acid Bacteria

The gram-staining characteristics, physiological characteristics and 16S rDNA sequences of the strains isolated
25 from kimchi or human feces were analyzed to identify the

species of the strains, and names were given to the strains. Table 10 below the control numbers and strain names of the lactic acid bacteria isolated from Chinese cabbage kimchi, radish kimchi and green onion kimchi, and Table 11 below shows
5 the control numbers and strain names of the lactic acid bacteria isolated from human feces.

Table 10

Control No.	Strain name	Control No.	Strain name
1	<i>Lactobacillus plantarum</i> LC1	26	<i>Lactobacillus plantarum</i> LC26
2	<i>Lactobacillus plantarum</i> LC2	27	<i>Lactobacillus plantarum</i> LC27
3	<i>Lactobacillus plantarum</i> LC3	28	<i>Lactobacillus plantarum</i> LC28
4	<i>Lactobacillus plantarum</i> LC4	29	<i>Lactobacillus plantarum</i> LC29
5	<i>Lactobacillus plantarum</i> LC5	30	<i>Lactobacillus plantarum</i> LC30
6	<i>Lactobacillus plantarum</i> LC6	31	<i>Lactobacillus plantarum</i> LC31
7	<i>Lactobacillus plantarum</i> LC7	32	<i>Lactobacillus plantarum</i> LC32
8	<i>Lactobacillus plantarum</i> LC8	33	<i>Lactobacillus plantarum</i> LC33
9	<i>Lactobacillus plantarum</i> LC9	34	<i>Lactobacillus plantarum</i> LC34
10	<i>Lactobacillus plantarum</i> LC10	35	<i>Lactobacillus plantarum</i> LC35
11	<i>Lactobacillus plantarum</i> LC11	36	<i>Lactobacillus plantarum</i> LC36
12	<i>Lactobacillus plantarum</i> LC12	37	<i>Lactobacillus plantarum</i> LC37
13	<i>Lactobacillus plantarum</i> LC13	38	<i>Lactobacillus plantarum</i> LC38
14	<i>Lactobacillus plantarum</i> LC14	39	<i>Lactobacillus plantarum</i> LC39
15	<i>Lactobacillus plantarum</i> LC15	40	<i>Lactobacillus plantarum</i> LC40
16	<i>Lactobacillus plantarum</i> LC16	41	<i>Lactobacillus plantarum</i> LC41
17	<i>Lactobacillus plantarum</i> LC17	42	<i>Lactobacillus plantarum</i> LC42
18	<i>Lactobacillus plantarum</i> LC18	43	<i>Lactobacillus plantarum</i> LC43
19	<i>Lactobacillus plantarum</i> LC19	44	<i>Lactobacillus plantarum</i> LC44
20	<i>Lactobacillus plantarum</i> LC20	45	<i>Lactobacillus plantarum</i> LC45
21	<i>Lactobacillus plantarum</i> LC21	46	<i>Lactobacillus plantarum</i> LC46
22	<i>Lactobacillus plantarum</i> LC22	47	<i>Lactobacillus plantarum</i> LC47
23	<i>Lactobacillus plantarum</i> LC23	48	<i>Lactobacillus plantarum</i> LC48
24	<i>Lactobacillus plantarum</i> LC24	49	<i>Lactobacillus plantarum</i> LC49
25	<i>Lactobacillus plantarum</i> LC25	50	<i>Lactobacillus plantarum</i> LC50

Table 11

Control No.	Strain name	Control No.	Strain name
51	<i>Bifidobacterium longum</i> LC51	76	<i>Bifidobacterium longum</i> LC76
52	<i>Bifidobacterium longum</i> LC52	77	<i>Bifidobacterium longum</i> LC77
53	<i>Bifidobacterium longum</i> LC53	78	<i>Bifidobacterium longum</i> LC78
54	<i>Bifidobacterium longum</i> LC54	79	<i>Bifidobacterium longum</i> LC79
55	<i>Bifidobacterium longum</i> LC55	80	<i>Bifidobacterium longum</i> LC80
56	<i>Bifidobacterium longum</i> LC56	81	<i>Bifidobacterium longum</i> LC81
57	<i>Bifidobacterium longum</i> LC57	82	<i>Bifidobacterium longum</i> LC82
58	<i>Bifidobacterium longum</i> LC58	83	<i>Bifidobacterium longum</i> LC83
59	<i>Bifidobacterium longum</i> LC59	84	<i>Bifidobacterium longum</i> LC84
60	<i>Bifidobacterium longum</i> LC60	85	<i>Bifidobacterium longum</i> LC85
61	<i>Bifidobacterium longum</i> LC61	86	<i>Bifidobacterium longum</i> LC86
62	<i>Bifidobacterium longum</i> LC62	87	<i>Bifidobacterium longum</i> LC87
63	<i>Bifidobacterium longum</i> LC63	88	<i>Bifidobacterium longum</i> LC88
64	<i>Bifidobacterium longum</i> LC64	89	<i>Bifidobacterium longum</i> LC89
65	<i>Bifidobacterium longum</i> LC65	90	<i>Bifidobacterium longum</i> LC90
66	<i>Bifidobacterium longum</i> LC66	91	<i>Bifidobacterium longum</i> LC91
67	<i>Bifidobacterium longum</i> LC67	92	<i>Bifidobacterium longum</i> LC92
68	<i>Bifidobacterium longum</i> LC68	93	<i>Bifidobacterium longum</i> LC93
69	<i>Bifidobacterium longum</i> LC69	94	<i>Bifidobacterium longum</i> LC94
70	<i>Bifidobacterium longum</i> LC70	95	<i>Bifidobacterium longum</i> LC95
71	<i>Bifidobacterium longum</i> LC71	96	<i>Bifidobacterium longum</i> LC96
72	<i>Bifidobacterium longum</i> LC72	97	<i>Bifidobacterium longum</i> LC97
73	<i>Bifidobacterium longum</i> LC73	98	<i>Bifidobacterium longum</i> LC98
74	<i>Bifidobacterium longum</i> LC74	99	<i>Bifidobacterium longum</i> LC99
75	<i>Bifidobacterium longum</i> LC75	100	<i>Bifidobacterium longum</i> LC100

It was shown that *Lactobacillus plantarum* LC5 shown in Table 10 above was a gram-positive anaerobic bacillus and the 16S rDNA thereof had a nucleotide sequence of SEQ ID NO: 4. The 16S rDNA nucleotide sequence of *Lactobacillus plantarum* LC5 was identified by BLAST in Genbank (<http://www.ncbi.nlm.nih.gov/>), and as a result, a

Lactobacillus plantarum strain having the same 16S rDNA nucleotide sequence as that of *Lactobacillus plantarum* LC5 was not found, and *Lactobacillus plantarum* LC5 showed a homology of 99% with the 16S rDNA sequence of *Lactobacillus plantarum* strain KF9. Furthermore, it was shown that *Lactobacillus plantarum* LC27 shown in Table 10 above was a gram-positive anaerobic bacillus and the 16S rDNA thereof had a nucleotide sequence of SEQ ID NO: 5. The 16S rDNA nucleotide sequence of *Lactobacillus plantarum* LC27 was identified by BLAST in Genebank (<http://www.ncbi.nlm.nih.gov/>), and as a result, a *Lactobacillus plantarum* strain having the same 16S rDNA nucleotide sequence as that of *Lactobacillus plantarum* LC27 was not found, and *Lactobacillus plantarum* LC27 showed a homology of 99% with the 16S rDNA sequence of *Lactobacillus plantarum* strain JL18. In addition, it was shown that *Lactobacillus plantarum* LC28 shown in Table 10 above was a gram-positive anaerobic bacillus and the 16S rDNA thereof had a nucleotide sequence of SEQ ID NO: 6. The 16S rDNA nucleotide sequence of *Lactobacillus plantarum* LC28 was identified by BLAST in Genebank (<http://www.ncbi.nlm.nih.gov/>), and as a result, a *Lactobacillus plantarum* strain having the same 16S rDNA nucleotide sequence as that of *Lactobacillus plantarum* LC28 was not found, and *Lactobacillus plantarum* LC28 showed a homology of 99% with the 16S rDNA sequence of *Lactobacillus plantarum* strain USIM01.

It was shown that *Bifidobacterium longum* LC67 shown in Table 11 above was a gram-positive anaerobic bacillus and the 16S rDNA thereof had a nucleotide sequence of SEQ ID NO: 7. The 16S rDNA nucleotide sequence of *Bifidobacterium longum* LC67 was identified by BLAST in Genbank (<http://www.ncbi.nlm.nih.gov/>), and as a result, a *Bifidobacterium longum* strain having the same 16S rDNA nucleotide sequence as that of *Bifidobacterium longum* LC67 was not found, and *Bifidobacterium longum* LC67 showed a homology of 99% with the 16S rDNA sequence of *Bifidobacterium longum* strain CBT-6. Furthermore, it was shown that *Bifidobacterium longum* LC68 shown in Table 11 above was a gram-positive anaerobic bacillus and the 16S rDNA thereof had a nucleotide sequence of SEQ ID NO: 8. The 16S rDNA nucleotide sequence of *Bifidobacterium longum* LC68 was identified by BLAST in Genbank (<http://www.ncbi.nlm.nih.gov/>), and as a result, a *Bifidobacterium longum* strain having the same 16S rDNA nucleotide sequence as that of *Bifidobacterium longum* LC68 was not found, and *Bifidobacterium longum* LC68 showed a homology of 99% with the 16S rDNA sequence of *Bifidobacterium longum* strain IMAUFB067.

In addition, among the physiological characteristics of *Lactobacillus plantarum* LC5, *Lactobacillus plantarum* LC27, *Bifidobacterium longum* LC67 and *Bifidobacterium longum* LC68, the carbon source utilization was analyzed using a sugar

fermentation by an API kit (model: API 50 CHL; manufactured by BioMerieux's, USA). Table 12 below shows the results of analyzing the carbon source utilization of *Lactobacillus plantarum* LC5 and *Lactobacillus plantarum* LC27, and Table 13 below shows the results of analyzing the carbon source utilization of *Bifidobacterium longum* LC67 and *Bifidobacterium longum* LC68. In Tables 12 and 13 below, "+" indicates the case in which carbon source utilization is positive; "-" indicates the case in which carbon source utilization is negative; and "+" indicates the case in which carbon source utilization is ambiguous. As shown in 12 and 13 below, *Lactobacillus plantarum* LC5, *Lactobacillus plantarum* LC27, *Bifidobacterium longum* LC67 and *Bifidobacterium longum* LC68 showed carbon source utilization different from that of known strains of the same species with respect to some carbon sources.

Table 12

Carbon source	Strain name		Carbon source	Strain name	
	<i>L. plantarum</i> LC5	<i>L. plantarum</i> LC27		<i>L. plantarum</i> LC5	<i>L. plantarum</i> LC27
glycerol	-	-	salicin	+	+
erythritol	-	-	cellobiose	+	+
D-arabinose	-	-	maltose	+	+
L-arabinose	-	+	lactose	-	+
D-ribose	+	+	melibiose	+	+
D-xylose	-	-	sucrose	+	+
L-xylose	-	-	trehalose	+	+
D-adonitol	-	-	inulin	-	-
methyl- β -D-xylopyranoside	-	-	melezitose	+	+

D-galactose	+	±	raffinose	±	-
D-glucose	+	+	starch	-	-
D-fructose	+	+	glycogen	-	-
D-mannose	+	+	xylitol	-	-
L-sorbose	-	-	gentiobiose	+	+
L-rhamnose	-	±	D-turanose	-	+
dulcitol	-	-	D-lyxose	-	-
inositol	-	-	D-tagatose	-	-
mannitol	+	+	D-fucose	-	-
sorbitol	+	+	L-fucose	-	-
α-methyl-D-mannoside	-	±	D-arabitol	-	-
α-methyl-D-glucoside	-	-	L-arabitol	-	-
N-acetyl-glucosamine	+	+	gluconate	-	-
amygdalin	+	+	2-keto-gluconate	-	-
arbutin	+	+	5-keto-gluconate	-	-
esculin	+	+			

Table 13

Carbon source	Strain name		Carbon source	Strain name	
	<i>B. longum</i> LC67	<i>B. longum</i> LC68		<i>B. longum</i> LC67	<i>B. longum</i> LC68
glycerol	-	-	salicin	-	-
erythritol	-	-	cellobiose	-	-
D-arabinose	-	-	maltose	+	+
L-arabinose	+	+	lactose	+	+
D-ribose	-	-	melibiose	+	+
D-xylose	+	±	sucrose	+	±
L-xylose	-	-	trehalose	-	±
D-adonitol	-	-	inulin	-	-
methyl-β-D-xylopyranoside	-	-	melezitose	-	+
D-galactose	±	+	raffinose	+	+
D-glucose	+	+	starch	-	-
D-fructose	+	+	glycogen	-	-
D-mannose	-	-	xylitol	-	-
L-sorbose	-	-	gentiobiose	+	±

L-rhamnose	-	-	D-turanose	±	±
dulcitol	-	-	D-lyxose	-	-
inositol	-	-	D-tagatose	-	-
mannitol	±	+	D-fucose	-	-
sorbitol	±	+	L-fucose	-	-
α-methyl-D-mannoside	-	-	D-arabitol	-	-
α-methyl-D-glucoside	±	±	L-arabitol	-	-
N-acetyl-glucosamine	-	-	gluconate	-	-
amygdalin	-	±	2-keto-gluconate	-	-
arbutin	-	-	5-keto-gluconate	-	-
esculin	+	+			

(4) Information on Deposition of Lactic Acid Bacteria

The present inventors deposited *Lactobacillus plantarum* LC5 with the Korean Culture Center of Microorganisms (address: Yurim Building, 45, Hongjenae 2ga-gil, Seodaemun-gu, Seoul, Korea), an international depositary authority, on January 11, 2016 under accession number KCCM 11800P. Furthermore, the present inventors deposited *Lactobacillus plantarum* LC27 with the Korean Culture Center of Microorganisms (address: Yurim Building, 45, Hongjenae 2ga-gil, Seodaemun-gu, Seoul, Korea), an international depositary authority, on January 11, 2016 under accession number KCCM 11801P. Furthermore, the present inventors deposited *Bifidobacterium longum* LC67 with the Korean Culture Center of Microorganisms (address: Yurim Building, 45, Hongjenae 2ga-gil, Seodaemun-gu, Seoul, Korea), an international depositary authority, on January 11, 2016 under accession number KCCM 11802P.

2. Evaluation of the Effect of Lactic Acid Bacteria on Alleviation of Intestinal Damage or Intestinal Permeability

In order to evaluate the effect of the lactic acid bacteria isolated from kimchi or human feces, on the alleviation of intestinal damage or internal permeability, the antioxidant activity, lipopolysaccharide (LPS) production inhibitory activity, β -glucuronidase (harmful intestinal enzyme) inhibitory activity and tight junction protein expression-inducing activity of the lactic acid bacteria were measured.

(1) Experimental Methods

*** Antioxidant activity**

DPPH (2,2-diphenyl-1-picrylhydrazyl) was dissolved in ethanol to a concentration of 0.2 mM to prepare a DPPH solution. A lactic acid bacteria suspension (1×10^8 CFU/ml) or a vitamin C solution (1 g/ml) was added to 0.1 ml of the DPPH solution and cultured at 37°C for 20 minutes. The culture was centrifuged at 3000 rpm for 5 minutes, and the supernatant was collected. Next the absorbance of the supernatant at 517 nm was measured, and the antioxidant activity of the lactic acid bacteria was calculated.

*** Lipopolysaccharide (LPS) production inhibitory activity**

0.1 g of human fresh feces was suspended in 0.9 ml of sterile physiological saline and diluted 100-fold with general anaerobic medium to prepare a fecal suspension. 0.1 ml of the

fecal suspension and 0.1 ml of lactic acid bacteria (1×10^4 or 1×10^5 CFU) were added to 9.8 ml of sterile anaerobic medium (Nissui Pharmaceuticals, Japan) and cultured anaerobically for 24 hours. Next, the culture was sonicated for about 1 hour to
5 disrupt the outer cell membrane of the bacteria, and centrifuged at $5000 \times g$ and the supernatant was collected. Next, the content of LPS (lipopolysaccharide) (which is a typical endotoxin) in the supernatant was measured by a LAL (Limulus Amoebocyte Lysate) assay kit (manufactured by Cape Cod Inc.,
10 USA). In addition, in order to evaluate the *E. coli* proliferation inhibitory activity of the lactic acid bacteria, the culture obtained through the same experiment as described above was diluted 1000-fold and 100000-fold and cultured in DHL medium, and then the number of *E. coli* cells was counted.

15 * β -glucuronidase inhibitory activity

0.1 ml of 0.1 mM p-nitrophenyl- β -D-glucuronide solution, 0.2 ml of 50 mM phosphate buffered saline and 0.1 ml of a lactic acid bacteria suspension (prepared by suspending of a lactic acid bacteria culture in 5 ml of physiological saline)
20 were placed in a reactor and subjected to β -glucuronidase enzymatic reaction, and 0.5 ml of 0.1 mM NaOH solution was added to stop the reaction. Next, the reaction solution was centrifuged at 3000 rpm for 5 minutes, and the supernatant was collected. Then, the absorbance of the supernatant at 405 nm
25 was measured.

* Tight junction protein expression-inducing activity

Caco2 cells obtained from the Korean Cell Line Bank were cultured in RPMI 1640 medium for 48 hours, and then the cultured Caco2 cells were dispensed to each well of a 12-well plate at a density of 2×10^6 cells/well. Next, each well was treated with 1 μ g of LPS (lipopolysaccharide) or a combination of 1 μ g of LPS (lipopolysaccharide) and 1×10^3 CFU of lactic acid bacteria and incubated for 24 hours. Next, the cultured cells were collected from each well, and the expression level of tight junction protein ZO-1 in the cells was measured by an immunoblotting method.

(2) Experimental Results

The antioxidant activity, lipopolysaccharide (LPS) production inhibitory activity, β -glucuronidase inhibitory activity and tight junction protein expression-inducing activity of the lactic acid bacteria isolated from kimchi or human feces were measured, and the results of the measurement are shown in Tables 14 to 16 below. As shown in Tables 14 to 16 below, *Lactobacillus plantarum* LC5, *Lactobacillus plantarum* LC15, *Lactobacillus plantarum* LC17, *Lactobacillus plantarum* LC25, *Lactobacillus plantarum* LC27, *Lactobacillus plantarum* LC28, *Bifidobacterium longum* LC55, *Bifidobacterium longum* LC65, *Bifidobacterium longum* LC67 and *Bifidobacterium longum* LC68 had excellent antioxidant activity, strongly inhibited lipopolysaccharide (LPS) production and β -glucuronidase

activity, and strongly induced the expression of tight junction protein. In particular, *Bifidobacterium longum* LC67 showed the best tight junction protein expression-inducing activity. These lactic acid bacteria have an excellent antioxidant effect, have an excellent effect of inhibiting the enzymatic activity of intestinal flora's harmful bacteria associated with inflammation and carcinogenesis, inhibit the production of endotoxin LPS (lipopolysaccharide) produced by intestinal flora's harmful bacteria, and induce the expression of tight junction protein. Thus, these lactic acid bacteria can improve intestinal permeability syndrome.

Table 14

Control No.	Strain name	Antioxidant activity	Beta-glucuronidase inhibitory activity	LPS production inhibitory activity	Tight junction protein expression inducing activity
1	<i>Lactobacillus plantarum</i> LC1	++	+	+	-
2	<i>Lactobacillus plantarum</i> LC2	++	++	+	-
3	<i>Lactobacillus plantarum</i> LC3	+++	++	+	-
4	<i>Lactobacillus plantarum</i> LC4	+++	++	+	+
5	<i>Lactobacillus plantarum</i> LC5	+++	+++	++	++
6	<i>Lactobacillus plantarum</i> LC6	++	+++	+	-
7	<i>Lactobacillus plantarum</i> LC7	+++	++	+	-
8	<i>Lactobacillus plantarum</i> LC8	++	+++	+	-
9	<i>Lactobacillus plantarum</i> LC9	++	++	+	+
10	<i>Lactobacillus plantarum</i> LC10	++	+++	+	+
11	<i>Lactobacillus plantarum</i> LC11	++	++	+	-
12	<i>Lactobacillus plantarum</i> LC12	++	+++	+	+
13	<i>Lactobacillus plantarum</i> LC13	++	++	+	+
14	<i>Lactobacillus plantarum</i> LC14	++	++	+	-

15	<i>Lactobacillus plantarum</i> LC15	+++	+++	++	++
16	<i>Lactobacillus plantarum</i> LC16	+	+++	+	-
17	<i>Lactobacillus plantarum</i> LC17	+++	+++	++	++
18	<i>Lactobacillus plantarum</i> LC18	++	++	+	+
19	<i>Lactobacillus plantarum</i> LC19	++	+++	+	+
20	<i>Lactobacillus plantarum</i> LC20	++	++	+	-
21	<i>Lactobacillus plantarum</i> LC21	++	++	+	-
22	<i>Lactobacillus plantarum</i> LC22	++	+++	-	-
23	<i>Lactobacillus plantarum</i> LC23	+++	++	-	-
24	<i>Lactobacillus plantarum</i> LC24	+++	+	-	-
25	<i>Lactobacillus plantarum</i> LC25	+++	+++	++	++
26	<i>Lactobacillus plantarum</i> LC26	++	+	+	+
27	<i>Lactobacillus plantarum</i> LC27	+++	+++	++	++
28	<i>Lactobacillus plantarum</i> LC28	+++	+++	++	++
29	<i>Lactobacillus plantarum</i> LC29	++	+	-	-
30	<i>Lactobacillus plantarum</i> LC30	++	+	+	-
31	<i>Lactobacillus plantarum</i> LC31	+++	++	+	-
32	<i>Lactobacillus plantarum</i> LC32	+++	++	+	-
33	<i>Lactobacillus plantarum</i> LC33	+++	++	+	-
34	<i>Lactobacillus plantarum</i> LC34	++	++	+	+
35	<i>Lactobacillus plantarum</i> LC35	++	++	+	+

Table 15

Control No.	Strain name	Antioxidant activity	Beta-glucuronidase inhibitory activity	LPS production inhibitory activity	Tight junction protein expression inducing activity
36	<i>Lactobacillus plantarum</i> LC36	++	++	++	-
37	<i>Lactobacillus plantarum</i> LC37	+++	++	+	+
38	<i>Lactobacillus plantarum</i> LC38	++	++	+	-
39	<i>Lactobacillus plantarum</i> LC39	++	+	+	-
40	<i>Lactobacillus plantarum</i> LC40	+++	+	-	-
41	<i>Lactobacillus plantarum</i> LC41	++	++	-	+
42	<i>Lactobacillus plantarum</i> LC42	+++	+	-	+
43	<i>Lactobacillus plantarum</i> LC43	++	+	-	+
44	<i>Lactobacillus plantarum</i> LC44	++	+	-	+

45	<i>Lactobacillus plantarum</i> LC45	++	++	-	+
46	<i>Lactobacillus plantarum</i> LC46	++	+	-	+
47	<i>Lactobacillus plantarum</i> LC47	+++	+	-	+
48	<i>Lactobacillus plantarum</i> LC48	++	++	-	+
49	<i>Lactobacillus plantarum</i> LC49	++	+++	+	+
50	<i>Lactobacillus plantarum</i> LC50	+++	++	+	-
51	<i>Bifidobacterium longum</i> LC51	++	++	+	-
52	<i>Bifidobacterium longum</i> LC52	+++	+++	+	-
53	<i>Bifidobacterium longum</i> LC53	++	+++	-	+
54	<i>Bifidobacterium longum</i> LC54	+++	++	+	+
55	<i>Bifidobacterium longum</i> LC55	+++	+++	++	++
56	<i>Bifidobacterium longum</i> LC56	+++	++	+	+
57	<i>Bifidobacterium longum</i> LC57	++	+	-	+
58	<i>Bifidobacterium longum</i> LC58	+++	+	-	+
59	<i>Bifidobacterium longum</i> LC59	++	+	-	-
60	<i>Bifidobacterium longum</i> LC60	+++	+	-	-
61	<i>Bifidobacterium longum</i> LC61	++	+	+	-
62	<i>Bifidobacterium longum</i> LC62	+++	+	+	-
63	<i>Bifidobacterium longum</i> LC63	++	++	++	-
64	<i>Bifidobacterium longum</i> LC64	+++	+	-	-
65	<i>Bifidobacterium longum</i> LC65	+++	+++	++	++
66	<i>Bifidobacterium longum</i> LC66	++	+	+	+
67	<i>Bifidobacterium longum</i> LC67	+++	+++	++	+++
68	<i>Bifidobacterium longum</i> LC68	+++	+++	++	++
69	<i>Bifidobacterium longum</i> LC69	+++	+	-	-
70	<i>Bifidobacterium longum</i> LC70	++	+	-	+

Table 16

Control No.	Strain name	Antioxidant activity	Beta-glucuronidase inhibitory activity	LPS production inhibitory activity	Tight junction protein expression inducing activity
71	<i>Bifidobacterium longum</i> LC71	++	+	-	+
72	<i>Bifidobacterium longum</i> LC72	+++	++	-	+
73	<i>Bifidobacterium longum</i> LC73	++	++	+	-
74	<i>Bifidobacterium longum</i> LC74	++	+++	+	-

75	<i>Bifidobacterium longum</i> LC75	+++	+	-	+
76	<i>Bifidobacterium longum</i> LC76	++	+	-	+
77	<i>Bifidobacterium longum</i> LC77	++	++	+	+
78	<i>Bifidobacterium longum</i> LC78	++	+	+	+
79	<i>Bifidobacterium longum</i> LC79	+++	+	+	+
80	<i>Bifidobacterium longum</i> LC80	++	+	+	+
81	<i>Bifidobacterium longum</i> LC81	++	+	+	+
82	<i>Bifidobacterium longum</i> LC82	++	++	-	+
83	<i>Bifidobacterium longum</i> LC83	+++	+	-	+
84	<i>Bifidobacterium longum</i> LC84	++	++	-	-
85	<i>Bifidobacterium longum</i> LC85	+++	++	-	+
86	<i>Bifidobacterium longum</i> LC86	++	+	+	-
87	<i>Bifidobacterium longum</i> LC87	++	++	+	-
88	<i>Bifidobacterium longum</i> LC88	++	+++	+	+
89	<i>Bifidobacterium longum</i> LC89	++	++	+	+
90	<i>Bifidobacterium longum</i> LC90	++	++	+	+
91	<i>Bifidobacterium longum</i> LC91	+++	+++	+	+
92	<i>Bifidobacterium longum</i> LC92	+++	++	+	+
93	<i>Bifidobacterium longum</i> LC93	++	++	+	+
94	<i>Bifidobacterium longum</i> LC94	++	++	-	+
95	<i>Bifidobacterium longum</i> LC95	++	+++	-	-
96	<i>Bifidobacterium longum</i> LC96	++	+	-	-
97	<i>Bifidobacterium longum</i> LC97	++	+	-	-
98	<i>Bifidobacterium longum</i> LC98	++	++	-	-
99	<i>Bifidobacterium longum</i> LC99	++	++	-	-
100	<i>Bifidobacterium longum</i> LC100	++	++	-	+

* The final concentration of lactic acid bacteria in measurement of antioxidant activity: 1×10^4 CFU/ml; the concentration of lactic acid bacteria added for measurement of beta-glucuronidase inhibitory activity and lipopolysaccharide (LPS) production inhibitory activity: 1×10^4 CFU/ml; the concentration of lactic acid bacteria in measurement of tight junction protein expression-inducing activity: 1×10^4 CFU/ml.

* Criteria for measurement of various activities of lactic acid bacteria: very strongly (+++; >90%); strongly (++; >60-90%); weakly (+; >20-60%); not or less than 20% (-; <20%).

3. Evaluation of the Effect of Lactic Acid Bacteria on

5 Alleviation of Liver injury

Based on evaluation of the effect of the lactic acid bacteria on the alleviation of intestinal damage or intestinal permeability syndrome, the following ten strains were selected:

Lactobacillus plantarum LC5, *Lactobacillus plantarum* LC15,
10 *Lactobacillus plantarum* LC17, *Lactobacillus plantarum* LC25
Lactobacillus plantarum LC27, *Lactobacillus plantarum* LC28,
Bifidobacterium longum LC55, *Bifidobacterium longum* LC65,
Bifidobacterium longum LC67 and *Bifidobacterium longum* LC68.
The effect of each of these selected lactic acid bacteria
15 strains or a mixture of these strains on the alleviation of
liver injury was evaluated using model animals having liver
injury induced by tert-butylperoxide.

1) Experimental Method

Mice (C57BL/6, male) were divided into several groups,
20 each consisting of 6 animals. Tert-butylperoxide was
administered intraperitoneally to the test animals of groups
other than a normal group at a dose of 2.5 mmol/kg to induce
liver injury. From 2 hours after administration of tert-
butylperoxide, 2×10^9 CFU of lactic acid bacteria were
25 administered orally to the test animals of groups other than

the normal group and the negative control group, once a day for 3 days. In addition, silymarin in place of lactic acid bacteria was administered orally to the test animals of the positive control group at a dose of 100 mg/kg, once a day for 3 days. At 6 hours after the last administration of the drug, blood was taken from the heart. The taken blood was allowed to stand at room temperature for 60 minutes, and centrifuged at 3,000 rpm for 15 minutes to separate serum. The GPT (glutamic pyruvate transaminase) and GOT (glutamic oxalacetic transaminase) levels in the separated serum were measured using a blood assay kit (ALT & AST measurement kit; Asan Pharm. Co., Korea). In addition, 1 g of the liver tissue dissected from each test animal was added to saline and homogenized using a homogenizer, and the supernatant was analyzed by an ELISA kit to measure the level of TNF- α .

(2) Experimental Results

Table 17 below shows the changes in GOT, GPT and TNF- α values when lactic acid bacteria were administered to model animals having liver injury induced by tert-butylperoxide. As shown in Table 17 below, *Lactobacillus plantarum* LC5, *Lactobacillus plantarum* LC27, *Lactobacillus plantarum* LC28, *Bifidobacterium longum* LC67 and *Bifidobacterium longum* LC68 showed excellent effects on the alleviation of liver injury compared to silymarin, and mixtures of these lactic acid

bacteria showed better effects on the alleviation of liver injury.

Table 17

Test groups	GOT (IU/L)	GPT (IU/L)	TNF- α (pg/g)
Normal group	42.4	6.2	140.4
Negative control group	103.1	28.0	298.0
Group administered with LC5	36.9	5.4	115.7
Group administered with LC15	60.3	6.2	154.3
Group administered with LC17	65.8	6.8	136.7
Group administered with LC25	64.6	11.3	132.4
Group administered with LC27	35.3	3.3	157.1
Group administered with LC28	42.0	1.0	185.7
Group administered with LC55	55.6	17.6	251.4
Group administered with LC65	61.4	17.3	127.6
Group administered with LC67	50.8	3.8	150.5
Group administered with LC68	40.8	5.7	82.4
Group administered with LC5+LC67	32.7	3.1	115.9
Group administered with LC5+LC68	36.8	5.6	105.4
Group administered with LC27+LC67	30.5	2.3	121.2
Group administered with LC27+LC68	35.4	3.2	112.8
Group administered with LC28+LC67	32.5	2.8	128.2
Group administered with silymarin	52.9	5.9	93.8

5 In Table 17 above, "LC5" indicates *Lactobacillus plantarum* LC5; "LC15" indicates *Lactobacillus plantarum* LC15; "LC17" indicates *Lactobacillus plantarum* LC17; "LC25" indicates *Lactobacillus plantarum* LC25; "LC27" indicates *Lactobacillus plantarum* LC27; "LC28" indicates *Lactobacillus plantarum* LC28;

10 "LC55" indicates *Bifidobacterium longum* LC55; "LC65" indicates *Bifidobacterium longum* LC65; "LC67" indicates *Bifidobacterium longum* LC67; "LC68" indicates *Bifidobacterium longum* LC68;

"LC5+LC67" indicates a lactic acid bacteria mixture prepared by mixing *Lactobacillus plantarum* LC5 and *Bifidobacterium longum* LC67 in the same amount; "LC5+LC68" indicates a lactic acid bacteria mixture prepared by mixing *Lactobacillus plantarum* LC5 and *Bifidobacterium longum* LC68 in the same amount; "LC27+LC67" indicates a lactic acid bacteria mixture prepared by mixing *Lactobacillus plantarum* LC27 and *Bifidobacterium longum* LC67 in the same amount; "LC27+LC68" indicates a lactic acid bacteria mixture prepared by mixing *Lactobacillus plantarum* LC27 and *Bifidobacterium longum* LC68 in the same amount; and "LC28+LC67" indicates a lactic acid bacteria mixture prepared by mixing *Lactobacillus plantarum* LC28 and *Bifidobacterium longum* LC67 in the same amount. In the following Tables showing the experimental results, the same symbols are used for single lactic acid bacteria or lactic acid bacteria mixtures.

4. Evaluation of the Effect of Lactic Acid Bacteria on Alleviation of Allergy

(1) Measurement of the Inhibition of Degranulation by Lactic Acid Bacteria

The RBL-2H3 cell line (rat mast cell line, the Korean Cell Line Bank, Cat. No.22256) was cultured with DMEM (Dulbeccos' modified Eagle's medium, Sigma, 22256) containing 10% FBS (fetal bovine serum) and L-glutamine in a humidified 5% CO₂ incubator at 37°C. The cells contained in the culture medium were floated using trypsin-EDTA solution, and the floated cells

were isolated, collected and used in the experiment. The collected RBL-2H3 cells were dispensed into a 24-well plate at a density of 5×10^5 cells/well and sensitized by incubation with 0.5 $\mu\text{g/ml}$ of mouse monoclonal IgE for 12 hours. The sensitized
5 cells were washed with 0.5 ml of siraganian buffer (119mM NaCl, 5mM KCl, 0.4mM MgCl_2 , 25mM PIPES, 40mM NaOH, pH 7.2), and then incubated with 0.16 ml of siraganian buffer (supplemented with 5.6mM glucose, 1mM CaCl_2 , 0.1% BSA) at 37°C for 10 minutes. Next, lactic acid bacteria as a test drug were added to the
10 cell culture to a concentration of 1×10^4 CFU/ml, or 0.04 ml of DSCG (disodium cromoglycate) as a control drug was added to the cell culture, and after 20 minutes, the cells were activated with 0.02 ml of antigen (1 $\mu\text{g/ml}$ DNP-BSA) at 37°C for 10 minutes. Next, the cell culture was centrifuged at 2000 rpm
15 for 10 minutes, and the supernatant was collected. 0.025 ml of the collected supernatant was transferred to a 96-well plate, and then 0.025 ml of 1mM p-NAG (a solution of p-nitrophenyl-N-acetyl- β -D-glucosamide in 0.1M citrate buffer, pH 4.5) was added thereto, and then the mixture was allowed to react at
20 37°C for 60 minutes. Next, the reaction was stopped by addition of 0.2 ml of 0.1M $\text{Na}_2\text{CO}_3/\text{NaHCO}_3$, and the absorbance at 405 nm was measured by an ELISA analyzer.

(2) Experimental Results

Table 18 below shows the results of measuring of the
25 inhibition (%) of degranulation by lactic acid bacteria. As

shown in Table 18, *Lactobacillus plantarum* LC5, *Lactobacillus plantarum* LC27, *Lactobacillus plantarum* LC28, *Bifidobacterium longum* LC67, *Bifidobacterium longum* LC68 and mixtures thereof effectively inhibited the degranulation of basophils. Thus, these lactic acid bacteria or mixtures thereof can very effectively alleviate allergic atopy, asthma, pharyngitis, chronic dermatitis or the like.

Table 18

Drug	Degranulation inhibition (%)
None	0
LC5	65
LC15	45
LC17	43
LC25	48
LC27	52
LC28	54
LC55	38
LC65	42
LC67	65
LC68	61
LC5+LC67	65
LC5+LC68	60
LC27+LC67	65
LC27+LC68	59
LC28+LC67	62
DSCG(disodium cromoglycate)	62

10 5. In Vitro Evaluation of the Anti-inflammatory and Immune Regulatory Effects of Lactic Acid Bacteria

(1) Isolation of Macrophages and Measurement of Inflammatory Marker

6-Week-old C57BL/6J male mice (20-23g) were purchased from RaonBio Co., Ltd. 2 ml of 4% sterile thioglycolate was administered into the abdominal cavity of each mouse, after 96 hours, the mice were anesthetized and 8 ml of RPMI 1640 medium was administered into the abdominal cavity of each mouse. After 5-10 minutes, the RPMI medium (including macrophages) in the abdominal cavity of the mice was taken out, centrifuged at 1000 rpm for 10 minutes, and then washed twice with RPMI 1640 medium. The macrophages were seeded on a 24-well plate at a density of 0.5×10^6 cells/well and treated with the test substance lactic acid bacteria and the inflammation inducer LPS (lipopolysaccharide) for 2 hours or 24 hours, and then the supernatant and the cells were collected. In this case, the lactic acid bacteria were used at a concentration of 1×10^4 CFU/ml for treatment of the cells. The collected cells were homogenized in buffer (Gibco). Using the collected supernatant, the expression levels of cytokines such as TNF- α were measured by an ELISA kit. In addition, using the collected cells, the expression levels of p65 (NF-kappa B), p-p65 (phosphor-NF-kappa B) and β -actin were measured by an immunoblotting method. Specifically, 50 μ g of the supernatant was taken and electrophoresed on SDS 10% (w/v) polyacrylamide gel for 1 hour and 30 minutes. The electrophoresed sample was transferred to a nitrocellulose membrane under the conditions of 100 V and 400 mA for 1 hour and 10 minutes. The sample-transferred

nitrocellulose membrane was blocked with 5% skim milk for 30 minutes, and then washed three times with PBS-Tween for 5 minutes each time, and incubated with a 1:100 dilution of primary antibody (Santa Cruz Biotechnology, USA) overnight.

5 Next, the membrane was washed three times for 10 minutes each time, and incubated with a 1:1000 dilution of secondary antibody (Santa Cruz Biotechnology, USA) for 1 hour and 20 minutes. Next, the membrane was washed three times for 15 minutes each time, and it was developed by fluorescence and
10 visualized. The intensity of the developed band was measured, and then inhibition (%) was calculated using the following equation. In the following equation, the normal group indicates a group in which macrophages were treated with saline alone; the group treated with LPS indicates a group in which
15 macrophages were treated with LPS alone; and the group treated with lactic acid bacteria indicates a group in which macrophages were treated with both lactic acid bacteria and LPS.

Inhibition (%) = (expression level in group treated with LPS - expression level in group treated with lactic acid
20 bacteria) / (expression level in group treated with LPS - expression level in normal group) x 100

Table 19 below shows the inhibition of NF-kappa B activation and the inhibition of TNF- α expression when macrophages having inflammation induced by LPS
25 (lipopolysaccharide) were treated with the lactic acid bacteria.

As shown in Table 19 below, *Lactobacillus plantarum* LC5, *Lactobacillus plantarum* LC27, *Lactobacillus plantarum* LC28, *Bifidobacterium longum* LC67, *Bifidobacterium longum* LC68 and mixtures thereof effectively inhibited inflammation induced by LPS (lipopolysaccharide).

Table 19

Lactic acid bacteria used for treatment	Inhibition (%) of TNF- α expression	Inhibition (%) of p-p65/p65 activation
LC5	71	73
LC15	54	55
LC17	61	55
LC25	52	65
LC27	70	72
LC28	74	71
LC55	63	62
LC65	65	68
LC67	76	77
LC68	75	71
LC5+LC67	78	72
LC5+LC68	76	72
LC27+LC67	81	75
LC27+LC68	77	73
LC28+LC67	77	73

(2) Isolation of T cells from Spleen and Measurement of Differentiation into Th17 Cells or Treg Cells

Spleen was separated from C56BL/6J mice, crushed suitably and suspended in 10% FCS-containing RPMI 1640 medium, and CD4 T cells were isolated therefrom using a CD4 T cell isolation kit (MiltenyiBiotec, Bergisch Gladbach, Germany). The isolated CD4 T cells were seeded in a 12-well plate at a density of 5×10^5

cells/well, and anti-CD3 (1 µg/ml, MiltenyiBiotec, Bergisch Gladbach, Germany) and anti-CD28(1 µg/ml, MiltenyiBiotec, Bergisch Gladbach, Germany) were added thereto, or anti-CD3 (1 µg/ml, MiltenyiBiotec, Bergisch Gladbach, Germany), anti-CD28
5 (1 µg/ml, MiltenyiBiotec, Bergisch Gladbach, Germany), recombinant IL-6 (20 ng/ml, MiltenyiBiotec, Bergisch Gladbach, Germany) and recombinant transforming growth factor beta (1 ng/ml, MiltenyiBiotec, Bergisch Gladbach, Germany) were added. While the cells were cultured, 1×10^3 or 1×10^5 CFU of the
10 lactic acid bacteria were added thereto, and the cells were cultured for 4 days. Next, the cells of the culture were stained with anti-FoxP3 or anti-IL-17A antibody, and the distribution of Th17 cells and Treg cells was analyzed using a FACS (fluorescence-activated cell sorting) system (C6 Flow
15 Cytometer® System, San Jose, CA, USA).

Table 20 below shows the level of differentiation of T cells (isolated from spleen) into Th17 cells when the T cells were treated with anti-CD3, anti-CD28, IL-6 and TGF-β, and Table 21 below shows the level of differentiation of T cells
20 (isolated from spleen) into Treg cells when the T cells were treated with anti-CD3 and anti-CD28. As shown in Tables 20 and 21 below, *Lactobacillus plantarum* LC5, *Lactobacillus plantarum* LC27, *Lactobacillus plantarum* LC28, *Bifidobacterium longum* LC67, *Bifidobacterium longum* LC68 and mixtures thereof inhibited the
25 differentiation of T cells into Th17 cells (T helper 17 cells)

and promoted the differentiation of T cells into Treg cells. These results suggest that the lactic acid bacteria or mixtures thereof can effectively alleviate inflammatory diseases such as colitis or arthritis.

5 Table 20

T-cell treatment method		Differentiation (%) into Th17 cells
Treatment with anti-CD3, anti-CD28, IL-6 and TGF- β	Treatment with lactic acid bacteria	
Not treated	Not treated	12.2
Treated	Not treated	25.6
Treated	Treated with LC5	14.2
Treated	Treated with LC15	19.6
Treated	Treated with LC17	17.9
Treated	Treated with LC25	18.2
Treated	Treated with LC27	15.1
Treated	Treated with LC28	14.9
Treated	Treated with LC55	18.8
Treated	Treated with LC65	17.9
Treated	Treated with LC67	15.9
Treated	Treated with LC68	15.7
Treated	Treated with LC5+LC67	14.2
Treated	Treated with LC5+LC68	14.5
Treated	Treated with LC27+LC67	13.9
Treated	Treated with LC27+LC68	14.4
Treated	Treated with LC28+LC67	14.1

Table 21

T-cell treatment method		Differentiation (%) into Treg cells
Treatment with anti-CD3 and anti-CD28	Treatment with lactic acid bacteria	
Not treated	Not treated	9.1
Treated	Not treated	11.4
Treated	Treated with LC5	22.9
Treated	Treated with LC15	15.8

Treated	Treated with LC17	16.9
Treated	Treated with LC25	18.4
Treated	Treated with LC27	21.8
Treated	Treated with LC28	21.4
Treated	Treated with LC55	19.5
Treated	Treated with LC65	19.2
Treated	Treated with LC67	21.6
Treated	Treated with LC68	20.5
Treated	Treated with LC5+LC67	21.8
Treated	Treated with LC5+LC68	21.8
Treated	Treated with LC27+LC67	22.0
Treated	Treated with LC27+LC68	21.5
Treated	Treated with LC28+LC67	21.9

6. In Vivo Evaluation of the Anti-inflammatory and Colitis-Alleviating Effects of Lactic Acid Bacteria

(1) Test Animals

5 5-Week-old C57BL/6 male mice (24-27g) were purchased from OrientBio, and housed under controlled environmental conditions (humidity: 50±10%, temperature: 25±2°C, 12-hr light/12-hr dark cycle), and then used in the experiment. As feed, standard experimental feed (Samyang, Korea) was used, and the animals
10 had access to drinking water *ad libitum*. In all the experiments, one group consisted of 6 animals.

(2) Colitis Induction by TNBS and Sample Administration

One group of the test animals was used as a normal group, and the test animals of the other groups were treated with
15 2,4,6-trinitrobenzenesulfonic acid (TNBS) to induce acute colitis. Specifically, the test animals were lightly

anesthetized with ether, and then a mixture solution of 2.5 g of TNBS (2,4,6-trinitrobenzene sulfonic acid) in 100 ml of 50% ethanol was administered into the colon through the anal in an amount of 0.1 ml each time by use of a 1-ml round-tip syringe, and lifted vertically and maintained for 30 seconds, thereby inducing inflammation. On the other hand, the normal group was orally administered with 0.1 ml of saline. On the next day, the lactic acid bacteria or the lactic acid bacteria mixture as a test sample was suspended in saline and administered orally to each mouse in an amount of 2.0×10^9 CFU, once a day for three days. On the next day following the end of sample administration, the animals were killed with carbon dioxide, and a colon portion ranging from the cecum to the site just before the anus was dissected and used. Meanwhile, the test animals of the normal group were orally administered with saline alone instead of the lactic acid bacteria. In addition, the test animals of the negative control group were orally administered with saline alone instead of the lactic acid bacteria after the induction of colitis by TNBS. Furthermore, the test animals of the positive control group were orally administered with 50 mg/kg of sulfasalazine, which is a drug for treating colitis, instead of the lactic acid bacteria.

(3) Macroscopic Analysis of Colon

The length and appearance of the dissected colon were observed, and the appearance was analyzed by scoring according

to the criteria (Hollenbach et al., 2005, Criteria for Degree of Colitis) shown in Table 22 below. After complete removal of colon contents, the colon tissue was washed with saline. A portion of the washed colon tissue was fixed with 4% formaldehyde solution in order to use it as a pathological tissue sample, and the remainder was freeze-stored at -80°C for molecular biological analysis.

Table 22

Macroscopic score	Criteria
0	Any ulcer and inflammation are not found.
1	Edema without bleeding is found.
2	Ulcer with edema is found.
3	Ulcer and inflammation are found at only one site.
4	Ulcer and inflammation are found at two or more sites.
5	Ulcer has an increased size of 2 cm or more.

10 (4) Measurement of Myeloperoxidase (MPO) Activity

100 mg of colon tissue was homogenized in 200 µl of 10 mM potassium phosphate buffer (pH 7.0) containing 0.5% hexadecyl trimethyl ammonium bromide. The homogenized tissue was centrifuged at 10,000×g and 4°C for 10 minutes, and the supernatant was collected. 50 µl of the supernatant was added to 0.95 ml of a reaction solution (containing 1.6mM tetramethyl benzidine and 0.1mM H₂O₂) and allowed to react at 37°C, and the absorbance at 650 nm was measured at various time points during the reaction. To calculate myeloperoxidase (MPO) activity, 1 µmol/ml of peroxide produced by the reaction was used as 1 unit.

(5) Measurement of Inflammatory Marker

Using a Western blotting method, inflammatory markers such as p-p65, p65, iNOS, COX-2 and β -actin were measured. Specifically, according to the same method as the experiment
5 for measurement of myeloperoxidase (MPO) activity, a supernatant was obtained. 50 μ g of the supernatant was taken and electrophoresed on SDS 10% (w/v) polyacrylamide gel for 1 hour and 30 minutes. The electrophoresed sample was transferred to a nitrocellulose membrane under the conditions
10 of 100 V and 400 mA for 1 hour and 10 minutes. The sample-transferred nitrocellulose membrane was blocked with 5% skim milk for 30 minutes, and then washed three times with PBS-Tween for 5 minutes each time, and incubated with a 1:100 dilution of primary antibody (Santa Cruz Biotechnology, USA) overnight.
15 Next, the membrane was washed three times for 10 minutes each time, and incubated with a 1:1000 dilution of secondary antibody (Santa Cruz Biotechnology, USA) for 1 hour and 20 minutes. Next, the membrane was washed three times for 15 minutes each time, and it was developed by fluorescence and
20 visualized.

In addition, inflammation-related cytokines such as TNF- α , IL-17, IL-10 and the like were measured using an ELISA kit.

(6) Analysis of Immune Regulatory Markers

Dissected colon was washed twice with 2.5 mM EDTA solution.
25 The washed colon was agitated in RPMI medium containing 1 mg/ml

collagenase type VIII (Sigma) at 30°C for 20 minutes and was filtered to separate the Lamina propria. Next, the Lamina propria was treated with 30-100% percoll solution and centrifuged to separate T cells. The separated T cells were
5 stained with anti-FoxP3 or anti-IL-17A antibody, and the distribution of Th17 and Treg cells was analyzed using a FACS (fluorescence-activated cell sorting) system (C6 Flow Cytometer® System, San Jose, CA, USA).

(7) Experimental Results

10 Table 23 below shows the effects of the lactic acid bacteria on the weight of the colon, the appearance of the colon, myeloperoxidase (MPO) activity and inflammation-related cytokine contents when the lactic acid bacteria were administered to the model animals having TNBS-induced acute
15 colitis. As shown in Table 23 below, the model animals having acute colitis induced by TNBS showed reduced weight, reduced macroscopic score of the colon, reduced colon length and increased MPO activity. However, when the lactic acid bacteria were administered to the model animals having acute colitis
20 induced by TNBS, all these markers were improved. In particular, administration of *Bifidobacterium longum* LC67 alone or administration of a mixture of *Bifidobacterium longum* LC67 and *Lactobacillus plantarum* LC5 showed a very excellent effect on the alleviation of colitis. In addition, the model animals
25 having acute colitis induced by TNBS showed increased TNF- α and

IL-17 levels and decreased IL-10 levels. However, when the lactic acid bacteria were administered to the model animals having acute colitis induced by TNBS, all these markers were improved. In particular, when *Bifidobacterium longum* LC67 was administered alone or a mixture of *Bifidobacterium longum* LC67 and *Lactobacillus plantarum* LC5 was administered, TNF- α and IL-17 levels greatly decreased, and IL-10 levels greatly increased.

Table 23

Test groups	Weight gain (g)	Macroscopic score	Colon length (cm)	MPO activity (μ U/mg)	TNF- α (pg/mg)	IL-17 (pg/mg)	IL-10 (pg/mg)
Normal group	0.64	5.9	0.14	0.42	35.1	18.4	61.2
Negative control group	-2.46	4.2	2.32	1.54	95.5	65.2	30.7
Group administered with LC5	-1.90	4.65	1.30	0.91	75.5	52.8	43.9
Group administered with LC27	-1.0	4.56	1.08	0.82	67.2	50.4	44.0
Group administered with LC67	-0.28	4.92	0.50	0.43	48.5	38.5	54.6
Group administered with LC 68	-1.02	4.5	1.34	1.04	54.4	50.5	48.1
Group administered with LC5+LC67	-0.3	5.08	0.84	0.42	45.1	37.3	55.3
Group administered	-1.15	4.8	1.17	0.78	59.8	45.0	50.0

with LC27+LC68							
Positive control group	-0.91	4.58	1.43	0.95	58.2	48.5	45.5

FIG. 22 shows the differentiation patterns of T cells into Th17 cells, which indicate the effect of lactic acid bacteria on model animals having acute colitis induced by TNBS, and FIG. 23 shows the differentiation patterns of T cells into Treg cells, which indicate the effect of lactic acid bacteria on model animals having acute colitis induced by TNBS. In FIGS. 22 and 23, "NOR" indicates a normal group; "TNBS" indicates a negative control group; "LC5" indicates a group administered with *Lactobacillus plantarum* LC5; "LC27" indicates a group administered with *Lactobacillus plantarum* LC27, "LC67" indicates a group administered with *Bifidobacterium longum* LC67; "LC68" indicates a group administered with *Bifidobacterium longum* LC68; "LC5+LC67" indicates a group administered with a lactic acid bacteria mixture prepared by mixing *Lactobacillus plantarum* LC5 and *Bifidobacterium longum* LC67 in the same amount; "LC27+LC68" indicates a group administered with a lactic acid bacteria mixture prepared by mixing *Lactobacillus plantarum* LC27 and *Bifidobacterium longum* LC68 in the same amount; and "SS" indicates a group administered with sulfasalazine. As shown in FIGS. 22 and 23, in the case of the animals having acute colitis induced by TNBS,

the differentiation of T cells into Th17 cells was promoted, and the differentiation of T cells into Treg cells was inhibited. However, when the lactic acid bacteria were administered to the animals having acute colitis induced by TNBS, the differentiation of T cells into Th17 cells was inhibited, and the differentiation of T cells into Treg cells was promoted. In particular, when *Bifidobacterium longum* LC67 was administered alone or a mixture of *Bifidobacterium longum* LC67 and *Lactobacillus plantarum* LC5 was administered, the differentiation of T cells into Th17 cells was significantly inhibited, and the differentiation of T cells into Treg cells was significantly promoted.

FIG. 24 shows inflammatory response markers indicating the effect of lactic acid bacteria on model animals having acute colitis induced by TNBS. In FIG. 24, "Nor" indicates a normal group; "T" indicates a negative control group; "LC5" indicates a group administered with *Lactobacillus plantarum* LC5; "LC27" indicates a group administered with *Lactobacillus plantarum* LC27; "LC67" indicates a group administered with *Bifidobacterium longum* LC67; "LC68" indicates a group administered with *Bifidobacterium longum* LC68; "LC5+LC67" indicates a group administered with a lactic acid bacteria mixture prepared by mixing *Lactobacillus plantarum* LC5 and *Bifidobacterium longum* LC67 in the same amount; "LC27+LC68" indicates a group administered with a lactic acid bacteria

mixture prepared by mixing *Lactobacillus plantarum* LC27 and *Bifidobacterium longum* LC68 in the same amount, and "SS" indicates a group administered with sulfasalazine. As shown in FIG. 24, in the case of the model animals having acute colitis induced by TNBS, NF- κ B was activated (p-p65) and the expression levels of COX-2 and iNOS increased. However, when the lactic acid bacteria were administered, the activation of NF- κ B (p-p65) was inhibited, and the expression levels of COX-2 and iNOS also decreased. In particular, administration of *Bifidobacterium longum* LC67 alone or administration of a mixture of *Bifidobacterium longum* LC67 and *Lactobacillus plantarum* LC5 exhibited excellent effects on the inhibition of NF- κ B activation (p-p65) and on the inhibition of expression of COX-2 and iNOS.

7. In Vivo Evaluation of the Effect of Lactic Acid Bacteria on Alleviation of Alcohol-Induced Gastric Ulcer

(1) Test Animals

5-Week-old C57BL/6 male mice (24-27g) were purchased from OrientBio, and housed under controlled environmental conditions (humidity: 50 $\pm$ 10%, temperature: 25 $\pm$ 2°C, 12-hr light/12-hr dark cycle), and then used in the experiment. As feed, standard experimental feed (Samyang, Korea) was used, and the animals had access to drinking water *ad libitum*. In all the experiments, one group consisted of 6 animals.

(2) Induction of Gastric Ulcer by Alcohol and Administration of Sample

To one test group, 1×10^9 CFU of *Lactobacillus plantarum* LC27 suspended in saline was orally administered once a day for 3 days. To another test group, 1×10^9 CFU of *Bifidobacterium longum* LC67 suspended in saline was orally administered once a day for 3 days. To still another test group, 1×10^9 CFU of a lactic acid bacteria mixture prepared by mixing *Lactobacillus plantarum* LC27 and *Bifidobacterium longum* LC67 in the same amount was orally administered once a day for 3 days, after it was suspended in saline. In addition, to a positive control group, ranitidine, a commercial agent for treating gastric ulcer, was orally administered once a day for 3 days in an amount of 50 mg/kg. In addition, to a normal group and a negative control group, 0.2 ml of saline was orally administered one a day for 3 days. After the sample was orally administered for 3 days, the test mice were fasted and water-deprived for 18 hours. On day 4 of the experiment, at 1 hour after administration of saline, 0.2 ml of 99% pure ethanol was administered orally to the mice of all the test groups other than the normal group to induce gastric ulcer. In addition, to the normal group, 0.2 ml of saline was administered instead of ethanol.

(3) Measurement of Macroscopic Marker Related to Gastric Injury

3 Hours after administration of ethanol, the test mice were sacrificed and gastric tissue was dissected split longitudinally and washed with PBS (phosphate buffer saline) solution, and then the degree of gastric injury was observed
5 visually or microscopically and scored (see Park, S.W., Oh, T.Y., Kim, Y.S., Sim, H., et al., *Artemisia asiatica* extracts protect against ethanol-induced injury in gastric mucosa of rats. *J. Gastroenterol. Hepatol.* 2008, 23, 976-984).

(4) Measurement of Myeloperoxidase (MPO) Activity

10 100 mg of the gastric tissue was homogenized in 200 μ l of 10 mM potassium phosphate buffer (pH 7.0) containing 0.5% hexadecyl trimethyl ammonium bromide. Then, the tissue solution was centrifuged at 10,000 $\times$ g and 4°C for 10 minutes, and the supernatant was collected. 50 μ l of the supernatant
15 was added to 0.95 ml of a reaction solution (containing 1.6mM tetramethyl benzidine and 0.1mM H₂O₂) and allowed to react at 37°C, and the absorbance at 650 nm was measured at various time points during the reaction. To calculate myeloperoxidase (MPO) activity, 1 μ mol/ml of peroxide produced by the reaction was
20 used as 1 unit.

(5) Measurement of Inflammatory Markers

2 μ g of mRNA was isolated from gastric tissue by a Qiagen RNeasy Mini Kit and synthesized into cDNA using Takara Prime Script Rtase. Next, the expression levels of CXCL4 [chemokine
25 (C-X-C motif) ligand 4] and TNF- α (tumor necrosis factor-alpha)

were measured using a quantitative real time polymerase chain reaction (Qiagen thermal cycler, Takara SYBER premix agent, Thermal cycling conditions: activation of DNA polymerase for 5 min at 95°C, followed by 40 cycles of amplification for 10 s at 95°C and for 45 s at 60°C). Table 24 below shows the primer sequences used to analyze each cytokine in the quantitative real time polymerase chain reaction.

Table 24

Cytokine to be analyzed	Kind of primer	Primer nucleotide sequence
TNF- α	Forward	5'-CTGTAGCCCACGTCGTAGC-3'
	Reverse	5'-TTGAGATCCATGCCGTTG-3'
CXCL4	Forward	5'-AGTCCTGAGCTGCTGCTTCT-3'
	Reverse	5'-GATCTCCATCGCTTTCTTCG-3'

10 (6) Experimental Results

FIG. 25 depicts images showing the effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in the second experiment of the present invention; FIG. 26 shows the gross gastric lesion score indicating the effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in the second experiment of the present invention; FIG. 27 shows the ulcer index indicating the effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in the second experiment of the present invention; and FIG. 28 shows the histological activity index indicating the

effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in the second experiment of the present invention. Furthermore, FIG. 29 shows the myeloperoxidase (MPO) activity indicating the effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in the second experiment of the present invention. In addition, FIG. 30 shows CXCL4 expression levels indicating the effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in the second experiment of the present invention; and FIG. 31 shows TNF- α expression levels indicating the effect of lactic acid bacteria on the stomach mucosa of mice having gastric ulcer induced by ethanol, in the second experiment of the present invention. In FIGS. 30 and 31, the CXCL4 expression levels and TNF- α expression levels in the test groups other than the normal group are expressed fold-changes relative to the expression levels in the normal group. In FIGS. 25 to 31, "Nor" indicates a normal group; "Ethanol" indicates a negative control group having ethanol-induced gastric ulcer and administered with saline as a sample; "Ethanol+Ranitidine" indicates a test group having ethanol-induced gastric ulcer and administered with Ranitidine as a sample; "Ethanol+LC27" indicates a test group having ethanol-induced gastric ulcer and administered with *Lactobacillus plantarum* LC27 as a sample; "Ethanol+LC67" indicates a test group having ethanol-induced

gastric ulcer and administered with *Bifidobacterium longum* LC67 as a sample; and "Ethanol+LC27/LC67" indicates a test group having ethanol-induced gastric ulcer and administered with a lactic acid bacteria mixture, prepared by mixing *Lactobacillus plantarum* LC27 and *Bifidobacterium longum* LC67 in the same amount, as a sample. As shown in FIGS. 25 to 29, *Bifidobacterium longum* LC67, *Lactobacillus plantarum* LC27 or a mixture thereof effectively alleviated the gastric injury or gastric ulcer induced by ethanol. Furthermore, as shown in 10 FIGS. 30 and 31, *Bifidobacterium longum* LC67, *Lactobacillus plantarum* LC27 or a mixture thereof greatly reduced the inflammatory marker levels in the mice having ethanol-induced gastric injury or gastric ulcer.

8. In Vivo Evaluation of the Effect of Lactic Acid

Bacteria on Alleviation of Alcohol-Induced Liver injury

(1) Test Animals

5-Week-old C57BL/6 male mice (24-27g) were purchased from OrientBio, and housed under controlled environmental conditions (humidity: 50±10%, temperature: 25±2°C, 12-hr light/12-hr dark 20 cycle), and then used in the experiment. As feed, standard experimental feed (Samyang, Korea) was used, and the animals had access to drinking water *ad libitum*. In all the experiments, one group consisted of 6 animals.

(2) Induction of Liver Injury by Alcohol and

Administration of Sample

To one test group, 1×10^9 CFU of *Lactobacillus plantarum* LC27 suspended in saline was orally administered once a day for 3 days. To another test group, 1×10^9 CFU of *Bifidobacterium longum* LC67 suspended in saline was orally administered once a day for 3 days. To still another test group, 1×10^9 CFU of a lactic acid bacteria mixture prepared by mixing *Lactobacillus plantarum* LC27 and *Bifidobacterium longum* LC67 in the same amount was orally administered once a day for 3 days, after it was suspended in saline. To a positive control group, silymarin, a commercial agent for treating liver injury, was orally administered once a day for 3 days in an amount of 50 mg/kg. In addition, to a normal group and a negative control group, 0.1 ml of saline was orally administered once a day for 3 days. 3 hours after 3 days of oral administration of the sample or saline, ethanol was administered intraperitoneally to the mice of all the test groups other than the normal group in an amount of 6 ml/kg in order to induce liver injury. In addition, to the normal group, saline in place of ethanol was administered intraperitoneally in an amount of 6 ml/kg. Next, the test mice were fasted and water-deprived for 12 hours, and then sacrificed, and blood was taken from the heart.

(3) Measurement of Liver Function Markers and Results

The taken blood was allowed to stand at room temperature for 60 minutes and centrifuged at 3,000 rpm for 15 minutes to separate serum. The GPT (glutamic pyruvate transaminase) and

GOT (glutamic oxalacetic transaminase) levels in the separated serum were measured using a blood assay kit (ALT & AST measurement kit; Asan Pharm. Co., Korea), and the results of the measurement are shown in Table 25 below. As shown in Table 5 25 below, *Bifidobacterium longum* LC67, *Lactobacillus plantarum* LC27 or a mixture thereof effectively alleviated ethanol-induced liver injury. In particular, *Bifidobacterium longum* LC67 showed a better effect than silymarin which is a commercial agent for treating liver injury.

10 Table 25

Test groups	GOT (IU/L)	GPT (IU/L)
Normal group	52.1	42.2
Negative control group	107.7	156.3
Group administered with ethanol and LC27	82.3	95.4
Group administered with ethanol and LC67	62.5	65.8
Group administered with ethanol and LC27/LC67	71.4	78.3
Group administered with ethanol and silymarin	79.5	87.5

* LC27: *Lactobacillus plantarum* LC27
 * LC67: *Bifidobacterium longum* LC67
 * LC27/LC67: lactic acid bacteria mixture prepared by mixing *Lactobacillus plantarum* LC27 and *Bifidobacterium longum* LC67 in 15 the same amount.

Although the present invention has been described above with reference to the examples, the scope of the present invention is not limited to these examples, and various modifications are possible without departing from the scope and 20 idea of the present invention. Therefore, the scope of

protection of the present invention should be interpreted to include all embodiments falling within the appended claims.

Claims

Claim 1

Isolated *Lactobacillus plantarum* LC27 (accession number:
5 KCCM 11801P), a culture thereof, a lysate thereof or an extract thereof.

Claim 2

The isolated *Lactobacillus plantarum* LC27 according to
claim 1, wherein the isolated *Lactobacillus plantarum* LC27
10 having one or more characteristics selected from antioxidant activity, beta-glucuronidase inhibitory activity, lipopolysaccharide (LPS) production inhibitory activity and tight junction protein expression-inducing activity.

Claim 3

15 The isolated *Lactobacillus plantarum* LC27 according to claim 1, wherein the isolated *Lactobacillus plantarum* LC27 comprising a 16S rDNA nucleotide sequence represented by SEQ ID NO: 5.

Claim 4

20 A pharmaceutical composition for preventing or treating one or more diseases selected from a group consisting of intestinal damage, liver injury, allergic disease and inflammatory disease comprising the *Lactobacillus plantarum* LC27, a culture thereof, a lysate thereof or an extract thereof
25 of claim 1.

Claim 5

A food composition for preventing or alleviating one or more diseases selected from a group consisting of intestinal

damage, liver injury, allergic disease and inflammatory disease comprising the *Lactobacillus plantarum* LC27, a culture thereof, a lysate thereof or an extract thereof of claim 1.

Claim 6

5 The food composition according to claim 5, wherein the food composition further comprising one or more lactic acid bacteria selected from a group consisting of *Lactobacillus brevis* CH23 (accession number: KCCM 11762P), *Bifidobacterium longum* CH57 (accession number: KCCM 11764P), *Lactobacillus*
10 *plantarum* LC5 (accession number: KCCM 11800P) and *Bifidobacterium longum* LC67 (accession number: KCCM 11802P).

Claim 7

 The food composition according to claim 5, wherein the food composition further comprising *Bifidobacterium longum* LC67
15 (accession number: KCCM 11802P), a culture thereof, a lysate thereof or an extract thereof.

Claim 8

 The food composition according to claim 5, wherein the intestinal damage is intestinal permeability syndrome.

Claim 9

 The food composition according to claim 5, wherein the liver injury is selected from a group consisting of hepatitis, fatty liver and liver cirrhosis.

Claim 10

25 The food composition according to claim 5, wherein the allergic disease is selected from a group consisting of atopic dermatitis, asthma, pharyngitis and chronic dermatitis.

Claim 11

The food composition according to claim 5, wherein the inflammatory disease is selected from a group consisting of gastritis, gastric ulcer, colitis and arthritis.

Claim 12


5 A method of preventing or treating one or more diseases selected from a group consisting of intestinal damage, liver injury, allergic disease and inflammatory disease comprising administering a composition comprising *Lactobacillus plantarum* LC27 (accession number: KCCM 11801P), a culture thereof, a
10 lysate thereof or an extract thereof to a subject.

Claim 13


Use of a composition comprising *Lactobacillus plantarum* LC27 (accession number: KCCM 11801P), a culture thereof, a lysate thereof or an extract thereof for manufacture of a
15 medicament for preventing or treating one or more diseases selected from a group consisting of intestinal damage, liver injury, allergic disease and inflammatory disease.

【DRAWINGS】


【Figure 1】




【Figure 2】




【Figure 3】




【Figure 4】




【Figure 5】




【Figure 6】



【Figure 7】



【Figure 8】




【Figure 9】

NOR


TNBS

CH57


SS50




【Figure 10】




【Figure 11】




【Figure 12】




【Figure 13】




【Figure 14】




【Figure 15】




【Figure 16】




【Figure 17】




【Figure 18】




【Figure 19】




【Figure 20】




【Figure 21】




【Figure 22】




【Figure 23】




【Figure 24】




【Figure 25】




【Figure 26】




【Figure 27】




【Figure 28】




【Figure 29】



【Figure 30】



【Figure 31】



P17010- NVP_Sequence list.txt

<110> University-Industry Cooperation Group of Kyung Hee University
 <120> Novel lactic acid bacteria with various functionality and use thereof
 <130> PCT- 16- 130
 <150> KR 10- 2015- 0130124
 <151> 2015- 09- 15
 <150> KR 10- 2016- 0005018
 <151> 2016- 01- 15
 <160> 8
 <170> Kopat ent l n 2. 0
 <210> 1
 <211> 1132
 <212> DNA
 <213> Unknown
 <220>
 <223> 16S rDNA of Lactobacillus brevis CH23

<400> 1
 agggat aaca ct t ggaaaca ggt gct aat a ccgt at aaca acaaaat ccg cat ggat t t t 60
 gt t t gaaagg t ggct t cggc t at cact t ct ggat gat ccc gcggcgt at t agt t agt t gg 120
 t gaggt aaag gccaccaag acgat gat ac gt agccgacc t gagagggt a at cggccaca 180
 t t gggact ga gacacggccc aaact cct ac gggaggcagc agt agggaat ct t ccacaat 240
 ggacgaaagt ct gat ggagc aat gccgct gagt gaagaa ggggt t t cggc t cgt aaaact 300
 ct gt t gt t aa agaagaacac ct t t gagagt aact gt t caa ggggt t gacgg t at t t aacca 360
 gaaagccacg gct aact acg t gccagcagc cgcggt aat a cgt aggt ggc aagcgt t gt c 420
 cggat t t at t gggcgt aaag cgagcgcagg cgggt t t t t a agt ct gat gt gaaagcct t c 480
 ggct t aaccg gagaagt gca t cggaaact g ggagact t ga gt gcagaaga ggacagt gga 540
 act ccat gt g t agcgggt gga at gcgt agat at at ggaaga acaccagt gg cgaaggcggc 600
 t gt ct agt ct gt aact gacg ct gaggct cg aaagcat ggg t agcgaacag gat t agat ac 660
 cct gst agt c cat gccgt aa acgat gagt g ct aagt gt t g gagggt t t cc gccct t cagt 720
 gct gcagct a acgcat t aag cact ccgcct ggggagt acg accgcaaggt t gaaact caa 780
 aggaat t gac gggggcccg c acaagcgggt g gagcat gt gg t t t aat t cga agct acgcga 840
 agaacct t ac cagggt ct t ga cat ct t ct gc caat ct t aga gat aagacgt t ccct t cggg 900
 gacagaat ga cagggt ggt gc at ggt t gt cg t cagct cgt g t cgt gagat g t t ggggt t aag 960
 t cccgcaacg agcgcaaccc t t at t at cag t t gccagcat t cagt t gggc act ct ggt ga 1020
 gact gccggt gacaaaccgg aggaagggt gg ggat gacgt c aaat cat cat gccct t at g 1080
 acct gggct a cacacgt gct acaat ggacg gt acaacgag t cgcaaggt c gt 1132

<210> 2
 <211> 1258

P17010- NVP_Sequence l i s t . t x t

<212> DNA
<213> Unknown

<220>
<223> 16S rDNA of Lactobacillus johnsonii CH32

<400> 2
gt aact t gcc caagagact g ggat aacacc t ggaaacaga t gct aat acc ggat aacaac 60
act agacgca t gt ct agagt t t gaaagat g gt t ct gct at cact ct t gga t ggacct gcg 120
gt gcat t agc t agt t ggt aa ggt aacggct t accaaggca at gat gcat a gccgagt t ga 180
gagact gat c ggccacat t g ggact gagac acggcccaaa ct cct acggg aggcagcagt 240
agggaat ct t ccacaat gga cgaaagt ct g at ggagcaac gccgcgt gag t gaagaaggg 300
t t t cggct cg t aaagct ct g t t ggt agt ga agaaagat ag aggt agt aac t ggcct t gat 360
t t gacggg aa t t cccagaa agt cacggct aact acgt gc cagcagccgc ggt aat acgt 420
aggt ggcaag cgt t gt ccgg at t t at t ggg cgt aaagcga gt gcaggcgg t t caat aagt 480
ct gat gt gaa agcct t cggc t caaccggag aat t gcat ca gaaact gt t g aact t gagt g 540
cagaagagga gagt ggaact ccat gt gt ag cggt ggaat g cgt agat at a t ggaagaaca 600
ccagt ggcga aggcggct ct ct ggt ct gca act gacgct g aggcg caaaa gcat gggg ag 660
cgaacaggat t agat accct ggt agt ccat gccgt aaacg at gagt gct a agt gt t ggga 720
ggg t t t ccgcc t ct cagt gct gcagct aacg cat t aagcac t ccgcct ggg gagt acgacc 780
gcaagggt t ga aact caaagg aat t gacggg ggcccgcaca agcggg ggag cat gt ggt t t 840
aat t cgaagc aacgcgaaga acct t accag gt ct t gacat ccagt gcaaa cct aagagat 900
t aggt gt t cc ct t cggggac gct gagacag gt ggt gcat g gct gt cgt ca gct cgt gt cg 960
t gagat gt t g ggt t aagt cc cgcaacgagc gcaaccct t g t cat t agt t g ccat cat t aa 1020
gt t gggcact ct aat gagac t gccggg gac aaaccggagg aaggt gggga t gacgt caag 1080
t cat cat gcc cct t at gacc t gggct acac acgt gct aca at ggacggg a caacgagaag 1140
cgaacct gcg aaggcaagcg gat ct ct t aa agccgt t ct c agt t cggact gt aggct gca 1200
act cgcct ac acgaagct gg aat cgct agt aat cgcgga cagt acgccg cggt gaat 1258

<210> 3
<211> 1348
<212> DNA
<213> Unknown

<220>
<223> 16S rDNA of Bifidobacterium longum CH57

<400> 3
cgggct t t gc t t ggt ggt ga gagt ggcgaa cgggt gagt a at gcgt gacc gacct gcccc 60
at acaccgga at agct cct g gaaacgggt g gt aat gccgg at gct ccagt t gat cgcat g 120
gt ct t ct ggg aaagct t t cg cggt at ggga t ggggt cgcg t cct at cagc t t gacggcgg 180
ggg aacggcc caccgt ggct t cgacgggt a gccggcct ga gagggcgacc ggccacat t g 240

P17010- NVP_Sequence l i s t . t x t

```

ggact gagat  acggcccaga  ct cct acggg  aggcagcagt  ggggaat at t  gcacaat ggg      300
cgcaagcct g  at gcagcgac  gccgcgt gag  ggat ggaggc  ct t cgggt t g  t aaacct ct t      360
t t at cgggga  gcaagcgaga  gt gagt t t ac  ccgt t gaat a  agcaccggct  aact acgt gc      420
cagcagccgc  ggt aat acgt  agggg gcaag  cgt t at ccgg  aat t at t ggg  cgt aaagggc      480
t cgt aggcgg  t t cgt cgcgt  ccggt gt gaa  agt ccat cgc  t t aacggg gg  at ccgcgccg      540
ggg acgggcg  ggct t gagt g  cggg agggga  gact ggaat t  cccggg gt aa  cggg ggaat g      600
t gt agat at c  gggaagaaca  ccaat ggcga  aggcaggt ct  ct gggccgt t  act gacgct g      660
aggagcgaaa  gcgt ggggag  cgaacaggat  t agat accct  ggt agt ccac  gccgt aaacg      720
gt ggat gct g  gat gt ggggc  ccgt t ccacg  ggt t ccgt gt  cggagct aac  gcgt t aagca      780
t cccgcct gg  gggagt acgg  ccgcaaggct  aaaact caaa  gaaat t gacg  ggggcccgca      840
caagcgggcg  agcat gcgga  t t aat t cgt  gcaacgcgaa  gaacct t acc  t gggct t cac      900
at gt t cccga  cggg cgt aga  gat acggct t  ccct t cgggg  cgggt t caca  ggt ggt gcat      960
ggg cgt cgt c  agct cgt gt c  gt gagat gt t  gggg t aagt c  ccgcaacgag  cgcaaccct c     1020
gccccgt gt t  gccagcggat  t at gccggga  act cacgggg  gaccgccggg  gt t aact cgg     1080
aggaagggt gg  ggat gacgt c  agat cat cat  gcccct t acg  t ccagggt t  cacgcat gct     1140
acaat ggccg  gt acaacggg  at gcgacgcg  gcgacgcgga  gcggat ccct  gaaaaccggt     1200
ct cagt t cgg  at cgcagt ct  gcaact cgac  t gcgt gaagg  cggagt cgct  agt aat cgcg     1260
aat cagcaac  gt cgcggt ga  at gcgt t ccc  gggcct t gt a  cacaccgccc  gt caagt cat     1320
gaaagt gggc  agcacccgaa  gccggg gg      1348

```

<210> 4
 <211> 1421
 <212> DNA
 <213> Unknown

<220>
 <223> 16S rDNA of Lactobacillus plantarum LC5

```

<400> 4
ct ggt at t ga  t t ggt gct t g  cat cat gat t  t acat t t gag  t gagt ggcga  act ggt gagt      60
aacacgt ggg  aaacct gccc  agaagcgggg  gat aacacct  ggaaacagat  gct aat accg      120
cat aacaact  t ggaccgcat  ggt ccgagct  t gaaagat gg  ct t cggct at  cact t t t gga      180
t ggt cccgcg  gcgt at t agc  t agat ggt gg  ggt aacggct  caccat ggca  at gat acgt a      240
gccgacct ga  gagggg aat c  ggccacat t g  ggact gagac  acggcccaaa  ct cct acggg      300
aggcagcagt  agggaa ct t  ccacaat gga  cgaaagt ct g  at ggagcaac  gccgcgt gag      360
t gaagaaggg  t t t cggct cg  t aaaact ct g  t t gt t aaaga  agaaca at c  t gagagt aac      420
t gt t caggt a  t t gacggg at  t t aaccagaa  agccacggct  aact acgt gc  cagcagccgc      480
ggg aat acgt  aggt ggcaag  cgt t gt ccgg  gat t t at t gg  gcgt aaagcg  agcgcaggcg      540

```

P17010- NVP_Sequence l i s t . t x t

gt t t t t t aag t ct gat gt ga aagcct t cgg ct caaccgaa gaagt gcat c ggaaact ggg	600
aaact t gagt gcagaagagg acagt ggaac t ccat gt gt a gcggt gaaat gcgt agat at	660
at ggaagaac accagt ggcg aaggcggct g t ct ggt ct gt aact gacgct gaggct cgaa	720
agt at ggg t a gcaaacagga t t agat accc t ggt agt cca t accgt aaac gat gaat gct	780
aagt gt t gga ggg t t t ccgc cct t cagt gc t gcagct aac gcat t aagca t t ccgcct gg	840
ggagt acggc cgcaaggct g aaact caaag gaat t gacgg gggcccgcac aagcggg gga	900
gcat gt ggt t t aat t cgaag ct acgcgaag aacct t acca ggt ct t gaca t act at gcaa	960
at ct aagaga t t agacgt t c cct t cgggga cat ggat aca ggt ggt gcat ggt t gt cgt c	1020
agct cgt gt c gt gagat gt t ggg t t aagt c ccgcaacgag cgcaaccct t at t at cagt t	1080
gccagcat t a agt t gggcac t ct ggt gaga ct gccggt ga caaaccggag gaaggt gggg	1140
at gacgt caa at cat cat gc ccct t at gac ct gggct aca cacgt gct ac aat ggat ggt	1200
acaacgagt t gcgaact cgc gagagt aagc t aat ct ct t a aagccat t ct cagt t cggat	1260
t gt aggct gc aact cgcct a cat gaagt cg gaat cgct ag t aat cgcgga t cagcat gcc	1320
gcggt gaat a cgt t cccggg cct t gt acac accgcccgt c acaccat gag agt t t gt aac	1380
acccaaagt c ggt ggggt aa cct t t t agga accagccgcc t	1421

<210> 5
 <211> 1356
 <212> DNA
 <213> Unknown

<220>
 <223> 16S rDNA of Lactobacillus plantarum LC27

<400> 5 t t t acat t t g at t gagt ggc gaact ggt ga gt aacacgt g ggaaacct gc ccagaagcgg	60
gggat aacac ct ggaaacag at gct aat ac cgc at aacaa ct t ggaccgc at ggt ccgag	120
t t t gaaagat ggct t cggct at cact t t t g gat ggt cccg cggcgt at t a gct agat ggt	180
ggggt aacgg ct caccat gg caat gat acg t agccgacct gagagggt aa t cggccacat	240
t gggact gag acacggccca aact cct acg ggaggcagca gt agggaat c t t ccacaat g	300
gacgaaagt c t gat ggagca acgcccgt g agt gaagaag ggt t t cggct cgt aaaact c	360
t gt t gt t aaa gaagaacat a t ct gagagt a act gt t cagg t at t gacggt at t t aaccag	420
aaagccacgg ct aact acgt gccagcagcc gcggt aat ac gt aggt ggca agcgt t gt cc	480
ggat t t at t g ggcgt aaagc gagcgcaggc ggt t t t t aa gt ct gat gt g aaagcct t cg	540
gct caaccga agaagt gcat cggaaact gg gaaact t gag t gcagaagag gacagt ggaa	600
ct ccat gt gt agcggg gaaa t gcgt agat a t at ggaagaa caccagt ggc gaaggcggct	660
gt ct ggt ct g t aact gacgc t gaggct cga aagt at ggg t agcaaacagg at t agat acc	720
ct ggt agt cc at accgt aaa cgat gaat gc t aagt gt t gg aggg t t ccg ccct t cagt g	780
ct gcagct aa cgc at t aagc at t ccgcct g gggagt acgg ccgcaaggct gaaact caaa	840

P17010- NVP_Sequence l i s t . t x t

ggaat t gacg ggggcccga caagcggg gg agcat gt ggt t t aat t cgaa gct acgcgaa	900
gaacct t acc aggt ct t gac at act at gca aat ct aagag at t agacgt t ccct t cgggg	960
acat ggat ac aggt ggt gca t ggt t gt cgt cagct cgt gt cgt gagat gt t gggg t aagt	1020
cccgcaacga gcgcaaccct t at t at cagt t gccagcat t aagt t gggca ct ct ggt gag	1080
act gccggg g acaaaccgga ggaaggt ggg gat gacgt ca aat cat cat g ccct t at ga	1140
cct gggct ac acacgt gct a caat ggat gg t acaacgagt t gcgaact cg cgagagt aag	1200
ct aat ct ct t aaagccat t c t cagt t cgga t t gt aggct g caact cgcct acat gaagt c	1260
ggaat cgct a gt aat cgcgg at cagcat gc cgcggg gaat acgt t cccgg gcct t gt aca	1320
caccgcccgt cacaccat ga gagt t t gt aa caccga	1356

<210> 6
 <211> 1325
 <212> DNA
 <213> Unknown

<220>
 <223> 16S rDNA of Lact obaci l l u s pl ant ar um LC28

<400> 6	
cat cat gat t t acat t t gag t gagt ggcga act ggt gagt aacacgt ggg aaacct gccc	60
agaagcgggg gat aacacct ggaaacagat gct aat accg cat aacaact t ggaccgcat	120
ggg ccgagt t t gaaagat gg ct t cggct at cact t t t gga t ggt cccgcg gcgt at t agc	180
t agat ggt gg ggt aacggct caccat ggca at gat acgt a gccgacct ga gagggg aat c	240
ggccacat t g ggact gagac acggcccaaa ct cct acggg aggcagcagt aggggaat ct t	300
ccacaat gga cgaaagt ct g at ggagcaac gccgcgt gag t gaagaaggg t t t cggct cg	360
t aaaact ct g t t gt t aaaga agaacat at c t gagagt aac t gt t caggt a t t gacggg at	420
t t aaccagaa agccacggct aact acgt gc cagcagccgc ggt aat acgt aggt ggcaag	480
cgt t gt ccgg at t t at t ggg cgt aaagcga gcgcaggcgg t t t t t aagt ct gat gt gaa	540
agcct t cggc t caaccgaag aagt gcat cg gaaact ggga aact t gagt g cagaagagga	600
cagt ggaact ccat gt gt ag cggg gaaat g cgt agat at a t ggaagaaca ccagt ggcga	660
aggcggct gt ct ggt ct gt a act gacgct g aggct cgaaa gt at gggg ag caaacaggat	720
t agat accct ggt agt ccat accgt aaacg at gaat gct a agt gt t ggag ggt t t ccgcc	780
ct t cagt gct gcagct aacg cat t aagcat t ccgcct ggg gagt acggcc gcaaggct ga	840
aact caaagg aat t gacggg ggcccgcaca agcggg ggag cat gt ggt t t aat t cgaagc	900
t acgcgaaga acct t accag gt ct t gacat act at gcaaa t ct aagagat t agacgt t cc	960
ct t cgggggac at ggat acag gt ggt gcat g gt t gt cgt ca gct cgt gt cg t gagat gt t g	1020
ggg t aagt cc cgcaacgagc gcaaccct t a t t at cagt t g ccagcat t aa gt t gggcact	1080
ct ggt gagac t gccggg gac aaaccggagg aagggt gggga t gacgt caaa t cat cat gcc	1140

P17010- NVP_Sequence l i s t . t x t

cct t at gacc t gggct acac acgt gct aca at ggat ggt a caacgagt t g cgaact cgcg	1200
agagt aagct aat ct ct t aa agccat t ct c agt t cggat t gt aggct gca act cgcct ac	1260
at gaagt cgg aat cgct agt aat cgcggat cagcat t ccg cggg gaat ac gt t cccgggc	1320
ct t gt	1325

<210> 7
 <211> 1342
 <212> DNA
 <213> Unknown

<220>
 <223> 16S rDNA of Bi f i dobact eri um l ongum LC67

<400> 7	
ggct t t gct t ggt ggt gaga gt ggcgaacg ggt gagt aat gcgt gaccga cct gccccat	60
acaccggaat agct cct gga aacgggt ggt aat gccggat gct ccagt t g at cgcct ggt	120
ct t ct gggaa agct t t cgcg gt at gggat g gggg cgcgt c ct at cagct t gacggcgggg	180
t aacggccca ccgt ggct t c gacgggt agc cggcct gaga gggcgaccgg ccacat t ggg	240
act gagat ac ggcccagact cct acgggag gcagcagt gg ggaat at t gc acaat gggcg	300
caagcct gat gcagcgacgc cgcgt gaggg at ggaggcct t cgggt t gt a aacct ct t t t	360
at cggggagc aagcgagagt gagt t t accc gt t gaat aag caccggct aa ct acgt gccca	420
gcagccgcgg t aat acgt ag ggt gcaagcg t t at cccgga at t at t gggc gt aaagggt	480
cgt aggcgt t cgt cgcgt c cgggt gt gaaa gt ccat cgct t aacggg gga t ccgcgcgg	540
gt acgggcgg gct t gagt gc ggt aggggag act ggaat t c ccgggt gt aac ggt ggaat gt	600
gt agat at cg ggaagaacac caat ggcgaa ggcaggt ct c t gggccgt t a ct gacgt ga	660
ggagcgaaag cgt ggggagc gaacaggat t agat accct g gt agt ccacg ccgt aaacgg	720
t ggat gct gg at gt ggggcc cgt t ccacgg gt t ccgt gt c ggagct aacg cgt t aagcat	780
ccgcct ggg ggagt acggc cgcaaggct a aaact caaag aaat t gacgg gggcccgcac	840
aagcggcgga gcat gcggat t aat t cgat g caacgcgaag aacct t acct gggct t gaca	900
t gt t cccgac ggt cgt agag at acggct t c cct t cggggc ggggt t cacag gt ggt gcat g	960
gt cgt cgt ca gct cgt gt cg t gagat gt t g ggt t aagt cc cgcaacgagc gcaaccct cg	1020
ccccgt gt t g ccagcggat t at gccgggaa ct cacggggg accgccgggg t t aact cgga	1080
ggaaggt ggg gat gacgt ca gat cat cat g cccct t acgt ccagggt t c acgcat gct a	1140
caat ggccgg t acaacggga t gcgacgcgg cgacgcggag cggat ccct g aaaaccggt c	1200
t cagt t cgga t cgcagt ct g caact cgact gcgt gaaggc ggagt cgct a gt aat cgcga	1260
at cagcaacg t cgcgggt gaa t gcgt t cccg ggcct t gt ac acaccgcccg t caagt cat g	1320
aaagt gggca gcacccgaag cg	1342

<210> 8
 <211> 1343

P17010- NVP_Sequence l i s t . t x t

<212> DNA
<213> Unknown

<220>
<223> 16S rDNA of Bifidobacterium longum LC68

```

<400>      8
cggtt cggg t gct gcccac t t t cat gact t gacggg cgg t gt gt acaag gcccgggaac      60
gcat t caccg cgacgt t gct gat t cgcat t act agcgac t ccgcct t ca cgacgt cgag      120
t t gcagact g cgat ccgaac t gagaccggt t t t cagggat ccgct ccgcg t cgccgcgt c      180
gcat cccgt t gt accggcca t t gt agcat g cgt gaagccc t ggacgt aag gggcat gat g      240
at ct gacgt c at cccacct t cct ccgagt t aaccccggc ggt ccccggt gagt t cccgg      300
cat aat ccgc t ggcaacacg gggcgagggt t gcgct cgt t gcgggact t a acccaacat c      360
t cacgacacg agct gacgac gacat gcac cacct gt gaa cccgccccga aggaagccg      420
cat ct ct acg accgt cggga acat gt caag cccaggt aag gt t ct t cgcg t t gcat cgaa      480
t t aat ccgca t gct ccgccg ct t gt gcggg ccccggt caa t t t ct t t gag t t t t agcct t      540
gcggccgt ac t cccagggc ggat gct t aa cgct t agct ccgacacgga acccggt ggaa      600
cgggccccac at ccagcat c caccgt t t ac ggcgt ggact accagggt at ct aat cct gt      660
t cgct ccca cgct t t cgct cct cagcgt c agt aacggcc cagagacct g cct t cgccat      720
t ggt gt t ct t cccgat at ct acacat t cca ccgt t acacc gggaat t cca gt ct cccct a      780
ccgcact caa gcccggccgt acccggcgcg gat ccaccgt t aagcgat gg act t t cacac      840
cggacgcgac gaaccgcct a cgagccct t t acgccaat a at t ccggat a acgct t gcac      900
cct acgt at t accgcggt g ct ggcacgt a gt t agccggt gct t at t caa cgggt aaact      960
cact ct cgct t gct cccga t aaaagaggt t t acaaccg aaggcct cca t ccct cacgc      1020
ggcgt cgct g cat caggct t gcgccccat t g t gcaat at t c cccact gct g cct cccgt ag      1080
gagt ct gggc cgt at ct cag t cccaat gt g gccggt cgcc ct ct caggcc ggct acccg      1140
cgaagccacg gt gggccgt t accccgccgt caagct gat a ggacgcgacc ccat cccat a      1200
ccgcgaaagc t t t cccagaa gacat gcga t caact ggag cat ccggcat t accaccgt      1260
t t ccaggagc t at t ccggt g t at ggggcag gt cggt cacg cat t act cac ccgt t cgcca      1320
ct ct caccac caagcaaagt ct c      1343

```