



(51) International Patent Classification:

C07D 311/78 (2006.01) **A61P 3/04** (2006.01)
A61K 31/353 (2006.01)

(21) International Application Number:

PCT/US2013/021914

(22) International Filing Date:

17 January 2013 (17.01.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/587,825 18 January 2012 (18.01.2012) US

(71) Applicant: **ZAFGEN, INC.** [US/US]; One Broadway, 8th Floor, Cambridge, MA 02142 (US).

(72) Inventors: **PALLIN, Thomas, David**; 8/9 Spire Green Centre, Flex Meadow, Harlow, Essex CM19 5TR (GB). **DYKE, Hazel, Joan**; 8/9 Spire Green Centre, Flex Meadow, Harlow, Essex CM19 5TR (GB). **CRAMP, Susan, Mary**; 8/9 Spire Green Centre, Flex Meadow, Harlow, Essex CM19 5TR (GB). **ZAHLER, Robert**; One Broadway, 8th Floor, Cambridge, MA 02142 (US).

(74) Agents: **KAVANAUGH, Theresa, C.** et al.; GOODWIN PROCTER LLP, Exchange Place, Boston, MA 02109 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: TRICYCLIC SULFONE COMPOUNDS AND METHODS OF MAKING AND USING SAME

(57) Abstract: The invention provides tricyclic sulfone compounds and their use in treating medical disorders, such as obesity. Pharmaceutical compositions and methods of making various tricyclic compounds are provided. The compounds are contemplated to have activity against methionyl aminopeptidase 2.



- 1 -

TRICYCLIC SULFONE COMPOUNDS AND METHODS OF MAKING AND USING SAME

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application No. 61/587,825, filed January 18, 2012, hereby incorporated by reference in its entirety.

BACKGROUND

[0002] Over 1.1 billion people worldwide are reported to be overweight. Obesity is
5 estimated to affect over 90 million people in the United States alone. Twenty-five percent of the population in the United States over the age of twenty is considered clinically obese. While being overweight or obese presents problems (for example restriction of mobility, discomfort in tight spaces such as theater or airplane seats, social difficulties, etc.), these conditions, in particular clinical obesity, affect other aspects of health, i.e., diseases and other adverse health
10 conditions associated with, exacerbated by, or precipitated by being overweight or obese. The estimated mortality from obesity-related conditions in the United States is over 300,000 annually (O'Brien et al. *Amer J Surgery* (2002) 184:4S-8S; and Hill et al. (1998) *Science*, 280:1371).

[0003] There is no curative treatment for being overweight or obese. Traditional
15 pharmacotherapies for treating an overweight or obese subject, such as serotonin and noradrenergic re-uptake inhibitors, noradrenergic re-uptake inhibitors, selective serotonin re-uptake inhibitors, intestinal lipase inhibitors, or surgeries such as stomach stapling or gastric banding, have been shown to provide minimal short-term benefits or significant rates of relapse, and have further shown harmful side-effects to patients.

20 [0004] MetAP2 encodes a protein that functions at least in part by enzymatically removing the amino terminal methionine residue from certain newly translated proteins such as glyceraldehyde-3-phosphate dehydrogenase (Warder *et al.* (2008) *J Proteome Res* 7:4807). Increased expression of the MetAP2 gene has been historically associated with various forms of cancer. Molecules inhibiting the enzymatic activity of MetAP2 have been identified and

- 2 -


have been explored for their utility in the treatment of various tumor types (Wang et al. (2003) Cancer Res. 63:7861) and infectious diseases such as microsporidiosis, leishmaniasis, and malaria (Zhang et al. (2002) J. Biomed. Sci. 9:34). Notably, inhibition of MetAP2 activity in obese and obese-diabetic animals leads to a reduction in body weight in part by increasing the oxidation of fat and in part by reducing the consumption of food (Rupnick et al. (2002) Proc. Natl. Acad. Sci. USA 99:10730).

[0005] Such MetAP2 inhibitors may be useful as well for patients with excess adiposity and conditions related to adiposity including type 2 diabetes, hepatic steatosis, and cardiovascular disease (*via* e.g. ameliorating insulin resistance, reducing hepatic lipid content, and reducing cardiac workload). Accordingly, compounds capable of modulating MetAP2 are needed to address the treatment of obesity and related diseases as well as other ailments favorably responsive to MetAP2 modulator treatment.

SUMMARY

[0006] The invention provides, for example, compounds which may be modulators of MetAP2, and their use as medicinal agents, processes for their preparation, and pharmaceutical compositions containing them as an active ingredient both alone or in combination with other agents, as well as provides for their use as medicaments and/or in the manufacture of medicaments for the inhibition of MetAP2 activity in warm-blooded animals such as humans. In particular this invention relates to compounds useful for the treatment of obesity, type 2 diabetes, and other obesity-associated conditions. Also provided are pharmaceutical compositions comprising at least one disclosed compound and a pharmaceutically acceptable carrier.

[0007] In an embodiment, provided herein are compounds represented by formula I:



Formula I

or pharmaceutically acceptable salts, stereoisomers, esters or prodrugs thereof, where A, D, R^{A1} , R^{A2} , R^{B1} , R^{B2} , R^{B3} , R^{B4} , X, and n are as defined herein.

DETAILED DESCRIPTION

[0008] The features and other details of the disclosure will now be more particularly described. Before further description of the present invention, certain terms employed in the specification, examples and appended claims are collected here. These definitions should be read in light of the remainder of the disclosure and as understood by a person of skill in the art.

5 Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by a person of ordinary skill in the art.

Definitions

[0009] “Treating” includes any effect, e.g., lessening, reducing, modulating, or eliminating, that results in the improvement of the condition, disease, disorder and the like.

10 [0010] The term “alkenyl” as used herein refers to an unsaturated straight or branched hydrocarbon having at least one carbon-carbon double bond. Exemplary alkenyl groups include, but are not limited to, a straight or branched group of 2-6 or 3-4 carbon atoms, referred to herein as C₂₋₆alkenyl, and C₃₋₄alkenyl, respectively. Exemplary alkenyl groups include, but are not limited to, vinyl, allyl, butenyl, pentenyl, etc.

15 [0011] The term “alkoxy” as used herein refers to a straight or branched alkyl group attached to oxygen (alkyl-O-). Exemplary alkoxy groups include, but are not limited to, alkoxy groups of 1-6 or 2-6 carbon atoms, referred to herein as C₁₋₆alkoxy, and C₂₋₆alkoxy, respectively. Exemplary alkoxy groups include, but are not limited to methoxy, ethoxy, isopropoxy, etc.

20 [0012] The term “alkoxyalkyl” as used herein refers to a straight or branched alkyl group attached to oxygen, attached to a second straight or branched alkyl group (alkyl-O-alkyl-). Exemplary alkoxyalkyl groups include, but are not limited to, alkoxyalkyl groups in which each of the alkyl groups independently contains 1-6 carbon atoms, referred to herein as C₁₋₆alkoxy-C₁₋₆alkyl. Exemplary alkoxyalkyl groups include, but are not limited to
25 methoxymethyl, 2-methoxyethyl, 1-methoxyethyl, 2-methoxypropyl, ethoxymethyl, 2-isopropoxyethyl etc.

[0013] The term “alkyoxycarbonyl” as used herein refers to a straight or branched alkyl group attached to oxygen, attached to a carbonyl group (alkyl-O-C(O)-). Exemplary alkyoxycarbonyl groups include, but are not limited to, alkyoxycarbonyl groups of 1-6 carbon

- 4 -

atoms, referred to herein as C₁₋₆alkoxycarbonyl. Exemplary alkoxycarbonyl groups include, but are not limited to, methoxycarbonyl, ethoxycarbonyl, t-butoxycarbonyl, etc.

[0014] The term “alkenyloxy” used herein refers to a straight or branched alkenyl group attached to oxygen (alkenyl-O-). Exemplary alkenyloxy groups include, but are not limited to, groups with an alkenyl group of 3-6 carbon atoms, referred to herein as C₃₋₆alkenyloxy. Exemplary “alkenyloxy” groups include, but are not limited to allyloxy, butenyloxy, etc.

[0015] The term “alkynyloxy” used herein refers to a straight or branched alkynyl group attached to oxygen (alkynyl-O). Exemplary alkynyloxy groups include, but are not limited to, groups with an alkynyl group of 3-6 carbon atoms, referred to herein as C₃₋₆alkynyloxy. Exemplary alkynyloxy groups include, but are not limited to, propynyloxy, butynyloxy, etc.

[0016] The term “alkyl” as used herein refers to a saturated straight or branched hydrocarbon. Exemplary alkyl groups include, but are not limited to, straight or branched hydrocarbons of 1-6, 1-4, or 1-3 carbon atoms, referred to herein as C₁₋₆alkyl, C₁₋₄alkyl, and C₁₋₃alkyl, respectively. Exemplary alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, 2-methyl-1-butyl, 3-methyl-2-butyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, butyl, isobutyl, t-butyl, pentyl, isopentyl, neopentyl, hexyl, etc.

[0017] The term “alkylcarbonyl” as used herein refers to a straight or branched alkyl group attached to a carbonyl group (alkyl-C(O)-). Exemplary alkylcarbonyl groups include, but are not limited to, alkylcarbonyl groups of 1-6 atoms, referred to herein as C₁₋₆alkylcarbonyl groups. Exemplary alkylcarbonyl groups include, but are not limited to, acetyl, propanoyl, isopropanoyl, butanoyl, etc.

[0018] The term “alkynyl” as used herein refers to an unsaturated straight or branched hydrocarbon having at least one carbon-carbon triple bond. Exemplary alkynyl groups include, but are not limited to, straight or branched groups of 2-6, or 3-6 carbon atoms, referred to herein as C₂₋₆alkynyl, and C₃₋₆alkynyl, respectively. Exemplary alkynyl groups include, but are not limited to, ethynyl, propynyl, butynyl, pentynyl, hexynyl, methylpropynyl, etc.

[0019] The term “carbonyl” as used herein refers to the radical -C(O)-.

[0020] The term “cyano” as used herein refers to the radical -CN.

- 5 -

[0021] The term “cycloalkoxy” as used herein refers to a cycloalkyl group attached to oxygen (cycloalkyl-O-). Exemplary cycloalkoxy groups include, but are not limited to, cycloalkoxy groups of 3-6 carbon atoms, referred to herein as C₃₋₆cycloalkoxy groups. Exemplary cycloalkoxy groups include, but are not limited to, cyclopropoxy, cyclobutoxy, cyclohexyloxy, etc

[0022] The terms “cycloalkyl” or a “carbocyclic group” as used herein refers to a saturated or partially unsaturated hydrocarbon group of, for example, 3-6, or 4-6 carbons, referred to herein as C₃₋₆cycloalkyl or C₄₋₆cycloalkyl, respectively. Exemplary cycloalkyl groups include, but are not limited to, cyclohexyl, cyclopentyl, cyclopentenyl, cyclobutyl or cyclopropyl.

[0023] The terms “halo” or “halogen” as used herein refer to F, Cl, Br, or I.

[0024] The terms “heteroaryl” or “heteroaromatic group” as used herein refers to a monocyclic aromatic 5-6 membered ring system containing one or more heteroatoms, for example one to three heteroatoms, such as nitrogen, oxygen, and sulfur. Where possible, said heteroaryl ring may be linked to the adjacent radical through carbon or nitrogen. Examples of heteroaryl rings include but are not limited to furan, thiophene, pyrrole, thiazole, oxazole, isothiazole, isoxazole, imidazole, pyrazole, triazole, pyridine or pyrimidine etc.

[0025] The terms “heterocyclyl” or “heterocyclic group” are art-recognized and refer to saturated or partially unsaturated 4-7 membered ring structures, whose ring structures include one to three heteroatoms, such as nitrogen, oxygen, and sulfur. Where possible, heterocyclyl rings may be linked to the adjacent radical through carbon or nitrogen. Examples of heterocyclyl groups include, but are not limited to, pyrrolidine, piperidine, morpholine, thiomorpholine, piperazine, oxetane, azetidine, tetrahydrofuran or dihydrofuran etc.

[0026] The term “heterocyclyloxy” as used herein refers to a heterocyclyl group attached to oxygen (heterocyclyl-O-).

[0027] The term “heteroaryloxy” as used herein refers to a heteroaryl group attached to oxygen (heteroaryl-O-).

[0028] The terms “hydroxy” and “hydroxyl” as used herein refers to the radical -OH.

[0029] The term “oxo” as used herein refers to the radical =O.

- 6 -

[0030] “Pharmaceutically or pharmacologically acceptable” include molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, or a human, as appropriate. For human administration, preparations should meet sterility, pyrogenicity, and general safety and purity standards as required by FDA Office of Biologics standards.

[0031] The term “pharmaceutically acceptable carrier” or “pharmaceutically acceptable excipient” as used herein refers to any and all solvents, dispersion media, coatings, isotonic and absorption delaying agents, and the like, that are compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. The compositions may also contain other active compounds providing supplemental, additional, or enhanced therapeutic functions.

[0032] The term “pharmaceutical composition” as used herein refers to a composition comprising at least one compound as disclosed herein formulated together with one or more pharmaceutically acceptable carriers.

[0033] “Individual,” “patient,” or “subject” are used interchangeably and include any animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans. The compounds of the invention can be administered to a mammal, such as a human, but can also be administered to other mammals such as an animal in need of veterinary treatment, *e.g.*, domestic animals (*e.g.*, dogs, cats, and the like), farm animals (*e.g.*, cows, sheep, pigs, horses, and the like) and laboratory animals (*e.g.*, rats, mice, guinea pigs, and the like). The mammal treated in the methods of the invention is desirably a mammal in which treatment of obesity or weight loss is desired. “Modulation” includes antagonism (*e.g.*, inhibition), agonism, partial antagonism and/or partial agonism.

[0034] In the present specification, the term “therapeutically effective amount” means the amount of the subject compound that will elicit the biological or medical response of a tissue, system or animal, (*e.g.* mammal or human) that is being sought by the researcher, veterinarian, medical doctor or other clinician. The compounds of the invention are administered in therapeutically effective amounts to treat a disease. Alternatively, a therapeutically effective amount of a compound is the quantity required to achieve a desired therapeutic and/or prophylactic effect, such as an amount which results in weight loss.

- 7 -

[0035] The term "pharmaceutically acceptable salt(s)" as used herein refers to salts of acidic or basic groups that may be present in compounds used in the compositions.

Compounds included in the present compositions that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable acid addition salts of such basic compounds are those that form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, including, but not limited to, malate, oxalate, chloride, bromide, iodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate and pamoate (i.e., 1,1'-methylene-*bis*-(2-hydroxy-3-naphthoate)) salts. Compounds included in the present compositions that are acidic in nature are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include alkali metal or alkaline earth metal salts, particularly calcium, magnesium, sodium, lithium, zinc, potassium, and iron salts. Compounds included in the present compositions that include a basic or acidic moiety may also form pharmaceutically acceptable salts with various amino acids. The compounds of the disclosure may contain both acidic and basic groups; for example, one amino and one carboxylic acid group. In such a case, the compound can exist as an acid addition salt, a zwitterion, or a base salt.

[0036] The compounds of the disclosure may contain one or more chiral centers and, therefore, exist as stereoisomers. The term "stereoisomers" when used herein consist of all enantiomers or diastereomers. These compounds may be designated by the symbols "(+)," "(-)," "R" or "S," depending on the configuration of substituents around the stereogenic carbon atom, but the skilled artisan will recognize that a structure may denote a chiral center implicitly. The present invention encompasses various stereoisomers of these compounds and mixtures thereof. Mixtures of enantiomers or diastereomers may be designated "(±)" in nomenclature, but the skilled artisan will recognize that a structure may denote a chiral center implicitly.

[0037] The compounds of the disclosure may contain one or more double bonds and, therefore, exist as geometric isomers resulting from the arrangement of substituents around a carbon-carbon double bond. The symbol $\equiv$ denotes a bond that may be a single, double or triple bond as described herein. Substituents around a carbon-carbon double bond are

- 8 -

designated as being in the “*Z*” or “*E*” configuration wherein the terms “*Z*” and “*E*” are used in accordance with IUPAC standards. Unless otherwise specified, structures depicting double bonds encompass both the “*E*” and “*Z*” isomers. Substituents around a carbon-carbon double bond alternatively can be referred to as “*cis*” or “*trans*,” where “*cis*” represents substituents on the same side of the double bond and “*trans*” represents substituents on opposite sides of the double bond.

[0038] Compounds of the disclosure may contain a carbocyclic or heterocyclic ring and therefore, exist as geometric isomers resulting from the arrangement of substituents around the ring. The arrangement of substituents around a carbocyclic or heterocyclic ring are designated as being in the “*Z*” or “*E*” configuration wherein the terms “*Z*” and “*E*” are used in accordance with IUPAC standards. Unless otherwise specified, structures depicting carbocyclic or heterocyclic rings encompass both “*Z*” and “*E*” isomers. Substituents around a carbocyclic or heterocyclic rings may also be referred to as “*cis*” or “*trans*”, where the term “*cis*” represents substituents on the same side of the plane of the ring and the term “*trans*” represents substituents on opposite sides of the plane of the ring. Mixtures of compounds wherein the substituents are disposed on both the same and opposite sides of plane of the ring are designated “*cis/trans*.”

[0039] Individual enantiomers and diastereomers of compounds of the present invention can be prepared synthetically from commercially available starting materials that contain asymmetric or stereogenic centers, or by preparation of racemic mixtures followed by resolution methods well known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and liberation of the optically pure product from the auxiliary, (2) salt formation employing an optically active resolving agent, (3) direct separation of the mixture of optical enantiomers on chiral liquid chromatographic columns or (4) kinetic resolution using stereoselective chemical or enzymatic reagents. Racemic mixtures can also be resolved into their component enantiomers by well known methods, such as chiral-phase liquid chromatography or crystallizing the compound in a chiral solvent. Stereoselective syntheses, a chemical or enzymatic reaction in which a single reactant forms an unequal mixture of stereoisomers during the creation of a new stereocenter or during the transformation of a pre-existing one, are well known in the art. Stereoselective syntheses encompass both enantio- and diastereoselective

- 9 -

transformations, and may involve the use of chiral auxiliaries. For examples, see Carreira and Kvaerno, *Classics in Stereoselective Synthesis*, Wiley-VCH: Weinheim, 2009.

[0040] The compounds disclosed herein can exist in solvated as well as unsolvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like, and it is intended that the invention embrace both solvated and unsolvated forms. In one embodiment, the compound is amorphous. In one embodiment, the compound is a single polymorph. In another embodiment, the compound is a mixture of polymorphs. In another embodiment, the compound is in a crystalline form.

[0041] The invention also embraces isotopically labeled compounds of the invention which are identical to those recited herein, except that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, sulfur, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively. For example, a compound of the invention may have one or more H atom replaced with deuterium.

[0042] Certain isotopically-labeled disclosed compounds (*e.g.*, those labeled with ^3H and ^{14}C) are useful in compound and/or substrate tissue distribution assays. Tritiated (*i.e.*, ^3H) and carbon-14 (*i.e.*, ^{14}C) isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium (*i.e.*, ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability (*e.g.*, increased *in vivo* half-life or reduced dosage requirements) and hence may be preferred in some circumstances. Isotopically labeled compounds of the invention can generally be prepared by following procedures analogous to those disclosed in the examples herein by substituting an isotopically labeled reagent for a non-isotopically labeled reagent.

[0043] The term "prodrug" refers to compounds that are transformed *in vivo* to yield a disclosed compound or a pharmaceutically acceptable salt, hydrate or solvate of the compound. The transformation may occur by various mechanisms (such as by esterase, amidase, phosphatase, oxidative and or reductive metabolism) in various locations (such as in the intestinal lumen or upon transit of the intestine, blood or liver). Prodrugs are well known in the art (for example, see Rautio, Kumpulainen, *et al*, Nature Reviews Drug Discovery 2008, 7, 255). For example, if a compound of the invention or a pharmaceutically acceptable salt,

- 10 -

hydrate or solvate of the compound contains a carboxylic acid functional group, a prodrug can comprise an ester formed by the replacement of the hydrogen atom of the acid group with a group such as (C₁₋₈)alkyl, (C₂₋₁₂)alkylcarbonyloxymethyl, 1-(alkylcarbonyloxy)ethyl having from 4 to 9 carbon atoms, 1-methyl-1-(alkylcarbonyloxy)-ethyl having from 5 to 10 carbon atoms, alkoxycarbonyloxymethyl having from 3 to 6 carbon atoms, 1-(alkoxycarbonyloxy)ethyl having from 4 to 7 carbon atoms, 1-methyl-1-(alkoxycarbonyloxy)ethyl having from 5 to 8 carbon atoms, N-(alkoxycarbonyl)aminomethyl having from 3 to 9 carbon atoms, 1-(N-(alkoxycarbonyl)amino)ethyl having from 4 to 10 carbon atoms, 3-phthalidyl, 4-crotonolactonyl, gamma-butyrolacton-4-yl, di-N,N-(C₁₋₂)alkylamino(C₂₋₃)alkyl (such as β-dimethylaminoethyl), carbamoyl-(C₁₋₂)alkyl, N,N-di(C₁₋₂)alkylcarbamoyl-(C₁₋₂)alkyl and piperidino-, pyrrolidino- or morpholino(C₂₋₃)alkyl.


[0044] Similarly, if a compound of the invention contains an alcohol functional group, a prodrug can be formed by the replacement of the hydrogen atom of the alcohol group with a group such as (C₁₋₆)alkylcarbonyloxymethyl, 1-((C₁₋₆)alkylcarbonyloxy)ethyl, 1-methyl-1-((C₁₋₆)alkylcarbonyloxy)ethyl (C₁₋₆)alkoxycarbonyloxymethyl, N-(C₁₋₆)alkoxycarbonylaminomethyl, succinoyl, (C₁₋₆)alkylcarbonyl, α-amino(C₁₋₄)alkylcarbonyl, arylalkylcarbonyl and α-aminoalkylcarbonyl, or α-aminoalkylcarbonyl-α-aminoalkylcarbonyl, where each α-aminoalkylcarbonyl group is independently selected from the naturally occurring L-amino acids, P(O)(OH)₂, -P(O)(O(C₁₋₆)alkyl)₂ or glycosyl (the radical resulting from the removal of a hydroxyl group of the hemiacetal form of a carbohydrate).

[0045] If a compound of the invention incorporates an amine functional group, a prodrug can be formed, for example, by creation of an amide or carbamate, an N-alkylcarbonyloxyalkyl derivative, an (oxodioxolenyl)methyl derivative, an N-Mannich base, imine or enamine. In addition, a secondary amine can be metabolically cleaved to generate a bioactive primary amine, or a tertiary amine can be metabolically cleaved to generate a bioactive primary or secondary amine. For examples, see Simpicio, *et al.*, *Molecules* 2008, 13, 519 and references therein.

I. Tricyclic Compounds

[0046] In certain embodiments, the present invention provides compounds of Formula I:

- 11 -



Formula I

wherein

D may be a 5-7 membered heterocyclic or carbocyclic ring;

X may be selected from the group consisting of: $^+-C(R^{D1}R^{D2})-C(R^{D5}R^{D6})-^*$, $^+-W^1-$
 5 $C(R^{D5}R^{D6})-^*$, $^+-W^1-C(O)-^*$, $^+-C(R^{D1}R^{D2})-W^3-^*$, $^+-C(R^{D1}R^{D2})-C(R^{D3}R^{D4})-C(R^{D5}R^{D6})-^*$, $^+-W^1-$
 $C(R^{D3}R^{D4})-C(R^{D5}R^{D6})-^*$, $^+-W^1-C(O)-C(R^{D5}R^{D6})-^*$, $^+-C(R^{D1}R^{D2})-W^2-C(R^{D5}R^{D6})-^*$, $^+-$
 $C(R^{D1}R^{D2})-W^2-C(O)-^*$, $^+-C(R^{D1}R^{D2})-C(R^{D3}R^{D4})-W^3-^*$, $^+-C(R^{D1}R^{D2})-C(O)-W^3-^*$, in which the
 attachment points are indicated by $^+$ and * in X and in Formula I;

W^1 may be selected from the group consisting of: O or $N(R^{N1})$

10 W^2 may be selected from the group consisting of: O or $N(R^{N2})$

W^3 may be selected from the group consisting of: O or $N(R^{N3})$

R^{B1} and R^{B2} may be independently selected from the group consisting of hydrogen,
 fluorine, hydroxyl, CN, C_{1-3} alkyl or C_{1-2} alkoxy; wherein the C_{1-3} alkyl and C_{1-2} alkoxy groups
 may be optionally substituted by one or more fluorine atoms or a cyano, hydroxyl or C_{1-2} alkoxy
 15 group.

R^{B3} and R^{B4} may be independently selected from the group consisting of hydrogen,
 fluorine or C_{1-2} alkyl group; wherein the C_{1-2} alkyl group may be optionally substituted by one or
 more fluorine atoms.

A may be a ring selected from the group consisting of: phenyl, a 5-6 membered
 20 heteroaryl having 1, 2 or 3 heteroatoms each selected from S, N or O, and a 4-7 membered
 heterocycle having 1, 2 or 3 heteroatoms each selected from N or O;

R^{A1} may be selected, independently for each occurrence, from the group consisting of:
 hydrogen, hydroxyl, cyano, halogen, C_{1-4} alkyl or C_{1-3} alkoxy; wherein C_{1-4} alkyl, or C_{1-3} alkoxy
 may be optionally substituted by one or more fluorines;

25 n may be 1 or 2;

R^{A2} may be selected from the group consisting of: hydrogen, R^iR^jN- , heterocyclyl,
 heterocyclyloxy, heterocyclyl- $(NR^a)-$; wherein said heterocyclyl may optionally be substituted

- 12 -

by one or more substituents selected from R^g and wherein if said heterocyclyl contains a $-NH$ moiety that nitrogen may optionally be substituted by one or more groups R^h ; or

R^{A2} may be selected from the group consisting of: C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{3-6} alkenyloxy, C_{3-6} alkynyloxy, C_{3-6} cycloalkoxy, C_{1-6} alkyl- $S(O)_w$ - (wherein w is 0, 1 or 2), C_{1-6} alkyl- $N(R^a)$ -, C_{1-6} alkyl- $N(R^a)$ -carbonyl-, C_{1-6} alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkyl- $N(R^a)$ -carbonyl- $N(R^a)$ -, C_{1-6} alkyl- $N(R^a)$ - SO_2 -, C_{1-6} alkyl- SO_2 - $N(R^a)$ -, C_{1-6} alkoxycarbonyl- $N(R^a)$ -, C_{1-6} alkylcarbonyl- $N(R^a)$ - C_{1-6} alkyl-, C_{1-6} alkyl- $N(R^a)$ -carbonyl- C_{1-6} alkyl-, C_{1-6} alkoxy- C_{1-6} alkyl-; wherein C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{3-6} alkenyloxy, C_{3-6} alkynyloxy, C_{3-6} cycloalkoxy, C_{1-6} alkyl- $S(O)_w$ -, C_{1-6} alkyl- $N(R^a)$ -, C_{1-6} alkyl- $N(R^a)$ -carbonyl-, C_{1-6} alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkyl- $N(R^a)$ -carbonyl- $N(R^a)$ -, C_{1-6} alkyl- $N(R^a)$ - SO_2 -, C_{1-6} alkyl- SO_2 - $N(R^a)$ -, C_{1-6} alkoxycarbonyl- $N(R^a)$ -, C_{1-6} alkylcarbonyl- $N(R^a)$ - C_{1-6} alkyl-, C_{1-6} alkyl- $N(R^a)$ -carbonyl- C_{1-6} alkyl-, C_{1-6} alkoxy- C_{1-6} alkyl may optionally be substituted by R^p , phenyl, phenoxy, heteroaryl, heteroaryloxy, heteroaryl- (NR^a) -, heterocyclyl, heterocyclyloxy or heterocyclyl- $N(R^a)$ -, and wherein said heteroaryl or phenyl may optionally be substituted with one or more substituents selected from R^f ; and wherein said heterocyclyl may optionally be substituted by one or more substituents selected from R^g ; and wherein if said heterocyclyl contains a $-NH$ moiety that nitrogen may optionally be substituted by one or more groups R^h ;

R^{D1} and R^{D2} may be each independently selected from the group consisting of: hydrogen, fluorine, hydroxyl, C_{1-2} alkyl or C_{1-2} alkoxy; wherein the C_{1-2} alkyl and C_{1-2} alkoxy may optionally be substituted by one or more fluorine atoms or a group selected from cyano or hydroxyl.

R^{D3} and R^{D4} may be each independently selected from the group consisting of: hydrogen, fluorine, hydroxyl, cyano, C_{1-3} alkyl or C_{1-3} alkoxy; wherein the C_{1-3} alkyl and C_{1-3} alkoxy may optionally be substituted by one or more fluorine atoms or a group selected from cyano, hydroxyl or $N(R^aR^b)$;

R^{D5} and R^{D6} may be each independently selected from the group consisting of: hydrogen, fluorine, hydroxyl, cyano, C_{1-2} alkyl or C_{1-2} alkoxy; wherein the C_{1-2} alkyl and C_{1-2} alkoxy may optionally be substituted by one or more fluorine atoms or a group selected from cyano, hydroxyl or $N(R^aR^b)$;

R^{N1} may be selected from the group consisting of: hydrogen or C_{1-2} alkyl;

- 13 -

R^{N2} may be selected from the group consisting of: hydrogen, C_{1-3} alkyl or C_{1-2} alkylcarbonyl; wherein the C_{1-3} alkyl and C_{1-2} alkylcarbonyl may optionally be substituted by one or more fluorine atoms or a group selected from cyano, hydroxyl or $N(R^aR^b)$;

R^{N3} may be selected from the group consisting of: hydrogen, C_{1-3} alkyl or C_{1-2} alkylcarbonyl; wherein the C_{1-3} alkyl and C_{1-2} alkylcarbonyl may optionally be substituted by one or more fluorine atoms or a group selected from cyano, hydroxyl or $N(R^aR^b)$;

R^a and R^b may be independently selected, for each occurrence, from the group consisting of: hydrogen and C_{1-3} alkyl; wherein C_{1-3} alkyl may optionally be substituted by one or more substituents selected from: fluorine, cyano, oxo and hydroxyl;

or R^a and R^b , together with the nitrogen to which they are attached, may form a 4-6 membered heterocyclic ring which may have an additional heteroatom selected from O, S, or N; and wherein the 4-6 membered heterocyclic ring may optionally be substituted by one or more substituents selected from the group consisting of: fluorine, cyano, oxo or hydroxyl;

R^f may be independently selected, for each occurrence, from the group consisting of: R^P , hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl-S(O)_w- (wherein w is 0, 1 or 2), C_{1-6} alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkoxycarbonyl- $N(R^a)$ -, and wherein C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl-S(O)_w-, C_{1-6} alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkoxycarbonyl- $N(R^a)$ - may be optionally substituted by one or more substituents selected from R^P ;

R^g may be independently selected for each occurrence from the group consisting of: R^P , hydrogen, oxo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl-S(O)_w- (wherein w is 0, 1 or 2), C_{1-6} alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkoxycarbonyl- $N(R^a)$ -, wherein C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl-S(O)_w-, C_{1-6} alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkoxycarbonyl- $N(R^a)$ - may be optionally substituted by one or more substituents selected from R^P ;

R^h may be independently selected for each occurrence from the group consisting of: hydrogen, C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkyl-S(O)₂-, C_{1-6} alkoxycarbonyl-, R^iR^jN -carbonyl-, R^iR^jN -SO₂-; wherein C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkyl-S(O)₂-, C_{1-6} alkylcarbonyl- may optionally be substituted by one or more substituents selected from R^P ;

- 14 -

R^i and R^j , may be selected independently for each occurrence from the group consisting of: hydrogen, C_{1-4} alkyl and C_{3-6} cycloalkyl; wherein C_{1-4} alkyl and C_{3-6} cycloalkyl may be optionally substituted by one or more substituents selected from fluorine, hydroxyl, cyano, R^aR^bN -, R^aR^bN -carbonyl-, C_{1-3} alkoxy;

5 or R^i and R^j taken together with the nitrogen to which they are attached form a 4-7 membered heterocyclic ring which may have an additional heteroatom selected from O, S, or N, optionally substituted on carbon by one or more substituents selected from the group consisting of: fluorine, hydroxyl, oxo, cyano, C_{1-6} alkyl, C_{1-6} alkoxy, R^aR^bN -, R^aR^bN -SO₂-, R^aR^bN -carbonyl-; and wherein said C_{1-6} alkyl or C_{1-6} alkoxy may optionally be substituted by
 10 fluorine, hydroxyl or cyano; and optionally substituted on nitrogen by one or more substituents selected from the group consisting of: C_{1-6} alkyl, R^aR^bN -carbonyl-; and wherein said C_{1-6} alkyl may be optionally substituted by fluorine, hydroxyl, cyano;

R^P may be independently selected, for each occurrence, from the group consisting of: halogen, hydroxyl, cyano, C_{1-6} alkoxy, R^iR^jN -, R^iR^jN -carbonyl-, R^iR^jN -SO₂-, R^iR^jN -carbonyl-
 15 $N(R^a)$ -;

and pharmaceutically acceptable salts, stereoisomers, esters and prodrugs thereof.

[0047] In some embodiments, X may be selected from the group consisting of:
 $^+C(R^{D1}R^{D2})-C(R^{D5}R^{D6})-^*$, $^+O-C(R^{D5}R^{D6})-^*$, $^+N(R^{N2})-C(R^{D5}R^{D6})-^*$, and
 $^+O-C(R^{D3}R^{D4})-C(R^{D5}R^{D6})-^*$; wherein the $^+$ and * indicate the attachment points of X as
 20 indicated in Formula I. Exemplary X moieties may be selected from the group consisting of:
 $^+CH_2-CH_2-^*$, $^+O-CH_2-^*$, and $^+NH-CH_2-^*$; wherein the $^+$ and * indicate the attachment points of X as indicated in Formula I.

[0048] In one embodiment, R^{D1} , R^{D2} , R^{N1} and R^{N2} that form part of X may be independently selected from the group consisting of hydrogen and methyl. For example, R^{D1} ,
 25 R^{D2} , R^{N1} and R^{N2} that form part of X may be hydrogen.


[0049] In certain embodiments, R^{D3} , R^{D4} , R^{D5} and R^{D6} that form part of X may be independently selected for each occurrence from the group consisting of hydrogen, fluorine, cyano, and C_{1-2} alkyl. For example, R^{D3} , R^{D4} , R^{D5} and R^{D6} that form part of X may be hydrogen.

30 **[0050]** In an embodiment, R^{B1} and R^{B2} may be independently selected from the group consisting of H and methyl.

- 15 -

[0051] In another embodiment, R^{B3} and R^{B4} may be independently selected from the group consisting of H and fluorine.


[0052] Provided herein, for example, are tricyclic compounds represented by formulas Ia, Ib, Ic, Id and Ie:



5

[0053] In certain emodiments, A may be phenyl.

[0054] Also provided herein is a compound represented by Formula II:



10 wherein:

R^{A1} may be selected from the group consisting of: hydrogen, halogen, C_{1-2} alkyl, C_{1-2} alkoxy; wherein C_{1-2} alkyl may optionally be substituted by one or more fluorines;

15 R^{A2} may be selected from the group consisting of: hydrogen, R^iR^jN , heterocyclyl, C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy; wherein said heterocyclyl may optionally be substituted by one or more groups R^g , and wherein if said heterocyclyl contains a -NH moiety, that nitrogen may optionally be substituted by on or more groups R^h ; and wherein said C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} cycloalkyl and C_{1-6} alkoxy may optionally be substituted by one or more groups R^p ;

n may be 1 or 2;

20 R^g may be independently selected for each occurrence from the group consisting of: R^p , hydrogen, oxo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl-S(O)_w-

- 16 -

(wherein w is 0, 1 or 2), C₁₋₆alkylcarbonyl-N(R^a)-, C₁₋₆alkoxycarbonyl-N(R^a)-; wherein C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₆cycloalkyl, C₁₋₆alkoxy, C₁₋₆alkyl-S(O)_w-, C₁₋₆alkylcarbonyl-N(R^a)-, C₁₋₆alkoxycarbonyl-N(R^a)- may be optionally substituted by one or more substituents selected from R^P; R^h is independently selected for each occurrence from the group consisting of: hydrogen, C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, C₃₋₆cycloalkyl, C₁₋₆alkyl-S(O)₂-, C₁₋₆alkoxycarbonyl-, RⁱR^jN-carbonyl-, RⁱR^jN-SO₂-; wherein C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, C₃₋₆cycloalkyl, C₁₋₆alkyl-S(O)₂-, C₁₋₆alkylcarbonyl- may optionally be substituted by one or more substituents selected from R^P;

R^h may be independently selected for each occurrence from the group consisting of: hydrogen, C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, C₃₋₆cycloalkyl, C₁₋₆alkyl-S(O)₂-, C₁₋₆alkoxycarbonyl-, RⁱR^jN-carbonyl-, RⁱR^jN-SO₂-; wherein C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, C₃₋₆cycloalkyl, C₁₋₆alkyl-S(O)₂-, C₁₋₆alkylcarbonyl- may optionally be substituted by one or more substituents selected from R^P;

Rⁱ and R^j, may be selected independently for each occurrence from the group consisting of: hydrogen, C₁₋₄alkyl and C₃₋₆cycloalkyl; wherein C₁₋₄alkyl and C₃₋₆cycloalkyl may be optionally substituted by one or more substituents selected from fluorine, hydroxyl, cyano, R^aR^bN-, R^aR^bN-carbonyl-, C₁₋₃alkoxy;

or Rⁱ and R^j taken together with the nitrogen to which they are attached form a 4-7 membered heterocyclic ring which may have an additional heteroatom selected from O, S, or N, optionally substituted on carbon by one or more substituents selected from the group consisting of: fluorine, hydroxyl, oxo, cyano, C₁₋₆alkyl, C₁₋₆alkoxy, R^aR^bN-, R^aR^bN-SO₂-, R^aR^bN-carbonyl-; and wherein said C₁₋₆alkyl or C₁₋₆alkoxy may optionally be substituted by fluorine, hydroxyl or cyano; and optionally substituted on nitrogen by one or more substituents selected from the group consisting of: C₁₋₆alkyl, R^aR^bN-carbonyl-; and wherein said C₁₋₆alkyl may be optionally substituted by fluorine, hydroxyl, cyano;

R^P may be independently selected, for each occurrence, from the group consisting of: halogen, hydroxyl, cyano, C₁₋₆alkoxy, RⁱR^jN-, RⁱR^jN-carbonyl-, RⁱR^jN-SO₂-, RⁱR^jN-carbonyl-N(R^a)-;

R^a and R^b may be independently selected, for each occurrence, from the group consisting of: hydrogen and C₁₋₃alkyl; wherein C₁₋₃alkyl may optionally be substituted by one or more substituents selected from: fluorine, cyano, oxo and hydroxyl;

- 17 -

or R^a and R^b, together with the nitrogen to which they are attached, may form a 4-6 membered heterocyclic ring which may have an additional heteroatom selected from O, S, or N; and wherein the 4-6 membered heterocyclic ring may optionally be substituted by one or more substituents selected from the group consisting of: fluorine, cyano, oxo or hydroxyl;

5 and pharmaceutically acceptable salts, stereoisomers, esters and prodrugs thereof.

[0055] In certain embodiments, R^{A1} of the tricyclic compound of Formula II may be selected from the group consisting of hydrogen and fluorine.

[0056] Procedures for making compounds described herein are provided below with reference to Schemes 1-2. In the reactions described below, it may be necessary to protect reactive functional groups (such as hydroxyl, amino, thio or carboxyl groups) to avoid their
10 unwanted participation in the reactions. The incorporation of such groups, and the methods required to introduce and remove them are known to those skilled in the art [for example, see Greene, Wuts, Protective Groups in Organic Synthesis, 2nd Ed. (1999)]. The deprotection step may be the final step in the synthesis such that the removal of protecting groups affords
15 compounds of Formula I, as disclosed herein, or as exemplified in, for example, General Formula I, below. Starting materials used in the following schemes can be purchased or prepared by methods described in the chemical literature, or by adaptations thereof, using methods known by those skilled in the art. The order in which the steps are performed can vary depending on the groups introduced and the reagents used, but would be apparent to those
20 skilled in the art.


[0057] The general synthetic strategy used to prepare the tricyclic compounds of General Formula I is depicted in Scheme 1. The tricyclic system may be assembled in a variety of ways, starting from an appropriately substituted and protected phenyl ring 1A. The group G' is a suitably protected carboxylic acid, such as a methyl- or *tert*-butyl carboxylate or is a
25 functional group that may be readily converted into a carboxylic acid, such as a nitrile or aldehyde. The group G is a thioether, sulfoxide or sulfone, or a functional group that may be subsequently converted into a thioether, sulfoxide or sulfone group such as a halide or mesylate. The cyclopropane ring can be directly attached to the substituted phenyl ring to give intermediate 1B, and then the D'-ring can be formed by an intra-molecular reaction to give
30 intermediate 1E. Alternatively, the cyclopropane ring can be attached to the substituted phenyl ring 1A via a linker, X', to give intermediate 1C, and then the D'-ring can be formed by an intra-molecular reaction to give intermediate 1E. Alternatively, the D'-ring can be built up onto

- 18 -

the substituted phenyl ring to give intermediate 1D, and then the cyclopropane ring formed to give intermediate 1E. Modifications to the cyclopropane and D' rings may be necessary to provide the required ring systems and this may be carried out prior to the formation of the tricyclic core or after it. For example, a dibromocyclopropane ring may be formed followed by
 5 removal of the bromines to give the unsubstituted cyclopropane ring.

[0058] Compounds of Formula I can be prepared from intermediate 1E by removal of any protecting groups. Alternatively, further modifications may be made to 1E, such as modifications at G, before the removal of any protecting groups to give compounds of General Formula I. Specific steps in the synthetic process are described in more detail below.

10

SCHEME 1

[0059] In Scheme 1, the carbon-carbon bond between the phenyl ring in 1A and the cyclopropane ring in 1B' in Step (i) may be formed under a range of coupling conditions to
 15 give compounds of structure 1B. The introduction of the cyclopropane ring may require a number of steps and the preparation of a number of intermediates. Protecting groups may also be required. If R¹ is a suitable leaving group (such as a halide or triflate), and R³ is a borane, boronate or boronic acid group, then the carbon-carbon bond may be formed by treatment with
 20 a palladium catalyst (such as palladium chloride dppf or *tris*-(dibenzylideneacetone)-dipalladium, in the presence of a base (such as cesium carbonate) and a suitable reagent (such

- 19 -

as a phosphine, for example tri-*tert*-butylphosphonium tetrafluoroborate) in an appropriate solvent (such as dioxane or water, or mixtures thereof), at a temperature between 50°C and the reflux temperature of the solvent or alternatively by irradiating in the microwave at a temperature up to 180°C. A wide range of appropriate reagents and conditions are known to those skilled in the art to couple organoboranes, boronates and boronic acids to compounds such as 1A. [For example, see Miyaura, Suzuki, Chem. Rev. 1995, 95, 2457; Suzuki, Modern Arene Chemistry 2002, 53-106].

[0060] Alternatively, If R¹ is a suitable group (such as a halide or triflate), and R³ is a trialkyl stannane (such as a tri-*n*-butyl stannane), then the carbon-carbon bond may be formed by treatment with a palladium catalyst (such as palladium chloride dppf), in an appropriate solvent (such as dimethoxyethane or tetrahydrofuran) at a temperature between 50°C and the reflux temperature of the solvent or alternatively by irradiation in the microwave at a temperature up to 180°C to give 1B. A wide range of appropriate reagents and conditions are known to those skilled in the art to couple stannanes to aryl halides such as 1A. [For example, see Smith, March, March's Advanced Organic Chemistry, 5th Edition, Wiley: New York, 2001, pp.931-932; De Souza, Current Organic Synthesis 2006, 3(3), 313-326.].

[0061] Alternatively compounds of structure 1A, where R¹ is a suitable group (such as a halide or triflate), can be treated with, for example, a diboronate (such as bis-pinacolatodiboron) in the presence of a palladium catalyst (such as palladium chloride dppf) and a base (such as potassium acetate or diisopropylamine) in an appropriate solvent (such as a mixture of dioxane and water) at a temperature between 50°C and the reflux temperature of the solvent or alternatively by irradiation in the microwave at a temperature up to 180°C to give a compound of structure 1A, where R¹ is a boronate. A wide range of appropriate reagents and conditions are known to those skilled in the art to convert an aryl halide (or aryl triflate) to an arylboronate (or an arylborane) [for example, see Marshall, Chemtracts 2000, 13(4), 219-222]. The arylboronate (or arylborane) thus formed, can then be treated with compounds of structure 1B' (where R³ is a halogen or triflate) in the presence of suitable reagents such as a phosphine (for example tri-*tert*-butyl-phosphonium tetrafluoroborate), a base (such as cesium carbonate) and a catalyst (such as *tris*-(dibenzylideneacetone)-dipalladium) in an appropriate solvent (such as a mixture of water and dioxane) at a temperature between 50°C and the reflux temperature of the solvent or alternatively by irradiation in the microwave at a temperature up to 180°C to afford compounds of structure 1B.

- 20 -

[0062] In Scheme 1, Step (iv), the groups R^2 and R^4 of compound 1B can be coupled together to give the group X' , which forms the D' -ring. R^2 or R^4 may have been masked by protecting groups during Step (i), and may require deprotection before the group X' can be formed. Alternatively, R^2 or R^4 may require chemical modification before the group X' can be formed. For example if R^2 or R^4 is a nitro group, that group may be reduced, for example using hydrogen in the presence of a suitable catalyst (such as palladium on a solid support, such as carbon); or by treatment with an inorganic reducing agent (such as tin (II) chloride in DMF) to give an amino group. For example, if R^2 or R^4 is a hydroxyalkyl group, that group may be treated with an oxidising agent (such as Jones reagent or manganese dioxide) to give an aldehyde; or with a different oxidising agent (such as potassium permanganate) to give a carboxylic acid. For example, if R^2 or R^4 is an aldehyde, that group may be treated with an oxidising agent (such as potassium permanganate) to give a carboxylic acid or with a reducing agent (such as sodium borohydride) to give an alcohol. For example, if R^2 or R^4 is a ketone, that group may be treated with a reducing agent (such as sodium borohydride) to give a secondary alcohol. For example, if R^2 or R^4 is a carboxylic acid or ester, that group may be treated with a reducing agent (such as lithium aluminium hydride) to give an alcohol. For example, if R^2 or R^4 is an alkene group, that group may be treated with a borane (such as 9-borobicyclononane) and converted to a primary or secondary alcohol.

[0063] Formation of the linker X' may be carried out in a number of ways known to those skilled in the art. For example, if the group R^2 is a hydroxyl and the group R^4 is a substituted alkylalcohol then 1B can be treated with a dehydrating agent (such as diisopropyl azodicarboxylate) in the presence of a phosphine, (such as triphenylphosphine) to give 1E, where X' is an ether. Alternatively, if one of the two groups R^2 or R^4 is a hydroxyl and the other group is an alkyl group substituted with a leaving group (such as a halogen, tosylate or triflate) 1B can be treated with a base (such as diisopropylethylamine, potassium carbonate or sodium hydride) to form 1E, where X' is an ether.

[0064] Alternatively, if the group R^4 is a carboxylic acid and R^2 is an alkyl halide or sulfonate, then 1B can be treated with a base such as diisopropyl-ethylamine, potassium carbonate or sodium hydride to form 1E, where X' is an ester.

[0065] Alternatively, if the group R^2 is a hydroxyl, or substituted alkylalcohol and R^4 is a carboxylic acid or ester or an alkyl group substituted by a carboxylic acid or ester, then 1B

- 21 -

can be treated with an acid (such as hydrochloric acid) or dehydrating agent (such as dicyclohexylcarbodiimide or acetic anhydride) to form 1E, where X' is an ester.

[0066] Alternatively, if the group R^2 on 1B is a hydroxyl or substituted alkylalcohol and R^4 is a carboxylic acid or an alkyl group substituted by a carboxylic acid, then the carboxylic acid can first be converted to a mixed anhydride (for example by treatment with 2,4,6-trichlorobenzoyl chloride) or to an activated ester (for example by treatment with HATU in the presence of a base such as diisopropylethylamine or pyridine), and the resulting mixed anhydride or activated ester can be further treated with a base (such as diisopropylethylamine, pyridine or potassium carbonate) to form 1E, where X' is an ester.

10 [0067] Alternatively, if the group R^2 on 1B is an amine or a substituted alkylamine and the group R^4 is a carboxylic acid or an alkyl group substituted by a carboxylic acid, the carboxylic acid can be converted to an activated ester (for example by treatment with HATU and a base such as diisopropylethylamine or pyridine or TBTU in the presence of N-methylmorpholine), and the resulting activated ester can be further treated with a base to form
15 1E where X' is an amide.

[0068] Alternatively, if the group R^2 on 1B is an amine, or a substituted alkylamine and the group R^4 is a carboxylic acid or an alkyl group substituted by a carboxylic acid, then 1B can then be treated with a dehydrating agent (such as such as diisopropylcarbodiimide) to form 1E, where X' is an amide.

20 [0069] Alternatively, if one of the two groups R^2 or R^4 is an amine, or substituted alkylamine and the other group is an alkyl group substituted with a leaving group (such as a halogen, tosylate or triflate) then 1B can be treated with a base (such as diisopropylethylamine, pyridine or potassium carbonate) to form 1E, where X' is a substituted amine.

[0070] In Scheme 1, Step (ii), compounds of the structure 1A can be reacted with 1C' to form the linker X' and give compounds of the structure 1C. The formation of the linker X' in compounds with the structure 1C may require a number of steps and the preparation of a number of intermediates, and the use of protecting groups may also be required. The linker X' may be prepared using one of the methods described for Step (iv) in the formation of 1E.

[0071] In Scheme 1, Step (v), compounds of structure 1E may be prepared from
30 compounds of structure 1C by reaction of the groups R^1 and R^5 under a range of conditions to

- 22 -

form a carbon-carbon bond. The carbon-carbon bond may be formed using one of the methods described for Step (i) in the formation of 1B.

[0072] In Scheme 1, Step (iii), compounds of structure 1A may be reacted under a range of conditions with intermediates of the type 1D' to give compounds of structure 1D where D' is a six- or seven-membered fused carbocyclic or heterocyclic ring. The groups R¹ and R⁷ may be reacted together to form a carbon-carbon bond, and the groups R² and R⁸ may be reacted together to form the group X'. Methods to form bicyclic compounds of structure 1D from substituted phenyl rings of structure 1A are well known to those skilled in the art (see Comprehensive Heterocyclic Chemistry Ed.: Katritzky, Ramsden, Scriven, and Taylor, Elsevier, 2008).

[0073] For example, a compound of structure 1A, where R² is a hydroxyl and R¹ is hydrogen, can be treated with a propargyl halide or tosylate or a substituted propargyl halide or tosylate in the presence of a base (such as potassium carbonate or cesium carbonate) in a solvent (such as acetone) at a temperature between room temperature and the reflux temperature of the solvent to give a compound of type 1A in which R² is a propargyloxy or substituted propargyloxy group. This intermediate may be heated to ~200°C or treated with an appropriate catalyst (such as a gold catalyst, for example triphenylphosphine gold triflamide) in an appropriate solvent (such as toluene) at a temperature between 80°C and the reflux temperature of the solvent to give a compound of structure 1D wherein X' is -OCH₂-.

[0074] In Scheme 1, Step (vi), compounds of structure 1D, may be treated with diethyl zinc and di-iodomethane optionally in the presence of additional reagents (such as trifluoroacetic acid or zinc iodide) in a solvent (such as 1,2-dichloroethane) at a temperature between -78°C and room temperature to give compounds of structure 1E. Alternatively, compounds of structure 1D, may be treated with trimethyl sulfoxonium iodide or trimethyl sulphonium iodide in the presence of a base (such as sodium hydride) in a solvent (such as dimethyl sulfoxide) at a temperature between room temperature and 100°C. Alternatively, compounds of structure 1D, may be treated with diazomethane in the presence of a catalyst (such as palladium acetate) in a solvent such as diethyl ether at a temperature between 0°C and room temperature.

[0075] Alternatively, compounds of structure 1E may be prepared from compounds of structure 1D in a two step procedure. For example, compounds of structure 1D, may be treated with trimethylsilyl diazomethane in the presence of a catalyst (such as palladium acetate) in a

- 23 -

solvent (such as diethyl ether) at a temperature between 0°C and room temperature to give an intermediate of structure 1E in which R¹³ is H and R¹⁴ is a trimethyl silyl group. This intermediate may be converted to compounds of structure 1E in which R¹³ and R¹⁴ are both H, by treatment with a source of fluoride, (for example tetrabutyl ammonium fluoride) in a solvent (such as tetrahydrofuran) at a temperature between 0°C and room temperature.

[0076] Alternatively, compounds of structure 1D may be treated with a haloform (such as chloroform or bromoform) in the presence of a base such as sodium hydroxide or potassium t-butoxide in the presence of a catalyst (such as triethylbenzylammonium chloride or tetrabutyl ammonium bromide) in a mixture of water and a halogenated solvent (such as dichloromethane or dichloroethane) or alternatively in excess of the haloform, at a temperature between room temperature and 80°C to give an intermediate compound of structure 1E in which R¹³ and R¹⁴ are both chlorine or bromine. This intermediate may be converted to compounds of structure 1E in which R¹³ and R¹⁴ are both H by treatment with a base (such as sodium methoxide or sodium t-butoxide) in a solvent (such as tetrahydrofuran) at a temperature between room temperature and the reflux temperature of the solvent. Alternatively, the intermediate of structure 1E in which R¹³ and R¹⁴ are both chlorine or bromine may be converted to compounds of structure 1E in which R¹³ and R¹⁴ are both H by reduction; for example, by treatment with lithium aluminium hydride in a solvent (such as tetrahydrofuran) at a temperature between 0°C and the reflux temperature of the solvent or alternatively, for example by hydrogenation using a catalyst (such as palladium on carbon) in a solvent (such as methanol or ethanol), optionally in the presence of a base (such as triethylamine) or alternatively by treatment with a metal reducing agent such as zinc, in the presence of a catalyst such as ammonium chloride. Alternatively, the intermediate of structure 1E in which R¹³ and R¹⁴ are both chlorine or bromine may be converted to compounds of structure 1E, in which R¹³ and R¹⁴ are both H, under radical conditions; for example, by treatment with a tin hydride (such as tributyl tin hydride) optionally in the presence of a radical initiator (such as azo-bis-isobutyronitrile) in a solvent (such as toluene) at a temperature between room temperature and the reflux temperature of the solvent.

[0077] Alternatively, compounds of structure 1E, in which R¹³ and R¹⁴ are both fluorine may be prepared by the treatment of compounds of structure 1D by treatment with a source of difluorocarbene. For example, by treatment with sodium difluorochloroacetate or

- 24 -

difluorobromoacetate in an ether solvent such as diethyleneglycol dimethyl ether while heating at a temperature between 130°C and 180°C.

[0078] In Scheme 1, Step (vii), compounds of general structure 1E may be converted to compounds of General Formula I by the conversion of the group G' to a carboxylic acid. If the group G' is a carboxylic ester (such as a methyl, *tert*-butyl or benzyl ester) then a variety of reagents and conditions can be used to convert 1E into a compound of the General Formula I. For example, if G' is a methyl, ethyl or benzyl ester, it may be converted to a carboxylic acid by treatment with an inorganic base (such as lithium hydroxide or sodium hydroxide) in a solvent (such as methanol, dioxane or water, or mixtures thereof) at a temperature between room temperature and the reflux temperature of the solvent, or alternatively by microwave irradiation at 120-180°C for 10 minutes to 1 hour. Alternatively if G' is a benzyl ester it may be converted to a carboxylic acid by hydrogenation in the presence of a catalyst (such as palladium on a solid support such as carbon) in a solvent (such as dioxane or ethyl acetate). Alternatively if G' is a *tert*-butyl ester, it may be converted to a carboxylic acid by treatment with an acid (such as trifluoroacetic acid or hydrogen chloride) in a solvent (such as dichloromethane or dioxane).


[0079] Alternatively, if the group G' is a nitrile, it may be converted into a carboxylic acid by treatment with aqueous acid (such as a mineral acid, for example hydrochloric acid) under appropriate conditions (such as heating, for example to reflux); or by treatment with aqueous base (such as an aqueous hydroxide, for example aqueous sodium hydroxide) under appropriate conditions (such as heating, for example to reflux).

[0080] Alternatively, if the group G' is an aldehyde or a hydroxymethyl moiety then it may be converted into a carboxylic acid by treatment with a suitable oxidising reagent (such as potassium permanganate or chromic acid).

[0081] The general synthetic strategy to modify the group G is depicted in Scheme 2. The G group may be introduced and/or modified either before, during or after the assembly of the tricyclic ring system. It may be that the group G is protected during the sequence to form the tricyclic ring system and these protecting groups will need to be removed or modified prior to conversion to the sulfone moiety. For example, the group G may be a methoxy group during the earlier sequence. Conversion of this methoxy group to an appropriate leaving group such as a halogen may be achieved by treatment with acetyl chloride in the presence of, for example, zinc chloride in a solvent (such as diethyl ether) at a temperature between room temperature

and the reflux temperature of the solvent. Specific steps used to assemble sulfone are described in more detail below.

SCHEME 2



5 [0082] In Scheme 2, the asterisks denote either the presence of the groups R¹ and R² (as shown in Scheme 1) or the presence of the tricyclic ring system, or intermediates towards the ring system (as shown in Scheme 1).

[0083] In Scheme 2, Step (i), if G is a suitable leaving group, such as a halide (chloride or bromide) or triflate, it may be reacted with an appropriately substituted thiol in the presence of a base (such as potassium carbonate) in a solvent (such as acetone or THF) at a temperature between room temperature and the reflux temperature of the solvent. A wide range of reagents and conditions are known to one skilled in the art for the conversion of compounds of structure 2A in which G is a leaving group to compounds of structure 2B [for example, see Peach in Patai *The Chemistry of the Thiol Group*, pt2; Wiley: New York, 1974, pp.721].

15 [0084] In Scheme 2, Step (ii) compounds of structure 2B may be converted to compounds of structure 2C by oxidation. The oxidation may be carried out by treatment of compounds of structure 2B with hydrogen peroxide in a solvent (such as acetic acid) at a temperature between room temperature and the reflux temperature of the solvent. Alternative oxidizing agents may be used to convert compounds of structure 2B to compounds of structure 2C such as 3-chloroperbenzoic acid in a chlorinated solvent (such as dichloromethane) at a temperature between 0°C and the reflux temperature of the solvent. A wide range of reagents and conditions are known to one skilled in the art for the oxidation of thiols to sulfones. [for example, see Drabowicz, Kielbasinski, Mikolajczyk in Patai, Rappoport, Stirling *The Chemistry of Sulphones and Sulphoxides*; Wiley: New York, 1988, pp.233-378, pp235-255.

20 Madesclaire *Tetrahedron* **1986**, 42, 549-5495. Oae *The Organic Chemistry of Sulfur*; Plenum: New York 1977, pp.385-390. Smith, March, *March's Advanced Organic Chemistry*, 5th Edition, Wiley: New York, 2001, pp.1541-1542].

- 26 -

[0085] Alternatively in Scheme 2, Step (iii), compounds of structure 2A may be converted directly to compounds of structure 2C by treatment with an appropriately substituted sulfinate in the presence of a base (such as sodium bicarbonate, potassium bicarbonate or di-isopropylethylamine) in a solvent (such as a mixture of dimethyl acetamide and water) at a temperature between room temperature and the reflux temperature of the solvent.

[0086] Compounds of any of Formula I or, for example, General Formula I as depicted above, or any of the intermediates described in the schemes above, can be further derivatised by using one or more standard synthetic methods known to those skilled in the art. Such methods can involve substitution, oxidation or reduction reactions. These methods can also be used to obtain or modify compounds of General Formula I or any preceding intermediates by modifying, introducing or removing appropriate functional groups. Particular substitution approaches include alkylation, arylation, heteroarylation, acylation, thioacylation, halogenation, sulfonylation, nitration, formylation, hydrolysis and coupling procedures. These procedures can be used to introduce a functional group onto the parent molecule (such as the nitration or sulfonylation of aromatic rings) or to couple two molecules together (for example to couple an amine to a carboxylic acid to afford an amide; or to form a carbon-carbon bond between two heterocycles). For example, alcohol or phenol groups can be converted to ether groups by coupling a phenol with an alcohol in a solvent (such as tetrahydrofuran) in the presence of a phosphine (such as triphenylphosphine) and a dehydrating agent (such as diethyl, diisopropyl or dimethyl azodicarboxylate). Alternatively, ether groups can be prepared by deprotonation of an alcohol, using a suitable base (such as sodium hydride) followed by the addition of an alkylating agent (such as an alkyl halide or an alkyl sulfonate).

[0087] In another example, a primary or secondary amine can be alkylated using a reductive alkylation procedure. For example, the amine can be treated with an aldehyde and a borohydride (such as sodium triacetoxyborohydride, or sodium cyanoborohydride) in a solvent (such as a halogenated hydrocarbon, for example dichloromethane, or an alcohol, for example ethanol) and, where necessary, in the presence of an acid (such as acetic acid).

[0088] In another example, hydroxy groups (including phenolic OH groups) can be converted into leaving groups, such as halogen atoms or sulfonyloxy groups (such as alkylsulfonyloxy, for example trifluoromethanesulfonyloxy, or aryl sulphonyloxy, for example p-toluenesulfonyloxy) using conditions known to those skilled in the art. For example, an aliphatic alcohol can be reacted with thionyl chloride in a halogenated hydrocarbon (such as

- 27 -

dichloromethane) to afford the corresponding alkyl chloride. A base (such as triethylamine) can also be used in the reaction.

[0089] In another example, ester groups can be converted to the corresponding carboxylic acid by acid- or base-catalysed hydrolysis depending on the nature of the ester group. Acid catalysed hydrolysis can be achieved by treatment with an organic or inorganic acid (such as trifluoroacetic acid in an aqueous solvent, or a mineral acid such as hydrochloric acid in a solvent such as dioxane). Base catalysed hydrolysis can be achieved by treatment with an alkali metal hydroxide (such as lithium hydroxide in an aqueous alcohol, for example methanol).

10 [0090] In another example, aromatic halogen substituents in the compounds may be subjected to halogen-metal exchange by treatment with a base (such as a lithium base, for example n-butyl or t-butyl lithium) optionally at a low temperature (such as -78°C) in a solvent (such as tetrahydrofuran) and the mixture may then be quenched with an electrophile to introduce a desired substituent. Thus, for example, a formyl group can be introduced by using dimethylformamide as the electrophile. Aromatic halogen substituents can also be subjected to palladium catalysed reactions to introduce groups such as carboxylic acids, esters, cyano or amino substituents.

[0091] In another example, an aryl, or heteroaryl ring substituted with an appropriate leaving group (such as a halogen or sulfonyl ester, for example a triflate) can undergo a palladium catalysed coupling reaction with a wide variety of substrates to form a carbon-carbon bond. For example, a Heck reaction can be used to couple such a ring system to an alkene (which may, or may not, be further substituted) by treatment with an organopalladium complex (such as tetrakis(triphenylphosphine)palladium(0), palladium (II) acetate or palladium (II) chloride) in the presence of a ligand (such as a phosphine, for example triphenylphosphine) in the presence of a base (such as potassium carbonate or a tertiary amine, for example, triethylamine), in an appropriate solvent (such as tetrahydrofuran or DMF), under appropriate conditions (such as heating to, for example, 50-120°C). In another example, a Sonogashira reaction can be used to couple such a ring system to an alkyne (which may, or may not be further substituted) by treatment with a palladium complex (such as tetrakis(triphenylphosphine)palladium(0)) and a halide salt of copper (I) (such as copper (I) iodide), in the presence of a base (such as a potassium carbonate or a tertiary amine, for example, triethylamine), in an appropriate solvent (such as tetrahydrofuran or

- 28 -

dimethylformamide), under appropriate conditions (such as heating to, for example, 50-120°C). In another example, a Stille reaction can be used to couple such a ring system to an alkene, by treatment with an organotin compound (such as an alkynyltin or alkenyltin reagent, for example an alkenyltributylstannane) in the presence of a palladium complex (such as

5 tetrakis(triphenylphosphine)palladium(0)), with, or without the presence of a salt (such as a copper (I) halide), in an appropriate solvent (such as dioxane or dimethylformamide), under appropriate conditions (such as heating to, for example, 50-120°C).

[0092] Particular oxidation approaches include dehydrogenations and aromatisation, decarboxylation and the addition of oxygen to certain functional groups. For example,

10 aldehyde groups can be prepared by oxidation of the corresponding alcohol using conditions well known to those skilled in the art. For example, an alcohol can be treated with an oxidising agent (such as Dess-Martin periodinane) in a solvent (such as a halogenated hydrocarbon, for example dichloromethane). Alternative oxidising conditions can be used, such as treatment with oxalyl chloride and an activating amount of dimethylsulfoxide and subsequent quenching

15 by the addition of an amine (such as triethylamine). Such a reaction can be carried out in an appropriate solvent (such as a halogenated hydrocarbon, for example dichloromethane) and under appropriate conditions (such as cooling below room temperature, for example to -78°C followed by warming to room temperature). In another example, sulfur atoms can be oxidised to the corresponding sulfoxide or sulfone using an oxidising agent (such as a peroxy acid, for

20 example 3-chloroperoxybenzoic acid) in an inert solvent (such as a halogenated hydrocarbon, for example dichloromethane) at around ambient temperature.

[0093] Particular reduction approaches include the removal of oxygen atoms from particular functional groups or saturation (or partial saturation) of unsaturated compounds including aromatic or heteroaromatic rings. For example, primary alcohols can be generated

25 from the corresponding ester or aldehyde by reduction, using a metal hydride (such as lithium aluminium hydride or sodium borohydride in a solvent such as methanol). Alternatively, CH₂OH groups can be generated from the corresponding carboxylic acid by reduction, using a metal hydride (such as lithium aluminium hydride in a solvent such as tetrahydrofuran). In another example, a nitro group may be reduced to an amine by catalytic hydrogenation in the

30 presence of a metal catalyst (such as palladium on a solid support such as carbon) in a solvent (such as an ether, for example tetrahydrofuran, or an alcohol, such as methanol), or by chemical reduction using a metal (such as zinc, tin or iron) in the presence of an acid (such as acetic acid

- 29 -

or hydrochloric acid). In a further example an amine can be obtained by reduction of a nitrile, for example by catalytic hydrogenation in the presence of a metal catalyst (such as palladium on a solid support such as carbon), or Raney nickel in a solvent (such as tetrahydrofuran) and under suitable conditions (such as cooling to below room temperature, for example to -78°C, or heating, for example to reflux).

[0094] Salts of compounds of General Formula I can be prepared by the reaction of a compound of General Formula I with an appropriate acid or base in a suitable solvent, or mixture of solvents (such as an ether, for example, diethyl ether, or an alcohol, for example ethanol, or an aqueous solvent) using conventional procedures. Salts of compound of General Formula I can be exchanged for other salts by treatment using conventional ion-exchange chromatography procedures.

[0095] Where it is desired to obtain a particular enantiomer of a compound of General Formula I, this may be produced from a corresponding mixture of enantiomers by employing any suitable conventional procedure for resolving enantiomers. For example, diastereomeric derivatives (such as salts) can be produced by reaction of a mixture of enantiomers of a compound of General Formula I (such a racemate) and an appropriate chiral compound (such as a chiral base). The diastereomers can then be separated by any conventional means such as crystallisation, and the desired enantiomer recovered (such as by treatment with an acid in the instance where the diastereomer is a salt). Alternatively, a racemic mixture of esters can be resolved by kinetic hydrolysis using a variety of biocatalysts (for example, see Patel Stereoselective Biocatalysts, Marcel Decker; New York 2000).

[0096] In another resolution process a racemate of compounds of General Formula I can be separated using chiral High Performance Liquid Chromatography. Alternatively, a particular enantiomer can be obtained by using an appropriate chiral intermediate in one of the processes described above. Chromatography, recrystallisation and other conventional separation procedures may also be used with intermediates or final products where it is desired to obtain a particular geometric isomer of the invention.

II. Methods

[0097] Another aspect of the invention provides methods of modulating the activity of MetAP2. Such methods comprise exposing said receptor to a compound described herein. In some embodiments, the compound utilized by one or more of the foregoing methods is one of

- 30 -

the generic, subgeneric, or specific compounds described herein, such as a compound of Formula I, Ia, Ib, Ic, Id, or II. The ability of compounds described herein to modulate or inhibit MetAP2 can be evaluated by procedures known in the art and/or described herein. Another aspect of the invention provides methods of treating a disease associated with expression or activity of MetAP2 in a patient. For example, a contemplated method includes administering a disclosed compound in an amount sufficient to establish inhibition of intracellular MetAP2 effective to increase thioredoxin production in the patient and to induce multi organ stimulation of anti-obesity processes in the subject, for example, by administering a disclosed compound in an amount insufficient to reduce angiogenesis in the patient.

10 **[0098]** In certain embodiments, the invention provides a method of treating and or ameliorating obesity in a patient by administering an effective amount of a disclosed compound. Also provided herein are methods for inducing weight loss in a patient in need thereof. Contemplated patients include not only humans, but other animals such as companion animals (e.g., dogs, cats).

15 **[0099]** Other contemplated methods of treatment include method of treating or ameliorating an obesity-related condition or co-morbidity, by administering a compound disclosed herein to a subject. For example, contemplated herein are methods for treating type 2 diabetes in a patient in need thereof.

[00100] Exemplary co-morbidities include cardiac disorders, endocrine disorders, 20 respiratory disorders, hepatic disorders, skeletal disorders, psychiatric disorders, metabolic disorders, and reproductive disorders.

[00101] Exemplary cardiac disorders include hypertension, dyslipidemia, ischemic heart disease, cardiomyopathy, cardiac infarction, stroke, venous thromboembolic disease and pulmonary hypertension. Exemplary endocrine disorders include type 2 diabetes and latent 25 autoimmune diabetes in adults. Exemplary respiratory disorders include obesity-hypoventilation syndrome, asthma, and obstructive sleep apnea. An exemplary hepatic disorder is nonalcoholic fatty liver disease. Exemplary skeletal disorders include back pain and osteoarthritis of weight-bearing joints. Exemplary metabolic disorders include Prader-Willi Syndrome and polycystic ovary syndrome. Exemplary reproductive disorders include sexual 30 dysfunction, erectile dysfunction, infertility, obstetric complications, and fetal abnormalities. Exemplary psychiatric disorders include weight-associated depression and anxiety.

[00102] In particular, in certain embodiments, the invention provides a method of treating the above medical indications comprising administering to a subject in need thereof a therapeutically effective amount of a compound described herein, such as a compound of Formula I, Ia, Ib, Ic, Id, or II.

5 [00103] Obesity or reference to “overweight” refers to an excess of fat in proportion to lean body mass. Excess fat accumulation is associated with increase in size (hypertrophy) as well as number (hyperplasia) of adipose tissue cells. Obesity is variously measured in terms of absolute weight, weight:height ratio, distribution of subcutaneous fat, and societal and esthetic norms. A common measure of body fat is Body Mass Index (BMI). The BMI refers to the
10 ratio of body weight (expressed in kilograms) to the square of height (expressed in meters). Body mass index may be accurately calculated using either of the formulas: $\text{weight}(\text{kg}) / \text{height}^2(\text{m}^2)$ (SI) or $703 \times \text{weight}(\text{lb}) / \text{height}^2(\text{in}^2)$ (US).

[00104] In accordance with the U.S. Centers for Disease Control and Prevention (CDC), an overweight adult has a BMI of 25 kg/m^2 to 29.9 kg/m^2 , and an obese adult has a BMI of 30
15 kg/m^2 or greater. A BMI of 40 kg/m^2 or greater is indicative of morbid obesity or extreme obesity. Obesity can also refer to patients with a waist circumference of about 102 cm for males and about 88 cm for females. For children, the definitions of overweight and obese take into account age and gender effects on body fat. Patients with differing genetic background may be considered “obese” at a level differing from the general guidelines, above.

20 [00105] The compounds of the present invention also are useful for reducing the risk of secondary outcomes of obesity, such as reducing the risk of left ventricular hypertrophy. Methods for treating patients at risk of obesity, such as those patients who are overweight, but not obese, e.g. with a BMI of between about 25 and 30 kg/m^2 , are also contemplated. In certain embodiments, a patient is a human.

25 [00106] BMI does not account for the fact that excess adipose can occur selectively in different parts of the body, and development of adipose tissue can be more dangerous to health in some parts of the body rather than in other parts of the body. For example, “central obesity”, typically associated with an “apple-shaped” body, results from excess adiposity especially in the abdominal region, including belly fat and visceral fat, and carries higher risk of co-
30 morbidity than “peripheral obesity”, which is typically associated with a “pear-shaped” body resulting from excess adiposity especially on the hips. Measurement of waist/hip circumference ratio (WHR) can be used as an indicator of central obesity. A minimum WHR indicative of

- 32 -

central obesity has been variously set, and a centrally obese adult typically has a WHR of about 0.85 or greater if female and about 0.9 or greater if male.

[00107] Methods of determining whether a subject is overweight or obese that account for the ratio of excess adipose tissue to lean body mass involve obtaining a body composition of the subject. Body composition can be obtained by measuring the thickness of subcutaneous fat in multiple places on the body, such as the abdominal area, the subscapular region, arms, buttocks and thighs. These measurements are then used to estimate total body fat with a margin of error of approximately four percentage points. Another method is bioelectrical impedance analysis (BIA), which uses the resistance of electrical flow through the body to estimate body fat. Another method is using a large tank of water to measure body buoyancy. Increased body fat will result in greater buoyancy, while greater muscle mass will result in a tendency to sink.

[00108] In another aspect, the invention provides methods for treating an overweight or obese subject involving determining a level of at least one biomarker related to being overweight or obese in the subject, and administering an effective amount of a disclosed compound to achieve a target level in the subject. Exemplary biomarkers include body weight, Body Mass Index (BMI), Waist/Hip ratio WHR, plasma adipokines, and a combination of two or more thereof.

[00109] In certain embodiments, the compound utilized by one or more of the foregoing methods is one of the generic, subgeneric, or specific compounds described herein, such as a compound of Formula I, Ia, Ib, Ic, Id, or II.

[00110] The compounds of the invention may be administered to patients (animals and humans) in need of such treatment in dosages that will provide optimal pharmaceutical efficacy. It will be appreciated that the dose required for use in any particular application will vary from patient to patient, not only with the particular compound or composition selected, but also with the route of administration, the nature of the condition being treated, the age and condition of the patient, concurrent medication or special diets then being followed by the patient, and other factors which those skilled in the art will recognize, with the appropriate dosage ultimately being at the discretion of the attendant physician. For treating clinical conditions and diseases noted above, a compound of this invention may be administered orally, subcutaneously, topically, parenterally, by inhalation spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants

- 33 -

and vehicles. Parenteral administration may include subcutaneous injections, intravenous or intramuscular injections or infusion techniques.

[00111] Treatment can be continued for as long or as short a period as desired. The compositions may be administered on a regimen of, for example, one to four or more times per day. A suitable treatment period can be, for example, at least about one week, at least about two weeks, at least about one month, at least about six months, at least about 1 year, or indefinitely. A treatment period can terminate when a desired result, for example a weight loss target, is achieved. A treatment regimen can include a corrective phase, during which dose sufficient to provide reduction of weight is administered, and can be followed by a maintenance phase, during which a e.g. a lower dose sufficient to prevent weight gain is administered. A suitable maintenance dose is likely to be found in the lower parts of the dose ranges provided herein, but corrective and maintenance doses can readily be established for individual subjects by those of skill in the art without undue experimentation, based on the disclosure herein. Maintenance doses can be employed to maintain body weight in subjects whose body weight has been previously controlled by other means, including diet and exercise, bariatric procedures such as bypass or banding surgeries, or treatments employing other pharmacological agents.

III. Pharmaceutical Compositions and Kits

[00112] Another aspect of the invention provides pharmaceutical compositions comprising compounds as disclosed herein formulated together with a pharmaceutically acceptable carrier. In particular, the present disclosure provides pharmaceutical compositions comprising compounds as disclosed herein formulated together with one or more pharmaceutically acceptable carriers. These formulations include those suitable for oral, rectal, topical, buccal, parenteral (e.g., subcutaneous, intramuscular, intradermal, or intravenous) rectal, vaginal, or aerosol administration, although the most suitable form of administration in any given case will depend on the degree and severity of the condition being treated and on the nature of the particular compound being used. For example, disclosed compositions may be formulated as a unit dose, and/or may be formulated for oral or subcutaneous administration.

[00113] Exemplary pharmaceutical compositions of this invention may be used in the form of a pharmaceutical preparation, for example, in solid, semisolid or liquid form, which contains one or more of the compound of the invention, as an active ingredient, in admixture with an organic or inorganic carrier or excipient suitable for external, enteral or parenteral

- 34 -

applications. The active ingredient may be compounded, for example, with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, suppositories, solutions, emulsions, suspensions, and any other form suitable for use. The active object compound is included in the pharmaceutical composition in an amount sufficient to produce the
5 desired effect upon the process or condition of the disease.

[00114] For preparing solid compositions such as tablets, the principal active ingredient may be mixed with a pharmaceutical carrier, *e.g.*, conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, *e.g.*, water, to form a solid
10 preformulation composition containing a homogeneous mixture of a compound of the invention, or a non-toxic pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules.

[00115] In solid dosage forms for oral administration (capsules, tablets, pills, dragees, powders, granules and the like), the subject composition is mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin,
20 polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, acetyl alcohol and glycerol monostearate; (8) absorbents, such as kaolin and
25 bentonite clay; (9) lubricants, such a talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and (10) coloring agents. In the case of capsules, tablets and pills, the compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular
30 weight polyethylene glycols and the like.

[00116] A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin

- 35 -

or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the subject composition moistened with an inert liquid diluent. Tablets, and other
5 solid dosage forms, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art.

[00117] Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders.

10 Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the subject composition, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate,
15 propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, cyclodextrins and mixtures thereof.

[00118] Suspensions, in addition to the subject composition, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan
20 esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

[00119] Formulations for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing a subject composition with one or more suitable non-irritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol,
25 a suppository wax or a salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the body cavity and release the active agent.

[00120] Dosage forms for transdermal administration of a subject composition include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active component may be mixed under sterile conditions with a pharmaceutically acceptable
30 carrier, and with any preservatives, buffers, or propellants which may be required.

- 36 -

[00121] The ointments, pastes, creams and gels may contain, in addition to a subject composition, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

5 [00122] Powders and sprays may contain, in addition to a subject composition, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays may additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

10 [00123] Compositions and compounds of the present invention may alternatively be administered by aerosol. This is accomplished by preparing an aqueous aerosol, liposomal preparation or solid particles containing the compound. A non-aqueous (e.g., fluorocarbon propellant) suspension could be used. Sonic nebulizers may be used because they minimize exposing the agent to shear, which may result in degradation of the compounds contained in the
15 subject compositions. Ordinarily, an aqueous aerosol is made by formulating an aqueous solution or suspension of a subject composition together with conventional pharmaceutically acceptable carriers and stabilizers. The carriers and stabilizers vary with the requirements of the particular subject composition, but typically include non-ionic surfactants (Tweens, Pluronic, or polyethylene glycol), innocuous proteins like serum albumin, sorbitan esters, oleic
20 acid, lecithin, amino acids such as glycine, buffers, salts, sugars or sugar alcohols. Aerosols generally are prepared from isotonic solutions.

[00124] Pharmaceutical compositions of this invention suitable for parenteral administration comprise a subject composition in combination with one or more pharmaceutically-acceptable sterile isotonic aqueous or non-aqueous solutions, dispersions,
25 suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

[00125] Examples of suitable aqueous and non-aqueous carriers which may be employed
30 in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate and

- 37 -

cyclodextrins. Proper fluidity may be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants

[00126] In another aspect, the invention provides enteral pharmaceutical formulations including a disclosed compound and an enteric material; and a pharmaceutically acceptable carrier or excipient thereof. Enteric materials refer to polymers that are substantially insoluble in the acidic environment of the stomach, and that are predominantly soluble in intestinal fluids at specific pHs. The small intestine is the part of the gastrointestinal tract (gut) between the stomach and the large intestine, and includes the duodenum, jejunum, and ileum. The pH of the duodenum is about 5.5, the pH of the jejunum is about 6.5 and the pH of the distal ileum is about 7.5. Accordingly, enteric materials are not soluble, for example, until a pH of about 5.0, of about 5.2, of about 5.4, of about 5.6, of about 5.8, of about 6.0, of about 6.2, of about 6.4, of about 6.6, of about 6.8, of about 7.0, of about 7.2, of about 7.4, of about 7.6, of about 7.8, of about 8.0, of about 8.2, of about 8.4, of about 8.6, of about 8.8, of about 9.0, of about 9.2, of about 9.4, of about 9.6, of about 9.8, or of about 10.0. Exemplary enteric materials include cellulose acetate phthalate (CAP), hydroxypropyl methylcellulose phthalate (HPMCP), polyvinyl acetate phthalate (PVAP), hydroxypropyl methylcellulose acetate succinate (HPMCAS), cellulose acetate trimellitate, hydroxypropyl methylcellulose succinate, cellulose acetate succinate, cellulose acetate hexahydrophthalate, cellulose propionate phthalate, cellulose acetate maleate, cellulose acetate butyrate, cellulose acetate propionate, copolymer of methylmethacrylic acid and methyl methacrylate, copolymer of methyl acrylate, methylmethacrylate and methacrylic acid, copolymer of methylvinyl ether and maleic anhydride (Gantrez ES series), ethyl methacrylate-methylmethacrylate-chlorotrimethylammonium ethyl acrylate copolymer, natural resins such as zein, shellac and copal colophonium, and several commercially available enteric dispersion systems (e. g. , Eudragit L30D55, Eudragit FS30D, Eudragit L100, Eudragit S100, Kollicoat EMM30D, Estacryl 30D, Coateric, and Aquateric). The solubility of each of the above materials is either known or is readily determinable *in vitro*. The foregoing is a list of possible materials, but one of skill in the art with the benefit of the disclosure would recognize that it is not comprehensive and that there are other enteric materials that would meet the objectives of the present invention.

- 38 -

[00127] Advantageously, the invention also provides kits for use by a e.g. a consumer in need of weight loss. Such kits include a suitable dosage form such as those described above and instructions describing the method of using such dosage form to mediate, reduce or prevent inflammation. The instructions would direct the consumer or medical personnel to administer
5 the dosage form according to administration modes known to those skilled in the art. Such kits could advantageously be packaged and sold in single or multiple kit units. An example of such a kit is a so-called blister pack. Blister packs are well known in the packaging industry and are being widely used for the packaging of pharmaceutical unit dosage forms (tablets, capsules, and the like). Blister packs generally consist of a sheet of relatively stiff material covered with
10 a foil of a preferably transparent plastic material. During the packaging process recesses are formed in the plastic foil. The recesses have the size and shape of the tablets or capsules to be packed. Next, the tablets or capsules are placed in the recesses and the sheet of relatively stiff material is sealed against the plastic foil at the face of the foil which is opposite from the direction in which the recesses were formed. As a result, the tablets or capsules are sealed in
15 the recesses between the plastic foil and the sheet. Preferably the strength of the sheet is such that the tablets or capsules can be removed from the blister pack by manually applying pressure on the recesses whereby an opening is formed in the sheet at the place of the recess. The tablet or capsule can then be removed via said opening.

[00128] It may be desirable to provide a memory aid on the kit, *e.g.*, in the form of
20 numbers next to the tablets or capsules whereby the numbers correspond with the days of the regimen which the tablets or capsules so specified should be ingested. Another example of such a memory aid is a calendar printed on the card, *e.g.*, as follows “First Week, Monday, Tuesday, . . . etc. . . . Second Week, Monday, Tuesday, . . . “ etc. Other variations of memory aids will be readily apparent. A “daily dose” can be a single tablet or capsule or several pills or
25 capsules to be taken on a given day. Also, a daily dose of a first compound can consist of one tablet or capsule while a daily dose of the second compound can consist of several tablets or capsules and vice versa. The memory aid should reflect this.

[00129] Also contemplated herein are methods and compositions that include a second active agent, or administering a second active agent. For example, in addition to being
30 overweight or obese, a subject or patient can further have overweight- or obesity-related co-morbidities, *i.e.*, diseases and other adverse health conditions associated with, exacerbated by, or precipitated by being overweight or obese. Contemplated herein are disclosed compounds in

- 39 -

combination with at least one other agent that has previously been shown to treat these overweight- or obesity-related conditions.

[00130] For example, Type II diabetes has been associated with obesity. Certain complications of Type II diabetes, e.g., disability and premature death, can be prevented, ameliorated, or eliminated by sustained weight loss (Astrup, A. Pub Health Nutr (2001) 4:499-515). Agents administered to treat Type II diabetes include sulfonylureas (e.g., Chlorpropamide, Glipizide, Glyburide, Glimepiride); meglitinides (e.g., Repaglinide and Nateglinide); biguanides (e.g., Metformin); thiazolidinediones (Rosiglitazone, Troglitazone, and Pioglitazone); dipeptidylpeptidase-4 inhibitors (e.g., Sitagliptin, Vildagliptin, and Saxagliptin); glucagon-like peptide-1 mimetics (e.g., Exenatide and Liraglutide); and alpha-glucosidase inhibitors (e.g., Acarbose and Miglitol).

[00131] Cardiac disorders and conditions, for example hypertension, dyslipidemia, ischemic heart disease, cardiomyopathy, cardiac infarction, stroke, venous thromboembolic disease and pulmonary hypertension, have been linked to overweight or obesity. For example, hypertension has been linked to obesity because excess adipose tissue secretes substances that are acted on by the kidneys, resulting in hypertension. Additionally, with obesity there are generally higher amounts of insulin produced (because of the excess adipose tissue) and this excess insulin also elevates blood pressure. A major treatment option of hypertension is weight loss. Agents administered to treat hypertension include Chlorthalidone; Hydrochlorothiazide; Indapamide, Metolazone; loop diuretics (e.g., Bumetanide, Ethacrynic acid, Furosemide, Lasix, Torsemide); potassium-sparing agents (e.g., Amiloride hydrochloride, benzamil, Spironolactone, and Triamterene); peripheral agents (e.g., Reserpine); central alpha-agonists (e.g., Clonidine hydrochloride, Guanabenz acetate, Guanfacine hydrochloride, and Methyldopa); alpha-blockers (e.g., Doxazosin mesylate, Prazosin hydrochloride, and Terazosin hydrochloride); beta-blockers (e.g., Acebutolol, Atenolol, Betaxolol, Bisoprolol fumarate, Carteolol hydrochloride, Metoprolol tartrate, Metoprolol succinate, Nadolol, Penbutolol sulfate, Pindolol, Propranolol hydrochloride, and Timolol maleate); combined alpha- and beta-blockers (e.g., Carvedilol and Labetalol hydrochloride); direct vasodilators (e.g., Hydralazine hydrochloride and Minoxidil); calcium antagonists (e.g., Diltiazem hydrochloride and Verapamil hydrochloride); dihydropyridines (e.g., Amlodipine besylate, Felodipine, Isradipine, Nicardipine, Nifedipine, and Nisoldipine); ACE inhibitors (benazepril hydrochloride, Captopril, Enalapril maleate, Fosinopril sodium, Lisinopril, Moexipril, Quinapril

- 40 -

hydrochloride, Ramipril, Trandolapril); Angiotensin II receptor blockers (e.g., Losartan potassium, Valsartan, and Irbesartan); Renin inhibitors (e.g., Aliskiren); and combinations thereof. These compounds are administered in regimens and at dosages known in the art.

[00132] Carr et al. (The Journal of Clinical Endocrinology & Metabolism (2004) Vol.

5 89, No. 6 2601-2607) discusses a link between being overweight or obese and dyslipidemia. Dyslipidemia is typically treated with statins. Statins, HMG-CoA reductase inhibitors, slow down production of cholesterol in a subject and/or remove cholesterol buildup from arteries. Statins include mevastatin, lovastatin, pravastatin, simvastatin, velostatin, dihydrocompactin, fluvastatin, atorvastatin, dalvastatin, carvastatin, crilvastatin, bevastatin, cefvastatin,

10 rosuvastatin, pitavastatin, and glenvastatin. These compounds are administered in regimens and at dosages known in the art. Eckel (Circulation (1997) 96:3248-3250) discusses a link between being overweight or obese and ischemic heart disease. Agents administered to treat ischemic heart disease include statins, nitrates (e.g., Isosorbide Dinitrate and Isosorbide Mononitrate), beta-blockers, and calcium channel antagonists. These compounds are
15 administered in regimens and at dosages known in the art.

[00133] Wong et al. (Nature Clinical Practice Cardiovascular Medicine (2007) 4:436-443) discusses a link between being overweight or obese and cardiomyopathy. Agents

administered to treat cardiomyopathy include inotropic agents (e.g., Digoxin), diuretics (e.g., Furosemide), ACE inhibitors, calcium antagonists, anti-arrhythmic agents (e.g., Sotalol,

20 Amiodarone and Disopyramide), and beta-blockers. These compounds are administered in regimens and at dosages known in the art. Yusef et al. (Lancet (2005) 366(9497):1640-1649)

discusses a link between being overweight or obese and cardiac infarction. Agents administered to treat cardiac infarction include ACE inhibitors, Angiotensin II receptor blockers, direct vasodilators, beta blockers, anti-arrhythmic agents and thrombolytic agents

25 (e.g., Alteplase, Retaplast, Tenecteplase, Anistreplase, and Urokinase). These compounds are administered in regimens and at dosages known in the art.

[00134] Suk et al. (Stroke (2003) 34:1586-1592) discusses a link between being overweight or obese and strokes. Agents administered to treat strokes include anti-platelet agents (e.g., Aspirin, Clopidogrel, Dipyridamole, and Ticlopidine), anticoagulant agents (e.g.,

30 Heparin), and thrombolytic agents. Stein et al. (The American Journal of Medicine (2005)

18(9):978-980) discusses a link between being overweight or obese and venous

thromboembolic disease. Agents administered to treat venous thromboembolic disease include

anti-platelet agents, anticoagulant agents, and thrombolytic agents. Sztrymf et al. (Rev Pneumol Clin (2002) 58(2):104-10) discusses a link between being overweight or obese and pulmonary hypertension. Agents administered to treat pulmonary hypertension include inotropic agents, anticoagulant agents, diuretics, potassium (e.g., K-dur), vasodilators (e.g.,
5 Nifedipine and Diltiazem), Bosentan, Epoprostenol, and Sildenafil. Respiratory disorders and conditions such as obesity-hypoventilation syndrome, asthma, and obstructive sleep apnea, have been linked to being overweight or obese. Elamin (*Chest* (2004) 125:1972-1974) discusses a link between being overweight or obese and asthma. Agents administered to treat asthma include bronchodilators, anti-inflammatory agents, leukotriene blockers, and anti-Ige
10 agents. Particular asthma agents include Zafirlukast, Flunisolide, Triamcinolone, Beclomethasone, Terbutaline, Fluticasone, Formoterol, Beclomethasone, Salmeterol, Theophylline, and Xopenex.

[00135] Kessler et al. (Eur Respir J (1996) 9:787-794) discusses a link between being overweight or obese and obstructive sleep apnea. Agents administered to treat sleep apnea
15 include Modafinil and amphetamines.

[00136] Hepatic disorders and conditions, such as nonalcoholic fatty liver disease, have been linked to being overweight or obese. Tolman et al. (Ther Clin Risk Manag (2007) 6:1153-1163) discusses a link between being overweight or obese and nonalcoholic fatty liver disease. Agents administered to treat nonalcoholic fatty liver disease include antioxidants (e.g.,
20 Vitamins E and C), insulin sensitizers (Metformin, Pioglitazone, Rosiglitazone, and Betaine), hepatoprotectants, and lipid-lowering agents.

[00137] Skeletal disorders and conditions, such as, back pain and osteoarthritis of weight-bearing joints, have been linked to being overweight or obese. van Saase (J Rheumatol (1988) 15(7):1152-1158) discusses a link between being overweight or obese and osteoarthritis
25 of weight-bearing joints. Agents administered to treat osteoarthritis of weight-bearing joints include Acetaminophen, non-steroidal anti-inflammatory agents (e.g., Ibuprofen, Etodolac, Oxaprozin, Naproxen, Diclofenac, and Nabumetone), COX-2 inhibitors (e.g., Celecoxib), steroids, supplements (e.g. glucosamine and chondroitin sulfate), and artificial joint fluid.

[00138] Metabolic disorders and conditions, for example, Prader-Willi Syndrome and polycystic ovary syndrome, have been linked to being overweight or obese. Cassidy (Journal of Medical Genetics (1997) 34:917-923) discusses a link between being overweight or obese
30 and Prader-Willi Syndrome. Agents administered to treat Prader-Willi Syndrome include

- 42 -

human growth hormone (HGH), somatropin, and weight loss agents (e.g., Orlistat, Sibutramine, Methamphetamine, Ionamin, Phentermine, Bupropion, Diethylpropion, Phendimetrazine, Benzphetamine, and Topamax).

[00139] Hoeger (Obstetrics and Gynecology Clinics of North America (2001) 28(1):85-

5 97) discusses a link between being overweight or obese and polycystic ovary syndrome.

Agents administered to treat polycystic ovary syndrome include insulin-sensitizers, combinations of synthetic estrogen and progesterone, Spironolactone, Eflornithine, and Clomiphene. Reproductive disorders and conditions such as sexual dysfunction, erectile

10 dysfunction, infertility, obstetric complications, and fetal abnormalities, have been linked to

being overweight or obese. Larsen et al. (Int J Obes (Lond) (2007) 8:1189-1198) discusses a link between being overweight or obese and sexual dysfunction. Chung et al. (Eur Urol (1999) 36(1):68-70) discusses a link between being overweight or obese and erectile dysfunction.

Agents administered to treat erectile dysfunction include phosphodiesterase inhibitors (e.g., Tadalafil, Sildenafil citrate, and Vardenafil), prostaglandin E analogs (e.g., Alprostadil),

15 alkaloids (e.g., Yohimbine), and testosterone. Pasquali et al. (Hum Reprod (1997) 1:82-87) discusses a link between being overweight or obese and infertility. Agents administered to

treat infertility include Clomiphene, Clomiphene citrate, Bromocriptine, Gonadotropin-releasing Hormone (GnRH), GnRH agonist, GnRH antagonist, Tamoxifen/nolvadex, gonadotropins, Human Chorionic Gonadotropin (HCG), Human Menopausal Gonadotropin

20 (HmG), progesterone, recombinant follicle stimulating hormone (FSH), Urofollitropin, Heparin, Follitropin alfa, and Follitropin beta.

[00140] Weiss et al. (American Journal of Obstetrics and Gynecology (2004)

190(4):1091-1097) discusses a link between being overweight or obese and obstetric

complications. Agents administered to treat obstetric complications include Bupivacaine

25 hydrochloride, Dinoprostone PGE2, Meperidine HCl, Ferro-folic-500/iberet-folic-500,

Meperidine, Methylegonovine maleate, Ropivacaine HCl, Nalbuphine HCl, Oxymorphone HCl, Oxytocin, Dinoprostone, Ritodrine, Scopolamine hydrobromide, Sufentanil citrate, and Oxytocic.

[00141] Psychiatric disorders and conditions, for example, weight-associated depression

30 and anxiety, have been linked to being overweight or obese. Dixon et al. (Arch Intern Med (2003) 163:2058-2065) discusses a link between being overweight or obese and depression.

Agents administered to treat depression include serotonin reuptake inhibitors (e.g., Fluoxetine,

- 43 -

Escitalopram, Citalopram, Paroxetine, Sertraline, and Venlafaxine); tricyclic antidepressants (e.g., Amitriptyline, Amoxapine, Clomipramine, Desipramine, Dosulepin hydrochloride, Doxepin, Imipramine, Iprindole, Lofepramine, Nortriptyline, Opipramol, Protriptyline, and Trimipramine); monoamine oxidase inhibitors (e.g., Isocarboxazid, Moclobemide, Phenelzine, 5 Tranlycypromine, Selegiline, Rasagiline, Nialamide, Iproniazid, Iproclozide, Toloxatone, Linezolid, Dienolide kavapyrone desmethoxyangonin, and Dextroamphetamine); psychostimulants (e.g., Amphetamine, Methamphetamine, Methylphenidate, and Arecoline); antipsychotics (e.g., Butyrophenones, Phenothiazines, Thioxanthenes, Clozapine, Olanzapine, Risperidone, Quetiapine, Ziprasidone, Amisulpride, Paliperidone, Symbyax, Tetrabenazine, 10 and Cannabidiol); and mood stabilizers (e.g., Lithium carbonate, Valproic acid, Divalproex sodium, Sodium valproate, Lamotrigine, Carbamazepine, Gabapentin, Oxcarbazepine, and Topiramate).

[00142] Simon et al. (Archives of General Psychiatry (2006) 63(7):824-830) discusses a link between being overweight or obese and anxiety. Agents administered to treat anxiety 15 include serotonin reuptake inhibitors, mood stabilizers, benzodiazepines (e.g., Alprazolam, Clonazepam, Diazepam, and Lorazepam), tricyclic antidepressants, monoamine oxidase inhibitors, and beta-blockers.

[00143] Another aspect of the invention provides methods for facilitating and maintaining weight loss in a subject involving administering to the subject an amount of a 20 disclosed compound effective to result in weight loss in the subject; and administering a therapeutically effective amount of a different weight loss agent to maintain a reduced weight in the subject. Weight loss agents include serotonin and noradrenergic re-uptake inhibitors; noradrenergic re-uptake inhibitors; selective serotonin re-uptake inhibitors; and intestinal lipase inhibitors. Particular weight loss agents include orlistat, sibutramine, methamphetamine, 25 ionamin, phentermine, bupropion, diethylpropion, phendimetrazine, benzphetamine, bromocriptine, lorcaserin, topiramate, or agents acting to modulate food intake by blocking ghrelin action, inhibiting diacylglycerol acyltransferase 1 (DGAT1) activity, inhibiting stearoyl CoA desaturase 1 (SCD1) activity, inhibiting neuropeptide Y receptor 1 function, activating neuropeptide Y receptor 2 or 4 function, or inhibiting activity of sodium-glucose cotransporters 30 1 or 2. These compounds are administered in regimens and at dosages known in the art.

- 44 -

EXAMPLES

[00144] The compounds described herein can be prepared in a number of ways based on the teachings contained herein and synthetic procedures known in the art. In the description of the synthetic methods described below, it is to be understood that all proposed reaction conditions, including choice of solvent, reaction atmosphere, reaction temperature, duration of the experiment and workup procedures, can be chosen to be the conditions standard for that reaction, unless otherwise indicated. It is understood by one skilled in the art of organic synthesis that the functionality present on various portions of the molecule should be compatible with the reagents and reactions proposed. Substituents not compatible with the reaction conditions will be apparent to one skilled in the art, and alternate methods are therefore indicated. The starting materials for the examples are either commercially available or are readily prepared by standard methods from known materials.

[00145] At least some of the compounds identified as "Intermediates" herein are contemplated as compounds of the invention.

[00146] ¹H NMR spectra were recorded at ambient temperature using a Varian Unity Inova (400MHz) spectrometer with a triple resonance 5mm probe for Example compounds, and either a Bruker Avance DRX (400MHz) spectrometer or a Bruker Avance DPX (300MHz) spectrometer for Intermediate compounds. Chemical shifts are expressed in ppm relative to tetramethylsilane. The following abbreviations have been used: br = broad signal, s = singlet, d = doublet, dd = double doublet, dt = double triplet, t = triplet, q = quartet, m = multiplet.

[00147] Mass Spectrometry (LCMS) experiments to determine retention times and associated mass ions were performed using the following methods:

[00148] Method A: Experiments were performed on a Waters V G Platform II quadrupole spectrometer linked to a Hewlett Packard 1050 LC system with a diode array detector. The spectrometer has an electrospray source operating in positive and negative ion mode. Additional detection was achieved using a Sedex 85 evaporative light scattering detector. LC was carried out using a Luna 3micron 30 x 4.6mm C18 column and a 2mL/minute flow rate. The initial solvent system was 95% water containing 0.1% formic acid (solvent A) and 5% acetonitrile containing 0.1% formic acid (solvent B) for the first 0.3 minute followed by a gradient up to 5% solvent A and 95% solvent B over the next 4 minutes. The final solvent system was held constant for a further 1 minute.

- 45 -

[00149] Method B: Experiments were performed on a Waters Micromass ZQ2000 quadrupole mass spectrometer linked to a Waters Acquity UPLC system with a PDA UV detector. The spectrometer has an electrospray source operating in positive and negative ion mode. LC was carried out using an Acquity BEH 1.7micron C18 column, an Acquity BEH
5 Shield 1.7micron RP18 column or an Acquity HSST 1.8micron column. Each column has dimensions of 100 x 2.1mm and was maintained at 40°C with a flow rate of 0.4mL/minute. The initial solvent system was 95% water containing 0.1% formic acid (solvent A) and 5% acetonitrile containing 0.1% formic acid (solvent B) for the first 0.4 minute followed by a gradient up to 5% solvent A and 95% solvent B over the next 6 minutes. The final solvent
10 system was held constant for a further 0.8 minutes.

[00150] Microwave experiments were carried out using a Biotage Initiator™, which uses a single-mode resonator and dynamic field tuning. Temperatures from 40-250 °C can be achieved, and pressures of up to 20 bars can be reached. A facility exists to apply air cooling during the irradiation.

15 [00151] Preparative HPLC purification was carried out using either a C18-reverse-phase column from Genesis (C18) or a C6-phenyl column from Phenomenex (C6-phenyl) (100 × 22.5 mm i.d. with 7 micron particle size, UV detection at 230 or 254 nm, flow 5-15mL/min), eluting with gradients from 100-0 to 0-100 % water/acetonitrile or water/methanol containing 0.1% formic acid. Fractions containing the required product (identified by LCMS analysis) were
20 pooled, the organic fraction removed by evaporation, and the remaining aqueous fraction lyophilised, to give the product.


[00152] Compounds which required column chromatography were purified manually or fully automatically using either a Biotage SP1™ Flash Purification system with Touch Logic Control™ or a Combiflash Companion® with pre-packed silica gel Isolute® SPE cartridge,
25 Biotage SNAP cartridge or Redisep® Rf cartridge respectively.

[00153] Abbreviations:

DCM: Dichloromethane
DMSO: Dimethylsulfoxide
THF: Tetrahydrofuran
30 TBME: *tert*-Butyl methyl ether
DMAP: 4-Dimethylaminopyridine

- 46 -

Example 1: Cis-(1aRS,7bSR)-5-[2-((Z)-3-diethylaminoprop-1-enyl)-4-fluoro-benzenesulfonylmethyl]-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylic acid



Mixture of enantiomers

- 5 **[00154]** A mixture of methyl cis-(1aRS,7bSR)-5-[2-((Z)-3-diethylaminoprop-1-enyl)-4-fluoro-benzenesulfonylmethyl]-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate (Intermediate 1, 0.72g) and lithium hydroxide (0.622g) in dioxane (20mL) and water (5mL) was stirred and heated at 80°C for 5 hours. After cooling, the mixture was filtered and the filtrate was concentrated to low volume under vacuum. The aqueous residue was acidified with
- 10 formic acid to pH 5 and extracted with DCM. The organic layer was dried (MgSO₄) and filtered. The filtrate was evaporated to dryness and the residue was purified by reverse phase chromatography using a C-18 5 micron cartridge and eluting with a mixture of methanol and water, with a gradient of 10-98%. The product was repurified by HPLC (C18) eluting with a mixture of acetonitrile and water (containing 0.1% formic acid) to give cis-(1aRS,7bSR)-5-[2-
- 15 ((Z)-3-diethylaminoprop-1-enyl)-4-fluorobenzenesulfonylmethyl]-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylic acid (0.125g) as a white solid.

[00155] ¹H NMR (DMSO) δ: 7.45 (1H, dd), 7.36 (1H, d), 7.21 (2H, d), 7.06 (1H, d), 6.87 (1H, d), 6.22 (1H, m), 4.61 (2H, s), 4.08 (1H, d), 3.81 (2H, d), 3.50 (1H, d), 2.99 (4H, q), 1.91 (1H, m), 1.70 (1H, m), 1.09 (6H, t), 0.93 (1H, m), 0.72 (1H, m).

- 20 **[00156]** LCMS (Method B): r/t 2.96 (M+H) 474.

Examples 2 and 3: Separation of enantiomers from Example 1.

[00157] Sample from Example 1 was subjected to chiral separation using a Chiralpak IC column, 10mm x 250mm, particle size 5micron. Eluting solvent ethanol : TBME (20%).

- 25 **Example 2:** First eluting enantiomer: r/t on analytical column (4.6mm x 250mm): 8.5 minutes >98% ee

[00158] ¹H NMR (CDCl₃) δ: 7.77 (1H, d), 7.46 (1H, dd), 7.11 (1H, d), 7.03 (1H, d), 6.89 (1H, dt), 6.79 (1H, dd), 5.99 (1H, m), 5.2-4.8 (1H, br s), 4.7-4.3 (1H, br s), 4.3-3.0 (2H, br),

- 47 -

4.22 (1H, d), 3.67 (1H, br s), 3.12 (4H, m), 1.87 (1H, m), 1.62 (1H, m), 1.24 (6H, t), 1.06 (1H, m), 0.92 (1H, m).


[00159] LCMS (Method B): r/t 2.95 (M+H) 474.

Example 3: Second eluting enantiomer: r/t on analytical column (4.6mm x 250mm): 11.8 minutes >98% ee

[00160] ¹H NMR (CDCl₃) δ: 7.77 (1H, d), 7.46 (1H, dd), 7.11 (1H, d), 7.03 (1H, d), 6.89 (1H, dt), 6.79 (1H, dd), 5.99 (1H, m), 5.2-4.8 (1H, br s), 4.7-4.3 (1H, br s), 4.3-3.0 (2H, br) 4.22 (1H, d), 3.67 (1H, br s), 3.12 (4H, m), 1.87 (1H, m), 1.62 (1H, m), 1.24 (6H, t), 1.06 (1H, m), 0.92 (1H, m).

10 [00161] LCMS (Method B): r/t 2.96 (M+H) 474.

Intermediate 1: Methyl cis-(1aRS,7bSR)-5-[2-((Z)-3-diethylaminoprop-1-enyl)-4-fluorobenzenesulfonylmethyl]-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate




Mixture of enantiomers

15 [00162] A mixture of methyl cis-(1aRS,7bSR)-5-(2-bromo-4-fluorobenzenesulfonylmethyl)-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate (Intermediate 5, 0.66g), N,N-diethyl-N-((Z)-1-tributylstannanylprop-1-en-3-yl)amine (Intermediate 2, 1.17g), tri-*tert*-butylphosphonium tetrafluoroborate (0.145g), *tris*-(dibenzylideneacetone)-dipalladium (0.066g) in dioxane (16mL) and DMSO (1.6mL) was
20 degassed and purged with nitrogen, then heated at 100°C for 1.5 hours. After cooling, the mixture was poured into brine and extracted with ethyl acetate. The organic layer was dried (MgSO₄) and filtered. The filtrate was evaporated to dryness and the residue was purified by chromatography on silica, eluting with a mixture of methanol and DCM with a gradient of 0-15% to give methyl cis-(1aRS,7bSR)-5-[2-((Z)-3-diethylaminoprop-1-enyl)-4-
25 fluorobenzenesulfonylmethyl]-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate (0.72g) as a glass.

[00163] LCMS (Method A) r/t 2.20 (M+H) 488.

- 48 -


Intermediate 2: N,N-Diethyl-N-((Z)-1-tributylstannanylprop-1-en-3-yl)-amine



[00164] Diethylamine (19mL) was added to a solution of ((Z)-3-bromoprop-1-enyl)tributyl-stannane (Intermediate 3, 7.52g) in THF (60mL) and the mixture was stirred for 3 hours. The reaction mixture was evaporated to dryness and the residue was purified by chromatography on a silica column which had been pre-washed with 20% triethylamine in acetonitrile. The column was eluted with a mixture of ethyl acetate and pentane with a gradient of 0-10% to give N,N-diethyl-N-((Z)-1-tributylstannanylprop-1-en-3-yl)amine (4.75 g) as an orange oil.

[00165] ^1H NMR (CDCl_3) δ : 6.59 (1H, dt), 5.97 (1H, dt), 3.08 (2H, dd), 2.53 (4H, q), 1.49 (6H, m), 1.37-1.24 (6H, m), 1.04 (6H, t), 0.92-0.89 (15H, m).


Intermediate 3: ((Z)-3-Bromoprop-1-enyl)tributylstannane



[00166] A solution of triphenylphosphine (5.32g) in DCM (60mL) was added to a solution of (Z)-3-tributylstannanylprop-2-en-1-ol (Intermediate 4, 6.4g) and carbon tetrabromide (9.18g) in DCM (60mL) and the mixture was stirred for 2.5 hours. The mixture was evaporated to low volume and pentane was added. The solids were removed by filtration and the filtrate was evaporated to dryness. Pentane was added and the solids were again removed by filtration and the filtrate was evaporated to dryness to give ((Z)-3-bromoprop-1-enyl)tributylstannane (12.14 g) as an oil.

[00167] ^1H NMR (CDCl_3) δ : 6.71 (1H, dt), 6.11 (1H, d), 3.88 (2H, d), 1.52-1.50 (6H, m), 1.37-1.27 (6H, m), 0.99-0.97 (6H, m), 0.90 (9H, t).

Intermediate 4: (Z)-3-Tributylstannanylprop-2-en-1-ol




- 49 -

[00168] Propargyl alcohol (5mL) was added to a solution of lithium aluminium hydride (1M in THF, 43mL) in THF (70mL) at -78°C. The resultant mixture was warmed to room temperature and stirred for 18 hours. It was re-cooled to -78°C and a solution of tri-n-butyl tin chloride (8.32mL) in diethyl ether (50mL) was added and the mixture was stirred for 3 hours whilst gradually warming to room temperature. The reaction mixture was cooled to -5°C and quenched by addition of water and 15% aqueous sodium hydroxide solution then warmed to room temperature. Ethyl acetate was added and the mixture was stirred for 1 hour. The mixture was filtered through Celite and the filtrate was evaporated to dryness. The residue was purified by chromatography on a silica column which had been pre-washed with 20% triethylamine in acetonitrile. The column was eluted with a mixture of ethyl acetate and pentane with a gradient of 0-10% to give (Z)-3-tributylstannanyl-prop-2-en-1-ol (5.06 g) as a clear oil.

[00169] ¹H NMR (CDCl₃) δ: 6.70 (1H, dt), 6.08 (1H, dt), 4.12 (2H, dd), 1.49 (6H, m), 1.31 (6H, m), 0.98-0.84 (15H, m).

Intermediate 5: Methyl cis-(1aRS,7bSR)-5-(2-bromo-4-fluorobenzenesulfonyl-methyl)-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate




Mixture of enantiomers

[00170] A mixture of methyl cis-(1aRS,7bSR)-5-chloromethyl-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate (Intermediate 6, 0.486g), 2-bromo-4-fluorothiophenol (0.478g), potassium carbonate (0.828g) and potassium iodide (0.01g) in THF (20mL) was stirred and heated at 60°C for 2 hours. After cooling, the mixture was partitioned between ethyl acetate and brine. The organic layer was dried (MgSO₄) and filtered. The filtrate was evaporated to dryness and the residue was dissolved in acetic acid (20mL) and treated with hydrogen peroxide (50%, 5mL). The resultant mixture was stirred at room temperature for 48 hours. Water was added and the mixture was extracted with ethyl acetate, washed with saturated aqueous sodium bicarbonate solution, dried (MgSO₄) and filtered. The filtrate was evaporated to dryness and the residue was purified by chromatography on silica, eluting with a mixture of ethyl acetate and pentane with a gradient of 0-100% to give methyl cis-(1aRS,7bSR)-5-(2-bromo-4-fluorobenzenesulfonylmethyl)-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate (0.781g) as a colourless glass.

- 50 -

[00171] ^1H NMR (CDCl_3) δ : 7.87 (1H, dd), 7.50 (1H, dd), 7.17 (1H, d), 7.07 (1H, dt), 6.74 (1H, d), 4.85 (1H, d), 4.75 (1H, d), 4.34 (1H, d), 3.89 (3H, s), 3.83 (1H, d), 1.95 (1H, m), 1.76 (1H, m), 1.08 (2H, m).

- 5 **Intermediate 6:** Methyl cis-(1aRS,7bSR)-5-chloromethyl-1,1a,2,7b-tetrahydro-cyclopropa[c]chromene-4-carboxylate




Mixture of enantiomers

- [00172] Zinc chloride (0.014g) was added to a solution of methyl cis-(1aRS,7bSR)-5-methoxymethyl-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate (Intermediate 7, 0.53g) and acetyl chloride (0.77mL) in anhydrous diethyl ether (50mL) and the resultant mixture was stirred for 1.5 hours. The mixture was washed with water, dried (MgSO_4) and filtered. The filtrate was evaporated to dryness to give methyl cis-(1aRS,7bSR)-5-chloromethyl-1,1a,2,7b-tetrahydro-cyclopropa[c]chromene -4-carboxylate (0.486g) as a gum.

- [00173] ^1H NMR (CDCl_3) δ : 7.25 (1H, d), 6.94 (1H, d), 4.61 (1H, d), 4.56 (1H, d), 4.37 (1H, d), 3.93 (3H, s), 3.87 (1H, d), 1.99 (1H, m), 1.76 (1H, m), 1.12 (1H, m), 1.07 (1H, m).

Intermediate 7: Methyl cis-(1aRS,7bSR)-5-methoxymethyl-1,1a,2,7b-tetrahydro-cyclopropa[c]chromene-4-carboxylate



Mixture of enantiomers


- 20 [00174] A mixture of methyl cis-(1aRS,7bSR)-1,1-dibromo-5-methoxymethyl-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate (Intermediate 8, 1.06g), zinc dust (4.14g), and ammonium chloride (1.75g) in absolute ethanol (100mL) was stirred and heated at reflux for 6 hours. Further zinc dust (2.0g) and ammonium chloride (0.9g) were added and the mixture was stirred and heated at reflux overnight. After cooling, the mixture was filtered through Celite and

- 51 -

the filtrate was evaporated to dryness. The residue was partitioned between DCM and water and the organic layer was dried (MgSO₄) and filtered. The filtrate was evaporated to dryness to give methyl cis-(1aRS,7bSR)-5-methoxymethyl-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate (0.63g) as an oil.

- 5 [00175] ¹H NMR (CDCl₃) δ: 7.24 (1H, d), 6.89 (1H, d), 4.44 (1H, d), 4.40 (1H, d), 4.36 (1H, d), 3.89 (3H, s), 3.87 (1H, d), 3.31 (3H, s), 1.98 (1H, m), 1.74 (1H, m), 1.10 (1H, m), 1.04 (1H, m).

10 **Intermediate 8:** Methyl cis-(1aRS,7bSR)-1,1-dibromo-5-methoxymethyl-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate




Mixture of enantiomers

- [00176] Aqueous sodium hydroxide solution (50% w/v, 1.63mL) was added slowly to a vigorously stirred mixture of methyl 7-methoxymethyl-2H-chromene-8-carboxylate (Intermediate 9, 0.875g), benzyltriethylammonium chloride (0.173g) and bromoform (15mL).
- 15 The resultant mixture was vigorously stirred and heated at 60°C for 3 hours. After cooling, the mixture was poured into a mixture of brine and ethyl acetate. The aqueous layer was further extracted with ethyl acetate and the combined organic layers were dried (MgSO₄) and filtered. The filtrate was evaporated to dryness and the residue was purified by chromatography on silica, eluting with a mixture of TBME and cyclohexane with a gradient of 0-35% to give
- 20 methyl cis-(1aRS,7bSR)-1,1-dibromo-5-methoxymethyl-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylate (1.06g) as a white solid.

[00177] ¹H NMR (CDCl₃) δ: 7.37 (1H, d), 6.98 (1H, d), 4.50-4.40 (3H, m), 4.35 (1H, dd), 3.87 (3H, s), 3.33 (3H, s), 2.91 (1H, d), 2.45 (1H, m).

- 25 **Intermediate 9:** Methyl 7-methoxymethyl-2H-chromene-8-carboxylate


- 52 -



[00178] A mixture of methyl 2-methoxymethyl-6-propargyloxybenzoate (Intermediate 10, 1.4g) and [*Bis*-(trifluoromethanesulfonyl)imidate] (triphenyl phosphine) gold 2:1 complex with toluene (0.188g) in toluene (40mL) was stirred and heated under an atmosphere of
 5 nitrogen overnight. After cooling, the mixture was filtered through Celite and the filtrate was evaporated to dryness. The residue was purified by chromatography on silica, eluting with a mixture of TBME and cyclohexane with a gradient of 0-30% to give methyl 7-methoxymethyl-2H-chromene-8-carboxylate (0.875g) as an oil.

[00179] ¹H NMR (CDCl₃) δ: 6.96 (1H, d), 6.86 (1H, d), 6.40 (1H, dt), 5.79 (1H, m), 4.86
 10 (2H, dd), 4.43 (2H, s), 3.89 (3H, s), 3.32 (3H, s).

Intermediate 10: Methyl 2-methoxymethyl-6-propargyloxybenzoate




[00180] A mixture of methyl 2-hydroxy-6-methoxymethylbenzoate (Intermediate 11, 1.46g), propargyl bromide (80% solution in toluene, 1.24mL) and potassium carbonate (1.55g) in acetone (50mL) was stirred and heated at reflux for 2 hours. Further propargyl bromide (80% solution in toluene, 0.9mL) was added and the mixture was stirred and heated at reflux for 2 hours. After cooling, the mixture was filtered and the filtrate was evaporated to dryness. The residue was purified by chromatography on silica, eluting with a mixture of ethyl acetate and pentane with a gradient of 0-30% to give methyl 2-methoxymethyl-6-propargyloxybenzoate (1.34g) as an oil.

[00181] ¹H NMR (CDCl₃) δ: 7.35 (1H, t), 7.04 (2H, d), 4.72 (2H, d), 4.47 (2H, s), 3.91 (3H, s), 3.34 (3H, s), 2.50 (1H, t).

25 **Intermediate 11:** Methyl 2-hydroxy-6-methoxymethylbenzoate


- 53 -



[00182] A solution of methyl 2-bromomethyl-6-(*tert*-butyldimethylsilyloxy)benzoate (Intermediate 12, 4.26g) in methanol (63mL) was stirred and heated at reflux overnight. After cooling, the mixture was basified by addition of solid sodium bicarbonate and then evaporated to dryness. The residue was triturated with DCM and filtered. The filtrate was evaporated to dryness and the residue was purified by chromatography on silica, eluting with a mixture of ethyl acetate and cyclohexane with a gradient of 0-25% to give methyl 2-hydroxy-6-methoxymethylbenzoate (1.46g) as a white crystalline solid.

[00183] ¹H NMR (CDCl₃) δ: 11.10 (1H, s), 7.42 (1H, t), 7.11 (1H, d), 6.94 (1H, d), 4.71 (2H, s), 3.98 (3H, s), 3.44 (3H, s).

Intermediate 12: Methyl 2-bromomethyl-6-(*tert*-butyldimethylsilyloxy)benzoate




[00184] N-Bromosuccinimide (7.6g) was added to a solution of methyl 2-(*tert*-butyldimethylsilyloxy)-6-methylbenzoate (Intermediate 13, 11.9g) in carbon tetrachloride (150mL) and the mixture was stirred and heated to reflux. Azo-*bis*-isobutyronitrile (0.1g) was added and the mixture was illuminated with a 500W tungsten filament lamp for 2 hours. The mixture was cooled and left to stand overnight then filtered through a 25g silica cartridge eluting with a mixture of ethyl acetate and pentane (20%). The eluent was evaporated to dryness and the residue was redissolved in carbon tetrachloride (150mL) and N-bromosuccinimide (6.96g) was added. The mixture was heated to reflux and azo-*bis*-isobutyronitrile (0.1g) was added. The resultant mixture was illuminated with a 500W tungsten filament lamp overnight. After cooling, the mixture was filtered and the solid was washed with pentane. The combined organic solutions were evaporated to dryness and the residue was purified by chromatography on silica, eluting with a mixture of TBME and pentane with a gradient of 0-10% to give methyl 2-bromomethyl-6-(*tert*-butyldimethylsilyloxy)benzoate (4.99g) as a yellow oil.

- 54 -

[00185] ^1H NMR (CDCl_3) δ : 7.23 (1H, t), 6.99 (1H, d), 6.80 (1H, d), 4.51 (2H, s), 3.92 (3H, s), 0.97 (9H, s), 0.22 (6H, s).

Intermediate 13: Methyl 2-(*tert*-butyldimethylsilyloxy)-6-methylbenzoate



[00186] A mixture of methyl 2-hydroxy-6-methylbenzoate (prepared according to Doehner *et al*, WO 01/05739, 10.0g), *tert*-butyldimethylsilyl chloride (11.0g), *N,N*-diisopropylethylamine (32mL) and DMAP (0.75g) in DCM (400mL) was stirred at room temperature overnight. The mixture was washed with water and brine, dried (MgSO_4) and filtered. The filtrate was evaporated to dryness and the residue was purified by chromatography on silica eluting with a mixture of ethyl acetate and pentane with a gradient of 0-10% to give methyl 2-(*tert*-butyldimethylsilyloxy)-6-methylbenzoate (11.9g) as an oil.

[00187] ^1H NMR (CDCl_3) δ : 7.13 (1H, t), 6.78 (1H, d), 6.66 (1H, d), 3.87 (3H, s), 2.27 (3H, s), 0.96 (9H, s), 0.21 (6H, s).

Biological Example:

[00188] Compounds are tested for their capacity to inhibit recombinant human MetAP2 activity using the following assay.

[00189] Human recombinant MetAP2 expressed in Sf9 cells followed by affinity purification and EDTA treatment to remove endogenous active site cation was dialysed against MnCl_2 to produce the manganese enzyme used in the assay. The assay was carried out for 30 minutes at 25°C in 50mM HEPES buffer containing 100mM NaCl, at pH 7.5. The presence of 0.75mM Methionine-Alanine-Serine (MAS) substrate and 50 $\mu\text{g}/\text{mL}$ amino acid oxidase using a dilution of purified MetAP2 gives > 3-fold signal : noise. Cleavage of the substrate by MetAP2 and oxidation of free methionine by amino acid oxidase was detected and quantified using fluorescence generated by Amplex red (10-acetyl-3,7-dihydroxyphenoxazine) in combination with horseradish peroxidase which detects H_2O_2 released during the oxidation step. The fluorescent signal was detected using a multiwell fluorimeter. Compounds were diluted in DMSO prior to addition to assay buffer, the final DMSO concentration in the assay being 1%.

- 55 -

[00190] The IC₅₀ is defined as the concentration at which a given compound achieves 50% inhibition of control. IC₅₀ values are calculated using the XLfit software package (version 2.0.5).

[00191] Compounds of the invention demonstrated activity in the assay of this Example
5 as indicated in the following table, wherein A represents < 0.1 μM, B represents IC₅₀ > 0.1 μM.

Compound name	Activity
Cis-(1aRS,7bSR)-5-[2-((Z)-3-diethylaminoprop-1-enyl)-4-fluoro-benzenesulfonylmethyl]-1,1a,2,7b-tetrahydrocyclopropa[c]-chromene-4-carboxylic	B
First eluting enantiomer of cis-(1aRS,7bSR)-5-[2-((Z)-3-diethylaminoprop-1-enyl)-4-fluoro-benzenesulfonylmethyl]-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylic	B
Second eluting enantiomer of cis-(1aRS,7bSR)-5-[2-((Z)-3-diethylaminoprop-1-enyl)-4-fluoro-benzenesulfonylmethyl]-1,1a,2,7b-tetrahydrocyclopropa[c]chromene-4-carboxylic	A

INCORPORATION BY REFERENCE

[00192] All publications and patents mentioned herein, including those items listed
10 below, are hereby incorporated by reference in their entirety for all purposes as if each individual publication or patent was specifically and individually incorporated by reference. In case of conflict, the present application, including any definitions herein, will control.

EQUIVALENTS

[00193] While specific embodiments of the subject invention have been discussed, the above specification is illustrative and not restrictive. Many variations of the invention will
15 become apparent to those skilled in the art upon review of this specification. The full scope of the invention should be determined by reference to the claims, along with their full scope of equivalents, and the specification, along with such variations.

[00194] Unless otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as
20 being modified in all instances by the term “about.” Accordingly, unless indicated to the


- 56 -

contrary, the numerical parameters set forth in this specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention.

[00195] What is claimed it:

- 57 -

1 1. A tricyclic compound represented by:



Formula I

3 wherein:

4 D is a 5-7 membered heterocyclic or carbocyclic ring;

5 X is selected from the group consisting of: $^{+}\text{-C(R}^{D1}\text{R}^{D2}\text{)-C(R}^{D5}\text{R}^{D6}\text{)-*}$, $^{+}\text{-W}^1\text{-C(R}^{D5}\text{R}^{D6}\text{)-}$
 6 * , $^{+}\text{-W}^1\text{-C(O)-*}$, $^{+}\text{-C(R}^{D1}\text{R}^{D2}\text{)-W}^3\text{-*}$, $^{+}\text{-C(R}^{D1}\text{R}^{D2}\text{)-C(R}^{D3}\text{R}^{D4}\text{)-C(R}^{D5}\text{R}^{D6}\text{)-*}$, $^{+}\text{-W}^1\text{-C(R}^{D3}\text{R}^{D4}\text{)-}$
 7 $\text{C(R}^{D5}\text{R}^{D6}\text{)-*}$, $^{+}\text{-W}^1\text{-C(O)-C(R}^{D5}\text{R}^{D6}\text{)-*}$, $^{+}\text{-C(R}^{D1}\text{R}^{D2}\text{)-W}^2\text{-C(R}^{D5}\text{R}^{D6}\text{)-*}$, $^{+}\text{-C(R}^{D1}\text{R}^{D2}\text{)-W}^2\text{-C(O)-}$
 8 * , $^{+}\text{-C(R}^{D1}\text{R}^{D2}\text{)-C(R}^{D3}\text{R}^{D4}\text{)-W}^3\text{-*}$, $^{+}\text{-C(R}^{D1}\text{R}^{D2}\text{)-C(O)-W}^3\text{-*}$, in which the attachment points are
 9 indicated by $^{+}$ and * in X and in Formula I;

10 W^1 is selected from the group consisting of: O or $\text{N(R}^{N1}\text{)}$

11 W^2 is selected from the group consisting of: O or $\text{N(R}^{N2}\text{)}$

12 W^3 is selected from the group consisting of: O or $\text{N(R}^{N3}\text{)}$

13 R^{B1} and R^{B2} are each independently selected from the group consisting of hydrogen,
 14 fluorine, hydroxyl, CN, C_{1-3} alkyl or C_{1-2} alkoxy; wherein the C_{1-3} alkyl and C_{1-2} alkoxy groups
 15 may be optionally substituted by one or more fluorine atoms or a cyano, hydroxyl or C_{1-2} alkoxy
 16 group.

17 R^{B3} and R^{B4} are each independently selected from the group consisting of hydrogen,
 18 fluorine or C_{1-2} alkyl group; wherein the C_{1-2} alkyl group may be optionally substituted by one or
 19 more fluorine atoms.

20 A is a ring selected from the group consisting of: phenyl, a 5-6 membered heteroaryl
 21 having 1, 2 or 3 heteroatoms each selected from S, N or O, and a 4-7 membered heterocycle
 22 having 1, 2 or 3 heteroatoms each selected from N or O;

23 R^{A1} is selected, independently for each occurrence, from the group consisting of:
 24 hydrogen, hydroxyl, cyano, halogen, C_{1-4} alkyl or C_{1-3} alkoxy; wherein C_{1-4} alkyl, or C_{1-3} alkoxy
 25 may be optionally substituted by one or more fluorines;

- 58 -

26 n is 1 or 2;

27 R^{A2} is selected from the group consisting of: hydrogen, R^iR^jN -, heterocyclyl,
 28 heterocyclyloxy, heterocyclyl- (NR^a) -; wherein said heterocyclyl may optionally be substituted
 29 by one or more substituents selected from R^g and wherein if said heterocyclyl contains a $-NH$
 30 moiety that nitrogen may optionally be substituted by one or more groups R^h ; or

31 R^{A2} is selected from the group consisting of: C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6}
 32 cycloalkyl, C_{1-6} alkoxy, C_{3-6} alkenyloxy, C_{3-6} alkynyloxy, C_{3-6} cycloalkoxy, C_{1-6} alkyl- $S(O)_w$ -
 33 (wherein w is 0, 1 or 2), C_{1-6} alkyl- $N(R^a)$ -, C_{1-6} alkyl- $N(R^a)$ -carbonyl-, C_{1-6} alkylcarbonyl- $N(R^a)$ -,
 34 C_{1-6} alkyl- $N(R^a)$ -carbonyl- $N(R^a)$ -, C_{1-6} alkyl- $N(R^a)$ - SO_2 -, C_{1-6} alkyl- SO_2 - $N(R^a)$ -, C_{1-6}
 35 alkoxy carbonyl- $N(R^a)$ -, C_{1-6} alkylcarbonyl- $N(R^a)$ - C_{1-6} alkyl-, C_{1-6} alkyl- $N(R^a)$ -carbonyl- C_{1-6}
 36 alkyl-, C_{1-6} alkoxy- C_{1-6} alkyl-; wherein C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6}
 37 alkoxy, C_{3-6} alkenyloxy, C_{3-6} alkynyloxy, C_{3-6} cycloalkoxy, C_{1-6} alkyl- $S(O)_w$ -, C_{1-6} alkyl- $N(R^a)$ -,
 38 C_{1-6} alkyl- $N(R^a)$ -carbonyl-, C_{1-6} alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkyl- $N(R^a)$ -carbonyl- $N(R^a)$ -, C_{1-6}
 39 alkyl- $N(R^a)$ - SO_2 -, C_{1-6} alkyl- SO_2 - $N(R^a)$ -, C_{1-6} alkoxy carbonyl- $N(R^a)$ -, C_{1-6} alkylcarbonyl-
 40 $N(R^a)$ - C_{1-6} alkyl-, C_{1-6} alkyl- $N(R^a)$ -carbonyl- C_{1-6} alkyl-, C_{1-6} alkoxy- C_{1-6} alkyl may optionally be
 41 substituted by R^p , phenyl, phenoxy, heteroaryl, heteroaryloxy, heteroaryl- (NR^a) -, heterocyclyl,
 42 heterocyclyloxy or heterocyclyl- $N(R^a)$ -; and wherein said heteroaryl or phenyl may optionally
 43 be substituted with one or more substituents selected from R^f ; and wherein said heterocyclyl
 44 may optionally be substituted by one or more substituents selected from R^g ; and wherein if said
 45 heterocyclyl contains a $-NH$ moiety that nitrogen may optionally be substituted by one or more
 46 groups R^h ;

47 R^{D1} and R^{D2} are each independently selected from the group consisting of: hydrogen,
 48 fluorine, hydroxyl, C_{1-2} alkyl or C_{1-2} alkoxy; wherein the C_{1-2} alkyl and C_{1-2} alkoxy may
 49 optionally be substituted by one or more fluorine atoms or a group selected from cyano or
 50 hydroxyl.

51 R^{D3} and R^{D4} are each independently selected from the group consisting of: hydrogen,
 52 fluorine, hydroxyl, cyano, C_{1-3} alkyl or C_{1-3} alkoxy; wherein the C_{1-3} alkyl and C_{1-3} alkoxy may
 53 optionally be substituted by one or more fluorine atoms or a group selected from cyano,
 54 hydroxyl or $N(R^aR^b)$;

55 R^{D5} and R^{D6} are each independently selected from the group consisting of: hydrogen,
 56 fluorine, hydroxyl, cyano, C_{1-2} alkyl or C_{1-2} alkoxy; wherein the C_{1-2} alkyl and C_{1-2} alkoxy may

- 59 -

57 optionally be substituted by one or more fluorine atoms or a group selected from cyano,
58 hydroxyl or $N(R^aR^b)$;

59 R^{N1} is selected from the group consisting of: hydrogen or C_{1-2} alkyl;

60 R^{N2} is selected from the group consisting of: hydrogen, C_{1-3} alkyl or C_{1-2} alkylcarbonyl;
61 wherein the C_{1-3} alkyl and C_{1-2} alkylcarbonyl may optionally be substituted by one or more
62 fluorine atoms or a group selected from cyano, hydroxyl or $N(R^aR^b)$;

63 R^{N3} is selected from the group consisting of: hydrogen, C_{1-3} alkyl or C_{1-2} alkylcarbonyl;
64 wherein the C_{1-3} alkyl and C_{1-2} alkylcarbonyl may optionally be substituted by one or more
65 fluorine atoms or a group selected from cyano, hydroxyl or $N(R^aR^b)$;

66 R^a and R^b are independently selected, for each occurrence, from the group consisting of:
67 hydrogen and C_{1-3} alkyl; wherein C_{1-3} alkyl may optionally be substituted by one or more
68 substituents selected from: fluorine, cyano, oxo and hydroxyl;

69 or R^a and R^b , together with the nitrogen to which they are attached, may form a 4-6
70 membered heterocyclic ring which may have an additional heteroatom selected from O, S, or
71 N; and wherein the 4-6 membered heterocyclic ring may optionally be substituted by one or
72 more substituents selected from the group consisting of: fluorine, cyano, oxo or hydroxyl;

73 R^f is independently selected, for each occurrence, from the group consisting of: R^P ,
74 hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6}
75 alkyl-S(O)_w- (wherein w is 0, 1 or 2), C_{1-6} alkylcarbonyl- $N(R^a)$ -; C_{1-6} alkoxycarbonyl- $N(R^a)$ -;
76 and wherein C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6}
77 alkyl-S(O)_w-, C_{1-6} alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkoxycarbonyl- $N(R^a)$ - may be optionally
78 substituted by one or more substituents selected from R^P ;

79 R^g is independently selected for each occurrence from the group consisting of: R^P ,
80 hydrogen, oxo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl-S(O)_w-
81 (wherein w is 0, 1 or 2), C_{1-6} alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkoxycarbonyl- $N(R^a)$ -; wherein C_{1-6}
82 alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl-S(O)_w-, C_{1-6}
83 alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkoxycarbonyl- $N(R^a)$ - may be optionally substituted by one or
84 more substituents selected from R^P ;

85 R^h is independently selected for each occurrence from the group consisting of:
86 hydrogen, C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkyl-S(O)₂-, C_{1-6}

- 60 -

alkoxycarbonyl-, R^iR^j N-carbonyl-, R^iR^j N-SO₂-; wherein C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, C₃₋₆cycloalkyl, C₁₋₆alkyl-S(O)₂-, C₁₋₆alkylcarbonyl- may optionally be substituted by one or more substituents selected from R^P;

Rⁱ and R^j, are selected independently for each occurrence from the group consisting of: hydrogen, C₁₋₄alkyl and C₃₋₆cycloalkyl; wherein C₁₋₄alkyl and C₃₋₆cycloalkyl may be optionally substituted by one or more substituents selected from fluorine, hydroxyl, cyano, R^aR^bN-, R^aR^bN-carbonyl-, C₁₋₃alkoxy;

or Rⁱ and R^j taken together with the nitrogen to which they are attached form a 4-7 membered heterocyclic ring which may have an additional heteroatom selected from O, S, or N, optionally substituted on carbon by one or more substituents selected from the group consisting of: fluorine, hydroxyl, oxo, cyano, C₁₋₆alkyl, C₁₋₆alkoxy, R^aR^bN-, R^aR^bN-SO₂-, R^aR^bN-carbonyl-; and wherein said C₁₋₆alkyl or C₁₋₆alkoxy may optionally be substituted by fluorine, hydroxyl or cyano; and optionally substituted on nitrogen by one or more substituents selected from the group consisting of: C₁₋₆alkyl, R^aR^bN-carbonyl-; and wherein said C₁₋₆alkyl may be optionally substituted by fluorine, hydroxyl, cyano;

R^P is independently selected, for each occurrence, from the group consisting of: halogen, hydroxyl, cyano, C₁₋₆alkoxy, RⁱR^jN-, RⁱR^jN-carbonyl-, RⁱR^jN-SO₂-, RⁱR^jN-carbonyl-N(R^a)-;

and pharmaceutically acceptable salts, stereoisomers, esters and prodrugs thereof.

2. The tricyclic compound of claim 1, wherein X is selected from the group consisting of $^{+}\text{-C(R}^{D1}\text{R}^{D2}\text{)-C(R}^{D5}\text{R}^{D6}\text{)-*}$, $^{+}\text{-O-C(R}^{D5}\text{R}^{D6}\text{)-*}$, $^{+}\text{-N(R}^{N2}\text{)-C(R}^{D5}\text{R}^{D6}\text{)-*}$, and $^{+}\text{-O-C(R}^{D3}\text{R}^{D4}\text{)-C(R}^{D5}\text{R}^{D6}\text{)-*}$;

wherein the ⁺ and * indicate the attachment points of X as indicated in Formula I.

3. The tricyclic compound of any one of claims 1-2, X is selected from the group consisting of $^{+}\text{-CH}_2\text{-CH}_2\text{-*}$, $^{+}\text{-O-CH}_2\text{-*}$, and $^{+}\text{-NH-CH}_2\text{-*}$;


wherein the ⁺ and * indicate the attachment points of X as indicated in Formula I.

4. The tricyclic compound of any one of claims 1-3, wherein R^{D1}, R^{D2}, R^{N1} and R^{N2} that form part of X are independently selected from the groups consisting of hydrogen and methyl.


5. The tricyclic compound of any one of claims 1-4, wherein R^{D1}, R^{D2}, R^{N1} and R^{N2} that form part of X are hydrogen.

- 61 -

- 1 6. The tricyclic compound of any one of claims 1-5, wherein R^{D3} , R^{D4} , R^{D5} and R^{D6} that form
 2 part of X are independently selected from the group consisting of hydrogen, fluorine, cyano,
 3 and C_{1-2} alkyl.
- 1 7. The tricyclic compound of any one of claims 1-6, wherein R^{D3} , R^{D4} , R^{D5} and R^{D6} that form
 2 part of X are hydrogen.
- 1 8. The tricyclic compound of any one of claims 1-7, wherein R^{B1} and R^{B2} are independently
 2 selected from the group consisting of H and methyl.
- 1 9. The tricyclic compound of any one of claims 1-8, wherein R^{B3} and R^{B4} are independently
 2 selected from the group consisting of H and fluorine.
- 1 10. The tricyclic compound of claim 1, wherein the compound is represented by:



- 2
- 1 11. The tricyclic compound of claim 10, wherein A is phenyl.
- 1 12. A tricyclic compound represented by:



Formula II;

- 2
- 3 wherein:
- 4 R^{A1} is selected from the group consisting of: hydrogen, halogen, C_{1-2} alkyl, C_{1-2} alkoxy;
- 5 wherein C_{1-2} alkyl may optionally be substituted by one or more fluorines;
- 6 R^{A2} is selected from the group consisting of: hydrogen, R^iR^jN , heterocyclyl, C_{1-6} alkyl,
- 7 C_{3-6} alkenyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy; wherein said heterocyclyl may optionally be substituted

- 62 -

by one or more groups R^g , and wherein if said heterocyclyl contains a $-NH$ moiety, that nitrogen may optionally be substituted by one or more groups R^h ; and wherein said C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} cycloalkyl and C_{1-6} alkoxy may optionally be substituted by one or more groups R^P ;

n is 1 or 2;

R^g is independently selected for each occurrence from the group consisting of: R^P , hydrogen, oxo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl- $S(O)_w$ - (wherein w is 0, 1 or 2), C_{1-6} alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkoxycarbonyl- $N(R^a)$ -; wherein C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl- $S(O)_w$ -, C_{1-6} alkylcarbonyl- $N(R^a)$ -, C_{1-6} alkoxycarbonyl- $N(R^a)$ - may be optionally substituted by one or more substituents selected from R^P ; R^h is independently selected for each occurrence from the group consisting of: hydrogen, C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkyl- $S(O)_2$ -, C_{1-6} alkoxycarbonyl-, R^iR^jN -carbonyl-, R^iR^jN - SO_2 -; wherein C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkyl- $S(O)_2$ -, C_{1-6} alkylcarbonyl- may optionally be substituted by one or more substituents selected from R^P ;

R^h is independently selected for each occurrence from the group consisting of: hydrogen, C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkyl- $S(O)_2$ -, C_{1-6} alkoxycarbonyl-, R^iR^jN -carbonyl-, R^iR^jN - SO_2 -; wherein C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-6} cycloalkyl, C_{1-6} alkyl- $S(O)_2$ -, C_{1-6} alkylcarbonyl- may optionally be substituted by one or more substituents selected from R^P ;

R^i and R^j , are selected independently for each occurrence from the group consisting of: hydrogen, C_{1-4} alkyl and C_{3-6} cycloalkyl; wherein C_{1-4} alkyl and C_{3-6} cycloalkyl may be optionally substituted by one or more substituents selected from fluorine, hydroxyl, cyano, R^aR^bN -, R^aR^bN -carbonyl-, C_{1-3} alkoxy;

or R^i and R^j taken together with the nitrogen to which they are attached form a 4-7 membered heterocyclic ring which may have an additional heteroatom selected from O, S, or N, optionally substituted on carbon by one or more substituents selected from the group consisting of: fluorine, hydroxyl, oxo, cyano, C_{1-6} alkyl, C_{1-6} alkoxy, R^aR^bN -, R^aR^bN - SO_2 -, R^aR^bN -carbonyl-; and wherein said C_{1-6} alkyl or C_{1-6} alkoxy may optionally be substituted by fluorine, hydroxyl or cyano; and optionally substituted on nitrogen by one or more substituents

- 63 -

38 selected from the group consisting of: C₁₋₆alkyl, R^aR^bN-carbonyl-; and wherein said C₁₋₆alkyl
39 may be optionally substituted by fluorine, hydroxyl, cyano;

40 R^p is independently selected, for each occurrence, from the group consisting of:
41 halogen, hydroxyl, cyano, C₁₋₆alkoxy, RⁱR^jN-, RⁱR^jN-carbonyl-, RⁱR^jN-SO₂-, RⁱR^jN-carbonyl-
42 N(R^a)-;

43 R^a and R^b are independently selected, for each occurrence, from the group consisting of:
44 hydrogen and C₁₋₃alkyl; wherein C₁₋₃alkyl may optionally be substituted by one or more
45 substituents selected from: fluorine, cyano, oxo and hydroxyl;

46 or R^a and R^b, together with the nitrogen to which they are attached, may form a 4-6
47 membered heterocyclic ring which may have an additional heteroatom selected from O, S, or
48 N; and wherein the 4-6 membered heterocyclic ring may optionally be substituted by one or
49 more substituents selected from the group consisting of: fluorine, cyano, oxo or hydroxyl;

50 and pharmaceutically acceptable salts, stereoisomers, esters and prodrugs thereof.

1 13. The tricyclic compound of claim 12, wherein R^{Al} is selected from the group consisting of
2 hydrogen and fluorine.

1 14. A compound selected from the group consisting of: cis-(1aRS,7bSR)-5-[2-((Z)-3-
2 diethylaminoprop-1-enyl)-4-fluoro-benzenesulfonylmethyl]-1,1a,2,7b-
3 tetrahydrocyclopropa[c]chromene-4-carboxylic acid; cis-(1aR,7bS)-5-[2-((Z)-3-
4 diethylaminoprop-1-enyl)-4-fluoro-benzenesulfonylmethyl]-1,1a,2,7b-
5 tetrahydrocyclopropa[c]chromene-4-carboxylic acid; cis-(1aS,7bR)-5-[2-((Z)-3-
6 diethylaminoprop-1-enyl)-4-fluoro-benzenesulfonylmethyl]-1,1a,2,7b-
7 tetrahydrocyclopropa[c]chromene-4-carboxylic acid and pharmaceutically acceptable salts,
8 stereoisomers, esters and prodrugs thereof.

1 15. A pharmaceutically acceptable composition comprising a compound of any one of claims
2 1-14 and a pharmaceutically acceptable excipient.

1 16. A method of treating and/or controlling obesity, comprising administering to a patient in
2 need thereof an effective amount of a compound of any one of claims 1-14.

1 17. A method of inducing weight loss in a patient in need thereof, comprising administering to
2 said patient an effective amount of a compound of any one of claims 1-14.

- 64 -

- 1 18. The method of claim 16 or 17, wherein the patient is a human.
- 1 19. The method of claim 16 or 17, wherein the patient is a cat or dog.
- 1 20. The method of any one of claims 16-19, wherein the patient has a body mass index greater
2 than or equal to about 30 kg/m² before the administration.
- 1 21. The method of any one of claims 16-20, wherein the compound is administered orally.
- 1 22. The composition of claim 15, wherein the composition is formulated as a unit dose.
- 1 23. The composition of claim 15, wherein the composition is formulated for oral
2 administration.
- 1 24. The composition of claim 15, wherein the composition is formulated for intravenous or
2 subcutaneous administration.
- 1 25. The method of claim 16 or 17, comprising administering said compound in an amount
2 sufficient to establish inhibition of intracellular MetAP2 effective to increase thioredoxin
3 production in the patient and to induce multi organ stimulation of anti-obesity processes in the
4 subject.
- 1 26. The method of claim 25, comprising administering said compound in an amount
2 insufficient to reduce angiogenesis in the patient.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2013/021914

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D311/78 A61K31/353 A61P3/04
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2011/044506 A2 (ZAFGEN CORP [US]; DYKE HAZEL JOAN [GB]; CRAMP SUSAN MARY [GB]; PALLIN) 14 April 2011 (2011-04-14) page 69; example 5 page 226, lines 41-46 abstract	1-26
A	WO 2010/065883 A2 (ZAFGEN CORP [US]; HUGHES THOMAS E [US]; VATH JAMES E [US]) 10 June 2010 (2010-06-10) claim 1 page 14; compound (H) ----- -/--	1,12, 14-17



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

11 April 2013

Date of mailing of the international search report

18/04/2013

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Fitz, Wolfgang

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2013/021914

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WANG ET AL: "Lead optimization of methionine aminopeptidase-2 (MetAP2) inhibitors containing sulfonamides of 5,6-disubstituted anthranilic acids", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, PERGAMON, ELSEVIER SCIENCE, GB, vol. 17, no. 10, 27 April 2007 (2007-04-27), pages 2817-2822, XP022049594, ISSN: 0960-894X, DOI: 10.1016/J.BMCL.2007.02.062 abstract tables 1-4 -----	1,12, 14-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2013/021914

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2011044506 A2	14-04-2011	AU 2010303270 A1	03-05-2012
		CA 2777108 A1	14-04-2011
		CN 102639494 A	15-08-2012
		EP 2486004 A2	15-08-2012
		JP 2013507392 A	04-03-2013
		KR 20120083905 A	26-07-2012
		US 2013053364 A1	28-02-2013
		WO 2011044506 A2	14-04-2011

WO 2010065883 A2	10-06-2010	US 2012034233 A1	09-02-2012
		US 2012322867 A1	20-12-2012
		WO 2010065883 A2	10-06-2010
