



US 20050158342A1

(19) **United States**

(12) **Patent Application Publication**

Kemble et al.

(10) **Pub. No.: US 2005/0158342 A1**

(43) **Pub. Date:**

Jul. 21, 2005

(54) **MULTI PLASMID SYSTEM FOR THE PRODUCTION OF INFLUENZA VIRUS**

(75) Inventors: **George Kemble**, Saratoga, CA (US); **Gregory Duke**, Redwood City, CA (US)

Correspondence Address:
JOHNATHAN KLEIN-EVANS
ONE MEDIMMUNE WAY
GAITHERSBURG, MD 20878 (US)

(73) Assignee: **MEDIMMUNE, INC.**, Gaithersburg, MD

(21) Appl. No.: **11/018,624**

(22) Filed: **Dec. 22, 2004**

Related U.S. Application Data

(60) Provisional application No. 60/532,164, filed on Dec. 23, 2003.

Publication Classification

(51) **Int. Cl.⁷** **A61K 39/12; A61K 39/145;**

C12N 7/02


(52) **U.S. Cl.** **424/209.1; 435/239**

(57)

ABSTRACT

Vectors and methods for the production of influenza viruses suitable as recombinant influenza vaccines in cell culture are provided. Bi-directional expression vectors for use in a multi-plasmid influenza virus expression system are provided. Additionally, the invention provides methods of producing influenza viruses with enhanced ability to replicate in embryonated chicken eggs and/or cells (e.g., Vero and/or MDCK) and further provides influenza viruses with enhanced replication characteristics. In addition, the present invention includes an improved method of rescue, wherein animal cells (e.g., SF Vero cells) are electroporated with plasmids and vectors of the invention.

pAD3000



pAD3000




Fig. 1




Fig. 2




Fig. 3




Fig. 4

Fig. 5A





Fig. 5B






Fig. 5C

Fig. 5D

2. Sequence in Genbank-format

LOCUS pAD3000 2836 bp DNA circular 14-JAN-2002
 DEFINITION Derivative of pHW2000 with SV40 PolyA Signal replacing BGH

FEATURES Location/Qualifiers
 promoter 2420..2799
 /vntifkey="29"
 /label=pCMV
 /note="truncated CMV promoter (corresponding to 484-863
 region of pcDNA3)"
 misc_marker 1422..2282
 /vntifkey="22"
 /label=bla
 /note="beta lactamase"
 rep_origin 612..1172
 /vntifkey="33"
 /label=Col\Elori
 /note="Col E1 replication origin"
 terminator 11..45
 /vntifkey="43"
 /label=tI
 /note="Pol I terminator"
 promoter complement(65..276)
 /vntifkey="29"
 /label=PolI
 /note="Human Pol I Promoter"
 exon 296..430
 /vntifkey="61"
 /label=pA
 /note="pA(SV40)"
 BASE COUNT 717 a 734 c 703 g 682 t
 ORIGIN
 1 ctagcgtta accggagtag tggtcgacct ccgaagttgg gggggaggag acggtaccgt
 61 ctccaataac ccggcgcccc aaaatgccga ctccggagcgaa aagatataacc tccccccgggg
 121 cccggaggcgc gctgtacccga ccacgcccgc ggcggcaggcg acgcgcgaca cggacacac
 181 tcccccaaaaaa cgcaccatc gcagccacac acggagcgcgc cggggccctc tggtcaaccc
 241 caggacacac cggggaggcgc cgcggggccgc gggacgcgc cccggccgtc acctcagaca
 301 tgataagata cattgtatgg tttggacaaa ccacaacttag aatgcagtga aaaaaatgct
 361 ttattttgtga aattttgtat gctattgtct tattttgtaa cattataagc tgcaataaaac
 421 aaggatctgc attaatgaat cggccaacgc gcgggggagag gcggttgcgc tattgggcgc
 481 tcttcgcctt cctcgctcac tgactcgctg cgctcggtcg ttcggctgcgc gcgagcggta
 541 tcagctcact caaaggccgt aatacggta tccacagaat caggggataa cgccggaaag
 601 aacatgttag caaaaggccca gcaaaaggccca aggaacccgtaa aaaaaggccgc gttgctggcg
 661 tttttccata ggctccggcc ccctgacgag catcacaaaaa atcgacgctc aagttagagg
 721 tggcgaaacc cgacaggact ataaagatac caggcgttcc cccctgaaag ctccctcggt
 781 cgctctcctg ttccgaccc tccgcgttacc ggatacctgt ccgcctttct cccttcggga
 841 agcgtggcgc tttctcaatg ctcacgctgt aggtatctca gttcggtgtaa ggtcggttcgc
 901 tccaagctgg gctgtgtca cgaacccccc gttcagcccg accgctgcgc cttatccgt
 961 aactatcgtc ttgagtccaa cccgtaaaga caccacttat cgccactggc agcagccact
 1021 ggttaacagga ttagcagacg gaggtatgtt ggcgggtgtaa cagagttctt gaagtgggtgg
 1081 cctaactacg gctacactag aaggacagta tttggtatct gcgctctgtt gaagccagtt
 1141 accttcgaa aaagagtgg tagctttgtt cccggcaaac aaaccaccgc tggtagcggt
 1201 ggtttttttt ttttgcacca gcaagattacg cgcagaaaaa aaggatctca agaagatcct
 1261 ttgatctttt ctacgggtc tgacgcttag tggaaacgaaa actcacgttta aggattttt
 1321 gtcatgat tataaaaaag gattttccacc tagatcctt taaattaaaa atgaagttt

Fig. 6

1381 aaatcaatct aaagtataata ttagttaaact tggctcgaca gttaccaatg cttaatcagt
1441 gaggcaccta tctcagcgat ctgtcttattt cgttcatccaa tagttgcctg actccccgtc
1501 gtgttagataa ctacgataacg ggagggctta ccatctggcc ccagtgcgtc aatgataccg
1561 cgagacccac gctcaccggc tccagattt tcaagaaata accagccagc cgaaggggcc
1621 gagcgcagaa gtggctctgc aactttatcc gcctccatcc agtctattaa ttgttgcgg
1681 gaagctagag taagttagtcc gccagttaaat agtttgcgca acgttgttgc cattgttaca
1741 ggcacatcggtt tgcacgctc gtcgttggg atggcttcat tcagctccgg tttccaaacga
1801 tcaaggcgag ttacatgatc ccccatgttg tgcaaaaaaag cggttagctc ctccggctt
1861 ccgatcggtt tcagaagtaa gttggccgca gtgttatcac tcatggttat ggcagcactg
1921 cataattctc ttactgtcat gccatccgta agatgtttt ctgtgactgg tgagtactca
1981 accaagtcat tctgagaata gtgtatgcgg cgaccgagttt gctctgccc ggcgtcaata
2041 cgggataata ccgcgccaca tagcagaact taaaagtgc tcataattgg aaaaacgttct
2101 tcggggcgaa aactctcaag gatcttaccg ctgttggat ccagtcgat gtaacccact
2161 cgtgcaccca actgatcttc agcatctttt actttcacca gcggttctgg gtgagcaaaa
2221 acaggaaggc aaaaatgcgc aaaaaggga ataaggcgaa cacggaaatg ttgaatactc
2281 atactttcc ttttcaata ttatttgaagc attttatcagg gttattgtct catgagcgga
2341 tacatatttg aatgtattta gaaaaataaa caaatagggg ttccgcgcac atttccccga
2401 aaagtgcac ctgacgtcga tatgccaagt acgccccctt ttgacgtcaa tgacggtaaa
2461 tggcccgctt ggcattatgc ccagttacatg accttattggg actttcttac ttggcagttac
2521 atctacgtat tagtcatcgc tattaccatg gtgtatgcggg tttggcagta catcaatggg
2581 cgtggatagc gggttgactc acggggattt ccaagtctcc accccattga cgtcaatggg
2641 agttgtttt ggcaccaaaa tcaacgggac tttccaaaat gtcgtaacaa ctccggccca
2701 ttgacgcaaa tgggcggtag gcgtgtacgg tgggaggtct atataagcag agctctctgg
2761 ctaactagag aacccactgc ttactggctt atcggaaatattt atacgactca ctataggag
2821 acccaagctg ttaacg

//

Fig. 6 Cont.

**ALIGNMENT OF CONSENSUS SEQUENCE OF MDV-B WITH CDNA IN THE
EIGHT PLASMIDS (PAB12[N-SEGMENT])**

PB1

	* 20 * 40 *				
pAB121-PB1 :	: 50
MDV-B-PB1 :	: 50
	AGCAGAACGGAGCCTTAAGATGAATATAATCCTTATTTCTCTCAT				
	60 * 80 * 100				
pAB121-PB1 :	: 100
MDV-B-PB1 :	: 100
	AGATGTACCCATACAGGCAGCAATTCAACAAACATTCCATACACCGGTG				
	* 120 * 140 *				
pAB121-PB1 :	: 150
MDV-B-PB1 :	: 150
	TTCCCCCTTATTCCCATGGAACGGGAACAGGCTACACAAATAGACACCGTG				
	160 * 180 * 200				
pAB121-PB1 :	: 200
MDV-B-PB1 :	: 200
	ATTAGAACACATGAGTACTAAACAAGGGAAAACAATACATTCTGATGT				
	* 220 * 240 *				
pAB121-PB1 :	: 250
MDV-B-PB1 :	: 250
	TACAGGATGTGCAATGGTAGATCCAACAAATGGCCATTACCGAAGATA				
	260 * 280 * 300				
pAB121-PB1 :	: 300
MDV-B-PB1 :	: 300
	ATGAGCCAGTGCCTATGCACAAATTGGATTGCGTTCTGGAGGCTTGGAT				
	* 320 * 340 *				
pAB121-PB1 :	: 350
MDV-B-PB1 :	: 350
	AGAATGGATGAAGAACATCCAGGTCTGTTCAAGCAGCCTCACAGAATGC				
	360 * 380 * 400				
pAB121-PB1 :	: 400
MDV-B-PB1 :	: 400
	CATGGAGGCATAATGGTCACAACACTGTAGACAAATTAAACCCAGGGAGAC				
	* 420 * 440 *				
pAB121-PB1 :	: 450
MDV-B-PB1 :	: 450
	AGACTTTGATTGGACAGTGTGCAGAAACCAACCTGCTGCAACGGCACTG				
	460 * 480 * 500				
pAB121-PB1 :	: 500
MDV-B-PB1 :	: 500
	AACACAAACAATAACCTCTTTAGGTTGAATGATTGAATGGAGCCGACAA				

Fig. 7

	* 520		* 540		* 550
pAB121-PB1				:
MDV-B-PB1				:
	GGGTGGATTAGTACCCCTTGGCAAGATATCATTGATTCAATTGGACAAAC				
	560		* 580		* 600
pAB121-PB1				:
MDV-B-PB1				:
	CTGAAATGACTTCTCGGAAAGAATATAAAGAAAAATTGCCTGCT				
	* 620		* 640		* 650
pAB121-PB1				:
MDV-B-PB1				:
	AAAAACAGAAAGGGTTTCCTCATAAAGAGAATACCAATGAAGGTAAAAGA				
	660		* 680		* 700
pAB121-PB1				:
MDV-B-PB1				:
	CAGAATAACCAGAGTGGAAACATCAAAAGAGCATTATCATTAAACACAA				
	* 720		* 740		* 750
pAB121-PB1				:
MDV-B-PB1				:
	TGACAAAAGATGCTGAAAGAGGCAAACATAAAAGAAGAGCAATTGCCACC				
	760		* 780		* 800
pAB121-PB1				:
MDV-B-PB1				:
	GCTGGGATAACAAATCAGAGGGTTGTATTAGTAGTTGAAAACCTGGCTAA				
	* 820		* 840		* 850
pAB121-PB1				:
MDV-B-PB1				:
	AAATATCTGTGAAAATCTAGAACAAAGTGGTTGCCAGTAGGTGGAACG				
	860		* 880		* 900
pAB121-PB1				:
MDV-B-PB1				:
	AGAAGAAGGCCAAACTGTCAAATGCAGTGGCCAAATGCTCAGTAACG				
	* 920		* 940		* 950
pAB121-PB1G.....				:
MDV-B-PB1				:
	CCACCAGGAGGGATCAGCATGACAGTGACAGGAGACAATACTAAATGGAA				
	960		* 980		* 1000
pAB121-PB1				:
MDV-B-PB1				:
	TGAATGCTTAAATCCAAGAATTTGGCTATGACTGAAAGAATAACCA				
	* 1020		* 1040		* 1050
pAB121-PB1				:
MDV-B-PB1				:
	GAGACAGCCCAATTGGTCCGGATTTGTAGTATAGCACCGGTCTTG				
	1060		* 1080		* 1100

Fig. 7 Cont.

pAB121-PB1	:	:	1100
MDV-B-PB1	:	:	1100
TTCTCCAATAAAATGCCAGATTGGGAAAGGGTTCATGATAACAAGCAA							
* 1120 * 1140 *							
pAB121-PB1	:	:	1150
MDV-B-PB1	:	:	1150
AACAAAAAGACTGAAGGCTCAAATACCTTGTCCCGATCTGTTAATATAC							
1160 * 1180 * 1200							
pAB121-PB1	:	:	1200
MDV-B-PB1	:	:	1200
CATTAGAAAGATATAATGAAGAAACAAGGGCAAAATTAAAAAGCTGAAA							
* 1220 * 1240 *							
pAB121-PB1	:	:	1250
MDV-B-PB1	:	:	1250
CCATTCTTCAATGAAGAAGGAACGGCATCTTGTGCCTGGGATGATGAT							
1260 * 1280 * 1300							
pAB121-PB1	:	:	1300
MDV-B-PB1	:	:	1300
GGGAATGTTAACATGCTATCTACCGTGTGGAGTAGCCGCACTAGGGA							
* 1320 * 1340 *							
pAB121-PB1	:	:	1350
MDV-B-PB1	:	:	1350
TCAAAACATTGAAACAAAGAACATTTATGGGATGGACTGCAATCTTCT							
1360 * 1380 * 1400							
pAB121-PB1	:	:	1400
MDV-B-PB1	:	:	1400
GATGATTTGCTCTGTTGTTAACATGCAAAAGATGAAGAGACATGTATGGA							
* 1420 * 1440 *							
pAB121-PB1	:	:	1450
MDV-B-PB1	:	:	1450
AGGAATAAACGATTTTACCGAACATGTAAGCTATTGGAAATAAACATGA							
1460 * 1480 * 1500							
pAB121-PB1	:	:	1500
MDV-B-PB1	:	:	1500
GCAAAAGAAAAGTTACTGTAATGAAACTGGAATGTTGAATTACAAGC							
* 1520 * 1540 *							
pAB121-PB1	:	:	1550
MDV-B-PB1	:	:	1550
ATGTTCTACAGAGATGGATTGTATCTAATTGCAATGGAACCTCCCTC							
1560 * 1580 * 1600							
pAB121-PB1	:	:	1600
MDV-B-PB1	:	:	1600
ATTGGGAGTTGCTGGAGTAAATGAATCAGCAGATATGGCAATAGGAATGA							

Fig. 7 Cont.

	* 1620	* 1640	* 1650
pAB121-PB1	:	:
MDV-B-PB1	:	CAATAATAAGAACAAATATGATCAACAATGGGATGGGCCAGAACAGCA	: 1650
	1660	* 1680	* 1700
pAB121-PB1	:	:
MDV-B-PB1	:	CAAACAGCCATACAATTATTAGCTGATTAGATAACACCTACAAATG	: 1700
	* 1720	* 1740	* 1750
pAB121-PB1	:	T.....	:
MDV-B-PB1	:	CCACAGGGAGATTCAAAGTGGAAAGGAAAGAGAATGAAAATTATAAAGG	: 1750
	1760	* 1780	* 1800
pAB121-PB1	:	:
MDV-B-PB1	:	AGCTATGGGAAAACACTAAAGGAAGAGATGGTCTGTTAGTAGCAGATGGT	: 1800
	* 1820	* 1840	* 1850
pAB121-PB1	:	:
MDV-B-PB1	:	GGGCCTAACATTACAATTGAGAAACTGCATATCCAGAAATAGTATT	: 1850
	1860	* 1880	* 1900
pAB121-PB1	:	:
MDV-B-PB1	:	AAAGTACAACTTAATGGACCTGAATACAAAGGGCGGTTACTGCATCCTC	: 1900
	* 1920	* 1940	* 1950
pAB121-PB1	:	:
MDV-B-PB1	:	AAAATCCCTTGAGGACATTGTCTATTGAGGGCATCAAAGAGGCAGAT	: 1950
	1960	* 1980	* 2000
pAB121-PB1	:	:
MDV-B-PB1	:	ATAACCCCAGCACATGGCCAGTAAAGAAAATGGACTATGATGCGGTATC	: 2000
	* 2020	* 2040	* 2050
pAB121-PB1	:	:
MDV-B-PB1	:	TGGAACTCATAGTGGAGAACCAAAAGGAACAGATCTATACTAAACACTG	: 2050
	2060	* 2080	* 2100
pAB121-PB1	:	:
MDV-B-PB1	:	ATCAGAGGAACATGATTCTGAGGAACAATGCTACGCTAAGTGTGCAAC	: 2100
	* 2120	* 2140	* 2150
pAB121-PB1	:	:
MDV-B-PB1	:	CTTTTGAGGCCTGTTAACAGTGCATCATAAGGAAACCAGTAGGTCA	: 2150
	2160	* 2180	* 2200

Fig. 7 Cont.

```

pAB121-PB1 : ..... : 2200
MDV-B-PB1  : ..... : 2200
                      GCACAGCATGCTTGAGGCTATGCCACAGATTAGAATGGATGCACGAC

                      *      2220      *      2240      *
pAB121-PB1 : ..... : 2250
MDV-B-PB1  : ..... : 2250
                      TAGATTATGAATCAGGAAGAATGTCAAAGGATGATTTGAGAAAGCAATG

                      2260      *      2280      *      2300
pAB121-PB1 : ..... : 2300
MDV-B-PB1  : ..... : 2300
                      GCTCACCTTGGTGAGATTGGGTACATATAAGCTCGAAGATGTCTATGGG

                      *      2320      *      2340      *
pAB121-PB1 : ..... : 2350
MDV-B-PB1  : ..... : 2350
                      GTTATTGGTCATCATTGAATACATGCGGTACACAAATGATTTAAATGAAA

                      2360
pAB121-PB1 : ..... : 2369
MDV-B-PB1  : ..... : 2369
                      AAAGGCTCGTGTCTACT

```

Fig. 7 Cont.

PB2

	*	20	*	40	*		
pAB122-PB2	:	:	50
MDV-B-PB2	:	:	50
		AGCAGAACGGAGCGTTTCAAGATGACATTGCCAAATTGAATTGTTA					
	60	*	80	*	100		
pAB122-PB2	:	:	100
MDV-B-PB2	:	:	100
		AAACAACTGTTAAGGGACAATGAAGCCAAACGGTATTGAAACAAACAC					
	*	120	*	140	*		
pAB122-PB2	:	:	150
MDV-B-PB2	:	:	150
		GGTAGACCAATATAACATAATAAGAAAATTCAATACATCAAGAATTGAAA					
	160	*	180	*	200		
pAB122-PB2	:	:	200
MDV-B-PB2	:	:	200
		AGAACCCCTCATTAAGGATGAAGTGGGCCATGTGTTCTAATTTCCCTG					
	*	220	*	240	*		
pAB122-PB2	:	:	250
MDV-B-PB2	:	:	250
		GCTCTGACCAAGGGTGATATGGCAAATAGAACATCCCCTTGGAAATACAAGGG					
	260	*	280	*	300		
pAB122-PB2	:	:	300
MDV-B-PB2	:	:	300
		AATAACAACCTAAAACAAATGCTGAAGACATAGGAACCAAGGCCAAATGT					
	*	320	*	340	*		
pAB122-PB2	:	:	350
MDV-B-PB2	:	:	350
		GCTCAATAGCAGCAGTTACCTGGTGGAAATACATATGGACCAATAGGAGAT					
	360	*	380	*	400		
pAB122-PB2	:	:	400
MDV-B-PB2	:	:	400
		ACTGAAGGTTTCGAAAAGGTCTACGAAAGCTTTCTCAGAAAGATGAG					
	*	420	*	440	*		
pAB122-PB2	:	:	450
MDV-B-PB2	:	:	450
		ACTTGACAAATGCCACTTGGGCCGAATACTTTGGCCAGTTGAAAGAG					
	460	*	480	*	500		
pAB122-PB2	:	:	500
MDV-B-PB2	:	:	500
		TGAGAAAAAGGGTACTGCTAACCCCTCTCACCAAGGAAATGCCTCCAGAT					
	*	520	*	540	*		
pAB122-PB2	:	:	550
MDV-B-PB2	:	:	550
		GAAGCGAGCAATGTGATAATGAAATATTGTTCCCTAAAGAACAGGAAT					
	560	*	580	*	600		

Fig. 7 Cont.

Fig. 7 Cont.

	* 1120	* 1140	*	
pAB122-PB2 :	: 1150
MDV-B-PB2 :	: 1150
	GACGGAGAAGAGGAGTTCCATGTAAGATGTGGTAATGCAGGGAAATATT			
	1160	* 1180	* 1200	
pAB122-PB2 :	: 1200
MDV-B-PB2 :	: 1200
	AAAAAAAGAGCAAAATGAGAATGGAAAAACTACTAATAAATTCAAGCAAAA			
	* 1220	* 1240	*	
pAB122-PB2 :	: 1250
MDV-B-PB2 :	: 1250
	AGGAGGACATGAAAGATTTAATAATCTTGTGCATGGTATTTCTCAAGAC			
	1260	* 1280	* 1300	
pAB122-PB2 :	: 1300
MDV-B-PB2 :	: 1300
	ACTAGGATGTTCCAAGGAGTGAGAGGAGAAATAAATTCTTAATCGAGC			
	* 1320	* 1340	*	
pAB122-PB2 :	: 1350
MDV-B-PB2 :	: 1350
	AGGCCAACTTTATCTCAATGTACCAACTCCAGCGATATTTTTGAATA			
	1360	* 1380	* 1400	
pAB122-PB2 :	: 1400
MDV-B-PB2 :	: 1400
	GGAGCAACGACCTTTGATCAATGGGGTATGAGGAATCACCAAAGCA			
	* 1420	* 1440	*	
pAB122-PB2 :	: 1450
MDV-B-PB2 :	: 1450
	AGTGAACATACATGGATAATGAATTAATGAATGCATCTGACTATACGTT			
	1460	* 1480	* 1500	
pAB122-PB2 :	: 1500
MDV-B-PB2 :	: 1500
	GAAAGGGTTGTAGTAACAAAAATGTGATTGACTTAGTTACTG			
	* 1520	* 1540	*	
pAB122-PB2 :	: 1550
MDV-B-PB2 :	: 1550
	AAACAGAAAAAGTATCTATAACAAAAATCTAGTTAATAAAAGGACT			
	1560	* 1580	* 1600	
pAB122-PB2 :	: 1600
MDV-B-PB2 :	: 1600
	GGGAAAGTCATAATGGGGCTAATGACGTAAGTGAATTAGAATCACAGC			
	* 1620	* 1640	*	
pAB122-PB2 :	: 1650
MDV-B-PB2 :	: 1650
	ACAGCTAATGATAACATATGATACACCTAACAGATGTGGAGATGGGAACAA			
	1660	* 1680	* 1700	

Fig. 7 Cont.

Fig. 7 Cont.

	*	2220	*	2240	*	
pAB122-PB2	:	: 2250
MDV-B-PB2	:	: 2250
		AAACATCTTACTTATCAAGGAAAGCCCGTTAAAGTAGTTAAAAGGAAAA				
	2260	*	2280	*	2300	
pAB122-PB2	:	: 2300
MDV-B-PB2	:	: 2300
		GATATAGTGCTTATCCAATGACATTCAAGGAATTAAGAGACAAAGA				
	*	2320	*	2340	*	
pAB122-PB2	:	: 2350
MDV-B-PB2	:	: 2350
		ATGACAGTTGAGTCCATGGGGTGGGCCTTGAGCTAATATAAATTATCCA				
	2360	*	2380	*	*	
pAB122-PB2	:	: 2396
MDV-B-PB2	:	: 2396
		TTAATTCAATAGACACAATTGAGTGAAAAATGCTCGTCTACT				

Fig. 7 Cont.

PA

	*	20	*	40	*		
pAB123-PA	:	:	50
MDV-B-PA	:	:	50
		AGCAGAACGGTGCCTTGATTGCCATAATGGATACTTTATTACAAGA					
	60	*	80	*	100		
pAB123-PA	:	:	100
MDV-B-PA	:	:	100
		AACTTCCAGACTACAATAATAACAAAAGGCCAAAACACAATGGCAGAATT					
	*	120	*	140	*		
pAB123-PA	:	:	150
MDV-B-PA	:	:	150
		TAGTGAAGATCCTGAATTACAACCAGCAATGCTATTCAACATCTGCGTCC					
	160	*	180	*	200		
pAB123-PA	:	:	200
MDV-B-PA	:	:	200
		ATCTGGAGGTCTGCTATGTAATAAGTGTATGAATTCTTGATGAAGAA					
	*	220	*	240	*		
pAB123-PA	:	:	250
MDV-B-PA	:	:	250
		GGAAAAACATATACAGCATTAGAAGGACAAGGAAAAGAACAAAACTTGAG					
	260	*	280	*	300		
pAB123-PA	:	:	300
MDV-B-PA	:	:	300
		ACCACAATATGAAGTGATTGAGGGATGCCAAGAACATAGCATGGATGG					
	*	320	*	340	*		
pAB123-PA	:	:	350
MDV-B-PA	:	:	350
		TTCAAAGATCCTAGGCCAAGAGCATGGAATAGAGACTCCAAGGTATCTG					
	360	*	380	*	400		
pAB123-PA	:	:	400
MDV-B-PA	:	:	400
		GCTGATTGTTGATTATAAACCAAGAGGTTATAGAAGTTGGAATAAC					
	*	420	*	440	*		
pAB123-PA	:	:	450
MDV-B-PA	:	:	450
		AAAGGGATTGGCTGACGATTACTTTGGAAAAAGAAAAGCTGGGA					
	460	*	480	*	500		
pAB123-PA	:	:	500
MDV-B-PA	:	:	500
		ATAGCATGGAACTGATGATATTCTAGCTACAATCAAGACTATTGTTAAGT					
	*	520	*	540	*		
pAB123-PA	:	:	550
MDV-B-PA	:	:	550
		AATGAATCCTCATGGATGAGGAAGGAAAGGGAGAGTGCTAACGAGACT					
	560	*	580	*	600		
pAB123-PA	:	:	600

Fig. 7 Cont.

Fig. 7 Cont.

	* 1120		* 1140		*
pAB123-PA :		: 1150
MDV-B-PA :		: 1150
	AAAACCAATTATGCCAAGTGGGCCACAGGAGATGGATTAACATACCAAGAA				
	1160		* 1180		* 1200
pAB123-PA :		: 1200
MDV-B-PA :		: 1200
	AATAATGAAAGAAGTAGCAATAGATGACGAAACAATGTACCAAGAAGAGC				
	* 1220		* 1240		*
pAB123-PA :		: 1250
MDV-B-PA :		: 1250
	CCAAAATACCTAACAAATGTAGAGTGGCTGCTGGTTAACACAGAGATG				
	1260		* 1280		* 1300
pAB123-PA :		: 1300
MDV-B-PA :		: 1300
	AATCTATTGAGCACTCTGACAAGTAAAGGGCCCTGGATCTACCAGAAAT				
	* 1320		* 1340		*
pAB123-PA :		: 1350
MDV-B-PA :		: 1350
	AGGGCCAGACGTAGCACCATGGAGCATGTAGGGAGTGAAAGAAGGAAAT				
	1360		* 1380		* 1400
pAB123-PA :		: 1400
MDV-B-PA :		: 1400
	ACTTTGTTAATGAAATCAACTACTGTAAGGCCTCTACCGTTATGATGAAG				
	* 1420		* 1440		*
pAB123-PA :		: 1450
MDV-B-PA :		: 1450
	TATGTACTTTTCACACTTCATTATTAAATGAAAGCAATGCCAGCATGGG				
	1460		* 1480		* 1500
pAB123-PA :		: 1500
MDV-B-PA :		: 1500
	AAAATATAAAGTAATACCAATAACCAACAGAGTAGTAAATGAAAAAGGAG				
	* 1520		* 1540		*
pAB123-PA :		: 1550
MDV-B-PA :		: 1550
	AAAGTTTGACATGCTCATGGCTGGCGGTTAAAGGGCAATCTCATCTG				
	1560		* 1580		* 1600
pAB123-PA :		: 1600
MDV-B-PA :		: 1600
	AGGGGAGATACTGATGTTAACAGAGTGTGACTTCGAATTAGTAGTAC				
	* 1620		* 1640		*
pAB123-PA :		: 1650
MDV-B-PA :		: 1650
	AGATCCCAGAGTGGACTCAGGAAAGTGGCCAAATATACTGTATTTAGAA				
	1660		* 1680		* 1700

Fig. 7 Cont.

Fig. 7 Cont.

	*	2220		*	2240		*	
pAB123-PA :	: 2250
MDV-B-PA :	: 2250
	GGATGAATGAAAGAAGGGCATAGCGCTCAATTGGTACTATTTGTTCAT							
	2260		*	2280		*	2300	
pAB123-PA :	: 2300
MDV-B-PA :	: 2300
	TATGTATCTAACATCCAATAAAAGAATTGAGAATTAAAAATGCACGTG							
pAB123-PA :	: 2308
MDV-B-PA :	: 2308
	TTTCTACT							

Fig. 7 Cont.

HA

	*	20	*	40	*		
MDV-B-HA	:	:	50
pAB124-HA	:	:	50
	AGCAGAAGCAGAGCATTTCTAATATCCACAAAATGAAGGCAATAATTGT						
	60	*	80	*	100		
MDV-B-HA	:	:	100
pAB124-HA	:	:	100
	ACTACTCATGGTAGTAACATCCAATGCAGATCGAATCTGCACTGGGATAA						
	*	120	*	140	*		
MDV-B-HA	:	t	150
pAB124-HA	:	:	150
	CATCGTCAAACTCACCCATGTGGTCAAAACTGCTACTCAAGGGAAAGTC						
	160	*	180	*	200		
MDV-B-HA	:	..t	:	200
pAB124-HA	:	:	200
	AACGTGACTGGTGTGATACCACTGACAACACACCTACCAAATCTCATT						
	*	220	*	240	*		
MDV-B-HA	:	:	250
pAB124-HA	:	:	250
	TGCAAATCTCAAAGGAACACAGACCAGAGGGAAACTATGCCAAACTGTC						
	260	*	280	*	300		
MDV-B-HA	:	:	300
pAB124-HA	:	:	300
	TCAACTGCACAGATCTGGACGTGGCCTGGGCAGACCAAAGTGTATGGGG						
	*	320	*	340	*		
MDV-B-HA	:	:	350
pAB124-HA	:	:	350
	ACCATACTTCGGCAAAAGCTCAATACTCCACGAAGTCAAACCTGTTAC						
	360	*	380	*	400		
MDV-B-HA	:	:	400
pAB124-HA	:	:	400
	ATCTGGGTGTTTCTATAATGCACGACAGAACAAAAATCAGACAGCTAC						
	*	420	*	440	*		
MDV-B-HA	:	:	450
pAB124-HA	:	:	450
	CCAATCTTCTCAGAGGATATGAAAATATCAGGTTATCAGCCGTAACGTT						
	460	*	480	*	500		
MDV-B-HA	:	:	500
pAB124-HA	:	:	500
	ATCAACCGCAGAACGGCACCAAGGAGGACCCATAGTTGGAACCTCAGG						
	*	520	*	540	*		
MDV-B-HA	:	:	550
pAB124-HA	:	:	550
	ATCTTGCCCTAACGTTACCAATGGAAAGGATTCTCGCAACAATGGCTT						
	560	*	580	*	600		
MDV-B-HA	:	:	600

Fig. 7 Cont.

Fig. 7 Cont.

	* 1120	* 1140	*	
MDV-B-HA :	: 1150
pAB124-HA :	: 1150
	AAAACATTTAAAGGAAAGGGGTTCTCGGAGCTATTGCTGGTTCTGG			
	1160	* 1180	* 1200	
MDV-B-HA :	: 1200
pAB124-HA :	: 1200
	AAGGAGGATGGGAAGGAATGATTGCAGGTTGGCACGGATAACACATCTCAT			
	* 1220	* 1240	*	
MDV-B-HA :	: 1250
pAB124-HA :	: 1250
	GGAGCACATGGAGTGGCAGTGGCAGCAGACCTTAAGAGTACGCAAGAAC			
	1260	* 1280	* 1300	
MDV-B-HA :	: 1300
pAB124-HA :	: 1300
	TATAAACAAAGATAACAAAAATCTCAATTCTTAAGTGAGCTAGAAGTAA			
	* 1320	* 1340	*	
MDV-B-HA :	: 1350
pAB124-HA :	: 1350
	AGAATCTCAAAGACTAAGCGGTGCAATGGATGAACCTCCACAAACGAAATA			
	1360	* 1380	* 1400	
MDV-B-HA :	: 1400
pAB124-HA :	: 1400
	CTCGAGCTGGATGAGAAAGTGGATGATCTCAGAGCTGATAACAATAAGCTC			
	* 1420	* 1440	*	
MDV-B-HA :	: 1450
pAB124-HA :	: 1450
	GCAAATAGAGCTTGCAGTCTGCTTCAACGAAGGAATAATAAACAGTG			
	1460	* 1480	* 1500	
MDV-B-HA :	: 1500
pAB124-HA :	: 1500
	AAGATGAGCATCTCTGGCACTTGAAAGAAAATGAAGAAAATGCTGGC			
	* 1520	* 1540	*	
MDV-B-HA :	: 1550
pAB124-HA :	: 1550
	CCCTCTGCTGTAGACATAGGGATGGATGCTTCGAAACCAAACACAAATG			
	1560	* 1580	* 1600	
MDV-B-HA :	: 1600
pAB124-HA :	: 1600
	CAACCAGACTTGCCTAGACAGGATAGCTGCTGGCACCTTAATGCAGGAG			
	* 1620	* 1640	*	
MDV-B-HA :	: 1650
pAB124-HA :	: 1650
	AATTTCTCTCCACTTTGATTCACTAAATATTACTGCTGCATCTTA			
	1660	* 1680	* 1700	

Fig. 7 Cont.

Fig. 7 Cont.

NP	10	20	30	40	50	
pAB125-NP :	: 50
MDV-B-NP :	: 50
	AGCAGAACGACAGCATTTCTTGTGAACCTCAAGTACCAACAAAAACTGA					
	60	70	80	90	100	
pAB125-NP :	: 100
MDV-B-NP :	: 100
	AAATCAAAATGTCCAACATGGATATTGACGGCATCAACACTGGAACAATT					
	110	120	130	140	150	
pAB125-NP :	: 150
MDV-B-NP :	: 150
	GACAAAACACCAGAACAGAACATAACTTCCGGAACCAAGTGGGGCAACCAGACC					
	160	170	180	190	200	
pAB125-NP :	: 200
MDV-B-NP :	: 200
	AATCATCAAACCAGAACCCCTGCCCCACCAAGCAACAAACGAACCCGAA					
	210	220	230	240	250	
pAB125-NP :	: 250
MDV-B-NP :	: 250
	ACCCATCCCCGGAAAGGGCAGGCCACAAGCAGTGAAGCTGATGTCGGAAGG					
	260	270	280	290	300	
pAB125-NP :	: 300
MDV-B-NP :	: 300
	AGAACCCAAAAGAACAAACCCCCGACAGAGATAAACAGAGAGCTACAA					
	310	320	330	340	350	
pAB125-NP :	: 350
MDV-B-NP :	: 350
	TATGGTAGTGAAACTGGGTGAATTCTACAACCAGATGATGGTCAAAGCTG					
	360	370	380	390	400	
pAB125-NP :	: 400
MDV-B-NP :	: 400
	GACTCAACGATGACATGGAGAGAACCTAATCCAAATGCACATGCTGCG					
	410	420	430	440	450	
pAB125-NP :	: 450
MDV-B-NP :	: 450
	GAAAGAATTCTATTGGCTGCTACTGATGACAAGAAAAGTGAATTCCAAA					
	460	470	480	490	500	
pAB125-NP :	: 500
MDV-B-NP :	: 500
	GAAAAAGAATGCCAGAGATGTCAAAGAAGGGAAAGAAGAAATAGACCACA					
	510	520	530	540	550	
pAB125-NP :	: 550
MDV-B-NP :	: 550
	ACAAAACAGGAGGCACCTTTACAAGATGGTAAGAGATGATAAAACCATC					
	560	570	580	590	600	
pAB125-NP :	: 600

Fig. 7 Cont.

Fig 7. Cont.

	1110	1120	1130	1140	1150	
pAB125-NP	:	: 1150
MDV-B-NP	:	: 1150
	ATGTTGAAGAATACTCTATGGTGGTATGAAGCCATGGCTTTATAAT					
	1160	1170	1180	1190	1200	
pAB125-NP	:	: 1200
MDV-B-NP	:	: 1200
	ATGGCAACACCTGTTCCATATTAAGAATGGGAGACGATGCAAAAGATAA					
	1210	1220	1230	1240	1250	
pAB125-NP	:	: 1250
MDV-B-NP	:	: 1250
	ATCACAAATTATTCTTCATGTCTGCTTCGGAGCTGCCTATGAAGACCTAA					
	1260	1270	1280	1290	1300	
pAB125-NP	:	: 1300
MDV-B-NP	:	: 1300
	GAGTTTGTCTGCACTAACAGGCACAGAACATCAAGCATAGGTCAAGCATTAA					
	1310	1320	1330	1340	1350	
pAB125-NP	:	: 1350
MDV-B-NP	:	: 1350
	AAGTGCAAGGGTTCCACGTTCCAGCAAAGGAGCAAGTGGAAAGGAATGGG					
	1360	1370	1380	1390	1400	
pAB125-NP	:	: 1400
MDV-B-NP	:	: 1400
	GGCAGCTCTGATGTCCATCAAGCTCCAGTTGGCTCCAATGACCAGAT					
	1410	1420	1430	1440	1450	
pAB125-NP	:	: 1450
MDV-B-NP	:	: 1450
	CTGGGGGAATGAAGTAGGTGGAGACGGAGGGTCTGGTCAAATAAGTTGC					
	1460	1470	1480	1490	1500	
pAB125-NP	:	: 1500
MDV-B-NP	:	: 1500
	AGCCCCGTGTTGCAGTAGAAAGACCTATTGCTCTAACAGCAAGCTGT					
	1510	1520	1530	1540	1550	
pAB125-NP	:	: 1550
MDV-B-NP	:	: 1550
	AAGAAGAATGCTGTCAATGAATATTGAGGGACGTGATGCAGATGTCAAAG					
	1560	1570	1580	1590	1600	
pAB125-NP	:	: 1600
MDV-B-NP	:	: 1600
	GAAATCTACTCAAGATGATGAATGATTCAATGACTAACAAAACCAATGGA					
	1610	1620	1630	1640	1650	
pAB125-NP	:	: 1650
MDV-B-NP	:	: 1650
	AATGCTTCATTGGGAAGAAAATGTTCAAATATCAGACAAAAACAAAC					
	1660	1670	1680	1690	1700	
pAB125-NP	:	: 1700

Fig 7. Cont.

MDV-B-NP : : 1700
CAATCCCATGAGATTCCAATTAAAGCAGACCATCCCCAATTCTCTTTG

1710 1720 1730 1740 1750 : 1750
pAB125-NP : : 1750
MDV-B-NP : : 1750
GGAGGGACACAGCAGAGGATTATGATGACCTCGATTATAAGCAACAAA

1760 1770 1780 1790 1800 : 1800
pAB125-NP : : 1800
MDV-B-NP : : 1800
ATAGACACTATGGCTGTGACTGTTCACTACGTTGGAATGTGGGTGTT

1810 1820 1830 1840 1850 : 1842
pAB125-NP : : 1842
MDV-B-NP : : 1842
ACTTTTATTGAAATAAATGTAAAAAATGCTGTTGTTCTACT

pAB125-NP : ----- : -
MDV-B-NP : ----- : -

Fig. 7 Cont.

NA

	* 20	* 40	* 50	
pAB126-NA :	: 50
MDV-B-NA :	: 50
	AGCAGAAGCAGAGCATCTCTCAAAACTGAAGCAAATAGGCCAAAATGA			
	60	* 80	* 100	
pAB126-NA :	: 100
MDV-B-NA :	: 100
	ACAATGCTACCTTCAACTATAACAAACGTTAACCTATTCTCACATCAGG			
	* 120	* 140	* 150	
pAB126-NA :	: 150
MDV-B-NA :	: 150
	GGGAGTGTATTATCACTATATGTGTCAGCTTCACTGTCATACTATTGT			
	160	* 180	* 200	
pAB126-NA :	: 200
MDV-B-NA :	: 200
	ATTCGGATATATTGCTAAAATTTCACCAACAAAAATAACTGCACCAACA			
	* 220	* 240	* 250	
pAB126-NA :	: 250
MDV-B-NA :	: 250
	ATGTCATTGGATTGCGCGAACGTATCAAATGTTAGGCTGTGAACCGTTC			
	260	* 280	* 300	
pAB126-NA :	: 300
MDV-B-NA :	: 300
	TGCAACAAAAGAGATGACATTCTCTCCAGAGCCGGAGTGGACATACC			
	* 320	* 340	* 350	
pAB126-NA :	: 350
MDV-B-NA :	: 350
	CTCGTTATCTGCCAGGGCTAACCTTCAGAAAGCACTCCTAATTAGC			
	360	* 380	* 400	
pAB126-NA :	: 400
MDV-B-NA :	: 400
	CCTCATAGGTCGGAGAACCAAGAGGAAACTCAGCTCCCTGATAATAAG			
	* 420	* 440	* 450	
pAB126-NA :	: 450
MDV-B-NA :	: 450
	GGAACCCTTGTTGCTGTGGACCAAAGGAATGCAGACACTTGTCTAA			
	460	* 480	* 500	
pAB126-NA :	: 500
MDV-B-NA :	: 500
	CCCATTATGCGCTAACCAAGGGGGATACTACAATGGAACAAGAAAGGAC			
	* 520	* 540	* 550	
pAB126-NA :	: 550
MDV-B-NA :	: 550
	AGAAACAAAGCTGAGGCATCTGATTCAGTCAAATTAGGCAAATCCAAC			
	560	* 580	* 600	

Fig. 7 Cont.

Fig. 7 Cont.

	* 1120		* 1140		*
pAB126-NA :		: 1150
MDV-B-NA :		: 1150
	GGAGGATTGTCCATCAAAGAATGGCATCTAAGATTGGAAGATGGTACTC				
	1160		* 1180		* 1200
pAB126-NA :		: 1200
MDV-B-NA :		: 1200
	CCGAACGATGTCTAAACTGAAAGAATGGGATGGAACGTATGTCAAGT				
	* 1220		* 1240		*
pAB126-NA :		: 1250
MDV-B-NA :		: 1250
	ATGATGGAGACCCATGGACTGACAGTGACGCCCTGCTCTAGTGGAGTA				
	1260		* 1280		* 1300
pAB126-NA :		: 1300
MDV-B-NA :		: 1300
	ATGGTTCAATGAAAGAACCTGGTGGTATTCTTGGCTTCGAAATAAA				
	* 1320		* 1340		*
pAB126-NA :		: 1350
MDV-B-NA :		: 1350
	AGATAAGAAATGTGATGTCCCCGTATTGGGATAGAGATGGTACACGATG				
	1360		* 1380		* 1400
pAB126-NA :		: 1400
MDV-B-NA :		: 1400
	GTGGAAAAGAGACTTGGCACTCAGCAGAACAGCCATTACTGTTGATG				
	* 1420		* 1440		*
pAB126-NA :		: 1450
MDV-B-NA :		: 1450
	GGCTCAGGACAATTGCTATGGGACACTGTCACAGGTGTTGATATGGCTCT				
	1460		* 1480		* 1500
pAB126-NA :		: 1500
MDV-B-NA :		: 1500
	GTAATGGAGGAATGGTGAATCTGTTCAAACCTTGTCTATTGT				
	* 1520		* 1540		*
pAB126-NA :		: 1550
MDV-B-NA :		: 1550
	TTGAACAAATTGTCCTACTGGACTTAATTGTTCTGAAAATGCTCTGT				
pAB126-NA :				: 1557
MDV-B-NA :				: 1557
	TACTACT				

Fig 7. Cont.

M

	*	20	*	40	*		
pAB127-M	:	:	50
MDV-B-M	:	:	50
		AGCAGAACGCACGCAC	TTCTTAAATGTCGCTGTTGGAGACACAATTGC				
		60	*	80	*	100	
pAB127-M	:	:	100
MDV-B-M	:	:	100
		CTACCTGCTTCAC	TAACAGAACATGGAGAAGGCAAAGCAGAACTAGCAG				
	*	120	*	140	*		
pAB127-M	:	:	150
MDV-B-M	:	:	150
		AAAAATTACACTGTTGGTCGGTGGAAAGAATTGACCTAGACTCTGCT					
	160	*	180	*	200		
pAB127-M	:	:	200
MDV-B-M	:	:	200
		TTGGAATGGATAAAAAACAAAAGATGCCTA	ACTGATATAACAAAAGCACT				
	*	220	*	240	*		
pAB127-M	:	:	250
MDV-B-M	:	:	250
		AATTGGTGCCTCTATCTGCTTTAAACCCAAAGACCAAGAAAGAAAAAA					
	260	*	280	*	300		
pAB127-M	:	:	300
MDV-B-M	:	:	300
		GAAGATTCATCACAGAGCCCTGTCAGGAATGGGAACAACAGCAACAAAAA					
	*	320	*	340	*		
pAB127-M	:	:	350
MDV-B-M	:	:	350
		AAGAAAGGCCTGATTCTAGCTGAGAGAAAATGAGAACATGTGTGAGTTT					
	360	*	380	*	400		
pAB127-M	:	:	400
MDV-B-M	:	:	400
		TCATGAAGCATTGAAATAGCAGAAGGCCATGAAAGCTCAGCACTACTAT					
	*	420	*	440	*		
pAB127-M	:	:	450
MDV-B-M	:	:	450
		ATTGTCTCATGGTCATGTACCTGAACCTGGAAATTATTCAATGCAAGTA					
	460	*	480	*	500		
pAB127-M	:	:	500
MDV-B-M	:	:	500
		AAACTAGGAACGCTCTGTGCTTATGCGAGAACACAAGCATCACATTACA					
	*	520	*	540	*		
pAB127-M	:	:	550
MDV-B-M	:	:	550
		AAGAGCTCATAGCAGAGCAGCAAGATCTCAGTGCCTGGAGTGAGGCGAG					
	560	*	580	*	600		
pAB127-M	:	:	600

Fig. 7 Cont.

Fig. 7 Cont.

	*	1120	*	1140	*	
pAB127-M	:	: 1150
MDV-B-M	:	: 1150
		ACCCAATTTACCGTATTCTTGCTATGCATTAAAGCAAATTGTAATCA				
	1160		*	1180	*	
pAB127-M	:	: 1190
MDV-B-M	:	: 1190
		ATGTCAGCAAATAAACTGGAAAAAGTGCCTGTTCTACT				

Fig. 7 Cont.

NS

	10	20	30	40	50	
pAB128-NS	:	:
MDV-B-NS	:	:
	AGCAGAACGAGGAGTTGTTAGTCAGTGGCAAACGGAAAAAAATGGCG					
	60	70	80	90	100	
pAB128-NS	:	:
MDV-B-NS	:	:
	GACAACATGACCACAAACACAAATTGAGGTAGGTCCGGGAGCAACCAATGC					
	110	120	130	140	150	
pAB128-NS	:	:
MDV-B-NS	:	:
	CACCATAAACTTGAAGCAGGAATTCTGGAGTGCTATGAAAGGCTTCAT					
	160	170	180	190	200	
pAB128-NS	:	:
MDV-B-NS	:	:
	GGCAAAGAGCCCTTGACTACCCCTGGTCAAGACCCCTAACAGACTAAAG					
	210	220	230	240	250	
pAB128-NS	:	:
MDV-B-NS	:	:
	AGAAAATTAGAATCAAGAATAAGACTCACACACAAAGTGAGCCTGAAAG					
	260	270	280	290	300	
pAB128-NS	:	:
MDV-B-NS	:	:
	TAAAAGGATGTCTCTTGAAGAGAGAAAAGCAATTGGGTAAAAATGATGA					
	310	320	330	340	350	
pAB128-NS	:	:
MDV-B-NS	:	:
	AAGTGCTCCTATTATGAATCCATCTGCTGGATTGAAGGGTTGAGCCA					
	360	370	380	390	400	
pAB128-NS	:	:
MDV-B-NS	:	:
	TACTGTATGAAAATTCCCTCAAATAGCAACTGTCCAAACTGCAATTGGAC					
	410	420	430	440	450	
pAB128-NS	:	G.....	:
MDV-B-NS	:	:
	CGATTACCTCCAACACCCAGGAAAGTGCCTTGATGACATAGAAGAAGAAC					
	460	470	480	490	500	
pAB128-NS	:	:
MDV-B-NS	:	:
	CGGAGAATGTTGATGACCAACTGAAATAGTATTGAGGGACATGAACAAAC					
	510	520	530	540	550	
pAB128-NS	:	:
MDV-B-NS	:	:
	AAAGATGCAAGGCAAAAGATAAAGGAGGAAGTAAACACTCAGAAAGAAGG					
	560	570	580	590	600	
pAB128-NS	:	:

Fig. 7 Cont.

Fig. 7 Cont.



Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12A




Fig. 12B

PA	NP	M1	MDCK				PCK			
			log	37°C	37°C	Δlog	log	37°C	37°C	Δlog
431	497		55	114	410	509	159	183		
M H		A A H T		Q V		ts	6.6	<2	>3	5.6
V Y		T V P A	H	M	non-ts	7.6	6.6	1.0	8.1	7.4
V Y		A V P A	H	M	non-ts	7.6	7.1	0.5	7.4	6.5
V Y		A V P A	H	M	non-ts	8.1	7.1	1.0	7.7	6.5
M H		A A H T		Q V		ts	7.1	4.0	7.1	3.5
V Y		T V P A	H	M	non-ts	8.1	7.1	1.0	8.7	7.8
V Y		A V P A	H	M	non-ts	8.1	7.2	0.9	8.5	7.8

Fig. 13

PA	NP	M1				MDCK log pfu/ml	PCK log TCID ₅₀ /ml	
		33°C	37°C	Δlog	33°C			
M H	AA HT	H M	ts	7.1	3.2	3.9	6.2	3.3 2.9
M H	AV PA	Q V	ts	n.d.			5.8	2.9 2.9
V Y	AA HT	Q V	ts	6.2	3.2	3.0	6.1	2.7 3.4
V Y	AA HT	H M	ts	7.4	4.4	3.0	7.5	3.4 4.1
V Y	AA HT	H M	ts	7.6	4.2	3.4	8.3	4.3 4.0
M H	AV PA	H M	ts	7.4	4.4	3.0	8.1	4.3 3.8
M H	TV PA	H M	ts	8.0	6.0	2.0	8.4	4.3 4.1
V Y	TV PA	Q V	non-ts	5.6	6.0	-0.4	6.4	4.5 1.9
V Y	TV PA	Q V	non-ts	6.6	5.8	0.8	6.8	4.8 2.0

Fig. 14

PA	NP	M1	MDCK			PCK		
			log pfu/ml	37°C	Δlog	log pfu/ml	37°C	Δlog
431	497	55 114 410 509	159 183					
V Y	A V P T	Q V	non-ts	6.2	5.2 1.0	6.8	5.5	1.4
V Y	A A P T	Q V	non-ts	6.8	6.4 0.4	7.2	6.1	1.1
V Y	A A P T	Q V	non-ts	6.4	6.2 0.2	7.1	5.7	1.4
V Y	T A H T	Q V	ts	6.6	4.4 2.2	6.6	3.4	3.2
V Y	A A P T	H M	non-ts	7.4	6.8 0.6	8.3	7.0	1.3
V Y	T A P T	H M	non-ts	n.d.		8.0	7.2	0.8

Fig. 15




Fig. 16




Fig. 17




Fig. 18




Fig. 19




Fig. 20A




Fig. 20B





Fig. 21

Antigenically Drifted H3N2 strains Caused Influenza Outbreak in 2003-2004 Season



Flu season started uncommonly early, from October 2003

Mortality: 10.3% P & I in USA

144 Flu associated deaths in 18 years and younger

Influenza A predominant (99.4%), mostly H3N2 strains
Flu Shot was not effective (MMWR January 16, 2004)

Fig. 22




Fig. 23

Amino Acid Difference Between HAs of A/Panama/99 and A/Fujian/02





Fig. 24

卷之二

Site E

Site D
W202D
S219F/
W222R
C225D
V226I

Site C


ste B

e A

Sit
m

A/Panai
a/2007/9

Generation of Cold-adapted Live Attenuated Influenza Vaccines by Plasmids



ca A/AA/6/60 (MDV-A) *ts* and *att* loci :
PB1: K391E, E581G and G661T
PB2: N265S
NP: D34G

Molecular Basis of Antigenic Drift of Epidemic A/Fujian/02-like Viruses

Ag site	21	25	50	75	83	128	131	155,6	183	186	202	219	222,5,6	D	Egg growth
A/Panama/99	S	L	R	H	E	T	A	H	Q	L	S	V	S	WGV	Yes
A/Wyoming/03	P	I	G	Q	K	A	T	T	H	H	V	I	Y	RDI	Yes
• Flu274				G	Q	K			T						Yes
• Flu275									T	T	H			RDI	Yes
• Flu276									T	T	H			RDI	Yes
• Flu277									T						Yes
• Flu278														RDI	Yes
• Flu279										H	V				No Yes (HA-V182F)
• Flu280										T	H				No Yes (HA-P185L)

Fig. 26

Immunogenicity of Antigenically Modified Mutants




Fig. 27

Anti-A/Wyoming/03 ferret serum

Minimal Genetic Change for Antigenic Drift of Epidemic H3N2 Strains






Fig. 28


Features of A/Fujian/411/2002-like Strains

Virus strain	A/Sendai H/F4962/02	A/Fujian /411/02	A/Wyoming ⁽¹⁾ /03/03
Isolated	12/4/02	8/11/02	2/13/04
Passages	CXE8/E2	C1/C1	spfPCK2E2/E8
Egg Growth	Yes	No	Yes
Ferret Nose	Poor	ND	Good
HA	128 186 219 226	 	 
NA	119 136 347	 	 

⁽¹⁾ Vaccine strain for 2004-2005
⁽²⁾ Neuraminidase inhibitors resistant site.

Fig. 29.

Fig. 30 NA Neuraminidase Activities and Virus Replication



119E and 136Q are Critical for NA Activity




Fig. 31

Effect of HA Residues on Virus Replication in Eggs

HA		NA		Wy-NA 119E/136Q /347H		119E/136Q	
128	186	219	226				
T	G	S	V	<1.5	<1.5		
T	V	S	V	4.95	4.39		
T	G	S	V	5.20	3.85		
T	V	S	V	7.38	7.30		
T	V	Y	V	7.40	7.40		
A	V	Y	V	7.75	7.18		

Four residues: HA-186V, 226I and HA-119E, 136Q are sufficient to restore virus replication in eggs.

Fig. 32

Adaptation of rA/Fujian/02 in Eggs





Fig. 33

HA Receptor-Binding Sites



Ha et al. Virology 309: 209-218, 2003

Fig. 34

Balance Between HA and NA Activities is
Critical for Influenza Virus Replication




Fig. 35




Fig. 36




Fig. 37




Fig. 38

MULTI PLASMID SYSTEM FOR THE PRODUCTION OF INFLUENZA VIRUS

[0001] This application claims the benefit under 35 U.S.C. § 119 (e) of U.S. Provisional Application No. 60/532,164 filed Dec. 23, 2003, which is incorporated by reference herein.

BACKGROUND OF THE INVENTION

[0002] Influenza viruses are made up of an internal ribonucleoprotein core containing a segmented single-stranded RNA genome and an outer lipoprotein envelope lined by a matrix protein. Influenza A and B viruses each contain eight segments of single stranded RNA with negative polarity. The influenza A genome encodes at least eleven polypeptides. Segments 1-3 encode the three polypeptides, making up the viral RNA-dependent RNA polymerase. Segment 1 encodes the polymerase complex protein PB2. The remaining polymerase proteins PB1 and PA are encoded by segment 2 and segment 3, respectively. In addition, segment 1 of some influenza A strains encodes a small protein, PB1-F2, produced from an alternative reading frame within the PB1 coding region. Segment 4 encodes the hemagglutinin (HA) surface glycoprotein involved in cell attachment and entry during infection. Segment 5 encodes the nucleocapsid nucleoprotein (NP) polypeptide, the major structural component associated with viral RNA. Segment 6 encodes a neuraminidase (NA) envelope glycoprotein. Segment 7 encodes two matrix proteins, designated M1 and M2, which are translated from differentially spliced mRNAs. Segment 8 encodes NS1 and NS2 (NEP), two nonstructural proteins, which are translated from alternatively spliced mRNA variants.

[0003] The eight genome segments of influenza B encode 11 proteins. The three largest genes code for components of the RNA polymerase, PB1, PB2 and PA. Segment 4 encodes the HA protein. Segment 5 encodes NP. Segment 6 encodes the NA protein and the NB protein. Both proteins, NB and NA, are translated from overlapping reading frames of a bicistronic mRNA. Segment 7 of influenza B also encodes two proteins: M1 and BM2. The smallest segment encodes two products: NS1 is translated from the full length RNA, while NS2 is translated from a spliced mRNA variant.

[0004] Vaccines capable of producing a protective immune response specific for influenza viruses have been produced for over 50 years. Vaccines can be characterized as whole virus vaccines, split virus vaccines, surface antigen vaccines and live attenuated virus vaccines. While appropriate formulations of any of these vaccine types is able to produce a systemic immune response, live attenuated virus vaccines are also able to stimulate local mucosal immunity in the respiratory tract.

[0005] FluMist™ is a live, attenuated vaccine that protects children and adults from influenza illness (Belshe et al. (1998) The efficacy of live attenuated, cold-adapted, trivalent, intranasal influenza virus vaccine in children *N Engl J Med* 338:1405-12; Nichol et al. (1999) Effectiveness of live, attenuated intranasal influenza virus vaccine in healthy, working adults: a randomized controlled trial *JAMA* 282:137-44). FluMist™ vaccine strains contain HA and NA gene segments derived from the currently circulating wild-type strains along with six gene segments, PB1, PB2, PA, NP, M and NS, from a common master donor virus (MDV). The MDV for influenza A strains of FluMist (MDV-A), was

created by serial passage of the wt A/Ann Arbor/6/60 (A/AA/6/60) strain in primary chicken kidney tissue culture at successively lower temperatures (Maassab (1967) Adaptation and growth characteristics of influenza virus at 25 degrees C. *Nature* 213:612-4). MDV-A replicates efficiently at 25° C. (ca, cold adapted), but its growth is restricted at 38 and 39° C. (ts, temperature sensitive). Additionally, this virus does not replicate in the lungs of infected ferrets (att, attenuation). The ts phenotype is believed to contribute to the attenuation of the vaccine in humans by restricting its replication in all but the coolest regions of the respiratory tract. The stability of this property has been demonstrated in animal models and clinical studies. In contrast to the ts phenotype of influenza strains created by chemical mutagenesis, the ts property of MDV-A did not revert following passage through infected hamsters or in shed isolates from children (for a recent review, see Murphy & Coelingh (2002) Principles underlying the development and use of live attenuated cold-adapted influenza A and B virus vaccines *Viral Immunol* 15:295-323).

[0006] Clinical studies in over 20,000 adults and children involving 12 separate 6:2 reassortant strains have shown that these vaccines are attenuated, safe and efficacious (Belshe et al. (1998) The efficacy of live attenuated, cold-adapted, trivalent, intranasal influenza virus vaccine in children *N Engl J Med* 338:1405-12; Boyce et al. (2000) Safety and immunogenicity of adjuvanted and unadjuvanted subunit influenza vaccines administered intranasally to healthy adults *Vaccine* 19:217-26; Edwards et al. (1994) A randomized controlled trial of cold adapted and inactivated vaccines for the prevention of influenza A disease *J Infect Dis* 169:68-76 ; Nichol et al. (1999) Effectiveness of live, attenuated intranasal influenza virus vaccine in healthy, working adults: a randomized controlled trial *JAMA* 282:137-44). Reassortants carrying the six internal genes of MDV-A and the two HA and NA gene segments of the wt virus (6:2 reassortant) consistently maintain ca, ts and att phenotypes (Maassab et al. (1982) Evaluation of a cold-recombinant influenza virus vaccine in ferrets *J Infect Dis* 146:780-900).

[0007] To date, all commercially available influenza vaccines in the United States have been propagated in embryonated hen's eggs. Although influenza virus grows well in hen's eggs, production of vaccine is dependent on the availability of eggs. Supplies of eggs must be organized, and strains for vaccine production selected months in advance of the next flu season, limiting the flexibility of this approach, and often resulting in delays and shortages in production and distribution. Unfortunately, some influenza vaccine strains, such as the prototype A/Fujian/411/02 strain that circulated during the 2003-04 season, do not replicate well in embryonated chicken eggs, and have to be isolated by cell culture a costly and time consuming procedure. The present invention further provides a new technology to increase the ability of vaccine strains to replicate in embryonated chicken eggs. Furthermore, the present invention allows for more efficient and cost effective production of influenza vaccines.

[0008] Systems for producing influenza viruses in cell culture have also been developed in recent years (See, e.g., Furminger. *Vaccine Production*, in Nicholson et al. (eds) *Textbook of Influenza* pp. 324-332; Merten et al. (1996) Production of influenza virus in cell cultures for vaccine preparation, in Cohen & Shafferman (eds) *Novel Strategies*

in *Design and Production of Vaccines* pp. 141-151). Typically, these methods involve the infection of suitable immortalized host cells with a selected strain of virus. While eliminating many of the difficulties related to vaccine production in hen's eggs, not all pathogenic strains of influenza grow well and can be produced according to established tissue culture methods. In addition, many strains with desirable characteristics, e.g., attenuation, temperature sensitivity and cold adaptation, suitable for production of live attenuated vaccines, have not been successfully grown in tissue culture using established methods.

[0009] Production of influenza viruses from recombinant DNA would significantly increase the flexibility and utility of tissue culture methods for influenza vaccine production. Recently, systems for producing influenza A viruses from recombinant plasmids incorporating cDNAs encoding the viral genome have been reported (See, e.g., Neumann et al. (1999) Generation of influenza A virus entirely from cloned cDNAs. *Proc Natl Acad Sci USA* 96:9345-9350; Fodor et al. (1999) Rescue of influenza A virus from recombinant DNA. *J. Virol.* 73:9679-9682; Hoffmann et al. (2000) A DNA transfection system for generation of influenza A virus from eight plasmids *Proc Natl Acad Sci USA* 97:6108-6113; WO 01/83794). These systems offer the potential to produce recombinant viruses, and reassortant viruses expressing the immunogenic HA and NA proteins from any selected strain. However, unlike influenza A virus, no reports have been published describing plasmid-only systems for influenza B virus.

[0010] Additionally, none of the currently available plasmid only systems are suitable for generating attenuated, temperature sensitive, cold adapted strains suitable for live attenuated vaccine production. The present invention provides an eight plasmid system for the generation of influenza B virus entirely from cloned cDNA, and methods for the production of attenuated live influenza A and B virus suitable for vaccine formulations, such as live virus vaccine formulations useful for intranasal administration, as well as numerous other benefits that will become apparent upon review of the specification.

SUMMARY OF THE INVENTION

[0011] The present invention relates to a multi-vector system for the production of influenza viruses in cell culture, and to methods for producing recombinant and reassortant influenza viruses, including, e.g., attenuated (att), cold adapted (ca) and/or temperature sensitive (ts) influenza viruses, suitable as vaccines, including live attenuated influenza vaccines, such as those suitable for administration in an intranasal vaccine formulation.

[0012] In a first aspect the invention provides vectors and methods for producing recombinant influenza B virus in cell culture, e.g., in the absence of helper virus (i.e., a helper virus free cell culture system). The methods of the invention involve introducing a plurality of vectors, each of which incorporates a portion of an influenza B virus into a population of host cells capable of supporting viral replication. The host cells are cultured under conditions permissive for viral growth, and influenza viruses are recovered. In some embodiments, the influenza B viruses are attenuated viruses, cold adapted viruses and/or temperature sensitive viruses. For example, in an embodiment, the vector-derived recom-

binant influenza B viruses are attenuated, cold adapted, temperature sensitive viruses, such as are suitable for administration as a live attenuated vaccine, e.g., in a intranasal vaccine formulation. In an exemplary embodiment, the viruses are produced by introducing a plurality of vectors incorporating all or part of an influenza B/Ann Arbor/1/66 virus genome, e.g., a ca B/Ann Arbor/1/66 virus genome.

[0013] For example, in some embodiments, the influenza B viruses are artificially engineered influenza viruses incorporating one or more amino acid substitutions which influence the characteristic biological properties of influenza strain ca B/Ann Arbor/1/66. Such influenza viruses include mutations resulting in amino acid substitutions at one or more of positions PB1³⁹¹, PB1⁵⁸¹, PB1⁶⁶¹, PB2²⁶⁵ and NP³⁴, such as: PB1³⁹¹, (K391E), PB1⁵⁸¹ (E581G), PB1⁶⁶¹ (A661T), PB2²⁶⁵ (N265S) and Np34 (D34G). Any mutation (at one or more of these positions) which individually or in combination results in increased temperature sensitivity, cold adaptation or attenuation relative to wild type viruses is a suitable mutation in the context of the present invention.

[0014] In some embodiments, a plurality of vectors incorporating at least the 6 internal genome segments of a one influenza B strain along with one or more genome segments encoding immunogenic influenza surface antigens of a different influenza strain are introduced into a population of host cells. For example, at least the 6 internal genome segments of a selected attenuated, cold adapted and/or temperature sensitive influenza B strain, e.g., a ca, att, ts strain of B/Ann Arbor/1/66 or an artificially engineered influenza B strain including an amino acid substitution at one or more of the positions specified above, are introduced into a population of host cells along with one or more segments encoding immunogenic antigens derived from another virus strain. Typically the immunogenic surface antigens include either or both of the hemagglutinin (HA) and/or neuraminidase (NA) antigens. In embodiments where a single segment encoding an immunogenic surface antigen is introduced, the 7 complementary segments of the selected virus are also introduced into the host cells.

[0015] In certain embodiments, a plurality of plasmid vectors incorporating influenza B virus genome segments are introduced into a population of host cells. For example, 8 plasmids, each of which incorporates a different genome segment are utilized to introduce a complete influenza B genome into the host cells. Alternatively, a greater number of plasmids, incorporating smaller genomic subsequences can be employed.

[0016] Typically, the plasmid vectors of the invention are bi-directional expression vectors. A bi-directional expression vector of the invention typically includes a first promoter and a second promoter, wherein the first and second promoters are operably linked to alternative strands of the same double stranded cDNA encoding the viral nucleic acid including a segment of the influenza virus genome. Optionally, the bi-directional expression vector includes a polyadenylation signal and/or a terminator sequence. For example, the polyadenylation signal and/or the terminator sequence can be located flanking a segment of the influenza virus genome internal to the two promoters. One favorable polyadenylation signal in the context of the invention is the SV40 polyadenylation signal. An exemplary plasmid vector of the invention is the plasmid pAD3000, illustrated in FIG. 1.

[0017] The vectors are introduced into host cells capable of supporting the replication of influenza virus from the vector promoters. Favorable examples of host cells include Vero cells, Per.C6 cells, BHK cells, PCK cells, MDCK cells, MDBK cells, 293 cells (e.g., 293T cells), and COS cells. In combination with the pAD3000 plasmid vectors described herein, Vero cells, 293 cells, and COS cells are particularly suitable. In some embodiments, co-cultures of a mixture of at least two of these cell lines, e.g., a combination of COS and MDCK cells or a combination of 293T and MDCK cells, constitute the population of host cells.

[0018] The host cells including the influenza B vectors are then grown in culture under conditions permissive for replication and assembly of viruses. Typically, host cells incorporating the influenza B plasmids of the invention are cultured at a temperature below 37° C., preferably at a temperature equal to, or less than, 35° C. Typically, the cells are cultured at a temperature between 32° C. and 35° C. In some embodiments, the cells are cultured at a temperature between about 32° C. and 34° C., e.g., at about 33° C. Following culture for a suitable period of time to permit replication of the virus to high titer, recombinant and/or reassortant viruses are recovered. Optionally, the recovered viruses can be inactivated.

[0019] The invention also provides broadly applicable methods of producing recombinant influenza viruses in cell culture by introducing a plurality of vectors incorporating an influenza virus genome into a population of host cells capable of supporting replication of influenza virus, culturing the cells at a temperature less than or equal to 35° C., and recovering influenza viruses.

[0020] In certain embodiments, a plurality of plasmid vectors incorporating influenza virus genome segments are introduced into a population of host cells. In certain embodiments, 8 plasmids, each of which incorporates a different genome segment are utilized to introduce a complete influenza genome into the host cells. Typically, the plasmid vectors of the invention are bi-directional expression vectors. An exemplary plasmid vector of the invention is the plasmid pAD3000, illustrated in **FIG. 1**.

[0021] In some embodiments, the influenza viruses correspond to an influenza B virus. In some embodiments, the influenza viruses correspond to an influenza A virus. In certain embodiments, the methods include recovering recombinant and/or reassortant influenza viruses capable of eliciting an immune response upon administration, e.g., intranasal administration, to a subject. In some embodiments, the viruses are inactivated prior to administration, in other embodiments, live-attenuated viruses are administered. Recombinant and reassortant influenza A and influenza B viruses produced according to the methods of the invention are also a feature of the invention.

[0022] In certain embodiments, the viruses include an attenuated influenza virus, a cold adapted influenza virus, a temperature sensitive influenza virus, or a virus with any combination of these desirable properties. In one embodiment, the influenza virus incorporates an influenza B/Ann Arbor/1/66 strain virus, e.g., a cold adapted, temperature sensitive, attenuated strain of B/Ann Arbor/1/66. In another embodiment, the influenza virus incorporates an influenza A/Ann Arbor/6/60 strain virus, e.g., a cold adapted, temperature sensitive, attenuated strain of A/Ann Arbor/6/60. In

another embodiment of the invention, the viruses are artificially engineered influenza viruses incorporating one or more substituted amino acid which influences the characteristic biological properties of, e.g., ca A/Ann Arbor/6/60 or ca B/Ann Arbor/1/66. Such substituted amino acids favorably correspond to unique amino acids of ca A/Ann Arbor/6/60 or ca B/Ann Arbor/1/66, e.g., in an A strain virus: PB1³⁹¹ (K391E), PB1⁵⁸¹ (E581G), PB1⁶⁶¹ (A661T), PB2²⁶⁵ (N265S) and NP³⁴ (D34G); and, in a B strain virus: PB2⁶³⁰ (S630R); PA⁴³¹ (V431M); PA⁴⁹⁷ (Y497H); NP⁵⁵ (T55A); NP¹¹⁴ (V114A); NP⁴¹⁰ (P410H); NP⁵⁰⁹ (A509T); M1¹⁵⁹ (H159Q) and M1¹⁸³ (M183V). Similarly, other amino acid substitutions at any of these positions resulting in temperature sensitivity, cold adaptation and/or attenuation are encompassed by the viruses and methods of the invention.

[0023] Optionally, reassortant viruses are produced by introducing vectors including the six internal genes of a viral strain selected for its favorable properties regarding vaccine production, in combination with the genome segments encoding the surface antigens (HA and NA) of a selected, e.g., pathogenic strain. For example, the HA segment is favorably selected from a pathogenically relevant H1, H3 or B strain, as is routinely performed for vaccine production. Similarly, the HA segment can be selected from an emerging pathogenic strain such as an H2 strain (e.g., H2N2), an H5 strain (e.g., H5N1) or an H7 strain (e.g., H7N7). Alternatively, the seven complementary gene segments of the first strain are introduced in combination with either the HA or NA encoding segment. In certain embodiments, the internal gene segments are derived from the influenza B/Ann Arbor/1/66 or the A/Ann Arbor/6/60 strain.

[0024] Additionally, the invention provides methods for producing novel influenza viruses with desirable properties relevant to vaccine production, e.g., temperature sensitive, attenuated, and/or cold adapted, influenza viruses, as well as influenza vaccines including such novel influenza viruses. In certain embodiments, novel influenza A strain virus is produced by introducing mutations that result amino acid substitutions at one or more specified positions demonstrated herein to be important for the temperature sensitive phenotype, e.g., PB1³⁹¹, PB1⁵⁸¹, PB1⁶⁶¹, PB2²⁶⁵ and NP³⁴. For example, mutations are introduced at nucleotide positions PB1¹¹⁹⁵, PB1¹⁷⁶⁶, PB1²⁰⁰⁵, PB2⁸²¹ and NP¹⁴⁶, or other nucleotide positions resulting in an amino acid substitution at the specified amino acid position. Any mutation (at one or more of these positions) which individually or in combination results in increased temperature sensitivity, cold adaptation or attenuation relative to wild type viruses is a suitable mutation in the context of the present invention. For example; mutations selected from among PB1³⁹¹ (K391E), PB1⁵⁸¹ (E581G), PB1⁶⁶¹ (A661T), PB2²⁶⁵ (N265S) and NP³⁴ (D34G) are favorably introduced into the genome of a wild type influenza A strain, e.g., PR8, to produce a temperature sensitive variant suitable for administration as a live attenuated vaccine. To increase stability of the desired phenotype, a plurality of mutations are typically introduced. Following introduction of the selected mutation(s) into the influenza genome, the mutated influenza genome is replicated under conditions in which virus is produced. For example, the mutated influenza virus genome can be replicated in hens' eggs. Alternatively, the influenza virus genome can be replicated in cell culture. In the latter case, the virus is optionally further amplified in hens' eggs to increase the titer. Temperature sensitive, and optionally,

attenuated and/or cold adapted viruses produced according to the methods of the invention are also a feature of the invention, as are vaccines including such viruses. Similarly, novel recombinant viral nucleic acids incorporating one or more mutations at positions PB1³⁹¹, PB1⁵⁸¹, PB1⁶⁶¹, PB2²⁶⁵ and NP³⁴, e.g., mutations selected from among PB1³⁹¹ (K391E), PB1⁵⁸¹ (E581G), PB1⁶⁶¹ (A661T), PB2²⁶⁵ (N265S) and NP³⁴ (D34G), and polypeptides with such amino acid substitutions are a feature of the invention.

[0025] Likewise, the methods presented herein are adapted to producing novel influenza B strains with temperature sensitive, and optionally attenuated and/or cold adapted phenotypes by introducing one or more specified mutations into an influenza B genome. For example, one or more mutations resulting in an amino acid substitution at a position selected from among PB2⁶³⁰, PA⁴³¹, PA⁴⁹⁷, NP⁵⁵; NP¹¹⁴, NP⁴¹⁰; NP509; M1¹⁵⁹ and M1¹⁸³ are introduced into an influenza B strain genome to produce a temperature sensitive influenza B virus. Exemplary amino acid substitutions include the following: : PB2⁶³⁰ (S630R); PA⁴³¹ (V431M); PA⁴⁹⁷ (Y497H); NP⁵⁵ (T55A); NP¹¹⁴ (V114A); NP⁴¹⁰ (P410H); NP⁵⁰⁹ (A509T); M1¹⁵⁹ (H159Q) and M1¹⁸³ (M183V). As indicated above, vaccines incorporating such viruses as well as nucleic acids and polypeptides incorporating these mutations and amino acid substitutions are all features of the invention.

[0026] Accordingly, influenza viruses incorporating the mutations of the invention are a feature of the invention regardless of the method in which they are produced. That is, the invention encompasses influenza strains including the mutations of the invention, e.g., any influenza A virus with an amino acid substitution relative to wild type at one or more positions selected from among: PB1³⁹¹, PB1⁵⁸¹, PB1⁶⁶¹, PB2²⁶⁵ and NP³⁴ or any influenza B virus with an amino acid substitution relative to wild type at one or more positions selected from among: PB2⁶³⁰, PA⁴³¹, PA⁴⁹⁷, NP⁵⁵; NP¹¹⁴, NP⁴¹⁰, NP⁵⁰⁹; M1¹⁵⁹ and M1¹⁸³, with the proviso that the strains ca A/Ann Arbor/6/60 and B/Ann Arbor/1/66 are not considered a feature of the present invention. In certain preferred embodiments, the influenza A viruses include a plurality of mutations selected from among PB1³⁹¹ (K391E), PB1⁵⁸¹ (E581G), PB1⁶⁶¹ (A661T), PB2²⁶⁵ (N265S) and NP³⁴ (D34G); and the influenza B viruses include a plurality of mutations selected from among PB2⁶³⁰ (S630R); PA⁴³¹ (V431M); PA⁴⁹⁷ (Y497H); NP⁵⁵ (T55A); NP¹¹⁴ (V114A); NP⁴¹⁰ (P410H); NP⁵⁰⁹ (A509T); M1¹⁵⁹ (H159Q) and M1¹⁸³ (M183V), respectively.

[0027] In one embodiment, a plurality of plasmid vectors incorporating the influenza virus genome are introduced into host cells. For example, segments of an influenza virus genome can be incorporated into at least 8 plasmid vectors. In one preferred embodiment, segments of an influenza virus genome are incorporated into 8 plasmids. For example, each of 8 plasmids can favorably incorporate a different segment of the influenza virus genome.

[0028] The vectors of the invention can be bi-directional expression vectors. A bi-directional expression vector of the invention typically includes a first promoter and a second promoter, wherein the first and second promoters are operably linked to alternative strands of the same double stranded viral nucleic acid including a segment of the influenza virus genome. Optionally, the bi-directional

expression vector includes a polyadenylation signal and/or a terminator sequence. For example, the polyadenylation signal and/or the terminator sequence can be located flanking a segment of the influenza virus genome internal to the two promoters. One favorable polyadenylation signal in the context of the invention is the SV40 polyadenylation signal. An exemplary plasmid vector of the invention is the plasmid pAD3000, illustrated in FIG. 1.

[0029] Any host cell capable of supporting the replication of influenza virus from the vector promoters is suitable in the context of the present invention. Favorable examples of host cells include Vero cells, Per.C6 cells, BHK cells, PCK cells, MDCK cells, MDBK cells, 293 cells (e.g., 293T cells), and COS cells. In combination with the pAD3000 plasmid vectors described herein, Vero cells, 293 cells, COS cells are particularly suitable. In some embodiments, co-cultures of a mixture of at least two of these cell lines, e.g., a combination of COS and MDCK cells or a combination of 293T and MDCK cells, constitute the population of host cells.

[0030] A feature of the invention is the culture of host cells incorporating the plasmids of the invention at a temperature below 37° C., preferably at a temperature equal to, or less than, 35° C. Typically, the cells are cultured at a temperature between 32° C. and 35° C. In some embodiments, the cells are cultured at a temperature between about 32° C. and 34° C., e.g., at about 33° C.

[0031] Another aspect of the invention relates to novel methods for rescuing recombinant or reassortant influenza A or influenza B viruses (i.e., wild type and variant strains of influenza A and/or influenza viruses) from Vero cells in culture. A plurality of vectors incorporating an influenza virus genome is electroporated into a population of Vero cells. The cells are grown under conditions permissive for viral replication, e.g., in the case of cold adapted, attenuated, temperature sensitive virus strains, the Vero cells are grown at a temperature below 37° C., preferably at a temperature equal to, or less than, 35° C. Typically, the cells are cultured at a temperature between 32° C. and 35° C. In some embodiments, the cells are cultured at a temperature between about 32° C. and 34° C., e.g., at about 33° C. Optionally (e.g., for vaccine production), the Vero cells are grown in serum free medium without any animal-derived products.

[0032] In the methods of the invention described above, viruses are recovered following culture of the host cells incorporating the influenza genome plasmids. In some embodiments, the recovered viruses are recombinant viruses. In some embodiments, the viruses are reassortant influenza viruses having genetic contributions from more than one parental strain of virus. Optionally, the recovered recombinant or reassortant viruses are further amplified by passage in cultured cells or in hens' eggs.

[0033] Optionally, the recovered viruses are inactivated. In some embodiments, the recovered viruses comprise an influenza vaccine. For example, the recovered influenza vaccine can be a reassortant influenza viruses (e.g., 6:2 or 7:1 reassortant viruses) having an HA and/or NA antigen derived from a selected strain of influenza A or influenza B. In certain favorable embodiments, the reassortant influenza viruses have an attenuated phenotype. Optionally, the reassortant viruses are cold adapted and/or temperature sensitive, e.g., an attenuated, cold adapted or temperature sensi-

tive influenza B virus having one or more amino acid substitutions selected from the substitutions of Table 17. Such influenza viruses are useful, for example, as live attenuated vaccines for the prophylactic production of an immune response specific for a selected, e.g., pathogenic influenza strain. Influenza viruses, e.g., attenuated reassortant viruses, produced according to the methods of the invention are a feature of the invention.

[0034] In another aspect, the invention relates to methods for producing a recombinant influenza virus vaccine involving introducing a plurality of vectors incorporating an influenza virus genome into a population of host cells capable of supporting replication of influenza virus, culturing the host cells at a temperature less than or equal to 35° C., and recovering an influenza virus capable of eliciting an immune response upon administration to a subject. The vaccines of the invention can be either influenza A or influenza B strain viruses. In some embodiments, the influenza vaccine viruses include an attenuated influenza virus, a cold adapted influenza virus, or a temperature sensitive influenza virus. In certain embodiments, the viruses possess a combination of these desirable properties. In an embodiment, the influenza virus contains an influenza A/Ann Arbor/6/60 strain virus. In another embodiment, the influenza virus incorporates an influenza B/Ann Arbor/1/66 strain virus. Alternatively, the vaccine includes artificially engineered influenza A or influenza B viruses incorporating at least one substituted amino acid which influences the characteristic biological properties of ca A/Ann Arbor/6/60 or ca/B/Ann Arbor/1/66, such as a unique amino acid of these strains. For example, vaccines encompassed by the invention include artificially engineered recombinant and reassortant influenza A viruses including at least one mutation resulting in an amino acid substitution at a position selected from among PB1³⁹¹, PB1⁵⁸¹, PB1⁶⁶¹, PB2²⁶⁵ and NP³⁴ and artificially engineered recombinant and reassortant influenza B viruses including at least one mutation resulting in an amino acid substitution at a position selected from among PB2⁶³⁰, PA⁴³¹, PA⁴⁹⁷, NP⁵⁵, NP¹¹⁴, NP⁴¹⁰, NP⁵⁰⁹, M1¹⁵⁹ and M1¹⁸³.

[0035] In some embodiments, the virus includes a reassortant influenza virus (e.g., a 6:2 or 7:1 reassortant) having viral genome segments derived from more than one influenza virus strain. For example, a reassortant influenza virus vaccine favorably includes an HA and/or NA surface antigen derived from a selected strain of influenza A or B, in combination with the internal genome segments of a virus strain selected for its desirable properties with respect to vaccine production. Often, it is desirable to select the strain of influenza from which the HA and/or NA encoding segments are derived based on predictions of local or worldwide prevalence of pathogenic strains (e.g., as described above). In some cases, the virus strain contributing the internal genome segments is an attenuated, cold adapted and/or temperature sensitive influenza strain, e.g., of A/Ann Arbor/6/60, B/Ann Arbor/1/66, or an artificially engineered influenza strain having one or more amino acid substitutions resulting in the desired phenotype, e.g., influenza A viruses including at least one mutation resulting in an amino acid substitution at a position selected from among PB1³⁹¹, PB1⁵⁸¹, PB1⁶⁶¹, PB2²⁶⁵ and NP³⁴ and influenza B viruses including at least one mutation resulting in an amino acid substitution at a position selected from among PB2⁶³⁰, PA⁴³¹, PA⁴⁹⁷, NP⁵⁵, NP¹¹⁴, NP⁴¹⁰, NP⁵⁰⁹, M1¹⁵⁹ and M1¹⁸³. For example, favorable reassortant viruses include artifi-

cially engineered influenza A viruses with one or more amino acid substitution selected from among PB1³⁹¹ (K391E), PB1⁵⁸¹ (E581G), PB1⁶⁶¹ (A661T), PB2²⁶⁵ (N265S) and NP³⁴ (D34G); and influenza B viruses including one or more amino acid substitutions selected from among PB2⁶³⁰ (S630R); PA⁴³¹ (V431M); PA⁴⁹⁷ (Y497H); NP⁵⁵ (T55A); NP¹¹⁴ (V114A); NP⁴¹⁰ (P410H); NP⁵⁰⁹ (A509T); M1¹⁵⁹ (H159Q) and M1¹⁸³ (M183V).

[0036] If desired, the influenza vaccine viruses are inactivated upon recovery.

[0037] Influenza virus vaccines, including attenuated live vaccines, produced by the methods of the invention are also a feature of the invention. In certain favorable embodiments the influenza virus vaccines are reassortant virus vaccines.

[0038] Another aspect of the invention provides plasmids that are bi-directional expression vectors. The bi-directional expression vectors of the invention incorporate a first promoter inserted between a second promoter and a polyadenylation site, e.g., an SV40 polyadenylation site. In an embodiment, the first promoter and the second promoter can be situated in opposite orientations flanking at least one cloning site. An exemplary vector of the invention is the plasmid pAD3000, illustrated in **FIG. 1**.

[0039] In some embodiments, at least one segment of an influenza virus genome is inserted into the cloning site, e.g., as a double stranded nucleic acid. For example, a vector of the invention includes a plasmid having a first promoter inserted between a second promoter and an SV40 polyadenylation site, wherein the first promoter and the second promoter are situated in opposite orientations flanking at least one segment of an influenza virus.

[0040] Kits including one or more expression vectors of the invention are also a feature of the invention. Typically, the kits also include one or more of: a cell line capable of supporting influenza virus replication, a buffer, a culture medium, an instruction set, a packaging material, and a container. In some embodiments, the kit includes a plurality of expression vectors, each of which includes at least one segment of an influenza virus genome. For example, kits including a plurality of expression vectors each including one of the internal genome segments of a selected virus strain, e.g., selected for its desirable properties with respect to vaccine production or administration, are a feature of the invention. For example, the selected virus strain can be an attenuated, cold adapted and/or temperature sensitive strain, e.g., A/Ann Arbor/6/60 or B/Ann Arbor/1/66, or an alternative strain with the desired properties, such as an artificially engineered strain having one or more amino acid substitutions as described herein, e.g., in Table 17. In an embodiment, the kit includes expression vectors incorporating members of a library of nucleic acids encoding variant HA and/or NA antigens.

[0041] Productively growing cell cultures including at least one cell incorporating a plurality of vectors including an influenza virus genome, at a temperature less than or equal to 35° C., is also a feature of the invention. The composition can also include a cell culture medium. In some embodiments, the plurality of vectors includes bi-directional expression vectors, e.g., comprising a first promoter inserted between a second promoter and an SV40 polyadenylation site. For example, the first promoter and the second pro-

motor can be situated in opposite orientations flanking at least one segment of an influenza virus. The cell cultures of the invention are maintained at a temperature less than or equal to 35° C., such as between about 32° C. and 35° C., typically between about 32° C. and about 34° C., for example, at about 33° C.

[0042] The invention also includes a cell culture system including a productively growing cell culture of at least one cell incorporating a plurality of vectors comprising a an influenza virus genome, as described above, and a regulator for maintaining the culture at a temperature less than or equal to 35° C. For example, the regulator favorably maintains the cell culture at a temperature between about 32° C. and 35° C., typically between about 32° C. and about 34° C., e.g., at about 33° C.

[0043] Another feature of the invention are artificially engineered recombinant or reassortant influenza viruses including one or more amino acid substitutions which influence temperature sensitivity, cold adaptation and/or attenuation. For example, artificially engineered influenza A viruses having one or more amino acid substitution at a position selected from among: PB1³⁹¹, PB1⁵⁸¹, PB1⁶⁶¹, PB2²⁶⁵ and NP³⁴ and artificially engineered influenza B viruses having one or more amino acid substitutions at a position selected from among PB2⁶³⁰, PA⁴³¹, PA⁴⁹⁷, NP⁵⁵, NP¹¹⁴, NP⁴¹⁰, NP⁵⁰⁹, M1¹⁵⁹ and M1¹⁸³ are favorable embodiments of the invention. Exemplary embodiments include influenza A viruses with any one or more of the following amino acid substitutions: PB1³⁹¹ (K391E), PB1⁵⁸¹ (E581G), PB1⁶⁶¹ (A661T), PB2²⁶⁵ (N265S) and NP³⁴ (D34G); and influenza B viruses with any one or more of the following amino acid substitutions: PB2⁶³⁰ (S630R); PA⁴³¹ (V431M); PA⁴⁹⁷ (Y497H); NP⁵⁵ (T55A); NP¹¹⁴ (V114A); NP⁴¹⁰ (P410H); NP⁵⁰⁹ (A509T); M1¹⁵⁹ (H159Q) and M1¹⁸³ (M183V). In certain embodiments, the viruses include a plurality of mutations, such as one, two, three, four, five, six, seven, eight or nine amino acid substitutions at positions identified above. Accordingly, artificially engineered influenza A viruses having amino acid substitutions at all five positions indicated above, e.g., PB1³⁹¹ (K391E), PB1⁵⁸¹ (E581G), PB1⁶⁶¹ (A661T), PB2²⁶⁵ (N265S) and NP³⁴ (D34G) and artificially engineered influenza B viruses having amino acid substitutions at eight or all nine of the positions indicated above, e.g., PB2⁶³⁰ (S630R); PA⁴³¹ (V431M); PA⁴⁹⁷ (Y497H); NP⁵⁵ (T55A); NP¹¹⁴ (V114A); NP⁴¹⁰ (P410H); NP⁵⁰⁹ (A509T); M1¹⁵⁹ (H159Q) and M1¹⁸³ (M183V), are encompassed by the invention. In addition, the viruses can include one or more additional amino acid substitutions not enumerated above.

[0044] In certain embodiments, the artificially engineered influenza viruses are temperature sensitive influenza viruses, cold adapted influenza viruses and/or attenuated influenza viruses. For example, a temperature sensitive influenza virus according to the invention typically exhibits between about 2.0 and 5.0 log₁₀ reduction in growth at 39° C. as compared to a wild type influenza virus. For example, a temperature sensitive virus favorably exhibits at least about 2.0 log₁₀, at least about 3.0 log₁₀, at least about 4.0 log₁₀, or at least about 4.5 log₁₀ reduction in growth at 39° C. relative to that of a wild type influenza virus. Typically, but not necessarily, a temperature sensitive influenza virus retains robust growth characteristics at 33° C. An attenuated influenza virus of the invention typically exhibits between about a 2.0 and a 5.0

log₁₀ reduction in growth in a ferret attenuation assay as compared to a wild type influenza virus. For example, an attenuated influenza virus of the invention exhibits at least about a 2.0 log₁₀, frequently about a 3.0 log₁₀, and favorably at least about a 4.0 log₁₀ reduction in growth in a ferret attenuation assay relative to wild type influenza virus.

[0045] The present invention also relates to the identification and manipulation of amino acid residues in HA and NA which affect influenza virus replication in cells and embryonated chicken eggs. The present invention further relates to the use of reverse genetics technology to generate HA and NA influenza virus vaccine variants with improved replication in embryonated chicken eggs and/or cells. The invention further relates to methods for modulating HA receptor binding activity and/or NA neuraminidase activity. Additionally, the invention provides influenza viruses with enhanced ability to replicate in embryonated chicken eggs and/or cells.

[0046] In one embodiment the invention provides methods for manipulating the amino acid residues of HA and/or NA to increase the ability of an influenza virus to replicate in embryonated chicken eggs and/or cells. The method involves the introduction of amino acid residues substitutions in HA and/or NA and makes use of methods of producing influenza virus in cell culture by introducing a plurality of vectors incorporating an influenza virus genome into a population of host cells capable of supporting replication of influenza virus, culturing the cells and recovering influenza virus. Preferably, the recovered influenza virus has increase ability to replicate in embryonated chicken eggs and/or cells. In another embodiment, the present invention provides influenza virus variants with increase ability to replicate in embryonated chicken eggs (referred to herein as “replication enhanced influenza variant(s)”) when compared to unmodified influenza viral strains.

[0047] The present invention further includes an improved method of rescue, wherein electroporated animal (e.g., SF Vero) cells (electroporated with, e.g., polynucleotides (e.g., plasmids and vectors) of the invention) are co-cultivated with another cell selected from the group including, but not limited to: chicken embryo kidney (CEK) cells, chicken embryo fibroblasts, primary chick kidney cells, and cells isolated from the chorioallantoic membrane of embryonated chicken eggs. Other cells useful for this rescue method may include any cell that supports replication of influenza virus and meets acceptable standards for regulatory approval. Sources of cells include, for example, chicken flocks from SPF chicken flocks. See, Examples 9 and 10 herein.

[0048] In one preferred embodiment of the invention, rescue efficiency of virus is improved by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 2-fold, or at least 3-fold, or at least 5-fold.

[0049] In another preferred embodiment of the invention, rescue efficiency of virus is at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 99%. Efficiency can be determined, for example, by measuring how many eggs injected with the rescued viruses (X) have subsequent detectable HA titers (Y) and dividing Y/X.

[0050] The methods described supra as Examples 9 and 10 may be used to electroporate polynucleotides (e.g., plasmids and vectors) described herein or, e.g., in U.S. patent application Ser. Nos. 09/396,539, 09/844,517, PCT/US0113656, PCT/US00/09021, US03012728; U.S. Pat. No. 6,649,372; WO 03/091401, US200201677, which are incorporated by reference herein.

[0051] A preferred embodiment of the invention is a method of rescue of influenza virus, wherein animal cells (e.g., Vero cells) are electroporated with plasmids that encode an influenza RNA polymerase and nucleoprotein and wherein the electroporated animal cells are co-cultivated with another cell type.

[0052] A preferred embodiment of the invention is a method of rescue of influenza virus (e.g., influenza A virus, cold adapted viruses, an attenuated viruses), wherein animal cells (e.g., Vero cells) are electroporated with plasmids that encode an influenza RNA polymerase and nucleoprotein. The number of plasmids electroporated may be, for example, eight or twelve.

[0053] A preferred embodiment of the invention is a method of rescue of influenza virus (e.g., influenza A virus, cold adapted viruses, an attenuated viruses), wherein animal cells (e.g., Vero cells) are electroporated with plasmids that encode an influenza RNA polymerase and nucleoprotein and wherein the electroporated animal cells are co-cultivated with another cell type (e.g., CEK cells). The number of plasmids electroporated may be, for example, eight or twelve.

[0054] Another preferred embodiment of the invention is a method of rescue of influenza virus, wherein (a) animal cells are electroporated with cell expression vectors which direct the expression in said cells of genomic or antigenomic vRNA segments, and a nucleoprotein, and an RNA-dependent polymerase, such that ribonucleoprotein complexes can be formed and viral particles can be assembled (with or without a helper virus); and (b) culturing said cells wherein viral particles are packaged and rescued.

[0055] Another preferred embodiment of the invention is a method of rescue of influenza virus, wherein animal cells are electroporated with expression plasmids (see, e.g., U.S. patent application Ser. Nos. 09/396,539, 09/844,517, PCT/US0113656, PCT/US00/09021, US03012728; U.S. Pat. No. 6,649,372; WO 03/091401, US200201677, which are incorporated by reference herein), for example, comprising viral cDNA corresponding to the genomic segment of an influenza virus, wherein the cDNA is inserted between an RNA polymerase I (polI) promoter and a regulatory element for the synthesis of vRNA or cRNA with an exact 3' end, which are in turn inserted between an RNA polymerase II (polII) promoter and a polyadenylation signal, and wherein the cDNA only encodes an influenza viral protein.

[0056] Other embodiments of the invention include influenza viruses produced by the methods described herein (e.g., Examples 9 and 10) and vaccines comprising the same.

[0057] Other preferred embodiments of the invention include compositions which generates infectious influenza viruses from cloned viral cDNA comprising SF Vero electroporated with a set of plasmids wherein each plasmid comprises one viral genomic segment, and wherein viral cDNA corresponding to the genomic segment is inserted

between an RNA polymerase I (polI) promoter and a regulatory element for the synthesis of vRNA or cRNA with an exact 3' end, which results in expression of viral mRNA and a corresponding viral protein, wherein the expression of the full set of vRNAs or cRNAs and viral proteins results in the assembly of an infectious influenza virus.

BRIEF DESCRIPTION OF THE DRAWINGS

[0058] **FIG. 1:** Illustration of pAD3000 plasmid (SEQ ID NO: 90).

[0059] **FIG. 2:** Micrographs of infected cells

[0060] **FIG. 3:** Genotyping analysis of rMDV-A and 6:2 H1N1 reassortant virus from plasmid transfection.

[0061] **FIG. 4:** Illustration of eight plasmid system for the production of influenza B virus.

[0062] **FIG. 5:** A and B. Characterization of recombinant MDV-B virus by RT-PCR; C and D. Characterization of recombinant B/Yamanashi/166/98 by RT PCR.

[0063] **FIG. 6:** Sequence of pAD3000 in GeneBank format.

[0064] **FIG. 7:** Sequence alignment with MDV-B and eight plasmids (SEQ ID NOS: 91-98, respectively)

[0065] **FIG. 8:** RT-PCR products derived from simultaneous amplification of HA and NA segments of influenza B strains.

[0066] **FIG. 9:** Bar graph illustrating relative titers of recombinant and reassortant virus.

[0067] **FIG. 10:** Bar graph illustrating relative titers of reassortant virus under permissive and restrictive temperatures (temperature sensitivity).

[0068] **FIG. 11:** Graphic representation of reassortant viruses incorporating specific mutations (knock-in) correlating with temperature sensitivity (left panel) and relative titers at permissive and restrictive temperatures (temperature sensitivity) (right panel).

[0069] **FIG. 12:** Determination of ts mutations in a mini-genome assay. A. HEp-2 cells were transfected with PB1, PB2, PA, NP and pFlu-CAT, incubated at 33 or 39° C. for 18 hr and cell extracts were analyzed for CAT reporter gene expression. B. CAT mRNA expression by primer extension assay.

[0070] **FIG. 13:** Schematic illustration of triple-gene recombinants with wild type residues in PA, NP, and M1 proteins.

[0071] **FIG. 14:** Tabulation of growth of single-gene and double-gene recombinant viruses.

[0072] **FIG. 15:** Tabulation of amino acid residue of the nucleoprotein corresponding to non-ts phenotype.

[0073] **FIG. 16:** Schematic diagram of recombinant PR8 mutants. The mutations introduced in PB1 and/or PB2 genes are indicated by the filled dots.

[0074] **FIG. 17:** Bar graph illustrating relative titers at 33° C. and 39° C.

[0075] **FIG. 18:** Photomicrographs illustrating plaque morphology of PR8 mutants at various temperatures. MDCK cells were infected with virus as indicated and incubated at 33, 37 and 39° C. for three days. Virus plaques were visualized by immunostaining and photographed.

[0076] **FIG. 19:** Protein synthesis at permissive and non-permissive temperatures. MDCK cells were infected with viruses as indicated and incubated at 33 or 39° C. overnight. Radiolabeled labeled polypeptides were electrophoresed on an SDS-PAGE and autoradiographed. Viral proteins, HA, NP, M1 and NS are indicated.

[0077] **FIG. 20:** A. Line graphs illustrating differential replication of MDV-A and MDV-B in Per.C6 cells relative to replication in MDCK cells; B. Line graph illustrating differential replication of MDV-A single gene reassortants in Per.C6 cells.

[0078] **FIG. 21:** Bar graphs illustrating differential replication of reassortant viruses. Gray boxes represent wild type amino acid residues. The dotted line represents the shut-off temperature (ts) of $2.0 \log_{10}$.

[0079] FIGS. 22-23: Antigenically compare A/Panama/99 (H3N2) and A/Fujian/411/02-like (H3N2).

[0080] FIGS. 24-28: Show molecular basis for antigenic drift from A/Panama/99 to A/Fujian/02-like.

[0081] FIGS. 29-35: Detail modifications in strains to produce increased virus growth in embryonated eggs.

[0082] **FIG. 36:** HA receptor binding affinity of recombinant viruses. 6:2 A/Fujian, A/Sendai, A/Wyoming, and A/Fujian variants with V186 and 1226 or L183 and A226 changes were adsorbed to MDCK cells at an moi of 1.0 at 4° C. or 33° C. for 30 min, and the infected cells were washed three times (+) or left untreated (-). After 6 hr of incubation at 33° C., the cells were processed for immunofluorescence staining. The percentage of infected cells (mean \pm SD) indicated in each image was an average of six images.

[0083] **FIG. 37:** Growth kinetics of recombinant viruses in MDCK cells. MDCK cells were infected at an moi of 1.0 at either 33° C. or 4° C. for 30 min, washed 3x with PBS. The infected cells were incubated at 33° C. and at the indicated time intervals the culture supernatants were collected and the virus amount was determined by plaque assay.

[0084] **FIG. 38:** receptor-binding sites in HA and NA of H3N2 subtypes. The residues that were shown to increase the HA receptor-binding affinity and to decrease the NA enzymatic activity in relation to sialic acid (SIA) binding sites are indicated. The HA monomer was modeled using SHMG and the NA monomer was modeled based on 2BAT using WebLab ViewerLite 3.10 (Accelrys, San Diego, Calif.).

DETAILED DESCRIPTION

[0085] Many pathogenic influenza virus strains grow only poorly in tissue culture, and strains suitable for production of live attenuated virus vaccines (e.g., temperature sensitive, cold adapted and/or attenuated influenza viruses) have not been successfully grown in cultured cells for commercial production. The present invention provides a multi-plasmid transfection system which permits the growth and recovery

of influenza virus strains which are not adapted for growth under standard cell culture conditions. An additional challenge in developing and producing influenza vaccines is that one or more of the circulating influenza strains may not replicate well in embryonic chicken eggs. The present invention identifies several amino acid residues which influence the activities of the HA and NA proteins and have identified specific amino acid substitutions which can modulate these activities. The present invention discloses that modulation of the HA receptor binding activity and/or the NA neuraminidase activity can enhance the replication of influenza in eggs and/or host cells (e.g., Vero or MDCK cells). Specifically the present invention discloses combinations of amino acid substitutions in HA and/or NA can enhance viral replication in eggs and/or cells and demonstrates that these amino acid substitutions have no significant impact on antigenicity of these recombinant influenza viruses. Thus, the present invention provides for the use of reverse genetic technology to improve the manufacture of influenza virus vaccines.

[0086] In a first aspect, the methods of the invention provide vectors and methods for producing recombinant influenza B virus in cell culture entirely from cloned viral DNA. In another aspect, the methods of the present invention are based in part on the development of tissue culture conditions which support the growth of virus strains (both A strain and B strain influenza viruses) with desirable properties relative to vaccine production (e.g., attenuated pathogenicity or phenotype, cold adaptation, temperature sensitivity, etc.) in vitro in cultured cells. Influenza viruses are produced by introducing a plurality of vectors incorporating cloned viral genome segments into host cells, and culturing the cells at a temperature not exceeding 35° C. When vectors including an influenza virus genome are transfected, recombinant viruses suitable as vaccines can be recovered by standard purification procedures. Using the vector system and methods of the invention, reassortant viruses incorporating the six internal gene segments of a strain selected for its desirable properties with respect to vaccine production, and the immunogenic HA and NA segments from a selected, e.g., pathogenic strain, can be rapidly and efficiently produced in tissue culture. Thus, the system and methods described herein are useful for the rapid production in cell culture of recombinant and reassortant influenza A and B viruses, including viruses suitable for use as vaccines, including live attenuated vaccines, such as vaccines suitable for intranasal administration.

[0087] Typically, a single Master Donor Virus (MDV) strain is selected for each of the A and B subtypes. In the case of a live attenuated vaccine, the Master Donor Virus strain is typically chosen for its favorable properties, e.g., temperature sensitivity, cold adaptation and/or attenuation, relative to vaccine production. For example, exemplary Master Donor Strains include such temperature sensitive, attenuated and cold adapted strains of A/Ann Arbor/6/60 and B/Ann Arbor/1/66, respectively. The present invention elucidates the underlying mutations resulting in the ca, ts and att phenotypes of these virus strains, and provides methods for producing novel strains of influenza suitable for use as donor strains in the context of recombinant and reassortant vaccine production.

[0088] For example, a selected master donor type A virus (MDV-A), or master donor type B virus (MDV-B), is produced from a plurality of cloned viral cDNAs constituting the viral genome. In an exemplary embodiment, recombinant viruses are produced from eight cloned viral cDNAs. Eight viral cDNAs representing either the selected MDV-A or MDV-B sequences of PB2, PB1, PA, NP, HA, NA, M and NS are cloned into a bi-directional expression vector, such as a plasmid (e.g., pAD3000), such that the viral genomic RNA can be transcribed from an RNA polymerase I (pol I) promoter from one strand and the viral mRNAs can be synthesized from an RNA polymerase II (pol II) promoter from the other strand. Optionally, any gene segment can be modified, including the HA segment (e.g., to remove the multi-basic cleavage site).

[0089] Infectious recombinant MDV-A or MDV-B virus is then recovered following transfection of plasmids bearing the eight viral cDNAs into appropriate host cells, e.g., Vero cells, co-cultured MDCK/293T or MDCK/COS7 cells. Using the plasmids and methods described herein, the invention is useful, e.g., for generating 6:2 reassortant influenza vaccines by co-transfection of the 6 internal genes (PB1, PB2, PA, NP, M and NS) of the selected virus (e.g., MDV-A, MDV-B) together with the HA and NA derived from different corresponding type (A or B) influenza viruses. For example, the HA segment is favorably selected from a pathogenically relevant H1, H3 or B strain, as is routinely performed for vaccine production. Similarly, the HA segment can be selected from a strain with emerging relevance as a pathogenic strain such as an H2 strain (e.g., H2N2), an H5 strain (e.g., H5N1) or an H7 strain (e.g., H7N7). Reassortants incorporating seven genome segments of the MDV and either the HA or NA gene of a selected strain (7:1 reassortants) can also be produced. In addition, this system is useful for determining the molecular basis of phenotypic characteristics, e.g., the attenuated (att), cold adapted (ca), and temperature sensitive (ts) phenotypes, relevant to vaccine production.

[0090] In another aspect the invention provides methods for manipulating the amino acid residues of HA and/or NA to increase the ability of an influenza virus to replicate in embryonated chicken eggs and/or cells. For example, the methods of the present invention can be used to modulate HA receptor binding activity and/or NA neuraminidase activity to increase the ability of an influenza virus to replicate in eggs and/or cells. Additionally, the invention provides influenza viruses with enhanced ability to replicate in embryonated chicken eggs and/or cells.

DEFINITIONS

[0091] Unless defined otherwise, all scientific and technical terms are understood to have the same meaning as commonly used in the art to which they pertain. For the purpose of the present invention the following terms are defined below.

[0092] The terms "nucleic acid," "polynucleotide," "poly-nucleotide sequence" and "nucleic acid sequence" refer to single-stranded or double-stranded deoxyribonucleotide or ribonucleotide polymers, or chimeras or analogues thereof. As used herein, the term optionally includes polymers of analogs of naturally occurring nucleotides having the essential nature of natural nucleotides in that they hybridize to

single-stranded nucleic acids in a manner similar to naturally occurring nucleotides (e.g., peptide nucleic acids). Unless otherwise indicated, a particular nucleic acid sequence of this invention encompasses complementary sequences, in addition to the sequence explicitly indicated.

[0093] The term "gene" is used broadly to refer to any nucleic acid associated with a biological function. Thus, genes include coding sequences and/or the regulatory sequences required for their expression. The term "gene" applies to a specific genomic sequence, as well as to a cDNA or an mRNA encoded by that genomic sequence.

[0094] Genes also include non-expressed nucleic acid segments that, for example, form recognition sequences for other proteins. Non-expressed regulatory sequences include "promoters" and "enhancers," to which regulatory proteins such as transcription factors bind, resulting in transcription of adjacent or nearby sequences. A "Tissue specific" promoter or enhancer is one which regulates transcription in a specific tissue type or cell type, or types.

[0095] The term "vector" refers to the means by which a nucleic can be propagated and/or transferred between organisms, cells, or cellular components. Vectors include plasmids, viruses, bacteriophage, pro-viruses, phagemids, transposons, and artificial chromosomes, and the like, that replicate autonomously or can integrate into a chromosome of a host cell. A vector can also be a naked RNA polynucleotide, a naked DNA polynucleotide, a polynucleotide composed of both DNA and RNA within the same strand, a poly-lysine-conjugated DNA or RNA, a peptide-conjugated DNA or RNA, a liposome-conjugated DNA, or the like, that are not autonomously replicating. Most commonly, the vectors of the present invention are plasmids.

[0096] An "expression vector" is a vector, such as a plasmid, which is capable of promoting expression, as well as replication of a nucleic acid incorporated therein. Typically, the nucleic acid to be expressed is "operably linked" to a promoter and/or enhancer, and is subject to transcription regulatory control by the promoter and/or enhancer.

[0097] A "bi-directional expression vector" is typically characterized by two alternative promoters oriented in the opposite direction relative to a nucleic acid situated between the two promoters, such that expression can be initiated in both orientations resulting in, e.g., transcription of both plus (+) or sense strand, and negative (-) or antisense strand RNAs. Alternatively, the bi-directional expression vector can be an ambisense vector, in which the viral mRNA and viral genomic RNA (as a cRNA) are expressed from the same strand.

[0098] In the context of the invention, the term "isolated" refers to a biological material, such as a nucleic acid or a protein, which is substantially free from components that normally accompany or interact with it in its naturally occurring environment. The isolated material optionally comprises material not found with the material in its natural environment, e.g., a cell. For example, if the material is in its natural environment, such as a cell, the material has been placed at a location in the cell (e.g., genome or genetic element) not native to a material found in that environment. For example, a naturally occurring nucleic acid (e.g., a coding sequence, a promoter, an enhancer, etc.) becomes isolated if it is introduced by non-naturally occurring means

to a locus of the genome (e.g., a vector, such as a plasmid or virus vector, or amplicon) not native to that nucleic acid. Such nucleic acids are also referred to as "heterologous" nucleic acids.

[0099] The term "recombinant" indicates that the material (e.g., a nucleic acid or protein) has been artificially or synthetically (non-naturally) altered by human intervention. The alteration can be performed on the material within, or removed from, its natural environment or state. Specifically, when referring to a virus, e.g., an influenza virus, the virus is recombinant when it is produced by the expression of a recombinant nucleic acid.

[0100] The term "reassortant," when referring to a virus, indicates that the virus includes genetic and/or polypeptide components derived from more than one parental viral strain or source. For example, a 7:1 reassortant includes 7 viral genomic segments (or gene segments) derived from a first parental virus, and a single complementary viral genomic segment, e.g., encoding hemagglutinin or neuraminidase, from a second parental virus. A 6:2 reassortant includes 6 genomic segments, most commonly the 6 internal genes from a first parental virus, and two complementary segments, e.g., hemagglutinin and neuraminidase, from a different parental virus.

[0101] The term "introduced" when referring to a heterologous or isolated nucleic acid refers to the incorporation of a nucleic acid into a eukaryotic or prokaryotic cell where the nucleic acid can be incorporated into the genome of the cell (e.g., chromosome, plasmid, plastid or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (e.g., transfected mRNA). The term includes such methods as "infection," "transfection," "transformation" and "transduction." In the context of the invention a variety of methods can be employed to introduce nucleic acids into prokaryotic cells, including electroporation, Calcium phosphate precipitation, lipid mediated transfection (lipofection), etc.

[0102] The term "host cell" means a cell which contains a heterologous nucleic acid, such as a vector, and supports the replication and/or expression of the nucleic acid, and optionally production of one or more encoded products including a polypeptide and/or a virus. Host cells can be prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, avian or mammalian cells, including human cells. Exemplary host cells in the context of the invention include Vero (African green monkey kidney) cells, Per.C6 cells (human embryonic retinal cells), BHK (baby hamster kidney) cells, primary chick kidney (PCK) cells, Madin-Darby Canine Kidney (MDCK) cells, Madin-Darby Bovine Kidney (MDBK) cells, 293 cells (e.g., 293T cells), and COS cells (e.g., COS1, COS7 cells). The term host cell encompasses combinations or mixtures of cells including, e.g., mixed cultures of different cell types or cell lines.

[0103] The terms "temperature sensitive," "cold adapted" and "attenuated" are well known in the art. For example, the term "temperature sensitive" ("ts") indicates that the virus exhibits a 100 fold or greater reduction in titer at 39° C. relative to 33° C. for influenza A strains, and that the virus exhibits a 100 fold or greater reduction in titer at 37° C. relative to 33° C. for influenza B strains. For example, the term "cold adapted" ("ca") indicates that the virus exhibits growth at 25° C. within 100 fold of its growth at 33° C. For

example, the term "attenuated" ("att") indicates that the virus replicates in the upper airways of ferrets but is not detectable in lung tissues, and does not cause influenza-like illness in the animal. It will be understood that viruses with intermediate phenotypes, i.e., viruses exhibiting titer reductions less than 100 fold at 39° C. (for A strain viruses) or 37° C. (for B strain viruses), exhibiting growth at 25° C. that is more than 100 fold than its growth at 33° C. (e.g., within 200 fold, 500 fold, 1000 fold, 10,000 fold less), and/or exhibit reduced growth in the lungs relative to growth in the upper airways of ferrets (i.e., partially attenuated) and/or reduced influenza like illness in the animal, which possess one or more of the amino acid substitutions described herein are also useful viruses encompassed by the invention. Growth indicates viral quantity as indicated by titer, plaque size or morphology, particle density or other measures known to those of skill in the art.

[0104] The expression "artificially engineered" is used herein to indicate that the virus, viral nucleic acid or virally encoded product, e.g., a polypeptide, a vaccine, comprises at least one mutation introduced by recombinant methods, e.g., site directed mutagenesis, PCR mutagenesis, etc. The expression "artificially engineered" when referring to a virus (or viral component or product) comprising one or more nucleotide mutations and/or amino acid substitutions indicates that the viral genome or genome segment encoding the virus (or viral component or product) is not derived from naturally occurring sources, such as a naturally occurring or previously existing laboratory strain of virus produced by non-recombinant methods (such as progressive passage at 25° C.), e.g., a wild type or cold adapted A/Ann Arbor/6/60 or B/Ann Arbor/1/66strain.

[0105] Influenza Virus

[0106] The genome of Influenza viruses is composed of eight segments of linear (-) strand ribonucleic acid (RNA), encoding the immunogenic hemagglutinin (HA) and neuraminidase (NA) proteins, and six internal core polypeptides: the nucleocapsid nucleoprotein (NP); matrix proteins (M); non-structural proteins (NS); and 3 RNA polymerase (PA, PB1, PB2) proteins. During replication, the genomic viral RNA is transcribed into (+) strand messenger RNA and (-) strand genomic cRNA in the nucleus of the host cell. Each of the eight genomic segments is packaged into ribonucleoprotein complexes that contain, in addition to the RNA, NP and a polymerase complex (PB1, PB2, and PA).

[0107] In the present invention, viral genomic RNA corresponding to each of the eight segments is inserted into a recombinant vector for manipulation and production of influenza viruses. A variety of vectors, including viral vectors, plasmids, cosmids, phage, and artificial chromosomes, can be employed in the context of the invention. Typically, for ease of manipulation, the viral genomic segments are inserted into a plasmid vector, providing one or more origins of replication functional in bacterial and eukaryotic cells, and, optionally, a marker convenient for screening or selecting cells incorporating the plasmid sequence. An exemplary vector, plasmid PAD3000 is illustrated in **FIG. 1**.

[0108] Most commonly, the plasmid vectors of the invention are bi-directional expression vectors capable of initiating transcription of the inserted viral genomic segment in either direction, that is, giving rise to both (+) strand and (-) strand viral RNA molecules. To effect bi-directional tran-

scription, each of the viral genomic segments is inserted into a vector having at least two independent promoters, such that copies of viral genomic RNA are transcribed by a first RNA polymerase promoter (e.g., Pol I), from one strand, and viral mRNAs are synthesized from a second RNA polymerase promoter (e.g., Pol II). Accordingly, the two promoters are arranged in opposite orientations flanking at least one cloning site (i.e., a restriction enzyme recognition sequence) preferably a unique cloning site, suitable for insertion of viral genomic RNA segments. Alternatively, an "ambisense" vector can be employed in which the (+) strand mRNA and the (-) strand viral RNA (as a cRNA) are transcribed from the same strand of the vector.

[0109] Expression Vectors

[0110] The influenza virus genome segment to be expressed is operably linked to an appropriate transcription control sequence (promoter) to direct mRNA synthesis. A variety of promoters are suitable for use in expression vectors for regulating transcription of influenza virus genome segments. In certain embodiments, e.g., wherein the vector is the plasmid pAD3000, the cytomegalovirus (CMV) DNA dependent RNA Polymerase II (Pol II) promoter is utilized. If desired, e.g., for regulating conditional expression, other promoters can be substituted which induce RNA transcription under the specified conditions, or in the specified tissues or cells. Numerous viral and mammalian, e.g., human promoters are available, or can be isolated according to the specific application contemplated. For example, alternative promoters obtained from the genomes of animal and human viruses include such promoters as the adenovirus (such as Adenovirus 2), papilloma virus, hepatitis-B virus, polyoma virus, and Simian Virus 40 (SV40), and various retroviral promoters. Mammalian promoters include, among many others, the actin promoter, immunoglobulin promoters, heat-shock promoters, and the like. In addition, bacteriophage promoters can be employed in conjunction with the cognate RNA polymerase, e.g., the T7 promoter.

[0111] Transcription is optionally increased by including an enhancer sequence. Enhancers are typically short, e.g., 10-500 bp, cis-acting DNA elements that act in concert with a promoter to increase transcription. Many enhancer sequences have been isolated from mammalian genes (hemoglobin, elastase, albumin, alpha-fetoprotein, and insulin), and eukaryotic cell viruses. The enhancer can be spliced into the vector at a position 5' or 3' to the heterologous coding sequence, but is typically inserted at a site 5' to the promoter. Typically, the promoter, and if desired, additional transcription enhancing sequences are chosen to optimize expression in the host cell type into which the heterologous DNA is to be introduced (Scharf et al. (1994) *Heat stress promoters and transcription factors* *Results Probl Cell Differ* 20:125-62; Kriegler et al. (1990) *Assembly of enhancers, promoters, and splice signals to control expression of transferred genes* *Methods in Enzymol* 185: 512-27). Optionally, the amplicon can also contain a ribosome binding site or an internal ribosome entry site (IRES) for translation initiation.

[0112] The vectors of the invention also favorably include sequences necessary for the termination of transcription and for stabilizing the mRNA, such as a polyadenylation site or a terminator sequence. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of

eukaryotic or viral DNAs or cDNAs. In one embodiment, e.g., involving the plasmid pAD3000, the SV40 polyadenylation sequences provide a polyadenylation signal.

[0113] In addition, as described above, the expression vectors optionally include one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells, in addition to genes previously listed, markers such as dihydrofolate reductase or neomycin resistance are suitable for selection in eukaryotic cell culture.

[0114] The vector containing the appropriate DNA sequence as described above, as well as an appropriate promoter or control sequence, can be employed to transform a host cell permitting expression of the protein. While the vectors of the invention can be replicated in bacterial cells, most frequently it will be desirable to introduce them into mammalian cells, e.g., Vero cells, BHK cells, MDCK cell, 293 cells, COS cells, for the purpose of expression.

[0115] Additional Expression Elements

[0116] Most commonly, the genome segment encoding the influenza virus protein includes any additional sequences necessary for its expression, including translation into a functional viral protein. In other situations, a minigene, or other artificial construct encoding the viral proteins, e.g., an HA or NA protein, can be employed. In this case, it is often desirable to include specific initiation signals which aid in the efficient translation of the heterologous coding sequence. These signals can include, e.g., the ATG initiation codon and adjacent sequences. To insure translation of the entire insert, the initiation codon is inserted in the correct reading frame relative to the viral protein. Exogenous transcriptional elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers appropriate to the cell system in use.

[0117] If desired, polynucleotide sequences encoding additional expressed elements, such as signal sequences, secretion or localization sequences, and the like can be incorporated into the vector, usually, in-frame with the polynucleotide sequence of interest, e.g., to target polypeptide expression to a desired cellular compartment, membrane, or organelle, or into the cell culture media. Such sequences are known to those of skill, and include secretion leader peptides, organelle targeting sequences (e.g., nuclear localization sequences, ER retention signals, mitochondrial transit sequences), membrane localization/anchor sequences (e.g., stop transfer sequences, GPI anchor sequences), and the like.

[0118] Influenza Virus Vaccine

[0119] Historically, influenza virus vaccines have been produced in embryonated hens' eggs using strains of virus selected based on empirical predictions of relevant strains. More recently, reassortant viruses have been produced that incorporate selected hemagglutinin and neuraminidase antigens in the context of an approved attenuated, temperature sensitive master strain. Following culture of the virus through multiple passages in hens' eggs, influenza viruses are recovered and, optionally, inactivated, e.g., using formaldehyde and/or β -propiolactone. However, production of influenza vaccine in this manner has several significant drawbacks. Contaminants remaining from the hens' eggs are highly antigenic, pyrogenic, and frequently result in signifi-

cant side effects upon administration. More importantly, strains designated for production must be selected and distributed, typically months in advance of the next flu season to allow time for production and inactivation of influenza vaccine. Attempts at producing recombinant and reassortant vaccines in cell culture have been hampered by the inability of any of the strains approved for vaccine production to grow efficiently under standard cell culture conditions.

[0120] The present invention provides a vector system, and methods for producing recombinant and reassortant viruses in culture which make it possible to rapidly produce vaccines corresponding to one or many selected antigenic strains of virus. In particular, conditions and strains are provided that result in efficient production of viruses from a multi plasmid system in cell culture. Optionally, if desired, the viruses can be further amplified in Hens' eggs.

[0121] For example, it has not been possible to grow the influenza B master strain B/Ann Arbor/1/66 under standard cell culture conditions, e.g., at 37° C. In the methods of the present invention, multiple plasmids, each incorporating a segment of an influenza virus genome are introduced into suitable cells, and maintained in culture at a temperature less than or equal to 35° C. Typically, the cultures are maintained at between about 32° C. and 35° C., preferably between about 32° C. and about 34° C., e.g., at about 33° C.

[0122] Typically, the cultures are maintained in a system, such as a cell culture incubator, under controlled humidity and Co₂, at constant temperature using a temperature regulator, such as a thermostat to insure that the temperature does not exceed 35° C.

[0123] Reassortant influenza viruses can be readily obtained by introducing a subset of vectors corresponding to genomic segments of a master influenza virus, in combination with complementary segments derived from strains of interest (e.g., antigenic variants of interest). Typically, the master strains are selected on the basis of desirable properties relevant to vaccine administration. For example, for vaccine production, e.g., for production of a live attenuated vaccine, the master donor virus strain may be selected for an attenuated phenotype, cold adaptation and/or temperature sensitivity. In this context, Influenza A strain ca A/Ann Arbor/6/60; Influenza B strain ca B/Ann Arbor/1/66; or another strain selected for its desirable phenotypic properties, e.g., an attenuated, cold adapted, and/or temperature sensitive strain, such as an artificially engineered influenza A strain as described in Example 4; or an artificially engineered influenza B strain incorporating one or more of the amino acid substitutions specified in Table 17 are favorably selected as master donor strains.

[0124] In one embodiment, plasmids incorporating the six internal genes of the influenza master virus strain, (i.e., PB1, PB2, PA, NP, NB, M1, BM2, NS1 and NS2) are transfected into suitable host cells in combination with hemagglutinin and neuraminidase segments from an antigenically desirable strain, e.g., a strain predicted to cause significant local or global influenza infection. Following replication of the reassortant virus in cell culture at appropriate temperatures for efficient recovery, e.g., equal to or less than 35° C., such as between about 32° C. and 35° C., for example between about 32° C. and about 34° C., or at about 33° C., reassortant

viruses is recovered. Optionally, the recovered virus can be inactivated using a denaturing agent such as formaldehyde or β-propiolactone.

[0125] Attenuated, Temperature Sensitive and Cold Adapted Influenza Virus Vaccines

[0126] In one aspect, the present invention is based on the determination of the mutations underlying the ts phenotype in preferred Master Donor Strains of virus. To determine the functional importance of single nucleotide changes in the MDV strain genome, reassortant viruses derived from highly related strains within the A/AA/6/60 lineage were evaluated for temperature sensitivity. The isogenic nature of the two parental strains enables the evaluation of single nucleotide changes on the ts phenotype. Accordingly, the genetic basis for the ts phenotype of MDV-A is mapped at the nucleotide level to specific amino acid residues within PB1, PB2, and NP.

[0127] Previous attempts to map the genetic basis of the ts phenotype of ca A/AA/6/60 utilized classical coinfection/reassortant techniques to create single and multiple gene reassortants between A/AA/6/60 and an unrelated wt strain. These studies suggested that both PB2, and PB1 contributed to the ts phenotype (Kendal et al. (1978) Biochemical characteristics of recombinant viruses derived at sub-optimal temperatures: evidence that ts lesions are present in RNA segments 1 and 3, and that RNA 1 codes for the virion transcriptase enzyme, p. 734-743. In B. W. J. Mahy, and R. D. Barry (ed.) *Negative Strand Viruses*, Academic Press; Kendal et al. (1977) Comparative studies of wild-type and cold mutant (temperature sensitive) influenza viruses: genealogy of the matrix (M) and the non-structural (NS) proteins in recombinant cold-adapted H3N2 viruses *J Gen Virol* 37:145-159; Kendal et al. (1979) Comparative studies of wild-type and cold-mutant (temperature sensitive) influenza viruses. independent segregation of temperature-sensitivity of virus replication from temperature-sensitivity of virion transcriptase activity during recombination of mutant A/Ann Arbor/6/60 with wild-type H3N2 strains *J Gen Virol* 44:443-456; Snyder et al. (1988) Four viral genes independently contribute to attenuation of live influenza A/Ann Arbor/6/60 (H2N2) cold-adapted reassortant virus vaccines *J Virol* 62:488-95). Interpretation of these studies, however, was confounded by constellation effects, which were caused by mixing gene segments from two divergent influenza A strains. Weakened interactions could have occurred through changes between the A/AA/6/60 and wt gene segments other than those specifically involved in expression of the ts phenotype from the A/AA/6/60 background. Constellation effects were also shown to confound the interpretation of association of the M gene segment with the att phenotype (Subbarao et al. (1992) The attenuation phenotype conferred by the M gene of the influenza A/Ann Arbor/6/60 cold-adapted virus (H2N2) on the A/Korea/82 (H3N2) reassortant virus results from a gene constellation effect *Virus Res* 25:37-50).

[0128] In the present invention, mutations resulting in amino acid substitutions at positions PB1³⁹¹, PB1⁵⁸¹, PB1⁶⁶¹, PB2²⁶⁵ and NP³⁴ are identified as functionally important in conferring the temperature sensitive phenotype on the MDV-A strain virus. As will be understood by those of skill in the art, mutations in nucleotides at positions PB1¹¹⁹⁵, PB1¹⁷⁶⁶, PB1²⁰⁰⁵, PB2⁸²¹ and NP¹⁴⁶ designate amino acid substitutions at PB1³⁹¹, PB1⁵⁸¹, PB1⁶⁶¹, PB2²⁶⁵ and NP³⁴, respectively. Thus, any nucleotide substitutions

resulting in substituted amino acids at these positions are a feature of the invention. Exemplary mutations PB1³⁹¹ (K391E), PB1⁵⁸¹ (E581G), PB1⁶⁶¹ (A661T), PB2²⁶⁵ (N265S) and NP³⁴ (D34G), singly, and more preferably in combination, result in a temperature sensitive phenotype. Simultaneous reversion of these mutations to wild type abolishes the ts phenotype, while introduction of these mutations onto a wild-type background results in virus with a ts phenotype. Consistent with the stability of these phenotypes during passage of the virus, no single change can individually revert the temperature sensitivity profile of the resulting virus to that of wild-type. Rather, these changes appear to act in concert with one another to fully express the ts phenotype. This discovery permits the engineering of additional strains of temperature sensitive influenza A virus suitable for master donor viruses for the production of live attenuated influenza vaccines.

[0129] Similarly, substitutions of individual amino acids in a Master Donor Virus-B strain are correlated with the ts phenotype as illustrated in Table 17. Thus, the methods presented herein are adapted to producing novel influenza B strains with temperature sensitive, and optionally attenuated and/or cold adapted phenotypes by introducing one or more specified mutations into an influenza B genome. For example, one or more mutations resulting in an amino acid substitution at a position selected from among PB2⁶³⁰; PA⁴³¹; PA⁴⁹⁷; N⁵⁵; NP¹¹⁴; NP⁴¹⁰; NP⁵⁰⁹; M1¹⁵⁹ and M1¹⁸³ are introduced into an influenza B strain genome to produce a temperature sensitive influenza B virus. Exemplary amino acid substitutions include the following: PB2⁶³⁰ (S630R); PA⁴³¹ (V431M); PA⁴⁹⁷ (Y497H); NP⁵⁵ (T55A); NP¹¹⁴ (V114A); NP⁴¹⁰ (P410H); NP⁵⁰⁹ (A509T); M1¹⁵⁹ (H159Q) and M1¹⁸³ (M183V).

[0130] Influenza viruses incorporating the mutations of the invention are a feature of the invention regardless of the method in which they are produced. That is, the invention encompasses influenza strains including the mutations of the invention, e.g., any influenza A virus with an amino acid substitution relative to wild type at one or more positions selected from among: PB1³⁹¹, PB1⁵⁸¹, PB1⁶⁶¹, PB2²⁶⁵ and NP³⁴ or any influenza B virus with an amino acid substitution relative to wild type at one or more positions selected from among: PB2⁶³⁰; PA⁴³¹; PA⁴⁹⁷; NP⁵⁵; NP¹¹⁴; NP⁴¹⁰; NP⁵⁰⁹; M1¹⁵⁹ and M1¹⁸³, with the proviso that the strains ca A/Ann Arbor/6/60 and B/Ann Arbor/1/66 are not considered a feature of the present invention. In certain preferred embodiments, the influenza A viruses include a plurality of mutations (e.g., two, or three, or four, or five, or more mutations) selected from among PB1³⁹¹ (K391E), PB1⁵⁸¹ (E581G), PB1⁶⁶¹ (A661T), PB2²⁶⁵ (N265S) and NP³⁴ (D34G); and the influenza B viruses include a plurality of mutations selected from among PB2⁶³⁰ (S630R); PA⁴³¹ (V431M); PA⁴⁹⁷ (Y497H); NP⁵⁵ (T55A); NP¹¹⁴ (V114A); NP⁴¹⁰ (P410H); NP⁵⁰⁹ (A509T); M1¹⁵⁹ (H159Q) and M1¹⁸³ (M183V), respectively. For example, in addition to providing viruses with desired phenotypes relevant for vaccine production, viruses with a subset of mutations, e.g., 1, or 2, or 3, or 4, or 5 selected mutations, are useful in elucidating the contribution of additional mutations to the phenotype of the virus. In certain embodiments, the influenza viruses include at least one additional non-wild type nucleotide (e.g., possibly resulting in an additional amino acid substitution), which optionally refines the desired phenotype or confers a further desirable phenotypic attribute.

[0131] Enhanced Viral Replication

[0132] The present invention also provides a method of introducing of at least one amino acid residue substitution in HA and/or NA to increase the ability of an influenza virus to replicate in embryonated chicken eggs and/or host cells. The invention further provides influenza virus variants with increased ability to replicate in embryonated chicken eggs and/or host cells (referred to herein as “replication enhanced variants”) when compared to HA and/or NA unsubstituted influenza virus. It is specifically contemplated that the method of the invention can be utilized to enhance the replication of an influenza virus in a host cell and that replication enhanced variants may have enhanced replication in chicken eggs and/or host cells. Suitable host cells for the replication of influenza virus include, e.g., Vero cells, Per.C6 cells, BHK cells, MDCK cells, 293 cells and COS cells, including 293T cells, COS7 cells.

[0133] In one embodiment, the method of the invention introduces at least one amino acid substitution into HA and/or NA which will enhance the ability of an influenza virus to replicate in eggs and/or host cells by at least 10%, or by at least 20%, or by at least 30%, or by at least 40%, or by at least 50%, or by at least 60%, or by at least 70%, or by at least 80%, or by at least 90%, or by at least 100%, or by at least 200%, or by at least 300%, or by at least 400%, or by at least 500% when compared to the unmodified influenza virus. It is specifically contemplated that amino acid substitutions may be made in both HA and NA. Preferably, the method of the invention does not significantly alter the antigenicity of the substituted influenza virus when compared to the unsubstituted virus. In a specific embodiment, the method of the invention reduces the antigenicity of the substituted influenza virus when compared to the unsubstituted virus by less than 10%, or by less than 20%, or by less than 30%, or by less than 40%, or by less than 50%, or by less than 60%, or by less than 70%, or by less than 80%, or by less than 90%, or by less than 100%. Methods to determine viral antigenicity are well known in the art (also see, “Example 11” supra).

[0134] In one embodiment, the method of the invention further incorporates an attenuated influenza virus, a cold adapted influenza virus, a temperature sensitive influenza virus, or a virus with any combination of these desirable properties. Preferably, the viruses incorporated by the method of the invention include but are not limited to, influenza B/Ann Arbor/1/66 strain virus, influenza A/Ann Arbor/6/60 strain virus. In another embodiment, the method of the invention introduces vectors including the six internal genes of a viral strain selected for its favorable properties regarding vaccine production, in combination with the genome segments encoding the desired manipulated HA and NA surface antigens to produce influenza viruses with enhanced ability to replicate in embryonated chicken eggs and/or host cells (see, supra and “Example 11”). In another embodiment, the method of the invention further incorporates a non-attenuated influenza virus.

[0135] In one embodiment, the method of the invention introduces at least one amino acid substitution which modulates the receptor binding activity of HA. Receptor binding activity of HA includes but is not limited to the binding of HA to sialic acid residues (e.g., 2,6-linked sialyl-galactosyl moieties [Siaα(2,6)Gal] and 2,3-linked sialyl-galactosyl

moieties [Sia α (2,3)Gal]) present on the cell surface glycoproteins or glycolipids. One method to assay HA binding is presented in "Example 11" (infra), other methods are well known in the art. In another embodiment, the method of the invention introduces amino acid substitutions which modulate the receptor binding specificity of HA for [Sia α (2,6)Gal] and/or [Sia α (2,3)Gal] moieties. Preferably, the method will enhance the binding of HA to [Sia α (2,3)Gal] moieties.

[0136] In a one embodiment, the method of the invention introduces at least one amino acid substitution which enhances the receptor binding activity of HA. Preferably, the receptor binding activity is increased by at least 10%, or by at least 20%, or by at least 30%, or by at least 40%, or by at least 50%, or by at least 60%, or by at least 70%, or by at least 80%, or by at least 90%, or by at least 100%, or by at least 200%.

[0137] In another embodiment, the method of the invention introduces at least one amino acid substitution which reduces the receptor binding activity of HA. Preferably, the receptor binding activity is reduced by at least 10%, or by at least 20%, or by at least 30%, or by at least 40%, or by at least 50%, or by at least 60%, or by at least 70%, or by at least 80%, or by at least 90%, or by at least 100%, or by at least 200%.

[0138] In a preferred embodiment, the method introduces at least one amino acid substitution in HA at positions 183, 186 and/or 226. Preferably, amino acid substitutions are made at positions 183 and 226 or at positions 186 and 226. Most preferably, amino acid substitutions are made such that position 183 is a leucine and position 226 is an alanine or such that position 186 is a valine and position 226 is an isoleucine.

[0139] In one embodiment, the method of the invention introduces at least one amino acid substitution which modulates the neuraminidase activity of NA. Neuraminidase activity of NA includes but is not limited to, the hydrolysis of substrates which contain alpha-ketosidically linked N-acetylneurameric acid (Neu5Ac). Methods to determine the neuraminidase activity are well known in the art (see also, "Example 11" infra).

[0140] In a one embodiment, the method of the invention introduces at least one amino acid substitution which enhances the neuraminidase activity of NA. Preferably, the receptor binding activity is increased by at least 10%, or by at least 20%, or by at least 30%, or by at least 40%, or by at least 50%, or by at least 60%, or by at least 70%, or by at least 80%, or by at least 90%, or by at least 100%, or by at least 200%.

[0141] In another embodiment, the method of the invention introduces at least one amino acid substitution which reduces the neuraminidase activity of NA. Preferably, the neuraminidase activity is reduced by at least 10%, or by at least 20%, or by at least 30%, or by at least 40%, or by at least 50%, or by at least 60%, or by at least 70%, or by at least 80%, or by at least 90%, or by at least 100%, or by at least 200%.

[0142] In a preferred embodiment, the method introduces at least one amino acid substitution in NA at positions 119 and/or 136. Preferably, amino acid substitutions are made such that position 119 is a glutamate and position 136 is a glutamine.

[0143] One skilled in the art would appreciate that in some cases the HA and/or NA protein will already have the preferred amino acid residues at one or more of the aforementioned positions. In this situation, substitution(s) will only be introduced at the remaining non-matching positions.

[0144] It is specifically contemplated that conservative amino acid substitutions may be made for said amino acid substitutions at positions 183, 186 and/or 226 of HA and positions 119 and/or 136 of NA, described supra.

[0145] It is well known in the art that "conservative amino acid substitution" refers to amino acid substitutions that substitute functionally-equivalent amino acids. Conservative amino acid changes result in silent changes in the amino acid sequence of the resulting peptide. For example, one or more amino acids of a similar polarity act as functional equivalents and result in a silent alteration within the amino acid sequence of the peptide. Substitutions that are charge neutral and which replace a residue with a smaller residue may also be considered "conservative substitutions" even if the residues are in different groups (e.g., replacement of phenylalanine with the smaller isoleucine). Families of amino acid residues having similar side chains have been defined in the art. Families of conservative amino acid substitutions include but are not limited to, non-polar (e.g., Trp, Phe, Met, Leu, Ile, Val, Ala, Pro), uncharged polar (e.g., Gly, Ser, Thr, Asn, Gln, Tyr, Cys), acidic/negatively charged (e.g., Asp, Glu), basic/positively charged (e.g., Arg, Lys, His), Beta-branched (e.g., Thr, Val, Ile), residues that influence chain orientation (e.g., Gly, Pro) and aromatic (e.g., Trp, Tyr, Phe, His). The term "conservative amino acid substitution" also refers to the use of amino acid analogs or variants. Guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," (1990, *Science* 247:1306-10).

[0146] In one embodiment, the present invention provides modified influenza viruses, referred to herein as "replication enhanced influenza variant(s)", which incorporate at least one amino acid substitution in HA and/or NA which enhances their replication in embryonated chicken eggs and/or host cells when compared to the unmodified influenza virus. Preferably, the ability of an replication enhanced influenza variant to replicate in eggs and/or host cells has been enhanced by at least 10%, or by at least 20%, or by at least 30%, or by at least 40%, or by at least 50%, or by at least 60%, or by at least 70%, or by at least 80%, or by at least 90%, or by at least 100%, or by at least 200%, or by at least 300%, or by at least 400%, or by at least 500% when compared to the unmodified influenza virus.

[0147] In certain embodiment, a replication enhanced influenza variant further incorporates an attenuated influenza virus, a cold adapted influenza virus, a temperature sensitive influenza virus, or a virus with any combination of these desirable properties. Preferably, the virus incorporated into a replication enhanced influenza variant includes but is not limited to, influenza B/Ann Arbor/1/66 strain virus, influenza A/Ann Arbor/6/60 strain virus. It is specifically contemplated that a replication enhanced influenza variant is produced by introducing vectors including the six internal genes of a viral strain selected for its favorable properties regarding vaccine production, in combination with the genome segments encoding the desired substituted HA and NA surface antigens (see, supra and "Example 11").

[0148] In one embodiment, a replication enhanced influenza variant incorporates at least one amino acid substitution in HA which modulates the receptor binding activity of HA (see supra). Preferably, the method will enhance the binding of HA to [SiaC(2,3)Gal] moieties.

[0149] In a specific embodiment, a replication enhanced influenza variant incorporates at least one amino acid substitution which enhances the receptor binding activity of HA. Preferably, the receptor binding activity is increased by at least 10%, or by at least 20%, or by at least 30%, or by at least 40%, or by at least 50%, or by at least 60%, or by at least 70%, or by at least 80%, or by at least 90%, or by at least 100%, or by at least 200%. It is specifically contemplated that an egg enhance influenza variant does not have significantly altered viral antigenicity when compared to the unsubstituted influenza virus. In a specific embodiment, a replication enhanced influenza variant has an antigenicity that is reduced by less than 10%, or by less than 20%, or by less than 30%, or by less than 40%, or by less than 50%, or by less than 60%, or by less than 70%, or by less than 80%, or by less than 90%, or by less than 100% when compared to the unsubstituted virus. Methods to determine viral antigenicity are well known in the art (also see, "Example 11" supra).

[0150] In another embodiment, a replication enhanced influenza variant incorporates incorporate at least one amino acid substitution which reduces the receptor binding activity of HA. Preferably, the receptor binding activity is reduced by at least 10%, or by at least 20%, or by at least 30%, or by at least 40%, or by at least 50%, or by at least 60%, or by at least 70%, or by at least 80%, or by at least 90%, or by at least 100%, or by at least 200%.

[0151] In a preferred embodiment, a replication enhanced influenza variant incorporates incorporate at least one amino acid substitution in HA at positions 183, 186 and/or 226. Preferably, amino acid substitutions are present at positions 183 and 226 or at positions 186 and 226. Most preferably, amino acid substitutions are present such that position 183 is a leucine and position 226 is an alanine or such that position 186 is a valine and position 226 is an isoleucine.

[0152] In one embodiment, a replication enhanced influenza variant incorporates at least one amino acid substitution which modulates the neuraminidase activity of NA (see supra).

[0153] In a one embodiment, a replication enhanced influenza variant incorporates at least one amino acid substitution which enhances the neuraminidase activity of NA. Preferably, the receptor binding activity is increased by at least 10%, or by at least 20%, or by at least 30%, or by at least 40%, or by at least 50%, or by at least 60%, or by at least 70%, or by at least 80%, or by at least 90%, or by at least 100%, or by at least 200%.

[0154] In a another embodiment, a replication enhanced influenza variant incorporates at least one amino acid substitution which reduces the neuraminidase activity of NA. Preferably, the neuraminidase activity is reduced by at least 10%, or by at least 20%, or by at least 30%, or by at least 40%, or by at least 50%, or by at least 60%, or by at least 70%, or by at least 80%, or by at least 90%, or by at least 100%, or by at least 200%.

[0155] In a preferred embodiment, a replication enhanced influenza variant incorporates at least one amino acid substitution in NA at positions 119 and/or 136. Preferably, amino acid substitutions are made such that position 119 is a is a glutamate and position 136 is a glutamine.

[0156] Cell Culture

[0157] Typically, propagation of the virus is accomplished in the media compositions in which the host cell is commonly cultured. Suitable host cells for the replication of influenza virus include, e.g., Vero cells, Per.C6 cells, BHK cells, MDCK cells, 293 cells and COS cells, including 293T cells, COS7 cells. Commonly, co-cultures including two of the above cell lines, e.g., MDCK cells and either 293T or COS cells are employed at a ratio, e.g., of 1:1, to improve replication efficiency. Typically, cells are cultured in a standard commercial culture medium, such as Dulbecco's modified Eagle's medium supplemented with serum (e.g., 10% fetal bovine serum), or in serum free medium, under controlled humidity and CO₂ concentration suitable for maintaining neutral buffered pH (e.g., at pH between 7.0 and 7.2). Optionally, the medium contains antibiotics to prevent bacterial growth, e.g., penicillin, streptomycin, etc., and/or additional nutrients, such as L-glutamine, sodium pyruvate, non-essential amino acids, additional supplements to promote favorable growth characteristics, e.g., trypsin, β-mercaptoethanol, and the like.

[0158] Procedures for maintaining mammalian cells in culture have been extensively reported, and are known to those of skill in the art. General protocols are provided, e.g., in Freshney (1983) *Culture of Animal Cells: Manual of Basic Technique*, Alan R. Liss, New York; Paul (1975) *Cell and Tissue Culture*, 5th ed., Livingston, Edinburgh; Adams (1980) *Laboratory Techniques in Biochemistry and Molecular Biology-Cell Culture for Biochemists*, Work and Burdon (eds.) Elsevier, Amsterdam. Additional details regarding tissue culture procedures of particular interest in the production of influenza virus in vitro include, e.g., Merten et al. (1996) Production of influenza virus in cell cultures for vaccine preparation. In Cohen and Shafferman (eds) *Novel Strategies in Design and Production of Vaccines*, which is incorporated herein in its entirety. Additionally, variations in such procedures adapted to the present invention are readily determined through routine experimentation.

[0159] Cells for production of influenza virus can be cultured in serum-containing or serum free medium. In some case, e.g., for the preparation of purified viruses, it is desirable to grow the host cells in serum free conditions. Cells can be cultured in small scale, e.g., less than 25 ml medium, culture tubes or flasks or in large flasks with agitation, in rotator bottles, or on microcarrier beads (e.g., DEAE-Dextran microcarrier beads, such as Dormacell, Pfeiffer & Langen; Superbead, Flow Laboratories; styrene copolymer-tri-methylamine beads, such as Hillex, SoloHill, Ann Arbor) in flasks, bottles or reactor cultures. Microcarrier beads are small spheres (in the range of 100-200 microns in diameter) that provide a large surface area for adherent cell growth per volume of cell culture. For example a single liter of medium can include more than 20 million microcarrier beads providing greater than 8000 square centimeters of growth surface. For commercial production of viruses, e.g., for vaccine production, it is often desirable to culture the cells in a bioreactor or fermenter. Bioreactors are available

in volumes from under 1 liter to in excess of 100 liters, e.g., Cyto3 Bioreactor (Osmonics, Minnetonka, Minn.); NBS bioreactors (New Brunswick Scientific, Edison, N.J.); laboratory and commercial scale bioreactors from B. Braun Biotech International (B. Braun Biotech, Melsungen, Germany).

[0160] Regardless of the culture volume, in the context of the present invention, it is important that the cultures be maintained at a temperature less than or equal to 35° C., to insure efficient recovery of recombinant and/or reassortant influenza virus using the multi plasmid system described herein. For example, the cells are cultured at a temperature between about 32° C. and 35° C., typically at a temperature between about 32° C. and about 34° C., usually at about 33° C.

[0161] Typically, a regulator, e.g., a thermostat, or other device for sensing and maintaining the temperature of the cell culture system is employed to insure that the temperature does not exceed 35° C. during the period of virus replication.

[0162] Introduction of Vectors into Host Cells

[0163] Vectors comprising influenza genome segments are introduced (e.g., transfected) into host cells according to methods well known in the art for introducing heterologous nucleic acids into eukaryotic cells, including, e.g., calcium phosphate co-precipitation, electroporation, microinjection, lipofection, and transfection employing polyamine transfection reagents. For example, vectors, e.g., plasmids, can be transfected into host cells, such as COS cells, 293T cells or combinations of COS or 293T cells and MDCK cells, using the polyamine transfection reagent TransIT-LT1 (Mirus) according to the manufacturer's instructions. Approximately 1 μ g of each vector to be introduced into the population of host cells with approximately 2 μ l of TransIT-LT1 diluted in 160 μ l medium, preferably serum-free medium, in a total vol. of 200 μ l. The DNA:transfection reagent mixtures are incubated at room temperature for 45 min followed by addition of 800 μ l of medium. The transfection mixture is added to the host cells, and the cells are cultured as described above. Accordingly, for the production of recombinant or reassortant viruses in cell culture, vectors incorporating each of the 8 genome segments, (PB2, PB1, PA, NP, M, NS, HA and NA) are mixed with approximately 20 μ l TransIT-LT1 and transfected into host cells. Optionally, serum-containing medium is replaced prior to transfection with serum-free medium, e.g., Opti-MEM I, and incubated for 4-6 hours.

[0164] Alternatively, electroporation can be employed to introduce vectors incorporating influenza genome segments into host cells. For example, plasmid vectors incorporating an influenza A or influenza B virus are favorably introduced into Vero cells using electroporation according to the following procedure. In brief, 5×10^6 Vero cells, e.g., grown in Modified Eagle's Medium (MEM) supplemented with 10% Fetal Bovine Serum (FBS) are resuspended in 0.4 ml Opti-MEM and placed in an electroporation cuvette. Twenty micrograms of DNA in a volume of up to 25 μ l is added to the cells in the cuvette, which is then mixed gently by tapping. Electroporation is performed according to the manufacturer's instructions (e.g., BioRad Gene Pulser II with Capacitance Extender Plus connected) at 300 volts, 950 microFarads with a time constant of between 28-33 msec. The cells are remixed by gently tapping and approximately

1-2 minutes following electroporation 0.7 ml MEM with 10% FBS is added directly to the cuvette. The cells are then transferred to two wells of a standard 6 well tissue culture dish containing 2 ml MEM, 10% FBS or OPTI-MEM without serum. The cuvette is washed to recover any remaining cells and the wash suspension is divided between the two wells. Final volume is approximately 3.5 mls. The cells are then incubated under conditions permissive for viral growth, e.g., at approximately 33° C. for cold adapted strains.

[0165] Recovery of Viruses

[0166] Viruses are typically recovered from the culture medium, in which infected (transfected) cells have been grown. Typically crude medium is clarified prior to concentration of influenza viruses. Common methods include filtration, ultrafiltration, adsorption on barium sulfate and elution, and centrifugation. For example, crude medium from infected cultures can first be clarified by centrifugation at, e.g., 1000-2000xg for a time sufficient to remove cell debris and other large particulate matter, e.g., between 10 and 30 minutes. Alternatively, the medium is filtered through a 0.8 μ m cellulose acetate filter to remove intact cells and other large particulate matter. Optionally, the clarified medium supernatant is then centrifuged to pellet the influenza viruses, e.g., at 15,000xg, for approximately 3-5 hours. Following resuspension of the virus pellet in an appropriate buffer, such as STE (0.01 M Tris-HCl; 0.15 M NaCl; 0.0001 M EDTA) or phosphate buffered saline (PBS) at pH 7.4, the virus is concentrated by density gradient centrifugation on sucrose (60%-12%) or potassium tartrate (50%-10%). Either continuous or step gradients, e.g., a sucrose gradient between 12% and 60% in four 12% steps, are suitable. The gradients are centrifuged at a speed, and for a time, sufficient for the viruses to concentrate into a visible band for recovery. Alternatively, and for most large scale commercial applications, virus is elutriated from density gradients using a zonal-centrifuge rotor operating in continuous mode. Additional details sufficient to guide one of skill through the preparation of influenza viruses from tissue culture are provided, e.g., in Furminger. *Vaccine Production*, in Nicholson et al. (eds) *Textbook of Influenza* pp. 324-332; Merten et al. (1996) Production of influenza virus in cell cultures for vaccine preparation, in Cohen & Shafferman (eds) *Novel Strategies in Design and Production of Vaccines* pp. 141-151, and U.S. Pat. No. 5,690,937. If desired, the recovered viruses can be stored at -80° C. in the presence of sucrose-phosphate-glutamate (SPG) as a stabilizer

[0167] Methods and Compositions for Prophylactic Administration of Vaccines

[0168] Recombinant and reassortant viruses of the invention can be administered prophylactically in an appropriate carrier or excipient to stimulate an immune response specific for one or more strains of influenza virus. Typically, the carrier or excipient is a pharmaceutically acceptable carrier or excipient, such as sterile water, aqueous saline solution, aqueous buffered saline solutions, aqueous dextrose solutions, aqueous glycerol solutions, ethanol, allantoic fluid from uninfected Hens' eggs (i.e., normal allantoic fluid "NAF") or combinations thereof. The preparation of such solutions insuring sterility, pH, isotonicity, and stability is effected according to protocols established in the art. Generally, a carrier or excipient is selected to minimize allergic

and other undesirable effects, and to suit the particular route of administration, e.g., subcutaneous, intramuscular, intranasal, etc.

[0169] Generally, the influenza viruses of the invention are administered in a quantity sufficient to stimulate an immune response specific for one or more strains of influenza virus. Preferably, administration of the influenza viruses elicits a protective immune response. Dosages and methods for eliciting a protective immune response against one or more influenza strains are known to those of skill in the art. For example, inactivated influenza viruses are provided in the range of about 1-1000 HID₅₀ (human infectious dose), i.e., about 10⁵-10⁸ pfu (plaque forming units) per dose administered. Alternatively, about 10-50 µg, e.g., about 15 µg HA is administered without an adjuvant, with smaller doses being administered with an adjuvant. Typically, the dose will be adjusted within this range based on, e.g., age, physical condition, body weight, sex, diet, time of administration, and other clinical factors. The prophylactic vaccine formulation is systemically administered, e.g., by subcutaneous or intramuscular injection using a needle and syringe, or a needleless injection device. Alternatively, the vaccine formulation is administered intranasally, either by drops, large particle aerosol (greater than about 10 microns), or spray into the upper respiratory tract. While any of the above routes of delivery results in a protective systemic immune response, intranasal administration confers the added benefit of eliciting mucosal immunity at the site of entry of the influenza virus. For intranasal administration, attenuated live virus vaccines are often preferred, e.g., an attenuated, cold adapted and/or temperature sensitive recombinant or reassortant influenza virus. While stimulation of a protective immune response with a single dose is preferred, additional dosages can be administered, by the same or different route, to achieve the desired prophylactic effect.

[0170] Alternatively, an immune response can be stimulated by ex vivo or in vivo targeting of dendritic cells with influenza viruses. For example, proliferating dendritic cells are exposed to viruses in a sufficient amount and for a sufficient period of time to permit capture of the influenza antigens by the dendritic cells. The cells are then transferred into a subject to be vaccinated by standard intravenous transplantation methods.

[0171] Optionally, the formulation for prophylactic administration of the influenza viruses, or subunits thereof, also contains one or more adjuvants for enhancing the immune response to the influenza antigens. Suitable adjuvants include: saponin, mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil or hydrocarbon emulsions, bacille Calmette-Guerin (BCG), *Corynebacterium parvum*, and the synthetic adjuvants QS-21 and MF59.

[0172] If desired, prophylactic vaccine administration of influenza viruses can be performed in conjunction with administration of one or more immunostimulatory molecules. Immunostimulatory molecules include various cytokines, lymphokines and chemokines with immunostimulatory, immunopotentiating, and pro-inflammatory activities, such as interleukins (e.g., IL-1, IL-2, IL-3, IL-4, IL-12, IL-13); growth factors (e.g., granulocyte-macrophage (GM)-colony stimulating factor (CSF)); and other immunostimulatory molecules, such as macrophage inflammatory

factor, Flt3 ligand, B7.1; B7.2, etc. The immunostimulatory molecules can be administered in the same formulation as the influenza viruses, or can be administered separately. Either the protein or an expression vector encoding the protein can be administered to produce an immunostimulatory effect.

[0173] In another embodiment, the vectors of the invention including influenza genome segments can be employed to introduce heterologous nucleic acids into a host organism or host cell, such as a mammalian cell, e.g., cells derived from a human subject, in combination with a suitable pharmaceutical carrier or excipient as described above. Typically, the heterologous nucleic acid is inserted into a non-essential region of a gene or gene segment, e.g., the M gene of segment 7. The heterologous polynucleotide sequence can encode a polypeptide or peptide, or an RNA such as an antisense RNA or ribozyme. The heterologous nucleic acid is then introduced into a host or host cells by producing recombinant viruses incorporating the heterologous nucleic, and the viruses are administered as described above.

[0174] Alternatively, a vector of the invention including a heterologous nucleic acid can be introduced and expressed in a host cells by co-transfected the vector into a cell infected with an influenza virus. Optionally, the cells are then returned or delivered to the subject, typically to the site from which they were obtained. In some applications, the cells are grafted onto a tissue, organ, or system site (as described above) of interest, using established cell transfer or grafting procedures. For example, stem cells of the hematopoietic lineage, such as bone marrow, cord blood, or peripheral blood derived hematopoietic stem cells can be delivered to a subject using standard delivery or transfusion techniques.

[0175] Alternatively, the viruses comprising a heterologous nucleic acid can be delivered to the cells of a subject in vivo. Typically, such methods involve the administration of vector particles to a target cell population (e.g., blood cells, skin cells, liver cells, neural (including brain) cells, kidney cells, uterine cells, muscle cells, intestinal cells, cervical cells, vaginal cells, prostate cells, etc., as well as tumor cells derived from a variety of cells, tissues and/or organs. Administration can be either systemic, e.g., by intravenous administration of viral particles, or by delivering the viral particles directly to a site or sites of interest by a variety of methods, including injection (e.g., using a needle or syringe), needleless vaccine delivery, topical administration, or pushing into a tissue, organ or skin site. For example, the viral vector particles can be delivered by inhalation, orally, intravenously, subcutaneously, subdermally, intradermally, intramuscularly, intraperitoneally, intrathecally, by vaginal or rectal administration, or by placing the viral particles within a cavity or other site of the body, e.g., during surgery.

[0176] The above described methods are useful for therapeutically and/or prophylactically treating a disease or disorder by introducing a vector of the invention comprising a heterologous polynucleotide encoding a therapeutically or prophylactically effective polypeptide (or peptide) or RNA (e.g., an antisense RNA or ribozyme) into a population of target cells in vitro, ex vivo or in vivo. Typically, the polynucleotide encoding the polypeptide (or peptide), or

RNA, of interest is operably linked to appropriate regulatory sequences as described above in the sections entitled "Expression Vectors" and "Additional Expression Elements." Optionally, more than one heterologous coding sequence is incorporated into a single vector or virus. For example, in addition to a polynucleotide encoding a therapeutically or prophylactically active polypeptide or RNA, the vector can also include additional therapeutic or prophylactic polypeptides, e.g., antigens, co-stimulatory molecules, cytokines, antibodies, etc., and/or markers, and the like.

[0177] The methods and vectors of the present invention can be used to therapeutically or prophylactically treat a wide variety of disorders, including genetic and acquired disorders, e.g., as vaccines for infectious diseases, due to viruses, bacteria, and the like.

[0178] Kits

[0179] To facilitate use of the vectors and vector systems of the invention, any of the vectors, e.g., consensus influenza virus plasmids, variant influenza polypeptide plasmids, influenza polypeptide library plasmids, etc., and additional components, such as, buffer, cells, culture medium, useful for packaging and infection of influenza viruses for experimental or therapeutic purposes, can be packaged in the form of a kit. Typically, the kit contains, in addition to the above components, additional materials which can include, e.g., instructions for performing the methods of the invention, packaging material, and a container.

[0180] Manipulation of Viral Nucleic Acids and Proteins

[0181] In the context of the invention, influenza virus nucleic acids and/or proteins are manipulated according to well known molecular biology techniques. Detailed protocols for numerous such procedures, including amplification, cloning, mutagenesis, transformation, and the like, are described in, e.g., in Ausubel et al. *Current Protocols in Molecular Biology* (supplemented through 2000) John Wiley & Sons, New York ("Ausubel"); Sambrook et al. *Molecular Cloning—A Laboratory Manual* (2nd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1989 ("Sambrook"), and Berger and Kimmel *Guide to Molecular Cloning Techniques, Methods in Enzymology* volume 152 Academic Press, Inc., San Diego, Calif. ("Berger").

[0182] In addition to the above references, protocols for in vitro amplification techniques, such as the polymerase chain reaction (PCR), the ligase chain reaction (LCR), Q β -replicase amplification, and other RNA polymerase mediated techniques (e.g., NASBA), useful e.g., for amplifying cDNA probes of the invention, are found in Mullis et al. (1987) U.S. Pat. No. 4,683,202; *PCR Protocols A Guide to Methods and Applications* (Innis et al. eds) Academic Press Inc. San Diego, Calif. (1990) ("Innis"); Arnheim and Levinson (1990) *C&EN* 36; *The Journal Of NIH Research* (1991) 3:81; Kwok et al. (1989) *Proc Natl Acad Sci USA* 86, 1173; Guatelli et al. (1990) *Proc Natl Acad Sci USA* 87:1874; Lomell et al. (1989) *J Clin Chem* 35:1826; Landegren et al. (1988) *Science* 241:1077; Van Brunt (1990) *Biotechnology* 8:291; Wu and Wallace (1989) *Gene* 4: 560; Barringer et al. (1990) *Gene* 89:117, and Sooknanan and Malek (1995) *Biotechnology* 13:563. Additional methods, useful for cloning nucleic acids in the context of the present invention,

include Wallace et al. U.S. Pat. No. 5,426,039. Improved methods of amplifying large nucleic acids by PCR are summarized in Cheng et al. (1994) *Nature* 369:684 and the references therein.

[0183] Certain polynucleotides of the invention, e.g., oligonucleotides can be synthesized utilizing various solid-phase strategies including mononucleotide- and/or trinucleotide-based phosphoramidite coupling chemistry. For example, nucleic acid sequences can be synthesized by the sequential addition of activated monomers and/or trimers to an elongating polynucleotide chain. See e.g., Caruthers, M. H. et al. (1992) *Meth Enzymol* 211:3.

[0184] In lieu of synthesizing the desired sequences, essentially any nucleic acid can be custom ordered from any of a variety of commercial sources, such as The Midland Certified Reagent Company (mcrc@oligos.com), The Great American Gene Company (www.genco.com), ExpressGen, Inc. (www.expressgen.com), Operon Technologies, Inc. (www.operon.com), and many others.

[0185] In addition, substitutions of selected amino acid residues in viral polypeptides can be accomplished by, e.g., site directed mutagenesis. For example, viral polypeptides with amino acid substitutions functionally correlated with desirable phenotypic characteristic, e.g., an attenuated phenotype, cold adaptation, temperature sensitivity, can be produced by introducing specific mutations into a viral nucleic acid segment encoding the polypeptide. Methods for site directed mutagenesis are well known in the art, and described, e.g., in Ausubel, Sambrook, and Berger, *supra*. Numerous kits for performing site directed mutagenesis are commercially available, e.g., the Chameleon Site Directed Mutagenesis Kit (Stratagene, La Jolla), and can be used according to the manufacturers instructions to introduce, e.g., one or more amino acid substitutions described in Table 6 or Table 17, into a genome segment encoding a influenza A or B polypeptide, respectively.

EXAMPLES

Example 1

Construction of pAD3000

[0186] The plasmid pHW2000 (Hoffmann et al. (2000) A DNA transfection system for generation of influenza A virus from eight plasmids *Proc Natl Acad Sci USA* 97:6108-6113) was modified to replace the bovine growth hormone (BGH) polyadenylation signals with a polyadenylation signal sequences derived from Simian virus 40 (SV40).

[0187] Sequences derived from SV40 were amplified with Taq MasterMix (Qiagen) using the following oligonucleotides, designated in the 5' to 3' direction:

(SEQ ID NO:1)

polyA.1:
AACAAATTGAGATCTCGGTACACCTCAGACATGATAAGATACTTGTGAGT

(SEQ ID NO:2)

polyA.2:
TATAACTGCAGACTAGTGTATCCTTGTATTGCAGCTTATAATGGTTA

[0188] The plasmid pSV2His was used as a template. A fragment consistent with the predicted 175 bp product was obtained and cloned into pcDNA3.1, using a Topo TA cloning vector (Invitrogen) according to the manufacturer's directions. The desired 138 bp fragment containing the SV40 polyadenylation signals was excised from the resulting plasmid with EcoRV and BstEII, isolated from an agarose gel, and ligated between the unique PvuII and BstEI sites in pHW2000 using conventional techniques (see, e.g., Ausubel, Berger, Sambrook). The resulting plasmid, pAD3000 (FIG. 1), was sequenced and found to contain the SV40 polyadenylation site in the correct orientation. Nucleotides 295-423 in pAD3000 correspond to nucleotides 2466-2594, respectively, in SV40 strain 777 (AF332562).

Example 2

Eight Plasmid System for Production of MDV-A

[0189] A cold-adapted influenza virus type A strain A/AA/6/60 variant has commonly been used as a master donor virus for the production of nasally administered Influenza A vaccines. This strain is an exemplary Master Donor Virus (MDV) in the context of the present invention. For simplicity, this strain A/AA/6/60 variant is designated herein MDV-A. MDV-A viral RNA was extracted using the RNeasy mini kit (Qiagen) and the eight corresponding cDNA fragments were amplified by RT-PCR using the primers listed in Table 1.

TABLE 1

Sequence of the primers used for cloning MDV-A eight segments		
SEQ ID.	Primer	Sequence (5'-3')
<u>MDV-A FORWARD PRIMERS</u>		
SEQ ID AarI PB2	long	CAC TTA TAT TCA CCT GCC TCA NO:3 GGG AGC GAA AGC AGG TC
SEQ ID BsmBI-PB1		TAT TCG TCT CAG GGA GCG AAA NO:4 GCA GGC AAA
SEQ ID BsmBI-PA		TAT TCG TCT CAG GGA GCG AAA NO:5 GCA GGT ACT
SEQ ID BsmBI-NP		TAT TCG TCT CAG GGA GCA AAA NO:6 GCA GGG TAG A
SEQ ID AarI HA-long		CAC TTA TAT TCA CCT GCC TCA NO:7 GGG AGC AAA AGC AGG GG
SEQ ID BsmBI-NA		TAT TCG TCT CAG GGA GCA AAA NO:8 GCA GGA GTG A
SEQ ID BsmBI-M		TAT TCG TCT CAG GGA GCA AAA NO:9 GCA GGT AGA T
SEQ ID BsmBI-NS		TAT TCG TCT CAG GGA GCA AAA NO:10 GCA GGG TGA
<u>MDV-A REVERSE PRIMERS</u>		
SEQ ID AarI PB2-long		CCT AAC ATA TCA CCT GCC TCG NO:11 TAT TAG TAG AAA CAA GGT CGT TT
SEQ ID BsmBI-PB1		ATA TCG TCT CGT ATT AGT AGA NO:12 AAC AAG GCA TTT

TABLE 1-continued

Sequence of the primers used for cloning MDV-A eight segments		
SEQ ID.	Primer	Sequence (5'-3')
<u>MDV-A FORWARD PRIMERS</u>		
SEQ ID BsmBI-PA		ATA TCG TCT CGT ATT AGT AGA NO:13 AAC AAG GTA CTT
SEQ ID BsmBI-NP		ATA TCG TCT CGT ATT AGT AGA NO:14 AAC AAG GGT ATT
SEQ ID AarI HA-long		CCT AAC ATA TCA CCT GCC TCG NO:15 TAT TAG TAG AAA CAA GGG TGT T
SEQ ID BsmBI-NA		ATA TCG TCT CGT ATT AGT AGA NO:16 AAC AAG GAG TTT
SEQ ID BsmBI-M		ATA TCG TCT CGT ATT AGT AGA NO:17 AAC AAG GTA GTT
SEQ ID BsmBI-NS		ATA TCG TCT CGT ATT AGT AGA NO:18 AAC AAG GGT GTT

[0190] With the exception of the influenza genome segments encoding HA and PB2, which were amplified using the primers containing Aar I restriction enzyme recognition site, the remaining 6 genes were amplified with primers containing the BsmB I restriction enzyme recognition site. Both AarI and BsmB I cDNA fragments were cloned between the two BsmB I sites of the pAD3000 vector.

[0191] Sequencing analysis revealed that all of the cloned cDNA fragments contained mutations with respect to the consensus MDV-A sequence, which were likely introduced during the cloning steps. The mutations found in each gene segment are summarized in Table 2.

TABLE 2

Mutations introduced into the MDV-A clones in pAD3000		
Gene segment	Mutation positions (nt)	Amino acid changes
PB2	A954(G/C/T), G1066A, T1580C, T1821C	Silent, Gly to Ser, Val to Ala, Silent
PB1	C1117T	Arg to Stop
PA	G742A, A1163G, A1615G, T1748C, C2229del	Gly to Ser, Asp to Gly, Arg to Gly, Met to Thr, non-coding
HA	A902C, C1493T	Asn to His, Cys to Arg
NP	C113A, T1008C	Thr to Asn, silent
NA	C1422T	Pro to Leu
M	A191G	Thr to Ala
NS	C38T	Silent

[0192] All the mutations were corrected back to the consensus MDV-A sequence using a QuikChange Site-directed Mutagenesis Kit (Stratagene) and synthetic oligonucleotide primers as shown in Table 3.

TABLE 3

Primers used for correcting the mutations in the MDV-A clones			
HJ67	PB2A954G	5/P/gcaagctgtggaaatatgcaaggc (SEQ ID NO:19)	
HJ68	PB2A954G.as	gccttgcataattccacagcttgc (SEQ ID NO:20)	
HJ69	PB2G1066A	5/P/gaagtgcitacgggcaatttcaaac (SEQ ID NO:21)	
PB2	HJ70	PB2G1066A.as	gtttgaqaattggccgtaaagcacttc (SEQ ID NO:22)
	HJ71	PB2T1580A	5/P/cctgaggaggctcgtgaaacac (SEQ ID NO:23)
	HJ72	PB2T1580A.as	gtgtttcactgacctccctcagg (SEQ ID NO:24)
	HJ73	PB21821C	5/P/gtttgttagacttccaaac (SEQ ID NO:25)
	HJ74	PB21821C.as	gttggaaatagactcataacaac (SEQ ID NO:26)
PB1	HJ75	PB1C1117T	gacagtaagctccgaacacaaaatac (SEQ ID NO:27)
	HJ76	PB1C1117T.as	gtatttgttgcggagcttcatgc (SEQ ID NO:28)
	HJ77	PA-G742A	5/P/cgaaccgaaacggctacattgggg (SEQ ID NO:29)
	HJ78	PA-G742A.as	ccctcaatgtagccgttgcgttcg (SEQ ID NO:30)
	HJ79	PA-A1163G	5/P/cagagaaggtagattgcgactg (SEQ ID NO:31)
	HJ80	PA-A1163G.as	cagtcgtcaaaagtctacccctctg (SEQ ID NO:32)
PA	HJ81	PA-A1615G	5/P/caactgacccaagacttgagccac (SEQ ID NO:33)
	HJ82	PA-A1615G.as	gtggctcaagtcttggctcgtg (SEQ ID NO:34)
	HJ83	PA-T1748C	5/P/caaagattaaatgaaatqggaaatg (SEQ ID NO:35)
	HJ84	PA-T1748C.as	cattccccatttcattttatctttg (SEQ ID NO:36)
	HJ85	PA-C2229	5/P/gtaccttgcattactaataaccgg (SEQ ID NO:37)
	HJ86	PA-C2230.as	ccgggttattatgttagaaacaaggatc (SEQ ID NO:38)
	HJ87	HA-A902C	5/P/ggaacacttgagaactgtgagacc (SEQ ID NO:39)
HA	HJ88	HA-A902C.as	ggtctcacagtttcaagtgttcc (SEQ ID NO:40)
	HJ89	HA-C1493T	5/P/gaattttatcacaatgtgtatgaaatg (SEQ ID NO:41)
	HJ90	HA-C1493T.as	cattcatcatcataattgtgataaaaattc (SEQ ID NO:42)
	HJ91	NP-C113A	5/P/gccaaatgcaactgaaatcagagc (SEQ ID NO:43)
NP	HJ92	NP-C113A.as	gctctgatttcagtttcatctggc (SEQ ID NO:44)
	HJ93	NP-T1008C	5/P/ccgaatgagaatccagcacacaag (SEQ ID NO:45)
	HJ94	NP-T1008C.as	cttgcgtgtgattctcattcgg (SEQ ID NO:46)
	HJ95	NA-C1422T	catcaatttcatgcctatataagcttgc (SEQ ID NO:47)
NS	HJ96	NA-C1422T.as	gaaagcttatataaggcatgaaatttgatg (SEQ ID NO:48)
	HJ97	NS-C38T	cataatggatcctaacaactgtgtcaagc (SEQ ID NO:49)
	HJ98	NS-C38T.as	gcttgacacagtgttagatccattatg (SEQ ID NO:50)
PA	HJ99	PAGC375T	ggagaatagattcatcgagattggag (SEQ ID NO:51)
	HJ100	PA6C375T.as	ctccaaatctcgatgaatcttcc (SEQ ID NO:52)

Example 3

Generation of Infectious Recombinant MDV-A and Reassorted Influenza Virus

[0193] Madin-Darby canine kidney (MDCK) cells and human COS7 cells were maintained in modified Eagle Medium (MEM) containing 10% fetal bovine serum (FBS). Human embryonic kidney cells (293T) were maintained in Opti-MEM I (Life Technologies) containing 5% FBS. MDCK and either COS7 or 293T cells were co-cultured in 6-well plates at a ratio of 1:1 and the cells were used for transfection at a confluence of approximately 80%. 293T and COS7 cells have a high transfection efficiency, but are not permissive for influenza virus replication. Co-culture with MDCK cells ensures efficient replication of the recombinant viruses. Prior to transfection, serum-containing media were replaced with serum free medium (Opti-MEM I) and incubated for 4-6 hours. Plasmid DNA transfection was performed using TransIT-LT1 (Mirus) by mixing 1 μ g of each of the 8 plasmid DNAs (PB2, PB1, PA, NP, M, NS, HA and NA) with 20 μ l of TransIT-LT1 diluted in 160 μ l Opti-MEM I in a total volume of 200 μ l. The DNA:transfection reagent mixtures were incubated at room temperature for 45 min followed by addition of 800 μ l of Opti-MEM I. The transfection mixture was then added to the co-cultured MDCK/293T or MDCK/COS7 cells. The transfected cells

were incubated at 35° C. or 33° C. for between 6 hours and 24 hours, e.g., overnight, and the transfection mixture was replaced with 1 ml of Opti-MEM I in each well. After incubation at 35° C. or 33° C. for 24 hours, 1 ml of Opti-MEM I containing 1 μ g/ml TPCK-trypsin was added to each well and incubated for an additional 12 hours. The recovered virus was then amplified in confluent MDCK cells or directly amplified in embryonated chick eggs. MDCK cells in 12-well plate were infected with 0.2 ml of the transfection mixture for 1 hour at room temperature, the mixture was then removed and replaced with 2 ml of Opti-MEM I containing 1 μ g/ml TPCK-trypsin. The cells were incubated at 35° C. or 33° C. for 3-4 days. The amplified viruses were stored at -80° C. in the presence of SPG stabilizer or plaque-purified and amplified in MDCK cells or chicken embryonic eggs.

[0194] Functional Expression of MDV-A Polymerase Proteins

[0195] Functional activity of the four MDV-A polymerase proteins, PB2, PB1, PA and NP, were analyzed by their ability to replicate an influenza virus minigenome encoding an EGFP reporter gene. A set of 8 expression plasmids (see, e.g., Table 4) (Hoffmann et al. (2001) Eight plasmid rescue system for influenza A virus; Options for the control of influenza International Congress Series 1219:1007-1013) that contained the cDNAs of A/PR/8/34 strain (H1N1) and

an influenza virus minigenome containing a reporter gene encoding the enhanced green fluorescent protein (EGFP), pHW72-EGFP).

[0196] The MDV-A PB1, PB2, PA and NP or PB1, PA, NP (-PB2 as a negative control) were transfected into the co-cultured MDCK/293T cells together with a plasmid representing an influenza A virus EGFP minigenome (pHW72-EGFP)(Hoffmann et al. (2000) "Ambisense" approach for the generation of influenza A virus. vRNA and mRNA synthesis from one template *Virology* 15:267(2):310-7). The transfected cells were observed under phase contrast microscope or fluorescence microscope at 48 hours post-transfection. Alternatively, flow cytometry can be employed to detect EGFP expression.

[0197] As shown in FIG. 2, green fluorescence, indicating expression of the EGFP minigenome was observed in the cells transfected with PB2, PB1, PA and NP of MDV-A, but not in the cells transfected with only three polymerase proteins. This indicated that the MDV-A polymerase proteins in pAD3000 were functional.

[0198] In other assays a minigenome including the chloramphenicol acetyl transferase (CAT) gene, designated pFlu-CAT is utilized to measure polymerase activity. In such an assay, CAT expression is measured at the protein (e.g., by ELISA) or RNA level, as an indicator of minigenome replication.

[0199] Analysis of the MDV-A Plasmids by Single Gene Reassortant Experiment

[0200] Each of the 8 MDV-A genome segments cloned in pAD3000 was shown to be functionally expressed in a reassortant experiment by co-transfected a single gene segment from MDA-A together with the complementary seven segments from control A/PR/8/34 strain. All eight single genome segment plasmids in combination with complementary control segments generated infectious reassortant virus, which caused cytopathic effects in infected MDCK cells, indicating that all eight plasmids encode functional MDV-A proteins.

TABLE 4

Recovery of 7 + 1 reassortants by plasmids				
Virus gene segment	PB2	PB1	PA	NP
1	PMDV-A-PB2	pHW191-PB2	pHW191-PB2	pHW191-PB2
2	PHW192-PB1	pMDV-A-PB1	pHW192-PB1	pHW192-PB1
3	PHW193-PA	pHW193-PA	pMDV-A-PA	pHW193-PA
4	PHW195-NP	pHW195-NP	pHW195-NP	pMDV-A-NP
5	PHW197-M	pHW197-M	pHW197-M	pHW197-M
6	PHW198-NS	pHW198-NS	pHW198-NS	pHW198-NS
7	PHW194-HA	pHW194-HA	pHW194-HA	pHW194-HA
8	PHW-196-NA	pHW-196-NA	pHW-196-NA	pHW-196-NA
CPE (+)	(+)	(+)	(+)	(+)

TABLE 4-continued

Recovery of 7 + 1 reassortants by plasmids

Virus gene segment M	NS	HA	NA
1 PHW191-PB2	pHW191-PB2	pHW191-PB2	pHW191-PB2
2 PHW192-PB1	pHW192-PB1	pHW192-PB1	pHW192-PB1
3 PHW193-PA	pHW193-PA	pHW193-PA	pHW193-PA
4 PHW195-NP	pHW195-NP	pHW195-NP	pHW195-NP
5 PMDV-A-M	pHW197-M	pHW197-M	pHW197-M
6 PHW198-NS	pMDV-A-NS	pHW198-NS	pHW198-NS
7 PHW194-HA	pHW194-HA	pMDV-A-HA	pHW194-HA
8 PHW-196-NA	pHW-196-NA	pHW-196-NA	pMDV-A-NA
CPE (+)	(+)	(+)	(+)

[0201] To further determine the packaging constraints of influenza A virus, the NS segment was separated into two separate gene segments: one encoding the NS1 genomic segment and the other encoding the NS2 genomic segment. The nine plasmids incorporating the genomic segments of influenza A were transfected into MDCK/COS cells as described above, and the recovered viruses were amplified in embryonated chicken eggs prior to titration on MDCK cells. Reduced plaque size was observed for the nine-plasmid system as compared to the eight-plasmid system described above. RT-PCR analysis demonstrated that only the S2 segment was present in the virions, and that the NS1 gene segment was not packaged.

[0202] Recovery of MDV-A and 6:2 Reassortant Viruses

[0203] Following the procedures described above, three days post transfection with either the 8 MDV-A plasmids (recombinant), or with plasmids incorporating the 6 MDV-A internal genes, and HA and NA derived from A/PR/8/34 (6:2 reassortant), transfected culture supernatants were used to infect fresh MDCK cells, and the infected cells were incubated at 33° C. for three days in the presence of 1 µg/ml TPCK-trypsin. The cytoplasmic effect of the recombinant virus on infected MDCK cells was observed using a microscope. Expression of viral hemagglutinin was monitored using a standard hemagglutination assay (HA). HA assays were performed by mixing 50 µl of serially 2-fold diluted culture supernatants with 50 µl of 1% chick red blood cells in 96-well plates. A HA titer of approximately 1:254-1:1024 was detected for the amplified viruses derived from either the transfected 8 MDV-A plasmids, or the 6:2 reassortant virus. The transfection reaction using the 8 A/PR/8/34 plasmid obtained from Dr. E. Hoffman was used as a positive control. Infectious influenza viruses were produced from these three transfection reactions as indicated in Table 5.

TABLE 5

Plasmids used for recovery of A/PR/8/34, MDV-A and 6:2 reassortant

Virus gene segment	A/PR/8/34 (H1N1)	rMDV-A(H2N2)	6:2 reassortant
1	pHW191-PB2 (AD731)	pMDV-A-PB2#2 (AD760)	pMDV-A-PB2#2 (AD760)
2	pHW192-PB1(AD732)	pMDV-A-PB1 (AD754)	pMDV-A-PB1 (AD754)

TABLE 5-continued

Plasmids used for recovery of A/PR/8/34, MDV-A and 6:2 reassortant			
Virus gene segment	A/PR/8/34 (H1N1)	rMDV-A(H2N2)	6:2 reassortant
3	pHW193-PA (AD733)	pMDV-A-PA (AD755)	pMDV-A-PA (AD755)
4	pHW195-NP (AD735)	pMDV-A-NP#1 (AD757)	pMDV-A-NP#1 (AD757)
5	pHW197-M (AD737)	pMDV-A-M (AD752)	pMDV-A-M (AD752)
6	pHW198-NS (AD738)	pMDV-A-NS (AD750)	pMDV-A-NS (AD750)
7	pHW194-HA (AD734)	pMDV-A-HA (AD756)	pHW194-HA (AD734)
8	pHW196-NA(AD735)	pMDV-A-NA#4 (AD759)	pHW196-NA (AD736)
CPE	+	+	+

[0204] RT-PCR was performed to map the genotypes of the recovered viruses. Viral RNA was isolated from the infected cell culture supernatant using the RNeasy mini Kit (Qiagen) and the eight influenza virus segments were amplified by RT-PCR using primers specific to each MDV-A gene segment and H1- and N1-specific primers. As shown in FIG. 3, rMDV-A contained PB2, PB1, NP, PA, M and NS that were specific to MDV-A and HA and NA specific to the H2 and N2 subtype. The 6:2 reassortant contained the 6 internal genes derived from MDV-A, and the HA and NA derived from A/PR/8/34 (H1N1). This confirmed that viruses generated from the transfected plasmids had the correct genotypes.

[0205] The rescued viruses were titrated by plaque assay on MDCK cells and the plaques were confirmed to be influenza virus by immunostaining using chicken serum raised against MDV-A. MDCK cells at 100% confluence on 12-well plates were infected with 100 μ l of 10-fold serially diluted virus at RT for 1 hour with gentle rocking. The inoculum was removed and the cells were overlaid with 1x L15 containing 0.8% agarose and 1 μ g/ml TPCK-trypsin. The plates were incubate at 35°C. or 33°C. for three days, fixed with 100% methanol, blocked by 5% milk in PBS, and incubated with 1:2000 diluted chicken anti-MDV-A antiserum for 1 hour followed by incubation with HRP-conjugated rabbit anti-chicken IgG for 1 hr. The plaques were visualized by addition of the HRP substrate solution (DAKO). All the recovered viruses exhibited positive immunostaining.

Example 4

Mapping the Genetic Basis of CA, TS, ATT Phenotypes of MDV-A

[0206] The MDV-A influenza virus vaccine strain has several phenotypes relevant to the production of vaccines, e.g., live attenuated vaccines: cold adaptation (ca), temperature sensitivity (ts) and attenuation (att). Sequence comparison of the MDV-A strain with the non-ts virulent wt A/AA/6/60 strain revealed that a minimal of 17 nt differences between these two strains (Table 6). Several of the changes in the MDV-A sequence are unique to this strain as compared to all the available influenza type A viruses in the GeneBank database, suggesting that one or more of these amino acid substitutions is functionally related to the att, ca and ts phenotype(s). The single amino acid change at PB2⁸²¹ was the only nucleotide position that had been previously reported as a determinant in the ts phenotype of MDV-A (Subbarao et al. (1995) Addition of Temperature-Sensitive Missense Mutations into the PB2 Gene of Influenza A

Transfected Viruses Can Effect an Increase in Temperature Sensitivity and Attenuation and Permits the Rational Design of a Genetically Engineered Live Influenza A Virus Vaccine *J. Virol.* 69:5969-5977).

[0207] In order to pinpoint the minimal substitutions involved in the MDV-A phenotypes, the nucleotides in the MDV-A clone that differ from wt A/AA/6/60 were individually changed to those of wt A/AA/6/60 (i.e., "reverted"). Each reverted gene segment was then introduced into host cells in combination with complementary segments of MDV-A to recover the single gene reassortants. In addition, the reverted gene segment and the corresponding MDV-A segment can also be transfected in combination with segments derived from other wild type strains, e.g., strain A/PR/8/34, to assess the contribution of each gene segment to the virus phenotypes. Using the recombinant MDV-A plasmid system described above, site-directed mutagenesis was performed to further modify the six internal genes to produce a non-ts reassortant. A total of 15 nucleotides substitution mutations were introduced into the six MDV-A plasmids to represent the recombinant wild type A/AA/6/60 genome (rWt, Flu064) as listed in Table 6. Madin-Darby canine kidney (MDCK) cells and COS-7 cells were maintained and transfected as described above. The recovered virus was then passaged in MDCK cells once, followed by amplification in the allantoic cavities of embryonic chicken eggs. Transfection and virus growth in MDCK and eggs were performed at 33°C., a temperature permissive for both ca and wt viruses to minimize any temperature selection pressures. Virus genotype was confirmed by sequence analysis of cDNA fragments amplified from viral RNA.

TABLE 6

Sequence Comparisons of "wt" A/AA/6/60 and MDV-A				
RNA Segment	Base (amino acid) Position	E10SE2	MDV-A	rWT (Flu044)
PB2	141	A	G	A
	821 (265)	A (Asn)	G(Ser)	A
	1182	A	T	T
	1212	C	T	T
	1933	T	C	T
	123	A	G	G
PB1	1195 (391)	A (Lys)	G (Glu)	A
	1395 (457)	G (Glu)	T (Asp)	G
	1766 (581)	A (Glu)	G (Gly)	A
	2005 (661)	G (Ala)	A (Thr)	A
	2019	C	T	C
PA	20	T	C	T
	1861 (613)	A (Lys)	G (Glu)	G
	2167/8 (715)	TT (Leu)	CC (Pro)	TT

TABLE 6-continued

Sequence Comparisons of "wt" A/AA/6/60 and MDV-A				
RNA Segment	Base (amino acid)	E10SE2	MDV-A	rWT (Flu044)
NP	146 (34) A (Asp)	A (Asp)	G (Gly)	G
	1550 '5A'	'5A'	'6A'	'6A'
M	969 (M2-86) G (Ala)	G (Ala)	T (Ser)	G
NS	483 (NS1-153) G (Ala)	G (Ala)	A (Thr)	G

Numbers in bold represent the differences between rMDV-A and rWT.

Words in bold (15) are the changes between rmdv-a and rwt.

[0208] Phenotypic characteristics were determined by procedures known in the art, e.g., as previously described in U.S. Pat. No. 6,322,967 to Parkin entitled "Recombinant tryptophan mutants of influenza," which is incorporated herein in its entirety. Briefly, temperature sensitivity of the recombinant viruses was determined by plaque assay on MDCK cells at 33, 38 and 39° C. MDCK cells in 6-well plates were infected with 400 μ l of 10-fold serially diluted virus and adsorbed at room temperature for 60 min. The inoculants were removed and replaced with 1 \times L15/MEM containing 1% agarose and 1 μ g/ml TPCK-trypsin. The infected cells were incubated at 33° C. in a CO₂ incubator or in water-tight containers containing 5% CO₂ submerged in circulating water baths maintained at 38 \pm 0.1° C. or 39 \pm 0.1° C. (Parkin et al. (1996) Temperature sensitive mutants of influenza A virus generated by reverse genetics and clustered charged to alanine mutagenesis. *Vir. Res.* 46:31-44). After three days' incubation, the monolayers were immunostained using chicken anti-MDV polyclonal antibodies and the plaques were enumerated. Plaque counts obtained at each of the temperatures were compared to assess the ts phenotype of each virus and each assay was performed a minimum of three times. The shut-off temperature was defined as the lowest temperature that had a titer reduction of 100-fold or greater compared to 33° C.

[0209] Infectious virus obtained from the cocultured COS-7/MDCK cells transfected with the eight plasmids (pMDV-PB2, pMDV-PB1, pMDV-PA, pMDV-NP, pMDV-HA, pMDV-NA, pMDV-M, and pMDV-NS) was amplified in chicken embryonated eggs, and was shown to exhibit the characteristic ts phenotype of nonrecombinant, biological derived MDV-A (Table 7). Neither MDV-A nor rMDV-A formed distinct plaques at 39° C., although both formed easily visualized plaques at 33° C.

TABLE 7

Replication of MDV/Wt reassortants at various temperatures					
Virus with Wt genes	33° C.	38° C.	33° C./38° C.	39° C.	33° C./39° C.
MDV	8.91	6.10	2.82	<4.0 [†]	>4.91
rMDV-A	8.72	6.19	2.53	<4.0	>4.72
Wt (E10SE2)	8.86	8.87	-0.01	8.87	-0.01
rWT (Flu064)	9.02	9.07	-0.05	8.96	0.06
Wt-PB2	8.46	7.87	0.59	5.80*	2.66
Wt-PB1	8.92	8.74	0.18	7.86*	1.06
Wt-NP	8.40	7.24	1.15	<4.0	>4.40
Wt-PA	8.57	6.10	2.48	<4.0	>4.57
Wt-M	8.80	6.68	2.12	<4.0	>4.80
Wt-NS	8.72	6.10	2.62	<4.0	>4.72

TABLE 7-continued

Replication of MDV/Wt reassortants at various temperatures					
Virus with Wt genes	33° C.	38° C.	33° C./38° C.	39° C.	33° C./39° C.
Wt-PB1/PB2	8.94	8.89	0.05	8.10*	0.85
Wt-PB1/PB2/NP	8.52	8.38	0.14	8.41	0.1

*Indicates reduction in plaque size compared to rWT.

[†]The underlined indicates that no plaques were detected at 10⁻⁴-fold dilution

[0210] In order to perform a systematic, detailed analysis of the genetic basis of the ts phenotype of MDV-A, the sequences of several closely related non-ts, non-att wt A/AA/6/60 strains with 17-48 nt differences from the ca A/AA/6/60, including the highly related isolate, wt A/AA/6/60 E10SE2, were utilized for comparison. A total of 19 nt differences exist between E10SE2 and MDV-A (Table 6). E10SE2 was shown to be non-ts (Table 7) and non-att in ferrets. In order to generate a recombinant non-ts virus, the MDV-A plasmids were altered by site directed mutagenesis to incorporate 15 of the 19 differences representing 10 amino acids changes. Four of the nucleotide positions, PB2-1182, 1212, PB1-123, and NP-1550, that differed between MDV-A and E10SE2 were not altered from the MDV-A sequence, since these nucleotides were observed in other non-ts isolates of A/AA/6/60 and, therefore, not expected to have a role in expression of the ts phenotype (Herlocher et al. (1996) Sequence comparisons of A/AA/6/60 influenza viruses: mutations which may contribute to attenuation. *Virus Research* 42:11-25). Recombinant virus (rWT, Flu064), encoding the 15 nucleotide changes, was obtained from the cocultured COS-7/MDCK cells transfected with a set of 8 plasmids, pWt-PB2, pWt-PB1, pWt-PA, pWt-NP, pWt-M, pWt-NS, pMDV-HA, and pMDV-NA. Sequencing analysis indicated that rWT contained the designed genetic changes and was non-ts at 39° C., identical to the biologically derived wt A/AA/6/60. These observations demonstrated that the ts phenotype mapped to a subset of these 15 nt changes.

[0211] Contribution of the Six Internal Gene Segments to Virus ts Phenotype

[0212] The effect of each wt gene segment on the MDV-A ts phenotype was assessed by creating recombinant, single-gene reassortants (Table 7). Introduction of wt PB2 into rMDV-A resulted in a virus that was only non-ts at 38° C.; however, it remained ts at 39° C. The reduction in virus titer at 38° C. and 39° C. (relative to 33° C.) was 0.6 log₁₀ and 2.7 log₁₀, respectively, as measured by plaque assay in MDCK cells. The reassortant containing the wt PB1 gene segment was non-ts, with respect to its ability to form plaques at both 38 and 39° C. The plaque size of this recombinant, however, was influenced by increased temperature and was significantly reduced at 39° C. as compared to rWT. Introduction of the wt NP gene segment into rMDV-A resulted in a virus that was also non-ts at 38° C., but in contrast to the wt PB2 recombinant, the virus containing the wt NP gene segment did not form plaques at 39° C. Introduction of wt PA, M or NS gene segments independently into rMDV-A did not alter the ts phenotype, indicating that these three gene segments had minimal role in maintenance of this phenotype.

[0213] Because neither wt PB1, wt PB2 or wt NP expressed individually on the MDV-A background could create a plaque efficiency and plaques size profile identical to non-ts rWT, these gene segments were introduced into MDV-A in various combinations. The combination of wt PB1 and wt PB2 resulted in a virus that was non-ts at both 38 and 39° C. (Table 7). Although the plaque size was larger than that of either single gene reassortant, it was significantly smaller than rWT. The triple combination of wt PB1/PB2/NP in rMDV-A resulted in a virus that was similar or identical to rWT in its plaquing efficiency and plaque size at 39° C. Therefore, whereas the wt PB2, PB1 and NP gene segments only partially reverted the ts phenotype when introduced individually, the combination of all three wt gene segments was able to fully revert the ts phenotype to a non-ts behavior identical to rWT.

[0214] In order to determine whether these 3 gene segments were capable of imparting the characteristic MDV-A ts phenotype to rWT, the six internal gene segments derived

only nt 821 resulted in an amino acid change (N265S, Table 6) and presumably represented the ts locus located in the PB2 gene segment. The PB1 gene of MDV-A differed from wt PB1 at 6 nt positions, of which 4 were coding changes (Table 6). Each of the wt amino acid residue substitutions was substituted individually into the PB1 gene segment of rMDV-A to assess their role in the ts phenotype. 1395G (Glu-457) and 2005G (Ala) did not affect the MDV-A ts phenotype. 1195A (Lys-391) and 1766A (Glu-581) each resulted in a slight reduction in the ts phenotype at 38° C., but had no effect at 39° C. (Table 8). These data indicated that 1195A and 1766A were the likely ts loci in the PB1 gene segment. However, combination of both 1195A and 1766A did not produce a ts phenotype similar to wt PB1 (Table 6). Addition of 2005G but not 1395A to PB1-1195A/1766A further decreased the virus ts phenotype at 39° C., demonstrating that 2005A also had a role in the expression of the ts phenotype specified by the PB1 segment of MDV-A.

TABLE 8

Virus with Wt sequence	Mapping the residues in PB1 that determine ts phenotype				
	33° C.	38° C.	33° C./38° C. \log_{10} PFU/mL	39° C.	33° C./39° C.
rMDV-A	8.67	6.00	2.67	<u>≤4.0</u> [†]	>4.67
rWT	9.04	9.01	0.03	9.03	0.01
PB1-1195A	8.06	6.68	1.38	<u>≤4.0</u>	>4.06
PB1-1395G	8.72	5.88	2.85	<u>≤4.0</u>	>4.72
PB1-1766A	8.07	6.70	1.37	<u>≤4.0</u>	>4.07
PB1-2005G	8.76	6.31	2.45	<u>≤4.0</u>	>4.76
PB1-1195A1766A	8.65	7.60	1.05	5.98*	2.68
PB1-1195A1395G1766A	8.84	8.13	0.71	6.38*	2.46
PB1-1195A1766A2005G	8.79	8.12	0.66	7.14*	1.64
PB1/PB2/NP	8.26	8.63	0.12	8.59	0.16
PB2/NP	8.81	8.21	0.59	7.56*	1.25
PB1-1195A/PB2/NP	8.86	8.81	0.05	7.60*	1.26
PB1-1766A/PB2/NP	9.33	8.84	0.50	8.71*	0.62
PB1-1766A2005G/PB2/NP	8.30	8.22	0.08	8.11*	0.18
PB1-1766A1395G/PB2/NP	8.88	8.85	0.03	8.39*	0.49
PB1-1195A1766A/PB2/NP	8.45	8.48	0.06	8.10	0.35

*Indicates reduction in plaque size compared to rWT.

[†]The underlined indicates that no plaques were detected at 10⁻⁴-fold dilution.

from MDV-A were introduced into rWT individually or in combination. Introduction of single PB1, PB2, or NP gene segment into rWT resulted in a reduction of virus titer at 38° C. and a greater reduction at 39° C., however, none of these single gene reassortants was as restricted at high temperature as rMDV-A (**FIG. 10**). The PA, M and NS gene segments derived from MDV-A did not influence the non-ts phenotype of rWT. Consistent with the previous reassortants, it was demonstrated that introduction of both MDV-A PB1 and PB2 genes into rWT backbone greatly increased virus ts phenotype at 38° C.; however, complete reversion of virus ts phenotype required addition of the NP gene. Thus, the PB1, PB2 and NP gene segments derived from MDV-A were important in conferring the complete ts phenotype.

[0215] Mapping the Genetic Loci that Determined MDV-A ts Phenotype.

[0216] The specific differences between the PB1, PB2 and NP gene segments of rWT and rMDV-A were addressed systematically to identify those changes that played a significant role in the ts phenotype. The NP gene of rMDV-A differed from rWT NP only at nt 146 (G34D, Table 6). The PB2 gene of rMDV-A differed from rWT at three sites, but

[0217] PB1 single site mutations were then introduced together with wt PB2 and wt NP into rMDV-A. Wt PB2/NP and rMDV-A reassortant was non-ts at 38° C. and had a titer reduction of 1.25 \log_{10} at 39° C. but its plaque size was much reduced compared to rWT. Addition of either PB1-1195A or 1766A did not significantly change the phenotype of wt PB2/NP reassortant. Only the combination of PB1-1195A and 1766A, together with a wt PB2 and wt NP, resulted in a virus that had the same non-ts phenotype as wt PB1/PB2/NP and rMDV-A reassortant (Table 8). Addition of PB1-1395G or 2005G to wt PB1-1766/PB2/NP did not convert the virus to a characteristic rWT non-ts phenotype. These data, therefore, demonstrated that the four amino acids distributed in the three PB1, PB2 and NP genes could completely revert the MDV-A ts phenotype.

[0218] Host Cell Restriction of MDV-A and Reassortant Viruses

[0219] In addition to the temperature sensitivity and attenuation phenotypes exhibited by the MDV-A virus and reassortant viruses with one or more MDV-A derived segment as described above, the MDV-A virus exhibited host cell restriction as indicated by reduced growth in Per.C6

cells relative to growth in MDCK cells. MDV-A and reassortant viruses with MDV-A derived PB1 and PB2 segments exhibited significantly reduced growth in Per.C6 cells relative to their growth in MDCK cells, as shown in **FIGS. 20A** and **B**.

[0220] Engineering of a Temperature Sensitive, Attenuated Virus Strain

[0221] To determine whether the five amino acids identified in the PB1, PB2 and NP gene segments of MDV-A would reproduce the ts and att phenotypes of MDV-A, PB1-391E, 581G, 661T, PB2-265S, NP-34G were introduced into a divergent wild type virus strain (A/PR/8/34; "PR8"), and the resulting virus exhibited 1.9 \log_{10} reduction in virus titer at 38° C. and 4.6 \log_{10} reduction at 39° C., which was very similar to that of rMDV-A (**FIG. 11**).

[0222] Sequence comparison between the PB1, PB2 and NP genes of ca A/AA/6/60 (MDV-A) and A/PR/8/34 revealed that the four substituted amino acids identified in the PB1 and PB2 genes of MDV-A are unique. NP³⁴ is conserved between MDV-A and PR8. Therefore, the three ts sites, PB1³⁹¹ (K391 E), PB1⁵⁸¹ (E581G) and PB1⁶⁶¹ (A661T), identified in the PB1 gene of MDV-A were introduced into PB1 of A/PR/8/34 and the PB2²⁶⁵ (N265S) was introduced into PB2 of A/PR/8/34 by site-directed mutagenesis. The mutations introduced into the PB1 and PB2 genes were verified by sequencing analysis. The primer pairs used for mutagenesis reaction are listed as in Table 9. These viruses are shown schematically in **FIG. 16**.

excess of DNA primer (5'-ATGTTCTTACGATGCGATTGGG, SEQ ID NO:89) labeled at its 5' end with [γ -³²P]-ATP and T4 polynucleotide kinase in 6 μ l of water. Following denaturing at 95° C. for 3 min, primer extension was performed after addition of 50 U of superscript reverse transcriptase (Invitrogen) in the reaction buffer provided with the enzyme containing 0.5 mM dNTP for 1 hr at 42° C. Transcription products were analyzed on 6% polyacrylamide gels containing 8M urea in TBE buffer and were detected by autoradiograph.

[0225] As shown in **FIGS. 12A** and **B**, the PB1 gene carrying three amino acid substitutions (PR8-3s), PB1³⁹¹ (K391E), PB1⁵⁸¹ (E581G) and PB1⁶⁶¹ (A661T), had reduced activity at 33° C. compared to PR8 control. A greater reduction in CAT protein expression (**FIG. 12A**) was observed for this mutant at 39° C., indicating PB1 gene with the three introduced MDV-A ts sites exhibited temperature sensitive replication in this in vitro assay. Introduction of PB2²⁶⁵ (N265S) into PR8 had very little effect on its activity at both permissive (33° C.) and nonpermissive temperatures (39° C.). Combination of both PB1-3s and PB2-1s resulted in greater reduction in protein activity (PR8-4s), which appeared to be even more ts than MDV-A. As expected, a low level activity (15%) was detected in cells transfected with PB1, PB2, PA, NP genes derived from MDV-A at 39° C. compared to wt A/AA/6/60 (wt A/AA).

[0226] PR8 mutant viruses were generated and recovered as described above. In brief, co-cultured cos7 and MDCK cells were transfected with eight plasmids encoding PR8 HA, NA, PB1, PB2, PA, NP, M and NS genes derived from

TABLE 9

Primers used for introducing ts mutations into PR8 PB1 and PB2 genes		
HJ240 PR8-PB1A1195G	5'	GAAAGAAGATTGAAGAAATCCGACCGCTC (SEQ ID NO:79)
HJ241 PR8-PB1A1195G.as	5'	GAGCGGTCGGATTCTTCAATCTCTTTC (SEQ ID NO:80)
HJ242 PR8-PB1A1766G	5'	GAAATAAGAAACTGTGGGGCAAACCCGTTCC (SEQ ID NO:81)
HJ243 PRS-PB1A1766G.as	5'	GGAACGGGTTGCCCAACAGTTCTTATTTC (SEQ ID NO:82)
HJ244 PR8-PB1G2005A	5'	GTATGATGCTGTTACAACAAACACTCC (SEQ ID NO:83)
HJ245 PR8-PB1G2005A.as	5'	GGAGTGTGTTGTTGAAACAGCATCATAC (SEQ ID NO:84)
HJ246 PR8-PB2A821G	5'	ATTGCTGCTAGGAGCATAGTGAGAAGAGC (SEQ ID NO:85)
HJ247 PR8-PB2A821G.as	5'	GCTCTCTCACTATGCTCCTAGCAGCAAT (SEQ ID NO:86)

[0223] To examine if the ts mutations introduced into PB1 and PB2 genes of PR8 confer the ts phenotype in vitro, a minigenome assay was performed. The influenza minigenome reporter, designated pFlu-CAT, contained the negative sense CAT gene cloned under the control of the pol I promoter. Expression of the CAT protein depended on the expression of influenza PB1, PB2, PA, and NP proteins.

[0224] Briefly, HEp-2 cells were transfected with 1 μ g of each of PB1, PB2, PA, NP and pFlu-CAT minigenome by lipofectamine 2000 (Invitrogen). After overnight (approximately 18 hour) incubation at 33° C. or 39° C., the cell extracts were analyzed for CAT protein expression by CAT ELISA kit (Roche Bioscience). The level of CAT mRNA was measured by primer extension assay. At 48 hr post-transfection, total cellular RNA was extracted by TRIzol reagent (Invitrogen) and $\frac{1}{3}$ of RNA was mixed with an

PR8. To make a virus carrying four ts loci (PR8-4s), PB1-3s containing three changes in PB1 at positions nt 1195 (K391E), nt 1766 (E581G) and nt 2005 (A661T) and PB1-1s containing one change in PB2 at position 821 (N265S) were used. In addition, PR8 virus carrying either three mutations in PB1 (PR8-3s) or one mutation in PB2 (PR8-1s) was also recovered separately. These viruses are shown schematically in **FIG. 16**. All four of the recombinant mutant PR8 viruses grew to very high titer in embryonic eggs, reaching a titer of 9.0 \log_{10} pfu/ml or greater as shown in Table 10.

[0227] To examine viral protein synthesis in infected cells, MDCK cells were infected with virus at an m.o.i of 5 and cells were labeled with ³⁵S-Trans at 7 hr post-infection for 1 hr. The labeled cell lysate was electrophoresed on 1.5% polyacrylamide gel containing SDS and autoradiographed. Protein synthesis was also studied by Western blotting. Virus

infected cells were harvested at 8 hr postinfection and electrophoresed on 4-15% gradient gel. The blot was probed with anti-M1 antibody or chicken anti-MDV-A polyclonal antibody, followed by incubation with HRP-conjugated secondary antibody. The antibody-conjugated protein bands were detected by the Chemiluminescent Detection System (Invitrogen) followed by exposure to X-ray film.

[0228] As shown in **FIG. 19**, all had a similar level of protein synthesis at 33° C., however, at 39° C. the level of protein synthesis was reduced slightly for PR8-1s but greatly reduced in PR8-3s and PR8-4s infected cells. Western blotting analysis also showed that reduced protein synthesis in the order of PR8-4s>PR8-3s>PR8-1s. Thus, the reduced replication of the ts mutants was likely the result of their reduced replication at the nonpermissive temperatures.

[0229] Temperature sensitivity of the PR8 mutant viruses was determined by plaque assay on MDCK cells at 33° C., 37° C., 38° C. and 39° C. The recovered viruses were amplified in embryonic eggs and introduced into cells as described above. After incubation of virus-infected cells for three days at the designated temperatures, cell monolayers were immunostained using chicken anti-MDV polyclonal antibodies and the plaques were enumerated. Plaque counts obtained at each of the temperatures were compared to assess the ts phenotype of each virus. The shut-off temperature was defined as the lowest temperature that had a titer reduction of 100-fold or greater compared to 33° C.

[0230] As shown in Table 10 and **FIG. 17**, all mutants replicated well at 33° C. although a slight reduction in virus titer was observed. At 38° C., a significant reduction in virus titer was observed for all the mutants. At 39° C., a reduction in virus titer greater than 4.0 \log_{10} was observed for viruses carrying the three ts loci in the PB1 gene (PR8-3s and PR8-4s). PR8-1s was also ts at 39° C. The ts phenotype of PR8-4s was very similar to that of MDV-A that had a reduction of 4.6 \log_{10} at 39° C. compared to 33° C. Although all the three PR8 mutants did not have greater than 2.0 \log_{10} reduction in virus titer at 37° C., their plaque morphology was different from those at 33° C. As shown in **FIG. 18**, the plaque size for each mutant was only slightly reduced at 33° C. compared to PR8. A significant reduction in plaque size at 37° C. was observed for PR8-3s and greater for PR8-4s. PR8-1s did not have significant reduction in plaque size at 37° C. At 39° C., only a few pin-point sized plaques were observed for both PR8-3s and PR8-4s. The plaque size of approximately 30% of that wt PR8 was observed for PR8-1s.

TABLE 10

Virus	Temperature sensitivity of PR8 with the introduced ts loci			
	33° C.	37° C.	38° C.	39° C.
MDV-A	8.6	7.0	6.4	4*
Wt A/AA	8.7	8.7	8.9	8.3
PR8	9.6	9.5	9.5	9
PR8-1s	9.4	8.9	7.7	7.4
PR8-3s	9.2	8.8	7.8	5.2
PR8-4s	9.5	7.8	7.1	4.4

A titer of 4.0 was assigned when no virus was detected at 10,000 dilutions.

[0231] Attenuation of the mutant PR8 viruses was examined in ferrets. In brief, male ferrets 9-10 weeks old were used to assess virus replication in the respiratory tracts of an

animal host. Ferrets were housed individually and inoculated intranasally with 8.5 \log_{10} pfu of virus. Three days after infection, ferrets were sedated with ketamine-HCL, lungs and nasal turbinates (NT) were harvested. The lung tissue homogenates were serially diluted and titrated in 10-day-old embryonated chicken eggs. Virus titer (\log_{10} EID₅₀/ml) in lungs was calculated by the Karber methods. Virus replication in NT was determined by plaque assay and expressed as \log_{10} pfu/ml.

[0232] The levels of virus replication in lungs and nasal turbinates were measured by EID50 or plaque assays (Table 11). Three days after infection, PR8 replicated to a level of 5.9 \log_{10} EID50/gram lung tissues. However, PR8-1s exhibited a 3.0 \log_{10} reduction in replication of ferret lungs and very little replication was detected for PR8-3s. No replication was detected for PR8-4s that was studied in two virus groups infected with virus obtained independently. Virus detection limit in ferret lungs by EID50 assay is 1.5 \log_{10} and thus a titer of 1.5 \log_{10} EID50 was assigned for PR8-4s. As a control, MDV-A did not replicate in ferret lungs and wt A/AA/6/60 replicated to a titer of 4.4 \log_{10} . Virus replication in nasal turbinates (NT) was examined by plaque assay on MDCK cells. PR8 replicated to a titer of 6.6 \log_{10} pfu/g in the nose. Only slight reductions in virus titer were observed for PR8-1s and PR8-3s. A reduction of 2.2 \log_{10} was observed for PR8-4s (A), whereas a 4.3 \log_{10} reduction was observed for PR8-4s (B), which carried a change in the PB1 gene (E390G). The greatly reduced replication of PR8-4s (B) correlates well with its ts phenotype at 37° C. An infectious dose of 8.5 \log_{10} pfu was used here instead of 7.0 \log_{10} pfu that was usually used for evaluating the attenuation phenotype of MDV-A derived influenza vaccines. This result indicated that PR8 carrying the four ts loci derived from MDV-A was attenuated in replication in the lower respiratory tracts of ferrets.

TABLE 11

Virus	Ferrets	Dose (\log_{10} pfu)	Replication of PR8 mutants in ferrets	
			Virus titer in lungs (\log_{10} EID ₅₀ /g \pm SE)	Virus titer in nasal turbinates (\log_{10} pfu/g \pm SE)
PR8	4	8.5	5.9 \pm 0.3	6.6 \pm 0.1
PR8-1s	4	8.5	3.8 \pm 0.4	5.9 \pm 0.2
PR8-3s	4	8.5	1.7 \pm 0.1	5.8 \pm 0.3
PR8-4s (A)	4	8.5	1.5 \pm 0.0 ^a	4.6 \pm 0.2
PR8-4s (B) ^b	4	8.5	1.5 \pm 0.0	2.3 \pm 0.3
MDV-A	4	8.5	1.5 \pm 0.0	4.6 \pm 0.1
Wt A/AA	4	8.5	4.4 \pm 0.1	5.4 \pm 0.1

no virus was detected and a titer of 1.5 \log_{10} EID50/g was assigned. The virus contains an additional change in PB1-1193 (E390G)

[0233] In both the ts and att assays, the PR8 mutant virus exhibited both ts and att phenotypes that were very similar to that of MDV-A. These data indicate that introduction of the unique amino acid substitutions of the MDV-A into a divergent influenza virus strain results in a virus exhibiting the temperature sensitive and attenuated phenotypes desirable for producing, e.g., live attenuated, vaccines. Additionally, the ts, att, PR-8 virus grew to a high titer that suitable for use as a master donor virus for the production of live attenuated or inactivated influenza vaccines. These results indicate that the five MDV-A mutations: PB1-391E, PB1-

581G, PB1-661T, PB2-265S, and NP-34G can impart the ts and att phenotypes to any influenza A strains. Similarly, novel ts, att B strains suitable for vaccine production can be produced by introducing the mutations of the MDV-B strain into influenza B strain viruses. In addition to producing live attenuated virus vaccines, introduction of these mutations into donor strains will lead to the production of safer inactivated vaccines.

Example 5

Eight Plasmid System for Production of MDV-B

[0234] Viral RNA from a cold adapted variant of influenza B/Ann Arbor/1/66 (ca/Master Ann Arbor/1/66 P1 Aviron 10/2/97), an exemplary influenza B master donor strain (MDV-B) was extracted from 100 μ l of allantoic fluid from infected embryonated eggs using the RNeasy Kit (Qiagen, Valencia, Calif.), and the RNA was eluted into 40 μ l H₂O. RT-PCR of genomic segments was performed using the One Step RT-PCR kit (Qiagen, Valencia, Calif.) according to the protocol provided, using 1 μ l of extracted RNA for each reaction. The RT-reaction was performed 50 min at 50° C., followed by 15 min at 94° C. The PCR was performed for 25 cycles at 94° C. for 1 min, 54° C. for 1 min, and 72° C. for 3 min. The P-genes were amplified using segment specific primers with BsmBI-sites that resulted in the generation of two fragments (Table 12).

[0235] Cloning of Plasmids

[0236] PCR fragments were isolated, digested with BsmBI (or BsaI for NP) and inserted into pAD3000 (a derivative of pHW2000 which allows the transcription of negative sense vRNA and positive mRNA) at the BsmBI site as described above. Two to four each of the resultant plasmids were sequenced and compared to the consensus sequence of MDV-B based on sequencing the RT-PCR fragments directly. Plasmids which had nucleotide substitutions resulting in amino acid changes different from the consensus sequence were “repaired” either by cloning of plasmids or by utilizing the Quikchange kit (Stratagene, La Jolla, Calif.). The resultant B/Ann Arbor/1/66 plasmids were designated pAB121-PB1, pAB122-PB2, pAB123-PA, pAB124-HA, pAB125-NP, pAB126-NA, pAB127-M, and pAB 128-NS. Using this bi-directional transcription system all viral RNAs and proteins are produced intracellularly, resulting in the generation of infectious influenza B viruses (FIG. 4).

[0237] It is noteworthy that pAB121-PB1 and pAB124-HA had 2 and pAB128-NS had 1 silent nucleotide substitution compared to the consensus sequence (Table 13). These nucleotide changes do not result in amino acid alterations, and are not anticipated to affect viral growth and rescue. These silent substitutions have been retained to facilitate genotyping of the recombinant viruses.

TABLE 12

RT-PCR primers for amplification of the eight vRNAs of influenza ca B/Ann Arbor/1/66.

	Forward primer	Reverse primer
PB1	Bm-PB1b-1: (SEQ ID NO:53) [1A] TATTCGTCAGGGAGCAGAACGGAGCCTTAAGATG	Bm-PB1b-1200R: (SEQ ID NO:54) TATTCGTCAGCCGTTCTCTTCATTGAAGAATGG
PB1	Bm-PB1b-1220: (SEQ ID NO:55) [1B] TATTCGTCAGGGAGCAGAACGGAGCCTGGATGATGATG	Bm-PB1b-2369R: (SEQ ID NO:56) ATATCGTCGTTAGTAGAAACACGAGCCTT
PB2	Bm-PB2b-1: (SEQ ID NO:57) [2A] TATTCGTCAGGGAGCAGAACGGAGCCTTCAGATG	Bm-PB2b-1145R: (SEQ ID NO:58) TATTCGTCCTCTCATTTGCTCTTTAATATTCCCC
PB2	Bm-PB2b-1142: (SEQ ID NO:59) [2B] TATTCGTCATGAGAATGGAAAACACTAATAAATTCAAGC	Bm-PB2b-2396R: (SEQ ID NO:60) ATATCGTCGTTAGTAGAAACACGAGCATT
PA	Bm-Pab-1: (SEQ ID NO:61) [3A] TATTCGTCAGGGAGCAGAACGGAGCTTGA	Bm-Pab-1261R: (SEQ ID NO:62) TATTCGTCAGGGCCCTTTACTTGTCAAGAGTGC
PA	Bm-Pab-1283: (SEQ ID NO:63) [3B] TATTCGTCCTGGATCTACCAAGAAATAGGGCCAGAC	Bm-Pab-2308R: (SEQ ID NO:64) ATATCGTCGTTAGTAGAAACACGAGCATT
HA	MDV-B 5' BsmBI-HA: (SEQ ID NO:65) TATTCGTCAGGGAGCAGAACGGAGCATTTCTAAATTC	MDV-B 3' BsmBI-HA: (SEQ ID NO:66) ATATCGTCGTTAGTAGAAACAGCATT
NP	Ba-NPb-1: (SEQ ID NO:67) TATTCGTCAGGGAGCAGAACGGAGCATTTCTTGT	Ba-NPb-1842R: (SEQ ID NO:68) ATATGGTCGTTAGTAGAAACACAGCATT
NA	MDV-B 5' BsmBI-NA: (SEQ ID NO:69) TATTCGTCAGGGAGCAGAACGGAGCATCTCTCAAAC	MDV-B 3' BsmBI-NA: (SEQ ID NO:70) ATATCGTCGTTAGTAGAAACAGCATT
M	MDV-B 5' BsmBI-M: (SEQ ID NO:71) TATTCGTCAGGGAGCAGAACGGAGCATCTCTAAATG	MDV-B 3' BsmBI-M: (SEQ ID NO:72) ATATCGTCGTTAGTAGAAACACAGCATT
NS	MDV-B 5' BsmBI-NS: (SEQ ID NO:73) TATTCGTCAGGGAGCAGAACGGAGCATCTCTTGTAGTC	MDV-B 3' BsmBI-NS: (SEQ ID NO:74) ATATCGTCGTTAGTAGAAACAGGAGATT

The sequences complementary to the influenza sequences are shown in bold. The 5'-ends have recognition sequences for the restriction endonucleases BsmBI (Bm) or BsaI (Ba).

TABLE 13

Plasmid set representing the eight segments of B/Ann Arbor/1/66 (MDV-B)			
Seg.	plasmids	nucleotides	protein
PB1	PAB121-PB1	A924 > G924; C1701 > T1701	silent
PB2	PAB122-PB2	consensus	—
PA	PAB123-PA	consensus	—
HA	PAB124-HA	T150 > C150; T153 > C153	silent
NP	PAB125-NP	consensus	—
NA	PAB126-NA	consensus	—
M	PAB127-M	consensus	—
NS	PAB128-NS	A416 > G416	NS1: silent

[0238] For construction of the plasmids with nucleotide substitution in PA, NP, and M1 genes the plasmids pAB123-PA, pAB 125-NP, pAB127-M were used as templates. Nucleotides were changed by Quikchange kit (Stratagene, La Jolla, Calif.). Alternatively, two fragments were amplified by PCR using primers which contained the desired mutations, digested with BsmBI and inserted into pAD3000-BsmBI in a three fragment ligation reaction. The generated plasmids were sequenced to ensure that the cDNA did not contain unwanted mutations.

[0239] The sequence of template DNA was determined by using Rhodamine or dRhodamine dye-terminator cycle sequencing ready reaction kits with AmpliTaq® DNA polymerase FS (Perkin-Elmer Applied Biosystems, Inc, Foster City, Calif.). Samples were separated by electrophoresis and analyzed on PE/ABI model 373, model 373 Stretch, or model 377 DNA sequencers.

[0240] In a separate experiment, viral RNA from influenza B/Yamanshi/166/98 was amplified and cloned into pAD3000 as described above with respect to the MDV-B strain, with the exception that amplification was performed for 25 cycles at 94° C. for 30 seconds, 54° C. for 30 seconds and 72° C. for 3 minutes. Identical primers were used for amplification of the B/Yamanashi/166/98 strain segments, with the substitution of the following primers for amplification of the NP and NA segments: MDV-B 5'BsmBI-NP: TATTCGTCTCAGGGAGCAGAACAGACAG-CATTTCTGTG (SEQ ID NO:75) and MDV-B 3'BsmBI-NP: ATATCGTCCTCGTATTAGTAGAAACAA-CAGCATTTTTAC (SEQ ID NO:76) and Bm-NAb-1: TATTCGTCTCAGGGAGCAGAACAGAGCA (SEQ ID NO:77) and Bm-NAb-1557R:ATATCGTCTCGTATTAG-TAGTAACAAAGAGCATT (SEQ ID NO:78), respectively. The B/Yamanashi/166/98 plasmids were designated pAB251-PB1, pAB252-PB2, pAB253-PA, pAB254-HA, pAB255-NP, pAB256-NA, pAB257-M, and pAB258-NS. Three silent nucleotide differences were identified in PA facilitating genotyping of recombinant and reassortant B/Yamanashi/166/98 virus.

Example 6

Generation of Infectious Recombinant Influenza B and Reassorted Influenza Virus

[0241] To overcome the obstacles encountered in attempting to grow influenza B in a helper virus free cell culture system, the present invention provides novel vectors and protocols for the production of recombinant and reassortant

B strain influenza viruses. The vector system used for the rescue of influenza B virus is based on that developed for the generation of influenza A virus (Hoffmann et al. (2000) A DNA transfection system for generation of influenza A virus from eight plasmids *Proc Natl Acad Sci USA* 97:6108-6113; Hoffmann & Webster (2000) Unidirectional RNA polymerase I-polymerase II transcription system for the generation of influenza A virus from eight plasmids *J Gen Virol* 81:2843-7). 293T or COS-7 cells (primate cells with high transfection efficiency and polI activity) were co-cultured with MDCK cells (permissive for influenza virus), 293T cells were maintained in OptiMEM I-AB medium containing 5% FBS cells, COS-7 cells were maintained in DMEM I-AB medium containing 10% FBS. MDCK cells were maintained in 1× MEM, 10% FBS with the addition of antibiotic and antimycotic agents. Prior to transfection with the viral genome vectors, the cells were washed once with 5 ml PBS or medium without FBS. Ten ml trypsin-EDTA was added to confluent cells in a 75 cm² flask (MDCK cells were incubated for 20-45 min, 293T cells were incubated for 1 min). The cells were centrifuged, and resuspended in 10 ml OptiMEM I-AB. One ml of each suspended cell line was then diluted into 18 ml OptiMEM I-AB, and mixed. The cells were then aliquoted into a 6 well plate at 3 ml/well. After 6-24 hours, 1 μ g of each plasmid was mixed in an 1.5 ml Eppendorf tube with OptiMEM I-AB to the plasmids (x μ l plasmids+x μ l OptiMEM I-AB+x μ l TransIT-LT1=200 μ l); 2 μ l TransIT-LT1 per μ g of plasmid DNA. The mixture was incubated at room temperature for 45 min. Then 800 μ l of OptiMEM I-AB was added. The medium was removed from the cells, and the transfection mixture was added to the cells (t=0) at 33° C. for 6-15 hours. The transfection mixture was slowly removed from the cells, and 1 ml of OptiMEM I-AB was added, and the cells were incubated at 33° C. for 24 hours. Forty-eight hours following transfection, 1 ml of OptiMEM I-AB containing 1 μ g/ml TPCK-trypsin was added to the cells. At 96 hours post-transfection, 1 ml of OptiMEM I-AB containing 1 μ g/ml TPCK-trypsin was added to the cells.

[0242] Between 4 days and 7 days following transfection 1 ml of the cell culture supernatant was withdrawn and monitored by HA or plaque assay. Briefly, 1 ml of supernatant was aliquoted into an Eppendorf tube and centrifuge at 5000 rpm for 5 min. Nine hundred μ l of supernatant was transferred to a new tube, and serial dilutions were performed at 500 μ l/well to MDCK cells (e.g., in 12 well plates). The supernatant was incubated with the cells for 1 hour then removed, and replaced with infection medium (1×MEM) containing 1 μ g/ml of TPCK-trypsin. HA assay or plaque assays were then performed. For example, for the plaque assays supernatants were titrated on MDCK cells which were incubated with an 0.8% agarose overlay for three days at 33° C. For infection of eggs the supernatant of transfected cells were harvested six or seven days after transfection, 100 μ l of the virus dilutions in Opti-MEM I were injected into 11 days old embryonated chicken eggs at 33° C. The titer was determined three days after inoculation by TCID₅₀ assay in MDCK cells.

[0243] To generate MDV-B, either co-cultured 293T-MDCK or COS-7-MDCK cells were transfected with 1 μ g of each plasmid. When examined at 5 to 7 days post-transfection the co-cultured MDCK cells showed cytopathic effects (CPE), indicating the generation of infectious MDV-B virus from cloned cDNA. No CPE was observed in

cells transfected with seven plasmids (Table 14). To determine the efficiency of the DNA transfection system for virus generation, supernatants of cells were titrated seven days after transfection on MDCK cells and the virus titer was determined by plaque assay. The virus titer of the supernatant of co-cultured 293T-MDCK was 5.0×10^6 pfu/ml and 7.6×10^6 pfu/ml in COS7-MDCK cells.

TABLE 14

Generation of infectious Influenza-B virus from eight plasmids				
segment				
1	2	3	4	
PB1	pAB121-PB1	—	PAB121-PB1	—
PB2	pAB122-PB2	pAB122-PB2	PAB122-PB2	pAB122-PB2
PA	pAB123-PA	pAB123-PA	pAB123-PA	pAB123-PA
HA	pAB124-HA	pAB124-HA	pAB124-HA	pAB124-HA
NP	pAB125-NP	pAB125-NP	pAB125-NP	pAB125-NP
NA	pAB126-NA	pAB126-NA	pAB126-NA	pAB126-NA
M	pAB127-M	pAB127-M	pAB127-M	pAB127-M
NS	pAB128-NS	pAB128-NS	pAB128-NS	pAB128-NS
co-cultured 293T-MDCK cells co-cultured COS-7-MDCK cells				
CPE	+	—	+	—
pfu/ml	5.0×10^6	0	7.6×10^6	0

[0244] Transiently co-cultured 293T-MDCK (1, 2) or co-cultured COS7-MDCK cells (3, 4) were transfected with seven or eight plasmids. Cytopathic effect (CPE) was monitored seven days after transfection in the co-cultured MDCK cells. Seven days after transfection the supernatants of transfected cells were titrated on MDCK cells. The data of pfu/ml represent the average of multiple, (e.g., three or four) transfection experiments.

[0245] Comparable results were obtained in transfection experiments utilizing the B/Yamanashi/166/98 plasmid vectors. These results show that the transfection system allows the reproducible de novo generation of influenza B virus from eight plasmids.

[0246] Genotyping of Recombinant Influenza B

[0247] After a subsequent passage on MDCK cells, RT-PCR of the supernatant of infected cells was used to confirm the authenticity of the generated virus. RT-PCR was performed with segment specific primers for all eight segments (Table 12). As shown in FIG. 5A, PCR products were generated for all segments. Direct sequencing of the PCR products of the PB1, HA, and NS segments revealed that the four nucleotides analyzed were the same as found in the plasmid pAB121-PB1, pAB124-HA, and pAB128-NS. These results confirmed that the generated virus was generated from the designed plasmids and exclude (in addition to the negative controls) any possible laboratory contamination with the parent virus (FIG. 5B).

[0248] Similarly, following transfection with the B/Yamanashi/166/98 plasmid vectors, virus was recovered and the region encompassing nucleotides 1280-1290 of the PA segment were amplified. Sequencing confirmed that the recovered virus corresponded to the plasmid-derived recombinant B/Yamanashi/166/98 (FIGS. 5C and D).

[0249] Phenotyping of rMDV-B

[0250] The MDV-B virus shows two characteristic phenotypes: temperature sensitivity (ts) and cold adaptation (ca). By definition a 2 log (or higher) difference in virus titer at 37° C. compared to 33° C. defines ts, ca is defined by less than 2 log difference in virus growth at 25° C. compared to 33° C. Primary chicken kidney (PCK) cells were infected with the parent virus MDV-B and with the transfected virus derived from plasmids to determine the viral growth at three temperatures.

[0251] For plaque assay confluent MDCK cells (ECACC) in six well plates were used. Virus dilutions were incubated for 30-60 min. at 33° C. The cells were overlayed with an 0.8% agarose overlay. Infected cells were incubated at 33° C. or 37° C. Three days after infection the cells were stained with 0.1% crystal violet solution and the number of plaques determined.

[0252] The ca-ts phenotype assay was performed by TCID₅₀ titration of the virus samples at 25, 33, and 37° C. This assay format measures the TCID₅₀ titer by examining the cytopathic effect (CPE) of influenza virus on primary chick kidney cell monolayers in 96-well cell culture plates at different temperatures (25° C., 33° C., 37° C.). This assay is not dependent on the plaque morphology, which varies with temperature and virus strains; instead it is dependent solely on the ability of influenza virus to replicate and cause CPE. Primary chicken kidney (PCK) cell suspension, prepared by trypsinization of the primary tissue, were suspended in MEM (Earl's) medium containing 5% FCS. PCK cells were seeded in 96 well cell culture plates for 48 hours in order to prepare monolayer with >90% confluence. After 48 hrs, the PCK cell monolayer were washed for one hour with serum free MEM medium containing 5 mM L-Glutamine, antibiotics, non-essential amino acid, referred as Phenotype Assay Medium (PAM). Serial ten-fold dilution of the virus samples were prepared in 96 well blocks containing PAM. The diluted virus samples were then plated onto the washed PCK monolayer in the 96 well plates. At each dilution of the virus sample, replicates of six wells were used for infection with the diluted virus. Un-infected cells as cell control were included as replicate of 6 wells for each sample. Each virus sample was titrated in 2-4 replicates. Phenotype control virus with pre-determined titers at 25° C., 33° C., and 37° C. is included in each assay. In order to determine the ts phenotype of the virus samples, the plates were incubated for 6 days at 33° C. and 37° C. in 5% CO₂ cell culture incubators. For ca-phenotype characterization the plates were incubated at 25° C. for 10 days. The virus titer was calculated by the Karber Method and reported as Log₁₀ Mean (n=4) TCID₅₀ Titer/ml±Standard Deviation. The standard deviations of the virus titers presented in FIG. 1-3 ranged from 0.1 to 0.3. The difference in virus titer at 33° C. and 37° C. were used to determine the ts phenotype and difference in titer at 25° C. and 33° C. of the virus were used to determine the ca phenotype.

[0253] The plasmid derived recombinant MDV-B (rec-MDV-B) virus expressed the two characteristic phenotypes in cell culture, ca and ts, as expected. The ca phenotype, efficient replication at 25° C., is functionally measured as a differential in titer between 25° C. and 33° C. of less than or equal to 2 log10 when assayed on PCK cells. Both the parental MDV-B and recMDV-B expressed ca; the difference between 25° C. and 33° C. was 0.3 and 0.4 log10, respectively (Table 15). The ts phenotype is also measured by observing the titers at two different temperatures on PCK

cells; for this phenotype, however, the titer at 37° C. should be less than the titer at 33° C. by 2 log₁₀ or more. The difference between 33° C. and 37° C. for the parental MDV-B and recMDV-B was 3.4 and 3.7 log₁₀, respectively (Table 15). Thus, the recombinant plasmid-derived MDV-B virus expressed both the ca and ts phenotypes.

[0254] The recombinant virus had a titer of 7.0 log₁₀ TCID₅₀/ml at 33° C. and 3.3 TCID₅₀/ml at 37° C. and 8.8 log₁₀ TCID₅₀/ml at 25° C. (Table 15). Thus, the recombinant virus derived from transfection with the eight influenza MDV-B genome segment plasmids has both the ca and ts phenotype.

TABLE 15

Virus	Temperature (0 C.)			Phenotype
	25	33	37	
	Log10 TCID50/ml (Mean + SD)			
ca B/Ann Arbor/01/66 (MDV-B)	8.8 + 0.3	8.5 + 0.05	5.1 + 0.1	ca, ts
RecMDV-B	7.4 + 0.3	7.0 + 0.13	3.3 + 0.12	ca, ts
Rec53-MDV-B	5.9 + 0.1	5.7 + 0.0	5.3 + 0.1	ca, non-ts

Primary chicken kidney cells were infected with the parent virus MDV-B and the plasmid-derived recombinant virus (recMDV-B). The virus titer was determined at three different temperatures.

Example 7

Production of Reassortant B/Yamanashi/166/98 Virus

[0255] The HA and NA segments of several different strains representing the major lineages of influenza B were amplified and cloned into pAD3000, essentially as described

CA (SEQ ID NO:87); Bm-NAb-1557R: ATA TCG TCT CGT ATT AGT AGT AAC AAG AGC ATT TT (SEQ ID NO:88) was synthesized and used to simultaneously amplify the HA and NA genes from various influenza B strains (FIG. 8). The HA and NA PCR-fragments of B/Victoria/504/2000, B/Hawaii/10/2001, and B/Hong Kong/330/2001 were isolated, digested with BsmBI and inserted into pAD3000. These results demonstrated the applicability of these primers for the efficient generation of plasmids containing the influenza B HA and NA genes from several different wild type viruses representing the major lineages of influenza B. The RT-PCR products can be used for sequencing and/or cloning into the expression plasmids.

[0256] In order to demonstrate the utility of B/Yamanashi/166/98 (a B/Yamagata/16/88-like virus) to efficiently express antigens from various influenza B lineages, reassortants containing PB1, PB2, PA, NP, M, NS from B/Yamanashi/166/98 and the HA and NA from strains representing both the Victoria and Yamagata lineages (6+2 reassortants) were generated. Transiently cocultured COS7-MDCK cells were cotransfected with six plasmids representing B/Yamanashi/166/98 and two plasmids containing the cDNA of the HA and NA segments of two strains from the B/Victoria/2/87 lineage, B/Hong Kong/330/2001 and B/Hawaii/10/2001, and one strain from the B/Yamagata/16/88 lineage, B/Victoria/504/2000, according to the methods described above. Six to seven days after transfection the supernatants were titrated on fresh MDCK cells. All three 6+2 reassortant viruses had titers between 4-9×10⁶ pfu/ml (Table 16). These data demonstrated that the six internal genes of B/Yamanashi/166/98 could efficiently form infectious virus with HA and NA gene segments from both influenza B lineages.

[0257] Supernatants of cocultured COS7-MDCK cells were titrated six or seven days after transfection and the viral titer determined by plaque assays on MDCK cells.

TABLE 16

Plasmid set used for the generation of B/Yamanashi/166/98 and 6 + 2 reassortants.

segment						
1	—	pAB251-PB1	pAB251-PB1	pAB251-PB1	pAB251-PB1	pAB251-PB1
2	pAB252-PB2	pAB252-PB2	pAB252-PB2	pAB252-PB2	pAB252-PB2	pAB252-PB2
3	pAB253-PA	pAB253-PA	pAB253-PA	pAB253-PA	pAB253-PA	pAB253-PA
4	pAB254-HA	pAB254-HA	pAB281-HA	pAB285-HA	pAB287-HA	pAB287-HA
5	pAB255-NP	pAB255-NP	pAB255-NP	pAB255-NP	pAB255-NP	pAB255-NP
6	pAB256-NA	pAB256-NA	pAB291-NA	pAB295-NA	pAB297-NA	pAB297-NA
7	pAB257-M	pAB257-M	pAB257-M	pAB257-M	pAB257-M	pAB257-M
8	pAB258-NA	pAB258-NA	pAB258-NA	pAB258-NA	pAB258-NA	pAB258-NA
Recombinant virus	8	6 + 2	6 + 2	6 + 2	6 + 2	6 + 2
B/Yamanashi/166/98		B/Victoria/504/2000	B/Hawaii/10/2001	B/Hong Kong/330/2001		
pfu/ml ^a	0	4 × 10 ⁶	9 × 10 ⁶	6 × 10 ⁶	7 × 10 ⁶	

above. The primers were optimized for simultaneous RT-PCR amplification of the HA and NA segments. Comparison of the terminal regions of the vRNA representing the non coding region of segment 4 (HA) and segment 6 (NB/NA) revealed that the 20 terminal nucleotides at the 5' end and 15 nucleotides at the 3' end were identical between the HA and NA genes of influenza B viruses. A primer pair for RT-PCR (underlined sequences are influenza B virus specific) Bm-NAb-1: TAT TCG TCT CAG GGA GCA GAA GCA GAG

[0258] Relatively high titers are obtained by replication of wild type B/Yamanashi/166/98 in eggs. Experiments were performed to determine whether this property was an inherent phenotype of the six "internal" genes of this virus. To evaluate this property, the yield of wild type B/Victoria/504/2000, which replicated only moderately in eggs, was compared to the yield of the 6+2 reassortant expressing the B/Victoria/504/2000 HA and NA. These viruses in addition

to wild type and recombinant B/Yamanashi/166/98 were each inoculated into 3 or 4 embryonated chicken eggs, at either 100 or 1000 pfu. Three days following infection, the allantoic fluids were harvested from the eggs and the TCID₅₀ titers determined on MDCK cells. The 6+2 reassortants produced similar quantities of virus in the allantoic fluid to the wt and recombinant B/Yamanashi/166/98 strain (FIG. 9). The difference in titer between B/Victoria/504/2000 and the 6+2 recombinant was approximately 1.6 log₁₀ TCID₅₀ (0.7-2.5 log₁₀ TCID₅₀/mL, 95% CI). The difference between B/Victoria/504/2000 and the 6+2 recombinant were confirmed on three separate experiments (P<0.001). These results demonstrated that the egg growth properties of B/Yamanashi/166/98 could be conferred to HA and NA antigens that are normally expressed from strains that replicated poorly in eggs.

Example 8

Molecular Basis for Attenuation of CA B/Ann Arbor/1/66

[0259] The MDV-B virus (ca B/Ann Arbor/1/66) is attenuated in humans, shows an attenuated phenotype in ferrets and shows a cold adapted and temperature sensitive phenotype in cell culture. The deduced amino acid sequences of the internal genes of MDV-B were compared with sequences in the Los Alamos influenza database (on the world wide web at: flu.lanl.gov) using the BLAST search algorithm. Eight amino acids unique to MDV-B, and not present in any other strain were identified (Table 17). Genome segments encoding PB1, BM2, NS1, and NS2 show no unique substituted residues. The PA and M1 proteins each have two, and the NP protein has four unique substituted amino acids (Table 17). One substituted amino acid is found in PB2 at position 630 (an additional strain B/Harbin/7/94 (AF170572) also has an arginine residue at position 630).

[0260] These results suggested that the gene segments PB2, PA, NP and M1 may be involved in the attenuated phenotype of MDV-B. In a manner analogous to that described above for MDV-A, the eight plasmid system can be utilized to generate recombinant and reassortant (single and/or double, i.e., 7:1; 6:2 reassortants) in a helper independent manner simply by co-transfection of the relevant plasmids into cultured cells as described above with respect to MDV-A. For example, the 6 internal genes from B/Lee/40 can be used in conjunction with HA and NA segments derived from MDV-B to generate 6+2 reassortants.

TABLE 17

Unique substituted amino acids of B/Ann Arbor/1/66				Number of aligned sequences
ca B/Ann Arbor/1/66	Aligned sequences (wild type viruses)			
Nr. pos.	amino acid	codon	amino acid	codon
PB1 0	—			23
PB2 1	630 Arg630 AGA	Ser630	AGC	23
PA 2	431 Met431 ATG	Val431	GTG	23
	497 His497 CAT	Tyr497	TAT	
NP 4	55 Ala55 GCC	Thr55	ACC	26
	114 Ala114 GCG	Val114	GTG	

TABLE 17-continued

Unique substituted amino acids of B/Ann Arbor/1/66				Number of aligned sequences
ca B/Ann Arbor/1/66	Aligned sequences (wild type viruses)			
Nr. pos.	amino acid	codon	amino acid	codon
M1 2	410 His410 CAT	Pro410	CCT, CCC	
	509 Thr509 GAC	Ala509	GGC	
	159 Gln159 CAA	His159	CAT	24
	183 Val183 GTG	M183	ATG	
BM2 0	—	—	—	24
NS1 0	—	—	—	80
NS2 0	—	—	—	80

The deduced amino acid sequence of eight proteins of ca B/Ann Arbor was used in a BLAST search. Amino acid position which were different between MDV-B and the aligned sequences are shown. The nucleotides in the codons that are underlined represent the substituted positions.

[0261] In order to determine whether the 8 unique amino acid differences had any impact on the characteristic MDV-B phenotypes, a recombinant virus was constructed in which all eight nucleotide positions encoded the amino acid reflecting the wt influenza genetic complement. A set of plasmids was constructed in which the eight residues of the PA, NP, and M1 genes were changed by site directed mutagenesis to reflect the wild type amino acids (as indicated in Table 17). A recombinant with all eight changes, designated rec53-MDV-B, was generated by cotransfection of the constructed plasmids onto cocultured COS7-MDCK cells. The coculturing of MDCK cells and growth at 33° C. ensured that the supernatant contained high virus titers six to seven days after transfection. The supernatants of the transfected cells were titrated and the titer determined on MDCK cells by plaque assay and PCK cells at 33° C. and 37° C.

[0262] As shown in FIG. 13, in two different independent experiments, recMDV-B expressed the ts-phenotype in both MDCK cells and PCK cells. The triple reassortant virus rec53-MDV-B designed harboring all eight amino acid changes expressed the non-ts-phenotype, the difference in titer between 33° C. and 37° C. was only 0.7 log₁₀ in PCK cells. This titer was less than the required 2 log₁₀ difference characteristic of the ts definition and significantly lower than the ~3 log₁₀ difference observed with recMDV-B. These results show that the alteration of the eight amino acids within PA, NP, and M1 proteins was sufficient to generate a non-ts, wild type-like virus with both homologous and heterologous glycoproteins.

[0263] The contribution of each gene segment to the ts phenotype was then determined. Plasmid derived recombinants harboring either the PA, NP, or M gene segment with the wild-type amino acid complement were generated by the DNA cotransfection technique. All single gene recombinants exhibited growth restriction at 37° C. in MDCK cells and in PCK cells (FIG. 14), indicating that changes in no one gene segment were capable of reverting the ts phenotype. In addition, recombinant viruses that carried both the NP and M or PA and M gene segments together also retained the ts-phenotype. In contrast, recombinant viruses that harbored

both the PA and NP gene segments had a difference in titer between 37° C. and 33° C. of 2.0 log₁₀ or less, similar to the rec53-MDV-B. These results show that the NP and PA genes have a major contribution to the ts-phenotype.

[0264] To determine whether all of the four amino acids in the NP protein and two in the PA protein contribute to non-ts, triple gene and double-gene recombinants with altered NP and PA genes were generated (FIG. 15). The substitution of two amino acids in the NP protein, A114→V114 and H410→P410 resulted in non-ts phenotype. Viruses with single substitution H410→P410 in the nucleoprotein showed non-ts phenotype in MDCK and PCK. On the other hand, the single substitution A55→T55 showed a ts-phenotype, as did the single substitution at position 509. These results indicate that amino acid residues V114 and P410 in NP are involved in efficient growth at 37° C. (FIG. 21A). A similar strategy was employed to dissect the contribution of the two amino acids in the PA gene. A set of recombinants was constructed, each harboring an NP gene segment with four wild-type consensus amino acids and a PA gene with only one of the two consensus wild type amino acids. Substitution of H497→Y497 remained ts (FIG. 21B), demonstrating that this locus had little impact on expression of the phenotype. In contrast, substitution of M431 with V431 resulted in reversion of the ts phenotype. These results show that amino acids A114 and H410 in NP and M431 in PA are the major determinants for temperature sensitivity of MDV-B.

[0265] Based on prior evidence, a ts-phenotype and an attenuated phenotype are highly correlated. It is well established that ca B/Ann Arbor/1/66 virus is not detectable in lung tissue of infected ferrets, whereas non attenuated influenza B viruses viruses are detectable in lungs after intranasal infection. To determine whether identical mutation underlie the ts and att phenotypes, the following studies were performed.

[0266] Recombinant viruses obtained after transfection were passaged in embryonated chicken eggs to produce a virus stock. Nine week old ferrets were inoculated intranasally with 0.5 ml per nostril of viruses with titers of 5.5, 6.0 or 7.0 log₁₀ pfu/ml. Three days after infection ferrets were sacrificed and their lungs and turbinates were examined as described previously.

[0267] Ferrets (four animals in each group) were infected intranasally with recMDV-B or rec53-MDV-B. Three days after infection virus nasal turbinates and lung tissue were harvested and the existence of virus was tested. No virus was detected in lung tissues of ferrets infected with 7.0 log₁₀ pfu recMDV-B. From the four animals infected with rec53-MDV-B virus with 7.0 log₁₀ pfu in three animals virus was

detected in lung tissue (one animal in this group for unknown reasons). In two out of four lung tissues of ferrets infected with rec53-MDV-B at a lower dose (5.5 log pfu/ml) virus could be isolated from lung tissue. Thus, the change of the eight unique amino acids in PA, NP, and M1 protein into wild type residues were sufficient to convert a att phenotype into a non-ts phenotype.

[0268] Since the data in cell culture showed that PA and NP are main contributors to the ts-phenotype, in a second experiment, ferrets were infected with rec53-MDV-B (PA, NP, M), rec62-MDV-B (PA), NP rec71-MDV-B (NP) with 6 log pfu. Two out of four animals infected with rec53-MDV-B had virus in the lung. None of the lung tissues of ferrets infected with single and double reassortant viruses had detectable levels of virus. Thus, in addition to the amino acids in the PA and NP proteins, the M1 protein is important for the att phenotype. Virus with wt PA and NP did not replicate in ferret lung, indicating that a subset of the mutations involved in attenuation are involved in the ts phenotype.

[0269] Thus, the ts and att phenotypes of B/Ann Arbor/1/66 are determined by at most three genes. The conversion of eight amino acids in the PA, NP, and M1 protein into wild type residues resulted in a recombinant virus that replicated efficiently at 37° C. Similarly, a 6+2 recombinant virus representing the six internal genes of MDV-B with the HA and NA segments from B/HongKong/330/01 showed a ts-phenotype and the triple recombinant was non-ts.

[0270] Our results using the MDV-B backbone indicated that six amino acids were sufficient to convert a ts/att phenotype into a non-ts/non-att phenotype. Therefore, we were interested in determining whether the introduction of those six 'attenuation' residues would transfer these biological properties to a heterologous wildtype, non attenuated influenza B virus, such as B/Yamanashi/166/98.

[0271] Recombinant wildtype B/Yamanashi/166/98 (recyam) (7) and a recombinant virus (rec6-Yam): with six amino acid changes PA (V431→M431, H497→Y497), NP (V114→A114, P410→H410), and M1 (H159→Q159, M183→V183) were produced. RecYam showed a 0.17 log₁₀ titer reduction in titer at 37° C. compared to 33° C., whereas rec6Yam was clearly ts, the difference in viral titer between 37° C. and 33° C. was 4.6 log₁₀. Virus was efficiently recovered from ferrets infected with recYam, as expected for a typical wildtype influenza B virus. When rec6Yam was inoculated into ferrets, no virus was detected in the lung tissues (Table 18). Thus, the transfer of the ts/att loci from MDV-B are sufficient to transfer the ts- and att-phenotypes to a divergent virus.

TABLE 18

Attenuation studies in ferrets

Recombinant virus	wt components ^a	Ts-phenotype	ferrets	Dose [log10pfu]	Nasal turbinates ^b [log10pfu/g]	Lung tissue [log10EID50/g] ^c
rMDV-B	none	ts	4	6.0	4.01	<1.5
rec53-B	NP, PA, M	Non-ts	4	6.0	4.65	3.81
rec62-B	NP, PA	Non-ts	4	6.0	4.69	<1.5
rec71NP-B	NP	ts	4	6.0	4.13	<1.5

TABLE 18-continued

Attenuation studies in ferrets						
Recombinant virus	wt components ^a	Ts-phenotype	Dose ferrets	Nasal turbinates ^b	Lung tissue	
rec71M-B	M	ts	4	6.0	4.17	<1.5
RecYam		Non-ts	4	6.0	4.92	3.31
rec6Yam		ts	4	6.0	4.02	<1.5

^aRecombinant viruses with MDV-B backbone that differed in wildtype amino acids (for details see table 2) were used to infected ferrets intranasally. RecYam is recombinant B/Yamanashi/166/98 and Rec6Yam represents a virus that has six 'MDV-B-attenuation' amino acid changes in NP, PA, and M1 with a B/Yamanashi backbone.

^bThree days after infection the virus titer of the nasal turbinates and lung tissue was determined, the average titer of four infected ferrets is shown.

^c<1.5 indicates that no virus was detected.

[0272] As described above with respect to influenza A strains, substitution of the residues indicated above, e.g., PB2 (S630R); PA⁴³¹ (V431M); PA⁴⁹⁷ (Y497H); NP⁵⁵ (T55A); NP¹¹⁴ (V114A); NP⁴¹⁰ (P410H); NP⁵⁰⁹ (A509T); M1¹⁵⁹ (H159Q) and M1¹⁸³ (M183V), confers the ts and att phenotypes. Accordingly, artificially engineered variants of influenza B strain virus having one or more of these amino acid substitutions exhibit the ts and att phenotypes and are suitable for use, e.g., as master donor strain viruses, in the production of attenuated live influenza virus vaccines.

Example 9

Rescue of Influenza from Eight Plasmids by Electroporation of Vero Cells

[0273] Previously it has been suggested that recombinant influenza A can be rescued from Vero cells (Fodor et al. (1999) *Rescue of influenza A virus from recombinant DNA J. Virol.* 73:9679-82; Hoffmann et al. (2002) Eight-plasmid system for rapid generation of influenza virus vaccine *Vaccine* 20:3165-3170). The reported method requires the use of lipid reagents and has only been documented for a single strain of a highly replication competent laboratory strains of influenza A (A/WSN/33 and A/PR/8/34), making it of limited application in the production of live attenuated virus suitable for vaccine production. The present invention provides a novel method for recovering recombinant influenza virus from Vero cells using electroporation. These methods are suitable for the production of both influenza A and influenza B strain viruses, and permit the recovery of, e.g., cold adapted, temperature sensitive, attenuated virus from Vero cells grown under serum free conditions facilitating the preparation of live attenuated vaccine suitable for administration in, e.g., intranasal vaccine formulations. In addition to its broad applicability across virus strains, electroporation requires no additional reagents other than growth medium for the cell substrate and thus has less potential for undesired contaminants. In particular, this method is effective for generating recombinant and reassortant virus using Vero cells adapted to growth under serum free condition, such as Vero cell isolates qualified as pathogen free and suitable for vaccine production. This characteristic supports the choice of electroporation as an appropriate method for commercial introduction of DNA into cell substrates.

[0274] Electroporation was compared to a variety of methods for introduction of DNA into Vero cells, including transfection using numerous lipid based reagents, calcium phosphate precipitation and cell microinjection. Although

some success was obtained using lipid based reagents for the rescue of influenza A, only electroporation was demonstrated to rescue influenza B as well as influenza A from Vero cells.

[0275] One day prior to electroporation, 90-100% confluent Vero cells were split, and seeded at a density of 9×10⁶ cells per T225 flask in MEM supplemented with pen/strep, L-glutamine, nonessential amino acids and 10% FBS (MEM, 10% FBS). The following day, the cells were trypsinized and resuspend in 50 ml phosphate buffered saline (PBS) per T225 flask. The cells are then pelleted and resuspend in 0.5 ml OptiMEM I per T225 flask. Optionally, customized OptiMEM medium containing no human or animal-derived components can be employed (this can be obtained from the manufacturer of OptiMEM I upon request). Following determination of cell density, e.g., by counting a 1:40 dilution in a hemocytometer, 5×10⁶ cells were added to a 0.4 cm electroporation cuvette in a final volume of 400 µl OptiMEM I. Twenty µg DNA consisting of an equimolar mixture of eight plasmids incorporating either the MDV-A or MDV-B genome in a volume of no more than 25 µl was then added to the cells in the cuvette. The cells were mixed gently by tapping and electroporated at 300 volts, 950 microFarads in a BioRad Gene Pulser II with Capacitance Extender Plus connected (BioRad, Hercules, Calif.). The time constant should be in the range of 28-33 msec.

[0276] The contents of the cuvette were mixed gently by tapping and 1-2 min after electroporation, 0.7 ml MEM, 10% FBS was added with a 1 ml pipet. The cells were again mixed gently by pipetting up and down a few times and then split between two wells of a 6 well dish containing 2 ml per well MEM, 10% FBS. The cuvette was then washed with 1 ml MEM, 10% FBS and split between the two wells for a final volume of about 3.5 ml per well.

[0277] In alternative experiments, Vero cells adapted to serum free growth conditions, e.g., in OptiPro (SFM) (Invitrogen, Carlsbad, Calif.) were electroporated as described above except that following electroporation in OptiMEM I, the cells were diluted in OptiPro (SFM) in which they were subsequently cultured for rescue of virus. Subsequent experiments have shown that, following electroporation, cells may be diluted in OptiMEM I or customized OptiMEM medium containing no human or animal-derived components.

[0278] The electroporated cells were then grown under conditions appropriate for replication and recovery of the introduced virus, i.e., at 33° C. for the cold adapted Master Donor Strains. The following day (e.g., approximately 19 hours after electroporation), the medium was removed, and the cells were washed with 3 ml per well OptiMEM I or OptiPro (SFM). One ml per well OptiMEM I or OptiPro (SFM) containing pen/strep was added to each well, and the supernatants were collected daily by replacing the media. Supernatants were stored at -80° C. in SPG. Peak virus production was typically observed between 2 and 3 days following electroporation.

[0279] Therefore, the present invention includes an improved method of rescue, wherein animal cells (e.g., SF Vero cells) are electroporated with polynucleotides (e.g., plasmids and vectors) of the invention.

TABLE 19

Results of 8 Plasmid Rescue of MDV strains on Different Cell Types and by Different Transfection Methods

Substrate	Method	No of Test	Result (Infectious Virus Recovered)	
			MDV-B	
COS-7/MDCK	Lipo	3	positive	
COS-7/MDCK	CaPO4	2	positive	
MRC-5	Lipo	5	negative	
MRC-5	CaPO4	3	negative	
MRC-5	Electroporation	2	negative	
WI-38	Lipo	2	negative	
WI-38	Electroporation	4	negative	
WI-38	Microinjection	1	negative	
LF1043	Lipo	1	negative	
LF1043	CaPO4	2	negative	
Vero	Lipo	7	negative	
Vero	CaPO4	2	negative	
Vero/MDCK	Lipo	1	negative	
Vero (serum)	Electroporation	5	positive (5/5)	
Vero (serum free)	Electroporation	4	positive (4/4)	
MDV-A				
Vero (serum)	Electroporation	3	positive (3/3)	
Vero (serum Free)	Electroporation	3	positive (3/3)	

Example 10

Co-Cultivation of Electroporated SF Vero Cells Improves Efficiency of Rescue

[0280] As discussed above, influenza virus can be rescued from SF vero cells by electroporation of the cells with plasmids that encode each of the eight segments of the viral genome. This method can be used to make 6:2 viruses composed of the HA and NA from wild type strains of influenza and the PB1, PB2, PA, NP, NS, and M from a MDV strain, e.g., a cold-adapted MDV strain or PR8. For some wild type HA and NA segments, rescue in SF vero cells is inefficient. To this end, it has been found that co-cultivation of the electroporated SF vero cells with Chicken Embryo Kidney (CEK) cells improved the efficiency of the plasmid rescue. For example, when electroporation of SF vero cells was performed to rescue an A/Panama 6:2 virus, none of the 30 eggs tested (5 eggs/day, days 2-7 post-electroporation) had detectable HA titers. However, when an equal sample of the same electroporated SF vero cells was co-cultivated with CEK cells, 27 out of 30 eggs

had detectable HA titers (90% efficiency) and these titers were 100 or better. In addition, this improved rescue efficiency was also observed for MDV A. Further, A/Sendai (another 6:2 virus which is difficult to rescue from SF vero cells) has been rescued by the co-cultivation method.

[0281] Therefore, the present invention includes an improved method of rescue, wherein electroporated SF vero cells are co-cultivated with another cell selected from the group including, but not limited to: chicken embryo kidney (CEK) cells, chicken embryo fibroblasts, primary chick kidney cells, and cells isolated from the chorioallantoic membrane of embryonated chicken eggs. Other cells useful for this rescue method may include any cell that supports replication of influenza virus and meets acceptable standards for regulatory approval. Sources of cells include, for example, chicken flocks from SPF chicken flocks.

[0282] In one preferred embodiment of the invention, rescue efficiency of virus is improved by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 90%, or at least 2-fold, or at least 3-fold, or at least 5-fold.

[0283] In another preferred embodiment of the invention, rescue efficiency of virus is at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 99%. Efficiency can be determined, for example, by measuring how many eggs injected with the rescued viruses (X) have subsequent detectable HA titers (Y) and dividing Y/X.

Example 11

Influenza Virus Vector System for Gene Delivery

[0284] The vectors of the present invention can also be used as gene delivery systems and for gene therapy. For such applications, it is desirable to generate recombinant influenza virus, e.g., recombinant influenza A or B virus expressing a foreign protein. For example, because segment 7 of the influenza B virus is not spliced, it provides a convenient genetic element for the insertion of heterologous nucleic acid sequences. The mRNA contains two cistrons with two open reading frames encoding the M1 and BM2 proteins. The open reading frame of BM2 or M1 is substituted by the heterologous sequence of interest, e.g., a gene encoding the enhanced green fluorescent protein (EGFP). Using the plasmid based vector system of the present invention, the cDNA encoding the open reading frame of M1-EGFP and BM2 are cloned on two different plasmids. The open reading frame is flanked by the non coding region of segment 7, which contains the signals required for replication and transcription. Alternatively, two plasmids are constructed: one containing M1 ORF and the other containing EGFP-BM2. Co-transfection of the resultant nine plasmids results in the generation of a recombinant influenza B virus containing the heterologous gene sequence. Similarly, EGFP can be expressed from the NS1 segment of influenza A.

[0285] The exemplary "green" influenza B virus can be used for standardization in virus assays, such as micro neutralization assays. The combination of the plasmid based technology and the simple detection of protein expression (fluorescence derived from EGFP can be monitored by microscopy, as illustrated in FIG. 2), permits the optimization of protein expression.

Example 12

Genetic Studies of Recent H3N2 Influenza Vaccine Strains

[0286] The live attenuated cold-adapted influenza A/AA/6/60 strain, in typical preferred embodiments, is the master donor virus (MDV-A) for influenza A FluMist™ vaccines. The 6 internal genes of MDV-A confer the cold-adapted (ca) temperature sensitive (ts) and attenuated (att) phenotypes to each of the vaccine strains. Using reverse genetics, it is demonstrated that multiple amino acids segregated among three gene segments: PB1-K391E, E581G, A661T, PB2-N265S, and NP-D34G which control expression of the ts and att phenotypes of MDV-A. Plasmid rescue of 6:2 vaccine strains allows more efficient generation of influenza vaccines than classical reassortment techniques.

[0287] The inactivated influenza vaccines for the 2003-04 season contained the A/Panama/99 (H3N2) antigen and were unable to elicit robust antibody responses in seronegative children to the drifted A/Fujian/411/02-like H3N2 strains that circulated during this season. See FIGS. 22 and 23. Unfortunately, A/Fujian/411/02 did not replicate well in embryonated chicken eggs and, thus, prohibited its use for vaccine manufacture. Using the reverse genetics technology, we showed that the loss in the balance of the HA and NA activities was responsible for poor replication of the prototype A/Fujian/411/02 strain in eggs. See FIGS. 29 through 34. A/Fujian virus could gain its efficient replication in eggs by either increasing its HA activity or by reducing its NA activity. Specifically, we demonstrate that while a several different single amino acid substitution were able to slightly enhance the replication of A/Fujian/411/02 strain in eggs several combination gave a much more robust enhancement. See FIGS. 35 through 38. This work has demonstrated the feasibility of improving influenza virus growth in embryonated chicken eggs and/or host cells by introducing specific changes in the HA or NA genes without affecting virus antigenicity.

[0288] To produce a strain viable in eggs, a set of related H3N2 6:2 reassortants of the A/Fujian/411/02 lineage were evaluated for their replication in MDCK cells, embryonated eggs and ferrets. While A/Fujian/411/02 did not grow in eggs, an egg-adaptation of this virus resulted in two amino acid substitutions in HA, H183L and V226A which allowed for virus growth in embryonated eggs. Additionally, an egg-adapted A/Wyoming/03/2003 strain that grew well in eggs and ferrets and the A/Sendai/H-F4962/02 vaccine that grew well in eggs, but replicated poorly in ferrets, were compared in terms of sequence. It was determined that G186V and V226I in HA, and/or Q119E and K136Q in NA were required for efficient virus replication in vitro and in vivo. Nevertheless, these amino acid changes had no effect on virus antigenicity. Adoption of such techniques to produce strains capable of growth in eggs (for strains that are difficult/problematic to grow in eggs) or to produce strains more capable of growth in eggs (for strains that can already grow in eggs) for other influenza viruses is contemplated and expected.

[0289] The molecular basis for the antigenic drift from A/Panama/99 to A/Fujian/02-like strains was studied by changing clusters of HA residues from A/Panama/99 to those of A/Wyoming/03. See FIG. 24. Antigenicity of the

modified 6:2 reassortants were examined by HAI and micro-neutralization assays using ferret sera from animals immunized with either A/Panama/99 or A/Wyoming/03. See FIGS. 25 through 28. It was determined that only a few changes were responsible for antigenic drift while others had a more dramatic impact on virus replication. Thus, as indicated by the data, reverse genetics are optionally used to modify vaccine strains to increase vaccine yields without affecting virus antigenicity.

[0290] Materials and Methods

[0291] Virus strains, cells and antibodies: Wild-type (wt) influenza A virus strains, A/Fujina/411/02 (A/Fujian), A/Sendai-H/F4962/02 (A/Sendai) and A/Wyoming/03/03 (A/Wyoming), were obtained from the Center for Disease Control (Atlanta, Ga.) and amplified once in MDCK cells or in embryonated chicken eggs (eggs). The modified vaccinia virus Ankara strain expressing the bacteriophage T7 RNA polymerase (MVA-T7) was grown in CEK cells. HEp-2, COS-7 and MDCK cells (obtained from American Type Culture Collections, ATCC) were maintained in minimal essential medium (MEM) containing 5% fetal bovine serum (FBS). Polyclonal antisera against A/Ann Arbor/6/60, A/Sendai-H/F4962/02 and A/Wyoming/03/03 were produced in chicken. Monoclonal antibodies against the NP protein of influenza A were obtained from BioDesign (Saco, Mich.).

[0292] Generation of recombinant 6:2 reassortants: Recombinant 6:2 reassortants that contained the HA and NA RNA segments of the H3N2 strains reassorted into MDV-A, were generated according to the previously described procedures. Briefly, a set of six plasmids containing the internal genes of MDV-A together with the HA and NA expression plasmids were transfected into the co-cultured COS-7/MDCK cells using TRANSIT LT1 reagents (Mirus, Madison, Wis.). The transfected cell culture supernatant was collected at 3 days post transfection and used to infect fresh MDCK cells and 10-day-old embryonated chicken eggs. The infected MDCK cells were incubated at 33° C. until 80-90% cells exhibited cytopathic effect. The infected embryonated chicken eggs were incubated at 33° C. for three days and the allantoic fluids were collected and stored at -80° C. in the presence of the SPG stabilizer (0.2 M sucrose, 3.8 mM KH₂PO₄, 7.2 mM K₂HPO₄, 5.4 mM monosodium glutamate). Virus titer was determined by plaque assay on MDCK cells incubated under an overlay that consisted of 1× L15/MEM, 1% agarose and 1 µg/ml TPCK-trypsin at 33° C. for 3 days. The plaques were enumerated by immunostaining using chicken anti-MDV-A polyclonal antibodies.

[0293] Cloning of HA and NA expression plasmids: To make recombinant 6:2 reassortant viruses containing the HA and NA segments of H3N2 subtype and the six internal MDV-A RNA segments, the HA and NA cDNAs of wt A/Sendai-H/F4962/02 and A/Wyoming/03/03 were amplified by RT-PCR using SuperscriptIII reverse transcriptase (Invitrogen, Carlsbad, Calif.) and pfu DNA polymerase (Stratagene, La Jolla, Calif.), the extracted vRNA as template and the H3 and N2 specific primers. HA-AarI5 (5'cacttatattacatcgctcaggagcaaaggcagggtt3') and HA-AarI3 (5'cttaacatcatcacctgcctcgatttagtagaaacaagggtt3') primers were used to amplify the HA segment. N2-AarI5 (5'cacttatattacatcgctcaggagcaaaggcagggtt3') and N2-AarI3 (5'cttaacatcatcacctgcctcgatttagtagaaacaagggtt3') primers

were used to amplify the NA segment. Both the HA and NA primer pairs contained the Aar I restriction sites that was designed to be comparable to the BsmB I sites present in the pAD3000 pol I/pol II expression plasmid. The HA and NA cDNA clones were sequenced and compared to the consensus HA and NA sequences that were obtained by direct sequencing of the HA and NA RT-PCR amplified cDNA products. Any mutations introduced into the cDNA clones during the cloning process were corrected by QuickChange site-directed mutagenesis kit (Stratagene, La Jolla, Calif.).

[0294] HAI assay (Hemagglutination Inhibition Assay for Influenza Virus): Reagents: 0.5% cRBC (washed three times with PBS-, can be used within 2-3 days); 96-well U bottom microplate; PBS- (without Ca and Mg); Tips; Influenza virus; Serum samples and positive control serum of high and low titer Preparations: Determine HA titer of virus by HA assay (Use virus titer at 1:8 for HAI. If HA titer of a given virus is 1:256, divide it by 8. Thus, need to dilute virus 1:32. Prepare 2.5 ml of virus for each 96 well plate); Treat serum with RDE (receptor destroy enzyme) optional for ferrets samples; Prepare RDE as instructed by manufacturer; Combine RDE and serum sample at 1:4 dilution. For example, add 100 μ l of serum to 300 μ l of RDE. Vortex the mix and incubate overnight (18-20 hr) in 37° C. incubator. Heat mixture at 56° C. for 45-50 min. Screen serum for non-specific agglutinins; Mix 25 μ l of RDE-treated serum with 25 μ l of PBS- by pipetting up and down 3x; Add 50 μ l of 0.5% cRBC to the mix and to the control well with only PBS-; Incubate at RT for 30-45 min (+: indicates partial or complete non-specific hemagglutination -: indicates no hemagglutination); Non-specific cRBC agglutinins can be removed by pre-incubation of serum with packed RBC at 20:1 ratio at 4° C. for 1 hr, followed by centrifugation at 2000 rpm for 10 min at 4° C. 4) Controls can typically include the following: cRBC cell control; Virus back titration: 2-fold dilution of 8 units/50 μ l virus diluted from 1:2 to 1:32 to make sure that virus used is at the correct concentrations; Positive serum control: dilute known titer serum 2-fold serially together with the test serum samples. A typical HAI protocol can comprise: Dilute serum samples two-fold serially; Add 25 μ l of PBS- to each well; Add 25 μ l of virus to well 1A (e.g., 1:2), mix by pipetting up and down 3x; Transfer 25 μ l from well A to well B (e.g., 1:4) and mix as above 3x, repeat dilution until well H (e.g., 1:256); Add virus 25 μ l (8 unit/50 μ l) to diluted serum samples, mix up and down 3x and incubate at RT for 30-40 min; Add 50 μ l of 0.5% cRBC, mix well by pipetting up and down 3x; Incubate at RT for 30-45 min.; Record hemagglutination. The HAI titer is defined as the highest dilution of the serum that completely inhibits hemagglutination. If no inhibition is observed, the titer is <1:4. If all wells display inhibition, the titer is >1:256.

[0295] Measurement of the neuraminidase activity of the transiently expressed NA protein: To measure the neuraminidase activity of the NA proteins, wt NA and its modified derivatives were expressed from the plasmid transfected cells. To obtain a high level of expression of the NA proteins, the NA RNA was transcribed from the T7 and CMV promoters as the gene was inserted downstream of these dual promoters. HEp-2 cells in 10 cm dishes were infected with MVA-T7 at moi of 5.0 for 1 hr followed by transfection of 5 μ g of the NA plasmid using Lipofectamine 2000 reagent (Invitrogen, Carlsbad, Calif.). The transfected cells were incubated at 35° C. for 48 hr. After washing with

phosphate-buffered saline (PBS), the cells were scraped from the dishes and lysed in 100 μ l of 0.125M NaOAc, pH 5.0. The neuraminidase activity in the transfected cells was determined by a fluorimetric assay. After one time of freezing-thawing, 50 μ l of cell lysates were 2-fold serially diluted and incubated with 150 μ l of 1.2 mM 2'-(4-methylumbelliferyl)- α -D-N-Acetylneuraminic Acid (MU-NANA) substrate (Sigma, St. Louis, Mo.) at 37° C. for 1 hr and stopped by 75 μ l of 1.0 M Glycine (pH 5.5). The fluorescence level of the released chromophore 4-methylumbelliferon was determined at 362 nm on a SpectroMAX plate reader. The level of each NA protein expressed in the transfected cells was monitored by Western blotting using chicken anti-A/Wyoming antisera. The neuraminidase activities of wt A/Sendai and A/Wyoming viruses containing 6.0 \log_{10} PFU in 100 μ l were also measured by the fluorimetric assay.

[0296] Receptor binding and replication of 6:2 recombinants in MDCK cells: HA receptor-binding and growth kinetics of recombinant 6:2 reassortants were determined in MDCK cells. MDCK cells in six-well plates were infected with 6:2 A/Fujian, A/Sendai, A/Wyoming and two modified recombinant viruses at a moi of 1.0. After 30 min of adsorption at either 33° C. or 4° C., the infected cells were either washed three times with PBS, or directly overlaid with 3 ml of Opti-MEM I containing 1 μ g/ml TPCK-trypsin and incubated at 33° C. One set of the infected plates was fixed with 1% paraformaldehyde at 6 hr post infection for 15 min at room temperature, and permeabilized with 0.2% Triton X-100 in PBS for 15 min followed by immunofluorescence analysis using anti-NP monoclonal antibodies. The cell images captured by ORCA-100 digital camera were analyzed by Compix image capture and dynamic intensity analysis software, Version 5.3 (Cranberry Township, PA) to calculate the percentage of the infected cells. Another set of plates was incubated at 33° C. At various times of intervals, 250 μ l of culture supernatant was collected and stored at -80° C. in the presence of SPG prior to virus titration. After each aliquot was removed, an equal amount of fresh medium was added to the cells. The virus titer in these aliquots was determined by plaque assay on MDCK cells at 33° C.

[0297] To determine whether the binding difference between these viruses affected virus growth kinetics in MDCK cells, the infected MDCK cells were incubated at 33° C. and the culture supernatants were collected at various times for virus titration. When adsorbed at 33° C., 6:2 A/Fujian had slower growth kinetics and lower titer (FIG. 2), 6:2 A/Sendai, A/Fujian with HA-V1861226 or HA-L183A226 behaved similarly to 6:2 A/Wyoming. When adsorption was done at 4° C., 6:2 A/Fujian as well as 6:2 A/Sendai had slower growth kinetics. 6:2 A/Wyoming and the two A/Fujian variants grew similarly. These results were consistent with the virus-binding assay whereas the washing step reduced efficient infection of A/Fujian at both temperatures.

[0298] Antigenicity of 6:2 recombinant viruses: Antigenicity of each virus was analyzed by hemagglutinin inhibition (HAI) assay using ferret anti-A/Sendai and anti-A/Wyoming sera. Aliquots of 25 μ l of 2-fold serially diluted ferret antisera were incubated with 25 μ l virus containing 4 HA units of 6:2 reassortant viruses at 37° C. for 1 hr followed by incubation with 50 μ l of 0.5% turkey red blood cells

(RBC) at 25° C. for 45 min. The HAI titer was defined as the reciprocal of the highest serum dilution that inhibited hemagglutination.

[0299] Generation of 6:2 A/Fujian, A/Sendai, and A/Wyoming Vaccine Strains

[0300] Wild-type (wt) influenza A virus strains, A/Fujian/411/02, A/Sendai-H/F4962/02 and A/Wyoming/03/03 were obtained from the Center for Disease Control (Atlanta, Ga.) and amplified once in MDCK cells or in embryonated chicken eggs. As indicated in Table 20, A/Fujian was only passaged for three times in cell culture, whereas A/Sendai and A/Wyoming went through 11 passages in eggs. The HA and NA sequences of these three strains were determined by sequencing of the RT-PCR products using vRNA extracted from these viruses. The difference in the HA and NA sequence of these three H3N2 strains is listed in Table 1. A/Sendai was identical to A/Fujian in its HA1 amino acid sequence but differed in the NA sequence at three amino

[0302] To confirm that the HA and NA segments of these H3N2 strains controlled virus replication in eggs and cells, the HA and NA gene segments were reassorted with the internal gene segments of the cold adapted A/Ann Arbor/6/60 strain, the master donor virus for live attenuated influenza FluMist vaccines (MDV-A) to generate three 6:2 reassortant viruses. Replication of these three viruses was evaluated in MDCK cells and embryonated chicken eggs. 6:2 A/Fujian ($6.2 \log_{10}$ PFU/ml) showed a lower titer than 6:2 A/Sendai ($7.1 \log_{10}$ PFU/ml) and A/Wyoming ($7.0 \log_{10}$ PFU/ml) in MDCK cells. Similar to wt A/Fujian, 6:2 A/Fujian replicated poorly in embryonated chicken eggs with a titer of $4.1 \log_{10}$ PFU/ml. Both 6:2 A/Sendai and A/Wyoming replicated to higher titers of 8.7 and $8.1 \log_{10}$ PFU/ml, respectively. Thus, the transfer of the wt HA and NA gene segments into MDV-A did not change the capability of each virus to replicate in eggs.

TABLE 20

Comparison of wt and recombinant 6:2 A/Fujian/411/02-like strains in HA and NA sequence and their replication in MDCK cells and eggs.								
Virus strains	Amino acid positions							
	HA1		HA2		NA			
	128	186	219	226	150	119	136	347
A/Fujian/411/02 ⁽¹⁾ (C1/C2)	T	G	S	V	G	E	Q	H
A/Sendai-H/F4962/02 (Cx8/E3)	—	—	—	—	E	Q	K	Y
A/Wyoming/03/03 (ck2E2/E9)	A	V	Y/F	I	E	—	—	—

Virus strains	Virus titer (\log_{10} PFU/ml \pm SE) ⁽³⁾			
	MDCK		Eggs	
(Passage history)	wt	6:2	wt	6:2
A/Fujian/411/02 ⁽¹⁾ (C1/C2)	6.1 ± 0.3	6.2 ± 0.3 ⁽²⁾	4.1 ± 0.6	4.2 ± 0.5
A/Sendai-H/F4962/02 (Cx8/E3)	8.1 ± 0.2	7.1 ± 0.1	9.0 ± 0.3	8.7 ± 0.2
A/Wyoming/03/03 (ck2E2/E9)	6.7 ± 0.5	7.0 ± 0.4	8.9 ± 0.3	8.1 ± 0.1

⁽¹⁾wt A/Fujian had the H183L change after one time passage in MDCK cells and eggs.

⁽²⁾Recombinant 6:2 A/Fujian contained E150 in HA2.

⁽³⁾Virus titers were expressed as mean \log_{10} PFU/ml \pm SE from two or more samples.

acids at positions 119, 146 and 347. A/Wyoming had the NA sequence identical to that of A/Fujian, but differed from A/Fujian and A/Sendai in HA1 by four amino acids. In addition, both A/Sendai and A/Wyoming had Glu-150 instead of Gly-150 in the HA2. After one time of amplification in MDCK cells, the 183 residue in HA1 of wt A/Fujian mutated from His-183 to Leu-183 and it was difficult to isolate the wt A/Fujian virus with His-183, indicating that the virus with His-183 had growth advantage *in vitro*.

[0301] These three wt viruses grew differently in MDCK cells, reaching titers of 6.1 , 8.1 and $6.7 \log_{10}$ PFU/ml for wt A/Fujian, wt A/Sendai and wt A/Wyoming, respectively. wt A/Fujian replicated poorly in eggs, reaching a titer of $4.1 \log_{10}$ PFU/ml (Table 20). The virus isolated from eggs had the H183L change in the HA. In contrast, wt A/Sendai and wt A/Wyoming grew well in eggs having titers of 9.0 and $8.9 \log_{10}$ PFU/ml, respectively.

[0303] Effect of Amino Acid Changes in the NA on Neuraminidase Activities and Virus Replication

[0304] A/Fujian differed from A/Sendai by three amino acids in NA, E119Q, Q136K and H347Y (Table 20), it is hypothesized that one or more of these changes enabled A/Sendai to replicate in embryonated chicken eggs to a higher titer than A/Fujian. Substitutions of E119 by G, D, A or V residues have been reported for several anti-neuraminidase drug resistant strains that resulted in the reduced neuraminidase activity. To determine whether the E119Q or either of the other two changes in the NA had an effect on the NA activity of A/Fujian and on its ability to replicate in embryonated chicken eggs, single and double substitution mutations were introduced into A/Fujian NA expression plasmids and the NA activity in the transfected HEp-2 cells was measured. In addition, recombinant 6:2 recombinant viruses bearing mutations in the A/Fujian NA were also recovered and their growth in MDCK cells and eggs were

compared (Table 21). A/Fujian (E119Q136H147) had approximately 80% higher NA activity compared to that of A/Sendai (Q119K136Y147). Single Q119 mutation had 66% of NA activity, Y347 change had minimal effect on NA activity but K136 only had 25% activity. Double mutations, K136Y347, Q119Y347, and Q119K136 had reduced NA activity at levels of 29%, 52% and 25% of that A/Fujian, respectively. These data indicated that these three NA residues affected the NA activity in the order of K136>Q119>Y347.

[0305] The correlation of the NA activity of the NA mutants with virus replication in embryonated chicken eggs was examined (Table 21). The six modified viruses were shown to replicate well in MDCK cells reaching titers ranging from 6.2 to 6.9 \log_{10} PFU/ml, but replicated significantly different in eggs. FJ-Q119 and FJ-347 that had 66% and 99% NA activity of A/Fujian were unable to grow in eggs. FJ-K136 with 25% NA activity was able to grow to a titer of 4.8 \log_{10} PFU/ml in eggs, but 4.0 \log_{10} lower than that of A/Sendai (8.8 \log_{10} PFU/ml). Unexpectedly, although K136Y347 significantly decreased the NA activity in vitro, the recombinant virus carrying these two mutations (FJ-K136Y347) was not able to replicate in embryonated chicken eggs. Q119Y347 that had 52% of NA activity replicated in eggs to a titer of 4.5 \log_{10} PFU/ml. Q119K136 that had the NA activity slightly higher than that of A/Sendai replicated to a titer of 6.2 \log_{10} PFU/ml but was still 2.6 \log_{10} lower than A/Sendai. These results indicated that each of the three NA residues differed between A/Fujian and A/Sendai impacted virus replication differently. Although several NA mutations could reduce the NA activity to the level close to that A/Sendai, only Q136K and E119Q changes could result in significant improvement in virus replication in embryonated chicken eggs. Since the Q119K136 double mutations did not replicate as efficiently as A/Sendai virus in eggs, the Y347 residue might also affect virus replication in eggs.

TABLE 21

Effects of NA residues on virus replication in MDCK cells and embryonated eggs.

NA	NA residues			NA activity ⁽¹⁾ (Mean \pm SE)	Virus ⁽²⁾ titer (\log_{10} PFU/ml)	
	119	136	347		MDCK	Eggs
A/Fujian	E	Q	H	100 66 \pm 3	6.5	<1.5
FJ-Q119	Q	—	—	99 \pm 1	6.7	<1.5
FJ-Y347	—	—	Y	25 \pm 1	6.6	<1.5
FJ-K136	—	K	—	29 \pm 3	6.6	4.8
FJ-K136Y347	—	K	Y	52 \pm 4	6.6	4.5
FJ-Q119Y347	Q	—	Y	25 \pm 1	6.2	6.2
FJ-Q119K136	Q	K	—	21 \pm 1	6.9	8.8
A/SENDAI	Q	K	Y	—	—	—

⁽¹⁾The NA activities in NA cDNA-transfected HEp-2 cells are expressed as the percentage of that of A/Fujian (mean \pm standard error) from four independent experiments.

⁽²⁾Recombinant 6:2 viruses were generated using A/Fujian HA and NA or A/Fujian NA with mutations indicated.

[0306] Effects of HA Residues on Virus Replication

[0307] The changes of the four HA1 residues in A/Wyoming/03/03 that differed from A/Fujian were investigated for their roles in virus replication. The single and multiple substitution mutations were introduced into A/Fujian HA cDNA and the modified HA plasmids were introduced into MDV-A together with either A/Fujian NA. All of the 6:2 reassortant virus mutants replicated well in MDCK cells but

grew differently in embryonated chicken eggs (Table 33). The 6:2 reassortants with A/Fujian HA (T128G186S219V226) were unable to replicate in eggs. A single T128A change did not improve virus growth in eggs. However, single G186V or V226I change resulted in increased virus replication in eggs. Double G186V and V226I changes in HA replicated efficiently in eggs. Additional substitutions at residues 128 and/or 219 did not significantly increase virus replication. Thus, a minimal of two G186V and V226I changes enabled 6:2 A/Fujian to grow efficiently in embryonated chicken eggs.

TABLE 22

EFFECTS OF HA RESIDUES ON VIRUS REPLICATION IN EMBRYONATED EGGS.

Virus ⁽¹⁾	HA residues				Virus titer in eggs (\log_{10} PFU/ml)
	128	186	219	226	
A/Fujian	T	G	S	V	<1.5
HA-A128	A	—	—	—	<1.5
HA-V186	—	V	—	—	4.9
HA-I226	—	—	—	I	5.2
HA-V186I226	—	V	—	I	7.6
HA-V186Y219I226	—	V	Y	I	7.5
A/Wyoming	A	V	Y	I	7.3

⁽¹⁾Virus recovered from the transfected cells contained A/Fujian NA and HA with the indicated amino acid changes.

[0308] Adaptation of 6:2 A/Fujian/411/02

[0309] To determine whether 6:2 A/Fujian strain could be adapted to grow in embryonated chicken eggs, the virus was amplified in MDCK cells followed by passage in eggs (Table 23). When 3.0 \log_{10} PFU of virus was inoculated into an egg, less than 2.0 \log_{10} PFU/ml of virus was detected in the harvested allantonic fluid. Infectious virus could not be recovered following passages of this material. During the second passage experiment, the amount of virus inoculated into embryonated chicken eggs was increased to 5.9 \log_{10} PFU. A titer of 3.9 \log_{10} PFU/ml was detected in the harvested allantonic fluid (FJ-EP1) and an additional passage in eggs increased virus titer to 6.2 \log_{10} PFU/ml (FJ-EP2). A further passage in eggs (FJ-EP3) increased virus titer to 8.2 \log_{10} PFU/ml. Sequence analysis of the FJ-EP2 virus revealed an A to U mutation at nt 625 in the HA RNA segment which resulted in H183L change in the HA protein. Further analysis showed this change also occurred during virus amplification in MDCK cells. The H183L mutation was also found in the wt A/Fujian HA during its replication in MDCK and eggs as described previously. An additional U to C mutation at nt 754 of HA resulting in V226A substitution was found in the FJ-EP3 amplified virus (Table 23). No changes were detected in the NA segment.

[0310] To confirm that H183L and V226A mutations in HA were indeed responsible for the increased replication of 6:2 A/Fujian in eggs, H183L and V226A were introduced into A/Fujian HA singly or in combination. Three recombinant viruses were obtained and they grew to a titer of 7.4 \log_{10} PFU/ml for FJ-H183L, 7.9 \log_{10} PFU/ml for FJ-V226A and 8.4 \log_{10} PFU/ml for FJ-H183L/V226A (Table 23). Therefore, H183L and V226A independently contributed to the improved replication of A/Fujian virus in embryonated chicken eggs.

TABLE 23

Mutations in the HA of egg-adapted 6:2 A/Fujian revertants and their replication in embryonated eggs.

Virus	Mutations at nucleotide (amino acid)	Virus titers (\log_{10} PFU/ml)
<u>Egg-passaged</u>		
FJ-EP1	ND ¹	3.9
FJ-EP2	A625U (H183L)	6.2
FJ-EP3	A625U (H183L), U745C (V226A)	8.2
<u>Recombinants</u>		
FJ-183L	A625T (H183L)	7.4
FJ-226A	T745C (V226A)	7.9
FJ-183L/226A	A625U (H183L), U745C (V226A)	8.4

¹Not determined.

[0311] Receptor-Binding Properties and Replication of Recombinant Viruses

[0312] From the above studies, the NA changes that reduced the NA activity of A/Fujian were shown to be sufficient for this virus to grow in eggs. On the other hand, the HA changes (G186V and V226I or H183L and V226A)

a washing step. These data indicated that A/Fujian and A/Sendai HA had such a low binding affinity that the bound viruses at 4° C. could be readily washed off from the cells. The binding and virus entry kinetics were faster at 33° C., thus, the washing step had a minimal impact on 6:2 A/Sendai virus infection. However, the majority of the bound 6:2 A/Fujian was washed off at the similar condition because its higher NA activity prevented efficient virus binding at 33° C. (data not shown).

[0313] Antigenicity of Recombinant Viruses

[0314] To examine whether viruses with the modified HA and NA residues affected virus antigenicity, haemagglutination inhibition assay (HAI) was performed using ferret anti-A/Wyoming and anti-A/Sendai sera (Table 24). Anti-A/Wyoming or anti-A/Sendai ferret sera had a similar HAI titer when measured with either 6:2 A/Fujian or A/Sendai virus. A slightly higher HAI titer was detected with 6:2 A/Wyoming virus, probably due to the tighter binding of A/Wyoming HA to the cell receptor on the red blood cells. The two modified viruses (A/FujianHA-V186I226 and A/Fujian HA-L183A226) had HAI titer similar to A/Wyoming when measured by either serum. These results indicated that the amino acid difference between A/Sendai and A/Wyoming and the modified HA viruses generated in this study did not alter virus antigenicity.

TABLE 24

Virus ⁽¹⁾	Antigenicity of modified 6:2 A/Fujian viruses									
	HA					NA				
	128	183	186	219	226	119	136	347	anti-A/WY	anti-A/SD
A/Fujian	T	H	G	S	V	E	Q	H	9	9
A/Wyoming A	—	V	Y	I	—	—	—	—	11	10
HA-V186I226	—	—	V	—	I	—	—	Y	11	11
HA-L183A226	—	L	—	—	A	—	—	—	11	11

⁽¹⁾A/Fujian was grown in MDCK cells and the rest of viruses were grown in eggs.

⁽²⁾Antigenicity was measured by HAI assay using A/Wyoming (anti-A/WY) or A/Sendai (anti-A/SD) immunized ferret serum with the indicated virus antigens

might have increased receptor-binding affinity to compensate for the higher NA activity of A/Fujian. To determine whether the changes in the HA protein of A/Fujian increased its receptor-binding ability, adsorption of 6:2 A/Fujian carrying HA-V186I226 change and egg-adapted 6:2 A/Fujian that contained HA-L183A226 changes were compared to 6:2 A/Fujian, A/Sendai, and A/Wyoming. Each virus was adsorbed onto MDCK cells at moi of 1.0 for 30 min at 4° C. or 33° C., the inoculum was removed and the infected cells were washed three times or without the washing step. After 6 hr of incubation at 33° C., the percentage of the infected cells was determined by immunofluorescence analysis using anti-NP antibody. As shown in **FIG. 36**, 6:2 A/Fujian and A/Sendai infected 26-27% of cells when adsorption was performed at 4° C., but the majority of viruses were readily removed by the washing step. At 33° C., washing greatly reduced infection of 6:2 A/Fujian virus (6.2% compared to 37.8%) but did not have significant effect on the infection of 6:2 A/Sendai (42.8% compared to 51.7%). In contrast, 6:2 A/Wyoming, A/Fujian with HA-V186I226 or HA-L183A226 had similar infection rate no matter whether the cells were adsorbed at 4° C. or 33° C. and with or without

[0315] While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be clear to one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention. For example, all the techniques and apparatus described above may be used in various combinations. All publications, patents, patent applications, or other documents cited in this application are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication, patent, patent application, or other document were individually indicated to be incorporated by reference for all purposes.

[0316] In particular, the following patent applications are incorporated by reference in their entirety: U.S. Provisional Application Nos. 60/574,117, filed May 24, 2004; 60/578,962 file Jun. 12, 2004; 60/532,164 filed Dec. 23, 2003; PCT Application No. US03/12728, filed Apr. 25, 2003; and U.S. application Ser. No.10/423,828, filed Apr. 25, 2003.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 98

<210> SEQ ID NO 1
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer polyA.1

<400> SEQUENCE: 1

aacaatttgag atctcggta cctcagacat gataagatac attgtatgatg 50

<210> SEQ ID NO 2
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer polyA.2

<400> SEQUENCE: 2

tataactgca gactagtat atccttgttt attgcagctt ataatggta 50

<210> SEQ ID NO 3
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 3

cacttatattt cacctgcctc agggagcgaa agcaggctc 38

<210> SEQ ID NO 4
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 4

tattcgtctc agggagcgaa agcaggctt 30

<210> SEQ ID NO 5
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 5

tattcgtctc agggagcgaa agcaggctt 30

<210> SEQ ID NO 6
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 6

tattcgtctc agggagctaa agcaggctt a 31

-continued

<210> SEQ_ID NO 7
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 7

cacttatatt cacctgcctc agggagcaaa agcaggggg 38

<210> SEQ_ID NO 8
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 8

tattcgtctc agggagcaaa agcaggagtg a 31

<210> SEQ_ID NO 9
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 9

tattcgtctc agggagcaaa agcaggtaga t 31

<210> SEQ_ID NO 10
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 10

tattcgtctc agggagcaaa agcaggggtga 30

<210> SEQ_ID NO 11
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 11

cctaacatat cacctgcctc gtatttagtag aaacaaggta gttt 44

<210> SEQ_ID NO 12
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 12

atatcgtctc gtatttagtag aaacaaggca ttt 33

<210> SEQ_ID NO 13
<211> LENGTH: 33
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 13

atatcgctc gtattagtag aaacaaggta ctt

33

<210> SEQ ID NO 14
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 14

atatcgctc gtattagtag aaacaagggt att

33

<210> SEQ ID NO 15
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 15

cctaacatat cacctgcctc gtattagtag aaacaagggt gtt

43

<210> SEQ ID NO 16
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 16

atatcgctc gtattagtag aaacaaggag ttt

33

<210> SEQ ID NO 17
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 17

atatcgctc gtattagtag aaacaaggta gtt

33

<210> SEQ ID NO 18
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A RT-PCR

<400> SEQUENCE: 18

atatcgctc gtattagtag aaacaagggt gtt

33

<210> SEQ ID NO 19
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

-continued

<400> SEQUENCE: 19
gcaagctgtg gaaatatgca aggc 24

<210> SEQ ID NO 20
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 20
gccttgcata tttccacagc ttgc 24

<210> SEQ ID NO 21
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 21
gaagtgccta cgggcaatct tcaaac 26

<210> SEQ ID NO 22
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 22
gtttgaagat tgcccgtaag cacttc 26

<210> SEQ ID NO 23
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 23
cctgaggagg tcagtgaaac ac 22

<210> SEQ ID NO 24
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 24
gtgtttcact gacccctca gg 22

<210> SEQ ID NO 25
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:

-continued

<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 25

gtttgttagg actctattcc aac

23

<210> SEQ ID NO 26

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 26

gttggaaatag agtcctaaca aac

23

<210> SEQ ID NO 27

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 27

gacagtaagc tccgaacaca aatac

25

<210> SEQ ID NO 28

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 28

gtatttgtgt tcggagcttc atgc

24

<210> SEQ ID NO 29

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 29

cgaaccgaac ggctacattg aggg

24

<210> SEQ ID NO 30

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 30

ccctcaatgt agccgttcgg ttcg

24

<210> SEQ ID NO 31

<211> LENGTH: 24

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 31

cagagaaggt agatggacg actg

24

<210> SEQ ID NO 32
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 32

cagtcgtcaa agtctaccc ttctcg

25

<210> SEQ ID NO 33
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 33

cactgaccca agacttgagc cac

23

<210> SEQ ID NO 34
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 34

gtggctcaag tcttgggtca gtg

23

<210> SEQ ID NO 35
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 35

caaagattaa aatgaaatgg ggaatg

26

<210> SEQ ID NO 36
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 36

cattcccat ttcatttaa tctttg

26

-continued

<210> SEQ ID NO 37
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 37

gtaccttgtt tctactaata acccg

26

<210> SEQ ID NO 38
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 38

ccgggttatt agtagaaaca aggtac

26

<210> SEQ ID NO 39
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 39

ggaacacttg agaactgtga gacc

24

<210> SEQ ID NO 40
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 40

ggtctcacag ttctcaagtg ttcc

24

<210> SEQ ID NO 41
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 41

gaattttatc acaaatgtga tgatgaatg

29

<210> SEQ ID NO 42
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 42

cattcatcat cacatttgtg ataaaattc

29

-continued

<210> SEQ ID NO 43
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 43

gccagaatgc aactgaaatc agagc

25

<210> SEQ ID NO 44
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 44

gctctgattt cagtttcatt ctggc

25

<210> SEQ ID NO 45
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 45

ccgaatgaga atccagcaca caag

24

<210> SEQ ID NO 46
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 46

cttgcgtgtc ggattctcat tcgg

24

<210> SEQ ID NO 47
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 47

catcaatttc atgcctatat aagcttgc

28

<210> SEQ ID NO 48
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

-continued

<400> SEQUENCE: 48
gaaagcttat ataggcatga aattgatg 28

<210> SEQ ID NO 49
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 49
cataatggat cctaacaactg tgtcaagc 28

<210> SEQ ID NO 50
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 50
gcttgacaca gtgttaggat ccattatg 28

<210> SEQ ID NO 51
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 51
ggagaataga ttcatcgaga ttggag 26

<210> SEQ ID NO 52
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for MDV-A mutation correction

<400> SEQUENCE: 52
ctccaatctc gatgaatcta ttctcc 26

<210> SEQ ID NO 53
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann Arbor/1/66 RT-PCR

<400> SEQUENCE: 53
tattcgtctc agggagcaga agcggagcct ttaagatg 38

<210> SEQ ID NO 54
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial

-continued

<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
Arbor/1/66 RT-PCR

<400> SEQUENCE: 54

tattcgtctc gatgccgttc cttcttcatt gaagaatgg 39

<210> SEQ ID NO 55
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
Arbor/1/66 RT-PCR

<400> SEQUENCE: 55

tattcgtctc ggcatcttg tcgcctggga ttagtgatg 38

<210> SEQ ID NO 56
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
Arbor/1/66 RT-PCR

<400> SEQUENCE: 56

atatcgtctc gtatttagtag aaacacgagc ctt 33

<210> SEQ ID NO 57
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
Arbor/1/66 RT-PCR

<400> SEQUENCE: 57

tattcgtctc agggagcaga agcggagcgt tttcaagatg 40

<210> SEQ ID NO 58
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
Arbor/1/66 RT-PCR

<400> SEQUENCE: 58

tattcgtctc tctcattttg ctcttttta atattcccc 39

<210> SEQ ID NO 59
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
Arbor/1/66 RT-PCR

<400> SEQUENCE: 59

tattcgtctc atgagaatgg aaaaactact aataaattca gc 42

<210> SEQ ID NO 60
<211> LENGTH: 33

-continued

```
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
    Arbor/1/66 RT-PCR
```

```
<400> SEQUENCE: 60
```

```
atatcgcttc gtattagtag aaacacgagc att
```

```
33
```

```
<210> SEQ ID NO 61
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
    Arbor/1/66 RT-PCR
```

```
<400> SEQUENCE: 61
```

```
tattcgcttc agggagcaga agcggtgcgt ttga
```

```
34
```

```
<210> SEQ ID NO 62
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
    Arbor/1/66 RT-PCR
```

```
<400> SEQUENCE: 62
```

```
tattcgcttc ccagggccct tttacttgtc agagtgc
```

```
37
```

```
<210> SEQ ID NO 63
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
    Arbor/1/66 RT-PCR
```

```
<400> SEQUENCE: 63
```

```
tattcgcttc tcctggatct accagaaata gggccagac
```

```
39
```

```
<210> SEQ ID NO 64
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
    Arbor/1/66 RT-PCR
```

```
<400> SEQUENCE: 64
```

```
atatcgcttc gtattagtag aaacacgtgc att
```

```
33
```

```
<210> SEQ ID NO 65
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann
    Arbor/1/66 RT-PCR
```

```
<400> SEQUENCE: 65
```

```
tattcgcttc agggagcaga agcagagcat tttctaataat c
```

```
41
```

-continued

<210> SEQ ID NO 66
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann Arbor/1/66 RT-PCR

<400> SEQUENCE: 66

atatcgcttc gtatttagtag taacaagagc atttttc 37

<210> SEQ ID NO 67
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann Arbor/1/66 RT-PCR

<400> SEQUENCE: 67

tattggcttc agggagcaga agcacagcat tttcttgt 38

<210> SEQ ID NO 68
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann Arbor/1/66 RT-PCR

<400> SEQUENCE: 68

atatggcttc gtatttagtag aaacaacagc attttt 36

<210> SEQ ID NO 69
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann Arbor/1/66 RT-PCR

<400> SEQUENCE: 69

tattcgcttc agggagcaga agcagagcat cttctaaaaa c 41

<210> SEQ ID NO 70
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann Arbor/1/66 RT-PCR

<400> SEQUENCE: 70

atatcgcttc gtatttagtag taacaagagc atttttcag 39

<210> SEQ ID NO 71
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann Arbor/1/66 RT-PCR

<400> SEQUENCE: 71

tattcgcttc agggagcaga agcacgcact ttctaaaaat g 41

-continued

<210> SEQ ID NO 72
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann Arbor/1/66 RT-PCR

<400> SEQUENCE: 72

atatcgcttc gtatttagtag aaacaacgca cttttccag 40

<210> SEQ ID NO 73
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann Arbor/1/66 RT-PCR

<400> SEQUENCE: 73

tattcgcttc agggagcaga agcagaggat ttgttttagtc 40

<210> SEQ ID NO 74
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for influenza ca B/Ann Arbor/1/66 RT-PCR

<400> SEQUENCE: 74

atatcgcttc gtatttagtag taacaagagg atttttat 38

<210> SEQ ID NO 75
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for B/Yamanashi/166/98 NP amplification

<400> SEQUENCE: 75

tattcgcttc agggagcaga agcacagcat tttcttg 39

<210> SEQ ID NO 76
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for B/Yamanashi/166/98 NP amplification

<400> SEQUENCE: 76

atatcgcttc gtatttagtag aaacaacagc attttttac 39

<210> SEQ ID NO 77
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for B/Yamanashi/166/98 NA amplification

-continued

<400> SEQUENCE: 77
tattcgtctc agggagcaga agcagagca 29

<210> SEQ ID NO 78
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for B/Yamanashi/166/98
NA amplification

<400> SEQUENCE: 78
atatcgtctc gtatttagtag taacaagagc atttt 35

<210> SEQ ID NO 79
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for introducing ts
mutations into PR8 PB1 and PB2 genes

<400> SEQUENCE: 79
gaaagaagat tgaagaaatc cgaccgctc 29

<210> SEQ ID NO 80
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for introducing ts
mutations into PR8 PB1 and PB2 genes

<400> SEQUENCE: 80
gagcgggtcgg atttcttcaa tcttcttcc 29

<210> SEQ ID NO 81
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for introducing ts
mutations into PR8 PB1 and PB2 genes

<400> SEQUENCE: 81
gaaataaaaga aactgtgggg gcaaaccctgt tcc 33

<210> SEQ ID NO 82
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for introducing ts
mutations into PR8 PB1 and PB2 genes

<400> SEQUENCE: 82
ggaacgggtt tgcccccaca gtttctttat ttc 33

<210> SEQ ID NO 83
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial

-continued

<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for introducing ts mutations into PR8 PB1 and PB2 genes

<400> SEQUENCE: 83

gtatgatgct gttacaacaa cacactcc

28

<210> SEQ ID NO 84
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for introducing ts mutations into PR8 PB1 and PB2 genes

<400> SEQUENCE: 84

ggagtgtgtt gttgttaacag catcatac

28

<210> SEQ ID NO 85
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for introducing ts mutations into PR8 PB1 and PB2 genes

<400> SEQUENCE: 85

attgctgcta ggagcatagt gagaagagc

29

<210> SEQ ID NO 86
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for introducing ts mutations into PR8 PB1 and PB2 genes

<400> SEQUENCE: 86

gctcttctca ctagctcct agcagcaat

29

<210> SEQ ID NO 87
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for RT-PCR of HA and NA

<400> SEQUENCE: 87

tattcgtctc agggagcaga agcagagca

29

<210> SEQ ID NO 88
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for RT-PCR of HA and NA

<400> SEQUENCE: 88

atatcgtctc gtatttagtag taacaagagc atttt

35

<210> SEQ ID NO 89
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial

-continued

<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for primer extension

<400> SEQUENCE: 89

atgttcttta cgatgcgatt ggg 23

<210> SEQ ID NO 90
<211> LENGTH: 2836
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: pAD3000

<400> SEQUENCE: 90

ctagcagtttta accggagttac tggtcgaccc tccgaagttgg gggggaggag acggtaccgt 60
ctccaataaac ccggccggccc aaaatgcgca ctcggagcga aagatataacc tccccccgggg 120
ccgggagggtc gcgtcaccga ccacgcccgc ggcccaggcg acgcgcgaca cggacacactg 180
tccccaaaaa cgccaccatc gcagccacac acggagcgcgc cggggccctc tggtaacacc 240
caggacacac cggggagcag cggccggccg gggacgcctt cccgggggtc acctaagaca 300
tgataagata cattgtatgg tttggacaaa ccacaacttag aatgcagttga aaaaaatgtct 360
ttatgttgc aattttgtat gctattgtt tattttgttaac cattataaagc tgcaataaac 420
aaggatctgc attaatgaat cggccaaacgc gcggggagag gcgggttgcg tattggcgc 480
tcttcgcctt cctcgctcac tgactcgctg cgctcggtcg ttccggctgcg gcgagcggt 540
tcagctcaact caaaggcggt aatacggtttca tccacacaaat caggggatataa cgcaggaaag 600
aacatgttag caaaaggccca gcaaaaggcc aggaaccgtt aaaaaggccgc gttgtggcg 660
ttttccata ggctccgcggcc ccctgacgag catcacaaaa atcgacgcctc aagtcaagagg 720
tggcgaaacc cggacaggact ataaagatac caggcggttc cccctggaaag ctccctcg 780
cgctctccgtt ttccggaccct gccgcttacc ggataccctgt ccgcctttct cccttggga 840
agcgtggcgc ttctcaatg ctcacgctgtt aggtatctca gttcggtgtt ggtcggttc 900
tccaagctgg gctgtgtca cgaacccccc gttcagcccg accgcgtgcgc cttatccgtt 960
aactatcgcc ttgagtccaa cccggtaaga cacgacttat cgccactggc agcagccact 1020
gttaacagga tttagcagagc gaggtatgtt ggcgggtgttca cagagttttt gaagtgggtt 1080
cctaactacg gctacactacg aaggacagta tttggtatct ggcgtctgtt gaagccagtt 1140
accttcggaa aaagagttgg tagctcttgc tccggcaaac aaaccaccgc tggtagcggt 1200
gttttttttgg tttgcaagca gcagattacg cgcagaaaaa aaggatctca agaagatccct 1260
ttgatctttt ctacgggttc tgacgcttagtggaaacgaaa actcacgttta agggatttt 1320
gtcatgagat tatcaaaaag gatcttacc tagatcttt taaattaaaa atgaagtttt 1380
aaatcaatct aaagtatata ttagttaact tggctgtaca gttaccaatg cttaatcagt 1440
gaggcaccta tctcagcgat ctgtcttattt cggtcatcca tagttgcgtgc actccccgtc 1500
gtgttagataa ctacgatacg ggagggttta ccatctggcc ccagtgtgtc aatgataaccg 1560
cgagacccac gtcacccggc tccagattta tcagcaataa accagccagc cggaaaggcc 1620
gagcgcagaa gtgggtcctgc aactttatcc gcctccatcc agtctattaa ttgttgcgg 1680
gaagctagag taagtagttc gccagttaaat agtttgcgtca acgttgcgttgc cattgttaca 1740
ggcatcggttgg tggcatcgctc gtcgtttggt atggcttcat tcagctccgg ttcccaacgaa 1800

-continued

tcaaggcgag ttacatgatc ccccatgttg tgcaaaaaag cggttagctc cttcggtcct	1860
ccgatcgttg tcagaagtaa gttggccgca gtgttatcac tcatggttat ggcagactg	1920
cataattctc ttactgtcat gccatccgta agatgcttt ctgtgactgg tgagtactca	1980
accaagtcat tctgagaata gtgtatgcgg cgaccgagtt gctcttgcgg ggcgtcaata	2040
cggataata ccgcgccaca tagcagaact taaaagtg tcatcatgg aaaacgttct	2100
tcggggcgaa aactctcaag gatcttaccc ctgtttagat ccagttcgat gtaaccact	2160
cgtgcaccca actgatcttc agcatcttt actttcacca gcgtttctgg gtgagcaaaa	2220
acaggaaggc aaaatgccgc aaaaaaggga ataaggcgca cacggaaatg ttgaataactc	2280
atactcttcc ttttcaata ttattgaagc atttatcagg gttattgtct catgagcgga	2340
tacatatttg aatgtattta gaaaaataaaa caaatagggg ttccgcgcac atttccccga	2400
aaagtgccac ctgacgtcga tatgccaagt acgcccccta ttgacgtcaa tgacggtaaa	2460
tggccgcct ggcattatgc ccagtagatc accttatggg actttccatc ttggcagttac	2520
atctacgtat tagtcatcgc tattaccatg gtgtatgggt tttggagta catcaatggg	2580
cgtggatagc ggtttactc acggggattt ccaagtctcc accccatgtc cgtcaatggg	2640
agtttgggggg ggcaccaaaa tcaacgggac tttccaaaat gtcgtaccaa ctccgcggca	2700
ttgacgcaaa tggggcgtag gcgtgtacgg tgggaggctt atataaggag agctctctgg	2760
ctaacttagag aacccactgc ttactggctt atcgaaattha atacgactca ctatagggag	2820
acccaagctg ttaacg	2836

<210> SEQ_ID NO 91
 <211> LENGTH: 2369
 <212> TYPE: DNA
 <213> ORGANISM: Influenza B virus

<400> SEQUENCE: 91

agcagaagcg gaggcttaa gatgaatata aatccttatt ttctcttcatt agatgtaccc	60
atacaggcag caatttcaac aacattccca tacaccgggt ttcccccattt ttcccatggaa	120
acgggaacag gctacacaat agacaccgtt attagaacac atgagtactc aaacaaggga	180
aaacaataca ttctgtatgt tacaggatgt gcaatggtag atccaacaaa tggggcattt	240
cccgaaagata atgagccgag tgcctatgca caattggatt gcgttctggaa ggctttggat	300
agaatggatg aagaacatcc aggtctgttta caagcagcct cacagaatgc catggaggca	360
ctaatggta caactgtaga caaattaacc caggggagac agacttttgaa ttggacagt	420
tgcagaaacc aacctgctgc aacggcactt aacacaacaa taacctttt taggttgaat	480
gatttgaatg gagccgacaa gggtggatta gtaccctttt gccaagatata cattgattca	540
ttggacaaac ctgaaatgac ttcttctcg gtaaagaata taaagaaaaa attgcctgt	600
aaaaacagaa agggtttcct cataaagaga ataccaatga aggtaaaaga cagaataacc	660
agagtggaaat acatcaaaag agcattatca taaaacacaa tgacaaaaga tgctgaaaga	720
ggcaaactaa aaagaagagc aattgccacc gctgggatc aaatcagagg gtttggat	780
gtagttgaaa acttggctaa aaatatctgt gaaaatcttag aacaaaatggg tttggccagta	840
ggtgggaaacg agaagaaggc caaactgtca aatgcgtgg cccaaatgct cagtaactgc	900
ccaccaggag ggatcagcat gacagtgaca ggagacaata ctaaatggaa tgaatgttta	960

-continued

aatccaagaa	tcttttggc	tatgactgaa	agaataacca	gagacagccc	aatttggttc	1020
cgggattttt	gtagtatagc	accggcttgc	ttctccaata	aatatagccag	attggaaaaa	1080
gggttcatga	taacaagcaa	aacaaaaaga	ctgaaggctc	aaataccctg	tcccgatctg	1140
ttaataatac	cattagaaag	atataatgaa	gaaacaaggg	caaattaaa	aaagctgaaa	1200
ccattcttca	atgaagaagg	aacggcatct	ttgtcgctg	ggatgtatgt	ggaaatgttt	1260
aatatgctat	ctaccgtgtt	gggagtagcc	gcactaggaa	tcaaaaacat	tggaaacaaa	1320
gaatacttat	gggatggact	gcaatcttct	gatgattttg	ctctgtttgt	taatgcaaaa	1380
gatgaagaga	catgtatgg	aggaataaac	gatttttacc	gaacatgtaa	gctattggaa	1440
ataaacatga	gcaaaaagaa	aagttactgt	aatgaaactg	aatgtttga	atttacaagc	1500
atgttctaca	gagatggatt	tgtatctaat	tttgcaatgg	aacttccctc	atttggagtt	1560
gctggagtaa	atgaatcagc	agatatggca	ataggaatga	caataataaa	gaacaatatg	1620
atcaacaatg	ggatgggtcc	agcaacagca	caaacagcc	tacaattatt	catagctgat	1680
tatagataca	cctacaaatg	ccacagggaa	gattccaaag	tggaggaaa	gagaatgaaa	1740
attataaagg	agctatggaa	aaacactaa	ggaagagatg	gtctgttagt	agcagatggt	1800
gggcctaaca	tttacaattt	gagaaacttg	catatcccag	aaatagtatt	aaagtacaac	1860
ctaatggacc	ctgataacaa	agggcggtt	ctgcacccctc	aaaatccctt	tgtaggacat	1920
ttgtctattt	agggcatcaa	agggcagat	ataacccctg	cacatggtcc	agtaaagaaa	1980
atggactatg	atcggttac	tggactcat	agttggagaa	ccaaaaggaa	cagatctata	2040
ctaaacactg	atcagaggaa	catgatttct	gaggaacaat	gctacgctaa	gtgttgcac	2100
ctttttgagg	cctgttttaa	cagtgcata	tacaggaaac	cagtaggtca	gcacagcatg	2160
cttgaggcta	tggcccacag	attaagaatg	gatgcacgac	tagattatga	atcaggaaga	2220
atgtcaaaagg	atgattttga	gaaagcaatg	gctcacctt	gtgagatgg	gtacatataa	2280
gcttcgaaga	tgtctatggg	gttattggtc	atcattgaat	acatgcggta	cacaaatgat	2340
taaaatgaaa	aaaggcttgt	gtttctact				2369

<210> SEQ_ID NO 92
 <211> LENGTH: 2396
 <212> TYPE: DNA
 <213> ORGANISM: Influenza B virus

<400> SEQUENCE: 92

agcagaagcg	gagcgtttc	aaagatgacat	tggccaaat	tgaattgtta	aaacaactgt	60
taagggacaa	tgaagccaa	acggatttga	aacaaacaac	ggttagacca	tataacataa	120
taagaaaatt	caatacatca	agaattgaaa	agaacccttc	attaaggatg	aagtggccca	180
tgtgttctaa	ttttcccttgc	gctctgacca	agggtgatat	ggcaaataga	atccccttgg	240
aataacaagg	aataacaactt	aaaacaaatg	ctgaagacat	aggaacccaa	ggccaaatgt	300
gctcaatagc	agcagttacc	tggtggata	catatggacc	aataggatg	actgaagggt	360
tcgaaaaggt	ctacgaaagc	tttttctca	gaaagatgag	acttgacaat	gccacttgg	420
gccgaataac	tttggccca	gttggaaagag	tgagaaaaag	ggtactgcta	aaccctctca	480
ccaaaggaaat	gcctccagat	gaagcgagca	atgtgataat	ggaaatattt	ttccctaaag	540
aagcaggaat	accaagagaa	tctacttgg	tacataggaa	actgataaaa	gaaaaaagag	600

-continued

<210> SEQ_ID NO 93
<211> LENGTH: 2308
<212> TYPE: DNA
<213> ORGANISM: Influenza B virus

```
<400> SEQUENCE: 93  
agcagaagcg gtgcgttga tttgccataa tggatacttt tattacaaga aacttccaga 60  
ctacaataat acaaaaaggcc aaaaacacaa tggcagaatt tagtgaagat cctgaattac 120  
aaccagcaat gctattcaac atctgcgtcc atctggaggt ctgctatgta ataagtgata 180  
tgaattttct tcatgttggaa ggaaaaacat atacagcatt agaaggacaa ggaaaaagac 240
```

-continued

aaaacctgag	accacaatat	gaagtgattt	agggaatgcc	aagaaaacata	gcatggatgg	300
ttcaaagatc	cttagccaa	gagcatggaa	tagagactcc	aaggtatctg	gctgatttgc	360
tcgattataa	aaccaagagg	tttatagaag	tttggataaac	aaagggattt	gctgacgatt	420
acttttggaa	aaagaaagaa	aagctgggg	atagcatgga	actgtatgata	ttcagctaca	480
atcaagacta	ttcgtaagt	aatgaatcct	cattggatga	ggaaggaaaa	gggagagtgc	540
taagcagact	cacagaactt	caggctgatgt	taagtctgaa	aatctatgg	caagttctca	600
taggagaaga	agatattgaa	aaaggaattt	acttcaaact	tggacaaaca	atatctaaac	660
taaggatata	atctgttcca	gctggtttct	ccaattttga	aggaatgagg	agctacatag	720
acaatataga	tcctaaagga	gcaatagaga	gaaatctagc	aaggatgtct	cccttagtat	780
cagttacacc	taaaaatgtt	aaatgggg	acctaagacc	aatagggct	cacatttaca	840
accatgagct	accagaagtt	ccatataatg	cctttttct	aatgtctgt	gagttggggc	900
tggctaata	gactgaaggg	aagtccaa	aaccgaagac	cttagccaa	aatgtcttag	960
aaaagtactc	aacactacgg	gatcaaactg	acccaatatt	aataatgaaa	agcggaaaag	1020
ctaacgaaaa	cttcttatgg	aagctgttgg	gggactgtgt	aaatacaata	agtaatgagg	1080
aaacaagtaa	cgaattacag	aaaaccaatt	atgccaagtg	ggccacagga	gatggattaa	1140
cataccagaa	aataatgaaa	gaagtagcaa	tagatgacga	aacaatgtac	caagaagac	1200
ccaaaatacc	taacaaatgt	agagtggctg	cttgggttca	aacagagatg	aatctattga	1260
gcactctgac	aagtaaaaagg	gccctggatc	taccagaaat	agggccagac	gtggcaccca	1320
tggagcatgt	agggagtgaa	agaaggaaat	actttgttaa	tgaatcaac	tactgttaagg	1380
cctctaccgt	tatgtgaag	tatgtacttt	ttcacacttc	attattaaat	gaaagoatg	1440
ccagcatggg	aaaatataaa	gtaataccaa	taaccaacag	agtagtaat	aaaaaggag	1500
aaagtttga	catgcttcat	ggctggcgg	ttaaaggggca	atctcatctg	agggggagata	1560
ctgatgttgt	aacagttgtg	actttcaat	ttagtagtac	agatcccaga	gtggactcag	1620
gaaagtggcc	aaaatatact	gtatgtttagaa	ttggctccctt	atttgtgatgt	ggaaggaaaa	1680
aatctgttta	cctatattgc	cgagtgaatg	gtacaaataa	gatccaaatg	aatggggaa	1740
tggaaagctag	aagatgtctg	cttcaatcaa	tgcaacaaat	ggaagcaatt	gttgaacaag	1800
aatcatcgat	acaaggatata	gacatgacca	aagcttgtt	caaggagac	agagtgaata	1860
gtccccaaac	tttcagtatt	gggactcaag	aaggaaaact	agtaaaaagg	tcctttggga	1920
aagcactaag	agtaatattc	accaaatgtt	tgtatgcacta	tgtatgttgg	aatgccaat	1980
tggagggttt	tagtggccaa	tcttaggagac	ttctactgtt	aattcaggca	ttaaaggaca	2040
gaaaggggccc	ttgggtattt	gacttagagg	gaatgtattt	tggatagaa	aatgttatta	2100
gtaacaaccc	ttgggttata	cagagtgcatt	actggtttaa	tgaatgggtt	ggctttgaaa	2160
aagagggggag	taaagtatta	gaatcaatag	atgaaataat	ggatgaatga	aagaaggc	2220
tagcgctcaa	tttggtacta	tttgttcat	tatgtatctt	aacatccaaat	aaaaagaattt	2280
gagaattaaa	aatgcacgtg	tttctact				2308

<210> SEQ ID NO 94

<211> LENGTH: 1884

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

-continued

<400> SEQUENCE: 94

agcagaagca	gagcattttc	taatatccac	aaaatgaagg	caataattgt	actactcatg	60
gtagtaacat	ccaatgcaga	tcgaatctgc	actggataa	catcgtaaa	ctcacccat	120
gtggtaaaaa	ctgctactca	aggggaagtc	aacgtgactg	gtgtgatacc	actgacaaca	180
acacctacca	aatctcattt	tgcaaatactc	aaaggaacac	agaccagagg	gaaactatgc	240
ccaaactgtc	tcaactgcac	agatctggac	gtggccttgg	gcagacaaa	gtgtatgggg	300
accatacctt	cggcaaaagc	ttcaataactc	cacgaagtca	aacctgttac	atctgggtgc	360
tttcctataa	tgcacgcac	aacaaaaatc	agacagctac	ccaaatcttct	cagaggatat	420
gaaaatatac	ggttatcagc	ccgtaacgtt	atcaacgcag	aaacggcacc	aggaggaccc	480
tacatagttt	gaacctcagg	atcttgcctt	aacgttacca	atggaaagg	attcttcgca	540
acaatggctt	gggctgtccc	aaaaaacaac	aaaaccaaaa	cagcaacgaa	cccattaaca	600
gtagaagtac	catacatttgc	tacaaaagga	gaagacaaa	ttactgtttg	ggggttccat	660
tctgatgacg	aaacccaaat	ggtaacactc	tatggagact	cgaaggctca	aaagttcacc	720
tcatctgcca	acggagtaac	cacacattat	gtttctcaga	ttgggtggctt	cccaaatacaa	780
acagaagacg	aaggctacc	acaaagcggc	agaattgttgc	ttgattacat	ggtgcaaaaa	840
cctggaaaaa	caggaacaat	tgtctatcaa	agaggtgttt	tattgcctca	aaaagtgtgg	900
tgcgcaagtg	gcaggagcaa	ggtataaaaa	ggggccttgc	ctttaattgg	tgaagcagat	960
tgcctccacg	aaaaatacgg	tggattaaac	aaaagcaagc	cttactacac	aggagaacat	1020
gc当地agcca	taggaatttgc	ccaaatatgg	gtgaaaacac	ccttgaagct	ggccaaatgg	1080
accaaataata	gacctcctgc	aaaactatta	aaggaaaggg	gtttcttgcg	agctattgt	1140
gggttcttgg	aaggaggatg	ggaaggaatg	attgcagggt	ggcacggata	cacatctcat	1200
ggagcacatg	gagtggcagt	ggcagcagac	cttaagagta	cgcaagaagc	tataaacaag	1260
ataacaaaaaa	atctcaatttgc	tttaagttag	ctagaagtaa	agaatcttca	aagactaagc	1320
ggtgcaatgg	atgaaacttca	caacgaaata	ctcgagctgg	atgagaaagt	ggatgatctc	1380
agagctgata	caataagctc	gcaaatagag	cttgcagttc	tgctttccaa	cgaaggaata	1440
ataaaacagtg	aagatgagca	tctcttggca	cttggaaagaa	aactgaagaa	aatgctggc	1500
ccctctgctg	tagacatagg	gaatggatgc	ttcgaaacca	aacacaaatg	caaccagact	1560
tgccttagaca	ggatagctgc	tggcaccttt	aatgcaggag	aattttctct	tccactttt	1620
gattcactaa	atattactgc	tgcatacttta	aatgtatgt	gattggataa	tcatactata	1680
ctgctctact	actcaactgc	tgcttcttagt	ttggctgtaa	cattgtatgt	agctatcttt	1740
attgtttata	tggctccag	agacaatgtt	tcttgctcca	tctgtctata	aggaaaatta	1800
agccctgtat	tttcctttat	tgttagtgctt	gtttgcttgc	caccattaca	aaaaacgtta	1860
ttgaaaaatg	ctcttgttac	tact				1884

<210> SEQ ID NO 95

<211> LENGTH: 1842

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 95

agcagaagca	cagcattttc	ttgtgaactt	caagtaccaa	aaaaaactga	aaatcaaaaat	60
------------	------------	------------	------------	------------	-------------	----

-continued

gtccaacatg gatattgacg gcatcaacac tggacaatt gacaaaacac cagaagaaat	120
aacttccgga accagtgggg caaccagacc aatcatcaa ccagcaaccc ttgccccacc	180
aagcaacaaa cgaacccgaa acccatcccc ggaaaggcga gccacaagca gtgaagctga	240
tgtcggagg agaaccaaa agaaacaaac cccgacagag ataaagaaga gcgtctacaa	300
tatggtagtg aaactgggtg aattctacaa ccagatgatg gtcaaagctg gactcaacga	360
tgacatggag agaaaccta tccaaaatgc acatgctgca gaaagaattc tattggctgc	420
tactgatgac aagaaaactg aattccaaaaa gaaaaagaat gccagagatg tcaaagaagg	480
gaaagaagaa atagaccaca acaaaacagg aggcacccctt tacaagatgg taagagatga	540
taaaaccatc tacttcagcc ctataagaat tacctttta aaagaagagg tgaaaacaat	600
gtacaaaacc accatggggta gtatggttt cagtggacta aatcacatca tgattggca	660
ttcacagatg aacgatgtct gttccaaag atcaaaggca ctaaaaagag ttggacttga	720
cccttcatta atcagttactt ttgcaggaaag cacactcccc agaagatcg gtcaactgg	780
tgttgcgatc aaaggagggtg gaacttttagt ggcagaagcc attcgatttta taggaagagc	840
aatggcagac agagggctat tgagagacat cagagccaag acggcctatg aaaagattct	900
tctgaatctg aaaaacaagt gctctgcgcc ccaacaaaag gctctagttg atcaagtgtat	960
cggaaagtata aatccaggga ttgcagacat agaagaccta accctgcttg cccgaagcat	1020
ggtcgttgctc aggcctctg tagcgagcaa agtggtgctt cccataagca tttatgcca	1080
aataacctaa ctagggttca atggttgaaga atactctatg gttgggtatg aagccatggc	1140
tctttataat atggcaacac ctgtttccat attaagaatg ggagacgtatg caaaagataa	1200
atcacaatta ttcttcatgt ctgcgttgcg agctgcctat gaagaccta gagttttgtc	1260
tgcactaaca ggcacagaat tcaagcatac gtcaagcatat aagtgcagg gtttccacgt	1320
tccagcaaag gagcaagtgg aaggaatggg ggcagctctg atgtccatca agctccagtt	1380
ttgggctcca atgaccagat ctggggggaa tgaagtaggt ggagacggag ggtctggtca	1440
aataagttgc agccccgtgt ttgcagtaga aagacctatt gctctaagca agcaagctgt	1500
aagaagaatg ctgtcaatga atattgaggg acgtgatgca gatgtcaaag gaaatctact	1560
caagatgtatg aatgattcaa tgactaagaa aaccaatgga aatgcattca ttggaaagaa	1620
aatgtttcaa atatcagaca aaaacaaaac caatcccatt gagattccaa ttaagcagac	1680
catccccat ttcttcatttgg ggagggacac agcagaggat tatgtatgacc tcgatttata	1740
aagcaacaaa atagacacta tggctgtac tgtttcagta cgtttggaaat gtgggtgttt	1800
acttttatttgg aaataaatgt aaaaaatgt gttttttctatct	1842

<210> SEQ ID NO 96

<211> LENGTH: 1557

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 96

agcagaagca gagcatcttc tcaaaaactga agcaaatagg caaaaatga acaatgctac	60
cttcaactat acaaacgtta accctatttc tcacatcagg gggagtgtta ttatcactat	120
atgtgtcagc ttcaactgtca tacttattgt attcgatat attgctaaaa ttttcaccaa	180
caaaaataac tgcaccaaca atgtcattgg attgcgcgaa cgtatcaa atgtcaggctg	240

-continued

tgaaccgttc tgcaacaaaa gagatgacat ttcttctccc agagccggag tggacatacc	300
ctcgtttatac ttgccagggc tcaaccttcc agaaagcaact cctaatttgc cctcataggt	360
tcggagaaac cagagggaaac tcagctccct tgataataag ggaacccttt gttgcttgc	420
gaccaaaagga atgcagacac tttgctctaa cccattatgc agctcaacca gggggataact	480
acaatggaaac aagaaaggac agaaacaaggc tgaggcatct gatttcagtc aaattaggca	540
aaatcccaac ttagaaaaac tccattttcc acatggcagc ttggagttgg tccgcattgc	600
atgatggtag agaatggaca tatatcgag ttgatggccc tgacagtaat gcactgtca	660
aaataaaaata tggagaagca tatactgaca cataccatc ctatgcacaa acatctaa	720
gaacacaaga aagtgcctgc aattgcattcg ggggagatgt ttatcttgcataactgtatg	780
gctcagcttc aggaattatgt aaatgcagat ttcttaaaat tcgagagggt cgaataataa	840
aagaaatatt tccaaacagga agagtagagc atactgaaga atgcacatgc gggttcgcca	900
gcaataaaaac catagaatgt gcctgttagatcataacatgttgcacagaaaa agaccctttg	960
tcaaattaaa tgtggagact gatacagctg aaataagatt gatgtgcaca gagacttatt	1020
tggacacccc cagaccagat gatggagca taacagggcc ttgcgaatct aatggggaca	1080
aagggcttgg aggcattcaaa ggaggatttg tccatcaaag aatggcatct aagattggaa	1140
gatggtaactc ccgaacatgc tctaaaactg aaagaatggg gatggactg tatgtcaatgt	1200
atgatggaga cccatggact gacagtgcacg cccttgcattc tagtggagta atggttcaaa	1260
tgaaagaacc tgggtggat tcttttgc tgcataaaaa agataagaaa tgtgatgtcc	1320
cctgtattgg gatagagatg gtacacatgc gtggaaaaga gacttggcac tcagcagcaa	1380
cagccattta ctgtttgtatggc aattgtatgc ggacactgtc acaggtgttgc	1440
atatggctct gtaatggagg aatggttgaa tctgttctaa accctttgtt cctattttgt	1500
ttgaacaattt gtccttactg gacttaattt tttctgaaaa atgctctgt tactact	1557

<210> SEQ ID NO 97
 <211> LENGTH: 1190
 <212> TYPE: DNA
 <213> ORGANISM: Influenza B virus

<400> SEQUENCE: 97

agcagaagca cgcactttct taaaatgtcg ctgtttggag acacaattgc ctacctgctt	60
tcaactaacat aagatggaga aggcaaaagca gaactagcag aaaaattaca ctgttgc	120
gttggaaaag aatttgcact agactctgc ttggatgaa taaaaacaa aagatgccta	180
actgatatac aaaaagcact aattggcgc tctatctgc ttttttttacc caaagaccaa	240
gaaagaaaaaa gaagattcat cacagagccc ctgtcaggaa tggaaacaac agcaacaaaa	300
aagaaaggcc tgattctgc tgagagaaaa atgagaagat gtgtgatgtt tcatgaagca	360
tttggaaatag cagaaggcca taaaatgtcg gactactat attgtctcat ggtcatgtac	420
ctgaaccctgc gaaatttattc aatgcagat aacttagaa cgctctgc tttatgcgag	480
aaacaagcat cacattcaca aagagctcat agcagagcag caagatcttc agtgcctgg	540
gtgaggcag aatgcagat ggttgcgtat gtgaaacacag caaaaacaat gaatggatgt	600
ggaaaggcc aagacgtcca aaaactggca gaagagctgc aaagcaacat tggatgttgc	660
agatctctgg gggcaagtca aaagaatggaa gaaggaatgtt caaaggatgt aatggaaatgt	720

-continued

ctaaaggcaga gctctatggg aaattcagct cttgtgaaga aataccatata atgctcgAAC	780
catttcagat tctttcaatt tggttcttca ttttatcagc tctccatttc atggcttggA	840
caataaggcata ttgttatcaa ataaaaagag gagtaaacct gaaaatacga ataagaaatc	900
caaaataaaga gacaataaac agagaggtat caatTTTgag acacagttac caaaaagaaaa	960
tccaaaggccaa agaaacaatg aaggaagtac tctctgacaa catggagata ttgagtgacc	1020
acatagataat tgaggggctt tctgtgttcaag agataataaa aatgggtgaa acagttttgg	1080
aggtagaaga attgcagtaa acccaatttt caccgtatTTT cttgttatgc atttaagcaa	1140
attgttaatca atgttcgqaa ataaacttqqa aaaatgtcqgt tttttctact	1190

<210> SEQ_ID NO 98
<211> LENGTH: 1098
<212> TYPE: DNA
<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 98

agcagaagca gaggatttgt tttagtcactg gcaaacggaa aaaaatggcg gacaacatga 60
ccacaacaca aattgggatg ggtccggag caaccaatgc caccataaac tttgaagcag 120
gaattctgga gtgctatgaa aggcttcat ggcaaagagc ccttgactac cctggtaag 180
accgcctaaa cagactaaag agaaaattag aatcaagaat aaagactcac aacaaaatg 240
agcctgaag taaaaggatg tctcttgaaag agagaaaagc aattgggta aaaatgtga 300
aagtgcctt atttatgaat ccatactgtg gaattgaagg gtttgagcca tactgtatga 360
aaaattcctc aaatagcaac tgtccaaact gcaattggac cgattaccct ccaacaccag 420
gaaaagtgcct tgatgacata gaagaagaac cggagaatgt tgatgaccca actaaaatag 480
tattgggaa catgaacaac aaagatgca ggc当地aaatgggaa gt当地aaactc 540
agaaagaagg gaagttccgt ttgacaataa aaagggatatacgtatgtg ttgttcttga 600
gagtgttggtaa acggaaaca ttcccaagc accctaatacg atacaagtcc ttataactc 660
tgcatagatt gaatgcatat gaccagatg ggaggcttgt tgctaaactt gttgtactg 720
atgatcttac agtggaggat gaagaagatg gcatcgat cctcaactca ctcttcgagc 780
gttttaatgaa aggacattca aagccaaatc gagcagctgaaactgcgggtggagtttat 840
cccaatttgg tcaagagcac cgattatcac cagaggaggg agacaatttag actggttacg 900
gaagaactt atcttttaag taaaagaatt gatgataaca tattgttcca caaaacagta 960
atagctaaca gctccataat agctgacatg attgtatcat tatcattatttggaaacattt 1020
tatgaaatgaaaggatgtgttggatgaaatgtac agcaggcagt gttgtgaat taaaataaa 1080
aattccctttttaactact 1098

1. A method of rescue of influenza virus, wherein animal cells are electroporated with plasmids that encode an influenza RNA polymerase and nucleoprotein and wherein the electroporated animal cells are co-cultivated with another cell type.
2. The method of claim 1, wherein the animal cells are Vero cells.
3. The method of claim 1, wherein the animal cells are SF Vero cells.
4. The method of claim 1, wherein said another cell type is CEK cells.
5. The method of claim 1, wherein the influenza virus is an influenza A virus.
6. The method of claim 1, wherein the influenza virus is an influenza B virus.
7. The method of claim 1, wherein the influenza virus is a cold adapted virus.

- 8.** The method of claim 1, wherein the influenza virus is an attenuated virus.
- 9.** The method of claim 1, wherein the number of plasmids electroporated is eight.
- 10.** The method of claim 1, wherein the number of plasmids electroporated is twelve.
- 11.** The method of claim 1, wherein the efficiency of said rescue of influenza virus is at least 90%.
- 12.** An influenza virus produced by the method of claim 1.
- 13.** The influenza virus of claim 1, wherein said rescued influenza virus comprises vRNA segments derived from A/PR/8/34.
- 14.** The influenza virus of claim 1, wherein said rescued influenza virus comprises vRNA segments derived from MDV-A.
- 15.** A vaccine composition comprising the influenza virus of claim 1.
- 16.** A method for producing influenza viruses in cell culture, the method comprising:
 - i) introducing a plurality of vectors comprising an influenza virus genome into a population of Vero cells by electroporation;
 - ii) co-cultivating the population of Vero cells with another cell type under conditions permissive for viral replication; and,
 - iii) recovering a plurality of influenza viruses.
- 17.** A method of rescue of influenza virus, wherein (a) animal cells are electroporated with cell expression vectors which direct the expression in said cells of genomic or antigenomic vRNA segments, and a nucleoprotein, and an RNA-dependent polymerase, such that ribonucleoprotein complexes can be formed and viral particles can be assembled; and (b) culturing said cells wherein viral particles are packaged and rescued.
- 18.** The method of claim 17, wherein assembly does not require a helper virus.

19-25. (canceled)

* * * * *