



(86) Date de dépôt PCT/PCT Filing Date: 2012/12/14
(87) Date publication PCT/PCT Publication Date: 2013/06/20
(45) Date de délivrance/Issue Date: 2021/01/19
(85) Entrée phase nationale/National Entry: 2014/05/22
(86) N° demande PCT/PCT Application No.: GB 2012/053140
(87) N° publication PCT/PCT Publication No.: 2013/088164
(30) Priorité/Priority: 2011/12/14 (GB1121513.4)

(51) Cl.Int./Int.Cl. *C07K 16/36* (2006.01)
(72) Inventeurs/Inventors:
HUNTINGTON, JAMES ANDREW, GB;
BAGLIN, TREVOR, GB;
LANGDOWN, JONATHAN, GB
(73) Propriétaire/Owner:
JANSSEN PHARMACEUTICALS, INC., US
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : MOLECULES D'ANTICORPS SE LIANT A LA THROMBINE ET LEURS UTILISATIONS
(54) Title: THROMBIN-BINDING ANTIBODY MOLECULES AND USES THEREOF

(57) **Abrégé/Abstract:**

This invention relates to isolated antibodies which recognise the exosite 1 epitope of thrombin and selectively inhibit thrombin without promoting bleeding. These antibody molecules may be useful in the treatment and prevention of thrombosis, embolism and other conditions mediated by thrombin.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(10) International Publication Number
WO 2013/088164 A1(43) International Publication Date
20 June 2013 (20.06.2013)(51) International Patent Classification:
C07K 16/36 (2006.01)(21) International Application Number:
PCT/GB2012/053140(22) International Filing Date:
14 December 2012 (14.12.2012)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
1121513.4 14 December 2011 (14.12.2011) GB(71) Applicant: CAMBRIDGE ENTERPRISE LIMITED
[GB/GB]; The Old Schools, Trinity Lane, Cambridge,
Cambridgeshire CB2 1TN (GB).(72) Inventors: HUNTINGTON, James Andrew; Department
of Haematology, CIMR, Hills Road, Cambridge, Cam-
bridgeshire CB2 2XY (GB). BAGLIN, Trevor; School of
Clinical Medicine, PO Box 111, Addenbrookes Hospital,
Hills Road, Cambridge Cambridgeshire CB2 0SP (GB).
LANGDOWN, Jonathan; School of Clinical Medicine,
PO Box 111, Addenbrookes Hospital, Hills Road, Cam-
bridge Cambridgeshire CB2 0SP (GB).(74) Agents: ROBERTS, Michael Austin et al.; Reddie &
Grose LLP, 16 Theobalds Road, London WC1X 8PL
(GB).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).**Published:**

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))



WO 2013/088164 A1

(54) Title: THROMBIN-BINDING ANTIBODY MOLECULES AND USES THEREOF

(57) Abstract: This invention relates to isolated antibodies which recognise the exosite 1 epitope of thrombin and selectively inhibit thrombin without promoting bleeding. These antibody molecules may be useful in the treatment and prevention of thrombosis, embolism and other conditions mediated by thrombin.

Thrombin-Binding Antibody Molecules and Uses Thereof

This invention relates to antibody molecules that inhibit thrombin.

5

Blood coagulation is a key process in the prevention of bleeding from damaged blood vessels (haemostasis). However, a blood clot that obstructs the flow of blood through a vessel (thrombosis) or breaks away to lodge in a vessel elsewhere in the body

10 (thromboembolism) can be a serious health threat.

A number of anticoagulant therapies are available to treat pathological blood coagulation. A common drawback of these therapies is an increased risk of bleeding (Mackman (2008) *Nature* 15 451(7181): 914-918). Many anticoagulant agents have a narrow therapeutic window between the dose that prevents thrombosis and the dose that induces bleeding. This window is often further restricted by variations in the response in individual patients.

20 The present invention relates to the unexpected finding that antibody molecules which recognise the exosite 1 epitope of thrombin selectively inhibit thrombin without promoting bleeding. These antibody molecules may be useful in the treatment and prevention of thrombosis, embolism and other conditions mediated 25 by thrombin.

An aspect of the invention provides an isolated antibody molecule that specifically binds to exosite 1 of thrombin.

30 Isolated anti-exosite 1 antibody molecules may inhibit thrombin *in vivo* without promoting or substantially promoting bleeding or haemorrhage, i.e. the antibody molecules do not inhibit or substantially inhibit normal physiological responses to vascular injury (i.e. haemostasis). For example, haemostasis may not be 35 inhibited or may be minimally inhibited by the antibody molecules (i.e. inhibited to an insignificant extent which does not affect the well-being of patient or require further intervention). Bleeding may not be increased or may be minimally increased by the antibody molecules.

40

Exosite 1 (also known as 'anion binding exosite 1' and the 'fibrinogen recognition exosite') is a well-characterised secondary binding site on the thrombin molecule (see for example James A. Huntington, 2008, *Structural Insights into the Life*

History of Thrombin, in *Recent Advances in Thrombosis and Hemostasis 2008*, editors; K. Tanaka and E.W. Davie, Springer Japan KK, Tokyo, pp. 80-106) . Exosite 1 is formed in mature thrombin but is not formed in prothrombin (see for example Anderson et al (2000) JBC 2775 16428-16434).

Exosite 1 is involved in recognising thrombin substrates, such as fibrinogen, but is remote from the catalytic active site. Various thrombin binding factors bind to exosite 1, including the anticoagulant dodecapeptide hirugen (Naski et al 1990 JBC 265 13484-13489), factor V, factor VIII, thrombomodulin (cofactor for protein C and TAFI activation) , fibrinogen, PARI and fibrin (the co-factor for factor XIII activation).

Summary

In one aspect of the invention, it is provided an isolated antibody molecule that specifically binds to the exosite 1 region of thrombin, wherein the antibody molecule comprises: a VH domain comprising an HCDR1, HCDR2 and HCDR3 having the sequences of SEQ ID NOs 3, 4 and 5, respectively; and a VL domain comprising an LCDR1, LCDR2 and LCDR3 having either the sequences of SEQ ID NOs 7, 8 and 9, respectively, or an LCDR1 having the sequence of SEQ ID NO: 7 with a glycosylation site mutated out with a substitution at an amino acid residue corresponding to N28 or S30 in SEQ ID NO: 6.

This summary of the invention does not necessarily describe all features of the invention.

Detailed Description

An anti-exosite 1 antibody may bind to exosite 1 of mature human thrombin. The sequence of human prothrombin is set out in SEQ ID NO: 1. Human prothrombin has the sequence of residues 44 to 622 of SEQ ID NO: 1. Mature human thrombin has the sequence of residues 314-363 (light chain) and residues 364 to 622 (heavy chain).

In some embodiments, an anti-exosite 1 antibody may also bind to exosite 1 of mature thrombin from other species. Thrombin sequences from other species are known in the art and available on public databases such as Genbank . The corresponding residues in thrombin sequences from other species may be easily identified using sequence alignment tools.

The numbering scheme for thrombin residues set out herein is conventional in the art and is based on the chymotrypsin template (Bode W et al EMBO J. 1989 Nov; 8 (11): 3467-75) . Thrombin has insertion loops relative to chymotrypsin that are lettered sequentially using lower case letters.

Exosite 1 of mature human thrombin is underlined in SEQ ID NO: 1 and may include the following residues: M32, F34, R35, K36, S36a, P37, Q38, E39, L40, L65, R67, S72, R73, T74, R75, Y76, R77a, N78, E80, K81, 182, S83, M84, K109, K110, K149e, G150, Q151, S153 and V154. In some embodiments, other thrombin residues which are located close to (i.e. within 0.5 nm or within 1 nm) of any one of these residues may also be considered to be part of exosite 1.

An anti-exosite 1 antibody may bind to an epitope which comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more than 20 residues of exosite 1. Preferably, an anti exosite 1 antibody binds to an epitope which consists
5 entirely of exosite 1 residues.

For example, an anti-exosite 1 antibody may bind to an epitope which comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or all 16 residues selected from the group consisting of M32,
10 F34, S36a, P37, Q38, E39, L40, L65, R67, R73, T74, R75, Y76, R77a, I82 and Q151 of human thrombin or the equivalent residues in thrombin from another species. In some preferred embodiments, the epitope may comprise the thrombin residues Q38, R73, T74, Y76 and R77a and optionally one or more additional residues.

15 Anti-exosite 1 antibody molecules as described herein are specific for thrombin exosite 1 and bind to this epitope with high affinity relative to other epitopes, for example epitopes from mammalian proteins other than mature thrombin. For example,
20 an anti-exosite 1 antibody molecule may display a binding affinity for thrombin exosite 1 which is at least 500 fold, at least 1000 fold or at least 2000 fold greater than other epitopes.

25 Preferably, an antibody molecule as described herein which is specific for exosite 1 may bind to mature thrombin but display no binding or substantially no binding to prothrombin.

Without being bound by any theory, anti-exosite 1 antibodies may
30 be unable to access thrombin within the core of a haemostatic clot, and are therefore unable to affect haemostasis by interrupting normal thrombin function at sites of vascular injury. However, because the anti-exosite 1 antibodies still bind to thrombin on the surface of the clot and in the outer shell of
35 the clot, thrombosis is prevented, i.e. non-haemostatic clot extension is prevented.

An anti-exosite 1 antibody molecule may have a dissociation constant for exosite 1 of less than 50nM, less than 40nM, less
40 than 30nM, less than 20nM, less than 10nM, or less than 1nM. For example, an antibody molecule may have an affinity for exosite 1 of 0.1 to 50 nM, e.g. 0.5 to 10 nM. A suitable anti-exosite 1

antibody molecule may, for example, have an affinity for thrombin exosite 1 of about 1 nM.

5 Binding kinetics and affinity (expressed as the equilibrium dissociation constant, K_d) of the anti-exosite 1 antibody molecules may be determined using standard techniques, such as surface plasmon resonance e.g. using BIAcore analysis.

10 An anti-exosite 1 antibody molecule as described herein may be an immunoglobulin or fragment thereof, and may be natural or partly or wholly synthetically produced, for example a recombinant molecule.

15 Anti-exosite 1 antibody molecules may include any polypeptide or protein comprising an antibody antigen-binding site, including Fab, Fab₂, Fab₃, diabodies, triabodies, tetrabodies, minibodies and single-domain antibodies, including nanobodies, as well as whole antibodies of any isotype or sub-class. Antibody molecules and methods for their construction and use are described, in for example Holliger & Hudson, Nature Biotechnology 23(9):1126-1136
20 (2005).

In some preferred embodiments, the anti-exosite 1 antibody molecule may be a whole antibody. For example, the anti-exosite
25 1 antibody molecule may be an IgG, IgA, IgE or IgM or any of the isotype sub-classes, particularly IgG1 and IgG4. The anti-exosite 1 antibody molecules may be monoclonal antibodies. In other preferred embodiments, the anti-exosite 1 antibody molecule may be an antibody fragment.

30 Anti-exosite 1 antibody molecules may be chimeric, humanised or human antibodies.

35 Anti-exosite 1 antibody molecules as described herein may be isolated, in the sense of being free from contaminants, such as antibodies able to bind other polypeptides and/or serum components. Monoclonal antibodies are preferred for some purposes, though polyclonal antibodies may also be employed.

40 Anti-exosite 1 antibody molecules may be obtained using techniques which are standard in the art. Methods of producing antibodies include immunising a mammal (e.g. mouse, rat, rabbit, horse, goat, sheep or monkey) with the protein or a fragment

thereof. Antibodies may be obtained from immunised animals using any of a variety of techniques known in the art, and screened, preferably using binding of antibody to antigen of interest. For instance, Western blotting techniques or immunoprecipitation may
5 be used (Armitage et al., 1992, Nature 357: 80-82). Isolation of antibodies and/or antibody-producing cells from an animal may be accompanied by a step of sacrificing the animal.

As an alternative or supplement to immunising a mammal with a
10 peptide, an antibody specific for a protein may be obtained from a recombinantly produced library of expressed immunoglobulin variable domains, e.g. using lambda bacteriophage or filamentous bacteriophage which display functional immunoglobulin binding domains on their surfaces; for instance see WO92/01047. The
15 library may be naive, that is constructed from sequences obtained from an organism which has not been immunised with any of the proteins (or fragments), or may be one constructed using sequences obtained from an organism which has been exposed to the antigen of interest.

20

Other anti-exosite 1 antibody molecules may be identified by screening patient serum for antibodies which bind to exosite 1.

In some embodiments, anti-thrombin antibody molecules may be
25 produced by any convenient means, for example a method described above, and then screened for differential binding to mature thrombin relative to thrombin with an exosite 1 mutation, gamma thrombin (exosite 1 defective due to autolysis at R75 and R77a) or prothrombin. Suitable screening methods are well-known in the
30 art.

An antibody which displays increased binding to mature thrombin, relative to non-thrombin proteins, thrombin with an exosite 1 mutation, gamma-thrombin or prothrombin, for example an antibody
35 which binds to mature thrombin but does not bind to thrombin with an exosite I mutation, gamma thrombin or prothrombin, may be identified as an anti-exosite 1 antibody molecule.

After production and/or isolation, the biological activity of an
40 anti-exosite 1 antibody molecule may be tested. For example, the ability of the antibody molecule to inhibit thrombin substrate, cofactor or inhibitor binding and/or cleavage by thrombin may be

determined and/or the ability of the antibody molecule to inhibit thrombosis without promoting bleeding may be determined.

5 Suitable antibody molecules may be tested for activity using a fibrinogen clotting or thrombin time assay. Suitable assays are well-known in the art.

10 The effect of an antibody molecule on coagulation and bleeding may be determined using standard techniques. For example, the effect of an antibody molecule on thrombosis may be determined in an animal model, such as a mouse model with ferric chloride induced clots in blood vessels. Effects on haemostasis may also be determined in an animal model, for example, by measuring tail bleed of a mouse.

15 Antibody molecules normally comprise an antigen binding domain comprising an immunoglobulin heavy chain variable domain (VH) and an immunoglobulin light chain variable domain (VL), although antigen binding domains comprising only a heavy chain variable domain (VH) are also possible (e.g. camelid or shark antibodies).

20 Each of the VH and VL domains typically comprise three complementarity determining regions (CDRs) responsible for antigen binding, interspersed by framework regions.

25 In some embodiments, binding to exosite 1 may occur wholly or substantially through the VHCDR3 of the anti-exosite 1 antibody molecule.

30 For example, an anti-exosite 1 antibody molecule may comprise a VH domain comprising a HCDR3 having the amino acid sequence of SEQ ID NO: 5 or the sequence of SEQ ID NO: 5 with 1 or more, for example 2, 3, 4 or 5 or more amino acid substitutions, deletions or insertions. The substitutions may be conservative
35 substitutions. In some embodiments, the HCDR3 may comprise the amino acid residues at positions 4 to 9 of SEQ ID NO: 5 (SEFEPF), or more preferably the amino acid residues at positions 2, and 4 to 10 of SEQ ID NO: 5 (D and SEFEPFS) with substitutions,
40 deletions or insertions at one or more other positions in SEQ ID NO:5. The HCDR3 may be the only region of the antibody molecule that interacts with a thrombin exosite 1 epitope or substantially the only region. The HCDR3 may therefore determine the

specificity and/or affinity of the antibody molecule for the exosite 1 region of thrombin.

The VH domain of an anti exosite 1 antibody molecule may additionally comprise an HCDR2 having the amino acid sequence of SEQ ID NO: 4 or the sequence of SEQ ID NO: 4 with 1 or more, for example 2, 3, 4 or 5 or more amino acid substitutions, deletions or insertions. In some embodiments, the HCDR2 may comprise the amino acid residues at positions 3 to 7 of SEQ ID NO: 4 (DPQDG) or the amino acid residues at positions 2 and 4 to 7 of SEQ ID NO: 4 (L and PQDG) of SEQ ID NO: 4, with substitutions, deletions or insertions at one or more other positions in SEQ ID NO: 4.

The VH domain of an anti-exosite 1 antibody molecule may further comprise an HCDR1 having the amino acid sequence of SEQ ID NO: 3 or the sequence of SEQ ID NO: 3 with 1 or more, for example 2, 3, 4 or 5 or more amino acid substitutions, deletions or insertions. In some embodiments, the HCDR1 may comprise amino acid residue T at position 5 of SEQ ID NO: 3 with substitutions, deletions or insertions at one or more other positions in SEQ ID NO: 3.

In some embodiments, an antibody molecule may comprise a VH domain comprising a HCDR1, a HCDR2 and a HCDR3 having the sequences of SEQ ID NOs 3, 4 and 5 respectively. For example, an antibody molecule may comprise a VH domain having the sequence of SEQ ID NO: 2 or the sequence of SEQ ID NO: 2 with 1 or more, for example 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acid substitutions, deletions or insertions in SEQ ID NO: 2.

The anti-exosite 1 antibody molecule may further comprise a VL domain, for example a VL domain comprising LCDR1, LCDR2 and LCDR3 having the sequences of SEQ ID NOs 7, 8 and 9 respectively, or the sequences of SEQ ID NOs 7, 8 and 9 respectively with, independently, 1 or more, for example 2, 3, 4 or 5 or more amino acid substitutions, deletions or insertions. The substitutions may be conservative substitutions. For example, an antibody molecule may comprise a VL domain having the sequence of SEQ ID NO: 6 or the sequence of SEQ ID NO: 6 with 1 or more, for example 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acid substitutions, deletions or insertions in SEQ ID NO: 6.

In some embodiments, the VL domain may comprise Tyr49.

The anti-exosite 1 antibody molecule may for example comprise one or more amino acid substitutions, deletions or insertions which improve one or more properties of the antibody, for example affinity, functional half life, on and off rates.

5

The techniques that are required in order to introduce substitutions, deletions or insertions within amino acid sequences of CDRs, antibody VH or VL domains and antibodies are generally available in the art. Variant sequences may be made, with substitutions, deletions or insertions that may or may not be predicted to have a minimal or beneficial effect on activity, and tested for ability to bind exosite 1 of thrombin and/or for any other desired property.

15 In some embodiments, anti-exosite 1 antibody molecule may comprise a VH domain comprising a HCDR1, a HCDR2 and a HCDR3 having the sequences of SEQ ID NOs 3, 4, and 5, respectively, and a VL domain comprising a LCDR1, a LCDR2 and a LCDR3 having the sequences of SEQ ID NOs 7, 8 and 9, respectively.

20

For example, the VH and VL domains may have the amino acid sequences of SEQ ID NO: 2 and SEQ ID NO: 6 respectively; or may have the amino acid sequences of SEQ ID NO: 2 and SEQ ID NO: 6 comprising, independently 1 or more, for example 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acid substitutions, deletions or insertions. The substitutions may be conservative substitutions.

25

In some embodiments, an antibody may comprise one or more substitutions, deletions or insertions which remove a glycosylation site. For example, a glycosylation site in VL domain of SEQ ID NO 6 may be mutated out by introducing a substitution at either N28 or S30.

30

The anti-exosite 1 antibody molecule may be in any format, as described above, In some preferred embodiments, the anti-exosite 1 antibody molecule may be a whole antibody, for example an IgG, such as IgG1 or IgG4, IgA, IgE or IgM.

35

An anti-exosite 1 antibody molecule of the invention may be one which competes for binding to exosite 1 with an antibody molecule described above, for example an antibody molecule which

40

(i) binds thrombin exosite 1 and

(ii) comprises a VH domain of SEQ ID NO: 2 and/or VL domain of SEQ ID NO: 6; an HCDR3 of SEQ ID NO: 5; an HCDR1, HCDR2, LCDR1, LCDR2, or LCDR3 of SEQ ID NOS: 3, 4, 7, 8 or 9 respectively; a VH domain comprising HCDR1, HCDR2 and HCDR3 sequences of SEQ ID NOS: 3, 4 and 5 respectively; and/or a VH domain comprising HCDR1, HCDR2 and HCDR3 sequences of SEQ ID NOS: 3, 4 and 5 and a VL domain comprising LCDR1, LDR2 and LCDR3 sequences of SEQ ID NOS: 7, 8 and 9 respectively.

10 Competition between antibody molecules may be assayed easily *in vitro*, for example using ELISA and/or by tagging a specific reporter molecule to one antibody molecule which can be detected in the presence of one or more other untagged antibody molecules, to enable identification of antibody molecules which bind the same epitope or an overlapping epitope. Such methods are readily known to one of ordinary skill in the art. Thus, a further aspect of the present invention provides an antibody molecule comprising a antibody antigen-binding site that competes with an antibody molecule, for example an antibody molecule comprising a VH and/or VL domain, CDR e.g. HCDR3 or set of CDRs of the parent antibody described above for binding to exosite 1 of thrombin. A suitable antibody molecule may comprise an antibody antigen-binding site which competes with an antibody antigen-binding site for binding to exosite 1 wherein the antibody antigen-binding site is composed of a VH domain and a VL domain, and wherein the VH and VL domains comprise HCDR1, HCDR2 and HCDR3 sequences of SEQ ID NOS: 3, 4, and 5 and LCDR1, LDR2 and LCDR3 sequences of SEQ ID NOS: 7, 8, and 9 respectively, for example the VH and VL domains of SEQ ID NOS: 2 and 6.

30 An anti-exosite 1 antibody molecule as described herein may inhibit the binding of thrombin-binding factors, including factors which bind to exosite 1. For example, an antibody molecule may competitively or non-competitively inhibit the binding of one or more of fV, FVIII, thrombomodulin, fibrinogen or fibrin, PAR1 and/or hirugen and hirudin analogues to thrombin.

40 An anti-exosite 1 antibody molecule as described herein may inhibit one or more activities of thrombin. For example, an anti-exosite 1 antibody molecule may inhibit the hydrolytic cleavage of one or more thrombin substrates, such as fibrinogen, platelet receptor PAR-1 and coagulation factor FVIII. For example, binding of the antibody molecule to thrombin may result

in an at least 5-fold, at least 10-fold, or at least 15-fold decrease in the hydrolysis of fibrinogen, PAR-1, coagulation factor FVIII and/or another thrombin substrates, such as factor V, factor XIII in the presence of fibrin, and protein C and/or
5 TAFI in the presence of thrombomodulin. In some embodiments, binding of thrombin by the anti-exosite 1 antibody molecule may result in no detectable cleavage of the thrombin substrate by thrombin.

10 Techniques for measuring thrombin activity, for example by measuring the hydrolysis of thrombin substrates *in vitro* are standard in the art and are described herein.

Anti-exosite 1 antibody molecules may be further modified by
15 chemical modification, for example by PEGylation, or by incorporation in a liposome, to improve their pharmaceutical properties, for example by increasing *in vivo* half-life.

The effect of an anti-exosite 1 antibody molecule on coagulation
20 and bleeding may be determined using standard techniques. For example, the effect of an antibody on a thrombosis model may be determined. Suitable models include ferric chloride clot induction in blood vessels in a murine model, followed by a tail bleed to test normal haemostasis. Other suitable thrombosis
25 models are well known in the art (see for example Westrick et al ATVB (2007) 27:2079-2093)

Anti-exosite 1 antibody molecules may be comprised in
30 pharmaceutical compositions with a pharmaceutically acceptable excipient.

A pharmaceutically acceptable excipient may be a compound or a combination of compounds entering into a pharmaceutical composition which does not provoke secondary reactions and which
35 allows, for example, facilitation of the administration of the anti-exosite 1 antibody molecule, an increase in its lifespan and/or in its efficacy in the body or an increase in its solubility in solution. These pharmaceutically acceptable vehicles are well known and will be adapted by the person skilled
40 in the art as a function of the mode of administration of the anti-exosite 1 antibody molecule.

In some embodiments, anti-exosite 1 antibody molecules may be provided in a lyophilised form for reconstitution prior to administration. For example, lyophilised antibody molecules may be reconstituted in sterile water and mixed with saline prior to administration to an individual.

Anti-exosite 1 antibody molecules will usually be administered in the form of a pharmaceutical composition, which may comprise at least one component in addition to the antibody molecule. Thus pharmaceutical compositions may comprise, in addition to the anti-exosite 1 antibody molecule, a pharmaceutically acceptable excipient, carrier, buffer, stabilizer or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the anti-exosite 1 antibody molecule. The precise nature of the carrier or other material will depend on the route of administration, which may be by bolus, infusion, injection or any other suitable route, as discussed below.

For parenteral, for example sub-cutaneous or intra-venous administration, e.g. by injection, the pharmaceutical composition comprising the anti-exosite 1 antibody molecule may be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles, such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection. Preservatives, stabilizers, buffers, antioxidants and/or other additives may be employed as required including buffers such as phosphate, citrate and other organic acids; antioxidants, such as ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride; benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens, such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3'-pentanol; and m-cresol); low molecular weight polypeptides; proteins, such as serum albumin, gelatin or immunoglobulins; hydrophilic polymers, such as polyvinylpyrrolidone; amino acids, such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides and other carbohydrates including glucose, mannose or dextrans; chelating agents, such as EDTA; sugars, such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions, such as sodium; metal complexes (e.g. Zn-

protein complexes); and/or non-ionic surfactants, such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

5 A pharmaceutical composition comprising an anti exosite 1 antibody molecule may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated.

10 An anti-exosite 1 antibody molecule as described herein may be used in a method of treatment of the human or animal body, including prophylactic or preventative treatment (e.g. treatment before the onset of a condition in an individual to reduce the risk of the condition occurring in the individual; delay its onset; or reduce its severity after onset). The method of
15 treatment may comprise administering an anti-exosite 1 antibody molecule to an individual in need thereof.

Administration is normally in a "therapeutically effective amount", this being sufficient to show benefit to a patient.
20 Such benefit may be at least amelioration of at least one symptom. The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of
25 the disorder, the site of delivery of the composition, the method of administration, the scheduling of administration and other factors known to medical practitioners. Prescription of treatment, e.g. decisions on dosage etc, is within the responsibility of general practitioners and other medical doctors
30 and may depend on the severity of the symptoms and/or progression of a disease being treated. Appropriate doses of antibody molecules are well known in the art (Ledermann J.A. et al. (1991) Int. J. Cancer 47: 659-664; Bagshawe K.D. et al. (1991) Antibody, Immunoconjugates and Radiopharmaceuticals 4: 915-922). Specific
35 dosages may be indicated herein or in the Physician's Desk Reference (2003) as appropriate for the type of medicament being administered may be used. A therapeutically effective amount or suitable dose of an antibody molecule may be determined by comparing its *in vitro* activity and *in vivo* activity in an animal
40 model. Methods for extrapolation of effective dosages in mice and other test animals to humans are known. The precise dose will depend upon a number of factors, including whether the antibody is for prevention or for treatment, the size and

location of the area to be treated, the precise nature of the antibody (e.g. whole antibody, fragment) and the nature of any detectable label or other molecule attached to the antibody.

- 5 A typical antibody dose will be in the range 100 µg to 1 g for systemic applications, and 1 µg to 1 mg for topical applications. An initial higher loading dose, followed by one or more lower doses, may be administered. Typically, the antibody will be a whole antibody, e.g. the IgG1 or IgG4 isotype. This is a dose
- 10 for a single treatment of an adult patient, which may be proportionally adjusted for children and infants, and also adjusted for other antibody formats in proportion to molecular weight. Treatments may be repeated at daily, twice-weekly, weekly or monthly intervals, at the discretion of the physician.
- 15 The treatment schedule for an individual may be dependent on the pharmacokinetic and pharmacodynamic properties of the antibody composition, the route of administration and the nature of the condition being treated.
- 20 Treatment may be periodic, and the period between administrations may be about two weeks or more, e.g. about three weeks or more, about four weeks or more, about once a month or more, about five weeks or more, or about six weeks or more. For example, treatment may be every two to four weeks or every four to eight weeks.
- 25 Treatment may be given before, and/or after surgery, and/or may be administered or applied directly at the anatomical site of surgical treatment or invasive procedure. Suitable formulations and routes of administration are described above.
- 30 In some embodiments, anti-exosite 1 antibody molecules as described herein may be administered as sub-cutaneous injections. Sub-cutaneous injections may be administered using an auto-injector, for example for long term prophylaxis/treatment.
- 35 In some preferred embodiments, the therapeutic effect of the anti-exosite 1 antibody molecule may persist for several half-lives, depending on the dose. For example, the therapeutic effect of a single dose of anti-exosite 1 antibody molecule may persist in an individual for 1 month or more, 2 months or more, 3 months
- 40 or more, 4 months or more, 5 months or more, or 6 months or more.

Anti-exosite 1 antibody molecules described herein inhibit thrombin and may be useful in the treatment of thrombin-mediated conditions.

5 Haemostasis is the normal coagulation response i.e. the prevention of bleeding or haemorrhage, for example from a damaged blood vessel. Haemostasis arrests bleeding and haemorrhage from blood vessels in the body.

10 Anti-exosite 1 antibody molecules may have no effect or substantially no effect on haemostasis i.e. they do not promote bleeding or haemorrhage.

15 Aspects of the invention provide; an anti-exosite 1 antibody molecule as described herein for use in a method of treatment of the human or animal body; an anti-exosite 1 antibody molecule as described herein for use in a method of treatment of a thrombin-mediated disorder; the use of an anti-exosite 1 antibody molecule as described herein in the manufacture of a medicament for the
20 treatment of a thrombin-mediated condition; and a method of treatment of a thrombin-mediated condition comprising administering an anti-exosite 1 antibody molecule as described herein to an individual in need thereof.

25 Inhibition of thrombin by anti-exosite 1 antibodies as described herein may be of clinical benefit in the treatment of any thrombin-mediated condition. A thrombin-mediated condition may include disorders associated with the formation or activity of thrombin.

30 Thrombin plays a key role in haemostasis, coagulation and thrombosis. Thrombin-mediated conditions include thrombotic conditions, such as thrombosis and embolism.

35 Thrombosis is coagulation which is in excess of what is required for haemostasis (i.e. excessive coagulation), or which is not required for haemostasis (i.e. extra-haemostatic or non-haemostatic coagulation).

40 Thrombosis is blood clotting within the blood vessel lumen. It is characterised by the formation of a clot (thrombus) that is in excess of requirement or not required for haemostasis. The clot may impede blood flow through the blood vessel leading to medical

complications. A clot may break away from its site of formation, leading to embolism elsewhere in the circulatory system. In the arterial system, thrombosis is typically the result of atherosclerotic plaque rupture.

5

In some embodiments, thrombosis may occur after an initial physiological haemostatic response, for example damage to endothelial cells in a blood vessel. In other embodiments, thrombosis may occur in the absence of any physiological haemostatic response.

10

Thrombosis may occur in individuals with an intrinsic tendency to thrombosis (i.e. thrombophilia) or in 'normal' individuals with no intrinsic tendency to thrombosis, for example in response to an extrinsic stimulus.

15

Thrombosis and embolism may occur in any vein, artery or other blood vessel within the circulatory system and may include microvascular thrombosis.

20

Thrombosis and embolism may be associated with surgery (either during surgery or afterwards) or the insertion of foreign objects, such as coronary stents, into a patient.

25

For example, anti-exosite 1 antibodies as described herein may be useful in the surgical and other procedures in which blood is exposed to artificial surfaces, such as open heart surgery and dialysis.

30

Thrombotic conditions may include thrombophilia, thrombotic stroke and coronary artery occlusion.

Patients suitable for treatment as described herein include patients with conditions in which thrombosis is a symptom or a side-effect of treatment or which confer an increased risk of thrombosis or patients who are predisposed to or at increased risk of thrombosis, relative to the general population. For example, an anti-exosite 1 antibody molecule as described herein may also be useful in the treatment or prevention of venous thrombosis in cancer patients, and in the treatment or prevention of hospital-acquired thrombosis, which is responsible for 50% of cases of venous thromboembolism.

40

Anti-exosite 1 antibody molecules as described herein may exert a therapeutic or other beneficial effect on thrombin-mediated conditions, such as thrombotic conditions, without substantially inhibiting or impeding haemostasis. For example, the risk of haemorrhage in patients treated with anti-exosite 1 antibody molecules may not be increased or substantially increased relative to untreated individuals.

Individuals treated with conventional anticoagulants, such as natural and synthetic heparins, warfarin, direct serine protease inhibitors (e.g. argatroban, dabigatran, apixaban, and rivaroxaban), hirudin and its derivatives (e.g. lepirudin and bivalirudin), and anti-platelet drugs (e.g. clopidogrel, ticlopidine and abciximab) cause bleeding. The risk of bleeding in patients treated with anti-exosite 1 antibody molecules as described herein may be reduced relative to individuals treated with conventional anticoagulants.

Thrombin-mediated conditions include non-thrombotic conditions associated with thrombin activity, including inflammation, infection, tumour growth and metastasis, organ rejection and dementia (vascular and non-vascular, e.g. Alzheimer's disease) (Licari et al J Vet Emerg Crit Care (San Antonio). 2009 Feb;19(1):11-22; Tsopanoglou et al Eur Cytokine Netw. 2009 Dec 1;20(4):171-9).

Anti-exosite 1 antibody molecules as described herein may also be useful in *in vitro* testing, for example in the analysis and characterisation of coagulation, for example in a sample obtained from a patient.

Anti-exosite 1 antibody molecules may be useful in the measurement of thrombin generation. Assays of thrombin generation are technically problematic because the conversion of fibrinogen to fibrin causes turbidity, which precludes the use of a simple chromogenic end-point.

The addition of an anti-exosite 1 antibody molecule as described herein to a sample of blood prevents or inhibits fibrin formation and hence turbidity and permits thrombin generation to be measured using a chromogenic substrate, without the need for a defibrination step.

For example, a method of measuring thrombin generation may comprise contacting a blood sample with a chromogenic thrombin substrate in the presence of an anti-exosite 1 antibody molecule as described herein and measuring the chromogenic signal from the
5 substrate;

wherein the chromogenic signal is indicative of thrombin generation in the sample.

The chromogenic signal may be measured directly without
10 defibrination of the sample.

Suitable substrates are well known in the art and include S2238 (H-D-Phe-Pip-Arg-pNa), β -Ala-Gly-Arg-p-nitroanilide diacetate (Prasa, D. et al. (1997) *Thromb. Haemost.* 78, 1215; Sigma Aldrich
15 Inc) and Tos-Gly-Pro-Arg-pNa.ACOH (Biophen CS-01(81); Aniara Inc OH USA).

Anti-exosite 1 antibody molecules may also be useful in inhibiting or preventing the coagulation of blood as described
20 above in extracorporeal circulations, such as haemodialysis and extracorporeal membrane oxygenation.

For example, a method of inhibiting or preventing blood coagulation *in vitro* or *ex vivo* may comprise introducing an anti-exosite 1 antibody molecule as described herein to a blood
25 sample. The blood sample may be introduced into an extracorporeal circulation system before, simultaneous with or after the introduction of the anti-exosite 1 antibody and optionally subjected to treatment such as haemodialysis or oxygenation. In
30 some embodiments, the treated blood may be subsequently administered to an individual. Other embodiments provide an anti-exosite 1 antibody molecule as described herein for use in a method of inhibiting or preventing blood coagulation in a blood sample *ex vivo* and the use of an anti-exosite 1 antibody molecule
35 as described herein in the manufacture of a medicament for use in a method of inhibiting or preventing blood coagulation in a blood sample *ex vivo*.

Other aspects of the invention relate to the production of
40 antibody molecules which bind to the exosite 1 epitope of thrombin and may be useful, for example in the treatment of pathological blood coagulation or thrombosis.

A method for producing an antibody antigen-binding domain for the exosite 1 epitope of thrombin, may comprise;

providing, by way of addition, deletion, substitution or insertion of one or more amino acids in the amino acid sequence of a parent VH domain comprising HCDR1, HCDR2 and HCDR3, wherein HCDR1, HCDR2 and HCDR3 have the amino acid sequences of SEQ ID NOS: 3, 4 and 5 respectively, a VH domain which is an amino acid sequence variant of the parent VH domain, and;

optionally combining the VH domain thus provided with one or more VL domains to provide one or more VH/VL combinations; and

testing said VH domain which is an amino acid sequence variant of the parent VH domain or the VH/VL combination or combinations to identify an antibody antigen binding domain for the exosite 1 epitope of thrombin.

A VH domain which is an amino acid sequence variant of the parent VH domain may have the HCDR3 sequence of SEQ ID NO: 5 or a variant with the addition, deletion, substitution or insertion of one, two, three or more amino acids.

The VH domain which is an amino acid sequence variant of the parent VH domain may have the HCDR1 and HCDR2 sequences of SEQ ID NOS: 3 and 4 respectively, or variants of these sequences with the addition, deletion, substitution or insertion of one, two, three or more amino acids.

A method for producing an antibody molecule that specifically binds to the exosite 1 epitope of thrombin may comprise:

providing starting nucleic acid encoding a VH domain or a starting repertoire of nucleic acids each encoding a VH domain, wherein the VH domain or VH domains either comprise a HCDR1, HCDR2 and/or HCDR3 to be replaced or lack a HCDR1, HCDR2 and/or HCDR3 encoding region;

combining said starting nucleic acid or starting repertoire with donor nucleic acid or donor nucleic acids encoding or produced by mutation of the amino acid sequence of an HCDR1, HCDR2, and/or HCDR3 having the amino acid sequences of SEQ ID NOS: 3, 4 and 5 respectively, such that said donor nucleic acid is or donor nucleic acids are inserted into the CDR1, CDR2 and/or CDR3 region in the starting nucleic acid or starting repertoire, so as to provide a product repertoire of nucleic acids encoding VH domains;

expressing the nucleic acids of said product repertoire to produce product VH domains;

optionally combining said product VH domains with one or more VL domains;

5 selecting an antibody molecule that binds exosite 1 of thrombin, which antibody molecule comprises a product VH domain and optionally a VL domain; and

recovering said antibody molecule or nucleic acid encoding it.

10

Suitable techniques for the maturation and optimisation of antibody molecules are well-known in the art.

15 Antibody antigen-binding domains and antibody molecules for the exosite 1 epitope of thrombin may be tested as described above. For example, the ability to bind to thrombin and/or inhibit the cleavage of thrombin substrates may be determined.

20 The effect of an antibody molecule on coagulation and bleeding may be determined using standard techniques. For example, a mouse thrombosis model of ferric chloride clot induction in a blood vessel, such as the femoral vein or carotid artery, followed by a tail bleed to test normal haemostasis, may be employed.

25 Various further aspects and embodiments of the present invention will be apparent to those skilled in the art in view of the present disclosure.

30 All documents mentioned in this specification are incorporated herein by reference in their entirety.

Unless stated otherwise, antibody residues are numbered herein in accordance with the Kabat numbering scheme.

35 "and/or" where used herein is to be taken as specific disclosure of each of the two specified features or components with or without the other. For example "A and/or B" is to be taken as specific disclosure of each of (i) A, (ii) B and (iii) A and B, just as if each is set out individually herein.

40

Unless context dictates otherwise, the descriptions and definitions of the features set out above are not limited to any particular aspect or embodiment of the invention and apply

equally to all aspects and embodiments which are described. Thus, the features set out above are disclosed in all combinations and permutations.

5 Certain aspects and embodiments of the invention will now be illustrated by way of example and with reference to the figures and tables described below.

10 Figure 1 shows the binding and elution of the IgA on human thrombin-Sepharose column. Figure 1A shows an elution profile for IgA (narrow peak) from a thrombin-Sepharose column using a pH gradient (neutral to low, indicated by upward sloping line). Figure 1B shows a native blue gel showing total IgA load, flow-through from the human thrombin column and eluate following
15 elution at low pH.

Figure 2 shows a non-reducing SDS-PAGE gel which indicates that the IgA binds thrombin but not prothrombin. In this pull-down assay, lectin agarose is used to bind to IgA in the presence of
20 thrombin or prothrombin. The supernatant is then run on an SDS gel. Lane 1 is size standards; lane 2 shows a depletion of thrombin from the supernatant; Lane 3 shows that depletion is dependent on the presence of the IgA; Lanes 3 and 4 show that prothrombin is not depleted, and therefore does not bind to the
25 IgA.

Figure 3 shows the relative rate of S2238 cleavage by thrombin in the presence or absence of IgA (i.e. a single slope of Abs405 with time for S2238 hydrolysis). This indicates that the IgA
30 does not bind at the thrombin active site.

Figure 4 shows the results of binding studies which indicate that the IgA competes with the fluorescently labelled dodecapeptide hirugen for binding to thrombin.
35

Figure 5 shows the effect of the IgA on the cleavage of S2238 by thrombin. This analysis allows the estimate of Kd for the IgA-thrombin interaction of 12nM.

40 Figure 6 shows an SDS-PAGE gel of whole IgA and Fab fragments under reducing and non-reducing (ox) conditions. The non-reduced IgA is shown to have a molecular weight of between 100-200 kDa and the non-reduced Fab has a molecular weight of about 50kDa.

Figure 7 shows the crystal structure of Thrombin-Fab complex showing interaction between the exosite 1 of thrombin and HCDR3 of the Fab fragment.

5

Figure 8 shows detail of crystal structure showing interaction between specific residues of thrombin exosite 1 and HCDR3 of the Fab fragment.

10 Figure 9 shows fluorescence microscopy images of FeCl₃ induced blood clots in femoral vein injuries in C57BL/6 mice injected with FITC labelled fibrinogen taken at between 2 and 30 minutes. 100ul of PBS was administered (vehicle control).

15 Figure 10 shows fluorescence microscopy images of FeCl₃ induced blood clots in femoral vein injuries in C57BL/6 mice injected with FITC labelled fibrinogen and 40nM (final concentration in mouse blood, equivalent to a dose of approximately 0.6 mg/Kg) anti-exosite 1 IgA (100ul in PBS).

20

Figure 11 shows fluorescence microscopy images of FeCl₃ induced blood clots in femoral vein injuries in C57BL/6 mice injected with FITC labelled fibrinogen and 80nM (final concentration in mouse blood, equivalent to a dose of approximately 1.2 mg/Kg) anti-exosite 1 IgA(100ul in PBS), and a region outside of injury site for comparison.

25

Figure 12 shows fluorescence microscopy) images of FeCl₃ induced blood clots in femoral vein injuries in C57BL/6 mice injected with FITC labelled fibrinogen and 200nM (final concentration in mouse blood, equivalent to a dose of approximately 3 mg/Kg) anti-exosite 1 IgA (100ul in PBS), and a region outside of injury site for comparison.

30

35 Figure 13 shows fluorescence microscopy images of FeCl₃ induced blood clots in femoral vein injuries in C57BL/6 mice injected with FITC labelled fibrinogen and 400nM (final concentration in mouse blood, equivalent to a dose of approximately 6 mg/Kg) anti-exosite 1 IgA (100ul in PBS).

40

Figure 14 shows fluorescence microscopy) images of FeCl₃ induced blood clots in femoral vein injuries in C57BL/6 mice treated with FITC labelled fibrinogen and 4μM (final concentration in mouse

blood, equivalent to a dose of approximately 60 mg/Kg) anti-exosite 1 IgA (100 μ l in PBS).

Figure 15 shows a quantitation of the dose response to anti
5 exosite 1 IgA from the fluorescent images shown in figures 9 to 13.

Figure 16 shows tail bleed times in control C57BL/6 mice and in
10 mice treated with increasing amounts of anti-exosite 1 IgA. The second average excludes the outlier.

Figure 17 shows the results of tail clip assays on wild-type male
C57BL/6 mice (n=5) after injection into tail vein with either IgA
or PBS. 15 mins after injection, tails were cut at diameter of
15 3mm and blood loss monitored over 10min.

Figure 18A to 18D show the results of an FeCl₃ carotid artery
occlusion model on 9 week old WT C57BL/6 male mice injected as
previously with 400nM anti-thrombin IgA (final concentration in
20 blood, equivalent to a dose of approximately 6 mg/Kg) or PBS 15
min prior to injury with 5% FeCl₃ for 2 min. Figure 18A shows
results for a typical PBS-injected mice (occlusion in 20min) and
figures 18B, 18C and 18D show examples of results for mice
treated with 400nM anti-thrombin IgA (no occlusion).

25

Figure 19 shows thrombin times (i.e. clotting of pooled plasma)
with increasing concentrations of IgG and IgA of the invention,
upon addition of 20nM human thrombin.

30 Figure 20 shows the binding of synthetic IgG to immobilized
thrombin (on ForteBio Octet Red instrument).

Figure 21 shows a typical Octet trace for the binding of 24nM
S195A thrombin to immobilized IgG showing the on phase, followed
35 by an off phase. The black line is the fit.

Figure 22 shows an Octet trace of 500nM prothrombin with a tip
loaded with immobilized IgG. The same conditions were used as the
experiment with thrombin in fig 21. There is no evidence of
40 binding, even at this high concentration.

Experiments

1. Antibody Isolation and Characterisation

Coagulation screening was carried out on a blood plasma sample from a patient. The coagulation tests were performed on a patient who suffered subdural haematoma following head injury. The haematoma spontaneously resolved without intervention. There
5 was no previous history of bleeding and in the 4 years since the patient presented, there have been no further bleeding episodes. The results are shown in Table 1.

The prothrombin time (PT), activated partial thromboplastin time (APTT), and thrombin time (TT) were all prolonged in the patient
10 compared to controls, but reptilase time was normal.

Thrombin time was not corrected by heparinase, indicating that heparin treatment or contamination was not responsible.
15 Fibrinogen levels were normal in the patient, according to ELISA and Reptilase assays. The Clauss assay gave an artifactually low fibrinogen level due to the presence of the thrombin inhibitor. The PT and APTT clotting times were found to remain prolonged following a mixing test using a 50:50 mix with pooled plasma from
20 normal individuals. This showed the presence of an inhibitor in the sample from the patient.

The patient's blood plasma was found to have a high titre of an IgA. This IgA molecule was found to bind to a human thrombin
25 column (Figure 1). IgA binding lectin-agarose pulled down thrombin in the presence but not the absence of the IgA. Prothrombin was not pulled down by the lectin-agarose in the presence of the IgA, indicating that the IgA specifically binds to thrombin but not prothrombin (Figure 2).

30 The binding site of the IgA on the thrombin molecule was then investigated.

A slightly higher rate of cleavage of S2238 by thrombin was
35 measured in the presence of the IgA, indicating that the IgA does not block the active site of thrombin (Figure 3).

The binding of fluorescently labelled hirugen to thrombin is
40 inhibited by the presence of 700 nM of the IgA, indicating that the epitope for the antibody overlaps with the binding site of hirugen on thrombin, namely the exosite 1 of thrombin (Figure 4).

The effect of the IgA on the hydrolysis of some of thrombin's procoagulant substrates was tested. The results are shown in Table 2. These results demonstrate that the IgA molecule isolated from the patient sample inhibits multiple procoagulant activities of thrombin.

Inhibition of thrombin by antithrombin (AT) in the presence of the IgA was only marginally affected in both the absence and presence of heparin (Table 3).

The dissociation constant (K_d) of the IgA for thrombin was initially estimated based on rate of S2238 hydrolysis to be approximately 12nM (Figure 5). The K_d for the binding of the IgA to S195A thrombin (inactivated by mutation of the catalytic serine) was determined to be 2nM using the ForteBio Octet Red instrument (Table 4).

The purified IgA was cleaved with papain (Figure 6), and the Fab fragment was isolated and combined with human PPACK-Thrombin (PPACK is a covalent active site inhibitor). The human PPACK-Thrombin-FAB complex was crystallized and used for structural analysis. The statistics of the structure obtained were as follows: resolution is 1.9Å; Rfactor = 19.43%; Rfree = 23.42%; one complex in the asymmetric unit; Ramachandran: favoured = 97.0%, outliers = 0%. The crystal structure revealed a close association between the HCDR3 of the IgA Fab and the exosite 1 of thrombin (Figure 7).

In particular, residues M32, F34, Q38, E39, L40, L65, R67, R73, T74, R75, Y76, R77a and I82 of the exosite 1 all directly interact with the HCDR3 loop of the IgA Fab (Figure 8).

PISA analysis of the antibody-thrombin interface showed that the total buried surface area in the complex is 1075 Å². The contact residues in the IgA heavy chain were (Kabat numbering): 30, 51, 52a, 53-55, 96, 98, 99, 100, 100a, 100b, 100c, 100d). These are all in CDRs: CDRH1- GYTLTEAAIH; CDRH2- GLDPQDGETVYAQQFKG; CDRH3- GDFSEFEPFSMDYFHF (underlined residues contacting). CDRH3 was found to be the most important, providing 85% of the buried surface area on the antibody. The light chain made one marginal contact with Tyr49, right before CDRL2 (with Ser36a of thrombin). Some individual contributions to buried surface were: Glu99 54Å², Phe100 134.8 Å², Glu100a 80.6 Å², Phe100c 141.7 Å².

The contact residues in thrombin were found to be (chymotrypsin numbering): 32, 34, 36a-40, 65, 67, 73-76, 77a, 82, and 151. The most important individual contributors to the buried surface
5 were: Gln38 86.4 Å², Arg73 44.5 Å², Thr74 60.1 Å², Tyr76 78.4 Å², Arg77a 86.9 Å².

The patient did not display increased or abnormal bleeding or haemorrhage, in spite of 3g/l circulating levels of this IgA,
10 demonstrating that the antibody inhibits thrombin without affecting normal haemostasis.

2. The effect of IgA on Animal Thrombosis Models

C57BL/6 mice were anaesthetized. A catheter was inserted in the
15 carotid artery (for compound injection). FITC labelled fibrinogen (2mg/ml) was injected via the carotid artery. PBS (control) or IgA was also injected via the carotid artery. The femoral vein was exposed and 10% FeCl₃ applied (saturated blotting paper 3mm in length) for 3 min to induce clotting.

20

Fluorescence microscopy images were taken along the length of injury site at 0, 5, 10, and 20 min post FeCl₃ injury using fluorescence microscopy techniques.

25 Clots (fibrin deposits) in the femoral vein were clearly visible as bright areas (figure 9). The lowest dose of the antibody was observed to cause significant inhibition of clotting but as the dose increased, clotting was abolished (figures 10 to 15).

30 The bleeding times of the mice were also measured. Bleeding times were assessed as time to cessation of blood flow after a tail cut. Despite the presence of a single outlier sample, the bleeding time was found to be unaffected by treatment with anti-exosite 1 IgA (figure 16).

35

These results show that the anti-exosite 1 IgA antibody is a potent inhibitor of thrombosis but has no effect on bleeding time.

3. Tail clip assays

40 A tail clip assay was performed on wild-type male C57BL/6 mice injected with either 400nM IgA (final concentration in blood,

equivalent to a dose of approximately 6 mg/Kg) or PBS. Blood loss was monitored over 10mins after the tail was cut at 3mm diameter 15 minutes after the injection. Total blood loss was found to be unaffected by treatment with anti exosite 1 IgA (figure 17).

5

4. FeCl₃ injury carotid artery occlusion

FeCl₃ injury carotid artery occlusion studies were performed on 9 week old WT C57BL/6 male mice. Mice were injected with 400nM anti-IIa IgA (final concentration in blood, equivalent to a dose of approximately 6 mg/Kg) or PBS 15 min prior to injury with 5% FeCl₃ for 2 min. Blood flow was then monitored by Doppler and the time to occlusion measured. A "clot" was defined as stable occlusive thrombus where blood flow was reduced to values typically less than 0.1ml/min and stayed reduced. In the control mice, a stable clot was observed to form about 20mins after injury (Figure 18A). However, the majority of mice treated with 400nM anti-IIa IgA were unable to form stable clots and gave traces in which the clots were quickly resolved, repeatedly resolved or never formed. Three representative traces are shown in Figures 18B to 18D.

10
15
20

5. Anti-exosite 1 IgG

The IgA molecule identified in the patient described above was re-formatted as an IgG using standard techniques.

25

The clotting time of pooled human plasma spiked with increasing amounts of the original IgA and the new IgG was tested upon addition of human thrombin to 20nM (Figure 19). Both parent IgA and the synthetic IgG increased time to clot formation in an identical concentration-dependent manner, implying identical affinities for thrombin.

30

This was confirmed by measuring the binding of synthetic IgG to immobilized S195A thrombin using a ForteBio™ Octet Red instrument. Thrombin was attached to the probe and the binding of the antibodies (at various concentrations) was monitored. On-rates and off-rates were determined. Both antibodies gave similar on-rates of approximately $3 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$ and off-rates of approximately $5 \times 10^{-4} \text{ s}^{-1}$, and dissociation constants (Kd) of approximately 2nM. Kds of approximately 2nM were also obtained for the IgA and the IgG by steady-state analysis (Table 4). A representative steady state curve is shown in Figure 20. The

35
40

properties of the IgA were therefore reproduced on an IgG framework.

Binding of prothrombin to the IgG antibody was tested using the
5 Octet system by immobilizing IgG. Thrombin bound to the
immobilized IgG with comparable rates and affinities as those
obtained using immobilized thrombin (Table 4); prothrombin did
not bind to the IgG. Figure 21 is a trace of 24nM thrombin
binding to and dissociating from the immobilized IgG. Figure 22
10 is the same experiment using 500nM prothrombin, and shows no
evidence of binding.

Sequences

Amino acid sequence of human preprothrombin (SEQ ID NO: 1;
GeneID: 2147; NP_000497.1 CI: 4503635; exosite 1 residues

5 underlined)

```

1 mahvrglqlp gclalaaals lvhsqhvfla pqqarsllqr vrrantflee vrkgnlerec
6 veetcsyeea fealesstat dvfwakytac etartprckl aaclegnae glgtnyrghv
12 nitrsgiecq lwrsryphkp einstthpga dlqenfcrrp dssttgpcwy ttdptvrrqe
10 18 csipvcqqdq vtvamtprse gssvnlsppl eqcvpdrqqq yqgrlavtth glpclawasa
24 qakalskhqd fnsavqlven fcrnpgddee gwvcyvagkp gdfgycdlny ceeaveeetg
30 dgldegsdra iegrtatsey qtffnprtfg sgeadcglrp lfekksledk terellesyi
36 dgrivegsda eigmspwqvm lfrkspqell cgaslisdrw vltaahclly ppwdknften
42 dllvrigkhs rtryerniek ismlekiyih prynwrenld rdialmklkk pvafsdyyhp
15 48 vclpdretaa sllqagykgr vtgwgnlket wtanvgkqgp svlqvvnlpi verpvckdst
54 riritdnmfc agykpdegkr gdacegdsgg pfvmkspfnn rwyqmgivsw gegcdrdgyk
60 gfythvfrlk kwiqkvidqf ge

```

Amino acid sequence of anti-exosite 1 IgA and IgG VH domain with
20 Kabat Numbering (CDRs underlined): (SEQ ID NO: 2).

```

QVQLIQSGSAVKKPGASVRVSVCKVSGYTLTEAAIHWVRQAPGKGLEWMGG
      10      20      30      40      50

```

```

25 LDPQDGETVYAQQFKGRVIMTEDRSTDTAYMEVNNLRSEDTATYYCTTGD
52a      60      70      8082abc      90

```

```

FSEFEPFSMDYFHFWGQGTVVTVAS
100abcdefgh      110

```

30

Amino acid sequence of anti-exosite 1 IgA and IgG HCDR1 (SEQ ID
NO: 3).

GYTLTEAAIH

35

Amino acid sequence of anti-exosite 1 IgA and IgG HCDR2 (SEQ ID
NO: 4).

GLDPQDGETVYAQQFKG

40

Amino acid sequence of anti-exosite 1 IgA and IgG HCDR3 (SEQ ID
NO: 5).

GDFSEFEPFSMDYFHF

45

WO 2013/088164

PCT/GB2012/053140

Amino acid sequence of anti-exosite 1 IgA and IgG VL domain with Kabat Numbering: (SEQ ID NO: 6).

EIVLTQSPATLSLSPGERATLSCRASQNVSSFLAWYQHKPCQAPRLLIYD
 5 10 20 30 40 50

ASSRATDIPIRFSGSGSGTDFLTISGLEPEDFAVYYCQRRSWPPLTFG
 60 70 80 90 95a

10 GGTKVEIKR
 100 108

Amino acid sequence of anti-exosite 1 IgA and IgG LCDR1 (SEQ ID NO: 7).

15 RASQNVSSFLA

Amino acid sequence of anti-exosite 1 IgA and IgG LCDR2 (SEQ ID NO: 8).

20 DASSRAT

Amino acid sequence of anti-exosite 1 IgA and IgG LCDR3 (SEQ ID NO: 9).

25 QQRRSWPPLT

| Test | | Result | Control/Normalised Ratio (NR) |
|--------------------------------|------------------|-----------|-------------------------------|
| Prothrombin Time | | 43 sec. | NR = 11-13 sec. |
| | 50:50 correction | 35 sec. | |
| Act. part. Thromboplastin Time | | 157 sec. | NR = 22-23 sec. |
| | 50:50 correction | 105 sec. | |
| Thrombin Time | | >150 sec. | NR = 10-13 sec. |
| Reptilase Time | | 16 sec. | Control = 15 sec. |
| Fibrinogen | Clauss | 0.7 g/l | NR = 1.5-4.5 g/l |
| | Antigenic | 5.0 g/l | |

5 Table 1 - Coagulation Screening Results

| Thrombin substrate | Activity | Antibody Effect |
|---------------------------------|---|--------------------------------|
| Fibrinogen | Formation of fibrin clot | No detectable cleavage |
| Platelet receptor PAR-1 peptide | Activation of platelets | 15-fold decrease in hydrolysis |
| FVIII | Feedback activation of thrombin <i>via</i> Xase complex | 7-fold decrease in hydrolysis |

10 Table 2 - Effect of anti-exosite 1 IgA on thrombin hydrolysis of procoagulant substrates

| | Rate of Inhibition($M^{-1}s^{-1}$) | Heparin effect |
|------------|--------------------------------------|----------------|
| AT | $4.8 \pm 0.2 \times 10^3$ | 2.4-fold |
| AT+Hep | $11.8 \pm 0.3 \times 10^3$ | |
| AT+Fab | $1.7 \pm 0.1 \times 10^3$ | 3.3-fold |
| AT+Hep+Fab | $5.6 \pm 0.3 \times 10^3$ | |

5 Table 3 - Effect of saturating concentration of anti-exosite 1 IgA (Fab) on thrombin inhibition by antithrombin (AT) in the absence and presence of 1nM heparin (Hep).

| | Kd (nM) * | k_{on} ($M^{-1}s^{-1}$) | k_{off} (s^{-1}) | Kd (nM)# |
|----------------------|---------------|-----------------------------|------------------------------|---------------|
| IgA | 1.8 | 3.3×10^5 | 3.7×10^{-4} | 1.2 |
| IgG | 1.5 ± 0.3 | $3.3 \pm 0.5 \times 10^5$ | $6.8 \pm 1.1 \times 10^{-4}$ | 2.1 ± 0.3 |
| IgG FAB | ND | 5.0×10^5 | 2.7×10^{-3} | 5.3 |
| IgG FAB ⁺ | 3.3 ± 0.3 | 4.3×10^5 | 2.1×10^{-3} | 4.9 |

10 Table 4 - Binding constants of anti-exosite 1 IgA (n=1 under this precise condition), IgG (n=3) antibodies, and IgG-derived FAB to S195A thrombin (active site free, recombinant thrombin). * Kd determined from steady-state analysis of response vs. concentration. # Kd calculated from rates. + Determined using
15 immobilised FAB.

Claims

1. An isolated antibody molecule that specifically binds to the exosite 1 region of thrombin, wherein the antibody molecule
5 comprises: a VH domain comprising an HCDR1, HCDR2 and HCDR3 having the sequences of SEQ ID NOs 3, 4 and 5, respectively; and a VL domain comprising an LCDR1, LCDR2 and LCDR3 having the sequences of SEQ ID NOs 7, 8 and 9, respectively, or an LCDR1 having the sequence of SEQ ID NO: 7 with a glycosylation site
10 mutated out with a substitution at an amino acid residue corresponding to N28 or S30 in SEQ ID NO: 6.
2. The antibody molecule according to claim 1, wherein the antibody inhibits thrombin activity.
15
3. The antibody molecule according to claim 1 or claim 2, wherein the antibody molecule comprises a VH domain having the amino acid sequence of SEQ ID NO: 2.
- 20 4. The antibody molecule according to any one of claims 1 to 3, wherein the antibody molecule comprises a VL domain having the amino acid sequence of SEQ ID NO: 6 or the amino acid sequence of SEQ ID NO: 6 wherein a glycosylation site is mutated out with a substitution at an amino acid residue corresponding to N28 or S30
25 in SEQ ID NO: 6.
5. The antibody molecule according to any one of claims 1 to 4, comprising a VH domain having the amino acid sequence of SEQ ID NO: 2, a VL domain having the amino acid sequence of SEQ ID
30 NO: 6, or a VL domain having the amino acid sequence of SEQ ID NO: 6 wherein a glycosylation site is mutated out with a substitution at N28 or S30.
6. The antibody molecule according to claim 5, comprising a
35 VL domain having the amino acid sequence of SEQ ID NO: 6 wherein the glycosylation site is mutated out with a substitution at S30.
7. The antibody molecule according to any one of claims 1 to 6, which is a whole antibody.
40
8. The antibody molecule according to claim 7, which is an IgA or IgG.

9. The antibody molecule according to any one of claims 1 to 8, which is a monoclonal antibody.

10. The antibody molecule according to any one of claims 1 to 6 which is an antibody fragment.

11. A pharmaceutical composition comprising the antibody molecule according to any one of claims 1 to 10 and a pharmaceutically acceptable excipient.

10

12. An antibody molecule according to any one of claims 1 to 10 for use in treating a thrombin-mediated condition selected from thrombosis and embolism.

15


13. Use of the antibody molecule according to any one of claims 1 to 10 in the manufacture of a medicament for use in treating a thrombin-mediated condition selected from thrombosis and embolism.

20

14. Use of the antibody molecule according to any one of claims 1 to 12 for treating a thrombin-mediated condition selected from thrombosis and embolism.

1/19

A



B



Figure 1




Figure 2

3/19




Figure 3

4/19




Figure 4

5/19




Figure 5

6/19




Figure 6

7/19




Figure 7




Figure 8

8/19




Figure 9




Figure 10

9/19




Figure 11




Figure 12

10/19




Figure 13




Figure 14

11/19




Figure 15

12/19




Figure 16




Figure 17




Figure 18A




Figure 18B




Figure 18C




Figure 18D




Figure 19




Figure 20




Figure 21




Figure 22