



- (51) **International Patent Classification:**
C07D 401/06 (2006.01)
- (21) **International Application Number:**
PCT/US2017/022570
- (22) **International Filing Date:**
15 March 2017 (15.03.2017)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
15/070,938 15 March 2016 (15.03.2016) US
- (71) **Applicant:** **CELGENE QUANTICEL RESEARCH, INC.** [US/US]; 9393 Towne Centre Drive, San Diego, California 92121 (US).
- (72) **Inventors:** **CHEN, Young**; 746 Santa Barbara Drive, San Marcos, California 92078 (US). **NIE, Zhe**; 17104 Monterey Ridge Way, San Diego, California 92127 (US). **STAFFORD, Jeffrey**; 12752 Sandy Crest Court, San Diego, California 92130 (US). **VEAL, James**; 8916 Weaver Crossing Road, Apex, California 27502 (US).
- (74) **Agent:** **SYLVIA, Mary**; Wiley Rein, LLP, Patent Administration, 1776 K Street, NW, Washington, District of Columbia 20006 (US).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

(54) **Title:** HISTONE DEMETHYLASE INHIBITORS

(57) **Abstract:** The present invention relates generally to compositions and methods for treating cancer and neoplastic diseases. Provided herein are substituted imidazole-pyridine derivative compounds and pharmaceutical compositions comprising said compounds. The subject compounds and compositions are useful for inhibition of histone demethylase enzymes. Furthermore, the subject compounds and compositions are useful for the treatment of cancer, such as pancreatic cancer, prostate cancer, breast cancer, bladder cancer, lung cancer, gastric cancer, leukemia and/or melanoma and the like.



HISTONE DEMETHYLASE INHIBITORS

CROSS-REFERENCE

[0001] This application is a continuation-in-part of U.S. Patent Application No. 14/855,269, filed on September 15, 2015, which claims the benefit of U.S. Provisional Application 62/051,268, filed September 16, 2014, the contents of both of which are hereby incorporated by reference in their entirety for all purposes.


BACKGROUND

[0002] A need exists in the art for an effective treatment of cancer and neoplastic diseases.

BRIEF SUMMARY OF THE INVENTION

[0003] Provided herein are substituted imidazole-pyridine derivative compounds and pharmaceutical compositions comprising said compounds. The subject compounds and compositions are useful for the inhibition of histone demethylases. Furthermore, the subject compounds and compositions are useful for the treatment of cancer, such as pancreatic cancer, prostate cancer, breast cancer, bladder cancer, lung cancer, gastric cancer, leukemia and/or melanoma and the like. The substituted imidazole-pyridine derivative compounds described herein are based upon a substituted 2-(1H-imidazol-4-yl)pyridine ring system bearing a fused or linked triazole ring system.

[0004] One embodiment provides a compound having the structure of Formula (I),



or a pharmaceutically acceptable salt thereof, wherein

R^1 is hydrogen, halogen, -OH, -OR⁵, -N(R⁵)₂, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclalkyl, heterocyclalkyl, aralkyl, or heteroarylalkyl;


R^2 is alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclalkyl, heterocyclalkyl, aralkyl, or heteroarylalkyl;

R^3 is hydrogen, halogen, -OH, -NH₂, -NH(C₁-C₃alkyl) or C₁-C₃alkyl;

R^4 is hydrogen, C₁-C₄alkyl, C₁-C₄alkene, C₁-C₄alkyne, halogen, or -CN; and

each R^5 is independently hydrogen, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclylalkyl, heterocyclylalkyl, aralkyl, or heteroarylalkyl.

[0005] One embodiment provides a compound having the structure of Formula (II),



or a pharmaceutically acceptable salt thereof, wherein

R^1 is hydrogen, halogen, -OH, -OR⁴, -N(R⁴)₂, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclylalkyl, heterocyclylalkyl, aralkyl, or heteroarylalkyl;

R^2 is alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclylalkyl, heterocyclylalkyl, aralkyl, or heteroarylalkyl;

R^3 is hydrogen, halogen, -OH, -NH₂, -NH(C1-C3alkyl) or C1-C3alkyl; and

each R^4 is independently hydrogen, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclylalkyl, heterocyclylalkyl, aralkyl, or heteroarylalkyl.

[0006] One embodiment provides a pharmaceutical composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient. One embodiment provides a pharmaceutical composition comprising a compound of Formula (II), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.

[0007] One embodiment provides a method for inhibiting a histone demethylase enzyme comprising contacting the histone demethylase enzyme with a compound of Formula (I). One embodiment provides a method for inhibiting a histone demethylase enzyme comprising contacting the histone demethylase enzyme with a compound of Formula (II).

[0008] One embodiment provides a method of treating cancer in a patient in need thereof, comprising administering to the patient a compound of Formula (I), or a pharmaceutically acceptable salt thereof. One embodiment provides a method of treating cancer in a patient in need thereof, comprising administering to the patient a compound of Formula (II), or a pharmaceutically acceptable salt thereof.

INCORPORATION BY REFERENCE

[0009] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

DETAILED DESCRIPTION OF THE INVENTION

[0010] As used herein and in the appended claims, the singular forms “a,” “and,” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “an agent” includes a plurality of such agents, and reference to “the cell” includes reference to one or more cells (or to a plurality of cells) and equivalents thereof known to those skilled in the art, and so forth. When ranges are used herein for physical properties, such as molecular weight, or chemical properties, such as chemical formulae, all combinations and subcombinations of ranges and specific embodiments therein are intended to be included. The term “about” when referring to a number or a numerical range means that the number or numerical range referred to is an approximation within experimental variability (or within statistical experimental error), and thus the number or numerical range may vary between 1% and 15% of the stated number or numerical range. The term “comprising” (and related terms such as “comprise” or “comprises” or “having” or “including”) is not intended to exclude that in other certain embodiments, for example, an embodiment of any composition of matter, composition, method, or process, or the like, described herein, may “consist of” or “consist essentially of” the described features.

Definitions

[0011] As used in the specification and appended claims, unless specified to the contrary, the following terms have the meaning indicated below.

- [0012] “Amino” refers to the -NH_2 radical.
- [0013] “Cyano” refers to the -CN radical.
- [0014] “Nitro” refers to the -NO_2 radical.
- [0015] “Oxa” refers to the -O- radical.
- [0016] “Oxo” refers to the =O radical.
- [0017] “Thioxo” refers to the =S radical.
- [0018] “Imino” refers to the =N-H radical.
- [0019] “Oximo” refers to the =N-OH radical.
- [0020] “Hydrazino” refers to the =N-NH_2 radical.

[0021] “Alkyl” refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, having from one to fifteen carbon atoms (e.g., C₁-C₁₅ alkyl). In certain embodiments, an alkyl comprises one to thirteen carbon atoms (e.g., C₁-C₁₃ alkyl). In certain embodiments, an alkyl comprises one to eight carbon atoms (e.g., C₁-C₈ alkyl). In other embodiments, an alkyl comprises one to five carbon atoms (e.g., C₁-C₅ alkyl). In other embodiments, an alkyl comprises one to four carbon atoms (e.g., C₁-C₄ alkyl). In other embodiments, an alkyl comprises one to three carbon atoms (e.g., C₁-C₃ alkyl). In other embodiments, an alkyl comprises one to two carbon atoms (e.g., C₁-C₂ alkyl). In other embodiments, an alkyl comprises one carbon atom (e.g., C₁ alkyl). In other embodiments, an alkyl comprises five to fifteen carbon atoms (e.g., C₅-C₁₅ alkyl). In other embodiments, an alkyl comprises five to eight carbon atoms (e.g., C₅-C₈ alkyl). In other embodiments, an alkyl comprises two to five carbon atoms (e.g., C₂-C₅ alkyl). In other embodiments, an alkyl comprises three to five carbon atoms (e.g., C₃-C₅ alkyl). In other embodiments, the alkyl group is selected from methyl, ethyl, 1-propyl (*n*-propyl), 1-methylethyl (*iso*-propyl), 1-butyl (*n*-butyl), 1-methylpropyl (*sec*-butyl), 2-methylpropyl (*iso*-butyl), 1,1-dimethylethyl (*tert*-butyl), 1-pentyl (*n*-pentyl). The alkyl is attached to the rest of the molecule by a single bond. Unless stated otherwise specifically in the specification, an alkyl group is optionally substituted by one or more of the following substituents: halo, cyano, nitro, oxo, thioxo, imino, oximo, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -OC(O)-N(R^a)₂, -N(R^a)C(O)R^a, -N(R^a)S(O)_tR^a (where t is 1 or 2), -S(O)_tOR^a (where t is 1 or 2), -S(O)_tR^a (where t is 1 or 2) and -S(O)_tN(R^a)₂ (where t is 1 or 2) where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl.

[0022] “Alkoxy” refers to a radical bonded through an oxygen atom of the formula -O-alkyl, where alkyl is an alkyl chain as defined above.

[0023] “Alkenyl” refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one carbon-carbon double bond, and having from two to twelve carbon atoms. In certain embodiments, an alkenyl comprises two to eight carbon atoms. In other embodiments, an alkenyl comprises two to four carbon atoms. The alkenyl is attached to the rest of the molecule by a single bond, for example, ethenyl (i.e., vinyl), prop-1-enyl (i.e., allyl), but-1-enyl, pent-1-enyl, penta-1,4-dienyl, and the like. Unless stated otherwise specifically in the specification, an alkenyl group is optionally substituted by one or more of the following substituents: halo, cyano, nitro, oxo, thioxo, imino, oximo, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -OC(O)-N(R^a)₂,

$-N(R^a)C(O)R^a$, $-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-S(O)_tOR^a$ (where t is 1 or 2), $-S(O)_tR^a$ (where t is 1 or 2) and $-S(O)_tN(R^a)_2$ (where t is 1 or 2) where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl.

[0024] “Alkynyl” refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one carbon-carbon triple bond, having from two to twelve carbon atoms. In certain embodiments, an alkynyl comprises two to eight carbon atoms. In other embodiments, an alkynyl has two to four carbon atoms. The alkynyl is attached to the rest of the molecule by a single bond, for example, ethynyl, propynyl, butynyl, pentynyl, hexynyl, and the like. Unless stated otherwise specifically in the specification, an alkynyl group is optionally substituted by one or more of the following substituents: halo, cyano, nitro, oxo, thioxo, imino, oximo, trimethylsilanyl, $-OR^a$, $-SR^a$, $-OC(O)-R^a$, $-N(R^a)_2$, $-C(O)R^a$, $-C(O)OR^a$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)OR^a$, $-OC(O)-N(R^a)_2$, $-N(R^a)C(O)R^a$, $-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-S(O)_tOR^a$ (where t is 1 or 2), $-S(O)_tR^a$ (where t is 1 or 2) and $-S(O)_tN(R^a)_2$ (where t is 1 or 2) where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl.

[0025] “Alkylene” or “alkylene chain” refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, containing no unsaturation and having from one to twelve carbon atoms, for example, methylene, ethylene, propylene, *n*-butylene, and the like. The alkylene chain is attached to the rest of the molecule through a single bond and to the radical group through a single bond. The points of attachment of the alkylene chain to the rest of the molecule and to the radical group can be through one carbon in the alkylene chain or through any two carbons within the chain. In certain embodiments, an alkylene comprises one to eight carbon atoms (e.g., C_1 - C_8 alkylene). In other embodiments, an alkylene comprises one to five carbon atoms (e.g., C_1 - C_5 alkylene). In other embodiments, an alkylene comprises one to four carbon atoms (e.g., C_1 - C_4 alkylene). In other embodiments, an alkylene comprises one to three carbon atoms (e.g., C_1 - C_3 alkylene). In other embodiments, an alkylene comprises one to two carbon atoms (e.g., C_1 - C_2 alkylene). In other embodiments, an alkylene comprises one carbon atom (e.g., C_1 alkylene). In other embodiments, an alkylene comprises five to eight carbon atoms (e.g., C_5 - C_8 alkylene). In other embodiments, an alkylene comprises two to five carbon atoms (e.g., C_2 - C_5 alkylene). In other embodiments, an alkylene comprises three to five carbon atoms (e.g., C_3 - C_5 alkylene). Unless stated otherwise specifically in the specification, an alkylene chain is optionally substituted by one or more of the

following substituents: halo, cyano, nitro, oxo, thioxo, imino, oximo, trimethylsilyl, $-OR^a$, $-SR^a$, $-OC(O)-R^a$, $-N(R^a)_2$, $-C(O)R^a$, $-C(O)OR^a$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)OR^a$, $-OC(O)-N(R^a)_2$, $-N(R^a)C(O)R^a$, $-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-S(O)_tOR^a$ (where t is 1 or 2), $-S(O)_tR^a$ (where t is 1 or 2) and $-S(O)_tN(R^a)_2$ (where t is 1 or 2) where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl.

[0026] “Aryl” refers to a radical derived from an aromatic monocyclic or multicyclic hydrocarbon ring system by removing a hydrogen atom from a ring carbon atom. The aromatic monocyclic or multicyclic hydrocarbon ring system contains only hydrogen and carbon from five to eighteen carbon atoms, where at least one of the rings in the ring system is fully unsaturated, i.e., it contains a cyclic, delocalized $(4n+2)$ π -electron system in accordance with the Hückel theory. The ring system from which aryl groups are derived include, but are not limited to, groups such as benzene, fluorene, indane, indene, tetralin and naphthalene. Unless stated otherwise specifically in the specification, the term “aryl” or the prefix “ar-” (such as in “aralkyl”) is meant to include aryl radicals optionally substituted by one or more substituents independently selected from alkyl, alkenyl, alkynyl, halo, fluoroalkyl, cyano, nitro, optionally substituted aryl, optionally substituted aralkyl, optionally substituted aralkenyl, optionally substituted aralkynyl, optionally substituted carbocyclyl, optionally substituted carbocyclalkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^b-OR^a$, $-R^b-OC(O)-R^a$, $-R^b-OC(O)-OR^a$, $-R^b-OC(O)-N(R^a)_2$, $-R^b-N(R^a)_2$, $-R^b-C(O)R^a$, $-R^b-C(O)OR^a$, $-R^b-C(O)N(R^a)_2$, $-R^b-O-R^c-C(O)N(R^a)_2$, $-R^b-N(R^a)C(O)OR^a$, $-R^b-N(R^a)C(O)R^a$, $-R^b-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-R^b-S(O)_tOR^a$ (where t is 1 or 2), $-R^b-S(O)_tR^a$ (where t is 1 or 2) and $-R^b-S(O)_tN(R^a)_2$ (where t is 1 or 2), where each R^a is independently hydrogen, alkyl, fluoroalkyl, cycloalkyl, cycloalkylalkyl, aryl (optionally substituted with one or more halo groups), aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl, each R^b is independently a direct bond or a straight or branched alkylene or alkenylene chain, and R^c is a straight or branched alkylene or alkenylene chain, and where each of the above substituents is unsubstituted unless otherwise indicated.

[0027] “Aralkyl” refers to a radical of the formula $-R^c$ -aryl where R^c is an alkylene chain as defined above, for example, methylene, ethylene, and the like. The alkylene chain part of the aralkyl radical is optionally substituted as described above for an alkylene chain. The aryl part of the aralkyl radical is optionally substituted as described above for an aryl group.

[0028] “Aralkenyl” refers to a radical of the formula $-R^d$ -aryl where R^d is an alkenylene chain as defined above. The aryl part of the aralkenyl radical is optionally substituted as described above for an aryl group. The alkenylene chain part of the aralkenyl radical is optionally substituted as defined above for an alkenylene group.

[0029] “Aralkynyl” refers to a radical of the formula $-R^e$ -aryl, where R^e is an alkynylene chain as defined above. The aryl part of the aralkynyl radical is optionally substituted as described above for an aryl group. The alkynylene chain part of the aralkynyl radical is optionally substituted as defined above for an alkynylene chain.

[0030] “Aralkoxy” refers to a radical bonded through an oxygen atom of the formula $-O-R^c$ -aryl where R^c is an alkylene chain as defined above, for example, methylene, ethylene, and the like. The alkylene chain part of the aralkyl radical is optionally substituted as described above for an alkylene chain. The aryl part of the aralkyl radical is optionally substituted as described above for an aryl group.


[0031] “Carbocyclyl” refers to a stable non-aromatic monocyclic or polycyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, which may include fused or bridged ring systems, having from three to fifteen carbon atoms. In certain embodiments, a carbocyclyl comprises three to ten carbon atoms. In other embodiments, a carbocyclyl comprises five to seven carbon atoms. The carbocyclyl is attached to the rest of the molecule by a single bond. Carbocyclyl may be saturated, (i.e., containing single C-C bonds only) or unsaturated (i.e., containing one or more double bonds or triple bonds.) A fully saturated carbocyclyl radical is also referred to as “cycloalkyl.” Examples of monocyclic cycloalkyls include, e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. An unsaturated carbocyclyl is also referred to as “cycloalkenyl.” Examples of monocyclic cycloalkenyls include, e.g., cyclopentenyl, cyclohexenyl, cycloheptenyl, and cyclooctenyl. Polycyclic carbocyclyl radicals include, for example, adamantyl, norbornyl (i.e., bicyclo[2.2.1]heptanyl), norbornenyl, decalanyl, 7,7-dimethyl-bicyclo[2.2.1]heptanyl, and the like. Unless otherwise stated specifically in the specification, the term “carbocyclyl” is meant to include carbocyclyl radicals that are optionally substituted by one or more substituents independently selected from alkyl, alkenyl, alkynyl, halo, fluoroalkyl, oxo, thioxo, cyano, nitro, optionally substituted aryl, optionally substituted aralkyl, optionally substituted aralkenyl, optionally substituted aralkynyl, optionally substituted carbocyclyl, optionally substituted carbocyclylalkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^b-OR^a$, $-R^b-OC(O)-R^a$, $-R^b-OC(O)-OR^a$, $-R^b-OC(O)-N(R^a)_2$, $-R^b-N(R^a)_2$,

$-R^b-C(O)R^a$, $-R^b-C(O)OR^a$, $-R^b-C(O)N(R^a)_2$, $-R^b-O-R^c-C(O)N(R^a)_2$, $-R^b-N(R^a)C(O)OR^a$, $-R^b-N(R^a)C(O)R^a$, $-R^b-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-R^b-S(O)_tOR^a$ (where t is 1 or 2), $-R^b-S(O)_tR^a$ (where t is 1 or 2) and $-R^b-S(O)_tN(R^a)_2$ (where t is 1 or 2), where each R^a is independently hydrogen, alkyl, fluoroalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, each R^b is independently a direct bond or a straight or branched alkylene or alkenylene chain, and R^c is a straight or branched alkylene or alkenylene chain, and where each of the above substituents is unsubstituted unless otherwise indicated.

[0032] “Carbocyclylalkyl” refers to a radical of the formula $-R^c$ -carbocyclyl where R^c is an alkylene chain as defined above. The alkylene chain and the carbocyclyl radical is optionally substituted as defined above.

[0033] “Carbocyclylalkoxy” refers to a radical bonded through an oxygen atom of the formula $-O-R^c$ -carbocyclyl where R^c is an alkylene chain as defined above. The alkylene chain and the carbocyclyl radical is optionally substituted as defined above.

[0034] As used herein, “carboxylic acid bioisostere” refers to a functional group or moiety that exhibits similar physical, biological and/or chemical properties as a carboxylic acid moiety. Examples of carboxylic acid bioisosteres include, but are not limited to,



[0035] “Halo” or “halogen” refers to bromo, chloro, fluoro or iodo substituents.

[0036] “Fluoroalkyl” refers to an alkyl radical, as defined above, that is substituted by one or more fluoro radicals, as defined above, for example, trifluoromethyl, difluoromethyl, fluoromethyl, 2,2,2-trifluoroethyl, 1-fluoromethyl-2-fluoroethyl, and the like. The alkyl part of the fluoroalkyl radical may be optionally substituted as defined above for an alkyl group.

[0037] “Heterocyclyl” refers to a stable 3- to 18-membered non-aromatic ring radical that comprises two to twelve carbon atoms and from one to six heteroatoms selected from nitrogen, oxygen and sulfur. Unless stated otherwise specifically in the specification, the heterocyclyl radical is a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems. The heteroatoms in the heterocyclyl radical may be optionally oxidized. One or more nitrogen atoms, if present, are optionally quaternized. The heterocyclyl radical is partially or fully

saturated. The heterocyclyl may be attached to the rest of the molecule through any atom of the ring(s). Examples of such heterocyclyl radicals include, but are not limited to, dioxolanyl, thienyl[1,3]dithianyl, decahydroisoquinolyl, imidazolyl, imidazolidinyl, isothiazolidinyl, isoxazolidinyl, morpholyl, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidinyl, piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidinyl, quinuclidinyl, thiazolidinyl, tetrahydrofuryl, trithianyl, tetrahydropyranyl, thiomorpholyl, thiamorpholyl, 1-oxo-thiomorpholyl, and 1,1-dioxo-thiomorpholyl. Unless stated otherwise specifically in the specification, the term “heterocyclyl” is meant to include heterocyclyl radicals as defined above that are optionally substituted by one or more substituents selected from alkyl, alkenyl, alkynyl, halo, fluoroalkyl, oxo, thioxo, cyano, nitro, optionally substituted aryl, optionally substituted aralkyl, optionally substituted aralkenyl, optionally substituted aralkynyl, optionally substituted carbocyclyl, optionally substituted carbocyclylalkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^b-OR^a$, $-R^b-OC(O)-R^a$, $-R^b-OC(O)-OR^a$, $-R^b-OC(O)-N(R^a)_2$, $-R^b-N(R^a)_2$, $-R^b-C(O)R^a$, $-R^b-C(O)OR^a$, $-R^b-C(O)N(R^a)_2$, $-R^b-O-R^c-C(O)N(R^a)_2$, $-R^b-N(R^a)C(O)OR^a$, $-R^b-N(R^a)C(O)R^a$, $-R^b-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-R^b-S(O)_tOR^a$ (where t is 1 or 2), $-R^b-S(O)_tR^a$ (where t is 1 or 2) and $-R^b-S(O)_tN(R^a)_2$ (where t is 1 or 2), where each R^a is independently hydrogen, alkyl, fluoroalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, each R^b is independently a direct bond or a straight or branched alkylene or alkenylene chain, and R^c is a straight or branched alkylene or alkenylene chain, and where each of the above substituents is unsubstituted unless otherwise indicated.

[0038] “*N*-heterocyclyl” or “*N*-attached heterocyclyl” refers to a heterocyclyl radical as defined above containing at least one nitrogen and where the point of attachment of the heterocyclyl radical to the rest of the molecule is through a nitrogen atom in the heterocyclyl radical. An *N*-heterocyclyl radical is optionally substituted as described above for heterocyclyl radicals. Examples of such *N*-heterocyclyl radicals include, but are not limited to, 1-morpholyl, 1-piperidinyl, 1-piperazinyl, 1-pyrrolidinyl, pyrazolidinyl, imidazolyl, and imidazolidinyl.

[0039] “*C*-heterocyclyl” or “*C*-attached heterocyclyl” refers to a heterocyclyl radical as defined above containing at least one heteroatom and where the point of attachment of the heterocyclyl radical to the rest of the molecule is through a carbon atom in the heterocyclyl radical. A *C*-heterocyclyl radical is optionally substituted as described above for heterocyclyl radicals.

Examples of such C-heterocyclyl radicals include, but are not limited to, 2-morpholinyl, 2- or 3- or 4-piperidinyl, 2-piperazinyl, 2- or 3-pyrrolidinyl, and the like.

[0040] “Heterocyclalkyl” refers to a radical of the formula $-R^c$ -heterocyclyl where R^c is an alkylene chain as defined above. If the heterocyclyl is a nitrogen-containing heterocyclyl, the heterocyclyl is optionally attached to the alkyl radical at the nitrogen atom. The alkylene chain of the heterocyclalkyl radical is optionally substituted as defined above for an alkylene chain. The heterocyclyl part of the heterocyclalkyl radical is optionally substituted as defined above for a heterocyclyl group.

[0041] “Heterocyclalkoxy” refers to a radical bonded through an oxygen atom of the formula $-O-R^c$ -heterocyclyl where R^c is an alkylene chain as defined above. If the heterocyclyl is a nitrogen-containing heterocyclyl, the heterocyclyl is optionally attached to the alkyl radical at the nitrogen atom. The alkylene chain of the heterocyclalkoxy radical is optionally substituted as defined above for an alkylene chain. The heterocyclyl part of the heterocyclalkoxy radical is optionally substituted as defined above for a heterocyclyl group.

[0042] “Heteroaryl” refers to a radical derived from a 3- to 18-membered aromatic ring radical that comprises two to seventeen carbon atoms and from one to six heteroatoms selected from nitrogen, oxygen and sulfur. As used herein, the heteroaryl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, wherein at least one of the rings in the ring system is fully unsaturated, *i.e.*, it contains a cyclic, delocalized $(4n+2)$ π -electron system in accordance with the Hückel theory. Heteroaryl includes fused or bridged ring systems. The heteroatom(s) in the heteroaryl radical is optionally oxidized. One or more nitrogen atoms, if present, are optionally quaternized. The heteroaryl is attached to the rest of the molecule through any atom of the ring(s). Examples of heteroaryls include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzindolyl, 1,3-benzodioxolyl, benzofuranyl, benzooxazolyl, benzo[d]thiazolyl, benzothiadiazolyl, benzo[b][1,4]dioxepinyl, benzo[b][1,4]oxazinyl, 1,4-benzodioxanyl, benzonaphthofuranyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyranyl, benzopyranonyl, benzofuranyl, benzofuranonyl, benzothienyl (benzothiophenyl), benzothieno[3,2-d]pyrimidinyl, benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl, carbazolyl, cinnolinyl, cyclopenta[d]pyrimidinyl, 6,7-dihydro-5H-cyclopenta[4,5]thieno[2,3-d]pyrimidinyl, 5,6-dihydrobenzo[h]quinazolinyl, 5,6-dihydrobenzo[h]cinnolinyl, 6,7-dihydro-5H-benzo[6,7]cyclohepta[1,2-c]pyridazinyl, dibenzofuranyl, dibenzothiophenyl, furanyl, furanonyl, furo[3,2-c]pyridinyl, 5,6,7,8,9,10-hexahydrocycloocta[d]pyrimidinyl, 5,6,7,8,9,10-hexahydrocycloocta[d]pyridazinyl, 5,6,7,8,9,10-hexa-

hydrocycloocta[d]pyridinyl, isothiazolyl, imidazolyl, indazolyl, indolyl, indazolyl, isoindolyl, indolinyl, isoindolinyl, isoquinolyl, indoliziny, isoxazolyl, 5,8-methano-5,6,7,8-tetrahydroquinazolinyl, naphthyridinyl, 1,6-naphthyridinonyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl, 5,6,6a,7,8,9,10,10a-octahydrobenzo[h]quinazolinyl, 1-phenyl-1*H*-pyrrolyl, phenazinyl, phenothiazinyl, phenoxazinyl, phthalazinyl, pteridinyl, purinyl, pyrrolyl, pyrazolyl, pyrazolo[3,4-d]pyrimidinyl, pyridinyl, pyrido[3,2-d]pyrimidinyl, pyrido[3,4-d]pyrimidinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, quinazolinyl, quinoxalinyl, quinoliny, isoquinoliny, tetrahydroquinoliny, 5,6,7,8-tetrahydroquinazolinyl, 5,6,7,8-tetrahydrobenzo[4,5]thieno[2,3-d]pyrimidinyl, 6,7,8,9-tetrahydro-5*H*-cyclohepta[4,5]thieno[2,3-d]pyrimidinyl, 5,6,7,8-tetrahydropyrido[4,5-c]pyridazinyl, thiazolyl, thiadiazolyl, triazolyl, tetrazolyl, triazinyl, thieno[2,3-d]pyrimidinyl, thieno[3,2-d]pyrimidinyl, thieno[2,3-c]pyridinyl, and thiophenyl (i.e. thienyl). Unless stated otherwise specifically in the specification, the term “heteroaryl” is meant to include heteroaryl radicals as defined above which are optionally substituted by one or more substituents selected from alkyl, alkenyl, alkynyl, halo, fluoroalkyl, haloalkenyl, haloalkynyl, oxo, thioxo, cyano, nitro, optionally substituted aryl, optionally substituted aralkyl, optionally substituted aralkenyl, optionally substituted aralkynyl, optionally substituted carbocyclyl, optionally substituted carbocyclylalkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^b-OR^a$, $-R^b-OC(O)-R^a$, $-R^b-OC(O)-OR^a$, $-R^b-OC(O)-N(R^a)_2$, $-R^b-N(R^a)_2$, $-R^b-C(O)R^a$, $-R^b-C(O)OR^a$, $-R^b-C(O)N(R^a)_2$, $-R^b-O-R^c-C(O)N(R^a)_2$, $-R^b-N(R^a)C(O)OR^a$, $-R^b-N(R^a)C(O)R^a$, $-R^b-N(R^a)S(O)_tR^a$ (where *t* is 1 or 2), $-R^b-S(O)_tOR^a$ (where *t* is 1 or 2), $-R^b-S(O)_tR^a$ (where *t* is 1 or 2) and $-R^b-S(O)_tN(R^a)_2$ (where *t* is 1 or 2), where each R^a is independently hydrogen, alkyl, fluoroalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, each R^b is independently a direct bond or a straight or branched alkylene or alkenylene chain, and R^c is a straight or branched alkylene or alkenylene chain, and where each of the above substituents is unsubstituted unless otherwise indicated.

[0043] “*N*-heteroaryl” refers to a heteroaryl radical as defined above containing at least one nitrogen and where the point of attachment of the heteroaryl radical to the rest of the molecule is through a nitrogen atom in the heteroaryl radical. An *N*-heteroaryl radical is optionally substituted as described above for heteroaryl radicals.

[0044] “*C*-heteroaryl” refers to a heteroaryl radical as defined above and where the point of attachment of the heteroaryl radical to the rest of the molecule is through a carbon atom in the


heteroaryl radical. A *C*-heteroaryl radical is optionally substituted as described above for heteroaryl radicals.

[0045] “Heteroarylalkyl” refers to a radical of the formula $-R^c$ -heteroaryl, where R^c is an alkylene chain as defined above. If the heteroaryl is a nitrogen-containing heteroaryl, the heteroaryl is optionally attached to the alkyl radical at the nitrogen atom. The alkylene chain of the heteroarylalkyl radical is optionally substituted as defined above for an alkylene chain. The heteroaryl part of the heteroarylalkyl radical is optionally substituted as defined above for a heteroaryl group.

[0046] “Heteroarylalkoxy” refers to a radical bonded through an oxygen atom of the formula $-O-R^c$ -heteroaryl, where R^c is an alkylene chain as defined above. If the heteroaryl is a nitrogen-containing heteroaryl, the heteroaryl is optionally attached to the alkyl radical at the nitrogen atom. The alkylene chain of the heteroarylalkoxy radical is optionally substituted as defined above for an alkylene chain. The heteroaryl part of the heteroarylalkoxy radical is optionally substituted as defined above for a heteroaryl group.

[0047] The compounds disclosed herein may contain one or more asymmetric centers and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)-. Unless stated otherwise, it is intended that all stereoisomeric forms of the compounds disclosed herein are contemplated by this disclosure. When the compounds described herein contain alkene double bonds, and unless specified otherwise, it is intended that this disclosure includes both *E* and *Z* geometric isomers (*e.g.*, *cis* or *trans*.) Likewise, all possible isomers, as well as their racemic and optically pure forms, and all tautomeric forms are also intended to be included. The term “geometric isomer” refers to *E* or *Z* geometric isomers (*e.g.*, *cis* or *trans*) of an alkene double bond. The term “positional isomer” refers to structural isomers around a central ring, such as *ortho*-, *meta*-, and *para*- isomers around a benzene ring.

[0048] A “tautomer” refers to a molecule wherein a proton shift from one atom of a molecule to another atom of the same molecule is possible. The compounds presented herein may, in certain embodiments, exist as tautomers. In circumstances where tautomerization is possible, a chemical equilibrium of the tautomers will exist. The exact ratio of the tautomers depends on several factors, including physical state, temperature, solvent, and pH. Some examples of tautomeric equilibrium include:



[0049] “Optional” or “optionally” means that a subsequently described event or circumstance may or may not occur and that the description includes instances when the event or circumstance occurs and instances in which it does not. For example, “optionally substituted aryl” means that the aryl radical may or may not be substituted and that the description includes both substituted aryl radicals and aryl radicals having no substitution.

[0050] “Pharmaceutically acceptable salt” includes both acid and base addition salts. A pharmaceutically acceptable salt of any one of the substituted imidazole-pyridine derivative compounds described herein is intended to encompass any and all pharmaceutically suitable salt forms. Preferred pharmaceutically acceptable salts of the compounds described herein are pharmaceutically acceptable acid addition salts and pharmaceutically acceptable base addition salts.

[0051] “Pharmaceutically acceptable acid addition salt” refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, hydroiodic acid, hydrofluoric acid, phosphorous acid, and the like. Also included are salts that are formed with organic acids such as aliphatic mono- and

dicarboxylic acids, phenyl-substituted alkanolic acids, hydroxy alkanolic acids, alkanedioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, etc. and include, for example, acetic acid, trifluoroacetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like. Exemplary salts thus include sulfates, pyrosulfates, bisulfates, sulfites, bisulfites, nitrates, phosphates, monohydrogenphosphates, dihydrogenphosphates, metaphosphates, pyrophosphates, chlorides, bromides, iodides, acetates, trifluoroacetates, propionates, caprylates, isobutyrate, oxalates, malonates, succinate suberates, sebacates, fumarates, maleates, mandelates, benzoates, chlorobenzoates, methylbenzoates, dinitrobenzoates, phthalates, benzenesulfonates, toluenesulfonates, phenylacetates, citrates, lactates, malates, tartrates, methanesulfonates, and the like. Also contemplated are salts of amino acids, such as arginates, gluconates, and galacturonates (see, for example, Berge S.M. et al., "Pharmaceutical Salts," *Journal of Pharmaceutical Science*, 66:1-19 (1997), which is hereby incorporated by reference in its entirety). Acid addition salts of basic compounds may be prepared by contacting the free base forms with a sufficient amount of the desired acid to produce the salt according to methods and techniques with which a skilled artisan is familiar.

[0052] "Pharmaceutically acceptable base addition salt" refers to those salts that retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Pharmaceutically acceptable base addition salts may be formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Salts derived from inorganic bases include, but are not limited to, sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, for example, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, diethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, *N,N*-dibenzylethylenediamine, chlorprocaine, hydrabamine, choline, betaine, ethylenediamine, ethylenedianiline, *N*-methylglucamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, *N*-ethylpiperidine, polyamine resins and the like. See Berge et al., *supra*.

[0053] As used herein, "treatment" or "treating," or "palliating" or "ameliorating" are used interchangeably herein. These terms refers to an approach for obtaining beneficial or desired results

including but not limited to therapeutic benefit and/or a prophylactic benefit. By “therapeutic benefit” is meant eradication or amelioration of the underlying disorder being treated. Also, a therapeutic benefit is achieved with the eradication or amelioration of one or more of the physiological symptoms associated with the underlying disorder such that an improvement is observed in the patient, notwithstanding that the patient may still be afflicted with the underlying disorder. For prophylactic benefit, the compositions may be administered to a patient at risk of developing a particular disease, or to a patient reporting one or more of the physiological symptoms of a disease, even though a diagnosis of this disease may not have been made.

[0054] “Prodrug” is meant to indicate a compound that may be converted under physiological conditions or by solvolysis to a biologically active compound described herein. Thus, the term “prodrug” refers to a precursor of a biologically active compound that is pharmaceutically acceptable. A prodrug may be inactive when administered to a subject, but is converted *in vivo* to an active compound, for example, by hydrolysis. The prodrug compound often offers advantages of solubility, tissue compatibility or delayed release in a mammalian organism (*see, e.g.*, Bundgard, H., Design of Prodrugs (1985), pp. 7-9, 21-24 (Elsevier, Amsterdam).

[0055] A discussion of prodrugs is provided in Higuchi, T., et al., “Pro-drugs as Novel Delivery Systems,” A.C.S. Symposium Series, Vol. 14, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated in full by reference herein.


[0056] The term “prodrug” is also meant to include any covalently bonded carriers, which release the active compound *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of an active compound, as described herein, may be prepared by modifying functional groups present in the active compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent active compound. Prodrugs include compounds wherein a hydroxy, amino or mercapto group is bonded to any group that, when the prodrug of the active compound is administered to a mammalian subject, cleaves to form a free hydroxy, free amino or free mercapto group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol or amine functional groups in the active compounds and the like.

Substituted Imidazole-Pyridine Derivative Compounds

[0057] Substituted imidazole-pyridine derivative compounds are described herein that inhibit a histone demethylase enzyme. These compounds, and compositions comprising these

compounds, are useful for the treatment of cancer and neoplastic diseases. The compounds described herein may, therefore, be useful for treating pancreatic cancer, prostate cancer, breast cancer, bladder cancer, lung cancer, gastric cancer, leukemia and/or melanoma and the like.

[0058] One embodiment provides a compound having the structure of Formula (I),



or a pharmaceutically acceptable salt thereof, wherein

R^1 is hydrogen, halogen, -OH, -OR⁵, -N(R⁵)₂, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclylalkyl, heterocyclylalkyl, aralkyl, or heteroarylalkyl;

R^2 is alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclylalkyl, heterocyclylalkyl, aralkyl, or heteroarylalkyl;

R^3 is hydrogen, halogen, -OH, -NH₂, -NH(C1-C3alkyl) or C1-C3alkyl;

R^4 is hydrogen, C₁-C₄alkyl, C₁-C₄alkene, C₁-C₄alkyne, halogen, or -CN; and

each R⁵ is independently hydrogen, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclylalkyl, heterocyclylalkyl, aralkyl, or heteroarylalkyl.

[0059] Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R³ is hydrogen.

[0060] Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R² is alkyl. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R² is methyl.

[0061] Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R² is carbocyclyl or carbocyclylalkyl. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R² is -(C₁-C₆alkylene)carbocyclyl. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R² is -(C₁-C₆alkylene)carbocyclyl and the (C₁-C₆alkylene) is a C₁alkylene, or a C₂alkylene.

[0062] Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R² is -(C₁-C₆alkylene)carbocyclyl and the carbocyclyl is 1,2,3,4-tetrahydronaphthyl optionally substituted with one or more groups selected from halogen,

hydroxy, -CN, alkyl, alkoxy, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclalkyl, and heterocyclalkyl.

[0063] Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^2 is heterocyclyl or heterocyclalkyl.

[0064] Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^2 is heteroaryl or heteroarylalkyl.

[0065] Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^2 is aryl or aralkyl. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^2 is aralkyl, and the aralkyl is - $(C_1-C_6\text{alkylene})\text{aryl}$. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^2 is aralkyl, the aralkyl is $-(C_1-C_6\text{alkylene})\text{aryl}$ and the $(C_1-C_6\text{alkylene})$ is $C_1\text{alkylene}$, or a $C_2\text{alkylene}$.

[0066] Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^2 is aryl or aralkyl, and the aryl is a phenyl optionally substituted with one or more groups selected from halogen, hydroxy, -CN, alkyl, alkoxy, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclalkyl, and heterocyclalkyl.


[0067] Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^2 is aryl or aralkyl, and the aryl is naphthyl optionally substituted with one or more groups selected from halogen, hydroxy, -CN, alkyl, alkoxy, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclalkyl, and heterocyclalkyl.

[0068] Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^4 is hydrogen, halogen, -CN, or an alkyl optionally substituted with at least one fluoro. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^4 is hydrogen. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^4 is fluoro. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^4 is chloro. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^4 is iodo. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^4 is -CN. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^4 is $-\text{CF}_3$.

[0069] Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^1 is hydrogen or aryl. Another embodiment provides a compound

of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^1 is hydrogen. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^1 is aryl. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^1 is aryl and the aryl is a phenyl optionally substituted with one or more groups selected from halogen, hydroxy, -CN, alkyl, alkoxy, -O-(cycloalkylalkyl), alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclalkyl, and heterocyclalkyl. Another embodiment provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^1 is aryl, and the aryl is a phenyl is substituted with one or more groups selected from halogen, alkoxy, or -O-(cycloalkylalkyl).

[0070] One embodiment provides a compound having the structure of Formula (Ia),



or a pharmaceutically acceptable salt thereof, wherein

R^1 is aryl, or heteroaryl; wherein R^1 is optionally substituted with halogen, -OH, -CN, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclalkyl, heterocyclalkyl, aralkyl, heteroarylalkyl, alkyl-O-, carbocycl-O-, heterocycl-O-, aryl-O-, heteroaryl-O-, carbocyclalkyl-O-, heterocyclalkyl-O-, aralkyl-O-, or heteroarylalkyl-O-;

R^2 is methyl;

R^3 is hydrogen; and


R^4 is hydrogen, C_1 - C_4 alkyl, C_1 - C_4 alkene, C_1 - C_4 alkyne, halogen, or -CN.

[0071] Another embodiment provides the compound of Formula (Ia), or a pharmaceutically acceptable salt thereof, wherein R^4 is C_1 - C_4 alkyl. Another embodiment provides the compound of Formula (Ia), or a pharmaceutically acceptable salt thereof, wherein R^4 is C_1 - C_4 alkyl, and the alkyl is substituted with at least one fluoro. Another embodiment provides the compound of Formula (Ia), or a pharmaceutically acceptable salt thereof, wherein R^4 is CH_2F , CHF_2 , or CF_3 . Another embodiment provides the compound of Formula (Ia), or a pharmaceutically acceptable salt thereof, wherein R^4 is CF_3 .

[0072] Another embodiment provides the compound of Formula (Ia), or a pharmaceutically acceptable salt thereof, wherein R^1 is aryl. Another embodiment provides the compound of Formula (Ia), or a pharmaceutically acceptable salt thereof, wherein R^1 is heteroaryl. Another

embodiment provides the compound of Formula (Ia), or a pharmaceutically acceptable salt thereof, wherein R^1 is aryl, and the aryl is optionally substituted with halogen, alkyl-O-, carbocyclyl-O-, heterocyclyl-O-, aryl-O-, heteroaryl-O-, carbocyclylalkyl-O-, heterocyclylalkyl-O-, aralkyl-O-, or heteroarylalkyl-O-. Another embodiment provides the compound of Formula (Ia), or a pharmaceutically acceptable salt thereof, wherein R^1 is aryl, and the aryl is optionally substituted with halogen, alkyl-O-, carbocyclyl-O-, heterocyclyl-O-, aryl-O-, heteroaryl-O-, carbocyclylalkyl-O-, heterocyclylalkyl-O-, aralkyl-O-, or heteroarylalkyl-O-.

[0073] One embodiment provides a compound having the structure of Formula (Ib),



or a pharmaceutically acceptable salt thereof, wherein

R^1 is hydrogen;

R^2 is chosen from $-(C_1-C_6\text{alkylene})-(1,2,3,4\text{-tetrahydronaphthyl})$, $-(C_1-C_6\text{alkylene})-(\text{indanyl})$, $-(C_1-C_6\text{alkylene})-(\text{chromanyl})$, $-(C_1-C_6\text{alkylene})-(\text{dihydrobenzofuranyl})$

R^3 is hydrogen; and

R^4 is hydrogen, $C_1-C_4\text{alkyl}$, $C_1-C_4\text{alkene}$, $C_1-C_4\text{alkyne}$, halogen, or $-\text{CN}$.

[0074] Another embodiment provides the compound of Formula (Ib), or a pharmaceutically acceptable salt thereof, wherein R^2 is chosen from $-(C_1-C_6\text{alkylene})-(1,2,3,4\text{-tetrahydronaphthyl})$, optionally substituted with one or more groups selected from halogen, hydroxy, $-\text{CN}$, alkyl, alkoxy, alkylamino, aryl, carbocyclyl, carbocyclyloxy, heterocyclyl, carbocyclylalkyl, carbocyclylalkyloxy, and heterocyclylalkyl.

[0075] Another embodiment provides the compound of Formula (Ib), or a pharmaceutically acceptable salt thereof, wherein R^2 is chosen from $-(C_1-C_6\text{alkylene})-(\text{indanyl})$, optionally substituted with one or more groups selected from halogen, hydroxy, $-\text{CN}$, alkyl, alkoxy, alkylamino, aryl, carbocyclyl, carbocyclyloxy, heterocyclyl, carbocyclylalkyl, carbocyclylalkyloxy, and heterocyclylalkyl.


[0076] Another embodiment provides the compound of Formula (Ib), or a pharmaceutically acceptable salt thereof, wherein R^2 is chosen from $-(C_1-C_6\text{alkylene})-(\text{chromanyl})$, optionally substituted with one or more groups selected from halogen, hydroxy, $-\text{CN}$, alkyl, alkoxy,

alkylamino, aryl, carbocyclyl, carbocycloxy, heterocyclyl, carbocyclalkyl, carbocyclalkyloxy, and heterocyclalkyl.

[0077] Another embodiment provides the compound of Formula (Ib), or a pharmaceutically acceptable salt thereof, wherein R^2 is chosen from $-(C_1-C_6\text{alkylene})-(\text{dihydrobenzofuranyl})$, optionally substituted with one or more groups selected from halogen, hydroxy, $-\text{CN}$, alkyl, alkoxy, alkylamino, aryl, carbocyclyl, carbocycloxy, heterocyclyl, carbocyclalkyl, carbocyclalkyloxy, and heterocyclalkyl.

[0078] Another embodiment provides the compound of Formula (Ib), or a pharmaceutically acceptable salt thereof, wherein R^4 is C_1-C_4 alkyl. Another embodiment provides the compound of Formula (Ib), or a pharmaceutically acceptable salt thereof, wherein R^4 is C_1-C_4 alkyl, and the alkyl is substituted with at least one fluoro. Another embodiment provides the compound of Formula (Ib), or a pharmaceutically acceptable salt thereof, wherein R^4 is CH_2F , CHF_2 , or CF_3 . Another embodiment provides the compound of Formula (Ib), or a pharmaceutically acceptable salt thereof, wherein R^4 is CF_3 .

[0079] One embodiment provides a compound having the structure of Formula (Ic),



or a pharmaceutically acceptable salt thereof, wherein

R^1 is hydrogen, halogen, $-\text{OH}$, $-\text{OR}^5$, $-\text{N}(\text{R}^5)_2$, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclalkyl, heterocyclalkyl, aralkyl, or heteroarylalkyl;

R^2 is alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclalkyl, heterocyclalkyl, aralkyl, or heteroarylalkyl;

R^3 is hydrogen;

R^4 is hydrogen, C_1-C_4 alkyl, C_1-C_4 alkene, C_1-C_4 alkyne, halogen, or $-\text{CN}$; and

each R^5 is independently hydrogen, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclalkyl, heterocyclalkyl, aralkyl, or heteroarylalkyl.

[0080] Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^2 is alkyl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^2 is methyl.

[0081] Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^2 is carbocyclyl or carbocyclylalkyl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^2 is $-(C_1-C_6\text{alkylene})\text{carbocyclyl}$. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein carbocyclyl is 1,2,3,4-tetrahydronaphthyl optionally substituted with one or more groups selected from halogen, hydroxy, -CN, alkyl, alkoxy, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein the $(C_1-C_6\text{alkylene})$ is a $C_1\text{alkylene}$, or a $C_2\text{alkylene}$. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^2 is heterocyclyl or heterocyclylalkyl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^2 is heteroaryl or heteroarylalkyl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^2 is aryl or aralkyl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein the aralkyl is $-(C_1-C_6\text{alkylene})\text{aryl}$.


[0082] Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein the aryl is a phenyl optionally substituted with one or more groups selected from halogen, hydroxy, -CN, alkyl, alkoxy, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein $(C_1-C_6\text{alkylene})$ is $C_1\text{alkylene}$, or a $C_2\text{alkylene}$. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein aryl is naphthyl optionally substituted with one or more groups selected from halogen, hydroxy, -CN, alkyl, alkoxy, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein $(C_1-C_6\text{alkylene})$ is $C_1\text{alkylene}$, or a $C_2\text{alkylene}$.

[0083] Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^1 is hydrogen or aryl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^1 is hydrogen. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^1 is aryl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein the aryl is a phenyl optionally substituted with one or more groups selected from halogen, hydroxy, -CN, alkyl, alkoxy, -O-(cycloalkylalkyl),

alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclalkyl, and heterocyclalkyl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein the phenyl is substituted with one or more groups selected from halogen, alkoxy, or $-O-(\text{cycloalkylalkyl})$.

[0084] Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^4 is hydrogen. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^4 is $-\text{CN}$. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^4 is $\text{C}_1\text{-C}_4$ alkene, or $\text{C}_1\text{-C}_4$ alkyne. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^4 is halogen. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^4 is $\text{C}_1\text{-C}_4$ alkyl. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^4 is $\text{C}_1\text{-C}_4$ alkyl, and the alkyl is substituted with at least one fluoro. Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^4 is CH_2F , CHF_2 , or CF_3 . Another embodiment provides the compound of Formula (Ic), or a pharmaceutically acceptable salt thereof, wherein R^4 is CF_3 .

[0085] One embodiment provides a compound having the structure of Formula (II),



or a pharmaceutically acceptable salt thereof, wherein

R^1 is hydrogen, halogen, $-\text{OH}$, $-\text{OR}^4$, $-\text{N}(\text{R}^4)_2$, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclalkyl, heterocyclalkyl, aralkyl, or heteroarylalkyl;

R^2 is alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclalkyl, heterocyclalkyl, aralkyl, or heteroarylalkyl;

R^3 is hydrogen, halogen, $-\text{OH}$, $-\text{NH}_2$, $-\text{NH}(\text{C}_1\text{-C}_3\text{alkyl})$ or $\text{C}_1\text{-C}_3\text{alkyl}$; and

each R^4 is independently hydrogen, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclalkyl, heterocyclalkyl, aralkyl, or heteroarylalkyl.

[0086] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^3 is hydrogen.

[0087] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is alkyl. Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is methyl.

[0088] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is carbocyclyl or carbocyclylalkyl. Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is $-(C_1-C_6\text{alkylene})\text{carbocyclyl}$. Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is $-(C_1-C_6\text{alkylene})\text{carbocyclyl}$, and the $(C_1-C_6\text{alkylene})$ is a $C_1\text{alkylene}$, or a $C_2\text{alkylene}$.

[0089] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is $-(C_1-C_6\text{alkylene})\text{carbocyclyl}$, and the carbocyclyl is 1,2,3,4-tetrahydronaphthyl optionally substituted with one or more groups selected from halogen, hydroxy, $-\text{CN}$, alkyl, alkoxy, $-\text{O}-(\text{cycloalkylalkyl})$, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl.

[0090] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is heterocyclyl or heterocyclylalkyl. Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is $-(C_1-C_6\text{alkylene})\text{heterocyclyl}$.

[0091] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is heteroaryl or heteroarylalkyl.

[0092] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is heteroaryl or heteroarylalkyl, and the heteroaryl is pyridine or pyrimidine optionally substituted with one or more groups selected from halogen, hydroxy, $-\text{CN}$, alkyl, alkoxy, $-\text{O}-(\text{cycloalkylalkyl})$, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl.

[0093] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is heteroaryl, and the heteroaryl is a chromanyl optionally substituted with one or more groups selected from halogen, hydroxy, $-\text{CN}$, alkyl, alkoxy, $-\text{O}-(\text{cycloalkylalkyl})$, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl.

[0094] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^2 is heteroarylalkyl, and the heteroarylalkyl comprises a chromanyl optionally substituted with one or more groups selected from halogen, hydroxy, $-\text{CN}$,

alkyl, alkoxy, -O-(cycloalkylalkyl), alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl; and a C₁alkylene, or a C₂alkylene.

[0095] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R² is aryl or aralkyl.

[0096] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R² is aryl or aralkyl, and the aryl is a phenyl optionally substituted with one or more groups selected from halogen, hydroxy, -CN, alkyl, alkoxy, -O-(cycloalkylalkyl), alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl. Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R² is aryl or aralkyl, and the aryl is phenyl optionally substituted with one or more groups selected from halogen, alkoxy, and alkyl.

[0097] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R² is aralkyl, and the aralkyl is a -(C₁-C₆alkylene)aryl.

[0098] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R² is aralkyl, the aralkyl is a -(C₁-C₆alkylene)aryl, and the aryl is a phenyl optionally substituted with one or more groups selected from halogen, hydroxy, -CN, alkyl, alkoxy, -O-(cycloalkylalkyl), alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl.

[0099] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R² is aralkyl, the aralkyl is a -(C₁-C₆alkylene)aryl, and the (C₁-C₆alkylene) is a C₁alkylene, or a C₂alkylene.

[0100] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R² is aralkyl, and the aralkyl comprises a naphthyl optionally substituted with one or more groups selected from halogen, hydroxy, -CN, alkyl, alkoxy, -O-(cycloalkylalkyl), alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl. Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein the aralkyl further comprises a C₁alkylene, or a C₂alkylene.




[0101] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R¹ is hydrogen or aryl. Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R¹ is hydrogen.






[0102] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R¹ is aryl.






[0103] Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^1 is aryl, and the aryl is a phenyl optionally substituted with one or more groups selected from halogen, hydroxy, -CN, alkyl, alkoxy, -O-(cycloalkylalkyl), alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl. Another embodiment provides a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^1 is aryl, and the aryl is a phenyl optionally substituted with one or more groups selected from halogen, alkoxy, or -O-(cycloalkylalkyl).





[0104] In some embodiments, the compound disclosed herein has the structure provided in Table 1.






TABLE 1






Chemical Synthesis Example	Structure	Name
1		2-(1-methylimidazol-4-yl)-4-(1H-triazol-4-yl)pyridine
2		2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
3		2-[5-(4-fluorophenyl)-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine





Chemical Synthesis Example	Structure	Name
4		2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
5		2-[1-(1-phenylethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
6		2-[1-[2-(2-chlorophenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
7		2-[1-[2-(2-methoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
8		2-[1-(1,2,3,4-tetrahydronaphthalen-1-ylmethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine





Chemical Synthesis Example	Structure	Name
9		2-[1-[2-(2-ethoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
10		4-(1H-triazol-4-yl)-2-[1-[2-[2-(2,2,2-trifluoroethoxy)phenyl]ethyl]imidazol-4-yl]pyridine
11		2-[1-[2-[2-(cyclopropylmethoxy)phenyl]ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
12		2-[1-(3,4-dihydro-2H-chromen-4-ylmethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
13		4-[2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile





Chemical Synthesis Example	Structure	Name
14		4-[2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile
15		4-[2-[1-(1,2,3,4-tetrahydronaphthalen-1-ylmethyl)imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile
16		4-[2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile
17		2-[1-[2-(2-phenylmethoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine






Chemical Synthesis Example	Structure	Name
18		2-[1-[2-(2-phenoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
19		2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-(5-iodo-1H-triazol-4-yl)pyridine
20		2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-(5-fluoro-1H-triazol-4-yl)pyridine
21		2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
22		2-[1-[[2-chloro-3-(trifluoromethyl)phenyl]methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine






Chemical Synthesis Example	Structure	Name
23		2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
24		2-[5-(4-fluoro-3-methoxyphenyl)-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
25		2-[5-(3-ethoxy-4-fluorophenyl)-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
26		2-[1-[[2-fluoro-3-(trifluoromethoxy)phenyl]methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
27		2-[1-[2-(2-phenylphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine

Chemical Synthesis Example	Structure	Name
28		2-[1-[(2-fluoro-3-methylphenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
29		2-[1-[(3-chloro-2-fluorophenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
30		2-[1-[(2-fluoro-3-methoxyphenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
31		4-(1H-triazol-4-yl)-2-[1-[2-[2-(trifluoromethyl)phenyl]ethyl]imidazol-4-yl]pyridine

Chemical Synthesis Example	Structure	Name
32		2-[1-[2-(2-chlorophenyl)-2-methylpropyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
33		2-[1-(1-phenylpropan-2-yl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine
34		4-(1H-triazol-4-yl)-2-[1-[2-[2-(trifluoromethoxy)phenyl]ethyl]imidazol-4-yl]pyridine
35		4-(5-fluoro-1H-triazol-4-yl)-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine

Chemical Synthesis Example	Structure	Name
36		4-(5-chloro-1H-triazol-4-yl)-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine
37		2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-triazol-4-yl]pyridine
38		4-(5-iodo-1H-triazol-4-yl)-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine
39		4-[2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile






Chemical Synthesis Example	Structure	Name
40		4-(5-chloro-1H-triazol-4-yl)-2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridine
41		2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-(5-fluoro-1H-triazol-4-yl)pyridine
42		1-(cyclopropylmethyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole
43		4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1-(2,2,2-trifluoroethyl)-1H-imidazole
44		1-benzyl-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole






Chemical Synthesis Example	Structure	Name
45		1-[(2-chlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole
46		1-[(3-chlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole
47		1-[(4-chlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole
48		1-[(3,4-dichlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole
49		1-(4-chlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole






Chemical Synthesis Example	Structure	Name
50		1-(2-chlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole
51		1-(3-chlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole
52		1-(3,5-dichlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole
53		5-(4-fluorophenyl)-1-methyl-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole
54		5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methyl-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole
55		2-(5-bromo-1-(2-chlorobenzyl)-1H-imidazol-4-yl)-4-(2H-1,2,3-triazol-4-yl)pyridine






Chemical Synthesis Example	Structure	Name
56		2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-(5-(trifluoromethyl)-1H-triazol-4-yl)pyridine
57		2-{1-[2-(2-chlorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
58		2-{1-[(1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
59		2-{1-[2-(2-ethoxyphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
60		4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-(1-{2-[2-(trifluoromethyl)phenyl]ethyl}-1H-imidazol-4-yl)pyridine

Chemical Synthesis Example	Structure	Name
61		2-{1-[(4-fluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
62		2-{1-[(4-fluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
63		2-{1-[(4-chlorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
64		2-{1-[(2-chlorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
65		2-{1-[(4-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine

Chemical Synthesis Example	Structure	Name
66		2-{1-[(2-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
67		2-{1-[(4-methoxyphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
68		2-{1-[(2,4-difluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
69		2-{1-[(2,6-difluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
70		2-{1-[(4-fluoro-2-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine







Chemical Synthesis Example	Structure	Name
71		2-{1-[(2-fluoro-4-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
72		2-(1-benzyl-1H-imidazol-4-yl)-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
73		2-[1-(1-phenylethyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
74		2-[1-(1-phenylpropyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
75		2-[1-(2-methyl-1-phenylpropyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine

Chemical Synthesis Example	Structure	Name
76		2-(1-{[2-(2,2,2-trifluoroethoxy)phenyl]methyl}-1H-imidazol-4-yl)-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
77		2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
78		2-{1-[2-(3-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
79		2-{1-[2-(2,3-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
80		2-{1-[2-(2-chloro-6-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine



Chemical Synthesis Example	Structure	Name
81		4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-{1-[2-(2,4,5-trifluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine
82		4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-{1-[2-(2,4,6-trifluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine
83		2-{1-[(2,3-dihydro-1H-inden-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
84		2-{1-[2-(2-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
85		2-[1-(6-fluoro-2,3-dihydro-1H-inden-1-yl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine

Chemical Synthesis Example	Structure	Name
86		2-{1-[2-(2,4-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
87		2-{1-[2-(2,6-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
88		2-{1-[2-(2,5-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
89		2-{1-[(3R)-1-benzoylpiperidin-3-yl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
90		2-{1-[(3S)-1-benzoylpiperidin-3-yl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine

Chemical Synthesis Example	Structure	Name
91		2-{1-[2-(4-methylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
92		2-{1-[2-(4-chlorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
93		2-{1-[2-(4-methoxyphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
94		4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-(1-{2-[4-(trifluoromethyl)phenyl]ethyl}-1H-imidazol-4-yl)pyridine
95		2-{1-[2-(4-ethylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine





Chemical Synthesis Example	Structure	Name
96		2-{1-[2-(4-tert-butylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
97		2-[5-(4-fluorophenyl)-1-methyl-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
98		2-{1-[(3,4-dihydro-2H-1-benzopyran-4-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
99		2-{1-[(5-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
100		2-{1-[(6-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
101		2-{1-[(7-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine


Chemical Synthesis Example	Structure	Name
102		2-{1-[(6-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
103		2-{1-[(7-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
104		2-{1-[(5-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
105		2-{1-[(6-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
106		2-{1-[(7-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine
107		2-{1-[(8-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine


Chemical Synthesis Example	Structure	Name
108		4-(5-fluoro-1H-1,2,3-triazol-4-yl)-2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine
109		4-(5-chloro-1H-1,2,3-triazol-4-yl)-2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine


[0105] In some embodiments, the compound disclosed herein has the structure provided in Table 2, where R=CF₃, F or Cl.











TABLE 2


	
	











	
	
	
	
	
















































	
	
	
	
	

















	
	
	
	
	
	





	
	
	
	
	
	




























	
	
	
	
	






























Preparation of the Substituted Imidazole-Pyridine Derivative Compounds

[0106] The compounds used in the reactions described herein are made according to organic synthesis techniques known to those skilled in this art, starting from commercially available chemicals and/or from compounds described in the chemical literature. “Commercially available chemicals” are obtained from standard commercial sources including Acros Organics (Pittsburgh, PA), Aldrich Chemical (Milwaukee, WI, including Sigma Chemical and Fluka), Apin Chemicals Ltd. (Milton Park, UK), Avocado Research (Lancashire, U.K.), BDH Inc. (Toronto, Canada), Bionet (Cornwall, U.K.), Chemservice Inc. (West Chester, PA), Crescent Chemical Co. (Hauppauge, NY), Eastman Organic Chemicals, Eastman Kodak Company (Rochester, NY), Fisher Scientific Co. (Pittsburgh, PA), Fisons Chemicals (Leicestershire, UK), Frontier Scientific (Logan, UT), ICN Biomedicals, Inc. (Costa Mesa, CA), Key Organics (Cornwall, U.K.), Lancaster Synthesis (Windham, NH), Maybridge Chemical Co. Ltd. (Cornwall, U.K.), Parish Chemical Co. (Orem, UT), Pfaltz & Bauer, Inc. (Waterbury, CN), Polyorganix (Houston, TX), Pierce Chemical Co. (Rockford, IL), Riedel de Haen AG (Hanover, Germany), Spectrum Quality Product, Inc. (New Brunswick, NJ), TCI America (Portland, OR), Trans World Chemicals, Inc. (Rockville, MD), and Wako Chemicals USA, Inc. (Richmond, VA).


[0107] Methods known to one of ordinary skill in the art are identified through various reference books and databases. Suitable reference books and treatise that detail the synthesis of reactants useful in the preparation of compounds described herein, or provide references to articles that describe the preparation, include for example, “Synthetic Organic Chemistry”, John Wiley & Sons, Inc., New York; S. R. Sandler et al., “Organic Functional Group Preparations,” 2nd Ed., Academic Press, New York, 1983; H. O. House, “Modern Synthetic Reactions”, 2nd Ed., W. A. Benjamin, Inc. Menlo Park, Calif. 1972; T. L. Gilchrist, “Heterocyclic Chemistry”, 2nd Ed., John Wiley & Sons, New York, 1992; J. March, “Advanced Organic Chemistry: Reactions, Mechanisms and Structure”, 4th Ed., Wiley-Interscience, New York, 1992. Additional suitable reference books and treatise that detail the synthesis of reactants useful in the preparation of compounds described

herein, or provide references to articles that describe the preparation, include for example, Fuhrhop, J. and Penzlin G. "Organic Synthesis: Concepts, Methods, Starting Materials", Second, Revised and Enlarged Edition (1994) John Wiley & Sons ISBN: 3-527-29074-5; Hoffman, R.V. "Organic Chemistry, An Intermediate Text" (1996) Oxford University Press, ISBN 0-19-509618-5; Larock, R. C. "Comprehensive Organic Transformations: A Guide to Functional Group Preparations" 2nd Edition (1999) Wiley-VCH, ISBN: 0-471-19031-4; March, J. "Advanced Organic Chemistry: Reactions, Mechanisms, and Structure" 4th Edition (1992) John Wiley & Sons, ISBN: 0-471-60180-2; Otera, J. (editor) "Modern Carbonyl Chemistry" (2000) Wiley-VCH, ISBN: 3-527-29871-1; Patai, S. "Patai's 1992 Guide to the Chemistry of Functional Groups" (1992) Interscience ISBN: 0-471-93022-9; Solomons, T. W. G. "Organic Chemistry" 7th Edition (2000) John Wiley & Sons, ISBN: 0-471-19095-0; Stowell, J.C., "Intermediate Organic Chemistry" 2nd Edition (1993) Wiley-Interscience, ISBN: 0-471-57456-2; "Industrial Organic Chemicals: Starting Materials and Intermediates: An Ullmann's Encyclopedia" (1999) John Wiley & Sons, ISBN: 3-527-29645-X, in 8 volumes; "Organic Reactions" (1942-2000) John Wiley & Sons, in over 55 volumes; and "Chemistry of Functional Groups" John Wiley & Sons, in 73 volumes.

[0108] Specific and analogous reactants may also be identified through the indices of known chemicals prepared by the Chemical Abstract Service of the American Chemical Society, which are available in most public and university libraries, as well as through on-line databases (the American Chemical Society, Washington, D.C., may be contacted for more details). Chemicals that are known but not commercially available in catalogs may be prepared by custom chemical synthesis houses, where many of the standard chemical supply houses (*e.g.*, those listed above) provide custom synthesis services. A reference for the preparation and selection of pharmaceutical salts of the substituted imidazole-pyridine derivative compounds described herein is P. H. Stahl & C. G. Wermuth "Handbook of Pharmaceutical Salts", Verlag Helvetica Chimica Acta, Zurich, 2002.


[0109] The substituted 4-triazolylpyridine derivative compounds are prepared by the general synthetic route described below in Scheme 1-4.

Scheme 1




[0110] Referring to Scheme 1, 2-chloro-4-pyridinenitrile (1-0) undergoes Stille coupling with tin reagent, N-methyl-4-(tributylstannyl)imidazole or trityl protected tributyltin imidazole, in presence of a catalyst, such as tetrakis(triphenylphosphine) palladium (0), under heating condition, i.e., 130 °C, to give the coupling products 1-1 and 1-2. Deprotection of the trityl protective group is performed using acids, such as acetic acid and TFA, at room temperature to give compound 1-3, which is then alkylated with halogenated alkyl derivative or mesylated derivative to give 1-4. Reaction of the nitrile intermediates 1-1 and 1-4 with TMS diazomethane, in presence of n-butyl lithium, in an anhydrous organic solvent, such as THF, provides TMS protected 4-triazolopyridine intermediate 1-5, which undergoes deprotection using a base, such as NaOH, in an organic solvent, such as MeOH, under heating condition (i.e. 50 °C) to give the final product, compound 1-6.

Scheme 2




[0111] Synthesis of substituted 2-imidazole 4-triazolopyridine derivatives is described in Scheme 2. 2-(1-Methyl-1H-imidazol-4-yl)pyridine-4-carbonitrile is brominated by using stoichiometric amount of NBS in an organic solvent, such as methylene chloride, to give brominated intermediate 2-1. Compound 2-1 then undergoes Suzuki coupling with an aromatic boronic acid or ester to give 2-2. The nitrile group is then hydrolyzed to the acid (2-3) by refluxing in a solvent mixture of concentrated NaOH (i.e. 5N or 6N) and EtOH. The acid is then reduced to the alcohol (2-4) by using LiAlH_4 in an organic solvent, such as THF. The primary alcohol is then treated with Dess-Martin oxidant to provide the aldehyde 2-5, which subsequently reacts with 2-(benzenesulfonyl)acetonitrile in presence of a base, such as NaHCO_3 , in an alcoholic solvent, such as EtOH, to give the intermediate 2-6, which then reacts with NaN_3 in DMF under heating condition, i.e., 100 °C, to afford the 4-triazolopyridine compound 2-7.

Scheme 3



[0112] Referring to Scheme 3, similar to Scheme 1, intermediate 1-4 can also be synthesized from 4-amide-2-imidazole pyridine (3-0). Alkylation using either halogenated RX or mesylate $ROMs$ gives 3-1, which can be converted to nitrile intermediate 1-4 using trifluoroacetic anhydride, in presence of excess of pyridine (i.e. 3 equiv.) in an organic solvent, such as methylene chloride. Final compounds 3-6 can then be obtained using similar procedures as described in Scheme 2.


Scheme 4




[0113] Synthesis of substituted triazole analogs 4-4 is shown in Scheme 4. Starting from the aldehyde 3-7, (1-diazo-2-oxo-propyl)-phosphonic acid dimethyl ester is added to a mixture of the aldehyde and a base, such as potassium carbonate, in an alcoholic solvent, i.e. MeOH, to give the alkyne intermediate (4-1). It is then added to a mixture of PMB protected azide and KI in THF, in presence of a copper complex and a base, such as triethylamine, to give the iodo-triazole intermediate 4-2. Conversion of the iodo to various substitutions can be accomplished by treating 4-2 with KF, KCl or TMSCF₃ in a solvent mixture, such as CH₃CN and H₂O, and heated at elevated temp., i.e. 160 °C, in a microwave oven, followed by deprotection of the PMB group using TFA under heating condition, i.e. 60 °C, to give the final compounds 4-4.

[0114] The substituted azabenzotriazole derivative compounds are prepared by the general synthetic route described below in Scheme 5.

Scheme 5



[0115] Referring to Scheme 5, treatment of diaminopyridine 5-1 with diazotization reagent afforded azabenzotriazole 5-2, which was subsequently protected with a SEM protecting group. Carbonylation of 5-3 in the presence of methanol and Palladium catalyst furnished 5-4 which was reduced to alcohol 5-5 and re-oxidized to corresponding aldehyde 5-6. Treatment with Ts-MIC reagent followed by ammonia workup furnished imidazole 5-7. Alkylation with alkylhalide under basic conditions or coupling with a boronic acid under Chan-Lam reaction conditions afforded 5-8. Removal of SEM protecting group under strongly acidic conditions furnished 5-9. Alternatively, substituted 4-triazolylpyridine derivative compounds could be prepared by the general synthetic route described below in Scheme 6.



[0116] As shown in Scheme 6, 4-iodo-2-(1-methoxyethenyl)pyridine first reacts with the tin reagent, tetrabutyl(3,3,3-trifluoroprop-1-yn-1-yl)stannane, in presence of a catalytic amount of $\text{Pd}(0)$, in toluene, in elevated temperature, i.e. 120°C , in a microwave oven to give 6-2. It is then heated with 1-azidomethyl-4-methoxy-benzene at reflux in an organic solvent, such as t-butyl alcohol or toluene, to give a mixture of PMB protected trifluoromethyl triazole intermediates 6-4a and 6-4b. The mixture is then treated with a reducing agent, such as sodium borohydride, to give the

alcohols, 6-5a and 6-5b. Oxidation by Dess-Martin reagent or MnO_2 gives the aldehyde, 6-6a and 6-6b, which can be separated on flash column. Either the mixture or the separated intermediate is then treated with a mixture of Tos-Mic and KCN in an alcoholic solvent, such as ethanol, at room temperature, for a few hours, followed by evaporation and then treated with NH_3/MeOH in a sealed tube at elevated temperature, i.e. 125°C , for an extended period, such as 20 hr. Isolation by flash column chromatography gives an intermediate 6-7, which is subjected to alkylation with either alkyl bromide or mesylate in presence of an inorganic base, such as K_2CO_3 or Cs_2CO_3 , in DMF, at room temperature or elevated temperature, i.e. 90°C , to give PMB protected product, which is treated with TFA at room temperature overnight or at 50°C for shorter period of time to give the final product 6-8.

[0117] In each of the above reaction procedures or schemes, the various substituents may be selected from among the various substituents otherwise taught herein.

Pharmaceutical Compositions

[0118] In certain embodiments, the substituted imidazole-pyridine derivative compound as described herein is administered as a pure chemical. In other embodiments, the substituted imidazole-pyridine derivative compound described herein is combined with a pharmaceutically suitable or acceptable carrier (also referred to herein as a pharmaceutically suitable (or acceptable) excipient, physiologically suitable (or acceptable) excipient, or physiologically suitable (or acceptable) carrier) selected on the basis of a chosen route of administration and standard pharmaceutical practice as described, for example, in *Remington: The Science and Practice of Pharmacy* (Gennaro, 21st Ed. Mack Pub. Co., Easton, PA (2005)), the disclosure of which is hereby incorporated herein by reference in its entirety.

[0119] Accordingly, provided herein is a pharmaceutical composition comprising at least one substituted imidazole-pyridine derivative compound, or a stereoisomer, pharmaceutically acceptable salt, hydrate, solvate, or N-oxide thereof, together with one or more pharmaceutically acceptable carriers. The carrier(s) (or excipient(s)) is acceptable or suitable if the carrier is compatible with the other ingredients of the composition and not deleterious to the recipient (*i.e.*, the subject) of the composition.

[0120] One embodiment provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

[0121] One embodiment provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of Formula (II), or a pharmaceutically acceptable salt thereof.

[0122] In certain embodiments, the substituted imidazole-pyridine derivative compound as described by Formula (I) or (II) is substantially pure, in that it contains less than about 5%, or less than about 1%, or less than about 0.1%, of other organic small molecules, such as contaminating intermediates or by-products that are created, for example, in one or more of the steps of a synthesis method.

[0123] Suitable oral dosage forms include, for example, tablets, pills, sachets, or capsules of hard or soft gelatin, methylcellulose or of another suitable material easily dissolved in the digestive tract. Suitable nontoxic solid carriers can be used which include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like. (*See, e.g., Remington: The Science and Practice of Pharmacy* (Gennaro, 21st Ed. Mack Pub. Co., Easton, PA (2005)).

[0124] The dose of the composition comprising at least one substituted imidazole-pyridine derivative compound as described herein may differ, depending upon the patient's (e.g., human) condition, that is, stage of the disease, general health status, age, and other factors that a person skilled in the medical art will use to determine dose.

[0125] Pharmaceutical compositions may be administered in a manner appropriate to the disease to be treated (or prevented) as determined by persons skilled in the medical arts. An appropriate dose and a suitable duration and frequency of administration will be determined by such factors as the condition of the patient, the type and severity of the patient's disease, the particular form of the active ingredient, and the method of administration. In general, an appropriate dose and treatment regimen provides the composition(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit (*e.g.*, an improved clinical outcome, such as more frequent complete or partial remissions, or longer disease-free and/or overall survival, or a lessening of symptom severity). Optimal doses may generally be determined using experimental models and/or clinical trials. The optimal dose may depend upon the body mass, weight, or blood volume of the patient.

[0126] Oral doses can typically range from about 1.0 mg to about 1000 mg, one to four times, or more, per day.

Histone Demethylase

[0127] Chromatin is the complex of DNA and protein that makes up chromosomes. Histones are the major protein component of chromatin, acting as spools around which DNA winds. Changes in chromatin structure are affected by covalent modifications of histone proteins and by non-histone binding proteins. Several classes of enzymes are known which can covalently modify histones at various sites.

[0128] Proteins can be post-translationally modified by methylation on amino groups of lysines and guanidino groups of arginines or carboxymethylated on aspartate, glutamate, or on the C-terminus of the protein. Post-translational protein methylation has been implicated in a variety of cellular processes such as RNA processing, receptor mediated signaling, and cellular differentiation. Post-translational protein methylation is widely known to occur on histones, such reactions known to be catalyzed by histone methyltransferases, which transfer methyl groups from S-adenyosyl methionine (SAM) to histones. Histone methylation is known to participate in a diverse range of biological processes including heterochromatin formation, X-chromosome inactivation, and transcriptional regulation (Lachner et al., (2003) *J. Cell Sci.* 116:2117-2124; Margueron et al., (2005) *Curr. Opin. Genet. Dev.* 15:163-176).

[0129] Unlike acetylation, which generally correlates with transcriptional activation, whether histone methylation leads to transcription activation or repression depends on the particular site of methylation and the degree of methylation (e.g., whether a particular histone lysine residue is mono-, di-, or tri-methylated). However, generally, methylation on H3K9, H3K27 and H4K20 is linked to gene silencing, while methylation on H3K4, H3K36, and H3K79 is generally associated with active gene expression. In addition, tri- and di-methylation of H3K4 generally marks the transcriptional start sites of actively transcribed genes, whereas mono-methylation of H3K4 is associated with enhancer sequences.

[0130] A “demethylase” or “protein demethylase,” as referred to herein, refers to an enzyme that removes at least one methyl group from an amino acid side chain. Some demethylases act on histones, e.g., act as a histone H3 or H4 demethylase. For example, an H3 demethylase may demethylate one or more of H3K4, H3K9, H3K27, H3K36 and/or H3K79. Alternately, an H4 demethylase may demethylate histone H4K20. Demethylases are known which can demethylate either a mono-, di- and/or a tri-methylated substrate. Further, histone demethylases can act on a methylated core histone substrate, a mononucleosome substrate, a dinucleosome substrate and/or an oligonucleosome substrate, peptide substrate and/or chromatin (e.g., in a cell-based assay).

[0131] The first lysine demethylase discovered was lysine specific demethylase 1 (LSD1/KDM1), which demethylates both mono- and di-methylated H3K4 or H3K9, using flavin as a cofactor. A second class of Jumonji C (JmjC) domain containing histone demethylases were predicted, and confirmed when a H3K36 demethylase was found using a formaldehyde release assay, which was named JmjC domain containing histone demethylase 1 (JHDM1/KDM2A).

[0132] More JmjC domain-containing proteins were subsequently identified and they can be phylogenetically clustered into seven subfamilies: JHDM1, JHDM2, JHDM3, JMJD2, JARID, PHF2/PHF8, UTX/UTY, and JmjC domain only.

FBXL10 and FBXL11

[0133] F-box and leucine-rich repeat protein 10 (FBXL10) and F-box and leucine-rich repeat protein 11 (FBXL11) are multifunctional F-box family proteins that demethylate histone H3 through a hydroxylation based mechanism. FBXL10, also known as lysine (K)-specific demethylase 2B (KDM2B) or Jumonji C domain-containing histone demethylase 1B (JHDM1B), preferentially demethylates trimethylated K4 and dimethylated K36 of histone H3, but contains weak or no activity for mono- and tri-methylated H3K36. FBXL10 contains three domains, a catalytic JMJC domain, an F-box domain and a CXXC DNA-binding domain. The N-terminal JMJC domain coordinates iron and α -ketoglutarate to catalyze demethylation through the hydroxylation based mechanism. The CXXC DNA-binding domain allows FBXL10 to preferentially bind to transcribed region of the ribosomal RNA, leading to repression of the ribosomal RNA gene transcription and ultimately leading to inhibition of cell growth and proliferation. FBXL10 has been found to be overexpressed in acute myeloid leukemia, bladder carcinoma and pancreatic ductal adenocarcinoma. Recently, it has been demonstrated that FBXL10 regulates the expression of Polycomb target genes, those proteins are epigenetic regulators essential for stem cell differentiation. This regulation implicates FBXL10's involvement in tumorigenesis through the regulation of these Polycomb target genes.

[0134] FBXL11, also known as KDM2A or JHDM1A, demethylates mono- and di-methylated K36 of histone H3. The CXXC DNA-binding domain recognizes non-methylated DNA and targets CpG island regions where it specifically removes H3K36 methylation. Further, FBXL11 is required to maintain a heterochromatic state, sustain centromeric integrity and genomic stability during mitosis. In addition, FBXL11 is a key negative regulator of NF- κ B. Overexpression of FBXL11 has been observed in non-small cell lung cancer (NSCLC) where FBXL11 upregulates phosphor-ERK1/2 by repressing DUSP3 expression in NSCLC cell lines. Negative regulation of

gluconeogenic gene expression by FBXL11 results in suppression of two rate-limiting gluconeogenic enzymes, critical for maintaining blood glucose homeostasis.

[0135] In an additional embodiment is a method for inhibiting a histone-demethylase enzyme comprising contacting a histone demethylase enzyme with a compound of Formula (I) or Formula (II).

[0136] In an additional embodiment is the method for inhibiting a histone-demethylase enzyme, wherein the histone-demethylase enzyme comprises a JmjC domain. In an additional embodiment is the method for inhibiting a histone-demethylase enzyme, wherein the histone-demethylase enzyme is selected from FBXL10 or FBXL11.

Methods of Treatment

[0137] Disclosed herein are methods of modulating demethylation in a cell or in a subject, either generally or with respect to one or more specific target genes. Demethylation can be modulated to control a variety of cellular functions, including without limitation: differentiation; proliferation; apoptosis; tumorigenesis, leukemogenesis or other oncogenic transformation events; hair loss; or sexual differentiation. For example, in particular embodiments, the invention provides a method of treating a disease regulated by histone methylation and/or demethylation in a subject in need thereof by modulating the activity of FBXL10 or FBXL11.

[0138] In an additional embodiment is a method for treating cancer in subject comprising administering a composition comprising a compound of Formula (I) or (II), or a pharmaceutically acceptable salt thereof.

[0139] In a further embodiment is the method for treating cancer in a subject wherein the cancer is selected from pancreatic cancer, prostate cancer, breast cancer, gastric cancer, leukemia, bladder cancer, lung cancer or melanoma.

[0140] Other embodiments and uses will be apparent to one skilled in the art in light of the present disclosures. The following examples are provided merely as illustrative of various embodiments and shall not be construed to limit the invention in any way.


EXAMPLES

I. Chemical Synthesis

[0141] Unless otherwise noted, reagents and solvents were used as received from commercial suppliers. Anhydrous solvents and oven-dried glassware were used for synthetic transformations sensitive to moisture and/or oxygen. Yields were not optimized. Reaction times are


approximate and were not optimized. Column chromatography and thin layer chromatography (TLC) were performed on silica gel unless otherwise noted. Spectra are given in ppm (δ) and coupling constants, J are reported in Hertz. For proton spectra the solvent peak was used as the reference peak.

Preparation 1A: 2-(1-methylimidazol-4-yl)pyridine-4-carbonitrile




[0142] A mixture of 2-chloro-4-pyridinenitrile (1.85 g, 13.4 mmol), N-methyl-4-(tributylstannyl)imidazole (5 g, 13.4 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (1.42 g, 1.34 mmol) in DMF (50 mL) was stirred for 3 hr at 130 °C under N_2 . The mixture was concentrated and purified by flash column chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}=20/1$) to afford the title compound (2.0 g, 80%). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{10}\text{H}_8\text{N}_4$, 185; Found, 185.

Preparation 1B: trimethyl-[4-[2-(1-methylimidazol-4-yl)pyridin-4-yl]-1H-triazol-5-yl]silane




[0143] To a solution of TMSCHN_2 (0.76 mL, 1.52 mmol) in THF was added n-BuLi (0.60 mL, 1.52 mmol) at 0 °C, stirred for 20 min, then a solution of 2-(1-methylimidazol-4-yl)pyridine-4-carbonitrile (200 mg, 1.09 mmol) in THF was added, and the reaction mixture was stirred overnight at rt. LC/MS showed the reaction was completed, sat. NH_4Cl was added and extracted with EtOAc, dried, concentrated to give the title compound (296 mg, 90%). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{14}\text{H}_{18}\text{N}_6\text{Si}$, 299; Found, 299.

Example 1: 2-(1-methylimidazol-4-yl)-4-(1H-triazol-4-yl)pyridine




[0144] A mixture of trimethyl-[4-[2-(1-methylimidazol-4-yl)pyridin-4-yl]-1H-triazol-5-yl]silane (296 mg, 0.99 mmol) and NaOH (2.2 mL, 4.46 mmol, 2M) in MeOH was stirred overnight at 50 °C. LC/MS showed the reaction was completed, H₂O was added, extracted with ethyl acetate, dried, purified by HPLC to give the title compound (38 mg, 18%) as a yellow solid. ¹H NMR (300 MHz, CD₃OD): δ 3.99 (3H, s), 7.96 (1H, dd, *J* = 5.4 Hz, *J* = 1.2 Hz), 8.21 (1H, d, *J* = 0.6 Hz), 8.46 (1H, s), 8.47-8.64 (1H, m), 8.65 (1H, s), 8.69 (1H, d, *J* = 5.4 Hz). [M+H] Calc'd for C₁₁H₁₀N₆, 227; Found, 227.

Preparation 2A: 2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridine-4-carboxamide



[0145] A mixture of 2-(1H-imidazol-4-yl)pyridine-4-carboxamide (1 g, 5.32 mmol), 1-bromomethyl-2-chloro-benzene (2.17 g, 10.64 mmol) and K₂CO₃ (1.47 g, 10.64 mmol) in DMF (50 mL) was stirred overnight at rt. LC/MS showed the reaction was completed. The mixture was concentrated and purified by flash column chromatography on silica gel (CH₂Cl₂/MeOH=20/1) to afford the title compound (1.2 g, 66%). [M+H] Calc'd for C₁₆H₁₃ClN₄O, 313; Found, 313.


Preparation 2B: 2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridine-4-carbonitrile



[0146] To a solution of 2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridine-4-carboxamide (1.2 g, 3.54 mmol) and pyridine (0.91 g, 10.64 mmol) in CH₂Cl₂ was added trifluoroacetic acid anhydride (1.62 g, 7.08 mmol) at 0°C, which was then stirred for 2 hr at 0°C.


LC/MS showed the reaction was completed. H₂O was added and extracted with CH₂Cl₂, the organic layer was washed by aqueous NaHCO₃ and brine, dried, concentrated and purified by flash column chromatography on silica gel (PE/EA=1/3) to give the title compound (844 mg, 74%). [M+H] Calc'd for C₁₆H₁₁ClN₄, 295; Found, 295.

Preparation 2C: [4-[2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridin-4-yl]-1H-triazol-5-yl]-trimethylsilane




[0147] The title compound was prepared in 98% yield according to the procedure of Preparation 1B. [M+H] Calc'd for C₂₀H₂₁ClN₆Si, 409; Found, 409.

Example 2: 2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0148] The title compound was prepared in 44% yield according to the procedure of last step of Example 1. ¹H NMR (300 MHz, CD₃OD): δ 5.58 (2H, s), 7.44-7.54 (4H, m), 8.03 (1H, d, *J* = 5.7 Hz), 8.22 (1H, s), 8.55-8.65 (4H, m). [M+H] Calc'd for C₁₇H₁₃ClN₆, 337; Found, 337.

Preparation 3A: 2-(5-bromo-1-methylimidazol-4-yl)pyridine-4-carbonitrile




[0149] To a solution of 2-(1-methylimidazol-4-yl)pyridine-4-carbonitrile (600 mg, 3.26 mmol, Preparation 1A) in CH₂Cl₂ (30 mL), was added NBS (610 mg, 3.42 mmol) and it was stirred for 3 hr at rt. Washed by H₂O, dried and concentrated to afford the title compound (810 mg, 95%). [M+H] Calc'd for C₁₀H₇BrN₄, 263; Found, 263.

Preparation 3B: 2-[5-(4-fluorophenyl)-1-methylimidazol-4-yl]pyridine-4-carbonitrile




[0150] A mixture of 2-(5-bromo-1-methylimidazol-4-yl)pyridine-4-carbonitrile (1 eq), 4-fluorophenylboronic acid (1.2 eq), Pd(dppf)Cl₂ (0.1 eq) and 2M Na₂CO₃ (2 eq) in dioxane was refluxed overnight under N₂, concentrated and purified by flash column chromatography (DCM/MeOH =20/1) to give the title compound (140 mg, 50%). [M+H]⁺ Calc'd for C₁₆H₁₁FN₄, 279; Found, 279.

Preparation 3C: [4-[2-[5-(4-fluorophenyl)-1-methylimidazol-4-yl]pyridin-4-yl]-1H-triazol-5-yl]-trimethylsilane




[0151] The title compound was prepared in 87% yield according to the procedure of Preparation 1B. [M+H]⁺ Calc'd for C₂₀H₂₁FN₆Si, 393; Found, 393.

Example 3: 2-[5-(4-fluorophenyl)-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0152] The title compound was prepared in 12% yield according to the procedure of last step of Example 1. ¹H NMR (300 MHz, DMSO + CD₃OD): δ 3.61 (3H, s), 7.44-7.85 (6H, m), 8.30 (1H, s), 8.64 (1H, d, *J* = 5.4 Hz), 8.84-8.85 (1H, m). [M+H]⁺ Calc'd for C₁₇H₁₃FN₆, 321; Found, 321.

Preparation 4A: 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0153] The title compound was prepared in 25% yield according to the procedure of Preparation 3B. [M+H] Calc'd for C₂₀H₁₇FN₄O, 349; Found, 349.

Preparation 4B: [4-[2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridin-4-yl]-1H-triazol-5-yl]-trimethylsilane




[0154] The title compound was prepared in 25% yield according to the procedure of Preparation 1B. [M+H] Calc'd for C₂₄H₂₇FN₆OSi, 463; Found, 463.

Example 4: 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0155] The title compound was prepared in 13% yield according to the procedure of last step of Example 1. ¹H NMR (300 MHz, DMSO + CD₃OD): δ 0.10-0.36 (4H, m), 0.91-0.98 (1H, m), 3.37 (3H, s), 3.72-3.82 (2H, m), 6.80-6.81 (1H, m), 6.97 (1H, d, *J* = 10.5 Hz), 7.28 (1H, t, *J* = 3.6 Hz), 7.52 (1H, d, *J* = 4.8 Hz), 7.86 (1H, s), 8.25-8.33 (3H, m). [M+H] Calc'd for C₂₁H₁₉FN₆O, 391; Found, 391.

Example 5: 2-[1-(1-phenylethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0156] The title compound was prepared in 6% overall yield according to the procedure of Example 2 starting from 2-(1H-imidazol-4-yl)pyridine-4-carboxamide. ¹H NMR (300 MHz, CD₃OD): δ 1.94 (3H, d, *J* = 14.1 Hz), 5.58-5.65 (1H, m), 7.32-7.44 (5H, m), 7.75 (1H, d, *J* = 5.1 Hz), 7.83 (1H, s), 7.94 (1H, s), 8.39 (2H, s), 8.52 (1H, d, *J* = 5.1 Hz). [M+H] Calc'd for C₁₈H₁₆N₆, 317; Found, 317.

Preparation 6A: 2-[1-[2-(2-chlorophenyl)ethyl]imidazol-4-yl]pyridine-4-carboxamide




[0157] A mixture of 2-(1H-imidazol-4-yl)pyridine-4-carboxamide (500 mg, 2.66 mmol), 1-(2-bromo-ethyl)-2-chloro-benzene (2.32 g, 10.64 mmol) and K₂CO₃ (1.47 g, 10.64 mmol) in DMF (20 mL) was stirred overnight at 100 °C. LC/MS showed the reaction was completed. The reaction mixture was purified by flash column chromatography to give the title compound (300 mg, 34%) as a yellow solid. [M+H] Calc'd for C₁₇H₁₅ClN₄O, 327; Found, 327.

Preparation 6B: 2-[1-[2-(2-chlorophenyl)ethyl]imidazol-4-yl]pyridine-4-carbonitrile




[0158] The title compound was prepared in 77% yield according to the procedure of Preparation 2B. [M+H] Calc'd for C₁₇H₁₃ClN₄, 309; Found, 309.

Preparation 6C: [4-[2-[1-[2-(2-chlorophenyl)ethyl]imidazol-4-yl]pyridin-4-yl]-1H-triazol-5-yl]-trimethylsilane




[0159] The title compound was prepared in 100% yield according to the procedure of Preparation 1B. [M+H] Calc'd for $C_{21}H_{23}ClN_6Si$, 423; Found, 423.

Example 6: 2-[1-[2-(2-chlorophenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0160] The title compound was prepared in 42% yield according to the procedure of last step of Example 1. ^1H NMR (300 MHz, CD_3OD): δ 3.23 (2H, t, $J = 7.2$ Hz), 4.31 (2H, t, $J = 7.2$ Hz), 7.25-7.28 (3H, m), 7.44-7.47 (1H, m), 7.63-7.64 (2H, m), 7.79 (1H, s), 8.30 (1H, s), 8.53 (1H, d, $J = 5.1$ Hz), 8.62 (1H, s). [M+H] Calc'd for $C_{18}H_{15}ClN_6$, 351; Found, 351.

Example 7: 2-[1-[2-(2-methoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0161] The title compound was prepared in 12% overall yield according to the procedure of Example 6 starting from 2-(1H-imidazol-4-yl)pyridine-4-carboxamide. ^1H NMR (300 MHz, CD_3OD): δ 3.14 (2H, t, $J = 6.9$ Hz), 3.82 (3H, s), 4.32 (2H, t, $J = 6.9$ Hz), 6.81-7.22 (4H, m), 7.52 (1H, s), 7.68 (1H, s), 7.73 (1H, d, $J = 5.4$ Hz), 8.32 (1H, s), 8.37 (1H, s), 8.52 (1H, d, $J = 5.4$ Hz). [M+H] Calc'd for $C_{19}H_{18}N_6O$, 347; Found, 347.

Example 8: 2-[1-(1,2,3,4-tetrahydronaphthalen-1-ylmethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0162] The title compound was prepared in 12% overall yield according to the procedure of Example 6 starting from 2-(1H-imidazol-4-yl)pyridine-4-carboxamide. ¹H NMR (300 MHz, DMSO): δ 1.45-1.85 (4H, m), 2.69-2.74 (2H, m), 3.28-3.34 (1H, m), 4.11-4.35 (2H, m), 7.09-7.16 (3H, m), 7.28-7.31 (1H, m), 7.65 (1H, d, *J* = 5.4 Hz), 7.77 (1H, s), 7.88 (1H, s), 8.34 (1H, s), 8.84 (1H, d, *J* = 5.4 Hz), 8.34 (1H, s). [M+H] Calc'd for C₂₁H₂₀N₆, 357; Found, 357.

Preparation 9A: 2-[1-[2-(2-ethoxyphenyl)ethyl]imidazol-4-yl]pyridine-4-carboxamide




[0163] A mixture of 2-(1H-imidazol-4-yl)pyridine-4-carboxamide (1 eq), ROMs (1.2 eq) and K₂CO₃ (2 eq) in DMF (50 mL) was stirred overnight at 80 °C. LC/MS showed the reaction was completed. The mixture was concentrated and purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to afford the title compound. [M+H] Calc'd for C₁₉H₂₀N₄O₂, 337; Found, 337.

Preparation 9B: 2-[1-[2-(2-ethoxyphenyl)ethyl]imidazol-4-yl]pyridine-4-carbonitrile




[0164] The title compound was prepared in 60% yield according to the procedure of Preparation 2B. [M+H] Calc'd for C₁₉H₁₈N₄O, 319; Found, 319.

Preparation 9C: [4-[2-[1-[2-(2-ethoxyphenyl)ethyl]imidazol-4-yl]pyridin-4-yl]-1H-triazol-5-yl]-trimethylsilane




[0165] The title compound was prepared in 100% yield according to the procedure of Preparation 1B. [M+H] Calc'd for $C_{23}H_{28}N_6OSi$, 433; Found, 433.

Example 9: 2-[1-[2-(2-ethoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine



[0166] The title compound was prepared in 54% yield according to the procedure of the last step of Example 1. ^1H NMR (300 MHz, DMSO): δ 1.36 (3H, t, $J = 6.9$ Hz), 3.06 (2H, t, $J = 6.9$ Hz), 4.02 (2H, q, $J = 6.9$ Hz), 4.25 (2H, t, $J = 6.9$ Hz), 6.79-7.22 (4H, m), 7.59 (1H, s), 7.63 (1H, d, $J = 4.2$ Hz), 7.72 (1H, s), 8.30 (1H, s), 8.53 (1H, d, $J = 5.4$ Hz), 8.62 (1H, s). [M+H] Calc'd for $C_{20}H_{20}N_6O$, 361; Found, 361.


Example 10: 4-(1H-triazol-4-yl)-2-[1-[2-[2-(2,2,2-trifluoroethoxy)phenyl]ethyl]imidazole-4-yl]pyridine



[0167] The title compound was prepared in 11% overall yield according to the procedure of Example 9 starting from 2-(1H-imidazol-4-yl)pyridine-4-carboxamide. ^1H NMR (300 MHz, DMSO): δ 3.10 (2H, t, $J = 7.2$ Hz), 4.25 (2H, t, $J = 7.2$ Hz), 4.81 (2H, q, $J = 9.0$ Hz), 6.93-7.26


(4H, m), 7.56 (1H, s), 7.63 (1H, d, $J = 4.5$ Hz), 7.71 (1H, s), 8.30 (1H, s), 8.53 (1H, d, $J = 4.8$ Hz), 8.64 (1H, m). [M+H] Calc'd for $C_{20}H_{17}F_3N_6O$, 415; Found, 415.

Example 11: 2-[1-[2-[2-(cyclopropylmethoxy)phenyl]ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0168] The title compound was prepared in 3% overall yield according to the procedure of Example 9 starting from 2-(1H-imidazol-4-yl)pyridine-4-carboxamide. ^1H NMR (300 MHz, DMSO): δ 0.36-0.37 (2H, m), 0.57-0.60 (2H, m), 1.30-1.35 (1H, m), 3.07 (2H, t, $J = 6.9$ Hz), 3.85 (2H, d, $J = 5.7$ Hz), 4.47 (2H, t, $J = 6.6$ Hz), 6.81-7.18 (4H, m), 7.60-7.63 (2H, m), 7.74 (1H, s), 8.30 (1H, s), 8.52-8.53 (2H, m). [M+H] Calc'd for $C_{22}H_{22}N_6O$, 387; Found, 387.

Example 12: 2-[1-(3,4-dihydro-2H-chromen-4-ylmethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0169] The title compound was prepared in 4% overall yield according to the procedure of Example 9 starting from 2-(1H-imidazol-4-yl)pyridine-4-carboxamide. ^1H NMR (300 MHz, DMSO): δ 1.64-1.83 (2H, m), 3.30-3.33 (1H, m), 4.13-4.46 (4H, m), 6.77-6.90 (2H, m), 7.11-7.26 (2H, m), 7.65 (1H, d, $J = 5.7$ Hz), 7.80 (1H, s), 7.92 (1H, s), 8.34 (1H, s), 8.55 (1H, d, $J = 5.7$ Hz), 8.64 (1H, m). [M+H] Calc'd for $C_{20}H_{18}N_6O$, 359; Found, 359.

Preparation 13A: 2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridine-4-carboxylic acid




[0170] A mixture of 2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridine-4-carbonitrile (430 mg, 1.46 mmol, Preparation 2B) and NaOH (1.5 mL, 7.31 mmol, 5M) in EtOH was refluxed overnight. LC/MS showed the reaction was completed, cooled to rt and acidified to pH=3-4 by 1N HCl, the solid was collected, dried to give the title compound (424 mg, 92%). [M+H] Calc'd for C₁₆H₁₂ClN₃O₂, 314; Found, 314.

Preparation 13B: [2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridin-4-yl]methanol




[0171] To a solution of 2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridine-4-carboxylic acid (424 mg, 1.34 mmol) in THF was added a solution of LiAlH₄ in THF (1.8 mL, 4.43 mmol, 2.4 M) at 0 °C, then stirred for 2 hr at rt. LC/MS showed the reaction was completed. 0.1 mL H₂O, 0.1 mL NaOH, and 0.3 mL H₂O was added subsequently, filtered, concentrated to give the title compound (405 mg, 100%). [M+H] Calc'd for C₁₆H₁₄ClN₃O, 300; Found, 300.

Preparation 13C: 2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridine-4-carbaldehyde




[0172] To a solution of Dess-Martin oxidant (640 mg, 1.5 mmol) in THF (20 mL) was added a solution of [2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridin-4-yl]methanol (300 mg, 1.0 mmol) at 0 °C. The reaction mixture was stirred for 2 hr at rt. LC/MS showed the reaction was completed. NaOH (42 mL, 42 mmol) was added to quench the reaction. It was then extracted by EtOAc, dried and concentrated to give the title compound (300 mg, 61%). [M+H] Calc'd for C₁₆H₁₂ClN₃O, 298; Found, 298.

Preparation 13D: 2-(benzenesulfonyl)-3-[2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridin-4-yl]prop-2-enenitrile




[0173] A mixture of 2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridine-4-carbaldehyde (100 mg, 0.33 mmol), 2-(benzenesulfonyl)acetonitrile (60 mg, 0.33 mmol) and NaHCO_3 (1.64 mL, 0.41 mmol, 0.25 M) in EtOH was stirred overnight at rt. LC/MS showed the reaction was completed, concentrated and purified by preparative TLC (DCM/MeOH=20/1) to give the title compound (40 mg, 26%). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{24}\text{H}_{17}\text{ClN}_4\text{O}_2\text{S}$, 461; Found, 461.

Example 13: 4-[2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile




[0174] A mixture of 2-(benzenesulfonyl)-3-[2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridin-4-yl]prop-2-enenitrile (40 mg, 0.086 mmol, Preparation 13D) and NaN_3 (6 mg, 0.086 mmol) in DMF was stirred for 2 hr at 100 °C. LC/MS showed the reaction was completed, acidified to pH=3-4 by 1N HCl, and stirred for 30 min, then adjusted to pH=7-8 by 1N NaOH, concentrated and purified by prep-HPLC to give the title compound (12 mg, 39%). ^1H NMR (400 MHz, CD_3OD): δ 5.64 (2H, s), 7.46-7.53 (4H, m), 7.92 (1H, d, $J = 5.6$ Hz), 8.20 (1H, s), 8.35 (1H, s), 8.74 (1H, d, $J = 5.2$ Hz), 9.01 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{18}\text{H}_{12}\text{ClN}_7$, 362; Found, 362.

Example 14: 4-[2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile




[0175] The title compound was prepared in 3% overall yield according to the procedure of Example 13 starting from 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]pyridine-4-carbonitrile. ¹H NMR (300 MHz, DMSO): δ 5.43 (2H, s), 7.12-7.15 (1H, m), 7.37-7.43 (1H, m), 7.60-7.93 (4H, m), 8.41-8.49 (2H, m). [M+H] Calc'd for C₁₈H₁₁Cl₂N₇, 396; Found, 396.

Example 15: 4-[2-[1-(1,2,3,4-tetrahydronaphthalen-1-ylmethyl)imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile




[0176] The title compound was prepared in 2% overall yield according to the procedure of Example 13 starting from 2-[1-(1,2,3,4-tetrahydronaphthalen-1-ylmethyl)imidazol-4-yl]pyridine-4-carbonitrile. ¹H NMR (400 MHz, CD₃OD): δ 1.77-1.95 (4H, m), 2.84-2.86 (2H, m), 3.10-3.11 (1H, m), 4.49-4.60 (2H, m), 7.13-7.20 (4H, m), 8.06 (1H, d, *J* = 4.8 Hz), 8.41 (1H, s), 8.46 (1H, s), 8.90 (1H, d, *J* = 5.2 Hz), 9.01 (1H, s). [M+H] Calc'd for C₂₂H₁₉N₇, 382; Found, 382.

Example 16: 4-[2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile




[0177] The title compound was prepared in 2% overall yield according to the procedure of Example 13 starting from 2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine-4-carbonitrile. ¹H NMR (300 MHz, CD₃OD): δ 3.75-3.77 (2H, m), 4.70-4.72 (2H, m), 7.31-7.57 (4H, m), 7.82-8.14 (4H, m), 8.27 (1H, s), 8.35 (1H, s), 8.72 (1H, s), 8.85 (1H, s). [M+H] Calc'd for C₂₃H₁₇N₇, 392; Found, 392.

Preparation 17a: 2-(2-phenylmethoxyphenyl)ethanol




[0178] A mixture of 2-(2-hydroxyethyl)phenol (1g, 7.2 mmol), bromomethylbenzene (1.35 g, 7.92 mmol) and K_2CO_3 (2 g, 14.5 mmol) was stirred overnight at 90 °C. H_2O was added, extracted with ethylacetate, purified by flash column chromatography on silica gel (PE/EA=1/1) to afford the title compound (1.48 g, 90%). [M+H] Calc'd for $C_{15}H_{16}O_2$, 229; Found, 229.

Preparation 17b: 2-(2-phenylmethoxyphenyl)ethyl methanesulfonate




[0179] To a solution of 2-(2-phenylmethoxyphenyl)ethanol (1.48 g, 6.48 mmol) and triethylamine (1.31 g, 13 mmol) in CH_2Cl_2 (20 mL) was added $MsCl$ (1.11g, 9.7 mmol) at 0 °C, then stirred for 2 hr at rt, washed by H_2O , dried, concentrated to give the title compound (1.6 g, 81%) which was used directly for next step. [M+H] Calc'd for $C_{16}H_{18}O_4S$, 307; Found, 307.

Example 17: 2-[1-[2-(2-phenylmethoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0180] The title compound was prepared in 4% overall yield according to the procedure of Example 9 starting from 2-(2-phenylmethoxyphenyl)ethyl methanesulfonate and 2-(1H-imidazol-4-yl)pyridine-4-carboxamide. ^1H NMR (300 MHz, DMSO): δ 3.11 (2H, t, J = 6.9 Hz), 4.26 (2H, t, J = 6.9 Hz), 5.16 (2H, s), 6.83-7.74 (12H, m), 8.30 (1H, s), 8.54-8.64 (2H, m). [M+H] Calc'd for $C_{25}H_{22}N_6O$, 423; Found, 423.

Preparation 18a: 2-(2-phenoxyphenyl)ethanol




[0181] A mixture of 2-(2-hydroxyethyl)phenol (1g, 7.2 mmol), bromobenzene (1.70 g, 10.8 mmol), 1-pyridin-2-yl-propan-2-one (194 mg, 1.44 mmol), CuBr (103 mg, 0.72 mmol) and Cs₂CO₃ (4.73 g, 14.5 mmol) in DMSO (20 mL) was stirred overnight at 80 °C, H₂O was added, extracted with EtOAc, purified by flash column chromatography on silica gel (PE/EA=1/1) to afford the title compound (850 mg, 54%). [M+H]⁺ Calc'd for C₁₄H₁₄O₂, 215; Found, 215.

Preparation 18b: 2-(2-phenoxyphenyl)ethyl methanesulfonate




[0182] The title compound was prepared in 100% yield according to the procedure of Preparation 17b. [M+H]⁺ Calc'd for C₁₅H₁₆O₄S, 293; Found, 293.

Example 18: 2-[1-[2-(2-phenoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0183] The title compound was prepared in 8% overall yield according to the procedure of Example 9 starting from 2-(2-phenoxyphenyl)ethyl methanesulfonate and 2-(1H-imidazol-4-yl)pyridine-4-carboxamide. ¹H NMR (300 MHz, DMSO): δ 3.10 (2H, t, *J* = 6.9 Hz), 4.29 (2H, t, *J* = 6.9 Hz), 6.83-7.34 (9H, m), 7.59 (1H, s), 7.63 (1H, d, *J* = 5.1 Hz), 7.72 (1H, s), 8.30 (1H, s), 8.52 (1H, d, *J* = 5.1 Hz), 8.64 (1H, s). [M+H]⁺ Calc'd for C₂₄H₂₀N₆O, 409; Found, 409.

Preparation 19A: 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]pyridine-4-carbaldehyde




[0184] The title compound was prepared in 19% overall yield according to the procedures of Preparation 13A-13C starting from 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]pyridine-4-carboxamide. [M+H] Calc'd for $C_{16}H_{11}Cl_2N_3O$, 332; Found, 332.

Preparation 19B: 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-ethynylpyridine




[0185] A mixture of 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]pyridine-4-carbaldehyde (330 mg, 1.0 mmol) and K_2CO_3 (276 mg, 2.0 mmol) in MeOH (20 mL) was stirred for 15 min at rt, then (1-diazo-2-oxo-propyl)-phosphonic acid dimethyl ester (470 mg, 2.4 mmol) was added and stirred for 2 hr at rt. LC/MS showed the reaction was completed, concentrated and purified by flash column chromatography on silica gel ($CH_2Cl_2/MeOH=20/1$) to afford the title compound (260 mg, 79%). [M+H] Calc'd for $C_{17}H_{11}Cl_2N_3$, 328; Found, 328.

Preparation 19C: 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-[5-iodo-1-[(4-methoxyphenyl)methyl]triazol-4-yl]pyridine



[0186] A mixture of 1-(azidomethyl)-4-methoxybenzene (310 mg, 1.86 mmol), KI (617 mg, 3.72 mmol) and $Cu(ClO_4)_2 \cdot 6H_2O$ (1.376 g, 3.72 mmol) in THF was stirred for 5 min at rt. Then triethylamine (375 mg, 3.72 mmol) and 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-ethynylpyridine (610 mg, 1.86 mmol) were added sequentially and stirred for 1 h at rt. LC/MS showed the reaction was completed, diluted with EtOAc, washed by 25% $NH_3 \cdot H_2O$, dried and concentrated to give the title compound (916 mg, 80%). [M+H] Calc'd for $C_{25}H_{19}Cl_2IN_6O$, 617; Found, 617.


Example 19: 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-(5-iodo-1H-triazol-4-yl)pyridine



[0187] A solution of 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-[5-iodo-1-[(4-methoxyphenyl)methyl]triazol-4-yl]pyridine (50 mg, 0.08 mmol) in TFA (3 ml) was stirred for 5 hr at 65 °C. LC/MS showed the reaction was completed, concentrated and purified by *prep*-HPLC to give the title compound (10 mg, 25%). ¹H NMR (400 MHz, CD₃OD): δ 5.62 (2H, s), 7.40-7.42 (2H, m), 7.63-7.66 (1H, m), 8.14-8.16 (1H, m), 8.23 (1H, s), 8.57 (1H, s), 8.69-8.72 (2H, m).


[M+H] Calc'd for C₁₇H₁₁Cl₂IN₆, 497; Found, 497.

Preparation 20A: 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-[5-fluoro-1-[(4-methoxyphenyl)methyl]triazol-4-yl]pyridine




[0188] A mixture of 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-[5-iodo-1-[(4-methoxyphenyl)methyl]triazol-4-yl]pyridine (70 mg, 0.11 mmol, Preparation 19C) and KF (33 mg, 0.55 mmol) in ACN/H₂O (2ml/2ml) was stirred for 10 min at 160 °C in a microwave oven. LC/MS showed the reaction was completed, H₂O was added, extracted with EtOAc, dried to give the title compound (46 mg, 80%) as a yellow solid. [M+H] Calc'd for C₂₅H₁₉Cl₂FN₆O, 509; Found, 509.

Example 20: 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-(5-fluoro-1H-triazol-4-yl)pyridine




[0189] The title compound was prepared in 13% yield according to the procedure of last step of Example 19. ^1H NMR (300 MHz, CD_3OD): δ 5.58 (2H, s), 7.36-7.41 (2H, m), 7.60-7.62 (1H, m), 7.81-7.82 (1H, m), 8.18-8.25 (2H, m), 8.39 (1H, s), 8.41 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{17}\text{H}_{11}\text{Cl}_2\text{FN}_6$, 389; Found, 389.

Preparation 21A: 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile




[0190] A mixture of 2-chloro-4-pyridinenitrile (661 mg, 4.77 mmol), 4-(tributylstannyl)-1-tritylimidazole (3 g, 5 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (276 mg, 0.24 mmol) in toluene (20 mL) was heated in a microwave oven for 2 hr at 120 $^\circ\text{C}$. The mixture was concentrated and purified by flash column chromatography on silica gel (EtOAc/hexane) to give the trityl protected product. It was then treated with HOAc/TFA (4 mL/5 mL) for 2 hrs. Purification by flash column chromatography gave the title compound (800 mg, 98%). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_9\text{H}_6\text{N}_4$, 171; Found, 171.

Example 21: 2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0191] The title compound was prepared in 40% yield starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A) according to the procedure of Example 2. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 5.45 (2H, s), 7.13 (1H, d, $J = 6.7$ Hz), 7.40 (1H, t, $J = 7.9$ Hz), 7.63 (1H, br s), 7.65 (1H, d, $J = 8$ Hz), 7.82 (1H, s), 7.93 (1H, s), 8.63 (1H, d, $J = 5.1$ Hz), 8.96 (1H, br s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{17}\text{H}_{12}\text{Cl}_2\text{N}_6$, 371; Found, 371.

Example 22: 2-[1-[[2-chloro-3-(trifluoromethyl)phenyl]methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0192] The title compound was prepared in 4% yield starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A) according to the procedure of Example 2. ¹H NMR (400 MHz, DMSO-d₆): δ 5.52 (2H, s), 7.43 (1H, d, J = 8.0 Hz), 7.59 (1H, t, J = 7.9 Hz), 7.65 (1H, d, J = 3.4 Hz), 7.86 (1H, d, J = 6.7 Hz), 7.95 (1H, s), 8.35 (1H, s), 8.53 (2H, d, J = 5.0 Hz). [M+H] Calc'd for C₁₈H₁₂ClF₃N₆, 405; Found, 405.

Example 23: 2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0193] The title compound was prepared in 11% yield starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A) according to the procedure of Example 6. ¹H NMR (400 MHz, DMSO-d₆): δ 3.60 (2H, t, J = 6.8 Hz), 4.39 (2H, t, J = 6.9 Hz), 7.36 (1H, d, J = 7.0 Hz), 7.42 (1H, t, J = 7.0 Hz), 7.53-7.64 (3H, m), 7.69 (1H, s), 7.82 (1H, d, J = 8.3 Hz), 7.91 (1H, s), 7.94 (1H, d, J = 7.8 Hz), 8.24 (1H, d, J = 8.4 Hz), 8.31 (1H, s), 8.53 (1H, d, J = 4.9 Hz), 8.62 (1H, s). [M+H] Calc'd for C₂₂H₁₈N₆, 367; Found, 367.

Example 24: 2-[5-(4-fluoro-3-methoxyphenyl)-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0194] The title compound was prepared in 20% yield according to the procedure of Example 3 starting from 2-(5-bromo-1-methylimidazol-4-yl)pyridine-4-carbonitrile (Preparation 3A). ¹H NMR (400 MHz, DMSO-d₆): δ 3.53 (3H, s), 3.82 (3H, s), 6.98-7.01 (1H, m), 7.23-7.30 (2H, m), 7.56 (1H, dd, J = 5.0 and 1.6 Hz), 7.86 (1H, s), 8.18 (1H, s), 8.32 (1H, d, J = 5.1 Hz), 8.35 (1H, s), 8.56 (1H, s). [M+H] Calc'd for C₁₈H₁₅FN₆O, 351; Found, 351.

Example 25: 2-[5-(3-ethoxy-4-fluorophenyl)-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0195] The title compound was prepared in 16% yield according to the procedure of Example 3 starting from 2-(5-bromo-1-methylimidazol-4-yl)pyridine-4-carbonitrile (Preparation 3A). ^1H NMR (400 MHz, DMSO- d_6): δ 1.32 (3H, t, $J = 7.0$ Hz), 3.50 (3H, s), 4.08 (2H, q, $J = 6.9$ Hz), 6.96-7.00 (1H, m), 7.23-7.28 (2H, m), 7.56 (1H, dd, $J = 5.1$ and 1.7 Hz), 7.86 (1H, s), 8.18 (1H, s), 8.32 (1H, d, $J = 5.1$ Hz), 8.34 (1H, s), 8.56 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{19}\text{H}_{17}\text{FN}_6\text{O}$, 365; Found, 365.

Example 26: 2-[1-[[2-fluoro-3-(trifluoromethoxy)phenyl]methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0196] The title compound was prepared in 17% yield starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A) according to the procedure of Example 2. ^1H NMR (400 MHz, DMSO- d_6): δ 5.45 (2H, s), 7.34-7.36 (2H, m), 7.55-7.57 (1H, m), 7.64-7.66 (1H, m), 7.81 (1H, s), 7.92 (1H, s), 8.25 (1H, s), 8.33 (1H, s), 8.52 (1H, d, $J = 5.1$ Hz), 8.61 (1H, br s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{18}\text{H}_{12}\text{F}_4\text{N}_6\text{O}$, 405; Found, 405.

Example 27: 2-[1-[2-(2-phenylphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0197] The title compound was prepared in 4% overall yield according to the procedure of Example 6 starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A). ¹H NMR (400 MHz, DMSO-d₆): δ 3.04 (2H, t, *J* = 7.3 Hz), 4.13 (2H, t, *J* = 7.2 Hz), 7.20 (1H, m), 7.21-7.31 (5H, m), 7.33-7.47 (4H, m), 7.54 (1H, dd, *J* = 5.4 and 1.6 Hz), 7.62 (1H, dd, *J* = 5.1 and 1.6 Hz), 8.27 (1H, s), 8.32 (1H, s), 8.59 (1H, s). [M+H]⁺ Calc'd for C₂₄H₂₀N₆, 393; Found, 393.

Example 28: 2-[1-[(2-fluoro-3-methylphenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0198] The title compound was prepared in 7% yield starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A) according to the procedure of Example 2. ¹H NMR (400 MHz, DMSO-d₆): δ 2.14 (3H, s), 5.41 (2H, s), 7.11 (1H, t, *J* = 7.8 Hz), 7.18 (1H, m), 7.28 (1H, m), 7.63 (1H, m), 7.74 (1H, s), 7.87 (1H, s), 8.32 (1H, s), 8.52 (1H, d, *J* = 7.3 Hz). [M+H]⁺ Calc'd for C₁₈H₁₅FN₆, 335; Found, 335.

Example 29: 2-[1-[(3-chloro-2-fluorophenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0199] The title compound was prepared in 5% yield starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A) according to the procedure of Example 2. ¹H NMR (400 MHz, DMSO-d₆): δ 5.41 (2H, s), 7.24-7.33 (2H, m), 7.59 (1H, m), 7.65 (1H, dd, *J* = 5.1 and 1.7 Hz), 7.79 (1H, s), 7.91 (1H, s), 8.32 (1H, s), 8.52 (1H, d, *J* = 5.1 Hz). [M+H]⁺ Calc'd for C₁₇H₁₂ClFN₆, 355; Found, 355.

Example 30: 2-[1-[(2-fluoro-3-methoxyphenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0200] The title compound was prepared in 6% yield starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A) according to the procedure of Example 2. ^1H NMR (400 MHz, DMSO- d_6): δ 3.84 (3H, s), 5.34 (2H, s), 6.90 (1H, m), 7.15 (2H, 2s), 7.64 (1H, d, $J = 5.0$ Hz), 7.74 (1H, s), 7.87 (1H, s), 8.32 (1H, s), 8.53 (1H, d, $J = 5.2$ Hz), 8.63 (1H, br s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{18}\text{H}_{15}\text{FN}_6\text{O}$, 351; Found, 351.

Example 31: 4-(1H-triazol-4-yl)-2-[1-[2-[2-(trifluoromethyl)phenyl]ethyl]imidazole-4-yl]pyridine



[0201] The title compound was prepared in 3% overall yield according to the procedure of Example 6 starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A). ^1H NMR (400 MHz, DMSO- d_6): δ 3.27 (2H, t, $J = 6.4$ Hz), 4.32 (2H, t, $J = 7.8$ Hz), 7.46 (2H, m), 7.60-7.81 (6H, m), 8.47-8.63 (2H, m). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{19}\text{H}_{15}\text{F}_3\text{N}_6$, 385; Found, 385.


Example 32: 2-[1-[2-(2-chlorophenyl)-2-methylpropyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine



[0202] The title compound was prepared in 7% overall yield according to the procedure of Example 6 starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A). ^1H NMR


(400 MHz, DMSO-d₆): δ 1.51 (6H, s), 4.61 (2H, s), 7.23-7.37 (5H, m), 7.53 (1H, d, J = 7.5 Hz), 8.22 (1H, s), 8.42 (1H, s), 8.52 (1H, m). [M+H]⁺ Calc'd for C₂₀H₁₉ClN₆, 379; Found, 379.

Example 33: 2-[1-(1-phenylpropan-2-yl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine




[0203] The title compound was prepared in 1% overall yield according to the procedure of Example 6 starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A). ¹H NMR (400 MHz, DMSO-d₆): δ 0.86 (3H, t, J = 7.2 Hz), 2.33 (2H, m), 5.31 (1H, m), 7.33 (1H, d, J = 7.3 Hz), 7.38 (1H, t, J = 7.2 Hz), 7.46 (1H, s), 7.48 (1H, s), 7.73 (1H, dd, J = 5.0 and 1.6 Hz), 7.93 (1H, s), 7.98 (1H, s), 8.47 (1H, s), 8.54 (1H, d), 8.60 (1H, s). [M+H]⁺ Calc'd for C₁₉H₁₈N₆, 331; Found, 331.

Example 34: 4-(1H-triazol-4-yl)-2-[1-[2-[2-(trifluoromethoxy)phenyl]ethyl]imidazole-4-yl]pyridine



[0204] The title compound was prepared in 5% overall yield according to the procedure of Example 6 starting from 2-(1H-imidazol-4-yl)pyridine-4-carbonitrile (Preparation 21A). ¹H NMR (400 MHz, DMSO-d₆): δ 3.18 (2H, t, J = 7.0 Hz), 4.32 (2H, t, J = 7.2 Hz), 7.20 (1H, d, J = 8.3 Hz), 7.27 (1H, s), 7.62 (1H, dd, J = 5.1 and 1.6 Hz), 7.64 (1H, d, J = 4.5 Hz), 7.83 (1H, d, J = 4.5 Hz), 8.28 (1H, s), 8.56 (1H, d, J = 5.2 Hz), 8.58 (1H, d, J = 5.2 Hz). [M+H]⁺ Calc'd for C₁₉H₁₅F₃N₆O, 401; Found, 401.


Preparation 35A: 4-ethynyl-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine



[0205] A mixture of 2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine-4-carbaldehyde (1.08 g, 3.32 mmol) and K_2CO_3 (0.92 g, 6.64 mmol) in MeOH (20 mL) was stirred for 15 min at rt, then (1-diazo-2-oxo-propyl)-phosphonic acid dimethyl ester (2.4 eq) was added and stirred for 2 hr at rt. LC/MS showed the reaction was completed, concentrated and purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to afford the title compound (0.8 g, 75%).


[M+H] Calc'd for $C_{22}H_{17}N_3$, 324; Found, 324.

Preparation 35B: 4-[5-iodo-1-[(4-methoxyphenyl)methyl]triazol-4-yl]-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine




[0206] A mixture of 1-(azidomethyl)-4-methoxybenzene (310 mg, 1.86 mmol), KI (617 mg, 3.72 mmol) and $Cu(ClO_4)_2 \cdot 6H_2O$ (1.376 g, 3.72 mmol) in THF was stirred for 5 min at rt. Then TEA (375 mg, 3.72 mmol) and 4-ethynyl-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine (600 mg, 1.86 mmol) were added sequentially and stirred for 1 h at rt. LC/MS showed the reaction was completed, diluted with EtOAc, washed by 25% $NH_3 \cdot H_2O$, dried, concentrated to give the title compound (1.1 g, 97%). [M+H] Calc'd for $C_{30}H_{25}IN_6O$, 613; Found, 613.

Preparation 35C: 4-[5-fluoro-1-[(4-methoxyphenyl)methyl]triazol-4-yl]-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine




[0207] A mixture of 4-[5-iodo-1-[(4-methoxyphenyl)methyl]triazol-4-yl]-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine (100 mg, 0.25 mmol) and KF (77 mg, 1.23 mmol) in ACN/H₂O (2ml/2ml) was stirred for 10 min at 160 °C in a microwave oven. LC/MS showed the reaction was completed, H₂O was added, extracted with EtOAc, dried to give the title compound (100 mg, 80%) as a yellow solid. [M+H]⁺ Calc'd for C₃₀H₂₅FN₆O, 505; Found, 505.

Example 35: 4-(5-fluoro-1H-triazol-4-yl)-2-[1-(2-naphthalen-1-ylethyl)imidazole-4-yl]pyridine




[0208] A solution of 4-[5-fluoro-1-[(4-methoxyphenyl)methyl]triazol-4-yl]-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine (100 mg) in TFA (3 mL) was stirred for 5 hr at 65 °C. LC/MS showed the reaction was completed. The reaction mixture was concentrated and purified by prep-HPLC to give the title compound (23 mg, 30%). ¹H NMR (400 MHz, CD₃OD): δ 3.75 (2H, m), 4.95 (2H, m), 7.30-7.58 (4H, m), 7.82-7.93 (3H, m), 8.10-8.25 (3H, m), 8.71 (1H, m), 8.73 (1H, m). [M+H]⁺ Calc'd for C₂₂H₁₇FN₆, 385; Found, 385.

Preparation 36A: 4-[5-chloro-1-[(4-methoxyphenyl)methyl]triazol-4-yl]-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine




[0209] The title compound was prepared in 70% yield according to the procedure of Preparation 35C using KCl. [M+H]⁺ Calc'd for C₃₀H₂₅ClN₆O, 521; Found, 521.

Example 36: 4-(5-chloro-1H-triazol-4-yl)-2-[1-(2-naphthalen-1-ylethyl)imidazole-4-yl]pyridine




[0210] The title compound was prepared in 10% yield according to the general procedure for the preparation of Example 35. ^1H NMR (400 MHz, CD_3OD): δ 3.62-3.64 (2H, m), 4.55-4.57 (2H, m), 7.15-7.44 (4H, m), 7.70-8.25 (6H, m), 8.45-8.47 (1H, m), 8.61-8.63 (1H, m). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{22}\text{H}_{17}\text{ClN}_6$, 401; Found, 401.

Preparation 37A: 4-[1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)triazol-4-yl]-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine



[0211] To a mixture of 4-[5-iodo-1-[(4-methoxyphenyl)methyl]triazol-4-yl]-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine (100 mg, 0.26 mmol), CuI (50 mg, 0.26 mmol), KF (45 mg, 0.78 mmol), Ag_2CO_3 (144 mg, 0.52 mmol) and 1,10-phenanthroline in DMF was added TMSCF_3 (111 mg, 0.78 mmol) at rt. The reaction mixture was stirred for 2 hr at 100 °C in a sealed tube. It was then filtered and separated between water and CH_2Cl_2 , dried and concentrated to give the crude compound (97 mg, 70%). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{31}\text{H}_{25}\text{F}_3\text{N}_6\text{O}$, 555; Found, 555.


Example 37: 2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-triazol-4-yl]pyridine



[0212] The title compound was prepared in 3% yield according to the general procedure for the preparation of Example 35. ^1H NMR (400 MHz, CD_3OD): δ 3.65 (2H, t, J = 6.8 Hz), 4.59 (2H, t,


$J=6.8$ Hz), 7.18-7.48 (4H, m), 7.62-7.64 (1H, m), 7.72 (1H, d, $J=8.0$ Hz), 7.81 (1H, d, $J=8.0$ Hz), 7.89-8.02 (2H, m), 8.10 (1H, s), 8.58 (1H, s), 8.69 (1H, d, $J=4.8$ Hz). [M+H] Calc'd for $C_{23}H_{17}F_3N_6$, 435; Found, 435.

Example 38: 4-(5-iodo-1H-triazol-4-yl)-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine




[0213] The title compound was prepared in 30% yield according to the general procedure for the preparation of Example 35 using 4-[5-iodo-1-[(4-methoxyphenyl)methyl]triazol-4-yl]-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine. ^1H NMR (400 MHz, CD_3OD): δ 3.77 (2H, t, $J=6.8$ Hz), 4.68 (2H, t, $J=6.8$ Hz), 7.32-7.59 (4H, m), 7.84 (1H, d, $J=8.0$ Hz), 7.93 (1H, d, $J=8.0$ Hz), 8.11-8.16 (2H, m), 8.20 (1H, s), 8.42 (1H, s), 8.53 (1H, s), 8.75 (1H, d, $J=4.2$ Hz). [M+H] Calc'd for $C_{22}H_{17}IN_6$, 493; Found, 493.

Preparation 39A: 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridine-4-carboxylic acid




[0214] A mixture of 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine (348 mg, 1 mmol, Preparation 4A) and NaOH (1 mL, 5 mmol) in EtOH (10 mL) was refluxed overnight. LC/MS showed the reaction was completed, cooled to rt and acidified to pH=3-4 by 1N HCl, the solid was filtered and dried to give the title compound (256 mg, 70%). [M+H] Calc'd for $C_{20}H_{18}FN_3O_3$, 368; Found, 368.

Preparation 39B: [2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridin-4-yl]methanol




[0215] To a solution of 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridine-4-carboxylic acid (256 mg, 0.7 mmol) in THF was added a solution of LiAlH_4 in THF (0.96 mL, 2.31 mmol, 2.4 M) at 0 °C, then stirred for 2 hr at rt. LC/MS showed the reaction was completed, 0.1 mL H_2O , 0.1 mL NaOH, and 0.3 mL H_2O were added, filtered, concentrated and purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to give the title compound (123 mg, 50%). [M+H] Calc'd for $\text{C}_{20}\text{H}_{20}\text{FN}_3\text{O}_2$, 354; Found, 354.

Preparation 39C: 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridine-4-carbaldehyde




[0216] To a solution of Dess-Martin reagent (221 mg, 0.52 mmol) in THF (10 mL) was added a solution of 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridine-4-carbaldehyde (123 mg, 0.35 mmol) at 0 °C, then stirred for 2 hr at rt. LC/MS showed the reaction was completed. NaOH (14.7 mL, 14.7 mmol, 1M) was added to quench the reaction. The reaction mixture was extracted by EtOAc, dried and concentrated to give the crude compound (165 mg, 90%). [M+H] Calc'd for $\text{C}_{20}\text{H}_{18}\text{FN}_3\text{O}_2$, 352; Found, 352.

Preparation 39D: 2-(benzenesulfonyl)-3-[2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridin-4-yl]prop-2-enenitrile




[0217] A mixture of 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridine-4-carbaldehyde (165 mg, 0.47 mmol), 2-(benzenesulfonyl)acetonitrile (85 mg, 0.47 mmol) and NaHCO_3 (2.35 mL, 0.58 mmol, 0.25 M) in EtOH (10 mL) was stirred overnight at rt. LC/MS showed the reaction was completed, concentrated and purified by prep-TLC (DCM/MeOH=20/1) to give the title compound (170 mg, 70%). [M+H] Calc'd for $\text{C}_{28}\text{H}_{23}\text{FN}_4\text{O}_3\text{S}$, 515; Found, 515.

Example 39: 4-[2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile




[0218] A mixture of 2-(benzenesulfonyl)-3-[2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridin-4-yl]prop-2-enenitrile (170 mg, 0.33 mmol) and NaN_3 (22 mg, 0.33 mmol) in DMF (10 mL) was stirred for 2 h at 100 °C. LC/MS showed the reaction was completed, acidified to pH=3-4 by 1N HCl, and stirred for 30 min, then adjusted to pH=7-8 by 1N NaOH, concentrated and purified by HPLC to give the title compound (28 mg, 20 %). ^1H NMR (300 MHz, DMSO): δ 0.04-0.35 (4H, m), 0.92-0.93 (1H, m), 3.55 (3H, s), 3.74-3.92 (2H, m), 6.94-6.95 (1H, m), 7.15 (1H, d, $J = 10.2$ Hz), 7.48 (1H, t, $J = 7.5$ Hz), 7.80 (1H, d, $J = 5.4$ Hz), 7.87 (1H, s), 8.73 (1H, d, $J = 5.4$ Hz), 8.85 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{22}\text{H}_{18}\text{FN}_7\text{O}$, 416; Found, 416.

Preparation 40A: 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-ethynylpyridine




[0219] A mixture of 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridine-4-carbaldehyde (1.18 g, 3.32 mmol, Preparation 39C) and K_2CO_3 (0.92 g, 6.64 mmol) in MeOH (20 mL) was stirred for 15 min at rt, then (1-diazo-2-oxo-propyl)-phosphonic acid dimethyl ester (2.4 eq) was added and stirred for 2 hr at rt. LC/MS showed the reaction was completed, concentrated and purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to afford the title compound (617 mg, 75%). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{21}\text{H}_{18}\text{FN}_3\text{O}$, 348; Found, 348.

Preparation 40B: 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-[5-iodo-1-[(4-methoxyphenyl)methyl]triazol-4-yl]pyridine




[0220] To a mixture of 1-(azidomethyl)-4-methoxybenzene (415 mg, 2.49 mmol), KI (826 mg, 4.98 mmol) and $\text{Cu}(\text{ClO}_4)_2 \cdot 6 \text{H}_2\text{O}$ (1.842 g, 4.98 mmol) in THF which was stirred for 5 min at rt, added TEA (503 mg, 4.98 mmol) and 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-ethynylpyridine (617 mg, 2.49 mmol) sequentially. The reaction mixture was stirred for 1 h at rt. LC/MS showed the reaction was completed. It was diluted with EtOAc, washed by 25% $\text{NH}_3 \cdot \text{H}_2\text{O}$, dried, and concentrated to give the title compound (1.1 g, 70%). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{29}\text{H}_{26}\text{FIN}_6\text{O}_2$, 637; Found, 637.

Preparation 40C: 4-[5-chloro-1-[(4-methoxyphenyl)methyl]triazol-4-yl]-2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridine




[0221] A mixture of 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-[5-iodo-1-[(4-methoxyphenyl)methyl]triazol-4-yl]pyridine (230 mg, 0.35 mmol) and KCl (130 mg, 1.75 mmol) in ACN/ H_2O (2ml/2ml) was stirred for 10 min at 160 °C in a microwave oven. LC/MS showed the reaction was completed, H_2O was added, extracted with EtOAc, dried to give the title compound (128 mg, 67%) as a yellow solid. $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{29}\text{H}_{26}\text{ClFN}_6\text{O}_2$, 545; Found, 545.

Example 40: 4-(5-chloro-1H-triazol-4-yl)-2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridine




[0222] A solution of 4-[5-chloro-1-[(4-methoxyphenyl)methyl]triazol-4-yl]-2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridine (128 mg, 0.235 mmol) in TFA (3 ml) was stirred for 5 h at 65 °C. LC/MS showed the reaction was completed, concentrated, purified by HPLC to give the title compound (7 mg, 7%). ¹H NMR (400 MHz, CD₃OD): δ 0.02-0.34 (4H, m), 0.94-0.97 (1H, m), 3.54 (3H, s), 3.77-3.87 (2H, m), 6.88-6.93 (1H, m), 7.06 (1H, d, *J* = 12.8 Hz), 7.44 (1H, t, *J* = 8.0 Hz), 7.78 (1H, s), 7.91 (1H, d, *J* = 4.2 Hz), 8.69 (1H, d, *J* = 4.2 Hz), 8.94 (1H, s). [M+H]⁺ Calc'd for C₂₁H₁₈ClFN₆O, 425; Found, 425.

Preparation 41A: 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-[5-fluoro-1-[(4-methoxyphenyl)methyl]triazol-4-yl]pyridine




[0223] The title compound was prepared in 70% yield according to the procedure of Preparation 35C. [M+H]⁺ Calc'd for C₂₉H₂₆F₂N₆O₂, 529; Found, 529.

Example 41: 2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-(5-fluoro-1H-triazol-4-yl)pyridine



[0224] The title compound was prepared in 10% yield according to the general procedure for the preparation of Example 35. ¹H NMR (400 MHz, CD₃OD): δ 0.04-0.28 (4H, m), 0.83-0.87 (1H, m), 3.43-3.80 (5H, m), 6.81-7.01 (2H, m), 7.34-7.68 (3H, m), 7.98-8.98 (2H, m). [M+H] Calc'd for C₂₁H₁₈F₂N₆O, 409; Found, 409.

Example 42: 1-(cyclopropylmethyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole



[0225] To a solution of 6-chloro-3,4-pyridinediamine (12.5 g, 87 mmol) in HCl (Conc, 100 mL) was added NaNO₂ (6.6 g, 96 mmol) in water (20 mL) dropwise at 0 °C for 1 h. Upon completion, the pH was adjusted to 10 with Na₂CO₃ solution. The slurry was filtered, the solid was washed with PE and dried *in vacuo* to afford 6-chloro-1H-[1,2,3]triazolo[4,5-c]pyridine (10.0 g, 75%) as a yellow solid. [M+H] Calc'd for C₅H₃ClN₄, 155; Found, 155.

[0226] To a mixture of 6-chloro-1H-[1,2,3]triazolo[4,5-c]pyridine (4.62 g, 30 mmol), DIEA (7.8 g, 60 mmol), in DMF (50 mL) was added SEMCl (6.0 g, 36 mmol) at 0 °C and the reaction was stirred at ambient temperature for 16 hours. The mixture was concentrated *in vacuo* and purified by chromatography (ethyl acetate) to afford 6-chloro-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridine (5.1 g, 61%) as a yellow solid. [M+H] Calc'd for C₁₁H₁₇ClN₄OSi, 285; Found, 285.

[0227] To a solution of compound 6-chloro-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridine (6.2 g, 2.2 mmol) in MeOH (200 mL), was added DPPP (943 mg, 2.2 mmol), Et₃N (4.4 g, 44 mmol), and Pd(OAc)₂ (490 mg, 2.2 mmol) at room temperature. The mixture was heated at 100 °C for 48 h under 5 MPa carbon monoxide. The reaction was filtered, and the filtrate concentrated *in vacuo*. The residue was purified by column chromatography (EA/PE = 1/1) to afford methyl 1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridine-6-carboxylate (2.8 g, 41%) as a yellow solid. [M+H] Calc'd for C₁₃H₂₀N₄O₃Si, 309; Found, 309.

[0228] To a solution of methyl 1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridine-6-carboxylate (2.8 g, 9 mmol) in MeOH (30 mL) was added NaBH₄ (1.4 g, 36 mmol) at 0 °C. The mixture was stirred at ambient temperature overnight. Upon completion, the reaction was concentrated *in vacuo* and purified by chromatograph (EA/PE = 2/1) to afford (1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)methanol (2.4 g, 95%) as a yellow oil. [M+H] Calc'd for C₁₂H₂₀N₄O₂Si, 281; Found, 281.


[0229] To a solution of 1-([2-(trimethylsilyl)ethoxy]methyl)-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)methanol (2.4 g, 8.6 mmol) in DCM (100 mL) was added MnO₂ (4.6 g, 36 mmol) at 0°C. The mixture was stirred at ambient temperature overnight. Upon completion, the reaction was concentrated *in vacuo* and the residue purified by column chromatography (EA/PE = 1/1) to afford 1-([2-(trimethylsilyl)ethoxy]methyl)-1H-[1,2,3]triazolo[4,5-c]pyridine-6-carbaldehyde (1.0 g, 40%) as a yellow oil. [M+H] Calc'd for C₁₂H₁₈N₄O₂Si, 279; Found, 279.

[0230] To a solution of 1-([2-(trimethylsilyl)ethoxy]methyl)-1H-[1,2,3]triazolo[4,5-c]pyridine-6-carbaldehyde (0.9 g, 3.2 mmol) and TosMic (0.6 g, 3.2 mmol) in EtOH (50 mL) was added KCN (30 mg, 3.2 mmol) at ambient temperature. The reaction was allowed to stir for 20 min. Upon completion, the mixture was concentrated *in vacuo* and a solution of NH₃ in MeOH (1 g/20 mL) was added. The reaction was stirred for 16 h at 125 °C. The reaction was concentrated *in vacuo* and the residue purified by column chromatography (EA=100) to give 4-(1-([2-(trimethylsilyl)ethoxy]methyl)-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-1H-imidazole (350 mg, 34.5%). [M+H] Calc'd for C₁₄H₂₀N₆OSi, 317; Found, 317.

[0231] A mixture of 4-(1-([2-(trimethylsilyl)ethoxy]methyl)-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-1H-imidazole (100 mg, 0.3 mmol), 1-(bromomethyl)cyclopropane (81 mg, 0.6 mmol) and K₂CO₃ (85 mg, 0.6 mmol) in DMF (10 mL) was allowed to stir overnight at 70 °C. The reaction mixture was filtered and the filtrate concentrated *in vacuo*. The residue was purified by column chromatography (EA/PE= 1/1) to give 1-(cyclopropylmethyl)-4-(1-([2-(trimethylsilyl)ethoxy]methyl)-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-1H-imidazole as a yellow solid in good yield (80 mg, 72%): [M+H] Calc'd for C₁₈H₂₆N₆OSi, 371; Found, 371.


[0232] To a vial charged with 1-(cyclopropylmethyl)-4-(1-([2-(trimethylsilyl)ethoxy]methyl)-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-1H-imidazole (60 mg, 0.16 mmol) in DCM (5 mL) was added TFA (1 mL) and allowed to stir at ambient temperature for 2 h. The reaction mixture was concentrated *in vacuo* and the residue purified by *prep*-HPLC to afford 1-(cyclopropylmethyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole (17 mg, 44%): ¹H NMR (400 MHz, CD₃OD): δ 0.59-0.61 (2H, m), 0.81-0.83 (2H, m), 1.47-1.49 (1H, m), 4.20 (2H, d, *J* = 7.6 Hz), 8.36-8.39 (2H, m), 9.06 (1H, s), 9.49 (1H, s). LCMS (mobile phase: 5%-95% Acetonitrile-Water- 0.02% NH₄Ac): purity is >95%, Rt = 2.694 min. [M+H] Calc'd for C₁₂H₁₂N₆, 241; Found, 241.

Example 43: 4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1-(2,2,2-trifluoroethyl)-1H-imidazole




[0233] The title compound was prepared in 50% yield according to the procedure for example 42. ¹H NMR (400 MHz, CD₃OD): δ 5.15-5.17 (2H, m), 8.20 (1H, s), 8.36 (1H, s), 8.51 (1H, s), 9.49 (1H, s). LCMS (mobile phase: 5%-95% Acetonitrile-Water- 0.02% NH₄Ac): purity is >95%, Rt = 2.639 min. [M+H] Calc'd for C₁₀H₇F₃N₆, 267; Found, 267.

Example 44: 1-benzyl-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole




[0234] The title compound was prepared in 44% yield according to the procedure for example 42. ¹H NMR (400 MHz, CD₃OD): δ 5.47 (2H, s), 7.43-7.49 (5H, m), 8.16 (1H, s), 8.28 (1H, s), 8.79 (1H, s), 9.43 (1H, s). LCMS (mobile phase: 5%-95% Acetonitrile-Water- 0.02% NH₄Ac): purity is >95%, Rt = 2.439 min. [M+H] Calc'd for C₁₅H₁₂N₆, 277; Found, 277.

Example 45: 1-[(2-chlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole




[0235] The title compound was prepared in 43% yield according to the procedure for example 42. ¹H NMR (400 MHz, DMSO-*d*₆): δ 5.46 (2H, s), 7.37-7.45 (3H, m), 7.51-7.56 (1H, m), 8.06 (1H, s), 8.26 (1H, s), 8.55 (1H, s), 9.44 (1H, s). LCMS (mobile phase: 5%-95% Acetonitrile-Water- 0.02% NH₄Ac): purity is >95%, Rt = 2.718 min. [M+H] Calc'd for C₁₅H₁₁ClN₆, 311; Found, 311.

Example 46: 1-[(3-chlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole




[0236] The title compound was prepared in 52% yield according to the procedure for example 42. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 5.49 (2H, s), 7.42-7.46 (3H, m), 7.54 (1H, s), 8.21 (1H, s), 8.30 (1H, s), 8.87 (1H, s), 9.44 (1H, s). LCMS (mobile phase: 5%-95% Acetonitrile-Water-0.02% NH_4Ac): purity is >95%, R_t = 2.815 min. $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{15}\text{H}_{11}\text{ClN}_6$, 311; Found, 311.

Example 47: 1-[(4-chlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole



[0237] The title compound was prepared in 48% yield according to the procedure for example 42. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 5.39 (2H, s), 7.46-7.51 (4H, m), 8.19 (1H, s), 8.27 (1H, s), 8.71 (1H, s), 9.47 (1H, s). LCMS (mobile phase: 5%-95% Acetonitrile-Water-0.02% NH_4Ac): purity is >95%, R_t = 2.682 min. $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{15}\text{H}_{11}\text{ClN}_6$, 311; Found, 311.


Example 48: 1-[(3,4-dichlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole



[0238] The title compound was prepared in 48% yield according to the procedure for example 42. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 5.53 (2H, s), 7.45 (1H, J = 8.0 Hz, d), 7.65 (1H, J = 8.0 Hz, d), 7.74 (1H, s), 8.31 (1H, s), 8.33 (1H, s), 9.11 (1H, s), 9.47 (1H, s). LCMS (mobile phase:


5%-95% Acetonitrile-Water- 0.02% NH₄Ac): purity is >95%, Rt = 2.682 min. [M+H] Calc'd for C₁₅H₁₀Cl₂N₆, 346; Found, 346.

Example 49: 1-(4-chlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole




[0239] A mixture of compound 4-(1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-1H-imidazole (100 mg, 0.3 mmol), 4-chlorophenylboronic acid (56 mg, 0.6 mmol), Cu(OAc)₂ (82 mg), pyridine (0.2 mL) in DCM (10 mL) was allowed to stir overnight at ambient temperature. Upon completion, the reaction was filtered and concentrated *in vacuo* and the residue purified by column chromatography (EA/PE= 1/1) to give crude 4-(1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-1H-imidazole. To a mixture of the crude intermediate in DCM (5 mL) was added TFA (1 mL) and stirred at room temperature for 2 hours. The mixture was concentrated *in vacuo* and purified by *prep*-HPLC to afford title compound (4 mg, 10%). ¹H NMR (300 MHz, DMSO-*d*₆): δ 7.60-7.63 (2H, d, *J* = 9.0 Hz), 7.84-7.87 (2H, d, *J* = 9.0 Hz), 8.19 (1H, s), 8.38 (1H, s), 8.46 (1H, s), 9.43 (1H, s). LCMS (mobile phase: 5%-95% Acetonitrile-Water- 0.02% NH₄Ac): purity is >95%, Rt = 2.972 min. [M+H] Calc'd for C₁₄H₉ClN₆, 297; Found, 297.

Example 50: 1-(2-chlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole




[0240] The title compound was prepared in 6% yield according to the procedure for example 49. ¹H NMR (300 MHz, CD₃OD): δ 7.61-7.69 (2H, m), 7.74-7.79 (2H, m), 8.49 (1H, s), 8.53 (1H, s), 9.15 (1H, s), 9.58 (1H, s). LCMS (mobile phase: 5%-95% Acetonitrile-Water- 0.02% NH₄Ac): purity is >95%, Rt = 2.982 min. [M+H] Calc'd for C₁₄H₉ClN₆, 297; Found, 297.

Example 51: 1-(3-chlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole




[0241] The title compound was prepared in 8% yield according to the procedure for example 49. ¹H NMR (300 MHz, DMSO-*d*₆): δ 7.45-7.61 (2H, m), 7.79-7.84 (1H, m), 8.04 (1H, s), 8.20-8.23 (1H, m), 8.48-8.51 (1H, m), 8.59-8.61 (1H, m), 9.45-9.46 (1H, s). LCMS (mobile phase: 5%-95% Acetonitrile-Water- 0.02% NH₄Ac): purity is >95%, Rt = 3.105 min. [M+H] Calc'd for C₁₄H₉ClN₆, 297; Found, 297.

Example 52: 1-(3,5-dichlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole



[0242] The title compound was prepared in 6% yield according to the procedure for example 49. ¹H NMR (400 MHz, DMSO-*d*₆): δ 7.63 (1H, s), 8.05 (2H, s), 8.19 (1H, s), 8.52-8.59 (2H, m), 9.45 (1H, br). LCMS (mobile phase: 5%-95% Acetonitrile-Water- 0.02% NH₄Ac): purity is >95%, Rt = 2.682 min. [M+H] Calc'd for C₁₄H₈Cl₂N₆, 331; Found, 331.

Example 53: 5-(4-fluorophenyl)-1-methyl-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole




[0243] A mixture of 6-chloro-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridine (1.4 g, 5.0 mmol), 1-methyl-4-(tributylstannyl)-1H-imidazole (2.5 g, 5.0 mmol), and Pd(PPh₃)₄ (530 mg, 0.5 mmol) in DMF (20 mL) was heated at 130 °C for 5 h. Upon completion, the reaction mixture was concentrated *in vacuo* and the residue purified by column chromatography (DCM/MeOH= 10/1) to give 1-methyl-4-(1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-1H-imidazole (1.5 g, 90% yield).

[0244] A round-bottom flask charged with 1-methyl-4-(1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-1H-imidazole (660 mg, 2.0 mmol), and NBS (356 mg, 2.1 mmol) in DCM (10 mL) was allow to stir for 2 h at ambient temperature. Upon completion, water was added and the aqueous layer extracted by DCM. The combined organic layers were successively washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. The resulting crude 5-bromo-1-methyl-4-(1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-1H-imidazole (700 mg) was used in the next step without further purification.


[0245] A mixture of 5-bromo-1-methyl-4-(1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-1H-imidazole (0.5 mmol), 4-fluorophenylboronic acid (140 mg, 1.0 mmol), Pd(dppf)Cl₂ (37 mg, 0.05 mmol) and 2M Na₂CO₃ (1.0 mmol, 2.0 eq) in 5 mL dioxane was kept at 130 °C overnight. Upon completion, the reaction mixture was concentrated *in vacuo* and the residue was purified by column chromatography (DCM/MeOH=50/1) to afford the title compound (50 mg, 30%). ¹H NMR (300 MHz, DMSO-*d*₆): δ 9.17 (1H, s), 7.90 (1H, s), 7.80 (1H, s), 7.44-7.48 (2H, m), 7.25 (t, *J* = 8.7 Hz, 2H), 3.61 (3H, s). LCMS (mobile phase: 5%-95% Acetonitrile-Water-0.1% TFA) purity is >95%, Rt = 2.427min. [M+H] Calc'd for C₁₅H₁₁FN₆, 295; Found, 295.

Example 54: 5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methyl-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole




[0246] The title compound was prepared in 28% yield according to the procedure for example 53. ¹H NMR (300 MHz, CD₃OD): δ 9.47 (1H, s), 9.18 (1H, s), 7.52-7.57 (1H, m), 7.49 (1H, s), 7.14-7.18 (1H, m), 6.96-7.02 (1H, m), 3.89-3.94 (m, 2H), 3.82 (s, 3H), 1.02-1.07 (m, 1H), 0.06-0.43 (m, 4H). LCMS (mobile phase: 5%-95% Acetonitrile-Water- 0.1% NH₄Ac): purity is >95%, Rt = 2.855 min. [M+H] Calc'd for C₁₉H₁₇FN₆O, 365; Found, 365.

Preparation 55A: 2-(5-bromo-1-(2-chlorobenzyl)-1H-imidazol-4-yl)isonicotinonitrile




[0247] The title compound was prepared according to the procedure of Preparation 3A starting from 2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridine-4-carbonitrile. [M+H]⁺ Calc'd for C₁₆H₁₀BrClN₄, 373; Found, 373.

Example 55: 2-(5-bromo-1-(2-chlorobenzyl)-1H-imidazol-4-yl)-4-(2H-1,2,3-triazol-4-yl)pyridine




[0248] The title compound was prepared in 12% yield according to the procedure of Example 1. ¹H NMR (400 MHz, DMSO-*d*₆): δ 5.43 (2H, s), 6.75 (1H, dd, *J* = 7.0 and 2.0 Hz), 7.34-7.40 (2H, m), 7.55 (1H, dd, *J* = 7.8 and 1.8 Hz), 7.73 (1H, dd, *J* = 5.1 and 1.6 Hz), 8.16 (1H, s), 8.46 (1H, s), 8.63-8.68 (2H, m). [M+H]⁺ Calc'd for C₁₇H₁₂BrClN₆, 417; Found, 417.

Example 56: 2-[1-[(2,3-Dichlorophenyl)methyl]imidazol-4-yl]-4-(5-trifluoromethyl-1H-triazol-4-yl)pyridine




[0249] The title compound was prepared in 1% yield according to the procedure of Example 37. ¹H NMR (400 MHz, CD₃OD): δ 5.60 (2H, s), 7.39-7.41 (2H, m), 7.61-7.68 (2H, m), 8.16-8.28 (2H, m), 8.72-8.77 (2H, m). [M+H]⁺ Calc'd for C₁₈H₁₁Cl₂F₃N₆, 439; Found, 439.

Preparation 57A: Methyl 4-(3,3,3-trifluoroprop-1-yn-1-yl)pyridine-2-carboxylate




[0250] A mixture of 4-iodo-2-(1-methoxyethenyl)pyridine (1 g, 3.80 mmol), tetrabutyl(3,3,3-trifluoroprop-1-yn-1-yl)stannane (1.7 g, 3.99 mmol, 90%) and $\text{Pd}(\text{PPh}_3)_4$ (402 mg, 0.38 mmol) in toluene (15 mL) was heated for 2 hr at 120 °C under N_2 in a microwave oven. The mixture was concentrated and purified by flash column chromatography on silica gel (PE/EA=5/1) to afford the title compound (426 mg, 49%). [M+H] Calc'd for $\text{C}_{10}\text{H}_6\text{F}_3\text{NO}_2$, 230; Found, 230.

Preparation 57B: 4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine 2-carboxylate and 4-{1-[(4-methoxyphenyl)methyl]-4-(trifluoromethyl)-1H-1,2,3-triazol-5-yl}pyridine-2-carboxylate




[0251] A solution of methyl 4-(3,3,3-trifluoroprop-1-yn-1-yl)pyridine-2-carboxylate (2.1 g, 9.17 mmol) and 1-azidomethyl-4-methoxy-benzene (1.53 g, 9.17 mmol) in toluene (20 mL) was refluxed for 6 hr. The solution was concentrated and purified by flash column chromatography on silica gel (PE/EA=1/1) to afford a mixture of the title compounds (3.2 g, 89%). [M+H] Calc'd for $\text{C}_{18}\text{H}_{15}\text{F}_3\text{N}_4\text{O}_3$, 393; Found, 393.

Preparation 57C: (4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridin-2-yl)methanol and (4-{1-[(4-methoxyphenyl)methyl]-4-(trifluoromethyl)-1H-1,2,3-triazol-5-yl}pyridin-2-yl)methanol




[0252] A mixture of compound 4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine-2-carboxylate and 4-{1-[(4-methoxyphenyl)methyl]-4-(trifluoromethyl)-1H-1,2,3-triazol-5-yl}pyridine-2-carboxylate (3.2 g, 8.16 mmol) and CaCl_2 (3.624 g, 32.65 mmol) in MeOH/THF (30/15 mL) was added NaBH_4 (620 mg, 16.327 mmol) at 0 °C under N_2 . It was then stirred for 2 hr at rt. LC/MS showed the reaction was completed. The mixture was quenched with H_2O and concentrated, the residue was dissolved in ethyl acetate, washed by H_2O , dried over Na_2SO_4 , concentrated and purified by flash column chromatography on silica gel (PE/EA=1/1) to afford a mixture of the title compounds (2.67 g, 90%). [M+H] Calc'd for $\text{C}_{17}\text{H}_{15}\text{F}_3\text{N}_4\text{O}_2$, 365; Found, 365.

Preparation 57D: 4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine-2-carbaldehyde and 4-{1-[(4-methoxyphenyl)methyl]-4-(trifluoromethyl)-1H-1,2,3-triazol-5-yl}pyridine-2-carbaldehyde




[0253] To a solution of (4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridin-2-yl)methanol and (4-{1-[(4-methoxyphenyl)methyl]-4-(trifluoromethyl)-1H-1,2,3-triazol-5-yl}pyridin-2-yl)methanol (2.67 g, 7.33 mmol) in DCM (50 mL) was added MnO_2 (6.38 g, 73.3 mmol) at rt and stirred for 48 hr. After filtration, the solvent was removed under vacuum and the residue was purified by flash column chromatography on silica gel (PE/EA=2/1) to afford a mixture of the title compounds (1.01/0.94 g, 75%); these two isomers could be separated at this step. [M+H] Calc'd for $\text{C}_{17}\text{H}_{13}\text{F}_3\text{N}_4\text{O}_2$, 363; Found, 363.

Preparation 57E: 2-(1H-imidazol-4-yl)-4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine




[0254] To a solution of 4-{1-[4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine-2-carbaldehyde (2.48 g, 6.85 mmol) and Tos-Mic (1.34 g, 6.85 mmol) in EtOH (100 mL) was added KCN (46 mg, 0.685 mmol) at rt. The reaction mixture was stirred for 2 hr. Then it was added to a solution of NH₃ in MeOH (50 g/50 mL) and stirred overnight at 125°C in a pressure tube. The reaction mixture was concentrated and purified by flash column chromatography (DCM/MeOH/NH₃=10/1/ 0.05) to give the title compound (1.9 g, 61%). [M+H]⁺ Calc'd for C₁₉H₁₅F₃N₆O, 401; Found, 401.

Example 57: 2-{1-[2-(2-chlorophenyl)ethyl]-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0255] A mixture of 2-(1H-imidazol-4-yl)-4-{1-[4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine (80 mg, 0.2 mmol), 2-(2-chlorophenyl)ethyl methanesulfonate (94 mg, 0.4 mmol) and K₂CO₃ (55 mg, 0.4 mmol) in DMF (10 mL) was stirred at room temperature overnight under N₂. The reaction mixture was concentrated and purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to afford an intermediate which was dissolved in TFA/DCM (5/5 mL) and stirred overnight at 50 °C. After the solvent was removed under vacuum, the residue was dissolved in NH₃/MeOH (10 mL), stirred for 20 min, after concentration, the residue was purified by *prep*-TLC (DCM/MeOH/NH₃ =10/1/ 0.05) to give the title compound (7 mg, 8.4 %). ¹H NMR (300 MHz, CD₃OD): δ 3.19-3.20 (2H, m), 4.38-4.44 (2H, m), 6.99-7.21 (5H, m), 7.54-7.82 (2H, m), 8.08-8.22 (2H, m). Calc'd for C₁₉H₁₄ClF₃N₆, 419; Found, 419.

Example 58: 2-{1-[1,2,3,4,4a,8a-hexahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0256] The title compound was prepared in 6% yield according to the general procedure for the preparation of Example 57. ^1H NMR (300 MHz, CD_3OD): δ 1.63-1.85 (4H, m), 2.57-2.64 (2H, m), 3.25-3.30 (1H, m), 4.25-4.26 (2H, m), 7.03-7.20 (4H, m), 7.54-7.86 (2H, m), 8.19-8.24 (2H, m), 8.58-8.63 (1H, m). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{22}\text{H}_{19}\text{F}_3\text{N}_6$, 425; Found, 425.

Example 59: 2-{1-[2-(6-ethoxycyclohexa-2,4-dien-1-yl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0257] The title compound was prepared in 18% yield according to the general procedure for the preparation of Example 57. ^1H NMR (300 MHz, CD_3OD): δ 1.23-1.27 (3H, m), 3.09-3.12 (2H, m), 3.74-3.75 (2H, m), 4.37-4.39 (2H, m), 6.63-7.10 (5H, m), 7.59-7.73 (2H, m), 8.09-8.21 (2H, m). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{21}\text{H}_{19}\text{F}_3\text{N}_6\text{O}$, 429; Found, 429.

Example 60: 4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-(1-{2-[6-(trifluoromethyl)cyclohexa-2,4-dien-1-yl]ethyl}-1H-imidazol-4-yl)pyridine




[0258] The title compound was prepared in 15% yield according to the general procedure for the preparation of Example 57. ^1H NMR (300 MHz, CD_3OD): δ 3.31 (2H, t, $J = 7.2$ Hz), 4.37 (2H, t, $J = 7.2$ Hz), 7.28-7.30 (1H, m), 7.39-7.73 (6H, m), 8.18 (1H, s), 8.58 (1H, d, $J = 5.1$ Hz). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{20}\text{H}_{14}\text{F}_6\text{N}_6$, 453; Found, 453.

Example 61: 2-{1-[(4-fluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0259] A mixture of 2-(1H-imidazol-4-yl)-4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine (60 mg, 0.15 mmol), 4-fluorobenzylbromide (34 mg, 0.18 mmol) and K₂CO₃ (41.4 mg, 0.3 mmol) in DMF (10 mL) was stirred at rt under N₂ overnight. The solvent was removed under vacuum, and the residue was purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to afford the intermediate, which was dissolved in TFA/DCM (5/5 mL) and stirred overnight at 50 °C. The reaction mixture was then concentrated and the residue was dissolved in NH₃/MeOH (10 mL), stirred for 20 min, then concentrated and purified by prep TLC (DCM/MeOH/NH₃ =10/1/ 0.05) to give the title compound (19 mg, 32 %). ¹H NMR (300 MHz, CD₃OD): δ 5.49 (2H, s), 7.18 (2H, t, *J* = 8.7 Hz), 7.52-7.57 (2H, m), 7.71 (1H, d, *J* = 4.8 Hz), 8.22 (1H, s), 8.32 (1H, s), 8.79 (1H, d, *J* = 5.1 Hz), 9.12 (1H, s). [M+H]⁺ Calc'd for C₁₈H₁₂F₄N₆, 389; Found, 389.

Example 62: 2-{1-[(2-fluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0260] The title compound was prepared in 34% yield according to the general procedure for the preparation of Example 61. ¹H NMR (300 MHz, CD₃OD): δ 5.36 (2H, s), 7.13-7.50 (5H, m), 7.77-7.88 (2H, m), 8.16-8.20 (1H, m), 8.53-8.59 (1H, m). [M+H]⁺ Calc'd for C₁₈H₁₂F₄N₆, 389; Found, 389.

Example 63: 2-{1-[(4-chlorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0261] The title compound was prepared in 42% yield according to the general procedure for the preparation of Example 61. ¹H NMR (300 MHz, CD₃OD): δ 5.52 (2H, s), 7.43-7.52 (4H, m), 7.71 (1H, d, *J* = 4.8 Hz), 8.23 (1H, s), 8.35 (1H, s), 8.79 (1H, d, *J* = 5.1 Hz), 9.15 (1H, s). [M+H]⁺ Calc'd for C₁₈H₁₂ClF₃N₆, 405; Found, 405.

Example 64: 2-{1-[(2-chlorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0262] The title compound was prepared in 34% yield according to the general procedure for the preparation of Example 61. ^1H NMR (300 MHz, CD_3OD): δ 5.42 (2H, s), 7.34-7.48 (5H, m), 7.74 (1H, s), 7.86 (1H, s), 8.20 (1H, s), 8.51 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{18}\text{H}_{12}\text{ClF}_3\text{N}_6$, 405; Found, 405.

Example 65: 2-{1-[(4-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0263] The title compound was prepared in 45% yield according to the general procedure for the preparation of Example 61. ^1H NMR (300 MHz, CD_3OD): δ 2.33 (3H, s), 5.45 (2H, s), 7.24-7.38 (4H, m), 7.69 (1H, d, $J = 4.8$ Hz), 8.20 (1H, s), 8.27 (1H, s), 8.78 (1H, d, $J = 5.1$ Hz), 9.09 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{19}\text{H}_{15}\text{F}_3\text{N}_6$, 385; Found, 385.

Example 66: 2-{1-[(2-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0264] The title compound was prepared in 63% yield according to the general procedure for the preparation of Example 61. ^1H NMR (300 MHz, CD_3OD): δ 2.17 (3H, s), 5.52 (2H, s), 7.28-7.34 (4H, m), 7.71 (1H, d, $J = 4.0$ Hz), 8.17-8.22 (2H, m), 8.77 (1H, d, $J = 5.2$ Hz), 8.94 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{19}\text{H}_{15}\text{F}_3\text{N}_6$, 385; Found, 385.

Example 67: 2-{1-[(4-methoxyphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0265] The title compound was prepared in 55% yield according to the general procedure for the preparation of Example 61. ^1H NMR (300 MHz, CD_3OD): δ 3.79 (3H, s), 5.42 (2H, s), 6.97 (2H, d, $J = 8.7$ Hz), 7.44 (2H, d, $J = 9.0$ Hz), 7.68 (1H, d, $J = 4.5$ Hz), 8.20 (1H, s), 8.27 (1H, s), 8.74 (1H, d, $J = 5.4$ Hz), 9.08 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{19}\text{H}_{15}\text{F}_3\text{N}_6\text{O}$, 401; Found, 401.

Example 68: 2-{1-[(2,4-difluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0266] The title compound was prepared in 67% yield according to the general procedure for the preparation of Example 61. ^1H NMR (300 MHz, CD_3OD): δ 5.57 (2H, s), 7.04 (2H, t, $J = 8.7$ Hz), 7.65-7.71 (2H, m), 8.24 (1H, s), 8.30 (1H, s), 8.76 (1H, d, $J = 5.4$ Hz), 9.09 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{18}\text{H}_{11}\text{F}_5\text{N}_6$, 407; Found, 407.

Example 69: 2-{1-[(2,6-difluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0267] The title compound was prepared in 30% yield according to the general procedure for the preparation of Example 61. ^1H NMR (300 MHz, CD_3OD): δ 5.63 (2H, s), 7.09-7.15 (2H, m), 7.50-7.55 (1H, m), 7.74 (1H, d, $J = 4.8$ Hz), 8.26 (2H, d, $J = 6.9$ Hz), 8.77 (1H, d, $J = 4.8$ Hz), 9.05 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{18}\text{H}_{11}\text{F}_5\text{N}_6$, 407; Found, 407.

Example 70: 2-{1-[(4-fluoro-2-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0268] The title compound was prepared in 53% yield according to the general procedure for the preparation of Example 61. ^1H NMR (400 MHz, CD_3OD): δ 2.38 (3H, s), 5.53 (2H, s), 7.02-7.10 (2H, m), 7.39-7.43 (1H, m), 7.38-7.50 (1H, m), 8.21 (1H, s), 8.25 (1H, s), 8.80 (1H, d, $J = 5.2$ Hz), 8.97 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{19}\text{H}_{14}\text{F}_4\text{N}_6$, 403; Found, 403.

Example 71: 2-{1-[(2-fluoro-4-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0269] The title compound was prepared in 48% yield according to the general procedure for the preparation of Example 61. ^1H NMR (300 MHz, CD_3OD): δ 2.37 (3H, s), 5.54 (2H, s), 7.03-7.12 (2H, m), 7.47 (1H, t, $J = 8.1$ Hz), 7.70 (1H, d, $J = 4.8$ Hz), 8.24 (1H, s), 8.31 (1H, s), 8.79 (1H, d, $J = 5.1$ Hz), 9.12 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{19}\text{H}_{14}\text{F}_4\text{N}_6$, 403; Found, 403.

Example 72: 2-(1-benzyl-1H-imidazol-4-yl)-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0270] The title compound was prepared in 6% yield according to the general procedure for the preparation of Example 61. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 5.43 (2H, s), 7.15 (1H, d, $J = 8.0$ Hz), 7.24 (1H, d, $J = 8.1$ Hz), 7.26 (5H, m), 7.52 (1H, t, $J = 4.8$ Hz), 7.63 (1H, br s), 7.91 (1H, br s), 7.98 (1H, s), 8.18 (1H, br s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{18}\text{H}_{13}\text{F}_3\text{N}_6$, 371; Found, 371.

Example 73: 2-[1-(1-phenylethyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0271] The title compound was prepared in 6% yield according to the general procedure for the preparation of Example 71. [M+H] Calc'd for C₁₉H₁₅F₃N₆, 385; Found, 385.

Example 74: 2-[1-(1-phenylpropyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0272] The title compound was prepared in 69% yield according to the general procedure for the preparation of Example 61. ¹H NMR (400 MHz, DMSO-d₆): δ 0.87 (3H, s), 2.30 (1H, t, J = 8.8 Hz), 2.40 (1H, t, J = 6.4 Hz), 5.44 (2H, s), 7.36-7.57 (6H, m), 8.22 (1H, s), 8.41 (1H, br s), 8.76 (1H, br s). [M+H] Calc'd for C₂₀H₁₇F₃N₆, 398; Found, 398.

Example 75: 2-[1-(2-methyl-1-phenylpropyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0273] The title compound was prepared in 6% yield according to the general procedure for the preparation of Example 61. ¹H NMR (400 MHz, DMSO-d₆): δ 0.81 (3H, d, J = 5.1 Hz), 0.92 (3H, d, J = 3.6 Hz), 2.80 (1H, m), 5.05 (1H, d, J = 10.2 Hz), 7.34-7.63 (6H, m), 8.19 (1H, s), 8.37 (1H, br s), 8.72 (1H, br s). [M+H] Calc'd for C₂₁H₁₉F₃N₆, 413; Found, 413.

Example 76: 2-(1-{[2-(2,2,2-trifluoroethoxy)phenyl]methyl}-1H-imidazol-4-yl)-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0274] The title compound was prepared in 31% yield according to the general procedure for the preparation of Example 61. ^1H NMR (400 MHz, DMSO- d_6): δ 4.84 (2H, q, $J = 17.2$ Hz), 5.35 (2H, s), 7.10 (1H, m), 7.18 (1H, m), 7.41 (2H, m), 7.55 (1H, s), 8.07 (1H, s), 8.18 (1H, s), 8.47 (1H, br s), 8.74 (1H, s). [M+H] Calc'd for $\text{C}_{20}\text{H}_{14}\text{F}_6\text{N}_6\text{O}$, 469; Found, 469.

Example 77: 2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0275] A mixture of 2-(1H-imidazol-4-yl)-4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine (50 mg, 0.12 mmol), 1-(2-bromoethyl)-4-fluorobenzene (60 mg, 0.3 mmol) and Cs_2CO_3 (196 mg, 0.6 mmol) in DMF (5 mL) was heated at 90 °C for 2 h under N_2 . The reaction mixture was concentrated and purified by flash column chromatography on silica gel (EtOAc/Hexane, 0-100%) to afford an intermediate which was dissolved in TFA/DCM (5/5 mL) and stirred overnight at 50 °C. The reaction mixture was concentrated and the residue was purified by flash column chromatography to give the title compound (21.5 mg, 44 %). ^1H NMR (400 MHz, DMSO- d_6): δ 3.16 (2H, s), 4.38 (2H, s), 7.13 (2H, m), 7.26 (2H, s), 7.57 (1H, s), 8.16 (1H, s), 8.21 (1H, s), 8.38 (1H, br s), 8.76 (1H, s). [M+H] Calc'd for $\text{C}_{19}\text{H}_{14}\text{F}_4\text{N}_6$, 403; Found, 403.

Example 78: 2-{1-[2-(3-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0276] The title compound was prepared in 68% yield according to the general procedure for the preparation of Example 77. ¹H NMR (400 MHz, DMSO-d₆): δ 3.20 (2H, s), 4.43 (2H, s), 7.05 (2H, m), 7.12 (1H, d, J = 7.4 Hz), 7.33 (1H, d, J = 7.4 Hz), 7.58 (1H, s), 8.17 (1H, s), 8.24 (1H, s), 8.46 (1H, br s), 8.77 (1H, s). [M+H] Calc'd for C₁₉H₁₄F₄N₆, 403; Found, 403.

Example 79: 2-{1-[2-(2,3-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0277] The title compound was prepared in 32% yield according to the general procedure for the preparation of Example 77. ¹H NMR (400 MHz, DMSO-d₆): δ 3.22 (2H, t, J = 6.7 Hz), 4.35 (2H, t, J = 5.2 Hz), 6.97-7.13 (3H, m), 7.29 (1H, m), 7.48 (1H, m), 7.89 (1H, s), 8.14 (1H, s), 8.66 (1H, s). [M+H] Calc'd for C₁₉H₁₃F₅N₆, 421; Found, 421.

Example 80: 2-{1-[2-(2-chloro-6-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0278] The title compound was prepared in 27% yield according to the general procedure for the preparation of Example 77. ¹H NMR (400 MHz, DMSO-d₆): δ 3.20 (2H, t, J = 6.3 Hz), 4.35 (2H, t, J = 6.1 Hz), 6.96 (1H, m), 7.18 (2H, m), 7.22 (2H, m), 7.36 (1H, m), 7.92 (1H, s), 8.14 (1H, s), 8.69 (1H, s). [M+H] Calc'd for C₁₉H₁₃ClF₄N₆, 437; Found, 437.

Example 81: 4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-{1-[2-(2,4,5-trifluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine




[0279] The title compound was prepared in 23% yield according to the general procedure for the preparation of Example 77. ^1H NMR (400 MHz, DMSO- d_6): δ 3.08 (2H, t, J = 6.7 Hz), 4.32 (2H, t, J = 6.7 Hz), 7.41-7.55 (3H, m), 7.81 (1H, m), 7.91 (1H, s), 7.95 (1H, s), 8.14 (1H, s), 8.67 (1H, d, J = 4.9 Hz). [M+H] Calc'd for $\text{C}_{19}\text{H}_{12}\text{F}_6\text{N}_6$, 439; Found, 439.

Example 82: 4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-{1-[2-(2,4,6-trifluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine




[0280] The title compound was prepared in 23% yield according to the general procedure for the preparation of Example 77. ^1H NMR (400 MHz, DMSO- d_6): δ 3.13 (2H, t, J = 6.5 Hz), 4.30 (2H, t, J = 5.9 Hz), 7.15 (2H, t, J = 8.5 Hz), 7.47 (1H, d, J = 4.0 Hz), 7.55 (1H, m), 7.83 (1H, m), 7.95 (1H, s), 8.13 (1H, s), 8.66 (1H, d, J = 4.3 Hz). [M+H] Calc'd for $\text{C}_{19}\text{H}_{12}\text{F}_6\text{N}_6$, 439; Found, 439.

Example 83: 2-{1-[(2,3-dihydro-1H-inden-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0281] The title compound was prepared in 5% yield according to the general procedure for the preparation of Example 77. ^1H NMR (400 MHz, DMSO- d_6): δ 1.85 (2H, m), 2.05 (1H, m), 2.95 (2H, m), 4.28 (2H, m), 7.16 (3H, m), 7.26 (1H, m), 7.40 (1H, s), 7.80 (1H, s), 8.22 (1H, s), 8.37 (1H, s). [M+H] Calc'd for $\text{C}_{21}\text{H}_{17}\text{F}_3\text{N}_6$, 411; Found, 411.

Example 84: 2-{1-[2-(2-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0282] The title compound was prepared in 9% yield according to the general procedure for the preparation of Example 77. ¹H NMR (400 MHz, DMSO-d₆): δ 3.26 (2H, br s), 4.48 (2H, br s), 7.17 (2H, m), 7.29 (2H, m), 7.66 (1H, s), 8.19 (1H, s), 8.49 (1H, s), 8.85 (1H, s), 8.97 (1H, br s). [M+H]⁺ Calc'd for C₁₉H₁₄F₄N₆, 403; Found, 403.

Example 85: 2-[1-(6-fluoro-2,3-dihydro-1H-inden-1-yl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine



[0283] The title compound was prepared in 28% yield according to the general procedure for the preparation of Example 61. ¹H NMR (400 MHz, DMSO-d₆): δ 2.33 (2H, m), 2.80 (1H, m), 2.99 (1H, m), 6.10 (1H, m), 7.19 (2H, m), 7.46 (1H, m), 7.62 (1H, m), 8.26 (1H, s), 8.33 (1H, s), 8.82 (1H, s), 9.04 (1H, br s). [M+H]⁺ Calc'd for C₂₀H₁₄F₄N₆, 415; Found, 415.


Example 86: 2-{1-[2-(2,4-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine



[0284] A mixture of 2-(1H-imidazol-4-yl)-4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine (60 mg, 0.15 mmol), 1-(2-bromoethyl)-2,4-difluorobenzene (40 mg, 0.18 mmol) and K₂CO₃ (41.4 mg, 0.3 mmol) in DMF (10 mL) was stirred overnight at 80 °C under N₂. The solvent was removed under vacuum and the residue was purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to afford the PBM protected


intermediate which was dissolved in TFA/DCM (5/5 mL) and stirred overnight at 50 °C. After concentration, the residue was dissolved in NH₃/MeOH (10 mL), stirred for 20 min. The mixture was then concentrated and purified by *prep*-TLC (DCM/MeOH/NH₃ = 10/1/ 0.05) to give the title compound (10 mg, 15%). ¹H NMR (300 MHz, CD₃OD): δ 3.30 (2H, t, *J* = 6.9 Hz), 5.54 (2H, t, *J* = 6.9 Hz), 6.93-6.99 (2H, m), 7.25-7.27 (1H, m), 7.73 (1H, d, *J* = 4.2 Hz), 8.18 (1H, s), 8.28 (1H, s), 8.80 (1H, d, *J* = 5.1 Hz), 8.89 (1H, s). [M+H] Calc'd for C₁₉H₁₃F₅N₆, 421; Found, 421.

Example 87: 2-{1-[2-(2,6-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0285] The title compound was prepared in 11% yield according to the general procedure for the preparation of Example 86. ¹H NMR (300 MHz, CD₃OD): δ 3.38 (2H, t, *J* = 6.8 Hz), 4.59 (2H, t, *J* = 6.8 Hz), 6.96-6.99 (2H, m), 7.32-7.37 (1H, m), 7.73-7.75 (1H, m), 8.17 (1H, s), 8.28 (1H, s), 8.80 (1H, d, *J* = 5.2 Hz), 8.98 (1H, s). [M+H] Calc'd for C₁₉H₁₃F₅N₆, 421; Found, 421.

Example 88: 2-{1-[2-(2,5-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine



[0286] The title compound was prepared in 21% yield according to the general procedure for the preparation of Example 86. ¹H NMR (300 MHz, CD₃OD): δ 3.38 (2H, t, *J* = 6.8 Hz), 4.59 (2H, t, *J* = 6.8 Hz), 7.04-7.13 (3H, m), 7.74 (1H, dd, *J* = 5.2 Hz, *J* = 1.2 Hz), 8.20 (1H, s), 8.31 (1H, s), 8.81 (1H, d, *J* = 5.2 Hz), 8.98 (1H, s). [M+H] Calc'd for C₁₉H₁₃F₅N₆, 421; Found, 421.


Preparation 89A: tert-butyl (3S)-3-(methanesulfonyloxy)piperidine-1-carboxylate



[0287] To a solution of tert-butyl (3S)-3-hydroxypiperidine-1-carboxylate (1 g, 4.98 mmol) and TEA (754 mg, 7.46 mmol) in DCM was added MsCl (684 mg, 5.97 mmol) at 0 °C, then stirred


for 2 hr at rt. The reaction mixture was washed by H₂O, dried over Na₂SO₄; after concentration the crude title compound (1.3 g, 94%) was used directly for the next step.

Preparation 89B: 4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}-2-{1-[(3R)-piperidin-3-yl]-1H-imidazol-4-yl}pyridine



[0288] A mixture of 2-(1H-imidazol-4-yl)-4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine (110 mg, 0.275 mmol), tert-butyl (3S)-3-(methanesulfonyloxy)piperidine-1-carboxylate (154 mg, 0.552 mmol), and Cs₂CO₃ (173 mg, 0.552 mmol) in DMF (10 mL) was stirred overnight at 100 °C under N₂. After concentration, the residue was purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to afford the Boc protected intermediate which was dissolved in TFA/DCM (1/10 mL) and stirred for 3 hr at rt, after concentration the residue was dissolved in NH₃/MeOH (10 mL), stirred for 20 min, then concentrated and purified by *prep*-TLC (DCM/MeOH/NH₃ =10/1/ 0.05) to give the title compound (30 mg, 23%). [M+H] Calc'd for C₂₄H₂₄F₃N₇O, 484; Found, 484.


Example 89: 2-{1-[(3R)-1-benzoylpiperidin-3-yl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine



[0289] To a solution of 4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}-2-{1-[(3R)-piperidin-3-yl]-1H-imidazol-4-yl}pyridine (30 mg, 0.062 mmol) and TEA (13 mg, 0.124 mmol) in DCM was added benzoyl chloride (11 mg, 0.0745 mmol) at rt. The resulted solution was stirred for 2 hr at rt, then washed by H₂O, dried over Na₂SO₄, concentrated to give the crude intermediate which was dissolved in TFA/DCM (5/5 mL) and stirred overnight at 50 °C. After concentration, the residue was dissolved in NH₃/MeOH (10 mL), stirred for 20 min, then concentrated and purified by *prep*-TLC (DCM/MeOH/NH₃ =10/1/ 0.05) to give the title compound (16 mg, 55%). ¹H NMR (300 MHz, CD₃OD): δ 1.84-2.02 (2H, m), 2.18-2.32 (2H, m), 3.64-3.68


(2H, m), 4.65-4.66 (2H, m), 5.28-5.33 (1H, m), 7.46-7.49 (5H, m), 7.73-7.74 (1H, m), 8.25-8.27 (1H, m), 8.49-8.50 (1H, m), 8.80-8.82 (1H, m), 9.16-9.18 (1H, m). [M+H] Calc'd for $C_{23}H_{20}F_3N_7O$, 468; Found, 468.

Example 90: 2-{1-[(3S)-1-benzoylpiperidin-3-yl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0290] The title compound was prepared in 15% yield according to the general procedure for the preparation of Example 89. ^1H NMR (300 MHz, CD_3OD): δ 1.86-2.05 (2H, m), 2.29-2.43 (2H, m), 3.62-3.63 (2H, m), 4.63-4.64 (2H, m), 5.27-5.32 (1H, m), 7.46-7.49 (5H, m), 7.72-7.73 (1H, m), 8.25-8.27 (1H, m), 8.49-8.50 (1H, m), 8.80-8.82 (1H, m), 9.17-9.19 (1H, m). [M+H] Calc'd for $C_{23}H_{20}F_3N_7O$, 468; Found, 468.

Example 91: 2-{1-[2-(4-methylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0291] A mixture of 2-(1H-imidazol-4-yl)-4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine (60 mg, 0.15 mmol), 1-(2-bromoethyl)-4-methylbenzene (63 mg, 0.30 mmol) and Cs_2CO_3 (147 mg, 0.45 mmol) in DMF (10 mL) was stirred overnight at 90 °C under N_2 , the solvent was removed under vacuum and the residue was purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to afford the intermediate. It was dissolved in TFA/DCM (5/5 mL) and stirred overnight at 50 °C; after concentration, the residue was dissolved in NH_3 /MeOH (10 mL), stirred for 20 min, then concentrated and purified by *prep*-TLC (DCM/MeOH/ NH_3 =10/1/0.05) to give the title compound (20 mg, 33%). ^1H NMR (400 MHz, CD_3OD): δ 2.29 (3H, s), 3.22 (2H, t, J = 6.8 Hz), 4.56 (2H, t, J = 6.8 Hz), 7.07-7.14 (4H, m), 7.72 (1H, d, J = 5.2 Hz), 8.16 (1H, s), 8.23 (1H, s), 8.81 (2H, s). [M+H] Calc'd for $C_{20}H_{17}F_3N_6$, 399; Found, 399.

Example 92: 2-{1-[2-(4-chlorophenyl)ethyl]-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0292] The title compound was prepared in 39.8% yield according to the general procedure for the preparation of Example 91. ^1H NMR (400 MHz, CD_3OD): δ 3.27 (2H, t, $J = 6.8$ Hz), 4.59 (2H, t, $J = 6.8$ Hz), 7.23 (2H, d, $J = 8.0$ Hz), 7.31 (2H, d, $J = 8.0$ Hz), 7.73 (1H, d, $J = 4.8$ Hz), 8.20 (1H, s), 8.31 (1H, s), 8.81 (1H, d, $J = 5.2$ Hz), 8.88 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{19}\text{H}_{14}\text{ClF}_3\text{N}_6$, 419; Found, 419.

Example 93: 2-{1-[2-(4-methoxyphenyl)ethyl]-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine



[0293] The title compound was prepared in 37% yield according to the general procedure for the preparation of Example 91. ^1H NMR (300 MHz, CD_3OD): δ 3.19 (2H, t, $J = 6.4$ Hz), 3.75 (3H, s), 4.54 (2H, t, $J = 6.8$ Hz), 6.86 (2H, d, $J = 8.0$ Hz), 7.11 (2H, d, $J = 8.0$ Hz), 7.72 (1H, d, $J = 5.2$ Hz), 8.17 (1H, s), 8.23 (1H, s), 8.79-8.80 (2H, m). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{20}\text{H}_{17}\text{F}_3\text{N}_6\text{O}$, 415; Found, 415.


Example 94: 4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-(1-{2-[4-(trifluoromethyl)phenyl]ethyl}-1H-imidazol-4-yl)pyridine



[0294] The title compound was prepared in 22% yield according to the general procedure for the preparation of Example 91. ^1H NMR (300 MHz, CD_3OD): δ 3.38 (2H, t, $J = 6.8$ Hz), 4.63 (2H, t, $J = 7.2$ Hz), 7.45 (2H, d, $J = 8.0$ Hz), 7.62 (2H, d, $J = 7.6$ Hz), 7.74 (1H, d, $J = 4.8$ Hz), 8.20


(1H, s), 8.32 (1H, s), 8.81 (1H, d, $J = 5.2$ Hz), 8.90 (1H, s). [M+H] Calc'd for $C_{20}H_{14}F_6N_6$, 453; Found, 453.

Example 95: 2-{1-[2-(4-ethylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0295] A mixture of 2-(1H-imidazol-4-yl)-4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine (100 mg, 0.25 mmol), 2-(4-ethylphenyl)ethyl methanesulfonate (134 mg, 0.50 mmol) and Cs_2CO_3 (245 mg, 0.75 mmol) in DMF (10 mL) was stirred overnight at 100 °C under N_2 , the solvent was removed under vacuum and the residue was purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to afford the intermediate. It was then dissolved in TFA/DCM (5/5 mL) and stirred overnight at 50 °C; after concentration, the residue was dissolved in NH_3 /MeOH (10 mL), stirred for 20 min, then concentrated and purified by *prep*-TLC (DCM/MeOH/ NH_3 =10/1/ 0.05) to give the title compound (37 mg, 36%). ^1H NMR (400 MHz, CD_3OD): δ 1.14 (3H, t, $J = 7.6$ Hz), 2.56 (2H, q, $J = 8.0$ Hz), 3.20 (2H, t, $J = 6.8$ Hz), 4.54 (2H, t, $J = 6.8$ Hz), 7.06-7.13 (4H, m), 7.69 (1H, dd, $J = 4.8$ Hz, $J = 1.2$ Hz), 8.13 (1H, s), 8.16 (1H, s), 8.75-8.77 (2H, m). [M+H] Calc'd for $C_{21}H_{19}F_3N_6$, 413; Found, 413.

Example 96: 2-{1-[2-(4-tert-butylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0296] The title compound was prepared in 34% yield according to the general procedure for the preparation of Example 95. ^1H NMR (300 MHz, CD_3OD): δ 1.27 (9H, s), 3.24 (2H, t, $J = 6.8$ Hz), 4.58 (2H, t, $J = 7.2$ Hz), 7.13 (2H, d, $J = 8.4$ Hz), 7.35 (2H, d, $J = 8.0$ Hz), 7.72 (1H, dd, $J = 4.8$ Hz, $J = 1.2$ Hz), 8.16 (1H, s), 8.18 (1H, s), 8.79-8.82 (2H, m). [M+H] Calc'd for $C_{23}H_{23}F_3N_6$, 441; Found, 441.

Preparation 97A: 4-[1-benzyl-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-(1-methyl-1H-imidazol-4-yl)pyridine




[0297] A mixture of 2-(1H-imidazol-4-yl)-4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine (300 mg, 0.75 mmol) in DMF (10 mL) was added NaH (33 mg, 0.83 mmol, 60%) at rt, then stirred for 1 hr at rt under N₂. CH₃I (107 mg, 0.75 mmol) was added and stirred for 2 hr at rt, then MeOH was added to quench the reaction. The solvent was removed under vacuum and the residue was purified by flash chromatography on silica gel (DCM/MeOH=20/1) to afford compound 2 (243 mg, 62%). [M+H]⁺ Calc'd for C₂₀H₁₇F₃N₆O, 415; Found, 415.

Preparation 97B: 2-(5-bromo-1-methyl-1H-imidazol-4-yl)-4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine




[0298] To a mixture of 4-[1-benzyl-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-(1-methyl-1H-imidazol-4-yl)pyridine (243 mg, 0.58 mmol) in DCM (10 mL), was added NBS (104 mg, 0.58 mmol) at rt then stirred overnight. LC/MS showed the reaction was completed. The solvent was removed under vacuum and the residue was purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to afford the title compound (208 mg, 72%). [M+H]⁺ Calc'd for C₂₀H₁₆BrF₃N₆O, 493; Found, 493.

Example 97C: 2-[5-(4-fluorophenyl)-1-methyl-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0299] A mixture of 2-(5-bromo-1-methyl-1H-imidazol-4-yl)-4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine (100 mg, 0.20 mmol), 4-fluorophenyl boronic acid (57 mg, 0.41 mmol), Pd(dppf)Cl₂ (33 mg, 0.04 mmol) and Na₂CO₃ (0.2 mL, 0.40 mmol, 2 M) in dioxane (10 mL) was stirred for 2 hr at 140 °C in a microwave oven. After which the solvent was removed under vacuum and the residue was purified by flash column chromatography (DCM/MeOH =20/1) to give the PMB protected intermediate. This compound was dissolved in TFA/DCM (5/5 mL) and stirred overnight at 50 °C, concentrated, the residue was dissolved in NH₃/MeOH (10 mL), stirred for 20 min, then concentrated and purified by prep-TLC (DCM/MeOH/NH₃ =10/1/ 0.05) to give the title compound (22 mg, 28%). ¹H NMR (400 MHz, CD₃OD): δ 3.73 (3H, t), 7.39-7.45 (3H, m), 7.64-7.68 (2H, m), 7.75 (1H, *J* = 5.2Hz, *J* = 1.2Hz, dd), 8.82 (1H, *J* = 5.2Hz, d), 9.12 (1H, s). [M+H]⁺ Calc'd for C₁₈H₁₂F₄N₆, 389; Found, 389.

Example 98: 2-{1-[(3,4-dihydro-2H-1-benzopyran-4-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0300] The title compound was prepared in 5.6% yield according to the general procedure for the preparation of Example 95. ¹H NMR (400 MHz, CD₃OD): δ 1.89-2.10 (2H, m), 3.49-3.51 (1H, m), 4.22-4.26 (2H, m), 4.49-4.72 (2H, m), 6.82-6.88 (2H, m), 7.07-7.18 (2H, m), 7.75 (1H, dd, *J* = 5.2 Hz, *J* = 1.2 Hz), 8.25 (1H, s), 8.44 (1H, s), 8.83 (1H, d, *J* = 5.2 Hz), 9.07 (1H, s). [M+H]⁺ Calc'd for C₂₁H₁₇F₃N₆O, 427; Found, 427.

Example 99A: 5-fluoro-1,2,3,4-tetrahydronaphthalene-1-carboxylic acid




[0301] To a solution of 5-fluoro-1,2,3,4-tetrahydronaphthalen-1-one (1 g, 6.09 mmol) in DCM was added ZnI_2 (59 mg, 0.183 mmol), TMSCN (905 mg, 9.15 mmol) at rt, then stirred overnight at rt. aq NaHCO_3 was added, extracted with DCM, dried, concentrated. The residue was dissolved in HCl/AcOH (10/10 mL), $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ was added to the reaction and stirred for 4 days at 100°C . Then the mixture was cooled to rt and H_2O was added, the mixture was extracted with DCM. The organic layer was washed by 2N NaOH, the water layer was acidified to $\text{pH} < 3$, extracted with DCM, dried, concentrated to give the title compound (517 mg, 44%), which was used directly for the next step without further purification.

Example 99B: (5-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methanol




[0302] To a solution of 5-fluoro-1,2,3,4-tetrahydronaphthalene-1-carboxylic acid (517 mg, 2.66 mmol) in THF (10 mL) was added $\text{BH}_3 \cdot \text{THF}$ (14 mL, 14 mmol, 1M) at rt. The reaction mixture was refluxed for 6 hr, cooled to rt. MeOH was added slowly to the mixture, after concentration the residue was purified by flash column chromatography on silica gel (PE/EA=3/1) to afford the title compound (230 mg, 48%).

Example 99C: (5-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl methanesulfonate




[0303] To a mixture of (5-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methanol (230 mg, 1.27 mmol) and TEA (258 mg, 2.56 mmol) in DCM was added MsCl (220 mg, 1.91 mmol) at 0°C , then stirred for 2 hr at rt. After the reaction was completed, the mixture was washed by H_2O , dried, concentrated to give the title compound (320 mg, 100%), which was used directly for the next step.

Example 99: 2-{1-[(5-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0304] A mixture of 2-(1H-imidazol-4-yl)-4-{1-[(4-methoxyphenyl)methyl]-5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl}pyridine (100 mg, 0.25 mmol), (5-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl methanesulfonate (140 mg, 0.50 mmol) and Cs_2CO_3 (245 mg, 0.75 mmol) in DMF (10 mL) was stirred overnight at 100 °C under N_2 . The solvent was removed under vacuum and the residue was purified by flash column chromatography on silica gel (DCM/MeOH=20/1) to afford the PMB protected intermediate which was then dissolved in TFA/DCM (5/5 mL) and stirred overnight at 50 °C. After concentration, the residue was dissolved in NH_3/MeOH (10 mL), stirred for 20 min, then concentrated and purified by *prep*-TLC (DCM/MeOH/ NH_3 = 10/1/ 0.05) to give the title compound (13 mg, 11%). ^1H NMR (400 MHz, CD_3OD): δ 1.86-1.90 (4H, m), 2.84-2.87 (2H, m), 3.50-3.51 (1H, m), 4.45-4.61 (2H, m), 6.93-6.97 (2H, m), 7.13-7.16 (1H, m), 7.75 (1H, dd, $J = 5.2$ Hz, $J = 1.2$ Hz), 8.24 (1H, s), 8.41 (1H, s), 8.83 (1H, d, $J = 5.2$ Hz), 9.03 (1H, s). [M+H]⁺ Calc'd for $\text{C}_{22}\text{H}_{18}\text{F}_4\text{N}_6$, 443; Found, 443.

Example 100: 2-{1-[(6-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0305] The title compound was prepared in 23.5% yield according to the general procedure for the preparation of Example 99. ^1H NMR (300 MHz, CD_3OD): δ 1.71-1.92 (4H, m), 2.79-2.87 (2H, m), 3.46-3.47 (1H, m), 4.41-4.46 (2H, m), 6.84-6.90 (2H, m), 7.11-7.15 (1H, m), 7.75 (1H, d, $J = 5.1$ Hz), 8.24 (1H, s), 8.38 (1H, s), 8.82 (1H, d, $J = 5.1$ Hz), 9.01 (1H, s). [M+H]⁺ Calc'd for $\text{C}_{22}\text{H}_{18}\text{F}_4\text{N}_6$, 443; Found, 443.

Example 101: 2-{1-[(7-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0306] The title compound was prepared in 31% yield according to the general procedure for the preparation of Example 99. ^1H NMR (400 MHz, CD_3OD): δ 1.67-1.92 (4H, m), 2.75-2.82 (2H, m), 3.48-3.51 (1H, m), 4.43-4.65 (2H, m), 6.90-7.06 (2H, m), 7.14-7.18 (1H, m), 7.75 (1H, dd, $J = 5.2$ Hz, $J = 1.2$ Hz), 8.25 (1H, s), 8.42 (1H, s), 8.83 (1H, d, $J = 5.2$ Hz), 9.08 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{22}\text{H}_{18}\text{F}_4\text{N}_6$, 443; Found, 443.

Example 102: 2-{1-[(6-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine



[0307] The title compound was prepared in 30% yield according to the general procedure for the preparation of Example 99. ^1H NMR (300 MHz, CD_3OD): δ 1.68-1.94 (4H, m), 2.23 (3H, s), 2.70-2.82 (2H, m), 3.43-3.44 (1H, m), 4.41-4.60 (2H, m), 6.93-7.00 (3H, m), 7.73 (1H, d, $J = 5.1$ Hz), 8.22 (1H, s), 8.34 (1H, s), 8.83 (1H, d, $J = 5.1$ Hz), 9.00 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{23}\text{H}_{21}\text{F}_3\text{N}_6$, 439; Found, 439.

Example 103: 2-{1-[(7-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0308] The title compound was prepared in 27% yield according to the general procedure for the preparation of Example 99. ^1H NMR (400 MHz, CD_3OD): δ 1.68-1.94 (4H, m), 2.24 (3H, s), 2.73-2.77 (2H, m), 3.43-3.44 (1H, m), 4.43-4.59 (2H, m), 6.89 (1H, s), 6.98-7.04 (2H, m), 7.74

(1H, dd, $J = 5.2$ Hz, $J = 1.2$ Hz), 8.23 (1H, s), 8.34 (1H, s), 8.83 (1H, d, $J = 5.2$ Hz), 8.99 (1H, s).


[M+H] Calc'd for $C_{23}H_{21}F_3N_6$, 439; Found, 439.

Example 104: 2-{1-[(5-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0309] The title compound was prepared in 7% yield according to the general procedure for the preparation of Example 99. ^1H NMR (400 MHz, CD_3OD): δ 1.84-1.87 (4H, m), 2.50-2.86 (2H, m), 3.43-3.44 (1H, m), 3.81 (3H, s), 4.47-4.59 (2H, m), 6.67 (1H, d, $J = 8.0$ Hz), 6.82 (1H, d, $J = 8.0$ Hz), 7.09-7.12 (1H, m), 7.74 (1H, dd, $J = 5.2$ Hz, $J = 0.8$ Hz), 8.22 (1H, s), 8.35 (1H, s), 8.83 (1H, d, $J = 5.2$ Hz), 8.98 (1H, s). [M+H] Calc'd for $C_{23}H_{21}F_3N_6O$, 455; Found, 455.

Example 105: 2-{1-[(6-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0310] The title compound was prepared in 13% yield according to the general procedure for the preparation of Example 99. ^1H NMR (300 MHz, CD_3OD): δ 1.84-1.87 (4H, m), 2.50-2.86 (2H, m), 3.38-3.41 (1H, m), 3.76 (3H, s), 4.41-4.54 (2H, m), 6.69-6.72 (2H, m), 6.97-7.00 (1H, m), 7.73 (1H, d, $J = 5.1$ Hz), 8.21 (1H, s), 8.32 (1H, s), 8.81 (1H, d, $J = 5.1$ Hz), 8.96 (1H, s). [M+H] Calc'd for $C_{23}H_{21}F_3N_6O$, 455; Found, 455.

Example 106: 2-{1-[(7-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0311] The title compound was prepared in 13% yield according to the general procedure for the preparation of Example 99. ^1H NMR (300 MHz, CD_3OD): δ 1.66-1.87 (4H, m), 2.70-2.74 (2H, m), 3.42-3.44 (1H, m), 3.69 (3H, s), 4.41-4.61 (2H, m), 6.21-6.30 (1H, m), 6.74-6.78 (1H, m), 7.03-7.07 (1H, m), 7.74 (1H, d, $J = 5.1$ Hz), 8.22 (1H, s), 8.37 (1H, s), 8.82 (1H, d, $J = 5.1$ Hz), 8.99 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{23}\text{H}_{21}\text{F}_3\text{N}_6\text{O}$, 455; Found, 455.

Example 107: 2-{1-[(8-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine




[0312] The title compound was prepared in 4.4% yield according to the general procedure for the preparation of Example 99. ^1H NMR (400 MHz, CD_3OD): δ 1.78-1.90 (4H, m), 2.77-2.84 (2H, m), 3.66-3.67 (1H, m), 3.78 (3H, s), 4.35-4.59 (2H, m), 6.76-6.79 (2H, m), 7.17 (1H, t, $J = 8.0$ Hz), 7.74 (1H, d, $J = 5.2$ Hz), 8.24-8.25 (2H, m), 8.83 (1H, d, $J = 5.2$ Hz), 9.02 (1H, s). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{23}\text{H}_{21}\text{F}_3\text{N}_6\text{O}$, 455; Found, 455.

Example 108: 4-(5-fluoro-1H-1,2,3-triazol-4-yl)-2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine



[0313] The title compound was prepared in 0.5% overall yield according to the general procedure for the preparation of Example 35. ^1H NMR (400 MHz, CD_3OD): δ 3.25 (2H, t, $J = 6.8$ Hz), 4.53 (2H, t, $J = 6.8$ Hz), 7.04 (2H, t, $J = 8.4$ Hz), 7.22-7.25 (2H, m), 7.86 (1H, d, $J = 5.2$ Hz), 8.29 (1H, s), 8.33 (1H, s), 8.70-8.73 (2H, m). $[\text{M}+\text{H}]$ Calc'd for $\text{C}_{18}\text{H}_{14}\text{F}_2\text{N}_6$, 353; Found, 353.

Example 109: 4-(5-chloro-1H-1,2,3-triazol-4-yl)-2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine



[0314] The title compound was prepared in 0.5% overall yield according to the general procedure for the preparation of Example 36. ¹H NMR (400 MHz, CD₃OD): δ 3.25 (2H, t, *J* = 6.8 Hz), 4.54 (2H, t, *J* = 7.2 Hz), 7.04 (2H, t, *J* = 8.4 Hz), 7.22-7.25 (2H, m), 8.03 (1H, dd, *J* = 5.2 Hz, *J* = 1.6 Hz), 8.30 (1H, s), 8.41 (1H, s), 8.74-8.76 (2H, m). [M+H]⁺ Calc'd for C₁₈H₁₄ClFN₆, 369; Found, 369.

II. Biological Evaluation

Example 1: *In Vitro* Enzyme Inhibition Assay

[0315] This assay determines the ability of a test compound to inhibit FBXL11, FBXL10, and PHF8 demethylase activity. Baculovirus expressed FBXL11 (GenBank Accession #NM.012308, AA1-1162) was purchased from BPS Bioscience (Cat#50102). Baculovirus expressed FBXL10 (GenBank Accession #NM_032590, AA 1-650) was purchased from BPS Bioscience (Cat #50120). Baculovirus expressed PHF8 (GenBank Accession NP_055922.1) was purchased from Active Motif (Cat#31435).

FBXL11 Assay

[0316] The ability of test compounds to inhibit the activity of FBXL11 was determined in 384-well plate format under the following reaction conditions: 0.15 nM FBXL11, 30 nM H3K36me2-biotin labeled peptide (Anaspec cat # 64442), 0.2 μM alpha-ketoglutaric acid in assay buffer of 50 mM HEPES, pH7.3, 0.005% Brij35, 0.5 mM TCEP, 0.2 mg/ml BSA, 50 μM sodium L-ascorbate, 5 μM ammonium iron(II) sulfate. Reaction product is determined quantitatively by AlphaScreen detection after the addition of detection reagents anti-H3K36me1 antibody, AlphaScreen® Streptavidin-coated Donor beads, and AlphaScreen® Protein A Acceptor beads in 50 mM HEPES, pH7.3, 10 mM NaCl, 0.005% Brij35, 5 mM EDTA, 2 mg/ml BSA to final 10 μg/ml beads.

[0317] The assay reaction was initiated by the following: 3 μl of the mixture of 90 nM H3K36me2-biotin labeled peptide and 0.6 μM alpha-ketoglutaric acid with 3 μl of 11-point serial

diluted inhibitor in 3% DMSO are added to each well of 384 well Proxiplate (Perkin Elmer), followed by the addition of 3 μ l of 0.45 nM FBXL11 to initiate the reaction. The reaction mixture is incubated at room temperature for 1 hour, and terminated by the addition of 3 μ l of appropriate dilution of anti H3K36me1 antibody in 50 mM HEPES, pH7.3, 10 mM NaCl, 0.005% Brij35, 5 mM EDTA, 2 mg/ml BSA. Plates will then incubated at room temperature for 40 minutes, followed by addition of 3 μ l of 50 μ g/ml AlphaScreen® Streptavidin-coated Donor beads and AlphaScreen® Protein A Acceptor beads in 50 mM HEPES, pH7.3, 10 mM NaCl, 0.005% Brij35, 5 mM EDTA, 2 mg/ml BSA. Plates will then be read by EnVision Multilabel Reader in AlphaScreen mode after minimum 2 hour incubation at room temperature. The AlphaScreen signal for each well is used to determine inhibition constant (IC_{50}).

FBXL10 Assay

[0318] The ability of test compounds to inhibit the activity of FBXL10 was determined in 384-well plate format under the following reaction conditions: 0.3 nM FBXL10, 30 nM H3K36me2-biotin labeled peptide (Anaspec cat # 64442), 0.2 μ M alpha-ketoglutaric acid in assay buffer of 50 mM HEPES, pH7.3, 0.005% Brij35, 0.5 mM TCEP, 0.2 mg/ml BSA, 50 μ M sodium L-ascorbate, 5 μ M ammonium iron(II) sulfate. Reaction product is determined quantitatively by AlphaScreen detection after the addition of detection reagents anti-H3K36me1 antibody, AlphaScreen® Streptavidin-coated Donor beads, and AlphaScreen® Protein A Acceptor beads in 50 mM HEPES, pH7.3, 10 mM NaCl, 0.005% Brij35, 5 mM EDTA, 2 mg/ml BSA to final 10 μ g/ml beads.

[0319] The assay reaction was initiated by the following: 3 μ l of the mixture of 90 nM H3K36me2-biotin labeled peptide and 0.6 μ M alpha-ketoglutaric acid with 3 μ l of 11-point serial diluted inhibitor in 3% DMSO are added to each well of 384 well Proxiplate (Perkin Elmer), followed by the addition of 3 μ l of 0.9 nM FBXL10 to initiate the reaction. The reaction mixture is incubated at room temperature for 1 hour, and terminated by the addition of 3 μ l of appropriate dilution of anti H3K36me1 antibody in 50 mM HEPES, pH7.3, 10 mM NaCl, 0.005% Brij35, 5 mM EDTA, 2 mg/ml BSA. Plates will then incubated at room temperature for 40 minutes, followed by addition of 3 μ l of 50 μ g/ml AlphaScreen® Streptavidin-coated Donor beads and AlphaScreen® Protein A Acceptor beads in 50 mM HEPES, pH7.3, 10 mM NaCl, 0.005% Brij35, 5 mM EDTA, 2 mg/ml BSA. Plates will then be read by EnVision Multilabel Reader in AlphaScreen mode after minimum 2 hour incubation at room temperature. The AlphaScreen signal for each well is used to determine inhibition constant (IC_{50}).

PHF8 Assay

[0320] The ability of test compounds to inhibit the activity of PHF8 was determined in 384-well plate format under the following reaction conditions: 3 nM PHF8, 200 nM H3K9me1-biotin labeled peptide (Anaspec cat #64358), 0.5 μ M alpha-ketoglutaric acid in assay buffer of 50 mM HEPES, pH7.3, 0.005% Brij35, 0.5 mM TCEP, 0.2 mg/ml BSA, 50 μ M sodium L-ascorbate, and 5 μ M ammonium iron(II) sulfate. Reaction product was determined quantitatively by TR-FRET after the addition of detection reagent Phycolink Streptavidin-allophycocyanin (Prozyme) and Europium-anti-unmodified-histone H3 lysine 9/lysine27 (H3K9/K27) antibody (PerkinElmer) in the presence of 5 mM EDTA in LANCE detection buffer (PerkinElmer) at a final concentration of 25 nM and 0.5 nM, respectively.

[0321] The assay reaction was initiated by the following: 2 μ l of the mixture of 600 nM H3K9me1-biotin labeled peptide and 1.5 μ M alpha-ketoglutaric acid with 2 μ l of 11-point serial diluted inhibitor in 3% DMSO were added to each well of the plate, followed by the addition of 2 μ l of 9 nM PHF8 to initiate the reaction. The reaction mixture was incubated at room temperature for 15 minutes, and terminated by the addition of 6 μ l of 5 mM EDTA in LANCE detection buffer containing 50 nM Phycolink Streptavidin-allophycocyanin and 1 nM Europium-anti-unmodified H3K9/K27 antibody. Plates were read by EnVision Multilabel Reader in TR-FRET mode (excitation at 320 nm, emission at 615 nm and 665 nm) after 1 hour incubation at room temperature. A ratio was calculated (665/615) for each well and fitted to determine inhibition constant (IC_{50}).

[0322] The ability of the compounds disclosed herein to inhibit demethylase activity was quantified and the respective IC_{50} value was determined. Table 3 provides the IC_{50} values of various compounds disclosed herein.

TABLE 3

Chemical Synthesis Example	Name	FBXL10 IC_{50} (μ M)	FBXL11 IC_{50} (μ M)	PHF8 IC_{50} (μ M)
1	2-(1-methylimidazol-4-yl)-4-(1H-triazol-4-yl)pyridine	B		B
2	2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A	B	B

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	FBXL11 IC ₅₀ (μM)	PHF8 IC ₅₀ (μM)
3	2-[5-(4-fluorophenyl)-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A		B
4	2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A	A	C
5	2-[1-(1-phenylethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A	A	B
6	2-[1-[2-(2-chlorophenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A	A	A
7	2-[1-[2-(2-methoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A	A	B
8	2-[1-(1,2,3,4-tetrahydronaphthalen-1-ylmethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A	A	B
9	2-[1-[2-(2-ethoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A	A	B
10	4-(1H-triazol-4-yl)-2-[1-[2-[2-(2,2,2-trifluoroethoxy)phenyl]ethyl]imidazol-4-yl]pyridine	A		B
11	2-[1-[2-[2-(cyclopropylmethoxy)phenyl]ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A		B
12	2-[1-(3,4-dihydro-2H-chromen-4-ylmethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A		A

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	FBXL11 IC ₅₀ (μM)	PHF8 IC ₅₀ (μM)
13	4-[2-[1-[(2-chlorophenyl)methyl]imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile	A		C
14	4-[2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile	A		C
15	4-[2-[1-(1,2,3,4-tetrahydronaphthalen-1-ylmethyl)imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile	A		B
16	4-[2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile	A		B
17	2-[1-[2-(2-phenylmethoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A		C
18	2-[1-[2-(2-phenoxyphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A		C
19	2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-(5-iodo-1H-triazol-4-yl)pyridine	A		B
20	2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-(5-fluoro-1H-triazol-4-yl)pyridine	A		B
21	2-[1-[(2,3-dichlorophenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A	A	B

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	FBXL11 IC ₅₀ (μM)	PHF8 IC ₅₀ (μM)
22	2-[1-[[2-chloro-3-(trifluoromethyl)phenyl]methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A	A	B
23	2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A	A	B
24	2-[5-(4-fluoro-3-methoxyphenyl)-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A		B
25	2-[5-(3-ethoxy-4-fluorophenyl)-1-methylimidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A	B	C
26	2-[1-[[2-fluoro-3-(trifluoromethoxy)phenyl]methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	B		C
27	2-[1-[2-(2-phenylphenyl)ethyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	B		C
28	2-[1-[(2-fluoro-3-methylphenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A		B
29	2-[1-[(3-chloro-2-fluorophenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A		A
30	2-[1-[(2-fluoro-3-methoxyphenyl)methyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A		B

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	FBXL11 IC ₅₀ (μM)	PHF8 IC ₅₀ (μM)
31	4-(1H-triazol-4-yl)-2-[1-[2-[2-(trifluoromethyl)phenyl]ethyl]imidazol-4-yl]pyridine	A		B
32	2-[1-[2-(2-chlorophenyl)-2-methylpropyl]imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	A		B
33	2-[1-(1-phenylpropan-2-yl)imidazol-4-yl]-4-(1H-triazol-4-yl)pyridine	B		C
34	4-(1H-triazol-4-yl)-2-[1-[2-[2-(trifluoromethoxy)phenyl]ethyl]imidazol-4-yl]pyridine	A		B
35	4-(5-fluoro-1H-triazol-4-yl)-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine	A		B
36	4-(5-chloro-1H-triazol-4-yl)-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine	A		C
37	2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-triazol-4-yl]pyridine	A		D
38	4-(5-iodo-1H-triazol-4-yl)-2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]pyridine	A		D
39	4-[2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridin-4-yl]-1H-triazole-5-carbonitrile	A		C

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	FBXL11 IC ₅₀ (μM)	PHF8 IC ₅₀ (μM)
40	4-(5-chloro-1H-triazol-4-yl)-2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]pyridine	A		D
41	2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-(5-fluoro-1H-triazol-4-yl)pyridine	A		C
42	1-(cyclopropylmethyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	A		C
43	4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1-(2,2,2-trifluoroethyl)-1H-imidazole	B		C
44	1-benzyl-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	A	A	B
45	1-[(2-chlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	A	A	C
46	1-[(3-chlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	A	A	C
47	1-[(4-chlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	A	A	C
48	1-[(3,4-dichlorophenyl)methyl]-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	A	A	C

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	FBXL11 IC ₅₀ (μM)	PHF8 IC ₅₀ (μM)
49	1-(4-chlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	A	A	C
50	1-(2-chlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	A	A	C
51	1-(3-chlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	B	B	C
52	1-(3,5-dichlorophenyl)-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	B		C
53	5-(4-fluorophenyl)-1-methyl-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	B		C
54	5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methyl-4-{1H-[1,2,3]triazolo[4,5-c]pyridin-6-yl}-1H-imidazole	B		C
55	2-(5-bromo-1-(2-chlorobenzyl)-1H-imidazol-4-yl)-4-(2H-1,2,3-triazol-4-yl)pyridine	A		B

Note: Biochemical assay IC₅₀ data are designated within the following ranges:

A: ≤ 0.10 μM

B: > 0.10 μM to ≤ 1.0 μM

C: > 1.0 μM to ≤ 10 μM

D: > 10 μM

Example 2: *In Vitro* Cell-based Assay

Mia Paca-2 Pancreatic Cell line Expression Assay for KDM2B-related Genes

[0323] Gene expression assay to assess the ability of KDM2B small molecule inhibitors to activate the expression of KDM2B-bound genes of the established Mia Paca-2 cancer cell line.

Assay Background

[0324] The KDM2B protein has been shown to regulate the proliferation of AML, pancreatic and breast cancer. To demonstrate the specificity of small molecules for KDM2B's enzymatic activity, an assay to measure the relief of suppression caused by KDM2B in the established pancreatic cancer cell line model Mia Paca-2 was employed.

Assay Principle

[0325] This gene expression assay is a real-time PCR assay which quantifies the amount of messenger RNA from genes that are normally repressed by KDM2B interacting with EZH2 compared with a control GAPDH gene whose expression is not regulated by KDM2B. The amount of gene expression is correlated to the amount of small molecule inhibition of KDM2B's enzymatic activity.

Assay Method

[0326] The established cancer cell line Mia Paca-2 was purchased from *American Type Culture Collection* (ATCC) and routinely passaged according to ATCC published protocols. Cells were seeded at a density of 40,000 per 96-well. 24 hours after plating, cells received final concentrations of 20nM, 40nM, 80nM and 320nM of test compound. Time-matched control wells of 0.1% DMSO treatment were also included. Cells were incubated with test compound for 3, 6, 24 and 48hrs at 37 °C, 5% CO₂. At the end of each of the compound incubation period, media was removed and cells were trypsinized according to the ATCC published protocol. mRNA was then harvested from the cell samples using a RNeasy kit (Qiagen). mRNA was quantified and qualified using a Nanodrop machine (Thermo Scientific). mRNA was converted to cDNA using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems). Equal amounts of cDNA were subjected to real-time PCR using Taqman Universal Master Mix (Thermo Scientific). The following Taqman gene expression assays were used for detection on a Vii7 Real-time PCR system: Hs00224960_m1, Hs00164982_m1, Hs00907496_m1 and Hs02758991_g1 (Thermo Scientific). Data was analyzed using the delta-delta Ct method to calculate fold enrichment over DMSO sample gene expression.

[0327] Table 4 provides the cellular IC₅₀ values of various substituted heterocyclic compounds disclosed herein.

Table 4

Chemical Synthesis Example	Name	Cellular IC ₅₀ (μM)
37	2-[1-(2-naphthalen-1-ylethyl)imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-triazol-4-yl]pyridine	B
41	2-[5-[2-(cyclopropylmethoxy)-4-fluorophenyl]-1-methylimidazol-4-yl]-4-(5-fluoro-1H-triazol-4-yl)pyridine	C

Note: Cellular assay IC₅₀ data are designated within the following ranges:

A: ≤ 0.10 μM B: > 0.10 μM to ≤ 1.0 μM

C: > 1.0 μM to ≤ 10 μM D: > 10 μM

Example 3: *In Vitro* Enzyme Inhibition Assay

[0328] This assay determines the ability of a test compound to inhibit Jarid1B, JMJD2C and JMJD3 demethylase activity. Baculovirus expressed Jarid1B (GenBank Accession #NM—006618, AA 2-751) was purchased from BPS Bioscience (Cat #50121) or custom made by MolecularThroughput. Baculovirus expressed JMJD2C (GenBank Accession #BC143571, AA 2-372) was purchased from BPS Bioscience (Cat#50105). Baculovirus expressed JMJD3 (GenBank Accession #NM—001080424, AA1043-end) was purchased from BPS Bioscience (Cat#50115).

Jarid1B Assay

[0329] The ability of test compounds to inhibit the activity of Jarid1B was determined in 384-well plate format under the following reaction conditions: 0.8 nM Jarid1B, 300 nM H3K4me3-biotin labeled peptide (Anaspec cat #64357), 2 μM alpha-ketoglutaric acid in assay buffer of 50 mM HEPES, pH7.3, 0.005% Brij35, 0.5 mM TCEP, 0.2 mg/ml BSA, 50 μM sodium L-ascorbate, and 2 μM ammonium iron(II) sulfate. Reaction product was determined quantitatively by TR-FRET after the addition of detection reagent Phycolink Streptavidin-allophycocyanin (Prozyme) and Europium-anti-mono-or di-methylated histone H3 lysine 4 (H3K4me1-2) antibody (PerkinElmer) in the presence of 5 mM EDTA in LANCE detection buffer (PerkinElmer) at a final concentration of 25 nM and 1 nM, respectively.

[0330] The assay reaction was initiated by the following: 2 μ l of the mixture of 900 nM H3K4me3-biotin labeled peptide and 6 μ M alpha-ketoglutaric acid with 2 μ l of 11-point serial diluted inhibitor in 3% DMSO was added to each well of the plate, followed by the addition of 2 μ l of 2.4 nM Jarid1B to initiate the reaction. The reaction mixture was incubated at room temperature for 30 minutes, and terminated by the addition of 6 μ l of 5 mM EDTA in LANCE detection buffer containing 50 nM Phycolink Streptavidin-allophycocyanin and 2 nM Europium-anti-H3K4me1-2 antibody. Plates were read by EnVisionMultilabel Reader in TR-FRET mode (excitation at 320 nm, emission at 615 nm and 665 nm) after 1 hour incubation at room temperature. A ratio was calculated (665/615) for each well and fitted to determine inhibition constant (IC_{50}).

JMJD2C Assay

[0331] The ability of test compounds to inhibit the activity of JMJD2C was determined in 384-well plate format under the following reaction conditions: 0.3 nM JMJD2C, 300 nM H3K9me3-biotin labeled peptide (Anaspec cat #64360), 2 μ M alpha-ketoglutaric acid in assay buffer of 50 mM HEPES, pH7.3, 0.005% Brij35, 0.5 mM TCEP, 0.2 mg/ml BSA, 50 μ M sodium L-ascorbate, and 2 μ M ammonium iron(II) sulfate. Reaction product was determined quantitatively by TR-FRET after the addition of detection reagent Phycolink Streptavidin-allophycocyanin (Prozyme) and Europium-anti-di-methylated histone H3 lysine 9 (H3K9me2) antibody (PerkinElmer) in the presence of 5 mM EDTA in LANCE detection buffer (PerkinElmer) at a final concentration of 50 nM and 1 nM, respectively.

[0332] The assay reaction was initiated by the following: 2 μ l of the mixture of 900 nM H3K9me3-biotin labeled peptide and 6 μ M alpha-ketoglutaric acid with 2 μ l of 11-point serial diluted inhibitor in 3% DMSO were added to each well of the plate, followed by the addition of 2 μ l of 0.9 nM JMJD2C to initiate the reaction. The reaction mixture was incubated at room temperature for 30 minutes, and terminated by the addition of 6 μ l of 5 mM EDTA in LANCE detection buffer containing 100 nM Phycolink Streptavidin-allophycocyanin and 2 nM Europium-anti-H3K9me2 antibody. Plates were read by EnVisionMultilabel Reader in TR-FRET mode (excitation at 320 nm, emission at 615 nm and 665 nm) after 1 hour incubation at room temperature. A ratio was calculated (665/615) for each well and fitted to determine inhibition constant (IC_{50}).

JMJD3 Assay

[0333] The ability of test compounds to inhibit the activity of JMJD3 was determined in 384-well plate format under the following reaction conditions: 1 nM JMJD3, 250 nM H3K27me3-biotin labeled peptide (Anaspec cat # 64367), 1 μ M alpha-ketoglutaric acid in assay buffer of 50 mM HEPES, pH7.3, 0.005% Brij35, 0.5 mM TCEP, 0.2 mg/ml BSA, 50 μ M sodium

L-ascorbate, 5 μ M ammonium iron(II) sulfate. Reaction product is determined quantitatively by AlphaScreen detection after the addition of detection reagents anti-H3K27me1 antibody, 5 μ g/ml AlphaScreen® Streptavidin-coated Donor beads, and 5 μ g/ml AlphaScreen® Protein A Acceptor beads in 50 mM HEPES, pH7.3, 10 mM NaCl, 0.005% Brij35, 10 mM EDTA, 2 mg/ml BSA.

[0334] The assay reaction was initiated by the following: 3 μ l of the mixture of 750 nM H3K27me3-biotin labeled peptide and 3 μ M alpha-ketoglutaric acid with 3 μ l of 11-point serial diluted inhibitor in 3% DMSO are added to each well of 384 well Proxiplate (Perkin Elmer), followed by the addition of 3 μ l of 3 nM JMJD3 to initiate the reaction. The reaction mixture is incubated at room temperature for 20 minutes, and terminated by the addition of 3 μ l of appropriate dilution of anti H3K27me1 antibody in 50 mM HEPES, pH7.3, 10 mM NaCl, 0.005% Brij35, 10 mM EDTA, 2 mg/ml BSA. Plates will then incubated at room temperature for 1 hour, followed by addition of 3 μ l of 25 μ g/ml AlphaScreen® Streptavidin-coated Donor beads and AlphaScreen® Protein A Acceptor beads in 50 mM HEPES, pH7.3, 10 mM NaCl, 0.005% Brij35, 10 mM EDTA, 2 mg/ml BSA. Plates will then be read by EnVision Multilabel Reader in AlphaScreen mode after minimum 2 hour incubation at room temperature. The AlphaScreen signal for each well is used to determine inhibition constant (IC_{50}).

[0335] The ability of the compounds disclosed herein to inhibit demethylase activity was quantified and the respective IC_{50} value was determined. Table 5 provides the IC_{50} values of various compounds disclosed herein.

TABLE 5

Chemical Synthesis Example	Name	FBXL10 IC_{50} (μ M)	JARID1B IC_{50} (μ M)	JMJD2C IC_{50} (μ M)	JMJD3 IC_{50} (μ M)
57	2-{1-[2-(2-chlorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	D
58	2-{1-[(1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	D

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μ M)	JARID1B IC ₅₀ (μ M)	JMJD2C IC ₅₀ (μ M)	JMJD3 IC ₅₀ (μ M)
59	2-{1-[2-(2-ethoxyphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	A	D	D
60	4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-(1-{2-[2-(trifluoromethyl)phenyl]ethyl}-1H-imidazol-4-yl)pyridine	B	A	C	
61	2-{1-[(4-fluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
62	2-{1-[(4-fluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
63	2-{1-[(4-chlorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
64	2-{1-[(2-chlorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
65	2-{1-[(4-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	D	D	D
66	2-{1-[(2-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	JARID1B IC ₅₀ (μM)	JMJD2C IC ₅₀ (μM)	JMJD3 IC ₅₀ (μM)
67	2-{1-[(4-methoxyphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	C	D
68	2-{1-[(2,4-difluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
69	2-{1-[(2,6-difluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
70	2-{1-[(4-fluoro-2-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
71	2-{1-[(2-fluoro-4-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
72	2-(1-benzyl-1H-imidazol-4-yl)-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
73	2-[1-(1-phenylethyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	JARID1B IC ₅₀ (μM)	JMJD2C IC ₅₀ (μM)	JMJD3 IC ₅₀ (μM)
74	2-[1-(1-phenylpropyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
75	2-[1-(2-methyl-1-phenylpropyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	B	D	D
76	2-(1-{[2-(2,2,2-trifluoroethoxy)phenyl]methyl}-1H-imidazol-4-yl)-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	B	D	D
77	2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	C
78	2-{1-[2-(3-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	C
79	2-{1-[2-(2,3-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	D
80	2-{1-[2-(2-chloro-6-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	B	D	D

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	JARID1B IC ₅₀ (μM)	JMJD2C IC ₅₀ (μM)	JMJD3 IC ₅₀ (μM)
81	4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-{1-[2-(2,4,5-trifluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine	A	A	D	D
82	4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-{1-[2-(2,4,6-trifluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine	A	B	D	D
83	2-{1-[(2,3-dihydro-1H-inden-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	A	C	
84	2-{1-[2-(2-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	
85	2-[1-(6-fluoro-2,3-dihydro-1H-inden-1-yl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	C	
86	2-{1-[2-(2,4-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	D
87	2-{1-[2-(2,6-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	JARID1B IC ₅₀ (μM)	JMJD2C IC ₅₀ (μM)	JMJD3 IC ₅₀ (μM)
88	2-{1-[2-(2,5-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	D
89	2-{1-[(3R)-1-benzoylpiperidin-3-yl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	D
90	2-{1-[(3S)-1-benzoylpiperidin-3-yl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	D
91	2-{1-[2-(4-methylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
92	2-{1-[2-(4-chlorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	B	D	D
93	2-{1-[2-(4-methoxyphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	D
94	4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-(1-{2-[4-(trifluoromethyl)phenyl]ethyl}-1H-imidazol-4-yl)pyridine	B	B	D	D

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	JARID1B IC ₅₀ (μM)	JMJD2C IC ₅₀ (μM)	JMJD3 IC ₅₀ (μM)
95	2-{1-[2-(4-ethylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	B	D	D
96	2-{1-[2-(4-tert-butylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	B	D	D
97	2-[5-(4-fluorophenyl)-1-methyl-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	D
98	2-{1-[(3,4-dihydro-2H-1-benzopyran-4-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	A	A	D	D
99	2-{1-[(5-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	B	D	D
100	2-{1-[(6-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	A	D	D

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μM)	JARID1B IC ₅₀ (μM)	JMJD2C IC ₅₀ (μM)	JMJD3 IC ₅₀ (μM)
101	2-{1-[(7-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	A	D	D
102	2-{1-[(6-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	C	B	D	D
103	2-{1-[(7-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	A	D	D
104	2-{1-[(5-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	B	D	D
105	2-{1-[(6-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	B	D	D

Chemical Synthesis Example	Name	FBXL10 IC ₅₀ (μ M)	JARID1B IC ₅₀ (μ M)	JMJD2C IC ₅₀ (μ M)	JMJD3 IC ₅₀ (μ M)
106	2-{1-[(7-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	A	D	D
107	2-{1-[(8-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine	B	B	D	D

Note: Biochemical assay IC₅₀ data are designated within the following ranges:

A: $\leq 0.10 \mu\text{M}$

B: $> 0.10 \mu\text{M}$ to $\leq 1.0 \mu\text{M}$

C: $> 1.0 \mu\text{M}$ to $\leq 10 \mu\text{M}$

D: $> 10 \mu\text{M}$

III. Preparation of Pharmaceutical Dosage Forms


Example 1: Oral Tablet

[0336] A tablet is prepared by mixing 48% by weight of a compound of Formula (I) or (III), or a pharmaceutically acceptable salt thereof, 45% by weight of microcrystalline cellulose, 5% by weight of low-substituted hydroxypropyl cellulose, and 2% by weight of magnesium stearate. Tablets are prepared by direct compression. The total weight of the compressed tablets is maintained at 250-500 mg.

CLAIMS

We Claim:

1. A compound having the structure of Formula (Ic),



or a pharmaceutically acceptable salt thereof, wherein

R^1 is hydrogen, halogen, -OH, -OR⁵, -N(R⁵)₂, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclylalkyl, heterocyclylalkyl, aralkyl, or heteroarylalkyl;

R^2 is alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclylalkyl, heterocyclylalkyl, aralkyl, or heteroarylalkyl;

R^3 is hydrogen;

R^4 is hydrogen, C₁-C₄alkyl, C₁-C₄alkene, C₁-C₄alkyne, halogen, or -CN; and

each R^5 is independently hydrogen, alkyl, carbocyclyl, heterocyclyl, aryl, heteroaryl, carbocyclylalkyl, heterocyclylalkyl, aralkyl, or heteroarylalkyl.

2. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein R^2 is alkyl.

3. The compound of claim 2, or a pharmaceutically acceptable salt thereof, wherein R^2 is methyl.

4. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein R^2 is carbocyclyl or carbocyclylalkyl.

5. The compound of claim 4, or a pharmaceutically acceptable salt thereof, wherein R^2 is carbocyclyl, and the carbocyclyl is 1,2,3,4-tetrahydronaphthyl optionally substituted with at least one of halogen, hydroxy, -CN, alkyl, alkoxy, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, or heterocyclylalkyl.

6. The compound of claim 4, or a pharmaceutically acceptable salt thereof, wherein R^2 is $-(C_1-C_6\text{alkylene})\text{carbocyclyl}$.
7. The compound of claim 6, or a pharmaceutically acceptable salt thereof, wherein the $(C_1-C_6\text{alkylene})$ is a $C_1\text{alkylene}$, or a $C_2\text{alkylene}$.
8. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein R^2 is heterocyclyl or heterocyclylalkyl.
9. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein R^2 is heteroaryl or heteroarylalkyl.
10. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein R^2 is aryl or aralkyl.
11. The compound of claim 10, or a pharmaceutically acceptable salt thereof, wherein the aralkyl is $-(C_1-C_6\text{alkylene})\text{aryl}$.
12. The compound of claim 11, or a pharmaceutically acceptable salt thereof, wherein $(C_1-C_6\text{alkylene})$ is $C_1\text{alkylene}$, or a $C_2\text{alkylene}$.
13. The compound of claim 10, or a pharmaceutically acceptable salt thereof, wherein the aryl is phenyl or naphthyl, wherein said phenyl or naphthyl is optionally substituted with at least one of halogen, hydroxy, $-\text{CN}$, alkyl, alkoxy, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, or heterocyclylalkyl.
14. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein R^1 is hydrogen or aryl.
15. The compound of claim 14, or a pharmaceutically acceptable salt thereof, wherein R^1 is an aryl, and wherein the aryl is a phenyl optionally substituted with at least one of halogen, hydroxy, $-\text{CN}$, alkyl, alkoxy, $-\text{O}-(\text{cycloalkylalkyl})$, alkylamino, aryl, carbocyclyl, heterocyclyl, carbocyclylalkyl, or heterocyclylalkyl.

16. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein R⁴ is C₁-C₄alkyl, and the alkyl is substituted with at least one fluoro.

17. The compound of claim 16, or a pharmaceutically acceptable salt thereof, wherein R⁴ is CH₂F, CHF₂, or CF₃.

18. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein said compound is selected from the group consisting of

2-{1-[2-(2-chlorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(2-ethoxyphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-(1-{2-[2-(trifluoromethyl)phenyl]ethyl}-1H-imidazol-4-yl)pyridine,

2-{1-[(4-fluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(4-fluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(4-chlorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(2-chlorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(4-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(2-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(4-methoxyphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(2,4-difluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(2,6-difluorophenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(4-fluoro-2-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(2-fluoro-4-methylphenyl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-(1-benzyl-1H-imidazol-4-yl)-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-[1-(1-phenylethyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-[1-(1-phenylpropyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-[1-(2-methyl-1-phenylpropyl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-(1-{[2-(2,2,2-trifluoroethoxy)phenyl]methyl}-1H-imidazol-4-yl)-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(3-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(2,3-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(2-chloro-6-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-{1-[2-(2,4,5-trifluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine,

4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-{1-[2-(2,4,6-trifluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine,

2-{1-[(2,3-dihydro-1H-inden-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(2-fluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-[1-(6-fluoro-2,3-dihydro-1H-inden-1-yl)-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(2,4-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(2,6-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(2,5-difluorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(3R)-1-benzoylpiperidin-3-yl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(3S)-1-benzoylpiperidin-3-yl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(4-methylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(4-chlorophenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(4-methoxyphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]-2-(1-{2-[4-(trifluoromethyl)phenyl]ethyl}-1H-imidazol-4-yl)pyridine,

2-{1-[2-(4-ethylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[2-(4-tert-butylphenyl)ethyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-[5-(4-fluorophenyl)-1-methyl-1H-imidazol-4-yl]-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(3,4-dihydro-2H-1-benzopyran-4-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(5-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(6-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(7-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,

2-{1-[(6-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,
 2-{1-[(7-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,
 2-{1-[(5-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,
 2-{1-[(6-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,
 2-{1-[(7-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,
 2-{1-[(8-methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl]-1H-imidazol-4-yl}-4-[5-(trifluoromethyl)-1H-1,2,3-triazol-4-yl]pyridine,
 4-(5-fluoro-1H-1,2,3-triazol-4-yl)-2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine, and
 4-(5-chloro-1H-1,2,3-triazol-4-yl)-2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine.

19. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein said compound is selected from the group consisting of 4-(5-fluoro-1H-1,2,3-triazol-4-yl)-2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine, and 4-(5-chloro-1H-1,2,3-triazol-4-yl)-2-{1-[2-(4-fluorophenyl)ethyl]-1H-imidazol-4-yl}pyridine.

20. A pharmaceutical composition comprising the compound of claim 1, or a pharmaceutically acceptable salt thereof.

21. A method for inhibiting histone demethylase enzyme comprising contacting the histone demethylase enzyme with a compound of claim 1.

22. A method for treating cancer in a subject in need thereof comprising administering to the subject a pharmaceutical composition comprising the compound of claim 1, or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 17/22570

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - C07D 401/06 (2017.01)

CPC - C07D 401/14, C07D 401/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History Document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

See Search History Document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History Document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2014/275084 A1 (Kanouni et al.) 18 September 2014 (18.09.2014); para [0023], [0035], [0136], claim 28	1-15, 20-22
--		-----
A		16-19
A	US 2015/0038534 A1 (Baloglu et al.) 05 February 2015 (05.02.2015); para [0002], [0414]	1-22
A	US 2016/0009711 A1 (Wu et al.) 14 January 2016 (14.01.2016); para [0268], abstract	1-22
A	US 9,107,916 B2 (Nie et al.) 18 August 2015 (18.08.2015); entire document	1-22
P, X	US 2016/0108033 A1 (Chen et al.) 21 April 2016 (21.04.2016); entire document	1-22

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

04 May 2017

Date of mailing of the international search report

07 JUN 2017

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774