



US010668094B2

(12) **United States Patent**
Karlish et al.

(10) **Patent No.:** **US 10,668,094 B2**
(45) **Date of Patent:** **Jun. 2, 2020**

(54) **SELECTIVE INHIBITORS OF ALPHA2-CONTAINING ISOFORMS OF NA,K-ATPASE AND USE THEREOF FOR REDUCTION OF INTRAOCULAR PRESSURE**

3,850,578 A 11/1974 McConnell
3,850,752 A 11/1974 Schuurs et al.
3,853,987 A 12/1974 Dreyer
3,867,517 A 2/1975 Ling et al.
3,879,262 A 4/1975 Schuurs et al.
3,901,654 A 8/1975 Gross
3,925,355 A * 12/1975 Piasio C07J 43/003
536/6.2

(71) Applicant: **Yeda Research and Development Co. Ltd.**, Rehovot (IL)

3,935,074 A 1/1976 Rubinstein et al.
3,984,533 A 10/1976 Uzgiris
3,996,345 A 12/1976 Ullman et al.
4,034,074 A 7/1977 Miles
4,098,876 A 7/1978 Piasio et al.
4,190,496 A 2/1980 Rubenstein et al.

(72) Inventors: **Steven J. D. Karlish**, Rehovot (IL); **Adriana Katz**, Rehovot (IL); **Daniel M. Tal**, Rehovot (IL); **Arie Marcovich**, Rehovot (IL)

4,666,828 A 5/1987 Gusella
4,683,202 A 7/1987 Mullis
4,801,531 A 1/1989 Frossard
4,879,219 A 11/1989 Wands et al.

(73) Assignee: **Yeda Research and Development Co. Ltd.**, Rehovot (IL)

5,011,771 A 4/1991 Bellet et al.
5,192,659 A 3/1993 Simons
5,272,057 A 12/1993 Smulson et al.
5,281,521 A 1/1994 Trojanowski et al.
5,972,630 A 10/1999 Cromer et al.

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 115 days.

9,938,316 B2 * 4/2018 Karlish C07J 43/003
2003/0207431 A1 11/2003 Ghoshal et al.
2004/0210044 A1 10/2004 Slattum et al.
2008/0199895 A1 8/2008 Karlish et al.
2016/0244479 A1 8/2016 Karlish et al.

(21) Appl. No.: **15/745,441**

(22) PCT Filed: **Jul. 19, 2016**

(86) PCT No.: **PCT/IL2016/050785**

§ 371 (c)(1),
(2) Date: **Jan. 17, 2018**

(87) PCT Pub. No.: **WO2017/013648**

PCT Pub. Date: **Jan. 26, 2017**

(65) **Prior Publication Data**

US 2018/0207189 A1 Jul. 26, 2018

FOREIGN PATENT DOCUMENTS

DE e 224037 6/1985
EP 0532187 3/1993
FR 2262670 9/1975
WO WO 98/52961 11/1998
WO WO 2006/044916 4/2006
WO WO 2007/016656 2/2007
WO WO 2015/029035 3/2015
WO WO 2017/013637 1/2017
WO WO 2017/013648 1/2017

Related U.S. Application Data

(63) Continuation-in-part of application No. PCT/IL2015/050741, filed on Jul. 19, 2015.

(60) Provisional application No. 62/302,226, filed on Mar. 2, 2016.

(51) **Int. Cl.**

C07H 17/00 (2006.01)
C07J 43/00 (2006.01)
C07J 19/00 (2006.01)
A61K 31/7052 (2006.01)
A61P 27/06 (2006.01)
A61P 9/00 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 31/7052** (2013.01); **A61P 9/00** (2018.01); **A61P 27/06** (2018.01); **C07H 17/00** (2013.01); **C07J 43/003** (2013.01); **C07J 19/005** (2013.01)

(58) **Field of Classification Search**

None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,791,932 A 2/1974 Schuurs et al.
3,839,153 A 10/1974 Schuurs et al.

Wolff, Manfred E. "Burger's Medicinal Chemistry, 5ed, Part I", John Wiley & Sons, 1995, pp. 975-977 (Year: 1995).
Banker, G.S. et al, "Modern Pharmaceuticals, 3ed.", Marcel Dekker, New York, 1996, p. 596 (Year: 1996).
Adamczyk et al., Journal of Organic Chemistry, 1995, vol. 60(11), pp. 3557-3560 (Year: 1995).
Hara et al., Journal of Labelled Compounds and Radiopharmaceuticals—1977, vol. 13(2), pp. 222-223 (Year: 1977).
Communication Relating to the Results of the Partial International Search dated Oct. 17, 2016 From the International Searching Authority Re. Application No. PCT/IL2016/050785.
Communication Relating to the Results of the Partial International Search dated Oct. 27, 2015 From the International Searching Authority Re. Application No. PCT/IL2015/050741.

(Continued)

Primary Examiner — Traviss C McIntosh, III

(57) **ABSTRACT**

Provided herein are alpha2-selective Na,K-ATPase inhibitors and prodrugs thereof, characterized by having a cyclic moiety attached to a digoxin or digitoxin derivative, as well as uses thereof in lowering intraocular pressure and in treating glaucoma and heart conditions.

17 Claims, 4 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

- International Preliminary Report on Patentability dated Feb. 1, 2018 From the International Bureau of WIPO Re. Application No. PCT/IL2015/050741. (16 Pages).
- International Preliminary Report on Patentability dated Feb. 1, 2018 From the International Bureau of WIPO Re. Application No. PCT/IL2016/050785. (15 Pages).
- International Preliminary Report on Patentability dated Mar. 10, 2016 From the International Bureau of WIPO Re. Application No. PCT/IL2014/050773.
- International Search Report and the Written Opinion dated Nov. 18, 2014 From the International Searching Authority Re. Application No. PCT/IL2014/050773.
- International Search Report and the Written Opinion dated Jan. 22, 2016 From the International Searching Authority Re. Application No. PCT/IL2015/050741.
- International Search Report and Written Opinion dated Dec. 19, 2016 From the International Searching Authority Re. Application No. PCT/IL2016/050785. (23 Pages).
- Notice of Allowance dated Jan. 11, 2018 From the US Patent and Trademark Office Re. U.S. Appl. No. 14/913,366. (7 pages).
- Office Action and Search Report dated Oct. 17, 2016 From the State Intellectual Property Office of the People's Republic of China Re. Application No. 201480059614.5 and Its Translation Into English. Office Action dated May 2, 2017 From the State Intellectual Property Office of the People's Republic of China Re. Application No. 201480059614.5 and Its Translation Into English. (10 Pages).
- Official Action dated Dec. 1, 2009 From the US Patent and Trademark Office Re. U.S. Appl. No. 11/905,833.
- Official Action dated Apr. 2, 2010 From the US Patent and Trademark Office Re. U.S. Appl. No. 11/905,833.
- Official Action dated Jul. 14, 2017 From the US Patent and Trademark Office Re. U.S. Appl. No. 14/913,366. (28 Pages).
- Restriction Official Action dated Apr. 5, 2017 From the US Patent and Trademark Office Re. U.S. Appl. No. 14/913,366. (12 pages).
- Supplementary European Search Report and the European Search Opinion dated Mar. 3, 2017 From the European Patent Office Re. Application No. 14839412.5. (10 Pages).
- Adameczyk et al. "Digoxin Dialdehyde Reductive Aminations. Structure Proof of the Perhydro-1,4-Oxazepine Product", *Steroids*, XP004026467, 60(11): 753-758, Nov. 1, 1995. Compound 3, Preparation Thereof in Scheme 1, p. 754, 1st col.
- Adameczyk et al. "Unexpectedly Facile Hydrolysis of Digoxin Esters. The Importance of Appropriate Controls in Lipase-Mediated Hydrolysis", *The Journal of Organic Chemistry*, XP055221120, 60(11): 3557-3560, Jun. 1, 1995. Table 1, Compounds 7a-7e, Preparation Thereof on p. 3559, 1st col.
- Bachrach et al. "Attachment of Drugs to Polydimethylsiloxanes", *European Polymer Journal*, XP024053540, 20(5): 493-500, Jan. 1984. Abstract, p. 498, col. 1, Para 1-2, Compound I.
- Bers "Cardiac Excitation-Contraction Coupling", *Nature*, 415: 198-205, Jan. 10, 2002.
- Bitter "Heterologous Gene Expression in Yeast", *Methods in Enzymology*, 152(Chap.70): 673-684, 1987.
- Bitter et al. "Expression and Secretion Vectors for Yeast", *Methods in Enzymology*, 153(Chap.33): 516-544, 1987.
- Cohen et al. "Purification of Na⁺,K⁺-ATPase Expressed in *Pichia Pastoris* Reveals an Essential Role of Phospholipid-Protein Interactions", *The Journal of Biological Chemistry*, 280(17): 16610-16618, Apr. 29, 2005.
- Cornelius "Cholesterol Modulation of Molecular Activity of Reconstituted Shark Na⁺,K⁺-ATPase", *Biochimica et Biophysica Acta*, 1235: 205-212, 1995.
- Cornelius "Modulation of Na⁺,K⁺-ATPase and Na-ATPase Activity by Phospholipids and Cholesterol. I. Steady-State Kinetics", *Biochemistry*, 40(30): 8842-8851, 2001.
- Crambert et al. "New Molecular Determinants Controlling the Accessibility of Ouabain to Its Binding Site in Human Na⁺,K⁺-ATPase α Isoforms", *Molecular Pharmacology*, 65(2): 335-341, 2004.
- Crambert et al. "Transport and Pharmacological Properties of Nine Different Human Na⁺,K⁺-ATPase Isozymes", *The Journal of Biological Chemistry*, 275(3): 1976-1986, Jan. 21, 2000.
- Ferraiolo et al. "Digoxin-Induced Decrease in Intraocular Pressure in the Cat", *European Journal of Pharmacology*, XP023839155, 55(1): 19-22, Apr. 1, 1979. Abstract.
- Ferrari et al. "PST2238: A New Antihypertensive Compound That Antagonizes the Long-Term Pressor Effect of Ouabain", *The Journal of Pharmacology and Experimental Therapeutics*, JPET, 285(1): 83-94, 1998.
- Ferrari et al. "Rostafuroxin: An Ouabain Antagonist That Corrects Renal and Vascular Na⁺-K⁺-ATPase Alterations in Ouabain and Aldosterone-Dependent Hypertension", *American Journal of Physiology—Regulatory, Integrative and Comparative Physiology*, 290: R529-R535, 2006.
- Hara et al. "Heart Uptake of Radioiodine Labeled Cardenolides", *Journal of Labelled Compounds and Radiopharmaceuticals*, XP055348816, 13(2): 222-223, Jan. 1977. p. 222, Para 3-5, p. 223, Compound I.
- Hardt et al. "Effect of Systemic Digitalis Application on Intraocular Pressure [Zum Einfluss von Systemischer Digitalis-Applikation auf den Intraocularen Druck]", *Graefes's Archive for Clinical and Experimental Ophthalmology*, 219(2): 76-79, Aug. 1982. Abstract.
- Hartmanis et al. "Solubilization of a Membrane-Bound Diol Dehydratase With Retention of EPR G=2.02 Signal by Using 2-(N-Cyclohexylamino)Ethanesulfonic Acid Buffer", *Proc. Natl. Acad. Sci. USA*, 84: 76-79, Jan. 1987.
- Haviv et al. "Stabilization of Na⁺,K⁺-ATPase Purified From *Pichia Pastoris* Membranes by Specific Interactions With Lipids", *Biochemistry*, 46: 12855-12867, Published on Web Oct. 16, 2007.
- Hayashi et al. "Minimum Enzyme Unit for Na⁺/K⁺-ATPase is the $\alpha\beta$ -Promoter. Determination by Low-Angle Laser Light Scattering Photometry Coupled With High-Performance Gel Chromatography for Substantially Simultaneous Measurement of ATPase Activity and Molecular Weight", *Biochimica et Biophysica Acta*, 983: 217-229, 1989.
- Jørgensen "Purification of Na⁺,K⁺-ATPase: Enzyme Sources, Preparative Problems, and Preparation From Mammalian Kidney", *Methods in Enzymology*, 156(2): 29-43, 1988.
- Jørgensen et al. "Role of Conserved TGDGVND-Loop in Mg²⁺ Binding, Phosphorylation, and Energy Transfer in Na⁺,K⁺-ATPase", *Journal of Bioenergetics and Biomembranes*, 35(5): 367-377, Oct. 2001.
- Juhaszova et al. "Distinct Distribution of Different Na⁺ Pump α Subunit Isoforms in Plasmalemma", *Annals New York Academy of Sciences*, 834: 524-536, 1997.
- Kapri-Pardes et al. "Stabilization of the Alpha2 Isoform of Na⁺,K⁺-ATPase by Mutations in a Phospholipid Binding Pocket", *The Journal of Biological Chemistry*, 286(50): 42888-42899, Dec. 16, 2011.
- Katz et al. "Digoxin Derivatives With Enhanced Selectivity for the Alpha2 Isoform of Na⁺,K⁺-ATPase. Effects on Intraocular Pressure in Rabbits", *The Journal of Biological Chemistry*, XP055241264, 289(30): 21153-21162, Jul. 25, 2014. Abstract, p. 21153, Left col., Para 2, p. 21155, Right col., Lines 17-21, p. 21162, Right col., Para 2, p. 15, Table 1, p. 21157, Table 1, Table S1.
- Katz et al. "Selectivity of Digitalis Glycosides for Isoforms of Human Na⁺,K⁺-ATPase", *The Journal of Biological Chemistry*, 285(25): 19582-19592, JBC Papers in Press Apr. 13, 2010.
- Keating et al. "Potentiometric Digoxin Antibody Measurements With Antigen-Ionophore Based Membrane Electrodes", *Analytical Chemistry*, XP055221088, 56(4): 801-806, Apr. 1, 1984. Fig.2, Final Compound, Compounds Analyzed in Figs.3-4, Preparation.
- Laursen et al. "Structures and Characterization of Digoxin- and Bufalin-Bound Na⁺, K⁺-ATPase Compared With the Ouabain-Bound Complex", *Proc. Natl. Acad. Sci. USA*, PNAS, 112(6): 1755-1760, Feb. 10, 2015.
- Lifshitz et al. "Functional Interactions of Phospholemman (PLM) (FXFD1) With Na⁺,K⁺-ATPase. Purification of $\alpha 1/\beta 1$ /PLM Complexes Expressed in *Pichia Pastoris*", *The Journal of Biological Chemistry*, 281(23): 15790-15799, Jun. 9, 2006.

(56)

References Cited

OTHER PUBLICATIONS

Lifshitz et al. "Purification of the Human Alpha2 Isoform of Na,K-ATPase Expressed in *Pichia Pastoris*. Stabilization by Lipids and FXYD1", *Biochemistry*, 46: 14937-14950, Published on Web Dec. 2, 2007.

Lingrel "NA,K-ATPase: Isoform Structure, Function, and Expression", *Journal of Bioenergetics and Biomembrane*, 24(3): 263-270, 1992.

Mishra et al. "FXYD Proteins Stabilize Na,K-ATPase. Amplification of Specific Phosphatidylserine-Protein Interactions", *The Journal of Biological Chemistry*, 286(11): 9699-9712, Mar. 18, 2011.

Müller-Ehmsen et al. "Ouabain and Substrate Affinities of Human Na⁺-K⁺-ATPase α 1 β 1, α 2 β 1 and α 3 β 1 When Expressed Separately in Yeast Cells", *American Journal of Physiology—Cell Physiology*, 281: C1355-C1364, Oct. 2001.

Roeder et al. "Radioiodination of Hydroxyphenyl-Ethylamine Derivatives of Some Digitalisglycosides and Their Aglycones [Zur Radiojodmarkierung von Tyraminderivativen Einiger Digitalisglykoside und Deren Aglyka]", *Journal of Labelled Compounds and Radiopharmaceuticals*, XP055221078, 15: 197-214, Jan. 1978. Compounds 19, 20, 25, 26, Preparation.

Serrano-Wu et al. "Sordarin Oxazepine Derivatives as Potent Antifungal Agents", *Bioorganic and Medicinal Chemistry Letters*, 12: 2757-2760, 2002.

Specht et al. "Two Different Na,K-ATPases in the Optic Nerve: Cells of Origin and Axonal Transport", *Proc. Natl. Acad. Sci. USA*, 81: 1234-1238, Feb. 1984.

Strugatsky et al. "Expression of Na⁺,K⁺-ATPase in *Pichia Pastoris*. Analysis of Wild Type and D369N Mutant Proteins by Fe²⁺-Catalyzed Oxidative Cleavage", *The Journal of Biological Chemistry*, 278(46): 46064-46073, Nov. 14, 2003.

Sweadner "Isozymes of the Na⁺/K⁺-ATPase", *Biochimica et Biophysica Acta*, 988: 185-220, 1989.

Sweadner "Two Molecular Forms of (Na⁺ + K⁺)-Stimulated ATPase in Brain. Separation, and Difference in Affinity for Strophanthidin", *The Journal of Biological Chemistry*, 254(13): 6060-6067, Jul. 10, 1979.





Tian et al. "Signal-Transducing Function of Na⁺-K⁺-ATPase is Essential for Ouabain's Effect on [Ca²⁺]_i in Rat Cardiac Myocytes", *American Journal of Physiology—Heart and Circulation Physiology*, 281: H1899-H1907, 2001.





Xie et al. "Na⁺/K⁺-ATPase as A Signal Transducer", *European Journal of Biochemistry*, 269: 2434-2439, Feb. 2002.

Yeagle et al. "Effects of Cholesterol on (Na⁺,K⁺)-ATPase ATP Hydrolyzing Activity in Bovine Kidney", *Biochemistry*, 27: 6449-6452, 1988.

Office Action dated Jun. 21, 2018 From the Israel Patent Office Re. Application No. 244359 and Its Translation Into English. (6 Pages).

* cited by examiner











FIG. 4



**SELECTIVE INHIBITORS OF
ALPHA2-CONTAINING ISOFORMS OF
NA,K-ATPASE AND USE THEREOF FOR
REDUCTION OF INTRAOCULAR PRESSURE**

RELATED APPLICATIONS

This application is a National Phase of PCT Patent Application No. PCT/IL2016/050785 having International filing date of Jul. 19, 2016, which claims the benefit of priority under 35 USC § 119(e) of U.S. Provisional Patent Application No. 62/302,226, filed on Mar. 2, 2016. PCT Patent Application No. PCT/IL2016/050785 is also a Continuation-in-Part (CIP) of PCT Patent Application No. PCT/IL2015/050741, having international filing date of Jul. 19, 2015. The contents of the above applications are all incorporated by reference as if fully set forth herein in their entirety.

SEQUENCE LISTING STATEMENT

The ASCII file, entitled 72096SequenceListing.txt, created on Jan. 17, 2018, comprising 73,501 bytes, submitted concurrently with the filing of this application is incorporated herein by reference. The sequence listing submitted herewith is identical to the sequence listing forming part of the international application.

FIELD AND BACKGROUND OF THE
INVENTION

The present invention, in some embodiments thereof, relates to pharmaceutical agents and, more particularly, but not exclusively, to digoxin and digitoxin derivatives exhibiting selective inhibition of $\alpha 2$ -containing isoforms of Na,K-ATPase, and uses thereof to reduce intraocular pressure (IOP), and/or as cardiotoxic agents in a subject in need thereof.

Glaucoma is a disease leading to irreversible blindness. Control of intraocular pressure (IOP) is the mainstay of current therapy of glaucoma, and is achieved by various drugs, such as β -blockers, prostaglandin analogues, $\alpha 2$ adrenergic receptor agonists, cholinergic agonists and carbonic anhydrase inhibitors given topically or systemically. The topical route is preferable, provided the drug effectively permeates the cornea, because this minimizes systemic side-effects. Despite the selection of drugs available, uncontrolled IOP in many patients eventually makes surgical intervention necessary. Thus, fresh approaches to drug treatment of glaucoma are highly desirable.

The Na,K-ATPase is the motor for production of the aqueous humour (bodily fluid) in the ciliary body epithelium and, in principle, inhibition of the Na,K-ATPase can suppress the production of aqueous humour, and control IOP. Control of IOP is the mainstay of glaucoma therapy; however, the available drugs suffer from a variety of shortcomings, particularly due to systemic adverse effects and low therapeutic index. Previously, intra-venous digoxin, a classical inhibitor of the Na,K-pump, typically used primarily to treat congestive heart failure, was considered for this role but was discarded due to systemic toxicity.

Isoforms of the Na,K-ATPase ion pump consists of α and β subunits (α/β) and accessory FXYD regulatory subunits. There are four isoforms of the $\alpha 1$ subunit ($\alpha 1-4$) and three isoforms of the β subunit ($\beta 1-3$) expressed in a tissue-specific fashion. The $\alpha 1$ isoform is the common isoform that maintains Na and gradients in all tissues, $\alpha 2$ is expressed

mainly in muscle and astrocytes, and $\alpha 3$ is expressed mainly in nerve cells. For example, human heart expresses $\alpha 1$ (about 70%) and both $\alpha 2$ and $\alpha 3$ isoforms (about 30%) and $\beta 1$.

The ciliary epithelium in the eye is a functional syncytium consisting of apical pigmented cells (PE) oriented towards the blood and baso-lateral non-pigmented (NPE) cells oriented towards the anterior chamber of the eye. It is known that the primary Na,K-ATPase isoform of the PE is $\alpha 1\beta 1$ while that of the NPE is $\alpha \beta 3$. The Na,K-ATPase in the NPE powers the production of the aqueous humor and controls intraocular pressure.

Thus, in principle, topically applied $\alpha 2$ -selective cardiac glycosides that penetrate the intact eye and reach the ciliary epithelium could effectively reduce IOP. A potential advantage of topical application could be that systemic toxic effects typical of cardiac glycosides should be minimal.

Another possible application of an $\alpha 2$ -selective cardiac glycoside could be as an effective cardiotoxic drug, with reduced cardiotoxicity, compared to known drugs such as digoxin. Digitalis drugs such as digoxin have been used to treat heart failure for over two hundred years but are dangerous drugs with multiple side effects. There is now good evidence that selective inhibition of $\alpha 2$ is especially effective in enhancing cardiac excitation-contraction coupling and mediating cardiac glycoside-mediated positive inotropy. Inhibition of $\alpha 2$, which is a minor isoform and is located largely in T-tubules, may mediate the positive cardiotoxic effects, but $\alpha 2$ -selective cardiac glycosides should only minimally inhibit $\alpha 1$, located primarily in the outer sarcolemma membrane, and thus avoid cellular Ca overload, the hallmark of cardiac toxicity.

The isoform selectivity of a large number of known cardiac glycosides has been previously studied, using the yeast *P. pastoris* expressing Na,K-ATPase isoforms ($\alpha 1\beta 1$, $\alpha 2\beta 1$, $\alpha 3\beta 1$), and purified detergent-soluble isoform complexes of Na,K-ATPase [Cohen E. et al., 2005, *J Biol Chem*, 280(17), pp. 16610-16618; Flaviv H, et al., 2007, *Biochemistry*, 46(44), pp. 12855-12867; Lifshitz Y, et al., 2007, *Biochemistry*, 46(51), pp. 14937-14950; Mishra N K, et al., 2011, *J Biol Chem*, 286(11), pp. 9699-9712; and Kapri-Pardes E, et al., 2011, *J Biol Chem*, 286(50), pp. 42888-42899].

Dissociation constants, K_D , for digitalis glycosides, digoxin and digitoxin, measured by ^3H -ouabain displacement assays in membranes, showed moderate selectivity (3-4-fold) for $\alpha 2/\alpha 3$ over $\alpha 1$. By contrast to the digitalis glycosides, the K_D of ouabain showed some preference for $\alpha 1$ over $\alpha 2$ and similar K_i values for all three isoforms. In assays of inhibition of Na,K-ATPase activity, measured with the purified isoform protein complexes, digoxin and digitoxin showed 3-4-fold lower K_i (inhibition) values for $\alpha 2$ compared to $\alpha 1$, with $\alpha 3$ more similar to $\alpha 1$. No aglycones of any cardiac glycosides tested showed isoform selectivity. For digoxin derivatives, with one to four digitoxose moieties the maximal $\alpha 2/\alpha 1$ selectivity was found for digoxin itself, with three digitoxose sugars [Katz, A. et al., *J Biol Chem*, 2010, 285(25), pp. 19582-19592].

Based on recent studies [Laursen, M. et al., *Prot Natl Acad Sci USA*, 2015, 112(6):1755-60], it was inferred that the sugar moiety of digoxin likely determines isoform selectivity, which is generally consistent with recent structures of Na,K-ATPase with bound ouabain, bufalin or digoxin. The unsaturated lactone ring and steroid portion of ouabain are bound between trans-membrane segments M1, M4, M5 of the α subunit, in which there are no amino-acid differences between isoforms. Assuming that the aglycones of all cardiac glycosides bind similarly, the implication is


3

that isoforms cannot discriminate between any of the aglycones, as found experimentally. By contrast, the sugar is bound near extracellular loops, where there are a number of amino-acid differences between the isoforms. These residues might interact with the sugars of bound digoxin in an isoform-selective way.

Additional background art include WO 2015/029035, WO 2007/079128, WO 2010/053771, U.S. Patent Application No. 2005/0032138 and U.S. Pat. Nos. 7,888,059 and 7,851,145; these documents are hereby incorporated by reference.

SUMMARY OF THE INVENTION

According to an aspect of some embodiments of the present invention there is provided a compound represented by general Formula I:




Formula I

including any pharmaceutically acceptable salt, prodrug, hydrate, solvate, enantiomer and diastereomer thereof, and any mixtures thereof,

wherein:

X is H or OH;

R is represented by general Formula II;



Formula II

A is a spacer moiety or a covalent bond; and

B is a cyclic moiety;

or

B is selected from the group consisting of an alkylsulfonyl, an arylsulfonyl and a sulfonamide;

or

B is $-NR_1R_2$, wherein R_1 and R_2 are each independently H or a C_1 - C_4 alkyl provided that at least one of R_1 and R_2 is a C_1 - C_4 alkyl.

According to some embodiments of the invention, A is selected from the group consisting of a covalent bond, an unsubstituted C_1 - C_6 alkyl, a substituted C_1 - C_6 alkyl, an unsubstituted C_1 - C_6 alkyl interrupted by one or more heteroatom and a substituted C_1 - C_6 alkyl interrupted by one or more heteroatom.

According to some embodiments of the invention, B is a cyclic moiety selected from the group consisting of an

4

unsubstituted alicyclic moiety, a substituted alicyclic moiety, an unsubstituted heterocyclic moiety, a substituted heterocyclic moiety, an unsubstituted aryl moiety, a substituted aryl moiety, an unsubstituted heteroaryl moiety and a substituted heteroaryl moiety.

According to some embodiments of the invention, B is an unsubstituted alicyclic moiety.

According to some embodiments of the invention, the unsubstituted alicyclic moiety is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

According to some embodiments of the invention, B is a substituted alicyclic moiety.

According to some embodiments of the invention, the substituted alicyclic moiety is selected from the group

consisting of 2,3-dimethylcyclopropane-1-yl, 3,3-dimethylcyclobutane-1-yl, 3,4-dimethylcyclopentane-1-yl and 3,5-dimethylcyclohexane-1-yl.

According to some embodiments of the invention, B is an unsubstituted heterocyclic moiety.

According to some embodiments of the invention, the unsubstituted heterocyclic moiety is selected from the group consisting of oxiranyl, aziridinyl, oxetanyl, azetidiny, thietanyl, tetrahydrofuranyl, pynrolidinyl, tetrahydropyranyl and piperidinyl.

According to some embodiments of the invention, B is an unsubstituted aryl moiety.

According to some embodiments of the invention, the unsubstituted aryl moiety is selected from the group consisting of cyclopentadienyl, phenyl and naphthyl.

According to some embodiments of the invention, B is an unsubstituted heteroaryl moiety.

According to some embodiments of the invention, the unsubstituted heteroaryl moiety is imidazolyl.

According to some embodiments of the invention, the alkylsulfonyl is selected from the group consisting of methylsulfonyl, ethylsulfonyl and isopropylsulfonyl.

According to some embodiments of the invention, the arylsulfonyl is selected from the group consisting of phenylsulfonyl, benzylsulfonyl and tosyl.

According to some embodiments of the invention, the sulfonamide is selected from the group consisting of methylsulfonamide, N-methylmethanesulfonamide and N,N-dimethylmethanesulfonamide.

5

According to some embodiments of the invention, B is —N(Et)₂.

According to some embodiments of the invention, X is H.

According to some embodiments of the invention, A is a covalent bond and B is cyclobutyl.

According to some embodiments of the invention, A is —CH₂— and B is cyclopropyl.

According to some embodiments of the invention, X is OH.

According to some embodiments of the invention, A is a covalent bond and B is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl.

According to some embodiments of the invention, A is —CH₂— and B is selected from the group consisting of cyclopropyl, 3,3-dimethylcyclobutane-1-yl and phenyl.

According to some embodiments of the invention, A is —(CH₂)₂— and B is cyclopropyl.

According to some embodiments of the invention, R is selected from the group consisting of cyclopropyl, methylcyclopropane, ethylcyclopropane, propylcyclopropane, cyclobutyl, methylcyclobutane, methyl-3,3-dimethylcyclobutane, ethylcyclobutane, propylcyclobutane, cyclopentyl, methylcyclopentane, ethylcyclopentane, propylcyclopentane, cyclohexyl, azetidiny, oxetanyl, thietanyl, histaminy and benzyl.

According to some embodiments of the invention, R is selected from the group consisting of cyclopropyl, methylcyclopropane and cyclobutyl.

According to some of any of the embodiments of the invention, the compound is having an affinity to at least one isoform of Na,K-ATPase.

According to some embodiments of the invention, the isoform is selected from the group consisting of $\alpha 1\beta 1$, $\alpha 1\beta 2$, $\alpha 1\beta 3$, $\alpha 2\beta 1$, $\alpha 2\beta 2$, $\alpha 2\beta 3$, $\alpha 3\beta 1$, $\alpha 3\beta 2$, $\alpha 3\beta 3$, $\alpha 4\beta 1$, $\alpha 4\beta 2$ and $\alpha 4\beta 3$.

According to some embodiments of the invention, the affinity of the compound to any one of $\alpha 2\beta 1$, $\alpha 2\beta 2$ and $\alpha 2\beta 3$ is higher than the affinity to $\alpha 1\beta 1$, $\alpha 1\beta 2$, $\alpha 1\beta 3$, $\alpha 3\beta 1$, $\alpha 3\beta 2$, $\alpha 3\beta 3$, $\alpha 4\beta 1$, $\alpha 4\beta 2$ and $\alpha 4\beta 3$ by at least 100%.

According to some embodiments of the invention, the affinity of the compound to any one of $\alpha 2\beta 1$, $\alpha 2\beta 2$ and $\alpha 2\beta 3$ is higher than the affinity to $\alpha 1\beta 1$ by at least 300% (4-fold).

According to some embodiments of the invention, the affinity of the compound to $\alpha 2\beta 3$ is higher than the affinity to $\alpha 1\beta 1$ by at least 500 (6-fold).

According to some embodiments of the invention, the compound is having a $\alpha 2\beta 3$ inhibition constant (K_i) lower than 10 nM.

6

According to an aspect of some embodiments of the present invention there is provided a pharmaceutical composition that includes as an active ingredient a compound according to any of the embodiments of the invention and a pharmaceutically acceptable carrier.

According to some embodiments of the invention, the pharmaceutical composition is packaged in a packaging material and identified in print, or on the packaging material, for use in reducing intraocular pressure (IOP).

According to some embodiments of the invention, the pharmaceutical composition is packaged in a packaging material and identified in print, or on the packaging material, for use in a treatment of a heart condition.

According to an aspect of some embodiments of the present invention there is provided a method of reducing intraocular pressure (IOP) in a subject in need thereof, which includes administering to the subject a therapeutically effective amount of a compound according to any of the embodiments of the invention, or a pharmaceutical composition according to some of the embodiments of the invention.

According to an aspect of some embodiments of the present invention there is provided a method of treating a heart condition in a subject in need thereof, that includes administering to the subject a therapeutically effective amount of a compound according to any of the embodiments of the invention, or a pharmaceutical composition according to some of the embodiments of the invention.


According to an aspect of some embodiments of the present invention there is provided a use of a compound according to any of the embodiments of the invention, or a pharmaceutical composition according to some of the embodiments of the invention, for the manufacture of a medicament for reducing intraocular pressure (IOP).

According to an aspect of some embodiments of the present invention there is provided a use of a compound according to any of the embodiments of the invention, or a pharmaceutical composition according to some of the embodiments of the invention, for the manufacture of a medicament for treating a heart condition.

According to some embodiments of the invention, the heart condition is selected from the group consisting of atrial fibrillation, atrial flutter, mitral stenosis, chronic heart failure and congestive heart failure.

According to an aspect of some embodiments of the present invention there is provided a pharmaceutical composition that includes as active ingredients:

at least one ingredient selected from the group consisting of a prostaglandin analog, a β -blocker, an adrenergic agent, an $\alpha 2$ -adrenergic receptor agonist, a miotic agent, a carbonic anhydrase inhibitor and a cholinergic agonist; and a compound represented by Formula III:



7

including any pharmaceutically acceptable salt, prodrug, hydrate, solvate, enantiomer and diastereomer thereof, and any mixtures thereof, and a pharmaceutically acceptable carrier,

wherein:

X is H or OH;

R' is selected from the group consisting of OH, C₁-C₆ alkyl, C₁-C₆ haloalkyl (C₁-C₆ alkyl substituted with at least one halo), $-(CR^bR^c)_nSi(R^a)_3$, $-(CR^bR^c)_n-C(=Y)-NR_1R_2$, $-(CR^bR^c)_n-C(=Y)-NHOH$, $-(CR^dR^e)_n-C(=Y)-COOR_3$, $-NHC(=Y)NR_1R_2$ and $-(CR^bR^c)_n-NH_2$;

Y is O or S;

R₁, R₂ and R₃ are each independently H or a C₁-C₄ alkyl;


R_a is a C₁-C₄ alkyl;

R_b, R_c and R_d are each independently selected from H, a C₁-C₄ alkyl and a C₁-C₄ hydroxyalkyl;

R_e is selected from a C₁-C₄ alkyl and a C₁-C₄ hydroxyalkyl; and

n is 0, 1 or 2;

or R' is represented by general Formula II:



Formula II

25

A is a spacer moiety or a covalent bond; and


B is a cyclic moiety, or B is selected from the group consisting of an alkylsulfonyl, an arylsulfonyl and a sulfonamide, or B is $-NR_1R_2$, wherein R₁ and R₂ are each independently H or a C₁-C₄ alkyl provided that at least one of R₁ and R₂ is a C₁-C₄ alkyl,

and a pharmaceutically acceptable carrier.

According to some embodiments of the invention, the pharmaceutical composition is packaged in a packaging material and identified in print, or on the packaging material, for use in reducing intraocular pressure (IOP).

According to an aspect of some embodiments of the present invention there is provided a use of an agent selected from the group consisting of a prostaglandin analog, a β-blocker, an adrenergic agent, an α₂-adrenergic receptor agonist, a miotic agent, a carbonic anhydrase inhibitor and a cholinergic agonist, and

a compound represented by Formula III.



Formula III

8

including any pharmaceutically acceptable salt, prodrug, hydrate, solvate, enantiomer and diastereomer thereof, and any mixtures thereof,

wherein:

5 X is H or OH;

R' is selected from the group consisting of OH, C₁-C₆ alkyl, C₁-C₆ haloalkyl (C₁-C₆ alkyl substituted with at least one halo), $-(CR^bR^c)_nSi(R^a)_3$, $-(CR^bR^c)_n-C(=Y)-NR_1R_2$, $-(CR^bR^c)_n-C(=Y)-NHOH$, $-(CR^dR^e)_n-C(=Y)-COOR_3$, $-NHC(=Y)NR_1R_2$ and $-(CR^bR^c)_n-NH_2$;

Y is O or S;

R₁, R₂ and R₃ are each independently H or a C₁-C₄ alkyl;


R_a is a C₁-C₄ alkyl;

15 R_b, R_c and R_d are each independently selected from H, a C₁-C₄ alkyl and a C₁-C₄ hydroxyalkyl;

R_e is selected from a C₁-C₄ alkyl and a C₁-C₄ hydroxyalkyl; and

20 n is 0, 1 or 2;

or R' is represented by general Formula II:



Formula II

25

30 A is a spacer moiety or a covalent, bond; and

B is a cyclic moiety, or B is selected from the group consisting of an alkylsulfonyl, an arylsulfonyl and a sulfonamide, or B is $-NR_1R_2$, wherein R₁ and R₂ are each independent r a C₁-C₄ alkyl provided that at least one of R₁ and R₂ is a C₁-C₄ alkyl,


35

for the manufacture of a medicament for reducing intraocular pressure (IOP).

40 According to an aspect of some embodiments of the present invention there is provided a method of reducing intraocular pressure (IOP) in a subject in need thereof, which includes co-administering to the subject a therapeutically effective amount of:

45 an agent selected from the group consisting of a prostaglandin analog, a (β-blocker, an adrenergic agent, an α₂-adrenergic receptor agonist, a miotic agent, a carbonic anhydrase inhibitor and a cholinergic agonist; and

a compound represented by Formula III:



Formula III

including any pharmaceutically acceptable salt, prodrug, hydrate, solvate, enantiomer and diastereomer thereof, and any mixtures thereof,

wherein:

X is H or OH;

R' is selected from the group consisting of OH, C₁-C₆ alkyl, C₁-C₆ haloalkyl (C₁-C₆ alkyl substituted with at least one halo), —(CR^bR^c)_nSi(R^a)₃, —(CR^bR^c)_n-C(=Y)—NR₁R₂, —(CR^bR^c)_n-C(=Y)—NHOH, —(CR^dR^e)_n-C(=Y)—COOR₃, —NHC(=Y)NR₁R₂ and —(CR^bR^c)_n-NH₂;

Y is O or S;

R₁, R₂ and R₃ are each independently H or a C₁-C₄ alkyl;


R_a is a C₁-C₄ alkyl;

R_b, R_c and R_d are each independently selected from H, a C₁-C₄ alkyl and a C₁-C₄ hydroxyalkyl;

Re is selected from a C₁-C₄ alkyl and a C₁-C₄ hydroxyalkyl; and

n is 0, 1 or 2;

or R' is represented by general Formula II:



Formula II

A is a spacer moiety or a covalent bond; and

B is a cyclic moiety, or B is selected from the group consisting of an alkylsulfonyl, an arylsulfonyl and a sulfo-

namide, or B is —NR₁R₂, wherein R₁ and R₂ are each independently H or a C₁-C₄ alkyl provided that at least one of R₁ and R₂ is a C₁-C₄ alkyl.

According to some embodiments of the invention, the mode of administration is effected topically, extraocularly, intraocularly and/or intravitreally.

According to some embodiments of the invention, the composition according to some embodiments of the invention is formulated as an ophthalmic composition suitable for topical, extraocular, intraocular and/or intravitreal administration to the eye of the subject.


According to some embodiments of the invention, the composition according to some embodiments of the invention is in the form selected from the group consisting of an eye-drop solution, a spray, an eye wash solution, an ointment, a suspension, a gel, a cream and an injectable solution.

According to some embodiments of the invention, the intraocular pressure (IOP) is associated with glaucoma, low-tension glaucoma and normal-tension glaucoma.

According to an aspect of some embodiments of the present invention there is provided a method of treating a heart condition in a subject in need thereof, that includes co-administering to the subject a therapeutically effective amount of:

an agent selected from the group consisting of a β-blocker, an anticoagulation agent, an angiotensin-converting-enzyme inhibitor and an angiotensin II receptor antagonist; and

a compound represented by Formula III:



Formula III

including any pharmaceutically acceptable salt, prodrug, hydrate, solvate, enantiomer and diastereomer thereof, and any mixtures thereof,

wherein:

X is H or OH;

R' is selected from the group consisting of OH, C₁-C₆ alkyl, C₁-C₆ haloalkyl (C₁-C₆ alkyl substituted with at least one halo), $-(CR^bR^c)_nSi(R^a)_3$, $-(CR^bR^c)_n-C(=Y)-NR_1R_2$, $-(CR^bR^c)_n-C(=Y)-NHOH$, $-(CR^aR^e)_n-C(=Y)-COOR_3$, $-NHC(=Y)NR_1R_2$ and $-(CR^bR^c)_n-NH_2$;

Y is O or S;

R₁, R₂ and R₃ are each independently H or a C₁-C₄ alkyl;


R_a is a C₁-C₄ alkyl;

R_b, R_c and R_d are each independently selected from H, a C₁-C₄ alkyl and a C₁-C₄ hydroxyalkyl;

R_e is selected from a C₁-C₄ alkyl and a C₁-C₄ hydroxyalkyl; and

n is 0, 1 or 2;

or R' is represented by general Formula II:



Formula II

A is a spacer moiety or a covalent bond; and

B is a cyclic moiety, or B is selected from the group consisting of an alkylsulfonyl, an arylsulfonyl and a sulfonamide, or B is $-NR_1R_2$, wherein R₁ and R₂ are each independently H or a C₁-C₄ alkyl provided that at least one of R₁ and R₂ is a C₁-C₄ alkyl.

According to an aspect of some embodiments of the present invention there is provided a process of preparing a compound according any of the embodiments of the invention, the process includes:

converting the third digitoxose moiety of digoxin or digitoxin into a dialdehyde; and

reacting the dialdehyde with a reagent represented by general formula IV:



Formula IV

A is a spacer moiety or a covalent bond; and

B is a cyclic moiety, or B is selected from the group consisting of an alkylsulfonyl, an arylsulfonyl and a sulfonamide, or B is $-NR_1R_2$, wherein R₁ and R₂ are each independently H or a C₁-C₄ alkyl provided that at least one of R₁ and R₂ is a C₁-C₄ alkyl.

According to some embodiments of the invention, converting the third digitoxose moiety of digoxin or digitoxin into a dialdehyde is effected by sodium periodate (NaIO₄).

According to some embodiments of the invention, reacting the dialdehyde with a reagent represented by general formula IV is effected in the presence of NaCNBH₃.

According to an aspect of some embodiments of the present invention there is provided a method of determining an affinity of a compound, according to any embodiment of the invention, to at least one isoform of Na,K-ATPase, the method includes contacting the isoform of Na,K-ATPase with the compound in an affinity measurement setup and determining the affinity.

According to an aspect of some embodiments of the present invention there is provided a method of isolating an isoform of Na,K-ATPase of a mammal, that includes:

transforming yeast cells with a clone that includes an a chain sequence and a β chain sequence of the Na,K-ATPase;

expressing the clone in the yeast cells; and

isolating the isoform,

wherein:

the α chain sequence is selected from a group consisting of α1, α2, α3 and α4; and

the β chain sequence is selected from the group consisting of β1, β2 and β3.

According to some embodiments of the invention, the isoform is selected from the group consisting of α1β1, α1β2, α1β3, α2β1, α2β2, α2β3, α3β1, α3β2, α3β3, α4β1, α4β2 and α4β3.

According to some embodiments of the invention, the isoform is α2β2.

According to some embodiments of the invention, the isoform is α2β3.

According to an aspect of some embodiments of the present invention there is provided an isolated isoform of Na,K-ATPase of a mammal having at least 70% purity, wherein the isoform is α2β2.

According to an aspect of some embodiments of the present invention there is provided an isolated isoform of Na,K-ATPase of a mammal having at least 70% purity, wherein the isoform is α2β3.


13

According to an aspect of some embodiments of the present invention there is provided an isolated isoform of Na,K-ATPase of a mammal having a yeast-characterizing glycosylation pattern, wherein the isoform is $\alpha 2\beta 2$.

According to an aspect of some embodiments of the present invention there is provided an isolated isoform of Na,K-ATPase of a mammal having a yeast-characterizing glycosylation pattern, wherein the isoform is $\alpha 2\beta 3$.

According to some embodiments of the invention, the isolated isoform, according to any embodiment of the invention, is human.

According to some embodiments of the invention a prodrug of the compound is represented by Formula V:



Formula V

wherein X is H or PD₃, and

each of PD₁-PD₄ is H or independently selected from the group consisting of a methoxymethyl ether, a tetrahydropyranyl ether, a t-butyl ether, an allyl ether, a benzyl ether, a t-butyldimethylsilyl ether, a t-butyldiphenylsilyl ether, an acetic acid ester (Ac), ethyl, propyl, butyl, t-butyl or pivalic acid ester and a benzoic acid ester,

provided that at least one of PD₁-PD₄ is not H.

According to some embodiments, each of PD₁-PD₄ is an acetic acid ester (Ac).

According to some embodiments, each of PD₁ and PD₂ is an acetic acid ester (Ac).

According to some embodiments, each of PD₁-PD₃ is an acetic acid ester (Ac).

Unless otherwise defined, all technical and/or scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of embodiments of the invention, exemplary methods and/or materials are described below. In case of conflict, the patent specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and are not intended to be necessarily limiting.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

Some embodiments of the invention are herein described, by way of example only, with reference to the accompanying drawings. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of

14

embodiments of the invention. In this regard, the description taken with the drawings makes apparent to those skilled in the art how embodiments of the invention may be practiced.

In the drawings:

FIGS. 1A, 1B, 1C and 1D present comparative plots of IOP as a function of time, showing the dose response of $\alpha 2$ -inhibitor compounds, according to some embodiments of the present invention, in lowering IOP in live rabbits, wherein FIG. 1A shows the results obtained for DiB, FIG. 1B shows the duration of the effect of DiB while 4AP is added every 2 hours so as to maintain the raised IOP, FIG. 1C shows the results obtained for DMcP, and FIG. 1D shows the results obtained for DcB;

FIGS. 2A, 2B, 2C and 2D present comparative plots of IOP as a function of time, demonstrating the capacity of the $\alpha 2$ -inhibitor compounds, according to some embodiments of the present invention, to lower IOP below basal levels compared to a buffer control when administered topically to one eye of a rabbit, while the other eye received PBS as a control, wherein FIG. 2A shows the lack of effect of digoxin, FIG. 2B shows the lack of effect of DiB (a non-cyclic moiety inhibitor), FIG. 2C shows the notable effect of DMcP, and FIG. 2D shows the notable effect of DcB;

FIGS. 3A, 3B and 3C present comparative plots of IOP as a function of time, demonstrating the effect of $\alpha 2$ -inhibitor compounds, according to some embodiments of the present invention, to potentiate the drug Latanoprost in lowering IOP below basal levels, wherein FIG. 3A shows the effect of DcB alone, FIG. 3B shows the effect of Latanoprost alone, and FIG. 3C shows the effect of co-administering DcB with Latanoprost; and

FIG. 4 presents comparative plots of IOP as a function of time, demonstrating the capacity of the trisAcDcB prodrug of the $\alpha 2$ -inhibitor compound DcB, according to some embodiments of the present invention, to lower IOP below basal levels compared to a buffer control when administered topically to one eye of a rabbit, while the other eye received PBS as a control.

DESCRIPTION OF SPECIFIC EMBODIMENTS OF THE INVENTION

The present invention, in some embodiments thereof, relates to a pharmaceutical agents and, more particularly, but not exclusively, to digoxin and digitoxin derivatives exhibiting selective inhibition of $\alpha 2$ -containing isoforms of Na,K-

15

ATPase, and uses thereof to reduce intraocular pressure (IOP), and/or as cardiotoxic agents in a subject in need thereof.

The principles and operation of the present invention may be better understood with reference to the figures and accompanying descriptions.


Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not limited in its application to the details set forth in the following description or exemplified by the Examples. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of to description and should not be regarded as limiting.

As discussed hereinabove, Na,K-ATPase inhibitors that exhibit selectivity towards tissue-specific isoforms of the protein, present pharmaceutical advantages such as broader therapeutic window and wider scope of modes of administration. For example, Na,K-ATPase inhibitors exhibiting selectivity towards protein isoforms containing the $\alpha 2$ subunit offer this advantages over unselective inhibitors in treating medical conditions wherein lowering the intraocular pressure is called for.

While searching for $\alpha 2$ -selective inhibitors, the present inventors have surprisingly found that certain derivatives of digoxin and digitoxin, wherein the perhydro-1-4-oxazepine moiety thereof is N-substituted with a cyclic moiety, exhibit a notable selectivity towards $\alpha 2$ -containing isoform of Na,K-ATPase.

Compounds:


According to an aspect of some embodiments of the present invention there is provided a compound represented by general Formula I:



wherein:

X is H or OH, whereas derivatives having X=H are referred to as digitoxin derivatives, and derivatives having X=OH are referred to as digoxin derivatives;

R is represented by general Formula II:




Formula II

16

the wiggled line represents the N-link to the compound; A is a spacer moiety or a covalent bond; and B is a cyclic moiety;

or

R is represented by general Formula II:




Formula II

the wiggled line represents the N-link to the compound; A is a spacer moiety or a covalent bond; and B is selected from the group consisting of an alkylsulfonyl, an arylsulfonyl and a sulfonamide;

or

R is represented by general Formula II:



Formula II

the wiggled line represents the N-link to the compound; A is a spacer moiety or a covalent bond; and B is $-\text{NR}_1\text{R}_2$, wherein R_1 and R_2 are each independently H or a C_1 - C_4 alkyl provided that at least one of R_1 and R_2 is a C_1 - C_4 alkyl, namely B is a secondary amine or tertiary amine.

As used herein, the term "cyclic moiety" refers to a group of atoms that are covalently attached to one another so as to

Formula I

form at least one ring of atoms. Non-limiting examples of cyclic moieties include unsubstituted alicyclic moieties, substituted alicyclic moieties, unsubstituted heterocyclic moieties, substituted heterocyclic moieties, unsubstituted aryl moieties, substituted aryl moieties, unsubstituted heteroaryl moiety and substituted heteroaryl moieties.

A substituted cyclic moiety has one or more chemical group or atom attached to one of the atoms in the ring of atoms. Examples of such chemical groups or atoms include, without limitation, C_1 - C_6 alkyl, hydroxyl, amine, halo, alkoxy, carboxyl, amide and the like, or a second cyclic moiety attached by a covalent bond(s) to one or two of the ring atoms of the cyclic moiety.

17

The terms “hydroxyl” or “hydroxy”, as used herein, refer to an —OH group.

As used herein, the term “amine” describes a —NR¹R² group where each of R¹ and R² is independently hydrogen, alkyl, cycloalkyl, heteroalicyclic, aryl or heteroaryl, as these terms are defined herein.

As used herein, the term “alkyl” describes an aliphatic hydrocarbon including straight chain and branched chain groups. The alkyl may have 1 to 20 carbon atoms, or 1-10 carbon atoms, and may be branched or unbranched. According to some embodiments of the present invention, the alkyl is a low (or lower) alkyl, having 1-4 carbon atoms (namely, methyl, ethyl, propyl and butyl).

Whenever a numerical range; e.g., “1-10”, is stated herein, it implies that the group, in this case the alkyl group, may contain 1 carbon atom, 2 carbon atoms, 3 carbon atoms, etc., up to and including 10 carbon atoms. In some embodiments, the alkyl is a lower alkyl, including 1-6 or 1-4 carbon atoms.

A C₁-C₆ alkyl group refers to any one of the moieties methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, i-butyl, t-butyl, n-pentyl, t-pentyl, neopentyl, i-pentyl, s-pentyl, 3-pentyl, n-hexyl, 2-methylpentyl, 3-methylpentyl, 2,2-dimethylbutyl and 2,3-dimethylbutyl.

The term “halide”, as used herein, refers to the anion of a halo atom, i.e. F⁻, Cl⁻, Br⁻ and I⁻.

The term “halo” refers to F, Cl, Br and I atoms as substituents.

The term “alkoxy” refers to an —OR¹ group, wherein R¹ is as defined herein, but other than hydrogen.

The term “amide” as used herein encompasses C-amide and N-amide.

The term “C-amide” describes a —C(=O)—NR¹R² group, where R¹ and R² are as defined herein.

The term “N-amide” describes a R¹C(=O)—NR²— group, where R¹ and R² are as defined herein.

The term alkylsulfonyl and arylsulfonyl refers to an R¹—S(=O)₂— group, wherein R¹ is as defined herein, but other than hydrogen. Examples of arylsulfonyl groups include p-toluenesulfonyl (tosyl; Ts), p-bromobenzenesulfonyl (brosyl; Bs), 2- or 4-nitrobenzenesulfonyl (nosyl; Ns), methanesulfonyl (mesyl; Ms), trifluoromethanesulfonyl (triflyl; Tf), and 5-(dimethylamino)naphthalene-1-sulfonyl (Dansyl; Ds).

The term solfonamide refers to an R³—S(=O)₂— group, wherein R³ is amine as defined herein.

The terms “Acyclic” and “cycloalkyl”, refer to an all-carbon monocyclic or fused ring (i.e., rings which share an adjacent pair of carbon atoms), branched or unbranched group containing 3 or more carbon atoms where one or more of the rings does not have a completely conjugated pi-electron system, and may further be substituted or unsubstituted. The cycloalkyl can be substituted or unsubstituted.

Examples of alicyclic moieties include, without limitation, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclododecyl.

Examples of substituted alicyclic is selected from the group consisting of 2,3-dimethylcyclopropane-1-yl, 3,3-dimethylcyclobutane-1-yl, 3,4-dimethylcyclopentane-1-yl and 3,5-dimethylcyclohexane-1-yl.

The terms “heterocyclic” or “heteroalicyclic”, as used herein, describe a monocyclic or fused ring group having in the ring(s) one or more atoms such as nitrogen, oxygen and sulfur. The rings may also have one or more double bonds. However, the rings do not have a completely conjugated pi-electron system. The heteroalicyclic may be substituted

18

or unsubstituted. Representative examples are morpholine, piperidine, piperazine, tetrahydrofuran, tetrahydropyran and the like.


In some embodiments, the heterocyclic moiety is selected from the group consisting of oxiranyl, aziridinyl, oxetanyl, azetidinyl, tetrahydrofuranyl, pyrrolidinyl, tetrahydropyranyl and piperidinyl.

The term “aryl” describes an all-carbon monocyclic or fused-ring polycyclic (i.e., rings which share adjacent pairs of carbon atoms) groups having a completely conjugated pi-electron system, as in the example of phenyl. The aryl group may be unsubstituted or substituted by one or more substituents. Examples of aryls include cyclopentadienyl, phenyl and naphthyl.


The term “heteroaryl” describes a monocyclic or fused ring (i.e., rings which share an adjacent pair of atoms) group having in the ring(s) one or more atoms, such as, for example, nitrogen, oxygen and sulfur and, in addition, having a completely conjugated pi-electron system. Examples, without limitation, of heteroaryl moieties include pyrrole, furane, thiophene, imidazole, oxazole, thiazole, pyrazole, pyridine, pyrimidine, quinoline, isoquinoline, purine, thiazazole, indole and the like. The heteroaryl group may be unsubstituted or substituted by one or more substituents.

According to some embodiments, the spacer moiety A can be a covalent bond, an unsubstituted, C₁-C₆ alkyl, a substituted, C₁-C₆ alkyl, an unsubstituted C₁-C₆ alkyl interrupted by one or more heteroatom (e.g., O, N or S) and a substituted C₁-C₆ alkyl interrupted by one or more heteroatom. A spacer moiety may be substituted with one or more C₁-C₆ alkyl, hydroxyl, amine, halo, alkoxy, carboxyl, amide and the like.

In some embodiments, A is a covalent bond and B an alicyclic moiety such as, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the corresponding R in Formula I is:




In some embodiments, A is a covalent bond and B a heteroalicyclic moiety such as, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the corresponding R in Formula I is:





19

-continued




Thietanyl (Th)

In some embodiments, A is $-\text{CH}_2-$ and B a cyclic moiety such as, for example, cyclopropyl, 3,3-dimethylcyclobutane-1-yl and phenyl, and the corresponding R in Formula I is:

methylcyclopropyl
(McP)methyl-3,3-
dimethylcyclobutane
(MDMcB)


benzyl (Bz)

In some embodiments, A is $-(\text{CH}_2)_2-$ and B a cyclic moiety such as, for example, cyclopropyl, and the corresponding R in Formula I is:



Ethylcyclopropane (EcP)

In some embodiments, A is $-(\text{CH}_2)_2-$ and B a heteroaryl cyclic moiety such as, for example, imidazolyl, and the corresponding R in Formula I is:



Histaminyl (His)

20

In some embodiment, X is H, A is a covalent bond and B is cyclobutyl,

In some embodiment, X is H, A is $-\text{CH}_2-$ and B is cyclopropyl.


In some embodiment, X is OH, A is a covalent bond and B is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

In some embodiment, X is OH, A is $-\text{CH}_2-$ and B is cyclopropyl, 3,3-dimethylcyclobutane-1-yl and phenyl.


In some embodiment, X is OH, A is $-\text{CH}_2-$ and B is cyclopropyl.

In some embodiments, R of general Formula I is cyclopropyl, methylcyclopropane, ethylcyclopropane, propylcyclopropane, cyclobutyl, methylcyclobutane, methyl-3,3-dimethylcyclobutane, ethylcyclobutane, propylcyclobutane, cyclopentyl, methylcyclopentane, ethylcyclopentane, propylcyclopentane, cyclohexyl and benzyl. Alternatively, R is cyclopropyl, methylcyclopropane and cyclobutyl.

In some embodiments, A is $-\text{CH}_2-$ and B an alkylsulfonyl such as, for example, methylsulfonyl, and the corresponding R in Formula I is:


(methylsulfonyl)methyl
(MSM)

In some embodiments, A is $-(\text{CH}_2)_2-$ and B an alkylsulfonyl such as, for example, methylsulfonyl, or a solfonamide, and the corresponding R in Formula I is:




(methylsulfonyl)ethyl (ESM) (sulfonamide)ethyl (ESA)


In some embodiments, A is $-(\text{CH}_2)_2-$ and B a tertiary amine such as, for example, N,N-dimethylamine, and the corresponding R in Formula I is:

(N,N-dimethylamine)ethyl
(EDA)

In some embodiments, the compound is any one of the structures presented below:




DcP or 4-((3S,5R,8R,9S,10S,12R,13S,14S,17R)-3-(((2R, 4S,5S,6R)-5-(((2S,4S,5S,6R)-5((4-cyclopropyl-2-methyl-1, 4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-12,14-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)furan-2(5H)-one; 25



DMcP or 4-((3S,5R,8R,9S,10S,12R,13S,14S,17R)-3-(((2R,4S,5S,6R)-5-(((2S,4S,5S,6R)-5((4-cyclopropylmethyl)-2-methyl-1,4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-12,14-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)furan-2(5H)-one; 60


23

24



DEcP or 4((3S,5R,8R,9S,10S,12R,13S,14S,17R)-3-(((2S,4S,5R,6R)-5-(((2S,4S,5S,6R)-5-(2-cyclopropylethyl)-2-methyl-1,4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-12,14-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl) furan-2(5H)-one;

25




DcB or 4((3S,5R,8R,9S,10S,12R,13S,14S,17R)-3-(((2R,4S,5S,6R)-5-4(2S,4S,5S,6R)-5-((4-cyclobutyl-2-methyl-1,4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-12,14-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl) furan-2(5H)-one;

65


25

26



DcPe or 4-((3S,5R,8R,9S,10S,12R,13S,14S,17R)-3-(((2S,4S,5R,6R)-5-(((2S,4S,5S,6R)-5-((4-cyclopentyl-2-methyl-1,4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-12,14-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)furan-2(5H)-one;

25




DcHe or 4-((3S,5R,8R,9S,10S,12R,13S,14S,17R)-3-(((2S,4S,5R,6R)-5-(((2S,4S,5S,6R)-5-((4-cyclohexyl-2-methyl-1,4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-12,14-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)furan-2(5H)-one;

60


65

27

28




DBz or 4-(3S,5R,8R,9S,10S,12R,13S,14S,17R)-3-(((2S,4S,5R,6R)-5-(((2S,4S,5S,6R)-5-((4-benzyl-2-methyl-1,4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yloxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-12,14-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)furan-2(5H)-one;




DMDMcB or 4-((3S,5R,8R,9S,10S,12R,13S,14S,17R)-3-(((2S,4S,5R,6R)-5-(((2S,4S,5S,6R)-5-(((3,3-dimethylcyclobutyl)methyl)-2-methyl-1,4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-12,14-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)furan-2(5H)-one;

29

30




DtxMcP or 4-((3S,5R,8R,9S,10S,13R,14S,17R)-3-(((2S,4S, 5R,6R)-5-(((2S,4S,5S,6R)-5((4-(cyclopropylmethyl)-2- 25
methyl-1,4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetra-
hydro-2H-pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-
2H-pyran-2-yl)oxy)-14-hydroxy-10,13-
dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-
yl)furan-2(5H)-one;



DtxcB or 4-((3S,5R,8R,9S,10S,13R,14S,17R)-3-(((2S,4S, 5R,6R)-5-(((2S,5S,6R)-5((4-cyclobutyl-2-methyl-1,4-ox- 65
azepan-7-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-
pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-
2-yl)oxy)-14-hydroxy-10,13-dimethylhexadecahydro-1H-
cyclopenta[a]phenanthren-17-yl)furan-2(5H)-one;


31

32



DAz or 4-((3S,5R,8R,9S,10S,12R,13S,14S,17R)-3-
 (((2R,4S,5S,6R)-5-(((2S,4S,5S,6R)-5-((4-(azetidin-3-yl)-2-
 methyl-1,4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetra-
 hydro-2H-pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-
 2H-pyran-2-yl)oxy)-12,14-dihydroxy-10,13-
 dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-
 yl)furan-2(5H)-one;

25




DOx or 4-((3S,5R,8R,9S,10S,12R,13S,14S,17R)-12,14-di-
 hydroxy-3-(((2R,4S,5S,6R)-4-hydroxy-5-(((2S,4S,5S,6R)-
 4-hydroxy-6-methyl-54((2-methyl-4-(oxetan-3-yl)-1,4-ox-
 azepan-7-yl)oxy)tetrahydro-2H-pyran-2-yl)oxy)-6-
 methyltetrahydro-2H-pyran-2-yl)oxy)-10,13-
 dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-
 yl)furan-2(5H)-one;

60

65


33

34



DTh or 4-((3S,5R,8R,9S,10S,12R,13S,14S,17R)-12,14-dihydroxy-3-(((2R,4S,5S,6R)-4-hydroxy-5-4(((2S,4S,5S,6R)-4-hydroxy-6-methyl-5-((2-methyl-4-(thietan-3-yl)-1,4-oxazepan-7-yl)oxy)tetrahydro-2H-pyran-2-yl)oxy)-6-methyltetrahydro-2H-pyran-2-yl)oxy)-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)furan-2(5H)-one; and

25



DHis or 4-((3S,5R,8R,9S,10S,12R,13S,14S,17R)-3-(2R,4S,5S,6R)-5-(((2S,4S,5S,6R)-5-((4-(2-(1H-imidazol-5-yl)ethyl)-2-methyl-1,4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-12,14-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)furan-2(5H)-one.


60

65


35

36

In some embodiments, the compound is any one of the structures presented below:




DMSM or 4-((3S,5R,8R,9S,10S,12R,13S,14S,17R)-12, 25
14-dihydroxy-3-(((2S,4S,5R,6R)-4-hydroxy-5-(((2S,4S,5S,
6R)-4-hydroxy-6-methyl-5-((2-methyl-4-((methylsulfonyl
methyl)-1,4-oxazepan-7-yl)oxy)tetrahydro-2H-pyran-2-yl)
oxy)-6-methyltetrahydro-2H-pyran-2-yl)oxy)-10,13-
dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-
yl)furan-2(5H)-one;



DESM or 4-((3S,5R,8R,9S,10S,12R,13S,14S,17R)-12, 60
14-dihydroxy-3-(((2S,4S,5R,6R)-4-hydroxy-5-(((2S,4S,5S,
6R)-4-hydroxy-6-methyl-5-((2-methyl-4-(2-(methylsulfo-
nyl)ethyl)-1,4-oxazepan-7-yl)oxy)tetrahydro-2H-pyran-2-
yl)oxy)-6-methyltetrahydro-2H-pyran-2-yl)oxy)-10,13- 65
dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-
yl)furan-2(5H)-one; and

37


38



DESA or 2-7-(((2R,3S,4S,6S)-6-(((2R,3R,4S,6S)-6-(((3S,5R,8R,9S,10S,12R,13S,14S,17R)-12,14-dihydroxy-10,13-dimethyl-17-(5-oxo-2,5-dihydrofuran-3-yl)hexadecahydro-1H-cyclopenta[a]phenanthren-3-yl)oxy)-4-hydroxy-2-methyltetrahydro-2H-pyran-3-yl)oxy)-4-hydroxy-2-methyltetrahydro-2H-pyran-3-yl)oxy)-2-methyl-1,4-oxazepan-4-yl)ethanesulfonamide.

In some embodiments, the compound is any one of the structures presented below:

organic and inorganic anions and cations. The term “organic or inorganic cation” refers to counter-ions for an acid. The counter-ions can be chosen from the alkali and to alkaline earth metals, (such as lithium, sodium, potassium, barium, aluminum and calcium), ammonium and the like. Furthermore, the term includes salts that form by standard acid-base reactions of basic groups and organic or inorganic acids. Such acids include hydrochloric, hydrofluoric, hydrobromic, trifluoroacetic, sulfuric, phosphoric, acetic, succinic, citric,



DEDA or 4-(((3S,5R,8R,9S,10S,12R,13S,14S,17R)-3-(((2R,4S,5S,6K)-5-(((2S,4S,5S,6R)-5-((4-(2-(dimethylamino)ethyl)-2-methyl-1,4-oxazepan-7-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-4-hydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-12,14-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)furan-2(5H)-one.

The present embodiments further encompass any enantiomers, diastereomers, optical isomers, prodrugs, solvates, hydrates, polymorphs, geometrical isomers and/or pharmaceutically acceptable salts of the compounds described herein.

Any one or more of the compounds presented herein may be present as a salt. The term “salt” encompasses both basic and acid addition salts, and include salts formed with

lactic, maleic, fumaric, cholic, pamoic, mucic, D-camphoric, phthalic, tartaric, salicylic, methanesulfonic, benzenesulfonic, p-toluenesulfonic, sorbic, picric, benzoic, cinnamic, and like acids.

As used herein, the phrase “pharmaceutically acceptable salt” refers to a charged species of the parent compound and its counter-ion, which is typically used to modify the solubility characteristics of the parent compound and/or to reduce any significant irritation to an organism by the parent compound, while not abrogating the biological activity and properties of the administered compound. A pharmaceutically acceptable salt of a compound as described herein can alternatively be formed during the synthesis of the compound, e.g., in the course of isolating the compound from a reaction mixture or re-crystallizing the compound.

In the context of some of the present embodiments, a pharmaceutically acceptable salt of the compounds described herein may optionally be an acid addition salt comprising at least one basic (e.g., amine and/or guanidine) group of the compound which is in a positively charged form (e.g., wherein the basic group is protonated), in combination with at least one counter-ion, derived from the selected base, that forms a pharmaceutically acceptable salt.

The acid addition salts of the compounds described herein may therefore be complexes formed between one or more basic groups of the compound and one or more equivalents of an acid.

Depending on the stoichiometric proportions between the charged group(s) in the compound and the counter-ion in the salt, the acid additions salts can be either mono-addition salts or poly-addition salts.

The phrase “mono-addition salt”, as used herein, refers to a salt in which the stoichiometric ratio between the counterion and charged form of the compound is 1:1, such that the addition salt includes one molar equivalent of the counterion per one to molar equivalent of the compound.

The phrase “poly-addition salt”, as used herein, refers to a salt in which the stoichiometric ratio between the counterion and the charged form of the compound is greater than 1:1 and is, for example, 2:1, 3:1, 4:1 and so on, such that the addition salt includes two or more molar equivalents of the counter-ion per one molar equivalent of the compound.

An example, without limitation, of a pharmaceutically acceptable salt would be an ammonium cation or guanidinium cation and an acid addition salt thereof.

The acid addition salts may include a variety of organic and inorganic acids, such as, but not limited to, hydrochloric acid which affords a hydrochloric acid addition salt, hydrobromic acid which affords a hydrobromic acid addition salt, acetic acid which affords an acetic acid addition salt, ascorbic acid which affords an ascorbic acid addition salt, benzenesulfonic acid which affords a besylate addition salt, camphorsulfonic acid which affords a camphorsulfonic acid addition salt, citric acid which affords a citric acid addition salt, maleic acid which affords a maleic acid addition salt, malic acid which affords a malic acid addition salt, methanesulfonic acid which affords a methanesulfonic acid (mesylate) addition salt, naphthalenesulfonic acid which affords a naphthalenesulfonic acid addition salt, oxalic acid which affords an oxalic acid addition salt, phosphoric acid which affords a phosphoric acid addition salt, toluenesulfonic acid which affords a p-toluenesulfonic acid addition salt, succinic acid which affords a succinic acid addition salt, sulfuric acid which affords a sulfuric acid addition salt, tartaric acid which affords a tartaric acid addition salt and trifluoroacetic acid which affords a trifluoroacetic acid addition salt. Each of these acid addition salts can be either a mono-addition salt or a poly-addition salt, as these terms are defined herein.

As used herein, the term “enantiomer” refers to a stereoisomer of a compound that is superposable with respect to its counterpart only by a complete inversion/reflection (mirror image) of each other. Enantiomers are said to have “handedness” since they refer to each other like the right and left hand. Enantiomers have identical chemical and physical properties except when present in an environment which by itself has handedness, such as all living systems. In the context of the present embodiments, a compound may exhibit one or more chiral centers, each of which exhibiting an R- or an S-configuration and any combination, and compounds according to some embodiments of the present invention, can have any their chiral centers exhibit an R- or an S-configuration.

The term “diastereomers”, as used herein, refers to stereoisomers that are not enantiomers to one another. Diastereomerism occurs when two or more stereoisomers of a compound have different configurations at one or more, but not all of the equivalent (related) stereocenters and are not mirror images of each other. When two diastereoisomers differ from each other at only one stereocenter they are epimers. Each stereo-center (chiral center) gives rise to two different configurations and thus to two different stereoisomers. In the context of the present invention, embodiments of the present invention encompass compounds with multiple chiral centers that occur in any combination of stereo-configuration, namely any diastereomer.

All stereoisomers, optical and geometrical isomers of the compounds of the present invention are contemplated, either in admixture or in pure or substantially pure form. The compounds of the present invention can have asymmetric centers at one or more of the atoms. Consequently, the compounds can exist in enantiomeric or diastereomeric forms or in mixtures thereof. The present invention contemplates the use of any racemates (i.e. mixtures containing equal amounts of each enantiomers), enantiomerically enriched mixtures (i.e., mixtures enriched for one enantiomer), pure enantiomers or diastereomers, or any mixtures thereof. The chiral centers can be designated as R or S or R,S or d,D, l,L or d,l, D,L.

The term “prodrug” refers to an agent, which is converted into the active compound (the active parent drug) in vivo. Prodrugs are typically useful for facilitating the administration of the parent drug. They may, for instance, be bioavailable by oral administration whereas the parent drug is not. A prodrug may also have improved solubility as compared with the parent drug in pharmaceutical compositions. Prodrugs are also often used to achieve a sustained release of the active compound in vivo. An example, without limitation, of a prodrug would be a compound of the present invention, having one or more carboxylic acid moieties, which is administered as an ester (the “prodrug”). Such a prodrug is hydrolyzed in vivo, to thereby provide the free compound (the parent drug). The selected ester may affect both the solubility characteristics and the hydrolysis rate of the prodrug.


The term “solvate” refers to a complex of variable stoichiometry (e.g., di-, tri-, tetra-, penta-, hexa-, and so on), which is formed by a solute (the compound of the present invention) and a solvent, whereby the solvent does not interfere with the biological activity of the solute. Suitable solvents include, for example, ethanol, acetic acid and the like.

The term “hydrate” refers to a solvate, as defined hereinabove, where the solvent is water.

The present invention also includes solvates of the compounds of the present invention and salts thereof. “Solvate” means a physical association of a compound of the invention with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation. “Solvate” encompasses both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include ethanulates, methanulates and the like. “Hydrate” is a solvate wherein the solvent molecule is water.

The present invention also includes polymorphs of the compounds of the present invention and salts thereof. The term “polymorph” refers to a particular crystalline state of a substance, which can be characterized by particular physical properties such as X-ray diffraction, IR spectra, melting point, and the like.

According to some aspects of some embodiments of the present invention, the compound is represented by general Formula III:



Formula III

wherein:

X is H or OH;

R' is selected from the group consisting of OH, C₁-C₆ alkyl, C₁-C₆ haloalkyl (C₁-C₆ alkyl substituted with at least one halo), $-(CR^bR^c)_nSi(R^a)_3$, $-(CR^bR^c)_n-C(=Y)-NR_1R_2$, $-(CR^bR^c)_n-C(=Y)-NHOH$, $-(CR^dR^e)_n-C(=Y)-COOR_3$, $-NHC(=Y)NR_1R_2$ and $-(CR^bR^c)_n-NH_2$;

Y is O or S;

R₁, R₂ and R₃ are each independently H or a C₁-C₄ alkyl;


R_a is a C₁-C₄ alkyl;

R^b, R^c, R^d and R^e are each independently selected from H, a C₁-C₄ alkyl and a C₁-C₄ hydroxyalkyl; and

n is 0, 1 or 2;

or

R' is represented by general Formula



Formula II


the wiggled line represents the N-link to the compound;

A is a spacer moiety or a covalent bond; and

B is a cyclic moiety;

or

R' is represented by general Formula



Formula II


the wiggled line represents the N-link to the compound;

A is a spacer moiety or a covalent bond; and

B is selected from the group consisting of an alkylsulfonyl, arylsulfonyl and a sulfonamide;

or

R' is represented by general Formula II:



Formula II

the wiggled line represents the N-link to the compound;

A is a spacer moiety or a covalent bond; and

B is $-NR_1R_2$, wherein R₁ and R₂ are each independently H or a C₁-C₄ alkyl provided that at least one of R₁ and R₂ is a C₁-C₄ alkyl, including any pharmaceutically acceptable salt, prodrug, hydrate, solvate, enantiomer and diastereomer thereof, and any mixtures thereof, and a pharmaceutically acceptable carrier.

Additional alternative structures of R' are described in WO2015/029035, which is hereby specifically incorporated by reference as if fully set forth herein, thereby describing each and all such additional alternative structures.

Process of Preparing the Compounds:

According to an aspect of some embodiments of the present invention, there is provided a process for preparing the compounds represented by general Formula I, the process includes:

converting the third digitoxose moiety of digoxin or digitoxin into a dialdehyde; and

reacting said dialdehyde with a reagent represented by general formula, IV:




Formula IV

A is a spacer moiety or a covalent bond as described hereinabove; and

B is selected from the group consisting of a cyclic moiety, an alkylsulfonyl, arylsulfonyl, a sulfonamide, a secondary amine and a tertiary amine, as described hereinabove.

Scheme 1



As can be seen in Scheme 1, the conversion of the third digitoxose moiety of digoxin or digitoxin is typically effected by selective oxidation thereof into a dialdehyde by reacting digoxin or digitoxin with a reagent that breaks apart 1,2-diols (vicinal diols) to afford aldehydes and/or ketones, such as sodium periodate (NaIO_4);

The reaction of the dialdehyde is effected by reductive amination of the dialdehyde using a free amine of R ($\text{R}-\text{NH}_2$) in the presence of, e.g., NaCNBH_3 .

Selective Affinity and Inhibition:

The results presents in the Examples section below (see, e.g., Table 2) show clearly that compounds encompassed by general Formula I, according to embodiments of the present invention, exhibit high selectivity for $\alpha 2\beta 3$ over $\alpha 1\beta 1$ compared to digoxin and digitoxin. The results obtained for cyclic R-substituents (R comprising a cyclic moiety), some

with 4 carbon atoms, provide a clear indication for selective structural interactions of the modified sugar of digoxin with the $\beta 3$ subunit. This effect may also be combined with an enhanced interaction with the $\alpha 2$ subunit. The detailed enzymological study, presented hereinbelow, has been obtained with the purified detergent-soluble human Na,K-ATPase isolated isoform complexes, as well as with 1.5 intact bovine ciliary NPE cells, i.e., selective inhibition of human Na,K-ATPase isolated isoform complexes provided corroborating indication of inhibition by the compounds provided herein also of isoform mixture ($\alpha 1\beta 1$ plus $\alpha 2\beta 3$) as these are expressed in native NPE cell membranes.

According to some embodiments, the compounds presented herein, such as a compound represented by general Formula I or by general Formula III, exhibit an affinity to at least one isoform of Na,K-ATPase.

In the context of embodiments of the present invention, isoforms of Na,K-ATPase include any combination of $\alpha 1$, $\alpha 2$, $\alpha 3$ and $\alpha 4$ complexed with $\beta 1$, $\beta 2$ and $\beta 3$; hence, isoforms of Na,K-ATPase include $\alpha 1\beta 1$, $\alpha 1\beta 2$, $\alpha 1\beta 3$, $\alpha 2\alpha 1$, $\alpha 2\beta 2$, $\alpha 2\beta 3$, $\alpha 3\beta 1$, $\alpha 3\beta 2$, $\alpha 3\beta 3$, $\alpha 4\beta 1$, $\alpha 4\beta 2$ and $\alpha 4\beta 3$.

According to some embodiments of the present invention, a compound represented by general Formula I or by general Formula III, has a higher affinity to isoforms containing an $\alpha 2$ subunit, compared to its affinity to isoforms containing one of the α subunits, such as $\alpha 1$, as demonstrated in the Examples section that follows below. The phrase "higher affinity to", as used herein, is a relative term that means that a given ligand molecule (compound, inhibitor, drug, etc.) is attracted and can bind to and (form a) complex with a target entity (protein, enzyme, drug-target) more strongly compared to its binding to another target entity. Affinity can be measured directly and indirectly by a number of methodologies. In some embodiments of the present invention, the affinity is referred to in terms of K_i (inhibition constant) or K_D (dissociation constant), as these terms are known in the art. For example, a small value for K_i means that an inhibitor has a higher effective affinity to the enzyme relative to an affinity of the same inhibitor to another enzyme or another inhibitor to the same enzyme.

According to some embodiments of the present invention, a compound represented by general Formula I or by general Formula III, is a selective inhibitor of $\alpha 2$ -containing isoforms of Na,K-ATPase, compared to $\alpha 1$ -containing isoforms of Na,K-ATPase, as demonstrated in the Examples section that follows below.

The terms "selective inhibitor of $\alpha 2$ -containing isoforms of Na,K-ATPase" or "selective inhibitor of the $\alpha 2$ isoform of Na,K-ATPase" or " $\alpha 2$ -selective inhibitor", as used herein interchangeably, refer to a compound that inhibits $\alpha 2$ -containing isoforms of Na,K-ATPase to a greater degree than the compound inhibits other isoforms of Na,K-ATPase, such as those containing $\alpha 1$. In some embodiments, the compounds described herein are selective for the $\alpha 2\beta 1$, $\alpha 2\beta 2$ and/or $\alpha 2\beta 3$ isoforms of Na,K-ATPase over the $\alpha 1\beta 1$ isoform thereof. In some embodiments, the selectivity of the compound for the $\alpha 2$ -containing isoform of Na,K-ATPase (e.g., $\alpha 2\beta 1$, $\alpha 2\beta 2$ and/or $\alpha 2\beta 3$ isoform) is at least about 4-fold (300% more selective) over other isoforms, or at least 5-fold (400%), at least 6-fold (500%), at least 8-fold (700%), at least 10-fold, at least 16-fold, at least 20-fold, at least 30-fold, or at least 50-fold greater inhibition of the $\alpha 2$ -containing isoform of Na,K-ATPase over other isoforms of Na,K-ATPase.

Uses of $\alpha 2$ -Selective Inhibitors:

The experimental work presented in the Examples section below demonstrates a clear correlation between increased $\alpha 2\beta 3$ -selectivity of the compounds presented herein and potency and duration in reducing IOP of either pharmacologically raised or basal (normal) IOP. Thus the results support a central role of $\alpha 2\beta 3$ in production of aqueous humour. The mechanism of the IOP reduction has been shown to involve inhibition of active Na and K fluxes via NPE cells and reduction of inflow of aqueous humour after topical administration of compound(s) and permeation thereof via the cornea.

Without being bound by any particular theory, it is assumed that the low K_i values for inhibition of $\alpha 2\beta 3$ (K_i of about 4 nM) and hydrophobic properties of the compounds according to some embodiments of the present invention, suggest that both traits contribute to the potency and long duration of action exhibited by the compounds provided herein. The finding that the compounds provided herein are

effective in reducing basal (normal) IOP as low as 25-30% (see, e.g., FIGS. 2A-D and FIGS. 3A-C), as well as the reduction of acute 4AP-induced raised IOP (see, e.g., FIGS. 1A-D), provides an insight into the mechanism of action of the compounds.

The compounds presented herein are useful as topical ophthalmological agents for treatment of glaucoma, since they exhibit at least one of efficacy and low levels of side-effects. In relations to alternative drugs, the compounds presented herein are useful as topical ophthalmological agents for treatment of glaucoma, since they have been shown to exhibit at least one of improved efficacy, extended duration of desired effects and reduced side-effects, compared to currently available drugs, exemplified in the Examples section that follows below by the first-line drug latanoprost, and iopidine, used in short-term adjunctive therapy of chronic glaucoma. In general, currently available drugs include β -adrenergic antagonists and carbonic anhydrase inhibitors, that reduce the rate of aqueous humor production, or prostaglandin analogs, cholinergic agonists and sympathomimetics, that increase the rate of outflow through the trabecular meshwork and uveoscleral pathway. In this respect, the experimental results presented hereinbelow, show 25-30% higher reduction in basal IOP effected by the compounds presented herein in the rabbits, compared to Latanoprost, the current first-line drug for treatment of glaucoma.

In principle, the ability of the compounds presented herein, according to some embodiments, to reduce the basal IOP, could be relevant not only to optical hypertension and primary open angle glaucoma but also to normotensive glaucoma for which reduction of IOP below the basal level is required.

Regarding local toxicity, the rationale for making $\alpha 2\beta 3$ -selective inhibitors included not only the expectation of high potency but also of minimal local side-effects. In the Examples section below it is shown that corneal swelling, used to assess such adverse effect, is not observed at least over several days of topical administration of the compounds presented herein (see, e.g., Table 4). Similar evidence for safety was observed by inspection of local redness and irritation.

Concerning possible systemic toxicity of the $\alpha 2\beta 3$ -selective compounds presented herein, it is expected that cardiotoxicity, such as is associated with clinical use of digoxin, would be minimal, because the $\alpha 2$ -selective compounds are likely to be intrinsically non-cardiotoxic, compared to a non-selective cardiac glycoside, and also because the pharmacokinetic concentration is likely to be low (see, e.g., inhibition results in Table 2).

The results presented hereinbelow indicate that after repeated administration to over several weeks of the compounds according to embodiments of the present invention, no signs of adverse effects have been observed, indicating that compounds represented by general Formula I or III are not toxic and exhibit essentially no adverse effects.

Thus, the $\alpha 2$ -selective inhibitors presented herein (compounds represented by general Formula I or by general Formula III) can be used to reduce intraocular pressure in a subject in need thereof, while taking advantage of the selective affinity these inhibitors exhibit towards $\alpha 2$ -containing isoforms of Na,K-ATPase, which minimize adverse effects associated with unselective inhibition of the other isoforms.

According to an aspect of some embodiments of the present invention, there is provided a pharmaceutical composition which includes as an active ingredient any one of

the compounds represented by general Formula I and a pharmaceutically acceptable carrier.

Hereinafter, the term "pharmaceutically acceptable carrier" refers to a carrier or a diluent that does not cause significant irritation to an organism and does not abrogate the biological activity and properties of the administered compound. Examples, without limitations, of carriers are: propylene glycol, saline, emulsions and mixtures of organic solvents with water, as well as solid (e.g., powdered) and gaseous carriers.

Herein the term "excipient" refers to an inert substance added to a pharmaceutical composition to further facilitate administration of a compound. Examples, without limitation, of excipients include calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils and polyethylene glycols.

Techniques for formulation and administration of drugs may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton, Pa., latest edition, which is incorporated herein by reference.

Pharmaceutical compositions of the present invention may be manufactured by processes well known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more pharmaceutically acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the compounds presented herein into preparations which, can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

According to some embodiments, the administration is effected topically, extraocularly, intraocularly and/or intravitreally. In some embodiments, the pharmaceutical composition is formulated as an ophthalmic composition suitable for topical, extraocular, intraocular and/or intravitreal administration to the eye of the subject. According to some embodiments, the pharmaceutical composition of the invention is an ophthalmic composition which is administered topically onto the eye of a subject for facilitating effective intraocular levels of the compound and for preventing unnecessary and unintentional levels of the compound in other tissues and/or organs. Such a non-systemic, site-specific administration reduces the side effects associated with the compounds.

In the context of some embodiments of the present invention, topical and/or extraocular administration is effected by applying the compound(s), or compositions and medicaments comprising the compound(s) to the eye or a bodily surface near the eye. According to some embodiments, the composition may take the form of an eye-drop solution, a spray, an eye wash solution, an ointment, a lotion, a suspension, a gel or a cream. The topical pharmaceutical compositions may be in the form of eye-drops to be applied by instillation into the eye or may be in the form of a viscous ointment, gel or cream to be applied by an ointment onto the ocular surface and may contain control release means for facilitating sustained release over a prolonged period of time.

In the context of some embodiments of the present invention, intraocular and/or intravitreal administration is effected by injecting the compound(s), or compositions and medicaments comprising the compound(s) into the eye or

into a bodily tissue near the eye. According to some embodiments, the composition may take the form of an injectable solution.

According to some embodiments, oral or otherwise systemic administration in a dosage effective for reducing the intraocular pressure is also possible. For example, the composition may be administered by a dermal patch for extended release.

According to some embodiments, the administration is effected orally. For oral administration, the compounds presented herein can be formulated readily by combining the compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds presented herein to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for oral ingestion by a patient. Pharmacological preparations for oral use can be made using a solid excipient, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carbomethylcellulose; and/or physiologically acceptable polymers such as polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Pharmaceutical compositions, which can be used orally, include push-fit capsules made of gelatin as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules may contain the active ingredients in admixture with filler such as lactose, binders such as starches, lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the compounds presented herein may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for the chosen route of administration.

For injection, the compounds presented herein may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological saline buffer with or without organic solvents such as propylene glycol, polyethylene glycol.

For transmucosal administration, penetrants are used in the formulation. Such penetrants are generally known in the art.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active aminoglycoside compounds doses.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds presented herein are conveniently delivered in the form of an aerosol spray presentation (which typically includes powdered, liquefied and/or gaseous carriers) from a pressurized

pack or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichloro-tetrafluoroethane or carbon dioxide. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compounds presented herein and a suitable powder base such as, but not limited to, lactose or starch.

The compounds presented herein may be formulated for parenteral administration, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multidose containers with optionally, an added preservative. The compositions may be suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical compositions for parenteral administration include aqueous solutions of the compounds preparation in water-soluble form. Additionally, suspensions of the compounds presented herein may be prepared as appropriate oily injection suspensions and emulsions (e.g., water-in-oil, oil-in-water or water-in-oil in oil emulsions). Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acids esters such as ethyl oleate, triglycerides or liposomes. Aqueous injection suspensions may contain substances, which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol or dextran. Optionally, the suspension may also contain suitable stabilizers or agents, which increase the solubility of the compounds presented herein to allow for the preparation of highly concentrated solutions.

Alternatively, the compounds presented herein may be in powder form for to constitution with a suitable vehicle, e.g., sterile, pyrogen-free water, before use.

The compounds presented herein may also be formulated in rectal compositions such as suppositories or retention enemas, using, e.g., conventional suppository bases such as cocoa butter or other glycerides.

The pharmaceutical compositions herein described may also comprise suitable solid of gel phase carriers or excipients. Examples of such carriers or excipients include, but are not limited to, calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin and polymers such as polyethylene glycols.

Pharmaceutical compositions suitable for use in context of the present invention include compositions wherein the active ingredients are contained in an amount effective to achieve the intended purpose. More specifically, a therapeutically effective amount means an amount of compounds presented herein effective to prevent, alleviate or ameliorate symptoms of the disorder, or prolong the survival of the subject being treated.

Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

For any compounds presented herein used in the methods of the present embodiments, the therapeutically effective amount or dose can be estimated initially from activity assays in animals. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes the mutation suppression levels as determined by activity assays (e.g., the concentration of the test compounds which achieves a substantial read-through of the truncation mutation). Such information can be used to more accurately determine useful doses in humans.

Toxicity and therapeutic efficacy of the compounds presented herein can be determined by standard pharmaceutical procedures in experimental animals, e.g., by determining the EC_{50} (the concentration of a compound where 50% of its maximal effect is observed) and the LD_{50} (lethal dose causing death in 50% of the tested animals) for a subject compound. The data obtained from these activity assays and animal studies can be used in formulating a range of dosage for use in human.

The dosage may vary depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See e.g., Fingl et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p. 1).

Dosage amount and interval may be adjusted individually to provide plasma levels of the compounds presented herein which are sufficient to maintain the desired effects, termed the minimal effective concentration (MEC). The MEC will vary for each preparation, but can be estimated from *in vitro* data; e.g., the concentration of the compounds necessary to achieve 50-90% expression of the whole gene having a truncation mutation, i.e. read-through of the mutation codon. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. HPLC assays or bioassays can be used to determine plasma concentrations.

Dosage intervals can also be determined using the MEC value. Preparations should be administered using a regimen, which maintains plasma levels above the MEC for 10-90% of the time, preferable between 30-90% and most preferably 50-90%.

Depending on the severity and responsiveness of the chronic condition to be treated, dosing can also be a single periodic administration of a slow release composition described hereinabove, with course of periodic treatment lasting from several days to several weeks or until sufficient amelioration is effected during the periodic treatment or substantial diminution of the disorder state is achieved for the periodic treatment.

The amount of a composition to be administered will, of course, be dependent on the subject being treated, the severity of the affliction, the manner of administration, the judgment of the prescribing physician, etc. Compositions of the present invention may, if desired, be presented in a pack or dispenser device, such as an FDA (the U.S. Food and Drug Administration) approved kit, which may contain one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as, but not limited to a blister pack or a pressurized container (for inhalation). The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accompanied by a notice associated with the container in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the compositions for human or veterinary administration. Such notice, for example, may be of labeling approved by the U.S. Food and Drug Administration for prescription drugs or of an approved product insert. Compositions comprising a compound according to the present embodiments, formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition or diagnosis, as is detailed hereinabove.

In accordance with other embodiments, the compounds presented herein may be loaded into a drug-delivery device

to be inserted or implanted into the eye of the subject for allowing releasing of the compound in a controlled and continuous rate, by dissolving, diffusion or leaching, thus maintaining effective therapeutic concentration over a prolonged period of time. The drug-delivery device may be for example a biocompatible thin film loaded with the active agent, inserted for example beneath the lower eyelid. In some embodiments, the drug-delivery device is a contact lens or any other ophthalmic device, as these are known in the art.

In some embodiments, the pharmaceutical composition is packaged in a packaging material and identified in print, or on the packaging material, for use in reducing intraocular pressure (IOP).

According to an aspect of some embodiments of the present invention, there is provided a use of the compounds represented by general Formula I, or a pharmaceutical composition comprising the same, for the manufacture of a medicament for reducing intraocular pressure (IOP).

According to an aspect of some embodiments of the present invention, there is provided a method of reducing intraocular pressure (IOP) in a subject, the method includes administering to a subject in need thereof a therapeutically effective amount of a compound represented by general Formula I, or a pharmaceutical composition comprising the same.

According to some embodiments of the present invention, intraocular pressure (IOP) is associated with glaucoma, low-tension glaucoma and normal-tension glaucoma. Hence, the compounds, compositions and medicaments presented herein are useful in treating, without limitation, glaucoma, low-tension glaucoma and normal-tension glaucoma.

As used herein, the term "treating" includes abrogating, substantially inhibiting, slowing or reversing the progression of a condition, substantially ameliorating clinical or aesthetic symptoms of a condition or substantially preventing the appearance of clinical or aesthetic symptoms of a condition.

As used herein, the phrase "therapeutically effective amount" describes an amount of the polymer being administered which will relieve to some extent one or more of the symptoms of the condition being treated.

In some embodiments, the concentration of the compounds presented herein in the pharmaceutical compositions or medicaments presented herein, is in the range of about 1 to about 5,000 $\mu\text{g/ml}$ of composition, preferably from about 80 to about 800 $\mu\text{g/ml}$ and the formulation is preferably applied in one to four doses per day wherein each dose contains about 1 to 125 μg of the compound, or from about 2 to about 20 μg of the compound.

Co-Administration with Other IOP Reducing Agents:

In the experimental section presented below, it has been demonstrated that a combination of a compound according to some embodiments of the present invention and Latanoprost resulted in a notable increase in the duration of the effects for the combination compared to the individually administered agents.

In the context of some embodiments of the present invention, an active agent which is not a compound represented by general Formula I or general Formula III, is referred to herein as "another agent" or "other agent".

According to an aspect of some embodiments of the present invention, there is provided a pharmaceutical composition which includes as active ingredients at least one other agent (active ingredient) which is in use in reducing IOP and is not a compound represented by general Formula

I or general Formula III, a compound represented by general Formula I or general Formula III, and a pharmaceutically acceptable carrier.

Non-limiting examples of active ingredients which are in use in reducing IOP and which are referred to herein other agents, include prostaglandin analogs, β -blockers, adrenergic agents, α_2 -adrenergic receptor agonists, miotic agents, carbonic anhydrase inhibitors and cholinergic agonists.

Non-limiting examples of prostaglandin analogs include Latanoprost (Xalatan), Bimatoprost (Lumigan) and Travoprost (Travatan).

Non-limiting examples of β -blockers include Timolol (Timoptic) and Betaxolol (Betoptic).

A non-limiting example of adrenergic agents is Brimonidine (Alphagan).

A non-limiting example of miotic agents is Pilocarpine (Isotocarpine, Pilocar).

Non-limiting examples of carbonic anhydrase inhibitors include Dorzolamide (Trusopt), Brinzolamide (Azopt) and Acetazolamide (Diamox).

Non-limiting examples of cholinergic agonists include carbachol (Miostat), echothiophate (Phospholine) and pilocarpine (Isopto Carpine, Pilopine).

Administration of more than one active agent to a subject is generally known in the art as co-administration. The term "co-administration" as used herein, refers to a concomitant administration of more than one active agent (active ingredient) to a subject, whereas in the context of embodiments presented herein, the term "concomitant" means that the co-administered active agents are present in the subject (PK), or otherwise exert an effect (PD), at similar, identical, consecutive or partially overlapping periods of time.

In the context of co-administration of more than one active agent, the terms "substantially simultaneous" and "rapid succession" correspond to the term "concomitant" as used herein, namely meaning that the period of time between a first administration and a second administration of more than one active agent is sufficiently short to be regarded as a single administration event, and/or a number of administrations of different active agents takes place within 5-60 minutes or less. Optionally, each administration in such "rapid succession" delivers to the user a different amount or composition of one or more pharmaceutically active agents. Alternatively, two or more of the administrations provide the same composition and amount of the one or more pharmaceutically active agents. In some embodiments, the later administration of the second active agent is performed at such timing that the first active agent of a previous administration within the same rapid succession does not yet have a significant pharmacodynamic effect or before it can be measured (pharmacokinetically) by, e.g., blood concentration thereof the first active agent.

According to some embodiments of the present invention, the co-administration of at least one other agent which is in use in reducing IOP and is not a compound represented by general Formula I or general Formula III, and a compound represented by general Formula I or general Formula III, exhibits a potentiating effect, namely that the effect of the co-administered active ingredients is greater in at least one parameter, such as magnitude or duration, compared to the effects exhibited by each of the active ingredients when administered alone (separately).

Accordingly, there is provided a use of at least one other agent which is in use in reducing IOP, and a compound represented by general Formula I or general Formula III, for the manufacture of a medicament for reducing IOP.

According to an aspect of some embodiments of the present invention, there is provided a method of reducing IOP in a subject in need thereof, the method includes co-administering to the subject a therapeutically effective amount of at least one other agent which is in use in reducing IOP, and a therapeutically effective amount of a compound represented by general Formula I or general Formula III.

According to an aspect of some embodiments of the present invention, there is provided a method of reducing IOP in a subject in need thereof, the method includes co-administering to the subject a synergistically effective amount of at least one other agent which is in use in reducing IOP, and a synergistically effective amount of a compound represented by general Formula I or general Formula III. It is noted herein that a synergistically effective amount is also a therapeutically effective amount in the sense of providing the desired therapeutic effect, and is smaller than the therapeutically effective amount of a singly-administered active ingredient.

In the context of the herein provided uses and method of co-administration of other agents and the compounds presented herein, the other agent is a prostaglandin analog.

In the context of the herein provided uses and method of co-administration of other agents and the compounds presented herein, the other agent is a β -blocker.

In the context of the herein provided uses and method of co-administration of other agents and the compounds presented herein, the other agent is an adrenergic agent.

In the context of the herein provided uses and method of co-administration of other agents and the compounds presented herein, the other agent is an α 2-adrenergic receptor agonist.

In the context of the herein provided uses and method of co-administration of other agents and the compounds presented herein, the other agent is a miotic agent.

In the context of the herein provided uses and method of co-administration of other agents and the compounds presented herein, the other agent is a carbonic anhydrase inhibitor.

In the context of the herein provided uses and method of co-administration of other agents and the compounds presented herein, the other agent is a cholinergic agonist.

Cardiotonic Agent:

According to some embodiments, the selectivity of the compounds presented herein can be utilized in the treatment of other medical conditions. For example, an α 2-selective inhibitor, such as the compounds presented herein and represented by general Formula I, can be used as an effective cardiotonic agent, with reduced cardiotoxicity, compared to known agents such as digoxin.

According to an aspect of embodiments of the present invention, there is provided a method of treating a heart condition which is carried out by administering to a subject in need thereof a therapeutically effective amount of a compound represented by general Formula I.

According to an aspect of embodiments of the present invention, there is provided a pharmaceutical composition which includes as an active ingredient a compound represented by general Formula I and a pharmaceutically acceptable carrier, identified in print, or on a packaging material, for use in a treatment of a heart condition.

According to an aspect of embodiments of the present invention, there is provided a uses of a compound represented by general Formula I or a pharmaceutical composition comprising the same, for the manufacture of a medicament for treating a heart condition.

Examples of heart conditions which are relevant in the context of embodiments of the present invention, include, without limitation, atrial fibrillation, atrial flutter, mitral stenosis, chronic heart failure and congestive heart failure.

In some embodiments, the present invention provides cardiotonic compositions comprising a therapeutically effective amount of the compounds represented by general Formula I, or a pharmaceutical composition comprising the same. In accordance with such embodiments, the compounds may be formulated for oral, buccal, topical, intravenous, parenteral or rectal administration.

A compound represented by general Formula I or a compound represented by general Formula III can be used according to embodiments of the present invention to treat a heart condition in combination with one or more other drugs for treating a heart condition, such as, but not limited to, selective and nonselective β -blocker agents, anticoagulation agents, angiotensin-converting-enzyme inhibitors (ACE inhibitors) and angiotensin II receptor antagonists.

According to an aspect of embodiments of the present invention, there is provided a method of treating a heart condition in a subject in need thereof, which includes co-administering to the subject a therapeutically effective amount of an agent selected from the group consisting of a β -blocker, an anticoagulation agent, an angiotensin-converting-enzyme inhibitor and an angiotensin II receptor antagonist; and a compound represented by Formula III, as these are described herein, and a pharmaceutically acceptable carrier.

According to an aspect of embodiments of the present invention, there is provided a pharmaceutical composition that includes, as active ingredients, at least one ingredient selected from the group consisting of a β -blocker, an anticoagulation agent, an angiotensin-converting-enzyme inhibitor and an angiotensin II receptor antagonist; and a compound represented by Formula III, as these are described herein.

According to an aspect of embodiments of the present invention, there is provided a use of an agent selected from the group consisting of a β -blocker, an anticoagulation agent, an angiotensin-converting-enzyme inhibitor and an angiotensin II receptor antagonist, and a compound represented by Formula III, as these are described herein, for the manufacture of a medicament for treating a heart condition.

Nonselective β -blocker agents include propranolol, bucindolol, carteolol, carvedilol, labetalol nadolol, oxprenolol, penbutolol, pindolol, sotalol, timolol, eucommia bark (herb); β 1-selective agents include acebutolol, atenolol, betaxolol, bisoprolol, celiprolol, esmolol, metoprolol, nebivolol; β 2-selective agents include butaxamine and ICI-118, 551; and β 3-selective agents include SR 59230A.

Anticoagulation agents include heparin, dicumarol, coumadin (warfarin) and aspirin.

ACE inhibitors include captopril, zofenopril, enalapril (vasotec, renitec), ramipril (altace, prilace, ramace, ramiwin, triatec, tritace), quinapril (accupril), perindopril (coversyl, aceon, perindo), lisinopril (listril, lopril, novatec, prinitiv, zestril), benazepril (lotensin), imidapril (tanatril), trandolapril (mavik, odrik, gopten), cilazapril (inhibace) and fosinopril (fositen, monopril).

Angiotensin II receptor antagonists include losartan, EXP 3174, candesartan, valsartan, irbesartan, telmisartan, eprosartan, olmesartan, azilsartan and fimasartan.

According to some embodiments, the compound represented by general Formula I or the compound represented by general Formula III, according to embodiments of the present invention, and any one of the other drugs for treating a

heart a condition can be co-formulated in a single composition, or be formulated into individual compositions.

High-Risk Pharmacokinetics:

When plasma concentrations of active drug depend exclusively on a single metabolic pathway, any condition that inhibits that pathway (be it disease-related, genetic, or due to a drug interaction) can lead to dramatic changes in drug concentrations and marked variability in drug action. This problem of high-risk pharmacokinetics is especially pronounced in drug elimination that relies on a single pathway. In this case, inhibition of the elimination pathway leads to striking elevation of drug concentration. For drugs with a narrow therapeutic window, this leads to an increased likelihood of dose-related toxicity. An example is digoxin, whose elimination is dependent on P-glycoprotein; many drugs inhibit P-glycoprotein activity (amiodarone, quinidine, erythromycin, cyclosporine, itraconazole) and coadministration of these with digoxin reduces digoxin clearance, and increases toxicity unless maintenance doses are lowered. Drugs with a high risk of generating pharmacokinetic interactions with digoxin include antacids and bile acid sequestrants, which can cause reduced absorption; inhibitors of CYPs and of P-glycoprotein such as amiodarone, quinidine, amiodarone, verapamil, cyclosporine, itraconazole and erythromycin which can cause decreased clearance.

Since compounds encompassed under Formula III are in essence digoxin derivatives that exhibit lower digoxin toxicity, these compounds can be used with any of the drugs described above which exhibit adverse drug-drug interaction with digoxin.

According to an aspect of embodiments of the present invention, there is provided use of the compounds represented by general Formula III for the treatment of a medical condition or a combination of medical conditions, with is treatable by digoxin and a drug having an adverse interaction with digoxin.

Prodrugs:

Prodrugs corresponding to the compounds presented herein are contemplated in the context of the present invention in order to provide active agents that exhibit improved pharmacokinetic and/or pharmacodynamics profile, and/or a reduction in adverse effects. As known in the art, a prodrug is typically a chemical derivative of the corresponding drug,


being chemically modified such that a naturally occurring metabolic pathway in the subject's system can convert it to the parent drug molecule during the time in which the prodrug/drug is still present in the system. For instance, drugs that exhibit poor absorption due to high hydrophobicity, can be modified to exhibit improved bioavailability by introducing chemical functionalities that increase the solubility of the prodrug compared to the parent active compound. In other cases where the active compound exhibits adverse effects in the GI-tract, the prodrug is a modified parent active compound that is metabolized by enzymes back to the parent compound substantially at the target tissues, organs or cells.

In the context of some embodiments of the present invention, the prodrugs are metabolized to afford the corresponding (parent) compound by naturally occurring metabolic agents, such as enzymes. In some embodiments, the chemical group of the prodrug is bioliable or biodegradable, such as in the case of an ester, which can be hydrolyzed by esterases to afford a hydroxyl of the parent compound.

According to some embodiments, the compounds presented herein exhibit several structural positions that can offer potential locations for chemical modifications on route to becoming suitable prodrugs. For example, a digoxin skeleton exhibits four hydroxyl groups which can potentially be converted into, e.g., esters, under various conditions. Similarly, the digitoxin skeleton exhibits three hydroxyl moieties. Since each hydroxyl is located at a different chemical environment, is it chemically converted under varying conditions, thereby allowing the production of a single or a multiple hydroxyl-to-ester conversion. Thus, in the case of digoxin, a mono-, a bis-, a tris- or a tetra-modified parent compound can be afforded, offering a variety of prodrugs of the same parent compound, each exhibiting a different pharmacokinetic/pharmacodynamics profile.

In addition, R and R' substituents (see, Formula I and Formula III respectively) may exhibit a chemical position which is readily converted into another, bioliable (biodegradable) moiety, and thus can be used to afford prodrugs from the corresponding parent compound.

According to some embodiments, prodrugs of the compounds having the general Formula I or general Formula III, are represented by general Formula V:



wherein X is H or PD₃, and each of PD₁-PD₄ is H or independently represents a modified hydroxyl group turned into a bioliable functionality, provided that at least one of PD₁-PD₄ is not H.

In some embodiments, the compounds presented herein are converted into a prodrug by modifying any one of the hydroxyl groups on the digoxin or digitoxin moiety of the compounds into any one of a methoxymethyl ether, a tetrahydropyranyl ether, a t-butyl ether, an allyl ether, a benzyl ether, a t-butyl dimethylsilyl ether, a t-butyl diphenylsilyl ether, an acetic acid ester (Ac), ethyl, propyl, butyl, t-butyl or pivalic acid ester and a benzoic acid ester, in any combination. Each of the converted hydroxyl functionalities may be metabolized (biodegraded) back to the parent hydroxyl by one or more metabolic and/or enzymatic systems.

According to some embodiments, any one of PD₁-PD₄ in general Formula V is selected from the group consisting of a methoxymethyl ether, a tetrahydropyranyl ether, a t-butyl ether, an allyl ether, a benzyl ether, a t-butyl dimethylsilyl ether, a t-butyl diphenylsilyl ether, an acetic acid ester (Ac), ethyl, propyl, butyl, t-butyl or pivalic acid ester and a benzoic acid ester.

As can be seen in the Examples section that follows below, the exemplary digoxin-derived compound DcB, according to some embodiments of the present invention, has been converted successfully into the corresponding bit-acetyl (BisAcDcB) prodrug and the tris-acetyl (TrisAcDcB) prodrug (see, Example 6).

Isolation and Use of Na,K-ATPase Isoforms:

According to an aspect of some embodiments of the present invention, there is provided a method of determining an apparent affinity of the compound represented by general Formula I to at least one isoform of Na,K-ATPase, the method includes contacting an isoform of Na,K-ATPase with the compound in an activity measurement setup and determining the apparent affinity of the compound to the isoform.

According to an aspect of some embodiments of the present invention, there is provided a method of isolating an isoform of Na,K-ATPase of a mammal, the method includes: transforming yeast cells with a clone that comprises an α chain sequence and a β chain sequence of the Na,K-ATPase; expressing the clone in the yeast cells; and isolating the isoform,

wherein:

the α chain sequence is selected from the group consisting of $\alpha 1$, $\alpha 2$, $\alpha 3$ and $\alpha 4$; and

the β chain sequence is selected from the group consisting of $\beta 1$, $\beta 2$ and $\beta 3$.

In some embodiments, the isolated isoform is $\alpha 2\beta 2$.

In some embodiments, the isolated isoform is $\alpha 2\beta 3$.

According to an aspect of some embodiments of the present invention, there is provided an isolated $\alpha 2\beta 2$ isoform of Na,K-ATPase of a mammal having at least 70% purity.

According to an aspect of some embodiments of the present invention, there is provided an isolated $\alpha 2\beta 3$ isoform of Na,K-ATPase of a mammal having at least 70% purity.

The yeast-expressed α/β subunits have distinct levels of glycosylation compared to those in the human-expressed subunits. The distinct yeast-expressed glycosylation pattern does not substantially affect the activity of the enzyme but may increase its stability.

According to an aspect of some embodiments of the present invention, there is provided an isolated $\alpha 2\beta 2$ iso-

form of Na,K-ATPase of a mammal having a yeast-characterizing glycosylation pattern.

According to an aspect of some embodiments of the present invention, there is provided an isolated $\alpha 2\beta 3$ isoform of Na,K-ATPase of a mammal having a yeast-characterizing glycosylation pattern.

In some embodiments, the isolated isoforms presented herein have an amino acid sequence which identical, substantially similar or derived from any one of the isoform of human Na,K-ATPase.

SEQ ID NOS.

Description SEQ ID No.

- | | |
|----|--|
| 15 | Amino acid sequence of $\alpha 1$ subunit of human Na,K-ATPase (P05023) 1 |
| | Amino acid sequence of $\alpha 2$ subunit of human Na,K-ATPase (P50993) 2 |
| | Amino acid sequence of $\alpha 3$ subunit of human Na,K-ATPase (P13637) 3 |
| 20 | Amino acid sequence of $\alpha 4$ subunit of human Na,K-ATPase (Q13733) 4 |
| | Amino acid sequence of $\beta 1$ subunit of human Na,K-ATPase (P05026) 5 |
| 25 | Amino acid sequence of HIS tagged $\beta 1$ subunit of human Na,K-ATPase (P05026) 6 |
| | Amino acid sequence of $\beta 2$ subunit of human Na,K-ATPase (P14415) 7 |
| | Amino acid sequence of HIS tagged $\beta 1$ subunit of human Na,K-ATPase (P14415) 8 |
| 30 | Amino acid sequence of $\beta 3$ subunit of human Na,K-ATPase (P54709) 9 |
| | Amino acid sequence of HIS tagged $\beta 3$ subunit of human Na,K-ATPase (P54709) 10 |
| 35 | Nucleotide sequence of $\alpha 1$ subunit of human NaX-ATPase (ATP1A1) 11 |
| | Nucleotide sequence of $\alpha 2$ subunit of human Na,K-ATPase (ATP1A2) 12 |
| | Nucleotide sequence of $\alpha 3$ subunit of human Na,K-ATPase (ATP1A3) 13 |
| 40 | Nucleotide sequence of $\alpha 4$ subunit of human Na,K-ATPase (ATP1A4) 14 |
| | Nucleotide sequence of HIS tagged $\beta 1$ subunit of human Na,K-ATPase (ATP1B1) 15 |
| 45 | Nucleotide sequence of HIS tagged $\beta 2$ subunit of human Na,K-ATPase (ATP1B2) 16 |
| | Nucleotide sequence of HIS tagged $\beta 3$ subunit of human Na,K-ATPase (ATP1B3) 17 |
| | Nucleotide sequence of HIS tagged human FXYD1 18 |
| 50 | Amino acid sequence of HIS tagged human FXYD1 19 |

As used herein the term "about" refers to 10%.

The terms "comprises", "comprising", "includes", "including", "having" and their conjugates mean "including but not limited to".

The term "consisting of" means "including and limited to".

The term "consisting essentially of" means that the composition, method or structure may include additional ingredients, steps and/or parts, but only if the additional ingredients, steps and/or parts do not materially alter the basic and novel characteristics of the claimed composition, method or structure.

As used herein, the singular form "a", "an" and "the" include plural references unless the context clearly dictates otherwise. For example, the term "a compound" or "at least one compound" may include a plurality of compounds, including mixtures thereof. Throughout this application,

various embodiments of this invention may be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

Whenever a numerical range is indicated herein, it is meant to include any cited numeral (fractional or integral) within the indicated range. The phrases "ranging/ranges between" a first indicate number and a second indicate number and "ranging/ranges from" a first indicate number "to" a second indicate number are used herein interchangeably and are meant to include the first and second indicated numbers and all the fractional and integral numerals therebetween.

As used herein the term "method" refers to manners, means, techniques and procedures for accomplishing a given task including, but not limited to, those manners, means, techniques and procedures either known to, or readily developed from known manners, means, techniques and procedures by practitioners of the chemical, pharmacological, biological, biochemical and medical arts.

It is expected that during the life of a patent maturing from this application many relevant inhibitors exhibiting selectivity towards α_2 -containing isoforms of Na,K-ATPase, as defined herein, will be uncovered and the scope of this term is intended to include all such new selective inhibitors a priori.

It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention, which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable subcombination or as suitable in any other described embodiment of the invention. Certain features described in the context of various embodiments are not to be considered essential features of those embodiments, unless the embodiment is inoperative without those elements.

Various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below find experimental support in the following examples.

EXAMPLES

Reference is now made to the following examples, which together with the above descriptions illustrate some embodiments of the invention in a non-limiting fashion.

Example 1

Compound Synthesis

Compounds represented by general Formula I, according to embodiments of the present invention, were synthesized as a series of perhydro-1-4-oxazepine derivatives of the third digitoxose of digoxin or digitoxin following a general pro-

cedure described elsewhere [Adamczyk, M. et al., *Steroids*, 1995, 60(11), pp. 753-758] by selective oxidation with NaIO_4 and reductive amination with NaCNBH_3 . The compounds were purified by HPLC and masses, $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra were determined to verify both correctness of the structure and purity.

The structure and names based on of the amine substituents (R-NH_2), as well as their theoretical and measured masses, are presented in Table 1 below, which includes other digoxin derivatives with non-cyclic moieties (last eight compounds in Table 1).

TABLE 1

R	Name	Theoretical mass	Measured mass (with Na+ ion)
	cyclopropyl	803.48	826.45
	methylcyclopropane	817.50	840.44
	ethylcyclopropane	831.51	854.58
	cyclobutyl	817.50	840.43
	cyclopentyl	831.51	854.51
	cyclohexyl	845.53	868.65
	benzyl	853.50	876.44
	methyl(3,3-dimethylcyclobutane)	859.54	882.59
	methylcyclopropane	801.50	824.48
	cyclobutyl	801.50	824.44
	methyl	777.47	800.57
	ethyl	791.48	814.52
	2,2,2-trifluoroethyl	845.45	868.14
	propyl	805.50	828.27
	iso-propyl	805.50	828.41
	iso-butyl	819.51	842.66
	tert-butyl	819.51	842.41
	methyl(trimethylsilyl)	849.51	872.77
	(methylsulfonyl)methyl	855.44	878.49
	(methylsulfonyl)ethyl	869.46	892.29
	(sulfonamide)ethyl	870.45	893.47
	azetidiny	818.49	841.55
	oxetanyl	819.48	842.41
	thietanyl	835.45	858.37
	histaminyl	857.50	880.45
	(N,N-dimethylamine)ethyl	834.52	857.62
	aminoethyl	806.49	829.55
	(methylsulfonyl)methyl	855.44	878.49

Names of compounds encompassed under Formula I wherein $\text{X}=\text{OH}$ (digoxin derivatives) include Dcp, DMcP, DEcP, DcB, DcB, DcB, DcB and DMDMcB; names of compounds encompassed under Formula I wherein $\text{X}=\text{H}$ (digitoxin derivatives) include, DtxMcP and DtxcB.

Example 2

Expression, Purification and Characterization of Recombinant Human Na,K-ATPase Isolated Isoforms

Plasmid construction for the expression of $\alpha_1\beta_1$, $\alpha_1\beta_2$, $\alpha_2\beta_1$, $\alpha_2\beta_2$ and $\alpha_2\beta_3$ Na,K-ATPase was conducted by generation of pHi1-D2 expression vector containing cDNA of human α_1 and His10 tagged human β_1 was described previously [Lifshitz Y, et al., 2007, *Biochemistry*, 46(51), pp. 14937-14950]. cDNA of human β_2 and β_3 in pSD5 were a gift from K. Geering, University Lausanne, Switzerland. Open reading frames and flanking regions of human β_2 and human β_3 were amplified by PCR using primers containing BglII and SalI cleavage sites. The resulting fragments were subcloned into pHi1-D2- $\text{h}\alpha_2$ /His10-p β_1 to create pHi1-D2- $\text{h}\alpha_2$ /His10-h β_2 and pHi1-D2- $\text{h}\alpha_2$ /His10-h β_3 , respectively. Correct integration and sequence was confirmed by sequencing.

Yeasts were grown in BMG (100 mM potassium phosphate pH 6, 1.34% yeast nitrogen base, $4 \times 10^{-5}\%$ biotin, 0.3% glycerol) to OD 6-8 and expression was induced in BMM (100 mM potassium phosphate pH 6, 1.34% yeast nitrogen base, $4 \times 10^{-5}\%$ biotin, 0.5% methanol added daily).

Pichia pastoris transformation, yeast growth, membrane preparation and His-tag purification of recombinant human $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha\beta_2$ and $\alpha_2\beta_3$ Na,K-ATPase were carried out essentially as described previously [Katz A, et al., 2010, *J Biol Chem*, 285(25), pp. 19582-19592; and Katz A, et al., 2014, *J Biol Chem*, 289(30), pp. 21153-21162].

Expression and purification of $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha\beta_2$ and $\alpha_2\beta_3$ complexes was conducted in small scale whole cell lysates that were prepared as describe in Loayza, D. et al., *Mol. Cell. Biol.*, 1998, 18, p. 779-789. Yeast membrane production and expression of recombinant Na,K-ATPase as detergent soluble complexes was performed as described in Habeck, M. et al., *J. Biol. Chem.*, 2015, 290, pp. 4829-4842, using modified lipid content (C12E8, 0.1 mg/ml; SOPS, 0.07 mg/ml; cholesterol, 0.01 mg/ml).

Na,K-ATPase $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_2\beta_1$ and $\alpha_2\beta_3$ complexes were reconstituted with FXYD1 and purified in a mixture of 0.1 mg/ml. C12E8, 0.07 mg/ml SOPS and 0.01 mg/ml cholesterol.

The purified isoform complexes (0.3-0.5 mg/ml) were eluted from the BD-Talon beads in a solution containing Imidazole 170 mM, NaCl 100 mM; Tricine.HCl 20 mM pH 7.4; C12E8, 0.1 mg/ml; SOPS 0.07 mg/ml, cholesterol 0.01 mg/ml, glycerol 25%, by gravity-column. The proteins were stored at -80°C . Protein purity was determined by gel electrophoresis and protein concentration was determined with BCA (B9643 Sigma).

The specific Na,K-ATPase activity was highest for $\alpha_1\beta_1$ (16.4 ± 0.7 $\mu\text{mol}/\text{mg}/\text{min}$) followed by $\alpha_2\beta_1$ (10.9 ± 0.6) and $\alpha_2\beta_3$ (10.7 ± 1.9). The $\alpha_2\beta_2$ isoform had the lowest activity (8.4 ± 1.4). The second significantly different parameter is the apparent K^+ -affinity. $\text{K}_{0.5} \text{K}^+$ for $\alpha_2\beta_1$ was 2.7 ± 0.14 mM compared to 1.47 ± 0.06 mM for $\alpha_1\beta_1$. $\alpha_2\beta_2$ and $\alpha_2\beta_3$ had an even lower affinity than $\alpha_2\beta_1$ with apparent $\text{K}_{0.5}$ values of 7.4 ± 0.19 mM and 6.4 ± 0.50 mM, respectively. Na-titrations revealed that the affinity for Na^+ -ions was not different between $\alpha_1\beta_1$ and $\alpha_2\beta_1$ whereas $\alpha_2\beta_2$ and $\alpha_2\beta_3$ had a somewhat higher Na-affinity.

The reduced apparent affinity for K^+ together with an increased affinity for Na^+ indicates that the conformational equilibrium of $\alpha_2\beta_2$ and $\alpha_2\beta_3$ might be shifted towards E1. In order to test this hypothesis, the apparent affinity of vanadate was determined for all four isoform complexes. Vanadate is a phosphate analogue that binds to the E2 conformation, mimicking the transition state E2PK₂ during dephosphorylation, thus inhibiting the enzyme. An three α_2 isoforms had a lower vanadate affinity compared to $\alpha_1\beta_1$ (0.48 μM). $\alpha_2\beta_2$ had the lowest vanadate affinity (34 μM)

followed by $\alpha_2\beta_3$ (1.9 μM) and $\alpha_2\beta_1$ (3.5 μM). Thus, the order of inhibition by vanadate equals the order of potassium activation ($\text{K}_{0.5}\text{K}^+$ and K_i vanadate $\alpha_1\beta_1 < \alpha_2\beta_1 < \alpha_2\beta_3 < \alpha_2\beta_2$) supporting the hypothesis proposed above.

Example 3

Selective Inhibition Assays of Isolated Na,K-ATPase

To screen for isoform selectivity of the digoxin derivatives we compared inhibition of Na,K-ATPase activity of purified detergent-soluble human isoform complexes $\alpha_1\beta_1\text{FXYD1}$, $\alpha_2\beta_1\text{FXYD1}$, $\alpha_2\beta_2\text{FXTD1}$ and $\alpha_2\beta_3\text{FXYD1}$. Although all the preparations and assays were conducted with FXYD1 in order to stabilize the complexes, the FXYD1 suffix is omitted in naming of isoform complexes for simplicity.

Na,K-ATPase activity of $\alpha/\beta\text{PFXYD1}$ complexes was measured over one hour at 37°C in a medium containing 130 mM NaCl, 5 mM KCl, 3 mM MgCl_2 , 1 mM EGTA, 25 mM Histidine, pH 7.4 and 1 mM ATP using the PiColor Lock gold malachite green assay (Inova Biosciences).

The Na,K-ATPase activities were $\alpha_1\beta_1$, 21.5 ± 5.3 $\mu\text{moles}/\text{min}/\text{mg}$; $\alpha_2\beta_1$, 18.7 ± 1.8 $\mu\text{moles}/\text{min}/\text{mg}$, and $\alpha_2\beta_3$, 10.7 ± 1.9 $\mu\text{moles}/\text{min}/\text{mg}$ protein. As discussed below, an important kinetic property in relation to inhibition by cardiac glycosides is $\text{K}_{0.5}$ for activation by K^+ : $\alpha_1\beta_1$ 1.25 ± 0.05 mM, $\alpha_2\beta_1$ 2.7 ± 0.14 mM and $\alpha_2\beta_3$ 6.4 ± 0.50 mM, respectively.

Selectivity of the compounds for various isolated isoforms of human Na,K-ATPase was determined essentially as described before [Katz, A. et al., *J Biol Chem.*, 2010, 285(25), pp. 19582-19592].

ATPase activity assays as well as titrations with NaCl, KCl and vanadate were performed as described in Lifshitz-2007 and Loayza-1998 using PiColorLock™ malachite green assay (Inova Bioscience). Inhibitor assays were performed as described in Katz-2010. [³H]ouabain binding and K^+ -[³H]digoxin displacement assays were performed as described in Katz-2010.

The percent inhibition VCG/V0 was calculated and K_i values were obtained by fitting the data to the function $\text{VCG}/\text{V0} = \text{K}_i / ([\text{CG}] + \text{K}_i) + c$ (CG stands for cardiac glycoside). Inhibition was estimated in 3-5 separate experiments and average K_i values \pm standard error of the mean (SEM) were calculated. The ratios $\text{K}_i \alpha_1\beta_1/\alpha_2\beta_1$, $\alpha_1\beta_1/\alpha_2\beta_2$ and $\alpha_1\beta_1/\alpha_2\beta_3$ was calculated for each compound.

Table 2 shows the K_i values and selectivity ratios ($\text{K}_i \alpha_1\beta_1:\alpha_2\beta_1$, $\text{K}_i \alpha_1\beta_1:\alpha_2\beta_2$, and $\alpha_1\beta_1:\alpha_2\beta_3$) for inhibition of Na,K-ATPase activity of compounds according to embodiments of the present invention, as well as some digoxin derivatives having a non-cyclic moiety, compared to digoxin and digitoxin. Table 2 is arbitrarily sorted according to column "K_i ratio $\alpha_1\beta_1/\alpha_2\beta_3$ " marked by "***".

TABLE 2

Compound Name	No. of C atoms in R	K _i in nM \pm SEM				Selectivity		
		$\alpha_1\beta_1$	$\alpha_2\beta_1$	$\alpha_2\beta_2$	$\alpha_2\beta_3$	K _i ratio $\alpha_1\beta_1/\alpha_2\beta_1$	K _i ratio $\alpha_1\beta_1/\alpha_2\beta_2$	* K _i ratio $\alpha_1\beta_1/\alpha_2\beta_3$
DeB (cyclic)	4	135 \pm 11	8 \pm 1.25	6 \pm 1	4 \pm 0.15	16.9	22.2	33.6
DMcP (cyclic)	4	95.8 \pm 13.7	18.3 \pm 1.6	8.0 \pm 0.8	4.3 \pm 0.6	5.2	12	22.2
DESM (sulfonyl)	2	464 \pm 14	49.2 \pm 1.9	31.7 \pm 3.2	24.7 \pm 2.1	9.4	14.6	18.8
DiB	4	92 \pm 8.9	20.6 \pm 1.4	10 \pm 0.8	5.8 \pm 0.6	4.4	9.0	16
DESA (sulfonamide)	2	301 \pm 23	38.9 \pm 2.2	31.5 \pm 4.4	20.1 \pm 0.9	7.7	9.5	15
DiP	3	149 \pm 20.7	28.9 \pm 1.7	16.7 \pm 1.9	10.3 \pm 1.8	5.1	8.9	14.4

TABLE 2-continued

Compound Name	No. of C atoms in R	Ki in nM \pm SEM				Selectivity		
		$\alpha 1\beta 1$	$\alpha 2\beta 1$	$\alpha 2\beta 2$	$\alpha 2\beta 3$	Ki ratio $\alpha 1\beta 1/\alpha 2\beta 1$	Ki ratio $\alpha 1\beta 1/\alpha 2\beta 2$	* Ki ratio $\alpha 1\beta 1/\alpha 2\beta 3$
DcP (cyclic)	3	109 \pm 6.2	14.6 \pm 11.6	13.0 \pm 1.3	8.1 \pm 1.36	7.5	8.5	13.4
DEcP (cyclic)	5	86.1 \pm 7	14.3 \pm 2	12.1 \pm 1.9	7.2 \pm 1	6.02	7.1	12.02
DMSM (sulfonyl)	3	944 \pm 123	137 \pm 9.8	123 \pm 7.3	89 \pm 8.7	6.9	7.7	10.6
DBz (cyclic)	6	57.9 \pm 15.5	10.1 \pm 2.2	6.8 \pm 1.2	5.6 \pm 1.6	5.7	8.5	10.3
DcH (cyclic)	6	70.4 \pm 4.1	15.2 \pm 3.7	15.3 \pm 2.9	11.7 \pm 4.5	4.6	4.6	10.1
DCF ₃	2	119 \pm 15.0	28.6 \pm 0.9	18.1 \pm 1.9	12.4 \pm 1.5	4.1	6.5	9.6
DEt	2	137.9 \pm 12.6	23.2 \pm 0.9	16.4 \pm 1.6	14.4 \pm 1.27	5.9	8.3	9.5
DMe	1	103 \pm 5.6	15.3 \pm 1.2	20.36 \pm 1.8	10.8 \pm 0.6	6.7	5.1	9.5
DP	3	87.7 \pm 7.9	18.3 \pm 1.68	10.5 \pm 1.8	9.8 \pm 1.1	4.8	8.3	8.8
DtB	4	135 \pm 12.1	21.6 \pm 5.6	18.4 \pm 1.1	16.3 \pm 0.28	6.2	7.3	8.2
DDMcB (cyclic)	7	31.6 \pm 0.5	8.6 \pm 1.4	5.1 \pm 0.5	3.9 \pm 0.7	3.69	6.16	8.19
DtxcB (cyclic)	4	30.7 \pm 7.2	5.4 \pm 0.5	5.3 \pm 0.8	4.3 \pm 0.6	5.6	5.8	7.1
DtxMcP (cyclic)	4	25 \pm 4	4.2 \pm 0.4	5.4 \pm 1.2	3.7 \pm 0.5	5.9	4.6	6.7
DTMS (silyl)	4	72.6 \pm 17.6	24.3 \pm 4.0	14.3 \pm 1.3	11.1 \pm 1.6	3	5.1	6.5
Digoxin	0	268 \pm 13.8	58.7 \pm 5.4	58 \pm 1.9	42.8 \pm 3.0	4.5	4.6	6.2
DcPe (cyclic)	5	138 \pm 21	33.4 \pm 7.5	33.5 \pm 11.9	27.6 \pm 9.5	4.1	4.1	5
Digitoxin	0	89 \pm 15.8	29.5 \pm 2.7	40.7 \pm 6.7	28.8 \pm 5.9	3	2.1	3.1

As can be seen in Table 2, some compounds show an even greater selectivity ratio towards $\alpha 2\beta 2$ and $\alpha 2\beta 3$, particularly $\alpha 2\beta 3$. Based on the ratios of Ki values, several derivatives show significantly improved selectivity for $\alpha 2\beta 3$ compared to $\alpha 2\beta 1$ over $\alpha 1\beta 1$, in particular DMcP and DcB. DMcP and DcB show exceptional selectivity for $\alpha 2\beta 2$ and $\alpha 2\beta 3$ over $\alpha 1\beta 1$ of 22-fold and 33-fold, respectively, and very low Ki values for inhibition of $\alpha 2\beta 3$ (Ki about 4 nM). The full inhibition curves of DMcP and DcB emphasize the extent of the difference between $\alpha 1\beta 1$ and $\alpha 2\beta 3$.

Bearing in mind the large differences in $K_{0.5}$ potassium ions, a point to be aware of in analyzing the results presented in Table 2 is the well-known K-cardiac glycoside antagonism. Digoxin itself has moderate selectivity for $\alpha 2\beta 1$ over $\alpha 1\beta 1$ (about 4-fold) and the data in Table 2 shows increased selectivity for $\alpha 2\beta 3$ over $\alpha 1\beta 1$ (about 6-fold). The $\alpha 2\beta 1$ -selectivity of digoxin is attributed to a combination of increased binding affinity for $\alpha 2$ over $\alpha 1$ and also reduced K-digoxin antagonism in the Na,K-ATPase reaction conditions (with K, 5 mM). Similarly, the increased selectivity of digoxin for $\alpha 2\beta 3$ compared to $\alpha 2\beta 1$ is attributable to reduced K-cardiac glycoside antagonism due to the higher $K_{0.5}$ K of $\alpha 2\beta 3$ compared to $\alpha 2\beta 1$.

The compounds presented herein show a notable increase of the ratio of Ki's $\alpha 1\beta 1:\alpha 2\beta 3$ compared to $\alpha 1\beta 1:\alpha 2\beta 1$, which must be partly due to the reduced K-cardiac glycoside antagonism. However, the difference for the most $\alpha 2\beta 3$ -selective compounds is significantly greater than that for digoxin and cannot be explained only by this factor. In particular, there is a distinct structural effect in that the maximal $\alpha 2\beta 3$ -selectivity is seen for R-substituents with cyclic moiety, as well as those with four carbon atoms. In the case of DMcP and DcB, the Ki for $\alpha 1\beta 1$ is about 2-3-fold lower than for digoxin itself but the Ki for $\alpha 2\beta 3$ is about 10-fold lower than for digoxin, thus raising selectivity for $\alpha 2\beta 3$ over $\alpha 1\beta 1$ to more than 22-fold and more than 33-fold, respectively. It is also noticeable that for derivatives with five or a higher number of carbon atoms in the R-substituents, such as DEcP, DcPe, DcH, DBz and DDMcB, although the Ki values for all isoforms are all lower than for digoxin itself, the selectivity for $\alpha 2\beta 3$ over $\alpha 1\beta 1$, 10-12-fold, is also lower than the selectivity observed in four carbon R-substituents, DMcP and DcB. The same is true for compounds with R-substituent three carbons or shorter.

Another structural insight comes from the results obtained for two digitoxin derivatives (DtxMcP and DtxcB) compared with the digoxin derivatives DMcP and DcB. As seen in Table 2, the Ki values are relatively low for all three isoform complexes $\alpha 1\beta 1$, $\alpha 2\beta 1$ and $\alpha 2\beta 3$, being reduced about 10-fold compared to digoxin. Consequently, these digitoxin derivatives showed little increase in selectivity for $\alpha 2\beta 3$ compared to digoxin itself. Thus, the absence of a single OH group in position 12 of the steroid moiety of digitoxin reduced the effect of the sugar modification on $\alpha 2\beta 3$ -selectivity.

Thus, modification of the third digitoxose moiety of digoxin, but not of digitoxin, with cyclic substituents such as McP and cB confers notable $\alpha 2\beta 3$ -selectivity, presumably due to selective interaction with $\beta 3$.

Example 4

Inhibition of Na,K-ATPase Activity Permeabilized Bovine NPE Cells

In order to prove the concept of selective inhibition of the compounds presented herein in Na,K-pumps of native ciliary epithelium, inhibition assays using Na,K-ATPase in NPE cells isolated from the ciliary body dissected out of bovine eyes were conducted.

Isolation of ciliary epithelium PE and NPE cells was carried out as previously described [Edelman, J L. et al., 1994, *Am J. Physiol*, 266(5 Pt 1), pp. C1210-1221], with some modifications, using 15-20 fresh bovine eyes. Ciliary bodies were isolated from bovine eyes and washed in ringer solution. The tissue was treated with trypsin and homogenized followed by separation on a density gradient of Metrizamide, which separates between the NPE and PE cells.

Using isolated human isoforms to calibrate the response of the antibodies, NPE cell lysates were shown to contain about 70% $\alpha 2$ and 30% $\alpha 1$, while PE were shown to contain about 90% $\alpha 1$ and 5-10% $\alpha 2$. After unmasking the Na,K-ATPase by treating the cells with alamethicin, Na,K-ATPase activity was measured and was found to be 0.195 ± 0.027 and 0.035 ± 0.008 nmoles/mg protein/min in NPE and PE cells, respectively. Ouabain-sensitive fractions of total ATPase activity were about 65% and 35% for NPE and PE cells, respectively.

To determine NaK-ATPase activity in the cells, the cells were incubated with 0.8 mg/ml alamecithin for 30 minutes at room temperature prior to transfer to the reaction medium containing 130 mM NaCl, 5 mM KCl, 3 mM MgCl₂, 25 mM histidine, pH 7.4, 1 mM EGTA, 1 nM sodium azide, 0.5 mM ATP, and were then incubated for 45 minutes at 37° C., with or without the tested inhibitors as indicated, or 0.5 mM ouabain to determine the ouabain insensitive ATPase activity. The data was fitted to one or two sites inhibition model.

Table 3 presents Ki values for inhibition of NPE Na,K-ATPase activity by digoxin, DMe, DMcP and DcB, fitted to a single site inhibition model.

TABLE 3

Compound	Ki, nM ± SEM	n
1 site model		
Digoxin	91.7 ± 10.2	4
DMe	15.6 ± 1.3	4
DMcP	7.9 ± 2.2	5
DcB	17.3 ± 2.5	4
2 sites model		
DcB	Ki α2 6.9 ± 2 Ki α1 151 ± 7.6 (A α2 0.66 ± 0.090 A α1 0.34 ± 0.097)	4

As can be seen in Table 3, the Ki values of the derivatives are all lower than that of digoxin. Since NPE cells contain about 70% α2 and 30% α1, Na,K-ATPase activity and inhibition should reflect the properties of the isoform mixture. Indeed, the detailed inhibition curve for the most α2β3-selective compound, DcB, was fitted better by a two site model, compared to fitting according to a one site model.

As can be seen in Table 3, the two site model provides the best fit parameters of 66% α2, Ki 6.9±2 nM; 34% α1, Ki 151±7.6 nM (Ki α1/α2=22), which are quite close to the proportions of α2β1 estimated in the immunoassays, and the selectivity ratio Ki α1β1/α2β3 is about 33.

Thus, it can be concluded that the selectivity properties of the digoxin derivative DcB observed with purified human isoforms is corroborated by the results obtained using intact NPE cells.

Example 5

Reduction of Intraocular Pressure

These experiments examined the effects of topically administered α2β3-selective digoxin derivatives, according to some embodiments of the present invention, DiB, DMcP and DcB, on IOP in rabbits. Due to the lower Ki for inhibition of α2β3, compared to digoxin, and the high hydrophobicity, these compounds were predicted to both permeate the cornea well and efficiently inhibit the α2β3 in the NPE ciliary epithelium, thus reducing inflow of aqueous humour and IOP.

New Zealand white rabbits (3-3.5 kg) about 1 year old, of either sex, were housed in pairs in cage in animal room conditions on a reversed, 12-hour dark/light cycle. For the experiments the animals were transferred to rabbit restrainers in a quiet and calm atmosphere. No ocular abnormalities were detected prior or during the experiments.

IOP measurements were made with a pneumatometer (Model 30, Reichert technologies) either after raising IOP with 4-aminopyridine (4AP; 1 drop 40 mg/ml), or on basal IOP after addition of one drop of 1 mM solution of digoxin

derivatives to the right eye (RE) and one drop of PBS to the left eye (LE) that served as control.

For comparison of effects of digoxin derivatives, such as DcB with a known glaucoma drug Latanoprost, three groups of five rabbits were used. Rabbits treated with Latanoprost, received the medication every day for 5 days before the start of the experiment. On the day of the experiment rabbits were treated at 5 minutes interval with one drop of 1 mM DcB, one drop of 0.005% Latanoprost (Xalatan™, Pfizer) or one drop each of DcB and Latanoprost (RE), or normal saline (LE, Control). IOP was measured every hour for 12 hours (DcB and Latanoprost alone) and after 24 hours (DcB with Latanoprost). Basal IOPs in both eyes, without any medication, were measured 5 days before and on the day of the experiment. All eyes were examined routinely by ophthalmic examinations and were free of any abnormalities. Corneal thickness (μm) was measured using an ultrasonic pachometer (Sonogage pachometer, Cleveland, USA).

Stock solutions of the tested compounds were dissolved in ethanol, and freshly diluted in phosphate buffer (PBS) for each experiment, such that the final ethanol concentration did not exceed 1%.

The first set of experiments examined the effects of the α2-inhibitor compounds, according to embodiments of the present invention, when applied just before 4AP, used as a pharmacological tool to transiently raise IOP. One drop of the tested compound (0.01-0.3 mM) was applied topically to the rabbit's eyes prior to the 4AP, and IOP was then measured over 5 hours.

FIGS. 1A-D present comparative plots of IOP as a function of time, showing the dose response of α2-inhibitor compounds, according to some embodiments of the present invention, in lowering IOP in live rabbits, wherein FIG. 1A shows the results obtained for DiB, FIG. 1B shows the duration of the effect of DiB while 4AP is added to every 2 hours so as to maintain the raised control IOP, FIG. 1C shows the results obtained for DMcP, and FIG. 1D shows the results obtained for DcB.

As can be seen in FIG. 1A, DiB, having Ki α1β1:α2β3 ratio of 16-fold) prevents the rise of the 4AP-induced IOP at concentrations of more than 0.030 mM while at 10 μM it is still effective. As can be seen in FIG. 1B, the duration of the DiB effect was shows that 1 mM DiB is effective for about 8 hours before the IOP begins to rise back up to control levels, which is a significantly long effect.

As can be seen in FIGS. 1B and 1C, DMcP, having a Ki α1β1:α2β3 ratio of 22-fold, and DcB, having a Ki α1β1:α2β3 ratio of 33-fold, exhibit similar IOP reduction as DiB (FIG. 1A), but in even lower concentrations. As can be seen in FIGS. 1B and 1C, low concentrations of 0.01 mM of DMcP or DcB are sufficient to prevent the 4AP-induced rise in IOP.

As can further be seen in FIGS. 1B and 1C, at higher concentrations of 0.1-0.3 mM, the IOP is reduced to levels which are significantly lower than the starting IOP. The latter observation implies that these compounds could reduce basal IOP even in the absence of 4AP.

FIGS. 2A-D present comparative plots of IOP as a function of time, demonstrating the capacity of the α2-inhibitor compounds, according to some embodiments of the present invention, to lower IOP below basal levels compared to a buffer control when administered topically to one eye of a rabbit, while the other eye received PBS as a control, wherein FIG. 2A shows the lack of effect of digoxin, FIG. 2B shows the lack of effect of MB (a non-cyclic moiety inhibitor), FIG. 2C shows the notable of effect of DMcP, and FIG. 2D shows the notable of effect of DcB.

67

As can be seen in FIG. 2A-B, neither digoxin nor DiB had a significant effect on lowering the basal IOP. This observation coincides with other observations with digoxin or other non-cyclic moiety digoxin derivatives, DMe, DGlyN, the latter two exhibiting enhanced selectivity for $\alpha 2\beta 1$.

As can be seen in FIGS. 2C-D, both DMcP and DcB significantly reduced the basal IOP by 20-25% (about 4 mm Hg for rabbit with a basal IOP of 17 mm Hg) over the test period of 4-5 hours. A higher concentration of DcB (2 mM) reduces the IOP similarly to 1 mM DcB.

This observation indicates that the $\alpha 2$ -inhibitor compounds, according to some embodiments of the present invention, can be used effectively to treat medical conditions where there is a need to lower IOP below what is considered to be a normal pressure, such as cases of low-tension glaucoma and normal-tension glaucoma.

A final set of experiments compared the effectiveness and duration of the effects of topical administration of DcB on basal IOP, with those of a widely used anti-glaucoma drug, Latanoprost, applied either alone or in combination (co-administration).

Groups of 5 rabbits were treated once a day for 5 days with Latanoprost and on the sixth day with DcB, Latanoprost, or DcB/Latanoprost combination. IOP measurements were made for the next 12 hours or over 24 hours for the group treated with both DcB and Latanoprost.

FIGS. 3A-C present comparative plots of IOP as a function of time, demonstrating the effect of $\alpha 2$ -inhibitor compounds, according to some embodiments of the present invention, to potentiate the drug Latanoprost in lowering IOP below basal levels, wherein FIG. 3A shows the effect of DcB alone, FIG. 3B shows the effect of Latanoprost alone, and FIG. 3C shows the effect of co-administering DcB with Latanoprost.

As can be seen in FIGS. 3A-C, compared to the basal IOP values of 17-18 mmHg, after 3-8 hours the steady-state IOP was lower by 3.5 ± 0.15 , 2.6 ± 0.11 and 3.44 ± 0.39 mmHg with DcB, Latanoprost and DcB/Latanoprost, respectively, corresponding to steady-state IOP's of 75-80%, 85% and 75-80% of the unchanged control values.

As can further be seen in FIGS. 3A-C, DcB, $\alpha 2$ -inhibitor compound according to some embodiments of the present invention, was about 25% more effective than Latanoprost in reducing the basal (normal) levels of IOP.

The combination of DcB and Latanoprost was similar in effect to DcB alone; however, with respect to the duration of the effect, for either DcB and Latanoprost, applied alone, the IOP returned to the control (normal) value after 12 hours, but with the combined DcB/Latanoprost treatment the low IOP was maintained for a significantly longer period and it returned to the control value only after 24 hours.

Latanoprost was applied for 5 days prior to the day of measurement, due to reports that this pre-treatment produces optimal effects on IOP in humans, although in these experiments with rabbits it seems that this was unnecessary since

68

the effects of Latanoprost were observed acutely and had dissipated completely after 24 hours.

The animals which received the treatment presented in FIGS. 3A-C were used for corneal thickness measurements after topically administering drops of DcB or Latanoprost to 4 rabbits once a day for 5 additional days, totaling 6 days of treatment with DcB or Latanoprost, and the results are presented in Table 4 below.

TABLE 4

Rabbit	Treatment to RE	Pachymetry (mm)	
		RE	LE
1	DcB	455	456
2	DcB	480	456
3	DcB	400	396
4	DcB	382	374
5	Latanoprost	467	471
6	Latanoprost	489	498

As seen in Table 4, there was no detectable effect of either drug on corneal thickness. In addition, by inspection, no significant redness or ocular irritation was observed.

In order to assess whether topical application of DcB damaged the tissues of the eye including cornea, iris, lens, ciliary body, retina, choroid and sclera, a histological examination was conducted after topical application, using one drop of 1 mM DcB daily for one week in one eye of each animal, with the other eye treated topically with one drop of PBS and serving as the control. Animals were sacrificed, eyes were removed, fixed in 10% neutral buffered formalin, processed routinely for histological examination, trimmed at 4 μ m, and stained with hematoxylin and eosin. No significant histological differences were observed in a comparative analysis of the treated and untreated eyes, indicating that topical treatment with DcB for a week does not cause tissue damage in the eye.

Example 6

Prodrug Synthesis and Characterization

Prodrugs of the compounds presented herein were developed in order to improve the pharmacokinetic profile of the active compounds, e.g., to avoid potential toxic effects in the cornea. The synthetic strategy included preparation of substantially inactive derivatives of the compounds, which are ineffective as an inhibitor of the major isoform of the Na,K-ATPase in the cornea (mainly the $\alpha 1\beta 1$ isoform), and which can penetrate into the eye and thereafter are converted into the active form by biodegradation processes. The biodegraded and active form of the prodrug can then inhibit the $\alpha 2\beta/3$ complex in the ciliary NPE cells and reduce inflow of aqueous humor.



Synthesis:

The synthesis of an exemplary prodrug compound, according to some embodiments of the present invention, 3',3"-bisacetyl digoxin cyclobutane (bisAcDcB), from an exemplary compound according to some embodiments of the present invention, DcB, is illustrated in Scheme 2 below.

69

70

Scheme 2



55

In this example, the bisacetyl derivative (IUPAC name (2R,3R,4S,6R)-3-(((2S,4S,5R,6R)-4-acetoxy-5-((4-cyclobutyl-2-methyl-1,4-oxazepan-7-yl)oxy)-6-methyltetrahydro-2H-pyran-2-yl)oxy)-6-(((3S,5R,8R,9S,10S,12R,13S,14S,17R)-12,14-dihydroxy-10,13-dimethyl-17-(5-oxo-2,5-dihydrofuran-3-yl)hexadecahydro-1H-cyclopenta[a]phenanthren-3-yl)oxy)-2-methyltetrahydro-2H-pyran-4-yl acetate) was afforded by stirring for 6 hours at room temperature a solution of DcB (10 mg, 12.3 μ mol, MW=817) and 4-dimethylaminopyridine (DMAP) catalyst (0.3 mg, 2.4 μ mol, MW=122) in 2 ml toluene, 2 ml pyridine and 0.7 ml acetic anhydride.

60

65


Thereafter, the resulting mixture, rich in the bis-acetylated derivative, was extracted with water and dichloromethane, and subjected to final wash with HCl-acidified water pH of about 4. The resulting residue was dried over MgSO₄, and purified in normal phase HPLC, using a silica gel column.

The synthesis of an exemplary prodrug compound, according to some embodiments of the present invention, 12,3',3''-trisacetyl digoxin cyclobutane (trisAcDcB), from an exemplary compound according to some embodiments of the present invention, DcB, is illustrated in Scheme 3 below.

71

72

Scheme 3



In this example, the trisacetyl derivative (IUPAC name (2R,3R,4S,6R)-6-(((3S,5R,8R,9S,10S,12R,13S,14S,17R)-12-acetoxy-14-hydroxy-10,13-dimethyl-17-(5-oxo-2,5-dihydrofuran-3-yl)hexadecahydro-1H-cyclopenta[a]phenanthren-3-yl)oxy)-3-(((2S,4S,5R,6R)-4-acetoxy-5-((4-cyclobutyl-2-methyl-1,4-oxazepam-7-yl)oxy)-6-methyltetrahydro-2H-pyran-2-yl)oxy)-2-methyltetrahydro-2H-pyran-4-yl acetate) was afforded by stirring for 24 hours at room temperature a solution of DcB3 (10 mg, 12.3 μ mol, MW=817) and 4-dimethylaminopyridine (DMAP) catalyst (0.3 mg, 2.4 μ mol, MW=122) in 2 ml toluene, 2 ml pyridine and 0.7 ml acetic anhydride.

Thereafter, the resulting mixture, rich in the tris-acetylated derivative, was extracted with water and dichloromethane, and subjected to final wash with HCl-acidified water pH of about 4. The resulting residue was dried over MgSO₄, and purified in normal phase HPLC using a silica gel column.

Calculated octanol-water partition coefficient (ClogP) for DcB is 3.43, for a mono-acetyl derivative of DcB is 3.93, for the bisAcDcB derivative is 4.37, and for the trisAcDcB derivative is 4.85 (CLogP software ALOGPS 2.1 [Tetko, I. V. et al., *J. Chem. Inf. Comput. Sci.*, 2002, 42, 1136-45])

In Vitro Inhibition Activity:

The results of the inhibition assay of purified human Na,K-ATPase (α 1 β 1FXVD1 and α 2 β 1FXVD1) by DcB, bisAcDcB and trisAcDcB are presented as K_i values in Table 5 below.

TABLE 5

Compound	K _i of α 1 β 1FXVD1 [nM]	K _i of α 2 β 1FXVD1 [nM]
DcB	135 \pm 11	8 \pm 1.25
BisAcDcB	1180 \pm 280	395 \pm 50
TrisAcDcB	13200 \pm 6500	2500 \pm 490

As can be seen in Table 5, the K_i values of α 1 β 1 by the exemplary bisAcDcB prodrug derivative and the exemplary trisAcDcB prodrug derivative are about 9-fold and 100-fold higher than that observed for the corresponding exemplary compound DcB, respectively, while the K_i values of α 2 β 1

by bisAcDcB and trisAcDcB are about 50-fold and 300-fold higher than that observed for the corresponding DcB, respectively.

In Vivo Activity—Lowering of basal IOP:

One 30 μ l drop of trisAcDcB or vehicle was applied to the right or left eye of two rabbits, respectively, at 0.05, 0.1, and 0.2 mM concentrations. The IOP was then measured every 1-2 hours over 12 hours.

FIG. 4 presents comparative plots of IOP as a function of time, demonstrating the capacity of the trisAcDcB prodrug of the α 2-inhibitor compound DcB, according to some embodiments of the present invention, to lower IOP below basal levels compared to a buffer control when administered topically to one eye of a rabbit, while the other eye received PBS as a control.

As can be seen in FIG. 4, no effect was observed at the first four hours, and after that the IOP in the drug-treated samples dropped from 15-16 mmHg to a minimum of 12 mmHg within 5-6 hours, before returning to the control level after 12 hours. In addition, the maximal effect of the trisAcDcB on TOP was essentially equal to that produced by the parent DcB (see, for example, FIG. 2D). The comparison with the parent DcB compound shows that significantly lower doses of the prodrug are required to produce the maximal effect.

Thus, it can be concluded that acetyl prodrugs of the compounds presented herein exhibit low affinity and weak inhibition of α 1 β 1 isoform, which is the major isoform in

the cornea, making it unlikely that topical application of the prodrug in human eye would cause corneal swelling or other toxic adverse effects. The lag of four hours prior to a detectable effect of the prodrug suggests that this time is required for the intra-ocular esterases to hydrolyze the acetyl groups and regenerate DcB within the to anterior chamber of the eye. Moreover, significantly lower doses of prodrug are required to achieve a comparable effect of the parent compound. This finding is consistent with a higher permeability through the cornea compared to DcB (ClogP 3.43) due to a higher lipophilicity of the esters (triAcDcB ClogP 4.85).

Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims.

All publications, patents and patent applications mentioned in this specification are herein incorporated in their entirety by reference into the specification, to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated herein by reference. In addition, citation or identification of any reference in this application shall not be construed as an admission that such reference is available as prior art to the present invention. To the extent that section headings are used, they should not be construed as necessarily limiting.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 19

<210> SEQ ID NO 1

<211> LENGTH: 1023

<212> TYPE: PRT

<213> ORGANISM: homo sapiens

<400> SEQUENCE: 1

```

Met Gly Lys Gly Val Gly Arg Asp Lys Tyr Glu Pro Ala Ala Val Ser
 1           5           10           15

Glu Gln Gly Asp Lys Lys Gly Lys Lys Gly Lys Lys Asp Arg Asp Met
 20           25           30

Asp Glu Leu Lys Lys Glu Val Ser Met Asp Asp His Lys Leu Ser Leu
 35           40           45

Asp Glu Leu His Arg Lys Tyr Gly Thr Asp Leu Ser Arg Gly Leu Thr
 50           55           60

Ser Ala Arg Ala Ala Glu Ile Leu Ala Arg Asp Gly Pro Asn Ala Leu
 65           70           75           80

Thr Pro Pro Pro Thr Thr Pro Glu Trp Ile Lys Phe Cys Arg Gln Leu
 85           90           95

Phe Gly Gly Phe Ser Met Leu Leu Trp Ile Gly Ala Ile Leu Cys Phe
 100          105          110

Leu Ala Tyr Ser Ile Gln Ala Ala Thr Glu Glu Glu Pro Gln Asn Asp
 115          120          125

Asn Leu Tyr Leu Gly Val Val Leu Ser Ala Val Val Ile Ile Thr Gly
 130          135          140

Cys Phe Ser Tyr Tyr Gln Glu Ala Lys Ser Ser Lys Ile Met Glu Ser
 145          150          155          160

Phe Lys Asn Met Val Pro Gln Gln Ala Leu Val Ile Arg Asn Gly Glu
 165          170          175

```

-continued

Lys Met Ser Ile Asn Ala Glu Glu Val Val Val Gly Asp Leu Val Glu
 180 185 190
 Val Lys Gly Gly Asp Arg Ile Pro Ala Asp Leu Arg Ile Ile Ser Ala
 195 200 205
 Asn Gly Cys Lys Val Asp Asn Ser Ser Leu Thr Gly Glu Ser Glu Pro
 210 215 220
 Gln Thr Arg Ser Pro Asp Phe Thr Asn Glu Asn Pro Leu Glu Thr Arg
 225 230 235 240
 Asn Ile Ala Phe Phe Ser Thr Asn Cys Val Glu Gly Thr Ala Arg Gly
 245 250 255
 Ile Val Val Tyr Thr Gly Asp Arg Thr Val Met Gly Arg Ile Ala Thr
 260 265 270
 Leu Ala Ser Gly Leu Glu Gly Gly Gln Thr Pro Ile Ala Ala Glu Ile
 275 280 285
 Glu His Phe Ile His Ile Ile Thr Gly Val Ala Val Phe Leu Gly Val
 290 295 300
 Ser Phe Phe Ile Leu Ser Leu Ile Leu Glu Tyr Thr Trp Leu Glu Ala
 305 310 315 320
 Val Ile Phe Leu Ile Gly Ile Ile Val Ala Asn Val Pro Glu Gly Leu
 325 330 335
 Leu Ala Thr Val Thr Val Cys Leu Thr Leu Thr Ala Lys Arg Met Ala
 340 345 350
 Arg Lys Asn Cys Leu Val Lys Asn Leu Glu Ala Val Glu Thr Leu Gly
 355 360 365
 Ser Thr Ser Thr Ile Cys Ser Asp Lys Thr Gly Thr Leu Thr Gln Asn
 370 375 380
 Arg Met Thr Val Ala His Met Trp Phe Asp Asn Gln Ile His Glu Ala
 385 390 395 400
 Asp Thr Thr Glu Asn Gln Ser Gly Val Ser Phe Asp Lys Thr Ser Ala
 405 410 415
 Thr Trp Leu Ala Leu Ser Arg Ile Ala Gly Leu Cys Asn Arg Ala Val
 420 425 430
 Phe Gln Ala Asn Gln Glu Asn Leu Pro Ile Leu Lys Arg Ala Val Ala
 435 440 445
 Gly Asp Ala Ser Glu Ser Ala Leu Leu Lys Cys Ile Glu Leu Cys Cys
 450 455 460
 Gly Ser Val Lys Glu Met Arg Glu Arg Tyr Ala Lys Ile Val Glu Ile
 465 470 475 480
 Pro Phe Asn Ser Thr Asn Lys Tyr Gln Leu Ser Ile His Lys Asn Pro
 485 490 495
 Asn Thr Ser Glu Pro Gln His Leu Leu Val Met Lys Gly Ala Pro Glu
 500 505 510
 Arg Ile Leu Asp Arg Cys Ser Ser Ile Leu Leu His Gly Lys Glu Gln
 515 520 525
 Pro Leu Asp Glu Glu Leu Lys Asp Ala Phe Gln Asn Ala Tyr Leu Glu
 530 535 540
 Leu Gly Gly Leu Gly Glu Arg Val Leu Gly Phe Cys His Leu Phe Leu
 545 550 555 560
 Pro Asp Glu Gln Phe Pro Glu Gly Phe Gln Phe Asp Thr Asp Asp Val
 565 570 575
 Asn Phe Pro Ile Asp Asn Leu Cys Phe Val Gly Leu Ile Ser Met Ile
 580 585 590

-continued

Asp Pro Pro Arg Ala Ala Val Pro Asp Ala Val Gly Lys Cys Arg Ser
 595 600 605

Ala Gly Ile Lys Val Ile Met Val Thr Gly Asp His Pro Ile Thr Ala
 610 615 620

Lys Ala Ile Ala Lys Gly Val Gly Ile Ile Ser Glu Gly Asn Glu Thr
 625 630 635 640

Val Glu Asp Ile Ala Ala Arg Leu Asn Ile Pro Val Ser Gln Val Asn
 645 650 655

Pro Arg Asp Ala Lys Ala Cys Val Val His Gly Ser Asp Leu Lys Asp
 660 665 670

Met Thr Ser Glu Gln Leu Asp Asp Ile Leu Lys Tyr His Thr Glu Ile
 675 680 685

Val Phe Ala Arg Thr Ser Pro Gln Gln Lys Leu Ile Ile Val Glu Gly
 690 695 700

Cys Gln Arg Gln Gly Ala Ile Val Ala Val Thr Gly Asp Gly Val Asn
 705 710 715 720

Asp Ser Pro Ala Leu Lys Lys Ala Asp Ile Gly Val Ala Met Gly Ile
 725 730 735

Ala Gly Ser Asp Val Ser Lys Gln Ala Ala Asp Met Ile Leu Leu Asp
 740 745 750

Asp Asn Phe Ala Ser Ile Val Thr Gly Val Glu Glu Gly Arg Leu Ile
 755 760 765

Phe Asp Asn Leu Lys Lys Ser Ile Ala Tyr Thr Leu Thr Ser Asn Ile
 770 775 780

Pro Glu Ile Thr Pro Phe Leu Ile Phe Ile Ile Ala Asn Ile Pro Leu
 785 790 795 800

Pro Leu Gly Thr Val Thr Ile Leu Cys Ile Asp Leu Gly Thr Asp Met
 805 810 815

Val Pro Ala Ile Ser Leu Ala Tyr Glu Gln Ala Glu Ser Asp Ile Met
 820 825 830

Lys Arg Gln Pro Arg Asn Pro Lys Thr Asp Lys Leu Val Asn Glu Arg
 835 840 845

Leu Ile Ser Met Ala Tyr Gly Gln Ile Gly Met Ile Gln Ala Leu Gly
 850 855 860

Gly Phe Phe Thr Tyr Phe Val Ile Leu Ala Glu Asn Gly Phe Leu Pro
 865 870 875 880

Ile His Leu Leu Gly Leu Arg Val Asp Trp Asp Asp Arg Trp Ile Asn
 885 890 895

Asp Val Glu Asp Ser Tyr Gly Gln Gln Trp Thr Tyr Glu Gln Arg Lys
 900 905 910

Ile Val Glu Phe Thr Cys His Thr Ala Phe Phe Val Ser Ile Val Val
 915 920 925

Val Gln Trp Ala Asp Leu Val Ile Cys Lys Thr Arg Arg Asn Ser Val
 930 935 940

Phe Gln Gln Gly Met Lys Asn Lys Ile Leu Ile Phe Gly Leu Phe Glu
 945 950 955 960

Glu Thr Ala Leu Ala Ala Phe Leu Ser Tyr Cys Pro Gly Met Gly Val
 965 970 975

Ala Leu Arg Met Tyr Pro Leu Lys Pro Thr Trp Trp Phe Cys Ala Phe
 980 985 990

Pro Tyr Ser Leu Leu Ile Phe Val Tyr Asp Glu Val Arg Lys Leu Ile
 995 1000 1005

Ile Arg Arg Arg Pro Gly Gly Trp Val Glu Lys Glu Thr Tyr Tyr

-continued

1010 1015 1020

<210> SEQ ID NO 2
 <211> LENGTH: 1020
 <212> TYPE: PRT
 <213> ORGANISM: homo sapiens

<400> SEQUENCE: 2

Met Gly Arg Gly Ala Gly Arg Glu Tyr Ser Pro Ala Ala Thr Thr Ala
 1 5 10 15

Glu Asn Gly Gly Gly Lys Lys Lys Gln Lys Glu Lys Glu Leu Asp Glu
 20 25 30

Leu Lys Lys Glu Val Ala Met Asp Asp His Lys Leu Ser Leu Asp Glu
 35 40 45

Leu Gly Arg Lys Tyr Gln Val Asp Leu Ser Lys Gly Leu Thr Asn Gln
 50 55 60

Arg Ala Gln Asp Val Leu Ala Arg Asp Gly Pro Asn Ala Leu Thr Pro
 65 70 75 80

Pro Pro Thr Thr Pro Glu Trp Val Lys Phe Cys Arg Gln Leu Phe Gly
 85 90 95

Gly Phe Ser Ile Leu Leu Trp Ile Gly Ala Ile Leu Cys Phe Leu Ala
 100 105 110

Tyr Gly Ile Gln Ala Ala Met Glu Asp Glu Pro Ser Asn Asp Asn Leu
 115 120 125

Tyr Leu Gly Val Val Leu Ala Ala Val Val Ile Val Thr Gly Cys Phe
 130 135 140

Ser Tyr Tyr Gln Glu Ala Lys Ser Ser Lys Ile Met Asp Ser Phe Lys
 145 150 155 160

Asn Met Val Pro Gln Gln Ala Leu Val Ile Arg Glu Gly Glu Lys Met
 165 170 175

Gln Ile Asn Ala Glu Glu Val Val Val Gly Asp Leu Val Glu Val Lys
 180 185 190

Gly Gly Asp Arg Val Pro Ala Asp Leu Arg Ile Ile Ser Ser His Gly
 195 200 205

Cys Lys Val Asp Asn Ser Ser Leu Thr Gly Glu Ser Glu Pro Gln Thr
 210 215 220

Arg Ser Pro Glu Phe Thr His Glu Asn Pro Leu Glu Thr Arg Asn Ile
 225 230 235 240

Cys Phe Phe Ser Thr Asn Cys Val Glu Gly Thr Ala Arg Gly Ile Val
 245 250 255

Ile Ala Thr Gly Asp Arg Thr Val Met Gly Arg Ile Ala Thr Leu Ala
 260 265 270

Ser Gly Leu Glu Val Gly Arg Thr Pro Ile Ala Met Glu Ile Glu His
 275 280 285

Phe Ile Gln Leu Ile Thr Gly Val Ala Val Phe Leu Gly Val Ser Phe
 290 295 300

Phe Val Leu Ser Leu Ile Leu Gly Tyr Ser Trp Leu Glu Ala Val Ile
 305 310 315 320

Phe Leu Ile Gly Ile Ile Val Ala Asn Val Pro Glu Gly Leu Leu Ala
 325 330 335

Thr Val Thr Val Cys Leu Thr Leu Thr Ala Lys Arg Met Ala Arg Lys
 340 345 350

Asn Cys Leu Val Lys Asn Leu Glu Ala Val Glu Thr Leu Gly Ser Thr
 355 360 365

-continued

Ser Thr Ile Cys Ser Asp Lys Thr Gly Thr Leu Thr Gln Asn Arg Met
 370 375 380
 Thr Val Ala His Met Trp Phe Asp Asn Gln Ile His Glu Ala Asp Thr
 385 390 395 400
 Thr Glu Asp Gln Ser Gly Ala Thr Phe Asp Lys Arg Ser Pro Thr Trp
 405 410 415
 Thr Ala Leu Ser Arg Ile Ala Gly Leu Cys Asn Arg Ala Val Phe Lys
 420 425 430
 Ala Gly Gln Glu Asn Ile Ser Val Ser Lys Arg Asp Thr Ala Gly Asp
 435 440 445
 Ala Ser Glu Ser Ala Leu Leu Lys Cys Ile Glu Leu Ser Cys Gly Ser
 450 455 460
 Val Arg Lys Met Arg Asp Arg Asn Pro Lys Val Ala Glu Ile Pro Phe
 465 470 475 480
 Asn Ser Thr Asn Lys Tyr Gln Leu Ser Ile His Glu Arg Glu Asp Ser
 485 490 495
 Pro Gln Ser His Val Leu Val Met Lys Gly Ala Pro Glu Arg Ile Leu
 500 505 510
 Asp Arg Cys Ser Thr Ile Leu Val Gln Gly Lys Glu Ile Pro Leu Asp
 515 520 525
 Lys Glu Met Gln Asp Ala Phe Gln Asn Ala Tyr Met Glu Leu Gly Gly
 530 535 540
 Leu Gly Glu Arg Val Leu Gly Phe Cys Gln Leu Asn Leu Pro Ser Gly
 545 550 555 560
 Lys Phe Pro Arg Gly Phe Lys Phe Asp Thr Asp Glu Leu Asn Phe Pro
 565 570 575
 Thr Glu Lys Leu Cys Phe Val Gly Leu Met Ser Met Ile Asp Pro Pro
 580 585 590
 Arg Ala Ala Val Pro Asp Ala Val Gly Lys Cys Arg Ser Ala Gly Ile
 595 600 605
 Lys Val Ile Met Val Thr Gly Asp His Pro Ile Thr Ala Lys Ala Ile
 610 615 620
 Ala Lys Gly Val Gly Ile Ile Ser Glu Gly Asn Glu Thr Val Glu Asp
 625 630 635 640
 Ile Ala Ala Arg Leu Asn Ile Pro Met Ser Gln Val Asn Pro Arg Glu
 645 650 655
 Ala Lys Ala Cys Val Val His Gly Ser Asp Leu Lys Asp Met Thr Ser
 660 665 670
 Glu Gln Leu Asp Glu Ile Leu Lys Asn His Thr Glu Ile Val Phe Ala
 675 680 685
 Arg Thr Ser Pro Gln Gln Lys Leu Ile Ile Val Glu Gly Cys Gln Arg
 690 695 700
 Gln Gly Ala Ile Val Ala Val Thr Gly Asp Gly Val Asn Asp Ser Pro
 705 710 715 720
 Ala Leu Lys Lys Ala Asp Ile Gly Ile Ala Met Gly Ile Ser Gly Ser
 725 730 735
 Asp Val Ser Lys Gln Ala Ala Asp Met Ile Leu Leu Asp Asp Asn Phe
 740 745 750
 Ala Ser Ile Val Thr Gly Val Glu Glu Gly Arg Leu Ile Phe Asp Asn
 755 760 765
 Leu Lys Lys Ser Ile Ala Tyr Thr Leu Thr Ser Asn Ile Pro Glu Ile
 770 775 780
 Thr Pro Phe Leu Leu Phe Ile Ile Ala Asn Ile Pro Leu Pro Leu Gly

-continued

Ser Lys Ile Met Glu Ser Phe Lys Asn Met Val Pro Gln Gln Ala Leu
 145 150 155 160

Val Ile Arg Glu Gly Glu Lys Met Gln Val Asn Ala Glu Glu Val Val
 165 170 175

Val Gly Asp Leu Val Glu Ile Lys Gly Gly Asp Arg Val Pro Ala Asp
 180 185 190

Leu Arg Ile Ile Ser Ala His Gly Cys Lys Val Asp Asn Ser Ser Leu
 195 200 205

Thr Gly Glu Ser Glu Pro Gln Thr Arg Ser Pro Asp Cys Thr His Asp
 210 215 220

Asn Pro Leu Glu Thr Arg Asn Ile Thr Phe Phe Ser Thr Asn Cys Val
 225 230 235 240

Glu Gly Thr Ala Arg Gly Val Val Val Ala Thr Gly Asp Arg Thr Val
 245 250 255

Met Gly Arg Ile Ala Thr Leu Ala Ser Gly Leu Glu Val Gly Lys Thr
 260 265 270

Pro Ile Ala Ile Glu Ile Glu His Phe Ile Gln Leu Ile Thr Gly Val
 275 280 285

Ala Val Phe Leu Gly Val Ser Phe Phe Ile Leu Ser Leu Ile Leu Gly
 290 295 300

Tyr Thr Trp Leu Glu Ala Val Ile Phe Leu Ile Gly Ile Ile Val Ala
 305 310 315 320

Asn Val Pro Glu Gly Leu Leu Ala Thr Val Thr Val Cys Leu Thr Leu
 325 330 335

Thr Ala Lys Arg Met Ala Arg Lys Asn Cys Leu Val Lys Asn Leu Glu
 340 345 350

Ala Val Glu Thr Leu Gly Ser Thr Ser Thr Ile Cys Ser Asp Lys Thr
 355 360 365

Gly Thr Leu Thr Gln Asn Arg Met Thr Val Ala His Met Trp Phe Asp
 370 375 380

Asn Gln Ile His Glu Ala Asp Thr Thr Glu Asp Gln Ser Gly Thr Ser
 385 390 395 400

Phe Asp Lys Ser Ser His Thr Trp Val Ala Leu Ser His Ile Ala Gly
 405 410 415

Leu Cys Asn Arg Ala Val Phe Lys Gly Gly Gln Asp Asn Ile Pro Val
 420 425 430

Leu Lys Arg Asp Val Ala Gly Asp Ala Ser Glu Ser Ala Leu Leu Lys
 435 440 445

Cys Ile Glu Leu Ser Ser Gly Ser Val Lys Leu Met Arg Glu Arg Asn
 450 455 460

Lys Lys Val Ala Glu Ile Pro Phe Asn Ser Thr Asn Lys Tyr Gln Leu
 465 470 475 480

Ser Ile His Glu Thr Glu Asp Pro Asn Asp Asn Arg Tyr Leu Leu Val
 485 490 495

Met Lys Gly Ala Pro Glu Arg Ile Leu Asp Arg Cys Ser Thr Ile Leu
 500 505 510

Leu Gln Gly Lys Glu Gln Pro Leu Asp Glu Glu Met Lys Glu Ala Phe
 515 520 525

Gln Asn Ala Tyr Leu Glu Leu Gly Gly Leu Gly Glu Arg Val Leu Gly
 530 535 540

Phe Cys His Tyr Tyr Leu Pro Glu Glu Gln Phe Pro Lys Gly Phe Ala
 545 550 555 560

Phe Asp Cys Asp Asp Val Asn Phe Thr Thr Asp Asn Leu Cys Phe Val

-continued

565				570				575							
Gly	Leu	Met	Ser	Met	Ile	Asp	Pro	Pro	Arg	Ala	Ala	Val	Pro	Asp	Ala
			580							585				590	
Val	Gly	Lys	Cys	Arg	Ser	Ala	Gly	Ile	Lys	Val	Ile	Met	Val	Thr	Gly
		595					600						605		
Asp	His	Pro	Ile	Thr	Ala	Lys	Ala	Ile	Ala	Lys	Gly	Val	Gly	Ile	Ile
	610					615					620				
Ser	Glu	Gly	Asn	Glu	Thr	Val	Glu	Asp	Ile	Ala	Ala	Arg	Leu	Asn	Ile
	625				630					635				640	
Pro	Val	Ser	Gln	Val	Asn	Pro	Arg	Asp	Ala	Lys	Ala	Cys	Val	Ile	His
			645							650				655	
Gly	Thr	Asp	Leu	Lys	Asp	Phe	Thr	Ser	Glu	Gln	Ile	Asp	Glu	Ile	Leu
		660								665				670	
Gln	Asn	His	Thr	Glu	Ile	Val	Phe	Ala	Arg	Thr	Ser	Pro	Gln	Gln	Lys
		675					680						685		
Leu	Ile	Ile	Val	Glu	Gly	Cys	Gln	Arg	Gln	Gly	Ala	Ile	Val	Ala	Val
	690					695					700				
Thr	Gly	Asp	Gly	Val	Asn	Asp	Ser	Pro	Ala	Leu	Lys	Lys	Ala	Asp	Ile
	705				710					715				720	
Gly	Val	Ala	Met	Gly	Ile	Ala	Gly	Ser	Asp	Val	Ser	Lys	Gln	Ala	Ala
			725							730				735	
Asp	Met	Ile	Leu	Leu	Asp	Asp	Asn	Phe	Ala	Ser	Ile	Val	Thr	Gly	Val
		740								745				750	
Glu	Glu	Gly	Arg	Leu	Ile	Phe	Asp	Asn	Leu	Lys	Lys	Ser	Ile	Ala	Tyr
		755					760							765	
Thr	Leu	Thr	Ser	Asn	Ile	Pro	Glu	Ile	Thr	Pro	Phe	Leu	Leu	Phe	Ile
	770					775					780				
Met	Ala	Asn	Ile	Pro	Leu	Pro	Leu	Gly	Thr	Ile	Thr	Ile	Leu	Cys	Ile
	785				790					795				800	
Asp	Leu	Gly	Thr	Asp	Met	Val	Pro	Ala	Ile	Ser	Leu	Ala	Tyr	Glu	Ala
			805							810				815	
Ala	Glu	Ser	Asp	Ile	Met	Lys	Arg	Gln	Pro	Arg	Asn	Pro	Arg	Thr	Asp
			820							825				830	
Lys	Leu	Val	Asn	Glu	Arg	Leu	Ile	Ser	Met	Ala	Tyr	Gly	Gln	Ile	Gly
		835					840							845	
Met	Ile	Gln	Ala	Leu	Gly	Gly	Phe	Phe	Ser	Tyr	Phe	Val	Ile	Leu	Ala
	850					855					860				
Glu	Asn	Gly	Phe	Leu	Pro	Gly	Asn	Leu	Val	Gly	Ile	Arg	Leu	Asn	Trp
	865				870					875				880	
Asp	Asp	Arg	Thr	Val	Asn	Asp	Leu	Glu	Asp	Ser	Tyr	Gly	Gln	Gln	Trp
			885							890				895	
Thr	Tyr	Glu	Gln	Arg	Lys	Val	Val	Glu	Phe	Thr	Cys	His	Thr	Ala	Phe
		900								905				910	
Phe	Val	Ser	Ile	Val	Val	Val	Gln	Trp	Ala	Asp	Leu	Ile	Ile	Cys	Lys
		915					920							925	
Thr	Arg	Arg	Asn	Ser	Val	Phe	Gln	Gln	Gly	Met	Lys	Asn	Lys	Ile	Leu
	930					935					940				
Ile	Phe	Gly	Leu	Phe	Glu	Glu	Thr	Ala	Leu	Ala	Ala	Phe	Leu	Ser	Tyr
	945					950				955				960	
Cys	Pro	Gly	Met	Asp	Val	Ala	Leu	Arg	Met	Tyr	Pro	Leu	Lys	Pro	Ser
			965							970				975	
Trp	Trp	Phe	Cys	Ala	Phe	Pro	Tyr	Ser	Phe	Leu	Ile	Phe	Val	Tyr	Asp
			980							985				990	

-continued

Glu Ile Arg Lys Leu Ile Leu Arg Arg Asn Pro Gly Gly Trp Val Glu
 995 1000 1005

Lys Glu Thr Tyr Tyr
 1010

<210> SEQ ID NO 4
 <211> LENGTH: 1029
 <212> TYPE: PRT
 <213> ORGANISM: homo sapiens

<400> SEQUENCE: 4

Met Gly Leu Trp Gly Lys Lys Gly Thr Val Ala Pro His Asp Gln Ser
 1 5 10 15
 Pro Arg Arg Arg Pro Lys Lys Gly Leu Ile Lys Lys Lys Met Val Lys
 20 25 30
 Arg Glu Lys Gln Lys Arg Asn Met Glu Glu Leu Lys Lys Glu Val Val
 35 40 45
 Met Asp Asp His Lys Leu Thr Leu Glu Glu Leu Ser Thr Lys Tyr Ser
 50 55 60
 Val Asp Leu Thr Lys Gly His Ser His Gln Arg Ala Lys Glu Ile Leu
 65 70 75 80
 Thr Arg Gly Gly Pro Asn Thr Val Thr Pro Pro Pro Thr Thr Pro Glu
 85 90 95
 Trp Val Lys Phe Cys Lys Gln Leu Phe Gly Gly Phe Ser Leu Leu Leu
 100 105 110
 Trp Thr Gly Ala Ile Leu Cys Phe Val Ala Tyr Ser Ile Gln Ile Tyr
 115 120 125
 Phe Asn Glu Glu Pro Thr Lys Asp Asn Leu Tyr Leu Ser Ile Val Leu
 130 135 140
 Ser Val Val Val Ile Val Thr Gly Cys Phe Ser Tyr Tyr Gln Glu Ala
 145 150 155 160
 Lys Ser Ser Lys Ile Met Glu Ser Phe Lys Asn Met Val Pro Gln Gln
 165 170 175
 Ala Leu Val Ile Arg Gly Gly Glu Lys Met Gln Ile Asn Val Gln Glu
 180 185 190
 Val Val Leu Gly Asp Leu Val Glu Ile Lys Gly Gly Asp Arg Val Pro
 195 200 205
 Ala Asp Leu Arg Leu Ile Ser Ala Gln Gly Cys Lys Val Asp Asn Ser
 210 215 220
 Ser Leu Thr Gly Glu Ser Glu Pro Gln Ser Arg Ser Pro Asp Phe Thr
 225 230 235 240
 His Glu Asn Pro Leu Glu Thr Arg Asn Ile Cys Phe Phe Ser Thr Asn
 245 250 255
 Cys Val Glu Gly Thr Ala Arg Gly Ile Val Ile Ala Thr Gly Asp Ser
 260 265 270
 Thr Val Met Gly Arg Ile Ala Ser Leu Thr Ser Gly Leu Ala Val Gly
 275 280 285
 Gln Thr Pro Ile Ala Ala Glu Ile Glu His Phe Ile His Leu Ile Thr
 290 295 300
 Val Val Ala Val Phe Leu Gly Val Thr Phe Phe Ala Leu Ser Leu Leu
 305 310 315 320
 Leu Gly Tyr Gly Trp Leu Glu Ala Ile Ile Phe Leu Ile Gly Ile Ile
 325 330 335
 Val Ala Asn Val Pro Glu Gly Leu Leu Ala Thr Val Thr Val Cys Leu

-continued

Glu Glu Gly Arg Leu Ile Phe Asp Asn Leu Lys Lys Ser Ile Met Tyr
 770 775 780
 Thr Leu Thr Ser Asn Ile Pro Glu Ile Thr Pro Phe Leu Met Phe Ile
 785 790 795 800
 Ile Leu Gly Ile Pro Leu Pro Leu Gly Thr Ile Thr Ile Leu Cys Ile
 805 810 815
 Asp Leu Gly Thr Asp Met Val Pro Ala Ile Ser Leu Ala Tyr Glu Ser
 820 825 830
 Ala Glu Ser Asp Ile Met Lys Arg Leu Pro Arg Asn Pro Lys Thr Asp
 835 840 845
 Asn Leu Val Asn His Arg Leu Ile Gly Met Ala Tyr Gly Gln Ile Gly
 850 855 860
 Met Ile Gln Ala Leu Ala Gly Phe Phe Thr Tyr Phe Val Ile Leu Ala
 865 870 875 880
 Glu Asn Gly Phe Arg Pro Val Asp Leu Leu Gly Ile Arg Leu His Trp
 885 890 895
 Glu Asp Lys Tyr Leu Asn Asp Leu Glu Asp Ser Tyr Gly Gln Gln Trp
 900 905 910
 Thr Tyr Glu Gln Arg Lys Val Val Glu Phe Thr Cys Gln Thr Ala Phe
 915 920 925
 Phe Val Thr Ile Val Val Val Gln Trp Ala Asp Leu Ile Ile Ser Lys
 930 935 940
 Thr Arg Arg Asn Ser Leu Phe Gln Gln Gly Met Arg Asn Lys Val Leu
 945 950 955 960
 Ile Phe Gly Ile Leu Glu Glu Thr Leu Leu Ala Ala Phe Leu Ser Tyr
 965 970 975
 Thr Pro Gly Met Asp Val Ala Leu Arg Met Tyr Pro Leu Lys Ile Thr
 980 985 990
 Trp Trp Leu Cys Ala Ile Pro Tyr Ser Ile Leu Ile Phe Val Tyr Asp
 995 1000 1005
 Glu Ile Arg Lys Leu Leu Ile Arg Gln His Pro Asp Gly Trp Val
 1010 1015 1020
 Glu Arg Glu Thr Tyr Tyr
 1025

<210> SEQ ID NO 5
 <211> LENGTH: 303
 <212> TYPE: PRT
 <213> ORGANISM: homo sapiens

<400> SEQUENCE: 5

Met Ala Arg Gly Lys Ala Lys Glu Glu Gly Ser Trp Lys Lys Phe Ile
 1 5 10 15
 Trp Asn Ser Glu Lys Lys Glu Phe Leu Gly Arg Thr Gly Gly Ser Trp
 20 25 30
 Phe Lys Ile Leu Leu Phe Tyr Val Ile Phe Tyr Gly Cys Leu Ala Gly
 35 40 45
 Ile Phe Ile Gly Thr Ile Gln Val Met Leu Leu Thr Ile Ser Glu Phe
 50 55 60
 Lys Pro Thr Tyr Gln Asp Arg Val Ala Pro Pro Gly Leu Thr Gln Ile
 65 70 75 80
 Pro Gln Ile Gln Lys Thr Glu Ile Ser Phe Arg Pro Asn Asp Pro Lys
 85 90 95
 Ser Tyr Glu Ala Tyr Val Leu Asn Ile Val Arg Phe Leu Glu Lys Tyr

-continued

100					105					110					
Lys	Asp	Ser	Ala	Gln	Arg	Asp	Asp	Met	Ile	Phe	Glu	Asp	Cys	Gly	Asp
	115						120					125			
Val	Pro	Ser	Glu	Pro	Lys	Glu	Arg	Gly	Asp	Phe	Asn	His	Glu	Arg	Gly
	130					135					140				
Glu	Arg	Lys	Val	Cys	Arg	Phe	Lys	Leu	Glu	Trp	Leu	Gly	Asn	Cys	Ser
145				150					155						160
Gly	Leu	Asn	Asp	Glu	Thr	Tyr	Gly	Tyr	Lys	Glu	Gly	Lys	Pro	Cys	Ile
			165						170					175	
Ile	Ile	Lys	Leu	Asn	Arg	Val	Leu	Gly	Phe	Lys	Pro	Lys	Pro	Pro	Lys
		180						185					190		
Asn	Glu	Ser	Leu	Glu	Thr	Tyr	Pro	Val	Met	Lys	Tyr	Asn	Pro	Asn	Val
	195						200					205			
Leu	Pro	Val	Gln	Cys	Thr	Gly	Lys	Arg	Asp	Glu	Asp	Lys	Asp	Lys	Val
	210					215					220				
Gly	Asn	Val	Glu	Tyr	Phe	Gly	Leu	Gly	Asn	Ser	Pro	Gly	Phe	Pro	Leu
225					230					235					240
Gln	Tyr	Tyr	Pro	Tyr	Tyr	Gly	Lys	Leu	Leu	Gln	Pro	Lys	Tyr	Leu	Gln
			245						250					255	
Pro	Leu	Leu	Ala	Val	Gln	Phe	Thr	Asn	Leu	Thr	Met	Asp	Thr	Glu	Ile
		260						265					270		
Arg	Ile	Glu	Cys	Lys	Ala	Tyr	Gly	Glu	Asn	Ile	Gly	Tyr	Ser	Glu	Lys
		275					280					285			
Asp	Arg	Phe	Gln	Gly	Arg	Phe	Asp	Val	Lys	Ile	Glu	Val	Lys	Ser	
	290					295					300				

<210> SEQ ID NO 6

<211> LENGTH: 319

<212> TYPE: PRT

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Amino Acid sequence encoding His10-ATP1B1

<400> SEQUENCE: 6

Met	Ala	Arg	Ser	His	Pro	Arg									
1				5						10				15	
Arg	Ser	Arg	Gly	Lys	Ala	Lys	Glu	Glu	Gly	Ser	Trp	Lys	Lys	Phe	Ile
			20						25				30		
Trp	Asn	Ser	Glu	Lys	Lys	Glu	Phe	Leu	Gly	Arg	Thr	Gly	Gly	Ser	Trp
		35					40					45			
Phe	Lys	Ile	Leu	Leu	Phe	Tyr	Val	Ile	Phe	Tyr	Gly	Cys	Leu	Ala	Gly
	50					55					60				
Ile	Phe	Ile	Gly	Thr	Ile	Gln	Val	Met	Leu	Leu	Thr	Ile	Ser	Glu	Phe
65					70					75					80
Lys	Pro	Thr	Tyr	Gln	Asp	Arg	Val	Ala	Pro	Pro	Gly	Leu	Thr	Gln	Ile
				85					90					95	
Pro	Gln	Ile	Gln	Lys	Thr	Glu	Ile	Ser	Phe	Arg	Pro	Asn	Asp	Pro	Lys
		100						105					110		
Ser	Tyr	Glu	Ala	Tyr	Val	Leu	Asn	Ile	Val	Arg	Phe	Leu	Glu	Lys	Tyr
	115						120					125			
Lys	Asp	Ser	Ala	Gln	Arg	Asp	Asp	Met	Ile	Phe	Glu	Asp	Cys	Gly	Asp
	130					135					140				
Val	Pro	Ser	Glu	Pro	Lys	Glu	Arg	Gly	Asp	Phe	Asn	His	Glu	Arg	Gly
145					150					155					160
Glu	Arg	Lys	Val	Cys	Arg	Phe	Lys	Leu	Glu	Trp	Leu	Gly	Asn	Cys	Ser

-continued

Leu Met Tyr Phe Pro Tyr Tyr Gly Lys Lys Phe His Val Asn Tyr Thr
 225 230 235 240

Gln Pro Leu Val Ala Val Lys Phe Leu Asn Val Thr Pro Asn Val Glu
 245 250 255

Val Asn Val Glu Cys Arg Ile Asn Ala Ala Asn Ile Ala Thr Asp Asp
 260 265 270

Glu Arg Asp Lys Phe Ala Gly Arg Val Ala Phe Lys Leu Arg Ile Asn
 275 280 285

Lys Thr
 290

<210> SEQ ID NO 8
 <211> LENGTH: 306
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Amino Acid sequence encoding His10-ATP1B2

<400> SEQUENCE: 8

Met Ala Arg Ser His His His His His His His His His Pro Arg
 1 5 10 15

Arg Ser Arg Gly Lys Glu Lys Lys Ser Cys Gly Gln Val Val Glu Glu
 20 25 30

Trp Lys Glu Phe Val Trp Asn Pro Arg Thr His Gln Phe Met Gly Arg
 35 40 45

Thr Gly Thr Ser Trp Ala Phe Ile Leu Leu Phe Tyr Leu Val Phe Tyr
 50 55 60

Gly Phe Leu Thr Ala Met Phe Thr Leu Thr Met Trp Val Met Leu Gln
 65 70 75 80

Thr Val Ser Asp His Thr Pro Lys Tyr Gln Asp Arg Leu Ala Thr Pro
 85 90 95

Gly Leu Met Ile Arg Pro Lys Thr Glu Asn Leu Asp Val Ile Val Asn
 100 105 110

Val Ser Asp Thr Glu Ser Trp Asp Gln His Val Gln Lys Leu Asn Lys
 115 120 125

Phe Leu Glu Pro Tyr Asn Asp Ser Ile Gln Ala Gln Lys Asn Asp Val
 130 135 140

Cys Arg Pro Gly Arg Tyr Tyr Glu Gln Pro Asp Asn Gly Val Leu Asn
 145 150 155 160

Tyr Pro Lys Arg Ala Cys Gln Phe Asn Arg Thr Gln Leu Gly Asn Cys
 165 170 175

Ser Gly Ile Gly Asp Ser Thr His Tyr Gly Tyr Ser Thr Gly Gln Pro
 180 185 190

Cys Val Phe Ile Lys Met Asn Arg Val Ile Asn Phe Tyr Ala Gly Ala
 195 200 205

Asn Gln Ser Met Asn Val Thr Cys Ala Gly Lys Arg Asp Glu Asp Ala
 210 215 220

Glu Asn Leu Gly Asn Phe Val Met Phe Pro Ala Asn Gly Asn Ile Asp
 225 230 235 240

Leu Met Tyr Phe Pro Tyr Tyr Gly Lys Lys Phe His Val Asn Tyr Thr
 245 250 255

Gln Pro Leu Val Ala Val Lys Phe Leu Asn Val Thr Pro Asn Val Glu
 260 265 270

Val Asn Val Glu Cys Arg Ile Asn Ala Ala Asn Ile Ala Thr Asp Asp
 275 280 285

-continued

Glu Arg Asp Lys Phe Ala Gly Arg Val Ala Phe Lys Leu Arg Ile Asn
 290 295 300

Lys Thr
 305

<210> SEQ ID NO 9
 <211> LENGTH: 279
 <212> TYPE: PRT
 <213> ORGANISM: homo sapiens

<400> SEQUENCE: 9

Met Thr Lys Asn Glu Lys Lys Ser Leu Asn Gln Ser Leu Ala Glu Trp
 1 5 10 15
 Lys Leu Phe Ile Tyr Asn Pro Thr Thr Gly Glu Phe Leu Gly Arg Thr
 20 25 30
 Ala Lys Ser Trp Gly Leu Ile Leu Leu Phe Tyr Leu Val Phe Tyr Gly
 35 40 45
 Phe Leu Ala Ala Leu Phe Ser Phe Thr Met Trp Val Met Leu Gln Thr
 50 55 60
 Leu Asn Asp Glu Val Pro Lys Tyr Arg Asp Gln Ile Pro Ser Pro Gly
 65 70 75 80
 Leu Met Val Phe Pro Lys Pro Val Thr Ala Leu Glu Tyr Thr Phe Ser
 85 90 95
 Arg Ser Asp Pro Thr Ser Tyr Ala Gly Tyr Ile Glu Asp Leu Lys Lys
 100 105 110
 Phe Leu Lys Pro Tyr Thr Leu Glu Glu Gln Lys Asn Leu Thr Val Cys
 115 120 125
 Pro Asp Gly Ala Leu Phe Glu Gln Lys Gly Pro Val Tyr Val Ala Cys
 130 135 140
 Gln Phe Pro Ile Ser Leu Leu Gln Ala Cys Ser Gly Met Asn Asp Pro
 145 150 155 160
 Asp Phe Gly Tyr Ser Gln Gly Asn Pro Cys Ile Leu Val Lys Met Asn
 165 170 175
 Arg Ile Ile Gly Leu Lys Pro Glu Gly Val Pro Arg Ile Asp Cys Val
 180 185 190
 Ser Lys Asn Glu Asp Ile Pro Asn Val Ala Val Tyr Pro His Asn Gly
 195 200 205
 Met Ile Asp Leu Lys Tyr Phe Pro Tyr Tyr Gly Lys Lys Leu His Val
 210 215 220
 Gly Tyr Leu Gln Pro Leu Val Ala Val Gln Val Ser Phe Ala Pro Asn
 225 230 235 240
 Asn Thr Gly Lys Glu Val Thr Val Glu Cys Lys Ile Asp Gly Ser Ala
 245 250 255
 Asn Leu Lys Ser Gln Asp Asp Arg Asp Lys Phe Leu Gly Arg Val Met
 260 265 270
 Phe Lys Ile Thr Ala Arg Ala
 275

<210> SEQ ID NO 10
 <211> LENGTH: 296
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Amino Acid sequence encoding His10-ATP1B3

<400> SEQUENCE: 10

Met Ala Arg Ser His Pro Arg

-continued

1	5	10	15
Arg Ser Arg Gly Asn Glu Lys Lys Ser Leu Asn Gln Ser Leu Ala Glu	20	25	30
Trp Lys Leu Phe Ile Tyr Asn Pro Thr Thr Gly Glu Phe Leu Gly Arg	35	40	45
Thr Ala Lys Ser Trp Gly Leu Ile Leu Leu Phe Tyr Leu Val Phe Tyr	50	55	60
Gly Phe Leu Ala Ala Leu Phe Ser Phe Thr Met Trp Val Met Leu Gln	65	70	80
Thr Leu Asn Asp Glu Val Pro Lys Tyr Arg Asp Gln Ile Pro Ser Pro	85	90	95
Gly Leu Met Val Phe Pro Lys Pro Val Thr Ala Leu Glu Tyr Thr Phe	100	105	110
Ser Arg Ser Asp Pro Thr Ser Tyr Ala Gly Tyr Ile Glu Asp Leu Lys	115	120	125
Lys Phe Leu Lys Pro Tyr Thr Leu Glu Glu Gln Lys Asn Leu Thr Val	130	135	140
Cys Pro Asp Gly Ala Leu Phe Glu Gln Lys Gly Pro Val Tyr Val Ala	145	150	160
Cys Gln Phe Pro Ile Ser Leu Leu Gln Ala Cys Ser Gly Met Asn Asp	165	170	175
Pro Asp Phe Gly Tyr Ser Gln Gly Asn Pro Cys Ile Leu Val Lys Met	180	185	190
Asn Arg Ile Ile Gly Leu Lys Pro Glu Gly Val Pro Arg Ile Asp Cys	195	200	205
Val Ser Lys Asn Glu Asp Ile Pro Asn Val Ala Val Tyr Pro His Asn	210	215	220
Gly Met Ile Asp Leu Lys Tyr Phe Pro Tyr Tyr Gly Lys Lys Leu His	225	230	240
Val Gly Tyr Leu Gln Pro Leu Val Ala Val Gln Val Ser Phe Ala Pro	245	250	255
Asn Asn Thr Gly Lys Glu Val Thr Val Glu Cys Lys Ile Asp Gly Ser	260	265	270
Ala Asn Leu Lys Ser Gln Asp Asp Arg Asp Lys Phe Leu Gly Arg Val	275	280	285
Met Phe Lys Ile Thr Ala Arg Ala	290	295	

<210> SEQ ID NO 11

<211> LENGTH: 3072

<212> TYPE: DNA

<213> ORGANISM: homo sapiens

<400> SEQUENCE: 11

```

atggggaagg ggggtggacg tgataagtat gagcctgcag ctgtttcaga acaaggtgat      60
aaaaagggca aaaagggcaa aaaagacagg gacatggatg aactgaagaa agaagtttct      120
atggatgatc ataaacttag ccttgatgaa cttcatcgta aatatggaac agacttgagc      180
cggggattaa catctgctcg tgcagctgag atcctggcgc gagatgggcc caacgcctc      240
actccccctc ccontactcc tgaatggatc aagttttgtc ggcagctctt tgggggggtc      300
tcaatgttac tgtggattgg agcgattctt tgtttcttgg cttatagcat ccaagetget      360
acagaagagg aacctcaaaa cgataatctg tacctgggtg tgggtctatc agccgttgta      420
atcataactg gttgcttctc ctactatcaa gaagctaaaa gttcaaagat catggaatcc      480

```

-continued

ttcaaaaaca	tggtccctca	gcaagccctt	gtgattcgaa	atggtgagaa	aatgagcata	540
aatgctggagg	aagttgtggt	tggggatctg	gtggaagtaa	aaggaggaga	ccgaattcct	600
gctgacctca	gaatcatatc	tgcaaatggc	tgcaaggtag	ataactcctc	gctcactggt	660
gaatcagaac	cccagactag	gtctccagat	ttcacaatg	aaaacccctc	ggagacgagg	720
aacattgcct	tcttttcaac	aaattgtgtt	gaaggcaccg	cacgtggtat	tgttgtctac	780
actggggatc	gcactgtgat	gggaagaatt	gccacacttg	cttctgggct	ggaaggaggc	840
cagaccccca	ttgctgcaga	aattgaacat	tttatccaca	tcatcacggg	tgtggctgtg	900
ttcctgggtg	tgtctttctt	catcctttct	ctcatccttg	agtacacctg	gcttgaggct	960
gtcatcttcc	tcatcggtat	catcgtagcc	aatgtgccgg	aaggtttgc	ggccactgtc	1020
acggtctgtc	tgacacttac	tgccaacgc	atggcaagga	aaaactgctt	agtgaagaac	1080
ttagaagctg	tggagacctt	ggggctccacg	tccaccatct	gctctgataa	aactggaact	1140
ctgactcaga	accggatgac	agtggcccac	atgtggtttg	acaatcaaat	ccatgaagct	1200
gatacgacag	agaatcagag	tgtgtctctt	tttgacaaga	cttcagctac	ctggcttget	1260
ctgtccagaa	ttgcaggtct	ttgtaacagg	gcagtgttcc	aggctaacca	ggaaaactta	1320
cctattctta	agcgggcagt	tgcaaggagat	gcctctgagt	cagcactctt	aaagtgcata	1380
gagctgtgct	gtggttccgt	gaaggagatg	agagaaagat	acgcaaaaat	cgctcgagata	1440
cccttcaact	ccaccaacaa	gtaccagttg	tctattcata	agaaccccaa	cacatcggag	1500
ccccaacacc	tgttggatg	gaagggcgcc	ccagaaagga	tcctagaccg	ttgcagctct	1560
atcctcctcc	acggcaagga	gcagccctg	gatgaggagc	tgaagagcgc	ctttcagaac	1620
gcctatattg	agctgggggg	cctcgagaaa	cgagtcctag	gtttctgcca	cctctttctg	1680
ccagatgaac	agtttctcga	agggttccag	tttgacactg	acgatgtgaa	ttccctatc	1740
gataatctgt	gctttgttgg	gctcatctcc	atgattgacc	ctccacgggc	ggcgttctct	1800
gatgccgtgg	gcaaatgtcg	aagtgtctga	attaaggtca	tcatggtcac	aggagaccat	1860
ccaatcacag	ctaaagctat	tgccaaggt	gtgggcatca	tctcagaagg	caatgagacc	1920
gtggaagaca	ttgctgccc	cctcaacatc	ccagtcagcc	agggtgaacc	cagggatgcc	1980
aaggcctcgc	tagtacacgg	cagtgatcta	aaggacatga	cctccgagca	gctggatgac	2040
atthtgaagt	accacactga	gatagtgttt	gccaggacct	cccctcagca	gaagctcatc	2100
attgtggaag	gctgccaag	acaggggtgct	atcgtggctg	tgactggtga	cggtgtgaat	2160
gactctccag	ctttgaagaa	agcagacatt	ggggttgcta	tggggattgc	tggtccagat	2220
gtgtccaagc	aagctgtgta	catgattctt	ctggatgaca	actttgcctc	aattgtgact	2280
ggagtagagg	aaggtctgtc	gatctttgat	aacttgaaga	aatccattgc	ttatacctta	2340
accagtaaca	ttcccagat	caccccgctc	ctgatattta	ttattgcaaa	cattccacta	2400
ccactgggga	ctgtcaccat	cctctgcatt	gacttgggca	ctgacatggt	tcctgccatc	2460
tccttgctt	atgagcaggc	tgagagtgtg	atcatgaaga	gacagcccag	aaatcccaaa	2520
acagacaaac	ttgtgaatga	gcgctgtgac	agcatggcct	atgggcagat	tggaatgatc	2580
caggccctgg	gaggtctctt	tacttacttt	gtgattctgg	ctgagaacgg	cttcctccca	2640
attcacctgt	tgggcctccg	agtggactgg	gatgaccgct	ggatcaacga	tgtggaagac	2700
agctacgggc	agcagtggtg	ctatgagcag	aggaaaatcg	tggagttcac	ctgccacaca	2760
gccttcttcg	tcagtatcgt	ggtggtgcag	tgggcccact	tggtcatctg	taagaccagg	2820

-continued

aggaattcgg tcttccagca ggggatgaag aacaagatct tgatatttgg cctctttgaa	2880
gagacagccc tggctgcttt cctttcctac tgccctggaa tgggtgttgc tcttaggatg	2940
tatcccctca aacctacctg gtggttctgt gccttccct actctcttct catcttctga	3000
tatgacgaag tcagaaaact catcatcagg cgacgcctg gcggtgggt ggagaaggaa	3060
acctactatt ag	3072

<210> SEQ ID NO 12

<211> LENGTH: 3063

<212> TYPE: DNA

<213> ORGANISM: homo sapiens

<400> SEQUENCE: 12

atgggccgtg gggctggcgg tgagtactca cctgcccga ccacggcaga gaatgggggc	60
ggcaagaaga aacagaagga gaaggaactg gatgagctga agaaggaggt ggcaatggat	120
gaccacaagc tgtccttggg tgagctgggc cgcaataacc aagtggacct gtccaagggc	180
ctcaccaacc agcgggctca ggacgttctg gctcgagatg ggcccaacgc cctcacacca	240
cctcccacaa cccctgagtg ggtcaagttc tgccgtcagc ttttcggggg gttctccatc	300
ctgctgtgga ttgggctat cctctgcttc ctggcctacg gcatccagge tgccatggag	360
gatgaacat ccaacgacaa tctatatctg ggtgtggtgc tggcagctgt ggtcattgtc	420
actggctgct tctcctacta ccaggaggcc aagagctcca agatcatgga ttccttcaag	480
aacatggtac ctcagcaagc ccttgtgatc cgggaggag agaagatgca gatcaacgca	540
gaggaagtgg tgggtggaga cctggtggag gtgaagggtg gagaccgct ccctgctgac	600
ctccggatca tctcttctca tggctgtaag gtggataact catcctaac aggagagtcg	660
gagccccaga cccgctcccc cgagttcacc catgagaacc ccctggagac ccgcaatate	720
tgtttcttct ccaccaactg tgttgaagge actgccaggg gcattgtgat tgccacagga	780
gaccggacgg tgatgggccc catagctact ctgcctcag gcctggaggt tgggcccaca	840
cccatagcaa tggagattga acacttcate cagctgatca cagggtgcgc tgtattctg	900
ggggtctcct tcttctgctc ctcccctate ctgggctaca gctggctgga ggcagtcate	960
ttcctcatcg gcatcatagt ggccaactg cctgaggggc ttctggccac tgtcaactgt	1020
tgctgaccc tgacagccaa gcgcatggca cggagaact gcctggtgaa gaacctggag	1080
gcggtggaga cgctgggctc cacgtccacc atctgctcgg acaagacggg cacctcacc	1140
cagaaccgca tgaccctgc ccacatgtgg ttcgacaacc aaatccatga ggctgacacc	1200
accgaagatc agtctggggc cacttttgac aaacgatccc ctacgtggac ggcctgtct	1260
cgaattgctg gtctctgcaa ccgcccctc ttcaaggcag gacaggagaa catctccgtg	1320
tctaagcggg acacagctgg tgatgcctct gagtcagctc tgctcaagtg cattgagctc	1380
tcctgtggct cagtgaggaa aatgagagac agaaaaccca aggtggcaga gattccttct	1440
aactctacca acaagtaacca gctgtctatc caccagcag aagacagccc ccagagccac	1500
gtgctggtga tgaagggggc ccagagcgc attctggacc ggtgctccac catcctggtg	1560
cagggcaagg agatcccgc ctgacaagg atgcaagatg cctttcaaaa tgctacatg	1620
gagctggggg gacttgggga gcgtgtgctg ggattctgtc aactgaatct gccatctgga	1680
aagtttctc ggggcttcaa attcgacac gatgagctga actttcccac ggagaagctt	1740
tgctttgtgg ggctcatgtc tatgattgac cctccccggg ctgctgtgcc agatgctgtg	1800
ggcaagtgc gaagcgcagg catcaagtg atcatggtaa ccgggatca ccctatcaca	1860

-continued

gccaaggcca	ttgccaagg	cgtgggcac	atatcagagg	gtaacgagac	tgtggaggac	1920
attgcagccc	ggctcaacat	tcccagagt	caagtcaacc	ccagagaagc	caaggcatgc	1980
gtggtgcacg	gctctgacct	gaaggacatg	acatcggagc	agctcogatga	gatcctcaag	2040
aaccacacag	agatcgtcct	tgctegaacg	tctcccagc	agaagctcat	cattgtggag	2100
ggatgtcaga	ggcagggagc	cattgtggcc	gtgacgggtg	acggggtgaa	cgactcccct	2160
gcattgaaga	aggctgacat	tggcattgcc	atgggcacat	ctggctctga	cgtctctaag	2220
caggcagccg	acatgatcct	gctggatgac	aactttgcct	ccatcgtcac	gggggtggag	2280
gagggccgcc	tgatcttga	caacttgaag	aaatccatcg	cctacacct	gaccagcaac	2340
atccccgaga	tcacccccct	cctgctgttc	atcattgcca	acatccccct	acctctgggc	2400
actgtgacca	tcctttgcat	tgacctgggc	acagatatgg	tcctgccat	ctcctggcc	2460
tatgaggcag	ctgagagtga	tatcatgaag	cggcagccac	gaaactccca	gacggacaag	2520
ctggtgaatg	agaggctcat	cagcatggcc	tacggacaga	tcgggatgat	ccaggcactg	2580
ggtggctctt	tcacctactt	tgtgatcctg	gcagagaacg	gtttcctgcc	atcacggcta	2640
ctgggaatcc	gcctcgaactg	ggatgaccgg	accatgaatg	atctggagga	cagctatgga	2700
caggagtgga	cctatgagca	gcggaagggtg	gtggagtcca	cgtgccacac	ggcattcttt	2760
gccagcatcg	tggtggtgca	gtgggtgac	ctcatcatct	gcaagacccg	ccgcaactca	2820
gtcttcacgc	agggcatgaa	gaacaagatc	ctgatttttg	ggctcctgga	ggagacggcg	2880
ttggctgect	ttctctctta	ctgcccaggc	atgggtgtag	ccctccgcat	gtacccgctc	2940
aaagtcacct	ggtggtctcg	cgctctccc	tacagctccc	tcattctcat	ctatgatgag	3000
gtccgaaagc	tcctcctgcg	gcggtatcct	ggtgctggg	tggagaagga	gacatactac	3060
tga						3063

<210> SEQ ID NO 13

<211> LENGTH: 3042

<212> TYPE: DNA

<213> ORGANISM: homo sapiens

<400> SEQUENCE: 13

atgggggaca	agaaagatga	caaggactca	cccaagaaga	acaagggcaa	ggagcgcg	60
gacctggatg	acctcaagaa	ggaggtggct	atgacagagc	acaagatgtc	agtggaagag	120
gtctgccgga	aatacaacac	agactgtgtg	cagggtttga	cccacagcaa	agcccaggag	180
atcctggccc	gggatgggccc	taacgcactc	acgccaccgc	ctaccacccc	agagtgggtc	240
aagttttgcc	ggcagctcct	cgggggcttc	tccatcctgc	tgtggatcgg	ggctatcctc	300
tgtctcctgg	cctacggtat	ccagggcggc	accgaggacg	accctctgg	tgacaacctg	360
tacctgggca	tcgtgctggc	ggcgtgggtg	atcatcactg	gctgcttctc	ctactaccag	420
gaggccaaga	gctccaagat	catggagtcc	ttcaagaaca	tggtgcccca	gcaagccctg	480
gtgatccggg	aaggtgagaa	gatgcagggtg	aacgctgagg	aggtgggtgg	cggggacctg	540
gtggagatca	aggggtggaga	ccgagtgcc	gctgacctgc	ggatcatctc	agcccacggc	600
tgcaaggtgg	acaactcctc	cctgactggc	gaatccgagc	cccagactcg	ctctcccagc	660
tgcacgcacg	acaacccctt	ggagactcgg	aacatcacct	tcttttccac	caactgtgtg	720
gaaggcacgg	ctcggggcgt	ggtggtggcc	acgggcgacc	gcaactgcat	gggcccgtatc	780
gccaccctgg	catcagggtc	ggaggtgggc	aagacgccc	tcgccatcga	gattgagcac	840

-continued

```

ttcatccagc tcatacccg cgtggctgtc ttectgggtg tctccttctt catcctctcc 900
ctcattctcg gatacacctg gcttgaggct gtcactctcc tcatcggeat catcgtggcc 960
aatgtcccag agggctctgt ggccaactgt actgtgtgtc tgacgctgac cgccaagcgc 1020
atggcccgga agaactgect ggtgaagaac ctggaggctg tagaaacctt gggtccacg 1080
tccaccatct gctcagataa gacagggacc ctcaactcaga accgcatgac agtcgcccac 1140
atgtggtttg acaaccagat ccacgaggct gacaccactg aggaccagtc agggacctca 1200
tttgacaaga gttcgcacac ctgggtggcc ctgtctcaca tcgctgggct ctgcaatcgc 1260
gctgtcttca aggggtgtca ggacaacatc cctgtgtctc agagggatgt ggctggggat 1320
gcgctgagtg ctgccctgct caagtgcacg gagctgtcct ctggctccgt gaagctgatg 1380
cgtgaacgca acaagaaagt ggctgagatt cccttcaatt ccaccaacaa ataccagctc 1440
tccatccatg agaccgagga ccccaacgac aaccgatacc tgctgggtgat gaagggtgcc 1500
cccgagcgca tctcggaccg ctgctccacc atcctgctac agggcaagga gcagcctctg 1560
gacgaggaaa tgaaggagge cttccagaat gcctaccttg agctcgggtg cctgggcgag 1620
cgctgtcttg gtttctgcca ttattacctg cccgaggagc agttcccaca gggctttgcc 1680
ttcgactgtg atgacgtgaa cttcaccacg gacaacctct gctttgtggg cctcatgtcc 1740
atgatcgacc caccctgggc agcctgcctt gacgcgggtg gcaagtgtcg cagcgcaggc 1800
atcaaggtca tcattggctac cggcgatcac cccatcacgg ccaaggccat tgccaagggc 1860
gtgggcatca tctctgaggg caacgagact gtggaggaca tcgcccaccg gctcaacatt 1920
cccgtcagcc aggttaacc cgggatgcc aaggcctgct tgatccacgg caccgacctc 1980
aaggacttca cctccgagca aatcgacgag atcctgcaga atcacaccga gatcgtcttc 2040
gccccacat cccccagca gaagctcacc attgtggagg gctgtcagag acaggggtgca 2100
attgtggctg tgaccgggga tgggtgtaac gactcccccg ctctgaagaa ggccgacatt 2160
ggggtggcca tgggcatcgc tggctctgac gtctccaagc aggcagctga catgatcctg 2220
ctggacgaca actttgctc catcgtcaca ggggtggagg agggccgctt gatctctgac 2280
aacctaaaga agtccattgc ctacacctg accagcaata tcccggagat cagcccttc 2340
ctgctgttca tcattggcaa catcccctg cccctgggca ccatcaccat cctctgcatc 2400
gatctgggca ctgacatggt cctgcccac tcaactggct acgaggctgc cgaagcgcac 2460
atcatgaaga gacagcccag gaaccgcgg acggacaaat tggtaaatga gagactcatc 2520
agcatggcct acgggcagat tggaaatgat caggctctcg gtggcttctt ctcttacttt 2580
gtgatcctgg cagaaaatgg cttcttgccc ggcaacctgg tgggcatccg gctgaaatgg 2640
gatgaccgca ccgtcaatga cctggaagac agttacgggc agcagtggac atacgagcag 2700
aggaaggtgg tggagtccac ctgccacacg gccttctttg tgagcatcgt tgtcgtccag 2760
tgggcccgatc tgatcatctg caagaccgg aggaactcgg tcttcagca gggcatgaag 2820
aacaagatcc tgatctcgg gctggttgag gagacggccc tggctgcctt cctgtcctac 2880
tgccccgca tggacgtggc cctgcgcgatg taccctctca agcccagctg gtggttctgt 2940
gccttcccct acagtttctt catctctgct tacgacgaaa tccgcaaacct catcctgcgc 3000
aggaaccacg ggggttgggt ggagaaggaa acctactact ga 3042

```

<210> SEQ ID NO 14

<211> LENGTH: 3090

<212> TYPE: DNA

<213> ORGANISM: homo sapiens

-continued

<400> SEQUENCE: 14

atggggcttt gggggaagaa agggacagt gctccccatg accagagtcc aagacgaaga 60
cctaaaaaag ggcttatcaa gaaaaaatg gtgaagaggg aaaaacagaa gcgcaatatg 120
gaggaactga agaaggaagt ggtcatggat gatcacaaat taaccttggga agagctgagc 180
accaagtact ccgtggacct gacaaagggc catagccacc aaagggcaaa ggaaatcctg 240
actcgaggtg gacccaatac tgttaccca cccccacca ctccagaatg ggtcaaattc 300
tgtaagcaac tgttcggagg cttctccctc ctactatgga ctggggccat tctctgcttt 360
gtggcctaca gcatccagat atatttcaat gaggagccta ccaaagacaa cctctactg 420
agcatcgtac tgtccgtcgt ggtcatcgtc actggctgct tctcctatta tcaggaggcc 480
aagagctcca agatcatgga gtcttttaag aacatgggtc ctcagcaagc tctggtaatt 540
cgaggaggag agaagatgca aattaatgta caagaggtgg tgttgggaga cctggtggaa 600
atcaagggtg gagaccgagt cctctctgac ctccggctta tctctgcaca aggatgtaag 660
gtggacaact catccttgac tggggagtca gaacccaga gccgctcccc tgacttcacc 720
catgagaacc ctctggagac ccgaaacatc tgcttctttt ccaccaactg tgtggaagga 780
accgcccggg gtatttgtat tctacggga gactccacag tgatgggcag aattgcctcc 840
ctgacgtcag gccctggcgt tggccagaca cctatcgtg ctgagatoga acacttcac 900
catctgatca ctgtggtggc cgtcttctt ggtgtcactt tttttgcgt ctcacttctc 960
ttgggctatg gttggctgga ggctatcatt tttctcattg gcatcattgt ggccaatgtg 1020
cctgaggggc tgttggctac agtcaactgt tgcctgacct tcacagccaa gcgcatggca 1080
cggaagaact gcctggtgaa gaacctggag gcggtggaga cgctgggctc cacgtccacc 1140
atctgctcag acaagacggg caccctcacc cagaaccgca tgaccgtcgc ccacatgtgg 1200
tttgatatga ccgtgtatga ggccgacacc actgaagaac agactggaaa aacatttacc 1260
aagagctctg atacctggtt tatgctggcc cgaatcgtg gcctctgcaa ccgggctgac 1320
ttaaaggcta atcaggagat cctgcccatt gctaagaggg ccacaacagg tgatgcttcc 1380
gagtcagccc tcctcaagtt catcgagcag tcttacagct ctgtggcggga gatgagagag 1440
aaaaaccca aggtggcaga gattcccttt aattctacca acaagtacca gatgtccatc 1500
caccttcggg aggacagctc ccagacccac gtactgatga tgaaggggtc tccggagagg 1560
atcttgaggt tttgtctac ctttctctg aatgggcagg agtactcaat gaacgatgaa 1620
atgaaggaag ccttccaaaa tgcctattta gaactgggag gtctggggga acgtgtgcta 1680
ggcttctgct tcttgaatct gcctagcagc ttctccaagg gattccatt taatacagat 1740
gaaataaatt tcccattgga caaccttgt tttgtgggcc tcatatccat gattgaccct 1800
ccccgagctg cagtcctga tctgtgagc aagtgtcga gtgcaggaat taaggatgatc 1860
atggtaacag gagatcatcc cattacagct aaggccattg ccaaggggtg gggcatcatc 1920
tcagaaggca ctgagacggc agaggaagtc gctgccggc ttaagatccc tatcagcaag 1980
gtcgatgcca gtgctgcaa agccattgtg gtgcatggtg cagaactgaa ggacatacag 2040
tccaagcagc ttgatcagat cctccagaac caccctgaga tcgtgtttgc tcggacctcc 2100
cctcagcaga agctcatcat tctcagagga tctcagaggc tgggagccgt tgtggccgtg 2160
acaggtgacg ggggtaacga ctcccctgcg ctgaagaagg ctgacattgg cattgccatg 2220
ggcatctctg gctctgacgt ctctaagcag gcagccgaca tgatcctgct ggatgacaac 2280

-continued

```

tttgcctcca tcgtcacggg ggtggaggag gcccgcctga tctttgacaa cctgaagaaa 2340
tccatcatgt acaccctgac cagcaacatc cccgagatca cgcccttctc gatgttcac 2400
atcctcggtg tacccctgcc tctgggaacc ataaccatcc tctgcattga tctcgccact 2460
gacatggtcc ctgccatctc cttggcttat gagtcagctg aaagcgacat catgaagagg 2520
cttccaagga acccaaagac ggataatctg gtgaaccacc gtctcattgg catggcctat 2580
ggacagattg ggatgatcca ggtctgggct ggattcttta cctactttgt aatcctggct 2640
gagaatggtt ttaggcctgt tgatctgctg ggcatccgcc tccactggga agataaatac 2700
ttgaatgacc tggaggacag ctacggacag cagtggacct atgagcaacg aaaagtgtg 2760
gagttcacat gccaaacggc cttttttgtc accatcgtgg ttgtgcagtg ggcggatctc 2820
atcatctcca agactcgccg caactcactt ttccagcagg gcatgagaaa caaagtctta 2880
atatttggga tcctggagga gacactcttg gctgcatttc tgctctacac tccaggcatg 2940
gacgtggccc tgcgaatgta cccactcaag ataacctggg ggctctgtgc cattccctac 3000
agtattctca tcttcgtcta tgatgaaatc agaaaactcc tcatccgtca gcaccggat 3060
ggctgggtgg aaaggagac gtactactaa 3090

```

<210> SEQ ID NO 15

<211> LENGTH: 963

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic Acid sequence encoding His10-ATP1B1

<400> SEQUENCE: 15

```

atggccagat cacatcatca ccatcaccac catcaccatc atcctaggag atctgcccgc 60
gggaaagcca aggaggaggg cagctggaag aaattcatct ggaactcaga gaagaaggag 120
tttctgggca ggaccggtgg cagttggttt aagatccttc tattctacgt aatattttat 180
ggctgcctgg ctggcatctt catcggaacc atccaagtga tgctgctcac catcagtgaa 240
tttaagccca catatcagga ccgagtggcc ccgccaggat taacacagat tcctcagatc 300
cagaagactg aaatttcctt tcgtccta atgatccaaga gctatgagggc atatgtactg 360
aacatagtta ggttcctgga aaagtacaaa gattcagccc agagggatga catgattttt 420
gaagattgtg gcgatgtgcc cagtgaaccg aaagaacgag gagactttaa tcatgaacga 480
ggagagcgaa aggtctgcag attcaagctt gaatggctgg gaaattgctc tggattaaat 540
gatgaaactt atggctacaa agagggcaaa ccgtgcatta ttataaagct caaccgagtt 600
ctaggcttca aacctaaagc tccaagaat gagtccttgg agacttacc agtgatgaag 660
tataacccaa atgctcctcc cgttcagtgc actggcaagc gagatgaaga taaggataaa 720
gttgaaaatg tggagtattt tggactgggc aactcccctg gttttcctct gcagtattat 780
ccgtactatg gcaaacctct gcagcccaaa tacctgcagc ccctgctggc cgtacagttc 840
accaatctta ccatggacac tgaattcgc atagagtgta aggcgtacgg tgagaacatt 900
gggtacagtg agaagaccg ttttcaggga cgttttgatg taaaaattga agttaagagc 960
tga 963

```

<210> SEQ ID NO 16

<211> LENGTH: 924

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic Acid sequence encoding His10-ATP1B2

-continued

<400> SEQUENCE: 16

```

atggccagat cacatcatca ccatcaccac catcaccatc atcctaggag atctgtcacc 60
cagaaagaga agaagagctg cgggcagggtg gttgaggagt ggaaggagtt cgtgtggaac 120
ccgaggagcgc accagtttat gggccgcacc gggaccagct gggcctttat cctcctcttc 180
tacctcgttt tttatgggtt ccccacgcc atgttcaccc tcaccatgtg ggtgatgctg 240
cagactgtct ccgaccatac ccccaagtac caggaccgac tggccacacc gggcttgatg 300
attcgcccca agactgagaa ccttgatgtc attgtcaatg tcagtgcacc tgaagctgg 360
gaccagcatg ttcagaagct caacaagttc ttggagcctt acaacgactc tatgcaagcc 420
caaaagaatg atgtctgccc ccttggggcgc tattacgaac agccagataa tggagtccctc 480
aactacccca aactggcctg ccaattcaac cggaccaccg tgggcaactg ctccggcatt 540
ggggactcca cccactatgg ttacagcact gggcagccct gtgtcttcat caagatgaac 600
cgggtcatca acttctatgc aggagcaaac cagagcatga atgttacctg tgctgggaa 660
cgagatgaag atgctgagaa tctcggaac ttcgtcatgt tccccgcaa cggcaacatc 720
gacctcatgt acttccccta ctatggcaaa aagttccacg tgaactacac acagccctg 780
gtggctgtga agttcctgaa tgtgaccccc aacgtggagg tgaatgtaga atgtcgcac 840
aacgccgcca acatcgccac agacgatgag cgagacaagt tcgccggccg cgtggccttc 900
aaactccgca tcaacaaaac ctga 924

```

<210> SEQ ID NO 17

<211> LENGTH: 891

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic Acid sequence encoding His10-ATP1B3

<400> SEQUENCE: 17

```

atggccagat cacatcatca ccatcaccac catcaccatc atcctaggag atctacgaag 60
aacgagaaga agtccctcaa ccagagcctg gccgagtgga agctcttcat ctacaacccg 120
accaccggag aattcctggg ggcacccgcc aagagctggg gtttgatctt gctcttctac 180
ctagtttttt atgggttctt ggctgcactc ttctcattca cgatgtgggt tatgcttcag 240
actctcaacg atgaggttcc aaaataccgt gaccagatcc ctagcccagg actcatgggt 300
tttccaaaac cagtgcaccg attggaatat acattcagta ggtctgatcc aacttcgcat 360
gcagggtaca ttgaagacct taagaagttt ctaaaacat atactttaga agaacagaag 420
aacctcacag tctgtcctga tggagcactt tttgaacaga agggctccagt ttatgttgca 480
tgtcagtttc ctatttcatt acttcaagca tgcagtggta tgaatgatcc tgattttggc 540
tattctcaag gaaacccttg tattcttgtg aaaatgaaca gaataattgg attaaagcct 600
gaaggagtgc caaggataga ttgtgtttca aagaatgaag atataccaaa tgtagcagtt 660
tatcctcata atggaatgat agacttaaaa tatttccat attatgggaa aaaactgcat 720
gttgggtatc tacagccatt ggttctgtt caggtcagct ttgctcctaa caaacctggg 780
aaagaagtaa cagttgagtg caagattgat ggatcagcca acctaaaaag tcaggatgat 840
cgtgacaagt ttttgggacg agttatgttc aaaatcacag cacgtgcata g 891

```

<210> SEQ ID NO 18

<211> LENGTH: 288

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Nucleic Acid sequence encoding Human His6-FXYD1

<400> SEQUENCE: 18

```

atgggcagca gccatcatca tcacatcac tccgcgggtg aaaacctgta cttccagggt    60
accaaggcag aaagtccaaa ggaacacgac ccgttcactt acgactacca gtccttcgag    120
atcgagggcc tcgtcatgac cgggatcctc ttcactctgg gcactcctcat cgtgctgagc    180
agaagatgcc ggtgcaagtt caaccagcag cagaggactg gggaaccgca tgaagaggag    240
ggaactttcc gcagctccat ccgcccgtctg tccaccgcga ggcggtag                288
  
```

<210> SEQ ID NO 19
 <211> LENGTH: 95
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Amino Acid sequence encoding Human His6-FXYD1



<400> SEQUENCE: 19

```

Met Gly Ser Ser His His His His His Ser Ala Gly Glu Asn Leu
 1         5         10        15
Tyr Phe Gln Gly Thr Lys Ala Glu Ser Pro Lys Glu His Asp Pro Phe
 20        25        30
Thr Tyr Asp Tyr Gln Ser Leu Gln Ile Gly Gly Leu Val Ile Ala Gly
 35        40        45
Ile Leu Phe Ile Leu Gly Ile Leu Ile Val Leu Ser Arg Arg Cys Arg
 50        55        60
Cys Lys Phe Asn Gln Gln Gln Arg Thr Gly Glu Pro Asp Glu Glu Glu
 65        70        75        80
Gly Thr Phe Arg Ser Ser Ile Arg Arg Leu Ser Thr Arg Arg Arg
 85        90        95
  
```

What is claimed is:

1. A compound represented by general Formula I:




including any pharmaceutically acceptable salt, hydrate, solvate, enantiomer and diastereomer thereof, and any mixtures thereof,

wherein:

X is H or OH;

R is represented by general Formula II:



A is $-\text{CH}_2-$, $-(\text{CH}_2)_2-$ or a covalent bond; and

B is a cyclic moiety selected from the group consisting of an unsubstituted alicyclic moiety, a substituted alicyclic moiety, an unsubstituted heterocyclic moiety, a substituted heterocyclic moiety, an unsubstituted heteroaryl moiety and a substituted heteroaryl moiety;

or

B is selected from the group consisting of an alkylsulfonyl, an arylsulfonyl and a sulfonamide;

or

B is $-\text{N}_1\text{R}_2$, wherein R_1 and R_2 are each independently H or a C_1 - C_4 alkyl provided that at least one of R_1 and R_2 is a C_1 - C_4 alkyl.

2. The compound of claim 1, wherein A is selected from the group consisting of a covalent bond, an unsubstituted C_1 - C_6 alkyl, a substituted C_1 - C_6 alkyl, an unsubstituted

121

C_1 - C_6 alkyl interrupted by one or more heteroatom and a substituted C_1 - C_6 alkyl interrupted by one or more heteroatom.

3. The compound of claim 1, wherein said unsubstituted alicyclic moiety is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

4. The compound of claim 1, wherein said unsubstituted heteroaryl moiety is imidazolyl.

5. The compound of claim 1, wherein said alkylsulfonyl is selected from the group consisting of methylsulfonyl, ethylsulfonyl and isopropylsulfonyl.

6. The compound of claim 1, wherein said arylsulfonyl is selected from the group consisting of phenylsulfonyl, benzylsulfonyl and tosyl.

7. The compound of claim 1, wherein said sulfonamide is selected from the group consisting of methylsulfonamide, N-methylmethanesulfonamide and N,N-dimethylmethanesulfonamide.

8. The compound of claim 1, wherein B is $-N(Et)_2$.

9. The compound of claim 1, wherein R is selected from the group consisting of cyclopropyl, methylcyclopropane, ethylcyclopropane, propylcyclopropane, cyclobutyl, methylcyclobutane, methyl-3,3-dimethylcyclobutane, ethylcyclobutane, propylcyclobutane, cyclopentyl, methylcyclopentane, ethylcyclopentane, propylcyclopentane, cyclohexyl, azetidiny, oxetanyl, thietanyl, histaminy and benzyl.

10. The compound of claim 1, having an affinity to at least one isoform of Na,K-ATPase, wherein said isoform is selected from the group consisting of $\alpha 1\beta 1$, $\alpha 1\beta 2$, $\alpha 1\beta 3$, $\alpha 2\beta 1$, $\alpha 2\beta 2$, $\alpha 2\beta 3$, $\alpha 3\beta 1$, $\alpha 3\beta 2$, $\alpha 3\beta 3$, $\alpha 4\beta 1$, $\alpha 4\beta 2$ and $\alpha 4\beta 3$.

11. The compound of claim 10, wherein said affinity to any one of $\alpha 2\beta 1$, $\alpha 2\beta 2$ and $\alpha 2\beta 3$ is higher than the affinity to $\alpha 1\beta 1$, $\alpha 1\beta 2$, $\alpha 1\beta 3$, $\alpha 3\beta 1$, $\alpha 3\beta 2$, $\alpha 3\beta 3$, $\alpha 4\beta 1$, $\alpha 4\beta 2$ and $\alpha 4\beta 3$ by at least 100%.

12. A pharmaceutical composition comprising as an active ingredient a compound of claim 1 and a pharmaceutically acceptable carrier.

13. A method of reducing intraocular pressure (TOP) in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of claim 1.


122

14. A method of treating a heart condition in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of claim 1.

15. A pharmaceutical composition comprising as active ingredients:

at least one ingredient selected from the group consisting of a prostaglandin analog, a β -blocker, an adrenergic agent, an $\alpha 2$ -adrenergic receptor agonist, a miotic agent, a carbonic anhydrase inhibitor and a cholinergic agonist; and

a compound represented by Formula III:




Formula III

including any pharmaceutically acceptable salt, hydrate, solvate, enantiomer and diastereomer thereof, and any mixtures thereof, and a pharmaceutically acceptable carrier,

wherein:

X is H or OH;

R' is represented by general Formula II:



Formula II

A is $-CH_2-$, $-(CH_2)_2-$ or a covalent bond; and

B is a cyclic moiety selected from the group consisting of an unsubstituted alicyclic moiety, a substituted alicyclic moiety, an unsubstituted heterocyclic moiety, a substituted heterocyclic moiety, an unsubstituted heteroaryl moiety and a substituted heteroaryl moiety, or B is selected from the group consisting of an alkylsulfonyl, an arylsulfonyl and a sulfonamide, or B is $-N_1R_2$, wherein R_1 and R_2 are each independently H or a C_1 - C_4 alkyl provided that at least one of R_1 and R_2 is a C_1 - C_4 alkyl, and a pharmaceutically acceptable carrier.

16. The pharmaceutical composition of claim 15, being packaged in a packaging material and identified in print, or on said packaging material, for use in reducing intraocular pressure (TOP).

17. A method of treating a heart condition in a subject in need thereof, comprising co-administering to the subject a therapeutically effective amount of:


an agent selected from the group consisting of a β -blocker, an anticoagulation agent, an angiotensin-converting-enzyme inhibitor and an angiotensin II receptor antagonist; and

123

a compound represented by Formula III:

124

Formula III



including any pharmaceutically acceptable salt, hydrate, solvate, enantiomer and diastereomer thereof, and any mixtures thereof,

wherein:

X is H or OH;


R' is represented by general Formula II:

25

A is a spacer moiety or a covalent bond; and

B is a cyclic moiety, or B is selected from the group consisting of an alkylsulfonyl, an arylsulfonyl and a sulfonamide, or B is —N₁R₂, wherein R₁ and R₂ are each independently H or a C₁-C₄ alkyl provided that at least one of R₁ and R₂ is a C₁-C₄ alkyl.

Formula II 30



* * * * *