

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 199748284 B2**
(10) Patent No. **731045**

(54) Title
Method for promoting ovulation parturition and lactation in mammals

(51)⁶ International Patent Classification(s)
A61K 031/445

(21) Application No: **199748284** (22) Application Date: **1997.12.10**

(30) Priority Data

(31) Number (32) Date (33) Country
08/768981 1996.12.18 US

(43) Publication Date : **1998.06.25**
(43) Publication Journal Date : **1998.06.25**
(44) Accepted Journal Date : **2001.03.22**

(71) Applicant(s)
Clemson University

(72) Inventor(s)
Dee L Cross

(74) Agent/Attorney
F B RICE and CO,605 Darling Street,BALMAIN NSW 2041

(56) Related Art
US 5372818
EUR. J. OBSTET.GYNECOL.REPROD.BIOL (1985)19:281-287

Abstract of the Disclosure

The present invention generally relates to various methods for promoting ovulation, parturition and lactation in female mammals. These benefits are obtained by administering to the mammals a composition containing a D₂ receptor antagonist that does not substantially cross the blood brain barrier. In one embodiment, the D₂ receptor antagonist is domperidone. The domperidone can be administered to the mammal either orally or subcutaneously and can be used to treat, for instance, anestrous mammals, mammals that have problems bearing offspring and mammals suffering from agalactia. Unexpectedly, it has also been discovered that the D₂ receptor antagonist may also stimulate feed intake, causing the mammal to eat more and gain weight faster.

AUSTRALIA

Patents Act 1990

CLEMSON UNIVERSITY

ORIGINAL

COMPLETE SPECIFICATION
STANDARD PATENT

Invention Title:

*Method for promoting ovulation parturition and lactation in
mammals*

The following statement is a full description of this invention including
the best method of performing it known to us:-

5


Field of the Invention

The present invention generally relates to a method for promoting ovulation, parturition and lactation in female mammals. More particularly, the present invention is directed to a method of administering a dopamine D₂ receptor antagonist, such as domperidone, to a female mammal for altering hormonal levels in the animal in order to promote ovulation, parturition or lactation.

10


**Background of the Invention**

15



Breeders of various animals, such as horses and other livestock, face many problems in getting the animals to breed properly. For instance, some female mammals fail to ovulate and fall into heat in regular intervals. Mammals that exhibit a prolonged period of inactivity between two periods of heat are described as being anestrus. Anestrus female mammals will not accept the male and are incapable of conceiving offspring.

20



25

Another problem breeders experience is the inability of some mammals to prepare for and give birth after the mammals have become pregnant and the fetus is ready to be born. In preparation for and during the act of giving birth, a process known as parturition, the pregnant mammal should experience cervical relaxation, swelling of the vulva, and relaxation of ligaments around the pelvis. Without these events occurring, the mammal is not capable of giving natural, unassisted birth. Further, should these events not occur, the health of the mother and of the unborn offspring are at grave risk.

30

35

Another problem commonly experienced by breeders

is the inability of mammals to produce and secrete milk after giving birth. The condition of failing to lactate properly after child birth is referred to as agalactia, and is especially prevalent in mares and other livestock. Should the mammal not lactate properly, the offspring must then be bottle fed which is time consuming, labor intensive, and significantly adds to the cost of raising the livestock.

Each of the problems mentioned above can be caused to a great extent by hormonal imbalance or by hormonal irregularities. Hormones released in the body are primarily responsible for initiating ovulation, parturition, and lactation in mammals. Thus, if hormones are not released in the body at particular critical times, the above described problems can be experienced.

For instance, hormones can be prevented from being released in the body by various chemical agents. One such known chemical agent is dopamine. Dopamine is a decarboxylated form of dopa and is found and produced in the adrenal glands of mammals. Dopamine is believed to be produced by the body when it is necessary to suppress hormone secretion. Dopamine suppresses hormone release by binding to and tying up receptors on the anterior pituitary, an endocrine gland located at the base of the brain not far from the hypothalamus. By binding to the anterior pituitary, the gland is prevented from receiving a stimulus hormone that causes it to release other hormones such as those necessary for ovulation, parturition, and lactation.

Although dopamine is necessary during particular periods for keeping hormone levels in the body within controlled ranges, excess levels of dopamine can adversely interfere with the process of reproduction.

Also, besides dopamine, there are other chemical agents that can interfere with or prevent hormone secretion, adversely affecting biological processes.

Thus, a need exists for a method of promoting follicular growth and ovulation, parturition, and lactation in female mammals. A need also exists for treating anestrus mammals, agalactic mammals, and mammals that fail to prepare for parturition when a fetus is ready to be born. A further need exists for a chemical agent that antagonizes dopamine and other chemicals that act in a similar manner in order to counteract hormonal imbalance and irregularities.

Summary of the Invention

It is desirable to provide a process for promoting ovulation and for treating anestrus mammals.

It is also desirable to provide a method for promoting parturition in a pregnant mammal that is at the end of the pregnancy cycle.

It is also desirable to provide a process for treating agalactia in mammals that have just given birth.

It is also desirable to provide a process for controlling hormonal release in the body.

It is also desirable to provide a method for altering hormone levels in the body of a mammal by administering to the mammal a dopamine antagonist.

It is also desirable to provide a process for treating a mammal that has excess levels of dopamine and other similar acting agents within its body.

The present invention provides a method for promoting follicular growth and ovulation, for preparing a mammal for parturition, and for promoting lactation by administering to a female mammal a composition. The composition contains a dopamine D₂ receptor antagonist. For instance, in one embodiment, the composition can contain domperidone.



The present inventor has used domperidone in the past for treating animals suffering from fescue toxicosis. For instance, the present inventor's prior work is disclosed in U.S. Patent No. 5,372,818 entitled "Method of Treating Fescue Toxicosis with Domperidone", which is incorporated herein by reference in its entirety. In U.S. Patent No. 5,372,818, it was discovered not only that domperidone is an effective agent for treating fescue toxicosis, but that domperidone does not substantially cross the blood brain barrier.



Therefore, domperidone can be administered to animals while avoiding substantial adverse behavioral and neurological side effects. Such neuroleptic side effects have been observed in animals exposed to other D_2 receptor antagonists such as the drugs, metoclopramide and sulpiride.

5 Although U.S. Patent No. 5,372,818 provides great advances in the art for treating animals infected with fescue toxicosis, various advantages, aspects and features of the present invention remain absent from the present inventor's prior patent.

10 According to the present invention, ovulation, parturition or lactation can be promoted by administering to a mammal a composition containing a



dopamine D₂ receptor antagonist, such as domperidone. The D₂ receptor antagonist can be administered to the mammal in an amount from about 0.08 mg to about 3.3 mg per kilogram of body weight of the mammal. The D₂ receptor antagonist can be administered to the mammal either orally or subcutaneously.

Specifically, when using domperidone to promote follicular growth and ovulation, domperidone can be orally administered at a concentration of from about 0.2 mg to about 3.3 mg per kilogram weight of said mammal. In one preferred embodiment, the dosage of domperidone is about 0.55 mg per kilogram weight of the mammal.

When administered subcutaneously, the dosage can be from about 0.08 mg to about 1.32 mg and particularly around 0.22 mg per kilogram weight of the mammal.

When attempting to promote parturition, udder development, and lactation, on the other hand, domperidone can be administered orally to a mammal at a dosage of from about 0.2 mg to about 3.3 mg and particularly at about 1.1 mg per kilogram weight of the mammal. When administered subcutaneously, the dosage of domperidone can be from about 0.08 mg to about 1.32 mg and particularly at about 0.44 mg per kilogram weight of the mammal.

All of the above dosage levels according to the present invention can be given daily. Although unknown, it is believed that the treatments cause hormonal levels in the mammal to change for promoting either ovulation, parturition or lactation. Unexpectedly, it has also been discovered that a dopamine D₂ receptor antagonist such as domperidone may also cause an increase in feed intake in the mammal resulting in weight increases.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

Other features and aspects of the present invention are discussed in greater detail below.

Detailed Description of Preferred Embodiments

It is to be understood by one of ordinary skill in the art that the present discussion is a description of exemplary embodiments only, and is not intended as limiting the broader aspects of the present invention, which broader aspects are embodied in the exemplary construction.

The present invention is generally directed to various processes and methods for promoting and stimulating follicular growth and ovulation, for preparing a mammal for parturition, and for promoting and stimulating udder development and lactation. The process of the present invention can be used to treat anestrus mammals, agalactic mammals, and more generally any mammal experiencing problems during the reproductive cycle. The process of the present invention can also be used to manipulate and control ovulation and lactation. For instance, in mammals that only ovulate seasonally, the present invention can be used to promote ovulation at times when ovulation would not normally occur.

In general, the advantages of the present invention are achieved by administering to a female mammal a composition containing a dopamine D₂ receptor antagonist. A D₂ receptor antagonist should be chosen but does not substantially cross the blood brain barrier in order to ensure that the mammal does not suffer any adverse behavioral or neurological side effects. For instance, in a preferred embodiment, the dopamine D₂ receptor antagonist is domperidone.

Although unknown, it is believed that domperidone ties up receptor cells in the anterior pituitary and



other places in the body preventing various chemical agents, such as dopamine, from binding to the receptor cells and altering the release of hormones throughout the body. By altering the effects of dopamine and
5 other similar agents, domperidone allows the anterior pituitary and other glands of the body to secrete hormones necessary for ovulation, parturition, and lactation as will be described in greater detail hereinafter.

As discussed above, one embodiment of the present invention is directed to a process for stimulating and promoting follicular growth and ovulation. The process can be used to treat anestrus mammals, to cause a mammal to ovulate out of season, or to
15 otherwise control when ovulation occurs.

The process of the present invention can be used to treat any mammal, but is particularly applicable to livestock. For instance, many problems have been experienced in attempting to get horses to breed. Further, mares only ovulate and fall into heat during particular times of the year. The process of the present invention cannot only be used to treat horses that experience problems ovulating but can also be used to breed horses during times of the year when it
20 was not before possible.

According to the present invention, follicular growth and ovulation is promoted by administering to a mammal a dopamine D_2 receptor antagonist. In one preferred embodiment, the dopamine D_2 receptor
30 antagonist is domperidone. The D_2 receptor antagonist can be taken orally by the mammal or can be injected subcutaneously.

In one embodiment, when using domperidone, it has been found that ovulation is promoted when the
35 domperidone is taken orally by a mammal in an amount

from about 0.2 mg/kg (mg of domperidone per kg of body weight) to about 3.3 mg/kg. At concentrations greater than 3.3 mg/kg, no further benefits have been observed. Thus far, the best results have been obtained when domperidone has been taken orally at a concentration of about 0.55 mg/kg.

If the domperidone is injected subcutaneously into the mammal, the dosages listed above can be reduced to about 40% of the oral dose. Thus, if injected into a mammal, domperidone can be administered at a concentration of from about 0.08 mg/kg to about 1.32 mg/kg, with a preferred concentration of about 0.22 mg/kg.

Ovulation is the process by which an egg is discharged from a follicle of an ovary and is released into the fallopian tube and the uterus. More particularly, in mammals, as eggs mature in the ovary, the eggs are surrounded by a layer of follicular cells creating a fluid filled follicle. As the egg reaches maturity, the fluid filled follicle bulges from the surface of the ovary. Stimulated by a hormone, ovulation occurs when the follicle ruptures, releasing the egg.

Ovarian cycles are initiated and controlled by a variety of hormones secreted throughout the body. Specifically, these hormones include follicle stimulating hormone (FSH) which is secreted by the pituitary gland and which stimulates follicular and egg growth. Luteinizing hormone (LH), which is also released from the pituitary gland, stimulates the follicle to secrete estrogen which causes the walls of the uterine to thicken. LH also triggers ovulation causing the follicle to rupture.

The levels and frequency of release of these hormones can be altered in the body by various

chemical agents such as dopamine. Although unknown, it is believed that the D₂ receptor antagonists of the present invention, such as domperidone, neutralize the effects of dopamine and other related chemical agents.

5 By administering to a mammal a D₂ receptor antagonist, LH and FSH can be released by the body without inhibition causing ovulation to occur.

Besides promoting ovulation, the present invention is also directed to a process for preparing a mammal for parturition. This process is for treating mammals and particularly livestock that have problems giving birth once pregnant.

Parturition is stimulated in a mammal according to the present invention by administering to the mammal a composition containing a dopamine D₂ receptor antagonist. For instance, in one embodiment, the composition can contain domperidone. Once administered to the mammal, it is believed that the composition neutralizes the effects of dopamine and other similar chemical agents from altering the release of hormones that prepare a mammal for giving birth. Thus, the composition of the present invention allows the body to secrete hormones that may affect cervical dilation and general broodiness.

25 By administering the composition to pregnant mammals, the mammal is more likely to give birth on time. The composition also promotes natural births and prevents against having to deliver the offspring by C-sections. The treatment not only protects the unborn offspring during delivery but also protects the pregnant mother from being harmed during birth.

30 In order to promote normal parturition, in one embodiment, a pregnant mammal can be orally fed domperidone at a concentration of from about 0.2 mg/kg of body weight to about 3.3 mg/kg. In one preferred

35

embodiment, the oral dosage can be about 1.1 mg/kg.

When injected subcutaneously into the mammal, domperidone has been found to produce effective results at concentrations of from about 0.08 mg/kg to about 1.32 mg/kg, with best results being obtained at about 0.44 mg/kg. Amounts greater than 1.32 mg/kg can be administered to the animal without any adverse side effects. Additional benefits, however, have not been observed at higher dosage levels.

Whether administered orally or subcutaneously, treatments of the D₂ receptor antagonist administered to the mammal can begin at any time during the pregnancy. For larger livestock such as cattle and horses, the treatment should begin about fifteen to twenty days prior to the expected birth date.

A further embodiment of the present invention is directed to a process for promoting udder development and lactation in female mammals. The process can be used to prevent or treat agalactia or any other ailments regarding the low level or non-production of milk. For instance, lactation problems have been particularly observed in mares that have recently given birth. After giving birth, many mares either fail to produce milk or do not produce enough milk to adequately nurture the foal. When this occurs, the foal must be bottle fed or fed in some other manner which significantly adds to the expense of raising the livestock.

The process of the present invention can be used in these circumstances to stimulate or increase milk production in non-milking mares and other mammals. Further, the process can be used simply to increase milk production without any adverse effects. For instance, the process of the present invention can also be used to increase the quantity of milk obtained

from cows for human consumption.

Lactation is stimulated and promoted according to the present invention by administering to mammals a composition containing a dopamine D₂ receptor antagonist. For instance, the composition can contain domperidone in an amount sufficient to stimulate milk production.

In one embodiment, milk production was stimulated in horses by orally administering to the horses domperidone in a concentration of from about 0.2 mg/kg of body weight to about 3.3 mg/kg of body weight, and particularly at a concentration of about 1.1 mg/kg. At oral dosages greater than 3.3 mg/kg, no further beneficial results were observed.

Domperidone can also be administered to the animal subcutaneously. If injected into the mammal, about 40% of the dosage amounts given above can be used. Thus, when injected subcutaneously, the domperidone concentration can be from about 0.08 mg/kg to about 1.32 mg/kg.

Although unknown, it is believed that the dopamine D₂ receptor antagonist, such as domperidone, influences hormonal levels within the animal. Domperidone is believed to alter hormone levels by neutralizing the effect of dopamine-like agents that prevent particular hormones from being secreted. With respect to lactation, it is believed that the dopamine D₂ receptor antagonist increases the levels of progesterone, estrogen and prolactin within the body of the mammal. These hormones are believed to be primarily responsible for promoting udder development and lactation.

Besides promoting ovulation, preparing a mammal for parturition, and stimulating lactation, it has also been unexpectedly discovered that the composition

of the present invention can cause a mammal to increase its feed intake and thus gain weight faster. This process is particularly beneficial in raising livestock for human consumption.

5 In order to increase feed intake according to the present invention, a dopamine D₂ receptor antagonist, such as domperidone, is administered to a mammal. The treatment can be given to the mammal orally or subcutaneously. If domperidone is used and
10 administered orally, the domperidone should be fed to the mammal at a concentration of from about 0.2 mg/kg of body weight to about 3.3 mg/kg of body weight and particularly at about 1.1 mg/kg. If domperidone is injected into the mammal, the dosage should be from
15 about 0.08 mg/kg to about 1.32 mg/kg.

At the present time, it is unknown why domperidone causes increases in feed intake. It is believed, however, that domperidone may stimulate feed intake by increasing levels of the hormone prolactin
20 in the body of the mammal. It may also be possible that domperidone causes levels of hormones to be altered that firm up the gut of the mammal. If the muscles of the gut were to be stimulated, then food passage may increase in the gut allowing the animal to
25 eat more.

In general, in delivering an effective dosage of a dopamine D₂ receptor antagonist to a mammal according to the present invention, various vehicles may be used. For instance, when taken orally, the D₂
30 receptor antagonist may be combined with a feed or feed supplement material as the carrier. If injected, the drug may be mixed with any suitable carrier. Additionally, the D₂ receptor antagonist, such as domperidone, may be added to salt blocks or mineral
35 blocks during casting or mixed directly into feed.

Further, various other administration techniques well known in the art may be employed. It is to be understood that the present invention is not to be limited to any particular vehicle.

5 It also should be appreciated that although the above description and following examples relate primarily to livestock such as horses and cattle, it is believed that the drug will work as described with any mammal. For instance, although untested at the present time, it is believed that a dopamine D₂ receptor antagonist that does not substantially cross the blood brain barrier such as domperidone may be administered to humans for promoting ovulation, facilitating childbirth, or stimulating lactation.

15 The present invention may be better understood with reference to the following examples.

EXAMPLE NO. 1

The following tests were performed in order to demonstrate the ability of domperidone to stimulate and promote ovulation in mares.

20 Ten mares were fed a composition containing domperidone at a concentration of 0.55 mg/kg of body weight. The composition was administered daily. Specifically, the domperidone was mixed with molasses and fed to the mares with a 5 cc syringe. Nine additional mares were used as a control. These mares were fed the molasses carrier not containing any domperidone.

25 Of importance, the treatments were carried out in January and February during a time when mares typically do not ovulate. The following results were obtained:

TABLE 1
Effect of Domperidone on
Follicular Growth and Ovulation

Test No.	Treated with Domperidone at .55 mg/kg	No. of Days of Treatment	Date of Ovulation	No. of Days of Treatment Before Ovulation
1	No	45	d/n ovulate	d/n ovulate
2	No	45	d/n ovulate	d/n ovulate
3	No	45	d/n ovulate	d/n ovulate
4	No	25	Feb. 26	25
5	No	45	d/n ovulate	d/n ovulate
6	No	45	d/n ovulate	d/n ovulate
7	No	45	d/n ovulate	d/n ovulate
8	No	37	Mar. 10	37
9	No	45	d/n ovulate	d/n ovulate
10	Yes	18	Feb. 9	18
11	Yes	15	Feb. 6	18
12	Yes	21	Feb. 12	21
13	Yes	17	Feb. 8	17
14	Yes	24	Feb. 15	24
15	Yes	14	Feb. 5	14
16	Yes	16	Feb. 7	16
17	Yes	22	Feb. 13	22
18	Yes	45	d/n ovulate	d/n ovulate
19	Yes	19	Feb. 10	19

As shown above, only two of the control mares ovulated. In comparison, nine of the ten mares treated with domperidone ovulated. Also of significance, the treated mares ovulated during the

early part of February. No neurological side effects were observed in any of the mares treated with domperidone during the study.

EXAMPLE NO. 2

5 The following tests were performed to demonstrate the ability of domperidone to prepare a mammal for parturition and to stimulate udder development and lactation.

10 Twelve known agalactic mares who were either pregnant or had just given birth were treated with domperidone. The domperidone was administered orally at a dosage of 1.1 mg/kg of body weight per day. All of the mares treated were not suffering from fescue toxicosis. The following results were obtained:

0010374009

TABLE 2
Effect of Domperidone on Preparation for Parturition, Udder Development and Lactation in Mares

Test No.	Days Before (-) or After Expected Foaling Dosing Was Initiated	Days Before (-) or After Expected Foaling that Mare Foaled	Mare's Udder Development Before Treatment Began	Mare's Udder Development After Treatment Began	Was Mare Lactating Normally After Foaling?	Did Mare Have A Live Foal?
1	+5	+12	Very Little	Dripping Milk 5th Day of Treatment	Yes	Yes
2	-28	-1	Very Little	Increased Weekly	Yes	Yes
3	No data	+8	No Development	After 2 Days, Made Udder Sack	Yes	Yes
4	+6	+15	Very Little	Developed Slowly But Natural	Yes	Yes
5	+4	+14	Virtually No Udder	Developed Adequately	No, but drug helped some	Yes
6	-21	0	Normal	Normal	Yes	Yes
7	+5	No data	Incomplete Small Udder	Udder Developed As Soon As Treatment Started	Yes	Yes
8	-24	-3	No Development	Very Slight Development	Yes	Yes
9	No data	+3	Minimal	Gradual Development	Yes	Yes
10	-13	No data	Slight Base, No Filling of Teat	Normal Development	Yes	Yes
11	-20	-3	No Development	No Response	Yes	Yes
12	-10	-2	None to Minimal	Immediate Development After Treatment	Yes	Yes

EXAMPLE NO. 3

It has also been unexpectedly discovered that treating a mammal with a D₂ receptor antagonist, such as domperidone, increases feed intake. The following results demonstrate this phenomenon.

Ten quarter horse mares were housed in individual pens and fed a standard feed concentrate weighing 0.5% of each mares' initial body weight at approximately 8:00 A.M. each day during the study. In addition, hay was fed to the mare ad libitum each day. During the night, the mares were placed in a dry lot having no access to food.

After seven days on the above feed schedule (control), the mares were orally fed at 8:00 A.M. each day domperidone at a dosage of 1.1 mg/kg of body weight. The domperidone was fed to the mares in a molasses carrier. During treatment with domperidone, the mares were fed the concentrate feed and hay in the same manner as during the control period. Domperidone was fed to the mares for two weeks. The following results were obtained:

TABLE 3
Effect of Domperidone On
The Daily Feed Intake of Mares

Mare ID	Initial Body Weight (lbs)	Concentrate Feed (lbs)	Avg. Hay Consumption Before Treatment (lbs)	Avg. Hay Consumption After Treatment (lbs)
1	1025	5.1	11.2	13.0
2	980	4.9	10.3	12.2
3	1110	5.6	12.2	14.1
4	900	4.5	11.1	13.2
5	1050	5.3	12.0	14.1
6	1000	5.0	11.0	14.2
7	980	4.9	9.8	12.3
8	1010	5.1	11.0	13.5
9	1100	5.5	12.5	15.0
10	950	4.8	10.2	12.1

As shown above, daily hay consumption increased for each of the ten mares after treatment with domperidone began.

These and other modifications and variations to the present invention may be practiced by those of ordinary skill in the art, without departing from the spirit and scope of the present invention, which is more particularly set forth in the appended claims. In addition, it should be understood that aspects of the various embodiments may be interchanged both in whole or in part. Furthermore, those of ordinary skill in the art will appreciate that the foregoing description is by way of example only, and is not intended to limit the invention so further described



in such appended claims.

- Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken
- 5 as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed in Australia before the priority date of each claim of this application.



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method for promoting follicular growth and ovulation in mammals comprising the step of:
administering to a female mammal a composition comprising a
5 dopamine D₂ receptor antagonist that does not substantially cross the blood brain barrier, said dopamine D₂ receptor antagonist being administered to said female mammal in an amount sufficient to promote follicular growth and ovulation.
2. A method as defined in claim 1, wherein said dopamine D₂ receptor
10 antagonist comprises domperidone.
3. A method as defined in claim 1 or claim 2, wherein said dopamine D₂ receptor antagonist is administered to said female mammal in an amount from 0.08 mg to 3.3 mg per kilogram weight of said mammal.
4. A method as defined in any one of claims 1 to 3, wherein said
15 composition is administered orally.
5. A method as defined in any one of claims 1 to 3, wherein said composition is administered subcutaneously.
6. A method as defined in claim 2, wherein said domperidone is
20 administered to said female mammal orally in an amount from 0.2 mg to 3.3 mg per kilogram weight of said mammal.
7. A method as defined in claim 2, wherein said domperidone is administered to said female mammal subcutaneously in an amount from 0.08 mg to 1.32 mg per kilogram weight of said mammal.
8. A method as defined in any one of claims 1 to 7, wherein said female
25 mammal is an animal selected from the group consisting of a mare, or a cow.
9. A method for promoting parturition in mammals comprising the step of:
administering to a female mammal a composition comprising a
30 dopamine D₂ receptor antagonist that does not substantially cross the blood brain barrier, said dopamine D₂ receptor antagonist being administered to said pregnant mammal in an amount sufficient to promote parturition.
10. A method as defined in claim 9, wherein said dopamine D₂ receptor antagonist comprises domperidone.
11. A method as defined in claim 9 or claim 10, wherein said dopamine D₂
35 receptor antagonist is administered to said pregnant mammal in an amount from 0.08 mg to 3.3 mg per kilogram weight of said mammal.



12. A method as defined in any one of claims 9 to 11, wherein said composition is administered orally.
13. A method as defined in any one of claims 9 to 11, wherein said composition is administered subcutaneously.
- 5 14. A method as defined in claim 10, wherein said composition is administered orally to said pregnant mammal in an amount from 0.2 mg to 3.3 mg per kilogram weight of said mammal.
15. A method as defined in claim 10, wherein said composition is administered subcutaneously to said pregnant mammal in an amount from
- 10 0.08 mg to 1.32 mg per kilogram weight of said mammal.
16. A method as defined in any one of claims 9 to 15, wherein said pregnant mammal is an animal selected from the group consisting of a mare or a cow.
17. A method for adjusting hormonal levels in female mammals in order to
- 15 promote ovulation or parturition comprising the step of:
 administering to a female mammal a composition comprising domperidone, said domperidone being administered to said female mammal in an amount from 0.08 mg to 3.3 mg per kilogram weight of said female mammal.

20

Dated this fourth day of January 2001

Clemson University
 Patent Attorneys for the Applicant:

F B RICE & CO

