




(86) Date de dépôt PCT/PCT Filing Date: 2014/10/28
(87) Date publication PCT/PCT Publication Date: 2015/05/07
(45) Date de délivrance/Issue Date: 2019/04/02
(85) Entrée phase nationale/National Entry: 2016/04/25
(86) N° demande PCT/PCT Application No.: JP 2014/078644
(87) N° publication PCT/PCT Publication No.: 2015/064580
(30) Priorité/Priority: 2013/10/29 (JP2013-224694)

(51) Cl.Int./Int.Cl. *C07D 239/54* (2006.01),
A61K 31/513 (2006.01), *A61K 31/517* (2006.01),
A61K 31/519 (2006.01), *A61P 1/04* (2006.01),
A61P 11/06 (2006.01), *A61P 17/06* (2006.01),
A61P 19/02 (2006.01), *A61P 25/00* (2006.01),
A61P 29/00 (2006.01), *A61P 3/10* (2006.01),
A61P 35/00 (2006.01), ...

(72) Inventeurs/Inventors:
UENO, HIROKAZU, JP;
YAMAMOTO, TAKASHI, JP;
MIYAZAWA, TOMOKO, JP;
SHINKAI, KENJI, JP;
...

(54) Titre : DERIVE DE SULFONAMIDE ET SON UTILISATION MEDICINALE
(54) Title: SULFONAMIDE DERIVATIVE AND PHARMACEUTICAL USE THEREOF



(1)

(57) Abrégé/Abstract:

Provided is a sulfonamide derivative represented by the following general formula (1) and having an $\alpha 4$ integrin inhibitory effect with high selectivity with a low effect on $\alpha 4\beta 1$ and a high effect on $\alpha 4\beta 7$, or a pharmaceutically acceptable salt thereof (in the general

(51) Cl.Int./Int.Cl. (suite/continued) *A61P 35/04* (2006.01), *A61P 37/00* (2006.01), *A61P 37/06* (2006.01), *A61P 37/08* (2006.01), *A61P 43/00* (2006.01), *A61P 9/00* (2006.01), *A61P 9/10* (2006.01), *C07D 471/04* (2006.01), *C07D 487/04* (2006.01)

(72) Inventeurs(suite)/Inventors(continued): ARISAKA, HARUMI, JP; TAKANOHASHI, TOSHIYUKI, JP

(73) Propriétaires(suite)/Owners(continued): AJINOMOTO CO., INC., JP

(74) Agent: NORTON ROSE FULBRIGHT CANADA LLP/S.E.N.C.R.L., S.R.L.

(57) Abrégé(suite)/Abstract(continued):

formula (1), A, B, D, E, R₄₁, and a to h are as described in the description).

Abstract

Provided is a sulfonamide derivative represented by the following general formula (1) and having an α -4 integrin inhibitory effect with high selectivity with a low effect on α 4 β 1 and a high effect on α 4 β 7, or a pharmaceutically acceptable salt thereof (in the general formula (1), A, B, D, E, R₄₁, and a to h are as described in the description).

Specification

Title of Invention: SULFONAMIDE DERIVATIVE AND
PHARMACEUTICAL USE THEREOF

5

Technical Field

[0001]



The present invention relates to a sulfonamide derivative or a
pharmaceutically acceptable salt thereof, and a pharmaceutical composition
10 containing such a compound as an active ingredient. In particular, the
present invention relates to a compound that is useful as a therapeutic
medicine or a preventive medicine for inflammatory disease in which an $\alpha 4$
integrin-dependent adhesion process is involved in pathological conditions.

15

Background Art

[0002]


Orally available compounds having an $\alpha 4$ integrin inhibitory effect
that are useful as a therapeutic medicine or a preventive medicine for
inflammatory disease in which an $\alpha 4$ integrin-dependent adhesion process is
20 involved in pathological conditions have already been known. For example,
Patent Literature 1 discloses a phenylalanine derivative represented by the
general formula (1) or a pharmaceutically acceptable salt thereof, and a
typical compound thereof has the following chemical structure.



[0003]

Further, Patent Literature 1 shows the results of VCAM inhibitory activity (VCAM-1/ α 4 β 1 binding assay) and (VCAM-1/ α 4 β 7 binding assay).

Further, Patent Literature 2 also discloses a phenylalanine derivative represented by the general formula (1) below and having R₁₂ (R₁₃) N-X₁- group at an end or a pharmaceutically acceptable salt thereof.



It is shown that this compound has high VCAM-1/ α 4 β 1 integrin inhibitory activity in the presence of serum as compared with the compound of Example 1 of Patent Literature 1. Further, Patent Literatures 3 and 4 also disclose compounds having an α 4 integrin inhibitory effect.

Citation List

Patent Literature

[0004]

Patent Literature 1: WO 02/016329 A1

5 Patent Literature 2: WO 05/061466 A1

Patent Literature 3: WO 03/070709 A1

Patent Literature 4: WO 01/042225 A1

Summary of Invention

10 [0005]

The present invention aims to provide a new compound having a chemical structural formula that is unknown so far and having excellent $\alpha 4$ integrin inhibitory effect.

15 In particular, the present invention aims to provide a new compound having an $\alpha 4$ integrin inhibitory effect with high selectivity with a low effect on $\alpha 4\beta 1$ and a high effect on $\alpha 4\beta 7$.

The present invention also aims to provide an orally available compound having excellent $\alpha 4$ integrin inhibitory effect.

20 The present invention also aims to provide a pharmaceutical composition containing such a new compound as described above and a pharmaceutically acceptable carrier.

The present invention also aims to provide a medicine containing such a new compound as described above.

25 The present invention also aims to provide a therapeutic agent or a preventive agent for inflammatory disease in which an $\alpha 4\beta 7$ integrin-dependent adhesion process is involved in pathological conditions.

The present invention also aims to provide an $\alpha 4$ integrin inhibitor.


[0006]

30 The present invention has been accomplished based on the finding that a sulfonamide derivative having a specific chemical structure and

containing, at an end, a sulfonamide group having, as a substituent, a hetero
 ring group containing a heteroatom as a constituent element or a phenyl
 group, or a pharmaceutically acceptable salt thereof has excellent α 4
 integrin inhibitory activity, and use of such a compound can solve the
 5 above-described problems.

That is, the present invention includes the following aspects [1] to
 [17].

[1] A sulfonamide derivative represented by the general formula (1) below or
 a pharmaceutically acceptable salt thereof:




(1)


10

wherein


A represents a group represented by the general formula (2-1), (2-2), or (2-3)
 below:




(2-1)



(2-2)



(2-3)



(2-4)

[0007]

wherein

Arm is a cyclic alkyl group or an aromatic ring containing 0, 1, 2, 3, or 4
 5 heteroatoms selected from oxygen atoms, sulfur atoms, and nitrogen atoms,
 R1, R11, R21, R51, and R52 each independently represent any one of a
 hydrogen atom, a halogen atom, a hydroxyl group, a lower alkyl group, a
 lower alkenyl group, a lower alkynyl group, a lower alkoxy group, a lower
 alkylthio group, a hydroxy-lower alkyl group, a hydroxy-lower alkenyl group,
 10 a hydroxy-lower alkoxy group, a halogeno-lower alkyl group, a
 halogeno-lower alkoxy group, a halogeno-lower alkylthio group, a
 halogeno-lower alkenyl group, a nitro group, a cyano group, an amino group,
 a lower alkylamino group, a lower alkylamino-lower alkyl group, a carboxyl
 group, a lower alkyloxycarbonyl group, a carbamoyl group, a lower alkanoyl
 15 group, an aroyl group, a lower alkylsulfonyl group, a sulfamoyl group, and an
 ammonium group,

R12, R13, and R14 each independently represent any one of a hydrogen atom, a halogen atom, a hydroxyl group, a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a lower alkoxy group, a lower alkylthio group, a hydroxy-lower alkyl group, a hydroxy-lower alkenyl group, a hydroxy-lower alkoxy group, a halogeno-lower alkyl group, a halogeno-lower alkoxy group, a halogeno-lower alkylthio group, a halogeno-lower alkenyl group, a nitro group, a cyano group, an amino group, a lower alkylamino group, a lower alkoxy-lower alkylene group, a lower alkylthio-lower alkylene group, a lower alkylamino-lower alkylene group, a lower alkylamino-lower alkyl group, a carboxyl group, a lower alkyloxycarbonyl group, a carbamoyl group, a lower alkanoyl group, an aroyl group, a lower alkylsulfonyl group, a sulfamoyl group, and an ammonium group, and

R2, R3, R22, R23, R24, R25, and R53 each independently represent any one of a hydrogen atom, a halogen atom, a hydroxyl group, a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a lower alkoxy group, a lower alkylthio group, a hydroxy-lower alkyl group, a hydroxy-lower alkenyl group, a hydroxy-lower alkoxy group, a halogeno-lower alkyl group, a halogeno-lower alkoxy group, a halogeno-lower alkylthio group, a halogeno-lower alkenyl group, a nitro group, a cyano group, an amino group, a lower alkylamino group, a lower alkylamino-lower alkyl group, a carboxyl group, a lower alkyloxycarbonyl group, a carbamoyl group, a lower alkanoyl group, an aroyl group, a lower alkylsulfonyl group, a sulfamoyl group, and an ammonium group,

B represents any one of an alkoxy group having 1 to 10 carbon atoms, a hydroxyl group, or a hydroxyamino group, these groups being optionally substituted with a substituent selected from the group consisting of an aryl group, a hydroxyl group, an alkyl group having 1 to 10 carbon atoms, a lower alkoxy group, a lower alkylamino group, a halogen atom, and a heterocyclic group,

[0008]

R41 represents a hydrogen atom or a lower alkyl group,

a, b, c, and d independently represent C-R31, C-R32, C-R33, or C-R34, respectively, wherein one or two of a, b, c, and d may represent a nitrogen atom,

- 5 R31, R32, R33, and R34 each independently represent any one of a hydrogen atom, a halogen atom, a lower alkyl group, a lower alkoxy group, and a nitro group, wherein any one of R31, R32, R33, and R34 is a halogen atom or a lower alkyl group,

- e, f, g, and h independently represent C-R35, C-R36, C-R37, or C-R38,
10 respectively, wherein one or two of e, f, g, and h may represent a nitrogen atom,

R35, R36, R37, and R38 each independently represent any one of a hydrogen atom, a halogen atom, a lower alkyl group, a lower alkoxy group, and a nitro group,

- 15 D represents a phenyl group or a heterocyclic group, optionally having a substituent selected from the group consisting of a halogen atom, a hydroxyl group, a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a lower alkoxy group, a lower alkylthio group, a hydroxy-lower alkyl group, a hydroxy-lower alkenyl group, a hydroxy-lower alkoxy group, a
20 halogeno-lower alkyl group, a halogeno-lower alkoxy group, a halogeno-lower alkylthio group, a halogeno-lower alkenyl group, a nitro group, a cyano group, an amino group, a carboxyl group, a lower alkyloxycarbonyl group, a carbamoyl group, a lower alkanoyl group, an aroyl group, a lower alkylsulfonyl group, a sulfamoyl group, a lower alkoxy-lower alkylene group,
25 a lower alkylthio-lower alkylene group, a lower alkylamino-lower alkylene group, and an ammonium group,

[0009]

- when A represents a group represented by the general formula (2-1), (2-2), or (2-3), E represents: a 5- or 6-membered heterocyclic group substituted with a
30 substituent selected from the group consisting of a 3- to 8-membered

saturated ring group containing a nitrogen atom connected by a carbon atom, a 3- to 8-membered saturated or unsaturated ring group containing a heteroatom selected from an oxygen atom and a sulfur atom, a lower alkoxy-lower alkylene group, a lower alkylthio-lower alkylene group, a lower alkylamino-lower alkylene group, a lower alkylamino group, and a lower alkenylamino group; or a 5- or 6-membered cyclic ketone group containing 1 to 4 nitrogen atoms as atoms constituting the ring, the cyclic ketone group being substituted with a lower alkyl group or a lower alkenyl group, and when A represents a group represented by the general formula (2-4), E represents: a phenyl group or a 5- or 6-membered heterocyclic group optionally having a substituent selected from the group consisting of a halogen atom, a hydroxyl group, a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a lower alkoxy group, a lower alkylthio group, a hydroxy-lower alkyl group, a hydroxy-lower alkenyl group, a hydroxy-lower alkoxy group, a halogeno-lower alkyl group, a halogeno-lower alkoxy group, a halogeno-lower alkylthio group, a halogeno-lower alkenyl group, a nitro group, a cyano group, an amino group, a 4- to 6-membered cyclic amino group, a carboxyl group, a lower alkyloxycarbonyl group, a carbamoyl group, a lower alkanoyl group, an aroyl group, a lower alkylsulfonyl group, a sulfamoyl group, and an ammonium group; an aminocarbonyl group optionally having a substituent selected from the group consisting of a hydroxyl group, a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a lower alkoxy group, a hydroxy-lower alkyl group, a hydroxy-lower alkenyl group, a hydroxy-lower alkoxy group, a halogeno-lower alkyl group, a halogeno-lower alkoxy group, a halogeno-lower alkenyl group, an amino group, a lower alkylamino group, an aryl group, a heterocyclic group, a heterocyclic lower alkyl group, a lower alkylsulfonyl group, and a sulfamoyl group; a hydrogen atom; a halogen atom; a hydroxyl group; a lower alkyl group; a lower alkenyl group; a lower alkynyl group; a lower alkoxy group; a lower alkylthio group; a hydroxy-lower alkyl group; a hydroxy-lower alkenyl


group; a hydroxy-lower alkoxy group; a halogeno-lower alkyl group; a halogeno-lower alkoxy group; a halogeno-lower alkylthio group; a halogeno-lower alkenyl group; a nitro group; a cyano group; an amino group; a carboxyl group; a dihydroxyboryl group; a lower alkylcarbonyl group; a lower alkyloxycarbonyl group; a carbamoyl group; a lower alkanoyl group; an aroyl group; a lower alkylsulfonyl group; a sulfamoyl group; an ammonium group; a lower alkylaminoalkylene group; a 5- or 6-membered heterocyclic group substituted with a substituent selected from the group consisting of a 3- to 8-membered saturated ring group containing a nitrogen atom connected by a carbon atom, a 3- to 8-membered saturated or unsaturated ring group containing a heteroatom selected from an oxygen atom and a sulfur atom, a lower alkoxy-lower alkylene group, a lower alkylthio-lower alkylene group, a lower alkylamino-lower alkylene group, a lower alkylamino group, and a lower alkenylamino group; or a 5- or 6-membered cyclic ketone group containing 1 to 4 nitrogen atoms as atoms constituting the ring, the cyclic ketone group being substituted with a lower alkyl group or a lower alkenyl group, wherein the lower alkylcarbonyl group and the lower alkyloxycarbonyl group may form a fused ring by being bonded to the phenyl group of D,

provided that the sulfonamide derivative excludes sulfonamide derivatives selected from the group consisting of (a) to (h) below:

[0010]


(a)

(2S)-2-[[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)phenyl]propanoic acid




(b) cyclohexyl

(2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)phenyl]propionate



(c) cyclohexyl

(2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoic acid




[0011]

(d) cyclohexyl


(2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoic acid

-yl)-3-pyridyl]propanoate



(e)


(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
5 amino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)p
henyl]propanoic acid



(f)

cyclohexyl


(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
10 amino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)p
henyl]propanoate



(g)

(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
15 amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-


3-pyridyl]propanoic acid



and

(h) cyclohexyl

- 5 (2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate.



[0012]

- 10 [2] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to [1], wherein D is a phenyl group optionally having a substituent selected from the group consisting of a lower alkyl group, a halogen atom, a hydroxyl group, and a lower alkoxy group.

- [3] The sulfonamide derivative or a pharmaceutically acceptable salt thereof
15 according to [1], wherein D is a 6-membered aromatic heterocyclic group optionally having a substituent selected from the group consisting of a lower alkyl group, a halogen atom, and a lower alkoxy group, and having a nitrogen atom as an atom constituting the ring.

- [4] The sulfonamide derivative or a pharmaceutically acceptable salt thereof
20 according to [4], wherein the heterocyclic group of D is a pyridyl group or a

pyrrole group optionally having a substituent selected from the group consisting of a lower alkyl group, a halogen atom, and a lower alkoxy group.

[5] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [4], wherein the 5- or 6-membered heterocyclic group of E is selected from the group consisting of a pyridyl group, a pyridazyl group, a pyrimidyl group, a pyrazinyl group, a pyrrolyl group, a triazolyl group, and a tetrazole group, and the cyclic ketone group of E is a 5- or 6-membered cyclic ketone group containing 2 or 3 nitrogen atoms as atoms constituting the ring.

[6] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [4], wherein A represents a group represented by the general formula (2-4), and E is a 5- or 6-membered aromatic heterocyclic group containing 1, 2, 3, or 4 heteroatoms selected from oxygen atoms, sulfur atoms, or nitrogen atoms and optionally having a substituent selected from the group consisting of a lower alkyl group, a lower alkoxy group, a lower alkoxy-lower alkylene group, and a halogen atom.

[0013]

[7] The phenylalanine derivative or a pharmaceutically acceptable salt thereof according to [6], wherein the aromatic heterocyclic group is selected from the group consisting of a pyridyl group, a pyridazyl group, a pyrimidyl group, a pyrazinyl group, a pyrrolyl group, a furanyl group, a thiophenyl group, a thiazolyl group, an oxazolyl group, a triazolyl group, and a tetrazole group.

[8] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [4], wherein A represents a group represented by the general formula (2-4), and E is an aminocarbonyl group optionally substituted with a lower alkyl group, a heterocyclic group, or a heterocyclic lower alkyl group, or a lower alkylaminoalkylene group.

[9] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [4], and [5], wherein the heterocyclic group of E

is a pyridyl group, a pyrimidyl group, a triazolyl group, and a tetrazole group optionally having a substituent selected from the group consisting of a 3- to 8-membered saturated or unsaturated ring group containing a heteroatom selected from an oxygen atom and a sulfur atom, a lower alkoxy-lower alkylene group, or a lower alkylamino-lower alkylene group.

[10] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [5] and [9], wherein A represents a group represented by the general formula (2-2), and R11, R12, and R13 are selected from the group consisting of a lower alkyl group, a lower alkoxy group, a lower alkoxy-lower alkylene group, a lower alkylamino-lower alkylene group, and a halogen atom.

[0014]

[11] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [10], wherein B is a lower alkoxy group optionally substituted with a substituent selected from the group consisting of a lower alkyl group, a lower alkoxy group, a lower alkylamino group, a halogen atom, and a heterocyclic group; or a hydroxyl group.

[12] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [6], [8], [9], [10], and [11], wherein the lower alkyl group is a straight-chain, branched-chain, or cyclic alkyl group.

[13] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [12], wherein R1, R11, R21, R51, and R52 are lower alkyl groups.

[14] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [13], wherein Arm is selected from the group consisting of a phenyl group, a pyridyl group, a pyrimidyl group, and an imidazolyl group.

[15] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [4], wherein A represents a group represented by the general formula (2-2), and

E represents a 5- or 6-membered heterocyclic group containing 1 to 4 nitrogen atoms, the 5- or 6-membered heterocyclic group being substituted with a substituent selected from the group consisting of a lower alkyl group, a lower alkoxy-lower alkylene group, and a 5- or 6-membered heterocyclic group containing an oxygen atom; or a cyclic ketone group containing 1 to 4 nitrogen atoms, the cyclic ketone group being substituted with a lower alkyl group or a lower alkenyl group.

[16] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to [15], wherein

10 B is optionally substituted with a substituent selected from the group consisting of an alkyl group having 1 to 10 carbon atoms, a lower alkoxy group, and a heterocyclic group containing 1 to 4 heteroatoms selected from oxygen atoms and nitrogen atoms, and

Arm is a pyridyl group.

15 [17] The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [4], wherein

A represents a group represented by the general formula (2-4),

E is a 5- or 6-membered heterocyclic group containing an oxygen atom, or a 5- or 6-membered heterocyclic group containing 1 to 4 nitrogen atoms, and

20 E optionally has a substituent selected from a lower alkyl group, a lower alkoxy-lower alkylene group, and a 5- or 6-membered heterocyclic group containing an oxygen atom.

[18] A pharmaceutical composition comprising the sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [17].

25 [19] A therapeutic agent or a preventive agent for inflammatory disease in which an $\alpha 4\beta 7$ integrin-dependent adhesion process is involved in pathological conditions, the agent comprising: the sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of [1] to [17] as an active ingredient.

30 [20] An $\alpha 4\beta 7$ integrin inhibitor comprising: the sulfonamide derivative or a

pharmaceutically acceptable salt thereof according to any one of [1] to [17] as an active ingredient.

Description of Embodiments

5 [0015]

The term "lower" of the lower alkyl group or the like in this description means a group having 1 to 6 carbon atoms, preferably a group having 1 to 4 carbon atoms. An alkyl group, an alkenyl group, and an alkynyl group serving as a component of an alkyl group, an alkenyl group, an alkynyl group, an alkoxy group, an alkylthio group, an alkanoyl group, an alkylamino group, or the like can be straight chain or branched. Alternatively, they may be cyclic. Examples of the alkyl group include a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a secondary butyl group, a tertiary butyl group, a pentyl group, a 2-
10 pentyl group, a 3-pentyl group, a hexyl group, a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, a cyclopropylmethyl group, and a cyclopropylethyl group, preferably having 1 to 6 carbon atoms, more preferably having 1 to 4 carbon atoms. Examples of the alkenyl group include a vinyl group, a propenyl group, a butenyl group, and a pentenyl group, preferably having 2 to 6 carbon atoms, more preferably having 2 to 4 carbon atoms. Examples of the alkynyl group include an ethynyl group, a propynyl group, and a butynyl group, preferably having 2 to 6 carbon atoms, more preferably having 2 to 4 carbon atoms.

Examples of the alkoxy group include a methoxy group, an ethoxy
25 group, a propyloxy group, and an isopropyloxy group, preferably having 1 to 6 carbon atoms, more preferably having 1 to 4 carbon atoms. Examples of the heteroatom include nitrogen, oxygen, and sulfur. Examples of the halogen atom include fluorine, chlorine, bromine, and iodine. Examples of the halogeno-alkyl group include a chloromethyl group, a trichloromethyl
30 group, a trifluoromethyl group, a trifluoroethyl group, and a

pentafluoromethyl group. Examples of the halogeno-alkoxy group include a trichloromethoxy group, and a trifluoromethoxy group. Examples of the hydroxyalkyl group include a hydroxymethyl group, and a hydroxyethyl group.

5 [0016]

In this description, the aryl group means a substituted or unsubstituted aryl group, and examples thereof include a phenyl group, a 1-naphthyl group, and a 2-naphthyl group, in which a phenyl group and a substituted phenyl group are preferable, where a halogen atom, an alkoxy group, an alkyl group, a hydroxyl group, a halogeno-alkyl group, and a halogeno-alkoxy group are particularly preferable as a substituent. The heterocyclic group means a 4- to 7-membered hetero ring group that consists of one to three rings and that is composed of carbon and nitrogen, oxygen, sulfur, or the like, and specific examples thereof include a pyridyl group, a dihydropyranyl group, a tetrahydropyranyl group, a pyridazyl group, a pyrimidyl group, a pyrazyl group, a pyrrolyl group, a furyl group, a thienyl group, a thienyl group, a thiazolyl group, a triazolyl group, an isoxazolyl group, an isothiazolyl group, an oxazolyl group, an indolyl group, a quinolyl group, an isoquinolyl group, a benzoimidazolyl group, a pyrazolyl group, an imidazolyl group, a thiadiazolyl group, a pyrrolidyl group, a piperidyl group, a piperazyl group, a morpholyl group, an oxetanyl ring, an isoindolyl group, a benzofuryl group, an isobenzofuryl group, a benzothienyl group, a benzopyrazolyl group, a benzoimidazolyl group, a benzoxazolyl group, and a benzothiazolyl group, in which a pyridyl group, a pyrazyl group, a pyrimidyl group, a furyl group, and a thienyl group are preferable.

25

[0017]

In the general formula (1) of the present invention, it is preferable that (i) D be a phenyl group optionally having a substituent selected from the group consisting of a lower alkyl group, a halogen atom, a hydroxy group, and a lower alkoxy group, or (ii) D be a 6-membered aromatic heterocyclic

30

group optionally having a substituent selected from the group consisting of a lower alkyl group, a halogen atom, and a lower alkoxy group and having a nitrogen atom as an atom constituting the ring. It is particularly preferable that the heterocyclic group of D be a pyridyl group.

5 Further, in the general formula (1) of the present invention, B represents any one of an alkoxy group having 1 to 10 carbon atoms, a hydroxyl group, or a hydroxyamino group, and these groups are optionally substituted with a substituent selected from the group consisting of an aryl group, a hydroxyl group, an alkyl group having 1 to 10 carbon atoms, a lower
10 alkoxy group, a lower alkylamino group, a halogen atom, and a heterocyclic group, in which a lower alkoxy group optionally substituted with a substituent selected from the group consisting of a lower alkyl group, a lower alkoxy group, a lower alkylamino group, a halogen atom, and a heterocyclic group, or a hydroxyl group is preferable. The heterocyclic group when B is
15 any one of an alkoxy group substituted with a heterocyclic group and having 1 to 10 carbon atoms, a hydroxyl group, or a hydroxyamino group is preferably a saturated ring group containing a heteroatom selected from a nitrogen atom, an oxygen atom, and a sulfur atom, and specific examples thereof include a morpholyl group, a piperazyl group, a tetrahydropyranyl
20 group, and an oxetanyl group, in which a tetrahydropyranyl group is preferable.

[0018]

Further, in the general formula (1) of the present invention, when A represents a group represented by the general formula (2-1), (2-2), or (2-3), E
25 preferably represents a 5- or 6-membered heterocyclic group substituted with a substituent selected from the group consisting of a 3- to 8-membered saturated ring group containing a nitrogen atom connected by a carbon atom, a 3- to 8-membered saturated or unsaturated ring group containing a heteroatom selected from an oxygen atom and a sulfur atom, a lower
30 alkoxyalkylene group, a lower alkylthioalkylene group, a lower

alkylaminoalkylene group, a lower alkylamino group, and a lower alkenylamino group (preferably, a 5- or 6-membered cyclic amino group containing 1 to 4 nitrogen atoms having the above-described substituent); or a 5- or 6-membered cyclic ketone group containing 1 to 4 nitrogen atoms as
 5 atoms constituting the ring, the cyclic ketone group being substituted with a lower alkyl group or a lower alkenyl group, in combination with the above-described (i) or (ii).

It is preferable that the 5- or 6-membered heterocyclic group of E be selected from the group consisting of a pyridyl group, a pyridazyl group, a
 10 pyrimidyl group, a pyrazyl group, a pyrrolyl group, a triazolyl group, and a tetrazole group, and the cyclic ketone group be a 5- or 6-membered cyclic ketone group containing 2 or 3 nitrogen atoms as atoms constituting the ring.

Further, the lower alkylene group of the lower alkoxy-lower alkylene
 15 group, the lower alkylthio-lower alkylene group, or the lower alkylamino-lower alkylene group preferably has 1 to 6 carbon atoms, more preferably has 1 to 3 carbon atoms, and the lower alkyl group preferably has 1 to 3 carbon atoms.

Further, the 3- to 8-membered saturated or unsaturated ring group
 20 containing a heteroatom selected from an oxygen atom and a sulfur atom is preferably a 5- or 6-membered saturated or unsaturated ring group containing one heteroatom selected from an oxygen atom and a sulfur atom, and specifically, tetrahydropyran, tetrahydrofuran, oxetane, furan, thiophene, or the like is preferable.

25 The 3- to 8-membered saturated ring group containing a nitrogen atom is preferably a 5- or 6-membered saturated ring group containing one nitrogen atom, and specifically, pyrrolidine, piperidine, or the like is preferable.

Further, the cyclic ketone group is specifically preferably a
 30 pyrimidinone group, and the lower alkyl group or the lower alkenyl group is

preferably bonded to the nitrogen atoms constituting the cyclic ketone group.
[0019]

When A represents a group represented by the general formula (2-4),
E preferably represents a phenyl group or a 5- or 6-membered heterocyclic
5 group optionally having a substituent selected from the group consisting of a
halogen atom, a hydroxyl group, a lower alkyl group, a lower alkenyl group, a
lower alkynyl group, a lower alkoxy group, a lower alkylthio group, a
hydroxy-lower alkyl group, a hydroxy-lower alkenyl group, a hydroxy-lower
alkoxy group, a halogeno-lower alkyl group, a halogeno-lower alkoxy group, a
10 halogeno-lower alkylthio group, a halogeno-lower alkenyl group, a nitro
group, a cyano group, an amino group, a carboxyl group, a lower
alkyloxycarbonyl group, a carbamoyl group, a lower alkanoyl group, an aroyl
group, a lower alkylsulfonyl group, a sulfamoyl group, and an ammonium
group; an aminocarbonyl group optionally having a substituent selected from
15 the group consisting of a hydroxyl group, a lower alkyl group, a lower alkenyl
group, a lower alkynyl group, a lower alkoxy group, a hydroxy-lower alkyl
group, a hydroxy-lower alkenyl group, a hydroxy-lower alkoxy group, a
halogeno-lower alkyl group, a halogeno-lower alkoxy group, a halogeno-lower
alkenyl group, an amino group, a lower alkylamino group, an aryl group, a
20 heterocyclic group, a lower alkylsulfonyl group, and a sulfamoyl group; a
hydrogen atom; a halogen atom; a hydroxyl group; a lower alkyl group; a
lower alkenyl group; a lower alkynyl group; a lower alkoxy group; a lower
alkylthio group; a hydroxy-lower alkyl group; a hydroxy-lower alkenyl group;
a hydroxy-lower alkoxy group; a halogeno-lower alkyl group; a
25 halogeno-lower alkoxy group; a halogeno-lower alkylthio group; a
halogeno-lower alkenyl group; a nitro group; a cyano group; an amino group;
a carboxyl group; a dihydroxyboryl group; a lower alkyloxycarbonyl group; a
carbamoyl group; a lower alkanoyl group; an aroyl group; a lower
alkylsulfonyl group; a sulfamoyl group; an ammonium group; a lower
30 alkylaminoalkylene group; or a 5- or 6-membered heterocyclic group

substituted with a substituent selected from the group consisting of a 3- to 8-membered saturated ring group containing a nitrogen atom connected by a carbon atom, a 3- to 8-membered saturated or unsaturated ring group containing a heteroatom selected from an oxygen atom and a sulfur atom, a lower alkoxyalkylene group, a lower alkylthioalkylene group, a lower alkylaminoalkylene group, a lower alkylamino group, and a lower alkenylamino group (preferably, a 5- or 6-membered cyclic amino group containing 1 to 4 nitrogen atoms having the above-described substituent); or a 5- or 6-membered cyclic ketone group containing 1 to 4 nitrogen atoms as atoms constituting the ring, the cyclic ketone group being substituted with a lower alkyl group or a lower alkenyl group.

Note that, preferable examples of E are the same as described for the case where A represents a group represented by the general formula (2-1), (2-2), or (2-3).

[0020]

Further, when A represents a group represented by the general formula (2-4), E is preferably a 5- or 6-membered aromatic heterocyclic group containing 1, 2, 3, or 4 heteroatoms selected from oxygen atoms, sulfur atoms, or nitrogen atoms and optionally having a substituent selected from the group consisting of a lower alkyl group, a lower alkoxy group, and a halogen atom, in combination with the above-described (i) or (ii). It is preferable that the aromatic heterocyclic group be selected from the group consisting of a pyridyl group, a pyridazyl group, a pyrimidyl group, a pyrazinyl group, a pyrrolyl group, a furanyl group, a thiophenyl group, a thiazolyl group, an oxazolyl group, a triazolyl group, and a tetrazole group.

When A represents a group represented by the general formula (2-4), E is preferably an aminocarbonyl group optionally substituted with a lower alkyl or a heterocyclic or heterocyclic lower alkyl group; or a lower alkylaminoalkylene group, in combination with the above-described (i) or (ii). Further, E is also preferably a pyridyl group, a pyrimidyl group, a triazolyl

group, or a pyrrolyl group optionally having a lower alkyl group or a 4 to 6-membered cyclic amino group; or an aminocarbonyl group optionally substituted with a lower alkyl or heterocyclic lower alkyl group. Examples of the 4 to 6-membered cyclic amino group include an azetidiny group, and
 5 examples of the heterocyclic lower alkyl group include a tetrapyranyl lower alkyl group.

Further, when E represents a lower alkylcarbonyl group or a lower alkyloxycarbonyl group, examples of a fused ring formed by such a group bonding to the phenyl group of D include a 1-oxoindanyl group and a
 10 3-oxo-1H-isobenzofuranyl group.
 [0021]

Further, in the general formula (1) of the present invention, or in any one of the above-described preferable aspects, R1, R11, R21, R51, and R52 are preferably lower alkyl groups.

15 Further, in the general formula (1) of the present invention, or in any one of the above-described preferable aspects, Arm is preferably selected from the group consisting of a phenyl group, a pyridyl group, a pyrimidyl group, and an imidazolyl group.


Further, in the general formula (1) of the present invention, or in any
 20 one of the above-described preferable aspects, it is preferable that one of R31, R32, R33, and R34 be a halogen atom with the others being hydrogen atoms, two of them be halogen atoms with the others being hydrogen atoms, or four of them be halogen atoms.

Further, in the general formula (1) of the present invention, or in any
 25 one of the above-described preferable aspects, it is preferable that g be a carbon or nitrogen atom, and e be a carbon or nitrogen atom.
 [0022]

In the present invention, compounds represented by the following formulae or pharmaceutically acceptable salts thereof are particularly
 30 preferable. Further, prodrugs of these compounds or pharmaceutically


acceptable salts thereof are also preferable.

(2S)-2-[[[2,5-Difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propionic acid



Cyclohexyl


(2S)-2-[[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate



Isopropyl

(2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate


15



1-Ethylpropyl

(2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl


amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate



[0024]


5 (4,4-Dimethylcyclohexyl)

(2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl]amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate



10 Cyclopentyl

(2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl]amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate




15 [0025]

Cycloheptyl


(2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl]amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate

3-pyridyl]propanoate



2,2-Dimethylpropyl


- (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
 5 amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
 3-pyridyl]propanoate




[0026]

Cyclobutylmethyl

- 10 (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
 amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
 3-pyridyl]propanoate




- (2S)-2-[[2,5-Difluoro-4-(2-furysulfonylamino)benzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoic acid
 15



[0027]

(2S)-2-[[[2,5-Difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoic acid



[0028]

Further, when A represents a group represented by the general formula (2-2), E preferably represents a 5- or 6-membered heterocyclic group containing 1 to 4 nitrogen atoms, the 5- or 6-membered heterocyclic group being substituted with a substituent selected from a lower alkyl group, a lower alkoxy-lower alkylene group, and a 5- or 6-membered heterocyclic group containing an oxygen atom; or a cyclic ketone group containing 1 to 4 nitrogen atoms, the cyclic ketone group being substituted with a lower alkyl group or a lower alkenyl group.

Further, E preferably represents a pyrimidyl group substituted with a lower alkoxy-lower alkylene group or a tetrahydropyranyl group; or a pyrimidinone group substituted with a lower alkenyl group.

Further, B preferably represents an alkoxy group having 1 to 10 carbon atoms, a hydroxyl group, or a hydroxyamino group, wherein B is

optionally substituted with a substituent selected from the group consisting of an alkyl group having 1 to 10 carbon atoms, a lower alkoxy group, and a heterocyclic group containing 1 to 4 heteroatoms selected from oxygen atoms and nitrogen atoms.

5 Further, Arm is preferably a pyridyl group.

[0029]

Further, when A represents a group represented by the general formula (2-4), E is preferably a 5- or 6-membered heterocyclic group containing an oxygen atom, or a 5- or 6-membered heterocyclic group
10 containing 1 to 4 nitrogen atoms, wherein E optionally has a substituent selected from the group consisting of a lower alkyl group, a lower alkoxy-lower alkylene group, and a 5- or 6-membered heterocyclic group containing an oxygen atom.


Further, E is preferably a tetrahydropyranyl group or a pyrimidyl
15 group, wherein E optionally has a substituent selected from the group consisting of a lower alkyl group and a tetrahydropyranyl group.





Further, B is preferably a lower alkoxy group or a hydroxyl group.





Further, R₅₁, R₅₂, and R₅₃ are each preferably a lower alkyl group.





[0030]





20 Further, in the present invention, compounds represented by the following formulae or pharmaceutically acceptable salts thereof are particularly preferable. Further, prodrugs of these compounds or pharmaceutically acceptable salts thereof are also preferable.















[0031]

When the compound represented by the general formula (1) of the
 5 present invention can take the form of a salt, the salt thereof may be
 pharmaceutically acceptable, and examples thereof for the acidic group such
 as the carboxyl group in the formula can include ammonium salts, salts with
 alkali metals such as sodium and potassium, salts with alkaline earth
 metals such as calcium and magnesium, aluminum salts, zinc salts, salts
 10 with organic amines such as triethylamine, ethanolamine, morpholine,
 piperidine, and dicyclohexylamine, and salts with basic amino acids such as
 arginine and lysine. Examples thereof for the basic group when the basic
 group is present in the formula can include salts with inorganic acids such as
 hydrochloric acid, sulfuric acid, and phosphoric acid, salts with organic
 15 carboxylic acids such as acetic acid, citric acid, benzoic acid, maleic acid,
 fumaric acid, tartaric acid, and succinic acid, and salts with organic sulfonic
 acids such as methanesulfonic acid and p-toluenesulfonic acid. As a method
 for forming salts, salts can be obtained by mixing the compound of the

general formula (1) with a necessary acid or base at a suitable quantitative ratio in a solvent or dispersant, or performing cation exchange or anion exchange from other forms of salts.

The compound of the present invention may contain a solvate of the compound represented by the general formula (1), such as hydrates and alcohol adducts.

[0032]

The compound of the present invention includes embodiments of prodrugs of the compound represented by the general formula (1). A prodrug of the compound of the present invention means a compound converted into the compound represented by the general formula (1) by reaction with enzyme, gastric acid, or the like under physiological conditions in vivo, that is, a compound transformed into the compound represented by the general formula (1) by undergoing enzymatic oxidation, reduction, hydrolysis, or the like, or a compound transformed into the compound represented by the general formula (1) by undergoing hydrolysis by gastric acid or the like. The prodrug of the compound represented by the general formula (1) is exemplified by the compounds of Examples, but is not limited thereto, and examples thereof to be used include, when the compound represented by the general formula (1) has amino, compounds obtained by acylation, alkylation, or phosphorylation of the amino (e.g., compounds obtained by eicosanoylation, alanylation, pentylaminocarbonylation, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methoxycarbonylation, tetrahydrofuranylation, pyrrolidylmethylation pivaloyloxymethylation, or tert-butylation of the amino of the compound represented by the general formula (1)); when the compound represented by the general formula (1) has hydroxy, compounds obtained by acylation, alkylation, phosphorylation, or boration of the hydroxy (e.g., compounds obtained by acetylation, palmitoylation, propanoylation, pivaloylation, succinylation, fumarylation, alanylation, or dimethylaminomethylcarbonylation of the hydroxy of the

compound represented by the general formula (1)); and when the compound represented by the general formula (1) has carboxyl, compounds obtained by esterification or amidation of the carboxyl (e.g., compounds obtained by ethyl esterification, phenyl esterification, isopropyl esterification, isobutyl esterification, cyclopentyl esterification, cyclohexyl esterification, cycloheptyl esterification, cyclobutylmethyl esterification, cyclohexylmethyl esterification, n-hexyl esterification, sec-butyl esterification, tert-butyl esterification, (4-tetrahydropyranyl)methyl esterification, (4-tetrahydropyranyl) esterification, carboxymethyl esterification, dimethylaminomethyl esterification, pivaloyloxymethyl esterification, ethoxycarbonyloxyethyl esterification, phthalidyl esterification, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl esterification, cyclohexyloxycarbonylethyl esterification, or methylamidation of the carboxyl of the compound represented by the general formula (1)). These compounds can be produced from the compound represented by the general formula (1) by known methods.

Further, the prodrug of the compound (1) may be a compound transformed into the compound (1) under physiological conditions as described in "IYAKUHIN no KAIHATSU (Development of Pharmaceuticals)", Vol. 7, Design of Molecules, p. 163 to 198, published by HIROKAWA SHOTEN in 1990.


[0033]

The present invention includes all isotopes of the compound represented by formula (1). An isotope of the compound of the present invention has at least one atom substituted with an atom having the same atomic number (the number of protons) and a different mass number (the sum of the number of protons and neutrons). Examples of the isotope contained in the compound of the present invention include a hydrogen atom, a carbon atom, a nitrogen atom, an oxygen atom, a phosphorus atom, a sulfur atom, a fluorine atom, and a chlorine atom, which respectively contain

^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , ^{36}Cl , or the like. In particular, unstable radioactive isotopes that emit radiation to emit neutrons, such as ^3H and ^{14}C , are useful for distribution examination of pharmaceutical products or compounds in body tissues. Stable isotopes can be used with safety, since they do not disintegrate, their abundance is less likely to change, and they are not radioactive. The isotope of the compound of the present invention can be converted according to conventional methods by replacing a reagent used for synthesis with a reagent containing a corresponding isotope.

10 [0034]

The compound represented by the general formula (1) of the present invention can be produced, for example, by subjecting an intermediate represented by the general formula (M-I) and having an amino group at an end and an intermediate represented by the general formula (M-II) and having a carboxyl group at an end to dehydration reaction.



Of these intermediates, the intermediate represented by the general formula (M-II) and having a carboxyl group at an end can be produced, for example, by the following method.


20 [0035]

A typical method for producing a compound of the intermediate represented by the general formula (M-II) and having a carboxyl group at an end, which is the compound of the present invention, is shown below. In the following description, the symbols in the formulae are the same as defined in the formula (I), unless otherwise specified.

(1) In the general formula (M-II), an intermediate (S6) having a carboxyl group at an end in which D is a phenyl group or a pyridyl group optionally having a substituent selected from the group consisting of a lower alkyl group and a halogen atom, and E is a 5- or 6-membered aromatic heterocyclic group containing 1, 2, 3, or 4 heteroatoms selected from oxygen atoms, sulfur atoms, or nitrogen atoms and optionally having a substituent can be synthesized, for example, using methods described below (production methods A, B, C, and D).

[0036]

10 <Production method A>



[0037]

D' in the formula represents the above-described substituent represented by D or a substituent that can be easily converted into D by an operation such as deprotection, and E' in the formula represents the above-described substituent represented by E or a substituent that can be easily derivatized to E by an operation such as deprotection. Further, R51 represents a common substituent of ester such as a lower alkyl group.

A sulfonamide derivative (S3) can be synthesized by allowing a sulfonyl chloride derivative (S1) (X in the formula represents a halogen atom such as chlorine, bromine, and iodine, or a leaving group such as a trifluoromethanesulfonyloxy group) to react with an amine derivative (S2) in


a solvent having no adverse effect on the present reaction such as methylene chloride in the presence of a base such as pyridine. The resulting sulfonamide derivative (S3) can be derivatized to a corresponding sulfonamide derivative (S5) by Suzuki coupling reaction with a boronic acid derivative (S4). Subsequently, the target intermediate (S6) having a carboxyl group at an end can be produced by subjecting the resulting sulfonamide derivative (S5) to alkaline hydrolysis using a base such as sodium hydroxide or deprotection such as acid hydrolysis, for example, using hydrochloric acid or trifluoroacetic acid, in a solvent having no adverse effect on the present reaction such as tetrahydrofuran, methanol, and ethanol.

[0038]

Suzuki coupling reaction is known, and is performed by allowing a boronic acid derivative or a boronic acid ester derivative to react with a halogen derivative, trifluoromethanesulfonate, or methanesulfonate in a solvent having no adverse effect on the present reaction such as 1,4-dioxane, tetrahydrofuran, toluene, dimethylsulfoxide, and 1,2-dimethoxyethane in the presence or absence of a cosolvent such as water using a transition metal catalyst such as tetrakis(triphenylphosphine)palladium(0), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), and tris(dibenzylideneacetone)dipalladium(0), or using a transition metal catalyst such as palladium(II) acetate and a ligand such as triphenylphosphine in the presence of a base such as potassium carbonate, sodium carbonate, potassium phosphate, sodium methoxide, and sodium hydroxide.

[0039]

<Production method B>



[0040]


D' in the formula represents the above-described substituent represented by D, or a substituent that can be easily converted into D by an operation such as deprotection, and E' in the formula represents the above-described substituent represented by E or a substituent that can be easily derivatized to E by an operation such as deprotection. Further, R₅₁ represents a common substituent of ester such as a lower alkyl group.

A corresponding boronic acid derivative (S7) can be synthesized by coupling reaction of the sulfonamide derivative (S3) that is the intermediate described in <Production method A> with a borane derivative such as bis(pinacolato)diborane in a solvent having no adverse effect on the present reaction such as N,N-dimethylformamide in the presence of a base such as potassium acetate using a metal catalyst such as [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) to lead it to a corresponding boronic acid ester derivative, and subsequently subjecting the resulting boronic acid ester derivative to treatment, for example, by adding sodium periodate, ammonium acetate, and water in a solvent having no adverse effect on the present reaction such as acetone to remove boronic acid ester. The resulting boronic acid derivative (S7) can be derivatized to the corresponding sulfonamide derivative (S5) by Suzuki coupling reaction

(described above) with a halogen derivative (S8) (X in the formula represents a halogen atom such as chlorine, bromine, and iodine, or a leaving group such as a trifluoromethanesulfonyloxy group). The target intermediate (S6) having a carboxyl group at an end can be produced by subjecting the sulfonamide derivative (S5) to deprotection by hydrolysis or the like using the method described in <Production method A>.

[0041]

<Production method C>



[0042]


D' in the formula represents the above-described substituent represented by D or a substituent that can be easily converted into D by an operation such as deprotection, and E' in the formula represents the above-described substituent represented by E or a substituent that can be easily derivatized to E by an operation such as deprotection. Further, R₅₁ represents a common substituent of ester such as a lower alkyl group.

The sulfonamide derivative (S3) that is the intermediate described in <Production method A> (X in the formula represents a halogen atom such as chlorine, bromine, and iodine, or a leaving group such as a trifluoromethanesulfonyloxy group) can be derivatized to the corresponding sulfonamide derivative (S5) by coupling reaction with an aromatic heterocycle (S9) in a solvent having no adverse effect on the present reaction such as N-methylpyrrolidone in the presence of a ligand that is commonly used for organic synthesis such as trans-N,N'-dimethylcyclohexane-1,2-diamine and a base such as potassium phosphate using a metal catalyst such as copper(I) iodide. Subsequently,

the target intermediate (S6) having a carboxyl group at an end can be produced by subjecting the resulting sulfonamide derivative (S5) to deprotection by hydrolysis or the like using the method described in <Production method A>.

5 [0043]

<Production method D>



[0044]

D' in the formula represents the above-described substituent represented by D or a substituent that can be easily converted into D by an operation such as deprotection, and E' in the formula represents the above-described substituent represented by E or a substituent that can be easily derivatized to E by an operation such as deprotection. Further, R₅₁ represents a common substituent of ester such as a lower alkyl group.

15 A sulfonamide derivative (S10) (X in the formula represents a halogen atom such as chlorine, bromine, and iodine, or a leaving group such as a trifluoromethanesulfonyloxy group) can be derivatized to the corresponding sulfonamide derivative (S5) by coupling reaction with a halogen derivative (S11) in a solvent having no adverse effect on the present reaction such as 1,4-dioxane in the presence of a metal catalyst such as tris(dibenzylideneacetone)dipalladium(0), a ligand commonly used for organic synthesis such as XantPHOS, and a base such as cesium carbonate. Subsequently, the target intermediate (S6) having a carboxyl group at an end can be produced by subjecting the resulting sulfonamide derivative (S5) to deprotection such as hydrolysis, for example, by the method described in <Production method A>.

25 [0045]

$$\begin{array}{c}
 \text{HO-C(=O)-D'-S(=O)}_2\text{-Cl} \quad + \quad \text{HN(R}_{41}\text{)-C(d)=C(a)-C(b)=C(c)-C(=O)OR}_{51} \quad \longrightarrow \quad \text{HO-C(=O)-D'-S(=O)}_2\text{-N(R}_{41}\text{)-C(d)=C(a)-C(b)=C(c)-C(=O)OR}_{51} \quad \xrightarrow{\text{(S14)}} \quad \text{R}_{71}\text{-N(R}_{61}\text{)-C(=O)-D'-S(=O)}_2\text{-N(R}_{41}\text{)-C(d)=C(a)-C(b)=C(c)-C(=O)OR}_{51} \\
 \text{(S12)} \qquad \qquad \qquad \text{(S2)} \qquad \qquad \qquad \text{(S13)} \qquad \qquad \qquad \text{(S15)} \\
 \\
 \longrightarrow \quad \text{R}_{71}\text{-N(R}_{61}\text{)-C(=O)-D'-S(=O)}_2\text{-N(R}_{41}\text{)-C(d)=C(a)-C(b)=C(c)-C(=O)OH} \\
 \text{(S16)}
 \end{array}$$

D' in the formula represents the above-described substituent represented by D or a substituent that can be easily converted into D by an operation such as deprotection. Further, R51 represents a common substituent of ester such as a lower alkyl group. Further, R61 and R71 represent common substituents on the amine such as a lower alkyl group, a heterocyclic group, or a lower alkyl group substituted with a heterocycle.

41

Subsequently, an intermediate (S16) having a carboxyl group at an end can be produced by subjecting the resulting amide derivative (S15) to deprotection such as hydrolysis using the method described in <Production method A>.

5 [0047]

The amidation reaction is known, and examples thereof include (1) a method using a condensing agent and (2) a method using an acid halide.

(1) The method using a condensing agent is performed by allowing carboxylic acid to react with amine or a salt thereof in a solvent having no
 10 adverse effect on the present reaction such as dichloromethane, tetrahydrofuran, 1,4-dioxane, N,N-dimethylformamide, or acetonitrile in the presence or absence of a base such as pyridine, triethylamine, or N-ethyl-diisopropylamine in the presence or absence of a condensation aid such as 1-hydroxybenzotriazole (HOBt), 1-hydroxy-7-azabenzotriazole
 15 (HOAt), or N-hydroxysuccinimide (HOSu) using a condensing agent such as 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide (WSC), 1,3-dicyclohexylcarbodiimide (DCC), or O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU).

20 [0048]


(2) The method using an acid halide is performed by allowing carboxylic acid to react with thionyl chloride, oxalyl chloride, thionyl bromide, or the like, in a solvent having no adverse effect on the present reaction such as dichloromethane or in the absence of the solvent, in the presence or
 25 absence of a catalyst such as N,N-dimethylformamide to obtain an acid halide, and allowing the acid halide to react with amine or a salt thereof in a solvent having no adverse effect on the present reaction such as dichloromethane or tetrahydrofuran in the presence of a base such as pyridine, triethylamine, or N-ethyl-diisopropylamine.

30 In each step, the synthesis can be performed using generally

replacable reaction conditions, which should be appropriately selected depending on the types or the like of raw material compounds. The compound of the present invention obtained by the above-described methods can be purified by techniques that are generally used for organic synthesis, such as extraction, distillation, crystallization, and column chromatography. [0049]

The intermediate represented by the general formula (M-I) and having an amino group at an end, which is the compound of the present invention, can be synthesized, for example, using the method disclosed in Patent Literature 1 or the production methods (production methods F and G) described below. In the following description, the symbols in the formulae are the same as defined in the formula (I), unless otherwise specified. [0050]

<Production method F>



[0051]


In the formula, R₈₁ represents a common substituent of amine that can be removed by an operation such as deprotection, such as a tert-butoxycarbonyl group and a benzyloxycarbonyl group, and Y in the formula represents a halogen atom such as chlorine, bromine, and iodine, or a leaving group such as a trifluoromethanesulfonyloxy group. Y' in the formula represents a substituent that can be easily derivatized to Y by an

operation such as trifluoromethanesulfonylation.

A corresponding amino acid derivative (S18) can be synthesized by allowing an amino acid derivative (S17) to react, for example, with N-phenyl bis(trifluoromethanesulfonamide) in a solvent having no adverse effect on the present reaction such as tetrahydrofuran in the presence of a base such as diisopropylethylamine. A corresponding boronic acid derivative (S19) can be synthesized by subjecting the resulting amino acid derivative (S18) to coupling reaction with a boronic acid derivative such as bis(2,2,3,3-tetramethyl-2,3-butanedionate)diboron in a solvent having no adverse effect on the present reaction such as 1,4-dioxane in the presence of a base such as potassium acetate using a metal catalyst such as [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) to lead it to a corresponding boronic acid ester derivative, and subsequently subjecting the resulting boronic acid ester derivative to treatment, for example, by adding sodium periodate, ammonium acetate, and water in a solvent having no adverse effect on the present reaction such as acetone to hydrolyze the boronic acid ester. An amino acid derivative (S21) can be produced by coupling reaction of the resulting boronic acid derivative (S19) with a compound (S20) in a solvent having no adverse effect on the present reaction such as methylene chloride in the presence of a base such as triethylamine using a metal catalyst such as copper acetate (II), or coupling reaction thereof in a solvent having no adverse effect on the present reaction such as 1,4-dioxane in the presence of a base such as sodium carbonate using a metal catalyst such as [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II). A target intermediate (S22) having an amino group at an end can be produced by subjecting the resulting amino acid derivative (S21) to acid hydrolysis, for example, using hydrochloric acid or trifluoroacetic acid, or deprotection by hydrogenation reaction or the like in the presence of hydrogen using a catalyst such as palladium carbon.

[0052]

<Production method G>



In the formula, R81 represents a common substituent of amine that can be removed by an operation such as deprotection, such as a tert-butoxycarbonyl group and a benzyloxycarbonyl group, and X' and X" in the formula each independently represent a halogen atom such as chlorine, bromine, and iodine, or a leaving group such as a trifluoromethanesulfonyloxy group.

A corresponding halogen derivative (S24) can be synthesized by subjecting an amine derivative (S23) to the method disclosed, for example, in Patent Literature 1, or by subjecting the same substrate as disclosed in Patent Literature 1 to ureation with triphosgene or the like in a solvent having no adverse effect on the present reaction such as methylene chloride, subsequently to cyclization with potassium carbonate and p-toluenesulfonic acid methyl, methyl iodide, or the like in a solvent having no adverse effect on the present reaction such as N,N-dimethylformamide, and subsequently to N-methylation. An amino acid derivative (S25) can be derivatized to the corresponding amino acid derivative (S21) by coupling reaction with the halogen derivative (S24) in a solvent having no adverse effect on the present reaction such as N,N-dimethylformamide in the presence of metal such as zinc using a metal catalyst such as bis(triphenylphosphine)palladium(II) dichloride. A target intermediate (S22) having an amino group at an end can be produced by subjecting the resulting amino acid derivative (S21) to

acid hydrolysis, for example, using hydrochloric acid or trifluoroacetic acid, or deprotection by hydrogenation or the like in the presence of hydrogen using a catalyst such as palladium carbon.

A compound in which A has a group represented by the general
5 formula (2-4) also can be produced according to the above-described method or the method disclosed in Patent Literature 4.

[0053]

The compound represented by the general formula (1) or a salt thereof is administered as it is or as various pharmaceutical compositions.
10 Examples of the dosage forms of the pharmaceutical compositions include tablets, powders, pills, granules, capsules, suppositories, solutions, dragees, depot preparations, or syrups, and the pharmaceutical compositions can be produced according to conventional methods using common formulation aids.

For example, a tablet can be obtained by mixing the phenylalanine
15 derivative that is the active ingredient of the present invention with known auxiliary substances, e.g., inert diluents such as lactose, calcium carbonate, or calcium phosphate, binders such as gum arabic, corn starch, or gelatin, swelling agents such as alginic acid, corn starch, or pregelatinized starch, sweeteners such as sucrose, lactose, or saccharin, flavoring agents such as
20 peppermint, Gaultheria adenothrix (Akamono) oil, or cherry, wetting agents such as magnesium stearate, talc, or carboxymethylcellulose, excipients for soft gelatin capsules and suppositories such as fats, waxes, semi-solid and liquid polyols, natural oils, or hardened oils, or excipients for solutions such as water, alcohol, glycerol, polyol, sucrose, invert sugar, glucose, or vegetable
25 oils.

An inhibitor containing the compound represented by the general formula (1) or a salt thereof as an active ingredient can be used as a therapeutic agent or a preventive agent for any one of inflammatory disease in which an $\alpha 4$ integrin-dependent adhesion process is involved in
30 pathological conditions, rheumatoid arthritis, inflammatory bowel disease,

systemic lupus erythematosus, multiple sclerosis, Sjogren's syndrome, asthma, psoriasis, allergy, diabetes, cardiovascular disease, arteriosclerosis, restenosis, tumor growth, tumor metastasis, and transplantation rejection.

The dose used for the above-described object is determined depending on the target therapeutic effect, the administration method, the treatment period, the age, the body weight, or the like, but the adult dose per day by a route of oral or parenteral administration may be generally 1 μ g to 5 g in the case of oral administration, and 0.01 μ g to 1 g in the case of parenteral administration.

10

Examples


[0054]

The present invention will be described in detail below with reference to the following Examples. The Examples are preferred embodiments of the present invention, and the present invention will not be limited to the Examples.

15

Example 1: Synthesis of M-1

(Step 1) Methyl 3-[(5-bromo-2-pyridyl)carbamoylamino]pyridine-4-carboxylate



20

Methyl 3-aminopyridine-4-carboxylate (31.8 g, 209 mmol) was dissolved in methylene chloride (1.1 L), a solution (50 ml) of triphosgene (20.7 g, 69.8 mmol) in methylene chloride was added thereto, and the resulting mixture was stirred at 0°C for 3 hours. A solution (50 ml) of 5-bromopyridin-2-amine (30.0 g, 174 mmol) in methylene chloride was added to the reaction solution, and the resulting mixture was stirred at room temperature for 12 hours. The precipitated solid was filtered, washed with methylene chloride, and then dried under reduced pressure to obtain the

25

title compound (30 g, 49%).


¹H NMR (400 MHz, DMSO-d₆): δ 11.02 (s, 1H), 10.36 (s, 1H), 9.44 (s, 1H), 8.39 (s, 2H), 7.98 (d, J = 8.0 Hz, 1H), 7.74 - 7.73 (m, 1H), 7.48 (d, J = 8.0 Hz, 1H), 3.92 (s, 3H).

5 [0055]

(Step

2)

3-(5-Bromo-2-pyridyl)-1-methyl-pyrido[3,4-d]pyrimidine-2,4-dione



Methyl

10 3-[(5-bromo-2-pyridyl)carbamoylamino]pyridine-4-carboxylate <see (step 1)>
(30 g, 85.7 mmol) was dissolved in N,N-dimethylformamide (600 ml), an
aqueous solution (80 ml) of potassium carbonate (23.7 g, 171 mmol) was
added thereto, and the resulting mixture was stirred at room temperature
for 3 hours. To the resulting solution, potassium carbonate (23.7 g, 171
15 mmol) and p-toluenesulfonic acid methyl ester (31.9 g, 171 mmol) were
further added, and the resulting mixture was stirred at room temperature
for 90 minutes. The reaction liquid was cooled to 0°C, and then diluted with
water and extracted with ethyl acetate (100 ml × 4). The extracts were
combined, washed with a saturated saline solution, dried over anhydrous
20 sodium sulfate, and then the solvent was removed under reduced pressure.
The obtained residue was stirred in petroleum ether for 20 minutes, and the
obtained solid was filtered and dried under reduced pressure to obtain the
title compound (9.8 g, 34%).

¹H NMR (400 MHz, DMSO-d₆): δ 9.00 (s, 1H), 8.77 (m, 1H), 8.58 (d, J = 4.8
25 Hz, 1H), 8.32 - 8.29 (m, 1H), 7.91 (d, J = 4.8 Hz, 1H), 7.52 (d, J = 8.4 Hz, 1H),
3.61 (s, 3H).


[0056]

(Step

3)

Cyclohexyl

(2S)-3-benzyloxy-2-(tert-butoxycarbonylamino)propanoate



(2S)-3-Benzoyloxy-2-(tert-butoxycarbonylamino)propionic acid (300 g,
 5 1.02 mol) was dissolved in N,N-dimethylformamide (2.0 L), EDCL (189 g,
 1.22 mmol), cyclohexanol (210 g, 2.10 mol), and
 N,N-dimethyl-4-aminopyridine (12.4 g, 102 mmol) were added thereto, and
 the resulting mixture was stirred in the presence of nitrogen gas at room
 temperature for 90 minutes. The reaction solution was diluted with water
 10 (5.0 L) and then extracted with ethyl acetate (1.0 L $\times$ 2). The extracts were
 combined and washed with water (1.0 L), 1 N hydrochloric acid (900 ml), a
 saturated aqueous solution of sodium hydrogen carbonate (1.0 L), and a
 saturated saline solution (900 ml), dried over anhydrous sodium sulfate, and
 then the solvent was removed under reduced pressure to obtain the title
 15 compound (360 g, 94%).

^1H NMR (400 MHz, DMSO- d_6): δ 7.35 - 7.25 (m, 5H), 7.06 (d, J = 8.4 Hz, 1H),
 4.73 - 4.69 (m, 1H), 4.51 - 4.39 (m, 2H), 4.26 - 4.21 (m, 1H), 3.73 - 3.64 (m, 2H),
 1.47 - 1.10 (m, 19H).

[0057]


20

(Step

4)

Cyclohexyl

(2S)-2-(tert-butoxycarbonylamino)-3-hydroxy-propanoate



Cyclohexyl (2S)-2-(tert-butoxycarbonylamino)-3-hydroxy-propanoate
 (360 g, 954 mmol) <see (step 3)> was dissolved in ethanol (1.6 L), palladium

hydroxide 20% on carbon (40 g) was added thereto, and the resulting mixture was stirred in the presence of hydrogen gas (50 psi) at 70°C for 3 days. The reaction solution was filtered, and the obtained filtrate was concentrated and then dried under reduced pressure to obtain the title compound (250 g, 91%).

5 ¹H NMR (400 MHz, DMSO-d₆): δ 6.80 (d, J = 8.0 Hz, 1H), 4.82 (s, 1H), 4.72 - 4.67 (m, 1H), 4.37 (s, 1H), 4.01 - 3.99 (m, 1H), 3.46 - 3.38 (m, 1H), 1.48 - 1.35 (m, 19H).


[0058]

(Step

5)

Cyclohexyl

10 (2R)-2-(tert-butoxycarbonylamino)-3-iodine-propanoate




Triphenylphosphine (275 g, 1.05 mol) and imidazole (75.0 g, 1.05 mol) were dissolved in methylene chloride (3.0 L), the resulting solution was cooled to 0°C, then iodine (270 g, 1.05 mol) was added thereto, and the resulting mixture was stirred in the presence of nitrogen gas at room temperature for 30 minutes. The reaction solution was cooled to 0°C, then a solution (500 ml) of cyclohexyl (2S)-2-(tert-butoxycarbonylamino)-3-hydroxy-propanoate <see (step 4)> (250 g, 871 mmol) in methylene chloride was slowly added dropwise over 1 hour, and the reaction liquid was stirred at room temperature for 2 hours. The reaction liquid was filtered, the obtained residue was washed with hexane/diethyl ether (1:1), and the obtained filtrate was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 1:1 to 1:2) to obtain the title compound (200 g, 58%).

25 ¹H NMR (400 MHz, DMSO-d₆): δ 7.24 (d, J = 8.0 Hz, 1H), 4.75 - 4.69 (m, 1H), 4.21 - 4.16 (m, 1H), 3.51 - 3.47 (m, 1H), 3.36 - 3.33 (m, 1H), 1.75 - 1.10 (m, 19H).

[0059]

(Step 6) Cyclohexyl
(2S)-2-(tert-butoxycarbonylamino)-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate



5 Zinc (9.50 g, 146 mmol) was heated at 210°C for 10 minutes, cooled to 70°C, then heated to 210°C again, and stirred for 10 minutes. The resultant was cooled to room temperature, then N,N-dimethylformamide (20 ml) and a solution (5.0 ml) of dibromoethane (2.10 g, 11.2 mmol) in N,N-dimethylformamide were added thereto, and the resulting mixture was stirred at 90°C for 30 minutes. The reaction mixture was cooled to room temperature, then trimethylsilyl chloride (243 mg, 2.25 mmol) was added thereto, and the resulting mixture was stirred at room temperature for 10 minutes. A solution (15 ml) of cyclohexyl (2R)-2-(tert-butoxycarbonylamino)-3-iodine-propanoate <see (step 5)> (8.90 g, 22.5 mmol) in N,N-dimethylformamide was added to the reaction liquid, and the resulting mixture was stirred at 35°C for 90 minutes. This zinc derivative was added to a suspension of 3-(5-bromo-2-pyridyl)-1-methyl-pyrido[3,4-d]pyrimidine-2,4-dione <see (step 2)> (2.50 g, 7.49 mmol) and Pd(PPh₃)₂Cl₂ (788 mg, 1.12 mmol) in N,N-dimethylformamide (20 ml), and the resulting mixture was stirred in the presence of nitrogen gas at 80°C for 2 hours. The reaction mixture was cooled to room temperature, then the reaction solution was filtered, the obtained filtrate was diluted with water (150 ml) and extracted with ethyl acetate (100 ml × 3). The extracts were combined, washed with a saturated saline solution, dried over anhydrous sodium sulfate, and then the solvent was removed under reduced pressure. The obtained residue was purified by

10

15

20


25

silica gel column chromatography (petroleum ether:ethyl acetate = 1:1 to 1:2) to obtain the title compound (1.88 g, 48%).

¹H NMR (400 MHz, DMSO-d₆): δ 8.99 (s, 1H), 8.56 (d, J = 4.8 Hz, 1H), 8.46 - 8.45 (m, 1H), 7.92 - 7.89 (m, 2H), 7.43 - 7.40 (m, 2H), 4.69 - 4.67 (m, 1H), 4.26 - 4.21 (m, 1H), 3.60 (s, 3H), 3.13 - 2.96 (m, 2H), 1.76 - 1.20 (m, 19H).

[0060]

(Step 7) Cyclohexyl 3-[6-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)pyridin-3-yl]-L-alaninate (M-1)



10

Cyclohexyl

(2S)-2-(tert-butoxycarbonylamino)-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate <see (step 6)> (7.50 g, 14.3 mmol) was dissolved in ethyl acetate (20 ml), 4 N hydrochloric acid/ethyl acetate (25 ml) was added thereto, and the resulting mixture was stirred at room temperature for 1 hour. The solvent was removed under reduced pressure, then ethyl acetate (30 ml) was added thereto, and the resulting mixture was stirred for 5 minutes. The obtained white solid was filtered and dried under reduced pressure to obtain the title compound (5.77 g, 87%) as a hydrochloride.


20

¹H NMR (400 MHz, CD₃OD): δ 9.29 (s, 1H), 8.75 (d, J = 6.0 Hz, 1H), 8.61 (d, J = 2.0 Hz, 1H), 8.58 (d, J = 5.6 Hz, 1H), 8.11 (dd, J = 8.0, 2.4 Hz, 1H), 7.61 (d, J = 8.0 Hz, 1H), 4.87 - 4.83 (m, 1H), 4.47 (t, J = 7.6 Hz, 1H), 3.73 (s, 3H), 3.40 - 3.38 (m, 2H), 1.87 - 1.29 (m, 10H); MS (ESI) m/z 424 (M+H)⁺

25 [0061]

Example 2: Synthesis of M-2

(Step 1) 5-Iodine-6-methyl-1H-pyrimidine-2,4-dione




Iodobenzene diacetate (38.0 g, 119 mmol) was dissolved in dichloromethane (100 ml), iodine (5.00 g, 19.8 mmol) was added thereto, and the resulting mixture was stirred for 30 minutes. 6-Methyl-1H-pyrimidine-2,4-dione (5.01 g, 39.7 mmol) was added to the reaction solution, and the resulting mixture was stirred. The reaction solution was concentrated under reduced pressure and then purified by silica gel column chromatography (petroleum ether:ethyl acetate = 1:1 to 1:3) to obtain the title compound (3.50 g, 35%).

¹H NMR (DMSO-d₆, 300 MHz): δ 11.3 (s, 2H), 2.27 (s, 3H).

[0062]

(Step 2) 5-Iodine-1,3,6-trimethyl-1H-pyrimidine-2,4-dione



5-Iodine-6-methyl-1H-pyrimidine-2,4-dione (3.50 g, 13.9 mmol) <see (step 1)> and potassium carbonate (5.70 g, 41.7 mmol) were dissolved in N,N-dimethylformamide (50 ml), iodomethane (5.02 g, 34.7 mmol) was added thereto, and the resulting mixture was stirred at room temperature for 12 hours. The reaction liquid was diluted with water and then extracted with ethyl acetate (50 ml × 3). The organic layers were combined, then washed with a saturated saline solution and dried over magnesium sulfate. The solvent was removed under reduced pressure to obtain the title compound (3.15 g, 81%).

¹H NMR (DMSO-d₆, 300 MHz): δ 3.41 (s, 3H), 3.22 (s, 3H), 2.61 (s, 3H)


[0063]

(Step

3)

Methyl

N-(tert-butoxycarbonyl)-O-[(trifluoromethyl)sulfonyl]-L-tyrosinate



Methyl N-(tert-butoxycarbonyl)-L-tyrosinate (60.0 g, 203 mmol) was
 5 dissolved in tetrahydrofuran, N-phenylbis(trifluoromethanesulfonamide)
 (79.8 g, 224 mmol) and diisopropylethylamine (90.0 g, 714 mmol) were added
 thereto, and the resulting mixture was stirred at room temperature for 12
 hours. The reaction solution was concentrated under reduced pressure,
 then the obtained residue was purified by silica gel column chromatography
 10 (petroleum ether:ethyl acetate = 3:1) to obtain the title compound (74.0 g,
 85%).

¹H NMR (CDCl₃, 400 MHz): δ 7.25 - 7.19 (m, 4H), 5.08 - 5.06(m, 1H), 4.61 -
 4.59 (m, 1H), 3.71 (3H, s), 3.19 - 3.15 (m, 1H), 3.06 - 3.01(m, 1H), 1.41 (s, 9H);
 MS (ESI) m/z 328 (M+H)⁺


15 [0064]

(Step

4)

Methyl

N-(tert-butoxycarbonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-L-ph
 enylalaninate



20 Methyl

N-(tert-butoxycarbonyl)-O-[(trifluoromethyl)sulfonyl]-L-tyrosinate <see (step
 3)> (74.0 g, 173 mmol) and bis(pinacolato)diborane (66.0 g, 260 mmol) were
 dissolved in N,N-dimethylformamide (1.0 L).
 [1,1'-bis(Diphenylphosphino)ferrocene]dichloropalladium(II) (13.6 g, 18.6
 25 mmol) and potassium acetate (60.0 g, 612 mmol) were added to the solution,


and the resulting mixture was stirred at 95°C for 12 hours. The reaction mixture was cooled to room temperature, then the reaction solution was filtered through Celite, and the filtrate was concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 4:1) to obtain the title compound (45.0 g, 65%).

¹H NMR (CDCl₃, 400 MHz): δ 7.73 (d, J = 8 Hz, 2H), 7.12 (d, J = 8 Hz, 2H), 4.59 - 4.57 (m, 1H), 3.70 (3H, s), 3.11 - 3.06 (m, 2H), 1.41 (s, 9H), 1.13 (s, 12H); MS (ESI) m/z 306 (M+H)⁺

[0065]

(Step 5)

[4-[(2S)-2-(tert-Butoxycarbonylamino)-3-methoxy-3-oxo-propyl]phenyl]boronic acid



Methyl

N-(tert-butoxycarbonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-L-phenylalaninate <see (step 4)> (67.0 g, 165 mmol) was dissolved in acetone (700 ml), sodium periodate (71.0 g, 330 mmol), ammonium acetate (25.0 g, 330 mmol), and water (300 ml) were added thereto, and the resulting mixture was stirred at room temperature for 55 hours. The reaction solution was filtered through Celite, then the filtrate was concentrated under reduced pressure, ethyl acetate was added, washed with water and a saturated saline solution, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the precipitated solid was washed with a mixed solvent of petroleum ether:ethyl acetate (1:10) and then dried to obtain the title compound (29.0 g, 55%).


¹H NMR (CDCl₃, 400 MHz): δ: 7.71 - 7.56 (m, 2H), 7.25 - 7.16 (m, 2H), 4.39 -

4.37 (m, 1H), 3.71 (s, 3H), 3.14 - 3.10 (m, 1H), 2.96 - 2.90 (m, 1H), 1.40 (s, 9H); MS (ESI) m/z 224 (M+H)⁺

[0066]

(Step 6) Methyl

5 (2S)-2-(tert-butoxycarbonylamino)-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoate



5-Iodine-1,3,6-trimethyl-1H-pyrimidine-2,4-dione <see (step 2)> (0.52

g, 1.8 mmol),

10 [4-[(2S)-2-(tert-butoxycarbonylamino)-3-methoxy-3-oxo-propyl]phenyl]boronic acid <see (step 5)> (0.50 g, 1.5 mmol), sodium carbonate (0.49 g, 4.65 mmol), and Pd(dppf)Cl₂ (57 mg, 0.078 mmol) were dissolved in dioxane (100 ml) and water (3.0 ml), and the resulting mixture was stirred in the presence of nitrogen gas at 100°C for 3 hours. The reaction solution was concentrated

15 under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 3:1 to 1:1) to obtain the title compound (0.35 g, 51%).


¹H NMR (CDCl₃, 300 MHz): δ 7.20 - 7.13 (m, 4H), 5.04 - 5.02 (m, 1H), 4.63 - 4.60 (m, 1H), 3.74 (s, 3H), 3.51 (s, 3H), 3.41 (s, 3H), 3.14 - 3.07 (m, 2H), 2.05

20 (s, 3H), 1.35 (s, 9H); MS (ESI) m/z 432 (M+H)⁺

[0067]

(Step 7) Methyl

(2S)-2-amino-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoate (M-2)



Methyl


(2S)-2-(tert-butoxycarbonylamino)-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoate <see step 6> (1.30 g, 2.80 mmol) was dissolved in dichloromethane (10 ml), the reaction solution was cooled to 0°C, then 3 N hydrochloric acid/dichloromethane (10.0 ml, 30.0 mmol) was added thereto, and the resulting mixture was stirred at room temperature for 1 hour. The reaction solution was concentrated under reduced pressure, then dichloromethane (10.0 ml) was added thereto, and the resulting mixture was stirred at room temperature for 30 minutes. The precipitated solid was filtered and dried to obtain the title compound (0.84 g, 82%) as a white solid.

¹H NMR (CD₃OD, 400 MHz): δ 7.35 (d, J = 10.0 Hz, 2H), 7.24 (t, J = 11.2 Hz, 2H), 4.39 (dd, J = 11.2, 7.6 Hz, 1H), 3.88 (s, 3H), 3.53 (s, 3H), 3.36 (s, 3H), 3.34 - 3.15 (m, 2H), 2.21 (s, 3H); MS (ESI) m/z 332 (M+H)⁺

[0068]

Example 3: Synthesis of M-3

(Step 1) 5-(Methylaminomethyl)-2-nitro-benzoic acid




5-Methyl-2-nitro-benzoic acid (50.0 g, 276 mmol) and 5-dimethylhydantoin (36.6 g, 128 mmol) were suspended in chlorobenzene (100 ml), and the suspension was heated at 40°C in the presence of argon gas. Azobis(2-methylpropionitrile) (1.26 g, 7.69 mmol) was added thereto, and the

resulting mixture was stirred at 80°C for 1 hour. The reaction liquid was cooled to 40°C once, then 5-dimethylhydantoin (18.3 g, 64.1 mmol) and azobis(2-methylpropionitrile) (1.26 g, 7.69 mmol) were added again, and the resulting mixture was stirred at 80°C for 2 hours. The reaction liquid was cooled to 10°C, and then the resulting mixture was stirred for 12 hours. The crystallized solid was filtered off, the obtained filtrate was added dropwise to a 40% methylamine/methanol solution (260 ml, 2.56 mol) that had been cooled to 10°C over 80 minutes, and the resulting mixture was stirred at 25°C. The reaction liquid was concentrated under reduced pressure, then 2-propanol (100 ml) was added thereto, the resulting mixture was stirred at 50°C for 1 hour and then cooled to 9°C, and the precipitated solid was filtered off, washed with 2-propanol (100 ml) and dried to obtain the title compound (42.2 g, 52%) as a white solid.

¹H NMR (400 MHz, D₂O): δ 8.03 (d, J = 8.44 Hz, 1H), 7.55 (dd, J = 8.44 Hz, 1.92 Hz, 1H), 7.49 (t, 1H), 4.25 (s, 2H), 2.70 (s, 3H); MS (ESI) m/z 210.8 (M+H)⁺

[0069]

(Step 2) 5-[[Formyl(methyl)aminomethyl]-2-nitro-benzoic acid




5-(Methylaminomethyl)-2-nitro-benzoic acid <see (step 1)> (5.58 g) and sodium formate (0.65 g, 9.52 mmol) were dissolved in formic acid (4.00 ml) and acetic anhydride (2.70 mmol, 14.3 mmol), and the resulting mixture was stirred at room temperature for 1 hour. The reaction liquid was diluted with water (32 ml), a crystal was precipitated, and a slurry was obtained. The slurry was stirred at 9°C for 12 hours, and the precipitated solid was filtered off and dried under reduced pressure to obtain the title compound (3.95 g, 87%) as a white solid.

¹H-NMR (400 MHz, DMSO-*d*₆): δ8.32 (s, 1H), 8.19 (s, 1H), 8.01 (d, *J* = 8.24 Hz, 1H), 7.98 (d, *J* = 8.24 Hz, 1H), 7.73 (d, 1H, *J* = 1.88 Hz), 7.66 (d, *J* = 1.92 Hz, 1H), 7.65 (dd, *J* = 8.24 Hz, 1.72 Hz, 1H), 7.58 (dd, *J* = 8.24 Hz, 1.92 Hz, 1H), 4.62 (s, 2H), 4.57 (s, 2H), 2.89 (s, 3H), 2.65 (s, 3H); MS (ESI) *m/z* 239

(M+H)⁺

[0070]

(Step 3) 2-Amino-5-[[formyl(methyl)amino]methyl]benzoic acid



5-[[Formyl(methyl)amino]methyl]-2-nitro-benzoic acid <see (step 2)>

(6.96 g, 25.19 mmol) was suspended in methanol (48.0 ml), and a 6 N aqueous solution of sodium hydroxide (3.5 ml) was added thereto and dissolved. 5% Pd/C (1.03 g, 0.250 mmol) was added, and the resulting mixture was stirred in the presence of hydrogen gas at 40°C for 5 hours. The reaction liquid was cooled to room temperature and filtered, and the solvent was removed under reduced pressure. Water (48 ml) and a 2 N aqueous solution of hydrochloric acid (11.0 ml) were added to the obtained residue, the precipitated solid was filtered off, washed with water (24.0 ml), and dried under reduced pressure to obtain the title compound (4.84 g, 92.3%) as a white solid.

¹H NMR (400 MHz, DMSO-*d*₆): δ8.49 (bs, 2H), 8.25 (s, PhCCH₂NMeCH_aO), 8.08 (s, PhCCH₂NMeCH_bO), 7.60 and 7.59 (s, 1H), 7.12 (m, 1H), 6.74 (m, 1H), 4.27 (s, 2H), 2.77 and 2.58 (s, 3H); MS (ESI) *m/z* 231 (M+H)⁺


[0071]

(Step

4)

Methyl


(2S)-2-[tert-butoxycarbonylamino]-3-[4-[6-[[formyl(methyl)amino]methyl]-1-methyl-2,4-dioxo-quinazolin-3-yl]phenyl]propanoate



Carbodiimidazole (16.4 g, 101 mmol) was dissolved in N,N-dimethylformamide (50 ml), methyl (2S)-3-(4-aminophenyl)-2-(tert-butoxycarbonylamino)propanoate (25.0 g, 84.9 mmol) and 2-amino-5-[[formyl(methyl)amino]methyl]benzoic acid <see (step 3)> (17.6 g, 84.5 mmol) were added thereto, and the resulting mixture was stirred at 65°C. The reaction liquid was cooled to room temperature, then carbodiimidazole (16.4 g, 101 mmol) was added again, and the resulting mixture was stirred at 65°C. After the reaction was completed, the reaction liquid was diluted with water (2,000 ml) and extracted with methylene chloride (500 ml, 100 ml). The extracts were combined, washed with a saturated saline solution, and dried. The obtained compound was used in (step 5).

[0072]

(Step 5) Methyl (2S)-2-amino-3-[4-[6-[[formyl(methyl)amino]methyl]-1-methyl-2,4-dioxo-quinazolin-3-yl]phenyl]propanoate (M-3)



The compound obtained in (step 4) was dissolved in N,N-dimethylformamide (150 ml), potassium carbonate (21.7 g) and methyl p-toluenesulfonate (17.8 ml) was added thereto, and the resulting mixture was stirred at 40°C for 4 hours. Acetic acid (9.0 ml) was added, then the


reaction liquid was diluted with water (1,500 ml) and extracted with methylene chloride (1,000 ml), and the solvent was removed under reduced pressure. The obtained residue was purified by silica gel column chromatography (methylene chloride:methanol = 9:1). The obtained
 5 compound was dissolved in isopropyl alcohol (150 ml), 4 N hydrochloric acid/dioxane (150 ml) was added thereto, and the resulting mixture was stirred at room temperature for 1 hour. The precipitated solid was filtered, washed with isopropyl alcohol (100 ml) and dried to obtain a hydrochloride of the title compound (28.0 g).

10 ¹H NMR (400 MHz, DMSO-d₆) δ 8.70 (s, 3H), 8.35 (s, PhCCH₂NMeCH_aO), 8.16 (s, PhCCH₂NMeCH_bO), 7.93 (dd, J = 15.8, 2.1 Hz, 1H), 7.77 - 7.66 (m, 1H), 7.53 (dd, J = 11.7, 8.6 Hz, 1H), 7.38 - 7.33 (m, 2H), 7.30 - 7.25 (m, 2H), 4.55 and 4.53 (s, 2H), 4.36 (dd, J = 7.3, 6.0 Hz, 1H), 3.70 (s, 3H), 3.54 and 3.53 (s, 3H), 3.33 (s, 3H), 3.25 (dd, J = 14.1, 5.9 Hz, 1H), 3.16 (dd, J = 14.1, 7.3 Hz,
 15 1H); MS (ESI) m/z 425.48 (M+H)⁺

[0073]

Example 4: Synthesis of M-4

(Step 1) 2-Chloro-5-nitro-pyridine-4-carboxylic acid




20 2-Chloro-4-methyl-5-nitro-pyridine (20.5 g, 119 mmol) was dissolved in concentrated sulfuric acid (200 ml), the resulting solution was cooled to 0°C, then chromium(VI) oxide (40.0 g, 400 mmol) was added thereto, and the resulting mixture was stirred at 0°C for 1 hour and then at room temperature for 12 hours. The reaction liquid was diluted with water
 25 (2,000 ml), and the precipitated solid was filtered and dried to obtain the title compound (18.0 g, 75%).

¹H NMR (400 MHz, CD₃OD): δ 10.8 (br, s, 1H), 9.13 (s, 1H), 7.70 (s, 1H).

[0074]


(Step 2) Methyl 2-chloro-5-nitro-pyridine-4-carboxylate



2-Chloro-5-nitro-pyridine-4-carboxylic acid <see (step 1)> (20.2 g, 100
 5 mmol) was dissolved in ethyl acetate (100 ml), the resulting solution was
 cooled to 0°C, then a diazomethane/diethyl ether solution was added thereto,
 and the resulting mixture was stirred at 0°C for 1 hour. The reaction liquid
 was concentrated, then the obtained residue was purified by silica gel
 column chromatography (petroleum ether:ethyl acetate = 6:1 to 2:1) to obtain
 10 the title compound (20.0 g, 92%).

[0075]


(Step 3) Methyl 2-methoxy-5-nitro-pyridine-4-carboxylate



Methyl 2-chloro-5-nitro-pyridine-4-carboxylate <see (step 2)> (10.8 g,
 15 50.0 mmol) was dissolved in methanol (100 ml), sodium methoxide (8.10 g,
 150 mmol) was added thereto, and the resulting mixture was stirred at 65°C
 for 4 hours. The reaction liquid was concentrated under reduced pressure,
 and then the obtained residue was purified by silica gel column
 chromatography (petroleum ether:ethyl acetate = 8:1 to 4:1) to obtain the
 20 title compound (8.0 g, 75%).

[0076]

(Step 4) Methyl 5-amino-2-methoxy-pyridine-4-carboxylate




Methyl 2-methoxy-5-nitro-pyridine-4-carboxylate <see (step 3)> (8.0 g, 37 mmol) was dissolved in methanol (100 ml), a saturated aqueous solution of ammonium chloride (50 ml) and iron (10.6 g, 175 mmol) were added thereto, and the resulting mixture was stirred at 65°C for 2 hours. The reaction liquid was cooled to room temperature, then the pH was adjusted to > 7.0 with an aqueous solution of sodium hydrogen carbonate, and the reaction solution was filtered and extracted with ethyl acetate (100 ml × 3). The extracts were combined, washed with a saturated saline solution, dried over anhydrous sodium sulfate, and then the solvent was removed under reduced pressure. The obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 4:1 to 1:1) to obtain the title compound (5.80 g, 84%).

¹H NMR (400 MHz, CDCl₃): δ 7.80 (s, 1H), 7.15 (s, 1H), 5.10 (s, 2H), 3.90 (s, 3H), 3.84 (s, 3H).

[0077]

(Step 5)

3-(6-Bromo-3-pyridyl)-6-methoxy-1-methyl-pyrido[3,4-d]pyrimidine-2,4-dione



20

Two steps similar to Example 1 (step 1), (step 2) were performed to methyl 5-amino-2-methoxy-pyridine-4-carboxylate <see (step 4)> (21.0 g, 115 mmol) and 6-bromopyridin-3-amine (19.9 g, 115 mmol) to obtain the title compound (12.9 g, 31%).

¹H NMR (400 MHz, DMSO-d₆): δ 8.58 (s, 1H), 8.40 (s, 1H), 7.87 - 7.78 (m, 2H),


25

7.30 (s, 1H), 3.93 (s, 3H), 3.57 (s, 3H).

[0078]

(Step 6) Methyl (2S)-2-(tert-butoxycarbonylamino)-3-[5-(6-methoxy-1-methyl-2,4-dioxo-pyrid

5 o[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate




A method similar to Example 1 (step 6) was performed to 3-(6-bromo-3-pyridyl)-6-methoxy-1-methyl-pyrido[3,4-d]pyrimidine-2,4-dione <see (step 5)> (12.5 g, 34.6 mmol) to obtain the title compound (10.0 g, 60%).

10 ¹H NMR (400 MHz, DMSO-d₆): δ 8.57 (s, 1H), 8.45 (s, 1H), 7.43 (d, J = 17.2 Hz, 1H), 7.34 (d, J = 8 Hz, 1H), 7.29 (s, 1H), 4.55 - 4.49 (m, 1H), 3.93 (s, 3H), 3.62 (s, 3H), 3.56 (s, 3H), 3.33 - 3.07 (m, 2H), 1.37 (s, 9H).

[0079]

(Step 7) Methyl (2S)-2-amino-3-[5-(6-methoxy-1-methyl-2,4

15 -dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate (M-4)



A method similar to Example 1 (step 7) was performed to methyl (2S)-2-(tert-butoxycarbonylamino)-3-[5-(6-methoxy-1-methyl-2,4-dioxo-pyrid

o[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate <see (step 6)> (10.0 g, 20.0


20 mmol) to obtain a hydrochloride of the title compound. A saturated aqueous solution of sodium hydrogen carbonate (100 ml) was added to the obtained hydrochloride, then the resulting mixture was extracted with ethyl acetate

(100 ml $\times$ 2), the extracts were combined, the resulting mixture was washed with a saturated saline solution, dried over anhydrous sodium sulfate, and then the solvent was removed under reduced pressure to obtain the title compound (4.20 g, 53%).

5 ^1H NMR (400 MHz, CD_3OD): δ 8.49 (s, 1H), 8.46 (d, $J = 2.0$ Hz, 1H), 7.75 (dd, $J = 8.4, 2.4$ Hz, 1H), 7.49 (d, $J = 8.4$ Hz, 1H), 7.41 (s, 1H), 4.02 - 3.98 (m, 4H), 3.73 (s, 3H), 3.67 (s, 3H), 3.33 - 3.30 (m, 1H), 3.22 - 3.16 (m, 1H); MS (ESI) m/z 386 ($\text{M}+\text{H}$)⁺
[0080]

10 Example 5: Synthesis of M-5

(Step 1) Methyl N-(tert-butoxycarbonyl)-4-(3-methyl-2,6-dioxo-3,6-dihydropyrimidine-1(2H)-yl)-L-phenylalaninate




15 Pyrimidine-2,4(1H,3H)-dione (30.0 g, 268 mmol), 1,1,1,3,3,3-hexamethyldisilazane (550 ml), and trimethylchlorosilane (55 ml) were stirred at 130°C for 5 hours. The reaction solution was cooled to 60°C, then methyl iodide (200 ml) was added thereto, and the resulting mixture was stirred at 60°C for 30 hours. The reaction solution was cooled to 0°C,
20 acetic acid (500 ml) was slowly added thereto, and the solvent was removed under reduced pressure. 2-Propanol (1,600 ml) was added to the residue, the resulting mixture was intensely stirred, and then the obtained solid was filtered and washed with water (500 ml) to obtain a white solid (22 g, 67%).
[4-[(2S)-2-(tert-Butoxycarbonylamino)-3-methoxy-3-oxo-propyl]phenyl]boronic acid <see Example 2 (step 5)> (49.0 g, 152 mmol) was dissolved in
25 methylene chloride (500 ml), then the obtained white solid (22.0 g, 175 mmol),

copper(II) acetate (18.0 g, 98.9 mmol), and triethylamine (40 ml) were added thereto, and the resulting mixture was stirred at room temperature for 60 hours. The reaction solution was concentrated under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 1:1) to obtain the title compound (6.0 g, 10%) as a white crystal.

¹H NMR (CDCl₃, 400 MHz): δ 7.20 - 7.15 (m, 3H), 7.08 (d, J = 8.0 Hz, 2H), 5.76 (d, J = 8.0 Hz, 1H), 4.55 - 4.51 (m, 1H), 3.60 (s, 3H), 3.35 (s, 3H), 3.06 (d, J = 4.0 Hz, 1H), 1.36 (s, 9H); MS (ESI) m/z 421 (M+H)⁺

[0081]

(Step 2) Methyl 4-(3-methyl-2,6-dioxo-3,6-dihydropyrimidine-1(2H)-yl)-L-phenylalaninate (M-5)



• HCl

15 Methyl


N-(tert-butoxycarbonyl)-4-(3-methyl-2,6-dioxo-3,6-dihydropyrimidine-1(2H)-yl)-L-phenylalaninate <see (step 1)> (6.00 g, 15.0 mmol) was dissolved in a 3 N hydrochloric acid/ethyl acetate solution (100 ml), and the resulting mixture was stirred at room temperature for 1 hour. The solvent was removed under reduced pressure, then ethyl acetate (50 ml) was added, and the resulting mixture was further stirred at room temperature for 0.5 hours. The suspension was filtered, and the obtained solid was dried to obtain the title compound (4.0 g, 80%) as a white solid.

¹H NMR (CD₃OD, 500 MHz): δ 7.72 (d, J = 8 Hz, 1H), 7.45 (d, J = 8 Hz, 2H), 7.29 (d, J = 8 Hz, 2H), 5.85 (d, J = 8 Hz, 1H), 4.44 - 4.41 (m, 1H), 3.91 (s, 3H), 3.49 - 3.45 (m, 4H), 3.23 - 3.17 (m, 1H); MS (ESI) m/z 304 (M+H)⁺

[0082]

Example 6: Synthesis of M-6

(Step 1) Methyl 4-aminopyrimidine-5-carboxylate




5 4-Aminopyrimidine-5-carboxylic acid (10.0 g, 71.9 mmol) was dissolved in methanol (100 ml), the reaction solution was cooled to 0°C, then concentrated sulfuric acid (25 ml) was slowly added dropwise, and the resulting mixture was stirred at 85°C for 12 hours. The reaction liquid was concentrated under reduced pressure, then the resulting solution was
 10 diluted with water, and sodium hydrogen carbonate (10.0 g) was added. The resulting mixture was extracted with ethyl acetate (80 ml × 4), the extracts were combined, washed with a saturated saline solution, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, the obtained residue was stirred in a mixed solvent (petroleum
 15 ether:ethyl acetate = 8:1), and the precipitated solid was filtered and dried to obtain the title compound (9.90 g, 90%).

¹H-NMR (400 MHz, DMSO-d₆): δ 8.72 (s, 1H), 8.53 (s, 1H), 8.04 (s, 1H), 7.62 (s, 1H), 3.83 (s, 3H); MS (ESI) m/z 154 (M+H)⁺

[0083]

20 (Step 2)
 3-(6-Bromo-3-pyridyl)-1-methyl-pyrimido[4,5-d]pyrimidine-2,4-dione




Two steps similar to Example 1 (step 1), (step 2) were performed (methyl iodide was used instead of p-toluenesulfonic acid methyl ester) to
 25 methyl 4-aminopyrimidine-5-carboxylate <see (step 1)> (4.50 g, 29.4 mmol)

and 6-bromopyridin-3-amine (4.20 g, 24.7 mmol) to obtain the title compound (3.10 g, 38%).

¹H NMR (400 MHz, DMSO-d₆): δ 9.30 (s, 1H), 9.23 (s, 1H), 8.40 (s, 1H), 7.89 (d, J = 11.2 Hz, 1H), 7.81-7.77 (m, 1H), 3.57 (s, 3H).

5 [0084]

(Step 3) Methyl (2S)-2-(tert-butoxycarbonylamino)-3-[5-(1-methyl-2,4-dioxo-pyrimido[4,5-d]pyrimidin-3-yl)-2-pyridyl]propanoate



10 A method similar to Example 1 (step 6) was performed to 3-(6-bromo-3-pyridyl)-1-methyl-pyrimido[4,5-d]pyrimidine-2,4-dione <see (step 2)> (1.50 g, 4.50 mmol) to obtain the title compound (1.33 g, 65%).


¹H NMR (400 MHz, DMSO-d₆): δ 9.29 (s, 1H), 9.20 (s, 1H), 8.44 (s, 1H), 7.73 - 7.70 (m, 1H), 7.44 (d, J = 8.0 Hz, 1H), 7.38 (d, J = 7.6 Hz, 1H), 4.55 - 4.50 (m, 1H), 3.62 (s, 3H), 3.56 (s, 3H), 3.22 - 3.06 (m, 2H), 1.36 (s, 9H); MS (ESI) m/z 457 (M+H)⁺

15

[0085]

(Step 4) Methyl (2S)-2-amino-3-[5-(1-methyl-2,4-dioxo-pyrimido[4,5-d]pyrimidin-3-yl)-2-pyridyl]propanoate (M-6)

20




A method similar to Example 1 (step 7) was performed to methyl

(2S)-2-(tert-butoxycarbonylamino)-3-[5-(1-methyl-2,4-dioxo-pyrimido[4,5-d]p
 yrimidin-3-yl)-2-pyridyl]propanoate <see (step 3)> (1.30 g, 2.85 mmol) to
 obtain a hydrochloride of the title compound (1.10 g, 98%).

¹H NMR (400 MHz, DMSO-d₆): δ 9.36 (s, 1H), 9.33 (s, 1H), 8.70 (d, J = 2.4 Hz,
 1H), 8.09 (dd, J = 8.4, 2.4 Hz, 1H), 7.78 (d, J = 8.4 Hz, 1H), 4.70 (t, J = 6.4 Hz,
 1H), 3.86 (s, 3H), 3.76 (s, 3H), 3.68 - 3.65 (m, 2H); MS (ESI) m/z 357 (M+H)⁺
 [0086]

Example 7: Synthesis of M-7

(Step 1) Methyl
 10 (2S)-2-(tert-butoxycarbonylamino)-3-[4-[2,4-dioxo-6-(trifluoromethyl)-1H-pyr
 imidin-3-yl]phenyl]propanoate



6-(Trifluoromethyl)-1H-pyrimidine-2,4-dione (1.00 g, 5.60 mmol) and
 [4-[(2S)-2-(tert-butoxycarbonylamino)-3-methoxy-3-oxo-propyl]phenyl]boroni
 15 c acid <see Example 2 (step 5)> (2.00 g, 6.20 mmol) were dissolved in
 methylene chloride (10 ml), copper(II) acetate (3.01 g, 15.1 mmol) and
 triethylamine (2.40 g, 2.40 mmol) were added thereto, and the resulting
 mixture was stirred at room temperature for 12 hours. The reaction liquid
 was filtered and the filtrate was concentrated under reduced pressure. The
 20 obtained residue was purified by silica gel column chromatography
 (petroleum ether:ethyl acetate = 5:1 to 1:1) to obtain the title compound (750
 mg, 29%) as a yellow crystal.


¹H NMR (CD₃OD, 400 MHz): δ 7.26 (d, J = 8.0 Hz, 2H), 7.22 (d, J = 8.0 Hz,
 2H), 6.27 (s, 1H), 4.45 - 4.41 (m, 1H), 3.73 (s, 3H), 3.21 - 3.16 (m, 1H), 3.07 -
 25 3.02 (m, 1H), 1.43 (s, 9H)
 [0087]

(Step

2)

Methyl

(2S)-2-(tert-butoxycarbonylamino)-3-[4-[3-methyl-2,6-dioxo-4-(trifluoromethyl)pyrimidin-1-yl]phenyl]propanoate



5

Methyl

(2S)-2-(tert-butoxycarbonylamino)-3-[4-[2,4-dioxo-6-(trifluoromethyl)-1H-pyrimidin-3-yl]phenyl]propanoate <see (step 1)> (700 mg, 1.53 mmol) was dissolved in N,N-dimethylformamide (15 ml), potassium carbonate (400 mg, 2.90 mmol) and methyl iodide (6.60 g, 46.0 mmol) were added thereto, and the resulting mixture was stirred at room temperature for 90 minutes. The reaction liquid was concentrated under reduced pressure, diluted with water (20 ml), and then extracted with ethyl acetate. The extracts were combined, washed with a saturated saline solution, dried over anhydrous sodium sulfate, and the solvent was removed under reduced pressure to obtain the title compound (700 mg, 97%) as a yellow solid.

15

¹H NMR (CD₃OD, 400 MHz): δ 7.37 (d, J = 8.4 Hz, 2H), 7.21 (d, J = 8.0 Hz, 2H), 6.42 (s, 1H), 4.45 - 4.42 (m, 1H), 3.73 (s, 3H), 3.34 (s, 3H), 3.21 - 3.16 (m, 1H), 3.07 - 3.02 (m, 1H), 1.43 (s, 9H)

[0088]


20

(Step

3)

Methyl

(2S)-2-amino-3-[4-[3-methyl-2,6-dioxo-4-(trifluoromethyl)pyrimidin-1-yl]phenyl]propanoate (M-7)




Methyl

(2S)-2-(tert-butoxycarbonylamino)-3-[4-[3-methyl-2,6-dioxo-4-(trifluoromethyl)pyrimidin-1-yl]phenyl]propanoate <see (step 2)> (604 mg, 1.28 mmol) was
 5 dissolved in 6 N hydrochloric acid/ethyl acetate (10.0 ml), and the resulting solution was stirred at room temperature for 1 hour. The reaction liquid was concentrated under reduced pressure, ethyl acetate (30.0 ml) was added to the obtained residue, and the resulting mixture was stirred at room temperature for 30 minutes. The precipitated solid was filtered and dried
 10 to obtain a hydrochloride of the title compound (470 mg, 90%) as a white solid.

¹H NMR (CD₃OD, 400 MHz): δ 7.34 (d, J = 8.4 Hz, 2H), 7.20 (d, J = 8.4 Hz, 2H), 6.34 (s, 1H), 4.33 - 4.29 (m, 1H), 3.78 (s, 3H), 3.43 (s, 3H), 3.36 - 3.31 (m, 1H), 3.13 - 3.07 (m, 1H); MS (ESI) m/z 372 (M+H)⁺
 15 [0089]

Example 8: Synthesis of M-8

(Step 1) Methyl
 N-(tert-butoxycarbonyl)-4-(2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)-L-phenylalaninate



20


Methyl 3-aminoisonicotinate (10.0 g, 65.0 mmol) was dissolved in methylene chloride (100 ml), diisopropylethylamine (17.0 g, 130 mmol) and

triphosgene (6.40 g, 22.0 mmol) were added thereto, and the resulting mixture was stirred at 0°C for 3 hours. Methyl 4-amino-N-(tert-butoxycarbonyl)-L-phenylalaninate (19.0 g, 65.0 mmol) was added to the solution, and then the resulting mixture was stirred for 12 hours while the temperature was slowly raised from 0°C to room temperature. The reaction solution was concentrated under reduced pressure, then ethyl acetate was added, and the resulting solution was washed with water and a saturated saline solution and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the resulting product was purified by silica gel column chromatography (petroleum ether/ethyl acetate = 1:1). The obtained compound was dissolved in N,N-dimethylformamide (200 ml), an aqueous solution of potassium carbonate (1.20 g, 8.70 mmol) was added thereto, and the resulting mixture was stirred at room temperature for 3 hours. The reaction solution was concentrated under reduced pressure, then ethyl acetate was added, and the resulting solution was washed with water and a saturated saline solution and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 1:1) to obtain the title compound (12.0 g, 43% for two steps).

¹H NMR (CD₃Cl₃, 400 MHz): δ 10.07 (1H, s), 8.54 (d, J = 5.2 Hz, 1H), 8.14 (s, 1H), 7.92 (d, J = 8.8 Hz, 1H), 7.35 (d, J = 7.6 Hz, 2H), 7.23 (d, J = 8.0 Hz, 2H), 6.12 - 6.10 (m, 1H), 4.73 - 4.69 (m, 1H), 3.71 (s, 3H), 3.22 - 3.21 (m 2H), 1.44 (s, 9H)

[0090]

(Step 2) Methyl N-(tert-butoxycarbonyl)-4-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)-L-phenylalaninate



Methyl

N-(tert-butoxycarbonyl)-4-(2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)-L-phenylalaninate <see (step 1)> (8.00 g, 18.5 mmol) was dissolved in
 5 N,N-dimethylformamide (100 ml), potassium carbonate (5.10 g, 37.0 mmol) and methyl iodide (1.9 ml, 37 mmol) were added thereto, and the resulting mixture was stirred at room temperature for 3 hours. The suspension was filtered, and the filtrate was concentrated under reduced pressure and then purified by silica gel column chromatography (petroleum ether:ethyl acetate
 10 = 1:1) to obtain the title compound (3.5 g, 43%).

¹H NMR (CD₃Cl₃, 400 MHz): δ 8.81 (1H, s), 8.60 (d, J = 4.8 Hz, 1H), 8.03 (d, J = 5.6 Hz, 1H), 7.30 (d, J = 8.0 Hz, 2H), 7.19 (d, J = 8.4 Hz, 2H), 5.05 - 5.03 (m, 1H), 4.65 - 4.63 (m, 1H), 3.73 (s, 3H), 3.71 (s, 3H), 3.18 - 3.16 (m 2H), 1.44 (s, 9H); MS (ESI) m/z 355 (M+H·Boc)⁺


15 [0091]

(Step

3)

Methyl

4-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)-L-phenylalaninate (M-8)



• HCl

20 Methyl

N-(tert-butoxycarbonyl)-4-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrim


idin-3(2H)-yl)-L-phenylalaninate (3.5 g, 7.7 mmol) was dissolved in a 4 N hydrochloric acid/ethyl acetate solution (40 ml), and the resulting mixture was stirred at room temperature for 1 hour. The reaction liquid was filtered, then the obtained solid was dried under reduced pressure to obtain the title compound (3.0 g, 94%).

¹H NMR (CD₃OD, 400 MHz): δ 9.28 (s, 1H), 8.76 (d, J = 0.8 Hz, 1H), 8.56 (d, J = 1.6 Hz, 1H), 7.51 (d, J = 8.4 Hz, 2H), 7.41 (d, J = 8.4 Hz, 2H), 4.48 - 4.44 (m, 1H), 3.90 (s, 3H), 3.76 (s, 3H), 3.49 - 3.44 (m 1H), 3.30 - 3.24 (m, 1H); MS (ESI) m/z 355 (M+H)⁺

[0092]

Example 9: Synthesis of M-9

(Step 1) 6-Bromopyridin-3-amine



2-Bromo-5-nitro-pyridine (202 g, 1.0 mol) was dissolved in methanol (2.0 L), a saturated aqueous solution of ammonium chloride (2.0 L) was added thereto, the resulting mixture was stirred at 50°C, then iron (224 g, 4.0 mol) was slowly added thereto, and the resulting mixture was stirred at 50°C for 6 hours. The reaction solution was cooled to room temperature and then filtered and washed with ethyl acetate. The filtrate was diluted with water and then extracted with ethyl acetate (1.0 L × 6). The extracts were combined, washed with a saturated aqueous solution of sodium hydrogen carbonate (2.0 L) and a saturated saline solution, dried over anhydrous sodium sulfate, and then the solvent was removed under reduced pressure. The obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 1:1) to obtain the title compound (126 g, 73%).


¹H NMR (400 MHz, DMSO-d₆): δ 7.69 (d, J = 2.8 Hz, 1H), 7.19 (d, J = 8.4 Hz, 1H), 6.90 (dd, J = 8.4 Hz, 2.8 Hz, 1H), 5.50 (s, 2H)

[0093]

(Step

2)

3-(6-Bromo-3-pyridyl)-1-methyl-pyrido[3,4-d]pyrimidine-2,4-dione



5 A method similar to Example 1 (step 1), (step 2) was performed to 6-bromopyridin-3-amine <see (step 1)> (67.7 g, 394 mmol) to obtain the title compound (32.5 g, 24.8% for two steps).

1H NMR (400 MHz, DMSO-d₆): δ 9.00 (s, 1H), 8.59 (d, J = 8.8 Hz, 1H), 8.41 (d, J = 4.8 Hz, 1H), 7.92 (d, J = 4.8 Hz, 1H), 7.86 (d, J = 8.0 Hz, 1H), 7.80 (dd, J = 8.4 Hz, 2.0 Hz, 1H), 3.62 (s, 3H)

10


[0094]

(Step

3)

Cyclohexyl

(2S)-2-(tert-butoxycarbonylamino)-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate



15


A method similar to Example 1 (step 6) was performed to 3-(6-bromo-3-pyridyl)-1-methyl-pyrido[3,4-d]pyrimidine-2,4-dione <see (step 2)> (23.0 g, 69.2 mmol) to obtain the title compound (21.7 g, 60%).

1H NMR (400 MHz, DMSO-d₆): δ 8.99 (s, 1H), 8.56 (d, J = 4.8 Hz, 1H), 8.46 (d, J = 1.6 Hz, 1H), 7.91 (d, J = 5.2 Hz, 1H), 7.72 (dd, J = 8.0, 2.4 Hz, 1H), 7.42 (d, J = 8.0 Hz, 1H), 7.31 (d, J = 8.0 Hz, 1H), 4.67 - 4.65 (m, 1H), 4.48 - 4.47 (m, 1H), 3.61 (s, 3H), 3.17 - 3.13 (m, 2H), 1.71 - 1.25 (m, 19H)

20

[0095]

(Step 4) Cyclohexyl
3-[5-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)pyridin-2-yl]-L-alaninate (M-9)




5 A method similar to Example 1 (step 7) was performed to cyclohexyl (2S)-2-(tert-butoxycarbonylamino)-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate <see (step 3)> (35.0 g, 66.9 mmol) to obtain the title compound (27.9 g, 91%) as a hydrochloride.

¹H NMR (400 MHz, CD₃OD): δ 9.16 (s, 1H), 8.65 (d, J = 5.6 Hz, 1H), 8.52 (d, J = 2.0 Hz, 1H), 8.44 (d, J = 5.6 Hz, 1H), 7.89 (dd, J = 8.4, 2.4 Hz, 1H), 7.61 (d, J = 8.4 Hz, 1H); 4.80 (m, 1H), 4.54 (t, J = 6.0 Hz, 1H), 3.65 (s, 3H), 3.46 - 3.59 (m, 2H), 1.70 - 1.20 (m, 10H); MS (ESI) m/z 424 (M+H)⁺
[0096]

Example 10: Synthesis of M-10

15 (Step 1) tert-Butyl
4-(5-bromopyrimidin-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate




5-Bromo-2-iodine-pyrimidine (210 mg, 0.74 mmol) and tert-butyl 4-(5-bromopyrimidin-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate (275 mg, 0.890 mmol) were dissolved in N,N-dimethylformamide (50 ml) and water (10 ml), potassium carbonate (255 mg, 1.85 mmol) and Pd(PPh₃)₄ (805 mg, 1.10 mmol) were added thereto, and the resulting mixture was stirred in the presence of nitrogen gas at 100°C for 4 hours. The reaction liquid was

concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 5:1 to 1:1) to obtain the title compound (78 mg, 31%).

¹H NMR (300 MHz, CDCl₃): δ 8.72 (s, 2H), 7.23 - 7.20 (m, 1H), 4.19 - 4.17 (m, 2H), 3.66 - 3.62 (m, 2H), 2.72 - 2.66 (m, 2H), 8.72 (s, 9H)
[0097]

(Step 2) 4-Bromo-2,5-difluorobenzoic acid




1,4-Dibromo-2,5-difluorobenzene (51.2 g, 188 mmol) was dissolved in 1,2-diethoxyethane (400 ml), and a 2.5 M n-butyllithium/hexane solution (76.0 ml, 190 mmol) was slowly added thereto dropwise at -78°C in the presence of nitrogen gas. The reaction solution was stirred at -78°C for 30 minutes, then dry ice was added, and the resulting mixture was further stirred for 30 minutes. The temperature was gradually raised to room temperature, and then water (200 ml) was added to the reaction liquid. The reaction liquid was diluted with ethyl acetate, the resulting solution was washed with a 10% aqueous solution of sodium carbonate (200 ml × 2), then the obtained aqueous layers were combined, 1 N hydrochloric acid was added for adjustment to acidic pH, and the precipitated yellow solid was filtered and dried to obtain the title compound (30.0 g, 67%).

¹H NMR (DMSO-d₆, 400 MHz): δ 7.90 - 7.87 (m, 1H), 7.79 - 7.75 (m, 1H); MS (ESI) m/z 191 (M+H-44)⁺

[0098]

(Step 3) Methyl 4-bromo-2,5-difluorobenzoate




4-Bromo-2,5-difluorobenzoic acid <see (step 2)> (30.0 g, 127 mmol) was dissolved in ethyl acetate (500 ml), the reaction solution was cooled to 0°C, then a diazomethane/ether solution was added thereto. The reaction liquid was stirred at 0°C for 1 hour, and the solvent was removed under reduced pressure. The obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 6:1 to 2:1) to obtain the title compound (25.0 g, 78%).

¹H NMR (DMSO-d₆, 400 MHz): δ 7.93 - 7.89 (m, 1H), 7.80 - 7.77 (m, 1H), 3.86 (s, 3H)

[0099]

(Step 4) Methyl 4-amino-2,5-difluorobenzoate



Methyl 4-bromo-2,5-difluorobenzoate <see (step 3)> (25.0 g, 99.6 mmol) and BINAP (1.86 g, 3.00 mmol) were dissolved in toluene (500 ml). Palladium(II) acetate (672 mg, 3.00 mmol), cesium carbonate (52.0 g, 160 mmol), and benzophenone imine (25.3 g, 140 mmol) were added to the solution, and the resulting mixture was stirred at 110°C for 12 hours. The reaction mixture was cooled to room temperature, and then filtered through Celite and washed with ethyl acetate. The obtained filtrate was concentrated under reduced pressure. The obtained compound was dissolved in water (30 ml) and tetrahydrofuran (80 ml), concentrated hydrochloric acid (30 ml) was added thereto, and the resulting mixture was stirred at room temperature for 2 hours. The precipitated solid was filtered

to obtain a white solid. The filtrate was also concentrated under reduced pressure until a white solid was precipitated, the obtained solid was filtered, the white solids were combined, and the resulting product was dried to obtain the title compound (9.8 g, 53% for two steps).

5 ^1H NMR (CD_3OD , 400 MHz): δ 7.53 - 7.49 (m, 1H), 6.60 - 6.56 (m, 1H), 3.85 (s, 3H); MS (ESI) m/z 188 ($\text{M}+\text{H}$)⁺


[0100]

(Step

5)

Methyl

4-[(4-bromophenyl)sulfonylamino]-2,5-difluoro-benzoate




Methyl 4-amino-2,5-difluoro-benzoate <see (step 4)> (24.4 g, 130 mmol) was dissolved in methylene chloride (1.0 L), 4-bromobenzenesulfonyl chloride (50.0 g, 196 mmol) and pyridine (100 ml) were added thereto, and the resulting mixture was stirred at room temperature for 12 hours. The reaction liquid was concentrated under reduced pressure, then 6 N hydrochloric acid was added to the obtained residue for adjustment to pH = 1.0. The obtained yellow solid was filtered, washed with water and methylene chloride, and dried to obtain the title compound (36.5 g, 69%) as a yellow solid.

20 MS (ESI) m/z 406 ($\text{M}+\text{H}$)⁺

(Step

6)

[4-[(2,5-Difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]phenyl]boronic acid



Methyl 4-[(4-bromophenyl)sulfonylamino]-2,5-difluoro-benzoate <see (step 5)> (40.5 g, 100 mmol) and bis(pinacolato)diborane (35.5 g, 140 mmol)

were dissolved in N,N-dimethylformamide (800 ml). [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II) (2.20 g, 3.00 mmol) and potassium acetate (29.4 g, 300 mmol) were added to the solution, and the resulting mixture was stirred in the presence of nitrogen gas at 90°C
 5 for 12 hours. The reaction liquid was cooled to room temperature, and then diluted with water and extracted with ethyl acetate (600 ml $\times$ 3). The extracts were combined, washed with a saturated saline solution, dried over anhydrous sodium sulfate, and then the solvent was removed under reduced pressure. The obtained residue was purified by silica gel column
 10 chromatography (petroleum ether:ethyl acetate = 4:1 to 2:1) to obtain methyl 2,5-difluoro-4-[[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]sulfonylamino]benzoate (40.0 g, 88%).

^1H NMR (CD_3OD , 400 MHz): δ 7.89 (s, 4H), 7.62 - 7.58 (m, 1H), 7.44 - 7.40 (m, 1H), 3.87 (s, 3H), 1.36 (s, 12H); MS (ESI) m/z 454 ($\text{M}+\text{H}$)⁺
 15 [0101]


The obtained methyl 2,5-difluoro-4-[[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]sulfonylamino]benzoate (59.0 g, 130 mmol) was dissolved in acetone (450 ml), sodium periodate (83.6 g, 390 mmol), ammonium acetate (30.0 g, 390 mmol),
 20 and water (150 ml) were added thereto, and the resulting mixture was stirred at room temperature for 72 hours. The reaction liquid was concentrated under reduced pressure, and then diluted with water and extracted with ethyl acetate (500 ml $\times$ 3). The extracts were combined, washed with a saturated saline solution, dried over anhydrous sodium
 25 sulfate, and then the solvent was removed under reduced pressure. The obtained residue was washed with a mixed solvent (petroleum ether:ethyl acetate = 1:10) and dried under reduced pressure to obtain the title compound (43.0 g, 89%).

^1H NMR: (400 MHz, $\text{DMSO}-d_6$): δ 11.05 (s, 1H), 8.37 (s, 2H), 7.94 (d, J = 8.4
 30 Hz, 2H), 7.81 (d, J = 8.0 Hz, 2H), 7.65 - 7.61 (m, 1H), 7.27 - 7.23 (m, 1H), 3.80

(s, 3H); MS (ESI) m/z 372 (M+H)⁺

[0102]

(Step 7) tert-Butyl
4-[5-[4-[(2,5-difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]phenyl]pyrimidin
5 -2-yl]-3,6-dihydro-2H-pyridine-1-carboxylate




tert-Butyl

4-(5-bromopyrimidin-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate <see (step
1)> (100 mg, 0.290 mmol),
10 [4-[(2,5-difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]phenyl]boronic acid
<see (step 6)> (131 mg, 0.350 mmol), sodium carbonate (93.8 mg, 0.885
mmol), and Pd(dppf)Cl₂ (11.0 mg, 0.030 mmol) were dissolved in dioxane
(10.0 ml) and water (0.50 ml), and the resulting mixture was stirred in the
15 presence of nitrogen gas at 100°C for 8 hours. The reaction liquid was
concentrated under reduced pressure, and then the obtained residue was
purified by silica gel column chromatography (petroleum ether:ethyl acetate
= 3:1 to 1:1) to obtain the title compound (50.0 mg, 29%).

MS (ESI) m/z 587 (M+H)⁺

[0103]

20 (Step 8) tert-Butyl
4-[5-[4-[(2,5-difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]phenyl]pyrimidin
-2-yl]piperidine-1-carboxylate



tert-Butyl


4-[5-[4-[(2,5-difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]phenyl]pyrimidin-2-yl]-3,6-dihydro-2H-pyridine-1-carboxylate <see (step 7)> (120 mg, 0.20 mmol) and 10% Pd/C (wet, 50 mg) were suspended in methanol (10 ml), and the resulting mixture was stirred in the presence of hydrogen gas (50 psi) at 50°C for 24 hours. The reaction liquid was cooled to room temperature and then filtered, and the filtrate was concentrated under reduced pressure to obtain the title compound (31.0 mg, 27%).

MS (ESI) m/z 589 (M+H)⁺
[0104]

(Step

9)

4-[[4-[2-(1-tert-Butoxycarbonyl-4-piperidyl)pyrimidin-5-yl]phenyl]sulfonylaminol-2,5-difluoro-benzoic acid (M-10)



15

tert-Butyl

4-[5-[4-[(2,5-difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]phenyl]pyrimidin-2-yl]piperidine-1-carboxylate <see (step 8)> (60 mg, 0.10 mmol) was dissolved in methanol (10 ml), a 2 N aqueous solution of lithium hydroxide (3.0 ml) was added thereto, and the resulting mixture was stirred at room


temperature for 2 hours. The completion of the reaction was verified with TLC, and 4 N hydrochloric acid was added for adjustment to pH = 5.0. The precipitated solid was filtered, washed with a mixed solvent of methylene chloride:methanol (10:1), and dried to obtain the title compound (45 mg, 78%) as a white solid.

^1H NMR (CD_3OD , 400 MHz): δ 8.93 (s, 2H), 7.94 - 7.91 (m, 2H), 7.80 (d, J = 8.4 Hz, 2H), 7.50 - 7.45 (m, 1H), 7.36 - 7.32 (m, 1H), 4.10 - 4.07 (m, 2H), 3.05 - 2.99 (m, 1H), 2.90 - 2.81 (m, 2H), 1.91 - 1.89 (m, 2H), 1.76 - 1.69 (m, 2H), 1.43 (s, 9H); MS (ESI) m/z 573 ($M-1$)

[0105]

Example 11: Synthesis of M-11


(Step 1) 5-Bromo-2-(3,6-dihydro-2H-pyran-4-yl)pyrimidine



5-Bromo-2-iodopyrimidine (5.87 g, 20.7 mmol), 2-(3,6-dihydro-2H-pyran-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborane (4.35 g, 20.7 mmol), potassium carbonate (5.70 g, 41.4 mmol), and $\text{Pd}(\text{dppf})\text{Cl}_2$ (805 mg, 1.10 mmol) were suspended in N,N -dimethylformamide (50 ml) and water (10 ml), and the resulting mixture was stirred under a nitrogen gas atmosphere at 100°C for 4 hours. The reaction liquid was concentrated under reduced pressure, and the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 5:1 to 1:1) to obtain the title compound (1.44 g, 29%).

^1H NMR (400 MHz, CDCl_3): δ 8.72 (s, 2H), 7.25 (s, 1H), 4.42 - 3.38 (m, 2H), 3.94 - 3.91 (m, 2H), 2.68 - 2.66 (m, 2H)

(Step 2) Methyl 4-[[4-[2-(3,6-dihydro-2H-pyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluorobenzoate




5-Bromo-2-(3,6-dihydro-2H-pyran-4-yl)pyrimidine <see (step 1)>
 (1.44 g, 6.00 mmol),
 [4-[(2,5-difluoro-4-methoxycarbonylphenyl)sulfonyl]phenyl]boronic acid
 (see Example 10 (step 6)> (2.20 g, 6.00 mmol), sodium carbonate (1.27 g, 12.0
 mmol), and Pd(dppf)Cl₂ (220 mg, 0.3 mmol) were suspended in dioxane (20
 ml) and water (5.0 ml), and the resulting mixture was stirred at 100°C for 8
 hours. The reaction liquid was concentrated under reduced pressure, and
 the obtained residue was purified by silica gel column chromatography
 (petroleum ether:ethyl acetate = 3:1 to 1:1) to obtain the title compound (1.20
 g, 41%).

MS (ESI) m/z 488 (M+H)⁺

[0106]

(Step 3) Methyl
 2,5-difluoro-4-[[4-(2-tetrahydropyran-4-ylpyrimidin-5-yl)phenyl]sulfonylami
 nol]benzoate



Methyl

4-[[4-[2-(3,6-dihydro-2H-pyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,
 5-difluorobenzoate <see (step 2)> (1.20 g, 2.46 mmol) was dissolved in
 methanol (10 ml), Pd/C (10%, 500 mg) was added thereto, and the resulting
 mixture was stirred under hydrogen atmosphere (50 psi) at 50°C for 24 hours.


The reaction liquid was cooled to room temperature and then filtered, and the solvent was removed under reduced pressure to obtain the title compound (843 mg, 70%).

MS (ESI) m/z 490 ($M+H$)⁺

5 [0107]

(Step 4)

2,5-Difluoro-4-[[4-(2-tetrahydropyran-4-ylpyrimidin-5-yl)phenyl]sulfonylaminobenzoic acid (M-11)



10 Methyl


2,5-difluoro-4-[[4-(2-tetrahydropyran-4-ylpyrimidin-5-yl)phenyl]sulfonylaminobenzoate <see (step 3)> (600 mg, 1.22 mmol) was dissolved in methanol (10 ml), a 2 N aqueous solution of lithium hydroxide (3.0 ml) was added thereto, and the resulting mixture was stirred at room temperature for 2 hours. The completion of the reaction was verified with TLC, 4 N hydrochloric acid was added for adjustment of the pH to 4 to 5, and the precipitated white solid was filtered off, dried, and washed with dichloromethane:methanol = 10:1 to obtain the title compound (512 mg, 88%) as a white solid.

20 ¹H NMR (CD₃OD, 400 MHz): δ 9.07 (s, 2H), 8.06 (d, J = 8.4 Hz, 2H), 7.93 (d, J = 8.4 Hz, 2H), 7.63 - 7.59 (m, 1H), 7.50 - 7.45 (m, 1H), 4.10 - 4.07 (m, 2H), 3.66 - 3.59 (m, 2H), 3.25 - 3.19 (m, 1H), 2.04 - 1.94 (m, 4H); MS (ESI) m/z 476 ($M+H$)⁺

[0108]

25 Example 12: Synthesis of M-12

(Step 1) Chloromethyl benzoate




Paraformaldehyde (4.5 g) and zinc chloride (catalyst amount) were mixed at 0°C. Benzoyl chloride (20 g, 0.14 mol) was added dropwise thereto over 1 hour. The temperature of the reaction liquid was raised to room temperature, and then the reaction liquid was stirred at 55°C for 10 hours. The reaction liquid was cooled and then purified by silica gel column chromatography (petroleum ether:ethyl acetate = 30:1 to 10:1) to obtain the title compound (9.7 g, 40%).

¹H NMR (CDCl₃, 400 MHz): δ 8.09 - 8.07 (m, 2H), 7.63 - 7.60 (m, 1H), 7.49 - 7.44 (m, 2H), 5.96 (s, 2H)

[0109]

(Step 2) Iodomethyl benzoate




Chloromethyl benzoate <see (step 1)> (10.0 g, 58.8 mmol) was dissolved in acetonitrile (70.0 ml), sodium iodide (17.6 g, 117 mmol) was added thereto, and the resulting mixture was stirred at room temperature for 24 hours. Acetonitrile was distilled off under reduced pressure, and diethyl ether was added. The precipitated solid was filtered off, washed with diethyl ether, and dried under reduced pressure, and then purified by silica gel column chromatography (petroleum ether:ethyl acetate = 30:1 to 10:1) to obtain the title compound (14.5 g, 94%).

¹H NMR (CDCl₃, 400 MHz): δ 8.06 - 8.04 (m, 2H), 7.64 - 7.62 (m, 1H), 7.49 - 7.44 (m, 2H), 6.17 (s, 2H)

[0110]

(Step 3) (5-Bromopyrimidin-2-yl)methyl benzoate




Zinc (21.6 g, 332 mmol) was heated at 210°C for 10 minutes, then cooled to 70°C, heated to 210°C again, stirred for 10 minutes, and cooled to room temperature. N,N-dimethylformamide (100 ml) and a
 5 dibromomethane (7.72 g, 41.2 mmol)/N,N-dimethylformamide (20 ml) solution was added thereto, and the resulting mixture was stirred at 90°C for 30 minutes. The reaction solution was cooled to room temperature, chlorotrimethylsilane (900 mg, 8.30 mmol) was added, and the resulting
 10 mixture was stirred at room temperature for 10 minutes. Iodomethyl benzoate <see (step 2)> (14.5 g, 55.3 mmol)/N,N-dimethylformamide solution (60 ml) was added dropwise at 35°C, and the resulting mixture was stirred for 90 minutes. 5-Bromo-2-iodopyrimidine (7.90 g, 28.0 mmol) and Pd(PPh₃)₂Cl₂ (3.0 g, 4.1 mmol) were suspended in N,N-dimethylformamide
 15 (80 ml), the zinc reagent obtained in the above step was added with a syringe, and the resulting mixture was stirred under a nitrogen atmosphere at 80°C for 2 hours. The reaction liquid was cooled to room temperature and filtered, and then the filtrate was diluted with water (600 ml) and extracted with ethyl acetate (200 ml × 3). The extracts were combined, washed with a
 20 saturated saline solution, and dried over sodium sulfate. The solvent was removed under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 6:1 to 2:1) to obtain the title compound (7.38 g, 90%).

¹H NMR (CDCl₃, 400 MHz): δ 8.79 (s, 2H), 8.15 - 8.13 (m, 2H), 7.63 - 7.57 (m, 1H), 7.47 - 7.45 (m, 2H), 5.63 (s, 2H)

[0111]

(Step 4) (5-Bromopyrimidin-2-yl)methanol




(5-Bromopyrimidin-2-yl)methyl benzoate <see (step 3)> (7.30 g, 25.0 mmol) was dissolved in methanol (15 ml), a 1 N sodium methoxide/methanol solution (50.0 ml, 0.500 mmol) was added thereto, and the resulting mixture was stirred at room temperature. The solvent was distilled off after deprotection was completed, and the obtained residue was purified by silica gel column chromatography (dichloromethane:methanol = 30:1 to 15:1) to obtain the title compound (3.58 g, 76%).

¹H NMR (CDCl₃, 400 MHz): δ 8.80 (s, 2H), 4.82 (d, J = 4.0 Hz, 2H), 3.39 (s, 1H)

[0112]


(Step 5) 5-Bromo-2-(methoxymethyl)pyrimidine



(5-Bromopyrimidin-2-yl)methanol <see (step 4)> (3.72 g, 19.8 mmol) was dissolved in tetrahydrofuran (10 ml), the resulting solution was cooled to 0°C, then sodium hydride (60% in oil, 1.19 g, 29.7 mmol) and methyl iodide (4.20 g, 29.7 mmol) were added thereto, and the resulting mixture was stirred at room temperature for 1 hour. The reaction liquid was concentrated under reduced pressure, and the obtained residue was purified by silica gel column chromatography to obtain the title compound (2.40 g, 60%).

[0113]

(Step 6) Methyl 2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoate



5-Bromo-2-(methoxymethyl)pyrimidine <see (step 5)> (2.30 g, 11.3 mmol), [4-[(2,5-difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]phenyl]boronic acid <see Example 10 (step 6)> (4.20 g, 11.3 mmol), sodium carbonate (2.40 g, 22.6 mmol), and Pd(dppf)Cl₂ (415 mg, 0.57 mmol) were suspended in dioxane (50 ml) and water (10 ml), and the resulting mixture was stirred at 100°C for 8 hours. The reaction liquid was concentrated under reduced pressure, and the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 2:1 to 1:4) to obtain the title compound (2.95 g, 58%).


MS (ESI) m/z 450 (M+H)⁺

[0114]

(Step

7)

2,5-Difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoic acid (M-12)



Methyl

2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoate <see (step 5)> (700 mg, 1.56 mmol) was dissolved in methanol (10 ml), a 2 N aqueous solution of lithium hydroxide (3.0 ml) was added thereto, and the resulting mixture was stirred at room temperature for 2 hours. The completion of the reaction was verified with TLC, and then 4 N hydrochloric acid was added for adjustment of the pH to 4 to 5. The precipitated white solid was filtered, dried, and washed with a mixed solvent (methylene


chloride:methanol = 10:1) to obtain the title compound (617 mg, 91%) as a white solid.

¹H NMR (CD₃OD, 400 MHz): δ 9.06 (s, 2H), 8.01 (d, J = 8.4 Hz, 2H), 7.90 (d, J = 8.4 Hz, 2H), 7.56 - 7.52 (m, 1H), 7.43 - 7.39 (m, 1H), 4.68 (s, 2H), 3.47 (s,

[0115]

Example 13: Synthesis of M-13

(Step 1) 4-Methoxy-2-butyric acid




3-Methoxypropyne (1.00 g, 14.3 mmol) was dissolved in tetrahydrofuran (100 ml), the resulting solution was cooled to -78°C, and then a 2.5 M n-butyllithium/hexane solution (24.4 ml, 61.0 mmol) was added dropwise. The resulting mixture was stirred at -78°C, then dry ice was added to the reaction liquid, and the temperature of the reaction liquid was gradually raised to room temperature. The reaction liquid was diluted with water, and then 4 N hydrochloric acid was added for adjustment of the pH to 5.0. The resulting liquid was extracted with methylene chloride (50 ml × 3), the organic layers were combined and dried over sodium sulfate, and the solvent was removed under reduced pressure to obtain the title compound (1.50 g, 92%) as a yellow oil.

¹H NMR (CDCl₃, 400 MHz): δ 8.97 (s, 1H), 4.27 (s, 2H), 3.44 (s, 3H)

[0116]

(Step 2) Methyl 2,5-difluoro-4-[(4-iodophenyl)sulfonylamino]benzoate




A method similar to Example 10 (step 5) was performed to methyl

4-amino-2,5-difluoro-benzoate <see Example 10 (step 4)> (6.50 g, 34.8 mmol) and 4-iodobenzenesulfonyl chloride (21.0 g, 69.5 mmol) to obtain the title compound (14.0 g, 89%) as a yellow solid.

MS (ESI) m/z 452 (M+1)

5 [0117]

(Step 3) Methyl 2,5-difluoro-4-[[4-[4-(methoxymethyl)triazol-1-yl]phenyl]sulfonylamino]benzoate



10 4-Methoxy-2-butynoic acid <see (step 1)> (502 mg, 4.40 mmol) and methyl 2,5-difluoro-4-[(4-iodophenyl)sulfonylamino]benzoate <see (step 2)> (2.00 g, 4.40 mmol) were dissolved in dimethylsulfoxide (25.0 ml) and water (3.0 ml), L-proline (102 mg, 0.880 mmol), copper(II) sulfate pentahydrate (110 mg, 0.440 mmol), sodium ascorbate (175 mg, 0.880 mmol), sodium azide
15 (345 mg, 5.30 mmol), and potassium carbonate (731 mg, 5.30 mmol) were sequentially added, and the resulting mixture was stirred at 65°C for 12 hours. Ethyl acetate (50 ml), ammonium water (50 ml), and water (100 ml) were added to the reaction liquid, then the resulting mixture was extracted with ethyl acetate (60 ml $\times$ 10), the organic layers were combined and
20 washed with a saturated saline solution (300 ml), dried over sodium sulfate, and the solvent was removed under reduced pressure to obtain the title compound (510 mg, 26%).


^1H NMR (CD_3OD , 400 MHz): δ 8.48 (s, 1H), 7.98 (d, J = 8.4 Hz, 2H), 7.86 (d, J = 8.0 Hz, 2H), 7.30 - 7.26 (m, 1H), 6.94 - 6.88 (m, 1H), 4.50 (s, 2H), 3.68 (s,
25 3H), 3.31 (s, 3H)

[0118]

(Step

4)

2,5-Difluoro-4-[[4-[4-(methoxymethyl)triazol-1-yl]phenyl]sulfonylamino]benzoic acid (M-13)



5

Methyl


2,5-difluoro-4-[[4-[4-(methoxymethyl)triazol-1-yl]phenyl]sulfonylamino]benzoate <see (step 3)> (500 mg, 1.14 mmol) was dissolved in methanol (10 ml), a 2 N aqueous solution of lithium hydroxide (5.0 ml) was added thereto, and the resulting mixture was stirred at room temperature for 2 hours. The completion of the reaction was verified with TLC, and then 4 N hydrochloric acid was added for adjustment of the pH to 5.0. The precipitated solid was filtered and dried to obtain the title compound (450 mg, 93%) as a yellow solid.

¹H NMR (CD₃OD, 400 MHz): δ 8.66 (s, 1H), 8.14 - 8.08 (m, 4H), 7.56 - 7.47 (m, 2H), 4.64 (s, 2H), 3.44 (s, 3H); MS (ESI) m/z 425 (M+H)⁺ [0119]

15

Example 14: Synthesis of M-14

(Step 1) 2-Allyloxy-5-bromo-pyrimidine



20


5-Bromo-2-chloro-pyrimidine (30.0 g, 0.150 mol) was dissolved in tetrahydrofuran (300 ml), potassium-tert butoxide (20.1 g, 0.180 mol) and allyl alcohol (19.9 g, 0.170 mol) were added thereto, and the resulting mixture was stirred at 0°C for 2 hours. The reaction liquid was diluted with water and extracted with diethyl ether (300 ml × 3). The extracts were

combined and washed with a saturated saline solution, dried over sodium sulfate, and the solvent was removed under reduced pressure to obtain the title compound (23.3 g, 70%).

^1H NMR (CDCl_3 , 300 MHz): δ 8.53 (s, 2H), 6.11 - 6.02 (m, 1H), 5.45 - 5.39 (m, 1H), 5.30 - 5.26 (m, 1H), 4.89 - 4.87 (m, 2H)

[0120]

(Step 2) Methyl 4-[[4-(1-allyl-2-oxo-pyrimidin-5-yl)phenyl]sulfonylamino]-2,5-difluoro-benzoate




2-Allyloxy-5-bromo-pyrimidine <see (step 1)> (7.12 g, 3.30w mmol), 4-[(2,5-difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]phenyl]boronic acid <see Example 10 (step 6)> (8.19 g, 22.1 mmol), sodium carbonate (7.03 g, 66.1 mmol), and $\text{Pd}(\text{dppf})\text{Cl}_2$ (2.40 g, 3.30 mmol) were dissolved in N,N-dimethylformamide (200 ml) and water (20 ml), and the resulting mixture was stirred in the presence of nitrogen gas at 100°C for 2 hours. The reaction liquid was concentrated under reduced pressure, and then the obtained residue was purified by HPLC (acetonitrile in water) to obtain the title compound (3.56 g, 35%).

MS (ESI) m/z 462 ($\text{M}+\text{H}^+$)

[0121]

(Step 3) 4-[[4-(1-Allyl-2-oxo-pyrimidin-5-yl)phenyl]sulfonylamino]-2,5-difluoro-benzoic acid



Methyl


4-[[4-(1-allyl-2-oxo-pyrimidin-5-yl)phenyl]sulfonylamino]-2,5-difluorobenzoate <see (step 2)> (3.51 g, 7.60 mmol) was dissolved in methanol (30 ml), a 2 N aqueous solution of lithium hydroxide (10.0 ml) was added thereto, and the resulting mixture was stirred at room temperature for 2 hours. The completion of the reaction was verified with TLC, and then 4 N hydrochloric acid was added for adjustment of the pH to 5.0. The precipitated solid was filtered and dried to obtain the title compound (2.72 g, 80%) as a yellow solid.

¹H NMR (DMSO-d₆, 400 MHz): δ 13.42 - 13.40 (br, 1H), 11.13 - 11.05 (br, 1H), 9.04 (d, J = 3.3 Hz, 1H), 8.72 (d, J = 3.3 Hz, 1H), 7.93 - 7.85 (m, 4H), 7.67 - 7.57 (m, 1H), 6.07 - 5.95 (m, 1H), 5.25 - 5.18 (m, 2H), 4.58 - 4.51 (m, 2H); MS (ESI) m/z 448 (M+H)⁺

[0122]

Example 15: Synthesis of M-15

(Step 1) 5-Bromo-2-ethoxymethylpyrimidine




(5-Bromopyrimidin-2-yl)methanol <see Example 12 (step 4)> (3.72 g, 19.8 mmol) was dissolved in tetrahydrofuran (10.0 ml), the reaction solution was cooled to 0°C, then sodium hydride (60% wt in mineral oil, 1.19 g, 29.7 mmol) and ethane iodide (4.63 mmol, 29.7 mmol) were sequentially added, and the resulting mixture was stirred at room temperature for 1 hour. The reaction liquid was concentrated under reduced pressure, and the obtained

residue was purified by silica gel column chromatography (petroleum ether) to obtain the title compound (2.58 g, 60%).

[0123]

(Step 2) Methyl

5 4-[[4-[2-(ethoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-benzoate




5-Bromo-2-(ethoxymethyl)pyrimidine <see (step 1)> (2.45 g, 11.3 mmol), 4-[(2,5-difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]phenyl]boronic acid <see Example 10 (step 6)> (4.20 g, 11.3 mmol), sodium carbonate (2.40 g, 22.6 mmol), and Pd(dppf)Cl₂ (415 mg, 0.57 mmol) were dissolved in dioxane (50.0 ml) and water (10.0 ml), and the resulting mixture was stirred in the presence of nitrogen at 100°C for 8 hours. The reaction liquid was concentrated under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 2:1 to 1:4) to obtain the title compound (2.94 g, 56%).

MS (ESI) m/z 464 (M+H)⁺

[0124]

(Step 3)

20 4-[[4-[2-(Ethoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-benzoic acid (M-15)



Methyl


4-[[4-[2-(ethoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-be

nzoate <see (step 2)> (800 mg, 1.72 mmol) was dissolved in methanol (10 ml), a 2 N aqueous solution of lithium hydroxide (3.0 ml) was added thereto, and the resulting mixture was stirred at room temperature for 2 hours. The completion of the reaction was verified with TLC, and then 4 N hydrochloric acid was added for adjustment of the pH to 5.0. The precipitated solid was filtered, washed with a mixed solution of methylene chloride:methanol (10:1), and dried to obtain the title compound (698 mg, 90%) as a white solid.

¹H NMR (DMSO-d₆, 400 MHz): δ 13.35 (br, 1H), 11.19 (br, 1H), 9.25 (d, J = 7.6 Hz, 2H), 8.16 - 7.96 (m, 4H), 7.63 - 7.56 (m, 1H), 7.31 - 7.27 (m, 1H), 4.67 (s, 2H), 3.63 - 3.48 (m, 2H), 1.23 - 1.15 (m, 3H); MS (ESI) m/z 450 (M+H)⁺ [0125]

Example 16: Synthesis of M-16

(Step 1) 4-Nitro-2-tetrahydropyran-4-yl-pyridine




Zinc (19.2 g, 293 mmol) was heated at 210°C for 10 minutes, cooled to 70°C, then heated to 210°C again, and the resulting mixture was stirred for 10 minutes. The reaction mixture was cooled to room temperature, then a solution of N,N-dimethylformamide (100 ml) and dibromoethane (6.87 g, 33.6 mmol) in N,N-dimethylformamide (10.0 ml) was added thereto, and the resulting mixture was stirred at 90°C for 30 minutes. The reaction mixture was cooled to room temperature, then trimethylsilyl chloride (800 mg, 7.30 mmol) was added thereto, and the resulting mixture was stirred at room temperature for 10 minutes. A solution of 4-iodine tetrahydropyran (10.4 g, 49.2 mmol) in N,N-dimethylformamide (60.0 ml) was added to the reaction liquid, and the resulting mixture was stirred at 35°C for 90 minutes. This zinc derivative was added to a suspension of 2-bromo-4-nitro-pyridine (5.00 g, 24.6 mmol) and Pd(PPh₃)₂Cl₂ (2.60 g, 3.70 mmol) in N,N-dimethylformamide

(80.0 ml), and the resulting mixture was stirred in the presence of nitrogen gas at 90°C for 2 hours. The reaction mixture was cooled to room temperature, then the reaction solution was filtered, and the obtained filtrate was diluted with water (600 ml) and extracted with ethyl acetate (200 ml × 3). The extracts were combined, washed with a saturated saline solution, dried over anhydrous sodium sulfate, and then the solvent was removed under reduced pressure. The obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 10:1 to 5:1) to obtain the title compound (900 mg, 17%).

¹H NMR (CDCl₃, 400 MHz): δ 8.85 (d, J = 5.6 Hz, 1H), 7.90-7.86 (m, 2H), 4.15-4.11 (m, 2H), 3.61-3.54 (m, 2H), 3.16-3.10 (m, 1H), 2.00-1.91 (m, 4H) [0126]


(Step 2) 2-Tetrahydropyran-4-ylpyridin-4-amine



4-Nitro-2-tetrahydropyran-4-yl-pyridine <see (step 1)> (900 mg, 4.30 mmol) and 10% Pd/C (wet, 100 mg) were suspended in methanol (5.0 ml), and the resulting mixture was stirred in the presence of hydrogen at room temperature for 24 hours. The reaction liquid was filtered, and the filtrate was concentrated under reduced pressure to obtain the title compound (700 mg, 91%).

¹H NMR (CD₃OD, 300 MHz): δ 7.89 (d, J = 5.7 Hz, 1H), 6.49 - 6.41 (m, 2H), 4.07 - 4.01 (m, 2H), 3.59 - 3.50 (m, 2H), 2.80 - 2.71 (m, 1H), 1.84 - 1.76 (m, 4H) [0127]


(Step 3) 4-Bromo-2-tetrahydropyran-4-yl-pyridine



2-Tetrahydropyran-4-ylpyridin-4-amine <see (step 2)> (700 mg, 3.90 mmol) was dissolved in bromoform (30 ml), amyl nitrite (20 ml) was added thereto, and the resulting mixture was stirred in the presence of nitrogen gas at 85°C for 4 hours. The reaction mixture was cooled to room temperature, then the reaction liquid was concentrated under reduced pressure, and the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 10:1 to 5:1) to obtain the title compound (350 mg, 37%).

¹H NMR (CDCl₃, 300 MHz): δ 8.37 (d, 1H, J = 5.4 Hz), 7.36 - 7.27 (m, 2H), 4.13 - 4.07 (m, 2H), 3.59 - 3.50 (m, 2H), 2.97 - 2.92 (m, 1H), 1.92 - 1.84 (m, 4H) [0128]

(Step 4) 2-Benzylsulfanyl-5-bromo-pyridine




5-Bromo-2-chloro-pyridine (80.0 g, 645 mmol) was dissolved in tetrahydrofuran (600 ml), sodium hydride (60% wt in mineral oil, 45.0 g, 1.13 mol) was added thereto over 20 minutes, and the resulting mixture was stirred at room temperature for 30 minutes. Benzyl thiol (123 g, 640 mmol) was added to the reaction liquid, and the resulting mixture was stirred at room temperature for 3 hours. The reaction liquid was diluted with water, extracted with diethyl ether, the extracts were combined, and washed with a saturated aqueous solution of sodium hydrogen carbonate, and a saturated saline solution. The obtained residue was dried over magnesium sulfate, and the solvent was removed under reduced pressure to obtain the title

compound (148 g, 82%).

[0129]


(Step 5) 5-Bromopyridine-2-sulfonyl chloride



2-Benzylsulfanyl-5-bromo-pyridine <see (step 4)> (250 g, 0.89 mol) was suspended in acetic acid (2,250 ml) and water (750 ml), NCS (340 g, 2.60 mol) was added thereto, and the resulting mixture was stirred at room temperature for 2 hours. The reaction liquid was diluted with water and extracted with methylene chloride. The extracts were combined, washed with a saturated aqueous solution of sodium hydrogen carbonate and a saturated saline solution, and dried over magnesium sulfate. The obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 30:1) to obtain the title compound (60.4 g, 26%).

[0130]

15	(Step	6)	Methyl
	4-[(5-bromo-2-pyridyl)sulfonylamino]-2,5-difluoro-benzoate		



Methyl 4-amino-2,5-difluoro-benzoate <see Example 10 (step 4)> (14.1 g, 75.0 mmol) and 5-bromopyridine-2-sulfonyl chloride <see (step 5)> (29.0 g, 113 mmol) were dissolved in methylene chloride (250 ml), pyridine (23 ml) was added thereto, and the resulting mixture was stirred at room temperature for 12 hours. The reaction liquid was concentrated under reduced pressure, 6 N hydrochloric acid was added for adjustment of the pH to 1.0, and the precipitated yellow solid was filtered. The obtained solid was

washed with water and dried to obtain the title compound (21.4 g, 70%) as a yellow solid.


¹H NMR (400 MHz, DMSO-d₆): δ 11.30 (s, 1H), 8.91 (d, J = 2.0 Hz, 1H), 8.40 (dd, J = 8.4, 2.4 Hz, 1H), 8.00 (d, J = 8.4 Hz, 1H), 7.69 - 7.65 (m, 1H), 7.45 - 7.39 (m, 1H), 3.83 (s, 3H); MS (ESI) m/z 407, 409 (M+H)⁺

[0131]

(Step

7)

[6-[(2,5-Difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]-3-pyridyl]boronic acid



10

Two steps similar to Example 10 (step 6) were performed to methyl 4-[(5-bromo-2-pyridyl)sulfonylamino]-2,5-difluoro-benzoate <see (step 6)> (41 g, 0.10 mol) to obtain the title compound (30 g, 81%) as a black oil.

MS (ESI) m/z 371 (M-1)


[0132]

(Step

8)

Methyl

2,5-difluoro-4-[[5-(2-tetrahydropyran-4-yl-4-pyridyl)-2-pyridyl]sulfonylamino]benzoate



20


4-Bromo-2-tetrahydropyran-4-yl-pyridine <see (step 3)> (350 mg, 1.45 mmol), [6-[(2,5-difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]-3-pyridyl]boronic acid

<see (step 7)> (647 mg, 1.74 mmol), sodium carbonate (462 mg, 4.83 mmol), and Pd(dppf)Cl₂ (106 mg, 0.14 mmol) were dissolved in dioxane (10.0 ml) and water (2.0 ml), and the resulting solution was stirred in the presence of nitrogen at 105°C for 2 hours. The reaction liquid was concentrated under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 10:1 to 1:1) to obtain the title compound (210 mg, 29%).

[0133]

(Step 9)

2,5-Difluoro-4-[[5-(2-tetrahydropyran-4-yl-4-pyridyl)-2-pyridyl]sulfonylamino]benzoic acid (M-16)



Methyl

2,5-difluoro-4-[[5-(2-tetrahydropyran-4-yl-4-pyridyl)-2-pyridyl]sulfonylamino]benzoate <see (step 8)> (210 mg, 0.43 mmol) was dissolved in methanol (5.0 ml), a 2 N aqueous solution of lithium hydroxide (3.0 ml) was added thereto, and the resulting mixture was stirred at room temperature for 2 hours. The completion of the reaction was verified with TLC, and then 4 N hydrochloric acid was added for adjustment of the pH to 5.0. The precipitated solid was filtered and dried to obtain the title compound (140 mg, 69%) as a white solid.


¹H NMR (DMSO-d₆, 300 MHz): δ 13.32 (br s, 1H), 11.27 (br s, 1H), 9.18 (s, 1H), 8.65 (d, J = 3.6 Hz, 1H), 8.53 (dd, J = 6.0, 1.2 Hz, 1H), 8.15 (d, J = 6.0 Hz, 1H), 7.77 (s, 1H), 7.69 (d, J = 3.6 Hz, 1H), 7.61 (dd, J = 8.1, 5.1 Hz, 1H), 7.45 (dd, J = 9.3, 5.1 Hz, 1H), 3.98 - 3.96 (m, 2H), 3.49 - 3.43 (m, 2H), 3.04 - 2.99

(m, 1H), 1.87 - 1.78 (m, 4H); MS (ESI) m/z 476 (M+H)⁺

[0134]

Example 17: Synthesis of M-17

(Step 1) (5-Bromopyrimidin-2-yl)methyl methanesulfonate



5

(5-Bromopyrimidin-2-yl)methanol <see Example 12 (step 4)> (3.72 g, 19.7 mmol) was dissolved in tetrahydrofuran (10 ml), triethylamine (3.0 g, 30 mmol), mesyl chloride (3.38 g, 29.6 mmol) was added at 0°C, and the resulting mixture was stirred at room temperature for 1 hour. The reaction liquid was concentrated under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 10:1) to obtain the title compound (4.54 g, 86%).


10

¹H NMR (CDCl₃, 400 MHz): δ 8.83 (s, 2H), 5.39 (s, 2H), 3.20 (s, 3H)

[0135]

15

(Step 2) N-[(5-Bromopyrimidin-2-yl)methyl]propan-2-amine



20


(5-Bromopyrimidin-2-yl)methyl methanesulfonate <see (step 1)> (200 mg, 0.750 mmol) was dissolved in ethanol (10 ml), isopropylamine (88.5 mg, 1.50 mmol) and triethylamine (151 mg, 1.50 mmol) were added thereto, and the resulting mixture was stirred at 80°C for 1 hour. The reaction liquid was concentrated under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 3:1 to 1:1) to obtain the title compound (144 mg, 84%).

25

¹H NMR (CDCl₃, 400 MHz): δ 8.77 (s, 2H), 4.08 (s, 2H), 2.89 - 2.83 (m, 1H), 1.12 (d, J = 6.4 Hz, 6H)

[0136]


(Step 3) Methyl 2,5-difluoro-4-[[4-[2-[(isopropylamino)methyl]pyrimidin-5-yl]phenyl]sulfonyl amino]benzoate



5 N-[(5-Bromopyrimidin-2-yl)methyl]propan-2-amine <see (step 2)> (0.260 g, 1.13 mmol), [4-[(2,5-difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]phenyl]boronic acid <see Example 10 (step 6)> (0.420 g, 1.13 mmol), sodium carbonate (0.24 g, 2.2 mmol), and Pd(dppf)Cl₂ (41.5 mg, 0.057 mmol) were dissolved in dioxane 10 (5.0 ml) and water (1.0 ml), and the resulting mixture was stirred in the presence of nitrogen gas at 100°C for 8 hours. The reaction liquid was concentrated under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 2:1 to 1:4) to obtain the title compound (0.32 g, 59%).

15 MS (ESI) m/z 477 (M+H)⁺ [0137]

(Step 4) 2,5-Difluoro-4-[[4-[2-[(isopropylamino)methyl]pyrimidin-5-yl]phenyl]sulfonyl amino]benzoic acid (M-17)



20


Methyl 2,5-difluoro-4-[[4-[2-[(isopropylamino)methyl]pyrimidin-5-yl]phenyl]sulfonyl amino]benzoate <see (step 3)> (476 mg, 1.00 mmol) was dissolved in

methanol (10 ml), a 2 N aqueous solution of lithium hydroxide (3.0 ml) was added thereto, and the resulting mixture was stirred at room temperature for 2 hours. The completion of the reaction was verified with TLC, and then 4 N hydrochloric acid was added for adjustment of the pH to 5.0. The precipitated solid was filtered, washed with a mixed solution of methylene chloride:methanol (10:1), and dried to obtain the title compound (406 mg, 88%) as a white solid.

¹H NMR (CD₃OD, 400 MHz): δ 9.06 (s, 2H), 7.88 (d, J = 8.8 Hz, 2H), 7.80 (d, J = 8.4 Hz, 2H), 7.63 - 7.56 (m, 2H), 4.44 (s, 2H), 3.48 - 3.45 (m, 1H), 1.33 (d, J = 6.4 Hz, 6H); MS (ESI) m/z 463 (M+H)⁺ [0138]

Example 18: Synthesis of M-18

(Step 1) 5-Bromo-2-vinyl-pyrimidine




5-Bromo-2-iodine-pyrimidine (28.4 g, 0.100 mol) and Pd(PPh₃)₄ (1.73 g, 1.50 mmol) were dissolved in tetrahydrofuran (350 ml), a 1.5 M solution of vinyl magnesium bromide/tetrahydrofuran (75.0 ml, 0.110 mol) was added thereto, and the resulting mixture was stirred at room temperature for 16 hours. The reaction liquid was diluted with water (200 ml) and extracted with ethyl acetate (200 ml × 3). The extracts were combined, washed with a saturated saline solution, dried over anhydrous sodium sulfate, and then the solvent was removed under reduced pressure to obtain the title compound.

MS (ESI) m/z 185, 187 (M+H)⁺

[0139]

(Step 2) 2-(5-Bromopyrimidin-2-yl)-N,N-dimethylethanamine




5-Bromo-2-vinyl-pyrimidine <see (step 1)> was dissolved in methanol (50 ml), dimethylamine (a 40% aqueous solution, 19.0 ml, 0.250 mol) was added thereto, and the resulting mixture was stirred at room temperature for 90 minutes. The reaction liquid was concentrated under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (methylene chloride:methanol = 100:1 to 10:1) to obtain the title compound (6.90 g, 30%, two steps).

¹H NMR (CD₃OD, 300 MHz): δ 8.82 (s, 2H), 3.33 - 3.32 (m, 2H), 2.88 - 2.83 (m, 2H), 2.30 (s, 6H)

10 [0140]

(Step 3) Methyl 4-[[4-[2-(2-dimethylaminoethyl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluorobenzoate




15 2-(5-Bromopyrimidin-2-yl)-N,N-dimethylethanamine <see (step 2)> (5.20 g, 22.6 mmol), methyl 2,5-difluoro-4-[[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]sulfonylamino]benzoate <see Example 10 (step 6)> (10.2 g, 22.6 mmol), sodium carbonate (3.60 g, 33.9 mmol), and Pd(dppf)Cl₂ (830 mg, 1.14 mmol) were dissolved in dioxane (60 ml) and water (20 ml), and the resulting mixture was stirred in the presence of nitrogen at 100°C for 8 hours. The reaction liquid was concentrated under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (methylene chloride:methanol = 100:1 to 10:1) to obtain the title compound (5.90 g, 55%).

25 MS (ESI) m/z 477 (M+H)⁺

[0141]

(Step 4)

4-[[4-[2-(2-Dimethylaminoethyl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-benzoic acid (M-18)



5 Methyl


4-[[4-[2-(2-dimethylaminoethyl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-benzoate <see (step 3)> (5.90 g, 12.4 mmol) was dissolved in methanol (10 ml), a 2 N aqueous solution of lithium hydroxide (30 ml) was added thereto, and the resulting mixture was stirred at room temperature for 2
10 hours. The completion of the reaction was verified with TLC, and then 4 N hydrochloric acid was added for adjustment of the pH to 5.0. The precipitated solid was filtered and dried to obtain the title compound (4.6 g, 81%) as a white solid.

¹H NMR (DMSO-d₆, 300 MHz): δ 9.11 (s, 2H), 7.86 (s, 4H), 7.30 - 7.23 (m, 1H),
15 6.99 - 6.92 (m, 1H), 3.55 - 3.51 (m, 2H), 3.38 - 3.34 (m, 2H), 2.81 (s, 6H); MS (ESI) m/z 463 (M+H)⁺

[0142]

Example 19: Synthesis of M-19

(Step 1) 5-Bromo-2-(isopropoxymethyl)pyrimidine



20


Isopropanol (10 ml) was added to sodium hydride (60% wt in mineral oil, 225 mg, 5.60 mmol), and the resulting mixture was stirred at 60°C for 30 minutes. The reaction liquid was cooled to room temperature, then

(5-bromopyrimidin-2-yl)methyl methanesulfonate <see Example 17 (step 1)> (1.0 g, 3.7 mmol) a solution of isopropanol (15 ml) was added thereto, and the resulting mixture was stirred at 0°C for 3 hours. The reaction liquid was diluted with water (50 ml), extracted with diethyl ether (50 ml × 3), the
 5 extracts were combined, washed with a saturated saline solution, dried over anhydrous sodium sulfate, then the solvent was removed under reduced pressure, and the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 10:1) to obtain the title compound (657 mg, 76%).

10 ¹H NMR (CDCl₃, 400 MHz): δ 8.80 (s, 2H), 4.71 (s, 2H), 3.82 - 3.79 (m, 1H), 1.28 - 1.27 (d, J = 6.0 Hz, 6H)

[0143]

(Step 2) Methyl
 2,5-difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino
 15]benzoate



5-Bromo-2-(isopropoxymethyl)pyrimidine <see (step 1)> (508 mg, 2.20 mmol),
 2.20 [4-[(2,5-difluoro-4-methoxycarbonyl-phenyl)sulfamoyl]phenyl]boronic acid
 20 <see Example 10 (step 6)> (820 mg, 2.20 mmol), sodium carbonate (700 mg, 6.60 mmol), and Pd(dppf)Cl₂ (48 mg, 0.066 mmol) were dissolved in dioxane (20 ml) and water (7.0 ml), and the resulting mixture was stirred in the presence of nitrogen at 105°C for 12 hours. The reaction liquid was concentrated under reduced pressure, then the obtained residue was washed
 25 with diethyl ether (20 ml) and dried to obtain the title compound (1.50 g).


MS (ESI) m/z 478 (M+H)⁺

[0144]

(Step

3)

2,5-Difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino
]benzoic acid (M-19)



Methyl


2,5-difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino
]benzoate <see (step 2)> (1.50 g) was dissolved in methanol (10 ml), a 2 N
aqueous solution of lithium hydroxide (6.0 ml) was added thereto, and the
10 resulting mixture was stirred at room temperature for 2 hours. The
completion of the reaction was verified with TLC, and then 4 N hydrochloric
acid was added for adjustment of the pH to 4.0. The precipitated solid was
filtered off, washed with acetonitrile, and dried to obtain the title compound
(763 mg, 75% for two steps) as a yellow solid.

15 ¹H NMR (DMSO-d₆, 400 MHz): δ 13.41 (br, 1H), 11.15 (br, 1H), 9.19 (s, 2H),
8.07 - 7.97 (m, 4H), 7.62 (dd, J = 10.8, 6.4 Hz, 1H), 7.31 (dd, J = 12.0, 6.4 Hz,
1H), 4.66 (s, 2H), 3.81 - 3.75 (m, 1H), 1.24 (d, J = 6.0 Hz, 6H); MS (ESI) m/z
464 (M+H)⁺

[0145]

20 Example 20: Synthesis of M-20

(Step 1) 2-(Azetidin-1-yl)-5-bromo-pyrimidine




5-Bromo-2-chloro-pyrimidine (29.9 g, 0.150 mol) was dissolved in
N,N-dimethylformamide (300 ml), azetidine (11.4 g, 0.200 mol) and

potassium carbonate (42.5 g, 0.310 mol) were added thereto, and the resulting mixture was stirred at 50°C for 12 hours. The reaction liquid was diluted with water, extracted with diethyl ether (300 ml × 3), the extracts were combined, washed with a saturated saline solution, and dried over
 5 anhydrous sodium sulfate. The solvent was removed under reduced pressure to obtain the title compound (30 g, 93%).

¹H NMR (CDCl₃, 400 MHz): δ 8.26 (s, 2H), 4.12 (t, J = 7.4 Hz, 4H), 2.40-2.33 (m, 2H)

[0146]

10 (Step 2) Methyl
 4-[[4-[2-(azetidin-1-yl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-benzoate




A method similar to Example 10 (step 7) was performed to
 15 2-(Azetidin-1-yl)-5-bromo-pyrimidine <see (step 1)> (10.7 g, 50.4 mmol) to obtain the title compound (15.0 g, 64%).

MS (ESI) m/z 461 (M+H)⁺

[0147]

(Step 3)
 20 4-[[4-[2-(Azetidin-1-yl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-benzoic acid (M-20)



Methyl


4-[[4-[2-(azetidin-1-yl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-benzoate <see (step 2)> (15.0 g, 32.6 mmol) was dissolved in methanol (180 ml), a 2 N aqueous solution of lithium hydroxide (60 ml) was added thereto, and the resulting mixture was stirred at room temperature for 2 hours. The completion of the reaction was verified with TLC, and then 4 N hydrochloric acid was added for adjustment of the pH to 4.0. The precipitated solid was filtered, washed with a mixed solvent (methylene chloride:methanol = 10:1), and dried to obtain the title compound (10.0 g, 69%) as a white solid.

¹H NMR (DMSO-d₆, 400 MHz): δ 13.43 - 13.40 (br, 1H), 11.13 - 11.05 (br, 1H), 8.78 (d, J = 10.0 Hz, 2H), 7.98 - 7.88 (m, 4H), 7.62 (dd, J = 10.4, 6.8 Hz, 1H), 7.29 (dd, J = 12.0, 6.4 Hz, 1H), 4.12 (t, J = 7.2 Hz, 4H), 2.52 - 2.32 (m, 2H); MS (ESI) m/z 447 (M+H)⁺

[0148]

Example 21: Synthesis of M-21

(Step 1) Methyl 2,5-difluoro-4-[[5-(4-methyltriazol-1-yl)-2-pyridyl]sulfonylamino]benzoate



2-Butynoic acid (3.70 g, 44.0 mmol) and methyl 2,5-difluoro-4-[(5-iodine-2-pyridyl)sulfonylamino]benzoate <synthesized by a method similar to Example 16 (step 6)> (20.0 g, 44.0 mmol) were dissolved in dimethylsulfoxide (250 ml) and water (30 ml), L-proline (1.00 g, 8.80 mmol), copper(II) sulfate pentahydrate (1.10 g, 4.40 mmol), sodium ascorbate (1.70 g, 8.80 mmol), sodium azide (4.29 g, 66.0 mmol), and potassium carbonate (7.30 g, 53.0 mmol) were sequentially added, and the resulting mixture was stirred at 65°C for 12 hours. Ethyl acetate (500 ml), ammonium water (500 ml), and water (1,000 ml) were added to the reaction liquid, then the reaction

liquid was extracted with ethyl acetate (600 ml $\times$ 15), the organic layers were combined and washed with a saturated saline solution (3,000 ml), dried over sodium sulfate, and the solvent was removed under reduced pressure. The obtained residue was purified by HPLC separation to obtain the title compound (4.70 g, 26%).


^1H NMR (CD_3OD , 300 MHz): δ 9.11 (s, 1H), 8.43 - 8.42 (m, 2H), 8.22 - 8.19 (m, 1H), 7.43 - 7.30 (m, 2H), 3.83 (s, 3H), 2.43 (s, 3H)

[0149]

(Step

2)

2,5-Difluoro-4-[[5-(4-methyltriazol-1-yl)-2-pyridyl]sulfonylamino]benzoic acid (M-21)



Methyl

2,5-difluoro-4-[[5-(4-methyltriazol-1-yl)-2-pyridyl]sulfonylamino]benzoate


<see (step 1)> (4.50 g, 11.0 mmol) was dissolved in methanol (60 ml), a 2 N aqueous solution of lithium hydroxide (15 ml) was added thereto, and the resulting mixture was stirred at room temperature for 30 minutes. The completion of the reaction was verified with TLC, and then 4 N hydrochloric acid was added for adjustment of the pH to 4.0. The precipitated solid was filtered off, washed with a saturated saline solution, and dried to obtain the title compound (3.90 g, 90%) as a white solid.

^1H NMR (CD_3OD , 400 MHz): δ 9.22 (d, J = 2.0 Hz, 1H), 8.58 - 8.47 (d, J = 9.2, 2.8 Hz, 1H), 8.47 (s, 1H), 8.29 (d, J = 8.8 Hz, 1H), 7.66 - 7.58 (m, 2H), 2.44 (s, 3H); MS (ESI) m/z 396 ($\text{M}+\text{H}$) $^+$

[0150]

Example 22: Synthesis of M-22


(Step 1) Furan-2-sulfonyl chloride



Furan (3.67 g, 53.9 mmol) was dissolved in diethyl ether (50 ml), the reaction solution was cooled to 0°C, then a solution of t-butyllithium/hexane (1.3 mol/l, 50 ml) was slowly added dropwise, and the resulting mixture was stirred for 15 minutes. Sulfur dioxide was added to the reaction liquid, and the resulting mixture was further stirred for 15 minutes. Then N-chlorosuccinimide (8.65 g, 64.8 mmol) was added thereto, and the resulting mixture was stirred at room temperature for 30 minutes. The reaction liquid was diluted with water and extracted with ethyl acetate (100 ml × 2), then the extracts were combined, washed with a saturated saline solution, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and then the residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 10:1 to 5:1) to obtain the title compound (3.2 g, 36%) as a yellow oil.

[0151]

(Step 2) Methyl 2,5-difluoro-4-[(2-furylsulfonyl)amino]benzoate




Methyl 4-amino-2,5-difluorobenzoate (1.40 g, 7.48 mmol) <see Example 10 (step 4)> was dissolved in methylene chloride (60 ml), furan-2-sulfonyl chloride <see (step 1)> (3.00 g, 18.0 mmol) and pyridine (10 ml) were added thereto, and the resulting mixture was stirred at room temperature for 12 hours. The reaction liquid was diluted with water and extracted with ethyl acetate. The extracts were combined, washed with a saturated saline solution, dried over anhydrous sodium sulfate, the solvent was removed under reduced pressure, and then the obtained residue was

purified by silica gel column chromatography (petroleum ether:ethyl acetate = 10:1 to 5:1) to obtain the title compound (800 mg, 34%) as a gray solid.

^1H NMR ($\text{DMSO}-d_6$, 400 MHz): δ 8.02 (s, 1H), 7.71 - 6.66 (m, 1H), 7.32 - 7.25 (m, 2H), 6.70 - 6.66 (m, 1H), 3.82 (s, 3H); MS (ESI) m/z 318 ($\text{M}+\text{H}$) $^+$

5 [0152]

(Step 3) 2,5-Difluoro-4-[(2-furysulfonyl)amino]benzoic acid (M-22)



Methyl 2,5-difluoro-4-[(2-furysulfonyl)amino]benzoate <see (step 2)> (800 mg, 2.52 mmol) was dissolved in methanol (10 ml), a 6 N aqueous solution of sodium hydroxide (10.0 ml) was added thereto, and the resulting mixture was stirred at room temperature for 30 minutes. The completion of the reaction was verified with TLC, and then 4 N hydrochloric acid was added for adjustment of the pH to 4.0. The precipitated solid was filtered and dried to obtain the title compound (600 mg, 79%).

15 ^1H NMR (CD_3OD , 400 MHz): δ 7.74 (dd, J = 2.4, 1.6 Hz, 1H), 7.61 (dd, J = 14.4, 8.4 Hz, 1H), 7.38 (dd, J = 15.6, 8.4 Hz, 1H), 7.18 (dd, J = 4.8, 1.6 Hz, 1H), 6.58 (dd, J = 4.8, 2.4 Hz, 1H); MS (ESI) m/z 304 ($\text{M}+\text{H}$) $^+$

[0153]

Example 23: Synthesis of M-23

20 (Step 1) Methyl 4-amino-2,3,5,6-tetrafluoro-benzoate




4-Amino-2,3,5,6-tetrafluoro-benzoic acid (10.0 g, 47.8 mmol) was dissolved in methanol (80 ml), the reaction solution was cooled to 0°C, then thionyl chloride (15 ml) was added dropwise, and the resulting mixture was stirred at 65°C for 12 hours. The reaction liquid was concentrated under

reduced pressure, and then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 5:1 to 2:1) to obtain the title compound (9.28 g, 87%).

NMR (CDCl₃, 400 MHz): δ 4.38 (br s, 2H), 3.92 (s, 3H).

5 [0154]

(Step 2) Methyl 4-[(4-bromophenyl)sulfonylamino]-2,3,5,6-tetrafluoro-benzoate




Methyl 4-amino-2,3,5,6-tetrafluoro-benzoate <see (step 1)> (4.60 g, 20.6 mmol) was dissolved in tetrahydrofuran (50 ml), sodium hydride (60% in oil, 1.64 g, 41.0 mmol) was added thereto, and the resulting mixture was stirred in the presence of nitrogen gas at room temperature for 30 minutes. 4-Bromobenzenesulfonyl chloride (5.70 g, 22.3 mmol) was added to the reaction liquid, and the resulting mixture was stirred at 60°C for 2 hours. A saturated aqueous solution of sodium hydrogen carbonate was added to the reaction liquid and the resulting mixture was extracted with ethyl acetate (50 ml $\times$ 3). The extracts were combined, washed with a saturated saline solution, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 5:1 to 1:1) to obtain the title compound (6.29 g, 69%).

¹H NMR (CDCl₃, 400 MHz): δ 7.75 (d, J = 7.8 Hz, 2H), 7.50 (d, J = 7.8 Hz, 2H), 3.90 (s, 3H)

[0155]

(Step 3) Methyl 4-[[4-(2-ethylpyrimidin-5-yl)phenyl]sulfonylamino]-2,3,5,6-tetrafluoro-benzoate



Methyl

4-[(4-bromophenyl)sulfonylamino]-2,3,5,6-tetrafluorobenzoate <see (step 2)>
(587 mg, 1.33 mmol) and

5 2-ethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrimidine (311 mg, 1.33 mmol) were dissolved in N,N-dimethylformamide (25 ml) and water (4.0 ml), sodium carbonate (421 mg, 3.98 mmol) and Pd(dppf)Cl₂ (194 mg, 0.266 mmol) were added thereto, and the resulting mixture was stirred in the presence of nitrogen gas at 100°C for 4 hours. The reaction liquid was
10 concentrated under reduced pressure, and then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 5:1 to 1:1) to obtain the title compound (541 mg, 87%).


¹H NMR (CDCl₃, 400 MHz): δ 8.91 (s, 2H), 8.03 (d, J = 7.8 Hz, 2H), 7.73 (d, J = 7.8 Hz, 2H), 3.98 (s, 3H), 3.08 (q, J = 8.0 Hz, 2H), 1.42 (t, J = 8.0 Hz, 3H)

15 [0156]

(Step

4)

4-[[4-(2-Ethylpyrimidin-5-yl)phenyl]sulfonylamino]-2,3,5,6-tetrafluorobenzoic acid (M-23)



20 Methyl

4-[[4-(2-ethylpyrimidin-5-yl)phenyl]sulfonylamino]-2,3,5,6-tetrafluorobenzoate <see (step 3)> (541 mg, 1.15 mmol) was dissolved in methanol (5.0 ml), a 2 N aqueous solution of lithium hydroxide (1.0 ml) was added thereto, and the resulting mixture was stirred at room temperature for 30 minutes. The


completion of the reaction was verified with TLC, and then 4 N hydrochloric acid was added for adjustment of the pH to 4.0. The precipitated solid was filtered and dried to obtain the title compound (260 mg, 49%) as a white solid.

5 ^1H NMR ($\text{DMSO}-d_6$, 400 MHz): δ 9.31 (s, 2H), 8.11 (d, $J = 8.0$ Hz, 2H), 7.92 (d, $J = 8.0$ Hz, 2H), 3.04 (q, $J = 8.0$ Hz, 2H), 1.38 (t, $J = 8.0$ Hz, 3H); MS (ESI) m/z 456 ($\text{M}+\text{H}$)⁺

[0157]

Example 24: Synthesis of M-24

10 (Step 1) Methyl 4-[(4-bromo-2-methyl-phenyl)sulfonylamino]-2-fluoro-benzoate




Methyl 4-amino-2-fluoro-benzoate (2.00 g, 12.0 mmol) and 4-bromo-2-methyl-benzensulphonyl chloride (4.70 g, 17.0 mmol) were
15 dissolved in methylene chloride (50 ml), pyridine (10 ml) was added thereto, and the resulting mixture was stirred at room temperature for 12 hours. The reaction solution was concentrated under reduced pressure, then the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 4:1) to obtain the title compound (3.28 g,
20 69%).

MS (ESI) m/z 402, 404 ($\text{M}+\text{H}$)⁺

[0158]

(Step 2) Methyl 2-fluoro-4-[[2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]sulfonylamino]benzoate

25




Methyl

4-[(4-bromo-2-methyl-phenyl)sulfonylamino]-2-fluorobenzoate <see (step 1)>
 (2.30 g, 5.70 mmol) and bis(pinacolato)diborane (1.70 g, 6.90 mmol) were
 5 dissolved in N,N-dimethylformamide (30 ml),
 [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (22 mg, 0.030
 mmol) and potassium acetate (1.70 g, 17.0 mmol) were added thereto, and
 the resulting mixture was stirred in the presence of nitrogen gas at 85°C for
 12 hours. The reaction liquid was cooled to room temperature and then
 10 filtered, and the filtrate was extracted with methylene chloride (30 ml $\times$ 4).
 The extracts were combined and concentrated under reduced pressure, the
 obtained residue was diluted with water, the precipitated solid was filtered
 off, washed with a mixed solvent (petroleum ether:methylene chloride = 5:2),
 and dried to obtain the title compound (1.90 g, 74%) as a brown solid.
 15 ¹H NMR (DMSO-d₆, 400 MHz): δ 8.00 (d, J = 8.0 Hz, 1H), 7.95 (s, 1H), 7.92 (s,
 1H), 7.76 (t, J = 8.8 Hz, 1H), 7.68 - 7.68 (m, 1H), 6.97 (dd, J = 8.8, 2.0 Hz, 1H),
 6.90 (dd, J = 8.8, 2.0 Hz, 1H), 3.79 (s, 3H), 2.60 (s, 3H), 1.28 (s, 12H)
 [0159]

(Step 3)

20 [4-[(3-Fluoro-4-methoxycarbonyl-phenyl)sulfamoyl]-3-methyl-phenyl]boronic
 acid



Methyl

2-fluoro-4-[[2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]sulfonylamino]benzoate <see (step 2)> (1.50 g, 3.30 mmol) was dissolved in acetone (15 ml), sodium periodate (1.40 g, 6.60 mmol), ammonium acetate (500 mg, 6.60 mmol), and water (6.0 ml) were added thereto, and the
 5 resulting mixture was stirred at room temperature for 48 hours. The reaction liquid was concentrated under reduced pressure, then diluted with water, and extracted with ethyl acetate (500 ml $\times$ 3). The extracts were combined, washed with a saturated saline solution, and dried over anhydrous sodium sulfate, and then the solvent was removed under reduced
 10 pressure. The obtained residue was washed with a mixed solvent (petroleum ether:ethyl acetate = 1:10) and dried under reduced pressure to obtain the title compound (1.00 g, 82%).

^1H NMR (DMSO- d_6 , 400 MHz): δ 8.70 (d, J = 4.4 Hz, 1H), 7.76 (t, J = 8.4 Hz, 1H), 7.65 - 7.57 (m, 2H), 6.93 - 6.87 (m, 2H), 3.80 (s, 3H), 2.64 (s, 3H)


15 [0160]

(Step

4)

Methyl

2-fluoro-4-[[2-methyl-4-(2-pyridyl)phenyl]sulfonylamino]benzoate




[4-[(3-Fluoro-4-methoxycarbonyl-phenyl)sulfamoyl]-3-methyl-phenyl]

boronic acid <see (step 3)> (800 mg, 2.18 mmol) and 2-bromopyridine (680 mg, 4.35 mmol) were dissolved in dioxane (20 ml) and water (5.0 ml), sodium carbonate (530 mg, 5.00 mmol) and Pd(dppf)Cl₂ (162 mg, 0.200 mmol) were added thereto, and the resulting mixture was stirred in the presence of nitrogen at 110°C for 4 hours. The reaction liquid was concentrated under
 20 reduced pressure, and then the obtained residue was purified by silica gel column chromatography (n-hexane:ethyl acetate = 10:1 to 1:1) to obtain the
 25 title compound (530 mg, 61%).

¹H NMR (CD₃OD, 400 MHz): δ 8.70 (d, J = 8.0 Hz, 1H), 8.12 - 8.03 (m, 3H), 8.00 (d, J = 8.0 Hz, 1H), 7.96 - 7.90 (m, 2H), 7.76 (t, J = 8.4 Hz, 1H), 7.44 - 7.41 (m, 1H), 7.02 - 6.91 (m, 2H), 3.76 (s, 3H), 2.69 (s, 3H)

[0161]

5 (Step 5)
2-Fluoro-4-[[2-methyl-4-(2-pyridyl)phenyl]sulfonylamino]benzoic acid (M-24)



Methyl


2-fluoro-4-[[2-methyl-4-(2-pyridyl)phenyl]sulfonylamino]benzoate <see (step
10 4)> (630 mg, 1.57 mmol) was dissolved in methanol (10 ml), a 6 N aqueous
solution of sodium hydroxide (5.0 ml) was added thereto, and the resulting
mixture was stirred at room temperature for 30 minutes. The completion of
the reaction was verified with TLC, and then 4 N hydrochloric acid was
added for adjustment of the pH to 4.0. The precipitated solid was filtered
15 and dried to obtain the title compound (560 mg, 92%) as a white solid.

¹H NMR (CD₃OD, 400 MHz): δ 8.67 (d, J = 4.8 Hz, 1H), 8.18 (d, J = 8.0 Hz, 1H), 7.99 - 7.96 (m, 4H), 7.81 (t, J = 8.4 Hz, 1H), 7.47 - 7.45 (m, 1H), 6.99 - 6.94 (m, 2H), 2.77 (s, 3H); MS (ESI) m/z 385 (M-1)

[0162]

20 Example 25: Synthesis of M-25

(Step 1) 4-({[3,5-Difluoro-4-(methoxycarbonyl)phenyl]amino}sulfonyl)benzoic acid




4-(Chlorosulfonyl)benzoic acid (25.0 g, 113 mmol) and methyl

4-amino-2,6-difluorobenzoate (19.0 g, 101 mmol) were dissolved in methylene chloride (500 ml), pyridine (25.0 ml, 285 mmol) was added thereto, and the resulting mixture was stirred at room temperature for 12 hours. The reaction solution was concentrated under reduced pressure, the obtained
 5 residue was diluted with water, a 6 N aqueous solution of hydrochloric acid was added for adjustment of the pH to 1.0, and the precipitated solid was filtered and washed with water. The obtained solid was suspended in water again, washed with a saturated aqueous solution of sodium hydrogen carbonate, and then extracted with ethyl acetate (100 ml $\times$ 2). A 6 N
 10 aqueous solution of hydrochloric acid was added to the obtained aqueous layer for adjustment of the pH to 6.0, and ethyl acetate was added (100 ml $\times$ 2) for extraction. The extracts were combined, washed with a saturated saline solution, dried over anhydrous sodium sulfate, and then the solvent was removed to obtain the title compound (15.0 g, 36%).

15 ^1H NMR (d-DMSO, 400 MHz): δ 11.50 (s, 1H), 8.14 (d, J = 8.4 Hz, 2H), 8.01 (d, J = 8.4 Hz, 2H), 6.67 (d, J = 10.4 Hz, 2H), 3.87 (s, 3H); MS (ESI) m/z 372 (M+H)⁺
 [0163]

(Step 2)

20 4-[(4-[(Cyclopropylamino)carbonyl]phenyl)sulfonyl]amino]-2,6-difluorobenzoic acid (M-25)



4-[(3,5-Difluoro-4-(methoxycarbonyl)phenyl)amino]sulfonylbenzoic acid <see (step 1)> (1.5 g, 4.0 mmol) was dissolved in thionyl chloride (40 ml),
 25 and the resulting mixture was stirred at 75°C for 4 hours. The reaction solution was cooled to room temperature, and then concentrated under reduced pressure. The obtained residue was dissolved in methylene

chloride (30 ml), cyclopropylamine (680 mg, 12.0 mmol), and the resulting mixture was stirred at room temperature for 30 minutes. The solvent was removed under reduced pressure, and the obtained residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate = 2:1 to 1:1).

5 A part of the obtained compound (740 mg, 1.8 mmol) was dissolved in methanol (15 ml), a 3 N aqueous solution of sodium hydroxide (5.0 ml) was added thereto, and the resulting mixture was stirred at room temperature for 30 minutes. A 6 N aqueous solution of hydrochloric acid was added for adjustment of the pH to 4.0, and the precipitated solid was filtered and then
10 dried under reduced pressure to obtain the title compound (670 mg, 51% for two steps) as a white solid.


¹H NMR (d-DMSO, 400 MHz): δ 7.79 - 7.72 (m, 4H), 6.62 (d, J = 10 Hz, 2H), 2.67 - 2.65 (m, 1H), 0.63 - 0.59 (m, 2H), 0.45 - 0.43 (m, 2H); MS (ESI) m/z 397 (M+H)⁺

15 [0164]

The following compounds (Examples 26 to 47) were synthesized by using methods similar to Example 28 (step 1), (step 2) performed to intermediates corresponding to M-1 to M-25 and compounds similar thereto.

Example 26: Synthesis of A-1

20 (Step 1)
(2S)-2-[[2,5-Difluoro-4-[[4-[2-(4-piperidyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoic acid (A-1)




• 3TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.99 (s, 1H), 9.16 (s, 2H), 9.00 (s, 1H), 8.75 (dd, J = 7.8, 3.8 Hz, 1H), 8.57 (d, J = 4.9 Hz, 1H), 8.47 (d, J = 2.5 Hz, 1H), 8.07 - 7.94 (m, 4H), 7.90 (d, J = 5.0 Hz, 1H), 7.75 (dd, J = 8.2, 2.5 Hz, 1H), 7.47 (d, J = 8.3 Hz, 1H), 7.38 (dd, J = 10.3, 6.4 Hz, 1H), 7.29 (dd, J = 11.4, 6.3 Hz, 1H), 4.89 (td, J = 8.2, 4.9 Hz, 1H), 3.61 (s, 3H), 3.42 - 2.99 (m, 8H), 2.20 - 2.12 (m, 2H), 2.05 - 1.92 (m, 2H); MS (ESI) m/z 798.26 (M+H)⁺ [0165]

Example 27: Synthesis of A-2 and B-2

(Step 1)
(2S)-2-[[2,5-Difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propionic acid (A-2)




• 1TFA

Methyl
3-[5-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)pyridin-

2-yl]-L-alaninate (a methyl ester of M-9) <synthesized by a method similar to Example 9> (67.0 mg, 0.190 mmol) and 2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoic acid (M-11) <see Example 11 (step 4)> (75.0 mg, 0.160 mmol) were
 5 suspended in methylene chloride (2.0 ml), HATU (91.0 mg, 0.240 mmol) and diisopropylethylamine (0.167 ml, 0.960 mmol) were added thereto, and the resulting mixture was stirred at room temperature for 12 hours. The reaction liquid was concentrated under reduced pressure, then a 4 N solution of hydrochloric acid/dioxane (3.0 ml) and water (1.0 ml) were added thereto,
 10 and the resulting mixture was stirred at 60°C for 2 hours. The reaction liquid was concentrated under reduced pressure and then purified by reverse phase HPLC (an H₂O/CH₃CN system including 0.1% TFA) to obtain a TFA salt of the title compound (30.7 mg, 21%).

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.13 (s, 2H), 9.00 (s, 1H), 8.73 (dd, J = 7.9, 3.8 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.46 (d, J = 2.4 Hz, 1H), 8.05 - 7.99 (m, 2H), 7.98 - 7.93 (m, 2H), 7.90 (d, J = 4.9 Hz, 1H), 7.73 (dd, J = 8.2, 2.5 Hz, 1H), 7.45 (d, J = 8.2 Hz, 1H), 7.38 (dd, J = 10.4, 6.4 Hz, 1H), 7.28 (dd, J = 11.4, 6.3 Hz, 1H), 4.88 (td, J = 8.2, 5.0 Hz, 1H), 3.95 (ddd, J = 11.3, 4.4, 2.1 Hz, 2H), 3.61 (s, 3H), 3.48 (td, J = 11.6, 2.5 Hz, 2H), 3.42 - 3.23 (m, 2H), 3.13 (tt, J = 11.2, 4.2 Hz, 1H), 1.97 - 1.75 (m, 4H); MS (ESI) m/z 799.51 (M+H)⁺
 20 [0166]

(Step 2) Cyclohexyl
 (2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate (B-2)
 25



• 2TFA

Cyclohexyl

3-[5-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)pyridin-2-yl]-L-alaninate (M-9) <see Example 9 (step 4)> (107 mg, 0.250 mmol) and
 5 2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylami-
 no]benzoic acid (M-11) <see Example 11 (step 4)> (100 mg, 0.210 mmol) were
 suspended in methylene chloride (3.0 ml), HATU (120 mg, 0.320 mmol) and
 diisopropylethylamine (0.219 ml, 1.26 mmol) were added thereto, and the
 resulting mixture was stirred at room temperature for 12 hours. The
 10 reaction liquid was concentrated under reduced pressure and then purified
 by reverse phase HPLC (an H₂O/CH₃CN system including 0.1% TFA) to
 obtain a TFA salt of the title compound (48.7 mg, 21%).

¹H NMR (400 MHz, DMSO-d₆) δ 10.99 (s, 1H), 9.14 (s, 2H), 9.00 (s, 1H), 8.85
 (dd, J = 7.6, 3.4 Hz, 1H), 8.57 (d, J = 4.9 Hz, 1H), 8.45 (d, J = 2.4 Hz, 1H), 8.05
 15 - 8.00 (m, 2H), 7.99 - 7.94 (m, 2H), 7.90 (d, J = 4.9 Hz, 1H), 7.73 (dd, J = 8.2,
 2.5 Hz, 1H), 7.46 (d, J = 8.2 Hz, 1H), 7.38 (dd, J = 10.3, 6.3 Hz, 1H), 7.29 (dd,
 J = 11.3, 6.3 Hz, 1H), 4.91 (td, J = 7.6, 5.9 Hz, 1H), 4.68 (tt, J = 8.2, 3.7 Hz,
 1H), 4.01 - 3.90 (m, 2H), 3.61 (s, 3H), 3.38 - 3.25 (m, 2H), 3.19 - 3.07 (m, 1H),
 1.97 - 1.77 (m, 4H), 1.75 - 1.52 (m, 4H), 1.50 - 1.17 (m, 7H). ; MS (ESI)
 20 m/z881.56 (M+H)⁺


[0167]

Example 28: Synthesis of A-3 and B-3

(Step

1)

(2S)-2-[[2,5-Difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoic acid (A-3)



• 1TFA


5 ¹H NMR (400 MHz, DMSO-d₆) δ 10.99 (s, 1H), 9.18 (s, 2H), 9.00 (s, 1H), 8.73 (dd, J = 7.8, 3.8 Hz, 1H), 8.57 (d, J = 4.9 Hz, 1H), 8.46 (d, J = 2.4 Hz, 1H), 8.07 - 8.02 (m, 2H), 7.99 - 7.95 (m, 2H), 7.90 (d, J = 4.9 Hz, 1H), 7.73 (dd, J = 8.2, 2.5 Hz, 1H), 7.46 (d, J = 8.2 Hz, 1H), 7.38 (dd, J = 10.4, 6.4 Hz, 1H), 7.28 (dd, J = 11.3, 6.2 Hz, 1H), 4.93 - 4.84 (m, 1H), 4.63 (s, 2H), 3.61 (s, 3H), 3.40 (s, 3H), 3.38 - 3.23 (m, 2H); MS (ESI) m/z 759.46 (M+H)⁺
10 [0168]

(Step

2)

Cyclohexyl

(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate (B-3)
15



• 2TFA


¹H NMR (400 MHz, DMSO-d₆) δ 11.00 (s, 1H), 9.18 (s, 2H), 9.00 (s, 1H), 8.85 (dd, J = 7.6, 3.4 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.45 (d, J = 2.4 Hz, 1H), 8.08

· 8.02 (m, 2H), 8.01 · 7.95 (m, 2H), 7.90 (d, $J = 4.9$ Hz, 1H), 7.73 (dd, $J = 8.2$, 2.5 Hz, 1H), 7.46 (d, $J = 8.2$ Hz, 1H), 7.38 (dd, $J = 10.3$, 6.3 Hz, 1H), 7.29 (dd, $J = 11.3$, 6.3 Hz, 1H), 4.97 · 4.87 (m, 1H), 4.63 (s, 3H), 3.61 (s, 3H), 3.40 (s, 3H), 3.38 · 3.25 (m, 2H), 1.77 · 1.51 (m, 4H), 1.46 · 1.18 (m, 6H); MS (ESI) m/z
 5 841.55 (M+H)⁺
 [0169]

Example 29: Synthesis of A-4 and B-4

(Step 1)


(2S)-2-[[2,5-Difluoro-4-[[4-[4-(methoxymethyl)triazol-1-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoic acid (A-4)



· 1TFA

¹H NMR (400 MHz, DMSO- d_6) δ 10.97 (s, 1H), 9.00 (d, $J = 0.8$ Hz, 1H), 8.93 (s, 1H), 8.74 (dd, $J = 7.9$, 3.7 Hz, 1H), 8.58 (d, $J = 5.0$ Hz, 1H), 8.49 · 8.47 (m, 1H),
 15 8.20 · 8.11 (m, 2H), 8.05 · 8.00 (m, 2H), 7.90 (dd, $J = 4.9$, 0.7 Hz, 1H), 7.76 (dd, $J = 8.2$, 2.0 Hz, 1H), 7.48 (d, $J = 8.2$ Hz, 1H), 7.37 (dd, $J = 10.3$, 6.4 Hz, 1H), 7.27 (dd, $J = 11.2$, 6.2 Hz, 1H), 4.94 · 4.85 (m, 1H), 4.54 (s, 2H), 3.61 (s, 3H), 3.43 · 3.23 (m, 5H). ; MS (ESI) m/z 748.46 (M+H)⁺
 [0170]

(Step 2) Cyclohexyl
 (2S)-2-[[2,5-difluoro-4-[[4-[4-(methoxymethyl)triazol-1-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate (B-4)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.00 (s, 1H), 8.93 (s, 1H), 8.86 (dd, J = 7.7, 3.3 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.46 (d, J = 2.4 Hz, 1H), 8.21 - 8.12 (m, 2H), 8.07 - 7.99 (m, 2H), 7.90 (d, J = 4.9 Hz, 1H), 7.74 (dd, J = 8.2, 2.5 Hz, 1H), 7.47 (d, J = 8.2 Hz, 1H), 7.37 (dd, J = 10.3, 6.3 Hz, 1H), 7.28 (dd, J = 11.1, 6.2 Hz, 1H), 4.97 - 4.87 (m, 1H), 4.74 - 4.63 (m, 1H), 4.55 (s, 2H), 3.61 (s, 3H), 3.38 - 3.28 (m, 5H), 1.79 - 1.52 (m, 4H), 1.50 - 1.08 (m, 6H); MS (ESI) m/z 830.63 (M+H)⁺ [0171]

Example 30: Synthesis of A-5 and B-5

(Step 1)
(2S)-2-[[4-[[4-(1-Allyl-2-oxo-pyrimidin-5-yl)phenyl]sulfonylamino]-2,5-difluorophenyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoic acid (A-5)




• 1TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.91 - 10.88 (m, 1H), 9.05 (d, J = 3.3 Hz, 1H), 9.00 (d, J = 0.7 Hz, 1H), 8.78 - 8.72 (m, 2H), 8.58 (d, J = 4.9 Hz, 1H), 8.51 - 8.47 (m, 1H), 7.93 - 7.84 (m, 5H), 7.79 - 7.74 (m, 1H), 7.49 (d, J = 8.3 Hz, 1H), 7.37 (dd, J = 10.3, 6.4 Hz, 1H), 7.26 (dd, J = 11.4, 6.3 Hz, 1H), 6.07 - 5.95 (m, 1H), 5.27 - 5.19 (m, 2H), 4.93 - 4.86 (m, 1H), 4.58 - 4.53 (m, 2H), 3.62 (s, 3H), 3.42 - 3.26 (m, 2H). ; MS (ESI) m/z 771.58 (M+H)⁺

[0172]

(Step 2) Cyclohexyl
(2S)-2-[[4-[[4-(1-allyl-2-oxo-pyrimidin-5-yl)phenyl]sulfonylamino]-2,5-difluor
o-benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyr
idyl]propanoate (B-5)



• 2TFA


¹H NMR (400 MHz, DMSO-d₆) δ 10.90 (s, 1H), 9.04 (d, J = 3.4 Hz, 1H), 9.00 (s, 1H), 8.85 (dd, J = 7.6, 3.4 Hz, 1H), 8.72 (d, J = 3.4 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.45 (d, J = 2.3 Hz, 1H), 7.94 - 7.83 (m, 5H), 7.73 (dd, J = 8.2, 2.5 Hz, 1H), 7.46 (d, J = 8.2 Hz, 1H), 7.37 (dd, J = 10.3, 6.3 Hz, 1H), 7.26 (dd, J = 11.3, 6.3 Hz, 1H), 6.07 - 5.94 (m, 1H), 5.27 - 5.17 (m, 2H), 4.96 - 4.87 (m, 1H), 4.73 - 4.63 (m, 1H), 4.57 - 4.50 (m, 2H), 3.61 (s, 3H), 3.33 - 3.28 (m, 2H), 1.76 - 1.52 (m, 4H), 1.46 - 1.17 (m, 6H); MS (ESI) m/z 853.63 (M+H)⁺

[0173]

Example 31: Synthesis of A-6 and B-6

(Step 1)
(2S)-2-[[4-[[4-[2-(Ethoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-dif


luoro-benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoic acid (A-6)



• 1TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.99 (s, 1H), 9.17 (s, 2H), 9.00 (s, 1H), 8.74
 5 (dd, J = 7.9, 3.8 Hz, 1H), 8.58 (d, J = 5.0 Hz, 1H), 8.48 (dd, J = 2.6, 1.1 Hz, 1H),
 8.08 - 8.02 (m, 2H), 8.00 - 7.95 (m, 2H), 7.90 (dd, J = 5.0, 0.7 Hz, 1H), 7.76
 (ddd, J = 9.0, 3.3, 1.9 Hz, 1H), 7.48 (d, J = 8.2 Hz, 1H), 7.38 (dd, J = 10.4, 6.4
 Hz, 1H), 7.29 (dd, J = 11.3, 6.2 Hz, 1H), 4.89 (td, J = 8.2, 4.9 Hz, 1H), 4.66 (s,
 2H), 3.66 - 3.55 (m, 5H), 3.43 - 3.25 (m, 2H), 1.17 (t, J = 7.0 Hz, 3H). ; MS
 10 (ESI) m/z 773.58 (M+H)⁺
 [0174]

(Step 2) Cyclohexyl
 (2S)-2-[[4-[[4-[2-(ethoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-
 15 pyridyl]propanoate (B-6)



• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 11.00 (s, 1H), 9.18 (s, 2H), 9.00 (s, 1H), 8.86
 (dd, J = 7.6, 3.4 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.46 (d, J = 2.4 Hz, 1H), 8.09

- 8.02 (m, 2H), 8.00 - 7.96 (m, 2H), 7.91 (dd, $J = 5.0, 0.7$ Hz, 1H), 7.79 - 7.71 (m, 1H), 7.47 (d, $J = 8.3$ Hz, 1H), 7.38 (dd, $J = 10.3, 6.3$ Hz, 1H), 7.29 (dd, $J = 11.3, 6.3$ Hz, 1H), 4.98 - 4.87 (m, 1H), 4.66 (s, 3H), 3.65 - 3.57 (m, 5H), 3.39 - 3.24 (m, 2H), 1.80 - 1.52 (m, 4H), 1.50 - 1.21 (m, 6H), 1.17 (t, $J = 7.0$ Hz, 3H);

5 MS (ESI) m/z 855.67 ($M+H$)⁺


[0175]

Example 32: Synthesis of A-7 and B-7

(Step

1)

(2S)-2-[[2,5-Difluoro-4-[[4-[2-[(isopropylamino)methyl]pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoic acid (A-7)



• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 11.03 (s, 1H), 9.32 (s, 2H), 9.14 (s, 2H), 9.00 (s, 1H), 8.75 (dd, $J = 7.8, 4.0$ Hz, 1H), 8.58 (d, $J = 5.0$ Hz, 1H), 8.46 (d, $J = 2.5$ Hz, 1H), 8.13 - 8.08 (m, 2H), 8.02 - 7.99 (m, 2H), 7.90 (d, $J = 5.0$ Hz, 1H), 7.73 (dd, $J = 8.2, 2.5$ Hz, 1H), 7.45 (d, $J = 8.2$ Hz, 1H), 7.39 (dd, $J = 10.3, 6.4$ Hz, 1H), 7.31 (dd, $J = 11.4, 6.3$ Hz, 1H), 4.89 (td, $J = 8.1, 5.1$ Hz, 1H), 4.54 (t, $J = 6.2$ Hz, 2H), 3.61 (s, 3H), 3.47 (dq, $J = 11.6, 5.8, 5.4$ Hz, 1H), 3.40 - 3.24 (m, 2H), 1.32 (d, $J = 6.5$ Hz, 6H); MS (ESI) m/z 786.74 ($M+H$)⁺


20 [0176]

(Step

2)

Cyclohexyl

(2S)-2-[[2,5-difluoro-4-[[4-[2-[(isopropylamino)methyl]pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate (B-7)



• 3TFA


¹H NMR (400 MHz, DMSO-d₆) δ 11.04 (s, 1H), 9.32 (s, 2H), 9.12 (d, J = 6.9 Hz, 2H), 9.00 (d, J = 0.7 Hz, 1H), 8.87 (dd, J = 7.6, 3.5 Hz, 1H), 8.58 (d, J = 5.0 Hz, 1H), 8.45 (dd, J = 2.4, 0.7 Hz, 1H), 8.14 - 8.07 (m, 2H), 8.03 - 7.99 (m, 2H),
 5 7.91 (dd, J = 5.0, 0.7 Hz, 1H), 7.73 (dd, J = 8.2, 2.5 Hz, 1H), 7.49 - 7.44 (m, 1H), 7.40 (dd, J = 10.3, 6.4 Hz, 1H), 7.32 (dd, J = 11.3, 6.3 Hz, 1H), 4.96 - 4.88 (m, 1H), 4.72 - 4.63 (m, 1H), 4.57 - 4.51 (m, 2H), 3.61 (s, 3H), 3.53 - 3.42 (m, 1H), 3.38 - 3.24 (m, 2H), 1.75 - 1.52 (m, 4H), 1.46 - 1.21 (m, 12H); MS (ESI) m/z 868.71 (M+H)⁺

10 [0177]

Example 33: Synthesis of A-8 and B-8

(Step 1)

(2S)-2-[[4-[[4-[2-(2-Dimethylaminoethyl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoic acid (A-8)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 11.00 (s, 1H), 9.44 (s, 1H), 9.18 (s, 2H), 9.00 (s, 1H), 8.74 (dd, J = 7.8, 3.9 Hz, 1H), 8.58 (d, J = 5.0 Hz, 1H), 8.45 (dd, J = 2.4, 0.7 Hz, 1H), 8.07 - 8.02 (m, 2H), 8.00 - 7.95 (m, 2H), 7.90 (dd, J = 4.9, 0.8 Hz, 1H), 7.72 (dd, J = 8.2, 2.5 Hz, 1H), 7.45 (d, J = 8.2 Hz, 1H), 7.39 (dd, J = 10.4, 6.4 Hz, 1H), 7.29 (dd, J = 11.4, 6.3 Hz, 1H), 4.92 - 4.86 (m, 1H), 3.61 (s, 5H), 3.40 (dd, J = 8.2, 6.9 Hz, 2H), 3.38 - 3.24 (m, 2H), 2.88 (d, J = 4.4 Hz, 6H); MS (ESI) m/z 786.75 (M+H)⁺

[0178]

(Step 2) Cyclohexyl
 10 (2S)-2-[[4-[[4-[2-(2-dimethylaminoethyl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate (B-8)



• 3TFA

¹H NMR (400 MHz, DMSO-d₆) δ 11.01 (s, 1H), 9.31 (s, 1H), 9.18 (s, 2H), 9.00 (s, 1H), 8.87 (dd, J = 7.6, 3.4 Hz, 1H), 8.58 (d, J = 4.9 Hz, 1H), 8.47 (d, J = 2.4 Hz, 1H), 8.08 - 8.02 (m, 2H), 8.01 - 7.96 (m, 2H), 7.91 (dd, J = 5.0, 0.8 Hz, 1H), 7.76 (dd, J = 8.2, 2.5 Hz, 1H), 7.48 (d, J = 8.2 Hz, 1H), 7.39 (dd, J = 10.3, 6.4 Hz, 1H), 7.30 (dd, J = 11.3, 6.3 Hz, 1H), 4.98 - 4.86 (m, 1H), 4.73 - 4.64 (m, 1H), 3.65 - 3.57 (m, 5H), 3.47 - 3.25 (m, 4H), 2.88 (d, J = 4.9 Hz, 6H), 1.76 - 1.52 (m, 4H), 1.48 - 1.16 (m, 6H); MS (ESI) m/z 868.71 (M+H)⁺


[0179]

Example 34: Synthesis of A-9 and B-9

(Step

1)

(2S)-2-[[[2,5-Difluoro-4-[[5-(2-tetrahydropyran-4-yl-4-pyridyl)-2-pyridyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoic acid (A-9)



• 2TFA


- 5 ^1H NMR (400 MHz, DMSO- d_6) δ 11.15 (s, 1H), 9.25 (d, J = 2.2 Hz, 1H), 9.00 (s, 1H), 8.75 (dd, J = 7.2, 3.9 Hz, 2H), 8.63 - 8.55 (m, 2H), 8.48 (d, J = 2.4 Hz, 1H), 8.18 (d, J = 8.3 Hz, 1H), 8.07 - 7.87 (m, 3H), 7.78 - 7.74 (m, 1H), 7.52 - 7.37 (m, 3H), 4.95 - 4.86 (m, 1H), 4.02 - 3.95 (m, 2H), 3.61 (s, 3H), 3.51 - 3.42 (m, 2H), 3.42 - 3.26 (m, 2H), 3.17 - 3.08 (m, 1H), 1.95 - 1.80 (m, 4H); MS (ESI) m/z
- 10 799.70 ($M+H$)⁺
[0180]

(Step

2)

Cyclohexyl

- (2S)-2-[[[2,5-difluoro-4-[[5-(2-tetrahydropyran-4-yl-4-pyridyl)-2-pyridyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate (B-9)
- 15



• 3TFA


¹H NMR (400 MHz, DMSO-d₆) δ 11.15 (s, 1H), 9.24 (dd, J = 2.4, 0.8 Hz, 1H), 9.00 (s, 1H), 8.87 (dd, J = 7.6, 3.5 Hz, 1H), 8.74 (d, J = 5.4 Hz, 1H), 8.63 - 8.55 (m, 2H), 8.47 (dd, J = 2.4, 0.7 Hz, 1H), 8.17 (dd, J = 8.3, 0.8 Hz, 1H), 7.97 (s, 1H), 7.89 (ddd, J = 7.1, 5.2, 1.2 Hz, 2H), 7.75 (dd, J = 8.2, 2.5 Hz, 1H), 7.51 - 7.37 (m, 3H), 4.93 (td, J = 7.6, 5.9 Hz, 1H), 4.68 (td, J = 8.2, 3.9 Hz, 1H), 3.98 (ddd, J = 11.4, 4.2, 2.0 Hz, 2H), 3.61 (s, 3H), 3.46 (td, J = 11.4, 2.8 Hz, 2H), 3.40 - 3.26 (m, 2H), 3.16 - 3.05 (m, 1H), 1.97 - 1.78 (m, 4H), 1.76 - 1.53 (m, 4H), 1.33 (dddt, J = 34.0, 24.1, 17.5, 9.8 Hz, 6H); MS (ESI) m/z 881.71 (M+H)⁺

[0181]

Example 35: Synthesis of A-10 and B-10

(Step 1)


(2S)-2-[[[2,5-Difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoic acid (A-10)



• 1TFA

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.00 (s, 1H), 9.17 (s, 2H), 9.00 (d, J = 0.8 Hz, 1H), 8.74 (dd, J = 7.8, 3.8 Hz, 1H), 8.58 (d, J = 4.9 Hz, 1H), 8.49 (dd, J = 2.4, 0.7 Hz, 1H), 8.07 - 8.03 (m, 2H), 8.00 - 7.96 (m, 2H), 7.91 (dd, J = 4.9, 0.7 Hz, 1H), 7.78 (dd, J = 8.2, 2.5 Hz, 1H), 7.49 (d, J = 8.2 Hz, 1H), 7.39 (dd, J = 10.4, 6.4 Hz, 1H), 7.29 (dd, J = 11.4, 6.3 Hz, 1H), 4.90 (td, J = 8.3, 4.9 Hz, 1H), 4.65 (s, 2H), 3.83 - 3.72 (m, 1H), 3.61 (s, 3H), 3.43 - 3.25 (m, 2H), 1.16 (d, J = 6.1 Hz, 6H); MS (ESI) m/z 787.62($\text{M}+\text{H}$)⁺ [0182]

10 (Step 2) Cyclohexyl
(2S)-2-[[[2,5-difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoate (B-10)




• 2TFA

15 ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.00 (s, 1H), 9.17 (s, 2H), 9.00 (s, 1H), 8.86 (dd, J = 7.6, 3.4 Hz, 1H), 8.57 (d, J = 4.9 Hz, 1H), 8.47 (dd, J = 2.5, 0.7 Hz, 1H),

8.08 - 8.02 (m, 2H), 8.01 - 7.94 (m, 2H), 7.91 (dd, $J = 5.0, 0.8$ Hz, 1H), 7.75 (dd, $J = 8.2, 2.5$ Hz, 1H), 7.47 (d, $J = 8.3$ Hz, 1H), 7.38 (dd, $J = 10.3, 6.3$ Hz, 1H), 7.29 (dd, $J = 11.3, 6.3$ Hz, 1H), 4.97 - 4.88 (m, 1H), 4.72 - 4.62 (m, 3H), 3.77 (hept, $J = 6.1$ Hz, 1H), 3.61 (s, 3H), 3.39 - 3.25 (m, 2H), 1.76 - 1.53 (m, 4H),
 5 1.48 - 1.19 (m, 6H), 1.16 (d, $J = 6.1$ Hz, 6H); MS (ESI) m/z 869.67 (M+H)⁺
 [0183]

Example 36: Synthesis of A-11


(Step 1)
 (2S)-2-[[4-[[4-[2-(Azetidin-1-yl)pyrimidin-5-yl]phenyl]sulfonylamino]-2,5-difluoro-benzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoic acid (A-11)



¹H NMR (400 MHz, DMSO- d_6) δ 10.86 (s, 1H), 8.75 (s, 2H), 8.52 (dd, $J = 8.0, 2.7$ Hz, 1H), 7.87 (s, 4H), 7.30 - 7.20 (m, 4H), 7.10 - 7.05 (m, 2H), 4.60 (ddd, $J = 9.9, 7.9, 4.7$ Hz, 1H), 4.14 - 4.06 (m, 4H), 3.38 (s, 3H), 3.20 (s, 4H), 3.00 (dd, $J = 13.9, 9.9$ Hz, 1H), 2.40 - 2.29 (m, 2H), 2.04 (s, 3H); MS (ESI) m/z 746.66 (M+H)⁺
 [0184]

Example 37: Synthesis of A-12

(Step 1)
 (2S)-2-[[2,5-Difluoro-4-[[5-(4-methyltriazol-1-yl)-2-pyridyl]sulfonylamino]benzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoic acid (A-12)




¹H NMR (400 MHz, DMSO-d₆) δ 11.10 (s, 1H), 9.28 (dd, J = 2.5, 0.7 Hz, 1H), 8.75 (d, J = 0.9 Hz, 1H), 8.59 (dd, J = 8.6, 2.5 Hz, 1H), 8.55 (dd, J = 8.0, 2.6 Hz, 1H), 8.22 (dd, J = 8.6, 0.7 Hz, 1H), 7.36 (dd, J = 11.1, 6.3 Hz, 1H), 7.31 - 7.23 (m, 3H), 7.11 - 7.05 (m, 2H), 4.62 (ddd, J = 9.9, 7.9, 4.7 Hz, 1H), 3.39 (s, 3H), 3.20 (s, 4H), 3.01 (dd, J = 13.9, 9.8 Hz, 1H), 2.36 (d, J = 0.9 Hz, 3H), 2.06 (s, 3H); MS (ESI) m/z 695.56 (M+H)⁺

[0185]

Example 38: Synthesis of A-13

10 (Step 1) 1)
(2S)-2-[[2,5-Difluoro-4-(2-furysulfonylamino)benzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoic acid (A-13)




15 ¹H NMR (400 MHz, DMSO-d₆) δ 11.10 (s, 1H), 8.61 (dd, J = 8.0, 2.4 Hz, 1H), 8.02 (dd, J = 1.8, 0.9 Hz, 1H), 7.31 - 7.25 (m, 3H), 7.24 - 7.16 (m, 2H), 7.11 - 7.07 (m, 2H), 6.68 (dd, J = 3.6, 1.8 Hz, 1H), 4.62 (ddd, J = 9.9, 7.9, 4.7 Hz, 1H), 3.40 (s, 3H), 3.21 (s, 4H), 3.02 (dd, J = 13.9, 9.9 Hz, 1H), 2.07 (s, 3H); MS (ESI) m/z 603.49 (M+H)⁺

[0186]

20 Example 39: Synthesis of A-14

(Step 1)
 (2S)-2-[[2,5-Difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonylaminobenzoyl]amino]-3-[4-[1-methyl-6-(methylaminomethyl)-2,4-dioxo-quinazolin-3-yl]phenyl]propanoic acid (A-14)



• TFA


5

MS (ESI) m/z 800.99 (M+H)⁺

[0187]

Example 40: Synthesis of A-15

(Step 1)
 10 (2S)-2-[[2,5-Difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylaminobenzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoic acid (A-15)



¹H NMR (400 MHz, DMSO-d₆) δ 10.96 (s, 1H), 9.17 (s, 2H), 8.54 (dd, J = 7.9, 2.6 Hz, 1H), 8.07 - 8.03 (m, 2H), 7.98 - 7.94 (m, 2H), 7.30 - 7.22 (m, 4H), 7.09 - 7.05 (m, 2H), 4.68 - 4.57 (m, 3H), 3.78 (hept, J = 6.1 Hz, 1H), 3.38 (s, 3H), 3.20 (s, 4H), 3.00 (dd, J = 13.9, 9.9 Hz, 1H), 2.05 (s, 3H), 1.16 (d, J = 6.1 Hz, 6H);
 15 MS (ESI) m/z 763.58 (M+H)⁺


[0188]

Example 41: Synthesis of A-16

(Step

1)

(2S)-2-[[[2,5-Difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(6-methoxy-1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyridyl]propanoic acid (A-16)



• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.16 (s, 2H), 8.73 (dd, J = 7.9, 3.8 Hz, 1H), 8.56 (d, J = 0.8 Hz, 1H), 8.46 (dd, J = 2.4, 0.7 Hz, 1H), 8.06 - 8.01 (m, 2H), 7.99 - 7.95 (m, 2H), 7.73 (dd, J = 8.2, 2.5 Hz, 1H), 7.45 (d, J = 8.3 Hz, 1H), 7.38 (dd, J = 10.4, 6.4 Hz, 1H), 7.32 - 7.25 (m, 2H), 4.88 (td, J = 8.2, 5.0 Hz, 1H), 4.65 (s, 2H), 3.93 (s, 3H), 3.76 (dq, J = 12.1, 6.1 Hz, 1H), 3.55 (s, 3H), 3.40 - 3.24 (m, 2H), 1.15 (d, J = 6.1 Hz, 6H); MS (ESI) m/z 817.59 (M+H)⁺


[0189]

Example 42: Synthesis of A-17

(Step

1)


(2S)-2-[[[2,5-Difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[4-(3-methyl-2,6-dioxo-pyrimidin-1-yl)phenyl]propanoic acid (A-17)



¹H NMR (400 MHz, DMSO-d₆) δ 10.96 (s, 1H), 9.17 (s, 2H), 8.59 (dd, J = 7.8, 2.6 Hz, 1H), 8.07 - 8.02 (m, 2H), 7.99 - 7.95 (m, 2H), 7.74 (d, J = 7.9 Hz, 1H), 7.35 - 7.23 (m, 4H), 7.12 - 7.06 (m, 2H), 5.73 (dd, J = 7.9, 2.2 Hz, 1H), 4.66 (s, 2H), 4.63 - 4.55 (m, 1H), 3.83 - 3.72 (m, 1H), 3.29 (s, 3H), 3.19 (dd, J = 14.1, 4.6 Hz, 1H), 3.04 (dd, J = 14.1, 9.8 Hz, 1H), 1.16 (d, J = 6.1 Hz, 6H); MS (ESI) m/z 735.50 (M+H)⁺
[0190]

Example 43: Synthesis of A-18

10 (Step 1)
(2S)-2-[[[2,5-Difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrimido[4,5-d]pyrimidin-3-yl)-2-pyridyl]propanoic acid (A-18)



• 2TFA

15 ¹H NMR (400 MHz, DMSO-d₆) δ 10.99 (s, 1H), 9.28 (s, 1H), 9.19 (s, 1H), 9.16 (s, 2H), 8.73 (dd, J = 7.9, 3.7 Hz, 1H), 8.46 (d, J = 2.5 Hz, 1H), 8.07 - 8.02 (m, 2H), 7.99 - 7.95 (m, 2H), 7.73 (dd, J = 8.2, 2.5 Hz, 1H), 7.47 (d, J = 8.2 Hz, 1H), 7.37 (dd, J = 10.4, 6.4 Hz, 1H), 7.28 (dd, J = 11.4, 6.3 Hz, 1H), 4.93 - 4.85 (m,


1H), 4.65 (s, 2H), 3.77 (hept, J = 6.1 Hz, 1H), 3.55 (s, 3H), 3.41 - 3.23 (m, 2H), 1.15 (d, J = 6.1 Hz, 6H); MS (ESI) m/z 788.63 (M+H)⁺

[0191]

Example 44: Synthesis of A-19

5 (Step 1)

(2S)-2-[[2,5-Difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[4-[3-methyl-2,6-dioxo-4-(trifluoromethyl)pyrimidin-1-yl]phenyl]propanoic acid (A-19)



10 ¹H NMR (400 MHz, DMSO-d₆) δ 10.96 (s, 1H), 9.17 (s, 2H), 8.59 (dd, J = 7.9, 2.5 Hz, 1H), 8.07 - 8.02 (m, 2H), 7.99 - 7.94 (m, 2H), 7.37 - 7.32 (m, 2H), 7.32 - 7.23 (m, 2H), 7.18 - 7.12 (m, 2H), 6.48 (s, 1H), 4.68 - 4.56 (m, 3H), 3.78 (hept, J = 6.1 Hz, 1H), 3.38 - 3.35 (m, 3H), 3.21 (dd, J = 14.0, 4.6 Hz, 1H), 3.04 (dd, J = 14.1, 9.9 Hz, 1H), 1.16 (d, J = 6.1 Hz, 6H); MS (ESI) m/z 803.62 (M+H)⁺


15 [0192]

Example 45: Synthesis of A-20

(Step 1)

(2S)-2-[[4-[[4-(2-Ethylpyrimidin-5-yl)phenyl]sulfonylamino]-2,3,5,6-tetrafluorobenzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoic acid (A-20)

20




^1H NMR (400 MHz, DMSO-d_6) δ 10.83 (s, 1H), 9.33 (d, $J = 8.1$ Hz, 1H), 9.14 (s, 2H), 8.08 - 8.02 (m, 2H), 7.93 - 7.87 (m, 2H), 7.27 (d, $J = 7.8$ Hz, 2H), 7.11 - 7.06 (m, 2H), 4.73 - 4.65 (m, 1H), 3.38 (s, 3H), 3.25 - 3.18 (m, 4H), 3.00 - 2.91 (m, 3H), 2.06 (s, 3H), 1.32 (t, $J = 7.6$ Hz, 3H); MS (ESI) m/z 755.50 ($\text{M}+\text{H}$)⁺ [0193]

Example 46: Synthesis of A-21

(Step

1)

(2S)-2-[[[2-Fluoro-4-[[2-methyl-4-(2-pyridyl)phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoic acid (A-21)




^1H NMR (400 MHz, DMSO-d_6) δ 11.12 (s, 1H), 8.72 - 8.68 (m, 1H), 8.28 (dd, $J = 7.9, 3.1$ Hz, 1H), 8.12 - 8.03 (m, 4H), 7.95 (td, $J = 7.8, 1.8$ Hz, 1H), 7.47 - 7.36 (m, 2H), 7.26 - 7.21 (m, 2H), 7.07 - 7.02 (m, 2H), 6.96 (dd, $J = 8.5, 2.1$ Hz, 1H), 6.89 (dd, $J = 12.4, 2.1$ Hz, 1H), 4.62 - 4.53 (m, 1H), 3.38 (s, 3H), 3.22 - 3.12 (m, 4H), 2.98 (dd, $J = 13.9, 9.8$ Hz, 1H), 2.68 (s, 3H), 2.04 (s, 3H); MS (ESI) m/z 686.62 ($\text{M}+\text{H}$)⁺ [0194]

Example 47: Synthesis of A-22

(Step 1)

(2S)-2-[[4-[[4-(Cyclopropylcarbamoyl)phenyl]sulfonylamino]-2,6-difluorobenzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoic acid (A-22)



^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.13 (s, 1H), 8.97 (d, $J = 8.0$ Hz, 1H), 8.63 (d, $J = 4.3$ Hz, 1H), 7.97 - 7.89 (m, 4H), 7.27 - 7.23 (m, 2H), 7.09 - 7.04 (m, 2H), 6.76 (d, $J = 9.0$ Hz, 2H), 4.57 (ddd, $J = 10.1, 8.0, 4.6$ Hz, 1H), 3.41 (s, 3H), 3.21 (s, 3H), 3.14 (dd, $J = 14.0, 4.6$ Hz, 1H), 2.97 - 2.79 (m, 2H), 2.08 (s, 3H), 0.72 - 0.66 (m, 2H), 0.58 - 0.52 (m, 2H); MS (ESI) m/z 696.58 ($\text{M}+\text{H}$)⁺


The following compounds (Examples 48 to 79) were synthesized by using methods similar to Example 51 (step 1), (step 2) and Example 52 (step 1) performed to intermediates corresponding to M-1 to M-25 and compounds similar thereto.

[0195]

Example 48: Synthesis of C-1

(Step 1) Tetrahydropyran-4-yl

(2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)phenyl]propanoate (C-1)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.14 (s, 2H), 8.97 (s, 1H), 8.81 (dd, J = 7.3, 1.9 Hz, 1H), 8.54 (d, J = 5.0 Hz, 1H), 8.07 - 8.01 (m, 2H), 8.01 - 7.94 (m, 2H), 7.88 (d, J = 5.0 Hz, 1H), 7.40 - 7.25 (m, 4H), 7.24 - 7.19 (m, 2H),
 4.94 - 4.84 (m, 1H), 4.68 - 4.61 (m, 1H), 4.00 - 3.91 (m, 2H), 3.79 - 3.66 (m, 2H),
 3.59 (s, 3H), 3.53 - 3.44 (m, 4H), 3.22 - 3.07 (m, 3H), 1.95 - 1.71 (m, 6H), 1.57 - 1.39 (m, 2H); MS (ESI) m/z 882.68 (M+H)⁺
 [0196]

Example 49: Synthesis of C-2

(Step 1) Tetrahydropyran-4-ylmethyl (2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-ylpyrimidin-5-yl)phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)phenyl]propanoate (C-2)



0.03 cm

• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 11.01 - 10.91 (m, 1H), 9.14 (s, 2H), 8.97 (s, 1H), 8.80 (dd, J = 7.5, 2.0 Hz, 1H), 8.55 (d, J = 5.0 Hz, 1H), 8.07 - 7.94 (m, 4H),


Example 50: Synthesis of C-3

O=C1OC(=O)C[C@H](N1C(=O)c2cc(F)c(S(=O)(=O)Nc3ccc(cc3c4nc5cnc(C6CCOCC6)c5n4)cc4)cc2)c2cc(F)c(S(=O)(=O)Nc3ccc(cc3c4nc5cnc(C6CCOCC6)c5n4)cc4)cc2

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.14 (s, 2H), 8.99 (s, 1H), 8.88 (dd, J = 7.6, 1.7 Hz, 1H), 8.56 (d, J = 4.9 Hz, 1H), 8.47 (d, J = 2.3 Hz, 1H), 8.05 - 8.01 (m, 2H), 7.99 - 7.95 (m, 2H), 7.93 (dd, J = 8.2, 2.4 Hz, 1H), 7.89 (d, J = 4.9 Hz, 1H), 7.41 (d, J = 8.0 Hz, 1H), 7.29 (ddd, J = 10.8, 6.3, 3.9 Hz, 2H), 4.97 - 4.84 (m, 1H), 4.76 - 4.66 (m, 1H), 3.99 - 3.90 (m, 2H), 3.79 - 3.66 (m, 2H), 3.59 (s, 3H), 3.54 - 3.42 (m, 4H), 3.26 (dd, J = 14.1, 5.8 Hz, 1H), 3.19 - 3.07 (m, 2H), 1.96 - 1.72 (m, 6H), 1.58 - 1.39 (m, 2H); MS (ESI) m/z 883.72 (M+H)⁺ [0198]

(Step	1)	Cyclohexyl
(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl		
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-		

3-pyridyl]propanoate



• 2TFA

Cyclohexyl

3-[6-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)pyridin-
 5 3-yl]-L-alaninate (M-1) <see Example 1> (127 mg, 0.280 mmol) and
 2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]b
 enzoic acid (M-11) <see Example 11> (100 mg, 0.230 mmol) were suspended
 in methylene chloride (4.0 ml), HATU (131 mg, 0.350 mmol) and
 diisopropylethylamine (0.160 ml, 0.920 mmol) were added thereto, and the
 10 resulting mixture was stirred at room temperature for 12 hours. The
 reaction liquid was concentrated under reduced pressure and then purified
 by reverse phase HPLC (an H₂O/CH₃CN system including 0.1% TFA) to
 obtain a TFA salt of the title compound (130 mg, 53%) as a white solid.


¹H NMR (400 MHz, DMSO-d₆): δ 10.97 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.84
 15 (dd, J = 7.7, 1.7 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.49 - 8.43 (m, 1H), 8.10 -
 8.01 (m, 2H), 8.01 - 7.95 (m, 2H), 7.95 - 7.85 (m, 2H), 7.40 (dd, J = 8.0, 0.8 Hz,
 1H), 7.33 - 7.23 (m, 2H), 4.76 - 4.64 (m, 2H), 4.63 (s, 2H), 3.59 (s, 3H), 3.40 (s,
 3H), 3.24 (dd, J = 14.1, 5.6 Hz, 1H), 3.12 (dd, J = 14.1, 9.7 Hz, 1H), 1.78 - 1.58
 (m, 4H), 1.50 - 1.19 (m, 6H); MS(ESI) m/z 841.59(M+H)⁺

20 [0199]

(Step

2)

(2S)-2-[[[2,5-Difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
 amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
 3-pyridyl]propanoic acid




• 1TFA

A 4 N solution of hydrochloric acid/dioxane (3.0 ml) and water (2.0 ml) were added to cyclohexyl (2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate <see (step 1)> (70.0 mg, 65.0 mmol), and the resulting mixture was stirred at 60°C for 4 hours. The reaction liquid was concentrated under reduced pressure and then purified by reverse phase HPLC (an H₂O/CH₃CN system including 0.1% TFA) to obtain a TFA salt of the title compound (42.3 mg, 66%) as a white solid.

¹H NMR (400 MHz, DMSO-d₆): δ 10.96 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.69 (dd, J = 8.1, 2.2 Hz, 1H), 8.56 (d, J = 4.9 Hz, 1H), 8.46 (d, J = 2.3 Hz, 1H), 8.09 - 8.01 (m, 2H), 8.01 - 7.93 (m, 2H), 7.93 - 7.85 (m, 2H), 7.40 (d, J = 8.0 Hz, 1H), 7.34 - 7.23 (m, 2H), 4.74 - 4.64 (m, 1H), 4.62 (s, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.29 (dd, J = 14.0, 4.6 Hz, 1H), 3.09 (dd, J = 14.1, 10.2 Hz, 1H); MS(ESI) m/z 759.5(M+H)⁺
[0200]

Example 52: Synthesis of C-4

(Step 1) Tetrahydropyran-4-yl
(2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-4)



• 2TFA


A 4 N solution of hydrochloric acid/dioxane (3.0 ml) and tetrahydro-4-pyranol (1.0 ml) were added to (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate <see (step 2)> (100 mg, 101 mmol), and the resulting mixture was stirred at 60°C for 12 hours. The reaction liquid was concentrated under reduced pressure and then purified by reverse phase HPLC (an H₂O/CH₃CN system including 0.1% TFA) to obtain a TFA salt of the title compound (39.6 mg, 37%) as a white solid.

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.91 - 8.86 (m, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.47 (d, J = 2.3 Hz, 1H), 8.07 - 8.03 (m, 2H), 8.00 - 7.96 (m, 2H), 7.93 (dd, J = 8.2, 2.4 Hz, 1H), 7.89 (d, J = 4.9 Hz, 1H), 7.41 (d, J = 8.1 Hz, 1H), 7.33 - 7.24 (m, 2H), 4.96 - 4.86 (m, 1H), 4.76 - 4.67 (m, 1H), 4.63 (s, 2H), 3.80 - 3.67 (m, 2H), 3.59 (s, 3H), 3.49 - 3.44 (m, 2H), 3.40 (s, 3H), 3.26 (dd, J = 14.1, 5.8 Hz, 1H), 3.14 (dd, 1H), 1.88 - 1.72 (m, 2H), 1.59 - 1.40 (m, 2H); MS (ESI) m/z 843.59 (M+H)⁺

[0201]

Example 53: Synthesis of C-5

(Step 1) Tetrahydropyran-4-ylmethyl (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-5)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.87 (dd, J = 7.7, 1.6 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.47 (d, J = 2.3 Hz, 1H), 8.09 - 8.02 (m, 2H), 8.00 - 7.96 (m, 2H), 7.94 - 7.86 (m, 2H), 7.40 (d, J = 8.0 Hz, 1H), 7.32 - 7.24 (m, 2H), 4.79 - 4.70 (m, 1H), 4.63 (s, 2H), 3.99 - 3.89 (m, 2H), 3.84 - 3.76 (m, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.30 - 3.20 (m, 3H), 3.13 (dd, J = 14.1, 9.9 Hz, 1H), 1.89 - 1.74 (m, 1H), 1.56 - 1.47 (m, 2H), 1.29 - 1.12 (m, 2H); MS (ESI) m/z 857.75 (M+H)⁺

[0202]

Example 54: Synthesis of C-6

(Step 1) Cyclohexylmethyl (2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl]amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-6)



• 2TFA


¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.87 - 8.83 (m, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.48 - 8.46 (m, 1H), 8.08 - 8.02 (m, 2H), 8.00 - 7.95 (m, 2H), 7.93 - 7.87 (m, 2H), 7.42 - 7.38 (m, 1H), 7.31 - 7.23 (m, 2H),

4.79 - 4.71 (m, 1H), 4.63 (s, 2H), 3.96 - 3.84 (m, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.28 (dd, J = 14.1, 5.0 Hz, 1H), 3.22 - 3.06 (m, 1H), 1.68 - 1.54 (m, 5H), 1.22 - 0.87 (m, 6H); MS (ESI) m/z 855.67 (M+H)⁺

[0203]

5 Example 55: Synthesis of C-7

(Step 1) Isopropyl (2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-7)



10

• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.19 (s, 2H), 8.99 (d, J = 0.8 Hz, 1H), 8.83 (dd, J = 7.7, 1.8 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.46 (dd, J = 2.5, 0.8 Hz, 1H), 8.08 - 8.04 (m, 2H), 8.01 - 7.96 (m, 2H), 7.95 - 7.87 (m, 2H), 7.40 (dd, J = 8.1, 0.7 Hz, 1H), 7.34 - 7.22 (m, 2H), 4.91 (hept, J = 6.3 Hz, 1H), 4.72 - 4.59 (m, 3H), 3.59 (s, 3H), 3.40 (s, 3H), 3.28 - 3.06 (m, 2H), 1.16 (dd, J = 18.9, 6.2 Hz, 6H); MS (ESI) m/z 801.63 (M+H)⁺


15

[0204]

Example 56: Synthesis of C-8

(Step 1) 2-Ethylbutyl (2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-8)

20




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.84 (dd, J = 8.0, 1.8 Hz, 1H), 8.57 (d, J = 4.9 Hz, 1H), 8.47 (d, J = 2.3 Hz, 1H), 8.08 - 8.03 (m, 2H), 8.00 - 7.96 (m, 2H), 7.93 - 7.87 (m, 2H), 7.42 - 7.38 (m, 1H),
 5 7.31 - 7.22 (m, 2H), 4.79 - 4.71 (m, 1H), 4.63 (s, 2H), 4.01 (d, J = 5.6 Hz, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.28 (dd, J = 14.2, 4.9 Hz, 1H), 3.12 (dd, J = 14.1, 10.3 Hz, 1H), 1.46 (hept, J = 6.2 Hz, 1H), 1.32 - 1.23 (m, 4H), 0.81 (t, J = 7.4 Hz, 6H); MS (ESI) m/z 843.71 (M+H)⁺

[0205]

10 Example 57: C-9

(Step 1) 2-Adamantyl
 (2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
 amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
 3-pyridyl]propanoate (C-9)



15

• 2TFA


¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.19 (s, 2H), 8.99 (s, 1H), 8.86 (dd, J = 7.8, 1.7 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.48 (d, J = 2.3 Hz, 1H), 8.07 - 8.03 (m, 2H), 7.99 - 7.96 (m, 2H), 7.93 (dd, J = 8.2, 2.4 Hz, 1H), 7.88 (d, J =

4.9 Hz, 1H), 7.40 (d, J = 8.0 Hz, 1H), 7.31 - 7.23 (m, 2H), 4.86 (d, J = 3.2 Hz, 1H), 4.82 - 4.72 (m, 1H), 4.63 (s, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.30 (dd, J = 14.1, 5.1 Hz, 1H), 3.14 (dd, J = 14.1, 10.2 Hz, 1H), 1.99 - 1.40 (m, 14H); MS (ESI) m/z 893.72 (M+H)⁺

5 [0206]

Example 58: C-10

(Step 1) (1-Methyl-4-piperidyl)
(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
amino]benzoylamino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3
10 -pyridyl]propanoate (C-10)




• 3TFA

MS (ESI) m/z 856.67 (M+H)⁺

[0207]

Example 59: Synthesis of C-11

15 (Step 1) 1-Ethylpropyl
(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
3-pyridyl]propanoate (C-11)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.97 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.85 (dd, J = 8.0, 1.6 Hz, 1H), 8.56 (d, J = 5.0 Hz, 1H), 8.48 (d, J = 2.3 Hz, 1H), 8.07 - 8.04 (m, 2H), 8.00 - 7.96 (m, 2H), 7.93 (dd, J = 8.2, 2.4 Hz, 1H), 7.88 (d, J = 4.9 Hz, 1H), 7.40 (d, J = 8.1 Hz, 1H), 7.31 - 7.21 (m, 2H), 4.78 - 4.66 (m, 2H), 4.63 (s, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.27 (dd, J = 14.1, 5.3 Hz, 1H), 3.13 (dd, J = 14.1, 10.2 Hz, 1H), 1.62 - 1.41 (m, 4H), 0.86 - 0.76 (m, 6H); MS (ESI) m/z 829.63 (M+H)⁺

[0208]

Example 60: Synthesis of C-12

(Step 1) (4,4-Dimethylcyclohexyl) (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-12)



• 2TFA


15

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.83 (dd, J = 7.6, 1.8 Hz, 1H), 8.57 (d, J = 4.9 Hz, 1H), 8.46 (d, J = 2.3 Hz, 1H), 8.08 - 8.02 (m, 2H), 8.00 - 7.96 (m, 2H), 7.94 - 7.87 (m, 2H), 7.40 (d, J = 8.1 Hz, 1H),

7.32 - 7.25 (m, 2H), 4.74 - 4.66 (m, 2H), 4.63 (s, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.25 (dd, J = 14.1, 5.4 Hz, 1H), 3.13 (dd, J = 14.1, 9.8 Hz, 1H), 1.73 - 1.11 (m, 8H), 0.85 (d, J = 15.3 Hz, 6H); MS (ESI) m/z 869.71 (M+H)⁺
[0209]

5 Example 61: Synthesis of C-13

(Step 1) 1-Propylbutyl
(2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
3-pyridyl]propanoate (C-13)




10 • 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.97 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.84 (dd, J = 7.8, 1.6 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.48 (dd, J = 2.3, 0.8 Hz, 1H), 8.07 - 8.03 (m, 2H), 8.00 - 7.96 (m, 2H), 7.93 (dd, J = 8.2, 2.4 Hz, 1H), 7.88 (dd, J = 4.9, 0.7 Hz, 1H), 7.40 (dd, J = 8.1, 0.7 Hz, 1H), 7.25 (ddd, J = 17.9, 10.5, 6.2 Hz, 2H), 4.86 (p, J = 6.3 Hz, 1H), 4.77 - 4.68 (m, 1H), 4.63 (s, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.27 (dd, J = 14.2, 5.0 Hz, 1H), 3.11 (dd, J = 14.1, 10.4 Hz, 1H), 1.52 - 1.42 (m, 4H), 1.35 - 1.17 (m, 4H), 0.87 - 0.81 (m, 6H); MS (ESI) m/z 857.71 (M+H)⁺
[0210]

20 Example 62: Synthesis of C-14

(Step 1) (4,4-Difluorocyclohexyl)
(2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
3-pyridyl]propanoate (C-14)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.99 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.90 (dd, J = 7.5, 1.6 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.47 (d, J = 2.3 Hz, 1H), 8.08 - 8.03 (m, 2H), 8.00 - 7.96 (m, 2H), 7.93 (dd, J = 8.1, 2.4 Hz, 1H), 7.89 (dd, J = 5.0, 0.7 Hz, 1H), 7.43 - 7.39 (m, 1H), 7.32 - 7.25 (m, 2H), 4.96 - 4.88 (m, 1H), 4.77 - 4.69 (m, 1H), 4.63 (s, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.27 (dd, J = 14.1, 5.6 Hz, 1H), 3.14 (dd, J = 14.1, 9.7 Hz, 1H), 2.05 - 1.63 (m, 8H); MS (ESI) m/z 877.72 (M+H)⁺

[0211]

Example 63: Synthesis of C-15

(Step 1) Ethyl (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-15)



• 2TFA


15

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.19 (s, 2H), 8.99 (s, 1H), 8.86 (dd, J = 7.8, 1.7 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.47 (d, J = 2.3 Hz, 1H), 8.08 - 8.03 (m, 2H), 8.00 - 7.96 (m, 2H), 7.94 - 7.87 (m, 2H), 7.42 - 7.39 (m, 1H),

7.33 - 7.25 (m, 2H), 4.75 - 4.68 (m, 1H), 4.63 (s, 2H), 4.11 (q, $J = 7.1$ Hz, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.25 (dd, $J = 14.0, 5.6$ Hz, 1H), 3.13 (dd, $J = 14.0, 9.7$ Hz, 1H), 1.16 (t, $J = 7.1$ Hz, 3H); MS (ESI) m/z 787.63 (M+H)⁺
[0212]

5 Example 64: Synthesis of C-16


(Step 1) Cyclopentyl
(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
3-pyridyl]propanoate (C-16)



10 • 2TFA
¹H NMR (400 MHz, DMSO- d_6) δ 10.98 (s, 1H), 9.19 (s, 2H), 8.99 (s, 1H), 8.83 (dd, $J = 7.7, 1.7$ Hz, 1H), 8.57 (d, $J = 4.9$ Hz, 1H), 8.46 (d, $J = 2.3$ Hz, 1H), 8.08 - 8.03 (m, 2H), 8.01 - 7.95 (m, 2H), 7.93 - 7.87 (m, 2H), 7.40 (d, $J = 8.1$ Hz, 1H), 7.32 - 7.24 (m, 2H), 5.14 - 5.07 (m, 1H), 4.72 - 4.59 (m, 3H), 3.59 (s, 3H), 3.40
15 (s, 3H), 3.24 (dd, $J = 14.1, 5.5$ Hz, 1H), 3.11 (dd, $J = 14.2, 9.7$ Hz, 1H), 1.86 - 1.46 (m, 8H); MS (ESI) m/z 827.71 (M+H)⁺
[0213]

Example 65: Synthesis of C-17

(Step 1) Hexyl
20 (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
3-pyridyl]propanoate (C-17)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.84 (dd, J = 8.0, 1.8 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.46 (d, J = 2.3 Hz, 1H), 8.08 - 8.03 (m, 2H), 8.00 - 7.95 (m, 2H), 7.93 - 7.86 (m, 2H), 7.40 (d, J = 8.1 Hz, 1H),
 5 7.31 - 7.24 (m, 2H), 4.76 - 4.69 (m, 1H), 4.63 (s, 2H), 4.09 - 4.02 (m, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.27 (dd, J = 14.1, 5.2 Hz, 1H), 3.12 (dd, J = 14.1, 10.1 Hz, 1H), 1.58 - 1.48 (m, 2H), 1.32 - 1.18 (m, 6H), 0.83 - 0.77 (m, 3H); MS (ESI) m/z 843.71 (M+H)⁺

[0214]

Example 66: Synthesis of C-18

(Step 1) Cycloheptyl (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyridol[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-18)




• 2TFA

15

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.83 (dd, J = 7.5, 1.7 Hz, 1H), 8.57 (d, J = 4.9 Hz, 1H), 8.46 (d, J = 2.3 Hz, 1H), 8.08 - 8.03 (m, 2H), 8.00 - 7.96 (m, 2H), 7.93 - 7.87 (m, 2H), 7.40 (d, J = 8.1 Hz, 1H),


(Step	1)	Methyl
(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl		
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-		
3-pyridyl]propanoate (C-19)		



• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 11.00 (s, 1H), 9.19 (s, 2H), 8.98 (s, 1H), 8.86 (dd, J = 8.1, 2.0 Hz, 1H), 8.56 (d, J = 5.0 Hz, 1H), 8.49 (d, J = 2.3 Hz, 1H), 8.09 - 8.03 (m, 2H), 8.02 - 7.98 (m, 2H), 7.93 (dd, J = 8.2, 2.4 Hz, 1H), 7.90 - 7.87 (m, 1H), 7.43 (d, J = 8.1 Hz, 1H), 7.31 (td, J = 10.8, 10.4, 6.3 Hz, 2H), 4.82 - 4.75 (m, 1H), 4.63 (s, 2H), 3.68 (s, 3H), 3.60 (s, 3H), 3.40 (s, 3H), 3.30 (dd, J = 14.0, 5.1 Hz, 1H), 3.15 (dd, J = 14.1, 10.1 Hz, 1H); MS (ESI) m/z 773.62 (M+H)⁺

20	(Step 1)	2,2-Dimethylpropyl (2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)- 3-pyridyl]propanoate (C-20)
----	----------	---




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.98 (s, 1H), 8.86 (dd, J = 7.9, 1.7 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.47 (d, J = 2.3 Hz, 1H), 8.07 - 8.03 (m, 2H), 7.99 - 7.96 (m, 2H), 7.92 (dd, J = 8.2, 2.4 Hz, 1H), 7.88 (d, J = 5.0 Hz, 1H), 7.40 (d, J = 8.0 Hz, 1H), 7.30 - 7.23 (m, 2H), 4.83 - 4.76 (m, 1H), 4.62 (s, 2H), 3.79 (d, J = 1.6 Hz, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.31 (dd, J = 14.2, 5.0 Hz, 1H), 3.14 (dd, J = 14.1, 10.4 Hz, 1H), 0.88 (s, 9H); MS (ESI) m/z

[0217]

Example 69: Synthesis of C-21

(Step 1) Cyclopentylmethyl (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-21)



• 2TFA

15


¹H NMR (400 MHz, DMSO-d₆) δ 10.99 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.85 (dd, J = 7.9, 1.8 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.47 (d, J = 2.3 Hz, 1H), 8.08 - 8.04 (m, 2H), 8.00 - 7.96 (m, 2H), 7.94 - 7.87 (m, 2H), 7.43 - 7.39 (m, 1H),

7.31 - 7.24 (m, 2H), 4.78 - 4.70 (m, 1H), 4.63 (s, 2H), 3.97 (d, $J = 6.9$ Hz, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.32 - 3.23 (m, 1H), 3.13 (dd, $J = 14.1, 10.1$ Hz, 1H), 2.13 (hept, $J = 7.4$ Hz, 1H), 1.72 - 1.40 (m, 6H), 1.25 - 1.15 (m, 2H); MS (ESI) m/z 841.67 (M+H)⁺

5 [0218]

Example 70: Synthesis of C-22

(Step 1) 1-Adamantylmethyl
(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
10 3-pyridyl]propanoate (C-22)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.19 (s, 2H), 8.99 (s, 1H), 8.84 (dd, $J = 8.0, 1.8$ Hz, 1H), 8.57 (d, $J = 4.9$ Hz, 1H), 8.48 (d, $J = 2.3$ Hz, 1H), 8.08 - 8.03 (m, 2H), 7.99 - 7.95 (m, 2H), 7.94 - 7.86 (m, 2H), 7.43 - 7.37 (m, 1H),
15 7.32 - 7.22 (m, 2H), 4.83 - 4.74 (m, 1H), 4.63 (s, 2H), 3.67 (d, $J = 1.8$ Hz, 3H), 3.59 (s, 3H), 3.39 (s, 3H), 3.31 (dd, $J = 14.1, 4.7$ Hz, 1H), 3.14 (dd, $J = 14.1, 10.5$ Hz, 1H), 2.00 - 1.40 (m, 14H); MS (ESI) m/z 907.76 (M+H)⁺
[0219]

Example 71: Synthesis of C-23

20 (Step 1) Cyclobutylmethyl
(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
3-pyridyl]propanoate (C-23)




• 2TFA

^1H NMR (400 MHz, DMSO- d_6) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.99 (d, J = 0.7 Hz, 1H), 8.86 (dd, J = 7.7, 1.7 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.47 (dd, J = 2.4, 0.8 Hz, 1H), 8.09 - 8.04 (m, 2H), 8.00 - 7.95 (m, 2H), 7.94 - 7.87 (m, 2H), 7.40 (dd, J = 8.0, 0.8 Hz, 1H), 7.31 - 7.25 (m, 2H), 4.79 - 4.70 (m, 1H), 4.63 (s, 2H), 4.05 (d, J = 6.5 Hz, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.27 (dd, J = 14.1, 5.2 Hz, 1H), 3.13 (dd, J = 14.1, 10.0 Hz, 1H), 2.61 - 2.52 (m, 1H), 2.00 - 1.91 (m, 2H), 1.87 - 1.66 (m, 4H); MS (ESI) m/z 827.71 ($M+H$)⁺
[0220]

Example 72: Synthesis of C-24

(Step 1) 2-Cyclohexylethyl (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-24)



• 2TFA


^1H NMR (400 MHz, DMSO- d_6) δ 10.99 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.84 (dd, J = 7.8, 1.9 Hz, 1H), 8.57 (d, J = 4.9 Hz, 1H), 8.47 (dd, J = 2.4, 0.8 Hz, 1H), 8.09 - 8.03 (m, 2H), 8.01 - 7.95 (m, 2H), 7.93 - 7.86 (m, 2H), 7.40 (dd, J = 7.9,

0.8 Hz, 1H), 7.34 - 7.24 (m, 2H), 4.76 - 4.68 (m, 1H), 4.63 (s, 2H), 4.14 - 4.08 (m, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.31 - 3.22 (m, 1H), 3.13 (dd, $J = 14.1, 10.1$ Hz, 1H), 1.74 - 0.74 (m, 13H); MS (ESI) m/z 869.75 (M+H)⁺

[0221]

5 Example 73: Synthesis of C-25

(Step 1) (1-Isobutyl-3-methyl-butyl)
(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
3-pyridyl]propanoate (C-25)



• 2TFA


10

MS (ESI) m/z 885.80 (M+H)⁺

[0222]

Example 74: Synthesis of C-26

(Step 1) 4,4,4'-Trifluorobutyl
15 (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
3-pyridyl]propanoate (C-26)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.89 (dd, J = 7.5, 1.7 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.48 - 8.46 (m, 1H), 8.07 - 8.03 (m, 2H), 7.99 - 7.96 (m, 2H), 7.92 (dd, J = 8.1, 2.4 Hz, 1H), 7.88 (dd, J = 4.9, 0.8 Hz, 1H), 7.41 (dd, J = 8.0, 0.7 Hz, 1H), 7.31 - 7.24 (m, 2H), 4.78 - 4.70 (m, 1H), 4.63 (s, 2H), 4.21 - 4.06 (m, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.29 (dd, J = 14.1, 5.1 Hz, 1H), 3.13 (dd, J = 14.1, 10.1 Hz, 1H), 2.39 - 2.23 (m, 2H), 1.84 - 1.74 (m, 2H); MS (ESI) m/z 869.63 (M+H)⁺
[0223]

Example 75: Synthesis of C-27

(Step 1) Cyclobutyl
(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
3-pyridyl]propanoate (C-27)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.19 (s, 2H), 8.99 (d, J = 0.8 Hz, 1H), 8.85 (dd, J = 7.7, 1.7 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.47 (dd, J = 2.3, 0.8 Hz, 1H), 8.08 - 8.02 (m, 2H), 8.01 - 7.96 (m, 2H), 7.94 - 7.87 (m, 2H), 7.41 (dd, J = 8.1, 0.7 Hz, 1H), 7.33 - 7.25 (m, 2H), 4.98 - 4.87 (m, 1H), 4.74 - 4.67 (m, 1H), 4.63 (s, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.25 (dd, J = 14.2, 5.7 Hz, 1H), 3.12 (dd, J = 14.0, 9.7 Hz, 1H), 2.31 - 2.19 (m, 2H), 2.06 - 1.87 (m, 2H), 1.79 - 1.68 (m, 1H), 1.65 - 1.50 (m, 1H); MS (ESI) m/z 813.67 (M+H)⁺
[0224]

Example 76: Synthesis of C-28

(Step 1) 2,2-Difluoroethyl
 (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
 amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
 3-pyridyl]propanoate (C-28)




• 2TFA

5
¹H NMR (400 MHz, DMSO-d₆) δ 10.99 (s, 1H), 9.16 (s, 2H), 9.00 (s, 1H), 8.75
 (dd, J = 7.8, 3.8 Hz, 1H), 8.57 (d, J = 4.9 Hz, 1H), 8.47 (d, J = 2.5 Hz, 1H), 8.07
 - 7.94 (m, 4H), 7.90 (d, J = 5.0 Hz, 1H), 7.75 (dd, J = 8.2, 2.5 Hz, 1H), 7.47 (d,
 J = 8.3 Hz, 1H), 7.38 (dd, J = 10.3, 6.4 Hz, 1H), 7.29 (dd, J = 11.4, 6.3 Hz, 1H),
 10 4.89 (td, J = 8.2, 4.9 Hz, 1H), 3.61 (s, 3H), 3.42 - 2.99 (m, 8H), 2.20 - 2.12 (m,
 2H), 2.05 - 1.92 (m, 2H); MS (ESI) m/z 823.63 (M+H)⁺
 [0225]

Example 77: Synthesis of C-29

(Step 1) 3-Cyclohexylpropyl
 15 (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl
 amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-
 3-pyridyl]propanoate (C-29)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.99 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.84 (dd, J = 7.8, 1.8 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.46 (d, J = 2.3 Hz, 1H), 8.09 - 8.02 (m, 2H), 8.01 - 7.95 (m, 2H), 7.94 - 7.86 (m, 2H), 7.40 (d, J = 8.1 Hz, 1H), 7.32 - 7.24 (m, 2H), 4.76 - 4.67 (m, 1H), 4.63 (s, 2H), 4.08 - 4.01 (m, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.31 - 3.21 (m, 1H), 3.17 - 3.07 (m, 1H), 1.68 - 1.49 (m, 7H), 1.22 - 0.97 (m, 6H), 0.85 - 0.70 (m, 2H); MS (ESI) m/z 883.76 (M+H)⁺ [0226]

Example 78: Synthesis of C-30

(Step 1) Isopentyl
 10 (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-30)




• 2TFA

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.99 (d, J = 0.7 Hz, 1H), 8.86 - 8.83 (m, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.47 - 8.45 (m, 1H), 8.07 - 8.04 (m, 2H), 8.00 - 7.96 (m, 2H), 7.93 - 7.86 (m, 2H), 7.40 (dd, J = 8.1, 0.7 Hz, 1H), 7.31 - 7.23 (m, 2H), 4.76 - 4.68 (m, 1H), 4.63 (s, 2H), 4.10 (t, J = 6.7 Hz, 2H), 3.59 (s, 3H), 3.40 (s, 3H), 3.26 (dd, J = 14.0, 5.2 Hz, 1H), 3.12 (dd, J = 14.1, 10.0 Hz, 1H), 1.71 - 1.56 (m, 1H), 1.45 (q, J = 6.8 Hz, 2H), 0.85 (d, J = 6.6 Hz, 6H); MS (ESI) m/z 829.71 (M+H)⁺ [0227]

Example 79: Synthesis of C-31

(Step 1) 2-Cyclopentylethyl
 (2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl

aminobenzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-31)



• 2TFA


MS (ESI) m/z 855.99 (M+H)⁺

5 [0228]

Further, by the following methods, intermediate compounds M-26 to M-30, compounds A-24 to A-30 and compounds C-32 to C-46 were synthesized.

[0229]

10 Example 80: Synthesis of
4-[[4-(4-ethyltriazol-1-yl)phenyl]sulfonylamino]-2,5-difluorobenzoic acid
(M-26)
(Step 1) Methyl
4-[[4-(4-ethyltriazol-1-yl)phenyl]sulfonylamino]-2,5-difluoro-benzoate



15


L-proline (1.6 g, 20 mol %, 14 mmol), CuSO₄·5H₂O (1.7 g, 10 mol %, 6.8 mmol), sodium ascorbate (2.7 g, 20 mol %, 14 mmol), sodium azide (6.50 g, 100 mmol), potassium carbonate (11.4 g, 82.6 mmol), methyl 2-pentynoate (6.76 g, 69 mmol), methyl
20 2,5-difluoro-4-[(4-iodophenyl)sulfonylamino]benzoate obtained in Example

13 (Step 2) (31.2 g, 69 mmol), DMSO (225 mL), and water (25 mL) were sequentially added in a 1 L vial. The resulting mixture was gently stirred at 65°C overnight, and a mixture of concentrated ammonium water (500 mL), water (1.0 L), and ethyl acetate (400 mL) was added. The aqueous phase
 5 was extracted with ethyl acetate (600 mL $\times$ 10), the organic phase was washed with a saturated saline solution (3.0 L), dried over sodium sulfate and filtered, and the solvent was distilled off under reduced pressure to obtain the title compound (14.2 g, 49%).

^1H NMR (CD_3OD , 400 MHz): δ 8.37 (s, 1H), 8.13-8.06 (m, 2H), 8.10-7.96 (m,
 10 2H), 7.45- 7.04 (m, 2H), 3.27 (s, 3H), 1.72-1.64 (m, 2H), 1.04-1.02 (m, 3H).

[0230]

(Step 2)
 4-[[4-(4-Ethyltriazol-1-yl)phenyl]sulfonylamino]-2,5-difluorobenzoic acid
 (M-26)




The compound obtained in step 1 (14.1 g, 33.4 mmol) was dissolved in methanol (60 mL), a 2 N aqueous solution of lithium hydroxide (30 mL) was added, then the resulting mixture was stirred at room temperature for 30 minutes, then completion of the reaction was verified, and 4 N hydrochloric
 20 acid was added for adjustment of the pH to 4 to 5. The precipitated white solid was filtered and the obtained solid was dried to obtain the title compound as a white solid.

^1H NMR ($\text{DMSO}-d_6$, 400 MHz): δ 13.41 (br s, 1H), 11.10 (br s, 1H), 8.70 (s, 1H), 8.13 (d, J = 8.8 Hz, 2H), 8.04 (d, J = 8.8 Hz, 2H), 7.63-7.59 (m, 1H),
 25 7.30-7.25 (m, 1H), 2.73 (q, J = 7.2 Hz, 2H), 1.26 (t, J = 7.2 Hz, 3H); MS (ESI)

m/z 409 (M+H)⁺.

[0231]

Example 81: Synthesis of
2,5-Difluoro-4-[(4-tetrahydropyran-4-ylphenyl)sulfonylamino]benzoic acid
5 (M-27)
(Step 1) Methyl
4-[[4-(3,6-dihydro-2H-pyran-4-yl)phenyl]sulfonylamino]-2,5-difluorobenzoate




10 Methyl 2,5-difluoro-4-[(4-iodophenyl)sulfonylamino]benzoate
obtained in step 2 of Example 13 (0.2 g, 0.44 mmol) was dissolved in
N,N'-dimethylformamide (5 mL) and water (1 mL),
3,6-dihydro-2H-pyran-4-boronic acid pinacol ester (0.11 g, 0.53 mmol),
[1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (18 mg, 0.022
15 mmol), and potassium acetate (0.15 g, 1.1 mmol) were added thereto, and the
resulting mixture was stirred at 100°C for 4 hours. The reaction solution
was cooled to room temperature and filtered through Celite, and the filtrate
was concentrated under reduced pressure. The obtained residue was
purified by silica gel column chromatography (hexane/ethyl acetate).

20 MS (ESI) m/z 410 (M+H)⁺.

[0232]

(Step 2)
2,5-Difluoro-4-[(4-tetrahydropyran-4-ylphenyl)sulfonylamino]benzoic acid
(M-27)




The compound obtained in step 1 was dissolved in methanol (20 mL), 10% palladium/carbon of a catalyst amount was added thereto, and the resulting mixture was stirred in the presence of hydrogen gas at room temperature overnight. The reaction solution was filtered, and the filtrate was concentrated. A 2 N solution of sodium hydroxide (4 mL) and methanol were added to the obtained residue, and the resulting mixture was stirred at room temperature for 3 hours. The reaction liquid was neutralized with 2 N hydrochloric acid, purified by reverse phase HPLC (an H₂O/CH₃CN system including 0.1% TFA), and freeze-dried to obtain the title compound.

¹H NMR (400 MHz, DMSO-*d*₆) δ 10.97 (s, 1H), 7.84 - 7.77 (m, 2H), 7.60 (dd, J = 10.7, 6.6 Hz, 1H), 7.53 - 7.47 (m, 2H), 7.24 (dd, J = 11.8, 6.4 Hz, 1H), 3.98 - 3.88 (m, 2H), 3.46 - 3.38 (m, 2H), 2.93 - 2.80 (m, 1H), 1.77 - 1.54 (m, 4H); MS (ESI) *m/z* 398 (M+H)⁺.

[0233]

Example	82:	Synthesis	of
2,5-difluoro-4-{{4-(2-isopropylpyrimidin-5-yl)phenyl}sulfonylamino}Benzoic acid (M-28)			



2-Isopropylpyrimidin-5-ylboronic acid was used instead of 3,6-dihydro-2H-pyran-4-boronic acid pinacol ester obtained in step 1 of


Example 81, and the title compound was synthesized by a similar method.

MS (ESI) m/z 434 (M+H)⁺.

[0234]

Example 83: Synthesis of methyl
 5 (2S)-2-amino-3-[5-(1,3,4-trimethyl-2,6-dioxypyrimidin-5-yl)-2-pyridyl]propanoate (M-29)

(Step 1) Methyl
 (2S)-2-(tert-butoxycarbonylamino)-3-[(5-(4,4,5,5-tetramethyl-1,3,2-dioxaboron-2-yl)-2-pyridyl)]propanoate



10

Methyl


(2S)-3-(5-bromo-2-pyridyl)-2-(tert-butoxycarbonylamino)propanoate (556 mg, 1.55 mmol) was dissolved in N,N'-dimethylformamide (10 mL), bis(pinacolato)diborane (588 mg, 2.32 mmol),
 15 [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (57 mg, 0.078 mmol), and potassium acetate (456 mg, 4.64 mmol) were added thereto, and the resulting mixture was stirred at 80°C overnight. The reaction solution was cooled to room temperature and filtered through Celite, and the filtrate was concentrated under reduced pressure. Ethyl acetate was added to the
 20 obtained residue, and the mixture was sequentially washed with water and a saturated saline solution. The solvent was distilled off under reduced pressure, and the obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) to obtain the title compound.

MS (ESI) m/z 407 (M+H)⁺.

25 [0235]

(Step 2)

(2S)-2-Amino-3-[5-(1,3,4-trimethyl-2,6-dioxypyrimidin-5-yl)-2-pyridyl]propanoate (M-29)




The compound obtained in step 1 was used instead of N-(tert-butoxycarbonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-L-phenylalaninate obtained in step 5 of Example 2, and the title compound was synthesized by a similar method.

MS (ESI) m/z 333 (M+H)⁺.

[0236]

Example 84: Synthesis of Methyl 3-[6-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)pyridin-3-yl]-L-alaninate (M-30)

(Step 1) Methyl 3-(((5-bromopyridin-2-yl)amino)carbonyl)amino)isonicotinate




Methyl 3-aminoisonicotinate (4.80 g, 32.0 mmol) and diisopropylethylamine (8.20 g, 64.0 mmol) were dissolved in methylene chloride (100 mL), a solution of triphosgene (3.10 g, 10.4 mmol) in methylene chloride (20 mL) was added thereto, and the resulting mixture was stirred at 0°C for 3 hours. 5-Bromopyridin-2-amine (4.60 g, 26.6 mmol) was added to this solution, and the resulting mixture was further stirred at room temperature for 12 hours. The reaction solution was concentrated under reduced pressure, then ethyl acetate was added thereto, the resulting mixture was washed with a saturated aqueous solution of sodium hydrogen

carbonate, water, and a saturated saline solution, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the obtained residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate = 1:1 to 1:2) to obtain the title compound (6.2 g, 67%).

^1H NMR (400 MHz, DMSO- d_6): δ 11.0 (s, 1H), 10.3 (s, 1H), 9.4 (s, 1H), 8.39 (d, J = 5.2 Hz, 2H), 8.0 (d, J = 8.8 Hz, 1H), 7.54 (d, J = 4.8 Hz, 1H), 7.48 (d, J = 8.8 Hz, 1H), 3.92 (s, 3H).

[0237]

(Step 2)
3-(5-Bromopyridin-2-yl)-1-methylpyrido[3,4-d]pyrimidine-2,4(1H,3H)-dione



Methyl

3-((5-bromopyridin-2-yl)amino)carbonyl)amino)isonicotinate (6.20 g, 17.7 mmol) was dissolved in N,N'-dimethylformamide (60 mL), an aqueous solution (4.0 mL) of potassium carbonate (600 mg, 4.34 mmol) was added thereto, and the resulting mixture was stirred at room temperature for 3 hours. p-Toluenesulfonic acid methyl ester (4.40 g, 23.6 mmol) and potassium carbonate (3.00 g, 21.7 mmol) were added to the reaction solution, and the resulting mixture was stirred at room temperature for 12 hours. The reaction solution was diluted with water (100 mL) and extracted with ethyl acetate (100 mL $\times$ 3). The extracts were combined, washed with a saturated saline solution, and dried over anhydrous sodium sulfate, and then the solvent was removed under reduced pressure. The obtained residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate = 1:1) to obtain the title compound (5.2 g, 88%).

^1H NMR (400 MHz, DMSO- d_6): δ 9.00 (s, 1H), 8.77 (d, J = 2.4 Hz, 1H), 8.58 (d,

$J = 4.8$ Hz, 1H), 8.32-8.30 (m, 1H), 7.91 (d, $J = 4.8$ Hz, 1H), 7.52 (d, $J = 8.8$ Hz, 1H), 3.61 (s, 3H).


[0238]

(Step

3)

Methyl

- 5 N-(tert-butoxycarbonyl)-3-[6-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)pyridin-3-yl]-L-alaninate




- Zinc (2.40 g, 36.7 mmol) was heated at 200°C for 30 minutes and cooled to room temperature, this operation was repeated 3 times, then a solution of dibromoethane (0.340 g, 1.82 mmol) in N,N'-dimethylformamide (5.0 mL) was added. An operation of heating the obtained reaction mixture to 90°C and cooling the reaction mixture to room temperature was repeated twice, then chlorotrimethylsilane (40.0 mg, 0.364 mmol) was added, and the resulting mixture was stirred at room temperature for 30 minutes. A solution of methyl N-(tert-butoxycarbonyl)-3-iodin-L-alaninate (2.00 g, 6.08 mmol) in N,N'-dimethylformamide (5.0 mL) was added dropwise, the resulting mixture was stirred at 35°C for 2.5 hours, then 3-(5-bromopyridin-2-yl)-1-methylpyrido[3,4-d]pyrimidine-2,4(1H,3H)-dione (2.60 g, 7.80 mmol) and bis(triphenylphosphine)palladium(II) chloride (230 mg, 0.328 mmol) were further added to the reaction solution, and the resulting mixture was stirred at 70°C for 2 hours. The reaction solution was cooled to room temperature, diluted with water, and extracted with ethyl acetate (20 mL $\times$ 3). The extracts were combined, washed with a saturated saline solution, dried over anhydrous sodium sulfate, and then the solvent was removed. The obtained residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate = 1:1 to 1:2) to obtain

the title compound (300 mg, 11%).

¹H NMR (400 MHz, CDCl₃): δ 8.83 (s, 1H), 8.62 (d, J = 3.6 Hz, 1H), 8.46 (s, 1H), 8.05 (d, J = 3.2 Hz, 1H), 7.75 (d, J = 4.4 Hz, 1H), 7.29-7.27 (m, 1H), 5.16-5.12 (m, 1H), 4.69-4.67 (m, 1H), 3.76 (s, 3H), 3.72 (s, 3H), 3.23-3.20 (m, 2H), 1.45 (s, 9H).

[0239]

(Step 4) Methyl 3-[6-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)pyridin-3-yl]-L-alaninate (M-30)




4 N Hydrochloric acid/ethyl acetate (5.0 mL) was added to methyl N-(tert-butoxycarbonyl)-3-[6-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)pyridin-3-yl]-L-alaninate (300 mg, 0.659 mmol), and the resulting mixture was stirred at room temperature for 1 hour. The obtained white solid was filtered and dried under reduced pressure to obtain a hydrochloride of the title compound (M-30) (253 mg, 98%).

¹H NMR (400 MHz, CD₃OD): δ 9.29 (s, 1H), 8.77 (d, J = 6.0 Hz, 1H), 8.62 (d, J = 2.0 Hz, 1H), 8.57 (d, J = 5.6 Hz, 1H), 8.09 (dd, J = 8.0, 2.0 Hz, 1H), 7.64 (dd, J = 15.6, 8.0 Hz, 1H), 4.55-4.51 (m, 1H), 3.88 (s, 3H), 3.76 (s, 3H), 3.52-3.48 (m, 1H), 3.46-3.41 (m, 1H); MS (ESI) m/z 356 (M+H)⁺.

[0240]

Example 85: (2S)-2-[[2,5-difluoro-4-[[4-(2-isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)phenyl]propanoic acid (A-24)




The title compound was obtained by a method similar to step 1 of Example 27 by using methyl 4-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)-L-phenylalaninate (M-8) and 2,5-difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoic acid (M-19).

¹H NMR (400 MHz, DMSO-d₆) δ 10.97 (s, 1H), 9.16 (s, 2H), 8.96 (s, 1H), 8.60 (dd, J = 8.0, 2.6 Hz, 1H), 8.55 (d, J = 4.9 Hz, 1H), 8.09 - 8.01 (m, 2H), 8.00 - 7.94 (m, 2H), 7.87 (d, J = 4.9 Hz, 1H), 7.36 (d, J = 8.3 Hz, 2H), 7.34 - 7.23 (m, 2H), 7.21 (d, J = 8.3 Hz, 1H), 4.68 - 4.58 (m, 4H), 3.77 (sep, J = 6.1 Hz, 1H), 3.58 (s, 3H), 3.25 - 3.18 (m, 1H), 3.12 - 3.01 (m, 1H), 1.15 (d, J = 6.1 Hz, 6H); MS (ESI) m/z 786 (M+H)⁺.

[0241]

Example 86: (2S)-2-[[[2,5-Difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxypyrimidin-5-yl)phenyl]propanoic acid (A-25)



The title compound was obtained by a method similar to step 1 of Example 27 by using methyl (2S)-2-amino-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoate (M-2) and


5 2,5-difluoro-4-[[4-(2-tetrahydropyran-4-ylpyrimidin-5-yl)phenyl]sulfonylamino]benzoic acid (M-11).

¹H NMR (400 MHz, DMSO-d₆) δ 10.95 (s, 1H), 9.14 (s, 2H), 8.54 (dd, J = 8.0, 2.6 Hz, 1H), 8.13 - 7.99 (m, 2H), 7.99 - 7.81 (m, 2H), 7.40 - 7.16 (m, 4H), 7.16 - 6.65 (m, 2H), 4.85 - 4.47 (m, 1H), 4.22 - 3.89 (m, 2H), 3.89 - 3.54 (m, 2H), 3.38
10 (s, 3H), 3.29 - 3.08 (m, 5H), 3.00 (dd, J = 13.9, 9.9 Hz, 1H), 2.05 (s, 3H), 2.00 - 1.64 (m, 4H) ; MS (ESI) m/z 775 (M+H)⁺.

[0242]

Example 87:

(2S)-2-[[2,5-Difluoro-4-[(4-tetrahydropyran-4-ylphenyl)sulfonylamino]benzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxypyrimidin-5-yl)phenyl]propanoic acid
15 (A-26)




The title compound was obtained by a method similar to step 1 of Example 27 by using methyl (2S)-2-amino-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoate (M-2) and
20 2,5-difluoro-4-[(4-tetrahydropyran-4-ylphenyl)sulfonylamino]benzoic acid (M-27).

¹H NMR (400 MHz, DMSO-d₆) δ 10.72 (s, 1H), 8.45 (dd, J = 8.0, 2.7 Hz, 1H),
25 7.70 (d, J = 8.4 Hz, 2H), 7.42 (d, J = 8.4 Hz, 2H), 7.30 - 7.17 (m, 4H), 7.17 -

[0243]

5	Example (2S)-2-[[2,5-Difluoro-4-[[4-(2-isopropylpyrimidin-5-yl)phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxypyrimidin-5-yl)phenyl]propanoic acid (A-27)	88:
---	---	-----




The title compound was obtained by a method similar to step 1 of Example 27 by using methyl (2S)-2-amino-3-[4-(1,3,4-trimethyl-2,6-dioxo-pyrimidin-5-yl)phenyl]propanoate (M-2) and 2,5-difluoro-4-{[4-(2-isopropylpyrimidin-5-yl)phenyl]sulfonylamino}Benzoic acid (M-28).

¹H NMR (400 MHz, DMSO-d₆) δ 10.95 (s, 1H), 9.11 (s, 2H), 8.53 (dd, J = 8.0, 2.6 Hz, 1H), 8.02 (d, J = 8.6 Hz, 2H), 7.99 - 7.84 (m, 2H), 7.40 - 7.16 (m, 4H), 7.16 - 6.94 (m, 2H), 4.76 - 4.35 (m, 1H), 3.37 (s, 3H), 3.28 - 3.10 (m, 5H), 3.00 (dd, J = 13.9, 9.9 Hz, 1H), 2.05 (s, 3H), 1.31 (d, J = 6.9 Hz, 6H); MS (ESI) m/z 733 (M+H)⁺.

[0244]

Example 89:
(2S)-2-[[[2,5-Difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[5-(1,3,4-trimethyl-2,6-dioxypyrimidin-5-yl)-2-pyridyl]propanoic acid (A-28)




The title compound was obtained by a method similar to step 1 of Example 27 by using methyl (2S)-2-amino-3-[5-(1,3,4-trimethyl-2,6-dioxypyrimidin-5-yl)-2-pyridyl]propanoate (M-29) and 2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonamino]benzoic acid (M-11).

¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.13 (s, 2H), 8.72 (dd, J = 7.9, 3.6 Hz, 1H), 8.41 (d, J = 2.2 Hz, 1H), 8.11 - 8.00 (m, 2H), 8.00 - 7.89 (m, 2H), 7.73 (d, J = 7.9 Hz, 1H), 7.47 (d, J = 8.0 Hz, 1H), 7.34 (dd, J = 10.3, 6.4 Hz, 1H), 7.27 (dd, J = 11.4, 6.3 Hz, 1H), 5.04 - 4.70 (m, 1H), 4.13 - 3.89 (m, 2H), 3.46 (s, 2H), 3.41 (s, 3H), 3.35 - 3.24 (m, 1H), 3.22 (s, 3H), 3.18 - 3.02 (m, 2H), 2.11 (s, 3H), 1.98 - 1.60 (m, 4H) ; MS (ESI) m/z 776 (M+H)⁺.

[0245]

Reference Example 1:
(2S)-2-[[2,5-Difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoic acid (A-29)




The title compound was obtained by a method similar to step 1 of

Example 27 by using methyl
 3-[6-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)pyridin-
 3-yl]-L-alaninate (M-30) and
 2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylami
 5 no]benzoic acid (M-11).
¹H NMR (400 MHz, DMSO-d₆): δ 10.95 (s, 1H), 9.13 (s, 2H), 8.99 (s, 1H), 8.69
 (dd, J = 8.2, 2.2 Hz, 1H), 8.56 (d, J = 5.0 Hz, 1H), 8.46 (d, J = 2.3 Hz, 1H), 8.06
 - 7.99 (m, 2H), 7.99 - 7.92 (m, 2H), 7.92 - 7.85 (m, 2H), 7.43 - 7.36 (m, 1H),
 7.33 - 7.22 (m, 2H), 4.74 - 4.63 (m, 1H), 4.00 - 3.90 (m, 2H), 3.59 (s, 3H), 3.54 -
 10 3.42 (m, 2H), 3.29 (dd, J = 14.1, 4.6 Hz, 1H), 3.19 - 3.03 (m, 2H), 1.95 - 1.74
 (m, 4H); MS(ESI) m/z 799.51(M+H)⁺.

[0246]

Reference	Example	2:
	(2S)-2-[[2,5-Difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl	
15	aminol]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)p	
	henyl]propanoic acid (A-30)	



The title compound was obtained by a method similar to step 1 of


Example	27	by	using	methyl
20	4-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)-L-phenyla			
	laninate	(M-8)		and
	2,5-difluoro-4-[[4-[2-(isopropoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino			
	]benzoic acid (M-19).			

¹H NMR (400 MHz, DMSO-d₆): δ 10.97 (s, 1H), 9.18 (s, 2H), 8.96 (s, 1H), 8.60
 25 (dd, J = 8.0, 2.6 Hz, 1H), 8.55 (d, J = 5.0 Hz, 1H), 8.10 - 8.02 (m, 2H), 8.02 -

7.94 (m, 2H), 7.88 (d, $J = 5.0$ Hz, 1H), 7.40 - 7.24 (m, 4H), 7.24 - 7.17 (m, 2H), 4.63 (s, 3H), 3.59 (s, 3H), 3.40 (s, 3H), 3.23 (dd, $J = 14.0, 4.5$ Hz, 1H), 3.06 (dd, $J = 14.0, 9.9$ Hz, 1H); MS(ESI) m/z 758.54(M+H)⁺.

[0247]

5 Example 90: Isopropyl
(2S)-2-[[4-[[4-(1-allyl-2-oxopyrimidin-5-yl)phenyl]sulfonylamino]-2,5-difluoro-
-benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyri-
dyl]propanoate (C-32)



10 Isopropanol alcohol (4 mL) and 4 N hydrochloric acid/dioxane (2 mL)
were added to A-5 (80 mg, 0.10 mmol), and the resulting mixture was stirred
at 60°C for 2 hours. The reaction liquid was concentrated under reduced
pressure, the obtained residue was purified by reverse phase HPLC (an
H₂O/CH₃CN system including 0.1% TFA) to obtain a TFA salt of the title
15 compound (53 mg, 57%).

¹H NMR (400 MHz, DMSO-d₆) δ 10.90 (s, 1H), 9.04 (d, $J = 3.4$ Hz, 1H), 9.00
(s, 1H), 8.85 (dd, $J = 7.6, 3.3$ Hz, 1H), 8.74 (d, $J = 3.4$ Hz, 1H), 8.58 (d, $J = 4.9$
Hz, 1H), 8.46 (dd, $J = 2.4, 0.7$ Hz, 1H), 7.93 - 7.84 (m, 5H), 7.74 (dd, $J = 8.2,$
2.5 Hz, 1H), 7.47 (d, $J = 8.2$ Hz, 1H), 7.38 (dd, $J = 10.3, 6.3$ Hz, 1H), 7.26 (dd,
20 $J = 11.3, 6.3$ Hz, 1H), 6.07 - 5.95 (m, 1H), 5.26 - 5.17 (m, 2H), 4.94 - 4.79 (m,
2H), 4.54 (dt, $J = 5.9, 1.4$ Hz, 2H), 3.61 (s, 3H), 3.36 - 3.23 (m, 2H), 1.15 (d, $J =$
6.2 Hz, 3H), 1.10 (d, $J = 6.2$ Hz, 3H); MS (ESI) m/z 813 (M+H)⁺.


[0248]

A method similar to Example 90 was performed by using A-5, A-20,

A-29, or A-30 as carboxylic acid and methanol, ethanol, isopropyl alcohol, 3-pentanol, 2-methoxyethanol, or N-(2-hydroxy)morpholine as alcohol to synthesize C-33 to C-42.

[0249]


5 Example 91: 1-Ethylpropyl
(2S)-2-[[4-[[4-(1-allyl-2-oxopyrimidin-5-yl)phenyl]sulfonylamino]-2,5-difluoro-
-benzoyl]amino]-3-[5-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-2-pyri-
dyl]propanoate (C-33)



10 ¹H NMR (400 MHz, DMSO-d₆) δ 10.90 (s, 1H), 9.04 (d, J = 3.4 Hz, 1H), 9.00
(s, 1H), 8.88 (dd, J = 7.6, 3.3 Hz, 1H), 8.73 (d, J = 3.4 Hz, 1H), 8.57 (d, J = 4.9
Hz, 1H), 8.46 (d, J = 2.4 Hz, 1H), 7.94 - 7.83 (m, 5H), 7.74 (dd, J = 8.2, 2.5 Hz,
1H), 7.47 (d, J = 8.2 Hz, 1H), 7.35 (dd, J = 10.3, 6.3 Hz, 1H), 7.26 (dd, J = 11.3,
6.2 Hz, 1H), 6.08 - 5.93 (m, 1H), 5.27 - 5.17 (m, 2H), 5.02 - 4.91 (m, 1H), 4.72 -
15 4.61 (m, 1H), 4.57 - 4.50 (m, 2H), 3.61 (s, 3H), 3.41 - 3.24 (m, 2H), 1.63 - 1.32
(m, 4H), 0.86 - 0.68 (m, 6H); MS (ESI) m/z 841 (M+H)⁺.

[0250]


 Example 92: Methyl
(2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulf
20 onylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-
-yl)-3-pyridyl]propanoate (C-34)



^1H NMR (400 MHz, DMSO- d_6) δ 10.99 (s, 1H), 9.14 (s, 2H), 8.99 (s, 1H), 8.86 (dd, J = 8.0, 2.0 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.47 (d, J = 2.0 Hz, 1H), 8.07 - 8.00 (m, 2H), 8.00 - 7.94 (m, 2H), 7.94 - 7.85 (m, 2H), 7.41 (d, J = 8.0 Hz, 1H),
 5 7.37 - 7.23 (m, 2H), 4.87 - 4.67 (m, 1H), 4.05 - 3.88 (m, 2H), 3.67 (s, 3H), 3.61 (s, 3H), 3.54 - 3.43 (m, 2H), 3.33 - 3.22 (m, 1H), 3.22 - 3.03 (m, 2H), 2.02 - 1.72 (m, 4H); MS(ESI) m/z 813.47 ($M+H$) $^+$.

[0251]

	Example	93:	Ethyl
10	(2S)-2-[[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-35)		




^1H NMR (400 MHz, DMSO- d_6) δ 10.98 (s, 1H), 9.14 (s, 2H), 8.99 (s, 1H), 8.86 (dd, J = 7.7, 1.7 Hz, 1H), 8.57 (d, J = 4.9 Hz, 1H), 8.47 (d, J = 2.3 Hz, 1H), 8.06 - 8.00 (m, 2H), 7.99 - 7.94 (m, 2H), 7.94 - 7.87 (m, 2H), 7.41 (d, J = 8.1 Hz, 1H),
 15 7.34 - 7.24 (m, 2H), 4.78 - 4.63 (m, 1H), 4.11 (q, J = 7.1 Hz, 2H), 4.00 - 3.90 (m, 2H), 3.59 (s, 3H), 3.49 - 3.41 (m, 2H), 3.25 (dd, J = 14.1, 5.6 Hz, 1H), 3.18 -

3.07 (m, 2H), 1.98 - 1.70 (m, 4H), 1.16 (t, $J = 7.1$ Hz, 3H); MS (ESI) m/z 827 (M+H)⁺.

[0252]

	Example	94:	Isopropyl
5	(2S)-2-[[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-36)		




¹H NMR (400 MHz, DMSO-d₆) δ 10.97 (s, 1H), 9.14 (s, 2H), 8.99 (s, 1H), 8.87 - 8.79 (m, 1H), 8.56 (d, $J = 5.0$ Hz, 1H), 8.46 (d, $J = 2.3$ Hz, 1H), 8.03 (d, $J = 8.6$ Hz, 2H), 7.97 (d, $J = 8.6$ Hz, 2H), 7.94 - 7.85 (m, 2H), 7.40 (d, $J = 8.1$ Hz, 1H), 7.28 (ddd, $J = 11.1, 6.3, 4.5$ Hz, 2H), 4.91 (sep, $J = 6.2$ Hz, 1H), 4.72 - 4.61 (m, 1H), 4.02 - 3.90 (m, 2H), 3.59 (s, 3H), 3.52 - 3.39 (m, 2H), 3.28 - 3.03 (m, 3H), 1.98 - 1.71 (m, 4H), 1.18 (d, $J = 6.2$ Hz, 3H), 1.13 (d, $J = 6.2$ Hz, 3H);

15 MS (ESI) m/z 841 (M+H)⁺.

[0253]


	Example	95:	1-Ethylpropyl
20	(2S)-2-[[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-37)		



¹H NMR (400 MHz, DMSO-d₆) δ 10.97 (s, 1H), 9.14 (s, 2H), 8.99 (s, 1H), 8.86 (d, J = 7.8 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.49 (s, 1H), 8.03 (d, J = 8.1 Hz, 2H), 7.99 - 7.91 (m, 3H), 7.89 (d, J = 4.9 Hz, 1H), 7.40 (d, J = 8.0 Hz, 1H), 7.32 - 7.20 (m, 2H), 4.82 - 4.62 (m, 2H), 4.01 - 3.89 (m, 2H), 3.60 (s, 3H), 3.53 - 3.42 (m, 2H), 3.33 - 3.22 (m, 1H), 3.19 - 3.02 (m, 2H), 1.98 - 1.74 (m, 4H), 1.63 - 1.40 (m, 4H), 0.90 - 0.73 (m, 6H); MS (ESI) m/z 869 (M+H)⁺.

[0254]

Example	96:	2-Methoxyethyl
10	(2S)-2-[[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-38)	




¹H NMR (400 MHz, DMSO-d₆) δ 10.97 (s, 1H), 9.14 (s, 2H), 8.99 (s, 1H), 8.87 (dd, J = 7.9, 1.7 Hz, 1H), 8.56 (d, J = 5.0 Hz, 1H), 8.47 (d, J = 2.4 Hz, 1H), 8.06 - 7.99 (m, 2H), 7.98 - 7.85 (m, 4H), 7.40 (d, J = 8.1 Hz, 1H), 7.33 - 7.22 (m, 2H), 4.81 - 4.69 (m, 1H), 4.24 - 4.15 (m, 2H), 4.00 - 3.90 (m, 2H), 3.59 (s, 3H), 3.54 - 3.42 (m, 4H), 3.30 - 3.25 (m, 1H), 3.24 (s, 3H), 3.19 - 3.04 (m, 2H), 1.94 - 1.74

(m, 4H); MS (ESI) m/z 857 (M+H)⁺.

[0255]


Example 97: 2-Morpholinoethyl
(2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-ylpyrimidin-5-yl)phenyl]sulfonylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate (C-39)



¹H NMR (400 MHz, DMSO-d₆) δ 11.00 (s, 1H), 9.14 (s, 2H), 9.03 - 8.93 (m, 2H), 8.57 (d, J = 4.9 Hz, 1H), 8.49 (d, J = 2.3 Hz, 1H), 8.07 - 7.85 (m, 6H), 7.46 - 7.40 (m, 1H), 7.39 - 7.21 (m, 2H), 4.91 - 4.78 (m, 1H), 4.53 - 4.35 (m, 2H), 4.00 - 3.64 (m, 8H), 3.59 (s, 3H), 3.54 - 3.35 (m, 4H), 3.24 - 3.02 (m, 5H), 1.96 - 1.72 (m, 4H); MS (ESI) m/z 912 (M+H)⁺.

[0256]

Example 98: Isopropyl
(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)phenyl]propanoate (C-40)




¹H NMR (400 MHz, DMSO-d₆) δ 10.97 (s, 1H), 9.19 (s, 2H), 8.97 (s, 1H), 8.76

(dd, $J = 7.3, 2.0$ Hz, 1H), 8.55 (d, $J = 5.0$ Hz, 1H), 8.16 - 8.02 (m, 2H), 8.02 - 7.93 (m, 2H), 7.88 (dd, $J = 5.0, 0.8$ Hz, 1H), 7.42 - 7.35 (m, 2H), 7.35 - 7.24 (m, 2H), 7.24 - 7.12 (m, 2H), 5.02 - 4.81 (m, 1H), 4.70 - 4.51 (m, 3H), 3.59 (s, 3H), 3.40 (s, 3H), 3.26 - 3.00 (m, 2H), 1.18 (d, $J = 6.2$ Hz, 3H), 1.12 (d, $J = 6.2$ Hz, 3H); MS (ESI) m/z 800 ($M+H$)⁺.

[0257]


Example 99: 1-Ethylpropyl (2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)phenyl]propanoate (C-41)



¹H NMR (400 MHz, DMSO-d₆) δ 10.98 (s, 1H), 9.18 (s, 2H), 8.96 (s, 1H), 8.79 (dd, $J = 7.7, 1.9$ Hz, 1H), 8.55 (d, $J = 5.0$ Hz, 1H), 8.13 - 8.02 (m, 2H), 8.02 - 7.93 (m, 2H), 7.88 (d, $J = 5.0$ Hz, 1H), 7.48 - 7.35 (m, 2H), 7.33 - 7.24 (m, 2H), 7.25 - 7.16 (m, 2H), 4.77 - 4.59 (m, 4H), 3.59 (s, 3H), 3.47 - 3.37 (m, 3H), 3.27 - 3.16 (m, 1H), 3.16 - 3.03 (m, 1H), 1.64 - 1.37 (m, 4H), 0.90 - 0.73 (m, 6H); MS(ESI) m/z 828.47 ($M+H$)⁺.

[0258]

Example 100: Methyl (2S)-2-[[[4-[[4-(2-ethylpyrimidin-5-yl)phenyl]sulfonylamino]-2,3,5,6-tetrafluorobenzoyl]amino]-3-[4-(1,3,4-trimethyl-2,6-dioxypyrimidin-5-yl)phenyl]propanoate (C-42)




¹H NMR (400 MHz, DMSO-d₆) δ 10.85 (s, 1H), 9.46 (d, J = 7.9 Hz, 1H), 9.14 (s, 2H), 8.05 (d, J = 8.5 Hz, 2H), 7.90 (d, J = 8.4 Hz, 2H), 7.31 - 7.21 (m, 2H), 7.14 - 6.91 (m, 2H), 5.03 - 4.58 (m, 1H), 3.50 - 3.26 (m, 4H), 3.26 - 3.11 (m, 6H),
 5 3.11 - 2.82 (m, 3H), 2.20 - 1.87 (m, 3H), 1.32 (t, J = 7.6 Hz, 3H) ; MS (ESI) m/z 769 (M+H)⁺.

[0259]

Example

101:

2,5-Difluoro-N-[(1S)-2-(hydroxyamino)-1-[[6-(1-methyl-2,4-dioxo-pyrido[3,4-d
 10]pyrimidin-3-yl)-3-pyridyl]methyl]-2-oxo-ethyl]-4-[[4-[2-(methoxymethyl)pyri
 midin-5-yl]phenyl]sulfonylamino]benzamide (C-43)



(2S)-2-[[[2,5-Difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]
 sulfonylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimid
 15 in-3-yl)-3-pyridyl]propanoic acid (A-23) (50.2 mg, 0.0662 mmol),
 1-hydroxy-7-azabenzotriazole (10.8 mg, 0.0794 mmol), and hydroxyamine
 hydrochloride (13.8 mg, 0.199 mmol) were dissolved in acetonitrile (2 mL),
 diisopropylethylamine (45.0 μL, 0.265 mmol) and
 O-(7-azabenzotriazol-1-yl)-N,N,N',N',-tetramethyluronium
 20 hexafluorophosphate (30.2 mg, 0.0794 mmol) were added thereto, and the

resulting mixture was stirred at room temperature overnight. The reaction liquid was concentrated under reduced pressure, the obtained residue was purified by reverse phase HPLC (an H₂O/CH₃CN system including 0.1% TFA) to obtain a TFA salt of the title compound (26.5 mg, 45%).

5 ¹H NMR (400 MHz, DMSO-d₆) δ 10.94 (s, 1H), 10.86 (s, 1H), 9.18 (s, 2H), 8.99 (s, 1H), 8.70 - 8.61 (m, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.54 - 8.41 (m, 1H), 8.09 - 8.02 (m, 2H), 8.00 - 7.94 (m, 2H), 7.92 - 7.83 (m, 2H), 7.40 (d, J = 8.1 Hz, 1H), 7.26 (ddd, J = 13.0, 10.6, 6.3 Hz, 2H), 4.67 - 4.58 (m, 3H), 3.59 (s, 3H), 3.40 (s, 3H), 3.11 (dd, J = 14.0, 5.0 Hz, 1H), 3.02 (dd, J = 14.0, 9.8 Hz, 1H);
10 MS (ESI) m/z 774 (M+H).

[0260]


A method similar to Example 101 was performed by using A-29 as carboxylic acid and hydroxyamine hydrochloride or methoxyamine hydrochloride as amine to synthesize C-44 and C-45.

15 [0261]

Example

102:

2,5-Difluoro-N-[(1S)-2-(hydroxyamino)-1-[[6-(1-methyl-2,4-dioxypyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]methyl]-2-oxoethyl]-4-[[4-(2-tetrahydropyran-4-yl)pyrimidin-5-yl]phenyl]sulfonylamino]benzamide (C-44)



20

MS(ESI) m/z 814.63 (M+H)⁺.


[0262]

Example

103:

2,5-Difluoro-N-[(1S)-2-(methoxyamino)-1-[[6-(1-methyl-2,4-dioxypyrido[3,4-d]


[pyrimidin-3-yl]-3-pyridyl)methyl]-2-oxoethyl]-4-[[4-(2-tetrahydropyran-4-yl
pyrimidin-5-yl)phenyl]sulfonylamino]benzamide (C-45)



¹H NMR (400 MHz, DMSO-d₆) δ 11.43 (s, 1H), 10.96 (s, 1H), 9.14 (s, 2H),
5 8.99 (s, 1H), 8.70 (d, J = 8.0 Hz, 1H), 8.57 (d, J = 5.0 Hz, 1H), 8.45 (d, J = 2.4
Hz, 1H), 8.04 (d, J = 8.3 Hz, 2H), 7.97 (d, J = 8.3 Hz, 2H), 7.93 - 7.82 (m, 2H),
7.41 (d, J = 8.0 Hz, 1H), 7.38 - 7.16 (m, 2H), 4.67 - 4.47 (m, 1H), 4.05 - 3.87 (m,
2H), 3.60 (s, 3H), 3.51 (s, 3H), 3.50 - 3.42 (m, 2H), 3.23 - 2.98 (m, 3H), 2.03 -
1.69 (m, 4H); MS(ESI) m/z 828.71 (M+H)⁺.

10 [0263]

Example 104: Isopropyl
(2S)-2-[[4-[[4-(4-ethyltriazol-1-yl)phenyl]sulfonylamino]-2,5-difluorobenzoyl]
amino]-3-[4-(1-methyl-2,4-dioxopyrido[3,4-d]pyrimidin-3-yl)phenyl]propanoa
te (C-46)



15

A method similar to step 1 of Example 27 was performed by using methyl
4-(1-methyl-2,4-dioxo-1,4-dihydropyrido[3,4-d]pyrimidin-3(2H)-yl)-L-phenyla
laninate (M-8) and

4-[[4-(4-ethyltriazol-1-yl)phenyl]sulfonylamino]-2,5-difluorobenzoic acid (M-26) to obtain carboxylic acid, and then an isopropyl ester of the obtained product was produced by a method similar to Example 90 to obtain the title compound.

5 ¹H NMR (400 MHz, DMSO-d₆) δ 10.93 (s, 1H), 8.97 (s, 1H), 8.76 (dd, J = 7.4, 1.8 Hz, 1H), 8.69 (s, 1H), 8.55 (d, J = 5.0 Hz, 1H), 8.23 - 8.09 (m, 2H), 8.09 - 7.93 (m, 2H), 7.88 (d, J = 5.0 Hz, 1H), 7.36 (d, J = 8.1 Hz, 2H), 7.34 - 7.23 (m, 2H), 7.21 (d, J = 8.2 Hz, 2H), 5.06 - 4.72 (m, 1H), 4.72 - 4.53 (m, 1H), 3.59 (s, 3H), 3.26 - 3.00 (m, 2H), 2.72 (q, J = 7.6 Hz, 2H), 1.25 (t, J = 7.5 Hz, 3H), 1.18
10 (d, J = 6.2 Hz, 3H), 1.12 (d, J = 6.2 Hz, 3H) ; MS (ESI) m/z 773 (M+H)⁺.

[0264]

(1) Evaluation test for VCAM-1/α4β1 integrin binding inhibitory activity

The ability of test substances to inhibit the binding of Jurkat cell strain, a human T-cell line known to express α4β1 integrin, to VCAM-1 was
15 examined.

A 96-well microtiter plate was coated overnight at 4°C with a recombinant human VCAM-1/Fc (R&D systems) solution diluted with a buffer A (carbonate buffer, pH 9.6). After washing once with PBS, 150 μL/well of Block Ace (manufactured by Snow Brand Milk Products Co., Ltd.)
20 was added, and it was incubated at room temperature for 2 hours. After removal, it was washed once with PBS.

100 μL of each of test substances and Jurkat cells diluted with a binding buffer solution (DMEM containing 40 mM HEPES, 0.2% BSA, and 4 mM MnCl₂) to various concentrations was added to the plate coated with the
25 VCAM-1/Fc, and it was incubated at 30°C for 15 minutes to 60 minutes. After the cells were bound to the wells, unbound cells were removed by washing with PBS. 50 μL/well of a buffer solution C (PBS containing 1.5% Triton X-100) was added to the plate to dissolve the bound Jurkat cells. 30 μL of substrate buffer (Promega, CytoTox 96 Non-Radioactive Cytotoxicity
30 Assay) was added to 30 μL of a cell lysate to allow a reaction for 30 minutes

at room temperature in a dark place. 30 μ L of stop solution (Promega, CytoTox 96 Non-Radioactive Cytotoxicity Assay) was added thereto to determine the absorbance at 490 nm using a plate reader. The absorbance was obtained herein by detecting the activity of lactate dehydrogenase (LDH) eluted in a supernatant of each well, that is, the absorbance is proportional to the number of Jurkat cells remaining on the plate by being bound to VCAM-1. The test was duplicated, and the cell binding rate at various concentrations with the absorbance of the well containing no test substance being taken as 100% was determined, to calculate the concentration IC_{50} at which 50% of binding was inhibited. As test compounds, carboxylic acid compounds (compounds A-1 to A-28) among the compounds synthesized in concerned Examples were used. The same applies to the following description.

[0265]

(2) Evaluation test for MAdCAM-1/ $\alpha 4\beta 7$ integrin binding inhibitory activity

The ability of test substances to inhibit the binding of RPMI-8866 cell strain, a human B-cell line known to express $\alpha 4\beta 7$ integrin, to MAdCAM-1 was examined.

A 96-well microtiter plate was coated overnight at 4°C with a recombinant mouse MAdCAM-1/Fc (R&D systems) solution diluted with the buffer A (carbonate buffer, pH 9.6). After washing once with PBS, 150 μ L/well of Block Ace (manufactured by Snow Brand Milk Products Co., Ltd.) was added, and it was incubated at room temperature for 2 hours. After removal, it was washed once with PBS.

100 μ L of each of test substances and RPMI-8866 cells diluted with a binding buffer solution (DMEM containing 40 mM HEPES, 0.2% BSA, and 4 mM $MnCl_2$) to various concentrations was added to the plate coated with MAdCAM-1/Fc, and it was incubated at 30°C for 15 minutes to 60 minutes. After the cells were bound to the wells, unbound cells were removed by washing with PBS. 50 μ L/well of the buffer solution C (PBS containing

1.5% Triton X-100) was added to the plate to dissolve the bound RPMI-8866 cells. 30 μ L of substrate buffer (Promega, CytoTox 96 Non-Radioactive Cytotoxicity Assay) was added to 30 μ L of a cell lysate to allow a reaction for 30 minutes at room temperature in a dark place. 30 μ L of stop solution
 5 (Promega, CytoTox 96 Non-Radioactive Cytotoxicity Assay) was added thereto to determine the absorbance at 490 nm using a plate reader. The absorbance was obtained herein by detecting the activity of lactate dehydrogenase (LDH) eluted in a supernatant of each well, that is, the absorbance is proportional to the number of RPMI-8866 cells remaining on
 10 the plate by being bound to MAdCAM-1. The test was duplicated, and the cell binding rate at various concentrations with the absorbance of the well containing no test substance being taken as 100% was determined, to calculate the concentration IC₅₀ at which 50% of binding was inhibited.

The selectivity was determined by dividing the IC₅₀ value in the
 15 evaluation test for VCAM-1/ α 4 β 1 integrin binding inhibitory activity by the IC₅₀ value in the evaluation test for MAdCAM-1/ α 4 β 7 integrin binding inhibitory activity.

[0266]

(3) Evaluation test (1) for MAdCAM-1/ α 4 β 7 integrin binding inhibitory
 20 activity in the presence of serum

The ability of test substances to inhibit the binding of RPMI-8866 cell line, a human B-cell line known to express α 4 β 7 integrin, to MAdCAM-1 was examined.

A 96-well microtiter plate was coated overnight at 4°C with a
 25 recombinant mouse MAdCAM-1/Fc (R&D systems) solution diluted with the buffer A (carbonate buffer, pH 9.6). After washing once with PBS, 150 μ L/well of Block Ace (manufactured by Snow Brand Milk Products Co., Ltd.) was added, and it was incubated at room temperature for 2 hours. After removal, it was washed once with PBS.

30 100 μ L of each of test substances and RPMI-8866 cells diluted with a

binding buffer solution (DMEM containing 40 mM HEPES, 0.2% BSA, and 4 mM MnCl_2) to various concentrations was added to the plate coated with MAdCAM-1/Fc, so that human serum was contained at a final concentration of 50%, and it was incubated at 30°C for 15 minutes to 60 minutes. After the cells were bound to the wells, unbound cells were removed by washing with PBS. 50 μL /well of the buffer solution C (PBS containing 1.5% Triton X-100) was added to the plate to dissolve the bound RPMI-8866 cells. 30 μL of substrate buffer (Promega, CytoTox 96 Non-Radioactive Cytotoxicity Assay) was added to 30 μL of a cell lysate to allow a reaction for 30 minutes at room temperature in a dark place. 30 μL of stop solution (Promega, CytoTox 96 Non-Radioactive Cytotoxicity Assay) was added thereto to determine the absorbance at 490 nm using a plate reader. The absorbance was obtained herein by detecting the activity of lactate dehydrogenase (LDH) eluted in a supernatant of each well, that is, the absorbance is proportional to the number of RPMI-8866 cells remaining on the plate by being bound to MAdCAM-1. The test was duplicated, and the cell binding rate at various concentrations with the absorbance of the well containing no test substance being taken as 100% was determined, to calculate the concentration IC_{50} at which 50% of binding was inhibited.

[0267]

Table 1 and Table 2 show the results. It was confirmed that all the compounds described herein had selectivity, which was determined by dividing the IC_{50} value in the evaluation test for VCAM-1/ $\alpha 4\beta 1$ integrin binding inhibitory activity by the IC_{50} value in the evaluation test for MAdCAM-1/ $\alpha 4\beta 7$ integrin binding inhibitory activity, of 100 times or more and thus had preferable properties. The evaluation test for VCAM-1/ $\alpha 4\beta 1$ integrin binding inhibitory activity and the evaluation test for MAdCAM-1/ $\alpha 4\beta 7$ integrin binding inhibitory activity were conducted in accordance with the aforementioned methods described in the tests (1) and (2).

[0268]

Table 1

Example No.	$\alpha 4\beta 7$ Serum Activity rank when $\alpha 4\beta 7$ serum was added 1-10 nM: A, 10-50 nM: B, 50-200 nM: C, >200 nM: D
Example 26	B
Example 27	C
Example 28	B
Example 29	C
Example 30	B
Example 33	B
Example 34	B
Example 36	C
Example 37	B
Example 38	B
Example 39	B
Example 40	C
Example 41	D
Example 42	C
Example 43	D
Example 44	C
Example 45	B
Example 46	D
Example 47	B
Example 51	B

Table 2


Example No.	$\alpha 4\beta 7$ Serum Activity rank when $\alpha 4\beta 7$ serum was added 1-10 nM: A, 10-50 nM: B, 50-200 nM: C, >200 nM: D
Example 85	B
Example 86	B
Example 87	D
Example 88	C
Example 89	D

[0269]

As is obvious from the results of the selectivity determined by dividing the IC₅₀ value in the evaluation test for VCAM-1/ α 4 β 1 integrin binding inhibitory activity by the IC₅₀ value in the evaluation test for
5 MAdCAM-1/ α 4 β 7 integrin binding inhibitory activity, the compounds of the present invention, as compared with the compound of Patent Literature 1, has high selectivity with a low effect on α 4 β 1 and a high effect on α 4 β 7, and it can be seen from the results of Table 1 that, in particular, the
10 MAdCAM-1/ α 4 β 7 integrin binding inhibitory activity in the presence of serum is high. The high selectivity with a low effect on α 4 β 1 and a high effect on α 4 β 7 can reduce the action on α 4 β 1 that suppresses infiltration of lymphocytes circulating throughout the whole body, and can significantly suppress the action on α 4 β 7 that specifically acts on the intestinal tract, and thus has an advantage of being capable of treating adaptation disease
15 efficiently.

What is claimed is:


1. A sulfonamide derivative represented by the general formula (1) below or a pharmaceutically acceptable salt thereof:




(1)

wherein


A represents a group represented by the general formula (2-1), (2-2), (2-3) or (2-4) below:




(2 - 1)



(2 - 2)



(2 - 3)



(2 - 4)

wherein

Arm is a cyclic alkyl group or an aromatic ring containing 0, 1, 2, 3, or 4 heteroatoms selected from the group consisting of oxygen atoms, sulfur atoms and nitrogen atoms,

R₁, R₁₁, R₂₁, R₅₁, and R₅₂ each independently represent a hydrogen atom, a halogen atom, a hydroxyl group, a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a lower alkoxy group, a lower alkylthio group, a hydroxy-lower alkyl group, a hydroxy-lower alkenyl group, a hydroxy-lower alkoxy group, a halogeno-lower alkyl group, a halogeno-lower alkoxy group, a halogeno-lower alkylthio group, a halogeno-lower alkenyl group, a nitro group, a cyano group, an amino group, a lower alkylamino group, a lower alkylamino-lower alkyl group, a carboxyl group, a lower alkyloxycarbonyl group, a carbamoyl group, a lower alkanoyl group, an aroyl group, a lower alkylsulfonyl group, a sulfamoyl group, or an ammonium group,

R₁₂, R₁₃, and R₁₄ each independently represent a hydrogen atom, a halogen atom, a hydroxyl group, a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a lower alkoxy group, a lower alkylthio group, a hydroxy-lower alkyl group, a hydroxy-lower alkenyl group, a hydroxy-lower alkoxy group, a halogeno-lower alkyl group, a halogeno-lower alkoxy group, a halogeno-lower alkylthio group, a halogeno-lower alkenyl group, a nitro group, a cyano group, an amino group, a lower alkylamino group, a lower alkoxy-lower alkylene group, a lower alkylthio-lower alkylene group, a lower alkylamino-lower alkylene group, a lower alkylamino-lower alkyl group, a carboxyl group, a lower alkyloxycarbonyl group, a carbamoyl group, a lower alkanoyl group, an aroyl group, a lower alkylsulfonyl group, a sulfamoyl group, or an ammonium group, and

R₂, R₃, R₂₂, R₂₃, R₂₄, R₂₅, and R₅₃ each independently represent a hydrogen atom, a halogen atom, a hydroxyl group, a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a lower alkoxy group, a lower alkylthio group, a hydroxy-lower alkyl group, a hydroxy-lower alkenyl group, a hydroxy-lower alkoxy group, a halogeno-lower alkyl group, a halogeno-lower alkoxy group, a halogeno-lower alkylthio group, a halogeno-lower alkenyl group, a nitro group, a cyano group, an amino group, a lower alkylamino group, a lower alkylamino-lower alkyl group, a carboxyl group, a lower alkyloxycarbonyl group, a carbamoyl group, a lower alkanoyl group, an aroyl group, a lower alkylsulfonyl group, a sulfamoyl group, or an ammonium group,

R₆₀ represents an alkoxy group having 1 to 10 carbon atoms, a hydroxyl group, or a hydroxyamino group, these groups being optionally substituted with a substituent selected from the group consisting of an aryl group, a hydroxyl group, an alkyl group having 1 to 10 carbon atoms, a lower alkoxy group, a lower alkylamino group, a halogen atom, and a heterocyclic group,

R₄₁ represents a hydrogen atom or a lower alkyl group,

a, b, c, and d independently represent C-R₃₁, C-R₃₂, C-R₃₃, or C-R₃₄, respectively, wherein one or two of a, b, c, and d may represent a nitrogen atom,

R₃₁, R₃₂, R₃₃, and R₃₄ each independently represent a hydrogen atom, a halogen atom, a lower alkyl group, a lower alkoxy group, or a nitro group, wherein any one of R₃₁, R₃₂, R₃₃, and R₃₄ is a halogen atom or a lower alkyl group,

e, f, g, and h independently represent C-R₃₅, C-R₃₆, C-R₃₇, or C-R₃₈, respectively, wherein one or two of e, f, g, and h may represent a nitrogen atom,

R₃₅, R₃₆, R₃₇, and R₃₈ each independently represent a hydrogen atom, a halogen atom, a lower alkyl group, a lower alkoxy group, or a nitro group,

D represents a phenylene group or a heterocyclene group, optionally having a substituent selected from the group consisting of a halogen atom, a hydroxyl group, a lower alkyl group, a lower alkenyl group, a lower alkynyl group, a lower alkoxy group, a lower alkylthio group, a hydroxy-lower alkyl group, a hydroxy-lower alkenyl group, a hydroxy-lower alkoxy group, a halogeno-lower alkyl group, a halogeno-lower alkoxy group, a halogeno-lower alkylthio group, a halogeno-lower alkenyl group, a nitro group, a cyano group, an amino group, a carboxyl group, a lower alkyloxycarbonyl group, a carbamoyl group, a lower alkanoyl group, an aroyl group, a lower alkylsulfonyl group, a sulfamoyl group, a lower alkoxy-lower alkylene group, a lower alkylthio-lower alkylene group, a lower alkylamino-lower alkylene group, or an ammonium group,

when A represents a group represented by the general formula (2-1), (2-2), or (2-3), E represents:

a 5- or 6-membered heterocyclic group substituted with a substituent selected from the group consisting of: a 3- to 8-membered saturated or unsaturated ring group containing a heteroatom selected from the group consisting of an oxygen atom and a sulfur atom; a lower

alkoxy-lower alkylene group; a lower alkylthio-lower alkylene group; a lower alkylamino-lower alkylene group; a lower alkylamino group; and a lower alkenylamino group; or

a 5- or 6-membered cyclic ketone group containing 1 to 4 nitrogen atoms as atoms constituting the ring, the cyclic ketone group being substituted with a lower alkyl group or a lower alkenyl group, and

when A represents a group represented by the general formula (2-4), E represents:

(1) a phenyl group or a 5- or 6-membered heterocyclic group optionally having a substituent selected from the group consisting of (1-1) a halogen atom, (1-2) a hydroxyl group, (1-3) a lower alkyl group, (1-4) a lower alkenyl group, (1-5) a lower alkynyl group, (1-6) a lower alkoxy group, (1-7) a lower alkylthio group, (1-8) a hydroxy-lower alkyl group, (1-9) a hydroxy-lower alkenyl group, (1-10) a hydroxy-lower alkoxy group, (1-11) a halogeno-lower alkyl group, (1-12) a halogeno-lower alkoxy group, (1-13) a halogeno-lower alkylthio group, (1-14) a halogeno-lower alkenyl group, (1-15) a 3- to 8-membered saturated ring group containing a nitrogen atom connected by a carbon atom, (1-16) a 3- to 8-membered saturated or unsaturated ring group containing a heteroatom selected from the group consisting of an oxygen atom and a sulfur atom, (1-17) a lower alkoxy-lower alkylene group, (1-18) a lower alkylthio-lower alkylene group, (1-19) a lower alkylamino-lower alkylene group, (1-20) a lower alkylamino group, (1-21) a lower alkenylamino group, a nitro group, (1-22) a cyano group, (1-23) an amino group, (1-24) a 4- to 6-membered cyclic amino group, (1-25) a carboxyl group, (1-26) a lower alkyloxycarbonyl group, (1-27) a carbamoyl group, (1-28) a lower alkanoyl group, (1-29) an aroyl group, (1-30) a lower alkylsulfonyl group, (1-31) a sulfamoyl group, and (1-32) an ammonium group;

(2) an aminocarbonyl group optionally having a substituent selected from the group consisting of (2-1) a hydroxyl group, (2-2) a lower

alkyl group, (2-3) a lower alkenyl group, (2-4) a lower alkynyl group, (2-5) a lower alkoxy group, (2-6) a hydroxy-lower alkyl group, (2-7) a hydroxy-lower alkenyl group, (2-8) a hydroxy-lower alkoxy group, (2-9) a halogeno-lower alkyl group, (2-10) a halogeno-lower alkoxy group, (2-11) a halogeno-lower alkenyl group, (2-12) an amino group, (2-13) a lower alkylamino group, (2-14) an aryl group, (2-15) a heterocyclic group, (2-16) a heterocyclic lower alkyl group, (2-17) a lower alkylsulfonyl group, and (2-18) a sulfamoyl group;

- (3) a hydrogen atom;
- (4) a halogen atom;
- (5) a hydroxyl group;
- (6) a lower alkyl group;
- (7) a lower alkenyl group;
- (8) a lower alkynyl group;
- (9) a lower alkoxy group;
- (10) a lower alkylthio group;
- (11) a hydroxy-lower alkyl group;
- (12) a hydroxy-lower alkenyl group;
- (13) a hydroxy-lower alkoxy group;
- (14) a halogeno-lower alkyl group;
- (15) a halogeno-lower alkoxy group;
- (16) a halogeno-lower alkylthio group;
- (17) a halogeno-lower alkenyl group;
- (18) a nitro group;
- (19) a cyano group;
- (20) an amino group;
- (21) a carboxyl group;
- (22) a dihydroxyboryl group;
- (23) a lower alkylcarbonyl group;
- (24) a lower alkyloxycarbonyl group;
- (25) a carbamoyl group;

- (26) a lower alkanoyl group;
- (27) an aroyl group;
- (28) a lower alkylsulfonyl group;
- (29) a sulfamoyl group;
- (30) an ammonium group;
- (31) a lower alkylaminoalkylene group; or
- (32) a 5- or 6-membered cyclic ketone group containing 1 to 4


nitrogen atoms as atoms constituting the ring, the cyclic ketone group being substituted with a lower alkyl group or a lower alkenyl group,

wherein the (23) lower alkylcarbonyl group and the (24) lower alkyloxycarbonyl group may form a fused ring by being bonded to the phenylene group of D,

provided that the sulfonamide derivative excludes sulfonamide derivatives selected from the group consisting of (a) to (h) below:


(a)

(2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-ylpyrimidin-5-yl)phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)phenyl]propanoic acid




(b) cyclohexyl

(2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-ylpyrimidin-5-yl)phenyl]sulfonylamino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)phenyl]propionate




(c)

(2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-ylpyrimidin-5-yl)phenyl]sulfonylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoic acid



(d) cyclohexyl


(2S)-2-[[2,5-difluoro-4-[[4-(2-tetrahydropyran-4-ylpyrimidin-5-yl)phenyl]sulfonylamino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate



(e)


(2S)-2-[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl

amino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)phenyl]propanoic acid




(f) cyclohexyl

(2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[4-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)phenyl]propanoate



(g)


(2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoic acid



and

(h) cyclohexyl

(2S)-2-[[[2,5-difluoro-4-[[4-[2-(methoxymethyl)pyrimidin-5-yl]phenyl]sulfonyl amino]benzoyl]amino]-3-[6-(1-methyl-2,4-dioxo-pyrido[3,4-d]pyrimidin-3-yl)-3-pyridyl]propanoate



2. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to claim 1, wherein D is a phenylene group optionally having a substituent selected from the group consisting of a lower alkyl group, a halogen atom, a hydroxyl group, and a lower alkoxy group.
3. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to claim 1, wherein D is a 6-membered aromatic heterocyclene group optionally having a substituent selected from the group consisting of a lower alkyl group, a halogen atom, and a lower alkoxy group, and having a nitrogen atom as an atom constituting the ring.
4. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to claim 3, wherein the heterocyclene group of D is a pyridylene group optionally having a substituent selected from the group consisting of a lower alkyl group, a halogen atom, and a lower alkoxy group.
5. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 4, wherein the 5- or 6-membered heterocyclic group of E is selected from the group consisting of a pyridyl group, a pyridazyl group, a pyrimidyl group, a pyrazinyl group, a pyrrolyl

group, a triazolyl group, and a tetrazole group, and the cyclic ketone group of E is a 5- or 6-membered cyclic ketone group containing 2 or 3 nitrogen atoms as atoms constituting the ring.

6. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 4, wherein A represents a group represented by the general formula (2-4), and E is a 5- or 6-membered aromatic heterocyclic group containing 1, 2, 3, or 4 heteroatoms selected from the group consisting of oxygen atoms, sulfur atoms and nitrogen atoms and optionally having a substituent selected from the group consisting of a lower alkyl group, a lower alkoxy group, a lower alkoxy-lower alkylene group, and a halogen atom.

7. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to claim 6, wherein the aromatic heterocyclic group is selected from the group consisting of a pyridyl group, a pyridazyl group, a pyrimidyl group, a pyrazinyl group, a pyrrolyl group, a furanyl group, a thiophenyl group, a thiazolyl group, an oxazolyl group, a triazolyl group, and a tetrazole group.

8. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 4, wherein

A represents a group represented by the general formula (2-4), and

E is an aminocarbonyl group optionally substituted with a lower alkyl group, a heterocyclic group, or a heterocyclic lower alkyl group, or a lower alkylaminoalkylene group.

9. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 5, wherein the heterocyclic group of E is a pyridyl group, a pyrimidyl group, a triazolyl group, or a tetrazole

group optionally having a substituent selected from the group consisting of a 3- to 8-membered saturated or unsaturated ring group containing a heteroatom selected from an oxygen atom and a sulfur atom, a lower alkoxy-lower alkylene group, and a lower alkylamino-lower alkylene group.

10. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 5 and 9, wherein A represents a group represented by the general formula (2-2), and R₁₂, R₁₃, and R₁₄ are selected from the group consisting of a lower alkyl group, a lower alkoxy group, a lower alkoxy-lower alkylene group, a lower alkylamino-lower alkylene group, and a halogen atom.

11. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 10, wherein R₆₀ is a lower alkoxy group or hydroxyl group, optionally substituted with a substituent selected from the group consisting of a lower alkyl group, a lower alkoxy group, a lower alkylamino group, a halogen atom, and a heterocyclic group.

12. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 6, 8, 9, 10, and 11, wherein the lower alkyl group is a straight-chain, branched-chain, or cyclic alkyl group.

13. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 12, wherein R₁, R₁₁, R₂₁, R₅₁, and R₅₂ are lower alkyl groups.

14. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 13, wherein Arm is selected from the group consisting of a phenyl group, a pyridyl group, a pyrimidyl group, and an imidazolyl group.

15. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 4, wherein

A represents a group represented by the general formula (2-2), and

E represents a 5- or 6-membered heterocyclic group containing 1 to 4 nitrogen atoms, the 5- or 6-membered heterocyclic group being substituted with a substituent selected from the group consisting of a lower alkoxy-lower alkylene group, and a 5- or 6-membered heterocyclic group containing an oxygen atom; or a cyclic ketone group containing 1 to 4 nitrogen atoms, the cyclic ketone group being substituted with a lower alkyl group or a lower alkenyl group.

16. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to claim 15, wherein

R₆₀ is optionally substituted with a substituent selected from the group consisting of: an alkyl group having 1 to 10 carbon atoms, a lower alkoxy group, and a heterocyclic group containing 1 to 4 heteroatoms selected from the group consisting of oxygen atoms and nitrogen atoms, and

Arm is a pyridyl group.

17. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 4, wherein

A represents a group represented by the general formula (2-4),




E is a 5- or 6-membered heterocyclic group containing an oxygen atom, or a 5- or 6-membered heterocyclic group containing 1 to 4 nitrogen atoms, and





E optionally has a substituent selected from a group consisting of: a lower alkyl group; a lower alkoxy-lower alkylene group; and a 5- or 6-membered heterocyclic group containing an oxygen atom.





E represents: a 5- or 6-membered heterocyclic group substituted with a 3- to 8-membered saturated or unsaturated ring group containing a heteroatom selected from the group consisting of an oxygen atom and a sulfur atom; or a 5- or 6-membered cyclic ketone group containing 1 to 4 nitrogen atoms as atoms constituting the ring, the cyclic ketone group being substituted with a lower alkyl group or a lower alkenyl group.


20. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to claim 19, wherein two of R₃₁, R₃₂, R₃₃, and R₃₄ are halogen atoms and the others are hydrogen atoms.

O=C1C(=O)N(C1Cc2ccnc3cc(NC(=O)c4cc(F)c(S(=O)(=O)Nc5ccc6c(c4)nc7c6CCOCC7)c5)cc32)c8ccncc8N











,



,



.

22. A pharmaceutical composition comprising the sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 21 and a pharmaceutically acceptable carrier.


23. The sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 21 for use in the treatment or

prevention of an inflammatory disease in which an $\alpha 4\beta 7$ integrin-dependent adhesion process is involved in pathological conditions.

24. Use of the sulfonamide derivative or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 21 for inhibiting an $\alpha 4\beta 7$ integrin.

25. The composition of claim 22 for inhibiting an $\alpha 4\beta 7$ integrin.

26. The composition of claim 22 for use in the treatment or prevention of an inflammatory disease in which an $\alpha 4\beta 7$ integrin-dependent adhesion process is involved in pathological conditions.



(1)