

(10) International Publication Number
WO 2014/001057 A1(43) International Publication Date
3 January 2014 (03.01.2014)

(51) International Patent Classification:

A01P 1/00 (2006.01) *A01N 31/04* (2006.01)
A01N 31/08 (2006.01) *A61K 8/34* (2006.01)
A01N 31/06 (2006.01) *A61K 31/05* (2006.01)

(21) International Application Number:

PCT/EP2013/061705

(22) International Filing Date:

6 June 2013 (06.06.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12173845.4 27 June 2012 (27.06.2012) EP

(71) **Applicant** (for all designated States except AE, AG, AU, BB, BH, BN, BZ, CA, EG, GD, IL, IN, KN, LC, LK, MY, NG, NZ, OM, PG, QA, SC, SG, TT, US, VC, ZA): **UNILEVER N.V.** [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).

(71) **Applicant** (for all designated States except AM, AO, AZ, BF, BJ, BR, BY, CF, CG, CI, CL, CM, CN, CO, CR, CU, DM, DO, DZ, EC, GA, GE, GN, GQ, GT, GW, HN, ID, JP, KG, KM, KP, KR, KZ, LA, LY, MA, MD, ME, MG, ML, MN, MR, MX, NE, NI, PA, PE, PH, RU, SN, ST, SV, SY, TD, TG, TH, TJ, TM, TN, UA, US, UZ, VN): **UNILEVER PLC** [GB/GB]; a company registered in England and Wales under company no. 41424 of, Unilever House, 100 Victoria Embankment, London EC4Y 0DY (GB).

(71) **Applicant** (for US only): **CONOPCO, INC., D/B/A UNILEVER** [US/US]; 800 Sylvan Avenue, AG West, S. Wing, Englewood Cliffs, New Jersey 07632 (US).

(72) **Inventors**: **CORNELL, Robert, Joseph**; Unilever R&D Port Sunlight, Quarry Road East, Bebington, Wirral, Merseyside CH63 3JW (GB). **GOLDING, Stephen**; Unilever R&D Port Sunlight, Quarry Road East, Bebington, Wirral, Merseyside CH63 3JW (GB). **STOTT, Ian, Peter**; Unilever R&D Port Sunlight, Quarry Road East, Bebington, Wirral, Merseyside CH63 3JW (GB). **THOMPSON,**

Katherine, Mary; Unilever R&D Port Sunlight, Quarry Road East, Bebington, Wirral, Merseyside CH63 3JW (GB).

(74) **Agent**: **ROSEN JACOBSON, Frans, L, M**; Unilever Patent Group, Olivier van Noortlaan 120, NL-3133 AT Vlaarding (NL).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) **Title**: ANTIMICROBIAL COMPOSITION

(57) **Abstract**: The present invention relates to an antimicrobial composition and a method for disinfection involving the antimicrobial composition. It particularly relates to an antimicrobial composition for personal cleaning, oral care or hard surface cleaning applications. It was found that compositions comprising halogenated isopropylmethylphenols, terpineol and a carrier provide synergistic antimicrobial action. In a preferred aspect the composition also comprises 1 to 80 %-wt of one or more surfactants.



WO 2014/001057 A1

ANTIMICROBIAL COMPOSITION

FIELD OF THE INVENTION

The present invention relates to an antimicrobial composition and a method for
5 disinfection involving the antimicrobial composition. It particularly relates to an
antimicrobial composition for personal cleaning, oral care or hard surface cleaning
applications.

BACKGROUND TO THE INVENTION

- 10 Sanitising and disinfecting soap or cleaning compositions are of great benefit to
individuals, since proper use generally may reduce the number of germs and
pathogens the individual is exposed to. Thus, such compositions may for instance play
an important role in reducing the occurrence and spread of infectious diseases.
- 15 Sanitising and disinfecting soap compositions comprising chlorine-based antimicrobial
agent such as triclosan are known. Such compositions require a rather long contact
time to provide efficacious antimicrobial action. In practice, users, in particular children,
do not spend a long time on cleansing and as a result cleaning with such compositions
does not provide adequate prevention from surface or topical infection or adequate
20 protection against diseases. The user, in spite of cleaning his hands, is generally likely
to end up with relatively inadequate bacterial removal from his skin. Therefore, he may
cause contamination of further animate and/or inanimate surfaces and contribute to the
spreading of pathogens and consequent diseases. Users in general and children in
particular who wash contaminated hands before meals with slow-acting antimicrobial
25 compositions for relatively short time are at risk of contracting diseases.

Similarly in the area of hard surface cleaning, e.g. cleaning of floors, table tops or
utensils, the antimicrobial in the compositions are in contact with the substrate for less
than a few minutes after which the surface is either wiped off or rinsed with water.

- 30 These short time scales of cleaning action are ineffective in providing the desired
benefit since most known antimicrobials commonly used in such products take many
minutes to hours to provide the desired kill of microbes.

Therefore, there is a need of providing a composition that –upon application– provides relatively more efficacious antimicrobial action during a relatively short cleaning period, preferably about 30 seconds or less.

- 5 A well-established class of antimicrobially active compounds are phenolic compounds [P A Goddard and K A McCue in “*Disinfection, Sterilisation and Preservation*”, ed. S S Block, 5th edition, Lippincott, Williams and Wilkins, Philadelphia, 2001 pp. 255-282]. However, not every phenolic compound is suitable as an antimicrobial agent. Moreover, many phenols – even if they are antimicrobially active – may exhibit
- 10 undesirable side-effects, such as corrosiveness, malodour and irritating or sensitising effects when applied on the human or animal skin.

WO 2010/046238 A1 describes an effective antimicrobial composition which provides rapid kill of pathogenic bacteria and which comprises 0.01 to 5% by weight of thymol,

15 0.01 to 5% by weight of terpineol and a carrier. WO 2010/046238 A1 also discloses a method of disinfecting a surface including the step of applying the above composition to the surface.

WO 04/035723 is directed to concentrated cleaning and/or disinfecting compositions

20 which bloom when diluted in water. It discloses a hard surface cleaning concentrate composition comprising: a) from about 0.05 to about 10 wt% of a non-cationic antimicrobial agent; b) from about 1 to about 20 wt% of a water soluble organic solvent; c) from about 1 to about 20 wt% of an anionic soap surfactant; d) from about 1 to about 15 wt% of a hydrocarbon diluent; e) from about 0.001 to about 20 wt% of pine oil which

25 is at least 60% terpene alcohols; f) optionally, from about 0 to about 10 wt% of optional materials selected from dyes, colorants, pH stabilizers and buffers, non-ionic surfactants, fragrance/fragrance enhancers, viscosity modifiers, insect repellants, and light stabilizers; and g) the balance being water. Phenol based non-cationic antimicrobials are preferred. Important constituents of the pine oil include terpineol. It is

30 shown that a composition comprising 1.50 wt% of aromatic hydrocarbons (Aromatic 200), 1.00 wt% of Pine Oil 80, 1.50 wt% of PCMX (*p*-chloro-*m*-xylenol), 2.00 wt% of isopropyl alcohol, 20 wt% of NaCOS (sodium castor oil soap – 40%), 0.40 wt% of insect repellent, 73.60 wt% water and added dye provides a Log10 reduction value of greater than 3 in a microbial reduction suspension test, after a contact time of 15

seconds, using 1:60 to 1:100 dilutions of the composition in water and one part of organism suspension (*Staphylococcus aureus* or *Salmonella choleraesuis*).

WO 99/36033 describes a non-aqueous antimicrobial nail coating composition
5 comprising: (a) an antimicrobial agent comprising benzalkonium chloride or a phenol
(b) at least one film-forming polymer or resin; and (c) an organic solvent.

US 2007/202177 discloses an antimicrobial composition comprising: at least one
poloxamer; at least 10 weight percent of at least one alcohol on a basis of total weight
10 of the antimicrobial composition; and at least one biocidal agent which is not an
alcohol.

FR 2 697 133 discloses biocidal and/or biostatic compositions, comprising mono-
oxygenated sesquiterpenes of general formula $C_{15}H_xO$, wherein x is between 20 and
15 26 and aromatic compounds.

Despite the general availability of antimicrobial compounds and compositions, there
remains a continuous need to find alternative antimicrobial compositions and active
compounds that are suitable for use in such compositions. In particular, alternative
20 compositions providing fast antimicrobial action remain highly desirable in view of
current consumer habits. Such alternatives may reduce the dependency on current
raw materials. Moreover, in the field of antimicrobials, the availability of alternatives
may reduce the risk of development of microbial resistance or insensitivity to particular
antimicrobial compounds.

25

In addition, there is a continued need to reduce the total amount of active ingredients
required in such an antimicrobial composition. This need may for instance be driven by
the desire for cost-efficiency, because such compositions are particularly relevant to
developing countries. Moreover, reducing the amounts may also be beneficial for
30 environmental reasons.

A particular problem of certain antimicrobial actives, and especially some phenolic
compounds such as thymol is that they are generally well-perceptible due to their
olfactory properties. Although the latter may – at least for some species – be
35 appreciated in certain fragrance compositions, they are considered too intense by

some users when they are applied at concentrations efficacious in rapid disinfection. Additionally, a lower concentration of odoriferous compounds or the availability of antimicrobial compounds that are less or not odoriferous allows greater flexibility to the manufacturer in providing alternative scents to his composition at lower doses. Hence
5 there is a need to provide alternative antimicrobial compositions and methods that preferably require lower concentrations and/or have a more acceptable sensory profile.

In view of the above-observed problems and drawbacks of the prior art, it is an object of the present invention to provide alternative antimicrobial compositions.
10
It is a particular object of the invention to provide such compositions, requiring a lower dose of antimicrobial compounds.

Similarly, it is an object of the present invention to provide an antimicrobial composition
15 in which the olfactory contribution of the antimicrobially active compounds is reduced or in which the active compound contributes to providing a consumer-acceptable or even consumer-appreciated scent.

It is another particular object of the invention to provide an antimicrobial composition
20 that contributes to reducing the required contact time in a method for disinfection of a surface.

In particular, it is an object of the invention to provide an antimicrobial composition which gives improved disinfection during cleansing of surfaces of the human body,
25 such as the skin and the oral cavity.

It is yet another object of the present invention to provide an alternative method for sanitising and/or disinfecting, in particular of surfaces.

30 It is a further object of the invention to provide a method for disinfection with a reduced disinfection time. More specifically, it is an object of the invention to provide a method, wherein the disinfection time of the method is less than 300 seconds, preferably less than 60 seconds, and more preferably less than 15 seconds.

In particular, it is an object of the invention to provide a method for disinfection that gives improved disinfection during cleansing of surfaces, in particular hard surfaces, or surfaces of the human body, such as the skin and the oral cavity.


5 SUMMARY OF THE INVENTION

We have now found that one or more of the above objects are met by the present invention. Thus, we have found that compositions comprising selected halogenated isopropylmethylphenols and terpineol provide synergistic antimicrobial action. In particular, we found that combinations of halogenated isopropylmethylphenols and
10 terpineol according to this invention are capable of very fast antimicrobial action. For instance, we found that complete microbial inactivation could be effected with compositions according to the present invention after a contact time of only 15 seconds.

15 Accordingly, in a first aspect the invention provides an antimicrobial composition comprising:

- i. 0.001 to 5% by weight of one or more halogenated isopropylmethylphenols,
- ii. 0.001 to 5% by weight of terpineol, and
- iii. a carrier;

20 wherein the halogenated isopropylmethylphenols have the following structure



wherein the substituent X is selected from the group consisting of fluoro, chloro, bromo, and iodo;

and wherein the terpineol is selected from the group consisting of alpha-terpineol, beta-
25 terpineol, gamma-terpineol, delta-terpineol, 4-terpineol, and mixtures thereof.

According to a second aspect of the invention, there is provided a method of disinfecting a surface comprising the steps of

- a. applying a composition according to the invention on to the surface; and
- 30 b. removing the composition from the surface.

In a third aspect, the invention provides the use of a composition according to the invention for improved hygiene.

DETAILED DESCRIPTION OF THE INVENTION

For the avoidance of doubt, any feature of one aspect of the present invention may be utilised in any other aspect of the invention. The word "comprising" is intended to
5 mean "including" but not necessarily "consisting of" or "composed of." Thus, the term "comprising" is meant not to be limiting to any subsequently stated elements but rather to optionally also encompass non-specified elements of major or minor functional importance. In other words, the listed steps or options need not be exhaustive. Whenever the words "including" or "having" are used, these terms are meant to be
10 equivalent to "comprising" as defined above. It is noted that the examples given in the description below are intended to clarify the invention and are not intended to limit the invention to those examples per se.

Except in the examples, or where otherwise explicitly indicated, all numbers in this
15 description indicating amounts of material or conditions of reaction, physical properties of materials and/or use are to be understood as modified by the word "about". Unless specified otherwise, numerical ranges expressed in the format "from x to y" are understood to include x and y. When for a specific feature multiple preferred ranges are described in the format "from x to y", it is understood that all ranges combining the
20 different endpoints are also contemplated.

Phenol in its strict sense refers to hydroxybenzene (C_6H_5OH), but where appropriate the word phenol may in its wider meaning as understood by the skilled person also refer to a member of the class of compounds which comprise at least one such a
25 characteristic hydroxybenzene moiety.

Throughout this description, the term disinfection refers to reduction of the number of viable microorganisms in a given medium or on a given surface by physical or chemical means. Typically, disinfection involves the destruction or inactivation of said
30 microorganisms. Both animate and inanimate media and surfaces are contemplated.

The term "microbicide" refers to a compound capable of killing, inhibiting the growth of or controlling the growth of microorganisms at a locus; microbicides include bactericides, fungicides and algacides. The term "microorganism" includes, for
35 example, fungi (such as yeast and mould), bacteria and algae.

The antimicrobial composition comprises one or more halogenated isopropylmethylphenols, terpineol and a carrier. Various components of the antimicrobial composition are described below.

- 5 The compositions of the present invention are preferred for non-therapeutic use, and more particularly preferred for use in cleaning surfaces of human body including skin, hair or oral cavity or for hard surface cleaning applications.

halogenated isopropylmethylphenols

- 10 The antimicrobial composition according to the invention comprises 0.001 to 5% by weight of one or more halogenated isopropylmethylphenols. The composition comprises preferably 0.005 to 4.5 wt-%, more preferably 0.01 to 4 wt-%, even more preferably 0.02 to 3 wt-%, yet more preferably 0.03 to 2 wt-%, still more preferably 0.04 to 1 wt-%, even more preferably 0.05 to 0.75 wt-% and still more preferably 0.1 to 0.5
- 15 wt-% of the one or more halogenated isopropylmethylphenols. In compositions intended to be diluted before application, the minimum preferred concentrations of the one or more halogenated isopropylmethylphenols can be higher. For example, when washing hands with water and a composition according to the invention, the lather produced, typically is a 50 wt% dilution of the original composition. Similarly, in body
- 20 wash situations, soap bars or soap liquids are typically diluted until about 8 wt% soap in water, corresponding to an approximately tenfold dilution of the product. Therefore, compositions according to the invention intended for dilution upon use preferably comprise 0.05 to 4.5 wt-%, more preferably 0.1 to 4 wt-%, even more preferably 0.2 to 3 wt-%, still more preferably 0.4 to 1 wt-%, and still more preferably 0.5 to 1 wt-% of the
- 25 one or more halogenated isopropylmethylphenols. Thus, the concentration of the one or more halogenated isopropylmethylphenols in the antimicrobial composition is preferably such that, when the composition is diluted or dissolved with a suitable medium during use, the concentration in the diluted or dissolved mixture is still sufficient to be antimicrobially efficacious.


30

- The halogenated isopropylmethylphenol can be a single compound or can be a mixture of the halogenated isopropylmethylphenols as detailed below. In certain preferred embodiments, mixtures of halogenated isopropylmethylphenols are preferred, since such mixtures may show increased antimicrobial activity against a wider range of
- 35 microbes. On the other hand, for reasons including e.g. control over the formulation, it

is preferred that in case the composition according to the invention comprises a mixture of such halogenated isopropylmethylphenols, the mixture preferably comprises at least 30%, more preferably at least 50%, even more preferably at least 70% and still more preferably at least 90% by weight of one halogenated isopropylmethylphenol with
5 respect to the total weight of the halogenated isopropylmethylphenols.

At concentration ranges of the halogenated isopropylmethylphenols below their lower concentration limits, the desired fast acting antimicrobial kinetics in combination with terpeneol would not be met. At concentrations higher than the higher preferred
10 concentrations of halogenated isopropylmethylphenols, when in combination with a terpeneol, while the kinetics of action would not be compromised, the present inventors have found that unlike in therapeutic/pesticidal/herbicidal applications where sensorial aspects are not critical, in the present application, which is preferably a personal cleaning, oral care or hard surface cleaning applications, the product is in contact with
15 hands, mouth or other body parts, the sensorial aspects including smell and skin feel would be compromised.


The halogenated isopropylmethylphenols have the following structure



20 wherein the substituent X is selected from the group consisting of fluoro, chloro, bromo, and iodo.

All positional isomers of the halogenated isopropylmethylphenols are contemplated. For example, the methyl substituent can occupy the 2, 3, 4, 5 or 6 position with respect
25 to the hydroxyl group, and the isopropyl and halo-substituent X can occupy any of these positions not occupied by the methyl substituents. Therefore, the halogenated isopropylmethylphenols can for example be selected from the group consisting of 4-fluoro-2-isopropyl-5-methylphenol, 4-chloro-2-isopropyl-5-methylphenol, 4-chloro-5-methyl-6-isopropylphenol, 6-chloro-2-isopropyl-3-methylphenol, 4-chloro-2-methyl-5-
30 isopropylphenol, 2-chloro-3-methyl-6-isopropylphenol, 2-chloro-4-isopropyl-3-methylphenol, 4-chloro-2-isopropyl-6-methylphenol, 4-bromo-2-isopropyl-5-methylphenol, 3-bromo-6-isopropyl-2-methylphenol, 2-bromo-6-isopropyl-3-methylphenol, and 4-iodo-2-isopropyl-5-methylphenol.

Preferably, the one or more halogenated isopropylmethylphenols have the following structure:



5 Therefore, the halogenated isopropylmethylphenols can for example be selected from the group consisting of 4-chloro-2-isopropyl-5-methylphenol, 4-chloro-2-methyl-5-isopropylphenol, 2-chloro-3-methyl-6-isopropylphenol, 4-bromo-2-isopropyl-5-methylphenol, 2-bromo-6-isopropyl-3-methylphenol, and 4-iodo-2-isopropyl-5-methylphenol.

10

It is preferred that the substituent X is selected from the group consisting of chloro, bromo and iodo. More preferably, X is chlorine.

It is particularly preferred that the one or more halogenated isopropylmethylphenols are
 15 selected from the group consisting of 4-chloro-2-isopropyl-5-methylphenol, and 4-chloro-5-isopropyl-2-methylphenol. Even more preferably, the halogenated isopropylmethylphenol is 4-chloro-2-isopropyl-5-methylphenol. 4-Chloro-2-isopropyl-5-methylphenol is also known as 4-chlorothymol. 4-Chloro-5-isopropyl-2-methylphenol is also known as 4-chlorocarvacrol. The structures of these compounds are provided in
 20 Table 1.

Table 1

4-chloro-2-isopropyl-5-methylphenol	
4-chloro-5-isopropyl-2-methylphenol	

In case the carrier or the dissolution medium during later application is water-based, it
 25 can be advantageous if the one or more halogenated isopropylmethylphenols are

sufficiently water-soluble. The halogenated isopropylmethylphenols are sufficiently water-soluble if they are soluble to at least the minimum concentration required in the antimicrobial composition according to the invention.

- 5 Advantageously, some compounds according to the invention have a weaker odour, when compared to that of thymol, or an odour which can be more appreciable to the consumer, when dosed into the compositions according to the invention at efficacious levels. This benefit especially applies to for instance 4-chloro-2-isopropyl-5-methylphenol.

10

Mixtures of preferred halogenated isopropylmethylphenols are also preferred.

- Without wishing to be bound by theory, it is believed that the synergistic mode of antimicrobial action of the halogenated isopropylmethylphenols according to the
15 present invention in combination with terpineol is similar for the different respective halogenated isopropylmethylphenols.

- The extent to which the halogenated isopropylmethylphenols according to the present invention are soluble in water or in organic solvents can be suitably expressed by the
20 logarithm of the octanol/water partitioning coefficient, $\log\{P\}$. This partition coefficient is a measure for the distribution of a solute between water and octanol at equilibrium. The value of the partitioning coefficient can also be predicted using computational methods. A commonly used predicted value is the so-called AlogP value. A definition and preferred method of calculation of AlogP values is provided in A.K. Ghose, V.N.
25 Viswanadhan, and J.J. Wendoloski, *J.Phys.Chem. A*, vol **102**, pag. 3762 (1998). Due to the standardised calculation as described by Ghose *et al*, AlogP values of different compounds can be compared with one another and used to assess the similarity of or difference between different compounds. For instance, phenol has an AlogP value of 1.59 and thymol has an AlogP value of 3.27. Such comparison may also extend to
30 other properties that are believed to be related to the same molecular properties that are believed to be related to the same molecular properties that underlie the value of AlogP. Though not wishing to be bound by theory in this respect, it is believed that there is some correlation between AlogP and the antimicrobial efficacy of the halogenated isopropylmethylphenols according to the present invention. Therefore, the
35 halogenated isopropylmethylphenols according to the present invention preferably

have an AlogP value of between 2.0 and 4.5, more preferably between 2.2 and 4.4, even more preferably between 2.4 and 4.1 and still more preferably between 2.5 and 4.0.

- 5 Suitable halogenated isopropylmethylphenols according to the present invention may be commercially sourced or obtained via synthetic chemical methods. Such methods are generally well-known to the person skilled in the art.

Terpineol

- 10 The antimicrobial composition according to the invention comprises 0.001 to 5% by weight of terpineol. The composition comprises preferably 0.005 to 4.5 wt-%, more preferably 0.01 to 4 wt-%, even more preferably 0.02 to 3 wt-%, yet more preferably 0.03 to 2 wt-%, still more preferably 0.04 to 1 wt-%, still more preferably 0.05 to 0.75 wt-% and even more preferably 0.1 to 0.5 wt-% of terpineol. In compositions intended
15 to be diluted before application, the minimum preferred concentrations of the terpineol can be higher, for the same reasons as for the halogenated isopropylmethylphenols. Therefore, compositions according to the invention intended for dilution upon use preferably comprise 0.05 to 4.5 wt-%, more preferably 0.1 to 4 wt-%, even more preferably 0.2 to 3 wt-%, still more preferably 0.4 to 1 wt-%, and still more preferably
20 0.5 to 1 wt-% of the terpineol. Any of the concentrations ranges for the terpineol is preferably combined with any of the concentration ranges for the one or more halogenated isopropylmethylphenols specified above. Therefore, the antimicrobial composition according to the invention for example comprises:
- a. 0.01 to 0.4% by weight of the one or more halogenated
 - 25 isopropylmethylphenols;
 - b. 0.05 to 1% by weight of terpineol.

- The terpineol can be a single compound or a mixture of the terpineol isomers as detailed below. Mixtures of terpineols are preferred, since such mixtures may show
30 increased antimicrobial activity against a wider range of microbes. In case the composition according to the invention comprises such a mixture of terpineols, the mixture preferably comprises at least 30%, more preferably at least 50%, even more preferably at least 70% and still more preferably at least 90% by weight of one terpineol isomer with respect to the total weight of the terpineol. The preferred concentrations
35 ranges of the terpineol are important for the same reasons as the preferred






concentration ranges of the one or more halogenated isopropylmethylphenols in meeting the desired fast acting antimicrobial kinetics while not being sensorially unpleasant when used in products for personal cleaning, oral care or hard surface cleaning applications.

5

Terpineol isomers

Terpineol exists as different isomers, which all are considered terpineols. The terpineol is selected from the group consisting of alpha-terpineol, beta- terpineol, gamma-terpineol, delta- terpineol, 4- terpineol, and mixtures thereof. More preferably, the
 10 terpineol is selected from the group consisting of alpha-terpineol, beta- terpineol, gamma- terpineol, delta- terpineol, and mixtures thereof. In nature, the isomeric compounds alpha-terpineol, beta-terpineol and gamma-terpineol are among the most abundant terpineols and often occur together in mixtures. Therefore even more preferably, the terpineol is selected from the group consisting of alpha-terpineol, beta-
 15 terpineol, gamma-terpineol, and mixtures thereof. The structures of these isomers are schematically depicted in Table 2 below.

Table 2

Terpineol isomers		
 alpha-terpineol	 beta- terpineol	 gamma- terpineol
 delta- terpineol	 4-terpineol	

20 Where applicable, the different stereoisomers of these terpineol isomers are contemplated. For instance, cis-beta-terpineol and trans-beta-terpineol are both

contemplated. Thus, compositions comprising enantiomerically pure radicals, racemic mixtures and other mixtures of different stereoisomers are equally preferable.

These terpeneols are all members of the menthenol class of compounds [A L

- 5 Gunatilaka, Natural products in Plants: Chemical Diversity in the Wiley Encyclopedia of Chemical Biology, pp 1-17 and E Breitmaier, Terpenes: flavors, fragrances, pharmaca, pheromones; p. 17, Wiley-VCH, 2006], also termed oxytetrahydrocymenes [F Heusler, The Chemistry of the Terpenes, trans. F J Pond, P Blakistons's son & Co, Philadelphia, 1902, p. 21] which may be defined as monohydroxy alcohol derivatives of p-menthene,
10 with the generic formula $C_{10}H_{17}OH$. Combinations of these terpeneol isomers are often found together in nature, because it is generally believed that their biosyntheses proceed via closely related synthetic pathways. Without wishing to be bound by theory, it is believed that the mode of antimicrobial action of these terpeneol isomers in combination with the halogenated isopropylmethylphenols according to the present
15 invention is similar.

The terpeneols may also be referred to by the alternative names as detailed below in Table 3.

Table 3

alpha-terpineol	<i>p</i> -menth-1-en-8-ol
beta-terpineol	<i>p</i> -menth-8-en-1-ol
gamma-terpineol	<i>p</i> -menth-4(8)-en-1-ol
delta-terpineol	<i>p</i> -menth-1(7)-en-8-ol
4-terpineol	<i>p</i> -menth-1-en-4-ol

20

Terpineol may be added to the antimicrobial composition in purified form. Alternatively pine oil comprising terpeneol may be added to the antimicrobial composition while ensuring that terpeneol is present in the desired concentration in the composition of the present invention.

25

Carrier

The antimicrobial composition according to the invention comprises a carrier. The carrier is preferably selected from the group consisting of water, oil, solvent, inorganic particulate material, starch, air and mixtures thereof. The carrier is preferably from 0.1
30 to 99% by weight of the composition. The antimicrobial composition may be in form of

a solid, liquid, gel, paste or soft solid and the carrier may be selected by a person skilled in the art depending on the format of the antimicrobial composition.

Examples of inorganic particulate materials include clay, talc, calcite, dolomite, silica,
5 and aluminosilicate. Examples of oils include mineral oils, oils of biological origin (e.g. vegetable oils), and petroleum-derived oils and waxes. The oils of biological origin are preferably triglyceride-based. Preferably, the carrier oil is not a perfume oil. Thus, the carrier oil preferably does not substantially contribute to the odour of the composition, more preferably it does not contribute to that odour. Examples of solvents include
10 alcohols, ethers and acetone. The starch may be natural starch obtained from food grains or may be a modified starch.

In certain preferred embodiments, suitable solvents include, for example, water; glycols, including ethylene glycol, propylene glycol, diethylene glycol, dipropylene
15 glycol, polyethylene glycol, and polypropylene glycol; glycol ethers; alcohols, such as methanol, ethanol, propanol, phenethyl alcohol and phenoxypropanol; ketones, including acetone and methyl ethyl ketone; esters, including ethyl acetate, butyl acetate, triacetyl citrate, and glycerol triacetate; carbonates, including propylene carbonate and dimethyl carbonate; and mixtures thereof. It is preferred that the solvent
20 is selected from water, glycols, glycol ethers, esters and mixtures thereof. In certain preferred embodiments, suitable solid carriers include, for example, cyclodextrin, silicas, diatomaceous earth, waxes, cellulosic materials, alkali and alkaline earth (e.g., sodium, magnesium, potassium) metal salts (e.g., chloride, nitrate, bromide, sulfate) and charcoal.

25

Air can for instance be used as a carrier when the halogenated isopropylmethylphenols according to the invention and/or the terpineol are atomised or otherwise dispersed as a fine mist.

30 Particularly preferred carriers are water or oil/solvent and even more preferred is a carrier that is a mixture of water and oil. Thus, in many of the envisaged applications like personal care/washing, oral care and hard surface cleaning, the antimicrobial composition may be formulated with either an aqueous base or a oil/solvent base. Compositions with an aqueous base (water being the carrier), may also for instance be

products in gel format. Compositions with a purely oil/solvent base may for instance be products in anhydrous stick form or propellant-containing products.

Thus, the antimicrobial composition may for instance, preferably be an antimicrobial anhydrous stick personal care composition on a purely oil/solvent base wherein the
5 composition has a water content of less than 0.01% by weight, and wherein the composition preferably is free of water. Alternatively, the antimicrobial composition may for instance, preferably be an antimicrobial propellant-drivable personal care composition, also comprising a propellant. Air can also be used as propellant, for instance in the form of compressed or liquefied air.

10

However, the most preferred product format has an emulsion base (water and/or oil being the carrier) or is capable of forming an emulsion upon dilution, e.g. soap products in liquid, solid, lotion or semisolid form for hand wash, face wash, body wash, or shaving applications; toothpaste/ dentifrices for oral care applications or products for
15 hard surface cleaning in bars or liquids form. If the product comprises an emulsion base, it preferably also comprises one or more surfactants as described below.

Surfactants

The antimicrobial composition according to the invention preferably comprises from 1
20 to 80% by weight of one or more surfactants. Surfactants may for instance advantageously contribute to the cleaning efficacy or the formulation stability of a composition.

Thus, the antimicrobial composition according to the invention preferably comprises

- 25 a. 0.001 to 5% by weight of the one or more halogenated
 isopropylmethylphenols according to the invention,
 b. 0.001 to 5% by weight of terpineol
 c. a carrier, and

from 1 to 80% by weight of one or more surfactants.

30

It is particularly preferred that the antimicrobial composition comprises from 1 to 80% by weight of one or more surfactants in combination with the one or more halogenated isopropylmethylphenols, and the terpineol at their more preferred concentrations as specified above.

35

In general, the surfactants may be chosen from the surfactants described in well-known textbooks like "Surface Active Agents" Vol. 1, by Schwartz & Perry, Interscience 1949, Vol. 2 by Schwartz, Perry & Berch, Interscience 1958, and/or the current edition of "McCutcheon's Emulsifiers and Detergents" published by Manufacturing Confectioners
5 Company or in "Tenside-Taschenbuch", H. Stache, 2nd Edn., Carl Hauser Verlag, 1981; "Handbook of Industrial Surfactants" (4th Edn.) by Michael Ash and Irene Ash; Synapse Information Resources, 2008. Any type of surfactant, i.e. anionic, cationic, nonionic, zwitterionic or amphoteric can be used. Preferably, the one or more surfactants are anionic, nonionic, or a combination of anionic and nonionic surfactants.
10 More preferably, the one or more surfactants are anionic.

A particularly preferred surfactant is soap. Soap is a suitable surfactant for personal washing applications of the antimicrobial composition of the invention. The soap is preferably C₈-C₂₄ soap, more preferably a C₁₀-C₂₀ soap and most preferably C₁₂-C₁₆
15 soap. The soap may or may not have one or more carbon-carbon double bonds or triple bonds. The cation of the soap can for instance be an alkali metal, alkaline earth metal or ammonium. Preferably, the cation of the soap is selected from sodium, potassium or ammonium. More preferably the cation of the soap is sodium or potassium.

20

The soap may be obtained by saponifying a fat and/or a fatty acid. The fats or oils may be fats or oils generally used in soap manufacture, such as tallow, tallow stearines, palm oil, palm stearines, soya bean oil, fish oil, castor oil, rice bran oil, sunflower oil, coconut oil, babassu oil, palm kernel oil, and others. In the above process the fatty
25 acids are derived from oils/fats selected from coconut, rice bran, groundnut, tallow, palm, palm kernel, cotton seed, soyabean, castor etc. The fatty acid soaps can also be synthetically prepared (e.g. by the oxidation of petroleum or by the hydrogenation of carbon monoxide by the Fischer-Tropsch process). Resin acids, such as those present in tall oil, may be used. Naphthenic acids are also suitable.

30

Tallow fatty acids can be derived from various animal sources. Other similar mixtures, such as those from palm oil and those derived from various animal tallow and lard are also included.

A typical fatty acid blend consists of 5 to 30%-wt coconut fatty acids and 70 to 95%-wt fatty acids ex hardened rice bran oil. Fatty acids derived from other suitable oils/fats such as groundnut, soybean, tallow, palm, palm kernel, etc. may also be used in other desired proportions. The soap, when present in solid forms of the present invention, is
5 preferably present in an amount of 30 to 80%, more preferably from 50 to 80%, and even more preferably 55 to 75% by weight of the composition. The soap, when present in liquid forms of the composition is preferably present in 0.5 to 20%, more preferably from 1 to 10% by weight of the composition.

10 Other preferred surfactants are fatty acid glycinates and fatty amphocarboxylates. The fatty acid glycinates are fatty acid amides of salts of glycine, including for example sodium cocoyl glycinate. The fatty amphocarboxylates are amphoteric surfactants including for example sodium lauroamphoacetate (i.e. sodium 2-[1-(2-hydroxyethyl)-2-undecyl-4,5-dihydroimidazol-1-ium-1-yl]acetate). Yet another example of suitable
15 surfactants are derivatives of isethionates, including acylisethionates.

The antimicrobial composition of the invention is also useful in hard surface cleaning applications. In such applications, preferred surfactants are nonionic surfactants, such as C_8-C_{22} , preferably C_8-C_{16} fatty alcohol ethoxylates, comprising between 1 and 8
20 ethylene oxide groups when the product is in the liquid form. When the product for hard surface cleaning applications is in the solid form, surfactants are preferably selected from primary alkyl sulphates, secondary alkyl sulphonates, alkyl benzene sulphonates, ethoxylated alkyl sulphates, or alcohol ethoxylate nonionic surfactants. The composition may further comprise an anionic surfactant, such as alkyl ether
25 sulphate preferably those having between 1 and 3 ethylene oxide groups, either from natural or synthetic source and/or sulphonic acid. Especially preferred are sodium lauryl ether sulphates. Alkyl polyglucoside may also be present in the composition, preferably those having a carbon chain length between C_6 and C_{16} . Other classes of useful surfactants include cationic surfactants, such as long chain quaternary
30 ammonium compounds and amphoteric surfactants such as betaines and alkyl dimethyl amine oxides. Suitable surfactant concentrations in liquid forms of hard surface cleaning application are generally from about from 0.5 to 10%, preferably from 1 to 5 % by weight of the composition. In solid compositions, surfactant is preferably present in 5 to 40%, preferably from 10 to 30% by weight of the composition.

The antimicrobial composition of the invention is useful in oral care compositions e.g. in a dentifrice/ toothpaste or an oral rinse product. In such applications, preferred surfactants are anionic, nonionic or amphoteric in nature, preferably anionic or amphoteric. The anionic surfactant is preferably an alkali metal alkyl sulphate, more preferably a sodium lauryl sulphate (SLS). Mixtures of anionic surfactants may also be employed. The amphoteric surfactant is preferably a betaine, more preferably an alkylamidopropyl betaine (wherein the alkyl group is a linear C₁₀–C₁₈ chain), and most preferably is cocoamidopropyl betaine (CAPB). Mixtures of amphoteric surfactants may also be employed. Suitable surfactant concentrations in oral care application are generally from about 2% to about 15%, preferably from about 2.2% to about 10%, more preferably from about 2.5 to about 5% by weight of the total composition.

Thus, it is highly preferred that the antimicrobial compositions include soap, alkyl sulphate or linear alkyl benzene sulphonate as the surfactants. More preferably, the one or more surfactants are selected from the group consisting of soaps, alkyl sulphates and linear alkyl benzene sulphonates.

Liquid and solid compositions

The antimicrobial composition may be in form of a solid, a liquid, a gel or a paste. A person skilled in the art can prepare compositions in various formats by choosing one or more carrier materials and/or surfactant. The antimicrobial compositions of the present invention are useful for cleansing and care, in particular for skin cleansing and skin care. It is envisaged that the antimicrobial composition can be used as a leave-on product or a wash-off product, preferably a wash-off product. The antimicrobial composition of the present invention can also be used for cleansing and care of hard surfaces such as glass, metal, plastic and the like.

A particularly preferred carrier is water. When water is present, it is preferably present in at least 1%, more preferably at least 2%, further more preferably at least 5% by weight of the composition. When water is the carrier, both liquid and solid compositions are possible. Different amounts of water may be preferred depending on the product format. When water is the carrier, a preferred liquid antimicrobial composition according to the invention comprises:

- a. 0.01 to 5% by weight of the one or more halogenated isopropylmethylphenols,

- b. 0.05 to 5% by weight of the terpineol
- c. 10 to 99.9% by weight of water, and;
- d. 1 to 30% by weight of surfactant.

- 5 The liquid antimicrobial composition is useful for skin cleansing, in particular for hand wash or a face wash.

When water is the carrier, a preferred solid antimicrobial composition according to the invention comprises:

- 10 a. 0.05 to 5% by weight of the one or more halogenated isopropylmethylphenols,
- b. 0.05 to 5% by weight of the terpineol,
- c. 5 to 30% by weight of water, and;
- d. 30 to 80% by weight of surfactant.

- 15 The solid antimicrobial composition is preferably in form of a shaped solid, more preferably a bar. The solid antimicrobial composition is particularly useful for skin cleansing in particular for hand wash or a face wash.

Such a bar-shaped solid antimicrobial composition may for instance be a soap bar.

- 20 Soap bar compositions are well-known and may be similar to the following non-limiting example composition, comprising 75.6 wt-% of anhydrous sodium soap, 1.0 wt-% of glycerine, 0.5 wt-% of sodium carbonate, 0.2 wt-% of EHDP (ethane-1-hydroxy-1,1-disphosphonate) acid, 0.04 wt-% of EDTA (ethylenediaminetetraacetic acid) tetrasodium salt, 8.5 wt-% of hydrated magnesium silicate (Talc), 0.7 wt-% of sodium
- 25 chloride, 0.05 wt-% of dyes, 0.75 wt-% perfume, 0.05 to 10 wt-% of antimicrobial agents including the halogenated isopropylmethylphenols and the terpineol according to the present invention, and water up to 100 wt-%.

Alternatively, inorganic particulate material is also a suitable carrier. When inorganic

- 30 particulate material is the carrier, the antimicrobial composition is in a solid form. Preferably the inorganic particulate material is talc. When the inorganic particulate material is talc, the solid antimicrobial composition is particularly useful as a talcum powder for application on face or body.

According to another alternative, a solvent different from water is a preferred carrier. Although any solvent can be used, alcohol is a preferred solvent. Short chain alcohols –in particular ethanol, propanol, and isopropanol– are particularly preferred as carrier for an antimicrobial wipe or an antimicrobial hand sanitiser composition.

5

Solvents like ethanol and isopropanol generally show antimicrobial efficacy themselves. However, they are also volatile and may readily evaporate during application of the composition. Thus, their levels on the surface that is treated might even reduce until below the minimum level required for antimicrobial action, before the
10 minimum period needed for disinfection has passed. In contrast, the terpeneol and the halogenated isopropylmethylphenols according to the present invention are much less volatile and may therefore yield prolonged antimicrobial action after applying them to the skin.

15 *Additional ingredients*

The composition may further comprise various additional ingredients known to a person skilled in the art. Such additional ingredients include but are not limited to: perfumes, pigments, preservative, emollients, sunscreens, emulsifiers, gelling agents, thickening agents, humectants (e.g. glycerine, sorbitol), sequestrants (e.g. EDTA) or
20 polymers (e.g. cellulose derivatives for structuring such as methyl cellulose)

Both terpeneol and some of the halogenated isopropylmethylphenols according to the invention may contribute to the olfactory properties of the composition. Although some of these compounds might be applied for instance in perfume compositions, the
25 present invention is directed towards antimicrobial compositions. Therefore, the composition is preferably not a perfume composition, although other perfume components can be present. Here, a perfume composition is defined as a composition comprising a plurality of olfactory components, wherein the composition is solely intended to provide a harmonious scent.

30

Synergistic effect of the invention

The inventors have surprisingly found that while one or more of the halogenated isopropylmethylphenols according to the present invention alone or terpeneol alone do not individually provide the fast antimicrobial kinetic action, a combination of one or
35 more halogenated isopropylmethylphenols and terpeneol at the selective concentrations

provides a synergistic antimicrobial action which is especially important in a wash off processes where the contact time of the antimicrobial actives with the surface is low, i.e. of the order of less than 5 minutes, preferably less than 2 minutes, further more preferably less than a minute and in many cases less than 15 seconds.

5

Synergistic combinations of halogenated isopropylmethylphenols and terpineol

The antimicrobial action of two or more active compounds is considered additive if the combined action merely results from the addition of the effects the individual components would have in isolation. In contrast, the antimicrobial action of two or more
10 active compounds is considered to be synergistic if the combined effect of the two or more compounds is stronger than expected based on the assumption of additivity. Without wishing to be bound by theory, it is believed that the antimicrobial action of the one compound may be enhanced by the action of the other compound and vice versa. Such enhancement may for instance originate from cooperative interplay between the
15 mechanisms of antimicrobial action at the molecular level. Such enhanced antimicrobial action may manifest itself for instance by the fact that lower concentrations of active compounds are required to obtain complete microbial kill, or alternatively, that the same extent of microbial kill is arrived at in a shorter time. Whether an antimicrobial composition comprising two or more active compounds is
20 capable of synergistic antimicrobial action may for instance be determined following the procedures and using the criteria as outlined in Example 1 below. Typically, evidence of synergistic antimicrobial action may be provided at concentrations below the minimum biocidal concentrations of each of the components when taken individually. However, it is generally believed that synergistic action can still occur when the
25 concentration of one or more of the active compounds is raised above its minimum biocidal concentration (when taken individually).

The antimicrobial composition according to the present invention preferably comprises the one or more halogenated isopropylmethylphenols and terpineol according to the
30 invention at concentrations at which they are capable of synergistic antimicrobial action. Thus, the concentrations of the one or more halogenated isopropylmethylphenols and of the terpineol in the antimicrobial composition are preferably such that, when the composition is diluted or dissolved with a suitable medium during use, (e.g. when washing hands with water and a composition according
35 to the invention) the concentration in the diluted or dissolved mixture is still sufficient to

be antimicrobially efficacious. That is, to be capable of synergistic antimicrobial action, the concentrations of the one or more halogenated isopropylmethylphenols and the terpeneol in the composition ($C_{\text{comp,phenol}}$, and $C_{\text{comp,terp}}$, respectively) are preferably such that upon application, at a given concentration of the one or more halogenated

5 isopropylmethylphenols in the application medium ($C_{\text{med,phenol}}$), the terpeneol is available at at least a minimum medium concentration ($C_{\text{med, terp}}$) or vice versa (i.e. such that at a given $C_{\text{med,terp}}$, a minimum $C_{\text{med,phenol}}$ is available, sufficient to provided synergistic antimicrobial action). Here, the application medium denotes the medium in which the antimicrobial action desirably takes place. For example, in personal care applications

10 like hand-washing, the composition may be a solid soap bar. In that case, C_{comp} refers to the concentration of the component in the soap bar, whereas C_{med} refers to the concentration in the lather. The minimum and optimum concentrations may for instance be determined by a protocol as described for Example 1 or by one of the standards as detailed below. It is generally preferred that the concentrations of the one or more

15 halogenated isopropylmethylphenols and the terpeneol in the composition according to the invention are equal to or higher than the optimal concentrations in the application medium, because in many typical applications, the composition is either used pure or is diluted to form the application medium.

20 Surprisingly, the synergy between the halogenated isopropylmethylphenols and terpeneol in compositions according to the invention, occurs over a wide range of concentrations and concentration ratios. Depending on factors including the type of antimicrobial composition, its intended application (for instance a hard surface cleaner, a skin cleanser, or a hand sanitiser) different concentration ranges and ratios will be

25 preferred.

Thus, when for instance antimicrobial action against E. Coli is desired, the data of Example 1 may be used to determine preferable medium concentrations C_{med} . For example, in case complete microbial inactivation is desired in the particular medium of

30 Example 1 and if the halogenated isopropylmethylphenol is 4-chloro-2-isopropyl-5-methylphenol, and $C_{\text{med,phenol}}$ is selected as 0.025%(w/v), $C_{\text{med, terp}}$ preferably is at least 0.15 %(w/v) and vice versa.

Alternatively, the desired antimicrobial effect may be obtained by selecting a ratio of the

35 respective concentrations of the one or more halogenated isopropylmethylphenols and

the terpineol. In view of the above-described considerations regarding the intended antimicrobial efficacy and other considerations, including for instance sensory properties, solubility, economic considerations, a concentration ratio of the halogenated isopropylmethylphenols to terpineol larger than one is preferred in some applications, 5 whereas a concentration ratio of the halogenated isopropylmethylphenols to terpineol smaller than one is preferred in others, whereby the concentrations are expressed in %-wt.

Thus, in case a concentration ratio of the halogenated isopropylmethylphenols to 10 terpineol smaller than one is desired, then the antimicrobial composition according to the invention preferably comprises the one or more halogenated isopropylmethylphenols and the terpineol in a concentration ratio (halogenated isopropylmethylphenols : terpineol) of between 1:2 and 1:12, wherein the concentration is expressed as weight percent.

15 A further additional advantage of the present invention is that it is observed that treatment of a surface with a composition according to the invention comprising one or more halogenated isopropylmethylphenols and terpineol, surprisingly enables continued protection of the surface against growth of microbes for a substantial period 20 of time thereafter.

Effect of including surfactant

Favourably, compositions suitable in wash-off processes as described above include a surfactant for the cleaning action. To the further surprise of the inventors, while the 25 surfactant alone does not provide the fast antimicrobial kill at the concentration present in wash off processes, it provides for further improvement in extent of reduction in viable microbial counts on the surface in the short period of time when surfaces are washed with a composition comprising one or more selected halogenated isopropylmethylphenols, terpineol and additionally surfactant. Thus, while surfactant is 30 generally known to be responsible for washing off dirt and also antimicrobial actives used in the composition, in the present invention, it provides a highly useful additional benefit in that it enhances the reduction of viable microbial count in a composition comprising a combination of halogenated isopropylmethylphenols and terpineol alone.

Method according to the invention

According to the second aspect, the invention relates to a method of disinfecting a surface comprising the steps of

- a. applying a composition according to the invention on to the surface; and
- 5 b. removing the composition from the surface.

Preferably, the surface is skin. Thus, for example, a surface like the hands, face, body, or the oral cavity is contacted with the composition of the invention. If the surface is a surface of a human or animal body, the method preferably is a non-therapeutic method
10 of disinfecting a surface. Alternatively, the surface is any hard surface. Typically, such hard surfaces are surfaces that commonly require cleaning and preferably also require sanitisation or disinfection. Such surfaces may be found in many household or industrial environments, and may include for example kitchen and bathroom surfaces, table tops, floors, walls, windows, utensils, cutlery, and crockery. Such surfaces may
15 be made from many different materials, including for instance plastics, wood, metal, ceramics, glass, concrete, marble, and painted surfaces.

The composition may be applied to the surface by any suitable means known to the skilled person. For instance, a suitable means may be pouring, dropping, spraying or
20 wiping in case of liquid compositions.

Preferably, the method includes diluting or dissolving the composition with a suitable solvent, preferably water, before or whilst applying the composition to the surface. Such dissolving is preferred in particular in case the composition is a solid composition.
25 Alternatively, solid compositions may also be directly spread, rubbed, or sprayed, e.g. in the form of a powder.

The method according to the first aspect of the present invention also includes the step of removing the composition from the surface. Here, removing the composition also
30 encompasses partially removing the composition, because traces of the composition may remain on the surface. In many typical situations, including washing of the skin or hard-surface cleaning, it is acceptable or sometimes even desirable if part of the composition – in particular certain active ingredients – remains on the surface.

Therefore, step b preferably involves removing at least 5%, more preferably at least
35 10%, even more preferably at least 25%, still more preferably at least 50% and yet

more preferably at least 75% of the composition by weight. Preferably, the step of removing the composition comprises rinsing the surface with a suitable solvent or wiping the surface with a suitable wipe, more preferably, this step consists of rinsing the surface with a suitable solvent or wiping the surface with a suitable wipe.

- 5 Alternatively, the removal step can also include evaporation of part of the composition, for example when the composition comprises volatile components, e.g. solvents.

A suitable medium for rinsing the surface is water but it could also be for example a mixture of water and alcohol. It is then rinsed preferably with sufficient amounts of
10 water after a pre-determined period of time to remove any visible or sensory residue of the composition. Alternatively, an alcohol wipe or a water/alcohol impregnated wipe may be used to wipe the surface to be visibly free of the anti-microbial composition. The step of removing the composition (e.g. by rinsing or wiping the surface) is preferably started less than 5 minutes, more preferably less than 2 minutes, even more
15 preferably less than 1 minute, still more preferably less than 30 seconds and yet more preferably less than 20 seconds after commencement of the step of applying the composition on the surface, because of the surprisingly fast antimicrobial action of the compositions according to the present invention. Even though partial microbial kill may be almost instantaneous upon application of the composition according to the
20 invention, it is preferred that the step of removing the composition from the surface is started out at least 5 seconds, preferably at least 10 seconds, more preferably at least 15 seconds after commencement of the step of applying the composition on the surface, in order to effect optimal antimicrobial action. Combinations of these times into time intervals are preferred too. Therefore, it is particularly preferred that the step
25 of removing the composition from the surface (i.e. step b) is started between 2 minutes and 5 seconds, more preferably between 1 minute and 10 seconds, even more preferably between 30 and 10 seconds and still more preferably between 20 and 15 seconds after commencement of the step of applying the composition on the surface (i.e. step a).

30

Disinfection time

These times between applying the composition and rinsing or wiping are preferably related to the disinfection time, in order to ensure optimal cleansing and sanitising of the surface. Therefore, the invention preferably relates to a method, wherein the
35 disinfection time T of said method is less than 300 seconds, preferably less than 60

seconds, and more preferably less than 15 seconds; wherein T is defined as the time that elapses from the moment of adding the composition to a microbial culture until the number of microbes per unit volume of the culture is reduced by a factor of 100 000; and wherein the initial number of microbes preferably exceeds about 100 000 000
5 microbes per millilitre and wherein the composition is preferably a liquid composition.

The disinfecting action of the method (as may be expressed in terms of the disinfection time T) is preferably determined according to the protocol of Example 1 as described hereinafter. This test relates to a standardised test environment in which the microbial
10 culture is kept in suspension. A similarly suitable test is the standard suspension method described in European Standard EN1276, with the proviso that the disinfection time is adapted to suit the above criteria as will be clear to a person skilled in the art. Alternatively, one of the test methods as described in WO 2010/046238 may for instance be applied to establish the disinfecting action.
15 Such test methods may preferably also be used by the skilled person to determine the optimal concentrations of the one or more halogenated isopropylmethylphenols and the terpineol in an antimicrobial composition according to the present invention.

Alternatively, since the method is directed towards surface disinfection, the disinfection
20 time may also be determined by test methods involving a surface. Therefore, the invention preferably relates to a method according to the present invention, wherein the surface disinfection time T₂ of said method is less than 60 seconds, preferably less than 15 seconds, wherein T₂ is defined as the time starting from the moment of applying the composition to the surface to be disinfected after which the number of
25 microbes per unit area is reduced by a factor of 10000 (i.e. a 4 log reduction), wherein the initial number of microbes preferably exceeds 10³, more preferably 10⁵, and even more preferably 10⁷ microbes per square centimetre. Such tests may for instance be performed as described in WO 2010/046238, or as described in European Standards EN 13697:2001 and EN 1500:1997.

30

Use according to the invention

The invention preferably provides for non-therapeutic benefits. Thus, for instance, the invention relates to use of an antimicrobial composition according to the present invention for faster reduction in viable microbial count.

35

Thus, according to the third aspect of the invention, there is provided use of a composition according to the invention for improved hygiene. Such use relates for example to use of an antimicrobial composition comprising the one or more halogenated isopropylmethylphenols, terpineol and a carrier, for reduction in viable microbial count, preferably fast reduction of viable microbial count. Thus, such use preferably is use in a method for disinfection. Fast reduction in viable microbial count therefore preferably relates to use for disinfection whereby the disinfection time is less than 300 seconds, preferably less than 60 seconds, and more preferably less than 15 seconds. Here, the disinfection is preferably defined similar to the disinfection times T and T₂ as described above.

Thus, there is provided use of a composition according to the invention for improved hygiene of surfaces of the human body. Such surfaces include e.g. skin, hands and the oral cavity. According to a preferred aspect, the invention relates to use of a composition according to the invention for improved hand hygiene. According to another preferred aspect, the invention relates to use of a composition according to the invention for improved oral hygiene.

The microbicide compositions of the present invention can be used to inhibit the growth of microorganisms by introducing a microbicidally effective amount of the compositions onto, into or at a locus subject to attack. For instance, in the field of institutional and industrial applications, suitable loci include, for example: industrial process water including electrocoat deposition systems, cooling towers and air washers; gas scrubbers; wastewater treatment; ornamental fountains; reverse osmosis filtration; ultrafiltration; ballast water; evaporative condensers and heat exchangers; pulp and paper processing fluids and additives; mineral slurries; starch; plastics; emulsions; dispersions; paints; latices; coatings, such as varnishes; construction products, such as mastics, caulks, and sealants; construction adhesives, such as ceramic adhesives, carpet backing adhesives, and laminating adhesives; industrial or consumer adhesives; photographic chemicals; printing fluids; household and institutional products used in restaurants, healthcare facilities, schools, food processing facilities and farms including, cleaners, sanitizers and disinfectants, wipes, soaps, detergents, floor polishes and laundry rinse water; cosmetics; toiletries; shampoos; metalworking fluids; conveyor lubricants; hydraulic fluids; leather and leather processing products; textiles; textile and textile processing products; wood and wood processing products, such as

plywood, chipboard, wallboard, flakeboard, laminated beams, oriented strandboard, hardboard, and particleboard; oil and gas processing fluids such as injection fluids, fracture fluids, drilling muds and produced water; fuel transportation and storage systems; agriculture adjuvant preservation; surfactant preservation; medical devices; 5 diagnostic reagent preservation; food preservation, such as plastic or paper food wrap; food, beverage, and industrial process pasteurizers; toilet bowls; recreational water; pools; and spas.

Preferably, the microbicidal compositions of the present invention are used to inhibit 10 the growth of microorganisms at a locus selected from one or more of mineral slurries, pulp and paper processing fluids and additives, starch, emulsions, dispersions, paints, latices, coatings, construction adhesives (such as ceramic adhesives), carpet backing adhesives, photographic chemicals, printing fluids, household and institutional products such as cleaners, sanitizers, disinfectants, wipes, cosmetics, toiletries, shampoos, 15 soaps, detergents, floor polishes, laundry rinse water, metal working fluids, textile products, wood and wood products, agriculture adjuvant preservation, surfactant preservation, diagnostic reagent preservation, food preservation, food, beverage, and industrial process pasteurizers and oil and gas processing fluids.

20 *Fields of use*

The composition according to the invention can in view of the above be applied for disinfection, reduction in viable microbial count or improved hygiene, especially at a surface. In preferred embodiments, the composition is particularly suited for application to the skin. For example, a surface like the hands, face, body, or the oral cavity can 25 suitably be contacted with the composition of the invention. In other preferred embodiments, the surface is any hard surface. Typically, such hard surfaces are surfaces that commonly require cleaning and often also require sanitisation or disinfection. Such surfaces can be found in many household or industrial environments, and can include for example kitchen and bathroom surfaces, table tops, floors, walls, 30 windows, utensils, cutlery, and crockery. Such surfaces can be made from many different materials, for instance plastics, wood, metal, ceramics, glass, concrete, marble, and painted surfaces. In other preferred embodiments, the compositions can be used for such disinfection, reduction in viable microbial count or improved hygiene at loci other than the surfaces as described hereinbefore.

In preferred embodiments, the invention relates to compositions according to the invention for use as or incorporation in home care products and personal care products. More preferably, this embodiment of the invention relates to a composition according to the invention which is a home care product or a personal care product.

5

A "home care product" is a product for the treatment, cleaning, caring or conditioning of the home or any of its contents. The foregoing includes, but is not limited to, compositions, products, or combinations thereof relating to or having use or application in the treatment, cleaning, cleansing, caring or conditioning of surfaces, furniture and atmosphere of the home and household contents, such as clothes, fabrics and/or cloth fibres and the manufacture of all of the foregoing products. A "personal care product" is a product for the treatment, cleaning, caring or conditioning of the person. The foregoing includes, but is not limited to, chemicals, compositions, products, or combinations thereof relating to or having use or application in the treatment, cleaning, cleansing or conditioning of the person (including in particular the skin, hair and oral cavity), and the manufacture of all the foregoing. Home care products and personal care products are for example products marketed under mass market brands, non-limiting examples being soap bars, deodorants, shampoos, and home surface sanitisers/disinfectants.

20

Another preferred embodiment of the invention relates to compositions according to the invention for use as or incorporation in industrial and/or institutional products. More preferably, this embodiment of the invention relates to a composition according to the invention which is an industrial and/or an institutional product. Industrial and institutional products are for example products being marketed under professional brands, non-limiting examples being for industrial, institutional, janitorial, and medical cleaning, cleaning-in-place, food services, veterinary, and agricultural products. Industrial and/or institutional products also include products for cleaning of the person (such as hand sanitisers) for medical offices, hospitals and/or other institutions.

30

In another preferred embodiment, the invention also relates to a method or use according to the invention involving home care products or personal care products. For example, the method according to the invention – which comprises application of a composition according to the invention in step a – can be a method wherein that composition is a composition for use as or incorporation in home care products and

35

personal care products as described hereinabove. Similarly, in another preferred embodiment, the invention also relates to a method or use according to the invention involving industrial and/or institutional products. For example, the method according to the invention – which comprises application of a composition according to the invention

5 in step a – can be a method wherein that composition is a composition for use as or incorporation in industrial and/or institutional products as described hereinabove.

Products and/or methods for use in the home care or personal care field are generally distinct from products and/or methods for use in the industrial and/or institutional field.

10 Thus, for example, a product that is marketed as a home or personal care product will generally not be marketed as a product for industrial and/or institutional use and vice versa. Therefore, certain embodiments of the present invention, when carried forth into practice, will relate to the one field, but not the other.

15 **EXAMPLES**

The invention is illustrated by the following non-limiting examples.

Example 1: Assessment of antimicrobial efficacy

Materials

20 Halogenated isopropylmethylphenols were purchased as fine chemicals from suppliers such as Sigma Aldrich, Alfa Aesar and TCI Fine Chemicals. A terpeneol mixture comprising ca 88 wt-% of (S)-alpha-terpineol, and 12 wt-% of gamma-terpineol was purchased from Sigma Aldrich and used throughout the examples (being referred to as alpha-terpineol or terpeneol) unless specified otherwise.

25

General method for assessment of antimicrobial synergy

The efficacies of antimicrobial agents can be usefully compared by determining the Minimum Biocidal Concentration (MBC). The MBC is defined as the lowest absolute concentration of the particular active that provides complete kill (zero bacterial growth).

30

The differing behaviours of inhibitory antimicrobials in isolation and mixtures have been widely explored using the concept of the Fractional Concentration and Fractional Inhibitory Concentration (FIC). See for instance JRW Lambert and R Lambert, *J. Appl. Microbiol* **95**, 734 (2003); T. Jadavji, CG Prober and R Cheung, *Antimicrobial Agents*

and *Chemotherapy* **26**, 91 (1984), and WO 2004/006876. These parameters can be defined as follows:

$$\text{FC (component a)} = \frac{\text{Concentration of component a tested in the mixture}}{\text{MIC (component a tested as a single active)}}$$

$$\text{FIC (component a)} = \frac{\text{MIC (component a tested in the mixture)}}{\text{MIC (component a tested as a single active)}}$$

5

By analogy the Fractional Biocidal Concentration (FBC) is given by:

$$\text{FBC (component a)} = \frac{\text{MBC (component a tested in the mixture)}}{\text{MBC (component a tested as a single active)}}$$

The interactions between antimicrobials can be additive, synergistic or possibly
 10 antagonistic depending on whether the efficacy of the combination is equivalent to, greater than or less than that obtained for the same total concentration of the individual components when tested alone.

These relationships can be expressed mathematically by summing the fractional MBC
 15 values for all the components present in the mixture to give the "fractional biocidal index":

$$\Sigma\text{FBC} = \text{FBC}_{(\text{component 1})} + \text{FBC}_{(\text{component 2})} + \text{FBC}_{(\text{component 3})} + \dots \text{ etc}$$

such that

20 $\Sigma\text{FBC} \geq 1$ corresponds to additive or antagonistic bactericidal activity

$\Sigma\text{FBC} < 1$ corresponds to synergistic bactericidal activity

Experimental method

Antimicrobial efficacy is tested against a representative pathogenic bacterial organism,
 25 Gram negative *Escherichia coli*. Concentrations of actives are expressed in terms of the percentage weight/volume (%w/v) throughout Example 1.

Bacterial stock

An overnight culture of *Escherichia coli* (10536 strain) was prepared in 50 ml total volume of TSB broth, grown for ca. 18 hrs at 37°C and shaken at 150 rpm. 1 ml of this overnight *E. coli* culture was transferred to 50 ml of fresh TSB broth and incubated at 5 37°C at 150 rpm for ca. 4 hours. This culture was separated into equal volumes and centrifuged at 4000 rpm for 15 minutes, washed with sterile saline (0.85% NaCl), centrifuged once more and re-suspended in saline to give a final concentration of 0.8 OD₆₂₀ equivalent to about 10⁸ cells per millilitre for this particular organism. Here, OD₆₂₀ indicates the absorbance of a sample in a cuvette of 1.0 cm path length at a 10 wavelength of 620 nm. This bacterial stock was used for assaying against antimicrobial actives (in triplicate).

Protocol

The following assay describes the testing of 8 materials using 6 dilutions across half of 15 a 96-well micro titre plate (MTP). Using this approach it is possible to assay 16 actives (without replicates) with one full dilution plate, replicating this set up in two halves of the plate columns, 1-6 and 7-12.

1M solutions of the test actives were prepared in dimethylsulphoxide (DMSO). Stock 20 solutions of the actives at 1.11 times the desired final concentration were prepared by diluting the DMSO solutions in water, so that for example a 0.89% w/v solution was prepared for a desired "in test" concentration of 0.8% w/v in order to allow for the further dilution of the active when the bacterial suspension is added (dilution from 270µl to 300 µl), as described below.

25

Aliquots (270 µl) of the materials at 1.11 times the final concentration were dispensed into the wells of the MTP along one column (A1-H1). This MTP was labelled as the "Screening plate".

30 In another MTP, labelled as the "Dilution plate", 270µl of D/E neutralising solution from DIFCO Composition was added to column 1. The composition of the neutralising solution was as follows: pancreatic digest of casein, 5.0g/L; Yeast Extract, 2.5 g/L; Dextrose, 10 g/L, sodium thioglycollate, 1.0 g/L, sodium thiosulphate, 6.0 g/L; sodium bisulphite, 2.5 g/L; Polysorbate 80, 5.0 g/L; lecithin 7.0 g/L; bromocresol purple, 0.02 35 g/L with a pH in the range 7.6±0.2.

270µl of tryptone diluent solution was added to all the remaining wells of the Dilution MTP (columns 2-6).

- 5 Bacterial stock (30 µl) was then added to the prepared 270 µl of the solution of actives in the Screening Plate and mixed, using a multichannel pipette with 8 tips to aspirate and dispense the same volume of bacterial stock in parallel to 8 wells in rows A-H. After a contact time of 15 seconds, the mixtures were quenched by transferring 30µl volumes of the mixtures into the 270µl D/E neutralising solution in the prepared dilution
- 10 plate, using aspiration to mix. After exactly 5 minutes in the D/E neutralising solution, 30µl volumes were transferred from column 1 to column 2 of the Dilution MTP and mixed, before transferring further 30µl volumes from column 2 into column 3. This process was repeated serially diluting the bacteria across the plate to column 6.
- 15 30µl volumes from each well in the Dilution MTP were transferred onto pre-labelled segment of Tryptone Soya Agar (TSA) plates starting from the lowest bacterial concentration (highest dilution, column 6) to the highest bacterial concentration (column 1). The TSA plates were allowed to stand for ca. 2 hours so that the 30µl inocula spots could dry and the plates were then inverted and incubated overnight at
- 20 37°C before enumerating the bacterial colonies at the labelled dilutions to determine the effects of the actives on bacterial growth.

Calculation of results

- Mean bacterial survival numbers N_{MBS} (expressed in Log CFU/ml) are obtained by first
- 25 determining the segment of the TSA plate where the number of bacterial colonies is countable. From the colony number in this segment, N_{MBS} is calculated by the formula:

$$N_{MBS} = \log\{ N_{col} \cdot 10^{DF} \cdot 100 / 3 \}$$

- Here, N_{col} is the colony count, and DF is the dilution factor taken from the MTP-well corresponding to the TSA plate segment (i.e. DF may range from 1 for the quench, to 6
- 30 for the highest dilution). The factor 100/3 is a conversion factor from the volume of the inocula spot to one millilitre.

Every assay test was performed in triplicate. The reported mean bacterial survival results are the average of such a triplet, the error is the corresponding standard deviation.

Thus, a value of N_{MBS} of about 7 corresponds to a count of about 3 colonies from the fifth dilution well, i.e. with $DF = 5$. Such a count of about 7 is generally observed when bacteria are exposed to non-biocidal materials. In case no surviving colonies are observed in any segment of the TSA plate, this is interpreted as complete kill and a value of $N_{\text{MBS}}=0$ is reported.

Validation

All test results were validated by running every test assay in parallel with four control experiments on the same MTP. All control experiments are executed exactly according to the above protocol, but with the following active ingredients:

- A 0.025 % (w/v) thymol
- B 0.15 % (w/v) alpha-terpineol
- C 0.025 % (w/v) thymol + 0.15 % (w/v) alpha-terpineol
- D no active component

The control experiments A, B and D validate a test assay by not showing bacterial kill, whereas control experiment C, comprising a synergistic combination of thymol and alpha-terpineol according to WO 2010/046238 A1 validates a test assay by showing complete bacterial kill.

A reference experiment according to the above protocol, but without active component, showed that DMSO does not affect bacterial growth at the concentrations present in the test solutions in this protocol (<5 % (w/v)), as can be seen in Table 4.

Table 4

DMSO in water (% w/v)	Mean bacterial survival [log CFU/ml]	Standard deviation
4.5	8.2	0.1
3.6	8.4	0.2
2.7	8.2	0.1
1.8	8.5	0.2
0.9	8.6	0.1
0.0	8.5	0.1

Results

The above method was applied to assess the antibacterial efficacy of the halogenated isopropylmethylphenols according to the invention. Table 5 shows the antibacterial activities of 4-chloro-2-isopropyl-5-methylphenol, both alone and in conjunction with

5 terpineol.

Table 5: Antibacterial activities of thymol, 4-chloro-2-isopropyl-5-methylphenol, and triclosan, alone and/or in combination with terpineol

Example	Phenol concentration C_{phenol} (% w/v)	Concentration of terpineol C_{TERP} (% w/v)	N_{MBS}^a	S.D. ^b
1:1*	0.075 thymol	0	0	0
1:2*	0.05 thymol	0	0	0
1:3*	0.025 thymol	0	> 7	0.1
1:4*	0	0.5%	0	0
1:5*	0	0.4%	0	0
1:6*	0	0.3%	7	0.2
1:7*	0	0.15%	7	0.2
1:8*	0.075% 4-chloro-2-isopropyl-5-methylphenol	0	0	0
1:9*	0.05 % 4-chloro-2-isopropyl-5-methylphenol	0	7.6	0.1
1:10*	0.0375% 4-chloro-2-isopropyl-5-methylphenol	0	7.1	0.2
1:11*	0.025% 4-chloro-2-isopropyl-5-methylphenol	0	7.7	0.1
1:12	0.0375% 4-chloro-2-isopropyl-5-methylphenol	0.15%	0	0
1:13	0.025% 4-chloro-2-isopropyl-5-methylphenol	0.15%	0	0
1:14*	0.025% Triclosan ^c	0.15%	7.8	0.4

* Examples (1:1) to (1:11) and (1:14) are comparative examples

^a N_{MBS} in [log CFU/ml]

^b S.D. = standard deviation

^c Triclosan = 5-Chloro-2-(2,4-dichlorophenoxy)-phenol

Comparative examples

Determination of the parameter Σ FBC, which is used as measure of the synergistic antimicrobial action of compositions according to the present invention, requires

- 5 determination of the Minimum Biocidal Concentrations (MBCs) of the relevant actives first. As described above, the MBC for an active can be defined as the lowest concentration of the active that provides zero bacterial survival in the particular medium. Data for examples (1:1) to (1:3) demonstrate that the MBC value for thymol is 0.05% (w/v). For alpha-terpineol, compositions (1:4) to (1:7), show that the
- 10 MBC is 0.4% w/v. The same analysis has been carried out for 4-chloro-2-isopropyl-5-methylphenol and is summarised in Table 6 below. These MBC values constitute the upper boundaries to the respective MBCs in the particular medium used in these examples.

15 **Table 6:** Minimum biocidal concentrations of antimicrobial components

Component	MBC (% w/v)
thymol	0.05
alpha-terpineol	0.4
4-chloro-2-isopropyl-5-methylphenol	0.075

It is clear from the data in Table 6 that 4-chloro-2-isopropyl-5-methylphenol is antimicrobially efficacious.

20 *Synergistic interactions*

The tested combinations of 4-chloro-2-isopropyl-5-methylphenol with terpineol provide complete bacterial kill in the examples (1:12) and (1:13). Using the MBC values listed in Table 6 above, the fractional MBC values for the components present in these mixtures and the experimental Σ FBC of the compositions can be calculated in order to

- 25 discriminate between combinations providing evidence of synergistic effects, as opposed to additive biocidal effects. The results from this analysis are given in Table 7.

Table 7: Extent of synergistic interactions between binary compound mixtures for compositions providing complete bacterial kill


phenol				terpineol		Σ FBC	Evidence of Synergy ^c
compound	Ex.	MBC %(w/v)	FBC ^a	MBC %(w/v)	FBC ^b		
4-chloro-2-isopropyl-5-methylphenol	1:12	0.075	0.5	0.4	0.38	0.88	Yes
4-chloro-2-isopropyl-5-methylphenol	1:13	0.075	0.33	0.4	0.38	0.71	Yes
^a FBC of phenol: $C_{\text{phenol}}/MBC_{\text{phenol}}$ ^b FBC of terpineol: $C_{\text{terpineol}}/MBC_{\text{terpineol}}$ ^c Criterion for synergy: (Σ FBC <1)							

- 5 For Examples (1:12) and (1:13), the Σ FBC value is below 1, thus providing evidence for synergistic interactions, according to the set criteria. Therefore, these examples show how the antimicrobial efficacy of terpineol and the halogenated isopropylmethylphenols according to invention are enhanced when they are applied together. Such synergies allow for reductions in the concentrations of the antimicrobials required to achieve
- 10 complete kill. For example 0.4% w/v terpineol is required to achieve complete bacterial kill when tested in isolation but this can be reduced 2.7-fold to 0.15 % when used in combination with 0.25% w/v of 4-chloro-2-isopropyl-5-methylphenol.

CLAIMS

1. An antimicrobial composition comprising:
 - i. 0.001 to 5% by weight of one or more halogenated isopropylmethylphenols,
 - ii. 0.001 to 5% by weight of terpeneol, and
 - iii. a carrier;

wherein the halogenated isopropylmethylphenols have the following structure




wherein the substituent X is selected from the group consisting of

fluoro,
chloro,
bromo, and
iodo;

and wherein

the terpeneol is selected from the group consisting of alpha-terpineol, beta-terpineol, gamma-terpineol, delta-terpineol, 4-terpineol, and mixtures thereof.

2. An antimicrobial composition according to claim 1, wherein the one or more halogenated isopropylmethylphenols have the following structure



3. An antimicrobial composition according to claim 1 or 2 wherein X is chlorine.
4. An antimicrobial composition according to claim 3, wherein the one or more halogenated isopropylmethylphenols are selected from the group consisting of
 - 4-chloro-2-isopropyl-5-methylphenol, and
 - 4-chloro-5-isopropyl-2-methylphenol.
5. An antimicrobial composition according to claim 4, wherein the halogenated isopropylmethylphenol is 4-chloro-2-isopropyl-5-methylphenol.

6. An antimicrobial composition according to any one of claims 1 to 5 comprising
 - i. 0.05 to 0.3% by weight of the one or more halogenated isopropylmethylphenols, and
 - ii. 0.05 to 1% by weight of the terpeneol.
7. An antimicrobial composition according to any one of claims 1 to 6 comprising from 1 to 80% by weight of one or more surfactants.
8. An antimicrobial composition according to claim 7 wherein the one or more surfactants are anionic, nonionic, or a combination of anionic and nonionic surfactants.
9. An antimicrobial composition according to claim 7 or 8 wherein the one or more surfactants are selected from the group consisting of soaps, alkyl sulphates, and linear alkyl benzene sulphonates.
10. A solid antimicrobial composition according to any one of claims 7 to 9 comprising:
 - a. 0.05 to 5% by weight of the one or more halogenated isopropylmethylphenols,
 - b. 0.05 to 5% by weight of the terpeneol,
 - c. 5 to 30% by of weight water, and;
 - d. 30 to 80% by weight of the one or more surfactants.
11. A method of disinfecting a surface comprising the steps of
 - a. applying a composition according to any one of claims 1 to 10 on to the surface; and
 - b. removing the composition from the surface.
12. A method according to claim 11, wherein the disinfection time T of said method is less than 300 seconds, preferably less than 60 seconds, and more preferably less than 15 seconds; wherein T is defined as the time that elapses from the moment of adding the composition to a microbial culture until the number of microbes per unit volume of the culture is reduced by a factor of 100 000; and wherein the initial number of microbes preferably exceeds about 100 000 000 microbes per millilitre and wherein the composition is preferably a liquid composition.

13. Use of a composition according to any of claims 1 to 10 for improved hand hygiene.

14. Use of a composition according to any of claims 1 to 10 for improved oral hygiene.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/061705

A. CLASSIFICATION OF SUBJECT MATTER INV. A01P1/00 A01N31/08 A01N31/06 A01N31/04 A61K8/34 A61K31/05 ADD. According to International Patent Classification (IPC) or to both national classification and IPC											
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) A01N A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, CHEM ABS Data, WPI Data											
C. DOCUMENTS CONSIDERED TO BE RELEVANT <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 10%;">Category*</th> <th style="width: 70%;">Citation of document, with indication, where appropriate, of the relevant passages</th> <th style="width: 20%;">Relevant to claim No.</th> </tr> </thead> <tbody> <tr> <td style="text-align: center; vertical-align: top;">X</td> <td style="vertical-align: top;"> WO 2010/046238 A1 (UNILEVER NV [NL]; UNILEVER PLC [GB]; UNILEVER HINDUSTAN [IN]; CHAKRABO) 29 April 2010 (2010-04-29) cited in the application claims 1-14; table 1 ----- </td> <td style="text-align: center; vertical-align: top;">1-14</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">X</td> <td style="vertical-align: top;"> FR 2 697 133 A1 (TRANSMIOTECH [FR]; TERROM GERARD) 29 April 1994 (1994-04-29) pages 5,6,8; claims 2,5,6,13; tables 1,2 ----- </td> <td style="text-align: center; vertical-align: top;">1-14</td> </tr> </tbody> </table>			Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	X	WO 2010/046238 A1 (UNILEVER NV [NL]; UNILEVER PLC [GB]; UNILEVER HINDUSTAN [IN]; CHAKRABO) 29 April 2010 (2010-04-29) cited in the application claims 1-14; table 1 -----	1-14	X	FR 2 697 133 A1 (TRANSMIOTECH [FR]; TERROM GERARD) 29 April 1994 (1994-04-29) pages 5,6,8; claims 2,5,6,13; tables 1,2 -----	1-14
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.									
X	WO 2010/046238 A1 (UNILEVER NV [NL]; UNILEVER PLC [GB]; UNILEVER HINDUSTAN [IN]; CHAKRABO) 29 April 2010 (2010-04-29) cited in the application claims 1-14; table 1 -----	1-14									
X	FR 2 697 133 A1 (TRANSMIOTECH [FR]; TERROM GERARD) 29 April 1994 (1994-04-29) pages 5,6,8; claims 2,5,6,13; tables 1,2 -----	1-14									
☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex.											
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family											
Date of the actual completion of the international search 10 October 2013		Date of mailing of the international search report 18/10/2013									
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Kamdzhilov, Yavor									

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2013/061705

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2010046238	A1	29-04-2010	AR 073903 A1 09-12-2010
			AU 2009306592 A1 29-04-2010
			CA 2739843 A1 29-04-2010
			CN 102186341 A 14-09-2011
			EA 201100656 A1 31-10-2011
			EP 2348838 A1 03-08-2011
			ES 2414158 T3 18-07-2013
			JP 2012505851 A 08-03-2012
			KR 20110081198 A 13-07-2011
			US 2011223114 A1 15-09-2011
			WO 2010046238 A1 29-04-2010

FR 2697133	A1	29-04-1994	NONE
