




(86) Date de dépôt PCT/PCT Filing Date: 2013/09/13
(87) Date publication PCT/PCT Publication Date: 2014/03/20
(45) Date de délivrance/Issue Date: 2019/12/31
(85) Entrée phase nationale/National Entry: 2015/03/13
(86) N° demande PCT/PCT Application No.: EP 2013/068984
(87) N° publication PCT/PCT Publication No.: 2014/041112
(30) Priorité/Priority: 2012/09/14 (DE10 2012 108 621.4)

(51) Cl.Int./Int.Cl. *A01M 1/02* (2006.01),
A01M 1/20 (2006.01), *A01N 59/04* (2006.01),
A01N 63/00 (2006.01), *A01N 65/00* (2009.01)
(72) Inventeur/Inventor:
PATEL, ANANT, DE
(73) Propriétaire/Owner:
BIOCARE GESELLSCHAFT FUR BIOLOGISCHE
SCHUTZMITTEL MBH, DE
(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : SYSTEMES D'APPAT POUR PARASITES ET LEUR UTILISATION
(54) Title: ATTRACTION SYSTEMS FOR PESTS AND USE THEREOF



(57) **Abrégé/Abstract:**

The present application is directed to a system and device for attraction of pests. This system comprises CO₂-releasing microorganisms, and also nutrients specific to these microorganisms, and this system, or the device, comprises one or more biodegradable biopolymers and allows CO₂, and optionally other attractants, to be released over a period of more than 20 days. Furthermore, the present application provides for the use of such a system or such a device for the attraction of pests, more particularly of maize or potato pests, such as larvae of the Western corn rootworm or wire worm. The present application is also directed, lastly, to methods for attracting pests, more particularly pests of maize or potatoes, such as the larvae of the Western corn rootworm, or for wire worms, with the systems or devices of the invention being positioned in the immediate vicinity of the plants, but preferably not directly on the plants.

ABSTRACT

The present application is directed to a system and device for attraction of pests. This system comprises CO₂-releasing microorganisms, and also nutrients specific to these microorganisms, and this system, or the device, comprises one or more biodegradable biopolymers and allows CO₂, and optionally other attractants, to be released over a period of more than 20 days. Furthermore, the present application provides for the use of such a system or such a device for the attraction of pests, more particularly of maize or potato pests, such as larvae of the Western corn rootworm or wire worm. The present application is also directed, lastly, to methods for attracting pests, more particularly pests of maize or potatoes, such as the larvae of the Western corn rootworm, or for wire worms, with the systems or devices of the invention being positioned in the immediate vicinity of the plants, but preferably not directly on the plants.

Attraction systems for pests and use thereof

The present application is directed to a system and device for attracting pests. This system comprises CO₂-releasing microorganisms and also nutrients specific for these microorganisms, this system or device comprising one or more biodegradable biopolymers and allowing release of CO₂ and optionally other attractants over a period of more than 20 days.

The present application additionally provides for the use of such a system or device for attracting pests, more particularly corn or potato pests, such as, for example, larvae of the Western corn rootworm or wireworm. Lastly, the present application is also directed to methods of attracting pests, such as soil-dwelling pests, more particularly pests of corn or potatoes, such as the larvae of the Western corn rootworm, the black vine weevil, or for wireworms, the systems or devices of the invention being positioned in the immediate vicinity of the plants, but preferably not directly at the plants.

Prior Art

Carbon dioxide (CO₂) is a known attractant for various soil-dwelling organisms. A large number of pest insects make use of CO₂ and its concentration in order to locate the hosts. One important example of this is the Western corn rootworm (*Diabrotica virgifera virgifera*), whose larvae use CO₂ in order to locate the roots of living corn plants, in order then to use them as food, when they move in the direction of the ascending CO₂ gradient.

The destruction of the roots and the associated physiological stress on the plants by the feeding behavior of the larvae result in high losses in the case of these crop plants. Consequently, on a worldwide

WO 2014/041112

- 2 -

PCT/EP2013/068984

basis, the Western corn rootworm is an important pest, economically as well. There are studies from the early 2000s ascribing a loss of up to 1 billion U.S. dollars in the USA to damage by the larvae of the Western corn
5 rootworm.

In addition to various customary chemical or biological pesticides, attempts have been made to disrupt the location of the hosts by this pest - such location
10 taking place, as stated, through an ascending CO₂ gradient. For example, Bernklau E.A. et al., in Journal of Economic Entomology, 97(2), 330-339, 2004, describe methods and devices for attracting pests. There is a description of grains of baker's yeast and
15 corresponding nutrients therefor, with organic substrate as well, which then serve as a CO₂ source in order to keep the larvae of the Western corn rootworm away from the corn plant roots.

20 WO 01/32013 discloses methods and devices for attracting soil-dwelling insects. WO 03/028453 describes biodegradable solid preparations of a crop protection agent with delayed release of active ingredient. But a problem affecting the formulations
25 described in the prior art is that the CO₂ gradient and hence the activity of these systems are possible only for a short period; in the experiments, a period of 2 weeks at most was found for these formulations. In actual fact, however, effective control of the larvae
30 of the Western corn rootworm, and of other pests as well, such as wireworms in the case of potatoes, for example, requires a gradient over a longer period in order effectively to control the pests.

35 The aim of the present application is the provision of such systems and the use thereof in pest control, and also corresponding methods. In particular, the systems and devices of the invention permit control of soil

pests, more particularly corn and potato pests, such as the larvae of the Western corn rootworm or the wireworms, the larvae of beetles from the click beetle family. These larvae are known pests in forestry and agriculture, and in horticulture as well, and cause
5 considerable damage in both the softwood and hardwood sectors and also, in particular, in the potato sector.

Brief Description of the Drawings

Fig. 1 is a chart illustrating CO₂ release rate and CO₂ concentration in soil;

10 Fig. 2 is a chart illustrating CO₂ release rate and CO₂ concentration in soil; and

Fig. 3 is a chart illustrating number of wireworms per plant for different control methods.

Description of the Invention

15 Provided in accordance with the invention is a system or device for attracting pests. This system or device comprises CO₂-releasing microorganisms and nutrients specific for these microorganisms. A feature of the system or device is that it comprises one or more biodegradable polymers that surrounds, preferably completely, the
20 other constituents of the system, in the form of a capsule, for example, and permits release of CO₂ and optionally other attractants, released by the microorganisms or present additionally in the system, over a period of more than 20 days.

The release of the CO₂ is preferably over a period of more than
25 25 days, such as over 28 days or longer, as for example over 35 days or more.

The expressions "system" and "device" are presently used synonymously unless otherwise stated.

5 The expression "over a period of more than 20 days" means presently
that the CO₂-releasing microorganisms actively release CO₂ in the
system over a period of more than 20 days. This means more
particularly that the CO₂ concentration in the immediate
surroundings of the system is above the CO₂ concentration in a
region more remote from the system, thus forming a CO₂ gradient in
10 the direction of the system. This is so especially in the soil.

WO 2014/041112

- 4 -

PCT/EP2013/068984

In accordance with the invention, the system or device comprises particles including capsules, pellets, or granules, such as capsules, with at least the capsule shell or capsule matrix being formed by one or more biopolymers. The capsules in question may comprise those comprising biopolymers completely, in the form of pellets, for example. In one embodiment, these pellets or capsules may be fully surrounded by the biopolymer. Alternatively they may comprise granules, in which case the one or more biopolymers in the case of these granules surround - likewise preferably completely surround - the other constituents, or serve as a crosslinking and/or stabilizing matrix. Also possible are embodiments of the system of the invention wherein the biopolymers are interspersed through the particles, such as capsules, pellets, or granules, composed of the further constituents, and therefore provide the necessary stability to the system.

In particular, the CO₂-releasing microorganisms present in the system are fungi or bacteria. The microorganisms are preferably yeasts, including commercially available yeasts, such as baker's yeast, but also residual yeast from production of beer, wine, and bioethanol. One or more different microorganisms may be present, either likewise releasing CO₂ and/or having auxiliary functions.

In one embodiment an at least second microorganism, auxiliary organism, may be one having insecticidal or nematocidal activity. Suitable microorganisms are known to the skilled person. This at least second microorganism preferably comprises entomopathogenic microorganisms, such as entomopathogenic fungi or entomopathogenic nematodes. "Entomopathogenic" means presently that the organism possesses the capacity to

WO 2014/041112

- 5 -

PCT/EP2013/068984

infest, kill, immobilize, or otherwise negate the noxious effects of insects.

Of these, one preferred example of a second micro-
5 organism of this kind is the fungus *Beauveria bassiana*.
This fungus is known as a soil-dwelling fungus which
acts parasitically on various arthropod species and is
a known biological insecticide. A likewise-known
alternative to it is a fungus from the genus of the
10 *Metarhizia*, such as *M. anisopliae*.

The at least second microorganism, auxiliary organism,
may also preferably be one which is able to convert
nutrients present in the system in such a way that they
15 can be more effectively metabolized by the micro-
organisms responsible for CO₂ production. For example,
the fungus *Beauveria bassiana* or *M. anisopliae* allows
breakdown of starch, which is present as a specific
nutrient for the microorganism in the system. *Beauveria*
20 *bassiana* or *M. anisopliae* may therefore fulfill two
functions in this system - on the one hand, the
insecticidal or nematocidal activity as an entomo-
pathogenic fungus, and on the other hand the conversion
of nutrients in order to make them more effectively
25 metabolizable for the microorganisms responsible for CO₂
release, on the basis of its amylase activity, for
example.

The nutrients specific for these microorganisms are,
30 for example, nutrients selected from corn flour, corn
protein and/or other corn constituents, starch or
constituents containing starch, squash flour, potato
flour, or other polysaccharide-containing nutrients
suitable as nutrients, for example, for yeasts.

35

The skilled person knows of suitable nutrients for the
microorganisms present in the system.

WO 2014/041112

- 6 -

PCT/EP2013/068984

As already mentioned above, in a further embodiment, the system is one which additionally comprises a pesticide, more particularly an insecticide or nematocide. These pesticides may in one embodiment be
5 those from the group of the chemical insecticides, such as tefluthrin, those of plant extracts, such as neem oil or azadirachtin, quassia extract or the like, or else biological plant protection agents, such as the
aforementioned entomopathogenic fungi or entomo-
10 pathogenic nematodes.

It has been found that systems, more particularly capsules such as pellets, which comprise yeast, corn flour or other starch sources, and *Beauveria bassiana*
15 or *M. anisopliae*, the desired properties of the development of an appropriate CO₂ concentration which allows for the attraction of pests, such as the Western corn rootworm or wireworms over a period of more than
20 days.

20 The system may additionally have further constituents, such as fillers, drying assistants, glucose, sucrose, further attractants such as cucurbitacin or other known chemical attractants for the pests, and also cellulose,
25 lignin and/or swelling aids. The skilled person knows of suitable further constituents, which do not interact with microorganisms present in the system in such a way that these microorganisms present in the system are
killed or otherwise restricted in their function.

30 As stated, a feature of the systems of the invention is that they comprise one or more biodegradable biopolymers. These biopolymers are in a form, for example, in which they substantially, preferably fully,
35 surround the other constituents of the system. In the embodiment of capsules, accordingly, the biodegradable polymers form the capsule shell. In this case they are

WO 2014/041112

- 7 -

PCT/EP2013/068984

of a form such that exchange of nutrients, and more particularly emergence of CO₂, is possible.

5 In one alternative embodiment of a system of the invention, the biopolymers constitute the matrix. In that case they are of a form such that the biopolymers surround the other constituents preferably substantially, such as fully.

10 The biodegradable biopolymers are, in particular, those composed of alginate, carrageenan, cellulose, hemicellulose, starch, chitin, chitosan, pectinate, guar gum, acacia gum, poly-D,L-lactic acid, gelatin, poly-amino acids, lignin, and derivatives, and also
15 mixtures thereof. The skilled person knows of biopolymers that are appropriately suitable. Use may be made, for example, of alginate or of an alginate/gelatin mixture.

20 The biopolymers, which in particular are biodegradable, are those which are largely degraded in the soil preferably after the vegetation period or before the next vegetation period.

25 The present application is directed, furthermore, to the use of a system or device of the invention for attracting pests, more particularly corn pests or potato pests or, generally, pests of the plant constituents, such as the tuber-bearing stolons of the
30 potato, and the tubers themselves.

With particular preference the system or device of the invention is used in order to control corn pests or potato pests, here more particularly the Western corn
35 rootworm as a known corn pest, or the wireworm as a known potato pest. It is also used against the black vine weevil (in strawberries, for example). The system in this case is preferably one which also has

WO 2014/041112

- 8 -

PCT/EP2013/068984

insecticides or nematocides, such as *Beauveria bassiana*, for example, in order to kill the pests after attracting them. This approach is known generally in the art as "attract and kill" (alternatively: "lure and
5 kill"). The systems and devices of the invention may alternatively have active plant ingredients or active chemical ingredients for killing the pests.

The invention, furthermore, provides corresponding
10 methods for attracting and optionally controlling pests. This method is in particular for the control of pests of corn or potatoes, and in that context especially of those which attack the root region or the region located in the earth. The method is suitable
15 more particularly for attracting and controlling larvae of the Western corn rootworm, of the black vine weevil, or wireworms. The method of the invention is one which utilizes the systems and devices of the invention in order to attract the pests. The systems and devices, in
20 the form of capsules, for example, are positioned in the immediate vicinity of the plants, such as the corn or the potato, in the soil. It is preferred for this positioning to take place not directly at the plant, but rather at a predetermined distance in the vicinity
25 of the plant. The skilled person can determine the appropriate distance easily. The achievement thereby is that the pests are lured away from the plant. It is possible, for example, for the systems of the invention, in the form of capsules or granules, for
30 example, to be introduced between the sown seeds or seedlings. The systems in this case may take place simultaneously with the introduction of the corresponding plant seeds or plant seedlings. Alternatively or additionally the systems may be
35 introduced with appropriate measures in a subsequent period.

WO 2014/041112

- 9 -

PCT/EP2013/068984

The method of the invention is notable in particular in that it builds up an ascending CO₂ gradient from the plant in the direction of the system in the soil. As a result it is possible to attract the pests in question
5 via the gradient in the direction of the system and there, where appropriate, to kill them or otherwise immobilize them. The method is therefore suitable for attracting and killing the pests over a long period. In accordance with the invention, this period within which
10 the pests are attracted is at least 20 days, preferably at least 25 days, such as at least 30 days.

Particularly suitable embodiments are those in the form of capsules or pellets comprising a biopolymer or a
15 biopolymer mixture, at least one CO₂-releasing micro-organism, such as yeast, nutrients, such as starch-containing nutrients, auxiliary microorganisms, such as *B. bassiana* and/or *M. anisopliae*, and optionally further kill components such as neem or quassin.

20 One formulation of the system comprises biopolymer, yeast, corn flour or other flour, and *B. bassiana*. Another formulation of the system comprises biopolymer, yeast, corn flour or other flour, and *M. anisopliae*.

25 Further formulations for the system of the invention are those specified above and further comprising neem or quassin.

30 The systems and devices of the invention may include further attractants which attract the pests as well as or in addition to CO₂. More particularly these attractants are those which attract the pests in the direction of the system over a relatively short
35 distance, allowing them optionally to be killed there.

WO 2014/041112

- 10 -

PCT/EP2013/068984

In the text below, the system of the invention and the use in accordance with the invention are described in more detail, without being limited thereto.

5 **Examples:**

Example 1

Formulation of the system of the invention

10 *S. cerevisiae* is suspended in a predetermined amount, either as a cultivated strain or as commercially available baker's yeast mix, optionally with the other constituents, such as corn flour and *Beauveria*
15 *bassiana*, in 2% sodium alginate, introduced dropwise by means of a conventional encapsulating device into a 2% strength CaCl_2 solution, and subjected to postcrosslinking for 20 minutes. The capsules were produced with a diameter of 2.7 mm on average.

20 **Example 2**

Determining the CO_2 gradient and the duration of release

The capsules were placed in the earth in a pot in an amount of either 1 g or 10 g. CO_2 release rates and the
25 CO_2 concentration in the soil were measured using customary CO_2 meters. The CO_2 release rates were determined at the stated times. There were alginate capsules with no addition (active ingredient-free alginate capsules), alginate capsules with a fraction
30 of 16.7% baker's yeast mix, and alginate capsules with 16.7% baker's yeast mix, starch and, as an auxiliary microorganism exhibiting amylase activity, amylase source, in the present instance *Beauveria bassiana*.

35 Figure 1 shows the CO_2 release rate and the CO_2 concentration in the soil with the various capsules. Clearly evident are the increased release and the increased concentration even still after more than

WO 2014/041112

- 11 -

PCT/EP2013/068984

20 days. In figure 1, commercial baker's yeast was used, admixed with a starch and sugar mix with amylase activity.

5 In figure 2, the same measurements were carried out with pure *S. cerevisiae* and corn flour, and also *Beauveria bassiana*. The capsules used accordingly are shown. Particularly high release rates and CO₂ concentrations in the soil are observed for the
10 combination of yeast with starch as nutrient and also *B. bassiana* as amylase source.

Hence it is possible in accordance with the invention to provide systems and devices, in the form of
15 capsules, for example, which release CO₂ in the soil and so build up a gradient in order to attract pests and optionally to immobilize and/or kill them. This release takes place over a period of more than 20 days, such as more than 30 days, for example. This is a sufficient
20 period to keep the pests away from the plants.

Example 3:

Use of *M. anisopliae*

25 As observed in example 1, capsules of the invention are produced. Here, *B. bassiana* was replaced by *M. anisopliae*. The CO₂ gradient and the duration of CO₂ release were determined as in example 2.

30 It was found that the CO₂ gradient could be maintained over a period of more than 35 days. It was also found that the larvae of the corn rootworm were present to an increased extent around the capsules and in a smaller number around the plants.

35

It was apparent, moreover, that a reduction in wireworm density was achievable in the case of investigations with wireworms. This is clear from figure 3. CO₂ denotes

WO 2014/041112

- 12 -

PCT/EP2013/068984

capsules with yeast, CO₂ + MA denotes capsules with yeast and *M. anisopliae*, MA is *M. anisopliae* alone, and MAS denotes spores of the MA strain. ART2825 is the *M. anisopliae* strain used here.

5

Example 4:**Formulation of the system of the invention with plant-based pesticides**

10 As described in example 1, capsules are produced, with either Neem Azal or a quassin extract (Trifolio-M, Germany) being added instead of *B. bassiana*, in amounts of 1-10% w/w in the case of Neem Azal and 1-5% w/w for the quassin extract.

15

It was found that the added pesticides show no adverse effect on the yeast, and the yeast remained viable over a relatively long period. The build-up of the CO₂ gradient is comparable with the capsules described
20 before.

Claims:

1. A system for attracting pests, comprising CO₂-releasing microorganisms being yeast and nutrients specific for the CO₂-
5 releasing microorganisms; and
amylase or one or more auxiliary microorganisms with amylase activity being different from the CO₂-releasing microorganisms to support a supply of nutrients to the CO₂-releasing microorganisms
a biodegradable biopolymer which fully surrounds or
10 encapsulate the other constituents of the system and release CO₂ over a period of more than 20 days.
2. The system of claim 1, wherein the nutrients being selected from corn flour, corn protein, and other corn constituents, starch,
15 squash flour, potato flour.
3. The system of any one of claims 1 to 2, further comprising a pesticide.
- 20 4. The system of claim 3, wherein the pesticide is an insecticide or a nematicide.
5. The system of claim 4, wherein the insecticide or nematicide is at least one selected from the group of chemical insecticides,
25 plant extracts, and biological plant protection agents.
6. The system of claim 5, wherein at least one of the chemical insecticides is selected from tefluthrin, the plant extract is selected from neem oil or azadirachtin, quassia extract, and the
30 biological plant protection agent is selected from entomopathogenic fungi and entomopathogenic nematodes.

7. The system of any one of claims 1 to 6, further comprising as constituent fillers, drying assistants, glucose, sucrose, further attractants.

5

8. The system of claim 7, wherein the further attractants includes at least one of cucurbitacin, cellulose, lignin, and swelling aids.

10 9. The system of any one of claims 1 to 8, wherein the biodegradable biopolymers being selected from alginate, carrageenan, cellulose, hemicellulose, starch, chitin, chitosan, pectinate, guar gum, acacia gum, poly-D,L-lactic acid, gelatin, poly-amino acids, lignin, and derivatives, and also mixtures
15 thereof.

10. The system of any one of claims 1 to 9, wherein the system is provided in a form of capsules or granules.

20 11. The system of claim 1, wherein the auxiliary organisms are fungi.

12. The system of claim 11, wherein the fungi are *B. bassiana* or *M. anisopliae*.

25

13. The use of a system of any one of claims 1 to 12 for attracting pests including corn pests, potato pests, larvae of Western corn rootworm, larvae of wireworm, or larvae of black vine weevil.

14. A method for attracting pests including corn pests, potato pests, larvae of Western corn rootworm, larvae of wireworms, or larvae of black vine weevil, comprising:

5 providing a system of any one of claims 1 to 12; and
 positioning of said system in an immediate vicinity of plants including corn or potato.

15. The method of claim 14, wherein an ascending CO₂ gradient from the plant in a direction of the system is built up in soil.
10

16. The method of claim 14 or 15, wherein the system permits immobilization or killing of the larvae of the Western corn rootworm or of the wireworm or of the black vine weevil.




Figure 1




Figure 2




Figure 3

