



- (51) **International Patent Classification:**
G01N 33/00 (2006.01) G01N 33/574 (2006.01)
- (21) **International Application Number:**
PCT/US2014/022149
- (22) **International Filing Date:**
7 March 2014 (07.03.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/775,556 9 March 2013 (09.03.2013) US
- (72) **Inventors; and**
- (71) **Applicants :** STYLLI, Harry [GB/US]; 2452 Paseo Dorado, La Jolla, CA 92037 (US). KELLY, Colleen [US/US]; 7302 Golden Star Lane, Carlsbad, CA 92011 (US).
- (74) **Agents:** HALEY, James, F. et al.; Ropes & Gray LLP, 1211 Avenue of the Americas, New York, NY 10036 (US).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** METHODS OF DETECTING CANCER

Comparison		Macrophage/T-Cell	Neutrophil/T-Cell
Prostate cancer/Head and neck cancer	# of genes in signature	12	9
n=36, 8	average error	0.113	0.07
	sensitivity	75%	87.5%
	specificity	92.1%	94.4%
Prostate cancer/Head and neck cancer	# of genes in signature	10	9
n=58, 8	average error	0.112	0.085
	sensitivity	77.5%	88%
	specificity	90.7%	92.1%

Fig. 6

(57) **Abstract:** This invention provides a set of biological markers that are useful for detecting cancer. This invention further provides methods of using those biological markers for the diagnosis, prognosis, or monitoring of cancer.

1080340018W1

METHODS OF DETECTING CANCER

[0001] Related Application

[0002] This application claims priority and benefit from U.S. Provisional Patent
5 Application 61/775,556, filed March 9, 2013, the contents and disclosures of which
are incorporated herein by reference in their entirety.

[0003] Field of the Invention

[0004] This invention relates generally to using biological markers for the
diagnosis, prognosis, and monitoring of cancer.

10 [0005] Background of the Invention

[0006] Early diagnosis of a disease often increases the likelihood of successful
treatment or cure of such disease. Current diagnostic methods, however, depend
largely on population-derived average values obtained from healthy individuals.
Personalized diagnostic methods are needed that enable the diagnosis, especially
15 the early diagnosis, of the presence of a disease or a condition in individuals who
are not known to have the disease or who have recurrent disease. This is of
particular importance in cancer.

[0007] Leukocytes begin as pluripotent hematopoietic stem cells in the bone
marrow and develop along either the myeloid lineage (monocytes, macrophages,
20 neutrophils, eosinophils, and basophils) or the lymphoid lineage (T and B
lymphocytes and natural killer cells). The major function of the myeloid lineage
cells (e.g., neutrophils and macrophages) is the phagocytosis of infectious
organisms, live unwanted damaged cells, senescent and dead cells (apoptotic and
necrotic), as well as the clearing of cellular debris. Phagocytes from healthy
25 animals do not replicate and are diploid, i.e., have a DNA content of 2n. On
average, each cell contains <10 ng DNA, <20 ng RNA, and <300 ng of protein.
Non-phagocytic cells are also diploid and are not involved in the internalization of
dead cells or infectious organisms and have a DNA index of one.

[0008] The lifetime of various white blood cell subpopulations varies from a few
30 days (e.g., neutrophils) to several months (e.g., macrophages). Like other cell

types, leukocytes age and eventually die. During their aging process, human blood- and tissue-derived phagocytes (e.g., neutrophils) exhibit all the classic markers of programmed cell death (i.e., apoptosis), including caspase activation, pyknotic nuclei, and chromatin fragmentation. These cells also display a number of “eat-me” flags (e.g., phosphatidylserine, sugars) on the extracellular surfaces of their plasma membranes. Consequently, dying and dead cells and subcellular fragments thereof are cleared from tissues and blood by other phagocytic cells.

[0009] One object of the present invention is to provide diagnostic methods that can facilitate the detection of cancer markers, e.g., nucleic acids, proteins, and the like by using phagocytic cells alone, or in combination with non-phagocytic cells. Another object of this invention is to provide methods of identifying cancer markers and further use such markers alone or together with any known markers to diagnose cancer.

[0010] **Summary of the Invention**

[0011] In one aspect, the present invention provides methods for detecting or diagnosing cancer by using at least one or more markers selected from Tables 1- 8 (including C-Macro 1-200, C-Neutro 1-200, PC/HNC-Macro 1-200, and PC/HNC-Neutro 1-200). Levels (e.g., gene expression levels, protein expression levels, or activity levels) of the selected markers may be measured from macrophages or neutrophils, respectively, and from non-phagocytes, from a subject. Such levels then can be compared, e.g., the levels of the selected markers in the phagocytic cells and in the non-phagocytic cells to identify one or more differences between the measured levels, indicating whether the subject has cancer. The identified difference(s) can also be used for assessing the risk of developing cancer, prognosing cancer, monitoring cancer progression or regression, assessing the efficacy of a treatment for cancer, or identifying a compound capable of ameliorating or treating cancer.

[0012] In yet another aspect, the levels of the selected markers in the phagocytic cells may be compared to the levels of the selected markers in a control (e.g., a normal or healthy control subject, or a normal or healthy cell from the subject) to identify one or more differences between the measured levels, indicating whether the subject has cancer, the prognosis of the cancer and the monitoring of the

cancer. The identified difference(s) can also be used for assessing the risk of developing cancer, prognosing cancer, monitoring cancer progression or regression, assessing the efficacy of a treatment for cancer, or identifying a compound capable of ameliorating or treating cancer.

5 [0013] Some embodiments of this invention are as follows:

1. A method for diagnosing or aiding in the diagnosis of a cancer in a subject, the method comprising the steps of:
 - a) measuring the levels of one or more markers selected from the group
10 consisting of C-Macro 1-200 in a population of the subject's macrophage cells;
 - b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and
 - c) identifying a difference between the measured levels of the one or more selected C-Macro markers in steps a) and b),
15 wherein the identified difference indicates that the subject has said cancer.
2. A method for assessing the risk of developing a cancer in a subject, the method comprising the steps of:
 - a) measuring the levels of one or more markers selected from the group
20 consisting of C-Macro 1-200 in a population of the subject's macrophage cells;
 - b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and
 - c) identifying a difference between the measured levels of the one or more selected C-Macro markers in steps a) and b),
25 wherein the identified difference indicates that the subject has a risk of developing said cancer.
3. A method for prognosing or aiding in the prognosis of a cancer in a subject, the method comprising the steps of:
 - a) measuring the levels of one or more markers selected from the group
30 consisting of C-Macro 1-200 in a population of the subject's macrophage cells;

- b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and
- c) identifying a difference between the measured levels of the one or more selected C-Macro markers in steps a) and b),
- 5 wherein the identified difference is indicative of the prognosis of said cancer in the subject.
4. A method for assessing the efficacy of a treatment for a cancer in a subject comprising:
- 10 a) measuring the levels of one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells before the treatment;
- b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells before the treatment;
- 15 c) identifying a first difference between the measured levels of the one or more selected C-Macro markers in steps a) and b);
- d) measuring the levels of the one or more selected C-Macro markers in a population of the subject's macrophage cells after the treatment;
- e) measuring the levels of the one or more selected C-Macro markers in a
- 20 population of the subject's non-phagocytic cells after the treatment;
- f) identifying a second difference between the measured levels of the one or more selected C-Macro markers in steps d) and e); and
- g) identifying a difference between the first difference and the second difference,
- 25 wherein the difference identified in g) is indicative of the efficacy of the treatment for said cancer in the subject.
5. A method for monitoring the progression or regression of a cancer in a subject comprising:
- 30 a) measuring the levels of one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells at a first time point;

- b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells at the first time point;
- c) identifying a first difference between the measured levels of the one or more selected C-Macro markers in steps a) and b);
- 5 d) measuring the levels of the one or more selected C-Macro markers in a population of the subject's macrophage cells at a second time point;
- e) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells at the second time point;
- f) identifying a second difference between the measured levels of the one or more selected C-Macro markers in steps d) and e); and
- 10 g) identifying a difference between the first difference and the second difference,
- wherein the difference identified in g) is indicative of the progression or regression of said cancer in the subject.

15

6. A method for identifying a compound capable of ameliorating or treating a cancer in a subject comprising:
- a) measuring the levels of one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells
- 20 before administering the compound to the subject;
- b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells before administering the compound to the subject;
- c) identifying a first difference between the measured levels of the one or more selected C-Macro markers in steps a) and b);
- 25 d) measuring the levels of the one or more selected C-Macro markers in a population of the subject's macrophage cells after the administration of the compound;
- e) measuring the levels of the one or more selected C-Macro markers in a
- 30 population of the subject's non-phagocytic cells after the administration of the compound;

- f) identifying a second difference between the measured levels of the one or more selected C-Macro markers in steps d) and e); and
- g) identifying a difference between the first difference and the second difference,
- 5 wherein the difference identified in g) indicates that the compound is capable of ameliorating or treating said cancer in the subject.
7. A method for diagnosing or aiding in the diagnosis of a cancer in a subject, the method comprising the steps of:
- 10 a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells;
- b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells; and
- c) identifying a difference between the measured levels of the one or more
- 15 selected C-Neutro markers in steps a) and b),
- wherein the identified difference indicates that the subject has said cancer.
8. A method for assessing the risk of developing a cancer in a subject, the method comprising the steps of:
- 20 a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells;
- b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells; and
- c) identifying a difference between the measured levels of the one or more
- 25 selected C-Neutro markers in steps a) and b),
- wherein the identified difference indicates that the subject has a risk of developing said cancer.
9. A method for prognosing or aiding in the prognosis of a cancer in a
- 30 subject, the method comprising the steps of:
- a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells;

- b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells; and
- c) identifying a difference between the measured levels of the one or more selected C-Neutro markers in steps a) and b),
- 5 wherein the identified difference is indicative of the prognosis of said cancer in the subject.
10. A method for assessing the efficacy of a treatment for a cancer in a subject comprising:
- 10 a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells before the treatment;
- b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells before the treatment;
- 15 c) identifying a first difference between the measured levels of the one or more selected C-Neutro markers in steps a) and b);
- d) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's neutrophil cells after the treatment;
- e) measuring the levels of the one or more selected C-Neutro markers in a
- 20 population of the subject's non-phagocytic cells after the treatment;
- f) identifying a second difference between the measured levels of the one or more selected C-Neutro markers in steps d) and e); and
- g) identifying a difference between the first difference and the second difference,
- 25 wherein the difference identified in g) is indicative of the efficacy of the treatment for said cancer in the subject.
11. A method for monitoring the progression or regression of a cancer in a subject comprising:
- 30 a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells at a first time point;

- b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells at the first time point;
- c) identifying a first difference between the measured levels of the one or more selected C-Neutro markers in steps a) and b);
- 5 d) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's neutrophil cells at a second time point;
- e) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells at the second time point;
- f) identifying a second difference between the measured levels of the one or more selected C-Neutro markers in steps d) and e); and
- 10 g) identifying a difference between the first difference and the second difference,
- wherein the difference identified in g) is indicative of the progression or regression of said cancer in the subject.

15

12. A method for identifying a compound capable of ameliorating or treating a cancer in a subject comprising:

- a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells
- 20 before administering the compound to the subject;
- b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells before administering the compound to the subject;
- c) identifying a first difference between the measured levels of the one or more selected C-Neutro markers in steps a) and b);
- 25 d) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's neutrophil cells after the administration of the compound;
- e) measuring the levels of the one or more selected C-Neutro markers in a
- 30 population of the subject's non-phagocytic cells after the administration of the compound;

- f) identifying a second difference between the measured levels of the one or more selected C-Neutro markers in steps d) and e); and
- g) identifying a difference between the first difference and the second difference,
- 5 wherein the difference identified in g) indicates that the compound is capable of ameliorating or treating said cancer in the subject.

13. A method for diagnosing or aiding in the diagnosis of a cancer in a subject, the method comprising the steps of:
- 10 a) measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells, and
- measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil
- 15 cells;
- b) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and
- measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells;
- 20 c) identifying a difference between the measured levels of the at least one or more selected C-Macro markers in steps a) and b); and
- d) identifying a difference between the measured levels or activities the at least one or more selected C-Neutro markers in steps a) and b);
- wherein the differences identified in c) and d) indicate that the subject has
- 25 said cancer.

14. A method for assessing the risk of developing a cancer in a subject, the method comprising the steps of:
- a) measuring the levels of at least one or more markers selected from the
- 30 group consisting of C-Macro 1-200 in a population of the subject's macrophage cells, and

measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells;

- 5 b) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and
- measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells;
- c) identifying a difference between the measured levels of the at least one or more selected C-Macro markers in steps a) and b); and
- 10 d) identifying a difference between the measured levels of the at least one or more selected C-Neutro markers in steps a) and b);
- wherein the differences identified in c) and d) indicate that the subject has a risk of developing said cancer.

- 15 15. A method for prognosing or aiding in the prognosis of a cancer in a subject, the method comprising the steps of:

- a) measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells, and
- 20 measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells;
- b) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and
- 25 measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells;
- c) identifying a difference between the measured levels of the at least one or more selected C-Macro markers in steps a) and b); and
- d) identifying a difference between the measured levels of the at least one or more selected C-Neutro markers in steps a) and b);
- 30 wherein the differences identified in c) and d) are indicative of the prognosis of said cancer in the subject.

16. A method for assessing the efficacy of a treatment for a cancer in a subject comprising:
- a) measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells before the treatment, and
measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells before the treatment;
 - b) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells before the treatment; and
measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells before the treatment;
 - c) identifying a first difference between the measured levels of the at least one or more selected C-Macro markers in steps a) and b); and
identifying a second difference between the measured levels of the at least one or more selected C-Neutro markers in steps a) and b);
 - d) measuring the levels of the at least one or more selected C-Macro marker in a population of the subject's macrophage cells after the treatment, and
measuring the levels of the at least one or more selected C-Neutro marker in a population of the subject's neutrophil cells after the treatment;
 - e) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells after the treatment; and
measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells after the treatment;
 - f) identifying a third difference between the measured levels of the at least one or more selected C-Macro markers in steps d) and e); and
 - g) identifying a fourth difference between the measured levels of the at least one or more selected C-Neutro markers in steps d) and e);
 - h) identifying a difference between the first and second differences; and

i) identifying a difference between the third and fourth differences,
wherein the differences identified in h) and i) are indicative of the efficacy
of the treatment for said cancer in the subject.

5 17. A method for monitoring the progression or regression of a cancer in a
subject comprising:

 a) measuring the levels of at least one or more markers selected from the
group consisting of C-Macro 1-200 in a population of the subject's macrophage
cells at a first time point, and

10 measuring the levels of at least one or more markers selected from the
group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells
at the first time point;

 b) measuring the levels of the at least one or more selected C-Macro
markers in a population of the subject's non-phagocytic cells at the first time point;

15 and

 measuring the levels of the at least one or more selected C-Neutro markers
in a population of the subject's non-phagocytic cells at the first time point;

 c) identifying a first difference between the measured levels of the at least
one or more selected C-Macro markers in steps a) and b); and

20 identifying a second difference between the measured levels of the at least
one or more selected C-Neutro markers in steps a) and b);

 d) measuring the levels of the at least one or more selected C-Macro
markers in a population of the subject's macrophage cells at a second time point,
and

25 measuring the levels of the at least one or more selected C-Neutro markers
in a population of the subject's neutrophil cells at the second time point;

 e) measuring the levels of the at least one or more selected C-Macro
markers in a population of the subject's non-phagocytic cells at the second time
point; and

30 measuring the levels of the at least one or more selected C-Neutro markers
in a population of the subject's non-phagocytic cells at the second time point;

- f) identifying a third difference between the measured levels of the at least one or more selected C-Macro markers in steps d) and e); and
- g) identifying a fourth difference between the measured levels of the at least one or more selected C-Neutro markers in steps d) and e);
- 5 h) identifying a difference between the first and second differences; and
- i) identifying a difference between the third and fourth differences,
- wherein the differences identified in h) and i) are indicative of the progression or regression of said cancer in the subject.
- 10 18. A method for identifying a compound capable of ameliorating or treating a cancer in a subject comprising:
- a) measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells before administering the compound to the subject, and
- 15 measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells before administering the compound to the subject;
- b) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells before administering
- 20 the compound to the subject; and
- measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells before administering the compound to the subject;
- c) identifying a first difference between the measured levels of the at least
- 25 one or more selected C-Macro markers in steps a) and b); and
- identifying a second difference between the measured levels of the at least one or more selected C-Neutro markers in steps a) and b);
- d) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's macrophage cells after administering the
- 30 compound to the subject, and

- measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's neutrophil cells after administering the compound to the subject;
- e) measuring the levels of the at least one or more selected C-Macro
5 markers in a population of the subject's non-phagocytic cells after administering the compound to the subject; and
- measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells after administering the compound to the subject;
- 10 f) identifying a third difference between the measured levels of the at least one or more selected C-Macro markers in steps d) and e); and
- g) identifying a fourth difference between the measured levels of the at least one or more selected C-Neutro markers in steps d) and e);
- h) identifying a difference between the first and second differences; and
15 i) identifying a difference between the third and fourth differences,
- wherein the differences identified in h) and i) indicate that the compound is capable of ameliorating or treating said cancer in the subject.
19. The method of any one of the embodiments 1-18, further comprising
20 measuring at least one standard parameter associated with said cancer.
20. The method of embodiment 19, wherein the standard parameter is selected from the group consisting of tumor stage, tumor grade, tumor size, tumor visual characteristics, tumor growth, and tumor thickness, tumor progression, tumor
25 metastasis, tumor distribution within the body, odor, molecular pathology, genomics, tumor angiograms, or Gleason score.
21. The method of any one of the embodiments 13-18, wherein the selected C-Macro markers and the selected C-Neutro markers are measured from the same
30 population of non-phagocytic cells in steps b) or e).

22. The method of any one of the embodiments 13-18, wherein the selected C-Macro markers and the selected C-Neutro are from different populations of non-phagocytic cells in steps b) or e).
- 5
23. The method of any one of the embodiments 1-6 and 13-18, wherein at least two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty, twenty-one, twenty-two, twenty-three, twenty-four, or twenty-five markers are selected from C-Macro 1-200.
- 10
24. The method of any one of the embodiments 1-6 and 13-18, wherein the selected C-Macro markers comprise one or more markers selected from the group consisting of C-Macro 1-4 and C-Macro 101-104.
- 15
25. The method of any one of the embodiments 1-6 and 13-18, wherein the selected C-Macro markers are up-regulated or activated in the macrophage cells compared to the non-phagocytic cells.
- 20
26. The method of any one of the embodiments 1-6 and 13-18, wherein the selected C-Macro markers are up-regulated or activated in the macrophage cells compared to the non-phagocytic cells.
- 25
27. The method of any one of the embodiments 1-6 and 13-18, wherein the selected C-Macro markers are down-regulated or inhibited in the macrophage cells compared to the non-phagocytic cells.
- 30
28. The method of any one of the embodiments 1-6 and 13-18, wherein the selected C-Macro markers are down-regulated or inhibited in the macrophage cells compared to the non-phagocytic cells.

29. The method of any one of the embodiments 7-18, wherein at least two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty, twenty-one, twenty-two, twenty-three, twenty-four, or twenty-five markers are selected from C-Neutro 1-
5 200.
30. The method of any one of the embodiments 7-18, wherein the selected C-Neutro markers comprise one or more C-Neutro markers selected from the group consisting of C-Neutro 1-8 and C-Neutro 101-108.
10
31. The method of any one of the embodiments 7-18, wherein the selected C-Neutro markers comprise one or more markers selected from the group consisting of C-Neutro 1-200 and wherein the selected C-Neutro markers are down-regulated or inhibited in the neutrophil cells compared to the non-phagocytic cells.
15
32. The method of any one of the embodiments 7-18, wherein the selected C-Neutro markers are down-regulated or inhibited in the neutrophil cells compared to the non-phagocytic cells.
- 20 33. The method of any one of the embodiments 1-6 and 13-18, further comprising lysing the macrophage cells and the non-phagocytic cells before a).
34. The method of any one of the embodiments 1-6 and 13-18, further comprising extracting the cellular contents from the macrophage cells and the non-
25 phagocytic cells before a).
35. The method of any one of the embodiments 7-18, further comprising lysing the neutrophil cells and the non-phagocytic cells before a).
- 30 36. The method of any one of the embodiments 7-18, further comprising extracting the cellular contents from the neutrophil cells and the non-phagocytic cells before a).

37. The method of embodiment 34, wherein the cellular contents of the macrophage cells comprise viable diseased cells, dead diseased cells, apoptotic diseased cells, circulating tumor cells, infectious agents, fetal cells, trophoblasts, or
5 fragments thereof.

38. The method of embodiment 36, wherein the cellular contents of the neutrophil cells comprise viable diseased cells, dead diseased cells, apoptotic diseased cells, circulating tumor cells, infectious agents, fetal cells, trophoblasts, or
10 fragments thereof.

39. The method of embodiment 34, wherein the selected one or more markers are present in the cellular contents of the macrophage cells.

15 40. The method of embodiment 34, wherein the selected one or more markers are not present in the cellular contents of the non-phagocytic cells.

41. The method of any one of the embodiments 1-6 and 13-18, wherein the macrophage cells express the one or more selected C-Macro markers.
20

42. The method of embodiment 36, wherein the selected one or more markers are present in the cellular contents of the neutrophil cells.

43. The method of embodiment 36, wherein the selected one or more markers
25 are not present in the cellular contents of the non-phagocytic cells.

44. The method of any one of the embodiments 7-18, wherein the neutrophil cells express the one or more selected C-Neutro markers.

30 45. The method of any one of the embodiments 1-18, wherein the non-phagocytic cells are T cells, B cells, null cells, basophils, or mixtures thereof.

46. The method of any one of the embodiments 1-6 and 13-18, wherein the macrophage cells are isolated from a bodily fluid sample, tissues, or cells of the subject.
- 5 47. The method of any one of the embodiments 7-18, wherein the neutrophil cells are isolated from a bodily fluid sample, tissues, or cells of the subject.
48. The method of any one of the embodiments 1-18, wherein the non-phagocytic cells are isolated from a bodily fluid sample, tissues, or cells of the
10 subject.
49. The method of any one of the embodiments 46-48, wherein the bodily fluid sample is blood, urine, stool, saliva, lymph fluid, cerebrospinal fluid, synovial fluid, cystic fluid, ascites, pleural effusion, fluid obtained from a pregnant woman
15 in the first trimester, fluid obtained from a pregnant woman in the second trimester, fluid obtained from a pregnant woman in the third trimester, maternal blood, amniotic fluid, chorionic villus sample, fluid from a preimplantation embryo, maternal urine, maternal saliva, placental sample, fetal blood, lavage and cervical vaginal fluid, interstitial fluid, or ocular fluid.
- 20 50. The method of any one of the embodiments 1-6 and 13-18, wherein the macrophage cells are isolated using antibodies, using a ligand that binds to a molecular receptor expressed on the plasma membranes of white blood cells, or by flow cytometry, fluorescence activated cell sorting, filtration, gradient-based
25 centrifugation, elution, microfluidics, magnetic separation technique, fluorescent-magnetic separation technique, nanostructure, quantum dots, high throughput microscope-based platforms, or a combination thereof.
51. The method of any one of the embodiments 7-18, wherein the neutrophil
30 cells are isolated using antibodies, using a ligand that binds to a molecular receptor expressed on the plasma membranes of white blood cells, or by flow cytometry, fluorescence activated cell sorting, filtration, gradient-based centrifugation, elution,

microfluidics, magnetic separation technique, fluorescent-magnetic separation technique, nanostructure, quantum dots, high throughput microscope-based platforms, or a combination thereof.

- 5 52. The method of any one of the embodiments 1-18, wherein the non-phagocytic cells are isolated using antibodies, using a ligand that binds to a molecular receptor expressed on the plasma membranes of white blood cells, or by flow cytometry, fluorescence activated cell sorting, filtration, gradient-based centrifugation, elution, microfluidics, magnetic separation technique, fluorescent-
10 magnetic separation technique, nanostructure, quantum dots, high throughput microscope-based platforms, or a combination thereof.

53. The method of any one of the embodiments 1-6 and 13-18, wherein the macrophage cells are isolated using a product secreted by the macrophage cells.
15

54. The method of any one of the embodiments 7-18, wherein the neutrophil cells are isolated by using a product secreted by the neutrophil cells.

55. The method of any one the embodiments 1-6 and 13-18, wherein the
20 macrophage cells are isolated by using a cell surface target on the surface of macrophage cells.

56. The method of any one of the embodiments 7-18, wherein the neutrophil cells are isolated by using a cell surface target on the surface of neutrophil cells.
25

57. The method of embodiment 55, wherein the target is expressed by the macrophage cells.

58. The method of embodiment 55, wherein the target is not expressed by the
30 macrophage cells.

59. The method of embodiment 56, wherein the target is expressed by the neutrophil cells.
60. The method of embodiment 56, wherein the target is not expressed by the neutrophil cells.
61. The method of any one of the embodiments 55-60, wherein the target is a marker of said cancer.
62. The method of any one of the embodiments 1-18, wherein the measured levels are gene expression levels.
63. The method of any one of the embodiments 1-18, wherein the measured levels are protein expression levels.
64. The method of any one of the embodiment 1-18, wherein the levels or activities are measured by a qualitative assay, a quantitative assay, or a combination thereof.
65. The method of embodiment 64, wherein the quantitative assay uses sequencing, direct sequencing, RNA sequencing, whole transcriptome shotgun sequencing, random shotgun sequencing, Sanger dideoxy termination sequencing, whole-genome sequencing, sequencing by hybridization, pyrosequencing, capillary electrophoresis, gel electrophoresis, duplex sequencing, cycle sequencing, single-base extension sequencing, solid-phase sequencing, high-throughput sequencing, massively parallel signature sequencing, emulsion PCR, sequencing by reversible dye terminator, paired-end sequencing, near-term sequencing, exonuclease sequencing, sequencing by ligation, short-read sequencing, single-molecule sequencing, sequencing-by-synthesis, real-time sequencing, reverse-terminator sequencing, nanopore sequencing, 454 sequencing, Solexa Genome Analyzer sequencing, SOLiD® sequencing, MS-PET sequencing, mass spectrometry, matrix assisted laser desorption/ionization-time of flight (MALDI-TOF) mass

- spectrometry, electrospray ionization (ESI) mass spectrometry, surface-enhanced laser desorption/ionization-time of flight (SELDI-TOF) mass spectrometry, quadrupole-time of flight (Q-TOF) mass spectrometry, atmospheric pressure photoionization mass spectrometry (APPI-MS), Fourier transform mass spectrometry (FTMS), matrix-assisted laser desorption/ionization-Fourier transform-ion cyclotron resonance (MALDI-FT-ICR) mass spectrometry, secondary ion mass spectrometry (SIMS), polymerase chain reaction (PCR) analysis, quantitative PCR, real-time PCR, fluorescence assay, colorimetric assay, chemiluminescent assay, or a combination thereof.
66. The method of embodiment 62, wherein the gene expression levels are measured by polymerase chain reaction (PCR) analysis, sequencing analysis, electrophoretic analysis, restriction fragment length polymorphism (RFLP) analysis, Northern blot analysis, quantitative PCR, reverse-transcriptase-PCR analysis (RT-PCR), allele-specific oligonucleotide hybridization analysis, comparative genomic hybridization, heteroduplex mobility assay (HMA), single strand conformational polymorphism (SSCP), denaturing gradient gel electrophoresis (DGGE), RNAase mismatch analysis, mass spectrometry, tandem mass spectrometry, matrix assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry, electrospray ionization (ESI) mass spectrometry, surface-enhanced laser desorption/ionization-time of flight (SELDI-TOF) mass spectrometry, quadrupole-time of flight (Q-TOF) mass spectrometry, atmospheric pressure photoionization mass spectrometry (APPI-MS), Fourier transform mass spectrometry (FTMS), matrix-assisted laser desorption/ionization-Fourier transform-ion cyclotron resonance (MALDI-FT-ICR) mass spectrometry, secondary ion mass spectrometry (SIMS), surface plasmon resonance, Southern blot analysis, in situ hybridization, fluorescence in situ hybridization (FISH), chromogenic in situ hybridization (CISH), immunohistochemistry (IHC), microarray, comparative genomic hybridization, karyotyping, multiplex ligation-dependent probe amplification (MLPA), Quantitative Multiplex PCR of Short Fluorescent Fragments (QMPSF), microscopy, methylation specific PCR (MSP) assay, HpaII tiny fragment Enrichment by Ligation-mediated PCR (HELP) assay,

radioactive acetate labeling assays, colorimetric DNA acetylation assay, chromatin immunoprecipitation combined with microarray (ChIP-on-chip) assay, restriction landmark genomic scanning, Methylated DNA immunoprecipitation (MeDIP), molecular break light assay for DNA adenine methyltransferase activity,
5 chromatographic separation, methylation-sensitive restriction enzyme analysis, bisulfite-driven conversion of non-methylated cytosine to uracil, methyl-binding PCR analysis, or a combination thereof.

67. The method of embodiment 62, wherein the gene expression levels are
10 measured by a sequencing technique selected from the group consisting of direct sequencing, RNA sequencing, whole transcriptome shotgun sequencing, random shotgun sequencing, Sanger dideoxy termination sequencing, whole-genome sequencing, sequencing by hybridization, pyrosequencing, capillary electrophoresis, gel electrophoresis, duplex sequencing, cycle sequencing, single-
15 base extension sequencing, solid-phase sequencing, high-throughput sequencing, massively parallel signature sequencing, emulsion PCR, sequencing by reversible dye terminator, paired-end sequencing, near-term sequencing, exonuclease sequencing, sequencing by ligation, short-read sequencing, single-molecule sequencing, sequencing-by-synthesis, real-time sequencing, reverse-terminator
20 sequencing, nanopore sequencing, 454 sequencing, Solexa Genome Analyzer sequencing, SOLiD® sequencing, MS-PET sequencing, mass spectrometry, and a combination thereof.

68. The method of embodiment 63, wherein the protein expression levels are
25 measured by an immunohistochemistry assay, an enzyme-linked immunosorbent assay (ELISA), in situ hybridization, chromatography, liquid chromatography, size exclusion chromatography, high performance liquid chromatography (HPLC), gas chromatography, mass spectrometry, tandem mass spectrometry, matrix assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry,
30 electrospray ionization (ESI) mass spectrometry, surface-enhanced laser desorption/ionization-time of flight (SELDI-TOF) mass spectrometry, quadrupole-time of flight (Q-TOF) mass spectrometry, atmospheric pressure photoionization

mass spectrometry (APPI-MS), Fourier transform mass spectrometry (FTMS),
matrix-assisted laser desorption/ionization-Fourier transform-ion cyclotron
resonance (MALDI-FT-ICR) mass spectrometry, secondary ion mass spectrometry
(SIMS), radioimmunoassays, microscopy, microfluidic chip-based assays, surface
5 plasmon resonance, sequencing, Western blotting assay, or a combination thereof.

69. The method of any one the embodiments 1-68, wherein the subject is a mammal.
- 10 70. The method of embodiment 69, wherein the subject is a human.
71. The method of any one the embodiments 1-18, wherein the difference is greater than a 1-fold difference.
- 15 72. The method of embodiment 71, wherein the difference is at least 1.05-fold, 1.1-fold, 1.2-fold, 1.3-fold, 1.4-fold, 1.5-fold, 2-fold, 2.5-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, or 10-fold difference.
73. A kit for measuring the levels of at least one or more markers selected
20 from the group consisting of C-Macro 1-200, comprising reagents for specifically measuring the levels of the selected C-Macro marker.
74. A kit for measuring the levels of at least one or more markers selected
from the group consisting of C-Neutro 1-200, comprising reagents for specifically
25 measuring the levels of the selected C-Neutro marker.
75. A kit for measuring the levels of at least one or more markers selected
from the group consisting of C-Macro 1-200 and at least one or more markers
selected from the group consisting of C-Neutro 1-200, comprising reagents for
30 specifically measuring the levels of the selected C-Macro marker and reagents for specifically measuring the levels of the selected C-Neutro marker.

76. The kit of embodiment 73 or 75, wherein the selected C-Macro markers comprise one or more markers selected from the group consisting of C-Macro 1-4 and C-Macro 101-104.
- 5 77. The kit of embodiment 74 or 75, wherein the selected C-Neutro markers comprise one or more markers selected from the group consisting of C-Neutro 1-8 and C-Neutro 101-108.
78. The kit of any one of the embodiments 73-77, wherein the reagents
10 comprise one or more antibodies or fragments thereof, oligonucleotides, or aptamers.
79. A method of treating or preventing a cancer in a subject comprising administering to said subject an agent that modulates the activity or expression of
15 at least one or more markers selected from the group consisting of C-Macro 1-200.
80. A method of treating or preventing a cancer in a subject comprising administering to said subject an agent that modulates the activity or expression of at least one or more markers selected from the group consisting of C-Neutro 1-200.
20
81. The method of embodiment 79 or 80, wherein the agent is a small molecule modulator, siRNA, or an antibody or fragment thereof.
82. The method of any one of the above embodiments, wherein the cancer is a
25 prostate cancer, a head and neck cancer, a lung cancer, melanoma, a colon cancer, a non-small cell lung cancer, a CNS cancer, an ovarian cancer, a renal cancer, or a breast cancer.
- 30

[0014] Brief Description of the Drawings

[0015] Figure 1 depicts a diagram of a 3-fold cross validation method.

[0016] Figure 2 depicts a summary of cancer markers identified from macrophage vs. T cell and neutrophil vs. T cell comparisons in prostate cancer patients and head and neck cancer patients.

[0017] Figure 3 depicts a comparison of cancer detection using markers identified from macrophages and neutrophils vs. T cells, as compared to detection when the phagocyte gene expression is not compared to T cell gene expression.

[0018] Figure 4 depicts a purification method for validating methods of detecting cancer.

[0019] Figure 5 depicts a comparison of purification methods in validating a method of detecting cancer.

[0020] Figure 6 depicts a summary of cancer markers distinguishing between prostate cancer and head and neck cancer identified from macrophage vs. T cell and neutrophil vs. T cell comparisons.

[0021] Detailed Description of the Invention

[0022] The present invention provides biological markers and methods of using them to detect a cancer. More specifically, the present invention provides biomarkers that are specific for cancers.

[0023] As used here in, a “biomarker” or “marker” refers to an analyte (e.g., a nucleic acid, DNA, RNA, peptide, protein, or metabolite) that can be objectively measured and evaluated as an indicator for a biological process. In some embodiments, a marker is differentially detectable in phagocytes and is indicative of the presence or absence of cancer. An analyte is differentially detectable if it can be distinguished quantitatively or qualitatively in phagocytes compared to a control, e.g., a normal or healthy control or non-phagocytic cells.

[0024] The present invention is based on the discovery that one or more markers selected from Tables 1-8 are useful in diagnosing cancer. By measuring the levels of the biomarkers (e.g., gene expression levels, protein expression levels, or protein activity levels) in a population of phagocytes (e.g., macrophage or neutrophils) from a human subject, one can provide a reliable diagnosis for cancer.

[0025] As used herein, a “level” of a marker of this invention can be qualitative (e.g., presence or absence) or quantitative (e.g., amounts, copy numbers, or dosages). In some embodiments, a level of a marker at a zero value can indicate the absence of this marker. The levels of any marker of this invention can be
5 measured in various forms. For example, the level can be a gene expression level, a RNA transcript level, a protein expression level, a protein activity level, an enzymatic activity level.

[0026] The markers of this invention can be used in methods for diagnosing or aiding in the diagnosis of cancer by comparing levels (e.g., gene expression levels,
10 or protein expression levels, or protein activities) of one or more cancer markers (e.g., nucleic acids or proteins) between phagocytes (e.g., macrophages or neutrophils) and non-phagocytic cells taken from the same individual. This invention also provides methods for assessing the risk of developing cancer, prognosing said cancer, monitoring said cancer progression or regression,
15 assessing the efficacy of a treatment, or identifying a compound capable of ameliorating or treating said cancer.

[0027] Tables 1-8 provide the information for the markers of this invention, such as transcript cluster ID, gene names, and pattern of regulation. The skilled worker would readily identify the gene or protein name for the markers listed in Tables 1-
20 8, based on at least the transcript cluster ID information provided.

[0028] In a first aspect, the methods (e.g., diagnosis of cancer, prognosis of cancer, or assessing the risk of developing cancer) provided in the invention comprise: a) measuring the levels of one or more markers selected from Tables 1, 3, 5, and 7 (C-Macro markers 1-200 and PC/HNC-Macro markers 1-200) in a
25 population of a subject’s macrophage cells; b) measuring the levels of one or more of the selected markers in a population of a subject’s non-phagocytic cells (e.g., T-cells, B-cells, null cells, basophils or the mixtures of two more non-phagocytic cells); comparing the measured levels in step a) to the measured levels in step b) and further identify a difference between the measured levels of a) and b). The
30 identified difference is indicative of the diagnosis (e.g., presence or absence), prognosis (e.g., lethal outcome, or tumor stage), or the risk of developing cancer.

[0029] In a second aspect, the methods (e.g., diagnosis of cancer, prognosis of cancer, or assessing the risk of developing cancer) provided in the invention comprise: a) measuring the levels of one or more markers selected from Tables 1, 3, 5, and 7 (C-Macro markers 1-200 and PC/HNC-Macro markers 1-200) in a population of a subject's macrophage cells; identifying a difference between the measured levels of the selected markers in step a) and the levels of the selected markers in a control (e.g., a healthy control cell, or a control cell from a healthy subject). The identified difference is indicative of the diagnosis (e.g., presence or absence), prognosis (e.g., lethal outcome, or tumor stage), or the risk of developing cancer.

[0030] In a third aspect, the methods (e.g., diagnosis of cancer, prognosis of cancer, or assessing the risk of developing cancer) provided in the invention comprise: a) measuring the levels of one or more markers selected from Tables 2, 4, 6, and 8 (C-Neutro 1-200 and PC/HNC-Neutro 1-200) in a population of a subject's neutrophil cells; b) measuring the levels of one or more of the selected markers in a population of a subject's non-phagocytic cells (e.g., T-cells, B-cells, null cells, basophils or the mixtures of two more non-phagocytic cells); comparing the measured levels in step a) to the measured levels in step b) and further identify a difference between the measured levels of a) and b). The identified difference is indicative of the diagnosis (e.g., presence or absence), prognosis (e.g., lethal outcome, or tumor stage), or the risk of developing cancer.

[0031] In a fourth aspect, the methods (e.g., diagnosis of cancer, prognosis of cancer, or assessing the risk of developing cancer) provided in the invention comprise: a) measuring the levels of one or more markers selected from Tables 2, 4, 6, and 8 (C-Neutro 1-200 and PC/HNC-Neutro 1-200) in a population of a subject's neutrophil cells; identifying a difference between the measured levels of the selected markers in step a) and the levels of the selected markers in a control (e.g., a healthy control cell, or a control cell from a healthy subject). The identified difference is indicative of the diagnosis (e.g., presence or absence), prognosis (e.g., lethal outcome, or tumor stage), or the risk of developing cancer.

[0032] In a fifth aspect, the methods (e.g., diagnosis of cancer, prognosis of cancer, or assessing the risk of developing cancer) provided in the invention

- comprise: a) measuring the levels of one or more markers selected from Tables 2, 4, 6, and 8 (C-Neutro 1-200 and PC/HNC-Neutro 1-200) in a population of a subject's neutrophil cells and the levels of one or more markers selected from Tables 1, 3, 5, and 7 (C-Macro markers 1-200 and PC/HNC-Macro markers 1-200) in a population of a subject's macrophage cells; b) measuring the levels of one or more of the selected C-Macro (or PC/HNC-Macro) markers and the levels one or more of the selected C-Neutro (or PC/HNC-Neutro) markers in a population of a subject's non-phagocytic cells (e.g., T-cells, B-cells, null cells, basophils or the mixtures of two more non-phagocytic cells); identifying a difference between the measured levels of the selected C-Neutro (or PC/HNC-Neutro) markers of steps a) and b) and identifying a difference between the measured levels of the selected C-Macro (or PC/HNC-Macro) markers of steps a) and b). The identified differences are indicative of the diagnosis (e.g., presence or absence), prognosis (e.g., lethal outcome, or tumor stage), or the risk of developing cancer.
- 15 **[0033]** In a sixth aspect, the methods (e.g., diagnosis of cancer, prognosis of cancer, or assessing the risk of developing cancer) provided in the invention comprise: a) measuring the levels of one or more markers selected from Tables 2, 4, 6, and 8 (C-Neutro 1-200 and PC/HNC-Neutro 1-200) in a population of a subject's neutrophil cells and the levels of one or more markers selected from Tables 1, 3, 5, and 7 (C-Macro markers 1-200 and PC/HNC-Macro markers 1-200) in a population of a subject's macrophage cells; identifying a difference between the measured levels of the selected C-Neutro (or PC/HNC-Neutro) markers of steps a) and the levels of the selected C-Neutro (or PC/HNC-Neutro) markers in a control (e.g., a healthy control cell, or a control cell from a healthy subject) and identifying a difference between the measured levels of the selected C-Macro (or PC/HNC-Macro) markers of step a) and the levels of the selected C-Macro (or PC/HNC-Macro) markers in a control (e.g., a healthy control cell, or a control cell from a healthy subject). The identified differences are indicative of the diagnosis (e.g., presence or absence), prognosis (e.g., lethal outcome, or tumor stage), or the risk of developing cancer.
- 30 **[0034]** In a seventh aspect, the methods provided in this invention for assessing the efficacy of a treatment for cancer, monitoring the progression or regression of

cancer, or identifying a compound capable of ameliorating or treating cancer, respectively, in a subject comprising: a) measuring the levels of one or more markers selected from the group consisting of Tables 1, 3, 5, and 7 (C-Macro markers 1-200 and PC/HNC-Macro markers 1-200) in a population of the subject's macrophage cells before the treatment, or at a first time point, or before administration of the compound, respectively; b) measuring the levels of the one or more selected C-Macro (or PC/HNC-Macro) markers in a population of the subject's non-phagocytic cells before the treatment, or at the first time point, or before administration of the compound, respectively; c) identifying a first difference between the measured levels of the one or more selected C-Macro (or PC/HNC-Macro) markers in steps a) and b); d) measuring the levels of the one or more selected C-Macro (or PC/HNC-Macro) markers in a population of the subject's macrophage cells after the treatment, or at a second time point, or after administration of the compound, respectively; e) measuring the levels of the one or more selected C-Macro (or PC/HNC-Macro) markers in a population of the subject's non-phagocytic cells after the treatment, or at the second time point, or after administration of the compound, respectively; f) identifying a second difference between the measured levels of the one or more selected C-Macro (or PC/HNC-Macro) markers in steps d) and e); and g) identifying a difference between the first difference and the second difference, wherein the difference identified in g) is indicative of the efficacy of the treatment for the cancer, or the progression or regression of the cancer, or whether the compound is capable of ameliorating or treating the cancer, respectively, in the subject.

[0035] In a eighth aspect, the methods provided in this invention for assessing the efficacy of a treatment for cancer, monitoring the progression or regression of cancer, or identifying a compound capable of ameliorating or treating cancer, respectively, in a subject comprising: a) measuring the levels of one or more markers selected from the group consisting of Tables 1, 3, 5, and 7 (C-Macro markers 1-200 and PC/HNC-Macro markers 1-200) in a population of the subject's macrophage cells before the treatment, or at a first time point, or before administration of the compound, respectively; b) identifying a first difference between the measured levels of the one or more selected C-Macro (or PC/HNC-

Macro) markers in step (a) and the levels of the one or more selected C-Macro (or PC/HNC-Macro) markers in a control (e.g., a healthy control cell, or a control cell from a healthy subject) before the treatment, or at the first time point, or before administration of the compound, respectively; c) measuring the levels of the one or more selected C-Macro (or PC/HNC-Macro) markers in a population of the subject's macrophage cells after the treatment, or at a second time point, or after administration of the compound, respectively; d) identifying a second difference between the measured levels of the one or more selected C-Macro (or PC/HNC-Macro) markers in step c) and the levels of the one or more selected C-Macro (or PC/HNC-Macro) markers in a control after the treatment, or at the second time point, or after administration of the compound, respectively; and e) identifying a difference between the first difference and the second difference, wherein the difference identified in e) is indicative of the efficacy of the treatment for the cancer, or the progression or regression of the cancer, or whether the compound is capable of ameliorating or treating the cancer, respectively, in the subject.

[0036] In a ninth aspect, the methods provided in this invention for assessing the efficacy of a treatment for cancer, monitoring the progression or regression of cancer, or identifying a compound capable of ameliorating or treating cancer, respectively, in a subject comprising: a) measuring the levels of one or more markers selected from the group consisting of Tables 2, 4, 6, and 8 (C-Neutro 1-200 and PC/HNC-Neutro 1-200) in a population of the subject's neutrophil cells before the treatment, or at a first time point, or before administration of the compound, respectively; b) measuring the levels of the one or more selected C-Neutro (or PC/HNC-Neutro) markers in a population of the subject's non-phagocytic cells before the treatment, or at the first time point, or before administration of the compound, respectively; c) identifying a first difference between the measured levels of the one or more selected C-Neutro (or PC/HNC-Neutro) markers in steps a) and b); d) measuring the levels of the one or more selected C-Neutro (or PC/HNC-Neutro) markers in a population of the subject's neutrophil cells after the treatment, or at a second time point, or after administration of the compound, respectively; e) measuring the levels of the one or more selected C-Neutro (or PC/HNC-Neutro) markers in a population of the

subject's non-phagocytic cells after the treatment, or at the second time point, or after administration of the compound, respectively; f) identifying a second difference between the measured levels of the one or more selected C-Neutro (or PC/HNC-Neutro) markers in steps d) and e); and g) identifying a difference

5 between the first difference and the second difference, wherein the difference identified in g) is indicative of the efficacy of the treatment for the cancer, or the progression or regression of the cancer, or whether the compound is capable of ameliorating or treating the cancer, respectively, in the subject.

[0037] In a tenth aspect, the methods provided in this invention for assessing the

10 efficacy of a treatment for cancer, monitoring the progression or regression of cancer, or identifying a compound capable of ameliorating or treating cancer, respectively, in a subject comprising: a) measuring the levels of one or more markers selected from the group consisting of Tables 2, 4, 6, and 8 (C-Neutro 1-200 and PC/HNC-Neutro 1-200) in a population of the subject's neutrophil cells

15 before the treatment, or at a first time point, or before administration of the compound, respectively; b) identifying a first difference between the measured levels of the one or more selected C-Neutro (or PC/HNC-Neutro) markers in step (a) and the levels of the one or more selected C-Neutro (or PC/HNC-Neutro) markers in a control (e.g., a control cell from a healthy subject, or a normal or

20 healthy cell from the subject) before the treatment, or at the first time point, or before administration of the compound, respectively; c) measuring the levels of the one or more selected C-Neutro (or PC/HNC-Neutro) markers in a population of the subject's neutrophil cells after the treatment, or at a second time point, or after administration of the compound, respectively; d) identifying a second difference

25 between the measured levels of the one or more selected C-Neutro (or PC/HNC-Neutro) markers in step c) and the levels of the one or more selected C-Neutro (or PC/HNC-Neutro) markers in a control after the treatment, or at the second time point, or after administration of the compound, respectively; and e) identifying a difference between the first difference and the second difference, wherein the

30 difference identified in e) is indicative of the efficacy of the treatment for the cancer, or the progression or regression of the cancer, or whether the compound is capable of ameliorating or treating the cancer, respectively, in the subject.

[0038] In an eleventh aspect, the methods provided in this invention for assessing the efficacy of a treatment for cancer, monitoring the progression or regression of cancer, or identifying a compound capable of ameliorating or treating cancer, respectively, in a subject comprising:

- 5 a) measuring the levels of at least one or more markers selected from the group consisting of Tables 1, 3, 5, and 7 (C-Macro markers 1-200 and PC/HNC-Macro markers 1-200) in a population of the subject's macrophage cells before the treatment, at a first time point, or before administration of the compound, respectively, and
- 10 measuring the levels of at least one or more markers selected from the group consisting of Tables 2, 4, 6, and 8 (C-Neutro 1-200 and PC/HNC-Neutro 1-200) in a population of the subject's neutrophil cells before the treatment, at the first time point, or before administration of the compound, respectively;
- b) measuring the levels of the at least one or more selected C-Macro (or
- 15 PC/HNC-Macro) markers in a population of the subject's non-phagocytic cells before the treatment, at the first time point, or before administration of the compound, respectively; and
- measuring the levels of the at least one or more selected C-Neutro (or
- PC/HNC-Neutro) markers in a population of the subject's non-phagocytic cells
- 20 before the treatment, at the first time point, or before administration of the compound, respectively;
- c) identifying a first difference between the measured levels of the at least one or more selected C-Macro (or PC/HNC-Macro) markers in steps a) and b); and
- identifying a second difference between the measured levels of the at least
- 25 one or more selected C-Neutro (or PC/HNC-Neutro) markers in steps a) and b);
- d) measuring the levels of the at least one or more selected C-Macro (or
- PC/HNC-Macro) marker in a population of the subject's macrophage cells after the treatment, at a second time point, or after administration of the compound, respectively, and
- 30 measuring the levels of the at least one or more selected C-Neutro (or
- PC/HNC-Neutro) marker in a population of the subject's neutrophil cells after the

- treatment, at the second time point, or after administration of the compound, respectively;
- e) measuring the levels of the at least one or more selected C-Macro (or PC/HNC-Macro) markers in a population of the subject's non-phagocytic cells after the treatment, at the second time point, or after administration of the compound, respectively; and
- measuring the levels of the at least one or more selected C-Neutro (or PC/HNC-Neutro) markers in a population of the subject's non-phagocytic cells after the treatment, at the second time point, or after administration of the compound, respectively;
- f) identifying a third difference between the measured levels of the at least one or more selected C-Macro (or PC/HNC-Macro) markers in steps d) and e); and
- g) identifying a fourth difference between the measured levels of the at least one or more selected C-Neutro (or PC/HNC-Neutro) markers in steps d) and e);
- h) identifying a difference between the first and second differences; and
- i) identifying a difference between the third and fourth differences, wherein the differences identified in h) and i) are indicative of the efficacy of the treatment for the cancer, or the progression or regression of the cancer, or whether the compound is capable of ameliorating or treating the cancer, respectively, in the subject.
- [0039]** In an twelfth aspect, the methods provided in this invention for assessing the efficacy of a treatment for cancer, monitoring the progression or regression of cancer, or identifying a compound capable of ameliorating or treating cancer, respectively, in a subject comprising:
- a) measuring the levels of at least one or more markers selected from the group consisting of Tables 1, 3, 5, and 7 (C-Macro markers 1-200 and PC/HNC-Macro markers 1-200) in a population of the subject's macrophage cells before the treatment, at a first time point, or before administration of the compound, respectively, and
- measuring the levels of at least one or more markers selected from the group consisting of Tables 2, 4, 6, and 8 (C-Neutro 1-200 and PC/HNC-Neutro 1-

200) in a population of the subject's neutrophil cells before the treatment, at the first time point, or before administration of the compound, respectively;

b) identifying a first difference between the measured levels of the at least one or more selected C-Macro (or PC/HNC-Macro) markers in steps a) and the
5 levels of the at least one or more selected C-Macro (or PC/HNC-Macro) markers in a control before the treatment, at the first time point, or before administration of the compound, respectively; and

identifying a second difference between the measured levels of the at least one or more selected C-Neutro (or PC/HNC-Neutro) markers in steps a) and the
10 levels of the at least one or more selected C-Neutro (or PC/HNC-Neutro) markers in a control before the treatment, at the first time point, or before administration of the compound, respectively;

c) measuring the levels of the at least one or more selected C-Macro marker in a population of the subject's macrophage cells after the treatment, at a second
15 time point, or after administration of the compound, respectively, and

measuring the levels of the at least one or more selected C-Neutro marker in a population of the subject's neutrophil cells after the treatment, at the second time point, or after administration of the compound, respectively;

d) identifying a third difference between the measured levels of the at least
20 one or more selected C-Macro (or PC/HNC-Macro) markers in steps c) and the levels of the at least one or more selected C-Macro (or PC/HNC-Macro) markers in a control after the treatment, at the second time point, or after administration of the compound, respectively; and

e) identifying a fourth difference between the measured levels of the at
25 least one or more selected C-Neutro (or PC/HNC-Neutro) markers in steps c) and the levels of the at least one or more selected C-Neutro (or PC/HNC-Neutro) markers in a control after the treatment, at the second time point, or after administration of the compound, respectively;

f) identifying a difference between the first and second differences; and

30 g) identifying a difference between the third and fourth differences,

wherein the differences identified in f) and g) are indicative of the efficacy of the treatment for the cancer, or the progression or regression of the cancer, or

whether the compound is capable of ameliorating or treating the cancer, respectively, in the subject.

[0040] In various aspects of the present invention, the selected markers comprise one or more (e.g., two, three, four or more) C-Macro markers selected from the group consisting of C-Macro 1-4 and C-Macro 101-104. The selected markers may comprise one or more (e.g., two, three, four, five, six, seven, eight, nine, ten, eleven or twelve more) PC/HNC-Macro markers selected from the group consisting of PC/HNC-Macro 101-110 and PC/HNC-Macro 1-12. In some embodiments, the selected markers may comprise at least one or more C-Macro markers selected from the group consisting of C-Macro 1-4 and C-Macro 101-104 and at least one or more PC/HNC-Macro markers selected from the group consisting of PC/HNC-Macro 101-110 and PC/HNC-Macro 1-12. In some embodiments, the selected markers comprise C-Macro 1-4. In some embodiments, the selected markers comprise C-Macro 101-104. In some embodiments, the selected markers comprise P2RY10, TNFAIP3, TIPARP, and CXCR1. In some embodiments, the selected markers comprise markers that correspond to transcript cluster ID of 8168524, 8122265, 8058905, and 8034837, i.e., P2RY10, TNFAIP3, CXCR1, and DNAJB1. In some embodiments, the selected markers comprise PC/HNC-Macro 101-110. In some embodiments, the selected markers comprise PC/HNC-Macro 1-12. In some embodiments, the selected markers are up-regulated (see Tables 1, 3, 5, and 7 for up-regulated markers) in cancer patients. In some embodiments, the selected markers are down-regulated (see Tables 1, 3, 5, and 7 for down-regulated markers) in cancer patients. In some embodiments, the selected markers comprise at least one marker that is up-regulated and at least one marker that is down-regulated.

[0041] In various aspects of the present invention, the selected markers comprise one or more (e.g., two, three, four, five, six, seven, eight, or more) C-Neutro markers selected from the group consisting of C-Neutro1-8 and C-Neutro 101-108. In those aspect, the selected markers may also comprise one or more (e.g., two, three, four, five, six, seven, eight, nine, or more) PC/HNC-Neutro markers selected from the group consisting of PC/HNC-Neutro 101-109 and PC/HNC- Neutro 1-9. In some embodiments, the selected markers may comprise at least one or more C-

Neutro markers selected from the group consisting of C- Neutro 1-8 and C- Neutro 101-108 and at least one or more PC/HNC- Neutro markers selected from the group consisting of PC/HNC- Neutro 101-109 and PC/HNC- Neutro 1-9. In some embodiments, the selected markers comprise C- Neutro 1-8. In some
5 embodiments, the selected markers comprise C- Neutro 101-108. In some embodiments, the selected markers comprise EEF1A1, RPL23A, RPL14, and RPL3. In some embodiments, the selected markers comprise markers that correspond to transcript cluster ID of 8180410, 8158952, 8138531, 8091806, 8076209, 7956743, 8026440, and 8005943, for example, EEF1A1, RPL23A,
10 RPL3, and RPL14. In some embodiments, the selected markers comprise PC/HNC- Neutro 101-109. In some embodiments, the selected markers comprise PC/HNC- Neutro 1-9. In some embodiments, the selected markers are up-regulated (see Tables 2, 4, 6, and 8 for up-regulated markers) in cancer patients. In some embodiments, the selected markers are down-regulated (see Tables 2, 4, 6,
15 and 8 for down-regulated markers) in cancer patients. In some embodiments, the selected markers comprise at least one marker that is up-regulated and at least one marker that is down-regulated.

[0042] In various embodiments of the present invention, at least one or more of the selected markers may be substituted with a biological marker different from
20 any of the selected marker. In some embodiments, such biological marker may be a known marker for cancer. In some embodiments, such biological marker and the selected marker being substituted may belong to the same signaling or biological pathway (e.g., TGF-beta pathway, apoptosis pathway, programmed cell death-associated pathway), or may have similar biological function or activity (e.g.,
25 protein synthesis, cytokine production, Th1 cytokine production, nucleotide binding, receptor for purines coupled to G-proteins, inhibitor of programmed cell death, receptor of IL-8/activating neutrophils, or interacting with HSP70/ stimulating its ATPase activity), or may be regulated by a common protein, or may belong to the same protein complex (e.g., ribosome).

30 [0043] In various embodiments of the present invention, a population of the subject's macrophage cells is used as the selected phagocytic cells for measuring the levels of the selected markers and a population of the subject's T-cells is used

as the selected non-phagocytic cells for measuring the levels of the selected markers.

[0044] In various embodiments of the present invention, a population of the subject's neutrophil cells is used as the selected phagocytic cells for measuring the levels of the selected markers and a population of the subject's T-cells is used as the selected non-phagocytic cells for measuring the levels of the selected markers.

[0045] In some embodiments, two sub-populations of phagocytic cells are used in the methods of this invention, i.e., phagocytic cells that have a DNA content greater than 2n (the >2n phagocytic cells) and phagocytic cells that have a DNA content of 2n (the =2n phagocytic cells). In those embodiments, the levels of the selected markers in the >2n phagocytic cells are compared to the =2n phagocytic cells to identify one or more difference. The identified difference indicates whether the subject has cancer, or has a risk of developing cancer, or has a progressing or progressive cancer.

[0046] In some embodiments, the levels of two, three, four, five, six, seven, eight, nine, ten, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, or more markers selected from Tables 1-8 are measured. In some embodiments, the levels of two, three, four, five, six, seven, eight, nine, ten, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, or more markers selected from Tables 1, 3, 5, and 7 are measured. In some embodiments, the levels of two, three, four, five, six, seven, eight, nine, ten, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, or more markers selected from Tables 2, 4, 6, and 8 are measured. In some embodiments, one or more markers selected from Tables 1, 3, 5, and 7 and one or more markers selected from Tables 2, 4, 6, and 8 are measured.

[0047] The gene names/descriptions provided in Tables 1-8 are merely illustrative. The markers of this invention encompass all forms and variants of any specifically described markers, including, but not limited to, polymorphic or allelic variants, isoforms, mutants, derivatives, precursors including nucleic acids and pro-proteins, cleavage products, and structures comprised of any of the markers as constituent subunits of the fully assembled structure.

[0048] A "patient", "subject", or "individual" are used interchangeably and refer to either a human or a non-human animal. These terms include mammals, such as

humans, primates, livestock animals (e.g., bovines, porcines), companion animals (e.g., canines, felines) and rodents (e.g., mice and rats).

[0049] As used herein, the terms “control”, “normal control”, “healthy control”, and “not-diseased cells” likewise mean a sample (e.g., cells, serum, tissue) taken
5 from a source (e.g., subject, control subject, cell line) that does not have the condition or disease being assayed and therefore may be used to determine the baseline for the condition or disorder being measured. A control subject refers to any individual that has not been diagnosed as having the disease or condition being assayed. It is also understood that the control subject, normal control, and healthy
10 control, include data obtained and used as a standard, i.e. it can be used over and over again for multiple different subjects. In other words, for example, when comparing a subject sample to a control sample, the data from the control sample could have been obtained in a different set of experiments, for example, it could be an average obtained from a number of healthy subjects and not actually obtained at
15 the time the data for the subject was obtained.

[0050] The term “diagnosis” as used herein refers to methods by which the skilled artisan can estimate and/or determine whether or not a patient is suffering from a given disease or condition. In some embodiments, the term “diagnosis” also refers to staging (e.g., Stage I, II, III, or IV) of cancer. The skilled artisan often
20 makes a diagnosis on the basis of one or more diagnostic indicators, e.g., a marker, the presence, absence, amount, or change in amount of which is indicative of the presence, severity, or absence of the condition.

[0051] The term “prognosis” as used herein refers to is used herein to refer to the likelihood of cancer progression, including recurrence of cancer.

25 [0052] The disclosure of the International Applications PCT/US11/44969, PCT/US11/45018, and PCT/US09/31395 and U.S. Provisional Applications 61/660,518 and 61/660,427 are incorporated herein by reference for all purposes.

[0053] Each embodiment described herein may be combined with any other embodiment described herein.

30 [0054] Methods using the cancer markers described herein provide high specificity, sensitivity, and accuracy in detecting and diagnosing cancer. The methods also eliminate the “inequality of baseline” that is known to occur among

individuals due to intrinsic (e.g., age, gender, ethnic background, health status and the like) and temporal variations in marker expression. Additionally, by using a comparison of phagocytes and non-phagocytes from the same individual, the methods also allow detection, diagnosis, and treatment to be personalized to the individual. Accordingly, in some embodiments, the invention provides non-invasive assays for the early detection of cancer, i.e., before the cancer can be diagnosed by conventional diagnostic techniques, e.g., imaging techniques, and, therefore, provide a foundation for improved decision-making relative to the needs and strategies for intervention, prevention, and treatment of individuals with such disease or condition.

[0055] The methods described herein are supported by whole genome microarray data of total RNA samples isolated from macrophages and neutrophils and from non-phagocytic T cells. The samples were obtained from human subjects with and without cancer. The data from these microarray experiments demonstrate that macrophage-T cell and neutrophil-T cell comparisons easily and accurately differentiate between cancer patients and human subjects without cancer.

[0056] The methods of this invention can be used together with any known diagnostic methods, such as physical inspection, visual inspection, biopsy, scanning, histology, radiology, imaging, ultrasound, use of a commercial kit, genetic testing, immunological testing, analysis of bodily fluids, or monitoring neural activity.

[0057] Phagocytic cells that can be used in the methods of this invention include all types of cells that are capable of ingesting various types of substances (e.g., apoptotic cells, infectious agents, dead cells, viable cells, cell-free DNAs, cell-free RNAs, cell-free proteins). In some embodiments, the phagocytic cells are neutrophils, macrophages, monocytes, dendritic cells, foam cells, mast cells, eosinophils, or keratinocytes. In some embodiments, the phagocytic cells can be a mixture of different types of phagocytic cells. In some embodiments, the phagocytic cells can be activated phagocytic cells, e.g., activated macrophages or neutrophils. In some embodiments, a phagocyte is a histiocyte, e.g., a Langerhans cell.

[0058] The methods of this invention can be applied to any cancer, including without limitation, for example, carcinomas (*e.g.*, malignant tumors derived from epithelial cells such as, for example, common forms of breast, prostate, lung, and colon cancer), sarcomas (*e.g.*, malignant tumors derived from connective tissue or
5 mesenchymal cells), lymphomas and leukemias (*i.e.*, malignancies derived from hematopoietic cells), germ cell tumors (*i.e.*, tumors derived from totipotent cells). Specific examples of these cancers include, without limitation, cancers of: prostate, head and neck, breast, skin, bone, ovaries, uterus, cervix, liver, lung, brain, spine, larynx, gallbladder, pancreas, rectum, parathyroid, thyroid, adrenal gland, immune
10 system, head and neck, colon, stomach, bronchi, and kidneys.

[0059] As used herein, “treating” cancer refers to taking steps to obtain beneficial or desired results, including clinical results. Beneficial or desired clinical results include, but are not limited to, alleviation or amelioration of one or more symptoms associated with diseases or conditions.

15 [0060] As used herein, “administering” or “administration of” a compound or an agent to a subject can be carried out using one of a variety of methods known to those skilled in the art. For example, a compound or an agent can be administered, intravenously, arterially, intradermally, intramuscularly, intraperitoneally, intravenously, subcutaneously, ocularly, sublingually, orally (by ingestion),
20 intranasally (by inhalation), intraspinally, intracerebrally, and transdermally (by absorption, *e.g.*, through a skin duct). A compound or agent can also appropriately be introduced by rechargeable or biodegradable polymeric devices or other devices, *e.g.*, patches and pumps, or formulations, which provide for the extended, slow, or controlled release of the compound or agent. Administering can also be
25 performed, for example, once, a plurality of times, and/or over one or more extended periods. In some aspects, the administration includes both direct administration, including self-administration, and indirect administration, including the act of prescribing a drug. For example, as used herein, a physician who instructs a patient to self-administer a drug, or to have the drug administered by
30 another and/or who provides a patient with a prescription for a drug is administering the drug to the patient. In some embodiments, a compound or an agent is administered orally, *e.g.*, to a subject by ingestion, or intravenously, *e.g.*,

to a subject by injection. In some embodiments, the orally administered compound or agent is in an extended release or slow release formulation, or administered using a device for such slow or extended release.

[0061] In certain embodiments, markers used in the methods of invention are up-regulated or activated in phagocytes (e.g., macrophages or neutrophils) compared to non-phagocytes. In certain embodiments, markers used in the methods of invention are down-regulated or inhibited in phagocytes (e.g., macrophages or neutrophils) compared to non-phagocytes. As used herein, “up-regulation or up-regulated” can refer to an increase in expression levels (e.g., gene expression or protein expression), gene copy numbers, gene dosages, and other qualitative or quantitative detectable state of the markers. Similarly, “down-regulation or down-regulated” can refer to a decrease in expression levels, gene copy numbers, gene dosages, and other qualitative or quantitative detectable state of the markers. As used herein, “activation or activated” can refer to an active state of the marker, e.g., a phosphorylation state, a DNA methylation state, or a DNA acetylation state. Similarly, “inhibition or inhibited” can refer to a repressed state or an inactivated state of the marker, e.g., a de-phosphorylation state, a ubiquitination state, a DNA de-methylation state.

[0062] In certain embodiments, methods of this invention also comprise at least one of the following steps before determination of various levels: i) lysing the phagocytic or non-phagocytic cells; and ii) extracting cellular contents from the lysed cells. Any known cell lysis and extraction methods can be used herein. In certain embodiments, at least one or more cancer markers are present in the phagocytes. In certain embodiments, there is no marker present in the cellular contents of the non-phagocytic cells.

[0063] In certain embodiments, the phagocytic cells and/or non-phagocytic cells are isolated from a bodily fluid sample, tissues, or population of cells. Exemplary bodily fluid samples can be whole blood, urine, stool, saliva, lymph fluid, cerebrospinal fluid, synovial fluid, cystic fluid, ascites, pleural effusion, fluid obtained from a pregnant woman in the first trimester, fluid obtained from a pregnant woman in the second trimester, fluid obtained from a pregnant woman in the third trimester, maternal blood, amniotic fluid, chorionic villus sample, fluid

from a preimplantation embryo, maternal urine, maternal saliva, placental sample, fetal blood, lavage and cervical vaginal fluid, interstitial fluid, buccal swab sample, sputum, bronchial lavage, Pap smear sample, or ocular fluid. In some embodiments, the phagocytic cells or non-phagocytic cells are isolated from white
5 blood cells.

[0064] In the methods of this invention, cell separation/isolation/purification methods are used to isolate populations of cells from bodily fluid sample, cells, or tissues of a subject. A skilled worker can use any known cell separation/isolation/purification techniques to isolate phagocytic cells and non-
10 phagocytic cells from a bodily fluid. Exemplary techniques include, but are not limited to, using antibodies, flow cytometry, fluorescence activated cell sorting, filtration, gradient-based centrifugation, elution, microfluidics, magnetic separation technique, fluorescent-magnetic separation technique, nanostructure, quantum dots, high throughput microscope-based platform, or a combination thereof.

[0065] In certain embodiments, the phagocytic cells and/or non-phagocytic cells are isolated by using a product secreted by the cells. In certain embodiments, the phagocytic cells and/or non-phagocytic cells are isolated by using a cell surface target (e.g., receptor protein) on the surface of the cells. In some embodiments, the cell surface target is a protein that has been engulfed by phagocytic cells. In some
15 20 embodiments, the cell surface target is expressed by cells on their plasma membranes. In some embodiments, the cell surface target is an exogenous protein that is translocated on the plasma membranes, but not expressed by the cells (e.g., the phagocytic cells). In some embodiments, the cell surface target is a marker of cancer.

[0066] In certain aspects of the methods described herein, analytes include nucleic acids, proteins, or any combinations thereof. In certain aspects of the methods described herein, markers include nucleic acids, proteins, or any combinations thereof. As used herein, the term “nucleic acid” is intended to include DNA molecules (e.g., cDNA or genomic DNA), RNA molecules (e.g.,
25 30 mRNA), DNA-RNA hybrids, and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be a nucleotide, oligonucleotide, double-stranded DNA, single-stranded DNA, multi-stranded

DNA, complementary DNA, genomic DNA, non-coding DNA, messenger RNA (mRNAs), microRNA (miRNAs), small nucleolar RNA (snoRNAs), ribosomal RNA (rRNA), transfer RNA (tRNA), small interfering RNA (siRNA), heterogeneous nuclear RNAs (hnRNA), or small hairpin RNA (shRNA). In some
 5 embodiments, the nucleic acid is a transrenal nucleic acid. A transrenal nucleic acid is an extracellular nucleic acid that is excreted in the urine. See, e.g., U.S. Patent Publication No. 20100068711 and U.S. Patent Publication No. 20120021404.

[0067] As used herein, the term “amino acid” includes organic compounds
 10 containing both a basic amino group and an acidic carboxyl group. Included within this term are natural amino acids (e.g., L-amino acids), modified and unusual amino acids (e.g., D-amino acids and β -amino acids), as well as amino acids which are known to occur biologically in free or combined form but usually do not occur in proteins. Natural protein occurring amino acids include alanine,
 15 arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, serine, threonine, tyrosine, tryptophan, proline, and valine. Natural non-protein amino acids include arginosuccinic acid, citrulline, cysteine sulfuric acid, 3,4-dihydroxyphenylalanine, homocysteine, homoserine, ornithine, 3-monoiodotyrosine, 3,5-diiodotryosine, 3,
 20 5, 5-triiodothyronine, and 3,3',5,5'- tetraiodothyronine. Modified or unusual amino acids include D-amino acids, hydroxylysine, 4-hydroxyproline, N-Cbz-protected amino acids, 2,4-diaminobutyric acid, homoarginine, norleucine, N-methylaminobutyric acid, naphthylalanine, phenylglycine, .alpha.-phenylproline, tert-leucine, 4-aminocyclohexylalanine, N-methyl-norleucine, 3,4-dehydroproline,
 25 N,N-dimethylaminoglycine, N-methylaminoglycine, 4-aminopiperidine-4-carboxylic acid, 6-aminocaproic acid, trans-4-(aminomethyl)-cyclohexanecarboxylic acid, 2-, 3-, and 4-(aminomethyl)- benzoic acid, 1-aminocyclopentanecarboxylic acid, 1-aminocyclopropanecarboxylic acid, and 2-benzyl-5-aminopentanoic acid.

30 [0068] As used herein, the term “peptide” includes compounds that comprise two or more amino acids that are linked by means of a peptide bond. Peptides may have a molecular weight of less than 10,000 Daltons, less than 5,000 Daltons, or

less than 2,500 Daltons. The term “peptide” also includes compounds containing both peptide and non-peptide components, such as pseudopeptide or peptidomimetic residues or other non-amino acid components. Such compounds containing both peptide and non-peptide components may also be referred to as a
5 “peptide analog.”

[0069] As used herein, the term “protein” includes compounds that comprise amino acids arranged in a linear chain and joined together by peptide bonds between the carboxyl and amino groups of adjacent amino acid residues. Proteins used in methods of the invention include, but are not limited to, amino acids,
10 peptides, antibodies, antibody fragments, cytokines, lipoproteins, or glycoproteins.

[0070] As used herein, the term “antibody” includes polyclonal antibodies, monoclonal antibodies (including full length antibodies which have an immunoglobulin Fc region), antibody compositions with polyepitopic specificity, multispecific antibodies (e.g., bispecific antibodies, diabodies, and single-chain
15 molecules, and antibody fragments (*e.g.*, Fab or F(ab')₂, and Fv). For the structure and properties of the different classes of antibodies, see *e.g.*, Basic and Clinical Immunology, 8th Edition, Daniel P. Sties, Abba I. Terr and Tristram G. Parslow (eds), Appleton & Lange, Norwalk, Conn., 1994, page 71 and Chapter 6.

[0071] As used herein, the term “cytokine” refers to a secreted protein or active
20 fragment or mutant thereof that modulates the activity of cells of the immune system. Examples of cytokines include, without limitation, interleukins, interferons, chemokines, tumor necrosis factors, colony-stimulating factors for immune cell precursors, and the like.

[0072] As used herein, the term “lipoprotein” includes negatively charged
25 compositions that comprise a core of hydrophobic cholesteryl esters and triglyceride surrounded by a surface layer of amphipathic phospholipids with which free cholesterol and apolipoproteins are associated. Lipoproteins may be characterized by their density (*e.g.* very-low-density lipoprotein (VLDL), low-density lipoprotein (LDL) and high density lipoprotein (HDL)), which is
30 determined by their size, the relative amounts of lipid and protein. Lipoproteins may also be characterized by the presence or absence of particular modifications (*e.g.* oxidization, acetylation, or glycation).

[0073] As used herein, the term “glycoprotein” includes glycosides which have one or more oligo- or polysaccharides covalently attached to a peptide or protein. Exemplary glycoproteins can include, without limitation, immunoglobulins, members of the major histocompatibility complex, collagens, mucins, glycoprotein
5 IIb/IIIa, glycoprotein-41 (gp41) and glycoprotein-120 (gp12), follicle-stimulating hormone, alpha-fetoprotein, erythropoietin, transferrins, alkaline phosphatase, and lectins.

[0074] In some embodiments of the invention, a sample may comprise one or more stabilizers for a cell or an analyte such as DNA, RNA, and/or protein. For
10 example, a sample may comprise a DNA stabilizer, an RNA stabilizer, and/or a protein stabilizer. Stabilizers are well known in the art and include, for example, DNase inhibitors, RNase inhibitors, and protease inhibitors or equivalents thereof.

[0075] In some embodiments of the invention, levels of at least one or more
15 cancer markers are compared. This comparison can be quantitative or qualitative. Quantitative measurements can be taken using any of the assays described herein. For example, sequencing, direct sequencing, random shotgun sequencing, Sanger dideoxy termination sequencing, targeted sequencing, whole-genome sequencing, sequencing by hybridization, pyrosequencing, capillary electrophoresis, gel
20 electrophoresis, duplex sequencing, cycle sequencing, single-base extension sequencing, solid-phase sequencing, high-throughput sequencing, massively parallel signature sequencing, emulsion PCR, co-amplification at lower denaturation temperature-PCR (COLD-PCR), sequencing by reversible dye terminator, paired-end sequencing, near-term sequencing, exonuclease sequencing,
25 sequencing by ligation, short-read sequencing, single-molecule sequencing, sequencing-by-synthesis, real-time sequencing, reverse-terminator sequencing, nanopore sequencing, 454 sequencing, Solexa Genome Analyzer sequencing, SOLiD® sequencing, MS-PET sequencing, mass spectrometry, matrix assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry,
30 electrospray ionization (ESI) mass spectrometry, surface-enhanced laser desorption/ionization-time of flight (SELDI-TOF) mass spectrometry, quadrupole-time of flight (Q-TOF) mass spectrometry, atmospheric pressure photoionization

mass spectrometry (APPI-MS), Fourier transform mass spectrometry (FTMS), matrix-assisted laser desorption/ionization-Fourier transform-ion cyclotron resonance (MALDI-FT-ICR) mass spectrometry, secondary ion mass spectrometry (SIMS), polymerase chain reaction (PCR) analysis, quantitative PCR, real-time
5 PCR, fluorescence assay, colorimetric assay, chemiluminescent assay, or a combination thereof.

[0076] Quantitative comparisons can include statistical analyses such as t-test, ANOVA, Krustal-Wallis, Wilcoxon, Mann-Whitney, and odds ratio. Quantitative differences can include differences in the levels of markers between levels or
10 differences in the numbers of markers present between levels, and combinations thereof. Examples of levels of the markers can be, without limitation, gene expression levels, nucleic acid levels, and protein levels. Qualitative differences can include, but are not limited to, activation and inactivation, protein degradation, nucleic acid degradation, and covalent modifications.

15 [0077] In certain embodiments of the invention, the level is a nucleic acid level or a protein level, or a combination thereof. The level can be qualitatively or quantitatively determined.

[0078] A nucleic acid level can be, without limitation, a genotypic level, a single nucleotide polymorphism level, a gene mutation level, a gene copy number level, a
20 DNA methylation level, a DNA acetylation level, a chromosome dosage level, a gene expression level, or a combination thereof.

[0079] The nucleic acid level can be determined by any methods known in the art to detect genotypes, single nucleotide polymorphisms, gene mutations, gene copy numbers, DNA methylation states, DNA acetylation states, chromosome dosages.
25 Exemplary methods include, but are not limited to, polymerase chain reaction (PCR) analysis, sequencing analysis, electrophoretic analysis, restriction fragment length polymorphism (RFLP) analysis, Northern blot analysis, quantitative PCR, reverse-transcriptase-PCR analysis (RT-PCR), allele-specific oligonucleotide hybridization analysis, comparative genomic hybridization, heteroduplex mobility
30 assay (HMA), single strand conformational polymorphism (SSCP), denaturing gradient gel electrophoresis (DGGE), RNAase mismatch analysis, mass spectrometry, tandem mass spectrometry, matrix assisted laser

desorption/ionization-time of flight (MALDI-TOF) mass spectrometry, electrospray ionization (ESI) mass spectrometry, surface-enhanced laser desorption/ionization-time of flight (SELDI-TOF) mass spectrometry, quadrupole-time of flight (Q-TOF) mass spectrometry, atmospheric pressure photoionization mass spectrometry (APPI-MS), Fourier transform mass spectrometry (FTMS), matrix-assisted laser desorption/ionization-Fourier transform-ion cyclotron resonance (MALDI-FT-ICR) mass spectrometry, secondary ion mass spectrometry (SIMS), surface plasmon resonance, Southern blot analysis, in situ hybridization, fluorescence in situ hybridization (FISH), chromogenic in situ hybridization (CISH), immunohistochemistry (IHC), microarray, comparative genomic hybridization, karyotyping, multiplex ligation-dependent probe amplification (MLPA), Quantitative Multiplex PCR of Short Fluorescent Fragments (QMPSF), microscopy, methylation specific PCR (MSP) assay, HpaII tiny fragment Enrichment by Ligation-mediated PCR (HELP) assay, radioactive acetate labeling assays, colorimetric DNA acetylation assay, chromatin immunoprecipitation combined with microarray (ChIP-on-chip) assay, restriction landmark genomic scanning, Methylated DNA immunoprecipitation (MeDIP), molecular break light assay for DNA adenine methyltransferase activity, chromatographic separation, methylation-sensitive restriction enzyme analysis, bisulfite-driven conversion of non-methylated cytosine to uracil, co-amplification at lower denaturation temperature-PCR (COLD-PCR), multiplex PCR, methyl-binding PCR analysis, or a combination thereof.

[0080] As used herein, the term “sequencing” is used in a broad sense and refers to any technique known in the art that allows the order of at least some consecutive nucleotides in at least part of a nucleic acid to be identified, including without limitation at least part of an extension product or a vector insert. Exemplary sequencing techniques include targeted sequencing, single molecule real-time sequencing, whole transcriptome shotgun sequencing (“RNA-seq”), electron microscopy-based sequencing, transistor-mediated sequencing, direct sequencing, random shotgun sequencing, Sanger dideoxy termination sequencing, exon sequencing, whole-genome sequencing, sequencing by hybridization, pyrosequencing, capillary electrophoresis, gel electrophoresis, duplex sequencing,

cycle sequencing, single-base extension sequencing, solid-phase sequencing, high-throughput sequencing, massively parallel signature sequencing, emulsion PCR, co-amplification at lower denaturation temperature-PCR (COLD-PCR), multiplex PCR, sequencing by reversible dye terminator, paired-end sequencing, near-term
5 sequencing, exonuclease sequencing, sequencing by ligation, short-read sequencing, single-molecule sequencing, sequencing-by-synthesis, real-time sequencing, reverse-terminator sequencing, nanopore sequencing, 454 sequencing, Solexa Genome Analyzer sequencing, SOLiD® sequencing, MS-PET sequencing, mass spectrometry, and a combination thereof.. In some embodiments, sequencing
10 comprises an detecting the sequencing product using an instrument, for example but not limited to an ABI PRISM® 377 DNA Sequencer, an ABI PRISM® 310, 3100, 3100-Avant, 3730, or 3730xI Genetic Analyzer, an ABI PRISM® 3700 DNA Analyzer, or an Applied Biosystems SOLiD™ System (all from Applied Biosystems), a Genome Sequencer 20 System (Roche Applied Science), or a mass
15 spectrometer. In certain embodiments, sequencing comprises emulsion PCR. In certain embodiments, sequencing comprises a high throughput sequencing technique, for example but not limited to, massively parallel signature sequencing (MPSS).

[0081] In further embodiments of the invention, a protein level can be a protein
20 expression level, a protein activation level, or a combination thereof. In some embodiments, a protein activation level can comprise determining a phosphorylation state, an ubiquitination state, a myristoylation state, or a conformational state of the protein.

[0082] A protein level can be detected by any methods known in the art for
25 detecting protein expression levels, protein phosphorylation state, protein ubiquitination state, protein myristoylation state, or protein conformational state. In some embodiments, a protein level can be determined by an immunohistochemistry assay, an enzyme-linked immunosorbent assay (ELISA), in situ hybridization, chromatography, liquid chromatography, size exclusion
30 chromatography, high performance liquid chromatography (HPLC), gas chromatography, mass spectrometry, tandem mass spectrometry, matrix assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry,

electrospray ionization (ESI) mass spectrometry, surface-enhanced laser
 desorption/ionization-time of flight (SELDI-TOF) mass spectrometry, quadrupole-
 time of flight (Q-TOF) mass spectrometry, atmospheric pressure photoionization
 mass spectrometry (APPI-MS), Fourier transform mass spectrometry (FTMS),
 5 matrix-assisted laser desorption/ionization-Fourier transform-ion cyclotron
 resonance (MALDI-FT-ICR) mass spectrometry, secondary ion mass spectrometry
 (SIMS), radioimmunoassays, microscopy, microfluidic chip-based assays, surface
 plasmon resonance, sequencing, Western blotting assay, or a combination thereof.
 [0083] As used herein, the “difference” between different levels detected by the
 10 methods of this invention can refer to different gene copy numbers, different DNA,
 RNA, or protein expression levels, different DNA methylation states, different
 DNA acetylation states, and different protein modification states. The difference
 can be a difference greater than 1 fold. In some embodiments, the difference is a
 1.05-fold, 1.1-fold, 1.2-fold, 1.3-fold, 1.4-fold, 1.5-fold, 2-fold, 2.5-fold, 3-fold, 4-
 15 fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, or 10-fold difference. In some
 embodiments, the difference is any fold difference between 1-10, 2-10, 5-10, 10-
 20, or 10-100 fold.

[0084] In some embodiments, the difference is differential gene expression
 (DGE), e.g. DGE of phagocytes vs. non-phagocytes. DGE can be measured as $X =$
 20 $\log_2(Y_P) - \log_2(Y_{NP})$. The DGE may be any number, provided that it is
 significantly different between the phagocytes and the non-phagocytes. For
 example, a 2-fold increased in gene expression could be represented as $X =$
 $\log_2(Y_P) - \log_2(Y_{NP}) = \log_2(Y_P/Y_{NP}) = \log_2(2) = 1$, while a 2-fold decrease in gene
 expression could be represented as $X = \log_2(Y_P) - \log_2(Y_{NP}) = \log_2(Y_P/Y_{NP})$
 25 $= \log_2(1/2) = -1$. Down-regulated genes have $X < 0$, while up-regulated genes have
 $X > 0$. See, e.g., Efron, J Am Stat Assoc 104:1015-1028 (2009).

[0085] A general principle of assays to detect markers involves preparing a
 sample or reaction mixture that may contain the marker (e.g., one or more of DNA,
 RNA, or protein) and a probe under appropriate conditions and for a time sufficient
 30 to allow the marker and probe to interact and bind, thus forming a complex that
 can be removed and/or detected in the reaction mixture. These assays can be
 conducted in a variety of ways.

- [0086] For example, one method to conduct such an assay would involve anchoring the marker or probe onto a solid phase support, also referred to as a substrate, and detecting target marker/probe complexes anchored on the solid phase at the end of the reaction. In one embodiment of such a method, a sample from a subject, which is to be assayed for presence and/or concentration of marker, can be anchored onto a carrier or solid phase support. In another embodiment, the reverse situation is possible, in which the probe can be anchored to a solid phase and a sample from a subject can be allowed to react as an unanchored component of the assay.
- 10 [0087] There are many established methods for anchoring assay components to a solid phase. These include, without limitation, marker or probe molecules which are immobilized through conjugation of biotin and streptavidin. Such biotinylated assay components can be prepared from biotin-NHS(N-hydroxy-succinimide) using techniques known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). In certain embodiments, the surfaces with immobilized assay components can be prepared in advance and stored.
- 15 [0088] Other suitable carriers or solid phase supports for such assays include any material capable of binding the class of molecule to which the marker or probe belongs. Well known supports or carriers include, but are not limited to, glass, polystyrene, nylon, polypropylene, nylon, polyethylene, dextran, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite.
- 20 [0089] In order to conduct assays with the above mentioned approaches, the non-immobilized component is added to the solid phase upon which the second component is anchored. After the reaction is complete, uncomplexed components may be removed (e.g., by washing) under conditions such that any complexes formed will remain immobilized upon the solid phase. The detection of marker/probe complexes anchored to the solid phase can be accomplished in a number of methods outlined herein.
- 25 [0090] In certain exemplary embodiments, the probe, when it is the unanchored assay component, can be labeled for the purpose of detection and readout of the
- 30

assay, either directly or indirectly, with detectable labels discussed herein and which are well-known to one skilled in the art.

[0091] It is also possible to directly detect marker/probe complex formation without further manipulation or labeling of either component (marker or probe),
5 for example by utilizing the technique of fluorescence energy transfer (see, for example, U.S. Patent Nos. 5,631,169 and 4,868,103). A fluorophore label on the first, 'donor' molecule is selected such that, upon excitation with incident light of appropriate wavelength, its emitted fluorescent energy will be absorbed by a fluorescent label on a second 'acceptor' molecule, which in turn is able to
10 fluoresce due to the absorbed energy. Alternately, the 'donor' protein molecule may simply utilize the natural fluorescent energy of tryptophan residues. Labels are chosen that emit different wavelengths of light, such that the 'acceptor' molecule label may be differentiated from that of the 'donor'. Since the efficiency of energy transfer between the labels is related to the distance separating the
15 molecules, spatial relationships between the molecules can be assessed. In a situation in which binding occurs between the molecules, the fluorescent emission of the 'acceptor' molecule label in the assay should be maximal. An FET binding event can be conveniently measured through standard fluorometric detection means well known in the art (e.g., using a fluorimeter).

20 [0092] In another embodiment, determination of the ability of a probe to recognize a marker can be accomplished without labeling either assay component (probe or marker) by utilizing a technology such as real-time Biomolecular Interaction Analysis (BIA) (see, e.g., Sjolander, S. and Urbaniczky, C, 1991, Anal. Chem. 63:2338 2345 and Szabo et al, 1995, Curr. Opin. Struct. Biol. 5:699
25 705). As used herein, "BIA" or "surface plasmon resonance" is a technology for studying biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore). Changes in the mass at the binding surface (indicative of a binding event) result in alterations of the refractive index of light near the surface (the optical phenomenon of surface plasmon resonance (SPR)), resulting in
30 a detectable signal which can be used as an indication of real-time reactions between biological molecules.

[0093] Alternatively, in another embodiment, analogous diagnostic and prognostic assays can be conducted with marker and probe as solutes in a liquid phase. In such an assay, the complexed marker and probe are separated from uncomplexed components by any of a number of standard techniques, including
5 but not limited to: differential centrifugation, chromatography, electrophoresis and immunoprecipitation. In differential centrifugation, marker/probe complexes may be separated from uncomplexed assay components through a series of centrifugal steps, due to the different sedimentation equilibria of complexes based on their different sizes and densities (see, for example, Rivas and Minton (1993) Trends
10 Biochem. Sci. 18:284). Standard chromatographic techniques may also be utilized to separate complexed molecules from uncomplexed ones. For example, gel filtration chromatography separates molecules based on size, and through the utilization of an appropriate gel filtration resin in a column format, for example, the relatively larger complex may be separated from the relatively smaller
15 uncomplexed components. Similarly, the relatively different charge properties of the marker/probe complex as compared to the uncomplexed components may be exploited to differentiate the complex from uncomplexed components, for example through the utilization of ion-exchange chromatography resins. Such resins and chromatographic techniques are well known to one skilled in the art (see, e.g.,
20 Heegaard (1998) J. Mol. Recognit. 11 :141; Hage and Tweed (1997) J. Chromatogr. B. Biomed. Sci. Appl. 12:499). Gel electrophoresis may also be employed to separate complexed assay components from unbound components (see, e.g., Ausubel et al, ed., Current Protocols in Molecular Biology, John Wiley & Sons, New York, 1987 1999). In this technique, protein or nucleic acid
25 complexes are separated based on size or charge, for example. In order to maintain the binding interaction during the electrophoretic process, non-denaturing gel matrix materials and conditions in the absence of reducing agent are typically preferred. Appropriate conditions to the particular assay and components thereof will be well known to one skilled in the art.

30 [0094] In certain exemplary embodiments, the level of mRNA corresponding to the marker can be determined either by *in situ* and/or by *in vitro* formats in a biological sample using methods known in the art. Many expression detection

methods use isolated RNA. For in vitro methods, any RNA isolation technique that does not select against the isolation of mRNA can be utilized for the purification of RNA from blood cells (see, e.g., Ausubel et al, ed., Current Protocols in Molecular Biology, John Wiley & Sons, New York 1987 1999).

5 Additionally, large numbers of cells and/or samples can readily be processed using techniques well known to those of skill in the art, such as, for example, the single-step RNA isolation process of Chomczynski (1989, U.S. Patent No. 4,843,155).

[0095] Isolated mRNA can be used in hybridization or amplification assays that include, but are not limited to, Southern or Northern analyses, polymerase chain
10 reaction analyses and probe arrays. In certain exemplary embodiments, a diagnostic method for the detection of mRNA levels involves contacting the isolated mRNA with a nucleic acid molecule (probe) that can hybridize to the mRNA encoded by the gene being detected. The nucleic acid probe can be, for example, a full-length cDNA, or a portion thereof, such as an oligonucleotide of at
15 least 7, 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to an mRNA or genomic DNA encoding a marker of the present invention. Other suitable probes for use in the diagnostic assays of the invention are described herein. Hybridization of an mRNA with the probe indicates that the marker in question is being expressed.

20 [0096] In one format, the mRNA is immobilized on a solid surface and contacted with a probe, for example by running the isolated mRNA on an agarose gel and transferring the mRNA from the gel to a membrane, such as nitrocellulose. In an alternative format, the probe(s) are immobilized on a solid surface and the mRNA is contacted with the probe(s), for example, in a gene chip array. A skilled artisan
25 can readily adapt known mRNA detection methods for use in detecting the level of mRNA encoded by the markers of the present invention.

[0097] An alternative method for determining the level of mRNA corresponding to a marker of the present invention in a sample involves the process of nucleic acid amplification, e.g., by RT-PCR (the experimental embodiment set forth in
30 U.S. Patent Nos. 4,683,195 and 4,683,202), COLD-PCR (Li et al. (2008) Nat. Med. 14:579), ligase chain reaction (Barany, 1991, Proc. Natl. Acad. Sci. USA, 88:189), self sustained sequence replication (Guatelli et al., 1990, Proc. Natl.

Acad. Sci. USA 87:1874), transcriptional amplification system (Kwoh et al. (1989) Proc. Natl. Acad. Sci. USA 86:1173), Q- Beta Replicase (Lizardi et al. (1988) Bio/Technology 6:1197), rolling circle replication (U.S. Patent No. 5,854,033) or any other nucleic acid amplification method, followed by the

5 detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers. As used herein, amplification primers are defined as being a pair of nucleic acid molecules that can anneal to 5' or 3' regions of a gene (plus and minus strands, respectively,

10 or vice-versa) and contain a short region in between. In general, amplification primers are from about 10 to 30 nucleotides in length and flank a region from about 50 to 200 nucleotides in length. Under appropriate conditions and with appropriate reagents, such primers permit the amplification of a nucleic acid molecule comprising the nucleotide sequence flanked by the primers.

15 **[0098]** For *in situ* methods, mRNA does not need to be isolated from the sample (e.g., a bodily fluid (e.g., blood cells)) prior to detection. In such methods, a cell or tissue sample is prepared/processed using known histological methods. The sample is then immobilized on a support, typically a glass slide, and then contacted with a probe that can hybridize to mRNA that encodes the marker.

20 **[0099]** As an alternative to making determinations based on the absolute expression level of the marker, determinations may be based on the normalized expression level of the marker. Expression levels are normalized by correcting the absolute expression level of a marker by comparing its expression to the expression of a gene that is not a marker, e.g., a housekeeping gene that is

25 constitutively expressed. Suitable genes for normalization include housekeeping genes such as the actin gene, or epithelial cell- specific genes. This normalization allows the comparison of the expression level in a patient sample from one source to a patient sample from another source, e.g., to compare a population of phagocytic from an individual to a population of non-phagocytic cells from the

30 individual.

[0100] In one embodiment of this invention, a protein or polypeptide corresponding to a marker is detected. In certain embodiments, an agent for

detecting a protein or polypeptide can be an antibody capable of binding to the polypeptide, such as an antibody with a detectable label. As used herein, the term “labeled,” with regard to a probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin. Antibodies can be polyclonal or monoclonal. An intact antibody, or a fragment thereof (e.g., Fab or F(ab')₂) can be used. In one format, antibodies, or antibody fragments, can be used in methods such as Western blots or immunofluorescence techniques to detect the expressed proteins. In such uses, it is generally preferable to immobilize either the antibody or proteins on a solid support. Suitable solid phase supports or carriers include any support capable of binding an antigen or an antibody. Well known supports or carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, magnetite and the like.

[0101] A variety of formats can be employed to determine whether a sample contains a protein that binds to a given antibody. Examples of such formats include, but are not limited to, competitive and non-competitive immunoassay, enzyme immunoassay (EIA), radioimmunoassay (RIA), antigen capture assays, two-antibody sandwich assays, Western blot analysis, enzyme linked immunoabsorbant assay (ELISA), a planar array, a colorimetric assay, a chemiluminescent assay, a fluorescent assay, and the like. Immunoassays, including radioimmunoassays and enzyme-linked immunoassays, are useful in the methods of the present invention. A skilled artisan can readily adapt known protein/antibody detection methods for use in determining whether cells (e.g., bodily fluid cells such as blood cells) express a marker of the present invention.

[0102] One skilled in the art will know many other suitable carriers for binding antibody or antigen, and will be able to adapt such support for use with the present invention. For example, protein isolated from cells (e.g., bodily fluid cells such as

blood cells) can be run on a polyacrylamide gel electrophoresis and immobilized onto a solid phase support such as nitrocellulose. The support can then be washed with suitable buffers followed by treatment with the detectably labeled antibody. The solid phase support can then be washed with the buffer a second time to
5 remove unbound antibody. The amount of bound label on the solid support can then be detected by conventional means.

[0103] In certain exemplary embodiments, assays are provided for diagnosis, prognosis, assessing the risk of developing cancer, assessing the efficacy of a treatment, monitoring the progression or regression of cancer, and identifying a
10 compound capable of ameliorating or treating cancer. An exemplary method for these methods involves obtaining a bodily fluid sample from a test subject, isolating phagocytes and non-phagocytes, and contacting the phagocytes and non-phagocytes with a compound or an agent capable of detecting one or more of the markers of the disease or condition, *e.g.*, marker nucleic acid (*e.g.*, mRNA,
15 genomic DNA), marker peptide (*e.g.*, polypeptide or protein), marker lipid (*e.g.*, cholesterol), or marker metabolite (*e.g.*, creatinine) such that the presence of the marker is detected. In one embodiment, an agent for detecting marker mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to marker mRNA or genomic DNA. The nucleic acid probe can be, for example, a full-
20 length marker nucleic acid or a portion thereof. Other suitable probes for use in the diagnostic assays of the invention are described herein.

[0104] As used herein, a compound capable of ameliorating or treating cancer can include, without limitations, any substance that can improve symptoms or prognosis, prevent progression of the cancer, promote regression of the cancer, or
25 eliminate the cancer.

[0105] The methods of the invention can also be used to detect genetic alterations in a marker gene, thereby determining if a subject with the altered gene is at risk for developing cancer characterized by misregulation in a marker protein activity or nucleic acid expression. In certain embodiments, the methods include
30 detecting, in phagocytes, the presence or absence of a genetic alteration characterized by an alteration affecting the integrity of a gene encoding a marker peptide and/or a marker gene. For example, such genetic alterations can be

detected by ascertaining the existence of at least one of: 1) a deletion of one or more nucleotides from one or more markers genes; 2) an addition of one or more nucleotides to one or more markers genes; 3) a substitution of one or more nucleotides of one or more markers genes, 4) a chromosomal rearrangement of one or more markers genes; 5) an alteration in the level of a messenger RNA transcript of one or more markers genes; 6) aberrant modification of one or more markers genes, such as of the methylation pattern of the genomic DNA; 7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of one or more markers genes; 8) a non-wild type level of a one or more markers proteins; 9) allelic loss of one or more markers genes; and 10) inappropriate post-translational modification of one or more markers proteins. As described herein, there are a large number of assays known in the art which can be used for detecting alterations in one or more markers genes.

[0106] In certain embodiments, detection of the alteration involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g., U.S. Patent Nos. 4,683,195, 4,683,202 and 5,854,033), such as real-time PCR, COLD-PCR (Li et al. (2008) Nat. Med. 14:579), anchor PCR, recursive PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran et al. (1988) Science 241:1077; Prodromou and Pearl (1992) Protein Eng. 5:827; and Nakazawa et al. (1994) Proc. Natl. Acad. Sci. USA 91:360), the latter of which can be particularly useful for detecting point mutations in a marker gene (see Abravaya et al. (1995) Nucleic Acids Res. 23:675). This method can include the steps of collecting a sample of cell free bodily fluid from a subject, isolating nucleic acid (e.g., genomic, mRNA or both) from the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a marker gene under conditions such that hybridization and amplification of the marker gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

[0107] Alternative amplification methods include: self sustained sequence replication (Guatelli et al., (1990) Proc. Natl. Acad. Sci. USA 87:1874), transcriptional amplification system (Kwoh et al., (1989) Proc. Natl. Acad. Sci. USA 86:1173), Q Beta Replicase (Lizardi et al. (1988) Bio-Technology 6:1197), or
5 any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

[0108] In an alternative embodiment, mutations in one or more markers genes
10 from a sample can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, optionally amplified, digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA.
15 Moreover, the use of sequence specific ribozymes (see, for example, U.S. Pat. No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

[0109] In other embodiments, genetic mutations in one or more of the markers described herein can be identified by hybridizing a sample and control nucleic
20 acids, e.g., DNA or RNA, to high density arrays containing hundreds or thousands of oligonucleotides probes (Cronin et al. (1996) Human Mutation 7: 244; Kozal et al. (1996) Nature Medicine 2:753). For example, genetic mutations in a marker nucleic acid can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin, M. T. et al. supra. Briefly, a first
25 hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized
30 probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

- [0110] In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence a marker gene and detect mutations by comparing the sequence of the sample marker gene with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxam and Gilbert ((1977) Proc. Natl. Acad. Sci. USA 74:560) or Sanger ((1977) Proc. Natl. Acad. Sci. USA 74:5463). It is also contemplated that any of a variety of automated sequencing procedures can be utilized when performing the diagnostic assays ((1995) Biotechniques 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen et al. (1996) Adv. Chromatogr. 36:127-162; and Griffin et al. (1993) Appl. Biochem. Biotechnol. 38:147).
- [0111] Other methods for detecting mutations in a marker gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes (Myers et al. (1985) Science 230:1242).
- In general, the art technique of “mismatch cleavage” starts by providing heteroduplexes formed by hybridizing (labeled) RNA or DNA containing the wild-type marker sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent which cleaves single-stranded regions of the duplex such as which will exist due to base pair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA hybrids treated with S1 nuclease to enzymatically digesting the mismatched regions. In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. See, for example, Cotton et al. (1988) Proc. Natl. Acad. Sci. USA 85:4397; Saleeba et al. (1992) Methods Enzymol. 217:286. In one embodiment, the control DNA or RNA can be labeled for detection.
- [0112] In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called “DNA mismatch repair” enzymes) in defined systems for detecting and

- mapping point mutations in marker cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu et al. (1994) *Carcinogenesis* 15:1657). According to an exemplary embodiment, a
- 5 probe based on a marker sequence, e.g., a wild-type marker sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme, and the cleavage products, if any, can be detected from electrophoresis protocols or the like. See, for example, U.S. Patent No. 5,459,039.
- 10 **[0113]** In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in marker genes. For example, single strand conformation polymorphism (SSCP) may be used to detect differences in electrophoretic mobility between mutant and wild type nucleic acids (Orita et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:2766, see also Cotton (1993) *Mutat. Res.* 285:125; and
- 15 Hayashi (1992) *Genet. Anal. Tech. Appl.* 9:73). Single-stranded DNA fragments of sample and control marker nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or
- 20 detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In one embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen et al. (1991) *Trends Genet.* 7:5).
- 25 **[0114]** In yet another embodiment the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (DGGE) (Myers et al. (1985) *Nature* 313:495). When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp
- 30 of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to

identify differences in the mobility of control and sample DNA (Rosenbaum and Reissner (1987) Biophys. Chem. 265:12753).

[0115] Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification
5 or selective primer extension. For example, oligonucleotide primers may be prepared in which the known mutation is placed centrally and then hybridized to target DNA under conditions which permit hybridization only if a perfect match is found (Saiki et al. (1986) Nature 324:163; Saiki et al. (1989) Proc. Natl. Acad. Sci. USA 86:6230). Such allele specific oligonucleotides are hybridized to PCR
10 amplified target DNA or a number of different mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

[0116] Alternatively, allele specific amplification technology which depends on selective PCR amplification may be used in conjunction with the instant invention.
15 Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization) (Gibbs et al. (1989) Nucl. Acids Res. 17:2437) or at the extreme 3' end of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (Prossner (1993) Tibtech 11:238). In
20 addition it may be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection (Gasparini et al. (1992) Mol. Cell Probes 6:1). It is anticipated that in certain embodiments amplification may also be performed using Taq ligase for amplification (Barany (1991) Proc. Natl. Acad. Sci. USA 88:189). In such cases, ligation will occur only if there is a perfect
25 match at the 3' end of the 5' sequence making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

[0117] An exemplary method for detecting the presence or absence of an analyte (e.g., DNA, RNA, protein, polypeptide, or the like) corresponding to a marker of
30 the invention in a biological sample involves obtaining a bodily fluid sample (e.g., blood) from a test subject and contacting the bodily fluid sample with a compound or an agent capable of detecting one or more markers. Detection methods

described herein can be used to detect one or more markers in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of mRNA include Northern hybridizations and in situ hybridizations. *In vitro* techniques for detection of a polypeptide corresponding to a marker of the invention include

5 enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. *In vitro* techniques for detection of genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of a polypeptide corresponding to a marker of the invention include introducing into a subject a labeled antibody directed against the

10 polypeptide. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Because each marker is also an analyte, any method described herein to detect the presence or absence of a marker can also be used to detect the presence or absence of an analyte.

15 **[0118]** The markers useful in the methods of the invention can include any mutation in any one of the markers. Mutation sites and sequences can be identified, for example, by databases or repositories of such information, *e.g.*, The Human Gene Mutation Database (www.hgmd.cf.ac.uk), the Single Nucleotide Polymorphism Database (dbSNP, www.ncbi.nlm.nih.gov/projects/SNP), and the

20 Online Mendelian Inheritance in Man (OMIM) website (www.ncbi.nlm.nih.gov/omim).

[0119] The present invention also provides kits that comprise marker detection agents that detect at least one or more of the cancer markers described herein.

[0120] The present invention also provides methods of treating or preventing

25 cancer in a subject comprising administering to said subject an agent that modulates the activity or expression or disrupts the function of at least one or more of the markers of this invention.

[0121] The one or more markers identified by this invention (*e.g.*, markers in Tables 1-8) may be used in the treatment of cancer. For example, a marker (*e.g.*, a

30 protein or gene) identified by the present invention may be used as a molecular target for a therapeutic agent. A marker identified by the invention also may be used in any of the other methods of the invention, *e.g.*, for monitoring the

- progression or regression of a disease or condition. In certain embodiments, the one or more markers identified by the methods of this invention may have therapeutic potential. For example, if a marker is identified as being up-regulated (or down-regulated), see, for example, the up-regulated (or down-regulated) markers in Tables 1-8, or activated (or inhibited) in phagocytic cells from a subject having cancer, a compound or an agent that is capable of down-regulating (or up-regulating) or inhibiting (or activating) said marker may be useful in treating cancer. Similarly, a gene protein expression level, a protein expression level, or a combination thereof may be useful in this aspect of the invention.
- 5
- 10 **[0122]** Unless otherwise defined herein, scientific and technical terms used in this application shall have the meanings that are commonly understood by those of ordinary skill in the art. Generally, nomenclature used in connection with, and techniques of, cell and tissue culture, molecular biology, cell and cancer biology, neurobiology, neurochemistry, virology, immunology, microbiology,
- 15 pharmacology, genetics and protein and nucleic acid chemistry, described herein, are those well known and commonly used in the art.
- [0123]** All of the above, and any other publications, patents and published patent applications referred to in this application are specifically incorporated by reference herein. In case of conflict, the present specification, including its
- 20 specific definitions, will control.
- [0124]** Throughout this specification, the word “comprise” or variations such as “comprises” or “comprising” will be understood to imply the inclusion of a stated integer (or components) or group of integers (or components), but not the exclusion of any other integer (or components) or group of integers (or
- 25 components).
- [0125]** The singular forms “a,” “an,” and “the” include the plurals unless the context clearly dictates otherwise.
- [0126]** The term “including” is used to mean “including but not limited to.” “Including” and “including but not limited to” are used interchangeably.
- 30 **[0127]** It is to be understood that the embodiments of the present invention which have been described are merely illustrative of some of the applications of the principles of the present invention. Numerous modifications may be made by

those skilled in the art based upon the teachings presented herein without departing from the true spirit and scope of the invention.

[0128] The following examples are set forth as being representative of the present invention. These examples are not to be construed as limiting the scope of the invention as these and other equivalent embodiments will be apparent in view of the present disclosure and accompanying claims.

[0129] **Examples**

[0130] **Example 1: Microarray analysis of head and neck cancer patients**

[0131] Study Population

10 [0132] Blood samples were collected from a mixed population of prostate cancer and head and neck cancer patients and from blood donors without head and neck cancer. Approximately 10 ml of blood was collected from each patient into purple top blood collection EDTA tubes (BD Biosciences, CA). Within 3 hours, macrophages, neutrophils and T cells were isolated from each blood sample and
15 total RNA was extracted and purified on the same day. The healthy control blood samples were obtained from apheresis collars of anonymous platelet donors. Gender determination of the blood donors was performed by PCR using two sets of primers, SRY primers (Forward: 5'-CAG TGT GAA ACG GGA GAA AAC AG-3'; Reverse: 5'-ACT TCG CTG CAG AGT ACC GAA G-3') amplifying a 336 bp
20 fragment on Y chromosome and AR6 primers (Forward: 5'-CAA TCA GAG ACA TTC CCT CTG G-3'; Reverse: 5'-AGT GGT CCT CTC TGA ATC TC-3') amplifying a 267 bp fragment on X chromosome (males have both fragments amplified; females have only one). The PCR (36 cycles) was done under the conditions of 95 °C for 45 seconds, 56 °C for 45 seconds, and 72 °C for 45
25 seconds.

[0133] Isolation of Macrophages (M), Neutrophils (N), and T Cells (TC) from Whole Blood

[0134] 7 mL of 1X PBS containing 2% FBS and 2 mM EDTA were added to approximately 5 mL of whole blood, the sample centrifuged (2,000 RPM, 10
30 minutes at 20°C). The buffy coat was removed and centrifuged (2,000 RPM, 10 minutes at 20°C). The cell pellet was then suspended in ~1 mL of PBS and transferred to a 1.5 mL microfuge tube. Next, macrophages, neutrophils, and T

cells were isolated using magnetic beads coated with antibodies specific to each of the three cell types (positive cell depletion). Cells were separated from the buffy coat always in the following sequence: 1) macrophages; 2) neutrophils; and 3) T cells (changing the order did not alter the RNA yield and quality). The freshly
5 isolated white blood cell samples (in ~1 mL PBS) were incubated (25 min, 4°C, constant shaking) first with anti-monocyte coated Dynabeads® (CD14 - Cat. No. 11149D, Life Technologies), then with anti-neutrophil coated Dynabeads® (CD15 - Cat. No. 11137D, Life Technologies), and finally with anti- T cell coated Dynabeads® (CD2 Pan T - Cat. No. 11159D, Life Technologies). Following each
10 incubation, the bead-bound cells were separated using a magnet. The purity of these white blood cell subpopulations, which per manufacturer's specifications (Life Technologies) is >95%, was evident from the unique gene expression pattern obtained (cluster analysis). As soon as each white blood cell subpopulation was isolated, the magnetic bead bound cells were washed with 1x PBS and lysed in
15 Trizol®. The fractionation and subsequent lysis of all the three types of cells were completed in less than 2 hours after the isolation of the buffy coat.

[0135] Total RNA Isolation

[0136] Total RNA was extracted from cells and tissues with Trizol® and the Pure-Link RNA isolation kit (Cat. # 12183018A, Life Technologies). The quantity
20 and purity of the RNA samples were determined on a Bioanalyzer 2100 (Agilent Technologies) and the Degradometer software (version 1.41). In general, the RIN and 28s/18s ratios were always found to be in the satisfactory range, ≥ 9 and ≥ 1.9 , respectively.

[0137] Whole Genome Microarray Data Analysis

[0138] Total RNA from macrophages, neutrophils, T cells, tumor tissue (TT), and “normal” tissue (NT) of cancer patients, and from macrophages, neutrophils, and T cells extracted from healthy male blood donors were used in gene expression
25 profiling. Biotinylated cDNA probes were prepared from 100 ng of each RNA sample, fragmented, and hybridized with Human Gene 1.0 ST chip (Affymetrix).
30 Array signals of fluorescence were scanned, captured and recorded as CEL files. All the processing and analysis of the data were done using R 4.9 and Bioconductor software packages. To obtain the log₂ transformed expression levels, the raw data

files obtained in CEL file format were pre-processed using the oligo package and the RMA (robust multichip average algorithm) routine to background correct, quantile normalize and summarize at the core level.

[0139] Example 2: Statistical analysis of microarray data

5 **[0140]** Working with microarray data can be challenging because large numbers of genes can increase the likelihood of false positives, while a small number of samples can lead to overfitting. These issues can be overcome by using statistical methods to reduce the false rate of positives and using independent training and test data sets (e.g., cross-validation) to avoid overfitting. In particular, instead of
10 using a “typical” 5% significance level, the false discovery rate (FDR) can be controlled to ensure that only 5% of the genes that are discovered are false positives, and Empirical Bayesian estimates can be used to improve test statistics.

[0141] Because an overfit model will perform poorly on an independent test set, a good test of the fit of a model is how well it performs on an independent test set.
15 For small sample sizes, splitting data into test and training sets may leave too small of a data set for good training. This issue can be solved by using cross-validation, which splits the data into K-folds, trains the method on K-1 of the folds, and tests the method on the last fold. Figure 1 depicts a diagram of a three-fold cross validation, wherein the diagnostic accuracy is averaged from the three splits. The
20 ideal split for cross-validation is 10-fold for accurate and precise estimates of diagnostic accuracy. In a 10-fold cross validation, however, there are more than 10 splits because there are many choices for which data points go into the folds. For example, with the microarray data collected as described above, there are 50,979,600 ways to form 90% training/10% testing data sets.

25 **[0142]** The Empirical Bayesian method was used as follows:

1. The differential gene expression (DE) of phagocytes (macrophages or neutrophils) vs. T cells was calculated for each gene. DE is expressed as the log of the ratio of phagocyte to T cell expression: $DE = \log(GE_P/GE_{TC})$, where GE_P is phagocyte gene expression and GE_{TC} is T cell gene
30 expression.

2. The mean DE was compared in cancer and control patients with a two-sample t-test. Empirical Bayes estimates of the test statistics “shrink” these toward zero.
3. Calculate a diagnostic signature with K genes:

$$S = \sum_{i=1}^K w_i (DE_i - \mu_i)$$

5

If $S > 0$, then the patient was diagnosed with cancer.

4. The number of genes K to include in the signature was determined by comparing misclassification rates in independent test sets with cross-validation.
- 10 Errors were calculated using an average of 1-sensitivity and 1-specificity, and the cross-validated error was used to select markers.
- [0143]** Using the above methods, the markers associated with cancer in macrophages vs. T cells (Tables 1 and 3) and the markers associated with cancer in neutrophils vs. T cells (Tables 2 and 4) were identified. Of these, two specific
- 15 signatures of four markers (for macrophages) and two specific signatures of eight markers (for neutrophils) also were identified that give especially high sensitivity and specificity. For example, a four marker signature (C-Macro 1-4) from macrophages has a sensitivity of 99.86% and a specificity of 99.84%. The four genes identified were:

- 20
1. P2RY10;
 2. TNFAIP3;
 3. CXCR1; and
 4. DNAJB1.

- [0144]** A different four marker signature (C-Macro 101-104) from macrophages
- 25 has a sensitivity of 100% and a specificity of 98.1%. The four genes identified were:

- 30
1. P2RY10;
 2. TNFAIP3;
 3. TIPARP; and
 4. CXCR1.

[0145] An eight marker signature (C-Neutro 1-8) from neutrophils has a sensitivity of 97.7% or 99.2% and a specificity of 99.96% or 94.6% (dependent on data set analyzed). Exemplary genes identified were:

1. EEF1A1;
- 5 2. RPL23A;
3. RPL14; and
4. RPL3.

[0146] Figure 2 shows a summary of the cancer markers identified from macrophages and from neutrophils, as compared to T cells from the same individuals, for the C-Macro1-100, C-Macro 101-200, C-Neutro 1-100, and C-Neutro 101-200 markers. Specifically, average error, sensitivity, and specificity values are given for a four marker panel from C-Macro 1-100 (C-Macro 1-4), a four marker panel from C-Macro 101-200 (C-Macro 101-104), an eight marker panel from C-Neutro 1-100 (C-Neutro 1-8), and an eight marker panel from C-Neutro 101-200 (C-Neutro 101-108). Figure 3 demonstrates the power of a paired within-subject (phagocyte to non-phagocyte) comparison to detect cancer as compared to phagocytes not paired with T cell data for comparison. The paired approach (comparing macrophage or neutrophils to T cell expression) is better than the phagocyte gene expression alone.

[0147] Similar methods were used to distinguish between prostate cancer and head and neck cancer, indicating an ability to differentiate between various cancers (Figure 6). For example, a 12 marker panel (PC/HNC-Macro 1-12) was able to distinguish between the cancers with a sensitivity of 75% and a specificity of 92.1%. A 10 marker panel (PC/HNC-Macro 101-110) was able to distinguish between the cancers with a sensitivity of 77.5% and a specificity of 90.7%. A nine marker panel (PC/HNC-Neutro 1-9) was able to distinguish between the cancers with a sensitivity of 87.5% and a specificity of 94.4% or a sensitivity of 88% and a specificity of 92.1%, dependent on the data set used to calculate the specificity and sensitivity. Markers distinguishing the cancers are listed in Tables 5-8.

30 [0148] **Example 3: Additional validation**

[0149] Additional validation is performed by validating the gene signature on a new data set of ~50 cancer cases and 50 controls. Final validation is performed by

- estimating the sensitivity and specificity of the final gene signature on a large sample. For example, 195 cases and 195 controls can be used to estimate a sensitivity/specificity of at least 97.5% with a 95% margin of error no more than 5%. A challenge in designing a final validation study is that although cancer
- 5 patients are pure, controls may have up to 20% false negatives. A statistical issue is that, while estimating sensitivity is not a problem, specificity has an upper bound of 80%. The solution is to purify the control set of patients. A purification method uses secondary screening of all controls, wherein three methylated gene marker tests are used to purify the control test set:
- 10 1. GST-Pi (sensitivity = 95%, specificity =85%)
 2. RAR-2b (sensitivity = 95%, specificity =48%)
 3. APC (sensitivity = 95%, specificity =50%)
- in subjects with two serial negative biopsies. A second purification method is depicted in Figure 4, and a comparison of purification methods is shown in Figure
- 15 5.

Table 1. Cancer (C)-Macro Markers 1-100
(DR = down-regulated; UR= up-regulated)

C-Macro	Transcript Cluster ID	Gene Name	Cancer Mean	Control Mean	Pattern
1	8168524	P2RY10	0.142295261	0.029828375	Cancer UR
2	8122265	TNFAIP3	0.620558801	0.211725335	Cancer UR
3	8083569	TIPARP	1.066219276	0.464364876	Cancer UR
4	8058905	CXCR1	8.640447656	0.784600913	Cancer UR
5	7923547	CHI3L1	9.047300553	1.251169207	Cancer UR
6	7903592	KIAA1324	1.102801214	0.139725719	Cancer UR
7	8034837	DNAJB1	0.576263809	0.308941315	Cancer UR
8	7898693	ALPL	6.555677655	1.159450091	Cancer UR
9	7962516	SLC38A1	0.042303973	0.020268951	Cancer UR
10	8037205	CEACAM1	2.109889397	0.59390842	Cancer UR
11	8048227	CXCR2	10.18331766	1.417588198	Cancer UR
12	7961371	DUSP16	0.319711843	0.149852042	Cancer UR
13	7952036	MPZL3	1.225836577	0.541385722	Cancer UR
14	8016540	PHOSPHO1	4.298122261	1.192652558	Cancer UR
15	7904361	FAM46C	0.296616555	0.079575558	Cancer UR
16	8062927	PI3	5.243767336	1.068298221	Cancer UR
17	8174361	TSC22D3	0.976497207	0.618688293	Cancer UR
18	7996100	GPR97	4.671205034	1.128190892	Cancer UR
19	7899253	ZDHHC18	1.667577169	0.686773779	Cancer UR
20	8097461	CCRN4L	1.377673545	3.607902835	Cancer DR
21	8095728	EREG	1.74346798	10.7290871	Cancer DR
22	8070720	ICOSLG	1.478702161	3.309822542	Cancer DR
23	7972805	RAB20	2.240810322	5.494866343	Cancer DR
24	7972557	GPR183	0.328174362	1.175191296	Cancer DR
25	7955589	NR4A1	2.46791005	6.575336512	Cancer DR
26	8104492	ROPN1L	2.223391575	0.731058369	Cancer UR
27	7974870	SNAP	1.091337333	1.842512128	Cancer DR

C-Macro	Transcript Cluster ID	Gene Name	Cancer Mean	Control Mean	Pattern
28	7923917	FAIM3	0.055258086	0.02956543	Cancer UR
29	8179263	TNF	2.222279222	0.866313835	Cancer UR
30	8177983	TNF	2.222279222	0.866313835	Cancer UR
31	8118142	TNF	2.222279222	0.866313835	Cancer UR
32	8063382	SNAI1	1.195271879	2.439730089	Cancer DR
33	7930413	DUSP5	1.223995731	0.560750904	Cancer UR
34	8114572	HBEGF	3.822283637	12.5273771	Cancer DR
35	8156848	NR4A3	1.04205333	2.683137433	Cancer DR
36	8146550	SDCBP	4.492902766	2.677202226	Cancer UR
37	8083494	MME	7.078918953	1.175410257	Cancer UR
38	8105778	PIK3R1	0.427143974	0.269772089	Cancer UR
39	7961142	OLR1	1.180394865	4.96332356	Cancer DR
40	8019877	SMCHD1	1.11968151	0.652468814	Cancer UR
41	8061373	GZF1	1.433635431	2.4459341	Cancer DR
42	8073148	ATF4	1.494118435	1.063953313	Cancer UR
43	8026564	KLF2	0.789802831	0.526489836	Cancer UR
44	8063115	MMP9	3.928091465	1.325455323	Cancer UR
45	7920575	PBXIP1	0.59365049	0.399025072	Cancer UR
46	8075316	OSM	1.705251418	4.034924134	Cancer DR
47	8145244	TNFRSF10C	4.966074677	1.75691825	Cancer UR
48	7926916	ZEB1	0.280321783	0.11262711	Cancer UR
49	7929616	FRAT1	2.713037389	1.442319574	Cancer UR
50	7937335	IFITM1	0.366663584	0.116950226	Cancer UR
51	8092691	BCL6	8.936174395	4.206275916	Cancer UR
52	8052654	PELI	4.576614868	2.190339783	Cancer UR
53	8001317	N4BP1	1.307102127	0.784554396	Cancer UR
54	8072328	SEC14L2	0.765474863	0.403431114	Cancer UR
55	7922846	FAM129A	2.033587218	0.811233648	Cancer UR
56	7945169	TMEM45B	0.890862427	0.325128823	Cancer UR

C-Macro	Transcript Cluster ID	Gene Name	Cancer Mean	Control Mean	Pattern
57	8114010	IRF1	1.676395661	0.97821653	Cancer UR
58	8026456	CYP4F	5.840371569	1.394455051	Cancer UR
59	8119898	VEGFA	3.45619434	7.36028069	Cancer DR
60	8119016	MAPK13	0.641745677	0.368678501	Cancer UR
61	8117330	HIST1H3A	0.793010285	0.36466297	Cancer UR
62	8116983	CD83	2.133493613	4.212633506	Cancer DR
63	8000482	XPO6	1.281972043	0.730075877	Cancer UR
64	7998931	ZNF200	1.112174483	0.747138578	Cancer UR
65	8019885	SMCHD1	1.200633257	0.694017136	Cancer UR
66	8044049	IL18RAP	0.294549577	0.048739981	Cancer UR
67	8026047	JUNB	2.905720462	1.471000804	Cancer UR
68	7904465	HIST2H2BA	0.958354791	0.679665447	Cancer UR
69	7968035	SPATA13	0.600235304	0.958662754	Cancer DR
70	7946559	GNG10	3.147322264	2.040359059	Cancer UR
71	8128111	UBE2J1	2.281911738	1.434763478	Cancer UR
72	8116484	MGAT1	1.688726651	2.506886835	Cancer DR
73	7978595	BAZ1A	1.624425641	1.082164671	Cancer UR
74	7954711	C12orf35	0.842465662	0.489425283	Cancer UR
75	8064438	NSFL1C	2.046051239	1.405613346	Cancer UR
76	8011968	MED31	0.96067137	0.716049164	Cancer UR
77	8126839	TNFRSF21	1.114430756	2.098652381	Cancer DR
78	8008795	C17orf71	1.32003243	0.857441229	Cancer UR
79	7961365	MANSC1	4.930594711	1.75981802	Cancer UR
80	8060344	TRIB3	0.968959893	1.298214832	Cancer DR
81	7975453	SNORD56B	3.263124503	8.224312945	Cancer DR
82	8017850	WIP1	2.071214436	3.599837341	Cancer DR
83	8015133	KRT23	4.606031457	1.387347323	Cancer UR
84	7925048	EGLN1	1.750158869	1.20878267	Cancer UR
85	8177951	HCG27	1.285204235	0.794781361	Cancer UR

C-Macro	Transcript Cluster ID	Gene Name	Cancer Mean	Control Mean	Pattern
86	7922474	KIAA0040	1.227497745	0.772043684	Cancer UR
87	8127145	ELOVL5	1.232977586	0.890616673	Cancer UR
88	7992811	MMP25	3.552104643	1.519284767	Cancer UR
89	8095986	ANXA3	3.983011045	1.048862261	Cancer UR
90	8078014	SLC6	4.119936224	2.632320297	Cancer UR
91	8112220	PDE4D	0.557821444	0.904855095	Cancer DR
92	7974920	SYNE2	0.093612027	0.03658116	Cancer UR
93	8042283	HSPC159	2.314967309	0.966672965	Cancer UR
94	8079140	SNRK	0.797956307	0.592571613	Cancer UR
95	7987192	SLC12A6	2.049713493	1.421765397	Cancer UR
96	7945944	RHOG	2.161060473	1.613390851	Cancer UR
97	7910387	RHOU	2.175928457	4.143592241	Cancer DR
98	8144931	ATP6V1B2	5.296368584	3.704075834	Cancer UR
99	7955578	GRASP	1.070816788	1.67638387	Cancer DR
100	8091120	GK5	0.560158387	1.071261752	Cancer DR

Table 2. Cancer (C)-Neutro Markers 1-100
(DR = down-regulated; UR= up-regulated)

C- Neutro	Transcript Cluster ID	Gene Name	Cancer Mean	Control Mean	Pattern
1	8180410	Unknown	0.200117368	0.62620984	Cancer DR
2	8138531	EEF1A1	0.282512943	0.716120804	Cancer DR
3	8091806	RPL23A	0.209146764	0.601822054	Cancer DR
4	8026440	RPL23A	0.207439433	0.58982973	Cancer DR
5	8005943	RPL23A	0.193981776	0.55638321	Cancer DR
6	7956743	RPL14	0.152976283	0.476827764	Cancer DR
7	8076209	RPL3	0.095800723	0.394600877	Cancer DR
8	8158952	EEF1A1	0.263489836	0.692649185	Cancer DR
9	8109821	RPL10	0.225708831	0.694652859	Cancer DR
10	8107470	PTMA	0.350617242	0.80118868	Cancer DR
11	7946812	RPS13	0.141007364	0.569520819	Cancer DR
12	7961022	PTMA	0.344097164	0.753347849	Cancer DR
13	8026868	RPL18A	0.166113205	0.59749077	Cancer DR
14	8153903	RPL8	0.240816134	0.631773105	Cancer DR
15	8061136	PTMA	0.305795426	0.799922277	Cancer DR
16	8180297	Unknown	0.124290069	0.479274426	Cancer DR
17	8013348	RPS2	0.223049514	0.537889078	Cancer DR
18	7954006	PTMA	0.347107612	0.751137136	Cancer DR
19	7986323	GLTSCR2	0.110719193	0.411911366	Cancer DR
20	8030351	RPL13A	0.210885832	0.533772034	Cancer DR
21	8038086	RPL18	0.157146191	0.560978174	Cancer DR
22	7998655	RPS2	0.227101453	0.543477946	Cancer DR
23	8180355	Unknown	0.201008287	0.586677482	Cancer DR
24	8177003	SLC25A6	0.192013685	0.732041435	Cancer DR
25	8171111	SLC25A6	0.192013685	0.732041435	Cancer DR
26	8116929	RPL15	0.185391799	0.542468444	Cancer DR

C- Neutro	Transcript Cluster ID	Gene Name	Cancer Mean	Control Mean	Pattern
27	7983843	TCF12	0.189013624	0.541511564	Cancer DR
28	7948679	EEF1G	0.168199575	0.572470282	Cancer DR
29	8034416	RPL10	0.148652387	0.466314396	Cancer DR
30	7942824	RPS28	0.17826043	0.577772835	Cancer DR
31	7901038	RPS8	0.171855552	0.506397724	Cancer DR
32	8025395	RPS28	0.175134497	0.569300667	Cancer DR
33	8154394	SNAPC3	0.211303422	0.726268944	Cancer DR
34	8005471	RPS28	0.174230395	0.565217789	Cancer DR
35	8116520	GNB2L1	0.165907937	0.610016882	Cancer DR
36	7990965	RPS17	0.110561259	0.39582654	Cancer DR
37	7990916	RPS17	0.110561259	0.39582654	Cancer DR
38	7986765	RPL5	0.152304713	0.521955359	Cancer DR
39	8076511	RPL5	0.158788106	0.48242009	Cancer DR
40	8024299	RPS15	0.221491241	0.606758584	Cancer DR
41	8173513	RPS4X	0.105175669	0.411688846	Cancer DR
42	8007441	RPL27	0.316362607	0.642964949	Cancer DR
43	8022170	RPL6	0.201984973	0.515685725	Cancer DR
44	8164100	RPL35	0.0933485	0.377004089	Cancer DR
45	8085026	RPL35A	0.17574271	0.479603464	Cancer DR
46	7966534	RPL6	0.206485906	0.516323625	Cancer DR
47	8047635	RPL12	0.185139522	0.479381403	Cancer DR
48	7966996	RPLP0	0.135218504	0.435746306	Cancer DR
49	8174710	RPL39	0.217960245	0.509260022	Cancer DR
50	7917906	RPL7	0.227772593	0.55562462	Cancer DR
51	8109222	RPL7	0.223001632	0.548404866	Cancer DR
52	8051066	MPV17	0.230477076	0.772766442	Cancer DR
53	7900585	YBX1	0.376598391	0.834191228	Cancer DR
54	7929593	RPL13AP5	0.283564504	0.67977926	Cancer DR
55	7903010	RPL5	0.200350234	0.478097833	Cancer DR

C- Neutro	Transcript Cluster ID	Gene Name	Cancer Mean	Control Mean	Pattern
56	7920317	ILF2	0.172903097	0.537870922	Cancer DR
57	8172154	RPS2	0.307753383	0.60299509	Cancer DR
58	8125750	RPL12	0.168321141	0.449114014	Cancer DR
59	8043100	TMSB10	0.284664181	0.74343566	Cancer DR
60	8127526	RPL39	0.213299506	0.498657094	Cancer DR
61	7899160	CD52	0.140959424	0.610247056	Cancer DR
62	8171834	RPL	0.181018132	0.445097281	Cancer DR
63	8099887	RPL9	0.178515873	0.439813779	Cancer DR
64	7990898	RPL	0.176553431	0.439403367	Cancer DR
65	8154359	RPL18A	0.261579743	0.647185895	Cancer DR
66	8063473	RPL12	0.180284493	0.46028389	Cancer DR
67	7968872	DNAJC15	0.160001377	0.560505108	Cancer DR
68	8151376	RPL7	0.228059938	0.540641513	Cancer DR
69	7990949	RPL	0.173290283	0.432316094	Cancer DR
70	8118594	HLA-DPB1	0.2140086	1.32168702	Cancer DR
71	7984562	RPLP1	0.206090636	0.545118248	Cancer DR
72	8073799	ATXN10	0.139438169	0.465493834	Cancer DR
73	7933760	CCDC6	0.16481272	0.663958305	Cancer DR
74	8179519	HLA-DPB1	0.194723793	1.373055482	Cancer DR
75	8115234	ANXA6	0.0909779	0.316090919	Cancer DR
76	8009561	RPL38	0.28474799	0.593379311	Cancer DR
77	8178220	HLA-DPB1	0.206034919	1.254074325	Cancer DR
78	7973056	APEX1	0.130524364	0.465327418	Cancer DR
79	7905099	VPS45	0.206658639	0.611894043	Cancer DR
80	8115147	CD74	0.686654926	2.484644609	Cancer DR
81	8115158	RPS14	0.225993945	0.530386289	Cancer DR
82	7965515	NDUFA12	0.289356859	0.63334209	Cancer DR
83	8022972	RPL7A	0.249408078	0.547837095	Cancer DR
84	7944152	IL10RA	0.180917717	0.617370448	Cancer DR

C- Neutro	Transcript Cluster ID	Gene Name	Cancer Mean	Control Mean	Pattern
85	7999827	RPS15A	0.2290889	0.461445511	Cancer DR
86	8092457	ALG3	0.277479476	0.782108011	Cancer DR
87	8175177	MBNL3	0.154675537	0.537895141	Cancer DR
88	8036777	FBL	0.109732495	0.337880818	Cancer DR
89	8109750	RPLP0	0.076085091	0.355200772	Cancer DR
90	8036602	ECH1	0.187702656	0.625284322	Cancer DR
91	7912956	RCC2	0.474885591	0.960941494	Cancer DR
92	7939368	TRIM44	0.27317842	0.714981248	Cancer DR
93	8154727	LOC138412	0.209352409	0.645267809	Cancer DR
94	7899957	ZMYM4	0.329032305	0.697634114	Cancer DR
95	7996947	CYB5B	0.251686767	0.49172633	Cancer DR
96	8108954	TCERG1	0.218743912	0.578860239	Cancer DR
97	8097782	RPS3A	0.226550254	0.462605419	Cancer DR
98	8027778	FXVD5	0.312492986	0.629101591	Cancer DR
99	8178802	HLA-DRB3	0.527574488	2.183588812	Cancer DR
100	7948667	AHNAK	0.116700841	0.527579932	Cancer DR

Table 3. Cancer (C)-Macro Markers 101-200

(DR = down-regulated; UR = up-regulated)

C-Macro	Transcript Cluster ID	Cancer Mean	Control Mean	Pattern
101	8168524	0.153565123	0.02973116	Cancer UR
102	8122265	0.65451964	0.211659366	Cancer UR
103	8058905	8.159848429	0.786213943	Cancer UR
104	8034837	0.592830277	0.310069566	Cancer UR
105	7962516	0.043864659	0.020371106	Cancer UR
106	7903592	1.016691857	0.140023828	Cancer UR
107	7961371	0.334929877	0.149471052	Cancer UR
108	7952036	1.215665166	0.546920278	Cancer UR
109	8174361	0.982773283	0.614693417	Cancer UR
110	7923547	8.44988585	1.246796188	Cancer UR
111	8083569	1.038466184	0.467897834	Cancer UR
112	7904361	0.310213391	0.081850402	Cancer UR
113	8052654	4.997303691	2.175406803	Cancer UR
114	8072328	0.816130407	0.404105838	Cancer UR
115	8048227	9.32646217	1.400245526	Cancer UR
116	8037205	1.974724116	0.606234932	Cancer UR
117	8097461	1.373449889	3.614577473	Cancer DR
118	8095728	1.723937963	10.32404888	Cancer DR
119	8070720	1.406125133	3.319307189	Cancer DR
120	8063382	1.191086463	2.442584039	Cancer DR
121	7972805	2.374694892	5.471720102	Cancer DR
122	7955589	2.316706032	6.488994673	Cancer DR
123	8146550	4.686195231	2.695596257	Cancer UR
124	7898693	6.190410798	1.169632616	Cancer UR
125	7961142	1.159568077	4.879358896	Cancer DR
126	8075316	1.610268994	4.011305782	Cancer DR
127	8105778	0.43390843	0.27076014	Cancer UR
128	7926916	0.306607049	0.115655221	Cancer UR
129	7929616	2.738416061	1.431015336	Cancer UR
130	8116983	1.924800954	4.173442204	Cancer DR
131	7930413	1.211626549	0.561848213	Cancer UR
132	8156848	1.036529951	2.63646172	Cancer DR
133	8016540	4.043433052	1.188715166	Cancer UR

C-Macro	Transcript Cluster ID	Cancer Mean	Control Mean	Pattern
134	7974870	1.134909339	1.83735522	Cancer DR
135	8119898	3.248986904	7.317821066	Cancer DR
136	7978595	1.699976711	1.082763864	Cancer UR
137	7923917	0.057763804	0.029309611	Cancer UR
138	7900426	1.886251511	1.282061493	Cancer UR
139	8092691	8.618306924	4.197890572	Cancer UR
140	7945169	0.98233248	0.326074124	Cancer UR
141	7899253	1.638257144	0.692376881	Cancer UR
142	8062927	5.357679049	1.090689052	Cancer UR
143	8114010	1.697444502	0.97716878	Cancer UR
144	8019877	1.121857937	0.649599886	Cancer UR
145	7998931	1.116071169	0.752216983	Cancer UR
146	8099797	2.155278179	1.386902202	Cancer UR
147	8072346	0.845581633	0.598417483	Cancer UR
148	8114572	3.947945084	12.22822301	Cancer UR
149	8001317	1.363460231	0.792769423	Cancer UR
150	7972557	0.384337192	1.159820179	Cancer DR
151	8126839	1.082045018	2.076217645	Cancer DR
152	7922846	1.972746472	0.812013092	Cancer UR
153	8026564	0.810093826	0.531791285	Cancer UR
154	8061373	1.440273639	2.415653102	Cancer DR
155	8000482	1.302291393	0.728353956	Cancer UR
156	8019885	1.237616935	0.697370351	Cancer UR
157	7937335	0.347223275	0.115249042	Cancer UR
158	8036710	1.409486723	0.997985542	Cancer UR
159	7996100	4.242365648	1.129208861	Cancer UR
160	8117330	0.782966939	0.370185013	Cancer UR
161	7999642	1.172128698	0.85020082	Cancer UR
162	7946559	3.246861747	2.0341812	Cancer UR
163	7929032	0.960871798	0.582991036	Cancer UR
164	8017850	2.029468392	3.597053643	Cancer DR
165	8073148	1.50642594	1.064442258	Cancer UR
166	8104492	2.070359751	0.742388	Cancer UR
167	8026047	2.946231382	1.485086606	Cancer UR
168	8011968	0.976222262	0.717565997	Cancer UR
169	8179263	2.049647517	0.870811463	Cancer UR
170	8177983	2.049647517	0.870811463	Cancer UR
171	8118142	2.049647517	0.870811463	Cancer UR
172	7974920	0.101633362	0.037091839	Cancer UR

C-Macro	Transcript Cluster ID	Cancer Mean	Control Mean	Pattern
173	7961524	0.547581956	0.336258312	Cancer UR
174	8119016	0.65406192	0.374428114	Cancer UR
175	7909610	1.292924113	2.24858374	Cancer DR
176	7910387	2.083190219	4.109913744	Cancer DR
177	8094743	0.066620742	0.035169584	Cancer UR
178	8063115	3.60760332	1.32364083	Cancer UR
179	8063583	0.677459088	0.421285936	Cancer UR
180	7922474	1.279341367	0.777989719	Cancer UR
181	8124280	0.858619614	0.564538559	Cancer UR
182	7904465	0.957690902	0.686424857	Cancer UR
183	7950743	0.376342519	0.192646609	Cancer UR
184	7920575	0.588950106	0.398196223	Cancer UR
185	8064438	2.07633005	1.40433679	Cancer UR
186	8127145	1.265346585	0.897546589	Cancer UR
187	7987192	2.130050384	1.416637683	Cancer UR
188	8128111	2.215314765	1.429969971	Cancer UR
189	8025672	0.967268368	0.564381668	Cancer UR
190	8144931	5.367951956	3.726492662	Cancer UR
191	7945944	2.201257131	1.621397075	Cancer UR
192	8147206	2.389349073	3.927953313	Cancer DR
193	8067113	1.516532784	0.895542524	Cancer UR
194	8112202	1.198054022	2.831920376	Cancer DR
195	8083494	5.675927609	1.157097223	Cancer UR
196	7975453	3.029316639	8.237084986	Cancer DR
197	8124416	0.54131521	0.333145773	Cancer UR
198	7968035	0.609343962	0.950857801	Cancer DR
199	8078214	1.39328735	1.096726399	Cancer UR
200	8177951	1.2883823	0.784172225	Cancer UR

Table 4. Cancer (C)-Neutro Markers 101-200
(DR = down-regulated; UR= up-regulated)

C-Neutro	Transcript Cluster ID	Cancer mean	Control mean	Pattern
101	8180410	0.188422619	0.624612673	Cancer DR
102	8158952	0.234952625	0.686421334	Cancer DR
103	8138531	0.252644062	0.711316246	Cancer DR
104	8091806	0.194687503	0.597720357	Cancer DR
105	8076209	0.085043976	0.39237551	Cancer DR
106	7956743	0.141527881	0.473806633	Cancer DR
107	8026440	0.193099803	0.583462463	Cancer DR
108	8005943	0.180319775	0.550334281	Cancer DR
109	7983843	0.179815295	0.538902593	Cancer DR
110	7998655	0.218063184	0.537001903	Cancer DR
111	8013348	0.215628436	0.53311271	Cancer DR
112	8107470	0.326563344	0.798079666	Cancer DR
113	8180297	0.111743613	0.473432388	Cancer DR
114	8026868	0.156296284	0.59044297	Cancer DR
115	8180355	0.189798273	0.584095306	Cancer DR
116	8154394	0.192772688	0.727012806	Cancer DR
117	7986323	0.102313056	0.408935461	Cancer DR
118	7946812	0.13035898	0.563435992	Cancer DR
119	8153903	0.221302363	0.623162146	Cancer DR
120	8061136	0.280006301	0.794624343	Cancer DR
121	7942824	0.159541548	0.571170935	Cancer DR
122	8005471	0.155991628	0.558482978	Cancer DR
123	8025395	0.156722074	0.562233499	Cancer DR
124	7920317	0.154072165	0.5395029	Cancer DR
125	8173513	0.098448384	0.408782488	Cancer DR
126	8177003	0.16952303	0.728285779	Cancer DR
127	8171111	0.16952303	0.728285779	Cancer DR

C-Neutro	Transcript Cluster ID	Cancer mean	Control mean	Pattern
128	8085026	0.156368539	0.47693372	Cancer DR
129	8116929	0.176183303	0.539878645	Cancer DR
130	8030351	0.204267072	0.527648557	Cancer DR
131	8115158	0.19632498	0.529445703	Cancer DR
132	7948679	0.141781081	0.571490186	Cancer DR
133	7961022	0.341615207	0.748141158	Cancer DR
134	8051066	0.201946548	0.767347807	Cancer DR
135	7990965	0.102895902	0.391261856	Cancer DR
136	7990916	0.102895902	0.391261856	Cancer DR
137	7954006	0.343890276	0.743532451	Cancer DR
138	8115234	0.080336851	0.314830287	Cancer DR
139	8034416	0.135425646	0.463038573	Cancer DR
140	7900585	0.35279693	0.833649024	Cancer DR
141	8024299	0.212194413	0.607899485	Cancer DR
142	8038086	0.132433721	0.556045454	Cancer DR
143	8109821	0.204400245	0.686951642	Cancer DR
144	8043100	0.257335575	0.741689456	Cancer DR
145	8164100	0.087055859	0.365288977	Cancer DR
146	7899160	0.124157883	0.611959909	Cancer DR
147	8174710	0.217748847	0.509944666	Cancer DR
148	8116520	0.141135688	0.604419962	Cancer DR
149	8101429	0.072601984	0.441486372	Cancer DR
150	8127526	0.212672174	0.497601294	Cancer DR
151	8109750	0.061002259	0.352037718	Cancer DR
152	8009561	0.262817813	0.582845797	Cancer DR
153	8076511	0.151999944	0.478349935	Cancer DR
154	8092457	0.2563994	0.789063726	Cancer DR
155	7966996	0.133814518	0.433354821	Cancer DR
156	8007441	0.318916843	0.63569922	Cancer DR

C-Neutro	Transcript Cluster ID	Cancer mean	Control mean	Pattern
157	7948667	0.099027232	0.526841184	Cancer DR
158	8036602	0.167726762	0.625645255	Cancer DR
159	7944152	0.158272881	0.616576032	Cancer DR
160	7984562	0.202226921	0.545156929	Cancer DR
161	7973056	0.12273026	0.466766159	Cancer DR
162	7968872	0.152629271	0.559511625	Cancer DR
163	7954997	0.227293208	0.835336974	Cancer DR
164	7966534	0.204537211	0.514690813	Cancer DR
165	7912956	0.43944482	0.955570558	Cancer DR
166	8154727	0.189397438	0.644011951	Cancer DR
167	8172154	0.30567409	0.601073059	Cancer DR
168	8027778	0.294634415	0.625261076	Cancer DR
169	8118594	0.169194373	1.322799632	Cancer DR
170	7937476	0.091022022	0.433361614	Cancer DR
171	8043197	0.209947087	0.835958812	Cancer DR
172	8178220	0.167950937	1.257294898	Cancer DR
173	7986765	0.149388016	0.51818569	Cancer DR
174	8050215	0.276176146	0.545194096	Cancer DR
175	8099887	0.18299167	0.435941982	Cancer DR
176	7965515	0.277487295	0.635706106	Cancer DR
177	8036777	0.103814653	0.335518446	Cancer DR
178	7990898	0.181427017	0.435911815	Cancer DR
179	8022170	0.204421552	0.512559266	Cancer DR
180	8171834	0.186887426	0.444312549	Cancer DR
181	7990949	0.178419807	0.429876071	Cancer DR
182	8115147	0.588711197	2.543951919	Cancer DR
183	8125750	0.170693154	0.445978261	Cancer DR
184	7999520	0.094807209	0.434810338	Cancer DR
185	8047635	0.187992412	0.476720865	Cancer DR

C-Neutro	Transcript Cluster ID	Cancer mean	Control mean	Pattern
186	8179519	0.148520936	1.377666159	Cancer DR
187	8028916	0.203678078	0.582119546	Cancer DR
188	7899957	0.32647612	0.692408847	Cancer DR
189	7912385	0.222244737	0.515414934	Cancer DR
190	7929593	0.254483825	0.673379639	Cancer DR
191	7999827	0.22055311	0.458609083	Cancer DR
192	7942586	0.122846179	0.381746923	Cancer DR
193	7903010	0.204557858	0.476036067	Cancer DR
194	8180402	0.365236868	0.664121825	Cancer DR
195	8154359	0.270774973	0.649779593	Cancer DR
196	8160308	0.139708547	0.388974892	Cancer DR
197	7901038	0.17942811	0.505383487	Cancer DR
198	8026106	0.273759786	0.625401988	Cancer DR
199	8152764	0.290598102	1.494807117	Cancer DR
200	8084488	0.12996986	0.377518065	Cancer DR

Table 5. Prostate Cancer vs. Head and Neck Cancer (PC/HNC)-Macro Markers 1-

100

(DR=down-regulated; UR= up-regulated)

PC/HNC-Macro	Transcript Cluster ID	Prostate mean	HNC mean	Pattern
1	7900888	0.17371456	0.08817516	Prostate UR
2	7900639	1.802152255	1.15397654	Prostate UR
3	7898739	1.11408961	0.82749033	Prostate UR
4	7898413	0.67392556	0.49306004	Prostate UR
5	7897034	0.626201623	0.44690779	Prostate UR
6	7896961	0.630327274	0.40732072	Prostate UR
7	7899343	0.726076216	0.59157349	Prostate UR
8	7897960	0.356708994	0.25225113	Prostate UR
9	7897305	0.085317423	0.05232444	Prostate UR
10	7897119	0.0622508	0.04125291	Prostate UR
11	7897745	1.105341947	0.85257586	Prostate UR
12	7901048	0.893414334	0.55699338	Prostate UR
13	7900999	0.893414334	0.55699338	Prostate UR
14	7899562	0.920866373	0.7809056	Prostate UR
15	7899392	8.628724439	5.51200272	Prostate UR
16	7898998	0.839792645	0.66603122	Prostate UR
17	7898805	0.952125401	0.78215228	Prostate UR
18	7900438	2.940352474	3.89294087	Prostate DR
19	7899851	0.842252138	1.10398234	Prostate DR
20	7898736	1.346375231	1.89769295	Prostate DR
21	7898483	2.753599998	5.56947389	Prostate DR
22	7896937	0.977555399	1.2014229	Prostate DR
23	7896929	2.811611487	4.34483418	Prostate DR
24	7899323	0.932226468	0.74082024	Prostate UR
25	7900492	1.454019348	1.1539056	Prostate UR

PC/HNC-Macro	Transcript Cluster ID	Prostate mean	HNC mean	Pattern
26	7898881	1.466931081	1.17758359	Prostate UR
27	7900167	0.248306402	0.15711678	Prostate UR
28	7898020	1.664652638	1.35936858	Prostate UR
29	7900878	0.488076814	0.93299234	Prostate DR
30	7900609	9.440920803	6.580795	Prostate UR
31	7900409	2.146228006	1.60768619	Prostate UR
32	7899096	0.186776639	0.11669783	Prostate UR
33	7899534	1.152420579	1.01684571	Prostate UR
34	7900488	0.816735668	0.65919367	Prostate UR
35	7899407	0.822570685	0.69175622	Prostate UR
36	7899018	9.34362152	14.876575	Prostate UR
37	7899455	4.481504457	3.05597153	Prostate UR
38	7900228	0.565473717	0.44475451	Prostate UR
39	7897449	1.028962001	0.83599809	Prostate UR
40	7900654	1.18320818	0.9112754	Prostate UR
41	7897236	0.604752408	0.38075156	Prostate UR
42	7897849	1.308607915	1.02712991	Prostate UR
43	7898655	0.765326514	0.57697822	Prostate UR
44	7896865	0.981029132	0.85935166	Prostate UR
45	7897426	0.058940751	0.04164638	Prostate UR
46	7898431	1.612544635	1.14610124	Prostate UR
47	7899350	1.470736123	1.14874973	Prostate UR
48	7899448	1.138821395	0.9662266	Prostate UR
49	7899750	0.329695991	0.47917855	Prostate DR
50	7897691	1.377648768	1.10001944	Prostate UR
51	7900382	8.047259032	5.40447368	Prostate UR
52	7900192	1.59154501	1.77652546	Prostate DR
53	7898350	1.166464627	0.87088799	Prostate UR
54	7898296	1.166464627	0.87088799	Prostate UR

PC/HNC-Macro	Transcript Cluster ID	Prostate mean	HNC mean	Pattern
55	7896756	16.53743972	24.4383885	Prostate DR
56	7898124	1.314794095	0.88645691	Prostate UR
57	7898988	2.452450743	3.63039674	Prostate DR
58	7898337	0.632765821	0.45503236	Prostate UR
59	7898161	0.946204421	1.05104361	Prostate DR
60	7899829	2.145803014	1.73641438	Prostate UR
61	7900585	1.284762498	1.68441785	Prostate DR
62	7898594	2.122455762	1.65893241	Prostate UR
63	7899774	2.402309625	2.05018985	Prostate UR
64	7897257	1.013808953	0.75484435	Prostate UR
65	7897441	2.335757427	3.27479786	Prostate DR
66	7897522	2.270470627	1.91499556	Prostate UR
67	7899932	0.737050553	1.14729271	Prostate DR
68	7899898	0.544420785	0.65409018	Prostate DR
69	7900001	1.760395525	1.45748264	Prostate UR
70	7899921	2.464792372	3.98131153	Prostate DR
71	7897987	0.672103038	0.53917805	Prostate UR
72	7897890	1.720201525	1.30711874	Prostate UR
73	7900635	3.545016951	2.67473725	Prostate UR
74	7898833	1.779164948	1.37302263	Prostate UR
75	7897329	2.254239663	1.81517566	Prostate UR
76	7899790	5.863884634	4.13618926	Prostate UR
77	7900490	1.36827874	1.22792955	Prostate UR
78	7898916	0.987033618	0.8820278	Prostate UR
79	7897416	0.819651746	0.39507013	Prostate UR
80	7898192	0.971522039	0.84215204	Prostate UR
81	7898521	0.755178665	0.64866356	Prostate UR
82	7898256	1.152186796	0.96705766	Prostate UR
83	7898677	1.139329856	0.96095486	Prostate UR

PC/HNC-Macro	Transcript Cluster ID	Prostate mean	HNC mean	Pattern
84	7899016	0.594556425	0.49087673	Prostate UR
85	7898609	1.72946927	1.31260331	Prostate UR
86	7898516	1.435723474	1.20399467	Prostate UR
87	7898052	1.624343059	1.0990904	Prostate UR
88	7898750	1.002634449	1.1486647	Prostate DR
89	7898002	1.352879559	1.11186666	Prostate UR
90	7899882	0.913865556	0.80538181	Prostate UR
91	7897288	1.064532962	0.95246928	Prostate UR
92	7898263	1.19058275	1.09256179	Prostate UR
93	7901046	0.916598487	0.84402334	Prostate UR
94	7896754	0.866090536	0.79633452	Prostate UR
95	7898616	1.147030435	1.01524538	Prostate UR
96	7900076	0.805161154	0.69042447	Prostate UR
97	7898357	1.220596829	1.10802848	Prostate UR
98	7899688	1.611962379	2.33223295	Prostate DR
99	7898184	2.190196822	1.87918773	Prostate UR
100	7899727	2.187854482	2.01458398	Prostate UR

Table 6 Prostate Cancer vs. Head and Neck Cancer (PC/HNC)-Neutro Markers 1-

100

(DR=down-regulated; UR= up-regulated)

PC/HNC-Neutro	Transcript Cluster ID	Prostate Mean	HNC Mean	Pattern
1	7898314	0.472352149	0.295758103	Prostate UR
2	7898263	0.470346508	0.310072475	Prostate UR
3	7897277	0.403341785	0.246610932	Prostate UR
4	7897172	0.528467409	0.37306189	Prostate UR
5	7897044	0.146804612	0.092032212	Prostate UR
6	7896917	0.564875205	0.387977245	Prostate UR
7	7897955	0.188254956	0.113855395	Prostate UR
8	7897966	0.232515724	0.152024283	Prostate UR
9	7897179	0.578722421	0.340430311	Prostate UR
10	7898537	0.279292833	0.174491575	Prostate UR
11	7898184	0.226400032	0.14950648	Prostate UR
12	7896761	0.274708621	0.187070078	Prostate UR
13	7897714	0.520153631	0.363473911	Prostate UR
14	7899016	0.616979782	0.431180962	Prostate UR
15	7899343	0.777877153	0.343891728	Prostate UR
16	7899273	0.777877153	0.343891728	Prostate UR
17	7899029	0.179455931	0.111363025	Prostate UR
18	7897068	0.412604522	0.221994054	Prostate UR
19	7897849	0.246163341	0.171656261	Prostate UR
20	7897663	0.246163341	0.171656261	Prostate UR
21	7897728	0.160936511	0.122730523	Prostate UR
22	7898902	0.109896347	0.072605657	Prostate UR
23	7898124	0.758065325	0.433521341	Prostate UR
24	7897469	0.230680674	0.152019315	Prostate UR
25	7897416	0.230680674	0.152019315	Prostate UR
26	7897449	0.213629094	0.157994707	Prostate UR

PC/HNC-Neutro	Transcript Cluster ID	Prostate Mean	HNC Mean	Pattern
27	7897370	0.135066751	0.079756496	Prostate UR
28	7898750	0.654243362	0.327821118	Prostate UR
29	7897026	0.090788163	0.062728225	Prostate UR
30	7897691	0.194316525	0.117272696	Prostate UR
31	7899005	0.200247191	0.147403774	Prostate UR
32	7898413	0.472679058	0.322722967	Prostate UR
33	7899346	0.640398768	0.323867466	Prostate UR
34	7899284	0.640398768	0.323867466	Prostate UR
35	7899361	0.337035736	0.221863494	Prostate UR
36	7898431	0.156923062	0.09015974	Prostate UR
37	7896921	0.174899012	0.119106922	Prostate UR
38	7898448	0.135061146	0.109332089	Prostate UR
39	7897378	0.267292165	0.133889964	Prostate UR
40	7898516	0.098348954	0.062481326	Prostate UR
41	7898809	0.491677002	0.367987619	Prostate UR
42	7898736	0.113999161	0.082835856	Prostate UR
43	7897745	0.104572957	0.060987418	Prostate UR
44	7898052	1.445062229	0.629181447	Prostate UR
45	7897987	0.498426291	0.397985849	Prostate UR
46	7899023	0.214386673	0.148057449	Prostate UR
47	7898084	0.14555045	0.109512027	Prostate UR
48	7896740	0.618945016	0.477094838	Prostate UR
49	7898007	1.033298467	0.60491554	Prostate UR
50	7899087	0.489854074	0.403354804	Prostate UR
51	7898967	0.318853504	0.191403501	Prostate UR
52	7897132	0.110714756	0.059941241	Prostate UR
53	7896817	0.894991289	0.694908734	Prostate UR
54	7896985	0.516399994	0.384878562	Prostate UR
55	7898115	0.109725861	0.080915573	Prostate UR

PC/HNC-Neutro	Transcript Cluster ID	Prostate Mean	HNC Mean	Pattern
56	7898916	0.164417248	0.127708697	Prostate UR
57	7898602	0.067768264	0.038000029	Prostate UR
58	7899075	0.091420802	0.069853218	Prostate UR
59	7897974	0.664695996	0.509259275	Prostate UR
60	7899134	0.103951442	0.077074358	Prostate UR
61	7898910	0.545102973	0.397482411	Prostate UR
62	7898102	0.175886334	0.135446742	Prostate UR
63	7896861	0.230031409	0.190340424	Prostate UR
64	7898012	0.223460808	0.168124325	Prostate UR
65	7896822	0.372917775	0.21278994	Prostate UR
66	7899310	0.321550728	0.10832907	Prostate UR
67	7899192	0.321550728	0.10832907	Prostate UR
68	7898957	0.973685715	0.587477731	Prostate UR
69	7897824	0.223158535	0.17320102	Prostate UR
70	7898112	0.161852727	0.127616873	Prostate UR
71	7898655	0.253453749	0.177336112	Prostate UR
72	7898833	0.393403666	0.283521593	Prostate UR
73	7896878	0.506063167	0.342056413	Prostate UR
74	7896882	0.157090574	0.065476841	Prostate UR
75	7896961	0.182494175	0.100199558	Prostate UR
76	7897322	0.141715372	0.103369275	Prostate UR
77	7897441	0.186301612	0.146498074	Prostate UR
78	7897803	0.161327026	0.121522293	Prostate UR
79	7896754	0.173109995	0.130063698	Prostate UR
80	7898375	0.167664303	0.09775886	Prostate UR
81	7899220	0.195359832	0.085063774	Prostate UR
82	7899377	0.162661602	0.125660578	Prostate UR
83	7897685	0.156285092	0.117822	Prostate UR
84	7897426	0.18561995	0.155860307	Prostate UR

PC/HNC- Neutro	Transcript Cluster ID	Prostate Mean	HNC Mean	Pattern
85	7897034	0.297786106	0.198693521	Prostate UR
86	7898411	0.162384316	0.127798349	Prostate UR
87	7899167	0.181908563	0.123429432	Prostate UR
88	7899096	0.181908563	0.123429432	Prostate UR
89	7898653	0.261680692	0.206759226	Prostate UR
90	7898328	0.55280053	0.466282461	Prostate UR
91	7897295	0.289839522	0.228101347	Prostate UR
92	7898256	0.170890017	0.145553307	Prostate UR
93	7898799	0.218542702	0.188146131	Prostate UR
94	7897890	0.41916937	0.332096907	Prostate UR
95	7898483	0.161978892	0.133431594	Prostate UR
96	7897404	0.551161709	0.432147632	Prostate UR
97	7898988	0.283480243	0.220289746	Prostate UR
98	7896937	0.243352142	0.184139991	Prostate UR
99	7898725	0.190386924	0.099484518	Prostate UR
100	7898793	0.180311	0.133379341	Prostate UR

Table 7. Prostate Cancer vs. Head and Neck Cancer (PC/HNC)-Macro Markers
101-200

(DR = down-regulated; UR= up-regulated)

PC/HNC-Macro	Transcript Cluster ID	Gene Name	Prostate mean	HNC mean	Pattern
101	8168524	P2RY10	0.152005253	0.088175164	Prostate UR
102	8121257	PRDM1	0.25519801	0.157116777	Prostate UR
103	8026787	FAM129C	1.118594845	0.827490331	Prostate UR
104	8009685	SLC9A3R1	0.659954127	0.493060042	Prostate UR
105	7952126	RPL23AP64	1.121821816	0.852575862	Prostate UR
106	7920575	PBXIP1	0.61736145	0.446907787	Prostate UR
107	8031939	ZNF584	0.965208892	0.782152279	Prostate UR
108	8074695	MGC16703	0.839858433	0.691756222	Prostate UR
109	8087596	HYAL3	0.910617775	0.780905597	Prostate UR
110	8069332	MCM3AP-AS	0.706777639	0.591573489	Prostate UR
111	8070720	ICOSLG	1.531107114	1.148749732	Prostate UR
112	7923917	FAIM3	0.057531404	0.04125291	Prostate UR
113	7935660	DNMBP	2.425400218	1.815175659	Prostate UR
114	8144279	CLN8	1.43223828	1.153905598	Prostate UR
115	7928308	DDIT4	0.616394267	0.380751555	Prostate UR
116	8168028	STARD8	1.895181754	2.448347527	Prostate DR
117	8117377	HIST1H1E	0.581468103	0.696626006	Prostate DR
118	8101992	SLC39A8	0.299403929	0.479178554	Prostate DR
119	8026579	TMEM38A	1.378724037	1.897692954	Prostate DR
120	8012028	ASGR2	3.049693033	5.569473894	Prostate DR
121	7914184	PTAFR	3.028813569	4.344834176	Prostate DR
122	8132399	C7orf36	0.392349882	0.483849886	Prostate DR
123	8044766	INSIG2	0.75337043	0.86796442	Prostate DR
124	8121277	AIM1	0.827430258	1.014548905	Prostate DR
125	8137240	GIMAP7	0.402671799	0.505696085	Prostate DR

PC/HNC-Macro	Transcript Cluster ID	Gene Name	Prostate mean	HNC mean	Pattern
126	8137257	GIMAP5	0.210901215	0.280198715	Prostate DR
127	8112409	SGTB	0.782231488	0.950030056	Prostate DR
128	8112803	LHFPL2	2.616178494	3.981311526	Prostate DR
129	7898655	CDA	17.75853426	24.4383885	Prostate DR
130	8047419	CASP8	0.821464161	0.666031224	Prostate UR
131	8161701	TMEM2	0.931589162	1.197717741	Prostate DR
132	8046124	DHRS9	2.785100395	3.630396735	Prostate DR
133	8146533	FAM110B	1.309830556	1.684417848	Prostate DR
134	8112220	PDE4D	0.545705486	0.65409018	Prostate DR
135	7961371	DUSP16	0.330335584	0.252251134	Prostate UR
136	8072744	NCF4	8.116005803	5.512002723	Prostate UR
137	7942832	C11orf8	2.513452632	3.274797863	Prostate DR
138	7906400	IFI16	0.901058293	1.008070795	Prostate DR
139	7904465	HIST2H2BA	0.972877361	0.859351662	Prostate UR
140	7955606	C12orf4	1.312181584	1.02712991	Prostate UR
141	7944656	SC5DL	0.599262019	0.772958547	Prostate DR
142	8051396	NLRC4	10.46151626	14.87657498	Prostate DR
143	7917304	MCOLN3	0.555819644	0.407320715	Prostate UR
144	7949060	PPP1R14B	1.407220289	1.100019444	Prostate UR
145	8109086	ADRB2	0.905393742	1.103982342	Prostate DR
146	8113356	GPAM	0.723135275	1.147292709	Prostate DR
147	8114511	MGC29506	0.999375725	0.906642744	Prostate UR
148	7944152	IL10RA	1.001641777	0.835998092	Prostate UR
149	8027247	ZNF93	0.982313823	1.148664699	Prostate DR
150	8105778	PIK3R1	0.421185322	0.472940975	Prostate DR
151	7961142	OLR1	1.156078519	1.372673235	Prostate DR
152	8095728	EREG	1.674886156	2.332232946	Prostate DR
153	8134454	BRI3	3.27849217	3.892940871	Prostate DR
154	8144267	CLN8	1.389963945	1.227929549	Prostate UR

PC/HNC-Macro	Transcript Cluster ID	Gene Name	Prostate mean	HNC mean	Pattern
155	7982066	SNORD115-30	0.955490399	1.051043605	Prostate DR
156	8039766	ZNF837	1.001545409	0.882027802	Prostate UR
157	7896882	SCNN1D	1.012465552	0.933891434	Prostate UR
158	8121927	RNF146	1.652162902	1.776525457	Prostate DR
159	8078014	SLC6	4.293246551	3.055971532	Prostate UR
160	7962516	SLC38A1	0.04103982	0.052711457	Prostate DR
161	8017850	WIP1	2.135605918	1.658932412	Prostate UR
162	8003298	SLC7A5	0.630994213	0.455032363	Prostate UR
163	8029136	CD79A	1.212433912	0.638405917	Prostate UR
164	8105328	SNX18	2.134429359	1.736414377	Prostate UR
165	8117415	HIST1H3E	1.221646142	1.053585059	Prostate UR
166	8087374	C3orf62	2.239568296	1.740586151	Prostate UR
167	8124416	HIST1H3D	0.536020333	0.44475451	Prostate UR
168	8165672	RFC1	0.540896622	0.932992344	Prostate DR
169	8146934	LY96	8.750934207	6.580795003	Prostate UR
170	8143759	ATP6V0E2	0.779721234	0.659193671	Prostate UR
171	8076331	TOB2	1.122353474	0.966226601	Prostate UR
172	8153002	NDRG1	1.129775721	0.9112754	Prostate UR
173	7963280	SMAGP	0.680810703	0.539178053	Prostate UR
174	8009693	TMEM104	1.537379474	1.146101244	Prostate UR
175	8059854	ARL4C	0.165130728	0.116697835	Prostate UR
176	7940287	MS4A1	0.865271896	0.395070134	Prostate UR
177	8130768	RNASET2	2.001804778	1.607686191	Prostate UR
178	7915500	C1orf21	1.036426947	1.201422899	Prostate DR
179	8094226	FAM200B	1.426108596	1.724534154	Prostate DR
180	8130556	SOD2	7.561106712	5.404473683	Prostate UR
181	8179263	TNF	2.369373804	1.396352887	Prostate UR
182	8177983	TNF	2.369373804	1.396352887	Prostate UR

PC/HNC-Macro	Transcript Cluster ID	Gene Name	Prostate mean	HNC mean	Pattern
183	8118142	TNF	2.369373804	1.396352887	Prostate UR
184	8116484	MGAT1	1.723379292	1.457482638	Prostate UR
185	7961524	ERP27	0.510625121	0.604222101	Prostate DR
186	7989708	MTFMT	0.957190171	0.842152039	Prostate UR
187	8012896	PMP22	1.417750917	1.203994674	Prostate UR
188	8150225	RAB11FIP1	3.478582769	2.674737249	Prostate UR
189	7936041	ARL3	1.221836858	1.10635807	Prostate UR
190	8143564	FAM131B	0.99352351	0.901864091	Prostate UR
191	7926807	PDSS1	1.57365811	1.810161132	Prostate DR
192	7929032	FAS	0.97455032	0.754844347	Prostate UR
193	8076185	CBX7	0.630905883	0.470647702	Prostate UR
194	8038815	LIM2	1.007183131	0.902069895	Prostate UR
195	8058905	CXCR1	9.26228244	5.220575301	Prostate UR
196	7949754	CLCF1	0.932350828	0.823640044	Prostate UR
197	7998931	ZNF200	1.133830607	0.967057656	Prostate UR
198	7973352	LRP10	1.565492996	1.099090397	Prostate UR
199	7945944	RHOG	2.197395251	1.914995562	Prostate UR
200	8136388	TMEM140	1.551595593	1.19126752	Prostate UR

Table 8 Prostate Cancer vs. Head and Neck Cancer (PC/HNC)-Neutro Markers

101-200

(DR = down-regulated; UR= up-regulated)

PC/HNC-Neutro	Transcript Cluster ID	Gene Name	Prostate mean	HNC mean	Pattern
101	8066964	TMEM189	0.474044	0.295758	Prostate UR
102	8059222	DNPEP	0.463833	0.310072	Prostate UR
103	8043197	VAMP8	0.255407	0.149506	Prostate UR
104	7956220	OBFC2B	0.400177	0.246611	Prostate UR
105	7949206	MEN1	0.518189	0.373062	Prostate UR
106	7940372	TMEM109	0.157114	0.092032	Prostate UR
107	8008132	ATP5G1	0.195415	0.113855	Prostate UR
108	8092457	ALG3	0.295813	0.174492	Prostate UR
109	8012257	TP53	0.230245	0.152024	Prostate UR
110	7901601	MRPL37	0.282849	0.18707	Prostate UR
111	7920725	SCAMP3	0.523564	0.387977	Prostate UR
112	7949948	C11orf2	0.560081	0.34043	Prostate UR
113	7941936	GSTP1	0.437974	0.221994	Prostate UR
114	8154727	LOC138412	0.219733	0.147404	Prostate UR
115	8143054	AKR1B1	0.120667	0.072606	Prostate UR
116	8000507	EIF3C	0.258987	0.171656	Prostate UR
117	7994415	EIF3C	0.258987	0.171656	Prostate UR
118	8180022	HLA-DQB1	0.789467	0.343892	Prostate UR
119	8178826	HLA-DQB1	0.789467	0.343892	Prostate UR
120	8158714	EXOSC2	0.174407	0.111363	Prostate UR
121	8125537	HLA-DMA	0.724437	0.327821	Prostate UR
122	7937802	CD81	0.09944	0.062728	Prostate UR
123	8155327	ALDH1B1	0.581084	0.431181	Prostate UR
124	8180086	HLA-DMA	0.712986	0.323867	Prostate UR
125	8178884	HLA-DMA	0.712986	0.323867	Prostate UR
126	8037835	SLC1A5	0.764073	0.433521	Prostate UR

PC/HNC-Neutro	Transcript Cluster ID	Gene Name	Prostate mean	HNC mean	Pattern
127	8036602	ECH1	0.196343	0.135447	Prostate UR
128	7997099	SF3B3	0.161405	0.122731	Prostate UR
129	7996677	NUTF2	0.482693	0.363474	Prostate UR
130	7973056	APEX1	0.139701	0.079756	Prostate UR
131	8109750	RPLP0	0.083731	0.038	Prostate UR
132	7984562	RPLP1	0.213785	0.157995	Prostate UR
133	8036737	RPS16	0.176177	0.127617	Prostate UR
134	8085431	NUP210	0.106052	0.062481	Prostate UR
135	8164100	RPL35	0.097157	0.069853	Prostate UR
136	7948667	AHNAK	0.127933	0.059941	Prostate UR
137	8085026	RPL35A	0.182548	0.133432	Prostate UR
138	7950307	UCP2	0.237958	0.159583	Prostate UR
139	7986685	DEXI	0.223207	0.152019	Prostate UR
140	7982185	DEXI	0.223207	0.152019	Prostate UR
141	8180260	Unknown	0.345186	0.221863	Prostate UR
142	7994603	// SP	0.192445	0.117273	Prostate UR
143	7921637	CD84	0.177232	0.119107	Prostate UR
144	7999520	RSL1D1	0.109754	0.060987	Prostate UR
145	8119993	HSP90AB1	0.115186	0.082836	Prostate UR
146	8152764	MTSS1	0.348016	0.191404	Prostate UR
147	8079334	LIMD1	0.450301	0.322723	Prostate UR
148	8029136	CD79A	1.461863	0.629181	Prostate UR
149	8036777	FBL	0.114442	0.080916	Prostate UR
150	8132523	UBE2D4	0.488117	0.367988	Prostate UR
151	8115158	RPS14	0.234508	0.172846	Prostate UR
152	8173513	RPS4X	0.109783	0.077074	Prostate UR
153	7899134	CCDC21	0.622472	0.477095	Prostate UR
154	8150872	RPS20	0.165436	0.127709	Prostate UR
155	8051066	MPV17	0.23829	0.180993	Prostate UR

PC/HNC-Neutro	Transcript Cluster ID	Gene Name	Prostate mean	HNC mean	Pattern
156	8164373	CIZ1	0.477605	0.403355	Prostate UR
157	7906486	SLAMF8	0.901391	0.694909	Prostate UR
158	7936100	CALHM2	0.524725	0.384879	Prostate UR
159	7973067	PNP	0.252503	0.13389	Prostate UR
160	8081277	NIT2	0.146986	0.09016	Prostate UR
161	8177003	SLC25A6	0.204081	0.123429	Prostate UR
162	8171111	SLC25A6	0.204081	0.123429	Prostate UR
163	8036525	MAP4K1	0.137383	0.109512	Prostate UR
164	8024299	RPS15	0.230076	0.168124	Prostate UR
165	7920123	S100A1	0.188373	0.065477	Prostate UR
166	8014454	MYO19	0.472992	0.397986	Prostate UR
167	8118594	HLA-DPB1	0.237858	0.099485	Prostate UR
168	7937476	RPLP2	0.119264	0.074065	Prostate UR
169	8012464	LOC100128288	0.665447	0.509259	Prostate UR
170	8146649	MTFR1	0.548642	0.397482	Prostate UR
171	8178220	HLA-DPB1	0.232773	0.085064	Prostate UR
172	7906564	PEA15	0.39459	0.21279	Prostate UR
173	8152215	KLF10	1.026638	0.587478	Prostate UR
174	8000284	GGA2	0.230644	0.173201	Prostate UR
175	8158446	NUP188	0.201645	0.148057	Prostate UR
176	7912385	EXOSC10	0.232965	0.19034	Prostate UR
177	7984364	SMAD3	0.191893	0.146498	Prostate UR
178	8079019	RPS27	0.171457	0.127798	Prostate UR
179	8133721	HSPB1	0.403533	0.283522	Prostate UR
180	7983843	TCF12	0.194109	0.15586	Prostate UR
181	8180100	HLA-DPA1	0.310941	0.100245	Prostate UR
182	7952129	RPS25	0.181072	0.1281	Prostate UR
183	8179489	HLA-DQA1	0.33676	0.108329	Prostate UR
184	8178199	HLA-DQA1	0.33676	0.108329	Prostate UR

PC/HNC-Neutro	Transcript Cluster ID	Gene Name	Prostate mean	HNC mean	Pattern
185	7918593	RHOC	0.522543	0.342056	Prostate UR
186	8023415	TCF4	0.98316	0.604916	Prostate UR
187	8086148	Unknown	0.185502	0.142253	Prostate UR
188	8180287	Unknown	0.1684	0.125661	Prostate UR
189	8154394	SNAPC3	0.220226	0.15656	Prostate UR
190	7906662	UFC1	0.226711	0.184035	Prostate UR
191	7994565	RRN3	0.161065	0.117822	Prostate UR
192	8076209	RPL3	0.099145	0.074704	Prostate UR
193	8007302	TUBG1	0.432219	0.332097	Prostate UR
194	7965515	NDUFA12	0.299008	0.228101	Prostate UR
195	7900157	RPS27	0.176397	0.130064	Prostate UR
196	8000200	RRN3P1	0.162659	0.121522	Prostate UR
197	8178891	HLA-DPA1	0.316307	0.112019	Prostate UR
198	8125556	HLA-DPA1	0.316307	0.112019	Prostate UR
199	8114567	PFDN1	0.247425	0.177336	Prostate UR
200	8170859	RPL10	0.209331	0.145986	Prostate UR

What is Claimed is:

1. A method for diagnosing or aiding in the diagnosis of a cancer in a subject, the method comprising the steps of:
 - a) measuring the levels of one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells;
 - b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and
 - c) identifying a difference between the measured levels of the one or more selected C-Macro markers in steps a) and b),wherein the identified difference indicates that the subject has said cancer.
2. A method for assessing the risk of developing a cancer in a subject, the method comprising the steps of:
 - a) measuring the levels of one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells;
 - b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and
 - c) identifying a difference between the measured levels of the one or more selected C-Macro markers in steps a) and b),wherein the identified difference indicates that the subject has a risk of developing said cancer.
3. A method for prognosing or aiding in the prognosis of a cancer in a subject, the method comprising the steps of:
 - a) measuring the levels of one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells;
 - b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and
 - c) identifying a difference between the measured levels of the one or more selected C-Macro markers in steps a) and b),wherein the identified difference is indicative of the prognosis of said cancer in the subject.

4. A method for assessing the efficacy of a treatment for a cancer in a subject comprising:

- a) measuring the levels of one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells before the treatment;
- b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells before the treatment;
- c) identifying a first difference between the measured levels of the one or more selected C-Macro markers in steps a) and b);
- d) measuring the levels of the one or more selected C-Macro markers in a population of the subject's macrophage cells after the treatment;
- e) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells after the treatment;
- f) identifying a second difference between the measured levels of the one or more selected C-Macro markers in steps d) and e); and
- g) identifying a difference between the first difference and the second difference,

wherein the difference identified in g) is indicative of the efficacy of the treatment for said cancer in the subject.

5. A method for monitoring the progression or regression of a cancer in a subject comprising:

- a) measuring the levels of one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells at a first time point;
- b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells at the first time point;
- c) identifying a first difference between the measured levels of the one or more selected C-Macro markers in steps a) and b);
- d) measuring the levels of the one or more selected C-Macro markers in a population of the subject's macrophage cells at a second time point;

e) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells at the second time point;

f) identifying a second difference between the measured levels of the one or more selected C-Macro markers in steps d) and e); and

g) identifying a difference between the first difference and the second difference,

wherein the difference identified in g) is indicative of the progression or regression of said cancer in the subject.

6. A method for identifying a compound capable of ameliorating or treating a cancer in a subject comprising:

a) measuring the levels of one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells before administering the compound to the subject;

b) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells before administering the compound to the subject;

c) identifying a first difference between the measured levels of the one or more selected C-Macro markers in steps a) and b);

d) measuring the levels of the one or more selected C-Macro markers in a population of the subject's macrophage cells after the administration of the compound;

e) measuring the levels of the one or more selected C-Macro markers in a population of the subject's non-phagocytic cells after the administration of the compound;

f) identifying a second difference between the measured levels of the one or more selected C-Macro markers in steps d) and e); and

g) identifying a difference between the first difference and the second difference,

wherein the difference identified in g) indicates that the compound is capable of ameliorating or treating said cancer in the subject.

7. A method for diagnosing or aiding in the diagnosis of a cancer in a subject, the method comprising the steps of:
- a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells;
 - b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells; and
 - c) identifying a difference between the measured levels of the one or more selected C-Neutro markers in steps a) and b),
- wherein the identified difference indicates that the subject has said cancer.
8. A method for assessing the risk of developing a cancer in a subject, the method comprising the steps of:
- a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells;
 - b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells; and
 - c) identifying a difference between the measured levels of the one or more selected C-Neutro markers in steps a) and b),
- wherein the identified difference indicates that the subject has a risk of developing said cancer.
9. A method for prognosing or aiding in the prognosis of a cancer in a subject, the method comprising the steps of:
- a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells;
 - b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells; and
 - c) identifying a difference between the measured levels of the one or more selected C-Neutro markers in steps a) and b),
- wherein the identified difference is indicative of the prognosis of said cancer in the subject.

10. A method for assessing the efficacy of a treatment for a cancer in a subject comprising:

- a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells before the treatment;
 - b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells before the treatment;
 - c) identifying a first difference between the measured levels of the one or more selected C-Neutro markers in steps a) and b);
 - d) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's neutrophil cells after the treatment;
 - e) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells after the treatment;
 - f) identifying a second difference between the measured levels of the one or more selected C-Neutro markers in steps d) and e); and
 - g) identifying a difference between the first difference and the second difference,
- wherein the difference identified in g) is indicative of the efficacy of the treatment for said cancer in the subject.

11. A method for monitoring the progression or regression of a cancer in a subject comprising:

- a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells at a first time point;
- b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells at the first time point;
- c) identifying a first difference between the measured levels of the one or more selected C-Neutro markers in steps a) and b);
- d) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's neutrophil cells at a second time point;

- e) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells at the second time point;
 - f) identifying a second difference between the measured levels of the one or more selected C-Neutro markers in steps d) and e); and
 - g) identifying a difference between the first difference and the second difference,
- wherein the difference identified in g) is indicative of the progression or regression of said cancer in the subject.

12. A method for identifying a compound capable of ameliorating or treating a cancer in a subject comprising:

- a) measuring the levels of one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells before administering the compound to the subject;
 - b) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells before administering the compound to the subject;
 - c) identifying a first difference between the measured levels of the one or more selected C-Neutro markers in steps a) and b);
 - d) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's neutrophil cells after the administration of the compound;
 - e) measuring the levels of the one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells after the administration of the compound;
 - f) identifying a second difference between the measured levels of the one or more selected C-Neutro markers in steps d) and e); and
 - g) identifying a difference between the first difference and the second difference,
- wherein the difference identified in g) indicates that the compound is capable of ameliorating or treating said cancer in the subject.

13. A method for diagnosing or aiding in the diagnosis of a cancer in a subject, the method comprising the steps of:

a) measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells, and

measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells;

b) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and

measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells;

c) identifying a difference between the measured levels of the at least one or more selected C-Macro markers in steps a) and b); and

d) identifying a difference between the measured levels or activities the at least one or more selected C-Neutro markers in steps a) and b);

wherein the differences identified in c) and d) indicate that the subject has said cancer.

14. A method for assessing the risk of developing a cancer in a subject, the method comprising the steps of:

a) measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells, and

measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells;

b) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and

measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells;

c) identifying a difference between the measured levels of the at least one or more selected C-Macro markers in steps a) and b); and

d) identifying a difference between the measured levels of the at least one or more selected C-Neutro markers in steps a) and b);

wherein the differences identified in c) and d) indicate that the subject has a risk of developing said cancer.

15. A method for prognosing or aiding in the prognosis of a cancer in a subject, the method comprising the steps of:

a) measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells, and

measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells;

b) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells; and

measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells;

c) identifying a difference between the measured levels of the at least one or more selected C-Macro markers in steps a) and b); and

d) identifying a difference between the measured levels of the at least one or more selected C-Neutro markers in steps a) and b);

wherein the differences identified in c) and d) are indicative of the prognosis of said cancer in the subject.

16. A method for assessing the efficacy of a treatment for a cancer in a subject comprising:

a) measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells before the treatment, and

measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells before the treatment;

b) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells before the treatment; and

measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells before the treatment;

c) identifying a first difference between the measured levels of the at least one or more selected C-Macro markers in steps a) and b); and

identifying a second difference between the measured levels of the at least one or more selected C-Neutro markers in steps a) and b);

d) measuring the levels of the at least one or more selected C-Macro marker in a population of the subject's macrophage cells after the treatment, and

measuring the levels of the at least one or more selected C-Neutro marker in a population of the subject's neutrophil cells after the treatment;

e) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells after the treatment; and

measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells after the treatment;

f) identifying a third difference between the measured levels of the at least one or more selected C-Macro markers in steps d) and e); and

g) identifying a fourth difference between the measured levels of the at least one or more selected C-Neutro markers in steps d) and e);

h) identifying a difference between the first and second differences; and

i) identifying a difference between the third and fourth differences,

wherein the differences identified in h) and i) are indicative of the efficacy of the treatment for said cancer in the subject.

17. A method for monitoring the progression or regression of a cancer in a subject comprising:

a) measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells at a first time point, and

measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells at the first time point;

b) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells at the first time point; and

measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells at the first time point;

c) identifying a first difference between the measured levels of the at least one or more selected C-Macro markers in steps a) and b); and

identifying a second difference between the measured levels of the at least one or more selected C-Neutro markers in steps a) and b);

d) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's macrophage cells at a second time point, and

measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's neutrophil cells at the second time point;

e) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells at the second time point; and

measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells at the second time point;

f) identifying a third difference between the measured levels of the at least one or more selected C-Macro markers in steps d) and e); and

g) identifying a fourth difference between the measured levels of the at least one or more selected C-Neutro markers in steps d) and e);

h) identifying a difference between the first and second differences; and

i) identifying a difference between the third and fourth differences,

wherein the differences identified in h) and i) are indicative of the progression or regression of said cancer in the subject.

18. A method for identifying a compound capable of ameliorating or treating a cancer in a subject comprising:

a) measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200 in a population of the subject's macrophage cells before administering the compound to the subject, and

measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200 in a population of the subject's neutrophil cells before administering the compound to the subject;

b) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells before administering the compound to the subject; and

measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells before administering the compound to the subject;

c) identifying a first difference between the measured levels of the at least one or more selected C-Macro markers in steps a) and b); and

identifying a second difference between the measured levels of the at least one or more selected C-Neutro markers in steps a) and b);

d) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's macrophage cells after administering the compound to the subject, and

measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's neutrophil cells after administering the compound to the subject;

e) measuring the levels of the at least one or more selected C-Macro markers in a population of the subject's non-phagocytic cells after administering the compound to the subject; and

measuring the levels of the at least one or more selected C-Neutro markers in a population of the subject's non-phagocytic cells after administering the compound to the subject;

f) identifying a third difference between the measured levels of the at least one or more selected C-Macro markers in steps d) and e); and

g) identifying a fourth difference between the measured levels of the at least one or more selected C-Neutro markers in steps d) and e);

h) identifying a difference between the first and second differences; and

i) identifying a difference between the third and fourth differences,

wherein the differences identified in h) and i) indicate that the compound is capable of ameliorating or treating said cancer in the subject.

19. The method of any one of claims 1-18, further comprising measuring at least one standard parameter associated with said cancer.

20. The method of claim 19, wherein the standard parameter is selected from the group consisting of tumor stage, tumor grade, tumor size, tumor visual characteristics, tumor growth, and tumor thickness, tumor progression, tumor metastasis, tumor distribution within the body, odor, molecular pathology, genomics, tumor angiograms, or Gleason score.

21. The method of any one of claims 13-18, wherein the selected C-Macro markers and the selected C-Neutro markers are measured from the same population of non-phagocytic cells in steps b) or e).

22. The method of any one of claims 13-18, wherein the selected C-Macro markers and the selected C-Neutro are from different populations of non-phagocytic cells in steps b) or e).

23. The method of any one of claims 1-6 and 13-18, wherein at least two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen,

fifteen, sixteen, seventeen, eighteen, nineteen, twenty, twenty-one, twenty-two, twenty-three, twenty-four, or twenty-five markers are selected from C-Macro 1-200.

24. The method of any one of claims 1-6 and 13-18, wherein the selected C-Macro markers comprise one or more markers selected from the group consisting of C-Macro 1-4 and C-Macro 101-104.

25. The method of any one of claims 1-6 and 13-18, wherein the selected C-Macro markers are up-regulated or activated in the macrophage cells compared to the non-phagocytic cells.

26. The method of any one of claims 1-6 and 13-18, wherein the selected C-Macro markers are up-regulated or activated in the macrophage cells compared to the non-phagocytic cells.

27. The method of any one of claims 1-6 and 13-18, wherein the selected C-Macro markers are down-regulated or inhibited in the macrophage cells compared to the non-phagocytic cells.

28. The method of any one of claims 1-6 and 13-18, wherein the selected C-Macro markers are down-regulated or inhibited in the macrophage cells compared to the non-phagocytic cells.

29. The method of any one of claims 7-18, wherein at least two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty, twenty-one, twenty-two, twenty-three, twenty-four, or twenty-five markers are selected from C-Neutro 1-200.

30. The method of any one of claims 7-18, wherein the selected C-Neutro markers comprise one or more C-Neutro markers selected from the group consisting of C-Neutro 1-8 and C-Neutro 101-108.

31. The method of any one of claims 7-18, wherein the selected C-Neutro markers comprise one or more markers selected from the group consisting of C-Neutro 1-200 and wherein the selected C-Neutro markers are down-regulated or inhibited in the neutrophil cells compared to the non-phagocytic cells.

32. The method of any one of claims 7-18, wherein the selected C-Neutro markers are down-regulated or inhibited in the neutrophil cells compared to the non-phagocytic cells.

33. The method of any one of claims 1-6 and 13-18, further comprising lysing the macrophage cells and the non-phagocytic cells before a).

34. The method of any one of claims 1-6 and 13-18, further comprising extracting the cellular contents from the macrophage cells and the non-phagocytic cells before a).

35. The method of any one of claims 7-18, further comprising lysing the neutrophil cells and the non-phagocytic cells before a).

36. The method of any one of claims 7-18, further comprising extracting the cellular contents from the neutrophil cells and the non-phagocytic cells before a).

37. The method of claim 34, wherein the cellular contents of the macrophage cells comprise viable diseased cells, dead diseased cells, apoptotic diseased cells, circulating tumor cells, infectious agents, fetal cells, trophoblasts, or fragments thereof.

38. The method of claim 36, wherein the cellular contents of the neutrophil cells comprise viable diseased cells, dead diseased cells, apoptotic diseased cells, circulating tumor cells, infectious agents, fetal cells, trophoblasts, or fragments thereof.

39. The method of claim 34, wherein the selected one or more markers are present in the cellular contents of the macrophage cells.
40. The method of claim 34, wherein the selected one or more markers are not present in the cellular contents of the non-phagocytic cells.
41. The method of any one of claims 1-6 and 13-18, wherein the macrophage cells express the one or more selected C-Macro markers.
42. The method of claim 36, wherein the selected one or more markers are present in the cellular contents of the neutrophil cells.
43. The method of claim 36, wherein the selected one or more markers are not present in the cellular contents of the non-phagocytic cells.
44. The method of any one of claims 7-18, wherein the neutrophil cells express the one or more selected C-Neutro markers.
45. The method of any one of claims 1-18, wherein the non-phagocytic cells are T cells, B cells, null cells, basophils, or mixtures thereof.
46. The method of any one of claims 1-6 and 13-18, wherein the macrophage cells are isolated from a bodily fluid sample, tissues, or cells of the subject.
47. The method of any one of claims 7-18, wherein the neutrophil cells are isolated from a bodily fluid sample, tissues, or cells of the subject.
48. The method of any one of claims 1-18, wherein the non-phagocytic cells are isolated from a bodily fluid sample, tissues, or cells of the subject.

49. The method of any one of claims 46-48, wherein the bodily fluid sample is blood, urine, stool, saliva, lymph fluid, cerebrospinal fluid, synovial fluid, cystic fluid, ascites, pleural effusion, fluid obtained from a pregnant woman in the first trimester, fluid obtained from a pregnant woman in the second trimester, fluid obtained from a pregnant woman in the third trimester, maternal blood, amniotic fluid, chorionic villus sample, fluid from a preimplantation embryo, maternal urine, maternal saliva, placental sample, fetal blood, lavage and cervical vaginal fluid, interstitial fluid, or ocular fluid.

50. The method of any one of claims 1-6 and 13-18, wherein the macrophage cells are isolated using antibodies, using a ligand that binds to a molecular receptor expressed on the plasma membranes of white blood cells, or by flow cytometry, fluorescence activated cell sorting, filtration, gradient-based centrifugation, elution, microfluidics, magnetic separation technique, fluorescent-magnetic separation technique, nanostructure, quantum dots, high throughput microscope-based platforms, or a combination thereof.

51. The method of any one of claims 7-18, wherein the neutrophil cells are isolated using antibodies, using a ligand that binds to a molecular receptor expressed on the plasma membranes of white blood cells, or by flow cytometry, fluorescence activated cell sorting, filtration, gradient-based centrifugation, elution, microfluidics, magnetic separation technique, fluorescent-magnetic separation technique, nanostructure, quantum dots, high throughput microscope-based platforms, or a combination thereof.

52. The method of any one of claims 1-18, wherein the non-phagocytic cells are isolated using antibodies, using a ligand that binds to a molecular receptor expressed on the plasma membranes of white blood cells, or by flow cytometry, fluorescence activated cell sorting, filtration, gradient-based centrifugation, elution, microfluidics, magnetic separation technique, fluorescent-magnetic separation technique, nanostructure, quantum dots, high throughput microscope-based platforms, or a combination thereof.

53. The method of any one of claims 1-6 and 13-18, wherein the macrophage cells are isolated using a product secreted by the macrophage cells.

54. The method of any one of claims 7-18, wherein the neutrophil cells are isolated by using a product secreted by the neutrophil cells.

55. The method of any one of claims 1-6 and 13-18, wherein the macrophage cells are isolated by using a cell surface target on the surface of macrophage cells.

56. The method of any one of claims 7-18, wherein the neutrophil cells are isolated by using a cell surface target on the surface of neutrophil cells.

57. The method of claim 55, wherein the target is expressed by the macrophage cells.

58. The method of claim 55, wherein the target is not expressed by the macrophage cells.

59. The method of claim 56, wherein the target is expressed by the neutrophil cells.

60. The method of claim 56, wherein the target is not expressed by the neutrophil cells.

61. The method of any one of claims 55-60, wherein the target is a marker of said cancer.

62. The method of any one of claims 1-18, wherein the measured levels are gene expression levels.

63. The method of any one of claims 1-18, wherein the measured levels are protein expression levels.

64. The method of any one of claims 1-18, wherein the levels or activities are measured by a qualitative assay, a quantitative assay, or a combination thereof.

65. The method of claim 64, wherein the quantitative assay uses sequencing, direct sequencing, RNA sequencing, whole transcriptome shotgun sequencing, random shotgun sequencing, Sanger dideoxy termination sequencing, whole-genome sequencing, sequencing by hybridization, pyrosequencing, capillary electrophoresis, gel electrophoresis, duplex sequencing, cycle sequencing, single-base extension sequencing, solid-phase sequencing, high-throughput sequencing, massively parallel signature sequencing, emulsion PCR, sequencing by reversible dye terminator, paired-end sequencing, near-term sequencing, exonuclease sequencing, sequencing by ligation, short-read sequencing, single-molecule sequencing, sequencing-by-synthesis, real-time sequencing, reverse-terminator sequencing, nanopore sequencing, 454 sequencing, Solexa Genome Analyzer sequencing, SOLiD® sequencing, MS-PET sequencing, mass spectrometry, matrix assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry, electrospray ionization (ESI) mass spectrometry, surface-enhanced laser desorption/ionization-time of flight (SELDI-TOF) mass spectrometry, quadrupole-time of flight (Q-TOF) mass spectrometry, atmospheric pressure photoionization mass spectrometry (APPI-MS), Fourier transform mass spectrometry (FTMS), matrix-assisted laser desorption/ionization-Fourier transform-ion cyclotron resonance (MALDI-FT-ICR) mass spectrometry, secondary ion mass spectrometry (SIMS), polymerase chain reaction (PCR) analysis, quantitative PCR, real-time PCR, fluorescence assay, colorimetric assay, chemiluminescent assay, or a combination thereof.

66. The method of claim 62, wherein the gene expression levels are measured by polymerase chain reaction (PCR) analysis, sequencing analysis, electrophoretic analysis, restriction fragment length polymorphism (RFLP) analysis, Northern blot

analysis, quantitative PCR, reverse-transcriptase-PCR analysis (RT-PCR), allele-specific oligonucleotide hybridization analysis, comparative genomic hybridization, heteroduplex mobility assay (HMA), single strand conformational polymorphism (SSCP), denaturing gradient gel electrophoresis (DGGE), RNAase mismatch analysis, mass spectrometry, tandem mass spectrometry, matrix assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry, electrospray ionization (ESI) mass spectrometry, surface-enhanced laser desorption/ionization-time of flight (SELDI-TOF) mass spectrometry, quadrupole-time of flight (Q-TOF) mass spectrometry, atmospheric pressure photoionization mass spectrometry (APPI-MS), Fourier transform mass spectrometry (FTMS), matrix-assisted laser desorption/ionization-Fourier transform-ion cyclotron resonance (MALDI-FT-ICR) mass spectrometry, secondary ion mass spectrometry (SIMS), surface plasmon resonance, Southern blot analysis, in situ hybridization, fluorescence in situ hybridization (FISH), chromogenic in situ hybridization (CISH), immunohistochemistry (IHC), microarray, comparative genomic hybridization, karyotyping, multiplex ligation-dependent probe amplification (MLPA), Quantitative Multiplex PCR of Short Fluorescent Fragments (QMPSF), microscopy, methylation specific PCR (MSP) assay, HpaII tiny fragment Enrichment by Ligation-mediated PCR (HELP) assay, radioactive acetate labeling assays, colorimetric DNA acetylation assay, chromatin immunoprecipitation combined with microarray (ChIP-on-chip) assay, restriction landmark genomic scanning, Methylated DNA immunoprecipitation (MeDIP), molecular break light assay for DNA adenine methyltransferase activity, chromatographic separation, methylation-sensitive restriction enzyme analysis, bisulfite-driven conversion of non-methylated cytosine to uracil, methyl-binding PCR analysis, or a combination thereof.

67. The method of claim 62, wherein the gene expression levels are measured by a sequencing technique selected from the group consisting of direct sequencing, RNA sequencing, whole transcriptome shotgun sequencing, random shotgun sequencing, Sanger dideoxy termination sequencing, whole-genome sequencing, sequencing by hybridization, pyrosequencing, capillary electrophoresis, gel

electrophoresis, duplex sequencing, cycle sequencing, single-base extension sequencing, solid-phase sequencing, high-throughput sequencing, massively parallel signature sequencing, emulsion PCR, sequencing by reversible dye terminator, paired-end sequencing, near-term sequencing, exonuclease sequencing, sequencing by ligation, short-read sequencing, single-molecule sequencing, sequencing-by-synthesis, real-time sequencing, reverse-terminator sequencing, nanopore sequencing, 454 sequencing, Solexa Genome Analyzer sequencing, SOLiD® sequencing, MS-PET sequencing, mass spectrometry, and a combination thereof.

68. The method of claim 63, wherein the protein expression levels are measured by an immunohistochemistry assay, an enzyme-linked immunosorbent assay (ELISA), in situ hybridization, chromatography, liquid chromatography, size exclusion chromatography, high performance liquid chromatography (HPLC), gas chromatography, mass spectrometry, tandem mass spectrometry, matrix assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry, electrospray ionization (ESI) mass spectrometry, surface-enhanced laser desorption/ionization-time of flight (SELDI-TOF) mass spectrometry, quadrupole-time of flight (Q-TOF) mass spectrometry, atmospheric pressure photoionization mass spectrometry (APPI-MS), Fourier transform mass spectrometry (FTMS), matrix-assisted laser desorption/ionization-Fourier transform-ion cyclotron resonance (MALDI-FT-ICR) mass spectrometry, secondary ion mass spectrometry (SIMS), radioimmunoassays, microscopy, microfluidic chip-based assays, surface plasmon resonance, sequencing, Western blotting assay, or a combination thereof.

69. The method of any one of claims 1-68, wherein the subject is a mammal.

70. The method of claim 69, wherein the subject is a human.

71. The method of any one of claims 1-18, wherein the difference is greater than a 1-fold difference.

72. The method of claim 71, wherein the difference is at least 1.05-fold, 1.1-fold, 1.2-fold, 1.3-fold, 1.4-fold, 1.5-fold, 2-fold, 2.5-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, or 10-fold difference.

73. A kit for measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200, comprising reagents for specifically measuring the levels of the selected C-Macro marker.

74. A kit for measuring the levels of at least one or more markers selected from the group consisting of C-Neutro 1-200, comprising reagents for specifically measuring the levels of the selected C-Neutro marker.

75. A kit for measuring the levels of at least one or more markers selected from the group consisting of C-Macro 1-200 and at least one or more markers selected from the group consisting of C-Neutro 1-200, comprising reagents for specifically measuring the levels of the selected C-Macro marker and reagents for specifically measuring the levels of the selected C-Neutro marker.

76. The kit of claim 73 or 75, wherein the selected C-Macro markers comprise one or more markers selected from the group consisting of C-Macro 1-4 and C-Macro 101-104.

77. The kit of claim 74 or 75, wherein the selected C-Neutro markers comprise one or more markers selected from the group consisting of C-Neutro 1-8 and C-Neutro 101-108.

78. The kit of any one of claims 73-77, wherein the reagents comprise one or more antibodies or fragments thereof, oligonucleotides, or aptamers.

79. A method of treating or preventing a cancer in a subject comprising administering to said subject an agent that modulates the activity or expression of at least one or more markers selected from the group consisting of C-Macro 1-200.

80. A method of treating or preventing a cancer in a subject comprising administering to said subject an agent that modulates the activity or expression of at least one or more markers selected from the group consisting of C-Neuro 1-200.

81. The method of claim 79 or 80, wherein the agent is a small molecule modulator, siRNA, or an antibody or fragment thereof.

82. The method of any one of claims 1-81, wherein the cancer is a prostate cancer, a head and neck cancer, a lung cancer, melanoma, a colon cancer, a non-small cell lung cancer, a CNS cancer, an ovarian cancer, a renal cancer, or a breast cancer.




Fig. 1

Comparison		Macrophage/T-Cell	Neutrophil/T-Cell
Cancer/Controls	# of genes in signature	4	8
N=44, 44	average error	0.002	0.018
	sensitivity	99.86%	97.7%
	specificity	99.84%	99.96%
N=66, 44	average error	0.011	0.035
	sensitivity	100%	99.2%
	specificity	98.1%	94.6%

Fig. 2

Comparison		# of genes	Paired: Phagocyte/T-cell	Unpaired: Phagocyte
Cancer/Controls	Macrophage	4	0.002	0.012
n=44, 44	Neutrophil	8	0.018	0.024

Fig. 3




Fig. 4

Method	Average # of patients tested with marker test	Average size of purified control set	Upper bound on specificity (TNR)
1. Test all controls	250	172.5	98.6%
2. Test controls with positive gene signature signal ¹	53	202.85	98.3%
2. Test controls with positive gene signature signal ²	57.5	203.375	97.6%

- 1. Assuming 98% sensitivity and specificity of gene signature
- 2. Assuming 95% sensitivity and specificity of gene signature

Fig. 5

Comparison		Macrophage/T-Cell	Neutrophil/T-Cell
Prostate cancer/Head and neck cancer	# of genes in signature	12	9
n=36, 8	average error	0.113	0.07
	sensitivity	75%	87.5%
	specificity	92.1%	94.4%
Prostate cancer/Head and neck cancer	# of genes in signature	10	9
n=58, 8	average error	0.112	0.085
	sensitivity	77.5%	88%
	specificity	90.7%	92.1%

Fig. 6

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US14/22149

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - G01N 33/00, 33/574 (2014.01)

USPC - 435/7.23, 7.24, 7.2, 7.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8): G01N 33/00, 33/574 (2014.01)

USPC: 435/7.23, 7.24, 7.2, 7.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MicroPatent (US-G, US-A, EP-A, EP-B, WO, JP-bib, DE-C,B, DE-A, DE-T, DE-U, GB-A, FR-A); Google Scholar; PubMed; ScienceDirect; 'TNFAIP3,' 'SNAI1,' cancer, macrophage, neutrophil, 'CD83,' biomarker

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2012/0230953 A1 (GOLDENBERG, D et al.) September 13, 2012; paragraphs [0206], [0207]	79, 81/79
X --- Y	WO 2012/115885 A1 (SPETZLER, D et al.) August 30, 2012; paragraphs [0032], [001126]; Table 11	80, 81/80 ----- 1-18, 19/1-19/18, 20/19/1-20/19/18, 21/13-21/18, 22/13-22/18, 23/1-23/6, 23/13-23/18, 24/1-24/6, 24/13-24/18, 25/1-25/6, 25/13-25/18, 26/1-26/6, 26/13-16/18, 27/1-27/6, 27/13-27/18, 28/1-28/6, 28/13-28/18, 29/7-29/18, 30/7/1-30/18, 31/7-31/18, 32/7-32/18, 33/1-33/6, 33/13-33/18, 34/1-34/6, 34/13-34/18, 35/7-35/18, 36/7-36/18, 37/34/1-37/34/6, 37/34/13-37/34/18, 38/36/7-38/36/18, 39/34/1/39/34/6, 39/34/13-39/34/18, 40/34/1-40/34/6, ... Next Page ...



Further documents are listed in the continuation of Box C.



* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

16 June 2014 (16.06.2014)

Date of mailing of the international search report

09 JUL 2014

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Shane Thomas

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US14/22149

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	... Continued from Previous Page ...	40/34/13-40/34/18, 41/1-41/6, 41/13-41/18, 42/36/7-42/36/18, 43/36/7-43/36/18, 44/7-44/18, 45/1-45/18, 46/1-46/6, 46/13-46/18, 47/7-47/18, 48/1-48/18, 50/1-50/6, 50/13-50/18, 51/7-51/18, 52/1-52/18, 53/1-53/6, 53/13-53/18, 54/7-54/18, 55/1-55/6, 55/13-55/18, 56/7-56/18, 57/55/1-57/55/6, 57/55/13-57/55/18, 58/55/1-58/55/6, 58/55/13-58/55/18, 59/56/7-59/56/18, 60/56/7-60/56/18, 62/1-62/18, 63/1-63/18, 64/1-64/18, 65/64/1-65/64/18, 66/62/1-66/62/18, 67/62/1-67/62/18, 68/63/1-68/63/18, 71/1-71/18, 72/71/1-72/71/18, 73-75, 76/73, 76/75, 77/74, 77/75
Y	US 2013/0023426 A1 (KASSIS, AI) January 24, 2013; paragraphs [0014], [0018], [0023], [0025], [0029], [0125]	1-18, 19/1-19/18, 20/19/1-20/19/18, 21/13-21/18, 22/13-22/18, 23/1-23/6, 23/13-23/18, 24/1-24/6, 24/13-24/18, 25/1-25/6, 25/13-25/18, 26/1-26/6, 26/13-16/18, 27/1-27/6, 27/13-27/18, 28/1-28/6, 28/13-28/18, 29/7-29/18, 30/7/1-30/18, 31/7-31/18, 32/7-32/18, 33/1-33/6, 33/13-33/18, 34/1-34/6, 34/13-34/18, 35/7-35/18, 36/7-36/18, 37/34/1-37/34/6, 37/34/13-37/34/18, 38/36/7-38/36/18, 39/34/1/39/34/6, 39/34/13-39/34/18, 40/34/1-40/34/6, 40/34/13-40/34/18, 41/1-41/6, 41/13-41/18, 42/36/7-42/36/18, 43/36/7-43/36/18, 44/7-44/18, 45/1-45/18, 46/1-46/6, 46/13-46/18, 47/7-47/18, 48/1-48/18, 50/1-50/6, 50/13-50/18, 51/7-51/18, 52/1-52/18, 53/1-53/6, 53/13-53/18, 54/7-54/18, 55/1-55/6, 55/13-55/18, 56/7-56/18, 57/55/1-57/55/6, 57/55/13-57/55/18, 58/55/1-58/55/6, 58/55/13-58/55/18, 59/56/7-59/56/18, 60/56/7-60/56/18, 62/1-62/18, 63/1-63/18, 64/1-64/18, ... Next Page ...

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US14/22149

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	... Continued from Previous Page ...	65/64/1-65/64/18, 66/62/1-66/62/18, 67/62/1-67/62/18, 68/63/1-68/63/18, 71/1-71/18, 72/71/1-72/71/18, 73-75, 76/73, 76/75, 77/74, 77/75
Y	VENDRELL, J et al. A20/TNFAIP3, A New Estrogen-Regulated Gene That Confers Tamoxifen Resistance In Breast Cancer Cells. Oncogene. July 12, 2007, Vol. 26, No. 32; pages 4656-4667; page 4656, right column, second paragraph; page 4657, left column, first paragraph. DOI:10.1038/sj.onc.1210269.	25/1-25/6, 25/13-25/18, 26/1-26/6, 26/13-16/18
Y	MARTIN, T et al. Expression Of The Transcription Factors Snail, Slug, And Twist And Their Clinical Significance In Human Breast Cancer. Annals of Surgical Oncology. June 2005, Vol. 12, No. 6; pages 488-496; abstract. DOI: 10.1245/AO.2005.04.010.	27/1-27/6, 27/13-27/18, 28/1-28/6, 28/13-28/18, 31/7-31/18, 32/7-32/18

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US14/22149

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 49, 61, 69, 70, 78, 82
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.