

**(12) PATENT
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 199893162 B2
(10) Patent No. 759717**

(54) Title
April- a novel protein with growth effects

(51)⁷ International Patent Classification(s)
C07K 014/00

(21) Application No: **199893162** (22) Application Date: **1998.09.11**

(87) WIPO No: **WO99/12965**

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/058786	1997.09.12	US
60/079384	1998.03.26	US

(43) Publication Date : **1999.03.29**

(43) Publication Journal Date : **1999.05.27**

(44) Accepted Journal Date : **2003.04.17**

(71) Applicant(s)
Apotech S.A.

(72) Inventor(s)
Jurg Tschopp

(74) Agent/Attorney
CULLEN and CO, GPO Box 1074, BRISBANE QLD 4001

OPI DATE 29/03/99 APPLN. ID 93162/98
AOJP DATE 27/05/99 PCT NUMBER PCT/US98/19191



AU9893162

IN

(51) International Patent Classification 6 : C07K 14/00		A2	(11) International Publication Number: WO 99/12965 (43) International Publication Date: 18 March 1999 (18.03.99)
(21) International Application Number: PCT/US98/19191			
(22) International Filing Date: 11 September 1998 (11.09.98)			
(30) Priority Data: 60/058,786 12 September 1997 (12.09.97) US 60/079,384 26 March 1998 (26.03.98) US			
(71) Applicant (for all designated States except US): <i>Biogen, Inc.</i> [US/US]; 14 Cambridge Center, Cambridge, MA 02142 (US): 84 Rue du Rhone, CH-1204 Geneva, Switzerland.			
(72) Inventor; and (75) Inventor/Applicant (for US only): TSCHOPP, Jurg [CH/CH]; Chemin des Fontannins 10, CH-1066 Epalinges (CH).			
(74) Agent: FLYNN, Kerry; Biogen, Inc., 14 Cambridge Center, Cambridge, MA 02142 (US).			
(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).			
Published <i>Without international search report and to be republished upon receipt of that report.</i>			



(54) Title: APRIL- A NOVEL PROTEIN WITH GROWTH EFFECTS

(57) Abstract

APRIL, a novel member of the tumor necrosis factor family (TNF), modified APRILs, and pharmaceutical compositions comprising them.

APRIL- A NOVEL PROTEIN WITH GROWTH EFFECTS

BACKGROUND OF THE INVENTION

The present invention relates to novel ligand and polypeptides which are 5 members of the Tumor Necrosis Factor Family. The novel ligand is designated April for "A Proliferation Inducing Ligand." These proteins or their receptors may have anti-cancer and/or immunoregulatory applications. Furthermore, cells transfected with 10 the genes for these novel ligands may be used in gene therapy to treat tumors, autoimmune and inflammatory diseases or inherited genetic disorders, and blocking antitibodies to these proteins can have immunoregulatory applications.

BACKGROUND OF THE INVENTION

The tumor-necrosis factor (TNF)-related cytokines are mediators of host 15 defense and immune regulation. Members of this family exist in membrane-anchored forms, acting locally through cell-to-cell contact, or as secreted proteins capable of diffusing to more distant targets. A parallel family of receptors signals the presence of 20 these molecules leading to the initiation of cell death or cellular proliferation and differentiation in the target tissue. Presently, the TNF family of ligands and receptors has at least 13 recognized receptor-ligand pairs, including: TNF:TNF-R; LT- α :TNF-R; LT- α / β :LT- β -R; FasL:Fas; CD40L:CD40; CD30L:CD30; CD27L:CD27; OX40L:OX40 and 4-1BBL:4-1BB; tranne/rankL: Light and Tweak. The DNA sequences encoding these ligands have only about 25% to about 30% identity in even 25 the most related cases, although the amino acid relatedness is about 50%.

The defining feature of this family of cytokine receptors is found in the cysteine 30 rich extracellular domain initially revealed by the molecular cloning of two distinct TNF receptors.¹ This family of genes encodes glycoproteins characteristic of Type I transmembrane proteins with an extracellular ligand binding domain, a single membrane spanning region and a cytoplasmic region involved in activating cellular functions. The cysteine-rich ligand binding region exhibits a tightly knit disulfide linked core domain, which, depending upon the particular family member, is repeated multiple times. Most receptors have four domains, although there may be as few as three, or as many as six.

Proteins in the TNF family of ligands are characterized by a short N-terminal stretch of normally short hydrophilic amino acids, often containing several lysine or arginine residues thought to serve as stop transfer sequences. Next follows a

transmembrane region and an extracellular region of variable length, that separates the C-terminal receptor binding domain from the membrane. This region is sometimes referred to as the "stalk". The C-terminal binding region comprises the bulk of the protein, and often, but not always, contains glycosylation sites. These genes lack the 5 classic signal sequences characteristic of type I membrane proteins, type II membrane proteins with the C terminal domain lying outside the cell, and a short N-terminus residing in the cytoplasm. In some cases, e.g., TNF and LT- α , cleavage in the stalk region can occur early during protein processing and the ligand is then found primarily in secreted form. Most ligands, however, exist in a membrane form, mediating 10 localized signaling.

The structure of these ligands has been well-defined by crystallographic analyses of TNF, LT- α , and CD40L. TNF and lymphotoxin- α (LT- α) are both structured into a sandwich of two anti-parallel β -pleated sheets with the "jelly roll" or Greek key topology.ⁱⁱ The rms deviation between the C α and β residues is 0.61 C, 15 suggesting a high degree of similarity in their molecular topography. A structural feature emerging from molecular studies of CD40L, TNF and LT- α is the propensity to assemble into oligomeric complexes. Intrinsic to the oligomeric structure is the formation of the receptor binding site at the junction between the neighboring subunits creating a multivalent ligand. The quaternary structures of TNF, CD40L and LT- α 20 have been shown to exist as trimers by analysis of their crystal structures. Many of the amino acids conserved between the different ligands are in stretches of the scaffold β -sheet. It is likely that the basic sandwich structure is preserved in all of these molecules, since portions of these scaffold sequences are conserved across the various 25 family members. The quaternary structure may also be maintained since the subunit conformation is likely to remain similar.

TNF family members can best be described as master switches in the immune system controlling both cell survival and differentiation. Only TNF and LT α are currently recognized as secreted cytokines contrasting with the other predominantly membrane anchored members of the TNF family. While a membrane form of TNF has 30 been well-characterized and is likely to have unique biological roles, secreted TNF functions as a general alarm signaling to cells more distant from the site of the triggering event. Thus TNF secretion can amplify an event leading to the well-described changes in the vasculature lining and the inflammatory state of cells. In contrast, the membrane bound members of the family send signals through the TNF type

receptors only to cells in direct contact. For example T cells provide CD40 mediated "help" only to those B cells brought into direct contact via cognate TCR interactions. Similar cell-cell contact limitations on the ability to induce cell death apply to the well-studied Fas system.

5 It appears that one can segregate the TNF ligands into three groups based on their ability to induce cell death (Table III). First, TNF, Fas ligand and TRAIL can efficiently induce cell death in many lines and their receptors mostly likely have good canonical death domains. Presumably the ligand to DR-3 (TRAMP/WSL-1) would also fall into this category. Next there are those ligands which trigger a weaker death
10 signal limited to few cell types and TWEAK, CD30 ligand and LT α 1 β 2 are examples of this class. How this group can trigger cell death in the absence of a canonical death domain is an interesting question and suggests that a separate weaker death signaling mechanism exists. Lastly, there are those members that cannot efficiently deliver a death signal. Probably all groups can have antiproliferative effects on some cell types
15 consequent to inducing cell differentiation e.g. CD40 (Funakoshi et al., 1994)

The TNF family has grown dramatically in recent years to encompass at least 11 different signaling pathways involving regulation of the immune system. The widespread expression patterns of TWEAK and TRAIL indicate that there is still more functional variety to be uncovered in this family. This aspect has been especially
20 highlighted recently in the discovery of two receptors that affect the ability of rous sarcoma and herpes simplex virus to replicate as well as the historical observations that TNF has anti-viral activity and pox viruses encode for decoy TNF receptors (Brojatsch et al., 1996; Montgomery et al., 1996; Smith, 1994; Vassalli, 1992). TNF is a mediator of septic shock and cachexiaⁱⁱⁱ, and is involved in the regulation of hematopoietic cell
25 development.^{iv} It appears to play a major role as a mediator of inflammation and defense against bacterial, viral and parasitic infections^v as well as having antitumor activity.^{vi} TNF is also involved in different autoimmune diseases.^{vii} TNF may be produced by several types of cells, including macrophages, fibroblasts, T cells and natural killer cells.^{viii} TNF binds to two different receptors, each acting through
30 specific intracellular signaling molecules, thus resulting in different effects of TNF.^{ix} TNF can exist either as a membrane bound form or as a soluble secreted cytokine.^x

LT- α shares many activities with TNF, i.e. binding to the TNF receptors,^{xi} but unlike TNF, appears to be secreted primarily by activated T cells and some β -lymphoblastoid tumors.^{xii} The heteromeric complex of LT- α and LT- β is a membrane

bound complex which binds to the LT- β receptor.^{xiii} The LT system (LTs and LT-R) appears to be involved in the development of peripheral lymphoid organs since genetic disruption of LT- β leads to disorganization of T and B cells in the spleen and an absence of lymph nodes.^{xiv} The LT- β system is also involved in cell death of some adenocarcinoma cell lines.^{xv}

5 Fas-L, another member of the TNF family, is expressed predominantly on activated T cells.^{xvi} It induces the death of cells bearing its receptor, including tumor cells and HIV-infected cells, by a mechanism known as programmed cell death or apoptosis.^{xvii} Furthermore, deficiencies in either Fas or Fas-L may lead to 10 lymphoproliferative disorders, confirming the role of the Fas system in the regulation of immune responses.^{xviii} The Fas system is also involved in liver damage resulting from hepatitis chronic infection^{xix} and in autoimmunity in HIV-infected patients.^{xx} The Fas system is also involved in T-cell destruction in HIV patients.^{xxi} TRAIL, another 15 member of this family, also seems to be involved in the death of a wide variety of transformed cell lines of diverse origin.^{xxii}

CD40-L, another member of the TNF family, is expressed on T cells and induces the regulation of CD40-bearing B cells.^{xxiii} Furthermore, alterations in the CD40-L gene result in a disease known as X-linked hyper-IgM syndrome.^{xxiv} The CD40 system is also involved in different autoimmune diseases^{xxv} and CD40-L is 20 known to have antiviral properties.^{xxvi} Although the CD40 system is involved in the rescue of apoptotic B cells,^{xxvii} in non-immune cells it induces apoptosis^{xxviii}. Many additional lymphocyte members of the TNF family are also involved in costimulation.^{xxix}

25 Generally, the members of the TNF family have fundamental regulatory roles in controlling the immune system and activating acute host defense systems. Given the current progress in manipulating members of the TNF family for therapeutic benefit, it is likely that members of this family may provide unique means to control disease. Some of the ligands of this family can directly induce the apoptotic death of many transformed cells e.g. LT, TNF, Fas ligand and TRAIL (Nagata, 1997). Fas and 30 possibly TNF and CD30 receptor activation can induce cell death in nontransformed lymphocytes which may play an immunoregulatory function (Amakawa et al., 1996; Nagata, 1997; Sytwu et al., 1996; Zheng et al., 1995). In general, death is triggered following the aggregation of death domains which reside on the cytoplasmic side of the TNF receptors. The death domain orchestrates the assembly of various signal

transduction components which result in the activation of the caspase cascade (Nagata, 1997). Some receptors lack canonical death domains, e.g. LT_B receptor and CD30 (Browning et al., 1996; Lee et al., 1996) yet can induce cell death, albeit more weakly. It is likely that these receptors function primarily to induce cell differentiation and the 5 death is an aberrant consequence in some transformed cell lines, although this picture is unclear as studies on the CD30 null mouse suggest a death role in negative selection in the thymus (Amakawa et al., 1996). Conversely, signaling through other pathways such as CD40 is required to maintain cell survival. Thus, there is a need to identify and characterize additional molecules which are members of the TNF family thereby 10 providing additional means of controlling disease and manipulating the immune system.

It has been suggested that certain members of the TNF family may provide therapeutic anti-tumor benefits, for example, in combination with IL-2. (See, e.g. U.S. 5,425,940). However, to date, no completely satisfactory treatment for 15 cancer is known. Combination chemotherapy is commonly used in the clinic and in research, for example with antimetabolites, alkylating agents, antibiotics, general poisons, etc. Such drugs are administered alone or in combination in an attempt to obtain a cytotoxic effect on cancers, and/or to reduce or eliminate the emergence of drug-resistant cells, and to reduce side effects.

20 **SUMMARY OF THE INVENTION**

Accordingly, the present invention is directed to a novel polypeptide referred to as APRIL, which substantially obviates one or more of the problems due to the 25 limitations and disadvantages of the related art. The inventors have discovered a new member of the TNF family of cytokines, and defined both the human and murine amino acid sequence of the protein, as well as the DNA sequences encoding these proteins. The claimed invention may be used to identify new diagnostics and therapeutics for numerous diseases and conditions as discussed in more detail below, as well as to obtain information about, and manipulate, the immune system and its processes. Additionally, the invention may be involved in the induction of cell death in 30 carcinomas.

Additional features and advantages of the invention will be set forth in the description which follows, and in part will be apparent from the description, or may be learned by practice of the invention. The objectives and other advantages of the

invention will be realized and attained by the compositions and methods particularly pointed out in the written description and claims hereof, as well as in the appended drawings.

Thus, to achieve these and other advantages, and in accordance with the purpose 5 of the invention, as embodied and broadly described herein, the invention includes DNA sequences encoding APRIL. Specifically, the invention relates to DNA sequences which encode human APRIL, (SEQ. ID. NO. 1). Additionally, the claimed invention relates to the amino acid sequences of this novel ligand. The amino acid sequence of human APRIL is set forth in SEQ. ID. NO.: 2. Additionally, the inventors 10 have set forth herein the DNA and amino acid sequences for murine APRIL, set forth in SEQ.ID. NOS. 3 and 4 respectively. In other embodiments, the invention relates to sequences that have at least 50% homology with DNA sequences encoding the C terminal receptor binding domain of this ligand and when hybridize to the claimed DNA sequences or fragments thereof, and which encode APRIL having the sequence 15 identified in SEQ. ID. NO. 1 or a protein having similar biological activity.

The invention in certain embodiments furthermore relates to DNA sequences encoding APRIL where the sequences are operatively linked to an expression control sequence. Any suitable expression control sequences are useful in the claimed invention, and can easily be selected by one skilled in the art.

20 The invention also contemplates recombinant DNAs comprising a sequence encoding APRIL or fragments thereof, as well as hosts with stably integrated APRIL sequences introduced into their genome, or possessing episomal elements. Any suitable host may be used in the invention, and can easily be selected by one skilled in the art without undue experimentation.

25 In other embodiments, the invention relates to methods of producing substantially pure APRILs comprising the step of culturing transformed hosts. In yet other embodiments, the invention relates to APRIL essentially free of normally associated animal proteins.

The invention encompasses APRIL ligands having the amino acid sequence 30 identified in SEQ. ID. NO. 2, as well as fragments or homologs thereof. In various embodiments, the amino acid and/or the DNA sequences may comprise conservative insertions, deletions and substitutions, as further defined below or may comprise fragments of said sequences.

The invention relates in other embodiments to soluble constructs comprising APRIL, which may be used to directly trigger APRIL mediated pharmacological events. Such events may have useful therapeutic benefits in the stimulation of growth, treatment of cancer, tumors or the manipulation of the immune system to treat 5 immunologic diseases. Soluble forms of the claimed ligands could be genetically reengineered to incorporate an easily recognizable tag, thereby facilitating the identification of the receptors for these ligands.

Additionally, certain embodiments relate to antibodies against APRIL, and their use for the treatment of cancers, tumors, or manipulation of the immune system to 10 treat immunologic diseases.

In yet other embodiments the invention relates to methods of gene therapy using the genes for APRIL as disclosed and claimed herein.

The pharmaceutical preparations of the invention may, optionally, include pharmaceutically acceptable carriers, adjuvants, fillers, or other pharmaceutical 15 compositions, and may be administered in any of the numerous forms or routes known in the art.

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory, and are intended to provide further explanation of the invention as claimed.

20 The accompanying drawings are included to provide a further understanding of the invention, and are incorporated in, and constitute a part of this specification, illustrate several embodiments of the invention, and together with the description serve to explain the principles of the invention.

25 **BRIEF DESCRIPTION OF THE DRAWINGS**

Figure legends

Figure 1. (A) Predicted amino acid sequence of human APRIL. The predicted transmembrane region (TM, boxed), the potential N-linked glycosylation site (star) and 30 the N-terminus of the recombinant soluble APRIL (sAPRIL) are indicated. (B) Comparison of the extracellular protein sequence of APRIL and some members of the TNF ligand family. Identical and homologous residues are represented in black and shaded boxes, respectively. TNFa, tumor necrosis factor α , LT α , lymphotoxin α , FasL, Fas (CD95) ligand, TRANCE, RANK ligand.

Figure 2. Expression of APRIL. (A) Northern blots (2 μ g poly A⁺ RNA per lane) of various human tissues were probed with APRIL cDNA. (B) APRIL mRNA expression in various tumor cell lines: promyelocytic leukemia HL 60; HeLa Cell S3; chronic myelogenous leukemia K562; lymphoblastic leukemia Molt-4; Burkitt's lymphoma Raji; colorectal adenocarcinoma A459; melanoma G361. (C) APRIL mRNA expression in four different human tumors (T) and normal tissues (N). The 18S rRNA band shows equal loading. (D) APRIL mRNA expression in primary colon carcinoma. In situ hybridization revealed abundant APRIL message in human colon carcinoma as compared to normal colon tissue. Colon tumor tissue sections and adjacent normal colon tissue were hybridized to antisense APRIL 35 S-labeled cRNA, and as control, colon tumor tissue sections were also hybridized to sense APRIL 35 S cRNA (negative control). The upper panels are dark field micrographs, the lower panels are the corresponding light field micrographs.

15

Figure 3. APRIL stimulates cell growth. (A) Dose dependent increase of proliferation of Jurkat (human leukemia T cells), as determined 24 hrs after addition of soluble APRIL. Controls are Fas ligand (FasL), TWEAK and no ligand (Control) (left panel, cell viability; right panel, 3 H-Thymidine incorporation). (B) Influence of immunodepletion of FLAG-tagged APRIL on tumor cell growth. The proliferative effect of FLAG-tagged APRIL is neutralized by anti-FLAG antibodies, but not by anti-myc antibodies. (C) Effect of APRIL on the proliferation rate of Raji (human Burkitt lymphoma B cells), A20 cells (mouse B lymphoma), BJAB (human B lymphoma), COS (canine epithelial cells), MCF-7 (human breast adenocarcinoma), HeLa (human epitheloid carcinoma) and ME260 (human melanoma). (D) Influence of fetal calf serum concentration on APRIL-induced proliferation of Jurkat cells.

Figure 4. APRIL accelerates tumor growth. (A) Characterization of APRIL-transfected NIH-3T3 clones. FLAG-APRIL levels of the various clones were analyzed by Western blotting using an anti-FLAG antibody. The arrow points to the APRIL protein, the high molecular weight protein is detected non-specifically (B) APRIL-expressing NIH-3T3 clones grow faster than mock-transfected clones. (C) Increased

tumor growth of APRIL-expressing NIH-3T3 clones. NIH-3T3 cells (1×10^5 cells) and APRIL (NIH-AP, 2 different clones) transfectants (1×10^6 cells) were injected subcutaneously into nude mice, and tumor growth monitored.

Figure 5. An alignment of the human and mouse APRIL amino acid sequences showing the extensive identity between the two proteins. Identical residues are marked with the overlaying dot. The underlined residues represent a potential N-linked glycosylation site. The initiating methionine is considered a likely start site, however, it is possible that in frame methionines further upstream may serve as the actual start site, for example, in the human sequence.

10

DETAILED DESCRIPTION

Reference will now be made in detail to the present preferred embodiments of the invention. This invention relates to DNA sequences that code for human or mouse APRIL, fragments and homologs thereof, and expression of those DNA sequences in hosts transformed with them. The invention relates to uses of these DNA sequences and the peptides encoded by them. Additionally, the invention encompasses both human and mouse amino acid sequences for APRIL, or fragments thereof, as well as pharmaceutical compositions comprising or derived from them. The invention relates to methods of stimulating cell growth with APRIL, or, alternatively, methods of inhibiting tumorogenesis using antibodies directed against APRIL or a receptor of APRIL.

A. DEFINITIONS

“Homologous”, as used herein, refers to the sequence similarity between sequences of molecules being compared. When a position in both of the two compared sequences is occupied by the same base or amino acid monomer subunit, e.g., if a position in each of two DNA molecules is occupied by adenine, then the molecules are homologous at that position. The percent of homology between two sequences is a function of the number of matching or homologous positions shared by the two sequences divided by the number of positions compared x 100. For example, if 6 of 10 of the positions in two sequences are matched or homologous then the two sequences are 60% homologous. By way of example, the DNA sequences ATTGCC and TATGGC share 50% homology. Generally, a comparison is made when two sequences are aligned to give maximum homology.

As used herein, the term "cancer" refers to any neoplastic disorder, including such cellular disorders as, for example, renal cell cancer, Kaposi's sarcoma, chronic leukemia, breast cancer, sarcoma, ovarian carcinoma, rectal cancer, throat cancer, melanoma, colon cancer, bladder cancer, mastocytoma, lung cancer, mammary 5 adenocarcinoma, pharyngeal squamous cell carcinoma, and gastrointestinal or stomach cancer. Preferably, the cancer is leukemia, mastocytoma, melanoma, lymphoma, mammary adenocarcinoma, and pharyngeal squamous cell carcinoma

A "purified preparation" or a "substantially pure preparation" of a polypeptide, as used herein, means a polypeptide that has been separated from other proteins, lipids, 10 and nucleic acids with which it naturally occurs. Preferably, the polypeptide is also separated from other substances, e.g., antibodies, matrices, etc., which are used to purify it.

"Transformed host" as used herein is meant to encompass any host with stably integrated sequence, i.e. a sequence encoding APRIL, introduced into its genome.

15 A "treatment", as used herein, includes any therapeutic treatment, e.g., the administration of a therapeutic agent or substance, e.g., a drug.

A "substantially pure nucleic acid", e.g., a substantially pure DNA, is a nucleic acid which is one or both of: (1) not immediately contiguous with either one or both of the sequences, e.g., coding sequences, with which it is immediately contiguous (i.e., 20 one at the 5' end and one at the 3' end) in the naturally-occurring genome of the organism from which the nucleic acid is derived; or (2) which is substantially free of a nucleic acid sequence with which it occurs in the organism from which the nucleic acid is derived. The term includes, for example, a recombinant DNA which is incorporated into a vector, e.g., into an autonomously replicating plasmid or virus, or into the 25 genomic DNA of a prokaryote or eukaryote, or which exists as a separate molecule (e.g., a cDNA or a genomic DNA fragment produced by PCR or restriction endonuclease treatment) independent of other DNA sequences. Substantially pure DNA also includes a recombinant DNA which is part of a hybrid gene encoding APRIL.

30 The terms "peptides", "proteins", and "polypeptides" are used interchangeably herein.

"Biologically active" as used herein, means having an in vivo or in vitro activity which may be performed directly or indirectly. Biologically active fragments of APRIL may have, for example, 70% amino acid homology with the active site of

APRIL, more preferably at least 80%, and most preferably, at least 90% homology. Identity or homology with respect to APRIL is defined herein as the percentage of amino acid residues in the candidate sequence which are identical to the APRIL residues in SEQ. ID. NOS. 2 or 4.

5 “Ligand” as used herein generically refers to APRIL. The practice of the present invention will employ, unless otherwise indicated, conventional techniques of cell biology, cell culture, molecular biology, transgenic biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are described in the literature.

10 Introduction:

APRIL, a novel member of the TNF family, is described in detail herein. The inventors have found that while transcript of APRIL are of low abundance in normal tissues, high levels of mRNA are detected in several tumor cell lines, as well as in colon carcinomas, metastatic lymphomas and thyroid tumors. In vitro, the addition of recombinant 15 APRIL stimulates the proliferation of various cell lines. Moreover, transfection of APRIL into NIH-3T3 cells dramatically accelerated tumor growth in nude mice when compared to mock transfectants. The expression and growth stimulating effect of APRIL on tumor cells in vitro and in vivo suggests that APRIL is implicated in tumorigenesis.

20 APRIL appears to be unique among the members of the TNF family as it is both abundantly expressed in tumor cells and stimulates growth of many different tumor cell lines given the apparent role of APRIL in tumorigenesis, the antagonistic antibodies to APRIL, or the APRIL receptor, will provide novel approaches to cancer treatment.

B. DNA SEQUENCES OF THE INVENTION

25 As described herein, one aspect of the invention features a substantially pure (or recombinant) nucleic acid which includes a nucleotide sequence encoding APRIL, such as the DNA described in SEQ. ID. NO. 1 and/or equivalents of such nucleic acids. The term nucleic acid as used herein can include fragments and equivalents, such as, for example, sequences encoding functionally equivalent peptides. Equivalent nucleotide sequences may include sequences that differ by one or more nucleotide substitutions, 30 additions or deletions, such as allelic variants, mutations, etc. and include sequences

that differ from the nucleotide sequence encoding APRIL shown in SEQ. ID NO: 1 due to the degeneracy of the genetic code.

The invention will be described generally by reference to the human sequences, although one skilled in the art will understand that the mouse sequences or sequences 5 encoding APRIL from other species having a high level of homology with human, and are encompassed herein. The human proteins appear to have all of the characteristics of the TNF family, i.e., a type II membrane protein organization and conservation of the sequence motifs involved in the folding of the protein into the TNF anti-parallel β -sheet structure.

10 The sequences of the invention can be used to prepare a series of DNA probes that are useful in screening various collections of natural and synthetic DNAs for the presence of DNA sequences that are closely related to APRIL, or fragments or derivatives thereof. One skilled in the art will recognize that reference to APRIL as used herein, refers also to biologically active derivatives, fragments or homologs 15 thereof.

20 The DNA sequences of the invention coding on APRIL can be employed to produce the claimed peptides on expression in various prokaryotic and eukaryotic hosts transformed with them. These peptides may be used in anti-cancer, and immunoregulatory applications. In general, this comprises the steps of culturing a host 25 transformed with a DNA molecule containing the sequence encoding APRIL, operatively-linked to an expression control sequence.

25 The DNA sequences and recombinant DNA molecules of the present invention can be expressed using a wide variety of host/vector combinations. For example, useful vectors may consist of segments of chromosomal, non-chromosomal or synthetic DNA sequences. The expression vectors of the invention are characterized by at least one expression control sequence that may be operatively linked to the APRIL DNA sequence inserted in the vector, in order to control and to regulate the expression of the DNA sequence.

30 Furthermore, within each expression vector, various sites may be selected for insertion of a sequence of the invention. The sites are usually designated by a restriction endonuclease which cuts them, and these sites and endonucleases are well recognized by those skilled in the art. It is of course to be understood that an expression vector useful in this invention need not have a restriction endonuclease site for insertion of the desired DNA fragment. Instead, the vector may be cloned to the

fragment by alternate means. The expression vector, and in particular the site chosen therein for insertion of a selected DNA fragment, and its operative linking therein to an expression control sequence, is determined by a variety of factors. These factors include, but are not limited to, the size of the protein to be expressed, the susceptibility

5 of the desired protein to proteolytic degradation by host cell enzymes, number of sites susceptible to a particular restriction enzyme, contamination or binding of the protein to be expressed by host cell proteins which may prove difficult to remove during purification. Additional factors which may be considered include expression characteristics such as the location of start and stop codons relative to the vector
10 sequences, and other factors which will be recognized by those skilled in the art. The choice of a vector and insertion site for the claimed DNA sequences is determined by a balancing of these factors, not all selections being equally effective for a desired application. However, it is routine for one skilled in the art to analyze these parameters and choose an appropriate system depending on the particular application.

15 One skilled in the art can readily make appropriate modifications to the expression control sequences to obtain higher levels of protein expression, i.e. by substitution of codons, or selecting codons for particular amino acids that are preferentially used by particular organisms, to minimize proteolysis or to alter glycosylation composition. Likewise, cysteines may be changed to other amino acids
20 to simplify production, refolding or stability problems.

Thus, not all host/expression vector combinations function with equal efficiency in expressing the DNA sequences of this invention. However, a particular selection of a host/expression vector combination may be made by those of skill in the art. Factors one may consider include, for example, the compatibility of the host and vector,
25 toxicity to the host of the proteins encoded by the DNA sequence, ease of recovery of the desired protein, expression characteristics of the DNA sequences and expression control sequences operatively linked to them, biosafety, costs and the folding, form or other necessary post-expression modifications of the desired protein.

APRIL produced by hosts transformed with the sequences of the invention, as
30 well as native APRIL purified by the processes of this invention, or produced from the claimed amino acid sequences, are useful in a variety of compositions and methods for anticancer, antitumor and immunoregulatory applications. They are also useful in therapy and methods directed to other diseases.

This invention also relates to the use of the DNA sequences disclosed herein to express APRIL under abnormal conditions, i.e. in a gene therapy setting. Additionally, APRIL may be expressed in tumor cells under the direction of promoters appropriate for such applications. Such expression could enhance anti-tumor immune responses or 5 directly affect the survival of the tumor. APRIL is also likely to affect the survival of an organ graft by altering the local immune response. In this case, the graft itself or the surrounding cells would be modified with an engineered gene encoding APRIL.

Another aspect of the invention relates to the use of the isolated nucleic acid encoding either APRIL in "antigens" therapy. As used herein, "antisense" therapy 10 refers to administration or *in situ* generation of oligonucleotides or their derivatives which specifically hybridize under cellular conditions with the cellular mRNA and/or DNA encoding the APRIL sequence of interest, so as to inhibit expression of the encoded protein, i.e. by inhibiting transcription and/or translation. The binding may be by conventional base pair complementarity, or, for example, in the case of binding to 15 DNA duplexes, through specific interactions in the major groove of the double helix. In general, "antisense" therapy refers to a range of techniques generally employed in the art, and includes any therapy which relies on specific binding to oligonucleotide sequences.

An antisense construct of the present invention can be delivered, for example, as 20 an expression plasmid, which, when transcribed in the cell, produces RNA which is complementary to at least a portion of the cellular mRNA which encodes APRIL. Alternatively, the antisense construct can be an oligonucleotide probe which is generated *ex vivo*. Such oligonucleotide probes are preferably modified 25 oligonucleotides which are resistant to endogenous nucleases, and are therefore stable *in vivo*. Exemplary nucleic acids molecules for use as antisense oligonucleotides are phosphoramidates, phosphothioate and methylphosphonate analogs of DNA (See, e.g., 5,176,996; 5,264,564; and 5,256,775). Additionally, general approaches to constructing oligomers useful in antisense therapy have been reviewed, for example, by Van Der Krol et al., (1988) *Biotechniques* 6:958-976; and Stein et al. (1988) *Cancer Res* 48: 30 2659-2668, specifically incorporated herein by reference.

C. APRIL AND AMINO ACID SEQUENCES THEREFOR

APRIL, as discussed above, is a member of the TNF family. The protein, fragments or homologs of APRIL may have wide therapeutic and diagnostic applications as discussed in more detail below.

5 Although the precise three dimensional structure of APRIL is not known, it is predicted that, as a member of the TNF family, it may share certain structural characteristics with other members of the family.

10 Comparison of the claimed APRIL sequence with other members of the human TNF family reveals considerable structural similarity. All the proteins share several regions of sequence conservation in the extracellular domain. The overall sequence homology of the extracellular domain of APRIL show the highest homology to FasL (21% amino acid identities), TNF α (20%), LT- β (18%), followed by TRAIL, TWEAK and TRANCE (15%). Figure 2.

15 The novel polypeptides of the invention specifically interact with a receptor, which has not yet been identified. However, the peptides and methods disclosed herein enable the identification of receptors which specifically interact with APRIL or fragments thereof.

20 The claimed invention in certain embodiments includes peptides derived from APRIL which have the ability to bind to its receptors. Fragments of APRIL can be produced in several ways, e.g., recombinantly, by PCR, proteolytic digestion or by chemical synthesis. Internal or terminal fragments of a polypeptide can be generated by removing one or more nucleotides from one end or both ends of a nucleic acid which encodes the polypeptide. Expression of the mutagenized DNA produces polypeptide fragments.

25 Polypeptide fragments can also be chemically synthesized using techniques known in the art such as conventional Merrifield solid phase f- moc or t-boc chemistry. For example, peptides and DNA sequences of the present invention may be arbitrarily divided into fragments of desired length with no overlap of the fragment, or divided into overlapping fragments of a desired length. Methods such as these are described in more detail below.

30 D. Generation of Soluble Forms of APRIL

Soluble forms of APRIL can often signal effectively and hence can be administered as a drug which now mimics the natural membrane form. It is possible that APRIL as claimed herein is naturally secreted as a soluble cytokine, however, if not, one can reengineer the gene to force secretion. To create a soluble secreted form

of APRIL, one would remove at the DNA level the N-terminus transmembrane regions, and some portion of the stalk region, and replace them with a type I leader or alternatively a type II leader sequence that will allow efficient proteolytic cleavage in the chosen expression system. A skilled artisan could vary the amount of the stalk region retained in the secretion expression construct to optimize both receptor binding properties and secretion efficiency. For example, the constructs containing all possible stalk lengths, i.e. N-terminal truncations, could be prepared such that proteins starting at amino acids 81 to 139 would result. The optimal length stalk sequence would result from this type of analysis.

10 E. Generation of Antibodies Reactive with APRIL

The invention also includes antibodies specifically reactive with APRIL or its receptor. Anti-protein/anti-peptide antisera or monoclonal antibodies can be made by standard protocols (See, for example, *Antibodies: A Laboratory Manual* ed. by Harlow and Lane (Cold Spring Harbor Press: 1988)). A mammal such as a mouse, a hamster or 15 rabbit can be immunized with an immunogenic form of the peptide. Techniques for conferring immunogenicity on a protein or peptide include conjugation to carriers, or other techniques are well known in the art.

An immunogenic portion of APRIL or its receptor can be administered in the presence of an adjuvant. The progress of immunization can be monitored by detection 20 of antibody titers in plasma or serum. Standard ELISA or other immunoassays can be used with the immunogen as antigen to assess the levels of antibodies.

In a preferred embodiment, the subject antibodies are immunospecific for antigenic determinants of APRIL, or its receptor, e.g. antigenic determinants of a polypeptide of SEQ. ID. NO.: 2, or a closely related human or non-human mammalian 25 homolog (e.g. 70, 80 or 90 percent homologous, more preferably at least 95 percent homologous). In yet a further preferred embodiment of the present invention, the anti-APRIL or anti-APRIL-receptor antibodies do not substantially cross react (i.e. react specifically) with a protein which is e.g., less than 80 percent homologous to SEQ. ID. NO. 2; preferably less than 90 percent homologous with SEQ. ID. NO.: 2; and, most 30 preferably less than 95 percent homologous with SEQ. ID. NO.: 2. By "not substantially cross react", it is meant that the antibody has a binding affinity for a non-homologous protein which is less than 10 percent, more preferably less than 5 percent, and even more preferably less than 1 percent, of the binding affinity for a protein of SEQ. ID. NO. 2.

The term antibody as used herein is intended to include fragments thereof which are also specifically reactive with APRIL, or its receptor. Antibodies can be fragmented using conventional techniques and the fragments screened for utility in the same manner as described above for whole antibodies. For example, $F(ab')_2$ fragments 5 can be generated by treating antibody with pepsin. The resulting $F(ab')_2$ fragment can be treated to reduce disulfide bridges to produce Fab' fragments. The antibodies of the present invention are further intended to include biospecific and chimeric molecules having anti-APRIL or anti-APRIL -receptor activity. Thus, both monoclonal and polyclonal antibodies (Ab) directed against APRIL and its receptor, and antibody 10 fragments such as Fab' and $F(ab')_2$, can be used to block the action of APRIL and its respective receptor.

Various forms of antibodies can also be made using standard recombinant DNA techniques. (Winter and Milstein, *Nature* 349: 293-299 (1991) specifically incorporated by reference herein.) For example, chimeric antibodies can be constructed 15 in which the antigen binding domain from an animal antibody is linked to a human constant domain (e.g. Cabilly et al., U.S. 4,816,567, incorporated herein by reference). Chimeric antibodies may reduce the observed immunogenic responses elicited by animal antibodies when used in human clinical treatments.

In addition, recombinant "humanized antibodies" which recognize APRIL, or 20 its receptor can be synthesized. Humanized antibodies are chimeras comprising mostly human IgG sequences into which the regions responsible for specific antigen-binding have been inserted. Animals are immunized with the desired antigen, the corresponding antibodies are isolated, and the portion of the variable region sequences 25 responsible for specific antigen binding are removed. The animal-derived antigen binding regions are then cloned into the appropriate position of human antibody genes in which the antigen binding regions have been deleted. Humanized antibodies minimize the use of heterologous (i.e. inter species) sequences in human antibodies, and thus are less likely to elicit immune responses in the treated subject.

Construction of different classes of recombinant antibodies can also be 30 accomplished by making chimeric or humanized antibodies comprising variable domains and human constant domains (CH1, CH2, CH3) isolated from different classes of immunoglobulins. For example, antibodies with increased antigen binding site valencies can be recombinantly produced by cloning the antigen binding site into

vectors carrying the human chain constant regions. (Arulanandam et al., J. Exp. Med., 177: 1439-1450 (1993), incorporated herein by reference.)

In addition, standard recombinant DNA techniques can be used to alter the binding affinities of recombinant antibodies with their antigens by altering amino acid residues in the vicinity of the antigen binding sites. The antigen binding affinity of a humanized antibody can be increased by mutagenesis based on molecular modeling. (Queen et al., Proc. Natl. Acad. Sci. 86: 10029-33 (1989) incorporated herein by reference.

F. Generation of Analogs: Production of Altered DNA and Peptide Sequences

Analogs of APRIL can differ from the naturally occurring Ligands in amino acid sequence, or in ways that do not involve sequence, or both. Non-sequence modifications include in vivo or in vitro chemical derivatization of APRIL. Non-sequence modifications include, but are not limited to, changes in acetylation, methylation, phosphorylation, carboxylation or glycosylation.

Preferred analogs include, APRIL or biologically active fragments thereof, whose sequences differ from the sequence given in SEQ. ID NO. 2, by one or more conservative amino acid substitutions, or by one or more non-conservative amino acid substitutions, deletions or insertions which do not abolish the biological activity of APRIL. Conservative substitutions typically include the substitution of one amino acid for another with similar characteristics, e.g. substitutions within the following groups: valine, glycine; glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and, phenylalanine, tyrosine.

-19-

TABLE I
CONSERVATIVE AMINO ACID REPLACEMENTS

for amino Acid	code	replace with any of:
Alanine	A	D-Ala, Gly, Beta-Ala, L-Cys, D-Cys
Arginine	R	D-Arg, Lys, D-Lys, homo-Arg, D-homo-Arg, Met, Ile, D-Met, D-Ile, Orn, D-Orn
Asparagine	N	D-Asn, Asp, D-Asp, Glu, D-Glu, Gln, D-Gln
Aspartic Acid	D	D-Asp, D-Asn, Asn, Glu, D-Glu, Gln, D-Gln
Cysteine	C	D-Cys, S-Me-Cys, Met, D-Met, Thr, D-Thr
Glutamine	Q	D-Gln, Asn, D-Asn, Glu, D-Glu, Asp, D-Asp
Glutamic Acid	E	D-Glu, D-Asp, Asp, Asn, D-Asn, Gln, D-Gln
Glycine	G	Ala, D-Ala, Pro, D-Pro, -Ala, Acp
Isoleucine	I	D-Ile, Val, D-Val, Leu, D-Leu, Met, D-Met
Leucine	L	D-Leu, Val, D-Val, Leu, D-Leu, Met, D-Met

Lysine	K	D-Lys, Arg, D-Arg, Homo-arg, D-homo-Arg, Met, D-Met, Ile, D-Ile, Orn, D-Orn
Methionine	M	D-Met, S-Me-Cys, Ile, D- Ile, Leu, D-Leu, Val, D- Val
Phenylalanine	F	D-Phe, Tyr, D-Thr, L- Dopa, His, D-His, Trp, D- Trp, Trans-3, 4 or 5- phenylproline, cis-3, 4, or 5-phenylproline
Proline	P	D-Pro, L-L-thiazolidine-4- carboxylic acid, D-or L- oxazolidine-4-carboxylic acid
Serine	S	D-Ser, Thr, D-Thr, allo- Thr, Met, D-Met, Met(O), D-Met(O), L-Cys, D-Cys
Threonine	T	D-Thr, Ser, D-Ser, allo- Thr, Met, D-Met, Met(O), D-Met(O), Val, D-Val
Tyrosine	Y	D-Tyr, Phe, D-Phe, L- Dopa, His, D-His
Valine	V	D-Val, Leu, D-Leu, Ile, D- Ile, Met, D-Met

Useful methods for mutagenesis include PCR mutagenesis and saturation mutagenesis as discussed in more detail below. A library of random amino acid

sequence variants can also be generated by the synthesis of a set of degenerate oligonucleotide sequences.

-PCR Mutagenesis

In PCR mutagenesis, reduced Taq polymerase fidelity can be used to introduce 5 random mutations into a cloned fragment of DNA (Leung et al., 1989, *Technique* 1:11-15). This is a very powerful and relatively rapid method of introducing random mutations. The DNA region to be mutagenized can be amplified using the polymerase chain reaction (PCR) under conditions that reduce the fidelity of DNA synthesis by Taq 10 DNA polymerase, e.g., by using a dGTP/dATP ratio of five and adding Mn²⁺ to the PCR reaction. The pool of amplified DNA fragments can be inserted into appropriate cloning vectors to provide random mutant libraries.

-Saturation Mutagenesis

Saturation mutagenesis allows for the rapid introduction of a large number of 15 single base substitutions into cloned DNA fragments (Mayers et al., 1985, *Science* 229:242). This technique includes generation of mutations, e.g., by chemical treatment or irradiation of single-stranded DNA *in vitro*, and synthesis of a complimentary DNA strand. The mutation frequency can be modulated by modulating the severity of the treatment, and essentially all possible base substitutions can be obtained. Because this 20 procedure does not involve a genetic selection for mutant fragments both neutral substitutions, as well as of a protein can be prepared by random mutagenesis of DNA which those that alter function, can be obtained. The distribution of point mutations is not biased toward conserved sequence elements.

-Degenerate Oligonucleotides

A library of homologs can also be generated from a set of degenerate 25 oligonucleotide sequences. Chemical synthesis of degenerate sequences can be carried out in an automatic DNA synthesizer, and the synthetic genes then ligated into an appropriate expression vector. The synthesis of degenerate oligonucleotides is known in the art^{xxx}. Such techniques have been employed in the directed evolution of other proteins^{xxxi}.

30 Non-random or directed, mutagenesis techniques can be used to provide specific sequences or mutations in specific regions. These techniques can be used to create variants which include, e.g., deletions, insertions, or substitutions, of residues of the known amino acid sequence of a protein. The sites for mutation can be modified individually or in series, e.g., by (1) substituting first with conserved amino acids and

then with more radical choices depending upon results achieved, (2) deleting the target residue, or (3) inserting residues of the same or a different class adjacent to the located site, or combinations of options 1-3.

-Alanine Scanning Mutagenesis

5 Alanine scanning mutagenesis is a useful method for identification of certain residues or regions of the desired protein that are preferred locations or domains for mutagenesis, Cunningham and Wells (*Science* 244:1081-1085, 1989) specifically incorporated by reference. In alanine scanning, a residue or group of target residues are identified (e.g., charged residues such as Arg, Asp, His, Lys, and Glu) and replaced by 10 a neutral or negatively charged amino acid (most preferably alanine or polyalanine). Replacement of an amino acid can affect the interaction of the amino acids with the surrounding aqueous environment in or outside the cell. Those domains demonstrating functional sensitivity to the substitutions can then be refined by introducing further or other variants at or for the sites of substitution. Thus, while the site for introducing an 15 amino acid sequence variation is predetermined, the nature of the mutation per se need not be predetermined. For example, to optimize the performance of a mutation at a given site, alanine scanning or random mutagenesis may be conducted at the target codon or region and the expressed desired protein subunit variants are screened for the optimal combination of desired activity.

20 -Oligonucleotide-Mediated Mutagenesis

 Oligonucleotide-mediated mutagenesis is a useful method for preparing substitution, deletion, and insertion variants of DNA, see, e.g., Adelman et al., (*DNA* 2:183, 1983) incorporated herein by reference. Briefly, the desired DNA can be altered by hybridizing an oligonucleotide encoding a mutation to a DNA template, 25 where the template is the single-stranded form of a plasmid or bacteriophage containing the unaltered or native DNA sequence of the desired protein. After hybridization, a DNA polymerase is used to synthesize an entire second complementary strand of the template that will thus incorporate the oligonucleotide primer, and will code for the selected alteration in the desired protein DNA. Generally, oligonucleotides of at least 30 25 nucleotides in length are used. An optimal oligonucleotide will have 12 to 15 nucleotides that are completely complementary to the template on either side of the nucleotide(s) coding for the mutation. This ensures that the oligonucleotide will hybridize properly to the single-stranded DNA template molecule. The oligonucleotides are readily synthesized using techniques known in the art such as that

described by Crea et al. (*Proc. Natl. Acad. Sci. USA*, 75: 5765[1978]) incorporated herein by reference.

-Cassette Mutagenesis

Another method for preparing variants, cassette mutagenesis, is based on the 5 technique described by Wells et al. (*Gene*, 34:315[1985]) incorporated herein by reference. The starting material can be a plasmid (or other vector) which includes the protein subunit DNA to be mutated. The codon(s) in the protein subunit DNA to be mutated are identified. There must be a unique restriction endonuclease site on each side of the identified mutation site(s). If no such restriction sites exist, they may be 10 generated using the above-described oligonucleotide-mediated mutagenesis method to introduce them at appropriate locations in the desired protein subunit DNA. After the restriction sites have been introduced into the plasmid, the plasmid is cut at these sites to linearize it. A double-stranded oligonucleotide encoding the sequence of the DNA between the restriction sites but containing the desired mutation(s) is synthesized using 15 standard procedures. The two strands are synthesized separately and then hybridized together using standard techniques. This double-stranded oligonucleotide is referred to as the cassette. This cassette is designed to have 3' and 5' ends that are comparable with the ends of the linearized plasmid, such that it can be directly ligated to the plasmid. This plasmid now contains the mutated desired protein subunit DNA 20 sequence.

-Combinatorial Mutagenesis

Combinatorial mutagenesis can also be used to generate mutants. E.g., the amino acid sequences for a group of homologs or other related proteins are aligned, 25 preferably to promote the highest homology possible. All of the amino acids which appear at a given position of the aligned sequences can be selected to create a degenerate set of combinatorial sequences. The variegated library of variants is generated by combinatorial mutagenesis at the nucleic acid level, and is encoded by a variegated gene library. For example, a mixture of synthetic oligonucleotides can be enzymatically ligated into gene sequences such that the degenerate set of potential 30 sequences are expressible as individual peptides, or alternatively, as a set of larger fusion proteins containing the set of degenerate sequences.

Various techniques are known in the art for screening generated mutant gene products. Techniques for screening large gene libraries often include cloning the gene library into replicable expression vectors, transforming appropriate cells with the

resulting library of vectors, and expressing the genes under conditions in which detection of a desired activity, e.g., in this case, binding to APRIL or its receptor, facilitates relatively easy isolation of the vector encoding the gene whose product was detected. Each of the techniques described below is amenable to high through-put analysis for screening large numbers of sequences created, e.g., by random mutagenesis techniques.

5 The invention also provides for reduction of the protein binding domains of the claimed polypeptides or their receptors, to generate mimetics, e.g. peptide or non-peptide agents. The peptide mimetics are able to disrupt binding of APRIL with its
10 respective receptor. The critical residues of the APRIL involved in molecular recognition of a receptor polypeptide or of a downstream intracellular protein, can be determined and used to generate APRIL or its receptor-derived peptidomimetics which competitively or noncompetitively inhibit binding of APRIL with a receptor. (see, for
15 example, "Peptide inhibitors of human papilloma virus protein binding to retinoblastoma gene protein" European patent applications EP-412,762A and EP-B31,080A), specifically incorporated herein by reference.

20 By making available purified and recombinant APRIL, the present invention provides assays which can be used to screen for drug candidates which are either agonists or antagonists of the normal cellular function, in this case, of APRIL or its receptor. In one embodiment, the assay evaluates the ability of a compound to modulate binding between APRIL and a receptor. A variety of assay formats will suffice and, in light of the present inventions, will be comprehended by the skilled artisan.

25 In many drug screening programs which test libraries of compounds and natural extracts, high throughput assays are desirable in order to maximize the number of compounds surveyed in a given period of time. Assays which are performed in cell-free systems, such as may be derived with purified or semi-purified proteins, are often preferred as "primary" screens in that they can be generated to permit rapid development and relatively easy detection of an alteration in a molecular target which
30 is mediated by a test compound. Moreover, the effects of cellular toxicity and/or bioavailability of the test compound can be generally ignored in the *in vitro* system, the assay instead being focused primarily on the effect of the drug on the molecular target as may be manifest in an alteration of binding affinity with other proteins or change in enzymatic properties of the molecular target.

Isolation of a receptor binding to APRIL

Ligands of the TNF family can be used to identify and clone receptors. With the described APRIL sequences, one could fuse the 5' end of the extracellular domain which constitutes the receptor binding sequence to a marker or tagging sequence and then add a leader sequence that will force secretion of APRIL in any of a number of expression systems. One example of this technology is described by Browning et al., (1996) (JBC 271, 8618-8626) where the LT- β ligand was secreted in such a form. The VCAM leader sequence was coupled to a short myc peptide tag followed by the extracellular domain of the LT- β . The VCAM sequence is used to force secretion of the normally membrane bound LT- β molecule. The secreted protein retains a myc tag on the N-terminus which does not impair the ability to bind to a receptor. Such a secreted protein can be expressed in either transiently transfected Cos cells or a similar system, e.g., EBNA derived vectors, insect cell/baculovirus, picchia etc. The unpurified cell supernatant can be used as a source of the tagged ligand.

Cells expressing the receptor can be identified by exposing them to the tagged ligand. Cells with bound ligand are identified in a FACS experiment by labeling the myc tag with an anti-myc peptide antibody (9E10) followed by phycoerythrin (or a similar label) labeled anti-mouse immunoglobulin. FACS positive cells can be readily identified and would serve as a source of RNA encoding for the receptor. An expression library would then be prepared from this RNA via standard techniques and separated into pools. Pools of clones would be transfected into a suitable host cell and binding of the tagged ligand to receptor positive transfected cells determined via microscopic examination, following labeling of bound myc peptide tag with an enzyme labeled anti-mouse Ig reagent, i.e. galactosidase, alkaline phosphatase or luciferase labeled antibody. Once a positive pool has been identified, the pool size would be reduced until the receptor encoding cDNA is identified. This procedure could be carried out with either the mouse or human APRIL, as one may more readily lead to a receptor.

G. METHODS OF TREATMENT AND PHARMACEUTICAL COMPOSITIONS

The methods of the invention for the treatment of cancers involve the administration to a patient, preferably a mammalian host, such as a dog, cat, or human, an effective amount of a claimed composition comprising a blocking agent capable of interfering with the association between APRIL and its receptor. Such blocking agents 5 include, but are not limited to soluble APRIL, anti-APRIL antibodies, anti-APRIL receptor antibodies, or biologically active fragments thereof. Additionally, an inhibitory form of APRIL can be made by mutating APRIL, while maintaining the ability to block the association between APRIL and its receptor. Blocking agents may preferably comprise a receptor IG fusion protein, which can be constructed by methods 10 known to those of skill in the art.

The methods of the invention are useful for treating all cancers, including, but not limited to, cellular disorders as, for example, renal cell cancer, Kaposi's sarcoma, chronic leukemia, breast cancer, sarcoma, ovarian carcinoma, rectal cancer, throat cancer, melanoma, colon cancer, bladder cancer, mastocytoma, lung 15 cancer, mammary adenocarcinoma, pharyngeal squamous cell carcinoma, and gastrointestinal or stomach cancer. Additionally, such blocking agents are useful for the treatment of proliferative conditions that are not considered to be tumors, i.e. cellular hyperproliferation (hyperplasia), such as, for example, scleroderma, pannus formation in rheumatoid arthritis, postsurgical scarring and lung, liver and uterine 20 fibrosis.

Pharmaceutical compositions of the invention may comprise a therapeutically effective amount of APRIL, or its receptor, or fragments or mimetics thereof, and, optionally may include pharmaceutically acceptable carriers. Accordingly, this invention provides methods for treatment of cancer, and methods of 25 stimulating, or in certain instances, inhibiting the immune system, or parts thereof by administering a pharmaceutically effective amount of a compound of the invention or its pharmaceutically acceptable salts or derivatives. It should of course be understood that the compositions and methods of this invention can be used in combination with other therapies for various treatments.

30 The compositions can be formulated for a variety of routes of administration, including systemic, topical or localized administration. For systemic administration, injection is preferred, including intramuscular, intravenous, intraperitoneal, and subcutaneous for injection, the compositions of the invention can be formulated in

liquid solutions, preferably in physiologically compatible buffers such as Hank's solution or Ringer's solution. In addition, the compositions may be formulated in solid form and, optionally, redissolved or suspended immediately prior to use. Lyophilized forms are also included in the invention.

5 The compositions can be administered orally, or by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are known in the art, and include, for example, for transmucosal administration, bile salts, fusidic acid derivatives, and detergents. Transmucosal administration may be through nasal sprays
10 or using suppositories. For oral administration, the compositions are formulated into conventional oral administration forms such as capsules, tablets, and tonics. For topical administration, the compositions of the invention are formulated into ointments, salves, gels, or creams as known in the art.

15 The dose and dosing regimen will depend on the type of cancer, the patient and the patient's history. The amount must be effective to treat, suppress, or alter the progression of cancer. The doses may be single doses or multiple doses. If multiple doses are employed, as preferred, the frequency of administration will depend, for example, on the type of host and and type of cancer, dosage amounts etc. For some types of cancers or cancer lines, daily administration will be effective, whereas for others, 20 administration every other day or every third day will be effective. The amount of active compound administered at one time or over the course of treatment will depend on many factors. For example, the age and size of the subject, the severity and course of the disease being treated, the manner and form of administration, and the judgments of the treating physician. However, an effective dose may be in the range of from 25 about 0.005 to about 5 mg/kg/day, preferably about 0.05 to about 0.5 mg/kg/day. The dosage amount which will be most effective will be one which results in no tumor appearance or complete regression of the tumor, and is not toxic to the patient. One skilled in the art will recognize that lower and higher doses may also be useful.

30 Gene constructs according to the invention can also be used as a part of a gene therapy protocol to deliver nucleic acids encoding either an agonistic or antagonistic form of APRIL.

Expression constructs of the APRIL can be administered in any biologically effective carrier, e.g., any formulation or composition capable of effectively delivering the gene for APRIL to cells in vivo. Approaches include insertion of the gene in viral

vectors which can transfect cells directly, or delivering plasmid DNA with the help of, for example, liposomes, or intracellular carriers, as well as direct injection of the gene construct. Viral vector transfer methods are preferred.

A pharmaceutical preparation of the gene therapy construct can consist 5 essentially of the gene delivery system in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery system can be produced intact from recombinant cells, e.g. retroviral vectors, the pharmaceutical preparation can comprise one or more cells which produce the gene delivery system.

10 In addition to use in therapy, the oligomers of the invention may be used as diagnostic reagents to detect the presence or absence of the target DNA, RNA or amino acid sequences to which they specifically bind. In other aspects, the claimed invention may be used to evaluate a chemical entity for its ability to interact with, e.g., bind or physically associate with APRIL or a fragment thereof. The method includes 15 contacting the chemical entity with APRIL, and evaluating the ability of the entity to interact with APRIL. Additionally, APRIL can be used in methods of evaluating naturally occurring APRIL or receptors of APRIL, as well as to evaluate chemical entities which associate or bind with receptors of APRIL. It may be desirable to use tagged versions of APRIL to facilitate the detection of APRIL binding to its receptor, 20 or receptor positive cells, such as, for example, the purpose of screening for agents that block the APRIL ligand-APRIL receptor interaction. Additionally, one may use APRIL transfected cell lines that have increased growth rates as the basis for screening assays for molecules that block APRIL activity.

25 In certain aspects, the claimed invention features a method for evaluating a chemical entity for the ability to modulate the interaction between APRIL and its respective receptor. The method includes combining a receptor for APRIL, and APRIL under conditions wherein the pair is capable of interacting, adding the chemical entity to be evaluated and detecting the formation or dissolution of complexes. These 30 modulating agents may be further evaluated in vitro, e.g. by testing its activity in a cell free system, and then, optionally administering the compound to a cell or animal, and evaluating the effect.

I. EXAMPLES

Example 1

Northern blot analysis of APRIL revealed that the expression of APRIL was weak and restricted only to a few tissues (Fig. 2A). Two transcripts of 2.1 kb and 2.4 kb were found in the prostate, whereas PBLs revealed a shorter transcript of 1.8 kb.

5 Northern blot analysis was performed by using Human Multiple Tissue Northern Blots I and II (Clontech #7760-1 and #7759-1), Human Cancer Cell Line MTN Blot (Clontech #7757-1) and Human Tumor Panel Blot V (Invitrogen D3500-01). The membranes were incubated in ExpressHyb hybridization solution (Clontech #8015-1)

10 for at least 1 hour at 62°C. The random-primed cDNA probe (Boehringer Mannheim) was synthesized using cDNA corresponding to the extracellular domain of APRIL as template. The heat-denatured cDNA probe was added at 1.5×10^6 cpm/ml in fresh ExpressHyb. The membrane was hybridized 12-24 hr at 62°C, washed three times in 2xSSC containing 0.05% SDS and exposed at -70°C. Northern blot analysis of APRIL 15 revealed that the expression of APRIL was weak and restricted only to a few tissues.

Two transcripts of 2.1 kb and 2.4 kb were found in the prostate, whereas PBLs revealed a shorter transcript of 1.8 kb.

A longer exposure time revealed the 2.1 kb APRIL mRNA in colon, spleen, and pancreas (data not shown). This restricted distribution of the APRIL mRNA is 20 consistent with the origin of cDNA clones currently available in the EST database. Of the 23 clones identified only two were derived from normal tissues (pregnant uterus, pancreatic islands). Remarkably, the remainder of the EST-clones (21 clones, 91%) were present in cDNA libraries generated from tumors or tumor-derived cell lines (Ovary tumor, 11; prostate tumor, 3; Gessler Wilms tumor, 1; colon carcinoma, 1; 25 endometrial tumor, 1; parathyroid tumors, 1; pancreas tumor, 1; T-cell lymphoma, 1; LNCAP adenocarcinoma derived cell line, 1). This prompted us to test transformed cell lines for the expression of APRIL mRNA (Figure 2B), and indeed, all cell lines strongly expressed the 2.1 kb transcript of APRIL.

Highest APRIL-specific signals were detected in the colorectal 30 adenocarcinoma SW480, the Burkitt's lymphoma Raji and in the melanoma G361. To corroborate this finding, we measured APRIL mRNA expression levels in several tumors and compared them to normal tissues. APRIL mRNA was abundantly detected in thyroid carcinoma and in lymphoma, whereas in the corresponding normal tissues,

only weak or no hybridization signals were found (Fig. 2C). In the two other tumors analyzed by Northern blots (adrenal and paratoid tumors), APRIL mRNA was not elevated. However, *in situ* hybridization revealed abundant APRIL message in human colon adenocarcinoma as compared to normal colon tissue (Fig. 2D).

5 In order to explore possible activities of APRIL, we expressed a recombinant form of soluble extracellular domain of APRIL (sAPRIL) encompassing amino acids 110 to 250 in 293 cells (9). The full length APRIL gene was amplified from the EST-clone, using a specific 5' forward primer flanked by a EcoRI site (5'-CCAGCCTCATCTCCTTCTTGC-3') and a specific 3' reverse primer flanked by an 10 XbaI site (5'-TCACAGTTCACAAACCCCAGG-3'). The amplified fragment was cut with EcoRI/XbaI and cloned into a modified version of pCRIII (Invitrogen), in frame with an N-terminal Flag peptide (15). The soluble form of APRIL (sAPRIL) was generated using the two primers (5'-AACACAGAAGAAGCAGCACTCTG-3') and (5'-TCACAGTTCACAAACCCCAGG-3') containing a PstI and XbaI site, respectively, 15 and subsequently cloned into a modified pCRIII vector, containing both a HA signal for protein secretion in eukaryotic cells and an N-terminal Flag epitope (15).

Example 2

20 The widespread expression of APRIL in tumor cells and tissues suggested to us that APRIL may be associated with tumor growth, and we therefore incubated various tumor cell lines with purified recombinant Flag-tagged sAPRIL (10).

Human embryonic 293T cells, human leukemia Jurkat T-cells, human Burkitt lymphoma B-cells Raji and melanoma cell lines were grown as previously described (16, 17). Other cell lines referred in this paper are deposited in and described by the 25 American Type Culture Collection (Rockville, Maryland). All cell lines were cultured in RPMI or DMEM medium supplemented with 10% fetal calf serum.

Flag-tagged versions of the extracellular domain (residues 103-281) of human FasL and TRAIL (residues 95-281) were recently described (15). Flag-tagged soluble human TWEAK (residues 141-284) was produced in 293 cells (P. S. manuscript in preparation). The anti-Flag antibody M2 were obtained from Kodak International Biotechnologies. An increase in proliferation of the Jurkat T lymphoma cells in the presence of APRIL was observed in a dose dependent manner as detected by an increase in number (approximately 50%) (11) of viable cells 24 hrs after ligand

addition (Fig. 3A). The proliferation of cells was determined by incubating cells at 50,000 cells per well in 100 μ l medium with the indicated concentrations of recombinant APRIL, TWEAK, TRAIL, FasL and by determining the number of viable cells using the Celltiter 96 AQ proliferation assay (Promega) after 24 hrs, following the manufacturer's instructions, or by 3 H-thymidine incorporation. For the immunodepletion of Flag-APRIL, anti-Flag coupled to agarose was used.

The increase in proliferation was independent of a co-stimulatory signals such as anti-CD3 antibodies or other cytokines. As expected, the addition of identically produced and purified FasL to Jurkat cells decreased the number of viable cells, whereas TWEAK had no effect. The increased cell number correlated with augmented (40%) 3 H-thymidine incorporation in APRIL-treated cells (Fig. 3A). Immunodepletion of FLAG-tagged APRIL-containing conditioned medium by anti-FLAG antibodies, but not anti-myc antibodies, reduced the proliferative effect (Fig. 3B), indicating that the proliferative effect was specific and due to APRIL. Increased proliferation rates were also seen in some B lymphomas (human Raji, mouse A20 cells, but not human BJAB) and on cell lines of epithelial origin such as COS and HeLa, as well as melanomas (Fig. 3C). The breast carcinoma cell MCF-7 did not respond. The effect on Jurkat cells was even more pronounced when the fetal calf serum was reduced from 10% to 1% (Fig. 3D).

Recombinant sAPRIL has forms aggregates which may explain the rather high concentrations needed to detect a proliferative effect with sAPRIL. We therefore transfected NIH-3T3 cells with full-length human APRIL (12) and obtained several APRIL-expressing clones (Fig. 4A). NIH-3T3 APRIL clones were established using the calcium phosphate method of transfection and the full-length FLAG-tagged APRIL containing pCRIII expression vector. Cellular proteins of about 2×10^6 cells per lane were electrophoretically separated on a 12% polyacrylamide gel in the presence of SDS under reducing conditions and subsequently transferred to nitrocellulose. Immunoblot analysis of Flag-tagged APRIL was conducted using 5 μ g/ml of the rat monoclonal anti-Flag antibody M2 (Kodak International Biotechnologies). First antibodies were detected using affinity purified anti-peroxidase-conjugated donkey anti-mouse antibody (Dianova, Hamburg, Germany) followed by a chemiluminescence reaction using the ECL system (Amersham).

Interestingly, APRIL transfectants proliferated faster than mock-transfectants (Fig. 4B). We reasoned that the APRIL-transfected NIH-3T3 cells might also have a

growth advantage *in vivo*. When wild-type or mock-transfected NIH-3T3 cells were injected into nude mice, small palpable tumors were observed after 5-6 weeks (13). In contrast, two clones of NIH-3T3 cells stably transfected with APRIL both induced tumors after only 3-4 weeks. After 6 weeks, mice had to be killed due to the high 5 tumor burden (Fig. 4C). NIH/3T3 fibroblasts (American Type Culture Collection, Rockville, Maryland) and the various transfectants (1×10^5 cells) were suspended in 50 μ l PBS and injected subcutaneously into the flank region of BALB/c nude mice (Harlan, Zeist, Netherland). Tumor size was measured every three days. Mice were age-matched (3 animals per group).

10

Example 3:

Isolation of a receptor binding to APRIL.

Ligands of the TNF family can be used to identify and clone receptors. With the described APRIL sequences, one could fuse the 5' end of the extracellular domain 15 of APRIL which constitutes the receptor binding sequence to a marker or tagging sequence and then add a leader sequence that will force secretion of APRIL in any of a number of expression systems. One example of this technology is described by Browning et al., (1996) (JBC 271, 8618-8626) where the LT- β ligand was secreted in such a form. The VCAM leader sequence was coupled to a short myc peptide tag 20 followed by the extracellular domain of the LT- β . The VCAM sequence is used to force secretion of the normally membrane bound LT- β molecule. The secreted protein retains a myc tag on the N-terminus which does not impair the ability to bind to a receptor. Such a secreted protein can be expressed in either transiently transfected Cos 25 cells or a similar system, e.g., EBNA derived vectors, insect cell/baculovirus, picchia etc. The unpurified cell supernatant can be used as a source of the tagged ligand.

Cells expressing the receptor can be identified by exposing them to the tagged ligand. Cells with bound ligand are identified in a FACS experiment by labeling the myc tag with an anti-myc peptide antibody (9E10) followed by phycoerythrin (or a similar label) labeled anti-mouse immunoglobulin. FACS positive cells can be readily 30 identified and would serve as a source of RNA encoding for the receptor. An expression library would then be prepared from this RNA via standard techniques and separated into pools. Pools of clones would be transfected into a suitable host cell and binding of the tagged ligand to receptor positive transfected cells determined via

microscopic examination, following labeling of bound myc peptide tag with an enzyme labeled anti-mouse Ig reagent, i.e. galactosidase, alkaline phosphatase or luciferase labeled antibody. Once a positive pool has been identified, the pool size would be reduced until the receptor encoding cDNA is identified. This procedure could be
5 carried out with either the mouse or human APRIL, as one may more readily lead to a receptor.

It will be apparent to those skilled in the art that various modifications and variations can be made in APRIL, compositions and methods of the present invention
10 without departing from the spirit or scope of the invention. Thus, it is intended that the present invention cover the modifications and variations of this invention provided that they come within the scope of the appended claims and their equivalents.

EDITORIAL NOTE - NO 93162/98

Sequence listing pages 1-2 is part of the description.

The claims are to follow.

SEQ ID NO: 1

1 GGTACGAGGC TTCCCTAGAGG GACTGGAACC TAATTCTCCT GAGGCTGAGG
 5 51 GAGGGTGGAG GGTCTCAAGG CAACGCTGGC CCCACGACGG AGTGCCAGGA
 101 101 GCACTAACAG TACCCCTAGC TTGCTTCCT CCTCCCTCCT TTTTATTTC
 151 151 AAGTTCCCTT TTATTTCTCC TTGCGTAACA ACCTTCTTCC CTTCTGCACC
 201 201 ACTGCCCGTA CCCTTACCCG CCCCGCCACC TCCTTGCTAC CCCACTCTG
 251 251 AAACCACAGC TGTGGCAGG GTCCCCAGCT CATGCCAGCC TCATCTCCTT
 301 301 TCTTGCTAGC CCCAAAGGG CCTCCAGGCA ACATGGGGGG CCCAGTCAGA
 351 351 GAGCCGGCAC TCTCAGTTGC CCTCTGGTTG AGTTGGGGGG CAGCTCTGGG
 401 401 GGCCGTGGCT TGTGCCATGG CTCTGCTGAC CCAACAAACA GAGCTGCAGA
 451 451 GCCTCAGGAG AGAGGTGAGC CGGCTGCAGG GGACAGGAGG CCCTCCCGAG
 501 501 AATGGGGAAG GGTATCCCTG GCAGAGTCTC CCGGAGCAGA GTTCCGATGC
 551 551 CCTGGAAAGCC TGGGAGAATG GGGAGAGATC CCGGAAAAGG GAGCAGTGC
 601 601 TCACCCAAAA ACAGAAGAAG CAGCACTCTG TCCTGCACCT GGTTCCCATT
 651 651 AACGCCACCT CCAAGGATGA CTCCGATGTG ACAGAGGTGA TGTGGCAACC
 701 701 AGCTCTTAGG CGTGGGAGAG GCCTACAGGC CCAAGGATAT GGTGTCCGAA
 751 751 TCCAGGATGC TGGAGTTAT CTGCTGTATA GCCAGGTCTT GTTCAAGAC
 801 801 GTGACTTTCA CCATGGGTCA GGTGGTGTCT CGAGAAGGCC AAGGAAGGCA
 851 851 GGAGACTCTA TTCCGATGTA TAAGAAGTAT GCCCTCCAC CCGGACCGGG
 901 901 CCTACAAACAG CTGCTATAGC GCAGGTGTCT TCCATTACA CCAAGGGGAT
 951 951 ATTCTGAGTG TCATAATTCC CCGGCAAGG GCGAAACTTA ACCTCTCTCC
 1001 1001 ACATGGAAACC TTCTGGGGT TTGTGAAACT GTGATTGTGT TATAAAAAGT
 1051 1051 GGCTCCCAGC TTGGAAGACC AGGGTGGGT AACTGGAGACAGCCAAGA
 1101 1101 GCTGAGTATA TAAAGGAGAG GGAATGTGCA GGAAACAGAGGCATCTTCTG
 1151 1151 GGTTTGGCTC CCCGTTCCCT ACTTTCCCT TTTCATTCCC ACCCCCTAGA
 1201 1201 CTTTGATTTT ACGGATATCT TGCTTCTGTT CCCCATGGAG CTCCGAATT
 1251 1251 TTGCGTGTGT GTAGATGAGG GGCGGGGGAC GGGGCCAGG CATTGTTCA
 1301 1301 ACCTGGTCGG GCCCACTGG AAGCATCCAG AACAGCACCA CCATCTTA

30 SEQ ID NO: 2

1 MPASSPFLLA PKGPPGNMGG PVREPALSVA LWLSWGAALG AVACAMALLT
 5 51 QQTELQLSRR EVSRLQGTGG PSQNGEYPW QSLPEQSSDA LEAWENGERS
 101 101 RKRRAVLTQK QKQHCSVHL VPINATSKDD SDVTEVMWQP ALRRGRGLQA
 151 151 QGYGVRIQDA GVYLLYSQVL FQDVTFTMGQ VVSREGQGRQ ETLFRCIRSM
 201 201 PSHPDRAYNS CYSAGVFHLH QGDILSVIIP RARAKLNLSH HGTFLGFVKL

35 SEQ ID NO: 3

1 1 GAATTCCGCA CGAGGCTCCA GCCCACATGG GGGGCTCAGT CAGAGAGCCA
 40 51 51 GCCCTTTCGG TTGCTCTTG GTTGAGTTGG GGGGCAGTTC TGGGGCTGT
 101 101 GACTTGTGCT GTCGCACTAC TGATCCAACA GACAGAGCTG CAAAGCCTAA
 151 151 GGCAGGAGGT GAGCCGGCTG CAGCGGAGTG GAGGGCCTTC CCAGAACAG
 201 201 GGAGAGCGCC CATGGCAGAG CCTCTGGGAG CAGAGTCCTG ATGTCCTGG
 251 251 AGCCTGGAAG GATGGGGCGA AATCTCGGAG AAGGAGAGCA GTACTCACCC
 301 301 AGAACGACAA GAAGAACAC TCAGTCCTGC ATCTGTTCC AGTTAACATT
 351 351 ACCTCCAAGG ACTCTGACGT GACAGAGGTG ATGTGGCAAC CAGTACTTAG
 401 401 GCGTGGGAGA GGCCCTGGAG GCCCAGGGAG ACATTTGACG AGTCTGGAC
 451 451 ACTGGAATTAT ATCTGCTCTA TAGTCAGGTG CTGTTTCATG ATGTGACTTT
 501 501 CACAATGGGT CAGGTGGTAT CTCGGGAAGG ACAAGGGAGA AGAGAAACCTC
 551 551 TATTCCGATG TATCAGAAAGT ATGCCTTCCTG ATCCTGACCG TGCCTACAA
 601 601 AGCTGCTACA GTGCAGGTGT CTTTCATTAA CATCAAGGGG ATATTATCAC
 651 651 TGTCAAAATT CCACGGGCAA ACGCAAAACT TAGCCTTCT CCGCATGGAA
 701 701 CATTCCCTGGG GTTTGTGAAA CTATGATTGT TATAAAGGGG GTGGGGATTT
 751 751 CCCATTCCAA AAACCTGGCTA GACAAAGGAC AAGGAACGGT CAAGAACAGC
 801 801 TCTCCATGGC TTTGCCTTGA CTGTTGTTCC TCCCTTGCC TTTCCCGCTC
 851 851 CCACTATCTG GGCTTGACT CCATGGATAT TAAAAAAAGTA GAATATTTG
 901 901 TGTTTATCTC CCAAAAAA

SEQ ID NO: 4

5 1 MGGSVREPAL SVALWLSWGA VLGAVTCAVA LLIQQTELQS LRREVSRLQR
 51 SGGPSQKQGE RPWQSLWEQS PDVLEAWKDG AKSRRRRRAVL TQKHKKKHSV
 101 LHLVPVNITS KDSDVTEVMW QPVLRRGRGP GGQGDIVRVW DTGIYLLYSQ
 151 VLFHDVTFTM GQVVSREGQG RRETLFRCIR SMPSPDPDRAY NSCYSAGVFH
201 LHQGDIITVK IPRANAKLSL SPHGTFLGFV KL

The claims defining the invention are as follows:

1. A method of treating, suppressing or altering the progression of a cancer comprising administering to a patient an effective amount of an antibody directed to an APRIL ligand polypeptide capable of interfering with an interaction between said APRIL ligand polypeptide and an APRIL receptor.
5
2. The method of claim 1, wherein said antibody is administered to said patient in combination with a chemotherapeutic agent.
3. The method of claim 1, wherein said antibody is administered to said patient in combination with a radiation therapy.
- 10 4. A method of suppressing growth of a tumour cell that expresses an APRIL ligand polypeptide, comprising the step of exposing said cell to an effective amount of an agent selected from the group consisting of:
 - (a) an antibody that binds specifically to said APRIL ligand polypeptide comprising at least about 102 amino acids or a soluble ligand polypeptide thereof;
 - 15 (b) an antibody that binds specifically to said APRIL ligand polypeptide comprising an amino acid sequence found in amino acids 1-55 of SEQ ID NO:2, or a fragment thereof;
 - (c) an antibody that binds specifically to said APRIL ligand polypeptide comprising an amino acid sequence found in amino acids 157-250 of SEQ ID NO:2, a fragment thereof, or a soluble ligand polypeptide thereof;
 - 20 (d) an antibody that binds specifically to SEQ ID NO:2 or a soluble ligand polypeptide thereof; and
 - (e) an antibody that binds specifically to said APRIL ligand polypeptide and blocks an interaction between said APRIL ligand polypeptide and an APRIL receptor polypeptide.
5. A method of suppressing growth of a tumour cell that expresses an APRIL receptor polypeptide, comprising the step of exposing said cell to an effective amount of an agent selected from the group consisting of:
 - (a) an antibody that binds specifically to an APRIL ligand polypeptide comprising at least about 102 amino acids or a soluble ligand polypeptide thereof;
 - (b) an antibody that binds specifically to an APRIL ligand polypeptide comprising an amino acid sequence found in amino acids 1-55 of SEQ ID NO:2, or a fragment thereof;



(c) an antibody that binds specifically to an APRIL ligand polypeptide comprising an amino acid sequence found in amino acids 157-250 of SEQ ID NO:2, a fragment thereof, or a soluble ligand polypeptide thereof;

(d) an antibody that binds specifically to SEQ ID NO:2 or a soluble ligand polypeptide thereof; and

5 (e) an antibody that binds specifically to an APRIL ligand polypeptide and blocks an interaction between said APRIL ligand polypeptide and said APRIL receptor polypeptide.

6. A method for treating cancer comprising administering a therapeutically effective amount of a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of an antibody directed to an APRIL ligand polypeptide capable of interfering with an interaction between said APRIL ligand polypeptide and an APRIL receptor.

10 7. A method for suppressing growth of a tumour cell that expresses an APRIL ligand polypeptide, comprising the step of exposing said cell to an effective amount of an antibody directed to said APRIL ligand polypeptide capable of interfering with an interaction between said APRIL ligand polypeptide and an APRIL receptor.

20 8. A method for suppressing growth of a tumour cell that expresses an APRIL receptor polypeptide, comprising the step of exposing said cell to an effective amount of an antibody directed to an APRIL ligand polypeptide capable of interfering with an interaction between said APRIL ligand polypeptide and said APRIL receptor polypeptide.

9. A method for identifying an agent capable of suppressing the growth of a cell culture, comprising the steps of:

25 (a) identifying a cell that proliferates in response to the binding of a polypeptide encoded by SEQ ID NO:1 to its cell surface receptor;

(b) growing said cell in said cell culture;

(c) exposing said cell culture comprising said cell to a compound resulting in an exposed cell culture;

(d) concurrently or subsequently exposing said exposed cell culture to a polypeptide encoded by SEQ ID NO:1, or a fragment thereof;

30 (e) comparing the proliferation of said cell within said exposed cell culture to the



29106

2020090

proliferation of a substantially identical cell in a second cell culture that was not exposed to said compound; and

(f) determining whether said compound has suppressed the growth of said cell in said exposed cell culture.

5 10. The method of claim 9, wherein said compound is an analog of SEQ ID NO:2.

11. The method of claim 9, wherein said compound is an antibody directed to the polypeptide encoded by SEQ ID NO:1, or a fragment thereof.

12. The method of claim 9, wherein said cell is a tumour cell.

10 13. A method of treating, suppressing or altering the progression of a cancer, comprising the step of administering a therapeutically effective amount of an APRIL ligand polypeptide or a fragment thereof having substantially the same biological activity as APRIL ligand polypeptide.

15 14. A method of suppressing growth of a tumour cell that expresses an APRIL ligand polypeptide or an APRIL receptor polypeptide comprising the step of administering a therapeutically effective amount of an APRIL ligand polypeptide or a fragment thereof having substantially the same biological activity as APRIL ligand polypeptide.

20 15. A method of treating, suppressing or altering the progression of a cancer in a subject comprising the step of administering a therapeutically effective amount of a blocking agent that interferes with the binding of APRIL ligand polypeptide to its receptor, wherein said blocking agent comprises a modified inhibitory form of APRIL ligand polypeptide, or an anti-APRIL ligand polypeptide antibody or a biologically active fragment thereof.

25 16. A method of suppressing growth of a tumour cell that expresses an APRIL ligand polypeptide or an APRIL receptor polypeptide comprising the step of administering a therapeutically effective amount of a blocking agent that interferes with the binding of APRIL ligand polypeptide to its receptor, wherein said blocking agent comprises a modified inhibitory form of APRIL ligand polypeptide, or an anti-APRIL ligand polypeptide antibody or a biologically active fragment thereof.

17. The method of claim 15 or claim 16, wherein the blocking agent is administered to the subject in combination with at least one chemotherapeutic agent.



SEARCHED

002090

18. The method of claim 15, 16 or 17, wherein the subject is administered radiation therapy.

19. A method of treating a cancer or a tumour relating to APRIL ligand polypeptide in a mammal comprising the steps of:

(a) introducing into a cell of said mammal a therapeutically effective amount of a

5 vector comprising a gene encoding APRIL ligand polypeptide or a fragment thereof having substantially the same biological activity as APRIL ligand polypeptide; and

(b) expressing said gene in said mammalian cell.

20. The method of claim 19, wherein said mammal is a human.

21. The method of claim 19 or claim 20, wherein said vector is a virus.

10 22. A method of suppressing growth of a tumour cell that expresses an APRIL ligand polypeptide or an APRIL receptor polypeptide comprising the step of administering a therapeutically effective amount of a soluble form of APRIL ligand polypeptide, wherein said cell is contacted with said soluble polypeptide.

Dated this 4th day of February 2003

15

APOTECH S.A.

By its Patent Attorneys

CULLEN & CO



A

TM

MPASSPFL LAPKGGPNMGPVREPALS ~~IAWISAGA/CAVACAMALI~~ T 50

QQTELQSLRREVSRLQGTGGPSQN GEGYPWQLPEQSSDALEAWENGERS 100

RKRRAVL TQKQKQHSVLHLVPINATSKDDSDVTEVMQPALRRGRGLQA 150

QGYGVRIQDAGVYLLYSQVLFQDVTFIMQVVSREGQGRQETLFR CIRSM 200

PSHPDRAYNSCYSAGVFHLHQGDILSVIIPRARA KLNLSPHGTFLGFVKL 250

B

APRIL	117	VLHLVPINATSKDD DVTEVMQPA . LRRGRGLQA QGYGVRIQDAGVYLLYSQVLFQDVTF FTMGQV
TNFa	89	VAHVVAAPQAEQG LQVLNR RANALLANGVELRDNQLVVPSEGGLYLLYSQVLFRGQGCP THVLLTHTI
LT. a	64	AAHLIGDPSKQNS LLWRA TDRAFLQDGFSLSNNSSLVPTSGIYFVYSQVVFSGKAYSPKATSSPLYLAHEV
FASL	146	VAHLTOKSNSRSMP LEWEDT YGIVLLSGVKY . KKGCLVINETGLYFVYSKQVYFRGQSCN NLPLSHKV
TRAIL	123	AAHITGTRGRSNTLSSPNSKNEKALGRKINSSRESS . SHSFLSNLHL . RNCGLVIIHERGFYYIYSQTYFRFQEEIK ENTKNDKOM
TWEAK	143	AAHYEVHPRPGQDGAQAGV DGTVSGWEKARINSSSPRYNR . QIGEFIVTRAGLYLYCQVHFDEGKAVY LKLDLLVDGVL
TRANCE	164	FAHTINATDIPSGSH KVSLSSVYHD RCGWKLNSNMTF . SNGKLLVQDGFYLYANICFRHHETSGDATEYLOLMVYV
APRIL	182	VSREGQG RQETLFR CIRSM SHPDRAYNSCYSAGVFHLHQGDILSVIIPRARA KLNLSPHGTFLGFVKL
TNFa	157	SRIAVSY QTKVNLLSAIKSPCQRETPEGAEAKPWYERIYLGGVFOQEKGDRUSA EINRPDYLDFAESGQVYFGI IAL
LT. a	136	QLFSSQY PFHVPLLSSQKMV YPGLOEPWLHSMYHGAFAQLUTQGDQLSHTTDGIPHLVLSP . STVFFGAFA
FASL	212	YMRNSKY PQDLVMSQGKMS YCTTGQMWARS SYLGA VFNLTSA DHL YVNVSEL SIVN FEE . SOTFFGILYKL
TRAIL	189	VQYIYK YTSPDPILMKSARNS CWSKDAEYGLYSTYOGGIEELKENDRIFVSVTNEHLLIDMDH . EASFFGAFLV
TWEAK	223	ALRCLEE FSATAASSLGP QRLCQVSGLLALRPGSSLRIRTL PWAHLKAAP . FLTYFGLFQV
TRANCE	242	TKTSIKI PSSH TLMKGGSTK YMSGNSEFH FYSINVG GFFKLRSGEEISIEVS NPSL LDPDQ . DATYFGAFKV

Fig. 1

Fig. 2




Fig. 2D

D




Fig. 3




Fig. 4




Fig. 5

Comparison Human and Mouse APRIL

hAPRIL
mAPRIL
hAPRIL	QQTEQLSLRREVSRLQGTGGPSQN	PEQSSDALEAWENGERS
mAPRIL	QQTELQTLRREVSRLQRSGG	PSQKQGERPWQSLWEQSPDVLEAWKDGAKS
hAPRIL	RKRRAVLTQKQKKQH	SVLHLVPINATSKDDSDVTEVMWQPALRRGRGLQA
mAPRIL	RRRAVLTQKHKKKH	SVLHLVPVNITSKD-SDVTEVMWQPVLRGRGLEA
hAPRIL	QGYGVRIQDAGVYLLYSQVL	FQDVTFTMGQVSREGQGRQETLFRCIRSM
mAPRIL	QGDIVRVWDTGIYLLYSQVL	FHDVTFTMGQVSREGQGRRETLFRCIRSM
hAPRIL	PSHPDRAYNSCYSAGVFHL	HQGDILSIIIPRARA
mAPRIL	PSDPDRAYNSCYSAGVFHL	QGDIIITVKIPRANAKL