



US006083688A

United States Patent [19][11] **Patent Number:** **6,083,688****Lanza et al.**[45] **Date of Patent:** **Jul. 4, 2000**[54] **PLATELET GLYCOPROTEIN V GENE AND USES**[75] Inventors: **Francois Lanza**, Schiltigheim, France;
David R. Phillips, Oakland, Calif.;
Jean-Pierre Cazenave, Lampertheim, France[73] Assignee: **Cor Therapeutics, Inc**, South San Francisco, Calif.[21] Appl. No.: **08/195,006**[22] Filed: **Feb. 10, 1994****Related U.S. Application Data**

[63] Continuation-in-part of application No. 08/089,455, Jul. 9, 1993, abandoned.

[51] **Int. Cl.⁷** **C12N 15/12**[52] **U.S. Cl.** **435/6; 536/23.5; 536/24.1; 536/24.31; 435/252.3; 435/325**[58] **Field of Search** **536/23.5, 24.1, 536/24.31; 436/501; 435/69.1, 252.3, 320.1, 172.3, 6, 325**[56] **References Cited****PUBLICATIONS**Sambrook et al. *Molecular Cloning*. Cold Spring Harbor Press, pp. 8.4 and 8.76, 1989.Modderman et al. (1992) "Glycoproteins V and Ib-IX Form a Noncovalent Complex in the Platelet Membrane", *The Journal of Biological Chemistry*, 267:pp. 364-369.Berndt et al. (1981) "Purification and Preliminary Physicochemical Characterization of Human Platelet Membrane Glycoprotein V", *The Journal of Biological Chemistry*, 256(1):59-65.Phillips et al. (1977) "Platelet Plasma Membrane Glycoproteins", *Biochemical and Biophysical Research Communications*, 75(4):940-947.Sabatini et al. (1982) "Mechanisms for the Incorporation of Proteins in Membranes and Organelles", *The Journal of Cell Biology*, 92:1-22.Bienz et al. (1986) "Glycoprotein V is not the Thrombin Activation Receptor on Human Blood Platelets", *Blood*, 68(3):720-725.Zafar et al. (1989) "Platelet Membrane Glycoprotein V: Characterization of the Thrombin-Sensitive Glycoprotein from Human Platelets", *Thrombosis Research* 53:31-44.Wicki et al. (1989) "Isolation and Characterization of Human Blood Platelet mRNA and Construction of a cDNA Library in λ gt11", *Thromb. Haemostas.*, 61(5):448-453.Shimomura et al. (1990) "Rapid Purification and Characterization of Human Platelet Glycoprotein V: The Amino Acid Sequence Contains Leucine-Rich Repetitive Modules as in Glycoprotein Ib", *Blood*, 75(12):2349-2356.Roth (1991) "Developing Relationships: Arterial Platelet Adhesion, Glycoprotein Ib, and Leucine-Rich Glycoproteins", *Blood*, 77(1):5-19.Stubbs et al. (1993) "A Player of Many Parts: The Spotlight Falls on Thrombin's Structure", *Thrombosis Research* 69:1-58.George et al. (1984) "Molecular Defects in Interactions of Platelets with the Vessel Wall", *The New England Journal of Medicine* 311(17):1084-1098.Frohman et al. (1988) "Rapid Production of Full-Length cDNAs from Rare Transcripts: Amplification Using a Single Gene-Specific Oligonucleotide Primer", *Proc. Natl. Acad. Sci. USA* 85:8998-9002.Berndt et al. (1983) "Additional Glycoprotein Defects in Bernard-Soulier's Syndrome: Confirmation of Genetic Basis by Parental Analysis", *Blood* 62(4):800-807.Lanza et al. (1992) "A New Variant of Glanzmann's Thrombasthenia (Strasbourg I) Platelets with Functionally Defective Glycoprotein IIb-IIIa Complexes and a Glycoprotein IIIa ²¹⁴Arg \rightarrow ²¹⁴Trp Mutation", *J. Clin. Invest.*, 89:1995-2004.Faisst et al. (1992) "Compilation of Vertebrate-encoded Transcription Factors", *Nucleic Acids Research* 20(1):3-26.Roth et al. (1990) "Human Platelet Glycoprotein V: A Surface Leucine-Rich Glycoprotein Related to Adhesion", *Biochemical and Biophysical Research Communications* 170(1):153-161.Vu et al. (1991) "Molecular Cloning of a Functional Thrombin Receptor Reveals a Novel Proteolytic Mechanism of Receptor Activation", *Cell* 64:1057-1068.De Marco et al. (1991) "Function of Glycoprotein Ib α in Platelet Activation Induced by α -Thrombin", *J. Biol. Chem.* 266:23776-23783.Clemetson et al. (1982) "Characterization of the Platelet Membrane Glycoprotein Abnormalities in Bernard-Soulier Syndrome and Comparison with Normal by Surface-labeling Techniques and High-resolution Two-dimensional Gel Electrophoresis," *J. Clin. Invest.*, 70:304-311.Nurden et al. (1981) "Analysis of the Glycoprotein and Protein Composition of Bernard-Soulier Platelets by Single and Two-dimensional Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis," *J. Clin. Invest.*, 67:1431-1440.Vicente et al. (1990) "Identification of a Site in the α Chain of Platelet Glycoprotein Ib That Participates in von Willibrand Factor Binding," *J. Biol. Chem.*, 265:274-280.Lopez et al. (1987) "Cloning of the α chain of human platelet glycoprotein Ib: A transmembrane protein with homology to leucine-rich α_2 -glycoprotein," *Proc. Natl. Acad. Sci. USA*, 84:5615-5619.Wenger et al. (1988) "Structure of the Human Blood Platelet Membrane Glycoprotein Ib α Gene," *Biochem. Biophys. Res. Commun.*, 156(1):389-395.

(List continued on next page.)

Primary Examiner—Lorraine M. Spector
Assistant Examiner—Karen E. Brown
Attorney, Agent, or Firm—Morgan, Lewis & Bockius LLP

[57]

ABSTRACT

This invention relates to the glycoprotein v gene. Specifically, this invention discloses the sequence and structure of the glycoprotein v gene and the amino acid sequence of the glycoprotein v polypeptide. In addition, the evolutionary relationship of the glycoprotein v gene with other glycoproteins is described and several uses of the isolated glycoprotein v gene are shown.

12 Claims, 10 Drawing Sheets

OTHER PUBLICATIONS

- Lopez et al. (1988) "The α and β chains of human platelet glycoprotein Ib are both transmembrane proteins containing a leucine-rich amino acid sequence," *Proc. Natl. Acad. Sci. USA*, 85:2135–2139.
- Hickey et al. (1993), "Characterization of the Gene Encoding Human Platelet Glycoprotein IX," *J. Biol. Chem.*, 268(5):3438–3443.
- Hickey et al. (1989), "Human platelet glycoprotein IX: An adhesive prototype of leucine-rich glycoproteins with flank-center-flank structures," *Proc. Natl. Acad. Sci. USA*, 86:6733–6777.
- Hickey et al. (1993), "Human platelet glycoprotein V: Characterization of the polypeptide and the related Ib–V–IX receptor system of adhesive, leucine-rich glycoproteins," *Proc. Natl. Acad. Sci. USA*, 90:8327–8331.
- Lanza et al. (1993), "Cloning and Characterization of the Gene Encoding the Human Platelet Glycoprotein V," *J. Biol. Chem.*, 268(28):20801–20807.




FIG. 1.




FIG. 2A.




FIG. 2B.




FIG. 3.




FIG. 4.

5' - TGATCGGAAC	TGAAAGACCT	CCC GCGATAC	CTGGCAGAGG	CAGTGGCTCT	50
TRE					
TCCTGTGGT	CCAGGGCTGA	CTGACTTTGA	AGGTAATTTT	AGTCAACCCA	GCCTTACTG 110
GGCTCTGACT	GCATTAGGCT	GCATCAAAGG	GGATTGGATC	CCATGATTCT	TTATATCTT 170
TGACATTAAG	CCTTTGTGAG	CTATAGGTGT	TACAAATATC	TTTAGTTTGT	GGTTTATCTT 230
TTCCCTTTT	TTATGGTGTG	TTGAAGGATA	GAAGTCTTAA	TGCAGACAGC	ATTATCAGTG 290
TGTTCAAAG	ACAGCTAGAC	ACGTTTTGCC	TATAGACAAA	TGGGCAAAG	GAAACCCAGC 350
TTTCTCAAT	GAAGCACAAAG	TGGGCCTTAA	TTATGTGAAA	AGGTGTTCAA	GTTTCATCATT 410
AAACAGGAA	AGGAAAAGTT	AAAACCATGC	TGAGATATCT	TTCATAGAAA	TGGCAAAG 470
Ets-1		Ets-1			
CAGGAAGTGC	CACGTGTGGG	CAGAGAGGAA	GCACAGGAAC	TCTCACAAT	GGCAGGTGTC 530
ATCGTAGACC	AACACAACCA	CTTTGGAGAG	CAGTTTGACT	TTCCCGAGTT	AAACTGAACA 590
TGTGAGCGGC	CGGGCGTGGT	GGCTCATGCC	TGTAATCCCA	GCAGTTTGGG	AGGCCGAGGC 650
GGGGGATTG	CCTGAGCTCA	GGAGTTCAAG	AGCAGCCAGG	GCAAGACGGT	AAAACCCGGT 710
CTCTACTAAA	ATAGAAAAAA	TTAGCTGGGC	GTGATGGTGT	GTGCCTGTAA	TCCAGCTAC 770
TTGTGAGGCC	GAGGCAGGAG	AATGCTTGA	ACCAGGGAGC	AGGAGGTTGC	AGTGAGCCGA 830
GATCGACCA	CTGCACCCCA	GCCTGGCGAC	AGAGTCCCCC	TCCCCACCA	AAAAACAAC 890
Ets-1					
AAGTGACCAT	CCTGCAACCT	AGCAATGCCA	TTGTTGAACA	AGTTCAAAGA	TGTTCTTAGC 950
CTTATTAGTC	CCAAAAGGAA	GAAAAAATG	GAGGATTTGA	GAATGTTCTT	AGCTTTATTG 1010
CTAAGCGGAG	AAAGAAAAAC	AACACATACC	AAAAAAAAAA	AAAAAAAAAA	AAAAAACAA 1070
AAAACCTGGG	TGGAAATTA	GGGCCATGTG	GCATGAAAAG	GAAGACCCAG	GGGAAGTGTG 1130
Spl		Ets-1			
GCCCATCTAG	GGGTGTGGCT	ACTGCAGTGA	TCCAGCTGTA	TCACTGA	ACTTCCGTGGCAT 1190
TATA					
CATAGAGTTA	TATTGTGCCA	TTTATGAAA	AACTCTCCCC	ACTGCTCTTG	GCTTTGACAG 1250
TATA					
TAGGAATCAG	GTATATATG	GTCTCTGGT	TTGAAGATA	TTGTATTAA	AAACCAGAAG 1310
GATA					
AAGGGCTCTG	AGATAGGGTC	CTTCC TGAC	CTACTCTGGT	AAAGTCTTTA	TCCTCAGGAT 1370
Ets-1					
CAAGGATAC	CACCCTCTTC	CTG TGAAAAG	TGTCGAATCA	CATGCAGAGC	TCTAAGTCTT 1430
▽					
TCAGTTACTT	TGGAGTGCAG	AACCATTTCA	Ggtaaggcca	aataTTTTaa	acattagtat 1490
aggaattag	agggetcttt	agtcfgtg	tgcatgagaa	gtaaaattgc	acgagaagca 1550
atttatgtaa	aatttegett	aggaacatt	gittfggtag	gttagtagta	tggtgtgtat 1610
ttccagaaa	attcagtgcc	gtgagtatta	cctttagtta	agcaicttag	aaatagtagc 1670
tcttattggt	tatggctaa	tcagaaatac	ta cccctcaa	ttctatgtga	ccctagttaa 1730
actgttgogc	ctttctgtgc	ctcftgtcet	tcatecttga	atcggggata	atatacttac 1790
ctcctaagg	tattgttaagg	attaaatgca	tgtagtataa	ataaogagct	gagaa caatg 1850
catggcgta	agiga taggt	otta tta tat	gittttgttg	gctgttgatt	gaagggtttt 1910
gctgttttg	gggtgtcett	taatagagta	acttggtaet	gtggaaatag	catgattgtg 1970
agcaaaagaa	tcogotggg	gtggctgcag	actttgctgt	tccttcttg	actgttgggt 2030
atagccaatg	cagggttaagt	tataaagtca	agagcagagc	cgttttcaca	atgga catg 2090
ctttgtgatg	tctgtgaget	tgaatgtgag	aatgattatt	ttaattctet	atgtaagagc 2150
tttaaaagtat	tggctatfcg	gtagctfgat	ttctctgtaa	tctcatgctt	taaac tgaga 2210
gtggaaaate	aataaagca	aagc atgagg	ccacgcagtg	tagaattgagt	gtctttcac 2270
cacgtaggga	aatcftgagt	cctaagaaa	gagggagtg	gaattctggc	gaaaagattg 2330
tgcctctgca	caaagtgcag	gatcc caggg	ttcagta cag	gcgcgaacgc	tcctgtgtgt 2390
Met					
tgaccacact	cccacggttg	cttttttagA	CATGCTGAGG	GGGACTCTAC	TGTGGCGGGT 2450

FIG. 5A

GCTGGGGCTT	CTGCGCGCCC	AGCCCTTCCC	CTGTCCGCCA	GCTTGCAAGT	GTGTCTTCCG	2510
GGACGGCGCG	CAGTGCTCGG	GGGGCGACGT	GGCGCGCATC	TCCGGCGCTGG	GCCTGCCAC	2570
CAACCTCACG	CACATCCTGC	TCTTCGGAAT	GGGCCGCGGC	GTCCTGCAGA	GCCAGAGCTT	2630
CAGCGGCATG	ACCCTCTCTG	AGCGCCTCAT	GATCTCCGAC	AGCCACATTT	CCGCCGTTGC	2690
CCCCGGCACC	TTCAGTGACC	TGATAAAACT	GAAAACCCTG	AGGCTGTGGC	GCAACAAAAT	2750
CACGCATCTT	CCAGGTGCGC	TGCTGGATAA	GATGGTGCTC	CTGGAGCAGT	TGTTTTTGGG	2810
CCACAATGCG	CTAAGGGGCA	TTGACCAAAA	CATGTTTCAG	AAACTGGTTA	ACCTGCAGGA	2870
GCTCGCTCTG	AACCAGAATC	AGCTCGATTT	CCTTCCTCCC	AGTCTCTTCA	CGAATCTGGA	2930
GAACCTGAAG	TTGTTGGATT	TATCGGGAAA	CAACCTGACC	CACCTGCCCA	AGGGGTTGCT	2990
TGGAGCACAG	GCTAAGCTCG	AGAGACTTCT	GCTCCACTCG	AACCGCCTTG	TGTCTCTGGA	3050
TTCGGGGCTG	TTGAA CAGCC	TGGGCGCCCT	GACGGAGCTG	CAGTTCACCC	GAAATCACAT	3110
CCGTTCCATC	GCACCCGGGG	CCTTCGACCG	GCTCCCAAC	CTCAGTTCCT	TGACGCTTTC	3170
GAGAAAC CAC	CTTGCGTTTC	TCCCCTCTGC	GCTCTTTCTT	CATTG CACA	ATCTGACTCT	3230
GT T GACTCTG	TTGAGAACC	CGCTGGCAGA	GCTCCCGGGG	GTGCTCTTCG	GGGAGATGGG	3290
GGGCCTGCAG	GAGCTGTGGC	TGAACCGCAC	CCAGCTGCGC	ACCCTGCCCG	CCGCCGCCCT	3350
CGGCAACCTG	AGCCGCCTGC	GGTACTTAGG	GGTGACTCTG	AGCCCGCGGC	TGAGCGCGCT	3410
TCCGCA GGGC	GCCTTC CAGG	GCCTTGGCGA	GCTCCA GGTG	CTC GCCCTGC	ACTCCAACGG	3470
CCTGACC6CC	CTCCC6ACG	GCTTG CTGG	C6GCCTC6GC	AAGCTGCGCC	AGGTGTCCCT	3530
GGCCCGCAAC	AGGCTG C6CG	CCCTGCCCGG	TGCCCTC TTC	CGCAAT CTCA	GCAGCCTGGA	3590
GAGCCTCCAG	CTCGACCACA	ACCAGCTGGA	GACCCTGC CT	GGCGACG TGT	TTGGGGCTCT	3650
GCCCCGGCTG	ACGGAGGTCC	TGTTGGGGCA	CAACTCCTGG	CGCTGCGACT	GTGGCCTGGG	3710
GCCCTTCCTG	GGGTGGCTGC	GGCAG CACCT	AGG CCTCGTG	GGCGGGGAAAG	AGCCCCCAGG	3770
GTGCGCAGGC	CCTGGGGCGC	ACGC CGGCT	GCCGCTCTGG	GCCC TGCCGG	GGGGTGACCG	3830
CGAGTGCCCG	GGCCCCGGGG	GCCCCCTCC	CCGCCCGCT	GCCCACAGCT	CCTCGGAAGD	3890
CCCTGTCCAC	CCA GCCTTGG	CTCCCAACAG	CTCAGAACCC	TGGGTGT GGG	CCCAGCCGGT	3950
GACCACGGGC	AAAGGTCAAG	ATCATAG TCC	GTTCTGG GGG	TTTTATT TTC	TGCTTTAGC	4010
TGTTCAAGGCC	ATGATCACC	TGATCA TCGT	GTTTGCTA TG	ATTAAAT TC	GCCA ACTCTT	4070
STOP						
TCGAAAATT A	ATCAGA GAG A	GAGCCC TTGG	GTAACCAAT	GGGAAAATCT	TC TAATTACT	4130
TAGAACCTGA	CCAGATGTGG	CTCGGAGGGG	AATCCAGACC	CGGTGCTGTC	TTGCTCTCCC	4190
TCCCCTCCCC	ACTCCTCCTC	TCTTCTTCC	CTTCTCTCTC	ACTGCCACGC	CTTCCTTTCC	4250
CTCCTCCTCC	CCCTCTCCGC	TCTGTGCTCT	TCATTCTCAC	GGGCCCGCAA	CCCCCTCCTCT	4310
CTCTGT CCCC	GCCCGTCTCT	GGAAG CTGAG	CTTGACGTTT	GTAAACTG TG	GTTGCCTGCC	4370
TTCCAGCTC	CACGCGGTGT	GCGCTGACAC	TGCCGGGGGG	CTGGA CTG TG	TTGGACCCAT	4430
CCTTG CCCC	CTGTGCCTGG	CCTGGCCTCT	GGTGGAGAGA	GGGACCTCTT	CAGTGTCTAC	4490
TGAGTAAGGG	GACAGCTCCA	GGCCGGGGCT	GTCTCCTGCA	CAGAGTAAGC	CGGTAAATGT	4550
TTGTGAAATC	AATGCGTGGA	TAAAGGAACA	CATG CCA TCC	AAGTGATGAT	GGCTTTTCT	4610
GGAGGGAAAG	GATAGGCTGT	TGCTCTATCT	AATTTTTTGT	TTTTGTTTTT	GGACAGTCTA	4670
GCTCTGTGGC	CCAGGCTGGC	GTGCACTGGG	CCGTCTCAGT	TCACTGCAGC	CTCCGCCCTC	4730
CAGGTTCAAG	TGATTC TCAT	GCCTCA GCCT	TCT GAGTAGC	TGGGATTAGA	GGCGTGTGCC	4790
ACTACACCCG	GCTAATTTTT	GTACTTTTTA	AAGTAGAGAC	GGGCTTTGCC	ATA TTGGCCT	4850
G GCTGATCTC	AAACTCC TGG	TCTTGAACTC	CTGGCCACAA	GTGATCTGCC	CGCCTTAGCC	4910
TCCCAAAGTG	CTGGGATTAC	AGGCGCAAGC	CACTACACCT	GCCCTCTTCA	TGCAATTTTA	4970
TTTGAGAAGT	AGAGCTCTTG	CCATTTTTTC	CCTTGCTCCA	TTTTTCTCAC	TTTATGTCTC	5030
TCTGACC TAT	GGGCTACTTG	GGAGAGCACT	GGACTCCATT	CATGCATGAG	CATTTTCAGG	5090
ATAAGCGACT	TCTGTGA GGC	TGAGAGAGGA	AGAAAA CAGG	GAGCCTTCCC	TCCAGGTGCC	5150
CAGTGTAGGT	CCAGCGTGT	TCCTGAGCCT	CCTGTGAGTT	TGCACTTGCT	TTACATCCAT	5210
GCAACATGTC	ATTTTGAAC	TGATTGATT	TGCATTTCT	GGA ACTCTGC	CACCTCATTT	5270
CACAAGCATT	TATGGAGCAG	TTAACATGTG	ACTGGTATTC	ATGAATATAA	TGATAAGCTT	5330

FIG. 5B

GATTCTAGTT	CAGCTGCTGT	CACAGTCTCA	TTTGTCTTC	CAACTGAAAG	CCGTAAAACC	5390
TTTGTGCTT	TAATTGAATG	TCTGTGCTTA	TGAGAGGCAG	TGGTTAAAAC	ATTTTCTGGC	5450
GAGTTGACAA	CTGTGGGTTG	AAATCCCAGC	TCTACCACCT	ACTAACTGCA	TGGGACTTTG	5510
GGTAAGACAC	CTGCTTACAT	TCTCTAAGCC	TTGGTTTCCT	GAACCTTAAA	ACAGGATAAC	5570
ATAGTACCTG	CTTCATAGAG	TTTTGTGAGA	ATTAAGGCA	<u>ATAAAGCATA</u>	TAATGACTTA	5630
GCCCAGCGGC	CTGCAGACAA	TACATGTTAA	TGAATGTTAG	CTATTATTAC	TAAAGATGAG	5690
CAATTATTAT	TGGCATCATG	ATTTCTAAAG	AAGAGCTTTG	AGTTGGTATT	TTTCTCTGTG	5750
TATAAGGGTA	AGTCCGAACT	TTCTCATACT	GGAGGTTACA	TTCACATCAG	TCTGTCTTCC	5810
CCTGCGGATG	GCCTCAGCCC	TGGGTGGCCA	GGCTCTGTGC	TCACAGTCCA	GAGCAATGGA	5870
TCCTCCAACA	CCACCAGGTG	GATGTGGAGC	AGGAGAGCTG	GATCGTGGCA	TTTGTTCCTG	5930
GGTTCGCGAG	TTGGGAGTTG	GTTTCTGGGT	TCTCCATTGG	TCTACTTGTG	TAGTCCCATA	5990
CCAGACTCAC	GGTCTCCATT	ATTGGAGCTT	TAATAATTTT	TGGTATAGGG	TCATCTCTCC	6050
ACCTTGTTTT	TCTTCTATTC	TTGGTTCTTT	GCAATTCTAT	GAATATTTCA	GGGTCAGCAT	6110
GTCAACTCCA	TTGAAAACC	CTGCTGGGAT	TTTAATAGAA	<u>CTTACAGCTC</u>	<u>ACGCCTGTAA</u>	6170
<u>TCCAGCACT</u>	<u>TTGGGAGGCT</u>	<u>GAGGTGGGTG</u>	<u>GATCACAGGT</u>	<u>CAGGAGTTTG</u>	<u>AGAACAGCTG</u>	6230
<u>GCCAAGATGG</u>	<u>TGAAACCCCG</u>	<u>TCTCTACTAA</u>	<u>AAATACAAAA</u>	<u>ATTAGCTGGG</u>	<u>TGCGGTGGCA</u>	6290
<u>GGTGCCTGTA</u>	<u>GTCCCAGCTA</u>	<u>CTTGGGACAC</u>	<u>CGAGGCAGGA</u>	<u>GAATCACTTG</u>	<u>AACCCGGGAG</u>	6350
<u>GCGGAGGTTG</u>	<u>CAGTGAGCCG</u>	<u>AGATCGTGCC</u>	<u>ACTGCCTCT</u>	<u>AGCCTGGGCG</u>	<u>ACAGAGCGAG</u>	6410
<u>ACTCCATCTC</u>	<u>AAAAAAAAAG</u>	<u>AAAAAGAAA</u>	<u>TTGCAGTAA</u>	<u>TTTAAACTA</u>	<u>ATTTGGGGAA</u>	6470
GAATCTGTAT	TTTTACAATA	CCTAGTGTTC	TTGCCAGTAA	GCATGGTTCA	TCTTCCATT	6530
TATTTACGTC	ATTTAAATC	TTTCAGTGAT	GTTTTAGAAT	TTTTTTTATA	AAAACCTTCA	6590
CTATAAGAAC	AGAAAACCAA	ACACCGCATG	TTCTCACTCA	TAGGTGGGAA	TTGAACAATG	6650
AGAACACTTG	GACACAGGGC	GGGGAACGTC	ACACGCCTGG	ACTGTTGGGG	GGGTGGCTGG	6710
GAGAGGGATA	GTGTTAGGAG	AAATACCTAA	TGTAATGAC	GAGT TAATGG	TGCAGCCAAC	6770
CAACCTGGCA	CATGTATTCA	TATGTAACAA	ACCTGCACGT	TGTGCACATG	TACCCTAGAA	6830
CTTAAAGTAT	ATTAATAAAA	GAAACCTTGG	CACTGATTTT	GTTAGATTTA	TTCCTAGGTA	6890
TCCTTCTCT	TTTTTGATTT	GTCAATTGCTA	TTGTAGATGG	CATCTTTTTA	AAAAGTTATA	6950
TTTTCTAAAG	<u>CAAAAATAA</u>	<u>AAAAAGTTGT</u>	<u>ATTTCTAATT</u>	<u>TTTATTACCA</u>	<u>ATATATAAGA</u>	7010
ATGTAATTTA	TTTTTACATA	ATTATCTTAT	GTCTAGTAAT	AATTCTGATA	ATTTGCTTCT	7070
TCCTATTTAA	ACCTTACACC	CATTATTGAT	TTATTTTCT	GTTTTAAAT	ATCTTCCTGC	7130
ACTGGCTAAA	ACCTCCACTA	TAATGTTGAG	CAGAACAGTG	AGGCATCCTT	AGAACTATCT	7190
TGGTTGCAAA	GGGTA GGTCT	CTAATGTTTC	<u>ATCAATAAA</u>	GTGATGTTTC	TAGTCTGAGT	7250
TTGCTAAGTA	TATTTTAAAA	TAATCAGTAA	AGTTAGATTT	TATCCATTTT	TATCTTAACT	7310
ATTGAGATGC	TCATATCATT	TTTCTTCTTC	AATGTGTTAA	<u>AATGGTGAA</u>	<u>AAA</u> TTTATAG	7370
ATTTTGAAA	AGTAAATTCA	TTCTTGCAAT	CCCGAAGTAA	ACCAAGCCAT	GCTATGTGTA	7430
TTTAAATAT	ATTGCTGAAT	TC-3				7452

FIG. 5C

1 I M L R G T L L \textcircled{C} A V L G L L R A Q P F P \textcircled{C} P P A \textcircled{C} K \textcircled{C} V F R
 31 D A A Q \textcircled{C} S G G D V A R I S A L G L P T \downarrow N L T H I L L F G M
 61 G R G V L Q S Q S F S G M T V L Q R L M I S D S H I S A V A
 91 $\overset{m7}{P}$ G T F S D L I K L K T L R L S R N K I T H L P G A L L D K
 121 M V L L E Q L F L D H N A L R G I D Q N M F Q K L V N L Q E
 151 $\overset{k5}{L}$ A L N Q N Q L D F L P A S L F T N L E N L K L L D L S G N
 181 \downarrow N L T H L P K G L L G A Q A K L E R L L L H S N R L V S L D
 211 S G L L N S L G A L T E L Q F H R N H I R S I A P G A F D R
 241 L P N L S S L T L S R N H L A F L P S A L F L H S H N L T L
 271 L T L F E N P L A E L $\overset{m401}{P}$ G V L F G E M G G L Q E L W L N R T
 301 Q L R T L P A A A F R N L S R L R Y L G V T L S P R L S A L
 331 P Q G A F Q G L G E L Q V L A L H S N G L T A L P D G L L R
 361 G L G K L R Q V S L R R N R L R A L P R A L F R N L S S S L E
 391 S V Q L D H N Q L E T L P G D V F G A L P R L T E V L L G H
 421 N S W R \textcircled{C} D \textcircled{C} G L G P F L G W L R Q H L G L V G G E E P P R
 451 \textcircled{C} A G P G A H A G L P L W A L P G G D A E \textcircled{C} P G P R G P P P
 481 R P A A D S S S E A P V H P A L A P N S S E P W V W A Q P V
 511 T T G K G Q D H S P F W G F Y F L L L A V Q A M I T V I I V
 541 F A M I K I G Q L F R K L I R E R A L G 560

FIG. 5D




FIG. 6.




FIG. 7.




FIG. 8.

PLATELET GLYCOPROTEIN V GENE AND USES

This application is a Continuation-in-Part of U.S. application Ser. No. 08/089,455 filed Jul. 9, 1993 now abandoned.

BACKGROUND OF THE INVENTION

Platelets arise from the fragmentation of megakaryocytes, which are large polyploid bone marrow cells produced by several cycles of chromosomal duplication without cytoplasmic division (Handin (Wilson et. al., eds) in *Harrison's Principles of Internal Medicine*, 12th edition (1991)). Once free of the marrow space, approximately $\frac{2}{3}$ of the platelets circulate freely, while approximately $\frac{1}{3}$ are sequestered in the spleen. Circulating platelets last for 7 to 10 days, after which they are removed by phagocytic cells. A decrease in platelet mass stimulates megakaryocytopoiesis, resulting in an increase in the number, size and ploidy of the megakaryocytes.

Platelet receptors which mediate platelet adhesion and aggregation are located on the two major platelet surface glycoprotein complexes. These complexes are the glycoprotein Ib-IX complex which facilitates platelet adhesion by binding von Willebrand factor (vWF), and the glycoprotein IIb-IIIa complex which links platelets into aggregates by binding to fibrinogen. Patients with the Bernard-Soulier syndrome, a congenital bleeding disorder, show deficient platelet adhesion due to a deficiency in the glycoprotein Ib-IX complex which binds VWF, mild thrombocytopenia, and large lymphocoid platelets.

Glycoprotein v (GPV) is a major ($\approx 12,000$ molecules/platelet), heavily glycosylated platelet membrane protein (Mr 82,000) (Modderman et. al. *J. Biol. Chem.* 267:364-369). Earlier reports showing that GPV was a peripheral protein (Berndt and Phillips *J. Biol. Chem.* 256:59-65) were presumably due to the release of GPV from the membrane by calpain during the purification procedure. Exposure of platelets to thrombin liberates a 69 kDa soluble fragment termed GPVfl (Phillips and Poh-Agin, *Biochem. Biophys. Res. Commun.* 75:940-947). This, and its absence in the Bernard-Soulier syndrome (Clemetson et. al. *J. Clin. Invest.* 70:304-311 (1982); Nurden et. al., *J. Clin. Invest.* 67:1431 (1981); Berndt et. al., *Blood* 62:800-807 (1983)), led to the suggestion that GPV may be involved in the thrombin-induced activation response (Berndt and Phillips L. *J. Biol. Chem.* 256:59-65 (1981)). Recent experiments show that GPV can interact non-covalently with the GPIb-IX complex (Modderman et. al. *J. Biol. Chem.* 267:364-369 (1992)), a complex formed by the non-covalent association of GPIb (consisting of GPIb α , a 145 kDa protein, disulfide linked to GPIb β , a 24 kDa protein) with GPIX (a 22 kDa protein). The binding sites for von Willebrand factor and for thrombin on the GPIb-IX complex have been localized on GPIb α (Wicki and Clemetson *Eur. J. Biochem.* 153:1-11 (1985); Vicente et. al., *J. Biol. Chem.* 265:274-280 (1990)). Since thrombin is now known to activate platelets by cleaving the thrombin receptor (Vu et. al., *Cell* 64:1057-1068 (1990)), a G-protein coupled receptor, it is unknown whether thrombin cleaves GPV incidentally as a consequence of thrombin binding to GPIb α , or whether this cleavage has a physiological role.

The amino acid sequences of GPIb α , GPIb β , and GPIX have been deduced from their cDNA and genomic sequences (Lopez et. al., *Proc. Natl. Acad. Sci. USA* 84:5614-5619 (1987); Wenger et. al., *Biochem. Biophys. Res. Commun.*

156:389-395 (1988); Lopez et. al., *Proc. Natl. Acad. Sci. USA* 85:2135-2139 (1988); Hickey, et. al., *Proc. Natl. Acad. Sci. USA* 86:6733-6777 (1989); Hickey and Roth *J. Biol. Chem.* 268:3438-3443 (1993)). Analysis of the primary amino acid sequence of GPIb α , GPIb β , and GPIX has revealed a common evolutionary origin for the three proteins, as they contain one or more homologous 24 amino acid leucine-rich domains. These domains are also found in a large family of leucine-rich glycoproteins (LRG) including leucine-rich $\alpha 2$ GP, proteoglycan core, fibromodulin, human lutropin-chorio gonatropin receptor and RNase inhibitor, and toll protein and chaoptin found in *Drosophila* (reviewed in Roth *Blood* 77:5-19 (1991)). Recently, analysis of partial peptide sequences obtained from purified platelet GPV suggested that GPV is also a member of the LRG family (Shimomura et. al., *Blood* 75:2349-2356 (1990); Roth et. al., *Biochem. Biophys. Res. Commun.* 170:153-161 (1990)).

GPV is a very specific marker for the megakaryocytic cell lineage. A monoclonal antibody specific for GPV (SW16) was recently shown to bind exclusively to platelets (Modderman et. al., *J. Biol. Chem.* 267:364-369 (1992)). SW16 did not bind to red cells, leukocyte endothelial cells, or cell lines such as HEL or MEG-01 which are known to express platelet megakaryocyte markers.

SUMMARY OF THE INVENTION

The invention comprises an isolated DNA construct comprising the polynucleotide sequence of the glycoprotein v gene, including the polynucleotide sequence which has the sequence shown in FIGS. 5A-C. The polynucleotide sequence encodes a GPV polypeptide, including the amino acid sequence as shown in FIG. 5D. The polynucleotide sequence may lack introns, and may incorporate a heterologous promoter operably linked to the polynucleotide sequence which is capable of directing expression in a prokaryote or in a eukaryote.

The invention further comprises a DNA construct wherein the polynucleotide sequence encodes a full length glycoprotein V polypeptide.

The present application includes prokaryotic or eukaryotic a cell containing a glycoprotein v DNA construct.

The present application further provides for a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a glycoprotein v polypeptide. The polypeptide may have the sequence shown in FIG. 5D.

A further understanding of the nature and advantages of the invention will become apparent by reference to the remaining portions of the specification and drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 part (a) illustrates the cloning, sequencing strategy, and restriction map of a partial human platelet GPV cDNA, whereas part (b) illustrates the complete human GPV gene.

(a) The top line represents the coding region (open bar) and 5'-untranslated sequence (hatched bar) for a platelet GPV cDNA with a partial restriction map. The cloning strategy is indicated below. Overlapping clones (i to vi) covering 1,199 bp of cDNA were obtained after PCR amplification of platelet mRNA. The oligonucleotide primers used for the amplification are indicated and the corresponding sequences are listed in Table I. The GPV 8.1 kb genomic fragment (1) was obtained after screening a human genomic library in the λ Fix vector with a 748 bp 32 P-labelled GPV cDNA probe (indicated in (a) by a broken line). The top line is a partial restriction map of the gene. Exons are boxed: the

open box represents the coding sequence, the hatched box represents the 5'-untranslated sequences, and the shaded box represents the 3'-untranslated region. The vertical arrow indicates the beginning of the genomic sequence reported in FIGS. 5A–C. The open arrowhead indicates the 5'-end and the closed arrowhead indicates the 3'-end of the partial platelet cDNA obtained by PCR. A sequence with perfect consensus for a TATA box is indicated. The closed circles indicate AATAAA consensus sequences for polyadenylation signals. The restriction sites are indicated as follows A, Acc I; Be Bam HI; E, Eco RI; K, Ksp I; P, Pst I; Se Sac I; X, Xho I. FIG. 2A illustrates the detection of GPV mRNA in platelets and megakaryocytes by RT-PCR amplification. FIG. 2B illustrates the detection of GPV mRNA in HEL cells by RT-PCR amplification. Total RNA (25 ng) was subjected to reverse transcription followed by PCR with GPV cDNA based primers, and in a control reaction with primers for the housekeeping GAPDH gene. Ten μ l of the PCR reactions were separated on a 2% agarose gel stained with ethidium bromide and are shown together with Bgl I/Hinf I cut pBR328 DNA molecular-weight markers. FIG. 2A shows that platelet (PLT) and megakaryocyte (MGK) RNA were amplified with a mixture of two GPV primer pairs (nt 3,010–3,589 and 2,675–2,877) generating bands of 579 bp and 202 bp. FIG. 2B shows that HEL cells with (HEL+PMA) or without (HEL) stimulation with phorbol ester, HL60, and platelet (PLT) RNA were amplified with a GPV primer pair (nt 3,091–3,589) generating a 498 bp band.

FIG. 3: Northern blot analysis. Total RNA (10 μ g per lane) from human platelets (lanes a and b) and human monocytes (lane c) was electrophoretically separated on 1% agarose-formaldehyde gel, transferred to Zetaprobe, and probed with a 748 bp random prime 32 P-labelled cDNA probe. Lane d is an ethidium bromide stained gel of leukocyte total RNA showing the position of 28S and 18S ribosomal RNA. The molecular size was calibrated by reference to the migration of λ /Hind III DNA fragments indicated in kilo base pairs.

FIG. 4: Southern blot analysis. High molecular weight genomic DNA (10 vg) from human leukocytes was cut with an excess of Eco RI, Bam HI, and Bgl II restriction endonucleases, separated on a 0.7% agarose gel, and transferred to Hybond N⁺ nylon membranes. The filters were probed with a 748 bp 32 P-labelled GPV cDNA fragment. The size of the hybridizing bands in kilobase pairs was estimated by comparison with λ /Hind III DNA fragments.

FIGS. 5A–C (SEQ.ID.NO. 1) and 5D (SEQ.ID.NO. 2): Sequence of the human GPV gene FIG. 5A–C and deduced amino acid sequence FIG. 5D of the GPV protein. The GPV genomic sequence (SEQ.ID.NO. 1) FIGS. 5A–C are shown in the 5'- to 3'-orientation with the single intron sequence of 958 bp shown in lower case letters. The gt/ag donor and acceptor sites are in bold characters. Consensus sequences for putative cis-acting promoter elements are indicated as shaded areas. The closed circle indicates a possible Cap site. The ATG translation start and the in-frame TAA stop codon are boxed. The open arrowhead (nt 1,433) and closed arrowhead (nt 3,589) indicate the 5'- and 3'- end, respectively, of the partial cDNA sequence obtained by PCR amplification of platelet RNA. Two Alu repeats, nt 598–886 and nt 6,133–6,440, are underlined. Possible polyadenylation signal sequences (nt 5,610, nt 6,966, nt 7,224 and, nt 7,358) are double underlined. The GPV amino acid sequence FIG. 5D (SEQ.ID.NO. 2), indicated in single letter code, was deduced after translation of the cDNA and genomic sequences. The putative signal peptide is underlined. The putative transmembrane domain is double underlined. Cysteine residues are circled. Potential N-linked glycosylation

sites in the extracellular domain are indicated by a vertical arrowhead. N-glycosylation sites that had been identified by protein sequencing are indicated by a star. Internal peptide sequences that were obtained from purified platelet GPV (20, 21), indicated in italics, are underlined by a broken arrow. Differences between the DNA-derived and internal peptides sequences are indicated in parenthesis as lower case letters. (x) indicate a residue which had not been determined in the original peptide sequence.

FIG. 6: Alignment of the 15 tandem Leu-rich repeated structures for platelet GPV (SEQ.ID.NOs. 22–36). The alignment spans the sequences between residues 61 and 421 of the protein. Identical residues among the 15 segments are boxed. An overall consensus sequence for the GPV repetitive motifs is presented (SEQ.ID.NO. 37).

FIG. 7: comparison of the GPV thrombin cleavage site to other thrombin substrates. The GPV sequence around the RG thrombin cleavage peptide bond (SEQ.ID.NO. 38) was aligned with sequences of human fibrinogen (Fg) α (SEQ.ID.NOs. 39 and 40) and β (SEQ.ID.NO. 41) chains, to human plasma factor XIII (FXIII) (SEQ.ID.NO. 42), and to human chorionic gonatropin β -subunit (CG β) (SEQ.ID.NO. 43). Amino acid residues identical to GPV are boxed.

FIG. 8 Schematic representation of the GPV protein inserted in the platelet plasma membrane in comparison with the GPIb-IX complex. The proteins depicted as bars, are oriented with their NH₂— and COOH-termini oriented toward the outside and inside of the cell, respectively. Numbering of amino acids for the mature proteins is indicated. The transmembrane domains are represented as solid rectangles. The Leu-rich (LR) repetitive domains are represented as hatched rectangles. N-glycosylation sites are indicated as solid triangles. GPIb α contains a region rich in O-linked sugars (O—CHO) and is linked to GPIb β by a disulfide (S—S) bond. The location of the thrombin cleavage site in GPV is indicated by a double-headed arrow.

DETAILED DESCRIPTION OF SPECIFIC EMBODIMENTS

The present invention provides the primary structure of the human GPV gene and the structure of the GPV protein. The single-copy gene for GPV is contained within 6.5 kb of genomic sequence, and has a simple structure with a single intron of 958 bp in the 5'-untranslated sequence; the coding sequence is contained within a single exon. The promoter region contains a canonical TATA box, and putative GATA, Ets-1, and Spl cis-acting elements. RT-PCR analysis on RNAs from cells of different hematopoietic origins revealed that GPV was specifically transcribed from platelets and from calls of the megakaryocytic lineage (megakaryocytose HEL cells). A single transcript of 405 kb for GPV was detected in human platelets by Northern blot analysis, and the entire amino acid sequence of GPV was deduced from the cDNA and genomic sequences.

Mature GPV is composed of 543 amino acids which contain a single transmembrane domain, a short cytoplasmic domain (16 residues) and a large extracellular domain with 8 potential N-glycosylation sites. Analysis of the extracellular domain revealed the presence of 15 tandem Leu-rich repeats of 24 amino acids with homology to GPIb α , and identified a cleavage site for thrombin near the C-terminus with homology to the α chain of fibrinogen.

The predicted amino acid sequence of GPV accounts for the known features of the protein. First, it contains with one exception (peptide M4 Shimamura et. al. *Blood*

75:2349–2356 (1990) all of the partial peptide sequences which had been reported for purified platelet GPV (FIGS. 5A–C). Second, the predicted molecular weight of the polypeptide chain of 59,276 Da agrees with the 60 kDa value determined after SDS-PAGE analysis of the deglycosylated protein. Third, the predicted amino acid composition is very similar to that reported for purified GPV when the data are corrected for the 59,276 molecular mass. Fourth, the LRG repeats in GPV display significant similarity to those found in the subunits of the GPIb-IX complex, which GPV associates with in platelets. Finally, the translated protein contains a thrombin cleavage recognition site at a position which would generate a soluble cleavage fragment of the size of GPVfl, a fragment known to be generated after platelet treatment with thrombin (Phillips and Poh-Agin, *Biochem. Biophys. Res. Commun.* 75:940–947 (1977); Mosher et al., *Blood* 53:437–445 (1979)).

Analysis of the deduced primary amino acid sequence revealed several distinctive features for GPV. The protein contains an N-terminal signal peptide with a consensus cleavage site (Von Heijne, *J. Mol. Biol.* 173:243–251 (1984)) at a Gln residue. N-terminal glutamines are often cyclized to pyroglutamic acids, explaining the N-terminal blockade consistently observed with purified GPV. A second hydrophobic domain was located at the C-terminus of the protein suggesting that GPV is a transmembrane protein. This agrees with data showing that GPV was found in the hydrophobic phase of a Triton X-114 phase partition (Bienz et al., *Blood* 68:720–725 (1986)). GPV contains 8 potential N-glycosylation sites, located on the extracellular domain. The presence of O-linked carbohydrates and sialic acid has been suggested based upon a 10 kDa molecular weight reduction following neuraminidase treatment (Zafar and Walz *Thromb. Res.* 53:31–44 (1989)). One short region in the C-terminal region contains two Ser-rich segments and could contain O-linked sugars, but it is probable that the bulk of the carbohydrates are represented by N-sugars due to the observed 20,000 Da apparent molecular weight drop after treatment of GPV by N-glycanase (Zafar and Walz *Thromb. Res.* 53:31–44 (1989)). GPV has a very short intracellular domain which contains no potential phosphorylation site as it lacks any Tyr, Ser, or Thr residues. The C-terminal intracellular domain also lacks an unpaired cysteine residues which is a site for acylation by fatty acids which is found in GPIIb β and GPIIX (Lopez et al., *Proc. Natl. Acad. Sci. USA* 85:2135–2139 (1988); Hickey et al. *Proc. Natl. Acad. Sci. USA* 86:6773–6777 (1989)). Thus, most of the polypeptide chain (92%) is exposed to the outside of the platelet. This is consistent with the observed release of a GPV fragment slightly smaller (80 kDa) than intact membrane bound GPV after treatment of platelets with calpain (Bienz et al., *Blood* 68:720–725 (1986)). This observation shows that the cleavage site for calpain must lie in a region between the last C-terminal N-glycosylation site and the transmembrane domain. The eight cysteine residues are not evenly distributed in the protein: four are clustered in the H-terminal portion, and four are in the region between the Leu-rich domains and the membrane in the C-terminal part of the extracellular segments. The absence of an apparent molecular weight change upon reduction (Berndt and Phillips *J. Biol. Chem.* 256:59–65 (1981)) suggests that all the disulfide bonds are formed over short distances. The absence of cysteines in the middle portion of the molecule indicates that this region is susceptible to enzymatic cleavages accounting for its sensitivity to various enzymes such as calpain, chymotrypsin elastase and thrombin.

Analysis of the peptide sequence for a putative thrombin cleavage site revealed the presence of an Arg-Gly motif at

position 476–477. This appears to be the actual cleavage site based on the following observations: firsts the estimated molecular weight of the fragment liberated by thrombin would be 67,613 Da after correction for the presence of seven N-glycosylation sites, which is similar to the apparent molecular weight of the GPVfl fragment. Second, the amino acid sequence around the Arg-Gly peptide bond displayed significant similarity to sequences around known thrombin cleavage sites (Muszbek and Laki et al., (R. Nachovich, ed) in *The Thrombin* pp 83–90, CRC Press, Boca Raton, Fla. (1984)), and most notably to the α chain of fibrinogen. The sequence is also similar to other thrombin substrates where a high incidence of proline residues occur at the P2 subsite. Finally, the sequence immediately after the RG peptide corresponds to the N-terminal sequence of a peptide obtained after thrombin cleavage of purified GPV (Shimomura et al., *Blood* 75:2349–2356 (1990); Roth et al., *Biochem Biophys. Res. Commun.* 170:153–161 (1990)).

The prior art suggests that GPV has a high affinity binding site for thrombin. The GPVfl fragment is generated at concentrations of thrombin in the nM range: α -thrombin cleaves 100% of platelet GPV at concentrations less than 30 nM (Jandrot-Perrus et al. *Thromb. Haemostas.* 58:915–920 (1987)). In addition, direct interaction of GPV with thrombin was demonstrated by the selective retention of purified GPV on a thrombin-Sepharose column which could then be eluted with heparin (Bienz et al., *Blood* 68:720–725 (1986)). Other examples of platelet proteins known to interact with thrombin with high affinity are the newly cloned thrombin receptor, and GPIIb α (Vu, et al., *Cell* 64:1057–1068 (1991); Lopez et al., *Proc. Nat. Acad. Sci. USA* 84:5614–5619 (1987); De marco et al., *J. Biol. Chem.* 266:23776–23783 (1991)).

A distinctive feature of GPV is that it has the highest leucine content (comprising 20% of the amino acids) of the known platelet proteins. Most of the leucine residues in GPV are contained within 15 tandem Leu-rich repeats of 24 amino acids similar to repeats found in the LRG family of proteins (Roth, *Blood* 77:5–19 (1991)), and most noticeably to platelet GPIIb α (7 LRG repeats), GPIIb β (1 LRG repeat), and GPIIX (1 LRG repeat) (FIG. 8). The LRG domains, at least in some members of the family, mediate protein-protein, cell-cell, or cell-matrix interactions. For example, proteoglycan II (Krusius et al., *Proc. Natl. Acad. Sci. USA* 83:7683–7687 (1986)) and fibromodulin (Hashimoto et al., *Cell* 52:269–279 (1988)) bind to a specific type of collagen, and *Drosophila chaoptin* (Reinke et al., *Cell* 52:291–301 (1988)) and toll (Oldberg et al., *EMBO J.* 8:2601 (1989)) proteins orient cells during morphogenesis and embryogenesis, respectively.

Analysis by the sensitive RT-PCR amplification technique revealed the presence of GPV mRNA in platelets and megakaryocytes. A GPV transcript was also detected in HEL cells which were upregulated after treatment with a phorbol ester which is a known inducer of megakaryocyte differentiation in HEL cells. RT-PCR analysis did not reveal GPV mRNA in non megakaryocytic cells such as leukocytes, endothelial cells, HL60 and U937 cells. Northern analysis revealed a transcript of approximately 4.5 kb in platelets and also revealed a positive band of lower size in lymphocytes. Further analysis is needed to identify the nature of this transcript, but it could represent some related gene revealed by the long exposure times necessary to detect the minute amounts of mRNA present in platelets. The restricted distribution to platelets, coupled to a high sensitivity to thrombin cleavage makes GPV a useful marker for megakaryocytogenesis and for the detection of thrombin dependent platelet activation in thrombotic or prethrombotic states.

The present invention demonstrates that GPV is the product of a single gene. The GPV gene is interrupted by a single intron within the 5'-untranslated region with consensus GT/AG donor and acceptor sites. Several observations show that the isolated genomic clone was derived from the gene for GPV. First, the genomic sequence in exon 2 agrees completely with the cDNA sequence obtained from platelet mRNA. Second, the restriction map of the isolated clone is consistent with restriction fragments identified herein by Southern analysis of human chromosomal DNA. The structure of the GPV gene is very similar to that of the GPIB α gene (Wenger et. al., *Biochem. Biophys. Res. Commun.* 156:389-395 (1988), another platelet member of the LRG family, both have a single intron in the 5'-untranslated sequence and their entire coding sequence is contained within a single exon. The sequence of the GPIX gene was recently reported (Hickey et. al., *J. Biol. Chem.* 268:3438-3443 (1993)), and was shown to contain its entire coding region in a single exon and to have its 50-non coding region interrupted by two introns. The similar exon-intron distribution for the GPV, GPIB α , and GPIX genes suggests that these genes might have a common evolutionary origin within the LRG family of proteins. Analysis of the 5'-flanking region of the GPV gene for cis-acting elements, and comparison to available sequences from other megakaryocyte specific genes revealed significant differences and similarities. Unlike the PF4 (Doi et. al., *Mol. Cell. Biol.* 7:898-904 (1987)), GPIB α (Wenger et. al., *Biochem. Biophys. Res. Commun.* 156:389-395 (1988)), GPIIb (Prandini et. al., *Biochem. Biophys. Res. Commun.* 156:595-601 (1988); Heidenreich et. al., *Biochemistry* 29:1232-1244 (1990)), and GPIX (Hickey et. al., *J. Biol. Chem.* 268:3438-3443 (1993)) genes, the GPV gene contains a perfect consensus sequence for a canonical TATA box which is found in the majority of RNA polymerase II transcribed genes. Similar with the other megakaryocyte specific genes, the GPV gene lacks a CAAT sequences and contains putative binding sites for GATA-1, Ets-11 and Sp1 transactivating factors. Recent experiments support the association of GATA and Ets-1 cis-acting sequences in megakaryocyte-specific gene expression (Lemarchandel et. al., *Mol. Cell. Biol.* 16:668-676 (1993)) while Sp1 sites interact with more ubiquitous transcription factors.

The availability of the genomic sequence for GPV is useful in the characterization of patients with Bernard-Soulier syndrome. These patients are characterized by an absence of or defects in the GPIb-IX glycoprotein complex and the GPV platelet glycoprotein. The availability of the GPV cDNA sequence allows for the assessment of the role of GPV in the correct expression of the four proteins which are deficient in the Bernard-Soulier syndrome. The demonstration of a requirement of GPV for correct and efficient formation of the GPIb-IX complex indicates that a defect in the gene for GPV can cause certain types of Bernard-Soulier syndrome. The availability of the genomic sequence allows for the detection of possible alterations in the GPV gene of such patients.

As used herein the terms "GPV" or "glycoprotein V" refer to polypeptide sequences at least substantially similar to GPV sequence disclosed here. The terms also specifically refer to fragments such as GPVfl as well as the full-length protein. Typically polypeptides will consist of from about 50 to about 560 residues, preferably between about 75 and 500, more preferably between about 100 and about 480 residues. The GPV sequences of the present invention can be readily designed and manufactured utilizing various recombinant DNA techniques well known to those skilled in the art and

described in detail, below. For example, the chains can vary from the naturally-occurring sequence at the primary structure level by amino acid insertions, substitutions, deletions, and the like. These modifications can be used in a number of combinations to produce the final modified protein chain.

The amino acid sequence variants of GPV can be prepared with various objectives in mind, such as facilitating purification and preparation of the protein. The modified molecules are also useful for modifying plasma half life, improving therapeutic efficacy and lessening the severity or occurrence of side effects during therapeutic use. The amino acid sequence variants are usually predetermined variants not found in nature. The variants typically exhibit the same biological activity as naturally occurring GPV, such as the ability to form complexes with GPIb-IX. However, the variants and derivatives that are not capable of binding to ligands are useful nonetheless (a) as a reagent in diagnostic assays for GPV or antibodies to GPVE (b) as agents for purifying anti-GPV antibodies from antisera or hybridoma culture supernatants when insolubilized in accord with known methods, and (c) as immunogens for raising antibodies to GPV or as immunoassay kit components so long as at least one GPV epitope remains active.

In general modifications of the gene encoding the GPV may be readily accomplished by a variety of well-known techniques, such as site-directed mutagenesis (see, Gillman and Smith, *Gene* 8:81-97 (1979) and Roberts, S. et al., *Nature* 328:731-734 (1987), both of which are incorporated herein by reference). One of ordinary skill will appreciate that the effect of many mutations is difficult to predict. Thus, most modifications are evaluated by routine screening in a suitable assay for the desired characteristic. For instance, a change in the immunological character of the GPV can be detected by competitive immunoassay with an appropriate antibody. The effect of a modification on the ability of the GPV to promote platelet aggregation can be tested using in vitro assays, well known to those of skill in the art. Modifications of other properties such as redox or thermal stability, hydrophobicity, susceptibility to proteolysis, or the tendency to aggregate are all assayed according to standard techniques.

Insertional variants of the present invention are those in which one or more amino acid residues are introduced into a predetermined site in the protein and which displace the preexisting residues. For instances insertional variants can be fusions of heterologous proteins or polypeptides to the amino or carboxyl terminus of GPV. Such fusion proteins can be used to facilitate purification of the encoded protein.

Immunogenic fusions may also be produced by cross-linking in vitro or by recombinant cell culture using DNA encoding an immunogenic polypeptide linked to a nucleotide sequence encoding GPV. These immunogenic fusions are useful, for instance, to raise antibodies useful in diagnostics or in purification of GPV by immunoaffinity techniques well known to the skilled artisan.

Substitutional variants are those in which at least one residue has been removed and a different residue inserted in its place. Non-natural amino acid (i.e., amino acids not normally found in native proteins), as well as isosteric analogs (amino acid or otherwise) are also suitable for use in this invention.

Substantial changes in function or immunological identity are made by selecting substitute residues that differ in their effect on the structure of the polypeptide backbone (e.g., as a sheet or helical conformation), the charge or hydrophobicity of the molecule at the target site, or the bulk of the side

chain. The substitutions which in general are expected to produce the greatest changes in function will be those in which (a) a hydrophilic residue, e.g., serine or threonine, is substituted for (or by) a hydrophobic residue, e.g. leucine, isoleucine, phenylalanine, valine or alanine; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysine, arginine, or histidine, is substituted for (or by) an electronegative residue, e.g., glutamine or aspartate; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine.

Substitutional variants of the subunits also include variants in which functionally homologous (having at least about 70% similarity) domains of other proteins are substituted by routine methods for one or more of the GPV domains.

Another class of variants are deletional variants. Deletions are characterized by the removal of one or more amino acid residues from the GPV sequence. Deletions of cysteine or other labile residues also may be desirable for example in increasing the oxidative stability of the protein. Deletion or substitutions of potential proteolysis sites e.g., Arg Arg, is accomplished by deleting one of the basic residues or substituting one by glutaminyl or histidyl residues.

A preferred class of substitutional or deletional variants are those involving the transmembrane region of the protein. Inactivation of the transmembrane domain, typically by deletion or substitution of transmembrane domain hydroxylation residues, will facilitate recovery and formulation by reducing its cellular or membrane lipid affinity and improving its aqueous solubility. Alternatively, the transmembrane and cytoplasmic domains can be deleted to avoid the introduction of potentially immunogenic epitopes. Inactivation of the membrane binding function is accomplished by deletion of sufficient residues (not necessarily all the residues) to produce a substantially hydrophilic hydropathy profile at this site or by substituting with heterologous residues which accomplish the same result.

A principal advantage of the transmembrane inactivated GPV is that it may be secreted into the culture medium of recombinant hosts. This variant is soluble in body fluids such as blood and does not have an appreciable affinity for cell membrane lipids, thus considerably simplifying its recovery from recombinant cell culture. Deletional variants typically substantially lack a transmembrane domain and consist essentially of the effective portion of the extracellular domain of GPV. In some circumstances, the molecule may comprise sequences from the transmembrane region (up to about 10 amino acids), so long as solubility is not significantly affected.

The transmembrane domain may also be substituted by any amino acid sequence, e.g., a random or predetermined sequence of about 5 to 50 serine, threonine, lysine, arginine, glutamine, aspartic acid and like hydrophilic residues, which altogether exhibit a hydrophilic hydropathy profile. Like the deletional (truncated) variants these variants are secreted into the culture medium of recombinant hosts.

Glycosylation variants are included within the scope of this invention. They include variants completely lacking in glycosylation (unglycosylated) and variants having at least one less glycosylated site than the native form (deglycosylated) as well as variants in which the glycosylation has been changed. Included are deglycosylated and unglycosylated amino acid sequence variants, deglycosylated and unglycosylated subunits having the native, unmodified amino acid sequence. For example, substitu-

tional or deletional mutagenesis is employed to eliminate the N- or O-linked glycosylation sites of the subunit, e.g., the asparagine residue is deleted or substituted for by another basic residue such as lysine or histidine. Alternatively, flanking residues making up the glycosylation site are substituted or deleted, even though the asparagine residues remain unchanged, in order to prevent glycosylation by eliminating the glycosylation recognition site. Additionally, unglycosylated subunits which have the amino acid sequence of the native subunits are produced in recombinant prokaryotic cell culture because prokaryotes are incapable of introducing glycosylation into polypeptides.

Glycosylation variants are conveniently produced by selecting appropriate host cells or by in vitro methods. Yeast, for example, introduce glycosylation which varies significantly from that of mammalian systems. Similarly, mammalian cells from a different species (e.g., hamster, murine, insect, porcine, bovine or ovine) or tissue than the GPV source are routinely screened for the ability to introduce variant glycosylation as characterized for example by elevated levels of mannose or variant ratios of mannose, fucose, sialic acid, and other sugars typically found in mammalian glycoproteins. In vitro processing of the subunit typically is accomplished by enzymatic hydrolysis, e.g., neuraminidase digestion.

The polypeptides of the invention can consist of the full length GPV or a fragment thereof as described above. Particularly preferred polypeptides of the invention are those having a polypeptide sequence substantially identical to the sequence disclosed in FIG. 5D.

Two polynucleotides or polypeptides are said to be "identical" if the sequence of nucleotides or amino acid residues in the two sequences is the same when aligned for maximum correspondence. Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.) or by inspection. These references are incorporated herein by reference.

The percentage of sequence identity between two sequences is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity.

The term "substantial identity" means that a polypeptide comprises a sequence that has at least 80% sequence identity, preferably 90%, more preferably 95% or more compared to a reference sequence over a comparison window of about 20 residues to about 500 residues—typically about 50 to about 500 residues usually about 250 to 300 residues. The values of percent identity are determined using the programs above.

Another indication that polypeptide sequences are substantially identical is if one protein is immunologically reactive with antibodies raised against the other protein.

Thus, the polypeptides of the invention include polypeptides immunologically reactive with antibodies raised against GPV.

The present invention provides substantially pure preparation of GPV polypeptides, produced either by recombinant or synthetic means, or isolated from natural sources. The phrases "isolated" or "biologically pure" refer to material which is substantially or essentially free from components which normally accompany it as found in its native state. Thus, the binding domain polypeptides of this invention do not contain materials normally associated with their in situ environment, e.g., other proteins from a platelet membrane. However, even where a protein has been isolated to a homogenous or dominant band by PAGE, there can be trace contaminants in the range of 5–10% of native protein which co-purify with the desired protein. Isolated polypeptides of this invention do not contain such endogenous co-purified protein.

Protein purity or homogeneity may be indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a protein sample, followed by visualization upon staining. For certain purposes high resolution will be needed and HPLC or a similar means for purification utilized.

Much of the nomenclature and general laboratory procedures referred to in this application can be found in Sambrook et. al., *Molecular Cloning—A Laboratory Manual* (2nd Ed.), Vol. 1–3, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1989 or in Berger and Kimmel, *Guide to Molecular Cloning Techniques, Methods in Enzymology* 152 (Academic Press, Inc., San Diego, Calif.). The manuals are hereinafter referred to as "Sambrook" or "Berger" respectively.

Cloning

A variety of methods for cloning DNA sequences into prokaryotic cells are well known in the art. Organisms which are commonly utilized as hosts for the amplification of a vector include *Escherichia*, *Bacillus* and *Streptomyces*. The most common bacterial hosts are various commercially available strains of *E. coli*, due to the ease with which the organism may be cultured and the wealth of information which is available regarding the cell's life-cycle, genetics, viruses and developmental regulation. The vectors most commonly used in *E. coli* are those derived from the pBR322 plasmid and those derived from λ or M13 phage, although several vectors unrelated to any of these are also common. The Sambrook and Berger manuals contains methodology sufficient to direct persons of skill through most cloning exercises.

A number of vectors detailed in Sambrook and elsewhere may be initially cloned into *E. coli* and then subsequently transferred into a eukaryotic system without any necessity for re-cloning that part of the vector which is of interest to the person of skill. Vectors capable of replication in both prokaryotic and eukaryotic cells are generally termed "shuttle vectors" and must contain at a minimum a eukaryotic and a prokaryotic origin of replication. Several shuttle vectors are commercially available which contain polycloning sites, selectable markers for both bacterial and eukaryotic cells, promoters for both bacterial and eukaryotic expression of the gene(s) of interest, and integration sequences for insertion of the vector into the eukaryotic genome. A few examples of vectors which may be amplified in bacteria and used for transformation in eukaryotic cells include the family of P element vectors for *Drosophila*

melanogaster, a number of SV40-derived vectors for the transformation of COS cells, adenovirus-derived vectors for transformation in cells containing the appropriate transcription factor for RNA polymerase III, a variety of BPV-derived vectors and the YIp5-derived vectors of *Saccharomyces cerevisiae* (see Sambrook chapter 16 and Berger chapter 53 for an overview of different vectors which may be transferred between *E. coli* and eukaryotes). General techniques for shuttling DNA between prokaryotes and eukaryotes are also described in Cashion et. al., U.S. Pat. No. 5,017,478 and Kriegler, *Gene Transfer and Expression: A Laboratory Manual*, W. H. Freeman, N.Y., (1990) which are incorporated by reference.

Expression of Recombinant Proteins

Methods for expression of recombinant proteins may be found in Sambrook chapters 16 and 17.

Recombinant proteins may be expressed in either bacteria such as *E. coli* or in eukaryotic expression systems. In general, it is often necessary to express membrane proteins in eukaryotic systems to achieve proper post-translational modification of the proteins although it is sometimes possible to engineer the biologically active fragment of a polypeptide into an appropriate bacterial expression system, or to use the bacterial system for generating peptides which may be used for antibody generation. In these prokaryotic hosts, one can make expression vectors which will typically contain expression control sequences compatible with the host cell (e.g., an origin of replication) In addition, a variety of well-known promoters or promoter elements will be present, such as the lactose promoter system, a tryptophan (trp) promoter system, a beta-lactamase promoter system, or a promoter system from phage lambda (see Yanofsky, C., 1984, *J. Bacteriol.*, 158:1018–1024 and Herskowitz, I. and Hagen, D., 1980, *Ann. Rev. Genet.*, 14:399–445). The promoters will typically control expressions optionally with an operator sequence, and have ribosome binding site sequences and similar elements for initiating and completing transcription and translation.

Methods for expressing large amounts of a protein in a bacterial cell are often invaluable in determining the protein's function, or in generating simple methods of purifying a proteins such as by raising antibodies to a protein expressed in a bacterial cell for use in an immunopurification technique for isolation of a protein from a eukaryotic cell. During purification from *E. coli*, the expressed polypeptides may first be denatured and then renatured. This can be accomplished by solubilizing the bacterially produced proteins in a chaotropic agent such as guanidine HCl and reducing all the cysteine residues with a reducing agent such as beta-mercaptoethanol. The polypeptides are then renatured, either by slow dialysis or by gel filtration, U.S. Pat. No. 4,511,503. The most common of these techniques is the generation of fusion proteins which express a portion of the protein of interest fused to a known antigen which is not otherwise present in the bacterial cell (e.g., LacZ in *E. coli*), but for which antibodies are readily available. After purification using immunopurification methods directed against the known antigen, the fusion protein is used to raise antibodies via standard techniques.

Expression of genes in eukaryotic systems may be used for a number of purposes, including the following: to confirm the identity of a cloned gene, to express eukaryotic genes which require post-translational modifications to produce large quantities of proteins which are ordinarily available in small quantities from naturally-occurring biological

sources, to study the biosynthetic pathway of the gene product, to clarify the relationship between the structure and function of a protein through mutational analysis, to properly express proteins containing introns which prokaryotes cannot process, and to identify the gene's promoter elements. When choosing an expression vector several factors need to be taken into account including the size of the gene (some packaging viruses may incorporate only relatively small amounts of DNA), the type of host cell which is available (some cells such as CHO cells add more post-translational modifications than other cells such as NIH-3T3 cells), whether a permanent transformant or a transient expression system is desired and the presence of control elements in the vector. Eukaryotic expression vectors contain both prokaryotic origins of replication (generally derived from pBR322) and eukaryotic transcription units which are transcribed only in eukaryotes. The eukaryotic transcription unit consists of non-coding sequences and sequences coding for selectable markers such as thymidine kinase, aminoglycoside phosphotransferase or dihydrofolate reductase, as well as the portion of the gene of interest necessary for expression. In general the transcription unit is assembled from well-characterized viral or eukaryotic genes.

Introduction of the recombinant vectors into eukaryotic cells may be achieved by a variety of methods known in the art, including calcium phosphate or DEAE-mediated transfection, polybrene, protoplast fusion, electroporation, liposomes and direct microinjection.

Common vectors for mammalian replication systems include the Simian virus SV40e papilloma viruses such as bovine papilloma virus (BPV) and herpes viruses such as Epstein-Barr (EBV). Each of these vectors may be used to generate cell lines which contain multiple copies of the gene of interest. Cell lines with high levels of expression of the introduced gene may be selected by treating the cells with gradually increasing amounts of the toxin which the selectable marker provides resistance against. The DNA unit which is amplified under selective conditions is variable, but generally includes a substantial amount of flanking DNA, particularly in stably transfected lines in which the vector has integrated into the chromosome.

The DNA sequences will be expressed in hosts after the sequences have been operably linked to an expression control sequence. These expression vectors are typically replicable in the host organisms either as episomes or as an integral part of the host chromosomal DNA. Commonly, expression vectors will contain selection markers, e.g., tetracycline or neomycin, to permit detection of those cells transformed with the desired DNA sequences (see, e.g., U.S. Pat. No. 4,704,362, which is incorporated herein by reference).

Many standard purification techniques may be used to purify the gene product from the gene of interest which is expressed as described above. In the present invention which provides a direct means for antibody generation, it is possible to use an immunoprecipitation or immunochromatographic method in addition to or in conjunction with standard precipitation and chromatographic methods for purification of GPV or its cleavage products, without first generating antibodies using prokaryotic fusion proteins.

As indicated above, the vectors e.g., a plasmid, which is used to transform the host cell, preferably contains DNA sequences to initiate transcription and sequences to control the translation of the antigen gene sequence. These sequences are referred to as expression control sequences.

When the host cell is of insect or mammalian origins illustrative expression control sequences are obtained from the SV-40 promoter (*Science*, 222:524-527, 1983), the CMV I.E. Promoter (*Proc. Natl. Acad. Sci.* 81:659-6631, 1984) or the metallothionein promoter (*Nature* 296:39-42, 1982). The cloning vector containing the expression control sequences is cleaved using restriction enzymes and adjusted in size as necessary or desirable and ligated with DNA coding for the GPV polypeptide by means well known in the art.

As with yeast, when higher animal host cells are employed, polyadenylation or transcription terminator sequences from known mammalian genes need to be incorporated into the vector. An example of a terminator sequence is the polyadenylation sequence from the bovine growth hormone gene. Sequences for accurate splicing of the transcript may also be included. An example of a splicing sequence is the VP1 intron from SV40 (Sprague, J. et al., 1983, *J. Virol.* 45:773-781).

Additionally, gene sequences to control replication in the host cell may be incorporated into the vector such as those found in bovine papilloma virus type-vectors. Saveria-Campo, M., 1985, "Bovine Papilloma virus DNA a Eukaryotic Cloning Vector" in *DNA Cloning Vol. II a Practical Approach* Ed. D. M. Glover, IRL Press, Arlington, Va. pp. 213-238.

The host cells are competent or rendered competent for transformation by various means. There are several well-known methods of introducing DNA into animal cells. These include: calcium phosphate precipitation, fusion of the recipient cells with bacterial protoplasts containing the DNA, treatment of the recipient cells with liposomes containing the DNA, DEAE dextran, electroporation and microinjection of the DNA directly into the cells.

The transformed cells are cultured by means well known in the art. *Biochemical Methods in Cell Culture and Virology*, Kuchler, R. J., Dowden, Hutchinson and Ross, Inc., (1977). The expressed GPV polypeptides are isolated from cells grown as suspensions or as monolayers. The latter are recovered by well known mechanical, chemical or enzymatic means.

Production of GPV Peptides by Protein Chemistry Techniques

The polypeptides of the invention can be synthetically prepared in a wide variety of ways. For instance polypeptides of relatively short size can be synthesized in solution or on a solid support in accordance with conventional techniques. Various automatic synthesizers are commercially available and can be used in accordance with known protocols. See, for example, Stewart and Young, *Solid Phase Peptide Synthesis*, 2d. ed., Pierce Chemical Co. (1984). The peptides may be used to generate antibodies using standard methods, including those methods described in this application.

Alternatively, purified and isolated GPV may be treated with proteolytic enzymes in order to produce GPV polypeptides. The GPV protein sequence may be analyzed to select proteolytic enzymes to be used to generate polypeptides containing desired regions of the GPV protein. The desired polypeptides are then purified by using standard techniques for protein and peptide purification. For a review of standard techniques see, *Methods in Enzymology*, "Guide to Protein Purification", M. Deutscher, ed. Vol. 182 (1990), pages 619-626, which is incorporated herein by reference. Peptides generated by this strategy may be used to generate

antibodies using standard methods, including those described in this application.

Antibody Generation

The antibodies recognizing polypeptides of the present invention are suitable for modification using the multitude of techniques available to those skilled in the art for production and manipulation of various immunoglobulin molecules. Immunoglobulins are proteins consisting of one or more polypeptides substantially encoded by immunoglobulin genes. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as the myriad immunoglobulin variable region genes. For a discussion of immunoglobulin forms, see e.g., *Fundamental Immunology*, 2d Ed. W. E. Paul ed, Raven Press NY (1989), Huston et al., *Proc. Nat. Acad. Sci. USA* 85:5879-5883 (1988), Bird et al., *Science* 242:423-426 (1988), and Hunkapiller and Hood, *Nature* 323:15-16 (1986).

As used herein, "immunoglobulin," "antibody" or "antibody peptide(s)" refers to polyclonal antibodies, monoclonal antibodies, to an entire immunoglobulin or antibody or any functional fragment of an immunoglobulin molecule which binds to the target antigen. Examples of such peptides include complete antibody molecules, antibody fragments, such as Fab, F(ab')₂, complementarity determining regions (CDRs), V_L (light chain variable region), V_H (heavy chain variable region), and any combination of those or any other functional portion of an antibody peptide.

An F(ab')₂ fragment lacks the C-terminal portion of the heavy chain constant region, and has a molecular weight of approximately 110 kD. It retains the two antigen binding sites and the interchain disulfide bonds in the hinge region, but it does not have the effector functions of an intact IgG molecule. An F(ab')₂ fragment may be obtained from an IgG molecule by proteolytic digestion with pepsin at pH 3.0-3.5 using standard methods such as those described in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Pubs., N.Y. (1988).

An Fab fragment comprises a light chain and the N-terminus portion of the heavy chain to which it is linked by disulfide bonds. It has a molecular weight of approximately 50 kD and contains a single antigen binding site. Fab fragments may be obtained from F(ab')₂ fragments by limited reduction, or from whole antibody by digestion with papain in the presence of reducing agents. (See, Harlow and Lane, supra.)

A multitude of techniques available to those skilled in the art for production and manipulation of various immunoglobulin molecules can be readily applied to produce antibodies for use in the present invention. Antibodies which bind to GPV may be produced by a variety of means. The production of non-human monoclonal antibodies, e.g., murine lagomorpha, equine, etc., is well known and may be accomplished by, for example, immunizing the animal with a preparation containing cells bearing GPV or isolated GPV molecules. Antibody-producing cells obtained from the immunized animals are immortalized and screened, or screened first for the production of antibody which binds to GPV and then immortalized. For a discussion of general procedures of monoclonal antibody production see Harlow and Lane, supra.

The generation of human monoclonal antibodies to a human antigen is also known in the art. Generation of such human monoclonal antibodies may be difficult with conventional techniques. Thus, it may be desirable to isolate DNA

sequences which encode an anti-GPV human monoclonal antibody (or portions thereof) by screening a DNA library from human B cells according to the general protocol outlined by Huse et al., *Science* 246:1275-1281 (1989). The sequences which encode the antibody (or binding fragment) of the desired specificity are then cloned and amplified. Alternatively, one may transfer the antigen binding regions of non-human antibodies, e.g., the F(ab')₂ or hypervariable regions, to human constant regions (Fc) or framework regions by recombinant DNA techniques to produce substantially human molecules. Such methods are generally known in the art and are described below.

The invention also provides synthetic or recombinant immunoglobulins, including chimeric immunoglobulins, humanized antibodies or hybrid antibodies or derivatives of any of those. Chimeric immunoglobulins are typically the product of chimeric DNA, which is recombinant DNA containing genetic material from more than one eukaryotic species.

"Chimeric immunoglobulins" or "chimeric antibodies" refer to those antibodies or antibody peptides wherein one portion of the peptide has an amino acid sequence that is derived from, or is homologous to, a corresponding sequence in an antibody or peptide derived from a first gene source, while the remaining segment of the chain(s) is homologous to corresponding sequences of another gene source. For example, a chimeric antibody peptide may comprise an antibody heavy chain with a murine variable region and a human constant region. The two gene sources will typically involve two species, but will occasionally involve different sources from one species.

Chimeric antibodies or peptides are typically produced using recombinant molecular and/or cellular techniques. Typically, chimeric antibodies have variable regions of both light and heavy chains that mimic the variable regions of antibodies derived from one mammalian species, while the constant portions are homologous to the sequences in antibodies derived from a second different mammalian species. Methods for production of such antibodies are well known and are described in, for example, U.S. Pat. No. 4,816,397, and EP publications 173,494 and 239,400, which are incorporated herein by reference.

The definition of a chimeric immunoglobulin, however, is not limited to this example. A chimeric antibody is any antibody in which either or both of the heavy or light chains are composed of combinations of sequences mimicking the sequences in antibodies of different sources, whether these sources are differing classes, differing antigen responses, or differing species of origin, and whether or not the fusion point is at the variable/constant boundary.

The term "humanized" or "human-like immunoglobulin" refers to an immunoglobulin comprising a human-like framework region and a constant region that is substantially homologous to a human immunoglobulin constant region. Hence, most parts of a human-like immunoglobulin, except possibly the CDRs are substantially homologous to corresponding parts of one or more native human immunoglobulin sequences.

"Hybrid antibody" refers to an antibody wherein each chain is separately homologous with reference to a mammalian antibody chain, but the combination represents a novel assembly so that two different antigens are recognized by the antibody. In hybrid antibodies, one heavy and light chain pair is homologous to a pair found in an antibody raised against another epitope. This results in the property of multi-functional valency, i.e., ability to bind at least two

different epitopes simultaneously. Such hybrids may, of course, also be formed using chimeric chains.

Immunoglobulins may be fused to functional regions from other genes (e.g., those encoding enzymes) to produce fusion proteins (e.g., immunotoxins) having novel properties. In general, modifications of the genes may be readily accomplished by a variety of well-known techniques, such as site-directed mutagenesis (see, Gillman and Smith, *Gene*, 8:8197 (1979) and Roberts, S. et al, *Nature*, 328:731-734 (1987)).

For this invention, an immunoglobulin is specific for, or reactive with, a GPV molecule if the immunoglobulin binds GPV as measured or determined by standard antibody-antigen assays, for example, competitive binding assays, saturation assays, or standard immunoassays such as ELISA or RIA. This definition of specificity applies to single heavy and/or light chains, CDRs, fusion proteins or fragments of heavy and/or light chains, that are also specific for GPV if they bind GPV alone or if, when properly incorporated into immunoglobulin conformation with complementary variable regions and constant regions as appropriate, are then capable of binding GPV. Binding affinity is typically represented by the affinity constant (K_a) for equilibrium concentrations of associated and disassociated configurations, i.e., $K_a = [A-B]/[A][B]$ where $[A]$, $[B]$, and $[A-B]$ are the concentrations at equilibrium of the antibody (A), antigen (B) and antigen-antibody complex (A-B), respectively. Under physiological conditions, the affinity constant of a specific immunoglobulin of the present invention is typically about 10^{-3} to about 10^{-12} liters/mole, and preferably about 10^{-10} liters/mole or more. One of skill will recognize, however, that binding affinity between two molecules will be influenced by a number of factors such as temperature, pH, ionic strength, and the like.

Compositions of the present invention comprise immunoglobulins which selectively bind GPV molecules on platelet cells. The immunoglobulins and pharmaceutical compositions of this invention are particularly useful for parenteral administration, i.e., subcutaneous, intramuscular, or intravenous administration. A number of new drug delivery approaches are being developed, and the pharmaceutical compositions of the present invention are suitable for administration using these new methods, as well. See, Langer, *Science*, 249,1527-1533 (1990).

In one embodiment, the antibodies of the present invention can be used to target conventional drugs or other agents to platelets. By using an antibody to target a drug to cells bearing GPV, such drugs can achieve higher concentrations at sites of platelet aggregation. The immunoglobulins can be directly or indirectly coupled to the chemotherapeutic agent.

The antibodies of the present invention may also be used for diagnostic purposes, such as identifying areas of platelet aggregation. For diagnostic purposes, the antibodies may either be labeled or unlabeled. Unlabeled antibodies can be used in combination with other labeled antibodies (second antibodies) that are reactive with the antibody, such as antibodies specific for a particular immunoglobulin constant region. Alternatively, the antibodies can be directly labeled. A wide variety of labels may be employed, such as radionuclides, fluorescers, enzymes, enzyme substrates, enzyme cofactors, enzyme inhibitors, ligands (particularly haptens), etc. Numerous types of immunoassays are available and are well known to those skilled in the art.

In diagnostic applications, compositions containing the immunoglobulins or a cocktail thereof, are administered to a patient suspected of having a defect in platelet functions

Alternatively, the efficacy of a particular treatment can be monitored. An amount sufficient to accomplish this is defined to be a "diagnostically effective dose." In this use, the precise amounts will depend upon the patient's state of health and the binding constants for the specific antibodies employed.

Kits can also be supplied for use with the subject antibodies. Thus, the subject antibody composition of the present invention may be provided, usually in a lyophilized form in a container, either alone or in conjunction with additional antibodies specific for the desired cell type. The antibodies, which may be conjugated to a label or toxin, or unconjugated, are included in the kits with buffers, such as Tris, phosphate, carbonate, etc., stabilizers, biocides, inert proteins, e.g., serum albumin, or the like, and a set of instructions for use. Generally, these materials will be present in less than about 5% wt. based on the amount of active antibody, and usually present in total amount of at least about 0.001% wt. based again on the antibody concentration. Frequently, it will be desirable to include an inert extender or excipient to dilute the active ingredients, where the excipient may be present in from about 1 to 99% wt. of the total composition. Where a second antibody capable of binding to the chimeric antibody is employed in an assays this will usually be present in a separate vial. The second antibody is typically conjugated to a label and formulated in an analogous manner with the antibody formulations described above.

PCR

The use of PCR (Hullis et. al., U.S. Pat. No. 4,683,202 (1987)) in the field of molecular biology is well known. PCR has been adapted for many diverse purposes including cloning, sequencing, forensics, diagnostics and clastic analysis. The technique is detailed in several general sources which provide adequate guidance to one of skill to perform the technique, including Sambrook and *PCR Protocols: A Guide to Methods and Applications* (Innis et. al. eds) Academic Press Inc. San Diego, Calif. (1990) (hereinafter "Innis").

The following PCR protocol is provided as a starting point for a person of skill, who will readily recognize that it is usually necessary to optimize reaction conditions, and that optimization is especially important when the technique yields inadequate results, or when an essentially repetitive task utilizing the technique needs to be performed. In general, a 100 μ l reaction contains the following: 1 to 1×10^7 target molecules (generally about 1×10^5 to 1×10^6 target molecules); 1 pmol-100 pmol of each primer (generally about 20 pmol), the primer having a T_m of from about 30° C. to about 70° C. (preferably greater than about 50° C.); 20 mM Tris-Hcl (pH approximately 8.3 at 20° C.); 0.2 mM-5 mM $MgCl_2$ (generally about 1.5 mM $MgCl_2$; occasionally it may be helpful to substitute some of the $MgCl_2$ with $MnCl_2$); 25 mM KCl; 0.05% Tween 20; 100 μ g autoclaved gelatin or nuclease-free bovine serum albumin; 5-200 μ M of each dNTP (generally about 50 μ M of each dNTP), and from 0.25 to 5 units (generally about 2 units) of taq DNA polymerase. Many practitioners prefer to add an oil phase on top of the aqueous phase to prevent evaporation of the reaction mixture and to prevent the reaction components from being distributed unevenly in the reaction tube upon heating. The reaction mixture is cycled through 15-65 (usually 20-35) of the following temperature variations (generally using a commercially available thermal cycler, occasionally performed by hand with 3 temperature baths): "denaturation" at 96° C. for 0.25 min (on the first cycle it

is often better to leave the reaction mixture at 96° C. for 1–5 minutes), “primer annealing” at a temperature about 5° C. to 10° C. lower than the calculated T_m for 30 seconds, “primer extension” at 72° C. for 1–3 minutes depending on the length of the target sequence to be amplified. Cycling is generally concluded with a final 72° C. extension for about 5 minutes and the reaction is stopped by chilling the reactants to about 4° C. and/or by the addition of EDTA in an amount approximately 8-fold greater than the quantity of $MgCl_2$ plus $MnCl_2$ plus any other divalent cation in the mixture.

Once the complete sequence of a gene is known, it is entirely straightforward to design PCR experiments to detect the presence of abnormalities in the structure of the gene in individual organisms. After performing PCR on the individual's DNA or cDNA using primers designed from the known GPV gene sequence, the presence of gross defects in the gene or the gene's cDNA may be ascertained by standard agarose gel electrophoresis of restriction endonuclease-digested fragments of the DNA or cDNA. If desirable, all of the structural elements of a defective gene may be determined, either by direct sequencing of the PCR product, or by subcloning the PCR product into a sequencing vector for sequencing by standard methodologies; commercially available sequencing kits from a variety of sources are available for the sequencing of either PCR-generated or cloned DNA. In the present invention, the GPV gene may be examined in patients in order to diagnose Bernard-Soulier syndrome, or in order to determine any genetic predisposition towards the Bernard-Soulier syndrome in persons which may be at risk for the disease. The ability to detect genetic diseases in utero using PCR amplification of DNA from a developing fetus is also known in the art and it will be possible to detect abnormalities in the GPV gene using standard PCR methodologies. In addition, the GPV gene is a specific marker for cells of the megakaryocytic lineage and is generally useful as a morphological marker for tracking platelet development.

Southern Analysis and Northern Analysis

Southern analysis of genomic DNA and northern analysis of RNA using a cloned probe are basic to the art of molecular biology. Sambrook provides adequate guidance to perform most commonly used Southern and northern techniques including analysis of genomic DNA, mRNA and cDNA. The present invention provides an array of probes that may be constructed from the GPV gene for use in Southern analysis. These include synthetic oligonucleotide probes generated from the sequence of any region of the GPV gene, probes generated from cleavage products of the cloned gene using random-primer or terminal phosphate labeling methods and several other methods known to persons of skill. The probes may be used for a variety of purposes including isolation of homologous genes from other species by screening genomic or expression libraries or performing PCR, the identification of restriction fragment length polymorphism and the identification of tissues which express the GPV gene using in situ or northern analysis.

Pharmaceutical Composition and Therapeutic Uses of GPV Polypeptides

The recombinant proteins of the present invention may be used to boost the level of GPV and its cleavage products in a patient which exhibits a lower than normal level of the GPV or GPVfl, or at a specific site such as a wound in a patient with normal clotting. The pharmaceutical composi-

tions of the GPV gene are intended for parenteral, oral, topical, or local administration. For the pharmaceutical compositions which are administered parenterally (e.g. intravenously, subcutaneously, intradermally, or intramuscularly), the invention provides compositions for parenteral administration that comprise a solution of the GPV polypeptide isolated from an expression system as described above dissolved or suspended in an acceptable aqueous carrier. A variety of aqueous carriers may be used, e.g., water, buffered water, 0.4% saline, 0.3% glycine, hyaluronic acid and the like. These compositions may be sterilized by conventional, well known sterilization techniques or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile solution prior to administration. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, etc. For solid compositions, conventional nontoxic solid carriers may be used which include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium, saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like.

For oral administration, a pharmaceutically acceptable nontoxic composition is formed by incorporating any of the normally employed excipients, such as those carriers previously listed, and generally 10–95% of active ingredient and more preferably at a concentration of 25%–75%.

For aerosol administration, the GPV polypeptide(s) are preferably supplied in finely divided form along with a surfactant and propellant. The surfactant must be nontoxic, and preferably soluble in the propellant. Representative of such agents are the esters or partial esters of fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, oles-
teric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride. Mixed esters, such as mixed or natural glycerides may be employed. A carrier can also be included, as desired, as with, e.g., lecithin for intranasal delivery.

For topical administration, e.g., to a wound, the GPV polypeptides may be administered in an aqueous solution as described above, or may be applied in a salve composed of non-toxic carrying agents such as completely polymerized polyacrylamide, or long-chain esters or partial esters of fatty acids as described above. Additionally, the solution or salve may be applied to bandages in standard wound-dressings.

In therapeutic applications, GPV polypeptides are administered to a patient in an amount sufficient to affect platelet aggregation, an amount which constitutes a “therapeutically effective dose.” Amounts effective for this use will depend on several factors including the particular polypeptide, the manner of administration, the weight and general state of health of the patient, the presence of other blood disorders in the patient, the presence or severity of a wound and the judgment of the prescribing physician. This will typically be between about 1 $\mu g/kg$ and about 100 mg/kg , preferably about 3 mg/kg to about 15 mg/kg .

Gene Therapy

Strategies for gene therapy are reviewed in Friedmann, *Science* 2440:1275 (1989), which is incorporated herein by reference.

Delivery of the polynucleotide of interest may be accomplished *in vivo* by administering the therapy vector to an individual patient, typically by systemic administration (e.g., intravenous, intraperitoneal, intramuscular, subdermal, or intracranial infusion). Alternatively, the vector may be used to deliver polynucleotides to cells *ex vivo* such as cells explanted from an individual patient or universal donor hematopoietic stem cells, followed by reimplantation of the cells into a patient, usually after selection for cells which have incorporated the gene for the polynucleotide.

The vector may be used for gene therapy to treat congenital genetic diseases, acquired genetic diseases (e.g., cancer), viral diseases or to modify the genome of selected types of cells of a patient for any therapeutic benefit. A treatable disorder using the GPV gene of the present invention is the Bernard-Soulier syndrome. Polynucleotides which reverse or suppress the neoplastic phenotype (e.g. antisense inhibition of defective GPV expression) may be used to treat defective GPV expression as well as engineering the normal GPV gene into patients.

The following examples are offered by way of illustration, not limitation.

EXAMPLES

The following examples are offered by way of illustration and do not limit the claims. It will be apparent to one of skill that many of the experimental parameters may be altered.

Materials: The following applies to materials used in the Examples below: Restriction endonucleases, modifying enzymes, and the M13 cloning vector were purchased from Boehringer, Mannheim, Germany. The pBluescript KSII vector was obtained from Stratagene, San Diego, Calif. The Gene Clean II kit was from Bio 101, La Jolla, Calif. Radiolabeled nucleotides, Hybond N⁺ membranes and Hyperfilm X-ray films were obtained from Amersham Corp., Les Ulis, France. Nitrocellulose membranes were from Schleicher and Schuell, Ecquevilly, France. Synthese Oligonucleotides were obtained from the Service de Synthèse des oligonucleotides, INSERN U 184, LGME, Strasbourg, France or synthesized on a Beckmann Oligo 1000 oligonucleotide synthesizer (Beckmann, Gagny, France). All reagents were molecular biology grade.

Example 1

cDNA Cloning of Human Platelet Glycoprotein v via PCR Amplification of Platelet and Megakaryocyte cDNA

In order to clone the cDNA encoding glycoprotein v via PCR, a series of degenerate primers were designed based on published partial peptide sequences (Shimomura, et. al., *Blood* 75:2349-2356 (1990)) obtained from purified platelet GPV. Fresh human platelets were isolated and platelet total RNA was prepared according to previously described procedures (Lanza et. al., *J. Clin. Invest.* 89:1995-2004 (1992) and Wicki et. al., *Thromb. Haemostas* 61:448-453). Megakaryocyte RNA kindly provided by Dr. Nelly Kieffer, Laboratoire Franco-Luxembourgeois de Recherche Biomedicale, Luxembourg, was from a patient suffering from megakaryoblastic leukemia. Platelet or megakaryocyte poly^a RNA was used to synthesize cDNA with a commercial kit (Boehringer Mannheim). First strand synthesis was performed by priming with oligo dT or by priming with degenerate or exact primers specific for GPV and extending with 20 units of M-MLV reverse transcriptase (Gibco-BRL, Cergy Pontoise, France). Approximately 25 ng of platelet or

megakaryocyte cDNA was used in the PCR amplification reaction using a Gene Amp DNA amplification reaction kit (Perkin-Elmer Cetus, St. Quentin, France), a 0.2 μM concentration of each nucleotide primer, and 1 unit of Taq polymerase. The cDNA was denatured at 94° C. for 4 min, and amplification was performed for 30 cycles with extension at 72° C. for 2 min, denaturation at 94° C. for 1 min, and primer annealing between 45 to 60° C. for 1 min depending on the primers used. Degenerate primers 1 and 4 based on peptide sequence K5/6 and running on opposite strands were used successfully to amplify a 108 bp fragment (fragment i) from oligo dT primed platelet cDNA. Sequence analysis revealed that the cDNA fragment contained within primers 1 and 4 coded for a 20 amino acid peptide corresponding exactly to the published peptide sequence (amino acid residues 13 to 33). This demonstrated that the amplified fragment corresponded to GPV cDNA. In order to obtain additional cDNA sequence, exact oligonucleotide primers were generated in the (-) strand (primer 3) and in the (+) strand (primer 2) orientation. An additional 150 bp cDNA fragment (fragment ii) was obtained using primer 3 and degenerate primer 5 based on the M6 peptide sequence. Following PCR, 10 μl of the amplification mixture was analyzed on a 1 to 2% agarose gel.

Rapid amplification of cDNA ends (RACE) was used to extend the sequence in the 5'- and 3'-direction (Frohman, et. al., *Proc Natl. Acad. Sci. USA.* 85:8998-9002 (1988)). Using the 31-RACE procedure two additional overlapping fragments (iii and iv) covering 703 bp of cDNA were obtained in the 3'-direction. For the 3'-RACE, 25 ng of cDNA were subjected to a first round of PCR with the Adaptor-dTI7 primer and (+) strand primer 2 or 6, followed by a second PCR with the adaptor and primer 2 (SEQ.ID.NOS. 8-9) or 6 (SEQ.ID.NO. 16) (see Table 1 for a description of the primers). The 5'-RACE procedure using (-) strand primers 7 and 8 generated two more fragments of 260 and 150 bp (v and vi) in the 5'-direction. For the 5'-RACE, cDNA was prepared from 1 μg of RNA using a (-) strand specific primer and was detailed by incubation with 5 μM DGTP and 50 units of terminal transferase (Boehringer Mannheim) at 37° C. for 10 min in the buffer supplied by the manufacturer. After phenol-chloroform extraction, the reaction mixture was dialyzed over a Centricon 30 column (Amicon, Beverly, Mass.) and used in the PCR reaction. A first round of PCR was performed with the Adaptor-dC12 primer and primers 7 or 8 followed by a second round of PCR with the adaptor alone and primers 7 or 8.

The positive fragments obtained from regular PCR or from the RACE approach were end-cleaved with restriction enzymes (usually Eco RI and Sal I), isolated by electrophoresis on Sea Plaque agarose (FMC Bioproducts, Rockland, Me.), purified using the Gene Clean II kit, and subcloned into the M13 or pBluescript vectors. The inserts were sequenced using the Sequenase kit (United States Biochemical Corp., Cleveland, Ohio) with DATP 5'-α-[³⁵S]-thiophosphate. All the fragments obtained by the PCR approach were analyzed by sequencing on both strands and their identity to GPV was assessed by comparison to the published GPV partial sequences.

Using the strategy described above we were able to assemble 1,199 bp of GPV cDNA from 6 fragments amplified from platelet mRNA. Sequence analysis revealed the presence of 31 bp of 5' untranslated sequence followed by a 1,168 bp open reading frame starting with a methionine and coding for a total of 389 amino acids.

Example 2

Southern Analysis of Human Chromosomal DNA

In order to determine the complexity of the human GPV gene, a Southern blot analysis was performed under high

23

stringency on human chromosomal DNA using a 748 bp cDNA probe corresponding to the coding region. High molecular weight human leukocyte DNA was digested to completion with restriction endonucleases and subjected to electrophoresis on 0.7% agarose gels. The fragments were transferred to a Hybond N⁺ nylon membrane and were hybridized to a 748 bp ³²P-labelled GPV cDNA fragment at 45° C. overnight. The hybridization buffer was 50% (v/v) formamide, 0.9 M NaCl, 50 mM NaH₂PO₄, 2 mM EDTA, 1% (w/v) SDS, 5% (w/v) dextran sulfate, 0.02% (w/v) polyvinylpyrrolidone, 0.02% (w/v) Ficoll 400, and 50 µg/ml salmon sperm DNA. Membranes were washed in 0.5×SSC, 1% (w/v) SDS at 60° C. and autoradiographed.

Single positive bands of approximately 5 kb, 3.5 kb, and 12 kb were observed when the DNA was cut with Eco RI, Bam HI, and Bgl II restriction endonucleases, respectively (FIG. 4). Analysis of additional individuals revealed an extra polymorphic Bgl II band of 3.4 kb. This simple hybridization pattern was suggestive of a single copy gene of low complexity.

Example 3

Northern Analysis of Platelet mRNA

In order to determine the size of the platelet GPV mRNA, Northern blot analysis was performed on human total platelet RNA using a 748 bp cDNA probe corresponding to the coding region (FIG. 3). Total RNA (10 µg) from platelets (leukocyte contamination <10⁻⁷) or monocytes was electrophoresed on 1% formaldehyde-agarose gel, and transferred to a Zetaprobe (Biorad) membrane. A 748 bp GPV cDNA probe (50 ng) was labelled with [α -³²P]-dCTP using the random prime labelling method and cleaned on a Nick column (Pharmacia) (incorporation [α -³²P]-dCTP was 71%). The hybridization conditions were 0.5 M NaH₂PO₄, pH 7.2, 1 mM EDTA, and 7% SDS at 65° C.

Analysis of the gel revealed a single transcript of approximately 4.5 kb. A partially degraded RNA revealed a more complex pattern. A transcript of less than 2 kb was also revealed with monocyte RNA.

Example 4

Isolation and Characterization of GPV Genomic Clones

With the 748 bp fragment as a probe 15 genomic clones were isolated from a human fibroblast genomic library in the phage λ Fix vector. Approximately 8×10⁵ recombinants of a human commercial genomic library in the λ Fix vector (Stratagene) were plated on *E. coli* LE392, transferred to nitrocellulose membranes, and probed with a 748 bp ³²p labelled GPV cDNA fragment. The hybridization conditions were 50% (v/v) formamide, 5×SSC, 0.1% (w/v) SDS, 5×Denhardt's medium, and 0.1 mg/ml salmon sperm DNA at 42° C. overnight. The filters were washed in 0.1×SSC, 0.05% (w/v) SDS at 56° C., dried and exposed for autoradiography. Positive clones were subjected to two additional rounds of screening in order to obtain isolated clones. Phage DNA was purified using the liquid lysis procedure. The DNA was digested with EcoRI, separated on a 0.7% agarose gel, transferred to nitrocellulose, and hybridized to the ³²p-labelled GPV cDNA fragment to localize exon containing fragments. The positive fragments were subcloned into the pbluescript vector for further restriction enzyme analysis, and finally subcloned into the M13 sequencing vector. After characterization by restriction endonuclease

24

mapping and Southern blot analysis, clone G5a was chosen for further subcloning, restriction enzyme analysis, and nucleotide sequencing.

A 7.5 kb portion of the G5a clone, shown in FIG. 1 with a partial restriction map, was entirely sequenced on both strands. Comparison with the cDNA sequence obtained by PCR revealed that the 7.5 kb genomic fragment contained the entire 1,198 bp cDNA sequence in two exons (FIGS. 5A-C (SEQ.ID.NO. 1)) separated by a 958 bp intron. Exon 1 contained 29 bp of 5'-untranslated region and exon 2 was composed of 2 bp of 5'-untranslated sequence and 1,168 bp of coding sequence obtained by PCR. Exon 2 contained an additional 512 bp of coding sequence before reaching a TAA stop codon.

The sequence immediately adjacent to the 5'-end of the cDNA (exon 1) was examined for the presence of cis-regulatory elements. The analysis revealed the presence of a sequence which matched the consensus sequence for a TATA box (5'-TATATA-3') characteristic of RNA polymerase II transcribed genes, but did not reveal a consensus sequence for a CAAT box. The TATA box was followed 31 bp downstream by a putative Cap site. An additional sequence (TATAT) with similarity to the TATA box consensus was found at position 1,199. A 5'-AAGATA-3' and a 5'-AGATAG-3' sequence with similarity to the consensus 5'-(AT)GATA(AG)-3' motif for a GATA-1 binding site (Faissst and Mayer *Nucleic Acids Research* 20:3-26 (1992)) were located at position 1,285 and 1,321 respectively. The GATA motif has been found in the promoters and enhancers of all characterized erythroid and megakaryocyte specific genes. Other motifs for cis-acting elements include Ets-1 cis-acting sequences at positions 471 (5'-CAGGAAGT-3'), 493 (5'=GAGGAAGC-3'), 897 (5'-GCATCCTG-3', inverse), 1,178 (5'-ACTTCCC-3', inverse) and, 1,365 (5'-CAGGATGCAA-3') (SEQ.ID.NO 3) (consensus sequence: 5'-(GC)(AC)GGA(AT)G(TC)), and a Sp1 putative binding site at position 1,142 (5'-GGGGTGTGGC-3') (SEQ.ID.NO. 4), (consensus sequence: 5'-(GT)(GA)GGCG(GT)(GA)(CT)-3'). A putative TPA responsive element (TRE)(5'-TGACTGACT-3') was found at position 68. Analysis of 3,348 bp of genomic sequence 3'- of the TAA termination site revealed the presence of putative polyadenylation AATATA sites at positions, 5,610, 6,966, 7,224, and 7,358. Two Alu repetitive sequences (Schmid and Jelinek *Science* 216:1065-1070 (1982)) were located at positions 598-896, and 6,133-6,440.

Nucleotide sequence comparison and assembly was performed using the PC Gene software developed by Intelligenetics Inc., Palo Alto, Calif.

Example 5

Determination of The Primary Amino Acid Structure of GPV

The amino acid sequence of GPV as deduced from its cDNA and genomic sequences is shown in FIG. 5D (SEQ.ID.NO. 2). GPV was found to be composed of 560 amino acids, including a putative 16 amino acid signal peptide, and a putative C-terminal 25 amino acid transmembrane domain. Between the signal peptide and the transmembrane domain, is a sequence of 503 amino acids containing eight potential N-glycosylation sites (NXS, NXT) and eight cysteine residues. The putative transmembrane domain is followed by a 16 residue hydrophilic segment. The carboxy region of the transmembrane domain contains basic residues which are typically found on the cytoplasmic

side of the integral membrane proteins (Sabatini et. al., *J. Cell. Biol.* 92:1-22 (1982)). These features suggest that GPV is a type I integral membrane protein with most of its polypeptide chain located outside the cell (FIG. 8). The predicted molecular weight of the GPV polypeptide after removal of the signal peptide is 59,276 Da. Assuming a weight of 2,500 Da per oligosaccharide moiety, the addition of eight N-linked carbohydrates to the GPV polypeptide backbone would bring the weight to 79,276 Da, close to the reported 82 kDa apparent molecular weight estimated by SDS-PAGE analysis (Berndt and Phillips *J. Biol. Chem.* 256:59-65 (1981); Shimomura et. al., *Blood* 77:2349-2356 (1990); Zafar and Walz *Thromb. Res.* 53:31-44 (1989)).

Analysis of the GPV extracellular sequence revealed the presence of 15 leucine-rich tandem repeats of 24 amino acids (FIG. 6) (SEQ.ID.NOs. 22-36). These repeats are very similar to repeats found in platelet GPIB α , GPIB β and GPIX, and to the 24 amino acid consensus sequence based on the repeats found in other members of the LRG family (SEQ.ID.NO. 37). The last GPV Leu-rich repeat (SEQ.ID.NO. 36) is flanked on its C-terminal side by a sequence (NSWR CDCGL) (SEQ.ID.NO. 5) similar to sequences described at the C-terminal end of Leu-rich domains in other members of the LRG family (Hickey et. al., *Proc. Natl. Acad. Sci. USA* 86:6773-6777 (1989)).

Thrombin-induced cleavage of GPV results in the generation of a soluble fragment (GPVfl) of approximately 69 kDa. At position 476-477 a sequence was found containing an RG dipeptide which represents a potential cleavage site for thrombin (Stubbs and Bode *Thromb. Res.* 69:1-58 (1993)). Proteolytic cleavage at this RG site would cause a 67,613 Da loss in the molecular weight of GPV. Comparison of the sequence flanking the RG site to sequences of other known thrombin substrates revealed significant homology to the A α and B β chain of human fibrinogen, to human plasma FXIII, and to human chorionic gonatropin β -subunit (FIG. 7) (SEQ.ID.NOs. 38-43). Amino acid sequence comparison

and assembly was performed using the PC Gene software developed by Intelligenetics Inc., Palo Alto Calif.

The new N-terminal sequence revealed by the potential cleavage site matched that of the Th1 peptide obtained after N-terminal sequencing of thrombin-cleaved platelet GPV (Shimomura et. al., *Blood* 75:2349-2356 (1990); Roth et. al., *Biochem. Biophys. Res. Commun.* 170:152-161 (1990)). Inspection of the sequence surrounding the RG dipeptide did not reveal a cluster of negatively charged residues which are known to be responsible for the interaction of thrombin with the newly cloned thrombin receptor (Vu et. al., *Cell* 64 1057-1068 (1991)) or like those in GPIB α , another thrombin-binding membrane glycoprotein (Lopez et. al., *Proc. Natl. Acad. Sci. USA* 85:2135-2139 (1988)).

EXAMPLE 5

Determination of the Cellular Distribution of Human Platelet Glycoprotein v message via RT-PCR

The cellular distribution of GPV mRNA was assessed using the sensitive RT-PCR amplification technique using primers from the cDNA sequence. GPV mRNA was detected in platelets, megakaryocytes and HEL cells, and was increased in HEL cells after stimulation with phorbol ester, but was not detected in HL60 cells, K562, U937, or endothelial cells (FIG. 2).

All publications, patents, and patent applications herein are incorporated by reference to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference.

Although the present invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be apparent that certain changes and modifications may be practiced within the scope of the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 43

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7452 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(ix) FEATURE:

- (A) NAME/KEY: intron
- (B) LOCATION: 1462..2419

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 2422..4101

(ix) FEATURE:

- (A) NAME/KEY: misc_signal

-continued

-
- (B) LOCATION: 68..76
 - (D) OTHER INFORMATION: /function= "Putitive TPA responsive element"
/label= TRE
 - (ix) FEATURE:
 - (A) NAME/KEY: misc_signal
 - (B) LOCATION: 471..478
 - (D) OTHER INFORMATION: /function= "Ets-1 cis-acting sequence"
/label= Ets-1
 - (ix) FEATURE:
 - (A) NAME/KEY: misc_signal
 - (B) LOCATION: 493..502
 - (D) OTHER INFORMATION: /function= "Ets-1 cis-acting sequence"
/label= Ets-1
 - (ix) FEATURE:
 - (A) NAME/KEY: repeat_region
 - (B) LOCATION: 593..881
 - (D) OTHER INFORMATION: /rpt_type= "other"
/label= Alu
 - (ix) FEATURE:
 - (A) NAME/KEY: misc_signal
 - (B) LOCATION: 897..904
 - (D) OTHER INFORMATION: /function= "Ets-1 cis-acting sequence"
/label= Ets-1
 - (ix) FEATURE:
 - (A) NAME/KEY: misc_binding
 - (B) LOCATION: 1142..1149
 - (D) OTHER INFORMATION: /function= "Sp1 binding site"
/standard_name= "Sp1"
 - (ix) FEATURE:
 - (A) NAME/KEY: misc_signal
 - (B) LOCATION: 1178..1184
 - (D) OTHER INFORMATION: /function= "Ets-1 cis-acting sequences"
/label= Ets-1
 - (ix) FEATURE:
 - (A) NAME/KEY: TATA_signal
 - (B) LOCATION: 1199..1203
 - (ix) FEATURE:
 - (A) NAME/KEY: TATA_signal
 - (B) LOCATION: 1263..1269
 - (ix) FEATURE:
 - (A) NAME/KEY: misc_binding
 - (B) LOCATION: 1285..1289
 - (D) OTHER INFORMATION: /function= "GATA-1 binding site"
 - (ix) FEATURE:
 - (A) NAME/KEY: misc_binding
 - (B) LOCATION: 1321..1326
 - (D) OTHER INFORMATION: /function= "GATA-1 binding site"
 - (ix) FEATURE:
 - (A) NAME/KEY: misc_signal
 - (B) LOCATION: 1365..1372
 - (D) OTHER INFORMATION: /function= "Ets-1 cis-acting sequences"
/label= Ets-1
 - (ix) FEATURE:
 - (A) NAME/KEY: repeat_region
 - (B) LOCATION: 6133..6440
 - (D) OTHER INFORMATION: /rpt_type= "other"
/label= Alu
 - (ix) FEATURE:
 - (A) NAME/KEY: misc_signal
 - (B) LOCATION: 5610..5615

-continued

(D) OTHER INFORMATION: /standard_name= "Polyadenylation signal sequence"

(ix) FEATURE:
 (A) NAME/KEY: misc_signal
 (B) LOCATION: 6966..6971
 (D) OTHER INFORMATION: /standard_name= "Polyadenylation signal sequence"

(ix) FEATURE:
 (A) NAME/KEY: misc_signal
 (B) LOCATION: 7224..7229
 (D) OTHER INFORMATION: /standard_name= "Polyadenylation signal sequence"

(ix) FEATURE:
 (A) NAME/KEY: misc_signal
 (B) LOCATION: 7358..7363
 (D) OTHER INFORMATION: /standard_name= "Polyadenylation signal sequence"

(ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..7452
 (D) OTHER INFORMATION: /standard_name= "Nucleotide sequence containing the human GPV gene"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

TGATCGGAAC  TGAAAGACCT  CCCGCGATAC  CTGGCAGAGG  CAGTGGCTCT  TCCCTGTGGT    60
CCAGGGCTGA  CTGACTTTGA  AGGTAATTTT  AGTCAACCCA  GCCTTTACTG  GGCTCTGACT    120
GCATTAGGCT  GCATCAAAGG  GGATTGGATC  CCATGATTCT  TTATATCTTC  TGACATTAAG    180
CCTTTGTCAG  CTATAGGTGT  TACAAATATC  TTTAGTTTGT  GGTTTATCTT  TTCCCTTTT    240
TTATGGTGTC  TTGAAGGATA  GAAGTCTTAA  TGCAGACAGC  ATTATCAGTG  TGTTCAAAAG    300
ACAGCTAGAC  ACGTTTTGCC  TATAGACAAA  TGGGCAAAAG  GAAACCCAGC  TTTCTCAAAT    360
GAAGCACAAG  TGGGCCTTAA  TTATGTGAAA  AGGTGTTCAA  GTTCATCATT  AAACAGGGAA    420
AGGAAAAGTT  AAAACCATGC  TGAGATATCT  TTCATAGAAA  TGGCAAAAAG  CAGGAAGTGC    480
CACGTGTGGG  CAGAGAGGAA  GCACAGGAAC  TCTCACAAAT  GGCAGGTGTC  ATCGTAGACC    540
AACACAACCA  CTTTGGAGAG  CAGTTTGACT  TTCCCCAGTT  AAAGTGAACA  TGTGAGCGGC    600
CGGGCGTGGT  GGCTCATGCC  TGTAATCCCA  GCAGTTTGGG  AGGCCGAGGC  GGGCGGATTG    660
CCTGAGCTCA  GGAGTTCAAG  ACCAGCCAGG  GCAACACGGT  AAAACCCCGT  CTCTACTAAA    720
ATACAAAAAA  TTAGCTGGGC  GTGATGGTGT  GTGCCTGTAA  TCCAGCTAC  TTGTGAGGCC    780
GAGGCAGGAG  AATTGCTTGA  ACCAGGGAGC  AGGAGGTGTC  AGTGAGCCGA  GATCGCACCA    840
CTGCACCCCA  GCCTGGCGAC  AGAGTCCCCC  TCCCCACCA  AAAAAACAAC  AAGTGAGCAT    900
CCTGCAACCT  AGCAATGCCA  TTGTTGAACA  AGTTCAAAGA  TGTTCATTAGC  CTTATTAGTC    960
CCAAAAGGAA  GAAAAAATG  GAGGATTTGA  GAATGTTCTT  AGCTTTATTG  CTAAGCGGAG   1020
AAAGAAAAAC  AACACATACC  AAAAAAATAA  AAAAAAATAA  AAAAAACAA  AAAACCTGGG   1080
TGGGAAATTA  GGGCCATGTG  GCATGAAAAG  GAAGACCCAG  GGGGAAGTGTG  GCCCATCTAG   1140
GGGTGTGGCT  ACTGCAGTGA  TCCAGCTGTA  TCACTGAACT  TCCCTGGCAT  CATAGAGTTA   1200
TATTGTGCCA  TTTATGGAAA  AACTCCTCCC  ACTGCTCTTG  GCTTTGACAG  TAGGAATCAG   1260
GTTATATATG  GTCTCTCGGT  TTGAAGATAT  TTGTCATTAA  AAACCAGAAC  AAGGGCTCTG   1320
AGATAGGGTC  CTTTCTGAC  CTAAGTCTGGT  AAAGTCTTTA  TCCTCAGGAT  GCAAGGATAC   1380
CACCCCTCTC  CTGTGGAAG  TGTGGAATCA  CATGCAGAGC  TCTAAGTCTT  TCAGTTACTT   1440
TGGAGTGCAG  AACCATTTC  GGTAAGGCCA  AATATTTTAA  ACATTAGTAT  AGGAAATTAG   1500

```

-continued

AGGGCTCTTT AGTCTGTGTG TGCATGAGAA GTAAAATTGC ACGAGAAGCA ATTTATGTAA	1560
AATTTCCGCTT AGGAAACATT GTTTTGGCAG GTTAGTAGTA TGGTGTGCAT TTCCCAGAAA	1620
ATTCAGTGCC GTGAGTATTA CCTTTAGTTA AGCATCTTAG AAATAGTAGC TCTTATTGTT	1680
TATGGCTAAG TCAGAAATAC TACCCTCAAA TTCTATGTGA CCCTAGTTAT ACTGTTGAGC	1740
CTTTCTGTGC CTCTGTGCCT TCATCCTTGA ATCGGGGATA ATATACTTAC CTCCTAAGGT	1800
TATTGTAAAG ATTAATGCA TGTAGTATAA ATAAAGAGCT GAGAACAATG CATGGCGTAA	1860
AGTGATAGGT ATTATTATAT GCTTTTGTG GCTGTTGATT GAAGGTGTTT GCTGTTTGG	1920
GGGTGTCCTT TAATAGAGTA ACTTGGTACT GTGGAATAG CATGATGTG AGCAAAAGAA	1980
TCAGATGGTG GTGGCTGCAG ACTTTGCTGT TCCCTTCTTG ACTGTTGGTT ATAGCCAATG	2040
CAGGGTAAGT TATAAAGTCA AGAGCAGAGC CGTTTTTACA ATGGACATTG CTTTGTGATG	2100
TCTGTGAGCT TGAATGTGAG AATGATTATT TTAATTCCT ATGTAAAGAC TTAAAGTAT	2160
TGGCTATTTC GTAGCTTGAT TTCTCTGTAA TCTCATGCTT TAAACTGAGA GTGAAAAATC	2220
AATAAAGCAA AAGCATGAGG CCACGCAGTG TAGAATGAGT GTCTTTTAC CACGTAGGGA	2280
AATCTGTAGT CCTAAGAAAA GAGGGAGTGA GAATTCCTGGC GAAAAGATTG TGCCCTGCA	2340
CAAAGTGCAG GATCCCAGGG TTCAGTACAG GCGCGAACGC TCCTGTGTGT TGACCACACT	2400
CCCACGGTTG CTTTTTCAGA C ATG CTG AGG GGG ACT CTA CTG TGC GCG GTG	2451
Met Leu Arg Gly Thr Leu Leu Cys Ala Val	
1 5 10	
CTC GGG CTT CTG CGC GCC CAG CCC TTC CCC TGT CCG CCA GCT TGC AAG	2499
Leu Gly Leu Leu Arg Ala Gln Pro Phe Pro Cys Pro Pro Ala Cys Lys	
15 20 25	
TGT GTC TTC CGG GAC GCC GCG CAG TGC TCG GGG GGC GAC GTG GCG CGC	2547
Cys Val Phe Arg Asp Ala Ala Gln Cys Ser Gly Gly Asp Val Ala Arg	
30 35 40	
ATC TCC GCG CTG GGC CTG CCC ACC AAC CTC ACG CAC ATC CTG CTC TTC	2595
Ile Ser Ala Leu Gly Leu Pro Thr Asn Leu Thr His Ile Leu Leu Phe	
45 50 55	
GGA ATG GGC CGC GGC GTC CTG CAG AGC CAG AGC TTC AGC GGC ATG ACC	2643
Gly Met Gly Arg Gly Val Leu Gln Ser Gln Ser Phe Ser Gly Met Thr	
60 65 70	
GTC CTG CAG CGC CTC ATG ATC TCC GAC AGC CAC ATT TCC GCC GTT GCC	2691
Val Leu Gln Arg Leu Met Ile Ser Asp Ser His Ile Ser Ala Val Ala	
75 80 85 90	
CCC GGC ACC TTC AGT GAC CTG ATA AAA CTG AAA ACC CTG AGG CTG TCG	2739
Pro Gly Thr Phe Ser Asp Leu Ile Lys Leu Lys Thr Leu Arg Leu Ser	
95 100 105	
CGC AAC AAA ATC ACG CAT CTT CCA GGT GCG CTG CTG GAT AAG ATG GTG	2787
Arg Asn Lys Ile Thr His Leu Pro Gly Ala Leu Leu Asp Lys Met Val	
110 115 120	
CTC CTG GAG CAG TTG TTT TTG GAC CAC AAT GCG CTA AGG GGC ATT GAC	2835
Leu Leu Glu Gln Leu Phe Leu Asp His Asn Ala Leu Arg Gly Ile Asp	
125 130 135	
CAA AAC ATG TTT CAG AAA CTG GTT AAC CTG CAG GAG CTC GCT CTG AAC	2883
Gln Asn Met Phe Gln Lys Leu Val Asn Leu Gln Glu Leu Ala Leu Asn	
140 145 150	
CAG AAT CAG CTC GAT TTC CTT CCT GCC AGT CTC TTC ACG AAT CTG GAG	2931
Gln Asn Gln Leu Asp Phe Leu Pro Ala Ser Leu Phe Thr Asn Leu Glu	
155 160 165 170	
AAC CTG AAG TTG TTG GAT TTA TCG GGA AAC AAC CTG ACC CAC CTG CCC	2979
Asn Leu Lys Leu Leu Asp Leu Ser Gly Asn Asn Leu Thr His Leu Pro	
175 180 185	
AAG GGG TTG CTT GGA GCA CAG GCT AAG CTC GAG AGA CTT CTG CTC CAC	3027

-continued

Lys	Gly	Leu	Leu	Gly	Ala	Gln	Ala	Lys	Leu	Glu	Arg	Leu	Leu	Leu	His		
		190						195					200				
TCG	AAC	CGC	CTT	GTG	TCT	CTG	GAT	TCG	GGG	CTG	TTG	AAC	AGC	CTG	GGC	3075	
Ser	Asn	Arg	Leu	Val	Ser	Leu	Asp	Ser	Gly	Leu	Leu	Asn	Ser	Leu	Gly		
		205					210					215					
GCC	CTG	ACG	GAG	CTG	CAG	TTC	CAC	CGA	AAT	CAC	ATC	CGT	TCC	ATC	GCA	3123	
Ala	Leu	Thr	Glu	Leu	Gln	Phe	His	Arg	Asn	His	Ile	Arg	Ser	Ile	Ala		
		220				225					230						
CCC	GGG	GCC	TTC	GAC	CGG	CTC	CCA	AAC	CTC	AGT	TCT	TTG	ACG	CTT	TCG	3171	
Pro	Gly	Ala	Phe	Asp	Arg	Leu	Pro	Asn	Leu	Ser	Ser	Leu	Thr	Leu	Ser		
		235			240					245					250		
AGA	AAC	CAC	CTT	GCG	TTT	CTC	CCC	TCT	GCG	CTC	TTT	CTT	CAT	TCG	CAC	3219	
Arg	Asn	His	Leu	Ala	Phe	Leu	Pro	Ser	Ala	Leu	Phe	Leu	His	Ser	His		
				255					260					265			
AAT	CTG	ACT	CTG	TTG	ACT	CTG	TTC	GAG	AAC	CCG	CTG	GCA	GAG	CTC	CCG	3267	
Asn	Leu	Thr	Leu	Leu	Thr	Leu	Phe	Glu	Asn	Pro	Leu	Ala	Glu	Leu	Pro		
			270					275					280				
GGG	GTG	CTC	TTC	GGG	GAG	ATG	GGG	GGC	CTG	CAG	GAG	CTG	TGG	CTG	AAC	3315	
Gly	Val	Leu	Phe	Gly	Glu	Met	Gly	Gly	Leu	Gln	Glu	Leu	Trp	Leu	Asn		
		285				290						295					
CGC	ACC	CAG	CTG	CGC	ACC	CTG	CCC	GCC	GCC	GCC	TTC	CGC	AAC	CTG	AGC	3363	
Arg	Thr	Gln	Leu	Arg	Thr	Leu	Pro	Ala	Ala	Ala	Phe	Arg	Asn	Leu	Ser		
		300				305					310						
CGC	CTG	CGG	TAC	TTA	GGG	GTG	ACT	CTG	AGC	CCG	CGG	CTG	AGC	GCG	CTT	3411	
Arg	Leu	Arg	Tyr	Leu	Gly	Val	Thr	Leu	Ser	Pro	Arg	Leu	Ser	Ala	Leu		
		315				320				325					330		
CCG	CAG	GGC	GCC	TTC	CAG	GGC	CTT	GGC	GAG	CTC	CAG	GTG	CTC	GCC	CTG	3459	
Pro	Gln	Gly	Ala	Phe	Gln	Gly	Leu	Gly	Glu	Leu	Gln	Val	Leu	Ala	Leu		
				335					340					345			
CAC	TCC	AAC	GGC	CTG	ACC	GCC	CTC	CCC	GAC	GGC	TTG	CTG	CGC	GGC	CTC	3507	
His	Ser	Asn	Gly	Leu	Thr	Ala	Leu	Pro	Asp	Gly	Leu	Leu	Arg	Gly	Leu		
			350					355					360				
GGC	AAG	CTG	CGC	CAG	GTG	TCC	CTG	CGC	CGC	AAC	AGG	CTG	CGC	GCC	CTG	3555	
Gly	Lys	Leu	Arg	Gln	Val	Ser	Leu	Arg	Arg	Asn	Arg	Leu	Arg	Ala	Leu		
		365					370					375					
CCC	CGT	GCC	CTC	TTC	CGC	AAT	CTC	AGC	AGC	CTG	GAG	AGC	GTC	CAG	CTC	3603	
Pro	Arg	Ala	Leu	Phe	Arg	Asn	Leu	Ser	Ser	Leu	Glu	Ser	Val	Gln	Leu		
		380				385					390						
GAC	CAC	AAC	CAG	CTG	GAG	ACC	CTG	CCT	GGC	GAC	GTG	TTT	GGG	GCT	CTG	3651	
Asp	His	Asn	Gln	Leu	Glu	Thr	Leu	Pro	Gly	Asp	Val	Phe	Gly	Ala	Leu		
				400					405					410			
CCC	CGG	CTG	ACG	GAG	GTC	CTG	TTG	GGG	CAC	AAC	TCC	TGG	CGC	TGC	GAC	3699	
Pro	Arg	Leu	Thr	Glu	Val	Leu	Leu	Gly	His	Asn	Ser	Trp	Arg	Cys	Asp		
				415					420					425			
TGT	GGC	CTG	GGG	CCC	TTC	CTG	GGG	TGG	CTG	CGG	CAG	CAC	CTA	GGC	CTC	3747	
Cys	Gly	Leu	Gly	Pro	Phe	Leu	Gly	Trp	Leu	Arg	Gln	His	Leu	Gly	Leu		
			430					435					440				
GTG	GGC	GGG	GAA	GAG	CCC	CCA	CGG	TGC	GCA	GGC	CCT	GGG	GCG	CAC	GCC	3795	
Val	Gly	Gly	Glu	Glu	Pro	Pro	Arg	Cys	Ala	Gly	Pro	Gly	Ala	His	Ala		
			445				450					455					
GGC	CTG	CCG	CTC	TGG	GCC	CTG	CCG	GGG	GGT	GAC	GCG	GAG	TGC	CCG	GGC	3843	
Gly	Leu	Pro	Leu	Trp	Ala	Leu	Pro	Gly	Gly	Asp	Ala	Glu	Cys	Pro	Gly		
		460				465					470						
CCC	CGG	GGC	CCG	CCT	CCC	CGC	CCC	GCT	GCG	GAC	AGC	TCC	TCG	GAA	GCC	3891	
Pro	Arg	Gly	Pro	Pro	Pro	Arg	Pro	Ala	Ala	Asp	Ser	Ser	Ser	Glu	Ala		
		475			480					485				490			
CCT	GTC	CAC	CCA	GCC	TTG	GCT	CCC	AAC	AGC	TCA	GAA	CCC	TGG	GTG	TGG	3939	
Pro	Val	His	Pro	Ala	Leu	Ala	Pro	Asn	Ser	Ser	Glu	Pro	Trp	Val	Trp		
				495				500						505			

-continued

GCC CAG CCG GTG ACC ACG GGC AAA GGT CAA GAT CAT AGT CCG TTC TGG	3987
Ala Gln Pro Val Thr Thr Gly Lys Gly Gln Asp His Ser Pro Phe Trp	
510 515 520	
GGG TTT TAT TTT CTG CTT TTA GCT GTT CAG GCC ATG ATC ACC GTG ATC	4035
Gly Phe Tyr Phe Leu Leu Leu Ala Val Gln Ala Met Ile Thr Val Ile	
525 530 535	
ATC GTG TTT GCT ATG ATT AAA ATT GGC CAA CTC TTT CGA AAA TTA ATC	4083
Ile Val Phe Ala Met Ile Lys Ile Gly Gln Leu Phe Arg Lys Leu Ile	
540 545 550	
AGA GAG AGA GCC CTT GGG TAAACCAATG GGAAAATCTT CTAATTACTT	4131
Arg Glu Arg Ala Leu Gly	
555 560	
AGAACCTGAC CAGATGTGGC TCGGAGGGGA ATCCAGACCC GCTGCTGTCT TGCTCTCCCT	4191
CCCCTCCCCA CTCCTCTCTT CTCTCTCTCT TTCTCTCTCA CTGCCACGCC TTCCTTTCCC	4251
TCCTCTCTCC CCTCTCCGCT CTGTGCTCTT CATTCTCACG GGCCCGCAAC CCCTCTCTCT	4311
TCTGTCCCCG CCCGTCTCTG GAAACTGAGC TTGACGTTTG TAAACTGTGG TTGCCTGCCT	4371
TCCCAGCTCC ACGCGGTGTG CGCTGACACT GCCGGGGGGC TGGACTGTGT TGGACCCATC	4431
CTTGCCCCGC TGTGCCGTGC TTGGCCTCTG GTGGAGAGAG GGACCTCTTC AGTGTCTACT	4491
GAGTAAGGGG ACAGCTCCAG GCCGGGGCTG TCTCTGCAC AGAGTAAGCC GGTAAATGTT	4551
TGTGAAATCA ATGCGTGGAT AAAGGAACAC ATGCCATCCA AGTGATGATG GCTTTTCCTG	4611
GAGGGAAAGG ATAGGCTGTT GCTCTATCTA ATTTTGTGTT TTTGTTTGTG GACAGTCTAG	4671
CTCTGTGGCC CAGGCTGGCG TGCAGTGGGC CGTCTCAGTT CACTGCAGCC TCCGCCCTCC	4731
AGGTTCAGT GATTCTCATG CCTCAGCGTT CTGAGTAGCT GGGATTAGAG GCGTGTGCCA	4791
CTACACCCGG CTAATTTTTG TACTTTTTAA AGTAGAGACG GGCTTTGCCA TATTGGCCTG	4851
GCTGATCTCA AACTCCTGGT CTGAACTCC TGGCCACAAG TGATCTGCC GCCTTAGCCT	4911
CCCCAAGTGC TGGGATTACA GCGCAAGCC ACTACACCTG CCCTCTTCAT CGAATTTTAT	4971
TTGAGAAGTA GAGCTCTGC CATTTTTTCC CTGCTCCAT TTTTCTCACT TTATGTCTCT	5031
CTGACCTATG GGCTACTTGG GAGAGCAC TGACTCCATTC ATGCATGAGC ATTTTCAGGA	5091
TAAGCGACTT CTGTGAGGCT GAGAGAGGAA GAAAACACGG AGCCTTCCCT CCAGGTGCC	5151
AGTGTAGGTC CAGCGTGTCT CTTGAGCCTC CTGTGAGTTT CCACTTGCTT TACATCCATG	5211
CAACATGTCA TTTTGAAACT GGATTGATTT GCATTTCTCTG GAACTCTGCC ACCTCATTTT	5271
ACAAGCATTT ATGGAGCAGT TAACATGTGA CTGGTATTCA TGAATATAAT GATAAGCTTG	5331
ATTCTAGTTC AGCTGCTGTC ACAGTCTCAT TTGTTCTTCC AACTGAAAGC CGTAAAACCT	5391
TTGTTGCTTT AATTGAAATGT CTGTGCTTAT GAGAGGCAGT GGTAAAACA TTTTCTGGCG	5451
AGTTGACAAC TGTGGGTTC AATCCCAGCT CTACCACCTA CTAAGTGCAT GGGACTTTGG	5511
GTAAGACACC TGCTTACATT CTCTAAGCCT TGGTTTCTCG AACCTTAAAA CAGGATAACA	5571
TAGTACCTGC TTCATAGAGT TTTGTGAGAA TTAAAGGCAA TAAAGCATAT AATGACTTAG	5631
CCCAGCGGCC TGCAGACAA ACATGTTAAT GAATGTTAGC TATTATTACT AAAGATGAGC	5691
AATTATTATT GGCATCATGA TTTCTAAAGA AGAGCTTTGA GTTGGTATTT TTCTCTGTGT	5751
ATAAGGGTAA GTCCGAACCT TCTCATACTG GAGGTTACAT TCACATCAGT CTGTCTTCCC	5811
CTGCGGATGG CCTCAGCCCT GGGTGGCCAG GCTCTGTGCT CACAGTCCAG AGCAATGGAT	5871
CCTCCAACAC CACCAGTGG ATGTGGAGCA GGAGAGCTGG ATCGTGGCAT TTGTTTCTGG	5931
GTTCTGCAGT TGGGAGTTGG TTTCTGGGTT CTCCATTGGT CTAAGTGTCT AGTCCCATAC	5991
CAGACTCACG GTCTCCATTA TTGGAGCTTT AATAATTTTT GGTATAGGGT CATCTCTCCA	6051

-continued

CCTTGTTTTT CTTCTATTCT TGGTTCTTTG CAATTCCTATG AATATTTTCAG GGTCAGCATG 6111
 TCAACTCCAT TGAAAAACCC TGCTGGGATT TTAATAGAAC TTACAGCTCA CGCCTGTAAT 6171
 CCCAGCACTT TGGGAGGCTG AGGTGGGTGG ATCAGAGTC AGGAGTTTGA GAACAGCTGG 6231
 CCAAGATGGT GAAACCCCGT CTCTACTAAA AATACAAAAA TTAGCTGGGT GCGGTGGCAG 6291
 GTGCCTGTAG TCCCAGCTAC TTGGGACACC GAGGCAGGAG AATCACTTGA ACCCGGGAGG 6351
 CGGAGGTTGC AGTGAGCCGA GATCGTGCCA CTGCACTCTA GCCTGGGCGA CAGAGCGAGA 6411
 CTCCATCTCA AAAAAAAGA AAAAGAAAAT TGCAGTAAAT TTAAACTAA TTTGGGGAAG 6471
 AATCTGTATT TTTACAATAC CTAGTGTCTT TGCCAGTAAG CATGGTTCAT CTTCCCATTT 6531
 ATTTACGCTA TTTTAAATCT TTCAGTGATG TTTTAGAATT TTTTTATAA AAACCTTCAC 6591
 TATAAGAACA GAAACCCAAA CACCGCATGT TCTCACTCAT AGGTGGGAAT TGAACAATGA 6651
 GAACACTTGG ACACAGGGCG GGAACGCTCA CACGCCTGGA CTGTTGGGGG GGTGGCTGGG 6711
 AGAGGGATAG TGTTAGGAGA AATACCTAAT GTAATGACG AGTTAATGGT GCAGCCAACC 6771
 AACCTGGCAC ATGTATTTCAT ATGTAACAAA CCTGCACGTT GTGCACATGT ACCCTAGAAC 6831
 TTAAAGTATA TTAAAAAAG AAACCTTGGC ACTGATTTTG TTAGATTTAT TCCTAGGTAT 6891
 CCTTCTCTT TTTTGATTTG TCATTGCTAT TGTAGATGGC ATCTTTTAA AAAGTTATAT 6951
 TTTCTAAAGC AAAAAATAA AAAAGTTGTA TTTCTAATTT TTATTACCAA TATATAAGAA 7011
 TGTAATTTAT TTTTACATAA TTATCTTATG TCTAGTAATA ATTCTGATAA TTTGCTTCTT 7071
 CCTATTAATA CCTTACACCC ATTATGATT TATTTTCTG TTTTAAATA TCTTCTGCA 7131
 CTGGCTAAAA CCTCCACTAT AATGTTGAGC AGAACAGTGA GGCATCCTTA GAACTATCTT 7191
 GGTGCAAGG GGTAGTCTC TAATGTTTCA TCAATAAATG TGATGTTTCT AGTCTGAGTT 7251
 TGCTAAGTAT ATTTAAAAAT AATCAGTAAA GTTAGATTTT ATCCATTTT ATCTTAACTA 7311
 TTGAGATGCT CATATCATTT TTCTTCTCA ATGTGTTAAA ATGGTGAATA AATTTATAGA 7371
 TTTTGGAAAA GTAAATTCAT TCTTGCATTC CCGAAGTAAA CCAAGCCATG CTATGTGTAT 7431
 TTAAAAATA TTGCTGAATT C 7452

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 560 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Leu Arg Gly Thr Leu Leu Cys Ala Val Leu Gly Leu Leu Arg Ala
 1 5 10 15
 Gln Pro Phe Pro Cys Pro Pro Ala Cys Lys Cys Val Phe Arg Asp Ala
 20 25 30
 Ala Gln Cys Ser Gly Gly Asp Val Ala Arg Ile Ser Ala Leu Gly Leu
 35 40 45
 Pro Thr Asn Leu Thr His Ile Leu Leu Phe Gly Met Gly Arg Gly Val
 50 55 60
 Leu Gln Ser Gln Ser Phe Ser Gly Met Thr Val Leu Gln Arg Leu Met
 65 70 75 80
 Ile Ser Asp Ser His Ile Ser Ala Val Ala Pro Gly Thr Phe Ser Asp
 85 90 95

-continued

Leu Ile Lys Leu Lys Thr Leu Arg Leu Ser Arg Asn Lys Ile Thr His
 100 105 110

Leu Pro Gly Ala Leu Leu Asp Lys Met Val Leu Leu Glu Gln Leu Phe
 115 120 125

Leu Asp His Asn Ala Leu Arg Gly Ile Asp Gln Asn Met Phe Gln Lys
 130 135 140

Leu Val Asn Leu Gln Glu Leu Ala Leu Asn Gln Asn Gln Leu Asp Phe
 145 150 155 160

Leu Pro Ala Ser Leu Phe Thr Asn Leu Glu Asn Leu Lys Leu Leu Asp
 165 170 175

Leu Ser Gly Asn Asn Leu Thr His Leu Pro Lys Gly Leu Leu Gly Ala
 180 185 190

Gln Ala Lys Leu Glu Arg Leu Leu Leu His Ser Asn Arg Leu Val Ser
 195 200 205

Leu Asp Ser Gly Leu Leu Asn Ser Leu Gly Ala Leu Thr Glu Leu Gln
 210 215 220

Phe His Arg Asn His Ile Arg Ser Ile Ala Pro Gly Ala Phe Asp Arg
 225 230 235 240

Leu Pro Asn Leu Ser Ser Leu Thr Leu Ser Arg Asn His Leu Ala Phe
 245 250 255

Leu Pro Ser Ala Leu Phe Leu His Ser His Asn Leu Thr Leu Leu Thr
 260 265 270

Leu Phe Glu Asn Pro Leu Ala Glu Leu Pro Gly Val Leu Phe Gly Glu
 275 280 285

Met Gly Gly Leu Gln Glu Leu Trp Leu Asn Arg Thr Gln Leu Arg Thr
 290 295 300

Leu Pro Ala Ala Ala Phe Arg Asn Leu Ser Arg Leu Arg Tyr Leu Gly
 305 310 315 320

Val Thr Leu Ser Pro Arg Leu Ser Ala Leu Pro Gln Gly Ala Phe Gln
 325 330 335

Gly Leu Gly Glu Leu Gln Val Leu Ala Leu His Ser Asn Gly Leu Thr
 340 345 350

Ala Leu Pro Asp Gly Leu Leu Arg Gly Leu Gly Lys Leu Arg Gln Val
 355 360 365

Ser Leu Arg Arg Asn Arg Leu Arg Ala Leu Pro Arg Ala Leu Phe Arg
 370 375 380

Asn Leu Ser Ser Leu Glu Ser Val Gln Leu Asp His Asn Gln Leu Glu
 385 390 395 400

Thr Leu Pro Gly Asp Val Phe Gly Ala Leu Pro Arg Leu Thr Glu Val
 405 410 415

Leu Leu Gly His Asn Ser Trp Arg Cys Asp Cys Gly Leu Gly Pro Phe
 420 425 430

Leu Gly Trp Leu Arg Gln His Leu Gly Leu Val Gly Gly Glu Glu Pro
 435 440 445

Pro Arg Cys Ala Gly Pro Gly Ala His Ala Gly Leu Pro Leu Trp Ala
 450 455 460

Leu Pro Gly Gly Asp Ala Glu Cys Pro Gly Pro Arg Gly Pro Pro Pro
 465 470 475 480

Arg Pro Ala Ala Asp Ser Ser Ser Glu Ala Pro Val His Pro Ala Leu
 485 490 495

Ala Pro Asn Ser Ser Glu Pro Trp Val Trp Ala Gln Pro Val Thr Thr
 500 505 510

Gly Lys Gly Gln Asp His Ser Pro Phe Trp Gly Phe Tyr Phe Leu Leu

-continued

515	520	525
Leu Ala Val Gln Ala Met	Ile Thr Val Ile Ile	Val Phe Ala Met Ile
530	535	540
Lys Ile Gly Gln Leu Phe	Arg Lys Leu Ile Arg	Glu Arg Ala Leu Gly
545	550	555 560

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CAGGATGCAA

10

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGGGTGTGGC

10

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Asn Ser Trp Arg Cys Asp Cys Gly Leu
1 5

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 13 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Peptide
 (B) LOCATION: 1..13
 (D) OTHER INFORMATION: /note= "K5/6 peptide residues
 1-13."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Lys Met Val Leu Leu Glu Gln Leu Phe Leu Asp His Asn
1 5 10

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

-continued

- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..23
 (D) OTHER INFORMATION: /standard_name= "Primer 1 (+)"
 /note= "Oligonucleotide primer used for the PCR
 amplification of platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GARCARCTGT TYCTSGAYCA YAA

23

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 12 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

- (ix) FEATURE:
 (A) NAME/KEY: Peptide
 (B) LOCATION: 1..12
 (D) OTHER INFORMATION: /note= "K5/6 peptide residues
 13-24."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Asn Ala Leu Arg Gly Ile Asp Gln Asn Met Phe Gln
 1 5 10

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..32
 (D) OTHER INFORMATION: /standard_name= "Primer 2 (+)"
 /note= "Primer used for the PCR amplification of
 platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TGCGCTAAGG GGCATTGACC AAAACATGTT TC

32

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

- (ix) FEATURE:
 (A) NAME/KEY: Peptide
 (B) LOCATION: 1..10
 (D) OTHER INFORMATION: /note= "K5/6 peptide residues
 21-30."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Asn Met Phe Gln Lys Leu Val Asn Leu Gln
 1 5 10

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid

-continued

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 1..27
(D) OTHER INFORMATION: /standard_name= "Primer 3(-)"
/note= "Primer used for the PCR amplification of
platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GTACAAAGTC TTTGACCAAT TGGACGT 27

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:
(A) NAME/KEY: Peptide
(B) LOCATION: 1..11
(D) OTHER INFORMATION: /note= "K5/6 peptide residues
31-41."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Glu Leu Ala Leu Asn Gln Asn Gln Leu Asp Phe
1 5 10

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 1..21
(D) OTHER INFORMATION: /standard_name= "Primer 4 (-)"
/note= "Primer used for the PCR amplification of
platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TTRGTYTTRG TYGASCTRAA R 21

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:
(A) NAME/KEY: Peptide
(B) LOCATION: 1..7
(D) OTHER INFORMATION: /note= "M6 peptide residues
1-7."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Met Ile Ser Asp Ser His Ile
1 5

(2) INFORMATION FOR SEQ ID NO:15:

-continued

-
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..20
 (D) OTHER INFORMATION: /standard_name= "Primer 5 (+)"
 /note= "Primer used for the PCR amplification of
 platelet GPV."
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:
 ATGATCTSCG AYWSCCAYAT 20
- (2) INFORMATION FOR SEQ ID NO:16:
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..21
 (D) OTHER INFORMATION: /standard_name= "Primer 6 (+)"
 /note= "Primer used for the PCR amplification of
 platelet GPV."
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:
 GAGAGACTTC TGCTCCACTC G 21
- (2) INFORMATION FOR SEQ ID NO:17:
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..21
 (D) OTHER INFORMATION: /standard_name= "Primer 7 (-)"
 /note= "Primer used for the PCR amplification of
 platelet GPV."
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:
 TATCAGGTCA CTGAAGGTGC C 21
- (2) INFORMATION FOR SEQ ID NO:18:
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..18
 (D) OTHER INFORMATION: /standard_name= "Primer 8 (-)"
 /note= "Primer used for the PCR amplification of
 platelet GPV."
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:
 AAGACACACT TGCAAGCT 18
- (2) INFORMATION FOR SEQ ID NO:19:

-continued

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..35
 (D) OTHER INFORMATION: /standard_name= "Adaptor dT17"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:
- GACTCGAGTC GACATCGATT TTTTTTTTTT TTTT 35
- (2) INFORMATION FOR SEQ ID NO:20:
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..30
 (D) OTHER INFORMATION: /standard_name= "Adaptor dC12"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:
- GACTCGAGTC GACATCGACC CCCCCCCCCC 30
- (2) INFORMATION FOR SEQ ID NO:21:
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..18
 (D) OTHER INFORMATION: /standard_name= "Adaptor"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:
- GACTCGAGTC GACATCGA 18
- (2) INFORMATION FOR SEQ ID NO:22:
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown
- (ii) MOLECULE TYPE: peptide
- (ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 1..24
 (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:
- Gly Arg Gly Val Leu Gln Ser Gln Ser Phe Ser Gly Met Thr Val Leu
 1 5 10 15
- Gln Arg Leu Met Ile Ser Asp Ser
 20
- (2) INFORMATION FOR SEQ ID NO:23:

-continued

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 1..24
 (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

His Ile Ser Ala Val Ala Pro Gly Thr Phe Ser Asp Leu Ile Lys Leu
 1 5 10 15

Lys Thr Leu Arg Leu Ser Arg Asn
 20

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 1..24
 (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Lys Ile Thr His Leu Pro Gly Ala Leu Leu Asp Lys Met Val Leu Leu
 1 5 10 15

Glu Gln Leu Phe Leu Asp His Asn
 20

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 1..24
 (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Ala Leu Arg Gly Ile Asp Gln Asn Met Phe Gln Lys Leu Val Asn Leu
 1 5 10 15

Gln Glu Leu Ala Leu Asn Gln Asn
 20

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

-continued

(ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 1..24
 (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Gln Leu Asp Phe Leu Pro Ala Ser Leu Phe Thr Asn Leu Glu Asn Leu
 1 5 10 15

Lys Leu Leu Asp Leu Ser Gly Asn
 20

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 1..24
 (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Asn Leu Thr His Leu Pro Lys Gly Leu Leu Gly Ala Gln Ala Lys Leu
 1 5 10 15

Glu Arg Leu Leu Leu His Ser Asn
 20

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 1..24
 (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Arg Leu Val Ser Leu Asp Ser Gly Leu Leu Asn Ser Leu Gly Ala Leu
 1 5 10 15

Thr Glu Leu Gln Phe His Arg Asn
 20

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 1..24
 (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

-continued

His Ile Arg Ser Ile Ala Pro Gly Ala Phe Asp Arg Leu Pro Asn Leu
1 5 10 15

Ser Ser Leu Thr Leu Ser Arg Asn
20

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..24
- (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

His Leu Ala Phe Leu Pro Ser Ala Leu Phe Leu His Ser His Asn Leu
1 5 10 15

Thr Leu Leu Thr Leu Phe Glu Asn
20

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..24
- (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Pro Leu Ala Glu Leu Pro Gly Val Leu Phe Gly Glu Met Gly Gly Leu
1 5 10 15

Gln Glu Leu Trp Leu Asn Arg Thr
20

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..24
- (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Gln Leu Arg Thr Leu Pro Ala Ala Ala Phe Arg Asn Leu Ser Arg Leu
1 5 10 15

Arg Tyr Leu Gly Val Thr Leu Ser
20

-continued

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..24
- (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Pro Arg Leu Ser Ala Leu Pro Gln Gly Ala Phe Gln Gly Leu Gly Glu
 1 5 10 15
 Leu Gln Val Leu Ala Leu His Ser Asn
 20 25

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..24
- (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Gly Leu Thr Ala Leu Pro Asp Gly Leu Leu Arg Gly Leu Gly Lys Leu
 1 5 10 15
 Arg Gln Val Ser Leu Arg Arg Asn
 20

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..24
- (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Arg Leu Arg Ala Leu Pro Arg Ala Leu Phe Arg Asn Leu Ser Ser Leu
 1 5 10 15
 Glu Ser Val Gln Leu Asp His Asn
 20

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

-continued

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..24
- (D) OTHER INFORMATION: /note= "Tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Gln Leu Glu Thr Leu Pro Gly Asp Val Phe Gly Ala Leu Pro Arg Leu
 1 5 10 15

Thr Glu Val Leu Leu Gly His Asn
 20

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..24
- (D) OTHER INFORMATION: /note= "Consensus sequence for the tandem Leu-rich repeated structure for platelet GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Xaa Leu Xaa Xaa Leu Pro Xaa Xaa Leu Phe Xaa Xaa Leu Xaa Xaa Leu
 1 5 10 15

Xaa Xaa Leu Xaa Leu Xaa Xaa Asn
 20

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..11
- (D) OTHER INFORMATION: /note= "Amino acid sequence of the GPV thrombin cleavage site."

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..2
- (D) OTHER INFORMATION: /note= "Amino acid residues found in other thrombin substrates."

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 5..9
- (D) OTHER INFORMATION: /note= "Amino acid residues found in other thrombin substrates."

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 11
- (D) OTHER INFORMATION: /note= "Amino acid residue found in other thrombin substrates."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

-continued

Ala Glu Cys Pro Gly Pro Arg Gly Pro Pro Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..11
- (D) OTHER INFORMATION: /note= "Amino acid sequence of the human fibrinogen (Fg) A-alpha 1 chain thrombin cleavage site."

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..2
- (D) OTHER INFORMATION: /note= "Amino acid residues identical to GPV."

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 5
- (D) OTHER INFORMATION: /note= "Amino acid residue identical to GPV."

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 7..9
- (D) OTHER INFORMATION: /note= "Amino acid residues identical to GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

Ala Glu Gly Gly Gly Val Arg Gly Pro Arg Val
1 5 10

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..11
- (D) OTHER INFORMATION: /note= "Amino acid sequence of the human fibrinogen (Fg) A-alpha 2 chain thrombin cleavage site."

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 5..7
- (D) OTHER INFORMATION: /note= "Amino acid residues identical to GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

Gly Gly Val Arg Gly Pro Arg Val Val Glu Arg
1 5 10

(2) INFORMATION FOR SEQ ID NO:41:

-continued

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 11 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(ix) FEATURE:
 (A) NAME/KEY: Peptide
 (B) LOCATION: 1..11
 (D) OTHER INFORMATION: /note= "Amino acid sequence of the human fibrinogen (Fg) B-beta chain thrombin cleavage site."

(ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 7..8
 (D) OTHER INFORMATION: /note= "Amino acid residues identical to GPV."

(ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 11
 (D) OTHER INFORMATION: /note= "Amino acid residue identical to GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Glu Gly Phe Phe Ser Ala Arg Gly His Arg Pro
 1 5 10

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 11 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(ix) FEATURE:
 (A) NAME/KEY: Peptide
 (B) LOCATION: 1..11
 (D) OTHER INFORMATION: /note= "Amino acid sequence of the human plasma factor XIII (FXIII) thrombin cleavage site."

(ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 6..8
 (D) OTHER INFORMATION: /note= "Amino acid residues identical to GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

Glu Leu Gln Gly Val Pro Arg Gly Val Asp Leu
 1 5 10

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 11 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(ix) FEATURE:
 (A) NAME/KEY: Peptide
 (B) LOCATION: 1..11
 (D) OTHER INFORMATION: /note= "Amino acid sequence of the

-continued

human chorionic gonatropin beta-subunit (CGbeta)
thrombin cleavage site."

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 6..8
- (D) OTHER INFORMATION: /note= "Amino acid residues identical to GPV."

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 11
- (D) OTHER INFORMATION: /note= "Amino acid residue identical to GPV."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

Arg Leu Pro Gly Cys Pro Arg Gly Val Asn Pro
1 5 10

TABLE I

Peptide sequences and oligonucleotide primers used for the PCR amplification of platelet GPV																	
	K ¹	M	V	L	L	L	5'-	L	E	Q	L	F	L	D	H	N ¹³	(SEQ.ID.NO. 6) (SEQ.ID.NO. 7)
K5/6 peptide Primer 1 (+)																	
K5/6 peptide Primer 2 (+)	N ¹³ 5'-T	A GCG	L CTA	R AGG	G GGC	I AIT	D GAC	Q CAA	O GAC	L CAA	N AAC	F TIT	M ATG	F TIT	Q ²⁴ C-3'	AA-3'	(SEQ.ID.NO. 8) (SEQ.ID.NO. 9)
K5/6 peptide Primer 3 (-)	N ²¹ 3-G	M TAC	F AAA	Q GTC	K TIT	L GAC	V CAA	N TTG	L GAC	N GAC	L GAC	L GAC	L GAC	L GAC	L GAC	L GAC	(SEQ.ID.NO. 10) (SEQ.ID.NO. 11)
K5/6 peptide Primer 4 (-)	E ³¹	L	A	L 3-	N TTG	O T	N TTG	O T	N TTG	O T	L GAC	L GAC	L GAC	L GAC	L GAC	L GAC	(SEQ.ID.NO. 12) (SEQ.ID.NO. 13)
M6 peptide Primer 5 (+)	M ¹ 5'-ATG	I ATC	S TGC	D GAT	S TCC	H CAT	I ⁷ AT-3'										(SEQ.ID.NO. 14) (SEQ.ID.NO. 15)
Primer 6 (+)	5'-GAG	AGA	CTT	CTG	CTC	CAC	TCG-3'										(SEQ.ID.NO. 16)
Primer 7 (-)	5'-TAT	CAG	GTC	ACT	GAA	GGT	GCC-3'										(SEQ.ID.NO. 17)
Primer 8 (-)	5'-AAG	ACA	CAC	TTG	CAA	GCT											(SEQ.ID.NO. 18)
Adaptor-dIT7	5'-GAC	TCG	AGT	CGA	CAI	CGA	TTT	TTT	TTT	TTT	TTT	TTT	TTT	TTT	TT-3'		(SEQ.ID.NO. 19)
Adaptor dC12	5'-GAC	TCG	AGT	CGA	CAI	CGA	CCC	CCC	CCC	CCC	CCC	CCC	CCC-3'				(SEQ.ID.NO. 20)
Adaptor	5'-GAC	TCG	AGT	CGA	CAI	CGA-3'											(SEQ.ID.NO. 21)

The K/6 and M6 peptide sequences were taken from ref.X and numbered accordingly. Eco RI and Sal I restriction sites were added at the 5'-end of coding (+) and non coding (-) strand primers respectively to facilitate further subcloning of the PCR products.

What is claimed is:

1. An isolated DNA construct, comprising a polynucleotide sequence, wherein said polynucleotide sequence encodes a human glycoprotein V polypeptide having the amino acid sequence as shown in SEQ ID NO: 2.
2. The DNA construct of claim 1 wherein the polynucleotide sequence comprises the nucleotide sequence of SEQ ID NO: 1.
3. The DNA construct of claim 1 wherein the polynucleotide sequence comprises bases 2422 to 4101 of SEQ ID NO: 1.
4. The DNA construct of any one of claims 1 to 3, wherein the polynucleotide sequence lacks introns.
5. The DNA construct of any one of claims 1 to 3, further comprising a heterologous promoter operably linked to the polynucleotide sequence.
6. The DNA construct of claim 5, wherein the promoter is capable of directing expression of the polynucleotide sequence in a eukaryote.
7. The DNA construct of claim 5, wherein the promoter is capable of directing expression of the polynucleotide sequence in a prokaryote.
8. A recombinant host cell comprising the DNA construct of claim 5.
9. The recombinant host cell of claim 8, wherein the cell is a eukaryote.

10. The recombinant host cell of claim 8, wherein the cell is a prokaryote.

11. A method for detection of cells of megakaryoblastic lineage in humans, comprising:

- (a) providing a first primer, the sequence of said first primer corresponding to a portion of SEQ ID NO: 1 and having a T_M of about 30°C to about 70°C;
- (b) providing a second primer, the sequence of said second primer corresponding to the complement of a portion of SEQ ID NO: 1 and having a T_M of about 30°C to about 70°C;
- (c) isolating RNA from a cell; and
- (d) detecting the presence or absence of glycoprotein V RNA in said cellular RNA by the polymerase chain reaction utilizing said first and second primers, the presence of said glycoprotein V RNA indicating cells of a megakaryoblastic lineage.

12. An isolated DNA molecule comprising a polynucleotide sequence that is complementary to SEQ ID. NO. 1 and that is capable of producing a specific signal when hybridized to RNA isolated from a known megakaryoblastic cell under conditions of 0.5 M $\text{Na}_2\text{H}_2\text{PO}_4$, pH 7.2, 1 mM EDTA and 7% SDS at 65°C.

* * * * *