



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C11C	A2	(11) International Publication Number: WO 99/29814 (43) International Publication Date: 17 June 1999 (17.06.99)
(21) International Application Number: PCT/IT98/00351 (22) International Filing Date: 7 December 1998 (07.12.98) (30) Priority Data: RM97A000765 10 December 1997 (10.12.97) IT (71) Applicants (for all designated States except US): KEMINOVA ITALIANA S.R.L. [IT/IT]; Via G. Garibaldi, 19, I-25082 Botticino Sera (IT). MAYCOS ITALIANA S.A.S. [IT/IT]; Via N. Buonaparte, 9, I-46043 Castiglione Delle Stiviere (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): MORA, Valtiero [IT/IT]; Via Zamboni, 97, I-25100 Brescia (IT). COMINI, Palmiro [IT/IT]; Via Buonaparte, 9, I-46043 Castiglione Delle Stiviere (IT). RASTRELLI, Gianbattista [IT/IT]; Via Seselle, 37, I-25080 Manerba Del Garda (IT). (74) Agent: MASSARI, Marcello; Studio M. Massari S.r.l, Via di Campo Marzio, 46, I-00186 Roma (IT).		(81) Designated States: AU, BG, BR, CA, CN, CU, CZ, EE, HU, IL, IN, JP, KP, KR, MX, NO, NZ, PL, RO, RU, SG, SI, TR, UA, US, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHOD OF PREPARING LIPOPROTEINS, LIPOAMINOACIDS, LIPID ESTERS, AND GLUCOLIPIDS FROM OLIVE OIL AND/OR ITS CONSTITUENTS WHICH ARE CONDENSED WITH VEGETAL PROTEIN HYDROLYSATES OR AMINOACIDS OR ESTERIFICATED WITH FATTY ALCOHOLS OR SUGARS (57) Abstract A method of preparing lipoproteins, lipoaminoacids, lipid esters, and glucolipids from olive oil or one or more of its constituting fatty acids which are condensed with vegetal protein hydrolysates or aminoacids or esterificated with alcohols or sugars, said fatty acids being chlorinated fatty acids of the olive oil.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Method of preparing lipoproteins, lipoaminoacids, lipid esters, and glucolipids from olive oil and/or its constituents which are condensed with vegetal protein hydrolysates or aminoacids or esterificated with fatty alcohols or sugars

FIELD OF THE INVENTION

The present invention relates to a method of preparing a set of substances having new characteristics for use especially in the production of a very large range of products such as cosmetics, drugs, foods (emulsions, 5 creams, milks, ointments, shampoos, cleansing means for face/body, bath foams, etc.) industrial products (detergents for washing-machines, dishwashing machines, all-purpose cleansing agents), and chemical-pharmaceutical products (solvents, dispersing agents, 10 emulsifiers for aerosol, inhalations, environment deodorizers).

These preparations can be used under absolute safety conditions regarding both toxicological-allergic and 15 environment-friendly aspects since their constituents are already present in nature.

SUMMARY OF THE INVENTION

The products prepared by the method according to the invention can be considered to belong to four main 20 groups, namely lipoproteins, lipoaminoacids, lipid esters, and glucolipids.

The preparation of such groups starts from chlorides of fatty acids, particularly the individual fatty

acids forming the olive oil or a mixture of such fatty acids.

Among the fatty acids of the olive oil used individually or in combination in the method according to the invention oleic acid, palmitic acid, myristic acid, and stearic acid have the greatest importance. Such acids are condensed with vegetal protein hydrolysates to provide lipoproteins, with aminoacids to provide lipoaminoacids, with alcohols to provide lipid esters, and with sugars to provide glucolipids.

BEST MODE OF CARRYING OUT THE INVENTION

Some examples of the method according to the invention will be described herebelow.

In a preferred embodiment of the method of preparing a lipoprotein of fully vegetal nature with a large range of applications, as explained thereafter, the following steps are carried out by using a hydrolyzed vegetal protein, namely wheat gluten which is enzymatically hydrolyzed, in the condensation of the fatty acids of olive oil:

An enzymatic hydrolisate solution of wheat gluten (60-80% by weight referred to the finished product) is fed into a mixer of stainless steel and brought to a temperature between 45°C and 70°C; the following substances in the indicated proportions are added under stirring:

L.ascorbic acid	0.1 - 0.4%
-----------------	------------

glycerol	4 - 11%
----------	---------

and 2-4% potassium hydrate or sodium hydrate water	
--	--

solution having a concentration between 20 and 35% until a pH between 7.5 and 9.5 is reached while keeping the temperature at the initial value.

More particularly a substance with optimum characteristics is obtained by carrying out the following Example.

An amount of gluten hydrolisate equal to 70% by weight of the finished product is fed into a mixer of stainless steel and heated at 55°C under stirring.

When such temperature is reached, 0.3% L. ascorbic acid, 0.3% sodium EDTA, 10% glycerol, and 30% potassium hydrate (KOH) water solution are added until pH 8.5 is reached.

After about 15 minutes the reaction is started by adding 6% by weight of chlorinated fatty acids of olive oil still under stirring at constant temperature and the above pH 8.5 which is kept by a continuous, very gradual addition of 8% potassium hydrate solution (30% water solution). The reaction is continued under stirring at constant temperature for about 2 hours followed by a slow cooling. Finally, pH is adjusted to 7 by adding a total amount of 1% phosphoric acid and citric acid in a ratio of 1:1.

In order to obtain a lipoaminoacid to be used according to the applications of the present invention, the following method has been carried out by using a derivative of glutamic acid as aminoacid, namely sodium glutamate, and employing the following substances in the indicated weight proportion ranges:

An amount of 30-42% distilled water, 10-17% sodium

glutamate, 0.1-0.5% sodium EDTA, 8-11% 95°-ethyl alcohol are mixed in a reactor of stainless steel. Under stirring the mixture is brought to a temperature between 45°C and 75°C by adding very gradually an amount of 12-20% of a 30% potassium hydrate water solution and 16-25% chlorinated fatty acids of olive oil taking care of keeping pH between 8 and 12 and the temperature at the initial value. The reaction is continued for 2 to 5 hours followed by a cooling of the whole mixture and citric acid is added to adjust pH to 7-8. Finally, ethyl alcohol is distilled and the balance to 100 is formed by adding distilled water. Therefore, a lipoaminoacid having essentially the same characteristics as the lipoprotein both as far as its essential features and applications is concerned but capable of producing a persistent, creamy, more abundant foam has been prepared by the above-mentioned method.

A lipoaminoacid having the desired characteristics to an optimum extent has been prepared by reacting the above-mentioned substances as indicated above in the amounts of the following Example:

EXAMPLE of lipoaminoacid:

In a reactor of stainless steel the following substances are mixed:

Distilled water	36,7%
Sodium glutamate	14%
Sodium EDTA	0.3%
95°-Ethyl alcohol	10%

The mixture contained in the reactor is brought to

65°C under stirring and:

30% KOH water solution 17%

Chlorinated fatty acids of olive oil 21%

are gradually added still under stirring and keeping
pH to 9.5.

The reaction is continued for 3 hours and after
cooling citric acid is added until pH 7.5 is reached.
Finally, ethyl alcohol is distilled and the balance to
100 of the final product is formed, by adding distilled
water.

In order to obtain a lipid ester to be used according
to the applications of the invention, the following
method has been carried out by using a fatty alcohol,
namely oleic alcohol, and employing the following
substances in the indicated weight proportion ranges:

Oleic alcohol is fed into a reactor of stainless steel
provided with stirrer and distiller and, after having
brought the temperature to 60-96°C, chlorinated oleic
acid is added, the ratio between oleic alcohol and
oleic acid being 1:0.8-1:1.5. HCl fumes developed
during the reaction are then recovered by feeding them
to a separated alkali solution.

It should be noted that the addition of chlorinated
oleic acid must be gradual and very slow so as to be
concluded in a period of 2-5 hours. The reaction is
continued for 3-5 hours by keeping the initial
temperature constant. The distillation of the
hydrochloric acid should be complete. A washing is
carried out in the end by mixing with distilled water
in a proportion of about 20% of the total mass that is

eventually discharged from the bottom of the reactor.

A lipid ester having essentially the desired characteristics was prepared by the above-mentioned method.

5 A lipid ester essentially having the desired characteristics was prepared by reacting the above-mentioned substances according to the described method in the amounts listed in the following Example:

EXAMPLE of lipid ester:

10 Under the above-mentioned conditions oleic alcohol adjusted to 90° and chlorinated oleic alcohol in the ratio of 1:1.1 are fed by adding the latter very slowly during 4 hours into a reactor of stainless steel provided with stirrer and distillator. The
15 mixture is reacted for further 4 hours still at temperature of 90°C under distillation of the hydrochloric acid by conveying the HCl fumes to an alkali solution. The washing with 20% distilled water is carried out at the end of the distillation, as
20 indicated above.

By the Example described above a light, clear, colourless, stable to the oxidation, non-fatty oil is obtained so that quite soft, delicate cosmetics and dermatologic products, oils, emulsions, lipogels etc.
25 provided with excellent feel, softness and dermic compatibility can be prepared.

In the following method the chlorides of the olive oil acids are condensed with sugars derived, for example, from enzymatic hydrolysis of starch, barley, wheat and
30 the like.

In order to prepare a glucolipid to be used according to the applications of the present invention, the following method has been carried out by using tetraose malt as sugar and employing the following substances in the indicated weight proportion ranges:

An amount of 38-45% distilled water and 18-36% tetraose malt produced by enzymatic hydrolysis of a starch are fed into a reactor of stainless steel; an amount of 9-18% 95°-ethyl alcohol, referred to the finished product is added under stirring and the whole mixture is brought to a temperature between 45 and 75°C by adding 0.1-0.5% sodium EDTA and then very gradually 7-15% chlorinated oleic acid and 3-7% of a 30%-solution of sodium hydrate taking care of keeping pH between 7 and 9.

The reaction is continued for 2-5 hours still under stirring at the initial temperature. Final pH is adjusted between 6 and 8 and alcohol is distilled forming the balance to 100 by adding distilled water.

A glucolipid having the desired characteristics to an optimum extent has been prepared by reacting the above-mentioned substances as indicated above in the amounts of the following Example:

EXAMPLE of glucolipid:

In a reactor of stainless steel the following substances are mixed:

Distilled water	46%
Tetraose malt	24%
95°-Ethyl alcohol	15%
Sodium EDTA	0.3%

Still under stirring and bringing the content of the reactor to a temperature of 65°C an amount of 10% chlorinated oleic acid and 4,5% of a 30% NaOH water solution are gradually added keeping pH to 8.5.

5 The reaction is continued for 3 hours and thirty minutes and citric acid is added under constant temperature until pH 7.5 is reached. Finally, ethyl alcohol is distilled and the balance to 100 of the final product is formed by adding distilled water.

10 INDUSTRIAL APPLICABILITY

The finished product showed eudermic properties perfectly tolerated also by the most delicate skins and the mucosa. Furthermore it has hydrating,
15 emollient, protective properties and, as far as its inherent cleansing activity is concerned, it is also suitable for the preparation of cleansing creams and milks, shampoos, bath foams, personal cleanliness products and, more particularly, paedocosmetics.

Claims

1. A method of preparing lipoproteins, lipoaminoacids, lipid esters, and glucolipids from olive oil or one or more of its constituting fatty acids which are condensed with vegetal protein hydrolysates or aminoacids or esterificated with alcohols or sugars, wherein said fatty acids are chlorinated fatty acids of the olive oil.

2. The method according to claim 1, wherein said fatty acid is oleic acid.

3. The method according to claim 1, wherein said fatty acid is palmitic acid.

4. The method according to claim 1, wherein said fatty acid is myristic acid.

5. The method according to claim 1, wherein said fatty acid is stearic acid.

6. The method according to claim 1 for preparing a lipoprotein, wherein the following steps are provided: an enzymatic hydrolysis solution of wheat gluten (60-80% by weight referred to the finished product) is fed into a mixer of stainless steel and brought to a temperature between 45°C and 70°C; the following substances in the indicated proportions are added

under stirring:

L.ascorbic acid 0.1 - 0.4%

glycerol 4 - 11%

and 2-4% potassium hydrate or sodium hydrate water
5 solution having a concentration between 20 and 35%
until pH between 7.5 and 9.5 is reached while keeping
the temperature at the initial value.

7. The method according to claim 6, wherein an amount
10 of gluten hydrolisate equal to 70% by weight of the
finished product is fed into a mixer of stainless
steel and heated at 55°C under stirring; under
stirring 0.3% L.ascorbic acid, 0.3% sodium EDTA, 10%
glycerol, and 30% potassium hydrate (KOH) water
15 solution are then added until pH 8.5 is reached; after
about 15 minutes the reaction is started by adding 6%
by weight of chlorinated fatty acids of olive oil
still under stirring and keeping temperature and pH
constant by further adding continuously and gradually
20 8% of a 30% potassium hydrate water solution; the
reaction is continued under stirring at constant
temperature for about 2 hours followed by a slow
cooling; final pH is then adjusted to 7 by adding a
total amount of 1% phosphoric acid and citric acid in
25 a ratio of 1:1.

8. The method according to claim 1 for preparing a
lipoprotein, wherein sodium glutamate is used as
aminoacid and the following substances are employed in
30 the indicated weight proportion ranges:

an amount of 30-42% distilled water, 10-17% sodium glutamate, 0.1-0.5% sodium EDTA, 8-11% 95°-ethyl alcohol are mixed in a reactor of stainless steel; under stirring the mixture is brought to a temperature between 45°C and 75°C by adding an amount of 12-20% of a 30% potassium hydrate water solution and 16-25% chlorinated fatty acids of olive oil very slowly taking care of keeping pH between 8 and 12 and the temperature at the initial value; the reaction is then continued for 2 to 5 hours followed by a cooling of the whole mixture and citric acid is added to adjust pH to 7-8; finally, ethyl alcohol is distilled and the balance to 100 is formed by adding distilled water.

9. The method according to claim 8 for preparing a lipoaminoacid, wherein the following steps are provided:

in a reactor of stainless steel the following substances are mixed:

Distilled water	36,7%
Sodium glutamate	14%
Sodium EDTA	0.3%
95°-Ethyl alcohol	10%

the mixture contained in the reactor is brought to 65°C under stirring and:

30% KOH water solution	17%
Chlorinated fatty acids of olive oil	21%

are gradually added still under stirring and keeping pH to 9.5;

the reaction is continued for 3 hours and, after

cooling, citric acid is added to adjust pH to 7.5, and finally ethyl alcohol is distilled and the balance to 100 of the final product is formed by adding distilled water.

5

10. The method according to claims 1 and 2 for preparing a lipid ester, wherein a fatty alcohol, namely oleic alcohol, is used and the following substances are employed in the, indicated weight proportion ranges:

10

oleic alcohol is fed into a reactor of stainless steel provided with stirrer and distiller and, after having brought the temperature to 60-96°C, chlorinated oleic acid is added, the ratio between oleic alcohol and oleic acid being 1:0.8-1:1.5; hydrochloric acid fumes developed during the reaction are then recovered by feeding them to a separated alkali solution; it should be taken care that the addition of chlorinated oleic acid is gradual and very slow so as to be concluded in a period of 2-5 hours; the reaction is continued for 3-5 hours by keeping the initial temperature constant; the distillation of the hydrochloric acid should be complete; a washing is carried out in the end by mixing with distilled water in a proportion of about 20% of the total mass that is eventually discharged from the bottom of the reactor.

15

20

25

11. The method according to claim 10, wherein the following steps are provided:

30

oleic alcohol adjusted to 90° and chlorinated oleic

alcohol in the ratio of 1:1.1 are fed by adding the latter very slowly during 4 hours into a reactor of stainless steel provided with stirrer and distillator under the above-mentioned conditions; the mixture is reacted for further 4 hours still at temperature of 90°C under distillation of the hydrochloric acid by conveying the HCl fumes to an alkali solution; a washing with 20% distilled water is carried out at the end of the distillation, as indicated above.

12. The method according to claim 1, wherein the chlorides of the olive oil acids are condensed with sugars derived, for example, from enzymatic hydrolysis of starch, barley, wheat and the like.

13. The method according to claim 2 and 12, for preparing a glucolipid, wherein tetraose malt is used as sugar and the following substances are employed in the indicated weight proportion ranges:

an amount of 38-45% distilled water and 18-36% tetraose malt produced by enzymatic hydrolysis of a starch are fed into a reactor of stainless steel; an amount of 9-18% 95°-ethyl alcohol referred to the finished product is added under stirring and the whole mixture is brought to a temperature between 45 and 75°C by adding 0.1-0.5% sodium EDTA and then very gradually 7-15% chlorinated oleic acid and 3-7% of a 30%-solution of sodium hydrate taking care of keeping pH between 7 and 9; the reaction is continued for 2-5 hours still under stirring at the initial temperature

whereupon the final pH is adjusted between 6 and 8 and alcohol is distilled forming the balance to 100 by adding distilled water.

5 14. The method according to claim 13, wherein the following steps are provided:

in a reactor of stainless steel the following substances are mixed:

Distilled water	46%
10 Tetraose malt	24%
95°-Ethyl alcohol	15%
Sodium EDTA	0.3%

still under stirring and bringing the content of the reactor to a temperature of 65°C an amount of 10%
15 chlorinated oleic acid and 4,5% of a 30% NaOH water solution are gradually added keeping pH to 8.5; the reaction is continued for 3 hours and thirty minutes and citric acid is added under constant temperature until pH 7.5 is reached; finally, ethyl alcohol is
20 distilled and the balance to 100 of the final product is formed by adding distilled water.