



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2018/05/15 (87) Date publication PCT/PCT Publication Date: 2018/11/22 (85) Entrée phase nationale/National Entry: 2019/10/28 (86) N° demande PCT/PCT Application No.: US 2018/032726 (87) N° publication PCT/PCT Publication No.: 2018/213281 (30) Priorité/Priority: 2017/05/15 (US62/506,226)	(51) Cl.Int./Int.Cl. <i>A61K 31/135</i> (2006.01), <i>C07C 217/64</i> (2006.01) (71) Demandeur/Applicant: COGNITION THERAPEUTICS, INC., US (72) Inventeurs/Inventors: RISHTON, GILBERT M., US; LOOK, GARY C., US; CATALANO, SUSAN M., US (74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.
--	--

(54) Titre : COMPOSITIONS POUR LE TRAITEMENT DE MALADIES NEURODEGENERATIVES
(54) Title: COMPOSITIONS FOR TREATING NEURODEGENERATIVE DISEASES

(57) **Abrégé/Abstract:**

The present disclosure relates to novel compounds, pharmaceutical compositions containing the compounds and methods of using the compounds and pharmaceutical compositions for treating neurodegenerative diseases, including Alzheimer's disease and cognitive decline. Methods for inhibiting synapse number decline or membrane trafficking abnormalities associated with exposure of a neuronal cell to Abeta species are also disclosed.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau

(43) International Publication Date

22 November 2018 (22.11.2018)



(10) International Publication Number

WO 2018/213281 A1



- (51) International Patent Classification:
A61K 31/135 (2006.01) C07C 217/64 (2006.01)
- (21) International Application Number:
PCT/US2018/032726
- (22) International Filing Date:
15 May 2018 (15.05.2018)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
62/506,226 15 May 2017 (15.05.2017) US
- (71) Applicant: COGNITION THERAPEUTICS, INC.
[US/US]; 2403 Sidney Street, Suite 261, Pittsburgh, Penn-
sylvania 15203 (US).
- (72) Inventors: RISHTON, Gilbert M.; 401 S. Barrington
Avenue, Apt. 124, Los Angeles, California 90049 (US).
LOOK, Gary C.; 2576 Amethyst Drive, Santa Clara, Cal-
ifornia 95051 (US). CATALANO, Susan M.; 2536 A Sid-
ney Street, Pittsburgh, Pennsylvania 15203 (US).
- (74) Agent: MANDELBLATT, Jill M.; PEPPER HAMILTON
LLP, 500 Grant Street, Suite 5000, Pittsburgh, Pennsylvania
15219-2507 (US).
- (81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).
- (54) Title: COMPOSITIONS FOR TREATING NEURODEGENERATIVE DISEASES
- (57) Abstract: The present disclosure relates to novel compounds, pharmaceutical compositions containing the compounds and methods of using the compounds and pharmaceutical compositions for treating neurodegenerative diseases, including Alzheimer's disease and cognitive decline. Methods for inhibiting synapse number decline or membrane trafficking abnormalities associated with exposure of a neuronal cell to Abeta species are also disclosed.



WO 2018/213281 A1

COMPOSITIONS FOR TREATING NEURODEGENERATIVE DISEASES

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] The present application claims benefit of and priority to U.S. Provisional No. 62/506,226 entitled “COMPOSITIONS FOR TREATING NEURODEGENERATIVE DISEASES,” filed May 15, 2017, the contents of which is hereby incorporated by reference in its entirety.


GOVERNMENT INTEREST

[0002] This invention was made with support from the U. S. government under a grant from the National Institute On Aging of the National Institute of Health , grant number U01AG047059. The U. S. government has certain rights in this invention.

SUMMARY

[0003] Various embodiments provide novel compounds, pharmaceutical compositions comprising such compounds, and methods for inhibiting or restoring synapse loss in neuronal cells, modulating a membrane trafficking change in neuronal cells, and treating cognitive decline and neurodegenerative diseases and disorders.

[0004] Some embodiments of the present disclosure are directed to a compound of Formula I or pharmaceutically acceptable salt thereof:




wherein:

each of R^a , R^b , R^c , R^d and R^e is independently selected from the group consisting of H, hydroxyl, halo, alkyl, alkoxy, CF_3 , SO_2CH_3 , and morpholino;


R^1 is selected from the group consisting of hydrogen, alkyl, phenyl, or $-CH=C(CH_3)_2$; and

R^2 is an optionally substituted cyclic amino group.


[0005] Some embodiments of the present disclosure are directed to a compound selected from the group consisting of




,




,




,




,











,




,



,





[0006] Embodiments herein describe a pharmaceutical composition comprising: a compound according to any embodiment described herein, or a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable carrier or diluent

[0007] Some embodiments describe a method of treating Alzheimer's disease (AD) comprising administering to the subject a therapeutically effective amount of a compound or a pharmaceutical composition according to any embodiment described herein.

[0008] Some embodiments describe a method of inhibiting cognitive decline in a subject exhibiting, or at risk of exhibiting, cognitive decline, comprising administering a therapeutically effective amount of a compound or a pharmaceutical composition according to any embodiment described herein.

[0009] Some embodiments describe a method of inhibiting amyloid beta effect on a neuronal cell comprising administering to a subject in need thereof, a therapeutically effective amount of a pharmaceutical composition according to any embodiment described herein.

[0010] Some embodiments describe a method of treating mild cognitive impairment in Alzheimer's disease in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition according to any embodiment described herein.

[0011] Some embodiments describe use of a compound according to according to any embodiment described herein, in the manufacture of a medicament for the treatment of Alzheimer's disease.

[0012] Some embodiments describe a compound according to any embodiment described herein, for use in the treatment of Alzheimer's disease.

[0013] Some embodiments describe a compound according to any embodiment described herein, for use in medical therapy.

[0014] Some embodiments describe a pharmaceutical composition comprising a therapeutically effective amount of a compound according to any embodiment described herein, and pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.

DETAILED DESCRIPTION

[0015] This invention is not limited to the particular processes, compositions, or methodologies described, as these may vary. The terminology used in the description is for the purpose of describing the particular versions or embodiments only, and is not intended to limit the scope of the present invention. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art. All publications mentioned herein, are incorporated by reference in their entirety. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

Definitions

[0016] Where a range of values is provided, it is intended that each intervening value between the upper and lower limit of that range and any other stated or intervening value in that stated range is encompassed within the disclosure. For example, if a range of 1 μm to 8 μm is stated, it is intended that 2 μm , 3 μm , 4 μm , 5 μm , 6 μm , and 7 μm are also explicitly disclosed.

[0017] At various places in the present specification, substituents of compounds of the disclosure are disclosed in groups or in ranges. It is specifically intended that embodiments of the disclosure include each and every individual subcombination of the members of such groups and ranges. For example, the term "C₁₋₆ alkyl" is specifically intended to individually disclose, *e.g.* methyl (C₁ alkyl), ethyl (C₂ alkyl), propyl (C₃ alkyl), butyl (C₄ alkyl), pentyl (C₅ alkyl), and hexyl (C₆ alkyl) as well as, *e.g.* C₁-C₂ alkyl, C₁-C₃ alkyl, C₁-C₄ alkyl, C₂-C₃ alkyl, C₂-C₄ alkyl, C₃-C₆ alkyl, C₄-C₅ alkyl, and C₅-C₆ alkyl.

[0018] The articles "a" and "an" as used herein, mean "one or more" or "at least one," unless otherwise indicated. That is, reference to any element of the present invention by the indefinite article "a" or "an" does not exclude the possibility that more than one of the element is present.

[0019] As used herein, the term “about” means plus or minus 10% of the numerical value of the number with which it is being used. Therefore, about 50 mL means in the range of 45 mL-55 mL.


[0020] “Abeta species” or “A β ” shall include compositions comprising soluble amyloid peptide-containing components such as Abeta monomers, Abeta oligomers, or complexes of Abeta peptide (in monomeric, dimeric or polymeric form) with other soluble peptides or proteins as well as other soluble Abeta assemblies, including any processed product of amyloid precursor protein. Soluble A β oligomers are known to be neurotoxic. Even A β 1-42 dimers are known to impair synaptic plasticity in mouse hippocampal slices. In one theory known in the art, native A β 1-42 monomers are considered neuroprotective, and self-association of A β monomers into soluble Abeta oligomers is required for neurotoxicity. However, certain A β mutant monomers (arctic mutation (E22G)) are reported to be associated with familial Alzheimer’s Disease.

[0021] Unless specifically indicated, the term “active ingredient” is to be understood as referring to a compound according to any embodiment describe herein.

[0022] “Administering,” or “administration” and the like, when used in conjunction with the compounds of the disclosure refers to providing the compounds or pharmaceutical compositions according to any of the embodiments described herein, to a subject in need of treatment. Preferably the subject is a mammal, more preferably a human. The present invention comprises administering the pharmaceutical composition of the invention alone or in conjunction with another therapeutic agent. When a pharmaceutical composition of the invention is administered in conjunction with another therapeutic agent, the pharmaceutical composition of the invention and the other therapeutic agent. can be administered at the same time or different times.

[0023] The term “agonist” refers to a compound, the presence of which results in a biological activity of a receptor that is the same as the biological activity resulting from the presence of a naturally occurring ligand for the receptor.

[0024] The term “alkanoyl” or “alkylcarbonyl” as used herein, is meant to refer to

an alkyl group attached to a carbonyl radical. An example of an alkanoyl is .

[0025] As used herein, the term “alkyl” is meant to refer to a saturated hydrocarbon group which is straight-chained or branched. Example alkyl groups include, but are not limited to, methyl (Me), ethyl (Et), propyl (*e.g.* n-propyl and isopropyl), butyl (*e.g.* n-

butyl, isobutyl, t-butyl), pentyl (*e.g.* n-pentyl, isopentyl, neopentyl), and the like. An alkyl group can contain from 1 to about 20, from 2 to about 20, from 1 to about 10, from 1 to about 8, from 1 to about 6, from 1 to about 4, or from 1 to about 3 carbon atoms. "C₁-C₁₀ alkyl" or "C₁₋₁₀ alkyl", is intended to include C₁, C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, and C₁₀ alkyl groups. Additionally, for example, "C₁-C₆ alkyl" or "C₁₋₆ alkyl" denotes alkyl having 1 to 6 carbon atoms. The term "alkylene" refers to a divalent alkyl linking group. An example of alkylene is methylene (CH₂).

[0026] As used herein, "alkenyl" is intended to include hydrocarbon chains of either straight or branched configuration with one or more, preferably one to three, carbon-carbon double bonds that may occur in any stable point along the chain. For example, "C₂-C₆ alkenyl" or "C₂₋₆ alkenyl" (or alkenylene), is intended to include C₂, C₃, C₄, C₅, and C₆ alkenyl groups. Examples of alkenyl include, but are not limited to, ethenyl, 1-propenyl, 2-propenyl, 2-butenyl, 3-butenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 2-methyl-2-propenyl, and 4-methyl-3-pentenyl.

[0027] The term "alkoxy" or "alkyloxy" refers to an -O-alkyl group. "C₁-C₆ alkoxy" or "C₁₋₆ alkoxy" (or alkyloxy), is intended to include C₁, C₂, C₃, C₄, C₅, and C₆ alkoxy groups. Examples of alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy (*e.g.*, n-propoxy and isopropoxy), and t-butoxy.

[0028] The term "alkoxyalkoxy" refers to an alkoxy group attached to an alkoxy group. An example of an alkoxy group includes -O-(CH₂)₂-OCH₃.

[0029] As used herein, "alkynyl" is intended to include hydrocarbon chains of either straight or branched configuration having one or more, preferably one to three, carbon-carbon triple bonds that may occur in any stable point along the chain. For example, "C₂-C₆ alkynyl" is intended to include C₂, C₃, C₄, C₅, and C₆ alkynyl groups; such as ethynyl, propynyl, butynyl, pentynyl, and hexynyl.

[0030] As used herein, an "amyloid beta effect," for example, a "nonlethal amyloid beta effect," or "Abeta oligomer effect," refers to an effect, particularly a nonlethal effect, on a cell that is contacted with an Abeta species. For example, it has been found that when a neuronal cell is contacted with a soluble Amyloid-beta ("Abeta") oligomer, the oligomers bind to a subset of synapses on a subset of neuronal cells *in vitro*. This binding can be quantified in an assay measuring Abeta oligomer binding *in vitro* for example. Another documented effect of Abeta species is a reduction in synapse number, which has been reported to be about 18% in the human hippocampus (Scheff et al, 2007) and can be quantified (for example, in an assay measuring synapse number). As another example, it has

been found that, when a neuronal cell is contacted with an Amyloid-beta (“Abeta”) oligomer, membrane trafficking is modulated and alteration of membrane trafficking ensues. This abnormality can be visualized with many assays, including but not limited to, an MTT assay. For example, yellow tetrazolium salts are endocytosed by cells and the salts are reduced to insoluble purple formazan by enzymes located within vesicles in the endosomal pathway. The level of purple formazan is a reflection of the number of actively metabolizing cells in culture, and reduction in the amount of formazan is taken as a measure of cell death or metabolic toxicity in culture. When cells that are contacted with a yellow tetrazolium salt are observed through a microscope, the purple formazan is first visible in intracellular vesicles that fill the cell. Over time, the vesicles are exocytosed and the formazan precipitates as needle-shaped crystals on the outer surface of the plasma membrane as the insoluble formazan is exposed to the aqueous media environment. Still other effects of Abeta species include cognitive decline, such as a decline in the ability to form new memories and memory loss which can be measured in assays using animal models *in vivo*. In some embodiments, an Abeta effect is selected from Abeta oligomer-induced synaptic dysfunction, for example, as seen in an *in vitro* assay, such as a membrane trafficking assay, or a synapse loss assay, or Abeta oligomer mediated sigma-2 receptor activation of caspase-3, or Abeta induced neuronal dysfunction, Abeta mediated decrease in long term potentiation (LTP), or in cognitive decline in a behavioral assay, or in a patient in need thereof.

[0031] In some embodiments, a test compound is said to be effective to treat cognitive decline or a disease associated therewith when it can inhibit an effect associated with soluble Abeta oligomer species on a neuronal cell more than about 10%, preferably more than 15%, and preferably more than 20% as compared to a negative control. In some embodiments, a test agent is said to be effective when it can inhibit a processed product of amyloid precursor protein-mediated effect more than about 10%, preferably more than 15%, and preferably more than 20% as compared to a positive control. Although the present specification focuses on inhibition of nonlethal effects of Abeta species, such as abnormalities in neuronal metabolism and synapse number reduction, these are shown to correlate with cognitive function and are furthermore expected, over time, to result in reduction (compared to untreated subjects) of downstream measurable symptoms of amyloid pathology, notably clinical symptoms such as 1) fibril or plaque accumulation measured by amyloid imaging agents such as fluorbetapir, PittB or any other imaging agent, 2) synapse loss or cell death as measured by glucose hypometabolism detected with FDG-PET, 3) changes in protein expression or metabolite amount in the brain or body detectable by

imaging or protein/metabolite detection in cerebrospinal fluid, brain biopsies or plasma obtained from patients by ELISA, (such as changes in levels and or ratios of Abeta 42, phosphorylated tau, total tau measured by ELISA, or patterns of protein expression changes detectable in an ELISA panel), 4) cerebral vascular abnormalities as measured by the presence of vascular edema or microhemorrhage detectable by MRI and any other symptoms detectable by imaging techniques, and 5) cognitive loss as measured by any administered cognitive test such as ADAS-Cog, MMSE, CBIC or any other cognitive testing instrument.

[0032] The term “animal” as used herein, includes, but is not limited to, humans and non-human vertebrates such as wild, experimental, domestic and farm animals and pets.

[0033] The term “antagonist” refers to an entity, *e.g.* a compound, antibody or fragment, the presence of which results in a decrease in the magnitude of a biological activity of a receptor. In certain embodiments, the presence of an antagonist results in complete inhibition of a biological activity of a receptor. As used herein, the term “sigma-2 receptor antagonist” is used to describe a compound that acts as a “functional antagonist” at the sigma-2 receptor in that it blocks Abeta effects, for example, Abeta oligomer-induced synaptic dysfunction, for example, as seen in an *in vitro* assay, such as a membrane trafficking assay, or a synapse loss assay, or Abeta oligomer mediated sigma-2 receptor activation of caspase-3, or in a behavioral assay, or in a patient in need thereof. The functional antagonist may act directly by inhibiting binding of, for example, an Abeta oligomer to a sigma-2 receptor, or indirectly, by interfering with downstream signaling resultant from Abeta oligomer binding the sigma-2 receptor.

[0034] As used herein, “aryl” refers to monocyclic or polycyclic (*e.g.* having 2, 3 or 4 fused rings) aromatic hydrocarbons such as, for example, phenyl, naphthyl, anthracenyl, phenanthrenyl, indanyl, indenyl, and the like. In some embodiments, aryl groups have from 6 to about 20 carbon atoms. In some embodiments, aryl groups have from 5 to about 10 carbon atoms.

[0035] As used herein, “arylalkyl” refers to an aryl group attached to an alkyl radical. In preferred embodiments the alkyl is a C₁₋₆ alkyl.

[0036] The term “aroyl” or “arylcabonyl” as used herein, refers to an aryl group attached to a carbonyl radical. Examples of aroyl include but are not limited to benzoyl.

[0037] As used herein the term “brain penetrability” refers to the ability of a drug, antibody or fragment, to cross the blood-brain barrier. In some embodiments, an animal pharmacokinetic (pK) study, for example, a mouse pharmacokinetic/blood-brain barrier study can be used to determine or predict brain penetrability. In some embodiments various


concentrations of a compound or pharmaceutical composition according to any embodiment described herein, can be administered, for example at 3, 10 and 30 mg/kg, for example p.o. for 5 days and various pK properties are measured, e.g., in an animal model. In some embodiments, dose related plasma and brain levels are determined. In some embodiments, brain C_{max} > 100, 300, 600, 1000, 1300, 1600, or 1900 ng/mL. In some embodiments good brain penetrability is defined as a brain/plasma ratio of > 0.1, > 0.3, > 0.5, > 0.7, > 0.8, > 0.9, preferably >1, and more preferably > 2, >5, or > 10. In other embodiments, good brain penetrability is defined as greater than about 0.1%, 1%, 5%, greater than about 10%, and preferably greater than about 15% of an administered dose crossing the BBB after a predetermined period of time. In certain embodiments, the dose is administered orally (p.o.). In other embodiments, the dose is administered intravenously (i.v.), prior to measuring pK properties. Pharmacokinetic assays and brain penetrability are described in Example 7.

[0038] As used herein, “cognitive decline” can be any negative change in an animal’s cognitive function. For example cognitive decline, includes but is not limited to, memory loss (e.g. behavioral memory loss), failure to acquire new memories, confusion, impaired judgment, personality changes, disorientation, or any combination thereof. A compound that is effective to treat cognitive decline can be thus effective by restoring long term neuronal potentiation (LTP) or long term neuronal depression (LTD) or a balance of synaptic plasticity measured electrophysiologically; inhibiting, treating, and/or abatement of neurodegeneration; inhibiting, treating, and/or abatement of general amyloidosis; inhibiting, treating, abatement of one or more of amyloid production, amyloid assembly, amyloid aggregation, and amyloid oligomer binding; inhibiting, treating, and/or abatement of a nonlethal effect of one or more of Abeta species on a neuron cell (such as synapse loss or dysfunction and abnormal membrane trafficking); and any combination thereof. Additionally, that compound can also be effective in treating Abeta related neurodegenerative diseases and disorders including, but not limited to dementia, including but not limited to Alzheimer’s Disease (AD) including mild Alzheimer’s disease, Down’s syndrome, vascular dementia (cerebral amyloid angiopathy and stroke), dementia with Lewy bodies, HIV dementia, Mild Cognitive Impairment (MCI); Age-Associated Memory Impairment (AAMI); Age-Related Cognitive Decline (ARCD), preclinical Alzheimer’s Disease (PCAD); and Cognitive Impairment No Dementia (CIND).


[0039] As used herein, the term “contacting” refers to the bringing together or combining of molecules (or of a molecule with a higher order structure such as a cell or cell membrane) such that they are within a distance that allows for intermolecular interactions

such as the non-covalent interaction between two peptides or one protein and another protein or other molecule, such as a small molecule. In some embodiments, contacting occurs in a solution in which the combined or contacted molecules are mixed in a common solvent and are allowed to freely associate. In some embodiments, the contacting can occur at or otherwise within a cell or in a cell-free environment. In some embodiments, the cell-free environment is the lysate produced from a cell. In some embodiments, a cell lysate may be a whole-cell lysate, nuclear lysate, cytoplasm lysate, and combinations thereof. In some embodiments, the cell-free lysate is lysate obtained from a nuclear extraction and isolation wherein the nuclei of a cell population are removed from the cells and then lysed. In some embodiments, the nuclei are not lysed, but are still considered to be a cell-free environment. The molecules can be brought together by mixing such as vortexing, shaking, and the like.

[0040] The term “cyclic amino” or “cyclic amino group” as used herein, is a heterocycloalkyl or heteroaryl group containing a nitrogen radical, thus allowing bonding through the nitrogen atom. The group can be represented by the formula:



[0042] The term “cycloalkanoyl” or “cycloalkylcarbonyl” as used herein, is meant to describe a cycloalkyl group attached to a carbonyl radical. Examples of cycloalkanoyl



[0043] As used herein, “cycloalkyl” refers to non-aromatic cyclic hydrocarbons including cyclized alkyl, alkenyl, and alkynyl groups that contain up to 20 ring-forming carbon atoms. Cycloalkyl groups can include mono- or polycyclic (*e.g.* having 2, 3 or 4 fused rings) ring systems as well as spiro ring systems. A cycloalkyl group can contain from 3 to about 15, from 3 to about 10, from 3 to about 8, from 3 to about 6, from 4 to about 6, from 3 to about 5, or from 5 to about 6 ring-forming carbon atoms. Ring-forming carbon atoms of a cycloalkyl group can be optionally substituted by oxo or sulfido. Example of cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl,

cyclohexyl, cycloheptyl, cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptatrienyl, norbornyl, norpinyl, norcarnyl, adamantyl, and the like. Also included in the definition of cycloalkyl are moieties that have one or more aromatic rings fused (*i.e.* having a bond in common with) to the cycloalkyl ring, for example, benzo or thienyl derivatives of cyclopentane, cyclopentene, cyclohexane, and the like (*e.g.* 2,3-dihydro-1H-indene-1-yl, or 1H-inden-2(3H)-one-1-yl). Preferably, “cycloalkyl” refers to cyclized alkyl groups that contain up to 20 ring-forming carbon atoms. Examples of cycloalkyl preferably include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, and the like

[0044] The term “cycloalkylalkyl” refers to a cycloalkyl group attached to an alkyl radical. In preferred embodiments the alkyl is a C₁₋₆ alkyl.

[0045] The term “drug-like properties” is used herein, to describe the pharmacokinetic and stability characteristics of a compound upon administration; including brain penetrability, metabolic stability and/or plasma stability.

[0046] As used herein, the term “effective amount” refers to an amount that results in measurable inhibition of at least one symptom or parameter of a specific disorder or pathological process. For example, an amount of a disclosure compound according to any embodiment described herein, that provides a measurably lower synapse reduction in the presence of Abeta oligomer qualifies as an effective amount because it reduces a pathological process even if no clinical symptoms of amyloid pathology are altered, at least immediately.

[0047] As used herein, “halo” or “halogen” includes fluorine, chlorine, bromine, and iodine.

[0048] As used herein, “haloalkoxy” represents a haloalkyl group as defined herein, with the indicated number of carbon atoms, attached through an oxygen bridge. For example, “C₁-C₆ haloalkoxy” or “C₁₋₆ haloalkoxy”, is intended to include C₁, C₂, C₃, C₄, C₅, and C₆ haloalkoxy groups. An example haloalkoxy group is OCF₃. As used herein, “trihalomethoxy” refers to a methoxy group having three halogen substituents. Examples of trihalomethoxy groups include, but are not limited to, -OCF₃, -OCClF₂, -OCCl₃, and the like.


[0049] As used herein, “haloalkyl” is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with one or more halogens. Example haloalkyl groups include, but are not limited to, CF₃, C₂F₅, CHF₂, CCl₃, CHCl₂, C₂Cl₅, CH₂CF₃, and the like.

[0050] As used herein, “heteroaryl” groups refer to an aromatic heterocycle having up to 20 ring-forming atoms and having at least one heteroatom ring member (ring-forming atom) such as sulfur, oxygen, or nitrogen. In some embodiments, the heteroaryl

group has at least one or more heteroatom ring-forming atoms each independently selected from sulfur, oxygen, and nitrogen. Heteroaryl groups include monocyclic and polycyclic (*e.g.* having 2, 3 or 4 fused rings) systems. Examples of heteroaryl groups include without limitation, pyridyl (a.k.a. pyridinyl), pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, furyl, quinolyl, isoquinolyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrol (a.k.a. pyrrolyl), oxazolyl, benzofuryl, benzothienyl, benzthiazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, indazolyl, 1,2,4-thiadiazolyl, isothiazolyl, benzothienyl, purinyl, carbazolyl, benzimidazolyl, indolinyl, and the like. In some embodiments, the heteroaryl group has from 1 to about 20 carbon atoms, and in further embodiments from about 1 to about 5, from about 1 to about 4, from about 1 to about 3, from about 1 to about 2, carbon atoms as ring-forming atoms. In some embodiments, the heteroaryl group contains 3 to about 14, 3 to about 7, or 5 to 6 ring-forming atoms. In some embodiments, the heteroaryl group has 1 to about 4, 1 to about 3, or 1 to 2 heteroatoms.

[0051] The term “heterocycloalkoxy” as used herein, refers to an -O-

heterocycloalkyl group. An example of a heterocycloalkoxy group is



[0052] As used herein, “heterocycloalkyl” or “heterocyclyl” refers to a non-aromatic heterocyclyl group having up to 20 ring-forming atoms including cyclized alkyl, alkenyl, and alkynyl groups where one or more of the ring-forming carbon atoms is replaced by a heteroatom such as an O, N, or S atom. Heterocycloalkyl groups can be mono or polycyclic (*e.g.* both fused and spiro systems). For example, “heterocycloalkyl” groups include morpholino, thiomorpholino, piperazinyl, tetrahydrofuranyl, tetrahydrothienyl, 2,3-dihydrobenzofuryl, 1,3-benzodioxole, benzo-1,4-dioxane, piperidinyl, pyrrolidinyl, isoxazolidinyl, isothiazolidinyl, pyrazolidinyl, oxazolidinyl, thiazolidinyl, imidazolidinyl, pyrrolidin-2-one-3-yl, and the like. Ring-forming carbon atoms and heteroatoms of a heterocycloalkyl group can be optionally substituted by oxo or sulfido. For example, a ring-forming S atom can be substituted by 1 or 2 oxo (*i.e.* form a S(O) or S(O)₂). For example, a ring-forming C atom can be substituted by oxo (*i.e.* form carbonyl). Also included in the definition of heterocycloalkyl are moieties that have one or more aromatic rings fused (*i.e.* having a bond in common with) to the nonaromatic heterocyclic ring, for example pyridinyl, thiophenyl, phthalimidyl, naphthalimidyl, and benzo derivatives of heterocycles such as indoline, isoindoline, isoindolin-1-one-3-yl, 4,5,6,7-tetrahydrothieno[2,3-c]pyridine-5-yl, 5,6-dihydrothieno[2,3-c]pyridin-7(4H)-one-5-yl, and 3,4-dihydroisoquinolin-1(2H)-one-3yl

groups. Ring-forming carbon atoms and heteroatoms of the heterocycloalkyl group can be optionally substituted by oxo or sulfido. In some embodiments, the heterocycloalkyl group has from 2 to about 20 carbon atoms or 3 to 20 carbon atoms. In some embodiments, the heterocycloalkyl group contains 3 to about 14, 3 to about 7, or 5 to 6 ring-forming atoms. In some embodiments, the heterocycloalkyl group has 1 to 4 heteroatoms. In some embodiments, the heterocycloalkyl group contains 0 to 3 double bonds. In some embodiments, the heterocycloalkyl group contains 0 to 2 triple bonds.

[0053] In the present application, the term "high affinity" is intended to mean a compound which exhibits a K_i value of less than 600 nM, 500 nM, 400 nM, 300 nM, 200 nM, less than 150 nM, less than 100 nM, less than 80 nM, less than 60 nM, or preferably less than 50 nM in a sigma receptor binding assay, for example against [3H]-DTG, as disclosed by Weber et al., Proc. Natl. Acad. Sci (USA) 83: 8784-8788 (1986), incorporated herein by reference, which measures the binding affinity of compounds toward both the sigma-1 and sigma-2 receptor sites. Especially preferred compounds exhibit K_i values of less than about 150 nM, preferably less than 100 nM, less than about 60 nM, less than about 10 nM, or less than about 1 nM against [3H]-DTG.

[0054] The terms "hydroxyl" and "hydroxy" are used interchangeably to mean an OH group.

[0055] The term "improves" is used to convey that the disclosure changes either the characteristics and/or the physical attributes of the tissue to which it is being provided, applied or administered. The term "improves" may also be used in conjunction with a disease state such that when a disease state is "improved" the symptoms or physical characteristics associated with the disease state are diminished, reduced, eliminated, delayed or averted.

[0056] The term "inhibiting" includes the blockade, aversion of a certain result or process, or the restoration of the converse result or process. In terms of prophylaxis or treatment by administration of a compound of the disclosure, "inhibiting" includes protecting against (partially or wholly) or delaying the onset of symptoms, alleviating symptoms, or protecting against, diminishing or eliminating a disease, condition or disorder.

[0057] The term "inhibiting trafficking deficits" refers to the ability to block soluble A β oligomer-induced membrane trafficking deficits in a cell, preferably a neuronal cell. A compound capable of inhibiting trafficking deficits has an $EC_{50} < 20 \mu M$, less than 15 μM , less than 10 μM , less than 5 μM , and preferably less than 1 μM in the membrane

trafficking assay, and further is capable of at least 50%, preferably at least 60%, and more preferably at least 70% maximum inhibition of the Abeta oligomer effects of soluble Abeta oligomer-induced membrane trafficking deficits, for example, as described in Example 6.

[0058] The term “log P” refers to the partition coefficient of a compound. The partition coefficient is the ratio of concentrations of un-ionized compound in each of two solution phases, for example, octanol and water. To measure the partition coefficient of ionizable solute compounds, the pH of the aqueous phase is adjusted such that the predominant form of the compound is un-ionized. The logarithm of the ratio of concentrations of the un-ionized solute compound in the solvents is called log P. The log P is a measure of lipophilicity. For example,

$$\log P_{\text{oct/wat}} = \log ([\text{solute}]_{\text{octanol}}/[\text{solute}]_{\text{un-ionized, water}}).$$

[0059] As used herein the term “metabolic stability” refers to the ability of a compound to survive first-pass metabolism (intestinal and hepatic degradation or conjugation of a drug administered orally). This can be assessed, for example, *in vitro* by exposure of the compounds to mouse or human hepatic microsomes. In some embodiments, good metabolic stability refers to a $t_{1/2} > 5$ min, > 10 min, > 15 minutes, > 20 minutes, and preferably > 30 min upon exposure of a compound to mouse or human hepatic microsomes. In some embodiments, good metabolic stability refers to an Intrinsic Clearance Rate (Cl_{int}) of < 300 uL/min/mg, preferably ≤ 200 uL/min/mg, and more preferably ≤ 100 uL/min/mg.

[0060] The term “n-membered” where n is an integer typically describes the number of ring-forming atoms in a moiety where the number of ring-forming atoms is n. For example, pyridine is an example of a 6-membered heteroaryl ring and thiophene is an example of a 5-membered heteroaryl group.


[0061] As used herein, the term “natural ligand” refers to a ligand present in a subject that can bind to a protein, receptor, membrane lipid or other binding partner *in vivo* or that is replicated *in vitro*. The natural ligand can be synthetic in origin, but must also be present naturally and without human intervention in the subject. For example, Abeta oligomers are known to exist in human subjects. Therefore the Abeta oligomers found in a subject would be considered natural ligands. The binding of Abeta oligomers to a binding partner can be replicated *in vitro* using recombinant or synthetic techniques, but the Abeta oligomer would still be considered a natural ligand regardless of how the Abeta oligomer is prepared or manufactured. A synthetic small molecule that can also bind to the same binding partner is not a natural ligand if it does not exist in a subject. For example, compounds which

are described herein, are not normally present in a subject, and, therefore, would not be considered natural ligands.

[0062] As used herein, the term “a neuronal cell” can be used to refer to a single cell or to a population of cells. In some embodiments, the neuronal cell is a primary neuronal cell. In some embodiments, the neuronal cell is an immortalized or transformed neuronal cell or a stem cell. A primary neuronal cell is a neuronal cell that cannot differentiate into other types of neuronal cells, such as glia cells. A stem cell is one that can differentiate into neurons and other types of neuronal cells such as glia. In some embodiments, assays utilize a composition comprising at least one neuronal cell is free of glia cells. In some embodiments, the composition comprises less than about 30%, 25%, 20%, 15%, 10%, 5%, or 1% of glia cells, which are known to internalize and accumulate Abeta. The primary neuronal cell can be derived from any area of the brain of an animal. In some embodiments, the neuronal cell is a hippocampal or cortical cell. The presence of glia cells can be determined by any method. In some embodiments, glia cells are detected by the presence of GFAP and neurons can be detected by staining positively with antibodies directed against MAP2.

[0063] As used herein, the term “optionally substituted” means that substitution is optional and therefore includes both unsubstituted and substituted atoms and moieties. A “substituted” atom or moiety indicates that any hydrogen on the designated atom or moiety can be replaced with a selection from the indicated substituent group, provided that the normal valence of the designated atom or moiety is not exceeded, and that the substitution results in a stable compound. For example, if a methyl group (*i.e.* CH₃) is optionally substituted, then up to 3 hydrogen atoms on the carbon atom can be replaced with substituent groups. Substituent groups include, but are not limited to, alkanoyl, alkoxy, alkoxyalkyl, (alkoxy)alkoxyalkyl, alkoxycarbonyl, alkyl, aryloxy, aryloyl, cycloalkanoyl, substituted or unsubstituted C₃-C₁₀ cycloalkyl, -OC(O)NCH(CH₃)₂, (N,N-dimethylamino)pyridinyl, (N,N-dimethylamino)sulfonyl, halo, heterocyclyl, (heterocyclyl)alkoxyalkyl, heterocycloalkyl, hydroxyl, hydroxyalkyl, methylpiperidinyl, methylsulfonyl, methylsulfonylphenyl, morpholinylpyridinyl, optionally substituted C₁-C₁₀ alkyl, optionally substituted C₅-C₁₀ aryl, optionally substituted C₃-C₁₀ heteroaryl, perfluoroalkyl, phenyl, piperidinyl, pyrrolidinylpyridinyl, tetrahydropyranyl, CF₃. A substituted alkyl group for example indicates that one or more hydrogen atoms on the alkyl group is replaced with a substituent group, selected from but not limited to, halo, hydroxyl, alkoxy, heterocycloalkoxy, alkoxyalkoxy, C(O)OMe, and C(O)OEt. A substituted aryl group for example, indicates that one or more hydrogen atoms on the aryl group is replaced with a substituent group, selected

from but not limited to, $-\text{SO}_2\text{Me}$ or phenyl group. A substituted heteroaryl group for example, indicates that one or more hydrogen atoms on the heteroaryl group is replaced with a substituent group, selected from, but not limited to, heterocycloalkyl, heteroaryl, N,N-dimethylamino. A substituted heterocycloalkyl group for example, indicates that one or more hydrogen atoms on the heterocycloalkyl group is replaced with a substituent group, selected from, but not limited to, heterocycloalkyl, heteroaryl, N,N-dimethylamino, hydroxyl, alkoxy, alkoxycarbonyl, alkyl, aryl, sulfonyl, dimethylaminosulfonyl, aroyl, cycloalkanoyl, alkanoyl and $-\text{OC}(\text{O})\text{NCH}(\text{CH}_3)_2$. In some instances two hydrogen atoms on the same carbon of, for example, a heterocyclyl or alkyl group are replaced with a group to form a spiro compound

selected from but not limited to, for example,  and .

[0064] The term “partial agonist” refers to a compound the presence of which results in a biological activity of a receptor that is of the same type as that resulting from the presence of a naturally occurring ligand for the receptor, but of a lower magnitude.

[0065] The phrase “pharmaceutically acceptable” refers to molecular entities and compositions that are generally regarded as safe and nontoxic. In particular, pharmaceutically acceptable carriers, diluents or other excipients used in the pharmaceutical compositions of this disclosure are physiologically tolerable, compatible with other ingredients, and do not typically produce an allergic or similar untoward reaction (for example, gastric upset, dizziness and the like) when administered to a patient. Preferably, as used herein, the term “pharmaceutically acceptable” means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopoeia or other generally recognized pharmacopoeia for use in animals, and more particularly in humans.

[0066] The phrase “pharmaceutically acceptable salt(s)”, as used herein, includes those salts of compounds of the disclosure that are safe and effective for use in mammals and that possess the desired biological activity. Pharmaceutically acceptable salts include salts of acidic or basic groups present in compounds of the disclosure or in compounds identified pursuant to the methods of the disclosure. Pharmaceutically acceptable acid addition salts include, but are not limited to, hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, tartrate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucaronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzensulfonate, p-toluenesulfonate and pamoate (i.e., 1,1'-methylene-bis-(2-hydroxy-3-

naphthoate)) salts. Certain compounds of the disclosure can form pharmaceutically acceptable salts with various amino acids. Suitable base salts include, but are not limited to, aluminum, calcium, lithium, magnesium, potassium, sodium, zinc, iron and diethanolamine salts. Pharmaceutically acceptable base addition salts are also formed with amines, such as organic amines. Examples of suitable amines are N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine.

[0067] As used herein, the term "pharmaceutically acceptable carrier" includes any of the standard pharmaceutical carriers, such as a phosphate buffered saline solution, water, emulsions such as an oil/water or water/oil emulsion, and various types of wetting agents. The term also encompasses any of the agents approved by a regulatory agency of the US Federal government or listed in the US Pharmacopeia for use in animals, including humans.

[0068] The term "selectivity" or "selective" refers to a difference in the binding affinity of a compound (K_i) for a sigma receptor, for example, a sigma-2 receptor, compared to a non-sigma receptor. The compound possess high selectivity for a sigma receptor in synaptic neurons. The K_i for a sigma-2 receptor or both a sigma-2 and a sigma-1 receptor is compared to the K_i for a non-sigma receptor. In some embodiments, the compound is a selective sigma-2 receptor antagonist, or sigma-1 receptor ligand, and has at least 10-fold, 20-fold, 30-fold, 50-fold, 70-fold, 100-fold, or 500-fold higher affinity, or more, for binding to a sigma receptor compared to a non-sigma receptor as assessed by a comparison of binding dissociation constant K_i values, or IC_{50} values, or binding constant, at different receptors. Any known assay protocol can be used to assess the K_i or IC_{50} values at different receptors, for example, by monitoring the competitive displacement from receptors of a radiolabeled compound with a known dissociation constant, for example, by the method of Cheng and Prusoff (1973) (Biochem. Pharmacol. 22, 3099-3108), or specifically as provided herein.

[0069] As used herein the term "plasma stability" refers to the degradation of compounds in plasma, for example, by enzymes such as hydrolases and esterases. Any of a variety of *in vitro* assays can be employed. Test compounds are incubated in plasma over various time periods. The percent parent compound (analyte) remaining at each time point reflects plasma stability. Poor stability characteristics can tend to have low bioavailability. Good plasma stability can be defined as greater than 50% analyte remaining after 30 min, greater than 50% analyte remaining after 45 minutes, and preferably greater than 50% analyte remaining after 60 minutes.

[0070] “Sigma-2 ligand” refers to a compound that binds to a sigma-2 receptor and includes agonists, antagonists, partial agonists, inverse agonists and simply competitors for other ligands of this receptor or protein.

[0071] The term “sigma-2 receptor antagonist compound” refers to a compound that binds to a sigma-2 receptor in a measurable amount and acts as a functional antagonist with respect to Abeta effects oligomer induced synaptic dysfunction resultant from sigma-2 receptor binding.

[0072] The terms "subject," “individual” or “patient” are used interchangeably and as used herein, are intended to include human and non-human animals. Non-human animals includes all vertebrates, e.g. mammals and non-mammals, such as non-human primates, sheep, dogs, cats, cows, horses, chickens, amphibians, and reptiles, although mammals are preferred, such as non-human primates, sheep, dogs, cats, cows and horses. Preferred subjects include human patients. The methods are particularly suitable for treating human patients having a disease or disorder described herein.

[0073] A “test compound” is a compound according to any embodiment described herein that is being tested in any test. Tests include any *in vivo* or *in vitro* test, computer model or simulation, virtual drug trial, stem cell and genetic testing methods, non-invasive imaging techniques and the like.

[0074] As used herein, the term “therapeutic” means an agent utilized to treat, combat, ameliorate, protect against or improve an unwanted condition or disease of a subject.

[0075] A “therapeutically effective amount” of a compound, pharmaceutically acceptable salt thereof or pharmaceutical composition according to any embodiment described herein, is an amount sufficient to produce a selected effect on at least one symptom or parameter of a specific disease or disorder. The therapeutic effect may be objective (i.e., measurable by some test or marker) or subjective (i.e., subject gives an indication of or feels an effect or physician observes a change). A therapeutically effective amount of a compound, according to any embodiment described herein, may broadly range from 0.01 mg/kg to about 500 mg/kg, about 0.01 to about 250 mg/kg, about 0.01 to about 25 mg/kg, about 0.05 mg/kg to about 20 mg/kg, about 0.1 mg/kg to about 400 mg/kg, about 0.1 mg/kg to about 200 mg/kg, about 0.1 mg/kg to about 25 mg/kg, about 0.1 to about 10 mg/kg, about 0.2 to about 5 mg/kg, about 1 mg/kg to about 300 mg/kg, about 10 mg/kg to about 100 mg/kg, body weight. The effect contemplated herein, includes both medical therapeutic and/or prophylactic treatment, as appropriate. The specific dose of a compound administered according to this disclosure to obtain therapeutic and/or prophylactic effects is determined by the particular

circumstances surrounding the case, including, for example, the compound administered, the route of administration, the co-administration of other active ingredients, the condition being treated, the activity of the specific compound employed, the specific composition employed, the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed and the duration of the treatment;. The therapeutically effective amount administered will be determined by the physician in the light of the foregoing relevant circumstances and the exercise of sound medical judgment. A therapeutically effective amount of a compound, according to any embodiment described herein, is typically an amount such that when it is administered in a physiologically tolerable excipient composition, it is sufficient to achieve an effective systemic concentration or local concentration in the tissue. The total daily dose of the compounds according to any embodiment described herein administered to a human or other animal in single or in divided doses can be in amounts, for example, from about 0.01 mg/kg to about 500 mg/kg, about 0.01 to about 250 mg/kg, about 0.01 to about 25 mg/kg, about 0.05 mg/kg to about 20 mg/kg, about 0.1 mg/kg to about 400 mg/kg, about 0.1 mg/kg to about 200 mg/kg, about 0.1 mg/kg to about 25 mg/kg, about 0.1 to about 10 mg/kg, about 0.2 to about 5 mg/kg, about 1 mg/kg to about 300 mg/kg, about 10 mg/kg to about 100 mg/kg, body weight per day. Single dose pharmaceutical compositions of any embodiment described herein, may contain such amounts or submultiples thereof to make up the daily dose. For example, the compounds according to any embodiment described herein, may be administered on a regimen of 1 to 4 times per day, such as once, twice, three times or four times per day. In some embodiments, the therapeutically effective amount of a compound according to any embodiment disclosed herein, can range between about 0.01 and about 25 mg/kg/day. In some embodiments the therapeutically effective amount is between a lower limit of about 0.01 mg/kg of body weight, about 0.1 mg/kg of body weight, about 0.2 mg/kg of body weight, about 0.3 mg/kg of body weight, about 0.4 mg/kg of body weight, about 0.5 mg/kg of body weight, about 0.60 mg/kg of body weight, about 0.70 mg/kg of body weight, about 0.80 mg/kg of body weight, about 0.90 mg/kg of body weight, about 1 mg/kg of body weight, about 2.5 mg/kg of body weight, about 5 mg/kg of body weight, about 7.5 mg/kg of body weight, about 10 mg/kg of body weight, about 12.5 mg/kg of body weight, about 15 mg/kg of body weight, about 17.5 mg/kg of body weight, about 20 mg/kg of body weight, about 22.5 mg/kg of body weight, and about 25 mg/kg of body weight; and an upper limit of 25 mg/kg of body weight, about 22.5 mg/kg of body weight, about 20 mg/kg of body weight, about 17.5 mg/kg of body weight, about 15 mg/kg of body weight, about 12.5 mg/kg of body

weight, about 10 mg/kg of body weight, about 7.5 mg/kg of body weight, about 5 mg/kg of body weight, about 2.5 mg/kg of body weight, about 1 mg/kg of body weight, about 0.9 mg/kg of body weight, about 0.8 mg/kg of body weight, about 0.7 mg/kg of body weight, about 0.6 mg/kg of body weight, about 0.5 mg/kg of body weight, about 0.4 mg/kg of body weight, about 0.3 mg/kg of body weight, about 0.2 mg/kg of body weight, about 0.1 mg/kg of body weight, and about 0.01 mg/kg of body weight. In some embodiments, the therapeutically effective amount is about 0.1 mg/kg/day to about 10 mg/kg/day; in some embodiments the therapeutically effective amount is about 0.2 and about 5 mg/kg/day. In some embodiments, treatment regimens according to the disclosure comprise administration to a patient in need of such treatment will usually include from about 1 mg to about 5000 mg, about 10 mg to about 2000 mg, about 10 mg to about 200 mg, about 20 to about 1000 mg, about 20 to about 500 mg, about 20 to about 400 mg, about 40 to about 800 mg, about 50 mg to about 500 mg, about 80 to about 1600 mg and about 50 mg, of a compound according to any embodiment disclosed herein, or a pharmaceutically acceptable salt thereof, per day in single or multiple doses. In some embodiments the therapeutically effective amount is a total daily dose of 50 mg to 500 mg. In some embodiments, the daily dose is between a lower limit of about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg; about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg; about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, 300 mg, about 305 mg, about 310 mg, about 315 mg; about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395, about 400 mg, about 405 mg, about 410 mg, about 415 mg; about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, and about 500 mg and an upper limit of about 500 mg, about 495 mg, about 490 mg, about 485 mg, about 480 mg, about 475 mg, about 470 mg, about 465 mg, about 460 mg, about 455 mg, about 450 mg, about 445 mg, about 440 mg, about 435 mg, about 430 mg, about 425 mg, about 420 mg, about 415 mg, about 410 mg,

about 405 mg, about 400 mg, about 395 mg, about 390 mg, about 385 mg, about 380 mg, about 375 mg, about 370 mg, about 365 mg, about 360 mg, about 355 mg, about 350 mg, about 345 mg, about 340 mg, about 335 mg, about 330 mg, about 325 mg, about 320 mg, about 315 mg, about 310 mg, about 305 mg, about 300 mg, about 295 mg, about 290 mg, about 285 mg, about 280 mg, about 275 mg, about 270 mg, about 265 mg, about 260 mg, about 255 mg, about 250 mg, about 245 mg, about 240 mg, about 235 mg, about 230 mg, about 225 mg, about 220 mg, about 215 mg, about 210 mg, about 205 mg, 200 mg, about 195 mg, about 190 mg, about 185 mg, about 180 mg, about 175 mg, about 170 mg, about 165 mg, about 160 mg, about 155 mg, about 150 mg, about 145 mg, about 140 mg, about 135 mg, about 130 mg, about 125 mg, about 120 mg, about 115 mg, about 110 mg, about 105 mg, about 100 mg, about 95 mg, about 90 mg, about 85 mg, about 80 mg, about 75 mg, about 70 mg, about 65 mg, about 60 mg, about 55 mg, and about 50 mg of a compound according to any embodiment herein. In some embodiments, the total daily dose is about 50 mg to 150 mg. In some embodiments, the total daily dose is about 50 mg to 250 mg. In some embodiments, the total daily dose is about 50 mg to 350 mg. In some embodiments, the total daily dose is about 50 mg to 450 mg. In some embodiments, the total daily dose is about 50 mg. It will be understood that the pharmaceutical formulations of the disclosure need not necessarily contain the entire amount of the compound that is effective in treating the disorder, as such effective amounts can be reached by administration of a plurality of divided doses of such pharmaceutical formulations. The compounds may be administered on a regimen of 1 to 4 times per day, such as once, twice, three times or four times per day.

[0076] The term “therapeutic phenotype” is used to describe a pattern of activity for compounds in the *in vitro* assays that is predictive of behavioral efficacy. A compound that (1) selectively binds with high affinity to a sigma-2 receptor, and (2) acts as a functional antagonist with respect to Abeta oligomer-induced effects in a neuron, is said to have the “therapeutic phenotype” if (i) it blocks or reduces A β -induced membrane trafficking deficits; (ii) it blocks or reduces A β -induced synapse loss and (iii) it does not affect trafficking or synapse number in the absence of Abeta oligomer. This pattern of activity in the *in vitro* assays is termed the “therapeutic phenotype” and is predictive of behavioral efficacy.

[0077] The term “therapeutic profile” is used to describe a compound that meets the therapeutic phenotype, and also has good brain penetrability (the ability to cross the blood brain barrier), good plasma stability and good metabolic stability.

[0078] The term “tissue” refers to any aggregation of similarly specialized cells which are united in the performance of a particular function.

[0079] The terms “treat,” “treated,” or “treating” as used herein, refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to protect against (partially or wholly) or slow down (e.g., lessen or postpone the onset of) an undesired physiological condition, disorder or disease, or to obtain beneficial or desired clinical results such as partial or total restoration or inhibition in decline of a parameter, value, function or result that had or would become abnormal. For the purposes of this disclosure, beneficial or desired clinical results include, but are not limited to, alleviation of symptoms; diminishment of the extent or vigor or rate of development of the condition, disorder or disease; stabilization (i.e., not worsening) of the state of the condition, disorder or disease; delay in onset or slowing of the progression of the condition, disorder or disease; amelioration of the condition, disorder or disease state; and remission (whether partial or total), whether or not it translates to immediate lessening of actual clinical symptoms, or enhancement or improvement of the condition, disorder or disease. Treatment seeks to elicit a clinically significant response without excessive levels of side effects. Treatment also includes prolonging survival as compared to expected survival if not receiving treatment.

Human Amyloid Beta and Sigma -2 Antagonists

[0080] Overproduction and accumulation of amyloid beta is a pathologic feature of Alzheimer’s disease. Human amyloid beta (Abeta) is the main component of insoluble amyloid plaques-deposits found in the brain of patients with Alzheimer’s disease. The plaques are composed of fibrillar aggregates of Abeta. Amyloid beta fibrils have been associated with the advanced stages of Alzheimer's disease.

[0081] The cognitive hallmark of early Alzheimer’s disease is an extraordinary inability to form new memories. Early memory loss is considered a synapse failure caused by soluble A β oligomers. These oligomers block long-term potentiation, a classic experimental paradigm for synaptic plasticity, and they are strikingly elevated in AD brain tissue and transgenic AD models. It has been hypothesized that early memory loss stems from synapse failure before neuron death and that synapse failure derives from actions of soluble A β oligomers rather than fibrils. Lacor et al., *Synaptic targeting by Alzheimer’s-related amyloid β oligomers*, J. Neurosci. 2004, 24(45):10191-10200.

[0082] Abeta is a cleavage product of an integral membrane protein, amyloid precursor protein (APP), found concentrated in the synapses of neurons. Soluble forms of

Abeta are present in the brains and tissues of Alzheimer's patients, and their presence correlates with disease progression. Yu et al., 2009, *Structural characterization of a soluble amyloid beta-peptide oligomer*, Biochemistry, 48(9):1870-1877. Soluble amyloid β oligomers have been demonstrated to induce changes in neuronal synapses that block learning and memory.

[0083] Smaller, soluble A β oligomers interfere with a number of signaling pathways critical for normal synaptic plasticity, ultimately resulting in spine and synapse loss. Selkoe et al., 2008, *Soluble oligomers of the amyloid beta-protein impair synaptic plasticity and behavior*, Behav Brain Res 192(1): 106-113. Alzheimer's begins and persists as a synaptic plasticity disease.

[0084] The presence of soluble A β oligomers is believed to be to be responsible for early cognitive decline in the pre-Alzheimer's diseased brain. It is known that amyloid beta oligomers bind at neuronal synapses and that sigma-2 receptors are present in significant amounts in neurons and glia.

[0085] Sigma receptors are multifunctional adapter/chaperone proteins that participate in several distinct protein signaling complexes in a tissue and state-related manner. The sigma-2 receptor is expressed in brain and various peripheral tissues at low levels. (Walker et al., 1990 *Sigma receptors: biology and function*. Pharmacol. Rev. 42:355-402). Sigma-2 receptors are present in human hippocampus and cortex. The sigma-2 receptor was also previously validated as a biomarker for tumor cell proliferation. (Mach et al., *Sigma-2 receptors as potential biomarkers of proliferation in breast cancer*. Cancer Res. 57:156-161, 1997).

[0086] Sigma-2 receptors are implicated in many signaling pathways such as heme binding, Cytochrome P450 metabolism, cholesterol synthesis, progesterone signaling, apoptosis and membrane trafficking. Only a subset of sigma receptor binding sites/signaling pathways are relevant to oligomer signaling in AD. No sigma-2 receptor knock-outs are currently available and human mutations in sigma-2 sequence have not been studied in a neurodegeneration context.

[0087] A sigma-2 receptor was recently identified as the progesterone receptor membrane component 1 (PGRMC1) in rat liver by use of a photoaffinity probe WC-21, which irreversibly labels sigma-2 receptors in rat liver. Xu et al. *Identification of the PGRMC1 protein complex as the putative sigma-2 receptor binding site*. Nature Communications 2, article number 380, July 5, 2011, incorporated herein by reference.

PGRMC1 (progesterone receptor membrane component 1) was identified as the critical 25kDa component of sigma-2 receptor activity in August 2011 by Xu et al. PGRMC1 is a single transmembrane protein with no homology to sigma-1 protein; family members include PGRMC2 and neudesin. PGRMC1 contains a cytochrome b5 heme-binding domain. PGRMC1 is a single transmembrane protein with no homology to S1 protein; family members include PGRMC2 and neudesin. PGRMC1 contains a cytochrome b5 heme-binding domain. Endogenous PGRMC1 ligands include progesterone/steroids, cholesterol metabolites, glucocorticoids, and heme. PGRMC1 functions as chaperone/adaptor associated with different protein complexes in different subcellular locations (Cahill 2007. *Progesterone receptor membrane component 1: an integrative review*. J. Steroid Biochem. Mol. Biol. 105:16-36). PGRMC1 binds heme with reducing activity, complexes with CYP450 proteins (regulated redox reactions), associates with PAIRBP1 and mediates progesterone block of apoptosis, and associates with Insig-1 and SCAP to induce SRE-related gene transcription in response to low cholesterol. The *C. elegans* homolog VEM1 associates with UNC-40/DCC to mediate axon guidance. PGRMC1 contains two SH2 target sequences, an SH3 target sequence, a tyrosine kinase site, two acidophilic kinase sites (CK2), and consensus binding sites for ERK1 and PDK1. PGRMC1 contains several ITAM sequences involved in membrane trafficking (vesicle transport, clathrin-dependent endocytosis of calveolin-containing pits).

[0088] While not being bound by theory, it is proposed that the sigma-2 receptor is a receptor for Abeta oligomer in neurons. Various receptors have been proposed in the literature for soluble Abeta oligomers including prion protein, insulin receptor, beta adrenergic receptor and RAGE (receptor for advanced glycation end products). Laurén, J. et al, 2009, Nature, 457(7233): 1128-1132; Townsend, M. et al, J. Biol. Chem. 2007, 282:33305-33312; Sturchler, E. et al, 2008, J. Neurosci. 28(20):5149-5158. Indeed many investigators believe that Abeta oligomer may bind to more than one receptor protein. Without being bound by theory, the present inventors postulate an additional receptor for Abeta oligomer located (not necessarily exclusively) in neurons.

[0089] Without being bound by theory, Abeta oligomers are sigma receptor agonists that bind to sigma protein complexes and cause aberrant trafficking and synapse loss. It is demonstrated herein, that compounds described herein that antagonize this interaction and/or sigma receptor function in neurons will compete or otherwise interfere with Abeta oligomers and return neuronal responses to normal. Such compounds are considered functional sigma-2 receptor antagonists.

[0090] In some embodiments, a compound of any embodiment described herein, may act as a functional antagonist in a neuronal cell with respect to inhibiting soluble A β oligomer induced synapse loss, and inhibiting soluble A β oligomer induced deficits in a membrane trafficking assay; exhibiting high affinity at a sigma-2 receptor; as well as having high selectivity for one or more sigma receptors compared to any other non-sigma receptor; and exhibiting good drug-like properties.

[0091] In some embodiments, a compound according to any embodiment described herein, that acts as functional antagonist meeting certain *in vitro* assay criteria detailed herein, will exhibit behavioral efficacy, or be predicted to have behavioral efficacy, in one or more relevant animal behavioral models. In some embodiments, behavioral efficacy is determined at 10 mg/kg p.o., or less.

[0092] *In vitro* assay platforms predictive of behavioral efficacy useful in the invention described herein, are known in the art, in particular, in US 9,796,672, herein incorporated by reference in its entirety. In accordance with the *in vitro* assay platform, a compounds of any embodiment described herein, may bind with high affinity to a sigma-2 receptor; acts as a functional antagonist with respect to Abeta oligomer-induced effects in a neuron; inhibits Abeta oligomer-induced synapse loss in a central neuron or reduces Abeta oligomer binding to neurons to inhibit synapse loss; and does not affect trafficking or synapse number in the absence of Abeta oligomer. This pattern of activity in the *in vitro* assays is termed the “therapeutic phenotype”. The ability of a compound according to any embodiment described herein, to block Abeta oligomer effects in mature neurons without affecting normal function in the absence of Abeta oligomers meets the criteria for the therapeutic phenotype. A compounds of any embodiment described herein, having a therapeutic phenotype, can block Abeta oligomer-induced synaptic dysfunction.

[0093] In some embodiments, a compound according to any embodiment described herein, exhibits sigma-2 antagonist activity, high affinity for the sigma-2 receptor, and the ability to block soluble Abeta oligomer binding or Abeta oligomer-induced synaptic dysfunction.

[0094] In some embodiments, a compound according to any embodiment described herein, is designed to enhance the ability to cross the blood-brain barrier.


[0095] In some embodiments, a compound according to any embodiment described herein, blocks binding between soluble Abeta oligomers and a sigma-2 receptor.

[0096] In some embodiments, a compound according to any embodiment described herein, exhibits high affinity for the sigma-2 receptor.

[0097] Embodiments of the invention are directed to compounds according to any embodiment described herein, useful for treating neurodegenerative disease and cognitive decline, pharmaceutical compositions containing such compounds and pharmaceutically acceptable carries, excipients, or diluents, and methods for treating neurodegenerative disease and cognitive decline by administering such compounds and pharmaceutical compositions in a pharmaceutically acceptable amount.

Compounds of the Invention

[0098] Various embodiments are directed to a compound of Formula I:



or a pharmaceutically acceptable salt thereof.

[0099] Each of substituents R^a , R^b , R^c , R^d and R^e of Formula I is independently selected from the group consisting of, H, hydroxyl, halo, alkyl, alkoxy, CF_3 , SO_2CH_3 , and morpholino.

[0100] Substituent R^1 of Formula I is selected from the group consisting of hydrogen, alkyl, phenyl, or $-CH=C(CH_3)_2$.

[0101] Substituent R^2 of Formula I is an optionally substituted cyclic amino group.

[0102] In some embodiments, each of substituents R^a , R^b , R^c , R^d and R^e of Formula I is independently selected from the group consisting of, H, hydroxyl, Cl, F, methyl, $-OCH_3$, $-OC(CH_3)_3$, $O-CH(CH_3)_2$, CF_3 , SO_2CH_3 , and morpholino.


[0103] In some embodiments, each of substituents R^a , R^b , R^c , R^d and R^e of Formula I is independently selected from the group consisting of, H, Cl, F, and CF_3 .

[0104] In some embodiments, each of substituents R^a , R^b , R^d and R^e of Formula I is independently H and R^c , is selected from the group consisting of H, hydroxyl, halo, alkyl, alkoxy, CF_3 , SO_2CH_3 , and morpholino.

[0105] In some embodiments, each of substituents R^a , R^b , R^d and R^e of Formula I is independently H and R^c , is selected from the group consisting of H, hydroxyl, Cl, F, methyl, $-OCH_3$, $-OC(CH_3)_3$, $O-CH(CH_3)_2$, CF_3 , SO_2CH_3 , and morpholino.

[0106] In some embodiments, each of substituents R^a , R^b , R^d and R^e of Formula I is independently H and R^c , is selected from the group consisting of H, Cl, F, and CF_3 .


[0107] In various embodiments, R^2 is any heterocycloalkyl or heteroaryl containing a nitrogen in the ring that is bound to the aliphatic chain of Formula I through the nitrogen atom. In some embodiments, for example, R^2 is an optionally substituted cyclic amino group selected from:



and the like, wherein each nitrogen containing heterocycloalkyl or heteroaryl can be optionally substituted with one or more substituents selected from, hydroxyl, halo, CF_3 , alkoxy, aryloxy, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl, optionally substituted C_3 - C_{10} heteroaryl, substituted or unsubstituted C_3 - C_{10} cycloalkyl or heterocycloalkyl.


[0108] In various embodiments, R^2 is selected from the group consisting of optionally substituted aziridinyl, optionally substituted pyrrolidinyl, optionally substituted imidizolidinyl, optionally substituted piperidinyl, optionally substituted piperazinyl, optionally substituted oxopiperazinyl, and optionally substituted morpholinyl.

[0109] In some embodiments, when R^2 is a substituted cyclic amino, one or more of the hydrogen atoms in the cyclic amino group is replaced with a group selected from alkanoyl, alkoxy, alkoxyalkyl, (alkoxy)alkoxyalkyl, alkoxycarbonyl, alkyl, aryloxy, aryloyl, cycloalkanoyl, $-OC(O)NCH(CH_3)_2$, (N,N-dimethylamino)pyridinyl, (N,N-dimethylamino)sulfonyl, halo, heterocyclyl, (heterocyclyl)alkoxyalkyl, hydroxyl, hydroxyalkyl, methylpiperidinyl, methylsulfonyl, methylsulfonylphenyl, morpholinylpyridinyl, perfluoroalkyl, phenyl, piperidinyl, pyrrolidinylpyridinyl, tetrahydropyranyl, and CF_3 . In some embodiments two hydrogen atoms on the same carbon of the cyclic amino group are replaced with a compound selected from




[0110] In some embodiments, R^2 is a pyrrolidinyl or a substituted pyrrolidinyl substituted with one or more substituents selected from the group consisting of alkoxyalkyl, alkoxy carbonyl, alkyl, hydroxyl, and hydroxyalkyl. In some embodiments R^2 is a substituted pyrrolidinyl substituted with a single substituent selected from the group consisting of alkoxyalkyl, alkoxy carbonyl, alkyl, hydroxyl, and hydroxyalkyl. In some embodiments R^2 is a substituted pyrrolidinyl substituted with a single substituent selected from the group consisting of hydroxyl, hydroxymethyl, methoxymethyl, methoxy carbonyl and methyl.

[0111] In some embodiments, R^2 is a piperidinyl or a substituted piperidinyl substituted with one or more substituents selected from the group consisting of alkoxy, alkoxyalkyl, (alkoxy)alkoxyalkyl, alkoxy carbonyl, alkyl, aryloxy, $-\text{OC}(\text{O})\text{NCH}(\text{CH}_3)_2$, (N,N-dimethylamino)pyridinyl, halo, heterocyclyl, (heterocyclyl)alkoxyalkyl, hydroxy, hydroxyalkyl, methylpiperidinyl, methylsulfonylphenyl, morpholinylpyridinyl, perfluoroalkyl, phenyl, piperidinyl, pyrrolidinylpyridinyl, tetrahydropyranyl, and CF_3 . In some embodiments, R^2 is a piperidinyl or a substituted piperidinyl substituted with a single substituent selected from the group consisting of alkoxy, alkoxyalkyl, (alkoxy)alkoxyalkyl, alkoxy carbonyl, alkyl, aryloxy, $-\text{OC}(\text{O})\text{NCH}(\text{CH}_3)_2$, (N,N-dimethylamino)pyridinyl, halo, heterocyclyl, (heterocyclyl)alkoxyalkyl, hydroxyl, hydroxyalkyl, methylpiperidinyl, methylsulfonylphenyl, morpholinylpyridinyl, perfluoroalkyl, phenyl, piperidinyl, pyrrolidinylpyridinyl, tetrahydropyranyl, and CF_3 . In some embodiments, R^2 is a piperidinyl or a substituted piperidinyl substituted with a single substituent selected from the group consisting of methyl, isopropyl, isobutyl, CF_3 , hydroxymethyl, hydroxyethyl, (isopropoxy)ethyl, $-(\text{CH}_2)_2\text{O}(\text{CH}_2)_2\text{OCH}_3$, $-(\text{CH}_2)_3\text{OCH}_3$, $-\text{C}(\text{O})\text{OMe}$, $-\text{C}(\text{O})\text{OEt}$, hydroxyl, methoxy, isopropoxy, phenyloxy, F, ethoxy, phenyl,







In some embodiments, R² is a piperidinyl or a substituted piperidinyl substituted at the 4 position of the piperidinyl with a single substituent selected from the group consisting of alkoxy, alkoxyalkyl, (alkoxy)alkoxyalkyl, alkoxy carbonyl, alkyl, aryloxy, -OC(O)NCH(CH₃)₂, (N,N-dimethylamino)pyridinyl, halo, heterocyclyl, (heterocyclyl)alkoxyalkyl, hydroxyl, hydroxyalkyl, methylpiperidinyl, methylsulfonylphenyl, morpholinylpyridinyl, perfluoroalkyl, phenyl, piperidinyl, pyrrolidinylpyridinyl, tetrahydropyranyl, and CF₃. In some embodiments, R² is a piperidinyl or a substituted piperidinyl substituted at the 4 position of the piperidinyl with a single substituent selected from the group consisting of methyl, isopropyl, isobutyl, CF₃, hydroxymethyl, hydroxyethyl, (isopropoxy)ethyl, -(CH₂)₂O(CH₂)₂OCH₃, -(CH₂)₃OCH₃, -C(O)OMe, -C(O)OEt, hydroxyl,

methoxy, isopropoxy, phenyloxy, F, ethoxy, phenyl,




[0112] In some embodiments, R^2 is a piperidinyl or a substituted piperidinyl substituted with two substituent groups on the same carbon of the piperidinyl independently selected from the group consisting of alkoxyalkyl, alkyl, $-\text{OC}(\text{O})\text{NCH}(\text{CH}_3)_2$, hydroxyl, and phenyl. In some embodiments, R^2 is a piperidinyl or a substituted piperidinyl substituted with two substituent groups at the 4 position of the piperidinyl independently selected from the group consisting of alkoxyalkyl, alkyl, $-\text{OC}(\text{O})\text{NCH}(\text{CH}_3)_2$, hydroxyl, and phenyl. In some embodiments R^2 is a piperidinyl or a substituted piperidinyl substituted with two substituent groups at the 4 position selected from the group consisting of hydroxyl and methyl; hydroxyl and ethyl; hydroxyl and $-(\text{CH}_2)_2\text{OCH}_3$; hydroxyl and phenyl; methyl and phenyl; methyl and $-\text{OC}(\text{O})\text{NCH}(\text{CH}_3)_2$; and butyl and $-\text{OC}(\text{O})\text{NCH}(\text{CH}_3)_2$. In some embodiments two hydrogen atoms on the same carbon of the piperidinyl are replaced with a compound selected from




compound selected from ,  and  to form a spiro compound.


[0113] In some embodiments, R^2 is a piperazinyl or a substituted piperazinyl substituted with one or more substituents selected from the group consisting of alkanoyl, alkoxycarbonyl, aryloyl, cycloalkanoyl, (N,N-dimethylamino)sulfonyl, heterocyclyl, methylsulfonyl, and phenyl. In some embodiments, R^2 is a substituted piperazinyl substituted with a single substituent selected from the group consisting of alkanoyl, alkoxycarbonyl, aryloyl, cycloalkanoyl, (N,N-dimethylamino)sulfonyl, heterocyclyl, methylsulfonyl, and phenyl. In some embodiments, R^2 is a substituted piperazinyl substituted with a single substituent selected from the group consisting of $-C(O)OC(CH_3)_3$, $-C(O)OCH_2CH(CH_3)_2$, $-C(O)OCH_2CH_3$, $-C(O)OCH_3$, phenyl, $-C(O)CH_3$, $-C(O)Ph$, $-SO_2Me$, $-SO_2N(CH_3)_2$,



. In some embodiments, R^2 is a substituted piperazinyl substituted with a single substituent at the 4 position selected from the group consisting of $-C(O)OC(CH_3)_3$, $-C(O)OCH_2CH(CH_3)_2$, $-C(O)OCH_2CH_3$, $-C(O)OCH_3$, phenyl, $-C(O)CH_3$, $-C(O)Ph$, $-SO_2Me$, $-SO_2N(CH_3)_2$,




[0114] In certain embodiments, R^2 is a substituted piperdinyll of formula:




wherein, R^3 is hydrogen or C_1 - C_8 alkyl, and R^4 is hydrogen, hydroxyl, halogen, CF_3 , alkoxy, aryloxy, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl, optionally substituted C_3 - C_{10} heteroaryl, optionally substituted C_3 - C_{10} cycloalkyl or optionally substituted C_3 - C_{10} heterocycloalkyl.

[0115] In some embodiments, R^2 is




wherein each of R^5 and R^6 is independently, hydrogen, hydroxyl, sulfonyl, dialkylamino, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl optionally substituted C_3 - C_{10} heteroaryl, optionally substituted C_3 - C_{10} cycloalkyl or optionally substituted C_3 - C_{10} heterocycloalkyl. In some embodiments R^5 is hydrogen, dialkylamino, or C_3 - C_{10} heterocycloalkyl. In some embodiments R^5 is hydrogen, dialkylamino, pyrrolidinyl or morpholinyl. In some embodiments, R^6 is sulfonyl. In some embodiments, R^6 is methylsulfonyl.

[0116] In some embodiments, R^2 is:




34




wherein R^7 is hydrogen, hydroxyl, sulfonyl, dialkylaminosulfonyl, alkoxy carbonyl, acyl, benzoyl, cycloalkyl carbonyl, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl optionally substituted C_3 - C_{10} heteroaryl, optionally substituted C_3 - C_{10} cycloalkyl or optionally substituted C_3 - C_{10} heterocycloalkyl. In some embodiments R^7 is sulfonyl, dialkylaminosulfonyl, alkoxy carbonyl, acyl, benzoyl, cycloalkyl carbonyl, C_5 - C_{10} aryl or optionally substituted C_3 - C_{10} heterocycloalkyl.

[0120] In some embodiments R^2 is




[0121] In various embodiments, R^2 is optionally substituted pyrrolidinyl:




where R^8 is hydrogen, hydroxyl, sulfonyl, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl, optionally substituted C_3 - C_{10} heteroaryl, optionally substituted C_3 - C_{10} cycloalkyl or optionally substituted C_3 - C_{10} heterocycloalkyl. In some embodiments, R^8 is hydrogen, hydroxyl or optionally substituted C_1 - C_{10} alkyl.

[0122] In some embodiments, R^2 is:




[0123] In some embodiments, R^2 is an optionally substituted bicyclic ring or an optionally substituted fused ring. For example, in some embodiments, R^2 is selected from the group consisting of:



where R^9 is hydrogen, hydroxyl, sulfonyl, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl, optionally substituted C_3 - C_{10} heteroaryl, optionally substituted C_3 - C_{10} cycloalkyl or optionally substituted C_3 - C_{10} heterocycloalkyl.

[0124] In some embodiments, R^2 is




wherein each of R^{11a} , R^{11b} , R^{11c} , and R^{11d} , is, independently selected from, hydrogen, hydroxy, sulfonyl, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl, optionally substituted C_3 - C_{10} heteroaryl, optionally substituted C_3 - C_{10} cycloalkyl or optionally substituted C_3 - C_{10} heterocycloalkyl. In particular embodiments, R^2 is




[0125] Some embodiments disclosed herein describe a compound wherein each R^a , R^b , R^c , R^d and R^e is selected from any embodiment disclosed herein for each of R^a , R^b , R^c , R^d and R^e ; R^1 is selected from any embodiment disclosed herein for R^1 ; and R^2 is selected from any embodiment disclosed herein for R^2 .


[0126] Some embodiments are directed to a compound selected from




Example 1




Example 2




Example 3




Example 4




Example 5




Example 6




Example 7




Example 8




Example 9




Example 10




Example 11




Example 12




Example 13




Example 14




Example 15




Example 16




Example 17




Example 19




Example 20




Example 22




Example 23




Example 24




Example 25




Example 27




Example 28




Example 29




Example 30




Example 31




Example 32




Example 33




Example 34




Example 35




Example 36




Example 37




Example 38




Example 39




Example 40




Example 41




Example 42




Example 43




Example 44




Example 45




Example 46




Example 47




Example 49




Example 50




Example 51




Example 53




Example 55




Example 56




Example 58




Example 59




Example 60




Example 61




Example 62




Example 63




Example 67




Example 68




Example 70




Example 71




Example 72




Example 73




Example 75




Example 76




Example 77




Example 79




Example 81




Example 82




Example 83




Example 84




Example 85




Example 86




Example 87




Example 89




Example 90




Example 97




Example 98




Example 107




Example 116




Example 117




Example 118




Example 119




Example 120




Example 122




Example 123




Example 124




Example 125




Example 126




Example 127




Example 128




Example 129




Example 130




Example 131




Example 132




Example 133




Example 134




Example 135




Example 136




Example 137




Example 138




Example 139




Example 140




Example 141




Example 142




Example 143




Example 144




Example 145




Example 146




Example 147




Example 148




Example 149




Example 150




Example 151




Example 152




Example 153




Example 154




Example 155




Example 156




Example 157




Example 158




Example 159




Example 160




Example 161




Example 162




Example 163




Example 164




Example 167




Example 168




Example 169




Example 170




Example 171




Example 178




Example 179




Example 180




Example 181




Example 182




Example 183




Example 185




Example 187




Example 189




Example 191




Example 192




Example 193




Example 194




Example 195




Example 196




Example 197




Example 198




Example 199




Example 200




Example 201




Example 202




Example 203




Example 204




Example 205




Example 206




Example 207




Example 208




Example 209




Example 210




Example 211




Example 212




Example 213




Example 214




Example 215




Example 216




Example 217




Example 218




Example 219




Example 220




Example 221




Example 222




Example 223




Example 224




Example 225




Example 226




Example 227




Example 228




Example 229




Example 230




Example 231




Example 232




Example 233




Example 234




Example 235




Example 236




Example 237




Example 238




Example 239




Example 240




Example 241




Example 242




Example 243




Example 244




Example 245




Example 246




Example 247




Example 248




Example 249




Example 250




Example 251




Example 252




Example 253




Example 254




Example 255




Example 261




Example 262




Example 263




Example 264




Example 265




Example 266




Example 267




Example 268




Example 275




Example 276




Example 277




Example 278




Example 283




Example 284




Example 285




Example 286




Example 287




Example 288




Example 289




Example 290




Example 291




Example 292




Example 293




Example 294




Example 295




Example 296




Example 297




Example 298




Example 299




Example 300




Example 301




Example 302




Example 303




Example 304




Example 306




Example 307




Example 308




Example 310




Example 311




Example 312




Example 313




Example 314




Example 315




Example 316




Example 317




Example 318




Example 319




Example 320




Example 321




Example 322



Example 323




Example 324




Example 325

, and




Example 326


[0127] Some embodiments are directed to a compound selected from the group consisting of



Example 14




Example 17




Example 9

, and



Example 262

[0128] Further embodiments are directed to compounds of Formula II or pharmaceutically acceptable salt thereof:



II.


[0129] Each of substituents R^f , R^g , R^h , R^i and R^j of Formula II is independently selected from the group consisting of, H, hydroxyl, halo, alkyl, alkoxy, CF_3 , SO_2CH_3 , and morpholino.

[0130] Substituent R^{10} of Formula II is an optionally substituted cyclic amino group and m is an integer from 0 to 3.


[0131] In some embodiments each of substituents R^f , R^g , R^h , R^i and R^j of Formula II is independently selected from the group consisting of, H, hydroxyl, and alkoxy. In some embodiments each of substituents R^f , R^g , R^h , R^i and R^j of Formula II is independently selected from the group consisting of, H, hydroxyl, and methoxy. In some embodiments each of substituents R^f , R^g , and R^j is H and each of R^g , and R^h is independently selected from the hydroxyl, or methoxy.


[0132] In some embodiments, R^{10} is an optionally substituted aziridinyl, optionally substituted pyrrolidinyl, optionally substituted imidizolidinyl, optionally substituted piperidinyl, optionally substituted piperazinyl, optionally substituted oxopiperazinyl, or optionally substituted morpholinyl, and any of the individual substituted or unsubstituted piperdinyl, substituted or unsubstituted morpholinyl, substituted or unsubstituted piperazinyl, substituted or unsubstituted pyrrolidinyl, substituted or unsubstituted bicyclic, or substituted or unsubstituted fused rings described above in relation to Formula I.

[0133] In some embodiments, R^{10} is an optionally substituted fused ring, such as:




wherein each of R^{11e} , R^{11f} , R^{11g} , and R^{11h} is independently selected from, hydrogen, hydroxy, sulfonyl, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl optionally substituted C_3 - C_{10} heteroaryl, optionally substituted C_3 - C_{10} cycloalkyl or optionally substituted C_3 - C_{10} heterocycloalkyl. In certain embodiments R^{10} is not







[0135] Some embodiments describe a compound of Formula IIa:



IIa.


[0136] Each of substituents R^k and R^l is independently selected from the group consisting of H, hydroxyl, halo, alkyl, alkoxy, CF_3 , SO_2CH_3 , and morpholino.

[0137] Substituent R^{12} is selected from the group consisting of aryloxy, alkenyloxy, alkoxy, aminoalkyl, N,N-dimethylaminoalkyl, pyrrolidinyl, n-methylpyrrolidinyl, N-acylpyrrolidinyl, carboxyaminoalkyl, hydroxyalkyl, -




[0138] In some embodiments each of substituents R^k and R^l is independently selected from the group consisting of H, hydroxyl and methoxy. In some embodiments R^l is methoxy and R^k is hydroxyl.


[0139] In some embodiments substituent R^{12} is selected from the group consisting of phenyloxy, $-OCH_2CH=CH_2$, methoxy, $-CH_2NH_2$, $-CH(NH_2)CH_3$, $-CH_2N(Me)_2$, $-CH(CH_3)N(Me)_2$, $-CH_2NHC(O)CH_3$, $-CH(OH)CH_3$, $-O(CH_2)_2OC(O)CH_3$,




[0140] Some embodiments describe a compound selected from the group consisting of




Example 26




Example 256




Example 257




Example 258




Example 259




Example 260




Example 269




Example 270




Example 271




Example 272




Example 273




Example 274




Example 279





Example 280



Example 281




Example 282


and
Example 305

Example 309


[0141] Some embodiments describe a compound selected from the group consisting of




Example 18




Example 21




Example 48




Example 52




Example 54




Example 57




Example 64




Example 65




Example 66




Example 69




Example 74




Example 78




Example 80




Example 88




Example 91




Example 92




Example 93




Example 94




Example 95




Example 96




Example 99




Example 100




Example 101




Example 102




Example 103




Example 104




Example 105




Example 106




Example 108




Example 109




Example 110




Example 111




Example 112




Example 113




Example 114




Example 115




Example 121




Example 165




Example 166




Example 172




Example 173




Example 174




Example 175




Example 176




Example 177



Example 184




Example 186



Example 188

, and



Example 190

[0142] Additional embodiments include salts, solvates, stereoisomers, prodrugs, and active metabolites of the compounds according to any embodiment described herein.

[0143] Some embodiments are directed to free base forms of the compounds according to any embodiment described herein. Other embodiments include salts of such compounds including, for example, pharmaceutically acceptable acid addition salts or pharmaceutically acceptable addition salts of free bases. Examples of pharmaceutically acceptable acid addition salts include, but are not limited to, salts derived from nitric, phosphoric, sulfuric, or hydrobromic, hydroiodic, hydrofluoric, phosphorous, as well as salts derived from nontoxic organic acids such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanoic acids, hydroxyl alkanoic acids, alkanedioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, and acetic, maleic, succinic, or citric acids. Non-limiting examples of such salts include napadisylate, besylate, sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, nitrate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, trifluoroacetate, propionate, caprylate, isobutyrate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, mandelate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, phthalate, benzenesulfonate, toluenesulfonate, phenylacetate, citrate, lactate, maleate, tartrate, methanesulfonate, and the like. Additional salt forms of the compounds described above include salts of amino acids such as arginate and the like and gluconate, galacturonate (*see e.g.*, Berge, et al. "Pharmaceutical Salts," J. Pharma. Sci. 1977;66:1).

[0144] Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines include N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine. The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient

amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid.

[0145] Various embodiments include total and partial salts, *i.e.* salts with 1, 2 or 3, preferably 2, equivalents of base per mole of acid of a compound or salt described above, with 1, 2 or 3 equivalents, preferably 1 equivalent, of acid per mole of base of a compound of according to any embodiment described herein. Typically, a pharmaceutically acceptable salt of a compound according to any embodiment described herein, may be readily prepared by using a desired acid or base as appropriate. The salt may precipitate from solution and be collected by filtration or may be recovered by evaporation of the solvent. For example, an aqueous solution of an acid such as hydrochloric acid may be added to an aqueous suspension of a compound according to any embodiment described herein, and the resulting mixture evaporated to dryness (lyophilized) to obtain the acid addition salt as a solid. Alternatively, a compound according to any embodiment described herein, may be dissolved in a suitable solvent, for example an alcohol such as isopropanol, and the acid may be added in the same solvent or another suitable solvent. The resulting acid addition salt may then be precipitated directly, or by addition of a less polar solvent such as diisopropyl ether or hexane, and isolated by filtration.

[0146] Many organic compounds can form complexes with solvents in which they are reacted or from which they are precipitated or crystallized. These complexes are known as “solvates.” For example, a complex with water is known as a “hydrate.” Various embodiments include solvates of a compound according to any embodiment described herein. In some embodiments, salts of these compounds can form solvates.

[0147] Further embodiments include N-oxides of the compounds according to any embodiment described herein. N-oxides include heterocycles containing an otherwise unsubstituted sp^2 N atom. Examples of such N-oxides include pyridyl N-oxides, pyrimidyl N-oxides, pyrazinyl N-oxides and pyrazolyl N-oxides.

[0148] Compounds according to any embodiment described herein, may have one or more chiral centers and, depending on the nature of individual substituents, they can also have geometrical isomers. Thus, embodiments include stereoisomers, diastereomers, and enantiomers of the compounds according to any embodiment described herein. A chiral compound can exist as either an individual enantiomer or as a mixture of enantiomers. A mixture containing equal proportions of the enantiomers is called a “racemic mixture.” A mixture containing unequal portions of the enantiomers is described as having an “enantiomeric excess” (ee) of either the R or S compound. The excess of one enantiomer in a

mixture is often described with a % enantiomeric excess. The ratio of enantiomers can also be defined by “optical purity” wherein the degree at which the mixture of enantiomers rotates plane polarized light is compared to the individual optically pure R and S compounds. The compounds can also be a substantially pure (+) or (-) enantiomer of the compounds described herein. In some embodiments, a composition can include a substantially pure enantiomer that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% of one enantiomer. In certain embodiments, a composition may include a substantially pure enantiomer that is at least 99.5% one enantiomer.

[0149] The description above encompasses all individual isomers of the compounds according to any embodiment described herein, and the description or naming of a particular compound in the specification and claims is intended to include both individual enantiomers and mixtures thereof. Methods for the determination of stereochemistry and the resolution or stereotactic synthesis of stereoisomers are well-known in the art. Diastereomers differ in both physical properties and chemical reactivity. A mixture of diastereomers can be separated into enantiomeric pairs based on solubility, fractional crystallization or chromatographic properties, e.g., thin layer chromatography, column chromatography or HPLC. Purification of complex mixtures of diastereomers into enantiomers typically requires two steps. In a first step, the mixture of diastereomers is resolved into enantiomeric pairs, as described above. In a second step, enantiomeric pairs are further purified into compositions enriched for one or the other enantiomer or, more preferably resolved into compositions comprising pure enantiomers. Resolution of enantiomers typically requires reaction or molecular interaction with a chiral agent, e.g. solvent or column matrix. Resolution may be achieved, for example, by converting the mixture of enantiomers, e.g., a racemic mixture, into a mixture of diastereomers by reaction with a pure enantiomer of a second agent, i.e., a resolving agent. The two resulting diastereomeric products can then be separated. The separated diastereomers are then reconverted to the pure enantiomers by reversing the initial chemical transformation.

[0150] Resolution of enantiomers can also be accomplished by differences in their non-covalent binding to a chiral substance, e.g., by chromatography on homochiral adsorbants. The noncovalent binding between enantiomers and the chromatographic adsorbant establishes diastereomeric complexes, leading to differential partitioning in the mobile and bound states in the chromatographic system. The two enantiomers therefore move through the chromatographic system, e.g. column, at different rates, allowing for their separation

[0151] Further embodiments include prodrugs of the compounds according to any embodiment described herein, *i.e.* compounds which release an active compound according to any of the embodiments described herein, *in vivo* when administered to a mammalian subject. A prodrug is a pharmacologically active or more typically an inactive compound that is converted into a pharmacologically active agent by a metabolic transformation. Prodrugs of a compound according to any embodiment described herein, are prepared by modifying functional groups present in the compound according to any embodiment described herein, in such a way that the modifications may be cleaved *in vivo* to release the parent compound. *In vivo*, a prodrug readily undergoes chemical changes under physiological conditions (*e.g.* are hydrolyzed or acted on by naturally occurring enzyme(s)) resulting in liberation of the pharmacologically active agent. Prodrugs include compounds according to any embodiment described herein, wherein a hydroxyl, amino, or carboxy group is bonded to any group that may be cleaved *in vivo* to regenerate the free hydroxyl, amino or carboxy group, respectively. Examples of prodrugs include, but are not limited to esters (*e.g.*, acetate, formate, and benzoate derivatives) of compounds according to any embodiment described herein, or any other derivative which upon being brought to the physiological pH or through enzyme action is converted to the active parent drug. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described in the art (see, for example, Bundgaard. Design of Prodrugs. Elsevier, 1985).

[0152] The invention also embraces isolated compounds. An isolated compound refers to a compound which represents at least 10%, preferably at least 20%, more preferably at least 50% and most preferably at least 80% of the compound present in a mixture.

[0153] In some embodiments, one or more hydrogen atoms of a compound according to any embodiment described herein, is replaced by a deuterium. It is well established that deuteration of physiologically active compounds offer the advantage of retaining the pharmacological profile of their hydrogen counterparts while positively impacting their metabolic outcome. Selective replacement of one or more hydrogen with deuterium, in a compound according to any embodiment described herein, could improve the safety, tolerability and efficacy of the compound when compared to its all hydrogen counterpart.

[0154] Methods for incorporation of deuterium into compounds is well established. Using metabolic studies establish in the art, a compound according to any embodiment described herein, can be tested to identify sites for selective placement of a

deuterium isotope, wherein the isotope will not be metabolized. Moreover these studies identify sites of metabolism as the location where a deuterium atom would be placed.

Pharmaceutical Compositions

[0155] Some embodiments describe a pharmaceutical composition comprising: a compound according to any embodiment described herein, a pharmaceutically acceptable salt thereof, a solvate thereof, a stereoisomer thereof, a prodrug thereof, or an active metabolites thereof; and a pharmaceutically acceptable carrier or diluent. The pharmaceutical compositions can be prepared in a manner well known in the pharmaceutical art, and can be administered by a variety of routes, depending upon whether local or systemic treatment is desired and upon the area to be treated.

[0156] While it is possible that a compound as described in any embodiment herein, may be administered as the bulk substance, it is preferable to present the compound in a pharmaceutical formulation, *e.g.*, wherein the active agent is in an admixture with a pharmaceutically acceptable carrier selected with regard to the intended route of administration and standard pharmaceutical practice.

[0157] In particular, the disclosure provides a pharmaceutical composition comprising a therapeutically effective amount of at least one compound according to any embodiment described herein, and optionally, a pharmaceutically acceptable carrier.

Combinations

[0158] For the pharmaceutical compositions and methods of the disclosure, a compound according to any embodiment described herein, may be used in combination with other therapies and/or active agents.

[0159] In some embodiments, the compound according to any embodiment described herein, can be combined with one or more of a cholinesterase inhibitor, an N-methyl-D-aspartate (NMDA) glutamate receptor antagonist, a beta-amyloid specific antibody, a beta-secretase 1 (BACE1, beta-site amyloid precursor protein cleaving enzyme 1) inhibitor, a tumor necrosis factor alpha (TNF alpha) modulator, an intravenous immunoglobulin (IVIG), or a prion protein antagonist. In some embodiments the compound is combined with a cholinesterase inhibitor selected from tacrine (COGNEX®; Sciele), donepezil (ARICEPT®; Pfizer), rivastigmine (EXELON®; Novartis), or galantamine (RAZADYNE®; Ortho-McNeil-Janssen). In some embodiments, the compound is combined with a TNFalpha

modulator that is perispinal etanercept (ENBREL®, Amgen/Pfizer). In some embodiments, the compound is combined with a beta-amyloid specific antibody selected from bapineuzumab (Pfizer), solanezumab (Lilly), PF-04360365 (Pfizer), GSK933776 (GlaxoSmithKline), Gammagard (Baxter) or Octagam (Octapharma). In some embodiments, the compound is combined with an NMDA receptor antagonist that is memantine (NAMENDA®; Forest). In some embodiments, the BACE1 inhibitor is MK-8931 (Merck). In some embodiments, the compound is combined with IVIG as described in Magga et al., J Neuroinflamm 2010, 7:90, Human intravenous immunoglobulin provides protection against Ab toxicity by multiple mechanisms in a mouse model of Alzheimer's disease, and Whaley et al., 2011, Human Vaccines 7:3, 349-356, Emerging antibody products and Nicotiana manufacturing; each of which is incorporated herein by reference. In some embodiments, the compound is combined with a prion protein antagonist as disclosed in Strittmatter et al., US 2010/0291090, which is incorporated herein by reference.

[0160] Accordingly, the disclosure provides, in a further aspect, a pharmaceutical composition comprising at least one compound according to any embodiment described herein, or pharmaceutically acceptable derivative thereof; a second active agent; and, optionally a pharmaceutically acceptable carrier.

[0161] When combined in the same formulation it will be appreciated that the two or more compounds must be stable and compatible with each other and the other components of the formulation. When formulated separately they may be provided in any convenient formulation, in such manner as are known for such compounds in the art.

[0162] Preservatives, stabilizers, dyes and flavoring agents may be provided in any pharmaceutical composition described herein. Examples of preservatives include sodium benzoate, ascorbic acid and esters of p-hydroxybenzoic acid. Antioxidants and suspending agents may be also used.

[0163] With respect to combinations including biologics such as monoclonal antibodies or fragments, suitable excipients will be employed to prevent aggregation and stabilize the antibody or fragment in solution with low endotoxin, generally for parenteral administration, for example, intravenous, administration. For example, see Formulation and Delivery Issues for Monoclonal Antibody Therapeutics, Daugherty et al., in Current Trends in Monoclonal Antibody Development and Manufacturing, Part 4, 2010, Springer, New York pp 103-129.

[0164] The compounds according to any embodiment described herein, may be milled using known milling procedures such as wet milling to obtain a particle size

appropriate for tablet formation and for other formulation types. Finely divided (nanoparticulate) preparations of the compounds may be prepared by processes known in the art, for example see WO 02/00196 (SmithKline Beecham).

[0165] Compounds according to any embodiment described herein, or pharmaceutically acceptable salts thereof, a solvate thereof, a stereoisomer thereof, a prodrug thereof, or an active metabolites thereof, can be formulated for any route of administration.

Routes of Administration and Unit Dosage Forms

[0166] The routes for administration (delivery) include, but are not limited to, one or more of: oral (e.g., as a tablet, capsule, or as an ingestible solution), topical, mucosal (e.g., as a nasal spray or aerosol for inhalation), parenteral (e.g., by an injectable form), gastrointestinal, intraspinal, intraperitoneal, intramuscular, intravenous, intracerebroventricular, or other depot administration etc.

[0167] Therefore, the pharmaceutical compositions according to any embodiment described herein, include those in a form especially formulated for the mode of administration. In certain embodiments, the pharmaceutical compositions of the disclosure are formulated in a form that is suitable for oral delivery. In some embodiments, the compound is an orally bioavailable compound, suitable for oral delivery. In other embodiments, the pharmaceutical compositions of the disclosure are formulated in a form that is suitable for parenteral delivery.

[0168] The compounds according to any embodiment described herein, may be formulated for administration in any convenient way for use in human or veterinary medicine and the disclosure therefore includes within its scope pharmaceutical compositions comprising a compound according to any embodiment described herein, adapted for use in human or veterinary medicine. Such pharmaceutical compositions may be presented for use in a conventional manner with the aid of one or more suitable carriers. Acceptable carriers for therapeutic use are well-known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co. (A. R. Gennaro edit. 1985). The choice of pharmaceutical carrier can be selected with regard to the intended route of administration and standard pharmaceutical practice. The pharmaceutical compositions may comprise as, in addition to, the carrier any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), and/or solubilizing agent(s).

[0169] There may be different pharmaceutical composition/formulation requirements depending on the different delivery systems. It is to be understood that not all

of the compounds need to be administered by the same route. Likewise, if the pharmaceutical composition comprises more than one active component, then those components may be administered by different routes. By way of example, the pharmaceutical composition of the disclosure may be formulated to be delivered using a mini-pump or by a mucosal route, for example, as a nasal spray or aerosol for inhalation or ingestible solution, or parenterally in which the pharmaceutical composition is formulated by an injectable form, for delivery by, for example, an intravenous, intramuscular or subcutaneous route. Alternatively, the formulation may be designed to be delivered by multiple routes.

[0170] The combination of a compound according to any embodiment described herein, and an antibody or antibody fragment molecule can be formulated and administered by any of a number of routes and are administered at a concentration that is therapeutically effective in the indication or for the purpose sought. To accomplish this goal, the antibodies may be formulated using a variety of acceptable excipients known in the art. Typically, the antibodies are administered by injection, for example, intravenous injection. Methods to accomplish this administration are known to those of ordinary skill in the art. For example, Gokarn et al., 2008, J Pharm Sci 97(8):3051-3066, incorporated herein by reference, describe various high concentration antibody self buffered formulations. For example, monoclonal antibodies in self buffered formulation at e.g., 50 mg/mL mAb in 5.25% sorbitol, pH 5.0; or 60 mg/mL mAb in 5% sorbitol, 0.01% polysorbate 20, pH 5.2; or conventional buffered formulations, for example, 50 mg/mL mAb1 in 5.25% sorbitol, 25 or 50 mM acetate, glutamate or succinate, at pH 5.0; or 60 mg/mL in 10 mM acetate or glutamate, 5.25% sorbitol, 0.01% polysorbate 20, pH 5.2; other lower concentration formulations can be employed as known in the art..

[0171] Because some compounds of the disclosure cross the blood brain barrier they can be administered by a variety of methods including for example systemic (e.g., by iv, SC, oral, mucosal, transdermal route) or localized methods (e.g., intracranially). Where the compound according to any embodiment described herein, is to be delivered mucosally through the gastrointestinal mucosa, it should be able to remain stable during transit though the gastrointestinal tract; for example, it should be resistant to proteolytic degradation, stable at acid pH and resistant to the detergent effects of bile. For example, compounds according to any embodiment described herein, prepared for oral administration may be coated with an enteric coating layer. The enteric coating layer material may be dispersed or dissolved in either water or in a suitable organic solvent. As enteric coating layer polymers, one or more,

separately or in combination, of the following can be used; e.g., solutions or dispersions of methacrylic acid copolymers, cellulose acetate phthalate, cellulose acetate butyrate, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate succinate, polyvinyl acetate phthalate, cellulose acetate trimellitate, carboxymethylethylcellulose, shellac or other suitable enteric coating layer polymer(s). In some embodiments, the aqueous enteric coating layer is a methacrylic acid copolymer.

[0172] Where appropriate, the pharmaceutical compositions according to any embodiment described herein, can be administered by inhalation, by use of a skin patch, orally in the form of tablets containing excipients such as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs, solutions or suspensions containing flavoring or coloring agents, or they can be injected parenterally, for example intravenously, intramuscularly or subcutaneously. For buccal or sublingual administration the pharmaceutical compositions according to any embodiment described herein, may be administered in the form of tablets or lozenges, which can be formulated in a conventional manner.

[0173] Where the pharmaceutical composition according to any embodiment described herein, is to be administered parenterally, such administration includes without limitation: intravenously, intraarterially, intrathecally, intraventricularly, intracranially, intramuscularly or subcutaneously administering the compound of the disclosure; and/or by using infusion techniques. Antibodies or fragments are typically administered parenterally, for example, intravenously.

[0174] Pharmaceutical compositions according to any embodiment described herein, suitable for injection or infusion may be in the form of a sterile aqueous solution, a dispersion or a sterile powder that contains the active ingredient, adjusted, if necessary, for preparation of such a sterile solution or dispersion suitable for infusion or injection. This preparation may optionally be encapsulated into liposomes. In all cases, the final preparation must be sterile, liquid, and stable under production and storage conditions. To improve storage stability, such preparations may also contain a preservative to prevent the growth of microorganisms. Prevention of the action of micro-organisms can be achieved by the addition of various antibacterial and antifungal agents, e.g., paraben, chlorobutanol, or ascorbic acid. In many cases isotonic substances are recommended, e.g., sugars, buffers and sodium chloride to assure osmotic pressure similar to those of body fluids, particularly blood. Prolonged absorption of such injectable mixtures can be achieved by introduction of absorption-delaying agents, such as aluminum monostearate or gelatin.

[0175] Dispersions can be prepared in a liquid carrier or intermediate, such as glycerin, liquid polyethylene glycols, triacetin oils, and mixtures thereof. The liquid carrier or intermediate can be a solvent or liquid dispersive medium that contains, for example, water, ethanol, a polyol (e.g., glycerol, propylene glycol or the like), vegetable oils, non-toxic glycerine esters and suitable mixtures thereof. Suitable flowability may be maintained, by generation of liposomes, administration of a suitable particle size in the case of dispersions, or by the addition of surfactants.

[0176] For parenteral administration, the compound according to any embodiment described herein, is best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood. The aqueous solutions should be suitably buffered (preferably to a pH of from 3 to 9), if necessary. The preparation of suitable parenteral formulations under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art.

[0177] Sterile injectable solutions can be prepared by mixing a compound according to any embodiment described herein, with an appropriate solvent and one or more of the aforementioned carriers, followed by sterile filtering. In the case of sterile powders suitable for use in the preparation of sterile injectable solutions, preferable preparation methods include drying in vacuum and lyophilization, which provide powdery mixtures of the compounds and desired excipients for subsequent preparation of sterile solutions.

[0178] The compounds according to any embodiment described herein, may be formulated for use in human or veterinary medicine by injection (e.g., by intravenous bolus injection or infusion or via intramuscular, subcutaneous or intrathecal routes) and may be presented in unit dose form, in ampoules, or other unit-dose containers, or in multi-dose containers, if necessary with an added preservative. The pharmaceutical compositions for injection may be in the form of suspensions, solutions, or emulsions, in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing, solubilizing and/or dispersing agents. Alternatively the active ingredient may be in sterile powder form for reconstitution with a suitable vehicle, e.g., sterile, pyrogen-free water, before use.

[0179] The compounds according to any embodiment described herein, can be administered in the form of tablets, capsules, troches, ovules, elixirs, solutions or suspensions, for immediate-, delayed-, modified-, sustained-, pulsed- or controlled-release applications.

[0180] The compounds according to any embodiment described herein, may also be presented for human or veterinary use in a form suitable for oral or buccal administration, for example in the form of solutions, gels, syrups, or suspensions, or a dry powder for reconstitution with water or other suitable vehicle before use. Solid pharmaceutical compositions such as tablets, capsules, lozenges, troches, pastilles, pills, boluses, powder, pastes, granules, bullets or premix preparations may also be used. Solid and liquid pharmaceutical compositions for oral use may be prepared according to methods well-known in the art. Such pharmaceutical compositions may also contain one or more pharmaceutically acceptable carriers and excipients which may be in solid or liquid form.

[0181] The tablets may contain excipients such as microcrystalline cellulose, lactose, sodium citrate, calcium carbonate, dibasic calcium phosphate and glycine, disintegrants such as starch (preferably corn, potato or tapioca starch), sodium starch glycolate, croscarmellose sodium and certain complex silicates, and granulation binders such as polyvinylpyrrolidone, hydroxypropylmethylcellulose (HPMC), hydroxypropylcellulose (HPC), sucrose, gelatin and acacia.

[0182] Additionally, lubricating agents such as magnesium stearate, stearic acid, glyceryl behenate and talc may be included.

[0183] The pharmaceutical compositions according to any embodiment described herein, may be administered orally, in the form of rapid or controlled release tablets, microparticles, mini tablets, capsules, sachets, and oral solutions or suspensions, or powders for the preparation thereof. Oral preparations may optionally include various standard pharmaceutical carriers and excipients, such as binders, fillers, buffers, lubricants, glidants, dyes, disintegrants, odorants, sweeteners, surfactants, mold release agents, antiadhesive agents and coatings. Some excipients may have multiple roles in the pharmaceutical compositions, e.g., act as both binders and disintegrants.

[0184] Examples of pharmaceutically acceptable disintegrants for oral pharmaceutical compositions according to any embodiment described herein, include, but are not limited to, starch, pre-gelatinized starch, sodium starch glycolate, sodium carboxymethylcellulose, croscarmellose sodium, microcrystalline cellulose, alginates, resins, surfactants, effervescent compositions, aqueous aluminum silicates and cross-linked polyvinylpyrrolidone.

[0185] Examples of pharmaceutically acceptable binders for oral pharmaceutical compositions according to any embodiment described herein, include, but are not limited to, acacia; cellulose derivatives, such as methylcellulose, carboxymethylcellulose,

hydroxypropylmethylcellulose, hydroxypropylcellulose or hydroxyethylcellulose; gelatin, glucose, dextrose, xylitol, polymethacrylates, polyvinylpyrrolidone, sorbitol, starch, pre-gelatinized starch, tragacanth, xanthine resin, alginates, magnesium□aluminum silicate, polyethylene glycol or bentonite.

[0186] Examples of pharmaceutically acceptable fillers for oral pharmaceutical compositions according to any embodiment described herein, include, but are not limited to, lactose, anhydrolactose, lactose monohydrate, sucrose, dextrose, mannitol, sorbitol, starch, cellulose (particularly microcrystalline cellulose), dihydro- or anhydro-calcium phosphate, calcium carbonate and calcium sulphate.

[0187] Examples of pharmaceutically acceptable lubricants useful in the pharmaceutical compositions according to any embodiment described herein, include, but are not limited to, magnesium stearate, talc, polyethylene glycol, polymers of ethylene oxide, sodium lauryl sulphate, magnesium lauryl sulphate, sodium oleate, sodium stearyl fumarate, and colloidal silicon dioxide.

[0188] Examples of suitable pharmaceutically acceptable odorants for the oral pharmaceutical compositions according to any embodiment described herein, include, but are not limited to, synthetic aromas and natural aromatic oils such as extracts of oils, flowers, fruits (e.g., banana, apple, sour cherry, peach) and combinations thereof, and similar aromas. Their use depends on many factors, the most important being the organoleptic acceptability for the population that will be taking the pharmaceutical compositions.

[0189] Examples of suitable pharmaceutically acceptable dyes for the oral pharmaceutical compositions according to any embodiment described herein, include, but are not limited to, synthetic and natural dyes such as titanium dioxide, beta-carotene and extracts of grapefruit peel.

[0190] Examples of useful pharmaceutically acceptable coatings for the oral pharmaceutical compositions according to any embodiment described herein, typically used to facilitate swallowing, modify the release properties, improve the appearance, and/or mask the taste of the pharmaceutical compositions include, but are not limited to, hydroxypropylmethylcellulose, hydroxypropylcellulose and acrylate-methacrylate copolymers.

[0191] Suitable examples of pharmaceutically acceptable sweeteners for the oral pharmaceutical compositions according to any embodiment described herein, include, but are not limited to, aspartame, saccharin, saccharin sodium, sodium cyclamate, xylitol, mannitol, sorbitol, lactose and sucrose.

[0192] Suitable examples of pharmaceutically acceptable buffers include, but are not limited to, citric acid, sodium citrate, sodium bicarbonate, dibasic sodium phosphate, magnesium oxide, calcium carbonate and magnesium hydroxide.

[0193] Suitable examples of pharmaceutically acceptable surfactants include, but are not limited to, sodium lauryl sulphate and polysorbates.

[0194] Solid compositions of a similar type may also be employed as fillers in gelatin capsules. Preferred excipients in this regard include lactose, starch, a cellulose, milk sugar or high molecular weight polyethylene glycols. For aqueous suspensions and/or elixirs, the agent may be combined with various sweetening or flavoring agents, coloring matter or dyes, with emulsifying and/or suspending agents and with diluents such as water, ethanol, propylene glycol and glycerin, and combinations thereof.

[0195] As indicated, a compounds according to any embodiment described herein, can be administered intranasally or by inhalation and is conveniently delivered in the form of a dry powder inhaler or an aerosol spray presentation from a pressurized container, pump, spray or nebulizer with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, a hydrofluoroalkane such as 1,1,1,2-tetrafluoroethane (HFA 134AT) or 1,1,1,2,3,3,3-heptafluoropropane (HFA 227EA), carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurized container, pump, spray or nebulizer may contain a solution or suspension of the active compound, e.g., using a mixture of ethanol and the propellant as the solvent, which may additionally contain a lubricant, e.g., sorbitan trioleate.

[0196] Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated to contain a powder mix of a compound according to any embodiment described herein, and a suitable powder base such as lactose or starch.

[0197] For topical administration by inhalation a compounds according to any embodiment described herein, may be delivered for use in human or veterinary medicine via a nebulizer.

[0198] The pharmaceutical compositions of the disclosure may contain from 0.01 to 99% weight per volume of the active material. For topical administration, for example, the pharmaceutical composition will generally contain from 0.01-10%, more preferably 0.01-1% of the active material.

[0199] A compound according to any embodiment described herein, can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles,

large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

[0200] The pharmaceutical composition or unit dosage form, according to any embodiment described herein, may be administered according to a dosage and administration regimen defined by routine testing in the light of the guidelines given above in order to obtain optimal activity while minimizing toxicity or side effects for a particular patient. The dosage of the compounds or unit dosage form may vary according to a variety of factors such as underlying disease conditions, the individual's condition, weight, sex and age, and the mode of administration. The exact amount to be administered to a patient will vary depending on the state and severity of the disorder and the physical condition of the patient. A measurable amelioration of any symptom or parameter can be determined by a person skilled in the art or reported by the patient to the physician. It will be understood that any clinically or statistically significant attenuation or amelioration of any symptom or parameter is within the scope of the disclosure. Clinically significant attenuation or amelioration means perceptible to the patient and/or to the physician.

[0201] In some embodiments, the amount of the compound to be administered can range between about 0.01 and about 25 mg/kg/day. Generally, dosage levels of between 0.01 to 25 mg/kg of body weight daily are administered to the patient, e.g., humans. In some embodiments the therapeutically effective amount is between a lower limit of about 0.01 mg/kg of body weight, about 0.1 mg/kg of body weight, about 0.2 mg/kg of body weight, about 0.3 mg/kg of body weight, about 0.4 mg/kg of body weight, about 0.5 mg/kg of body weight, about 0.60 mg/kg of body weight, about 0.70 mg/kg of body weight, about 0.80 mg/kg of body weight, about 0.90 mg/kg of body weight, about 1 mg/kg of body weight, about 2.5 mg/kg of body weight, about 5 mg/kg of body weight, about 7.5 mg/kg of body weight, about 10 mg/kg of body weight, about 12.5 mg/kg of body weight, about 15 mg/kg of body weight, about 17.5 mg/kg of body weight, about 20 mg/kg of body weight, about 22.5 mg/kg of body weight, and about 25 mg/kg of body weight; and an upper limit of 25 mg/kg of body weight, about 22.5 mg/kg of body weight, about 20 mg/kg of body weight, about 17.5 mg/kg of body weight, about 15 mg/kg of body weight, about 12.5 mg/kg of body weight, about 10 mg/kg of body weight, about 7.5 mg/kg of body weight, about 5 mg/kg of body weight, about 2.5 mg/kg of body weight, about 1 mg/kg of body weight, about 0.9 mg/kg of body weight, about 0.8 mg/kg of body weight, about 0.7 mg/kg of body weight, about 0.6 mg/kg of body weight, about 0.5 mg/kg of body weight, about 0.4 mg/kg of body weight, about 0.3 mg/kg of body weight, about 0.2 mg/kg of body weight, about 0.1 mg/kg of

body weight, and about 0.01 mg/kg of body weight. In some embodiments, the therapeutically effective amount is about 0.1 mg/kg/day to about 10 mg/kg/day; in some embodiments the therapeutically effective amount is about 0.2 and about 5 mg/kg/day. It will be understood that the pharmaceutical formulations of the disclosure need not necessarily contain the entire amount of the compound that is effective in treating the disorder, as such effective amounts can be reached by administration of a plurality of divided doses of such pharmaceutical formulations. The compounds may be administered on a regimen of 1 to 4 times per day, such as once, twice, three times or four times per day.

[0202] In some embodiments of the disclosure, a compound according to any embodiment described herein, is formulated in capsules or tablets, usually containing about 10 to about 200 mg of the compounds. In some embodiments the capsule or tablet contains between a lower limit of about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg; about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, and about 200 mg, and an upper limit of about 200 mg, about 195 mg, about 190 mg, about 185 mg, about 180 mg, about 175 mg, about 170 mg, about 165 mg, about 160 mg, about 155 mg, about 150 mg, about 145 mg, about 140 mg, about 135 mg, about 130 mg, about 125 mg, about 120 mg, about 115 mg, about 110 mg, about 105 mg, about 100 mg, about 95 mg, about 90 mg; about 85 mg, about 80 mg, about 75 mg, about 70 mg, about 65 mg, about 60 mg, about 55 mg, about 50 mg, about 45 mg, about 40 mg, about 35 mg, about 30 mg, about 25 mg, about 20 mg, about 15 mg, and about 10 mg of a compound according to any embodiment herein.

[0203] In some embodiments, a compound according to any embodiment herein is administered to a patient at a total daily dose of 50 mg to 500 mg. In some embodiments, the daily dose is between a lower limit of about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg; about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg; about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg,

about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, 300 mg, about 305 mg, about 310 mg, about 315 mg; about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395, about 400 mg, about 405 mg, about 410 mg, about 415 mg; about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, and about 500 mg and an upper limit of about 500 mg, about 495 mg, about 490 mg, about 485 mg, about 480 mg, about 475 mg, about 470 mg, about 465 mg, about 460 mg, about 455 mg, about 450 mg, about 445 mg, about 440 mg, about 435 mg, about 430 mg, about 425 mg, about 420 mg, about 415 mg, about 410 mg, about 405 mg, about 400 mg, about 395 mg, about 390 mg, about 385 mg, about 380 mg, about 375 mg, about 370 mg, about 365 mg, about 360 mg, about 355 mg, about 350 mg, about 345 mg, about 340 mg, about 335 mg, about 330 mg, about 325 mg, about 320 mg, about 315 mg, about 310 mg, about 305 mg about 300 mg, about 295 mg, about 290 mg, about 285 mg, about 280 mg, about 275 mg, about 270 mg, about 265 mg, about 260 mg, about 255 mg, about 250 mg, about 245 mg, about 240 mg, about 235 mg, about 230 mg, about 225 mg, about 220 mg, about 215 mg, about 210 mg, about 205 mg 200 mg, about 195 mg, about 190 mg, about 185 mg, about 180 mg, about 175 mg, about 170 mg, about 165 mg, about 160 mg, about 155 mg, about 150 mg, about 145 mg, about 140 mg, about 135 mg, about 130 mg, about 125 mg, about 120 mg, about 115 mg, about 110 mg, about 105 mg, about 100 mg, about 95 mg, about 90 mg; about 85 mg, about 80 mg, about 75 mg, about 70 mg, about 65 mg, about 60 mg, about 55 mg, and about 50 mg of a compound according to any embodiment herein. In some embodiments, the total daily dose is about 50 mg to 150 mg. In some embodiments, the total daily dose is about 50 mg to 250 mg. In some embodiments, the total daily dose is about 50 mg to 350 mg. In some embodiments, the total daily dose is about 50 mg to 450 mg. In some embodiments, the total daily dose is about 50 mg.

[0204] A pharmaceutical composition for parenteral administration contains from about 0.01% to about 100% by weight of the active compound according to any embodiment described herein, based upon 100% weight of total pharmaceutical composition.

[0205] Generally, transdermal dosage forms contain from about 0.01% to about 100% by weight of the active compound according to any embodiment described herein, versus 100% total weight of the dosage form.

[0206] The pharmaceutical composition or unit dosage form may be administered in a single daily dose, or the total daily dosage may be administered in divided doses. In addition, co administration or sequential administration of another compound for the treatment of the disorder may be desirable. To this purpose, the combined active principles are formulated into a simple dosage unit.

[0207] Compounds according to any embodiment described herein, may be prepared by the general methods outlined in, for example, WO2013/029057, incorporated herein by reference, or as described hereinafter, said methods constituting a further aspect of the disclosure.

[0208] Compounds according to any embodiment disclosed herein can be synthesized in accordance with general methods provided herein and specific synthetic examples.

[0209] It will be appreciated by those skilled in the art that it may be desirable to use protected derivatives of intermediates used in the preparation of the compounds according to any embodiment described herein. Protection and deprotection of functional groups may be performed by methods known in the art (see, for example, Green and Wuts Protective Groups in Organic Synthesis. John Wiley and Sons, New York, 1999.). Hydroxy or amino groups may be protected with any hydroxy or amino protecting group. The amino protecting groups may be removed by conventional techniques. For example, acyl groups, such as alkanoyl, alkoxycarbonyl and aroyl groups, may be removed by solvolysis, e.g., by hydrolysis under acidic or basic conditions. Arylmethoxycarbonyl groups (e.g., benzyloxycarbonyl) may be cleaved by hydrogenolysis in the presence of a catalyst such as palladium-on-charcoal.

[0210] The synthesis of the target compounds is completed by removing any protecting groups which may be present in the penultimate intermediates using standard techniques, which are well-known to those skilled in the art. The deprotected final products are then purified, as necessary, using standard techniques such as silica gel chromatography, HPLC on silica gel and the like, or by recrystallization.

Methods of Use

[0211] In some embodiments, the disclosure provides methods of inhibiting synapse number decline or membrane trafficking abnormalities associated with exposure of a neuronal cell to Abeta species by administration of a compound according to any embodiment described herein.

[0212] In some embodiments the disclosure also provides methods for treating cognitive decline and/or a neurodegenerative disease, e.g. Alzheimer's disease or mild cognitive impairment (MCI) in a patient comprising administering to the patient a compound according to any embodiment described herein.

[0213] In some embodiments, the neurodegenerative disease is selected from Age-Associated Memory Impairment (AAMI), Age-Related Cognitive Decline (ARCD), agitation synucleinopathies, Alzheimer's disease (AD), Amyotrophic lateral sclerosis (ALS) dementia, autosomal-dominant Parkinson's disease, Cognitive Impairment No Dementia (CIND), dementia, Diffuse Lewy Body Disease (DLBD) also known as Dementia with Lewy Bodies (DLB), disorders or conditions characterized by the presence of Lewy bodies, Down syndrome, dyskinesia, HIV dementia, Huntington's disease, Incidental LBD, Inherited LBD, Lewy body dysphagia, Mild Cognitive Impairment (MCI), multiple sclerosis, multiple system atrophy (MSA), Olivopontocerebellar Atrophy, Parkinson's disease (PD), preclinical Alzheimer's Disease (PCAD), Psychosis, Pure Autonomic Failure, Shy-Drager Syndrome, Striatonigral Degeneration, synucleinopathies, combined Alzheimer's and Parkinson disease and/or MSA, vascular dementia, diseases, disorders or conditions associated with abnormal expression, stability, activities and/or cellular processing of α -synuclein, diseases, disorders or conditions characterized by the presence of Lewy bodies, and combinations thereof.

[0214] In some embodiments, the method of inhibiting, or treating, cognitive decline and/or a neurodegenerative disease, e.g. Alzheimer's disease, comprises inhibiting, or treating one or more symptoms of cognitive decline selected from the group consisting of memory loss, confusion, impaired judgment, personality changes, disorientation, and loss of language skills. In some embodiments, the method comprises inhibiting, or treating, diseases or disorders or conditions mediated by or associated with Abeta oligomers.

[0215] In some embodiments, the method of inhibiting, or treating, cognitive decline and/or a neurodegenerative disease, e.g. Alzheimer's disease, comprises one or more of: (i) restoration of long term potentiation (LTP), long term depression (LTD) or synaptic plasticity detectable by electrophysiological measurements or any of the other negative changes in cognitive function as mentioned in the definition of the term above; and/or (ii) inhibiting, or treating, neurodegeneration; and/or (iii) inhibiting, or treating, general amyloidosis; and/or (iv) inhibiting, or treating, one or more of amyloid production, amyloid assembly, amyloid aggregation, and amyloid oligomer binding, and amyloid deposition; and/or (v) inhibiting, treating, and/or abating an effect, notably a nonlethal effect, of one or more of Abeta oligomers on a neuron cell.

[0216] In some embodiments, the method of inhibiting, treating, and/or abating cognitive decline and/or a neurodegenerative disease, e.g. Alzheimer's disease, comprises inhibiting, treating, and/or abating one or more of amyloid production, amyloid assembly, the activity/effect of one or more of Abeta oligomers on a neuron cell, amyloid aggregation, amyloid binding, and amyloid deposition.

[0217] In some embodiments, the method of inhibiting, treating, and/or abating cognitive decline and/or a neurodegenerative disease, e.g. Alzheimer's disease, comprises inhibiting, treating, and/or abating one or more of the activity/effect of one or more of Abeta oligomers on a neuron cell.

[0218] In some embodiments, the activity/effect of one or more of Abeta oligomers on a neuron cell, amyloid aggregation and amyloid binding is the effect of Abeta oligomers on membrane trafficking or synapse number. In some embodiments, a compound according to any embodiment described herein, inhibits the Abeta oligomer effect on membrane trafficking or synapse number or Abeta oligomer binding.

[0219] In some embodiments, the disclosure provides methods of treating a proteopathic disease associated with Abeta oligomer toxicity, specifically nonlethal Abeta oligomer effects. In some embodiments, the method comprises contacting a subject with such a proteopathic disease with a compound according to any embodiment described herein, or a pharmaceutical composition containing the same that binds the sigma-2 receptor.

[0220] In some embodiments, the proteopathic disease is a CNS proteopathy, characterized by an increase in Abeta protein, such as MCI, Down's Syndrome, macular degeneration or Alzheimer's disease, and the like.

[0221] In some embodiments, the disclosure provides methods of treating one or more mild cognitive impairment (MCI), or dementia by administering a compound according to any embodiment described herein. In some embodiments, the disclosure provides methods of treating MCI, and dementia.

[0222] In some embodiments, the disclosure provides methods of treating Alzheimer's Disease by administering a compound according to any embodiment described herein.

[0223] In some embodiments, the disclosure provides methods of treating an individual with a compound according to any embodiment described herein, to restore, partially or totally, the subject's cells to a normal phenotype in terms of functions affected adversely by Abeta species, such as Abeta oligomers. Examples are synaptic number reduction and membrane trafficking abnormalities, which can be measured by various

methods including assays described herein. The normal phenotype can be, for example, normal membrane trafficking. In some embodiments, the normal phenotype is normal cognitive ability. The “normal” phenotype can be determined by comparing a subject’s results with a sample of normal subjects. The sample may be as small as 1 subject or 1 sample or may be more than 10 samples or subjects and the norm is an average that is calculated based upon a plurality of subjects.

[0224] In some embodiments, a compound according to any embodiment described herein, generally inhibits the Abeta effect on neurons. In some embodiments, the compounds describe above have an IC_{50} for inhibition of Abeta effect of less than about 100 μ M, about 50 μ M, about 20 μ M, about 15 μ M, about 10 μ M, about 5 μ M, about 1 μ M, about 500 nM, about 100 nM, about 50 nM, or about 10 nM on neurons (such as neurons in the brain), amyloid assembly or disruption thereof, and amyloid (including amyloid oligomer) binding, and amyloid deposition. In some embodiments, a compound according to any embodiment described herein, may have an IC_{50} for inhibition of the activity/effect of Abeta species such as oligomers of less than about 100 μ M, about 50 μ M, about 20 μ M, about 15 μ M, about 10 μ M, about 5 μ M, about 1 μ M, about 500 nM, about 100 nM, about 50 nM, or about 10 nM on neurons (such as central nervous system neurons).

[0225] A compound according to any embodiment described herein, may inhibit the Abeta effect by specifically binding to a sigma-2 receptor. A compound can be said to be “specific” for a sigma-2 receptor when it binds with a binding affinity that is at least 10% greater than to the sigma-1 receptor, even though the compound is capable of binding both sigma-1 and sigma-2 receptor. The compounds of such embodiments may exhibit a specificity of at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 200%, 300%, 400%, 500%, or 1000% greater for sigma-2 receptor than sigma-1 receptor.

[0226] In some embodiments, percentage inhibition by a compound according to any embodiment described herein, of one or more of the effects of Abeta species such as oligomers on neurons (such as neurons in the brain), such as amyloid (including amyloid oligomer) binding to synapses, and abnormalities in membrane trafficking mediated by Abeta oligomer can be about 1% to about 20%, about 20% to about 50%, about 1% to about 50%, or about 1% to about 80% as measured at a concentration of from 10 nM to 10 μ M. Inhibition can be assessed for example by quantifying synapse number of a neuron prior to and after exposure to an amyloid beta species or quantifying the number of synapses in the presence of both of the compound according to any embodiment described herein, and the Abeta species wherein the compound according to any embodiment described herein, is

simultaneous with, or precedes or follows, Abeta species exposure. As another example, inhibition can be assessed by determining membrane trafficking and comparing one or more parameters that measure exocytosis rate and extent, endocytosis rate and extent, or other indicators of cell metabolism in the presence and absence of an Abeta species and in the presence and absence of a compound according to any embodiment described herein.

[0227] In some embodiments, the disclosure provides methods of measuring beta-amyloid-associated cognitive decline in an animal using a labeled compound according to any embodiment described herein. In some embodiments, the method comprises contacting the animal with a labeled compound according to any embodiment described herein, and measuring sigma-2 activity or expression. In some embodiments, the method comprises comparing the sigma-2 activity or expression in the animal with an animal known to have beta-amyloid induced cognitive decline. If the activity or expression is the same as the animal known to have beta-amyloid induced cognitive decline the animal is said to have the same level of cognitive decline. The animals can be ranked according the similarities in known activity or expression of various stages of beta amyloid induced cognitive decline. Any of the a compound according to any embodiment described herein, can be labeled so that the labeled compound can be used *in vivo*.

[0228] In some embodiments, an assay is used to determine if a compound according to any embodiment described herein, can bind to a sigma-2 receptor. In some embodiments, the method further comprises determining whether the compound that binds to a sigma-2 receptor acts as a functional antagonist at a sigma-2 receptor by inhibiting soluble A β oligomer induced neurotoxicity with respect to inhibiting soluble A β oligomer induced synapse loss, and inhibiting soluble A β oligomer induced deficits in a membrane trafficking assay.

[0229] Any form of amyloid β may be used in the practice of the screening methods and of the assays according to the disclosure, including amyloid β monomers, oligomers, fibrils, as well as amyloid β associated with proteins (“protein complexes”) and more generally amyloid β assemblies. For example, screening methods can employ various forms of soluble amyloid β oligomers as disclosed, for example, in U.S. patent application serial number 13/021,872; U.S. Patent Publication 2010/0240868; International Patent Application WO/2004/067561; International Patent Application WO/2010/011947; U.S. Patent Publication 20070098721; U.S. Patent Publication 20100209346; International Patent Application WO/2007/005359; U.S. Patent Publication 20080044356; U.S. Patent

Publication 20070218491; WO/2007/126473; U.S. Patent Publication 20050074763; International Patent Application WO/2007/126473, International Patent Application WO/2009/048631, and U.S. Patent Publication 20080044406, U.S. Patent No. 7,902,328 and U.S. Patent No. 6,218,506, each of which is incorporated herein by reference.

[0230] Amyloid β forms, including monomers or oligomers of amyloid β may be obtained from any source. For example, in some embodiments, commercially available amyloid β monomers and/or amyloid β oligomers may be used in the aqueous solution, and in other embodiments, amyloid β monomers and/or amyloid β oligomers that are used in the aqueous protein solution can be isolated and purified by the skilled artisan using any number of known techniques. In general, the amyloid β monomers and/or amyloid β oligomers used in the preparation of the aqueous solution of proteins and amyloid β of various embodiments may be soluble in the aqueous solution. Therefore, both the proteins of the aqueous solution and the amyloid β may be soluble.

[0231] The amyloid β added may be of any isoform. For example, in some embodiments, the amyloid β monomers may be amyloid β 1-42, and in other embodiments the amyloid β monomers may be amyloid β 1-40. In still other embodiments, the amyloid β may be amyloid β 1-39 or amyloid β 1-41. Hence, the amyloid β of various embodiments may encompass any C-terminal isoform of amyloid β . Yet other embodiments include amyloid β in which the N-terminus has been frayed, and in some embodiments, the N-terminus of any of amyloid β C-terminal isomers described above may be amino acid 2, 3, 4, 5, or 6. For example, amyloid β 1-42 may encompass amyloid β 2-42, amyloid β 3-42, amyloid β 4-42, or amyloid β 5-42 and mixtures thereof, and similarly, amyloid β 1-40 may encompass amyloid β 2-40, amyloid β 3-40, amyloid β 4-40, or amyloid β 5-40.

[0232] The amyloid β forms used in various embodiments may be wild type, i.e. having an amino acid sequence that is identical to the amino acid sequence of amyloid β synthesized *in vivo* by the majority of the population, or in some embodiments, the amyloid β may be a mutant amyloid β . Embodiments are not limited to any particular variety of mutant amyloid β . For example, in some embodiments, the amyloid β introduced into the aqueous solution may include a known mutation, such as, for example, amyloid β having the “Dutch” (E22Q) mutation or the “Arctic” (E22G) mutation. Such mutated monomers may include naturally occurring mutations such as, for example, forms of amyloid β isolated from populations of individuals that are predisposed to, for example, Alzheimer’s disease, familial

forms of amyloid β . In other embodiments, mutant amyloid β monomers may be synthetically produced by using molecular techniques to produce an amyloid β mutant with a specific mutation. In still other embodiments, mutant amyloid β monomers may include previously unidentified mutations such as, for example, those mutants found in randomly generated amyloid β mutants. The term “amyloid β ” as used herein, is meant to encompass both wild type forms of amyloid β as well as any of the mutant forms of amyloid β .

[0233] In some embodiments, the amyloid β in the aqueous protein solution may be of a single isoform. In other embodiments, various C-terminal isoforms of amyloid β and/or various N-terminal isoforms of amyloid β may be combined to form amyloid β mixtures that can be provided in the aqueous protein solution. In yet other embodiments, the amyloid β may be derived from amyloid precursor protein (APP) that is added to the protein containing aqueous solution and is cleaved *in situ*, and such embodiments, various isoforms of amyloid β may be contained within the solution. Fraying of the N-terminus and/or removal of C-terminal amino acids may occur within the aqueous solution after amyloid β has been added. Therefore, aqueous solutions prepared as described herein, may include a variety of amyloid β isoforms even when a single isoform is initially added to the solution.

[0234] The amyloid β monomers added to the aqueous solution may be isolated from a natural source such as living tissue, and in other embodiments, the amyloid β may be derived from a synthetic source such as transgenic mice or cultured cells. In some embodiments, the amyloid β forms, including monomers, oligomers, or combinations thereof are isolated from normal subjects and/or patients that have been diagnosed with cognitive decline or diseases associated therewith, such as, but not limited to, Alzheimer’s disease. In some embodiments, the amyloid β monomers, oligomers, or combinations thereof are Abeta assemblies that have been isolated from normal subjects or diseased patients. In some embodiments, the Abeta assemblies are high molecular weight, e.g. greater than 100KDa. In some embodiments, the Abeta assemblies are intermediate molecular weight, e.g. 10 to 100KDa. In some embodiments, the Abeta assemblies are less than 10 kDa.

[0235] The amyloid β oligomers of some embodiments may be composed of any number of amyloid β monomers consistent with the commonly used definition of “oligomer.” For example, in some embodiments, amyloid β oligomers may include from about 2 to about 300, about 2 to about 250, about 2 to about 200 amyloid β monomers, and in other embodiments, amyloid β oligomers may be composed from about 2 to about 150, about 2 to

about 100, about 2 to about 50, or about 2 to about 25, amyloid β monomers. In some embodiments, the amyloid β oligomers may include 2 or more monomers. The amyloid β oligomers of various embodiments may be distinguished from amyloid β fibrils and amyloid β protofibrils based on the confirmation of the monomers. In particular, the amyloid β monomers of amyloid β oligomers are generally globular consisting of β -pleated sheets whereas secondary structure of the amyloid β monomers of fibrils and protofibrils is parallel β -sheets.

EXAMPLES

[0236] Examples 1 and 2 describe Abeta oligomer preparations that could be used for experiments described below. The particular preparations used in the membrane trafficking and oligomer binding/synapse reduction assays as well as those used in the *in vivo* assays described below are each described in the example to which they pertain.

Example 1: Preparation of Amyloid β Oligomers

[0237] The conditions in which amyloid β may oligomerize in nervous tissue, a milieu of aqueous-soluble proteins with which it may associate, were re-created to identify the more disease-relevant structural state of amyloid β oligomers and fibrils. Aqueous soluble proteins were prepared from rat brain by ultracentrifugation. Specifically, 5 volumes of TBS buffer (20mM Tris-HCL, pH 7.5, 34mM NaCl and a complete protease inhibitor cocktail (Santa Cruz) per gram of brain tissue was added to the rat brain tissue on ice. Dounce homogenization was then carried out with a tight-fitting pestle. The homogenized brain tissues were then centrifuged at 150,000 x g for 1 hour at 4° C (40,000 rpm Ty65). The infranatant (between floating myelin and a half cm above the pellet) was then removed and aliquots were frozen at -75° C. The pellets were then resuspended in TBS to the original volume and frozen in aliquots at -75° C. Synthetic, monomeric human amyloid β 1-42 was added to this mixture to provide a final concentration of 1.5uM amyloid β , and the solution was incubated for 24 hours at 4° C. Centrifugation of the mixture at 5,800g for 10 minutes was performed to remove fibrillar assemblies and then Immunoprecipitation was performed using 6E10 conjugated agarose spin columns (Pierce Chemical Company) for 24 hours at 4° C. The eluted amyloid β oligomers were then subject to MALDI -Tof mass spectroscopic analysis to identify the contents of the sample.

[0238] The amyloid β self-associated in the protein containing solution to form subunit assemblies of 22,599 Da, 5 subunit pentamers and 31,950 Da, 7 subunit, 7mers. Another peak at 49,291 Da may represent 12 subunit, 12mers, although this would not appear

to be an accurate molecular weight for amyloid β 12mers. Notably, no peaks are observed at either 4518 Da or 9036 Da which would represent amyloid β monomers and dimers. However, peaks at 9,882 Da and 14,731 Da could represent amyloid β dimers associated with a 786 Da (or 2 x 393 Da) lipids or proteins and amyloid β trimers associated with 3 x 393 Da lipids or proteins, respectively. In addition, the presence of a peak at 19,686 Da is indicative of an assembly state possibly involving a trimer complex and a rat amyloid β fragment of 4954 Da. Accordingly these data may reflect the association of small lipids or proteins with dimers and trimers of amyloid β which may direct the assembly of conformational states unique to physiological systems.

Example 2 Preparation of beta-amyloid oligomers

[0239] A solution of 1.5uM monomeric human amyloid β 1-42 in a mixture of rat brain soluble proteins was incubated for 24 hours at 4° C as described in Example 1. This solution was then treated with tri-fluoro ethanol (TFE) prior to taking the spectra. In TFE, assembled protein structures and non-covalently bound protein complexes dissociate into denatured proteins, and the peaks associated with assembled oligomers are expected to disappear. The majority of protein peaks observed in Example 1 disappeared including the 9822 Da, 14,731 Da, 31,950 Da, and 49,291 Da peaks identified above. However, an abundant peak is observed at 4518 Da which represents amyloid β monomer peak. A peak at 4954.7 is apparent which may represent a longer abeta fragment similar to amyloid β 1-46. An additional peak is observed at 7086 Da which was not present in the preparation described in Example 1, which may represent amyloid β monomers associated with a 2550 Da covalently bound protein.

Example 3: Isolation of beta-amyloid oligomers from human AD brain tissue.

[0240] TBS soluble extracts: Samples of post-mortem brain tissue from human patients characterized via histopathological analysis as Braak Stage V/VI Alzheimer's disease (AD) were obtained from a hospital brain tissue bank. Age and gender matched AD and normal tissue specimens were diluted to 0.15gm tissue/ml in 20mM Tris-HCL, 137mM NaCl, pH 7.6 containing 1mM EDTA and 1mg/ml complete protease inhibitor cocktail (Sigma P8340) and homogenized. Ultracentrifugation of the tissue homogenates was performed at 105,000g for 1 hour in a Beckman Optima XL-80K Ultracentrifuge. The resulting TBS soluble fractions were immunodepleted using protein-A and protein-G agarose columns (Pierce Chemical) and then size fractionated with Amicon Ultra 3, 10 & 100 kDa NMWCO filters (Millipore Corporation).

[0241] Immunoprecipitation: Size fractionated and immunodepleted TBS soluble extracts were concentrated to approximately 200ul in the appropriate NMWCO Amicon Ultra filters. The concentrated TBS soluble extracts were diluted up to 400ul with TBS sample buffer (Pierce Chemical) and centrifuged for 10 minutes at 5,800 g to remove fibrils. The resulting supernatant was then immunoprecipitated with 6E10-conjugated agarose beads overnight at 4 °C followed by antigen elution using high osmotic strength Gentle elution buffers (Pierce Chemical) to isolate Abeta containing protein species.

[0242] MALDI-mass spectrometry: Immunoisolated beta amyloid was subjected to mass spectroscopic analysis using an Applied Biosystems (ABI) Voyager DE-Pro MALDI-Tof instrument. Samples were analyzed using various matrix types such as α -Cyano-4-hydroxycinnamic acid (CHCA), Sinapic acid (SA), or 6-Aza-2-thiothymine (ATT) depending on the target molecular weight range of the analysis. The instrument was run in a linear-positive ion mode along with a variable extraction delay. Non-accumulated spectra represented 100 shots of a “hot spot” per acquisition while accumulated spectra were represented by 12 separate areas of each spot with 200 laser shots per acquisition.

[0243] Data analysis: Data acquisition and analysis was performed using Voyager’s Data Explorer software package. Standard processing of the mass spectra included smoothing of the spectrum and baseline subtraction functions in addition to variations in the signal to noise ratio.

[0244] ELISA for Ab quantification: Immunoprecipitated TBS soluble fractions were analyzed for both “total” Abeta and Abeta oligomer concentration using a modified sandwich ELISA technique. Briefly, 6E10 and 4G8 coated Nunc MaxiSorp 96-well plates were incubated with Abeta containing samples and then probed with a Biotinylated 4G8 detection antibody. Incubation with Streptavidin-HRP (Rockland) followed by development of a Tetramethyl benzidine (TMB) substrate allowed for colorimetric detection (OD 450) of abeta on a BioTEk Synergy HT plate reader. Monomeric Abeta 1-42 was used for generation of a standard curve and along with GEN 5 software allowed for quantification of Abeta levels in the immuno-precipitated samples.

Example 4: Receptor Binding Assays

[0245] Certain compounds were tested for interaction with several receptors by blocking the binding or action of their agonists or antagonists. Some compounds were tested to see whether they interact directly with known cellular receptor or signaling proteins. Compounds can be tested for the ability to displace binding of known agonists or antagonists

of a given human receptor that was overexpressed in cell lines or isolated from tissue. Compounds can also be tested for the ability to block downstream signaling induced by agonists or antagonists of a given human receptor. Compounds can be tested for action at 100 known receptors, and it is desirable that specific activity will occur at only a small subset of CNS-relevant receptors.

[0246] Using the same protocol, some compounds for which membrane trafficking data are given in Table 1 were tested for recognition of sigma-2 receptor. Certain compounds preferentially bind to the sigma-2 receptor.

Competitive Radioligand Binding Assay 1

[0247] For Sigma-1 binding, various concentrations of test compounds from 100 μ M to 1 nM were used to displace 8 nM [3H](+)pentazocine from endogenous receptors on Jurkat cell membranes (Ganapathy ME et al. 1991, J Pharmacol. Exp. Ther. 289:251-260). 10 μ M Haloperidol was used to define non-specific binding. For Sigma-2 receptors various concentrations of test compounds from 100 μ M to 1 nM were used to displace 5 nM [3H] 1,3-Di-(2-tolyl)guanidine from endogenous receptors on membranes from rat cerebral cortex in the presence of 300 nM (+)pentazocine to mask Sigma-1 receptors. (Bowen WD, et al. 1993, Mol. Neuropharmacol 3:117-126). 10 μ M Haloperidol was used to define non-specific binding. Reactions were terminated by rapid filtration through Whatman GF/C filters using a Brandel 12R cell harvester followed by two washes with ice-cold buffer. Radioactivity on the dried filter discs was measured using a liquid scintillation analyzer (Tri-Carb 2900TR; PerkinElmer Life and Analytical Sciences). The displacement curves were plotted and the K_i values of the test compounds for the receptor subtypes were determined using GraphPad Prism (GraphPad Software Inc., San Diego, CA). The percentage specific binding was determined by dividing the difference between total bound (disintegrations per minute) and nonspecific bound (disintegrations per minute) by the total bound (disintegrations per minute).

[0248] Affinities for Sigma-1 and Sigma-2 receptors are typically obtained from published studies using cerebral tissue homogenates with [3H](+)pentazocine to measure displacement from Sigma-1 receptors and [3H] 1,3-Di-(2-tolyl)guanidine in the presence of 300 nM (+)pentazocine to measure displacement from Sigma-2 receptors.

Competitive Radioligand Binding Assay 2

[0249] The affinity of test compounds at sigma-1 and sigma-2 receptors was also determined by displacement of different known labeled sigma-2 or sigma-1 ligands. Filtration

assays were conducted according the previously published procedure (Xu, et al., 2005). Test compounds were dissolved in N,N-Dimethylformamide (DMF), dimethyl sulfoxide (DMSO) or ethanol and then diluted in 50 mM Tris-HCl pH 7.4 buffer containing 150 mM NaCl and 100 mM EDTA. Membrane homogenates were made from guinea pig brain for sigma-1 binding assay and rat liver for sigma-2 binding assay. Membrane homogenates were diluted with 50 mM Tris-HCl buffer, pH 8.0 and incubated at 25°C in a total volume of 150 uL in 96 well plates with the radioligand and test compounds with concentrations ranging from 0.1 nM to 10 uM. After incubation was completed, the reactions were terminated by the addition of 150 uL of ice-cold wash buffer (10 mM Tris HCl, 150 mM NaCl, pH 7.4) using a 96 channel transfer pipette (Fisher Scientific, Pittsburgh,PA) and the samples harvested and filtered rapidly through 96 well fiber glass filter plate (Millipore, Billerica, MA) that had been presoaked with 100 uL of 50 mM Tris-HCl buffer. Each filter was washed four times with 200 uL of ice-cold wash buffer (10 mM Tris-HCl, 150 mM NaCl, pH 7.4). A Wallac 1450 MicroBeta liquid scintillation counter (Perkin Elmer, Boston, MA) was used to quantitate the bound radioactivity.

[0250] The sigma-1 receptor binding assays were conducted using guinea pig brain membrane homogenates (~300 ug protein) and ~5 nM [3H](+)-pentazocine (34.9 Ci/mmol, Perkin Elmer, Boston, MA), incubation time was 90 min at room temperature. Nonspecific binding was determined from samples that contained 10 µM of cold haloperidol.

[0251] The sigma-2 receptor binding assays were conducted using rat liver membrane homogenates (~300 ug protein) and ~2 nM sigma-2 highly selective radioligand [3H]RHM-1 only (no other blockers) (America Radiolabeled Chemicals Inc. St. Louis, MO), ~10 nM [3H]DTG (58.1 Ci/mmol, Perkin Elmer, Boston, MA) or ~10 nM [3H]Haloperidol (America Radiolabeled Chemicals Inc., St. Louis, MO) in the presence of 1uM (+)-pentazocine to block sigma-1 sites, incubation times were 6 minutes for [3H]RHM-1, 120 min for [3H]DTG and [3H]haloperidol at room temperature. Nonspecific binding was determined from samples that contained 10uM of cold haloperidol.

[0252] Data from the competitive inhibition experiments were modeled using nonlinear regression analysis to determine the concentration of inhibitor that inhibits 50% of the specific binding of the radioligand (IC₅₀ value). The binding affinity, K_i values was calculated using the method of Cheng and Prusoff. The K_d value used for [3H](+)-pentazocine in guinea pig brain was 7.89 nM, for [3H]RHM-1 and [3H]DTG in rat liver were 0.66 nM and 30.73 nM respectively. The standard compound haloperidol was used for quality

assurance. Affinity data at the sigma-2 receptor for exemplary compounds are shown in Table 1.

[0253] In some embodiments, compounds according to any embodiment herein, or pharmaceutically acceptable salts thereof, exhibit sigma-2 receptor binding affinity K_i of not more than 1,000 nM, not more than 750 nM, not more than 500 nM, not more than 250 nM, not more than 100 nM, not more than 50 nM, not more than 25 nM, or not more than 10 nM, when tested according to a sigma-2 receptor binding assay protocol provided herein.

Example 5: Inhibition of Abeta Oligomer Effect on Neuronal Cells in Membrane Trafficking Assay

[0254] Compounds according to any embodiment described herein, were tested for their ability to inhibit an amyloid beta effect on the cells. The compounds generally were able to inhibit the amyloid beta effect as measured by a membrane trafficking /exocytosis assay (MTT assay). The results are indicated in Table 1. The rationale for this assay was as follows:

[0255] Since synaptic and memory deficits, and not widespread cell death, predominate at the earliest stages of Alzheimer's disease, assays that measure these changes are particularly well suited to discovering small molecule inhibitors of oligomer activity. The MTT assay is frequently used as a measure of toxicity in cultures. Yellow tetrazolium salts are endocytosed by cells and reduced to insoluble purple formazan in the endosomal pathway. The level of purple formazan is a reflection of the number of actively metabolizing cells in culture, and reduction in the amount of formazan is taken as a measure of cell death or metabolic toxicity in culture. When observed through a microscope, the purple formazan is first visible in intracellular vesicles that fill the cell. Over time, the vesicles are exocytosed and the formazan precipitates as needle-shaped crystals on the outer surface of the plasma membrane as the insoluble formazan is exposed to the aqueous media environment. Liu and Schubert ('97) discovered that cells respond to sublethal levels of Abeta oligomers by selectively accelerating the exocytosis rate of reduced formazan, while leaving endocytosis rate unaffected. The inventors have replicated these observations in mature primary neurons in culture and quantified these morphological shifts via automated microscopy and image processing. Under these circumstances, there is no overall change in the total amount of reduced formazan, simply a shift in its morphology reflective of changes in rate of its formation and/or expulsion from the cell. The inventors have confirmed previous findings that this assay is sensitive to low levels of oligomers that do not cause cell death (Liu and

Schubert '04, Hong et al., '07). Indeed, low amounts of oligomers that lead to inhibition of LTP do not lead to cell death (Tong et al., '04) and are not expected to change total amounts of formazan in culture (or in brain slices).

[0256] Evidence adduced by other investigators suggests that Abeta oligomer-mediated reduction in neuronal surface receptor expression mediated by membrane trafficking is the basis for oligomer inhibition of electrophysiological measures of synaptic plasticity (LTP) and thus learning and memory (Kamenetz et al., '03, Hsieh et al., '06). Measuring membrane trafficking rate changes induced by oligomers via formazan morphological shifts has been used in cell lines to discover Abeta oligomer-blocking drugs (Maezawa et al., '06, Liu and Schubert '97, '04, '06, Rana et al., '09, Hong et al., '08) that lower Abeta brain levels in rodents *in vivo* (Hong et al., '09). Similar procedures for exocytosis assays/MTT assays can be found in the literature. See e.g., Liu Y, et. al., Detecting bioactive amyloid beta peptide species in Alzheimer's disease. J Neurochem. 2004 Nov;91(3):648-56; Liu Y, and Schubert D. "Cytotoxic amyloid peptides inhibit cellular 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction by enhancing MTT formazan exocytosis." J Neurochem. 1997 Dec;69(6):2285-93; and Liu Y, and Schubert D. "Treating Alzheimer's disease by inactivating bioactive amyloid beta peptide" Curr. Alzheimer Res. 2006 Apr;3(2):129-35. Therefore the approach is valid.

[0257] The present exocytosis assay was adapted for use with mature primary neuronal cultures grown for 3 weeks *in vitro*. See WO 2011/106785, which is incorporated herein by reference in its entirety. Abeta oligomers cause a dose-dependent decrease in the amount of intracellular vesicles (puncta) filled with reduced purple formazan as measured via image processing using a Cellomics VTI automated microscopy system. Photomicrographs for a cultured neuronal cell exposed to vehicle alone show vesicles filled with formazan; wherein a photomicrograph of a neuronal cell exposed to vehicle plus Abeta oligomer shows considerably fewer vesicles filled with formazan and instead shows exocytosed formazan which, when encountering the extracellular environment, precipitates into crystals. Increasing the amount of Abeta oligomers eventually results in overt toxicity. Thus, the concentration of neuroactive Abeta oligomers used in the assay is much lower than that causing cell death. The inventors confirmed that the assay is operative by showing that the effects of Abeta oligomer are blocked upon addition of anti-Abeta antibody but antibody alone has no effect on its own (data not shown). When configured in this manner, the assay is able to detect compounds that inhibit nonlethal effects of Abeta oligomer whether these compounds act via

disruption of oligomers, inhibition of oligomer binding to neurons, or counteraction of signal transduction mechanisms of action initiated by oligomer binding.

[0258] The methods used to generate the results were as follows in the Membrane Trafficking/Exocytosis (MTT) assay.

[0259] Primary hippocampal neurons from E18 Sprague-Dawley rat embryos were plated at optimized concentrations in 384 well plates in NB media (Invitrogen). Neurons were maintained in cultures for 3 weeks, with twice weekly feeding of NB media with N2 supplement (Invitrogen). These neurons express the full complement of synaptic proteins characteristic of neurons in the mature brain, and exhibit a complex network of activity-dependent electrical signaling. Neurons and glia in such cultures have molecular signaling networks exhibiting excellent registration with intact brain circuitry, and for this reason have been used for over two decades as a model system for learning and memory (See e.g. Kaech S, Banker G. Culturing hippocampal neurons. Nat Protoc. 2006;1(5):2406-15. Epub 2007 Jan 11; See also Craig AM, Graf ER, Linhoff MW. How to build a central synapse: clues from cell culture. Trends Neurosci. 2006 Jan;29(1):8-20. Epub 2005 Dec 7. Review).

[0260] A test compound was added to cells at concentrations ranging from 100uM to 0.001 nM followed by addition of vehicle or Abeta oligomer preparations (3 μ M total Abeta protein concentration), and incubated for 1 to 24 hr at 37 °C in 5% CO₂. MTT reagent (3-(4,5-dimethylthiazol-2-yl)-2,5diphenyl tetrazolium bromide) (Roche Molecular Biochemicals) was reconstituted in phosphate buffered saline to 5mg/mL. 10 μ L of MTT labeling reagent is added to each well and incubated at 37 °C for 1h, then imaged. Exocytosis was assessed by automated microscopy and image processing to quantify the amount of endocytosed and exocytosed formazan.

[0261] Each assay plate was formatted so that compounds are tested with and without Abeta oligomer on each plate. This design eliminates toxic or metabolically active compounds early on in the screening cascade (at the level of the primary screen). Reduced formazan was first visible in intracellular vesicles. Eventual formazan exocytosis was accelerated via Abeta oligomers.

[0262] In the presence of an effective concentration of active test compound, the membrane traffic changes are blocked and the cell is indistinguishable from a vehicle-treated neuron. Furthermore, in some cases this effect of test compound appears to be independent of whether test compound is added before or after exposure of the cells to Abeta oligomer, which indicates a therapeutic as well as a prophylactic effect. Adequate concentration of active test compound blocks membrane trafficking effects of Abeta oligomer seen in this

assay. Ascending doses of compounds according to any embodiment described herein, that are selective, high affinity agonists of the sigma-2 receptor, stop oligomer effects and make the cultures look more like vehicle-treated cultures.

[0263] Compounds according to any embodiment described herein, that are selective, high affinity agonists of the sigma-2 receptor that are effective for inhibiting Abeta oligomer toxicity are promising as therapeutic and prophylactic modalities for amyloid beta oligomer toxicity related cognitive decline such as that seen in Alzheimer's disease.

[0264] Synthetic Abeta oligomers were dosed in the membrane trafficking assay, where it exhibited an EC₅₀ of 820nM. Each concentration of Abeta was tested against several concentrations of each test compound. Active compounds caused a rightward shift in the EC₅₀ by almost two orders of magnitude. When the data were fitted to classical linear and non linear models, the data were linear with a Schild analysis (Hill slope nH of 1), which indicates that the compounds exhibit true pharmacological competition between oligomers and compound for targets that mediate membrane trafficking.

[0265] Abeta oligomers derived from Alzheimer's patient's brains can be dosed against test compounds, and a rightward shift is also expected to be exhibited by compound exposure. Specifically, at effective doses, the active test compounds exhibit pharmacological competition with both synthetic and human Alzheimer's patient-derived oligomers. Experimental controls

[0266] Abeta 1-42 oligomers made according to published methods were used as positive controls. [See e.g. Dahlgren et al., "Oligomeric and fibrillar species of amyloid-beta peptides differentially affect neuronal viability" J Biol Chem. 2002 Aug 30;277(35):32046-53. Epub 2002 Jun 10.; LeVine H 3rd. "Alzheimer's beta-peptide oligomer formation at physiologic concentrations" Anal Biochem. 2004 Dec 1;335(1):81-90; Shrestha et.al, "Amyloid beta peptide adversely affects spine number and motility in hippocampal neurons" Mol Cell Neurosci. 2006 Nov;33(3):274-82. Epub 2006 Sep 8; Puzzo et al., "Amyloid-beta peptide inhibits activation of the nitric oxide/cGMP/cAMP-responsive element-binding protein pathway during hippocampal synaptic plasticity" J Neurosci. 2005 Jul 20;25(29):6887-97; Barghorn et al., "Globular amyloid beta-peptide oligomer - a homogenous and stable neuropathological protein in Alzheimer's disease" J Neurochem. 2005 Nov;95(3):834-47. Epub 2005 Aug 31; Johansson et al., Physiochemical characterization of the Alzheimer's disease-related peptides A beta 1-42 Arctic and A beta 1-42wt. FEBS J. 2006 Jun;273(12):2618-30] as well as brain-derived Abeta oligomers (See e.g. Walsh et al., Naturally secreted oligomers of amyloid beta protein potently inhibit

hippocampal long-term potentiation *in vivo*. Nature (2002). 416, 535-539; Lesne et al., A specific amyloid-beta protein assembly in the brain impairs memory. Nature. 2006 Mar 16;440(7082):352-7; Shankar et al, Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. Nat Med. 2008 Aug;14(8):837-42. Epub 2008 Jun 22). It should be noted that any Abeta oligomer preparation can be used in this assay or as a control, including preparations described in the patent literature, cited above and incorporated by reference in their entirety.

[0267] Various different Abeta oligomer preparations were demonstrated to cause an Abeta effect in the membrane trafficking assay, including notably oligomer preparations isolated from the brain of Alzheimer's disease patients.

[0268] Oligomers were isolated from postmortem human hippocampus or prefrontal cortex without the use of detergents and inhibited membrane trafficking in a dose-dependent manner with a Kd of 6 pMolar. Human Alzheimer's disease patient-derived Abeta oligomers (137 pM) produce a statistically significant inhibition of membrane trafficking compared to vehicle. Compound 4-(3-(4-(trifluoromethyl)benzylamino)butyl)-2-methoxyphenol eliminates the membrane trafficking deficits induced by AD brain-derived Abeta oligomers, but does not affect trafficking when dosed in the absence of Abeta.

[0269] Although potencies of various Abeta oligomer preparations differ (for example native Alzheimer's isolates are more potent than any of the synthetic preparations tested—data not shown), the results are qualitatively the same: pathologies mediated by oligomers are countered by compounds according to any embodiment herein that act as sigma-2 functional antagonists.

Primary neuronal cultures

[0270] Optimal cell density is determined based on cellular response to Abeta oligomers using the exocytosis assay as a readout, and immunohistochemical analysis of the relative proportion of glia to neurons in the cultures. Cultures are monitored on a weekly basis with immunohistochemistry and image processing-based quantification to monitor the percentage of the cultures that are neurons vs. glia (Glial cells). Cultures containing more than 20% glia (positive for GFAP) vs. neurons (staining positively with (chicken polyclonal) antibodies (Millipore) directed against MAP2 at 1:5000 (concentration variable)) at the screening age of 21 days *in vitro* (21 DIV) are rejected.

Abeta Oligomer preparations

[0271] Human amyloid peptide 1-42 was obtained from a number of commercial vendors such as California Peptide, with lot-choice contingent upon quality control analysis. Quality controls of oligomer preparations consist of Westerns to determine oligomer size ranges and relative concentrations, and the MTT assay to confirm exocytosis acceleration without toxicity. Toxicity was monitored in each image-based assay via quantification of nuclear morphology visualized with the DNA binding blue dye DAPI (Invitrogen). Nuclei that are fragmented are considered to be in late stage apoptosis (Majno and Joris '95) and the test would be rejected. Peptide lots producing unusual peptide size ranges or significant toxicity at a standard 1.5 μ M concentration on neurons would also be rejected.

[0272] Plate-based controls –The assay optimization was considered complete when reformatted plates achieve a minimum of statistically significant two-fold separation between vehicle and Abeta oligomer-treated neurons ($p < 0.01$, Student's t-test, unequal variance) on a routine basis, with no more than 10% CV between plates.

Statistical software and Analysis

[0273] Data handling and analysis were accomplished by Cellomics VTI image analysis software and STORE automated database software. Because of the low dynamic range and neuronal well-to-well variability after three weeks in culture, statistical comparisons are made via pairwise Tukey-Kramer analysis to determine the significance of the separation between compound + Abeta oligomers from Abeta alone, and between compound alone from vehicle. The ability of mature primary neurons to more closely approximate the electrophysiologically mediated signal transduction network of the adult brain justifies this screening strategy. Power analysis was set for a number of replicate screening wells that minimized false negatives (e.g. $N=4$). Test compounds of the disclosure significantly reverse the effects of Abeta oligomers on membrane trafficking but do not affect neuronal metabolism themselves.

[0274] Selected compounds according to any embodiment described herein, were dosed in the MTT assay described herein, prior to Abeta oligomer addition and were shown to block the Abeta oligomer-induced membrane trafficking deficits with the indicated EC_{50} . Specifically, these results indicate that compounds block/abate the activity/effect of Abeta oligomer on membrane trafficking of neuron cells at micromolar concentrations. Certain compounds in Table 1 were shown to block the Abeta oligomer-induced acceleration of exocytosis with the indicated EC_{50} . Accordingly, the compounds in Table 1 significantly

blocked Abeta oligomer-mediated changes in membrane trafficking. These results indicate that compounds block/abate the activity/effect of Abeta oligomer on neuron cells and that compounds according to any embodiment described herein, can be used to block the Abeta oligomer induced membrane trafficking abnormalities.

[0275] In some embodiments, compounds according to any embodiment described herein, inhibit Abeta oligomer-induced membrane trafficking deficits, with an EC₅₀ of not more than 20 uM, not more than 15 uM, not more than 10 uM, not more than 5 uM, not more than 1 uM, not more than 0.5 uM, when tested according to the membrane trafficking assay protocol provided herein.

Example 6: Pharmacokinetic and Metabolic Stability Studies.

[0276] A first pharmacokinetic study was performed in microsomes of mice mouse liver microsomes, MLM). The studies were performed according to Obach, R.S et al.(1997) J. Pharmacol. Exp. Ther., 283: 46-58, which is incorporated herein by reference. The half-life (t_{1/2}) of the compounds in MLM assay is shown in Table 1.

[0277] In some embodiments, a compound according to any embodiment described herein, or pharmaceutically acceptable salts thereof, exhibits a half-life (t_{1/2}) in a mouse liver microsome (MLM) assay, as provided herein, of at least 5 minutes, at least 10 minutes, at least 25 minutes, at least 50 minutes, at least 100 minutes, or at least 200 minutes.

[0278] The results indicate that several of the compounds tested had a substantially longer half-life in mouse liver microsomes. This result portends greater bioavailability after oral administration for these compounds. The same compounds have been tested by the membrane trafficking assay described above and their activity as referred to herein.

[0279] If the rate of intrinsic clearance of test compound was rapid, it is suggestive of substantial first pass metabolism. In order to improve pharmacokinetic properties, compounds were designed to enhance metabolic stability and improve drug-like properties. Microsomal stability experiments and plasma stability experiments were performed to determine metabolic and hepatic stability of candidate compounds. In some embodiments, *in vitro* microsomal stability was normalized to standard compound, 4-(3-(4-(trifluoromethyl)benzylamino)butyl)-2-methoxyphenol, an early lead compound that had a mouse liver microsome t_{1/2} (min) of 16. Compounds of the invention are superior to the this early lead compound..

[0280] A second PK study can be conducted *in vivo* and involves measuring plasma levels and brain levels for test compounds administered by various routes and in an acute or chronic manner, as follows:

[0281] HPLC-MS Optimization

[0282] A solution of each test compound is prepared and infused into the TSQ Quantum spectrometer (Fisher Thermo Scientific) source via syringe pump at a constant rate. Full scan MS (mass spectroscopy) analysis is conducted and total ion current chromatograms and corresponding mass spectra are generated for each test compound in both positive and negative ionization modes. The precursor ions for MS/MS are selected from either the positive or the negative mass spectrum, as a function of the respective ion abundance. In addition, product ion MS/MS analysis is performed in order to determine the appropriate selected fragmentation reaction for use in quantitative analysis. The final reaction monitoring parameters are chosen to maximize the ability to quantify the test compound when present within a complex mixture of components. Following identification of the specific SRM transition to be used for each test compound, the detection parameters are optimized using the automated protocol in the TSQ Quantum Compound Optimization workspace. Finally, the chromatographic conditions to be used for LC-MS analysis are identified by injection and separation of the analyte on a suitable LC column and adjustment of the gradient conditions is performed as necessary.

[0283] Formulation for IV dosing:

[0284] The solubility of the test compound in phosphate-buffered saline, pH 7.4 (PBS) is first evaluated by visual inspection. PBS is used as the vehicle if the compound is soluble at the target concentration. (Other vehicles that are compatible with IV dosing may be evaluated if the compound is not completely soluble in PBS. Such vehicles include DMSO, polyethylene glycol (PEG 400), Solutol HS 15, and Cremophor EL among others.) In the experiments reported here a single bolus, 10 mg/kg, of test compound is administered IV.

[0285] Formulation for PO dosing: The solubility of the test compound in PBS is first evaluated. PBS is used as the vehicle if the compound is soluble at the target concentration. (DMSO/Solutol HS 15/PBS (5/5/90, v/v/v), or DMSO/1 % methylcellulose (5/95, v/v) may be used if the test compound is not completely soluble in PBS at the respective concentration.)

[0286] Linearity in Plasma

[0287] Aliquots of plasma are spiked with the test compounds at the specified concentrations. The spiked samples are processed using acetonitrile precipitation and analyzed by HPLC-MS or HPLC-MS/MS. A calibration curve of peak area versus concentration is constructed. The reportable linear range of the assay is determined, along with the lower limit of quantitation (LLQ).

[0288] Quantitative Bioanalysis of Plasma Samples

[0289] The plasma samples are processed using acetonitrile precipitation and analyzed by HPLC-MS or HPLC-MS/MS. A plasma calibration curve was generated. Aliquots of drug-free plasma are spiked with the test compound at the specified concentration levels. The spiked plasma samples are processed together with the unknown plasma samples using the same procedure. The processed plasma samples (dried extracts) are typically stored frozen (-20 °C) until the HPLC-MS or HPLC-MS/MS analysis. The dried extracts are reconstituted into a suitable solvent and after centrifugation were analyzed by HPLC-MS or HPLC-MS/MS. Peak areas are recorded, and the concentrations of the test compound in the unknown plasma samples are determined using the respective calibration curve. The reportable linear range of the assay is determined, along with the lower limit of quantitation (LLQ).

[0290] Animals used in the study are typically male C57BL/6 mice weighing 20-30 g each or male Sprague-Dawley rats weighing 180-250 g. Three animals are treated for each administration condition and each time point, so that each animal is subjected to only one blood draw. Subcutaneous compound administration was accomplished by intraperitoneal injection. Per oral administration is accomplished by gastric gavage. Intravenous administration is accomplished via jugular catheter.

[0291] Following compound administration at various concentrations, plasma samples are collected at, e.g., 10, 30, 60, 120, 240, 360, 480 and 1440 min.

[0292] Plasma Sample Collection from Mice and Rats

[0293] Animals are sedated under general inhalant anesthesia (3 % isoflurane) for blood collection by cardiac puncture (mice) or jugular catheter (rats). Blood aliquots (300-400 µL) are collected in tubes coated with lithium heparin, mixed gently, and are kept on ice and centrifuged at 2,500 ×g for 15 minutes at 4°C, within 1 hour of collection. The plasma is then harvested and kept frozen at -20 °C until further processing.

[0294] Animal Dosing Design - *In vivo* PK - Non cannulated, nonfasted animals

[0295] Group 1: SC, n=3 animals per time point (24 animals total) or

[0296] IV, n=3 animals per time point (24 animals total)

[0297] Group 2: PO, n=3 animals per time point (24 animals total)

[0298] Group 3: Control animals (for drug-free blood), n=5 mice

[0299] Each animal is subject to one blood draw and one brain collection.

[0300] Brain sample collection from animals

[0301] Immediately after blood sampling, animals are decapitated and the whole brains are quickly removed, rinsed with cold saline (0.9% NaCl, g/mL), surface vasculature ruptured, blotted dry with gauze, weighted, kept on ice until further processing within one hour of collection. Each brain is homogenized in 1.5 mL cold phosphate buffered saline, pH 7.4 (mice =1.5 mL, rats =), for 10 seconds on ice using the Power Gen 125. The brain homogenate from each brain is then stored at -20 °C until further processing.

[0302] Linearity in Brain samples

[0303] Aliquots of brain homogenate are spiked with the test compound at the specified concentrations. To each brain aliquot an equal volume of chilled 26% (g/mL) neutral Dextran (average molecular Weight 65,000-85,000 from Sigma, catalog number D-1390) solution is added to obtain a final Dextran concentration of 13%. The homogenate is centrifuged at 54000 x g for 15 minutes at 4°C. The supernatants are subsequently processed using acetonitrile precipitation and analyzed by HPLC-MS/MS. A calibration curve of peak versus concentration is constructed. The reportable linear range of the assay is determined, along with the lower limit of quantitation (LLQ).

[0304] Quantitative analysis of Brain Samples

[0305] To each brain homogenate aliquot an equal volume of chilled 26% (g/mL) neutral Dextran (average molecular Weight 65,000-85,000 from Sigma, catalog number D-1390) solution is added to obtain a final Dextran concentration of 13%. The homogenate is centrifuged at 54000 x g for 15 minutes at 4°C. The supernatants are subsequently processed using acetonitrile precipitation and analyzed by HPLC-MS/MS. A brain calibration curve is generated. Aliquots of drug-free brain homogenate are spiked with the test compound at specified concentration levels. The spiked brain homogenate samples are processed together with the unknown brain homogenate samples using the same procedure. The processed brain samples are stored at -20°C until the LC-MS/MS analysis, at which time peak areas were recorded, and the concentrations of test compound in the unknown brain samples were determined using the respective calibration curve. The reportable linear range of the assay was determined along with the lower limit of quantitation (LLQ).

[0306] Brain penetrability

[0307] The concentrations of test compound in brain (ng/g tissue) and in plasma (ng/mL) as well as the ratio of the brain concentration and the plasma concentration at each time point are determined by LC-MS/MS and reported as described above.

[0308] Pharmacokinetics

[0309] Plots of plasma concentration of compound versus time are constructed. The fundamental pharmacokinetic parameters of compound after oral and SC dosing (AUC_{last}, AUC_{INF}, T_{1/2}, T_{max}, and C_{max}) are obtained from the non-compartmental analysis (NCA) of the plasma data using WinNonlin (Pharsight). Noncompartmental analysis does not require the assumption of a specific compartmental model for either drug or metabolite. NCA allows the application of the trapezoidal rule for measurements of the area under a plasma concentration-time curve (Gabrielsson, J. and Weiner, D. Pharmacokinetic and Pharmacodynamic Data Analysis: Concepts and Applications. Swedish Pharmaceutical Press. 1997).

[0310] Definitions of Terms Reported

[0311] Area Under the Curve (AUC) - Measure of the total amount of unchanged drug that reaches the systemic circulation. The area under the curve is a geometric measurement that was calculated by plotting concentration versus time and summing the incremental areas of each trapezoid.

[0312] WinNonlin has two computational methods for calculation of the area: the linear trapezoidal method and the linear-log trapezoidal method. Because the linear trapezoidal method may give biased results on the descending part of the concentration-time curve and overestimate the AUC, WinNonlin provides the linear-log option for calculation of AUC. By default, the log-linear trapezoidal method is used to measure the post-T_{max} area for the remainder of the plasma concentration-time curve.

[0313] AUC_{last}: area under the curve from the time of dosing to the time of last observation that was greater than the limit of quantitation.

[0314] AUC_{INF}: Area under the curve from the time of dosing extrapolated to infinity.

[0315] C_{max} - Maximum plasma drug concentration obtained after oral or non-IV administration of a drug between the time of doing and the final observed time point.

[0316] T_{max} - Time at maximum observed plasma concentration (C_{max}) noted in minutes after administration of drug.

[0317] T_{1/2} - Terminal elimination half-life from both IV and non-IV dosing.

[0318] where λ (z) is the first order rate constant associated with the terminal (log-linear) portion of the plasma concentration-time curve. z is estimated by linear regression of time versus log concentration.

[0319] The results are expected to show that certain test compounds exhibit good bioavailability and good brain penetrability when administered at doses ranging from 0.1 to 0.5 mg/kg acutely or chronically (daily over 5 days). Selected test compounds are evaluated for oral bioavailability in this manner.

Example 8: *In vitro* testing for hERG inhibition

[0320] *In vitro* testing for hERG inhibition was performed in a standard assay (See: Haverkamp W, Breithardt G, Camm AJ, Janse MJ, Rosen MR, Antzelevitch C, Escande D, Franz M, Malik M, Moss A and Shah R. (2000) *Eur Heart J* 21 (15); 1216-31). Results for test compounds for hERG inhibition (IC_{50} , nM) is shown in Table 1. In some embodiments, compounds according to any embodiment described herein, or pharmaceutically acceptable salts thereof, exhibit minimal hERG inhibition, with an IC_{50} of greater than 300 nM, greater than 500 nM, greater than 1,000 nM, greater than 3,000 nM, greater than 5,000 nM, greater than 10,000, or greater than 20,000 nM. In particular embodiments, compounds according to any embodiment described herein, or pharmaceutically acceptable salts thereof, exhibit minimal hERG inhibition, and exhibit an IC_{50} of greater than 5,000 nM, greater than 10,000, or greater than 20,000 nM.

[0321] Combined Results for particular compounds described herein, with respect to log P, membrane trafficking (uM), sigma-2 receptor affinity, sigma-1 receptor affinity microsomal stability in mouse liver microsomes (MLM) ($t_{1/2}$, min), $t_{1/2}$ normalized to CT)10914 and *in vitro* toxicity potassium channel hERG (IC_{50} , nM) are provided in Table 1:

Example Compound	MW g/mol	LC-MS [MH] ⁺	log P	EC ₅₀ Abeta (uM)	S1 K _i (nM)	S2 K _i (nM)	MLM T _{1/2} (min.)	T _{1/2} *	hERG IC ₅₀ (nM)
1	383.5	384.4	5.16		0.7	0.2			740
2	357.5	358.5	4.67	> 15	0.9	0.6			9200
3	383.6	384.5	6.22	0.25		0.76			
4	371.5	372.2	5.35		0.5	0.8			2200
5	323.9	324.7	4.4	> 15		0.8			75
6	343.4	344.4	4.49	> 15	0.3	1			750
7	343.4	344.6	4.31	> 15	0.7	1.5			1500
8	281.8	282.9	3.16	0.25	0.4	1.6			14000
9	437.6	438.5	6.58	0.5	13	1.8	114	166	1000
10	295.8	296.9	3.8	1.26	0.7	2.1			11000
11	404.0	405.2	6.31	3	14, 69	2.2	26,	87	1500
12	343.4	344.1	4.13		1	2.7			970
13	329.4	330.4	3.85		1	2.9			2500
14	329.4	330.2	3.71	3.7	1.9	3			18000
15	387.6	388.7	5.85	1.6	40	4.5	14	20	2200

Example Compound	MW g/mol	LC-MS [MH] ⁺	log P	EC ₅₀ Abeta (uM)	S1 K _i (nM)	S2 K _i (nM)	MLM T _{1/2} (min.)	T _{1/2} *	hERG IC ₅₀ (nM)
16	309.9	310.9	4.03		0.8	4.5			840
17	315.4	316.4	3.43	2.24	1.3	5.1			11000
18	293.4	294.4	2.72	0.25	2.6	5.6			29000
19	293.4	294.5	3.57	< 0.25	2.2	6			37000
20	295.8	296.9	3.57		1	6.1			1700
21	359.5	360.4	5.26	20	8	6.2	11	37	830
22	309.9	310.8	3.85		1	7			1700
23	392.0	393.2	6.49	0.95	140	9.2	8.2	25	2200
24	397.0	398.1	6.31	20		10			
25	279.4	280.5	3.11		2.3	10			4000
26	403.5	404.6	3.35	20		11			
27	376.6	377.7	6.22	0.4		15			
28	397.5	398.2	5.68	4	6.1	18	60	60	10200
29	392.6	393.7	5.55	20		19			
30	381.5	382.7	6.97	2.2	4.1	19	83	78	6500
31	348.0	349.2	6.7	0.7	41	28	21	20	10900
32	474.1	475.3	6.37	6.9		32			
33	341.6	342.6	7.04	1.92		33			
34	423.6	424.7	6.15	20	5.3	39	60	60	23000
35	383.5	384.5	5.31	20	56	39	60	60	13400
36	373.5	374.6	5.41	20	12	40	25	129	10000
37	466.6	467.8	6.18	0.2	8.8	40	38	36	26200
38	369.5	370.5	4.85	4.5	190	42	84	79	16200
39	383.5	384.6	5.49	0.6	25	43			8200
40	455.6	456.9	6.28	0.83	170	44			9600
41	390.0	391.2	5.87	1.6	16	67	35	60	4000
42	349.9	350.7	5.04	20	150	71	88	83	6900
43	417.1	418.3	6.37	0.3	61	74	29	88	4900
44	411.5	412.3	6.27	1.4	260	79			9200
45	356.6	357.7	6.26	2.6		85			
46	474.1	475.6	6.37	4.8		86			
47	404.6	405.4	5.55		1913	90			14000
48	349.9	350.7	5.3	1.8		93			
49	404.6	405.7	5.74		575	94			2400
50	313.5	314.7	6.31	1.7		96			
51	440.6	441.4	4.55	20		100			
52	327.5	328.6	6.9	4.8		110			
53	364.0	365.2	5.41	20	74	110	28	26	13300
54	335.5	336.3	7.02	3.2		120			
55	337.9	338.9	5.47	2.5	25	120	3.6	11	7200
56	362.5	362.8	4.49		571	139			1900
57	329.5	330.4	5.21	0.73		150			
58	347.5	348.6	4.94	0.5	64	160	39	79	14000
59	347.5	348.5	5.12	0.25	600	160			73000
60	376.6	377.8	6.91	1.0		210			
61	353.5	354.2	6.38	4.7	160	230	53	77	7300
62	333.5	334.2	4.4	0.8	130	230	60	122	41000
63	466.1	467.2	7.42	1.5		240			
64	318.4	319.6	3.52		535	268			11000
65	335.5	336.7	4.13		1200	270			2300
66	333.5	334.7	4.84	20	39	290	8.8	45	16000
67	482.1	483.2	6.8	13		300			
68	303.5	304.4	5.65	20	500	300	11	57	27000
69	335.5	336.7	6.93	4.2		320			
70	421.5	422.4	6.95	20	33	340	14	13	16300

Example Compound	MW g/mol	LC-MS [MH] ⁺	log P	EC ₅₀ Abeta (uM)	S1 K _i (nM)	S2 K _i (nM)	MLM T _{1/2} (min.)	T _{1/2} *	hERG IC ₅₀ (nM)
71	396.5	397.8	4.61		62	340			17000
72	355.4	356.6	5.32	20	770	400	2.1	11	80000
73	349.9	350.4	4.86	0.7	29	410	87	176	24000
74	345.5	346.5	4.54	12.9		430			
75	362.9	363.8	4.33	0.25	690	690			15000
76	371.5	371.7	6.21	0.3	370	790	3.5	7	18000
77	335.9	336.8	4.58	20	540	800	44	41	15200
78	335.5	336.7	6.93	0.56		940			
79	319.5	320.2	4.12	1.9	1200	940	74	150	29000
80	335.5	336.5	6.93	0.64		1300			
81	346.5	347.6	3.87		160	2500			32000
82	433.6	434.6	4.11	20	380	2800	60	60	100000
83	369.5	370.4	5.31	3.8	1000	1000			14000
84	309.9	310.5	4.67	0.25	1000	1000			3900
85	343.4	344.6	4.95	0.98	1000	1000			11000
86	357.6	358.6	6.37	0.22					
87	362.0	363.3	7.13	20					
88	393.6	394.5	3.54						
89	405.6	406.6	5.37						
90	440.1	441.3	7.02	20					
91	277.4	278.7	3.99	20					
92	275.4	276.4	5.06	20					
93	273.4	274.5	4.36	20			3.4	9	
94	271.4	272.6	5.43	20			1.8	5	
95	293.8	294.6	4.45	20			1.5	4	
96	291.9	292.3	5.52	1.6			6.2	16	
97	369.6	370.7	5.78	10			8.7	15	
98	385.6	386.4	5.11	20			1.7	4	
99	331.5	332.3	4.08	20			7.4	19	
100	315.5	316.9	4.75	13.6			4.5	12	
101	343.5	344.7	5.57	9.6			1.8	5	
102	359.5	360.5	4.9	20			1.5	4	
103	287.4	288.3	4.76	20			1.1	2	
104	289.4	290.4	3.69	20			1.3	2	
105	376.0	377.3	5.72	20			1.3	2	
106	373.6	374.7	5.35	20			3.9	7	
107	341.5	342.4	5.21	20			3.5	6	
108	345.5	344.5	4.46	20			7.3	12	
109	329.5	330.8	5.13	20			5.4	9	
110	387.6	388.5	5.73	20			2.6	4	
111	364.0	365.2	5.68	20			6.8	23	
112	343.5	344.4	5.59				4.9	16	
113	359.5	360.7	4.92	20			8.2	27	
114	355.6	356.8	5.63				1.8	6	
115	367.5	368.4	4.9	20					
116	337.5	338.7	5.71						
117	341.9	342.6	6.47						
118	339.5	340.9	4.64						
119	426.0	427.0	6.67						
120	343.9	344.6	5.4						
121	371.9	372.9	5.66						
122	421.6	422.4	5.91						
123	387.9	388.8	6.67	0.3					
124	395.6	396.7	6.45	0.2					

Example Compound	MW g/mol	LC-MS [MH] ⁺	log P	EC ₅₀ Abeta (uM)	S1 K _i (nM)	S2 K _i (nM)	MLM T _{1/2} (min.)	T _{1/2} *	hERG IC ₅₀ (nM)
125	372.6	373.4	5.23	0.25					
126	388.6	389.9	4.61	0.3					7400
127	436.6	437.7	6.41	0.3					3300
128	445.6	446.7	7.19	20					
129	383.5	384.4	5.13	1.3					24000
130	422.6	423.6	5.97	0.3					
131	331.5	332.8	6.24	20					
132	412.0	413.2	6.91	20					
133	386.6	387.6	5.68	2.2					
134	321.4	322.2	5.01	20					
135	389.0	390.2	5.69	20					
136	364.0	365.0	5.58	1.5					2100
137	438.6	439.8	5.34	0.15					
138	405.0	406.1	5.07	0.45					
139	375.6	376.9	6.03	20					
140	442.7	443.9	5.78	20					
141	371.5	372.5	5.74	1.6					
142	349.9	350.7	5.22	20					
143	438.6	439.6	5.96	20					
144	414.6	415.4	4.69	20					
145	333.5	334.9	4.76	20					
146	403.0	404.1	6.14	0.025					
147	440.7	441.8	5.16						
148	400.6	401.7	4.14						
149	454.7	455.8	5.59	20					
150	386.6	387.8	3.86	20					
151	372.5	373.3	4.65	20					
152	389.0	390.2	5.11	0.26					
153	408.6	409.7	5.73	0.5					
154	382.5	383.2	3.37						
155	458.6	459.6	6.46						
156	399.0	400.0	3.83	0.34					
157	425.0	426.3	6.19						
158	378.0	379.1	5.99	2.88					
159	389.6	390.4	6.36	10					
160	406.0	407.2	6.82	15					
161	432.5	433.5	4.1	0.27					
162	422.0	423.2	6	15					
163	405.6	406.3	5.54	15					
164	439.6	440.3	7.1						
165	351.9	352.9	4.6						
166	317.5	318.7	3.99						
167	321.9	322.9	5.1						
168	427.7	428.8	6.17						
169	387.6	388.3	5.15						
170	373.6	374.3	4.87						
171	415.7	416.5	6.5						
172	277.4	278.4	3.52						
173	275.4	276.4	4.81						
174	295.4	296.3	3.67						
175	311.8	312.8	4.13						
176	293.4	294.2	4.96						
177	309.9	310.7	5.42						
178	359.5	360.5	4.59						
179	417.6	418.6	5.49		430				4000

Example Compound	MW g/mol	LC-MS [MH] ⁺	log P	EC ₅₀ Abeta (uM)	S1 K _i (nM)	S2 K _i (nM)	MLM T _{1/2} (min.)	T _{1/2} *	hERG IC ₅₀ (nM)
180	373.6	374.2	4.87						20000
181	434.1	435.6	5.95						1800
182	467.6	468.6	6.22		160				2600
183	429.7	430.7	6.79						2100
184	275.4	276.2	2.57		220				62000
185	407.6	408.4	6.16						3000
186	309.8	310.9	3.18		42				2300
187	424.0	425.2	6.62						1000
188	337.9	338.9	4						400
189	457.6	458.5	6.89		37				5700
190	349.5	350.6	4.63						
191	329.4	330.2	4.07						
192	295.8	296.8	3.44						19000
193	349.9	350.7	4.89						
194	421.0	422.1	6.01						
195	411.6	412.7	3.36						
196	376.5	377.4	4.85						
197	378.9	379.7	4.95						
198	421.0	422.0	6.2						4800
199	309.9	310.8	4.22						1200
200	454.6	455.4	6.28						83000
201	301.5	302.5	3.97	15					15000
202	315.5	316.6	4.62						
203	332.5	333.6	3.75						
204	369.6	370.5	5.7						
205	445.6	446.7	6.5	0.59					3900
206	368.4	369.5	4.26						
207	461.6	462.7	4.09						
208	426.5	427.3	5.58						
209	454.6	455.8	6.47						
210	427.5	428.6	4.79	0.42					4400
211	412.0	413.0	6.23						
212	425.6	426.8	6.54	1.8					27000
213	315.5	316.7	4.25	6.6					6400
214	392.0	393.1	6.27						
215	418.6	419.6	5.5						
216	510.7	511.9	7.65						
217	394.0	395.0	4.51	4.6					7700
218	377.5	378.7	4.05						13000
219	359.5	360.3	3.91						
220	377.6	378.2	5.62						
221	395.6	396.7	5.76						
222	375.6	376.5	5.81						
223	357.6	357.9	5.67						
224	334.9	335.6	3.98						
225	477.1	478.1	7.37						
226	460.7	461.6	6.91	1.25					
227	300.4	301.3	3.38						
228	468.6	469.8	6.24						
229	295.8	296.8	4.03	15					21000
230	321.9	323.0	4.52	10					16000
231	295.8	296.6	4.03						11000
232	355.4	356.4	4.79	15					29000
233	315.4	316.9	3.79	0.26					24000
234	349.9	350.7	5.68						


Example Compound	MW g/mol	LC-MS [MH] ⁺	log P	EC ₅₀ Abeta (uM)	S1 K _i (nM)	S2 K _i (nM)	MLM T _{1/2} (min.)	T _{1/2} *	hERG IC ₅₀ (nM)
235	349.9	350.9	5.68						
236	369.5	370.5	5.31	15					20000
237	335.9	336.7	5.04						
238	309.9	310.7	4.67	15					13000
239	335.9	336.5	5.04						4600
240	335.9	336.9	5.04						
241	319.5	320.7	4.57						
242	319.5	320.3	4.57						14000
243	369.5	370.7	5.31						
244	333.5	334.6	5.22	8.5					4800
245	333.5	334.5	5.22	> 15					14000
246	349.9	350.9	5.68						
247	383.5	384.3	5.95						
248	392.0	393.0	5.84						
249	315.5	316.7	5.07	15					9400
250	281.8	282.3	3.51	15					28000
251	333.5	334.6	5.22						
252	315.5	316.7	5.07	15					7000
253	309.9	310.9	4.67	0.25					
254	343.4	344.4	4.95						5000
255	281.8	282.9	3.51	15					
256	543.7	544.3	5.01	15					
257	507.6	508.7	4.08						
258	508.7	509.6	2.62						
259	570.7	571.5	4.03						
260	520.7	521.7	2.46						
261	437.6	438.9	6.58	0.73					
262	437.6	438.7	6.58	1.85					
263	455.6	456.9	6.28						
264	455.6	456.4	6.28						
265	383.5	384.7	5.13						
266	383.5	384.6	5.13						
267	375.5	376.5	5.37						
268	351.9	352.9	4.83						
269	494.6	495.4	2.2						
270	480.6	481.5	1.81						
271	598.8	599.7	4.85						
272	481.6	482.6	3.35						
273	522.7	523.7	3.19						
274	534.7	535.9	3.24						
275	383.5	384.1	5.95						
276	343.4	344.6	4.95						
277	385.5	386.5	5.11						
278	357.5	358.5	5.23						
279	522.6	523.6	1.63						
280	612.8	613.9	3.85						
281	495.6	496.4	2.48						
282	562.7	563.9	2.47						
283	295.8	296.4	3.62						
284	309.9	310.7	3.98						
285	357.9	358.9	4.81						
286	425.5	426.5	6.11						
287	337.9	338.7	4.41						
288	267.8	267.9	3.1						
289	295.8	296.8	4.26						

Example Compound	MW g/mol	LC-MS [MH] ⁺	log P	EC ₅₀ Abeta (uM)	S1 K _i (nM)	S2 K _i (nM)	MLM T _{1/2} (min.)	T _{1/2} *	hERG IC ₅₀ (nM)
290	309.8	310.6	4.08						
291	323.9	324.9	4.06						
292	392.0	393.2	6.49	1.6					
293	275.4	276.5	3.48						
294	275.4	276.7	3.48						
295	289.4	290.5	3.45						
296	261.4	262.7	2.83						
297	321.5	322.4	5.81						
298	295.8	296.8	4.26						
299	303.4	304.4	3.81						
300	261.4	262.7	3.65						
301	233.3	234.5	2.49						
302	261.4	262.4	3.65						
303	368.0	369.0	4.58						
304	275.4	276.5	3.38						
305	553.6	554.6	3.1						
306	329.4	330.3	3.89						
307	391.5	392.6	5.08						
308	261.4	262.4	3.01						
309	579.6	579.6	3.96						
310	261.4	262.5	3.2						
311	371.4	372.3	4.69						
312	329.4	330.5	4.53						
313	315.5	316.4	4.28						
314	401.5	402.7	4.85						
315	301.3	302.3	3.37						
316	341.5	342.5	4.34						
317	329.4	330.6	4.53						
318	279.4	280.3	3.8						
319	279.4	280.6	3.8						
320	293.4	294.4	3.62						
321	321.5	322.7	4.61						
322	321.4	322.4	3.95						
323	249.4	250.4	4.15						
324	337.9	338.7	5.07						
325	303.5	304.6	4.47						
326	307.4	308.4	3.6						


*normalized to 4-(3-(4-(trifluoromethyl)benzylamino)butyl)-2-methoxyphenol

Synthetic Examples


Example Syn 1: Preparation of Example Compound 9:



Example 9




General Procedure for the Preparation of Compound S1-2



[0322] In three-neck flask, was placed magnesium (9.39 g, 391.09 mmol, 1.1 eq) and a grain of iodine. Then 15 percent volume of compound **S1-1** (80 g, 355.54 mmol, 1.0 eq) in THF (800 mL) was added into the mixture under nitrogen atmosphere, and the stirred mixture was heated to 70°C until yellow brown disappeared and then stirred at that temperature for another 6 h, to give a solution of compound **S1-2** (0.44 M) in THF, which was used directly in next step.

General Procedure for the Preparation of Compound S1-4




[0323] To a solution of compound **S1-2** (320 mmol, 1.0 eq) in THF (500 mL) was added CuCl (3.17 g, 32 mmol, 0.1 eq). When the solution was cooled to 0 °C the compound **S1-3** (64 g, 320 mmol, 1.0 eq) was added dropwise. The reaction mixture was stirred at rt for 3 h, quenched with saturated NH₄Cl solution, extracted with ethyl acetate. The organic layer was dried over Na₂SO₄, concentrated under vacuum. The residue was purified by column chromatography (PE/EA, 50:1~10:1) to afford the title compound **S1-4** (51 g, 46%). ¹H NMR (400 MHz, CDCl₃) δ 7.57-7.49 (m, 4H), 4.08-4.03 (m, 4H), 3.80 (d, J = 2.4 Hz, 1H), 1.62-1.58 (m, 6H), 1.13-1.09 (m, 6H)

TLC: PE/EA = 10:1, UV 254 nm

R_f (Compound **S1-4**) = 0.5

General Procedure for the Preparation of Compound S1-5




[0324] To a solution of compound **S1-4** (51 g, 147.25 mmol, 1.0 eq) in MeOH (300 mL) was added 1N aqueous solution of NaOH (736 mL, 736.25 mmol, 5.0 eq). The reaction mixture was stirred at 90 °C for 3 h, and then cooled to rt. The mixture was acidified with 1N HCl solution until pH 4-5 and extracted with ethyl acetate. The organic layer was dried over Na₂SO₄ and concentrated under vacuum to give a compound **S1-5** (40 g, 94%), which was used directly in next step without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.57-7.55 (m, 2H), 7.50-7.48 (m, 2H), 3.86 (s, 1H), 1.59 (s, 6H);

TLC: DCM/MeOH = 10:1, UV 254 nm

R_f (Compound **S1-5**) = 0.5

General Procedure for the Preparation of Compound S1-6



[0325] A solution of compound **S1-5** (40 g, 137.82 mmol, 1.0 eq) in DMSO (200 mL) was stirred at 150 °C for 2 h, diluted with ethyl acetate, washed with water, brine. Then the organic phase was dried over Na₂SO₄ and concentrated under vacuum to afford the title


compound **S1-6** (30 g, 88%), which was used directly without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.57-7.55 (m, 2H), 7.48-7.45 (m, 2H), 2.67 (s, 2H), 1.49-1.46 (m, 6H);

TLC: DCM/MeOH = 10:1, UV 254 nm

R_f (Compound **S1-5**) = 0.4

R_f (Compound **S1-6**) = 0.8

General Procedure for the Preparation of Compound S1-8




[0326] To a solution of compound **S1-6** (30 g, 121.84 mmol, 1.0 eq) in THF (500 mL) was added HOBt (19.8 g, 146.2 mmol, 1.2 eq), EDCI (28 g, 146.21 mmol, 1.2 eq), TEA (37 g, 365.52 mmol, 3.0 eq) and compound **S1-7** (23.77 g, 243.68 mmol, 2.0 eq). The reaction was stirred at rt overnight and then quenched with water, extracted with ethyl acetate. The organic layer was dried over Na₂SO₄ and concentrated to give the crude product, which was purified by column chromatography on a silica gel (PE/EA, 10:1~3:1) to give compound **S1-8** (31 g, 88%). ¹H NMR (400 MHz, CDCl₃) δ 7.56-7.54 (m, 2H), 7.50-7.47 (m, 2H), 3.59 (s, 3H), 3.06 (s, 3H), 2.78 (s, 2H), 1.49 (s, 6H);

TLC: PE/EA = 3:1, UV 254 nm

R_f (Compound **S1-6**) = 0.4


R_f (Compound **S1-8**) = 0.8

General Procedure for the Preparation of Compound S1-10



[0327] In three-neck flask, was placed magnesium (8.0 g, 331.39 mmol, 1.1 eq) and a grain of iodine. Then 15 percent volume of compound **S1-9** (40.67 g, 301.26 mmol, 1.0 eq) in THF(300 mL) was added into the mixture under nitrogen atmosphere, and the stirred mixture was heated to 70 °C until yellow brown disappeared and then the solution was stirred at that temperature for another 6 h, to give a solution of compound **S1-10** (1.0 M) in THF, which was used directly in next step.

General Procedure for the Preparation of Compound S-11




[0328] To a solution of compound **S1-8** (31 g, 107.16 mmol, 1.0 eq) in THF (200 mL) was added compound **S1-10** (1.0 M, 215 mL, 214.32 mmol, 2.0 eq) at 0 °C. The reaction was stirred at rt for 3 h. Then the mixture was quenched with saturated NH_4Cl solution, extracted with ether, and the organic layer was dried over Na_2SO_4 , concentrated in vacuum to give crude product, which was purified by column chromatography on a silica gel (PE) to give crude compound **S-11** (28.9 g, 95%). ^1H NMR (400 MHz, CDCl_3) δ 7.56-7.54 (m, 2H), 7.47-7.45 (m, 2H), 7.75 (m, 1H), 2.75 (s, 2H), 2.01-1.99 (m, 3H), 1.74-1.73 (m, 3H), 1.44-1.42 (m, 6H);

TLC: PE/EA = 10:1, UV 254 nm

R_f (Compound **S1-8**) = 0.5

R_f (Compound **S1-11**) = 0.9

General Procedure for the Preparation of Example 9




[0329] A mixture of compound **S1-11** (2.4 g, 8.44 mmol, 1.0 eq), compound **S1-12** (1.43 g, 8.44 mmol, 1.0 eq) and $\text{Ti}(\text{EtO})_4$ (7.7 g, 33.76 mmol, 4.0 eq) in THF (100 mL) was stirred at 70 °C overnight under nitrogen atmosphere. Then the mixture was allowed to cool to rt, NaBH_4 (1.23 g, 33.76 mmol, 4.0 eq) was added. After complete addition, the mixture was stirred at rt for 3 h. Then water was added and extracted with ethyl acetate, filtered. The organic phase was dried over Na_2SO_4 , concentrated in vacuum. The residue was purified by column chromatography on a silica gel (PE/EA, 10:1~1:1) to give compound **Example 9**, which was dissolved in HCl/EA (2.0 M, 10 mL). The mixture was stirred at rt for 2 h, concentrated to give **Example 9 HCl** (600 mg, 15%) as oil. ^1H NMR (400 MHz, CD_3OD) δ 7.60-7.52 (m, 4H), 4.70-4.67 (m, 1H), 3.95-3.92 (m, 2H), 3.79-3.73 (m, 1H), 3.61-


3.59 (m, 1H), 2.81-2.73 (m, 2H), 2.38-2.22 (m, 2H), 1.98 (s, 6H), 1.64-1.61 (m, 2H), 1.46-1.43 (m, 12H), 1.31-1.28 (m, 6H);

MS: [M+H]⁺ = 438.5

Example Syn 2: Preparation of Example Compound 262:




General Procedure for the Preparation of Compound S2-7



[0330] To a solution of compound **S2-6** (450 g, 1.83 mol, 1.0 eq) in anhydrous THF (4.5 L) cooled to 0 °C, a solution of 1 M BH₃ (2.75 L, 2.745 mol, 1.5 eq) was added

dropwise. The mixture was stirred at rt for 16 h. The reaction was monitored by TLC. The colorless homogeneous reaction mixture was cooled to 0 °C, and MeOH (2 L) was carefully added followed by water (1 L). MeOH and THF were then removed under *vacuum*. The mixture was extracted with DCM (3 x 1 L), the combined organic extracts were washed with brine (1 L), dried over MgSO₄, filtered, and concentrated to dryness under reduced pressure to give crude compound **S2-7** (340 g, 80%) as a colorless oil. ¹H NMR (400 MHz, CDCl₃): δ 7.57 (d, *J* = 8.0 Hz, 2H), 7.47 (d, *J* = 8.4 Hz, 2H), 3.47 (t, *J* = 7.2 Hz, 2H), 1.98 (t, *J* = 7.2 Hz, 2H), 1.29 (s, 6H).

General Procedure for the Preparation of Compound S2-9




[0331] A solution of compound **S2-7** (296 g, 1.27 mol, 1.0 eq) was dissolved in CH₃CN (3.6 L), NMI (10.3 g, 127 mmol, 0.1 eq), TEMPO (9.8 g, 63 mmol, 0.05 eq) and BPy (9.7 g, 63 mmol, 0.05 eq) were added, then Cu(PF₆)(MeCN)₄ (23.5 g, 63 mmol, 0.05 eq) was added. The mixture was stirred under oxygen atmosphere for 4 h. The reaction was monitored by TLC. The mixture was poured into water (2 L), extracted with DCM (3 x 2 L). The combined extracts were washed with water, dried over Na₂SO₄ and concentrated to give compound **S2-9** (281 g, 96%) as a dark green oil. ¹H NMR (400 MHz, CDCl₃): δ 9.54 (s, 1H), 7.61 (d, *J* = 7.6 Hz, 2H), 7.51 (d, *J* = 7.6 Hz, 2H), 2.66 (s, 2H), 1.49 (s, 6H).

TLC: PE/EA = 20:1, UV 254 nm

R_f (compound **S2-7**) = 0.1

R_f (**S2-9**) = 0.8

General Procedure for the Preparation of Compound S2-11R




[0332] To a solution of compound **S2-9** (150 g, 652.5 mmol, 1.0eq) in THF (750 mL), compound **S2-10R** (118.96 g, 978.77 mmol, 1.5eq) and $\text{Ti}(\text{OEt})_4$ (74.4 g, 326.2 mmol, 0.5 eq) were added. The mixture was stirred at rt overnight, and then quenched with water (24 g, 1.3 mol), filtered by celite pad and diluted with ethyl acetate, washed with brine, water. The organic layer was dried over Na_2SO_4 and concentrated to give crude product, which was purified by column chromatography on a silica gel (PE/EA, 5:1) to give compound **S2-11R** (165 g, 75.9%).

TLC: PE/EA = 3:1, UV 254 nm

R_f (Compound **S2-9**) = 0.6


R_f (Compound **S2-11R**) = 0.5

General Procedure for the Preparation of Compound S2-12



[0333] In three-neck flask, was placed magnesium (31.9 g, 1.3 mol, 1.1 eq) and a grain of iodine. Then 15 percent volume of compound **S2-31** (149 g, 1.2 mol, 1.0 eq) in THF (1.5 L) was added into the mixture under nitrogen atmosphere, and the stirred mixture was heated to 70 °C until yellow brown disappeared. Then the remaining solution was added dropwise and stirred for another 6 h to give a solution of compound **S2-12** (1.0 M) in THF, which was used directly in next step.

General Procedure for the Preparation of Compound S2-13R



[0334] To a solution of compound **S2-11R** (120 g, 359.9 mmol, 1.0 eq) in THF (80 mL) was added the compound **S2-12** (1080 mL, 539.9 mmol, 1.50 eq, 0.5 M) at 0-5°C.

After stirring at rt for overnight, the reaction was quenched by NH_4Cl , extracted with ethyl acetate. The organic layer was dried over Na_2SO_4 and concentrated to give crude product, which was purified by column chromatography on a silica gel (PE/EA, 10:1~3:1) to give compound **S2-13R** (69.1 g, 49.3%).


[0335] The compound **S2-13R** (69.1 g) was recrystallized with ether/hexane (1/1) (30mL) at 4~8 °C for up to two days, to give white solid (25.6 g, 37%), which was used to produce Example 262 in 99% ee. The mother liquor was recrystallized with ether/hexane (1/1) (20mL) to give white solid (15.1 g).

TLC: PE/EA = 3:1, UV 254 nm

R_f (Compound **S2-11R**) = 0.5

R_f (Compound **S2-13R**) = 0.3

General Procedure for the Preparation of Compound S2-14R




[0336] Compound **S2-13R** (34.6 g, 88.8 mmol, 1.0 eq) was added in the round bottom flask. HCl in EtOAc (90 mL, 2N, 2 eq) was then added. After stirring for two hours, about 80 mL of solvent was removed under vacuum. The mixture diluted with ether (20 mL), the white solid was filtered out as compound **S2-14R** hydrochloride. The hydrochloride was dissolved in H_2O (30 mL), alkalinized with saturated aqueous of Na_2CO_3 to pH 10. Then extracted with ether (30mL x 3), dried over (Na_2SO_4), concentrate to give colorless oil (20.2 g 79.7%).

TLC: PE/EA = 3:1, UV 254 nm

R_f (Compound **S2-13R**) = 0.3


R_f (Compound **S2-14R**) = 0.1

General Procedure for the Preparation of Example Compound 262




[0337] To a solution of compound **S2-14R** (4.8 g, 16.8 mmol, 1.0 eq) in ACN (48 mL) was added the compound **S2-15** (8.35 g, 16.8 mmol, 1.0 eq) and K_2CO_3 (4.46 g, 33.6 mmol, 2.0 eq). After stirring at 85 °C overnight, the reaction was quenched by H_2O (50 mL) and extracted with ethyl acetate (40 mL x 3). The organic layer was dried over Na_2SO_4 and concentrated to give crude product, which was purified by column chromatography on a silica gel (PE/EA, 10:1~3:1) to give compound **Example 262** (5.1 g 69%) as the free base. The ee% was determined to be 99% on chiral HPLC. The free base was dissolved in HCl/EA (1.3 M, 50 mL). The mixture was stirred at rt for 2 h, concentrated to give **Example 262 HCl** (5.5 g, 69%) as a white solid. ^1H NMR (400 MHz, CD_3OD) δ 7.60-7.52 (m, 4H), 4.70-4.67 (m, 1H), 3.95-3.92 (m, 2H), 3.79-3.73 (m, 1H), 3.61-3.59 (m, 1H), 2.81-2.73 (m, 2H), 2.38-2.22 (m, 2H), 1.98 (s, 6H), 1.64-1.61 (m, 2H), 1.46-1.43 (m, 12H), 1.31-1.28 (m, 6H); MS: $[M+H]^+ = 438.5$


[0338] General Procedure for the Preparation of Compound 2-15



[0339] To a solution of compound **S2-16** (5.0 g, 26.6 mmol, 1.0 eq) in DCM (50 mL) was added the compound TsCl (20.2 g, 106.2 mmol, 4.0 eq) and pyridine (8.4 g, 106.2 mmol, 4.0 eq). After stirring at rt overnight, the reaction was quenched by H_2O (100 mL) and extracted with ethyl acetate (40 mL x 3). The organic layer was dried over Na_2SO_4 and concentrated to give crude product, which was purified by column chromatography on a silica gel (PE/EA, 10:1~3:1) to give compound **S2-15** (5.4 g 40.9 %).

Example Syn 3: Preparation of Example Compound 14:

General Procedure for the Preparation of Example Compound 14




[0340] To a mixture of compound **S3-9** (60.0 g, 0.26 mol, 1.0 eq) and compound **S3-10** (30.0 g, 0.26 mol, 1.0 eq) in MeOH (600 mL) was added NaBH₄ (39.6 g, 1.04 mol, 4.0 eq). The mixture was stirred at rt for 3 h under nitrogen atmosphere. Then brine was added and the solution was extracted with ethyl acetate. The organic phase was combined, dried over Na₂SO₄, concentrated in vacuum. The residue was purified by column chromatography to give compound **Example 14** (45.1 g, 52%).

TLC: DCM: MeOH = 10:1

R_f(**Example 14**) = 0.2


General Procedure for the Preparation of Example Compound 14 HCl



[0341] A mixture of **Example 14** (65 g, 0.2 mol) in HCl/EA (2.5 M, 160 mL) was stirred at rt for 2 h. The cloudy mixture was concentrated to give **Example 14 HCl** (68 g, 94%) as white solid. $^1\text{H NMR}$ (400 MHz, CD_3OD): δ ppm 7.67-7.61 (m, 4H), 3.31 – 3.3 (m, 3H), 3.15 – 3.11 (m, 2H), 2.85 – 2.79 (m, 2H), 2.17 – 2.12 (m, 2H), 1.82 – 1.76 (m, 2H), 1.43 (s, 6H), 1.27 (s, 3H).


MS: $[\text{M}+\text{H}]^+=330.20$

Example Syn 4: Preparation of Example Compound 17:



[0342] A mixture of compound **S4-11** (300 mg, 1.3 mmol, 1.0 eq), compound **S4-12** (150 mg, 1.3 mmol, 1.0 eq) and NaBH_4 (197 mg, 5.2 mmol, 4.0 eq) in MeOH (10 mL) was stirred at rt for 3 h under nitrogen atmosphere. Then aq. NaCl was added and extracted with ethyl acetate, filtered. The organic phase was dried over Na_2SO_4 , concentrated in vacuum. The residue was purified by pre-TLC to give compound **Example 17**, which was dissolved in HCl/EA (1.3 M, 10 mL). The mixture was stirred at rt for 2 h, concentrated to give **Example 17 HCl** (144.1 mg, 31.4%) as white solid. $^1\text{H NMR}$ (400 MHz, CD_3OD): δ 7.75-7.60 (m, 4H), 4.07-4.03 (m, 1H), 3.63-3.57 (m, 1H), 3.34-3.30 (m, 6H), 3.16-3.09 (m, 2H), 2.87-2.83 (m, 2H), 2.19-2.14 (m, 2H), 1.43 (s, 6H);

MS: $[\text{M}+\text{H}]^+=316.4$

Example Syn 5: Preparation of Example Compound 307:


[0343] To a solution of **compound S5-9** (3.54 g, 15.38 mmol, 1.0 eq) in MeOH (60 mL) was added compound **S5-10** (3.0 g, 16.9 mmol, 1.1 eq) and 2 drops of AcOH. The reaction mixture was stirred at rt for 2 h. NaBH₄ (2.33 g, 61.55 mmol, 4.0 eq) was added and then the reaction mixture was stirred at rt for 2 h. The reaction was quenched with H₂O (20 mL), filtered, extracted with EA (20 mL x 3) and concentrated to get a residue, which was purified by column chromatography (PE:EA=1:1) to give compound **Example 307** (1.3 g, 21.6%).

TLC: DCM:EA:MeOH=1:1:0.1

R_f(**compound S5-9**) = 0.9

R_f(**compound Example 307**) = 0.3


General Procedure for the Preparation of Example 307 HCl



[0344] To a solution of compound **Example 307** (1.9 g, 4.85 mmol, 1.0 eq) in EA (5 mL) was added EA/HCl (2.5 M, 6 mL, 3.0 eq). The reaction mixture was stirred at rt for 30 min. The reaction mixture was concentrated to give **Example 307 HCl** (2.1 g, 100%). ¹H NMR (400 MHz, CDCl₃) δ 7.62-7.60 (m, 2H), 7.50-7.47 (m, 4H), 7.46-7.31 (m, 2H), 7.28 (m, 2H), 3.32 (m, 2H), 3.13 (m, 2H), 2.64 (m, 2H), 2.32 (m, 2H), 1.67 (m, 3H), 1.39 (m, 6H);

MS: [M+H]⁺= 392.6

Example Syn 6: Preparation of Example Compound 191:




[0345] To a solution of compound **S6-9** (5.4 g, 23.68 mmol, 1.0 eq) in MeOH (100 mL) was added compound **S6-12** (3.0 g, 26 mmol, 1.1 eq) and 2 drops of AcOH. The reaction mixture was stirred at rt for 2 h. NaBH₄ (3.58 g, 94.71 mmol, 4.0 eq) was added and then the reaction mixture was stirred at rt for 2 h. The reaction was quenched with H₂O (30 mL), filtered, extracted with EA (30 mL x 3) and concentrated to get a residue, which was purified by column chromatography (PE:EA=1:1) to give **Example 191** (3.4 g, 44%).

TLC: DCM:EA:MeOH=1:1:0.1

R_f(**compound S6-9**) = 0.9

R_f(**Example 191**) = 0.3


General Procedure for the Preparation of Example 191 HCl



[0346] To a solution of **Example 191** (3.4 g, 10.33 mmol, 1.0 eq) in EA (10 mL) was added **EA/HCl** (2.5 M, 8.9 mL, 2.0 eq). The reaction mixture was stirred at rt for 30 min. The reaction mixture was concentrated to give **Example 191 HCl** (3.5 g, 93%). ¹H NMR

(400 MHz, CDCl₃) δ 7.60-7.58 (m, 2H), 7.49-7.47 (m, 2H), 3.56 (s, 1H), 3.25 (s, 3H), 3.19 (m, 2H), 2.80 (m, 2H), 2.59 (m, 2H), 2.39 (m, 2H), 2.30 (m, 2H), 1.93 (m, 2H), 1.37 (s, 6H); MS: [M+H]⁺ = 330.2

Example Syn 7: Preparation of Example Compound 317:




[0347] To a solution of compound **S7-9** (5.46 g, 23.7 mmol, 1.0 eq) in MeOH (100 mL) was added compound **S7-14** (3.0 g, 26 mmol, 1.1 eq) and 2 drops of AcOH. The reaction mixture was stirred at rt for 2 h. NaBH₄ (3.58 g, 94.63 mmol, 4.0 eq) was added and then the reaction mixture was stirred at rt for 2 h. The reaction was quenched with H₂O (30 mL), filtered, extracted with EA (30 mL x 3) and concentrated to get a residue, which was purified by column chromatography (PE:EA=1:1) to give **Example 317** (2.1 g, 27%).

TLC: DCM:EA:MeOH=1:1:0.1

R_f(**compound S7-9**) = 0.9


R_f(**Example 317**) = 0.3

General Procedure for the Preparation of Example 317 HCl



[0348] To a solution of **Example 317** (2.1 g, 6.37 mmol, 1.0 eq) in EA (10 mL) was added **EA/HCl** (2.5 M, 6 mL, 2.0 eq). The reaction mixture was stirred at rt for 30 min. The reaction mixture was concentrated to give **Example 317 HCl** (2.1 g, 91%). ¹H NMR (400 MHz, CDCl₃) δ 12.17 (m, 1H), 7.61-7.59 (m, 2H), 7.50-7.48 (m, 2H), 4.16-4.13 (m, 1H), 3.90-3.86 (m, 1H), 3.55-3.53 (m, 1H), 3.33 (m, 1H), 3.23 (s, 3H), 2.63-2.59 (m, 2H), 2.51 (m, 1H), 2.23 (m, 1H), 2.19-1.90 (m, 4H), 1.39-1.37 (m, 6H); MS: [M+H]⁺= 330.6

Example Syn 8: Preparation of Example Compound 306:




[0349] To a solution of compound **S8-9** (3.08 g, 13.3 mmol, 1.0 eq) in MeOH (80 mL) was added compound **S8-16** (2.0 g, 17.36 mmol, 1.1 eq) and 2 drops of AcOH. The reaction mixture was stirred at rt for 2 h. NaBH₄ (2.02 g, 53.4 mmol, 4.0 eq) was added and then the reaction mixture was stirred at rt for 2 h. The reaction was quenched with H₂O (30 mL), filtered, extracted with EA (30 mL x 3) and concentrated to get a residue, which was purified by column chromatography (PE:EA=1:1) to give **Example 306** (1.2 g, 27%).

TLC: DCM:EA:MeOH=1:1:0.1

R_f(compound **S8-9**) = 0.9

R_f(**Example 306**) = 0.3


General Procedure for the Preparation of Example 306 HCl



[0350] To a solution of **Example 306** (2.0 g, 26.07 mmol, 1.0 eq) in EA (5 mL) was added **EA/HCl** (2.5 M, 5 mL, 2.0 eq). The reaction mixture was stirred at rt for 30 min. The reaction mixture was concentrated to give **Example 306 HCl** (2.28 g, 100%). ¹H NMR (400 MHz, CDCl₃) δ 12.75 (m, 1H), 7.60-7.58 (m, 2H), 7.49-7.47 (m, 2H), 3.53-3.51 (m, 2H), 2.63-2.60 (m, 2H), 2.53-2.50 (m, 2H), 2.34-2.30 (m, 2H), 2.05 (m, 4H), 2.00 (m, 1H), 1.97-1.88 (m, 2H), 1.38 (s, 6H); MS: [M+H]⁺= 330.3

CLAIMS:*What Is Claimed Is:*

1. A compound of Formula I or pharmaceutically acceptable salt thereof:




wherein:

each of R^a , R^b , R^c , R^d and R^e is independently selected from the group consisting of, H, hydroxyl, halo, alkyl, alkoxy, CF_3 , SO_2CH_3 , and morpholino;

R^1 is selected from the group consisting of hydrogen, alkyl, phenyl, or $-CH=C(CH_3)_2$; and

R^2 is an optionally substituted cyclic amino group.

2. The compound of claim 1, wherein R^2 is selected from the group consisting of optionally substituted aziridinyl, optionally substituted pyrrolidinyl, optionally substituted imidizolidinyl, optionally substituted piperidinyl, optionally substituted piperazinyl, optionally substituted oxopiperazinyl, and optionally substituted morpholinyl.
3. The compound of claim 1 wherein R^2 is optionally substituted piperidinyl.
4. The compound of claim 1, wherein R^2 is selected from the group consisting of




wherein:


R^3 is selected from the group consisting of hydrogen and C_1 - C_8 alkyl;

R^4 is selected from the group consisting of hydrogen, hydroxyl, halo, CF_3 , alkoxy, aryloxy, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl,

optionally substituted C₅-C₁₀ heteroaryl, optionally substituted C₃-C₁₀ cycloalkyl and optionally substituted C₃-C₁₀ heterocycloalkyl; and each of R⁵ and R⁶ is independently selected from the group consisting of hydrogen, hydroxyl, sulfonyl, dialkylamino, optionally substituted C₁-C₁₀ alkyl, optionally substituted C₅-C₁₀ aryl, optionally substituted C₅-C₁₀ heteroaryl, optionally substituted C₃-C₁₀ cycloalkyl and optionally substituted C₃-C₁₀ heterocycloalkyl.


5. The compound of claim 1, wherein R² is selected from the group consisting of:






wherein R^{3a} selected from the group consisting of hydrogen and C_1 - C_8 alkyl; and n is an integer selected from 0, 1 and 2.

6. The compound of claim 1 wherein R^2 is selected from




7. The compound of claim 1, wherein R^2 is selected from the group consisting of:




wherein R^7 is selected from the group consisting of hydrogen, hydroxyl, sulfonyl, dialkylaminosulfonyl, alkoxycarbonyl, acyl, benzoyl, cycloalkylcarbonyl, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl, optionally substituted C_5 - C_{10} heteroaryl, optionally substituted C_3 - C_{10} cycloalkyl and optionally substituted C_3 - C_{10} heterocycloalkyl.


8. The compound of claim 1, wherein R^2 is:



9. wherein R^8 is hydrogen, hydroxyl, sulfonyl, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl; optionally substituted C_5 - C_{10} heteroaryl, optionally substituted C_3 - C_{10} cycloalkyl and optionally substituted C_3 - C_{10} heterocycloalkyl. The compound of claim 1, wherein R^2 is selected from the group consisting of





10. The compound of claim 1, wherein R^2 is bicyclic.
11. The compound of claim 1, wherein R^2 is selected from the group consisting of




wherein R^9 is, independently selected from the group consisting of hydrogen, hydroxyl, sulfonyl, optionally substituted C_1 - C_{10} alkyl, optionally substituted C_5 - C_{10} aryl, optionally substituted C_5 - C_{10} heteroaryl, optionally substituted C_3 - C_{10} cycloalkyl and optionally substituted C_3 - C_{10} heterocycloalkyl.


12. A compound of claim 1 selected from the groups consisting of:









,




,




,




,




,




,




,




,




,




,




,















,




,




,







,




,




,




,




,




,




,




,




,




,




,












,




,









,




,




,




,




,




,




,




,




,




,




,




,




,

















,






,



,











,

,




,

,




,


,




,





,




,









,




,




,




,




,




,




,




,




,




,




,




,




,







,




,




,




,




,




,




,




,












,





,





,






,


,




,




,




,




,




,















,




,




,







,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,




,



,




, and




;

or pharmaceutically acceptable salt thereof.


13. The compound of claim 1, selected from the group consisting of




,



,




, and











;









or a pharmaceutically acceptable salt thereof.

14. A compound selected from the group consisting of



,






and

or a pharmaceutically acceptable salt thereof.

15. A pharmaceutical composition comprising a compound of any one of claims 1 to 14 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.

16. The pharmaceutical composition of claim 15 wherein the compound is selected from the group consisting of



or a pharmaceutically acceptable salt thereof.

17. A method of treating Alzheimer's disease comprising administering to a subject in need thereof, a therapeutically effective amount of a pharmaceutical composition according to claim 15 or 16.


18. A method for inhibiting cognitive decline in a subject exhibiting, or at risk of exhibiting, cognitive decline, comprising administering a therapeutically effective amount of a pharmaceutical composition according to claim 15 or 16.

19. A method of inhibiting amyloid beta effect on a neuronal cell comprising administering to a subject in need thereof, a therapeutically effective amount of a pharmaceutical composition according to claim 15 or 16.

20. A method of treating mild cognitive impairment in Alzheimer's disease in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition according to claim 15 or 16.

21. Use of a compound according to any one of claims 1 to 14 in the manufacture of a medicament for the treatment of Alzheimer's disease.


22. The use of claim 21 wherein the compound is selected from the group consisting of



or a pharmaceutically acceptable salt thereof.


23. A compound according to any one of claims 1 to 14 for use in the treatment of Alzheimer's disease.

24. The compound of claim 23 selected from the group consisting of



or a pharmaceutically acceptable salt thereof.

25. A compound according to any one of claims 1 to 14 for use in medical therapy.
26. The compound of claim 25 selected from the group consisting of



or a pharmaceutically acceptable salt thereof.

27. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any one of claims 1 to 14 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.