



(86) **Date de dépôt PCT/PCT Filing Date:** 2006/06/06
(87) **Date publication PCT/PCT Publication Date:** 2006/12/14
(45) **Date de délivrance/Issue Date:** 2017/01/17
(85) **Entrée phase nationale/National Entry:** 2007/12/03
(86) **N° demande PCT/PCT Application No.:** US 2006/021905
(87) **N° publication PCT/PCT Publication No.:** 2006/133184
(30) **Priorité/Priority:** 2005/06/07 (US60/688,409)

(51) **Cl.Int./Int.Cl. C12Q 1/68** (2006.01),
C12P 19/34 (2006.01), **C12Q 1/70** (2006.01)
(72) **Inventeur/Inventor:**
MOSER, MICHAEL JAMES, US
(73) **Propriétaire/Owner:**
LUMINEX CORPORATION, US
(74) **Agent:** BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) **Titre : METHODES DE DETECTION ET DE TYPAGE D'ACIDES NUCLEIQUES**
(54) **Title: METHODS FOR DETECTION AND TYPING OF NUCLEIC ACIDS**

(57) **Abrégé/Abstract:**

Disclosed are methods and kits for identifying and characterizing polynucleotide sequences in a sample which may include a heterogeneous sample. Some of the methods and kits are directed to the identification and characterization of a virus in a sample, which may include HIV capable of cause AIDS or AIDS-like symptoms. The virus may be HIV-1, and may also include drug resistant mutations. The methods may include reacting a mixture that includes, in addition to nucleic acid isolated from the sample, at least one oligonucleotide capable of specifically hybridizing to HIV nucleic acid where the oligonucleotide includes at least one non-natural base. In addition, the methods may include detection of one or more mutations in HIV nucleic acid that are associated with drug resistance.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 December 2006 (14.12.2006)

PCT

(10) International Publication Number
WO 2006/133184 A3

(51) International Patent Classification:
C12Q 1/68 (2006.01)

(21) International Application Number:

PCT/US2006/021905

(22) International Filing Date: 6 June 2006 (06.06.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/688,409

7 June 2005 (07.06.2005) US

(71) Applicant (for all designated States except US): **ERAGEN BIOSCIENCES, INC.** [US/US]; 918 Deming Way, Suite 201, Madison, Wisconsin 53717-1944 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **MOSER, Michael, James** [US/US]; 7406 Farmington Way, Madison, Wisconsin 53717 (US).

(74) Agent: **FOLEY & LARDNER LLP**; 150 East Gilman Street, Madison, Wisconsin 53202-1481 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,

KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

5 April 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS FOR DETECTION AND TYPING OF NUCLEIC ACIDS

(57) Abstract: Disclosed are methods and kits for identifying and characterizing polynucleotide sequences in a sample which may include a heterogeneous sample. Some of the methods and kits are directed to the identification and characterization of a virus in a sample, which may include HIV capable of cause AIDS or AIDS-like symptoms. The virus may be HIV-1, and may also include drug resistant mutations. The methods may include reacting a mixture that includes, in addition to nucleic acid isolated from the sample, at least one oligonucleotide capable of specifically hybridizing to HIV nucleic acid where the oligonucleotide includes at least one non-natural base. In addition, the methods may include detection of one or more mutations in HIV nucleic acid that are associated with drug resistance.



WO 2006/133184 A3

METHODS FOR DETECTION AND TYPING OF NUCLEIC ACIDS

FIELD OF THE INVENTION

[0002] The invention relates generally to the field of detecting and typing nucleic acids. More specifically, the invention relates to detecting and typing infectious agents, such as viruses. In particular, the invention relates to the detection of HIV, and to the detection of mutations in HIV that may lead to drug resistance.

BACKGROUND

[0003] Human Immunodeficiency Virus (“HIV”), an approximately 10-kb, enveloped, single-stranded RNA retrovirus, is the causative agent of Acquired Immunodeficiency Syndrome (“AIDS”). As the HIV epidemic continues to spread world-wide, the need for effective HIV detection methods remains paramount. Early detection of HIV infection is critical for preventing the spread of the virus and concomitant disease, and for determining effective treatments and therapies. However, a key obstacle to HIV detection and treatment has been – and remains to be – the incredible variability of HIV types, and the extent and swiftness of HIV mutation (Wain-Hobson in *The Evolutionary biology of Retroviruses*, SSB Morse Ed. Raven Press, NY, pgs 185 209 (1994)).

[0004] Molecular characterizations have shown that the HIV virus can be categorized into two broad types. HIV-1, discovered in 1984, is the main cause of AIDS around the world, particularly in the Western Hemisphere and in Europe. HIV-2, discovered two years later in 1986, is noted mainly in western Africa. Both HIV-1 and HIV-2 are additionally categorized into sub-types. For example, HIV-2 is broken down into sub-types A-G; sub-types A and B are considered “epidemic,” while C-G are considered “nonepidemic.” Similarly, molecular characterization of HIV-1 strains from around the world have identified three distinct groups, M, N and O. Group M viruses represent the majority of HIV-1 and based on sequence divergence Group M has been further subdivided into nine

different subtypes or clades, termed subtypes A, B, C, D, F, G, H, J, and K (Robertson, D. L. *et al.* In: Human Retroviruses and AIDS 1999-A Compilation and Analysis of Nucleic Acid and Amino Acid Sequences, Kuiken, C. *et al.* Eds., pgs. 492-505 (1999)).

[0005] The overall distribution of HIV-1 strains varies considerably in different geographic regions and is undergoing continual change. For example, while subtype B has been predominant in North America and Western Europe (*see e.g.*, McCutchan, F. E., AIDS 14 (suppl 3): S31-S44 (2000)), increasing numbers of non-subtype B infections are being observed in both Europe and the United States.

[0006] In addition to the great diversity of types and subtypes, drug-resistant and therapy-resistant mutants have also become prevalent. Due to the extreme mutability of the HIV virus (HIV does not employ a "proof-reading" mechanism during replication), the development of vaccines has been a major challenge, and the use of other drug therapies, such as anti-viral drugs, has been complicated by the rapid evolution of drug resistant strains.

[0007] Drug-resistance mutations have been identified in HIV-1 patients who have shown improvement under some type of drug treatment or therapy and have then experienced a "relapse" of HIV-1 viral growth and associated symptoms. Mutations associated with resistance to anti-viral drugs have been found in the gag, pol and env regions of the HIV viral genome, and have been shown to affect proteins such as reverse transcriptase, protease, and the GP41 envelope protein (*See e.g.*, Johnson, *et al.*, 2005 Special Contribution -Drug Resistance Mutations in HIV-1, 13:3, 125-131; Gingeras *et al.*, 1991, J. Infect. Dis. 164(6):1066-1074; Richman *et al.*, 1991, J. Infect. Dis. 164(6):1075-1081; Schinazi *et al.*, 1993, Antimicrob. Agents Chemother. 37(4):875-881; Najera *et al.*, 1994, AIDS Res. Hum. Retroviruses 10(11):1479-1488; Eastman *et al.*, 1995, J. AIDS Hum. Retrovirol. 9(3):264-273; Frenkel *et al.*, 1995, J. Clin. Microbiol. 33(2):342-347; Shiras *et al.*, 1995, Proc. Natl. Acad. Sci. USA 92(6):2398-2402; Leal *et al.*, 1996, Eur. J. Clin. Invest. 26(6):476-480; Cleland *et al.*, 1996, J. AIDS Hum. Retrovirol. 12(1):6-18; Schmit *et al.*, 1996, AIDS 10(9):995-999; Vasudevechari *et al.*, 1996, Antimicrob. Agents Chemother. 40(11):2535-2541; Winslow *et al.*, 1996, AIDS 10(11):1205-1209; Fontenot *et al.*, Virology 190(1):1-10; Cornelissen *et al.*, 1997, J. Virol. 71(9):6348-6358; Ives *et al.*, 1997, J. Antimicrob. Chemother. 39(6):771-779).

[0008] Determining whether an HIV-1 infected patient is carrying drug-resistant viral strains may be critical for proper treatment and therapy. For example, a clinician may be

able to decide whether to begin or maintain a particular antiretroviral therapy. Further, continued testing of patients for drug-resistant HIV mutants during treatment may be used to detect the emergence of a drug-resistant virus, thereby allowing the clinician to alter the therapy to something that may prove more effective. Thus, there is a need in the art for assays that provide information related to HIV drug-resistance.

SUMMARY

[0009] The methods and kits described herein relate to detecting the presence of nucleic acid in a sample. In particular, the methods and kits are useful for detecting a specific mutation in a heterogeneous population of polynucleotides. In some aspects, the methods and kits may be used to treat or diagnose viral infections in mammals. The methods and kits may relate to detecting and typing viruses that can cause Acquired Immunodeficiency Syndrome ("AIDS") or AIDS-like symptoms in mammals, such as HIV-1, using oligonucleotides having at least one non-natural nucleotide.

[0010] In some aspects, the methods and kits are used to detect a specific mutation in a heterogeneous population of polynucleotides, in which the specific mutation, if present, is located in a first region of the population. Typically, the population includes one or more additional mutations located in a second region.

[0011] The method may include amplifying the population of polynucleotides with a first set of primers to obtain an amplification product that includes the first region, in which method at least one primer of the first set of primers is capable of specifically hybridizing to the second region of the population. For example, the at least one primer of the first set of primers may be capable of specifically hybridizing to the second region of the entire population of polynucleotides as a universal primer. The method further may include amplifying the amplification product with a second set of primers to detect the mutation, in which at least one primer of the second primer set includes at least one non-natural base. In some embodiments, the primer of the second primer set may have a 3' nucleotide that is complementary to the first region if the mutation is present. Optionally, the second primer set may include a primer having a 3' nucleotide that is complementary to the first region if the mutation is absent. Typically, the at least one primer of the first primer set does not hybridize specifically to the first region of the population of polynucleotides. Typically, the at least one primer of the second primer set does hybridize specifically to the first region of the population.

[0012] In some embodiments the specific mutation includes a single base change. For example, the first region of the population of polynucleotides may consist of a single nucleotide.

[0013] In some embodiments, at least one primer of the first primer set or second primer set includes at least one non-natural base. Preferably, at least one primer of the second primer set includes at least one non-natural base. In some embodiments, the non-natural base may be selected from iC and iG. The at least one primer may include a label. For example, the label may be coupled to the non-natural base. Suitable labels include fluorophores and quenchers.

[0014] Amplification may be performed in a reaction mixture that includes at least one non-natural nucleotide having a non-natural base. The at least one non-natural nucleotide of the reaction mixture may base pair with the at least one non-natural base present in the primer of the first and/or second primer set. Optionally, the non-natural nucleotide is coupled to a label which may include fluorophores and quenchers. The quencher may quench a fluorophore present in the primer of the first and/or second primer set.

[0015] The method may be used to detect the mutation continuously during amplification or in real-time. The method may be used quantitatively.

[0016] In some embodiments, the population of polynucleotides includes a sequence of HIV-1. The mutation may be present in the HIV-1 polymerase or protease gene.

[0017] In some aspects, the methods and kits are used to detect a specific mutation in a heterogeneous population of polynucleotides, where the specific mutation, if present, is located in a first region of the population and the population includes one or more additional mutations located in a second region of the population. The method may include: (a) reacting the population and a mixture of oligonucleotides, where the mixture includes: (i) a first oligonucleotide capable of hybridizing to at least the first region of the population; and (ii) a pool of degenerate oligonucleotides capable of hybridizing to at least the second region of one or more polynucleotides of the population, where one or more oligonucleotides of the mixture include one or more non-natural bases and optionally a label; and (b) detecting the mutation. In additional aspects, the method may include: (a) reacting the population and a pool of degenerate oligonucleotides, where the oligonucleotides of the pool include one or more non-natural bases and where the pool includes: (i) at least one oligonucleotide capable of hybridizing to at least the first region of one or more polynucleotides of the population; and (ii) a plurality of oligonucleotides

capable of hybridizing to at least the second region of one or more polynucleotides of the population; and (b) detecting the mutation. In some embodiments, the non-natural bases are selected from the group consisting of iso-G, iso-C and a combination thereof. The degenerate oligonucleotides may include at least one label. In some embodiments, all of the degenerate oligonucleotides include a non-natural base and a label. Suitable labels include fluorophores and quenchers.

[0018] Detecting may include amplifying one or more polynucleotides of the population. For example, detecting may include amplifying one or more polynucleotides of the population in the presence of at least one non-natural nucleotide. The non-natural nucleotide may have a non-natural base (*e.g.*, iC and iG), which, optionally, is capable of base-pairing with the non-natural base of the mixture of oligonucleotides (*e.g.*, a non-natural base present in the degenerate oligonucleotides). The non-natural nucleotide may be coupled to a label. Suitable labels include fluorophores and quenchers. The methods may be used to detect mutations continuously during amplification or in real-time. The heterogeneous population may include polynucleotides of HIV-1.

[0019] In some aspects, the methods and kits are used to detect the presence, and optionally the absence, of a mutation in a polynucleotide at a specific nucleotide position. The method typically includes amplifying the polynucleotide with primers to detect the presence, and optionally the absence, of the mutation. In some embodiments, the primers include: (a) a first primer having a 3' nucleotide that is complementary to the polynucleotide (at a specific nucleotide position) if the mutation is present; (b) a second primer having a 3' nucleotide that is complementary to the polynucleotide (at a specific nucleotide position) if the mutation is absent. Typically, the first primer and the second primer are not complementary to the polynucleotide at one or more positions other than the 3' nucleotide and do not include identical nucleotides at the one or more positions. For example, the first and second primer may not be complementary to each other at the one or more positions other than the 3' nucleotide. In some embodiments, at least one of the first primer and the second primer includes at least one non-natural base (*e.g.*, iC and iG), which may be present at the one or more positions. At least one of the first primer and the second primer may include a label. Where both the first primer and second primer include a label, the label may be the same or different, preferably different. Suitable labels include fluorophores and quenchers. Optionally, the label may be coupled to the non-natural base. Amplification may be performed using a reaction mixture that includes at least one non-

natural nucleotide having a non-natural base, which optionally may base-pair with the non-natural base present in the first primer, the second primer, or both primers. The non-natural nucleotide of the reaction mixture may include a label. Suitable labels may include a fluorophore and a quencher, which optionally is capable of quenching a fluorophore, if present, in at least one of the first primer and second primer, preferably both. The methods may be used to detect mutations continuously during amplification or in real-time. The mutation may include a single nucleotide polymorphism present in HIV-1 nucleic acid, *e.g.*, in the polymerase gene or protease gene.

[0020] In some aspects, the methods and kits are used for identifying HIV-1 in a sample. The methods and kits may include (a) reacting a reaction mixture, where the reaction mixture includes: (i) the sample; (ii) at least one oligonucleotide comprising at least one non-natural base, where the oligonucleotide is capable of specifically hybridizing to HIV-1 nucleic acid; and (b) detecting HIV-1 nucleic acid if present in the sample. In some embodiments, the at least one oligonucleotide is selected from the group consisting of SEQ ID NO:2-97.

[0021] In some aspects, the methods and kits relate to detecting specific mutations in heterogeneous, (*e.g.*, polymorphic) populations of polynucleotides. By way of example, but not by way of limitation, the methods may be used to detect a specific mutation, such as a drug resistance mutation, in a heterogeneous population of HIV polynucleotides, such as HIV-1 polynucleotides. In some methods, the specific mutation may be a single base change, such as a transition or transversion. In other methods the specific mutation may be an insertion, a deletion or a rearrangement. Thus, the mutation if present, may be located in a first region of the population and the population may include one or more additional mutations located in a second region. In some methods, the "first region" may be only a single nucleotide.

[0022] To detect a mutation in such a polymorphic or heterogeneous population, the methods may include the steps of (a) amplifying the population with a first set of primers to obtain an amplification product, where at least one primer of the first set of primers may be capable of specifically hybridizing to the second region of the population; and (b) amplifying the product with a second set of primers to detect the mutation, where at least one primer of the second primer set may include at least one non-natural nucleotide.

[0023] In some embodiments, the at least one primer of the first primer set may not hybridize specifically to the first region of the population. In other methods, at least one primer of the first set of primers may include at least one non-natural nucleotide.

[0024] In other embodiments, the at least one primer of the second primer set may specifically hybridize to the first region of the population. In still other embodiments, the at least one primer of the second primer set, which may include at least one non-natural nucleotide, may also include a label. In further embodiments, the label may be a fluorophore. In some embodiments, amplifying the product with the second set of primers may include amplifying in the presence of at least one quencher coupled to a non-natural nucleotide. The methods of detecting a specific mutation may also include reverse transcription, amplification, and real-time detection. Some methods may include amplification in the presence of a quencher coupled to a non-natural nucleotide.

[0025] In some aspects, the methods and kits also relate to detecting a specific mutation in a heterogeneous population of polynucleotides using degenerate oligonucleotides. Such methods may include the steps of (a) reacting the population of heterogeneous polynucleotides, such as, for example HIV-1 polynucleotides, and a pool of degenerate oligonucleotides to detect the mutation. In some embodiments, the oligonucleotides of the pool may include one or more non-natural nucleotides, for example, iso-GTP ("iGTP"), iso-CTP ("iCTP") and combinations thereof. The degenerate oligonucleotides may further include at least one label, such as, for example, a fluorophore. The pool of degenerate oligonucleotides may include (i) at least one oligonucleotide capable of hybridizing to the first region of one or more polynucleotides of the population; and (ii) a plurality of oligonucleotides capable of hybridizing to the second region of one or more polynucleotides of the population.

[0026] In some embodiments, detecting the specific mutation may include reverse transcription, amplification and/or real time detection of one or more polynucleotides of the population. Amplification may be performed in the presence of one or more non-natural nucleotides and/or in the presence of at least one quencher coupled to a non-natural nucleotide. In some embodiments, the non-natural nucleotide coupled to the at least one quencher may be iCTP or iGTP.

[0027] The methods and kits also relate to detecting the presence or absence of a mutation, at a specific nucleotide, in a polynucleotide. By way of example, but not by way

of limitation, the methods may include detecting the presence or absence of a mutation, such as a drug resistance mutation, in an HIV polynucleotide such as HIV-1.

[0028] Such methods of detecting the presence or absence of a mutation may include amplifying the polynucleotide with primers. In these methods, the primers may include (a) a first primer having a 3' nucleotide that is complementary to the specific nucleotide where the mutation is present and capable of amplifying the polynucleotide; and (b) a second primer having a 3' nucleotide that is complementary to the specific nucleotide where the mutation is absent and capable of amplifying the polynucleotide. In some embodiments, the first primer and the second primer may include nucleotides that are non-complementary to the polynucleotide at one or more positions other than the 3' nucleotide. In other embodiments, the first and the second primers may include non-complementary nucleotides at different positions. In still other embodiments, the first and the second primers may include different non-complementary nucleotides at the same position. For example, in some embodiments, the first primer and the second primer are not complementary to the polynucleotide at a single position other than the 3' nucleotide and also do not include identical nucleotides at that single position.

[0029] In some embodiments, at least one of the first primer and the second primer may include at least one non-natural nucleotide. For example, the at least one non-natural nucleotide may be at the one or more non-complementary positions. Any non-natural nucleotide may be used; however, in some embodiments, iso-G, iso-C and combinations thereof may be preferred. In still other embodiments one or more of the primers may include a label; in some embodiments, the labels on each primer may be different. In some embodiments, at least one label may be a fluorophore. In some methods, detection may include reverse transcription, amplification and/or real time detection. In some embodiments, amplification may be performed in the presence of a quencher coupled to a non-natural base or non-natural base triphosphate.

[0030] Some methods may include an oligonucleotide that functions as an internal control nucleic acid. In other methods, the reaction mixture may include at least two oligonucleotides capable of hybridizing to an internal control nucleic acid and that may function as primers to amplify the internal control nucleic acid. In some methods, at least one of the two oligonucleotides used as a primer for the internal control may include at least one base or nucleotide other than A, C, G, T and U. For example, the nucleotide may include iso-cytosine and/or iso-guanine ("iC" and/or "iG," respectively). In some methods,

at least one of the oligonucleotides used as a primer for the internal control may include a second label. Suitable labels may include fluorophores and quenchers. In other methods, the reaction mixture may include a nucleotide covalently linked to a second quencher, which may be the same or different as the first quencher. For example, the reaction mixture may include a non-natural nucleotide (*e.g.*, a nucleotide having iC or iG as a base) covalently linked to a quencher.

[0031] In some methods, the first and second labels may be different. In some methods the first and second quencher may be different and may be capable of quenching two different fluorophores. In other methods, the first and second quenchers may be the same and may be capable of quenching two different fluorophores.

[0032] The methods and kits described herein also relate to detecting the presence or absence of one or more mutations in a nucleic acid sample, such as HIV. Some of these mutations may confer drug-resistance in HIV-infected mammals. By way of example, but not by way of limitation, drug resistance may include resistance to one or more of the following: azidothymidine (“AZT”), didanosine (“DDI”), tenofovir (“TDF”), amdoxovir (“DAPD”), lamivudine (“3TC”), emtricitabine (“FTC”), zalcitabine (“DOC”), saquinavir, nelfinavir, aprenavir, non-nucleoside reverse transcriptase inhibitors, multi-drug resistance, and a combination thereof. The methods and kits may include multiplex assays that are capable of detecting wild-type HIV nucleic acid and HIV nucleic acid having one or more mutations (*e.g.*, mutations in HIV reverse transcriptase nucleic acid and/or mutations in HIV protease nucleic which, in some embodiments, may correspond with drug resistance).

[0033] In some aspects, the methods and kits may be used to detect HIV nucleic acid. One method for detecting HIV nucleic acid and for detecting the presence or absence of one or more mutations in HIV nucleic acid may include reacting a mixture which includes: 1) HIV nucleic acid isolated from a sample, and 2) at least one oligonucleotide that is capable of specifically hybridizing to the viral nucleic acid, where the oligonucleotide includes at least one non-natural base. In some methods, the at least one oligonucleotide may specifically hybridize to HIV wild-type or HIV mutant nucleic acid sequence.

[0034] Other methods for detecting HIV may include reacting a mixture including HIV nucleic acid isolated from a sample, a control nucleic acid, and at least two pairs of oligonucleotides. In some methods, the first pair of oligonucleotides may be capable of hybridizing to the viral nucleic acid, and the second pair of oligonucleotides may be capable of hybridizing to the control nucleic acid, and at least one oligonucleotide of each pair of

oligonucleotides may include a label that is different from the label of the other oligonucleotide pairs, and at least one oligonucleotide of each pair of oligonucleotides may include at least one non-natural nucleotide. In another method, the viral nucleic acid and the control nucleic acid may be amplified and detected. In still another method, kits may be provided for the detection of HIV infection in a mammal.

[0035] The methods may be used to detect a viral agents such as HIV or a virus that is capable of causing AIDS or AIDS-like symptoms. For example, the viral agent may include HIV having the genomic sequence provided as GenBank Accession No. M19921, or natural or artificial variants thereof. For example, a natural or artificial variant may include a virus having at least about 95% genomic sequence identity to the genomic sequence, or regions of the genomic sequence (*e.g.*, the pol region, the reverse transcriptase sequence or the protease sequence) deposited as GenBank Accession No. M19921. A natural or artificial variant may include a virus whose genome, or regions of the genome (*e.g.*, the pol region, the reverse transcriptase sequence or the protease sequence) hybridizes to the genomic sequence deposited as GenBank Accession No. M19921 under stringent conditions.

[0036] For example, some embodiments of the methods may utilize at least one oligonucleotide including at least one non-natural nucleotide that is capable of hybridizing to the genomic sequence of HIV (*e.g.*, under stringent conditions). The oligonucleotide may be capable of specifically hybridizing to sequences in the HIV pol region, for example, the oligonucleotide may hybridize to the HIV reverse transcriptase sequence, or the HIV protease sequence. In some methods, the oligonucleotides of the present methods may be capable of hybridizing to a HIV nucleotide sequence that encodes the sequences represented by SEQ ID NO. 1. In other methods, the oligonucleotide may be capable of specifically hybridizing to a natural or artificial variant of the nucleotide sequence that encodes SEQ ID NO 1.

[0037] In some embodiments, the at least one oligonucleotide including at least one non-natural nucleotide may be capable of specifically hybridizing to mutant genomic sequences of HIV (*e.g.*, under stringent conditions). In some embodiments, the one or more mutations may be in the pol region of the HIV genome. For example, the one or more mutants may be in the reverse transcriptase gene, the protease gene or both. In other methods, the one or more mutations may include M41L, T215Y, T215F, K65R, L74V, T69D, E44K, V118I, M184V, M184I, L100I, K103N, Y181C, Y181I, Y188L, M46I, L90M, G48V, D30N, and

I50V. In some methods, at least one of the oligonucleotides may be SEQ ID NO: 4, 7, 10, 13, 16, 19, 22, 25, 28, 31, 34, 37, 40, 43, 46, 49, 52, 55, 58, 61, 64, 67, 70, 73 and 74. In some embodiments, the presence or absence of one or more mutations is determined relative to the HIV-1 encoded by Genbank Accession No. M19921.

[0038] Some methods for determining the presence or absence of one or more mutations in HIV may include reacting a reaction mixture which includes HIV nucleic acid isolated from a sample and a first oligonucleotide containing at least one non-natural nucleotide which may be capable of detecting a wild-type HIV sequence at a first location, and a second oligonucleotide containing at least one non-natural nucleotide which may be capable of detecting a mutant HIV sequence at the first location. In some methods, the oligonucleotide pairs may include SEQ ID NO: 3, 4; 6, 7; 9, 10; 12, 13; 15, 16; 18, 19; 21, 22; 24, 25; 27, 28; 30, 31; 33, 34; 36, 37; 39, 40; 42, 43; 45, 46; 48, 49; 51, 52; 54, 55; 57, 58; 60, 61; 63, 64; 66, 67; 69, 70; 72, 73; and degenerate oligonucleotide pairs, for example, oligonucleotides modeled from oligonucleotide SEQ ID NO: 74. In some methods, each oligonucleotide of the oligonucleotide pair may include at least one label. In some methods, the labels may be different.

[0039] In some methods, the reaction mixture may include at least two oligonucleotides that are capable of hybridizing to the HIV nucleic acid and that may function as primers for the amplification of HIV nucleic acid. In some methods, at least one of the two oligonucleotides is used as a primer and includes at least one base or nucleotide other than A, C, G, T and U; the base or nucleotide may include iC or iG (*e.g.*, diCTP or diGTP). In some methods, at least one of the oligonucleotides used as a primer may include a first label. Suitable labels may include fluorophores and quenchers. In some methods, the reaction mixture may include a nucleotide (*e.g.* a non-natural nucleotide) covalently linked to a first quencher. The non-natural nucleotide may include a non-natural base such as iC or iG.

[0040] In some embodiments, the methods may include performing reverse transcription of target RNA, for example, HIV RNA. In other embodiments, the methods may include performing an amplification (*e.g.*, PCR which may include RT-PCR). The methods may include hybridizing a probe to an amplified nucleic acid to detect an amplified target. For example, the methods may include performing RT-PCR followed by performing probe hybridization.

[0041] The methods described herein may include determining a melting temperature for an amplicon (*e.g.*, amplified nucleic acid of at least one of amplified nucleic acid of HIV and amplified control nucleic acid). The methods may include determining a melting temperature for a nucleic acid complex that includes a labeled probe hybridized to a target nucleic acid (which may include amplified target nucleic acid). The melting temperature may be determined by exposing the amplicon or nucleic acid complex to a gradient of temperatures and observing a signal from a label. Optionally, the melting temperature may be determined by (a) reacting an amplicon with an intercalating agent at a gradient of temperatures and (b) observing a detectable signal from the intercalating agent. The melting temperature of a nucleic acid complex may be determined by (1) hybridizing a probe to a target nucleic acid to form a nucleic acid complex, where at least one of the probe and the target nucleic acid includes a label; (2) exposing the nucleic acid complex to a gradient of temperatures; and (3) observing a signal from the label.

[0042] The methods may be performed in any suitable reaction chamber under any suitable conditions. For example, the methods may be performed in a reaction chamber without opening the reaction chamber. The reaction chamber may be part of an array of reaction chambers. In some embodiments, the steps of the methods may be performed separately in different reaction chambers.

[0043] In some aspect, the methods may utilize and/or the kits may include a first pair of oligonucleotides capable of hybridizing to a HIV nucleic acid. In some embodiments, at least one oligonucleotide of the first pair may include at least one non-natural nucleotide; in other embodiments, the oligonucleotide may also include at least one label. In further embodiments, a kit may also include control nucleic acid and a second pair of oligonucleotides capable of hybridizing to the control nucleic acid. In some embodiments, at least one oligonucleotide of the second pair may include at least one non-natural nucleotide and a second label. In some embodiments, the first and second label may be different.

[0044] In some embodiments, the methods may be capable of detecting no more than about 100 copies of the target nucleic acid in a sample (*e.g.*, in a sample having a volume of about 25 microliters). In other embodiments, the methods may be capable of detecting no more than about 500 copies, 1000 copies, 5000 copies, or 10,000 copies in a sample (*e.g.*, in a sample having a volume of about 25 microliters).

[0045] In other embodiments, the methods may be capable of detecting no more than about 100 copies of target nucleic acid in a sample (*e.g.*, in a sample having a volume of about 25 microliters) using real-time detection in no more than about 150 cycles of the PCR, no more than about 100 cycles, no more than about 90 cycles, no more than about 80 cycles, no more than about 70 cycles, no more than about 60 cycles, no more than about 50 cycles, no more than about 40 cycles, or no more than about 30 cycles of the PCR.

[0046] Typically, the methods may be capable of detecting a nucleic acid target such as a DNA or an RNA target in a sample, when the target represents about 1%-10% of the total molecules present in the sample. In some embodiments, the methods may be capable of detecting a nucleic acid target such as a DNA or an RNA target in a sample, when the target represents about 0.5% - 5% of the total molecules present in the sample (or about 0.1% to about 1%).

[0047] In some embodiments, the methods and kits related to detecting a target that includes a mutation. Detected mutations may include a single-base change, a deletion, insertion and/or rearrangement.

BRIEF DESCRIPTION OF THE DRAWINGS

[0048] FIG. 1 is an exemplary amino acid sequence of an HIV-1 pol polyprotein, Genbank Accession No. M19921, (SEQ ID NO:1).

[0049] FIG. 2 is a schematic representation of an exemplary HIV detection method ("MultiCode-RTx" genotyping and detection schematic) employing at least one oligonucleotide with at least one non-natural base. A. In this example, cDNA targets are amplified in the presence of iGTP-dabcyl (Q-iGTP), one standard reverse and two RTx forward primers. The two forward primers are bipartite. In this example, the 5'-parts contain single iCs, separable fluorescent reporters and target independent sequence tails that add 10°C to the annealing temperature of each primer. The 3'-parts are target-specific, contain a 3' mutation-specific base (A or G) and have an annealing temperature of 50°C to the cDNA targets. B. A single round of competitive priming at 50°C results in specific extension (gray arrow) creating the A:T target. C. Remaining cycles at 60°C inhibit annealing of the G-primer and place quenchers in close proximity to the reporter.

[0050] FIG. 3A and 3B show ΔC_t vs. fraction mutant plots for real-time PCR runs with 4 different cycling conditions and 2 different instruments.

[0051] FIG. 4 shows ΔC_t analysis of different total concentration fraction series.

[0052] FIGS. 5 and 6 show a ΔC_t comparison (“dCt”) of different mutant and wild-type M184 primer pairs having 0, 1 or 2 iso-Gs in the sequence. In FIGURE 5, the black bars represent samples tested with 100% wild-type target (M184M); the dark grey bars represents samples tested with 50% wild-type and 50% mutant target; the light gray bars represent samples tested with 100% mutant target (M184V). In FIGURE 6, the number of iso-Gs in the wild-type primer (“wt”) is on the top row of the X axis, the number of iso-Gs in the mutant primer (“mt”) is on the bottom row of the Y axis. In FIGURE 6, the black bars represent samples tested with 100% wild-type target (M184M); the dark grey bars indicate samples tested with 10% mutant target (M184V); and the light grey bars indicate samples tested with 100% mutant target (M184V).

[0053] FIG. 7 shows a schematic representation of the “healing primer” strategy.

DETAILED DESCRIPTION

[0054] Disclosed are methods and kits for detecting nucleic acids in a sample. Typically, the methods include detecting signals such as a signal emitted from a fluorophore. Also disclosed are oligonucleotides, especially primers and probes, which may be used for the detection of viruses capable of causing AIDS or AIDS-like symptoms. The methods, kits, and oligonucleotides disclosed herein may be used to detect HIV, which has been shown to be the causative agent of AIDS in humans. Additionally, some methods may be based on assay methods described in published international application WO 01/90417, U.S. published application 2002/0150900, and Moser *et al.*, Antimicrobial Agents and Chemotherapy, 2005, 49(8):3334-35.

[0055] As used herein, unless otherwise stated, the singular forms “a,” “an,” and “the” includes plural reference. Thus, for example, a reference to “an oligonucleotide” includes a plurality of oligonucleotide molecules, and a reference to “a nucleic acid” is a reference to one or more nucleic acids.

[0056] As used herein, the term “sample” is used in its broadest sense. A sample may include a bodily tissue or a bodily fluid including but not limited to blood (or a fraction of blood such as plasma or serum), lymph, mucus, tears, urine, and saliva. A sample may include an extract from a cell, a chromosome, organelle, or a virus. A sample may comprise DNA (*e.g.*, genomic DNA), RNA (*e.g.*, mRNA), and cDNA, any of which may be amplified to provide amplified nucleic acid. A sample may include nucleic acid in solution or bound to a substrate (*e.g.*, as part of a microarray). A sample may comprise material obtained

from an environmental locus (*e.g.*, a body of water, soil, and the like) or material obtained from a fomite (*i.e.*, an inanimate object that serves to transfer pathogens from one host to another).

[0057] The term "source of nucleic acid" refers to any sample which contains nucleic acids (RNA or DNA). Particularly preferred sources of target nucleic acids are biological samples including, but not limited to blood, plasma, serum, saliva, cerebral spinal fluid, pleural fluid, milk, lymph, sputum and semen.

[0058] As used herein, the term "limit of detection" refers to the lowest level or amount of an analyte, such as a nucleic acid, that can be detected and quantified. Limits of detection can be represented as molar values (*e.g.*, 2.0 nM limit of detection), as gram measured values (*e.g.*, 2.0 microgram limit of detection under, for example, specified reaction conditions), copy number (*e.g.*, 1×10^5 copy number limit of detection), or other representations known in the art.

[0059] As used herein, "HIV" is meant to include any retrovirus that is capable of infecting a mammal and that has been shown to be the causative agent of Acquired Immunodeficiency Syndrome ("AIDS") or AIDS-like symptoms. The terms "HIV-1" or "HIV-2" are meant to include all types and sub-types of HIV-1 and HIV-2. That is, if the term HIV-1 is used, it is meant to encompass all types (*e.g.*, M, N and O) and sub-types (*e.g.*, A-K), and natural and artificial variants of these type and subtypes. By way of example, but not by way of limitation, an exemplary HIV sequence is provided by Genbank Accession No. M19921. Natural and artificial variants of this sequence are contemplated in the methods and kits, and may include, for example, silent mutations and mutations that confer drug resistance.

[0060] As used herein, the term "pol region" or "pol gene" means that region of the HIV genome which includes the coding sequence for at least a reverse transcriptase and a protease. By way of example, but not by way of limitation, an exemplary pol region polyprotein amino acid sequence is shown in FIGURE 1, SEQ ID NO: 1. In some embodiments, the sequence of the detected HIV may have about 95% sequence identity to SEQ ID NO 1. In other embodiments, the sequence of the detected HIV may have about 90% sequence identity to SEQ ID NO: 1. In still other embodiments, the sequence of the detected HIV may have about 80% sequence identity to SEQ ID NO: 1.

[0061] As used herein the term "isolated" in reference to a nucleic acid molecule refers to a nucleic acid molecule which is separated from the organisms and biological materials

(e.g., blood, cells, serum, plasma, saliva, urine, stool, sputum, nasopharyngeal aspirates and so forth), which are present in the natural source of the nucleic acid molecule. An isolated nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. In some embodiments, nucleic acid molecules encoding polypeptides/proteins may also be isolated or purified. Methods of nucleic acid isolation are well known in the art and may include total nucleic acid isolation/purification methods, RNA-specific isolation/purification methods or DNA-specific isolation/purification methods.

[0062] As used herein, the term “isolated virus” refers to a virus which is separated from other organisms and biological materials which are present in the natural source of the virus, e.g., biological material such as cells, blood, serum, plasma, saliva, urine, stool, sputum, nasopharyngeal aspirates, and so forth. The isolated virus can be used to infect a subject.

[0063] As used herein, the term “subject” is refers to an animal, preferably a mammal, more preferably a human. The term “subject” and “patient” may be used interchangeably.

[0064] A “mutation” is meant to encompass at least a single nucleotide variation in a nucleic acid sequence relative to the normal sequence or wild-type sequence. A mutation may include a substitution, a deletion, an inversion or an insertion. With respect to an encoded polypeptide, a mutation may be “silent” and result in no change in the encoded polypeptide sequence or a mutation may result in a change in the encoded polypeptide sequence. For example, a mutation may result in a substitution in the encoded polypeptide sequence. A mutation may result in a frameshift with respect to the encoded polypeptide sequence. For example, the HIV virus of the present methods may be mutants as compared to the HIV virus nucleic acid of Genbank Accession No. M19921 or other known HIV virus strains.

[0065] Some mutations may confer drug resistance in HIV infected mammals, subjects or patients. As used herein the term “drug resistant” means the ability of an infectious agent to withstand a drug to which it was once sensitive and was either slowed in growth and proliferation or killed outright. By way of example, but not by way of limitation, drug resistance with respect to HIV infection may include resistance to one or more of the following: azidothymidine, didanosine, tenofovir, amdoxovir, lamivudine, emtricitabine, zalcitabine, saquinavir, nelfinavir, aprenavir, non-nucleoside reverse transcriptase inhibitors, multi-drug resistance, and a combination thereof.

[0066] Mutations, and drug resistance mutations, may be in any region of the HIV genome. For example, mutations may be in the pol region, and may include mutations in the reverse transcriptase gene and/or the protease gene. By way of example, but not by way of limitation, some exemplary mutations may include M41L, T215Y, T215F, K65R, L74V, T69D, E44K, V118I, M184V, M184I, L100I, K103N, Y181C, Y181I, Y188L, M46I, L90M, G48V, D30N, and I50V.

[0067] As used herein, the term “microarray” refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate. The terms “element” and “array element” refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

[0068] As used herein, an oligonucleotide is understood to be a molecule that has a sequence of bases on a backbone comprised mainly of identical monomer units at defined intervals. The bases are arranged on the backbone in such a way that they can enter into a bond with a nucleic acid having a sequence of bases that are complementary to the bases of the oligonucleotide. The most common oligonucleotides have a backbone of sugar phosphate units. A distinction may be made between oligodeoxyribonucleotides, made up of “dNTPs,” which do not have a hydroxyl group at the 2' position, and oligoribonucleotides, made up of “NTPs,” which have a hydroxyl group in this position. Oligonucleotides also may include derivatives, in which the hydrogen of the hydroxyl group is replaced with organic groups, *e.g.*, an allyl group.

[0069] In some embodiments, oligonucleotides as described herein may include a peptide backbone. For example, the oligonucleotides may include peptide nucleic acids or “PNA.” Peptide nucleic acids are described in WO 92/20702.

[0070] An oligonucleotide is a nucleic acid that includes at least two nucleotides. Oligonucleotides used in the methods disclosed herein typically include at least about ten (10) nucleotides and more typically at least about fifteen (15) nucleotides. Preferred oligonucleotides for the methods disclosed herein include about 10-25 nucleotides. An oligonucleotide may be designed to function as a “primer.” A “primer” is a short nucleic acid, usually a ssDNA oligonucleotide, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA or RNA strand by a polymerase enzyme, such as a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence (*e.g.*, by the

polymerase chain reaction (PCR)). An oligonucleotide may be designed to function as a “probe.” A “probe” refers to an oligonucleotide, its complements, or fragments thereof, which is used to detect identical, allelic or related nucleic acid sequences. Probes may include oligonucleotides which have been attached to a detectable label or reporter molecule. Typical labels include fluorescent dyes, quenchers, radioactive isotopes, ligands, scintillation agents, chemiluminescent agents, and enzymes.

[0071] An oligonucleotide may be designed to be specific for a target nucleic acid sequence in a sample. For example, an oligonucleotide may be designed to include “antisense” nucleic acid sequence of the target nucleic acid. As used herein, the term “antisense” refers to any composition capable of base-pairing with the “sense” (coding) strand of a specific target nucleic acid sequence. An antisense nucleic acid sequence may be “complementary” to a target nucleic acid sequence. As used herein, “complementarity” describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'. In some embodiments, primers or probes may be designed to include mismatches at various positions. As used herein, a “mismatch” means a nucleotide pair that does not include the standard Watson-Crick base pairs, or nucleotide pairs that do not preferentially form hydrogen bonds. The mismatch may include a natural nucleotide or a non-natural nucleotide substituted across from a particular base or bases in a target. For example, the probe or primer sequence 5'-AGT-3' has a single mismatch with the target sequence 3'-ACA-5'. The 5' “A” of the probe or primer is mismatched with the 3' “A” of the target. Similarly, the target sequence 5'-AGT-3' has a single mismatch with the probe or primer sequence 3'-(iC)CT-5'. Here an iso-C is substituted in place of the natural “T.” However, the sequence 3'-(iC)CT-5' is not mismatched with the sequence 5'-(iG)GA-3'.

[0072] Oligonucleotides may also be designed as degenerate oligonucleotides. As used herein “degenerate oligonucleotide” is meant to include a population, pool, or plurality of oligonucleotides comprising a mixture of different sequences where the sequence differences occur at a specified position in each oligonucleotide of the population. For example degenerate oligonucleotides may be represented as

GACATRGTYATCTATCARTAYR, where, for example R = A or G, Y = C or T.

Accordingly, the sequence of the population of oligonucleotides would be identical except for difference introduced at positions represented by “R” and “Y.” The various substitutions may include any natural or non-natural nucleotide, and may include any

number of different possible nucleotides at any given position. For example the above degenerate oligonucleotide may instead include **R** = iC or iG, or **R** = A or G or T or C or iC or iG.

[0073] Oligonucleotides as described herein typically are capable of forming hydrogen bonds with oligonucleotides having a complementary base sequence. These bases may include the natural bases such as A, G, C, T and U, as well as artificial, non-standard or non-natural bases such as iso-cytosine and iso-guanine. As described herein, a first sequence of an oligonucleotide is described as being 100% complementary with a second sequence of an oligonucleotide when the consecutive bases of the first sequence (read 5'→3') follow the Watson-Crick rule of base pairing as compared to the consecutive bases of the second sequence (read 3'→5'). An oligonucleotide may include nucleotide substitutions. For example, an artificial base may be used in place of a natural base such that the artificial base exhibits a specific interaction that is similar to the natural base.

[0074] An oligonucleotide that is specific for a target nucleic acid also may be specific for a nucleic acid sequence that has "homology" to the target nucleic acid sequence. As used herein, "homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences. The terms "percent identity" and "% identity" as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm (*e.g.*, BLAST).

[0075] An oligonucleotide that is specific for a target nucleic acid will "hybridize" to the target nucleic acid under suitable conditions. As used herein, "hybridization" or "hybridizing" refers to the process by which an oligonucleotide single strand anneals with a complementary strand through base pairing under defined hybridization conditions. "Specific hybridization" is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after any subsequent washing steps. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may occur, for example, at 65°C in the presence of about 6×SSC. Stringency of hybridization may be expressed, in part, with reference to the temperature under which the wash steps are carried out. Such temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined

ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Equations for calculating T_m , for example nearest-neighbor parameters, and conditions for nucleic acid hybridization are known in the art.

[0076] As used herein “target” or “target nucleic acid” refers to a nucleic acid molecule containing a sequence that has at least partial complementarity with an oligonucleotide, for example a probe or a primer. A “target” sequence may include a part of a gene or genome.

[0077] As used herein, “nucleic acid,” “nucleotide sequence,” or “nucleic acid sequence” refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof and to naturally occurring or synthetic molecules. These terms also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, or to any DNA-like or RNA-like material. An “RNA equivalent,” in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose. RNA may be used in the methods described herein and/or may be converted to cDNA by reverse-transcription for use in the methods described herein.

[0078] As used herein, “amplification” or “amplifying” refers to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies known in the art. The term “amplification reaction system” refers to any *in vitro* means for multiplying the copies of a target sequence of nucleic acid. The term “amplification reaction mixture” refers to an aqueous solution comprising the various reagents used to amplify a target nucleic acid. These may include enzymes (*e.g.*, a thermostable polymerase), aqueous buffers, salts, amplification primers, target nucleic acid, and nucleoside triphosphates, and optionally at least one labeled probe and/or optionally at least one agent for determining the melting temperature of an amplified target nucleic acid (*e.g.*, a fluorescent intercalating agent that exhibits a change in fluorescence in the presence of double-stranded nucleic acid).

[0079] The amplification methods described herein may include “real-time monitoring” or “continuous monitoring.” These terms refer to monitoring multiple times during a cycle of PCR, preferably during temperature transitions, and more preferably obtaining at least one data point in each temperature transition. The term “homogeneous detection assay” is used

to describe an assay that includes coupled amplification and detection, which may include "real-time monitoring" or "continuous monitoring."

[0080] Amplification of nucleic acids may include amplification of nucleic acids or subregions of these nucleic acids. For example, amplification may include amplifying portions of nucleic acids between 30 and 50, between 50 and 100, between 100 and 300 bases long by selecting the proper primer sequences and using the PCR.


[0081] The disclosed methods may include amplifying at least one or more nucleic acids in the sample. In the disclosed methods, amplification may be monitored using real-time methods.

[0082] Amplification mixtures may include natural nucleotides (including A, C, G, T, and U) and non-natural nucleotides (*e.g.*, including iC and iG). DNA and RNA are oligonucleotides that include deoxyriboses or riboses, respectively, coupled by phosphodiester bonds. Each deoxyribose or ribose includes a base coupled to a sugar. The bases incorporated in naturally-occurring DNA and RNA are adenosine (A), guanosine (G), thymidine (T), cytosine (C), and uridine (U). These five bases are "natural bases." According to the rules of base pairing elaborated by Watson and Crick, the natural bases can hybridize to form purine-pyrimidine base pairs, where G pairs with C and A pairs with T or U. These pairing rules facilitate specific hybridization of an oligonucleotide with a complementary oligonucleotide.

[0083] The formation of these base pairs by the natural bases is facilitated by the generation of two or three hydrogen bonds between the two bases of each base pair. Each of the bases includes two or three hydrogen bond donor(s) and hydrogen bond acceptor(s). The hydrogen bonds of the base pair are each formed by the interaction of at least one hydrogen bond donor on one base with a hydrogen bond acceptor on the other base. Hydrogen bond donors include, for example, heteroatoms (*e.g.*, oxygen or nitrogen) that have at least one attached hydrogen. Hydrogen bond acceptors include, for example, heteroatoms (*e.g.*, oxygen or nitrogen) that have a lone pair of electrons.

[0084] The natural or non-natural bases used herein can be derivatized by substitution at non-hydrogen bonding sites to form modified natural or non-natural bases. For example, a natural base can be derivatized for attachment to a support by coupling a reactive functional group (for example, thiol, hydrazine, alcohol, amine, and the like) to a non-hydrogen bonding atom of the base. Other possible substituents include, for example, biotin, digoxigenin, fluorescent groups, alkyl groups (*e.g.*, methyl or ethyl), and the like.

[0085] Non-natural bases, which form hydrogen-bonding base pairs, can also be constructed as described, for example, in U.S. Patents Nos. 5,432,272; 5,965,364; 6,001,983; 6,037,120; U.S. published Application No. 2002/0150900; and U.S. Patent No. 6,140,496. Suitable bases and their corresponding base pairs may include the following bases in base pair combinations (iso C/iso G, K/X, H/J, and M/N):



[0086] Where A is the point of attachment to the sugar or other portion of the polymeric backbone and R is H or a substituted or unsubstituted alkyl group. It will be recognized that other non-natural bases utilizing hydrogen bonding can be prepared, as well as modifications of the above-identified non-natural bases by incorporation of functional groups at the non-hydrogen bonding atoms of the bases.

[0087] The hydrogen bonding of these non-natural base pairs is similar to those of the natural bases where two or three hydrogen bonds are formed between hydrogen bond acceptors and hydrogen bond donors of the pairing non-natural bases. One of the differences between the natural bases and these non-natural bases is the number and

position of hydrogen bond acceptors and hydrogen bond donors. For example, cytosine can be considered a donor/acceptor/acceptor base with guanine being the complementary acceptor/donor/donor base. Iso-C is an acceptor/acceptor/donor base and iso-G is the complementary donor/donor/acceptor base, as illustrated in U.S. Patent No. 6,037,120.

[0088] Other non-natural bases for use in oligonucleotides include, for example, naphthalene, phenanthrene, and pyrene derivatives as discussed, for example, in Ren, *et al.*, J. Am. Chem. Soc. 1996, 118:1671 and McMinn *et al.*, J. Am. Chem. Soc. 1999, 121:11585. These bases do not utilize hydrogen bonding for stabilization, but instead rely on hydrophobic or van der Waals interactions to form base pairs.

[0089] The use of non-natural bases according to the methods disclosed herein is extendable beyond the detection and quantification of nucleic acid sequences present in a sample. For example, non-natural bases can be recognized by many enzymes that catalyze reactions associated with nucleic acids. While a polymerase requires a complementary nucleotide to continue polymerizing an extending oligonucleotide chain, other enzymes do not require a complementary nucleotide. If a non-natural base is present in the template and its complementary non-natural base is not present in the reaction mix, a polymerase will typically stall (or, in some instances, misincorporate a base when given a sufficient amount of time) when attempting to extend an elongating primer past the non-natural base.

However, other enzymes that catalyze reactions associated with nucleic acids, such as ligases, kinases, nucleases, polymerases, topoisomerases, helicases, and the like can catalyze reactions involving non-natural bases. Such features of non-natural bases can be taken advantage of, and are within the scope of the presently disclosed methods and kits.

[0090] For example, non-natural bases can be used to generate duplexed nucleic acid sequences having a single strand overhang. This can be accomplished by performing a PCR reaction to detect a target nucleic acid in a sample, the target nucleic acid having a first portion and a second portion, where the reaction system includes all four naturally occurring dNTP's, a first primer that is complementary to the first portion of the target nucleic acid, a second primer having a first region and a second region, the first region being complementary to the first portion of the target nucleic acid, and the second region being noncomplementary to the target nucleic acid. The second region of the second primer comprises a non-natural base. The first primer and the first region of the second primer

hybridize to the target nucleic acid, if present. Several rounds of PCR will produce an amplification product containing (i) a double-stranded region and (ii) a single-stranded region. The double-stranded region is formed through extension of the first and second primers during PCR. The single-stranded region includes the one or more non-natural bases. The single-stranded region of the amplification product results because the polymerase is not able to form an extension product by polymerization beyond the non-natural base in the absence of the nucleotide triphosphate of the complementary non-natural base. In this way, the non-natural base functions to maintain a single-stranded region of the amplification product.

[0091] A polymerase can, in some instances, misincorporate a base opposite a non-natural base. In some embodiments, the misincorporation takes place because the reaction mix does not include a complementary non-natural base. Therefore, if given sufficient amount of time, the polymerase can, in some cases, misincorporate a base that is present in the reaction mixture opposite the non-natural base.

[0092] The nucleotides disclosed herein, which may include non-natural nucleotides, may be coupled to a label (*e.g.*, a quencher or a fluorophore). Coupling may be performed using methods known in the art.

[0093] The oligonucleotides of the present methods may function as primers. In some embodiments, the oligonucleotides are labeled. For example, the oligonucleotides may be labeled with a reporter that emits a detectable signal (*e.g.*, a fluorophore). The oligonucleotides may include at least one non-natural nucleotide. For example, the oligonucleotides may include at least one nucleotide having a base that is not A, C, G, T, or U (*e.g.*, iC or iG). Where the oligonucleotide is used as a primer for PCR, the amplification mixture may include at least one nucleotide that is labeled with a quencher (*e.g.*, Dabcyl). The labeled nucleotide may include at least one non-natural nucleotide. For example, the labeled nucleotide may include at least one nucleotide having a base that is not A, C, G, T, or U (*e.g.*, iC or iG).

[0094] In some embodiments, the oligonucleotide may be designed not to form an intramolecular structure such as a hairpin. In other embodiments, the oligonucleotide may be designed to form an intramolecular structure such as a hairpin. For example, the oligonucleotide may be designed to form a hairpin structure that is altered after the oligonucleotide hybridizes to a target nucleic acid, and optionally, after the target nucleic acid is amplified using the oligonucleotide as a primer.

[0095] The oligonucleotide may be labeled with a fluorophore that exhibits quenching when incorporated in an amplified product as a primer. In other embodiments, the oligonucleotide may emit a detectable signal after the oligonucleotide is incorporated in an amplified product as a primer (*e.g.*, inherently, or by fluorescence induction or fluorescence dequenching). Such primers are known in the art (*e.g.*, LightCycler primers, Amplifluor® Primers, Scorpion® Primers and Lux™ Primers). The fluorophore used to label the oligonucleotide may emit a signal when intercalated in double-stranded nucleic acid. As such, the fluorophore may emit a signal after the oligonucleotide is used as a primer for amplifying the nucleic acid.

[0096] The oligonucleotides that are used in the disclosed methods may be suitable as primers for amplifying at least one nucleic acid in the sample and as probes for detecting at least one nucleic acid in the sample. In some embodiments, the oligonucleotides are labeled with at least one fluorescent dye, which may produce a detectable signal. The fluorescent dye may function as a fluorescence donor for fluorescence resonance energy transfer (FRET). The detectable signal may be quenched when the oligonucleotide is used to amplify a target nucleic acid. For example, the amplification mixture may include nucleotides that are labeled with a quencher for the detectable signal emitted by the fluorophore. Optionally, the oligonucleotides may be labeled with a second fluorescent dye or a quencher dye that may function as a fluorescence acceptor (*e.g.*, for FRET). Where the oligonucleotide is labeled with a first fluorescent dye and a second fluorescent dye, a signal may be detected from the first fluorescent dye, the second fluorescent dye, or both. Signals may be detected at a gradient of temperatures (*e.g.*, in order to determine a melting temperature for an amplicon or a complex that includes a probe hybridized to a target nucleic acid).

[0097] The disclosed methods may be performed with any suitable number of oligonucleotides. Where a plurality of oligonucleotides are used (*e.g.*, two or more oligonucleotides), different oligonucleotide may be labeled with different fluorescent dyes capable of producing a detectable signal. In some embodiments, oligonucleotides are labeled with at least one of two different fluorescent dyes. In further embodiments, oligonucleotides are labeled with at least one of three different fluorescent dyes.

[0098] In some embodiments, each different fluorescent dye emits a signal that can be distinguished from a signal emitted by any other of the different fluorescent dyes that are used to label the oligonucleotides. For example, the different fluorescent dyes may have

wavelength emission maximums all of which differ from each other by at least about 5 nm (preferably by least about 10 nm). In some embodiments, each different fluorescent dye is excited by different wavelength energies. For example, the different fluorescent dyes may have wavelength absorption maximums all of which differ from each other by at least about 5 nm (preferably by at least about 10 nm).

[0099] Where a fluorescent dye is used to determine the melting temperature of a nucleic acid in the method, the fluorescent dye may emit a signal that can be distinguished from a signal emitted by any other of the different fluorescent dyes that are used to label the oligonucleotides. For example, the fluorescent dye for determining the melting temperature of a nucleic acid may have a wavelength emission maximum that differs from the wavelength emission maximum of any other fluorescent dye that is used for labeling an oligonucleotide by at least about 5 nm (preferably by least about 10 nm). In some embodiments, the fluorescent dye for determining the melting temperature of a nucleic acid may be excited by different wavelength energy than any other of the different fluorescent dyes that are used to label the oligonucleotides. For example, the fluorescent dye for determining the melting temperature of a nucleic acid may have a wavelength absorption maximum that differs from the wavelength absorption maximum of any fluorescent dye that is used for labeling an oligonucleotide by at least about 5 nm (preferably by least about 10 nm).

[0100] The methods may include determining the melting temperature of at least one nucleic acid in a sample (e.g., an amplicon or a nucleic acid complex that includes a probe hybridized to a target nucleic acid), which may be used to identify the nucleic acid. Determining the melting temperature may include exposing an amplicon or a nucleic acid complex to a temperature gradient and observing a detectable signal from a fluorophore. Optionally, where the oligonucleotides of the method are labeled with a first fluorescent dye, determining the melting temperature of the detected nucleic acid may include observing a signal from a second fluorescent dye that is different from the first fluorescent dye. In some embodiments, the second fluorescent dye for determining the melting temperature of the detected nucleic acid is an intercalating agent. Suitable intercalating agents may include, but are not limited to SYBR™ Green 1 dye, SYBR dyes, Pico Green, SYTO dyes, SYTOX dyes, ethidium bromide, ethidium homodimer-1, ethidium homodimer-2, ethidium derivatives, acridine, acridine orange, acridine derivatives, ethidium-acridine heterodimer, ethidium monoazide, propidium iodide, cyanine monomers,

7-aminoactinomycin D, YOYO-1, TOTO-1, YOYO-3, TOTO-3, POPO-1, BOBO-1, POPO-3, BOBO-3, LOLO-1, JOJO-1, cyanine dimers, YO-PRO-1, TO-PRO-1, YO-PRO-3, TO-PRO-3, TO-PRO-5, PO-PRO-1, BO-PRO-1, PO-PRO-3, BO-PRO-3, LO-PRO-1, JO-PRO-1, and mixture thereof. In suitable embodiments, the selected intercalating agent is SYBRTM Green 1 dye.

[0101] Typically, an intercalating agent used in the method will exhibit a change in fluorescence when intercalated in double-stranded nucleic acid. A change in fluorescence may include an increase in fluorescence intensity or a decrease in fluorescence intensity. For example, the intercalating agent may exhibit an increase in fluorescence when intercalated in double-stranded nucleic acid, and a decrease in fluorescence when the double-stranded nucleic acid is melted. A change in fluorescence may include a shift in fluorescence spectra (*i.e.*, a shift to the left or a shift to the right in maximum absorbance wavelength or maximum emission wavelength). For example, the intercalating agent may emit a fluorescent signal of a first wavelength (*e.g.*, green) when intercalated in double-stranded nucleic acid and emit a fluorescent signal of a second wavelength (*e.g.*, red) when not intercalated in double-stranded nucleic acid. A change in fluorescence of an intercalating agent may be monitored at a gradient of temperatures to determine the melting temperature of the nucleic acid (where the intercalating agent exhibits a change in fluorescence when the nucleic acid melts).

[0102] In the disclosed methods, each of the amplified target nucleic acids may have different melting temperatures. For example, each of the amplified target nucleic acids may have a melting temperature that differs by at least about 1°C, more preferably by at least about 2°C, or even more preferably by at least about 4°C from the melting temperature of any of the other amplified target nucleic acids.

[0103] The methods disclosed herein may include transcription of RNA to DNA (*i.e.*, reverse transcription). For example, reverse transcription may be performed prior to amplification.

[0104] As used herein, “labels” or “reporter molecules” are chemical or biochemical moieties useful for labeling a nucleic acid, amino acid, or antibody. “Labels” and “reporter molecules” include fluorescent agents, chemiluminescent agents, chromogenic agents, quenching agents, radionuclides, enzymes, substrates, cofactors, scintillation agents, inhibitors, magnetic particles, and other moieties known in the art. “Labels” or “reporter

molecules” are capable of generating a measurable signal and may be covalently or noncovalently joined to an oligonucleotide.

[0105] As used herein, a “fluorescent dye” or a “fluorophore” is a chemical group that can be excited by light to emit fluorescence. Some suitable fluorophores may be excited by light to emit phosphorescence. Dyes may include acceptor dyes that are capable of quenching a fluorescent signal from a fluorescent donor dye. Dyes that may be used in the disclosed methods include, but are not limited to, the following dyes and/or dyes sold under the following tradenames: 1,5 IAEDANS; 1,8-ANS; 4-Methylumbelliferone; 5-carboxy-2,7-dichlorofluorescein; 5-Carboxyfluorescein (5-FAM); 5-Carboxytetramethylrhodamine (5-TAMRA) ; 5-FAM (5-Carboxyfluorescein); 5-HAT (Hydroxy Tryptamine) ; 5-Hydroxy Tryptamine (HAT); 5-ROX (carboxy-X-rhodamine); 5-TAMRA (5-Carboxytetramethylrhodamine); 6-Carboxyrhodamine 6G; 6-CR 6G; 6-JOE; 7-Amino-4-methylcoumarin; 7-Aminoactinomycin D (7-AAD); 7-Hydroxy-4-methylcoumarin; 9-Amino-6-chloro-2-methoxyacridine; ABQ; Acid Fuchsin; ACMA (9-Amino-6-chloro-2-methoxyacridine); Acridine Orange; Acridine Red; Acridine Yellow; Acriflavin; Acriflavin Feulgen SITSA; Alexa Fluor 350TM; Alexa Fluor 430TM; Alexa Fluor 488TM; Alexa Fluor 532TM; Alexa Fluor 546TM; Alexa Fluor 568TM; Alexa Fluor 594TM; Alexa Fluor 633TM; Alexa Fluor 647TM; Alexa Fluor 660TM; Alexa Fluor 680TM; Alizarin Complexon; Alizarin Red; Allophycocyanin (APC); AMC; AMCA-S; AMCA (Aminomethylcoumarin); AMCA-X; Aminoactinomycin D; Aminocoumarin; Aminomethylcoumarin (AMCA); Anilin Blue; Anthrocyl stearate; APC (Allophycocyanin); APC-Cy7; APTS; Astrazon Brilliant Red 4G; Astrazon Orange R; Astrazon Red 6B; Astrazon Yellow 7 GLL ; Atabrine; ATTO-TAGTM CBQCA; ATTO-TAGTM FQ; Auramine; Aurophosphine G; Aurophosphine; BAO 9 (Bisaminophenyloxadiazole); Berberine Sulphate; Beta Lactamase; BFP blue shifted GFP (Y66H); Blue Fluorescent Protein; BFP/GFP FRET; Bimane; Bisbenzamide; Bisbenzimidazole (Hoechst); Blancophor FFG; Blancophor SV; BOBOTM -1; BOBOTM -3; Bodipy 492/515; Bodipy 493/503; Bodipy 500/510; Bodipy 505/515; Bodipy 530/550; Bodipy 542/563; Bodipy 558/568; Bodipy 564/570; Bodipy 576/589; Bodipy 581/591; Bodipy 630/650-X; Bodipy 650/665-X; Bodipy 665/676; Bodipy FL; Bodipy FL ATP; Bodipy Fl-Ceramide; Bodipy R6G SE; Bodipy TMR; Bodipy TMR-X conjugate ; Bodipy TMR-X, SE; Bodipy TR; Bodipy TR ATP; Bodipy TR-X SE; BO-PROTM-1; BO-PROTM-3; Brilliant Sulphoflavin FF; Calcein; Calcein Blue ; Calcium CrimsonTM; Calcium Green; Calcium Orange; Calcofluor White; Carboxy-X-rhodamine (5-ROX); Cascade BlueTM; Cascade

Yellow; Catecholamine; CCF2 (GeneBlazer); CFDA; CFP - Cyan Fluorescent Protein; CFP/YFP FRET; Chlorophyll; Chromomycin A; CL-NERF (Ratio Dye, pH); CMFDA; Coelenterazine f; Coelenterazine fcp; Coelenterazine h; Coelenterazine hcp; Coelenterazine ip; Coelenterazine n; Coelenterazine O; Coumarin Phalloidin; C-phycocyanine; CPM Methylcoumarin; CTC; CTC Formazan; Cy2TM; Cy3.1 8; Cy3.5TM; Cy3TM; Cy5.1 8; Cy5.5TM; Cy5TM; Cy7TM; Cyan GFP; cyclic AMP Fluorosensor (FiCRhR); Dabcyl; Dansyl; Dansyl Amine; Dansyl Cadaverine; Dansyl Chloride; Dansyl DHPE; Dansyl fluoride; DAPI; Dapoxyl; Dapoxyl 2; Dapoxyl 3; DCFDA; DCFH (Dichlorodihydrofluorescein Diacetate); DDAO; DHR (Dihydorhodamine 123); Di-4-ANEPPS; Di-8-ANEPPS (non-ratio); DiA (4-Di-16-ASP); Dichlorodihydrofluorescein Diacetate (DCFH); DiD - Lipophilic Tracer; DiD (DiIC18(5)); DIDS ; Dihydorhodamine 123 (DHR); DiI (DiIC18(3)); Dinitrophenol; DiO (DiOC18(3)); DiR; DiR (DiIC18(7)); DNP; Dopamine; DsRed; DTAF; DY-630-NHS; DY-635-NHS; EBFP; ECFP; EGFP; ELF 97; Eosin; Erythrosin; Erythrosin ITC ; Ethidium Bromide; Ethidium homodimer -1 (EthD-1); Euchrysin; EukoLight; Europium (III) chloride; EYFP; Fast Blue; FDA; Feulgen (Pararosaniline); FITC; Flazo Orange; Fluo-3; Fluo-4; Fluorescein (FITC); Fluorescein Diacetate; Fluoro-Emerald; Fluoro-Gold (Hydroxystilbamidine); Fluor-Ruby; FluorX; FM 1-43TM; FM 4-46; Fura RedTM; Fura RedTM/Fluo-3; Fura-2; Fura-2/BCECF; Genacryl Brilliant Red B; Genacryl Brilliant Yellow 10GF; Genacryl Pink 3G; Genacryl Yellow 5GF; GeneBlazer (CCF2); GFP (S65T); GFP red shifted (rsGFP); GFP wild type, non-UV excitation (wtGFP); GFP wild type, UV excitation (wtGFP); GFPuv; Gloxalic Acid ; Granular Blue; Haematoporphyrin; Hoechst 33258; Hoechst 33342; Hoechst 34580; HPTS; Hydroxycoumarin; Hydroxystilbamidine (FluoroGold); Hydroxytryptamine; Indo-1; Indodicarbocyanine (DiD); Indotricarbocyanine (DiR); Intrawhite Cf; JC-1; JO-JO-1; JO-PRO-1; Laurodan; LDS 751 (DNA); LDS 751 (RNA); Leucophor PAF; Leucophor SF; Leucophor WS; Lissamine Rhodamine; Lissamine Rhodamine B ; Calcein/Ethidium homodimer; LOLO-1; LO-PRO-1; Lucifer Yellow; Lyso Tracker Blue; Lyso Tracker Blue-White; Lyso Tracker Green; Lyso Tracker Red; Lyso Tracker Yellow; LysoSensor Blue; LysoSensor Green; LysoSensor Yellow/Blue; Mag Green; Magdala Red (Phloxin B); Mag-Fura Red; Mag-Fura-2; Mag-Fura-5; Mag-Indo-1; Magnesium Green; Magnesium Orange; Malachite Green; Marina Blue; Maxilon Brilliant Flavin 10 GFF; Maxilon Brilliant Flavin 8 GFF; Merocyanin; Methoxycoumarin; Mitotracker Green FM; Mitotracker Orange; Mitotracker Red; Mitramycin ; Monobromobimane; Monobromobimane (mBBBr-GSH);

Monochlorobimane; MPS (Methyl Green Pyronine Stilbene); NBD; NBD Amine; Nile Red; NEDTM; Nitrobenzoxadidole; Noradrenaline; Nuclear Fast Red; Nuclear Yellow; Nylosan Brilliant Iavin E8G; Oregon Green; Oregon Green 488-X; Oregon GreenTM; Oregon GreenTM 488; Oregon GreenTM 500; Oregon GreenTM 514; Pacific Blue; Pararosanine (Feulgen); PBFI; PE-Cy5; PE-Cy7; PerCP; PerCP-Cy5.5; PE-TexasRed [Red 613]; Phloxin B (Magdala Red); Phorwite AR; Phorwite BKL; Phorwite Rev; Phorwite RPA; Phosphine 3R; Phycoerythrin B [PE]; Phycoerythrin R [PE]; PKH26 (Sigma); PKH67; PMIA; Pontochrome Blue Black; POPO-1; POPO-3; PO-PRO-1; PO-PRO-3; Primuline; Procion Yellow; Propidium Iodid (PI); PyMPO; Pyrene; Pyronine; Pyronine B; Pyrozal Brilliant Flavin 7GF; QSY 7; Quinacrine Mustard; Red 613 [PE-TexasRed]; Resorufin; RH 414; Rhod-2; Rhodamine; Rhodamine 110 ; Rhodamine 123; Rhodamine 5 GLD; Rhodamine 6G; Rhodamine B; Rhodamine B 200; Rhodamine B extra; Rhodamine BB; Rhodamine BG; Rhodamine Green; Rhodamine Phallicidine; Rhodamine Phalloidine; Rhodamine Red; Rhodamine WT ; Rose Bengal; R-phycocyanine; R-phycoerythrin (PE); RsGFP; S65A; S65C; S65L; S65T; Sapphire GFP; SBFI; Serotonin; Sevron Brilliant Red 2B; Sevron Brilliant Red 4G; Sevron Brilliant Red B; Sevron Orange; Sevron Yellow L; sgBFPTM; sgBFPTM (super glow BFP); sgGFPTM; sgGFPTM (super glow GFP); SITS; SITS (Primuline); SITS (Stilbene Isothiosulphonic Acid); SNAFL calcein; SNAFL-1; SNAFL-2; SNARF calcein; SNARF1; Sodium Green; SpectrumAqua; SpectrumGreen; SpectrumOrange; Spectrum Red; SPQ (6-methoxy-N-(3-sulfopropyl)quinolinium); Stilbene; Sulphorhodamine B can C; Sulphorhodamine G Extra; SYTO 11 ; SYTO 12; SYTO 13; SYTO 14; SYTO 15; SYTO 16; SYTO 17; SYTO 18; SYTO 20; SYTO 21; SYTO 22; SYTO 23; SYTO 24; SYTO 25; SYTO 40; SYTO 41; SYTO 42; SYTO 43; SYTO 44; SYTO 45; SYTO 59; SYTO 60; SYTO 61; SYTO 62; SYTO 63; SYTO 64; SYTO 80; SYTO 81; SYTO 82; SYTO 83; SYTO 84; SYTO 85; SYTOX Blue; SYTOX Green; SYTOX Orange; TETTM; Tetracycline; Tetramethylrhodamine (TRITC); Texas RedTM; Texas Red-XTM conjugate; Thiadicarbocyanine (DiSC3); Thiazine Red R; Thiazole Orange ; Thioflavin 5; Thioflavin S; Thioflavin TCN; Thiolyte; Thiozole Orange; Tinopol CBS (Calcofluor White); TMR; TO-PRO-1; TO-PRO-3; TO-PRO-5; TOTO-1; TOTO-3; TriColor (PE-Cy5); TRITC TetramethylRodamineIsoThioCyanate; True Blue; TruRed; Ultralite; Uranine B; Uvitex SFC; VIC®; wt GFP; WW 781; X-Rhodamine; XRITC; Xylene Orange; Y66F; Y66H; Y66W; Yellow GFP; YFP; YO-PRO-1; YO-PRO-3; YOYO-1; YOYO-3; and salts thereof.

[0106] Fluorescent dyes or fluorophores may include derivatives that have been modified to facilitate conjugation to another reactive molecule. As such, fluorescent dyes or fluorophores may include amine-reactive derivatives such as isothiocyanate derivatives and/or succinimidyl ester derivatives of the fluorophore.

[0107] The oligonucleotides and nucleotides of the disclosed methods may be labeled with a quencher. Quenching may include dynamic quenching (*e.g.*, by FRET), static quenching, or both. Suitable quenchers may include Dabcyl. Suitable quenchers may also include dark quenchers, which may include black hole quenchers sold under the tradename “BHQ” (*e.g.*, BHQ-0, BHQ-1, BHQ-2, and BHQ-3, Biosearch Technologies, Novato, CA). Dark quenchers also may include quenchers sold under the tradename “QXL™” (Anaspec, San Jose, CA). Dark quenchers also may include DNP-type non-fluorophores that include a 2,4-dinitrophenyl group.

[0108] The oligonucleotide of the present methods may be labeled with a donor fluorophore and an acceptor fluorophore (or quencher dye) that are present in the oligonucleotides at positions that are suitable to permit FRET (or quenching). Labeled oligonucleotides that are suitable for the present methods may include but are not limited to oligonucleotides designed to function as LightCycler primers or probes, Taqman® Probes, Molecular Beacon Probes, Amplifluor® Primers, Scorpion® Primers, and Lux™ Primers.

[0109] In some embodiments, the detection of viruses that are capable of causing AIDS or AIDS-like symptoms may be performed using MultiCode®-RTx PCR technology, which is disclosed in U.S. Patent Application Publication No. 2002-0150900 and WO/01/90417. The assays may be performed using real-time or continuous methods using any suitable commercial thermal cycler. The disclosed technology may be used to detect nucleic acid targets obtained from any source (*e.g.*, human, animal and infectious disease samples).

ILLUSTRATIVE EMBODIMENTS

[0110] The following list of embodiments is illustrative with respect to the disclosed methods and kits.

[0111] Embodiment 1. A method of detecting a specific mutation in a heterogeneous population of polynucleotides, where the specific mutation, if present, is located in a first region of the population, and the population comprises one or more additional mutations

located in a second region, the method comprising: (a) amplifying the population with a first set of primers to obtain an amplification product (*e.g.* which comprises the first region), where at least one primer of the first set of primers is capable of specifically hybridizing to the second region of the population; and (b) amplifying the product with a second set of primers to detect the mutation, where at least one primer of the second primer set includes at least one non-natural base.

[0112] Embodiment 2. The method of embodiment 1, where the at least one primer of the first primer set does not specifically hybridize to the first region of the population.

[0113] Embodiment 3. The method of embodiment 1, where the at least one primer of the second primer set specifically hybridizes to the first region of the population.

[0114] Embodiment 4. The method of embodiment 1, where the specific mutation comprises a single base change.

[0115] Embodiment 5. The method of embodiment 1, where the first region comprises a single nucleotide.

[0116] Embodiment 6. The method of embodiment 1, where at least one primer of the first set of primers includes at least one non-natural base.

[0117] Embodiment 7. The method of embodiment 1 or 6, where the at least one non-natural base is selected from iC and iG.

[0118] Embodiment 8. The method of embodiment 1, where the at least one primer of the second primer set which includes at least one non-natural base further comprise a label.

[0119] Embodiment 9. The method of embodiment 8, where the label is a fluorophore.

[0120] Embodiment 10. The method of embodiment 9, where amplifying the product with the second set of primers comprises, amplifying in the presence of at least one quencher coupled to a non-natural base.

[0121] Embodiment 11. The method of embodiment 1, where detecting the mutation comprise real-time detection.

[0122] Embodiment 12. The method of embodiment 1, where the population of polynucleotides comprises a sequence of HIV-1.

[0123] Embodiment 13. A method of detecting a specific mutation in a heterogeneous population of polynucleotides, where the specific mutation, if present, is located in a first region of the population and the population comprises one or more additional mutations located in a second region of the population, the method comprising: (a) reacting the population and a mixture of oligonucleotides, where the mixture comprises: (i) a first

oligonucleotide capable of hybridizing to the first region of the population; and (ii) a pool of degenerate oligonucleotides capable of hybridizing to the second region of one or more polynucleotides of the population, where the first oligonucleotide and/or the pool comprises one or more non-natural bases and optionally a label; and (b) detecting the mutation.

[0124] Embodiment 14. The method of embodiment 1, where detecting comprises amplifying one or more polynucleotides of the population.

[0125] Embodiment 15. A method of detecting a specific mutation in a heterogeneous population of polynucleotides, where the specific mutation, if present, is located in a first region of the population and the population comprises one or more additional mutations located in a second region of the population, the method comprising: (a) reacting the population and a pool of degenerate oligonucleotides, where the oligonucleotides of the pool comprise one or more non-natural bases and where the pool comprises: (i) at least one oligonucleotide capable of hybridizing to the first region of one or more polynucleotides of the population; and (ii) a plurality of oligonucleotides capable of hybridizing to the second region of one or more polynucleotides of the population; and (b) detecting the mutation.

[0126] Embodiment 16. The method of embodiment 14 or 16, where the non-natural bases are selected from the group consisting of iso-G, iso-C and a combination thereof.

[0127] Embodiment 17. The method of embodiment 14 or 16, where the degenerate oligonucleotides further comprise at least one label.

[0128] Embodiment 18. The method of embodiment 14 or 16, where at least one label is a fluorophore.

[0129] Embodiment 19. The method of embodiment 14 or 16, where detecting comprises amplifying one or more polynucleotides of the population.

[0130] Embodiment 20. The method of embodiment 19, where detecting comprises amplifying one or more polynucleotides in the presence of one or more non-natural nucleotides.

[0131] Embodiment 21. The method of embodiment 14 or 16, where detecting comprises amplifying one or more polynucleotides of the population in the presence of at least one quencher coupled to a non-natural base.

[0132] Embodiment 22. The method of embodiment 21, where the non-natural base coupled to the at least one quencher is selected from the group consisting of iso-C and iso-G.

[0133] Embodiment 23. The method of embodiment 14 or 16, where detecting comprises real-time detection.

[0134] Embodiment 24. The method of embodiment 14 or 16, where the heterogeneous population of polynucleotides comprises a sequence of HIV-1.

[0135] Embodiment 25. A method for identifying HIV-1 in a sample comprising: (a) reacting a reaction mixture, the mixture comprising: (i) the sample; (ii) at least one oligonucleotide comprising at least one non-natural base, where the oligonucleotide is capable of specifically hybridizing to HIV-1 nucleic acid; and (b) detecting HIV-1 nucleic acid if present in the sample.

[0136] Embodiment 26. The method of embodiment 25, where the at least one oligonucleotide is selected from the group consisting of SEQ ID NOs:2-97.

[0137] Embodiment 27. A method of detecting the presence or absence of a mutation in a polynucleotide at a specific nucleotide, comprising amplifying the polynucleotide with primers to detect the presence or absence of the mutation, the primers comprising: (a) a first primer having a 3' nucleotide that is complementary to the specific nucleotide where the mutation is present and capable of amplifying the polynucleotide; (b) a second primer having a 3' nucleotide that is complementary to the specific nucleotide where the mutation is absent and capable of amplifying the polynucleotide; where the first primer and the second primer are not complementary to the polynucleotide at one or more positions other than the 3' nucleotide and do not include identical nucleotides at the one or more positions, and where at least one of the first primer and the second primer includes at least one non-natural base, which optionally, is present at the one or more positions.

[0138] Embodiment 28. The method of embodiment 27, where the first primer and the second primer are not complementary to the polynucleotide at a single position other than the 3' nucleotide and do not include identical nucleotides at the single position.

[0139] Embodiment 29. The method of embodiment 27, where at least one of the first primer and the second primer has a non-natural base at the one or more positions.

[0140] Embodiment 30. The method of embodiment 27, where both the first primer and the second primer have a non-natural base at the one or more positions.

[0141] Embodiment 31. The method of embodiment 27, where the non-natural base is selected from the group consisting of iso-G, iso-C and a combination thereof.

[0142] Embodiment 32. The method of embodiment 27, where at least one of the primers further comprises a label.

[0143] Embodiment 33. The method of embodiment 27, where each primer further comprises a label, and where each label is different.

[0144] Embodiment 34. The method of embodiment 27, where amplification comprises amplification in the presence of a quencher coupled to a non-natural base.

[0145] Embodiment 35. The method of embodiment 27, where detecting further comprises real-time detection.

[0146] Embodiment 36. The method of embodiment 27, where the polynucleotide comprises a sequence of HIV-1.

EXAMPLES

[0147] The diversity and utility of the methods and kits are demonstrated in the following examples which are meant to be instructive and not limiting.

A. Detection and Quantification of 20 Antiretroviral-Resistant HIV-1 Mutations

[0148] Twenty different mutations that result in antiretroviral-drug resistant HIV-1 were targeted for detection. All twenty mutations selected for study were in the pol region of the HIV-1 genome, and were located in either the reverse transcriptase gene or the protease gene. Table 1 lists the HIV-1 gene carrying the mutation, the amino acid change caused by the mutation, and the drug or drugs to which resistance is conferred. It is understood that the present methods may be used to detect other HIV sequences or mutations; the following examples are illustrative only and not meant to be limiting.

Table 1. HIV-1 Mutations that confer drug resistance

Gene	Mutation	Drug(s)
reverse transcriptase ("RT")	M41L (TTG and CTG)	azidothymidine ("AZT")
RT	T215Y, T215F	AZT
RT	K65R	Didanosine ("DDI"); tenofovir ("TDF"); amdoxovir ("DAPD" or "AMDX")
RT	L74V	DDI; DAPD
RT	T69D	zalcitabine ("DDC")

Gene	Mutation	Drug(s)
RT	E44D	lamivudine ("3TC")
RT	V118I	3TC
RT	M184V, M184I	3TC; emtricitabine ("FTC")
RT	L100I	Multi-non-nucleoside reverse transcriptase inhibitors ("NNRTI")
RT	K103N	NNRTI
RT	Y181C, Y181I	NNRTI
RT	Y188L	NNRTI
protease	M46I	multi-drug resistance
protease	L90M	multi-drug resistance
protease	G48V	saquinavir
protease	D30N	nelfinavir
protease	I50V	aprenavir

1. Target Preparation

[0149] HIV-1 sequences corresponding to bp 2300-3285 of pNL4-3 (Genbank Accession No. M19921) were amplified from pNL4-3 plasmid DNA using standard PCR conditions. PCR products of 985 bp were cloned into pGEM-T vector (Promega, Madison, WI) using the T-A cloning method. Mutations were created using the Quik-Change mutagenesis kit (Stratagene, La Jolla, CA) according to manufacturer's instructions. Constructs and orientation were verified by DNA sequencing.

[0150] Plasmids were isolated using Plasmid Mini kits (QIAGEN, Valencia, CA). Plasmid concentration was determined by OD260 using an extinction coefficient of 50 µg/ml per OD (*see e.g.*, Maniatis *et al.*, Molecular Cloning. A Laboratory Manual. CSH Lab. N.Y. (1989)) and used as PCR targets and as templates for *in vitro* transcription. Plasmids were linearized immediately 3' of the HIV-1 insert using Sal I (New England Biolabs, Ipswich, MA). Following phenol extraction and ethanol precipitation, 1 µg of each linearized plasmid was transcribed into RNA using the Ampliscribe T7 Transcription Kit (Epicentre, Madison, WI) following the manufacturer's instructions. Transcription reactions were treated with RNase-free DNase to degrade plasmid template and then phenol extracted. Free nucleotides and pyrophosphate were removed by size exclusion

chromatography using NAP5 columns (Amersham, Piscataway, NJ) following the manufacturer's instructions. Following lithium chloride ethanol precipitation, RNA was treated again with DNase and then completely desalted with a second round of chromatography. RNA concentration was determined by OD260 and an extinction coefficient of 0.117 mM RNA nucleotide per OD. RNA and plasmid DNA targets were diluted in 10 mM MOPS pH 7.4, 0.1 mM EDTA.

2. Primer Design and Synthesis

[0151] Forward and reverse primers were designed to hybridize to the selected HIV-1 sequences in a background of plasmid pNL4-3 sequence (recombinant clone HIV-1 NY5/BRU (LAV-1) is cloned into pNL4-3; see Genbank # M19921). For each mutation, a wild-type and a mutant forward primer were designed. The wild-type and mutant forward primers for each target contained a different 5' label, either HEX (hexachlorofluorescein), FAM (6-carboxy-fluorescein) or CFO (CalFluor Orange). Primers were designed using Visual OMP software (DNA Software, MI) to have target specific T_m's of 50°C (forward primer) and 60°C (reverse primer).

[0152] The forward primers were constructed to have a 5' target-independent tail portion (*e.g.*, non-complementary to the HIV sequence of interest). PCR reactions with tailed primers were performed as follows. A first round of PCR is performed at a lower temperature to allow the target specific region of the primers to hybridize. Extension with polymerase creates a new template with an extended primer binding site: the target specific sequences and the non-specific tail sequence. Due to the longer primer binding sites, subsequent rounds of hybridization may be performed at higher annealing temperatures. The higher annealing temperatures will likely decrease the amount of non-specific primer hybridization, thereby diminishing the amplification of undesired products.

[0153] Table 2 shows the primer sequences and labels used in the HIV-1 mutation detection reactions.

Table 2. Primer Sequences ("RTx primers")

Protein	Site	Primers	Function	SEQ ID NO
Protease	30N	CGCAGATTTCTATGAGTATCTGAT	Reverse (cDNA)	2
		FAM(iC)GTTTAGCATACAGGAGCA GATG	WT Forward	3
		CFO(iC)GGCATGATACAGGAGCAG ATA	MUT Forward	4
Protease	46I	CTTCCAATTATGTTGACAGGTG	Reverse (cDNA)	5
		FAM(iC)GAGCGATGGAAACCAAAA ATG	WT Forward	6
		CFO(iC)CCTGACATAGATGGAAAC CAAAAATA	MUT Forward	7
Protease	48V	CTTCCAATTATGTTGACAGGTG	Reverse (cDNA)	8
		FAM(iC)GTATCAACGAAACCAAAA ATGATAGG	WT Forward	9
		CFO(iC)CGTCCTGAAACCAAAAAT GATAGT	MUT Forward	10

Protein	Site	Primers	Function	SEQ ID NO
Protease	50V	CTTCCAATTATGTTGACAGGTG	Reverse (cDNA)	11
		FAM(iC)ACTGATGAAATGATAGGG GGAG	WT Forward	12
		CFO(iC)ACTGCTAAAATGATAGGG GGAA	MUT Forward	13
Protease	90M	GTACAAATTTCTACTAATGCTTTT ATTTTT	Reverse (cDNA)	14
		FAM(iC)CGACAATAAAATTGGAAG AAATCTGT	WT Forward	15
		CFO(iC)CGCCATAATTGGAAGAAA TCTGA	MUT Forward	16
RT	41L-TTG	GCAAATACTGGAGTATTGTATGGA	Reverse (cDNA)	17
		FAM(iC)CCATTTACTGTAGAAATTT GTACAGAAA	WT Forward	18
		CFO(iC)GGCTGGTAGAAATTTGTAC AGAAT	MUT Forward	19
RT	41L-CTG	GCAAATACTGGAGTATTGTATGGA	Reverse (cDNA)	20
		FAM(iC)CCATTTACTGTAGAAATTT GTACAGAAA	WT Forward	21
		CFO(iC)GGTAAATGAGTAGAAATT TGTACAGAAC	MUT Forward	22

Protein	Site	Primers	Function	SEQ ID NO
RT	44D	ATTCCTAATTGAACTTCCCAGAAA	Reverse (cDNA)	23
		FAM(iC)CTTGCTCAGAAATGGAAA AGGAA	WT Forward	24
		CFO(iC)CTGGATGAGAAATGGAAA AGGAC	MUT Forward	25
RT	M184 V (1)	CTCAACAGATGTTGTCTCAGTTCC TCTA	Reverse (cDNA)	26
		FAMT(iC)GACAGGAGACATAGTCA TCTATCAATACA	WT Forward	27
		HEXT(iC)TGTCCTAATAGTCATCTAT CAATACG	MUT Forward	28
RT	M184 V (2)	TAAATCCCCACCTCAACAGAT	Reverse (cDNA)	29
		FAM(iC)GAGTAAGCTAGTCATCTA TCAATACATG	WT Forward	30
		CFO(iC)CAGCCATAGTCATCTATCA ATACG	MUT Forward	31
RT	69D	GAAAAATATGCATCGCCCAC	Reverse (cDNA)	32
		FAM(iC)TCGCTCATAAAGAAAAAA GACAGTA	WT Forward	33
		CFO(iC)CGGTTCATAAAGAAAAAA GACAGTC	MUT Forward	34
RT	74V	CATCGCCCACATCCAG	Reverse (cDNA)	35
		FAM(iC)GCTACGAGTACTAAATGG AGAAAAT	WT Forward	36
		CFO(iC)CCCTGTGTACTAAATGGAG AAAAG	MUT Forward	37

Protein	Site	Primers	Function	SEQ ID NO
RT	100L	CCCTGGTGTCTCATTGTTT	Reverse (cDNA)	38
		FAM(iC)TGTTGAAATCCTGCAGGG T	WT Forward	39
		CFO(iC)AGACGATCCTGCAGGGA	MUT Forward	40
RT	103N	CCCTGGTGTCTCATTGTTT	Reverse (cDNA)	41
		FAM(iC)AGGACCCAGGGTTAAAAC AGAAA	WT Forward	42
		CFO(iC)GTCTCGAGGGTTAAAACA GAAC	MUT Forward	43
RT	118I	TGTCATGCTACACTGGAATATTG	Reverse (cDNA)	44
		FAM(iC)TGCCCGATGCATATTTTTC AG	WT Forward	45
		CFO(iC)TCCGTCGATGCATATTTT CAA	MUT Forward	46
RT	181C	TAAATCCCCACCTCAACAGAT	Reverse (cDNA)	47
		FAM(iC)CCGTTCCAGACATAGTCA TCTA	WT Forward	48
		CFO(iC)AGGCACAGACATAGTCAT CTG	MUT Forward	49
RT	181I	TAAATCCCCACCTCAACAGAT	Reverse (cDNA)	50
		FAM(iC)CCGTTCCAGACATAGTCA TCTA	WT Forward	51
		CFO(iC)GGTTAGTCCAGACATAGTC ATCA	MUT Forward	52

Protein	Site	Primers	Function	SEQ ID NO
RT	M184 I	TAAATCCCCACCTCAACAGAT	Reverse (cDNA)	53
		FAM(iC)GAGTAAGCTAGTCATCTA TCAATACATG	WT Forward	54
		CFO(iC)CATTCGCATAGTCATCTAT CAATACATA	MUT Forward	55
RT	K65R (1)	ACCCTGCAGGATGTGG	Reverse (cDNA)	56
		FAM(iC)ACAGGTAGTATTTGCCAT AAAGAA	WT Forward	57
		CFO(iC)GACATCGTATTTGCCATAA AGAG	MUT Forward	58
RT	K65R (2)	CTGATTTTTTCTGTTTAAACCCTGC	Reverse (cDNA)	59
		FAM(iC)TCACGTAGTATTTGCCATA AAGAA	WT Forward	60
		CFO(iC)TGCTGGTATTTGCCATAAA GAG	MUT Forward	61
RT	188L- CTT	AAGGAATGGAGGTTCTTTCTG	Reverse (cDNA)	62
		FAM(iC)GACGGTAATACATGGATG ATTTGTA	WT Forward	63
		CFO(iC)GCCTAAGACATGGATGATT TGC	MUT Forward	64
RT	188L- TTA	AAGGAATGGAGGTTCTTTCTG	Reverse (cDNA)	65
		FAM(iC)GACGGTAATACATGGATG ATTTGTA	WT Forward	66
		CFO(iC)CCGCTATACATGGATGATT TGTT	MUT Forward	67

Protein	Site	Primers	Function	SEQ ID NO
RT	215F	AGCACTATAGGCTGTACTGTC	Reverse (cDNA)	68
		FAM(iC)ATCTGTTTGAGGTGGGGA TTTA	WT Forward	69
		CFO(iC)TGTGAGAGGTGGGGATTTT T	MUT Forward	70
RT	215Y	AGCACTATAGGCTGTACTGTC	Reverse (cDNA)	71
		FAM(iC)ATCTGTTTGAGGTGGGGA TTTA	WT Forward	72
		CFO(iC)CTAGACAGAGGTGGGGAT TTTA	MUT Forward	73

3. General Reaction and Cycling Conditions

[0154] For each assay, PCR primers were used at the following concentrations. Each reaction received 200 nM of a wild-type forward primer and 200 nM of a mutant forward primer. The M184V mutant primer was used at 150 nM; the K65K (wild-type) and K65R (mutant) primers were used at 100nM. A single reverse primer was used for each system at 400 nM.

[0155] Either 100 or 1000 copies of mutant RNA or DNA with 10-fold dilution series of wild-type targets (DNA or RNA) from 0 to 10⁶ targets were tested with each primer set.

[0156] PCR conditions were as follows: 25 uL reactions in 1x ISolution buffer (EraGen, Madison, WI) and Titanium Taq DNA polymerase (Clontech, Mountain View, CA) at manufacturers recommended concentration. For RT- PCR assays the conditions included the following: 0.5 Units/uL Maloney Murine Reverse Transcriptase (M-MLV RT) and 5 mM dithiothreitol and an additional 5 minutes at 50°C added prior to the denaturation step.

[0157] Cycling parameters for reactions performed on the ABI 7900 (Applied Biosystems, Foster City, CA) real-time thermal cycler were as follows:

2 minutes denature at 95°C, 1 cycle of 5 sec at 95°C, 5 sec at 45°C, 20 sec at 72°C, followed by 45 cycles of 5 sec at 95°C, 5 sec at 60°C, 20 sec

at 72°C with optical read. A thermal melt at 7% ramp rate with optical read from 60 to 95°C was performed directly following the last 72°C step of thermal cycling.

[0158] Cycling parameters for reactions performed on the Roche LightCycler (Roche, Indianapolis, IN) were as follows unless otherwise specified:

[0159] 2 minutes denature at 95°C, 1 cycle of 1 sec at 95°C, 1 sec at 45°C, 1°C/sec ramp to 20 sec at 72°C, followed by 50 cycles of 5 sec at 95°C, 5 sec at 55°C, 1°C/sec ramp to 20 sec at 72°C (SINGLE read), melt 60-95°C 0.4°C/sec ramp; (STEP read).

[0160] Cycle threshold data was analyzed to determine the sensitivity of each system for both mutant and wild-type as determined by the lowest copy number of targets that could consistently be detected. The slope of each wild-type standard curve was also determined as a measure of PCR efficiency.

B. Detection of HIV-1 Mutations in RNA Targets on the ABI 7900

[0161] HIV-1 drug resistant mutant RNA targets were mixed at 100 or 1000 copies with wild-type RNA target in a 10-fold dilution series from 0 to 10⁶ copies. Assays were performed using the ABI 7900. Targets were amplified using the cycling parameters described above. Cycle threshold data was analyzed to determine the sensitivity of each system for both mutant and wild-type as shown by the lowest copy number of targets that could consistently be detected. The slope of each wild-type standard curve was also determined as a measure of PCR efficiency.

[0162] Results are shown in Table 3. Sensitivity to both wild-type and mutant targets in copy numbers is presented; “not detected” indicates sensitivity greater than 1000 mutant RNA targets. The “Best Mixture” is the lowest percentage mutant target in a mixture with wild-type target that could be detected. “Slope” is the slope wild-type standard curve in log copy number vs. cycle threshold. A slope of -0.32 indicates 100% PCR efficiency with greater values indicating reduced PCR efficiency. “RT” indicates HIV-1 reverse transcriptase. Numerous systems can detect 100 copies of both wild-type and mutant RNA in a mixture of 1% or less.

Table 3. RNA detection results

Protein	Site	Wild-type		Mutant	
		Sensitivity	Slope	Sensitivity	Best Mixture
protease	30N	10	-0.23	100	0.1%
protease	46I	10000	-0.25	Not detected	MUT Fails
protease	48V	100000	-0.25	1000	0.1%
protease	50V	100	-0.24	100	0.1%
protease	90M	10000	-0.27	1000	0.1%
RT	41L-CTG	100	-0.22	100	0.01%
RT	44D	10	-0.23	100	0.01%
RT	65R (1)	100	-0.26	100	0.1%
RT	65R (2)	10	-0.22	100	0.01%
RT	69D	1000	-0.19	Not detected	MUT Fails
RT	74V	10000	-0.22	100	1%
RT	103N	10000	-0.25	100	1%
RT	118I	10000	-0.32	1000	MUT Fails
RT	181C	10	-0.23	Not detected	MUT Fails
RT	181I	10	-0.28	100	0.01%
RT	184I	100	-0.24	100	0.01%
RT	184V (1)	10	-0.24	100	0.01%
RT	184V (2)	10	-0.23	100	0.01%
RT	188L-CTT	10	-0.23	100	0.1%
RT	188L-TTA	10	-0.23	1000	0.1%
RT	215F	10	-0.22	100	1%
RT	215Y	10	-0.26	100	1%

C. Detection of HIV-1 Mutations in DNA Targets on the ABI 7900 and the LightCycler

[0163] HIV-1 drug resistant mutant plasmid DNA clone targets were mixed at 100 or 1000 copies with wild-type DNA target in a 10-fold dilution series from 0 to 10^6 copies. Assays were performed on the ABI 7900 and the Roche LightCycler (Roche, Indianapolis, IN) using cycling parameters described above in section A.3.

[0164] Cycle threshold data was analyzed to determine the sensitivity of each system for both mutant and wild-type as shown by the lowest copy number of targets that could consistently be detected. The slope of each wild-type standard curve was also determined as a measure of PCR efficiency.

[0165] Results are presented in Tables 4A and B. "Sensitivity" to both wild-type and mutant targets is presented as copy number. "Not detected" indicates a sensitivity greater than 1000 mutant RNA targets. "Best Mixture" is the lowest percentage mutant target in a mixture with wild-type target that was detected. "Slope" is the slope wild-type standard curve in log copy number vs. cycle threshold. A slope of -0.32 indicates 100% PCR

efficiency with greater values indicating reduced PCR efficiency. "RT" indicates reverse transcriptase. Again, numerous systems can detect 100 copies of both wild-type and mutant RNA in a mixture of 1% or less on either the ABI or the LightCycler. Note that two independent systems for the 65R and the 184V loci were evaluated.

Table 4A. DNA Detection Results

		ABI 7900			
		Wild-type		Mutant	
Protein	Site	Sensitivity	Slope	Sensitivity	Best Mixture
protease	30N	10	-0.29	1000	0.1%
protease	46I	10	-0.24	100	0.1%
protease	48V	100	-0.23	100	0.01%
protease	50V	1000	-0.24	1000	50%
protease	90M	100000	-0.17	Not detected	MUT Fails
RT	41L-TTG	100	-0.22	100	0.01%
RT	41L-CTG	100	-0.22	100	0.01%
RT	44D	1000	-0.21	1000	90%
RT	65R (1)	100	-0.28	100	0.01%
RT	65R (2)	10	-0.23	100	0.01%
RT	69D	100	-0.24	1000	0.01%
RT	74V	100	-0.27	100	0.1%
RT	100L	10000	-0.24	1000	0.1%
RT	103N	100	-0.24	100	0.1%
RT	118I	100	-0.26	100	50%
RT	181C	10	-0.23	Not detected	MUT Fails
RT	181I	100	-0.26	Not detected	MUT Fails
RT	184I	100	-0.24	100	0.01%
RT	184V (1)	1000	-0.25	1000	0.1%
RT	184V (2)	1000	-0.24	1000	0.1%
RT	188L-CTT	10	-0.24	1000	10%
RT	188L-TTA	100	-0.22	100	0.1%
RT	215F	10	-0.27	100	0.01%
RT	215Y	10	-0.26	100	1%

Table 4B. DNA Detection Results

		Light Cyclier			
		Wild-type		Mutant	
Protein	Site	Sensitivity	Slope	Sensitivity	Best Mixture
protease	30N	1000	-0.14	1000	0.1%
protease	46I	10	-0.12	100	0.01%
protease	48V	100	-0.09	100	0.01%
protease	50V	100	-0.16	1000	90%
protease	90M	1000	-0.08	100	0.1%
RT	41L-TTG	1000	-0.12	100	0.01%
RT	41L-CTG	10000	-0.09	100	0.01%
RT	44D	100	-0.09	100	0.1%
RT	65R (1)	100	-0.22	100	0.1%
RT	65R (2)	100	-0.08	100	0.01%
RT	69D	100	-0.08	100	0.01%
RT	74V	1000	-0.18	100	0.01%
RT	100L	1000	-0.15	100	0.01%
RT	103N	100	-0.16	100	0.1%
RT	118I	10	-0.12	1000	0.1%
RT	181C	1000	-0.12	Not detected	MUT Fails
RT	181I	100	-0.20	100	90%
RT	184I	100	-0.09	100	0.01%
RT	184V (1)	10	-0.23	1000	0.1%
RT	184V (2)	1000	-0.15	1000	0.1%
RT	188L-CTT	1000	-0.16	100	0.01%
RT	188L-TTA	100	-0.14	100	0.01%
RT	215F	100	-0.12	100	0.01%
RT	215Y	10	-0.14	100	0.01%

D. Detection of M184V and K65R in DNA Targets using Different Cycling Parameters and Different Instruments

[0166] To test for cycling robustness, the M184V and K65R assays were run and tested using the ABI 7900 instrument and the cycling parameters described in section A.3, and the Roche LightCycler using the three different sets of cycling parameters, with ramp rates of 20°C/sec (unless otherwise specified), described below.

- 1) 2 minute denature at 95°C, 1 cycle of 1 sec at 95°C, 1 sec at 45°C, 1°C/sec ramp to 20 sec at 72° C, 50 cycles 5 sec at 95°C, 5 sec at 55°C, 1°C/sec ramp to 20 sec at 72° C (SINGLE read), melt 60-95°C 0.4°C/sec ramp; (STEP read);

2) 2 min denature at 95°C, 1 cycle of 1 sec at 95°C, 1 sec at 45°C, 20 sec at 72°C, 50 cycles of 5 sec at 95°C, 5 sec at 55°C, 1°C/sec ramp to 20 sec at 72°C (SINGLE read), melt 60-95°C 0.4°C/sec ramp (STEP read);

3) 2 minute denature at 95°C, 1 cycle of 1 sec at 95°C, 1 sec at 45°C, 20s at 72°C, 100 cycles of 1 sec at 95°C, 1 sec at 55°C, to 20 sec at 72°C (SINGLE read), melt 60-95°C 0.4°C/sec ramp (STEP read).

[0167] Mixtures of wild-type and mutant cloned DNA targets were prepared containing 10^3 to 10^7 copies of mutant or wild-type targets at 10-fold intervals, with a total of 10^7 copies per reaction. This provided mixtures that varied in wild-type and mutant targets from 0.01 to 100%.

[0168] Linear regression analysis was performed to determine $\log[\text{copy number}]$ vs. C_t for the individual channels. Each condition for M184 afforded tight r^2 values which varied from 0.991 to 0.998 for the wild-type M184M channel and 0.990 to 0.997 for the mutant M184V channel. Linear regression for the K65 system had r^2 values which varied from 0.949 to 0.955 for the wild-type K65K channel and 0.977 to 0.994 for the mutant K65R channel. By measuring the differences between the C_t channels of wild-type and mutant (C_t of the mutant minus the C_t of the wild-type, defined as ΔC_t) ΔC_t standard curves were established for each condition. These standard curves may be used to determine the make-up of unknown sample mixtures.

[0169] Results are shown in FIGURE 3A and 3B, which shows ΔC_t vs. fraction mutant plots for real-time PCR with the four different cycling conditions. "Fraction Mut" is the fraction of mutant DNA template in a mixture with wild-type DNA, with fractions detected down to 1 in 10,000 copies (0.0001). A1-A4 show the curves for K65V detection and B1-B4 show the curves for M184V detection. A1-A3 and B1-B3 were performed on the Roche LightCycler, and A4 and B4 were performed on the ABI 7900.

E. Detection of Mutations Using Varied Concentrations of RNA Targets

[0170] RNA target mixtures were diluted in a 10-fold series from 10^3 to 10^7 total copies, using mixtures of targets from 1% to 99% M184V. Reactions were performed on the ABI

7900 as described in section A.3. ΔC_t values were determined for each sample and then plotted vs. fraction M184V in the mixtures. These different curves were then overlaid, and are shown in FIGURE 4. The results indicate that the detection methods allowed for percentages to be determined even as the overall concentration changes over four orders of magnitude (four logs of total viral particle input did not affect the overall curve). Accordingly, the method may be used to discriminate mixed populations even through a large range in overall concentration. Thus, quantification of the mixed viral populations need not be determined prior to mixed population analysis.

F. Detection of M184M, M184V and M184I DNA on LightCycler Using Triplex Detection System

[0171] Two different drug resistant mutants at codon 184 in the reverse transcriptase gene have been identified, M184V and M184I. Both are the result of transition mutations. In the case of M184V, the wild-type codon ATG, which codes for methionine becomes mutant GTG which codes for valine. In the case of the M184I mutant, the wild-type codon ATG becomes mutant ATA which codes for isoleucine.

[0172] Primers used for the detection of the three different alleles were labeled with three different fluorophores. The primers for the three different detection systems were combined in the reaction along with quenchers capable of quenching the different fluorophores. Five different targets mixes (containing 2×10^6 total copies) were tested: 100%M184M; 100%M184V; 100%M184I; 50%M184I : 50%M184M; 50%M184V : 50%M184M. Reactions were performed on the Roche LightCycler, using cycling parameters described in section A.3. Real-time and PCR melt data confirmed that such a triplex system is capable of detecting specific mutations in a mixed population. Probes specific for the M184M targets detected these targets in the 100% M184M mix, the 50%M184I : 50%M184M mix, and in the 50%M184I : 50%M184M mix. Probes specific for the M184V targets detected these targets in the 10%M184V mix and the 50%M184V : 50%M184M mix. Probes specific for the M184I targets detected these targets in the 100%M184I and the 50%M184I : 50%M184M mix.

G. Detection of HIV-1 Drug Resistant Mutants From Patient Samples

[0173] To further assess the detection methods, viral RNA extracted from 13 serum samples containing HIV-1 which had previously been genotyped by the line probe assay

(LiPA) was tested. These samples were obtained from subjects undergoing non-nucleoside reverse transcriptase inhibitor ("NRTI") therapy between the years 1994 and 1997 and displaying M184M, M184V or a mixed population of M and V by LiPA. The extracted RNA was tested in duplicate using the M184V system as described above using ABI 7900 and cycling parameters described above in A.3. Table 5 indicates that the detection methods were able to amplify and genotype all samples. The results from ten of the samples were completely concordant with LiPA. Of the three samples that were not 100% concordant (samples 2, 4 and 11), two samples displayed a mixture of M and V where LiPA only detected V (samples 4 and 11). For the third sample, the opposite was true; the present detection methods showed 100% M while LiPA displayed a mixture (sample 2). The range of allele fractions from the samples were tested at least twice in independent real-time PCR experiments (HIV Monitor, Roche, Indianapolis, IN) and compared to previous results determined by LiPA.

Table 5. Viral genotypes and load

LIPA^a	RTx	Viral Load/ml^b	Bleed Date	Sample
184M	100% M	6.5×10^5	12/15/94	1
184M/V	100% M	ND	02/27/95	2
184M	100% M	1.7×10^5	11/03/96	3
184V	64-92% V	1.6×10^4	08/23/96	4
184V	100% V	1.4×10^6	06/21/96	5
184M	100% M	1.1×10^6	02/15/96	6
184M	100% M	1.8×10^5	06/08/95	7
184V	100% V	2.8×10^5	06/21/97	8
184M	100% M	1.2×10^5	11/16/95	9
184V	100% V	1.0×10^5	04/26/96	10
184V	90-100% V	4.3×10^5	06/28/96	11
184M	100% M	3.3×10^5	12/29/95	12 ^c
184V	100% V	1.6×10^5	06/28/96	13 ^c

^b viral load determined via Roche Monitor™ system.

^c Samples only tested once.

H. Detection of Mutations in 13 HIV-1 Viral Clones Using Degenerate Primers

[0174] Plasmid DNA of 13 clones from the NIH Multidrug Resistant HIV-1 Reverse Transcriptase Panel and a wild-type clone (*i.e.*, no M184 mutation, pNL4-3 was used) were tested using the M184V assay. Numerous background polymorphisms exist within these clones, and the assay results indicated less than 50% of the clones could be genotyped.

Table 6.

[0175] To evaluate the extent of the polymorphisms present in the clones, the sequences of the discriminatory primers for each system were aligned with the sequences of the clones to determine the number of mismatched bases between the primer and different clone template sequences. The total number of mismatches was divided by the number of calls made to determine the average number of primer-target mismatches for both primers.

Table 6. Detection of M184V Viral Clones with Standard Primers

Mutation	Calls			Average Mismatches per Clone		
	Correct	Not Called	Miscalled	Correct	Not Called	Miscalled
M184V	6	7	0	2.8	5.4	NA

[0176] Notably, there were no incorrect calls for this system. It was also noted that when the number of mismatches increased, so did the inability to genotype.

I. Use of Degenerate Primers to Detect HIV-1 Mutations

[0177] Degenerate primers for the detection of the M184V mutation were synthesized as a pool using mixed phosphoramidites to insert multiple bases that corresponded to the most common substitutions in HIV-1 clades B, C, and E. Using the NIH panel plasmid clones described above, the RTx degenerate primers were evaluated using reaction conditions described in section A, above, and the ABI 7900 with the following cycling parameters:

2 minute denature at 95°C, followed by one cycle of 5s at
95°C, 30s at 40°C, 20s at 72°C, followed by 60 cycles of 5s at
95°C, 5s at 50°C, 20s at 72°C, melt 60°C to 95°C 7% ramp.

[0178] The results of the primer redesign are shown in Table 7 below. Bolding of letters within sequences indicates differences between targets, degenerate primer sequence and pNL4-3; **R** = A or G base, **Y** = C or T base. Note that the primer pairs are labeled with separate fluorophores. The sequence for each target within the priming region is shown, along with the T_m for both the forward and reverse non-degenerate primer. The results are shown on the right. Black, filled columns A-B indicate that the correct genotype was obtained in the detection reactions. Reactions: A, standard (non-degenerate) primers and 2 x10⁶ copies target used; B, degenerate primers and 2 x10⁶ copies of target DNA used. "For.Tm" is the predicted melting temperature in °C of the forward primer including

complementary 3' base on template under reaction conditions. "Rev.Tm" is the predicted melting temperature in °C of the reverse primer (sequence not shown) on template under reaction conditions; bold indicates lowest reverse primer Tm on samples tested.

[0179] Again, the standard primers genotyped 6 of 13 subject-derived clones (Column A). It was noted the T_m of the forward primer for the targets not genotyped was very low. Reactions using degenerate primers amplified and correctly genotyped 11 of the 13 clones (Column B). Additional clone 12-21 was isolated by cloning of cDNA from an HIV-1 infected subject sample for this study.

Table 7. Exemplary M184 Primer re-design

Degenerate Primers	GACATRGTYATCTATCARTAYR	SEQ IN NO 74			
Standard Primers	GACATAGTCATCTATCAATACR	SEQ ID NO 75			Reaction
Sample	Sequence	For. Tm	Rev. Tm	A	B
pNL4-3	GACATAGTCATCTATCAATACA	54.2	65.4		
7324-1	GAAATAGTTATCTATCAATATA	28.5	59.5		
7324-4	GAAATAGTTATCTATCAATATA	28.5	59.5		
10076-4	GAAATGGTTATCTATCAATACG	26.8	65.4		
7295-1	GACATAGTCATCTATCAATACG	55.4	61.2		
4755-1	GACATAGTTATCTATCAATATG	40.7	51.9		
6463-13	GAAATAGTTATCTATCAATACG	37.3	43.4		
7303-3	GACATAGTTATCTATCAATACA	45.5	43.4		
1617-1	GACATGGTTATCTATCAATATG	30.6	65.1		
35764-2	GACATAGTTATCTATCAATACA	45.5	61.5		
29129-2	GATATAGTCATCTATCAATACG	47.8	48.6		
52534-2	GACATAGTTATCTATCAATACG	47.0	61.9		
56252-1	GACATAGTTATCTATCAATACA	45.5	65.4		
12-21	GACATAGTTATTTACCAATACA	25.8	65.1		

J. Incorporating Mismatches Into Primer Sequences

[0180] RTx primers were designed with specific mismatches. In one embodiment, the mismatches were positioned adjacent to the 3' base of the forward RTx primer. As previously described, two different forward primers may be used in the RTx reactions. One forward primer has a 3' base that will hybridize to the wild-type nucleotide sequence of the

target, while the other primer has a 3' base that will hybridize to the mutant nucleotide sequence of the target. For example, one of the M184M forward primers described in Table 2 has the sequence 5'-FAMT(iC)GACAGGAGACATAGTCATCTATCAATACA, where the 3' "A" hybridizes to the wild-type "T." The M184V forward primer has the sequence 5'-HEXT(iC)TGTCCAATAGTCATCTATCAATACG, where the 3' "G" hybridizes to the mutant "C."

[0181] The sequence alignment of the 13 clones in Table 7 shows a polymorphism adjacent to M184 A/G mutation site. Some clones show a "T" while other show a "C" at this position. Primers were designed with a mismatch at this position having the following sequences. The forward mismatch primer specific for M184M is as follows: HEX(iC)GACAGGAGACATAGTTATCTATCAATATA (SEQ ID NO: 76). The forward mismatch primer specific for M184V is as follows:

FAM(iC)TGTCCAGACATAGTTATCTATCAATAGG (SEQ ID NO: 77).

[0182] Note that for the M184M forward primer, the "T" adjacent to the 3' end will not match either the "T" or the "C" in the any of the target sequences. Similarly, the "G" adjacent to the 3' end of the M184V primer will not match either the "C" or the "T" in any of the target sequences. In addition to the mismatch primers described above, the following non-mismatch primers were also tested:

[0183] 5'-HEX(iC)GACAGGAGACATAGTTATCTATCAATACA, (SEQ ID NO: 78) and 5'-FAM(iC)TGTCCAGACATAGTTATCTATCAATACG (SEQ ID NO: 79).

[0184] Reactions were prepared as follows: all odd numbered samples contained M184M target and all even numbered samples contained M184V target. Samples 1 and 2 contained both non-mismatched primers; samples 3 and 4 contained the mismatched M184M primer and the non-mismatched M184V primer; samples 5 and 6 contained the mismatched M184V primer and the non-mismatched M184M primer; samples 7 and 8 contained both the mismatched M184M and the M184V primers. PCR cycling conditions were as follows:

15m at 54°C, 2 minutes at 95°C, followed by 3 cycles of 5s at 95°C, 5s at 61°C, 1°C/sec, 10s at 72°C, followed by 1 cycle of 5s at 95°C, 5s at 45°C, 1°C/sec 20s at 72°C)1, followed by 110 cycles of (5s at 95°C, 5s at 55°C, 20s at 72°C), melt 60°C to 95°C 0.4°C STEP.

[0185] Real-time results demonstrate improved discrimination when both mismatched primers are used as compared to both non-mismatch or only one mismatch (data not shown).

K. Primer Mismatches Using Non-Natural Nucleotides

[0186] RTx primers with mismatches were designed with non-natural nucleotides, here either iso-G or iso-C, in the mismatch position. Using the HIV-1 M184 sequence as target, the primers designed and tested are shown in Table 8 below.

Table 8

SEQ ID NO:	PRIMER SEQUENCE	TARGET
80	HEX-XGACAGGAGACATAGTTATCTATCAATACA	HIV M184M
81	FAM-XTGTCCAGACATAGTTATCTATCAATACG	HIV M184V
82	HEX-XGACAGGAGACATAGTTATCTATCAATAXA	HIV M184M
83	FAM-XTGTCCAGACATAGTTATCTATCAATAXG	HIV M184V
84	HEX-XGACAGGAGACATAGTTATCTATCAATAYA	HIV M184M
85	FAM-XTGTCCAGACATAGTTATCTATCAATAYG	HIV M184V
86	HEX-XGACAGGAGACATAGTTATCTATCAATYCA	HIV M184M
87	FAM-XTGTCCAGACATAGTTATCTATCAATYCG	HIV M184V
88	HEX-XGACAGGAGACATAGTTATCTATCYATACA	HIV M184M
89	FAM-XTGTCCAGACATAGTTATCTATCYATACG	HIV M184V
90	HEX-XGACAGGAGACATAGTTATCTYTCAATYCA	HIV M184M
91	FAM-XTGTCCAGACATAGTTATCTYTCAATYCG	HIV M184V
92	HEX-XGACAGGAGACATAGTTYTCTATCYATACA	HIV M184M
93	FAM-XTGTCCAGACATAGTTYTCTATCYATACG	HIV M184V
94	HEX-XGACATAGTTATCTYTCAATYCA	90 No tail
95	FAM-XGACATAGTTATCTYTCAATYCG	91 No tail
96	HEX-XGACATAGTTYTCTATCYATACA	92 No tail
97	FAM-XGACATAGTTYTCTATCYATACG	97 No tail
target	3' - ... CTGTATCAGTAGATAGTTATGTAC	M184M

X = iC

Y = iG

[0187] Numerous primer combinations were tested including combinations in which each forward primer in the reaction (wild-type and mutant) has 0, 1 or 2 iso-G bases, mismatches at the same, or different positions. Exemplary primer combinations are shown below in Tables 9 and 10.

Table 9

M184M SEQ ID NO:	M184V SEQ ID NO:
80 (no non-natural bases)	81 (no non-natural bases)
80	87 (one non-natural base at 27)
80	89 (one non-natural base at 24)

M184M SEQ ID NO:	M184V SEQ ID NO:
80	91 (two non-natural bases: 21 and 27)
86 (one non-natural base at base 28)	81 (no non-natural bases)
86	87 (one non-natural base at 27)
86	89 (one non-natural base 24)
86	91 (two non-natural bases: 21 and 27)
88 (one non-natural base at base 25)	81 (no non-natural bases)
88	87 (one non-natural base at 27)
88	89 (one non-natural base 24)
88	91 (two non-natural bases: 21 and 27)
92 (one non-natural base at base 25)	81 (no non-natural bases)
92	87 (one non-natural base at 27)
92	89 (one non-natural base 24)
92	91 (two non-natural bases: 21 and 27)

Table 10

SEQ ID NO: M184M	SEQ ID NO: M184V
84 (no non-natural bases)	85 (no non-natural bases)
84	89 (one non-natural base at oligonucleotide position 28)
84	95 (two non-natural base: 21 and 27)
88 (one non-natural base at 29)	85 (no non-natural bases)
88	89 (one non-natural base at 28)
88	95 (two non-natural bases: 21 and 27)
96 (two non-natural bases: 18 and 25)	85 (no non-natural bases)

SEQ ID NO: M184M	SEQ ID NO: M184V
96	89 (one non-natural base at 28)
96	95 (two non-natural base: 21 and 27)

[0188] PCR reaction were performed in 1X ISOlution buffer (EraGen Biosciences, Madison, WI) containing 100 μ M dCTP, 200 nM forward primers, 400 nM reverse primer and 3×10^4 linearized plasmid targets. PCR was performed using the ABI 7900 with the following parameters:

2 minute denature at 95°C followed by 2 cycles of (5s at 95°C, 30s at 45°C, 20s at 72°C), followed by 70 cycles of (5s at 95°C, 10s at 55°C, 20s at 72°C), melt 60°C to 95°C with 7% gradient.

[0189] Results indicate that the addition of non-natural mismatches in different positions in the mutant and wild-type primers allows discrimination of mutant and wild-type targets. Data is shown in FIGURES 5 and 6. FIGURES 5 and 6 show the difference in cycle threshold (“dCt”) of Mutant Signal minus Wild-Type Signal for different primer pairs. Each set of three bars indicates results using (a) 100% wild-type M184M target (black bars); (b) 50% wild-type M184M and 50% mutant M184V target (dark grey bars in FIGURE 5) or 10% mutant target (dark grey bars in FIGURE 6); and (c) 100% mutant M184V target (light grey bars).

[0190] Note that primer concentration, salt, buffer conditions, temperature, cycling parameters, and enzyme identity may effect dCt values for any of the primer pairs tested.

L. Healing Primers

[0191] Another strategy employed to improve genotyping results involved “healing” primers as illustrated in FIGURE 7. Part A of FIGURE 7 illustrates long primers of high T_m (gray arrows) that can amplify polymorphic HIV-1 despite the presence of mismatches between primer and template. The forward primer terminates just upstream of the target drug-resistance mutation (filled circle). The healing primers may be designed to have a sequence that is highly homologous to the detection primers. Part B shows that the amplification reaction using the healing primers results in a bipartite template that contains perfect matches (gray lines) to the RTx primers which flank the mutation contained within

the original HIV-1 genome (black line). The healed HIV-1 target can now be amplified by the primers of the present method (black arrows).

[0192] The forward healing primer was designed with a 3' end that terminates just upstream of the drug-resistance mutation site of interest. Since the long healing primers have a higher T_m than that of the detection primers, hybridization and primer extension are expected to occur despite the presence of mismatches between the primer and template. Thus, amplification with the healing primers yields an amplification product from the polymorphic HIV-1 target that matches the detection primer binding sites but still contains the sequence context of the viral template drug-resistance mutation.

[0193] Reactions using healing primers were performed on the MJ PTC-200 as follows: a pre-PCR amplification with the healing primers used the following cycling parameters: 2 minutes at 95° C, followed by 40 cycles of 10s at 95° C, 10s at 55° C, 20s at 72° C. Samples were then transferred to the ABI 7900 and the following cycling parameters were used: 2 minutes at 95° C, followed by 1 cycle of 5s at 95° C, 5s at 45° C, 20s at 72° C, followed by 50 cycles of 5s at 95° C, 5s at 55° C, 20s at 72° C, melt 60° C to 95° C, 2% ramp.

[0194] Results using the healing primers and the same set of plasmid cDNA targets described in Table 7 above, are shown in Table 11 below. Black filled columns A-C indicate RTx genotyping reactions where the correct genotype was obtained. Reactions: A, standard RTx primers and 2×10^6 copies target used. B, healing forward primer combined with, standard RTx primers and on 2×10^6 copies target DNA; C, 40 rounds healing forward and reverse primer (pre-amplification) with 2×10^6 copies plasmid DNA. The PCR product was diluted to 2×10^5 copies for RTx with the standard primers.

[0195] As in section H, the standard primers could only genotype 6 of 13 subject-derived clones (Column A). Inclusion of an unlabeled 39 bp healing forward primer with sequence identical to pNL4-3 at 5 nM (and detection primers at 200 nM) allowed 9 of 13 samples to be amplified and genotyped with the initial pNL4-3 standard design (Column B). Finally, pre-amplification of all 13 clones tested with healing forward and reverse primers produced PCR amplification products that could be correctly genotyped by the standard pNL4-3 based RTx primers (Column C).

Table 12

Sample	Reaction		
	A	B	C
pNL4-3			
7324-1			
7324-4			
10076-4			
7295-1			
4755-1			
6463-13			
7303-3			
1617-1			
35764-2			
29129-2			
52534-2			
56252-1			
12-21			

M. Detection of HIV-1 Mutations from Additional Viral Isolates

[0196] Two additional panels of 20 wild-type patient viral RNA isolates were tested (data courtesy of Gilead Sciences). The first panel contained confirmed wild-type K65 isolates, the second panel contained confirmed wild-type M148 isolates. Some of these samples had low viral loads as determined by Amplicor (Amplicor® HIV Monitor, Roche, Indianapolis, IN). Additionally, complete population sequences of the target regions were known.

[0197] Amplification conditions using the LightCycler were as follows:

2 minutes at 95°C, followed by 3 cycles of 5s at 95°C, 5s at 61°C, 1°C/sec, 10s at 72°C, followed by one cycle of (5s at 95°C, 5s at 45°C, 1°C/sec 20s at 72°C, followed by 110 cycles of 5s at 95°C, 5s at 55°C, 20s at 72°C, melt 60°C to 95°C 0.4°C STEP.

[0198] Seventeen of 20 K65 viral RNA samples were successfully amplified using the standard K65 assay design. The failed samples had viral loads ranging from 3,900 to < 400 copies per mL. The three samples that failed to amplify with standard primer designs had viral loads ranging from 3900 to less than 400 copies per mL. All 17 genotypes were determined to be 100% K65K (wild-type).

[0199] Sixteen of 20 M184 viral RNA samples were successfully amplified using the standard M184 assay design. By including a forward healing primer into the M184V

reactions as described above, 19 of 20 samples were genotyped. The sample that failed had a very low viral load of < 400 copies per mL. Additionally, all of the sample K65 and M184 genotype results were correct, that is, seventeen of seventeen K65 and nineteen of nineteen M184 results were found to be 100% concordant with the previously determined genotypes.

[0200] It will be readily apparent to one skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention. The invention illustratively described herein suitably may be practiced in the absence of any element or elements, limitation or limitations which is not specifically disclosed herein. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention. Thus, it should be understood that although the present invention has been illustrated by specific embodiments and optional features, modification and/or variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention.

[0201] In addition, where features or aspects of the invention are described in terms of Markush groups or other grouping of alternatives, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group or other group.

[0202] Also, unless indicated to the contrary, where various numerical values are provided for embodiments, additional embodiments are described by taking any 2 different values as the endpoints of a range. Such ranges are also within the scope of the described invention.

CLAIMS:

1. A method for detecting a specific mutation in a heterogeneous population of polynucleotides, wherein the specific mutation, if present, is located in a first region of the polynucleotides of the population, the method comprising:

(a) amplifying the polynucleotides of the population with a first set of primers to obtain an amplification product that comprises the first region, wherein at least one primer of the first set of primers is a healing primer that specifically hybridizes to a region of the heterogeneous population of polynucleotides that comprises one or more additional mutations, and none of the primers of the first set of primers hybridizes to the first region; and

(b) amplifying the amplification product with a second set of primers to detect the specific mutation, wherein the second set of primers comprises a first primer having a nucleotide that is complementary to the first region if the specific mutation is present and the first primer includes at least one non-natural base, wherein the first primer specifically hybridizes to a region of the amplification product that is the complement of at least a portion of the healing primer, and further wherein one or more of the primers of the second set of primers comprises a label;

wherein the detection of a labeled amplicon indicates the presence of the specific mutation.

2. The method of claim 1, wherein the second set of primers further comprises a second primer having a nucleotide that is complementary to the first region if the mutation is absent and the second primer includes at least one non-natural base.

3. The method of claim 2, wherein the at least one non-natural base is selected from iC and iG.

4. The method of claim 3, wherein the first and second primers of the second primer set comprise the same non-natural base.


5. The method of claim 3, wherein the first and second primers of the second primer set comprise different non-natural bases.
6. The method of claim 2, wherein the first and second primers of the second primer set include a fluorophore.
7. The method of claim 6, wherein the first and second primers of the second primer set include different fluorophores.
8. The method of claim 7, wherein the amplification product is amplified in a reaction mixture that includes a non-natural nucleotide that is capable of base-pairing with the non-natural base of the first and second primers and is coupled to a quencher that is capable of quenching the different fluorophores.
9. The method of any one of claims 1 to 8, wherein the first primer of the second set of primers is fully complementary to the first region if the specific mutation is present, other than at the non-natural base of the first primer.
10. The method of any one of claims 1 to 9, wherein the specific mutation comprises a single base change.
11. The method of any one of claims 1 to 10, wherein the first region comprises a single nucleotide.
12. The method of any one of claims 1 to 11, wherein at least one primer of the first set of primers includes at least one non-natural base.
13. The method of any one of claims 1 to 12, wherein detecting occurs during amplification.
14. The method of any one of claims 1 to 13, wherein the population of polynucleotides comprises a nucleic acid sequence of HIV-1.

1 / 8





FIGURE 1**SEQ ID NO:1**

FFREDLAFPOGKAREFSSEQTRANSPTRRELQVWGRDNNLSLSEAGADRQGTVSFSFP
QITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMNLPGRWKPKMIGGIGGFIKVGQ
YDQILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPISPIETVPVKLKPGM
DGPVKQWPLTEEKIKALVEICTEMEKEGKISKIGPENPYNTPVFAIKKKDSTKWRK
LVDFRELNKRTQDFWEVQLGIPHPAGLKQKKSVTVLDVGDAYFSVPLDKDFRKYTAF
TIPSINNETPGIRYQYNVLPQGWKGSPAIFQCSMTKILEPFRKQNPDIVIYQYMDDL
YVGSDLEIGQHRTKIEELRQHLLRWGFTTPDKKHQKEPPFLWMGYELHPDKWTVQPI
VLPEKDSWTVNDIQKLVGKLNWASQIYAGIKVRQLCKLLRGTKALTEVVPLTEEAEL
ELAENREILKEPVHGVYYDPSKDLIAEIQKQGQGQWTYQIYQEPFKNLKTGKYARMK
GAHTNDVKQLTEAVQKIATESIVIWGKTPKFKLPIQKETWEAWWTEYWQATWIPEWE
FVNTPLVLKWLWYQLEKEPIIGAETFYVDGAANRETKLGKAGYVTDRGRQKVPLTDT
TNQKTELQAIHLALQDSGLEVNIVTDSQYALGIIQAQPKSESELVSQIIEQLIKKE
KVYLAWVPAHKGIGGNEQVDGLVSAGIRKVLFLDGIDKAQEEHEKYHSNWRAMASDF
NLPPVVAKEIVASCDKCQLKGEAMHGQVDCSPGIWQLDCTHLEGKVILVAVHVASGY
IEAEVIPAETGQETAYFLLKLAGRWPVKTVHTDNGSNFTSTTVKAACWWAGIKQEFG
IPYNPQSQGVIESMNKELKKIIGQVRDQAEHLKTAVQMAVFIHNFKRKGGIGGYSAG
ERIVDIIATDIQTKELQKQITKIQNFRVYYRDSRDPVWKGPALLWKGEAVVIQDN
SDIKVVP RRKAKIIRDYGKQ MAGDDCVASRQDED





FIGURE 2



3 / 8


FIGURE 3A**A1****A2****B1****B2**

4 / 8


FIGURE 3B**A3****A4****B3****B4**

5 / 8

FIGURE 4




6 / 8

FIGURE 5

7 / 8

FIGURE 6



8 / 8

FIGURE 7