




(86) Date de dépôt PCT/PCT Filing Date: 2009/04/21
(87) Date publication PCT/PCT Publication Date: 2009/10/29
(85) Entrée phase nationale/National Entry: 2010/10/19
(86) N° demande PCT/PCT Application No.: US 2009/041320
(87) N° publication PCT/PCT Publication No.: 2009/132050
(30) Priorités/Priorities: 2008/04/21 (US61/046,543);
2008/04/29 (US61/048,878); 2008/05/14 (US61/127,713);
2008/05/23 (US61/055,625); 2008/06/10 (US61/060,425);
2008/06/18 (US61/073,716); 2008/06/20 (US61/074,583);
2008/06/27 (US61/076,567); 2008/06/27 (US61/076,576);
2008/07/21 (US61/082,450); 2008/07/25 (US61/083,830);
2008/07/25 (US61/083,871); 2008/08/04 (US61/086,094);
2008/08/04 (US61/086,105); ...


(51) Cl.Int./Int.Cl. *A61K 39/395* (2006.01),
A61K 31/42 (2006.01), *A61K 31/423* (2006.01),
A61K 31/425 (2006.01), *A61K 31/426* (2006.01),
A61K 31/433 (2006.01), *A61K 36/38* (2006.01),
A61K 47/34 (2006.01), *A61K 9/06* (2006.01),
A61P 27/16 (2006.01)
(71) Demandeurs/Applicants:
OTONOMY, INC., US;
THE REGENTS OF THE UNIVERSITY OF CALIFORNIA,
US
(72) Inventeurs/Inventors:
LICHTER, JAY, US;
VOLLRATH, BENEDIKT, US;

(54) Titre : PREPARATIONS AURICULAIRES DE TRAITEMENT DE MALADIES ET ETATS OTIQUES
(54) Title: AURIS FORMULATIONS FOR TREATING OTIC DISEASES AND CONDITIONS



(57) Abrégé/Abstract:

Disclosed herein are compositions and methods for the treatment of otic disorders with immunomodulating agents and auris pressure modulators. In these methods, the auris compositions and formulations are administered locally to an individual afflicted



(30) **Priorités(suite)/Priorities(continued)**: 2008/08/11 (US61/087,951); 2008/08/11 (US61/087,905); 2008/08/12 (US61/088,275); 2008/09/04 (US61/094,384); 2008/09/29 (US61/101,112); 2008/12/22 (US61/140,033); 2009/03/13 (US61/160,233)

(72) **Inventeurs(suite)/Inventors(continued)**: TRAMMEL, ANDREW M., US; DURON, SERGIO G., US; PIU, FABRICE, US; DELLAMARY, LUIS A., US; YE, QIANG, US; LEBEL, CARL, US; SCAIFE, MICHAEL CHRISTOPHER, US; HARRIS, JEFFREY P., US

(74) **Agent**: GOWLING LAFLEUR HENDERSON LLP

(57) **Abrégé(suite)/Abstract(continued)**:

with an otic disorder, through direct application of the immunomodulating and/or auris pressure modulating compositions and formulations onto the auris media and/or auris interna target areas, or via perfusion into the auris media and/or auris interna structures.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 October 2009 (29.10.2009)


PCT

(10) International Publication Number
WO 2009/132050 A3

(51) International Patent Classification:			61/087,951	11 August 2008 (11.08.2008)	US
A61K 39/395 (2006.01)	A61K 31/433 (2006.01)		61/088,275	12 August 2008 (12.08.2008)	US
A61P 27/16 (2006.01)	A61K 31/426 (2006.01)		61/094,384	4 September 2008 (04.09.2008)	US
A61K 47/34 (2006.01)	A61K 31/425 (2006.01)		61/101,112	29 September 2008 (29.09.2008)	US
A61K 9/06 (2006.01)	A61K 31/423 (2006.01)		61/140,033	22 December 2008 (22.12.2008)	US
A61K 31/42 (2006.01)	A61K 36/38 (2006.01)		61/160,233	13 March 2009 (13.03.2009)	US
(21) International Application Number:			(71) Applicants (for all designated States except US): OTON- OMY, INC. [US/US]; 5626 Oberlin Drive, Suite 100, San Diego, CA 92121 (US). THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 1111 Franklin Street, 12th Floor, Oakland, CA 94607 (US).		
			PCT/US2009/041320		
(22) International Filing Date:					
			21 April 2009 (21.04.2009)		
(25) Filing Language:			English		
(26) Publication Language:			English		
(30) Priority Data:			(72) Inventors; and		
61/046,543	21 April 2008 (21.04.2008)	US	(75) Inventors/Applicants (for US only): LICHTER, Jay		
61/048,878	29 April 2008 (29.04.2008)	US	[US/US]; P.O. Box 676244, Rancho Santa Fe, CA 92067		
61/127,713	14 May 2008 (14.05.2008)	US	(US). VOLLRATH, Benedikt [DE/US]; 4704 Niagara		
61/055,625	23 May 2008 (23.05.2008)	US	Avenue, San Diego, CA 92107 (US). TRAMMEL, An-		
61/060,425	10 June 2008 (10.06.2008)	US	drew, M. [US/US]; 12485 South Alden Circle, Olathe,		
61/073,716	18 June 2008 (18.06.2008)	US	KS 66062 (US). DURÓN, Sergio, G. [US/US]; 1605		
61/074,583	20 June 2008 (20.06.2008)	US	Neale Street, San Diego, CA 92103 (US). PIU, Fabrice		
61/076,576	27 June 2008 (27.06.2008)	US	[FR/US]; 11859 Stonedale Ct., San Diego, CA 92131		
61/076,567	27 June 2008 (27.06.2008)	US	(US). DELLAMARY, Luis, A. [US/US]; 829 Quiet Hills		
61/082,450	21 July 2008 (21.07.2008)	US	Dr, San Marcos, CA 92069 (US). YE, Qiang [US/US];		
61/083,871	25 July 2008 (25.07.2008)	US	7395 Mannix Court, San Diego, CA 92129 (US).		
61/083,830	25 July 2008 (25.07.2008)	US	LEBEL, Carl [US/US]; 23256 Mariposa De Oro St.,		
61/086,105	4 August 2008 (04.08.2008)	US	Malibu, CA 90265 (US). SCAIFE, Michael, Christo-		
61/086,094	4 August 2008 (04.08.2008)	US	pher [US/US]; 60 Marvin Avenue, Los Altos, CA 94022		
61/087,905	11 August 2008 (11.08.2008)	US	(US). HARRIS, Jeffrey, P. [US/US]; 7717 Sierra Mar		
			Drive, La Jolla, CA 92037 (US).		

[Continued on next page]

(54) Title: AURIS FORMULATIONS FOR TREATING OTIC DISEASES AND CONDITIONS



(57) Abstract: Disclosed herein are compositions and methods for the treatment of otic disorders with immunomodulating agents and auris pressure modulators. In these methods, the auris compositions and formulations are administered locally to an individual afflicted with an otic disorder, through direct application of the immunomodulating and/or auris pressure modulating compositions and formulations onto the auris media and/or auris interna target areas, or via perfusion into the auris media and/or auris interna structures.

WO 2009/132050 A3

WO 2009/132050 A3

(74) **Agents:** AKHAVAN, Ramin et al.; 650 Page Mill Road, Palo Alto, CA 94304 (US).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,

ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

(88) **Date of publication of the international search report:**

14 January 2010

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 143

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 143

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

AURIS FORMULATIONS FOR TREATING OTIC DISEASES AND CONDITIONS**RELATED APPLICATIONS**

[0001] This patent application claims the benefit of U.S. Provisional Application Ser. Nos. 61/087,905, filed on August 11, 2008, 61/055,625 filed on May 23, 2008, 61/086,105 filed on August 04, 2008, 61/073,716 filed on June 18, 2008, 61/140,033 filed on December 22, 2008, 61/127,713 filed on May 14, 2008, 61/101,112 filed on September 29, 2008, 61/094,384 filed on September 04, 2008, 61/074,583 filed on June 20, 2008, 61/060,425 filed on June 10, 2008, 61/048,878 filed on April 29, 2008, 61/046,543 filed on April 21, 2008, 61/076,567 filed on June 27, 2008, 61/076,576 filed on June 27, 2008, 61/160,233 filed on March 13, 2009, 61/086,094 filed on August 04, 2008, 61/083,830 filed on July 25, 2008, 61/083,871 filed on July 25, 2008, 61/087,951 filed on August 11, 2008, 61/088,275 filed on August 12, 2008, the disclosures of all of which are herein incorporated by reference in their entirety.

BACKGROUND OF THE INVENTION

[0002] Described herein are formulations for enhanced drug delivery into the external, middle and/or inner ear, including the cochlea and vestibular labyrinth; preferably with little or no systemic release of the drug.

SUMMARY OF THE INVENTION

[0003] The auris formulations and therapeutic methods described herein have numerous advantages that overcome the previously-unrecognized limitations of formulations and therapeutic methods described in prior art.

Sterility

[0004] The environment of the inner ear is an isolated environment. The endolymph and the perilymph are static fluids and are not in contiguous contact with the circulatory system. The blood – labyrinth – barrier (BLB), which includes a blood-endolymph barrier and a blood-perilymph barrier, consists of tight junctions between specialized epithelial cells in the labyrinth spaces (i.e., the vestibular and cochlear spaces). The presence of the BLB limits delivery of active agents (e.g., immunomodulators, aural pressure modulators, antimicrobials) to the isolated microenvironment of the inner ear. Auris hair cells are bathed in endolymphatic or perilymphatic fluids and cochlear recycling of potassium ions is important for hair cell function. When the inner ear is infected, there is an influx of leukocytes and/or immunoglobins (e.g. in response to a microbial infection) into the endolymph and/or the perilymph and the delicate ionic composition of inner ear fluids is upset by the influx of leukocytes and/or immunoglobins. In certain instances, a change in the ionic composition of inner ear fluids results in hearing loss, loss of balance and/or ossification of auditory structures. In certain instances, even trace amounts of pyrogens and/or microbes can trigger infections and related physiological changes in the isolated microenvironment of the inner ear.

[0005] Due to the susceptibility of the inner ear to infections, auris formulations require a level of sterility that has not been recognized hitherto in prior art. Provided herein are auris formulations that are sterilized with stringent sterility requirements and are suitable for administration to the middle and/or inner ear. In some embodiments, the auris compatible compositions described herein are substantially free of pyrogens and/or microbes.

Compatibility with Inner Ear Environment

[0006] Described herein are otic formulations with an ionic balance that is compatible with the perilymph and/or the endolymph and does not cause any change in cochlear potential. In specific embodiments, osmolarity/osmolality of the present formulations is adjusted, for example, by the use of appropriate salt concentrations (e.g., concentration of sodium salts) or the use of tonicity agents which renders the formulations endolymph-compatible and/or perilymph-compatible (i.e. isotonic with the endolymph and/or perilymph). In some instances, the endolymph-compatible and/or perilymph-compatible formulations described herein cause minimal disturbance to the environment of the inner ear and cause minimum discomfort (e.g, vertigo) to a mammal (e.g., a human) upon administration. Further, the formulations comprise polymers that are biodegradable and/or dispersable, and/or otherwise non-toxic to the inner ear environment. In some embodiments, the formulations described herein are free of preservatives and cause minimal disturbance (e.g., change in pH or osmolarity, irritation) in auditory structures. In some embodiments, the formulations described herein comprise antioxidants that are non-irritating and/or non-toxic to otic structures.

Dosing Frequency

[0007] The current standard of care for auris formulations requires multiple administrations of drops or injections (e.g. intratympanic injections) over several days (e.g., up to two weeks), including schedules of receiving multiple injections per day. In some embodiments, auris formulations described herein are controlled release formulations, and are administered at reduced dosing frequency compared to the current standard of care. In certain instances, when an auris formulation is administered via intratympanic injection, a reduced frequency of administration alleviates discomfort caused by multiple intratympanic injections in individuals undergoing treatment for a middle and/or inner ear disease, disorder or condition. In certain instances, a reduced frequency of administration of intratympanic injections reduces the risk of permanent damage (e.g., perforation) to the ear drum. The formulations described herein provide a constant, sustained, extended, delayed or pulsatile rate of release of an active agent into the inner ear environment and thus avoid any variability in drug exposure in treatment of otic disorders.

Therapeutic Index

[0008] Auris formulations described herein are administered into the ear canal, or in the vestibule of the ear. Access to, for example, the vestibular and cochlear apparatus will occur through the auris media including the round window membrane, the oval window/stapes footplate, the annular ligament and through the otic capsule/temporal bone. Otic administration of the formulations described herein avoids toxicity associated with systemic administration (e.g., hepatotoxicity, cardiotoxicity, gastrointestinal side effects, renal toxicity) of the active agents. In some instances, localized administration in the ear allows an active agent to reach a target organ (e.g., inner ear) in the absence of systemic accumulation of the active agent. In some instances, local administration to the ear provides a higher therapeutic index for an active agent that would otherwise have dose-limiting systemic toxicity.

Prevention of Drainage into Eustachian Tube

[0009] In some instances, a disadvantage of liquid formulations is their propensity to drip into the eustachian tube and cause rapid clearance of the formulation from the inner ear. Provided herein, in certain embodiments, are auris formulations comprising polymers that gel at body temperature and remain in contact with the target auditory surfaces (e.g., the round window) for extended periods of time. In some embodiments, the formulations further comprise mucoadhesives that allow the formulations to adhere to otic mucosal surfaces. In some instances, the auris formulations described herein avoid attenuation of therapeutic benefit due to drainage or leakage of active agents via the eustachian tube.

Description of Certain Embodiments

[0010] Accordingly, provided herein, in some embodiments, are pharmaceutical formulations for use in the treatment of an otic disease or condition formulated to provide a therapeutically effective amount of an immunomodulating agent across the round window membrane into the cochlea, the formulation comprising:

between about 0.2% to about 6% by weight of an immunomodulating agent, or pharmaceutically acceptable prodrug or salt thereof;

between about 16% to about 21% by weight of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106;

sterile water, q.s., buffered to provide a perilymph-suitable pH between about 6.0 and about 7.6;

and substantially low degradation products of the immunomodulating agent;

wherein the pharmaceutical formulation has a perilymph-suitable osmolarity between about 250 and 320 mOsm/L,

less than about 50 colony forming units (cfu) of microbiological agents per gram of formulation, and

less than about 5 endotoxin units (EU) per kg of body weight of a subject.

[0011] Provided herein, in some embodiments, are pharmaceutical formulations for use in the treatment of an otic disease or condition formulated to provide a therapeutically effective amount of an immunomodulating agent across the round window membrane into the cochlea, the formulation comprising:

- between about 0.1 mg/mL to about 70 mg/mL of an immunomodulating agent, or pharmaceutically acceptable prodrug or salt thereof;
- between about 16% to about 21% by weight of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106;
- sterile water, q.s., buffered to provide a perilymph-suitable pH between about 6.0 and about 7.6;
- and substantially low degradation products of the immunomodulating agent;

wherein the pharmaceutical formulation has a perilymph-suitable osmolarity between about 250 and 320 mOsm/L,

less than about 50 colony forming units (cfu) of microbiological agents per gram of formulation, and less than about 5 endotoxin units (EU) per kg of body weight of a subject.

[0012] In some embodiments, the immunomodulating agent is released from the formulation for a period of at least 3 days. In some embodiments, the pharmaceutical formulation is an auris-acceptable thermoreversible gel. In some embodiments, the polyoxyethylene-polyoxypropylene triblock copolymer is biodegradable. In some embodiments, the formulations further comprise a mucoadhesive. In some embodiments, the formulations further comprise a penetration enhancer. In some embodiments, the formulations further comprise a thickening agent. In some embodiments, the formulations further comprise a dye.

[0013] In further embodiments, provided herein are formulations further comprising a drug delivery device selected from a needle and syringe, a pump, a microinjection device, a wick, an in situ forming spongy material or combinations thereof.

[0014] In some embodiments of the formulations described herein, the immunomodulating agent, or pharmaceutically acceptable salt thereof, has limited or no systemic release, systemic toxicity, poor PK characteristics, or combinations thereof. In some embodiments, the immunomodulating agent is in the form of a free base, salt, a prodrug, or a combination thereof. In some embodiments, the immunomodulating agent comprises multiparticulates. In some embodiments, the immunomodulating agent is essentially in the form of micronized particles.

[0015] In some embodiments, the immunomodulating agent is an anti-TNF agent, a calcineurin inhibitor, an IKK inhibitor, an interleukin inhibitor, a TNF- α converting enzyme (TACE) inhibitor, or a toll-like receptor inhibitor.

[0016] In some embodiments, the formulations further comprise an immunomodulating agent, or pharmaceutically acceptable salt thereof, as an immediate release agent.

[0017] In some embodiments, the formulations described herein further comprise an additional therapeutic agent. In some embodiments, the additional therapeutic agent is a Na/K ATPase modulator, a chemotherapeutic agent, a collagen, a gamma-globulin, an interferon, an anti-microbial agent, an antibiotic, a local acting anesthetic agent, a platelet activator factor antagonist, a nitric oxide synthase inhibitor, an anti-vertigo agent, a vasopressin antagonist, an anti-viral, an anti-emetic agent or combinations thereof.

[0018] In some embodiments, the pH of the composition is between about 6.0 to about 7.6. In some embodiments, the ratio of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106 to a thickening agent is from about 40:1 to about 10:1. In some embodiments, the thickening agent is carboxymethyl cellulose.

[0019] In some embodiments, the otic disease or condition is Meniere's disease, sudden sensorineural hearing loss, noise induced hearing loss, age-related hearing loss, auto immune ear disease or tinnitus.

[0020] Also provided herein is a method of treating an otic disease or condition comprising administering to an individual in need thereof an intratympanic composition comprising

between about 0.2% to about 6% by weight of an immunomodulating agent, or pharmaceutically acceptable prodrug or salt thereof;

between about 16% to about 21% by weight of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106;

sterile water, q.s., buffered to provide a perilymph-suitable pH between about 6.0 and about 7.6;

and substantially low degradation products of the immunomodulating agent;

wherein the pharmaceutical formulation has a perilymph-suitable osmolarity between about 250 and 320 mOsm/L,

less than about 50 colony forming units (cfu) of microbiological agents per gram of formulation, and less than about 5 endotoxin units (EU) per kg of body weight of a subject.

[0021] In some embodiments of the method, the immunomodulating agent is an anti-TNF agent, a calcineurin inhibitor, an IKK inhibitor, an interleukin inhibitor, a TNF- α converting enzyme (TACE) inhibitor, or a toll-like receptor inhibitor. In some embodiments of the method, the immunomodulating agent is released from the composition for a period of at least 3 days. In some embodiments of the method, the composition is administered across the round window. In some embodiments of the method, the otic disease or condition is Meniere's disease, sudden sensorineural

hearing loss, age-related hearing loss, noise induced hearing loss, auto immune ear disease or tinnitus.

[0022] Also provided herein, in some embodiments, are pharmaceutical formulations for use in the treatment of an otic disease or condition formulated to provide a therapeutically effective amount of an aural pressure modulating agent across the round window membrane into the cochlea, the formulation comprising:

between about 0.2% to about 6% by weight of an aural pressure modulating agent, or pharmaceutically acceptable prodrug or salt thereof;

between about 16% to about 21% by weight of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106;

sterile water, q.s., buffered to provide a perilymph-suitable pH between about 6.0 and about 7.6;

substantially low degradation of the aural pressure modulating agent;

wherein the pharmaceutical formulation has a perilymph-suitable osmolarity between about 250 and 320 mOsm/L,

less than about 50 colony forming units (cfu) of microbiological agents per gram of formulation, and less than about 5 endotoxin units (EU) per kg of body weight of a subject.

[0023] Also provided herein, in some embodiments, are pharmaceutical formulations for use in the treatment of an otic disease or condition formulated to provide a therapeutically effective amount of an aural pressure modulating agent across the round window membrane into the cochlea, the formulation comprising:

between about 0.1 mg/mL to about 70 mg/mL of an aural pressure modulating agent, or pharmaceutically acceptable prodrug or salt thereof;

between about 16% to about 21% by weight of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106;

sterile water, q.s., buffered to provide a perilymph-suitable pH between about 6.0 and about 7.6;

and substantially low degradation products of the aural pressure modulating agent;

wherein the pharmaceutical formulation has a perilymph-suitable osmolarity between about 250 and 320 mOsm/L,

less than about 50 colony forming units (cfu) of microbiological agents per gram of formulation, and less than about 5 endotoxin units (EU) per kg of body weight of a subject.

[0024] In some embodiments, the aural pressure modulating agent is released from the formulation for a period of at least 3 days. In some embodiments, the pharmaceutical formulation is an auris-acceptable thermoreversible gel. In some embodiments, the polyoxyethylene-polyoxypropylene

triblock copolymer is biodegradable. In some embodiments, the formulations further comprise a round window membrane mucoadhesive. In some embodiments, the formulations further comprise a round window membrane penetration enhancer. In some embodiments, the formulations further comprise thickening agent. In some embodiments, the formulations further comprise a dye.

5 [0025] In some embodiments of the formulations described herein, the formulations further comprise a drug delivery device selected from a needle and syringe, a pump, a microinjection device, a wick, an in situ forming spongy material or combinations thereof.

[0026] In some embodiments, the aural pressure modulating agent, or pharmaceutically acceptable salt thereof, has limited or no systemic release, systemic toxicity, poor PK characteristics, or
10 combinations thereof. In some embodiments, the aural pressure modulating agent is administered in the form of a free base, salt, a prodrug, or a combination thereof. In some embodiments, the aural pressure modulating agent comprises multiparticulates. In some embodiments, the aural pressure modulating agent is essentially in the form of micronized particles.

[0027] In some embodiments, the aural pressure modulating agent is a modulator of aquaporin, an
15 estrogen related receptor beta modulator, a gap junction protein modulator, an NMDA receptor modulator, an osmotic diuretic, a progesterone receptor modulator, a prostaglandin modulator, or a vasopressin receptor modulator.

[0028] In some embodiments, the formulations described herein further comprise an aural pressure modulating agent, or pharmaceutically acceptable salt thereof, as an immediate release agent.

20 [0029] In some embodiments, the formulations described herein further comprise an additional therapeutic agent. In some embodiments, the additional therapeutic agent is Na/K ATPase modulator, a chemotherapeutic agent, a collagen, a gamma-globulin, an interferon, an anti-microbial agent, an antibiotic, a local acting anesthetic agent, a platelet activator factor antagonist, a nitric oxide synthase inhibitor, an anti-vertigo medicine, a vasopressin antagonist, an anti-viral, an anti-
25 emetic agent or combinations thereof.

[0030] In some embodiments, the pH of the composition is between about 6.0 to about 7.6. In some embodiments, the ratio of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106 to a thickening agent is from about 40:1 to about 10:1. In some embodiments, the thickening agent is carboxymethyl cellulose.

30 [0031] In some embodiments, the otic disease or condition is Meniere's disease, sudden sensorineural hearing loss, age-related hearing loss, noise induced hearing loss, auto immune ear disease or tinnitus.

[0032] Also provided herein is a method of treating an otic disease or condition comprising administering to an individual in need thereof an intratympanic composition comprising

between about 0.2% to about 6% by weight of an aural pressure modulating agent, or pharmaceutically acceptable prodrug or salt thereof;

between about 16% to about 21% by weight of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106;

sterile water, q.s., buffered to provide a perilymph-suitable pH between about 6.0 and about 7.6;

and substantially low degradation products of the aural pressure modulating agent;

wherein the pharmaceutical formulation has a perilymph-suitable osmolarity between about 250 and 320 mOsm/L,

less than about 50 colony forming units (cfu) of microbiological agents per gram of formulation, and less than about 5 endotoxin units (EU) per kg of body weight of a subject.

[0033] In some embodiments, the aural pressure modulating agent is a modulator of aquaporin, an estrogen related receptor beta modulator, a gap junction protein modulator, an NMDA receptor modulator, an osmotic diuretic, a progesterone receptor modulator, a prostaglandin modulator, or a vasopressin receptor modulator.

[0034] In some embodiments of the method, the aural pressure modulating agent is released from the composition for a period of at least 3 days. In some embodiments of the method, the composition is administered across the round window.

[0035] In some embodiments of the method, the otic disease or condition is Meniere's disease, sudden sensorineural hearing loss, age-related hearing loss, noise induced hearing loss, auto immune ear disease or tinnitus.

[0036] In any of the aforementioned embodiments, the term "substantially low degradation products" means less than 5% by weight of the active agent are degradation products of the active agent. In further embodiments, the term means less than 3% by weight of the active agent are degradation products of the active agent. In yet further embodiments, the term means less than 2% by weight of the active agent are degradation products of the active agent. In further embodiments, the term means less than 1% by weight of the active agent are degradation products of the active agent.

[0037] Other objects, features, and advantages of the methods and compositions described herein will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments, are given by way of illustration only.

BRIEF DESCRIPTION OF THE FIGURES

[0038] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by

reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

[0039] **Figure 1** illustrates a comparison of non-sustained release formulations and sustained release formulations.

5 [0040] **Figure 2** illustrates the effect of concentration on viscosity of aqueous solutions of Blanos refined CMC

[0041] **Figure 3** illustrates the effect of concentration on viscosity of aqueous solutions of Methocel

DETAILED DESCRIPTION

10 [0042] Systemic administration of active agents is, in some instances, ineffectual in the treatment of diseases that affect inner ear structures. The cochlear canals and the cochlea, for example, are isolated from the circulatory system limiting systemic delivery of active agents to target sites in the inner ear. In some instances, systemic drug administration creates a potential inequality in drug concentration with higher circulating levels in the serum, and lower levels in the target auris interna
15 organ structures. In certain instances, large amounts of drug are required to overcome this inequality in order to deliver sufficient, therapeutically effective quantities of a drug to auditory structures. In some instances, systemic drug administration also increases the likelihood of secondary systemic accumulation and consequent adverse side effects.

[0043] Currently available treatment for inner ear diseases also carries the risk of attendant side
20 effects. For example, available methods require multiple daily doses (e.g., intratympanic injection or infusion) of drugs. In certain instances, multiple daily intratympanic injections cause patient discomfort and non-compliance. In certain instances, delivery of active agents to the inner ear via otic drops administered in the ear canal or via intratympanic injection is hindered by the biological barrier presented by the blood-labyrinth-barrier (BLB), the oval window membrane and/or the round
25 window membrane. In some instances, delivery of active agents to the inner ear via otic drops or intratympanic injection causes osmotic imbalance in inner ear structures, introduces infections or other immune disorders as a result of microbial or endotoxin presence, or results in permanent structural damage (e.g. perforation of the tympanic membrane), resulting in hearing loss and the like.

30 [0044] Clinical studies with steroids such as prednisolone or dexamethasone have demonstrated the benefit of having long term exposure of the steroids to the perilymph of the cochlea; this has been shown by improved clinical efficacy in improving sudden hearing loss when the steroid in question is given on multiple occasions.

[0045] U.S. Application Publication Nos. 2006/0063802 and 2005/0214338 disclose compositions
35 comprising arylecycloalkylamine NMDA antagonists for local administration to the inner ear. There is no disclosure of controlled release formulations, osmolality or pH requirements, or sterility

requirements for the compositions. WO 2007/038949 discloses compositions comprising arylecycloalkylamine NMDA antagonists in the treatment of inner ear disorders. No guidance is provided on pyrogenicity, sterility requirements, viscosity levels and/or controlled release characteristics of the formulation.

5 [0046] Fernandez et al. *Biomaterials*, 26: 3311-3318 (2005) describes compositions which comprise prednisolone useful to treat inner ear disease such as Meniere's disease. Fernandez et al. do not disclose osmolarity, pyrogenicity, pH, or sterility levels of the compositions described therein. Paulson et al. *The Laryngoscope*, 118: 706 (2008) describe sustained release compositions which comprise dexamethasone useful in treatment of, *inter alia*, inner ear diseases such as
10 Meniere's disease. Again, Paulson et al. do not disclose osmolarity, pyrogenicity, pH, or sterility requirements for the compositions described therein.

[0047] C. Gang et al., *J. Sichuan Univ.* 37:456-459 (2006) describe a dexamethasone sodium phosphate (DSP) preparation. The formulation described in Gang et al. comprises preservatives and adhesives and is sterilized under conditions that likely lead to breakdown of DSP. There is also no
15 disclosure regarding osmolarity, pyrogenicity, pH, or sterility requirements for the compositions described therein.

[0048] Feng et al., *Zhonghua Er Bi Yan Hou Tou Jing Wai Ke Za Zhi* 42:443-6 (June 2007) and Feng et al., *Zhonghua Yi Xue Za Zhi* 87:2289-91 (August 2007) describe 20% and 25% poloxamer 407 solutions as non-toxic to otic structures. There is no active agent in the solutions described
20 therein, and there is no disclosure regarding osmolarity, pyrogenicity, pH, or sterility requirements for the solutions described therein. J. Daijie et al., *J. Clin. Otorhinolaryngol Head Neck Surg* (China) 22(7) (April 2008), P. Yikun et al., *J. Clin. Otorhinolaryngol Head Neck Surg* (China) 22(10) (May 2008), and S. Wandong et al., *J. Clin. Otorhinolaryngol Head Neck Surg* (China) 22(19) (October 2008) describe intratympanic solution injections. However, Daijie et al, Yikun et al.
25 and Wandong et al. do not disclose any otic formulations that are polymer based, or any otic formulations that are sustained release formulations. There is also no disclosure regarding osmolarity, pyrogenicity, pH, or sterility requirements for the compositions described therein.

[0049] Intratympanic injection of therapeutic agents is the technique of injecting a therapeutic agent behind the tympanic membrane into the auris media and/or auris interna. Despite early success with
30 this technique (Schuknecht, *Laryngoscope* (1956) 66, 859-870) some challenges do remain. For example, access to the round window membrane, the site of drug absorption into the auris interna, can be challenging.

[0050] However, intra-tympanic injections create several *unrecognized* problems not addressed by currently available treatment regimens, such as changing the osmolarity and pH of the perilymph
35 and endolymph, and introducing pathogens and endotoxins that directly or indirectly damage inner ear structures. One of the reasons the art may not have recognized these problems is that there are no

approved intra-tympanic compositions: the inner ear provides sui generis formulation challenges. Thus, compositions developed for other parts of the body have little to no relevance for an intra-tympanic composition.

[0051] There is no guidance in the prior art regarding requirements (e.g., level of sterility, pH, osmolarity) for otic formulations that are suitable for administration to humans. There is wide anatomical disparity between the ears of animals across species. A consequence of the inter-species differences in auditory structures is that animal models of inner ear disease are often unreliable as a tool for testing therapeutics that are being developed for clinical approval.

[0052] Provided herein are otic formulations that meet stringent criteria for pH, osmolarity, ionic balance, sterility, endotoxin and/or pyrogen levels. The auris compositions described herein are compatible with the microenvironment of the inner ear (e.g., the perilymph) and are suitable for administration to humans. In some embodiments, the formulations described herein comprise dyes and aid visualization of the administered compositions obviating the need for invasive procedures (e.g., removal of perilymph) during preclinical and/or clinical development of intratympanic therapeutics.

[0053] Accordingly, provided herein, in certain embodiments, are controlled release auris-acceptable formulations and compositions that locally treat auris target structures and provide extended exposure of otic active agents to the target auris structures. In certain embodiments, the auris formulations described herein are polymer based formulations designed for stringent osmolarity and pH ranges that are compatible with auditory structures and/or the endolymph and perilymph. In some embodiments, the formulations described herein are controlled release formulations that provide extended release for a period of at least 3 days and meet stringent sterility requirements. In some instances, otic compositions described herein contain lower endotoxin levels (e.g. < 0.5 EU/mL when compared to typically acceptable endotoxin levels of 0.5 EU/mL. In some instances, the otic formulations described herein contain low levels of colony forming units (e.g., <50 CFUs) per gram of the formulation. In some instances, the auris formulations described herein are substantially free of pyrogens and/or microbes. In some instances the auris formulations described herein are formulated to preserve the ionic balance of the endolymph and/or the perilymph. The stringent requirement for sterility and compatibility with inner ear fluids for otic formulations has not been addressed hereto.

[0054] The formulations described herein represent an advantage over currently available therapeutics because they are sterile controlled release otic formulations that are compatible with auris structures (e.g., the perilymph) and are safe for long term administration to humans in need thereof. In some instances, by providing a slow extended release of an active agent, the formulations described herein prevent an initial burst release upon administration to the inner ear; i.e., the

formulations avoid causing a dramatic change in the pH of the endolymph or perilymph and subsequently reduce the impact on balance and/or hearing upon administration.

[0055] In some instances, local administration of the compositions described herein avoids potential adverse side effects as a result of systemic administration of active agents. In some instances, the locally applied auris-acceptable formulations and compositions described herein are compatible with auris structures, and administered either directly to the desired auris structure, e.g. the cochlear region, or administered to a structure in direct communication with areas of the auris structure; in the case of the cochlear region, for example, including but not limited to the round window membrane, the crista fenestrae cochleae or the oval window membrane.

[0056] In certain instances, an advantage of the controlled release formulations described herein is that they provide a constant rate of release of a drug from the formulation and provide a constant prolonged source of exposure of an otic active agent to the inner ear of an individual or patient suffering from an otic disorder, reducing or eliminating any variabilities associated with other methods of treatment (such as, e.g., otic drops and/or multiple intratympanic injections).

[0057] The drug formulations described herein provide extended release of the active ingredient(s) into the middle and/or inner ear (*auris interna*), including the cochlea and vestibular labyrinth. A further option includes an immediate or rapid release component in combination with a controlled release component.

Certain Definitions

[0058] The term "auris-acceptable" with respect to a formulation, composition or ingredient, as used herein, includes having no persistent detrimental effect on the auris media (or middle ear) and the auris interna (or inner ear) of the subject being treated. By "auris-pharmaceutically acceptable," as used herein, refers to a material, such as a carrier or diluent, which does not abrogate the biological activity or properties of the compound in reference to the auris media (or middle ear) and the auris interna (or inner ear), and is relatively or is reduced in toxicity to the auris media (or middle ear) and the auris interna (or inner ear), i.e., the material is administered to an individual without causing undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

[0059] As used herein, amelioration or lessening of the symptoms of a particular otic disease, disorder or condition by administration of a particular compound or pharmaceutical composition refers to any decrease of severity, delay in onset, slowing of progression, or shortening of duration, whether permanent or temporary, lasting or transient that is attributed to or associated with administration of the compound or composition.

[0060] As used herein, the terms "immunomodulating agent" or "immunomodulator" or "immunomodulator agent" or "immune-modulating agent" are used as synonyms.

[0061] The term “anti-TNF agent” or “anti tumor necrosis factor agent” or “TNF modulator” or “TNF modulating agent” or “TNF-alpha modulator” or “anti-TNF alpha agent” are used as synonyms. The term “anti-TNF agent” and its synonyms generally refer to agents that counteract the biological effect of TNF- α or the biological effect of pro-TNF- α stimulus including agents which bind to and antagonize the molecular target; here, tumor necrosis factor alpha or TNF-alpha (TNF- α), agents which inhibit release of TNF- α , or agents which interfere with TNF- α gene expression due to pro-TNF- α stimulus. Also included are agents that indirectly antagonize the biological activity of TNF- α by modulating targets in the general pathway of TNF- α activation, including but not limited to targets upstream of the pathway of TNF-alpha activation, including but not limited to agents which increase TNF-alpha expression, activity or function.

[0062] As used herein, the terms “aural pressure modulating agent” or “aural pressure modulator” are used as synonyms and do not define the degree of efficacy. The aural pressure modulator also includes compounds that modulate the expression or post-transcriptional processing of a fluid homeostasis protein, including vasopressin and estrogen-related receptor beta protein. Additionally, vasopressin receptor or estrogen-related receptor beta modulators include compounds that influence vasopressin receptor or estrogen-related receptor beta signalling or downstream functions under the control of the vasopressin receptor or estrogen-related receptor beta, such as aquaporin function. Vasopressin receptor or estrogen-related receptor beta modulating agents includes compounds that increase and/or decrease vasopressin receptor or estrogen-related receptor beta function, including antagonists, inhibitors, agonists, partial agonists and the like.

[0063] “Modulator of neuron and/or hair cells of the auris” and “auris sensory cell modulating agent” are synonyms. They include agents that promote the growth and/or regeneration of neurons and/or the hair cells of the auris, and agents that destroy neurons and/or hair cells of the auris.

[0064] As used herein, the term “antimicrobial agent” refers to compounds that inhibit the growth, proliferation, or multiplication of microbes, or that kill microbes. Suitable “antimicrobial agents” are antibacterial agents (effective against bacteria), antiviral agents (effective against viruses), antifungal agents (effective against fungi), antiprotozoal (effective against protozoa), and/or antiparasitic to any class of microbial parasites. “Antimicrobial agents” may work by any suitable mechanism against the microbes, including by being toxic or cytostatic.

[0065] The phrase “antimicrobial small molecule” refers to antimicrobial compounds that are of relatively low molecular weight, e.g., less than 1,000 molecular weight, that are effective for the treatment of otic disorders, particularly otic disorders caused by pathogenic microbes, and are suitable for use in the formulations disclosed herein. Suitable “antimicrobial small molecules” include antibacterial, antiviral, antifungal, antiprotozoal, and antiparasitic small molecules.

[0066] "Modulator of free-radicals" and "free-radical modulating agent" are synonyms. They refer to agents that modulate the production of and/or damage caused by free radicals, especially reactive oxygen species.

[0067] As used herein, the terms "ion channel modulating agent", "modulator of ion channels" or "ion channel modulator" are used as synonyms and do not define the degree of efficacy. The ion channel modulator also includes compounds that modulate the expression or post-transcriptional processing of a fluid homeostasis protein, including vasopressin and estrogen-related receptor beta protein. Additionally, vasopressin receptor or estrogen-related receptor beta modulators include compounds that influence vasopressin receptor or estrogen-related receptor beta signalling or downstream functions under the control of the vasopressin receptor or estrogen-related receptor beta, such as aquaporin function. Vasopressin receptor or estrogen-related receptor beta modulating agents includes compounds that increase and/or decrease vasopressin receptor or estrogen-related receptor beta function, including antagonists, inhibitors, agonists, partial agonists and the like.

[0068] As used herein, the term "otic agent" or "otic structure modulating agent" or "otic therapeutic agent" or "otic active agent" or "active agent" refers to compounds that are effective for the treatment of otic disorders, e.g., otitis media, otosclerosis, autoimmune diseases of the ear and cancer of the ear, and are suitable for use in the formulations disclosed herein. An "otic agent" or "otic structure modulating agent" or "otic therapeutic agent" or "otic active agent" or "active agent" includes, but is not limited to, compounds that act as an agonist, a partial agonist, an antagonist, a partial antagonist, an inverse agonist, a competitive antagonist, a neutral antagonist, an orthosteric antagonist, an allosteric antagonist, or a positive allosteric modulator of an otic structure modulating target, or combinations thereof.

[0069] "Balance disorder" refers to a disorder, illness, or condition which causes a subject to feel unsteady, or to have a sensation of movement. Included in this definition are dizziness, vertigo, disequilibrium, and pre-syncope. Diseases which are classified as balance disorders include, but are not limited to, Ramsay Hunt's Syndrome, Meniere's Disease, mal de débarquement, benign paroxysmal positional vertigo, and labyrinthitis.

[0070] "CNS modulator" and "CNS modulating agent" are synonyms. They refer to agents that decrease, diminish, partially suppress, fully suppress, ameliorate, antagonize, agonize, stimulate or increase the activity of the CNS. For example, they may increase the activity of GABA by, for example, increasing the sensitivity of the GABA receptors, or they may alter the depolarization in neurons.

[0071] "Local anesthetic" means a substance which causes a reversible loss of sensation and/or a loss of nociception. Often, these substances function by decreasing the rate of the depolarization and repolarization of excitable membranes (for example, neurons). By way of non-limiting example, local anesthetics include lidocaine, benzocaine, prilocaine, and tetracaine.

[0072] "Modulator of the GABA_A receptor," "modulator of the GABA receptor," "GABA_A receptor modulator," and "GABA receptor modulator," are synonyms. They refer to substances which modulate the activity of the GABA neurotransmitter, by, for example, increasing the sensitivity of the GABA receptor to GABA.

5 [0073] As used herein, the term "cytotoxic agent" refers to compounds that are cytotoxic (i.e., toxic to a cell) effective for the treatment of otic disorders, e.g., autoimmune diseases of the ear and cancer of the ear, and are suitable for use in the formulations disclosed herein.

[0074] The phrase "cytotoxic small molecule" refers to cytotoxic compounds that are of relatively low molecular weight, e.g., less than 1,000, or less than 600-700, or between 300-700 molecular
10 weight, that are effective for the treatment of otic disorders, e.g., autoimmune diseases of the ear and cancer of the ear, and are suitable for use in the formulations disclosed herein. Suitable "cytotoxic small molecules" include methotrexate, cyclophosphamide, and thalidomide, as well as metabolites, salts, polymorphs, prodrugs, analogues, and derivatives of methotrexate, cyclophosphamide, and thalidomide. In certain embodiments, preferred cytotoxic small molecules are the pharmaceutically
15 active metabolites of cytotoxic agents. For example, in the case of cyclophosphamide, preferred metabolites are pharmaceutically active metabolites of cyclophosphamide, including but not limited to 4-hydroxycyclophosphamide, aldophosphamide, phosphoramidate mustard, and combinations thereof.

[0075] "Antioxidants" are auris-pharmaceutically acceptable antioxidants, and include, for
20 example, butylated hydroxytoluene (BHT), sodium ascorbate, ascorbic acid, sodium metabisulfite and tocopherol. In certain embodiments, antioxidants enhance chemical stability where required. Antioxidants are also used to counteract the ototoxic effects of certain therapeutic agents, including agents that are used in combination with the otic agents disclosed herein.

[0076] "Auris interna" refers to the inner ear, including the cochlea and the vestibular labyrinth,
25 and the round window that connects the cochlea with the middle ear.

[0077] "Auris-interna bioavailability" or "Auris media bioavailability" refers to the percentage of the administered dose of compounds disclosed herein that becomes available in the inner or middle ear, respectively, of the animal or human being studied.

[0078] "Auris media" refers to the middle ear, including the tympanic cavity, auditory ossicles and
30 oval window, which connects the middle ear with the inner ear.

[0079] "Blood plasma concentration" refers to the concentration of compounds provided herein in the plasma component of blood of a subject.

[0080] "Auris-interna bioavailability" refers to the percentage of the administered dose of compounds disclosed herein that becomes available in the inner ear of the animal or human being
35 studied.

[0081] The term "auris-acceptable penetration enhancer" with respect to a formulation, composition or ingredient, as used herein, refers to the property of reducing barrier resistance.

[0082] "Carrier materials" are excipients that are compatible with the otic agent, the auris media, the auris interna and the release profile properties of the auris-acceptable pharmaceutical

formulations. Such carrier materials include, e.g., binders, suspending agents, disintegration agents, filling agents, surfactants, solubilizers, stabilizers, lubricants, wetting agents, diluents, and the like.

"Auris-pharmaceutically compatible carrier materials" include, but are not limited to, acacia, gelatin, colloidal silicon dioxide, calcium glycerophosphate, calcium lactate, maltodextrin, glycerine, magnesium silicate, polyvinylpyrrolidone (PVP), cholesterol, cholesterol esters, sodium caseinate, soy lecithin, taurocholic acid, phosphatidylcholine, sodium chloride, tricalcium phosphate, dipotassium phosphate, cellulose and cellulose conjugates, sugars sodium stearoyl lactylate, carrageenan, monoglyceride, diglyceride, pregelatinized starch, and the like.

[0083] The term "diluent" are chemical compounds that are used to dilute the otic agent prior to delivery and which are compatible with the auris media and/or auris interna.

[0084] "Dispersing agents," and/or "viscosity modulating agents" and/or "thickening agents" are materials that control the diffusion and homogeneity of the otic agent through liquid media.

Examples of diffusion facilitators/dispersing agents include but are not limited to hydrophilic polymers, electrolytes, Tween ® 60 or 80, PEG, polyvinylpyrrolidone (PVP; commercially known as Plasdone®), and the carbohydrate-based dispersing agents such as, for example, hydroxypropyl celluloses (e.g., HPC, HPC-SL, and HPC-L), hydroxypropyl methylcelluloses (e.g., HPMC K100, HPMC K4M, HPMC K15M, and HPMC K100M), carboxymethylcellulose, carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose phthalate, hydroxypropylmethylcellulose acetate stearate (HPMCAS), noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol (PVA), vinyl pyrrolidone/vinyl acetate copolymer (S630), 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde (also known as tyloxapol), poloxamers (e.g., Pluronic F68®, F88®, and F108®, which are block copolymers of ethylene oxide and propylene oxide); and poloxamines (e.g., Tetronic 908®, also known as Poloxamine 908®, which is a tetrafunctional block copolymer derived from sequential addition of propylene oxide and ethylene oxide to ethylenediamine (BASF Corporation, Parsippany, N.J.)), polyvinylpyrrolidone K12, polyvinylpyrrolidone K17, polyvinylpyrrolidone K25, or polyvinylpyrrolidone K30, polyvinylpyrrolidone/vinyl acetate copolymer (S-630), polyethylene glycol, e.g., the polyethylene glycol has a molecular weight of about 300 to about 6000, or about 3350 to about 4000, or about 7000 to about 5400, sodium carboxymethylcellulose, methylcellulose, polysorbate-80, sodium alginate, gums, such as, e.g., gum tragacanth and gum acacia, guar gum, xanthans, including xanthan gum, sugars, celluloses, such as, e.g., sodium carboxymethylcellulose, methylcellulose,

sodium carboxymethylcellulose, polysorbate-80, sodium alginate, polyethoxylated sorbitan monolaurate, polyethoxylated sorbitan monolaurate, povidone, carbomers, polyvinyl alcohol (PVA), alginates, chitosans and combinations thereof. Plasticizers such as cellulose or triethyl cellulose are also be used as dispersing agents. optional dispersing agents useful in liposomal dispersions and self-emulsifying dispersions of the otic agents disclosed herein are dimyristoyl phosphatidyl choline, natural phosphatidyl choline from eggs, natural phosphatidyl glycerol from eggs, cholesterol and isopropyl myristate.

[0085] "Drug absorption" or "absorption" refers to the process of movement of the otic agent from the localized site of administration, by way of example only, the round window membrane of the inner ear, and across a barrier (the round window membranes, as described below) into the auris interna or inner ear structures. The terms "co-administration" or the like, as used herein, are meant to encompass administration of the otic agent to a single patient, and are intended to include treatment regimens in which the otic agents are administered by the same or different route of administration or at the same or different time.

[0086] The terms "effective amount" or "therapeutically effective amount," as used herein, refer to a sufficient amount of the otic agent being administered that would be expected to relieve to some extent one or more of the symptoms of the disease or condition being treated. For example, the result of administration of the otic agents disclosed herein is reduction and/or alleviation of the signs, symptoms, or causes of AIED. For example, an "effective amount" for therapeutic uses is the amount of the otic agent, including a formulation as disclosed herein required to provide a decrease or amelioration in disease symptoms without undue adverse side effects. The term "therapeutically effective amount" includes, for example, a prophylactically effective amount. An "effective amount" of a otic agent composition disclosed herein is an amount effective to achieve a desired pharmacologic effect or therapeutic improvement without undue adverse side effects. It is understood that "an effective amount" or "a therapeutically effective amount" varies, in some embodiments, from subject to subject, due to variation in metabolism of the compound administered, age, weight, general condition of the subject, the condition being treated, the severity of the condition being treated, and the judgment of the prescribing physician. It is also understood that "an effective amount" in an extended-release dosing format may differ from "an effective amount" in an immediate-release dosing format based upon pharmacokinetic and pharmacodynamic considerations.

[0087] The terms "enhance" or "enhancing" refers to an increase or prolongation of either the potency or duration of a desired effect of the otic agent, or a diminution of any adverse symptomatology. For example, in reference to enhancing the effect of the otic agents disclosed herein, the term "enhancing" refers to the ability to increase or prolong, either in potency or duration, the effect of other therapeutic agents that are used in combination with the otic agents

disclosed herein. An “enhancing-effective amount,” as used herein, refers to an amount of an otic agent or other therapeutic agent that is adequate to enhance the effect of another therapeutic agent or otic agent in a desired system. When used in a patient, amounts effective for this use will depend on the severity and course of the disease, disorder or condition, previous therapy, the patient's health status and response to the drugs, and the judgment of the treating physician.

[0088] The term “penetration enhancer” refers to an agent that reduces barrier resistance (e.g., barrier resistance of the round window membrane, BLB or the like).

[0089] The term “inhibiting” includes preventing, slowing, or reversing the development of a condition, for example, AIED, or advancement of a condition in a patient necessitating treatment.

[0090] The terms “kit” and “article of manufacture” are used as synonyms.

[0091] The term “modulate” includes the interaction with a target, for example, with the TNF-alpha agents disclosed herein, the activity of TNF-alpha, or other direct or indirect targets that alter the activity of TNF-alpha, including, by way of example only, to inhibit the activity of TNF-alpha, or to limit the activity of the TNF-alpha.

[0092] “Pharmacodynamics” refers to the factors which determine the biologic response observed relative to the concentration of drug at the desired site within the auris media and/or auris interna.

[0093] “Pharmacokinetics” refers to the factors which determine the attainment and maintenance of the appropriate concentration of drug at the desired site within the auris media and/or auris interna.

[0094] In prophylactic applications, compositions containing the otic agents described herein are administered to a patient susceptible to or otherwise at risk of a particular disease, disorder or condition, for example, AIED, or patients that are suffering from diseases associated with AIED, including by way of example only, Ankylosing spondylitis, Systemic Lupus Erythematosus (SLE), Sjögren's Syndrome, Cogan's disease, ulcerative colitis, Wegener's granulomatosis, inflammatory bowel disease, rheumatoid arthritis, scleroderma and Behçet's disease. Such an amount is defined to be a “prophylactically effective amount or dose.” In this use, the precise amounts also depend on the patient's state of health, weight, and the like.

[0095] A “prodrug” refers to the otic agent that is converted into the parent drug *in vivo*. In certain embodiments, a prodrug is enzymatically metabolized by one or more steps or processes to the biologically, pharmaceutically or therapeutically active form of the compound. To produce a prodrug, a pharmaceutically active compound is modified such that the active compound will be regenerated upon *in vivo* administration. In one embodiment, the prodrug is designed to alter the metabolic stability or the transport characteristics of a drug, to mask side effects or toxicity, or to alter other characteristics or properties of a drug. Compounds provided herein, in some embodiments, are derivatized into suitable prodrugs.

[0096] “Solubilizers” refers to auris-acceptable compounds such as triacetin, triethylcitrate, ethyl oleate, ethyl caprylate, sodium lauryl sulfate, sodium doccusate, vitamin E TPGS,

dimethylacetamide, N-methylpyrrolidone, N-hydroxyethylpyrrolidone, polyvinylpyrrolidone, hydroxypropylmethyl cellulose, hydroxypropyl cyclodextrins, ethanol, n-butanol, isopropyl alcohol, cholesterol, bile salts, polyethylene glycol 200-600, glycofurol, transcutol®, propylene glycol, and dimethyl isosorbide and the like.

5 [0097] "Stabilizers" refers to compounds such as any antioxidation agents, buffers, acids, preservatives and the like that are compatible with the environment of the auris media and/or auris interna. Stabilizers include but are not limited to agents that will do any of (1) improve the compatibility of excipients with a container, or a delivery system, including a syringe or a glass bottle, (2) improve the stability of a component of the composition, or (3) improve formulation
10 stability.


[0098] "Steady state," as used herein, is when the amount of drug administered to the auris media and/or auris interna is equal to the amount of drug eliminated within one dosing interval resulting in a plateau or constant levels of drug exposure within the targeted structure.

[0099] As used herein, the term "subject" is used to mean an animal, preferably a mammal,
15 including a human or non-human. The terms patient and subject are used interchangeably.

[00100] "Surfactants" refers to compounds that are auris-acceptable, such as sodium lauryl sulfate, sodium docusate, Tween 60 or 80, triacetin, vitamin E TPGS, phospholipids, lecithins, phosphatidylcholines (c8-c18), phosphatidylethanolamines (c8-c18), phosphatidylglycerols (c8-c18), sorbitan monooleate, polyoxyethylene sorbitan monooleate, polysorbates, polaxomers, bile salts, glyceryl
20 monostearate, copolymers of ethylene oxide and propylene oxide, e.g., Pluronic® (BASF), and the like. Some other surfactants include polyoxyethylene fatty acid glycerides and vegetable oils, e.g., polyoxyethylene (60) hydrogenated castor oil; and polyoxyethylene alkylethers and alkylphenyl ethers, e.g., octoxynol 10, octoxynol 40. In some embodiments, surfactants are included to enhance physical stability or for other purposes.

25 [00101] The terms "treat," "treating" or "treatment," as used herein, include alleviating, abating or ameliorating a disease or condition, for example AIED, symptoms, preventing additional symptoms, ameliorating or preventing the underlying metabolic causes of symptoms, inhibiting the disease or condition, e.g., arresting the development of the disease or condition, relieving the disease or condition, causing regression of the disease or condition, relieving a condition caused by the disease
30 or condition, or controlling or stopping the symptoms of the disease or condition either prophylactically and/or therapeutically.

Anatomy of the Ear



[00102] The ear serves as both the sense organ that detects sound and the organ that maintains balance and body position. The ear is generally divided into three portions: the outer ear, middle ear and the inner ear (or auris interna). As shown in the illustration above, the outer ear is the external portion of the organ and is composed of the pinna (auricle), the auditory canal (external auditory meatus) and the outward facing portion of the tympanic membrane, also known as the ear drum. The pinna, which is the fleshy part of the external ear that is visible on the side of the head, collects sound waves and directs them toward the auditory canal. Thus, the function of the outer ear, in part, is to collect and direct sound waves towards the tympanic membrane and the middle ear.

[00103] The middle ear is an air-filled cavity, called the tympanic cavity, behind the tympanic membrane. The tympanic membrane, also known as the ear drum, is a thin membrane that separates the external ear from the middle ear. The middle ear lies within the temporal bone, and includes within this space the three ear bones (auditory ossicles): the malleus, the incus and the stapes. The auditory ossicles are linked together via tiny ligaments, which form a bridge across the space of the tympanic cavity. The malleus, which is attached to the tympanic membrane at one end, is linked to the incus at its anterior end, which in turn is linked to the stapes. The stapes is attached to the oval window, one of two windows located within the tympanic cavity. A fibrous tissue layer, known as the annular ligament connects the stapes to the oval window. Sound waves from the outer ear first cause the tympanic membrane to vibrate. The vibration is transmitted across to the cochlea through the auditory ossicles and oval window, which transfers the motion to the fluids in the auris interna. Thus, the auditory ossicles are arranged to provide a mechanical linkage between the tympanic membrane and the oval window of the fluid-filled auris interna, where sound is transformed and transduced to the auris interna for further processing. Stiffness, rigidity or loss of movement of the auditory ossicles, tympanic membrane or oval window leads to hearing loss, e.g. otosclerosis, or rigidity of the stapes bone.

[00104] The tympanic cavity also connects to the throat via the eustachian tube. The eustachian tube provides the ability to equalize the pressure between the outside air and the middle ear cavity. The round window, a component of the auris interna but which is also accessible within the tympanic

cavity, opens into the cochlea of the auris interna. The round window is covered by a membrane, which consists of three layers: an external or mucous layer, an intermediate or fibrous layer, and an internal membrane, which communicates directly with the cochlear fluid. The round window, therefore, has direct communication with the auris interna via the internal membrane.

5 [00105] Movements in the oval and round window are interconnected, i.e. as the stapes bone transmits movement from the tympanic membrane to the oval window to move inward against the auris interna fluid, the round window is correspondingly pushed out and away from the cochlear fluid. This movement of the round window allows movement of fluid within the cochlea, which eventually leads in turn to movement of the cochlear inner hair cells, allowing hearing signals to be
10 transduced. Stiffness and rigidity in the round window leads to hearing loss because of the lack of ability of movement in the cochlear fluid. Recent studies have focused on implanting mechanical transducers onto the round window, which bypasses the normal conductive pathway through the oval window and provides amplified input into the cochlear chamber.

[00106] Auditory signal transduction takes place in the auris interna. The fluid-filled inner ear, or
15 auris interna, consists of two major components: the cochlear and the vestibular apparatus.

[00107] The cochlea is the portion of the auris interna related to hearing. The cochlea is a tapered tube-like structure which is coiled into a shape resembling a snail. The inside of the cochlea is divided into three regions, which is further defined by the position of the vestibular membrane and the basilar membrane. The portion above the vestibular membrane is the scala vestibuli, which
20 extends from the oval window to the apex of the cochlea and contains perilymph fluid, an aqueous liquid low in potassium and high in sodium content. The basilar membrane defines the scala tympani region, which extends from the apex of the cochlea to the round window and also contains perilymph. The basilar membrane contains thousands of stiff fibers, which gradually increase in length from the round window to the apex of the cochlea. The fibers of the basement membrane
25 vibrate when activated by sound. In between the scala vestibuli and the scala tympani is the cochlear duct, which ends as a closed sac at the apex of the cochlea. The cochlear duct contains endolymph fluid, which is similar to cerebrospinal fluid and is high in potassium.

[00108] The Organ of Corti, the sensory organ for hearing, is located on the basilar membrane and extends upward into the cochlear duct. The Organ of Corti contains hair cells, which have hairlike
30 projections that extend from their free surface, and contacts a gelatinous surface called the tectorial membrane. Although hair cells have no axons, they are surrounded by sensory nerve fibers that form the cochlear branch of the vestibulocochlear nerve (cranial nerve VIII).

[00109] As discussed, the oval window, also known as the elliptical window communicates with the stapes to relay sound waves that vibrate from the tympanic membrane. Vibrations transferred to the
35 oval window increases pressure inside the fluid-filled cochlea via the perilymph and scala vestibuli/scala tympani, which in turn causes the membrane on the round window to expand in

response. The concerted inward pressing of the oval window/outward expansion of the round window allows for the movement of fluid within the cochlea without a change of intra-cochlear pressure. However, as vibrations travel through the perilymph in the scala vestibuli, they create corresponding oscillations in the vestibular membrane. These corresponding oscillations travel through the endolymph of the cochlear duct, and transfer to the basilar membrane. When the basilar membrane oscillates, or moves up and down, the Organ of Corti moves along with it. The hair cell receptors in the Organ of Corti then move against the tectorial membrane, causing a mechanical deformation in the tectorial membrane. This mechanical deformation initiates the nerve impulse which travels via the vestibulocochlear nerve to the central nervous system, mechanically transmitting the sound wave received into signals that are subsequently processed by the central nervous system.

[00110] The auris interna is located in part within the osseous or bony labyrinth, an intricate series of passages in the temporal bone of the skull. The vestibular apparatus is the organ of balance and consists of the three semi-circular canals and the vestibule. The three semi-circular canals are arranged relative to each other such that movement of the head along the three orthogonal planes in space can be detected by the movement of the fluid and subsequent signal processing by the sensory organs of the semi-circular canals, called the crista ampullaris. The crista ampullaris contains hair cells and supporting cells, and is covered by a dome-shaped gelatinous mass called the cupula. The hairs of the hair cells are embedded in the cupula. The semi-circular canals detect dynamic equilibrium, the equilibrium of rotational or angular movements.

[00111] When the head turns rapidly, the semicircular canals move with the head, but endolymph fluid located in the membranous semi-circular canals tends to remain stationary. The endolymph fluid pushes against the cupula, which tilts to one side. As the cupula tilts, it bends some of the hairs on the hair cells of the crista ampullaris, which triggers a sensory impulse. Because each semicircular canal is located in a different plane, the corresponding crista ampullaris of each semi-circular canal responds differently to the same movement of the head. This creates a mosaic of impulses that are transmitted to the central nervous system on the vestibular branch of the vestibulocochlear nerve. The central nervous system interprets this information and initiates the appropriate responses to maintain balance. Of importance in the central nervous system is the cerebellum, which mediates the sense of balance and equilibrium.

[00112] The vestibule is the central portion of the auris interna and contains mechanoreceptors bearing hair cells that ascertain static equilibrium, or the position of the head relative to gravity. Static equilibrium plays a role when the head is motionless or moving in a straight line. The membranous labyrinth in the vestibule is divided into two sac-like structures, the utricle and the saccule. Each structure in turn contains a small structure called a macula, which is responsible for maintenance of static equilibrium. The macula consists of sensory hair cells, which are embedded in

a gelatinous mass (similar to the cupula) that covers the macula. Grains of calcium carbonate, called otoliths, are embedded on the surface of the gelatinous layer.

[00113] When the head is in an upright position, the hairs are straight along the macula. When the head tilts, the gelatinous mass and otoliths tilts correspondingly, bending some of the hairs on the hair cells of the macula. This bending action initiates a signal impulse to the central nervous system, which travels via the vestibular branch of the vestibulocochlear nerve, which in turn relays motor impulses to the appropriate muscles to maintain balance.

[00114] The drug formulation will first be placed in the middle or inner ear, including the cochlea and vestibular labyrinth: one option is to use a syringe/needle or pump and inject the formulation across the tympanic membrane (the eardrum). For cochlear and vestibular labyrinth delivery, one option is to deliver the active ingredient across the round window membrane or even by microinjection directly into the auris interna also known as cochlear microperfusion.

Animal Models and Human Clinical Trials

[00115] There are, at present, no intratympanic therapeutics approved for administration to humans. In some instances, a lack of suitable animal models for inner ear diseases has hindered development of intratympanic therapeutics for human use.

[00116] In some instances, the use of animal models for inner ear diseases that are utilized for testing the efficacy of the formulations described herein is not accurately predictive of the efficacy of such formulations in humans. Rodent animal models for inner ear disease (e.g., inner ear disease models in guinea pigs) are not amenable to allometric scaling in humans because rodents are different anatomically in the organization of the middle and inner ear. The middle ear of the guinea pig (or bulla) is a cavity that contains all of the cochlea; the cochlea is anchored to the bulla via the basal turn, its apex residing in the cavity. In contrast, the human cochlea is imbedded into the temporal bone and the only access to the human cochlea is through the round window. In some instances, from a pharmacokinetics perspective, studies in guinea pigs that overfill the bulla and/or inject formulations towards the anterior quadrant of the tympani, or more generally away from the round window niche, will result in high perilymph exposure because of drug diffusion through the cochlea apex. This situation is not possible in humans because the human cochlea is imbedded into the temporal bone and as such the only access to the cochlea is on and/or through the round window or the elliptical/oval window. In addition, the ossicle chains in guinea pigs are adjacent to the round window. In some instances, the location of the ossicle chains next to the round window in guinea pig ears adversely affects the ABR threshold in experiments with guinea pigs. In contrast, the human ear is anatomically different from rodent ears; the ossicle chains and/or stapes are anatomically located away from the round window. In certain instances, an auris formulation injected intratympanically into a human ear does not make contact with the stapes and does not adversely affect the ABR threshold. Thus, in certain instances, the reliability of animal models of inner ear diseases as a

predictor of efficacy in human clinical trials is limited by the anatomical difference between the human ear and animal ears.

[00117] In some instances, a guinea pig animal model for inner ear disease utilizes an injection via a hole drilled into the bulla, i.e., the cavity surrounding the cochlear bones. In some instances, the bulla procedure leads to a local inflammatory reaction and a rapid accumulation of fluids within the bulla cavity, a condition that lasts for several days. In some instances, an accumulation of significant volumes of fluids in the bulla (about a 1/3-1/2 of the total bulla volume) seen with the bulla injection rapidly erodes any auris formulation injected, primarily by diluting the formulation and reverting a formulation (e.g., a gel formulation) to a liquid that drains away via the eustachian tube. For example, a gel formulation comprising a poloxamer will not form a gel at concentration below 12-14%, and at concentrations less than 15% concentration will gel at temperatures higher than 37 °C. In some instances, a guinea pig model is of limited utility for testing the efficacy of an auris formulation for administration to humans due the accelerated clearance of the gel from the bulla compartment of a guinea pig. For example, in some instances, a 17% Pluronic F-127 gel injection is cleared from the bulla of a guinea pig in less than 2 days.

[00118] In some instances, a guinea pig animal model for inner ear disease utilizes an injection through the tympanic membrane. In certain instances, in guinea pigs, an intratympanic injection is not associated with fluid accumulation at any of the time points evaluated (up to 10 days). In some instances, injection of an auris formulation described herein (e.g. a gel formulation) via the tympanic route allows for detectable amounts of the formulation (e.g., a gel) in the inner ear of a guinea pig up to at least 5 days.

[00119] In some instances, animal models (e.g., guinea pig models for inner ear diseases) utilizing intratympanic injections are limited by the volume that can be injected through the tympanic route. In the guinea pig, the round window niche and membrane are located just opposite the tympanic membrane in the posterior superior quadrant. In certain instances, about 50 mL can be injected within this quadrant in a 250-350g guinea pig. In some instances, a larger volume (up to 70 mL) can be injected in the posterior inferior quadrant; however most of the gel migrates towards the round window. In some instances larger volumes (100-120mcl) are injected in the anterior quadrant, but this action fills the bulla cavity and promotes drug transfer across the apical part of the cochlea (due to the bone structure thinness of the cochlea in rodents). In certain animal models, injection of larger volumes in any of these quadrants leads to tympanic perforation and presence of the gel in the external ear canal. In some instances, the volume injected has an impact on the hearing threshold (measured by ABR). In the guinea pig ear for example, intratympanic injections volumes up to 50 mL do not produce any shift in hearing threshold; but volumes of 90 and 120 mL produce an ABR threshold shift within 1 day. In some instances, the anatomical difference between human and animal ears and the variability in experimental outcomes lends a low predictive value to animal testing

data for use in subsequent human clinical trials. Further, the invasive procedures used in animal models of inner ear disease are not applicable in a clinical setting.

Visualization of otic formulations

[00120] Provided herein are otic formulations that comprise a dye (e.g., a Trypan blue dye, Evans blue dye) or other tracer compound. In some instances, addition of an auris-compatible dye to an otic formulation described herein aids visualization of any administered formulation in a ear (e.g., a rodent ear and/or a human ear). In certain embodiments, an otic composition comprising a dye or other tracer compound eliminates the need for invasive procedures that are currently used in animal models to monitor the concentrations of drugs in the endolymph and/or perilymph.

[00121] In some instances, intratympanic injections require the need of a specialist and the formulation needs to be delivered to a specific site of the ear to maximize efficiency of the medication delivered. In certain instances, a visualization technique for any formulation described herein allows for visualization of a dosing site (e.g., the round window) so that the medication is applied in the proper place. In some instances, a formulation comprising a dye allows visualization of the formulation during administration of the formulation to an ear (e.g., a human ear), ensures that the medication will be delivered at the intended site, and avoids any complications due to incorrect placement of a formulation. The inclusion of a dye to help enhance the visualization of the gel when applied, and the ability to visually inspect the location of the gel after administration without further intervention, represents an advance over currently available methods for testing intratympanic therapeutics in animal models and/or human trials. In some embodiments, dyes that are compatible with the otic compositions described herein include Evans blue (e.g., 0.5% of the total weight of an otic formulation), Methylene blue (e.g., 1% of the total weight of an otic formulation), Isosulfan blue (e.g., 1% of the total weight of an otic formulation), Trypan blue (e.g., 0.15% of the total weight of an otic formulation), and/or indocyanine green (e.g., 25mg/vial). Other common dyes, e.g., FD&C red 40, FD&C red 3, FD&C yellow 5, FD&C yellow 6, FD&C blue 1, FD&C blue2, FD&C green 3, fluorescence dyes (e.g., Fluorescein isothiocyanate, rhodamine, Alexa Fluors, DyLight Fluors) and/or dyes that are visualizable in conjunction with non-invasive imaging techniques such as MRI, CAT scans, PET scans or the like (e.g., Gadolinium-based MRI dyes, iodine-base dyes, barium-based dyes or the like) are also contemplated for use with any otic formulation described herein. Other dyes that are compatible with any formulation described herein are listed in the Sigma-Aldrich catalog under dyes (which is included herein by reference for such disclosure). In some embodiments, concentration of a dye in any otic formulation described herein is less than 2%, less than 1.5%, less than 1%, less than 0.5%, less than 0.25%, less than 0.1%, or less than 100 ppm of the total weight and/or volume of any formulation described herein.

[00122] In certain embodiments of such auris-compatible formulations that comprise a dye, the ability to visualize a controlled release otic formulation comprising a dye in an ear meets a long

inhibitors (by way of example only CEP/KT-7515, AS601245, SPC9766 and SP600125),
antioxidants, NSAIDs, neuroprotectants, glutamate modulators, interleukin 1 modulators,
interleukin-1 antagonists, including tumor necrosis factor- α converting enzyme (TACE) and
caspases, retinaldehyde modulator, notch modulator, gamma secretase modulator, thalidomide,
5 latanoprost (Xalatan[®]) for reducing internal pressure and combinations thereof.

Immunomodulating agents

Anti-TNF Agents

[00266] Contemplated for use with the formulations disclosed herein are agents which reduce or
ameliorate symptoms or effects as a result of an autoimmune disease and/or inflammatory disorder,
10 including AIED or OM. Accordingly, some embodiments incorporate the use of agents which block
the effects of TNF- α , including anti-TNF agents. By way of example only, anti-TNF agents include
protein-based therapeutics, such as etanercept (ENBREL[®]), infliximab (REMICADE[®]), adalimumab
(HUMIRA[®]) and golimumab (CNTO 148), and small molecule therapeutics, such as TACE
inhibitors, IKK inhibitors or calcineurin inhibitors or combinations thereof.

15 [00267] Infliximab and adalimumab are anti-TNF monoclonal antibodies, and etanercept is a fusion
protein designed to bind specifically to the TNF protein. All are currently approved for use in the
treatment of rheumatoid arthritis. Golimumab, which is currently in Phase 3 clinical trials for
rheumatoid arthritis, psoriatic arthritis and ankylosing spondylitis, is a fully-humanized anti-TNF-
alpha IgG1 monoclonal antibody that targets and neutralizes both the soluble and the membrane-
20 bound form of TNF- α . Other antagonists to TNF, by way of example only, include TNF receptors
(pegylated soluble TNF receptor type 1; Amgen); TNF binding factors (Onercept; Serono); TNF
antibodies (US Patent App. No. 2005/0123541; US Patent App. No. 2004/0185047); single domain
antibodies against the p55 TNF receptor (US Patent App. No. 2008/00088713); soluble TNF
receptors (US Patent App. No. 2007/0249538); fusion polypeptides binding to TNF (US Patent App.
25 No. 2007/0128177); and flavone derivatives (US Patent App. No. 2006/0105967), all of which are
incorporated by reference for such disclosure. The use of onercept, a soluble TNF p55 receptor, was
discontinued in 2005. Three phase-III clinical trials reported patients diagnosed with fatal sepsis. A
risk to benefit analysis was subsequently performed, resulting in the discontinuation of the clinical
trials. As discussed above, the embodiments herein specifically contemplate the use of anti-TNF
30 agents which have been previously shown to have limited or no systemic release, systemic toxicity,
poor PK characteristics of combinations thereof.

[00268] Although etanercept, infliximab and adalimumab are currently approved systemic therapies
for use in rheumatoid arthritis, these anti-TNF agents are not without serious adverse side effects. It
is contemplated that the localized application of the anti-TNF agents to the target otic structures for
35 treatment of autoimmune and/or inflammatory disorders will result in the reduction or elimination of
these adverse side effects experienced with systemic treatment. Moreover, localized treatment with

the anti-TNF agents contemplated herein will also reduce the amount of agent needed for effective treatment of the targeted disorder due, for example, to the existence of the biological blood barrier in the auris interna or to the lack of sufficient systemic access to the auris media.

[00269] Etanercept is a dimeric fusion protein consisting of the extracellular ligand-binding portion of the human 75 kilodalton (p75) tumor necrosis factor receptor (TNFR) linked to the Fc portion of human IgG1. The Fc component of etanercept contains the C_H2 domain, the C_H3 domain and hinge region, but not the C_H1 domain of IgG1. Etanercept is a recombinant protein consisting of 934 amino acids, with an apparent molecular weight of approximately 150 kilodaltons. Etanercept binds specifically to tumor necrosis factor (TNF), and acts by inhibiting the interaction of TNF with cell surface TNF receptors. Serious side effects with etanercept have been reported with systemic administration, including serious infections and sepsis that resulted in fatalities. Other side effects observed upon intravenous administration of etanercept include contraction of tuberculosis; onset or exacerbation of central nervous system disorders, including mental status changes, transverse myelitis, optic neuritis, multiple sclerosis and seizures resulting in permanent disability; adverse hematologic events, including pancytopenia, aplastic anemia with fatal outcomes, blood dyscrasias, persistent fever, bruising, bleeding and pallor, neutropenia and cellulitis. Treatment with etanercept may also result in the formation of autoantibodies, which may develop into a lupus-like syndrome, as well as development of malignant disorders. Moreover, over one-third of patients systemically treated with etanercept experience injection site reactions including mild to moderate erythema and/or itching, pain and/or swelling. Injection site bleeding and bruising has also been observed. Other side effects from the systemic administration of etanercept include headache, nausea, rhinitis, dizziness, pharyngitis, cough, asthenia, abdominal pain, rash, peripheral edema, respiratory disorder, dyspepsia, sinusitis, vomiting, mouth ulcer, alopecia and pneumonitis. Infrequent side effects include heart failure, myocardial infarction, myocardial ischemia, hypertension, hypotension, deep vein thrombosis, thrombophlebitis, cholecystitis, pancreatitis, gastrointestinal hemorrhage, bursitis, polymyositis, cerebral ischemia, depression, dyspnea, pulmonary embolism, and membranous glomerulonephropathy in rheumatoid arthritis patients. Varicella infections, gastroenteritis, depression/personality disorder, cutaneous ulcer, esophagitis/gastritis, group A streptococcal septic shock, type I diabetes mellitus, and soft tissue and post-operative wound infection was also seen in juvenile rheumatoid arthritis patients.

[00270] Infliximab is a chimeric human-mouse IgG1 κ monoclonal antibody with an approximate molecular weight of 149 kilodaltons. Infliximab binds specifically to TNF α with an association constant of 10^{10} M^{-1} . Infliximab is produced by a recombinant cell line cultured by continuous perfusion. Infliximab acts to neutralize the binding activity of TNF α by inhibiting binding of TNF to its cell surface receptors. Serious side effects as a result of systemic intravenous infusions or injections have been reported with the use of infliximab, including fatal sepsis and serious

infections. Cases of histoplasmosis, listeriosis, pneumocystosis and tuberculosis have also been observed. Hypersensitivity, including urticaria, dyspnea and hypotension have occurred upon treatment with infliximab. Infusion reactions include cardiopulmonary reactions (primarily chest pain, hypotension, hypertension or dyspnea), pruritus, and combined reactions. Other


5 hypersensitivity symptoms include fever, rash, headache, sore throat, myalgias, polyarthralgias, hand and facial edema and/or dysphagia, anaphylaxis, convulsions, erythematous rash, laryngeal/pharyngeal edema and severe bronchospasm. Neurologic adverse events include optic neuritis, seizure and new onset or exacerbation and/or radiographic evidence of central nervous system demyelinating disorders, including multiple sclerosis. The formation of autoantibodies have also been observed, including symptoms suggestive of a lupus-like syndrome following treatment. Other serious adverse events include worsening rheumatoid arthritis, rheumatoid nodules, abdominal hernia, asthenia, chest pain, diaphragmatic hernia, pancytopenia, splenic infarction, splenomegaly, syncope, cerebral hypoxia, convulsions, dizziness, encephalopathy, hemiparesis, spinal stenosis, upper motor neuron lesion, ceruminosis, endophthalmitis, and other infrequent-occurring side effects.

[00271] Adalimumab is a recombinant human IgG1 monoclonal antibody specific for human TNF. Adalimumab was created using phage display technology resulting in an antibody with human derived heavy and light chain variable regions and human IgG1:κ constant regions, and consists of 1330 amino acids with a molecular weight of approximately 148 kilodaltons. Adalimumab binds specifically to TNF-α and blocks its interaction with both the p55 and p75 TNF cell surface receptors. Adalimumab also lyses TNF expressing cells in vitro in the presence of complement. Adalimumab does not bind or inactivate lymphotoxin (TNF-β). Serious side effects from systemic administration have been reported with the intravenous administration or injection of adalimumab, including fatal sepsis and serious infections, including upper respiratory infections, bronchitis, urinary tract infections, pneumonia, septic arthritis, prosthetic and post-surgical infections, erysipelas cellulitis, diverticulitis, pyelonephritis, tuberculosis, and invasive opportunistic infections caused by histoplasma, aspergillus and nocardia. Other serious adverse reactions were neurologic events, including confusion, multiple sclerosis, paresthesia, subdural hematoma, and tremor, and the development of malignancies, including lymphoma development. The formation of autoantibodies has also been observed, including symptoms suggestive of a lupus-like syndrome following treatment. The most common adverse reaction was injection site reactions, with 20% of patients developing erythema and/or itching, hemorrhage, pain and/or swelling. Other adverse events as a result of systemic administration of adalimumab include clinical flare reaction, rash and pneumonia. Other adverse events included sinusitis, flu syndrome, nausea, abdominal pain, hypercholesterolemia, hyperlipidemia, hematuria, increased alkaline phosphatase levels, back pain, hypertension, as well as more infrequent serious adverse events, including pain, pelvic pain, thorax

pain, arrhythmia, atrial fibrillation, cardiovascular disorder, congestive heart failure, coronary artery disorder, heart arrest, hypertensive encephalopathy, myocardial infarct, palpitation, pericardial effusion, pericarditis, syncope, tachycardia, vascular disorders, and other disorders.

Calcineurin Inhibitors

- 5 [00272] Calcineurin inhibitors are a group of structurally diverse small molecule immunomodulators which function through the inhibition of calcineurin function. Calcineurin is a calcium-activated protein phosphatase which catalyses the dephosphorylation of cytoplasmic NFAT. Upon dephosphorylation, NFAT migrates to the nucleus and forms a regulatory complex involved in the transcription of cytokines, such as TNF- α , IL-2, IL-3 and IL-4. Inhibition of calcineurin function
10 blocks the dephosphorylation event and subsequent cytokine transcription. An unusual aspect of calcineurin inhibition is that cyclosporine, tacrolimus and pimecrolimus are required to form a complex with an immunophilin for the inhibitory properties to be realized (Schreiber et al, Immunol. Today (1992), 13:136-42; Liu et al, Cell (1991), 66:807-15). For cyclosporine the immunophilin is cyclophilin; tacrolimus and pimecrolimus bind to the FK506-binding protein (FKBP).



- 15 [00273] Cyclosporine is an 11-residue cyclic peptide produced as a metabolite of the fungus *Beauveria nivea* and has the chemical name cyclo[[*(E)*-(2*S*,3*R*,4*R*)-3-hydroxy-4-methyl-2-(methylamino)-6-octenoyl]-L-2-aminobutyryl-N-methylglycyl-N-methyl-L-leucyl-L-valyl-N-methyl-L-leucyl-L-alanyl-D-alanyl-N-methyl-L-leucyl-N-methyl-L-leucyl-N-methyl-L-valyl]. It is
20 provided in several formulations for both systemic or local administration. Sandimmune® provides cyclosporine in three different formulations: soft gelatin capsules, an oral solution or a formulation for injection. Sandimmune® is indicated for prevention of organ rejection in kidney, liver or heart transplants. Neoral® and Gengraf® provide cyclosporine in two formulations: soft gelatin capsules

and an oral solution. They are indicated for prevention of organ rejection in kidney, liver or heart transplants, for treatment of patients with severe active, rheumatoid arthritis, or for treatment of severe psoriasis. Compared to Sandimmune®, Neoral® and Gengraf® provide increased bioavailability of cyclosporine. Restasis® provides cyclosporine in an ophthalmic emulsion formulation. It is indicated to increase tear production in patients with reduced tear production due to ocular inflammation associated with keratoconjunctivitis sicca.

[00274] Tacrolimus, also known as FK-506 or fujimycin, is a 23-membered macrolide natural product produced by *Streptomyces tsukubaensis* and has the chemical name {3S-[3R*[E(1S*,3S*,4S*)], 4S*,5R*,8S*,9E,12R*,14R*,15S*,16R*,18S*,19S*,26aR*]}-5,6,8,11,12,13,14,15,16,17,18,19,24,25,26,26a-hexadecahydro-5,19-dihydroxy-3-[2-(4-hydroxy-3-methoxycyclohexyl)-1-methylethenyl]-14,16-dimethoxy-4,10,12,18-tetramethyl-8-(2-propenyl)-15,19-epoxy-3H-pyrido[2,1-39c][1,4]oxaazacyclotricosine-1,7,20,21(4H,23H)-tetrone monohydrate. It is provided in formulations suitable for systemic or topical administration. For systemic administration, the Prograf® formulation provides an oral capsule or a sterile solution for injection. Prograf® is indicated for prevention of organ rejection in liver, kidney or heart transplants. For topical administration, the Protopic® formulation is indicated for the treatment of moderate-to-severe atopic dermatitis.

[00275] Pimecrolimus is a semi-synthetic analog of tacrolimus and has the chemical name (1R,9S,12S,13R,14S,17R,18E,21S,23S,24R,25S,27R)-12-[(1E)-2-[(1R,3R,4S)-4-chloro-3-methoxycyclohexyl]-1-methylvinyl]-17-ethyl-1,14-dihydroxy-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-aza-tricyclo[22.3.1.04,9]octacos-18-ene-2,3,10,16-tetraone. It is provided in a formulation suitable for topical application and is indicated for the treatment of mild-to-moderate atopic dermatitis.

[00276] Studies have shown that tacrolimus and pimecrolimus do not suppress Langerhans' cells or dermal connective tissue and therefore do not cause atrophy of the skin, unlike corticosteroids (Stuetz et al, Int. Arch. Allergy Immunol. (2006), 141:199-212; Queille-Roussel et al, Br. J. Dermatol. (2001), 144:507-13). Because of the importance of calcineurin, systemic administration of calcineurin inhibitors leads to significant side effects. Systemic side effects are related to dose, exposure levels and duration of therapy. Prolonged elevated blood levels result in hypertension, nephrotoxicity, psychiatric disorders, hyperlipidemia, and profound immunosuppression. Topical application of tacrolimus or pimecrolimus has shown to afford very little, if any, systemic exposure, with tacrolimus having demonstrated less than 0.5 % bioavailability after topical application.

[00277] In one embodiment, the auris-acceptable controlled release immunomodulating formulation comprises a calcineurin inhibitor. In another embodiment, the auris-acceptable controlled release immunomodulating formulation comprises cyclosporine. In another embodiment, the auris-acceptable controlled release immunomodulating formulation comprises tacrolimus. In another

embodiment, the auris-acceptable controlled release immunomodulating formulation comprises pimecrolimus. In another embodiment, the auris-acceptable controlled release immunomodulating formulation comprises a calcineurin inhibitor which induces toxicity upon systemic administration.

[00278] Other pharmaceutical agents that are optionally used in combination with

immunomodulating- α agents for the treatment of autoimmune and/or inflammatory disorders include other agents that have been used to treat autoimmune and inflammatory disorders, including corticosteroids, local anesthetic agents, chemotherapeutic agents, including cytoxan, azathiaprine or methotrexate; treatment with collagen, gamma globulin, interferons, copaxone, or combinations thereof. Accordingly, also contemplated within the scope of the embodiments herein is the use of other pharmaceutical agents in combination with the immunomodulating compositions and formulations disclosed in the treatment of autoimmune otic disorders. In addition, other pharmaceutical agents are optionally used to treat attendant symptoms of AIED or other autoimmune disorder, including vomiting, dizziness and general malaise.


IKK Inhibitors

[00279] The transcription of TNF- α is dependent on the transcription factor NF- κ B. In unstimulated cells, NF- κ B is in the cytoplasm as part of a protein complex with the protein inhibitor of NF- κ B, also known as I κ B. Activation of NF- κ B depends on phosphorylation-induced ubiquitination of the I κ B. Once poly-ubiquitinated, the I κ B undergoes a rapid degradation through the 26S proteasome and the free NF- κ B migrates to the nucleus to activate pro-inflammatory gene transcription. The phosphorylation event which releases NF- κ B is mediated by the I κ B kinase (IKK) complex, composed of IKK kinases. Two IKK enzymes, generally referred to as IKK- α and IKK- β (Woroniec et al. Science (1997), 278:866; Zandi et al. Cell (1997), 91:243) or IKK-1 and IKK-2 (Mercurio et al. Science (1997), 278:860) have been discovered. Both forms of IKK can exist as homodimers and as IKK- α /IKK- β heterodimers. Another component of the I κ B kinase complex is a regulatory protein, known as IKK- γ or NEMO (NF- κ B-Essential Modulator) (Rothwarf et al. Nature (1998), 395:297). NEMO does not contain a catalytic domain, and thus it appears to have no direct kinase activity and it probably serves a regulatory function. Existing data suggests that the predominant form of IKK in cells is an IKK- α /IKK- β heterodimer associated with either a dimer or a trimer of NEMO (Rothwarf et al. Nature (1998) 395:297). Biochemical and molecular biology experiments have identified IKK- α and IKK- β as the most likely mediators of TNF- α -induced I κ B phosphorylation and degradation, which results in NF- κ B activation and upregulation of families of genes involved in inflammatory processes (Woroniec et al. Science (1997); Karin, Oncogene (1999) 18:6867; Karin, J. Biol. Chem. (1999) 274:27339).


[00280] Many IKK- β inhibitors have been identified. SPC-839 has been extensively studied. It

inhibits IKK- β with an IC_{50} of 62 nM and reduces paw edema in a rat arthritis model at 30 mg/kg.


Carboline PS-1145 inhibits the IKK complex with an IC_{50} of 150 nM and reduces the production of TNF- α in LPS-challenged mice. BMS-345541, an allosteric inhibitor, inhibits IKK- β with an IC_{50} of 0.3 μ M. In the mouse collagen-induced arthritis model it significantly reduced the severity of disease at a 30mg/kg dose. A scientific review of IKK inhibitors has been published (Karin *et al.*, Nature Reviews Drug Discovery (2004), 3, 17-26), incorporated herein by reference for such disclosure.



SPC-839



PS-1145



BMS-345541

[00281] In one embodiment, the auris-acceptable controlled release immunomodulating formulation comprises an IKK inhibitor. In a further embodiment, the auris-acceptable controlled release immunomodulating formulation comprises a IKK- β inhibitor. In another embodiment, the auris-acceptable controlled release immunomodulating formulation comprises a IKK inhibitor which induces toxicity upon systemic administration. In an additional embodiment, the auris-acceptable controlled release immunomodulating formulation comprises a IKK inhibitor which is not orally absorbed. In an additional embodiment, the auris-acceptable controlled release immunomodulating formulation comprises an IKK inhibitor selected from SPC-839, PS-1145, BMS-345541, or SC-514. In an additional embodiment, the auris-acceptable controlled release immunomodulating formulation comprises an IKK inhibitor selected from compounds disclosed in the following group of patent publications: WO199901441, WO2001068648, WO2002060386, WO2002030353, WO2003029242, WO2003010163, WO2001058890, WO2002044153, WO2002024679, WO2002046171, WO2003076447, WO2001030774, WO2001000610, WO2003024936, WO2003024935, WO2002041843, WO200230423, WO2002094265, WO2002094322, WO2005113544 and WO2006076318, all of which are incorporated by reference herein for such disclosure.

Interleukin Inhibitors

[00282] Interleukins are a class of cytokines. In certain instances, they are signaling molecules secreted by leukocytes having encountered a pathogen. In certain instances, the secretion of interleukins activates and recruits additional leukocytes to the site of infection. In certain instances, the recruitment of additional leukocytes to the site of infection results in inflammation (due to the increase in leukocyte containing lymph). IL-1 α , IL-1 β , IL-2, and IL-8 are found in middle ear effusions. In certain instances, IL-1 α and IL-1 β are also found in the epithelium of cholesteatomas. [00283] IL-1 is a class of interleukins comprised of IL-1 α , and IL-1 β . IL-1 is made by macrophages, B cells, monocytes, and dendritic cells (DC). It binds to receptors IL1R1/CD121a and

IL1R2/CD121b. The binding of IL-1 to its receptors results in an increase in cell-surface adhesion factors. This enables the migration of leukocytes to the site of infection.

[00284] IL-2 is made by TH-1 cells and binds to the receptors CD25/IL2Ra, CD122IL2Rb, and CD132/IL2Rg. IL-2 secretion is stimulated by the binding of an antigen to a TH-1 cell. The binding of IL-2 to a receptor stimulates the growth, and differentiation of memory T cells.

[00285] IL-8 is made by macrophages, lymphocytes, epithelial cells, and endothelial cells. It binds to CXCR1/IL8Ra and CXCR2/IL8Ra/CD128. Secretion of IL-8 initiates neutrophil chemotaxis to the site of infection.

[00286] In some embodiments, a subject in need thereof is administered an inhibitor of a pro-inflammatory interleukin. In some embodiments, the pro-inflammatory interleukin is IL-1 α , IL-1 β , IL-2, or IL-8. In some embodiments, the inhibitor of a pro-inflammatory interleukin is a WS-4 (an antibody against IL-8); [Ser IL-8]₇₂; or [Ala IL-8]₇₇ (See U.S. Patent No. 5,451,399 which is hereby incorporated by reference for disclosures relating to these peptides); IL-1RA; SB 265610 (N-(2-Bromophenyl)-N'-(7-cyano-1H-benzotriazol-4-yl)urea); SB 225002 (N-(2-Bromophenyl)-N'-(2-hydroxy-4-nitrophenyl)urea); SB203580 (4-(4-Fluorophenyl)-2-(4-methylsulfinyl phenyl)-5-(4-pyridyl) 1H-imidazole); SB272844 (GlaxoSmithKline); SB517785 (GlaxoSmithKline); SB656933 (GlaxoSmithKline); Sch527123 (2-hydroxy-N,N-dimethyl-3-{2-[[[(R)-1-(5-methyl-furan-2-yl)-propyl]amino]-3,4-dioxo-cyclobut-1-enylamino]-benzamide); PD98059(2-(2-amino-3-methoxyphenyl)-4H-1-Benzopyran-4-one); reparixin; N-[4-chloro-2-hydroxy-3-(piperazine-1-sulfonyl)phenyl]-N'-(2-chloro-3-fluorophenyl)urea p-toluenesulfonate (See WO/2007/150016 which is hereby incorporated by reference for disclosures relating to this compound); sivelestat; bG31P (CXCL8((3-74))K11R/G31P); basiliximab; cyclosporin A; SDZ RAD (40-O-(2-hydroxyethyl)-rapamycin); FR235222 (Astellas Pharma); daclizumab; anakinra; AF12198 (Ac-Phe-Glu-Trp-Thr-Pro-Gly-Trp-Tyr-Gln-L-azetidine-2-carbonyl-Tyr-Ala-Leu-Pro-Leu-NH₂); or combinations thereof.



Platelet Activating Factor Antagonists

[00287] Platelet activating factor antagonists are contemplated for use in combination with the immunomodulating formulations disclosed herein. Platelet activating factor antagonists include, by way of example only, kadsurenone, phomactin G, ginsenosides, apafant (4-(2-chlorophenyl)-9-methyl-2[3(4-morpholinyl)-3-propanol-1-yl][6H-thieno[3.2-f][1.2.4]triazolo[4,3-1]]1.4]diazepine), A-85783, BN-52063, BN-52021, BN-50730 (tetrahedra-4,7,8,10 methyl-1 (chloro-1 phenyl)-6 (methoxy-4 phenyl-carbamoyl)-9 pyrido [4',3'-4,5] thieno [3,2-f] triazolo-1,2,4 [4,3-a] diazepine-1,4), BN 50739, SM-12502, RP-55778, Ro 24-4736, SR27417A, CV-6209, WEB 2086, WEB 2170, 14-deoxyandrographolide, CL 184005, CV-3988, TCV-309, PMS-601, TCV-309 and combinations thereof.

TNF- α Converting Enzyme (TACE) Inhibitors

[00288] TNF- α is initially expressed on the cell surface as a 26 kDa, 233-amino acid, membrane-bound precursor protein. Proteolytic cleavage of the membrane-bound TNF- α by the matrix metalloproteinase TNF- α converting enzyme occurs between Ala-76 and Val-77 and results in a 17 kDa mature TNF- α which exists as a soluble trimer. Inhibition of the proteolytic cleavage could provide an alternative to the use of protein-based therapeutics in anti-inflammatory therapy. One potential complication, however, is that TACE is thought to be involved in the processing of other proteins in addition to TNF- α . For example, in a phase II clinical trial, indications of toxic effects in the liver occurred as a result of TACE inhibition. (Car et al, Society of Toxicology, 46th Annual Meeting, Charlotte, North Carolina, March 25-29, 2007). The hypothesis for this mechanism-based toxicity is that TACE also acts on other membrane bound proteins, such as TNFRI and TNFRII.

[00289] While toxicities following oral administration are problematic for a drug administered systemically, local delivery to the site of action overcomes this problem. Inhibitor GW3333 has a TACE IC₅₀ of 40 nM and an IC₅₀ of 0.97 μ M for inhibiting TNF- α production in the LPS-induced human PBMC cells (Conway et al, J. Pharmacol. Exp. Ther. (2001), 298:900). Nitroarginine analog A has an IC₅₀ TACE IC₅₀ of 4 nM and an IC₅₀ of 0.034 μ M for inhibiting TNF- α production in the LPS-induced MonoMac-6 cells (Musso et al, Bioorg. Med. Chem. Lett. (2001), 11:2147), but lacks oral activity. A scientific review of TNF- α converting enzyme inhibitors has been published (Skotnicki *et al.*, Annual Reports in Medicinal Chemistry (2003), 38, 153-162), incorporated by reference herein for such disclosure.



[00290] Accordingly, in one embodiment, the auris-acceptable controlled release anti-TNF formulation comprises a TACE inhibitor. In another embodiment, the auris-acceptable controlled release anti-TNF formulation comprises a TACE inhibitor which induces toxicity upon systemic administration. In additional embodiments, the auris-acceptable controlled release anti-TNF formulation comprise a TACE inhibitor which is not orally absorbed. In another embodiment, the auris-acceptable controlled release anti-TNF formulation comprises a TACE inhibitor selected from Nitroarginine analog A, GW3333, TMI-1, BMS-561392, DPC-3333, TMI-2, BMS-566394, TMI-005, apratastat, GW4459, W-3646, IK-682, GI-5402, GI-245402, BB-2983, DPC-A38088, DPH-067517, R-618, or CH-138.

Toll-like Receptor Inhibitors

[00291] Toll-like receptors (TLR) are a family of at least 12 pattern recognition cell-surface and intracellular receptors. The family is defined by the presence of two domains: a ligand-binding domain with multiple leucine-rich repeats, and a short Toll/Il-1 receptor domain; the latter controlling the initiation of downstream-signaling cascades. In certain instances, the receptors are activated by the binding of structurally conserved molecules (i.e. the "patterns") found on pathogens. Each receptor recognizes and binds to specific conserved molecules found on pathogens (e.g. TLR2 – lipopeptides; TLR3- viral dsRNA; TLR4 – LPS; TLR5 – flagellin; TLR9 – CpG DNA). In certain instances, the binding of a TLR to a pathogen, initiates the TLR signaling cascade which ultimately leads to the activation of various cytokines, chemokines, and antigen-specific and non-specific immune responses. In certain instances, the expression of TLR2 and/or TLR4 is up-regulated upon exposure to nontypeable *Hemophilus influenzae* (NTHi). Infection by NTHi is a common cause of otitis media.

[00292] Toll-like receptors belong to a class of single membrane-spanning non-catalytic receptors that recognize structurally conserved molecules derived from breached microbes are believed to play a key role in the innate immune system. Toll-like receptors thus recognize molecules that are broadly shared by pathogens, but are distinguishable from the host molecules. These receptors form a superfamily with Interleukin-1 receptors, and have in common a Toll-like receptor domain. Toll-like receptor agonists, such as CQ-07001, can stimulate Toll-like receptor 3 function, triggering anti-inflammatory and tissue regeneration activity. Toll-like receptor modulators, thus, have implication for use in both auris interna disorders, including AIED, and auris media diseases, including otitis media. In some embodiments, toll-like receptor modulators include toll-like receptor antagonist, partial agonist, inverse agonist, neutral or competitive antagonist, allosteric antagonist, and/or orthosteric antagonist. Other toll-like receptor modulators include but are not limited to polyinosinic-polycytidylic acid [poly(I:C)], polyAU, other nucleic acid molecules, including dsRNA agonists (such as AMPLIGEN®, Hemispherx, Inc., Rockville MD; and POLYADENUR®, Ipsen), and are also contemplated within the scope of the embodiments disclosed herein.

[00293] In some embodiments, the TLR inhibitor is an ST2 antibody; sST2-Fc (functional murine soluble ST2-human IgG1 Fc fusion protein; see Biochemical and Biophysical Research Communications, 29 December 2006, vol. 351, no. 4, 940-946 which is herein incorporated by reference for disclosures related to sST2-Fc); CRX-526 (Corixa); lipid IV_A; RSLA (*Rhodobacter sphaeroides* lipid A); E5531 ((6-O-{2-deoxy-6-O-methyl-4-O-phosphono-3-O-[(R)-3-Z-dodec-5-endoxyloxydecyl]-2-[3-oxo-tetradecanoylamino]-β-O-phosphono-α-D-glucopyranose tetrasodium salt); E5564 (α-D-Glucopyranose,3-O-decyl-2-deoxy-6-O-[2-deoxy-3-O-[(3R)-3-methoxydecyl]-6-O-methyl-2-[(11Z)-1-oxo-11-octadecenyl]amino]-4-O-phosphono-β-D-glucopyranosyl]-2-[(1,3-dioxotetradecyl)amino]-1-(dihydrogen phosphate), tetrasodium salt); compound 4a

(hydrocinnamoyl-L-valyl pyrrolidine; see PNAS, June 24, 2003, vol. 100, no. 13, 7971-7976 which is herein incorporated by reference for disclosures related to compound 4a); CPG 52364 (Coley Pharmaceutical Group); LY294002 (2-(4-Morpholinyl)-8-phenyl-4H-1-benzopyran-4-one); PD98059 (2-(2-amino-3-methoxyphenyl)-4H-1-Benzopyran-4-one); chloroquine; and an immune regulatory oligonucleotide (for disclosures relating to IROs see U.S. Patent Application Publication No. 2008/0089883).

Auto-Immune Agents

[00294] Also contemplated for use with the formulations disclosed herein are agents which reduce or ameliorate symptoms or effects as a result of autoimmune disease, including autoimmune inner ear disease (AIED). Accordingly, some embodiments may incorporate the use of agents which block the effects of TNF- α , including but not limited to anti-TNF agents. By way of example only, some anti-TNF agents include etanercept (ENBREL®), infliximab (REMICADE®) and adalimumab (HUMIRA®), or combinations thereof. Other pharmaceutical agents to treat autoimmune disorders include chemotherapeutic agents, including cytoxan, azathiaprine or methotrexate; treatment with collagen, gamma globulin, interferons, copaxone, or combinations thereof.

IL-1 Modulators

[00295] Interleukin-1 (IL-1) is a pleiotropic cytokine that plays a role in the modulation of local as well as systemic inflammation, immune regulation and hemopoiesis. IL-1 β , a member of the IL-1 family, has been implicated in angiogenesis processes, including tumor angiogenesis. In addition, IL-1 has been shown to stimulate the synthesis of inflammatory eicosanoids in macrophages, fibroblasts, synovial cells and chondrocytes, and is believed to contribute to leukocyte activation and tissue destruction in arthritic models. Interfering with IL-1 activity, therefore, is an approach for developing a disease modifying therapy for chronic inflammatory diseases, such as AIED and otitis media. In some embodiments, IL-1 modulators include an IL-1 antagonist, partial agonist, inverse agonist, neutral or competitive antagonist, allosteric antagonist, and/or orthosteric antagonist. In some embodiments, IL-1 modulators include but are not limited to antibodies that specifically recognize IL-1 subunits or its receptors, proteins, peptides, nucleic acids, and small molecule therapeutics. In some embodiments, IL-1 modulators are IL-1 antagonists, including, for example, AF12198, IL-1 natural antagonists, inactive receptor fragments that bind to IL-1 molecule, and antisense molecules or factors that block expression of IL-1 cytokine proteins. In some embodiments, IL-1 antagonists are IL-1 antibodies including, by way of example, anakinra (Kinaret®) and ACZ885 (Canakinumab®). In some embodiments, modulators of IL-1 are antibodies that modulate cytokines and/or growth factors that affect the release and/or expression of IL-1, including, by way of example, ranibizumab, tefibazumab, and bevacizumab. In some embodiments, IL-1 modulators are IL-1 traps that attach to IL-1 and neutralize IL-1 before it can bind to cell surface receptors and include, but are not limited to, riloncept (Arcalyst®).

RNAi

[00296] In some embodiments, where inhibition or down-regulation of a target is desired (e.g. genes encoding one or more calcineurins, IKKs, TACEs, TLRs, or cytokines), RNA interference are utilized. In some embodiments, the agent that inhibits or down-regulates the target is an siRNA molecule. In certain instances, the siRNA molecule inhibits the transcription of a target by RNA interference (RNAi). In some embodiments, a double stranded RNA (dsRNA) molecule with sequences complementary to a target is generated (e.g. by PCR). In some embodiments, a 20-25 bp siRNA molecule with sequences complementary to a target is generated. In some embodiments, the 20-25 bp siRNA molecule has 2-5 bp overhangs on the 3' end of each strand, and a 5' phosphate terminus and a 3' hydroxyl terminus. In some embodiments, the 20-25 bp siRNA molecule has blunt ends. For techniques for generating RNA sequences see Molecular Cloning: A Laboratory Manual, second edition (Sambrook et al., 1989) and Molecular Cloning: A Laboratory Manual, third edition (Sambrook and Russel, 2001), jointly referred to herein as "Sambrook"; Current Protocols in Molecular Biology (F. M. Ausubel et al., eds., 1987, including supplements through 2001); Current Protocols in Nucleic Acid Chemistry John Wiley & Sons, Inc., New York, 2000) which are hereby incorporated by reference for such disclosure.

[00297] In some embodiments, the dsRNA or siRNA molecule is incorporated into a controlled-release auris-acceptable microsphere or microparticle, hydrogel, liposome, actinic radiation curable gel, solvent-release gel, xerogel, paint, foam, in situ forming spongy material, or thermoreversible gel. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is injected into the inner ear. In some embodiments, the auris-acceptable microsphere or microparticle, actinic radiation curable gel, solvent-release gel, hydrogel, liposome, or thermoreversible gel is injected through the round window membrane. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, actinic radiation curable gel, solvent-release gel, nanocapsule or nanosphere or thermoreversible gel is injected into the cochlea, the Organ of Corti, the vestibular labyrinth, or a combination thereof.

[00298] In certain instances, after administration of the dsRNA or siRNA molecule, cells at the site of administration (e.g. the cells of cochlea, Organ of Corti, and/or the vestibular labyrinth) are transformed with the dsRNA or siRNA molecule. In certain instances following transformation, the dsRNA molecule is cleaved into multiple fragments of about 20-25 bp to yield siRNA molecules. In certain instances, the fragments have about 2bp overhangs on the 3' end of each strand.

[00299] In certain instances, an siRNA molecule is divided into two strands (the guide strand and the anti-guide strand) by an RNA-induced Silencing Complex (RISC). In certain instances, the guide strand is incorporated into the catalytic component of the RISC (i.e. argonaute). In certain instances,

the guide strand binds to a complementary target mRNA sequence. In certain instances, the RISC cleaves the target mRNA. In certain instances, the expression of the target gene is down-regulated.

[00300] In some embodiments, a sequence complementary to a target is ligated into a vector. In some embodiments, the sequence is placed between two promoters. In some embodiments, the promoters are orientated in opposite directions. In some embodiments, the vector is contacted with a cell. In certain instances, a cell is transformed with the vector. In certain instances following transformation, sense and anti-sense strands of the sequence are generated. In certain instances, the sense and anti-sense strands hybridize to form a dsRNA molecule which is cleaved into siRNA molecules. In certain instances, the strands hybridize to form an siRNA molecule. In some

[00301] In some embodiments, the vector is incorporated into a controlled-release auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is injected into the inner ear. In some

Aural Pressure Modulators

Aquaporin

[00302] Contemplated for use with the formulations disclosed herein are agents that treat disorders of the auris, and/or modulate the cells and structures of the auris. In certain instances, an aquaporin is involved in fluid homeostasis. In certain instances, AQP2 mRNA is elevated in rats treated with vasopressin above the levels observed in control animals. In certain instances, Aquaporin-1 is expressed in the cochlea and endolymphatic sac. In certain instances, Aquaporin-1 is expressed in the spiral ligament, the Organ of Corti, the scala tympani, and the endolymphatic sac. Aquaporin-3 is expressed in the stria vascularis, the spiral ligament, the Organ of Corti, the spiral ganglion and the endolymphatic sac. In certain instances, aquaporin 2 (AQP2) mRNA is elevated above normal levels in individuals with endolymphatic hydrops.

[00303] Accordingly, some embodiments incorporate the use of agents that modulate an aquaporin. In some embodiments, the aquaporin is aquaporin 1, aquaporin 2 and/or aquaporin 3. In some embodiments, the agent that modulates an aquaporin (e.g. aquaporin 1, aquaporin 2 or aquaporin 3) is an aquaporin antagonist, partial agonist, inverse agonist, neutral or competitive antagonist, allosteric antagonist, and/or orthosteric antagonist. In some embodiments, the aquaporin antagonist, partial agonist, inverse agonist, neutral or competitive antagonist, allosteric antagonist, and/or orthosteric antagonist includes, but is not limited to, substance P; RU-486; tetraethylammonium

(TEA); an anti-aquaporin antibody; a vasopressin and/or a vasopressin receptor antagonist, partial agonist, inverse agonist, neutral or competitive antagonist, allosteric antagonist, and/or orthosteric antagonist; or combinations thereof.

Estrogen-Related Receptor Beta Modulators

5 [00304] Estrogen-related receptor beta (ERR-beta; also known as Nr3b2), an orphan nuclear receptor, is specifically expressed in and controls the development of the endolymph-producing cells of the inner ear: the stria marginal cells in the cochlea and the vestibular dark cells in the ampulla and utricle. (Chen et al. *Dev. Cell.* (2007) 13:325-337). Nr3b2 expression has been localized in the endolymph-secreting stria marginal cells and vestibular dark cells of the cochlea and vestibular
10 apparatus, respectively. Studies in knockout mice have shown that stria marginal cells in these animals fail to express multiple ion channel and transporter genes, suggesting a role in the development and/or function of endolymph producing epithelia. Moreover, conditional knockout of the Nr3b2 gene results in deafness and diminished endolymphatic fluid volume.

[00305] Other studies suggest a role for estrogen-related receptor β /NR3B2 (ERR/Nr3b2) in
15 regulating endolymph production, and therefore pressure in the vestibular/cochlear apparatus. Treatment with antagonists to ERR/Nr3b2 may assist in reducing endolymphatic volume, and thus alter pressure in the auris interna structures. Accordingly, agents which antagonize ERR/Nr3b2 expression, protein production or protein function are contemplated as useful with the formulations disclosed herein.

GAP Junction Proteins

20 [00306] Contemplated for use with the formulations disclosed herein are agents that treat disorders of the auris, and/or modulate the cells and structures of the auris. Gap junctions are intracellular connections. In certain instances, a gap junction connects the cytoplasm of two cells. In certain instances, a gap junction facilitates the passage of small molecules (e.g. IP_3) and ions between the
25 cells. In certain instances, gap junctions are formed of connexins (e.g. six connexins form a connexon and two connexons form a gap junction). There are multiple connexins (e.g. Cx23, Cx25, Cx26, Cx29, Cx30, Cx30.2, Cx30.3, Cx31, Cx31.1, Cx31.9, Cx32, Cx33, Cx36, Cx37, Cx39, Cx40, Cx40.1, Cx43, Cx45, Cx46, Cx47, Cx50, Cx59, and Cx62). In certain instances, Cx26 and Cx43 are expressed in a spiral limbus, a spiral ligament, a stria vascularis, cells of the Organ of Corti. In
30 certain instances, non-syndromic deafness is associated with mutations in genes (e.g. GJB2) encoding connexins (e.g. Cx26). In certain instances, sensorineural hearing loss is associated with mutations in genes encoding connexins (e.g. Cx26). In certain instances, the expression of Cx26 and Cx43 is upregulated in a cholesteatoma. In certain instances, the expression of Cx26 is upregulated following acoustic trauma. In certain instances, gap junctions facilitate the movement of K^+ ions in
35 endolymph.

[00307] Accordingly, some embodiments disclosed herein incorporate the use of agents that modulate gap junction proteins. In some embodiments, the gap junction protein is a connexin. In some embodiments, the agent that modulates a connexin is a connexin agonist, partial agonist, and/or positive allosteric modulator of a connexin. In some embodiments, the connexin agonist, partial agonist, and/or positive allosteric modulator includes, but is not limited to, astaxanthin; rotigaptide; adenosine; corticotropin-releasing hormone; or combinations thereof.

Vasopressin and the Vasopressin Receptor

[00308] Vasopressin (VP) is a hormone that plays an important part in circulatory and water homeostasis. This hormone is synthesised by neurosecretory cells located predominantly in two specific hypothalamic nuclei—the supraoptic nucleus and the paraventricular nucleus. These neurons have axons that terminate in the neural lobe of the posterior pituitary gland (neurohypophysis) in which they release vasopressin. The three vasopressin receptor subtypes (VP1a, VP1b and VP2) all belong to the G-protein coupled receptor family and have differing tissue distributions. The VP1a receptor is predominantly located in the vascular smooth muscle, hepatocytes and blood platelets. The VP1b receptors are found in the anterior pituitary. The VP2 receptors are localized in the collecting duct of the kidney and regulate the presentation of aquaporin-2 channels at the apical cell surface. The effect of modulation of the VP2 subtype provides readily observed changes in urine volume and electrolyte levels to determine the pharmacological effects of anti-diuresis.

[00309] Vasopressin regulates systemic osmolality by controlling urinary volume and composition. Vasopressin is secreted in response to increases in plasma tonicity (very sensitive stimulus) or to decreases in plasma volume (less sensitive stimulus). Vasopressin mainly regulates urinary volume by binding to the VP receptor in the collecting duct of the kidney. The VP receptor also exists in the inner ear of rodents, and aquaporin-2 (AQP2), a VP mediated water channel protein, is also expressed (Kitano et al. Neuroreport (1997), 8:2289-92). Water homeostasis of the inner ear fluid was confirmed to be regulated using the VP-AQP2 system (Takeda et al. Hear Res (2000), 140:1-6; Takeda et al. Hear Res. (2003), 182:9-18). A recent study looked at tissue expression of VP2 and AQP2 in human endolymphatic sac by immunohistochemistry and noted that VP2 and AQP2 were located in the epithelial layer of the endolymphatic sac but not in surrounding connective tissue (Taguchi et al, Laryngoscope (2007), 117:695-698). Studies on the systemic administration of vasopressin in the guinea pig showed the development of endolymphatic hydrops (Takeda et al. Hear Res (2000), 140:1-6). Additionally, the aquaporin-4 knockout mouse, while otherwise healthy, is deaf (Beitz et al., Cellular and Molecular Neurobiology (2003) 23(3):315-29). This suggests that transport of water and solutes in a manner similar to that of the kidney may play a role in fluid homeostasis of the endolymphatic sac. A mutant human VP2 receptor protein (D136A) has been identified and characterized as constitutively active (Morin et al., FEBS Letters (1998) 441(3):470-

5). This hormone-independent activation of the VP2 receptor could play a role in the etiology of conditions such as Meniere's disease.

[00310] Contemplated for use with the formulations disclosed herein are agents that treat disorders of the auris, and/or modulate the cells (e.g., auris sensory cells) and structures of the auris. In certain instances, VP is involved in fluid homeostasis. In certain instances, VP is involved in endolymph and/or perilymph homeostasis. In certain instances, an increase in endolymph volume increases pressure in the vestibular and cochlear structures. In certain instances, plasma levels of VP are elevated above normal levels in endolymphatic hydrops and/or Meniere's Disease.

Vasopressin Receptor Modulators


[00311] Vasopressin receptor modulators can be differentiated based upon their efficacy relative to the vasopressin peptide hormone. A vasopressin receptor full agonist is a mimic of the native peptide. A vasopressin receptor antagonist blocks the effect of the native peptide. A partial agonist can serve as a mimic of the native peptide and induce a partial response, or in the presence of elevated levels of the native peptide, a partial agonist competes with the native peptide for receptor occupancy and provides a reduction in efficacy, relative to the native peptide alone. For a vasopressin receptor with constitutive activity, an inverse agonist serves to reverse the activity of the receptor.

[00312] Accordingly, some embodiments incorporate the use of agents that modulate vasopressin and/or a vasopressin receptor. In some embodiments, the agent that modulates vasopressin and/or a vasopressin receptor is a vasopressin and/or a vasopressin receptor antagonist, partial agonist, inverse agonist, neutral or competitive antagonist, allosteric antagonist, and/or orthosteric antagonist. In some embodiments, the vasopressin and/or a vasopressin receptor antagonist, partial agonist, inverse agonist, neutral or competitive antagonist, allosteric antagonist, and/or orthosteric antagonist includes, but is not limited to, an anti-vasopressin antibody; an anti-vasopressin receptor antibody; lithium; OPC-31260 ((±)-5-dimethylamino-1-(4-[2-methylbenzoylamino]benzoyl)-2,3,4,5-tetrahydro-1H-benzazepin hydrochloride); WAY-140288 (N-[4-[3-(Dimethylaminomethyl)-10,11-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-10-ylcarbonyl]-2-methoxyphenyl]biphenyl-2-carboxamide); CL-385004 (5-Fluoro-2-methyl-N-[5-(5H-pyrrolo[2,1-c][1,4]benzodiazepine-10(11H)-yl carbonyl)-2-pyridinyl]benzamide); relcovaptan, lixivaptan (VPA-985); tolvaptan; conivaptan; SR 121463A (1-(4-(N-tert-butylcarbonyl)-2-methoxybenzenesulfonyl)-5-ethoxy-3-spiro-(4-(2-morpholinoethoxy)cyclohexane)indol-2-one fumarate); SR-49059 ((2S)-1-[[[(2R,3S)-5-Chloro-3-(2-chlorophenyl)-1-[(3,4-dimethoxyphenyl)sulfonyl]-2,3-dihydro-3-hydroxy-1H-indol-2-yl]carbonyl]-2-pyrrolidinecarboxamide), Lixivaptan (VPA 985); AC-94544 (ACADIA Pharmaceuticals Inc.); AC-88324 (ACADIA Pharmaceuticals Inc.); AC-110484 (ACADIA Pharmaceuticals Inc.); or combinations thereof.

[00313] Recent studies have suggested a role for vasopressin in regulating auris interna pressure by regulating endolymph production, thereby mediating the pressure present in vestibular and cochlear structures. (Takeda et al. *Hearing Res.* (2006) 218:89-97). Treatment with vasopressin antagonists, including OPC-31260, resulted in the marked reduction of Meniere's disease symptoms.

5 Accordingly, vasopressin antagonists are contemplated as useful with the formulations disclosed herein. Examples of vasopressin antagonists include, but are not limited to OPC-31260, WAY-140288, CL-385004, tolvaptan, conivaptan, SR 121463A, VPA 985, valium (diazepam), benzodiazepines and combinations thereof. Testing of vasopressin antagonists may include testing and calculating hydrops reduction with treatment in a guinea pig animal model. See, e.g., Chi et al.
10 "The quantification of endolymphatic hydrops in an experimental animal model with guinea pigs", *J. Oto-Rhino-Larynol.* (2004) 66:56-61.

[00314] Agonists of the VP2 receptor are known, including OPC-51803 and related analogs (Kondo et al., *J. Med. Chem.* (2000) 43:4388; Nakamura et al., *Br. J. Pharmacol.* (2000) 129(8):1700; Nakamura et al., *J. Pharmacol. Exp. Ther.* (2000) 295(3):1005) and WAY-VNA-932 (Caggiano,
15 *Drugs Gut* (2002) 27(3):248). Antagonists of the VP2 receptor include lixivaptan, tolvaptan, conivaptan, SR-121463 and OPC-31260 (Martin et al., *J. Am. Soc. Nephrol.* (1999) 10(10):2165; Gross et al., *Exp. Physiol.* (2000) 85: Spec No 253S; Wong et al., *Gastroent* April 2000, vol 118, 4 Suppl. 2, Part 1); Norman et al., *Drugs Fut.* (2000), 25(11):1121; Inoue et al., *Clin. Pharm. Therap.* (1998) 63(5):561). In testing against the constitutively activated D136A mutant VP2 receptor, SR-
20 1211463 and OPC-31260 behaved as inverse agonist (Morin et al., *FEBS Letters* (1998) 441(3):470-75).



NMDA Receptor Modulators

[00315] Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing disorders such as tinnitus. Accordingly, some embodiments incorporate the use of agents which modulate NMDA receptors.

[00316] In certain instances, the over-activation of the NMDA glutamate receptors by the binding of excessive amounts of glutamate, results in the excessive opening of the ion channels under their control. In certain instances, this results in abnormally high levels of Ca^{2+} and Na^{+} entering the neuron. In certain instances, the influx of Ca^{2+} and Na^{+} into the neuron activates multiple enzymes including, but not limited to, phospholipases, endonucleases, and proteases. In certain instances, the over-activation of these enzymes results in tinnitus, and/or damage to the cytoskeleton, plasma membrane, mitochondria, and DNA of the neuron. In certain instances, the NMDA receptor modulator neramexane treats, and/or ameliorates the symptoms of tinnitus.

[00317] In some embodiments, the agent that modulates the NMDA receptor is an NMDA receptor antagonist. In some embodiments, the agent that modulates an NMDA receptor is an NMDA receptor antagonist, partial agonist, inverse agonist, neutral or competitive antagonist, allosteric

antagonist, and/or orthosteric antagonist. In some embodiments, the agent which antagonizes the NMDA receptor includes, but is not limited to, 1-aminoadamantane, dextromethorphan, dextrorphan, ibogaine, ketamine, nitrous oxide, phencyclidine, riluzole, tiletamine, memantine, neramexane, dizocilpine, aptiganel, remacimide, 7-chlorokynurenate, DCKA (5,7-dichlorokynurenic acid), kynurenic acid, 1-aminocyclopropanecarboxylic acid (ACPC), AP7 (2-amino-7-phosphonoheptanoic acid), APV (R-2-amino-5-phosphonopentanoate), CPPene (3-[(R)-2-carboxypiperazin-4-yl]-prop-2-enyl-1-phosphonic acid); (+)-(1S, 2S)-1-(4-hydroxy-phenyl)-2-(4-hydroxy-4-phenylpiperidino)-1-propanol; (1S, 2S)-1-(4-hydroxy-3-methoxyphenyl)-2-(4-hydroxy-4-phenylpiperidino)-1-propanol; (3R, 4S)-3-(4-(4-fluorophenyl)-4-hydroxypiperidin-1-yl)-chroman-4,7-diol; (1R*, 2R*)-1-(4-hydroxy-3-methylphenyl)-2-(4-(4-fluoro-phenyl)-4-hydroxypiperidin-1-yl)-propan-1-ol-mesylate; and/or combinations thereof.

ENaC Receptor Modulators

[00318] The epithelial sodium channel (ENaC, sodium channel non-neuronal 1 (SCNN1) or amiloride sensitive sodium channel (ASSC)) is a membrane-bound ion-channel that is permeable for Li^+ -ions, protons and Na^+ -ions. The ENaC is located in the apical membrane of polarized epithelial cells and is involved in transepithelial Na^+ -ion transport. Na^+/K^+ -ATPase is also involved in Na^+ transport and ion homeostasis.

[00319] ENaC plays a role in the Na^+ - and K^+ -ion homeostasis of blood, epithelia and extraepithelial fluids by resorption of Na^+ -ions. Modulators of the activity of ENaC modulate aural pressure and include, by way of example, the mineralcorticoid aldosterone, triamterene, and amiloride.

Osmotic Diuretics

[00320] Contemplated for use with the compositions disclosed herein, are agents which regulate aural pressure. Accordingly, some embodiments comprise osmotic diuretics. An osmotic diuretic is a substance that produces an osmotic gradient between two spaces. In certain instances, an osmotic diuretic produces an osmotic gradient between the endolymphatic and perilymphatic spaces. In certain instances, an osmotic gradient between the endolymphatic and perilymphatic spaces exerts a dehydrating effect on the endolymphatic space. In certain instances, dehydrating the endolymphatic space decreases aural pressure.

[00321] Accordingly, in some embodiments of the compositions and formulations disclosed herein, the aural pressure modulator is an osmotic diuretic. In some embodiments, the osmotic diuretic is erythritol, mannitol, glucose, isosorbide, glycerol; urea; or combinations thereof.

[00322] In some instances, contemplated for use in combination with the aural pressure modulating formulations disclosed herein are diuretic agents. A diuretic agent is a drug that elevates the rate of urination. Such diuretics include triamterene, amiloride, bendroflumethiazide, hydrochlorothiazide, furosemide, torsemide, bumetanide, acetazolamide, dorzolamide and combinations thereof.

Progesterone Receptors

[00323] Contemplated for use with the formulations disclosed herein are otic therapeutic agents that treat disorders (e.g., inflammation) of the auris, and/or modulate the cells and structures of the auris. Progesterone is a steroidal hormone. In certain instances, progesterone is a ligand for a progesterone receptor. In certain instances, progesterone is found in the brain. In certain instances, progesterone affects synaptic functioning. In certain instances, progesterone is associated with partial or complete loss of hearing. In certain instances, females taking progesterone and estrogen experienced greater hearing loss than females taking estrogen alone (e.g. about 10% to about 30%).

[00324] Accordingly, some embodiments incorporate the use of agents that modulate progesterone and/or a progesterone receptor. In some embodiments, the agent that modulates progesterone and/or a progesterone receptor is a progesterone and/or progesterone receptor antagonist, a partial agonist, an inverse agonist, a neutral or competitive antagonist, an allosteric antagonist, and/or an orthosteric antagonist. In other embodiments, the agent that modulates progesterone and/or a progesterone receptor includes, but is not limited to, RU-486 ((11b,17 b)-11-[4-(Dimethylamino)phenyl]-17-hydroxy-17-(1-propyn yl)-estra-4,9-dien-3-one); CDB-2914 (17 α -acetoxy-11 β -[4-N,N-dimethylaminophenyl]-19-norpregna-4,9-diene-3,20-dione); CDB-4124 (17 α -acetoxy-21-methoxy-11 β -[4-N,N-dimethylaminophenyl]-19-norpregna-4,9- diene-3,20-dione); CDB-4453 (17 α -acetoxy-21-methoxy-11 β -[4-N-methylaminophenyl]-19-norpregna-4,9-diene-3,20-dione); RTI 3021-022 (Research Triangle Institute); ZK 230211 (11-(4-acetylphenyl)-17-hydroxy-17-(1,1,2,2,2-pentafluoroethyl)estra-4,9-dien-3-one); ORG 31710 (11-(4-dimethylaminophenyl)-6-methyl-4',5'-dihydro(estra-4,9-diene-17,2'-(3H)-furan)-3-one); ORG 33628 (Organon); onapristone (ZK 98299); asoprisnil; ulipristal; a anti-progesterone antibody; an anti-progesterone receptor antibody; or combinations thereof.

Prostaglandins

[00325] Prostaglandins are members of a group of fatty-acid derived compounds and depending upon the subtype, participate in a variety of functions, including control of constriction or dilation in vascular smooth muscle cells, aggregation or disaggregation of platelets, sensitization of spinal neurons to pain, increase or decrease in intraocular pressure, regulation of inflammatory mediation, regulation of calcium movement, control of hormone regulation and control of hormonal regulation. Prostaglandins have both paracrine and autocrine functions, and are a subclass of eicosanoid compounds.

[00326] Prostaglandin analogues, such as latanoprost, travoprost, unoprostone, minprostin F2 alpha and bimatoprost, have been shown in reduce intra-ocular pressure in glaucoma patients by enhancing the uveoscleral outflow, possibly through vasodilation mechanisms, in addition to effects on the trabecular meshwork. In sensorineural hearing loss animal models, noise exposure induces 8-isoprostaglandin F2 α production in the cochlea, concomitant with an increase in vasoconstriction

and reduced blood flow. Treatment with SQ29548, a specific antagonist of 8-isoprostaglandin F2 α , prevents these noise-induced changes in cochlear blood flow and vascular conductance. Further, the prostaglandin analogue JB004/A improves hearing, and treats, and/or the symptoms of tinnitus and vertigo in patients suffering from Ménière's disease. Inhibition of prostaglandin F2 α function also reduces tinnitus in patients suffering from Meniere's disease, as well as improvements in hearing and vertigo. Finally, prostaglandins have been implicated in chronic inflammation associated with otitis media.

[00327] Accordingly, one embodiment disclosed herein is the use of prostaglandin modulators, including latanoprost, travoprost, unoprostone, minprostin F2-alpha, bimatoprost and SQ29548, and JB004/A (Synphora AB) to ameliorate or decrease inner ear and middle ear disorders, including Meniere's disease, tinnitus, vertigo, hearing loss and otitis media.

RNAi

[00328] In some embodiments, where inhibition or down-regulation of a target is desired (e.g. genes ERR, and Nr3b2), RNA interference are utilized. In some embodiments, the agent that inhibits or down-regulates the target is an siRNA molecule. In certain instances, the siRNA molecule is as described herein.

Cytotoxic Agents

[00329] In some instances, immunomodulators and/or aural pressure modulators are useful in treatment of inflammatory otic disorders.

[00330] Any cytotoxic agent useful for the treatment of otic disorders, e.g., inflammatory diseases of the ear or cancer of the ear, is suitable for use in the formulations and methods disclosed herein. In certain embodiments, the cytotoxic agent is an antimetabolite, an antifolate, an alkylating agent, a DNA intercalator, an anti-TNF agent, an anti-angiogenic agent, an anti-inflammatory agent, and/or an immunomodulatory agent. In some embodiments, the cytotoxic agent is a protein, a peptide, an antibody, DNA, a carbohydrate, an inorganic molecule, or an organic molecule. In certain embodiments, the cytotoxic agents are cytotoxic small molecules. Typically, cytotoxic small molecules are of relatively low molecular weight, e.g., less than 1,000, or less than 600-700, or between 300-700 molecular weight. In some embodiments, the cytotoxic small molecules will also have anti-inflammatory properties.

[00331] In certain embodiments, the cytotoxic agents are methotrexate (RHEUMATREX®, Amethopterin) cyclophosphamide (CYTOXAN®), and thalidomide (THALIDOMID®). All of the compounds can be used to treat cancer, including cancer of the ear. Further, all of the compounds have anti-inflammatory properties and can be used in the formulations and compositions disclosed herein for the treatment of inflammatory disorders of the ear, including AIED.

[00332] Although systemic administration of methotrexate, cyclophosphamide, and thalidomide is currently used to treat or is being investigated for the treatment of otic disorders, such as

inflammatory otic disorders, including AIED, Meniere's disease, and Behçet's disease, as well as cancer of the ear, the cytotoxic agents are not without the potential for serious adverse side effects. Moreover, cytotoxic agents which demonstrate efficacy but are otherwise not approvable because of safety considerations is also contemplated within the embodiments disclosed herein. It is

5 contemplated that localized application of the cytotoxic agents to the target otic structures for treatment of autoimmune and/or inflammatory disorders, as well as cancer of the ear, will result in the reduction or elimination of adverse side effects experienced with systemic treatment. Moreover, localized treatment with the cytotoxic agents contemplated herein will also reduce the amount of agent needed for effective treatment of the targeted disorder due, for example, to increased retention
10 of the active agents in the auris interna and/or media, to the existence of the biological blood barrier in the auris interna, or to the lack of sufficient systemic access to the auris media.

[00333] In some embodiments, cytotoxic agents used in the compositions, formulations, and methods disclosed herein are metabolites, salts, polymorphs, prodrugs, analogues, and derivatives of cytotoxic agents, including methotrexate, cyclophosphamide, and thalidomide. Particularly preferred
15 are metabolites, salts, polymorphs, prodrugs, analogues, and derivatives of cytotoxic agents, e.g., methotrexate, cyclophosphamide, and thalidomide, that retain at least partially the cytotoxicity and anti-inflammatory properties of the parent compounds. In certain embodiments, analogues of thalidomide used in the formulations and compositions disclosed herein are lenalidomide (REVLIMID®) and CC-4047 (ACTIMID®).

20 [00334] Cyclophosphamide is a prodrug that undergoes *in vivo* metabolism when administered systemically. The oxidized metabolite 4-hydroxycyclophosphamide exists in equilibrium with aldophosphamide, and the two compounds serve as the transport forms of the active agent phosphoramidate mustard and the degradation byproduct acrolein. Thus, in some embodiments, preferred cyclophosphamide metabolites for incorporation into the formulations and compositions
25 disclosed herein are 4-hydroxycyclophosphamide, aldophosphamide, phosphoramidate mustard, and combinations thereof.

[00335] Other cytotoxic agents used in the compositions, formulations, and methods disclosed herein, particularly for the treatment of cancer of the ear, are any conventional chemotherapeutic agents, including acridine carboxamide, actinomycin, 17-N-allylamino-17-demethoxygeldanamycin,
30 aminopterin, amsacrine, anthracycline, antineoplastic, antineoplaston, 5-azacytidine, azathioprine, BL22, bendamustine, biricodar, bleomycin, bortezomib, bryostatins, busulfan, calyculin, camptothecin, capecitabine, carboplatin, chlorambucil, cisplatin, cladribine, clofarabine, cytarabine, dacarbazine, dasatinib, daunorubicin, decitabine, dichloroacetic acid, discodermolide, docetaxel, doxorubicin, epirubicin, epothilone, eribulin, estramustine, etoposide, exatecan, exisulind,
35 ferruginol, floxuridine, fludarabine, fluorouracil, fosfestrol, fotemustine, gemcitabine, hydroxyurea, IT-101, idarubicin, ifosfamide, imiquimod, irinotecan, irrofulven, ixabepilone, laniquidar, lapatinib,

lenalidomide, lomustine, lurtotecan, mafosfamide, masoprocol, mechlorethamine, melphalan, mercaptopurine, mitomycin, mitotane, mitoxantrone, nelarabine, nilotinib, oblimersen, oxaliplatin, PAC-1, paclitaxel, pemetrexed, pentostatin, pipobroman, pixantrone, plicamycin, procarbazine, proteasome inhibitors (e.g., bortezomib), raltitrexed, rebeccamycin, rubitecan, SN-38, salinosporamide A, satraplatin, streptozotocin, swainsonine, tariquidar, taxane, tegafur-uracil, temozolomide, testolactone, thioTEPA, tioguanine, topotecan, trabectedin, tretinoin, triplatin tetranitrate, tris(2-chloroethyl)amine, troxacitabine, uracil mustard, valrubicin, vinblastine, vincristine, vinorelbine, vorinostat, and zosuquidar.

Auris Sensory Cell Modulators

[00336] In some instances, immunomodulators and/or aural pressure modulators modulate the function of neurons and/or auris sensory cells. Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and/or hair cells of the auris, promote the growth of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Accordingly, some embodiments incorporate the use of agents which promote the survival of neurons and otic hair cells, and/or the growth of neurons and otic hair cells. In some embodiments, the agent which promotes the survival of otic hair cells is a growth factor. In some embodiments, the growth factor modulator is a growth factor modulator antagonist, partial agonist, inverse agonist, neutral or competitive antagonist, allosteric antagonist, and/or orthosteric antagonist.

Amifostine

[00337] Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Accordingly, some embodiments incorporate the use of agents which rescue neurons and otic hair cells from cisplatin-induced ototoxicity.

[00338] Amifostine (also known as WR-2721, or ETHYOL®) is a cytoprotective agent. In certain instances, it prevents or ameliorates the damage to neuron and otic hair cells caused by cisplatin. In certain instances, doses at or above 40 mg/kg are needed to protect against or ameliorate the ototoxic effects of cisplatin.

Anti-intercellular adhesion molecule -1 antibody

[00339] Contemplated for use with the formulations disclosed herein are antibodies to anti-intercellular adhesion molecule (ICAM). In some instances, ICAM blocks the cascade of reactive oxygen species associated with exposure to noise. In some instances modulation of the cascade of reactive oxygen species associated with exposure to noise ameliorates or reduces the degeneration of

neurons and/or hair cells of the auris. Accordingly, some embodiments incorporate the use of agents that are antibodies to ICAMs (e.g., anti-ICAM-1 Ab, anti-ICAM-2 Ab or the like).

Modulation of Atoh/Math1

[00340] Contemplated for use with the formulations disclosed herein are agents that promote the growth and/or regeneration of neurons and/or otic hair cells. Atoh1 is a transcription factor which binds to an E-box. In certain instances, it is expressed during the development of the hair cells of the vestibular and auditory systems. In certain instances, mice with Atoh1 knocked-out did not develop otic hair cells. In certain instances, adenoviruses expressing Atoh1 stimulate the growth and/or regeneration of otic hair cells in guinea pigs treated with ototoxic antibiotics. Accordingly, some embodiments incorporate modulation of the Atoh1 gene.

[00341] In some embodiments, a subject is administered a vector engineered to carry the human Atoh1 gene (the "Atoh1 vector"). For disclosures of techniques for creating the Atoh1 vector see U.S. Pub. No. 2004/02475750, which is hereby incorporated by reference for those disclosures. In some embodiments, the Atoh1 vector is a retrovirus. In some embodiments, the Atoh1 vector is not a retrovirus (e.g. it is an adenovirus; a lentivirus; or a polymeric delivery system such as METAFFECTENE, SUPERFECT®, EFFECTENE®, or MIRUS TRANSIT).

[00342] In some embodiments, the Atoh1 vector is incorporated into a controlled-release auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is injected into the inner ear. In some embodiments, the auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is injected into the cochlea, the Organ of Corti, the vestibular labyrinth, or a combination thereof.

[00343] In certain instances, after administration of the Atoh1 vector, the Atoh1 vector infects the cells at the site of administration (e.g. the cells of cochlea, Organ of Corti, and/or the vestibular labyrinth). In certain instances the Atoh1 sequence is incorporated into the subject's genome (e.g. when the Atoh1 vector is a retrovirus). In certain instances the therapy will need to be periodically re-administered (e.g. when the Atoh1 vector is not a retrovirus). In some embodiments, the therapy is re-administered annually. In some embodiments, the therapy is re-administered semi-annually. In some embodiments, the therapy is re-administered when the subject's hearing loss is moderate (i.e. the subject cannot consistently hear frequencies less than 41 db to 55 dB) to profound (i.e. the subject cannot consistently hear frequencies less than 90 dB).

[00344] In some embodiments, a subject is administered the Atoh1 polypeptide. In some embodiments, the Atoh1 polypeptide is incorporated into controlled-release auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some embodiments,

the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel. In some embodiments, the auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is injected into the inner ear. In some
5 embodiments, the auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is injected into the cochlea, the Organ of Corti, the vestibular labyrinth, or a combination thereof. In
10 some embodiments, the auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is placed in contact with the round window membrane.

[00345] In some embodiments, a subject is administered a pharmaceutically acceptable agent which
15 modulates the expression of the *Atoh1* gene or activity of the *Atoh1* polypeptide. In some embodiments, the expression of the *Atoh1* gene or activity of the *Atoh1* polypeptide is up-regulated. In some embodiments, the expression of the *Atoh1* gene or activity of the *Atoh1* polypeptide is down-regulated.

[00346] In certain instances, a compound which agonizes or antagonizes *Atoh1* is identified (e.g. by
20 use of a high throughput screen). In some embodiments, a construct is designed such that a reporter gene is placed downstream of an E-box sequence. In some embodiments, the reporter gene is luciferase, CAT, GFP, β -lactamase or β -galactosidase. In certain instances, the *Atoh1* polypeptide binds to the E-box sequence and initiates transcription and expression of the reporter gene. In certain instances, an agonist of *Atoh1* aids or facilitates the binding of *Atoh1* to the E-box sequence, thus
25 increasing transcription and expression of the reporter gene relative to a pre-determined baseline expression level. In certain instances, an antagonist of *Atoh1* blocks the binding of *Atoh1* to the E-box, thus decreasing transcription and expression of the reporter gene relative to a pre-determined baseline expression level.

BRN-3 Modulators

[00347] Contemplated for use with the formulations disclosed herein are agents that promote the
30 growth and/or regeneration of neurons and/or otic hair cells. BRN-3 is a group of transcription factors that include, but are not limited to, BRN-3a, BRN-3b, and BRN-3c. In certain instances, they are expressed in postmitotic hair cells. In certain instances, the hair cells of mice with BRN-3c knocked-out did not develop stereocilia and/or underwent apoptosis. In certain instances, BRN3
35 genes regulate the differentiation of inner ear supporting cells into inner ear sensory cells. Accordingly, some embodiments incorporate modulation of the BRN3 genes, and/or polypeptides.

[00348] In some embodiments, a subject is administered a vector engineered to carry a human BRN-3 gene (the "BRN3 vector"). In some embodiments, the BRN3 vector is a retrovirus. In some embodiments, the BRN3 vector is not a retrovirus (e.g. it is an adenovirus; a lentivirus; or a polymeric delivery system such as METAFECTENE®, SUPERFECT®, EFFECTENE®, or
5 MIRUS' TRANSIT®).

[00349] In some embodiments, the subject is administered the BRN3 vector before, during, or after exposure to an ototoxic agent (e.g. an aminoglycoside or cisplatin), or a sound of sufficient loudness to induce acoustic trauma.

[00350] In some embodiments, the BRN3 vector is incorporated into a controlled-release auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some
10 embodiments, the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is injected into the inner ear. In some embodiments, the auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome,
15 paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is injected into the cochlea, the Organ of Corti, the vestibular labyrinth, or a combination thereof.

[00351] In certain instances, after administration of the BRN3 vector, the BRN3 vector infects the cells at the site of administration (e.g. the cells of cochlea, Organ of Corti, and/or the vestibular labyrinth). In certain instances the BRN3 sequence is incorporated into the subject's genome (e.g.
20 when the BRN3 vector is a retrovirus). In certain instances the therapy will need to be periodically re-administered (e.g. when the BRN3 vector is not a retrovirus).

[00352] In some embodiments, a subject is administered a BRN3 polypeptide. In some embodiments, the BRN3 polypeptide is incorporated into controlled-release auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some embodiments,
25 the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel. In some embodiments, the auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is injected into the inner ear. In some
30 embodiments, the auris-acceptable microsphere or microparticle, hydrogel, liposome, or thermoreversible gel. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome, paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is injected into the cochlea, the Organ of Corti, the vestibular labyrinth, or a combination thereof. In some embodiments, the auris-acceptable microsphere or microparticle, hydrogel, liposome, or
35 thermoreversible gel. In some embodiments, the auris-acceptable microsphere, hydrogel, liposome,

paint, foam, in situ forming spongy material, nanocapsule or nanosphere or thermoreversible gel is placed in contact with the round window membrane.

[00353] In some embodiments, a subject is administered a pharmaceutically acceptable agent which modulates the expression of the BRN3 gene or activity of the BRN3 polypeptide. In some
5 embodiments, the expression of the BRN3 gene or activity of the BRN3 polypeptide is up-regulated. In some embodiments, the expression of the BRN3 gene or activity of the BRN3 polypeptide is down-regulated.

[00354] In some embodiments, a compound which agonizes or antagonizes BRN3 is identified (e.g. by use of a high throughput screen). In some embodiments, a construct is designed such that a
10 reporter gene is placed downstream of a BRN3 binding site. In some embodiments, the BRN3 binding site has the sequence ATGAATTAAT (SBNR3). In some embodiments, the reporter gene is luciferase, CAT, GFP, β -lactamase or β -galactosidase. In certain instances, the BRN3 polypeptide binds to the SBNR3 sequence and initiates transcription and expression of the reporter gene. In certain instances, an agonist of BRN3 aids or facilitates the binding of BRN3 to the SBNR3
15 sequence, thus increasing transcription and expression of the reporter gene relative to a pre-determined baseline expression level. In certain instances, an antagonist of BRN3 blocks the binding of BRN3 to the SBNR3, thus decreasing transcription and expression of the reporter gene relative to a pre-determined baseline expression level.

Carbamates

[00355] Contemplated for use with the formulations disclosed herein are agents that modulate the
20 degeneration of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. In certain instances, carbamate compounds protect neurons and otic hair cells from glutamate-induced excitotoxicity. Accordingly, some embodiments incorporate the use of
25 carbamate compounds. In some embodiments, the carbamate compounds are 2-phenyl-1,2-ethanediol monocarbomates and dicarbamates, derivatives thereof, and/or combinations thereof.

Estrogen Receptors

[00356] In some embodiments, the agent that promotes the survival of otic hair cells is an Estrogen
30 Receptor agonist. In some embodiments, the estrogen receptor agonist is a partial agonist or inverse agonist.

[00357] In certain instances, Estrogen Receptor β (ER β) is expressed in an outer hair cell, an inner hair cell, a spiral ganglion neuron, or combinations thereof. In certain embodiments, agonism of ER α and/or ER β ameliorates hearing loss resulting from acoustic trauma. In certain embodiments, agonism of ER α and/or ER β increases and/or up-regulates the expression of a neurotroph gene
35 and/or the activity of a neurotroph polypeptide (e.g. BDNF). In certain embodiments, antagonism of ER α and/or ER β increases hearing loss resulting from acoustic trauma. In certain embodiments,

antagonism of ER α and/or ER β down-regulates the expression of a neurotroph gene and/or the activity of a neurotroph polypeptide (e.g. BDNF).

[00358] In some embodiments, the ER α agonist is PPT (4,4',4''-(4-Propyl-[1H]-pyrazole-1,3,5-triyl)trisphenol); SKF-82958 (6-chloro-7,8-dihydroxy-3-allyl-1-phenyl-2,3,4,5-tetrahydro-1H-3-benzazepine); estrogen; estradiol; estradiol derivatives, including but not limited to 17- β estradiol, estrone, estriol, synthetic estrogen compositions or combinations thereof. In some embodiments, the ER β agonist is ER β -131, phytoestrogen, MK 101 (bioNovo); VG-1010 (bioNovo); DPN (diarylpropionitrile); ERB-041; WAY-202196; WAY-214156; genistein; estrogen; estradiol; estradiol derivatives, including but not limited to 17- β estradiol, estrone, estriol, synthetic estrogen compositions or combinations thereof. Other ER β agonists include select benzopyrans and triazolo-tetrahydrofluorenones, disclosed in U.S. Patent No. 7,279,499, and Parker et al., Bioorg. & Med. Chem. Ltrs. 16: 4652-4656 (2006), each of which is incorporated herein by reference for such disclosure. In some embodiments, a neurotroph is administered before, after, or simultaneously with an Estrogen Receptor β (ER β) agonist. In some embodiments, the neurotroph is BDNF, CNTF, GDNF, neurotrophin-3, neurotrophin-4, and/or combinations thereof.

Fatty Acids

[00359] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris. Accordingly, some embodiments incorporate the use of fatty acids. In certain instances, the membrane surrounding auditory neurons and the vestibulocochlear nerve comprise fatty acids. In certain instances, a deficiency in omega-3 fatty acid results in a decreased response to auditory stimuli. In certain instances, maternal deficiency of alpha-linolenic acid (ALA) leads to offspring with hearing deficiency. In some embodiments, the fatty acid includes but is not limited to an omega-3 fatty acid, an omega-6 fatty acid, or combinations thereof. In some embodiments, the omega-3 fatty acid is α -Linolenic acid, Stearidonic acid, Eicosatrienoic acid, Eicosatetraenoic acid, Eicosapentaenoic acid, Docosapentaenoic acid, Clupanodonic acid, Docosahexaenoic acid, Tetracosapentaenoic acid, Tetracosahexaenoic acid (Nisinic acid), or combinations thereof. In some embodiments, the omega-3 fatty acid is α -Linolenic acid, docosahexaenoic acid, eicosapentaenoic acid, or combinations thereof. In some embodiments, the omega-6 fatty acid is Linoleic acid, Gamma-linolenic acid, Eicosadienoic acid, Dihomo-gamma-linolenic acid, Arachidonic acid, Docosadienoic acid, Adrenic acid, Docosapentaenoic acid, Calendic acid, or combinations thereof.

Gamma-Secretase Inhibitors

[00360] Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Accordingly, some embodiments incorporate the use of agents which inhibit

Notch1 signaling. Notch1 is a transmembrane polypeptide which participates in cell development. In some embodiments, the agents which inhibit Notch1 signaling are γ -secretase inhibitors. In certain instances, the inhibition of Notch1 by a γ -secretase inhibitor, following treatment with an ototoxic agent, results in the production of otic hair cells. In some embodiments, the γ -secretase inhibitor is

5 LY450139 (hydroxylvaleryl monobenzocaprolactam), L685458 (1S-benzyl-4R[1-[1-S-carbamoyl-2-phenethylcarbamoyl]-1S-3-methylbutylcarbamoyl]-2R-hydroxy-5-phenylpentyl} carbamic acid tert-butyl ester); LY411575 (N²-[(2S)-2-(3,5-difluorophenyl)-2-hydroxyethanoyl]-N¹[(7S)-5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[*bid*]azepin-7yl]-L-alaninamide), MK-0752 (Merck), tarenflurbil, and/or BMS-299897 (2-[(1R)-1-[[[(4-chlorophenyl) sulfonyl](2,5-difluorophenyl)amino]ethyl]-5-

10 fluorobenzenepropanoic acid).

Glutamate-Receptor Modulators

[00361] Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing

15 hairs in the inner ear. Accordingly, some embodiments incorporate the use of agents which modulate glutamate receptors. In some embodiments, the glutamate receptor is the AMPA receptor, the NMDA receptor, and/or a group II or III mGlu receptor.

[00362] In some embodiments, the agent that modulates the AMPA receptor is an AMPA receptor antagonist. In some embodiments, the agent which antagonizes the AMPA receptors is CNQX (6-

20 cyano-7-nitroquinoxaline-2,3-dione); NBQX (2,3-dihydroxy-6-nitro-7-sulfamoylbenzo[*f*]quinoxaline-2,3-dione); DNQX (6,7-dinitroquinoxaline-2,3-dione); kynurenic acid; 2,3-dihydroxy-6-nitro-7-sulfamoylbenzo-*[f]*quinoxaline; or combinations thereof.

[00363] In some embodiments, the agent that modulates the NMDA receptor is an NMDA receptor antagonist. In some embodiments, the agent which antagonizes the NMDA receptor is 1-

25 aminoadamantane, dextromethorphan, dextrorphan, ibogaine, ketamine, nitrous oxide, phencyclidine, riluzole, tiletamine, memantine, dizocilpine, aptiganel, remacimide, 7-chlorokynurenate, DCKA (5,7-dichlorokynurenic acid), kynurenic acid, 1-aminocyclopropanecarboxylic acid (ACPC), AP7 (2-amino-7-phosphonoheptanoic acid), APV (R-2-amino-5-phosphonopentanoate), CPPene (3-[(R)-2-carboxypiperazin-4-yl]-prop-2-enyl-1-phosphonic acid); (+)-(1S, 2S)-1-(4-hydroxy-phenyl)-2-(4-hydroxy-4-phenylpiperidino)-1-propanol; (1S, 2S)-1-(4-hydroxy-3-methoxyphenyl)-2-(4-hydroxy-4-phenylpiperidino)-1-propanol; (3R, 4S)-3-(4-(4-fluorophenyl)-4-hydroxypiperidin-1-yl)-chroman-4,7-diol; (1R*, 2R*)-1-(4-hydroxy-3-methylphenyl)-2-(4-(4-fluoro-phenyl)-4-hydroxypiperidin-1-yl)-propan-1-ol-mesylate; and/or combinations thereof.

30

[00364] In certain instances, the over-activation of the AMPA and NMDA glutamate receptors by the binding of excessive amounts of glutamate, results in the excessive opening of the ion channels

under their control. In certain instances, this results in abnormally high levels of Ca^{2+} and Na^+ entering the neuron. In certain instances, the influx of Ca^{2+} and Na^+ into the neuron activates multiple enzymes including, but not limited to, phospholipases, endonucleases, and proteases. In certain instances, the over-activation of these enzymes results in damage to the cytoskeleton, plasma membrane, mitochondria, and DNA of the neuron. Further, in certain instances, the transcription of multiple pro-apoptotic genes and anti-apoptotic genes are controlled by Ca^{2+} levels.

[00365] The mGlu receptors, unlike the AMPA and NMDA receptors, do not directly control an ion channel. However, in certain instances, they indirectly control the opening of ion channels by the activation of biochemical cascades. The mGlu receptors are divided into three groups. In certain instances, the members of groups II and III reduce or inhibit post-synaptic potentials by preventing or decreasing the formation of cAMP. In certain instances, this causes a reduction in the release of neurotransmitters, especially glutamate. GRM7 is the gene which encodes the mGlu7 receptor, a group III receptor. In certain instances, the agonism of mGlu7 results in a decrease in synaptic concentrations of glutamate. This ameliorates glutamate excitotoxicity.

[00366] In some embodiments, the glutamate receptor is a group II mGlu receptor. In some embodiments, the agent which modulates the group II mGlu receptor is a group II mGlu receptor agonist. In some embodiments, the group II mGlu receptor agonist is LY389795 ((-)-2-thia-4-aminobicyclo-hexane-4,6-dicarboxylate); LY379268 ((-)-2-oxa-4-aminobicyclo-hexane-4,6-dicarboxylate); LY354740 ((+)-2-aminobicyclo-hexane-2,6dicarboxylate); DCG-IV ((2S,2'R,3'R)-2-(2',3'-dicarboxycyclopropyl)glycine); 2R,4R-APDC (2R,4R-4-aminopyrrolidine-2,4-dicarboxylate), (S)-3C4HPG ((S)-3-carboxy-4-hydroxyphenylglycine); (S)-4C3HPG ((S)-4-carboxy-3-hydroxyphenylglycine); L-CCG-I ((2S,1'S,2'S)-2-(carboxycyclopropyl)glycine); and/or combinations thereof.

[00367] In some embodiments, the mGlu receptor is a group III mGlu receptor. In some embodiments, the group III mGlu receptor is mGlu7. In some embodiments, the agent which modulates the group III mGlu receptor is a group III mGlu receptor agonist. In some embodiments, the group III mGlu receptor agonist is ACPT-I ((1S,3R,4S)-1-aminocyclopentane-1,3,4-tricarboxylic acid); L-AP4 (L-(+)-2-Amino-4-phosphonobutyric acid); (S)-3,4-DCPG ((S)-3,4-dicarboxyphenylglycine); (RS)-3,4-DCPG ((RS)-3,4-dicarboxyphenylglycine); (RS)-4-phosphonophenylglycine ((RS)PPG); AMN082 (N'-bis(diphenylmethyl)-1,2-ethanediamine dihydrochloride); DCG-IV ((2S,2'R,3'R)-2-(2',3'-dicarboxycyclopropyl)glycine); and/or combinations thereof. In some embodiments, the mGlu receptor is mGlu7. In some embodiments, the agonist of mGlu7 is AMN082. In some embodiments, the mGlu receptor modulator is 3,5-Dimethyl pyrrole-2,4-dicarboxylic acid 2-propyl ester 4-(1,2,2-trimethyl-propyl) ester (3,5-dimethyl PPP); 3,3'-difluorobenzaldazine (DFB), 3,3'-dimethoxybenzaldazine (DMeOB), 3,3'-dichlorobenzaldazine (DCB) and other allosteric modulators of mGluR₅ disclosed in Mol.

Pharmacol. 2003, 64, 731-740; (E)-6-methyl-2-(phenyldiazenyl)pyridin-3-ol (SIB 1757); (E)-2-methyl-6-styrylpyridine (SIB 1893); 2-methyl-6-(phenylethynyl)pyridine (MPEP), 2-methyl-4-((6-methylpyridin-2-yl)ethynyl)thiazole (MTEP); 7-(Hydroxyimino)cyclopropa[b]chromen-1 α -carboxylate ethyl ester (CPCCOEt), N-cyclohexyl-3-methylbenzo[d]thiazolo[3,2-a]imidazole-2-carboxamide (YM-298198), tricyclo[3.3.3.1]nonanyl quinoxaline-2-carboxamide (NPS 2390); 6-methoxy-N-(4-methoxyphenyl)quinazolin-4-amine (LY 456239); mGluR1 antagonists disclosed in WO2004/058754 and WO2005/009987; 2-(4-(2,3-dihydro-1H-inden-2-ylamino)-5,6,7,8-tetrahydroquinazolin-2-ylthio)ethanol; 3-(5-(pyridin-2-yl)-2H-tetrazol-2-yl)benzonitrile, 2-(2-methoxy-4-(4-(pyridin-2-yl)oxazol-2-yl)phenyl)acetonitrile; 2-(4-(benzo[d]oxazol-2-yl)-2-methoxyphenyl)acetonitrile; 6-(3-methoxy-4-(pyridin-2-yl)phenyl)imidazo[2,1-b]thiazole; (S)-(4-fluorophenyl)(3-(3-(4-fluorophenyl)-1,2,4-oxadiazol-5-yl)piperidin-1-yl)methanone (ADX47273) and/or combinations thereof.

[00368] In some embodiments, a glutamate receptor modulator is a nootropic agent. Contemplated for use with the formulations disclosed herein are nootropic agents that modulate neuronal signalling by activating glutamate receptors. In some instances, nootropic agents treat or ameliorate hearing loss (e.g., NIHL) or tinnitus. Accordingly, some embodiments incorporate the use of nootropic agents including, and not limited to, piracetam, Oxiracetam, Aniracetam, Pramiracetam, Phenylpiracetam (Carphedon), Etiracetam, Levetiracetam, Nefiracetam, Nicoracetam, Rolziracetam, Nebracetam, Fasoracetam, Coluracetam, Dimiracetam, Brivaracetam, Seletacetam, and/or Rolipram for the treatment of NIHL or tinnitus.

Growth Factors

[00369] Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and/or hair cells of the auris, promote the survival and/or growth of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Accordingly, some embodiments incorporate the use of agents which promote the survival of neurons and otic hair cells, and/or the growth of neurons and otic hair cells. In some embodiments, the agent which promotes the survival of otic hair cells is a growth factor. In some embodiments, the growth factor is a neurotroph. In certain instances, neurotrophs are growth factors which prevent cells from initiating apoptosis, repair damaged neurons and otic hair cells, and/or induce differentiation in progenitor cells. In some embodiments, the neurotroph is brain-derived neurotrophic factor (BDNF), ciliary neurotrophic factor (CNTF), glial cell-line derived neurotrophic factor (GDNF), neurotrophin-3, neurotrophin-4, and/or combinations thereof. In some embodiments, the growth factor is a fibroblast growth factor (FGF), an insulin-like growth factor (IGF), an epidermal growth factor (EGF), a platelet-derived growth factor (PGF) and/or agonists thereof. In some embodiments, the growth factor is an agonist of the fibroblast growth factor (FGF) receptor,

standing need for suitable testing methods that are applicable to the development of intratympanic otic compositions suitable for human use. In certain embodiments of such auris-compatible formulations that comprise a dye, the ability to visualize a controlled release otic formulation comprising a dye allows for testing of any otic formulation described herein in human clinical trials.

5 **Diseases of the Ear**

[00123] The formulations described herein are suitable for the treatment and/or prevention of diseases or conditions associated with the middle and inner ear, including the cochlea, including vertigo, tinnitus, hearing loss, otosclerosis, balance disorders, and Ménière's disease (endolymphatic hydrops).

10 [00124] The formulations described herein reduce, reverse and/or ameliorate symptoms of otic disorders (e.g., auris interna disorders) which include but are not limited to hearing loss, nystagmus, vertigo, tinnitus, inflammation, swelling, infection and congestion. These disorders may have many causes, such as infection, injury, inflammation, tumors and adverse response to drugs or other chemical agents.

15 *Ménière's Disease*

[00125] Ménière's Disease is an idiopathic condition characterized by sudden attacks of vertigo, nausea and vomiting that may last for 3 to 24 hours, and may subside gradually. Progressive hearing loss, tinnitus and a sensation of pressure in the ears accompanies the disease through time. The cause of Ménière's disease is likely related to an imbalance of auris interna fluid homeostasis, including an increase in production or a decrease in resorption of auris interna fluid.

20 [00126] The cause of symptoms associated with Ménière's disease is likely an imbalance of inner ear fluid homeostasis, including an increase in production or a decrease in reabsorption of inner ear fluid.

[00127] Although the cause of Ménière's disease is unknown, certain evidence suggests a viral etiology for the disease. Specifically, histopathologic analysis of temporal bones in patients with Ménière's disease revealed viral ganglionitis. Also, viral DNA has been observed in the ganglia of patients with Ménière's disease at a higher rate than in healthy patients. Oliveira et al. *ORL* (2008) 70: 42-51. Based on these studies, a pilot study of intratympanic injection of the antiviral agent ganciclovir was conducted, resulting in an improvement of patients suffering from Ménière's disease. Guyot et al. *ORL* (2008) 70: 21-27. Accordingly, controlled release formulations disclosed herein comprising antiviral agents, e.g., ganciclovir, acyclovir, famovir, and valgancyclovir, is administered to the ear for localized treatment of Ménière's disease.

30 [00128] Recent studies of the vasopressin (VP)-mediated aquaporin 2 (AQP2) system in the auris interna suggest a role for VP in inducing endolymph production, thereby increasing pressure in the vestibular and cochlear structures. (Takeda et al. *Hearing Res.* (2006) 218:89-97). VP levels were found to be upregulated in endolymphatic hydrops (Ménière's Disease) cases, and chronic

administration of VP in guinea pigs was found to induce endolymphatic hydrops. Treatment with VP antagonists, including infusion of OPC-31260 (a competitive antagonist of V_2 -R) into the scala tympani resulted in a marked reduction of Meniere's disease symptoms. (Takeda et al. *Hearing Res.* (2003) 182:9-18). Other VP antagonists include WAY-140288, CL-385004, tolvaptan, conivaptan, SR 121463A and VPA 985. (Sanghi et al. *Eur. Heart J.* (2005) 26:538-543; Palm et al. *Nephrol. Dial Transplant* (1999) 14:2559-2562).

[00129] Other studies suggest a role for estrogen-related receptor β /NR3B2 (ERR/Nr3b2) in regulating endolymph production, and therefore pressure in the vestibular/cochlear apparatus. (Chen et al. *Dev. Cell.* (2007) 13:325-337). Knock-out studies in mice demonstrate the role of the protein product of the Nr3b2 gene in regulating endolymph fluid production. Nr3b2 expression has been localized in the endolymph-secreting strial marginal cells and vestibular dark cells of the cochlea and vestibular apparatus, respectively. Moreover, conditional knockout of the Nr3b2 gene results in deafness and diminished endolymphatic fluid volume. Treatment with antagonists to ERR/Nr3b2 may assist in reducing endolymphatic volume, and thus alter pressure in the auris interna structures.

[00130] Other treatments are aimed at dealing with the immediate symptoms and prevention of recurrence. Low-sodium diets, avoidance of caffeine, alcohol, and tobacco have been advocated. Medications that may temporarily relieve vertigo attacks include antihistamines (including meclizine (Antivert, Bonine, Dramamine, Driminate) and other antihistamines), and central nervous system agents, including barbiturates and/or benzodiazepines, including lorazepam or diazepam. Other examples of drugs that are useful in relieving symptoms include muscarinic antagonists, including scopolamine. Nausea and vomiting are relieved by suppositories containing antipsychotic agents, including the phenothiazine agent prochlorperazine (Compazine, Buccastem, Stemetil and Phenotil).

[00131] Surgical procedures have also been used to relieve symptoms of Meniere's disease, including destruction of vestibular function to relieve vertigo symptoms. These procedures aim to either reduce fluid pressure in the inner ear and/or to destroy inner ear balance function. An endolymphatic shunt procedure, which relieves fluid pressure, are placed in the inner ear to relieve symptoms of vestibular dysfunction. Severing of the vestibular nerve may also be employed, which may control vertigo while preserving hearing.

[00132] Another approach to destruction of vestibular function for the treatment of severe Meniere's disease is intratympanic application of an agent that destroys sensory hair cell function in the vestibular system, thereby eradicating inner ear balance function. Various antimicrobial agents are used in the procedure, including aminoglycosides such as gentamicin and streptomycin. The agents are injected through the tympanic membrane using a small needle, a tympanostomy tube with or without a wick, or surgical catheters. Various dosing regimens are used to administer the antimicrobial agents, including a low dose method in which less of the agents are administered over longer periods of time (e.g., one month between injections), and high dose methods in which more

of the agents are administered over a shorter time frame (e.g., every week). Although the high dose method is typically more effective, it is more risky, as it may result in hearing loss.

[00133] Accordingly, formulations disclosed herein are also useful for administration of antimicrobial agents, e.g., gentamicin and streptomycin, for disabling the vestibular apparatus to treat Meniere's disease. The formulations disclosed herein are used to maintain a steady release of the active agents inside the tympanic membrane, thereby avoiding the need for multiple injections or the insertion of a tympanostomy tube. Further, by keeping the active agents localized in the vestibular system, the formulations disclosed herein can also be used to administer higher doses of the antimicrobial agents with a decreased risk of hearing loss.

Meniere's Syndrome

[00134] Meniere's Syndrome, which displays similar symptoms as Meniere's disease, is attributed as a secondary affliction to another disease process, e.g. thyroid disease or auris interna inflammation due to syphilis infection. Meniere's syndrome, thus, are secondary effects to various process that interfere with normal production or resorption of endolymph, including endocrine abnormalities, electrolyte imbalance, autoimmune dysfunction, medications, infections (e.g. parasitic infections) or hyperlipidemia. Treatment of patients afflicted with Meniere's Syndrome is similar to Meniere's Disease.

Sensorineural Hearing Loss

[00135] Sensorineural hearing loss is a type of hearing loss which results from defects (congenital and acquired) in the vestibulocochlear nerve (also known as cranial nerve VIII), or sensory cells of the inner ear. The majority of defects of the inner ear are defects of otic hair cells.

[00136] Aplasia of the cochlea, chromosomal defects, and congenital cholesteatoma are examples of congenital defects which can result in sensorineural hearing loss. By way of non-limiting example, inflammatory diseases (e.g. suppurative labyrinthitis, meningitis, mumps, measles, viral syphilis, and autoimmune disorders), Meniere's Disease, exposure to ototoxic drugs (e.g. aminoglycosides, loop diuretics, antimetabolites, salicylates, and cisplatin), physical trauma, presbycusis, and acoustic trauma (prolonged exposure to sound in excess of 90 dB) can all result in acquired sensorineural hearing loss.

[00137] If the defect resulting in sensorineural hearing loss is a defect in the auditory pathways, the sensorineural hearing loss is called central hearing loss. If the defect resulting in sensorineural hearing loss is a defect in the auditory pathways, the sensorineural hearing loss is called cortical deafness.

[00138] In some instances, sensorineural hearing loss occurs when the components of the auris interna or accompanying neural components are affected, and may contain a neural, i.e. when the auditory nerve or auditory nerve pathways in the brain are affected, or sensory component. Sensory hearing loss are hereditary, or it are caused by acoustic trauma (i.e. very loud noises), a viral

infection, drug-induced or Meniere's disease. Neural hearing loss may occur as a result of brain tumors, infections, or various brain and nerve disorders, such as stroke. Some hereditary diseases, such as Refsum's disease (defective accumulation of branched fatty acids), may also cause neural disorders affecting hearing loss. Auditory nerve pathways are damaged by demyelinating diseases, e.g. idiopathic inflammatory demyelinating disease (including multiple sclerosis), transverse myelitis, Devic's disease, progressive multifocal leukoencephalopathy, Guillain-Barre syndrome, chronic inflammatory demyelinating polyneuropathy and anti-MAG peripheral neuropathy.

[00139] The incidence of sudden deafness, or sensorineural hearing loss, occurs in about 1 in 5000 individuals, and are caused by viral or bacterial infections, e.g. mumps, measles, influenza, chickenpox, cytomegalovirus, syphilis or infectious mononucleosis, or physical injury to the inner ear organ. In some cases, no cause can be identified. Tinnitus and vertigo may accompany sudden deafness, which subsides gradually. Oral corticosteroids are frequently prescribed to treat sensorineural hearing loss. In some cases, surgical intervention are necessary. Other treatments include AM-101 and AM-111, compounds under development for the treatment of auris interna tinnitus and acute sensorineural hearing loss. (Auris Medical AG, Basel, Switzerland).

Noise induced hearing loss

[00140] Noise induced hearing loss (NIHL) is caused upon exposure to sounds that are too loud or loud sounds that last a long time. Hearing loss may occur from prolonged exposure to loud noises, such as loud music, heavy equipment or machinery, airplanes or gunfire. Long or repeated or impulse exposure to sounds at or above 85 decibels can cause hearing loss. NIHL causes damage to the hair cells and/or the auditory nerve. The hair cells are small sensory cells that convert sound energy into electrical signals that travel to the brain. Impulse sound can result in immediate hearing loss that are permanent. This kind of hearing loss are accompanied by tinnitus—a ringing, buzzing, or roaring in the ears or head—which may subside over time. Hearing loss and tinnitus are experienced in one or both ears, and tinnitus may continue constantly or occasionally throughout a lifetime. Permanent damage to hearing loss is often diagnosed. Continuous exposure to loud noise also damages the structure of hair cells, resulting in hearing loss and tinnitus, although the process occurs more gradually than for impulse noise.

[00141] In some embodiments, an otoprotectant can reverse, reduce or ameliorate NIHL. Examples of otoprotectants that treat or prevent NIHL include, but are not limited to, D-methionine, L-methionine, ethionine, hydroxyl methionine, methioninol, amifostine, mesna (sodium 2-sulfanylethanesulfonate), a mixture of D and L methionine, normethionine, homomethionine, S-adenosyl-L-methionine), diethyldithiocarbamate, ebselen (2-phenyl-1, 2-benzisoselenazol-3(2H)-one), sodium thiosulfate, AM-111 (a cell permeable JNK inhibitor, (Laboratoires Auris SAS)), leucovorin, leucovorin calcium, dexrazoxane, or combinations thereof.

[00142] Although there is currently no treatment for noise-induced hearing loss, several treatment regimens have been experimentally developed, including treatment with insulin-like growth factor 1 (IGF-1) and antioxidant therapy, including treatment with alpha lipoic acid. (Lee et al. *Otol. Neurotol.* (2007) 28:976-981).

5 *Tinnitus*

[00143] Tinnitus is defined as the perception of sound in the absence of any external stimuli. It may occur in one or both ears, continuously or sporadically, and is most often described as a ringing sound. It is most often used as a diagnostic symptom for other diseases. There are two types of tinnitus: objective and subjective. The former is a sound created in the body which is audible to
10 anyone. The latter is audible only to the affected individual. Studies estimate that over 50 million Americans experience some form of tinnitus. Of those 50 million, about 12 million experience severe tinnitus.

[00144] In certain instances, tinnitus results from damage to otic structures (e.g. stereocillia), the dysfunction of one or more molecular receptors, and/or the dysfunction of one or more neural
15 pathways. In certain instances, tinnitus results from excitotoxicity caused by abnormal activity of an NMDA receptor. In certain instances, tinnitus results from by dysfunction of an $\alpha 9$ and/or $\alpha 10$ acetylcholine receptor. In certain instances, tinnitus results from damage to the vestibulocochlear nerve. In certain embodiments, a reduction in neurotransmitter reuptake (e.g. the increase in extracellular neurotransmitters) treats, and/or ameliorates the symptoms of tinnitus. In certain
20 embodiments, antagonism of an NK1 receptor treats, and/or ameliorates the symptoms of tinnitus. In certain embodiments, a reduction in neurotransmitter reuptake and antagonism of an NK1 receptor treats, and/or ameliorates the symptoms of tinnitus.

[00145] There are several treatments for tinnitus. Lidocaine, administered by IV, reduces or eliminates the noise associated with tinnitus in about 60-80% of sufferers. Selective neurotransmitter
25 reuptake inhibitors, such as nortriptyline, sertraline, and paroxetine, have also demonstrated efficacy against tinnitus. Benzodiazepines are also prescribed to treat tinnitus.

Autoimmune Inner Ear Disease

[00146] Autoimmune inner ear disease (AIED) is one of the few reversible causes of sensorineural hearing loss. It is a rare disorder appearing in both adults and children that often involves a bilateral
30 disturbance of the audio and vestibular functions of the auris interna. The origin of AIED is likely autoantibodies and/or immune cells attacking inner ear structures, but are associated with other autoimmune conditions. In many cases, AIED occurs without systemic autoimmune symptoms, but up to one-third of patients also suffer from a systemic autoimmune illness, such as inflammatory bowel disease, rheumatoid arthritis, Ankylosing spondylitis, Systemic Lupus Erythematosus (SLE),
35 Sjögren's Syndrome, Cogan's disease, ulcerative colitis, Wegener's granulomatosis and scleroderma. Behçet's disease, a multisystem disease, also commonly has audiovestibular problems.

There is some evidence for food-related allergies as a cause for cochlear and vestibular autoimmunity, but there is presently no agreement as to its importance in the aetiology of the disease. A classification scheme for AIED has been developed (Harris and Keithley, (2002) Autoimmune inner ear disease, in *Otorhinolaryngology Head and Neck Surgery*, 91, 18-32).

5 [00147] The immune system normally performs a crucial role in protecting the auris interna from invasive pathogens such as bacteria and viruses. However, in AIED the immune system itself begins to damage the delicate auris interna tissues. It is well established that the auris interna is fully capable of mounting a localized immune response to foreign antigens. (Harris, *Otolaryngol. Head Neck Surg.* (1983) 91, 18-32). When a foreign antigen enters the auris interna, it is first processed by
 10 immunocompetent cells which reside in and around the endolymphatic sac. Once the foreign antigen has been processed by these immunocompetent cells, these cells secrete various cytokines which modulate the immune response of the auris interna. One result of this cytokine release is to facilitate the influx of inflammatory cells which are recruited from the systemic circulation. These systemic inflammatory cells enter the cochlea via diapedesis through the spiral modiolar vein and its
 15 tributaries and begin to participate in antigen uptake and deregulation just as it occurs in other parts of the body (Harris, *Acta Otolaryngol.* (1990) 110, 357-365). Interleukin 1 (IL-1) plays an important role in modulating the innate (nonspecific) immune response and is a known activator of resting T helper cells and B-cells. T helper cells, once activated by IL-1, produce IL-2. IL-2 secretion results in differentiation of pluripotent T-cells into helper, cytotoxic and suppressor T-cell subtypes. IL-2
 20 also assists T helper cells in the activation of B lymphocytes and probably plays a pivotal role in the immunoregulation of the immune response of the auris interna. IL-2 has been identified within the perilymph of the auris interna as early as 6 h after antigen challenge with peak levels at 18 h after antigen challenge. The perilymphatic levels of IL-2 then dissipate, and it is no longer present within the perilymph at 120 hours post antigen challenge (Gloddek, *Acta Otolaryngol.* (1989) 108, 68-75).

25 [00148] Both IL-1 β and tumor necrosis factor- α (TNF- α) may play a key role in the initiation and amplification of the immune response. IL-1 β is expressed by the fibrocytes of the spiral ligament in the presence of trauma such as surgical trauma or acoustic trauma in a nonspecific response. TNF- α is expressed either by infiltrating systemic cells or by resident cells contained within the endolymphatic sac in the presence of antigen. TNF- α is released as part of the adaptive (specific)
 30 immune response in animal models. When antigen is injected into the auris internas of mice, IL-1 β and TNF- α are both expressed and a vigorous immune response occurs. However, when antigen is introduced to the auris interna via the cerebral spinal fluid without trauma to the auris interna, only TNF- α is expressed and the immune response is minimal (Sato, *J. Assoc. Res. Otolaryngol.* (2003), 4, 139-147). Importantly, cochlear trauma in isolation also results in a minimal immune
 35 response. These results suggest that both the nonspecific and specific components of the immune response may act in concert in the auris interna to achieve a maximal response.

[00149] Accordingly, if the cochlea is traumatized and an antigen is injected (or in the case of autoimmune disease, the patient has immune cells directed against auris interna antigens), both the nonspecific and the specific immune responses can be activated simultaneously. This results in the concurrent production of IL-1 β as well as THF- α which causes a greatly amplified level of inflammation leading to substantial damage to the auris interna. Subsequent experiments in animal models confirm that an important step in immune-mediated damage requires that the auris interna be conditioned by the non-specific innate immune response before the specific adaptive immune response can lead to enough inflammation to result in damage (Hashimoto, *Audiol. Neurotol.* (2005), 10, 35-43). As a result, agents which downregulate or block the specific immune response, and in particular the effect of TNF- α , might be able to prevent the excessive immune response seen when both the specific and nonspecific immune responses are simultaneously activated (Sato, *Laryngoscope* (2002), 112, 1627-1634).

[00150] Treatment of autoimmune ear disease, thus, may consist of anti-TNF agents. Trials using etanercept (ENBREL[®]), an anti-TNF drug, is emerging as a promising agent for treatment of autoimmune inner ear disease. (Rahmen et al., *Otol. Neurol.* (2001) 22:619-624; Wang et al., *Otology & Neurotology* (2003) 24:52-57). Additionally, the anti-TNF agents infliximab (REMICADE[®]) and adalimumab (HUMIRA[®]) may also be useful in treatment of autoimmune auris interna disorders. Trial protocols include injections of anti-TNF agents as an injection on a twice-weekly basis.

[00151] In addition, steroids have been used, e.g. prednisone or decadron, have also been tried with some success. Chemotherapeutic agents, e.g. cytoxan, azathiaprine or methotrexate are used on a long-term basis to treat autoimmune inner ear disorders. (Sismanis et al., *Laryngoscope* (1994) 104:932-934; Sismanis et al., *Otolaryngol* (1997) 116:146-152; Harris et al. *JAMA* (2003) 290:1875-1883). Plasmapheresis procedures have also been tried with some success. (Luetje et al. *Am. J. Otol.* (1997) 18:572-576). Treatment with oral collagen (Kim et al. *Ann. Otol. Rhinol. Laryngol.* (2001) 110: 646-654), gamma globulin infusions or other immune modulating drugs (e.g. beta-interferon, alpha interferon or copaxone) may also be used to treat autoimmune inner ear disorders.

[00152] Certain evidence suggests that viral infection is a factor in the initiation of the inflammatory response that results in AIED. Various autoimmune conditions are induced or enhanced by a variety of DNA and RNA virus infections. Acute or persistent viral infections induce or enhance autoimmune diseases in animal models as well. Similar antigenic determinants have also been observed on viruses and host components. Oldstone, M.B.A. *J. Autoimmun.* (1989) 2(suppl): 187-194. Further, serological tests have identified viral infection in at least one patient diagnosed with a systemic autoimmune disorder that is often associated with AIED (Cogan's syndrome). Garcia-Berrocal, et al. *O.R.L.* (2008) 70: 16-20.

[00153] Accordingly, in some embodiments, controlled release antimicrobial agent compositions and formulations disclosed herein are administered for the treatment of AIED. Particularly, in certain embodiments, formulations disclosed herein comprising antiviral agents are administered for treatment of AIED. In other embodiments, the antimicrobial agent formulations disclosed herein are administered for the treatment of AIED in conjunction with other pharmaceutical agents useful for treating the same conditions or symptoms of the same conditions, including steroids, cytotoxic agents, collagen, gamma globulin infusion, or other immune modulating drugs. Steroids include, e.g., prednisone or decadron. Cytotoxic agents for the treatment of AIED include, e.g., methotrexate, cyclophosphamide, and thalidomide. Plasmapheresis procedures are optionally used. Treatment with oral collagen, gamma globulin infusions, or other immune modulating drugs (e.g. beta-interferon, alpha-interferon or copaxone) is also optionally used in combination with the antimicrobial agent formulations disclosed herein. The additional pharmaceutical agents are optionally administered together with the controlled release formulations disclosed herein, or through other modes of administration, e.g., orally, by injection, topically, nasally or through any other suitable means. The additional pharmaceutical agents are optionally co-administered, or administered at different time periods.

Auditory Nerve Tumors

[00154] Auditory nerve tumors, including acoustic neuroma, acoustic neurinoma, vestibular schwannoma and eighth nerve tumor) are tumors that originate in Schwann cells, cells that wrap around a nerve. Auditory nerve tumors account for approximately 7-8% of all tumors originating in the skull, and are often associated with the diagnosis of neurofibromatosis in a patient. Depending upon the location of the tumor, some symptoms include hearing loss, tinnitus, dizziness and loss of balance. Other more serious symptoms may develop as the tumor becomes larger, which may compress against the facial or trigeminal nerve, which may affect connections between the brain and the mouth, eye or jaw. Smaller tumors are removed by microsurgery, or stereotactic radiosurgical techniques, including fractionated stereotactic radiotherapy. Malignant Schwannomas are treated with chemotherapeutic agents, including vincristine, adriamycin, cyclophosphamide and imidazole carboxamide.

Benign Paroxysmal Positional Vertigo

[00155] Benign paroxysmal positional vertigo is caused by the movement of free floating calcium carbonate crystals (otoliths) from the utricle to one of the semicircular canals, most often the posterior semicircular canal. Movement of the head results in the movement of the otoliths causing abnormal endolymph displacement and a resultant sensation of vertigo. The episodes of vertigo usually last for about a minute and are rarely accompanied by other auditory symptoms.

Cancer of the Ear

[00156] Although the cause is unknown, cancer of the ear is often associated with long-term and untreated otitis, suggesting a link between chronic inflammation and development of the cancer, at

least in some cases. Tumors in the ear can be benign or malignant, and they can exist in the external, middle, or inner ear. Symptoms of ear cancer include otorrhea, otalgia, hearing loss, facial palsy, tinnitus, and vertigo. Treatment options are limited, and include surgery, radiotherapy, chemotherapy, and combinations thereof. Also, additional pharmaceutical agents are used to treat symptoms or conditions associated with the cancer, including corticosteroids in the case of facial palsy, and antimicrobial agents when otitis is present.

[00157] Systemic administration of conventional cytotoxic agents have been used to treat cancer of the ear, including systemic administration of cyclophosphamide (in CHOP chemotherapy) in combination with radiotherapy and methotrexate, Merkus, P., et al. *J. Otorhinolaryngol. Relat. Spec.* (2000) 62:274-7, and perfusion of methotrexate through the external carotid artery, Mahindrakar, N. H. *J. Laryngol. Otol.* (1965) 79:921-5. However, treatments requiring systemic administration of the active agents suffer from the same drawbacks discussed above. Namely, relatively high doses of the agents are required to achieve the necessary therapeutic doses in the ear, which result in an increase of undesired, adverse side effects. Accordingly, local administration of the cytotoxic agents in the compositions and formulations disclosed herein results in treatment of cancer of the ear with lower effective doses, and with a decrease in the incidence and/or severity of side effects. Typical side effects of systemic administration of cytotoxic agents, e.g., methotrexate, cyclophosphamide, and thalidomide, for the treatment of cancer of the ear include anemia, neutropenia, bruising, nausea, dermatitis, hepatitis, pulmonary fibrosis, teratogenicity, peripheral neuropathy, fatigue, constipation, deep vein thrombosis, pulmonary edema, atelectasis, aspiration pneumonia, hypotension, bone marrow suppression, diarrhea, darkening of skin and nails, alopecia, changes in hair color and texture, lethargy, hemorrhagic cystitis, carcinoma, mouth sores, and decreased immunity.

[00158] In certain embodiments, the cytotoxic agents are methotrexate (RHEUMATREX®, Amethopterin) cyclophosphamide (CYTOXAN®), and thalidomide (THALIDOMID®). All of the compounds can be used to treat cancer, including cancer of the ear. Further, all of the compounds have anti-inflammatory properties and can be used in the formulations and compositions disclosed herein for the treatment of inflammatory disorders of the ear, including AIED.

[00159] Although systemic administration of methotrexate, cyclophosphamide, and thalidomide is currently used to treat or is being investigated for the treatment of otic disorders, such as inflammatory otic disorders, including AIED, Meniere's disease, and Behçet's disease, as well as cancer of the ear, the cytotoxic agents are not without the potential for serious adverse side effects. Moreover, cytotoxic agents which demonstrate efficacy but are otherwise not approvable because of safety considerations is also contemplated within the embodiments disclosed herein. It is contemplated that localized application of the cytotoxic agents to the target otic structures for treatment of autoimmune and/or inflammatory disorders, as well as cancer of the ear, will result in the reduction or elimination of adverse side effects experienced with systemic treatment. Moreover,

localized treatment with the cytotoxic agents contemplated herein will also reduce the amount of agent needed for effective treatment of the targeted disorder due, for example, to increased retention of the active agents in the auris interna and/or media, to the existence of the biological blood barrier in the auris interna, or to the lack of sufficient systemic access to the auris media.

5 [00160] In some embodiments, cytotoxic agents used in the compositions, formulations, and methods disclosed herein are metabolites, salts, polymorphs, prodrugs, analogues, and derivatives of cytotoxic agents, including methotrexate, cyclophosphamide, and thalidomide. Particularly preferred are metabolites, salts, polymorphs, prodrugs, analogues, and derivatives of cytotoxic agents, e.g., methotrexate, cyclophosphamide, and thalidomide, that retain at least partially the cytotoxicity and
10 anti-inflammatory properties of the parent compounds. In certain embodiments, analogues of thalidomide used in the formulations and compositions disclosed herein are lenalidomide (REVLIMID®) and CC-4047 (ACTIMID®).

[00161] Cyclophosphamide is a prodrug that undergoes *in vivo* metabolism when administered systemically. The oxidized metabolite 4-hydroxycyclophosphamide exists in equilibrium with
15 aldophosphamide, and the two compounds serve as the transport forms of the active agent phosphoramidate mustard and the degradation byproduct acrolein. Thus, in some embodiments, preferred cyclophosphamide metabolites for incorporation into the formulations and compositions disclosed herein are 4-hydroxycyclophosphamide, aldophosphamide, phosphoramidate mustard, and combinations thereof.

20 [00162] Other cytotoxic agents used in the compositions, formulations, and methods disclosed herein, particularly for the treatment of cancer of the ear, are any conventional chemotherapeutic agents, including acridine carboxamide, actinomycin, 17-N-allylamino-17-demethoxygeldanamycin, aminopterin, amsacrine, anthracycline, antineoplastic, antineoplaston, 5-azacytidine, azathioprine, BL22, bendamustine, biricodar, bleomycin, bortezomib, bryostatin, busulfan, calyculin,
25 camptothecin, capecitabine, carboplatin, chlorambucil, cisplatin, cladribine, clofarabine, cytarabine, dacarbazine, dasatinib, daunorubicin, decitabine, dichloroacetic acid, discodermolide, docetaxel, doxorubicin, epirubicin, epothilone, eribulin, estramustine, etoposide, exatecan, exisulind, ferruginol, floxuridine, fludarabine, fluorouracil, fosfestrol, fotemustine, gemcitabine, hydroxyurea, IT-101, idarubicin, ifosfamide, imiquimod, irinotecan, irofulven, ixabepilone, laniquidar, lapatinib,
30 lenalidomide, lomustine, lurtotecan, mafosfamide, masoprocol, mechlorethamine, melphalan, mercaptopurine, mitomycin, mitotane, mitoxantrone, nelarabine, nilotinib, oblimersen, oxaliplatin, PAC-1, paclitaxel, pemetrexed, pentostatin, pipobroman, pixantrone, plicamycin, procarbazine, proteasome inhibitors (e.g., bortezomib), raltitrexed, rebeccamycin, rubitecan, SN-38, salinosporamide A, satraplatin, streptozotocin, swainsonine, tariquidar, taxane, tegafur-uracil,
35 temozolomide, testolactone, thioTEPA, tioguanine, topotecan, trabectedin, tretinoin, triplatin

tetranitrate, tris(2-chloroethyl)amine, troxacitabine, uracil mustard, valrubicin, vinblastine, vincristine, vinorelbine, vorinostat, and zosuquidar.

Cholesteatoma

[00163] A cholesteatoma is a hyperproliferative cyst often found in the middle ear. Cholesteatoma are classified as congenital or acquired. Acquired cholesteatomas result from retraction of the ear drum (primary) and/or a tear in the ear drum (secondary).

[00164] The most common primary cholesteatoma results from the pars flaccida retracting into the epitympanum. As the pars flaccida continues to retract, the lateral wall of the epitympanum slowly erodes. This produces a defect in the lateral wall of the epitympanum that slowly expands. A less common type of primary acquired cholesteatoma results from the retraction of the posterior quadrant of the tympanic membrane retracts into the posterior middle ear. As the tympanic membrane retracts, squamous epithelium envelops the stapes and retracts into the sinus tympani. Secondary cholesteatomas result from injury to the tympanic membrane (e.g. a perforation resulting from otitis media; trauma; or a surgically-induced injury).

[00165] Complications associated with a growing cholesteatoma include injury to the osteoclasts and, in some cases, deterioration of the thin bone layer separating the top of the ear from the brain. Damage to the osteoclasts results from the persistent application of pressure to the bones resulting from the expansion of the cholesteatoma. Additionally, the presence of multiple cytokines (e.g. TNF- α , TGF- β 1, TGF- β 2, IL-1, and IL-6) in the epithelium of the cholesteatoma can result in further degradation of the surrounding bones.

[00166] Patients with a cholesteatoma often present with earache, hearing loss, mucopurulent discharge, and/or dizziness. Physical examination can confirm the presence of a cholesteatoma. Symptoms which can be identified upon physical examination include damage to the ossicles, and a canal filled with mucopus and granulation tissue.

[00167] There is currently no effective medical therapy for cholesteatomas. As a cholesteatoma has no blood supply, it cannot be treated with systemic antibiotics. Topical administration of antibiotics often fails to treat a cholesteatoma.

Drug-Induced Inner Ear Damage

[00168] Damage from the administration of drugs, including certain antibiotics, diuretics (e.g. ethacrynic acid and furosemide), aspirin, aspirin-like substances (e.g. salicylates) and quinine. Deterioration of the auris interna organ are hastened by impaired kidney function, which results in decreased clearance of the affecting drugs and their metabolites. The drugs may affect both hearing and equilibrium, but likely affects hearing to a greater extent.

[00169] For example, neomycin, kanamycin, amikacin have a greater effect on hearing than on balance. The antibiotics viomycin, gentamicin and tobramycin affect both hearing and equilibrium. Streptomycin, another commonly administered antibiotic, induces vertigo more than loss of hearing,

and can lead to Dandy's syndrome, where walking in the dark becomes difficult and induces a sensation of the environment moving with each step. Aspirin, when taken in very high doses, may also lead to temporary hearing loss and tinnitus, a condition where sound is perceived in the absence of external sound. Similarly, quinine, ethacrynic acid and furosemide can result in temporary or
5 permanent hearing loss.

Excitotoxicity

[00170] Excitotoxicity refers to the death or damaging of neurons and/or otic hair cells by glutamate and/or similar substances.

[00171] Glutamate is the most abundant excitatory neurotransmitter in the central nervous system.

10 Pre-synaptic neurons release glutamate upon stimulation. It flows across the synapse, binds to receptors located on post-synaptic neurons, and activates these neurons. The glutamate receptors include the NMDA, AMPA, and kainate receptors. Glutamate transporters are tasked with removing extracellular glutamate from the synapse. Certain events (e.g. ischemia or stroke) can damage the transporters. This results in excess glutamate accumulating in the synapse. Excess glutamate in
15 synapses results in the over-activation of the glutamate receptors.

[00172] The AMPA receptor is activated by the binding of both glutamate and AMPA. Activation of certain isoforms of the AMPA receptor results in the opening of ion channels located in the plasma membrane of the neuron. When the channels open, Na^+ and Ca^{2+} ions flow into the neuron and K^+ ions flow out of the neuron.

20 [00173] The NMDA receptor is activated by the binding of both glutamate and NMDA. Activation of the NMDA receptor, results in the opening of ion channels located in the plasma membrane of the neuron. However, these channels are blocked by Mg^{2+} ions. Activation of the AMPA receptor results in the expulsion of Mg^{2+} ions from the ion channels into the synapse. When the ion channels open, and the Mg^{2+} ions evacuate the ion channels, Na^+ and Ca^{2+} ions flow into the neuron, and K^+
25 ions flow out of the neuron.

[00174] Excitotoxicity occurs when the NMDA receptor and AMPA receptors are over-activated by the binding of excessive amounts of ligands, for example, abnormal amounts of glutamate. The over-activation of these receptors causes excessive opening of the ion channels under their control. This allows abnormally high levels of Ca^{2+} and Na^+ to enter the neuron. The influx of these levels of
30 Ca^{2+} and Na^+ into the neuron causes the neuron to fire more often, resulting in a rapid buildup of free radicals and inflammatory compounds within the cell. The free radicals eventually damage the mitochondria, depleting the cell's energy stores. Furthermore, excess levels of Ca^{2+} and Na^+ ions activate excess levels of enzymes including, but not limited to, phospholipases, endonucleases, and proteases. The over-activation of these enzymes results in damage to the cytoskeleton, plasma
35 membrane, mitochondria, and DNA of the sensory neuron.

Endolymphatic Hydrops

[00175] Endolymphatic hydrops refers to an increase in the hydraulic pressure within the endolymphatic system of the inner ear. The endolymph and perilymph are separated by thin membranes which contain multiple nerves. Fluctuation in the pressure stresses the membranes and the nerves they house. If the pressure is great enough, disruptions may form in the membranes. This results in a mixing of the fluids which can lead to a depolarization blockade and transient loss of function. Changes in the rate of vestibular nerve firing often lead to vertigo. Further, the Organ of Corti may also be affected. Distortions of the basilar membrane and the inner and outer hair cells can lead to hearing loss and/or tinnitus.

[00176] Causes include metabolic disturbances, hormonal imbalances, autoimmune disease, and viral, bacterial, or fungal infections. Symptoms include hearing loss, vertigo, tinnitus, and aural fullness. Nystagmus may also be present. Treatment includes systemic administration of benzodiazepine, diuretics (to decrease the fluid pressure), corticosteroids, and/or anti-bacterial, anti-viral, or anti-fungal agents.

Hereditary Disorders

[00177] Hereditary disorders, including Scheibe, Mondini-Michelle, Waardenburg's, Michel, Alexander's ear deformity, hypertelorism, Jervell-Lange Nielson, Refsum's and Usher's syndromes, are found in approximately 20% of patients with sensorineural hearing loss. Congenital ear malformations may result from defects in the development of the membranous labyrinthine, the osseous labyrinthine, or both. Along with profound hearing loss and vestibular function abnormalities, hereditary deformities may also be associated with other dysfunctions, including development of recurring meningitis, cerebral spinal fluid (CSF) leaks, as well as perilymphatic fistulas. Treatment of chronic infections are necessitated in hereditary disorder patients.

Inflammatory Disorders of the Auris Media

[00178] Otitis media (OM), which includes acute otitis media (AOM), otitis media with effusion (OME) and chronic otitis media as examples, is a condition affecting both adults and children. OM susceptibility is multifactorial and complex, including environmental, microbial and host factors. Bacterial infection accounts for a large percentage of OM cases, with more than 40% of cases attributed to *Streptococcus pneumoniae* infection. However, viral causes, as well as other microbial agents, may also account for OM conditions.

[00179] Regardless of the causative agent, increases in cytokine production, including interleukins and TNF, have been observed in the effluent media of individuals afflicted with OM. IL-1 β , IL-6 and TNF- α are acute-phase cytokines that promote acute inflammatory response after infection with viruses and bacteria. Genetic studies supports this link between cytokines and OM by demonstrating a correlation in the occurrence of TNF- α SNP (single-nucleotide polymorphisms) and an increased susceptibility for OM in pediatric patients suffering from AOM and with a subsequent need for placement of tympanostomy tubes. (Patel et al. *Pediatrics* (2006) 118:2273-2279). In animal models

of OM induced with pneumococci inoculations, TNF- α and interleukins levels were found to increase in early developmental phase of OM, with TNF- α levels steadily increasing 72 hours after inoculation. Moreover, higher TNF- α levels have been associated with a history of multiple tympanostomy tube placements, indicating a role for TNF- α in chronic OM cases. Finally, direct
5 injection of TNF- α and interleukins has been shown to induce middle ear inflammation in a guinea pig model. These studies support the role that cytokines may play in the origin and maintenance of OM in the auris media.

[00180] Because OM can be caused by a virus, bacteria or both, it is often difficult to identify the exact cause and thus the most appropriate treatment. Treatment options of OM in the auris media
10 include treatment with antibiotics, such as amoxicillin, clavulanate acid, trimethoprim-sulfamethoxazole, cefuroxime, clarithromycin and azithromycin and other cephalosporins, macrolides, penicillins or sulfonamides. Surgical intervention is also available, including a myringotomy, an operation to insert a tympanostomy tube through the tympanic membrane and into the patient's middle ear to drain the fluid and balance the pressure between the outer and middle ear.
15 Antipyretics and analgesics, including benzocaine, ibuprofen and acetaminophen, may also be prescribed to treat accompanying fever or pain effects. Pre-treatment with TNF- α inhibitors in experimental lipopolysaccharide (LPS)-induced OM animal models has been shown to suppress development of OM, suggesting a role in the treatment of OM or OME. In addition, treatment of such conditions include use of TNF- α inhibitors in combination with other inflammatory response
20 mediators, including platelet activating factor antagonists, nitric oxide synthase inhibitors and histamine antagonists.

[00181] As discussed above, methotrexate, cyclophosphamide, and thalidomide are all cytotoxic small molecule agents that are systemically administered to treat AIED. Thus, the compounds are useful in the compositions and formulations disclosed herein for the treatment of inflammatory
25 disorders of the auris media, including OM, by having a direct anti-inflammatory effect, particularly by interfering with TNF activity. In other embodiments, metabolites, salts, polymorphs, prodrugs, analogues, and derivatives of methotrexate, cyclophosphamide, and thalidomide that retain the ability of the parent cytotoxic agents to treat inflammatory disorders of the auris media, including OM, are useful in the formulations disclosed herein for the treatment of inflammatory disorders of
30 the auris media, including OM. In certain embodiments, preferred metabolites of cyclophosphamide for incorporation into the compositions and formulations disclosed herein include 4-hydroxycyclophosphamide, aldophosphamide, phosphoramidate mustard, or combinations thereof.

[00182] In addition, other otic disorders have inflammatory response aspects or are tangentially related to autoimmune conditions, including Meniere's disease and non-sudden hearing loss or noise
35 induced hearing loss. These disorders are also explicitly contemplated as benefiting from the

cytotoxic agent formulations disclosed herein, and therefore are within the scope of the embodiments disclosed.

Inflammatory Disorders of the Auris externa

[00183] Otitis externa (OE), also referred to as swimmer's ear, is an inflammation and/or infection of the external ear. OE is often caused by bacteria in the outer ear, which establish infection following damage to the skin of the ear canal. Primary bacterial pathogens that cause OE are *Pseudomonas aeruginosa* and *Staphylococcus aureus*, but the condition is associated with the presence of many other strains of gram positive and negative bacteria. OE is also sometimes caused by fungal infection in the outer ear, including *Candida albicans* and *Aspergillus*. Symptoms of OE include otalgia, swelling, and otorrhea. If the condition progresses significantly, OE may cause temporary conductive hearing loss as a result of the swelling and discharge.

[00184] Treatment of OE involves eliminating the aggravating pathogen from the ear canal and reducing inflammation, which is usually accomplished by administering combinations of antimicrobial agents, e.g., antibacterial and antifungal agents, with anti-inflammatory agents, e.g., steroids. Typical antibacterial agents for the treatment of OE include aminoglycosides (e.g., neomycin, gentamycin, and tobramycin), polymyxins (e.g., polymyxin B), fluoroquinolone (e.g., ofloxacin and ciprofloxacin), cephalosporins (e.g., cefuroxime, ceflacor, cefprozil, loracarbef, cefindir, cefixime, cefpodoxime proxetil, cefibuten, and ceftriaxone), penicillins (e.g., amoxicillin, amoxicillin-clavulanate, and penicillinase-resistant penicillins), and combinations thereof. Typical antifungal agents for the treatment of OE include clotrimazole, thimerasol, M-cresyl acetate, tolnaftate, itraconazole, and combinations thereof. Acetic acid is also administered to the ear, alone and in combination with other agents, to treat bacterial and fungal infections. Ear drops are often used as the vehicle for administration of the active agents. In the case that ear swelling has progressed substantially and ear drops do not penetrate significantly into the ear canal, a wick can be inserted into the ear canal to facilitate penetration of the treatment solutions. Oral antibiotics are also administered in the case of extensive soft tissue swelling that extends to the face and neck. When the pain of OE is extremely severe such that it interferes with normal activity, e.g., sleeping, pain relievers such as topical analgesics or oral narcotics are given until the underlying inflammation and infection are alleviated.

[00185] Notably, some types of topical ear drops, such as ear drops containing neomycin, are safe and effective for use in the ear canal, but can be irritating and even ototoxic to the auris media, prompting concern that such topical preparations should not be used unless the tympanic membrane is known to be intact. Utilization of the formulations disclosed herein for the treatment of OE allows for use of active agents that are potentially damaging to the auris media, even when the tympanic membrane is not intact. Specifically, the controlled release formulations disclosed herein can be applied locally in the external ear with improved retention time, thus eliminating concern that the

active agents will leak out of the ear canal into the auris media. Furthermore, otoprotectants can be added when ototoxic agents, such as neomycin, are used.

[00186] Treatment of severe OE with the compositions disclosed herein, particularly highly viscous and/or mucoadhesive formulations, also obviates the need for extended use of an ear wick.

5 Specifically, the compositions disclosed herein have increased retention time in the ear canal as a result of the formulation technology, thus eliminating the need for a device to maintain their presence in the outer ear. The formulations can be applied in the outer ear with a needle or an ear dropper, and the active agents can be maintained at the site of inflammation without the aid of an ear wick.

10 [00187] In some embodiments, the treatment of OE with antimicrobial formulations disclosed herein encompasses the treatment of granular myringitis, a specific form of OE characterized by chronic inflammation of the pars tensa of the tympanic membrane. The outer epithelial and underlying fibrous layers of the tympanic membrane are replaced by a proliferating granulation tissue. The predominant symptom is foul-smelling otorrhea. A variety of bacteria and fungi cause the condition,
15 including *Proteus* and *Pseudomonas* species. Accordingly, antimicrobial agent formulations disclosed herein comprising antibacterial or antifungal agents are useful for the treatment of granular myringitis.

[00188] In some embodiments, the treatment of OE with antimicrobial formulations disclosed herein encompasses the treatment of chronic stenosing otitis externa. Chronic stenosing otitis externa is
20 characterized by repeated infections, typically caused by bacteria or fungi. The primary symptoms are pruritus in the ear canal, otorrhea, and chronic swelling. Antimicrobial agent formulations disclosed herein comprising antibacterial or antifungal agents are useful for the treatment of chronic stenosing otitis externa.

[00189] In some embodiments, the treatment of OE with antimicrobial formulations disclosed herein
25 encompasses the treatment of malignant or necrotizing external otitis, an infection involving the temporal and adjacent bones. Malignant external otitis is typically a complication of external otitis. It occurs primarily in persons with compromised immunity, especially in older persons with diabetes mellitus. Malignant external otitis is often caused by the bacteria *Pseudomonas aeruginosa*. Treatment typically involves correction of immunosuppression when possible, in conjunction with
30 antibacterial therapy and pain relievers. According, antimicrobial agent formulations disclosed herein are useful for the treatment of malignant or necrotizing external otitis.

[00190] Otitis media (OM), which includes acute otitis media (AOM), chronic otitis media, otitis media with effusion (OME), secretory otitis media, and chronic secretory otitis media as examples, is a condition affecting both adults and children. OM susceptibility is multifactorial and complex,
35 including environmental, microbial and host factors. Bacterial infection accounts for a large

percentage of OM cases, with more than 40% of cases attributed to *Streptococcus pneumoniae* infection. However, viruses, as well as other microbes, may also account for OM conditions.

[00191] Because OM can be caused by a virus, bacteria or both, it is often difficult to identify the exact cause and thus the most appropriate treatment. Treatment options for OM include antibiotics, such as penicillins (e.g., amoxicillin and amoxicillin-clavulanate), clavulanate acid, trimethoprim-sulfamethoxazole, cephalosporins (e.g., cefuroxime, ceflacor, cefprozil, loracarbef, cefindir, cefixime, cefpodoxime proxetil, cefibuten, and ceftriaxone), macrolides and azalides (e.g., erythromycin, clarithromycin, and azithromycin), sulfonamides, and combinations thereof. Surgical intervention is also available, including myringotomy, an operation to insert a tympanostomy tube through the tympanic membrane and into the patient's middle ear to drain the fluid and balance the pressure between the outer and middle ear. Antipyretics and analgesics, including benzocaine, ibuprofen and acetaminophen, may also be prescribed to treat accompanying fever or pain effects.

[00192] Regardless of the causative agent, increases in cytokine production, including interleukins and TNF, have been observed in the effluent media of individuals afflicted with OM. IL-1 β , IL-6 and TNF- α are acute-phase cytokines that promote acute inflammatory response after infection with viruses and bacteria. Moreover, higher TNF- α levels have been associated with a history of multiple tympanostomy tube placements, indicating a role for TNF- α in chronic OM cases. Finally, direct injection of TNF- α and interleukins has been shown to induce middle ear inflammation in a guinea pig model. These studies support the role that cytokines may play in the origin and maintenance of OM in the auris media. Thus, treatment of OM includes the use of antimicrobial agents in conjunction with anti-inflammatory agents to eliminate the pathogen and treat the symptoms of inflammation. Such treatments include use of steroids, TNF- α inhibitors, platelet activating factor antagonists, nitric oxide synthase inhibitors, histamine antagonists, and combinations thereof in conjunction with the antimicrobial formulations disclosed herein.

[00193] Mastoiditis is an infection of the mastoid process, which is the portion of the temporal bone behind the ear. It is typically caused by untreated acute otitis media. Mastoiditis are acute or chronic. Symptoms include pain, swelling, and tenderness in the mastoid region, as well as otalgia, erythematous, and otorrhea. Mastoiditis typically occurs as bacteria spread from the middle ear to the mastoid air cells, where the inflammation causes damage to the bony structures. The most common bacterial pathogens are *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Staphylococcus aureus*, and gram-negative bacilli. Accordingly, antimicrobial agent formulations disclosed herein comprising antibacterial agents effective against the bacteria are useful for the treatment of mastoiditis, including acute mastoiditis and chronic mastoiditis.

[00194] Bullous myringitis is an infection of the tympanic membrane, caused by a variety of bacteria and viruses, including *Mycoplasma* bacteria. The infection leads to inflammation of the tympanic membrane and nearby canal, and causes the formation of blisters on the ear drum. The primary

symptom of Bullous myringitis is pain, which are relieved through the administration of analgesics. Antimicrobial formulations disclosed herein comprising antibacterial and antiviral agents are useful for the treatment of Bullous myringitis.

[00195] Eustachian tubal catarrh, or Eustachian salpingitis, is caused from inflammation and swelling of the Eustachian tubes, resulting in a build-up of catarrh. Accordingly, antimicrobial formulations disclosed herein are useful for the treatment of Eustachian salpingitis.

[00196] Labyrinthitis, e.g., serous labyrinthitis, is an inflammation of the inner ear that involves one or more labyrinths housing the vestibular system. The primary symptom is vertigo, but the condition is also characterized by hearing loss, tinnitus, and nystagmus. Labyrinthitis maybe acute, lasting for one to six weeks and being accompanied by severe vertigo and vomiting, or chronic, with symptoms lasting for months or even years. Labyrinthitis is typically caused by viral or bacterial infection. Accordingly, antimicrobial formulations disclosed herein comprising antibacterial and antiviral agents are useful for the treatment of labyrinthitis.

[00197] Facial nerve neuritis is a form of neuritis, an inflammation of the peripheral nervous system, afflicting the facial nerve. The primary symptoms of the condition are a tingling and burning sensation, and stabbing pains in the affected nerves. In severe cases, there are numbness, loss of sensation, and paralysis of the nearby muscles. The condition is typically caused by herpes zoster or herpes simplex viral infection, but has also been associated with bacterial infection, e.g., leprosy. Accordingly, antimicrobial formulations disclosed herein comprising antibacterial and antiviral agents are useful for the treatment of facial nerve neuritis.

[00198] In some embodiments, antimicrobial formulations disclosed herein are also useful for the treatment of temporal bone osteoradionecrosis.

Kinetosis

[00199] Kinetosis, also known as motion sickness, is a condition in which there is a disconnection between visually perceived movement and the vestibular system's sense of movement. Dizziness, fatigue, and nausea are the most common symptoms of kinetosis. Dimenhydrinate, cinnarizine, and meclizine are all systemic treatments for kinetosis. Additionally, benzodiazepines and antihistamines have demonstrated efficacy in treating kinetosis.

Labyrinthitis

[00200] Labyrinthitis is an inflammation of the labyrinths of the ear which contain the vestibular system of the inner ear. Causes include bacterial, viral, and fungal infections. It may also be caused by a head injury or allergies. Symptoms of labyrinthitis include difficulty maintaining balance, dizziness, vertigo, tinnitus, and hearing loss. Recovery may take one to six weeks; however, chronic symptoms are present for years.

[00201] There are several treatments for labyrinthitis. Prochlorperazine is often prescribed as an antiemetic. Serotonin-reuptake inhibitors have been shown to stimulate new neural growth within

the inner ear. Additionally, treatment with antibiotics is prescribed if the cause is a bacterial infection, and treatment with corticosteroids and antivirals is recommended if the condition is caused by a viral infection.

Mal de Debarquement

5 [00202] Mal de débarquement is a condition which usually occurs subsequent to a sustained motion event, for example, a cruise, car trip, or airplane ride. It is characterized by a persistent sense of motion, difficulty maintaining balance, fatigue, and cognitive impairment. Symptoms may also include dizziness, headaches, hyperacusis, and/or tinnitus. Symptoms often last in excess of a month. Treatment includes benzodiazepines, diuretics, sodium channel blockers, and tricyclic
10 antidepressants.

Microvascular compression syndrome

[00203] Microvascular compression syndrome (MCS), also called "vascular compression" or "neurovascular compression", is a disorder characterized by vertigo and tinnitus. It is caused by the irritation of Cranial Nerve VII by a blood vessel. Other symptoms found in subjects with MCS
15 include, but are not limited to, severe motion intolerance, and neuralgic like "quick spins". MCS is treated with carbamazepine, TRILEPTAL®, and baclofen. It can also be surgically treated.

Other Microbial Infections Causing Cochleovestibular Disorders

[00204] Other microbial infections are known to cause cochleovestibular disorders, including hearing loss. Such infections include rubella, cytomegalovirus, mononucleosis, varicella zoster
20 (chicken pox), pneumonia, *Borrelia* species of bacteria (Lyme disease), and certain fungal infections. Accordingly, controlled release antimicrobial agent formulations disclosed herein are also used for localized treatment of these infections in the ear.

Otic Disorders caused by Free Radicals

[00205] Free radicals are highly reactive atoms, molecules, or ions the reactivity of which results
25 from the presence of unpaired electrons. Reactive oxygen species ("ROS") form as a result of sequential reduction of molecular oxygen. Examples of reactive oxygen species of interest ("ROS") include, but are not limited to, superoxide, hydrogen peroxide, and hydroxyl radicals. ROS are naturally produced as a by-product of the production of ATP. ROS can also result from the use of cisplatin, and aminoglycosides. Further, stress to stereocilia caused by acoustic trauma results in otic
30 hair cells producing ROS.

[00206] ROS can damage cells directly by damaging nuclear DNA and mitochondrial DNA. Damage to the former can lead to mutations which impair the functioning of auditory cells and/or apoptosis. Damage to the latter often results in decreased energy production and increased ROS
35 production both of which can lead to impaired cellular functioning or apoptosis. Further, ROS can also damage or kill cells by oxidizing the polydesaturated fatty acids which comprise lipids,

oxidizing the amino acids which comprise proteins, and oxidizing co-factors necessary for the activity of enzymes. Antioxidants can ameliorate damage by caused by ROS by preventing their formation, or scavenging the ROS before they can damage the cell.

[00207] Damage to mitochondria by ROS is often seen in hearing loss, especially hearing loss due to aging. The loss of ATP correlates to a loss in neural functioning in the inner ear. It can also lead to physiological changes in the inner ear. Further, damage to mitochondria often results in an increased rate of cellular degradation and apoptosis of inner ear cells. The cells of the stria vascularis are the most metabolically active due to the vast energy requirements needed to maintain the ionic balance of fluids in the inner ear. Thus, the cells of the stria vascularis are most often damaged or killed due to damage of the mitochondria.

Otosclerosis

[00208] Otosclerosis is an abnormal growth of bone in the middle ear, which prevents structures within the ear from transducing vibration, which causes hearing loss. Otosclerosis usually effects the ossicles, in particular the stapes, which rests in the entrance to the cochlea in the oval window.

The abnormal bone growth fixates the stapes onto the oval window, preventing sound passing waves from traveling to the cochlea. Otosclerosis may cause a sensorineural hearing loss, i.e. damaged nerve fibers or hearing hair cells, or conductive hearing loss.

[00209] Treatment of otosclerosis may include surgery to remove the fixated stapes bone, called a stapedectomy. Fluoride treatment may also be used, which will not reverse the hearing loss but may slow the development of otosclerosis.

Ototoxicity

[00210] Ototoxicity refers to hearing loss caused by a toxin. The hearing loss are due to trauma to otic hair cells, the cochlea, and/or the cranial nerve VII. Multiple drugs are known to be ototoxic. Often ototoxicity is dose-dependent. It are permanent or reversible upon withdrawal of the drug.

[00211] Known ototoxic drugs include, but are not limited to, the aminoglycoside class of antibiotics (e.g. gentamicin, and amikacin), some members of the macrolide class of antibiotics (e.g. erythromycin), some members of the glycopeptide class of antibiotics (e.g. vancomycin), salicylic acid, nicotine, some chemotherapeutic agents (e.g. actinomycin, bleomycin, cisplatin, carboplatin and vincristine), and some members of the loop diuretic family of drugs (e.g. furosemide).

[00212] Cisplatin and the aminoglycoside class of antibiotics induce the production of reactive oxygen species ("ROS"). ROS can damage cells directly by damaging DNA, polypeptides, and/or lipids. Antioxidants prevent damage of ROS by preventing their formation or scavenging free radicals before they can damage the cell. Both cisplatin and the aminoglycoside class of antibiotics are also thought to damage the ear by binding melanin in the stria vascularis of the inner ear.

[00213] Salicylic acid is classified as ototoxic as it inhibits the function of the polypeptide prestin. Prestin mediates outer otic hair cell motility by controlling the exchange of chloride and carbonate

across the plasma membrane of outer otic hair cells. It is only found in the outer otic hair cells, not the inner otic hair cells. Accordingly, disclosed herein is the use of controlled release auris-compositions comprising antioxidants to prevent, ameliorate or lessen ototoxic effects of chemotherapy, including but not limited to cisplatin treatment, aminoglycoside or salicylic acid administration, or other ototoxic agents.

Postural Vertigo

[00214] Postural vertigo, otherwise known as positional vertigo, is characterized by sudden violent vertigo that is triggered by certain head positions. This condition are caused by damaged semicircular canals caused by physical injury to the auris interna, otitis media, ear surgery or blockage of the artery to the auris interna.

[00215] Vertigo onset in patients with postural vertigo usually develops when a person lies on one ear or tilts the head back to look up. Vertigo is accompanied by nystagmus. In severe cases of postural vertigo, the vestibular nerve is severed to the affected semicircular canal. Treatment of vertigo is often identical to Meniere's disease, and may include meclizine, lorazepam, prochlorperazine or scopolamine. Fluids and electrolytes may also be intravenously administered if the vomiting is severe.

Presbycusis (Age Related Hearing Loss)

[00216] Presbycusis (or presbycusis or age related hearing loss (ARHL)) is the progressive bilateral loss of hearing that results from aging. Most hearing loss occurs at higher frequencies (i.e.

frequencies above 15 or 16 Hz) making it difficult to hear a female voice (as opposed to male voice), and an inability to differentiate between high-pitched sounds (such as "s" and "th"). It is difficult to filter out background noise. The disorder is most often treated by the implantation of a hearing aid and/or the administration of pharmaceutical agents which prevent the build up of ROS.

[00217] The disorder is caused by changes in the physiology of the inner ear, the middle ear, and/or the VII nerve. Changes in the inner ear resulting in presbycusis include epithelial atrophy with loss of otic hair cells and/or stereocilia, atrophy of nerve cells, atrophy of the stria vascularis, and the thickening/stiffening of the basilar membrane. Additional changes which can contribute to presbycusis include the accumulation of defects in the tympanic membrane and the ossicles.

[00218] Changes leading to presbycusis can occur due to the accumulation of mutations in DNA, and mutations in mitochondrial DNA; however, the changes are exacerbated by exposure to loud noise, exposure to ototoxic agents, infections, and/or the lessening of blood flow to the ear. The latter is attributable to atherosclerosis, diabetes, hypertension, and smoking.

[00219] Presbycusis, or age-related hearing loss, occurs as a part of normal aging, and occurs as a result of degeneration of the receptor cells in the spiral Organ of Corti in the auris interna. Other causes may also be attributed to a decrease in a number of nerve fibers in the vestibulocochlear nerve, as well as a loss of flexibility of the basilar membrane in the cochlea. There is currently no

known cure for permanent hearing damage as a result of presbycusis or excessive noise, although treatment regimens have been proposed, including treatment with antioxidants such as alpha lipoic acid. (Seidman et al, *Am. J. Otol.* (2000) 21:161-167).

Ramsay Hunt's Syndrome (Herpes Zoster Infection)

5 [00220] Ramsay Hunt's syndrome is caused by a herpes zoster infection of the auditory nerve. The infection may cause severe ear pain, hearing loss, vertigo, as well as blisters on the outer ear, in the ear canal, as well as on the skin of the face or neck supplied by the nerves. Facial muscles may also become paralyzed if the facial nerves are compressed by the swelling. Hearing loss are temporary or permanent, with vertigo symptoms usually lasting from several days to weeks.

10 [00221] Treatment of Ramsay Hunt's syndrome includes administration of antiviral agents, including acyclovir. Other antiviral agents include famciclovir and valacyclovir. Combination of antiviral and corticosteroid therapy may also be employed to ameliorate herpes zoster infection. Analgesics or narcotics may also be administered to relieve the pain, and diazepam or other central nervous system agents to suppress vertigo. Capsaicin, lidocaine patches and nerve blocks may also
15 be used. Surgery may also be performed on compressed facial nerves to relieve facial paralysis.

Recurrent Vestibulopathy

[00222] Recurrent vestibulopathy is a condition wherein the subject experiences multiple episodes of severe vertigo. The episodes of vertigo may last for minutes or hours. Unlike Meniere's Disease, it is not accompanied by hearing loss. In some cases it may develop into Meniere's Disease or Benign
20 Paroxysmal Positional Vertigo. Treatment is similar to that of Meniere's Disease.

Syphilis Infection

[00223] Syphilis infection may also lead to congenital prenatal hearing loss, affecting approximately 11.2 per 100,000 live births in the United States, as well as sudden hearing loss in adults. Syphilis is a venereal disease, caused by the spirochete *Treponema pallidum*, which in its
25 secondary and tertiary stages may result in auditory and vestibular disorders due to membranous labyrinthitis, and secondarily include meningitis.

[00224] Both acquired and congenital syphilis can cause otic disorders. Symptoms of cochleovestibular disorders resulting from syphilis are often similar to those of other otic disorders, such as AIED and Meniere's disease, and include tinnitus, deafness, vertigo, malaise, sore throat,
30 headaches, and skin rashes. Syphilis infection may lead to congenital prenatal hearing loss, affecting approximately 11.2 per 100,000 live births in the United States, as well as sudden hearing loss in adults.

[00225] Treatment with steroids and antibiotics, including penicillins (e.g. benzathine penicillin G (BICILLIN LA[®]), are effective in eradicating the spirochete organism. However, *Treponemas* may
35 remain in the cochlear and vestibular endolymph even after eradication from other sites in the body.

Accordingly, long term treatment with penicillins are warranted to achieve complete eradication of the spirochete organism from the endolymph fluid.

[00226] Treatment of otosyphilis (syphilis presenting otic symptoms) typically includes a combination of steroids (e.g., prednisilone) and antibacterial agents (e.g., benzathine penicillin G (BICILLIN LA®), penicillin G procaine, doxycycline, tetracycline, ceftriaxone, azithromycin). Such treatments are effective in eradicating the spirochete organism. However, *Treponemas* may remain in the cochlear and vestibular endolymph even after eradication from other sites in the body.

Accordingly, long term treatment with penicillins are required to achieve complete eradication of the spirochete organism from the endolymph fluid. Also, in the case of a severe or advanced case of syphilis, a uricosuric drug, such as probenecid, are administered in conjunction with the antibacterial agent to increase its efficacy.

Temporal Bone Fractures

[00227] The temporal bone, which contains part of the ear canal, the middle ear and the auris interna, is subject to fractures from blows to the skull or other injuries. Bleeding from the ear or patchy bruising is symptomatic of a fracture to the temporal bone, and may a computed tomography (CT) scan for accurate diagnosis. Temporal bone fractures may rupture the eardrum, causing facial paralysis and sensorineural hearing loss.

[00228] Treatment of detected temporal bone fractures includes an antibiotic regimen to prevent meningitis, or an infection of brain tissue. In addition, surgery are performed to relieve any subsequent pressure on the facial nerve due to swelling or infection.

Temporomandibular Joint Disease

[00229] Some evidence exists for a relationship between temporomandibular joint disease (TMD) and auris interna disorders. Anatomical studies demonstrate the possible involvement of the trigeminal nerve, where trigeminal innervation of the vascular system has been shown to control cochlear and vestibular labyrinth function. (Vass et al. *Neuroscience* (1998) 84:559-567).

Additionally, projections of ophthalmic fibers of the trigeminal Gasser ganglion to the cochlea through the basilar and anterior inferior cerebellar arteries can play an important role in the vascular tone in quick vasoconstrictory response to metabolic stresses, e.g. intense noise. Auris interna diseases and symptoms, such as sudden hearing loss, vertigo and tinnitus, may originate from reduction of the cochlear blood flow due to the presence of abnormal activity in the trigeminal ganglion, for example from migraine or by the central excitatory effect originated in chronic or deep pain produced by TMD.

[00230] Similarly, other researchers have found that the trigeminal ganglion also innervates the ventral cochlear nucleus and the superior olivary complex, which may interfere with auditory pathways leading to the auditory cortex where constant peripheral somatic signals from the ophthalmic and mandibular trigeminal peripheral innervation occurs in TMD cases. (Shore et al. *J.*

Comp. Neurology (2000) 10:271-285). These somatosensory and auditory system interactions via the central nervous system may explain otic symptoms in the absence of existing disease in the ear, nose or throat.

[00231] Accordingly, forceful muscle contractions in TMD may elicit modulations in the neurological and auditory and equilibrium function. For example, the auditory and vestibular modulations may occur as a result of hypertonicity and muscular spasm, which in turn irritates nerves and blood vessels that affect auris interna function by muscular trapping. Relief of the affected nerve or muscular contractions may act to relieve auditory or vestibular symptoms. Medications, including barbiturates or diazepam, may thus relieve auditory or vestibular dysfunction in TMD patients.

Utricular Dysfunction

[00232] The utricle is one of the two otoliths found in the vestibular labyrinth. It is responsive to both gravity and linear acceleration along the horizontal plane. Utricular dysfunction is a disorder caused by damage to the utricle. It is often characterized by a subject's perception of tilting or imbalance.

Vertigo

[00233] Vertigo is described as a feeling of spinning or swaying while the body is stationary. There are two types of vertigo. Subjective vertigo is the false sensation of movement of the body. Objective vertigo is the perception that one's surrounding are in motion. It is often accompanied by nausea, vomiting, and difficulty maintaining balance.

[00234] While not wishing to be bound by any one theory, it is hypothesized that vertigo is caused by an over-accumulation of fluid in the endolymph. This fluid imbalance results in increased pressure on the cells of the inner ear which leads to the sensation of movement. The most common cause of vertigo is benign paroxysmal positional vertigo, or BPPV. It can also be brought on by a head injury, or a sudden change of blood pressure. It is a diagnostic symptom of several diseases including superior canal dehiscence syndrome.

Vestibular Neuritis

[00235] Vestibular neuronitis, or vestibular neuropathy, is an acute, sustained dysfunction of the peripheral vestibular system. It is theorized that vestibular neuronitis is caused by a disruption of afferent neuronal input from one or both of the vestibular apparatuses. Sources of this disruption include viral infection, and acute localized ischemia of the vestibular nerve and/or labyrinth. Vestibular neuronitis is characterized by sudden vertigo attacks, which may present as a single attack of vertigo, a series of attacks, or a persistent condition which diminishes over a matter of weeks. Symptoms typically include nausea, vomiting, and previous upper respiratory tract infections, although there are generally no auditory symptoms. The first attack of vertigo is usually

severe, leading to nystagmus, a condition characterized by flickering of the eyes involuntarily toward the affected side. Hearing loss does not usually occur.

[00236] In some instances, vestibular neuronitis is caused by inflammation of the vestibular nerve, the nerve that connects the inner ear to the brain, and is likely caused by viral infection. Diagnosis of vestibular neuronitis usually involves tests for nystagmus using electronystamography, a method of electronically recording eye movements. Magnetic resonance imaging may also be performed to determine if other causes may play a role in the vertigo symptoms.

[00237] Treatment of vestibular neuronitis typically involves alleviating the symptoms of the condition, primarily vertigo, until the condition clears on its own. Treatment of vertigo is often identical to Meniere's disease, and may include meclizine, lorazepam, prochlorperazine, or scopolamine. Fluids and electrolytes may also be intravenously administered if the vomiting is severe. Corticosteroids, such as prednisilone, are also given if the condition is detected early enough.

[00238] Compositions disclosed herein comprising an antiviral agent can be administered for the treatment of vestibular neuronitis. Further, the compositions are administered with other agents that are typically used to treat symptoms of the condition, including anticholinergics, antihistamines, benzodiazepines, or steroids. Treatment of vertigo is identical to Meniere's disease, and may include meclizine, lorazepam, prochlorperazine or scopolamine. Fluids and electrolytes may also be intravenously administered if the vomiting is severe.

[00239] The most significant finding when diagnosing vestibular neuronitis is spontaneous, unidirectional, horizontal nystagmus. It is often accompanied by nausea, vomiting, and vertigo. It is, generally, not accompanied by hearing loss or other auditory symptoms.

[00240] There are several treatments for vestibular neuronitis. H1-receptor antagonists, such as dimenhydrinate, diphenhydramine, meclizine, and promethazine, diminish vestibular stimulation and depress labyrinthine function through anticholinergic effects. Benzodiazepines, such as diazepam and lorazepam, are also used to inhibit vestibular responses due to their effects on the GABA_A receptor. Anticholinergics, for example scopolamine, are also prescribed. They function by suppressing conduction in the vestibular cerebellar pathways. Finally, corticosteroids (*i.e.* prednisone) are prescribed to ameliorate the inflammation of the vestibular nerve and associated apparatus.

Advantages of local otic administration

[00241] To overcome the toxic and attendant side effects of systemic delivery, disclosed herein are methods and compositions for local delivery of therapeutic agents to auris media and/or auris interna structures. Access to, for example, the vestibular and cochlear apparatus will occur through the auris media including the round window membrane, the oval window/stapes footplate, the annular ligament and through the otic capsule/temporal bone.

[00242] In addition, localized treatment of the auris media and/or auris interna also affords the use of previously undesired therapeutic agents, including agents with poor PK profiles, poor uptake, low systemic release and/or toxicity issues. Because of the localized targeting of the otic agent formulations and compositions, as well as the biological blood barrier present in the auris interna, the risk of adverse effects will be reduced as a result of treatment with previously characterized toxic or ineffective otic active agents, (e.g., immunomodulatory agents such as anti-TNF agents). Accordingly, also contemplated within the scope of the embodiments described herein is the use of active agents and/or agents that have been previously rejected by practitioners because of adverse effects or ineffectiveness of the otic agent.

[00243] By specifically targeting the auris media or auris interna structures, adverse side effects as a result of systemic treatment are avoided. Moreover, by providing a controlled release otic agent formulation (e.g., immunomodulating agent or auris pressure modulator formulation) or composition to treat otic disorders, a constant, variable and/or extended source of an otic agent is provided to the individual or patient suffering from an otic disorder, reducing or eliminating the variability of treatment. Accordingly, one embodiment disclosed herein is to provide a formulation that enables at least one therapeutic agent to be released in therapeutically effective doses either at variable or constant rates such as to ensure a continuous release of the at least one agent. In some embodiments, the auris active agents disclosed herein are administered as an immediate release formulation or composition. In other embodiments, the auris active agents are administered as a controlled release formulation, released either continuously or in a pulsatile manner, or variants of both. In still other embodiments, the active agent formulation is administered as both an immediate release and controlled release formulation, released either continuously or in a pulsatile manner, or variants of both. The release is optionally dependent on environmental or physiological conditions, for example, the external ionic environment (see, e.g. Oros[®] release system, Johnson & Johnson).

[00244] Also included within the embodiments disclosed herein is the use of additional auris media and/or auris interna agents in combination with the otic agent formulations and compositions disclosed herein. When used, such agents assist in the treatment of hearing or equilibrium loss or dysfunction as a result of an autoimmune disorder, including vertigo, tinnitus, hearing loss, balance disorders, infections, inflammatory response or combinations thereof. Accordingly, agents that ameliorate or reduce the effects of vertigo, tinnitus, hearing loss, balance disorders, infections, inflammatory response or combinations thereof are also contemplated to be used in combination with the otic agents described herein including steroids, anti-emetic agents, local anesthetic agents, corticosteroids, chemotherapeutic agents, including cytoxan, azathiaprime or methotrexate; treatment with collagen, gamma globulin, interferons, copaxone, central nervous system agents, antibiotics, platelet-activating factor antagonists, nitric oxide synthase inhibitors and combinations thereof.

[00245] In addition, the auris-compatible pharmaceutical compositions or formulations included herein also include carriers, adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure, and/or buffers. Such carriers, adjuvants, and other excipients will be compatible with the environment in the auris media and/or auris interna. Accordingly, specifically contemplated are carriers, adjuvants and excipients that lack ototoxicity or are minimally ototoxic in order to allow effective treatment of the otic disorders contemplated herein with minimal side effects in the targeted regions or areas. To prevent ototoxicity, otic pharmaceutical compositions or formulations disclosed herein are optionally targeted to distinct regions of the auris media and/or auris interna, including but not limited to the tympanic cavity, vestibular bony and membranous labyrinths, cochlear bony and membranous labyrinths and other anatomical or physiological structures located within the auris interna.

Treatment

[00246] Provided herein are otic compositions suitable for the treatment of any otic condition, disease or disorder (e.g., middle and/or inner ear disorder) described herein, comprising administration of an auris formulation described herein to an individual or patient in need thereof. The formulations described herein are suitable for the treatment of any disease described herein. In some instances, the treatment is long-term treatment for chronic recurring disease. In some instances, the treatment is prophylactic administration of an otic formulation described herein for the treatment of any otic disease or disorder described herein. In some instances, prophylactic administration avoids occurrence of disease in individuals suspected of having a disease or in individuals genetically predisposed to an otic disease or disorder. In some instances the treatment is preventive maintenance therapy. In some instances, preventive maintenance therapy avoids recurrence of a disease.

[00247] In some instances, because of their otic compatibility and improved sterility, the formulations described herein are safe for long-term administration. The auris compositions described herein have very low ototoxicity and provide a steady sustained release of a therapeutic agent for a period of at least one week, two weeks, three weeks or a month.

[00248] Provided herein are controlled release compositions and formulations to treat and/or prevent diseases associated with the ear, including the cochlea, the middle ear and inner ear, including autoimmune inner ear disorder (AIED), Ménière's disease (endolymphatic hydrops), noise induced hearing loss (NIHL), sensorineural hearing loss (SNL), tinnitus, otosclerosis, balance disorders, vertigo and the like.

[00249] The etiology of several ear diseases or disorders consists of a syndrome of progressive hearing loss, including noise induced hearing loss and age-related hearing loss, dizziness, nausea, nystagmus, vertigo, tinnitus, inflammation, swelling, infection and/or congestion. These disorders

may have many causes, such as infection, exposure to noise, injury, inflammation, tumors and/or adverse response to drugs or other chemical agents. Several causes of hearing and/or equilibrium impairment are attributed to inflammation and/or an autoimmune disorder and/or a cytokine-mediated inflammatory response.

5 [00250] Provided herein are controlled release immunomodulator compositions and formulations to treat inflammation or infection of the auris media, including otitis media. A few therapeutic products are available for the treatment of AIED, including anti-TNF agents; however, systemic routes via oral, intravenous or intramuscular routes are currently used to deliver these therapeutic agents.

[00251] Provided herein are controlled release aural pressure modulating compositions and
10 formulations to treat fluid homeostasis disorders of the inner ear, including Meniere's Disease, endolymphatic hydrops, progressive hearing loss, including noise induced hearing loss and age-related hearing loss, dizziness, vertigo, tinnitus and similar conditions.

[00252] In some embodiments, the compositions provided herein are CNS modulating compositions and formulations to treat tinnitus, progressive hearing loss, including noise induced hearing loss and
15 age-related hearing loss, and balance disorders. Balance disorders include benign paroxysmal positions vertigo, dizziness, endolymphatic hydrops, kinetosis, labyrinthitis, Mal de débarquement, Meniere's Disease, Meniere's Syndrome, myringitis, otitis media, Ramsay Hunt's Syndrome, recurrent vestibulopathy, tinnitus, vertigo, microvascular compression syndrome, utricular dysfunction, and vestibular neuronitis. A few therapeutic products are available for the treatment of
20 balance disorders, including GABA_A receptor modulators and local anesthetics.

[00253] In some embodiments, the compositions provided herein are cytotoxic agent compositions and formulations for the treatment of autoimmune diseases of the ear, including autoimmune inner ear disease (AIED). Also provided herein are controlled release cytotoxic agent compositions for the treatment of disorders of the auris media, including otitis media. The compositions disclosed herein
25 are also useful for the treatment of cancer, particularly cancer of the ear. A few therapeutic products are available for the treatment of AIED, including certain cytotoxic agents. Particularly, the cytotoxic agents methotrexate and cyclophosphamide have been tested and are used for systemic treatment of AIED. Also, thalidomide, while not currently administered for the treatment of AIED, has been used to treat Behçet's disease, which is often associated with AIED.

30 [00254] In some embodiments, the compositions provided herein comprise auris sensory cell modulators for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Further disclosed herein are controlled release auris sensory cell modulating agent compositions and formulations to treat ototoxicity, excitotoxicity, sensorineural hearing loss, Meniere's Disease/Syndrome, endolymphatic
35 hydrops, labyrinthitis, Ramsay Hunt's Syndrome, vestibular neuronitis and microvascular compression syndrome.

[00255] In some embodiments, the compositions provided herein are antimicrobial agent compositions and formulations for the treatment of otic disorders, including otitis externa, otitis media, Ramsay Hunt syndrome, otosyphilis, AIED, Meniere's disease, and vestibular neuronitis.

[00256] In some embodiments, the compositions provided herein prevent, relieve, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals and/or the dysfunction of the mitochondria.

[00257] Also provided herein are controlled release ion channel modulating compositions and formulations to treat fluid homeostasis disorders of the inner ear, including Meniere's Disease, endolymphatic hydrops, progressive hearing loss, including noise induced hearing loss and age-related hearing loss, dizziness, vertigo, tinnitus and similar conditions. Systemic routes via oral, intravenous or intramuscular routes are currently used to deliver ion channel modulating therapeutic agents.

Therapeutic Agents

[00258] Notwithstanding any therapeutic agent used in the formulations described herein, the otic compositions described herein will have pH and osmolarity that is auris-acceptable. Any otic composition described herein meets the stringent sterility requirements described herein and will be compatible with the endolymph and/or the perilymph. Pharmaceutical agents that are used in conjunction with the formulations disclosed herein include agents that ameliorate or lessen otic disorders, including auris interna disorders, and their attendant symptoms, which include but are not limited to hearing loss, nystagmus, vertigo, tinnitus, inflammation, swelling, infection and congestion. Otic disorders may have many causes, such as infection, injury, inflammation, tumors and adverse response to drugs or other chemical agents that are responsive to the pharmaceutical agents disclosed herein. A skilled practitioner would be familiar with agents that are useful in the amelioration or eradication of otic disorder; accordingly, agents which are not disclosed herein but are useful for the amelioration or eradication of otic disorders are expressly included and intended within the scope of the embodiments presented. In some embodiments, pharmaceutically active metabolites, salts, polymorphs, prodrugs, analogues, and derivatives of the otic agents disclosed herein that retain the ability of the parent antimicrobial agents to treat otic disorders are useful in the formulations.

[00259] Active ingredients or otic therapeutic agents include, but are not limited to, anti-inflammatory agents, anti-anti-oxidants, neuroprotective agents, glutamate modulators, TNF-alpha modulators, interleukin 1 beta modulators, retinaldehyde modulators, notch modulators, gamma-secretase modulators, thalidomide, ion and/or fluid (e.g., water) homeostasis modulators, vasopressin inhibitors, inhibitors of the vasopressin-mediated AQP2 (aquaporin 2) system, transcriptional regulators of the inner-ear transcriptional regulatory network (including, e.g., transcriptional regulators of estrogen-related receptor beta), inner ear hair cell growth factors,

including BDNF (brain derived and NF-3, and other therapeutic modalities. Agents explicitly include an agonist of an otic target, a partial agonist of an otic target, an antagonist of an otic target, a partial antagonist of an otic target, an inverse agonist of an otic target, a competitive antagonist of an otic target, a neutral antagonist of an otic target, an orthosteric antagonist of an otic target, an allosteric antagonist of an otic target a positive allosteric modulator of an otic target, or combinations thereof.

[00260] In addition, because the formulation is designed such that the active ingredient has limited or no systemic release, agents that produce systemic toxicities (e.g., liver toxicity) or have poor PK characteristics (e.g. short half-life) are also optionally used. Thus, pharmaceutical agents which have been previously shown to be toxic, harmful or non-effective during systemic application, for example through toxic metabolites formed after hepatic processing, toxicity of the drug in particular organs, tissues or systems, through high levels needed to achieve efficacy, through the inability to be released through systemic pathways or through poor PK characteristics, are useful in some embodiments herein. The formulations disclosed herein are contemplated to be targeted directly to otic structures where treatment is needed; for example, one embodiment contemplated is the direct application of the aural pressure modulating formulations disclosed herein onto the round window membrane or the crista fenestrae cochlea of the auris interna, allowing direct access and treatment of the auris interna, or inner ear components. In other embodiments, the aural pressure modulating formulation disclosed herein is applied directly to the oval window. In yet other embodiments, direct access is obtained through microinjection directly into the auris interna, for example, through cochlear microperfusion. Such embodiments also optionally comprise a drug delivery device, wherein the drug delivery device delivers the aural pressure modulating formulations through use of a needle and syringe, a pump, a microinjection device, an in situ forming spongy material or any combination thereof.

[00261] In still other embodiments, application of any otic agent formulation described herein is targeted to the auris media through piercing of the intratympanic membrane and applying the otic agent formulation directly to the auris media structures affected, including the walls of the tympanic cavity or auditory ossicles. By doing so, the auris active agent formulations disclosed herein are confined to the targeted auris media structure, and will not be lost, for example, through diffusion or leakage through the eustachian tube or pierced tympanic membrane. In some embodiments, the auris-compatible formulations disclosed herein are delivered to the auris externa in any suitable manner, including by cotton swab, injection or ear drops. Also, in other embodiments, the otic formulations described herein are targeted to specific regions of the auris externa by application with a needle and syringe, a pump, a microinjection device, an in situ forming spongy material or any combination thereof. For example, in the case of treatment of otitis externa, antimicrobial agent

formulations disclosed herein are delivered directly to the ear canal, where they are retained, thereby reducing loss of the active agents from the target ear structure by drainage or leakage.

[00262] In some embodiments, agents which may have been previously rejected as, for example, an antimicrobial agent, may find use herein because of the targeted nature of the embodiments which bypass systemic effects, including toxicity and harmful side effects. By way of example only, oncept, a previously rejected anti-TNF agent due to toxicity and safety issues, is useful as an anti-TNF agent in some of the embodiments disclosed herein. Also contemplated within the scope of embodiments described herein is the administration of higher doses of pharmaceutical agents, for example agents that have dose limiting toxicities, compared to currently approved doses for such pharmaceutical agents

[00263] Some pharmaceutical agents, either alone or in combination, are ototoxic. For example, some chemotherapeutic agents, including actinomycin, bleomycin, cisplatin, carboplatin and vincristine; and antibiotics, including erythromycin, gentamicin, streptomycin, dihydrostreptomycin, tobramycin, netilmicin, amikacin, neomycin, kanamycin, etiomycin, vancomycin, metronidazole, capreomycin, are mildly to very toxic, and may affect the vestibular and cochlear structures differentially. However, in some embodiments, the combination of an ototoxic drug, for example cisplatin, in combination with an antioxidant is protective and lessen the ototoxic effects of the drug. Moreover, the localized application of the potentially ototoxic drug lessens the toxic effects that might otherwise occur through systemic application through the use of lower amounts with maintained efficacy, or the use of targeted amounts for a shorter period of time. Accordingly, a skilled practitioner choosing a course of therapy for targeted otic disorder will have the knowledge to avoid or combine an ototoxic compound, or to vary the amount or course of treatment to avoid or lessen ototoxic effects.

[00264] In certain instances, pharmaceutical excipients, diluents or carriers are potentially ototoxic. For example, benzalkonium chloride, a common preservative, is ototoxic and therefore potentially harmful if introduced into the vestibular or cochlear structures. In formulating a controlled release otic formulation, it is advised to avoid or combine the appropriate excipients, diluents or carriers to lessen or eliminate potential ototoxic components from the formulation, or to decrease the amount of such excipients, diluents or carriers. In some instances, the ototoxicity of the pharmaceutical agents, excipients, diluents, carriers, or formulations and compositions disclosed herein can be ascertained using an accepted animal model. See, e.g., Maritini, A., et al. *Ann. N.Y. Acad. Sci.* (1999) 884:85-98. Optionally, a controlled release otic formulation includes otoprotective agents, such as antioxidants, alpha lipoic acid, calicum, fosfomycin or iron chelators, to counteract potential ototoxic effects that may arise from the use of specific therapeutic agents or excipients, diluents or carriers.

[00265] Other agents that are used in the embodiments disclosed herein, either alone or in combination with other auris interna agents, include anti-apoptotic agents, including caspases, JNK

the insulin-like growth factor (IGF) receptor, the epidermal growth factor (EGF) receptor, and/or the platelet-derived growth factor. In some embodiments, the growth factor is hepatocyte growth factor.

[00370] In some embodiments, the growth factor is an epidermal growth factor (EGF). In some embodiments, the EGF is heregulin (HRG). In certain instances, HRG stimulates the proliferation of utricular sensory epithelium. In certain instances, HRG-binding receptors are found in the vestibular and auditory sensory epithelium.

[00371] In some embodiments, the growth factor is an insulin-like growth factor (IGF). In some embodiments, the IGF is IGF-1. In some embodiments, the IGF-1 is mecasermin. In certain instances, IGF-1 attenuates the damage induced by exposure to an aminoglycoside. In certain instances, IGF-1 stimulates the differentiation and/or maturation of cochlear ganglion cells.

[00372] In some embodiments, the FGF receptor agonist is FGF-2. In some embodiments, the IGF receptor agonist is IGF-1. Both the FGF and IGF receptors are found in the cells comprising the utricle epithelium.

[00373] In some embodiments, the growth factor is hepatocyte growth factor (HGF). In some instances, HGF protects cochlear hair cells from noise-induced damage and reduces noise-exposure-caused ABR threshold shifts.

[00374] Also contemplated for use in the otic formulations described herein are growth factors including Erythropoietin (EPO), Granulocyte-colony stimulating factor (G-CSF), Granulocyte-macrophage colony stimulating factor (GM-CSF), Growth differentiation factor-9 (GDF9), Insulin-like growth factor (IGF), Myostatin (GDF-8), Platelet-derived growth factor (PDGF), Thrombopoietin (TPO), Transforming growth factor alpha (TGF- α), Transforming growth factor beta (TGF- β), Vascular endothelial growth factor (VEGF) or combinations thereof.

Neurotrophs

[00375] In some embodiments, the growth factor is a neurotroph. In certain instances, neurotrophs are growth factors which prevent cells from initiating apoptosis, repair damaged neurons and otic hair cells, and/or induce differentiation in progenitor cells. In some embodiments, the neurotroph is brain-derived neurotrophic factor (BDNF), ciliary neurotrophic factor (CNTF), glial cell-line derived neurotrophic factor (GDNF), neurotrophin-3, neurotrophin-4, and/or combinations thereof.

[00376] In some embodiments, the neurotroph is BDNF. In certain instances, BDNF is a neurotroph which promotes the survival of existing neurons (e.g. spiral ganglion neurons), and otic hair cells by repairing damaged cells, inhibiting the production of ROS, and inhibiting the induction of apoptosis. In certain embodiments, it also promotes the differentiation of neural and otic hair cell progenitors. Further, in certain embodiments, it protects the Cranial Nerve VII from degeneration. In some embodiments, BDNF is administered in conjunction with fibroblast growth factor.

[00377] In some embodiments, the neurotroph is neurotrophin-3. In certain embodiments, neurotrophin-3 promotes the survival of existing neurons and otic hair cells, and promotes the

differentiation of neural and otic hair cell progenitors. Further, in certain embodiments, it protects the VII nerve from degeneration.

[00378] In some embodiments, the neurotroph is CNTF. In certain embodiments, CNTF promotes the synthesis of neurotransmitters and the growth of neuritis. In some embodiments, CNTF is administered in conjunction with BDNF.

[00379] In some embodiments, the neurotroph is GDNF. In certain embodiments, GDNF expression is increased by treatment with ototoxic agents. Further, in certain embodiments, cells treated with exogenous GDNF have higher survival rates after trauma than untreated cells.

Immune System Cells

[00380] Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Accordingly, some embodiments incorporate the use of cells which participate in the repair of otic hair cells and neurons. In some embodiments, the cells which participate in the repair of otic hair cells and neurons are macrophages, microglia, and/or microglia-like cells. In certain instances, the concentration of macrophages and microglia increase in ears damaged by treatment with ototoxic agents. In certain instances, microglia-like cells eliminate waste from the Organ of Corti and participate in the structural repair of hair cells following treatment with the ototoxic antibiotic neomycin.

Ototoxic Agents

[00381] Contemplated for use with the formulations disclosed herein are agents that destroy neurons and/or otic hair cells. Accordingly, some embodiments incorporate the use of agents which fatally damage and/or induce apoptosis in the neurons and/or otic hair cells of the auris. In some embodiments, the agents which fatally damage and/or induce apoptosis in the neurons and/or otic hair cells of the auris are the aminoglycoside antibiotics (e.g. gentamicin, and amikacin), the macrolide antibiotics (e.g. erythromycin), the glycopeptide antibiotics (e.g. vancomycin), the loop diuretics (e.g. furosemide) salicylic acid, and nicotine.

Retinoblastoma Protein Modulation

[00382] Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and/or hair cells of the auris, promote the growth of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Further contemplated herein are agents that destroy neurons and/or otic hair cells. Accordingly, some embodiments incorporate the use of agents that modulate retinoblastoma protein (pRB). pRB is a member of the pocket protein family. It is encoded by the RB1 gene. In certain instances, it inhibits transition from G1 to S phase by binding to and inactivating the E2f family of transcription factors.

In certain instances, it also regulates differentiation, and survival of hair cells. In certain instances, pRB knock-out mice demonstrate increased proliferation of hair cells.

[00383] In some embodiments, the agent that modulates one or more of the pRB is an agonist of pRB. In some embodiments, the agent that modulates one or more of the pRB is an antagonist of pRB. In certain instances, a compound which agonizes or antagonizes pRB is identified (e.g. by use of a high throughput screen). In some embodiments, a construct is designed such that a reporter gene is placed downstream of an E2F binding sequence. In some embodiments, the binding sequence is TTTCGCGC. In some embodiments, the reporter gene is luciferase, CAT, GFP, β -lactamase or β -galactosidase. In certain instances, E2f binds to the binding sequence causing the transcription and expression of the reporter gene. In certain instances, an agonist of pRB causes an increase in the binding of pRB to E2f. In certain instances, the increase in binding of pRB and E2f results in a decrease in the transcription and expression of the reporter gene. In certain instances, an antagonist of pRB causes a decrease in the binding of pRB to E2f. In certain instances, the decrease in binding of pRB and E2f results in an increase in the transcription and expression of the reporter gene.

[00384] In some embodiments, the agent that modulates pRB is an siRNA molecule. In certain instances, the siRNA molecule is as described herein.

Salicylic acid

[00385] Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Accordingly, some embodiments incorporate the use of salicylic acid. In certain instances, when administered before treatment with an aminoglycoside, it protects otic hair cells and spiral ganglion neurons from aminoglycoside ototoxicity.

Sodium Channel Blockers

[00386] Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and hair cells, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. In certain instances, excitotoxicity causes the excessive opening of Na^+ channels. In certain instances, this results in excess Na^+ ions entering the neuron. In certain instances, the excess influx of Na^+ ions into the neuron causes the neuron to fire more often. In certain instances, this increased firing yields a rapid buildup of free radicals and inflammatory compounds. In certain instances, the free radicals damage the mitochondria, depleting the cell's energy stores. Further, in certain instances, excess levels of Na^+ ions activate excess levels of enzymes including, but not limited to, phospholipases, endonucleases, and proteases. In certain instances, the over-activation of these enzymes results in damage to the cytoskeleton, plasma membrane, mitochondria, and DNA of

the neuron. Accordingly, some embodiments incorporate the use of agents which antagonize the opening of Na⁺ channels. In some embodiments, sodium channel blockers are as described herein.

Stem Cells and Differentiated Auris Sensory Cells

[00387] Contemplated for use with the formulations disclosed herein are transplants of cells that

5 supplement and/or replace the pre-existing neurons and/or hair cells of the auris. In some embodiments, the agent is a stem cell. In some embodiments, the agent is a partially or fully differentiated auris sensory cell. In some embodiments, the differentiated auris sensory cell is derived from a human donor. In some embodiments, the differentiated auris sensory cell is derived from a stem cell, the differentiation of which was induced under artificial (e.g. laboratory)
10 conditions.

[00388] Stem cells are cells that possess the capacity to differentiate into multiple cell types.

Totipotent stem cells can differentiate into embryonic cells or extraembryonic cells. Pluripotent cells can differentiate into cells of any of endoderm, mesoderm, or ectoderm origin. Multipotent cells can differentiate into closely related cells (e.g. hematopoietic stem cells). Unipotent cells can
15 differentiate into only one type of cell, but like other stem cells have the characteristic of self-renewal. In some embodiments, the stem cell is totipotent, pluripotent, multipotent, or unipotent. Further, stem cells can undergo mitotic division without themselves differentiating (i.e. self-renewal).

[00389] Embryonic stem (ES) cells are stem cells derived from the epiblast tissue of the inner cell

20 mass of a blastocyst or earlier stage embryo. ES cells are pluripotent. In some embodiments, the stem cell is an ES cell. Adult stem cells (also known as somatic cells or germline cells) are cells isolated from a developed organism wherein the cells possess the characteristic of self-renewal, and the ability to differentiate into multiple cell types. Adult stem cells are pluripotent (for example, stem cells found in umbilical cord blood), multipotent or unipotent. In some embodiments, the stem
25 cell is an adult stem cell.

[00390] In some embodiments, a stem cell and/or a differentiated auris sensory cell is administered in combination with a differentiation stimulating agent. In some embodiments, the differentiation stimulating agent is a growth factor. In some embodiments, the growth factor is a neurotrophin (e.g. nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3),
30 neurotrophin-4 (NT-4), or novel neurotrophin-1 (NNT1). In some embodiments, the growth factor is FGF, EGF, IGF, PGF, or combinations thereof.

[00391] In some embodiments, a stem cell and/or a differentiated auris sensory cell is administered to a subject in need thereof as a controlled release agent. In some embodiments, a stem cell and/or a differentiated auris sensory cell is administered to a subject in need thereof as an immediate release
35 agent (e.g. in a cell suspension) in combination with a controlled release auris sensory cell modulating agent. In some embodiments, a controlled release auris sensory cell modulating agent is

a vector comprising an Atoh1 or BRN3 gene, an siRNA sequence targeting RB1, a growth factor, or combinations thereof.

[00392] In some embodiments, a stem cell and/or a differentiated auris sensory cell is administered to the cochlea or vestibular labyrinth. In some embodiments, a stem cell and/or a differentiated auris sensory cell is administered by via intratympanic injection, and/or a post-auricular incision. In some embodiments, a stem cell and/or a differentiated auris sensory cell is contacted with the Organ of Corti, vestibulocochlear nerve, and/or crista ampullaris.

Thyroid Hormone Receptor Modulation

[00393] Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and/or hair cells of the auris, promote the growth of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Accordingly, some embodiments incorporate the use of agents that modulate Thyroid Hormone (TH) receptors. The TH receptors are a family of nuclear hormone receptors. The family includes, but is not limited to TR α 1 and TR β . In certain instances, TR β knock-out mice demonstrate a decreased responsiveness to auditory stimuli, and a decrease in K⁺ current in hair cells.

[00394] In some embodiments, the agent that modulates one or more of the TH receptors is an agonist of the one or more TH receptors. In some embodiments, the agonist of one or more of the TH receptors is T₃ (3,5,3'-triiodo-L-thyronine); KB-141 (3,5-dichloro-4-(4-hydroxy-3-isopropylphenoxy)phenylacetic acid); GC-1 (3,5-dimethyl-4-(4'-hydroxy-3'-isopropylbenzyl)phenoxy acetic acid); GC-24 (3,5-dimethyl-4-(4'-hydroxy-3'-benzyl)benzylphenoxyacetic acid); sobetirome (QRX-431); 4-OH-PCB106 (4-OH-2',3,3',4',5'-pentachlorobiphenyl); MB07811 ((2R,4S)-4-(3-chlorophenyl)-2-[(3,5-dimethyl-4-(4'-hydroxy-3'-isopropylbenzyl)phenoxy)methyl]-2-oxido-[1,3,2]-dioxaphosphonane); MB07344 (3,5-dimethyl-4-(4'-hydroxy-3'-isopropylbenzyl)phenoxy)methylphosphonic acid); and combinations thereof. In certain instances, KB-141; GC-1; sobetirome; and GC-24 are selective for TR β .

TRPV Modulation

[00395] Contemplated for use with the formulations disclosed herein are agents that modulate the degeneration of neurons and hair cells, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Accordingly, some embodiments incorporate the use of agents that modulate TRPV receptors. The TRPV (Transient Receptor Potential Channel Vanilloid) receptors are a family of non-selective ion channels permeable to calcium, amongst other ions. There are six members of the family: TRPV1-6. In certain instances, following treatment with kanamycin, TRPV 1 is upregulated. Additionally, in certain instances, antagonism of the TRPV 4 receptor makes mice vulnerable to

acoustic trauma. Further, in certain instances, capsaicin, an agonist of TRPV 1, prevents hyperlocomotion following an ischemic event.

[00396] In some embodiments, the agent that modulates one or more of the TRPV receptors is an agonist of the one or more TRPV receptors. In some embodiments, the agonist of one or more of the TRPV receptors is capsaicin, resiniferatoxin, or combinations thereof. In some embodiments, TRPV modulating include the TRPV modulators disclosed in US application publications 2005/0277643, 2005/0215572, 2006/0194801, 2006/0205773, 2006/0194801, 2008/0175794, 2008/0153857, 2008/0085901, 20080015183, 2006/0030618, 2005/0277646, 2005/0277631, 2005/0272931, 2005/0227986, 2005/0153984, 2006/0270682, 2006/0211741, 2006/0205980, and 2006/0100490, and/or combinations thereof.

Sensory Hair Cell Restorative Agents

[00397] In some instances, immunomodulators and/or aural pressure modulators modulate the function of neurons and/or auris sensory cells. Therapeutic agents which assist in restoring sensory hair cell presence or function are also contemplated herein. These therapeutic agents assist in the treatment of hearing loss in patients, including sensorineural hearing loss, presbycusis and hearing loss from excessive noise. Recent studies have demonstrated the use of insulin-like growth factor 1 (IGF-1) in the restoration of auditory function for noise-induced hearing loss patients. (Lee et al. *Otol. Neurotol.* (2007) 28:976-981). Accordingly, agents IGF-1, IGF-1 agonists or agents which upregulate the expression, production or function of IGF-1 are optionally included with the formulations described herein.

Adenosine Modulators

[00398] Adenosine is comprised of adenine attached to ribofuranose via a β -N9-glycosidic bond. In certain instances, adenosine is an inhibitory neurotransmitter. In certain instances, it functions as a ligand for four GPCRs – adenosine receptor A₁, adenosine receptor A_{2A}, adenosine receptor A_{2B}, and adenosine receptor A₃. In certain instances, the binding of adenosine to an adenosine receptor results in (either partially or fully) an anti-inflammatory effect. In certain instances, the binding of adenosine to an adenosine receptor results in (either partially or fully) vasodilation. In certain instances, it is produced in response to cellular damage (e.g., hypoxia, and ischemia). For example, depolarization and asphyxia in the ear induce the release of adenosine into perilymph where it exerts a protective effect.

[00399] Accordingly, in some embodiment adenosine modulators are used in the treatment of cochlear and vestibular disorders. In some embodiments, the adenosine modulator is ATL313 (4-(3-(6-amino-9-(5-cyclopropylcarbonyl-3,4-dihydroxytetrahydrofuran-2-yl)-9H-purin-2-yl)prop-2-ynyl)piperidine-1-carboxylic acid methyl ester); GW328267X ((2R,3R,4S,5R)-2-{6-amino-2-[(1-benzyl-2-hydroxyethyl)amino]-9H-purin-9-yl}-5-(2-ethyl-2H-tetrazol-5-yl)tetrahydrofuran-3,4-diol); CGS 21680 hydrochloride (4-[2-[[6-Amino-9-(N-ethyl-b-D-ribofuranuronamidosyl)-9H -

purin-2-yl]amino]ethyl]benzenepropanoic acid hydrochloride); CV 1808 (2-Phenylaminoadenosine); p-DITC-APEC (2-[4-[2-[2-[(4-Isothiocyanatophenyl)thiocarbonylamino]ethylaminocarbonyl]ethyl]phenethylamino]-5'-N-ethylcarboxamidadenosine); SDZ WAG994 (N-Cyclohexyl-2'-O-methyladenosine); CVT-3146 (regadenoson; 1-(9-(3,4-dihydroxy-5-(hydroxymethyl)oxolan-2-yl)-6-aminopurin-2-yl)pyrazol-4-yl)-N-methylcarboxamide); ATL-146e (4-{3-[6-Amino-9-(5-ethylcarbamoyl-3,4-dihydroxy-tetrahydro-furan-2-yl)-9H-purin-2-yl]-prop-2-ynyl}-cyclohexanecarboxylic acid methyl ester); 5'-n- Ethyl-carboxamidoadenosine; tecadenoson; CVT-510 (N-(3(R)-tetrahydrofuranyl)-6-aminopurine riboside); CCPA (2-Chloro-N6-cyclopentyladenosine); CPA (N6-Cyclopentyladenosine); GR 79236 (N-[(1S,2S)-2-Hydroxycyclopentyl]adenosine); 2'-MeCCPA; PD 81723 ((2-Amino-4,5-dimethyl-3-thienyl)-[3-(trifluoromethyl)phenyl]methanone); PSB 36 (1-Butyl-8-(hexahydro-2,5-methanopentalen-3a(1H)-yl)-3,7 -dihydro-3-(3-hydroxypropyl)-1H-purine-2,6-dione); ribavirin; CHA (N6-cyclohexyladenosine); GW493838 (GSK); (-)-N6-(2-phenylisopropyl) adenosine; GW684067 ((2R,3R,4S,5R)-5-ethynyl-2-[6-tetrahydro-2H-pyran-4-ylamino)-9H-purin-9-yl]tetrahydrofuran-3,4-diol); CVT-3619 (2-(6-((2-hydroxycyclopentyl)amino)purin-9-yl)-5-((2-fluorophenylthio)methyl)oxolane-3,4-diol); 2-Cl-IB-MECA (CF102; 2-chloro-N⁶-(3-iodobenzyl)-5'-N-methylcarbamoyladenosine); HEMADO; IB-MECA (CF101; N⁶-(3-iodobenzyl)-5'-N-methylcarbamoyladenosine); CP-532903 (N⁶-(2,5-Dichlorobenzyl)-3'- aminoadenosine-5'-N-methylcarboxamide); CF502 (Can-Fite BioPharma); LJ-529 (2-chloro- N(6)- (3- iodobenzyl)- 5'-N- methylcarbamoyl- 4'- thioadenosine); BAA (8- butylaminoadenosine); 6-Amino-2-chloropurine riboside; 2-Chloroadenosine; NECA (5'-N- ethylcarboxamidoadenosine); APNEA (N6-2-(4-aminophenyl)ethyladenosine); or combinations thereof.

Modulators of Atoh 1

[00400] An additional sensory hair cell restorative agents are directed towards modulators to the products of the *Atoh1* (atonal; ATOH), *Neurod1* and *Neurog1* genes. *Atoh1* belongs to a family of basic Helix-Loop-Helix (bHLH) genes that are involved in cell fate determination across phyla and systems, typically being expressed in proliferating precursors. In mammals, at least three bHLH transcription factors are essential for sensory neuron development, including hair cells and sensory neurons of the ear: *Atoh1*, *Neurod1* and *Neurog1*. *Atoh1*, in particular, is essential for hair cell differentiation, and plays a role as a differentiation factor of postmitotic hair cells. Studies have also shown that expression of *Atoh1*, in combination with *Bdnf*, form afferent and efferent innervation in undifferentiated cells of epithelial origin.

[00401] Treatment of with ATOH protein supports the role of *Atoh1* in sensory hair cell development, inducing the formation of new sensory hair cells in cochlear structures, and restoring hearing and balance function. Gene therapy using vectors inserted with the *Atoh1* gene further supports ATOH's role in promoting and maintaining sensory hair cell function. Accordingly, one

embodiment disclosed herein is the use of ATOH proteins or manipulation of *Atoh1* expression to induce sensory hair cell development in hearing and balance disorders.

[00402] In additional embodiments, a neurotrophic growth factor is administered to the auris interna via the formulations described herein to stimulate inner ear hair cell neurotrophic growth factors.

5 The damage caused to spiral ganglion neurons removes not only neural activity, but also neurotrophin support that is normally supplied by hair cells, the absence of which leads to cell death via apoptosis.

[00403] In one embodiment, neurotrophic growth factor includes but is not limited to brain-derived neurotrophic factor, neurotrophin-3, glial-derived neurotrophic factor, neurotrophin-4/5, nerve growth factor, chlorphenylthio-cAMP (cptcAMP; a permeant cAMP analog), ciliary derived neurotrophic factor (CNTF) or combinations thereof. In another example, the sensory cell restorative agent is a brain-derived neurotrophic factor (BDNF). In yet another example, the neurotrophic growth factor is neurotrophin-3 (NT-3). In other examples, the neurotrophic growth factor is glial-derived neurotrophic factor (GDNF). In some examples, the neurotrophic growth factor is a peptide or protein. In other embodiments, the neurotrophic growth factor stimulates or enhances spiral ganglion neuron survival.

ERR/NR3B2 Antagonists

[00404] Studies have also suggested a role for the orphan receptor estrogen related receptor β /Nr3b2 in regulating endolymph production, thereby possibly playing a role in mediating cochlear and vestibular pressure in the endolymph fluid. (Chen et al. *Dev. Cell.* (2007) 13:325-337). Accordingly, agents which antagonize ERR/Nr3b2 expression, protein production or protein function are contemplated as useful with the formulations disclosed herein.

KCNQ Modulators

[00405] Modulators of KCNQ are also contemplated within the scope of the embodiments disclosed herein. KCNQ proteins form potassium channels, which play a role by preventing accumulation of potassium in hair cells. Potassium concentrations are high in the endolymph, giving the endocochlear fluid a high positive potential, which in turn provides a large drive force for potassium entry into the hair cell. KCNQ function is correlated with outer hair cell (OHC) survival; inhibition of KCNQ alters potassium homeostasis, resulting eventually in OHC degeneration. Accordingly, treatment of the auris interna with KCNQ modulators, in some cases activators, is contemplated within the scope of the embodiments disclosed herein as useful in the maintenance of sensory hair cell function in both vestibular and cochlear structures.

P2X Modulators

[00406] Modulators of P2X channel function are also contemplated within the scope of the embodiments, for use, for example, in auris interna disorders, such as cochlear inflammation and noise-induced hearing loss. P2X channels, which are gated by adenosine triphosphate, are present in

a broad distribution of tissues, and are thought to play a role in peripheral and central neuronal transmission, smooth muscle contraction and inflammation. Purine nucleotides are thought to play a role in cochlear disease, where ATP plays a cytotoxic role via both apoptosis and necrosis due to the activation of P2X receptors. For example, chronic perfusion of the perilymphatic space with ATP
5 causes the proliferation of fibrous tissue and neoostrogenesis in the scala tympani. Moreover, noise exposure and hypoxia cause a significant elevation of ATP concentration in the endolymphatic and perilymphatic compartments, which may represent an adaptive response of the cells to injury.

[00407] Accordingly, one embodiment is the use of modulators of P2X in the treatment of cochlear and vestibular disorders, including hearing and balance disorders. Antagonists and agonists to P2X
10 channels include BzATP, TNP-ATP, α,β -meATP, A-317491, PPADS, NF279, meSuramin, Reactive Blue II, RO-1, Adamantane amides, RO-3 and 4,5-diarylimidazolines.

CNS modulating agents

[00408] In some instances, immunomodulators and/or aural pressure modulators modulate central nervous system activity.

Anticholinergics

[00409] Contemplated for use with the formulations disclosed herein are agents which ameliorate otic disorders, including vestibular disorders and/or tinnitus, through local modulation of central nervous system (CNS) activity. Accordingly, some embodiments incorporate the use of agents which inhibit the release of the neurotransmitter acetylcholine in the CNS. Anticholinergic agents
20 are substances which block acetylcholine in the central and the peripheral nervous system. They treat balance disorders by suppressing conduction in vestibular cerebellar pathways, thus increasing motion tolerance.

[00410] In some embodiments, the anticholinergic is glycopyrrolate, homatropine, scopolamine or atropine. In some embodiments, the anticholinergic is glycopyrrolate. In some embodiments, the
25 anticholinergic is homatropine. In some embodiments, the anticholinergic is scopolamine. In some embodiments, the anticholinergic is atropine.

Antihistamines

[00411] Contemplated for use with the formulations disclosed herein are agents which ameliorate otic disorders, including vestibular disorders and/or tinnitus, through local modulation of central
30 nervous system (CNS) activity. Accordingly, some embodiments incorporate the use of agents which block the action of neurotransmitters in the CNS. Histamine is a neurotransmitter in the CNS. Accordingly, some embodiments incorporate the use of agents which modulate histamine receptors (e.g. the H₁ receptor, H₂ receptor, and/or the H₃ receptor). In some embodiments, antihistamines are as described herein.

Calcium Channel Blockers

[00412] Contemplated for use with the formulations disclosed herein are agents which ameliorate otic disorders, including vestibular disorders and/or tinnitus, through local modulation of central nervous system (CNS) activity. Accordingly, some embodiments incorporate the use of agents which block or antagonize Ca^{2+} channels. Calcium channels are channels formed in the plasma membrane of neurons (amongst other cells) by integral membrane proteins. These channels conduct Ca^{2+} through a cell's plasma membrane. In neurons, the flow of Ca^{2+} is partly responsible for creating and propagating action potentials in neurons. It can also be responsible for the release of certain neurotransmitters.

[00413] In some embodiments, the calcium channel antagonist is cinnarizine, flunarizine, or nimodipine. In some embodiments, the calcium channel antagonist is cinnarizine. In some embodiments, the calcium channel antagonist is flunarizine. In some embodiments, the calcium channel antagonist is nimodipine. Other calcium channel blockers include verapamil, diltiazem, omega-conotoxin, GVIA, amlodipine, felodipine, lacidipine, mibefradil, NPPB (5-Nitro-2-(3-phenylpropylamino)benzoic Acid), flunarizine, and/or combinations thereof

GABA Receptor Modulators

[00414] Contemplated for use with the formulations disclosed herein are agents which ameliorate otic disorders, including vestibular disorders and/or tinnitus, through local modulation of central nervous system (CNS) activity. Accordingly, some embodiments incorporate the use of agents which modulate the action of GABA receptors in the CNS. GABA, or γ -aminobutyric acid, is an inhibitory neurotransmitter in the CNS. It acts at inhibitory synapses of both pre- and postsynaptic neuronal processes. The binding of GABA to its receptors (the GABA_A receptor, the GABA_B receptor, and the GABA_C receptor) results in the opening of ion channels, and the flow of Cl^- into the cell and/or K^+ out of the neuron. The result is hyperpolarization of the neuron. Accordingly, some embodiments incorporate the use of agents which increase or decrease the sensitivity of the GABA receptors, or activate the GABA receptors by mimicking GABA.

[00415] The benzodiazepine class of therapeutic agents are agonists of the GABA_A receptor. When a benzodiazepine binds to the GABA_A receptor it induces a conformational change which increases the affinity of GABA for its receptor. The result of the increase in the binding of GABA is an increase in the frequency with which the Cl^- channels in the neurons open. This causes hyperpolarization of the neural membrane. In some embodiments, the benzodiazepine is selected from the group consisting of: alprazolam, bromazepam, brotizolam, chlordiazepoxide, clonazepam, clorazepate, diazepam, estazolam, flunitrazepam, flurazepam, loprazolam, lorazepam, lormetazepam, idazepam, nimetazepam, nitrazepam, oxazepam, prazepam, temazepam, triazolam or combinations thereof. In some embodiments, the benzodiazepine is clonazepam, diazepam, lorazepam, or combinations thereof. In some embodiments, the benzodiazepine is diazepam.

[00416] In some embodiments, the GABA receptor modulator is a loop diuretic. In some embodiments, the loop diuretic is furosemide, bumetanide, or ethacrynic acid. In some embodiments, the loop diuretic is furosemide. In some embodiments, the loop diuretic is bumetanide. In some embodiments, the loop diuretic is ethacrynic acid. Furosemide, for example, binds to the GABA_A receptor and reversibly antagonizes GABA-evoked currents of the $\alpha 6$, $\beta 2$, and $\gamma 2$ receptors. By way of example only, useful loop diuretics include, but are not limited to, furosemide, bumetanide, and ethacrynic acid.

[00417] In some embodiments, the modulator of a GABA receptor is a GABA analogue. GABA analogues mimic GABA. Thus, when they bind to a GABA receptor, the receptor acts as though GABA is binding to it and the receptor is activated. In some embodiments, the GABA analog is gabapentin, pregabalin, muscimol, or baclofen. In some embodiments, the GABA analog is gabapentin. In some embodiments, the GABA analog is pregabalin. In some embodiments, the GABA analog is muscimol. In some embodiments, the GABA analogue is baclofen. Baclofen is an analogue of GABA which binds to and activates the GABA_B receptor. Muscimol is also an analogue of GABA. It agonizes the GABA_A receptor.

Neurotransmitter Reuptake Inhibitors

[00418] Contemplated for use with the formulations disclosed herein are agents which ameliorate otic disorders, including vestibular disorders and/or tinnitus, through local modulation of central nervous system (CNS) activity. Accordingly, some embodiments incorporate the use of agents which inhibit the reuptake of neurotransmitters in the CNS. In some embodiments, the neurotransmitter reuptake modulator is an antagonist of a neurotransmitter reuptake target, partial agonist, inverse agonist, neutral or competitive antagonist, allosteric antagonist, and/or orthosteric antagonist. Neurotransmitter reuptake inhibitors inhibit the reuptake of neurotransmitters into presynaptic cells of the CNS. This increases the concentration of neurotransmitter available to stimulate post-synaptic cells of the CNS.

[00419] In some embodiments, the neurotransmitter reuptake inhibitors are tricyclic antidepressants. Tricyclic antidepressants work by inhibiting the re-uptake of the neurotransmitters norepinephrine and serotonin by pre-synaptic cells. This increases the level of serotonin and/or norepinephrine available to bind to the postsynaptic receptor. In some embodiments, the tricyclic antidepressant is amitriptyline, nortriptyline, or trimipramine. In some embodiments, the tricyclic antidepressant is amitriptyline. In some embodiments, the tricyclic antidepressant is nortriptyline. In some embodiments, the tricyclic antidepressant is trimipramine.

[00420] In some embodiments, the neurotransmitter reuptake inhibitor is a selective serotonin reuptake inhibitor. By inhibiting the reuptake of serotonin into the presynaptic cells, SSRIs increase the extracellular level of serotonin. This increases the level of serotonin available to bind to the postsynaptic receptor. SSRIs are hypothesized to stimulate new neural growth within the inner ear.

In some embodiments, the selective serotonin reuptake inhibitor is fluoxetine, paroxetine, or sertraline. In some embodiments, the selective serotonin reuptake inhibitor is fluoxetine. In some embodiments, the selective serotonin reuptake inhibitor is paroxetine. In some embodiments, the selective serotonin reuptake inhibitor is sertraline.

5 [00421] Contemplated for use with the formulations disclosed herein are agents that ameliorate otic disorders, including vestibular disorders and/or tinnitus, through local modulation of central nervous system (CNS) activity. Accordingly, some embodiments incorporate the use of agents that antagonize neurokinin receptors. There are at least three neurokinin receptors: NK1, NK2 and NK3. In certain embodiments, the binding of a ligand (e.g. a tachykinin peptide, substance P, neurokinin
10 A, and neurokinin B) to a neurokinin receptor induces the activation of phospholipase C. The activation of phospholipase C produces inositol triphosphate. In some embodiments, the neurokinin receptor is the NK1 receptor, the NK2 receptor, the NK3 receptor, or combinations thereof. In some embodiments, the neurokinin receptor is the NK1 receptor. In some embodiments, the antagonist of the NK1 receptor is vestipitant.

15 [00422] In some embodiments, the SSRI inhibitor is administered in combination with a neurokinin receptor antagonist. In some embodiments, the SSRI is paroxetine and the neurokinin receptor is NK1. In some embodiments, the NK1 receptor antagonist is vestipitant. In certain embodiments, the co-administration of paroxetine and vestipitant treats, and/or the symptoms of tinnitus.

Local Anesthetics

20 [00423] Contemplated for use with the formulations disclosed herein are agents which ameliorate otic disorders, including vestibular disorders and/or tinnitus, through local modulation of central nervous system (CNS) activity. Accordingly, some embodiments incorporate the use of agents which decrease the rate of the depolarization and repolarization of neurons by, for example, blocking the Na⁺ channels in cell membranes.

25 [00424] In some embodiments, the CNS modulator is a local anesthetic. In some embodiments, the local anesthetic is selected from the group consisting of: benzocaine, carticaine, cinchocaine, cyclomethycaine, lidocaine, prilocaine, propoxycaine, proparacaine, tetracaine, tocaïnide, and trimecaine. In some embodiments, the local anesthetic is lidocaine. In some embodiments, the local anesthetic is tocaïnide.

30 *Sodium Channel Blockers*

[00425] Contemplated for use with the formulations disclosed herein are agents which ameliorate otic disorders, including vestibular disorders and/or tinnitus, through local modulation of central nervous system (CNS) activity. Accordingly, some embodiments incorporate the use of agents which block or antagonize Na⁺ channels. Sodium channels are channels formed in the plasma
35 membrane of neurons (amongst other cells) by integral membrane proteins. These channels conduct

Na⁺ through a cell's plasma membrane. In neurons, the flow of Na⁺ is partly responsible for creating and propagating action potentials in the neurons.

[00426] In some embodiments, the sodium channel blocker is carbamazepine, oxcarbazepine, phenytoin, valproic acid, or sodium valproate. In some embodiments, the sodium channel blocker is carbamazepine. In some embodiments, the sodium channel blocker is oxcarbazepine. In some embodiments, the sodium channel blocker is phenytoin. In some embodiments, the sodium channel blocker is valproic acid. In some embodiments, the sodium channel blocker is sodium valproate.

[00427] In some embodiments, the Na⁺ channel blocker is vinpocetine ((3a,16a)-Eburnamenine-14-carboxylic acid ethyl ester); sipatrigine (2-(4-Methylpiperazin-1-yl)-5-(2,3,5-trichlorophenyl)-pyrimidin-4-amine); amiloride (3,5-diamino-N-(aminoiminomethyl)-6-chloropyrazinecarbox amide hydrochloride); carbamazepine (5H-dibenzo[b,f]azepine-5-carboxamide); TTX (octahydro-12-(hydroxymethyl)-2-imino-5,9:7,10a-dimethan o-10aH-[1,3]dioxocino[6,5-d]pyrimidine-4,7,10,11,12-pentol); RS100642 (1-(2,6-dimethyl-phenoxy)-2-ethylaminopropane hydrochloride); mexiletine ((1-(2,6-dimethylphenoxy)-2-aminopropane hydrochloride)); QX-314 (N-(2,6-Dimethylphenylcarbamoylmethyl)triethylammonium bromide); phenytoin (5,5-diphenylimidazolidine-2,4-dione); lamotrigine (6-(2,3-dichlorophenyl)-1,2,4-triazine-3,5-diamine); 4030W92 (2,4-diamino-5-(2,3-dichlorophenyl)-6-fluoromethylpyrimidine); BW1003C87 (5-(2,3,5-trichlorophenyl) pyrimidine-2,4- 1.1 ethanesulphonate); QX-222 (2-[(2,6-dimethylphenyl)amino]-N,N,N-trimethyl-2-oxoethaniminium chloride); ambroxol (trans-4-[(2-Amino-3,5-dibromophenyl)methyl]amino]cyclohexanol hydrochloride); R56865 (N-[1-(4-(4-fluorophenoxy)butyl]-4-piperidinyl-N-methyl-2-benzo-thiazolamine); lubeluzole; ajmaline ((17R,21alpha)-ajmalan-17,21-diol); procainamide (4-amino-N-(2-diethylaminoethyl)benzamide hydrochloride); flecainide; riluzole; or combinations thereof.

[00428] In some embodiments, agents which decrease the rate of the depolarization and repolarization of neurons by, for example, blocking the Na⁺ channels in cell membranes include local anesthetics. In some embodiments, the local anesthetic is selected from the group consisting of: benzocaine, articaine, cinchocaine, cyclomethycaine, lidocaine, prilocaine, propoxycaine, proparacaine, tetracaine, tocainide, and trimecaine. In some embodiments, the local anesthetic is lidocaine. In some embodiments, the local anesthetic is tocainide.

Thyrotropin-Releasing Hormone

[00429] Contemplated for use with the formulations disclosed herein are agents which ameliorate otic disorders, including vestibular disorders and/or tinnitus, through local modulation of central nervous system (CNS) activity. Accordingly, some embodiments incorporate the use of agents which modulate neurotransmitters. Thyrotropin-releasing hormone is a neurotransmitter which inhibits glutamate-induced excitation of neurons. In some embodiments, the CNS modulator is thyrotropin-releasing hormone.

Antimicrobial Agents

[00430] Any antimicrobial agent useful for the treatment of otic disorders, e.g., inflammatory diseases of the ear or cancer of the ear, is suitable for use in the formulations and methods disclosed herein. In some embodiments, the antimicrobial agent is an antibacterial agent, an antifungal agent, an antiviral agent, an antiprotozoal agent, and/or an antiparasitic agent. Antimicrobial agents include agents that act to inhibit or eradicate microbes, including bacteria, fungi, viruses, protozoa, and/or parasites. Specific antimicrobial agents are used to combat specific microbes. Accordingly, a skilled practitioner would know which antimicrobial agent would be relevant or useful depending on the microbe identified, or the symptoms displayed.

[00431] In some embodiments, the antimicrobial agent is a protein, a peptide, an antibody, DNA, a carbohydrate, an inorganic molecule, or an organic molecule. In certain embodiments, the antimicrobial agents are antimicrobial small molecules. Typically, antimicrobial small molecules are of relatively low molecular weight, e.g., less than 1,000, or less than 600-700, or between 300-700 molecular weight.

[00432] Antibacterial agents include amikacin, gentamicin, kanamycin, neomycin, netilmicin, streptomycin, tobramycin, paromomycin, geldanamycin, herbimycin, loracarbef, ertapenem, doripenem, imipenem, cilastatin, meropenem, cefadroxil, cefazolin, cefalotin, cefalexin, cefaclor, cefamandole, cefoxitin, defprozil, cefuroxime, cefixime, cefdinir, cefditoren, cefoperazone, cefotaxime, cefpodoxime, ceftazidime, ceftibuten, ceftizoxime, ceftriaxone, cefepime, ceftobiprole, teicoplanin, vancomycin, azithromycin, clarithromycin, dirithromycin, erythromycin, roxithromycin, troleandomycin, telithromycin, spectinomycin, aztreonam, amoxicillin, ampicillin, azlocillin, carbenicillin, cloxacillin, dicloxacillin, flucloxacillin, mezlocillin, meticillin, nafcillin, oxacillin, penicillin, piperacillin, ticarcillin, bacitracin, colistin, polymyxin B, ciprofloxacin, enoxacin, gatifloxacin, levofloxacin, lomefloxacin, moxifloxacin, norfloxacin, ofloxacin, trovfloxacin, mafenide, prontosil, sulfacetamide, sulfamethizole, sulfanilamide, sulfsalazine, sulfsioxazole, trimethoprim, demeclocycline, doxycycline, minocycline, oxtetracycline, tetracycline, arspenamine, chloramphenicol, clindamycin, lincomycin, ethambutol, fosfomycin, fusidic acid, furazolidone, isoniazid, linezolid, metronidazole, mupirocin, nitrofurantoin, platensimycin, pyrazinamide, quinupristin/dalfopristin, rifampin, tinidazole, AL-15469A (Alcon Research), AL-38905 (Alcon Research) and combinations thereof.

[00433] Antiviral agents include acyclovir, famciclovir and valacyclovir. Other antiviral agents include abacavir, aciclovir, adfovir, amantadine, amprenavir, arbidol, atazanavir, artipia, brivudine, cidofovir, combivir, edoxudine, efavirenz, emtricitabine, enfuvirtide, entecavir, fomvirsen, fosamprenavir, foscarnet, fosfonet, ganciclovir, gardasil, ibacitabine, immunovir, idoxuridine, imiquimod, indinavir, inosine, integrase inhibitors, interferons, including interferon type III, interferon type II, interferon type I, lamivudine, lopinavir, loviride, MK-0518, maraviroc,

moroxydine, nelfinavir, nevirapine, nexavir, nucleoside analogues, oseltamivir, penciclovir, peramivir, pleconaril, podophyllotoxin, protease inhibitors, reverse transcriptase inhibitors, ribavirin, rimantadine, rilonavir, saquinavir, stavudine, tenofovir, tenofovir disoproxil, tipranavir, trifluridine, trizivir, tromantadine, truvada, valganciclovir, vicriviroc, vidarabine, viramidine, zalcitabine,

zanamivir, zidovudine, and combinations thereof.

[00434] Antifungal agents include amroline, utenafine, naftifine, terbinafine, flucytosine, fluconazole, itraconazole, ketoconazole, posaconazole, ravuconazole, voriconazole, clotrimazole, econazole, miconazole, oxiconazole, sulconazole, terconazole, tioconazole, nikkomycin Z, caspofungin, micafungin, anidulafungin, amphotericin B, liposomal nystatin, pimaricin, griseofulvin, ciclopirox olamine, haloprogin, tolnaftate, undecylenate, clioquinol, and combinations thereof.

[00435] Antiparasitic agents include amitraz, amoscanate, avermectin, carbadox, diethylcarbamazine, dimetridazole, diminazene, ivermectin, macrofilaricide, malathion, mitaban, oxamniquine, permethrin, praziquantel, prantel pamoate, selamectin, sodium stibogluconate, thiabendazole, and combinations thereof.

[00436] In some embodiments, pharmaceutically active metabolites, salts, polymorphs, prodrugs, analogues, and derivatives of the antimicrobial agents discussed above that retain the ability of the parent antimicrobial agents to treat otic disorders of the ear are also useful in the formulations disclosed herein.

Free radical modulators

[00437] In some instances, immunomodulators and/or aural pressure modulators relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or the dysfunction of the mitochondria.

Antioxidants

[00438] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of agents which prevent and/or ameliorate the damage caused by free radicals. In some embodiments, the agents which prevent and/or ameliorate the damage caused by free radicals is an antioxidant.

[00439] Antioxidants, as disclosed herein, are also useful as protectants against ototoxic agents through the prevention of reactive oxygen species, neutralization of toxic products or blockage of the apoptosis pathway. Resveratrol (3,5,4'-Trihydroxystilbene), a representative example of an antioxidant, exerts its effects through a variety of pathways, including the inhibition of MnSOD, which reduces superoxide to H_2O_2 , which inhibits free radical chain reactions, reducing superoxide levels in the cell. Moreover, resveratrol has been implicated in preventing neuronal cell dysfunction and cell death. Other antioxidants include but are not limited to vitamin E (tocopherol), vitamin C

(ascorbic acid), glutathione, lipoic acid, alpha lipoic acid, uric acid, carotenes, ubiquinol, melatonin, tocotrienols, selenium, flavonoids, polyphenols, lycopene, lutein, lignan, butyl hydroxytoluene, coenzyme Q10, salicylate, or combinations thereof.

[00440] In certain embodiments, nitrones act synergistically with antioxidants. In certain

5 embodiments, nitrones trap free radicals. In some embodiments, a nitron (e.g. alpha-phenyl-tert-butyl nitron (PBN), allpurinol) is co-administered with an antioxidant. In certain embodiments, a nitron co-administered with an antioxidant treats acute acoustic noise-induced hearing loss.

[00441] In some embodiments, the antioxidant is N-acetylcysteine; vitamin E (tocopherols and tocotrienols); vitamin C; vitamin A; lutein; selenium glutathione; melatonin; a polyphenol; a

10 carotenoid (e.g. lycopene, carotenes); coenzyme Q-10; Ebselen (2-phenyl-1, 2-benzisoselenazol-3(2H)-one (also called PZ 51 or DR3305); L-methionine; azulenyl nitrones (e.g. stilbazulenyl nitron); L-(+)-Ergothioneine ((S)-a-Carboxy-2,3-dihydro-N,N,N-trimethyl-2-thioxo-1H-imidazole-4-ethanaminium inner salt); Caffeic Acid Phenyl Ester (CAPE); dimethylthiourea; dimethylsulfoxide; disulfenton sodium (NXY-059; disodium 4-[(Z)-(tert-butyl-oxidoazaniumylidene)methyl]benzene-1,3-disulfonate); pentoxifylline; MCI-186 (3-Methyl-1-phenyl-2-pyrazolin-5-one); Ambroxol (trans-4-(2-Amino-3,5-dibromobenzylamino)cyclohexane-HCl; U-83836E ((-)-2-((4-(2,6-di-1-Pyrrolidinyl-4-pyrimidinyl)-1-piperzainyl)methyl)-3,4-dihydro-2,5,7,8-tetramethyl-2H-1-benzopyran-6-ol•2HCl); MITOQ (mitoquinone mesylate, Antipodean Pharmaceuticals); Idebenone (2-(10-hydroxydecyl)-5,6-dimethoxy-3-methyl-cyclohexa-2,5-diene-1,4-dione); (+)-cyanidanol-3; or combinations thereof.

20

Glutamate-Receptor Modulators

[00442] Contemplated for use with the formulations disclosed herein are agents that modulate the production of free-radicals and/or inhibit damage to the mitochondria. Accordingly, some embodiments incorporate the use of agents which modulate glutamate receptors. In some

25 embodiments, the glutamate receptor is the AMPA receptor, the NMDA receptor, and/or a group II or III mGlu receptor. In some embodiments, a glutamate receptor modulator is as described herein.

Iron Chelators

[00443] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of agents which prevent and/or ameliorate the damage caused by free radicals. In some embodiments, the agents which prevent and/or ameliorate the damage caused by free radicals is an iron chelator. The iron chelator, deferoxamine, prevents ototoxic damage to the ear resulting from treatment with neomycin when it is co-administered with neomycin.

35 [00444] In some embodiments, the iron chelator is desferrioxamine (DFO); hydroxybenzyl ethylene diamine; fullerenol-1, pyrrolidine dithiocarbamate; desferal; Vk-28 (5-[4-(2-hydroxyethyl)

piperazine-1-ylmethyl]-quinoline-8-ol); clioquinol; echinochrome; PIH (pyridoxal isonicotinoyl hydrazone); deferasirox; HBED (N,N'-bis (2-hydroxybenzyl) ethylenediamine-N,N'-diacetic acid); SIIH (salicylaldehyde isonicotinoyl hydrazone); deferiprone; L1 (1,2-dimethyl-3-hydroxy-4-pyridone); Kojic acid (5-hydroxy-2-hydroxymethyl-4-pyrone); deferoxamine; 2,3-dihydroxybenzoate; or combinations thereof.

Mitochondrial Modulators

[00445] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of one or more agents that modulate the activity of the mitochondria. In some embodiments, the agent which modulates the activity of the mitochondria is acetylcarnitine; lipoic acid; or combinations thereof.

Nitric Oxide Synthase modulators

[00446] Contemplated for use with the compositions disclosed herein are agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Nitric oxide (NO) is a neurotransmitter. It is synthesized by multiple nitric oxide synthases (NOS) from arginine and oxygen. It is also derived from the reduction of inorganic nitrate. In certain instances, it induces vasodilation; thus, increasing blood flow. In certain instances, it increases cochlear blood flow. In certain instances, NO damages blood vessel walls. In certain instances, NO ameliorates vascular protein leakage in the cochlea. In certain instances, NO increases the sensitivity of hair cells. In certain instances, NO reacts with super-oxide to form the free radical peroxynitrite. Accordingly, some embodiments incorporate the use of agents that modulate nitric oxide and/or nitric oxide synthase (NOS).

[00447] In some embodiments, the agent that modulates NO and/or NOS is an antagonist of NO or NOS. In some embodiments, the antagonist of NO and/or NOS is aminoguanidine; 1-Amino-2-hydroxyguanidine *p*-toluensulfate; GED (guanidinoethyl disulfide); bromocriptine mesylate; dexamethasone; SDMA (symmetric N^G,N^G-Dimethyl-L-arginine); ADMA (asymmetric N^G,N^G-Dimethyl-L-arginine); L-NMMA (N^G-monomethyl-L-arginine); L-NMEA (N^G-monoethyl-L-arginine); D-MMA (N^G-monomethyl-D-arginine); L-NIL (N^G-(1-Iminoethyl)-L-lysine hydrochloride); L-NNA (N^G-nitro-L-arginine); L-NPA (N^G-propyl-L-arginine); L-NAME (N^G-nitro-L-arginine methyl ester dihydrochloride); L-VNIO (N^S-(1-imino-3-butenyl)-L-ornithine); diphenyleneiodonium chloride; 2-ethyl-2-thiopseudourea; haloperidol; L-NIO (L-N^S-(1-iminoethyl)ornithine); MEG (methylecgonidine); SMT (S-methylisothiourea sulfate); SMTC (S-methyl-L-thiocitrulline); 7-Ni (7-nitroindazole); nNOS inhibitor I ((4S)-N-(4-Amino-5[aminoethyl]aminopentyl)-N'-nitroguanidine); 1,3-PBITU (S,S'-1,3-Phenylene-bis(1,2-ethanediy)-bis-isothiourea); L-thiocitrulline; TRIM (1-(2-trifluoromethylphenyl) imidazole); MTR-

105 (S-ethylisothiuronium diethylphosphate); BBS-1; BBS-2; ONO-1714 ((1S,5S,6R,7R)-7chloro-3-amino-5methyl-2-azabicyclo[4.1.0]heptane hydrochloride); GW273629 (3-[[2-[(1-iminoethyl)amino]ethyl]sulphonyl]-L-alanine); GW 274150 ((S)-2-amino-(1-iminoethylamino)-5-thioheptanoic acid); PPA250 (3-(2,4-difluorophenyl)-6-{2-[4-(1H-imidazol-1-ylmethyl)phenoxy]ethoxy}-2-phenylpyridine); AR-R17477 ([N-(4-(2-((3-chlorophenylmethyl) amino) ethyl) phenyl)-2-thiophenecarboxamidine dihydrochloride); AR-R18512 (N(2-methyl-1,2,3,4-tetrahydroisoquinoline-7-yl)-2-thiophenecarboximidamide); spiroquinazalone; 1400W (N-[[3-(aminomethyl)phenyl]methyl]-ethanimidamide dihydrochloride); or combinations thereof.

[00448] In some embodiments, the agent that modulates NO and/or NOS is an agonist of NO and/or NOS, or a donor of NO. In some embodiments, the agonist of NO and/or NOS, or donor of NO, is S-NC (S-nitrosocysteine); NTG (nitroglycerine); SNP (sodium nitroprusside); thapsigargin; vascular endothelial growth factor (VEGF); bradykinin; ATP; sphingosine-1-phosphate; estrogen; angiopoietin; acetylcholine; SIN-1 (3-morpholiniosydnonimine); GEA 3162 (1,2,3,4-oxatriazolium, 5-amino-3-(3,4-dichlorophenyl)-,chloride); GEA 3175 (3-(3-chloro-2-methylphenyl)-5-[[4-methylphenyl]sulphonyl]amino]-hydroxide); GEA 5024 (1,2,3,4-oxatriazolium,5-amino-3-(3chloro-2-methyl-phenyl)chloride); GEA 5538 (,2,3,4-Oxatriazolium,3-(3-chloro-2-methylphenyl)-5-[[[cyanomethylamino]carbonyl]amino]-hydroxide inner salt); SNAP (S-nitroso-N-acetylpenicillamine); molsidomine; CNO-4 (1-[(4',5'-Bis(carboxymethoxy)-2'-nitrophenyl)methoxy]-2-oxo-3,3-diethyl-1-triazene dipotassium salt); CNO-5 ([1-(4',5'-Bis(carboxymethoxy)-2'-nitrophenyl)methoxy]-2-oxo-3,3-diethyl-1-triazine diacetoxymethyl ester); DEA/NO, IPA/NO, SPER/NO, SULFI/NO, OXI/NO, DETA/NO; or combinations thereof.

Sirtuin Modulators

[00449] The sirtuins (or Sir2 proteins) comprise class III of the histone deacetylases (HDACs).

While they are classified as protein deacetylases some also function as mono-ADP-

25 ribosyltransferases. Each sirtuin protein has a homologous core sequence of 250 amino acids. This sequence is highly conserved over multiple species. Further, in order to catalyze the deacetylation of a protein, each sirtuin requires NAD⁺ as a cofactor. There are seven members of the family: Sirt1, Sirt2, Sirt3, Sirt4, Sirt5, Sirt6, and Sirt7. Sirt1 and Sirt3 are protein deacetylases. Sirt2 is involved in mitosis.

30 [00450] Agonism of Sirt1 yields multiple benefits which have previously been identified in subjects undergoing caloric restriction. These benefits include, but are not limited to, decreased glucose levels and improved insulin sensitivity, increased mitochondrial activity, and decreased adiposity (due to the Sirt1 mediated repression of PPAR- γ). Decreases in glucose levels and adiposity can contribute to the amelioration of presbycusis as diabetes and atherosclerosis are both factors which
35 contribute to the development and progression of presbycusis.

[00451] Sirt1 can prevent apoptosis by deacetylating the pro-apoptotic genes p53 and Ku-70.

Additional substrates for Sirt1 include, but are not limited to, the transcription factors NFκB, Fox01, Fox03a, Fox04, Fox05; the transcription repressor Hic1; and Pgc-1α, which regulates, among other cellular functions, adaptive thermogenesis, glucose metabolism, and triglyceride metabolism.

5 Agonism of Sirt3 results in increased cellular respiration and a decrease in the production of reactive oxygen species (ROS).

[00452] The catalysis of deacetylation by sirtuins is NAD⁺ (nicotinamide adenine dinucleotide) dependent. Upon binding to an acetylated protein, the sirtuin hydrolyzes NAD⁺ by breaking the glycosidic bond between nicotinamide and ADP-ribose. The acetyl group of the acetylated protein is
10 then transferred to ADP-ribose. At the completion of the reaction nicotinamide, the deacetylated protein, and 2'-O-acetyl-ADP-ribose are released.

[00453] Multiple compounds modulate the sirtuin catalyzed deacetylation of proteins.

Administration of certain polyphenols such as, but not limited to, stilbenes, chalcones, flavones, isoflavones, flavanones, anthocyanidins, catechins, results in the decrease of the K_m of the
15 deacetylation reaction. Further, as free nicotinamide antagonizes the deacetylation reaction, compounds which inhibit the binding of nicotinamide to sirtuins will also agonize the activity of sirtuins.

[00454] Administration of the sirtuin agonizing agent resveratrol (*trans*-3,5,4'-trihydroxystilbene) decreases apoptosis. It also increases glutamate uptake and thus ameliorates excitotoxicity. Further,
20 administration of resveratrol results in lower levels of reactive oxygen species (ROS) and thus ameliorates damage caused by ischemia, excitotoxicity, ototoxicity caused by cisplatin and aminoglycosides, acoustic trauma and presbycusis.

[00455] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or
25 the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of one or more agents that modulate sirtuin catalyzed deacetylation reactions. In some embodiments, the agent which modulates sirtuin catalyzed deacetylation reactions is a stilbene. In some embodiments, the stilbene is *trans*-stilbene, *cis*-stilbene, resveratrol, piceatannol, rhapontin, deoxyrhapontin, butein, or combinations thereof.

30 [00456] In some embodiments, the stilbene is resveratrol. In some embodiments, the stilbene is an analog of resveratrol. In some embodiments, the analog of resveratrol is SRT-501 (RM-1821). For additional analogs of resveratrol see U.S. Patent App. Pub. No. 2006/0276393, which is hereby incorporated by reference for this disclosure.

[00457] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent,
35 reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of one or

more agents the modulate sirtuin catalyzed deacetylation reactions. In some embodiments, the agent which modulates sirtuin catalyzed deacetylation reactions is a chalcone. In some embodiments, the chalcone is chalcon; isoliquiritigen; butein; 4,2',4'-trihydroxychalcone; 3,4,2',4',6'-pentahydroxychalcone; or combinations thereof.

5 [00458] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of one or more agents the modulate sirtuin catalyzed deacetylation reactions. In some embodiments, the agent which modulates sirtuin catalyzed deacetylation reactions is a flavone. In some embodiments, the
10 flavone is flavone, morin, fisetin; luteolin; quercetin; kaempferol; apigenin; gossypetin; myricetin; 6-hydroxyapigenin; 5-hydroxyflavone; 5,7,3',4',5'-pentahydroxyflavone; 3,7,3',4',5'-pentahydroxyflavone; 3,6,3',4'-tetrahydroxyflavone; 7,3',4',5'-tetrahydroxyflavone; 3,6,2',4'-tetrahydroxyflavone; 7,4'-dihydroxyflavone; 7,8,3',4'-tetrahydroxyflavone; 3,6,2',3'-tetrahydroxyflavone; 4'-hydroxyflavone; 5-hydroxyflavone; 5,4'-dihydroxyflavone; 5,7-
15 dihydroxyflavone; or combinations thereof.

[00459] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of one or more agents the modulate sirtuin catalyzed deacetylation reactions. In some embodiments, the agent
20 which modulates sirtuin catalyzed deacetylation reactions is an isoflavone. In some embodiments, the isoflavone is daidzein, genistein, or combinations thereof.

[00460] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of one or
25 more agents the modulate sirtuin catalyzed deacetylation reactions. In some embodiments, the agent which modulates sirtuin catalyzed deacetylation reactions is a flavanone. In some embodiments, the flavanone is naringenin; flavanone; 3,5,7,3',4'-pentahydroxyflavanone; or combinations thereof.

[00461] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or
30 the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of one or more agents the modulate sirtuin catalyzed deacetylation reactions. In some embodiments, the agent which modulates sirtuin catalyzed deacetylation reactions is an anthocyanidin. In some embodiments, the anthocyanidin is pelargonidin chloride, cyanidin chloride, delphinidin chloride, or combinations thereof.

35 [00462] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or

the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of one or more agents that modulate sirtuin catalyzed deacetylation reactions. In some embodiments, the agent which modulates sirtuin catalyzed deacetylation reactions is a catechin. In some embodiments, the catechin is (–)-epicatechin (Hydroxy Sites: 3,5,7,3',4'); (–)-catechin (Hydroxy Sites: 3,5,7,3',4'); (–)-gallocatechin (Hydroxy Sites: 3,5,7,3',4',5') (+)-catechin (Hydroxy Sites: 3,5,7,3',4'); (+)-epicatechin (Hydroxy Sites: 3,5,7,3',4'); or combinations thereof.

[00463] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of one or more agents that modulate the catalytic rate of sirtuin catalyzed deacetylation reactions. In some embodiments, the agent which modulates the catalytic rate of sirtuin catalyzed deacetylation reactions is dipyrindamole, ZM 336372 (3-(dimethylamino)-N-[3-[(4-hydroxybenzoyl)-amino]-4-methylphenyl]benzamide), camptothecin, coumestrol, nordihydroguaiaretic acid, esculetin, SRT-1720 (Sirtris), SRT-1460 (Sirtris), SRT-2183 (Sirtris), or combinations thereof.

[00464] Contemplated for use with the formulations disclosed herein are agents that relieve, prevent, reverse or ameliorate the degeneration of neurons and/or hair cells of the auris due to free radicals or the dysfunction of the mitochondria. Accordingly, some embodiments incorporate the use of one or more agents that modulate sirtuin catalyzed deacetylation reactions. In some embodiments, the agent that modulates sirtuin catalyzed deacetylation reactions is a nicotinamide binding antagonist. In some embodiments, the nicotinamide binding antagonist is isonicotinamide or an analog of isonicotinamide. In some embodiments, the analog of isonicotinamide is β -1'-5-methyl-nicotinamide-2'-deoxyribose; β -D-1'-5-methyl-nicotinamide-2'-deoxyribofuranoside; β -1'-4,5-dimethyl-nicotinamide-2'-deoxyribose; or β -D-1'-4,5-dimethyl-nicotinamide-2'-deoxyribofuranoside. For additional analogs of isonicotinamide see U.S. Pat. Nos. 5,985,848; 6,066,722; 6,228,847; 6,492,347; 6,803,455; and U.S. Patent Publication Nos. 2001/0019823; 2002/0061898; 2002/0132783; 2003/0149261; 2003/0229033; 2003/0096830; 2004/0053944; 2004/0110772; and 2004/0181063, which are hereby incorporated by reference for that disclosure.

Ion channel modulators

Potassium Ion Channel Modulators

[00465] Contemplated for use with the formulations disclosed herein are agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs and neurons in the inner ear. Accordingly, some embodiments incorporate the use of agents that modulate potassium ion concentrations. In some embodiments, the agents that modulate potassium ion concentrations are agonists or antagonists of potassium ion channels.

Potassium ion channels are channels that regulate the flow of potassium ions into and out of cells. In the cochlea the transduction current through the sensory cells is carried by potassium ions and

depends on the high concentration of potassium ions in the endolymph. Mutations in the genes encoding potassium channel protein result in both acquired and congenital hearing loss.

[00466] The KCNQ family of potassium channels is a family of delayed rectifier voltage-gated potassium channels found in the cochlea. KCNQ1 subunits form potassium channels in vestibular dark cells and marginal cells of the stria vascularis. These channels regulate the level of potassium in endolymph. KCNQ4 subunits form channels hair cells. Mice with genes encoding KCNQ subunits knocked-out display a hearing loss during development, starting at four weeks of postnatal life.

[00467] In some embodiments, the agent that modulates a potassium channel is an agonist of a potassium channel (e.g. a potassium channel opener). In some embodiments, the agonist of a potassium channel is nicorandil; minoxidil; levcromakalim; lemakalim; cromakalim; L-735,334 (14-hydroxy CAF-603 oleate); retigabine; flupirtine; BMS-204352 (3S)-(+)-(5-Chloro-2-methoxyphenyl)-1,3-dihydro-3-fluoro-6-(trifluoromethyl)-2H-indole-2-one); DMP-543 (10,10-bis((2-fluoro-4-pyridinyl)methyl)-9(10H)-anthracenone); or combinations thereof.

[00468] In some embodiments, the agent that modulates a potassium channel is an antagonist of a potassium channel (e.g. a potassium channel blocker). In some embodiments, the antagonist of a potassium channel is linopirdine; XE991 (10,10-bis(4-pyridinylmethyl)-9(10H)-anthracenone); 4-AP (4-aminopyridine); 3,4-DAP (3,4-Diaminopyridine); E-4031 (4'-[[1-[2-(6-methyl-2-pyridyl)ethyl]-4-piperidinyl]carbonyl]-methanesulfonanilide); DIDS (4,4'-diisothiocyanostilbene-2,2'-disulfonic acid); Way 123,398 (N-methyl-N-(2-(methyl(1-methyl-1H-benzimidazol-2-yl)amino)ethyl)-4-((methylsulfonyl)amino) benzenesulfonamide HCl); CGS-12066A (7-Trifluoromethyl-4-(4-methyl-1-piperazinyl)pyrrolo-[1,2-a]quinoxaline); dofetilide; sotalol; apamin; amiodarone; azimilide; bretylium; clofilium; tedisamil; ibutilide; sematilide; nifekalant; tamulustoxin and combinations thereof.

Purigenic Receptor Modulators

[00469] Contemplated for use with the formulations disclosed herein are agents for modulating ion channels. Accordingly, some embodiments incorporate the use of agents that modulate the concentration of ions. In some embodiments, the agents that modulate the concentration of ions are agonists or antagonists of purigenic receptors.

[00470] Purigenic receptors are a family of plasma membrane-bound receptors. The family includes the P2X, P2Y, and P1 receptors. The P2X receptors comprise ion channels. When ATP binds to the receptor the channel opens. The P2Y receptors comprise G-coupled protein receptors. The ligands for these receptors are ATP, ADP, UTP, UDP, UDP-glucose. The P1 receptors comprise G-coupled protein receptors. The ligand for these receptors is adenosine. Purigenic receptors regulate ion homeostasis in the ear. Endolymph, for example, requires high potassium (K^+), low sodium (Na^+), and low calcium (Ca^{2+}) ion levels for normal auditory transduction.

[00471] In some embodiments, the agonist of a purigenic receptor is ATP; ADP; UTP; UDP; UDP-glucose; adenosine; 2-MeSATP; 2-MeSADP; $\alpha\beta$ meATP; dATP α S; ATP γ S; Bz-ATP; MRS2703 (2-MeSADP with the beta-phosphate group blocked by a 1-(3,4-dimethyloxyphenyl)eth-1-yl phosphoester); denufosol tetrasodium; MRS2365 ([[(1R,2R,3S,4R,5S)-4-[6-amino-2-(methylthio)-9H-purin-9-yl]-2,3-dihydroxybicyclo[3.1.0]hex-1-yl]methyl] diphosphoric acid mono ester trisodium salt); MRS 2690 (diphosphoric acid 1- α -D-glucopyranosyl ester 2-[(4'-methylthio)uridin-5''-yl] ester disodium salt); PSB 0474 (3-(2-Oxo-2-phenylethyl)-uridine-5'-diphosphate disodium salt); or combinations thereof.

[00472] In some embodiments, the antagonist of a purigenic receptor is A-317491 ((5-[(3-Phenoxybenzyl)[(1S)-1,2,3,4-tetrahydro-1-naphthalenyl]amino]carbonyl)-1,2,4-benzenetricarboxylic acid); RO-3 (Roche); suramin; PPADS (pyridoxalphosphate-6-azophenyl-2',4'-disulfonic acid); PPNDS (Pyridoxal-5'-phosphate-6-(2'-naphthylazo-6'-nitro-4',8'-disulfonate) tetrasodium salt); DIDS; pyridoxal-5-phosphate; 5-(3-bromophenyl)-1,3-dihydro-2H-benzofuro-[3,2-e]-1,4-diazepin-2-one; cibacron blue; basilen blue; ivermectin; A-438079 (3-[[5-(2,3-Dichlorophenyl)-1H-tetrazol-1-yl]methyl]pyridine hydrochloride); A-740003 ((N-(1-[(cyanoimino)(5-quinolinylamino) methyl]amino)-2,2-dimethylpropyl)-2-(3,4-dimethoxyphenyl)acetamide); NF449 (4,4',4'',4'''-(carbonylbis(imino-5,1,3-benzenetriylbis(carbonylimino)))tetrakis-benzene-1,3-disulfonic acid); NF110 (*para*-4,4',4'',4'''-(carbonylbis(imino-5,1,3-benzenetriylbis carbonylimino)))tetrakis-benzenesulfonic acid); MRS 2179 (2'-Deoxy-N6-methyladenosine 3',5'-bisphosphate tetrasodium salt); MRS 2211 (2-[(2-chloro-5-nitrophenyl)azo]-5-hydroxy-6-methyl-3-[(phosphonooxy)methyl]-4-pyridinecarboxaldehyde disodium salt); MRS 2279 ((1R,2S,4S,5S)-4-[2-chloro-6-(methylamino)-9H-purin-9-yl]-2-(phosphonooxy)bicyclo[3.1.0]hexane-1-methanol dihydrogen phosphate ester diammonium salt); MRS 2500 tetrasodium salt ((1R,2S,4S,5S)-4-[2-Iodo-6-(methylamino)-9H-purin-9-yl]-2-(phosphonooxy)bicyclo[3.1.0]hexane-1-methanol dihydrogen phosphate ester tetraammonium salt); NF157 (8,8'-[carbonylbis[imino-3,1-phenylenecarbonylimino(4-fluoro-3,1-phenylene)carbonylimino]]bis-1,3,5-naphthalene trisulfonic acid hexasodium salt); TNP-ATP; tetramethylpyrazine; Ip₅I; $\beta\gamma$ -carboxymethylene ATP; $\beta\gamma$ -chlorophosphomethylene ATP; KN-62 (4-[(2S)-2-[(5-isoquinolinylsulfonyl)methylamino]-3-oxo-3-(4-phenyl-1-piperazinyl)propyl] phenyl isoquinolinesulfonic acid ester); NF023 (8,8'-[carbonylbis(imino-3,1-phenylenecarbonylimino)]bis-1,3,5-naphthalene-trisulphonic acid, hexasodium salt); NF279 (8,8'-[Carbonylbis(imino-4,1-phenylenecarbonylimino-4,1-phenylenecarbonylimino)]bis-1,3,5-naphthalenetrisulfonic acid hexasodium salt); spinorphin; or combinations thereof.

RNAi

[00473] In some embodiments, where inhibition or down-regulation of a target is desired (e.g. genes encoding a component of a potassium channel, genes encoding a purigenic receptor), RNA

interference is optionally utilized. In some embodiments, the agent that inhibits or down-regulates the target is an siRNA molecule. In certain instances, the siRNA molecule is as described herein.

Combination therapy

5 [00474] In certain embodiments, any otic active agent (e.g., an immunomodulator or an auris pressure modulator) is administered in combination with one or more of any other otic active agent described herein. In some embodiments, an otic agent is administered with an anti-emetic agent (e.g., when a balance disorder is accompanied by nausea). In some embodiments, an otic agent is administered in combination with one or more otoprotectant (e.g., when the administration of a
10 cytotoxic agent is accompanied by ototoxicity). In certain embodiments, an otic agent is administered in combination with, for example, an anti-emetic, an antimicrobial agent, a nitric oxide synthase inhibitor, an antioxidant, a neurotransmitter reuptake inhibitor, an otoprotectant, a homeostasis modulator (e.g., ion/ fluid (e.g., water) homeostasis modulator) or the like.

Anti-Emetic Agents/Central Nervous System Agents

15 [00475] Anti-Emetic agents are optionally used in combination with any otic formulations disclosed herein. Anti-emetic agents include antihistamines and central nervous agents, including anti-psychotic agents, barbiturates, benzodiazepines and phenothiazines. Other anti-emetic agents include the serotonin receptor antagonists, which include dolasetron, granisetron, ondansetron, tropisetron, palonosetron, and combinations thereof; dopamine antagonists, including domperidone,
20 properidol, haloperidol, chlorpromazine, promethazine, prochlorperazine and combinations thereof; cannabinoids, including dronabinol, nabilone, sativex, and combinations thereof; anticholinergics, including scopolamine; and steroids, including dexamethasone; trimethobenzamine, emetrol, propofol, muscimol, and combinations thereof.

[00476] Optionally, Central Nervous System agents and barbiturates are useful in the treatment of
25 nausea and vomiting symptoms that accompany an autoimmune otic disorder. When used, an appropriate barbiturate and/or central nervous system agent is selected to relieve or ameliorate specific symptoms without possible side effects, including ototoxicity. Moreover, as discussed above, targeting of the drugs to the round window membrane of the auris interna reduces possible side effects and toxicity caused by systemic administration of these drugs. Barbiturates, which act as
30 a central nervous system depressant, include allobarbitol, alphenal, amobarbital, aprobarbital, barnexaclone, barbital, brallobarbitol, butabarbital, butalbital, butallylonal, butobarbital, corvalol, crotylbarbital, cyclobarbitol, cyclopal, ethallobarbitol, febarbamate, heptabarbitol, hexethal, hexobarbital, metharbital, methohexital, methylphenobarbital, narcobarbital, nealbarbital, pentobarbital, phenobarbital, primidone, probarbital, propallylonal, proxibarbitol, reposal,

secobarbital, sigmodal, sodium thiopental, talbutal, thialbarbital, thiamylal, thiobarbital, thiobutabarbital, tuinal, valofane, vinbarbital, vinylbital, and combinations thereof.

[00477] Other central nervous system agents which are optionally used in conjunction with the otic formulations disclosed herein include benzodiazepines or phenothiazines. Useful benzodiazepines include, but are not limited to diazepam, lorazepam, oxazepam, prazepam, alprazolam, bromazepam, chlordiazepoxide, clonazepam, clorazepate, brotizolam, estazolam, flunitrazepam, flurazepam, loprazolam, lormetazepam, midazolam, nimetazepam, nitrazepam, ternazepam, triazolam, and combinations thereof. Examples of phenothiazines include prochlorperazine, chlorpromazine, promazine, triflupromazine, levopromazine, methotrimeprazine, mesoridazine, thioridazine, fluphenazine, perphenazine, flupentixol, trifluoperazine, and combinations thereof.

[00478] Antihistamines, or histamine antagonists, act to inhibit the release or action of histamine. Antihistamines that target the H₁ receptor are useful in the alleviation or reduction of nausea and vomiting symptoms that are associated with AIED, other autoimmune disorders, as well as anti-inflammatory disorders. Accordingly, some embodiments incorporate the use of agents which modulate histamine receptors (e.g. the H₁ receptor, H₂ receptor, and/or the H₃ receptor).

[00479] Such antihistamines include, but are not limited to, meclizine, diphenhydramine, loratadine and quetiapine. Other antihistamines include mepyramine, piperoxan, antazoline, carbinoxamine, doxylamine, clemastine, dimenhydrinate, pheniramine, chlorphenamine, chlorpheniramine, dexchlorpheniramine, brompheniramine, triprolidine, cyclizine, chlorcyclizine, hydroxyzine, promethazine, alimemazine, trimeprazine, cyproheptadine, azatadine, ketotifen, oxatomide and combinations thereof.

[00480] In some embodiments, the H₁ receptor antagonist is meclizine hydrochloride. In some embodiments, the H₁ receptor antagonist is promethazine hydrochloride. In some embodiments, the H₁ receptor antagonist is dimenhydrinate. In some embodiments, the H₁ receptor antagonist is diphenhydramine. In some embodiments, the H₁ receptor antagonist is cinnarizine. In some embodiments, the H₁ receptor antagonist is hydroxyzine pamoate.

[00481] Antihistamines which target the H₃ receptor include, but are not limited to betahistine dihydrochloride.

Antimicrobial Agents

[00482] Antimicrobial agents are also contemplated as useful with the formulations disclosed herein. In some embodiments, the antimicrobial agent is as described herein.

Corticosteroids

[00483] Contemplated for use in combination with any otic formulation described herein (e.g, aural pressure modulating formulations, immunomodulator formulations described herein) are corticosteroid agents which reduce or ameliorate symptoms or effects as a result of an autoimmune disease and/or inflammatory disorder, including AIED. Such autoimmune response are a

contributing factor to otic disorders such as Meniere's disease. In some embodiments, corticosteroids modulate the degeneration of neurons and/or hair cells of the auris, and agents for treating or ameliorating hearing loss or reduction resulting from destroyed, stunted, malfunctioning, damaged, fragile or missing hairs in the inner ear. Accordingly, some embodiments incorporate the use of agents which protect otic hair cells from ototoxins. In some embodiments, the agent which protects otic hair cells from ototoxins is a corticosteroid. Such steroids include prednisolone, dexamethasone, dexamethasone phosphate, beclomethasone, 21-acetoxypregnenolone, alclometasone, algestone, amcinonide, beclomethasone, betamethasone, budesonide, chloroprednisone, clobetasol, clobetasone, clocortolone, cloprednol, corticosterone, cortisone, cortivazol, deflazacort, desonide, desoximetasone, diflorasone, diflucortolone, difluprednate, enoxolone, fluazacort, flucloronide, flumethasone, flunisolide, fluocinolone acetonide, fluocinonide, fluocortin butyl, fluocortolone, fluorometholone, fluperolone acetate, fluprednidene acetate, fluprednisolone, flurandrenolide, fluticasone propionate, formocortal, halcinonide, halobetasol propionate, halometasone, halopredone acetate, hydrocortamate, hydrocortisone, loteprednol etabonate, mazipredone, medrysone, meprednisone, methylprednisolone, mometasone furoate, paramethasone, prednicarbate, prednisolone, prednisolone 25-diethylamino-acetate, prednisolone sodium phosphate, prednisone, prednival, prednylidene, rimexolone, tixocortol, triamcinolone, triamcinolone acetonide, triamcinolone benetonide, triamcinolone hexacetonide and combinations thereof. In certain instances, triamcinolone actenoides and dexamethasone protect otic hair cells from damage caused by the naturally occurring toxin 4-hydroxy-2,3-nonenal (HNE), which is produced in the inner ear in response to oxidative stress.

Otoprotectants

[00484] In some embodiments, any otic formulation described herein (e.g. auris sensory cell modulating agent formulations disclosed herein) further comprise otoprotectants that reduce, inhibit or ameliorate the ototoxicity of agents such as chemotherapeutic agents and/or antibiotics as described herein, or reduce, inhibit or ameliorate the effects of other environmental factors, including excessive noise and the like. Examples of otoprotectants include, and are not limited to, thiols and/or thiol derivatives and/or pharmaceutically acceptable salts, or derivatives (e.g. prodrugs) thereof (e.g., D-methionine, L-methionine, ethionine, hydroxyl methionine, methioninol, amifostine, mesna (sodium 2-sulfanylethanesulfonate), a mixture of D and L methionine, normethionine, homomethionine, S-adenosyl-L-methionine), diethyldithiocarbamate, ebselen (2-phenyl-1, 2-benzisoselenazol-3(2H)-one), sodium thiosulfate, AM-111 (a cell permeable JNK inhibitor, (Laboratoires Auris SAS)), leucovorin, leucovorin calcium, dexrazoxane, piracetam, Oxiracetam, Aniracetam, Pramiracetam, Phenylpiracetam (Carphedon), Etiracetam, Levetiracetam, Nefiracetam, Nicoracetam, Rolziracetam, Nebracetam, Fasoracetam, Coluracetam, Dimiracetam, Brivaracetam, Seletracetam, Rolipram and or combinations thereof. Otoprotectants allow for the administration of

chemotherapeutic agents and/or antibiotics at doses that are higher than maximal toxic doses; the chemotherapeutic agents and/or antibiotics would otherwise be administered at lower doses due to ototoxicity. Otoprotectants, when optionally administered by itself, also allow for the amelioration, reduction or elimination of the effect of environmental factors that contribute to loss of hearing and attendant effects, including but not limited to noise-induced hearing loss and tinnitus.

[00485] The amount of otoprotectant in any formulation described herein on a mole:mole basis in relation to the ototoxic chemotherapeutic agent (e.g. cis platin) and/or an ototoxic antibiotic (e.g. gentamicin) is in the range of from about 5:1 to about 200:1, from about 5:1 to about 100:1, or from about 5:1 to about 20:1. The amount of otoprotectant in any formulation described herein on a molar basis in relation to the ototoxic chemotherapeutic agent (e.g. cis platin) and/or an ototoxic antibiotic (e.g. gentamicin) is about 50:1, about 20:1 or about 10:1. Any the auris sensory cell modulating agent formulation described herein comprises from about 10 mg/mL to about 50 mg/mL, from about 20 mg/mL to about 30 mg/mL, or from about 25 mg/mL of otoprotectant.

Chemotherapeutic Agents

[00486] Chemotherapeutic agents are also contemplated for use with the formulations disclosed herein. Chemotherapeutic agents act by killing cancer cells or microorganisms, and may include antineoplastic agents that target cancer or malignant cells. Some chemotherapeutic agents, either alone or in combination, are also ototoxic. For example, cisplatin is a known cochleotoxic agent. However, use of cisplatin in combination with antioxidants are protective and lessen the ototoxic effects of the chemotherapeutic agent. Moreover, the localized application of the cytotoxic drug may lessen the ototoxic effects that might otherwise occur through systemic application through the use of lower amounts with maintained efficacy, or the use of targeted amounts for a shorter period of time. Accordingly, a skilled practitioner choosing a course of therapy for tumor growth will have the knowledge to avoid or combine an ototoxic compound, or to vary the amount or course of treatment to avoid or lessen ototoxic effects.

[00487] Chemotherapeutic agents that are used in combination with the formulations disclosed herein include, for example, but are not limited to adriamycin, imidazole carboxamide, cyclophosphamide, mechlorethamine, chlorambucil, melphalan, daunorubicin, doxorubicin, epirubicin, idarubicin, mitoxanthrone, valrubicin, paclitaxel, docetaxel, etoposide, teniposide, tafluposide, azacitidine, azathioprine, capecitabine, cytarabine, doxifluridine, fluorouracil, gemcitabine, mercaptopurine, methotrexate, tioguanine, bleomycin, carboplatin, cisplatin, oxaliplatin, all-trans retinoic acid, vinblastine, vincristine, vindesine, vinorelbine, and combinations thereof.

Homeostasis modulators

[00488] Homeostatis modulators are contemplated as useful with the formulations described herein. Homeostasis modulators include ion and fluid (e.g. water) homeostasis modulators. In some

instances, homeostasis modulators include Na/K-ATPase modulators, ENaC modulators, vasopressin receptor modulators, diuretics or the like as described herein.

Na/K ATPase Modulators

[00489] Na/K-ATPase modulators are contemplated for use with the formulations disclosed herein.

- 5 Cochlear homeostasis is dependent on the electrolyte composition of the endolymph, which is regulated by an active exchange of Na⁺ and K⁺ via a ATPase. Examples of Na/K-ATPase modulators include, and are not limited to, nimodipine (a sodium-potassium adenosine triphosphatase stimulator), ouabain, and furosemide.

- 10 [00490] Presented below (**Table 1**) are examples of active agents contemplated for use with the formulations disclosed herein.

[00491] Active Agents (including pharmaceutically acceptable salts of these active agents) for Use with the Formulations Disclosed Herein

(TABLE 1)

Auris Condition	Therapeutic Agent
Benign Paroxysmal Positional Vertigo	Diphenhydramine
Benign Paroxysmal Positional Vertigo	Lorazepam
Benign Paroxysmal Positional Vertigo	Meclizine
Benign Paroxysmal Positional Vertigo	Oldansetron
Hearing Loss	Estrogen
Hearing Loss	Estrogen and progesterone (E+P)
Hearing Loss	Folic acid
Hearing Loss	Lactated Ringer's with 0.03% Ofloxacin
Hearing Loss	Methotrexate
Hearing Loss	Methylprednisolone sodium succinate
Hearing Loss	N-acetyl cysteine
Meniere's Disease	Betahistine
Meniere's Disease	Sildenafil
Middle Ear Effusion	Pneumococcal vaccine
Otitis Externa	Diclofenac sodium; dexote
Otitis Externa, Acute	AL-15469A/AL-38905
Otitis Media	Amoxicillin/clavulanate
Otitis Media	Amoxycillin
Otitis Media	Chlorpheniramine maleate
Otitis Media	Dornase alfa
Otitis Media	Echinacea purpurea
Otitis Media	Faropenem medoxomil
Otitis Media	Levofloxacin
Otitis Media	PNCRM9
Otitis Media	Pneumococcal vaccine

Auris Condition	Therapeutic Agent
Otitis Media	Telithromycin
Otitis Media	Triamcinolone acetonide
Otitis Media	Zmax
Otitis Media with Effusion	Lansoprazole
Otitis Media, Acute	AL-15469A; AL-38905
Otitis Media, Acute	Amoxicillin
Otitis Media, Acute	Amoxicillin-clavulanate
Otitis Media, Acute	Azithromycin
Otitis Media, Acute	Azithromycin SR
Otitis Media, Acute	Cefdinir
Otitis Media, Acute	Hyland's earache drops
Otitis Media, Acute	Montelukast
Otitis Media, Acute	Pneumococcal vaccine
Otitis Media, Acute with Tympanostomy Tubes	AL-15469A/AL38905
Otitis Media, Chronic	Sulfamethoxazole-trimethoprim
Otitis Media, Suppurative	Azithromycin
Otitis Media, Suppurative	Telithromycin
Otosclerosis	Acetylcysteine
Ototoxicity	Aspirin
Tinnitus	Acamprosate
Tinnitus	Gabapentin
Tinnitus	Modafinil
Tinnitus	Neramexane
Tinnitus	Neramexane mesylate
Tinnitus	Piribedil
Tinnitus	Vardenafil
Tinnitus	Vestipitant + Paroxetine
Tinnitus	Vestiplitant
Tinnitus	Zinc sulfate

Devices

[00492] Also contemplated herein are the use of devices for the delivery of the pharmaceutical formulations disclosed herein, or alternatively for the measurement or surveillance of the function of the auris formulations disclosed herein. For example, in one embodiment pumps, osmotic devices or other means of mechanically delivering pharmaceutical formulations are used for the delivery of the pharmaceutical formulations disclosed herein. Reservoir devices are optionally used with the pharmaceutical drug delivery units, and reside either internally along with the drug delivery unit, or externally of the auris structures.

[00493] Other embodiments contemplate the use of mechanical or imaging devices to monitor or survey the hearing, balance or other auris disorder. For example, magnetic resonance imaging (MRI)

devices are specifically contemplated within the scope of the embodiments, wherein the MRI devices (for example, 3 Tesla MRI devices) are capable of evaluating Meniere Disease progression and subsequent treatment with the pharmaceutical formulations disclosed herein. See, Carfrae et al. Laryngoscope 118:501-505 (March 2008). Whole body scanners, or alternatively cranial scanners, are contemplated, as well as higher resolution (7 Tesla, 8 Tesla, 9.5 Tesla or 11 Tesla for humans) are optionally used in MRI scanning.

General Methods of Sterilization

[00494] Provided herein are otic compositions that ameliorate or lessen otic disorders described herein. Further provided herein are methods comprising the administration of said otic compositions. In some embodiments, the compositions are sterilized. Included within the embodiments disclosed herein are means and processes for sterilization of a pharmaceutical composition disclosed herein for use in humans. The goal is to provide a safe pharmaceutical product, relatively free of infection causing micro-organisms. The U. S. Food and Drug Administration has provided regulatory guidance in the publication "Guidance for Industry: Sterile Drug Products Produced by Aseptic Processing" available at: <http://www.fda.gov/cder/guidance/5882fnl.htm>, which is incorporated herein by reference in its entirety. No specific guidelines are available for safe pharmaceutical products for treatment of the inner ear.

[00495] As used herein, sterilization means a process used to destroy or remove microorganisms that are present in a product or packaging. Any suitable method available for sterilization of objects and compositions is used. Available methods for the inactivation of microorganisms include, but are not limited to, the application of extreme heat, lethal chemicals, or gamma radiation. In some embodiments is a process for the preparation of an otic therapeutic formulation comprising subjecting the formulation to a sterilization method selected from heat sterilization, chemical sterilization, radiation sterilization or filtration sterilization. The method used depends largely upon the nature of the device or composition to be sterilized. Detailed descriptions of many methods of sterilization are given in Chapter 40 of Remington: The Science and Practice of Pharmacy published by Lippincott, Williams & Wilkins, and is incorporated by reference with respect to this subject matter.

Sterilization by Heat

[00496] Many methods are available for sterilization by the application of extreme heat. One method is through the use of a saturated steam autoclave. In this method, saturated steam at a temperature of at least 121 °C is allowed to contact the object to be sterilized. The transfer of heat is either directly to the microorganism, in the case of an object to be sterilized, or indirectly to the microorganism by heating the bulk of an aqueous solution to be sterilized. This method is widely practiced as it allows flexibility, safety and economy in the sterilization process.

[00497] Dry heat sterilization is a method which is used to kill microorganisms and perform depyrogenation at elevated temperatures. This process takes place in an apparatus suitable for heating HEPA-filtered microorganism-free air to temperatures of at least 130-180 °C for the sterilization process and to temperatures of at least 230-250 °C for the depyrogenation process.

5 Water to reconstitute concentrated or powdered formulations is also sterilized by autoclave.

Chemical Sterilization

[00498] Chemical sterilization methods are an alternative for products that do not withstand the extremes of heat sterilization. In this method, a variety of gases and vapors with germicidal properties, such as ethylene oxide, chlorine dioxide, formaldehyde or ozone are used as the anti-
10 apoptotic agents. The germicidal activity of ethylene oxide, for example, arises from its ability to serve as a reactive alkylating agent. Thus, the sterilization process requires the ethylene oxide vapors to make direct contact with the product to be sterilized.

Radiation Sterilization

[00499] One advantage of radiation sterilization is the ability to sterilize many types of products
15 without heat degradation or other damage. The radiation commonly employed is beta radiation or alternatively, gamma radiation from a ⁶⁰Co source. The penetrating ability of gamma radiation allows its use in the sterilization of many product types, including solutions, compositions and heterogeneous mixtures. The germicidal effects of irradiation arise from the interaction of gamma radiation with biological macromolecules. This interaction generates charged species and free
20 radicals. Subsequent chemical reactions, such as rearrangements and cross-linking processes, result in the loss of normal function for these biological macromolecules. The formulations described herein are also optionally sterilized using beta irradiation.

Filtration

[00500] Filtration sterilization is a method used to remove but not destroy microorganisms from
25 solutions. Membrane filters are used to filter heat-sensitive solutions. Such filters are thin, strong, homogenous polymers of mixed cellulosic esters (MCE), polyvinylidene fluoride (PVF; also known as PVDF), or polytetrafluoroethylene (PTFE) and have pore sizes ranging from 0.1 to 0.22 µm. Solutions of various characteristics are optionally filtered using different filter membranes. For example, PVF and PTFE membranes are well suited to filtering organic solvents while aqueous
30 solutions are filtered through PVF or MCE membranes. Filter apparatus are available for use on many scales ranging from the single point-of-use disposable filter attached to a syringe up to commercial scale filters for use in manufacturing plants. The membrane filters are sterilized by autoclave or chemical sterilization. Validation of membrane filtration systems is performed following standardized protocols (Microbiological Evaluation of Filters for Sterilizing Liquids, Vol
35 4, No. 3. Washington, D.C: Health Industry Manufacturers Association, 1981) and involve

challenging the membrane filter with a known quantity (ca. $10^7/\text{cm}^2$) of unusually small microorganisms, such as *Brevundimonas diminuta* (ATCC 19146).

[00501] Pharmaceutical compositions are optionally sterilized by passing through membrane filters. Formulations comprising nanoparticles (U.S. Pat No. 6,139,870) or multilamellar vesicles (Richard et al., International Journal of Pharmaceutics (2006), 312(1-2):144-50) are amenable to sterilization by filtration through 0.22 μm filters without destroying their organized structure.

[00502] In some embodiments, the methods disclosed herein comprise sterilizing the formulation (or components thereof) by means of filtration sterilization. In another embodiment the auris-acceptable otic therapeutic agent formulation comprises a particle wherein the particle formulation is suitable for filtration sterilization. In a further embodiment said particle formulation comprises particles of less than 300 nm in size, of less than 200 nm in size, of less than 100 nm in size. In another embodiment the auris-acceptable formulation comprises a particle formulation wherein the sterility of the particle is ensured by sterile filtration of the precursor component solutions. In another embodiment the auris-acceptable formulation comprises a particle formulation wherein the sterility of the particle formulation is ensured by low temperature sterile filtration. In a further embodiment, said low temperature sterile filtration occurs at a temperature between 0 and 30 °C, or between 0 and 20 °C, or between 0 and 10 °C, or between 10 and 20 °C, or between 20 and 30 °C. In another embodiment is a process for the preparation of an auris-acceptable particle formulation comprising: filtering the aqueous solution containing the particle formulation at low temperature through a sterilization filter; lyophilizing the sterile solution; and reconstituting the particle formulation with sterile water prior to administration.

[00503] In specific embodiments, filtration and/or filling procedures are carried out at about 5°C below the gel temperature (T_{gel}) of a formulation described herein and with viscosity below a theoretical value of 100cP to allow for filtration in a reasonable time using a peristaltic pump.

[00504] In another embodiment the auris-acceptable otic therapeutic agent formulation comprises a nanoparticle formulation wherein the nanoparticle formulation is suitable for filtration sterilization. In a further embodiment the nanoparticle formulation comprises nanoparticles of less than 300 nm in size, of less than 200 nm in size, or of less than 100 nm in size. In another embodiment the auris-acceptable formulation comprises a microsphere formulation wherein the sterility of the microsphere is ensured by sterile filtration of the precursor organic solution and aqueous solutions. In another embodiment the auris-acceptable formulation comprises a thermoreversible gel formulation wherein the sterility of the gel formulation is ensured by low temperature sterile filtration. In a further embodiment, the low temperature sterile filtration occurs at a temperature between 0 and 30 °C, or between 0 and 20 °C, or between 0 and 10 °C, or between 10 and 20 °C, or between 20 and 30 °C. In another embodiment is a process for the preparation of an auris-acceptable thermoreversible gel formulation comprising: filtering the aqueous solution containing the thermoreversible gel

components at low temperature through a sterilization filter; lyophilizing the sterile solution; and reconstituting the thermoreversible gel formulation with sterile water prior to administration.

[00505] In certain embodiments, the active ingredients are dissolved in a suitable vehicle (e.g. a buffer) and sterilized separately (e.g. by heat treatment, filtration, gamma radiation); the remaining excipients (e.g., fluid gel components present in auris formulations) are sterilized in a separate step by a suitable method (e.g. filtration and/or irradiation of a cooled mixture of excipients); the two solutions that were separately sterilized are then mixed aseptically to provide a final auris formulation.

[00506] In some instances, conventionally used methods of sterilization (e.g., heat treatment (e.g., in an autoclave), gamma irradiation, filtration) lead to irreversible degradation of polymeric components (e.g., thermosetting, gelling or mucoadhesive polymer components) and/or the active agent in the formulation. In some instances, sterilization of an auris formulation by filtration through membranes (e.g., 0.2 μ M membranes) is not possible if the formulation comprises thixotropic polymers that gel during the process of filtration.

[00507] Accordingly, provided herein are methods for sterilization of auris formulations that prevent degradation of polymeric components (e.g., thermosetting and/or gelling and/or mucoadhesive polymer components) and/or the active agent during the process of sterilization. In some embodiments, degradation of the active agent (e.g., any therapeutic otic agent described herein) is reduced or eliminated through the use of specific pH ranges for buffer components and specific proportions of gelling agents in the formulations. In some embodiments, the choice of an appropriate gelling agent and/or thermosetting polymer allows for sterilization of formulations described herein by filtration. In some embodiments, the use of an appropriate thermosetting polymer and an appropriate copolymer (e.g., a gelling agent) in combination with a specific pH range for the formulation allows for high temperature sterilization of formulations described with substantially no degradation of the therapeutic agent or the polymeric excipients. An advantage of the methods of sterilization provided herein is that, in certain instances, the formulations are subjected to terminal sterilization via autoclaving without any loss of the active agent and/or excipients and/or polymeric components during the sterilization step and are rendered substantially free of microbes and/or pyrogens.

Microorganisms

[00508] Provided herein are auris-acceptable compositions that ameliorate or lessen otic disorders described herein. Further provided herein are methods comprising the administration of said otic compositions. In some embodiments, the compositions are substantially free of microorganisms. Acceptable sterility levels are based on applicable standards that define therapeutically acceptable otic compositions, including but not limited to United States Pharmacopeia Chapters <1111> et seq. For example, acceptable sterility levels include 10 colony forming units (cfu) per gram of

formulation, 50 cfu per gram of formulation, 100 cfu per gram of formulation, 500 cfu per gram of formulation or 1000 cfu per gram of formulation. In addition, acceptable sterility levels include the exclusion of specified objectionable microbiological agents. By way of example, specified objectionable microbiological agents include but are not limited to *Escherichia coli* (*E. coli*),

5 *Salmonella* sp., *Pseudomonas aeruginosa* (*P. aeruginosa*) and/or other specific microbial agents.

[00509] Sterility of the auris-acceptable otic therapeutic agent formulation is confirmed through a sterility assurance program in accordance with United States Pharmacopeia Chapters <61>, <62> and <71>. A key component of the sterility assurance quality control, quality assurance and validation process is the method of sterility testing. Sterility testing, by way of example only, is

10 performed by two methods. The first is direct inoculation wherein a sample of the composition to be tested is added to growth medium and incubated for a period of time up to 21 days. Turbidity of the growth medium indicates contamination. Drawbacks to this method include the small sampling size of bulk materials which reduces sensitivity, and detection of microorganism growth based on a visual observation. An alternative method is membrane filtration sterility testing. In this method, a volume of product is passed through a small membrane filter paper. The filter paper is then placed into media to promote the growth of microorganisms. This method has the advantage of greater sensitivity as the entire bulk product is sampled. The commercially available Millipore Steritest sterility testing system is optionally used for determinations by membrane filtration sterility testing. For the filtration testing of creams or ointments Steritest filter system No. TLHVSL210 are used.

20 For the filtration testing of emulsions or viscous products Steritest filter system No. TLAREM210 or TDAREM210 are used. For the filtration testing of pre-filled syringes Steritest filter system No. TTHASY210 are used. For the filtration testing of material dispensed as an aerosol or foam Steritest filter system No. TTHVA210 are used. For the filtration testing of soluble powders in ampoules or vials Steritest filter system No. TTHADA210 or TTHADV210 are used.

25 [00510] Testing for *E. coli* and *Salmonella* includes the use of lactose broths incubated at 30 – 35 °C for 24-72 hours, incubation in MacConkey and/or EMB agars for 18-24 hours, and/or the use of Rappaport medium. Testing for the detection of *P. aeruginosa* includes the use of NAC agar. United States Pharmacopeia Chapter <62> further enumerates testing procedures for specified objectionable microorganisms.

30 [00511] In certain embodiments, any controlled release formulation described herein has less than about 60 colony forming units (CFU), less than about 50 colony forming units, less than about 40 colony forming units, or less than about 30 colony forming units of microbial agents per gram of formulation. In certain embodiments, the otic formulations described herein are formulated to be isotonic with the endolymph and/or the perilymph.

35 *Endotoxins*

[00512] Provided herein are otic compositions that ameliorate or lessen otic disorders described herein. Further provided herein are methods comprising the administration of said otic compositions. In some embodiments, the compositions are substantially free of endotoxins. An additional aspect of the sterilization process is the removal of by-products from the killing of microorganisms

5 (hereinafter, "Product"). The process of depyrogenation removes pyrogens from the sample. Pyrogens are endotoxins or exotoxins which induce an immune response. An example of an endotoxin is the lipopolysaccharide (LPS) molecule found in the cell wall of gram-negative bacteria. While sterilization procedures such as autoclaving or treatment with ethylene oxide kill the bacteria, the LPS residue induces a proinflammatory immune response, such as septic shock. Because the
10 molecular size of endotoxins can vary widely, the presence of endotoxins is expressed in "endotoxin units" (EU). One EU is equivalent to 100 picograms of E. coli LPS. Humans can develop a response to as little as 5 EU/kg of body weight. The sterility is expressed in any units as recognized in the art. In certain embodiments, otic compositions described herein contain lower endotoxin levels (e.g. < 4 EU/kg of body weight of a subject) when compared to conventionally acceptable endotoxin levels
15 (e.g., 5 EU/kg of body weight of a subject). In some embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 5 EU/kg of body weight of a subject. In other embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 4 EU/kg of body weight of a subject. In additional embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 3 EU/kg of body weight of a subject. In additional embodiments, the
20 auris-acceptable otic therapeutic agent formulation has less than about 2 EU/kg of body weight of a subject.

[00513] In some embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 5 EU/kg of formulation. In other embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 4 EU/kg of formulation. In additional embodiments, the auris-
25 acceptable otic therapeutic agent formulation has less than about 3 EU/kg of formulation. In some embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 5 EU/kg Product. In other embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 1 EU/kg Product. In additional embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 0.2 EU/kg Product. In some embodiments, the auris-acceptable otic
30 therapeutic agent formulation has less than about 5 EU/g of unit or Product. In other embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 4 EU/ g of unit or Product. In additional embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 3 EU/g of unit or Product. In some embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 5 EU/mg of unit or Product. In other embodiments, the auris-
35 acceptable otic therapeutic agent formulation has less than about 4 EU/ mg of unit or Product. In additional embodiments, the auris-acceptable otic therapeutic agent formulation has less than about

3 EU/mg of unit or Product. In certain embodiments, otic compositions described herein contain from about 1 to about 5 EU/mL of formulation. In certain embodiments, otic compositions described herein contain from about 2 to about 5 EU/mL of formulation, from about 3 to about 5 EU/mL of formulation, or from about 4 to about 5 EU/mL of formulation.

5 [00514] In certain embodiments, otic compositions described herein contain lower endotoxin levels (e.g. < 0.5 EU/mL of formulation) when compared to conventionally acceptable endotoxin levels (e.g., 0.5 EU/mL of formulation). In some embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 0.5 EU/mL of formulation. In other embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 0.4 EU/mL of formulation. In
10 additional embodiments, the auris-acceptable otic therapeutic agent formulation has less than about 0.2 EU/mL of formulation.

[00515] Pyrogen detection, by way of example only, is performed by several methods. Suitable tests for sterility include tests described in United States Pharmacopoeia (USP) <71> Sterility Tests (23rd edition, 1995). The rabbit pyrogen test and the Limulus amoebocyte lysate test are both specified in
15 the United States Pharmacopoeia Chapters <85> and <151> (USP23/NF 18, Biological Tests, The United States Pharmacopoeial Convention, Rockville, MD, 1995). Alternative pyrogen assays have been developed based upon the monocyte activation-cytokine assay. Uniform cell lines suitable for quality control applications have been developed and have demonstrated the ability to detect pyrogenicity in samples that have passed the rabbit pyrogen test and the Limulus amoebocyte lysate
20 test (Taktak et al, J. Pharm. Pharmacol. (1990), 43:578-82). In an additional embodiment, the auris-acceptable otic therapeutic agent formulation is subject to depyrogenation. In a further embodiment, the process for the manufacture of the auris-acceptable otic therapeutic agent formulation comprises testing the formulation for pyrogenicity. In certain embodiments, the formulations described herein are substantially free of pyrogens.

25 pH and Osmolarity

[00516] The main cation present in the endolymph is potassium. In addition the endolymph has a high concentration of positively charged amino acids. The main cation present in the perilymph is sodium. In certain instances, the ionic composition of the endolymph and perilymph regulate the electrochemical impulses of hair cells. In certain instances, any change in the ionic balance of the
30 endolymph or perilymph results in a loss of hearing due to changes in the conduction of electrochemical impulses along otic hair cells. In some embodiments, a composition disclosed herein does not disrupt the ionic balance of the perilymph. In some embodiments, a composition disclosed herein has an ionic balance that is the same as or substantially the same as the perilymph. In some embodiments, a composition disclosed herein does not disrupt the ionic balance of the
35 endolymph. In some embodiments, a composition disclosed herein has an ionic balance that is the same as or substantially the same as the endolymph. In some embodiments, an otic formulation

described herein is formulated to provide an ionic balance that is compatible with inner ear fluids (i.e., endolymph and/or perilymph).

[00517] The endolymph and the perilymph have a pH that is close to the physiological pH of blood. The endolymph has a pH range of about 7.2-7.9; the perilymph has a pH range of about 7.2 – 7.4.

5 The *in situ* pH of the proximal endolymph is about 7.4 while the pH of distal endolymph is about 7.9.

[00518] In some embodiments, the pH of a composition described herein is adjusted (e.g., by use of a buffer) to an endolymph-compatible pH range of about 7.0 to 8.0, and a preferred pH range of about 7.2 – 7.9. In some embodiments, the pH of the auris formulations described herein is adjusted
10 (e.g., by use of a buffer) to a perilymph-compatible pH of about 7.0 – 7.6, and a preferred pH range of about 7.2-7.4.

[00519] In some embodiments, useful formulations also include one or more pH adjusting agents or buffering agents. Suitable pH adjusting agents or buffers include, but are not limited to acetate, bicarbonate, ammonium chloride, citrate, phosphate, pharmaceutically acceptable salts thereof and
15 combinations or mixtures thereof.

[00520] In one embodiment, when one or more buffers are utilized in the formulations of the present disclosure, they are combined, e.g., with a pharmaceutically acceptable vehicle and are present in the final formulation, e.g., in an amount ranging from about 0.1% to about 20%, from about 0.5% to about 10%. In certain embodiments of the present disclosure, the amount of buffer included in the
20 gel formulations are an amount such that the pH of the gel formulation does not interfere with the body's natural buffering system. In some embodiments, from about 5 mM to about 200 mM concentration of a buffer is present in the gel formulation. In certain embodiments, from about a 20 mM to about a 100 mM concentration of a buffer is present. In other embodiments, the concentration of buffer is such that a pH of the formulation is between 3 and 9, between 5 and 8, or
25 alternatively between 6 and 7. In other embodiments, the pH of the gel formulation is about 7. In one embodiment is a buffer such as acetate or citrate at slightly acidic pH. In one embodiment the buffer is a sodium acetate buffer having a pH of about 4.5 to about 6.5. In another embodiment the buffer is a sodium acetate buffer having a pH of about 5.5 to about 6.0. In a further embodiment the buffer is a sodium acetate buffer having a pH of about 6.0 to about 6.5. In one embodiment the
30 buffer is a sodium citrate buffer having a pH of about 5.0 to about 8.0. In another embodiment the buffer is a sodium citrate buffer having a pH of about 5.5 to about 7.0. In one embodiment the buffer is a sodium citrate buffer having a pH of about 6.0 to about 6.5.

[00521] In some embodiments, the concentration of buffer is such that a pH of the formulation is between 6 and 9, between 6 and 8, between 6 and 7.6, between 7 and 8. In other embodiments, the
35 pH of the gel formulation is about 6.0, about 6.5, about 7 or about 7.5. In one embodiment is a buffer such as tris(hydroxymethyl)aminomethane, bicarbonate, carbonate or phosphate at slightly

basic pH. In one embodiment, the buffer is a sodium bicarbonate buffer having a pH of about 7.5 to about 8.5. In another embodiment the buffer is a sodium bicarbonate buffer having a pH of about 7.0 to about 8.0. In a further embodiment the buffer is a sodium bicarbonate buffer having a pH of about 6.5 to about 7.0. In one embodiment the buffer is a sodium phosphate dibasic buffer having a pH of about 6.0 to about 9.0. In another embodiment the buffer is a sodium phosphate dibasic buffer having a pH of about 7.0 to about 8.5. In one embodiment the buffer is a sodium phosphate dibasic buffer having a pH of about 7.5 to about 8.0.

[00522] In one embodiment, diluents are also used to stabilize compounds because they can provide a more stable environment. Salts dissolved in buffered solutions (which also can provide pH control or maintenance) are utilized as diluents in the art, including, but not limited to a phosphate buffered saline solution.

[00523] In a specific embodiment the pH of a composition described herein is between about between about 6.0 and about 7.6, between 7 and about 7.8, between about 7.0 and about 7.6, between about 7.2 and about 7.6, or between about 7.2 and about 7.4. In certain embodiments the pH of a composition described herein is about about 6.0, about 6.5, about 7.0, about 7.1, about 7.2, about 7.3, about 7.4, about 7.5, or about 7.6. In some embodiments, the pH of any formulation described herein is designed to be compatible with the targeted otic structure (e.g., endolymph, perilymph or the like).

[00524] In some embodiments, any gel formulation described herein has a pH that allows for sterilization (e.g. by filtration or aseptic mixing or heat treatment and/or autoclaving (e.g., terminal sterilization)) of a gel formulation without degradation of the otic agent or the polymers comprising the gel. In order to reduce hydrolysis and/or degradation of the otic agent and/or the gel polymer during sterilization, the buffer pH is designed to maintain pH of the formulation in the 7-8 range during the process of sterilization.

[00525] In specific embodiments, any gel formulation described herein has a pH that allows for terminal sterilization (e.g. by heat treatment and/or autoclaving) of a gel formulation without degradation of the otic agent or the polymers comprising the gel. For example, in order to reduce hydrolysis and/or degradation of the otic agent and/or the gel polymer during autoclaving, the buffer pH is designed to maintain pH of the formulation in the 7-8 range at elevated temperatures. Any appropriate buffer is used depending on the otic agent used in the formulation. In some instances, since pK_a of TRIS decreases as temperature increases at approximately $-0.03/^{\circ}\text{C}$ and pK_a of PBS increases as temperature increases at approximately $0.003/^{\circ}\text{C}$, autoclaving at 250°F (121°C) results in a significant downward pH shift (i.e. more acidic) in the TRIS buffer whereas a relatively much less upward pH shift in the PBS buffer and therefore much increased hydrolysis and/or degradation of an otic agent in TRIS than in PBS. Degradation of an otic agent is reduced by the use of an appropriate combination of a buffer and polymeric additives (e.g. P407, CMC) as described herein.

[00526] In some embodiments, a pH of between about 6.0 and about 7.6, between about 7 and about 7.8, between about 7.0 and about 7.6, between about 7.2 and 7.6, between about 7.2 and about 7.4 is suitable for sterilization (e.g., by filtration or aseptic mixing or heat treatment and/or autoclaving (e.g., terminal sterilization)) of auris formulations described herein. In specific
5 embodiments a formulation pH of about 6.0, about 6.5, about 7.0, about 7.1, about 7.2, about 7.3, about 7.4, about 7.5, or about 7.6 is suitable for sterilization (e.g., by filtration or aseptic mixing or heat treatment and/or autoclaving (e.g., terminal sterilization)) of any composition described herein.

[00527] In some embodiments, the formulations described herein have a pH between about 3 and about 9, or between about 4 and 8, or between about 5 and 8, or between about 6 and about 7, or
10 between about 6.5 and about 7, or between about 5.5 and about 7.5, or between about 7.1 and about 7.7, and have a concentration of active pharmaceutical ingredient between about 0.1 mM and about 100 mM. In some embodiments, the formulations described herein have a pH between about 5 and about 8, or between about 6 and about 7, or between about 6.5 and about 7, or between about 5.5 and about 7.5, or between about 7.1 and about 7.7, and have a concentration of active

15 pharmaceutical ingredient between about 1 and about 100 mM. In some embodiments, the formulations described herein have a pH between about 5 and about 8, or between about 6 and about 7, or between about 6.5 and about 7, or between about 5.5 and about 7.5, or between about 7.1 and about 7.7, and have a concentration of active pharmaceutical ingredient between about 50 and about 80 mM. In some embodiments, the concentration of active pharmaceutical ingredient between about
20 10 and about 100 mM. In other embodiments, the concentration of active pharmaceutical ingredient between about 20 and about 80 mM. In additional embodiments, the concentration of active pharmaceutical ingredient between about 10 and about 50 mM.

[00528] In some embodiments, the formulations have a pH as described herein, and include a thickening agent (i.e., a viscosity enhancing agent) such as, by way of non-limiting example, a
25 cellulose based thickening agent described herein. In some instances, the addition of a secondary polymer (e.g., a thickening agent) and a pH of formulation as described herein, allows for sterilization of a formulation described herein without any substantial degradation of the otic agent and/or the polymer components in the otic formulation. In some embodiments, the ratio of a thermoreversible poloxamer to a thickening agent in a formulation that has a pH as described herein,
30 is about 40:1, about 35:1, about 30:1, about 25:1, about 20:1, about 15:1 or about 10:1. For example, in certain embodiments, a sustained and/or extended release formulation described herein comprises a combination of poloxamer 407 (pluronic F127) and carboxymethylcellulose (CMC) in a ratio of about 40:1, about 35:1, about 30:1, about 25:1, about 20:1, about 15:1 or about 10:1. In some
35 embodiments, the amount of thermoreversible polymer in any formulation described herein is about 10%, about 15%, about 20%, about 25%, about 30%, or about 35% of the total weight of the formulation. In some embodiments, the amount of thermoreversible polymer in any formulation

described herein is about 14%, about 15%, about 16%, about 17%, about 18%, about 19%, about 20%, about 21%, about 22%, about 23%, about 24% or about 25% of the total weight of the formulation. In some embodiments, the amount of thickening agent (e.g., a gelling agent) in any formulation described herein is about 1%, 5%, about 10%, or about 15% of the total weight of the formulation. In some embodiments, the amount of thickening agent (e.g., a gelling agent) in any formulation described herein is about 0.5%, about 1%, about 1.5%, about 2%, about 2.5%, about 3%, about 3.5%, about 4%, about 4.5%, or about 5% of the total weight of the formulation.

[00529] In some embodiments, the pharmaceutical formulations described herein are stable with respect to pH over a period of any of at least about 1 day, at least about 2 days, at least about 3 days, at least about 4 days, at least about 5 days, at least about 6 days, at least about 1 week, at least about 2 weeks, at least about 3 weeks, at least about 4 weeks, at least about 5 weeks, at least about 6 weeks, at least about 7 weeks, at least about 8 weeks, at least about 1 month, at least about 2 months, at least about 3 months, at least about 4 months, at least about 5 months, or at least about 6 months. In other embodiments, the formulations described herein are stable with respect to pH over a period of at least about 1 week. Also described herein are formulations that are stable with respect to pH over a period of at least about 1 month.

Tonicity Agents

[00530] In general, the endolymph has a higher osmolality than the perilymph. For example, the endolymph has an osmolality of about 304 mOsm/kg H₂O while the perilymph has an osmolality of about 294 mOsm/kg H₂O. In some embodiments, auris compositions described herein are formulated to provide an osmolality of about 250 to about 320 mM (osmolality of about 250 to about 320 mOsm/kg H₂O); and preferably about 270 to about 320 mM (osmolality of about 270 to about 320 mOsm/kg H₂O). In specific embodiments, osmolality/osmolality of the present formulations is adjusted, for example, by the use of appropriate salt concentrations (e.g., concentration of potassium salts) or the use of tonicity agents which renders the formulations endolymph-compatible and/or perilymph-compatible (i.e. isotonic with the endolymph and/or perilymph. In some instances, the endolymph-compatible and/or perilymph-compatible formulations described herein cause minimal disturbance to the environment of the inner ear and cause minimum discomfort (e.g., vertigo and/or nausea) to a mammal upon administration.

[00531] In some embodiments, any formulation described herein is isotonic with the perilymph. Isotonic formulations are provided by the addition of a tonicity agent. Suitable tonicity agents include, but are not limited to any pharmaceutically acceptable sugar, salt or any combinations or mixtures thereof, such as, but not limited to dextrose, glycerin, mannitol, sorbitol, sodium chloride, and other electrolytes.

[00532] Useful auris compositions include one or more salts in an amount required to bring osmolality of the composition into an acceptable range. Such salts include those having sodium,

potassium or ammonium cations and chloride, citrate, ascorbate, borate, phosphate, bicarbonate, sulfate, thiosulfate or bisulfite anions; suitable salts include sodium chloride, potassium chloride, sodium thiosulfate, sodium bisulfite and ammonium sulfate.

[00533] In further embodiments, the tonicity agents are present in an amount as to provide a final osmolality of an otic formulation of about 100 mOsm/kg to about 500 mOsm/kg, from about 200 mOsm/kg to about 400 mOsm/kg, from about 250 mOsm/kg to about 350 mOsm/kg or from about 280 mOsm/kg to about 320 mOsm/kg. In some embodiments, the formulations described herein have a osmolality of about 100 mOsm/L to about 500 mOsm/L, about 200 mOsm/L to about 400 mOsm/L, about 250 mOsm/L to about 350 mOsm/L, or about 280 mOsm/L to about 320 mOsm/L.

In some embodiments, the osmolality of any formulation described herein is designed to be isotonic with the targeted otic structure (e.g., endolymph, perilymph or the like).

[00534] In some embodiments, the formulations described herein have a pH and osmolality as described herein, and have a concentration of active pharmaceutical ingredient between about 1 μ M and about 10 μ M, between about 1 mM and about 100 mM, between about 0.1 mM and about 100 mM, between about 0.1 mM and about 100 nM. In some embodiments, the formulations described herein have a pH and osmolality as described herein, and have a concentration of active pharmaceutical ingredient between about 0.2 -- about 20%, between about 0.2 -- about 10%,

between about 0.2 -- about 7.5%, between about 0.2 -- 5%, between about 0.2 -- about 3%, between about 0.1 -- about 2% of the active ingredient by weight of the formulation. In some embodiments, the formulations described herein have a pH and osmolality as described herein, and have a concentration of active pharmaceutical ingredient between about 0.1 -- about 70 mg/mL, between about 1 mg -- about 70 mg/mL, between about 1 mg -- about 50 mg/mL, between about 1 mg/mL and about 20 mg/mL, between about 1 mg/mL to about 10 mg/mL, between about 1 mg/mL to about 5 mg/mL, or between about 0.5 mg/mL to about 5 mg/mL of the active agent by volume of the formulation.

Particle size

[00535] Size reduction is used to increase surface area and/or modulate formulation dissolution properties. It is also used to maintain a consistent average particle size distribution (PSD) (e.g., micrometer-sized particles, nanometer-sized particles or the like) for any formulation described herein. In some instances, any formulation described herein comprises multiparticulates, i.e., a plurality of particle sizes (e.g., micronized particles, nano-sized particles, non-sized particles); i.e., the formulation is a multiparticulate formulation. In some embodiments, any formulation described herein comprises one or more multiparticulate (e.g., micronized) therapeutic agents. Micronization is a process of reducing the average diameter of particles of a solid material. Micronized particles are from about micrometer-sized in diameter to about picometer -- sized in diameter. In some

embodiments, the use of multiparticulates (e.g., micronized particles) of an otic agent allows for extended and/or sustained release of the otic agent from any formulation described herein compared to a formulation comprising non-multiparticulate (e.g., non-micronized) otic agent. In some instances, formulations containing multiparticulate (e.g., micronized) otic agents are ejected from a 1 mL syringe adapted with a 27G needle without any plugging or clogging.

[00536] In some instances, any particle in any formulation described herein is a coated particle (e.g., a coated micronized particle) and/or a microsphere and/or a liposomal particle. Particle size reduction techniques include, by way of example, grinding, milling (e.g., air-attrition milling (jet milling), ball milling), coacervation, high pressure homogenization, spray drying and/or supercritical fluid crystallization. In some instances, particles are sized by mechanical impact (e.g., by hammer mills, ball mill and/or pin mills). In some instances, particles are sized via fluid energy (e.g., by spiral jet mills, loop jet mills, and/or fluidized bed jet mills). In some embodiments formulations described herein comprise crystalline particles. In some embodiments, formulations described herein comprise amorphous particles. In some embodiments, formulations described herein comprise therapeutic agent particles wherein the therapeutic agent is a free base, or a salt, or a prodrug of a therapeutic agent, or any combination thereof.

[00537] In some instances, a combination of an otic agent and a salt of the otic agent is used to prepare pulsed release otic agent formulations using the procedures described herein. In some formulations, a combination of a micronized otic agent (and/or salt or prodrug thereof) and coated particles (e.g., nanoparticles, liposomes, microspheres) is used to prepare pulsed release otic agent formulations using any procedure described herein. Alternatively, a pulsed release profile is achieved by solubilizing up to 20% of the delivered dose of the otic agent (e.g., micronized otic agent, or free base or salt or prodrug thereof; multiparticulate otic agent, or free base or salt or prodrug thereof) with the aid of cyclodextrins, surfactants (e.g., poloxamers (407, 338, 188), tween (80, 60, 20,81), PEG-hydrogenated castor oil, cosolvents like N-methyl-2-Pyrrolidone or the like and preparing pulsed release formulations using any procedure described herein.

[00538] In some specific embodiments, any otic formulation described herein comprises one or more micronized otic agents. In some of such embodiments, a micronized otic agent comprises micronized particles, coated (e.g., with an extended release coat) micronized particles, or a combination thereof. In some of such embodiments, a micronized otic agent comprising micronized particles, coated micronized particles, or a combination thereof, comprises an otic agent as a free base, a salt, a prodrug or any combination thereof.

Controlled Release Otic Formulations

[00539] In certain embodiments, any controlled release otic formulation described herein increases the exposure of an otic agent and increases the Area Under the Curve (AUC) in otic fluids (e.g., endolymph and/or perilymph) by about 30%, about 40%, about 50%, about 60%, about 70%, about

80% or about 90% compared to a formulation that is not a controlled release otic formulation. In certain embodiments, any controlled release otic formulation described herein increases the exposure of an otic agent and decreases the C_{max} in otic fluids (e.g., endolymph and/or perilymph) by about 40%, about 30%, about 20%, or about 10%, compared to a formulation that is not a controlled release otic formulation. In certain embodiments, any controlled release otic formulation described herein alters (e.g. reduces) the ratio of C_{max} to C_{min} compared to a formulation that is not a controlled release otic formulation. In certain embodiments, the ratio of C_{max} to C_{min} is 10:1, 9:1, 8:1, 7:1, 6:1, 5:1, 4:1, 3:1, 2:1 or 1:1. In certain embodiments, any controlled release otic formulation described herein increases the exposure of an otic agent and increases the length of time that the concentration of an otic agent is above C_{min} by about 30%, about 40%, about 50%, about 60%, about 70%, about 80% or about 90% compared to a formulation that is not a controlled release otic formulation. In certain instances, controlled release formulations described herein delay the time to C_{max} . In certain instances, the controlled steady release of a drug prolongs the time the concentration of the drug will stay above the C_{min} . In some embodiments, auris compositions described herein prolong the residence time of a drug in the inner ear. In certain instances, once drug exposure (e.g., concentration in the endolymph or perilymph) of a drug reaches steady state, the concentration of the drug in the endolymph or perilymph stays at or about the therapeutic dose for an extended period of time (e.g., one day, 2 days, 3 days, 4 days, 5 days, 6 days, or 1 week).

[00540] The otic formulations described herein deliver an active agent to the external, middle and/or inner ear, including the cochlea and vestibular labyrinth. Local otic delivery of the auris compositions described herein allows for controlled release of active agents to auris structures and overcomes the drawbacks associated with systemic administration (e.g. low bioavailability of the drug in the endolymph or perilymph, variability in concentration of the drug in the middle and/or internal ear).

[00541] Controlled-release options include gel formulations, liposomes, cyclodextrins, biodegradable polymers, dispersable polymers, emulsions, microspheres or microparticles, hydrogels (e.g., a self-assembling hydrogel displaying thixotropic properties that also functions as an absorption enhancer; including instances in which the penetration enhancer is a surfactant comprising an alkyl-glycoside and/or a saccharide alkyl ester), other viscous media, paints, foams, in situ forming spongy materials, xerogels, actinic radiation curable gels, liposomes, solvent release gels, nanocapsules or nanospheres, and combinations thereof; other options or components include mucoadhesives, penetration enhancers, bioadhesives, antioxidants, surfactants, buffering agents, diluents, salts and preservatives. To the extent viscosity considerations potentially limit the use of a syringe/needle delivery system, thermoreversible gels or post-administration viscosity-enhancing options are also envisioned, as well as alternative delivery systems, including pumps, microinjection devices and the like.

[00542] In one embodiment of the auris-acceptable aural pressure modulating formulations described herein, the aural pressure modulator is provided in a gel formulation, also referred to herein as "auris acceptable gel formulations," "auris interna-acceptable gel formulations," "auris gel formulations" or variations thereof. All of the components of the gel formulation must be compatible with the auris interna. Further, the gel formulations provide controlled release of the aural pressure modulator to the desired site within the auris interna; in some embodiments, the gel formulation also has an immediate or rapid release component for delivery of the aural pressure modulator to the desired target site.

[00543] Provided herein, in some embodiments, are auris formulations that comprise thermoreversible gelling polymers and/or hydrogels. In some instances, the formulations are liquid at or below room temperature but gel at body temperatures. In some instances, intratympanic injection of cold formulations (e.g., formulation with temperatures of $< 20^{\circ}\text{C}$) causes a dramatic change in the inner ear environment and causes vertigo in individuals undergoing treatment for inner ear disorders. Preferably, the formulations described herein are designed to be liquids that are administered at or near room temperature and do not cause vertigo or other discomfort when administered to an individual or patient.

[00544] In some embodiments, the formulations are bimodal formulations and comprise an immediate release component and an extended release component. In some instances, bimodal formulations allow for a constant rate of release of an immediate release component (multiparticulate agent (e.g., micronized active agent)) from the gelled polymer and a constant rate of release of an extended release component (e.g., an encapsulated active agent that serves as a depot for extending the release of an active agent). In other embodiments, the otic compositions described herein are administered as a controlled release formulation, released either continuously or in a pulsatile manner, or variants of both. In still other embodiments, the active agent formulation is administered as both an immediate release and controlled release formulation, released either continuously or in a pulsatile manner, or variants of both. In certain embodiments, the formulations comprise penetration enhancers that allow for delivery of the active agents across the oval window or the round window of the ear.

[00545] In some embodiments, the auris gel formulations are biodegradable. In other embodiments, the auris gel formulations include a mucoadhesive excipient to allow adhesion to the external mucous membrane of the round window. In yet other embodiments, the auris gel formulations include a penetration enhancer excipient; in further embodiments, the auris gel formulation contains a viscosity enhancing agent. In other embodiments, the auris pharmaceutical formulations provide an auris-acceptable microsphere or microparticle; in still other embodiments, the auris pharmaceutical formulations provide an auris-acceptable liposome, in yet other embodiments, the auris pharmaceutical formulations provide an auris-acceptable paint, foam or xerogel. In other

embodiments, the auris pharmaceutical formulations provide an auris-acceptable in situ forming spongy material. Further embodiments include a thermoreversible gel or actinic radiation curable gel in the auris pharmaceutical formulation, such that upon preparation of the gel at room temperature or below, the formulation is a fluid, but upon application of the gel into or near the auris interna and/or
5 auris media target site, including the tympanic cavity, round window membrane or the crista fenestrae cochleae, the auris-pharmaceutical formulation stiffens or hardens into a gel-like substance. Some embodiments include the use of a combination of a mucoadhesive and a thermoreversible gel in any otic formulation described herein.

[00546] The formulations disclosed herein alternatively encompass an otoprotectant agent in
10 addition to the at least one active agent and/or excipients, including but not limited to such as antioxidants, alpha lipoic acid, calicum, fosfomycin or iron chelators, to counteract potential ototoxic effects that may arise from the use of specific therapeutic agents or excipients, diluents or carriers.

[00547] One aspect of the embodiments disclosed herein is to provide a controlled release aural
15 pressure modulating auris-acceptable composition or formulation for the treatment of fluid homeostasis disorders. The controlled release aspect of the compositions and/or formulations disclosed herein is imparted through a variety of agents, including but not limited to excipients, agents or materials that are acceptable for use in the auris interna or other otic structure. By way of example only, such excipients, agents or materials include an auris-acceptable polymer, an auris-
20 acceptable viscosity enhancing agent, an auris-acceptable gel, an auris-acceptable microsphere, an auris-acceptable hydrogel, an auris-acceptable liposome, an auris-acceptable nanocapsule or nanosphere, an auris-acceptable thermoreversible gel, or combinations thereof.

[00548] Thus, provided herein are pharmaceutical compositions that include at least one auris
25 therapeutic agent and auris-acceptable diluent(s), excipient(s), and/or carrier(s). In some embodiments, the pharmaceutical compositions include other medicinal or pharmaceutical agents, carriers, adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure, and/or buffers. In other embodiments, the pharmaceutical compositions also contain other therapeutic substances.

Auris-Acceptable Gel Formulations

[00549] Gels, sometimes referred to as jellies, have been defined in various ways. For example, the
30 United States Pharmacopoeia defines gels as semisolid systems consisting of either suspensions made up of small inorganic particles or large organic molecules interpenetrated by a liquid. Gels can further consist of a single-phase or a two-phase system. A single-phase gel consists of organic macromolecules distributed uniformly throughout a liquid in such a manner that no apparent
35 boundaries exist between the dispersed macromolecules and the liquid. Single-phase gels are usually prepared from synthetic macromolecules (e.g., Carbomer®) or from natural gums, (e.g., tragacanth).

In some embodiments, single-phase gels are generally aqueous, but will also be made using alcohols and oils. Two-phase gels consist of a network of small discrete particles.

[00550] Gels can also be classified as being hydrophobic or hydrophilic. The bases of a hydrophobic gel usually consists of a liquid paraffin with polyethylene or fatty oils gelled with colloidal silica, or aluminum or zinc soaps. In contrast, the bases of hydrophilic gels usually consists of water, glycerol, or propylene glycol gelled with a suitable gelling agent (e.g., tragacanth, starch, cellulose derivatives, carboxyvinylpolymers, and/or magnesium-aluminum silicates).

[00551] In certain embodiments, the rheology of the gel formulation is pseudo plastic, plastic, thixotropic, or dilatant.

Thermoreversible Gels

[00552] Polymers composed of polyoxypropylene and polyoxyethylene are known to form thermoreversible gels when incorporated into aqueous solutions. These polymers have the ability to change from the liquid state to the gel state at temperatures close to body temperature, therefore allowing useful topical formulations. The liquid state-to-gel state phase transition is dependent on the polymer concentration and the ingredients in the solution.

[00553] "ReGel™" is a tradename of MacroMed Incorporated for a class of low molecular weight, biodegradable block copolymers having reverse thermal gelation properties as described in U.S. Pat. Nos. 6,004,573, 6,117,949, 6,201,072, and 6,287,588. It also includes biodegradable polymeric drug carriers disclosed in pending U.S. patent application Ser. Nos. 09/906,041, 09/559,799 and 10/919,603. The biodegradable drug carrier comprises ABA-type or BAB-type triblock copolymers or mixtures thereof, wherein the A-blocks are relatively hydrophobic and comprise biodegradable polyesters or poly(ortho ester)s, and the B-blocks are relatively hydrophilic and comprise polyethylene glycol (PEG), said copolymers having a hydrophobic content of between 50.1 to 83% by weight and a hydrophilic content of between 17 to 49.9% by weight, and an overall block copolymer molecular weight of between 2000 and 8000 daltons. The drug carriers exhibit water solubility at temperatures below normal mammalian body temperatures and undergo reversible thermal gelation to then exist as a gel at temperatures equal to physiological mammalian body temperatures. The biodegradable, hydrophobic A polymer block comprises a polyester or poly(ortho ester), in which the polyester is synthesized from monomers selected from the group consisting of D,L-lactide, D-lactide, L-lactide, D,L-lactic acid, D-lactic acid, L-lactic acid, glycolide, glycolic acid, ε-caprolactone, ε-hydroxyhexanoic acid, γ-butyrolactone, γ-hydroxybutyric acid, δ-valerolactone, δ-hydroxyvaleric acid, hydroxybutyric acids, malic acid, and copolymers thereof and having an average molecular weight of between about 600 and 3000 daltons. The hydrophilic B-block segment is preferably polyethylene glycol (PEG) having an average molecular weight of between about 500 and 2200 daltons.

[00554] Additional biodegradable thermoplastic polyesters include AtriGel™ (provided by Atrix Laboratories, Inc.) and/or those disclosed, e.g., in U.S. Patent Nos. 5,324,519; 4,938,763; 5,702,716; 5,744,153; and 5,990,194; wherein the suitable biodegradable thermoplastic polyester is disclosed as a thermoplastic polymer. Examples of suitable biodegradable thermoplastic polyesters include

5 polylactides, polyglycolides, polycaprolactones, copolymers thereof, terpolymers thereof, and any combinations thereof. In some such embodiments, the suitable biodegradable thermoplastic polyester is a polylactide, a polyglycolide, a copolymer thereof, a terpolymer thereof, or a combination thereof. In one embodiment, the biodegradable thermoplastic polyester is 50/50 poly (DL-lactide-co-glycolide) having a carboxy terminal group; is present in about 30 wt. % to about 40

10 wt. % of the composition; and has an average molecular weight of about 23,000 to about 45,000. Alternatively, in another embodiment, the biodegradable thermoplastic polyester is 75/25 poly (DL-lactide-co-glycolide) without a carboxy terminal group; is present in about 40 wt. % to about 50 wt. % of the composition; and has an average molecular weight of about 15,000 to about 24,000. In further or alternative embodiments, the terminal groups of the poly(DL-lactide-co-glycolide) are


15 either hydroxyl, carboxyl, or ester depending upon the method of polymerization. Polycondensation of lactic or glycolic acid provides a polymer with terminal hydroxyl and carboxyl groups. Ring-opening polymerization of the cyclic lactide or glycolide monomers with water, lactic acid, or glycolic acid provides polymers with the same terminal groups. However, ring-opening of the cyclic monomers with a monofunctional alcohol such as methanol, ethanol, or 1-dodecanol provides a

20 polymer with one hydroxyl group and one ester terminal groups. Ring-opening polymerization of the cyclic monomers with a diol such as 1,6-hexanediol or polyethylene glycol provides a polymer with only hydroxyl terminal groups.

[00555] Additional embodiments include Poloxamer thermoreversible copolymers. Poloxamer 407 (PF-127) is a nonionic surfactant composed of polyoxyethylene-polyoxypropylene copolymers.

25 Other commonly used poloxamers include 188 (F-68 grade), 237 (F-87 grade), 338 (F-108 grade). Aqueous solutions of poloxamers are stable in the presence of acids, alkalis, and metal ions. PF-127 is a commercially available polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106, with an average molar mass of 13,000. It contains approximately 70% ethylene oxide, which accounts for its hydrophilicity. It is one of the series of poloxamer ABA block

30 copolymers, whose members share the chemical formula shown below.



[00556] P-F127 is of particular interest since concentrated solutions (>20% w/w) of the copolymer are transformed from low viscosity transparent solutions to solid gels on heating to body temperature. This phenomenon, therefore, suggests that when placed in contact with the body, the gel preparation will form a semi-solid structure and a controlled release depot. Furthermore, PF-127 has good solubilizing capacity, low toxicity and is, therefore, considered a good medium for drug delivery systems.

[00557] In an alternative embodiment, the thermogel is a PEG-PGLA-PEG triblock copolymer (Jeong et al, Nature (1997), 388:860-2; Jeong et al, J. Control. Release (2000), 63:155-63; Jeong et al, Adv. Drug Delivery Rev. (2002), 54:37-51). The polymer exhibits sol-gel behavior over a concentration of about 5% w/w to about 40% w/w. Depending on the properties desired, the lactide/glycolide molar ratio in the PGLA copolymer can range from about 1:1 to about 20:1. The resulting copolymers are soluble in water and form a free-flowing liquid at room temperature, but form a hydrogel at body temperature. A commercially available PEG-PGLA-PEG triblock copolymer is RESOMER RGP t50106 manufactured by Boehringer Ingelheim. This material is composed of a PGLA copolymer of 50:50 poly(DL-lactide-co-glycolide) and is 10% w/w of PEG and has a molecular weight of about 6000.

[00558] In some embodiments, a suitable combination of gelling agents and a thermoreversible gel is utilized in the controlled release formulations described herein. Suitable gelling agents for use in preparation of the gel formulation include, but are not limited to, celluloses, cellulose derivatives, cellulose ethers (e.g., carboxymethylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxymethylcellulose, hydroxypropylmethylcellulose, hydroxypropylcellulose, methylcellulose), guar gum, xanthan gum, locust bean gum, alginates (e.g., alginic acid), silicates, starch, tragacanth, carboxyvinyl polymers, carrageenan, paraffin, petrolatum and any combinations or mixtures thereof. In some other embodiments, hydroxypropylmethylcellulose (Methocel®) is utilized as the gelling agent. In certain embodiments, the thickening agents (i.e., viscosity enhancing agents) described herein are also utilized as the gelling agent for the gel formulations presented herein.

[00559] Suitable combinations of thermoreversible gels with a thickening agent include, by way of non-limiting example, a combination of poloxamer thermoreversible copolymers with cellulose based thickening agents described herein. In some instances, the addition of a secondary polymer (e.g., a thickening agent) introduces a diffusional barrier and reduces the rate of release of the otic agent. An appropriate thickening agent (e.g., a cellulose based polymer, e.g., CMC polymer) is selected based on the viscosity of a 2% solution of the secondary polymer (e.g., CMC); the selected secondary polymer (e.g., CMC) provides a 2% polymer solution with viscosity less than 15,000 cP. In specific formulations, the selected secondary polymer (e.g., CMC) provides a 2% polymer solution with viscosity from about 4,000 cP to about 10,000 cP. In some embodiments, the ratio of a thermoreversible poloxamer to a gelling agent is about 50:1, about 40:1, about 35:1, about 30:1,

about 25:1, about 20:1, about 15:1 or about 10:1. For example, in certain embodiments, a controlled release formulation described herein comprises a combination of poloxamer 407 (pluronic F127) and carboxymethylcellulose (CMC) in a ratio of about 50:1, 40:1, about 35:1, about 30:1, about 25:1, about 20:1, about 15:1 or about 10:1.

5 *Hydrogels*

[00560] Chitosan glycerophosphate (CGP) is a biodegradable matrix for the formation of hydrogels. CGP has been shown to be suitable for local delivery of dexamethasone to the inner ear, where 50% of the active agent was released after 24 hours, followed by a linear decline over 5 days of perilymph drug levels. In some embodiments, CGP is used as a biodegradable viscosity enhancing agent or gelling agent for controlled release of active agents from the formulations disclosed herein. In certain embodiments, when CGP is used as a viscosity enhancing agent or gelling agent, the compositions further comprise liposomes. Liposomes are added to further control the release of active agents from the formulations disclosed herein, whether they be hydrophobic or hydrophilic antimicrobial small molecules.

15 [00561] In some embodiments, other gel formulations are also contemplated to be useful depending upon the particular embodiment, and as such are considered to fall within the scope of the present disclosure. For example, other currently commercially-available glycerin-based gels, glycerin-derived compounds, conjugated, or crosslinked gels, matrices, hydrogels, and polymers, as well as gelatins and their derivatives, alginates, and alginate-based gels, and even various native and synthetic hydrogel and hydrogel-derived compounds are all expected to be useful in the formulations described herein. In some embodiments, gels include, but are not limited to, alginate hydrogels SAF-Gel (ConvaTec, Princeton, N.J.), Duoderm Hydroactive Gel (ConvaTec), Nu-gel (Johnson & Johnson Medical, Arlington, Tex.); Carrasyn (V) Acemannan Hydrogel (Carrington Laboratories, Inc., Irving, Tex.); glycerin gels Elta Hydrogel (Swiss-American Products, Inc., Dallas, Tex.) and K-Y Sterile (Johnson & Johnson). In one embodiment, a sterilized solution of sodium alginate is mixed with a sterilized solution of an auris-compatible calcium salt, the therapeutic agent(s), and a polysaccharide. Upon admixing, a gel is formed in a desired amount of time having a desired viscosity. In further embodiments, biodegradable biocompatible gels also represent compounds present in formulations disclosed and described herein. In some embodiments, a hardening agent (e.g., glutaraldehyde, formaldehyde) is added to a biodegradable hydrogel gel. Contemplated for use in formulations described herein are biodegradable hydrogels comprising, by way of example, 0.1, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90 or 100 mM glutaraldehyde (e.g., a gelatin gel and/or a glycerin gel and/or a chitosan hydrogel comprising 10 mM glutaraldehyde). In further embodiments, biodegradable biocompatible gels also represent compounds present in auris-acceptable formulations disclosed and described herein. For examples, of formulations and their characteristics see Table 1.

[00562] In some formulations developed for administration to a mammal, and for compositions formulated for human administration, the gel comprises substantially all of the weight of the composition. In other embodiments, the gel comprises as much as about 98% or about 99% of the composition by weight. In a further embodiment, this is desirable when a substantially non-fluid, or substantially viscous formulation is needed. In a further embodiment, when slightly less viscous, or slightly more fluid formulations are desired, the biocompatible gel portion of the formulation comprises at least about 50% by weight, at least about 60% by weight, at least about 70% by weight, or even at least about 80% or 90% by weight of the compound. Of course, all intermediate integers within these ranges are contemplated to fall within the scope of this disclosure, and in some embodiments, even more fluid (and consequently less viscous) gel compositions are formulated, such as for example, those in which the gel or matrix component of the mixture comprises not more than about 50% by weight, not more than about 40% by weight, not more than about 30% by weight, or even those that comprise not more than about 15% or about 20% by weight of the composition.

[00563] If desired, the gels may also contain preservatives, cosolvents, suspending agents, viscosity enhancing agents, ionic-strength and osmolality adjusters and other excipients in addition to buffering agents. Suitable water soluble preservatives which are employed in the drug delivery vehicle are sodium bisulfite, sodium thiosulfate, ascorbate, benzalkonium chloride, chlorobutanol, thimerosal, parabens, benzyl alcohol, phenylethanol and others. These agents are present, generally, in amounts of about 0.001% to about 5% by weight and, preferably, in the amount of about 0.01 to about 2% by weight.

[00564] Suitable water soluble buffering agents are alkali or alkaline earth metal carbonates, phosphates, bicarbonates, citrates, borates, acetates, succinates and the like, such as sodium phosphate, citrate, borate, acetate, bicarbonate, carbonate and tromethamine (TRIS). These agents are present in amounts sufficient to maintain the pH of the system at 7.4 ± 0.2 and preferably, 7.4. As such, the buffering agent can be as much as 5% on a weight basis of the total composition.

[00565] Cosolvents can be used to enhance drug solubility, however, some drugs are insoluble. These can often be suspended in the polymer vehicle with the aid of suitable suspending or viscosity enhancing agents.

[00566] Since the polymer systems of the thermoreversible gel dissolve more completely at reduced temperatures, the preferred methods of solubilization are to add the required amount of polymer to the amount of water to be used. Generally after wetting the polymer by shaking, the mixture is capped and placed in a cold chamber or in a thermostatic container at about 0-10 °C in order to dissolve the polymer. The mixture can be stirred or shaken to bring about a more rapid dissolution of the polymer. The active pharmaceutical ingredient and various additives such as buffers, salts, and preservatives can subsequently be added and dissolved. In some embodiments the

pharmacologically active substance is suspended if it is insoluble in water. The pH is modulated by the addition of appropriate buffering agents.

[00567] In certain embodiments, the polymer systems of the thermoreversible gels are designed to remain liquids up to temperatures of about 15 – 25 °C, about 18 – 22 °C, or about 20 °C. In some instances, the formulations described herein are manufactured under conditions such that the temperature of the manufacturing room is maintained below 25°C to retain the temperature of a polymer solution at about 25°C, about 23°C, about 21°C, or about 19°C. In certain instances, the formulations described herein are manufactured under conditions such that the temperature of a manufacturing room is maintained at about 19°C. In some of such instances, the temperature of the polymer solution is maintained at or below about 19°C up to 3 hours of the initiation of the manufacturing, without the need to chill/cool the container. In some instances, the temperature of the solution is maintained at or below about 19°C up to 3 hours of the initiation of the manufacturing by use of a jacketed container for the polymer solution.

Auris-Acceptable Actinic Radiation Curable Gel

[00568] In other embodiments, the gel is an actinic radiation curable gel, such that following administration to or near the targeted auris structure, use of actinic radiation (or light, including UV light, visible light, or infrared light) the desired gel properties are formed. By way of example only, fiber optics are used to provide the actinic radiation so as to form the desired gel properties. In some embodiments, the fiber optics and the gel administration device form a single unit. In other embodiments, the fiber optics and the gel administration device are provided separately.

Auris-Acceptable Solvent Release Gel

[00569] In some embodiments, the gel is a solvent release gel such that the desired gel properties are formed after administration to or near the targeted auris structure, that is, as the solvent in the injected gel formulation diffuses out the gel, a gel having the desired gel properties is formed. For example, a formulation that comprises sucrose acetate isobutyrate, a pharmaceutically acceptable solvent, one or more additives, and the auris therapeutic agent is administered at or near the round window membrane: diffusion of the solvent out of the injected formulation provides a depot having the desired gel properties. For example, use of a water soluble solvent provides a high viscosity depot when the solvent diffuses rapidly out of the injected formulation. On the other hand, use of a hydrophobic solvent (e.g., benzyl benzoate) provides a less viscous depot. One example of an auris-acceptable solvent release gel formulation is the SABER™ Delivery System marketed by DURECT Corporation.

[00570] Presented below are examples of potential controlled release excipients:

Example Formulation	Example Characteristics
Chitosan glycerophosphate (CGP)	- tunable degradation of matrix in vitro - tunable VP2 modulator release in vitro: e.g., ~50 % of drug released after 24 hrs

	• biodegradable • compatible with drug delivery to the inner ear • suitable for macromolecules and hydrophobic drugs
PEG-PLGA-PEG triblock polymers	• tunable high stability: e.g., maintains mechanical integrity > 1 month in vitro • tunable fast release of hydrophilic drugs: e.g., ~ 50 % of drug released after 24 hrs, and remainder released over ~ 5 days • tunable slow release of hydrophobic drugs: e.g., ~ 80 % released after 8 weeks • biodegradable • subcutaneous injection of solution: e.g., gel forms within seconds and is intact after 1 month
PEO-PPO-PEO triblock copolymers (e.g., Pluronic or Poloxameres) (e.g., F127)	• Tunable sol-gel transition temperature: e.g., decreases with increasing F127 concentration
Chitosan glycerophosphate with drug-loaded liposomes	• CGP formulation tolerates liposomes: e.g., up to 15 uM/mL liposomes. • liposomes tunably reduce drug release time (e.g., up to 2 weeks in vitro). • increase in liposome diameter optionally reduces drug release kinetics (e.g., liposome size between 100 and 300 nm) • release parameters are controlled by changing composition of liposomes

Auris Interna Mucoadhesive Excipients

[00571] Mucoadhesive characteristics may also be imparted to the gel or other auris-interna formulations disclosed herein, including a thermoreversible gel, by incorporation of mucoadhesive carbomers, such as Carbopol 934P, to the composition (Majithiya et al, AAPS PharmSciTech (2006), 7(3), p. E1; EP0551626).

[00572] The term 'mucoadhesion' is commonly used for materials that bind to the mucin layer of a biological membrane. To serve as mucoadhesive polymers, the polymers should possess some general physiochemical features such as predominantly anionic hydrophilicity with numerous hydrogen bond forming groups, suitable surface property for wetting mucus/mucosal tissue surfaces and sufficient flexibility to penetrate the mucus network. In some embodiments, mucoadhesive formulations described herein adhere to the round window and/or the oval window and/or any inner ear structure.

[00573] Mucoadhesive agents including, but not limited to, at least one soluble polyvinylpyrrolidone polymer (PVP); a water-swellaable, but water-insoluble, fibrous, cross-linked carboxy-functional polymer; a crosslinked poly(acrylic acid) (e.g. Carbopol 947P); a carbomer homopolymer; a carbomer copolymer; a hydrophilic polysaccharide gum, maltodextrin, a cross-linked alginate gum gel, a water-dispersible polycarboxylated vinyl polymer, at least two particulate components selected from the group consisting of titanium dioxide, silicon dioxide, and clay, or a mixture thereof. The mucoadhesive agent are used in combination with a viscosity increasing excipient, or

are used alone to increase the interaction of the composition with a mucosal layer. In one non-limiting example, the mucoadhesive agent is maltodextrin and/or an alginate gum. Those of ordinary skill in the art will recognize that the mucoadhesive character imparted to the composition should be at a level that is sufficient to deliver an effective amount of the composition to, for example, the mucosal membrane of the round window in an amount that may coat the mucosal membrane, and thereafter deliver the composition to the affected areas, including by way of example only, the vestibular and/or cochlear structures of the auris interna. Those of ordinary skill in the art can determine the mucoadhesive characteristics of the compositions provided herein, and may thus determine appropriate ranges. One method for determining sufficient mucoadhesiveness may include monitoring changes in the interaction of the composition with a mucosal layer, including but not limited to measuring changes in residence or retention time of the composition in the absence and presence of the excipient.

[00574] Mucoadhesive agents have been described, for example, in U.S. Patent Nos. 6,638,521, 6,562,363, 6,509,028, 6,348,502, 6,319,513, 6,306,789, 5,814,330, and 4,900,552, each of which is hereby incorporated by reference in its entirety.

[00575] In one non-limiting example, the mucoadhesive agent is maltodextrin. Maltodextrin is a carbohydrate produced by the hydrolysis of starch that are derived from corn, potato, wheat or other plant products. Maltodextrin are used either alone or in combination with other mucoadhesive agents to impart mucoadhesive characteristics on the compositions disclosed herein. In one embodiment, a combination of maltodextrin and a carbopol polymer are used to increase the mucoadhesive characteristics of the compositions disclosed herein.

[00576] In another non-limiting example, a mucoadhesive agent can be, for example, at least two particulate components selected from titanium dioxide, silicon dioxide, and clay, wherein the composition is not further diluted with any liquid prior to administration and the level of silicon dioxide, if present, is from about 3% to about 15%, by weight of the composition. Silicon dioxide, if present, are selected from the group consisting of fumed silicon dioxide, precipitated silicon dioxide, coacervated silicon dioxide, gel silicon dioxide, and mixtures thereof. Clay, if present, are kaolin minerals, serpentine minerals, smectites, illite or a mixture thereof. For example, clay can be laponite, bentonite, hectorite, saponite, montmorillonites or a mixture thereof.

Stabilizers

[00577] In one embodiment, stabilizers are selected from, for example, fatty acids, fatty alcohols, alcohols, long chain fatty acid esters, long chain ethers, hydrophilic derivatives of fatty acids, polyvinyl pyrrolidones, polyvinyl ethers, polyvinyl alcohols, hydrocarbons, hydrophobic polymers, moisture-absorbing polymers, and combinations thereof. In some embodiments, amide analogues of stabilizers are also used. In a further embodiment, the chosen stabilizer changes the hydrophobicity of the formulation (e.g., oleic acid, waxes), or improves the mixing of various components in the

formulation (e.g., ethanol), controls the moisture level in the formula (e.g., PVP or polyvinyl pyrrolidone), controls the mobility of the phase (substances with melting points higher than room temperature such as long chain fatty acids, alcohols, esters, ethers, amides etc. or mixtures thereof; waxes), and/or improves the compatibility of the formula with encapsulating materials (e.g., oleic acid or wax). In another embodiment some of these stabilizers are used as solvents/co-solvents (e.g., ethanol). In a further embodiment, stabilizers are present in sufficient amount to inhibit the degradation of the active pharmaceutical ingredient. Examples of such stabilizing agents, include, but are not limited to: (a) about 0.5% to about 2% w/v glycerol, (b) about 0.1% to about 1% w/v methionine, (c) about 0.1% to about 2% w/v monothioglycerol, (d) about 1 mM to about 10 mM EDTA, (e) about 0.01% to about 2% w/v ascorbic acid, (f) 0.003% to about 0.02% w/v polysorbate 80, (g) 0.001% to about 0.05% w/v. polysorbate 20, (h) arginine, (i) heparin, (j) dextran sulfate, (k) cyclodextrins, (l) pentosan polysulfate and other heparinoids, (m) divalent cations such as magnesium and zinc; or (n) combinations thereof.

[00578] Additional useful auris-acceptable formulations include one or more anti-aggregation additives to enhance stability of otic formulations by reducing the rate of protein aggregation. The anti-aggregation additive selected depends upon the nature of the conditions to which the otic agents, for example anti-TNF antibodies are exposed. For example, certain formulations undergoing agitation and thermal stress require a different anti-aggregation additive than a formulation undergoing lyophilization and reconstitution. Useful anti-aggregation additives include, by way of example only, urea, guanidinium chloride, simple amino acids such as glycine or arginine, sugars, polyalcohols, polysorbates, polymers such as polyethylene glycol and dextrans, alkyl saccharides, such as alkyl glycoside, and surfactants.

[00579] Other useful formulations include one or more antioxidants to enhance chemical stability where required. Suitable antioxidants include, by way of example only, ascorbic acid and sodium metabisulfite. In one embodiment, antioxidants are selected from metal chelating agents, thiol containing compounds and other general stabilizing agents.

[00580] Still other useful compositions include one or more surfactants to enhance physical stability or for other purposes. Suitable nonionic surfactants include polyoxyethylene fatty acid glycerides and vegetable oils, e.g., polyoxyethylene (60) hydrogenated castor oil; and polyoxyethylene alkylethers and alkylphenyl ethers, e.g., octoxynol 10, octoxynol 40.

[00581] In some embodiments, the pharmaceutical formulations described herein are stable with respect to compound degradation over a period of any of at least about 1 day, at least about 2 days, at least about 3 days, at least about 4 days, at least about 5 days, at least about 6 days, at least about 1 week, at least about 2 weeks, at least about 3 weeks, at least about 4 weeks, at least about 5 weeks, at least about 6 weeks, at least about 7 weeks, at least about 8 weeks, at least about 1 month, at least about 2 months, at least about 3 months, at least about 4 months, at least about 5 months, or at least

about 6 months. In other embodiments, the formulations described herein are stable with respect to compound degradation over a period of at least about 1 week. Also described herein are formulations that are stable with respect to compound degradation over a period of at least about 1 month.

5 [00582] In other embodiments, an additional surfactant (co-surfactant) and/or buffering agent is combined with one or more of the pharmaceutically acceptable vehicles previously described herein so that the surfactant and/or buffering agent maintains the product at an optimal pH for stability. Suitable co-surfactants include, but are not limited to: a) natural and synthetic lipophilic agents, e.g., phospholipids, cholesterol, and cholesterol fatty acid esters and derivatives thereof; b) nonionic
10 surfactants, which include for example, polyoxyethylene fatty alcohol esters, sorbitan fatty acid esters (Spans), polyoxyethylene sorbitan fatty acid esters (e.g., polyoxyethylene (20) sorbitan monooleate (Tween 80), polyoxyethylene (20) sorbitan monostearate (Tween 60), polyoxyethylene (20) sorbitan monolaurate (Tween 20) and other Tweens, sorbitan esters, glycerol esters, e.g., Myrj and glycerol triacetate (triacetin), polyethylene glycols, cetyl alcohol, cetostearyl alcohol, stearyl
15 alcohol, polysorbate 80, poloxamers, poloxamines, polyoxyethylene castor oil derivatives (e.g., Cremophor® RH40, Cremphor A25, Cremphor A20, Cremophor® EL) and other Cremophors, sulfosuccinates, alkyl sulphates (SLS); PEG glyceryl fatty acid esters such as PEG-8 glyceryl caprylate/caprate (Labrasol), PEG-4 glyceryl caprylate/caprate (Labrafac Hydro WL 1219), PEG-32 glyceryl laurate (Gelucire 444/14), PEG-6 glyceryl mono oleate (Labrafil M 1944 CS), PEG-6
20 glyceryl linoleate (Labrafil M 2125 CS); propylene glycol mono- and di-fatty acid esters, such as propylene glycol laurate, propylene glycol caprylate/caprate; Brij® 700, ascorbyl-6-palmitate, stearylamine, sodium lauryl sulfate, polyoxethyleneglycerol triiricinoleate, and any combinations or mixtures thereof; c) anionic surfactants include, but are not limited to, calcium carboxymethylcellulose, sodium carboxymethylcellulose, sodium sulfosuccinate, dioctyl, sodium
25 alginate, alkyl polyoxyethylene sulfates, sodium lauryl sulfate, triethanolamine stearate, potassium laurate, bile salts, and any combinations or mixtures thereof; and d) cationic surfactants such as quarternary ammonium compounds, benzalkonium chloride, cetyltrimethylammonium bromide, and lauryldimethylbenzyl-ammonium chloride.

[00583] In a further embodiment, when one or more co-surfactants are utilized in the formulations of
30 the present disclosure, they are combined, e.g., with a pharmaceutically acceptable vehicle and is present in the final formulation, e.g., in an amount ranging from about 0.1% to about 20%, from about 0.5% to about 10%. In one embodiment, the surfactant has an HLB value of 0 to 20. In additional embodiments, the surfactant has an HLB value of 0 to 3, of 4 to 6, of 7 to 9, of 8 to 18, of 13 to 15, of 10 to 18.

35 *Preservatives*

[00584] In some embodiments, an auris controlled release formulation described herein is free of preservatives. In some embodiments, a composition disclosed herein comprises a preservative. Suitable auris-acceptable preservatives for use in a composition disclosed herein include, but are not limited to benzoic acid, boric acid, p-hydroxybenzoates, benzyl alcohol, lower alkyl alcohols (e.g., ethanol, butanol or the like), quaternary compounds, stabilized chlorine dioxide, mercurials, such as merfen and thiomersal, mixtures of the foregoing and the like. Suitable preservatives for use with a formulation disclosed herein are not ototoxic. In some embodiments, a formulation disclosed herein does not include a preservative that is ototoxic. In some embodiments, a formulation disclosed herein does not include benzalkonium chloride or benzethonium chloride.

[00585] In certain embodiments, any controlled release formulation described herein has an endotoxin level of less than 0.5 EU/kg, less than 0.4 EU/kg or less than 0.3 EU/kg. In certain embodiments, any controlled release formulation described herein has less than about 60 colony forming units (CFU), has less than about 50 colony forming units, has less than about 40 colony forming units, has less than about 30 colony forming units of microbial agents per gram of formulation. In certain embodiments, any controlled release formulation described herein is substantially free of pyrogens.

[00586] In a further embodiment, the preservative is, by way of example only, an antimicrobial agent, within the formulation presented herein. In one embodiment, the formulation includes a preservative such as by way of example only, methyl paraben. In another embodiment, the methyl paraben is at a concentration of about 0.05% to about 1.0%, about 0.1% to about 0.2%. In a further embodiment, the gel is prepared by mixing water, methylparaben, hydroxyethylcellulose and sodium citrate. In a further embodiment, the gel is prepared by mixing water, methylparaben, hydroxyethylcellulose and sodium acetate. In a further embodiment, the mixture is sterilized by autoclaving at 120 °C for about 20 minutes, and tested for pH, methylparaben concentration and viscosity before mixing with the appropriate amount of the active pharmaceutical ingredient disclosed herein. In certain embodiments, the preservative employed in any auris-compatible formulation described herein is an antioxidant (e.g., butyl hydroxytoluene (BHT) or the like, as described herein). In certain embodiments, an antioxidant preservative is non-toxic and/or non-irritating to the inner ear environment.

Carriers

[00587] Suitable carriers for use in a formulation described herein include, but are not limited to, any pharmaceutically acceptable solvent. For example, suitable solvents include polyalkylene glycols such as, but not limited to, polyethylene glycol (PEG) and any combinations or mixtures thereof. In other embodiments, the base is a combination of a pharmaceutically acceptable surfactant and solvent.

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 143

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 143

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

WE CLAIM:

1. A pharmaceutical formulation for use in the treatment of an otic disease or condition formulated to provide a therapeutically effective amount of an immunomodulating agent, the
5 formulation comprising:
 - between about 0.2% to about 6% by weight of an immunomodulating agent, or pharmaceutically acceptable prodrug or salt thereof;
 - between about 16% to about 21% by weight of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106;
 - 10 sterile water, q.s., buffered to provide a perilymph-suitable pH between about 6.0 and about 7.6;
 - and substantially low degradation products of the immunomodulating agent;wherein the pharmaceutical formulation has a perilymph-suitable osmolarity between about 250 and 320 mOsm/L,
- 15 less than about 50 colony forming units (cfu) of microbiological agents per gram of formulation, and less than about 5 endotoxin units (EU) per kg of body weight of a subject.
2. The composition of claim 1, wherein the immunomodulating agent is released from the formulation for a period of at least 3 days.
3. The composition of claim 1 or claim 2, wherein the pharmaceutical formulation is an auris-
20 acceptable thermoreversible gel.
4. The compositions of any of claims 1-3, wherein the polyoxyethylene-polyoxypropylene triblock copolymer is biodegradable.
5. The composition of any of claims 1-4, further comprising a mucoadhesive.
6. The composition of any of claims 1-5, further comprising a penetration enhancer.
- 25 7. The composition of any of claims 1-6, further comprising a thickening agent.
8. The composition of any of claims 1-7, further comprising a dye.
9. The composition of any of claims 1-8, further comprising a drug delivery device selected from a needle and syringe, a pump, a microinjection device, a wick, an in situ forming spongy material or combinations thereof.
- 30 10. The composition of any of claims 1-8, wherein the immunomodulating agent, or pharmaceutically acceptable salt thereof, has limited or no systemic release, systemic toxicity, poor PK characteristics, or combinations thereof.
11. The composition of any of claims 1-8, wherein the immunomodulating agent is in the form of a free base, a salt, a prodrug, or a combination thereof.
- 35 12. The composition of any of claims 1-11, wherein the immunomodulating agent comprises multiparticulates.

13. The composition of claim 12, wherein the immunomodulating agent is essentially in the form of micronized particles.

14. The composition of any of claims 1-13, wherein the immunomodulating agent is an anti-TNF agent, a calcineurin inhibitor, an IKK inhibitor, an interleukin inhibitor, a TNF- α converting enzyme (TACE) inhibitor, or a toll-like receptor inhibitor.

15. The composition of any of claims 1-14, further comprising an immunomodulating agent, or pharmaceutically acceptable salt, free based, prodrug or combination thereof, as an immediate release agent.

16. The composition of any of claims 1-15 further comprising an additional therapeutic agent.

17. The composition of claim 16, wherein the additional therapeutic agent is a Na/K ATPase modulator, a chemotherapeutic agent, a collagen, a gamma-globulin, an interferon, an anti-microbial agent, an antibiotic, a local acting anesthetic agent, a platelet activator factor antagonist, a nitric oxide synthase inhibitor, an anti-vertigo agent, a vasopressin antagonist, an anti-viral, an anti-emetic agent or combinations thereof.

18. The composition of any of claims 1-17, wherein the pH of the composition is between about 6.0 to about 7.6.

19. The composition of any of claims 1-18, wherein the ratio of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106 to a thickening agent is from about 40:1 to about 10:1.

20. The composition of claim 19, wherein the thickening agent is carboxymethyl cellulose.

21. The composition of claim 1, wherein the otic disease or condition is Meniere's disease, sudden sensorineural hearing loss, noise induced hearing loss, age related hearing loss, auto immune ear disease or tinnitus.

22. A method of treating an otic disease or condition comprising administering to an individual in need thereof an intratympanic composition comprising

between about 0.2% to about 6% by weight of an immunomodulating agent, or pharmaceutically acceptable prodrug or salt thereof;

between about 16% to about 21% by weight of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106;

sterile water, q.s., buffered to provide a perilymph-suitable pH between about 6.0 and about 7.6;

and substantially low degradation products of the immunomodulating agent;

wherein the pharmaceutical formulation has a perilymph-suitable osmolarity between about 250 and 320 mM,

less than about 50 colony forming units (cfu) of microbiological agents per gram of formulation, and less than about 5 endotoxin units (EU) per kg of body weight of a subject.

23. The method of claim 22, wherein the immunomodulating agent is an anti-TNF agent, a calcineurin inhibitor, an IKK inhibitor, an interleukin inhibitor, a TNF- α converting enzyme (TACE) inhibitor, or a toll-like receptor inhibitor.

24. The method of claim 22 wherein the immunomodulating agent is released from the composition for a period of at least 3 days.

25. The method of claim 22, wherein the composition is administered across the round window.

26. The method of claim 22, wherein the otic disease or condition is Meniere's disease, sudden sensorineural hearing loss, noise induced hearing loss, age related hearing loss, auto immune ear disease or tinnitus.

27. A pharmaceutical formulation for use in the treatment of an otic disease or condition formulated to provide a therapeutically effective amount of an aural pressure modulating, the formulation comprising:

between about 0.2% to about 6% by weight of an aural pressure modulating agent, or pharmaceutically acceptable prodrug or salt thereof;

between about 16% to about 21% by weight of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106;

sterile water, q.s., buffered to provide a perilymph-suitable pH between about 6.0 and about 7.6;

and substantially low degradation products of the aural pressure modulating agent;

wherein the pharmaceutical formulation has a perilymph-suitable osmolarity between about 250 and 320 mOsm/L,

less than about 50 colony forming units (cfu) of microbiological agents per gram of formulation, and less than about 5 endotoxin units (EU) per kg of body weight of a subject.

28. The composition of claim 27, wherein the aural pressure modulating agent is released from the formulation for a period of at least 3 days.

29. The composition of claim 27 or 28, wherein the pharmaceutical formulation is an auris-acceptable thermoreversible gel.

30. The compositions of any of claims 27-29, wherein the polyoxyethylene-polyoxypropylene triblock copolymer is biodegradable.

31. The composition of any of claims 27-30, further comprising a mucoadhesive.

32. The composition of any of claims 27-31, further comprising a penetration enhancer.

33. The composition of any of claims 27-32, further comprising a thickening agent.

34. The composition of any of claims 27-33, further comprising a dye.

35. The composition of any of claims 27-34, further comprising a drug delivery device selected

from a needle and syringe, a pump, a microinjection device, a wick, an in situ forming spongy material or combinations thereof.

36. The composition of any of claims 27-35, wherein the aural pressure modulating agent, or pharmaceutically acceptable salt thereof, has limited or no systemic release, systemic toxicity, poor PK characteristics, or combinations thereof.

37. The composition of any of claims 27-36, wherein the aural pressure modulating agent is in the form of a free base, salt, a prodrug, or a combination thereof.

38. The composition of any of claims 27-37, wherein the aural pressure modulating agent comprises multiparticulates

39. The composition of claim 38, wherein the aural pressure modulating agent is essentially in the form of micronized particles.

40. The composition of any of claims 27-39, wherein the aural pressure modulating agent is a modulator of aquaporin, an estrogen related receptor beta modulator, a gap junction protein modulator, an NMDA receptor modulator, an osmotic diuretic, a progesterone receptor modulator, a prostaglandin modulator, or a vasopressin receptor modulator.

41. The composition of any of claims 27-40, further comprising an aural pressure modulating agent, or pharmaceutically acceptable salt thereof, as an immediate release agent.

42. The composition of any of claims 27-41, further comprising an additional therapeutic agent.

43. The composition of claim 42, wherein the additional therapeutic agent is a Na/K ATPase modulator, a chemotherapeutic agent, a collagen, a gamma-globulin, an interferon, an anti-microbial agent, an antibiotic, a local acting anesthetic agent, a platelet activator factor antagonist, a nitric oxide synthase inhibitor, an anti-vertigo medicine, a vasopressin antagonist, an anti-viral, an anti-emetic agent or combinations thereof.

44. The composition of any of claims 27-43, wherein the pH of the composition is between about 7.0 to about 7.6.

45. The composition of any of claims 27-44, wherein the ratio of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106 to a thickening agent is from about 40:1 to about 10:1.

46. The composition of claim 45, wherein the thickening agent is carboxymethyl cellulose.

47. The composition of claim 46, wherein the otic disease or condition is Meniere's disease, sudden sensorineural hearing loss, noise induced hearing loss, age related hearing loss, auto immune ear disease or tinnitus.

48. A method of treating an otic disease or condition comprising administering to an individual in need thereof an intratympanic composition comprising

between about 0.2% to about 6% by weight of an aural pressure modulating agent, or pharmaceutically acceptable prodrug or salt thereof;

between about 16% to about 21% by weight of a polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106;

sterile water, q.s., buffered to provide a perilymph-suitable pH between about 6.0 and about 7.6;

and substantially low degradation products of the aural pressure modulating agent;

wherein the pharmaceutical formulation has a perilymph-suitable osmolarity between about 250 and
5 320 mM,

less than about 50 colony forming units (cfu) of microbiological agents per gram of formulation, and less than about 5 endotoxin units (EU) per kg of body weight of a subject.

49. The method of claim 48, wherein the aural pressure modulating agent is a modulator of aquaporin, an estrogen related receptor beta modulator, a gap junction protein modulator, an NMDA
10 receptor modulator, an osmotic diuretic, a progesterone receptor modulator, a prostaglandin modulator, or a vasopressin receptor modulator.

50. The method of claim 48 wherein the aural pressure modulating agent is released from the composition for a period of at least 3 days.

51. The method of claim 48, wherein the composition is administered across the round window.

15 52. The method of claim 48, wherein the otic disease or condition is Meniere's disease, sudden sensorineural hearing loss, noise induced hearing loss, age related hearing loss, auto immune ear disease or tinnitus.

FIGURE 1




FIGURE 2




FIGURE 3

