



AU9673326

(12) PATENT ABRIDGMENT (11) Document No. AU-B-73326/96
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 700386

(54) Title
TRICICLIC COMPOUNDS

International Patent Classification(s)

(51) ⁶ C07C 235/16	A61K 031/35	A61K 031/38	A61K 031/44
C07D 221/06	C07D 307/93	C07D 313/12	C07D 333/78
C07D 337/12	C07D 495/04	A61K 031/435	C07D 491/044

(21) Application No. : 73326/96

(22) Application Date : 16.10.96

(87) PCT Publication Number : WO97/14672

(30) Priority Data

(31) Number	(32) Date	(33) Country
7-267232	16.10.95	JP JAPAN
7-324298	13.12.95	JP JAPAN

(43) Publication Date : 07.05.97

(44) Publication Date of Accepted Application : 07.01.99


(71) Applicant(s)
KYOWA HAKKO KOGYO CO., LTD.

(72) Inventor(s)
MAKOTO YOSHIDA; SHIN-ICHI SASAKI; SHIGERU AONO; SHIGEKI FUJIWARA; HARUKI TAKAI; TSUYOSHI YAMAGATA; KEN NAGASAHIMA; AKIRA KARASAWA

(74) Attorney or Agent
SPRUSON & FERGUSON, GPO Box 3898, SYDNEY NSW 2001

(57) Claim

1. A tricyclic compound represented by general formula (I):



wherein R¹ represents hydrogen, lower alkyl, lower alkoxy, or halogen; -X¹-X²-X³- represents -CR²=CR³-CR⁴=CR⁵- (wherein R², R³, R⁴, and R⁵, which may be the same or different, each represents hydrogen, lower alkyl, lower alkoxy, or halogen), -CR²=CR³-CR⁴=N- (wherein R², R³, and R⁴ have the same significances as defined above), -CR²=CR³-S- (wherein R² and R³ have the same significances as defined above), -CR²=CR³-O- (wherein R² and R³ have the same significances as defined above), -S-CR⁴=CR⁵- (wherein R⁴ and R⁵ have the same significances as defined above), or -O-CR⁴=CR⁵- (wherein R⁴ and R⁵ have the same significances as defined above); and Y represents -CH₂O-, -CH₂S-, -CH₂SO-, -CH=CH-, or -(CH₂)_n- (wherein n represents 0, 1, or 2).

(11) AU-B-73326/96
(10) 700386

-2-

89. A pharmaceutical composition comprising an effective amount of at least one compound according to any one of claims 1 to 88, together with a pharmaceutically acceptable carrier, diluent or adjuvant therefor.

90. A method for the treatment or prophylaxis of urinary incontinence in a mammal requiring said treatment or prophylaxis, which method comprises administering to said mammal an effective amount of at least one compound according to any one of claims 1 to 88, or of a composition according to claim 89.

PC

OPI DATE 07/05/97 APPLN. ID 73326/96
AOJP DATE 03/07/97 PCT NUMBER PCT/JP96/02997

AU9673326

(51) 国際特許分類6

C07C 235/16, C07D 221/06, 307/93,
313/12, 333/78, 337/12, 491/044, 495/04,
A61K 31/35, 31/38, 31/435, 31/44

A1

(11) 国際公開番号

WO97/14672

(43) 国際公開日

1997年4月24日(24.04.97)

(21) 国際出願番号

PCT/JP96/02997

(22) 国際出願日

1996年10月16日(16.10.96)

(30) 優先権データ

特願平7/267232

1995年10月16日(16.10.95)

JP

特願平7/324298

1995年12月13日(13.12.95)

JP

(71) 出願人 (米国を除くすべての指定国について)

協和醗酵工業株式会社

(KYOWA HAKKO KOGYO CO., LTD.)(JP/JP)

〒100 東京都千代田区大手町一丁目6番1号 Tokyo, (JP)

(72) 発明者: および

(75) 発明者/出願人 (米国についてのみ)

吉田 誠(YOSHIDA, Makoto)(JP/JP)

〒411 静岡県駿東郡長泉町納米里410-1 Shizuoka, (JP)

佐々木慎一(SASAKI, Shin-ichi)(JP/JP)

長島 健(NAGASHIMA, Ken)(JP/JP)

〒411 静岡県駿東郡長泉町下土狩1188 Shizuoka, (JP)

青野 滋(AONO, Shigeru)(JP/JP)

〒410 静岡県沼津市大岡2151-1 Shizuoka, (JP)

藤原繁喜(FUJIWARA, Shigeki)(JP/JP)

〒411 静岡県三島市老町田104-2-403 Shizuoka, (JP)

高井春樹(TAKAI, Haruki)(JP/JP)

〒411 静岡県駿東郡長泉町下土狩702-24 Shizuoka, (JP)

山形 強(YAMAGATA, Tsuyoshi)(JP/JP)

〒411 静岡県駿東郡長泉町下土狩103-7 Shizuoka, (JP)

唐沢 啓(KARASAWA, Akira)(JP/JP)

〒411 静岡県駿東郡長泉町納米里378-6 Shizuoka, (JP)


(81) 指定国 AU, BG, BR, CA, CN, CZ, HU, JP, KR, MX, NO,
NZ, PL, RO, SG, SI, SK, UA, US, VN, ユーラシア特許 (AM,
AZ, BY, KG, KZ, MD, RU, TJ, TM), 欧州特許 (AT, BE, CH, DE,
DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

添付公開書類

国際調査報告書

(54) Title: TRICICLIC COMPOUNDS

(54) 発明の名称 三環性化合物



(57) Abstract

Tricyclic compounds represented by general formula (I) which are useful as a remedy for urinary incontinence, wherein R^1 represents hydrogen, etc.; and $-X^1-X^2-X^3-$ represents $-CR^2=CR^3-CR^4=CR^5-$ (wherein R^2, R^3, R^4 and R^5 represent each hydrogen, etc.); and Y represents $-CH_2O-$ etc.

SPECIFICATION
TRICYCLIC COMPOUNDS

Technical Field


5 The present invention relates to tricyclic compounds which are useful as therapeutic agents for urinary incontinence.

Background Art

10 Japanese Published Unexamined Patent Application No. 286915/93 discloses N-substituted propanamide derivatives which are useful for the treatment of urinary incontinence.

Disclosure of the Invention

15 The present invention relates to tricyclic compounds represented by general formula (I):



wherein R^1 represents hydrogen, lower alkyl, lower alkoxy,
25 or halogen; $-X^1-X^2-X^3-$ represents $-CR^2=CR^3-CR^4=CR^5-$ (wherein R^2 , R^3 , R^4 , and R^5 , which may be the same or different, each represents hydrogen, lower alkyl, lower alkoxy, or halogen),
 $-CR^2=CR^3-CR^4=N-$ (wherein R^2 , R^3 , and R^4 have the same
significances as defined above), $-CR^2=CR^3-S-$ (wherein R^2 and
30 R^3 have the same significances as defined above), $-CR^2=CR^3-$
 $O-$ (wherein R^2 and R^3 have the same significances as defined
above), $-S-CR^4=CR^5-$ (wherein R^4 and R^5 have the same
significances as defined above), or $-O-CR^4=CR^5-$ (wherein R^4
and R^5 have the same significances as defined above); and Y
35 represents $-CH_2O-$, $-CH_2S-$, $-CH_2SO-$, $-CH=CH-$, or $-(CH_2)_n-$




(wherein n represents 0, 1, or 2) [which are hereinafter referred to as Compound (I), and the same applies to the compounds of other formula numbers].

In the definitions of the groups in general formula (I), the lower alkyl and the lower alkyl moiety of the lower alkoxy mean a straight-chain or branched alkyl group having 1 to 6 carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, and hexyl. The halogen includes fluorine, chlorine, bromine, and iodine.

The processes for preparing Compound (I) are described below.

Step 1

Compound (I) can be prepared according to the following reaction step.



(In the formulae, R^1 , $-X^1-X^2-X^3-$, and Y have the same significances as defined above.)

Compound (I) can be obtained by treating Compound (II) with an equivalent amount of halogenating agent such as thionyl chloride and oxalyl chloride in an inert solvent such as dichloromethane, dichloroethane, tetrachloroethane and dimethylacetamide at -30 to 0°C for 5 minutes to 12 hours, and then subjecting the obtained substance to reaction with Compound (III) at a temperature between 0°C and the boiling point of the solvent used for 5 minutes to 24 hours. This reaction is preferably carried out under an




atmosphere of dry inert gas such as argon gas and nitrogen gas.

The starting Compound (II) can be obtained according to the process described in J. Chem. Soc., 2329-2332 (1951).

- 5 The starting Compound (III) may be commercially available, or can be prepared according to the processes described in the following Steps 2 through 4.

Step 2

- 10 Compound (III) can be prepared according to the following reaction steps.



- 25 (In the formulae, R^1 , $-X^1-X^2-X^3-$, and Y have the same significances as defined above.)

The starting Compound (IV) can be obtained according to the known methods [J. Med. Chem., 19, 941 (1976); *ibid.*, 20, 66 (1977); *ibid.*, 20, 1499 (1977); *ibid.*, 20, 1557

- 30 (1977); *ibid.*, 21, 633 (1978); *ibid.*, 22, 1357 (1979); *ibid.*, 27, 372 (1984); *ibid.*, 29, 2347 (1986); *ibid.*, 29, 2074 (1992); *ibid.*, 38, 496 (1995), etc.] or similar methods thereto.

- Compound (V) can be obtained by treating Compound (IV) according to the methods described in Synthesis, 295 (1990), etc. in tert-butanol in the presence of an equivalent amount






of diphenylphosphoryl azide under an atmosphere of dry inert gas such as argon gas and nitrogen gas.

Compound (III) can be obtained by hydrolyzing Compound (V) in an appropriate solvent such as trifluoroacetic acid, hydrochloric acid/dioxane and hydrobromic acid/acetic acid at a temperature between 0°C and the boiling point of the solvent used for 1 to 24 hours.

Step 3

Compound (IIIa), which is Compound (III) in which Y is -CH₂O-, can also be prepared according to the following reaction steps.



(In the formulae, R^1 and $-X^1-X^2-X^3-$ have the same significances as defined above.)

The starting Compound (VI) can be obtained by heating under reflux the corresponding starting material which is commercially available such as methyl 2-methylbenzoate and methyl 2-methylnicotinate in a solvent such as carbon tetrachloride in the presence of a radical initiator such as azobisisobutyronitrile (AIBN) and an equivalent amount of a brominating agent such as N-bromosuccinimide (NBS) for 1 to 24 hours according to a known method.

Compound (VII) can be obtained by reacting Compound (VI) with commercially available acetamidophenol such as 3-acetamidophenol in an inert solvent such as tetrahydrofuran, dimethylformamide, dimethylacetamide and dimethyl sulfoxide in the presence of an appropriate base such as cesium carbonate at a temperature between 0°C and the boiling point of the solvent used for 1 to 24 hours.

Compound (VIII) can be obtained by hydrolyzing Compound (VII) in a mixed solvent of water and a solvent such as methanol, ethanol, dioxane and tetrahydrofuran in the presence of an appropriate base such as lithium hydroxide, sodium hydroxide and potassium hydroxide at a temperature between room temperature and the boiling point of the solvent used for 1 to 24 hours; treating the obtained carboxylic acid with an equivalent amount of trifluoroacetic anhydride in an inert solvent such as dichloromethane, tetrahydrofuran, dimethylformamide, dimethylacetamide and dimethyl sulfoxide at a temperature between -15°C and room temperature according to a known method (Japanese Published Unexamined Patent Application No. 91040/90 etc.) to give acid anhydride; and then without isolation subjecting the obtained acid anhydride to ring closure reaction in the same solvent in the presence of 0.1 to one equivalent of Lewis acid such as boron trifluoride diethyl etherate at a temperature between 0°C and the boiling point of the solvent used.





Compound (IIIa) can be obtained by heating Compound (VIII) under reflux in concentrated hydrochloric acid etc.

Compound (IIIb), which is Compound (III) in which Y is $-\text{CH}_2\text{S}-$, can also be obtained according to the method similar to that in Step 3 except for using acetamidothiophenol such as 3-acetamidothiophenol instead of acetamidophenol.

Step 4

Compound (IIIc), which is Compound (III) in which Y is $-(\text{CH}_2)_2-$, and Compound (IIId), which is Compound (III) in which Y is $-\text{CH}=\text{CH}-$, can also be prepared according to the following reaction steps.





(In the formulae, R^7 and R^8 represent lower alkyl; and R^1 and $-X^1-X^2-X^3-$ have the same significances as defined above.)

In the definitions of R^7 and R^8 , the lower alkyl has the same significance as the lower alkyl defined above.

The starting Compound (IX) can be obtained by reacting Compound (VI) with an equivalent amount of triphenylphosphine in a solvent such as toluene at a temperature between 0°C and the boiling point of the solvent used for 1 to 24 hours according to a known method.

Compound (X) can be obtained by reacting Compound (IX) with commercially available nitrobenzaldehyde such as 3-nitrobenzaldehyde in an inert solvent such as tetrahydrofuran, dioxane and ether in the presence of an appropriate base such as sodium hydride, potassium tert-butoxide and potassium carbonate at a temperature between 0°C and the boiling point of the solvent used for 1 to 24 hours.

Compound (XI) can be obtained as an E/Z mixture by reducing Compound (X) with an appropriate reducing agent such as reduced iron in a mixed solvent of water and a solvent such as methanol and ethanol at a temperature



between room temperature and the boiling point of the solvent used for 1 to 24 hours, and then reacting the obtained amine with an acylating agent such as acetic anhydride and acetyl chloride in an inert solvent such as
5 dichloromethane, tetrahydrofuran, dioxane and dimethylformamide in the presence of a base such as triethylamine and diisopropylethylamine at a temperature between 0°C and the boiling point of the solvent used for 1 to 24 hours.

10 Compound (XII) can be obtained by subjecting Compound (XI) to catalytic reduction with hydrogen in a solvent such as methanol, ethanol, dioxane and tetrahydrofuran in the presence of an appropriate catalyst such as palladium and platinum at a temperature between room temperature and the
15 boiling point of the solvent used for 1 to 24 hours, hydrolyzing the reduction product according to the method similar to that in Step 3 to give a carboxylic acid, and then subjecting the obtained carboxylic acid to ring closure reaction according to the method similar to that in Step 3.

20 Compound (IIIc) can be obtained by heating Compound (XII) under reflux in concentrated hydrochloric acid etc.

Compound (XIII) can be obtained by hydrolyzing Compound (XIa), which is Z-form within Compound (XI), according to the method similar to that in Step 3 to give a
25 carboxylic acid, and then subjecting the obtained carboxylic acid to ring closure reaction according to the method similar to that in Step 3.

Compound (IIId) can be obtained by heating Compound (XIII) under reflux in concentrated hydrochloric acid etc.

30 Compound (IIId) can also be prepared according to the known methods [J. Med. Chem., 20, 1557 (1977) etc.] or similar methods thereto.

Compound (XIII) can be obtained by brominating Compound (XII) by heating it under reflux in a solvent such
35 as carbon tetrachloride in the presence of a radical initiator such as AIBN and an equivalent amount of a

brominating agent such as NBS for 1 to 24 hours, and then
subjecting the obtained compound to dehydrobromination in an
appropriate inert solvent such as dimethylformamide in the
presence of 1,8-diazabicyclo[5.4.1]undec-7-ene (DBU) or the
5 like at a temperature between room temperature and 120°C.
Compound (IIId) can be obtained by hydrolyzing the obtained
Compound (XIII) with an acid in the same manner as described
above.


Compound (Ia), which is Compound (I) in which Y is -
10 $\text{CH}_2\text{SO}-$, can be obtained by treating Compound (Ib), which is
Compound (I) in which Y is $-\text{CH}_2\text{S}-$, with an equivalent amount
of an oxidizing agent such as 3-chloroperbenzoic acid in a
solvent such as dichloromethane at a temperature between 0°C
and the boiling point of the solvent used for 1 to 24 hours
15 according to a known method.

The intermediates and the desired compounds in the
processes described above can be isolated and purified by
purification methods conventionally used in synthetic
organic chemistry, for example, filtration, extraction,
20 washing, drying, concentration, recrystallization, and
various kinds of chromatography. The intermediates may be
subjected to the subsequent reaction without purification.
In the case where the intermediates are obtained in the form
of an E/Z mixture and its separation is desired, it may be
25 separated and purified by fractionation methods such as
fractional crystallization, fractional precipitation, and
fractional dissolution, various kinds of chromatography, and
the like.

Compound (I) may exist in the form of adducts with
30 water or various solvents, which are also within the scope
of the present invention.


Representative examples of Compound (I) are shown
below in Table 1.

Table 1



Compd. No.	R	Compd. No.	R
1		9	
2		10	
3		11	
4		12	
5		13	
6		14	
7		15	
8		16	


Table 1 (continued)



Compd. No.	R	Compd. No.	R
17		25	
18		26	
19		27	
20		28	
21		29	
22		30	
23		31	
24		32	



Table 1 (continued)




Compd. No.	R	Compd. No.	R
33*		41	
34**		42	
35*		43	
36**		44	
37*		45	
38**		46	
39*		47	
40		48	

*; (S) form, **; (R) form




Table 1 (continued)



Compd. No.	R	Compd. No.	R
49		57	
50		58	
51		59	
52		60	
53		61	
54		62	
55		63	
56		64	

Table 1 (continued)



Compd. No.	R	Compd. No.	R
65		72	
66		73	
67		74	
68		75	
69		76	
70		77	
71			



The activity of Compound (I) as a smooth muscle relaxant which is useful as a therapeutic agent for urinary incontinence can be observed, for example, using the following tests. In the following Test Examples 1 and 2,
5 "IC₅₀" means the concentration of a test compound which induces the decrease in the contraction of detrusor muscle by 50%.

Test Example 1

10 Male albino Hartley guinea pigs (350-550 g) were used as test animals. Each animal was exsanguinated to death under pentobarbital sodium anesthesia. The lower abdominal cavity was opened and its bladder was positioned. The connective tissue and the fat tissue around the bladder were
15 removed, and the bladder was cleaned. After two pelvic nerves on the surface of the abdominal side of the bladder were dissected, the bladder was taken out on the entrance of ureter. The bladder was washed with a Krebs-Henseleit buffer [composition (mM): NaCl 118, KCl 4.7, MgSO₄ 1.2,
20 KH₂PO₄ 1.2, CaCl₂ 2.5, NaHCO₃ 25, and D-glucose 11.1], and then put on a gauze patch soaked in the buffer in a Petri dish. The dome and the trigone of the bladder were dissected and discarded. The back of the bladder was cut vertically in the middle with scissors, and the bladder was
25 put flat on the gauze patch. The dome tip and the side of the trigone of the bladder were dissected and discarded, and the mucosa of the bladder was removed. The back of the bladder was cut vertically to get 6 strips which are ca. 2.0 mm wide and ca. 10 mm long.

30 One end of each strip was fixed on a plastic supporter rod, and the other end was fixed on a clip connected to a suture thread. The supporter was fixed in 20 ml of an organ bath and the suture thread connected to the clip was fixed on a force-displacement transducer (NIHON KODEN TB-611T).

35 The tissue was suspended in a Krebs-Henseleit buffer, and the bath solution was warmed to 37°C and bubbled with a gas



mixture of 5% CO₂ and 95% O₂. The pH of the solution was adjusted to about 7.4. The transducer was connected to a polygraph (NIHON KODEN AP-621G), and the change of tension was recorded on a recorder (YOKOGAWA type 3066). The polygraph was graduated by 0.5 g/cm.

After the tissue was incubated in the buffer for 15 minutes without preloading, tension was added thereto for about one hour. The preloading tension added was 1.5 g and it was relaxed to about 1 g. The tissue was washed at intervals of 15 minutes and the tension was adjusted to 1.5 g just before washing. The tension of the tissue was adjusted to be a steady state at 1 g. After this equilibrium time, 50 mM KCl (the whole content in the bath) was added, and the tissue was washed 10 minutes later. KCl was added to the tissue at intervals of 30 minutes, followed by washing 10 minutes later, and KCl was applied repeatedly until the developed tension induced by KCl becomes constant. When the tissue was confirmed to contract constantly by the application of KCl and relax to the steady state after washing, 50 mM KCl was applied again. After the tissue reached the steady state and the base line was obtained, a test compound was applied by the cumulative method (half-logarithmic increment). The contact time at each concentration was 30 minutes. The activity of the compound was expressed as the maximum relaxation rate (%) of the tension induced by the agonist.

The results are shown in Table 2.

The effect of a smooth muscle relaxant varies with the concentration of KCl based on the difference of an action mechanism. In the following Test Example, KCl was used at a concentration different from that in Test Example 1.

Test Example 2

A sample of the excised bladder was prepared in the same manner as in Test Example 1, and the tension of the preparation was measured. After the tension of the tissue



was adjusted to be a steady state at 1 g, 15 mM KCl was applied. After the rhythmical contraction was confirmed to appear by KCl, the tissue was washed. Then, 15 mM KCl was applied again. After the rhythmical contraction took place
5 constantly and the tissue reached the steady state, a test compound was applied by the cumulative method (half-logarithmic increment). The contact time at each concentration was 30 minutes. The activity of the compound was expressed as the maximum relaxation rate (%) of the
10 tension induced by the agonist.

The results are shown in Table 2.



Table 2

Compound			Compound		
IC ₅₀ (μM)			IC ₅₀ (μM)		
No.	KCl	KCl	No.	KCl	KCl
	50 mM	15 mM		50 mM	15 mM
1	1.7	2.4	35	31	1.0
2	3.6	3.4	36	8.3	1.3
3	2.2	1.8	37	4.3	0.2
4	2.6	8.4	38	13.4	0.9
5	8.0	2.1	39	>10.0	2.6
6	6.1	11.2	40	1.5	3.5
7	2.4	2.1	41	2.1	1.6
8	5.2	0.5	42	4.7	1.5
9	4.3	4.6	43	3.5	2.1
10	10.0	1.9	45	>10.0	2.5
11	2.0	2.2	47	NT	6.1
12	3.8	3.2	49	7.8	4.1
13	3.5	48	50	NT	6.1
14	5.7	2.3	51	NT	5.7
15	7.1	9.4	53	>10.0	2.9
16	>10.0	1.4	54	3.8	>10.0
18	>10.0	2.1	59	>10.0	5.0
20	2.3	6.0	60	>10.0	4.1
21	4.3	>10.0	61	3.2	2.3
22	4.2	>10.0	62	9.3	>10.0
23	>10.0	7.8	63	3.3	>10.0
25	2.2	>10.0	65	4.6	2.3
26	7.1	>10.0	66	1.4	2.1
27	4.8	>10.0	67	6.9	3.8
33	1.1	2.7	68	>10.0	3.5
34	1.6	2.2			

NT; not tested

The effect of the compounds of the present invention
 5 can also be proved by the standard examination method
 described below which is used to evaluate the smooth muscle
 relaxing activity.



Test Example 3

Female SD strain rats (120-180 g) were used as test animals. Each animal was anesthetized by subcutaneous administration of 1 g/kg urethane. A catheter was inserted into the trachea to secure respiration. A polyethylene cannula (PE50) was inserted into the bladder from the entrance of the urethra and the urethra was ligated at the base. A catheter for blood pressure measurement was inserted into the right carotid artery, and a cannula for drug administration was inserted into the right jugular vein. The arterial catheter and the vesical catheter were connected to pressure transducers to measure the blood pressure and the intravesical pressure. After the bladder was stabilized over about 15 minutes, physiological saline was injected into the bladder at a volume of 0.2-0.7 ml to induce the rhythmical contraction almost once a minute. After the contraction of the bladder started, the contraction pattern was stabilized for 30-45 minutes, followed by intravenous administration of a test compound.

By this examination method, the effect of the test compounds on both blood pressure and vesical contraction can be evaluated. The blood pressure was measured 5, 15, and 30 minutes after the test compound was injected into the vein. Micturition contraction was induced by injecting physiological saline into the bladder. The change of average contraction interval (time between contractions) over 20 minutes was expressed as the relative value based on the pre-administration value which was regarded as 100.

The results are shown in Table 3.

Table 3

Compound No.	Dose (mg/kg)	Change of average arterial blood pressure (%)			Change of contraction interval (%)
		5 min.	15 min.	30 min.	
1	1	+5	+5	+5	+150
7	1	-4	+1	+4	+186



The following is a test which is useful to prove that a test compound shows the selectivity to the bladder without exhibiting cardiovascular activity when orally administered.

5

Test Example 4

Male SD strain rats (200-300 g) were used as test animals. Each animal was anesthetized by intraperitoneal administration of 50 mg/kg Nembutal. The hair on the abdomen, the thigh, and the back of the neck was shaved, followed by disinfection with 70% ethanol. The left femoral artery of the rat was exposed by incising the thigh for the insertion of a catheter into the femoral artery. A catheter filled with physiological saline containing heparin (1000 units/ml) was introduced into the artery by 2-3 cm to position the tip of the catheter in the abdominal artery. The end of the catheter was exposed outside the body from the back of the neck, stoppered, and fixed on the skin tightly. A catheter for the bladder was inserted according to the method of Yaksh, T.L., Durant, P.A.C. and Brent, C.R. [(Micturition in rats: A chronic model for study of bladder function and effect of anesthesia. Am. J. Physiol., 251, R1177-1185 (1986))]. The bladder was exposed by middle incision. The bladder dome was given a small incision to make a small hole. A catheter filled with physiological saline was inserted into the bladder and was fixed thereto by ligation with a silk thread. The other end of the catheter was exposed outside the body from the back of the neck of the rat after being passed under the skin using a trocar. The exposed end of the catheter was stoppered and fixed on the skin tightly. The abdominal muscles and skin were sutured, and the test animal was recovered from anesthesia.

The rat was weighed 24-48 hours after the operation, put in a Bollman cage (NATSUME SEISAKUSHO), and subjected to the test under partial restraint. The arterial catheter was



connected to a pressure transducer (NIHON KODEN DX-300) used for the measurement of blood pressure. The vesical catheter was connected to a pump for injecting physiological saline and a pressure transducer with PE50 and a three-way stop-cock. Injection of physiological saline (0.15 ml/minute) through the bladder was started and continued during the test. The changes of blood pressure, heart rate, and intravesical pressure were recorded with a polygraph (NIHON KODEN). The test animal was equilibrated (ca. 60-120 minutes) until its micturition pattern was constant. At this point, the basal value of each test parameter was recorded, and the rat was compulsorily given a test compound (in 0.3% carboxymethyl cellulose-physiological saline) at a volume of 1 ml/kg (body weight) by oral administration. The effects of the compound on the test parameters were observed for 5 hours after the administration. As a control compound, cromakalim was used. The contraction interval, the systemic blood pressure, and the heart rate were measured for 5 hours after the administration of the compound at one-hour intervals and were expressed as the relative values based on the pre-administration values which were regarded as 100.

The results are shown in Tables 4 and 5. As for the compounds other than Compounds 1 and 7, the results are shown only in respect of the change of the contraction interval.



Table 4

Compound	Time (hr)	Contraction interval	Blood pressure	Heart rate
Solvent alone 1 ml/kg	0	100	100	100
	1	79	101	98
	2	100	101	100
	3	110	103	98
	4	104	100	99
	5	107	100	100

Compound	Time (hr)	Contraction interval	Blood pressure	Heart rate
Croma-kalim 1 mg/kg	0	100	100	100
	1	128	46	98
	2	133	51	93
	3	142	54	96
	4	133	60	97
	5	133	64	102

Compound	Time (hr)	Contraction interval	Blood pressure	Heart rate
Compound 1 3 mg/kg	0	100	100	100
	1	136	98	98
	2	148	99	98
	3	134	96	96
	4	122	91	93
	5	117	91	89

Compound	Time (hr)	Contraction interval	Blood pressure	Heart rate
Compound 7 3 mg/kg	0	100	100	100
	1	132	100	103
	2	141	96	103
	3	149	99	103
	4	154	101	106
	5	146	101	105

Compound	Dose (mg/kg)	Time after administ- ration (hr)	Contraction interval
Compound 9	1.0	1	123.4±6.8
		2	131.2±9.3
		3	132.9±9.1
		4	145.0±10.0
		5	154.9±12.2
Compound 10	1.0	1	132.4±11.2
		2	142.0±17.5
		3	158.4±22.3
		4	173.8±29.9
		5	182.7±42.7
Compound 8	1.0	1	130.3±9.3
		2	142.1±12.6
		3	144.3±8.8
		4	162.4±14.6
		5	189.4±19.6
Compound 33	0.1	1	135.1±21.7
		2	145.6±25.5
		3	143.4±24.4
		4	156.3±30.6
		5	160.6±37.4
Compound 37	1.0	1	113.5±10.7
		2	127.9±6.5
		3	126.5±6.5
		4	140.4±10.2
		5	143.6±11.1
Compound 35	1.0	1	123.2±8.8
		2	123.8±5.7
		3	130.7±9.8
		4	137.9±11.0
		5	143.6±8.9



Compound	Dose (mg/kg)	Time after administ- ration (hr)	Contraction interval
Compound 2	1.0	1	120.9±6.2
		2	135.4±12.3
		3	144.5±16.3
		4	140.8±15.6
		5	144.9±16.0
Compound 5	1.0	1	111.6±7.0
		2	135.5±16.5
		3	137.4±14.5
		4	131.5±9.5
		5	130.8±17.7
Compound 15	1.0	1	129.9±9.5
		2	127.8±5.3
		3	107.3±16.5
		4	120.7±11.4
		5	134.6±12.6
Compound 16	1.0	1	118.0±7.9
		2	132.3±14.2
		3	119.2±9.8
		4	126.9±7.1
		5	136.8±12.6
Compound 20	1.0	1	121.2±3.9
		2	146.6±9.9
		3	148.4±20.0
		4	155.8±19.6
		5	142.2±15.8
Compound 53	1.0	1	108.4±9.7
		2	134.5±12.0
		3	148.4±14.0
		4	155.3±11.9
		5	167.9±11.8



Compound (I) can be formulated into generally employed dose forms such as tablets, capsules and syrups, and administered orally or parenterally through intramuscular injection, intravenous injection, drip infusion, or rectal administration in the form of suppositories. For preparing these dose forms for oral or parenteral administration, generally known methods are applied. For example, the preparations may be formulated to contain various excipients, lubricants, binders, disintegrating agents, isotoning agents, emulsifiers, and the like.

Examples of the carriers which can be used are water, injectable distilled water, physiological saline, glucose, fructose, sucrose, mannitol, lactose, starch, cellulose, methyl cellulose, sulfoxymethyl cellulose, hydroxypropyl cellulose, alginic acid, talc, sodium citrate, calcium carbonate, calcium hydrogenphosphate, magnesium stearate, urea, silicone resins, sorbitan fatty acid esters, and glycerin fatty acid esters.

The effective dose and the administration schedule of Compound (I) vary depending upon the mode of administration, the age and body weight of a patient, and the type or degree of the disease to be treated, but generally, in the case of oral administration, Compound (I) is administered in a dose of 0.01 mg to 1 g/adult/day, preferably 0.05 to 50 mg/adult/day, in one to several parts. In the case of parenteral administration such as intravenous injection, Compound (I) is administered in a dose of 0.001 to 100 mg/adult/day, preferably 0.01 to 10 mg/adult/day, in one to several parts. It should, however, be noted that the dose may vary depending upon various conditions as given above.

Certain embodiments of the present invention are illustrated in the following Examples and Reference Examples.



Best Mode for Carrying Out the InventionExample 1

1-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
6,11-dihydrodibenz[b,e]oxepin-11-one (Compound 1)

5 In dimethylacetamide (3 ml) was dissolved 3,3,3-trifluoro-2-hydroxy-2-methylpropionic acid (0.17 g, 1.1 mmol), and thionyl chloride (80 μ l, 1.1 mmol) was added thereto at -15°C , followed by stirring at -15 to -5°C for one hour. To this reaction mixture was added 1-amino-6,11-dihydrodibenz[b,e]oxepin-11-one obtained in Reference
10 Example 4 (0.17 g, 0.77 mmol), and the mixture was stirred overnight at room temperature. After concentration of the reaction mixture under reduced pressure, the obtained oily residue was dissolved in ethyl acetate (20 ml). The organic
15 layer was washed successively with a 5% aqueous solution of sodium bicarbonate and a saturated aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure. The
20 obtained oily residue was purified by silica gel column chromatography (hexane/ethyl acetate = 5/1), followed by trituration with hexane to give Compound 1 (80 mg, 30%).

$^1\text{H-NMR}$ (CDCl_3) δ : 1.61(s, 3H), 4.47(br, 1H), 5.18(s, 2H), 6.89(d, 1H, $J=7.9\text{Hz}$), 7.34(d, 1H, $J=7.9\text{Hz}$),
25 7.45-7.54(m, 3H), 7.86(d, 1H, $J=7.9\text{Hz}$), 8.42(d, 1H, $J=7.9\text{Hz}$), 12.88(br, 1H)

In the following Examples 2 to 6, substantially the same procedure as in Example 1 was repeated except that the
30 corresponding tricyclic aromatic amine was used instead of 1-amino-6,11-dihydrodibenz[b,e]oxepin-11-one, to give the desired compound.



Example 2

3-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
6,11-dihydrodibenz[b,e]oxepin-11-one (Compound 2)
(yield: 45%)

5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.62(s, 3H), 3.69(br, 1H), 5.20(s,
2H), 7.19(d, 1H, $J=8.1\text{Hz}$), 7.37(d, 1H, $J=8.1\text{Hz}$),
7.45-7.60(m, 3H), 7.92(d, 1H, $J=8.1\text{Hz}$), 8.25(d,
1H, $J=8.1\text{Hz}$), 8.52(br, 1H)

10 Example 3

2-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
6,11-dihydrodibenz[b,e]oxepin-11-one (Compound 3)
(yield: 45%)

15 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.59(s, 3H), 5.27(s, 2H), 7.06(d,
1H, $J=7.9\text{Hz}$), 7.36(s, 1H), 7.50-7.56(m, 2H),
7.63-7.68(m, 1H), 7.79(d, 1H, $J=7.9\text{Hz}$), 7.86(s,
1H), 10.10(br, 1H)

Example 4

20 1-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
fluoren-9-one (Compound 4) (yield: 48%)

$^1\text{H-NMR}$ (DMSO-d_6) δ : 1.62(s, 3H), 7.39(t, 1H, $J=7.5\text{Hz}$),
7.50(d, 1H, $J=7.5\text{Hz}$), 7.57-7.63(m, 3H), 7.78(d,
1H, $J=7.5\text{Hz}$), 7.92(s, 1H), 8.32(d, 1H, $J=7.5\text{Hz}$),
25 11.36(br, 1H)

Example 5

2-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
fluoren-9-one (Compound 5) (yield: 90%)

30 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.62(s, 3H), 7.31(t, 1H, $J=7.8\text{Hz}$),
7.41(br, 1H), 7.53(d, 2H, $J=7.8\text{Hz}$), 7.67(d, 2H,
 $J=7.8\text{Hz}$), 7.93(d, 1H, $J=7.8\text{Hz}$), 8.09(s, 1H),
10.17(br, 1H)



Example 6

3-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
fluoren-9-one (Compound 6) (yield: 67%)

¹H-NMR (DMSO-d₆) δ: 1.64(s, 3H), 7.36(d, 1H, J=7.1Hz),
5 7.48-7.58(m, 3H), 7.62(d, 1H, J=7.1Hz), 7.74(d,
 1H, J=7.1Hz), 8.21(s, 1H), 10.21(br, 1H)

Example 7

9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
10 4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one
 (Compound 7)

In dimethylacetamide (3 ml) was dissolved 3,3,3-
trifluoro-2-hydroxy-2-methylpropionic acid (0.32 g, 2.03
mmol), and thionyl chloride (148 μl, 2.03 mmol) was added
15 thereto at -15°C, followed by stirring at -15 to -5°C for
one hour. To this reaction mixture was added 9-amino-4,10-
dihydrothieno[3,2-c][1]benzoxepin-10-one obtained in
Reference Example 11 (0.23 g, 1.02 mmol), and the mixture
was stirred overnight at room temperature. After
20 concentration of the reaction mixture under reduced
pressure, the obtained oily residue was dissolved in ethyl
acetate (20 ml). The organic layer was washed successively
with a 5% aqueous solution of sodium bicarbonate and a
saturated aqueous solution of sodium chloride, and then
25 dried over anhydrous magnesium sulfate. After the drying
agent was filtered off, the organic layer was concentrated
under reduced pressure. The obtained oily residue was
purified by silica gel column chromatography (hexane/ethyl
acetate = 5/1), followed by trituration with hexane to give
30 Compound 7 (0.34 g, 89%).

¹H-NMR (CDCl₃) δ: 1.77(s, 3H), 4.32(s, 1H), 5.16(s,
2H), 7.00-7.04(m, 2H), 7.54(t, 1H, J=8.4Hz),
7.72(d, 1H, J=5.0Hz), 8.41(d, 1H, J=8.4Hz),
12.24(br, 1H)



In the following Examples 8 to 32, 35 to 51, and 69 to 77, substantially the same procedure as in Example 7 was repeated except that the corresponding tricyclic aromatic amine was used instead of 9-amino-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one, to give the desired compound.

Example 8

7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one
(Compound 8) (yield: 64%)

¹H-NMR (DMSO-d₆) δ: 1.60(s, 3H), 5.29(s, 2H), 7.22(d, 1H, J=5.0Hz), 7.52(s, 1H), 7.68(d, 1H, J=8.6Hz), 7.75(s, 1H), 7.98-8.00(m, 2H), 10.31(br, 1H)

15 Example 9

5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrofuro[2,3-c][1]benzoxepin-4-one (Compound 9) (yield: 67%)

¹H-NMR (CDCl₃) δ: 1.77(s, 3H), 4.28(s, 1H), 5.17(s, 2H), 6.96(d, 1H, J=2.0Hz), 7.03(d, 1H, J=8.3Hz), 7.39(d, 1H, J=2.0Hz), 7.48(t, 1H, J=8.3Hz), 8.42(d, 1H, J=8.3Hz), 12.13(br, 1H)

Example 10

25 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrofuro[2,3-c][1]benzoxepin-4-one (Compound 10) (yield: 75%)

¹H-NMR (DMSO-d₆) δ: 1.62(s, 3H), 5.28(s, 2H), 6.90(d, 1H, J=2.0Hz), 7.40(s, 1H), 7.62-7.69(m, 2H), 7.76(s, 1H), 7.96(d, 1H, J=8.9Hz), 10.13(br, 1H)

Example 11

3-Chloro-2-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoylamino)fluoren-9-one (Compound 11) (yield:



59%)

$^1\text{H-NMR}$ (DMSO- d_6) δ : 1.63(s, 3H), 7.39(t, 1H, $J=7.7\text{Hz}$),
7.59-7.65(m, 2H), 7.82(d, 1H, $J=7.7\text{Hz}$), 8.07(s,
1H), 8.26(d, 1H, $J=7.7\text{Hz}$)

5

Example 12

3-Bromo-2-(3,3,3-trifluoro-2-hydroxy-2-methyl-
propanoylamino)fluoren-9-one (Compound 12) (yield:
90%)

10 $^1\text{H-NMR}$ (DMSO- d_6) δ : 1.64(s, 3H), 7.38(t, 1H, $J=7.3\text{Hz}$),
7.59-7.65(m, 2H), 7.82(d, 1H, $J=7.3\text{Hz}$), 8.22(s,
1H), 8.26(s, 1H), 9.83(br, 1H)

Example 13

15 7-Bromo-2-(3,3,3-trifluoro-2-hydroxy-2-methyl-
propanoylamino)fluoren-9-one (Compound 13) (yield:
98%)

$^1\text{H-NMR}$ (DMSO- d_6) δ : 1.59(s, 3H), 7.43(br, 1H), 7.66-
7.77(m, 4H), 7.96(d, 1H, $J=8.2\text{Hz}$), 8.10(s, 1H),
20 10.23(br, 1H)

Example 14

4-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
fluoren-9-one (Compound 14) (yield: 75%)

25 $^1\text{H-NMR}$ (DMSO- d_6) δ : 1.67(s, 3H), 7.36-7.42(m, 2H),
7.52-7.68(m, 5H), 10.24(br, 1H)

Example 15

30 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrofuro[3,2-c][1]benzoxepin-10-one
(Compound 15) (yield: 84%)

$^1\text{H-NMR}$ (CDCl_3) δ : 1.60(s, 3H), 4.32(br, 1H), 5.11(s,
2H), 6.53(d, 1H, $J=2.0\text{Hz}$), 7.03(d, 1H, $J=8.3\text{Hz}$),



7.54 (t, 1H, J=8.3Hz), 7.71 (d, 1H, J=2.0Hz), 8.46
(d, 1H, J=8.3Hz), 12.35 (br, 1H)

Example 16

5 8-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one
(Compound 16) (yield: 55%)
¹H-NMR (CDCl₃) δ: 1.60 (s, 3H), 5.26 (s, 2H), 7.15 (d, 1H,
J=8.9Hz), 7.21 (d, 1H, J=5.0Hz), 7.35 (br, 1H),
10 7.93 (d, 1H, J=8.9Hz), 7.98 (d, 1H, J=5.0Hz), 8.42
(s, 1H), 10.14 (br, 1H)

Example 17

15 5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrothieno[2,3-c][1]benzoxepin-4-one
(Compound 17) (yield: 65%)
¹H-NMR (CDCl₃) δ: 1.78 (s, 3H), 4.33 (s, 1H), 5.29 (s,
2H), 7.01 (dd, 1H, J=8.3, 1.3Hz), 7.20 (d, 1H, J=
5.3Hz), 7.53 (t, 1H, J=8.9Hz), 7.72 (d, 1H, J=
20 5.3Hz), 8.39 (dd, 1H, J=8.3, 1.3Hz), 12.30 (br, 1H)

Example 18

25 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrothieno[2,3-c][1]benzoxepin-4-one
(Compound 18) (yield: 27%)
¹H-NMR (DMSO-d₆) δ: 1.59 (s, 3H), 5.43 (s, 2H), 7.54 (s,
2H), 7.59 (br, 1H), 7.69 (m, 2H), 8.09 (d, 1H, J=
8.6Hz), 10.37 (br, 1H)

30 Example 19

6-Bromo-9-(3,3,3-trifluoro-2-hydroxy-2-methyl-
propanoylamino)-4,10-dihydrothieno[3,2-c][1]-
benzoxepin-10-one (Compound 19) (yield: 36%)



$^1\text{H-NMR}$ (CDCl_3) δ : 1.76(s, 3H), 4.19(br, 1H), 5.24(s, 2H), 7.01(d, 1H, $J=5.3\text{Hz}$), 7.75(d, 1H, $J=5.3\text{Hz}$), 7.78(d, 1H, $J=9.1\text{Hz}$), 8.34(d, 1H, $J=9.1\text{Hz}$), 11.94 (br, 1H)

5

Example 20

7-Fluoro-1-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one (Compound 20) (yield: 87%)

10 $^1\text{H-NMR}$ (CDCl_3) δ : 1.81(s, 3H), 4.37(br, 1H), 5.30(s, 2H), 6.92(d, 1H, $J=8.3\text{Hz}$), 7.31(t, 1H, $J=8.3\text{Hz}$), 7.34-7.48(m, 1H), 7.53(t, 1H, $J=8.3\text{Hz}$), 7.66(d, 1H, $J=8.3\text{Hz}$), 8.40(d, 1H, $J=8.3\text{Hz}$), 12.52(br, 1H)

15 Example 21

7-Fluoro-3-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one (Compound 21) (yield: 96%)

20 $^1\text{H-NMR}$ (CDCl_3) δ : 1.76(s, 3H), 3.83(br, 1H), 5.32(s, 2H), 7.22(d, 1H, $J=8.6\text{Hz}$), 7.26-7.34(m, 1H), 7.39-7.47(m, 1H), 7.51(s, 1H), 7.67(d, 1H, $J=8.6\text{Hz}$), 8.21(d, 1H, $J=8.6\text{Hz}$), 8.58(br, 1H)

Example 22

25 3-Methyl-1-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one (Compound 22) (yield: 92%)

30 $^1\text{H-NMR}$ (CDCl_3) δ : 1.81(s, 3H), 2.24(s, 3H), 4.46(br, 1H), 5.21(s, 2H), 7.34(d, 1H, $J=7.4\text{Hz}$), 7.39(d, 1H, $J=8.6\text{Hz}$), 7.45(t, 1H, $J=7.4\text{Hz}$), 7.58(t, 1H, $J=7.4\text{Hz}$), 7.87(d, 1H, $J=7.4\text{Hz}$), 8.26(d, 1H, $J=8.6\text{Hz}$), 12.57(br, 1H)



Example 23

3-Methoxy-1-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one
(Compound 23) (yield: 30%)

5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.80(s, 3H), 3.91(s, 3H), 4.50(br, 1H), 5.28(s, 2H), 7.13(d, 1H, $J=9.2\text{Hz}$), 7.32(t, 1H, $J=7.6\text{Hz}$), 7.47(t, 1H, $J=7.6\text{Hz}$), 7.57(t, 1H, $J=7.6\text{Hz}$), 7.85(d, 1H, $J=7.6\text{Hz}$), 8.31(d, 1H, $J=9.2\text{Hz}$), 12.52 (br, 1H)

10

Example 24

2-Methyl-1-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one
(Compound 24) (yield: 10%)

15 $^1\text{H-NMR}$ (CDCl_3) δ : 1.76(s, 3H), 2.19(s, 3H), 4.28(br, 1H), 5.24(s, 2H), 7.01(d, 1H, $J=8.3\text{Hz}$), 7.25(d, 1H, $J=7.3\text{Hz}$), 7.35(d, 1H, $J=8.3\text{Hz}$), 7.44(t, 1H, $J=7.6\text{Hz}$), 7.55(t, 1H, $J=7.6\text{Hz}$), 7.95(d, 1H, $J=7.6\text{Hz}$), 9.84(br, 1H)

20

Example 25

2-Methyl-3-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one
(Compound 25) (yield: 14%)

25 $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ : 1.76(s, 3H), 2.28(s, 3H), 3.92(br, 1H), 5.17(s, 2H), 7.36(d, 1H, $J=7.5\text{Hz}$), 7.47(t, 1H, $J=7.5\text{Hz}$), 7.56(t, 1H, $J=7.5\text{Hz}$), 7.91(d, 1H, $J=7.5\text{Hz}$), 7.97(s, 1H), 8.09(s, 1H), 8.54(br, 1H)

30 Example 26

6-Methyl-9-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-4,10-dihydrothieno[3,2-c][1]-benzoxepin-10-one (Compound 26) (yield: 90%)



$^1\text{H-NMR}$ (CDCl_3) δ : 1.75(s, 3H), 2.32(s, 3H), 4.34(br, 1H), 5.17(s, 2H), 6.99(d, 1H, $J=5.1\text{Hz}$), 7.42(d, 1H, $J=8.4\text{Hz}$), 7.70(d, 1H, $J=5.1\text{Hz}$), 8.25(d, 1H, $J=8.4\text{Hz}$), 11.92(br, 1H)

5

Example 27

6-Methoxy-9-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-4,10-dihydrothieno[3,2-c][1]-benzoxepin-10-one (Compound 27) (yield: 95%)

10 $^1\text{H-NMR}$ (CDCl_3) δ : 1.75(s, 3H), 3.91(s, 3H), 5.19(s, 2H), 6.97(d, 1H, $J=5.0\text{Hz}$), 7.15(d, 1H, $J=9.2\text{Hz}$), 7.69(d, 1H, $J=5.0\text{Hz}$), 8.27(d, 1H, $J=9.2\text{Hz}$), 11.62(br, 1H)

15 Example 28

8-Methyl-5-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-4,10-dihydrofuro[2,3-c][1]benzoxepin-4-one (Compound 28) (yield: 87%)

20 $^1\text{H-NMR}$ (CDCl_3) δ : 1.75(s, 3H), 2.33(s, 3H), 4.34(br, 1H), 5.13(s, 2H), 6.94(d, 1H, $J=2.0\text{Hz}$), 7.39(d, 1H, $J=2.0\text{Hz}$), 7.41(d, 1H, $J=8.6\text{Hz}$), 8.27(d, 1H, $J=8.6\text{Hz}$), 11.81(br, 1H)

Example 29

25 8-Methoxy-5-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-4,10-dihydrofuro[2,3-c][1]benzoxepin-4-one (Compound 29) (yield: 89%)

30 $^1\text{H-NMR}$ (CDCl_3) δ : 1.74(s, 3H), 3.93(s, 3H), 4.37(br, 1H), 5.17(s, 2H), 6.93(d, 1H, $J=2.0\text{Hz}$), 7.15(d, 1H, $J=9.2\text{Hz}$), 7.39(d, 1H, $J=2.0\text{Hz}$), 8.30(d, 1H, $J=9.2\text{Hz}$), 11.49(br, 1H)



Example 30

7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
5,11-dihydropyrido[2,3-c][1]benzoxepin-5-one (Compound
30) (yield: 47%)

5 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.65(s, 3H), 5.44(s, 2H), 7.22(d,
1H, $J=8.9\text{Hz}$), 7.50(br, 1H), 7.68(dd, 1H, $J=7.9$,
5.0Hz), 7.97(dd, 1H, $J=8.9$, 3.0Hz), 8.32(dd, 1H,
 $J=7.9$, 2.0Hz), 8.63(d, 1H, $J=3.0\text{Hz}$), 8.85(dd, 1H,
 $J=5.0$, 2.0Hz), 10.28(br, 1H)

10

Example 31

7-Methoxy-6-(3,3,3-trifluoro-2-hydroxy-2-methyl-
propanoylamino)-5,11-dihydropyrido[2,3-c][1]-
benzoxepin-5-one (Compound 31) (yield: 87%)

15 $^1\text{H-NMR}$ (CDCl_3) δ : 1.70(s, 3H), 3.90(s, 3H), 4.05(br,
1H), 5.41(s, 2H), 7.04(d, 1H, $J=8.9\text{Hz}$), 7.12(d,
1H, $J=8.9\text{Hz}$), 7.44(dd, 1H, $J=7.9$, 4.8Hz), 8.31
(dd, 1H, $J=7.9$, 1.6Hz), 8.69(dd, 1H, $J=4.8$,
1.6Hz), 8.75(br, 1H)

20

Example 32

7-Methyl-6-(3,3,3-trifluoro-2-hydroxy-2-methyl-
propanoylamino)-5,11-dihydropyrido[2,3-c][1]-
benzoxepin-5-one (Compound 32) (yield: 79%)

25 $^1\text{H-NMR}$ (CDCl_3) δ : 1.72(s, 3H), 2.23(s, 3H), 4.18(br,
1H), 5.38(s, 2H), 7.12(d, 1H, $J=8.2\text{Hz}$), 7.36-
7.44(m, 2H), 8.33(dd, 1H, $J=8.3$, 1.7Hz), 8.70(dd,
1H, $J=4.6$, 1.7Hz), 9.30(br, 1H)

30 Example 33

(S)-9-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl-
amino)-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one
(Compound 33)



Substantially the same procedure as in Example 1 was repeated using 9-amino-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one obtained in Reference Example 11 (0.840 g, 3.63 mmol) and (S)-3,3,3-trifluoro-2-hydroxy-2-methylpropionic acid obtained in Reference Example 27 (1.14 g, 7.26 mmol) to give Compound 33 (0.47 g, 35%).

¹H-NMR (CDCl₃) δ: 1.77(s, 3H), 4.28(br, 1H), 5.19(s, 2H), 7.00-7.04(m, 2H), 7.54(t, 1H, J=8.4Hz), 7.72(d, 1H, J=5.0Hz), 8.41(d, 1H, J=8.4Hz), 12.24(br, 1H)

Example 34

(R)-9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one (Compound 34) (yield: 46%)

Substantially the same procedure as in Example 1 was repeated using 9-amino-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one obtained in Reference Example 11 (0.951 g, 4.11 mmol) and (R)-3,3,3-trifluoro-2-hydroxy-2-methylpropionic acid obtained in Reference Example 28 (1.17 g, 7.40 mmol) to give Compound 34 (0.70 g, 46%).

¹H-NMR (CDCl₃) δ: 1.77(s, 3H), 4.28(br, 1H), 5.19(s, 2H), 7.00-7.04(m, 2H), 7.54(t, 1H, J=8.4Hz), 7.72(d, 1H, J=5.0Hz), 8.41(d, 1H, J=8.4Hz), 12.24(br, 1H)

Example 35

(S)-5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydrofuro[2,3-c][1]benzoxepin-4-one (Compound 35) (yield: 85%)

¹H-NMR (CDCl₃) δ: 1.56(s, 3H), 4.27(br, 1H), 5.17(s, 2H), 6.97(d, 1H, J=2.0Hz), 7.03(d, 1H, J=8.3Hz), 7.39(d, 1H, J=2.0Hz), 7.51(t, 1H, J=8.3Hz), 8.42(d, 1H, J=8.3Hz), 12.13(br, 1H)

Example 36

(R)-5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydrofuro[2,3-c][1]benzoxepin-4-one
(Compound 36) (yield: 74%)

5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.57(s, 3H), 4.29(br, 1H), 5.17(s, 2H), 6.96(d, 1H, $J=2.0\text{Hz}$), 7.04(d, 1H, $J=8.3\text{Hz}$), 7.39(d, 1H, $J=2.0\text{Hz}$), 7.52(t, 1H, $J=8.3\text{Hz}$), 8.42(d, 1H, $J=8.3\text{Hz}$), 12.12(br, 1H)

10 Example 37

(S)-7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydrofuro[2,3-c][1]benzoxepin-4-one
(Compound 37) (yield: 87%)

15 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.60(s, 3H), 5.31(s, 2H), 6.91(d, 1H, $J=2.0\text{Hz}$), 7.50(br, 1H), 7.70-7.77(m, 3H), 7.95(d, 1H, $J=8.9\text{Hz}$), 10.28(br, 1H)

Example 38

20 (R)-7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydrofuro[2,3-c][1]benzoxepin-4-one
(Compound 38) (yield: 56%)

25 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.60(s, 3H), 5.32(s, 1H), 6.92(d, 1H, $J=2.0\text{Hz}$), 7.58(br, 1H), 7.69-7.76(m, 3H), 7.95(d, 1H, $J=8.6\text{Hz}$), 10.30(br, 1H)

Example 39

(S)-8-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one
(Compound 39) (yield: 82%)

30 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.59(s, 3H), 5.26(s, 2H), 7.17'(d, 1H, $J=8.6\text{Hz}$), 7.24(d, 1H, $J=5.0\text{Hz}$), 7.41(br, 1H), 7.92(d, 1H, $J=8.6\text{Hz}$), 8.04(d, 1H, $J=5.0\text{Hz}$), 8.42(s, 1H), 10.19(br, 1H)



Example 40

1-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
6,11-dihydrodibenzo[b,e]thiepin-11-one (Compound 40)
(yield: 62%)

5 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.56(s, 3H), 4.26(s, 2H), 7.30(d, 1H, $J=7.9\text{Hz}$), 7.36(t, 1H, $J=7.9\text{Hz}$), 7.40-7.59(m, 3H), 7.56(d, 1H, $J=7.9\text{Hz}$), 7.66(br, 1H), 7.94(d, 1H, $J=7.9\text{Hz}$), 8.31(d, 1H, $J=9.2\text{Hz}$), 10.73(br, 1H)

10 Example 41

3-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
6,11-dihydrodibenzo[b,e]thiepin-11-one (Compound 41)
(yield: 60%)

15 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.58(s, 3H), 4.22(s, 2H), 7.36-7.40 (m, 2H), 7.48-7.52(m, 2H), 7.72(d, 1H, $J=8.9\text{Hz}$), 7.94(s, 1H), 8.11(d, 1H, $J=8.9\text{Hz}$), 8.31(d, 1H, $J=9.2\text{ Hz}$), 10.73(br, 1H)

Example 42

20 5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrofuro[2,3-c][1]benzothiepin-4-one
(Compound 42) (yield: 56%)

25 $^1\text{H-NMR}$ (CDCl_3) δ : 1.72(s, 3H), 3.97(s, 2H), 4.17(br, 1H), 6.87(d, 1H, $J=2.0\text{Hz}$), 7.28(d, 1H, $J=2.0\text{Hz}$), 7.43(t, 1H, $J=7.9\text{Hz}$), 7.55(d, 1H, $J=7.9\text{Hz}$), 8.32 (d, 1H, $J=7.9\text{Hz}$), 10.71(br, 1H)

Example 43

30 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrofuro[2,3-c][1]benzothiepin-4-one
(Compound 43) (yield: 70%)

$^1\text{H-NMR}$ (DMSO-d_6) δ : 1.60(s, 3H), 3.61(br, 1H), 4.24(s, 2H), 6.88(d, 1H, $J=2.0\text{Hz}$), 7.65(d, 1H, $J=2.0\text{Hz}$),

7.87(d, 1H, J=8.1Hz), 7.94(d, 1H, J=8.1Hz), 8.19
(s, 1H), 10.30(br, 1H)

Example 44

5 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrothieno[3,2-c][1]benzothiepin-10-one
(Compound 44) (yield: 62%)
¹H-NMR (CDCl₃) δ: 1.72(s, 3H), 3.98(s, 2H), 4.22(br,
1H), 6.97(d, 1H, J=5.0Hz), 7.43(t, 1H, J=8.0Hz),
10 7.51(d, 1H, J=8.0Hz), 7.62(d, 1H, J=5.0Hz), 8.27
(d, 1H, J=8.0Hz), 10.62(br, 1H)

Example 45

15 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrothieno[3,2-c][1]benzothiepin-10-one
(Compound 45) (yield: 99%)
¹H-NMR (DMSO-d₆) δ: 1.60(s, 3H), 4.19(s, 2H), 7.17(d,
1H, J=5.0Hz), 7.55(br, 1H), 7.87(d, 1H, J=
8.9Hz), 7.95(d, 1H, J=5.0Hz), 7.98(d, 1H, J=
20 8.9Hz), 8.17(s, 1H), 10.71(br, 1H)

Example 46

25 5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrothieno[2,3-c][1]benzothiepin-4-one
(Compound 46) (yield: 53%)
¹H-NMR (CDCl₃) δ: 1.81(s, 3H), 4.46(br, 1H), 5.21(s,
2H), 7.34(d, 2H, J=7.4Hz), 7.39(d, 1H, J=8.6Hz),
7.45(t, 1H, J=7.4Hz), 7.58(t, 1H, J=7.4Hz), 7.87
(d, 1H, J=7.4Hz), 8.26(d, 1H, J=8.6Hz), 12.57(br,
30 1H)

Example 47

7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,10-dihydrothieno[2,3-c][1]benzothiepin-4-one



(Compound 47) (yield: 28%)

¹H-NMR (DMSO-d₆) δ: 1.59(s, 3H), 4.39(s, 2H), 7.35(d, 1H, J=5.3Hz), 7.51(d, 1H, J=5.3Hz), 7.56(s, 1H), 7.86(dd, 1H, J=8.6, 2.0Hz), 8.00(d, 1H, J=8.6Hz), 8.14(d, 1H, J=2.0Hz), 10.37(br, 1H)

Example 48

5-Oxo-1-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl-amino)-6,11-dihydrodibenzo[b,e]thiepin-11-one

(Compound 48) (yield: 55%)

¹H-NMR (DMSO-d₆) δ: 1.53(s, 3H), 4.56(d, 1H, J=15.5Hz), 4.92(d, 1H, J=15.5Hz), 7.43(d, 1H, J=7.6Hz), 7.52(t, 1H, J=7.6Hz), 7.63-7.76(m, 3H), 7.89-8.02(m, 2H), 10.38(br, 1H)

Example 49

5-Oxo-3-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl-amino)-6,11-dihydrodibenzo[b,e]thiepin-11-one

(Compound 49) (yield: 60%)

¹H-NMR (DMSO-d₆) δ: 1.62(s, 3H), 4.57(d, 1H, J=13.5Hz), 4.98(d, 1H, J=13.5Hz), 7.49-7.68(m, 4H), 8.00(d, 1H, J=8.3Hz), 8.16(d, 1H, J=8.3Hz), 8.49(s, 1H), 10.61(br, 1H)

Example 50

9-Oxo-5-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydrothieno[2,3-c][1]benzothiepin-4-one
(Compound 50) (yield: 64%)

¹H-NMR (DMSO-d₆) δ: 1.55(s, 3H), 4.75(d, 1H, J=15.8Hz), 4.93(d, 1H, J=15.8Hz), 7.48(d, 1H, J=5.3Hz), 7.60(d, 1H, J=5.3Hz), 7.77(m, 3H), 8.19(d, 1H, J=8.6Hz), 10.80(br, 1H)



Example 51

9-Oxo-7-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydrothieno[2,3-c][1]benzothiepin-4-one
(Compound 51) (yield: 62%)

5 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.61(s, 3H), 4.83(d, 1H, $J=14.8\text{Hz}$),
4.96(d, 1H, $J=14.8\text{Hz}$), 7.61(m, 3H), 7.97(d, 1H,
 $J=8.6\text{Hz}$), 8.07(dd, 1H, $J=8.6, 2.0\text{Hz}$), 8.50(d, 1H,
 $J=2.0\text{Hz}$), 10.67(br, 1H)

10 Example 52

4-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-one
(Compound 52)

In dimethylacetamide (15 ml) was dissolved 3,3,3-
15 trifluoro-2-hydroxy-2-methylpropionic acid (0.370 g, 2.37
mmol), and thionyl chloride (0.206 ml, 2.83 mmol) was added
thereto at -5°C , followed by stirring at -5°C for 2 hours.
To this reaction mixture were added 4-amino-10,11-dihydro-
5H-dibenzo[a,d]cyclohepten-5-one obtained substantially in
20 the same manner as in Reference Examples 12 to 18 (0.210 g,
0.942 mmol) and triethylamine (0.394 ml, 2.83 mmol),
followed by stirring at room temperature for 4 hours. After
concentration of the reaction mixture under reduced
pressure, the obtained residue was dissolved in ethyl
25 acetate (20 ml). The organic layer was washed successively
with a saturated aqueous solution of sodium bicarbonate and
a saturated aqueous solution of sodium chloride, and then
dried over anhydrous magnesium sulfate. After the drying
agent was filtered off, the organic layer was concentrated
30 under reduced pressure. The obtained oily residue was
purified by silica gel column chromatography (hexane/ethyl
acetate = 5/1), followed by trituration with hexane to give
Compound 52 (0.210 g, 61%).

$^1\text{H-NMR}$ (CDCl_3) δ : 1.73(s, 3H), 3.10-3.35(br, 4H), 4.26
35 (br, 1H), 7.08(dd, 1H, $J=7.6, 0.8\text{Hz}$), 7.21-7.32



(m, 1H), 7.32-7.41(m, 1H), 7.41-7.52(m, 2H), 8.03
(dd, 1H, J=7.9, 1.7Hz), 8.16(dd, 1H, J=8.2,
0.8Hz), 10.28(br, 1H)

5 In the following Examples 53 to 68, substantially the
same procedure as in Example 52 was repeated except that the
corresponding tricyclic aromatic amine was used instead of
4-amino-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-one, to
give the desired compound.

10

Example 53

3-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-one
(Compound 53) (yield: 83%)

15 ¹H-NMR (CDCl₃) δ: 1.74(s, 3H), 3.20(br, 4H), 4.01(br,
1H), 7.21-7.28(m, 2H), 7.30-7.38(m, 1H), 7.42-
7.50(m, 1H), 7.89(d, 1H, J=2.3Hz), 7.95-8.03(m,
2H), 8.43(br, 1H)

20 Example 54

2-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-one
(Compound 54) (yield: 54%)

25 ¹H-NMR (CDCl₃) δ: 1.76(s, 3H), 3.20(br, 4H), 3.77(br,
1H), 7.23(dd, 1H, J=7.3, 1.0Hz), 7.31-7.50(m,
3H), 7.70(d, 1H, J=2.3Hz), 8.02(dd, 1H, J=7.9,
1.7Hz), 8.10(d, 1H, J=8.6Hz), 8.51(br, 1H)

Example 55

30 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,5-dihydro-10H-benzo[5,6]cyclohepta[1,2-b]
thiophen-10-one (Compound 55) (yield: 80%)

¹H-NMR (CDCl₃) δ: 1.75(s, 3H), 3.04-3.22(m, 4H), 4.39
(br, 1H), 6.96(d, 1H, J=5.2Hz), 7.11(dd, 1H, J=



8.0, 1.0Hz), 7.48(t, 1H, J=8.0Hz), 7.63(d, 1H, J=5.2Hz), 8.38(dd, 1H, J=8.0, 1.0Hz), 11.50(br, 1H)

Example 56

5 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4,5-dihydro-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-
10-one (Compound 56) (yield: 75%)

$^1\text{H-NMR}$ (CDCl_3) δ : 1.77(s, 3H), 3.06-3.21(m, 4H), 3.71

(br, 1H), 6.96(d, 1H, J=5.0Hz), 7.37(dd, 1H, J=

10 8.6, 2.2Hz), 7.58(d, 1H, J=5.0Hz), 7.77(d, 1H, J=2.2Hz), 8.04(d, 1H, J=8.6Hz), 8.52(br, 1H)

Example 57

15 5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
9,10-dihydro-4H-benzo[4,5]cyclohepta[1,2-b]thiophen-
4-one (Compound 57) (yield: 76%)

$^1\text{H-NMR}$ (CDCl_3) δ : 1.74(s, 3H), 3.19-3.30(m, 4H), 4.34

(br, 1H), 7.07(d, 1H, J=5.3Hz), 7.09(dd, 1H, J=

20 8.0, 1.0Hz), 7.45(t, 1H, J=8.0Hz), 7.57(d, 1H, J=5.3Hz), 8.31(dd, 1H, J=8.0, 1.0Hz), 11.09(br, 1H)

Example 58

25 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
9,10-dihydro-4H-benzo[4,5]cyclohepta[1,2-b]furan-4-one
(Compound 58) (yield: 65%)

$^1\text{H-NMR}$ (CDCl_3) δ : 1.76(s, 3H), 3.09-3.21(m, 4H), 3.98

(br, 1H), 6.91(d, 1H, J=2.0Hz), 7.30(d, 1H, J=

2.0Hz), 7.37(dd, 1H, J=8.4, 2.1Hz), 7.76(d, 1H, J=2.1Hz), 7.95(d, 1H, J=8.4 Hz), 8.55(br, 1H)

30

Example 59

8-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
10,11-dihydro-5H-benzo[4,5]cyclohepta[1,2-b]pyridin-5-
one (Compound 59) (yield: 25%)



¹H-NMR (CDCl₃) δ: 1.76(s, 3H), 3.21-3.28(m, 2H), 3.42-3.49(m, 2H), 5.00(br, 1H), 7.33(dd, 1H, J=7.9, 4.9Hz), 7.45(dd, 1H, J=8.6, 2.0Hz), 7.72(d, 1H, J=2.0Hz), 8.05(d, 1H, J=8.6Hz), 8.42(dd, 1H, J=7.9, 1.7Hz), 8.63(dd, 1H, J=4.9, 1.7Hz), 8.74(br, 1H)

Example 60

4-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-5H-dibenzo[a,d]cyclohepten-5-one (Compound 60) (yield: 87%)

¹H-NMR (CDCl₃) δ: 1.82(s, 3H), 4.35(br, 1H), 7.09(s, 2H), 7.39(dd, 1H, J=7.9, 1.2Hz), 7.46-7.69(m, 4H), 7.95(dd, 1H, J=7.6, 1.2Hz), 8.45(dd, 1H, J=8.2, 1.2Hz), 11.07(br, 1H)

Example 61

3-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-5H-dibenzo[a,d]cyclohepten-5-one (Compound 61) (yield: 47%)

¹H-NMR (CDCl₃) δ: 1.77(s, 3H), 3.95(br, 1H), 7.06(s, 2H), 7.55-7.65(m, 3H), 7.66-7.73(m, 1H), 8.12(d, 1H, J=2.3Hz), 8.23-8.31(m, 2H), 8.66(br, 1H)

Example 62

2-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-5H-dibenzo[a,d]cyclohepten-5-one (Compound 62) (yield: 91%)

¹H-NMR (CDCl₃) δ: 1.79(s, 3H), 3.65(br, 1H), 7.02(d, 1H, J=12.2Hz), 7.09(d, 1H, J=12.2Hz), 7.54-7.62(m, 1H), 7.64-7.71(m, 1H), 8.01(d, 1H, J=2.0Hz), 8.24-8.32(m, 2H), 8.59(br, 1H)



Example 63

9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one
(Compound 63) (yield: 88%)

5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.85(s, 3H), 4.50(br, 1H), 7.20-7.38
(m, 3H), 7.57(dd, 1H, $J=8.2$, 1.0Hz), 7.77(t, 1H,
 $J=8.2\text{Hz}$), 7.83(d, 1H, $J=5.3\text{Hz}$), 8.97(dd, 1H, $J=$
8.2, 1.0Hz), 13.84(br, 1H)

10 Example 64

8-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one
(Compound 64) (yield: 95%)

15 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.69(s, 3H), 7.49(d, 1H, $J=11.9\text{Hz}$),
7.55(d, 1H, $J=11.9\text{Hz}$), 7.66(d, 1H, $J=5.3\text{Hz}$), 8.03
(d, 1H, $J=8.6\text{Hz}$), 8.26(d, 1H, $J=5.3\text{Hz}$), 8.31(dd,
1H, $J=8.6$, 2.1Hz), 9.17(d, 1H, $J=2.1\text{Hz}$), 10.59
(br, 1H)

20 Example 65

7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one
(Compound 65) (yield: 84%)

25 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.69(s, 3H), 7.43(d, 1H, $J=11.6\text{Hz}$),
7.55(d, 1H, $J=11.6\text{Hz}$), 7.67(d, 1H, $J=5.3\text{Hz}$), 7.71
(br, 1H), 8.11-8.27(m, 2H), 8.47(d, 1H, $J=2.0\text{Hz}$),
8.67(d, 1H, $J=9.3\text{Hz}$), 10.58(br, 1H)

Example 66

30 5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
4H-benzo[4,5]cyclohepta[1,2-b]thiophen-4-one (Compound
66) (yield: 90%)

$^1\text{H-NMR}$ (CDCl_3) δ : 1.83(s, 3H), 4.48(br, 1H), 7.18(d,
1H, $J=11.9\text{Hz}$), 7.26(d, 1H, $J=11.9\text{Hz}$), 7.44(d, 1H,



$J=5.4\text{Hz}$), 7.51(dd, 1H, $J=7.9$, 1.3Hz), 7.71(t, 1H, $J=7.9\text{Hz}$), 7.83(d, 1H, $J=5.4\text{Hz}$), 8.75(dd, 1H, $J=7.9$, 1.3Hz), 12.83(br, 1H)

5 Example 67

9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10H-benzo[5,6]cyclohepta[1,2-b]furan-10-one (Compound 67) (yield: 81%)

$^1\text{H-NMR}$ (CDCl_3) δ : 1.87(s, 3H), 4.56(br, 1H), 6.84(d, 1H, $J=1.8\text{Hz}$), 7.17(d, 1H, $J=11.5\text{Hz}$), 7.43(d, 1H, $J=11.5\text{Hz}$), 7.61(dd, 1H, $J=8.1$, 1.3Hz), 7.80(t, 1H, $J=8.1\text{Hz}$), 7.91(d, 1H, $J=1.8\text{Hz}$), 9.05(dd, 1H, $J=8.1$, 1.3Hz), 14.23(br, 1H)

15 Example 68

7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10H-benzo[5,6]cyclohepta[1,2-b]furan-10-one (Compound 68) (yield: 85%)

$^1\text{H-NMR}$ (DMSO-d_6) δ : 1.63(s, 3H), 7.12(d, 1H, $J=1.8\text{Hz}$), 7.32(d, 1H, $J=11.8\text{Hz}$), 7.41(d, 1H, $J=11.8\text{Hz}$), 7.63(br, 1H), 8.17(dd, 1H, $J=8.9$, 2.0Hz), 8.25(d, 1H, $J=1.8\text{Hz}$), 8.41(d, 1H, $J=2.0\text{Hz}$), 8.71(d, 1H, $J=8.9\text{Hz}$), 10.47(br, 1H)

25 Example 69

8-Chloro-7-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydrothieno[3,2-c][1]-benzoxepin-10-one (Compound 69) (yield: 95%)

$^1\text{H-NMR}$ (DMSO-d_6) δ : 1.63(s, 3H), 5.35(s, 2H), 7.25(d, 1H, $J=5.0\text{Hz}$), 8.07(d, 1H, $J=5.0\text{Hz}$), 8.11(s, 1H), 8.13(s, 1H), 8.15(s, 1H), 9.85(br, 1H)



Example 70

7-Methyl-8-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydrothieno[3,2-c][1]-benzoxepin-10-one (Compound 70) (yield: 51%)

- 5 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.59(s, 3H), 2.22(s, 3H), 5.29(s, 2H), 7.12(s, 1H), 7.25(d, 1H, $J=5.0\text{Hz}$), 7.42(br, 1H), 7.94(s, 1H), 8.06(d, 1H, $J=5.0\text{Hz}$), 9.75(br, 1H)

10 Example 71

7-Methyl-9-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoylamino)-5,11-dihydropyrido[2,3-c][1]-benzoxepin-5-one (Compound 71) (yield: 64%)

- 15 $^1\text{H-NMR}$ (CDCl_3) δ : 1.77(s, 3H), 2.40(s, 3H), 5.48(s, 2H), 5.59(br, 1H), 7.53(dd, 1H, $J=7.9, 4.9\text{Hz}$), 7.81(d, 1H, $J=2.3\text{Hz}$), 8.37(dd, 1H, $J=7.9, 1.7\text{Hz}$), 8.57(d, 1H, $J=2.3\text{Hz}$), 8.72(dd, 1H, $J=4.9, 1.7\text{Hz}$), 9.30(br, 1H)

20 Example 72

9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,5-dihydro-10H-benzo[5,6]cyclohepta[1,2-b]furan-10-one (Compound 72) (yield: 63%)

- 25 $^1\text{H-NMR}$ (CDCl_3) δ : 1.77(s, 3H), 2.87-2.96(m, 2H), 3.13-3.22(m, 2H), 4.34(br, 1H), 6.46(d, 1H, $J=1.7\text{Hz}$), 7.10(dd, 1H, $J=8.0, 1.0\text{Hz}$), 7.48(t, 1H, $J=8.0\text{Hz}$), 7.64(d, 1H, $J=1.7\text{Hz}$), 8.48(dd, 1H, $J=8.0, 1.0\text{Hz}$), 11.94(br, 1H)

30 Example 73

7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,5-dihydro-10H-benzo[5,6]cyclohepta[1,2-b]furan-10-one (Compound 73) (yield: 88%)



¹H-NMR (CDCl₃) δ: 1.77(s, 3H), 2.89-2.98(m, 2H), 3.12-3.20(m, 2H), 4.04(br, 1H), 6.44(d, 1H, J=1.7Hz), 7.37(dd, 1H, J=8.6, 2.0Hz), 7.61(d, 1H, J=1.7Hz), 7.79(d, 1H, J=2.0Hz), 8.10(d, 1H, J=8.6Hz), 8.60
5 (br, 1H)

Example 74

7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
5H-benzo[4,5]cyclohepta[1,2-b]pyridin-5-one (Compound
10 74) (yield: 70%)

¹H-NMR (CDCl₃) δ: 1.77(s, 3H), 7.27(d, 1H, J=12.5Hz),
7.36(d, 1H, J=12.5Hz), 7.49(dd, 1H, J=7.9,
4.6Hz), 7.65(d, 1H, J=8.3Hz), 8.22(d, 1H, J=
2.4Hz), 8.27(dd, 1H, J=8.3, 2.4Hz), 8.56(dd, 1H,
15 J=7.9, 1.6Hz), 8.69(br, 1H), 8.91(dd, 1H, J=4.6,
1.6Hz)

Example 75

9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
20 5H-benzo[4,5]cyclohepta[1,2-b]pyridin-5-one (Compound
75) (yield: 65%)

¹H-NMR (CDCl₃) δ: 1.88(s, 3H), 7.40(d, 1H, J=12.7Hz),
7.49(d, 1H, J=12.7Hz), 7.57(dd, 1H, J=8.3,
5.0Hz), 7.65(t, 1H, J=7.9Hz), 7.98(dd, 1H, J=7.9,
25 1.3Hz), 8.24(dd, 1H, J=7.9, 1.3Hz), 8.42(dd, 1H,
J=8.3, 1.7Hz), 8.84(dd, 1H, J=5.0, 1.7Hz), 9.07
(br, 1H)

Example 76

7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-
10,11-dihydro-5H-benzo[4,5]cyclohepta[1,2-b]pyridin-
5-one (Compound 76) (yield: 40%)

¹H-NMR (DMSO-d₆) δ: 1.58(s, 3H), 3.13-3.27(m, 2H),
3.30-3.44(m, 2H), 7.35(d, 1H, J=8.3Hz), 7.45(br,



1H), 7.46(dd, 1H, J=7.9, 4.6Hz), 7.85(dd, 1H, J=8.3, 2.4Hz), 8.28(d, 1H, J=2.4Hz), 8.29(dd, 1H, J=7.9, 1.6Hz), 8.67(dd, 1H, J=4.6, 1.6Hz), 10.16(br, 1H)

5

Example 77

9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10,11-dihydro-5H-benzo[4,5]cyclohepta[1,2-b]pyridin-5-one (Compound 77) (yield: 45%)

10 ¹H-NMR (CDCl₃) δ: 1.80(s, 3H), 3.15-3.26(m, 2H), 3.44-3.54(m, 2H), 7.37-7.45(m, 2H), 7.78(dd, 1H, J=7.9, 1.3Hz), 8.04(dd, 1H, J=7.9, 1.3Hz), 8.29(dd, 1H, J=7.9, 1.7Hz), 8.60(dd, 1H, J=5.0, 1.7Hz), 8.84(br, 1H)

15

Reference Example 1

Methyl 2-(3-acetamidophenoxymethyl)benzoate

To a solution of 3-acetamidophenol (1.0 g, 6.6 mmol) in dimethylformamide (10 ml) was added cesium carbonate (1.3 g, 3.9 mmol), followed by stirring at room temperature for 30 minutes. To the resulting mixture was added methyl 2-bromomethylbenzoate (1.8 g, 7.9 mmol), and the mixture was stirred overnight at room temperature. After the reaction was completed, the reaction mixture was poured into a 5% aqueous solution of sodium bicarbonate, followed by extraction with ethyl acetate. The organic layer was washed with a saturated aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure. The obtained oily substance was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1), followed by trituration with hexane to give methyl 2-(3-acetamidophenoxymethyl)benzoate as a white solid (1.21 g, 61%).



¹H-NMR (CDCl₃) δ: 2.16(s, 3H), 3.90(s, 3H), 5.47(s, 2H), 6.73(d, 1H, J=7.8Hz), 7.09(d, 1H, J=7.8Hz), 7.07-7.26(m, 1H), 7.34(s, 1H), 7.38(d, 1H, J=7.8Hz), 7.55(t, 1H, J=7.8Hz), 7.73(d, 1H, J=7.8Hz), 8.15(d, 1H, J=7.8Hz)

Reference Example 2

2-(3-Acetamidophenoxymethyl)benzoic acid

The methyl ester obtained in Reference Example 1 (1.0 g, 3.4 mmol) was dissolved in methanol (25 ml), and a 2 N aqueous solution of sodium hydroxide (25 ml) was added thereto, followed by stirring at room temperature for 2 hours. After the reaction was completed, methanol was distilled off under reduced pressure, and the resulting mixture was adjusted to pH 3 with concentrated hydrochloric acid. The precipitated white solid was washed well with water and dried to give 2-(3-acetamidophenoxymethyl)benzoic acid (0.87 g, 90%).

¹H-NMR (DMSO-d₆) δ: 2.02(s, 3H), 5.42(s, 2H), 6.64(d, 1H, J=7.8Hz), 7.13-7.22(m, 2H), 7.31(s, 1H), 7.43(t, 1H, J=7.8Hz), 7.56-7.65(m, 2H), 7.94(d, 1H, J=7.8Hz), 9.92(br, 1H), 13.02(br, 1H)

Reference Example 3

1-Acetamido-6,11-dihydrodibenz[b,e]oxepin-11-one [1]

3-Acetamido-6,11-dihydrodibenz[b,e]oxepin-11-one [2]

The carboxylic acid obtained in Reference Example 2 (1.6 g, 5.6 mmol) was suspended in dichloromethane (20 ml), and trifluoroacetic anhydride (0.95 ml, 6.7 mmol) was added dropwise thereto under ice cooling, followed by stirring at room temperature for 2 hours. After the reaction mixture was ice cooled again, boron trifluoride diethyl etherate (0.5 equivalent) was added thereto, and the mixture was stirred overnight at room temperature. After the reaction was completed, the dichloromethane layer was washed



successively with water, a 5% aqueous solution of sodium bicarbonate, and a saturated aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer
5 was concentrated under reduced pressure. The obtained oily mixture was separated and purified by silica gel column chromatography (chloroform/methanol = 100/1) to give 1-acetamido-6,11-dihydrodibenz[b,e]oxepin-11-one [1] (0.31 g, 21%) and 3-acetamido-6,11-dihydrodibenz[b,e]oxepin-11-one
10 [2] (0.28 g, 19%), respectively.

[1] $^1\text{H-NMR}$ (CDCl_3) δ : 2.28(s, 3H), 5.15(s, 2H), 6.79(d, 1H, $J=7.8\text{Hz}$), 7.32(d, 1H, $J=7.8\text{Hz}$), 7.45-7.58(m, 3H), 7.84(d, 1H, $J=7.8\text{Hz}$), 8.41(d, 1H, $J=7.8\text{Hz}$)

[2] $^1\text{H-NMR}$ (CDCl_3) δ : 2.20(s, 3H), 5.19(s, 2H), 7.10(d, 1H, $J=8.0\text{Hz}$), 7.36(d, 1H, $J=8.0\text{Hz}$), 7.49-7.58(m, 3H), 7.91(d, 1H, $J=8.0\text{Hz}$), 8.21(d, 1H, $J=8.0\text{Hz}$)
15

Reference Example 4

1-Amino-6,11-dihydrodibenz[b,e]oxepin-11-one

20 In concentrated hydrochloric acid (10 ml), 1-acetamido-6,11-dihydrodibenz[b,e]oxepin-11-one obtained in Reference Example 3 (0.25 g, 0.94 mmol) was heated under reflux for 2 hours. The reaction mixture was poured into a saturated aqueous solution of sodium bicarbonate, followed
25 by extraction with ethyl acetate (25 ml). The organic layer was washed with a saturated aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure to give 1-amino-
30 6,11-dihydrodibenz[b,e]oxepin-11-one as a yellow solid (0.17 g, 81%).

$^1\text{H-NMR}$ (CDCl_3) δ : 5.12(s, 2H), 6.11(d, 1H, $J=8.1\text{Hz}$), 6.47(d, 1H, $J=8.1\text{Hz}$), 7.12(t, 1H, $J=8.1\text{Hz}$), 7.42-7.55(m, 3H), 7.73(d, 1H, $J=8.1\text{Hz}$)



Reference Example 5

3-Amino-6,11-dihydrodibenz[b,e]oxepin-11-one

Substantially the same procedure as in Reference Example 4 was repeated using 3-acetamido-6,11-dihydrodibenz[b,e]oxepin-11-one obtained in Reference Example 3 instead of 1-acetamido-6,11-dihydrodibenz[b,e]oxepin-11-one to give 3-amino-6,11-dihydrodibenz[b,e]oxepin-11-one as a white solid (yield: 95%).

¹H-NMR (CDCl₃) δ: 3.85(br, 2H), 5.05(s, 2H), 6.11(s, 1H), 6.31(d, 1H, J=7.9Hz), 7.23(d, 1H, J=7.9Hz), 7.33-7.45(m, 2H), 7.85(d, 1H, J=7.9Hz), 8.04(d, 1H, J=7.9Hz)

Reference Example 6

2-Amino-6,11-dihydrodibenz[b,e]oxepin-11-one

To tert-butanol (8.3 ml) were added diphenylphosphoryl azide (1.1 ml, 5.11 mmol), 2-carboxy-6,11-dihydrodibenz[b,e]oxepin-11-one (Japanese Published Unexamined Patent Application No. 91040/90) (1.0 g, 3.9 mmol), and triethylamine (0.71 ml, 5.11 mmol) under an atmosphere of argon gas, followed by heating under reflux for 2 hours. After the reaction mixture was cooled, it was poured into a 5% aqueous solution of sodium bicarbonate, followed by extraction with ethyl acetate. The organic layer was washed with a saturated aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure. The obtained oily substance was purified by silica gel column chromatography (hexane/ethyl acetate = 8/1), followed by trituration with hexane to give 2-tert-butoxycarbonylamino-6,11-dihydrodibenz[b,e]oxepin-11-one as a white solid (0.51 g, 41%).

¹H-NMR (CDCl₃) δ: 1.49(s, 9H), 5.23(s, 2H), 7.21-7.26(m, 2H), 7.48-7.54(m, 2H), 7.60-7.65(m, 1H), 7.80

(d, 1H, J=8.2Hz), 8.02(d, 1H, J=8.2Hz), 9.80(br, 1H)

To the obtained white solid (0.41 g) was added 4 N
5 hydrochloric acid/dioxane (10 ml), followed by stirring at
room temperature for one hour. The reaction mixture was
concentrated under reduced pressure to give 2-amino-6,11-
dihydrodibenz[b,e]oxepin-11-one as a white solid. This
product was used for the synthesis of Compound 3 without
10 purification.

Reference Example 7

Methyl 3-(3-acetamidophenoxymethyl)-2-thiophene-
carboxylate

15 To a solution of 3-acetamidophenol (4.79 g, 31.7 mmol)
in dimethylformamide (50 ml) was added cesium carbonate
(5.67 g, 17.4 mmol), followed by stirring at room
temperature for 30 minutes. To the resulting mixture was
added methyl 3-bromomethyl-2-thiophenecarboxylate (8.94 g,
20 38.0 mmol), and the mixture was stirred overnight at room
temperature. After the reaction was completed, the reaction
mixture was poured into a 5% aqueous solution of sodium
bicarbonate, followed by extraction with ethyl acetate. The
organic layer was washed with a saturated aqueous solution
25 of sodium chloride, and then dried over anhydrous magnesium
sulfate. After the drying agent was filtered off, the
organic layer was concentrated under reduced pressure. The
obtained oily substance was purified by silica gel column
chromatography (hexane/ethyl acetate = 5/1), followed by
30 trituration with hexane to give methyl 3-(3-acetamido-
phenoxymethyl)-2-thiophenecarboxylate as a white solid (7.27
g, 75%).

¹H-NMR (CDCl₃) δ: 2.17(s, 3H), 3.94(s, 3H), 5.45(s,
2H), 6.72(d, 1H, J=8.1Hz), 7.07(d, 1H, J=8.1Hz),
35 7.18-7.24(m, 2H), 7.29(d, 1H, J=5.1Hz), 7.48(d,
1H, J=5.1Hz)



Reference Example 8

3-(3-Acetamidophenoxymethyl)-2-thiophenecarboxylic acid

The methyl ester obtained in Reference Example 7 (6.27 g, 20.5 mmol) was dissolved in methanol (100 ml), and a 2 N aqueous solution of sodium hydroxide (50 ml) was added thereto, followed by stirring at room temperature for 2 hours. After the reaction was completed, methanol was distilled off under reduced pressure, and the resulting mixture was adjusted to pH 3 with concentrated hydrochloric acid. The precipitated white solid was washed well with water and dried to give 3-(3-acetamidophenoxymethyl)-2-thiophenecarboxylic acid (4.33 g, 73%).

$^1\text{H-NMR}$ (DMSO-d_6) δ : 2.02(s, 3H), 5.38(s, 2H), 6.62(d, 1H, $J=8.1\text{Hz}$), 7.13-7.20(m, 2H), 7.23(d, 1H, $J=5.1\text{Hz}$), 7.31(s, 1H), 7.78(d, 1H, $J=5.1\text{Hz}$)

Reference Example 9

7-Acetamido-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one [1]

9-Acetamido-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one [2]

The carboxylic acid obtained in Reference Example 8 (2.20 g, 7.55 mmol) was suspended in dichloromethane (20 ml), and trifluoroacetic anhydride (1.28 ml, 9.66 mmol) was added dropwise thereto under ice cooling, followed by stirring at room temperature for 2 hours. After the reaction mixture was ice cooled again, boron trifluoride diethyl etherate (0.5 equivalent) was added thereto, and the mixture was stirred overnight at room temperature. After the reaction was completed, the dichloromethane layer was washed successively with water, a 5% aqueous solution of sodium bicarbonate, and a saturated aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure. The



obtained oily mixture was separated and purified by silica gel column chromatography (chloroform/methanol = 100/1) to give 7-acetamido-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one [1] (0.42 g, 21%) and 9-acetamido-4,10-dihydrothieno-
 5 [3,2-c][1]benzoxepin-10-one [2] (0.28 g, 14%), respectively.

[1] $^1\text{H-NMR}$ (CDCl_3) δ : 2.10(s, 3H), 5.27(s, 2H), 7.22(d, 1H, $J=5.3\text{Hz}$), 7.36(d, 1H, $J=8.6\text{Hz}$), 7.58(s, 1H), 7.97-8.00(m, 2H), 10.33(br, 1H)

[2] $^1\text{H-NMR}$ (CDCl_3) δ : 2.23(s, 3H), 5.16(s, 2H), 6.92(d, 1H, $J=8.2\text{Hz}$), 6.98(d, 1H, $J=5.1\text{Hz}$), 7.47(t, 1H, $J=8.2\text{Hz}$), 7.69(d, 1H, $J=5.1\text{Hz}$), 8.40(d, 1H, $J=8.2\text{Hz}$), 11.28(br, 1H)

Reference Example 10

15 7-Amino-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one
 In concentrated hydrochloric acid (15 ml), 7-acetamido-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one obtained in Reference Example 9 (0.41 g, 1.51 mmol) was heated under reflux for 2 hours. The reaction mixture was
 20 poured into a saturated aqueous solution of sodium bicarbonate, followed by extraction with ethyl acetate (25 ml). The organic layer was washed with a saturated aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off,
 25 the organic layer was concentrated under reduced pressure to give 7-amino-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one as a yellow solid (0.28 g, 78%).

$^1\text{H-NMR}$ (CDCl_3) δ : 4.22(br, 2H), 5.14(s, 2H), 6.31(s, 1H), 6.48(d, 1H, $J=8.6\text{Hz}$), 6.99(d, 1H, $J=5.0\text{Hz}$),
 30 7.57(d, 1H, $J=5.0\text{Hz}$), 8.08(d, 1H, $J=8.6\text{Hz}$)



Reference Example 11

9-Amino-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one

Substantially the same procedure as in Reference Example 10 was repeated using 9-acetamido-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one obtained in Reference Example 9 (0.28 g, 1.02 mmol) instead of 7-acetamido-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one to give 9-amino-4,10-dihydrothieno[3,2-c][1]benzoxepin-10-one as a white solid (0.24 g, 99%).

10 $^1\text{H-NMR}$ (CDCl_3) δ : 5.11(s, 2H), 6.36(br, 2H), 6.41(d, 1H, $J=7.9\text{Hz}$), 6.47(d, 1H, $J=7.9\text{Hz}$), 6.95(d, 1H, $J=5.0\text{Hz}$), 7.17(t, 1H, $J=7.9\text{Hz}$), 7.58(d, 1H, $J=5.0\text{Hz}$)

15 Reference Example 12

2-Methoxycarbonylthiophen-3-ylmethyltriphenylphosphonium bromide

To a solution of methyl 3-bromomethyl-2-thiophenecarboxylate (7.31 g, 31.1 mmol) in toluene (90 ml) was added triphenylphosphine (7.92 g, 30.2 mmol) under ice cooling. The temperature of the mixture was raised to room temperature, followed by stirring for 2 days. The precipitated crystals were collected by filtration and dried to give 2-methoxycarbonylthiophen-3-ylmethyltriphenylphosphonium bromide (10.57 g, 70.4%). This product was subjected to the reaction of Reference Example 13 without purification.

Reference Example 13

30 Methyl (E)-3-(3-nitrostyryl)-2-thiophenecarboxylate [1]

Methyl (Z)-3-(3-nitrostyryl)-2-thiophenecarboxylate [2]

To a solution of the phosphonium salt obtained in Reference Example 12 (2.26 g, 4.54 mmol) in tetrahydrofuran



(30 ml) was added sodium hydride (0.131 g, 5.45 mmol) under an atmosphere of argon gas at 0°C, followed by stirring at room temperature for 3 hours. To the resulting mixture was added 3-nitrobenzaldehyde (0.527 g, 3.49 mmol) at 0°C, and the mixture was stirred overnight at room temperature. After the reaction was completed, the reaction mixture was poured into an aqueous solution of ammonium chloride, followed by extraction with ethyl acetate. The organic layer was washed successively with water and a saturated aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure. The obtained residue was separated and purified by silica gel column chromatography (hexane/ethyl acetate = 20/1) to give [1] (0.48 g, 47.5%) and [2] (0.37 g, 36.6%), respectively.

[1] $^1\text{H-NMR}$ (CDCl_3) δ : 3.93(s, 3H), 7.14(d, 1H, $J=16.5\text{Hz}$), 7.46(d, 1H, $J=5.4\text{Hz}$), 7.49(d, 1H, $J=5.4\text{Hz}$), 7.54(t, 1H, $J=8.2\text{Hz}$), 7.90(d, 1H, $J=8.2\text{Hz}$), 8.09-8.16(m, 1H), 8.25(d, 1H, $J=16.5\text{Hz}$), 8.32-8.37(m, 1H)

[2] $^1\text{H-NMR}$ (CDCl_3) δ : 3.91(s, 3H), 6.68(d, 1H, $J=5.1\text{Hz}$), 6.76(d, 1H, $J=12.2\text{Hz}$), 7.24(d, 1H, $J=12.2\text{Hz}$), 7.30(d, 1H, $J=5.1\text{Hz}$), 7.33-7.43(m, 2H), 8.00-8.10(m, 2H)

Reference Example 14

Methyl 3-(3-trimethylacetamidostyryl)-2-thiophene-carboxylate

A mixture of the (E) form and the (Z) form obtained in Reference Example 13 (2.56 g, 8.84 mmol) was suspended in a mixed solvent of ethanol (50 ml) and water (5 ml), and reduced iron (2.50 g) and iron (III) chloride (0.250 g) were added thereto, followed by stirring under heating under reflux for 20 minutes. After the reaction was completed,



the solid was filtered off, followed by washing with ethyl acetate. To the filtrate was added a 2 N aqueous solution of sodium hydroxide (pH 9), and the organic layer was washed with a saturated aqueous solution of sodium chloride, and
5 then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure to give methyl 3-(3-aminostyryl)-2-thiophenecarboxylate as a crude product. This product was dissolved in dichloromethane (50 ml)
10 without purification, and triethylamine (1.79 g, 17.7 mmol) and trimethylacetyl chloride (2.13 g, 17.7 mmol) were added thereto at 0°C. The temperature of the mixture was raised to room temperature, followed by stirring for 2 hours. The obtained reaction mixture was poured into an aqueous
15 solution of sodium bicarbonate. The organic layer was washed with a saturated aqueous solution of sodium bicarbonate, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure. The
20 obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate = 10/1) to give methyl 3-(3-trimethylacetamidostyryl)-2-thiophenecarboxylate as an E/Z mixture. This mixture was subjected to the reaction of Reference Example 15 without further purification.

25

Reference Example 15

Methyl 3-[2-(3-trimethylacetamidophenyl)ethyl]-2-thiophenecarboxylate

The E/Z mixture of the amide obtained in Reference
30 Example 14 was dissolved in a mixed solvent of ethanol (30 ml) and acetic acid (10 ml), and 10% palladium/carbon (moisture content: 50%, 1.5 g) was added thereto under an atmosphere of argon gas, followed by replacement by hydrogen and heating under reflux for 2 hours. After the reaction
35 mixture was cooled to room temperature, the solid was filtered off and washed with ethyl acetate. The obtained



organic layer was washed successively with water and a saturated aqueous solution of sodium bicarbonate, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure. The obtained residue was triturated with hexane to give methyl 3-[2-(3-trimethylacetamido-phenyl)ethyl]-2-thiophenecarboxylate (2.55 g, 3 steps, 83.5%).

¹H-NMR (CDCl₃) δ: 1.32(s, 9H), 2.84-2.93(m, 2H), 3.25-3.34(m, 2H), 3.87(s, 3H), 6.89(d, 1H, J=5.0Hz), 6.96(br d, 1H, J=7.6Hz), 7.23(t, 1H, J=7.6Hz), 7.30(br, 1H), 7.31-7.48(m, 3H)

Reference Example 16

3-[2-(3-Trimethylacetamidophenyl)ethyl]-2-thiophenecarboxylic acid

The methyl ester obtained in Reference Example 15 (2.40 g, 6.95 mmol) was dissolved in methanol (20 ml), and a 1 N aqueous solution of sodium hydroxide (20 ml) was added thereto, followed by heating under reflux for 150 minutes. After the reaction was completed, methanol was distilled off under reduced pressure and the resulting mixture was adjusted to pH 3 with 2 N hydrochloric acid. The resulting mixture was extracted three times with ethyl acetate, and the extract was washed with a saturated aqueous solution of sodium chloride and then dried over anhydrous sodium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure. The obtained residue was triturated with diisopropyl ether to give 3-[2-(3-trimethylacetamidophenyl)ethyl]-2-thiophenecarboxylic acid (2.06 g, 89.5%).

¹H-NMR (DMSO-d₆) δ: 1.28(s, 9H), 2.83-2.92(m, 2H), 3.23-3.32(m, 2H), 6.97(br d, 1H, J=7.8Hz), 7.15(d, 1H, J=5.0Hz), 7.25(t, 1H, J=7.8Hz), 7.55(br



d, 1H, J=7.8Hz), 7.60(br, 1H), 7.78(d, 1H, J=5.0Hz), 9.91(br, 1H), 13.00(br, 1H)

Reference Example 17

- 5 7-Trimethylacetamido-4,5-dihydro-10H-benzo[5,6]
 cyclohepta[1,2-b]thiophen-10-one [1]
 9-Trimethylacetamido-4,5-dihydro-10H-benzo[5,6]
 cyclohepta[1,2-b]thiophen-10-one [2]

The carboxylic acid obtained in Reference Example 16
10 (1.90 g, 5.73 mmol) was suspended in dichloromethane (10
ml), and trifluoroacetic anhydride (1.46 ml, 10.3 mmol) was
added dropwise thereto under ice cooling, followed by
stirring at room temperature for one hour. After this
reaction mixture was ice cooled again, boron trifluoride
15 diethyl etherate (0.7 equivalent) was added thereto, and the
mixture was stirred overnight at room temperature. The
reaction mixture was poured into 2 N hydrochloric acid and
the resulting mixture was stirred at room temperature for
one hour, followed by extraction with ethyl acetate. The
20 organic layer was washed twice with an aqueous solution of
sodium bicarbonate and once with a saturated aqueous
solution of sodium chloride successively, and then dried
over anhydrous magnesium sulfate. After the drying agent
was filtered off, the organic layer was concentrated under
25 reduced pressure. The obtained mixture was separated and
purified by silica gel column chromatography (hexane/ethyl
acetate = 20/1) to give [1] (0.56 g, 31.2%) and [2] (0.24 g,
13.4%), respectively.

[1] ¹H-NMR (CDCl₃) δ: 1.34(s, 9H), 3.03-3.20(m, 4H),
30 6.94(d, 1H, J=5.0Hz), 7.28(dd, 1H, J=8.6, 2.3Hz),
 7.47(br, 1H), 7.56(d, 1H, J=5.0Hz), 7.80(d, 1H,
 J=2.3Hz), 8.02(d, 1H, J=8.6Hz)

[2] ¹H-NMR (CDCl₃) δ: 1.32(s, 9H), 3.00-3.20(m, 4H),
 6.93(d, 1H, J=5.3Hz), 6.97(dd, 1H, J=8.0, 0.9Hz),



7.40(t, 1H, J=8.0 Hz), 7.57(d, 1H, J=5.3Hz), 8.40
(dd, 1H, J=8.0, 0.9Hz), 10.86(br, 1H)

Reference Example 18

5 7-Amino-4,5-dihydro-10H-benzo[5,6]cyclohepta[1,2-b]
 thiophen-10-one

 In ethanol (10 ml) was suspended 7-trimethylacetamido-
4,5-dihydro-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one
obtained in Reference Example 17 (0.40 g, 1.28 mmol), and 6
10 N hydrochloric acid (5 ml) was added thereto, followed by
heating under reflux for 6 hours. After the reaction was
completed, ethanol was distilled off under reduced pressure,
and the resulting mixture was adjusted to pH 10 with an
aqueous solution of sodium hydroxide. The mixture was
15 extracted with dichloromethane, and the organic layer was
washed with a saturated aqueous solution of sodium
bicarbonate and then dried over anhydrous magnesium sulfate.
After the drying agent was filtered off, the organic layer
was concentrated under reduced pressure to give 7-amino-4,5-
20 dihydro-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one (0.29
g, 98.8%).

¹H-NMR (CDCl₃) δ: 3.06(br, 4H), 6.48(d, 1H, J=2.3Hz),
6.62(dd, 1H, J=8.6, 2.3Hz), 6.92(d, 1H, J=5.1Hz),
7.51(d, 1H, J=5.1Hz), 7.96(d, 1H, J=8.6Hz)

25

Reference Example 19

9-Amino-4,5-dihydro-10H-benzo[5,6]cyclohepta[1,2-b]
thiophen-10-one

 Substantially the same procedure as in Reference
30 Example 18 was repeated using 9-trimethylacetamido-4,5-
dihydro-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one
obtained in Reference Example 17 to give 9-amino-4,5-
dihydro-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one
(99.9%).



$^1\text{H-NMR}$ (CDCl_3) δ : 3.00-3.18(m, 4H), 6.52(d, 1H, $J=7.8\text{Hz}$), 6.61(d, 1H, $J=7.8\text{Hz}$), 6.91(d, 1H, $J=5.1\text{Hz}$), 7.14(t, 1H, $J=7.8\text{Hz}$), 7.51(d, 1H, $J=5.1\text{Hz}$)

Reference Example 20

Methyl (Z)-3-(3-acetamidostyryl)-2-thiophene-carboxylate

Substantially the same procedure as in Reference Example 14 was repeated using methyl (Z)-3-(3-nitrostyryl)-2-thiophenecarboxylate obtained in Reference Example 13 (3.51g, 12.1 mmol), and also using acetic anhydride (3.43ml, 36.3 mmol) instead of trimethylacetyl chloride, to give methyl (Z)-3-(3-acetamidostyryl)-2-thiophenecarboxylate (3.33g, 2 steps, 91.3%).

$^1\text{H-NMR}$ (DMSO-d_6) δ : 2.06(s, 3H), 3.88(s, 3H), 6.77(d, 1H, $J=5.3\text{Hz}$), 6.82(d, 1H, $J=12.3\text{Hz}$), 6.88(br d, 1H, $J=8.0\text{Hz}$), 7.09(d, 1H, $J=12.3\text{Hz}$), 7.26(t, 1H, $J=8.0\text{Hz}$), 7.48(br, 1H), 7.57(br d, 1H, $J=8.0\text{Hz}$), 7.74(d, 1H, $J=5.3\text{Hz}$), 9.96(br, 1H)

Reference Example 21

(Z)-3-(3-Acetamidostyryl)-2-thiophenecarboxylic acid

Substantially the same procedure as in Reference Example 16 was repeated using the methyl ester obtained in Reference Example 20 (3.33g, 11.1 mmol) to give (Z)-3-(3-acetamidostyryl)-2-thiophenecarboxylic acid (2.95g, 92.9%).

$^1\text{H-NMR}$ (DMSO-d_6) δ : 2.05(s, 3H), 6.74(d, 1H, $J=5.1\text{Hz}$), 6.77(d, 1H, $J=11.9\text{Hz}$), 6.88(br d, 1H, $J=7.8\text{Hz}$), 7.12(d, 1H, $J=11.9\text{Hz}$), 7.25(t, 1H, $J=7.8\text{Hz}$), 7.47(br, 1H), 7.57(br d, 1H, $J=7.8\text{Hz}$), 7.66(d, 1H, $J=5.1\text{Hz}$), 9.96(br, 1H)

Reference Example 22

7-Acetamido-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one [1]

9-Acetamido-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one [2]

Substantially the same procedure as in Reference Example 17 was repeated using the carboxylic acid obtained in Reference Example 21 (2.80g, 9.75 mmol) to give [1] (0.94g, 42.9%) and [2] (0.78g, 29.7%), respectively.

[1] $^1\text{H-NMR}$ (DMSO-d_6) δ : 2.21(s, 3H), 7.41(d, 1H, $J=11.5\text{Hz}$), 7.51(d, 1H, $J=11.5\text{Hz}$), 7.66(d, 1H, $J=4.9\text{Hz}$), 7.99(dd, 1H, $J=9.0, 2.0\text{Hz}$), 8.23(d, 1H, $J=4.9\text{Hz}$), 8.25(d, 1H, $J=2.0\text{Hz}$), 8.67(d, 1H, $J=9.0\text{Hz}$), 10.59(d, 1H, $J=8.6\text{Hz}$)

[2] $^1\text{H-NMR}$ (CDCl_3) δ : 2.31(s, 3H), 7.21(d, 1H, $J=12.0\text{Hz}$), 7.30(d, 1H, $J=12.0\text{Hz}$), 7.31(d, 1H, $J=5.3\text{Hz}$), 7.45(dd, 1H, $J=8.3, 0.7\text{Hz}$), 7.71(t, 1H, $J=8.3\text{Hz}$), 7.79(d, 1H, $J=5.3\text{Hz}$), 8.99(dd, 1H, $J=8.3, 0.7\text{Hz}$), 12.87(br, 1H)

Reference Example 23

7-Amino-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one

Substantially the same procedure as in Reference Example 18 was repeated using 7-acetamido-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one obtained in Reference Example 22 to give 7-amino-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one (87.2%).

$^1\text{H-NMR}$ (CDCl_3) δ : 6.86(d, 1H, $J=2.3$ Hz), 6.99(dd, 1H, $J=8.9, 2.3\text{Hz}$), 7.08(d, 1H, $J=11.6\text{Hz}$), 7.19(d, 1H, $J=11.6\text{Hz}$), 7.30(d, 1H, $J=5.3\text{Hz}$), 7.69(d, 1H, $J=5.3\text{Hz}$), 8.71(d, 1H, $J=8.9\text{Hz}$)

Reference Example 24

5 9-Amino-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one

5
6
7
8
9
10

11
12
13
14

15
16
17



Substantially the same procedure as in Reference Example 18 was repeated using 9-acetamido-10H-benzo[5,6]-cyclohepta[1,2-b]thiophen-10-one obtained in Reference Example 22 to give 9-amino-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one (99.9%).

$^1\text{H-NMR}$ (CDCl_3) δ : 6.84(dd, 1H, $J=7.9$, 1.3Hz), 6.92(br d, 1H, $J=7.9\text{Hz}$), 6.97(d, 1H, $J=11.7\text{Hz}$), 7.10(d, 1H, $J=11.7\text{Hz}$), 7.20(d, 1H, $J=5.1\text{Hz}$), 7.38(t, 1H, $J=7.9\text{Hz}$), 7.66(d, 1H, $J=5.1\text{Hz}$)

10

Reference Example 25

2-Acetamido-5H-dibenzo[a,d]cyclohepten-5-one

In a mixed solvent of carbon tetrachloride (45 ml) and chloroform (25 ml) was dissolved 2-acetamido-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-one obtained substantially in the same manner as in Reference Examples 12 to 17 (2.60 g, 9.80 mmol), and NBS (1.92 g, 10.8 mmol) and AIBN (0.6 g) were added thereto, followed by heating under reflux for 5 hours. The reaction mixture was washed successively with water and a saturated aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure. The obtained residue was dissolved in dimethylformamide (100 ml), and DBU (2.60 ml) was added thereto, followed by stirring at 60°C for 2 hours. The reaction mixture was poured into ice-an aqueous solution of ammonium chloride, followed by extraction with ethyl acetate. The organic layer was washed successively with water and a saturated aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate. After the drying agent was filtered off, the organic layer was concentrated under reduced pressure. The obtained residue was triturated with diisopropyl ether to give 2-acetamido-5H-dibenzo[a,d]cyclohepten-5-one (0.97 g, 37.6%).

AUSTRALIAN

$^1\text{H-NMR}$ (DMSO-d_6) δ : 2.18(s, 3H), 7.18(d, 1H, $J=12.2\text{Hz}$), 7.28(d, 1H, $J=12.2\text{Hz}$), 7.63-7.74(m, 1H), 7.77-7.86(m, 3H), 8.02(d, 1H, $J=2.0\text{Hz}$), 8.13-8.22(m, 2H), 10.47(br, 1H)

Reference Example 26

5 2-Amino-5H-dibenzo[a,d]cyclohepten-5-one

Substantially the same procedure as in Reference Example 18 was repeated using 2-acetamido-5H-dibenzo[a,d]-cyclohepten-5-one obtained in Reference Example 25 (0.70g, 2.66 mmol) to give 2-amino-5H-dibenzo[a,d]cyclohepten-5-one(0.59g, 99.9%).

$^1\text{H-NMR}$ (CDCl_3) δ : 4.13(br, 2H), 6.71(d, 1H, $J=2.7\text{Hz}$), 6.83(dd, 1H, $J=8.6$, 10 2.7Hz), 6.89(d, 1H, $J=12.2\text{Hz}$), 7.00(d, 1H, $J=12.2\text{Hz}$), 7.48-7.65(m, 3H), 8.19(d, 1H, $J=8.6\text{Hz}$), 8.24-8.34(m, 1H)

Reference Example 27

(S)-3,3,3-trifluoro-2-hydroxy-2-methylpropionic acid

To a solution of 3,3,3-trifluoro-2-hydroxy-2-methyl-propionic acid (13.0g, 82.3 15 mmol) in ethanol (60ml) was added dropwise (S)-(-)-1-phenylethylamine (9.97g, 82.3 mmol) at 0°C , followed by stirring at room temperature for one hour. The resulting mixture was concentrated under reduced pressure to give a salt as white crystals. This salt was recrystallised from a 10% butanol/toluene solution (60ml) six times to give S-form of 3,3,3-trifluoro-2-hydroxy-2-methylpropionic acid phenylethylamine salt {(S,S) 20 salt} (3.85g, 13.8 mmol, 97% e.e.). The optical purity was determined with $^1\text{H-NMR}$. The obtained {(S,S) salt} was dissolved in 2 N hydrochloric acid, and diethyl ether was added thereto, followed by stirring at room temperature for one hour. The organic layer was washed successively with water and a saturated aqueous solution of sodium chloride, and then dried over anhydrous sodium sulfate. After the



drying agent was filtered off, the organic layer was concentrated under reduced pressure to give (S)-3,3,3-trifluoro-2-hydroxy-2-methylpropionic acid as white crystals (2.13 g, 13.5 mmol).

5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.65(s, 3H)

Melting point: 102-104°C

Reference Example 28

(R)-3,3,3-Trifluoro-2-hydroxy-2-methylpropionic acid

10 Substantially the same procedure as in Reference Example 27 was repeated using (R)-(-)-1-phenylethylamine instead of (S)-(-)-1-phenylethylamine to give (R)-3,3,3-trifluoro-2-hydroxy-2-methylpropionic acid.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.65(s, 3H)

15 Melting point: 101-105°C


Industrial Applicability

The present invention provides tricyclic compounds which are useful as therapeutic agents for urinary
20 incontinence.



The claims defining the invention are as follows:

1. A tricyclic compound represented by general formula (I):



- wherein R¹ represents hydrogen, lower alkyl, lower alkoxy, or halogen; -X¹-X²-X³- represents -CR²=CR³-CR⁴=CR⁵- (wherein R², R³, R⁴, and R⁵, which may be the same or different, each represents hydrogen, lower alkyl, lower alkoxy, or halogen), -CR²=CR³-CR⁴=N- (wherein R², R³, and R⁴ have the same significances as defined above), -CR²=CR³-S- (wherein R² and R³ have the same significances as defined above), -CR²=CR³-O- (wherein R² and R³ have the same significances as defined above), -S-CR⁴=CR⁵- (wherein R⁴ and R⁵ have the same significances as defined above), or -O-CR⁴=CR⁵- (wherein R⁴ and R⁵ have the same significances as defined above); and Y represents -CH₂O-, -CH₂S-, -CH₂SO-, -CH=CH-, or -(CH₂)_n- (wherein n represents 0, 1, or 2).

2. A tricyclic compound according to claim 1, wherein -X¹-X²-X³- is -CR²=CR³-CR⁴=CR⁵- (wherein R², R³, R⁴, and R⁵ have the same significances as defined above) or -CR²=CR³-CR⁴=N- (wherein R², R³, and R⁴ have the same significances as defined above).

3. A tricyclic compound according to claim 1, wherein -X¹-X²-X³- is -CR²=CR³-S- (wherein R² and R³ have the same significances as defined above) or -CR²=CR³-O- (wherein R² and R³ have the same significances as defined above).

4. A tricyclic compound according to claim 1, wherein -X¹-X²-X³- is -S-CR⁴=CR⁵- (wherein R⁴ and R⁵ have the same significances as defined above) or -O-CR⁴=CR⁵- (wherein R⁴ and R⁵ have the same significances as defined above).

5. A tricyclic compound according to claim 1, 2, 3, or 4, wherein Y is -CH₂O-.

6. A tricyclic compound according to claim 1, 2, 3, or 4, wherein Y is -(CH₂)_n- (wherein n has the same significance as defined above).

7. A tricyclic compound according to claim 6, wherein n is 0.

8. A tricyclic compound according to claim 6, wherein n is 2.

9. A tricyclic compound according to claim 1, 2, 3, or 4, wherein Y is -CH=CH-.

10. A tricyclic compound according to claim 1, 2, 3, or 9, wherein Y is -CH₂S- or -CH₂SO-.

11. 1-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-6,11-dihydro dibenz[b,e]oxepin-11-one

12. 3-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-6,11-dihydro dibenz[b,e]oxepin-11-one

5 13. 2-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-6,11-dihydro dibenz[b,e]oxepin-11-one

14. 1-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-fluoren-9-one

15. 2-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-fluoren-9-one

16. 3-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-fluoren-9-one

10 17. 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydro thieno[3,2-c][1]benzoxepin-10-one

18. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydro thieno[3,2-c][1]benzoxepin-10-one

19. 5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydrofuro [2,3-c][1]benzoxepin-4-one

20. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydrofuro [2,3-c][1]benzoxepin-4-one

21. 3-Chloro-2-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)fluoren-9-one

22. 3-Bromo-2-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)fluoren-9-one

23. 7-Bromo-2-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)fluoren-9-one

24. 4-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-fluoren-9-one

25. 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydrofuro [3,2-c][1]benzoxepin-10-one

26. 8-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydro thieno[3,2-c][1]benzoxepin-10-one

27. 5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydro thieno[2,3-c][1]benzoxepin-4-one

28. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydro thieno[2,3-c][1]benzoxepin-4-one

29. 6-Bromo-9-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-4,10-dihydro thieno[3,2-c][1]-benzoxepin-10-one

30. 7-Fluoro-1-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one

31. 7-Fluoro-3-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one

5 32. 3-Methyl-1-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one

33. 3-Methoxy-1-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one

10 34. 2-Methyl-1-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one

35. 2-Methyl-3-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-6,11-dihydrodibenz[b,e]oxepin-11-one

36. 6-Methyl-9-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-9,10-dihydrothieno[3,2-c][1]-benzoxepin-10-one

15 37. 6-Methoxy-9-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-4,10-dihydrothieno[3,2-c][1]-benzoxepin-10-one

38. 8-Methyl-5-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-4,10-dihydrofuro[2,3-c][1]benzoxepin-4-one

20 39. 8-Methoxy-5-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-4,10-dihydrofuro[2,3-c][1]benzoxepin-4-one

40. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-5,11-dihydro pyrido[2,3-c][1]benzoxepin-5-one

41. 7-Methoxy-6-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-5,11-dihydropyrido[2,3-c][1]-benzoxepin-5-one

25 42. 7-Methyl-6-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-5,11-dihydropyrido[2,3-c][1]-benzoxepin-5-one

43. (S)-9-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydro thieno[3,2-c][1]benzoxepin-10-one

30 44. (R)-9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydro thieno[3,2-c][1]benzoxepin-10-one

45. (S)-5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydro furo[2,3-c][1]benzoxepin-4-one

46. (R)-5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydro furo[2,3-c][1]benzoxepin-4-one

47. (S)-7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydro furo[2,3-c][1]benzoxepin-4-one

48. (R)-7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydro furo[2,3-c][1]benzoxepin-4-one

5 49. (S)-8-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydro thieno[3,2-c][1]benzoxepin-10-one

50. 1-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-6,11-dihydro dibenzo[b,e]thiepin-11-one

10 51. 3-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-6,11-dihydro dibenzo[b,e]thiepin-11-one

52. 5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydrofuro [2,3-c][1]benzothiepin-4-one

53. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydrofuro [2,3-c][1]benzothiepin-4-one

15 54. 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydro thieno[3,2-c][1]benzothiepin-10-one

55. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydro thieno[3,2-c][1]benzothiepin-10-one

20 56. 5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydro thieno[2,3-c][1]benzothiepin-4-one

57. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,10-dihydro thieno[2,3-c][1]benzothiepin-4-one

58. 5-Oxo-1-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl-amino)-6,11-dihydrodibenzo[b,e]thiepin-11-one

25 59. 5-Oxo-3-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl-amino)-6,11-dihydrodibenzo[b,e]thiepin-11-one

60. 9-Oxo-5-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydrothieno[2,3-c][1]benzothiepin-4-one

30 61. 9-Oxo-7-(3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl-amino)-4,10-dihydrothieno[2,3-c][1]benzothiepin-4-one

62. 4-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-one

63. 3-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-one

64. 2-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-one

65. 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,5-dihydro-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one

5 66. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,5-dihydro-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one

67. 5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-9,10-dihydro-4H-benzo[4,5]cyclohepta[1,2-b]thiophen-4-one

10 68. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-9,10-dihydro-4H-benzo[4,5]cyclohepta[1,2-b]furan-4-one

69. 8-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10,11-dihydro-5H-benzo[4,5]cyclohepta[1,2-b]pyridin-5-one

70. 4-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-5H-dibenzo[a,d]cyclohepten-5-one

15 71. 3-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-5H-dibenzo[a,d]cyclohepten-5-one

72. 2-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-5H-dibenzo[a,d]cyclohepten-5-one

20 73. 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one

74. 8-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one

75. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10H-benzo[5,6]cyclohepta[1,2-b]thiophen-10-one

25 76. 5-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4H-benzo[4,5]cyclohepta[1,2-b]thiophen-4-one

77. 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10H-benzo[5,6]cyclohepta[1,2-b]furan-10-one

30 78. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10H-benzo[5,6]cyclohepta[1,2-b]furan-10-one

79. 8-Chloro-7-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-4,10-dihydrothieno[3,2-c][1]-benzoxepin-10-one

80. 7-Methyl-8-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-4,10-dihydrothieno[3,2-c][1]-benzoxepin-10-one



81. 7-Methyl-9-(3,3,3-trifluoro-2-hydroxy-2-methyl-propanoylamino)-5,11-dihydropyrido[2,3-c][1]-benzoxepin-5-one

82. 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,5-dihydro-10H-benzo[5,6]cyclohepta[1,2-b]furan-10-one

83. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-4,5-dihydro-10H-benzo[5,6]cyclohepta[1,2-b]furan-10-one

84. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-5H-benzo[4,5]cyclohepta[1,2-b]pyridin-5-one

85. 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-5H-benzo[4,5]cyclohepta[1,2-b]pyridin-5-one

86. 7-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10,11-dihydro-5H-benzo[4,5]cyclohepta[1,2-b]pyridin-5-one

87. 9-(3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylamino)-10,11-dihydro-5H-benzo[4,5]cyclohepta[1,2-b]pyridin-5-one

88. A tricyclic compound active against urinary incontinence, substantially as hereinbefore described with reference to any one of the examples.

89. A pharmaceutical composition comprising an effective amount of at least one compound according to any one of claims 1 to 88, together with a pharmaceutically acceptable carrier, diluent or adjuvant therefor.

90. A method for the treatment or prophylaxis of urinary incontinence in a mammal requiring said treatment or prophylaxis, which method comprises administering to said mammal an effective amount of at least one compound according to any one of claims 1 to 88, or of a composition according to claim 89.

Dated 11 July 1997

KYOWA HAKKO KOGYO Co., LTD.


Patent Attorneys for the Applicant/Nominated Person

SPRUSON & FERGUSON



ABSTRACT

The present invention relates to tricyclic compounds represented by general formula (I) which are useful as therapeutic agents for urinary incontinence:



wherein R^1 represents hydrogen and the like; $-X^1-X^2-X^3-$ represents $-CR^2=CR^3-CR^4=CR^5-$ (wherein R^2 , R^3 , R^4 , and R^5 represent hydrogen and the like) and the like; and Y represents $-CH_2O-$ and the like.

