

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199892771 B2
(10) Patent No. 747150

(54) Title
Pyridyl compounds and pharmaceutical compositions containing them

(51)⁷ International Patent Classification(s)
C07D 213/79 C07D 213/80
A61K 031/44 C07D 405/12

(21) Application No: **199892771**

(22) Application Date: **1998.10.07**

(87) WIPO No: **WO99/19302**

(30) Priority Data

(31) Number	(32) Date	(33) Country
2364/97	1997.10.09	CH

(43) Publication Date : **1999.05.03**

(43) Publication Journal Date : **1999.07.01**

(44) Accepted Journal Date : **2002.05.09**

(71) Applicant(s)
Cermol S.A.


(72) Inventor(s)
**Pierre Statkow; Danielle Straumann; Shyam S. Chatterjee; Carlos Sunkel
Letelier; Miguel Fau De Casa-Juana Munoz; Julio Alvarez-Builla Gomez; Jose
M. Minguez Ortega**

(74) Agent/Attorney
**PHILLIPS ORMONDE and FITZPATRICK,367 Collins Street,MELBOURNE VIC
3000**



<p>(51) International Patent Classification ⁶ : C07D 213/79, 213/80, 405/12, A61K 31/44</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/19302 (43) International Publication Date: 22 April 1999 (22.04.99)</p>
<p>(21) International Application Number: PCT/IB98/01555 (22) International Filing Date: 7 October 1998 (07.10.98) (30) Priority Data: 2364/97 9 October 1997 (09.10.97) CH (71) Applicant (for all designated States except US): CERMOL S.A. [CH/CH]; CH-1902 Evionnaz (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): STATKOW, Pierre [CH/CH]; 6, rue du Roveray, CH-1206 Genève (CH). STRAUMANN, Danielle [CH/CH]; 6, rue Pré Borvey, CH-1920 Martigny (CH). CHATTERJEE, Shyam, S. [DE/DE]; Stettiner Strasse 1, D-76139 Karlsruhe (DE). SUNKEL LETELIER, Carlos [ES/ES]; Calle Almarza, 47, E-28033 Madrid (ES). FAU DE CASA-JUANA MUNOZ, Miguel [ES/ES]; Calle Torquemada, 17, E-28043 Madrid (ES). ALVAREZ-BUILLA GOMEZ, Julio [ES/ES]; Calle Condado de Trevino, 9, E-28033 Madrid (ES). MINGUEZ ORTEGA, José M. [ES/ES]; Calle Palestina, 17, E-28820 Coslada (ES). (74) Agent: LEGER, Jean-François; Kirker & Cie S.A., 122, rue de Genève, P.O. Box 324, CH-1226 Thônex (CH).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i></p>	

(54) Title: PYRIDYL COMPOUNDS AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM




(57) Abstract

The present invention is concerned with new pyridine double esters of formula (I), their acids, and pharmaceutically acceptable salts. These compounds can be obtained by oxydation of the corresponding 1,4-dihydropyridines, and they are useful as cardioprotective agents in pharmaceutical compositions.

PYRIDYL COMPOUNDS AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM.

The present invention relates to a series of new pyridine derivatives, their preparation procedures, their use as cardioprotective agents, and the pharmaceutical compositions containing said derivatives. More particularly, the object of the present invention is constituted by pyridine derivatives of general formula (I), in which



(I)

- X represents H, NO₂, OH, CN, F, Br, Cl, CF₃, -NH-CO-CH₃, 2,3-Cl₂, 2,3-(OCH₃)₂, 4-Cl, 3-NO₂ and 2-[CH=CH-COO-C(CH₃)₂], and the fused cycle 2,3-oxadiazole [2,3-(=N-O-N=)];
- Y represents an alcoxycarbonyl fragment R²OC-, where R² can be H, a small alkyl group, 2-méthoxyethyl, 2-tétrahydrofurfuryl or 5,5-diméthyl-1,3,2-dioxaphosphorinan-2-yl;
- R represents H, an small alkyl group or, alternatively, the following fragments: 2-methoxyethyl, 2-(benzyl-methylamino)ethyl, 2-tétrahydrofurfuryl, 5-oxotétrahydrofurfuryl, 3-[(4,4-diphenyl)-1-piperidinyl]propyl, (R)-1-benzyl-3-piperidinyl, 2-(N-benzyl-(N-phenylamino)ethyl), (S)-1-benzyl-3-pirrolidinyl, 2-[(4-diphenylmethyl)-1-piperazinylethyl, 3-phényl-2-propenyl, dually-amino-alkyl, 2-(N-morpholino)ethyl or 1,3-di(N-morpholino)isopropyl
- R' represents CH₃, CN or -CH₂-O-(CH₂)₂-NH₂, as well as the corresponding optical isomers, and the pharmaceutically acceptable salts of the said compounds.

As non-limiting examples, the following compounds corresponding to formula (1) of the present invention can be mentioned:

4-(2,3-Dichloro-phenyl)-2,6-dimethyl-pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(3-fluorophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

4-(3-Acetylamino-phenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester (hydrochloride)

2,6-Dimethyl-4-(3-hydroxyphenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester (hydrochloride)

2,6-Dimethyl-4-phenyl-pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(3-nitrophenyl)-pyridine-3, 5-dicarboxylic acid 3,5-di(tetrahydrofuran-2-ylmethyl) ester (hydrochloride)

4-(3-Cyanophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(3-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

4-(4-Chloro-3-nitrophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

4-(3-Bromophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3,5-di(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(3-trifluoromethylphenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

4-(2-Chlorophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(2-trifluoromethylphenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(2,3-dimethoxyphenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

4-(3-Chlorophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(5-oxotetrahydrofuran-2-ylmethyl) ester

(*S*)-2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

(*R*)-2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

4-(2-Bromophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(3-nitrophenyl)-pyridine-3, 5-dicarboxylic acid 3-isopropyl ester 5-(2-methoxyethyl) ester

2,6-Dimethyl-4-(2-nitrophenyl)-pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-isobutyl ester

2,6-Dimethyl-4-(3-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-ethyl ester 5-methyl ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid dimethyl ester

4-[2-(2-*tert*-Butoxycarbonylvinyl)-phenyl]-2,6-dimethyl pyridine-3, 5-dicarboxylic acid diethyl ester

2,6-Dimethyl-4-(3-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(*N*-methylbenzylamino)ethyl ester

4-(2,3-Dichlorophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-ethyl ester 5-methyl ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid

2-(2-Aminoethoxy)methyl-4-(2-chlorophenyl)-6-methyl pyridine-3, 5-dicarboxylic acid 3-ethyl ester 5-methyl ester

2,6-Dimethyl-4-(3-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-(2-methoxyethyl) ester 5-(3-phenyl)propen-2-yl ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-(2-dimethylaminoethyl) ester 5-methyl ester (dihydrobromide)

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-(2-diethylaminoethyl) ester 5-methyl ester (dihydrochloride)


2,6-Dimethyl-4-phenylpyridine-3, 5-dicarboxylic acid dimethyl ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-(3-dimethylaminopropyl) 5-methyl ester (dihydrochloride).

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(N-morpholino)ethyl ester (dihydrochloride)

2,6-Dimethyl-4-(2-nitrophenyl)pyridine-3,5-dicarboxylic acid 3-methyl ester 5-[1,3-di(N-morpholine)isopropyl] ester trihydrochloride

The compounds corresponding to formula (Ia, examples 1-30) could, in a general method, be prepared by oxydation of 1,4-dihydropyridines corresponding to formula (II). In some cases, when X=2-NO₂, many dihydropyridines II could be rather unstable, so products had to be obtained through Ib and III, (Scheme I), allowing the preparation of new derivatives Ic. In the scheme, X, R and R' are like have been defined before, and R'', although differentiated for clarity in the scheme, should be taken as defined for R.



The 1,4-dihydropyridine derivatives (II) are either already known, or could be prepared by the standard procedures described in the literature.

As oxydation agents for preparation of the compounds (Ia) it is possible to use all described for standard oxydation procedures, as oxygen, $\text{NaNO}_2/\text{HOAc}$, NaNO_2/HCl , HNO_3 , $\text{Fe}(\text{NO}_3)_3$ or $\text{Cu}(\text{NO}_3)_2$, $\text{Br}_2 + \text{NaAcO}$, CrO_3 , sulphur, KMnO_4 , chloranyl or o-chloranyl, 2,3-dichloro-5, 6-dicyano-1,4-benzoquinone (DDQ), Pd/C, pyridinium chlorochromate adsorbed on alumina ($\text{PCC}/\text{Al}_2\text{O}_3$), pyridinium dichromate, MnO_2 , etc. Controlled ester hydrolysis of Ia, to produce Ib, activation through III and esterification to Ic are described in the present report.

The pharmaceutically acceptable salts obtained by addition of acid to the compounds (I) are prepared by conventional methods, by treatment of a solution or suspension of the free base (I) with one or two equivalents of a pharmaceutically acceptable acid, either organic or inorganic. To give some examples, it is possible to mention the following acids: hydrochloric, hydrobromic, sulphuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, lactic, p-toluenesulphonic, gluconic, fumaric, succinic, ascorbic, maleic, methanesulphonic and benzenesulphonic. The obtained salts can show some advantages, particularly in relation with the higher solubility in polar solvents as water. This can facilitate the preparation of galenic forms requiring the administration of the product dissolved in water.

The mixtures of diastereomers or enantiomers can be separated using the differences of physico-chemical properties of the products, through the usual methods as fractional crystallisation, chromatography, reactions with asymmetric induction, or by the action of enzymes or micro-organisms.

The present inventors had put in evidence the fact that the pyridine compounds with formula (I), according the present invention, have a cardioprotective activity and thus, can be used as cardioprotective agents. In addition, another object of the present invention consists in pharmaceutical preparations containing at least, one compound of formula (I), as previously defined, or one of its pharmaceutically acceptable salts.

The compounds of formula (I) according the invention can be administered alone but, in general, they would be administered as a mixture with a selected pharmaceutical excipient, depending of the route of administration and the standard pharmaceutical practice. As an example, they can be administered by oral route, either in the form of tablets, containing excipients as starch or lactose, or in capsules, either alone or mixed with excipients, or in the form of syrups or suspensions, containing colour or aromatic substances. They can also be injected by parental route, as by example, by intramuscular, intravenous or subcutaneous route. When administered by parenteral route, they will be preferably used in the form of sterile aqueous solution, which can contain other solutes, as for instance, salt or glucose, to render the solution isotonic.

The pharmaceutical compositions according the invention could contain a quantity of any of the products with general formula (I) in the way the level of the administered dose being between 0,01 and 20 mg/kg. The daily dose of the active principle depends of the administration route. In general, an oral dose between 5 and 1000 mg/day will be used.

While when administered intramuscular route, the product can be given either in one dose or divided up to three doses, when intravenously administered the product can be included into a dropwise system, for continuous supply. There will be necessarily variations depending of the weight and conditions of the patient, as well as of the administration route chosen.

In that way, the pharmaceutical compositions according the present invention, can be used as new drugs for the prevention or treatment of cardiovascular diseases, as cardiomyopaties, myocardium infarction, angina, cardiac failures, coronary vasospasm, valvular heart disease, etc.


Throughout the description and claims of the specification the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.



The present invention will be illustrated in reference to some examples of preparation of compounds according the invention, as well as the results of the tests of toxicity and pharmacological activity.

Example 1

2,6-Dimethyl-4-phenyl pyridine-3,5-dicarboxylic acid dimethyl ester




Over a suspension of 119 g (0,1 mol) of pyridinium chlorochromate absorbed on alumina in 330 ml of CH_2Cl_2 , 10 g (0,033 mol) of 2,6-dimethyl-4-phenyl-1,4-dihydropyridine-3,5-dicarboxylic acid dimethyl ester were portionwise added with stirring. The mixture was maintained with stirring at room temperature for 8 hours. The remaining solid was eliminated by filtration, and the liquid was washed with water (3x500 ml), dried over anh. Na_2SO_4 , and concentrated at reduced pressure, to dryness. The product was obtained as a solid, (mp. 133-5°C, MeOH) with a yield of 75%.

Analysis calculated for $\text{C}_{17}\text{H}_{17}\text{NO}_4$

	%C	%H	%N
Calc.	68.22	5.72	4.68
Obs.	68.09	5.63	4.64

Example 2

2,6-Dimethyl-4-phenyl-pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester



A) 2,6-Dimethyl-4-phenyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A solution of 20 g (0.10 mol) of methyl benzylideneacetoacetate and 18.14 g (0.10 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 100 ml of 2-propanol was refluxed with stirring along 24 hours. The solvent was evaporated at reduced pressure, and the residue obtained was crystallised in 15 ml of diisopropylether. A white solid was obtained (mp 117-9°C, diisopropylether) with a yield of 82%.

B) 2,6-Dimethyl-4-phenyl-pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester


A mixture of 8 g (0.022 mol) of 2,6-dimethyl-4-phenyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester, obtained as described in A), and 2.15 g (0.025 mol) of MnO₂ suspended in 15 ml of methylisobutylketone was refluxed with stirring along 14 hours. The remaining solid was filtered of, and the solution obtained was concentrated at reduced pressure. The residue obtained was then dissolved in 15 ml of aq HCl 5%, washed with EtAcO (20 ml), and the aqueous phase was taken to pH 8 with aq K₂CO₃ 20%. Then, the aqueous phase was extracted with EtAcO (2x50 ml) and the organic extracts were dried (anh. Na₂SO₄) and concentrated to dryness under vacuum. An oil was obtained, which crystallised in diisopropylether (2 ml), producing a white solid (45-7°C, diisopropylether) with a yield of 59%.

Analysis calculated for C₂₁H₂₃NO₅

	%C	%H	%N
Calc.	68.28	6.28	3.79
Obs.	68.03	6.26	3.85

Example 3

2,6-Dimethyl-4-(3-fluorophényl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester



A) 2,6-Dimethyl-4-(3-fluorophényl) 1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A solution of 25 g (0.11 mol) of methyl 3-fluoro-benzylideneacetoacetate and 20.84 g (0.11 mol) de (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 110 ml of abs. ethanol was refluxed with stirring for 8 hours. Then, the solvent was evaporated under vacuum, up to 60 ml and the mixture was cooled to -10°C ; a precipitate was obtained and on filtration, the product was obtained as a clear yellow solid, with a m.p. of $118-20^{\circ}\text{C}$ (EtOH), and with a yield of 88%.

B) 2,6-Dimethyl-4-(3-fluorophényl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester


To a solution of 8 g (0.021 mol) of 2,6-dimethyl-4-(3-fluorophényl) 1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester, (obtained as indicated before) in 20 ml of acetic acid, and with stirring, 2.06 g (0.21 mol) of CrO_3 were slowly added, with stirring. Once the addition was complete, reflux with stirring was maintained for one hour. Then, the reaction mixture was cooled at room temperature and added to a mixture of 50 ml NH_4OH and ice. The resulting solution was then extracted with CH_2Cl_2 (3x100 ml), and the organic extracts were washed with water (250 ml) and dried with anh. Na_2SO_4 . Finally, the solvent was evaporated at reduced pressure, and an oil was obtained which crystallised with 5 ml of pentane. The product was obtained as a white solid, of m.p. $66-8^{\circ}\text{C}$ (EtOH), with a yield of 65%.

Analysis calculated for $\text{C}_{21}\text{H}_{22}\text{NFO}_5$

	%C	%H	%N
Calc.	65.11	5.72	3.62
Obs.	65.23	5.95	3.45

Example 4

4-(2-Chlorophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester



A) 4-(2-Chlorophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A solution of 15 g (0.06 mol) of methyl 2-chlorobenzylideneacetoacetate and 11,64 g (0,06 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 65 ml of abs ethanol, was refluxed with stirring in the absence of light, along 18 hours. Then 30 ml of solvent were evaporated, and the remaining solution was cooled to -10°C . In that way, a yellow solid was obtained (mp $165-7^{\circ}\text{C}$, EtOH) with a yield of 86%.

B) 4-(2-Chlorophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester


A mixture of 8 g (0,02 mol) of 4-(2-chlorophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester, obtained according A) and 71 g (0,059 mol) of pyridinium chlorochromate adsorbed on alumina, was suspended in 200 ml of CH_2Cl_2 , and the whole mixture was stirred at room temperature for 5 hours. The remaining solid was eliminated by filtration, and the liquid was washed with water (3x250 ml), dried over anh. Na_2SO_4 , and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 8:2) with a yield of 66%,

Analysis calculated for $\text{C}_{21}\text{H}_{22}\text{ClNO}_5$

	%C	%H	%N
Calc.	62.46	5.49	3.47
Obs.	62.43	5.57	3.59

Example 5

4-(3-Chlorophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester



A) 4-(3-Chlorophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A solution of 15 g (0,06 mol) of methyl 3-chlorobenzylideneacetoacetate and 11,64 g (0,06 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 65 ml of abs. ethanol, was refluxed with stirring and in the absence of light, for 8 hours. Then, the reaction mixture was cooled to -10°C and a white solid precipitated ($137-9^{\circ}\text{C}$, MeOH) with a yield of 87%.

B) 4-(3-Chlorophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester


A mixture of 8 g (0,02 mol) de 4-(3-chlorophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester, obtained according A), and 71 g (0,059 mol) of pyridinium chlorochromate absorbed on alumina, was suspended in 200 ml of CH_2Cl_2 . Then, the whole mixture was maintained with stirring at room temperature, along 8 hours. The remaining solid was eliminated by filtration, and the liquid was washed with water (3x250 ml), dried over anh. Na_2SO_4 , and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 8:2) with a yield of 70%.

Analysis calculated for $\text{C}_{21}\text{H}_{22}\text{ClNO}_5$

	%C	%H	%N
Calc.	62.46	5.49	3.47
Obs.	62.40	5.68	3.30

Example 6

4-(2,3-Dichlorophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester



A) 4-(2,3-Dichlorophenyl)-2,6-dimethyl 1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A solution of 20g (0.07mol) of methyl 2,3-dichlorobenzylidenacetoacetate and 13,56g (0.07mol) de (tetrahydrofuran-2-ylmethyl) 3-aminocrotonate in 75 ml of abs. ethanol was refluxed with stirring for 8 hours, in the absence of light. The reaction mixture was then cooled to -10°C , and the resulting precipitate was filtered. The product, with a m.p. of $172-3^{\circ}\text{C}$ (EtOH), was obtained with a yield of 79%.

B) 4-(2,3-Dichlorophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester


To a suspension of 82g (0.068 mol) of pyridinium chlorochromate on alumina in 230 ml of CH_2Cl_2 , 10g (0.023 mol) of 4-(2,3-dichlorophenyl)-2,6-dimethyl 1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester were added, and the mixture was maintained with strong stirring during 7 hours. The insoluble solid was filtered and the remaining liquid was washed with water (3x250 ml), dried with anh. Na_2SO_4 and concentrated at low pressure. Finally, the product was obtained as an oil, with a yield of 87%.

Analysis: Calculated for $\text{C}_{21}\text{H}_{21}\text{Cl}_2\text{NO}_5$

	%C	%H	%N
Calc.	57.55	4.83	3.20
Obs.	57.32	5.05	3.08

Example 7

4-(2-Bromophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester



A) 4-(2-Bromophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A solution of 15 g (0,05 mol) of methyl 2-bromobenzylideneacetoacetate and 9,81 g (0,05 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 55 ml of ethanol, was refluxed in the absence of light for 10 hours. The mixture was then cooled to -10°C , and a yellow solid was obtained (mp $170-2^{\circ}\text{C}$, EtOH) with a yield of 85%.

B) 4-(2-Bromophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A mixture of 10 g (0,022 mol) of 4-(2-bromophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester, obtained according A), and 80 g (0,067 mol) of pyridinium chlorochromate absorbed on alumina, was suspended on 220 ml of CH_2Cl_2 , and the mixture was stirred at room temperature for 6 hours.


The remaining solid was eliminated by filtration, and the liquid was washed with water (3x150 ml), dried over anh. Na_2SO_4 , and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 8:2) with a yield of 66%.

Analysis calculated for $\text{C}_{21}\text{H}_{22}\text{BrNO}_5$

	%C	%H	%N
Calc.	56.26	4.95	3.12
Obs.	56.21	5.17	3.08

Example 8

4-(3-Bromophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl 5-tetrahydrofuran-2-ylmethyl ester



A) 4-(3-Bromophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3,5-di(tetrahydrofuran-2-ylmethyl) ester

A solution of 15 g (0.05 mol) of methyl 3-bromobenzylideneacetoacetate and 9,81g (0,05 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 55 ml of abs ethanol, was refluxed with stirring, in the absence of light, for 10 hours. The reaction mixture was then cooled to -10°C , yielding a white solid (mp $138-40^{\circ}\text{C}$, EtOH) with a yield of 85%.

B) 4-(3-Bromophenyl)-2,6-dimethylpyridine-3,5-dicarboxylic acid 3,5-di(tetrahydrofuran-2-ylmethyl) ester

A mixture of 10 g (0,022 mol) de 4-(3-bromophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3,5-di(tetrahydrofuran-2-ylmethyl) ester obtained as described in A), and 80 g (0,067 mol) of pyridinium chlorochromate adsorbed on alumina, was suspended in 220 ml of CH_2Cl_2 , and the suspension was stirred at room temperature for 5 hours. The insoluble solid was eliminated by filtration, and the liquid was washed with water (3x200 ml), dried over anh. Na_2SO_4 , and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 8:2) with a yield of 74%.


Analysis calculated for $\text{C}_{21}\text{H}_{22}\text{BrNO}_5$

	%C	%H	%N
Calc.	56.26	4.95	3.12
Obs.	56.62	4.88	3.38

Example 9

2,6-Dimethyl-4-(2-trifluoromethylphenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A) 2,6-Dimethyl-4-(2-trifluoromethylphenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester



A solution of 90 ml (118,8 g; 0,68 mol) of 2-trifluoromethylbenzaldehyde, 75 ml (79,23 g; 0,68 mol) of methyl acetoacetate and 126,38 g (0,68 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 700 ml of 2-propanol, was refluxed in the absence of light along 10 hours.

Then, 400 ml of solvent were distilled off at reduced pressure, and the remaining solution was cooled at -10°C . In that way, a yellow solid was obtained (mp $147-9^{\circ}\text{C}$, MeOH) with a yield of 69%.

B) 2,6-Dimethyl-4-(2-trifluoromethylphenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A mixture of 7 g (0.016 mol) of 2,6-dimethyl-4-(2-trifluoromethylphenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester, obtained according A),


and 57,35 g (0,048 mol) of pyridinium chlorochromate adsorbed on alumina, were suspended on 150 ml of CH_2Cl_2 , then the whole mixture was maintained with stirring at room temperature for 8 hours. The remaining solid was eliminated by filtration, and the liquid was washed with water (3x150 ml), dried over anh. Na_2SO_4 , and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 8:2) with a yield of 69%.

Analysis calculated for $\text{C}_{22}\text{H}_{22}\text{F}_3\text{NO}_5$

	%C	%H	%N
Calc.	60.41	5.07	3.20
Obs.	60.11	5,31	3.29

Example 10

2,6-Dimethyl-4-(3-trifluoromethylphenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester



A) 2,6-Dimethyl-4-(3-trifluoromethylphenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A solution of 13,21 g (0,05 mol) of methyl 3-trifluoromethylbenzylideneacetoacetate and 8,99 g (0,05 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 50 ml of abs ethanol, was refluxed under stirring, and protected from light, for 10 hours. The solvent was then evaporated under reduced pressure, and the residue was crystallised in EtAcO (15 ml). The product was obtained as a white solid (mp 108-10°C, EtOH), with a yield of 88%.

B) 2,6-Dimethyl-4-(3-trifluoromethylphenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester


A mixture of 12 g (0,027 mol) of 2,6-dimethyl-4-(3-trifluoromethylphenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester, obtained as described in A), and 98g (0,082 mol) of pyridinium chlorochromate adsorbed on alumina, was suspended in 270 ml of CH₂Cl₂, and the suspension was maintained under stirring at room temperature for 6 hours. The remaining solid was eliminated by filtration, and the liquid was washed with water (3x300 ml), dried over anh. Na₂SO₄, and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 8:2) with a yield of 65%.

Analysis calculated for C₂₂H₂₂F₃NO₅

	%C	%H	%N
Calc.	60.41	5.07	3.20
Obs.	60.02	5.01	3.51

Example 11

4-(3-Cyanophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester



A) 4-(3-Cyanophenyl)-2,6-dimethyl 1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A solution of 30g (0.13 mol) of methyl 3-cyanobenzylideneacetoacetate and 24.24 g (0.13 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 130 ml of abs. ethanol was refluxed with stirring, in the absence of light, along 8 hours. The reaction mixture was then cooled to -10°C , and the product was obtained as a yellow solid (mp $127-9^{\circ}\text{C}$, EtOH) with a yield of 79%.

B) 4-(3-Cyanophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester


A mixture of 10 g (0.025 mol) of 4-(3-cyanophenyl)-2,6-dimethyl 1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester, obtained as described in A), and 91 g (0.076 mol) of pyridinium chlorochromate adsorbed on alumina, was suspended on 250 ml of CH_2Cl_2 , and the suspension was stirred at room temperature for 5 hours. The remaining solid was separated by filtration, and the organic solution was washed with water (3x150 ml), dried with anh. Na_2SO_4 , and concentrated to dryness at reduced pressure. The product was obtained as an oil, which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 8:2) with a yield of 78%.

Analysis calculated for $\text{C}_{22}\text{H}_{22}\text{N}_2\text{O}_5$

	%C	%H	%N
Calc.	66.99	5.62	7.10
Obs.	66.84	5.92	7.14

Example 12

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester




To a solution of 30 g (0.07 mol) of 2,6-dimethyl-4-(2-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester in 72 ml of acetic acid, heated to reflux, 7.2 g (0.07 mol) of CrO_3 were portionwise added. After the addition was complete, reflux was maintained for one additional hour. The reaction mixture was then slowly poured on ice-cooled aq NH_4OH (150 ml). The mixture separated in two phases, and it was extracted with CH_2Cl_2 (3x100 ml). The organic extract was dried with anhydrous Na_2SO_4 , and concentrated to dryness at reduced pressure. The resulting oil was crystallised in EtAcO-diethylether, yielding a white solid (mp 69-70°C, EtOH) with a yield of 79%.

Analysis calculated for $\text{C}_{21}\text{H}_{22}\text{N}_2\text{O}_7$

	%C	%H	%N
Calc.	60.86	5.35	6.76
Obs.	60.63	5.26	6.52

Example 13

(*S*)-2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester




Over a stirred solution of 1,183 g (2,84 mmol) of (*R/S*)-2,6-Dimethyl-4-(2-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(*S*-tetrahydrofuran-2-ylmethyl) ester in 4 ml of acetic acid, 588 mg (8,53 mmol) of sodium nitrite was portionwise added, at room temperature. Once the addition completed, the reaction mixture was maintained with stirring at 50°C, during 90 min. Then, the reaction mixture was poured over 20 ml of ice/water, and then it was neutralised (to pH 5) with 20% aq NaOH, and then (to pH 7) with NaHCO₃. The solution was then extracted with CH₂Cl₂ (2x25ml), and the organic extract was washed with a saturated solution of NaCl, dried over anh. Na₂SO₄, and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 1:1). In that way, 879 mg of the product were obtained (yield:74%). The hydrobromide was prepared by treating an ethanolic solution of the product with 48% HBr, obtaining a white solid (121-4°C, EtOH).

Mass Spectra:

M/e: 384(4); 368(20); 331(19); 315(25); 284(25); 270(15); 267(27); 253(34); 236(42); 235(25); 209(25); 180 (20); 153(28); 152(38); 140(33); 139(38); 128(35); 127(43); 126(30); 114(33); 102(23); 101(25); 85 (156); 84(543); 77(33); 76(26); 71(1000); 67(35); 55(38).

Example 14

(*R*)-2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester




Over a solution of 0.89 g (2.13 mmol) of (*R/S*)-2,6-dimethyl-4-(2-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(*R*-tetrahydrofuran-2-ylmethyl) ester on 4 ml of acetic acid, stirred and heated to 60°C, 442 mg (6.41 mmol) of sodium nitrite was portionwise added. Once the addition was finished, stirring and heating were maintained for 90 min. more. Then, the reaction mixture was poured over 20 ml of ice/water, and then it was neutralised (to pH 5) with 20% aq NaOH, and then (to pH 7) with NaHCO₃. The solution was then extracted with CH₂Cl₂ (2x25ml), and the organic extract was washed with a saturated solution of NaCl, dried over anh. Na₂SO₄, and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 6:4). In that way, 334 mg of the product were obtained (yield: 38%). The hydrobromide was prepared by treating an ethanolic solution of the product with 48% HBr, obtaining a white solid (114-6°C, EtOH).

Mass Spectra:

M/e: 383(3); 368(16); 331(13); 315(20); 284(21); 270(14); 253(27); 236(34); 235(20); 209(20); 180(19); 153(25); 152(33); 140(30); 139(33); 128(33); 127(37); 126(27); 102(22); 101(23); 85(149); 84(514); 77(29); 76 (27); 71 (1000); 55(46).

Example 15

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(5-oxotetrahydrofuran-2-ylmethyl) ester



A) 2,6-Dimethyl-4-(2-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(5-oxotetrahydrofuran-2-ylmethyl) ester

A solution of 10 g (0,03 mol) of (5-oxotetrahydrofuran-2-ylmethyl) 2-nitrobenzylideneacetoacetate and 3.5 g (0,03 mol) of methyl 3-aminocrotonate in 30 ml of abs. ethanol, was refluxed with stirring in the absence of light, along 10 hours. The solvent was then distilled off at reduced pressure, and the obtained residue was purified by column chromatography (Silicagel, toluene:EtAcO, 1:1). In that way, the product was isolated as an oil, which crystallised in diisopropylether (5 ml), yielding a yellow solid (mp 86-8°C, diisopropyl ether) with a yield of 68%.

B) 2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(5-oxotetrahydrofuran-2-ylmethyl) ester


Over a boiling solution of 15 g (0,035 mol) of 2,6-dimethyl-4-(2-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(5-oxotetrahydrofuran-2-ylmethyl) ester, obtained according A), in 35 ml of acetic acid, 3,5 g (0,035 mol) of CrO₃ was portionwise added with stirring. Once the addition completed, reflux and stirring were maintained for one additional hour. Then, the reaction mixture was poured over 375 ml of conc. NH₄OH with ice, and the mixture was left overnight at 5°C. In that way, a solid was obtained (mp 132-4°C, MeOH) with a yield of 69%.

Analysis calculated for C₂₁H₂₀N₂O₈

	%C	%H	%N
Calc.	58.88	4.71	6.54
Obs.	59.04	4.84	6.59

Example 16

2,6-Dimethyl-4-(3-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester




A mixture of 15 g (0.036 mol) of 2,6-dimethyl-4-(3-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester and 130 g (0.108 mol) of pyridinium chlorochromate adsorbed on alumina, was suspended on 360 ml of CH_2Cl_2 , and maintained with stirring at room temperature for 8 hours. The remaining solid was eliminated by filtration and the organic solution was washed with water (3x300 ml), dried over anh. Na_2SO_4 , and concentrated to dryness at reduced pressure. The product was obtained as an oil which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 8:2) with a yield of 72%.

Analysis calculated for $\text{C}_{21}\text{H}_{22}\text{N}_2\text{O}_7$

	%C	%H	%N
Calc.	60.86	5.35	6.76
Obs.	60.58	5.45	7.01

Example 17

2,6-Dimethyl-4-(3-nitrophenyl)-pyridine-3,5-dicarboxylic acid 3,5-di(tetrahydrofuran-2-ylmethyl) ester (hydrochloride)



A) 2,6-Dimethyl-4-(3-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3,5-di(tetrahydrofuran-2-ylmethyl) ester

A solution of 15 g (0.05 mol) of tetrahydrofuran-2-ylmethyl 3-nitrobenzylideneacetoacetate and 8.7 g (0.05 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 50 ml of abs ethanol, was refluxed with stirring, in the absence of light, along 8 hours. The reaction mixture was then cooled to -10°C . A yellow solid (mp $122-3^{\circ}\text{C}$, EtAcO) was obtained, with a yield of 80%.

B) 2,6-Dimethyl-4-(3-nitrophenyl)-pyridine-3,5-dicarboxylic acid 3,5-di(tetrahydrofuran-2-ylmethyl) ester (hydrochloride)

A mixture of 8 g (0.016 mol) of 2,6-Dimethyl-4-(3-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3,5-di(tetrahydrofuran-2-ylmethyl) ester, obtained as described in A), and 60 g (0.049 mol) of pyridinium chlorochromate adsorbed on alumina, was suspended in 160 ml of CH_2Cl_2 , and the mixture was maintained at room temp. with stirring for 6 hours. The solid was separated by filtration and the liquid was washed with water (3x200 ml), dried (anh. Na_2SO_4), and concentrated at reduced pressure. The residue, obtained as an oil, was converted into hydrochloride by dissolving it into ethanol (10 ml) and then, adding HCl saturated ethyl ether (20 ml). The product precipitated as a white solid (mp $75-77^{\circ}\text{C}$, EtOH-ether) with a yield of 67%.


Analysis calculated for $\text{C}_{25}\text{H}_{28}\text{N}_2\text{O}_8\cdot\text{HCl}$

	%C	%H	%N
Calc.	57.64	5.61	5.38
Obs.	57.75	5.65	5.27

Example 18

4-(4-Chloro-3-nitrophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A) 4-(4-Chloro-3-nitrophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester



A solution of 20 g (0.07 mol) of methyl 4-chloro-3-nitrobenzylideneacetoacetate and 13.06 g (0.07 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 70 ml of abs ethanol, was refluxed with stirring, in the absence of light, for 8 hours. The solution was then concentrated to dryness at reduced pressure, and the residue was crystallised in EtAcO. A yellow solid was obtained (mp 148-50°C, EtOH) with a yield of 85%.

B) 4-(4-Chloro-3-nitrophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester


A mixture of 9 g (0.020 mol) of 4-(4-chloro-3-nitrophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester, obtained as described in A) and 72 g (0.060 mol) of pyridinium chlorochromate adsorbed on alumina, were suspended on 200 ml of CH₂Cl₂ then, the mixture was maintained with stirring at room temperature for 9 hours. The remaining solid was filtered, and the filtrate was washed with water (3x200 ml), dried with anh. Na₂SO₄, and the solvent evaporated under reduced pressure. The product was isolated as an oil, which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 8:2) with a yield of 69%.

Analysis calculated for C₂₁H₂₁ClN₂O₇

	%C	%H	%N
Calc.	56.19	4.72	6.24
Obs.	56.17	4.78	6.59

Example 19

4-(3-Acetylaminophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester (hydrochloride)



A) 4-(3-Acetylaminophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A solution of 30 g (0.115 mol) of methyl 3-acetylaminobenzylideneacetoacetate and 21.27 g (0.115 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 120 ml of isopropanol was heated to reflux with stirring along 8 hours, protected from the light. The solvent was evaporated under vacuum, and the residue was crystallised in ethyl acetate. A yellow solid was obtained, (MP 158-60°C, meow) with a yield of 76%.

B) 4-(3-Acetylaminophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester (hydrochloride).


A mixture of 12 g (0.03 mol) de 4-(3-Acetylaminophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester, as indicated in A), and 4.87 g (0.056 mol) of MnO₂ suspended in 60 ml of methylisobutylketone was refluxed with stirring for 24 hours. The remaining solid was immediately filtered and the solution concentrated at low pressure. The residue obtained was dissolved in 250 ml of 5% aq HCl, the solution was extracted with EtAcO (3x10 ml) and then, pH of the aqueous layer was adjusted to 8 with 20% K₂CO₃. The aqueous phase then, was extracted with CH₂Cl₂ (3x100 ml), the organic extracts were dried over anh. Na₂SO₄, concentrated at reduced pressure, and a yellow oil was obtained, which was transformed in hydrochloride by dissolving it in ethanol (10 ml) and adding HCl saturated diethyl ether (20 ml). In that way, a white solid was obtained (mp 196-8°C, MeOH) with a yield of 59%.

Analysis calculated for C₂₃H₂₆N₂O₆.HCl

	%C	%H	%N
Calc.	59.67	5.88	6.05
Obs.	59.56	6.12	5.92

Example 20

2,6-Dimethyl-4-(3-hydroxyphenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester (hydrochloride)



A) 2,6-Dimethyl-4-(3-hydroxyphenyl) 1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

A solution of 15g (0.07 mol) of methyl 3-hydroxybenzylideneacetoacetate and 12.62 g (0.07 mol) of (tetrahydrofuran-2-ylmethyl)-3-aminocrotonate in 70 ml of abs. ethanol, was refluxed with stirring for 8 hours in the absence of light. The solvent was evaporated at reduced pressure, and the residue was crystallised in EtAcO. In that way, a yellow solid was obtained (mp 170-2°C, EtAcO) with a yield of 82%.

B) 2,6-Dimethyl-4-(3-hydroxyphenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester (hydrochloride)


A mixture of 10 g (0.026 mol) of 2,6-Dimethyl-4-(3-hydroxyphenyl) 1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester, obtained as indicated in A), and 2.6 g (0.030 mol) of MnO₂ in 50 ml of methylisobutylketone, was refluxed with stirring for 14 hours. The remaining solid was eliminated by filtration, and the liquid was concentrated to dryness under reduced pressure. The residue obtained was dissolved in 15 ml of 15% aq. HCl, then, the solution was washed with ethyl acetate (10 ml), the pH was adjusted to 8 with 20% aq. K₂CO₃. Finally, the solution was extracted with CH₂Cl₂ (2x50 ml), the organic extracts were dried over anhydrous Na₂SO₄, and concentrated to dryness at reduced pressure. The residue appeared as an oil, which crystallised in 2 ml of methanol, and was transformed into the hydrochloride by dissolution in ethanol (10 ml) and addition of HCl saturated diethyl ether (20 ml). A yellow solid was obtained (mp 95-7°C, MeOH) with a yield of 64%.

Analysis calculated for C₂₁H₂₃NO₆.HCl

	%C	%H	%N
Calc.	59.79	5.73	3.32
Obs.	59.64	5.66	3.23

Example 21

2,6-Dimethyl-4-(2,3-dimethoxyphenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester



A mixture of 11 g (0,025 mol) of 2,6-dimethyl-4-(2,3-dimethoxyphenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester and 92 g (0,076 mol) of pyridinium chlorochromate absorbed on alumina, was suspended on 250 ml of CH_2Cl_2 , and the whole mixture was maintained on stirring at room temperature for 6 hours. The remaining solid was eliminated by filtration, and the liquid was washed with water (3x300 ml), dried over anh. Na_2SO_4 , and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, n-hexane:EtAcO, 8:2) with a yield of 79%.


Analysis calculated for $\text{C}_{23}\text{H}_{27}\text{NO}_7$

	%C	%H	%N
Calc.	64.32	6.34	3.26
Obs.	64.34	6.36	3.28


The compounds described in the examples 22 to 30 have been obtained as previously described in the literature.

Example 22


2,6-Dimethyl-4-(3-nitrophenyl)-pyridine-3,5-dicarboxylic acid 3-isopropyl ester 5-(2-methoxyethyl) ester

Example 23

2,6-Dimethyl-4-(2-nitrophenyl)-pyridine-3,5-dicarboxylic acid 3-methyl ester 5-isobutyl ester




Example 24

2,6-Dimethyl-4-(3-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-ethyl ester 5-methyl ester




Example 25

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid dimethyl ester


Example 264-[2-(2-*tert*-Butoxycarbonylvinyl)-phenyl]-2,6-dimethyl pyridine-3,5-dicarboxylic acid diethyl esterExample 272,6-Dimethyl-4-(3-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(*N*-methylbenzylamino)ethyl ester

Example 28


4-(2,3-Dichlorophenyl)-2,6-dimethyl pyridine-3,5-dicarboxylic acid 3-ethyl ester
5-methyl ester

Example 29

2-(2-Aminoethoxy)methyl-4-(2-chlorophenyl)-6-methylpyridine-3,5-dicarboxylic acid 3-ethyl ester 5-methyl ester


Example 30

2,6-Dimethyl-4-(3-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-(2-methoxyethyl) ester 5-(3-phenyl)propen-2-yl ester



Example 31

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester



A) 2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3,5-dimethyl ester

Over a boiling solution of 3 kg (8.7 mol) of 2,6-dimethyl-4-(2-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylic acid 3,5-dimethyl ester in 8,25 l of acetic acid, 870 g (8.7 mol) of CrO_3 were portionwise added, with stirring. When the addition was complete, the reaction mixture was refluxed with stirring for one additional hour. When the mixture reached the room temperature, it was slowly poured, with stirring, over 22.5 l of a mixture of 25% aq. H_4NOH and ice (2:1), and the product precipitated. The solid was isolated by filtration, washed with water until neutral pH, and crystallised (mp 99-101°C, MeOH) with a yield of 83%.

B) 2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester


A solution of 500 g of 2,6-dimethyl-4-(2-nitrophenyl)pyridine-3,5-dicarboxylic acid 3,5-dimethyl ester (1.5 mol) and 87.5 g of KOH (1.32 mol) in 5 l of methanol and 1.5 l of water, was refluxed with stirring for 4 hours. Then, the solvent was eliminated at reduced pressure and the remaining residue was triturated with 1 l of water. The solid (mostly unreacted starting material) was eliminated by filtration, and the filtrate was treated, at 0°C, slowly and with stirring, with 100 ml of 35% aq. HCl, and the mixture was maintained with stirring for one additional hour with stirring, at room temperature. A solid was formed, which was washed with water (250 ml) and ethanol (250 ml), yielding a yellowish solid (mp 232-3°C, ethanol) with a yield of 70%. A small percentage (12%) of the diacid corresponding to example 32, was isolated from the mother liquids.

Analysis calculated for $\text{C}_{16}\text{H}_{14}\text{N}_2\text{O}_6$


	%C	%H	%N
Calc.	58.18	4.27	8.48
Obs.	58.30	4.49	8.62

Example 32

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid

Example 33

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-(2-dimethylaminoethyl) ester 5-methyl ester (dihydrobromide)



A) Synthesis of 5-chlorocarbonyl-2,6-dimethyl-4-(2-nitro-phenyl)pyridine-3-carboxylic acid methyl ester

Over a mixture of 2,5 g (7,5 mmol) of 2,6-dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester and 0,7 g (3,8 mmol) of cyanuryl chloride, in 20 ml of anh. acetone under argon atmosphere, 1.1 ml of triethylamine were added. The resulting mixture was maintained with stirring at room temperature along 7 hours. Then, the mixture was concentrated to dryness at reduced pressure, and the residue was purified by column chromatography (Silicagel, n-hexane:EtAcO, 1:1) with a yield of 80%.

B) 2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-(2-dimethylaminoethyl) ester 5-methyl ester (dihydrobromide)

Over a mixture of 0,169 (1,8 mmol) of N-dimethylaminoethanol and 0,20 ml of triethylamine in 8 ml of CH₂Cl₂, cooled to 0°C and in argon atmosphere, 0,5 g (1,5 mmol) of 5-chlorocarbonyl-2,6-dimethyl-4-(2-nitro-phenyl)pyridine-3-carboxylic acid methyl ester described in A), dissolved in 4 ml of CH₂Cl₂ were added. The reaction mixture was stirred overnight at room temperature. Then, the mixture was washed with water, dried over anh. Na₂SO₄, and


concentrated to dryness at reduced pressure. The product was obtained as an oil, which was converted in dihydrobromide by treatment with EtOH (10 ml) and conc. HBr (1 ml). A white solid was obtained (85-8°C, EtOH) with a yield of 73%.

Analysis calculated for $C_{20}H_{23}N_3O_6 \cdot HBr$

	%C	%H	%N
Calc.	49.89	5.03	8.73
Obs.	49.62	5.47	9.02

Example 34

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-(2-diethylaminoethyl) ester 5-methyl ester (dihydrochloride)




Over a solution of 0,37 g (3,1 mmol) of N-diethylaminoethanol and 0,29 g (2,86 mmol) of triethylamine in 10 ml of CH_2Cl_2 , cooled to 0°C, and under nitrogen atmosphere, 1 g (2,86 mmol) of 5-chlorocarbonyl-2,6-dimethyl-4-(2-nitro-phenyl)pyridine-3-carboxylic acid methyl ester, obtained as described in A) of the example 33, dissolved in 8 ml of CH_2Cl_2 , were added. The mixture was stirred at room temperature for 15 hours; then, it was washed with water (10 ml), dried over anh. Na_2SO_4 , and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, EtAcO) and converted into dihydrochloride by treatment with EtOH (10 ml) and conc. HCl (1 ml). A white solid was obtained (166-7°C, EtOH) with a yield of 71%.

Analysis calculated for $C_{22}H_{27}N_3O_6 \cdot 2HCl$

	%C	%H	%N
Calc.	52.68	5.83	8.38
Obs.	52.96	5.56	8.07

Example 35

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 5-(3-dimethylaminopropyl) 3-methyl ester (dihydrochloride).




Over a solution of 0,146 g (1,4 mmol) of 3-dimethylamino-1-propanol and 0,13 g (1,29 mmol) of triethylamine in 6 ml of CH_2Cl_2 , cooled to 0°C and under nitrogen atmosphere, 0,45 g (1,29 mmol) of 5-chlorocarbonyl-2,6-dimethyl-4-(2-nitro-phenyl)pyridine-3-carboxylic acid methyl ester, obtained as described in A) of the example 33, dissolved in 5 ml of CH_2Cl_2 , were added. The mixture was stirred at room temperature along 15 hours; then, it was washed with water (10 ml), dried over anhydrous Na_2SO_4 , and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, acetone) and converted into dihydrochloride by treatment with EtOH (10 ml) and conc. HCl (1 ml). A white solid was obtained ($164-6^\circ\text{C}$, EtOH) with a yield of 75 %.

Analysis calculated for $\text{C}_{21}\text{H}_{25}\text{N}_3\text{O}_6 \cdot 2\text{HCl}$

	%C	%H	%N
Calc.	51.73	5.59	8.62
Obs.	51.50	5.82	8.43

Example 36

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3,5-dicarboxylic acid 3-methyl ester 5-(N-morpholino)ethyl ester (dihydrochloride)




Over a solution of 0,21 g (1,57 mmol) of *N*-(2-hydroxyethyl)-4-morpholine and 0,15 g of triethylamine in 5 ml of CH₂Cl₂, cooled to 0°C and under argon atmosphere, 0,5 g (1,43 mmol) of 5-chlorocarbonyl-2,6-dimethyl-4-(2-nitro-phenyl)pyridine-3-carboxylic acid methyl ester, obtained as described in A) of the example 33, dissolved in 4 ml of CH₂Cl₂, were added. The reaction mixture was stirred at room temperature along 12 hours, ; then, it was washed with water (10 ml), dried over anh. Na₂SO₄, and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, EtAcO) and converted into dihydrochloride by treatment with EtOH (10 ml) and conc. HCl (1 ml). A white solid was obtained (mp 96-8°C, EtOH) with a yield of 51%.

Analysis calculated for C₂₂H₂₅N₃O₇.2HCl

	%C	%H	%N
Calc.	51.25	5.28	8.16
Obs.	51.48	5.11	8.00

Example 37

2,6-Dimethyl-4-(2-nitrophenyl)pyridine-3,5-dicarboxylic acid 3-methyl ester 5-[1,3-di(*N*-morpholine)isopropyl] ester trihydrochloride



A mixture of 0.34 g (0.86 mmol) of 5-chlorocarbonyl-2,6-dimethyl-4-(2-nitrophenyl)pyridine-3-carboxylic acid methyl ester, obtained as described in A) of the example 33, 0.22 g (0.95 mmol) of 1,3-dimorpholin-4-yl-propan-2-ol and 96 mg (0.86 mmol) of triethylamine, were stirred at room temperature for 12 hours. Then, the mixture was washed with water (10 ml), dried over anhydrous Na_2SO_4 , and concentrated at reduced pressure, to dryness. The product was obtained as an oil, which was purified by column chromatography (Silicagel, EtAcO) and converted into trihydrochloride by treatment with EtOH (10 ml) and conc. HCl (1 ml). A white solid was obtained (mp 187-92°C, EtOH) with a yield of 24%.

Analysis calculated for $\text{C}_{27}\text{H}_{34}\text{N}_4\text{O}_8 \cdot 3\text{HCl}$

	%C	%H	%N
Calc.	49.83	5.74	8.61
Obs.	49.48	5.61	8.95

PHARMACOLOGICAL AND TOXICOLOGICAL TESTS

The most active compounds of formula (I) of the present invention were submitted to pharmacological tests to put in evidence their *in vitro* and *in vivo* cardioprotective effects, as well as their hemodynamic properties and toxicity.

The methods used were the following:

In vitro cardioprotective effect

Isolated guinea-pig ventricular myocytes were obtained following the method described by Powell et al. (J. Physiol., 1980, 102: 131) as modified by Levi et al. (J. Physiol., 1990, 425:9P).

Cell hypercontracture and death was obtained either by incubation under N₂ atmosphere or by incubation with veratridine which induced important changes in the ionic homeostase of cardiomyocytes (Hiroko Hashizume et al., European Journal of Pharmacology, 1974, 271: 1-8).

Results expressed as percentage of cells protected from hypercontracture i.e. maintaining its rod-shaped morphology with reference to vehicle-treated cells are given in the tables I and II hereafter.

In order to confirm the cardioprotection, the activity of one of the most interesting compound (example 31 according to the invention) was tested on guinea-pig ventricular cardiomyocytes action potential (AP) in the presence of veratridine or in anoxic condition.

The action potential (AP) time courses of guinea-pig ventricular cardiomyocytes treated with veratridine or made anoxic by incubation under N₂ atmosphere were measured (patch clamp technique) in absence and presence of the compound of example 31 according to the invention. This compound does not influence significantly the normal action potential but it prevents (100-300nM) the prolongation of cardiomyocyte action potential duration induced by veratridine. In anoxic conditions, there is a shortening of cardiomyocyte action potential duration related mainly with an activation of K⁺-ATP dependent current. This effect is inhibited dose-dependently and strongly by the compound of example 31 according to the invention (100-300nM).

Table I: Percentage of cells protected from hypercontracture obtained by incubation under N₂ atmosphere:

Compounds (30nM) according to the examples:	% Protection
12	80
22	63
23	50
24	60
25	50
31	58

Table II: Percentage of cells protected from hypercontracture obtained by incubation with veratridine:

Compounds (1 μ M) according to the examples:	% Protection
12	57
22	37.5
23	41.3
24	37.5
25	34
31	64

In vivo cardioprotective effect

Cardiac death and arrhythmias were induced by the coronary occlusion and reperfusion method in anaesthetised rats according to the method of Seyle et al. (Angiology, 1960, 11, 398-407) modified by Krzeminski T. ("proceedings of 7th Freiburg Focus of Biomeasurement", Strasbourg, 16-19 Sept. 1991) Cardiovascular and Respiratory in vivo Studies, Biomesstechnik, Verlag March GmbH, Germany) and according to the LAMBETH conventions (Walker M.J. A. et al., Cardiovasc Re., 1988, 22, 447-455).

The products have been administered orally (one day and one hour before the experiment). Table III hereafter presents the results obtained. They are given in percentage of protection against the mortality and the arrhythmias (ventricular fibrillations).

Table III: Protective effect in vivo

COMPOUND (2x20mg/kg) ACCORDING TO EXAMPLE N°:	% PROTECTION AGAINST DEATH	% PROTECTION AGAINST VENTRICULAR FIBRILLATIONS
12	100	100
22	100	50
23	100	75
24	100	37.5
25	100	50
31	85.7	73.4

Hemodynamic effect

The most active compounds according to the invention have been tested in order to analyse their hemodynamic activity.

In anaesthetised rat the intravenous administration of different dosages (30, 100, 300 and 1000 mg/kg) of the mentioned compounds did not influence significantly the following parameters: systolic, diastolic and mean arterial blood pressure, heart rate, ejection time, left ventricular systolic pressure, left ventricular end diastolic pressure, dP/dt_{max} and dP/dt_{min} , contractility index and relaxation time.

Toxicity


The most active compounds according to the invention present a low per oral toxicity with LD50 more than 2000mg/kg in the rat.

Conclusion

The pyridyl compounds of formula (I) according to the invention are able to protect the cardiomyocytes against damages induced by ionic modifications (responsible for most of cardiac pathology) and in such a way, they are able to protect the treated animals (rats) against death related to cardiac dysfunction and arrhythmia. Consequently, they are potentially cardioprotective agents in different human cardiac diseases (for example cardiomyopathies, myocardium infarction, angina, cardiac failures, coronary vasospasm, valvular heart disease, etc.). Moreover, they have a low per oral toxicity and in therapeutic dosages, they do not present a deleterious hemodynamic activity.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. Pyridyl compounds of general formula (I), in which



5 - X represents H, NO₂, OH, CN, F, Br, Cl, CF₃, -NH-CO-CH₃, 2,3-Cl₂, 2,3-(OCH₃)₂, 4-Cl, 3-NO₂ and 2-[CH=CH-COO-C(CH₃)₃], and the fused cycle 2,3-oxadiazole [2,3-(=N-O-N=)];

- Y represents an alcoxycarbonyl group R²OOC-, where R² can be H, a lower alkyl group, 2-méthoxyethyl, 2-tétrahydrofurfuryl or 5,5-diméthyl-1,3,2-dioxaphosphorinan-2-yl;

10

- R represents H, a lower alkyl group or, alternatively, the following groups: 2-methoxyethyl, 2-(benzyl-methylamino)ethyl, 2-tétrahydrofurfuryl, 5-oxotétrahydrofurfuryl, 3-[(4,4-diphenyl)-1-piperidiny]propyl, (R)-1-benzyl-3-piperidiny, 2-(N-benzyl-(N-phenylamino)ethyl, (S)-1-benzyl-3-pirrolidiny, 2-[(4-diphenylmethyl)-1-piperazinylethyl, 3-phényl-2-propényl, lower dialkyl-amino-lower alkyl, 2-(N-morpholino)ethyl or 1,3-di(N-morpholino)isopropyl;

15

- R¹ represents CH₃, CN or -CH₂-O-(CH₂)₂-NH₂, as well as the corresponding optical isomers, and the pharmaceutically acceptable salts of the said compounds.


20

2. The following compounds of formula (I), according claim 1

4-(2,3-Dichloro-phenyl)-2,6-diméthyl-pyridine-3, 5-dicarboxylic acid 3-méthyl ester 5-(tetrahydrofuran-2-ylméthyl) ester

25 2,6-Diméthyl-4-(3-fluorophényl) pyridine-3, 5-dicarboxylic acid 3-méthyl ester 5-(tetrahydrofuran-2-ylméthyl) ester

4-(3-Acetylaminophényl)-2,6-diméthyl pyridine-3, 5-dicarboxylic acid 3-méthyl ester 5-(tetrahydrofuran-2-ylméthyl) ester (hydrochloride)



2,6-Dimethyl-4-(3-hydroxyphenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester (hydrochloride)

2,6-Dimethyl-4-phenyl-pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(3-nitrophenyl)-pyridine-3, 5-dicarboxylic acid 3,5-di(tetrahydrofuran-2-ylmethyl) ester (hydrochloride)

4-(3-Cyanophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(3-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester

4-(4-Chloro-3-nitrophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester

4-(3-Bromophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3,5-di(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(3-trifluoromethylphenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester

4-(2-Chlorophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(2-trifluoromethylphenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester


2,6-Dimethyl-4-(2,3-dimethoxyphenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester

4-(3-Chlorophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(5-oxotetrahydrofuran-2-ylmethyl) ester

(S)-2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester

(R)-2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5--(tetrahydrofuran-2-ylmethyl) ester



4-(2-Bromophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(tetrahydrofuran-2-ylmethyl) ester

2,6-Dimethyl-4-(3-nitrophenyl)-pyridine-3, 5-dicarboxylic acid 3-isopropyl ester 5-(2-methoxyethyl) ester

2,6-Dimethyl-4-(2-nitrophenyl)-pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-isobutyl ester

2,6-Dimethyl-4-(3-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-ethyl ester 5-methyl ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid dimethyl ester

4-[2-(2-*tert*-Butoxycarbonylvinyl)-phenyl]-2,6-dimethyl pyridine-3, 5-dicarboxylic acid diethyl ester

2,6-Dimethyl-4-(3-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(*N*-methylbenzylamino)ethyl ester

4-(2,3-Dichlorophenyl)-2,6-dimethyl pyridine-3, 5-dicarboxylic acid 3-ethyl ester 5-methyl ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid

2-(2-Aminoethoxy)methyl-4-(2-chlorophenyl)-6-methyl pyridine-3, 5-dicarboxylic acid 3-ethyl ester 5-methyl ester

2,6-Dimethyl-4-(3-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-(2-methoxyethyl) ester 5-(3-phenyl)propen-2-yl ester

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-(2-dimethylaminoethyl) ester 5-methyl ester (dihydrobromide)

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-(2-diethylaminoethyl) ester 5-methyl ester (dihydrochloride)

2,6-Dimethyl-4-phenyl pyridine-3, 5-dicarboxylic acid dimethyl ester


2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-(3-dimethylaminopropyl) 5-methyl ester (dihydrochloride).

2,6-Dimethyl-4-(2-nitrophenyl) pyridine-3, 5-dicarboxylic acid 3-methyl ester 5-(*N*-morpholino)ethyl ester (dihydrochloride)



2,6-Dimethyl-4-(2-nitrophenyl)pyridine-3,5-dicarboxylic acid 3-methyl ester 5-[1,3-di(N-morpholine)isopropyl] ester trihydrochloride

3. Process for the preparation of compounds of formula (I) according claim 1, by oxydation of a 1,4-dihydropyridine of formula (II)



in which X, R and R' are as defined in claim 1.

4. Procedure, according the claim 3, in which a 1,4-dihydropyridine (II) is oxidised by an oxidant chosen between oxygen, $\text{NaNO}_2/\text{AcOH}$, NaNO_2/HCl , HNO_3 , $\text{Fe}(\text{NO}_3)_3$, $\text{Cu}(\text{NO}_3)_2$, $\text{Br}_2 + \text{NaOAc}$, CrO_3 , sulphur, KMnO_4 , chloranyl or o-chloranyl, 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ), Pd/C, pyridinium chlorochromate absorbed on alumina ($\text{PCC}/\text{Al}_2\text{O}_3$), pyridinium dichromate and MnO_2 .

5. Pyridyl compounds of formula (I), according the claim 1, for use as cardioprotective agents.

6. Pharmaceutical compositions containing at least, one pyridyl compound of formula (I), according claim 1, or one of its pharmaceutically acceptable salts.

7. A pharmaceutical composition, according claim 6, in which the quantity of compound of formula (I) is suitable to produce administered doses between 0,01 et 20 mg/kg.

8. A compound according to claim 1 substantially as hereinbefore described with reference to any of the examples.

DATED: 28 November, 2000

PHILLIPS ORMONDE & FITZPATRICK

Attorneys for:

CERMOL S.A.

