



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2017/06/12	(51) Cl.Int./Int.Cl. <i>G01N 33/50</i> (2006.01), <i>C07K 14/725</i> (2006.01)
(87) Date publication PCT/PCT Publication Date: 2017/12/14	
(85) Entrée phase nationale/National Entry: 2018/12/10	(71) Demandeurs/Applicants: UMC UTRECHT HOLDING B.V., NL; GADETA B.V., NL; ALBERT-LUDWIGS-UNIVERSITAT FREIBURG, DE
(86) N° demande PCT/PCT Application No.: EP 2017/064325	
(87) N° publication PCT/PCT Publication No.: 2017/212074	
(30) Priorités/Priorities: 2016/06/10 (EP16173986.7); 2016/06/10 (EP16173970.1)	(72) Inventeurs/Inventors: KUBALL, JURGEN HERBERT ERNST, NL; JANSSEN, ANKE, NL; BERINGER, DENNIS, NL; FISCH, PAUL, DE; VILLACORTA HIDALGO, JOSE ALBERTO, DE
	(74) Agent: GOWLING WLG (CANADA) LLP

- (54) Titre : NOUVEAU PROCEDE D'IDENTIFICATION DE CHAINES DE RECEPTEURS DE LYMPHOCYTES T DELTA (OU DE LYMPHOCYTES T GAMMA) OU DE PARTIES DE CELLES-CI QUI CONDITIONNENT UNE REPONSE ANTI-TUMORALE OU UNE REPONSE ANTI-INFECTIEUS
- (54) Title: NOVEL METHOD FOR IDENTIFYING DELTAT-CELL (OR GAMMAT-CELL) RECEPTOR CHAINS OR PARTS THEREOF THAT MEDIATE AN ANTI-TUMOUR OR AN ANTI-INFECTIVE RESPONSE

(57) **Abrégé/Abstract:**

The present invention relates to a method for identifying δ T-cell (or YT-cell) receptors chains or parts thereof that mediate an anti-tumor or anti-infection response by identifying amino acid sequences comprising δ T-cells(or YT-cell) receptors chains or parts thereof that are shared between different donors.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2017/212074 A1

(43) International Publication Date
14 December 2017 (14.12.2017)

(51) International Patent Classification:

G01N 33/50 (2006.01) **C07K 14/725** (2006.01)

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/EP2017/064325

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(22) International Filing Date:

12 June 2017 (12.06.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

16173970.1	10 June 2016 (10.06.2016)	EP
16173986.7	10 June 2016 (10.06.2016)	EP

(71) Applicants: **UMC UTRECHT HOLDING B.V.** [NL/NL]; Yalelaan 40, 3584 CM Utrecht (NL). **GADETA B.V.** [NL/NL]; Yalelaan 62, 3584 CM Utrecht (NL). **ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG** [DE/DE]; Friedrichstrasse 39, 79098 Freiburg (DE).

(72) Inventors: **KUBALL, Jürgen Herbert Ernst**; Jacob Catsstraat 55, 1215 EX Hilversum (NL). **JANSSEN, Anke**; van Eysingalaan 393, 3527 VK Utrecht (NL). **BERINGER, Dennis**; Naberlaan 20, 3571 ZJ Utrecht (NL). **FISCH, Paul**; Sudermannstrasse 36, 79114 Freiburg im Breisgau (DE). **VILLACORTA HIDALGO, Jose Alberto**; Laerchenstrasse 2, 79194 Gundelfingen (DE).

(74) Agent: **NEDERLANDSCH OCTROOIBUREAU**; P.O. Box 29720, 2502 LS The Hague (NL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: NOVEL METHOD FOR IDENTIFYING DELTAT-CELL (OR GAMMAT-CELL) RECEPTOR CHAINS OR PARTS THEREOF THAT MEDIATE AN ANTI-TUMOUR OR AN ANTI-INFECTIVE RESPONSE

(57) Abstract: The present invention relates to a method for identifying δ T-cell (or γ T-cell) receptors chains or parts thereof that mediate an anti-tumor or anti-infection response by identifying amino acid sequences comprising δ T-cells(or γ T-cell) receptors chains or parts thereof that are shared between different donors.

WO 2017/212074 A1

Novel method for identifying δ T-cell (or γ T-cell) receptor chains or parts thereof that mediate an anti-tumour or an anti-infective response

Field of the invention

5 The present invention relates to a method for identifying δ T-cell (or γ T-cell) receptors chains or parts thereof that mediate an anti-tumour or an anti-infection response by identifying amino acid sequences comprising δ T-cell (or γ T-cell) receptors chains or parts thereof that are shared between different donors.

Background of the invention

10 Our immune system utilizes different lines of defenses to protect us from infections as well as cancer. In order to cover the magnitude of potential invaders and internal threats our adaptive immune system has the possibility to raise up to 10^{16} $\alpha\beta$ TCR combinations as well as 10^{11} variations in immunoglobulins (1). Although threats are frequently identical and many HLA types are shared between individuals, only little
15 overlap in the usage of defined CDR3 regions in the α and β TCR chains between individuals has been reported so far. "Private specificities", thus α or β TCRs with a very distinct sequences, which are only observed in one individual, are usually dominant (2). Occasionally also public specificities are observed. These consist of highly shared β TCR sequences and are only observed in the presence of the very same
20 peptide MHC complex in the context of chronic infections such as EBV, CMV, influenza, or allo-reactive immune responses (3, 4). Analyses of the diversity of immunoglobulin responses imply a similar diversity and random usage of CDR3 sequences. Though an increased correlation in antibody gene segment usages, junctional features, and mutation rates are observed in twins, antibody pools show little
25 similarity in clonal responses to acute stimuli against the very same antigen (5). These data suggest that protective immune responses are usually not shared in detail in sequences between different individuals in the highly diverse $\alpha\beta$ TCR and immunoglobulin repertoire. Consequently each individual needs to raise its very own army of α as well as β TCR chains and immunoglobulins. Analyses of sequences only
30 will therefore not allow the identification of sequences with large therapeutic interest. This is also reflected by the fact that protective α as well as β TCR chains (6) and immunoglobulins, which are currently used in the clinic are usually identified by functional analyses followed by sequence analysis and not vice versa.

Among all immune receptor chains, TCR δ s have even the highest potential diversity in the CDR3 loop (approximately 10^{16} combinations for murine TCR δ) owing to the presence of multiple D gene segments (two in mice, three in human, and up to five in cattle) that can join together. Each D gene segment can be read in all three open reading frames, and N nucleotides can be inserted into the junctions of the joining segments. Thus, despite the limited diversity at the VJ junctions of TCR γ chains, the potential diversity generated at the combined CDR3 junctions (approximately 10^{18} combinations) is still higher than that of $\alpha\beta$ TCRs ($\sim 10^{16}$) and immunoglobulins ($\sim 10^{11}$). (1) This suggests also a highly diverse usage of CDR3 regions for δ TCR chains in each individual and is again in line with mentioned considerations for the usage of $\alpha\beta$ TCR chains or immunoglobulins, that sequence analysis only will not result in the identification of therapeutically relevant δ TCR chains.

Surprisingly we were able to identify shared δ T-cell (or γ T-cell) receptor chains or parts thereof that mediate an anti-tumour or an anti-infection response.

Description of the figures

Figure 1. Correction of unique clones for overlapping unique molecular identifiers (UMI's). Ratio between percentage of overlapping unique clones versus percentage of overlapping UMI's based on three healthy donors (HD 18 19 20) is shown.

Figure 2. Frequency of overlapping clones when corrected for UMI's and size of repertoire. Ratio between percentage of overlapping unique clones versus percentage of overlapping UMI's divided by the average repertoire size is shown on a logarithmic scale. Data are based on three healthy donors (HD18 19 20)

Figure 3. Comparison of NGS sequencing results (amino acids) of the δ TCR chains from 4 different donors and the 7th "virtual" donor as indicated in a Venn diagram. Healthy donor 18, 19, 20, the "combined donor" (HD 11, 12, 15), and the 7th virtual donor are depicted.

Figure 4. The $\gamma\delta$ TCR FE11 critically depends on the CD8 co-receptor for tumour recognition. (A) $\gamma\delta$ T cell clone FE11 was generated by limiting dilution. To assess tumour reactivity, FE11 cells were incubated with Daudi, SW480 or EBV-LCL tumour targets in the presence of control antibody or antibodies blocking CD8 α or

CD8 β . IFN γ secretion was measured by ELISPOT. Healthy PBMCs served as negative control targets. (B) The TCR γ and δ chains of clone FE11 were sequenced and retrovirally transduced into $\alpha\beta$ T cells (left panel; $\gamma\delta$ TCR expression on mock-transduced (light curve) and $\gamma\delta$ TCR-transduced T cells is indicated). Transfer of

5 $\gamma\delta$ TCR-mediated tumour-reactivity was tested by co-incubating $\gamma\delta$ TCR- or mock-transduced T cells with indicated target cells in an IFN γ ELISPOT (right panel). (C) CD4 $^{+}$ and CD8 $^{+}$ $\alpha\beta$ T cells transduced with the FE11 $\gamma\delta$ TCR were sorted and co-cultured with indicated target cells. T cell activation was assessed by IFN γ ELISPOT. (D) CD4 $^{+}$ and CD8 $^{+}$ $\alpha\beta$ T cells expressing the FE11 $\gamma\delta$ T

10 CR were co-incubated with SW480 target cells as in (C) but now in the presence of a control antibody or blocking antibodies against CD8 α or CD8 β . Data are representative of three (A, D), two (B), and five (C) separate experiments. Error bars represent S.E.M. (*P < 0.01; ***P < 0.001).

Figure 5. Overview of shared TCR δ and TCR β sequences. (A) Overview of shared

15 TCR δ sequences found within healthy donors (dataset 1). The included sequences in the healthy donors all had a clonal frequency >0.1%. On the horizontal axis the number of shared sequences between donors. On the vertical axis the percentage of shared sequences of the total of sequences of those two donors. (B) Overview of shared TCR β sequences found within healthy donors. On the horizontal axis the number of shared

20 sequences between donors. On the vertical axis the percentage of shared sequences of the total of sequences of those two donors. (C) Overview of shared V δ 2 sequences between the different datasets. On the horizontal axis the number of shared sequences between donors. (D) Overview of shared V δ 1 sequences between the different datasets. On the horizontal axis the number of shared sequences between donors. (E) Overview

25 of shared V δ 3 sequences between the different datasets. On the horizontal axis the number of shared sequences between donor.

Figure 6. V δ gene distribution (A) The majority of the identified TCR δ sequences by NGS in the peripheral blood are of V δ 2 $\gamma\delta$ T cells. The frequencies and median of the V δ gene distribution in 14 healthy donors are shown.

30 **Figure 7. Activated $\gamma\delta$ T cells are in contact with the apoptotic tumors positive cells.** (A) Double Immunohistochemistry using TCR γ antibody and cleaved caspase 3 (Areas with double positive T cells are indicated with a circle). T in these images

indicates tumor tissue **(B)** Analyses of $\gamma\delta$ TILs in TNBC froze section with CD69 as activation marker. Arrow show positive cells, nuclei are stained with DAPI blue.

Figure 8. Shared TCR δ and TCR γ sequences from $\gamma\delta$ TIL. Overview of shared **(A)** TCR δ and **(B)** TCR γ sequences found in tumor samples of patients with triple
5 negative breast cancer. On the right, the total identified sequences and the unique sequences per patients are shown. On the left, the overlap between patients is represented by the numbers in the table.

Figure 9. Improved anti-tumor activity of TEG001 when compared to a bulk population of V γ 9V δ 2 T cells. A panel of tumor cell lines **(A)** or primary AML tumor
10 samples **(B)** was incubated with TEG001 or with a bulk population of primary $\gamma\delta$ T cells with or without 10 μ M pamidronate (PAM) for 20h and IFN γ secretion was measured by ELISPOT. IFN γ spots per 15.000 T cells is shown as mean of triplicates (+SD). Statistical significances were calculated by two-way anova; *p<0.05; **p<0.01; ***p<0.001.

Figure 10. $\gamma\delta$ T cell clones and TEGs with shared γ or δ TCR chains.

(A)V δ 1 clones isolated from the peripheral repertoire and TEGs engineered with shared δ TCR chains show reactivity against different tumor cell lines. V δ 2^{neg} $\gamma\delta$ T cell clones of a healthy donor were tested against a broad panel of tumor cell lines by an ELISPOT assay. The reactivity of Fe11 in TEG (TEG-011) format was measured by
20 ELISA. ^ Indicates shared Vdelta1 chain with D13 and D19.

(B) TEG's engineered with shared γ or δ TCR chains isolated from $\gamma\delta$ TILs show reactivity against tumor cell lines. Reactivity towards tumor cell lines of $\gamma\delta$ TIL TCRs in TEG format as measured was measured by ELISPOT. \$ Indicates shared V γ 4 sequence published by Lafarge et al. (51). % Indicates shared V δ 5 sequence within $\gamma\delta$
25 TILs and also published by Lafarge et al (51), & Indicates γ TCR chains sequence published by Uldirch et al. (32). ! Indicates shared V γ 8 sequence. § Indicates shared V γ 2 sequence. ¶ Indicates shared V γ 4 sequence θ Indicates shared V δ 1 within $\gamma\delta$ TILs.

Figure 11. Inhibition of tumor growth and increased overall survival in TEG001 treated tumor bearing mice. NSG mice were treated with TEG001 (n=17) or Mock cells
30 (TEG-LM1, n=7) at day 7 and day 14 after tumor engraftment. (A) Bioluminescence Imaging was used to monitor tumor growth every 7 days. Data represent mean of all mice per group (+/-SEM). (B) Overall survival of treated mice was monitored until the

end of the experiment and is presented in the right panel. Statistical significances were calculated by log-rank (Mantel-Cox) test; ****p<0.0001.

Figure 12. Tetramer staining of TEG-AU2.3 (CD1d-specific) and TEG-B9.

5

Description of the invention

Method

In a first aspect the invention provides a method for identifying δ T-cell receptors chains or parts thereof that mediate an anti-tumour or anti-infection response comprising the steps of:

- a) providing amino acid sequences obtained from a donor, comprising δ T-cell receptor chains or parts thereof each of said receptor chains or parts thereof comprising a CDR3 region;
- b) identifying amino acid sequences comprising δ T- cell receptors chains or parts thereof obtained in step a) that are shared between different donors;
- c) confirming the anti-tumour or anti-infection response of the δ T-cell receptors chains or parts thereof identified in step b) by assessing the anti-tumour or anti-infection response of a T cell expressing a nucleic acid molecule encoding the amino acid sequence provided in step a).

20

In a second aspect the invention provides a method for identifying γ T-cell receptors chains or parts thereof that mediate an anti-tumour or anti-infection response comprising the steps of:

- a) providing amino acid sequences obtained from a donor, comprising γ T-cell receptor chains or parts thereof each of said receptor chains or parts thereof comprising a CDR3 region;
- b) identifying amino acid sequences comprising γ T- cell receptors chains or parts thereof obtained in step a) that are shared between different donors;
- c) confirming the anti-tumour or anti-infection response of the γ T-cell receptors chains or parts thereof identified in step b) by assessing the anti-tumour or anti-infection response of a T cell expressing a nucleic acid molecule encoding the amino acid sequence provided in step a).

30

Unless otherwise indicated herein, the explanation provided for each feature of the method of the first aspect below also holds for each feature of the method of the second aspect; the only difference being that the method of the second aspect deals with the identification of γ T-cell receptor chains or parts thereof whereas the method of the first aspect deals with the identification of δ T-cell receptor chains or parts thereof.

Step a of the method of the first and second aspects

Step a) comprises the provision of the amino acid sequences from a donor, comprising δ T-cell (or γ T-cell for the method of the second aspect) receptor chains or parts thereof comprising a CDR3 region. In a first place, T cells or T lymphocytes should be first isolated.

T cells, or T lymphocytes, belong to a group of white blood cells named lymphocytes, which play a role in cell-mediated immunity. T cells originate from hematopoietic stem cells in the bone marrow, mature in the thymus (that is where the T is derived from), and gain their full function in peripheral lymphoid tissues. During T-cell development, $CD4^-CD8^-$ T-cells (negative for both the CD4 and CD8 co-receptor) are committed either to an $\alpha\beta$ or $\gamma\delta$ fate as a result of an initial β or δ TCR gene rearrangement. Cells that undergo early β chain rearrangement express a pre-TCR structure composed of a complete β chain and a pre-TCR α chain on the cell surface. Such cells switch to a $CD4^+CD8^+$ state, rearrange the TCR α chain locus, and express a mature $\alpha\beta$ TCR on the surface. $CD4^-CD8^-$ T cells that successfully complete the γ gene rearrangement before the β gene rearrangement express a functional $\gamma\delta$ TCR and remain $CD4^-CD8^-$. (Claudio Tripodo et al. Gamma delta T cell lymphomas Nature Reviews Clinical Oncology 6, 707-717 (December 2009)). The T cell receptor associates with the CD3 protein complex. Mature T cells, i.e. expressing a $\alpha\beta$ TCR or a $\gamma\delta$ TCR, express the T cell receptor complex on the cell surface. The $\gamma\delta$ T-cells, which constitute about 1 -5% of the total population of T cells, can be divided in further subpopulations. A subpopulation of $\gamma\delta$ T-cells constitutes $\gamma9\delta2$ T-cells, which express a $\gamma9\delta2$ TCR. Within the extracellular domain of a T cell receptor three complementarity determining regions (CDR1 , CDR2, CDR3) are located. These regions are in general the most variable domains and contribute significantly to the diversity among TCRs. CDR regions are composed during the development of a T-cell where so-called Variable-(V), Diverse-(D), and Joining-(J)-gene segments are randomly combined to generate diverse TCRs.

Of the three CDR regions CDR3, for both $\alpha\beta$ T-cells and $\gamma\delta$ T-cells, is the most variable one, and is therefore the key player in antigen/ligand recognition.

$\alpha\beta$ T cells may be defined with respect to function as T lymphocytes that express an $\alpha\beta$ TCR, which recognize peptides bound to MHC molecules (major histocompatibility complex), which are expressed on the surface of various cells. MHC molecules present peptides derived from the proteins of a cell. When for example a cell is infected with a virus, the MHC will present viral peptides, and the interaction between the $\alpha\beta$ TCR on the T cell and the MHC-complex on the target cell (i.e. the virus infected cell) activates specific types of T-cells which initiate and immune responses to eliminate the infected cell. Hence, $\alpha\beta$ T cells may be functionally defined as being cells capable of recognizing peptides bound to MHC molecules. $\alpha\beta$ T cells may be selected from peripheral blood for example via the CD3 antigen as described below and in the examples, as the large majority of T cells have the $\alpha\beta$ TCR. $\alpha\beta$ T cells may also be selected with an antibody specific for the $\alpha\beta$ TCR, such as described below. From such selected cells, the nucleic acid (or amino acid) sequence corresponding to the α T-cell receptor chain and the β T-cell receptor chain may be determined by sequencing, preferably as carried out in the experimental part. Hence, $\alpha\beta$ T-cells may also be defined as being cells comprising a nucleic acid (or amino acid) sequence corresponding to the α T-cell receptor chain and/or the β T-cell receptor chain.

$\gamma\delta$ T-cells may be functionally defined in that they are specifically and rapidly activated by e.g. a set of non-peptidic phosphorylated isoprenoid precursors, collectively named phosphoantigens or stress signals mediated by non classical HLA molecules like CD1. Phosphoantigens are produced by virtually all living cells, though the levels are usually very low in healthy cells, and increased in transformed / malignant cells or cells infected with e.g. mycobacterium tuberculosis, which deliver a derivate of phosphoantigens . The most common phosphoantigen found in human cells (including cancer cells) is isopentenyl pyrophosphate (IPP) and its isomer dimethylallyl pyrophosphate (DMAPP). Activation of $\gamma\delta$ T-cells comprises clonal expansion, cytotoxic activity and expression and release of cytokines. $\gamma\delta$ T-cells are also defined by expression of the $\gamma\delta$ T cell receptor. For example, cells may be selected using an antibody specific for the $\gamma\delta$ T cell receptor such as described below. From such selected

cells, the nucleic acid (or amino acid sequence) sequence corresponding to the γ T-cell receptor chain and/or the δ T-cell receptor chain may be determined by sequencing, preferably as carried out in the experimental part. Hence, $\gamma\delta$ T-cells may also be defined as being cells naturally comprising a nucleic acid (or amino acid) sequence
5 corresponding to a γ T-cell receptor chain and/or a δ T-cell receptor chain.

The person skilled in the art is well capable of selecting and/or identifying cell populations characterized by expression of an antigen or receptor on the surface of the cell such as described throughout herein. It is understood that with regard to expression
10 on the surface of cells, such as CD3, CD4, CD8, $\alpha\beta$ TCR, and $\gamma\delta$ TCR, this is typically done in a population of cells of which a portion of cells have a much higher level of expression of the antigen when compared to cells having a lower level of expression.

Hence, the terms positive or negative are to be understood as being relative, i.e.
15 positive cells have a much higher expression level as compared to cells being negative. Cells being negative in this sense may thus still have an expression level which may be detected.

Expression on the surface of cells may be analyzed using Fluorescence Activated Cell
20 Sorting (FACS), and many specific antibodies are commercially available, e.g. such as for CD3, CD4, CD8, $\alpha\beta$ TCR, and $\gamma\delta$ TCR that are suitable for such FACS analysis, such as described in the examples and as available. $\gamma\delta$ TCR may be $\gamma9\delta2$ TCR. As an example, $\alpha\beta$ T cells can also be defined and selected as being positive for $\alpha\beta$ TCR in FACS. The same holds for $\gamma\delta$ T cells. Antibodies suitable for FACS or similar
25 separation techniques (such as e.g. antibodies conjugated to magnetic beads) are widely available. Conditions are selected, such as provided by the antibody manufacturer that allows the selection of negative and/or positive cells. Examples of antibodies that may be suitable for selection of $\gamma\delta$ -T cells, or engineered $\gamma\delta$ T cells or $\gamma9\delta2$ T cells or engineered $\gamma9\delta2$ T cells such as available from BD Pharmingen (BD, 1 Becton Drive,
30 Franklin Lakes, NJ USA) are V γ 9-PE (clone B3, #555733), V δ 2-FITC (clone B6, #555738), $\gamma\delta$ TCR-APC (clone B1, #555718) or such as available from Beckman Coulter is pan- $\gamma\delta$ TCR-PE (clone IMM510, # IM 1418U) Similarly, suitable

antibodies for $\alpha\beta$ -T cell selection, such as anti-CD3 antibodies may be such as available from BD Pharmingen is CD3-FITC (#345763) or such as anti- $\alpha\beta$ TCR antibodies such as available from Beckman Coulter is pan- $\alpha\beta$ TCR-PE (#A39499) or pan- $\alpha\beta$ TCR-PC5 (#A39500). An alternative antibody that binds to the human endogenous $\alpha\beta$ T cell receptor is available commercially from Miltenyi (Miltenyi Biotec GmbH, Friedrich-Ebert-Straße 68, 51429 Bergisch Gladbach, Germany). This antibody is from cell clone BW242/412 which is of the mouse isotype IgG2b. A FITC labeled BW242/412 antibody is available from Miltenyi under order no. 130-098-688. The BW242/412 cell clone and the antibody expressed by BW242/412 is described in detail i.a. EP0403156B1 which is herein incorporated by reference.

Accordingly, in the method of the invention, first T-cells are provided. The T-cells may be primary cells, for example from a subject, such as described in the examples for a human subject. The T-cells may be $\alpha\beta$ or $\gamma\delta$ T-cells derived from a human subject. Alternatively, the T-cells may be T cell lines, such as SupT-1, Jurkat, or Raji cells or any other widely available cell line. Any cell type, being a primary cell or any other cell line will suffice, as long as the cell population, or a substantial part thereof, expresses the T-cell receptor, i.e. such as being positive for the $\alpha\beta$ T- cell or the $\gamma\delta$ TCR receptor in a FACS sorting or the like as described above, such a cell population may be contemplated. Also, any cell or cell population may be contemplated that, when provided with a $\gamma\delta$ TCR according to the invention is capable of forming a functional TCR complex and exerting e.g. a functional cytotoxic response and/or cytokine production. The cell that is provided may also be a progenitor cell, preferably a blood progenitor cell such as a thymocyte or a blood stem cell, which after it has been provided with the right stimuli can develop into T cells.

Preferably, T cells provided express or are able to express a $\gamma\delta$ TCR. T cells may have been transduced to express a $\gamma\delta$ TCR or already express a γ TCR and have been transduced to express a δ TCR (or respectively already express a δ TCR and have been transduced to express a γ TCR), comprising the nucleic acid sequences encoding the sequence obtained in step a). All theoretical combinations of a γ with a δ chain of the TCR is encompassed. In an embodiment, the $\gamma\delta$ TCR is $\gamma9\delta2$ TCR. In another embodiment the $\gamma\delta$ TCR is $\gamma5\delta1$ TCR. In another embodiment, the $\gamma\delta$ TCR is $\gamma8\delta5$ TCR. In another embodiment, the $\gamma\delta$ TCR is $\gamma4\delta5$ TCR. In another embodiment, the

$\gamma\delta$ TCR is $\gamma 2\delta 1$ TCR. In another embodiment the $\gamma\delta$ TCR is $\gamma 8\delta 1$ TCR.

Step a) comprising “providing amino acid sequences obtained from a donor, comprising δ T-cell receptor chains or parts thereof each of said receptor chains or parts thereof comprising a CDR3 region” may be replaced by “obtaining from a plurality of donors, δ T-cell receptor chains, γ T-cell receptor chains or parts thereof, said receptor chains or parts thereof comprising a CDR3 region, and determining amino acid sequences of said receptors chains or parts thereof or nucleic acid sequences encoding the same”.

10

In step a) at least two different donors are used. This is important in the method of the invention as one wishes to identify δ TCR (or γ TCR) chains or part thereof that are shared by different donors. A preferred δ TCR chain is a $\delta 2$ TCR chain. Another preferred δ TCR chain is a $\delta 1$ TCR chain. Another preferred δ TCR chain is a $\delta 3$ TCR chain. Another preferred δ TCR chain is a $\delta 4$ TCR chain. Another preferred δ TCR chain is a $\delta 5$ TCR chain. A preferred γ TCR chain is a $\gamma 2$ TCR chain. Another preferred γ TCR chain is a $\gamma 4$ TCR chain. Another preferred γ TCR chain is a $\gamma 5$ TCR chain. Another preferred γ TCR chain is a $\gamma 8$ TCR chain. Another preferred γ TCR chain is a $\gamma 9$ TCR chain.

20 In an embodiment, a $\gamma\delta$ TCR is a $\gamma 9\delta 2$ TCR. In another embodiment the $\gamma\delta$ TCR is $\gamma 5\delta 1$ TCR. In another embodiment the $\gamma\delta$ TCR is $\gamma 9\delta 4$ TCR. In another embodiment the $\gamma\delta$ TCR is $\gamma 4\delta 5$ TCR. In another embodiment the $\gamma\delta$ TCR is $\gamma 8\delta 5$ TCR. In another embodiment the $\gamma\delta$ TCR is $\gamma 2\delta 1$ TCR. In another embodiment the $\gamma\delta$ TCR is $\gamma 8\delta 1$ TCR. In another embodiment the $\gamma\delta$ TCR is $\gamma 4\delta 3$ TCR.

25 The number of different donors used may be as high as possible. At least 2 different donors are used. However it is preferred that 3, 4, 5, 6, 7, 8, 9, 10 or more different donors are used.

In an embodiment, at least one of the donors may be healthy. However, it is also encompassed that all donors are healthy.

30 In another embodiment, at least one of the donors may be diseased. However, it is also encompassed that all donors are diseased or elite controller of a disease. It means that a donor could be nor healthy nor diseased but got a cancer or an infection and was able to

control it. The TCR's of such a "controlled" infection are of particular interest.

A disease in this context may be cancer or any infection. Infection include infections immediate by viruses bacteria and fungi such as the Hepatitis virus, the Herpes Viruses (CMV, EBV, and more) or a Mycobacterium.

5

In another embodiment, it is also envisaged to have part of the donors being healthy and the remaining part of the donors being diseased.

A donor is preferably a human being.

Accordingly in a preferred embodiment of the step a) of the method of the invention:

10

- a. the donors are human beings,
- b. at least 2 different donors are used,
- c. at least one donor, preferably all donors are healthy and/or
- d. at least one donor is diseased.

Step b of the method of the first and second aspects

15

Step b) comprises the identification of amino acid sequences comprising δ T-cells (or γ T-cells) receptors chains or parts thereof obtained in step a) that are shared between different donors.

Such amino acid sequences are preferably defined by InMunoGeneTics information system (<http://www.imgt.org/IMGTScientificChart/Nomenclature/IMGT->

20

FRCDRdefinition.html).

It is to be understood that the expression "δT-cell receptors chains or parts thereof that are shared between different donors" means that it is not per se the whole "δT-cell receptors chains" that are shared or are identical between different donors. Part of such chain may be shared or may be identical between different donors. In theory each part of a δT-cell receptor may be shared between donors. The same holds for "γT-cell receptors chains or parts thereof that are shared between different donors".

25

It is also encompassed that the parts that are shared may not be identical but comprises conservative substitutions of a given amino acid. A list of amino acids that are considered to be a conservative substitution of another amino acid is provided in the

30

general part of the description dedicated to the definitions under identity/similarity. In an embodiment said shared part is comprised within a CDR3 region of a δT-cell (or γT-cell) receptor chain or comprises a CDR3 region of a δT-cell (or γT-cell) receptor chain

or consists of a δ T-cell (or γ T-cell) receptor chain. In another embodiment, said shared part is from 3 to 53 amino acids or from 5 and 40 or from 10 and 30. More preferably said shared part is comprised within a CDR3 region of a δ T-cell (or γ T-cell) receptor chain and is from 3 to 53 amino acids.

- 5 It is clear that the number of donors sharing a δ T-cell (or γ T-cell) receptor chain or part thereof depends on the number of donors used in step a). In a preferred embodiment, the sequences identified are shared between at least 2, 3, 4, 5 different donors.

Step c of the method of the first and second aspects

- 10 Step c) comprises the confirmation or the validation of sequences identified as shared in step b). In order to validate the biological relevance of such sequence, the anti-tumour or anti-infection response of a T-cell expressing a defined nucleic acid molecule encoding an amino acid sequence as provided in step a) is determined. The T-cell may already express a δ T- cell (or γ T-cell) receptor chain identified as shared in
- 15 step b). It is clear that even if step b) has led to the identification of a part of a δ T-cell (or γ T-cell) receptor chain being shared, in order to assess the biological relevance of the corresponding δ T-cell (or γ T-cell) receptor chain, a T cell is transduced with the corresponding δ T-cell (or γ T-cell) receptor chain comprising the part identified in step b). In an embodiment, an anti-tumour or anti-infection response of such sequence is
- 20 assessed in a T-cell that does not endogenously express a gamma or delta chain of the TCR. Such a cell may be an $\alpha\beta$ -T cell or a NK cell.

The nucleic acid sequences encoding the δ T-, preferably the δ 2-T cell receptor chain may be introduced into T-cells to provide an engineered T-cell as explained in the general part of the description dedicated to the definitions.

- 25 Alternatively in the method of the second aspect, the nucleic acid sequences encoding the γ T-, preferably the γ 9-T cell receptor chain may be introduced into T-cells to provide an engineered T-cell as explained in the general part of the description dedicated to the definitions.

- 30 It is clear to a skilled person that the T cells used should also express a γ T cell receptor chain in order to assess the biological relevance of a δ T cell receptor chain. In other words a $\gamma\delta$ TCR is preferably expressed in said T cells, the δ TCR being the one

identified in the method of the invention as being shared between different donors. In the method of the second aspect, it is also clear to a skilled person that the T cells used should also express a δ T cell receptor chain in order to assess the biological relevance of a γ T cell receptor chain. In other words a $\gamma\delta$ TCR is preferably expressed in
5 said T cells, the γ TCR being the one identified in the method of the invention as being shared between different donors.

In a preferred embodiment, the nucleic acid molecule encoding the δ T-cell (or γ T-cell) receptor chain or part thereof is provided in an expression vector or in a retroviral or
10 lentiviral vector in a T cell. This has been extensively explained in the general part of the description dedicated to the definitions.

T cells may be expanded before or after the transfer of the nucleic acids encoding the δ T-and/or γ T- cell receptor chain. Preferably, the expansion is after the transfer such
15 that the amount of nucleic acids that needs to be transferred is as low as possible. This expansion of T cells may be performed by stimulation with anti-CD3/CD28 Dynabeads in the presence of IL-2. The expanded cells comprising the engineered $\gamma\delta$ T-cell receptor, may be selected e.g. via a selectable marker and may be further selected for the presence of the CD4 antigen and the CD8 antigen, e.g. using the MACS separating
20 system as described in the examples. The engineered T-cells may be subsequently further expanded using the REP protocol as described by Riddel and Greenberg, 1990 J Immunol Methods. 128(2): 189-201, which is incorporated herein by reference, or using similar further expansion methods thereto. Briefly, the expansion method involves using antibodies directed against T cell activation molecules, such as TCR,
25 CD3 and CD28 and/or feeder cells and/or stimulating cytokines.

The anti-tumour response of the provided T-cell expressing a δ T-cell (or γ T-cell) receptor chain may be assessed using any technique known to the skilled person. A δ T-cell receptor chain may be a δ 2-T cell receptor chain. A γ T-cell receptor chain may be a
30 γ 9-T cell receptor chain.

In one embodiment, the step of determining an anti-tumour response or reactivity comprises contacting the T cells with tumour cells. The step of determining anti-

tumour reactivity may include any assay in which an anti-tumour effect may be determined, such as having an effect on tumour cell division rate, i.e. the speed with which the tumour cells divide, cell death, binding to the tumour cells, induction of the production of a cytokine such as IFN γ , IL-2 or TNF α .

5 Tumour cells may be any kind of tumour cells. For example, primary tumour cells from a patient. The tumour cells may be tumour cells from cell lines, such as the cell lines listed in the examples named Daudi, RPMI8226/S, OPM2, LME1, K562, Saos2, MZ1851 RC, SCC9, Fadu, MDA-MB231, MCF7, BT549, SW480, which are well known in the art. Tumour cell lines may easily be obtained from the American Type
10 Culture Collection (ATCC, Manassas, Virginia) and the like.

In a preferred embodiment, determining the anti-tumour responses includes contacting the T cell expressing a defined nucleic acid molecule encoding an amino acid comprising a δ T-cell (or γ T-cell) receptor chain identified as shared in step b) with a tumour cell and measuring its ability to lyse the tumour cell and/or induce the
15 production of a cytokine such as IFN- γ , IL-2 or TNF α . This contacting, culturing or incubation step may have a duration from 10 hours to 1, 2, 3, 4, 5 days. The ability to lyse the tumour cells include providing a fixed amount of tumour cells with which T cell expressing a defined nucleic acid molecule encoding an amino acid comprising a δ T-cell (or γ T-cell) receptor chain identified as shared in step b), is contacted and after
20 an incubation period the number of viable tumour cells is counted.

An anti-tumour response may have been identified or determined when the number of viable tumor cells at the end of the incubation step is less than 90%, less than 80%, less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20%, less than 10% of the number of initial tumour cells at the onset of the incubation step.
25 Alternatively, an anti-tumour response may have been identified or determined when the number of viable tumour cells at the end of the incubation step with the engineered T cells is lower than the number of tumour cells at the end of a similar incubation step with control T cells not engineered with sequences identified as shared. Lower in this context may mean at least 10% lower, at least 20% lower, at least 30% lower, at least
30 40% lower, at least 50% lower, at least 60% lower, at least 70% lower, at least 80% lower, at least 90% lower.

In addition or as alternative to the counting of the number of viable tumour cells at the

end of the incubation step, one may also perform a $^{51}\text{Chromium}$ -release assay which is known to the skilled person. The amount of $^{51}\text{Chromium}$ release is a measure of the number of cells that have been lysed.

- 5 Similarly, the production of a cytokine such as IFN- γ , IL-2 or TNF α or the secretion or the expression of activation markers may also be determined, e.g. via antibody staining, ELISA and/or quantitative PCR for the expressed mRNA. Assays for determining the production of a cytokine such as IFN- γ , IL-2 or TNF α are commercially widely available. When the production of a cytokine such as IL-2,
- 10 TNF α or IFN γ is detected at the end of the contacting step, the T cell expressing a defined nucleic acid molecule encoding an amino acid comprising a δT -cell (or γT -cell) receptor chain identified as shared in step b) is said to exhibit an anti-tumour response. Alternatively and preferably, when the amount of IFN γ , IL-2 or TNF α produced at the end of the contacting step with engineered T cells is higher (preferably
- 15 at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 100% or more) than the amount of IFN γ IL-2 or TNF α produced when tumour cells are contacted with control T cells, the T cells is said to exhibit an anti-tumour response. Control T cells do not express a defined nucleic acid molecule encoding an amino acid comprising a δT -cell (or γT -cell) receptor chain
- 20 identified as shared in step b).

- An anti-tumour response may also be determined by assessing the binding of the engineered T cells to the tumour cell at the end of the incubation step. When binding of the T cell expressing a defined nucleic acid molecule encoding an amino acid
- 25 comprising a δT -cell (or γT -cell) receptor chain identified as shared in step b) to the tumour cell is detected at the end of the contacting step, the T cell expressing a defined nucleic acid molecule encoding an amino acid comprising a δT -cell (or γT -cell) receptor chain identified as shared in step b) is said to exhibit an anti-tumour response. Alternatively and preferably, when the binding of the T cell expressing a defined
- 30 nucleic acid molecule encoding an amino acid comprising a δT -cell (or γT -cell) receptor chain identified as shared in step b) at the end of the contacting step is higher (preferably at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 100% or more) than the binding

of control T cells (see earlier definition) to the same tumour cell, the T cells is said to exhibit an anti-tumour response.

5 The contacting step may be carried out in the presence of a phosphoantigen, such as pamidronate.

In a preferred method the step of confirming the anti- tumour response comprises contacting the T cell expressing a defined nucleic acid molecule encoding an amino acid comprising a δ T-cell (or γ T-cell) receptor chain identified as shared in step b) with a tumour cell and measuring its ability to lyse the tumour cell and/or induce a cytokine such as IFN- γ , IL-2 or TNF α .

In a preferred embodiment, determining the anti-infection responses includes contacting the T cell expressing a defined nucleic acid molecule encoding an amino acid comprising a δ T-cell (or γ T-cell) receptor chain identified as shared in step b) with the infectious agent or cells comprising the infectious agent and measuring its ability to kill the infectious agent or cells comprising the infectious agent and/or induce the production of a cytokine such as IFN- γ , IL-2 or TNF α . This contacting, culturing or incubation step may have a duration from 10 hours to 1, 2, 3, 4, 5 days. The ability to kill the infectious agent or cells comprising the infectious agent includes providing a fixed amount of infectious agent or cells comprising the infectious agent or inducing indirectly an expression of a natural danger signal at the cell surface by the infectious agent with which the T-cell expressing a $\gamma\delta$ TCR as explained earlier herein (or a T cell expressing a defined nucleic acid molecule encoding an amino acid comprising a δ T-cell (or γ T-cell) receptor chain identified as shared in step b)), is contacted and after an incubation period the number of alive infectious agent or cells comprising the infectious agent is counted.

An anti-infectious response may have been identified or determined when the number of infectious agent or cells comprising the infectious agent at the end of the incubation step is less than 90%, less than 80%, less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20%, less than 10% of the number of initial infectious agent or cells comprising the infectious agent at the onset of the incubation step.

Alternatively, an anti-infectious response may have been identified or determined when

the number of viable infectious agent or cells comprising the infectious agent at the end of the incubation step with the T cells is lower than the number of infectious agent or cells comprising the infectious agent at the end of a similar incubation step with control T cells (earlier defined herein). Lower in this context may mean at least 10% lower, at least 20% lower, at least 30% lower, at least 40% lower, at least 50% lower, at least 60% lower, at least 70% lower, at least 80% lower, at least 90% lower.

In a preferred method the step of confirming the anti-infectious response comprises contacting the T cell expressing a defined nucleic acid molecule encoding an amino acid comprising a δ T-cell (or γ T-cell) receptor chain identified as shared in step b) with an infectious agent or cells comprising the infectious agent and measuring its ability to kill the infectious agent or cells comprising the infectious agent and/or induce a cytokine such as IFN- γ , IL-2 or TNF α .

15 δ T-cell receptor chain or a part thereof in general

Provided in certain aspects described herein are polypeptides comprising a δ T-cell receptor chain or a variant or functional portion thereof. In a further aspect the invention provides a δ T-cell receptor chain or a part thereof, comprising a CDR3 region, and which δ T-cell receptor chain or part thereof is represented by an amino acid sequence as defined in step 1 c) of the method of the invention and which is obtainable by the method of the first aspect. Each of these δ T-cell receptor chain or part thereof is represented by an amino acid sequence that could be identified using a SEQ ID NO. In an embodiment, a δ T-cell receptor chain is a δ 2T-cell receptor chain. In another embodiment, a δ T-cell receptor chain is a δ 1T-cell receptor chain or a δ 4T-cell receptor chain.

γ T-cell receptor chain or a part thereof in general

Provided in certain aspects described herein are polypeptides comprising a γ T-cell receptor chain or a variant or functional portion thereof. In a further aspect the invention provides a γ T-cell receptor chain or a part thereof, comprising a CDR3 region, and which γ T-cell receptor chain or part thereof is represented by an amino acid sequence as defined in step 1 c) of the method of the invention and which is obtainable

by the method of the first aspect. Each of these γ T-cell receptor chain or part thereof is represented by an amino acid sequence that could be identified using a SEQ ID NO. In an embodiment, a γ T-cell receptor chain is a γ 9T-cell receptor chain or a γ 5T-cell receptor chain.

5

Each of the δ T-cell receptor chain or part thereof comprising a CDR3 region identified herein may also be represented by its coding nucleic acid sequence instead of its amino acid sequence. Therefore the invention also relates to a nucleic acid molecule encoding said receptor chain or part thereof. The same holds for each of the γ T-cell receptor chain or part thereof comprising a CDR3 region identified herein. The same also holds for the TCR identified herein: it can be identified by the receptor chains it expresses or by the nucleic acid molecules encoding these chains it comprises. The same also holds for the T cell expressing said TCR: the T cell can be defined by reference to the receptor chains or parts thereof it expresses or by the nucleic acid molecules encoding these chains or parts thereof it comprises.

10
15

Preferred δ T-cell or γ T-cell receptor chain or a part thereof

Provided in certain aspects described herein are polypeptides comprising a δ T-cell receptor CDR or a variant or functional portion thereof.

20

In another preferred embodiment, there is provided a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 7 and/or 17 and/or 96.

25

In another preferred embodiment, there is provided a nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 50 and/or 62 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 50 and/or 62.

30

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,

98%, 99% or 100%.

Each of these preferred δ T-cell or γ T-cell receptor chains or parts thereof defined above by sequence identity and as encompassed by the invention are preferably considered to be able to exhibit an anti-tumour or anti-infective activity as assessed in step c) of the
5 method of the invention.

Provided in certain aspects described herein are polypeptides comprising a γ T-cell receptor CDR or a variant or functional portion thereof. There is also provided a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or
10 part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 10, 18, 64 and/or 97.

In another preferred embodiment, there is provided a nucleic acid molecule represented
15 by a nucleotide sequence comprising a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 51 and/or 63 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 51 and/or 63.

20 Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

In another preferred embodiment, there is provided a γ T-cell receptor chain or part
25 thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:18
Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

30 In another preferred embodiment, there is provided a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least

60% sequence identity or similarity with amino acid sequence SEQ ID NO: 19 and/or 30.

In another preferred embodiment, there is provided a nucleic acid molecule represented
5 by a nucleotide sequence comprising a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 41 and/or 53 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 41 and/or 53.

10 Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

In another preferred embodiment, there is provided a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being
15 represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 20 and/or 32.

In another preferred embodiment, there is provided a nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60%
20 sequence identity with SEQ ID NO: 43 and/or 55 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 43 and/or 55.

25 Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

In another preferred embodiment, there is provided a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least
30 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 21 and/or 34.

In another preferred embodiment, there is provided a nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60%

sequence identity with SEQ ID NO: 45 and/or 57 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 45 and/or 57.

- 5 Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

In another preferred embodiment, there is provided a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being
10 represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 40 and/or 70.

In another preferred embodiment, there is provided a nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60%
15 sequence identity with SEQ ID NO: 76 and/or 82 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 76 and/or 82.

- 20 Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

In another preferred embodiment, there is provided a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least
25 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 65 and/or 71.

In another preferred embodiment, there is provided a nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 77 and/or 83 and/or a nucleotide sequence that
30 encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 77 and/or 83.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,

98%, 99% or 100%.

In another preferred embodiment, there is provided a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being
5 represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 66 and/or 72.

In another preferred embodiment, there is provided a nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60%
10 sequence identity with SEQ ID NO: 78 and/or 84 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 78 and/or 84.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,
15 98%, 99% or 100%.

In another preferred embodiment, there is provided a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being
20 represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 67 and/or 73.

In another preferred embodiment, there is provided a nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60%
25 sequence identity with SEQ ID NO: 79 and/or 85 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 79 and/or 85.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,
30 98%, 99% or 100%.

In another preferred embodiment, there is provided a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being
represented by an amino acid sequence, said amino acid sequence comprising at least

60% sequence identity or similarity with amino acid sequence SEQ ID NO: 68 and/or 74.

In another preferred embodiment, there is provided a nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60%
5 sequence identity with SEQ ID NO: 80 and/or 86 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 80 and/or 86.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,
10 98%, 99% or 100%.

In another preferred embodiment, there is provided a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least
15 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 69 and/or 75.

In another preferred embodiment, there is provided a nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60%
20 sequence identity with SEQ ID NO: 81 and/or 87 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 81 and/or 87.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,
25 98%, 99% or 100%.

In another preferred embodiment, there is provided a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least
30 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 89 and/or 91.

In another preferred embodiment, there is provided a nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60%
sequence identity with SEQ ID NO: 93 and/or 95 and/or a nucleotide sequence that

encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 93 and/or 95.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,
5 98%, 99% or 100%.

Each of these preferred δ T-cell receptor chains or parts thereof defined above by sequence identity and as encompassed by the invention are preferably considered to be
10 able to exhibit an anti-tumour or anti-infective activity when expressed in a T cell already expressing a γ T-cell receptor as assessed in step c) of the method of the invention. The same holds for preferred γ T-cell receptor chains or parts thereof defined above by sequence identity, the T cell already expressing a δ T cell receptor.

15 Further aspects linked to the use of a δ T-cell (or γ T-cell) receptor chain or a part thereof

In this part, all aspects described below applied for any δ T-cell (or γ T-cell) receptor chain or variant or part thereof obtainable by the present invention, and especially for the preferred δ T cell (or γ T-cell) receptor chain or variant or part thereof identified
20 above.

In an embodiment, a variant or part of a δ T-cell (or γ T-cell) receptor chain described herein is a soluble polypeptide. Such a soluble polypeptide may also be called a binding unit. Such a soluble polypeptide can include various forms to binding entities such as a TCR, antibody, scFv, BCR, or any combination thereof. In some cases, at
25 least a portion of a TCR, such as a $V\gamma 9V\delta 2$ or $V\gamma 5V\delta 1$ or $V\gamma 4V\delta 5$ or $V\gamma 8V\delta 5$ or $V\gamma 2V\delta 1$ or $V\gamma 8V\delta 1$ can be generated and utilized in a pharmaceutical composition as described herein. For example, TCR-antibody chimeras can be generated and tested before arriving at a desired chimera. For example, $\gamma\delta$ -variable domains can replace heavy and light chain variable domains of an antibody. In addition to enhanced binding,
30 an Fc domain of an antibody can mediate cytotoxicity through Fc γ -receptor positive immune cells and/or a complementary system. In some cases, TCR-antibody chimeras can be generated using HEK293 cells and subsequently purified using protein A

affinity chromatography followed by size exclusion chromatography. A proper folding of chimeras can be probed using conformational-specific antibodies that can target γ and δ variable domains. Chimeras can be used in antibody dependent cell mediated cytotoxicity (ADCC) and complement dependent cytotoxicity (CDC) assays to
 5 determine functional efficacy. After performing *in vitro* assays, functional efficacy of TCR-antibody chimeras can be tested *in vitro* and/or *in vivo*.

In a further aspect, the invention also relates to a conjugate comprising (a part of) the δ T-cell (or γ T-cell) receptor chain as defined above which is linked to an agent. The
 10 type of agent used depends from the type of applications envisaged. Such conjugates may be linked to substrates (e.g. chemicals, nanoparticles) and may be used e.g. to deliver chemotherapy to a target of interest. In addition, in diagnostics expression of defined ligands may be tested by taking advantage of the soluble TCRs linked to fluorochromes which are then used as staining tool or for the biochemical isolation of
 15 the ligand.

In a further aspect, the invention relates to a nucleic acid construct comprising the δ T-cell (and/or γ T-cell) receptor chain or a part thereof represented by an amino acid sequence as identified in step c) of the method of the invention, preferably the δ T-cell
 20 (and/or γ T-cell) receptor chain or part thereof identified in the part entitled "Preferred δ T-cell (or γ T-cell) receptor chain or a part thereof".

In an aspect, there is therefore provided a $\gamma\delta$ TCR comprising:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell
 25 receptor chain or part thereof and
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof, each having been identified by a method as disclosed herein.

30 In another preferred embodiment, there is therefore provided a $\gamma\delta$ TCR comprising:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said

- amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 7 and/or 17 and/or 96, and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 10, 18 64 and/or 97.

In another preferred embodiment, there is provided a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 50 and/or 62 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 50 and/or 62, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 51 and/or 63 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 51 and/or 63.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

In another preferred embodiment, there is therefore provided a $\gamma\delta$ TCR comprising:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:22 and/or 35 , and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:39 and/or 36.

In another preferred embodiment, there is provided a nucleic acid molecule encoding a

$\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 46 and/or 58 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO:46 and/or 58, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 47 and/or 59 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 47 and/or 59.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

- 15 In another preferred embodiment, there is therefore provided a $\gamma\delta$ TCR comprising:
- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 23 and/or 29 and preferably
 - 20 - a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 19 and/or 30.

25 In another preferred embodiment, there is provided a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 26 and/or 52 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 26 and/or 52, and preferably
- 30 - a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO:

41 and/or 53 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 41 and/or 53.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,
5 98%, 99% or 100%.

In another preferred embodiment, there is therefore provided a $\gamma\delta$ TCR comprising:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said
10 amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 24 and/or 31, and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or
15 similarity with amino acid sequence SEQ ID NO: 20 and/or 32.

In another preferred embodiment, there is provided a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO:
20 42 and/or 54 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO:42 and/or 54, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO:
25 43 and/or 55 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 43 and/or 55.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,
30 98%, 99% or 100%.

In another preferred embodiment, there is therefore provided a $\gamma\delta$ TCR comprising:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell

receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 25 and/or 33 and preferably

- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 21 and/or 34

In another preferred embodiment, there is provided a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence

comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 44 and/or 56 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 44 and/or 56, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 45 and/or 57 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 45 and/or 57.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

In another preferred embodiment, there is therefore provided a $\gamma\delta$ TCR comprising:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:28 and/or 37, and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:27 and/or 38

In another preferred embodiment, there is provided a nucleic acid molecule encoding a

$\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 48 and/or 60 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 48 and/or 60, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 49 and/or 61 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 49 and/or 61.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

- 15 In another preferred embodiment, there is therefore provided a $\gamma\delta$ TCR comprising:
- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:40 and/or 70, and preferably
 - 20 - a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:65 and/or 71.

25 In another preferred embodiment, there is provided a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 76 and/or 82 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 76 and/or 82, and preferably
- 30 - a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO:

77 and/or 83 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 77 and/or 83.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,
5 98%, 99% or 100%.

In another preferred embodiment, there is therefore provided a $\gamma\delta$ TCR comprising:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said
10 amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:66 and/or 72, and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or
15 similarity with amino acid sequence SEQ ID NO:67 and/or 73.

In another preferred embodiment, there is provided a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO:
20 78 and/or 84 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 78 and/or 84, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO:
25 79 and/or 85 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 79 and/or 85.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,
30 98%, 99% or 100%.

In another preferred embodiment, there is therefore provided a $\gamma\delta$ TCR comprising:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell

receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:68 and/or 74, and preferably

- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:69 and/or 75.

In another preferred embodiment, there is provided a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence

comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 80 and/or 86 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO:80 and/or 86, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 81 and/or 87 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 81 and/or 87.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

In another preferred embodiment, there is therefore provided a $\gamma\delta$ TCR comprising:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:88 and/or 90, and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:89 and/or 91.

In another preferred embodiment, there is provided a nucleic acid molecule encoding a

$\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 92 and/or 94 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO:92 and/or 94, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 93 and/or 95 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 93 and/or 95.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

In a further aspect, the invention further relates to a vector comprising the nucleic acid construct defined above. A preferred vector is a retroviral vector or a lentiviral vector. In a further aspect, the invention provides a cell comprising the nucleic acid construct or the vector as defined above. This cell is preferably a T cell.

A preferred T cell expresses a δ T-cell (or γ T-cell) receptor chain or a part thereof identified in the part entitled “Preferred δ T-cell (or γ T-cell) receptor chain or a part thereof”.

The general part of the description dedicated to the definitions provides detailed explanation as to nucleic acid molecules and polypeptide encompassed by the invention, nucleic construct, viral vector and cells comprising said construct or vector.

Each of the δ T-cell (or γ T-cell) receptor chain or part thereof isolated using the method of the invention is expected to be biologically relevant for designing a medicament for preventing, treating, regressing, curing and/or delaying cancer or an infection since each of these chain or part thereof exhibits an anti-tumour or an anti-infectious activity (step c of the methods of the invention).

A δ T-cell (or γ T-cell) receptor chain, or a part thereof, a conjugate, a nucleic acid construct, a vector, a cell all as defined earlier herein are preferably for use as a

medicament. In an embodiment, a δ T-cell receptor chain is a δ 2T-cell, a δ 1T-cell, δ 5T-cell, or a δ 4T-cell receptor chain. In an embodiment, a γ T-cell receptor chain is a γ 9T-cell, γ 8T-cell, γ 4T-cell receptor chain or a γ 5T-cell receptor chain. The medicament is preferably for the prevention, suppression, treatment of cancer or an infection.

5 Accordingly the invention also relates to a composition, preferably a pharmaceutical composition comprising a δ T-cell (or γ T-cell) receptor chain, or a part thereof, a conjugate, a nucleic acid construct, a vector, a cell all as defined earlier herein.

In a further aspect, the invention relates to a method for preventing, treating, regressing, curing and/or delaying cancer or an infection in a subject wherein a δ T-cell (or γ T-cell)
10 receptor chain, or a part thereof, a conjugate, a nucleic acid construct, a vector, a cell all as defined earlier herein are administered to said subject. A preferred subject is a human being.

In a further aspect, the invention relates to a use of a δ T-cell (or γ T-cell) receptor chain, or a part thereof, a conjugate, a nucleic acid construct, a vector, a cell all as defined
15 earlier herein for the manufacture of a medicament for preventing, treating, regressing, curing and/or delaying cancer or an infection in a subject. A preferred subject is a human being.

A preferred T cell used as a medicament as explained is preferably a T cell expressing:

- 20
- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 7 and/or 17 and/or 96, and preferably
 - a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell
25 receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 10, 18, 64 and/or 97.

Another preferred T cell used as a medicament comprises a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide
30 sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 50 and/or 62 and/or a nucleotide sequence that encodes an amino acid sequence

that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 50 and/or 62, and preferably

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 51 and/or 63 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 51 and/or 63.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

Another preferred T cell used as a medicament as explained is preferably a T cell expressing:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:22 and/or 35, and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:39 and/or 36.

Another preferred T cell used as a medicament comprises a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 46 and/or 58 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 46 and/or 58, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 47 and/or 59 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid

sequence encoded by a nucleotide sequence SEQ ID NO: 47 and/or 59.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

5 Another preferred T cell used as a medicament as explained is preferably a T cell expressing:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity
10 with amino acid sequence SEQ ID NO: 23 and/or 29, and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 19 and/or 30.

15 Another preferred T cell used as a medicament comprises a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 26 and/or 52 and/or a nucleotide sequence that encodes an amino acid sequence
20 that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 26 and/or 52, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 41 and/or 53 and/or a nucleotide sequence that encodes an amino acid sequence
25 that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 41 and/or 53.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

30 Another preferred T cell used as a medicament as explained is preferably a T cell expressing:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell

receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 24 and/or 31, and preferably

- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 20 and/or 32.

Another preferred T cell used as a medicament comprises a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 42 and/or 54 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 42 and/or 54, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 43 and/or 55 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 43 and/or 55.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

Another preferred T cell used as a medicament as explained is preferably a T cell expressing:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 25 and/or 33 and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 21 and/or 34.

Another preferred T cell used as a medicament comprises a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 44 and/or 56 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 44 and/or 56, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 45 and/or 57 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 45 and/or 57.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

- Another preferred T cell used as a medicament as explained is preferably a T cell expressing:
- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:28 and/or 37 and preferably
 - a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:28 and/or 38.

Another preferred T cell used as a medicament comprises a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 48 and/or 60 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid

sequence encoded by a nucleotide sequence SEQ ID NO: 48 and/or 60, and preferably

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 49 and/or 61 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 49 and/or 61.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

Another preferred T cell used as a medicament as explained is preferably a T cell expressing:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:40 and/or 70 and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:65 and/or 71.

Another preferred T cell used as a medicament comprises a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 76 and/or 82 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 76 and/or 82, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 77 and/or 83 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 77 and/or 83.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%,

98%, 99% or 100%.

Another preferred T cell used as a medicament as explained is preferably a T cell expressing:

- 5 - a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:66 and/or 72 and preferably
- 10 - a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:67 and/or 73.

Another preferred T cell used as a medicament comprises a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide
15 sequence comprising:

- 20 - a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 78 and/or 84 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 78 and/or 84, and preferably
- 25 - a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 79 and/or 85 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 79 and/or 85.

25 Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

Another preferred T cell used as a medicament as explained is preferably a T cell expressing:

- 30 - a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity

with amino acid sequence SEQ ID NO:68 and/or 74 and preferably

- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or
5 similarity with amino acid sequence SEQ ID NO:69 and/or 75.

Another preferred T cell used as a medicament comprises a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO:
10 80 and/or 86 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 80 and/or 86, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO:
15 81 and/or 87 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 81 and/or 87.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

20

Another preferred T cell used as a medicament as explained is preferably a T cell expressing:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said
25 amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO:88 and/or 90 and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or
30 similarity with amino acid sequence SEQ ID NO:89 and/or 91.

Another preferred T cell used as a medicament comprises a nucleic acid molecule encoding a $\gamma\delta$ TCR, said nucleic acid molecule being represented by a nucleotide

sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 92 and/or 94 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 92 and/or 94, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 93 and/or 95 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 93 and/or 95.

Preferably, the identity is of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100%.

In the general part dedicated to definitions, more detailed information is provided as to the formulation of a pharmaceutical composition. The way the δ T-cell (or γ T-cell) receptor chain or part thereof or nucleic acid construct or viral vector or cells may be administered to a subject in a method of treatment or in a pharmaceutical use of each of these compounds has already been defined herein in the context of the method of the first aspect (see step c) and in the general part of the description dedicated to the definitions.

General part dedicated to definitions

Polypeptide/nucleic acid

In the context of the invention, a polypeptide is represented by an amino acid sequence. Preferred polypeptides are δ T-cell (or γ T-cell) receptor chains or parts thereof which mediates an anti-tumour response as explained herein.

In the context of the invention, a nucleic acid molecule as a nucleic acid molecule encoding such a δ T-cell (or γ T-cell) receptor chain or part thereof is represented by a nucleic acid or nucleotide sequence which encodes such a polypeptide. A nucleic acid molecule may comprise a regulatory region.

It is to be understood that each nucleic acid molecule or polypeptide or construct as identified herein by a given Sequence Identity Number (SEQ ID NO) is not limited to this specific sequence as disclosed. Throughout this application, each time one refers to

a specific nucleotide sequence SEQ ID NO (take SEQ ID NO: X as example) encoding a given polypeptide, one may replace it by:

- i. a nucleotide sequence comprising a nucleotide sequence that has at least 60% sequence identity or similarity with SEQ ID NO: X;
- 5 ii. a nucleotide sequences the complementary strand of which hybridizes to a nucleic acid molecule of sequence of (i);
- iii. a nucleotide sequence the sequence of which differs from the sequence of a nucleic acid molecule of (i) or (ii) due to the degeneracy of the genetic code; or,
- 10 iv. a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: X.

Throughout this application, each time one refers to a specific amino acid sequence SEQ ID NO (take SEQ ID NO: Y as example), one may replace it by: a polypeptide comprising an amino acid sequence that has at least 60% sequence identity or similarity
15 with amino acid sequence SEQ ID NO: Y.

Each nucleotide sequence or amino acid sequence described herein by virtue of its identity or similarity percentage (at least 60%) with a given nucleotide sequence or amino acid sequence respectively has in a further preferred embodiment an identity or a
20 similarity of at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or more identity or similarity with the given nucleotide or amino acid sequence respectively. In a preferred embodiment, sequence identity or similarity is determined by comparing the whole length of the sequences as identified herein. Unless otherwise indicated herein, identity or similarity with a given SEQ ID NO means identity or similarity based on the
25 full length of said sequence (*i.e.* over its whole length or as a whole).

Sequence identity

"Sequence identity" is herein defined as a relationship between two or more amino acid (polypeptide or protein) sequences or two or more nucleic acid
30 (polynucleotide) sequences, as determined by comparing the sequences. The identity between two amino acid or two nucleic acid sequences is preferably defined by assessing their identity within a whole SEQ ID NO as identified herein or part thereof. Part thereof may mean at least 50% of the length of the SEQ ID NO, or at least 60%, or

at least 70%, or at least 80%, or at least 90%.

In the art, "identity" also means the degree of sequence relatedness between amino acid or nucleic acid sequences, as the case may be, as determined by the match between strings of such sequences. "Similarity" between two amino acid sequences is
5 determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the sequence of a second polypeptide. "Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in (Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects,
10 Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part I, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heine, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., SIAM J. Applied Math., 48:1073
15 (1988).

Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include e.g. the GCG program
20 package (Devereux, J., et al., Nucleic Acids Research 12 (1): 387 (1984)), BestFit, BLASTP, BLASTN, and FASTA (Altschul, S. F. et al., J. Mol. Biol. 215:403-410 (1990). The BLAST X program is publicly available from NCBI and other sources (BLAST Manual, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., J. Mol. Biol. 215:403-410 (1990). The well-known Smith Waterman
25 algorithm may also be used to determine identity.

Preferred parameters for polypeptide sequence comparison include the following: Algorithm: Needleman and Wunsch, J. Mol. Biol. 48:443-453 (1970); Comparison matrix: BLOSSUM62 from Hentikoff and Hentikoff, Proc. Natl. Acad. Sci. USA. 89:10915-10919 (1992); Gap Penalty: 12; and Gap Length Penalty: 4. A program
30 useful with these parameters is publicly available as the "Ogap" program from Genetics Computer Group, located in Madison, WI. The aforementioned parameters are the default parameters for amino acid comparisons (along with no penalty for end gaps).

Preferred parameters for nucleic acid comparison include the following:

Algorithm: Needleman and Wunsch, J. Mol. Biol. 48:443-453 (1970); Comparison matrix: matches=+10, mismatch=0; Gap Penalty: 50; Gap Length Penalty: 3. Available as the Gap program from Genetics Computer Group, located in Madison, Wis. Given above are the default parameters for nucleic acid comparisons.

5 Optionally, in determining the degree of amino acid similarity, the skilled person may also take into account so-called "conservative" amino acid substitutions, as will be clear to the skilled person. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and
10 isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulphur-containing side
15 chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine. Substitutional variants of the amino acid sequence disclosed herein are those in which at least one residue in the disclosed sequences has been removed and a different residue inserted in its place. Preferably, the amino acid
20 change is conservative. Preferred conservative substitutions for each of the naturally occurring amino acids are as follows: Ala to Ser; Arg to Lys; Asn to Gln or His; Asp to Glu; Cys to Ser or Ala; Gln to Asn; Glu to Asp; Gly to Pro; His to Asn or Gln; Ile to Leu or Val; Leu to Ile or Val; Lys to Arg, Gln or Glu; Met to Leu or Ile; Phe to Met, Leu or Tyr; Ser to Thr; Thr to Ser; Trp to Tyr; Tyr to Trp or Phe; and Val to Ile or Leu.

25 *Conjugate*

A polypeptide comprising a δ T-cell (or γ T-cell) receptor chain or part thereof which mediates an anti-tumour response as explained herein may be coupled or linked to an agent to form a conjugate. The agent may be selected from the group consisting of a diagnostic agent, a therapeutic agent, an anti-cancer agent, a chemical, a nanoparticle, a
30 chemotherapeutic agent or a fluorochrome.

Gene or coding sequence

"Gene" or "coding sequence" or "nucleic acid" or "nucleic" refers to a DNA or RNA

region (the transcribed region) which “encodes” a particular polypeptide such as a δ T-cell receptor. A coding sequence is transcribed (DNA) and translated (RNA) into a polypeptide when placed under the control of an appropriate regulatory region, such as a promoter. A gene may comprise several operably linked fragments, such as a promoter, a 5' leader sequence, an intron, a coding sequence and a 3' nontranslated sequence, comprising a polyadenylation site or a signal sequence. A chimeric or recombinant gene (such as the one encoding a δ TCR or γ TCR chain comprising the polypeptide as identified herein and operably linked to a promoter) is a gene not normally found in nature, such as a gene in which for example the promoter is not associated in nature with part or all of the transcribed DNA region. “Expression of a gene” refers to the process wherein a gene is transcribed into an RNA and/or translated into an active protein.

Promoter

As used herein, the term “promoter” refers to a nucleic acid fragment that functions to control the transcription of one or more genes (or coding sequence), located upstream with respect to the direction of transcription of the transcription initiation site of the gene, and is structurally identified by the presence of a binding site for DNA-dependent RNA polymerase, transcription initiation sites and any other DNA sequences, including, but not limited to transcription factor binding sites, repressor and activator protein binding sites, and any other sequences of nucleotides known to one of skill in the art to act directly or indirectly to regulate the amount of transcription from the promoter. A “constitutive” promoter is a promoter that is active under most physiological and developmental conditions. An “inducible” promoter is a promoter that is regulated depending on physiological or developmental conditions. A “tissue specific” promoter is preferentially active in specific types of differentiated cells/tissues, such as preferably a T cell.

Operably linked

“Operably linked” is defined herein as a configuration in which a control sequence such as a promoter sequence or regulating sequence is appropriately placed at a position relative to the nucleotide sequence of interest, preferably coding for a δ TCR (or a γ TCR) chain comprising the polypeptide as identified such that the promoter or control or regulating sequence directs or affects the transcription and/or production or

expression of the nucleotide sequence of interest, preferably encoding a δ TCR (or γ TCR) chain comprising the polypeptide as identified in a cell and/or in a subject. For instance, a promoter is operably linked to a coding sequence if the promoter is able to initiate or regulate the transcription or expression of a coding sequence, in which case
5 the coding sequence should be understood as being "under the control of" the promoter.

Viral expression construct

An expression construct carries a genome that is able to stabilize and remain episomal in a cell. Within the context of the invention, a cell may mean to encompass a cell used
10 to make the construct or a cell wherein the construct will be administered. Alternatively a construct is capable of integrating into a cell's genome, *e.g.* through homologous recombination or otherwise. A particularly preferred expression construct is one wherein a nucleotide sequence encoding a δ TCR (or γ TCR) chain or part thereof is operably linked to a promoter as defined herein wherein said promoter is capable of
15 directing expression of said nucleotide sequence (*i.e.* coding sequence) in a cell. Such a preferred expression construct is said to comprise an expression cassette. An expression cassette as used herein comprises or consists of a nucleotide sequence encoding a δ TCR (or γ TCR) chain or part thereof. A viral expression construct is an expression construct which is intended to be used in gene therapy. It is designed to
20 comprise part of a viral genome as later defined herein.

Expression constructs disclosed herein could be prepared using recombinant techniques in which nucleotide sequences encoding said δ TCR (or γ TCR) chain or part thereof are expressed in a suitable cell, *e.g.* cultured cells or cells of a multicellular organism, such as described in Ausubel *et al.*, "Current Protocols in Molecular
25 Biology", Greene Publishing and Wiley-Interscience, New York (1987) and in Sambrook and Russell (2001, *supra*); both of which are incorporated herein by reference in their entirety. Also see, Kunkel (1985) Proc. Natl. Acad. Sci. 82:488 (describing site directed mutagenesis) and Roberts *et al.* (1987) Nature 328:731-734 or Wells, J.A., *et al.* (1985) Gene 34: 315 (describing cassette mutagenesis).

30 Typically, a nucleic acid or nucleotide sequence encoding a δ TCR (or γ TCR) chain is used in an expression construct or expression vector. The phrase "expression vector" generally refers to a nucleotide sequence that is capable of effecting expression of a gene in a host compatible with such sequences. These expression vectors typically

include at least suitable promoter sequences and optionally, transcription termination signals. An additional factor necessary or helpful in effecting expression can also be used as described herein. A nucleic acid or DNA or nucleotide sequence encoding a δ TCR (or γ TCR) chain is incorporated into a DNA construct capable of introduction
5 into and expression in an *in vitro* cell culture. Specifically, a DNA construct is suitable for replication in a prokaryotic host, such as bacteria, *e.g.*, *E. coli*, or can be introduced into a cultured mammalian, plant, insect, (*e.g.*, Sf9), yeast, fungi or other eukaryotic cell lines.

A DNA construct prepared for introduction into a particular host may include a
10 replication system recognized by the host, an intended DNA segment encoding a desired polypeptide, and transcriptional and translational initiation and termination regulatory sequences operably linked to the polypeptide-encoding segment. The term “operably linked” has already been defined herein. For example, a promoter or enhancer is operably linked to a coding sequence if it stimulates the transcription of the
15 sequence. DNA for a signal sequence is operably linked to DNA encoding a polypeptide if it is expressed as a preprotein that participates in the secretion of a polypeptide. Generally, a DNA sequence that is operably linked are contiguous, and, in the case of a signal sequence, both contiguous and in reading frame. However, enhancers need not be contiguous with a coding sequence whose transcription they
20 control. Linking is accomplished by ligation at convenient restriction sites or at adapters or linkers inserted in lieu thereof, or by gene synthesis.

The selection of an appropriate promoter sequence generally depends upon the host cell selected for the expression of a DNA segment. Examples of suitable promoter sequences include prokaryotic, and eukaryotic promoters well known in the art (see,
25 *e.g.* Sambrook and Russell, 2001, *supra*). A transcriptional regulatory sequence typically includes a heterologous enhancer or promoter that is recognised by the host. The selection of an appropriate promoter depends upon the host, but promoters such as the *trp*, *lac* and phage promoters, tRNA promoters and glycolytic enzyme promoters are known and available (see, *e.g.* Sambrook and Russell, 2001, *supra*). An expression
30 vector includes the replication system and transcriptional and translational regulatory sequences together with the insertion site for the polypeptide encoding segment can be employed. In most cases, the replication system is only functional in the cell that is used to make the vector (bacterial cell as *E. Coli*). Most plasmids and vectors do not

replicate in the cells infected with the vector. Examples of workable combinations of cell lines and expression vectors are described in Sambrook and Russell (2001, *supra*) and in Metzger *et al.* (1988) *Nature* 334: 31-36. For example, suitable expression vectors can be expressed in, yeast, *e.g. S.cerevisiae*, *e.g.*, insect cells, *e.g.*, Sf9 cells, mammalian cells, *e.g.*, CHO cells and bacterial cells, *e.g.*, *E. coli*. A cell may thus be a prokaryotic or eukaryotic host cell. A cell may be a cell that is suitable for culture in liquid or on solid media.

Alternatively, a host cell is a cell that is part of a multicellular organism such as a transgenic plant or animal.

Viral vector

A viral vector or a gene therapy vector is a vector that comprises a viral expression construct as defined above.

A viral vector or a gene therapy vector is a vector that is suitable for gene therapy. Vectors that are suitable for gene therapy are described in Anderson 1998, *Nature* 392: 25-30; Walther and Stein, 2000, *Drugs* 60: 249-71; Kay *et al.*, 2001, *Nat. Med.* 7: 33-40; Russell, 2000, *J. Gen. Virol.* 81: 2573-604; Amado and Chen, 1999, *Science* 285: 674-6; Federico, 1999, *Curr. Opin. Biotechnol.* 10: 448-53; Vigna and Naldini, 2000, *J. Gene Med.* 2: 308-16; Marin *et al.*, 1997, *Mol. Med. Today* 3: 396-403; Peng and Russell, 1999, *Curr. Opin. Biotechnol.* 10: 454-7; Sommerfelt, 1999, *J. Gen. Virol.* 80: 3049-64; Reiser, 2000, *Gene Ther.* 7: 910-3; and references cited therein.

A particularly suitable gene therapy vector includes an Adenoviral and Adeno-associated virus (AAV) vector. These vectors infect a wide number of dividing and non-dividing cell types including synovial cells and liver cells. The episomal nature of the adenoviral and AAV vectors after cell entry makes these vectors suited for therapeutic applications. (Russell, 2000, *J. Gen. Virol.* 81: 2573-2604; Goncalves, 2005, *Virol J.* 2(1):43) as indicated above. AAV vectors are even more preferred since they are known to result in very stable long term expression of transgene expression (up to 9 years in dog (Niemeyer *et al*, *Blood*, 2009 Jan 22;113(4):797-806) and ~ 2 years in human (Nathwani *et al*, *N Engl J Med.* 2011 Dec 22;365(25):2357-65, Simonelli *et al*, *Mol Ther.* 2010 Mar;18(3):643-50. Epub 2009 Dec 1.)). Preferred adenoviral vectors are modified to reduce the host response as reviewed by Russell (2000, *supra*). Method for gene therapy using AAV vectors are described by Wang *et*

al., 2005, J Gene Med. March 9 (Epub ahead of print), Mandel *et al.*, 2004, Curr Opin Mol Ther. 6(5):482-90, and Martin *et al.*, 2004, Eye 18(11):1049-55, Nathwani et al, N Engl J Med. 2011 Dec 22;365(25):2357-65, Apparailly et al, Hum Gene Ther. 2005 Apr;16(4):426-34.

5 Another suitable gene therapy vector includes a retroviral vector. A preferred retroviral vector for application in the present invention is a lentiviral based expression construct. Lentiviral vectors have the ability to infect and to stably integrate into the genome of dividing and non-dividing cells (Amado and Chen, 1999 Science 285: 674-6). Methods for the construction and use of lentiviral based expression constructs are
10 described in U.S. Patent No.'s 6,165,782, 6,207,455, 6,218,181, 6,277,633 and 6,323,031 and in Federico (1999, Curr Opin Biotechnol 10: 448-53) and Vigna *et al.* (2000, J Gene Med 2000; 2: 308-16).

Other suitable gene therapy vectors include a herpes virus vector, a polyoma virus vector or a vaccinia virus vector.

15 A gene therapy vector comprises a nucleotide encoding a δ TCR chain (or γ TCR), whereby each of said nucleotide sequence is operably linked to the appropriate regulatory sequences. Such regulatory sequence will at least comprise a promoter sequence. Suitable promoters for expression of such a nucleotide sequence from gene therapy vectors include *e.g.* cytomegalovirus (CMV) intermediate early promoter, viral
20 long terminal repeat promoters (LTRs), such as those from murine moloney leukaemia virus (MMLV) rous sarcoma virus, or HTLV-1, the simian virus 40 (SV 40) early promoter and the herpes simplex virus thymidine kinase promoter. Transposon or other non-viral delivery systems may also be used in this context. All systems can be used in vitro or in vivo.

25

A gene therapy vector may optionally comprise a further nucleotide sequence coding for a further polypeptide. A further polypeptide may be a (selectable) marker polypeptide that allows for the identification, selection and/or screening for cells containing the expression construct. Suitable marker proteins for this purpose are *e.g.*
30 the fluorescent protein GFP, and the selectable marker genes HSV thymidine kinase (for selection on HAT medium), bacterial hygromycin B phosphotransferase (for selection on hygromycin B), Tn5 aminoglycoside phosphotransferase (for selection on G418), and dihydrofolate reductase (DHFR) (for selection on methotrexate), CD20, the

low affinity nerve growth factor gene. Sources for obtaining these marker genes and methods for their use are provided in Sambrook and Russel (2001) "Molecular Cloning: A Laboratory Manual (3rd edition), Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, New York.

- 5 A gene therapy vector is preferably formulated in a pharmaceutical composition as defined herein. In this context, a pharmaceutical composition may comprise a suitable pharmaceutical carrier as earlier defined herein.

Transgene

- 10 A "transgene" is herein defined as a gene or a nucleic acid molecule (*i.e.* a molecule encoding a δ TCR (or a γ TCR) chain) that has been newly introduced into a cell, *i.e.* a gene that may be present but may normally not be expressed or expressed at an insufficient level in a cell. The transgene may comprise sequences that are native to the cell, sequences that naturally do not occur in the cell and it may comprise combinations
- 15 of both. A transgene may contain sequences coding for a δ TCR (or a γ TCR) chain and comprising the polypeptide as identified and/or additional proteins as earlier identified herein that may be operably linked to appropriate regulatory sequences for expression of the sequences coding for a δ TCR (or a γ TCR) chain. Preferably, the transgene is not integrated into the host cell's genome.

20

Transduction

- "Transduction" refers to the delivery of a δ TCR (or a γ TCR) chain or parts thereof into a recipient host cell by a viral vector. For example, transduction of a target cell by a retroviral or lentiviral vector of the invention leads to transfer of the genome contained
- 25 in that vector into the transduced cell.

Host cell/target cell

- "Host cell" or "target cell" refers to the cell into which the DNA delivery takes place, such as the T cells of a donor.

30

Engineered cells

"Engineered cells" refers herein to cells having been engineered, e.g. by the introduction of an exogenous nucleic acid sequence as defined herein. Such a cell has

been genetically modified for example by the introduction of for example one or more mutations, insertions and/or deletions in the endogenous gene and/or insertion of a genetic construct in the genome. An engineered cell may refer to a cell in isolation or in culture. Engineered cells may be "transduced cells" wherein the cells have been
5 infected with e.g. a modified virus, for example, a retrovirus may be used, such as described in the examples, but other suitable viruses may also be contemplated such as lentiviruses. Non-viral methods may also be used, such as transfections. Engineered cells may thus also be "stably transfected cells" or "transiently transfected cells". Transfection refers to non-viral methods to transfer DNA (or RNA) to cells such that a
10 gene is expressed. Transfection methods are widely known in the art, such as calciumphosphate transfection, PEG transfection, and liposomal or lipoplex transfection of nucleic acids. Such a transfection may be transient, but may also be a stable transfection wherein cells can be selected that have the gene construct integrated in their genome. In some cases genetic engineering systems such as CRISPR or
15 Argonaute maybe utilized to design engineered cells that express a polypeptide described herein.

Pharmaceutical composition/ method of treatment

In therapeutic applications, an effective amount of a δ TCR (or γ TCR) chain or parts
20 thereof or nucleic acid construct or viral vector or cell expressing these molecules as defined herein is administered to a subject.

The term "effective amount" as used herein is defined as the amount of the molecules of the present invention that are necessary to result in the desired physiological change in the cell or tissue to which it is administered. The term "therapeutically effective
25 amount" as used herein is defined as the amount of the molecules of the present invention that achieves a desired effect with respect to cancer. In this context, a "desired effect" is synonymous with "an anti-tumour activity" as earlier defined herein. A skilled artisan readily recognizes that in many cases the molecules may not provide a cure but may provide a partial benefit, such as alleviation or improvement of at least
30 one symptom or parameter. In some embodiments, a physiological change having some benefit is also considered therapeutically beneficial. Thus, in some embodiments, an amount of molecules that provides a physiological change is considered an "effective amount" or a "therapeutically effective amount."

Pharmaceutical compositions of the present invention comprise an effective amount of one or more molecules (i.e. a polypeptide comprising a δ TCR or γ TCR chain or variants or parts thereof or nucleic acid construct or viral vector or cell expressing these molecules as defined herein) optionally dissolved or dispersed in a pharmaceutically acceptable carrier. The phrases "pharmaceutical or pharmacologically acceptable" refers to molecular entities and compositions that do not produce or produce acceptable adverse, allergic or other untoward reaction when administered to an animal, such as, for example, a human, as appropriate. Whether certain adverse effects are acceptable is determined based on the severity of the disease. The preparation of a pharmaceutical composition that contains at least one active ingredient will be known to those of skill in the art in light of the present disclosure, as exemplified by Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, incorporated herein by reference. Moreover, for animal (e.g., human) administration, it will be understood that preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biological Standards.

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, surfactants, antioxidants, preservatives (e.g., antibacterial agents, antifungal agents), isotonic agents, absorption delaying agents, salts, preservatives, drugs, drug stabilizers, gels, binders, excipients, disintegration agents, lubricants, sweetening agents, flavoring agents, dyes, such like materials and combinations thereof, as would be known to one of ordinary skill in the art (see, for example, Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, pp. 1289-1329, incorporated herein by reference). Except insofar as any conventional carrier is incompatible with the active ingredient, its use in the therapeutic or pharmaceutical compositions is contemplated. In certain embodiments, a pharmaceutical composition described herein comprising a population of cells described herein, further comprises a suitable amount of an antifungal agent. In some cases, a pharmaceutical composition described herein comprises an antifungal agent in an amount sufficient for the the pharmaceutical composition to retain at least 20%, 30%, 40%, 50%, 60%, 70%, 80% or 90% of its desired activity for a period of at least 1 month, 2 months, 3 months, 6 months, 1 year, 1.5 years, 2 years, 2.5 years or 3 years.

The actual dosage amount of a composition of the present invention administered to an animal or a patient can be determined by physical and physiological factors such as body weight, severity of condition, the type of disease being treated, previous or concurrent therapeutic interventions, idiopathy of the patient and on the route of administration. The practitioner responsible for administration will, in any event, determine the concentration of active ingredient(s) in a composition and appropriate dose(s) for the individual subject.

T Cell as pharmaceutical composition

Prior to expansion and genetic modification of the T cells of the invention, a source of T cells is obtained from a subject. T cells can be obtained from a number of sources, including PBMCs, bone marrow, lymph node tissue, cord blood, thymus tissue, tissue from a site of infection, ascites, pleural effusion, spleen tissue, and tumors. In certain embodiments of the present invention, T cells can be obtained from a unit of blood collected from a subject using any number of techniques known to the skilled artisan, such as FicollTM separation. In one embodiment, cells from the circulating blood of an individual are obtained by apheresis. The apheresis product typically contains lymphocytes, including T cells, monocytes, granulocytes, B cells, other nucleated white blood cells, red blood cells, and platelets. In one embodiment, the cells collected by apheresis may be washed to remove the plasma fraction and to place the cells in an appropriate buffer or media for subsequent processing steps. In a particular embodiment, the engineered cell can be a T cell. The engineered cell can be an effector (T_{EFF}), effector-memory (T_{EM}), central-memory (T_{CM}), T memory stem (T_{SCM}), naïve (T_N), or CD4⁺ or CD8⁺ T cell. The T cells can also be selected from a bulk population, for example, selecting T cells from whole blood. The T cells can also be expanded from a bulk population. The T cells can also be skewed towards particular populations and phenotypes. The engineered cell can also be expanded *ex vivo*. The engineered cell can be formulated into a pharmaceutical composition. The engineered cell can be formulated into a pharmaceutical composition and used to treat a subject in need thereof as earlier explained herein. The engineered cell can be autologous to a subject in need thereof. The engineered cell can be allogeneic to a subject in need thereof. The engineered cell can also be a good manufacturing practices (GMP) compatible reagent. The engineered cell can be part of a combination therapy to treat a subject in need

thereof. The engineered cell can be a human cell. The subject that is being treated can be a human.

A method of attaining suitable cells can comprise sorting cells. In some cases, a cell
5 can comprise a marker that can be selected for the cell. For example, such marker can
comprise GFP, a resistance gene, a cell surface marker, an endogenous tag. Cells can
be selected using any endogenous marker. Suitable cells can be selected or sorted using
any technology. Such technology can comprise flow cytometry and/or magnetic
columns. The selected cells can then be infused into a subject. The selected cells can
10 also be expanded to large numbers. The selected cells can be expanded prior to
infusion.

Vectors can be delivered to cells *ex vivo*, such as cells explanted from an individual
patient (e.g., lymphocytes, T cells, bone marrow aspirates, tissue biopsy), followed by
re-implantation of the cells into a patient, usually after selection for cells which have
15 incorporated the vector. Prior to or after selection, the cells can be expanded.

Ex vivo cell transfection can also be used for diagnostics, research, or for gene therapy
(e.g. via re-infusion of the transfected cells into the host organism). In some cases, cells
are isolated from the subject organism, transfected with a nucleic acid (e.g., gene or
DNA), and re-infused back into the subject organism (e.g. patient). Further, also in vivo
20 cell transfection can be used for gene therapy, in order to reduced immune reactions of
the patient.

In some cases, populations of engineered T cells may be formulated for administration
to a subject using techniques known to the skilled artisan. Formulations comprising
populations of engineered T cells may include pharmaceutically acceptable
25 excipient(s). Excipients included in the formulations will have different purposes
depending, for example, on the subpopulation of T cells used and the mode of
administration. Examples of generally used excipients included, without limitation:
saline, buffered saline, dextrose, water-for-injection, glycerol, ethanol, and
combinations thereof, stabilizing agents, solubilizing agents and surfactants, buffers
30 and preservatives, tonicity agents, bulking agents, and lubricating agents. The
formulations comprising populations of engineered T cells will typically have been
prepared and cultured in the absence of any non-human components, such as animal
serum.

A formulation may include one population of engineered T cells, or more than one, such as two, three, four, five, six or more population of engineered T cells. The formulations comprising population(s) of engineered T cells may be administered to a subject using modes and techniques known to the skilled artisan. Exemplary modes
5 include, but are not limited to, intravenous injection. Other modes include, without limitation, intratumoral, intradermal, subcutaneous (s.c., s.q., sub-Q, Hypo), intramuscular (i.m.), intraperitoneal (i.p.), intra-arterial, intramedullary, intracardiac, intra-articular (joint), intrasynovial (joint fluid area), intracranial, intraspinal, and intrathecal (spinal fluids). Any known device useful for parenteral injection of infusion
10 of the formulations can be used to effect such administration. The formulations comprising population(s) of engineered T cells that are administered to a subject comprise a number of engineered T cells that is effective for the treatment and/or prophylaxis of the specific indication or disease. Thus, therapeutically-effective populations of engineered T cells are administered to subjects when the methods of the
15 present invention are practiced. In general, formulations are administered that comprise between about 1×10^4 and about 1×10^{10} engineered T cells. In most cases, the formulation will comprise between about 1×10^5 and about 1×10^9 engineered T cells, from about 5×10^5 to about 5×10^8 engineered T cells, or from about 1×10^6 to about 1×10^7 engineered T cells. However, the number of engineered T cells administered to a
20 subject will vary between wide limits, depending upon the location, source, identity, extent and severity of the cancer, the age and condition of the individual to be treated etc. A physician will ultimately determine appropriate dosages to be used.

General

25 In this document and in its claims, the verb "to comprise" and its conjugations is used in its non-limiting sense to mean that items following the word are included, but items not specifically mentioned are not excluded. In addition the verb "to consist" may be replaced by "to consist essentially of" meaning that a method as defined herein may comprise additional step(s), respectively component(s) than the ones specifically
30 identified, said additional step(s), respectively component(s) not altering the unique characteristic of the invention. In addition, reference to an element by the indefinite article "a" or "an" does not exclude the possibility that more than one of the element is present, unless the context clearly requires that there be one and only one of the

elements. The indefinite article "a" or "an" thus usually means "at least one".

The word "about" when used in association with an integer (about 10) preferably means that the value may be the given value of 10 more or less 1 of the value: about 10 preferably means from 9 to 11. The word "about" when used in association with a
5 numerical value (about 10.6) preferably means that the value may be the given value of 10.6 more or less 1% of the value 10.6.

All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety. The following examples are offered for
10 illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

Table 1: Overview of the sequences of the sequence listing

<u>SEQ ID NO</u>	<u>Name</u>	<u>Amino acid (aa) or DNA</u>
<u>1</u>	<u>Primer table 1</u>	<u>DNA</u>
<u>2</u>	<u>Primer table 1</u>	<u>DNA</u>
<u>3</u>	<u>Primer table 1</u>	<u>DNA</u>
<u>4</u>	<u>Primer table 1</u>	<u>DNA</u>
<u>5</u>	<u>Primer table 1</u>	<u>DNA</u>
<u>6</u>	<u>Primer table 1</u>	<u>DNA</u>
<u>7</u>	<u>CDR3 VD1 Fe11</u>	<u>aa</u>
<u>8</u>	<u>CDR3 VD2 cl3</u>	<u>aa</u>
<u>9</u>	<u>CD3 VD2 cl5</u>	<u>aa</u>
<u>10</u>	<u>CDR3 VG5 Fe11</u>	<u>aa</u>
<u>11</u>	<u>CDR3 VG cl3</u>	<u>aa</u>
<u>12</u>	<u>CDR3 VG9 cl5</u>	<u>aa</u>
<u>13</u>	<u>TRD cl3</u>	<u>aa</u>
<u>14</u>	<u>TRG cl3</u>	<u>aa</u>
<u>15</u>	<u>TRD cl5</u>	<u>aa</u>
<u>16</u>	<u>TRG cl5</u>	<u>aa</u>
<u>17</u>	<u>TRD Fe11</u>	<u>aa</u>
<u>18</u>	<u>TRG Fe11</u>	<u>aa</u>
<u>19</u>	<u>CDR3 VG4 E113</u>	<u>aa</u>
<u>20</u>	<u>CDR3 VG2 F4</u>	<u>aa</u>
<u>21</u>	<u>CDR3 VG8 Zi11</u>	<u>aa</u>
<u>22</u>	<u>CDR3 VD5 D37</u>	<u>aa</u>
<u>23</u>	<u>CDR3 VD5 E113</u>	<u>aa</u>
<u>24</u>	<u>CDR3 VD1 F4</u>	<u>aa</u>
<u>25</u>	<u>CDR3 VD1 Zi11</u>	<u>aa</u>
<u>26</u>	<u>TRD E113</u>	<u>DNA wild type</u>
<u>27</u>	<u>CDR3 VG4 C132</u>	<u>aa</u>
<u>28</u>	<u>CDR3 VD5 C132</u>	<u>aa</u>
<u>29</u>	<u>TRD E113</u>	<u>aa</u>
<u>30</u>	<u>TRG E113</u>	<u>aa</u>
<u>31</u>	<u>TRD F4</u>	<u>aa</u>
<u>32</u>	<u>TRG F4</u>	<u>aa</u>
<u>33</u>	<u>TRD Zi11</u>	<u>aa</u>
<u>34</u>	<u>TRG Zi11</u>	<u>aa</u>
<u>35</u>	<u>TRD D37</u>	<u>aa</u>
<u>36</u>	<u>TRG D37</u>	<u>aa</u>
<u>37</u>	<u>TRD C132</u>	<u>aa</u>
<u>38</u>	<u>TRG C132</u>	<u>aa</u>
<u>39</u>	<u>CDR3 VG8 D37</u>	<u>aa</u>
<u>40</u>	<u>CDR3 VD3 F2</u>	<u>aa</u>
<u>41</u>	<u>TRG E113</u>	<u>DNA wild type</u>
<u>42</u>	<u>TRD F4</u>	<u>DNA wild type</u>

<u>43</u>	<u>TRG F4</u>	<u>DNA wild type</u>
<u>44</u>	<u>TRD Zi11</u>	<u>DNA wild type</u>
<u>45</u>	<u>TRG Zi11</u>	<u>DNA wild type</u>
<u>46</u>	<u>TRDD37</u>	<u>DNA wild type</u>
<u>47</u>	<u>TRG D37</u>	<u>DNA wild type</u>
<u>48</u>	<u>TRD C132</u>	<u>DNA wild type</u>
<u>49</u>	<u>TRG C132</u>	<u>DNA wild type</u>
<u>50</u>	<u>TRD Fe11</u>	<u>DNA wild type</u>
<u>51</u>	<u>TRG Fe11</u>	<u>DNA wild type</u>
<u>52</u>	<u>TRD E113</u>	<u>DNA codon opt.</u>
<u>53</u>	<u>TRG E113</u>	<u>DNA codon opt.</u>
<u>54</u>	<u>TRD F4</u>	<u>DNA codon opt.</u>
<u>55</u>	<u>TRG F4</u>	<u>DNA codon opt.</u>
<u>56</u>	<u>TRD Zi11</u>	<u>DNA codon opt.</u>
<u>57</u>	<u>TRG Zi11</u>	<u>DNA codon opt.</u>
<u>58</u>	<u>TRD D37</u>	<u>DNA codon opt.</u>
<u>59</u>	<u>TRG D37</u>	<u>DNA codon opt.</u>
<u>60</u>	<u>TRD C132</u>	<u>DNA codon opt.</u>
<u>61</u>	<u>TRG C132</u>	<u>DNA codon opt.</u>
<u>62</u>	<u>TRD Fe11</u>	<u>DNA codon opt.</u>
<u>63</u>	<u>TRG Fe11</u>	<u>DNA codon opt.</u>
<u>64</u>	<u>Longer CDR3 VG5 Fe11</u>	<u>aa</u>
<u>65</u>	<u>CDR3 VG4 F2</u>	<u>Aa</u>
<u>66</u>	<u>CDR3 VD1 Ze11</u>	<u>aa</u>
<u>67</u>	<u>CDR3 VG8 Ze11</u>	<u>aa</u>
<u>68</u>	<u>CDR3 VD5 B23</u>	<u>aa</u>
<u>69</u>	<u>CDR3 VG8 B23</u>	<u>aa</u>
<u>70</u>	<u>TRD F2</u>	<u>aa</u>
<u>71</u>	<u>TRG F2</u>	<u>aa</u>
<u>72</u>	<u>TRD Ze11</u>	<u>aa</u>
<u>73</u>	<u>TRG Ze11</u>	<u>aa</u>
<u>74</u>	<u>TRD B23</u>	<u>aa</u>
<u>75</u>	<u>TRG B23</u>	<u>aa</u>
<u>76</u>	<u>TRD F2</u>	<u>DNA wild type</u>
<u>77</u>	<u>TRG F2</u>	<u>DNA wild type</u>
<u>78</u>	<u>TRD Ze11</u>	<u>DNA wild type</u>
<u>79</u>	<u>TRG Ze11</u>	<u>DNA wild type</u>
<u>80</u>	<u>TRD B23</u>	<u>DNA wild type</u>
<u>81</u>	<u>TRG B23</u>	<u>DNA wild type</u>
<u>82</u>	<u>TRD F2</u>	<u>DNA codon opt.</u>
<u>83</u>	<u>TRG F2</u>	<u>DNA codon opt.</u>
<u>84</u>	<u>TRD Ze11</u>	<u>DNA codon opt.</u>
<u>85</u>	<u>TRG Ze11</u>	<u>DNA codon opt.</u>
<u>86</u>	<u>TRD B23</u>	<u>DNA codon opt.</u>
<u>87</u>	<u>TRG B23</u>	<u>DNA codon opt.</u>

60

<u>88</u>	<u>CDR3 VD1 B9</u>	<u>aa</u>
<u>89</u>	<u>CDR3 VG5 B9</u>	<u>aa</u>
<u>90</u>	<u>TRD B9</u>	<u>aa</u>
<u>91</u>	<u>TRG B9</u>	<u>aa</u>
<u>92</u>	<u>TRD B9</u>	<u>DNA wild type</u>
<u>93</u>	<u>TRG B9</u>	<u>DNA wild type</u>
<u>94</u>	<u>TRD B9</u>	<u>DNA codon opt.</u>
<u>95</u>	<u>TRG B9</u>	<u>DNA codon opt.</u>
<u>96</u>	<u>TRD FE11 without signal pept.</u>	<u>aa</u>
<u>97</u>	<u>TRG FE11 without signal pept.</u>	<u>aa</u>
<u>98</u>	<u>TRD clone 3 without signal pept.</u>	<u>aa</u>
<u>99</u>	<u>TRG clone 3 without signal pept.</u>	<u>aa</u>
<u>100</u>	<u>TRD clone 5 without signal pept.</u>	<u>aa</u>
<u>101</u>	<u>TRG clone 5 without signal pept.</u>	<u>aa</u>

Examples

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims.

Unless specified, reagents employed in the examples are commercially available or can be prepared using commercially available instrumentation, methods, or reagents known in the art. The examples illustrate various aspects of the invention and practice of the methods of the invention. The examples are not intended to provide an exhaustive description of the many different embodiments of the invention. Thus, although the invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, those of ordinary skill in the art will realize readily that many changes and modifications can be made thereto without departing from the spirit or scope of the appended claims.

Example 1

Material & Methods

Frozen PBMC's from healthy donors were thawed and stained with the following antibodies: CD3 (eFluor450 clone okt 3 1:40 eBioscience), $\alpha\beta$ TCR (APC clone IP26 1:10, eBioscience), and pan $\gamma\delta$ TCR (PE clone IMMU510 1:10 Boeckman Coulter), or with CD3 (PE clone UCHT1 1:20 BD), CD4 (APC clone RPA-T4 1:100 Biolegend), CD8 (PerCP-Cy5.5 clone RPA-T8 1:1000 Biolegend), CD27 (APC eFluor780 clone 0323 1:20 eBioscience), and CD45RA (PB clone HI100 1:50 Biolegend). Samples were sorted with flow cytometry on the ARIAII (BD) in an $\alpha\beta$ -fraction (CD3⁺, $\alpha\beta$ ⁺), and a $\gamma\delta$ -fraction (CD3⁺, $\gamma\delta$ ⁺), or in different subsets of CD4⁺ and CD4⁻ T cells. RNA was isolated using Qiagen Rneasy Minikit for samples $\geq 0.5 \times 10^6$ cells or Qiagen Rneasy Microkit for samples $< 0.5 \times 10^6$ cells following manufacturer's instructions. Specific cDNA for TRB and TRD was synthesized with Superscript® II Reverse Transcriptase (ThermoFisher), using a specific primer at the 3' constant region of TRB or TRD. A universal template switch adaptor, containing a unique molecular identifier (UMI), was incorporated at the 5' end of the V region to be able to take PCR bias into account

during the analysis. An overview of the used primers can be found in Table 1. After cDNA synthesis the cDNA was purified using NucleoSpin® Gel and PCR Clean-UP (Machery-Nagel). cDNA was amplified during with PCR using Q5® High Fidelity DNA polymerase (New England Biolabs). A specific nested primer at the constant region of TRB or TRD, and a step-out primer which anneals to the switch adaptor were used (Table 1). PCR amplification was performed on a T100 Thermal Cycler (Biorad) using the following steps 90 s at 98°C, 35 cycles of 7 s at 98°C, 20 s at 62°C, 50 s at 72°C, followed by 10 m at 72°C. PCR products were purified using NucleoSpin® Gel and PCR Clean-UP (Machery-Nagel). PCR products were analyzed with QIAxcel Advanced System (Qiagen).

TruSeq Barcode adapters (Illumina) were ligated to the PCR products using the ClaSeek Ligation Mix (Thermo Scientific), according to the recommendations of the manufacturer. Cleanup of the samples was performed with The Agencourt AMPure XP system (Beckman Coulter). Next-generation sequencing was performed on an Illumina MiSeq system 500 (2x250 bp) (Illumina).

Sequencing data were analyzed with the MiXCR program (Bolotin, et al. (7)). To correct for PCR bias UMI's were extracted from the sequencing data using the MIGEC pipeline (Shugay, et al. (8)) Only sequences with a valid UMI were used for further analyses. To search for shared CDR3 sequence regions within the different donors an interactive webtool was used (<http://bioinformatics.psb.ugent.be/webtools/Venn/>).

Further methods for functional analyses have been described in Sebestyen et al (Cell Rep. 2016, 15: 1973-1985) and Scheper et al (Leukemia, 2013, 27: 1328-1338).

Table 2 Overview primers
cDNA
synthesis

Constant region TRB	CAGTATCTGGAGTCATTGA (SEQ ID NO:1)
Constant region TRD	CTTGGATGACACGAGAT (SEQ ID NO: 2)
Template switch adaptor with UMI	AAGCAGUGGTAUCAACGCAGAGUNNNNUNNNNUNNNNUCTT(rG) ₄ (SEQ ID NO:3)

PCR
amplification

Nested	TGCTTCTGATGGCTCAAACAC (SEQ ID NO:4)
--------	-------------------------------------

primer constant region TRB	
Nested primer constant region TRD	AACGGATGGTTTGGTATGAG (SEQ ID NO:5)
Step out primer (anneals to switch adaptor)	CACTCTATCCGACAAGCAGTGGTATCAACGCAG (SEQ ID NO:6)

rG= riboguanosine, necessary for the template switch adaptor to anneal. The backbone of this adaptor is cDNA, but in this part there are four RNA-G's. This template switch adapter is described in US5962272 and in Zhu Y et al (Zhu Y et al 2001, BioTechniques, 30: 892-897).

5

Results

We hypothesized that rapid sequencing of δ TCR chains by next generation sequencing followed by comparison of (dominant) clones between different individuals can be used to rapidly identify functionally relevant clones, which are shared between different individuals. In order to test this hypothesis, we isolated $\gamma\delta$ T cells from 6 different donors and analyzed the δ TCR repertoire by NGS as indicated in material and methods. For further analyses two strategies were used, either the individual analysis of three donors indicated as HD 18, 19, and 20 as well as a pool of three donors, depicted as “combined” (HD11, 12, and 15). In addition, we utilized as hypothetical 7th donor the randomly collected database of δ TCR chains obtained by single cell cloning and sequencing from many different donors, which are available in the public and our private database. After correction for the percentage of overlapping unique clones for overlapping UMI's a 7-fold higher amount of shared δ TCR has been observed when compared to β TCR repertoire (Figure 1). When further adjusting the corrected ratios for the average size of the found repertoire, the difference between the frequency of observed overlapping clones was even more pronounced with 1-2 log differences more overlapping clones observed for shared δ TCR chains (Figure 2). Thus, although the theoretical δ TCR repertoire is substantially bigger than the theoretical β TCR repertoire; we show that the chance of finding an overlapping clone in the δ TCR repertoire is 1-2

log-higher than when analyzing a β TCR repertoire. We hypothesized that these shared δ TCR chains must provide a specific survival advantage and could be therefore essential in function, such as daily tumor surveillance.

5 Comparing δ TCR sequences between different NGS-donors results in the identification of a limited number of clones, which have been shared at least between two different donors (Figure 3 and Table 2 and not shown). In addition, we compared our δ TCR sequences from HDs with the 7th hypothetical donor. Thereby we identified additional δ 2 negative TCR chain, which is the second most frequent sequence in donor HD19
10 (Tables 3 and 4) and δ 2 positive TCR chains, which are shared at least between HD15 and the 7th hypothetical donor (Table 3). Some of these relevant clones are within the top 30 clones of at least one the healthy donors, indicating the potential biological relevance of these clones. As proof of concept for the functional and potential therapeutic relevance of identified shared δ TCR sequences we further searched for
15 described complete $\gamma\delta$ TCR sequences harboring by us so far identified dominant and shared δ TCR sequences. We identified clone FE11 (Table 3), which has been described recently by our group (δ 2 negative δ TCR (9), Table 2). Clone Fe11 was the second most prevalent clone in HD19 (Table 4). Screening of tumor cell lines classified this clone as tumor reactive against malignant B cell and solid cancer cell lines ((10) and
20 Figure 4). In addition we could identify one overlapping δ 2 positive δ TCR sequence as part of the distinct clone 5, and one sequence corresponding to the so called clone 3, which have both also been extensively described by our group (Table 3 and (11)). Clone 5 was found in the top 20 of clones of HD15 (Table 4). This sequence has been reported to be very active against many types of hematological and solid cancer cells
25 ((11, 12) and Table 4).

In summary, we provide proof of concept that we can down scale a highly diverse immune repertoire to very limited number of sequences, which are tumor reactive and therefore therapeutically relevant. In addition to anti- tumor reactivity identified
30 sequences can also be active highly against infection as evidenced by the identification of shared V δ 2 positive TCR chain sequences. Thus, high throughput sequencing of δ TCR chains followed by the comparison of sequences between different donors can be

a very powerful tool and result in the rapid identification of δ TCR chains which can be interesting for therapeutic applications.

Table 3. Shared clonotypes		Vdelta	Previously described
V δ 2 negative			
CALGDSYGGGPLYTDKLIF (SEQ ID NO: 7)		Vd1	Fe11
V δ 2 positive			
CACDLLGYTDKLIF (SEQ ID NO: 8)		Vd2	Cl3
CACDALKRTDTDKLIF (SEQ ID NO: 9)		Vd2	Cl5

5

Table 4. Characteristics known clones		
Clone	CDR3 δ chain	CDR3 γ chain
Fe11	CALGDSYGGGPLYTDKLIF SEQ ID NO:7	ATWDRPEIYYKKL SEQ ID NO: 10
Cl3	CACDLLGYTDKLIF SEQ ID NO:8	CALWEEELGKKIKVF SEQ ID NO: 11
Cl5	CACDALKRTDTDKLIF SEQ ID NO: 9	CALWEIQELGKKIKVF SEQ ID NO:12

Discussion

Main finding of this study is that by comparing NGS data of β and δ TCR chains from different healthy donors we observe a substantial frequency of shared δ TCR chains between different individuals, though the β TCR repertoire is as expected seldomly shared. Data mining of CDR3 regions within identified δ TCR sequences and public data bases elucidated further that highly therapeutically interesting receptor chains such as the by us recently described chain of clone 5 from δ 2-positive δ TCRchains (11) was found among the dominant top 20 clones of HD15. Moreover, the δ 2-negative δ TCR chain of clone Fe11(9) was the second most prevalent clone in HD19. Main advantage is that such a strategy narrows down very quickly the potential very high diversity of δ TCR chains from 10^1 to litteraly less than 10^2 sequences of therapeutic interest. This has practical and therapeutical consequences as high through put screening of δ TCR chain sequences followed by the identification of shared sequences in healthy and diseased patients can serve as rapid hub for the identification of novel therapeutic tools which can be used for e.g. T cells engineered with defined $\gamma\delta$ TCRs, so called TEGs (13). In addition, repertoire analyses of diseased patients has the power to rapidly identify patients who might benefit from the addition of an immune repertoire with defined δ TCR sequences.

TRD Cl3 Amino acid sequence: SEQ ID NO:13

m e r i s s l i h l s l f w a g v m s a i e l v p e h q t v p v s i g v p a
 t l r c s m k g e a i g n y y i n w y r k t q g n t m t f i y r e k d i y g
 5 p g f k d n f q g d i d i a k n l a v l k i l a p s e r d e g s y y c a c d
 l l g y t d k l i f g k g t r v t v e p r s q p h t k p s v f v m k n g t n v
 a c l v k e f y p k d i r i n l v s s k k i t e f d p a i v i s p s g k y n a
 v k l g k y e d s n s v t c s v q h d n k t v h s t d f e v k t d s t d h v
 k p k e t e n t k q p s k s c h k p k a i v h t e k v n m m s l t v l g l r
 10 m l f a k t v a v n f l l t a k l f f l

TRG Cl3 Amino acid sequence: SEQ ID NO:14

m v s l l h a s t l a v l g a l c v y g a g h l e q p q i s s t k t l s k t a
 r l e c v v s g i t i s a t s v y w y r e r p g e v i q f l v s i s y d g t v r
 15 k e s g i p s g k f e v d r i p e t s t s t l t i h n v e k q d i a t y y c a l
 w e e e l g k k i k v f g p g t k l i i t d k q l d a d v s p k p t i f l p s
 i a e t k l q k a g t y l c l l e k f f p d v i k i h w e e k k s n t i l g s
 q e g n t m k t n d t y m k f s w l t v p e k s l d k e h r c i v r h e n n
 k n g v d q e i i f p p i k t d v i t m d p k d n c s k d a n d t l l l q l t
 20 n t s a y y m y l l l l l k s v v y f a i i t c c l l r r t a f c c n g e k s

TRD Cl5 Amino acid sequence: SEQ ID NO:15

m e r i s s l i h l s l f w a g v m s a i e l v p e h q t v p v s i g v p a
 t l r c s m k g e a i g n y y i n w y r k t q g n t m t f i y r e k d i y g
 25 p g f k d n f q g d i d i a k n l a v l k i l a p s e r d e g s y y c a c d a
 l k r t d t d k l i f g k g t r v t v e p r s q p h t k p s v f v m k n g t n
 v a c l v k e f y p k d i r i n l v s s k k i t e f d p a i v i s p s g k y n
 a v k l g k y e d s n s v t c s v q h d n k t v h s t d f e v k t d s t d h
 v k p k e t e n t k q p s k s c h k p k a i v h t e k v n m m s l t v l g l
 30 r m l f a k t v a v n f l l t a k l f f l

TRG Cl5 Amino acid sequence: SEQ ID NO:16

m v s l l h a s t l a v l g a l c v y g a g h l e q p q i s s t k t l s k t a
 r l e c v v s g i t i s a t s v y w y r e r p g e v i q f l v s i s y d g t v r
 35 k e s g i p s g k f e v d r i p e t s t s t l t i h n v e k q d i a t y y c a l
 w e i q e l g k k i k v f g p g t k l i i t d k q l d a d v s p k p t i f l p
 s i a e t k l q k a g t y l c l l e k f f p d v i k i h w e e k k s n t i l g
 s q e g n t m k t n d t y m k f s w l
 t v p e k s l d k e h r c i v r h e n n k n g v d q e i i f p p i k t d v i t
 40 m d p k d n c s k d a n d t l l l q l t n t s a y y m y l l l l l k s v v y f
 a i i t c c l l r r t a f c c n g e k s

TRD Fe11 Amino acid sequence: SEQ ID NO: 17

m v f s s l l c v f v a f s y s g s s v a q k v t q a q s s v s m p v r k a
 45 v t l n c l y e t s w w s y y i f w y k q l p s k e m i f l i r q g s d e q
 n a k s g r y s v n f k k a a k s v a l t i s a l q l e d s a k y f c a l g d
 s y g g g p l y t d k l i f g k g t r v t v e p r s q p h t k p s v f v m k
 n g t n v a c l v k e f y p k d i r i n l v s s k k i t e f d p a i v i s p s
 g k y n a v k l g k y e d s n s v t c s v q h d n k t v h s t d f e v k t d
 50 s t d h v k p k e t e n t k q p s k s c h k p k a i v h t e k v n m m s l t
 v l g l r m l f a k t v a v n f l l t a k l f f l

67

TRG Fe11 Amino acid sequence: SEQ ID NO: 18

m g w a l l v l l a f l s p a s q k s s n l e g g t k s v t r p t r s s a e i t
c d l t v i n a f y i h w y l h q e g k a p q r l l y y d v s n s k d v l e
s g l s p g k y y t h t p r r w s w i l i l r n l i e n d s g v y y c a t w d
5 r p e i y y k k l f g s g t t l v v t d k q l d a d v s p k p t i f l p s i a e
t k l q k a g t y l c l l e k f f p d v i k i h w e e k k s n t i l g s q e g
n t m k t n d t y m k f s w l t v p e k s l d k e h r c i v r h e n n k n g
v d q e i i f p p i k t d v i t m d p k d n c s k d a n d t l l l q l t n t s
a y y m y l l l l l k s v v y f a i i t c c l l r r t a f c c n g e k s
10

Example 2**Material & Methods****5 Patient, sample collection and cell lines**

All donor and patient material were collected according to GCP and Helsinki regulations. Peripheral blood mononuclear cell (PBMC) samples from healthy donors and acute myeloid leukemia (AML) patients were from the University Medical Center Utrecht Biobank. The use of the TNBC patient samples was approved by the ethics committee of the University of Freiburg Medical Center. From the archive of the Institute of Clinical Pathology, Freiburg we selected 16 formalin-fixed paraffin-embedded tissue specimens with the diagnosis of “TNBC”. The histopathological diagnosis was performed according to the Union for the International Cancer Control (UICC) criteria. All the tumors were Grade III in the modified Bloom-Richardson classification (Elston). Conforming to the recommendations for the evaluation of TILs (14 15), the H&E stained samples contained at least 50% tumor infiltration. The cohort of 11 tumors (5 medullary breast carcinoma’s and 6 invasive ductal carcinoma’s) that we studied further was selected on the basis that the correspondent frozen tissue samples were available in the tumor bank of the Comprehensive Cancer Center Freiburg (CCCF). All these samples were classified as basal like subtype according to the expression of CK 5/6 or 14 and EGFR (16). The median age of the patients was 59 years in a range between 43 and 82 years.

Breast cancer cell lines and frozen tumor material were tested for HCMV using IHC nested PCR according to the protocol previously published by Bender et al (17), and real-time PCR with the artus® CMV TM PCR kit (Qiagen, Hilden, Germany) in a 7900HT Fast real time PCR cycler (Applied Biosystems, CA, USA) according to the manufacturer instructions.

Next generation sequencing of TCR δ and TCR β repertoire

30 The protocol is adapted from Mamedov et al with modifications (18). Frozen PBMCs from healthy donors were thawed and stained with the following antibodies: CD3 eFluor 450, TCR $\alpha\beta$ APC, and TCR $\gamma\delta$ PE monoclonal antibodies (mAb). (Table 6. Samples were sorted on the ARIAII (BD) in an $\alpha\beta$ -fraction (CD3⁺, $\alpha\beta$ ⁺), and a $\gamma\delta$ -fraction (CD3⁺, $\gamma\delta$ ⁺). RNA was isolated using Qiagen Rneasy Minikit for samples

$\geq 0.5 \times 10^6$ cells or Qiagen Rneasy Microkit for samples $< 0.5 \times 10^6$ cells following manufacturer's instructions. Specific cDNA for TCR δ and TCR β was synthesized with Superscript® II Reverse Transcriptase (ThermoFisher), using a specific primer at the 3' constant region and an universal template switch adaptor was incorporated at the 5' end of the V region. cDNA was purified using NucleoSpin® Gel and PCR Clean-UP (Machery-Nagel) thereafter amplified with a first PCR amplification using Q5® High Fidelity DNA polymerase (New England Biolabs), on a T100 Thermal Cycler (Biorad) using the following steps 90 s at 98°C, 15 cycles of 7 s at 98°C, 20 s at 62°C, 50 s at 72°C, followed by 10 m at 72°C. A specific nested primer at the constant region, and a step-out primer which anneals to the switch adaptor were used (Table 5). PCR products were loaded on a 1.5% agarose gel for size selection of products between 400-600 base pairs. After purification of the gel with NucleoSpin® Gel and PCR Clean-UP (Machery-Nagel), the PCR product was used for a second PCR with a reverse nested primer on the constant region and a forward primer which annealed on the switch adaptor (Table 5), using the following steps: 90 s at 98°C, 20 cycles of 7 s at 98°C, 20 s at 62°C, 50 s at 72°C, followed by 10 m at 72°C. After purification with NucleoSpin® Gel and PCR Clean-UP (Machery-Nagel), the final PCR products were analyzed with QIAxcel Advanced System (Qiagen). Library preparation for NGS was done with NGSgo-LibrX kit with NGSgo-IndX indices from Gendx according to recommendations of the manufacturer. Cleanup of the samples was performed with HighPrep PCR beads from GC Biotech. Next-generation sequencing was performed on an Illumina MiSeq system 500 (2x250 bp) (Illumina). Sequencing data were analyzed with the MiXCR program (Bolotin, et al. (18)) and VDJ-tools for further analyses.

Immunohistochemistry (IHC) and image analysis of TNBC samples

Serial FFPE 2 μ m sections mounted on Superfrost plus glass slides (R Langenbrink, Germany), were dewaxed and rehydrated. After the proper antigen retrieval in a pressure cooker with citrate buffer (pH 6) and citrate buffer (pH 6,1) (Dako, Hamburg, Germany), blocking of nonspecific binding was performed using goat serum (5% in PBS). Mouse monoclonal anti-TCR γ -chain mAb and rabbit-anti-human cleaved caspase 3 (cC3) polyclonal antiserum (Table 6) as we previously reported (19). The HCMV detection was performed using the mouse anti-CMV mAb. Alkaline phosphatase-conjugated and Horseradish peroxidase-conjugated detection systems

were used to visualize the primary antibodies in a separate or sequential protocols for single or double staining test with red and brown chromogen (DakoREAL™ Alkaline Phosphatase/RED r/m and EnVision™ FLEX Systems, Dako, USA). Acidic hematoxylin was used as a counterstain.

- 5 IHC samples were analyzed and tiled scanned using an AxioObserver Z1 with Apotome2 system with an ERc5s digital camera. The initial analysis was performed using the AxioVision 4.8 and ZEN BLUE image software (all from Carl Zeiss MicroImaging, Jena, Germany). Colocalization and quantitate analysis were performed with ImageJ (NIH images, USA) and QuPath (GitHub, San Francisco CA, USA)
- 10 software with Bio-Formats, Stack Slicer and Cell counter plugins (20).

Immunofluorescence imaging of TNBC samples

- Sections at 5 µm were mounted on Superfrost plus Adhesion glass slides, air-dried for 3 hours and fixed with pre-cooled acetone (-20°C) for 10 minutes. Samples were rinsed
- 15 with TBST for 5 minutes (3X) and blocking for nonspecific binding using normal human serum 5% in PBS for 30 min. The samples were incubated with corresponding primary mouse-anti-human mAbs: anti-TCRγ and anti-CD69 mAb or goat-anti-human polyclonal anti-IFN- γ and anti-IL-17 antisera. Fluorescence-conjugated secondary antibodies used to visualized the primary antibodies were rabbit-anti-mouse AlexaFluor
- 20 488, donkey-anti-rabbit AlexaFluor 568, donkey-anti-goat AlexaFluor 594, donkey-anti-goat AlexaFluor 647 (Table 6). Samples were mounted using the Prolong® Diamond Antifade medium with DAPI (Thermo Fischer).

Tissue immunostaining and laser capture microdissection of TNBC samples

- 25 Frozen sections (8 µm thick) were air dried overnight on MembraneSlide 1.0 PEN™ membrane covered slides (Carl Zeiss, Munich, Germany), fixed in pre-cooled acetone (-20°C) for 10 min, washed twice with TBST and incubated for 30 min with 5% human serum in PBS. The samples were incubated for 30 min at room temperature with the mouse anti-human TCRγ mAb (Table 6). To detect the primary antibody, a biotinylated
- 30 anti-mouse secondary antibody with alkaline phosphatase detection system (Dako REAL™ Detection System Alkaline P/RED, rabbit-mouse). Mayer's hematoxylin was used as counterstain. Samples were dried at room temperature for 2 hours, examined under microscopy and stored at 4°C until processing.

Laser microdissection was performed using an Axiovert microscope equipped with a PALM MicroBeam system (ZEISS Microscopy, Oberkochen, Germany). Energy parameters of cutting and catapulting were established individually for each sample. Only infiltrating single lymphocytes in close contact with cancer cells were selected at 40x magnification and microdissected at 63X magnification. Single cells were catapulted into the cap of an Adhesive Cap 500 opaque™ 500 µl tube (Carl Zeiss, Göttingen, Germany). To improve isolation, the cap contained 5 µl of 1X Expand High Fidelity PCR buffer w/o MgCl₂ (Roche, Mannheim, Germany). Then, 15 µl of a 1:10 mix containing PCR buffer and Proteinase K (20 mg/ml, PCR grade, Roche, Mannheim, Germany) were added for the digestion. The tubes were incubated with the lid down for 4 h at 56°C, centrifuged for 2 min at 500 rpm and heat inactivated at 95°C for 10 min. Additional tubes containing only membrane were dissected from each sample and used as negative controls. All PCR tubes were overlaid with mineral oil under a laminar flow hood before adding the PCR master mix.

15

Single-cell PCR of TNBC γδ TILs

DNA isolation, the first and the second rounds of PCR were performed in different “single-cell rooms” under laminar flow cabinet previously decontaminated with UV lamp, ethanol and DNA Zap (Invitrogen, Bleiswijk, Netherlands). Separate sets of pipettes, consumables and reagents were used for each step and every two weeks all reagents were tested to prevent contamination. Corresponding to the single-cell PCR technique previously described for the analysis of rearranged immunoglobulin genes (21), a multiplex, semi-nested, hot-start PCR was prepared with 15 newly designed primers (Table 7). For the first round, a master mix was prepared with 5 µl dNTP (2 mM), 5 µl 10X PCR buffer (High Fidelity System), 5 µl primer mix (2.5 µM, forward and reverse primers), 3.2 µl 25 µM MgCl₂, 6.5 µl H₂O and 15 µl from the DNA digestion. A volume of 0.3 µl Expand High Fidelity enzyme mix (3.5 units/µl) was added after the first denaturation step to a final volume of 40 µl. The cyclor program was 95°C 2 min, 80°C pause (enzyme added), 72°C 1 min, 39x (95°C 50 s, 56°C 30 s, 72°C 60 s), 72°C 5 min and 10°C pause. For the second round, eight master mixes were prepared to detect TCRγ and δ chains: two mixes for TCRγ (Vγ1-8 and Vγ9) and six for TCRδ (Vδ1, Vδ2, Vδ3, Vδ4, Vδ5 and Vδ6) (Suppl. Table 4) with 2.5 µl dNTP (2 mM), 2.5 µl 10X PCR buffer, 1.25 µl of the respective Vγ and Vδ forward primers,

1.25 μ l of the respective joint mix primers, 2 μ l 25 mM $MgCl_2$, 12.2 μ l H_2O , 3 μ l of first round PCR product and 0.3 μ l Expand High Fidelity enzyme mix (3.5 units/ μ l). The cycler program was 95°C 5 min, 72°C 1 min, 35x (95°C 50 s, 55.5°C 30 s, 72°C 1 min), 72°C 5 min, 15°C 5 min and 4°C pause. The PCR products were analyzed by 2% agarose gel electrophoresis and the positive bands were cut under UV light and purified from the gel with the Qiaex II gel extraction kit (Qiagen, Hilden, Germany). The clean DNA was sequenced using the BigDye Terminator 3.1 system (Applied Biosystems, CA, USA), the sequencing reactions consisted of 1 μ l BigDye, 3.75 μ l 5X sequencing buffer, 0.75 μ l forward primer, 3-10 μ l template and water to a final volume of 20 μ l. The cycler conditions were 96°C 5 min, 24x (95°C 15 sec., 50°C 10 sec., 60°C 4 min), 10°C pause. The sequence sample was cleaned using the DyeEx 2.0 Spin Kit (Quiagen) and analyzed on the ABI 3130XL capillary sequencer (Applied Biosystems, Darmstadt, Germany). The sequences we compared and evaluated with the IMGT database (<http://www.imgt.org/>) and the IgBlast tool (<https://www.ncbi.nlm.nih.gov/igblast/>)

Retroviral expression of plasmids from TNBC $\gamma\delta$ TILs

Codon optimized DNA coding for the full length γ and δ TCR chains of the tumor infiltrating lymphocytes and for full length EPCR was ordered at Baseclear Inc. The synthetic genes were flanked with a 5' NcoI and a 3' BamHI site for subsequent cloning into a pBullet retroviral expression vector. The γ TCR genes were subcloned into pBullet-IRES-*neo* and the δ TCR genes as well as EPCR were subcloned into pBullet-IRES-puromycin.

25 Transduction of $\alpha\beta$ T cells.

For the generation of TEGs, PBMCs were transduced defined TCR γ and δ chains as described (22-24). In short, retroviral supernatant was produced by Phoenix-Ampho packaging cells, that were transfected with gag-pol (pHIT60), env (pCOLT-GALV) and pBullet retroviral constructs containing TCR γ or δ , using Fugene HD (Promega). PBMCs preactivated with α CD3 (30 ng/ml) (clone OKT3, Janssen-Cilag) and IL-2 (50 U/ml) were transduced twice with viral supernatant within 48 hours in the presence of 50 U/ml IL-2 and 4 μ g/ml polybrene (Sigma-Aldrich). Transduced T cells were expanded by stimulation with α CD3/CD28 Dynabeads (0.5×10^6 beads/ 10^6 cells)

(Invitrogen) and IL-2 (50 U/ml) and selected with 800 µg/ml geneticin (Gibco) and 5 µg/ml puromycin (Sigma-Aldrich) for one week. Following transduction, transduced T cells were stimulated biweekly with 1 µg/ml PHA-L (Sigma-Aldrich), 50 U/ml IL-2 (Novartis Pharma), 5 ng/ml IL-15 (R&D Systems), and irradiated allogeneic PBMCs, Daudi and LCL-TM cells. Fresh IL-2 was added twice a week. Transgenic TCR expression was routinely assessed by flow cytometry.

Functional assays of primary clones and TEG's

IFN γ ELISPOT was performed as previously described (25, 26). Briefly, 15,000 TCR-transduced or mock-transduced T cells and 50,000 target cells (ratio 0.3:1) were co-cultured for 24 hrs in nitrocellulose-bottomed 96-well plates (Millipore) precoated with anti-IFN γ antibody (clone 1-D1K) (Mabtech). Plates were washed and incubated with a second biotinylated anti-IFN γ antibody (clone 7-B6-1) (Mabtech) followed by streptavidin-HRP (Mabtech). IFN γ spots were visualized with TMB substrate (Sanquin) and the number of spots was quantified using ELISPOT Analysis Software (Aelvis). IFN γ ELISA was performed using ELISA-ready-go! Kit (eBioscience, San Diego, CA, USA) following manufacturer's instructions. Effector and target cells (E:T 1:1) were incubated for 24h in the presence of pamidronate when indicated.

CD1 Tetramer staining

CD1c and CD1d tetramers were produced as described before (27). TEGs were stained with anti- $\gamma\delta$ TCR-APC antibody (Table 5) and CD1 streptavidin-PE tetramers at a concentration of 50 nM in PBS + 0.5% BSA for 30' at room temperature. After two wash steps the cells were analyzed by flow cytometry on a BD FACSCanto II.

25

Mouse model for TEG001

The NOD.Cg-Prkcd^{scid} IL2rg^{tm1Wjl}/SzJ (NSG) mice, originally obtained from JAX (Bar Harbor, ME, USA), were bred and housed in the specific pathogen-free (SPF) breeding unit of the Central Animal Facility of Utrecht University. Experiments were conducted according to Institutional Guidelines after acquiring permission from the local Ethical Committee and in accordance with current Dutch laws on Animal Experimentation. For the experiments mice from 8 to 12 weeks of age were used. At day 0 mice received sublethal total body irradiation (175 cGy) followed by i.v. injection of 5x10⁶ OPM2-

30

Luc tumor cells at day 1. Mice were treated with 10^7 TEG001 cells or Mock TCR transduced T cells i.v. at day 7 and 14. Mice received 0.6×10^6 IU of IL-2 in IFA s.c. on day 7 and every 21 days till the end of the experiment. Pamidronate (10 mg/kg body weight) was applied at day 7 i.v. and every 21 days until the end of the experiment.

- 5 Tumors were visualized *in vivo* by bioluminescent imaging. Mice were anesthetized by isoflurane before they received an i.p. injection (100 μ l) of 25 mg/ml Beetle Luciferin (Promega, Madison WI). Bioluminescence images were acquired by using a third generation cooled GaAs intensified charge-coupled device camera, controlled by the Photo Vision software and analyzed with M³Vision software (all from Photon Imager;
- 10 Biospace Laboratory, Paris, France).

Results

Identification of shared TCR δ sequences with next generation sequencing.

In order to understand the functional implication of shared immune receptors within the TCR δ repertoire we analyzed TCR δ chains from 14 different healthy donors (dataset 1). After sorting $\gamma\delta$ T cells by flow cytometry, followed by TCR δ repertoire analyses by NGS, we found 9011 amino acids sequences of the TCR δ CDR3 (CDR3 δ) region. Further analysis on the most abundant sequences per donor, defined as sequences with a clonal frequency of $>0.1\%$ ($n=1478$), showed that 1.8% of sequences were shared between at least two donors. Of these shared sequences one was shared between three different donors (Table 8). Overlap of TCR δ sequences between 2 different donors differed from 0 to 3.9% of the sequences of the 2 donors (Figure 5A). In contrast in the TCR β repertoire only 0.7% of sequences were shared. We identified 6 shared sequences, and the overlap between 2 different donors ranged from 0 to 1.9% (Figure 5B). As expected for peripheral blood repertoires the majority of the identified sequences was V δ 2+ (Figure 6A). In line with this the majority of the shared sequences was also V δ 2. In addition we identified four shared V δ 1 sequences.

Analyses of TCR δ repertoires in the tumor infiltrating tissues

Immune repertoires of $\gamma\delta$ T cells in healthy individuals between peripheral blood and tissues differ substantially (28), although it is reported that TCR δ sequences identified in tissue could be traced back in the peripheral blood (28). To analyze whether the same holds true for $\gamma\delta$ tumor infiltrating lymphocytes ($\gamma\delta$ TILs) we compared the sequences of the peripheral TCR δ repertoire (dataset 1-4) with a fifth dataset consisting of published $\gamma\delta$ TIL TCR δ sequences and a publicly available dataset of Li et al. The dataset of Li et al consisted of CDR3 sequences of TILs of various tumors with 1060 complete CDR3 δ sequences. In line with the assumption that V δ 2^{neg} $\gamma\delta$ T cells are the dominant fraction in tissues, only 19% of TCR δ sequences were classified as V δ 2 TCR δ . The majority of the sequences were of V δ 1 origin, but compared to the healthy repertoires also the V δ 3, V δ 5, V δ 7 and V δ 8 fraction were increased. Although the percentage of V δ 2 TCR δ sequences in tumor tissue is relatively low, still 24 shared V δ 2 sequences could be identified (figure 5C). Interestingly, once peripheral T cells have been heavily enriched for V δ 1 $\gamma\delta$ T cells before NGS analyses (dataset 3) 10 V δ 1 sequences could be characterized as shared between the peripheral blood and $\gamma\delta$ TILs

(Figure 5D). Although these data demonstrate that it is possible to identify shared sequences between $\gamma\delta$ TILs and the peripheral blood, it also illustrates the limitation of this method because of the different composition of the TCR δ repertoire between peripheral blood and cancer tissues.

5

Single cell analyses of $\gamma\delta$ TILs in patients with triple negative breast cancer

In order to further analyze in depth $\gamma\delta$ TIL receptor sequences between individual cancer patients at the tumor site a previously characterized TNBC cohort was used. $\gamma\delta$ TILs in TNBC have been associated with elevated levels of the apoptotic marker cleaved caspase 3 at the tumor site (29). In order to identify $\gamma\delta$ TILs with close proximity to apoptotic tumor cells paraffin embedded tissues were co-stained for TCR $\gamma\delta$ and cleaved caspase-3 using IHC. $\gamma\delta$ TILs were observed at high frequency in all of the examined biopsies and apoptotic tumor cells were in contact with $\gamma\delta$ T cells (Figure 7A). Expression of CD69, as evident by IHC (Figure 7B), suggested a T cell receptor-mediated activation of $\gamma\delta$ TILs. Conflicting roles of $\gamma\delta$ TILs have been described such as cancer immune surveillance (30) or tumor promoting properties through IL-17 (31). Therefore, IFN γ and IL-17 expression was further assessed by IHC. Most $\gamma\delta$ TILs were IFN γ^{pos} and did not substantially stain for IL-17, suggesting that indeed investigated $\gamma\delta$ TILs are activated with a Th1-type and that $\gamma\delta$ T cells are an active participant of daily cancer immune surveillance.

20

Diversity and sharing of TCR γ and δ chains derived from $\gamma\delta$ TILs

Diversity and clonal expansion of $\gamma\delta$ TILs in individual TNBC patients was first analyzed by spectratyping of V γ and V δ genes. These data confirmed that also tumor infiltrating $\gamma\delta$ T cells V γ and V δ gene usage is polyclonal in most patients. These data were also confirmed for five of the breast cancer samples by NGS of the delta sequences (dataset 7, Figure 6B). However, with a mean of 68.9%, the V δ 2 T cell proportion appeared higher in the NGS breast cancer data than observed for other tumor tissues (Figure 6B). This is most likely due to the strong vascularization of investigated tissues. The V δ 3 and V δ 5 gene were again enriched in tumor tissue, suggesting that these subsets might be functionally important in cancer immune surveillance. Next we searched for public TCR δ sequences in the NGS breast cancer data. 52 V δ 2 sequences were shared with the healthy donor datasets and 3 sequences

25

30

were shared with the $\gamma\delta$ TIL dataset (figure 5C). Interestingly, a V δ 3 sequence which was also present in the $\gamma\delta$ TIL sequences, was also found within this dataset of breast cancer TILs (Figure 5E). No further V δ 1 shared sequences were observed (figure 5D). In summary, analysis of this rather unique and homogenous cohort indicates that the $\gamma\delta$ T cell repertoire in tumor tissue is quite diverse, and that despite some increased frequency for V δ 3 and V δ 5 in our TNBC cohort, but also in a dataset of $\gamma\delta$ TILs (32) of Li et al., no substantial clonal expansion can be observed among $\gamma\delta$ TILs.

Next, we aimed to assess whether shared individual γ or δ TCR chains were involved in a cognate $\gamma\delta$ T cell tumor interaction. Therefore, single cells were isolated from frozen TNBC sections using laser microdissection and transferred to a PCR tube to determine the TCR sequence using single cell sequencing. In total, 530 single $\gamma\delta$ T cells were isolated from 11 different tumors and 27 paired TCR γ and δ sequences from 9 different patients could be identified. (dataset 6, Table 9). The other single cell sequencing reactions did not result in reliable sequencing data for both the γ and/or δ TCR chain, however we were able to determine 63 additional non-paired γ and 28 additional non-paired δ CDR3 sequences. The single cell sequencing data again confirmed a polyclonal population of tumor infiltrating $\gamma\delta$ T cells, within patients some TCR δ and TCR γ sequences were seen multiple times. As expected from our analyses of TILs across many cancer patients, as well as our analysis tumor infiltrating $\gamma\delta$ T cells in patients suffering from TNBC. Non-V γ 9 and non-V δ 2 genes, were most prominent in the sequences obtained. The V δ 2 gene was used in 7 out of the 55 TCR δ sequences and the V γ 9 gene in 21 out of the 90 TCR γ sequences. Interestingly, 4 of the 27 clones used an identical V δ 5 sequence paired with a different γ TCR sequence (Table 9). Additionally, this V δ 5 sequence was also identified in 6 non-paired δ CDR3 sequences (Figure 8A). This sequence was found in 6 out of 11 patients, which classifies it as a public CDR3 sequence. This particular sequence has also been reported to associate with CMV- and tumor-reactivity (33). However, surprisingly none of the patients that had this particular shared TCRV δ 5 sequence were CMV positive. Two V δ 1 sequences were shared between 2 different patients. As additional quality control we could retrieve four of the single cell TRD sequences within NGS dataset from TNBC: three sequences of patient E and one sequence of patient F, indicating that many receptors

picked up by single cell sequencing might be of very low frequency and therefore not visible through NGS of TCR δ repertoires.

The TCR γ chain was more frequently shared between the isolated T cells out of the 90
 5 obtained sequences in total, 10 different sequences were shared amongst the patients. One sequence was shared between 5 patients and was identified in 10 different isolated cells. This sequence was found back in 4 of the 27 paired $\gamma\delta$ clones. One other CDR3 γ sequence was shared between 3 patients and some others were shared between two patients (Figure 8B). Additionally, there was a T cell that shared the γ TCR chain
 10 reported with a previously identified CD1d-restricted T cell clone (34). In summary, many shared TCR γ chains and V δ 2^{neg} TCR chains can be found within the local tumor microenvironment, however in contrast to $\gamma\delta$ T cell repertoires in the peripheral blood, repertoires are dominated by V δ 2^{neg} $\gamma\delta$ T cells and lack a substantial clonal expansion.

15 **Functional assessment of selected TCR $\gamma\delta$ from the peripheral and tumor infiltrating $\gamma\delta$ T cell repertoire**

In order to test our hypothesis that either shared γ or shared δ TCR chains from the peripheral and tissue residing immune repertoire functionally contribute to daily cancer immune surveillance, we completed first the corresponding TCR γ for selected shared
 20 TCR δ from the peripheral repertoire. This was done by single cell cloning for V δ 2⁺ (23) or V δ 2^{neg}(25) $\gamma\delta$ T cells and we sequenced both, γ and δ TCR chains. We identified a corresponding γ chain of the shared V δ 2TCR chain, Clone 5 as reported recently (23). The TCR δ chain of the TCRV γ 9 δ 2 expressing clone 5 was found in healthy donor 11 of dataset 1. This particular combination has been reported to be very
 25 active against many types of hematological and solid cancer cells (22, 23). In order to further extend analyses on activity of the TCR γ and δ chains isolated from clone 5 in comparison to bulk $\gamma\delta$ T cells $\alpha\beta$ T cells were engineered to express this defined TCR $\gamma\delta$ (TEG001) (23, 25, 26). TEG001 showed not only superior activity to different tumor cell lines when compared to bulk $\gamma\delta$ T cells (Figure 9A) but also to a variety of
 30 primary AML blasts (Figure 9B). These data suggest that the TCR V γ 9 δ 2 utilizing shared TCR δ chains can mediate a high functional activity and functionally outperform most $\gamma\delta$ T cells.

To further functionally characterize shared TCR δ chains from V δ 2^{neg} T cells from the peripheral repertoire healthy donors, V δ 2^{neg} $\gamma\delta$ T cells were isolated and cloned as reported (25) from an additional healthy donors, tested for reactivity by an IFN γ ELISPOT assay against a defined tumor cell panel, and sequenced for the

5 corresponding γ and δ TCR chain. From the 10 isolated V δ 2^{neg} $\gamma\delta$ T-cell clones, 3 clones showed reactivity against different cancer cell lines (Figure 10A). Sequencing of all functionally investigated clones identified clone FE11 as a clone, which shares its δ 1 TCR chain also with donor 13 and 19 of dataset 1. In order to investigate whether tumor reactivity of FE11 is mediated by the TCR $\gamma\delta$ chain only, we introduced γ and δ

10 TCR chains of clone FE11 into $\alpha\beta$ T cells (TEG-011), and tested activity of FE11 against a panel of tumor cell lines by an IFN γ ELISA assay, and confirmed the results as seen with the ELISPOT assay (Figure 10A). In summary, a variety of V δ 2^{neg} $\gamma\delta$ T-cell clones can be identified from the peripheral blood, which recognize in a complementary way different tumor cell lines. In addition, clone FE11 with a shared

15 TCR δ shows tumor reactivity.

In order to functionally confirm that also the TCR $\gamma\delta$ from $\gamma\delta$ TILs have the potential to mediate tumor reactivity and that this reactivity is at least partially mediated by their individual shared TCR γ or δ chains, we generated a series of 15 TEGs by expressing

20 paired γ and δ TCR chains derived from our breast cancer $\gamma\delta$ TILs (Table 9). In line with the observed reactivity of V δ 2^{neg} $\gamma\delta$ T cell clones from the peripheral blood 5 out of 10 TEGs with shared γ or δ TCR chains showed complementary reactivity against different types of cancer cell lines. One out of 15 TCR $\gamma\delta$ s with a shared receptor sequence showed a very broad reactivity, namely TEG-Zi11, while the 5 other clones

25 had a more restricted reactivity against a distinct cancer cell lines within the tested panel (Figure 10B). Notably, the broadly reactive TCR $\gamma\delta$ complex of TEG-Zi11 uses a shared γ chain with TEG-B23 and TEG-Ze11 which are both unresponsive to tested cancer cell lines, indicating in this case that the unique δ chain of TEG-Zi11 is required for the tumor-reactivity. In line with this observation, the TCR $\gamma\delta$ complex from TEG-

30 F4 which has a restricted reactivity to some tumor cell lines shared its γ chain with the non-reactive TCR from TEG-E20, and TCR $\gamma\delta$ complex from TEG-B9, which shares its γ chain with a previously reported CD1d-reactive TCR $\gamma\delta$ (34), did not bind to CD1d tetramer when the reported unique TCR δ chain was not present (Figure 12). Thus, a

shared TCR γ chain can be part of a tumor-reactive $\gamma\delta$ TCR, but that in itself is not sufficient for reactivity. Vice versa, TEG-C132, TEG-D37, TEG-E113 and TEG-B23, who share the very same δ 5 chain, were active against different cancer cell lines, namely K562 for TEG-B23, TEG-D37 and TEG-C132, U937 for TEG-B23, TEG-D37 and TEG-E113 (Figure 10B). Thus, TCR $\gamma\delta$ complexes comprised of shared V δ 2 and V δ 2^{neg} chains have a comprehensive and complementary ability to attack different types of cancer cells and chains comprised of V δ 5 could be in particular important participants in cancer immune surveillance.

10 **In vivo activity of public TCR δ chains within the therapeutic concept of TEGs**

Engineering immune cells for cancer immune therapy has been recognized as scientific breakthrough (35) and the next generation of chimeric antigen receptors (CARs) is urgently needed to overcome current limitations such as restricted tumor antigen availability (36). TEGs have most recently been appreciated as a new interesting twist on this strategy (36) and therefore we tested whether not only artificial TCR $\gamma\delta$ complexes (22, 23) but also natural TCR $\gamma\delta$ complexes with partly public δ TCR sequences can be part of the next generation of TEGs. Therefore, TEG001 activity was tested in vivo in a multiple myeloma mouse model utilizing OPM2 as target (22). As negative control TEGs expressing a non-functional TCR $\gamma\delta$ chain LM1 (23) were used. TEGs were infused 7 and 14 days after inoculation of OPM2. Treatment with TEG001 lead when compared to TEG-LM1 to a significant inhibition of tumor growth and associated in this treatment model also with a significant prolonged overall survival (Figure 11), indicating that TCR $\gamma\delta$ complexes comprised of partly public TCR chains are a valuable tool for cancer immune therapy with TEGs.

25

Discussion

Three types of immune cells undergo somatic DNA recombination during their ontogeny. Numerous functions of two of them, $\alpha\beta$ T cells and B cells, during immune responses have been elucidated to a substantial extent. In contrast, relatively very limited information is available about the repertoire and immunological function of the third type of cells undergoing somatic DNA recombination during their ontogeny, $\gamma\delta$ T cells. Here we attempted to illuminate this unknown territory by analyzing the $\gamma\delta$ TCR repertoires using combined high-throughput next generation and single-cell sequencing

techniques and found evidence of the involvement of public V δ 2⁺ and V δ 2^{neg} TCR chains in cancer immune surveillance.

While public β TCR chains have been reported to be mainly involved in infection and
5 allo-reactivity, our data suggest that public γ and δ TCR pairs are an important pillar of
daily cancer immune surveillance. Shared V δ 2 TCRs from the peripheral blood have
the ability to mediate highest functional activity against various tumor types and can
even outperform tumor reactivity of polyclonal $\gamma\delta$ T cells, as shown in the in vitro
experiments with TEG001. In addition also V δ 2^{neg} TCRs can mediate strong anti-tumor
10 reactivity against a set of different tumor cells, thus seem to cover surveillance against
a variety of solid and hematological malignancies. However, in depth analyses of $\gamma\delta$
TILs indicate that $\gamma\delta$ TILs with potentially tumor reactive $\gamma\delta$ TCR lack substantial
clonal proliferation within tumor tissues. In addition, to this basic biological
observation which indicates that public γ and δ TCR chains are an important but also
15 vulnerable part of early cancer immune surveillance, identified receptors are a valuable
tool for novel therapeutic compounds like TEGs.

Although clonal expansions of $\alpha\beta$ TILs have been described (38), no substantial clonal
dominance was observed within TNBC's $\gamma\delta$ TIL populations when analyzed by
20 spectratyping, NGS or single-cell PCR during our current study. This observed absence
of significant clonal expansions in TNBC's $\gamma\delta$ TILs is starkly contrasting the very
skewed TCR δ 1 repertoires in the adult peripheral blood (39, 40, 25) or in fetal CMV
infections *in utero* containing several high-frequency clonotypes (31). Interestingly,
proliferation proficiency of V δ 2⁺ cells noticed in infectious challenges (41), is
25 frequently absent in cancer patients (39). Also, elegant mouse model studies indicate
that $\gamma\delta$ T cells can prevent cancer development in the early disease stages (42, 43), but
once the disease becomes more advanced, $\gamma\delta$ TILs become rapidly proliferation-
deficient, while maintaining other functional activities such as IFN γ or CD69
upregulation. Since we could not detect any $\gamma\delta$ TILs with the Th17 phenotype, it's
30 possible that the tumor microenvironment may be particularly inhibitory for this subset.
The assumption of an early proliferation deficiency of $\gamma\delta$ TILs would also favor single
cell PCR instead of NGS for the identification of relevant receptors within the context
of $\gamma\delta$ TILs. However, a study as ours is technically challenging, and currently it is

difficult to perform single cell sequencing analysis of tumor tissue in a high throughput manner.

Our data are compatible with the earlier findings of low relapse incidence of leukemia following cord blood transplantations (25). In this scenario, the high frequency of public V δ 2^{neg} TCR δ chains in the cord blood cells could contribute substantially to a potential abundance of tumor-reactive receptors in the post-transplant repertoire (39, 40). In addition, CMV reactivation in patients who received non-T-cell-depleted peripheral blood stem cells after myeloablative conditioning regimens was associated with a reduced risk of leukemic relapse (44). We and others have suggested that this effect is partially mediated via $\gamma\delta$ T cells (25, 45).

Willcox *et al.* have identified endothelial protein C receptor (EPCR) as a CMV-induced stress ligand for V γ 4V δ 5 T cells (22). We now classify the receptors chains of CMV-reactive clones partially as public. In one patient, we detected public TCR γ and TCR δ chains with the EPCR reactivity to be present at the tumor side, suggesting that these chains, particularly the V δ 5, could form receptors with tumor-reactivity. Indeed, $\gamma\delta$ TILs in many TNBC patients expressed a shared V δ 5 TCR chain associated with anti-CMV responses. However, all TNBC patients with this particular shared V δ 5 TCR chain were CMV-negative. It is conceivable that CMV reactivations may not be clinically beneficial only in leukemias (46), but also could confer some protection against certain solid tumors. However, we also demonstrate with multiple TCR complexes comprised of public TCR γ or δ chains, that the corresponding chain is frequently private and key for anti-tumor activity. Therefore in these patients the corresponding counterpart might not mediate CMV or tumor reactivity when paired with a different opposing chain, as also suggested by our data and others (33).

Many clinical trials with ex vivo and in vitro expanded $\gamma\delta$ T cells were not very successful (47) and the functional diversity within the $\gamma\delta$ T cell repertoire may be a major factor contributing to these numerous failures of clinical translations of $\gamma\delta$ T cell-based therapeutics (48). The observation that public TCR δ chains are frequently involved in the observed anti-tumor reactivity points toward possible strategies for creating potentially clinically-important T cells with defined $\gamma\delta$ TCRs (TEGs) utilizing

shared TCR γ and/or δ chains. Indeed, high-throughput analyses of CDR3 regions combined with single-cell sequencing of $\gamma\delta$ TILs can narrow down very quickly the extremely diverse TCR δ repertoire from 10^{18} to literally perhaps less than 10^2 sequences of therapeutic interest for generating novel therapeutic TEGs (49). Such
5 receptors could be quickly analyzed functionally and custom made for each patient. Nevertheless, the absence of HLA restriction during the measured antitumor activities suggests the potential usage of these novel, freshly-created TEGs in numerous genetically unrelated patients.

10 For generating off the shelf products analyzing cord blood donors with a natural high frequency of public TCR γ and δ chains when compared to the adult repertoire could be a natural starting point. Shared TCR γ and δ chains have also been reported in patients with acute leukemia (50) and ovarian cancer (51) and such samples could also be an interesting source for further functional analyses. However, though the frequency and
15 potency of tumor reactive receptors was in our analyses always higher within complexes harboring shared as compared to non-shared receptor chains, the identification of interesting receptors will always require a functional confirmation of defined receptors against a comprehensive tumor panel. Many analyzed receptor complexes harboring shared TCR γ or δ chains were complementary in reactivity
20 against completely different types of cancer or not reactive at all suggesting that these receptors target complementary and so far unknown ligands.

Screening of TEGs against a limited panel of tumor cell lines and the potential absence of co-stimulatory signals that might be needed for the $\gamma\delta$ T cell response somewhat
25 restrict the interpretation of our data. Clearly, distinct TCR $\gamma\delta$ pairs mediated different patterns of tumor reactivity, indicating that each individual pair of public TCR $\gamma\delta$ chains may be reactive only against defined subsets of targets. One limitation of our studies clearly arises from screening of TEGs against a rather limited panel of tumor cell lines as we observed different patterns for tumor reactivity for different $\gamma\delta$ TCR pairs. In
30 addition, $\gamma\delta$ TCR pairs might depend on additional co-stimulation not present within the TEG format. However, all tested receptors were active within the TEG format and expression within NEG or GEGs did not show additional activity. This suggests that also for many V $\delta 2^{\text{neg}}$ $\gamma\delta$ T cells the TCR $\gamma\delta$ chain alone is sufficient for full T cell

activation but also that each individual pair with public TCR chains is apparently reactive only against defined subsets of targets. While an inside out mechanism involving CD277 (52, 53) and RhoB (24) has now been established for V γ 9V δ 2 TCR-mediated recognition, most ligands remain to be defined for many V δ 2^{neg} TCR $\gamma\delta$ combinations and testing of potentially interesting chains should therefore always include either a large and comprehensive tumor panels.

In summary, we describe frequencies of shared TCR δ chains within the peripheral blood $\gamma\delta$ TCRs and $\gamma\delta$ TILs. We also demonstrate that public TCR γ or δ chains are active and complementary participants in daily cancer immune surveillance. However, despite functional proofs of direct tumor-reactivity of involved receptors, we did not observe substantial clonal expansion of shared TCR δ repertoires within the local microenvironment, suggesting proliferation deficiency of $\gamma\delta$ TILs in advanced cancer patients. In contrast, we detected shared tumor-reactive TCR δ s within the peripheral blood repertoire of healthy individuals, which was compatible with the proposed role of the public tumor-reactive TCR δ s in cancer immune surveillance. Thus, the identification of therapeutically relevant TCRs might be facilitated by analyzing shared TCR $\gamma\delta$ repertoires in different compartments in healthy and cancer patients. This involvement of public V δ 2⁺ and V δ 2^{neg} TCR clonotypes in tumor immune surveillance could be exploited for augmenting tumor-specific immunity. In particular, clarifying the precise nature of the recognized ligands, the exact mode of their recognition and their tissue distribution and regulation of expression may lead to developing novel therapeutic approaches and thus improving future $\gamma\delta$ T cell-based cancer therapies.

Table 5
Overview used primers for NGS library preparation

cDNA synthesis	
TRB reverse primer constant	CAGTATCTGGAGTCATTGA
TRD reverse primer constant	CTTGGATGACACGAGAT
Template switch adaptor	AAGCAGUGGTAUCAACGCAGAGUNNNNUNNNNUNNNNUCTT(rG) ₄
1st PCR	
TRB nested reverse primer constant region	TGCTTCTGATGGCTCAAACAC
TRD nested reverse primer constant region	AACGGATGGTTTGGTATGAG
Forward primer (anneals to switch adaptor)	CACTCTATCCGACAAGCAGTGGTATCAACGCAG
2nd PCR	
TRB nested reverse primer constant region	ACACSTTKTTCAGGTCCTC
TRD nested reverse primer constant region	TTTGGTATGAGGCTGACTTC
Forward primer (anneals to switch adaptor)	CACTCTATCCGACAAGCAGT

Antibody	Fluorochrome/ color	Species	Diluti on	Isoty pe	Clone	Cat No	Source
TCR γ		Mouse monoclo nal	1:40	IgG	γ 3,20	TCR11 53	Thermo Fisher
TCR γ	FITC	Mouse monoclo nal	1:20	IgG1	IMMU5 10	IM1349	Beckma n Coulter
CD69		Mouse monoclo nal	1:30	IgG	CH11	NCLCD 69	Leica Biosyste ms
CMV		Mouse monoclo nal	Flex Syste m	IgG1, K+L	CCH2+ DDG9	IS752	Dako
Cleaved Caspase 3		Rabbit monoclo nal	1:700	IgG	5A1E	mAb96 64	Cell Signallin g
IFN γ		Goat polyclon al	1:200	IgG	polyclo nal	AF285 NA	R&D Systems
IL-17		Goat polyclon al	1:200	IgG	polyclo nal	AF317 NA	R&D Systems
anti-rabbit / Alexa Fluor 568 sec. Ab		Donkey	1:500	IgG	polyclo nal	Ab1754 70	Abcam
anti- Mouse IgG / Alexa Fluor 488 sec. Ab		Rabbit	1:700	IgG H+L	polyclo nal	A11059	Thermo Fisher
anti-goat IgG / Alexa		Donkey	1:500	IgG	polyclo nal	ab1501 31	Abcam

Fluor 647 sec. Ab							
anti-goat IgG / Alexa Fluor 594 sec. Ab		Donkey	1:500	IgG	polyclo nal	Ab1501 32	Abcam
CD3	eFluor450	Mouse monoclo nal	1:40	IgG2 a	OKT3	16- 0037- 81	eBioscie nce
TCR $\alpha\beta$	APC	Mouse monoclo nal	1:10	IgG1	IP26	14- 9986- 42	eBioscie nce
TCR $\gamma\delta$	PE	Mouse monoclo nal	1:10	IgG1	IMMU5 10	B49716	Beckma n Coulter
TCR $\gamma\delta$	APC	Mouse monoclo nal	1:5	IgG1	B1 (RUO)	555718	BD bioscien ces

Table 6: Overview of used antibodies in different experiments.

Primers Single cell PCR	
Forward primers	
Vg18-F1	AGGGGAAGGCCCCACAGCGTCTTC
Vg18-F2:	CAGCGTCTTCWGTACTATGAC
Vg9-F1	TGACGGCACTGTCAGAAAGGAATC
Vg9-F2	TGAGGTGGATAGGATACCTGAAAC
Vd1-F1(Biomed)	ATGCAAAAAGTGGTCGCTATTCTG
Vd1-F2	CAACTTCAAGAAAGCAGCGAAATC
Vd2-F1(Biomed)	ATACCGAGAAAAGGACATCTATG
Vd2-F2	CAAGGTGACATTGATATTGCAAAG
Vd3-F1	GGTTTTCTGTGAAACACATTCTGAC
Vd3-F2	CTTTCACCTTGGTGATCTCTCCAG
Vd4-F1(Biomed)	ATGACCAGCAAAATGCAACAGAAG
Vd4-F2	CGCTACTCATTGAATTTCCAGAAG
Vd5-F1(Biomed)	TACCCTGCTGAAGGTCCTACATTC
Vd5-F2	CTGTCTTCTTAAACAAAAGTGCCAAG
Vd6-F1 (Biomed)	CCCTGCATTATTGATAGCCATACG
Vd6-F2	TGCCAAGCAGTTCTCATTGCATATC
Reverse primers	
TRg-JP1/2-R1	TTACCAGGYGAAGTTACTATGAGC
TRg-J1/2-R1	AAGTGTTGTTCCACTGCCAAAGAG
TRg-JP-R1	AAGCTTTGTTCCGGGACCAAATAC
Jd1-R1	TTGGTTCCACAGTCACACGGGTTC
Jd2-R1	CTGGTTCCACGATGAGTTGTGTTC
Jd3-R1	CAACTCACGGGGCTCCACGAAGAG
Jd4-R1	TTGTACCTCCAGATAGGTTCTTTG

Table 7

Overview of used primers for single cell PCR

Shared TCRδ sequences (dataset 1) clonal frequency >0,1%	
Sequence CDR3δ	Identified in donor
Vδ1	
CALGDSYGGGPLYTDKLIF*	13, 19
Vδ2	
CACDVLGDTDKLIF**	18, 19, 20
CACDLLGDTGDKLIF**	25, 29
CACDTAGGSWDTRQMFF**	13, 19
CACDTLGAYTDKLIF**	20, 25
CACDTLGDTDKLIF**	19, 23
CACDTLLGDSSWDTRQMFF**	13, 19
CACDTPSSWDTRQMFF**	13, 19
CACDTTGGPSSWDTRQMFF**	12, 23
CACDTVGDTDKLIF**	11, 29
CACDTVGGTDKLIF**	15, 19
CACDTVGYTDKLIF**	13, 25
CACDTVLGDTRSWDTRQMFF**	20, 70
CACDTWGTDKLIF**	29, 31
CACDTWGYTDKLIF**	29, 31

Table 8.

Examples of shared sequences within 14 healthy donors were identified. CDR3δ sequences obtained with NGS of dataset 1 with a clonal frequency >0.1% were

5 compared to identify shared sequences between donors. *Indicates sequence of clone FE11 which has been isolated from an additional healthy donor. **Indicates sequence was also identified in another dataset.

P n t	Cln	TRG Rearrangement	TRG CDR3	TRD Rearrangement	TRD CDR3
B	9 [#]	TRGV5*GJ1**	CATWDRLYYKKLF&	TRDV1*DD3*D J1*	CALGNGNHIGYWRYTDKLIF
B	23 [#]	TRGV8*GJ1	CATWDNYKKLF	TRAV29/DV5DD3*D J1*	CAASSPIRGYTGSDKLIF%
C	132 [#] \$	TRGV4*GJ1*	CATWDGFPYYKKLF	TRAV29/DV5DJ1DD 2DD3*	CAASSPIRGYTGSDKLIF%
D	37 [#]	TRGV8*GJ1	CATWDNYMKLF	TRAV29/DV5 *DD2*DD3*D J1	CAASSPIRGYTGSDKLIF%
E	113 [#]	TRGV4*GJ1	CATWDGPPYYKKLF	TRAV29/DV5 *DD2DD3**DJ1	CAASSPIRGYTGSDKLIF%
F	2 [#]	TRGV4*GJ1	CATWDGPPYYKKLF	TRDV3 *DD2*DD3*D J1	CASSYTLKLGDTTPGRVRDWKLIF
F	4 [#]	TRGV2GJ1	CATWDGQKKLF	TRDV1*DD3*D J1	CALGELRYWGIVDKLIF
Z	e11 [#]	TRGV8*GJ1	CATWDNYKKLF	TRDV1*DD3*01DJ1	CALGDYLGDKYPSYDLLGDTTDLKIF
Z	i11 [#]	TRGV8*GJ1	CATWDNYKKLF	TRDV1*DD1*DD2*D D3*D J1	CALGELRGQISFLYLLGDTTDLKIF

Table 9
Overview of examples of paired sequences of $\gamma\delta$ TIL's found in patients with triple negative breast cancer. Grey scale colors indicate identical sequences. # Indicates that these clones were used to generate TEG's for functional essays. \$ Indicates clone published by Lafarge et al; doi:10.1002/eji.200425837. % Indicates delta chain published by Lafarge et al; doi:10.1002/eji.200425837. & Indicates gamma chain published by Uldirch et al.; doi:10.1038/ni.2713

Example 3: Stability testing:

A pharmaceutical composition described herein containing at least one polypeptide described herein or an engineered cell expressing a polypeptide described herein is stored in a sealed container at 25°C or 4°C and the container is placed in an atmosphere having 30%, 40%, 50%, 60%, 70%, 75%, 80%, 90% or 95% relative humidity. After 1 month, 2 months, 3 months, 6 months, 1 year, 1.5 years, 2 years, 2.5 years or 3 years, at least 50%, 60%, 70%, 80% or 90% of the pharmaceutical composition shall remain as determined by standard protocols.

While specific embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following

claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

List of references

1. Chien YH, Meyer C, Bonneville M. 2014. gammadelta T Cells: First Line of Defense and Beyond. *Annu.Rev.Immunol.*
- 5 2. Kim SK, Cornberg M, Wang XZ, Chen HD, Selin LK, Welsh RM. 2005. Private specificities of CD8 T cell responses control patterns of heterologous immunity. *J Exp Med* 201: 523-33
3. Miles JJ, Elhassen D, Borg NA, Silins SL, Tynan FE, Burrows JM, Purcell AW, Kjer-Nielsen L, Rossjohn J, Burrows SR, McCluskey J. 2005. CTL recognition
10 of a bulged viral peptide involves biased TCR selection. *J Immunol* 175: 3826-34
4. Venturi V, Price DA, Douek DC, Davenport MP. 2008. The molecular basis for public T-cell responses? *Nat Rev Immunol* 8: 231-8
5. Wang C, Liu Y, Cavanagh MM, Le Saux S, Qi Q, Roskin KM, Looney TJ, Lee
15 JY, Dixit V, Dekker CL, Swan GE, Goronzy JJ, Boyd SD. 2015. B-cell repertoire responses to varicella-zoster vaccination in human identical twins. *Proc Natl Acad Sci U S A* 112: 500-5
6. Kuball J, Dossett ML, Wolfl M, Ho WY, Voss RH, Fowler C, Greenberg PD. 2007. Facilitating matched pairing and expression of TCR chains introduced
20 into human T cells. *Blood* 109: 2331-8
7. Bolotin DA, Poslavsky S, Mitrophanov I, Shugay M, Mamedov IZ, Putintseva EV, Chudakov DM. 2015. MiXCR: software for comprehensive adaptive immunity profiling. *Nat Methods* 12: 380-1
8. Shugay M, Britanova OV, Merzlyak EM, Turchaninova MA, Mamedov IZ,
25 Tuganbaev TR, Bolotin DA, Staroverov DB, Putintseva EV, Plevova K, Linnemann C, Shagin D, Pospisilova S, Lukyanov S, Schumacher TN, Chudakov DM. 2014. Towards error-free profiling of immune repertoires. *Nat Methods* 11: 653-5
9. Scheper W, van Dorp S, Kersting S, Pietersma F, Lindemans C, Hol S,
30 Heijhuurs S, Sebestyen Z, Grunder C, Marcu-Malina V, Marchant A, Donner C, Plachter B, Vermijlen D, van Baarle D, Kuball J. 2013. gammadeltaT cells elicited by CMV reactivation after allo-SCT cross-recognize CMV and leukemia. *Leukemia* 27: 1328-38

10. Scheper w. 2014. Cancer immunotherapy using $\gamma\delta$ T cells: dealing with diversity. *Thesis*
11. Grunder C, van DS, Hol S, Drent E, Straetemans T, Heijhuurs S, Scholten K, Scheper W, Sebestyen Z, Martens A, Strong R, Kuball J. 2012. gamma9 and
5 delta2CDR3 domains regulate functional avidity of T cells harboring gamma9delta2TCRs. *Blood* 120: 5153-62
12. Straetemans T, Grunder C, Heijhuurs S, Hol S, Slaper-Cortenbach I, Bonig H, Sebestyen Z, Kuball J. 2015. Untouched GMP-ready purified engineered immune cells to treat cancer. *Clin Cancer Res*
- 10 13. de Witte MA, Kierkels GJ, Straetemans T, Britten CM, Kuball J. 2015. Orchestrating an immune response against cancer with engineered immune cells expressing alphabetaTCRs, CARs, and innate immune receptors: an immunological and regulatory challenge. *Cancer Immunol Immunother* 64: 893-902
- 15 14. Salgado R, Denkert C, Campbell C, Savas P, Nuciforo P, Aura C, de Azambuja E, Eidtmann H, Ellis CE, Baselga J, Piccart-Gebhart MJ, Michiels S, Bradbury I, Sotiriou C, Loi S. 2015. Tumor-Infiltrating Lymphocytes and Associations With Pathological Complete Response and Event-Free Survival in HER2-
20 Positive Early-Stage Breast Cancer Treated With Lapatinib and Trastuzumab: A Secondary Analysis of the NeoALTTO Trial. *JAMA Oncol* 1: 448-54
15. Loi S, Dushyanthen S, Beavis PA, Salgado R, Denkert C, Savas P, Combs S, Rimm DL, Giltane JM, Estrada MV, Sanchez V, Sanders ME, Cook RS, Pilkinton MA, Mallal SA, Wang K, Miller VA, Stephens PJ, Yelensky R, Doimi FD, Gomez H, Ryzhov SV, Darcy PK, Arteaga CL, Balko JM. 2016.
25 RAS/MAPK Activation Is Associated with Reduced Tumor-Infiltrating Lymphocytes in Triple-Negative Breast Cancer: Therapeutic Cooperation Between MEK and PD-1/PD-L1 Immune Checkpoint Inhibitors. *Clin Cancer Res* 22: 1499-509
16. Nielsen TO, Hsu FD, Jensen K, Cheang M, Karaca G, Hu Z, Hernandez-
30 Boussard T, Livasy C, Cowan D, Dressler L, Akslen LA, Ragaz J, Gown AM, Gilks CB, van de Rijn M, Perou CM. 2004. Immunohistochemical and clinical characterization of the basal-like subtype of invasive breast carcinoma. *Clin Cancer Res* 10: 5367-74

17. Bender C, Zipeto D, Bidoia C, Costantini S, Zamo A, Menestrina F, Bertazzoni U. 2009. Analysis of colorectal cancers for human cytomegalovirus presence. *Infect Agent Cancer* 4: 6
- 5 18. Mamedov IZ, Britanova OV, Zvyagin IV, Turchaninova MA, Bolotin DA, Putintseva EV, Lebedev YB, Chudakov DM. 2013. Preparing unbiased T-cell receptor and antibody cDNA libraries for the deep next generation sequencing profiling. *Front Immunol* 4: 456.
19. Hidalgo JV, Bronsert P, Orlowska-Volk M, Diaz LB, Stickeler E, Werner M, Schmitt-Graeff A, Kayser G, Malkovsky M, Fisch P. 2014. Histological Analysis of gammadelta T Lymphocytes Infiltrating Human Triple-Negative Breast Carcinomas. *Front Immunol* 5: 632
- 10 20. Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9: 671-5
- 15 21. Brauninger A, Hansmann ML, Strickler JG, Dummer R, Burg G, Rajewsky K, Kuppers R. 1999. Identification of common germinal-center B-cell precursors in two patients with both Hodgkin's disease and non-Hodgkin's lymphoma. *N Engl J Med* 340: 1239-47
22. Straetemans T, Grunder C, Heijhuurs S, Hol S, Slaper-Cortenbach I, Bonig H, Sebestyen Z, Kuball J. 2015. Untouched GMP-Ready Purified Engineered Immune Cells to Treat Cancer. *Clin Cancer Res* 21: 3957-68
- 20 23. Grunder C, van DS, Hol S, Drent E, Straetemans T, Heijhuurs S, Scholten K, Scheper W, Sebestyen Z, Martens A, Strong R, Kuball J. 2012. gamma9 and delta2CDR3 domains regulate functional avidity of T cells harboring gamma9delta2TCRs. *Blood* 120: 5153-62
- 25 24. Sebestyen Z, Scheper W, Vyborova A, Gu S, Rychnavska Z, Schiffler M, Cleven A, Cheneau C, van Noorden M, Peigne CM, Olive D, Lebbink RJ, Oostvogels R, Mutis T, Schuurhuis GJ, Adams EJ, Scotet E, Kuball J. 2016. RhoB Mediates Phosphoantigen Recognition by Vgamma9Vdelta2 T Cell Receptor. *Cell Rep* 15: 1973-85
- 30 25. Scheper W, van Dorp S, Kersting S, Pietersma F, Lindemans C, Hol S, Heijhuurs S, Sebestyen Z, Grunder C, Marcu-Malina V, Marchant A, Donner C, Plachter B, Vermijlen D, van Baarle D, Kuball J. 2013. gammadeltaT cells

elicited by CMV reactivation after allo-SCT cross-recognize CMV and leukemia. *Leukemia* 27: 1328-38

26. Marcu-Malina V, Heijhuurs S, van BM, Hartkamp L, Strand S, Sebestyen Z, Scholten K, Martens A, Kuball J. 2011. Redirecting alphabeta T cells against cancer cells by transfer of a broadly tumor-reactive gammadeltaT-cell receptor. *Blood* 118: 50-9
27. Holtmeier W, Chowers Y, Lumeng A, Morzycka-Wroblewska E, Kagnoff MF. 1995. The delta T cell receptor repertoire in human colon and peripheral blood is oligoclonal irrespective of V region usage. *J Clin Invest* 96: 1108-17
28. Chien YH, Meyer C, Bonneville M. 2014. gammadelta T Cells: First Line of Defense and Beyond. *Annu.Rev.Immunol.*
29. Hidalgo JV, Bronsert P, Orlowska-Volk M, Diaz LB, Stickeler E, Werner M, Schmitt-Graeff A, Kayser G, Malkovsky M, Fisch P. 2014. Histological Analysis of gammadelta T Lymphocytes Infiltrating Human Triple-Negative Breast Carcinomas. *Front Immunol* 5: 632
30. Gentles AJ, Newman AM, Liu CL, Bratman SV, Feng W, Kim D, Nair VS, Xu Y, Khuong A, Hoang CD, Diehn M, West RB, Plevritis SK, Alizadeh AA. 2015. The prognostic landscape of genes and infiltrating immune cells across human cancers. *Nat Med* 21: 938-45
31. Coffelt SB, Kersten K, Doornebal CW, Weiden J, Vrijland K, Hau CS, Verstegen NJ, Ciampricotti M, Hawinkels LJ, Jonkers J, de Visser KE. 2015. IL-17-producing gammadelta T cells and neutrophils conspire to promote breast cancer metastasis. *Nature* 522: 345-8
32. Li B, Li T, Pignon JC, Wang B, Wang J, Shukla SA, Dou R, Chen Q, Hodi FS, Choueiri TK, Wu C, Hacohen N, Signoretti S, Liu JS, Liu XS. 2016. Landscape of tumor-infiltrating T cell repertoire of human cancers. *Nat Genet*
33. Willcox CR, Pitard V, Netzer S, Couzi L, Salim M, Silberzahn T, Moreau JF, Hayday AC, Willcox BE, Chanet-Merville J. 2012. Cytomegalovirus and tumor stress surveillance by binding of a human gammadelta T cell antigen receptor to endothelial protein C receptor. *Nat.Immunol.* 13: 872-9

34. Uldrich AP, Le Nours J, Pellicci DG, Gherardin NA, McPherson KG, Lim RT, Patel O, Beddoe T, Gras S, Rossjohn J, Godfrey DI. 2013. CD1d-lipid antigen recognition by the gammadelta TCR. *Nat Immunol* 14: 1137-45
35. Couzin-Frankel J. 2013. Breakthrough of the year 2013. Cancer immunotherapy. *Science* 342: 1432-3
36. Lim WA, June CH. 2017. The Principles of Engineering Immune Cells to Treat Cancer. *Cell* 168: 724-40
37. Bouchie A, DeFrancesco L, Sheridan C, Webb S. 2017. Nature Biotechnology's academic spinouts of 2016. *Nat Biotechnol* 35: 322-33
38. Sittig SP, Kollgaard T, Gronbaek K, Idorn M, Hennenlotter J, Stenzl A, Gouttefangeas C, Thor Straten P. 2013. Clonal expansion of renal cell carcinoma-infiltrating T lymphocytes. *Oncoimmunology* 2: e26014
39. Davey MS, Willcox CR, Joyce SP, Ladell K, Kasatskaya SA, McLaren JE, Hunter S, Salim M, Mohammed F, Price DA, Chudakov DM, Willcox BE. 2017. Clonal selection in the human Vdelta1 T cell repertoire indicates gammadelta TCR-dependent adaptive immune surveillance. *Nat Commun* 8: 14760
40. Ravens S, Schultze-Florey C, Raha S, Sandrock I, Drenker M, Oberdorfer L, Reinhardt A, Ravens I, Beck M, Geffers R, von Kaisenberg C, Heuser M, Thol F, Ganser A, Forster R, Koenecke C, Prinz I. 2017. Human gammadelta T cells are quickly reconstituted after stem-cell transplantation and show adaptive clonal expansion in response to viral infection. *Nat Immunol*
41. Ding Y, Ma F, Wang Z, Li B. 2015. Characteristics of the Vdelta2 CDR3 Sequence of Peripheral gammadelta T Cells in Patients with Pulmonary Tuberculosis and Identification of a New Tuberculosis-Related Antigen Peptide. *Clin Vaccine Immunol* 22: 761-8
42. Dadi S, Chhangawala S, Whitlock BM, Franklin RA, Luo CT, Oh SA, Toure A, Pritykin Y, Huse M, Leslie CS, Li MO. 2016. Cancer Immunosurveillance by Tissue-Resident Innate Lymphoid Cells and Innate-like T Cells. *Cell* 164: 365-77
43. Xiang Z, Liu Y, Zheng J, Liu M, Lv A, Gao Y, Hu H, Lam KT, Chan GC, Yang Y, Chen H, Tsao GS, Bonneville M, Lau YL, Tu W. 2014. Targeted activation

- of human Vgamma9Vdelta2-T cells controls epstein-barr virus-induced B cell lymphoproliferative disease. *Cancer Cell* 26: 565-76
44. Elmaagacli AH, Koldehoff M. 2016. Cytomegalovirus replication reduces the relapse incidence in patients with acute myeloid leukemia. *Blood* 128: 456-9
- 5 45. Airoidi I, Bertaina A, Prigione I, Zorzoli A, Pagliara D, Cocco C, Meazza R, Loiacono F, Lucarelli B, Bernardo ME, Barbarito G, Pende D, Moretta A, Pistoia V, Moretta L, Locatelli F. 2015. gammadelta T-cell reconstitution after HLA-haploidentical hematopoietic transplantation depleted of TCR-alpha-beta+/CD19+ lymphocytes. *Blood* 125: 2349-58
- 10 46. Scheper W, Grunder C, Straetemans T, Sebestyen Z, Kuball J. 2013. Hunting for clinical translation with innate-like immune cells and their receptors. *Leukemia*
47. Deniger DC, Moyes JS, Cooper LJ. 2014. Clinical applications of gamma delta T cells with multivalent immunity. *Front Immunol* 5: 636
- 15 48. Scheper W, Sebestyen Z, Kuball J. 2014. Cancer Immunotherapy Using gammadeltaT Cells: Dealing with Diversity. *Front Immunol* 5: 601
49. de Witte MA, Kierkels GJ, Straetemans T, Britten CM, Kuball J. 2015. Orchestrating an immune response against cancer with engineered immune cells expressing alpha-betaTCRs, CARs, and innate immune receptors: an
- 20 immunological and regulatory challenge. *Cancer Immunol Immunother* 64: 893-902
50. Meeh PF, King M, O'Brien RL, Muga S, Buckhalts P, Neuberg R, Lamb LS, Jr. 2006. Characterization of the gammadelta T cell response to acute leukemia. *Cancer Immunol.Immunother.* 55: 1072-80
- 25 51. Xu C, Zhang H, Hu H, He H, Wang Z, Xu Y, Chen H, Cao W, Zhang S, Cui L, Ba D, He W. 2007. Gammadelta T cells recognize tumor cells via CDR3delta region. *Mol Immunol* 44: 302-10
52. Sandstrom A, Peigne CM, Leger A, Crooks JE, Konczak F, Gesnel MC, Breathnach R, Bonneville M, Scotet E, Adams EJ. 2014. The intracellular B30.2 domain of butyrophilin 3A1 binds phosphoantigens to mediate activation
- 30 of human Vgamma9Vdelta2 T cells. *Immunity* 40: 490-500

53. Harly C, Guillaume Y, Nedellec S, Peigne CM, Monkkonen H, Monkkonen J, Li J, Kuball J, Adams EJ, Netzer S, chanet-Merville J, Leger A, Herrmann T, Breathnach R, Olive D, Bonneville M, Scotet E. 2012. Key implication of CD277/butyrophilin-3 (BTN3A) in cellular stress sensing by a major human gammadelta T-cell subset. *Blood* 120: 2269-79
- 5

Claims

- 1 Method for identifying δ T-cell or γ T-cell receptors chains or parts thereof that
5 mediate an anti-tumor or anti-infection response comprising the steps of:
a) providing amino acid sequences obtained from a donor, comprising δ T-cell or
 γ T-cell receptor chains or parts thereof each of said receptor chains or parts thereof
comprising a CDR3 region;
b) identifying amino acid sequences comprising δ T-cell or γ T-cell receptors chains
10 or parts thereof obtained in step a) that are shared between different donors;
c) confirming the anti-tumor or anti-infection response of the δ T-cell or γ T-cell
receptors chains or parts thereof identified in step b) by assessing the anti-tumor or
anti-infectious response of a T-cell expressing a nucleic acid molecule encoding
the amino acid sequence provided in step a).
15
2. Method according to claim 1, wherein in step a):
a. the donors are human beings,
b. at least 2 different donors are used,
c. at least one donor, preferably all donors are healthy and/or
20 d. at least one donor is diseased.
3. Method according to claim 1 or 2, wherein in step b) the sequences identified are
shared between at least 2 different donors.
- 25 4. Method according to any one of claims 1 to 3, wherein the nucleic acid molecule
encoding the δ T-cell or γ T-cell receptor chain or part thereof identified in step c) of
claim 1 is provided in an expression vector or in a retroviral vector.
5. Method according to any one of claims 1 to 4, wherein the step of confirming the
30 anti-tumor response comprises contacting the T-cell of step c) with a tumor cell
and measuring its ability to lyse the tumor cell and/or induce IFN- γ , IL-2 or TNF α .
6. A δ T-cell receptor chain or a part thereof, each comprising a CDR3 region, and

which δ T-cell receptor chain or part thereof, said δ T-cell receptor or part thereof comprises an amino acid sequence as defined in step c) of claim 1 and is obtainable by the method of claims 1-5.

- 5 7. A γ T-cell receptor chain or a part thereof, each comprising a CDR3 region, and which γ T-cell receptor chain or part thereof, said γ T-cell receptor or part thereof comprises an amino acid sequence as defined in step c) of claim 1 and is obtainable by the method of claims 1-5.
- 10 8. A δ T-cell receptor chain or part thereof according to claim 6, comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 7 and/or 17 and/or 96.
- 15 9. A γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 10, 18, 64 and/or 97.
- 20 10. A nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 50 and/or 62 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 50 and/or 62.
- 25
11. A nucleic acid molecule represented by a nucleotide sequence comprising a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 51 and/or 63 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence
- 30 encoded by a nucleotide sequence SEQ ID NO: 51 and/or 63.

12. A $\gamma\delta$ TCR comprising:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 7 and/or 17 and/or 96, and preferably
- a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 10, 18, 64 and/or 97.

10

13. A nucleic acid molecule encoding a $\square\square$ TCR according to claim 12, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 50 and/or 62 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 50 and/or 62, and preferably
- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 51 and/or 63 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 51 and/or 63.

20

14. A conjugate comprising a part of the δ T-cell receptor chain of claim 6 or 8 or comprising a part of the γ T-cell receptor chain of claim 7 or 9 linked to an agent.

25

15. A conjugate according to claim 14, wherein the agent is selected from the group consisting of a diagnostic agent, a therapeutic agent, an anti-cancer agent, a chemical, a nanoparticle, a chemotherapeutic agent or a fluorochrome.

30

16. A nucleic acid construct comprising a nucleic acid molecule encoding the amino acid sequence as identified in step c) of claim 1 or 6 or 7 or 8 or 9 and/or being according to claim 10, 11 or 13.

17. A vector comprising the nucleic acid construct of claim 16.

18. A vector according to claim 17 which is a retroviral vector or lentiviral vector.

19. A cell comprising the nucleic acid construct of claim 16 or the vector of claim 17 or

5 18.

20. A cell according to claim 19, which is a T cell.

21. A T cell according to claim 20, which is preferably used as a medicament and

10 wherein said T cell expresses:

- a δ T-cell receptor chain or part thereof comprising a CDR3 region, said δ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 7 and/or 17 and/or 96, and preferably
- 15 - a γ T-cell receptor chain or part thereof comprising a CDR3 region, said γ T-cell receptor chain or part thereof being represented by an amino acid sequence, said amino acid sequence comprising at least 60% sequence identity or similarity with amino acid sequence SEQ ID NO: 10, 18, 64 and/or 97.

20 22. A T cell according to claim 20, which is preferably used as a medicament and comprises a nucleic acid molecule encoding a $\alpha\beta$ TCR, said nucleic acid molecule being represented by a nucleotide sequence comprising:

- a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 50 and/or 62 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 50 and/or 62, and preferably
- 25 - a nucleotide sequence that has at least 60% sequence identity with SEQ ID NO: 51 and/or 63 and/or a nucleotide sequence that encodes an amino acid sequence that has at least 60% amino acid identity or similarity with an amino acid sequence encoded by a nucleotide sequence SEQ ID NO: 51 and/or 63.

30

-

23. A δ T-cell receptor chain, or a part thereof, according to claim 6 or 8, a nucleic acid molecule according to claim 10, a conjugate according to any one of claims 14-15, a nucleic acid construct according to claim 16, a vector according to claim 17 or 18, a cell according to any one of claim 19 to 22 for use as a medicament.

5

24. A δ T-cell receptor chain, or a part thereof, according to claim 6 or 8, a nucleic acid molecule according to claim 10, a conjugate according to any one of claims 14-15, a nucleic acid construct according to claim 16, a vector according to claim 17 or 18, a cell according to any one of claim 19 to 22 for use as a medicament wherein the
10 medicament is for preventing, treating, regressing, curing and/or delaying cancer or an infection.

25. A γ T-cell receptor chain, or a part thereof, according to claim 7 or 9, a nucleic acid molecule according to claim 11, a conjugate according to any one of claims 14-15, a
15 nucleic acid construct according to claim 16, a vector according to claim 17 or 18, a cell according to any one of claim 19 to 22 for use as a medicament.

26. A γ T-cell receptor chain, or a part thereof, according to claim 7 or 9, a nucleic acid molecule according to claim 11, a conjugate according to any one of claims 14-15, a
20 nucleic acid construct according to claim 16, a vector according to claim 17 or 18, a cell according to any one of claim 19 to 22 for use as a medicament wherein the medicament is for preventing, treating, regressing, curing and/or delaying cancer or an infection.

25 27. A pharmaceutical composition comprising:
a polypeptide comprising an amino acid sequence with at least 60% sequence identity with the amino acid sequence of SEQ ID NO: 7 or SEQ ID NO: 17 or SEQ ID NO: 96, and a pharmaceutically acceptable adjuvant, diluent or carrier, wherein said polypeptide is expressed in an engineered cell.

30

28. The pharmaceutical composition of claim 27, wherein said polypeptide comprises an amino acid sequence with 65%, 70%, 75%, 80%, 85%, 90%, 95% or 99% sequence identity with the amino acid sequence of SEQ ID NO: 7.

29. The pharmaceutical composition of claim 27 or 28, wherein said polypeptide comprises the amino acid sequence of SEQ ID NO: 7.

5 30. The pharmaceutical composition of claim 27, wherein said polypeptide comprises an amino acid sequence with 65%, 70%, 75%, 80%, 85%, 90%, 95% or 99% sequence identity with the amino acid sequence of SEQ ID NO: 17 or SEQ ID NO: 96.

31. The pharmaceutical composition of claim 27 or 30, wherein said polypeptide
10 comprises the amino acid sequence of SEQ ID NO: 17 or SEQ ID NO: 96.

32. A pharmaceutical composition comprising:
a polypeptide comprising an amino acid sequence with at least 60% sequence identity
with the amino acid sequence of SEQ ID NO: 10 or SEQ ID NO: 18 or SEQ ID NO: 64
15 or SEQ ID NO: 97, and a pharmaceutically acceptable adjuvant, diluent or carrier,
wherein said polypeptide is expressed in an engineered cell.

33. The pharmaceutical composition of claim 32, wherein said polypeptide comprises
an amino acid sequence with 65%, 70%, 75%, 80%, 85%, 90%, 95% or 99% sequence
20 identity with the amino acid sequence of SEQ ID NO: 10 and/or 64.

34. The pharmaceutical composition of claim 32 or 33, wherein said polypeptide
comprises the amino acid sequence of SEQ ID NO: 10 and/or 64.

25 35. The pharmaceutical composition of claim 32, wherein said polypeptide comprises
an amino acid sequence with 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 99%
sequence identity with the amino acid sequence of SEQ ID NO: 18 or SEQ ID NO: 97.

36. The pharmaceutical composition of claim 32 or 35, wherein said polypeptide
30 comprises the amino acid sequence of SEQ ID NO: 18 or SEQ ID NO: 97.

37. The pharmaceutical composition of any one of claims 27 to 36, wherein the
composition comprises:

a polypeptide comprising an amino acid sequence with at least 60% sequence identity with the amino acid sequence of SEQ ID NO: 7 or SEQ ID NO: 17 or SEQ ID NO: 96, a polypeptide comprising an amino acid sequence with at least 60% sequence identity with the amino acid sequence of SEQ ID NO: 10 or SEQ ID NO: 18 or SEQ ID NO: 64 or SEQ ID NO: 97 and a pharmaceutically acceptable adjuvant, diluent or carrier, wherein each of said polypeptide is expressed in an engineered cell.

38. The pharmaceutical composition of any one of claims 27-37, wherein said engineered cell is an effector cell.

10

39. The pharmaceutical composition of claim 38, wherein said effector cell is a T cell.

40. The pharmaceutical composition of any one of claims 27-39, wherein said carrier is at least one of saline, buffered saline, dextrose, water, glycerol and ethanol.

15

41. The pharmaceutical composition of any one of claims 27-40, wherein said adjuvant comprises a stabilizer.

42. The pharmaceutical composition of any one of claims 27-41, wherein said adjuvant comprises a preservative.

20

43. An engineered cell comprising at least one nucleic acid sequence encoding a δ T-cell receptor chain or part thereof comprising a CDR3 region comprising an amino acid sequence comprising at least 60% sequence identity with the amino acid sequence of SEQ ID NO: 7 or SEQ ID NO: 17 or SEQ ID NO: 96, and a γ T-cell receptor chain or part thereof comprising a CDR3 region comprising an amino acid sequence comprising at least 60% sequence identity with the amino acid sequence of SEQ ID NO: 10 or SEQ ID NO: 18 or SEQ ID NO: 64 or SEQ ID NO: 97.

25

30

44. The engineered cell of claim 43, wherein said cell is an engineered T cell.

45. The engineered cell of any one of claims 43 or 44, wherein said at least one nucleic acid is introduced into said cell by at least one genomic alteration.

46. The engineered cell of claim 45, wherein said genomic alteration comprises a
5 knock in comprising a viral introduction of said nucleic acid sequence.

47. The engineered cell of claim 46, wherein said viral introduction comprises a virus selected from the group consisting of a lentivirus, retrovirus, adenovirus, and any combination thereof.

10

48. The engineered cell of claim 45, wherein said genomic alteration comprises a genome alteration technique selected from CRISPR, Argonaute and AttSite site-specific serine recombinase gene editing systems.

15 49. A population of cells comprising the engineered cell of any one of claims 43 to 48.

50. The population of claim 49, wherein from about 60% to about 100% of said engineered cells express at least one of said δ T-cell receptor chain or part thereof and
20 said γ T-cell receptor chain or part thereof.

51. The population of any one of claims 49 to 50, wherein said engineered cells are a plurality of $\alpha\beta$ T cells.

25 52. The population of any one of claims 49 to 51, wherein when said population is administered to a subject in need thereof a cancer is controlled, reduced, or eliminated.

53. The population of claim 52, wherein said administration is effective in reducing tumor size by at least 30% as measured by computerized tomography (CT) scan.

30

54. Method for identifying δ T-cell or γ T-cell receptors chains or parts thereof that mediate an anti-tumor or anti-infection response comprising the steps of:

a) obtaining from a plurality of donors, δ T-cell receptor chains, γ T-cell receptor chains

or parts thereof, said receptor chains or parts thereof comprising a CDR3 region, and determining amino acid sequences of said receptors chains or parts thereof or nucleic acid sequences encoding the same;

- b) identifying sequences obtained in step a) that are shared between different donors;
- 5 c) assessing the anti-tumor or anti-infectious response of T-cells expressing or encoding said sequences shared between different donors.

1/16

Fig. 1




Fig. 2




Fig. 3




Fig. 4A




Fig. 4B




Fig. 4C




Fig. 4D




Fig. 5B

[illegible]

Fig. 5C

Shared Vβ2 TCR sequences between different datasets							
	Source	Dataset 1	Dataset 2	Dataset 3	Dataset 4	Dataset 5	Dataset 6 Dataset 7
Dataset 1 - Healthy donors	Peripheral blood	1225					
Dataset 2 - Ravens et al	Peripheral blood	30	1206				
Dataset 3 - Davey et al	Peripheral blood	95	107	9800			
Dataset 4 - Published sequences	Peripheral blood and tissue (healthy)	4	4	23	320		
Dataset 5 - Published vδ TIL's	Various tumor tissue	5	6	20	0	184	
Dataset 6 - Breast cancer single cell	Breast cancer tissue	0	0	0	0	0	7
Dataset 7 - Breast cancer NGS	Breast cancer tissue	17	14	40	1	3	1 482

Fig. 5D

Shared Vδ1 TCR sequences between different datasets							
	Source	Dataset 1	Dataset 2	Dataset 3	Dataset 4	Dataset 5	Dataset 6 Dataset 7
Dataset 1 - Healthy donors	Peripheral blood	202					
Dataset 2 - Ravens et al	Peripheral blood	0	126				
Dataset 3 - Davey et al	Peripheral blood	0	0	27368			
Dataset 4 - Published sequences	Peripheral blood and tissue (healthy)	0	0	0	94		
Dataset 5 - Published vδTIL's	Various tumor tissue	0	0	11	0	624	
Dataset 6 - Breast cancer single cell	Breast cancer tissue	0	0	0	0	0	22
Dataset 7 - Breast cancer NGS	Breast cancer tissue	0	0	0	0	0	2 482

Fig. 5E

Shared Vβ3 TCR sequences between different datasets							
Source		Dataset 1	Dataset 2	Dataset 3	Dataset 4	Dataset 5	Dataset 6 Dataset 7
Dataset 1 - Healthy donors	Peripheral blood	14					
Dataset 2 - Ravens et al	Peripheral blood	0	26				
Dataset 3 - Davey et al	Peripheral blood	0	0	687			
Dataset 4 - Published sequences	Peripheral blood and tissue (healthy)	0	0	0	22		
Dataset 5 - Published vβTIL's	Various tumor tissue	0	0	0	0	81	
Dataset 6 - Breast cancer single cell	Breast cancer tissue	0	0	0	0	0	13
Dataset 7 - Breast cancer NGS	Breast cancer tissue	0	0	0	0	1	1 43

Fig. 6





Fig. 7A








Fig. 7B

Fig. 8A



Patient	Identified seq.	Unique seq.
A	2	1
B	8	6
C	9	8
D	5	3
E	10	6
F	8	8
G	5	5
H	2	2
I	0	0
K	2	2
Z	4	4
Total	55	41

Fig. 8B



Patient	Identified seq.	Unique seq.
A	18	16
B	13	11
C	16	16
D	8	7
E	12	10
F	8	7
G	9	9
H	0	0
I	2	2
K	1	1
Z	3	2
Total	90	66

Fig. 9A




Fig. 9B






Fig. 10A



^ Indicates shared Vdelta1 chain with D13 and D19

Fig. 10B



15/16

Fig. 11A




Fig. 11B




Fig. 12

