



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2185340 C 2007/07/17

(11)(21) **2 185 340**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1995/03/10
(87) Date publication PCT/PCT Publication Date: 1995/09/14
(45) Date de délivrance/Issue Date: 2007/07/17
(85) Entrée phase nationale/National Entry: 1996/09/11
(86) N° demande PCT/PCT Application No.: US 1995/002900
(87) N° publication PCT/PCT Publication No.: 1995/024492
(30) Priorité/Priority: 1994/03/11 (US08/209,951)

(51) Cl.Int./Int.Cl. *C12N 15/32* (2006.01),
A01H 5/00 (2006.01), *C12N 15/82* (2006.01),
C12N 5/10 (2006.01), *C12P 21/02* (2006.01)

(72) Inventeurs/Inventors:
MCBRIDE, KEVIN E., US;
MALIGA, PAL, US

(73) Propriétaires/Owners:
STATE UNIVERSITY OF NEW JERSEY
RUTGERS, US;
CALGENE INC., US

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : EXPRESSION DE PROTEINES DE BACILLUS THURINGIENSIS CRY DANS LES PLASTES
(54) Title: EXPRESSION OF BACILLUS THURINGIENSIS CRY PROTEINS IN PLANT PLASTIDS

(57) **Abrégé/Abstract:**

Novel compositions and methods useful for genetic engineering of plant cells to provide increased expression in the plastids of a plant or plant cell of the *Bacillus thuringiensis* insecticidal protein.





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 15/32, C07K 14/435, A01N 63/02, A01H 5/00	A1	(11) International Publication Number: WO 95/24492 (43) International Publication Date: 14 September 1995 (14.09.95)
(21) International Application Number: PCT/US95/02900 (22) International Filing Date: 10 March 1995 (10.03.95) (30) Priority Data: 08/209,951 11 March 1994 (11.03.94) US (71) Applicants: CALGENE INC. [US/US]; 1920 Fifth Street, Davis, CA 95616 (US). STATE UNIVERSITY OF NEW JERSEY RUTGERS [US/US]; Office of Corporate Liaison and Technology Transfer, Administrative Service Building Annex II, Piscataway, NJ 08855-1179 (US). (72) Inventors: McBRIDE, Kevin, E.; 1309 Marina Circle, Davis, CA 95616 (US). MALIGA, Pal; 51 Mitchell Avenue, East Brunswick, NJ 08816 (US). (74) Agents: LASSEN, Elizabeth et al.; Calgene Inc., 1920 Fifth Street, Davis, CA 95616 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: EXPRESSION OF <i>BACILLUS THURINGIENSIS</i> CRY PROTEINS IN PLANT PLASTIDS (57) Abstract Novel compositions and methods useful for genetic engineering of plant cells to provide increased expression in the plastids of a plant or plant cell of the <i>Bacillus thuringiensis</i> insecticidal protein.		

EXPRESSION OF BACILLUS THURINGIENSIS
CRY PROTEINS IN PLANT PLASTIDS

5

INTRODUCTION

Field of the Invention

10 This invention relates to the application of genetic engineering techniques to plants. More specifically, the invention relates to compositions and methods expressing insecticidal *Bacillus thuringiensis* toxin proteins in plant plastids.

15 Background

Plastids of higher plants, i.e. chloroplasts, amyloplasts and chromoplasts, have the same genetic content, and thus are believed to be derived from a common precursor, known as a proplastid. The plastid genome is circular and varies in size among plant species from about 120 to about 217 kilobase pairs (kb). The genome typically includes a large inverted repeat, which can contain up to about 76 kilobase pairs, but which is more typically in the range of about 20 to about 30 kilobase pairs. The inverted repeat present in the plastid genome of various organisms has been described (Palmer, J. D. (1990) *Trends Genet.* 6:115-120).

30 One advantage of plant plastid transformation over nuclear transformation is that the plastids of most plants are maternally inherited, and consequently heterologous plastid genes are not pollen disseminated. This feature is particularly attractive for transgenic plants having altered agronomic traits, as introduced resistance or tolerance to natural or chemical conditions will not be transmitted to wild-type relatives.

Plant plastids are also major biosynthetic centers. In addition to photosynthesis in chloroplasts, plastids are responsible for production of important compounds such as amino acids, complex carbohydrates, fatty acids, and
5 pigments.

Plastids can be present in a plant cell at a very high copy number, with up to 50,000 copies per cell present for the chloroplast genome (Bendich, A. J. (1987) *BioEssays*
10 6:279-282). Thus, through plastid transformation plant cells can be engineered to maintain an introduced gene of interest at a very high copy number.

For all of the above reasons, the plastids of higher
15 plants present an attractive target for genetic engineering. Stable transformation of plastids has been reported in the green algae *Chlamydomonas* (Boynton et al. (1988) *Science* 240:1534-1538) and more recently in higher plants (Svab et al. (1990) *Proc. Natl. Acad. Sci. USA*
20 87:8526-8530; Svab and Maliga (1993) *Proc. Natl. Acad. Sci. USA* 90:913-917); (Staub, J. M. and Maliga, P. (1993), *EMBO J.* 12:601-606). The method disclosed for plastid transformation in higher plants relies on particle gun delivery of DNA containing a selectable marker and
25 targeting of the DNA to the plastid genome through homologous recombination.

There is a continuing need to introduce newly discovered or alternative *Bacillus thuringiensis* genes into
30 crop plants. Cry proteins (d-endotoxins) from *Bacillus thuringiensis* have potent insecticidal activity against a number of Lepidopteran, Dipteran, and Coleopteran insects. These proteins are classified CryI to CryV, based on amino acid sequence homology and insecticidal activity. Most
35 CryI proteins are synthesized as protoxins (ca. 130-140 kDa) then solubilized and proteolytically processed into active toxin fragments (ca. 60-70 kDa).

The poor expression of the protoxin genes from the nucleus of plants has heretofore required the use of 'truncated' versions of these genes. The truncated versions code only for the active toxin fragments. Other attempts to increase the expression efficiency have included resynthesizing the *Bacillus thuringiensis* toxin genes to utilize plant preferred codons. Many problems can arise in such extensive reconstruction of these large cry genes (approximately 3.5 Kb), and the process is both laborious and expensive.

Problems can also arise as new insect pests become endemic, or as existing populations develop resistance to a particular level or type of *Bacillus thuringiensis* toxin. Thus, there is a particular need for producing higher and thereby more effective levels of the *Bacillus thuringiensis* toxin in plants, a need which will only increase with time.

20

SUMMARY OF THE INVENTION

Various embodiments of this invention provide a construct for expression in a plant plastid comprising (a) to (c) as follows, operably joined in the 5' to 3' direction: (a) a promoter functional in a plant plastid; (b) a DNA sequence encoding an insecticidal *Bacillus thuringiensis* toxin; (c) a transcription termination region capable of terminating transcription in a plant plastid; the construct further comprising: (d) a gene encoding a marker construct for selection of plant cells comprising a plastid expressing said marker; and (e) DNA regions of homology to the genome of said plastid; wherein (a), (b), (c), and (d) of said construct are between said regions of homology in (e). Also provided is a plant cell plastid containing such a construct as well as plant cells and progeny cells thereof containing the construct.

Various embodiments of this invention provide a method for enhancing the expression of an insecticidal *Bacillus thuringiensis* toxin in a plant cell, wherein said method comprises expressing said *Bacillus thuringiensis* toxin in plastids of said plant cell, and wherein said toxin is expressed from a construct of this invention.

25

Various embodiments of this invention provide a transgenic plant cell exhibiting enhanced expression of an insecticidal *Bacillus thuringiensis* toxin, wherein said toxin is expressed from a construct of this invention and wherein said toxin is expressed in plastids of said plant cell.

30

By this invention, plastid expression constructs are provided which are useful for genetic engineering of plant

cells and which provide for enhanced expression of the *Bacillus thuringiensis* cry proteins in plant cell plastids. The transformed plastids should be metabolically active plastids, and are preferably maintained at a high copy
5 number in the plant tissue of interest, most preferably the chloroplasts found in green plant tissues, such as leaves or cotyledons.

The plastid expression constructs for use in this
10 invention generally include a plastid promoter region, a DNA sequence encoding a *Bacillus thuringiensis* cry protein, and a transcription termination region capable of terminating transcription in a plant plastid.

15 The plastid expression construct of this invention is preferably linked to a construct having a DNA sequence

encoding a selectable marker which can be expressed in a plant plastid. Expression of the selectable marker allows the identification of plant cells comprising a plastid expressing the marker.

5

In a preferred embodiment, vectors for transfer of the construct into a plant cell include means for inserting the expression and selection constructs into the plastid genome. This preferably comprises regions of homology to the target plastid genome which flank the constructs.

10

The construct of the invention preferably comprise a native *Bacillus thuringiensis* DNA encoding sequence, which may be a sequence encoding the protoxin. In a preferred embodiment, the DNA encoding sequence is the cryIA(c) gene.

15

Plant cell plastids containing the construct are also contemplated in the invention, as are plants, plant seeds, plant cells or progeny thereof containing plastids comprising the construct. A preferred plant for this purpose is cotton.

20

The invention also includes a method for enhancing the expression of an insecticidal *Bacillus thuringiensis* toxin in a plant cell, by expressing the *Bacillus thuringiensis* toxin in plastids of the plant cell. By this method the expression is enhanced separate from that enhancement which occurs from the increased copy number in plastid transformation.

25

30

By this invention the insecticidal *Bacillus thuringiensis* toxin is produced in plastids of a plant cell from the native DNA encoding sequence, with enhanced levels of expression of an insect resistant phenotype, as measured by insect feeding assays.

35

DESCRIPTION OF THE FIGURES

Figure 1 shows integration of cry genes from vectors
5 pZS223 and pZS224 into the wild-type plastid genome (Nt-
ptDNA) to yield transplastomes Nt-pZS223 ptDNA and Nt-
pZS224 ptDNA, respectively.

DETAILED DESCRIPTION OF THE INVENTION

10

A plastid expression construct of this invention
generally comprises a promoter functional in a plant
plastid, a DNA sequence encoding a *Bacillus thuringiensis*
cry protein, and a transcription termination region capable
15 of terminating transcription in a plant plastid. These
elements are provided as operably joined components in the
5' to 3' direction of transcription.

In developing the constructs the various fragments
20 comprising the regulatory regions and open reading frame
may be subjected to different processing conditions, such
as ligation, restriction enzyme digestion, PCR, *in vitro*
mutagenesis, linkers and adapters addition, and the like.
Thus, nucleotide transitions, transversions, insertions,
25 deletions, or the like, may be performed on the DNA which
is employed in the regulatory regions or the DNA sequences
of interest for expression in the plastids. Methods for
restriction digests, Klenow blunt end treatments,
ligations, and the like are well known to those in the art
30 and are described, for example, by Maniatis *et al.* (in
Molecular cloning: a laboratory manual (1982) Cold Spring
Harbor Laboratory Press, Cold Spring Harbor, NY).

During the preparation of the constructs, the various
35 fragments of DNA will often be cloned in an appropriate
cloning vector, which allows for amplification of the DNA,
modification of the DNA or manipulation of the DNA by
joining or removing sequences, linkers, or the like.

Preferably, the vectors will be capable of replication to at least a relatively high copy number in *E. coli*. A number of vectors are readily available for cloning, including such vectors as pBR322, vectors of the pUC series, the M13 series vectors, and pBluescript vectors (Stratagene; La Jolla, CA).

In order to provide a means of selecting the desired plant cells, vectors for plastid transformation typically contain a construct which provides for expression of a selectable marker gene. Marker genes are plant-expressible DNA sequences which express a polypeptide which resists a natural inhibition by, attenuates, or inactivates a selective substance, i.e., antibiotic, herbicide etc..

Alternatively, a marker gene may provide some other visibly reactive response, i.e., may cause a distinctive appearance or growth pattern relative to plants or plant cells not expressing the selectable marker gene in the presence of some substance, either as applied directly to the plant or plant cells or as present in the plant or plant cell growth media.

In either case, the plants or plant cells containing such selectable marker genes will have a distinctive phenotype for purposes of identification, i.e., they will be distinguishable from non-transformed cells. The characteristic phenotype allows the identification of cells, cell groups, tissues, organs, plant parts or whole plants containing the construct.

Detection of the marker phenotype makes possible the selection of cells having a second gene to which the marker gene has been linked. This second gene typically comprises a desirable phenotype which is not readily identifiable in transformed cells, but which is present when the plant cell or derivative thereof is grown to maturity, even under

conditions wherein the selectable marker phenotype itself is not apparent.

The use of such a marker for identification of plant cells containing a plastid construct has been described. Svab *et al.* (1993 *supra*). In the examples provided below, a bacterial *aadA* gene is expressed as the marker under the regulatory control of chloroplast 5' promoter and 3' transcription termination regions, specifically the tobacco 16S rRNA promoter *rrn* region and *rps16* 3' termination region. Numerous additional promoter regions may also be used to drive expression of the selectable marker gene, including various plastid promoters and bacterial promoters which have been shown to function in plant plastids.

Expression of the *aadA* gene confers resistance to spectinomycin and streptomycin, and thus allows for the identification of plant cells expressing this marker. The *aadA* gene product allows for continued growth and greening of cells whose chloroplasts comprise the selectable marker gene product. Cells which do not contain the selectable marker gene product are bleached. Selection for the *aadA* gene marker is thus based on identification of plant cells which are not bleached by the presence of streptomycin, or more preferably spectinomycin, in the plant growth medium.

A number of markers have been developed for use with plant cells, such as resistance to chloramphenicol, the aminoglycoside G418, hygromycin, or the like. Other genes which encode a product involved in chloroplast metabolism may also be used as selectable markers. For example, genes which provide resistance to plant herbicides such as glyphosate, bromoxynil or imidazolinone may find particular use. Such genes have been reported (Stalker *et al.*, *J. Biol. Chem.* (1985) 260:4724-4728 (glyphosate resistant EPSP); Stalker *et al.*, *J. Biol. Chem.* (1985) 263:6310-6314 (bromoxynil resistant nitrilase gene); and Sathasivan *et*

al., *Nucl. Acids Res.* (1990) 18:2188 (AHAS imidazolinone resistance gene)).

5 Stable transformation of tobacco plastid genomes by particle bombardment is reported (Svab et.al. (1990 *supra*) and Svab et al. (1993 *supra*)). The methods described therein may be employed to obtain plants homoplasmic for plastid expression constructs.

10 Generally, bombarded tissue is cultured for approximately 2 days on a cell division-promoting media, after which the plant tissue is transferred to a selective media containing an inhibitory amount of the particular selective agent, as well as the particular hormones and
15 other substances necessary to obtain regeneration for that particular plant species. Shoots are then subcultured on the same selective media to ensure production and selection of homoplasmic shoots.

20 Homoplasmy is verified by southern analysis. In the examples provided below, *Bam*HI-digested total cellular DNA is tested with various probes, specifically, a part of the plastid targeting fragment, an *aadA* fragment, a 1.8 kb *cry1A* fragment and a 3.5 kb fragment of the *cry73* coding
25 region. Southern blot analysis with these probes confirms the integration of the chimeric *cry* genes in the tobacco plastid genome to yield transplastome lines.

As an alternative to a second round of shoot
30 formation, the initial selected shoots may be grown to mature plants and segregation relied upon to provide transformed plants homoplastic for the inserted gene construct.

35 Where transformation and regeneration methods have been adapted for a given plant species, either by *Agrobacterium*-mediated transformation, bombardment or some other method, the established techniques may be modified

for use in selection and regeneration methods to produce plastid-transformed plants. For example, the methods described herein for tobacco are readily adaptable to other solanaceous species, such as tomato, petunia and potato.

5

In *Brassica*, *Agrobacterium*-mediated transformation and regeneration protocols generally involve the use of hypocotyl tissue, a non-green tissue which might contain a low plastid content. Thus, for *Brassica*, preferred target
10 tissues would include microspore-derived hypocotyl or cotyledonary tissues (which are green and thus contain numerous plastids) or leaf tissue explants. While the regeneration rates from such tissues may be low, positional effects, such as seen with *Agrobacterium*-mediated
15 transformation, are not expected, thus it would not be necessary to screen numerous successfully transformed plants in order to obtain a desired phenotype.

For cotton, transformation of *Gossypium hirsutum* L. cotyledons by co-cultivation with *Agrobacterium tumefaciens*
20 has been described by Firoozabady et al., *Plant Mol. Bio.* (1987) 10:105-116 and Umbeck et al., *Bio/Technology* (1987) 5:263-266. Again, as for *Brassica*, this tissue may contain insufficient plastid content for chloroplast
25 transformation. Thus, as for *Brassica*, an alternative method for transformation and regeneration of alternative target tissue containing chloroplasts may be desirable, for instance targeting green embryogenic tissue.

30 Other plant species may be similarly transformed using related techniques. Alternatively, microprojectile bombardment methods, such as described by Klein et al. (*Bio/Technology* 10:286-291) may also be used to obtain nuclear transformed plants comprising the viral single
35 subunit RNA polymerase expression constructs described herein. Cotton transformation by particle bombardment is reported in WO 92/15675, published September 17, 1992.

The vectors for use in plastid transformation preferably include means for providing a stable transfer of the plastid expression construct and selectable marker construct into the plastid genome. This is most conveniently provided by regions of homology to the target plastid genome. The regions of homology flank the construct to be transferred and provide for transfer to the plastid genome by homologous recombination, via a double crossover into the genome. The complete DNA sequence of the plastid genome of tobacco has been reported (Shinozaki et al. (1986) *EMBO J.* 5:2043-2049). Complete DNA sequences of the plastid genomes from liverwort (Ohyama et al. (1986) *Nature* 322:572-574) and rice (Hiratsuka et al. (1989) *Mol. Gen. Genet.* 217:185-194), have also been reported.

Where the regions of homology are present in the inverted repeat regions of the plastid genome (known as IRA and IRB), two copies of the transgene are expected per transformed plastid. The regions of homology within the plastid genome are approximately 1kb in size. Smaller regions of homology may also be used, and as little as 100 bp can provide for homologous recombination into the plastid genome. However, the frequency of recombination and thus the frequency of obtaining plants having transformed plastids decreases with decreasing size of the homology regions.

Examples of constructs having regions of homology the plastid genome are described in Svab et al. (1990 *supra*) and Svab et al. (1993 *supra*). Regions useful for recombination into tobacco and *Brassica* plastid genomes are also identified in the following examples, but homologous recombination and selection constructs may be prepared using many plastid DNA sequences, and to any target plant species. In the examples provided herein, the flanking tobacco plastid homology regions of the plastid expression construct direct the insertion of a *Bacillus thuringiensis* transgene into the tobacco genome between *trnV* and the

rps12 operon. Since integration into the plastid genome occurs by homologous recombination and the target site is in an inverted repeat region of the plastid genome, two copies of the transgene per plastid genome are expected.

- 5 Selection is made for the spectinomycin resistance marker phenotype expressed by the *aadA* gene.

In the example the native cry gene, i.e., having an unmodified coding region to the protoxin, is placed into a
10 plastid expression construct for expression of *Bacillus thuringiensis* toxin from the plant plastid.

A synthetic *Bacillus thuringiensis* gene is placed in the same expression construct as the protoxin gene. The
15 synthetic gene is designed to have tobacco RuBPCO small subunit codon usage, with an overall increase in the guanine plus cytosine content from 39% to 55% with respect to the native gene, and has been truncated to leave only those sequences which encode the active fragment of the
20 toxin. Such a gene is known to provide optimal expression from the plant nuclear genome.

Both encoding sequences are introduced via a chloroplast transformation vector (Fig. 1). Tobacco lines
25 containing the native encoding sequence to the protoxin demonstrate strong insecticidal bioactivity, as measured by insect feeding assays. In transformed plants containing the native encoding sequence, the *Bacillus thuringiensis* toxin is present as a component of up to about 5% or
30 greater of the total leaf protein, a level which is much higher than is present in the leaf of plants resulting from nuclear transformation.

Tobacco lines having a synthetic cryIA(c) gene
35 demonstrate no observable bioactivity. In plants containing the gene resynthesized to approximate the preferred codons of the plant genome, the mRNA to the toxin

appears degraded, and little or no toxin protein appears present in the leaf.

It is now shown that the native *Bacillus thuringiensis* gene achieves an expression level which is much higher in plastid expression than is possible with resynthesized sequence to the same gene, thus demonstrating that a gene having native bacterial encoding sequence can achieve high levels of expression in a plant plastid. The above results eliminate the need to resynthesize the *Bacillus thuringiensis* toxin genes to achieve high level expression in plants.

The invention now being generally described, it will be more readily understood by reference to the following examples which are included for purposes of illustration only and are not intended to limit the present invention.

EXAMPLES

In the experimental disclosure which follows, all temperatures are given in degrees centigrade ($^{\circ}$), weights are given in grams (g), milligram (mg) or micrograms (μ g), concentrations are given as molar (M), millimolar (mM) or micromolar (μ M) and all volumes are given in liters (l), milliliters (ml) or microliters (μ l), unless otherwise indicated.

EXAMPLE 1. PLASTID TRANSFORMATION VECTORS

Constructs and methods for use in transforming the plastids of higher plants are described in Svab et al. (1990 *supra*), Svab et al. (1993 *supra*) and Staub et al. (1993 *supra*). The complete DNA sequences of the plastid genome of tobacco are reported by Shinozaki et al. (1986 *supra*). All plastid DNA references in the following description are to the nucleotide number from tobacco.

The *cryIA(c)* gene is obtained from plasmid pBtkHD73 (Toagosei Chemical Co., Japan). This gene is further processed by digestion with *SmaI*/*NsiI* and a synthetic adapter is inserted (top strand: 5'-

5 CCCGGATCCATGGATAACAATCCGA-ACATCAATGAATGCA-3'; bottom strand: 5'-TTCATTGATGTTCGGATT-GTTATCCATGGATCCGGG-3'). The entire 5' untranslated region from the *cryIA(c)* gene is then removed, and an *NcoI* site is introduced at the natural start codon (position 163 of the nucleotide sequence (Adang
10 et al. (1985) *Gene* 36;289-300). A *BamHI* site is introduced just upstream of the *NcoI* site. Oligonucleotide mutagenesis is performed to introduce *BglIII* and *SallI* sites directly adjacent to the stop codon of the *cryIA(c)* gene, to facilitate removal of unwanted DNA 3' of the coding
15 region. The remaining sequence includes the entire encoding region to the protoxin.

A synthetic *cryIA(c)* gene encoding the active toxin fragment is constructed by annealing and ligating 70 and 90
20 base oligonucleotides, in a method as described (Wosnick et al. (1987) *Gene* 60;115-127). The synthetic gene is designed to have tobacco RuBISCO small subunit codon usage, including a guanine and cytosine content of 55%, with an *NcoI* site at the start codon and a *SallI* site at the stop
25 codon, while still encoding the amino acid sequence of the toxin. This synthetic gene is also truncated, however, so that the encoding region only provides the amino acid sequence to the active fragment of the protoxin.

30 A plastid transformation vector is used which carries a passenger gene in a *Prn(L)rbcL(S)/Trps16* expression cassette, with polylinker restriction sites. The *Prn(L)rbcL(S)* fragments are described in Svab et al. (1993 *supra*). To further secure the stability of the mRNAs, the
35 *Trps16* fragment is cloned downstream of the passenger gene encoding region. The *Trps16* fragment comprises the *rps16* gene 3'-regulatory region from nucleotides 5,087 to 4,939 in the tobacco plasmid DNA.

Chimeric genes are preferably inserted into the vector to direct their transcription towards the *rrn* operon. Thus, in the plastid genome, chimeric genes are transcribed from the *Prrn(L)rbcL(S)* 5'-regulatory region comprising the long *rrn* operon promoter fragment from nucleotides 102,561 to 102,677 of the tobacco plastid genome, which is fused with a synthetic leader sequence designed after the *rbcL* gene leader between nucleotides 57,569 to 57,584 in the plastid DNA.

The plastid transformation vector also carries a selectable spectinomycin resistance gene (*aadA*) under control of *psbA* gene expression signals. The regulatory and encoding sequences are also flanked by plastid DNA homology regions whose limits are bp 138,447 (*EcoRI*) to 140,219 (*HincII*) and 140,219 (*HincII*) to 141,382 (*BglIII*) of the tobacco plastid genome (Shinozaki et al. (1986 *supra*)). This directs insertion of foreign genes located between the flanking regions into the plastid between the *trnV* gene and the *rps12/7* operon.

This plastid transformation vector is digested with the *NcoI/SalI* restriction endonucleases to remove the encoding region of the passenger gene, which is then replaced with a *NcoI/SalI* fragment containing the synthetic *cryIA(c)* coding region, yielding a vector which is designated pZS223 (Fig. 1). The wild type *cryIA(c)* protoxin gene is similarly cloned as an *NcoI/SalI* fragment, yielding a plasmid designated pZS224. By this approach *Bacillus thuringiensis* DNA 3' of the protein coding region is omitted for both plasmids, pZS223 and pZS224.

The insertion of the respective *cry* genes from vectors pZS223 and pZS224 into the wild-type plastid genome (Nt-ptDNA) to yield transplastomes Nt-pZS223 and Nt-pZS224, respectively, is shown in Fig. 1. The abbreviations used in Fig. 1 are as follows: 16S, 16S rRNA gene; *trnV*, *trnV*

gene; *aadA*, spectinomycin resistance gene; *cry1A* and *cry73* are synthetic and native *Bacillus thuringiensis* δ -endotoxin genes, respectively. The restriction endonuclease cleavage sites are designated as follows: B, *Bam*HI; Bg, *Bgl*II; H, *Hind*III; N, *Nco*I; RI, *Eco*RI, RV, *Eco*RV; S, *Sal*I.

EXAMPLE 2. PLANT PLASTID TRANSFORMATION

Stable transformation of tobacco plastid genomes by particle bombardment is reported in Svab et.al. (1990 *supra*) and Svab et al. (1993 *supra*). The methods described therein may be employed to obtain plants transformed with the plastid expression constructs described herein. Such methods generally involve DNA bombardment of a target host explant, preferably an explant made from a tissue which is rich in metabolically active plastids, such as green plant tissues including leaves or cotyledons.

Tobacco seeds (*N. tabacum* v. Xanthi N/C) are surface sterilized in a 50% chlorox™ solution (2.5% sodium hypochlorite) for 20 minutes and rinsed 4 times in sterile H₂O. These are plated aseptically on a 0.2x MS salts media and allowed to germinate. The seedlings are grown on agar solidified MS media with 30g/l sucrose (Murashige et al. (1962) *Physiol. Plant* 15:493-497).

Tungsten microprojectiles (1.0μm) are coated with plasmid DNA according to Maliga (Maliga, P. (1993) *Methods in Plant Molecular Biology - A Laboratory Manual*, eds. Pal Maliga, Daniel Klessig, Anthony Cashmore, Wilhelm Gruissem and Joseph Varner; Cold Spring Harbor Press) and used to bombard mature leaves, placed abaxial side up on RMOP media; MS salts, 1 mg/l BAP, 0.1 mg/l NAA, 30 g/l sucrose and 0.7% phytagar™. Svab et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:8526-8530 (using the Bio-Rad PDS 1000 He system (Sanford et al., An improved, helium-driven Biolistic device, *Technique* 3:3-16)). Plasmids pZS223 and pZS224 are used as the coating plasmid DNA.

The bombarded tissue is then cultured for approximately 2 days on a cell division-promoting media, after which the plant tissue is transferred to a selective media containing an inhibitory amount of the particular selective agent. Transformed explants form green shoots in approximately 3-8 weeks. Leaves from these shoots are then subcultured on the same selective media to ensure production and selection of homoplasmic shoots.

10

EXAMPLE 3. DNA GEL BLOT ANALYSIS OF TRANSPLASTOMIC LINES

Transformed plants selected for marker *aadA* marker gene expression are analyzed to determine whether the entire plastid content of the plant has been transformed (homoplastic transformants). Typically, following two rounds of shoot formation and spectinomycin selection, approximately 50% of the transgenic plantlets which are analyzed are homoplastic, as determined by Southern blot analysis of plastid DNA. Homoplasmic plantlets are selected for further cultivation.

Following a second round of shoot formation and spectinomycin selection, two transplastomic lines for each construct are obtained, Nt-pZS223 and Nt-pZS224. These lines are checked for homoplasmy. Southern blot analysis is used to confirm the integration of the chimeric *cry* genes in the tobacco plastid genome. Preparation, electrophoresis, and transfer of DNA to filters is as described (Svab et al., (1993 *supra*)).

The complete disappearance of the 3.3 Kb native tobacco *Bam*HI fragment in the Nt-pZS223 and Nt-pZS224 transformants with a probe covering the region of integration, and the appearance of expected sized bands for the inserted DNA fragments in those transformants, 5.5 kb and 7.3 kb, respectively (see Fig. 1), establishes that the transformed plants are homoplasmic for the intended

constructs. Probing identical filters with the *aadA*,
cryIA(c) protoxin, and synthetic *cryIA(c)* genes
demonstrated a linkage of the *aadA* and *cryIA(c)* genes to
the expected 5.5 and 7.3 Kb *Bam*HI fragments as well as the
5 lack of these genes in the negative control.

EXAMPLE 4. INSECT BIOASSAYS

As described, the development of transformed plant
10 lines Nt-pZS223 and Nt-pZS224 is accomplished on RMOP media
supplemented with 500 mg/l Spectinomycin dihydrochloride.
Plants are subcloned on the same selective medium, by the
method according Svab et al. (1990 supra). Selected plants
are then rooted in MS media containing 1 mg/l IBA, 500 mg/l
15 Spectinomycin dihydrochloride and 0.6% phytagar.

Helicoverpa zea and *Heliothis virescens* eggs are
obtained from the USDA-ARS in Stoneville, MS. and allowed
to hatch. Neonate larva are placed on Tobacco Budworm Diet
20 from Bioserve (Frenchtown, NJ), and incubated in a 16:8
photoperiod at 28°C for 5 days. The larva develop during
this time to late second or early third instar.

At 5 days, fully expanded leaves are excised from the
25 tobacco plants and placed on 3 ml of 2% agar in a 32 well
rearing tray from CD International (Pitman, NJ). The larva
are placed 1 per well, sealed and incubated for 5 days at
the same conditions. At day 10, the leaf material consumed
by the insect is estimated and insects checked for
30 mortality. The larva are considered dead if they showed no
movement after prodding with forceps.

EXAMPLE 5. INSECTICIDAL FEEDING ACTIVITY

35 To determine the presence and relative amount of
active *Bacillus thuringiensis* d-toxin in the tobacco lines
homoplasmic for native protoxin and synthetic 'truncated'
cryIA(c) gene expression constructs, efficacy of these

plants to third instar *Heliothis virescens* (tobacco budworm) and *Helicoverpa zea* (corn earworm/cotton bollworm) larvae is tested. Both test insects are sensitive to the cryIA(c) toxin with *H. zea* being 10-fold more resistant than *H. virescens* (MacIntosh et al. (1990) *J. Invertebr. Pathol.* 56:258-266.).

Third instar larvae are chosen for the bioassay since the insects are more resistant to the toxin at this stage than are first instar larvae thus allowing a more stringent comparison between the control and test plants. Tobacco lines designated 4083 and 4084, derived by nuclear transformation with the same synthetic cryIA(c) gene as used in pZS223 and shown to be highly toxic to third instar *H. virescens* larvae, are used as positive controls in the bioassay. *Nicotiana tabacum* var. 'Petite Havana' serves as the negative control since this is the genetic background used to generate the transplastomic lines.

Table 1 is a summary of *Bacillus thuringiensis* tobacco insect feeding assays. The data demonstrates that transplastomic line Nt-pZS224 is very toxic to both *H. virescens* and *H. zea* as it causes 100% mortality to these insects while sustaining less than 2% total leaf damage. This result compares favorably to the results for positive control 4083 and 4084 tobacco plants. The 4083-2-4 plant when assayed with *H. zea* causes 100% mortality but sustains a much greater level of leaf feeding damage than the Nt-pZS224 tobacco line indicating less toxin production. Tobacco line 4084-4-1 performed comparably to Nt-pZS224 tobacco in feeding, although it does not compare to the levels of toxin produced in Nt-pZS224 when measured as a component of total leaf protein.

Tobacco line Nt-pZS223 shows no detectable bioactivity.

5

TABLE 1
SUMMARY OF BT TOBACCO INSECT FEEDING ASSAYS

10

Chloroplast	Vector	plants tested	Heliothis virescens^^	% Leaf eaten	Helicoverpa zea^^	% Leaf eaten	
synthetic toxin gene	pZS223	223-3	NO mortality	100%	NO mortality	100%	
		223-5	NO mortality	75%	NO mortality	100%	
		223-12	NT*		NO mortality	100%	
		223-13	NO mortality	75%	NT*		
	wild type protoxin gene	pZS224	224-5	100% mortality	2%	100% mortality	2%
			224-9	100% mortality	2%	100% mortality	2%

15

Nuclear Controls	synthetic toxin gene	pCGN4083	4083-1-2	100% mortality	2%	NT*	
			4083-2-4	NT*		100% mortality	40%
		pCGN4084	4084-8-5	100% mortality	2%	NT*	
			4084-1-1	NT*		100% mortality	2%
Untransformed Controls							
			control 1	25% mortality	75%	NO mortality	100%
			control 2	NO mortality	100%	NT*	
			control 3	50% mortality	75%	NT*	

20

^^ 10 third instar larva were individually tested per plant

*NT: Plant not tested

25

All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains.

30

35

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claim.

CLAIMS

What is claimed is:

1. A construct for expression in a plant plastid comprising (a) to (c) as follows, operably joined in the 5' to 3' direction:

(a) a promoter functional in a plant plastid;

(b) a DNA sequence encoding an insecticidal *Bacillus thuringiensis* toxin;

(c) a transcription termination region capable of terminating transcription in a plant plastid;

the construct further comprising:

(d) a gene encoding a marker construct for selection of plant cells comprising a plastid expressing said marker; and

(e) DNA regions of homology to the genome of said plastid;

wherein (a), (b), (c), and (d) of said construct are between said regions of homology in (e).

2. The construct according to Claim 1 wherein said plant plastid is a chloroplast.

3. The construct according to Claim 1 or 2, wherein said DNA encoding sequence encodes a protoxin.

4. The construct according to Claim 1, 2, or 3, wherein said DNA encoding sequence comprises a native *Bacillus thuringiensis* DNA encoding sequence.

5. A plant cell plastid containing the construct according to any one of Claims 1 to 4.

6. A plant cell or progeny cell thereof containing the construct according to any one of Claims 1 to 4.

7. The plant cell or progeny cell thereof according to Claim 6, wherein said plant cell is a cotton cell.

8. A method for enhancing the expression of an insecticidal *Bacillus thuringiensis* toxin in a plant cell, wherein said method comprises expressing said *Bacillus thuringiensis* toxin in plastids of said plant cell, and

wherein said toxin is expressed from a construct as defined in any one of Claims 1 to 4.

9. The method according to Claim 8, wherein said plant plastids are chloroplasts.

5 10. A transgenic plant cell exhibiting enhanced expression of an insecticidal *Bacillus thuringiensis* toxin, wherein said toxin is expressed from a construct as defined in any one of Claims 1 to 4, and wherein said toxin is expressed in plastids of said plant cell.

10 11. A plant cell according to Claim 10, wherein said expressed toxin comprises about 5% or greater of the total protein of said cell.

12. A plant cell according to Claim 10 or 11, wherein said plant cell is cotton.




FIG. 1