



(86) Date de dépôt PCT/PCT Filing Date: 1998/07/01
 (87) Date publication PCT/PCT Publication Date: 1999/01/14
 (45) Date de délivrance/Issue Date: 2003/10/21
 (85) Entrée phase nationale/National Entry: 1999/12/17
 (86) N° demande PCT/PCT Application No.: US 1998/013783
 (87) N° publication PCT/PCT Publication No.: 1999/001542
 (30) Priorités/Priorities: 1997/07/01 (08/887,114) US;
 1997/07/10 (08/890,853) US

(51) Cl.Int.⁷/Int.Cl.⁷ C12N 15/54, C12Q 1/48, C12N 9/12
 (72) Inventeurs/Inventors:
 GOEDEL, DAVID V., US;
 WORONICZ, JOHN, US
 (73) Propriétaire/Owner:
 TULARIK INC., US
 (74) Agent: ADE & COMPANY

(54) Titre : PROTEINES IKK- β , ACIDES NUCLEIQUES ET PROCÉDES
 (54) Title: IKK- β PROTEINS, NUCLEIC ACIDS AND METHODS

(57) **Abrégé/Abstract:**

The invention provides methods and compositions relating to an I κ B kinase, IKK- β , and related nucleic acids. The polypeptides may be produced recombinantly from transformed host cells from the disclosed IKK- β encoding nucleic acids or purified from human cells. The invention provides isolated IKK- β hybridization probes and primers capable of specifically hybridizing with the disclosed IKK- β genes, IKK- β -specific binding agents such as specific antibodies, and methods of making and using the subject compositions in diagnosis, therapy and in the biopharmaceutical industry.



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ : C12N 9/12, C12Q 1/48	A1	(11) International Publication Number: WO 99/01542 (43) International Publication Date: 14 January 1999 (14.01.99)
(21) International Application Number: PCT/US98/13783 (22) International Filing Date: 1 July 1998 (01.07.98) (30) Priority Data: 08/887,114 1 July 1997 (01.07.97) US 08/890,853 10 July 1997 (10.07.97) US (71) Applicant: TULARIK INC. [US/US]; Two Corporate Drive, South San Francisco, CA 94080 (US). (72) Inventors: GOEDDEL, David, V.; Tularik Inc., Two Corporate Drive, South San Francisco, CA 94080 (US). WORONICZ, John; Tularik Inc., Two Corporate Drive, South San Fran- cisco, CA 94080 (US). (74) Agent: OSMAN, Richard, Aron; Science & Technology Law Group, 75 Denise Drive, Hillsborough, CA 94010 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HR, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: IKK- β PROTEINS, NUCLEIC ACIDS AND METHODS		
(57) Abstract The invention provides methods and compositions relating to an I κ B kinase, IKK- β , and related nucleic acids. The polypeptides may be produced recombinantly from transformed host cells from the disclosed IKK- β encoding nucleic acids or purified from human cells. The invention provides isolated IKK- β hybridization probes and primers capable of specifically hybridizing with the disclosed IKK- β genes, IKK- β -specific binding agents such as specific antibodies, and methods of making and using the subject compositions in diagnosis, therapy and in the biopharmaceutical industry.		

IKK- β Proteins, Nucleic Acids and Methods

INTRODUCTION

Field of the Invention

The field of this invention is proteins involved in transcription factor activation.

5

Background

Cytokines trigger changes in gene expression by modifying the activity of otherwise latent transcription factors (Hill and Treisman, 1995). Nuclear factor κ B (NF- κ B) is a prominent example of how such an external stimulus is converted into an active transcription factor (Verma et al., 1995). The NF- κ B system is composed of homo- and heterodimers of members of the Rel family of related transcription factors that control the expression of numerous immune and inflammatory response genes as well as important viral genes (Lenardo and Baltimore, 1989; Baeuerle and Henkel, 1994). The activity of NF- κ B transcription factors is regulated by their subcellular localization (Verma et al., 15 1995). In most cell types, NF- κ B is present as a heterodimer comprising of a 50 kDa and a 65 kDa subunit. This heterodimer is sequestered in the cytoplasm in association with I κ B α a member of the I κ B family of inhibitory proteins (Finco and Baldwin, 1995; Thanos and Maniatis, 1995; Verma et al., 1995). I κ B α masks the nuclear localization signal of NF- κ B and thereby prevents NF- κ B nuclear translocation. Conversion of NF- κ B into an active transcription factor that translocates into the nucleus and binds to cognate DNA sequences requires the phosphorylation and subsequent ubiquitin-dependent degradation of I κ B α in the 26s proteasome. Signal-induced phosphorylation of I κ B α occurs at serines 32 and 36. Mutation of one or both of these serines renders I κ B α resistant to ubiquitination and proteolytic degradation (Chen et al., 1995).

20

25

The pleiotropic cytokines tumor necrosis factor (TNF) and interleukin-1 (IL-1) are among the physiological inducers of I κ B phosphorylation and subsequent NF- κ B activation (Osborn et al., 1989; Beg et al., 1993). Although TNF and IL-1 initiate signaling cascades leading to NF- κ B activation via distinct families of cell-surface receptors (Smith et al., 1994; Dinarello, 1996), both pathways utilize members of the TNF receptor-associated factor (TRAF) family of adaptor proteins as signal transducers (Rothe 30 et al., 1995; Hsu et al., 1996; Cao et al., 1996b). TRAF proteins were originally found to

30

associate directly with the cytoplasmic domains of several members of the TNF receptor family including the 75 kDa TNF receptor (TNFR2), CD40, CD30, and the lymphotoxin- β receptor (Rothe et al., 1994; Hu et al., 1994; Cheng et al., 1995; Mosialos et al., 1995; Song and Donner, 1995; Sato et al., 1995; Lee et al., 1996; Gedrich et al., 1996; Ansieau et al., 1996). In addition, TRAF proteins are recruited indirectly to the 55 kDa TNF receptor (TNFR1) by the adaptor protein TRADD (Hsu et al., 1996). Activation of NF- κ B by TNF requires TRAF2 (Rothe et al., 1995; Hsu et al., 1996). TRAF5 has also been implicated in NF- κ B activation by members of the TNF receptor family (Nakano et al., 1996). In contrast, TRAF6 participates in NF- κ B activation by IL-1 (Cao et al., 1996b). Upon IL-1 treatment, TRAF6 associates with IRAK, a serine-threonine kinase that binds to the IL-1 receptor complex (Cao et al., 1996a).

The NF- κ B-inducing kinase (NIK) is a member of the MAP kinase kinase kinase (MAP3K) family that was identified as a TRAF2-interacting protein (Malinin et al., 1997). NIK activates NF- κ B when overexpressed, and kinase-inactive mutants of NIK comprising its TRAF2-interacting C-terminal domain (NIK₍₆₂₄₋₉₄₇₎) or lacking two crucial lysine residues in its kinase domain (NIK_(KK429-430AA)) behave as dominant-negative inhibitors that suppress TNF-, IL-1-, and TRAF2-induced NF- κ B activation (Malinin et al., 1997). Recently, NIK was found to associate with additional members of the TRAF family, including TRAF5 and TRAF6. Catalytically inactive mutants of NIK also inhibited TRAF5- and TRAF6-induced NF- κ B activation, thus providing a unifying concept for NIK as a common mediator in the NF- κ B signaling cascades triggered by TNF and IL-1 downstream of TRAFs.

Here, we disclose a novel kinase I κ B Kinase, IKK- β , as a NIK-interacting protein. IKK- β has sequence similarity to the conceptual translate of a previously identified open reading frame postulated to encode a serine-threonine kinase of unknown function ('Conserved Helix-loop-helix Ubiquitous Kinase' or CHUK, Connelly and Marcu, 1995; Mock et al., 1995). Catalytically inactive mutants of IKK- β are shown to suppress NF- κ B activation induced by TNF and IL-1 stimulation as well as by TRAF and NIK overexpression; transiently expressed IKK- β is shown to associate with the endogenous I κ B α complex; and IKK- β is shown to phosphorylate I κ B α on serines 32 and 36. As used herein, Ser32 and Ser36 of I κ B refers collectively to the two serine residues which are part of the consensus sequence DSGL/IXSM/L (e.g. ser 32 and 36 in I κ B α , ser 19 and 23 in

I κ B β , and ser 157 and 161, or 18 and 22, depending on the usage of methionines, in I κ B ϵ , respectively.

SUMMARY OF THE INVENTION

The invention provides methods and compositions relating to isolated IKK- β polypeptides, related nucleic acids, polypeptide domains thereof having IKK- β -specific structure and activity and modulators of IKK- β function, particularly I κ B kinase activity. IKK- β polypeptides can regulate NF κ B activation and hence provide important regulators of cell function. The polypeptides may be produced recombinantly from transformed host cells from the subject IKK- β polypeptide encoding nucleic acids or purified from mammalian cells. The invention provides isolated IKK- β hybridization probes and primers capable of specifically hybridizing with the disclosed IKK- β gene, IKK- β -specific binding agents such as specific antibodies, and methods of making and using the subject compositions in diagnosis (e.g. genetic hybridization screens for IKK- β transcripts), therapy (e.g. IKK- β kinase inhibitors to inhibit TNF signal transduction) and in the biopharmaceutical industry (e.g. as immunogens, reagents for isolating other transcriptional regulators, reagents for screening chemical libraries for lead pharmacological agents, etc.).

According to a first aspect of the invention, there is provided an isolated IKK β polypeptide comprising at least 31 consecutive residues of the amino acid sequence set forth as SEQ ID NO: 2.

According to a second aspect of the invention, there is provided an isolated IKK β polypeptide made by a method comprising the steps of: introducing a recombinant nucleic acid encoding the polypeptide described above into a host cell or cellular extract, incubating the host cell or cellular extract under conditions whereby the polypeptide is expressed; and isolating the polypeptide.

According to a third aspect of the invention, there is provided an isolated or recombinant nucleic acid comprising the sequence set forth as SEQ ID NO: 1.

According to a fourth aspect of the invention, there is provided an isolated or recombinant nucleic acid encoding the polypeptide described above.

According to a fifth aspect of the invention, there is provided an isolated cell comprising the recombinant nucleic acid described above.

According to a sixth aspect of the invention, there is provided a method of making an isolated polypeptide, said method comprising the steps of: introducing the recombinant nucleic acid described above into a host cell or cellular extract; incubating

said host cell or cellular extract under conditions whereby said polypeptide is expressed; and isolating said polypeptide.

According to a seventh aspect of the invention, there is provided a method of screening for an agent which modulates the interaction of an IKK β polypeptide to a binding target, the method comprising the steps of: incubating a mixture comprising: the polypeptide as described above, a binding target of the polypeptide, and a candidate agent, under conditions whereby, but for the presence of the agent, the polypeptide specifically binds said binding target at a reference affinity; and detecting the binding affinity of the polypeptide to the binding target to determine an agent-biased affinity, wherein a difference between the agent-biased affinity and the reference affinity indicates that the agent modulates the binding of the polypeptide to the binding target, wherein the binding target comprises (i) an I κ B polypeptide domain, (ii) an IKK polypeptide domain, (iii) a NIK polypeptide domain or (iv) a natural intracellular substrate and the reference and agent-biased binding affinity is detected as phosphorylation of the substrate.

DETAILED DESCRIPTION OF THE INVENTION

The nucleotide sequence of a natural cDNA encoding a human IKK- β polypeptide is shown as SEQ ID NO:1, and the full conceptual translate is shown as SEQ ID NO:2. The IKK- β polypeptides of the invention include incomplete translates of SEQ ID NO:1 and deletion mutants of SEQ ID NO:2, which translates and deletion mutants have IKK- β -specific amino acid sequence and binding specificity or function.

The IKK- β polypeptide domains of the invention have amino acid sequences distinguishable from IKK- α , generally at least 11, preferably at least 21, more preferably at least 31, more preferably at least 41 consecutive residues of SEQ ID NO:2, particularly of SEQ ID NO:2, residues 1-190 and residues 217-756 and provide IKK- β domain specific activity or function, such as IKK- β -specific kinase or kinase inhibitory activity, NIK-binding or binding inhibitory activity, I κ B-binding or binding inhibitory activity, NF κ B activating or inhibitory activity or antibody binding. Preferred domains phosphorylate at

least one and preferably both the serine 32 and 36 of IκB (Verma, I. M., et al. (1995)).

IKK-β-specific activity or function may be determined by convenient *in vitro*, cell-based, or *in vivo* assays: e.g. *in vitro* binding assays, cell culture assays, in animals (e.g. gene therapy, transgenics, etc.), etc. Binding assays encompass any assay where the molecular interaction of an IKK-β polypeptide with a binding target is evaluated. The binding target may be a natural intracellular binding target such as an IKK-β substrate, a IKK-β regulating protein or other regulator that directly modulates IKK-β activity or its localization; or non-natural binding target such a specific immune protein such as an antibody, or an IKK-β specific agent such as those identified in screening assays such as described below. IKK-β-binding specificity may assayed by kinase activity or binding equilibrium constants (usually at least about $10^7 M^{-1}$, preferably at least about $10^8 M^{-1}$, more preferably at least about $10^9 M^{-1}$), by the ability of the subject polypeptide to function as negative mutants in IKK-β-expressing cells, to elicit IKK-β specific antibody in a heterologous host (e.g a rodent or rabbit), etc. In any event, the IKK-β binding specificity of the subject IKK-β polypeptides necessarily distinguishes IKK-α (SEQ ID NO:4). Exemplary IKK-β polypeptides having IKK-β binding specificity are shown in Table 1.

TABLE 1. Exemplary IKK-β polypeptides having IKK-β binding specificity

hIKK-βΔ1 (SEQ ID NO:2, residues 1-9)	hIKK-βΔ1 (SEQ ID NO:2, residues217-229)
hIKK-βΔ1 (SEQ ID NO:2, residues 11-17)	hIKK-βΔ1 (SEQ ID NO:2, residues300-350)
hIKK-βΔ1 (SEQ ID NO:2, residues 21-29)	hIKK-βΔ1 (SEQ ID NO:2, residues419-444)
hIKK-βΔ1 (SEQ ID NO:2, residues 42-51)	hIKK-βΔ1 (SEQ ID NO:2, residues495-503)
hIKK-βΔ1 (SEQ ID NO:2, residues 73-89)	hIKK-βΔ1 (SEQ ID NO:2, residues565-590)
hIKK-βΔ1 (SEQ ID NO:2, residues 90-99)	hIKK-βΔ1 (SEQ ID NO:2, residues610-627)
hIKK-βΔ1 (SEQ ID NO:2, residues120-130)	hIKK-βΔ1 (SEQ ID NO:2, residues677-690)
hIKK-βΔ1 (SEQ ID NO:2, residues155-164)	hIKK-βΔ1 (SEQ ID NO:2, residues715-740)
hIKK-βΔ1 (SEQ ID NO:2, residues180-190)	hIKK-βΔ1 (SEQ ID NO:2, residues747-756)

The claimed IKK-β polypeptides are isolated or pure: an "isolated" polypeptide is unaccompanied by at least some of the material with which it is associated in its natural state, preferably constituting at least about 0.5%, and more preferably at least about 5% by weight of the total polypeptide in a given sample and a pure polypeptide constitutes at

least about 90%, and preferably at least about 99% by weight of the total polypeptide in a given sample. The IKK- β polypeptides and polypeptide domains may be synthesized, produced by recombinant technology, or purified from mammalian, preferably human cells. A wide variety of molecular and biochemical methods are available for biochemical synthesis, molecular expression and purification of the subject compositions, see e.g. Molecular Cloning, A Laboratory Manual (Sambrook, *et al.* Cold Spring Harbor Laboratory), Current Protocols in Molecular Biology (Eds. Ausubel, *et al.*, Greene Publ. Assoc., Wiley-Interscience, NY) or that are otherwise known in the art.

The invention provides binding agents specific to the subject kinase proteins including substrates, agonists, antagonists, natural intracellular binding targets, etc., methods of identifying and making such agents, and their use in diagnosis, therapy and pharmaceutical development. For example, specific binding agents are useful in a variety of diagnostic and therapeutic applications, especially where disease or disease prognosis is associated with improper utilization of a pathway involving the subject proteins, e.g. NF- κ B activation. Novel IKK- β -specific binding agents include IKK- β -specific receptors, such as somatically recombined polypeptide receptors like specific antibodies or T-cell antigen receptors (see, e.g. Harlow and Lane (1988) Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory) and other natural intracellular binding agents identified with assays such as one-, two- and three-hybrid screens, non-natural intracellular binding agents identified in screens of chemical libraries such as described below, etc. Agents of particular interest modulate IKK- β function, e.g. IKK- β -dependent transcriptional activation. For example, a wide variety of inhibitors of IKK- β I κ B kinase activity may be used to regulate signal transduction involving I κ B. Exemplary IKK- β I κ B kinase inhibitors include known classes of serine/threonine kinase (e.g. PKC) inhibitors such as competitive inhibitors of ATP and substrate binding, antibiotics, IKK- β -derived peptide inhibitors, etc., see Tables 2 and 3. IKK- β specificity and activity are readily quantified in high throughput kinase assays using panels of protein kinases (see cited references and Examples).

Preferred inhibitors include natural compounds such as staurosporine (Omura S, *et al.* J Antibiot (Tokyo) 1995 Jul;48(7):535-48), produced by a marine organism, and synthetic compounds such as PD 153035, which also potently inhibits the EGF receptor protein kinase (Fry DW *et al.* Science 1994 Aug 19;265(5175):1093-5). Members of the

tyrphostin family of synthetic protein kinase inhibitors are also useful; these include compounds which are pure ATP competitors, compounds which are pure substrate competitors, and compounds which are mixed competitors: compete with both ATP and substrate (Levitzki A and Gazit A, Science 1995 Mar 24;267(5205):1782-8). Additional IKK- β inhibitors include peptide-based substrate competitors endogenously made by the mammalian cell, e.g. PKI (protein kinase inhibitor, Seasholtz AF et al., Proc Natl Acad Sci USA 1995 Feb 28;92(5):1734-8), or proteins inhibiting cdc kinases (Correa-Bordes J and Nurse P, Cell 1995 Dec 15;83(6):1001-9). Additional small peptide based substrate competitive kinase inhibitors and allosteric inhibitors (inhibitory mechanisms independent of ATP or substrate competition) are readily generated by established methods (Hvalby O, et al. Proc Natl Acad Sci USA 1994 May 24;91(11):4761-5; Barja P, et al., Cell Immunol 1994 Jan;153(1):28-38; Villar-Palasi C, Biochim Biophys Acta 1994 Dec 30;1224(3):384-8; Liu WZ, et al., Biochemistry 1994 Aug 23;33(33):10120-6).

TABLE 2. Selected Small Molecule IKK- β Kinase Inhibitors

15	HA-100 ¹	Iso-H7 ¹²	A-3 ¹⁸
	Chelerythrine ²	PKC 19-31	HA1004 ^{19,20}
	Staurosporine ^{3,4,5}	H-7 ^{13,3,14}	K-252a ^{16,5}
	Calphostin C ^{6,7,8,9}	H-89 ¹⁵	KT5823 ¹⁶
	K-252b ¹⁰	KT5720 ¹⁶	ML-9 ²¹
20	PKC 19-36 ¹¹	cAMP-depPKinh ¹⁷	KT5926 ²²

Citations

1. Hagiwara, M., et al. Mol. Pharmacol. 32: 7 (1987)
2. Herbert, J. M., et al. Biochem Biophys Res Com 172: 993 (1990)
3. Schachtele, C., et al. Biochem Biophys Res Com 151: 542 (1988)
- 25 4. Tamaoki, T., et al. Biochem Biophys Res Com 135: 397 (1986)
5. Tischler, A. S., et al. J. Neurochemistry 55: 1159 (1990)
6. Bruns, R. F., et al. Biochem Biophys Res Com 176: 288 (1991)
7. Kobayashi, E., et al. Biochem Biophys Res Com 159: 548 (1989)
8. Tamaoki, T., et al. Adv2nd Mass Phosphoprotein Res 24:497(1990)
- 30 9. Tamaoki, T., et al. Biotechnology 8: 732 (1990)

10. Yasuzawa, T. J. Antibiotics 39: 1972 (1986)
11. House, C., et al. Science 238: 1726 (1987)
12. Quick, J., et al. Biochem. Biophys. Res. Com. 167: 657 (1992)
13. Bouli, N. M. and Davis, M. Brain Res. 525: 198 (1990)
14. Takahashi, I., et al. J. Pharmacol. Exp. Ther. 255: 1218 (1990)
- 5 15. Chijiwa, T., et al. J. Biol. Chem. 265: 5267 (1990)
16. Kase, H., et al. Biochem. Biophys. Res. Com. 142: 436 (1987)
17. Cheng, H. C., et al. J. Biol. Chem. 261: 989 (1986)
18. Inagaki, M., et al. Mol. Pharmacol. 29: 577 (1986)
19. Asano, T. and Hidaka, H. J Pharmaco. Exp Ther 231:141 (1984)
- 10 20. Hidaka, H., et al. Biochemistry 23: 5036 (1984)
21. Nagatsu, T., et al. Biochem Biophys Res Com 143:1045 (1987)
22. Nakanishi, S., et al. Mol. Pharmacol. 37: 482 (1990)

TABLE 3. Selected Peptidyl IKK- β Kinase Inhibitors

15	hI κ B α , residues 24-39, 32Ala	hIKK- β , Δ 5-203
	hI κ B α , residues 29-47, 36Ala	hIKK- β , Δ 1-178
	hI κ B α , residues 26-46, 32/36Ala	hIKK- β , Δ 368-756
	hI κ B β , residues 25-38, 32Ala	hIKK- β , Δ 460-748
	hI κ B β , residues 30-41, 36Ala	hIKK- α , Δ 1-289
20	hI κ B β , residues 26-46, 32/36Ala	hIKK- α , Δ 12-219
	hI κ B ϵ , residues 24-40, 32Ala	hIKK- α , Δ 307-745
	hI κ B ϵ , residues 31-50, 36Ala	hIKK- α , Δ 319-644
	hI κ B ϵ , residues 27-44, 32/36Ala	

Accordingly, the invention provides methods for modulating signal transduction involving I κ B in a cell comprising the step of modulating IKK- β kinase activity, e.g. by contacting the cell with a serine/threonine kinase inhibitor. The cell may reside in culture or in situ, i.e. within the natural host. Preferred inhibitors are orally active in mammalian hosts. For diagnostic uses, the inhibitors or other IKK- β binding agents are frequently labeled, such as with fluorescent, radioactive, chemiluminescent, or other easily detectable molecules, either conjugated directly to the binding agent or conjugated to a probe specific

for the binding agent.

The amino acid sequences of the disclosed IKK- β polypeptides are used to back-translate IKK- β polypeptide-encoding nucleic acids optimized for selected expression systems (Holler et al. (1993) Gene 136, 323-328; Martin et al. (1995) Gene 154, 150-166) or used to generate degenerate oligonucleotide primers and probes for use in the isolation of natural IKK- β -encoding nucleic acid sequences ("GCG" software, Genetics Computer Group, Inc, Madison WI). IKK- β -encoding nucleic acids used in IKK- β -expression vectors and incorporated into recombinant host cells, e.g. for expression and screening, transgenic animals, e.g. for functional studies such as the efficacy of candidate drugs for disease associated with IKK- β -modulated cell function, etc.

The invention also provides nucleic acid hybridization probes and replication / amplification primers having a IKK- β cDNA specific sequence contained in SEQ ID NO:1 and sufficient to effect specific hybridization thereto (i.e. specifically hybridize with SEQ ID NO:1 in the presence of IKK- α cDNA, SEQ ID NO:3). Such primers or probes comprise at least 12, preferably at least 24, more preferably at least 36 and most preferably at least 96 bases of SEQ ID NO:1, particularly of SEQ ID NO:1, nucleotides 1-567 and nucleotides 693-2268. Demonstrating specific hybridization generally requires stringent conditions, for example, hybridizing in a buffer comprising 30% formamide in 5 x SSPE (0.18 M NaCl, 0.01 M NaPO₄, pH7.7, 0.001 M EDTA) buffer at a temperature of 42°C and remaining bound when subject to washing at 42°C with 0.2 x SSPE; preferably hybridizing in a buffer comprising 50% formamide in 5 x SSPE buffer at a temperature of 42°C and remaining bound when subject to washing at 42°C with 0.2 x SSPE buffer at 42°C. IKK- β nucleic acids can also be distinguished using alignment algorithms, such as BLASTX (Altschul *et al.* (1990) Basic Local Alignment Search Tool, J Mol Biol 215, 403-410).

The subject nucleic acids are of synthetic/non-natural sequences and/or are isolated, i.e. unaccompanied by at least some of the material with which it is associated in its natural state, preferably constituting at least about 0.5%, preferably at least about 5% by weight of total nucleic acid present in a given fraction, and usually recombinant, meaning they comprise a non-natural sequence or a natural sequence joined to nucleotide(s) other than that which it is joined to on a natural chromosome. Recombinant nucleic acids comprising the nucleotide sequence of SEQ ID NO:1 or fragments thereof,

contain such sequence or fragment at a terminus, immediately flanked by a sequence other than that which it is joined to on a natural chromosome, or flanked by a native flanking region fewer than 10 kb, preferably fewer than 2 kb, which is at a terminus or is immediately flanked by a sequence other than that which it is joined to on a natural chromosome. While the nucleic acids are usually RNA or DNA, it is often advantageous to use nucleic acids comprising other bases or nucleotide analogs to provide modified stability, etc.

The subject nucleic acids find a wide variety of applications including use as translatable transcripts, hybridization probes, PCR primers, diagnostic nucleic acids, etc.; use in detecting the presence of IKK- β genes and gene transcripts and in detecting or amplifying nucleic acids encoding additional IKK- β homologs and structural analogs. In diagnosis, IKK- β hybridization probes find use in identifying wild-type and mutant IKK- β alleles in clinical and laboratory samples. Mutant alleles are used to generate allele-specific oligonucleotide (ASO) probes for high-throughput clinical diagnoses. In therapy, therapeutic IKK- β nucleic acids are used to modulate cellular expression or intracellular concentration or availability of active IKK- β .

The invention provides efficient methods of identifying agents, compounds or lead compounds for agents active at the level of a IKK- β modulatable cellular function. Generally, these screening methods involve assaying for compounds which modulate IKK- β interaction with a natural IKK- β binding target, in particular, IKK- β phosphorylation of I κ B-derived substrates, particularly I κ B and NIK substrates. A wide variety of assays for binding agents are provided including labeled *in vitro* protein-protein binding assays, immunoassays, cell based assays, etc. The methods are amenable to automated, cost-effective high throughput screening of chemical libraries for lead compounds. Identified reagents find use in the pharmaceutical industries for animal and human trials; for example, the reagents may be derivatized and rescreened in *in vitro* and *in vivo* assays to optimize activity and minimize toxicity for pharmaceutical development.

In vitro binding assays employ a mixture of components including an IKK- β polypeptide, which may be part of a fusion product with another peptide or polypeptide, e.g. a tag for detection or anchoring, etc. The assay mixtures comprise a natural intracellular IKK- β binding target. In a particular embodiment, the binding target is a substrate comprising I κ B serines 32 and/or 36. Such substrates comprise a I κ B α , β or ϵ

peptide including the serine 32 and/or 36 residue and at least 5, preferably at least 10, and more preferably at least 20 naturally occurring immediately flanking residues on each side (e.g. for serine 36 peptides, residues 26-46, 22-42, or 12-32 or 151-171 for $\text{I}\kappa\text{B}\alpha$, β or ϵ -derived substrates, respectively). While native full-length binding targets may be used, it is frequently preferred to use portions (e.g. peptides) thereof so long as the portion provides binding affinity and avidity to the subject IKK- β polypeptide conveniently measurable in the assay. The assay mixture also comprises a candidate pharmacological agent.

Candidate agents encompass numerous chemical classes, though typically they are organic compounds; preferably small organic compounds and are obtained from a wide variety of sources including libraries of synthetic or natural compounds. A variety of other reagents may also be included in the mixture. These include reagents like ATP or ATP analogs (for kinase assays), salts, buffers, neutral proteins, e.g. albumin, detergents, protease inhibitors, nuclease inhibitors, antimicrobial agents, etc. may be used.

The resultant mixture is incubated under conditions whereby, but for the presence of the candidate pharmacological agent, the IKK- β polypeptide specifically binds the cellular binding target, portion or analog with a reference binding affinity. The mixture components can be added in any order that provides for the requisite bindings and incubations may be performed at any temperature which facilitates optimal binding. Incubation periods are likewise selected for optimal binding but also minimized to facilitate rapid, high-throughput screening.

After incubation, the agent-biased binding between the IKK- β polypeptide and one or more binding targets is detected by any convenient way. For IKK- β kinase assays, 'binding' is generally detected by a change in the phosphorylation of a IKK- β substrate. In this embodiment, kinase activity may be quantified by the transfer to the substrate of a labeled phosphate, where the label may provide for direct detection as radioactivity, luminescence, optical or electron density, etc. or indirect detection such as an epitope tag, etc. A variety of methods may be used to detect the label depending on the nature of the label and other assay components, e.g. through optical or electron density, radiative emissions, nonradiative energy transfers, etc. or indirectly detected with antibody conjugates, etc.

A difference in the binding affinity of the IKK- β polypeptide to the target in the absence of the agent as compared with the binding affinity in the presence of the agent indicates that the agent modulates the binding of the IKK- β polypeptide to the IKK- β

binding target. Analogously, in the cell-based assay also described below, a difference in IKK- β -dependent transcriptional activation in the presence and absence of an agent indicates the agent modulates IKK- β function. A difference, as used herein, is statistically significant and preferably represents at least a 50%, more preferably at least a 90% difference.

5 The following experimental section and examples are offered by way of illustration and not by way of limitation.

EXPERIMENTAL

Identification of IKK- β

10 To investigate the mechanism of NIK-mediated NF- κ B activation, we initially identified proteins that associate directly with NIK by yeast two-hybrid protein interaction cloning (Fields and Song, 1989). An expression vector was generated that encodes NIK fused to the DNA-binding domain of the yeast transcription factor GAL4. This vector was used as bait in a two-hybrid screen of a human B cell cDNA library. From approximately six million
15 transformants, eight positive clones were obtained, as determined by activation of *his* and *lacZ* reporter genes. Of these clones, three encoded a member of the TRAF family, TRAF3 (Hu et al., 1994; Cheng et al., 1995; Mosialos et al., 1995; Sato et al., 1995) and one encoded a novel protein we call IKK- α . Retransformation into yeast cells verified the interaction between NIK and IKK- α . A full-length human IKK- α clone was isolated by screening a Jurkat cDNA
20 library with a probe generated from the 5'-end of the IKK- α two-hybrid clone. IKK- α comprises an N-terminal serine-threonine kinase catalytic domain, a C-terminal helix-loop-helix domain and a leucine zipper-like amphipathic α -helix juxtaposed in between the helix-loop-helix and kinase domain.

 To identify potential IKK- α -related genes, we searched a public database of human
25 expressed sequence tags (ESTs). We identified two ESTs (W68756 and AA128064), which we determined were capable of encoding distinct peptides with sequence similarity with IKK- α . IKK- α -related cDNA was cloned by probing a Jurkat cDNA library (human T cell) with an oligonucleotide probe corresponding to sequence from one of the ESTs. Sequence analysis demonstrated that the two ESTs included different fragments of the same gene. The longest
30 cDNA clones obtained had a ~3.2 kb insert (SEQ ID NO:1) and an open reading frame of 756 amino acids (SEQ ID NO:2). We have designed the protein encoded by this cDNA as IKK- β .

Interaction of IKK- β and NIK in Human Cells

The interaction of IKK- β with NIK was confirmed in mammalian cell coimmunoprecipitation assays. Human IKK- β containing an C-terminal Flag epitope tag was transiently coexpressed in 293 human embryonic kidney cells with Myc epitope-tagged NIK. Cell lysates were immunoprecipitated using a monoclonal antibody against the Flag epitope, and coprecipitating NIK was detected by immunoblot analysis with an anti-Myc monoclonal antibody. In this assay, IKK- β was able to coprecipitate NIK confirming the interaction between both proteins as detected for IKK- α by yeast two-hybrid analysis. Also, a deletion mutant IKK- β protein lacking most of the N-terminal kinase domain (IKK- $\beta_{(\Delta 5-203)}$, i.e. 1-4 & 204-756) was able to associate with NIK, indicating that the α -helical C-terminal half of IKK- β mediates the interaction with NIK.

Effect of IKK- β and IKK- β Mutants on NF- κ B Activation

To investigate a possible role for IKK- β in NF- κ B activation, we examined if transient overexpression of IKK- β might activate an NF- κ B-dependent reporter gene. An E-selectin-luciferase reporter construct (Schindler and Baichwal, 1994) and a IKK- β expression vector were cotransfected into HeLa cells. IKK- β expression activated the reporter gene in a dose-dependent manner, with a maximal induction of luciferase activity of about 20-fold compared to vector control. Similar results were obtained in 293 cells, where IKK- β overexpression induced reporter gene activity approximately 20-fold. Thus, IKK- β induces NF- κ B activation when overexpressed.

We next determined the effect of overexpression of kinase-inactive IKK- $\beta_{(\Delta 5-203)}$ that still associates with NIK on signal-induced NF- κ B activation in reporter gene assays in 293 cells. Overexpression of IKK- $\beta_{(\Delta 5-203)}$ blocked TNF- and IL-1-induced reporter gene activation similar to overexpression of NIK₍₆₂₄₋₉₄₇₎. IKK- $\beta_{(\Delta 5-203)}$ was also found to inhibit NF- κ B-dependent reporter gene activity elicited by overexpression of TRAF2, TRAF6 and NIK. Taken together these results demonstrate that a catalytically inactive IKK- β mutant is a dominant-negative inhibitor of TNF-, IL-1, TRAF- and NIK-induced NF- κ B activation. This indicates that IKK- β functions as a common mediator of NF- κ B activation by TNF and IL-1 downstream of NIK.

Parentetical References

- 30 Ansieau, S., et al. (1996). Proc. Natl. Acad. Sci. USA 93, 14053-14058.
Baeuerle, P. A., and Henkel, T. (1994). Annu. Rev. Immunol. 12, 141-179.

- Beg, A. A., et al. (1993). *Mol. Cell. Biol.* 13, 3301-3310.
- Cao, Z., Henzel, W. J., and Gao, X. (1996a). *Science* 271, 1128-1131.
- Cao, Z., et al. (1996b). *Nature* 383, 443-446.
- Chen, Z., et al. (1995). *Genes Dev.* 9, 1586-1597.
- Cheng, G., et al. (1995). *Science* 267, 1494-1498.
- 5 Connelly, M. A., and Marcu, K. B. (1995). *Cell. Mol. Biol. Res.* 41, 537-549.
- Dinarello, C. A. (1996). Biologic basis for interleukin-1 in disease. *Blood* 87, 2095-2147.
- Fields, S., and Song, O.-k. (1989). *Nature* 340, 245-246.
- Finco, T. S., and Baldwin, A. S. (1995). *Immunity* 3, 263-272.
- Gedrich, R. W., et al. (1996). *J. Biol. Chem.* 271, 12852-12858.
- 10 Hill, C. S., and Treisman, R. (1995). *Cell* 80, 199-211.
- Hsu, H., Shu, H.-B., Pan, M.-P., and Goeddel, D. V. (1996). *Cell* 84, 299-308.
- Hu, H. M., et al. (1994). *J. Biol. Chem.* 269, 30069-30072.
- Lee, S. Y., et al. (1996). *Proc. Natl. Acad. Sci. USA* 93, 9699-9703.
- Lenardo, M., and Baltimore, D. (1989). *Cell* 58, 227-229.
- 15 Malinin, N. L., et al. (1997). *Nature* 385, 540-544.
- Mock et al. (1995). *Genomics* 27, 348-351.
- Mosialos, G., et al. (1995). *Cell* 80, 389-399.
- Nakano, H., et al. (1996). *J. Biol. Chem.* 271, 14661-14664.
- Osborn, L., Kunkel, S., and Nabel, G. J. (1989). *Proc. Natl. Acad. Sci. USA* 86, 2336-2340.
- 20 Rothe, M., Sarma, V., Dixit, V. M., and Goeddel, D. V. (1995). *Science* 269, 1424-1427.
- Rothe, M., Wong, S. C., Henzel, W. J., and Goeddel, D. V. (1994). *Cell* 78, 681-692.
- Sato, T., Irie, S., and Reed, J. C. (1995). *FEBS Lett.* 358, 113-118.
- Schindler, U., and Baichwal, V. R. (1994). *Mol. Cell. Biol.* 14, 5820-5831.
- Smith, C. A., Farrah, T., and Goodwin, R. G. (1994). *Cell* 76, 959-962.
- 25 Song, H. Y., and Donner, D. B. (1995). *Biochem. J.* 809, 825-829.
- Thanos, D., and Maniatis, T. (1995). *Cell* 80, 529-532.
- Verma, I. M., et al. (1995). *Genes Dev.* 9, 2723-2735.

EXAMPLES

- 30 1. Protocol for at IKK- β - κ B α phosphorylation assay.
- A. Reagents:

- Neutralite Avidin: 20 µg/ml in PBS.
 - kinase: 10^{-8} - 10^{-5} M IKK-β (SEQ ID NO:2) at 20 µg/ml in PBS.
 - substrate: 10^{-7} - 10^{-4} M biotinylated substrate (21 residue peptide consisting of residues 26-46 of human IκBα) at 40 µg/ml in PBS.
- 5 - Blocking buffer: 5% BSA, 0.5% Tween 20™ in PBS; 1 hour at room temperature.
- Assay Buffer: 100 mM KCl, 10 mM MgCl₂, 1 mM MnCl₂, 20 mM HEPES pH 7.4, 0.25 mM EDTA, 1% glycerol, 0.5% NP-40, 50 mM BME, 1 mg/ml BSA, cocktail of protease inhibitors.
- [³²P]γ-ATP 10x stock: 2 x 10⁻⁵ M cold ATP with 100 µCi [³²P]γ-ATP. Place in the 4°C
- 10 microfridge during screening.
- Protease inhibitor cocktail (1000X): 10 mg Trypsin Inhibitor (BMB # 109894), 10 mg Aprotinin (BMB # 236624), 25 mg Benzamidine (Sigma # B-6506), 25 mg Leupeptin (BMB # 1017128), 10 mg APMSF (BMB # 917575), and 2mM NaVo₃ (Sigma # S-6508) in 10 ml of PBS.
- 15 B. Preparation of assay plates:
- Coat with 120 µl of stock N Avidin per well overnight at 4°C.
 - Wash 2 times with 200 µl PBS.
 - Block with 150 µl of blocking buffer.
 - Wash 2 times with 200 µl PBS.
- 20 C. Assay:
- Add 40 µl assay buffer/well.
 - Add 40 µl biotinylated substrate (2-200 pmoles/40 ul in assay buffer)
 - Add 40 µl kinase (0.1-10 pmoles/40 ul in assay buffer)
 - Add 10 µl compound or extract.
- 25 - Add 10 µl [³²P]γ-ATP 10x stock.
- Shake at 25°C for 15 minutes.
 - Incubate additional 45 minutes at 25°C.
 - Stop the reaction by washing 4 times with 200 µl PBS.
 - Add 150 µl scintillation cocktail.
- 30 - Count in Topcount.
- D. Controls for all assays (located on each plate):
- a. Non-specific binding

b. cold ATP at 80% inhibition.

2. Protocol for high throughput IKK- β -NIK binding assay.

A. Reagents:

- Neutralite Avidin: 20 μ g/ml in PBS.

5 - Blocking buffer: 5% BSA, 0.5% Tween 20 in PBS; 1 hour at room temperature.

- Assay Buffer: 100 mM KCl, 20 mM HEPES pH 7.6, 1 mM MgCl₂, 1% glycerol, 0.5% NP-40, 50 mM β -mercaptoethanol, 1 mg/ml BSA, cocktail of protease inhibitors.

10 - ³³P IKK- β polypeptide 10x stock: 10⁻⁸ - 10⁻⁶ M "cold" IKK- β supplemented with 200,000-250,000 cpm of labeled IKK- β (Beckman counter). Place in the 4°C microfridge during screening.

- Protease inhibitor cocktail (1000X): 10 mg Trypsin Inhibitor (BMB # 109894), 10 mg Aprotinin (BMB # 236624), 25 mg Benzamidine (Sigma # B-6506), 25 mg Leupeptin (BMB # 1017128), 10 mg APMSF (BMB # 917575), and 2mM NaVO₃ (Sigma # S-6508) in 10 ml of PBS.

15 - NIK: 10⁻⁷ - 10⁻⁵ M biotinylated NIK in PBS.

B. Preparation of assay plates:

- Coat with 120 μ l of stock N-Avidin per well overnight at 4°C.

- Wash 2 times with 200 μ l PBS.

- Block with 150 μ l of blocking buffer.

20 - Wash 2 times with 200 μ l PBS.

C. Assay:

- Add 40 μ l assay buffer/well.

- Add 10 μ l compound or extract.

- Add 10 μ l ³³P-IKK- β (20-25,000 cpm/0.1-10 pmoles/well = 10⁻⁹- 10⁻⁷ M final conc).

25 - Shake at 25°C for 15 minutes.

- Incubate additional 45 minutes at 25°C.

- Add 40 μ M biotinylated NIK (0.1-10 pmoles/40 μ l in assay buffer)

- Incubate 1 hour at room temperature.

- Stop the reaction by washing 4 times with 200 μ M PBS.

30 - Add 150 μ M scintillation cocktail.

- Count in Topcount.

- D. Controls for all assays (located on each plate):
 - a. Non-specific binding
 - b. Soluble (non-biotinylated NIK) at 80% inhibition.

- 5 Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT: Goeddel, David V.
Woronicz, John
- (ii) TITLE OF INVENTION: IKK- Proteins, Nucleic Acids and Methods
- 10 (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:
15 (A) ADDRESSEE: SCIENCE & TECHNOLOGY LAW GROUP
(B) STREET: 268 BUSH STREET, SUITE 3200
(C) CITY: SAN FRANCISCO
(D) STATE: CALIFORNIA
(E) COUNTRY: USA
(F) ZIP: 94104
- (v) COMPUTER READABLE FORM:
20 (A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- 25 (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:
- 30 (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: OSMAN, RICHARD A
(B) REGISTRATION NUMBER: 36,627
(C) REFERENCE/DOCKET NUMBER: T97-006-1
- 35 (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: (415) 343-4341
(B) TELEFAX: (415) 343-4342
- 40 (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 2268 base pairs
(B) TYPE: nucleic acid

WO 99/01542

PCT/US98/13783

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	ATGAGCTGGT CACCTTCCCT GACAACGCAG ACATGTGGGG CCTGGGAAAT GAAAGAGCGC	60
	CTTGGGACAG GGGGATTTGG AAATGTCATC CGATGGCACA ATCAGGAAAC AGGTGAGCAG	120
	ATTGCCATCA AGCAGTGCCG GCAGGAGCTC AGCCCCCGGA ACCGAGAGCG GTGGTGCCTG	180
	GAGATCCAGA TCATGAGAAG GCTGACCCAC CCCAATGTGG TGGCTGCCCC AGATGTCCCT	240
10	GAGGGGATGC AGAACTTGGC GCCCAATGAC CTGCCCCTGC TGGCCATGGA GTACTGCCAA	300
	GGAGGAGATC TCCGGAAGTA CCTGAACCAG TTTGAGAACT GCTGTGGTCT GCGGGAAGGT	360
	GCCATCCTCA CCTTGCTGAG TGACATTGCC TCTGCGCTTA GATACCTCA TGAAAACAGA	420
	ATCATCCATC GGGATCTAAA GCCAGAAAAC ATCGTCCTGC AGCAAGGAGA ACAGAGGTTA	480
	ATACACAAAA TTATTGACCT AGGATATGCC AAGGAGCTGG ATCAGGGCAG TCTTTGCACA	540
15	TCATTTCGTGG GGACCCTGCA GTACCTGGCC CCAGAGCTAC TGGAGCAGCA GAAGTACACA	600
	GTGACCGTCC ACTACTGGAG CTTCGGCACC CTGGCCTTTG AGTGCATCAC GGGCTTCCGG	660
	CCCTTCCTCC CCAACTGGCA GCCCGTGCAG TGGCATTCAA AAGTGCGGCA GAAGAGTGAG	720
	GTGGACATTG TTGTTAGCGA AGACTTGAAT GGAACGGTGA AGTTTTCAAG CTCTTTACCC	780
	TACCCCAATA ATCTTAACAG TGTCCTGGCT GAGCGACTGG AGAAGTGGCT GCAACTGATG	840
20	CTGATGTGGC ACCCCCGACA GAGGGGCACG GATCCCACGT ATGGGCCCAA TGGCTGCTTC	900
	AAGGCCCTGG ATGACATCTT AAACCTAAAG CTGGTTCATA TCTTGAACAT GGTACCGGGC	960
	ACCATCCACA CCTACCCTGT GACAGAGGAT GAGAGTCTGC AGAGCTTGAA GGCCAGAATC	1020
	CAACAGGACA CGGGCATCCC AGAGGAGGAC CAGGAGCTGC TGCAGGAAGC GGGCCTGGCG	1080
	TTGATCCCCG ATAAGCCTGC CACTCAGTGT ATTTGAGACG GCAAGTTAAA TGAGGGCCAC	1140
25	ACATTGGACA TGGATCTTGT TTTTCTCTTT GACAACAGTA AAATCACCTA TGAGACTCAG	1200
	ATCTCCCCAC GGCCCCAACC TGAAAGTGTC AGCTGTATCC TTCAAGAGCC CAAGAGGAAT	1260
	CTCGCCTTCT TCCAGCTGAG GAAGGTGTGG GGCCAGGTCT GGCACAGCAT CCAGACCCTG	1320
	AAGGAAGATT GCAACCGGCT GCAGCAGGGA CAGCGAGCCG CCATGATGAA TCTCCTCCGA	1380
	AACAACAGCT GCCTCTCCAA AATGAAGAAT TCCATGGCTT CCATGTCTCA GCAGCTCAAG	1440
30	GCCAAGTTGG ATTTCTTCAA AACCAGCATC CAGATTGACC TGGAGAAGTA CAGCGAGCAA	1500
	ACCGAGTTTG GGATCACATC AGATAAACTG CTGCTGGCCT GGAGGGAAAT GGAGCAGGCT	1560
	GTGGAGCTCT GTGGGCGGGA GAACGAAGTG AAACCTCCTGG TAGAACGGAT GATGGCTCTG	1620
	CAGACCGACA TTGTGGACTT ACAGAGGAGC CCCATGGGCC GGAAGCAGGG GGGAACGCTG	1680
	GACGACCTAG AGGAGCAAGC AAGGGAGCTG TACAGGAGAC TAAGGGAAAA ACCTCGAGAC	1740
35	CAGCGAACTG AGGGTGACAG TCAGGAAATG GTACGGCTGC TGCTTCAGGC AATTCAGAGC	1800
	TTCGAGAAGA AAGTGCGAGT GATCTATACG CAGCTCAGTA AAACCTGTGGT TTGCAAGCAG	1860
	AAGGCGCTGG AACTGTTGCC CAAGGTGGAA GAGGTGGTGA GCTTAATGAA TGAGGATGAG	1920
	AAGACTGTTG TCCGGCTGCA GGAGAAGCGG CAGAAGGAGC TCTGGAATCT CCTGAAGATT	1980
	GCTTGTAGCA AGGTCCGTGG TCCTGTCAGT GGAAGCCCGG ATAGCATGAA TGCCTCTCGA	2040
40	CTTAGCCAGC CTGGCAGCT GATGTCTCAG CCCTCCACGG CCTCCAACAG CTTACCTGAG	2100
	CCAGCCAAGA AGAGTGAAGA ACTGGTGGCT GAAGCACATA ACCTCTGCAC CCTGCTAGAA	2160
	AATGCCATAC AGGACACTGT GAGGGAACAA GACCAGAGTT TCACGGCCCT AGACTGGAGC	2220
	TGGTTACAGA CGGAAGAAGA AGAGCACAGC TGCCTGGAGC AGGCCTCA	2268

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 756 amino acids
- (B) TYPE: amino acid
- 5 (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

	Met	Ser	Trp	Ser	Pro	Ser	Leu	Thr	Thr	Gln	Thr	Cys	Gly	Ala	Trp	Glu
	1				5					10					15	
	Met	Lys	Glu	Arg	Leu	Gly	Thr	Gly	Gly	Phe	Gly	Asn	Val	Ile	Arg	Trp
				20					25					30		
15	His	Asn	Gln	Glu	Thr	Gly	Glu	Gln	Ile	Ala	Ile	Lys	Gln	Cys	Arg	Gln
			35					40					45			
	Glu	Leu	Ser	Pro	Arg	Asn	Arg	Glu	Arg	Trp	Cys	Leu	Glu	Ile	Gln	Ile
		50					55					60				
	Met	Arg	Arg	Leu	Thr	His	Pro	Asn	Val	Val	Ala	Ala	Arg	Asp	Val	Pro
20	65					70				75					80	
	Glu	Gly	Met	Gln	Asn	Leu	Ala	Pro	Asn	Asp	Leu	Pro	Leu	Leu	Ala	Met
				85					90					95		
	Glu	Tyr	Cys	Gln	Gly	Gly	Asp	Leu	Arg	Lys	Tyr	Leu	Asn	Gln	Phe	Glu
				100					105					110		
25	Asn	Cys	Cys	Gly	Leu	Arg	Glu	Gly	Ala	Ile	Leu	Thr	Leu	Leu	Ser	Asp
				115					120					125		
	Ile	Ala	Ser	Ala	Leu	Arg	Tyr	Leu	His	Glu	Asn	Arg	Ile	Ile	His	Arg
				130				135					140			
	Asp	Leu	Lys	Pro	Glu	Asn	Ile	Val	Leu	Gln	Gln	Gly	Glu	Gln	Arg	Leu
30	145					150					155				160	
	Ile	His	Lys	Ile	Ile	Asp	Leu	Gly	Tyr	Ala	Lys	Glu	Leu	Asp	Gln	Gly
				165						170					175	
	Ser	Leu	Cys	Thr	Ser	Phe	Val	Gly	Thr	Leu	Gln	Tyr	Leu	Ala	Pro	Glu
				180					185					190		
35	Leu	Leu	Glu	Gln	Gln	Lys	Tyr	Thr	Val	Thr	Val	Asp	Tyr	Trp	Ser	Phe
				195					200					205		
	Gly	Thr	Leu	Ala	Phe	Glu	Cys	Ile	Thr	Gly	Phe	Arg	Pro	Phe	Leu	Pro
				210				215				220				
	Asn	Trp	Gln	Pro	Val	Gln	Trp	His	Ser	Lys	Val	Arg	Gln	Lys	Ser	Glu
40	225					230						235			240	
	Val	Asp	Ile	Val	Val	Ser	Glu	Asp	Leu	Asn	Gly	Thr	Val	Lys	Phe	Ser
				245							250				255	
	Ser	Ser	Leu	Pro	Tyr	Pro	Asn	Asn	Leu	Asn	Ser	Val	Leu	Ala	Glu	Arg
				260					265					270		

WO 99/01542

PCT/US98/13783

	Leu	Glu	Lys	Trp	Leu	Gln	Leu	Met	Leu	Met	Trp	His	Pro	Arg	Gln	Arg
			275					280					285			
	Gly	Thr	Asp	Pro	Thr	Tyr	Gly	Pro	Asn	Gly	Cys	Phe	Lys	Ala	Leu	Asp
			290				295					300				
5	Asp	Ile	Leu	Asn	Leu	Lys	Leu	Val	His	Ile	Leu	Asn	Met	Val	Thr	Gly
	305					310					315					320
	Thr	Ile	His	Thr	Tyr	Pro	Val	Thr	Glu	Asp	Glu	Ser	Leu	Gln	Ser	Leu
					325					330					335	
	Lys	Ala	Arg	Ile	Gln	Gln	Asp	Thr	Gly	Ile	Pro	Glu	Glu	Asp	Gln	Glu
				340					345					350		
10	Leu	Leu	Gln	Glu	Ala	Gly	Leu	Ala	Leu	Ile	Pro	Asp	Lys	Pro	Ala	Thr
			355					360					365			
	Gln	Cys	Ile	Ser	Asp	Gly	Lys	Leu	Asn	Glu	Gly	His	Thr	Leu	Asp	Met
			370				375					380				
15	Asp	Leu	Val	Phe	Leu	Phe	Asp	Asn	Ser	Lys	Ile	Thr	Tyr	Glu	Thr	Gln
	385					390					395					400
	Ile	Ser	Pro	Arg	Pro	Gln	Pro	Glu	Ser	Val	Ser	Cys	Ile	Leu	Gln	Glu
					405					410					415	
	Pro	Lys	Arg	Asn	Leu	Ala	Phe	Phe	Gln	Leu	Arg	Lys	Val	Trp	Gly	Gln
					420				425						430	
20	Val	Trp	His	Ser	Ile	Gln	Thr	Leu	Lys	Glu	Asp	Cys	Asn	Arg	Leu	Gln
			435					440					445			
	Gln	Gly	Gln	Arg	Ala	Ala	Met	Met	Asn	Leu	Leu	Arg	Asn	Asn	Ser	Cys
			450				455					460				
25	Leu	Ser	Lys	Met	Lys	Asn	Ser	Met	Ala	Ser	Met	Ser	Gln	Gln	Leu	Lys
	465					470					475					480
	Ala	Lys	Leu	Asp	Phe	Phe	Lys	Thr	Ser	Ile	Gln	Ile	Asp	Leu	Glu	Lys
					485					490					495	
	Tyr	Ser	Glu	Gln	Thr	Glu	Phe	Gly	Ile	Thr	Ser	Asp	Lys	Leu	Leu	Leu
				500				505					510			
30	Ala	Trp	Arg	Glu	Met	Glu	Gln	Ala	Val	Glu	Leu	Cys	Gly	Arg	Glu	Asn
			515					520					525			
	Glu	Val	Lys	Leu	Leu	Val	Glu	Arg	Met	Met	Ala	Leu	Gln	Thr	Asp	Ile
			530				535					540				
35	Val	Asp	Leu	Gln	Arg	Ser	Pro	Met	Gly	Arg	Lys	Gln	Gly	Gly	Thr	Leu
	545					550					555					560
	Asp	Asp	Leu	Glu	Glu	Gln	Ala	Arg	Glu	Leu	Tyr	Arg	Arg	Leu	Arg	Glu
					565					570					575	
	Lys	Pro	Arg	Asp	Gln	Arg	Thr	Glu	Gly	Asp	Ser	Gln	Glu	Met	Val	Arg
					580					585				590		
40	Leu	Leu	Leu	Gln	Ala	Ile	Gln	Ser	Phe	Glu	Lys	Lys	Val	Arg	Val	Ile
			595					600					605			
	Tyr	Thr	Gln	Leu	Ser	Lys	Thr	Val	Val	Cys	Lys	Gln	Lys	Ala	Leu	Glu
			610				615					620				
	Leu	Leu	Pro	Lys	Val	Glu	Glu	Val	Val	Ser	Leu	Met	Asn	Glu	Asp	Glu

WO 99/01542

PCT/US98/13783

	625		630		635		640									
	Lys	Thr	Val	Val	Arg	Leu	Gln	Glu	Lys	Arg	Gln	Lys	Glu	Leu	Trp	Asn
					645					650					655	
	Leu	Leu	Lys	Ile	Ala	Cys	Ser	Lys	Val	Arg	Gly	Pro	Val	Ser	Gly	Ser
				660						665					670	
5	Pro	Asp	Ser	Met	Asn	Ala	Ser	Arg	Leu	Ser	Gln	Pro	Gly	Gln	Leu	Met
				675						680					685	
	Ser	Gln	Pro	Ser	Thr	Ala	Ser	Asn	Ser	Leu	Pro	Glu	Pro	Ala	Lys	Lys
				690						695					700	
	Ser	Glu	Glu	Leu	Val	Ala	Glu	Ala	His	Asn	Leu	Cys	Thr	Leu	Leu	Glu
10	705					710					715					720
	Asn	Ala	Ile	Gln	Asp	Thr	Val	Arg	Glu	Gln	Asp	Gln	Ser	Phe	Thr	Ala
						725					730					735
	Leu	Asp	Trp	Ser	Trp	Leu	Gln	Thr	Glu	Glu	Glu	Glu	His	Ser	Cys	Leu
						740					745					750
15	Glu	Gln	Ala	Ser												
						755										

(2) INFORMATION FOR SEQ ID NO:3:

- 20 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 2238 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

- 25 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

	ATGGAGCGGC	CCCCGGGGCT	GCGGCCGGGC	GCGGGCGGGC	CCTGGGAGAT	GCGGGAGCGG	60
30	CTGGGCACCG	GCGGCTTCGG	GAACGTCTGT	CTGTACCAGC	ATCGGGAACT	TGATCTCAAA	120
	ATAGCAATTA	AGTCTTGTCG	CCTAGAGCTA	AGTACCAAAA	ACAGAGAACG	ATGGTGCCAT	180
	GAAATCCAGA	TTATGAAGAA	GTTGAACCAT	GCCAATGTTG	TAAAGGCCTG	TGATGTTCTT	240
	GAAGAATTGA	ATATTTTGAT	TCATGATGTG	CCTCTTCTAG	CAATGGAATA	CTGTTCTGGA	300
	GGAGATCTCC	GAAAGCTGCT	CAACAAACCA	GAAAATTGTT	GTGGACTTAA	AGAAAGCCAG	360
35	ATACTTTCTT	TACTAAGTGA	TATAGGGTCT	GGGATTCGAT	ATTTGCATGA	AAACAAAATT	420
	ATACATCGAG	ATCTAAAACC	TGAAAACATA	GTTCTTCAGG	ATGTTGGTGG	AAAGATAATA	480
	CATAAAATAA	TTGATCTGGG	ATATGCCAAA	GATGTTGATC	AAGGAAGTCT	GTGTACATCT	540
	TTTGTGGGAA	CACTGCAGTA	TCTGGCCCCA	GAGCTCTTTG	AGAATAAGCC	TTACACAGCC	600
	ACTGTTGATT	ATTGGAGCTT	TGGGACCATG	GTATTTGAAT	GTATTGCTGG	ATATAGGCCT	660
40	TTTTTGCATC	ATCTGCAGCC	ATTTACCTGG	CATGAGAAGA	TTAAGAAGAA	GGATCCAAAG	720
	TGTATATTTG	CATGTGAAGA	GATGTCAGGA	GAAGTTCGGT	TTAGTAGCCA	TTTACCTCAA	780
	CCAAATAGCC	TTTGTAGTTT	AATAGTAGAA	CCCATGGAAA	ACTGGCTACA	GTTGATGTTG	840
	AATTGGGACC	CTCAGCAGAG	AGGAGGACCT	GTTGACCTTA	CTTTGAAGCA	GCCAAGATGT	900
	TTTGTATTAA	TGGATCACAT	TTTGAATTTG	AAGATAGTAC	ACATCCTAAA	TATGACTTCT	960

WO 99/01542

PCT/US98/13783

5 GCAAAGATAA TTTCTTTTCT GTTACCACCT GATGAAAGTC TTCATTCACT ACAGTCTCGT 1020
 ATTGAGCGTG AAACGTGAAT AAATACTGGT TCTCAAGAAC TTCTTTCAGA GACAGGAATT 1080
 TCTCTGGATC CTCGGAAACC AGCCTCTCAA TGTGTTCTAG ATGGAGTTAG AGGCTGTGAT 1140
 AGCTATATGG TTTATTTGTT TGATAAAAGT AAAACTGTAT ATGAAGGGCC ATTTGCTTCC 1200
 AGAAGTTTAT CTGATTGTGT AAATTATATT GTACAGGACA GCAAAATACA GCTTCCAATT 1260
 ATACAGCTGC GTAAAGTGTG GGCTGAAGCA GTGCACTATG TGTCTGGACT AAAAGAAGAC 1320
 TATAGCAGGC TCTTTCAGGG ACAAGGGCA GCAATGTAA GTCTTCTTAG ATATAATGCT 1380
 AACTTAACAA AAATGAAGAA CACTTTGATC TCAGCATCAC AACAACTGAA AGCTAAATTG 1440
 GAGTTTTTTC ACAAAGCAT TCAGCTTGAC TTGGAGAGAT ACAGCGAGCA GATGACGTAT 1500
 GGGATATCTT CAGAAAAAT GCTAAAAGCA TGGAAAGAAA TGGAAGAAAA GGCCATCCAC 1560
 10 TATGCTGAGG TTGGTGT CAT TGGATACCTG GAGGATCAGA TTATGTCTTT GCATGCTGAA 1620
 ATCATGGAGC TACAGAAGAG CCCCTATGGA AGACGTCAGG GAGACTTGAT GGAATCTCTG 1680
 GAACAGCGTG CCATTGATCT ATATAAGCAG TTA AACACA GACCTTCAGA TCACTCCTAC 1740
 AGTGACAGCA CAGAGATGGT GAAAATCATT GTGCACACTG TGCAGAGTCA GGACCGTGTG 1800
 CTCAAGGAGC TGTTTGGTCA TTTGAGCAAG TTGTTGGGCT GTAAGCAGAA GATTATTGAT 1860
 15 CTA CTACTCCCTA AGGTGGAAGT GGCCCTCAGT AATATCAAAG AAGCTGACAA TACTGT CATG 1920
 TTCATGCAGG GAAAAAGGCA GAAAGAAATA TGGCATCTCC TTAAAATTGC CTGTACACAG 1980
 AGTTCTGCC GGTCCCTTGT AGGATCCAGT CTAGAAGGTG CAGTAACCCC TCAGACATCA 2040
 GCATGGCTGC CCCC GACTTC AGCAGAACAT GATCATTCTC TGTCATGTGT GGTA ACTCCT 2100
 CAAGATGGGG A GACTTCAGC ACAAATGATA GAAGAAAATT TGA ACTGCCT TGGCCATT TA 2160
 20 AGCACTATTA TTCATGAGGC AAATGAGGAA CAGGGCAATA GTATGATGAA TCTTGATTGG 2220
 AGTTGGTTAA CAGAATGA 2238

(2) INFORMATION FOR SEQ ID NO:4:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 745 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

35 Met Glu Arg Pro Pro Gly Leu Arg Pro Gly Ala Gly Gly Pro Trp Glu
 1 5 10 15
 Met Arg Glu Arg Leu Gly Thr Gly Gly Phe Gly Asn Val Cys Leu Tyr
 20 25 30
 Gln His Arg Glu Leu Asp Leu Lys Ile Ala Ile Lys Ser Cys Arg Leu
 40 35 40 45
 Glu Leu Ser Thr Lys Asn Arg Glu Arg Trp Cys His Glu Ile Gln Ile
 50 55 60
 Met Lys Lys Leu Asn His Ala Asn Val Val Lys Ala Cys Asp Val Pro
 65 70 75 80

WO 99/01542

PCT/US98/13783

	Glu	Glu	Leu	Asn	Ile	Leu	Ile	His	Asp	Val	Pro	Leu	Leu	Ala	Met	Glu
					85					90					95	
	Tyr	Cys	Ser	Gly	Gly	Asp	Leu	Arg	Lys	Leu	Leu	Asn	Lys	Pro	Glu	Asn
				100					105					110		
5	Cys	Cys	Gly	Leu	Lys	Glu	Ser	Gln	Ile	Leu	Ser	Leu	Leu	Ser	Asp	Ile
			115					120					125			
	Gly	Ser	Gly	Ile	Arg	Tyr	Leu	His	Glu	Asn	Lys	Ile	Ile	His	Arg	Asp
		130					135					140				
	Leu	Lys	Pro	Glu	Asn	Ile	Val	Leu	Gln	Asp	Val	Gly	Gly	Lys	Ile	Ile
	145				150						155					160
10	His	Lys	Ile	Ile	Asp	Leu	Gly	Tyr	Ala	Lys	Asp	Val	Asp	Gln	Gly	Ser
				165						170					175	
	Leu	Cys	Thr	Ser	Phe	Val	Gly	Thr	Leu	Gln	Tyr	Leu	Ala	Pro	Glu	Leu
			180						185					190		
	Phe	Glu	Asn	Lys	Pro	Tyr	Thr	Ala	Thr	Val	Asp	Tyr	Trp	Ser	Phe	Gly
15			195					200						205		
	Thr	Met	Val	Phe	Glu	Cys	Ile	Ala	Gly	Tyr	Arg	Pro	Phe	Leu	His	His
		210					215					220				
	Leu	Gln	Pro	Phe	Thr	Trp	His	Glu	Lys	Ile	Lys	Lys	Lys	Asp	Pro	Lys
	225					230					235					240
20	Cys	Ile	Phe	Ala	Cys	Glu	Glu	Met	Ser	Gly	Glu	Val	Arg	Phe	Ser	Ser
				245						250					255	
	His	Leu	Pro	Gln	Pro	Asn	Ser	Leu	Cys	Ser	Leu	Ile	Val	Glu	Pro	Met
			260					265						270		
	Glu	Asn	Trp	Leu	Gln	Leu	Met	Leu	Asn	Trp	Asp	Pro	Gln	Gln	Arg	Gly
25			275					280						285		
	Gly	Pro	Val	Asp	Leu	Thr	Leu	Lys	Gln	Pro	Arg	Cys	Phe	Val	Leu	Met
		290					295					300				
	Asp	His	Ile	Leu	Asn	Leu	Lys	Ile	Val	His	Ile	Leu	Asn	Met	Thr	Ser
	305				310						315					320
30	Ala	Lys	Ile	Ile	Ser	Phe	Leu	Leu	Pro	Pro	Asp	Glu	Ser	Leu	His	Ser
				325						330					335	
	Leu	Gln	Ser	Arg	Ile	Glu	Arg	Glu	Thr	Gly	Ile	Asn	Thr	Gly	Ser	Gln
			340						345					350		
	Glu	Leu	Leu	Ser	Glu	Thr	Gly	Ile	Ser	Leu	Asp	Pro	Arg	Lys	Pro	Ala
35			355					360						365		
	Ser	Gln	Cys	Val	Leu	Asp	Gly	Val	Arg	Gly	Cys	Asp	Ser	Tyr	Met	Val
		370					375					380				
	Tyr	Leu	Phe	Asp	Lys	Ser	Lys	Thr	Val	Tyr	Glu	Gly	Pro	Phe	Ala	Ser
	385				390						395					400
40	Arg	Ser	Leu	Ser	Asp	Cys	Val	Asn	Tyr	Ile	Val	Gln	Asp	Ser	Lys	Ile
				405						410					415	
	Gln	Leu	Pro	Ile	Ile	Gln	Leu	Arg	Lys	Val	Trp	Ala	Glu	Ala	Val	His
			420						425				430			
	Tyr	Val	Ser	Gly	Leu	Lys	Glu	Asp	Tyr	Ser	Arg	Leu	Phe	Gln	Gly	Gln

WO 99/01542

PCT/US98/13783

	435		440		445														
	Arg	Ala	Ala	Met	Leu	Ser	Leu	Leu	Arg	Tyr	Asn	Ala	Asn	Leu	Thr	Lys			
	450						455					460							
	Met	Lys	Asn	Thr	Leu	Ile	Ser	Ala	Ser	Gln	Gln	Leu	Lys	Ala	Lys	Leu			
	465					470					475					480			
5	Glu	Phe	Phe	His	Lys	Ser	Ile	Gln	Leu	Asp	Leu	Glu	Arg	Tyr	Ser	Glu			
				485						490						495			
	Gln	Met	Thr	Tyr	Gly	Ile	Ser	Ser	Glu	Lys	Met	Leu	Lys	Ala	Trp	Lys			
				500					505							510			
	Glu	Met	Glu	Glu	Lys	Ala	Ile	His	Tyr	Ala	Glu	Val	Gly	Val	Ile	Gly			
10				515					520							525			
	Tyr	Leu	Glu	Asp	Gln	Ile	Met	Ser	Leu	His	Ala	Glu	Ile	Met	Glu	Leu			
	530						535									540			
	Gln	Lys	Ser	Pro	Tyr	Gly	Arg	Arg	Gln	Gly	Asp	Leu	Met	Glu	Ser	Leu			
	545					550					555					560			
15	Glu	Gln	Arg	Ala	Ile	Asp	Leu	Tyr	Lys	Gln	Leu	Lys	His	Arg	Pro	Ser			
				565							570					575			
	Asp	His	Ser	Tyr	Ser	Asp	Ser	Thr	Glu	Met	Val	Lys	Ile	Ile	Val	His			
				580							585					590			
	Thr	Val	Gln	Ser	Gln	Asp	Arg	Val	Leu	Lys	Glu	Leu	Phe	Gly	His	Leu			
20				595							600					605			
	Ser	Lys	Leu	Leu	Gly	Cys	Lys	Gln	Lys	Ile	Ile	Asp	Leu	Leu	Pro	Lys			
	610						615									620			
	Val	Glu	Val	Ala	Leu	Ser	Asn	Ile	Lys	Glu	Ala	Asp	Asn	Thr	Val	Met			
	625					630						635				640			
25	Phe	Met	Gln	Gly	Lys	Arg	Gln	Lys	Glu	Ile	Trp	His	Leu	Leu	Lys	Ile			
				645												655			
	Ala	Cys	Thr	Gln	Ser	Ser	Ala	Arg	Ser	Leu	Val	Gly	Ser	Ser	Leu	Glu			
				660												670			
	Gly	Ala	Val	Thr	Pro	Gln	Thr	Ser	Ala	Trp	Leu	Pro	Pro	Thr	Ser	Ala			
30				675												685			
	Glu	His	Asp	His	Ser	Leu	Ser	Cys	Val	Val	Thr	Pro	Gln	Asp	Gly	Glu			
	690						695									700			
	Thr	Ser	Ala	Gln	Met	Ile	Glu	Glu	Asn	Leu	Asn	Cys	Leu	Gly	His	Leu			
	705					710						715				720			
35	Ser	Thr	Ile	Ile	His	Glu	Ala	Asn	Glu	Glu	Gln	Gly	Asn	Ser	Met	Met			
				725												735			
	Asn	Leu	Asp	Trp	Ser	Trp	Leu	Thr	Glu										
				740												745			

CLAIMS

1. An isolated IKK β polypeptide comprising at least 31 consecutive residues of the amino acid sequence set forth as SEQ ID NO: 2.
2. An isolated IKK β polypeptide according to claim 1, comprising the amino
5 acid sequence set forth as SEQ ID NO: 2.
3. An isolated IKK β polypeptide made by a method comprising the steps of:
introducing a recombinant nucleic acid encoding the polypeptide of claims 1 or 2
into a host cell or cellular extract,
incubating the host cell or cellular extract under conditions whereby the
10 polypeptide is expressed; and
isolating the polypeptide.
4. An isolated IKK β polypeptide according to claim 3, wherein the method
comprises the steps of:
introducing a recombinant nucleic acid comprising SEQ ID NO: 1 or a fragment
15 thereof sufficient to encode the polypeptide of claim 1 or 2 into a host cell or cellular
extract;
incubating the host cell or cellular extract under conditions whereby the
polypeptide is expressed; and
isolating the polypeptide.
- 20 5. An isolated or recombinant nucleic acid comprising the sequence set forth
as SEQ ID NO: 1.
6. An isolated or recombinant nucleic acid encoding the polypeptide of claim 1
or 2.
7. An isolated cell comprising a recombinant nucleic acid according to claim 5
25 or 6.
8. A method of making an isolated polypeptide, said method comprising the
steps of:
introducing the recombinant nucleic acid of claim 5 or 6 into a host cell or cellular
extract;
30 incubating said host cell or cellular extract under conditions whereby said
polypeptide is expressed; and
isolating said polypeptide.
9. A method of screening for an agent which modulates the interaction of an
IKK β polypeptide to a binding target, the method comprising the steps of:

incubating a mixture comprising: the polypeptide of claim 1, 2, 3 or 4, a binding target of the polypeptide, and a candidate agent, under conditions whereby, but for the presence of the agent, the polypeptide specifically binds said binding target at a reference affinity; and

5 detecting the binding affinity of the polypeptide to the binding target to determine an agent-biased affinity, wherein a difference between the agent-biased affinity and the reference affinity indicates that the agent modulates the binding of the polypeptide to the binding target,

10 wherein the binding target comprises (i) an I κ B polypeptide domain, (ii) an IKK polypeptide domain, (iii) a NIK polypeptide domain or (iv) a natural intracellular substrate and the reference and agent-biased binding affinity is detected as phosphorylation of the substrate.

15 10. A method according to claim 9 wherein the binding target comprises (iv) a natural intracellular substrate and the reference and agent-biased binding affinity is detected as phosphorylation of the substrate, wherein the substrate comprises an I κ B polypeptide domain.

20 11. A method according to claim 9 wherein the binding target comprises (iv) a natural intracellular substrate and the reference and agent-biased binding affinity is detected as phosphorylation of the substrate, wherein the substrate comprises an I κ B polypeptide domain comprising at least one of serine 32 and serine 36, and at least five naturally occurring immediately flanking residues.

25 12. A method according to any one of claims 9, 10 or 11 wherein the binding target comprises (iv) a natural intracellular substrate and the reference and agent-biased binding affinity is detected as phosphorylation of the substrate, wherein the substrate is biotinylated and the detecting step comprises capturing the substrate on an avidin-coated surface.

30 13. A method according to any one of claims 9, 10 or 11 wherein the binding target comprises (iv) a natural intracellular substrate and the reference and agent-biased binding affinity is detected as phosphorylation of the substrate, wherein the mixture further comprises ATP having a labeled phosphate and the detecting step comprises detecting the presence of the labeled phosphate on the substrate.

14. A method according to any one of claims 9, 10 or 11 wherein the binding target comprises (iv) a natural intracellular substrate and the reference and agent-biased

binding affinity is detected as phosphorylation of the substrate, wherein the phosphorylation of the substrate is detected with an antibody.