



US008569210B2

(12) **United States Patent**
Fefer et al.

(10) **Patent No.:** **US 8,569,210 B2**
(45) **Date of Patent:** **Oct. 29, 2013**

(54) **TURFGRASS FUNGICIDE FORMULATION WITH PIGMENT**

(75) Inventors: **Michael Fefer**, Whitby (CA); **Jun Liu**, Oakville (CA); **Tomoki Ruo**, Oakville (CA); **Sonia Edith Hevia**, Mississauga (CA)

(73) Assignee: **Suncor Energy Inc.**, Calgary, Alberta (CA)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 501 days.

(21) Appl. No.: **12/492,863**

(22) Filed: **Jun. 26, 2009**

(65) **Prior Publication Data**

US 2009/0325922 A1 Dec. 31, 2009

Related U.S. Application Data

(60) Provisional application No. 61/075,821, filed on Jun. 26, 2008, provisional application No. 61/147,523, filed on Jan. 27, 2009.

(51) **Int. Cl.**
A01N 55/02 (2006.01)
A01N 25/00 (2006.01)
A61K 31/155 (2006.01)

(52) **U.S. Cl.**
USPC **504/191**; 424/405; 514/184; 514/185

(58) **Field of Classification Search**
USPC 504/191; 514/185, 184; 424/405
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,714,062 A 7/1955 Lockrey
2,786,821 A 3/1957 Gardner
2,870,037 A 1/1959 Converse
3,131,119 A 4/1964 Fordyce
3,615,799 A 10/1971 Gannon
3,689,574 A 9/1972 Engelhart
3,799,758 A 3/1974 Franz
3,950,265 A 4/1976 Albrecht
3,997,322 A 12/1976 Ratledge
4,002,628 A 1/1977 Benefiel
4,015,970 A 4/1977 Hennart
4,094,845 A 6/1978 De Long
4,124,720 A 11/1978 Wenmaekers
4,243,405 A 1/1981 Balasubramanyan
4,431,554 A 2/1984 Baur
4,584,013 A 4/1986 Brunner
4,618,360 A 10/1986 Brunner
4,693,745 A 9/1987 Brunner
4,698,334 A 10/1987 Horriere
4,734,432 A 3/1988 Szego
4,737,515 A 4/1988 Hallenbach et al.
4,761,423 A 8/1988 Szego
4,826,863 A 5/1989 Szego
4,834,908 A 5/1989 Hazen
4,853,026 A 8/1989 Frisch
4,902,333 A 2/1990 Quimby
4,971,840 A 11/1990 Boho

5,084,087 A 1/1992 Hazen
5,102,442 A 4/1992 Hazen et al.
5,137,726 A 8/1992 Ogawa
5,178,795 A 1/1993 Roberts
5,229,356 A 7/1993 Tocker
5,238,604 A 8/1993 Hazen
5,308,827 A 5/1994 Sakamoto
5,336,661 A 8/1994 Lucas
5,352,729 A 10/1994 Birkhofer
5,362,167 A 11/1994 Loftin
5,393,791 A 2/1995 Roberts
5,409,885 A 4/1995 Derian
5,504,054 A 4/1996 Murphy
5,547,918 A 8/1996 Newton
5,558,806 A * 9/1996 Policello et al. 516/204
5,580,567 A 12/1996 Roberts
5,599,768 A 2/1997 Hermansky
5,599,804 A 2/1997 Mudge
5,643,852 A 7/1997 Lucas
5,658,851 A 8/1997 Murphy
5,665,672 A 9/1997 Lucas

(Continued)

FOREIGN PATENT DOCUMENTS

CA 964482 3/1975
CA 2069311 A1 4/1991

(Continued)

OTHER PUBLICATIONS

Agnello, Petroleum-derived spray oils: chemistry, history, refining and formulation, in Beattie, G.A.C., Watson, D.M., Stevens, M., Spooner-Hart, R. and Rae, D.J. (eds). Spray Oils Beyond 2000—Sustainable Pest & Disease Management. University of Western Sydney, 2002.

Bakke, Analysis of Issues Surrounding the Use of Spray Adjuvants With Herbicides, 2002.

Blenis et al. Evaluation of Fungicides and Surfactants for Control of Fairy Rings Caused by Marasmius oreades (Bolt ex. Fr.) Fr., HortScience, 1997, 1077-1084, 32(6).

Burpee et al., Interactive Effects of Plant Growth Regulators and Fungicides on Epidemics of Dollar Spot in Creeping Bentgrass, Plant Disease, 1996, 1245-1250, 80(11).

Burpee and Latin, Reassessment of Fungicide Synergism for Control of Dollar Spot, Plant Disease, 2008, 601-606, 92(4).

Cline, OLR mating disruption just got easier, Western Farm Press, 2001, 1, 23(12).

(Continued)

Primary Examiner — Abigail Fisher

(74) *Attorney, Agent, or Firm* — Fish & Richardson P.C.

(57) **ABSTRACT**

Oil-in-water fungicidal formulations are prepared having pigment dispersed therein, the pigment being stable within the oil-in-water emulsion as a result of the addition of suitable silicone surfactants and suitable emulsifiers. The formulations can be prepared either as a 2-pack formulation or as a single formulation. In the case of the single formulation polyethylene glycol is also added. In either case, the formulations show a synergistic effect through the addition of the pigment, the resulting formulations having an increased efficacy. Further, the formulations show a synergistic effect when mixed with conventional chemical fungicides, both being added in reduced amounts compared to recommended rates.

17 Claims, 17 Drawing Sheets

(56)

References Cited

U.S. PATENT DOCUMENTS

5,703,016	A	12/1997	Magin	
5,739,371	A	4/1998	O'Lenick	
5,741,502	A	4/1998	Roberts	
5,919,858	A	7/1999	Loflin	
5,958,104	A	9/1999	Nonomura et al.	
6,033,647	A	3/2000	Touzan	
6,117,820	A	9/2000	Cutler	
6,146,652	A	11/2000	Gore	
6,159,900	A	12/2000	Bieringer	
6,210,656	B1	4/2001	Touzan	
6,221,811	B1	4/2001	Policello	
6,329,321	B2	12/2001	Okura	
6,403,061	B1	6/2002	Candau	
6,416,748	B1	7/2002	Candau	
6,432,877	B2	8/2002	Okura	
6,515,031	B2	2/2003	Fefer	
6,673,360	B2	1/2004	Fefer	
6,683,030	B2	1/2004	Kober	
6,713,518	B1	3/2004	Besette	
6,734,202	B2	5/2004	Cotter	
6,803,345	B2	10/2004	Herold	
6,878,674	B2	4/2005	Kobayashi	
7,135,435	B2	11/2006	Cooper	
7,799,343	B2	9/2010	Loughner	
2001/0019728	A1	9/2001	Basinger	
2003/0087764	A1	5/2003	Pallas	
2003/0194454	A1	10/2003	Besette	
2003/0198659	A1	10/2003	Hoffmann	
2003/0198696	A1	10/2003	Keen	
2004/0132621	A1	7/2004	Frisch	
2004/0167034	A1	8/2004	Coote	
2004/0192556	A1	9/2004	Schregenberger et al.	
2005/0026786	A1	2/2005	Deckwer	
2005/0181949	A1	8/2005	Norton	
2005/0202102	A1	9/2005	Miller	
2005/0233907	A1	10/2005	Nabors	
2005/0261379	A1 *	11/2005	Fefer	514/762
2005/0274164	A1	12/2005	Coates	
2006/0063676	A1	3/2006	Brigance	
2006/0068991	A1	3/2006	Norton	
2006/0276339	A1	12/2006	Windsor	
2006/0282961	A1	12/2006	Hughes	
2006/0293188	A1 *	12/2006	Norton et al.	504/284
2007/0184005	A1	8/2007	Policello	
2008/0085832	A1	4/2008	Fefer et al.	
2008/0112909	A1	5/2008	Faler	
2008/0153702	A1	6/2008	Voeste	
2008/0161367	A1	7/2008	Voeste	
2008/0280763	A1	11/2008	Hodge	
2009/0325922	A1	12/2009	Fefer	
2010/0016447	A1	1/2010	Fefer	

FOREIGN PATENT DOCUMENTS

CA	2434848	A1	8/2002
CA	2496142	A1	8/2005
CA	2472806	A1	11/2005
CA	2507482	A1	11/2005
CA	2562718	A1	4/2008
CA	2605092	A1	4/2008
DE	2511077	A1	9/1976
EP	0498231	B2	8/1992
EP	0598515	B1	5/1994
EP	0862857	A1	9/1998
EP	1173059	B1	11/2000
EP	1563734		8/2005
GB	745360	A	2/1956
GB	747909	A	4/1956
GB	748422	A	5/1956
GB	753976	A	8/1956
GB	758926	A	10/1956
GB	762866	A	12/1956
GB	763246	A	12/1956
GB	765459	A	1/1957

GB	792045	A	3/1958
GB	1044895	A	10/1966
GB	1168913	A	10/1969
GB	1249674	A	10/1971
GB	1417364	A	12/1975
GB	2123819	A	2/1984
GB	2176493	A	12/1986
JP	57028184	A	2/1982
JP	2138376	A	5/1990
JP	3183505	A	8/1991
JP	3221576	A	9/1991
JP	4128003	A	4/1992
JP	8218225	A	8/1996
JP	11137084	A	5/1999
JP	11349588	A	12/1999
WO	9007272	A1	7/1990
WO	9312175	A1	6/1993
WO	9632010	A2	10/1996
WO	9632011	A2	10/1996
WO	9835561	A1	8/1998
WO	0221913	A2	3/2002
WO	02089573	A1	11/2002
WO	02096199	A2	12/2002
WO	03101195	A1	12/2003
WO	03105587	A1	12/2003
WO	2004030641	A1	4/2004
WO	2004080177	A2	9/2004
WO	2005009132	A1	2/2005
WO	2005055716	A2	6/2005
WO	2005082137		9/2005
WO	2007117720	A2	10/2007
WO	2007136597	A2	11/2007
WO	2008049192	A1	5/2008
WO	2008073397	A1	6/2008
WO	2009155693		12/2009

OTHER PUBLICATIONS

Cockerham et al., Evaluation of Turfgrass Growth Retardant Chemicals, California Turfgrass Culture, 1971, 23-24, 21(3).

Colby, Calculating Synergistic and Antagonistic Responses of Herbicide Combinations, Weeds, 1967, 20-22, 20.

Coo-Ranger et al., Ionic Silicone Surfactants in Water-in-Silicone Oil Emulsions Containing Proteins, Polymer Preprints, 2004, 674-675, 45(1).

Cortes-Barco et al., Induced systemic resistance against three foliar diseases of *Agrostis stolonifera* by an isoparaffin mixture, 2nd European Turfgrass Society Congress Proceedings, 2010, vol. 2.

Cortes-Barco et al., Induced systemic resistance against three foliar diseases of *Agrostis stolonifera* by (2R,3R)-butanediol or an isoparaffin mixture, Annals of Applied Biology, 2010, 179-189, 157(2).

Cortes-Barco et al., Comparison of induced resistance activated by benzothiadiazole, (2R,3R)-butanediol and an isoparaffin mixture against anthracnose of *Nicotiana benthamiana*, Plant Pathology, 2010, 643-653, 59(4).

Cranmer et al., Controlled droplet application (CDA) of fluazifop and sethoxydim for annual and perennial weed control, 1983 Meeting of the Weed Science Society of America, 1983, 23-24, Weed Abstract vol. 033 Abs. (No. 00871).

Crocker, Pesticide Screening Test for the Southern Chinch Bug, Journal of Economic Entomology, 1981, p. 730-731, 74(6).

Erhan et al., Comparisons of volatile organic chemical content of news, sheeted, and heatset ink formulations, Journal of the American Oil Chemists' Society, 2001, 419-422, 78(4).

Findanza et al., Evaluation of fungicide and plant growth regulator tank-mix programmes on dollar spot severity of creeping bentgrass, Crop Protection, 2006, 1032-1038, 25(9).

Fidanza et al., Use of a Soil Surfactant with Fungicides for Control of Fairy Ring Disease in Turfgrass, Journal of ASTM International, 2007, 77-82, 4(4).

Furata, Strangers in a Strange Land, California Turfgrass Culture, 1971, 22-23, 21(3).

Gebhardt et al., Herbicide application with the controlled droplet applicator when using soybean oil, American Society of Agricultural Engineers Paper No. 83-1509, 1983, 12.

(56)

References Cited

OTHER PUBLICATIONS

- Guy et al., The performance of postemergence grass herbicides applied with sprinkler irrigation, Proceedings of the 39th annual meeting of the Southern Weed Science Society, 1986, 106, 8A (Abstract).
- Hartzler, Role of spray adjuvants with postemergence herbicides, ISU Weed Science Online, Mar. 7, 2001, <http://www.weeds.iastate.edu/mgmt/2001/additives.htm> (downloaded Aug. 19, 2011).
- Hill, Silicone surfactants—new developments, Current Opinion in Colloid & Interface Science, 2002, 255-261, 7(5-6).
- Hoffman, Analysis of Alcohol and Alkylphenol Polyethers via Packed Column Supercritical Fluid Chromatography, Ph.D. Thesis, VirginiaTech, Chapter 1.
- Hsiang et al., Baseline sensitivity and cross-resistance to demethylation-inhibiting fungicides in Ontario isolates of *Sclerotinia homoeocarpa*, European Journal of Plant Pathology, 1997, 409-416, 103(5).
- Hsiang et al., Sensitivity of *Sclerotinia homoeocarpa* to demethylation-inhibiting fungicides in Ontario, Canada, after a decade of use, Plant Pathology, 2007, 500-507, 56(3).
- Jordan, Enhanced post-emergence herbicide efficacy with ultra-low volume application, Proceedings of the 48th annual meeting of the Southern Weed Science Society, 1995, 208-212, 48.
- McCowan, Turf Herbicide Rx: Add Oil, Agricultural Chemicals, 1968, p. 18-21, 23(4).
- Nalewaja et al., Crop origin oils with grass control herbicides, Proceedings of the North Central Weed Control Conference, 1983, 3, 034 (Abstract).
- Ostmeyer, The color Green, Golf Course Management, 1994, 40, 44, August.
- Palla et al., Correlation of Dispersion Stability With Surfactant Concentration and Abrasive Particle Size for Chemical Mechanical Polishing (CMP) Slurries, Journal of Dispersion Science and Technology, 2000, 491-509, 21(5).
- Pavlista, Paraffin enhances yield and quality of the potato cultivar Atlantic, Journal of Production Agriculture, 1995, 40-42, 8(1).
- Perry, Ground Covers: Specifications and Costs, California Turfgrass Culture, 1971, 21-22, 21(3).
- Puterka, Fungal pathogens for arthropod pest control in orchard systems: mycoinsecticidal approach for pear psylla control, BioControl, 1999, 183-210, 44(2).
- Rieke, Thatchremoval, California Turfgrass Culture, 1971, 19-20, 21(3).
- "The Stylet-Oil User's Guide", <http://www.stylet-oil.com/> (downloaded Mar. 22, 2005).
- Schott et al., Effects of adjuvants on herbicidal action. III. Effects of petroleum and rapeseed oils on diclofop-methyl action on ryegrass, Agronomie, 1991, 27-34, 11(1).
- Shearman et al., Colorant Effects on Dormant Buffalograss Turf Performance, HortTechnology, 2005, 244-246, 15(2).
- Trathnigg et al., Molecular Characterization of Ethoxylates by Complementary Chromatographic Techniques. Evaluation of Efficiency and Reliability, Tenside Surf. Det. 2003, 148-154, 40(3).
- Tu et al., Weed Control Methods Handbook: Tools and Techniques for Use in Natural Areas, The Nature Conservancy Wildland Invasive Species Team, Apr. 2001.
- Van Dam et al., A Turfgrass Colorant Study, California Turfgrass Culture, 1971, 17-19, 21(3).
- Vincelli, Chemical Control of Turfgrass Diseases 2010, University of Kentucky College of Agriculture, <http://pest.ca.uky.edu/PSEP/Manuals/ppa1.pdf> (downloaded Aug. 22, 2011).
- Walsh et al., Biology and Management of Dollar Spot (*Sclerotinia homoeocarpa*), an Important Disease of Turfgrass, HortScience, 1999, 13-21, 34(1).
- Womack et al., A vegetable oil-based invert emulsion for mycoherbicide delivery, Biological Control, 1996, 23-28, 6(1).
- Product Bulletin for Caltex, Caltex Australia, http://www.caltex.com.au/products_oil_detail_print.asp?id=229 (downloaded Aug. 2, 2006).
- Yang et al., Infection of Leafy Spurge by *Alternaria alternata* and *A. angustiovoidea* in the Absence of Dew, Phytopathology, 1993, 953-958, 83(9).
- Youngner, Kikuyugrass, Pennisetum Clandestinum, and Its Control, Southern California Turfgrass Culture, 1958, 1-4, 8(1).
- Youngner, Gibberellic Acid on Zoysia Grasses, Southern California Turfgrass Culture, 1958, 5-6, 8(1).
- Youngner et al., Colorants for Dormant Bermuda and Other Subtropical Grasses, Southern California Turfgrass Culture, 1958, 7-8, 8(1).
- Material Safety Data Sheet for Broadcoat Spray Adjuvant, Caltex Australia Limited, Sep. 2003.
- Rhizoctonia Large Patch Disease of Zoysiagrass and Bermudagrass, University of Arkansas Division of Agriculture, http://www.uaex.edu/Other_Areas/publications/PDF/fsa-7527.pdf (downloaded Aug. 30, 2011).
- PureSpray Spray Oil 10E, Delaware Department of Agriculture Pesticide Database Searches (downloaded from DDA website Apr. 7, 2005).
- Pesticide Product Label System (PPLS)—Search Results for PureSpray Oil 10E—Approval dates Apr. 21, 2000, Jul. 23, 2004, Sep. 24, 2003, Mar. 5, 2004 (downloaded from EPA Office of Pesticide Programs website Apr. 27, 2005).
- Material Safety Data Sheet for AGRI-DEX, Helena Chemical Company, Apr. 29, 2005.
- Material Safety Data Sheet for BLENDEX VHC, Helena Chemical Company, Jul. 27, 2000.
- Material Safety Data Sheet for Chipco Signature, Bayer CropScience Pty Ltd, Oct. 21, 2002.
- Material Safety Data Sheet for Chipco Signature, Bayer CropScience Pty Ltd, May 1, 2007.
- Material Safety Data Sheet for Civitas, Petro-Canada Lubricants, Inc., Mar. 21, 2011.
- Material Safety Data Sheet for FORE 80 WP RAINSHIELD, Dow AgroSciences, Jun. 1, 2001.
- Material Safety Data Sheet for FORE Fungicide, Rohm and Haas Company, Oct. 16, 1995.
- Material Safety Data Sheet for Harmonizer, Petro-Canada Lubricants Inc., May 6, 2011.
- Material Safety Data Sheet for JMS Stylet-Oil (Mar. 1, 1994).
- Material Safety Data Sheet for PEPTOIL, Drexel Chemical Company, Jan. 7, 2005.
- Material Safety Data Sheet for SURF AC 820, Drexel Chemical Company, Jul. 22, 2005.
- Specimen Label for AGRI-DEX, Helena Chemical Company, 2005.
- Specimen Label for BLENDEX VHC, Helena Chemical Company, 2006.
- 2006 Pest Control Recommendations for Lawn and Turf Areas, Rutgers Cook College, <http://njaes.rutgers.edu/pubs/publication.asp?pid=e037> (downloaded Aug. 25, 2011).
- Brochure for Civitas, Petro-Canada, <http://www.civitasturf.com/pdf/CIVITAS-technical-brochure.pdf> (downloaded Aug. 22, 2011).
- Technical Bulletin for Civitas, Petro-Canada, <http://www.civitasturf.com/pdf/techBulletin.pdf> (downloaded Aug. 22, 2011).
- A Guide to Major Turfgrass Pests & Turfgrasses, NC State University, <http://www.turfinfo.ncsu.edu/PDFFiles/004041/ag348.pdf> (downloaded Aug. 25, 2011).
- Application of Fungicides for Suppression of Fusarium Head Blight (Scab), North Dakota State University, <http://www.ag.ndsu.edu/pubs/ageng/machine/ae1148w.htm> (downloaded Aug. 22, 2011).
- An Integrated Approach to Insect Management in Turfgrass: Black Cutworm, Richard J. Buckley et al., Rutgers, The State University of New Jersey, <http://snyderfarm.rutgers.edu/pdfs/BlackCutworms.pdf> (downloaded Aug. 26, 2011).
- An Integrated Approach to Insect Management in Turfgrass: Sod Webworms, Albrecht M. Koppenhofer et al., Rutgers, The State University of New Jersey, <http://snyderfarm.rutgers.edu/pdfs/SodWebworms.pdf> (downloaded Aug. 26, 2011).
- An Integrated Approach to Insect Management in Turfgrass: White Grubs, Albrecht M. Koppenhofer et al., Rutgers, The State University of New Jersey, http://www.co.somerset.nj.us/_pdffiles/JapBeetleFS.pdf (downloaded Aug. 26, 2011).

(56)

References Cited

OTHER PUBLICATIONS

- Armyworms and cutworms in turfgrass, Erin W. Hodgson, Utah State University, <http://extension.usu.edu/files/publications/factsheet/armyw-cutw-turf07.pdf> (downloaded Aug. 26, 2011).
- Bentgrass dead spot, University of Connecticut, http://www.turf.uconn.edu/pdf/research/factsheets/Disease_Bentgrass_Dead_Spot.pdf (downloaded Aug. 22, 2011).
- Bentgrass Deadspot, Cornell University, <http://plantclinic.cornell.edu/factsheets/bentgrassdeadspot.pdf> (downloaded Aug. 22, 2011).
- Biological/Biorational Products for Disease Management, University of Connecticut Integrated Pest Management, <http://www.ipm.uconn.edu/ipm/greenhs/htms/biofung.htm> (downloaded Aug. 22, 2011).
- Biology and Control of Dollar Spot Disease, Ontario Ministry of Agriculture Food & Rural Affairs, http://www.omafra.gov.on.ca/english/crops/facts/info_turfdollarspot.htm (downloaded Aug. 22, 2011).
- Black Cutworms, D.R. Smitley et al., Michigan State University Turfgrass Science, <http://www.turf.msu.edu/black-cutworms> (downloaded Aug. 26, 2011).
- Chemical Control of Turfgrass Diseases 2011, University of Kentucky College of Agriculture, <http://pest.ca.uky.edu/PSEP/Manuals/ppa1.pdf> (downloaded Aug. 25, 2011).
- Chemical Structures, The Bugwood Network, <http://www.bugwood.org/PAT/22chemicalstructures.html> (downloaded May 25, 2006).
- Chemical Trials for Dollar Spot Disease Control Summer 2006, Guelph Turfgrass Institute 2006 Annual Research Report, <http://131.104.104.3/06anrep/40-42.pdf> (downloaded Aug. 22, 2011).
- Clover and Other Mites of Turfgrass, W.S. Cranshaw, Colorado State University, <http://www.ext.colostate.edu/pubs/insect/05505.html> (downloaded Aug. 26, 2011).
- Cultural practices and their effects upon turf grass growth and stress tolerance, Greenkeeper International, <http://www.bigga.org.uk/about-us/magazine/back-issues/07-2006/cultural-pray-tim-butler/00919.html> (downloaded Aug. 24, 2011).
- Dead Spot Disease of Creeping Bentgrass, University of Maryland, <http://www.hgic.umd.edu/content/documents/TT-14DeadSpot.pdf> (downloaded Aug. 22, 2011).
- Dead Spot of Creeping Bentgrass and Hybrid Bermudagrass, Plant Management Network, <http://www.plantmanagementnetwork.org/pub/ats/diagnostic/2005/deadspot/> (downloaded Aug. 22, 2011).
- Dollar Spot on Turfgrass, Cornell University, http://counties.cce.cornell.edu/wyoming/agriculture/resources/ipd/dollar_spot_turfgrass.htm (downloaded Aug. 22, 2011).
- EPA: Pesticides—Inert (other) Pesticide Ingredients in Pesticide Products, U.S. Environment Protection Agency, <http://www.epa.gov/oppr001/inerts/lists.html> (downloaded Sep. 11, 2007).
- Fungicide Resistance Action Committee Code List: Fungicides sorted by mode of Action, Fungicide Resistance Action Committee, <http://www.frac.info/frac/publication/ahang/FRAC%20Code%20List%202011-final.pdf> (downloaded Aug. 22, 2011).
- Fungicide Synergy, Kansas State University, http://www.ksuturf.com/LISTServ/Archive/2009_02_26_Fungicide_Synergy.pdf (downloaded Aug. 22, 2011).
- Gray leaf spot of perennial ryegrass, Kansas State University Turfgrass Research, (downloaded Aug. 23, 2011).
- Gray Leaf Spot of Perennial Ryegrass, Plant Management Network, <http://www.plantmanagementnetwork.org/pub/php/diagnosticguide/2003/ryegrass/> (downloaded Aug. 23, 2011).
- Herbicide Recommendations for Turfgrass: Postemergence Broadleaf Herbicides, Ontario Ministry of Agriculture, Food and Rural Affairs, <http://www.omafra.gov.on.ca/english/crops/pub75/17turpbh.htm> (downloaded Sep. 10, 2001).
- Herbicide—Wikipedia, the free encyclopedia, <http://en.wikipedia.org/wiki/Herbicide> (downloaded Aug. 29, 2006).
- “Horticultural Oils”, <http://www.ipmofalaska.com/files/hortoils.html>, IPM of Alaska (downloaded Apr. 5, 2005).
- Insect Pest Management on Golf Courses, Eileen A. Buss, University of Florida, <http://edis.ifas.ufl.edu/in410> (downloaded Aug. 26, 2011).
- Insect Pest Management on Turfgrass, Eileen A. Buss, University of Florida, <http://edis.ifas.ufl.edu/ig001> (downloaded Aug. 26, 2011).
- Integrated Pest Management—Identification & Management of Turfgrass Disease, University of Missouri, <http://extension.missouri.edu/p/IPM1029> (downloaded Aug. 25, 2011).
- “It pays to be pure”, http://www.findarticles.com/p/articles/mi_qa3824/is_200405/ai_n9424665/print, Meister Media Worldwide, May 2004.
- Online Guide to Plant Disease Control of Oregon State University Extension (<http://plant-disease.ipcc.orst.edu/>) (downloaded May 16, 2005) and hardcopy version, “The 2004 PNW Plant Disease Management Handbook”.
- Performance of generic phosphite fungicides: A status report, AgNet Mar. 8, 2004, The Canadian Phytopathological Society, <http://www.cps-scp.ca/pathologynews/performanceofgenericfungicides.html> (downloaded Aug. 22, 2011).
- Propiconazole Pesticide Information Profile, Extension Toxicology Network, <http://pmep.cce.cornell.edu/profiles/extoxnet/metiram-propoxur/propiconazole-ext.html> (downloaded Aug. 19, 2011).
- Turf Tip, University of Arkansas, <http://turf.uark.edu/turfhelp/archives/030509.html> (downloaded Aug. 22, 2011).
- Turfgrass Disease Profiles Dollar Spot, Purdue University, <http://www.ces.purdue.edu/extmedia/BP/BP-105-W.pdf> (downloaded Aug. 22, 2011).
- Turfgrass Disease Profiles Gray Leaf Spot, Purdue University, <http://www.ces.purdue.edu/extmedia/BP/BP-107-W.pdf> (downloaded Aug. 23, 2011).
- Turfgrass Insects Sheet 1, D.E. Short et al., University of Florida, <http://edis.ifas.ufl.edu/in025> (downloaded Aug. 26, 2011).
- Turfgrass Insects Sheet 2, D.E. Short et al., University of Florida, <http://edis.ifas.ufl.edu/in026> (downloaded Aug. 26, 2011).
- Turfgrass Pest Control, West Virginia University, <http://www.wvu.edu/~exten/infores/pubs/pest/pcerti19.pdf> (downloaded Aug. 22, 2011).
- Understanding Bentgrass Dead Spot, USGA Turfgrass and Environmental Research Online, <http://turf.lib.msu.edu/tero/v02/n02.pdf> (downloaded Aug. 22, 2011).
- Heil, Induced Systemic Resistance (ISR) Against Pathogens in the Context of Induced Plant Defences, *Annals of Botany*, 2002, 503-512, 89(5).
- Lorbeer, Synergism, Antagonism, and Additive Action of Fungicides in Mixtures, *Phytopathology*, 1996, 1261-1262, 86(11).
- Samoucha et al., Synergism in fungicide mixtures against *Pseudoperonospora cubensis*, *Phytoparasitica*, 1988, 337-342, 16(4).
- Vallad et al., Systemic Acquired Resistance and Induced Systemic Resistance in Conventional Agriculture, *Crop Science*, 2004, 1920-1934, 44.
- Silicone Surface-Active Agents, Donna Perry, Dow Corning Corporation, <http://www.dowcorning.com/content/publishedlit/26-1365.pdf> (downloaded Aug. 30, 2011).
- Turfgrass Diseases: Leaf Spots and Tip Blights, Melting Out, Crown and Root Rots, University of Rhode Island Landscape Horticulture Program, <http://www.uri.edu/ce/factsheets/sheets/leafspotsetc.html> (downloaded Aug. 30, 2011).
- Leaf Spot and Melting-out (crown and root rot) Diseases—Center for Turfgrass Science, Penn State College of Agricultural Sciences, <http://cropsoil.psu.edu/turf/extension/factsheets/managing-diseases/leaf-spot> (downloaded Aug. 30, 2011).
- Turfgrass Disease Profiles Leaf Spot/Melting Out, Purdue University, <http://www.ces.purdue.edu/extmedia/bp/bp-103-w.pdf> (downloaded Aug. 30, 2011).
- Brown Patch, Center for Turfgrass Science, Penn State College of Agricultural Sciences, <http://cropsoil.psu.edu/turf/extension/factsheets/managing-diseases/brown-patch> (downloaded Aug. 30, 2011).
- Brown Patch, University of Guelph, <http://www.uoguelph.ca/pdc/Factsheets/PDFs/127TurfBrownPatch.pdf> (downloaded Aug. 30, 2011).

(56)

References Cited

OTHER PUBLICATIONS

- Turfgrass Disease Profiles Brown Patch, Purdue University, <http://www.ces.purdue.edu/extmedia/BP/BP-106-W.pdf> (downloaded Aug. 30, 2011).
- Brown Patch on Turfgrass, Cornell University Department of Plant Pathology and Plant-Microbe Biology, <http://plantclinic.cornell.edu/factsheets/brownpatch.pdf> (downloaded Aug. 30, 2011).
- Turfgrass Disease Profiles Rhizoctonia Large Patch, Purdue University, <http://www.extension.purdue.edu/extmedia/BP/BP-117-W.pdf> (downloaded Aug. 30, 2011).
- Material Safety Data Sheet for Banner MAXX, Syngenta Crop Protection, Inc., Aug. 30, 2010.
- Material Safety Data Sheet for Cleary 3336 Plus, Cleary Chemical Corporation, Feb. 1, 2005.
- Material Safety Data Sheet for Daconil 2787, Syngenta Crop Protection Canada, Inc., Dec. 31, 2008.
- Material Safety Data Sheet for Daconil Ultrex, Syngenta Crop Protection Canada, Inc., Aug. 1, 2009.
- Material Safety Data Sheet for Lambent MFF-199 SW, Lambent Technologies Corp., Jan. 31, 2005.
- Material Safety Data Sheet for Lambent MFF 159-100, Lambent Technologies Corp., Apr. 4, 2006.
- Material Safety Data Sheet for Rovral Green GT, Bayer CropScience Inc., Mar. 2, 2011.
- Material Safety Data Sheet for Silsurf A008-UP, Siltech Corporation, Aug. 21, 2009.
- Material Safety Data Sheet for Sylgard 309 Silicone Surfactant, Dow Corning Corporation, Apr. 5, 2001.
- Turfgrass Disease Profiles Gray Snow Mold, Purdue University, <http://www.ces.purdue.edu/extmedia/BP/BP-101-W.pdf> (downloaded Sep. 15, 2011).
- Turfgrass Disease Profiles Pink Snow Mold and Microdochium Patch, Purdue University, <http://www.ces.purdue.edu/extmedia/BP/BP-102-W.pdf> (downloaded Sep. 15, 2011).
- Office Action (Restriction Requirement) for U.S. Appl. No. 11/866,157 dated May 16, 2011.
- International Search Report for PCT International Application No. PCT/CA2007/001762, Jan. 5, 2011.
- Written Opinion for PCT International Application No. PCT/CA2007/001762, Jan. 5, 2011.
- International Preliminary Report on Patentability for PCT International Application No. PCT/CA2007/001762, Jan. 5, 2011.
- Office Action (Restriction Requirement) for U.S. Appl. No. 10/908,538 dated Feb. 26, 2009.
- Office Action for U.S. Appl. No. 10/908,538 dated Apr. 1, 2009.
- International Search Report for PCT International Application No. PCT/CA2009/000862, Sep. 20, 2009.
- Written Opinion for PCT International Application No. PCT/CA2009/000862, Sep. 20, 2009.
- International Preliminary Report on Patentability for PCT International Application No. PCT/CA2009/000862, Sep. 20, 2009.
- Examination Report for NZ Application No. NZ590318 dated May 6, 2011.
- Office Action for CA Application No. CA2,507,482 dated Jan. 18, 2011.
- Office Action for CA Application No. CA2,507,482 dated Aug. 11, 2009.
- Office Action for U.S. Appl. No. 11/866,157 dated Aug. 29, 2011.
- Cropper, Towards Reducing Fungicide Use in the Control of Dollar Spot (*Sclerotinia homoeocarpa* F.T. Bennett) Disease on Creeping Bentgrass (*Agrostis stolonifera* L.), Master of Science Thesis, University of Kentucky, <http://archive.uky.edu/handle/10225/1044> (downloaded Sep. 15, 2011).
- Kremer et al., Control of *Sclerotinia homoeocarpa* in Turfgrass Using Effective Microorganisms, EM World J., 2000, 16-21, 1(1).
- Pest Control for Professional Turfgrass Managers 2011, North Carolina State University, http://www.turffiles.ncsu.edu/PDFFiles/004176/AG408PestControl_Professionals.pdf (downloaded Sep. 15, 2011).
- Gilbert et al., Spring Greenup of Dormant Non-Overseeded Bermudagrass, University of Arizona College of Agriculture 2004 Turfgrass and Ornamental Research Report, <http://ag.arizona.edu/pubs/crops/az1359/az13593c11.pdf> (downloaded Sep. 16, 2011).
- Liu et al., Painting dormant bermudagrass putting greens, GCM, 86-91, Nov. 2007.
- Shaposhnikov et al., Carboxy-substituted Phthalocyanine Metal Complexes, Russian Journal of General Chemistry, 2005, 1480-1488, 75(9).
- Vol'Pin et al., Redox and fungicidal properties of phthalocyanine metal complexes as related to active oxygen, Journal of Inorganic Biochemistry, 2000, 285-292, 81(4).
- Material Safety Data Sheet for Grass Greenzit, W.A. Cleary Chemical Corporation, Oct. 1997.
- Material Safety Data Sheet for Green Lawnger, Becker Underwood, Inc., Feb. 25, 2009.
- Material Safety Data Sheet for Regreen, Precision Laboratories, Inc., Mar. 1, 2010.
- Pigment Identification Charts, Golden Artist Colors, <http://www.goldenpaints.com/technicaldata/pigment.php#> (downloaded Sep. 15, 2011).
- R.M. Goodwin et al., New Zealand Plant Protection 53:230-234 (2000).
- Burr RJ and Warren GF, Weed Science 19(6): 701-705 (1971).
- Grover et al., Weed Science 20(4): 320-324 (1972).
- Burt, Plantation Field Laboratory Mimeo Report PFL66-1, (Aug. 1966).
- Horn, Florida State Horticultural Society, pp. 494-499 (1966).
- Horn, Florida State Horticultural Society, pp. 499-509 (1966).
- Office Action dtd. Nov. 9, 2011 CA App No. 2,507,482.
- Christians, Creative Uses for Plant Growth Regulators, USGA Green Section Record, Sep. 11-13, 2001/ Oct. 2001.
- Danneberger et al., Turfgrass Growth Substances, Golf Course Management, 1990, 80, 82, 86, 88, 58(4).
- Grey et al., Timed Release of Flurprimidol from a Granular Formulation in Mulches and Sand, HortScience, 2009, 512-515, 44(2).
- Huang, Plant growth regulators: What and why, GCM golf course management, 2007, 157-160, Jan. 2007.
- Lickfeldt et al., Implications of Repeated Trinexapac-Ethyl Applications on Kentucky Bluegrass, Agronomy Journal, 2001, 1164-1168, 93(5).
- Mercier, Use of the growth regulator paclobutrazol in the management of dollar spot of creeping bentgrass in Minnesota, Phytoprotection, 1999, 65-70, 80(2).
- Nelson et al., 2,4-D and Mycoleptodiscus terrestris for Control of Eurasian Watermilfoil, J. Aquat. Plant Manage., 2005, 29-34, 43.
- Material Safety Data Sheet for Kiltex Lawn Weed Control Concentrate (Ortho), Scotts Canada Ltd., Sep. 13, 2005.
- Label for Killex, Scotts, Canada Ltd., Jul. 23, 2001.
- Specimen Label for Trimec Classic, PBI/Gordon Corporation, 1973.
- Specimen Label for Trimec Southern, PBI/Gordon Corporation, 1987.
- 2,4-Dichlorophenoxyacetic acid—Wikipedia, the free encyclopedia, <http://en.wikipedia.org/wiki/2%2C4-D> (downloaded Aug. 29, 2006).
- Scotts Canada Home: Killex Concentrate, <http://scottscanada.calindex.cfm/levent/ProductGuide/product/documentId/30B255B82B...> (downloaded Aug. 2, 2006).
- Notice for Mecoprop-P TGAC, Commonwealth of Australia Gazette No. NRA 3, Mar. 6, 2001.
- Quicksheet for SALVO herbicide, UAP Canada, 2006.
- International Search Report for PCT International Application No. PCT/CA2009/000862, 2009.
- Datapak for SALVO herbicide, United Agri Products Canada Inc., 2005.
- Product Information Bulletin for SALVO, Platte Chemical Co., 2001.
- Characteristics of Plant Growth Regulators used in Fine Turf, Clemson University, <http://www.clemson.edu/extension/horticulture/turf/pest-guidelines/2011-pest-guidelines/plant>, 2011.
- Chemical Update: Plant growth regulators, Grounds Maintenance, <http://grounds-mag.com/mag/grounds-maintenance-chemical-update-plant-6/> (downloaded Aug. 24, 2011).

(56)

References Cited

OTHER PUBLICATIONS

Ethephon and Trinexapac-ethyl Influence Creeping Bentgrass Growth, Quality, and Putting Green Performance, Plant Management Network, 2006.

Evaluation of Commercially Available Plant Growth Regulator Programs for Creeping Bentgrass Fairway Management, 2003, 2003 Annual Report—Purdue University Turfgrass.

Phytotoxicity, Food, Crop & Livestock Safety, British Columbia Ministry of Agriculture, <http://www.agf.gov.bc.ca/pesticides/e-10.htm> (downloaded Aug. 26, 2011).

Phytotoxicity on Foliage Ornamentals Caused by Bactericides and Fungicides, A.R. Chase et al., Plant Pathology Fact Sheet, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, <http://plantpath.ifas.ufl.edu/taxextpublFactSheetsippOO30.pdf> (downloaded Aug. 26, 2011).

Plant Growth Regulator Effects on Annual Bluegrass/Creeping Bentgrass Competition, Department of Crop & Soil Sciences Michigan State University, 1988.

Plant Growth Regulator Regimens Reduce *Poa annua* Populations in Creeping Bentgrass, Plant Management Network (downloaded Aug. 24, 2011).

Plant Growth Regulators, University of Florida (downloaded Aug. 24, 2011).

Plant Growth Regulators as a Turfgrass Management Tool, Greenkeeper (downloaded Aug. 24, 2011).

Plant Growth Regulators for Fine Turf, Clemson University (downloaded Aug. 24, 2011).

Plant Growth Regulators for Tuff, Landscape and Garden, Lawn Care Academy (downloaded Aug. 24, 2011).

Plant Growth Regulators: More Color, Less Clippings, Irrigation & green industry (downloaded Aug. 24, 2011).

Plant Growth Regulators Used in Turfgrass Management, Georgia Turf (downloaded Aug. 25, 2011).

Plant Growth Regulators Used in Turfgrass Management, Greenkeeper (downloaded Aug. 24, 2011).

Plant Growth Retardants for Fine Turf and Roadsides/Utilities, University of Florida (downloaded Aug. 24, 2011).

Putting the Nos. To PGRs, Grounds Maintenance (downloaded Aug. 25, 2011).

Repeat Applications of Paclobutrazole (TGR) Plant Growth Regulator on Overseeded Bermudagrass Tuff: Weed Control and Bermudagrass Transition, The 2009 Turfgrass, Landscape and Urban IPM Research Summary, The University of Arizona (downloaded Aug. 24, 2011).

The Effect of the Plant Growth Regulator Primo on Winter Hardiness Levels, Prairie Turfgrass Research Centre (downloaded Aug. 25, 2011).

Trinexapac-ethyl—PubChem Public Chemical Database (downloaded Aug. 25, 2011).

Turfgrass Growth Regulators for Professional Managers, Patrick E. McCullough, Extension Agronomist-Weed Science, Georgia Turf (downloaded Aug. 25, 2011).

Turfgrass Growth Regulators for Professional Managers, Tim R. Murphy, Extension Agronomist-Weed Science, Georgia Turf (downloaded Aug. 25, 2011).

Turfgrass quality and phytotoxicity affected by growth retardants, R.W. Duell (downloaded Aug. 24, 2011).

Use of Plant Growth Regulators to Retard Growth of Bermudagrass and Dallisgrass in the Landscape, Texas A&M University (downloaded Aug. 24, 2011).

Using plant growth regulators in turfgrass management. (Green Science), Golfscape (downloaded Aug. 24, 2011).

vol. 3.3—Plant Growth Regulators Mode of Action, Australian Golf Course Superintendents' Association (downloaded Aug. 24, 2011).

Beasley et al., Trinexapac-ethyl and Paclobutrazol Affect Kentucky Bluegrass Single-Leaf Carbon Exchange Rates and Plant Growth, Crop Science, 2007, 132-138, 47.

Deformulation of RD 7212 Grazz Greenzit, 2009.

Effect of Chloro-Aluminum-Phtahalocyanine on the Growth of *Lenma gibba* G3-Ben-Tal, 1989.

Effects of Turf Colorants and FES04 on Spring Greenup of Zoysiagrass-Diesburg, 1990.

Grass Greenzit specimen label, 1998.

Kannar Product Range Turf Enhancing Products, 2007.

Material Safety Data Sheet—Kannar Green KC 200, 2007.

Material Safety Data Sheet—Kannar Tufkare Green, 2007.

Quantification of Phosphorous in Water Based Green Pigments, 2009.

The Development of Solid Spectral Filters for the Regulation of Plant Growth, 2008.

Turf Grass Coloration Using Hexadentate Cobalt Phthalocyanine amine, 1976.

Use of Cleary's Grass Greenzit—Cleary Chem Corp, 2004.

A Guide to NTEP Turfgrass Ratings-Morris, 2011.

Better Creeping Bentgrass Through Electricity-Huang, 2003.

Cytokinin Effects on Creeping Bentgrass Responses to Heat Stress—Liu, 2002.

Grass Greenzit Material Safety Data Sheet—Cleary, 2011.

Heat Stress Study Using Greenzit Pigment—U. of Guelph, 2009.

Overseed Greens Performance Trials—Kopeck, 2004.

* cited by examiner


Fig. 1


Fig. 2


Fig. 3




Fig. 4

Fig. 5



**Fig. 6**




Fig. 7




Fig. 8

Fig. 9A




Fig. 9B







Fig. 10

Fig. 11

Typhula ishikariensis plots









Fig. 14



Fig. 15

Fig. 16

1

TURFGRASS FUNGICIDE FORMULATION WITH PIGMENT

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Patent application Ser. No. 61/075,821, filed Jun. 26, 2008 and U.S. Provisional Patent application Ser. No. 61/147,523, filed Jan. 27, 2009, the entirety of which are incorporated herein by reference.

FIELD OF THE INVENTION

Embodiments of the invention relate to oil-in water emulsions having fungicidal properties when applied to turfgrass and more particularly, to oil-in-water emulsions having stable dispersions of pigment therein for enhancing fungicidal activity and for imparting color to the turfgrass when applied thereon.

BACKGROUND OF THE INVENTION

In Applicant's previous applications, such as in co-pending US-2005-0261379-A1, the entirety of which is incorporated herein by reference, the Applicant has previously described the use of an oil-in-water emulsion as means of controlling turf diseases. A disadvantage may be that certain grasses such as varieties of bentgrass are sensitive to oil-in-water formulations which can often have the undesired effect of discolouring the grass especially under the heat of summer. Although the health of the plant is not negatively affected, the discolouration can be problematic from an aesthetic perspective.

Pigment colorants, such as phthalocyanine compounds, are known to have been used in the turf industry, both in the presence and the absence of fungicide. The known literature teaches the use of pigment color, such as non-chlorinated copper phthalocyanine as a means of colouring the grass or turf. One such example is US 2006/0293188 A1 to Norton (Bayer CropScience LP), the entirety of which is incorporated herein by reference. Additional references include U.S. Pat. No. 5,336,661 to Lucas, U.S. Pat. No. 5,643,852 to Lucas and U.S. Pat. No. 5,599,804 to Mudge, the entirety of which are incorporated herein by reference.

Others references of interest include German application DE 2511077 published in 1978, which teaches cobalt phthalocyanine as a colourant in the absence of an acid. In an English abstract for JP 03-221576 to Nippon Chemical Works, published Sep. 30, 1991, the disclosed formulation uses phthalocyanine "green", an anionic dispersant, and acrylic acid ester-styrene and water.

Unfortunately, none of the teachings provide a solution for use of pigment in oil-in-water emulsion systems as Applicant has found that pigments, such as polychlorinated copper phthalocyanine, appear to clump or coagulate and thereafter quickly fall out of suspension rendering the formulation unworkable.

Clearly, there is a need to provide an oil-in-water fungicidal formulation which contains a dispersion of pigment or colorant therein in which the colorant remains stably dispersed for application to turfgrass.

SUMMARY OF THE INVENTION

Applicant has discovered that one can generate stable compositions of pigment in a high oil-in-water emulsion environ-

2

ment by the addition of a small amount of a silicone surfactant of a specific chemistry in combination with a small amount of an emulsifier of specific chemistry. The silicone surfactant and emulsifier are thought to act as dispersants for the pigment within the oil-based formulation, prior to dilution for forming the oil-in-water emulsion and in the final oil-in-water emulsion. Formulations according to embodiments of the invention have an enhanced efficacy in treating turfgrass disease.


In addition, Applicant has found that the incorporation of a pigment dispersed into the oil-in-water emulsion disclosed herein has a synergistic impact and improves the overall efficacy of disease control of the active fungicidal components. At the same time, the amount of oil required to achieve adequate disease control can be reduced to about half the amount of oil compared to the amount required if the oil is used alone, thus further reducing the possibility of phytotoxicity. Further, discolouration issues which may occur on certain grasses, such as varieties of bentgrass that are sensitive to oil-in-water formulations especially under the heat of summer, are overcome by formulations prepared according to embodiments of the invention.

In addition, Applicant has determined that embodiments of the present invention have an unexpected synergistic effect when mixed with conventional chemical fungicides such as demethylation inhibitors (such as propiconazole), methyl benzimidazole carbamate (such as thiophanate-methyl) and dicarboximide (such as iprodione). When mixed with formulations according to embodiments of the invention, the dosage of such conventional chemical fungicides can be reduced significantly, such as to about 50% the recommended label rates, as well permitting significant reduction in the required dosage of the formulations of the present invention.

Advantageously, the present invention also acts to suppress certain turf insects such as fall armyworms and sod webworms.

In a broad aspect of the invention, a fungicidal formulation for application to turfgrass comprises: an oil-in water emulsion comprising a paraffinic oil, a suitable emulsifier and water; polychlorinated (Cu II) phthalocyanine; and a silicone surfactant selected from the group consisting of trisiloxanes and silicone polyethers of Formula I

(Formula I)



where: R=H, CH₃ or COCH₃; x=1 to 24; n=0 or ≥1; m=1, wherein the polychlorinated (Cu II) phthalocyanine is maintained in dispersion in the oil-in-water emulsion for delivery to turfgrass.

In another broad aspect, a method for preparing the fungicidal formulation comprises: preparing a first composition having the paraffinic oil; and the suitable emulsifier selected from the group consisting of natural and synthetic alcohol ethoxylates, including polyoxyethylene (4 to 12) lauryl ether (C₁₂), polyoxyethylene (10) cetyl ether (C₁₆), polyoxyethylene (10) stearyl ether (C₁₈), polyoxyethylene (10) oleyl ether (C₁₈ monounsaturated), polyoxyethylene (2 to 11) C₁₂-C₁₅ alcohols, polyoxyethylene (3 to 9) C₁₁-C₁₄ alcohols, polyoxyethylene (9) C₁₂-C₁₄ alcohols; polyoxyethylene (11) C₁₆-

C₁₈ alcohols, and polyoxyethylene (20) C₁₂-C₁₅ alcohols; alcohol alkoxylates including butyl polyoxyethylene/polyoxypropylene block copolymer; alkyl polysaccharides including C₈-C₁₁ alkylpolysaccharides; glycerol oleate; polyoxyethylene-polyoxypropylene block copolymers of MW 1100 to 11400 and 10 to 80% EO; nonyl phenol ethoxylates including polyoxyethylene (2 to 8) nonylphenol; polymeric surfactants including graft copolymer such as polymethacrylic acid and acrylate with polyoxyethylene chains and random copolymer with ester and ether group; polyethylene glycols including MW: 200 to 8000, MW: 400 PEG dioleate, and MW: 600 PEG dioleate; sorbitan fatty acid ester ethoxylates including polyoxyethylene (20) sorbitan tristearate, polyoxyethylene (20) sorbitan monooleate, polyoxyethylene (5) sorbitan monooleate, and polyoxyethylene (20) sorbitan trioleate; and mixtures thereof; preparing a second composition being an aqueous dispersion of the polychlorinated (Cu II) phthalocyanine; adding the silicone surfactant to one or both of the first and second compositions; and thereafter prior to use; and mixing an effective amount of the first composition with an effective amount of the second composition, the polychlorinated (Cu II) phthalocyanine being dispersed and stable therein.

In another broad aspect, a method for preparing the fungicidal formulation comprises: preparing a single composition having the paraffinic oil; the suitable emulsifier selected from the group consisting of natural or synthetic alcohol ethoxylates including polyoxyethylene (4 to 7) lauryl ether (C₁₂); polyoxyethylene (10) cetyl ether (C₁₆); polyoxyethylene (2 to 11) C₁₂-C₁₅ alcohols; polyoxyethylene (3 to 9) C₁₁-C₁₄ alcohols; polyoxyethylene (9) C₁₂-C₁₄ alcohols; polymeric surfactants including graft copolymer such as polymethacrylic acid and acrylate with polyoxyethylene chains; and random copolymer with ester and ether group; sorbitan fatty acid esters including sorbitan tristearate; and sorbitan trioleate; and mixtures thereof; the polychlorinated (Cu II) phthalocyanine being dispersed in oil; and the silicone surfactant, wherein the silicone surfactant further comprises polyethylene glycols (PEG) according to Formula (IV):



where: R₁=H or CH₂=CH-CH₂ or COCH₃.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a graphical representation of the results of Example 7 showing a synergistic effect resulting from the addition of pigment to an oil-in-water fungicidal formulation prepared as a 2-pack formulation, permitting use of the formulation at rates lower than predicted rates when treating turfgrass infected with dollar spot disease;

FIG. 2 is a graphical representation of the results of Example 8 showing the efficacy of the 2-pack fungicide applied at 5 gal/acre (non-aqueous portion) compared to conventional chemical fungicide, DACONIL® 2787, applied at the same rate for treating turfgrass infected with Dollar Spot disease;

FIG. 3 is a graphical representation of the results of Example 9 showing the efficacy of the 2-pack fungicide, applied at both 5 gal/acre (non-aqueous portion) and 10 gal/acre (non-aqueous portion), compared to the chemical fungicide, DACONIL® 2787, for treating turfgrass infected with Dollar Spot disease;

FIG. 4 is a graphical representation of the results of Example 10 showing the efficacy of weekly use of 2.5 gal/acre (non-aqueous portion) of 2-pack fungicide compared to an oil-in-water formulation without the addition of the poly-

chlorinated Cu (II) phthalocyanine pigment, emulsifier and silicone additives applied weekly at 5 gal/acre (non-aqueous portion);

FIG. 5 is a graphical representation of the results of Example 18 comparing the efficacy of the 2-pack formulation, a 1-pack formulation and conventional chemical fungicide DACONIL® Ultrex for treating turfgrass infected with Dollar Spot disease;

FIG. 6 is a picture of increased root length as a result of the application of a 2-Pack formulation prepared according to Example 6 when applied to a golf course fairway;

FIG. 7 is a picture of increased root length as a result of the application of a 2-Pack formulation, prepared according to Example 6, to areas of bentgrass;

FIG. 8 is a picture of increased root length as a result of the application of a 2-Pack formulation, prepared according to Example 6, to areas of bentgrass;

FIG. 9A is a graphical representation of the results of Example 21 comparing the efficacy of a 2-pack formulation and a 2-pack formulation tank mixed at 50:50 with chemical fungicide BANNER MAXX™ for treating turf infected with Dollar Spot disease;

FIG. 9B is a graphical representation of the results of Example 21 showing a synergistic effect of a 2-pack formulation tank mixed at 50:50 with chemical fungicide BANNER MAXX™ compared to the 2-pack formulation alone for treating turf infected with Dollar Spot disease, the residue period extending from 21 days to 28 days;

FIG. 10 is a graphical representation of the results of Example 22 showing a synergistic effect of a 1-pack formulation tank mixed 50:50 with chemical fungicide BANNER MAXX™ compared to the 1-pack formulation alone for treating turf infected with dollar spot disease;

FIG. 11 is a graphical representation of the results of Example 23 illustrating the effectiveness of a 2-pack formulation with 50% of the recommended label rate of BANNER MAXX™ compared to the 2-pack formulation alone, 50% of the recommended label rate of BANNER MAXX™ alone and an untreated control for treating turf infected with *Typhula ishkariensis*;

FIG. 12 is a graphical representation of the results of Example 24 illustrating the effectiveness of a 2-pack formulation with 50% of the recommended label rate of BANNER MAXX™ compared to the 2-pack formulation alone, 50% of the recommended label rate of BANNER MAXX™ alone and an untreated control for treating turf infected with *Typhula incarnata*;

FIG. 13 is a graphical representation of the results of Example 25 illustrating the effectiveness of a 2-pack formulation with 50% of the recommended label rate of BANNER MAXX™ compared to the 2-pack formulation alone, 50% of the recommended label rate of BANNER MAXX™ alone and an untreated control for treating turf infected with *Microdochium nivale*;

FIG. 14 is a graphical representation of the results of Example 26 comparing the efficacy of a 2-pack formulation enhanced with 50% of the recommended label rate of ROVRAL® Green GT to ROVRAL® Green GT at full label rate alone and ROVRAL® Green GT applied alone at 50% of the recommended label rate for the control of Fusarium Patch disease on bentgrass;

FIG. 15 is a graphical representation of the results of Example 27 comparing the efficacy of a 2-Pack formulation prepared according to Example 6 and applied at the full rate, conventional chemical fungicide Cleary 3336™ Plus at the full recommended label rate, the 2-pack formulation mixed 50:50 with Cleary 3336™ Plus, the 2-pack formulation alone

5

at 50% of the recommended rate, and Cleary 3336™ Plus alone at 50% the recommended label rate, when applied to perennial ryegrass for control of Gray Leaf Spot disease; and

FIG. 16 is a graphical representation of the results of Example 28 illustrating a synergistic effect of the addition of polychlorinated Cu (II) phthalocyanine when compared to oil/emulsifier alone and pigment alone, when compared to an untreated control.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

Prior Art

Applicant has found that the addition of colorants or pigments, based upon the teachings in the prior art, to oil-in-water emulsion formulations resulted in unstable dispersions of the pigment therein.

In the following Examples 1 through 3, polychlorinated Cu (II) phthalocyanine pigment (available from Sun Chemical as SUNSPERSE® Green 7, ~60% polychlorinated Cu (II) phthalocyanine dispersed in water or as Pigment Green 7 powder available from Hercules Exports, Mumbai, India) was added to a synthetic isoparaffinic oil (N65DW available from Petro-Canada, Calgary, AB, Canada), with or without polyoxyethylene lauryl ether, C₁₀ to C₁₆ alcohol ethoxylates, and glycerol oleate as an emulsifier (available as PC Emuls Green, from Petro-Canada, Calgary, AB, Canada) and was diluted in water to form the oil-in-water emulsion suitable for application to turfgrass.

In each of Examples 1-3, the pigment coagulated or formed clumps which rapidly separated out and/or resulted in phase separation of the formulations. Thus, the pigment did not remain dispersed in the formulations and therefore the formulations were not usable to reliably and evenly impart color to turfgrass.

Example 1

- 0.3 to 0.5 wt % polychlorinated Cu (II) phthalocyanine (SUNSPERSE® Green 7)
- 10 wt % synthetic isoparaffinic oil (N65DW)
- 89.5 wt % water

Example 2

- 0.3-0.5 wt % polychlorinated CU (II) phthalocyanine (SUNSPERSE® Green 7)
- 10 wt % isoparaffinic oil (N65DW)
- 0.1 wt % emulsifier (PC Emuls Green)
- 89.4 wt % water

Example 3

- 0.5 wt % polychlorinated Cu (II) phthalocyanine (Pigment Green 7 powder)
- 10 wt % isoparaffinic oil (N65DW)
- 0.1 wt % emulsifier (PC Emuls Green)
- 89.4 wt % water

Embodiments of the Invention

As shown in Example 4, Applicant has found that the incorporation of specific silicone surfactants and emulsifier dispersants added to an oil-in-water emulsion, having a significant portion of oil therein and containing pigment, results

6

in the pigment being stably dispersed therein for application, such as by spraying, to turfgrass.

The non-aqueous portion of the oil-in-water portion is typically applied at rates from about 1 gal/acre (0.093 L/100 m²) to about 15 gal/acre (1.395 L/100 m²). The total spray volume of the oil-in-water emulsion is typically from about 20 gal/acre (1.9 L/100 m²) to about 200 gal/acre (18.6 L/100 m²).

Example 4


A formulation according to an embodiment of the invention was prepared as follows:

- 0.5 wt % polychlorinated Cu (II) phthalocyanine (SUNSPERSE® Green 7)
- 10 wt % isoparaffinic oil (N65DW)
- 0.1 wt % emulsifier (PC Emuls Green)
- 89.3 wt % water
- 0.1 wt % silicone surfactant (Lambent MFF-199 SW)

An exemplary silicone super-wetting agent, such as Lambent MFF-199 SW (available from Lambent Technologies, a division of Petroferm, Inc., Gurnee, Ill., USA. MFF-199-SW) is a silicone copolyol, containing a hydrogen end group and one pendant polyethylene oxide group and has an average molecular weight between 600 to 1000 Daltons.

Lambent MFF-199 is a totally different class of silicone oil compared to common linear or cyclic polydimethyl-siloxane. Lambent MFF-199 is a trisiloxane with an ethoxylated alkyl group having a hydrogen end group (H-End). The number of ethoxylation group is in the range of 1-20.

The suggested structure is as follows:



where: n=1-20 and average n=8

2-Pack Formulation

In an embodiment of the invention, a first composition (Pack A) is prepared containing the paraffinic oil and a suitable emulsifier. A second composition (Pack B) is prepared comprising a polychlorinated Cu (II) phthalocyanine. The polychlorinated Cu (II) phthalocyanine can be provided as a powder dispersed in water or as a ready to use aqueous dispersion.


A silicone surfactant, such as Lambent MFF 159-100 or MFF 199 SW or other suitable silicone surfactant as described below can be added entirely to Pack A or to Pack B. Alternatively, the silicone surfactant can be split between Pack A and Pack B.

Immediately prior to use, an effective amount of each of Pack A and Pack B are mixed together to form the fungicidal dispersion which is further diluted in water to the desired concentration, as described below, for delivery to turfgrass at a predetermined dosage rate.

More particularly, in embodiments of the invention, the effective amount of Pack A is mixed with some or all of the water which would be required to obtain a desired concentration so as to form an emulsion. Thereafter, the effective amount of Pack B and any remaining water are added to the emulsion and the mixture is delivered to turfgrass as an oil-in-water emulsion at a predetermined dosage rate.

The silicone and emulsifier are selected to provide an inter-molecular hydrophilic and lipophilic balance upon mixing of Pack A and Pack B so as to substantially prevent the polychlorinated Cu (II) phthalocyanine from clumping and rapidly separating out of suspension in the presence of the oil phase during application to the turfgrass.

Suitable silicone surfactants comprise trisiloxanes or silicone polyethers having a suitable alkoxy group with hydrogen end groups (H-capped), methyl end groups (CH₃ capped) or acetyl end groups (COCH₃ capped) according to the following formula (I):



Where:

R=H; x=1 to 24; n=0; m=1; H-capped trisiloxane

R=H; x=1 to 24; n≥1; m=1; H-capped silicone polyethers

R=CH₃; x=1 to 24; n=0; m=1; CH₃-capped trisiloxane

R=CH₃; x=1 to 24; n≥1; m=1; CH₃-capped silicone polyethers


R=COCH₃; x=1 to 24; n=0; m=1; COCH₃-capped trisiloxane

R=COCH₃; x=1 to 24; n≥1; m=1; COCH₃-capped silicone polyethers

Commercial preparations of the silicone surfactants above may or may not contain small amounts of polyethylene glycols (PEG) or other low molecular weight polydimethyl siloxanes (PDMS).

In embodiments of the invention, silicone surfactant is added in a range of about 0.1 wt % to about 5 wt % in the non-aqueous portion of the oil-in-water emulsion.

In an embodiment of the invention, the silicone surfactant, added in an amount of about 2 wt % in the non-aqueous portion of the oil-in-water emulsion, is an H-capped dimethyl methyl (polyethylene oxide) silicone polymer having a molecular weight from 200-6000 as shown below in Formula (II):



where: n=2-70, the average n=44; m=2-16 and the average m=10.

Suitable emulsifiers are selected and added in amounts so as to generate a stable emulsion and to prevent phytotoxicity. The emulsifier is added in a range from about 0.5 wt % to about 5 wt % in the non-aqueous portion of the oil-in-water

emulsion. More particularly, in embodiments of the invention, the emulsifier is added in a range from about 1 wt % to about 3 wt % in the non-aqueous portion of the oil-in-water emulsion. In an embodiment of the invention, the emulsifier is added at about 2 wt % in non-aqueous portion of the oil-in-water emulsion.

The emulsifier is selected from the group consisting of:

Alcohol ethoxylates (natural and synthetic) including polyoxyethylene (4 to 12) lauryl ether (C12); polyoxyethylene (10) cetyl ether (C16); polyoxyethylene (10) stearyl ether (C18); polyoxyethylene (10) oleyl ether (C18 monounsaturated); polyoxyethylene (2 to 11) C12-C15 alcohols; polyoxyethylene (3 to 9) C11-C14 alcohols; polyoxyethylene (9) C12-C14 alcohols; polyoxyethylene (11) C16-C18 alcohols; and polyoxyethylene (20) C12-C15 alcohols;

Alcohol alkoxylates including butyl polyoxyethylene/polyoxypropylene block copolymer;

Alkyl polysaccharides including C8-C11 alkylpolysaccharides;

Glycerol oleate;

Polyoxyethylene-polyoxypropylene block copolymers of MW 1100 to 11400 and 10 to 80% EO;

Nonyl phenol ethoxylates including polyoxyethylene (2 to 8) nonylphenol;

Polymeric surfactants including graft copolymer such as polymethacrylic acid and acrylate with polyoxyethylene chains and random copolymers with ester and ether groups;

Polyethylene glycols including MW: 200 to 8000; MW: 400 PEG dioleate; and MW: 600 PEG dioleate; and

Sorbitan fatty acid ester ethoxylates including polyoxyethylene (20) sorbitan tristearate; polyoxyethylene (20) sorbitan monooleate; polyoxyethylene (5) sorbitan monooleate; and polyoxyethylene (20) sorbitan trioleate.

Examples of 2-Pack Embodiment

In embodiments of the invention, a sufficient volume of Pack A was added to a volume of water required to obtain a desired concentration in the final formulation and was mixed to form an emulsion. Sufficient Pack B was added to the emulsion to result in a fungicidal dispersion which comprised:

an effective amount of paraffinic oil, being about 5 wt %; about 0.1 wt % emulsifiers;

about 0.3 wt % polychlorinated Cu (II) phthalocyanine;

and

about 0.1 wt % silicone surfactant.

The fungicidal dispersion was sprayed onto turfgrass at a rate of from about 50 gal/acre (4.7 L/100 m²) to about 100 gal/acre (9.3 L/100 m²).

Example 5

In an example of the 2-Pack embodiment of the invention, the following compositions (Pack A and Pack B) were prepared and mixed together as described to achieve the sprayable dispersion.

Pack A

Components	Chemical Description	Supplier Name	% by weight	Purpose in Formulation
1	N65DW	Petro-Canada	96	Active Ingredient
	Highly saturated paraffinic oils with a carbon number distribution in the range of about C ₁₆ to about C ₃₅			

-continued

Components	Chemical Description	Supplier Name	% by weight	Purpose in Formulation
2 PC Emuls Green	1. Ethoxylated alcohols having primary C5-C20 carbon chains with an average of about 2 to about 7 ethoxylation groups 2. Glycerol Oleate	Petro-Canada	2	Emulsifier
3 Lambent MFF199 SW	Methyl (propylhydroxide, ethoxylated) bis (trimethylsiloxy) silane	Lambent Technologies Corp., Gurnee IL, USA	2	Wetting agent/dispersant

Pack B

Components	Chemical Description	Supplier Name	% by weight	Purpose in Formulation
1 SUNSPERSE® Green 7 (GCD9957)	58% polychlorinated Cu(II) phthalocyanine (C ₃₂ HCl ₁₅ CuN ₈) dispersed in water	Sun Chemical Corp Performance Pigment Cincinnati OH USA	100	Colorant

In the embodiment of the invention described in Example 5, all of the silicone surfactant was added to Pack A. ²⁵

Example 6

In an example of the 2-Pack embodiment of the invention, the following components or compositions were prepared and mixed together as described to achieve the sprayable dispersion. ³⁰
Pack A

Components	Chemical Description	Supplier Name	% by weight	Purpose in Formulation
1 N65DW	Highly saturated paraffinic oils with a carbon number distribution in the range of about C ₁₆ to about C ₃₅	Petro-Canada	98	Active Ingredient
2 PC Emuls Green	1. Ethoxylated alcohols having primary C ₅ -C ₂₀ carbon chains with an average of about 2 to about 7 ethoxylation groups 2. Glycerol Oleate	Petro-Canada	2	Emulsifier

Pack B

Components	Chemical Description	Supplier Name	% by weight	Purpose in Formulation
1 SUNSPERSE® Green 7 (GCD9957)	58% polychlorinated Cu(II) phthalocyanine (C ₃₂ HCl ₁₅ CuN ₈) dispersed in water	Sun Chemical Corp Performance Pigment Cincinnati, OH USA	83.8	Colorant
2 Lambent MFF159-10	Dimethyl, methyl (polyethylene oxide) silicone polymer	Lambent Technologies Corp., Gurnee IL USA	16.2	Wetting agent/dispersant

11

In the embodiment of the invention described in Example 6, all of the silicone surfactant was added to Pack B.

Examples 7 to 10

2-Pack Formulation

Examples 7 to 10 were performed using a formulation prepared according to Example 5.

Example 7

Comparison of the Efficacy of the 2-Pack Formulation to a Formulation Without Pigment

An embodiment of the invention was applied to creeping bentgrass (*Agrostis stolonifera*) at a close cut putting green height of 5 mm to determine the efficacy of an embodiment of the invention to control dollar spot. Dollar spot is caused by *Scierlinia Homeocarpa* and is the most common disease requiring control on golf courses throughout most of the world.

The embodiment of the invention, applied to the turfgrass at 5 gal/acre (0.5 L/100 m²) of the non-aqueous portion of the oil-in-water emulsion, was compared to the use a higher concentration of oil-in-water emulsion without pigment, applied at 10 gal/acre (1 L/100 m²) of the non-aqueous portion of the oil-in-water emulsion, and an inoculated control. The plots were inoculated with five strains of *Scierlinia Homeocarpa*, one week after initial chemical application.

The fungicides were applied in water at a total spray volume rate of about 129 gal/acre (12 L/100 m²) using a wheel-mounted compressed air boom sprayer at 140 kPa. Applications were made at 14 day intervals. Weekly counts of dollar spot infection were conducted over the 6 week period.

Chemical Treatment List

Treatment	Product per 100 m ²	No. of Applications	Treatment Interval
Fungicide with pigment (2-pack)	465 mL	3	14 days
Fungicide without pigment	930 mL	3	14 days

Conclusions

As shown in FIG. 1, acceptable control of dollar spot was achieved using the 2-pack fungicide at 5 gal/acre (0.5 L/100 m²) with the addition of the polychlorinated Cu (II) phthalocyanine pigment dispersion, emulsifier and silicone additives.

The use of 5 gal/acre (0.5 L/100 m²) of the non-aqueous portion of the 2-pack fungicide formulation performed substantially the same as using the oil-in-water emulsion alone at 10 gal/acre (1 L/100 m²) of the non-aqueous portion. The data suggests a synergistic effect resulting from the addition of pigment, permitting use of the formulation at rates lower than predicted rates.

An additional benefit of the lower treatment rate is the reduced propensity for phytotoxicity which may be observed at high oil rates.

Example 8

A 2-Pack embodiment of the invention was applied to creeping bentgrass (*Agrostis stolonifera*) at a close cut putting

12

green height of 5 mm to determine the efficacy of an embodiment of the invention to control dollar spot. Close cut turf is highly susceptible to dollar spot.

The 2-pack embodiment of the invention was compared to a chemical fungicide, DACONIL® 2787, also known as WEATHERSTIK™, available from Syngenta Crop Protection Canada, Inc. Guelph, Ontario, Canada, and an inoculated control. The trials were conducted on 8 year old PENNCROSS® creeping bentgrass (*Agrostis stolonifera*). The treatments were applied over a 7-week period and the turf and disease were monitored over a 9-week period to examine residual effects and grass recovery. A randomized complete block design having four replications was used and each treatment plot measure 1 m×2 m. The plots were inoculated with five strains of *Scierlinia Homeocarpa*, one day after initial chemical application.

Due to the close cutting of the turf in this example the disease pressure was extreme. The fungicides were applied in water at a total spray volume rate of 118 gal/acre (11 L/100 m²) using a wheel-mounted compressed air boom sprayer at 140 kPa. Applications were made at 14 day intervals for both the DACONIL® and the 2-pack formulation. An additional plot was treated with applications of the 2-pack formulation at 7 day intervals. Weekly counts of dollar spot infection were conducted over the 7 week period.

Chemical Treatment List

Treatment	Product per 100 m ²	No. of Applications	Treatment Interval
DACONIL® 2787	95 mL	4	14 days
2-pack fungicide	465 mL	4	14 days
2-pack fungicide	465 mL	7	7 days

Results

As shown in FIG. 2, low levels of dollar spot disease were present on all of the plots at the start of the trial. Nearing the end of the trial, the disease levels in the control exceeded 100 spots per plot on the inoculated areas. All three chemical treatments showed significant suppression of dollar spot disease when compared to the control. The suppression of disease continued for about two weeks following the last application.

It was noted that when applied weekly, the 2-pack formulation performed substantially better than the 2-pack formulation or the DACONIL® applied at the recommended rate of biweekly. One of skill in the art would recognize therefore that, in cases of extreme dollar spot disease, efficacy of the 2-pack formulation is improved if applied more frequently.

Phytotoxicity was not observed in any of the treatments.

Conclusions

The 2-pack fungicide, applied at 5 gal/acre (0.5 L/100 m²) of the non-aqueous portion performed substantially the same as the chemical fungicide, DACONIL® 2787.

In cases of extreme disease pressure the 2-pack formulation should be applied more frequently.

Example 9

An embodiment of the invention was applied to creeping bentgrass (*Agrostis stolonifera*) at a fairway height of 11 mm to determine the efficacy of an embodiment of the invention to control dollar spot. Dollar spot is caused by *Scierlinia Homeocarpa* and is the most common disease requiring control on golf courses throughout most of the world.

13

The embodiment of the invention was compared to a chemical fungicide, DAPONIL® 2878, also known as WEATHERSTIK™, available from Syngenta Crop Protection Canada, Inc. Guelph, Ontario, Canada, and an inoculated control. The trials were conducted on 13 year old PENNCROSS® creeping bentgrass (*Agrostis stolonifera*). The treatments were applied over a 6-week period and the turf and disease were monitored to examine residual effects and grass recovery. A randomized complete block design having four replications was used and each treatment plot measure 1 m×2 m. The plots were inoculated with five strains of *Sclerotinia Homeocarpa*, one week after initial chemical application.

The fungicides were applied in water at a total spray volume rate of 129 gal/acre (12 L/100 m²) using a wheel-mounted compressed air boom sprayer at 140 kPa. Applications were made at 14 day intervals. Weekly counts of dollar spot infection were conducted over the 6 week period.

Chemical Treatment List

Treatment	Product per 100 m ²	No. of Applications	Treatment Interval
DAPONIL® 2878	95 mL	3	14 days
2-pack fungicide	465 mL	3	14 days
2-pack fungicide	930 mL	3	14 days

Results

As shown in FIG. 3, dollar spot disease was present on all of the plots at the start of the trial. All three treatments showed significant suppression of dollar spot disease when compared to the control. The suppression of disease continued for about two weeks following the last application.

Phytotoxicity was not observed in any of the treatments.

Conclusions

The 2-pack fungicide, applied at both 5 gal/acre (0.5 L/100 m²) of the non-aqueous portion and 10 gal/acre (1 L/100 m²) of the non-aqueous portion, performed substantially the same as the chemical fungicide, DAPONIL® 2878.

Example 10

An embodiment of the 2-pack formulation was used to treat dollar Spot on *Poa Trivialis* (rough stalk bluegrass) over-seeded Bermudagrass plot cut at putting green height. The 2-pack formulation was applied at 2.5 gal/acre (0.2 L/100 m²) of the non-aqueous portion and was compared to a formulation prepared without the pigment, the silicone surfactant and the particular emulsifier and which was applied at 5 gal/acre (0.5 L/100 m²) of the non-aqueous portion and to an untreated control.

The formulations were applied using a CO₂ backpack boom sprayer calibrated to deliver products in 2 gallons of water per 1000 sq ft (8 L/100 m²) through two 8003 TEE-JET® flat fan nozzles.

The turfgrass plots were divided into 4 blocks and treatments were assigned in a randomized complete block design. The plots were inoculated with 1 L of wheat seed infested with *Sclerotinia homoeocarpa*. Fungicide formulations were applied weekly for 8 weeks.

Having reference to FIG. 4, dollar spot counts were made throughout the 8 week period.

Conclusions:

Acceptable control of dollar spot was obtained through weekly use of 2.5 gal/acre (0.2 L/10 m²) of the non-aqueous portion of 2-pack fungicide which represents approximately

14

one half the amount required to achieve the same result using a formulation without the addition of the polychlorinated Cu (II) phthalocyanine pigment dispersion, emulsifier and silicone additives (5 gal/acre (0.5 L/100 m²) of the non-aqueous portion, weekly).

Advantageously, the use of a lower treatment rate reduces the propensity for phytotoxicity which may be observed at high rate oil rates.

Example 11

Example 11 was performed using a formulation prepared according to Example 6.

Kentucky bluegrass was mowed three times per week at a cutting height of 2 inches. Applications of the 2-pack formulation were made at 14-day intervals beginning April 10. Assessment of Spring leaf spot or Melting-out disease was made using a "0 to 10" severity index, where 10 is equivalent to greater than 90% symptomatic turf area.

For the purposes of comparison to conventional treatment, DAPONIL® ULTREX, available from Syngenta Crop Protection Canada, Inc. Guelph, Ontario, Canada, was used as a control.

Results

Treatment	Application rate (oz/1000 ft ²)	Disease Severity			
		14-May	22-May	28-May	4-Jun
Untreated		2.7ab	4.3a	5.3a	6.7a
2-pack	21.75	0.0e	0.3ef	0.3ef	0.3hi
DAPONIL® Ultrex	3.2	0.0e	0.0f	0.0f	0.0i
LSD (P = 0.05)		1.45	0.96	1.38	1.28
Std Deviation		0.89	0.59	0.85	0.78

Means in a row followed by the same letter are not significantly different (alpha = 0.05) using LSD test.

No phytotoxicity was observed in any treatment plot.

Conclusions:

The 2-Pack formulation, used at 7.5 gal/acre (0.7 L/100 m²) of the non-aqueous portion) provided excellent disease control.

1-Pack Formulation

In another embodiment of the invention, Applicant has surprisingly found that pigment, such as polychlorinated Cu (II) phthalocyanine, can be prepared in a single, stable dispersion in an oil-based fungicidal composition, prior to dilution in water to form a sprayable dispersion. Thus, the user does not need to handle two separate components for preparing a stable fungicidal application.


More specifically, the pigment is dispersed in a compatible oil, such as a paraffinic oil or the same paraffinic oil as is used to provide the fungicidal properties as described in embodiments of the invention, for additional to the formulation. Use of specific silicone surfactant and emulsifier chemistries stabilizes the colorant in the oil-based composition.

In examples of the 1-pack formulation, polychlorinated Cu (II) phthalocyanine is dispersed in a paraffinic oil, such as N65DW (available from Petro-Canada) to provide about 18% polychlorinated Cu (II) phthalocyanine (SUNSPERSE® EXP 006-102, available from Sun Chemical Corp. Performance Pigments, Cincinnati, Ohio USA) prior to mixing with the remaining constituents.

15

In embodiments of the invention, the 1-pack formulation comprises silicone surfactant, emulsifier and polyethylene glycols which are selected to provide an intermolecular hydrophilic and lipophilic balance within the fungicidal formulation so as to substantially prevent the polychlorinated Cu (II) phthalocyanine from separating out of suspension during application to the turf grass.

Applicant believes that suitable silicone surfactants are those previously identified for the 2-pack formulation, as described in Formula (I). In embodiments of the invention the silicone surfactants are H-capped, CH₃-capped and COCH₃-capped trisiloxanes. In an embodiment of the invention, the silicone is an H-capped trisiloxane as shown below in Formula (III):



Where: n=1-24, average n=8-10.

Suitable emulsifiers for the 1-Pack formulation are selected from the following:

Alcohol ethoxylates (natural and synthetic) including polyoxyethylene (4 to 7) lauryl ether (C12); polyoxyethylene (10) cetyl ether (C16); polyoxyethylene (2 to 11) C12-C15 alcohols; polyoxyethylene (3 to 9) C11-C14 alcohols; polyoxyethylene (9) C12-C14 alcohols; Polymeric surfactants including graft copolymer such as polymethacrylic acid and acrylate with polyoxyethylene chains and random copolymers with ester and ether groups; and

Sorbitan fatty acid esters including sorbitan tristearate and sorbitan trioleate.

The emulsifier is added to the formulation in a range of about 0.5 wt % to about 5 wt % in the non-aqueous portion of the formulation.

In embodiments of the invention, Applicant has noted that the silicone surfactants used typically comprise 10-30% polyethylene glycols (PEG) according to Formula (IV) shown below.



Where:

R₁=H or CH₂=CH—CH₂ or COCH₃

R₂=H or CH₂=CH—CH₂ or COCH₃

n≥1

In embodiments of the invention, the PEG has a low molecular weight, typically about 300 to about 1500 Daltons. In embodiments of the invention, the PEG is a low molecular weight polyethylene glycol allyl ether. In embodiments of the invention, the PEG is a low molecular weight polyethylene glycol mono-allyl ether having an average molecular weight of from about 300 to about 600 Daltons and having from 1 to 20 moles of ethylene glycol with an average ethoxylation (EO) of 8 to 10.

Examples 12 TO 14

1-Pack Formulations

Example 12

28 wt % SUNSPERSE® EXP 006-102 containing 18 wt % polychlorinated Cu (II) phthalocyanine, dispersed in N65DW;

16

2 wt % silicone surfactant according to Formula (I) where: m=1, n=0, X=1-24 (average 8-10) and R=H; and PEG according to Formula (IV) where: R₁=CH₂=CH—CH₂, R₂=H and n=1-20 with an average n=8 (Lambert MFF199);

2 wt % polyoxyethylene (11) C₁₆₋₁₈ alcohols such as LUTENSOL® AT11 available from BASF; and

68 wt % N65DW.

Example 13

28 wt % SUNSPERSE® EXP 006-102 containing 18 wt % polychlorinated Cu (II) phthalocyanine, dispersed in N65DW;

2 wt % silicone surfactant according to Formula (I) where: m=1, n=0, X=1-24 (average 8-10) and R=H; and PEG according to Formula (IV) where: R₁=CH₂=CH—CH₂, R₂=H and n=1-20 with an average n=8 (Lambert MFF199);

2 wt % sorbitan tristearate such as SPAN65 available from Uniqema or S-MAZ® 65K available from BASF; and

68 wt % N65DW.

Example 14

28 wt % SUNSPERSE® EXP 006-102 containing 18 wt % polychlorinated Cu (II) phthalocyanine, dispersed in N65DW;

1.8 wt % silicone surfactant as described in Formula (I) where m=1, n=0, X=1-24 (average 8-10) and R=COCH₃, and Polyethylene Glycols as described in Formula (IV) where R₁=CH₂=CH—CH₂ or COCH₃, R₂=COCH₃ (SYL-GARD® 309, available from Dow Corning, USA)

0.2% Polyethylene Glycols as described in Formula (IV) where R₁=CH₂=CH—CH₂, R₂=H (Polyglykol A500, available from Clariant);

2 wt % polyoxyethylene (11) C₁₆₋₁₈ alcohols, such as LUTENSOL® AT11 available from BASF; and

68 wt % N65DW.

The formulations disclosed in Examples 12-14 were further diluted to 6% in water prior to application to the turfgrass. The resulting oil-in-water formulations comprised:

about 5 wt % paraffinic oil;

about 0.12 wt % emulsifier;

about 0.3 wt % polychlorinated Cu (II) phthalocyanine;

about 0.1 wt % silicone surfactant;

about 0.01-0.03 wt % polyethylene glycol; and

the balance being water.

The resulting oil-in-water emulsion/pigment dispersions were found to be stable until applied, typically within one day and can be applied to turfgrass. The oil-in-water emulsions were applied at a conventional total spray volume rate of about 50 gal/acre (4.7 L/100 m²) to about 100 gal/acre (9.3 L/100 m²).

Example 15-17

Additional testing was performed to determine the stability of 1-pack formulations without the presence of silicone surfactant and/or PEG.

In Examples 15-17, the formulations were further diluted to 6% in water to form the oil-in-water emulsion prior to application to turfgrass.

17

Example 15

A 1-pack formulation was prepared with neither silicone surfactant nor PEG according to the following formula:

- 28 wt % SUNSPERSE® EXP 006-102 containing 18 wt % polychlorinated Cu (II) phthalocyanine, dispersed in N65DW;
- 2 wt % polyoxyethylene (11) C₁₆₋₁₈ alcohols such as LUTENSOL AT11 available from BASF; and
- 70% N65DW.

Example 16

A 1-pack formulation was prepared with insufficient PEG according to the following formula:

- 28 wt % SUNSPERSE® EXP 006-102 containing 18 wt % polychlorinated Cu (II) phthalocyanine, dispersed in N65DW;
- 2 wt % silicone surfactant as described in Formula (I) where m=1, n=0, X=1-24 (average 8-10) and R=H (SILTECH® SILSURF A008-UP);
- 2 wt % polyoxyethylene (11) C₁₆₋₁₈ alcohols such as LUTENSOL® AT11 available from BASF; and
- 68% N65DW.

Example 17

A 1-pack formulation was prepared without silicone surfactant according to the following formula:

- 28 wt % SUNSPERSE® EXP 006-102 containing 18 wt % polychlorinated Cu (II) phthalocyanine, dispersed in N65DW;
- 2 wt % Polyethylene Glycols as described in Formula (IV) where R₁=CH₂=CH—CH₂, R₂=H (Polyglykol A500, available from Clariant);
- 2 wt % polyoxyethylene (11) C₁₆₋₁₈ alcohols such as LUTENSOL® AT11 available from BASF; and
- 68% N65DW.

Results:

The pigment aggregated and fell out of suspension in all of the formulations tested in Examples 15-17, rendering the formulations unusable.

Conclusions:

As shown in Examples 12-17, without both silicone surfactant and sufficient amounts of PEG, the resulting 1-pack formulations, diluted to form oil-in-water emulsions, are not stable and are therefore not usable for turf application.

As demonstrated in Examples 12-17, the presence of silicone surfactants and polyethylene glycols improve the stability and dispersibility of the 1-Pack formulation resulting in a commercially viable fungicidal dispersant for turf application and management of disease therein.

Example 18

Embodiments of the 1-pack and 2-pack formulations, according to the invention, and a conventional fungicide, DACONIL® Ultrex, were used to treat dollar Spot on *Poa Trivialis* (rough stalk bluegrass) overseeded Bermudagrass plot cut at putting green height. As with Example 10, the formulations were applied with a CO₂ backpack boom sprayer calibrated to deliver products in 2 gallons of water per 1000 sq ft (8 L/100 m²) through two 8003 TEEJET® flat fan nozzles.

The plots were divided into 4 blocks and treatments were assigned in a randomized complete block design. The plot

18

was inoculated with 1 L of wheat seed infested with *Sclerotinia homoeocarpa*. The formulations were applied biweekly for 8 weeks.

As shown in FIG. 5, dollar spot counts were made throughout the 8 week period.

Conclusions:

Both the 1-pack and 2-pack formulations had substantially the same efficacy as the DACONIL® Ultrex.

Example 19

A 2-Pack formulation was prepared according to Example 6 an applied on a fairway and an area of rough at the Saginaw Golf Course in Cambridge, Ontario, Canada for an 8 week period using a 14-day application interval.

At the end of 8 weeks, core samples of the treated and untreated areas were obtained to compare root development. Results

As shown in FIG. 6, the grass on the treated areas (left) was much denser and healthier than on the untreated areas (right). Further, the treated areas had a darker green color than the untreated areas.

It was found that the roots of the bentgrass on the treated area were longer by approximately twice the length of the roots of the untreated bentgrass. Further, the roots were observed to be denser than the roots of untreated bentgrass.

Conclusions:

The 2-Pack and 1-Pack formulations promote the growth of bentgrass and bluegrass.

Example 20

Having reference to FIGS. 7 and 8, areas of bentgrass were treated using a 2-Pack formulation prepared according to Example 6. Seven (7) core samples were randomly taken from treated areas and from untreated areas for comparison. The core samples were soaked overnight in water in a glass tray. Subsequently, soil from the roots was washed away with water to reveal the root structure, as shown in FIGS. 7 and 8 and summarized in Table A below.

Results

TABLE A

Core sample #	Untreated root length (inches)	Treated root length (inches)
1	3.5	5.5
2	3.0	4.5
3	3.75	4.0
4	2.75	5.25
5	2.75	4.5
6	2.75	4.75
7	4.25	4.75
Average	3.25	4.75

Conclusions:

On the average, the root length of the treated bentgrass is approximately 50% longer than that of the untreated bentgrass. Further it was noted that the root mass was considerably greater for the treated bentgrass.

Enhanced Formulation

Applicant has determined that embodiments of the present invention have a surprising synergistic effect when mixed with some conventional systemic chemical fungicides selected from the group consisting of demethylation inhibitors (such as propiconazole), methyl benzimidazole carbam-

19

ate (such as thiophanate-methyl) and dicarboximide (such as iprodione). As an example, a suitable propiconazole fungicide is BANNER MAXX™ (available from Syngenta Crop Protection Canada, Inc. Guelph, Ontario, Canada), a thiophanate-methyl fungicide is Cleary 3336™ (available from Cleary Chemical Corporation, Dayton, N.J., USA) and an iprodione fungicide is ROVRAL® Green (available from Bayer Environmental Science-Canada, Guelph, Ontario, Canada).

Applicant has found however that, despite predictions to the contrary, the synergistic effect was not observed with all conventional chemical fungicides, such as some contact fungicides, one of which was chlorothalonil (such as DICONIL® Ultrex, available from Syngenta Crop Protection Canada, Inc. Guelph, Ontario, Canada).

As shown in the following additional examples for treatment of Dollar Spot, snow moulds and gray leaf spot, mixing an embodiment of the present invention at half the predicated recommended label rate in equal parts with certain conventional chemical fungicides at half the amount of the label rate resulted in an equivalent or improved efficacy of the mixture when compared to each of the fungicides used alone at full label rate.

As one of skill would understand and as taught by Burpee and Latin, Plant Disease Vol. 92 No. 4, April 2008, pp 601-606, knowing that the efficacy of fungicides is not additive, the addition of half the label rate of each of the two fungicidal formulations would not conventionally be thought to result in high efficacy. Surprisingly this was not the case with embodiments of the present invention. Mixtures of formulations of the present invention and certain conventional chemical fungicides, at rates not thought to be useful, resulted synergistically in a highly efficacious formulation.

Applicant believes that the amount of formulations according to embodiments of the invention can be reduced to a range from about 25% to about 75% of the amount used if used alone, when mixed with the conventional chemical fungicide, also reduced to from about 25% to about 75% of the recommended label rate.

Examples 21 and 22

Examples 21 and 22 illustrate results of use of the 1-pack formulation, the two pack formulation and the enhanced formulation on the treatment of Dollar Spot on *Poa Trivialis* (rough stalk bluegrass) cut at putting green height.

Example 21

Having reference to FIGS. 9A and 9B, the 2-pack formulation was tank mixed at 50:50 with chemical fungicide BANNER MAXX™ and applied to turf infected with dollar spot disease.

The mixture of the 2-pack fungicide and the chemical fungicide exhibited a synergistic effect compared to using either of the fungicides alone. As shown in FIG. 9B, the residue period can be extended from 21 days to 28 days.

Example 22

Having reference to FIG. 10, the 1-pack formulation was tank mixed 50:50 with chemical fungicide BANNER MAXX™ and applied to turf infected with dollar spot disease.

20

The 1-pack formulation with the addition of chemical fungicide exhibited a synergistic effect when compared to use of the 1-pack formulation or BANNER MAXX™ alone, when applied at the full label rate.

Examples 23-25

Embodiments of the invention were mixed 50:50 with conventional chemical fungicide BANNER MAXX™ to form an enhanced formulation and were applied to creeping bentgrass (*Agrostis stolonifera*) at fairway height which was infected with a variety of snowmolds.

An enhanced formulation containing the 2-pack formulation and chemical fungicide was compared to use of the fungicides alone. BANNER MAXX™ was applied at 50% the recommended label rate for comparison to the enhanced formulation.

Example 23

As shown in FIG. 11, the effectiveness of the enhanced formulation was compared to the 2-pack formulation, 50% of the recommended label rate of BANNER MAXX™ alone and an untreated control for treating turf infected with *Typhula ishikariensis*.

Example 24

As shown in FIG. 12, the effectiveness of the enhanced formulation was compared to the 2-pack formulation, 50% of the recommended label rate of BANNER MAXX™ alone and an untreated control for treating turf infected with *Typhula incarnata*.

Example 25

As shown in FIG. 13, the effectiveness of the enhanced formulation was compared to the 2-pack formulation, 50% of the recommended label rate of BANNER MAXX™ alone and an untreated control for treating turf infected with *Microdochium nivale*.

Conclusions:

It is clear from Examples 23-25 that use of the enhanced formulation comprising 50% of the recommended rate of an embodiment of the invention and 50% of the recommended label rate of BANNER MAXX™ is as effective, or more effective, than using either the 2-pack formulation or the chemical fungicide alone.

Example 26

A 2-Pack formulation was prepared according to Example 6 using half the recommended rate (i.e. 232.5 ml per 100 m² for the non-aqueous portion) and was mixed with a commercial fungicide ROVRAL® Green GT (available from Bayer Environmental Science-Canada, Guelph, Ontario, Canada) at 50% the recommended label rate (i.e., applied at 125 ml per 100 m²) to form the enhanced formulation.

The enhanced formulation was applied to creeping bentgrass (*Agrostis stolonifera*) cut to greens height for the control of Fusarium Patch disease. ROVRAL® Green GT was also applied alone at 50% the recommended rate (i.e., applied at 125 ml per 100 m²) for comparison.

The treatments were applied over a 6-week period and disease was monitored weekly.

Conclusions:

As shown in FIG. 14, the enhanced formulation ("PC2 50%+ROVRAL® Green GT 50%") provided significant suppression of injury by Fusarium Patch disease.

Example 27

One half (50%) of the recommended rate of a 2-Pack formulation (i.e., at 2.5 gal per acre of the non-aqueous portion), prepared according to Example 6, was mixed 50:50 with a conventional chemical fungicide Cleary 3336™ Plus at 50% the recommended rate (i.e., at 2 oz per 1000 ft²).

The enhanced formulation (PC2, ½ rate+Cleary 3336™ Plus, ½ rate), as well as the individual formulations at full rate and at half rate, were applied to perennial ryegrass cut to fairway height for control of Gray Leaf Spot disease.

Conclusions:

As shown in FIG. 15, the enhanced formulation provided excellent control of Gray Leaf Spot disease on Ryegrass.

Synergistic Effect of the Oil/Emulsion and Pigment Dispersion

Example 28

As shown in FIG. 16, Applicant believes that in all formulations according to embodiments of the invention, the addition of the polychlorinated Cu (II) phthalocyanine has a synergistic effect when compared to oil/emulsifier alone or pigment alone, when compared to an untreated control.

Turf infected with Dollar Spot disease was treated with 10% oil/emulsifier alone, 0.5% SUNSPERSE® Green 7 alone and an embodiment of the invention comprising both 10% oil/emulsifier and 0.5% SUNSPERSE® Green 7.

Conclusions:

Clearly, it appears that addition of pigment to the fungicidal oil component enhances the fungicidal properties of the dispersion.

Use of Embodiments of the Invention for Controlling Additional Pests in Turfgrass

Example 29

Fall Armyworms

The efficacy of an embodiment of the 2-pack formulation against different larval stages of fall armyworms (*Spodoptera frugiperda*) was determined under laboratory conditions. Embodiments of the invention were compared to a conventional pesticide Cyfluthrin.

Methods:

1) Larvae were obtained from Benzon Research Inc.

2) 5 larvae/container and 3 containers=1 replication

3) Treatments=1) Control

2) 2-pack formulation at 5% dilution

3) Cyfluthrin (insecticide), AI=0.75%, 3 fl oz/gal H₂O (22 mL/L)

4) Container=760 ml plastic bowl with lid containing moist filter paper on bottom and grass clippings

5) St. Augustinegrass clippings dipped into respective fungicide treatment, drained to dry and put into container

6) Hold 4 days at 25° C. and measure survival

7) 7 replications of small larvae (2-10 mm) tested

8) 6 replications of medium larvae (11-20 mm) tested

9) 6 replications of large larvae (21-30 mm) tested

10) Statistical analysis=Least Significant Difference (LSD) test

Results:

Fall army worms	Mean survival ^a		
	control	Cyfluthrin	2-pack formulation
Larvae			
Small	3.86 A	0 B	0.43 B
Medium	3.50 A	0 B	1.17 B
Large	3.00 A	0 B	2.50 A

^aMeans in a row followed by the same letter are not significantly different (alpha = 0.05) using LSD test.

Conclusions:

The data indicates that embodiments of the 2-pack formulation are effective to kill small and medium larvae of fall army worms.

Tropical Sod Webworms

Similar testing was conducted to determine the efficacy of embodiments of the 2-pack formulation to kill all sizes larvae (small, medium and large) of tropical sod webworms.

Sodweb worms	Mean survival ^a		
	control	Cyfluthrin	PC
Larvae			
Small	3.0 A	0 B	0.3 B
Medium	4.8 A	0 B	0 B
Large	4.0 A	0.3 B	1.0 B

^aMeans in a row followed by the same letter are not significantly different (alpha = 0.05) using LSD test.

Conclusions:

The data indicates that embodiments of the 2-pack formulation are effective to kill small, medium and large larvae of tropical sod webworms.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A formulation for application to a turfgrass comprising:

a paraffinic oil;

one or more emulsifiers, each of which is independently selected from the group consisting of natural and synthetic alcohol ethoxylates, alcohol alkoxylates, glycerol oleate, and nonyl phenol ethoxylates;

a polychlorinated (Cu II) phthalocyanines;

a silicone surfactant; and

water,

wherein the silicone surfactant represents from about 0.1 wt % to about 5 wt % of the total weight of the paraffinic oil, the one or more emulsifiers, the polychlorinated (Cu II) phthalocyanine, and the silicone surfactant;

wherein the one or more emulsifiers represent from about 0.5 wt % to about 5 wt % of the total weight of the paraffinic oil, the one or more emulsifiers, the polychlorinated (Cu II) phthalocyanine, and the silicone surfactant;

wherein the polychlorinated (Cu II) phthalocyanines represents from about 1 wt % to about 10 wt % of the total weight of the paraffinic oil, the one or more emulsifiers, the polychlorinated (Cu II) phthalocyanine, and the silicone surfactant; and

wherein the formulation is an oil-in-water emulsion and the one or more emulsifiers and the silicone surfactants are selected such that the polychlorinated (Cu II) phthalocyanines is maintained in dispersion in the oil-in-water emulsion for delivery to the turfgrass.

23

2. The fungicidal formulation of claim 1 wherein the emulsifier is selected from the group consisting of polyoxyethylene(4 to 7) lauryl ether (C_{12}); polyoxyethylene (10) cetyl ether (C_{16}), polyoxyethylene (2 to 11) C_{12} - C_{15} alcohols, polyoxyethylene (3 to 9) C_{11} - C_{14} alcohols, and polyoxyethylene (9) C_{12} - C_{14} alcohols, and the silicone surfactant comprises polyethylene glycols (PEG) according to Formula (IV):



where: $R_1=H$ or $CH_2=CH-CH_2$ or $COCH_3$

$R_2=H$ or $CH_2=CH-CH_2$ or $COCH_3$


n is greater than or equal to 1.

3. The formulation of claim 1, wherein the non-aqueous portion of the oil-in-water emulsion is delivered to the turfgrass in a range of about 1 gal/acre (0.093 L/100 m²) to about 15 gal/acre (1.395 L/100 m²), a total spray volume of the oil-in-water emulsion being in a range of about 20 gal/acre (1.86 L/100 m²) to about 200 gal/acre (18.6 L/100 m²).

4. The formulation of claim 1, further comprising one or more chemical fungicides, each of which is independently selected from the group consisting of a demethylation inhibitor, a methyl benzimidazole carbamate and a dicarboximide.

5. The formulation of claim 1, wherein:

the silicone surfactants is selected from the group consisting of trisiloxanes and silicone polyethers of Formula I



where $R=H$, CH_3 , or $COCH_3$; $x=1$ to 24; $n=0$ or ≥ 1 ; and $m=1$.

6. The formulation of claim 1, wherein a non-aqueous portion of the oil-in-water emulsion is delivered to the turfgrass in a range of about 1 gal/acre (0.093 L/100 m²) to about 15 gal/acre (1.395 L/100 m²).

7. The formulation of claim 1, wherein each of the one or more emulsifiers is selected from the group consisting of polyoxyethylene(4 to 12) lauryl ether (C_{12}), polyoxyethylene (10) cetyl ether (C_{16}), polyoxyethylene (10) stearyl ether (C_{18}), polyoxyethylene (10) oleyl ether (C_{18} monounsaturated), polyoxyethylene (2 to 11) C_{12} - C_{15} alcohols, polyoxyethylene (3 to 9) C_{11} - C_{14} alcohols, polyoxyethylene (9) C_{12} - C_{14} alcohols, polyoxyethylene (11) C_{16} - C_{18} alcohols, polyoxyethylene (20) C_{12} - C_{15} alcohols, glycerol oleate, polyoxyethylene (2 to 8) nonylphenol, and mixtures thereof.

8. The formulation of claim 1, wherein the one or more emulsifiers is selected from a natural or synthetic alcohol ethoxylate and glycerol oleate.

9. The formulation of claim 1, wherein the one or more emulsifiers represent from about 1 wt % to about 3 wt % of the total weight of the paraffinic oil, one or more emulsifiers, polychlorinated (Cu II) phthalocyanine, and silicone surfactant.

10. The formulation of claim 1, wherein the one or more emulsifiers represent about 2 wt % of the total weight of the paraffinic oil, one or more emulsifiers, polychlorinated (Cu II) phthalocyanine, and silicone surfactant.

11. The formulation of claim 1, wherein each of the one or more emulsifiers is independently selected from the group

24

consisting of natural or synthetic alcohol ethoxylate, glycerol oleate, and nonyl phenol ethoxylates.

12. The formulation of claim 11, wherein each of the one or more emulsifiers is independently selected from the group consisting of polyoxyethylene (4 to 12) lauryl ether (C_{12}); polyoxyethylene (10) cetyl ether (C_{16}); polyoxyethylene (10) stearyl ether (C_{18}); polyoxyethylene (10) oleyl ether (C_{18} monounsaturated); polyoxyethylene (2 to 11) C_{12} - C_{15} alcohols; polyoxyethylene (3 to 9) C_{11} - C_{14} alcohols; polyoxyethylene (9) C_{12} C_{14} alcohols; polyoxyethylene (11) C_{16} - C_{18} alcohols; polyoxyethylene (20) C_{12} C_{15} alcohols, glycerol oleate, and polyoxyethylene (2 to 8) nonylphenol.

13. The formulation of claim 1, wherein each of the one or more emulsifiers is independently selected from the group consisting of natural or synthetic alcohol ethoxylate and glycerol oleate.


14. The formulation of claim 13, wherein each of the one or more emulsifiers is independently selected from the group consisting of polyoxyethylene (4 to 12) lauryl ether (C_{12}); polyoxyethylene (10) cetyl ether (C_{16}); polyoxyethylene (10) stearyl ether (C_{18}); polyoxyethylene (10) oleyl ether (C_{18} monounsaturated); polyoxyethylene (2 to 11) C_{12} - C_{15} alcohols; polyoxyethylene (3 to 9) C_{11} - C_{14} alcohols; polyoxyethylene (9) C_{12} C_{14} alcohols; polyoxyethylene (11) C_{16} - C_{18} alcohols; polyoxyethylene (20) C_{12} C_{15} alcohols, and glycerol oleate.

15. The formulation of claim 1, wherein each of the one or more paraffinic oils is a synthetic isoparaffinic oil.

16. The formulation of claim 15, wherein the paraffinic oil is a synthetic isoparaffinic oil having an independently selected carbon number distribution of from C_{16} to C_{35} .

17. A formulation for application to a turfgrass comprising: a synthetic isoparaffinic oil having an independently selected carbon number distribution of from C_{16} to C_{35} ; one or more emulsifiers, each of which is independently selected from the group consisting of natural or synthetic alcohol ethoxylate, glycerol oleate, and nonyl phenol ethoxylates;

a polychlorinated (Cu II) phthalocyanine; and a silicone surfactant, which is independently selected from the group consisting of trisiloxanes and silicone polyethers of Formula I



where $R=H$, CH_3 , or $COCH_3$; $x=1$ to 24; $n=0$ or ≥ 1 ; and $m=1$; and

water;

wherein the silicone surfactant represents from about 0.1 wt % to about 5 wt % of the total weight of the paraffinic oil, one or more emulsifiers, polychlorinated (Cu II) phthalocyanine, and silicone surfactant;

wherein the one or more emulsifiers represent from about 0.5 wt % to about 5 wt % of the total weight of the paraffinic oil, one or more emulsifiers, polychlorinated (Cu II) phthalocyanine, and silicone surfactant;

wherein the polychlorinated (Cu II) phthalocyanines represents from about 1 wt % to about 10 wt % of the total

25

weight of the paraffinic oil, one or more emulsifiers,
polychlorinated (Cu II) phthalocyanine, and silicone
surfactant; and

wherein the formulation is an oil-in-water emulsion and the
one or more emulsifiers and the silicone surfactant are
selected such that the polychlorinated (Cu II) phthalocyanine
is maintained in dispersion in the oil-in-water emulsion for
delivery to the turfgrass.

* * * * *

26

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 8,569,210 B2
APPLICATION NO. : 12/492863
DATED : October 29, 2013
INVENTOR(S) : Michael Fefer

Page 1 of 2

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page

Page 2, Col. 2, line 8, delete “rersistance” and insert -- resistance --

Page 4, Col. 2, line 56, delete “Landscae” and insert -- Landscape --

Page 5, Col. 2, line 35, delete “Fiurprimidol” and insert -- Flurprimidol --

Page 5, Col. 2, line 39, delete “Trinexapac-Ethyt” and insert -- Trinexapac-Ethyl --

Page 5, Col. 2, line 47, delete “Kiltex” and insert -- Killex --

Page 5, Col. 2, line 48, delete “Concentate” and insert -- Concentrate --

Page 5, Col. 2, line 51, delete “PBt/Gordon” and insert -- PBI/Gordon --

Page 6, Col. 1, line 5, delete “Bentrgrass” and insert -- Bentgrass --

Page 6, Col. 1, line 7, delete “Lifestock” and insert -- Livestock --

Page 6, Col. 1, line 40, delete “Paclobutrazole” and insert -- Paclobutrazol --

Page 6, Col. 2, line 25, delete “Phtahalocyanine” and insert -- Phthalocyanine --

Page 6, Col. 2, line 44, delete “Greass” and insert -- Grass --

In the Claims

Claim 1, Col. 22, line 46, delete “phthalocyanines;” and insert -- phthalocyanine; --

Signed and Sealed this
Eighteenth Day of March, 2014



Michelle K. Lee
Deputy Director of the United States Patent and Trademark Office

CERTIFICATE OF CORRECTION (continued)

Page 2 of 2

U.S. Pat. No. 8,569,210 B2

Claim 1, Col. 22, line 58, delete “phthalocyanines” and insert -- phthalocyanine --

Claim 1, Col. 22, line 64, delete “surfactants” and insert -- surfactant --

Claim 1, Col. 22, line 65-66, delete “phthalocyanines” and insert -- phthalocyanine --

Claim 5, Col. 23, line 25, delete “surfactants” and insert -- surfactant --

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 8,569,210 B2
APPLICATION NO. : 12/492863
DATED : October 29, 2013
INVENTOR(S) : Fefer et al.

Page 1 of 1


It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page:

The first or sole Notice should read --

Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b)
by 538 days.

Signed and Sealed this
Fourteenth Day of April, 2015

A handwritten signature in black ink, reading "Michelle K. Lee". The signature is written in a cursive, flowing style.

Michelle K. Lee
Director of the United States Patent and Trademark Office