

PCTWORLD INTELLECTUAL PROPERTY
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER

WO 9606106A1

(51) International Patent Classification⁶ :

C07H 21/02, C12Q 1/68

A1

(11) International Publication Number:

WO 96/06106

(43) International Publication Date:

29 February 1996 (29.02.96)

(21) International Application Number: PCT/US95/10721

(22) International Filing Date: 23 August 1995 (23.08.95)

(30) Priority Data:

08/294,450	23 August 1994 (23.08.94)	US
08/498,402	5 July 1995 (05.07.95)	US

(71) Applicant: UNIVERSITY OF MASSACHUSETTS MEDICAL CENTER [US/US]; 55 Lake Avenue North, Worcester, MA 01655 (US).

(72) Inventors: STERN, Seth; 183 Riverwood Road, Sterling, MA 01564 (US). PUROHIT, Prakash; 330 Sunderland Road #49, Worcester, MA 01604 (US).

(74) Agent: FASSE, J., Peter; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110 (US).


(81) Designated States: AU, CA, CN, CZ, FI, HU, JP, KR, MX, NO, NZ, PL, RO, RU, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published*With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*

(54) Title: HYBRIDIZATION AND FOOTPRINTING METHODS TO CHARACTERIZE THE INTERACTIONS OF 16S rRNA ANALOGUES FOR IDENTIFICATION OF NOVEL ANTIBIOTICS

(57) Abstract

The oligoribonucleotide analogs of the invention, as exemplified in the figure, are relatively small, three-dimensional structures derived from larger parental RNA molecules. The analogs include a first nucleic acid structure including one or more nucleotide sequences that are derived from a region of parental RNA, wherein in its native state, the region binds to a ligand, e.g., an aminoglycoside, with a parental RNA ligand binding pattern, and a second nucleic acid structure including one or more nucleotide sequences combined with the first nucleic acid structure to form the analog and provide the analog with a conformation that binds the ligand with a ligand binding pattern that is substantially identical to the parental RNA ligand binding pattern. These analogs can be used to identify novel therapeutic compounds.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

Hybridization and footprinting methods to characterize the interactions of 16S rRNA analogues for identification of novel antibiotics

Statement as to Federally Sponsored Research

This invention was made with Government support
5 under NIH grant RO1-GM48536. The Government has certain
rights in the invention.

Background of the Invention

The field of the invention is the evaluation of
ribonucleic acid (RNA)-interacting therapeutics, for
10 example, antibiotics that bind to 16S rRNA and inhibit
protein synthesis.

Ribosomes are large, multisubunit
ribonucleoproteins (RNPs) responsible for protein
synthesis, and are highly conserved across phyla, both
15 structurally and functionally. They include large (50S)
and small (30S) subunits assembled from ribosomal RNAs
(rRNAs) and proteins bound to the rRNA. The 30S
ribosomal subunit contains 16S rRNA, while the 50S
subunit contains 23S rRNA. Ribosomes synthesize proteins
20 when correctly bound to messenger RNA (mRNA) and transfer
RNA (tRNA).

The proper assembly of the various components
involved in protein synthesis is thought to be directed
by binding sites on the rRNAs. Two binding sites of
25 importance on the rRNA for protein synthesis are the so-
called A and P sites, which accommodate the incoming
aminoacyl-tRNA (A site) and the peptidyl-tRNA (P site),
respectively. In prokaryotes, these sites are composed
in part of highly ordered structures of the 16S rRNA,
30 probably in the cleft of the 30S subunit.

Ribosomes are structurally similar in all species
(including eukaryotes), although the primary nucleotide
sequences of rRNA molecules differ. For a review of
ribosomal RNA function, see Noller *In The RNA World*,

- 2 -

Gesteland and Atkins (eds.), 137-84 (CSHL Press, NY, 1993); Noller et al., *In The Ribosome: Structure, Function, and Evolution*, Hill, et al. (eds.), 73-92 (American Soc. for Microbiol., Washington, DC, 1990).

5 It is now generally accepted that 16S and 23S rRNAs (found in the 50S ribosomal subunit; analogous eukaryotic rRNAs are 28S, 5.8S, and 5S rRNA in the 60S ribosomal subunit, and 18S rRNA in the 40S ribosomal subunit) play important, if not critical, roles in the
10 decoding and peptidyl transferase activities of ribosomes (Noller et al., 1990, *supra*, Noller, *supra*).

 Most antibiotics that inhibit protein synthesis act directly on ribosomes. For example, some mutations in ribosomal proteins can result in antibiotic resistance
15 (Birge and Kurland (1969) *Science*, 166:1282; Ozaki et al. (1969) *Nature*, 222:333; Davies et al. (1965) *Science*, 149:1096). Work in this field has also demonstrated that aminoglycoside antibiotics interact with sites on ribosomal subunits, resulting in protection of RNA as
20 visualized by RNA footprint assays (e.g., Moazed and Noller (1987) *Nature*, 327:389; Woodcock et al. (1991) *EMBO J.*, 10:3099; Thompson and Cundliffe (1991) *Biochimie*, 73:1131-1135).

 Such research has shown that specific nucleotides
25 in 16S rRNA are the binding targets of aminoglycosides such as neomycin, streptomycin, hygromycin, gentamycin, and tetracycline. Similarly, specific nucleotides in 23S rRNA are targeted by numerous MLS compounds (macrolides, lincomycins, and streptogramins), including erythromycin.

30 Some antibiotics (e.g., edeine, pactamycin, apramycin, and neamine) inhibit protein synthesis by interfering with binding between tRNA and the A- or P-sites on the ribosome during translation (Woodcock et al. (1991) *EMBO J.* 10:3099). Interactions of many of these
35 compounds with 16S rRNA in the 30S ribosomal subunit have

- 3 -

been mapped to various functional sites, primarily by chemical footprinting assays. Other antibiotic compounds, such as the peptide antibiotic, thiostrepton, have been shown to similarly interact with 23S rRNA in 5 50S subunits (Thompson and Cundliffe (1991) *supra*).

Several factors, all related to the structural complexity of the ribosome, complicate screening assays that rely on binding of a potential drug candidate to a ribosomal target. First, obtaining large quantities of 10 purified ribosomes, even from common bacteria, is difficult. Second, ribosomes often degrade under typical screening conditions. Third, it is unclear to what degree a compound's ability to bind to ribosomes or RNA molecules is indicative of its antibiotic potential.

15 In particular, even small, well-defined RNA molecules are complex targets for such drug screening assays, because they support multiple binding sites for small molecules. For example, the polyanionic phosphodiester backbone of a nucleic acid presents an 20 attractive binding target for many compounds carrying a positive charge, as does the hydrophobic helical core for many aromatic compounds. Thus, compounds that bind to nucleic acids moieties not directly related to the targeted function are less likely to be useful drugs.

25 Summary of the Invention

The present invention is based on the discovery that relatively small RNA molecules, "oligoribonucleotide analogs," that are derived from portions of larger parental RNA structures (e.g., RNPs), surprisingly retain 30 important parental structure even without the rest of the parental RNA sequence(s) and associated proteins. Oligoribonucleotide analogs are useful in new methods of screening candidate compounds for antibiotic activity.

- 4 -

Accordingly, in one aspect, the invention features an oligoribonucleotide analog of a region of a parental ribonucleic acid (RNA) including (i) a first nucleic acid structure having one or more nucleotide sequences, the first structure being derived from the region of parental RNA, e.g., 16S ribosomal RNA (rRNA), wherein in its native state, the region binds to a ligand with a parental RNA ligand binding pattern, and (ii) a second nucleic acid structure having one or more nucleotide sequences combined with the first nucleic acid structure to form the analog and provide the analog with a conformation that binds the ligand with a ligand binding pattern that is substantially identical to the parental RNA ligand binding pattern.

As used herein, the terms "oligoribonucleotide analog" or "analog" mean a single-stranded molecule composed of ribonucleic acids that is smaller than the parental RNA (e.g., having a total of approximately 10 to several hundred nucleotides) that folds into a discreet three-dimensional conformation that mimics the structure of a subdomain of a larger parental RNA molecule. However, modified RNAs, e.g., phosphorothioates, deoxynucleotides, or 2'-O-methyl substitutions, can also be used to confer stability to the analogs.

The oligoribonucleotide analogs are produced *in vitro* either by transcription of DNA templates with RNA polymerase, e.g., T7 RNA polymerase, or by chemical synthesis with a commercially available DNA/RNA synthesizer. Generally, there are from approximately 10 nucleotides up to several hundred nucleotides (and preferably 20 to 50 nucleotides) in an oligoribonucleotide analog that are identical to nucleotides in a region of parental RNA (the number of parentally derived nucleotides is determined by the sequence necessary to mimic a functional subdomain).

- 5 -

The second nucleic acid structure can contain heterologous (artificial) nucleotide sequences, i.e., sequences that do not exist in the parental RNA, that, when combined with the first nucleic acid structure, stabilize the structure of the analog. As used herein, the term "combined" means that the nucleotide sequences of the first and second nucleic acid structures are linked, e.g., by covalent and non-covalent bonds (e.g., hydrogen bonds, ionic (electrostatic) interactions, and/or van der Waals forces), in such a way that the complete analog has a ligand binding pattern that is substantially identical to the ligand binding pattern of the parental RNA in the native state, e.g., in the intact ribosome.

As used herein, the term "parental RNA" molecule or structure refers to the naturally occurring (or "native"), intact RNA molecule or structure (including associated proteins and other components of the structure) from which the first nucleic acid structure of the oligoribonucleotide is derived. The parental RNA can be ribosomal RNA, viral RNA such as HIV RNA, messenger RNA, or specific cellular RNA regulatory elements. Preferred HIV RNA oligoribonucleotide analogs are derived from the Tat binding site (TAR) or the Rev Response Element (RRE), more preferably from a portion of the RRE having the sequence GCACUAUGGGCGCAGCGUCAAUGACGCU GACGGUACAGGCCAGACAAUUAUUGUCUGGUAUAGUGC (SEQ ID NO:8). Further, if the parental RNA-binding "ligand" is an aminoglycoside, such as neomycin, the rRNA binding pattern can also be referred to as the "aminoglycoside protection profile" of the parental RNA.

In specific embodiments, the region of parental RNA is a decoding region of 16S rRNA, e.g., including nucleotides 1398-1410 and 1490-1505 of 16S rRNA (of *Escherichia coli*). In other embodiments, the region is

- 6 -

the A site subdomain of the decoding region of 16S rRNA, e.g., including nucleotides 1404-1410 and 1490-1497 of 16S rRNA (of *E. coli*). The "decoding region" is the portion of the 16S rRNA which, in the intact ribosome, accurately aligns tRNA with mRNA for correct codon-anticodon base pairing during protein synthesis. The decoding region consists of the "A site," which in the intact ribosome accommodates the incoming aminoacyl-tRNA, and the "P site," which contains the peptidyl-tRNA complex (the tRNA still linked to all the amino acids added to the chain thus far). Other possible rRNAs which can be used in the invention are 23S rRNAs of prokaryotes; or 28S, 5.8S, 5S and 18S rRNAs of eukaryotes.

The second nucleic acid structure of the analog can include a stable stem loop such as tetraloop, e.g., having the nucleotide sequence 5'-CCUUCGGG-3', in which the nucleotides UUCG form the loop, and the nucleotides CC and GG are paired. The second nucleic acid structure can also include two nucleotide sequences forming a based-paired stable helix, also known as a "nucleotide clamp." For example, such a clamp can have the nucleotide sequence:

3'-CGUGUC-5' or 3'-CC-5'
5'-GCACAG-3' 5'-GG-3'.

Specific analogs of the invention include a first nucleic acid structure derived from a decoding region of 16S rRNA, and a second nucleic acid structure including a tetraloop and a base-paired nucleotide clamp. For example, the decoding region can include nucleotides 1398-1410 and 1490-1505 of 16S rRNA, the tetraloop can have nucleotide sequence 5'-CCUUCGGG-3', the base-paired nucleotide clamp can have the nucleotide sequence:

3'-CGUGUC-5'

- 7 -

5'-GCACAG-3',

and the complete linear nucleotide sequence of the combined first and second nucleotide structures of the analog would be 5'-

5 GCACAGACCGCCCGUCACACCUUCGGGUGAAGUCGUAACAAGGCUGUGC-3' (SEQ ID NO:1).

In another embodiment, the analog can be derived from a decoding region including nucleotides 1404-1410 and 1490-1497 of 16S rRNA, the tetraloop can have

10 nucleotide sequence 5'-CCUUCGGG-3', the base-paired nucleotide clamp can have the nucleotide sequence:

3'-CC-5'
5'-GG-3',

and the complete linear nucleotide sequence of the

15 combined first and second nucleotide structures of the analog would be 5'-GGCGUCACACCUUCGGGUGAAGUCGCC-3' (SEQ ID NO:11).

In another aspect, the invention features an affinity assay for determining the potential antibiotic
20 or therapeutic activity of a test compound, the assay including the steps of (i) mixing a test compound with an oligoribonucleotide analog of the invention under conditions that allow formation of a binding complex between the analog and the test compound, and (ii)
25 detecting the formation of a binding complex, wherein the presence of a binding complex indicates that the test compound has potential antibiotic activity.

It is preferred that the test compound show specific affinity, i.e., that its particular interaction
30 with the oligoribonucleotide analog is highly repeatable and affects the same nucleotide(s) in the analog. The higher the affinity of the test molecule for the oligoribonucleotide, the higher its potential usefulness as an therapeutic/ antibiotic compound.

35 In specific embodiments, the analog is labelled, e.g., with a fluorescent or radioactive label, and

- 8 -

immobilized on a surface, and the binding complex is detected by monitoring changes in the signal of the label when a test compound is bound to the analog, or the analog is immobilized on a surface, the test compound is labelled, and the binding complex is detected by
5 detecting any of the label bound to the surface via the analog.

In a preferred embodiment, the affinity assay is used to identify potentially useful compounds from a
10 mixture of compounds. It includes the steps of contacting the analog with numerous test compounds existing in a mixture, isolating the analog-test compound complexes, and determining the identity of the test compounds that bind with the analog. One example of such
15 a mixture of compounds is an encoded library of small molecules (see, e.g., Needleman et al. (1993) *PNAS* 90:10700-04).

In an encoded library, the molecular structure of synthetic small molecules, e.g., peptides or other
20 organic molecules, are encoded, e.g., by a DNA strand. For example, in an encoded peptide library, the specific DNA sequences for each peptide are attached to beads (with a unique sequence encoding the peptide sequence attached to each bead), as are the small peptide
25 molecules encoded by the DNA sequences. If one or more of these test molecules shows an affinity for the labelled oligoribonucleotide analog of the invention, it can be isolated, e.g., by biotinylation/ streptavidin interaction or fluorescently activated cell sorting of
30 labelled molecules, and identified by standard means, e.g., polymerase chain reaction (PCR) or dideoxy sequencing.

The invention also features a competitive binding assay for determining the potential antibiotic or
35 therapeutic activity of a test compound, the assay

- 9 -

including the steps of (i) mixing an analog with an analog-binding ligand under conditions that allow formation of a first binding complex between the analog and the ligand, (ii) mixing a test compound with the
5 first binding complex under conditions that allow the test compound to disrupt the first binding complex to form a second binding complex between the analog and the test compound, and (iii) detecting the disruption of the first binding complex, wherein the disruption of the
10 first binding complex indicates that the test compound has potential antibiotic activity.

In specific embodiments, the ligand is labelled, e.g., fluorescently or radioactively, the analog is immobilized on a surface, and the disruption of the first
15 binding complex is detected by monitoring any decrease in the signal of the label when a test compound displaces the ligand from the first binding complex.

Alternatively, the analog can be labelled, the ligand immobilized on a surface, and the disruption of the first
20 binding complex detected by monitoring any decrease in the signal of the label when a test compound displaces the analog from the first binding complex.

In a preferred embodiment, the ligand is an aminoglycoside, and is immobilized, e.g., covalently
25 cross-linked, to a solid support, e.g., a nylon or cellulose-derived membrane, microtiter plate, or other plastic support, and then incubated with a labelled (fluorescently tagged or radiolabeled) oligoribonucleotide analog. The aminoglycoside-analog
30 complexes are then challenged competitively with test compounds, or mixtures of compounds, cell extracts, etc., and those compounds that can effectively bind to the analog will displace the analog from the aminoglycoside-analog complex. Other embodiments include cross-linking

- 10 -

the analog to a solid support and challenging with a mixture of the aminoglycoside and test compounds.

In another aspect, the invention features an *in situ* footprinting assay for determining the potential antibiotic or therapeutic activity of a test compound, the assay including the steps of (i) mixing an oligoribonucleotide analog with a test compound under conditions that allow formation of a binding complex between the analog and the test compound, (ii) incubating the binding complex with a chemical probing reagent and monitoring for an effect of the reagent on the analog in the complex, (iii) in a separate control reaction, incubating the analog unbound to any test compound with the chemical probing reagent and monitoring for an effect of the reagent on the unbound analog, and (iv) comparing any effects of the probing reagent on the analog in the binding complex and on the unbound analog, wherein prevention of an effect of the reagent on the analog in the binding complex caused by the reagent on the unbound analog indicates that the test compound has potential antibiotic activity.

In specific embodiments, the chemical probing reagent is dimethyl sulfate (DMS), kethoxal (KE), or carbodiimide, e.g., 1-cyclohexyl-3-(2-morpholinoethyl) carbodiimide metho-*p*-toluene sulfonate (CMCT). These and other probing reagents covalently modify, e.g., methylate in the case of DMA, a nucleotide in the analog when unbound to any test compound. The effect of the probing reagent can be monitored by use of a labelled oligonucleotide, e.g., having the sequence CAGUGU, that is complementary to a portion of the analog, and thus hybridizes to the analog when the analog is protected by the test compound from modification, and does not hybridize to the analog when modified, e.g., methylated by the reagent, the presence of the label after

- 11 -

completion of the assay indicating that the test compound has potential antibiotic activity. Alternatively, the effect of the probing reagent is monitored by use of an oligonucleotide primer, e.g., having the sequence

5 TTCACCCGGAAGGTG (SEQ ID NO:12), that is complementary to a portion of the analog, a labelled nucleotide, and reverse transcriptase, wherein extension of the primer on the analog with the labelled nucleotide does not occur when the analog is methylated by the reagent, the
10 presence of the label after completion of the assay indicating that the test compound has potential antibiotic or therapeutic activity.

Other potential therapeutics or antibiotics that can be identified include human immunodeficiency virus
15 (HIV) therapeutics (e.g., molecules that inhibit viral replication or protein synthesis). Regions of interest include those involved in viral replication (e.g., Tat binding site (TAR) and the Rev Response Element (RRE) of RNA).

20 As used herein, "protection," e.g., by an aminoglycoside or thiostrepton, refers to the characteristic footprint or "profile," i.e., a pattern of bands on a polyacrylamide gel which results following various chemical treatments (e.g., dimethyl sulfate,
25 kethoxal, 1-cyclohexyl-3-(3-morpholinoethyl)-carbodiimide metho-p-toluene sulfonate) of RNA previously exposed to the protective molecule. These footprint assays are routine, using methods well known in the art (see, for example Moazed et al., 1986, *Cell*, 47:985-94; Stern et
30 al., (1988) *Meth. Enzymol.* 164:481-489; Peattie and Gilbert (1980) *PNAS* 77:4679-82).

A major advantage of the methods described herein is that these methods allow the discovery of new antibiotic compounds or molecules without the labor and
35 expense of standard antimicrobial activity assays. By

- 12 -

using the oligoribonucleotide analogs that mimic the aminoglycoside interacting site (ligand binding site) of a parent RNA, compounds which are likely to have antibiotic, protein synthesis-inhibiting, or viral-inhibiting properties can be quickly and inexpensively identified, even if very large-scale screening protocols are used with hundreds of test molecules. The assays are amenable to automation, which increases ease of large-scale screening methods.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the detailed description, and from the claims.

Description of the Drawings

Figs. 1A-1E are schematic depictions of 16S rRNA, the decoding region, two oligoribonucleotide analog, and probing data. Fig. 1A is a schematic diagram of 16S rRNA showing nucleotides outside the decoding region (boxed) implicated in tRNA-30S subunit interactions at the A- and P-sites (Noller et al., *In The Ribosome: Structure, Function, and Evolution* (Eds. Hill, Dahlberg, et al.) American Soc. for Microbiol., Washington, DC, 1990, pp.

- 13 -

73-92; Moazed et al., 1986, *Cell*, 47:985-94, Moazed et al., 1990, *J. Mol. Biol.*, 211:135-45). The position of a zero length cross-link between the tRNA anticodon loop and C1400 (cytosine at location 1400) in the decoding region (AC XL) is indicated (Prince et al., 1982, *Proc. Natl. Acad. Sci. USA*, 79:5450-54). Fig. 1B shows the decoding region (SEQ ID NOS:9 and 10) and the nucleotides protected from DMS modification (at N1 of A, N3 of C, and N7 of G) by neomycin-like aminoglycoside antibiotics (Moazed et al., 1987, *Nature*, 327:389-94, Woodcock et al., 1991, *EMBO J.*, 10:3099-103) (Neos) and A- and P-site tRNA (Moazed et al., 1986, *Cell*, *supra*, Moazed et al., 1990, *J. Mol. Biol.*, *supra*). The positions of the A- and P-site decoding region subdomains, and the identities of post-transcriptionally methylated nucleotides are indicated. The decoding region is drawn with the tertiary base pairs of Gutell and coworkers (Gutell et al., 1985, *Prog. Nucl. Acids Res. Mol. Biol.*, 32:153-216, Gutell, *In The Translational Apparatus*, (Eds. Nierhaus, Subramanian et al.) Plenum Publishing, NY, 1993, pp. 477-88): C1399-G1504, G1401-C1501, C1402-A1500, C1404-G1497, and G1405-C1496. Fig. 1C shows the decoding region oligoribonucleotide analog (SEQ ID NO:1) with nucleotides protected from DMS modification by neomycin and paromomycin (Neos) indicated. The analog contains 16S rRNA decoding region sequences from A1398 to A1410 on the proximal side and U1490 to G1505 on the distal side. The distal end of the analog terminates in the same stem-tetraloop found at the end of the penultimate stem of 16S rRNA (nucleotides 1448-1455). A heterologous clamping helix is shown at the top. Weak and strong protection are indicated by small and large symbols, respectively. Fig. 1D shows the decoding region oligoribonucleotide analog (SEQ ID NO:1) with nucleotides protected from DMS modification, or enhanced in reactivity toward DMS, in

- 14 -

response to poly U mRNA and tRNA anticodon stem-loop T7 transcripts. Fig. 1E shows an oligoribonucleotide analog (SEQ ID NO:11) of the A-site of the decoding region, with nucleotides protected from DMS modification by neomycins
5 indicated.

Figs. 2A-2B are photos of electrophoresis gels showing decoding region analog-antibiotic interactions. Fig. 2A shows results from DMS probing reactions with the oligoribonucleotide analog alone (lane 6) and in the
10 presence of 0.1, 1, 10, and 100 μ M neomycin (lanes 7-10), paromomycin (lanes 11-14), hygromycin (lanes 15-18), streptomycin (lanes 19-22), tetracycline (lanes 23-26), and erythromycin (lanes 27-30). Dideoxy sequencing reactions (lanes 1-4) and a control extension reaction
15 with unmodified RNA (K, lane 5) are to the left, and bands corresponding to nucleotides discussed below are indicated to the right. Fig. 2B shows DMS/N7 probing reactions with the oligoribonucleotide analog alone (lane 1) and in the presence of 0.1, 1, 10, and 100 μ M neomycin
20 (lanes 2-5), paromomycin (lanes 6-9).

Fig. 3 is an electrophoresis gel showing interactions of tRNA anticodon stem-loop transcripts and poly U with the decoding region oligoribonucleotide analog. DMS and DMS/N7 probing reactions with naked
25 analog (lanes 6 and 16), and 3, 6, 12, and 24 μ M anticodon stem-loop transcript (lanes 7-10 and 12-15). The same reactions were repeated in the presence of 4 μ g poly U (lanes 11-15 and 21-25).

Figs. 4A-4B are electrophoresis gels showing the
30 selectivity of decoding region analog interactions. In Fig. 4A, selectivity of mRNA interactions is shown by DMS probing reactions. Reactions with naked analog (lane 6), 3 μ M (lane 7), and 6 μ M (lane 8) anticodon stem-loop transcript were repeated in the presence of 2 μ g poly U
35 (lanes 9-11), polydeoxy U (lanes 12-14), and poly C

- 15 -

(lanes 15-17). Nucleotides protected by tRNA anticodon stem-loop or mRNA are indicated to the right. Fig. 4B shows selectivity of the P-site subdomain for stem-loop structures. DMS probing reactions with naked analog (lane 6), and 1.5, 3, 6, and 12 μ M *E. coli* tRNA^{Phe} anticodon stem-loop transcript (lanes 7-10), tRNA^{Pro} anticodon stem-loop transcript (lanes 11-14), scrambled tRNA^{Phe} (sctRNA^{Phe}, lanes 15-18), tetraloop element (lanes 19-22), and triUloop element (lanes 23-26). Nucleotides protected by the anticodon stem-loops are indicated to the right.

Figs. 5A and 5B are schematic diagrams of two different screening methods using the oligoribonucleotide analogs of the invention.

Fig. 6 is a schematic diagram of an analog of the A-site of the decoding region, and shows the potential nucleotide locations (boxes) for derivatization, e.g., to add a fluorescent label.

Figs. 7A and 7B are schematic diagrams of a high-throughput footprinting assay using the analogs of the invention and an oligonucleotide-based reporter system.

Figs. 8A and 8B are schematic diagrams of a high-throughput footprinting assay using the analogs of the invention and a reverse-transcriptase-based reporter system.

Detailed Description

Despite the complex structures and numerous associated proteins of complete ribosomes, we have discovered small oligoribonucleotide analogs that mimic small domains of parental RNAs and that can fold and function autonomously for purposes of the screening assays described herein. The invention includes all such autonomously functioning oligoribonucleotide analogs that can be used to identify novel therapeutic agents for

- 16 -

antibiotic, antiviral, anti-cancer, anti-proliferative, and anti-inflammatory use, particularly those specifically described below. Other oligoribonucleotide analogs useful as screening probes can be identified in
5 cellular RNA, including mRNA, and viral RNA.

One domain for use to derive the first nucleic acid structure of the analogs is the decoding region of 16S rRNA, which is located near the 3' end of 16S rRNA of *E. coli* (Figs. 1A and 1B). The experiments described
10 below show that an oligoribonucleotide analog of the decoding region interacts with both antibiotic and RNA ligands of the 30S subunit in a manner that correlates with normal subunit function. The activities of the decoding region analog suggest that the intimidating
15 structural complexity of the ribosome can be, to some degree, circumvented.

Experimental Methods

Preparing an analog

A first oligoribonucleotide analog RNA, derived
20 from the decoding region of 16S rRNA, was made by transcribing, with T7 RNA polymerase (Milligan et al., 1989, *Meth. Enzymol.*, 180:51-62), a linearized pGEM3 plasmid (Promega) as described in Zapp et al., 1993, *Cell*, 74:969-978, containing the analog sequence shown in
25 Fig. 1D (SEQ ID NO:1) flanked by EcoRI and BamHI restriction sites, and a 15 mer reverse primer annealing site immediately 3' of the BamHI site. This sequence includes both the first and second nucleic acid structures to form the complete analog.

30 Specifically, in the first analog (shown in Fig. 1D), the first nucleic acid structure includes the nucleotide sequences 5'-ACCGCCGUCACA-3' (SEQ ID NO:12) and 5'-UGAAGUCGUAACAAGG-3' (SEQ ID NO:13) derived from the complete decoding region. The second nucleic acid

- 17 -

structure includes a tetraloop 5'-CCUUCGGG-3', which is the minimum size of a tetraloop found to be sufficiently stable, and a nucleotide clamp made of the sequences 5'-GCACAG-3' and 3'-CGUGUC-5'. Any other stable helix can be
5 used in place of this particular nucleotide clamp. Likewise, any other stable stem loop can be used in place of this particular tetraloop.

The second analog, shown in Fig. 1E, was made in the same way, but is derived from the A-site subdomain of
10 the decoding region, and has the nucleotides sequences 5'-CGUCACA-3' and 5'-UGAAGGUCC-3'. In terms of the second nucleic acid structure, the second analog has the same tetraloop as in the first analog, but includes a shorter nucleotide clamp made of two nucleotide sequences
15 5'-GG-3' and 3'-CC-5'. The complete nucleotide sequence of the second analog is 5'-GGCGUCACACCUUCGGGUGAAGUCGCC-3' (SEQ ID NO:11).

Gel purified RNA was concentrated by ethanol precipitation, heated to 80°C for 1 minute, and
20 immediately placed at 37°C for 5 minutes.

Aminoglycoside protection experiments

Interaction (or "binding") reactions (12.5 µl) containing 125 ng oligoribonucleotide analog RNA and antibiotics in 80 mM K-Hepes (pH 7.9), 50mM NH₄Cl, and 5%
25 PEG buffer were annealed at 37°C for 15 minutes and incubated on ice for 1 hour. DMS (1 µl of 1:5 in ethanol) was added, and modification reactions were incubated for 40 minutes on ice. Reactions were stopped with DMS Stop (Peattie et al., 1980, *Proc. Natl. Acad. Sci. USA*, 77:4679-82) and the RNA was purified by ethanol
30 precipitation.

DMS/N7 reactions were performed according to standard protocols (Peattie et al., *supra*) except that lyophilization steps were replaced by acid-phenol

- 18 -

extraction and ethanol precipitation. For primer extension, 10 ng analog RNA was annealed to 0.75 ng end-labeled primer and extended with 10-15 U MoMuLV reverse transcriptase for 1 hour. Reactions were stopped by ethanol precipitation, pellets resuspended in 10 μ l 8M urea, 0.05X TBE loading buffer, and 2 μ l was loaded onto 8%, 19:1 acrylamide:bisacrylamide, 0.5X TBE sequencing gels.

trNA protection experiments

10 Binding reactions were performed as described above for aminoglycoside protection experiments, except that binding buffer contained 200 mM NH_4Cl and 80 mM MgCl_2 . *E. coli* tRNA^{phe} anticodon stem-loop (GGGGAUUGAAAAUCCCC; SEQ ID NO:3) was transcribed with T7 RNA polymerase, gel purified, concentrated by ethanol precipitation, and annealed with a brief heat step (80°C/1 minute followed by 10 minutes on ice).

Selectivity experiments

Reactions were as described for tRNA protection experiments. tRNA^{Pro} anticodon stem-loop (GGUCAUCUUGGGG UGAUGACC; SEQ ID NO:4), scrambled tRNA^{phe} (GGGAGCGUCAU CACAUA; SEQ ID NO:5), tetraloop element (GGGACUUCGGUCCC; SEQ ID NO:6), and triUloop element (GGCGCUUUGCGCC; SEQ ID NO:7) were transcribed and treated as described for the *E. coli* tRNA^{phe} anticodon stem-loop. The assay and results are described below and in Fig. 4B.

Experimental Design and Results

RNA structure analysis with chemical probes is a well established and powerful technique that allows individual atoms of nucleotide bases, or of the phosphodiester backbone, to be monitored for inter- or intra-molecular interactions. For example, dimethyl sulfate (DMS) methylates N1 of A, N7 of G, and N3 of C residues, kethoxal (KE) forms an adduct with N1 and N2 of

- 19 -

G, and a soluble carbodiimide (1-cyclohexyl-3-(2-morpholinoethyl) carbodiimide metho-p-toluene sulfonate, CMCT) modifies N3 of U residues. Nucleotide base modifications that interfere with formation of

5 Watson-Crick base pairs are typically monitored via reverse transcription, as the progress of the reverse transcriptase enzyme is strongly impeded when it encounters such modifications. This leads to production of a truncated reverse transcription product whose length
10 effectively maps the position of the modified base in the RNA chain when the products are run on standard DNA sequencing gels.

Other modifications that do not themselves directly interfere with formation of Watson-Crick base
15 pairs can be similarly monitored, by first inducing RNA strand scission. For example, methylation of N7 of G by DMS can be monitored by treating the methylated RNA with sodium borohydride and aniline to induce strand scission at the positions of N7 methylation. Reverse
20 transcriptase simply falls off the template at these points, again leading to a truncated product that maps the position of N7 methylation.

We used these principles and techniques to analyze the oligoribonucleotide analogs of the invention and to
25 develop screening assays using these analogs.

We first designed an oligoribonucleotide analog of the decoding region shown in Figs. 1C and 1D (SEQ ID NO:1) as described above, and asked if it could interact with aminoglycoside antibiotics such as neomycin, which
30 have previously been shown to protect N1 of A1408 (this notation indicates the adenine at location 1408) and N7 of G1491 and G1494 within the A-site subdomain of the decoding region (Moazed et al., 1987, *Nature*, *supra*; Woodcock et al., *supra*); similar interactions with the
35 analog would be a strong indicator of its proper folding

- 20 -

and functional potential. Conversely, other antibiotics, such as streptomycin, tetracycline, and erythromycin interact with other segments of 16S rRNA or 23S rRNA (Moazed et al., 1987, *Nature*, *supra*; Woodcock et al., 5 *supra*, Moazed et al., 1987, *Biochimie*, 69:879-84) and would not, therefore, be expected to interact with the analog. A second analog, derived from the A-site subdomain of the decoding region (and shown in Fig. 1E), was tested in the same way.

10 As summarized in Figs. 1C, 1D, and 1E, the interactions of aminoglycosides such as neomycin and paromomycin with decoding region oligoribonucleotide analogs bear a striking similarity to their counterpart interactions with the decoding region in subunits of 15 complete 16S rRNA in ribosomes. In particular, the decoding region of ribosomes (Fig. 1A) and a small analog containing just the A-site subdomain (Fig. 1E) exhibit virtually identical aminoglycoside interactions, which are abolished by a single G to U transversion at position 20 1491.

The analogs were probed with dimethyl sulfate (DMS; Stern et al., 1988, *Meth. Enzymol.*, 164:481-89) alone as naked RNA, or in the presence of increasing concentrations of neomycin, paromomycin, hygromycin, 25 streptomycin, tetracycline, and erythromycin. Fig. 2A shows that neomycin (lanes 7-10) and the closely related aminoglycoside paromomycin (lanes 11-14) strongly protected N1 of A1408 at concentrations of 10 and 1 μ M, respectively. In addition, hygromycin (lanes 15-18), a 30 structurally dissimilar aminoglycoside, weakly enhanced the reactivity of N1 of A1408. In contrast, streptomycin (lanes 19-22), tetracycline (lanes 23-26), and erythromycin (lanes 27-30), antibiotics that do not interact with the decoding region in 16S rRNA, did not 35 detectably interact with the analogs.

- 21 -

Fig. 2B shows that, in addition to N1 of A1408, N7 of G1405 and G1494 were strongly protected by neomycin and paromomycin, while N7 of G1491 and G1497 were weakly protected. These data are summarized in Fig. 1C.

5 Because these results closely match those previously obtained with antibiotics and intact, whole ribosomes (Moazed et al., 1987, *Nature*, supra, Woodcock et al., supra, Moazed et al., 1987, *Biochimie*, supra), we concluded that the A-site subdomain of our
10 oligoribonucleotide analog folds into a biologically relevant conformation.

We asked next if the analog could function similarly to the decoding region of 16S rRNA in the ribosome by interacting with tRNA and/or mRNA. In the
15 experiment shown in Fig. 3, increasing concentrations of a minimal tRNA for these purposes (Rose et al., 1983, *J. Mol. Biol.*, 167:103-17), a T7 transcript of the anticodon stem-loop of *E. coli* tRNA^{Phe}, were incubated either alone (lanes 7-10 and 17-20) or in the presence of cognate poly
20 U message (lanes 11-15 and 21-25). Fig. 3 shows that the anticodon stem-loop transcript protected N3 of C1402 and C1403 (compare lane 6 with lanes 7-10) and N7 of G1401 (compare lane 16 with lanes 17-20), while poly U protected N1 of A1408, A1492, A1493, A1499, and A1502
25 strongly, N1 of A1500, A1503 weakly, and N7 of G1494 weakly (compare lane 6 with lanes 11-15 and lane 16 with lanes 21-25). In addition, the reactivities of N1 of A1410 and A1493, and N3 of C1407 were enhanced by the stem-loop (compare lane 6 with lanes 7-10), effects which
30 were attenuated in the presence of poly U (compare lanes 7-10 with lanes 12-15).

Comparison of the present results with data from previous studies with ribosomes (Moazed et al., 1986, *Cell*, supra; Moazed et al., 1990, *J. Mol. Biol.*, supra)
35 shows that the nucleotides protected by the tRNA

- 22 -

anticodon stem-loop transcript in the analog (G1401, C1402, and C1403) are closely associated with, and in fact overlap, those protected by P-site bound tRNA in the presence or absence of message (C1399, C1400, and G1401).
5 Furthermore, within the A-site subdomain of the analog, nucleotides protected by mRNA (A1408, A1492-G1494) are identical to those associated with the mRNA-dependent interaction of tRNA with the A-site of ribosomes (Moazed et al., 1986, *Cell*, *supra*; Moazed et al., 1990, *J. Mol.*
10 *Biol.*, *supra*). The other nucleotides, protected by mRNA (A1499, A1500, A1502, and A1503), have not been previously identified with either A- or P-site function in chemical probing experiments.

To help determine whether poly U mRNA interactions
15 were mediated by Watson-Crick base pairs with A residues in the analog, we next tested poly C, an mRNA with no potential to form Watson-Crick base pairs with A residues, and with polydeoxy U, a 2'-deoxy-form of poly U, with similar Watson-Crick base pairing potential. The
20 experiment of Fig. 4A showed that while poly U protected A1408, A1492, and A1493 strongly as expected (compare lanes 6-8 with lanes 9-11), polydeoxy U failed to detectably interact with any nucleotides in the analog (compare lanes 6-8 with lanes 12-14). The interactions
25 of poly C were more complex, as the reactivities of N1 of A1410 and N3 of C1402 and C1403 were enhanced (compare lanes 6-8 with lanes 15-17). However, with the exception of A1492, poly C protected the same nucleotides as poly U, including A1408, A1493, A1499, A1500, A1502, and
30 A1503. Thus, despite the apparent disruption of analog structure induced by poly C, its protection pattern was virtually identical to that of poly U. Considered together, these results suggest that the interactions of mRNA with the decoding region are not mediated simply by
35 Watson-Crick base pairing.

- 23 -

C1402 and C1403 are unreactive in 16S rRNA (Moazed et al., 1986, *J. Mol. Biol.*, 187:399-416); their reactivity in the analog and their protection by the tRNA anticodon stem-loop transcript suggest that the structure of the P-site subdomain of the oligoribonucleotide analog differs from that of ribosomes.

To understand better the discriminatory capacity, or selectivity, of the P-site subdomain of the oligoribonucleotide analog, we asked next whether it differentiates between closely related RNA structures. In the experiment shown Fig. 4B, a second (sequence-divergent), proline-specific tRNA anticodon stem-loop (tRNA^{Pro}) (GGUCAUCUUGGGGUGAUGACC; SEQ ID NO:4), a scrambled-sequence *E. coli* tRNA^{Phe} with little potential for secondary structure formation (sctrNA^{Phe}) (GGGAGCGUCAUCAUA; SEQ ID NO:5), a highly stable tetraloop stem-loop element (tetraloop) (GGGACUUCGGUCCC; SEQ ID NO:6), and another, smaller stem-loop element with a loop composed of only three U residues (triUloop) (GGCGCUUUGCGCC; SEQ ID NO: 7) were each titrated into a binding reaction with the analog.

Fig. 4B shows that only the tRNA^{Phe} and tRNA^{Pro} anticodon stem-loop transcripts interacted, each strongly protecting C1402 and C1403. The failure of the scrambled tRNA^{Phe} and other stem-loop elements to interact detectably supports the hypothesis that, despite its altered conformation, the P-site subdomain interacts selectively with RNA ligands. These results suggest, therefore, that the P-site subdomain of the oligoribonucleotide analog exhibits significant discriminatory capacity that may be related to ribosomal P-site function.

In light of the omission of 1500 nucleotides of the complete 16S rRNA, all nucleotide modifications, and

- 24 -

the 21 small subunit proteins normally found in the 30S ribosomal subunit, the demonstration of interactions between the small decoding region oligoribonucleotide analog and 30S subunit ligands that correlate with normal 30S subunit activities was surprising and unexpected. Positive correlations are probably best exemplified by the interactions of the antibiotics with the A-site subdomain where a nearly perfect correspondence between the characteristics of the decoding region in ribosomes and the oligoribonucleotide analog was observed with a wide variety of compounds.

Uses/Advantages of the Invention

The oligoribonucleotides of the invention are diverse, in that they can be derived from any ribosomal RNA (e.g., 16S, 23S), human immunodeficiency virus (HIV) RNA (see Green (1993) *AIDS Res. Rev.* 3:41-55 for examples of desirable regions such as Tat or Rev interaction sites), or any other RNA which can be manipulated to retain a functionally relevant structure. Although the decoding region of 16S rRNA and aminoglycoside antibiotics are featured in the experiments described herein, they are not the only region or chemical class which can be used in this invention.

For example, messenger RNA regulatory elements, telomerase RNA (see, e.g., Bahattacharyya and Blackburn, EMBO J., 13:5721-31, 1994), oncogene mRNAs, cytokine and lymphokine mRNAs, thymidylate synthase mRNAs, and various viral elements such as the adenovirus late mRNA tripartite leader, hepatitis delta virus RNA, and picornavirus "ribosomal landing pads," or "internal ribosomal entry sites," located in the 5' untranslated region, can also be used to design oligoribonucleotide analogs of the invention.

- 25 -

Moreover, self-splicing group I introns, including the HDV group I intron, are inhibited by the binding of aminoglycosides, and thus can be used to design RNA oligoribonucleotide analogs. Furthermore, the HIV
5 Rev-Response-Element (RRE) is bound by some of the same antibiotics that bind to the decoding region of 16S rRNA. Neomycin binding to the RRE inhibits Rev function and viral replication in model cell culture systems. Thus, the RRE is a suitable target for designing
10 oligoribonucleotide analogs that are useful to assay antibiotics effective against the RRE.

Any known RNA structural motif can be used as parental RNA to design an oligonucleotide analog that can be used as a screening target, and because the transport,
15 processing, and translation of mRNA depend on such structures and their interactions with various binding partners, a large array of potential targets in viral and cellular mRNA exist. Extending these ideas even further, if one assumes that mutant cellular mRNA sequences
20 produce altered mRNA structures, it may even be possible to selectively target specific mutant mRNA sequences with small molecules.

In addition to the first nucleotide sequences derived from the parental RNA, the oligoribonucleotide
25 analogs of the invention also contain second, heterologous sequences (e.g., artificial stem loops and nucleotide clamps) that promote the appropriate three-dimensional conformation of the first nucleotide sequences for normal molecular interactions with
30 candidate drugs, nucleic acid molecules, etc. The small size of these candidate drugs and molecules makes them amenable to numerous inexpensive *in vitro* assays that use a variety of oligoribonucleotide analogs.

Protein synthesis inhibitors have two main
35 ribosomal targets: the decoding region of 16S rRNA, and

- 26 -

the peptidyl transferase region (Domain V) of 23S rRNA. Aminoglycosides interact with the decoding region while MLS antibiotics (macrolides, lincosamides, and streptogramins) interact with the peptidyl transferase region. Peptides, nucleic acid molecules, and a variety of other chemical compounds and molecules may be useful as therapeutics, and can easily be screened using the oligoribonucleotide analogs and methods of this invention.

10 The oligoribonucleotide analogs enable the use of improved methods to screen for novel therapeutic compounds, particularly antibiotic compounds that inhibit protein synthesis. Standard methods to screen antibiotics involve detecting antimicrobial or
15 bactericidal activity in cultured cells over long time periods (up to several days). Screening many potential antibiotic compounds under these conditions can be expensive and time consuming. On the other hand, screening assays using the oligoribonucleotide analogs of
20 the invention have several distinct advantages over these known screening methods.

 The new screening assays are much more rapid (an assay according to the invention can be carried out in about an hour), the individual assays require only small
25 amounts of materials (volumes of about 100 μ l), and numerous reactions can be carried out in parallel (e.g., in a 96 well microtiter plate). Using the methods of the invention, likely candidate compounds can be rapidly identified, and only the likely candidates then could be
30 further tested with cell culture assays.

 Some of the advantages derive from the analogs themselves compared to full-sized ribosomes. The oligoribonucleotide analogs are small enough to be easily produced, either enzymatically as T7 polymerase
35 transcripts, or chemically by automated RNA synthesis.

- 27 -

The analogs are far more stable than ribosomes because they are radically simpler. Furthermore, unlike ribosomes, their labile ribophosphodiester backbone can be stabilized by incorporation of 2'-O-methyl
5 ribonucleotides, deoxyribonucleotides, or phosphorothioates, either uniformly or at selected positions.

In addition, analogs can be readily derivatized during automated chemical synthesis. For example,
10 fluorescent moieties can be introduced at either termini (5' or 3'), or internally, and coupling moieties, such as primary amines or biotin, can be introduced at either termini (see discussion below). Further, the detection of interactions between new test compounds and analogs is
15 more likely to be relevant to potential antibiotic activity, because, unlike ribosomes, the new analogs present a small, well-defined target.

Screening Assays

No particular class of molecular structure is
20 selected by the following screening methods. Rather, the methods select only for the existence of highly specific interactions involving particular nucleotide atoms in the analog RNA target. Therefore, specific compounds can be tested to determine whether they are likely to have
25 antibiotic activity.

General Considerations

Aminoglycosides are polycations and charge-charge interactions with the negatively charged phosphodiester backbone of the oligoribonucleotide analogs are likely
30 important determinants of their binding specificity and affinity. Thus, important factors governing the formation and stability of aminoglycoside-RNA

- 28 -

interactions include mono- and divalent-cation (salt) concentrations, and the presence or absence of nonspecific nucleic acids. Accordingly, low salt concentrations (<100 mM K⁺, NH₄⁺, Na⁺, <5 mM Mg⁺⁺) favor
5 complex formation, and the presence of nonspecific nucleic acids disfavor it. Candidate compounds to be tested should, therefore, be prepared in low salt buffers largely free of contaminating nucleic acids.

To facilitate high throughput screening, a
10 standard microtiter plate format with a standard fluorescence or radiation detector is used. A robotic system is used to load, wash, and detect a signal in each well. Cation concentrations, nonspecific nucleic acids, and other variables including temperature and time can be
15 adjusted during binding and washing stages to adjust background binding levels.

Three Screening Methods

Figs. 5A and 5B illustrate two specific screening assays based on 1) binding of test compounds to an
20 immobilized (naked) oligoribonucleotide analog functioning as a reporter ("affinity assay"), and 2) displacement of a prebound reporter molecule from an immobilized analog by test compounds ("competitive binding assay"), respectively. These two methods, while
25 distinct, are related in their overall strategy and organization. A third method (Figs. 7A, 7B, 8A, and 8B), based on oligomer binding to an *in situ* footprinted analog, differs in its organization and strategy from the first two methods, and will be discussed separately.

30 **Method 1: Affinity Assay**

This screening method can be implemented with an immobilized, fluorescently labeled oligoribonucleotide analog, as shown in Fig. 5A. The method consists of (1)

- 29 -

incubation of the immobilized analog with test compounds, (2) wash step(s), and subsequent detection of complex formation relying on changes in the fluorescence properties of the immobilized RNA analog. The RNA analog can be derivatized, e.g., during automated chemical synthesis, both for immobilization and for labelling. Possible positions for derivatization on an oligoribonucleotide analog are indicated by boxes around the nucleotides in Fig. 6.

For example, for immobilization, the analog can be derivatized with either terminal biotin (using biotin phosphoramidite (5'-biotin) or CPG (3'-biotin)), or a free primary amino group (using amino modifier phosphoramidite (5'-NH₂) or CPG (3'-NH₂)).

Immobilization in microtiter plate wells can then be accomplished with polystyrene plates activated with streptavidin or maleic anhydride, respectively. After coupling, plates should be thoroughly washed to eliminate soluble analog RNA.

For labelling, analogs are derivatized to allow the addition of a fluorophore at one of the RNA termini with fluorescein phosphoramidites (3' labeling) or CPG (5' labeling), or the incorporation of fluorescent adenosine or cytosine nucleotides (with EthenoA and EthenoC phosphoramidites, respectively) at specific internal positions in the RNA. The latter approach can be much more sensitive, due to the proximity of the fluorophore and antibiotic binding site. Alternatively, a fluorescent dye such as ethidium bromide, or possibly Hoechst 33258, can be used with underivatized RNA. This labelling allows the detection of any changes in either the binding or fluorescent properties of the dye, or both, upon test compound-analog complex formation.

Typical (solution) binding reactions contain 25 mM neutral pH buffer (Tris or Hepes) and 50 mM monovalent

- 30 -

salt (KCl, NH₄Cl, etc). It is advantageous to carry out binding reactions at 30-37°C for approximately 30 minutes to facilitate the simultaneous annealing of the RNA. Washes can utilize the same buffer.

5 Test compound and salt concentrations should be adjusted empirically to optimize the assay. For example, the test compound concentration should be in the micromolar range, and salt concentrations should be in the range of 10 to 500 mM, but these numbers must be
10 adjusted depending on the particular assay conditions. In addition, nonspecific nucleic acids can be added to the binding step to reduce the effects of any contaminating compounds with nonspecific affinity for nucleic acids. For example, the nonspecific nucleic acid
15 could be total yeast tRNA, poly I-C, etc. Another potentially useful nonspecific nucleic acid would be (soluble) A-site analog carrying the G1491U point mutation.

Method 2: Competitive Binding Assay

20 As shown in Fig. 5B, this screening method is based on the hypothesis that useful test compounds will displace prebound soluble reporter ligands from their immobilized analog binding sites. For example, immobilized neomycin (an antibiotic ligand known to bind
25 to analogs derived from the decoding region of 16S rRNA) can be (1) pre-loaded with a labeled analog, and (2) test compounds added and allowed to compete with immobilized neomycin for the labeled analog, which serves as the reporter. After a wash step to remove any displaced
30 analog, still complexed with the test compound, detection of the absence of label in specific wells would indicate that specific test compounds bound the analog RNA efficiently, a positive result. The reciprocal configuration, with immobilized analog RNA and labeled
35 neomycin, can also be used.

- 31 -

Neomycin and other aminoglycosides contain numerous primary amino groups that are ideal targets for coupling to either amine-reactive fluorescent probes or activated polystyrene. For example, a soluble neomycin reporter molecule can be produced by coupling neomycin to NHS-Fluorescein. In the alternative, neomycin can be immobilized on maleic anhydride-activated polystyrene microtiter plates. As discussed above, the oligoribonucleotide analog can be similarly labeled or immobilized.

Method 3: Screening By *In situ* Footprinting

Figs. 7A and 7B, and 8A and 8B, illustrate a third screening method based directly on our chemical probing experience which has shown that the N1 atom of A1408 is strongly methylated by DMS in the absence of neomycin, and conversely, is strongly protected from DMS methylation in the presence of neomycin. In conventional footprinting experiments, this differential modification provides a method for detection of complex formation (monitored by inhibition of primer extension). Method 3 similarly exploits differential methylation, but monitors this methylation, e.g., via the differential annealing of a labeled complementary oligomer (e.g., CAGUGU) with a binding site overlapping the site of methylation, A1408 (Figs. 7A and 7B).

In this method, the N1 atom of A1408 is strongly methylated by dimethyl sulfate (DMS) in the absence of neomycin (Fig. 7A) and, conversely, strongly protected from DMS methylation in the presence of neomycin ("bound small molecule") (Fig. 7B). Thus, after complex formation (step 1) and DMS modification (step 2), the methylation state of A1408 reflects its (previous) involvement in complex formation with neomycin.

The mechanics of the assay are similar to Method 1 in that the first step involves forming complexes between

- 32 -

the immobilized (but here unlabeled) analog and test compounds (Step 1). After complex formation and washing steps, the plate is treated with DMS to differentially methylate the RNA (Step 2). DMS reactions are then
5 stopped with a 1M β ME wash, and labeled reporter oligo added and annealed (Step 3).

The technique for monitoring differential methylation illustrated in Figs. 7A and 7B relies on the inability of N1-methylated A to form Watson-Crick base
10 pairs with a short labeled oligomer. In Fig. 7A, with uncomplexed analog and methylated N1, the short oligo fails to anneal to the analog and is washed away, leaving no label in that well. In Fig. 7B, with complexed analog and unmethylated N1, the oligo anneals to the analog and
15 is effectively immobilized on it, leaving label in the well. Wells that retain the reporter oligo indicate that A1408 was not methylated, which in turn indicates that A1408 was protected by a bound antibiotic ligand. Thus, the presence of label in a well constitutes a positive
20 result.

An alternative technique for monitoring differential methylation shown in Figs. 8A and 8B relies on reverse transcription of the modified template. Steps 1 and 2 are identical to those shown in Figs. 7A and 7B,
25 except that the target analog (RNA) is no longer immobilized. Instead, a biotinylated primer (e.g., having the nucleotide sequence TTCACCCGGAAGGTG; SEQ ID NO:12) is added to each reaction (Figs. 8A and 8B), and primer extension is initiated in the presence of labeled
30 TTP (e.g., α -³²P-TTP) and reverse transcriptase.

In Fig. 8A, primer extension is inhibited by methylation of N1 of A1408 and no label is incorporated. In Fig. 8B, primer extension is successful, resulting in the incorporation of label into the primer. In both
35 cases the biotinylated primer is immobilized via

- 33 -

interaction with streptavidin-coated wells. After a wash step to remove unincorporated labeled TTP, only the scenario in Fig. 8B will allow retention of the label in the well, thus reporting the presence of a bound ligand
5 in step 1.

The reverse-transcriptase-based reporter methodology superficially resembles standard chemical probing methods. The main difference, aside from the immobilized primer, is that the primer extension
10 conditions, e.g., Mg concentration, reaction time, temperature, and amount of enzyme, are adjusted, using standard techniques, such that methylation of A1408 will completely stop extension of a high percentage (approximately 90%) of the transcripts. This is the
15 opposite of the desired situation in standard experiments, where conditions are adjusted such that greater than 90% of the transcripts are full-length.

Other Embodiments

It is to be understood that while the invention
20 has been described in conjunction with the detailed description thereof, that the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are
25 within the scope of the following claims.

- 34 -

SEQUENCE LISTING**(1) GENERAL INFORMATION:**

5 (i) APPLICANT: University of
 Massachusetts Medical
 Center

 (ii) TITLE OF INVENTION: OLIGORIBONUCLEOTIDE ASSAY
 FOR NOVEL ANTIBIOTICS

 (iii) NUMBER OF SEQUENCES: 14

 (iv) CORRESPONDENCE ADDRESS:

10 (A) ADDRESSEE: Fish & Richardson P.C.
 (B) STREET: 225 Franklin Street
 (C) CITY: Boston
 (D) STATE: Massachusetts
 (E) COUNTRY: U.S.A.
15 (F) ZIP: 02110-2804

 (v) COMPUTER READABLE FORM:

 (A) MEDIUM TYPE: 3.5" Diskette, 1.44 Mb
 (B) COMPUTER: IBM PS/2 Model 50Z or
 55SX
20 (C) OPERATING SYSTEM: MS-DOS (Version 5.0)
 (D) SOFTWARE: WordPerfect (Version 5.1)

 (vi) CURRENT APPLICATION DATA:

 (A) APPLICATION NUMBER:
 (B) FILING DATE:
25 (C) CLASSIFICATION:

 (vii) PRIOR APPLICATION DATA:

 (A) APPLICATION NUMBER: 08/294,450
 (B) FILING DATE: August 23, 1994

 (viii) ATTORNEY/AGENT INFORMATION:

30 (A) NAME: J. Peter Fasse
 (B) REGISTRATION NUMBER: 32,983
 (C) REFERENCE/DOCKET NUMBER: 04020/047WO1

 (ix) TELECOMMUNICATION INFORMATION:

35 (A) TELEPHONE: (617) 542-5070
 (B) TELEFAX: (617) 542-8906
 (C) TELEX: 200154

- 35 -

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 1:**(i) SEQUENCE CHARACTERISTICS:**

5 (A) LENGTH: 49
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GCACAGACCG CCCGUCACAC CUUCGGGUGA AGUCGUAACA AGGCUGUGC 49

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 2:

10 **(i) SEQUENCE CHARACTERISTICS:**

(A) LENGTH: 23
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 **(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:**

CGUCACACCU UCGGGUGAAG UCG 23

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 3:**(i) SEQUENCE CHARACTERISTICS:**

20 (A) LENGTH: 17
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GGGGAUUGAA AAUCCCC 17

25 **(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 4:**

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 21
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GGUCAUCUUG GGGUGAUGAC C 21

- 36 -

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 5:**(i) SEQUENCE CHARACTERISTICS:**

5 (A) LENGTH: 17
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GGGAGCGUCA UCACUA 17

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 6:

10 **(i) SEQUENCE CHARACTERISTICS:**

(A) LENGTH: 14
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 **(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:**

GGGACUUCGG UCCC 14

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 7:**(i) SEQUENCE CHARACTERISTICS:**

20 (A) LENGTH: 13
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

GGCGCUUUGC GCC 13

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 8:**(i) SEQUENCE CHARACTERISTICS:**

30 (A) LENGTH: 66
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 37 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GCACUAUGGG CGCAGCGUCA AUGACGCUGA CGGUACAGGC CAGACAAUUA UUGUCUGGUA 60
UAGUGC 66

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 9:**5 (i) SEQUENCE CHARACTERISTICS:**

(A) LENGTH: 13
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

ACCGCCCGUC ACA 13

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 10:**(i) SEQUENCE CHARACTERISTICS:**

15 (A) LENGTH: 16
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

UGAAGUCGUA ACAAGG 16

20 (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 11:**(i) SEQUENCE CHARACTERISTICS:**

25 (A) LENGTH: 27
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 38 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

GGCGUCACAC CUUCGGGUGA AGUCGCC 27

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 12:

(i) SEQUENCE CHARACTERISTICS:

5 (A) LENGTH: 13
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

10 ACCGCCCCGUC ACA 13

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 13:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 16
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

UGAAGUCGUA ACAAGG 16

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 14:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

TTCACCCGGA AGGTG 15

What is claimed is:

- 39 -

1. An oligoribonucleotide analog of a region of a parental ribonucleic acid (RNA) comprising

a first nucleic acid structure comprising one or more nucleotide sequences, said first structure being
5 derived from the region of parental RNA, wherein in its native state, the region binds to a ligand with a parental RNA ligand binding pattern, and

a second nucleic acid structure comprising one or more nucleotide sequences combined with said first
10 nucleic acid structure to form said analog and provide said analog with a conformation that binds said ligand with a ligand binding pattern that is substantially identical to said parental RNA ligand binding pattern.

2. An analog of claim 1, wherein said first
15 nucleic acid structure is derived from a region of a 16S ribosomal RNA (rRNA).

3. An analog of claim 2, wherein said region comprises a decoding region of 16S rRNA.

4. An analog of claim 3, wherein said region
20 comprises nucleotides 1398-1410 and 1490-1505 of 16S rRNA.

5. An analog of claim 3, wherein said region comprises the A site subdomain of the decoding region of 16S rRNA.

25 6. An analog of claim 5, wherein said region comprises nucleotides 1404-1410 and 1490-1497 of 16S rRNA.

7. An analog of claim 1, wherein said second nucleic acid structure comprises a tetraloop.

8. An analog of claim 7, wherein said tetraloop comprises the nucleotide sequence: 5'-CCUUCGGG-3'.

10. An analog of claim 9, wherein said base-paired nucleotide clamp comprises the nucleotide sequence:

3'-CGUGUC-5'
5'-GCACAG-3'.

3'-CC-5'
5'-GG-3'.

and the complete linear nucleotide sequence of said combined first and second nucleotide structures of said analog is

30 (SEQ ID NO:1).

- 41 -

14. An analog of claim 12, wherein said decoding region comprises nucleotides 1404-1410 and 1490-1497 of 16S rRNA, said tetraloop comprises the nucleotide sequence 5'-CCUUCGGG-3', said base-paired nucleotide
5 clamp comprises the nucleotide sequence: 3'-CC-5'
5'-GG-3',
and the complete linear nucleotide sequence of said combined first and second nucleotide structures of said analog is
10 5'-GGCGUCACACCUUCGGGUGAAGUCGCC-3' (SEQ ID NO:11).

15. An affinity assay for determining the potential therapeutic activity of a test compound, said assay comprising the steps of
mixing a test compound with an analog of claim 1
15 under conditions that allow formation of a binding complex between said analog and said test compound, and
detecting the formation of a binding complex,
wherein the presence of a binding complex indicates that said test compound has potential therapeutic activity.

20 16. A method of claim 15, wherein said analog is labelled and immobilized on a surface, and said binding complex is detected by monitoring changes in the signal of said label when a test compound is bound to said analog.

25 17. A method of claim 15, wherein said analog is immobilized on a surface, said test compound is labelled, and said binding complex is detected by detecting any of said label bound to said surface.

18. A method of claim 16, wherein said analog is
30 fluorescently or radioactively labelled.

- 42 -

19. A competitive binding assay for determining the potential therapeutic activity of a test compound, said assay comprising the steps of

5 mixing an analog of claim 1 with an analog-binding ligand under conditions that allow formation of a first binding complex between said analog and said ligand, mixing a test compound with the first binding complex under conditions that allow said test compound to disrupt said first binding complex to form a second
10 binding complex between said analog and said test compound, and

detecting the disruption of the first binding complex, wherein the disruption of the first binding complex indicates that said test compound has potential
15 therapeutic activity.

20. A method of claim 19, wherein said ligand is labelled, said analog is immobilized on a surface, and the disruption of the first binding complex is detected by monitoring any decrease in the signal of said label
20 when a test compound displaces said ligand from said first binding complex.

21. A method of claim 19, wherein said analog is labelled, said ligand is immobilized on a surface, and the disruption of the first binding complex is detected
25 by monitoring any decrease in the signal of said label when a test compound displaces said analog from said first binding complex.

22. A method of claim 21, wherein said analog is fluorescently or radioactively labelled.

30 23. A method of claim 21, wherein said ligand is fluorescently or radioactively labelled.

- 43 -

24. An *in situ* footprinting assay for determining the potential therapeutic activity of a test compound, said assay comprising the steps of

5 mixing an analog of claim 1 with a test compound
under conditions that allow formation of a binding
complex between said analog and said test compound,
incubating said binding complex with a chemical
probing reagent and monitoring for an effect of said
reagent on said analog in said complex,
10 in a separate control reaction, incubating said
analog unbound to any test compound with said chemical
probing reagent and monitoring for an effect of said
reagent on the unbound analog, and
comparing any effects of said probing reagent on
15 said analog in said binding complex and on said unbound
analog, wherein prevention of an effect of said reagent
on said analog in said binding complex caused by said
reagent on said unbound analog indicates that said test
compound has potential therapeutic activity.

20 25. An *in situ* footprinting assay of claim 24,
wherein said chemical probing reagent is dimethyl
sulfate, kethoxal, or carbodiimide.

26. An *in situ* footprinting assay of claim 24,
wherein said probing reagent covalently modifies a
25 nucleotide in said analog when unbound to any test
compound.

27. An *in situ* footprinting assay of claim 26,
wherein the effect of said probing reagent is monitored
by use of a labelled oligonucleotide that hybridizes to
30 said analog when protected by said test compound from
methylation, and does not hybridize to said analog when
methylated by said reagent, the presence of said label

- 44 -


after completion of said assay indicating that said test compound has potential therapeutic activity.

28. An *in situ* footprinting assay of claim 27, wherein said labelled oligonucleotide has the sequence
5 CAGUGU.

29. An *in situ* footprinting assay of claim 26, wherein the effect of said probing reagent is monitored by use of an oligonucleotide primer that is complementary to a portion of said analog, a labelled nucleotide, and
10 reverse transcriptase, wherein extension of said primer on said analog with said labelled nucleotide does not occur when said analog is methylated by said reagent, the presence of said label after completion of said assay
15 indicating that said test compound has potential therapeutic activity.

30. An *in situ* footprinting assay of claim 29, wherein said oligonucleotide primer has the sequence TTCACCCGGAAGGTG (SEQ ID NO:12).

31. An analog of claim 1, wherein said ligand is
20 an aminoglycoside.



1111


165 1774

1940	1941	1942	1943	1944	1945	1946	1947	1948	1949	1950	1951	1952	1953	1954	1955	1956	1957	1958	1959	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100																																																																																																																													
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	2

116-118

16:11

Page 110



Protection:

- Neomycins






Fig. 12

SEO 1A NO: 11

FIG. 2A



570. 13



4/10




FIG. 3




FIG. 4A




FIG. 4B

6/10




Fig. 5A

7/10




Fig. 53

8/10




Fig. 6

9/10




Fig. 7A




Fig. 7B




Fig. 2A




Fig. 2B

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/10721

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07H 21/02; C12Q 1/68

US CL : 536/22.1; 435/6

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/22.1; 435/6

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	Nature, Volume 370, issued 25 August 1994, Purohit et al, "Interactions of a small RNA with antibiotic and RNA ligands of the 30S subunit", pages 659-662, see entire document.	1-31
X	FEBS Letters, Volume 279, No. 2, issued February 1991, D. Leclerc et al, "A conformational switch involving the 915 region of <i>Escherichia coli</i> 16 S ribosomal RNA", pages 171-174, see entire document.	1-12
X ----- Y	J. Mol. Biol., Volume 212, issued 1990, E. A. De Stasio et al, "Effects of mutagenesis of a conserved base-paired site near the decoding region of <i>Escherichia coli</i> 16 S ribosomal RNA", pages 127-133, see entire document.	1-12 ----- 13-31



Further documents are listed in the continuation of Box C.



See patent family annex.

Special categories of cited documents:	
A document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
E earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O document referring to an oral disclosure, use, exhibition or other means	*Z* document member of the same patent family
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

31 OCTOBER 1995

Date of mailing of the international search report

20 DEC 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JEFFREY FREDMAN

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US95/10721

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, P	Nature, Volume 370, issued 25 August 1994, R. Schroeder, "Dissecting RNA function", pages 597-598.	1-31
A	Nature, Volume 353, issued 26 September 1991, U. V. Ahsen et al, "Antibiotic inhibition of group I ribozyme function", pages 368-370.	1-12
X	Biochimica et Biophysica Acta, Volume 1089, issued 1991, C. Bonny et al, "Analysis of streptomycin-resistance of <i>Escherichia coli</i> mutants", pages 213-219, see entire document.	1-12
-----		-----
Y		13-31
A	Biochemistry, Volume 26, issued 1987, M. Gravel et al, "Cross-linking of streptomycin to the 16S ribosomal RNA of <i>Escherichia coli</i> ", pages 6227-6232.	1-31

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/10721

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

MEDLINE, BIOSIS, CAPLUS, INPADOC, WPIDS, APS, GENBANK, EMBL

search terms: 16S rRNA, decoding, screening, antibiotic, aminoglycoside, neomycin, affinity, binding, ligand