




(86) Date de dépôt PCT/PCT Filing Date: 2003/07/14  
 (87) Date publication PCT/PCT Publication Date: 2004/01/22  
 (45) Date de délivrance/Issue Date: 2011/08/30  
 (85) Entrée phase nationale/National Entry: 2005/01/11  
 (86) N° demande PCT/PCT Application No.: JP 2003/008904  
 (87) N° publication PCT/PCT Publication No.: 2004/006956  
 (30) Priorités/Priorities: 2002/07/12 (JP2002-204725);  
 2003/01/16 (JP2003-8230)

(51) Cl.Int./Int.Cl. *A61K 31/53* (2006.01),  
*A61K 31/185* (2006.01), *A61K 31/381* (2006.01),  
*A61K 31/403* (2006.01), *A61K 31/4164* (2006.01),  
*A61K 31/4196* (2006.01), *A61K 31/454* (2006.01),  
*A61K 31/502* (2006.01), *A61K 45/00* (2006.01),  
*A61K 45/06* (2006.01), *A61P 25/00* (2006.01),  
*C12Q 1/00* (2006.01)

(72) Inventeurs/Inventors:  
 URADE, YOSHIHIRO, JP;  
 EGUCHI, NAOMI, JP;  
 ARITAKE, KOSUKE, JP;  
 SATO, YO, JP;  
 KADOYAMA, KEIICHI, JP; ...

(54) Titre : MEDICAMENTS AMELIORANT LE PRONOSTIC DE TROUBLE CEREBRAL ET PROCEDE DE CRIBLAGE ASSOCIE  
 (54) Title: DRUGS FOR IMPROVING THE PROGNOSIS OF BRAIN INJURY AND A METHOD OF SCREENING THE SAME



(57) Abrégé/Abstract:

The present invention is directed to a compound for treatment or prevention of brain injury caused by diseases such as cerebrovascular disorder, brain degenerative disease and demyelinating disease and a method for screening the same. Brain injury in which prostaglandin D<sub>2</sub> participates is treated or prevented by inhibition of hematopoietic prostaglandin D synthase induced in microglia cell or macrophage of a brain injury area by diseases such as cerebrovascular disorder, brain degenerative disease or

(72) Inventeurs(suite)/Inventors(continued): TANIKE, MASAKO, JP

(73) Propriétaires(suite)/Owners(continued): JAPAN SCIENCE AND TECHNOLOGY AGENCY, JP

(74) Agent: KIRBY EADES GALE BAKER

(57) Abrégé(suite)/Abstract(continued):

demyelinating disease or by the inhibition of activation of prostaglandin D receptor expressed in astroglia cell around the injured area. The invention also provides a method of testing those pharmaceutical substances using a transgenic mouse in which human hematopoietic prostaglandin D synthase is expressed in large amounts.

## ABSTRACT

The present invention is directed to a compound for treatment or prevention of brain injury caused by diseases such as cerebrovascular disorder, brain degenerative disease and demyelinating disease and a method for screening the same. Brain injury in which prostaglandin D<sub>2</sub> participates is treated or prevented by inhibition of hematopoietic prostaglandin D synthase induced in microglia cell or macrophage of a brain injury area by diseases such as cerebrovascular disorder, brain degenerative disease or demyelinating disease or by the inhibition of activation of prostaglandin D receptor expressed in astroglia cell around the injured area. The invention also provides a method of testing those pharmaceutical substances using a transgenic mouse in which human hematopoietic prostaglandin D synthase is expressed in large amounts.

DRUGS FOR IMPROVING THE PROGNOSIS OF BRAIN INJURY  
AND A METHOD OF SCREENING THE SAME

5

## TECHNICAL FIELD

This invention relates to a compound for treating or preventing brain injury and to a method of screening the same. More particularly, this invention relates to a compound which  
10 treats or prevents brain injury in which prostaglandin D<sub>2</sub> participates by inhibition of hematopoietic prostaglandin D synthase (hereinafter, may be referred to as "H-PGDS") induced in microglia cell or macrophage of a brain injury site by disease such as cerebrovascular accident, neurodegenerative disease or  
15 demyelinating disease or by inhibition of activation of prostaglandin D receptor (hereinafter, may be referred to as "DP receptor") expressed in astroglial cells around the injured site. This invention also relates to a method of testing those pharmaceutical substances using a human hematopoietic  
20 prostaglandin D synthase over-expressing transgenic mouse.

## BACKGROUND ART

As a result of development of medical technology, people narrowly escaping death even if the head suffers from  
25 a serious injury are increasing. There are many people who suffer from head injury caused by traffic accidents, work-related accidents, sports, etc. and, in fact, one half of the people who are killed by traffic accidents die because of head injury. They are caused by brain edema or a brain contusion generated

immediately beneath the cranium when it is pressed by external force. The cause of brain edema is breakage of a blood-brain barrier existing in cerebral blood vessels and is edema of the brain caused by leakage of plasma components outside the blood vessel. In recent years, hypothermia has been receiving public attention as a treating method therefor. It is a treating method where exacerbation of brain edema and increase in pressure in the cranium are suppressed by suppression of brain metabolism by induced hypothermia. Although that is an important treatment policy, there are some cases where problems occur by lowering of cardiopulmonary function and immunological function due to low body temperature (*N. Engl. J. Med.*, 2001; 344:556-563).

#### SUMMARY OF THE INVENTION

An object of the present invention, at least in preferred embodiments, is to provide a compound which is able to suppress the tissue injury by delay of local inflammation of the brain and to improve the prognosis.

Another object of the present invention, at least in preferred embodiments, is to provide a method of screening such a compound.

In order to achieve the above-mentioned objects, the present inventors have carried out intensive studies and have found the following and, on the basis thereof, the present invention has been accomplished.

1) When hereditary or traumatic brain injuries occur, expression of H-PGDS and DP receptor increases.

2) Expression of H-PGDS is induced in activated microglia cells or accumulated macrophage of a brain injury while DP receptor is induced in astroglia cells around the injured area.

3) In the injured area, accumulation of macrophage and activation of astroglia cells are significant.

4) When an inhibitor for H-PGDS or an antagonist for DP receptor is administered, expression of DP receptor decreases  
5 and activation of astroglia cell is suppressed.

5) In H-PGDS over-expressing transgenic mouse, brain injury is exacerbated as compared with a mouse of a wild type.

6) In an H-PGDS knockout mouse and DP receptor knockout mouse, bleeding at the injured site and activation of astroglia  
10 cell are slight as compared with a mouse of a wild type.

Thus, the gist of the present invention is a pharmaceutical composition used for treatment or prevention of a brain injury containing an inhibitor or suppressor for hematopoietic prostaglandin D synthase (H-PGDS) as an active ingredient.

15 The term "brain injury" includes not only traumatic injuries caused by traffic accidents or the like but also those caused by cerebrovascular disorder such as cerebral infarction and cerebral bleeding, neuronal degenerative disease including Alzheimer's disease, multiple sclerosis, etc. and  
20 demyelinating disease and the like. The term "treatment or prevention of brain injury" includes treatment or prevention of brain contusions, brain edema, cerebral infarction, cerebral bleeding, ischemic brain diseases, Alzheimer's disease, multiple sclerosis and demyelinating disease.

25 Another gist of the present invention is a pharmaceutical composition used for treatment or prevention of brain injury containing an antagonist for prostaglandin D receptor as an effective ingredient.

Still another gist of the present invention is a method

for treatment of brain injury including administration of a hematopoietic prostaglandin D synthase (H-PGDS) inhibitor of an effective dose.

Still another object of the present invention, at least in preferred embodiments, is the use of a hematopoietic prostaglandin D synthase (H-PGDS) inhibitor for the manufacture of a drug for treatment of a brain injury.

Still another object of the present invention, at least in preferred embodiments, is a method for treatment of a brain injury including administration of a prostaglandin D receptor antagonist of an effective dose.

Still another object of the present invention, at least in preferred embodiments, is the use of a prostaglandin D receptor antagonist for the manufacture of a drug for treatment of a brain injury.

Still another object of the present invention, at least in preferred embodiments, is a pharmaceutical composition to be used for treatment or prevention of a brain injury containing a hematopoietic prostaglandin D synthase (H-PGDS) inhibitor and a prostaglandin D receptor antagonist as effective ingredients.

Still another object of the present invention, at least in preferred embodiments, is a method of screening of a compound used for treatment or prevention of a brain injury including that

1) trauma is applied to the brain of a human H-PGDS over-expressing transgenic mouse,

2) a candidate compound is administered to the transgenic mouse before or after applying the trauma and

3) a state of the trauma in the mouse is compared with a state of a transgenic mouse to which no candidate compound is administered.

## BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the changes in expression level of H-PGDS and DP receptor mRNAs in the cerebrum and the cerebellum of Twitcher mice.

Fig. 2 shows H-PGDS locally existing in macrophage and microglia cells of Twitcher mice by an immunohistochemistry.

Fig. 3 shows DP receptors expressed in activated astroglia cells.

Fig. 4 shows the changes in expression level of H-PGDS and DP receptor mRNAs in mice suffering from experimental autoimmune encephalomyelitis.

Fig. 5 shows the changes in H-PGDS expression level in mice suffering from experimental autoimmune encephalomyelitis.

Fig. 6 shows the time course of H-PGDS mRNA expression in traumatic brain injury models.

Fig. 7 shows the time course of DP receptor mRNA expression in traumatic brain injury models.

Fig. 8 shows the time course of tissue edema in traumatic brain injury models.

Fig. 9 shows the time course of H-PGDS expression in traumatic brain injury models.

Fig. 10 shows the time course of astroglia-activation (GFAP activities) in traumatic brain injury models.

Fig. 11 shows expression of DP receptor in activated astroglia cells around the injured brain.

Fig.12 shows a comparison of a brain injury in a wild-type mouse after 4 days from a traumatic brain injury with that in a human H-PGDS over-expressing transgenic mouse.

Fig. 13 shows suppression of DP receptor expression and

activation of astroglia cells in a HQL-79-treated Twitcher mouse.

Fig. 14 shows an inhibitory effect of HQL-79 on DP receptor expression after a traumatic brain injury.

Fig. 15 shows a recovery-promoting effect of HQL-79 on a  
5 traumatic brain injury.

Fig. 16 shows a recovery-promoting effect of DP receptor antagonist on a traumatic brain injury.

Fig. 17 shows a comparison of a brain injury in a wild-type mouse after 4 days from a traumatic brain injury with that in a  
10 H-PGDS knockout.

Fig. 18 shows the structure of targeting vector used for the preparation of a transgenic mouse.

Fig. 19 shows the structure of mouse H-PGDS gene (upper column), the structure of mutant sequence in a targeting vector  
15 (middle column) and the structure of a mouse genome DNA after homologous recombination (lower column).

Fig. 20 shows the time course of brain edema in a traumatic brain injury model. The brain edema was evaluated by Evans Blue dye leakage into the tissues.

Fig. 21 shows an inhibitory effect of HQL-79 on tissue  
20 injury (leakage of dye) after a traumatic brain injury (Administration of HQL-79 was started before one hour from injury was applied).

Fig. 22 shows an inhibitory effect of HQL-79 on tissue  
25 injury after a traumatic brain injury (Administration of HQL-79 was started after 24 hours from injury was applied).

Fig. 23 shows an inhibitory effect of HQL-79 on tissue injury (leakage of dye) after a traumatic brain injury (Administration of HQL-79 was started after 24 hours from injury

was applied).

Fig. 24 shows an inhibitory effect of pinagladin on leakage of Evans Blue dye as a result of a traumatic brain injury.

Fig. 25 shows an inhibitory effect on progress of a  
5 traumatic brain injury by pinagladin.

Fig. 26 shows a comparison of a brain injury in a wild-type mouse after two days from a traumatic injury with that in an HPGDS KO or DPR KO mouse using leakage of dye into the injured site.

10 Fig. 27 shows inhibitory effects of BWA 868 and ramatroban on tissue injury (dye leakage) after a traumatic brain injury.

#### DETAILED DESCRIPTION OF THE INVENTION

Examples of an H-PGDS inhibitor include 4-benzhydryloxy-  
15 1-{3-(1H-tetrazol-5-yl)-propyl}piperidine (HQL-79), 1-amino-  
4-{4-[4-chloro-6-(2-sulfo-phenylamino)-[1,3,5]-triazine-2-yl  
methyl]-3-sulfo-phenylamino}-9,10-dioxo-9,10-  
dihydro-anthracene-2-sulfonic acid (Cibacron Blue), 1-amino-  
4-(4-sulfamoylanilino)-anthraquinone-2-sulfonic acid  
20 (PGD-042) or pharmaceutically acceptable salt thereof or  
hydrate thereof and 2-(2'-benzothiazolyl)-5-styryl-3-(4'-  
phthalhydrazidyl)tetrazolium chloride (PGD-016) or a hydrate  
thereof.

In the present specification, examples of the  
25 "pharmaceutically acceptable salt" in the case of salt with base  
include alkaline metal salt such as sodium salt and potassium  
salt; alkaline earth metal salt such as calcium salt and  
magnesium salt; ammonium salt; aliphatic amine salt such as  
trimethylamine salt, triethylamine salt, dicyclohexylamine


salt, ethanolamine salt, diethanolamine salt, triethanolamine salt and procaine salt; aralkylamine salt such as N,N-dibenzylethylenediamine salt; heterocyclic aromatic amine salt such as pyridine salt, picoline salt, quinoline salt and  
5 isoquinoline salt; quaternary ammonium salt such as tetramethylammonium salt, tetraethylammonium salt, benzyltrimethylammonium salt, benzyltriethylammonium salt, benzyltributylammonium salt, methyltrioctylammonium salt and tetrabutylammonium salt; and basic amino acid salt such as  
10 arginine salt and lysine salt. Examples in the case of salt with acid include inorganic acid salt such as hydrochloride, sulfate, nitrate, phosphate, carbonate, hydrogen carbonate and perchlorate; organic acid salt such as acetate, propionate, lactate, maleate, fumarate, tartrate, malate, citrate and  
15 ascorbate; sulfonate such as isothionate, benzenesulfonate and p-toluenesulfonate; and acidic amino acid salt such as aspartate and glutamate. Such a salt can be prepared by conventional methods. When a hydrate is formed, coordination with any number of water molecule(s) is acceptable.

20 Another embodiment of the present invention is a pharmaceutical composition comprising an antagonist for a prostaglandin D receptor as an active ingredient to be used for treatment or prevention of brain injury.


Examples of the antagonist for a prostaglandin D receptor  
25 are  $(\pm)$ -3-benzyl-5-(6-carboxyhexyl)-1-(2-cyclohexyl-2-hydroxyethylamino)-hydantoin (BW A868C), (+)-(3R)-3-(4-fluorobenzenesulfonamide)-1,2,3,4-tetrahydrocarbazol-9-propionic acid (ramatroban), (Z)-7-[(1R,2R,3S,5S)-2-(5-hydroxybenzo[b]thiophene-3-ylcarbonylamino)-10-norpinan-3-

yl]hepta-5-enoic acid, (Z)-7-[(1R,2R,3S,5S)-2-(benzo[b]-thiophene-3-ylcarbonylamino)-10-norpinan-3-yl]hepta-5-enoic acid (pinagladin) and a pharmaceutically acceptable salt thereof and hydrate thereof.

- 5 Further examples of the antagonist for a prostaglandin D receptor are a compound represented by the formula (I):




wherein



- 10 R is hydrogen, alkyl, alkoxy, halogen, hydroxyl, acyloxy or optionally substituted arylsulfonyloxy; X is hydrogen or alkyl; and a double bond of an  $\alpha$ -chain is in an E-configuration or a Z-configuration or a pharmaceutically acceptable salt or a hydrate thereof.


- 15 In a preferred embodiment, the antagonist for a prostaglandin D receptor is a compound represented by the formula (IA):



- 20 wherein R and X have the same meanings as defined above and a double bond of an  $\alpha$ -chain is in an E-configuration or a Z-configuration, or a pharmaceutically acceptable salt or a


hydrate thereof.

More preferably, the antagonist for a prostaglandin D receptor is a compound represented by the formula (IA-a):



5 wherein R and X have the same meanings as defined above and a double bond of an  $\alpha$ -chain is in an E-configuration or a Z-configuration, or a pharmaceutically acceptable salt or a hydrate thereof.

The compound represented by the formula (I) has been known  
10 and can be produced according to the following process:




wherein Y ring, X and R have the same meanings as defined above and a double bond of an  $\alpha$ -chain is in an E-configuration or a Z-configuration.

15 As shown in the above reaction formula, the compound represented by the formula (I) is able to be produced by the reaction of an amino compound represented by the formula (II) with a carboxylic acid or a reactive derivative thereof represented by the formula (III).

20 In the starting compound (II) in the present reaction method, a compound in which



is described in the specification of the Japanese Patent Publication No. 23,170 (1994) B while a compound in which



5 is described in the specifications of the Japanese Patent Laid-Open Nos. 49 (1986) A and 180,862 (1990) A.

The carboxylic acids represented by the formula (III) include

4-bromobenzo[b]thiophene-3-carboxylic acid,

5-bromobenzo[b]thiophene-3-carboxylic acid,

10 6-bromobenzo[b]thiophene-3-carboxylic acid,

7-bromobenzo[b]thiophene-3-carboxylic acid,

5-fluorobenzo[b]thiophene-3-carboxylic acid,

6-fluorobenzo[b]thiophene-3-carboxylic acid,

4-hydroxybenzo[b]thiophene-3-carboxylic acid,

15 5-hydroxybenzo[b]thiophene-3-carboxylic acid,

6-hydroxybenzo[b]thiophene-3-carboxylic acid,

7-hydroxybenzo[b]thiophene-3-carboxylic acid,

5-acetoxybenzo[b]thiophene-3-carboxylic acid,

benzo[b]thiophene-3-carboxylic acid,

20 5-benzosulfonyloxybenzo[b]thiophene-3-carboxylic acid,

5-methylbenzo[b]thiophene-3-carboxylic acid,

6-methylbenzo[b]thiophene-3-carboxylic acid,

5-methoxybenzo[b]thiophene-3-carboxylic acid and

6-methoxybenzo[b]thiophene-3-carboxylic acid.

Each of those carboxylic acids may have the above-defined substituent.

Those carboxylic acids can be produced in accordance with the methods described in *Nippon Kagaku Zasshi*, volume 88, no. 7, pages 758-763 (1967), *Nippon Kagaku Zasshi*, volume 86, no. 10, pages 1067-1072 (1965), *J. Chem. Soc. (c)*, pages 1899-1905 (1967), *J. Heterocyclic Chem.*, volume 10, pages 679-681 (1973), *J. Heterocyclic Chem.*, volume 19, pages 1131-1136 (1982) and *J. Med. Chem.*, volume 29, pages 1637-1643 (1986).

The reactive derivative of the carboxylic acid represented by the formula (III) means the corresponding acid halide (such as chloride, bromide and iodide), acid anhydride (such as that of formic acid or of mixed acid with acetic acid), activated ester (such as succinimide ester), etc. and includes an acylating agent which is usually used for acylation of amino group. In order to prepare an acid halide for example, acid may be reacted with thionyl halide (such as thionyl chloride), phosphorus halide (such as phosphorus trichloride and phosphorus pentachloride), oxalyl halide (such as oxalyl chloride), etc. according to a known method (such as *Shin Jikken Kagaku Koza* [New Experimental Chemistry], volume 14, page 1787 (1978); *Synthesis*, pages 852-854 (1986); and *Shin Jikken Kagaku Koza*, volume 22, page 115 (1992)).

The reaction may be carried out according to the condition for common acylation reaction for amino group. For example, in the case of a condensation reaction using an acid halide, the reaction may be carried out using ether type solvent (such as diethyl ether, tetrahydrofuran and dioxane), benzene type solvent (such as benzene, toluene and xylene), halogenated

hydrocarbon type solvent (such as dichloromethane, dichloroethane and chloroform) and others such as ethyl acetate, dimethylformamide, dimethyl sulfoxide and acetonitrile, etc. as a solvent. The reaction may be carried out under cooling to  
5 at room temperature or heating or, preferably, from -20°C to ice cooling or room temperature to a heating/refluxing temperature of the reaction system for several minutes to several tens of minutes, preferably 0.5 to 24 hour(s) or, more preferably, 1 to 12 hour(s), if necessary, in the presence of  
10 a base (such as an organic base [e.g., triethylamine, pyridine, N,N-dimethylaminopyridine and N-methylmorpholine] or an inorganic base [e.g., sodium hydroxide, potassium hydroxide and potassium carbonate]. When a carboxylic acid is not used as a reactive derivative but in a free form, the reaction is  
15 conducted in the presence of a condensing agent used for a condensation reaction of amine with carboxylic acid (such as dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide or N,N'-carbonyldiimidazole).

The pharmaceutical composition of the present invention  
20 may use both an inhibitor for hematopoietic prostaglandin D synthase (H-PGDS) and an antagonist for prostaglandin D receptor as active ingredients.

Although the compounds used for treatment or prevention of brain injury used in the present invention are able to be  
25 selected from H-PGDS inhibitors or prostaglandin D receptor antagonists as mentioned above, it is also possible to screen as follows.

Thus,

1) traumatic injury is applied to brain of transgenic

mouse in which human H-PGDS is expressed in large amounts,

2) a candidate compound is administered to the transgenic mouse before or after applying the traumatic injury and

3) the state of the traumatic injury in the mouse is compared with the state in a transgenic mouse to which no candidate compound is administered.

A method for the production of a transgenic mouse in which human H-PGDS is expressed in large amounts is disclosed in an international application PCT/JP00/06963 (WO 01/24627) filed on October 5, 2000.

### Examples

#### Preparation Example 1

15 Preparation of a human hematopoietic prostaglandin D synthase over-expressing transgenic mouse

A human hematopoietic prostaglandin D synthase over-expressing transgenic mouse was prepared according to a method disclosed in WO 01/24627.

20 From a cDNA library prepared from mRNA of human cells, cDNA of human h-PGDS (*Eur. J. Biochem.* 267:3315-3322, 2000; GenBank Accession No. NM-014485) was cloned using cDNA of a rat H-PGDS gene (*Cell* 90:1085-10975, 1997; GenBank Accession No. D 82071) as a probe. Then, cDNA of human H-PGDS was inserted  
25 into a cloning site (Sal I/Not I) of a vector pCAGGS (*Gene* 108:193-199 (1991)) to construct a transducing vector. Fig. 18 is a construction of transgene in this transducing vector. The transgene has a CMV enhancer and a chitin  $\beta$ -actin promoter at the upper stream of H-PGDS cDNA and, when it is transduced

into chromosome of a mouse, H-PGDS mRNA is over-expressed by the action of those enhancer and promoter. The transducing vector was infused into a fertilized egg of FVB mouse (obtained from the National Institute of Health Animal Genetic Resource) by a microinjection method. The fertilized egg into which gene was transduced was transplanted to oviduct of an acting parent by a common method, subjected to ontogeny and was born. DNA was extracted from the tail of the resulting mouse and, using a probe which was synthesized depending upon a sequence of the transgene, a transgenic mouse was selected by a Southern blot technique.

#### Preparation Example 2

#### Preparation of a hematopoietic prostaglandin D synthase knockout (H-PGDS KO) mouse

A hematopoietic prostaglandin D synthase knockout mouse was prepared according to a method taught in the Japanese Patent Application No. 2002/18666 filed on January 28, 2002.

A region including exon II (protein translation initiation region of H-PGDS) of known mouse H-PGDS gene was substituted with Neo<sup>r</sup> gene, then a mutant sequence was prepared by integration of thymidine kinase gene of herpes virus (HSV-tk gene) into about 7 Kb upstream of H-PGDS gene and the mutant sequence was integrated into a vector to prepare a targeting vector (refer to Fig. 19).

Targeting vector was transduced at the rate of 48 µg/ml into a non-differentiated incubated ES cells ( $1.2 \times 10^7$  cells) by electroporation to prepare ES cells into which gene was transduced. The cells were sown on a plate, G418 and ganciclovir were added to the medium after 2 days and incubation was conducted for 7 days more to prepare colonies showing a

resistance to G418 and ganciclovir. Those colonies were individually separated and further incubated, DNA was extracted and homologous recombinant ES cells were selected by a Southern blot technique.

5 After that, the homologous recombinant ES cell was infused into blastocyst of a mouse of C57BL/6 strain by a common method, transplanted to an acting parent and subjected to ontogeny.

10 As a result, 10 chimera mice were obtained. Among the resulting chimera mice, a female individual was crossed with a female wild-type mouse of C57BL/6 strain to give first generation (F<sub>1</sub>) mice. From those F<sub>1</sub> mice, individuals (male/female) where mutant sequence was confirmed in one of diploid chromosomes by a Southern blot analysis were selected  
15 and crossed to give secondary generation (F<sub>2</sub>) mice.

Finally, from those F<sub>2</sub> mice, individuals where mutant sequence was confirmed in both dipolar chromosomes (homozygotes) and individuals where mutant sequence was confirmed in one of dipolar chromosomes (heterozygotes) were  
20 selected by a Southern blot analysis to prepare H-PGDS knockout mice.

#### Example 1

Transduction of hematopoietic prostaglandin D synthase  
25 and DP receptor in hereditary demyelinating disease

Changes in mRNA of H-PGDS and DP receptor as a result of demyelination were quantified by a quantitative RT-PCR method using a model mouse Twitcher of human Krabbe diseases which is a disease where galactosylceramidase is deficient (Kobayashi

T, et al., *Brain Res.*, 202:479-483, 1980; Duchen LW, et al., *Brain*, 103:695-710, 1980; Sakai N, et al., *J. Neurochem.*, 66:1118-1124, 1996; Taniike M et al., *J. Neuropathol. Exp. Neurol.*, 58:644-653, 1999) (refer to Fig. 1). Expression of  
5 mRNA of both H-PGDS and DP receptor increased as a result of demyelination.

It was identified by an immunohistochemical staining that H-PGDS was expressed in microglia cells, macrophage cells and ameboid cells accumulated in local tissues where  
10 demyelination progressed (refer to Fig. 2). On the other hand, it was identified that DP receptor was expressed in activated astroglia cells distributed around the tissues where demyelination progressed (refer to Fig. 3).

#### Example 2

15 Transduction of hematopoietic prostaglandin D synthase and DP receptor in autoimmune demyelinating diseases

In an experimental autoimmune encephalomyelitis mouse which is a model of human multiple sclerosis (Ichikawa M., et al., *Cell Immunol.*, 191:97-104, 1999; Bernhard Hemmer, et al.,  
20 *Nature Review Neuroscience*, 3:291-301, 2002), expression of both H-PGDS and DP receptor as measured by a quantitative RT-PCR method also showed an increase correlated to the extent of demyelination (refer to Fig. 4).

In an observation by an immunohistochemistry, H-PGDS was  
25 expressed in microglia cells, macrophage and ameboid cells accumulated in local tissues where demyelination progressed (refer to Fig. 5).

#### Example 3

Transduction of hematopoietic prostaglandin D synthase

and DP receptor in traumatic brain injury

As a result of investigation of expression of mRNA of H-PGDS and DP receptor in a brain injury using a model of traumatic injury of the cerebral cortex (Stab would) (Salhia B, et al., *Brain Res.*, 888:87-97, 2000; Asahi M., et al., *J. Neurosci.*, 21:7724-7732, 2001; Garcia de Yebenes E., et al., *J. Neurochem.*, 73:812-820, 1999), H-PGDS showed its peaked at two days from injury (refer to Fig. 6) while DP receptor continuously increased from the second to the eighth days (refer to Fig. 7).

After 24 hours from the injury, transduction of H-PGDS took place in macrophage and microglia cells accumulated around the injured site (refer to Fig. 8 and Fig. 9) while, in astroglia cells around the injured site, expression of GFAP and DP receptor was enhanced and those phenomena continued until 8 days after being injured (refer to Fig. 10 and Fig. 11).

Degree of injury was quantified by a dye leakage (Evans Blue) to the injured site (Kakimura Y, et al., *Nature Medicine*, 4:1078-1080, 1998). After 2 days from injury, the maximum value was noted and, after that, it lowered as a result of recovery (refer to Fig. 20).

Example 4Suppression of activation of astroglia cells in hereditary demyelinating disease by administration of inhibitor for hematopoietic prostaglandin D synthase

HQL-79 (4-benzhydryloxy-1-{3-(1H-tetrazol-5-yl)-propyl}piperidine) which is an H-PGDS inhibitor was subcutaneously administered to Twitcher mice at the dose of 30 mg/kg/day for 14 days whereupon activation of astroglia cells was suppressed and, at the same time, expression of DP receptor

in astroglia cells lowered (refer to Fig. 13).

Example 5

Promotion of recovery of brain injury and suppression of transduction of DP receptor in traumatic brain injury by administration of inhibitor for hematopoietic prostaglandin D synthase

HQL-79, which is an H-PGDS inhibitor was orally administered to mice at the dose of 30 mg/kg/day for 4 days from one hour before injuring whereupon the amount of mRNA of DP receptor in the tissue injury region in a Stab wound model lowered (refer to Fig. 14) and promotion of recovery from a brain injury was noted (refer to Fig. 15). Such a therapeutic effect was also confirmed by an experiment using a leakage reaction of dye to the injured site as an index (refer to Fig. 21).

It was further confirmed that, even when administration of an H-PGDS inhibitor was initiated after an injury was applied, expansion of the injury was suppressed (refer to Fig. 22 and Fig. 23).

Example 6

Reduction in traumatic brain injury by administration of antagonist for prostaglandin D receptor

When BW A868C ((±)-3-benzyl-5-(6-carboxyhexyl)-1-(2-cyclohexyl-2-hydroxyethylamino)hydantoin), which is a DP receptor antagonist, was intravenously administered to Stab wound model mice at the dose of 1 mg/kg/day for 4 days from the initial day of injury, promotion of recovery from a brain injury was noted and activation of astroglia around the tissue injury site (refer to Fig. 16).

An effect of BW A868C or ramatroban ((±)-(3R)-3-(4-

fluorobenzenesulfonamide)-1,2,3,4-tetrahydrocarbazol-9-propionic acid), which is an antagonist for prostaglandin D receptor, to Stab wound model mice was evaluated using leakage amount of dye to the injured site as an index. Thus, Evans Blue dye was intravenously administered after 2 days from occurrence of a brain injury and the amount of leakage of the dye to the tissue during 2 hours thereafter was measured. When BW A868C (1 mg/kg) was intravenously administered after 3 hours and after one day from occurrence of the brain injury, leakage of the dye to the injured site was suppressed (refer to Fig. 27).

When ramatroban was orally administered (30 mg/kg) after 3 hours and after one day from the occurrence, leakage of the dye to the injured site was suppressed (refer to Fig. 27).

#### Example 7

An effect of pinagladin, which is a DP receptor antagonist, to a Stab wound model was evaluated by a leakage amount of dye to the injured site. Thus, Evans Blue was intravenously injected on the second day after occurrence of a brain injury and leakage amount of the dye into the tissue during 2 hours thereafter was measured. As a result, when pinagladin (10 mg/kg) was orally administered 1 hour before and one day after

the occurrence of the injury, an increase in leakage amount of Evans Blue dye noted as a result of brain injury was suppressed (refer to Fig. 24). In a further histopathological investigation, pinagladin suppressed the progress of brain injury in a Stab wound model in any of the administration routes (refer to Fig. 25).

#### Example 8

Exacerbation of traumatic brain injury by human

hematopoietic prostaglandin D synthase over-expression

In a Stab wound model using transgenic mice where human H-PGDS was expressed in large amounts prepared in Preparation Example 1, accumulation of macrophage in an injured site and  
5 activation of astroglia cells which was immunohistochemically tested were significant as compared with those in mild-type mice whereby recovery was delayed (refer to Fig. 12).

Example 9

Reduction in traumatic brain injury by hematopoietic  
10 prostaglandin D synthase gene deficiency

In a Stab wound model using hematopoietic prostaglandin D synthase knockout (H-PGDS KO) mice (homozygote) prepared in Preparation Example 2, bleeding in the injured site, activation of astroglia histoimmunochemically, checked using an anti-GFAP  
15 antibody, and dye leakage at the injured site were slight as compared with those in mice of a wild type (refer to Fig. 17 and Fig. 26).

On the other hand, in a Stab wound model using DP receptor knockout (DPR KO) mice (Matsuoka T, et al., *Science*,  
20 17;287(5460):2013-2017, 2000), leakage of the dye at the injured site was slight as compared with that in mice of a wild type.

When the inhibitor for hematopoietic prostaglandin D  
25 synthase (H-PGDS) and/or the antagonist for prostaglandin D receptor according to the present invention are/is used for the treatment, they/it are/is made into a pharmaceutical composition for oral or parenteral administration. A pharmaceutical composition comprising the inhibitor for hematopoietic

prostaglandin D synthase (H-PGDS) and/or the antagonist for prostaglandin D receptor according to the present invention may be in oral and parenteral dosage forms.

Thus, they may be made into formulation for oral  
5 administration such as tablets, capsules, granules, powder and syrup or into formulation for parenteral administration such as injection solution or suspension for intravenous injection, intramuscular injection, subcutaneous injection, etc., inhalant, eye drop, nose drop, suppository, preparation for  
10 percutaneous administration and percutaneous absorption such as ointment, poultice and cataplasm. Preferably, agents for oral administration or drugs for injection are used.

Those formulations can be manufactured using appropriate carriers, excipients, solvents, substrates, etc. which have been  
15 known by persons skilled in the art. In the case of tablets, for example, active ingredient and auxiliary components are compressed or molded together. As the auxiliary components, there may be used pharmaceutically acceptable excipients such as binder (e.g., corn starch), filler (e.g., lactose and  
20 microcrystalline cellulose), disintegrating agent (e.g., sodium starch glycolate) and/or lubricant (e.g., magnesium stearate). Tablets may be appropriately coated. In the case of liquid preparations such as syrup, liquid and suspension, there may be used suspending agent (e.g., methyl cellulose),  
25 emulsifier (e.g., lecithin), preservative, etc. In the case of the formulation for injection, any of the forms of solution, suspension and oily or aqueous emulsion may be used and they may contain, for example, a dispersing agent or a stabilizer for suspension. In the case of the formulation for percutaneous

administration and percutaneous absorption such as ointment, poultice and cataplasm, the formulation is prepared using an aqueous substrate (water, lower alcohol, polyol) or an oily substrate (higher fatty acid esters (isopropyl myristate), lipophilic alcohol).

Although the dose of the hematopoietic prostaglandin D synthase (H-PGDS) inhibitor and/or the prostaglandin D receptor antagonist to be administered according to the present invention may vary depending upon dosage form, symptom, age, body weight or sex of a patient or drug(s) used together (if any) and are/is finally subjected to the decision of medical doctors, it is administered, in the case of oral administration, in a dose of 0.01 to 100 mg, preferably 0.01 to 50 mg or, more preferably, 0.01 to 30 mg per kg of body weight while, in the case of parenteral administration, in a dose of 0.001 to 100 mg, preferably 0.001 to 5 mg or, more preferably, 0.001 to 3 mg per kg of body weight. Dosage may be administered by dividing into one to four time(s).

#### Formulation Examples

##### 20 Formulation 1

A tablet containing the following components was prepared.

Compound represented by the formula (I)	10 mg
Lactose	90 mg
25 Microcrystalline cellulose	30 mg
CMC-Na	15 mg
Magnesium stearate	5 mg
	150 mg

The compound represented by the formula (I), lactose,

microcrystalline cellulose and CMC-Na (carboxymethyl cellulose sodium salt) were passed through a sieve of 60 meshes and mixed. Magnesium stearate was mixed with the above mixture to give mixed powder for the manufacture of tablets. The mixed powder was directly compressed to give tablets each weighing 150 mg.

### Formulation 2

A suspension containing 50 mg of the active ingredient was prepared as follows:

10	Compound represented by the formula (I)	50 mg
	Carboxymethyl cellulose sodium	50 mg
	Syrup	1.25 ml
	Benzoic acid solution	0.10 ml
	Perfume	q.v.
15	Dye	q.v.
	Pure water was added to make	5 ml

The active ingredient was passed through a sieve of No. 45 mesh U. S. and mixed with carboxymethyl cellulose sodium and syrup to give a smooth paste. Then benzoic acid solution and perfume were diluted with a part of water and added thereto followed by stirring. After that, sufficient amount of water was added to give a necessary volume.

### Formulation 3

Formulation for intravenous administration was manufactured as follows.

	Compound represented by the formula (I)	100 mg
	Saturated fatty acid glyceride	1000 ml

Usually, a solution comprising the above components is intravenously administered to a patient at the rate of 1 ml per

minute.

Formulation 4

A gelatin hard capsule preparation of the following composition was prepared by a conventional method.

5	HQL-79	10 mg
	Starch	50 mg
	Magnesium stearate	10 mg

Formulation 5

10 The following tablet was prepared by a conventional method.

	HQL-79	10 mg
	Cellulose, microcrystalline	500 mg
	Silicon dioxide	10 mg
	Magnesium stearate	10 mg

15

As fully illustrated hereinabove, the composition of the present invention is able to be used for the treatment of diseases such as cerebrovascular disorder, brain degenerative disease and demyelinating disease.

20 When H-PGDS transduced by microglia cells or macrophage at the part affected by brain injury is inhibited or when activity of DP receptor expressed in astroglia cells around the injured area is inhibited, it is now possible to treat or prevent the brain injury in which prostaglandin D<sub>2</sub> participates.

25 In patients suffering from multiple sclerosis, biosynthesis of prostaglandin D<sub>2</sub> is active (Science 294:1731-11735, 2001) and there is a high possibility that it acts as an exacerbating factor during the onset step of experimental autoimmune encephalomyelitis in mice which is a

model thereof. Therefore, when a specific inhibitor for H-PGDS  
or an antagonist for receptor is used, it is able to be a useful  
treating method for treatment or prevention of multiple  
sclerosis as a substitute for steroid therapy and  
5 immunosuppressants which have been conducted/used at present.  
In addition, such a drug suppresses a local inflammation  
reaction as a result of autoimmune cell disorder and nerve cell  
death whereby it is also able to be applied for the treatment  
of intractable diseases accompanied by neurofibrillary  
10 degeneration including Alzheimer's disease.

## CLAIMS

1. A pharmaceutical composition used for treatment or prevention of brain injury comprising an inhibitor for hematopoietic prostaglandin D synthase (H-PGDS) as the  
5 active ingredient in admixture with a carrier or diluent.

2. The pharmaceutical composition according to claim 1, wherein the H-PGDS inhibitor is 4-benzhydryloxy-1-  
{3-(1H-tetrazol-5-yl)-propyl}piperidine, 1-amino-4-{4-[4-  
chloro-6-(2-sulfo-phenylamino)-[1,3,5]-triazine-2-ylmethyl-  
10 3-sulfo-phenylamino]-9,10-dioxo-9,10-dihydro-anthracene-2-  
sulfonic acid, 1-amino-4-(4-sulfamoylanilino)-  
anthraquinone-2-sulfonic acid or pharmaceutically  
acceptable salt thereof or hydrate thereof or  
2-(2'-benzothiazolyl)-5-styryl-3-(4'-phthalhydrazidyl)  
15 -tetrazolium chloride or a hydrate thereof.

3. Use of a hematopoietic prostaglandin D synthase (H-PGDS) inhibitor for treatment of a brain injury.

4. Use of a hematopoietic prostaglandin D synthase (H-PGDS) inhibitor for the manufacture of a drug for  
20 treatment of a brain injury.

5. A pharmaceutical composition to be used for treatment or prevention of a brain injury comprising a hematopoietic prostaglandin D synthase (H-PGDS) inhibitor and a prostaglandin D receptor antagonist the effective  
25 ingredients in admixture with a carrier or diluent.

Application number/numéro de demande: JP 2003 / 008904

Figures: 2, 3, 5, 8, 9, 10, 11, 12, 13, 15, 16, 17, 22, 25 <sup>20, 23, 26</sup>


Pages: \_\_\_\_\_

DRW - IP

Unscannable items  
received with this application  
(Request original documents in File Prep. Section on the 10th Floor)


Documents reçus avec cette demande ne pouvant être balayés  
(Commander les documents originaux dans la section de préparation des dossiers au  
10ième étage)

Fig. 1




\* : p < 0.05

Fig. 4



6/26

Fig. 6



7/26

Fig. 7




Fig. 14




Fig. 18





Fig. 19



F i g . 2 0




Fig. 24




Fig. 27

