



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2689633 A1 2008/12/11

(21) **2 689 633**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2008/03/24
(87) Date publication PCT/PCT Publication Date: 2008/12/11
(85) Entrée phase nationale/National Entry: 2009/12/04
(86) N° demande PCT/PCT Application No.: US 2008/058063
(87) N° publication PCT/PCT Publication No.: 2008/150565
(30) Priorité/Priority: 2007/05/31 (US60/941,210)

(51) Cl.Int./Int.Cl. *A61K 31/495* (2006.01),
A61P 9/10 (2006.01), *G01N 33/68* (2006.01)

(71) Demandeur/Applicant:
CV THERAPEUTICS, INC., US

(72) Inventeurs/Inventors:
BLACKBURN, BRENT, US;
JERLING, MARCUS, SE;
WOLFF, ANDREW, US

(74) Agent: SMART & BIGGAR

(54) Titre : UTILISATION DE RANOLAZINE POUR PEPTIDE CEREBRAL NATRIURETIQUE ELEVE
(54) Title: USE OF RANOLAZINE FOR ELEVATED BRAIN-TYPE NATRIURETIC PEPTIDE

(57) **Abrégé/Abstract:**

This invention is directed to the use of ranolazine to reduce the risk of adverse coronary events in a mammalian patient. Typically, the natriuretic peptide is associated with coronary disease, acute coronary syndrome, and/or diastolic dysfunction. Ranolazine may be administered to the patient as an intravenous solution or in an oral dose.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



PCT



(43) International Publication Date
11 December 2008 (11.12.2008)

(10) International Publication Number
WO 2008/150565 A3

(51) International Patent Classification:

A61K 31/495 (2006.01) **G01N 33/68** (2006.01)
A61P 9/10 (2006.01)

(21) International Application Number:

PCT/US2008/058063

(22) International Filing Date: 24 March 2008 (24.03.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/941,210 31 May 2007 (31.05.2007) US

(71) Applicant (*for all designated States except US*): **CV THERAPEUTICS, INC.** [US/US]; 3172 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BLACKBURN, Brent** [US/US]; 633 Almond Avenue, Los Altos, CA 94022 (US). **JERLING, Marcus** [SE/SE]; Stavgardsgatan 30, S-167 56 Bromma (SE). **WOLFF, Andrew** [US/US]; 2140 Taylor Street #803, San Francisco, CA 94133 (US).

(74) Agents: **HARTRUM, J., Elin** et al.; Cv Therapeutics, Inc., 3172 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

(88) Date of publication of the international search report:
2 April 2009

(54) Title: RANOLAZINE FOR ELEVATED BRAIN-TYPE NATRIURETIC PEPTIDE

(57) Abstract: This invention is directed to the use of ranolazine to reduce the risk of adverse coronary events in a mammalian patient. Typically, the natriuretic peptide is associated with coronary disease, acute coronary syndrome, and/or diastolic dysfunction. Ranolazine may be administered to the patient as an intravenous solution or in an oral dose.



WO 2008/150565 A3

USE OF RANOLAZINE FOR ELEVATED BRAIN-TYPE NATRIURETIC PEPTIDE

[0001] This invention claims priority to U.S. Provisional Patent Application Serial No. 60/941,210, filed May 31, 2007, the entirety of which is incorporated herein by reference.

FIELD OF THE INVENTION


[0002] This invention is directed to identifying the level of natriuretic peptides, such as brain-type natriuretic peptide (BNP) and N-terminal pro-brain natriuretic peptide (NT-proBNP), in the blood plasma of a mammalian patient and, if the level of BNP is elevated, administering ranolazine to the patient. The invention is also directed to methods for reducing the risk of adverse coronary events including, but not limited to, coronary artery disease (CAD), cardiovascular death, myocardial infarction, acute heart failure, ischemia, recurrent ischemia, acute coronary syndrome, diastolic dysfunction, and the like in patients with a plasma BNP of greater than about 80 pg/mL comprising administration of ranolazine.

BACKGROUND OF THE INVENTION

[0003] The natriuretic peptide system consists of structurally similar natriuretic peptides that compensate for changes in volume and pressure by promoting the removal of sodium ions from the blood. Elevated levels of arterial natriuretic peptide (ANP) and BNP may be found in patients with coronary disease. Clinical studies have indicated that the testing for elevated levels of BNP or NT-proBNP facilitates the diagnosis of heart failure and is validated by biomarker for increased risk. Persson, et al. "Diastolic Dysfunction in Heart Failure with Preserved Systolic Function: Need for Object Evidence" *JACC*, 49(6):687-694 (2007); Bibbins-Domingo et al. "N-Terminal Fragment of the Prohormone Brain-Type Natriuretic Peptide (NT-proBNP), Cardiovascular Events, and Mortality in Patients with Stable Coronary Heart Disease" *JAMA*, 297(2):169-176(2007).

[0004] Ranolazine, a late I_{Na} inhibitor, currently available in the U.S. under the trade name Ranexa® for angina, is suitable for delivery in the form of an oral dose, such as a compressed tablet, or an intravenous dose.

[0005] U.S. Patent No. 4,567,264, the specification of which is incorporated herein by reference in its entirety, discloses ranolazine, ($\pm$)-N-(2,6-dimethylphenyl)-4-[2-hydroxy-3-(2-methoxyphenoxy)-propyl]-1-piperazineacetamide, and its pharmaceutically acceptable salts, and their use in the treatment of cardiovascular diseases, including arrhythmias, variant and exercise-induced angina, and myocardial infarction. In its dihydrochloride salt form, ranolazine is represented by the formula:



[0006] U.S. Patent No. 5,506,229, which is incorporated herein by reference in its entirety, discloses the use of ranolazine and its pharmaceutically acceptable salts and esters for the treatment of tissues experiencing a physical or chemical insult, including cardioplegia, hypoxic or reperfusion injury to cardiac or skeletal muscle or brain tissue, and for use in transplants. Oral and parenteral formulations are disclosed, including controlled release formulations. In particular, Example 7D of U.S. Patent No. 5,506,229 describes a controlled release formulation in capsule form comprising microspheres of ranolazine and microcrystalline cellulose coated with release controlling polymers. This patent also discloses IV ranolazine formulations which at the low end comprise 5 mg ranolazine per milliliter of an IV solution containing about 5% by weight dextrose.

[0007] The presently preferred route of administration for ranolazine and its pharmaceutically acceptable salts is oral. A typical oral dosage form is a compressed tablet, a hard gelatin capsule filled with a powder mix or granulate, or a soft gelatin capsule (softgel) filled with a solution or suspension. U.S. Patent No. 5,472,707, the specification of which is incorporated herein by reference in its entirety, discloses a high-dose oral formulation employing supercooled liquid ranolazine as a fill solution for a hard gelatin capsule or softgel.

[0008] U.S. Patent Application Publication Number 2006/0177502, the specification of which is incorporated herein by reference in its entirety, discloses oral sustained release dosage forms in which the ranolazine is present in 35-50%, preferably 40-45% ranolazine. In one embodiment the ranolazine sustained release formulations of the invention include a pH dependent binder; a pH independent binder; and one or more pharmaceutically acceptable excipients. Suitable pH dependent binders include, but are not limited to, a methacrylic acid copolymer, for example Eudragit® (Eudragit® L100-55, pseudolatex of Eudragit® L100-55, and the like) partially neutralized with a strong base, for example, sodium hydroxide, potassium hydroxide, or ammonium hydroxide, in a quantity sufficient to neutralize the methacrylic acid copolymer to an extent of about 1-20%, for example about 3-6%. Suitable pH independent binders include, but are not limited to, hydroxypropylmethylcellulose (HPMC), for example Methocel® E10M Premium CR grade HPMC or Methocel® E4M Premium HPMC. Suitable pharmaceutically acceptable excipients include magnesium stearate and microcrystalline cellulose (Avicel® pH101).

SUMMARY OF THE INVENTION

[0009] It has been discovered that ranolazine, a late I_{Na} current inhibitor, is capable of reducing the risk of an adverse coronary event, including coronary artery disease (CAD), cardiovascular death, myocardial infarction, acute heart failure, ischemia, recurrent ischemia, acute coronary syndrome, diastolic dysfunction, and the like, in a patient exhibiting elevated levels of natriuretic peptides. By identifying the level of BNP in a patient suffering from a coronary disease, and, if the level of BNP is about 80 pg/mL or greater, it has been found that administering ranolazine reduces the risk of occurrence of an adverse coronary event in such a patient. Further, by administering ranolazine to a patient with a BNP level of about 80 pg/mL or greater, it has been found that adverse coronary events related to diastolic dysfunction and/or acute coronary syndrome are reduced.

[0010] One embodiment of the invention is directed to a method of reducing the risk of an adverse coronary event, including, but not limited to, coronary artery disease (CAD), cardiovascular death, myocardial infarction, acute heart failure, ischemia, recurrent ischemia, acute coronary syndrome, diastolic dysfunction, and the like in a patient exhibiting elevated levels of natriuretic peptides in blood plasma and/or heart tissue, the

method comprising identifying a patient exhibiting elevated levels of a natriuretic peptide; and administering the patient a therapeutically effective amount of ranolazine. The ranolazine may be administered as an oral dose or as an IV solution. In some cases, the patient may be administered an IV solution followed by an oral dose. Preferably, when administered as an oral dose, ranolazine is administered in a sustained release tablet.

[0011] In one embodiment, the elevated levels of natriuretic peptide are correlated with higher risks of adverse coronary events in a patient arising from diseases including, but not limited to, coronary artery disease (CAD), cardiovascular death, myocardial infarction, acute heart failure, ischemia, recurrent ischemia, acute coronary syndrome, diastolic dysfunction, and the like.

[0012] In one embodiment, the natriuretic peptide is BNP or NT-proBNP and the elevated levels are about 80 picograms or greater of natriuretic peptide per milliliter of blood.

[0013] In another embodiment, the invention is directed to kits of parts comprising an assay that detects levels of natriuretic peptides in blood plasma and/or heart tissue of a mammalian patient and a pharmaceutical dosage of ranolazine. The pharmaceutical dosage may in the form of an IV solution and/or an oral dose. The oral dose may be in the form of one or more sustained release tablets.

DESCRIPTION OF THE DRAWINGS

[0014] Figure 1 shows the cumulative incidence (%) of cardiovascular death, myocardial infarction, or recurrent ischemia for patients with positive BNP on placebo (curve A), with positive BNP on ranolazine (curve B), or negative BNP and placebo or ranolazine administered (2 curves C) versus days from randomization.

DETAILED DESCRIPTION OF THE INVENTION

[0015] As noted above, this invention relates to methods of reducing the risk of an adverse coronary event, including, but not limited to, coronary artery disease (CAD), cardiovascular death, myocardial infarction, acute heart failure, ischemia, recurrent ischemia, acute coronary syndrome, diastolic dysfunction, and the like in a patient exhibiting elevated levels of natriuretic peptides comprising administering ranolazine to

these patients. However, prior to describing this invention in more detail, the following terms will first be defined.

1. Definitions

[0016] In this specification and in the claims that follow, reference will be made to a number of terms that shall be defined to have the following meanings.

[0017] The term “natriuretic peptide” refers to peptides that causes natriuresis, the excretion of an excessively large amount of sodium in the urine. The natriuretic peptides are produced by the heart and vasculature:

A-type natriuretic peptide (or “ANP”) is secreted largely by the atrial myocardium in response to dilatation.

B-type natriuretic peptide (or “BNP”) is produced mainly by the ventricular myocardium.

C-type natriuretic peptide (or “CNP”) is produced by endothelial cells that line the blood vessels.

[0018] Preferably the natriuretic peptide is “BNP” and “NT-proBNP”. Both arise from a 108-amino acid pro form synthesized in the ventricles. In response to pressure overload, the pro form is cleaved into BNP (32 amino acids) and NT-proBNP (78 amino acids). Once released into the circulation, BNP plays a homeostatic role as a systemic vasodilator and diuretic. NT-proBNP appears to be a by-product. Of special interest for laboratory medicine, BNP has a shorter half-life than NT-proBNP. Thus, BNP fluctuates on a faster timescale, while NT-proBNP remains more stable in blood samples. Levels of BNP and NT-proBNP may be tested using commercially available assays as discussed below.

[0019] “Elevated levels of natriuretic peptide” refers to a concentration of endogenous natriuretic peptide that is predictive of a future adverse coronary event in the patient. For example, in the case of BNP, elevated levels refers to about 80 picograms (pg) or greater per milliliter of blood, preferably, 100 picograms or greater of natriuretic peptide per milliliter of blood. In another instance, the BNP concentration is between about 100 and about 300 picograms of natriuretic peptide per milliliter of blood. In another instance, the BNP concentration is between about 300 and about 600 picograms of natriuretic peptide per milliliter of blood. In another instance, the BNP concentration is between about 600 and about 900 picograms of natriuretic peptide per

milliliter of blood. In another instance, the BNP concentration is greater than about 900 picograms of natriuretic peptide per milliliter of blood. It has been found that identifying an elevated level of natriuretic peptide in a coronary patient and administering ranolazine to the patient reduces the risk of a future adverse coronary event in the patient.

[0020] An “adverse coronary event” or “adverse cardiac event” refers to a non-incremental decrease in the coronary performance of the patient due to a coronary or cardiac disease which event may involve either morbidity or hospitalization. Such adverse coronary/cardiac events include, by way of example, coronary artery disease (“CAD”), cardiovascular death (CD), myocardial infarction (MI), acute heart failure, ischemia, recurrent ischemia (RI), acute coronary syndrome (ACS), diastolic dysfunction and the like.

[0021] “Ranolazine” is the compound ($\pm$)-N-(2,6-dimethylphenyl)-4-[2-hydroxy-3-(2-methoxyphenoxy)propyl]-1-piperazine-acetamide, and its pharmaceutically acceptable salts, and mixtures thereof. Unless otherwise stated the ranolazine plasma concentrations used in the specification and examples refer to ranolazine free base. At pH of about 4, in an aqueous solution titrated with hydrogen chloride, ranolazine will be present in large part as its dihydrochloride salt.

[0022] “Physiologically acceptable pH” refers to the pH of an intravenous solution which is compatible for delivery into a human patient. Preferably, physiologically acceptable pH’s range is from about 4 to about 8.5 and preferably from about 4 to 7. Without being limited by any theory, the use of intravenous solutions having a pH of about 4 to 6 are deemed physiologically acceptable as the large volume of blood in the body effectively buffers these intravenous solutions.

[0023] “Coronary diseases” or “cardiovascular diseases” refer to diseases of the cardiovascular system arising from any one or more than one of, for example, heart failure, including congestive heart failure, acute heart failure, ischemia, recurrent ischemia, myocardial infarction, arrhythmias, angina (including exercise-induced angina, variant angina, stable angina, unstable angina), acute coronary syndrome, diabetes, and intermittent claudication. The treatment of such disease states is disclosed in various U.S. patents and patent applications, including U.S. Patent Nos. 6,503,911 and 6,528,511, U.S. Patent Application Serial Nos. 2003/0220344 and 2004/0063717, the complete disclosures of which are hereby incorporated by reference.

[0024] “Acute coronary syndrome” or “ACS” refers to a range of acute myocardial ischemic states. It encompasses unstable angina and non-ST-segment elevation myocardial infarction (UA/NSTEMI), and ST segment elevation myocardial infarction (STEMI).

[0025] The term “diastolic dysfunction” refers to and is characterized by an increase in diastolic filling pressure, which may be responsible for the occurrence of dyspnea. This symptom may occur during exercise ("latent" diastolic dysfunction) or may be present also at rest ("manifest" diastolic dysfunction).

[0026] “Optional” and “optionally” mean that the subsequently described event or circumstance may or may not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "optional pharmaceutical excipients" indicates that a formulation so described may or may not include pharmaceutical excipients other than those specifically stated to be present, and that the formulation so described includes instances in which the optional excipients are present and instances in which they are not.

[0027] "Treating" and "treatment" refer to any treatment of a disease in a patient and include: preventing the disease from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it; inhibiting the disease, i.e., arresting its further development; inhibiting the symptoms of the disease; relieving the disease, i.e., causing regression of the disease, or relieving the symptoms of the disease. The “patient” is a mammal, preferably a human.

[0028] The term "therapeutically effective amount" refers to that amount of ranolazine that is sufficient to effect treatment, as defined above, when administered to a mammal in need of such treatment. The therapeutically effective amount will vary depending upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art.

[0029] “Immediate release” (“IR”) refers to formulations or dosage units that rapidly dissolve *in vitro* and are intended to be completely dissolved and absorbed in the stomach or upper gastrointestinal tract. Conventionally, such formulations release at least 90% of the active ingredient within 30 minutes of administration.

[0030] “Sustained release” (“SR”) refers to formulations or dosage units used herein that are slowly and continuously dissolved and absorbed in the stomach and

gastrointestinal tract over a period of about six hours or more. Preferred sustained release formulations are those exhibiting plasma concentrations of ranolazine suitable for no more than twice daily administration with two or less tablets per dosing as described below.

[0031] "Intravenous (IV) infusion" or "intravenous administration" refers to solutions or dosage units used herein that are provided to the patient by intravenous route. Such IV infusions can be provided to the patient until for up to about 96 hours in order to stabilize the patient's cardiovascular condition. The method and timing for delivery of an IV infusion is within the skill of the attending medically trained person.

2. Methods of the Invention

[0032] In patients with a coronary disease, such as heart failure, it is believed that increased cardiac volume and increased pressure in the heart result in increased levels of natriuretic peptides. As stated above, the invention is directed to a method of identifying the level of natriuretic peptides, such as brain-type natriuretic peptide (BNP) and N-terminal pro-brain natriuretic peptide (NT-proBNP), in the blood plasma of a mammalian patient and, if the level of BNP is elevated, administering ranolazine to the patient. The invention is also directed to methods for reducing the risk of adverse coronary events including, but not limited to, coronary artery disease (CAD), cardiovascular death, myocardial infarction, acute heart failure, ischemia, recurrent ischemia, acute coronary syndrome, diastolic dysfunction, and the like in patients with a plasma BNP of about 80 pg/mL or greater comprising administration of ranolazine.

[0033] In the first step of the invention, a patient is identified that has elevated levels of a natriuretic peptide. Preferably, the levels of BNP and/or NT-proBNP is tested. Preferably, levels of NT-proBNP is tested as it tends to have a longer plasma half life and better *in vitro* stability than the chemically active BNP.

[0034] The testing of the peptides can take place in the hospital, the doctor's office, or other suitable environment as is typical in the industry for care of patients. Typically, a patient is asked to rest for 10 to 15 minutes prior to drawing between 5 cc and 25 cc of blood. For most natriuretic peptide testing, the blood is placed in a tube with a chelating agent, such as EDTA, and then centrifuged at below room temperatures. The blood plasma is then separated and run through an immunoassay. The assay provides data relating to the concentration of natriuretic peptide in the blood plasma.

[0035] A patient is suitable for treatment by the methods of this invention if the patient exhibits elevated levels of natriuretic peptide as defined above. For heart failure typically, the following schedule dictates the severity of the patient's cardiac condition:

- BNP levels below 100 pg/mL indicate little or no risk of heart failure.
- BNP levels of 100-300 pg/mL suggest a risk of heart failure is present.
- BNP levels above 300 pg/mL indicate a risk of mild heart failure.
- BNP levels above 600 pg/mL indicate a risk of moderate heart failure.
- BNP levels above 900 pg/mL indicate a risk of severe heart failure.

[0036] There are several tests available to one of skill in the art to measure these levels. For example, Elecsys® proBNP (Roche, Basel, Switzerland) test can rapidly test for BNP levels. Another example of the assay is the ADVIA® Centaur BNP Assay (Bayer Diagnostics Corporation, Tarrytown, New York), which can rapidly test for BNP levels. Yet another example of the BNP assay is the Triage® BNP Test (Biosite Inc., San Diego, California).

[0037] After the patient is identified to have elevated natriuretic peptide levels, the clinician can then administer ranolazine according to the discussion below.

3. Compositions of the Invention

[0038] As mentioned above, the methods of this invention employ either an IV solution of ranolazine and/or an oral dose of ranolazine. Both of these formulations are discussed below.

Oral Dose

[0039] In one embodiment, the ranolazine is administered an oral dose. In one embodiment, an oral formulation of ranolazine is a tablet. The tablet can be formulated as an instant release or a sustained release tablet. In one embodiment, the tablet of ranolazine is up to about 750 mg ranolazine. In another embodiment, the tablet of ranolazine is up to about 1000 mg ranolazine. In a preferred embodiment, the ranolazine tablet is 375 mg ranolazine, and/or 500 mg ranolazine. In another preferred embodiment, the ranolazine tablet is about 250 mg ranolazine.

[0040] The oral formulation of ranolazine is thoroughly discussed in U.S. Patent No. 6,303,607 and U.S. Publication No. 2003/0220344, which are both incorporated herein by reference in their entirety.

[0041] The oral sustained release ranolazine dosage formulations of this invention are administered one, twice, or three times in a 24 hour period in order to maintain a plasma ranolazine level above the threshold therapeutic level and below the maximally tolerated levels, which is preferably a plasma level of about 550 to about 7500 ng base/mL in a patient.

[0042] In a preferred embodiment, the plasma level of ranolazine ranges about 1500 to about 3500 ng base/mL in a patient.

[0043] In order to achieve the preferred plasma ranolazine level, it is preferred that the oral ranolazine dosage forms described herein are administered once or twice daily. If the dosage forms are administered twice daily, then it is preferred that the oral ranolazine dosage forms are administered at about twelve hour intervals.

[0044] In addition to formulating and administering oral sustained release dosage forms of this invention in a manner that controls the plasma ranolazine levels, it is also important to minimize the difference between peak and trough plasma ranolazine levels. The peak plasma ranolazine levels are typically achieved at from about 30 minutes to eight hours or more after initially ingesting the dosage form while trough plasma ranolazine levels are achieved at about the time of ingestion of the next scheduled dosage form. It is preferred that the sustained release dosage forms of this invention are administered in a manner that allows for a peak ranolazine level no more than 8 times greater than the trough ranolazine level, preferably no more than 4 times greater than the trough ranolazine level, preferably no more than 3 times greater than the trough ranolazine level, and most preferably no greater than 2 times trough ranolazine level.

[0045] The sustained release ranolazine formulations of this invention provide the therapeutic advantage of minimizing variations in ranolazine plasma concentration while permitting, at most, twice-daily administration. The formulation may be administered alone, or (at least initially) in combination with an immediate release formulation if rapid achievement of a therapeutically effective plasma concentration of ranolazine is desired or by soluble IV formulations and oral dosage forms.

Intravenous Solution

[0046] In one aspect, the invention employs an intravenous (IV) solution comprising a selected concentration of ranolazine. Specifically, the IV solution preferably comprises about 1.5 to about 3.0 mg of ranolazine per milliliter of a pharmaceutically acceptable aqueous solution, more preferably about 1.8 to about 2.2 mg and even more preferably about 2 mg. In order to allow for the rapid intravenous flow of ranolazine into the patient, the IV solution preferably contains no viscous components including by way of example as propylene glycol or polyethylene glycol (e.g., polyethylene glycol 400). It is understood that minor amounts of viscous components that do not materially alter the viscosity may be included in the intravenous formulations of this invention. In a particularly preferred embodiment, the viscosity of the IV solution is preferably less than 10 cSt (centistokes) at 20°C, more preferably less than 5 cSt at 20°C and even more preferably less than 2 cSt at 20°C.

[0047] In one embodiment, the IV solution comprises about 1.5 to about 3.0 mg of ranolazine per mL of IV solution; and either about 4.8 to about 5.0 weight percent dextrose or about 0.8 to about 1.0 weight percent sodium chloride.

[0048] In one embodiment, the IV solution comprises about 1.8 to about 2.2 mg of ranolazine per mL of IV solution; and either about 4.8 to about 5.0 weight percent dextrose or about 0.8 to about 1.0 weight percent sodium chloride.

[0049] In one embodiment, the IV solution of this invention comprises about 2 mg of ranolazine per mL of IV solution; and either about 4.8 to about 5.0 weight percent dextrose or about 0.9 weight percent sodium chloride.

[0050] The IV solutions described herein can be prepared from a stock solution comprising a 20 mL container for single use delivery which container comprises a sterile aqueous solution of ranolazine at a concentration of about 25 mg/mL; either about 36 mg/mL dextrose monohydrate or about 0.9 weight percent sodium chloride; and having a pH of about 4. Surprisingly, employing such high concentrations of ranolazine and dextrose monohydrate or ranolazine and sodium chloride in the stock solutions provide for compositions which are stable and have adequate shelf-lives, preferably of greater than 6 months.

[0051] Exemplary methods for preparing the stock solutions are described in Examples 1 and 2.

[0052] In a typical setting, two 20 mL containers described herein are injected into an IV container containing 460 mL of sterile saline (0.9 weight percent (w%) sodium chloride) or an aqueous dextrose solution (water containing 5 weight percent dextrose monohydrate) to provide for an IV solution of about 2 mg/mL of ranolazine maintained at physiologically acceptable pH. Containers useful herein include, but are not limited to, vials, syringes, bottles, IV bags, and the like.

[0053] In another embodiment, the intravenous formulation as set forth above, is diluted with a sterile diluent prior to use. In one embodiment, the sterile diluent is 5 % dextrose or a 0.9 weight percent saline solution. In one embodiment, the intravenous formulation is further diluted into bags of sterile diluent.

4. Kits of the Invention

[0054] One aspect of the invention is directed to a kit of parts comprising an assay to test for elevated levels of the natriuretic peptide and a pharmaceutical dosage of ranolazine.

[0055] The assay may be any of the assays discussed above. The pharmaceutical dosage of ranolazine may be either in the form of an IV solution or an oral dose or both. The IV solution can be packaged in a container as described above. The oral dose is preferably one or two compressed tablets as described above.

EXAMPLES

[0056] Unless stated otherwise, the following definitions are used in the examples.

μm	=	micrometers
cc	=	cubic centimeters
EDTA	=	Ethylene diamine tetraacetic acid
g	=	gram
mg	=	milligram
mL	=	milliliter
mm	=	millimeter
N	=	normal
rpm	=	revolutions per minute

EXAMPLE 1

Method of Preparing IV Solution of Ranolazine

[0057] 20-mL Type 1 flint vial of Ranolazine Injection filled to deliver 20 mL (at 1, 5, or 25 mg/mL ranolazine concentration).

Compositions

Ranolazine	1.0, 5.0, 25.0 mg/mL
Dextrose monohydrate	55.0, 52.0, 36.0 mg/mL
Hydrochloric acid	q.s. pH to 4.0 ± 0.2
Sodium hydroxide	q.s. pH to 4.0 ± 0.2
Water for Injection	q.s.

Container/Closure System

Vial:	Type 1 Flint, 20-cc, 20-mm finish
Stopper:	Rubber, 20-mm, West 4432/50, gray butyl, Teflon coated
Seal:	Aluminum, 20-mm, flip-top oversea

Method of Manufacture

[0058] The intravenous formulation of ranolazine is manufactured via an aseptic fill process as follows. In a suitable vessel, the required amount of dextrose monohydrate was dissolved in Water for Injection (WFI) at about 78% of the final batch weight. With continuous stirring, the required amount of ranolazine was added to the dextrose solution. To facilitate the dissolution of ranolazine, the solution pH was adjusted to a target of 3.88–3.92 with a 0.1 N or 1.0 N HCl solution. Additionally, 1 N NaOH may have been utilized to further adjust the solution to the target pH of 3.88–3.92. After ranolazine was dissolved, the batch was adjusted to the final weight with WFI. Upon confirmation that in-process specifications had been met, the ranolazine-formulated bulk solution was sterilized by sterile filtration through two 0.2 μ m sterile filters. Subsequently, the sterile ranolazine-formulated bulk solution was aseptically filled into sterile glass vials and aseptically stoppered with sterile stoppers. The stoppered vials were then sealed with clean flip-top aluminum overseals. The vials then went through a final inspection.

EXAMPLE 2

Method of Preparing IV Solution of Ranolazine

[0059] 20-mL Type 1 flint vial of Ranolazine Injection are filled to deliver 20 mL (25 mg/mL concentration).

Composition

Ranolazine	25.0 mg/mL
Dextrose monohydrate	36.0 mg/mL

Hydrochloric acid Adjust pH to 3.3–4.7

Water for Injection q.s.

Container/Closure System

Vial: Type 1 tubing, untreated, 20-mL, 20-mm finish

Stopper: Rubber, 20-mm, West 4432/50, gray butyl

Seal: Aluminum, 20-mm, blue flip-off overseal

Method of Manufacture

[0060] Water for Injection (WFI) is charged in a suitable vessel at about 90% of the final batch weight. About 90–95% of the required amount of 5 N HCl is added into the compounding vessel. With continuous stirring, the required amount of ranolazine is slowly added, followed by the addition of dextrose monohydrate into the ranolazine solution. To solubilize ranolazine, the solution pH is adjusted with 5 N HCl solution to a target of 3.9–4.1. The batch is subsequently adjusted to the final weight with WFI. Upon confirmation that in-process specifications have been met, the ranolazine-formulated bulk solution is sterilized by filtration through two redundant 0.22 μ m sterilizing filters. The sterile ranolazine-formulated bulk solution is then aseptically filled into 20 mL sterile/depyrogenated vials and aseptically stoppered with sterile/depyrogenated stoppers. The stoppered vials are sealed with clean flip-top aluminum overseals. The sealed vials are terminally sterilized by a validated terminal sterilization cycle at 121.1 °C for 30 minutes. After the terminal sterilization process, the vials go through an inspection. To protect the drug product from light, the vials are individually packaged into carton boxes.

EXAMPLE 3

Method of Preparing a Sustained Release Tablet

[0061] One sustained release formulation of ranolazine employed in this invention, includes a pH dependent binder and a pH independent binder. This formulation was prepared by combining Ranolazine (7500 g), Eudragit® L 100-55 (1000 g), hydroxypropyl methylcellulose (Methocel® E5-source) (200 g), and microcrystalline cellulose (Avicel®) (1060 g) by intimate mixing. The mixed powders were granulated with a solution of sodium hydroxide (40 g) in water (1900 to 2500 g). The granulate was dried and screened, mixed with magnesium stearate (200 g), and compressed for

example into tablets weighing 667 mg to achieve a dose of 500 mg of ranolazine free base per tablet. The tablets were spray coated in a 24 inch Accelacota® cylindrical pan coater with OPADRY film coating solution to a 2-4% weight gain. OPADRY film coating solutions are available in a variety of colors from Colorcon (West Point, PA).

[0062] The stepwise procedure for preparing this formulation is as follows:

- a) Blend together ranolazine, microcrystalline cellulose, methacrylate copolymer (Type C) and hydroxypropyl methyl cellulose using an appropriate blender.
- b) Dissolve sodium hydroxide in purified water.
- c) Using appropriate granulation equipment, slowly add the sodium hydroxide solution to the blend with constant mixing. Add a further aliquot of water, if necessary.
- d) Continue mixing to achieve additional massing. Add a further aliquot of water, if necessary.
- e) Dry granulated in a fluid bed dryer.
- f) Screen dried granules through an appropriate mill.
- g) Add magnesium stearate to the screened granules and blend together.
- h) Pass the granulated material through a chilsonator, if needed.
- i) Compress the granules into tablets using appropriately sized tooling.
- j) Disperse OPADRY powder in water and film-coat using appropriately sized coating equipment to a typical level of 2-4% by weight.
- k) Polish with carnauba wax using a typical level of 0.002-0.003% by weight.

EXAMPLE 4

Identifying Elevated Levels of BNP

[0063] The patient rests for about 10-15 minutes by sitting or lying. Blood is drawn in 10 mL aliquots and placed in tubes containing EDTA. The samples are centrifuged at about 0 °C to about 10 °C at about 1,000 to 2,000 rpm. The plasma is then removed and tested using Elecsys® proBNP sandwich immunoassay on an Elecsys® 2010 (Roche Diagnostics, Basel Switzerland). Levels of less about 100 pg/mL suggest little or no risk of heart failure is present. Levels between 100 pg/mL and 300 pg/mL suggest a risk of heart failure is present. Levels above 300 pg/mL indicate a risk of

mild heart failure. Levels above 600 pg/mL indicate a risk of moderate heart failure. Levels above 900 pg/mL indicate a risk of severe heart failure. Levels above about 80 pg/mL indicate that the patient may be at risk of an adverse coronary event.

Treating the patients with ranolazine

[0064] After identifying a patient with an elevated NT-proBNP level, a clinician administers ranolazine according to the protocol described above.

EXAMPLE 5

Effects of Ranolazine in Patients with non-ST Elevation Acute Coronary Syndromes in the MERLIN-TIMI 36 Trial

Background

[0065] Ranolazine is believed to exert anti-ischemic effects through a reduction in myocardial cellular sodium and calcium overload with a consequent reduction in wall stress. B-type natriuretic peptide (BNP) is released in response to increased wall stress and is a potent indicator of risk in acute coronary syndromes (ACS). This study was a prospective evaluation of the interaction between BNP and the effect of ranolazine as part of a randomized, double-blind, placebo-controlled trial.

[0066] Patients with a non-ST elevation acute coronary syndrome (NSTEMI), refers to patients with chest discomfort or anginal equivalent occurring at rest, lasting ≥ 10 minutes, and consistent with myocardial ischemia, and the presence of ischemic symptoms (≥ 5 minutes) at rest within 48 hours of admittance which may include index episode, and having at least one of the following indicators of moderate - high risk:

- Elevated cardiac troponin (above local MI limit) or CK-MB ($>ULN$)
- ST-depression (horizontal or down-sloping) ≥ 0.1 mV
- Diabetes mellitus (requiring insulin or oral therapy)
- A Risk Score of ≥ 3 wherein one point is assigned for each of the following variables and a total score calculated as the arithmetic sum:
 - Age ≥ 65 years;
 - Known CAD (prior MI, CABG, PCI or angiographic stenosis $\geq 50\%$);
 - Three or more cardiac risk factors (DM, elevated cholesterol, hypertension, family history);

- More than one episode of ischemic discomfort at rest in the prior 24 hours;
- Chronic aspirin use in the 7 days preceding onset of symptoms;
- ST segment depression ≥ 0.05 mV; and
- Elevated cardiac troponin or CK-MB.

Methods

[0067] Plasma BNP were measured (ADVIA® BNP) at baseline (N = 4543), 14 days (N = 4079), and the Final Visit (N = 3328) of the patients with non-ST elevation ACS randomized to ranolazine or placebo in the MERLIN-TIMI 36 trial. During the trial 200 mg of ranolazine (or matching placebo) was administered intravenously over 1 hour, followed by an 80 mg/hr intravenous infusion, which was reduced to 40 mg/hr for patients with an estimated creatinine clearance of less than 30 mL/min (<0.50 mL/s), and was continued for 12 to 96 hours. On or near the completion (about 1 hour before completion) of the infusion, transition from the IV solution to ranolazine extended release (or matching placebo) occurred and ranolazine extended release (or matching placebo) continued orally at a dose of 1000 mg twice daily until the end of the study. [Morrow, et al., J. Am. Med. Assoc'n. 297 (16), 1775-1783, 2007]. Patients were stratified using BNP >80 pg/mL based upon prior work with this assay. The primary endpoint of the trial was a composite of cardiovascular death, myocardial infarction, or recurrent ischemia (CVD/MI/RI).

Results

[0068] Patients with an elevated concentration of BNP (N = 1935) were at significantly higher risk of CVD/MK/RI (26.4% vs. 20.4%, $p<0.0001$), as well as CVD/MI (16.2% vs. 7.5%, $p<0.0001$) and CVD alone (9.0% vs. 2.4%, $p<0.0001$). In patients with BNP >80 pg/mL, ranolazine was associated with a significant reduction in the primary endpoint (HR 0.79; 95%CI 0.66-0.94, $p=0.009$) contrasting with the lack of detectable effect in those with a negative BNP result (see Figure 1, p -interaction = 0.052). This was driven by the endpoint component of worsening angina (HR 0.55; 0.35-0.84; $p = 0.005$ for BNP >80 pg/mL; HR 0.90; 95% CI 0.63-1.28, $p=0.55$ for BNP ≤ 80 pg/mL; p -interaction 0.079).

Conclusions

[0069] Elevated BNP is associated with worse outcomes in patients with ACS. The results of this planned exploratory analysis suggest that ranolazine may have enhanced efficacy in recurrent ischemia in patients with elevated BNP.

[0070] From the foregoing description, various modifications and changes in the composition and method will occur to those skilled in the art. All such modifications coming within the scope of the appended claims are intended to be included therein.

WHAT IS CLAIMED IS:

1. A method of reducing the risk of adverse coronary events in a mammalian patient, said method comprising:
 - a. identifying a patient exhibiting elevated levels of a natriuretic peptide;and
 - b. administering to said patient a therapeutically effective amount of ranolazine.
2. The method of claim 1, wherein the risk of adverse coronary events in a patient arises from adverse coronary events including, but not limited to, coronary artery disease (CAD), cardiovascular death, myocardial infarction, acute heart failure, ischemia, recurrent ischemia, acute coronary syndrome, diastolic dysfunction, and the like.
3. The method of claim 2, wherein the adverse coronary event is acute coronary syndrome.
4. The method of claim 2, wherein the adverse coronary event is cardiovascular death, myocardial infarction, or recurrent ischemia.
5. The method of claim 1, wherein the natriuretic peptide is brain-type natriuretic peptide (BNP) or N-terminal pro-brain natriuretic peptide (NT-proBNP).
6. The method of claim 1, wherein the patient is selected by performing a BNP assay.
7. The method of claim 1, wherein the patient exhibits about 80 picograms or greater of natriuretic peptide per milliliter of blood.
8. The method of claim 7, wherein the patient exhibits between about 100 and about 300 picograms of natriuretic peptide per milliliter of blood.

9. The method of claim 7, wherein the patient exhibits between about 300 and about 600 picograms of natriuretic peptide per milliliter of blood.
10. The method of claim 7, wherein the patient exhibits between about 600 and about 900 picograms of natriuretic peptide per milliliter of blood.
11. The method of claim 7, wherein the patient exhibits greater than about 900 picograms of natriuretic peptide per milliliter of blood.
12. The method claim 1, wherein the ranolazine is administered in an oral dose.
13. The method of claim 12, wherein the oral dose is a sustained release tablet.
14. The method of claim 13, wherein the patient is administered the tablet once a day, twice a day, or three times a day.
15. The method of claim 13, wherein the tablet comprises from about 350 to about 1000 milligrams of ranolazine
16. The method of claim 15, wherein the sustained release tablet comprises at least 50% by weight ranolazine, a pH dependent binder, and a pH independent binder.
17. The method of claim 16, wherein the sustained release table comprises at least 50% by weight ranolazine, from about 5 to about 12.5 % by weight methacrylic acid copolymer, and from about 1 to about 3 % by weight of hydroxypropyl methylcellulose, microcrystalline cellulose, sodium hydroxide, and magnesium stearate.
18. The method of claim 1, wherein the ranolazine is administered as an intravenous (IV) solution.

19. The method of claim 18, wherein the IV solution comprises from about 1.5 to about 3 milligrams of ranolazine per milliliter of solution.
20. The method of claim 19, wherein the IV solution is administered to the patient for a time sufficient to reduce the risk of adverse coronary events in the patient.
21. The method of claim 20, wherein the IV solution is administered at 200 mg intravenously over 1 hour.
22. The method of claim 21, wherein the IV solution is further administered as an infusion of 80 mg/hour.
23. The method of claim 20, wherein the IV solution is administered for up to about 96 hours.
24. The method of claim 20, wherein after administration of the IV solution to the patient, the patient is then transitioned from the IV solution to an oral dose by administering an oral sustained release formulation of ranolazine.
25. The method of claim 20, wherein 1 hour prior to completion of the administration of the IV solution, the patient is transitioned from the IV solution to an oral dose by administering an oral sustained release formulation of ranolazine.
26. The method of claim 24, wherein at the time of transition from the IV solution to the oral sustained release formulation of ranolazine, the IV solution is administering 60 mg/hour of ranolazine and the oral dose of ranolazine is 375 mg twice daily.
27. The method of claim 24, wherein at the time of transition from the IV solution to the oral sustained release dose of ranolazine, the IV solution is administering 80 mg/hour of ranolazine and the oral dose of ranolazine is 1000 mg twice daily.
28. The method of claim 20, wherein the IV solution of ranolazine is administered intravenously at 200 mg for 1 hour, followed by an IV infusion of 80 mg/hour for 12 to 96 hours.


29. The method of claim 28, wherein the patient is transitioned from IV solution to the oral sustained release dose of ranolazine, and the oral dose of ranolazine is 1000 mg twice daily.

30. A kit of parts comprising:

- a) an assay that detects levels of natriuretic peptides in blood plasma and/or heart tissue of a mammalian patient, and
- b) a pharmaceutical dosage of ranolazine.

31. The kit of claim 30, wherein the pharmaceutical dosage of ranolazine is an IV solution or and oral dose.

32. The kit of claim 31, wherein the pharmaceutical dosage is both an IV solution and an oral dose.

**Figure 1**