

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 2931902 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **A 01 H 5/06 (2006.01)** **C 12 N 15/82 (2006.01)** **A 01 H 3/04 (2006.01)**
A 01 H 4/00 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2018-09-03**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-08-01**
- (86) Europæisk ansøgning nr.: **13819001.2**
- (86) Europæisk indleveringsdag: **2013-12-13**
- (87) Den europæiske ansøgnings publiceringsdag: **2015-10-21**
- (86) International ansøgning nr.: **EP2013076618**
- (87) Internationalt publikationsnr.: **WO2014091021**
- (30) Prioritet: **2012-12-13 US 201261736817 P** **2012-12-13 EP 12196858**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **SESVanderHave N.V., Industriepark 15 , Soldatenplein Zone 2, No. 15, 3300 Tienen, Belgien**
- (72) Opfinder: **HAIN, Rüdiger, Dreieichstrasse 44, 60594 Frankfurt, Tyskland**
JOHANN, Gerhard, Bürgermeister-Schmidt-Strasse 24, 51399 Burscheid, Tyskland
Weyens, Guy, Dwersbos 25, 1650 Beersel, Belgien
LEFÈBVRE, Marc, Rue de la Chapelle Stevenaert 75, 1370 Jodoigne Souveraine, Belgien
- (74) Fuldmægtig i Danmark: **NORDIC PATENT SERVICE A/S, Bredgade 30, 1260 København K, Danmark**
- (54) Benævnelse: **FREMGANGSMÅDE TIL UDVIKLING AF HERBICIDRESISTENTE SUKKERROEPLANTER**
- (56) Fremdragne publikationer:
EP-A2- 0 257 993
WO-A1-95/10178
WO-A1-2012/049266
WO-A1-2012/049268
WO-A1-2012/150335
HALL R D ET AL: "SUGAR BEET GUARD CELL PROTOPLASTS DEMONSTRATE A REMARKABLE CAPACITYFOR CELL DIVISION ENABLING APPLICATIONS IN STOMATAL PHYSIOLOGY AND MOLECULAR BREEDING", JOURNAL OF EXPERIMENTAL BOTANY, OXFORD UNIVERSITY PRESS, GB, vol. 48, no. 307, 1 February 1997 (1997-02-01), pages 255-263, XP000893145, ISSN: 0022-0957, DOI: 10.1093/JXB/48.2.255
GUREL EKREM ET AL.: "Biotechnology applications for sugar beet", CRITICAL REVIEWS IN PLANT SCIENCES, vol. 27, no. 2, 2008, pages 108-140, XP009167608, DOI: 10.1080/07352680802202000
Dyer, Hess, Holt and Duke: "Potential Benefits and Risks of Herbicide-Resistant Crops Produced by Biotechnology" In: "Horticultural Reviews", 1993, John Wiley & Sons, XP002693215, ISBN: 978-0-471-57338-8 vol. 15, pages 367-408, page 379, line 6 - line 19
REY P ET AL: "ATRAZINE AND DIURON RESISTANT PLANTS FROM PHOTOAUTOTROPHIC PROTOPLAST-

Fortsættes ...

DERIVED CULTURES OF NICOTIANA-PLUMBAGINIFOLIA", PLANT CELL REPORTS, vol. 9, no. 5, 1990, pages 241-244, XP009167607, ISSN: 0721-7714

HORIKOSHI MAMORU ET AL: "Selection of tobacco cell lines resistant to photobleaching herbicides", JOURNAL OF PESTICIDE SCIENCE, vol. 24, no. 1, 1999, pages 13-16, XP009167605, ISSN: 0385-1559

TAN S ET AL: "IMIDAZOLINONE-TOLERANT CROPS: HISTORY, CURRENT STATUS AND FUTURE", PEST MANAGEMENT SCIENCE, WILEY & SONS, BOGNOR REGIS; GB, vol. 61, no. 3, 1 January 2005 (2005-01-01), pages 246-257, XP009058795, ISSN: 1526-498X, DOI: 10.1002/PS.993

DUGGLEBY ET AL: "Structure and mechanism of inhibition of plant acetohydroxyacid synthase", PLANT PHYSIOLOGY AND BIOCHEMISTRY, GAUTHIER-VILLARS, PARIS, FR, vol. 46, no. 3, 14 January 2008 (2008-01-14), pages 309-324, XP022550175, ISSN: 0981-9428

DUGGLEBY R G ET AL: "ACETOHYDROXYACID SYNTHASE", JOURNAL OF BIOCHEMISTRY AND MOLECULAR BIOLOGY, KOREAN SOCIETY FOR BIOCHEMISTRY AND MOLECULAR BIOLOGY, KR, vol. 33, no. 1, 1 January 2000 (2000-01-01), pages 1-36, XP001119823, ISSN: 1225-8687

DESCRIPTION

Field of the invention

[0001] The present invention relates to a method to generate sugar beet plants resistant to herbicides, for instance inhibitor(s) of the acetohydroxyacid synthase enzyme (ALS).

[0002] The present invention further relates to the plants that are obtained by this method.

[0003] Sugar beet is an important agricultural crop in temperate and subtropical regions.

[0004] In modern agriculture, herbicides are widely used to manage weeds proliferation.

[0005] The development of sugar beet plants resistant towards herbicide(s) such as ALS inhibitors can be undertaken by using transgenic approaches.

[0006] Indeed, the introduction of foreign DNA carrying a gene conferring resistance towards an herbicide has been successfully performed in a variety of field crops, including in sugar beet.

[0007] WO 95/10178 discloses a transgenic-induced resistance to the herbicide Bialaphos. The gene encoding the resistance is introduced in protoplasts of guard cells from sugar beet, then these protoplasts are regenerated into sugar beet plants. Those plants are further resistant to the chemical phosphinothricin and its derivative glufosinate.

Non transformed plants did not acquire resistance to Bialaphos.

[0008] At best, 28 transformed calli were regenerated from 83000 protoplasts; at worst, 1 callus was regenerated from 190 000 protoplasts. Subsequently, some of these calli can show somatic embryogenesis and can regenerate sugar beet plantlets, with an efficiency of 1% and, a characteristic found very advantageous, up to 30%.

[0009] This approach, is nevertheless not commonly used in the art, which widely relies on the more direct transformation of calli from explants obtained from sugar beet organs such as embryos and/or from leaf discs.

[0010] There still exists an important need to develop herbicide-resistant plants such as in the sugar beet crop and that without relying on DNA vectors and/or on the insertion of foreign genes.

[0011] On the other hand, the development of a sugar beet resistant plant towards herbicide(s), such as ALS inhibitors can theoretically be undertaken via classical breeding with a sugar beet plant which would contain a naturally-occurring resistance gene. However, such

approach is time consuming and, to the applicants' best knowledge, did not result into success, despite the fact that naturally-occurring ALS inhibitor-resistant plants were documented, at least for species other than sugar beet including a variety of weed species. Especially a double mutant (i.e a plant having two mutations in the same ALS gene) is very unlikely to occur in nature.

[0012] The patent application WO 98/02527 discloses a method for the manufacture of sugar beet plant resistant to some ALS inhibitors, such as sulfonylurea herbicides, which comprises the steps of exposing to sulfonylurea calli obtained from explants of *B. vulgaris*, and regenerating plants from the few spontaneous mutants that can grow in the presence of this herbicide.

[0013] This method yielded a plant having a mutation in the ALS gene, where the proline at position 188 of the encoded ALS enzyme (corresponding to the 197 position of the *Arabidopsis thaliana* ALS enzyme) is substituted by a serine. However, this mutant is not commercially used because treatments with the preferred modern sulfonylureas ALS herbicides (e.g. foramsulfuron) exhibit some phytotoxicity in field trials at the necessary dose rate.

[0014] The patent application WO 2012/049268 relies on the same method, except that calli obtained from explants of *B. vulgaris* are exposed to foramsulfuron, and thus resulted into sugar beet plant resistant to several ALS inhibitors, including to foramsulfuron.

[0015] This method yielded a plant having a mutation in the ALS gene, where the tryptophan at position 569 of the encoded ALS enzyme (corresponding to the 574 position of the *Arabidopsis thaliana* ALS enzyme) is substituted by a leucine.

[0016] Field trials of this homozygote 569/569 mutant showed good resistance to foramsulfuron, to iodosulfuron (another ALS inhibitor), as well as to mixtures of different ALS inhibitors.

[0017] Both these published methods take benefit from the well-established steps of isolation of calli from individual explants such as embryos. However, this is a time-consuming approach involving (i) isolation of large number of fresh embryos, (ii) their repeated culture on agar-solidified culture medium, and (iii) the selection of the regenerable calli using morphological selection approaches.

[0018] Other strategies to transfer genetic traits to sugar beet have been developed and rely on mesophyll protoplasts (which are different of stomatal guard cell protoplasts) as starting material (Krens et al., 1990, Theor. appl. Genet., vol 79, pages 390-396).

Gurel et al. 2008, Critical reviews in Plant Science, 27, 108-140, discloses biotechnology applications for sugar beet. Seven different *in vitro* culture techniques are disclosed. Among them is the culture of protoplasts either from stomatal guard cells for transformation purposes, or from friable callus from etiolated hypocotyl explants, the latter being much more efficient. This approach using protoplasts is however associated to major difficulties.

Hall et al. 1997, J. Exp. Botany, 48, 255-263 have used a culture of 500000 stomatal guard cell protoplasts for carrying a transformation experiment with foreign DNA. The transformation efficiency was higher than 2%. On the other hand, *in vitro* culture of plants is reported to be associated to stomatal failure, pointing to problems in the corresponding cells and thus to the production of very high amount of stomatal guard cell protoplasts, such as amounts needed for carrying a method involving a mutational event.

Summary of the invention

[0019] In a broad aspect, the present invention discloses a method for producing a mutant sugar beet plant being resistant to an herbicide comprising the steps of:

- obtaining protoplasts from stomatal guard cells isolated from a sugar beet plant;
- applying to an *in vitro* culture of these protoplasts a composition comprising this herbicide at a concentration that is lethal to more than 99% of the *in vitro* cultured cells; and
- regenerating sugar beet plants from the surviving cells of these *in vitro* cultured cells,
- possibly selecting regenerated sugar beet plants having a mutation in the gene encoding the peptide(s) targeted by this herbicide,

wherein these stomatal guard cells protoplasts are pre selected for their capacity to regenerate into a sugar beet plant, wherein the pre selected protoplasts have a probability of more than 10% (number of growing protoplasts:total protoplast number put into culture) to divide and regenerate into a viable callus, wherein calli obtained by the said protoplasts have more than 10% (number of calli producing shoots:total calli number) of capacity to develop shoots, and wherein this herbicide is applied to more than 20 000 000 of these protoplasts.

[0020] The herbicide used in the present method can be an herbicide not targeting ALS gene.

[0021] The preferred herbicides (not targeting ALS) are selected from the group consisting of:

- 4-HPPD inhibitors (such as mesotrione, isoxaflutole, pyrasulfotole, benzobicyclon, benzofenap, pyrazolynate, pyrazoxyfen, tembotrione, topramezone, sulcotrione and sulcotrion), inhibitors of the carotenoid biosynthesis (such as flurtamone, fluridone, flurochloridone, beflubutamid, norflurazon, picolinafen and diflufenican);
- inhibitors of EPSP synthase (such as glyphosate or glyphosate-trimesium);
- inhibitors of photosystem II (such as Phenyl-carbamates for instance phenmedipham or desmedipham, Pyridazinones (for instance chloridazon = pyrazon), Triazines (for instance cyanazine, remtal, eglinazine-ethyl, proglinazine-ethyl, ametryn, atrazine, desmetryn, dimethametryn, prometon, prometryn, propazine, simazine, simetryn, terbumeton, terbuthylazine, terbutryn, methoprotyn), Triazinones (for instance met amitron, metribuzin, hexazinone), Uracils (bromacil, lenacil, terbacil), Ureas (dimefuron, isoproturon, linuron, monolinuron, ethidimuron, methabenzthiazuron, tebuthiuron, diuron, fenuron, neburon, siduron, isouron, chlorobromuron, chlorotoluron,

chloroxuron, fluometuron, metobromuron, metoxuron, thiazafluron, monuron, cycluron, monolinuron), or amicarbazone, solan, propanil, bentazon, bromoxynil, ioxynil, bromofenoxim, pyridate, pyridafol;

- inhibitors of photosystem I (such as diquat or paraqua);
- inhibitors of cell division (such as carbetamide, chlorpropham, propham, naproanilide, diphenamid, napropamide, butenachlor, metazachlor, diethatyl-ethyl, acetochlor, alachlor, butachlor, propachlor Monsanto, propisochlor, dimethachlor, dimethenamid, metolachlor, pretilachlor, S-metolachlor, pethoxamid, thenylchlor, anilofos, cafenstrole, indanofan, bromobutide, piperophos, flufenacet, mefenacet, fentrazamide);
- inhibitors of microtubule assembly (such as propyzamide = pronamide, tebutam, chlorthal-dimethyl = DCPA, fluchloralin, pendimethalin, butralin, benefin=benfluralin, ethalfluralin, oryzalin, trifluralin, prodiamine, dinitramine, butamifos, dithiopyr, thiazopyr);
- inhibitors of protoporphyrinogen oxidase (such as Diphenylethers (acifluorfen-sodium, bifenox, ethoxyfen-ethyl, chlornitrofen, fluoroglycofen-ethyl, oxyfluorfen, chlomethoxyfen, fluordifen, fomesafen, lactofen, nitrofen, aclonifen), N-phenylphthalimides (cinidon-ethyl, flumiclorac-pentyl, flumioxazin), Oxadiazoles (oxadiargyl, oxadiazon) Oxazolidinediones, (pentoxazone), Phenylpyrazoles (fluazolate, pyraflufen-ethyl), Pyrimidindiones (saflufenacil, benzfendizone, butafenacil), Thiadiazoles (thidiazimin, fluthiacet-methyl), Triazolinones (azafenidin, carfentrazone-ethyl sulfentrazone), pyraclonil, proflunaz, flufenpyr-ethyl);
- inhibitors of Acetyl CoA carboxylase (such as Aryloxyphenoxypropionates (such as clodinafop-propargyl, cyhalofop-butyl, diclofop-methyl, fenoxaprop-P-ethyl, fluazifop-P-butyl, haloxyfop-etotyl, haloxyfop-methyl, haloxyfop-P-methyl, propaquizafop, quizalofop-P-ethyl or quizalofop-P-tefuryl), Cyclohexanediones (such as alloxymid, butoxydim, clethodim, cycloxydim, profoxydim, sethoxydim, tepraloxymid or tralkoxydim), Phenylpyrazoline (pinoxaden));
- cell wall synthesis inhibitors (such as indaziflam, isoxaben, chlorthiamid, dichlobenil, quinclorac, flupoxam);
- inhibitor of glutamine synth(et)ase (glufosinate-ammonium or bialaphos=bilanaphos) and
- synthetic auxin (such as TBA, dicamba, chloramben, benazolin-ethyl 4, dichlorprop-P, mecoprop-P, 2,4,5-T (weedare) 2,4-D (Weedare), 2,4-DB (Butyrol), dichlorprop, MCPB, mecoprop, MCPA-thioethyl, clomeprop, Agroxone 4, aminopyralid, clopyralid, fluroxypyr, halauxifen-methyl, picloram, triclopyr, quinclorac, quinmerac),

more preferably from the group consisting of 4-HPPD inhibitors, inhibitors of the carotenoid biosynthesis, inhibitors of EPSP synthase, inhibitors of photosystem II, inhibitors of photosystem I, inhibitors of microtubule assembly, inhibitors of protoporphyrinogen oxidase and synthetic auxin.

[0022] Another highly preferred herbicide is an ALS inhibitor.

[0023] Therefore, the present invention relates to a method for producing a mutant sugar beet plant being resistant to one or more inhibitor(s) of the acetohydroxyacid synthase enzyme (ALS) comprising the steps of:

- obtaining protoplasts from stomatal guard cells isolated from a sugar beet plant;
- applying to an *in vitro* culture of the said protoplasts a composition comprising one or more ALS inhibitor(s) at a concentration that is lethal to (more than 99% (preferably more than 99.9% or even more than 99.99%) of) the *in vitro* cultured cells (yet allowing some mutants to escape); and
- regenerating sugar beet plants from the surviving cells of the said *in vitro* cultured cells, wherein the said stomatal guard cells protoplasts are pre selected for their capacity to regenerate into a sugar beet plant, wherein the pre selected protoplasts have a probability of more than 10% (number of growing protoplasts:total protoplast number put into culture) to divide and regenerate into a viable callus, wherein calli obtained by the said protoplasts have more than 10% (number of calli producing shoots:total calli number) of capacity to develop shoots, and wherein the said ALS inhibitor(s) is/are applied to more than 20 000 000 or even more than 50 000 000 of the said protoplasts.

[0024] Preferably this method comprises the sub-steps of isolating stomatal guard cells protoplasts from sugar beet plants of different genotypes and measuring for each genotype the proportion of the said protoplasts that is growing when the said protoplasts are put in culture.

[0025] Preferably this method further comprises the step of sequencing the genome of the regenerated plants from the surviving *in vitro* cultured cells, advantageously for identifying a mutation in the ALS gene and/or for selecting regenerated sugar beet plants having one or more, preferably one, two, or more mutations in the ALS gene.

[0026] Advantageously, sugar beet having one or several mutation(s) in the ALS gene at positions encoding the amino acids selected from the group consisting of Glycine 112, Alanine 113, Methionine 115, Arginine 133, Valine 187, Arginine 190, Alanine 196, Phenylalanine 197, Lysine 247, Methionine 346, Histidine 347, Arginine 368, Aspartate 370, Aspartate 371, Arginine 372, Methionine 565, Valine 566, Phenylalanine 573, Serine 648 and Glycine 649, preferably selected from the group consisting of Alanine 196, Aspartate 371, Arginine 372, Serine 648 and Glycine 649, are obtained and/or selected by the method of the present invention. Another preferred sugar beet plant having a mutation of the Proline at position 188 and of the Tryptophan at position 569 is obtained and/or selected by the method of the present invention.

[0027] Preferably, sugar beet having two or more mutations has two mutations in one allele, meaning that the encoded peptide harbours two mutations synergizing the resistance towards ALS inhibitors (especially towards compositions comprising several ALS inhibitors). An examples of most preferred sugar beet is a sugar beet having a mutation of the Proline at position 188 and of the Tryptophan at position 569 in one allele (and possibly the same mutations in the second allele; alternatively, the second allele harbours different mutations).

[0028] Alternatively, sugar beet having two mutations has one mutations on each allele.

[0029] The method of the present invention allows to regenerate a sugar beet plant having one mutation in the ALS gene at positions encoding Proline 188 and one or more mutation(s) in the ALS gene at positions encoding Glycine 112, Alanine 113, Methionine 115, Arginine 133, Valine 187, Arginine 190, Alanine 196, Phenylalanine 197, Lysine 247, Methionine 346, Histidine 347, Arginine 368, Aspartate 370, Aspartate 371, Arginine 372, Methionine 565, Valine 566, Tryptophan 569 (preferably mutated into Glycine), Phenylalanine 573, Serine 648 and Glycine 649. An examples of preferred sugar beet is a sugar beet having a mutation of the Proline at position 188 and of another mutation (possibly not the tryptophan 569 or the Trp569Gly) in the same allele.

[0030] The method of the present invention allows to regenerate a sugar beet plant having one mutation in the ALS gene at position encoding Tryptophan 569 (preferably mutated into Leucine) and one or more mutation(s) in the ALS gene at positions encoding Glycine 112, Alanine 113, Methionine 115, Arginine 133, Valine 187, Proline 188 (preferably mutated into Threonine, Arginine, Leucine, Glutamine or Alanine), Arginine 190, Alanine 196, Phenylalanine 197, Lysine 247, Methionine 346, Histidine 347, Arginine 368, Aspartate 370, Aspartate 371, Arginine 372, Methionine 565, Valine 566, Phenylalanine 573, Serine 648 and Glycine 649. An examples of preferred sugar beet is a sugar beet having a mutation of the tryptophan at position 569 and of another mutation (possibly not the Proline 188 or the Pro188Ser) in the same allele.

[0031] The method of the present invention allows to regenerate a sugar beet plant having one or more mutation(s) in the ALS gene, wherein the said one or more mutation is selected from the group consisting of Alanine 113 Proline 188, Alanine 196, Aspartate 371, Arginine 372, Tryptophan 569, Serine 648 and Glycine 649, wherein the said Alanine 113 is mutated into Valine or Threonine, wherein the said Proline 188 is mutated into Threonine, Arginine, Leucine, Glutamine or Alanine, wherein the said Alanine 196 is mutated into Valine, wherein the said Aspartate 371 is mutated into Glutamate, wherein the said Arginine 372 is mutated into Histidine, wherein the said Tryptophan 569 is mutated into Glycine, wherein the said Serine 648 is mutated into Threonine and wherein the said and Glycine 649 is mutated into Aspartate.

[0032] Preferably, sugar beet having two or more of these specific mutations has two mutations in one allele, meaning that the encoded peptide harbours two mutations synergizing the resistance towards ALS inhibitors (especially towards compositions comprising several ALS inhibitors).

[0033] The method of the present invention allows to regenerate a sugar beet plant having one mutation in the ALS gene at position encoding Proline 188 and one mutation in the ALS gene at position encoding Tryptophan 569.

[0034] Advantageously, this method comprises a preliminary step of deducing the concentration of the one or more ALS inhibitor(s) at which the composition (comprising the one or more ALS inhibitor(s)) is lethal for at least 99% (preferably at least 99.9% or even 99.99%)

of the *in vitro* cultured cells (yet allowing some mutants to escape).

[0035] The preferred ALS inhibitor present in the composition (the said composition comprising in addition another ALS inhibitor, being preferably from another class than sulfonylurea ALS inhibitor, such as thien carbazon-methyl) to be added to the *in vitro* cultured stomatal guard cell protoplasts is foramsulfuron, preferably at a concentration of 10^{-9} mol/l to (more preferably) 10^{-6} mol/l.

[0036] Another suitable ALS inhibitor present in the composition to be added to the *in vitro* cultured stomatal guard cell protoplasts is ethoxysulfuron (possibly this composition further comprises other ALS inhibitors).

[0037] A related aspect of the present invention is a mutated sugar beet plant obtainable by the method of the present invention, such as a sugar beet having a mutation at tryptophan 569 and/or at proline 188.

[0038] Another related aspect of the present invention is a mutated sugar beet plant comprising SEQ.ID.NO:3 (or SEQ.ID.NO:4) and/or SEQ.ID.NO:5 (or SEQ.ID.NO:6).

[0039] A sugar beet plant disclosed herein corresponds to the deposit under NCIMB 42050 or NCIMB 42051.

[0040] The present invention is also related to tissue or plant part (for instance stomatal guard cells or leaf strips) or seeds derived from the mutated plant of the invention, as well as to their use for the introduction of another genetic trait.

[0041] The present invention is also related to the use of the (well regenerating) stomatal guard cell protoplasts developed in the present invention for the introduction of another genetic trait than resistance to ALS inhibitors.

Detailed description of the invention

[0042] The inventors have discovered that protoplasts from stomatal guard cell of sugar beet represent a good starting material for inducing resistance towards herbicides such as ALS inhibitors, despite the fact that regeneration of a sugar beet plant from a stomatal guard cell protoplast is very difficult, of low occurrence, and requires longer and more complex procedures than the direct regeneration from calli obtained from explants, as usually used in the art.

[0043] Indeed, callus (calli) from explants is (are) a mass of non-differentiated (or dedifferentiated) cells, which, under appropriate culture conditions, will differentiate (or re-differentiate) and regenerate a wholly functional sugar beet plant. In such method, seeds are

collected in large amounts, then easily sterilized and explants are thus obtained in large amounts. Such method is convenient as allowing to perform a large part of the work without the difficulties of a sterile environment.

On the other hand, the stomatal guard cells have a well-defined organization in the plant, and their isolation from the plant tissue results into individual cell(s), then after a treatment into individual protoplasts.

In the method of the present invention, plantlets must be grown *in vitro* and maintained under sterile conditions, then stomatal guard cells are isolated from these small plantlets and protoplasts are obtained while maintaining sterile conditions, then these individual protoplasts are induced to produce again their cell wall, to grow firstly into micro-calli, then into a sugar beet plant.

Stomatal guard cells protoplasts of sugar beet were found to be very vulnerable, reducing the scope of their application: for instance addition of a mutagen, instead of favouring the appearance of a desired genetic trait, was found to be lethal to the protoplasts in many cases.

In other words, it was completely unexpected to succeed in using stomatal guard cell protoplasts as a starting material for developing a sugar beet plant having acquired a desired genetic trait through mutation.

[0044] Furthermore, the inventors have found that the capacity of stomatal guard cell protoplasts to divide is very dependent on sugar beet genotypes: the stomatal guard cell protoplasts from the majority of the genotypes exhibit a very low (almost zero) capacity to divide (grow), while the protoplasts from some specific genotypes have a sizable capacity to grow *in vitro*.

[0045] The inventors have identified that the culture on solid medium (polymer-containing medium, such as alginate or agarose-like containing medium) of very large amounts of these stomatal guard cell protoplasts, then the exposure of the cultured material (normally in the form of regenerated calli) to herbicides such as ALS inhibitor(s) resulted into the selection of some mutated cells being resistant to this herbicide molecule, despite the fact that, in absence of added mutagens, spontaneous mutations are known to be very rare, and that the method cannot take benefit of the genetic diversity within a population, in selecting an already existing mutant plant exhibiting the desired resistance towards one or more ALS inhibitors.

[0046] Some of these mutated cells having acquired resistance towards herbicides such as ALS inhibition were further able to regenerate into viable sugar beet plant(s). Although protoplasts are very difficult to use in practice, the present method nevertheless yielded very rapidly stable mutants having acquired resistance towards the herbicide.

[0047] The method of the present invention allows the production of herbicide-resistant (ALS inhibitor-resistant) sugar beet plants having one or several mutations, including one or several mutations in the enzyme being the target of the herbicide, for instance an ALS inhibitor (such as a mutation of the ALS gene, wherein the ALS gene encodes an ALS protein containing an amino acid different from tryptophan at position 569 of the ALS protein; corresponding to position 574 of the *Arabidopsis thaliana* ALS), possibly in addition to at least another mutation

in this gene and/or to other mutations in other genes.

[0048] An aspect of the present invention is therefore a method for producing sugar beet plant (*Beta vulgaris*; thus also *Beta vulgaris maritima* ssp) resistant to one or more ALS inhibitor(s) comprising the steps of:

- obtaining (protoplasts from) stomatal guard cells isolated from a sugar beet plant (sensitive to ALS inhibitors);
- applying to these *in vitro* cultured cells a composition comprising one or more ALS inhibitors at a concentration that is lethal to these *in vitro* cultured cells (at least 99% of the (wild-type) cells are killed by the ALS inhibitor, but some mutants can escape the treatment); and

regenerating sugar beet plants from the surviving cells.

[0049] This method can be seen as an alternative to the introduction of a transgene (for instance disclosed in WO 95/10178). However, a method based on mutation of endogenous gene(s) is more laborious by comparison to a transgenic approach. Indeed, the introduction of a transgenic (mutated) gene is much more rapid, flexible and predictable.

[0050] The absence of transgene-induced resistance refers to the fact that the acquired resistance to the ALS inhibitor(s) is not directly caused by the *in vitro* insertion of a (foreign) DNA into the stomatal guard cells protoplasts (or into the cells obtained thereof), such as a (foreign) DNA encoding a protein directly providing resistance to the ALS inhibitor(s), such as a protein locally reducing the toxicity of the ALS inhibitor(s) (e.g. an enzyme degrading the ALS inhibitor(s) or reducing its intracellular concentration), or an ALS protein being resistant to ALS inhibitor(s), such as an ALS mutant enzyme that retains significant activity and functionality even in the presence of the inhibitor(s).

[0051] However, the plant obtained by the method of the invention can further be crossed with a transgenic plant in order to stack another genetic trait in the progeny. The plant (or a part thereof) obtained by the present invention can further be used in a subsequent transgenic method to introduce another genetic trait (different from the genetic trait obtained in the method of the present invention).

[0052] Possibly, this method further comprises the step of adding a mutagen agent to the culture of the isolated stomatal guard cell protoplasts.

Suitable mutagen agents are physical (such as UV or X-ray) exposure or exposure to chemical agents (such as Ethyl Methane Sulfonate (EMS), for instance at 0.05%, 0.1%, 0.15%, 0.2% or even at 2.5%). However, protoplasts viability was shown to be detrimentally affected by some mutagenesis treatments usually performed, such as treatment of more than 0.2% EMS.

[0053] Alternatively, this method thus does not comprises the step of adding a mutagen agent.

[0054] In the context of the present invention, an herbicide preferably refers to any molecule,

which, when applied at a given dose, is used for weed control.

[0055] Preferred herbicides used in the present method (for developing sugar beet plants resistant to this herbicide) have a specific known activity on one peptide (so that a single mutation in the corresponding gene can confer resistance towards this herbicide). In other words, the preferred herbicides are specific for one peptide target (usually inhibiting specifically the activity of one plant enzyme) and/or their target peptide is known (so as to be in a position to screen the resistant sugar beet for a mutation in the target gene)

[0056] In the context of the present invention, an ALS inhibitor refers to any molecule inhibiting the function of the ALS gene.

[0057] Preferably, the ALS inhibitor(s) used in the present invention (substantially) do not inhibit other (sugar beet) enzymes than ALS.

[0058] Advantageously, the ALS inhibitor(s) are selected and used in the present invention at a concentration where more than 90% (more than 95%, more than 99%, more than 99.9%) of the function of the wild-type ALS enzyme is inhibited, but substantially not affecting the function of non-related enzymes.

[0059] Suitable ALS inhibitors (for carrying the method) are selected from the group consisting of sulfonylurea herbicides, sulfonylaminocarbonyltriazolinone herbicides, imidazolinone herbicides, triazolopyrimidine herbicides and pyrimidinyl(thio)benzoate herbicides. Advantageously, the herbicide composition comprises at least one sulfonylurea herbicides and at least one triazolopyrimidine.

The preferred ALS inhibitors (for carrying the method) are foramsulfuron, amidosulfuron, thien carbazon-methyl, ethoxysulfuron and mixture thereof (especially a composition comprising thien carbazon-methyl and either foramsulfuron or amidosulfuron); however, the regenerated sugar beet mutant is advantageously resistant towards several ALS inhibitors.

[0060] Other ALS inhibitors (including mixtures of ALS inhibitors) can be used, and the skilled person knows which mutation provides strong resistance to a given ALS inhibitor (e.g. the mutation of the tryptophan at position 569 is known to be associated to foramsulfuron resistance); hence, given the flexibility and the efficacy of the present method, the skilled person can design sugar beet plants having acquired several mutations and thus broader resistance towards ALS inhibitors (e.g. mixtures of ALS inhibitors).

[0061] More preferably, this method comprises a preliminary step of selecting a sugar beet plant genotype (line) for its capacity of the stomatal guard cells protoplasts to regenerate into a fully functional sugar beet plant, and/or the method of the present invention is carried on stomatal guard cells protoplasts isolated from well-regenerating sugar beet genotypes (lines).

[0062] A suitable preliminary step (of selecting sugar beet plant genotypes for the ability of their stomatal guard cells protoplasts cells to regenerate into a sugar beet plant) involves the

comparison (independently of their possible advantageous features such as yield or resistance towards parasitic infections) of at least 10 different sugar beet plant genotypes (from different genotype backgrounds), preferably at least 15 different genotypes, or even at least 30 different genotypes, for the capacity of their stomatal guard cells protoplasts to regenerate into a sugar beet plant (much more preferably their capacity to grow *in vitro* and/or to form calli), and the selection of a well-regenerating genotype (line) for carrying the method of the present invention.

[0063] In the context of the present invention, well-regenerating stomatal guard cells protoplasts refer to protoplasts having a probability of more than 10% (growing:total) or even more than 20% (growing:total) or 50% (growing:total) to divide and/or to grow and/or regenerate into viable sugar beet callus (when grown in the suitable culture media and without exogenous selection pressure such as the toxic molecule/herbicide to be applied in the method of the present invention).

[0064] Callus (calli) refers to a mass of undifferentiated cells. In the art, calli can be obtained from explants such as embryos, or parenchyme-derived explants from leaves or cotyledon. However, in the context of the present invention, calli are the result of the growth of (well-regenerating) stomatal guard cells protoplasts.

The calli obtained by these well-regenerating protoplasts have more than 10% (number of calli producing shoots : total calli number; shoots:total), preferably more than 20% (shoots:total) or even more than 30% (shoots:total) of capacity to develop shoots.

[0065] Preferably, well-regenerating sugar beet stomatal guard cells protoplasts refer to protoplasts having more than 0.1% (sugar beet plant : total protoplast number) (more preferably more than 1%) of capacity to regenerate into a viable sugar beet plant.

[0066] In this method, the composition comprising the herbicide (e.g. one or more ALS inhibitor(s)) is applied to an *in vitro* culture of more than 20 000 000 of these (well-regenerating) sugar beet stomatal guard cells protoplasts.

[0067] Preferably, the composition comprising the (one or more) ALS inhibitor(s) is applied to an *in vitro* culture of more than 50 000 000, of these (well-regenerating) sugar beet stomatal guard cells protoplasts.

[0068] Preferably, at least 50 000, such as about 100 000, (well-regenerating) stomatal guard cells protoplasts per millilitre were grown onto polymer-containing medium (such as alginate- or agarose-containing medium).

[0069] Possibly, these, (well-regenerating) sugar beet stomatal guard cells protoplasts are grown for at least about one week (preferably about 3 weeks and/or of less than 4 weeks) on polymer (alginate)-containing medium, before the application of the composition comprising the (one or more) ALS inhibitor(s) (such as foramsulfuron and possibly thienicarbazone-methyl).

[0070] Preferably, this method further comprises the step of comparing the growth of mutated stomatal guard cell and the growth of wild-type stomatal guard cell (and/or naive and/or not yet herbicide-treated) on a medium that does not comprise ALS inhibitor, and possibly of selecting mutant(s) keeping at least 75% of the growth, preferably at least 90% of the growth of the corresponding wild-type cell.

[0071] Preferably, or in addition this method further comprises the step of comparing the growth and/or the yield of the regenerated sugar beet from the mutated cell and the growth and/or the yield of the wild-type (and/or naive and/or not yet treated by the ALS inhibitor) sugar beet in greenhouse assays and in agronomic conditions without any ALS inhibitor, and possibly of selecting ALS inhibitor-resistant mutant(s) sugar beet plants keeping at least 75% of the growth and/or of the yield, preferably at least 90% of the growth and/or of the yield of the wild-type sugar beet.

[0072] Preferably, this method further comprises the step of sequencing the regenerated plants from the surviving protoplasts and/or of identifying one or several mutation(s) that are (can be) associated to the resistance to the herbicide (e.g. one or more ALS inhibitor(s)).

[0073] In the context of the present invention, the term 'mutation' preferably refers to one (one single) change in the nucleotide sequence encoding the peptide targeted by the herbicide (e.g. the ALS protein) that causes one change in the corresponding amino acid, such that the resulting plant has acquired some resistance towards the herbicide such as ALS inhibitors.

In other words, in the context of the present invention, 'mutation' is preferably understood as equivalent to a 'point mutation' that allows some resistance to the herbicide (the ALS inhibitor). Accordingly, "several mutations" preferably refers, in the present invention, to (a stack of) multiple point mutations, each point mutation causing a change of the encoded amino acid so as to provide some resistance to an herbicide (e.g. an ALS inhibitor and/or an herbicide not an ALS inhibitor). Therefore, preferably, in the context of the present invention 'mutation' does not comprise change in the nucleotide sequence that do not modify the encoded protein (such as a change in the third amino acid of a coding triplet), change in amino acids that are not associated to herbicide (such as ALS inhibitor) resistance, nor multiple simultaneous changes in the nucleotide sequence.

[0074] Advantageously, this identification step of the mutation(s) associated to the resistance towards the (one or more) ALS inhibitor(s) (preferably one or two mutation(s) in the ALS gene) is coupled to the development of oligonucleotide primers spanning over this mutation.

[0075] Advantageously, this identification step of the mutation(s) in the ALS gene is followed by (*in vitro*) enzymatic activity measurements of the protein encoded by the wild-type and by the mutated ALS genes.

[0076] Preferably, these enzymatic measurements of the wild-type ALS enzyme and of the mutated ALS enzyme are performed in the presence of one or more ALS inhibitors (at one or

at several concentrations in order to derive an inhibition curve).

[0077] Possibly, these enzymatic measurement of the wild-type enzyme and of the mutated enzyme are (further) performed in absence of ALS inhibitors (to compare the enzymatic activity of the mutated enzyme; preferably, in absence of the ALS inhibitor, the mutated enzyme keeps at least 50% of the activity of the wild-type enzyme, more preferably at least 75%, still more preferably at least 90%, at least 95% or even at least 99%).

[0078] Preferably, the method of the present invention comprises a step of comparing compositions comprising one or more ALS inhibitors at different concentrations, and deducing the concentration at which the ALS inhibitor, and/or ALS inhibitor in a special formulation within this composition, is /are lethal for an *in vitro* culture of the stomatal guard cell protoplast isolated from the sugar beet plant (such as stomatal guard cell protoplasts grown on alginate for at least one week).

[0079] For instance, this step of deducing the concentration at which the (one or more) ALS inhibitor(s) is/are lethal for stomatal guard cell protoplast isolated from the sugar beet plant is performed on an *in vitro* culture of the stomatal guard cell protoplasts isolated from the wild-type sugar beet plant (and/or naive and/or not yet treated by the ALS inhibitor).

[0080] In the context of the present invention, the lethal concentration of the composition comprising (one or more) ALS inhibitor(s) refers to a concentration sufficient to kill at least 99%, preferably at least 99.9%, more preferably at least 99.99% of the cultured cells (yet allowing some mutants to escape this treatment).

[0081] Alternatively, or in addition, this step of deducing the concentration at which the (one or more) ALS inhibitor(s) is lethal for an *in vitro* culture of the stomatal guard cell protoplasts isolated from the sugar beet plant is (further) performed on an *in vitro* culture of mutated stomatal guard cells (on cells having acquired a mutation in the ALS gene and being resistant to ALS inhibitor(s)).

[0082] The comparison of the lethal concentration of the ALS inhibitor (in a composition comprising this ALS inhibitor) on the naive and on the mutated cells is advantageously expressed as a ratio (or as several ratios, one ratio per ALS inhibitor tested).

[0083] Preferably, for one ALS inhibitor, the lethal concentration on naive cell(s) is 50-fold lower than the lethal concentration on mutated cell(s), more preferably, the lethal concentration on naive cell(s) is 200-fold lower than the lethal concentration on mutated cell(s), still more preferably, the lethal concentration on naive cell(s) is 1000-fold lower than the lethal concentration on mutated cell(s).

[0084] The herbicide (used in the method of the present invention) can be a mixture (of inhibitors) comprising at least one ALS inhibitor, such as foramsulfuron.

[0085] Possibly, the ALS inhibitor used in the method of the invention is a mixture of ALS inhibitors, such as a sulfonylurea (e.g. foramsulfuron) and another ALS inhibitor selected from the group consisting of iodosulfuron, amidosulfuron and thien carbazone-methyl.

[0086] Preferably, the ALS inhibitor used in the method of the invention is (or comprises) foramsulfuron, such as foramsulfuron applied to a one-week old (or to a three-weeks old) *in vitro* culture of protoplasts (more particularly to the *in vitro* culture comprising calli regenerated from these cultured protoplasts) on alginate-containing medium and maintained during the *in vitro* culture of the cells at a concentration of 10^{-9} - 10^{-6} mol/l (or 10^{-9} - 10^{-6} mol/l).

[0087] A related aspect of the present invention is a mutated sugar beet plant obtainable by this method, (for instance when this methods comprises the use of one or more herbicide (not ALS inhibitors) or the use of one or more ALS inhibitors, or comprises the use of one ALS inhibitor and of one herbicide being not an ALS inhibitor).

[0088] Therefore, one aspect of the present invention is a sugar beet (obtainable by the method of the present invention) having one or several mutation(s) in the ALS gene at positions encoding the amino acids selected from the group consisting of Glycine 112, Alanine 113, Methionine 115, Arginine 133, Valine 187, Arginine 190, Alanine 196, Phenylalanine 197, Lysine 247, Methionine 346, Histidine 347, Arginine 368, Aspartate 370, Aspartate 371, Arginine 372, Methionine 565, Valine 566, Phenylalanine 573, Serine 648 and Glycine 649.

[0089] A preferred sugar beet (obtainable by the method of the present invention) has one or several mutation(s) in the ALS gene at positions encoding the amino acids selected from the group consisting of Alanine 113 (e.g. mutated into Valine or Threonine) Proline 188 mutated into Threonine, Arginine, Leucine, Glutamine or Alanine, Alanine 196 (e.g. mutated into Valine), Aspartate 371 (e.g. mutated into glutamate), Arginine 372 (e.g. mutated into Histidine), Tryptophan 569 mutated into Glycine, Serine 648 (e.g. mutated into Threonine) and Glycine 649 (e.g. mutated into Aspartate).

[0090] A related aspect of the present invention is a mutated sugar beet plant (or a mutated sugar beet plant cell such as a mutated stomatal guard cell isolated from sugar beet) comprising a mutation in the ALS gene where the tryptophan at position 569 in the encoded ALS enzyme (corresponding to position 574 in the Arabidopsis thaliana ALS enzyme) is substituted by another amino acid (such as a leucine), and possibly another (one or several) mutation, preferably another (one or several) mutation in the ALS gene, such as a mutation causing a further amino acid substitution in the ALS gene.

[0091] Another preferred a sugar beet plant has a mutation of the Tryptophan into Leucine at position 569 and one or several mutation(s) in the ALS gene at positions encoding the amino acids selected from the group consisting of Glycine 112, Alanine 113, Methionine 115, Arginine 133, Valine 187, Arginine 190, Alanine 196, Phenylalanine 197, Lysine 247, Methionine 346, Histidine 347, Arginine 368, Aspartate 370, Aspartate 371, Arginine 372, Methionine 565, Valine 566, Phenylalanine 573, Serine 648 and Glycine 649.

[0092] A preferred sugar beet (obtainable by the method of the present invention) has one mutation Tryptophan into Leucine mutation at position 569 and one or several mutation(s) in the ALS gene at positions encoding the amino acids selected from the group consisting of Alanine 113 (e.g. mutated into Valine or Threonine) Proline 188 mutated into Threonine, Arginine, Leucine, Glutamine or Alanine, Alanine 196 (e.g. mutated into Valine), Aspartate 371 (e.g. mutated into Glutamate), Arginine 372 (e.g. mutated into Histidine), Serine 648 (e.g. mutated into Threonine) and Glycine 649 (e.g. mutated into Aspartate).

[0093] This mutated sugar beet plant is resistant to one or several ALS inhibitor(s) used, such as a sulfonylurea (e.g. foramsulfuron) and advantageously to other ALS inhibitor(s), preferably selected from the group consisting of iodosulfuron, amidosulfuron and thienencarbazone-methyl.

[0094] A related aspect of the present invention is a mutated sugar beet plant (or a mutated sugar beet plant cell such as a mutated stomatal guard cell isolated from sugar beet) comprising a mutation in the ALS gene where the proline at position 188 in the encoded ALS enzyme (corresponding to position 197 in *Arabidopsis thaliana* ALS enzyme) is substituted by another amino acid (such as a serine).

[0095] Alternatively, a preferred sugar beet plant (obtainable by the method of the present invention) has a mutation of the Proline into Serine at position 188 and one or several mutation(s) in the ALS gene at positions encoding the amino acids selected from the group consisting of Glycine 112, Alanine 113, Methionine 115, Arginine 133, Valine 187, Arginine 190, Alanine 196, Phenylalanine 197, Lysine 247, Methionine 346, Histidine 347, Arginine 368, Aspartate 370, Aspartate 371, Arginine 372, Methionine 565, Valine 566, Phenylalanine 573, Serine 648 and Glycine 649.

[0096] A preferred sugar beet (obtainable by the method of the present invention) has one Proline into Serine mutation in the ALS gene at position 188 and one or more mutation(s) of the ALS gene at positions encoding Alanine 113 (e.g. mutated into Valine or Threonine), Aspartate 371 (e.g. mutated into glutamate), Arginine 372 (e.g. mutated into Histidine), Tryptophan 569 mutated into Glycine, Serine 648 (e.g. mutated into Threonine) and Glycine 649 (e.g. mutated into Aspartate).

[0097] Another related aspect of the present invention is a mutated sugar beet plant (or a mutated sugar beet plant cell such as a mutated stomatal guard cell isolated from sugar beet) comprising a mutation of tryptophan at position 569 in the ALS enzyme and a mutation of proline at position 188 in the ALS enzyme, as well as possibly another (one or several) mutation, preferably another (one or several) mutation in the ALS gene.

[0098] Preferably, (one allele of) the ALS gene of this mutated sugar beet plant corresponds to SEQ.ID.NO:3 or SEQ.ID.NO:5.

[0099] Advantageously, the mutated sugar beet plant of the present invention comprises

SEQ.ID.NO:3 (in one allele) and SEQ.ID.NO:5 (in the second allele).

[0100] Possibly, the mutated sugar beet plant of the present invention comprises SEQ.ID.NO:3 (in one allele) and either SEQ.ID.NO:1, or SEQ.ID.NO:7 (in the second allele).

[0101] Another related aspect of the present description is a nucleotide fragment (of at least 20 or at least 25 consecutive nucleotides, but of less than 200 consecutive nucleotides, preferably of less than 50 consecutive nucleotides) covering the one or more mutations; possibly this fragment is for use as a primer or a probe (including a nucleotide probe being further labelled e.g. by a non-nucleotidic moiety or using radioactivity, or a probe labelled with a nucleic acid sequence foreign to the ALS gene of sugar beet).

[0102] Still another related aspect of the present description is the use of this nucleotide fragment spanning over the mutation for marker-assisted selection of sugar beet plants having a resistance towards the toxic molecule (herbicide).

Examples

Comparative example

[0103] Because mutated sugar beet were successfully generated in the art (e.g. in WO 98/02527) upon the addition of ALS herbicide to calli being explants from wild-type sugar beet, the inventors firstly selected the sugar beet genotype (line) derived from the line of WO 98/02527 and isolated protoplasts from their stomatal guard cells.

[0104] Several millions of these protoplasts (on average about 2 to about 5 millions, and up to 11 millions by experiment; in total, ALS herbicide was applied to about 150 millions of protoplasts) were isolated as in WO 95/10178, placed in culture medium comprising alginate, and treated with MS culture medium comprising 10^{-9} to 10^{-6} mol/l foramsulfuron.

[0105] The inventors have then regenerated the sugar beet, following the protocol as described in WO 95/10178, and observed only a few calli surviving to the ALS inhibitor. However, to one exception, none of these regenerated calli were able to develop into a sugar beet plant. The only regenerated sugar beet plant showed no mutation in the ALS gene (encoding the target enzyme of foramsulfuron).

[0106] Therefore this sugar beet line, whose parental line was shown, on the basis of the direct exposition of calli (from an explant) to an herbicide, to acquire mutation-induced resistance to this herbicide, was not useful for the same purpose, when the method involves stomatal guard cells protoplasts.

Example 1 Selection of sugar beet genotypes (lines) for well-regenerating protoplasts

[0107] The inventors have then compared several sugar beet plant genotypes for their capacity of regeneration from stomatal guard cells protoplasts.

[0108] The inventors have found genotypes having about 0.01% (or even less) of capacity to regenerate and several genotypes having (much) more than 0.1% of capacity to regenerate.

[0109] The inventors have further established a distinction between the growth of the protoplasts (their capacity to growth and divide *in vitro*), the capacity of the grown calli to form shoots and the proportion of growing calli to regenerate a plant.

Table 1: comparison of some sugar beet genotypes

Genotype	protoplasts/gram	Percentage growing cells	Percentage shoot formation	Percentage obtained plants
F06R38309	1500000	0,02	55,67	3,43
F06R38313	500000	0,04	0,14	0,00
F06R38323	1000000	0,19	0,49	10,81
F07R38836	500000	0,26	10,51	0,73
REL1	1000000	0,07	70,00	44,00

[0110] Although the values reflecting the capacity of a stomatal guard cell protoplast to regenerate into a whole sugar beet plant, when taken as a whole were higher for the cell line "Rel1", this cell line was considered as not useful enough for running the present invention.

[0111] The inventors have further concluded that the "percentage of growing cells" parameter is much more important for running the present invention than the other parameters.

[0112] The inventors have selected a genotype having more than 0.25% of stomatal guard cells protoplasts that are able to grow *in vitro*.

Example 2 Herbicide treatment of protoplasts

[0113] The inventors have applied the same approach as in the comparative Example, but relying on well growing stomatal guard cells protoplasts (for instance identified as in Example 1; plants deposited as NCIMB 42050 or NCIMB 42051 can also be used, as well as other sugar beet plants having a high proportion of growing stomatal guard cell protoplasts).

[0114] In total, about 68 millions of well-growing stomatal guard cell protoplasts were treated

with an ALS herbicide composition comprising up to 10^{-6} M foramsulfuron.

[0115] The inventors obtained 46 calli.

[0116] Several regenerated plants are showing a mutation in the target gene, the ALS gene: in each case a mutation in the codon for tryptophan at position 569 (W569L; corresponding to tryptophan at position 574 in *Arabidopsis thaliana*). The two alleles of the ALS genes of this mutant are encoded by SEQ.ID.NO:3 and SEQ.ID.NO:7. Other grown calli were sequenced and have mutations in the ALS gene, (including mutations at other positions) but did not regenerate into a plant.

[0117] The inventors therefore conclude that the method of the present invention is very useful to develop plants having evolved mutations causing a resistance to an herbicide, especially since this method does not involve the use of foreign DNA and/or the introduction of DNA vectors encoding genetic elements already known to confer resistance to ALS inhibitors, and yielded positive results in only a few months.

[0118] The inventors then repeated this method and further applied a mutagen (0.05% to 0.2% EMS) to the protoplasts in order to increase the number of mutations.

Example 3 ALS inhibitor treatment of sugar beets

[0119] The inventors compared the behaviour of regenerated sugar beets having the mutated SEQ.ID.NO:3 (heterozygote for this mutation) and a wild-type (naïve) sugar beet commercial variety.

[0120] The (heterozygote) mutated variety showed good resistance to Foramsulfuron (12.5 g/ha; up to 3 applications), even when the herbicide has been combined with an organic compound (25 g/ha rapeseed oil methyl ester) to boost its effect.

[0121] As expected, the wild-type (naïve) plant was very sensitive to Foramsulfuron, even after the first application.

[0122] The same experiment was performed using amidosulfuron (15 g /ha), and yielded the same level of resistance in the mutated plants.

On the other hand, the wild-type (naïve) plants were very sensitive to amidosulfuron, especially when combined with the organic compound, and/or after several applications of amidosulfuron.

[0123] The same experiment was performed using iodosulfuron (3.5 g/ha), and demonstrated a good level of resistance in the mutated plants when iodosulfuron was added, but this resistance declined when iodosulfuron is applied together with the organic compound.

As expected, the wild-type (naïve) plant was very sensitive to iodosulfuron even after one application and without the organic compound.

[0124] The same experiment was performed using 7.5 g/ha thien carbazone-methyl, and yielded about the same level of resistance as for iodosulfuron in the mutated plants. The wild-type (naive) plant was very sensitive to thien carbazone-methyl at all the tested concentrations and regardless of the addition of the organic compound.

[0125] The inventors conclude that, by comparison to the wild-type, the mutated sugar beet plant comprising SEQ.ID.NO:3 (deposited under the Budapest Treaty NCIMB 42051) offers the best resistance against foramsulfuron.

[0126] The inventors further conclude that this (heterozygote) mutated plant has further acquired some (although partial) resistance towards other ALS inhibitors, including towards inhibitors belonging to other chemical classes.

Example 4 ALS inhibitor treatment of sugar beets having further mutations in the ALS gene

[0127] The inventors then developed a mutated sugar beet plant comprising SEQ.ID.NO:3 and SEQ.ID.NO:5 (on two different alleles). Such resulting dual mutant has been deposited under the Budapest Treaty under NCIMB 42050. A plant comprising both SEQ.ID.NO:3 and SEQ.ID.NO:5 can be generated by relying on several techniques, including, for instance, a subsequent mutagenesis step applied to the single mutant NCIMB 42051.

[0128] The inventors then compared the resistance of this dual mutant plant (a mutation in one allele at amino acid 569 and a mutation in the other allele at amino acid 188) with the single mutant (a mutation at position 569) sugar beet.

[0129] The dual mutant plant line at least keeps all the resistance features as in Example 3, and has also acquired a good resistance (compatible with field application) towards thien carbazone-methyl and towards amidosulfuron treatments, even when put in composition with organic compounds.

Therefore, this dual mutant plant displays improved, synergistic, resistance towards several ALS inhibitors by comparison to the resistance attributed to the single mutant plant (at position 569 in the ALS gene).

Example 5 Greenhouse trials: ALS inhibitor treatment of different sugar beets in direct comparison

[0130] Mutated sugar beet plants comprising SEQ.ID.NO:3 and SEQ.ID.NO:5 (on two different alleles) according to the present invention (as described in Example 4 above, "Line A") were treated with different ALS inhibitors in direct comparison with sugar beet plants where the

tryptophan at position 569 of the encoded ALS enzyme is substituted by a leucine ("Line B"), sugar beet plants described in WO 98/02527 where the proline at position 188 of the encoded ALS enzyme is substituted by a serine ("Line C"), and traditional variety (wild-type) sugar beet plants not having a mutation at positions 569 and 188 ("Line WT").

[0131] Several groups of seeds of the four different mentioned sugar beet plants were sown separately in the greenhouse and grew up to stage BBCH 14 for *Beta vulgaris* L. *ssp. vulgaris* (i.e. 4 leaves (the second pair) unfolded) according to the monographie "Entwicklungsstadien mono- und dikotyler Pflanzen", 2nd edition, 2001, ed. Uwe Meier, Biologische Bundesanstalt für Land und Forstwirtschaft. Subsequently, the resulting separate groups of sugar beet plants were each individually treated with an ALS inhibitor (ALS-in) in the amounts (g/ha) indicated in Table 2.

[0132] On day 14 after application of the respective ALS inhibitor, the damage (i.e. the phytotoxicity) for each sugar beet plant was rated on a scale from 0% (i.e. no damage, no phytotoxicity) to 100% (i.e. the plants were completely killed). The average rating for each group of plants is also shown in Table 2.

Table 2:

ALS-in	ALS-in g/ha	Line A	Line B	Line C	Line WT
Foramsulfuron	13	26.9%	45.6%	77.5%	80.0%
Iodosulfuron-methyl-Na	3.5	22.5%	38.8%	80.0%	82.5%
Amidosulfuron	15	6.3%	37.5%	51.9%	73.1%
Thiencarbazone-methyl	7.5	8.1%	35.6%	37.5%	84.4%
Bisbyribac-Na	50	17.5%	38.1%	71.7%	80.0%
Metosulam	15	13.1%	40.6%	69.4%	79.4%

[0133] Additionally, typical early phenotypes of each sugar beet plants were inspected after treatment with a mixture comprising thiencarbazone-methyl and foramsulfuron. A representative early phenotype of each Line is shown in Figure 1 (Fig.1).

[0134] Fig. 1 also demonstrates that the sugar beet plants according to the present invention ("Line A") show improved ALS inhibitor resistance, i.e. superior growth and less phytotoxic effects were observed in comparison to the other early phenotypes.

Example 6 Field trials: ALS inhibitor treatment of different sugar beets in direct comparison

Table 3:

[0135]

Table 3: effect of ALS inhibitors on sugar beet plant. The values represent the average percentage of measured damage.

		Sensitive	574 hetero	574&197	574 homo
1	UNTREATED	0	0	0	0
2	AE F 130360 00 WG50 A1 25g/HA (Foramsulfuron)	97	22	5	0
3	BYH18636 15g/Ha (Thiencarbazone)	97	39	5	0
4	ae f115008 00 wg 10 a2 7g/Ha (iodosulfuron)	98	65	23	28
5	AE F130060 00 WG75 A2 60g/Ha (mesosulfuron)	91	24	18	0
6	HOESTAR 30g/Ha (amidosulfuron)	97	34	0	0
7	AEF095404 00 WG60 A2 60 g/Ha (ethoxysulfuron)	99	39	0	0
8	RAPTOR 40 g/Ha (imazamox)	98	44	35	8
9	TACCO 30g/Ha (metosulam)	97	27	0	3
10	NOMINEE 50g/Ha (bispyribac)	98	78	70	28
11	MOTIVELL 60 g/Ha (nicosulfuron)	98	53	28	13
12	CROPPER SX 8 g/Ha (metosulfuron)	100	74	50	35
13	LEXUS 50 DF10g/Ha (flupyrsulfuron)	70	0	0	0
14	ATTRIBUT 70 g/Ha (propoxycarabazone)	91	25	0	5
15	SIMPLICITY 50g/Ha (pyroxysulam)	97	45	28	0
16	PRIMUS 10 g/Ha (florasulam)	99	55	38	0
17	POINTER SX 30g/Ha (tribenuron)	98	74	28	20
18	CATO 13 g/Ha (rimsulfuron)	68	8	0	0
19	MONITOR 80 WG 10g/Ha (sulfosulfuron)	93	23	0	0
20	DEBUT YX115 g/Ha (triflusulfuron)	0	0	0	0
21	EVEREST 40g/Ha (flucarbazone)	93	18	0	0
22	HARMONY 7.5 g/Ha (thiensulfuron)	98	39	0	0
23	# 2 & #3 (Foramsulfuron+Thiencarbazone)	100	65	35	5

		Sensitive	574 hetero	574&197	574 homo
	1 L/Ha				

[0136] The inventors have tested commercial compositions at the dose allowing destruction of the weeds.

[0137] The sensitive control (i.e. a sugar beet having no mutation in the ALS gene) was killed by all the herbicides but one. The inventors have measured small damages to control (untreated) plant, reaching sometimes 35% or even 40%. These 'damages' reflect the agronomic conditions of this field trial.

[0138] On the other hand, sugar beet plant being heterozygote at position 569 (574) have become partially resistant towards several herbicidal compositions. The plant having incorporated the 569 mutation (574) on both alleles and thus being (569/569) homozygote has a further increased resistance: only 7 herbicide composition are moderately toxic (from 5% to 35%).

[0139] A sugar beet plant having incorporated the mutation at position 569 (574) in one allele of the ALS gene and a mutation at position 188 (197) in the second allele of the ALS gene have also acquired improved resistance, since 9 herbicide compositions are moderately toxic, and only 3 are quite toxic. Surprisingly, such plant, where a mutation providing a strong resistance (569) has been lost and a mutation providing only a weak resistance (188) towards ALS inhibitor has been added, provides even better resistance than the homozygote (569/569) plant in 3 different conditions of this field test.

PCT

Print Out (Original in Electronic Form) (This sheet is not part of and does not count as a sheet of the international application)

[0140]

0-1	Form PCT/RO/134 (SAFE) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bis)	
0-1-1	Prepared Using	PCT Online Filing
		Version 3.5.000.235 MT/FOP
		20020701/0.20.5.20

0-2	International Application No.	PCT/EP2013/076618
0-3	Applicant's or agent's file reference	BPSESS0010PC
1	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
1-1	Paragraph number	8
1-3	Identification of deposit	
1-3-1	Name of depositary institution	NCIMBNCIMB Ltd.
1-3-2	Address of depositary institution	Ferguson Building, Craibstone Estate, Bucksburn, Aberdeen AB21 9YA, United Kingdom
1-3-3	Date of deposit	24 September 2012 (24.09.2012)
1-3-4	Accession Number	NCIMB 42050
1-4	Additional Indications	paragraphs 39, 114, 124, 127
1-5	Designated States for Which Indications are Made	All designations
2	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
2-1	Paragraph number	8
2-3	Identification of deposit	
2-3-1	Name of depositary institution	NCIMBNCIMB Ltd.
2-3-2	Address of depositary institution	Ferguson Building, Craibstone Estate, Bucksburn, Aberdeen AB21 9YA, United Kingdom
2-3-3	Date of deposit	24 September 2012 (24.09.2012)
2-3-4	Accession Number	NCIMB 42051
2-4	Additional Indications	paragraphs 39, 114, 124, 127
2-5	Designated States for Which Indications are Made	All designations

FOR RECEIVING OFFICE USE ONLY

[0141]

0-4	This form was received with the international	
------------	--	--

	application:	Yes
	(yes or no)	
0-4-1	Authorized officer	Aoustin, Isabelle

FOR INTERNATIONAL BUREAU USE ONLY

[0142]

0-5	This form was received by the international Bureau on:	
0-5-1	Authorized officer	

SEQUENCE LISTING

[0143]

<110> SES

<120> ALS resistant sugar beet

<130> SESS0010

<160> 8

<170> PatentIn version 3.5

<210> 1

<211> 1995

<212> DNA

<213> Beta vulgaris

<220>

<221> CDS

<222> (1)..(1995)

<223> 4D6834 WT all

<400> 1

```

atg gcg gct acc ttc aca aac cca aca ttt tcc cct tcc tca act cca      48
Met Ala Ala Thr Phe Thr Asn Pro Thr Phe Ser Pro Ser Ser Thr Pro
1           5           10           15

tta acc aaa acc cta aaa tcc caa tct tcc atc tct tca acc ctc ccc      96
Leu Thr Lys Thr Leu Lys Ser Gln Ser Ser Ile Ser Ser Thr Leu Pro
20          25          30

ttt tcc acc cct ccc aaa acc cca act cca ctc ttt cac cgt ccc ctc      144
Phe Ser Thr Pro Pro Lys Thr Pro Thr Pro Leu Phe His Arg Pro Leu

```

35	40	45	
caa atc tca tcc tcc caa tcc cac aaa tca tcc gcc att aaa aca caa			192
Gln Ile Ser Ser Ser Gln Ser His Lys Ser Ser Ala Ile Lys Thr Gln			
50	55	60	
act caa gca cct tct tct cca gct att gaa gat tca tct ttc gtt tct			240
Thr Gln Ala Pro Ser Ser Pro Ala Ile Glu Asp Ser Ser Phe Val Ser			
65	70	75	80
cga ttt ggc cct gat gaa ccc aga aaa ggg tcc gat gtc ctc gtt gaa			288
Arg Phe Gly Pro Asp Glu Pro Arg Lys Gly Ser Asp Val Leu Val Glu			
85	90	95	
gct ctt gag cgt gaa ggt gtt acc aat gtg ttt gct tac cct ggt ggt			336
Ala Leu Glu Arg Glu Gly Val Thr Asn Val Phe Ala Tyr Pro Gly Gly			
100	105	110	
gca tct atg gaa atc cac caa gct ctc aca cgc tct aaa acc atc cgc			384
Ala Ser Met Glu Ile His Gln Ala Leu Thr Arg Ser Lys Thr Ile Arg			
115	120	125	
aat gtc ctc cct cgc cat gaa caa ggc ggg gtt ttc gcc gcc gag gga			432
Asn Val Leu Pro Arg His Glu Gln Gly Gly Val Phe Ala Ala Glu Gly			
130	135	140	
tat gct aga gct act gga aag gtt ggt gtc tgc att gcg act tct ggt			480
Tyr Ala Arg Ala Thr Gly Lys Val Gly Val Cys Ile Ala Thr Ser Gly			
145	150	155	160
cct ggt gct acc aac ctc gta tca ggt ctt gct gac gct ctc ctt gat			528
Pro Gly Ala Thr Asn Leu Val Ser Gly Leu Ala Asp Ala Leu Asp			
165	170	175	
tct gtc cct ctt gtt gcc atc act ggc caa gtt cca cgc cgt atg att			576
Ser Val Pro Leu Val Ala Ile Thr Gly Gln Val Pro Arg Arg Met Ile			
180	185	190	
ggc act gat gct ttt cag gag act cca att gtt gag gtg aca agg tct			624
Gly Thr Asp Ala Phe Gln Glu Thr Pro Ile Val Glu Val Thr Arg Ser			
195	200	205	
att act aag cat aat tat tta gtt ttg gat gta gag gat att cct aga			672
Ile Thr Lys His Asn Tyr Leu Val Leu Asp Val Glu Asp Ile Pro Arg			
210	215	220	
att gtt aag gaa gcc ttt ttt tta gct aat tct ggt agg cct gga cct			720
Ile Val Lys Glu Ala Phe Phe Leu Ala Asn Ser Gly Arg Pro Gly Pro			
225	230	235	240
gtt ttg att gat ctt cct aaa gat att cag cag caa ttg gtt gtt cct			768
Val Leu Ile Asp Leu Pro Lys Asp Ile Gln Gln Gln Leu Val Val Pro			
245	250	255	
gat tgg gat agg cct ttt aag ttg ggt ggg tat atg tct agg ctg cca			816
Asp Trp Asp Arg Pro Phe Lys Leu Gly Gly Tyr Met Ser Arg Leu Pro			
260	265	270	
aag tcc aag ttt tcg acg aat gag gtt gga ctt ctt gag cag att gtg			864
Lys Ser Lys Phe Ser Thr Asn Glu Val Gly Leu Leu Glu Gln Ile Val			
275	280	285	
agg ttg atg agt gag tcg aag aag cct gtc ttg tat gtg gga ggt ggg			912
Arg Leu Met Ser Glu Ser Lys Lys Pro Val Leu Tyr Val Gly Gly Gly			
290	295	300	
tgt ttg aat tct agt gag gag ttg agg aga ttt gtt gag ttg aca ggg			960
Cys Leu Asn Ser Ser Glu Glu Leu Arg Arg Phe Val Glu Leu Thr Gly			
305	310	315	320
att ccg gtg gct agt act ttg atg ggg ttg ggg tct tac cct tgt aat			1008
Ile Pro Val Ala Ser Thr Leu Met Gly Leu Gly Ser Tyr Pro Cys Asn			
325	330	335	
gat gaa ctg tct ctt cat atg ttg gaa atg cac gaa act gtt tat acc			1056

Asp	Glu	Leu	Ser	Leu	His	Met	Leu	Gly	Met	His	Gly	Thr	Val	Tyr	Ala		
			340					345					350				
aat	tat	gcg	gtg	gat	aag	gcg	gat	ttg	ttg	ctt	gct	ttc	ggg	gtt	agg	1104	
Asn	Tyr	Ala	Val	Asp	Lys	Ala	Asp	Leu	Leu	Leu	Ala	Phe	Gly	Val	Arg		
		355					360					365					
ttt	gat	gat	cgt	gtg	acc	ggg	aag	ctc	gag	gcg	ttt	gct	agc	cgt	gct	1152	
Phe	Asp	Asp	Arg	Val	Thr	Gly	Lys	Leu	Glu	Ala	Phe	Ala	Ser	Arg	Ala		
	370					375					380						
aag	att	gtg	cat	att	gat	att	gac	tct	gct	gag	att	ggg	aag	aac	aag	1200	
Lys	Ile	Val	His	Ile	Asp	Ile	Asp	Ser	Ala	Glu	Ile	Gly	Lys	Asn	Lys		
	385				390				395					400			
cag	ccc	cat	gtg	tcc	att	tgt	gct	gat	gtt	aaa	ttg	gca	ttg	cgg	ggg	1248	
Gln	Pro	His	Val	Ser	Ile	Cys	Ala	Asp	Val	Lys	Leu	Ala	Leu	Arg	Gly		
			405					410					415				
atg	aat	aag	att	ctg	gag	tct	aga	ata	ggg	aag	ctg	aat	ttg	gat	ttc	1296	
Met	Asn	Lys	Ile	Leu	Glu	Ser	Arg	Ile	Gly	Lys	Leu	Asn	Leu	Asp	Phe		
			420					425					430				
tcc	aag	tgg	aga	gaa	gaa	tta	ggt	gag	cag	aag	aag	gaa	ttc	cca	ctg	1344	
Ser	Lys	Trp	Arg	Glu	Glu	Leu	Gly	Glu	Gln	Lys	Lys	Glu	Phe	Pro	Leu		
	435					440						445					
agt	ttt	aag	aca	ttt	ggg	gat	gca	att	cct	cca	caa	tat	gcc	att	cag	1392	
Ser	Phe	Lys	Thr	Phe	Gly	Asp	Ala	Ile	Pro	Pro	Gln	Tyr	Ala	Ile	Gln		
	450				455						460						
gtg	ctt	gat	gag	ttg	acc	aat	ggt	aat	gct	att	ata	agt	act	ggt	gtt	1440	
Val	Leu	Asp	Glu	Leu	Thr	Asn	Gly	Asn	Ala	Ile	Ile	Ser	Thr	Gly	Val		
	465				470				475					480			
ggg	cag	cac	caa	atg	tgg	gct	gcg	cag	cat	tac	aag	tac	aga	aac	cct	1488	
Gly	Gln	His	Gln	Met	Trp	Ala	Ala	Gln	His	Tyr	Lys	Tyr	Arg	Asn	Pro		
			485					490						495			
cgc	caa	tgg	ctg	acc	tct	ggt	ggg	ttg	ggg	gct	atg	ggg	ttt	ggg	cta	1536	
Arg	Gln	Trp	Leu	Thr	Ser	Gly	Gly	Leu	Gly	Ala	Met	Gly	Phe	Gly	Leu		
			500					505					510				
cca	gcc	gcc	att	gga	gct	gca	gtt	gct	cga	cca	gat	gca	gtg	gtt	gtc	1584	
Pro	Ala	Ala	Ile	Gly	Ala	Ala	Val	Ala	Arg	Pro	Asp	Ala	Val	Val	Val		
		515					520					525					
gat	att	gat	ggg	gat	ggc	agt	ttt	att	atg	aat	gtt	caa	gag	ttg	gct	1632	
Asp	Ile	Asp	Gly	Asp	Gly	Ser	Phe	Ile	Met	Asn	Val	Gln	Glu	Leu	Ala		
	530					535					540						
aca	att	agg	gtg	gaa	aat	ctc	cca	gtt	aag	ata	atg	ctg	cta	aac	aat	1680	
Thr	Ile	Arg	Val	Glu	Asn	Leu	Pro	Val	Lys	Ile	Met	Leu	Leu	Asn	Asn		
	545				550				555					560			
caa	cat	tta	ggt	atg	gtt	gtc	caa	tgg	gaa	gat	agg	ttc	tat	aaa	gct	1728	
Gln	His	Leu	Gly	Met	Val	Val	Gln	Trp	Glu	Asp	Arg	Phe	Tyr	Lys	Ala		
			565					570						575			
aac	cgg	gca	cat	aca	tac	ctt	gga	aac	cct	tcc	aaa	tct	gct	gat	atc	1776	
Asn	Arg	Ala	His	Thr	Tyr	Leu	Gly	Asn	Pro	Ser	Lys	Ser	Ala	Asp	Ile		
			580					585					590				
ttc	cct	gat	atg	ctc	aaa	ttc	gct	gag	gca	tgt	gat	att	cct	tct	gcc	1824	
Phe	Pro	Asp	Met	Leu	Lys	Phe	Ala	Glu	Ala	Cys	Asp	Ile	Pro	Ser	Ala		
		595					600					605					
cgt	gtt	agc	aac	gtg	gct	gat	ttg	agg	gcc	gcc	att	caa	aca	atg	ttg	1872	
Arg	Val	Ser	Asn	Val	Ala	Asp	Leu	Arg	Ala	Ala	Ile	Gln	Thr	Met	Leu		
	610						615				620						
gat	act	cca	ggg	cgg	tac	ctg	ctc	gat	gtg	att	gta	cgg	cat	caa	gag	1920	
Asp	Thr	Pro	Gly	Pro	Tyr	Leu	Leu	Asp	Val	Ile	Val	Pro	His	Gln	Glu		
	625				630				635					640			

cat gtg ttg cct atg att cca agt ggt gcc ggt ttc aag gat acc att 1968
 His Val Leu Pro Met Ile Pro Ser Gly Ala Gly Phe Lys Asp Thr Ile
 645 650 655

aca gag ggt gat gga aga acc tct taa 1995

Thr Glu Gly Asp Gly Arg Thr Ser
 660

<210> 2

<211> 664

<212> PRT

<213> Beta vulgaris

<400> 2

Met Ala Ala Thr Phe Thr Asn Pro Thr Phe Ser Pro Ser Ser Thr Pro
 1 5 10 15

Leu Thr Lys Thr Leu Lys Ser Gln Ser Ser Ile Ser Ser Thr Leu Pro
 20 25 30

Phe Ser Thr Pro Pro Lys Thr Pro Thr Pro Leu Phe His Arg Pro Leu
 35 40 45

Gln Ile Ser Ser Ser Gln Ser His Lys Ser Ser Ala Ile Lys Thr Gln
 50 55 60

Thr Gln Ala Pro Ser Ser Pro Ala Ile Glu Asp Ser Ser Phe Val Ser
 65 70 75 80

Arg Phe Gly Pro Asp Glu Pro Arg Lys Gly Ser Asp Val Leu Val Glu
 85 90 95

Ala Leu Glu Arg Glu Gly Val Thr Asn Val Phe Ala Tyr Pro Gly Gly
 100 105 110

Ala Ser Met Glu Ile His Gln Ala Leu Thr Arg Ser Lys Thr Ile Arg
 115 120 125

Asn Val Leu Pro Arg His Glu Gln Gly Gly Val Phe Ala Ala Glu Gly
 130 135 140

Tyr Ala Arg Ala Thr Gly Lys Val Gly Val Cys Ile Ala Thr Ser Gly
 145 150 155 160

Pro Gly Ala Thr Asn Leu Val Ser Gly Leu Ala Asp Ala Leu Leu Asp
 165 170 175

Ser Val Pro Leu Val Ala Ile Thr Gly Gln Val Pro Arg Arg Met Ile
 180 185 190

Gly Thr Asp Ala Phe Gln Glu Thr Pro Ile Val Glu Val Thr Arg Ser
 195 200 205

Ile Thr Lys His Asn Tyr Leu Val Leu Asp Val Glu Asp Ile Pro Arg
 210 215 220

Ile Val Lys Glu Ala Phe Phe Leu Ala Asn Ser Gly Arg Pro Gly Pro
 225 230 235 240
 Val Leu Ile Asp Leu Pro Lys Asp Ile Gln Gln Gln Leu Val Val Pro
 245 250 255
 Asp Trp Asp Arg Pro Phe Lys Leu Gly Gly Tyr Met Ser Arg Leu Pro
 260 265 270
 Lys Ser Lys Phe Ser Thr Asn Glu Val Gly Leu Leu Glu Gln Ile Val
 275 280 285
 Arg Leu Met Ser Glu Ser Lys Lys Pro Val Leu Tyr Val Gly Gly Gly
 290 295 300
 Cys Leu Asn Ser Ser Glu Glu Leu Arg Arg Phe Val Glu Leu Thr Gly
 305 310 315 320
 Ile Pro Val Ala Ser Thr Leu Met Gly Leu Gly Ser Tyr Pro Cys Asn
 325 330 335
 Asp Glu Leu Ser Leu His Met Leu Gly Met His Gly Thr Val Tyr Ala
 340 345 350
 Asn Tyr Ala Val Asp Lys Ala Asp Leu Leu Leu Ala Phe Gly Val Arg
 355 360 365
 Phe Asp Asp Arg Val Thr Gly Lys Leu Glu Ala Phe Ala Ser Arg Ala
 370 375 380
 Lys Ile Val His Ile Asp Ile Asp Ser Ala Glu Ile Gly Lys Asn Lys
 385 390 395 400
 Gln Pro His Val Ser Ile Cys Ala Asp Val Lys Leu Ala Leu Arg Gly
 405 410 415
 Met Asn Lys Ile Leu Glu Ser Arg Ile Gly Lys Leu Asn Leu Asp Phe
 420 425 430
 Ser Lys Trp Arg Glu Glu Leu Gly Glu Gln Lys Lys Glu Phe Pro Leu
 435 440 445
 Ser Phe Lys Thr Phe Gly Asp Ala Ile Pro Pro Gln Tyr Ala Ile Gln
 450 455 460
 Val Leu Asp Glu Leu Thr Asn Gly Asn Ala Ile Ile Ser Thr Gly Val
 465 470 475 480
 Gly Gln His Gln Met Trp Ala Ala Gln His Tyr Lys Tyr Arg Asn Pro
 485 490 495
 Arg Gln Trp Leu Thr Ser Gly Gly Leu Gly Ala Met Gly Phe Gly Leu
 500 505 510
 Pro Ala Ala Ile Gly Ala Ala Val Ala Arg Pro Asp Ala Val Val Val
 515 520 525

Asp Ile Asp Gly Asp Gly Ser Phe Ile Met Asn Val Gln Glu Leu Ala
530 535 540

Thr Ile Arg Val Glu Asn Leu Pro Val Lys Ile Met Leu Leu Asn Asn
545 550 555 560

Gln His Leu Gly Met Val Val Gln Trp Glu Asp Arg Phe Tyr Lys Ala
565 570 575

Asn Arg Ala His Thr Tyr Leu Gly Asn Pro Ser Lys Ser Ala Asp Ile
580 585 590

Phe Pro Asp Met Leu Lys Phe Ala Glu Ala Cys Asp Ile Pro Ser Ala
595 600 605

Arg Val Ser Asn Val Ala Asp Leu Arg Ala Ala Ile Gln Thr Met Leu
610 615 620

Asp Thr Pro Gly Pro Tyr Leu Leu Asp Val Ile Val Pro His Gln Glu
625 630 635 640

His Val Leu Pro Met Ile Pro Ser Gly Ala Gly Phe Lys Asp Thr Ile
645 650 655

Thr Glu Gly Asp Gly Arg Thr Ser
660

<210> 3

<211> 1995

<212> DNA

<213> Beta vulgaris

<220>

<221> misc_feature

<222> (1)..(1994)

<223> 4D6834 W574

<220>

<221> CDS

<222> (1)..(1995)

<400> 3

atg gcg gct acc ttc aca aac cca aca ttt tcc cct tcc tca act cca	48
Met Ala Ala Thr Phe Thr Asn Pro Thr Phe Ser Pro Ser Ser Thr Pro	
1 5 10 15	
tta acc aaa acc cta aaa tcc caa tot tcc atc tct tca acc ctc ccc	96
Leu Thr Lys Thr Leu Lys Ser Gln Ser Ser Ile Ser Ser Thr Leu Pro	
20 25 30	
ttt tcc acc cct ccc aaa acc cca act cca ctc ttt cac cgt ccc ctc	144
Phe Ser Thr Pro Pro Lys Thr Thr Pro Leu Phe His Arg Pro Leu	
35 40 45	
caa atc tca tcc tcc caa tcc cac aaa tca tcc gcc att aaa aca caa	192
Gln Ile Ser Ser Ser Gln Ser His Lys Ser Ser Ala Ile Lys Thr Gln	
50 55 60	

```

act caa gca cct tct tct cca gct att gaa gat tca tct ttc gtt tct      240
Thr Gln Ala Pro Ser Ser Pro Ala Ile Glu Asp Ser Ser Phe Val Ser
65              70              75              80

cga ttt ggc cct gat gaa ccc aga aaa ggg tcc gat gtc ctc gtt gaa      288
Arg Phe Gly Pro Asp Glu Pro Arg Lys Gly Ser Asp Val Leu Val Glu
85              90              95

gct ctt gag cgt gaa ggt gtt acc aat gtg ttt gct tac cct ggt ggt      336
Ala Leu Glu Arg Glu Gly Val Thr Asn Val Phe Ala Tyr Pro Gly Gly
100             105             110

gca tct atg gaa atc cac caa gct ctc aca cgc tct aaa acc atc cgc      384
Ala Ser Met Glu Ile His Gln Ala Leu Thr Arg Ser Lys Thr Ile Arg
115             120             125

aat gtc ctc cct cgc cat gaa caa ggc ggg gtt ttc gcc gcc gag gga      432
Asn Val Leu Pro Arg His Gln Gly Gly Val Phe Ala Ala Glu Gly
130             135             140

tat gct aga gct act gga aag gtt ggt gtc tgc att gcg act tct ggt      480
Tyr Ala Arg Ala Thr Gly Lys Val Gly Val Cys Ile Ala Thr Ser Gly
145             150             155             160

cct ggt gct acc aac ctc gta tca ggt ctt gct gac gct ctc ctt gat      528
Pro Gly Ala Thr Asn Leu Val Ser Gly Leu Ala Asp Ala Leu Leu Asp
165             170             175

tct gtc cct ctt gtt gcc atc act ggc caa gtt cca cgc cgt atg att      576
Ser Val Pro Leu Val Ala Ile Thr Gly Gln Val Pro Arg Arg Met Ile
180             185             190

ggc act gat gct ttt cag gag act cca att gtt gag gtg aca agg tct      624
Gly Thr Asp Ala Phe Gln Glu Thr Pro Ile Val Glu Val Thr Arg Ser
195             200             205

att act aag cat aat tat tta gtt ttg gat gta gag gat att cct aga      672
Ile Thr Lys His Asn Tyr Leu Val Leu Asp Val Glu Asp Ile Pro Arg
210             215             220

att gtt aag gaa gcc ttt ttt tta gct aat tct ggt agg cct gga cct      720

Ile Val Lys Glu Ala Phe Phe Leu Ala Asn Ser Gly Arg Pro Gly Pro
225             230             235             240

gtt ttg att gat ctt cct aaa gat att cag cag caa ttg gtt gtt cct      768
Val Leu Ile Asp Leu Pro Lys Asp Ile Gln Gln Gln Leu Val Val Pro
245             250             255

gat tgg gat agg cct ttt aag ttg ggt ggg tat atg tct agg ctg cca      816
Asp Trp Asp Arg Pro Phe Lys Leu Gly Gly Tyr Met Ser Arg Leu Pro
260             265             270

aag tcc aag ttt tcg acg aat gag gtt gga ctt ctt gag cag att gtg      864
Lys Ser Lys Phe Ser Thr Asn Glu Val Gly Leu Leu Glu Gln Ile Val
275             280             285

agg ttg atg agt gag tcg aag aag cct gtc ttg tat gtg gga ggt ggg      912
Arg Leu Met Ser Glu Ser Lys Lys Pro Val Leu Tyr Val Gly Gly Gly
290             295             300

tgt ttg aat tct agt gag gag ttg agg aga ttt gtt gag ttg aca ggg      960
Cys Leu Asn Ser Ser Glu Glu Leu Arg Arg Phe Val Glu Leu Thr Gly
305             310             315             320

att ccg gtg gct agt act ttg atg ggg ttg ggg tct tac cct tgt aat      1008
Ile Pro Val Ala Ser Thr Leu Met Gly Leu Gly Ser Tyr Pro Cys Asn
325             330             335

gat gaa ctg tct ctt cat atg ttg ggg atg cac ggg act gtt tat gcc      1056
Asp Glu Leu Ser Leu His Met Leu Gly Met His Gly Thr Val Tyr Ala
340             345             350

```

aat tat gcg gtg gat aag gcg gat ttg ttg ctt gct ttc ggg gtt agg Asn Tyr Ala Val Asp Lys Ala Asp Leu Leu Leu Ala Phe Gly Val Arg 355 360 365	1104
ttt gat gat cgt gtg acc ggg aag ctc gag gcg ttt gct agc cgt gct Phe Asp Asp Arg Val Thr Gly Lys Leu Glu Ala Phe Ala Ser Arg Ala 370 375 380	1152
aag att gtg cat att gat att gac tct gct gag att ggg aag aac aag Lys Ile Val His Ile Asp Ile Asp Ser Ala Glu Ile Gly Lys Asn Lys 385 390 395 400	1200
cag ccc cat gtg tcc att tgt gct gat gtt aaa ttg gca ttg cgg ggt Gln Pro His Val Ser Ile Cys Ala Asp Val Lys Leu Ala Leu Arg Gly 405 410 415	1248
atg aat aag att ctg gag tct aga ata ggg aag ctg aat ttg gat ttc Met Asn Lys Ile Leu Glu Ser Arg Ile Gly Lys Leu Asn Leu Asp Phe 420 425 430	1296
tcc aag tgg aga gaa gaa tta ggt gag cag aag aag gaa ttc cca ctg Ser Lys Trp Arg Glu Glu Leu Gly Glu Gln Lys Lys Glu Phe Pro Leu 435 440 445	1344
agt ttt aag aca ttt ggg gat gca att cct cca caa tat gcc att cag Ser Phe Lys Thr Phe Gly Asp Ala Ile Pro Pro Gln Tyr Ala Ile Gln 450 455 460	1392
gtg ctt gat gag ttg acc aat ggt aat gct att ata agt act ggt gtt Val Leu Asp Glu Leu Thr Asn Gly Asn Ala Ile Ile Ser Thr Gly Val 465 470 475 480	1440
ggg cag cac caa atg tgg gct gcg cag cat tac aag tac aga aac cct Gly Gln His Gln Met Trp Ala Ala Gln His Tyr Lys Tyr Arg Asn Pro 485 490 495	1488
cgc caa tgg ctg acc tct ggt ggg ttg ggg gct atg ggg ttt ggg cta Arg Gln Trp Leu Thr Ser Gly Gly Leu Gly Ala Met Gly Phe Gly Leu 500 505 510	1536
cca gcc gcc att gga gct gca gtt gct cga cca gat gca gtg gtt gtc Pro Ala Ala Ile Gly Ala Ala Val Ala Arg Pro Asp Ala Val Val Val 515 520 525	1584
gat att gat ggg gat ggc agt ttt att atg aat gtt caa gag ttg gct Asp Ile Asp Gly Asp Gly Ser Phe Ile Met Asn Val Gln Glu Leu Ala 530 535 540	1632
aca att agg gtg gaa aat ctc cca gtt aag ata atg ctg cta aac aat Thr Ile Arg Val Glu Asn Leu Pro Val Lys Ile Met Leu Leu Asn Asn 545 550 555 560	1680
caa cat tta ggt atg gtt gtc caa ttg gaa gat agg ttc tat aaa gct Gln His Leu Gly Met Val Val Gln Leu Glu Asp Arg Phe Tyr Lys Ala 565 570 575	1728
aac cgg gca cat aca tac ctt gga aac cct tcc aaa tct gct gat atc Asn Arg Ala His Thr Tyr Leu Gly Asn Pro Ser Lys Ser Ala Asp Ile 580 585 590	1776
ttc cct gat atg ctc aaa ttc gct gag gca tgt gat att cct tct gcc Phe Pro Asp Met Leu Lys Phe Ala Glu Ala Cys Asp Ile Pro Ser Ala 595 600 605	1824
cgt gtt agc aac gtg gct gat ttg agg gcc gcc att caa aca atg ttg Arg Val Ser Asn Val Ala Asp Leu Arg Ala Ala Ile Gln Thr Met Leu 610 615 620	1872
gat act cca ggg ccg tac ctg ctc gat gtg att gta ccg cat caa gag Asp Thr Pro Gly Pro Tyr Leu Leu Asp Val Ile Val Pro His Gln Glu 625 630 635 640	1920
cat gtg ttg cct atg att cca agt ggt gcc ggt ttc aag gat acc att Val Val Trp Phe Met Ile Phe Ser Gln Ala Gln Phe Thr Asn Thr Thr 645 650 655 660	1968

His Val Leu Pro Met Ile Pro Ser Gly Ala Gly Phe Lys Asp Thr Ile
645 650 655

aca gag ggt gat gga aga acc tct taa
Thr Glu Gly Asp Gly Arg Thr Ser
660

1995

$\langle 210 \rangle$ 4

<211> 664

<212> PRT

<213> Beta vulgaris

<400> 4

Met Ala Ala Thr Phe Thr Asn Pro Thr Phe Ser Pro Ser Ser Thr Pro
1 5 10 15

Leu Thr Lys Thr Leu Lys Ser Gln Ser Ser Ile Ser Ser Thr Leu Pro
20 25 30

Phe Ser Thr Pro Pro Lys Thr Pro Thr Pro Leu Phe His Arg Pro Leu
35 40 45

Gln Ile Ser Ser Ser Gln Ser His Lys Ser Ser Ala Ile Lys Thr Gln
50 55 60

Thr Gln Ala Pro Ser Ser Pro Ala Ile Glu Asp Ser Ser Phe Val Ser
65 70 75 80

Arg Phe Gly Pro Asp Glu Pro Arg Lys Gly Ser Asp Val Leu Val Glu
85 90 95

Ala Leu Glu Arg Glu Gly Val Thr Asn Val Phe Ala Tyr Pro Gly Gly
100 105 110

Ala Ser Met Glu Ile His Gln Ala Leu Thr Arg Ser Lys Thr Ile Arg
115 120 125

Asn Val Leu Pro Arg His Glu Gln Gly Gly Val Phe Ala Ala Glu Gly
130 135 140

Tyr Ala Arg Ala Thr Gly Lys Val Gly Val Cys Ile Ala Thr Ser Gly
145 150 155 160

Pro Gly Ala Thr Asn Leu Val Ser Gly Leu Ala Asp Ala Leu Leu Asp
165 170 175

Ser Val Pro Leu Val Ala Ile Thr Gly Gln Val Pro Arg Arg Met Ile
180 185 190

Gly Thr Asp Ala Phe Gln Glu Thr Pro Ile Val Glu Val Thr Arg Ser
195 200 205

Ile Thr Lys His Asn Tyr Leu Val Leu Asp Val Glu Asp Ile Pro Arg
210 215 220

Ile Val Lys Glu Ala Phe Phe Leu Ala Asn Ser Gly Arg Pro Gly Pro
225 230 235 240

Val Leu Ile Asp Leu Pro Lys Asp Ile Gln Gln Gln Leu Val Val Pro
245 250 255

Asp Trp Asp Arg Pro Phe Lys Leu Gly Gly Tyr Met Ser Arg Leu Pro
260 265 270

Lys Ser Lys Phe Ser Thr Asn Glu Val Gly Leu Leu Glu Gln Ile Val
275 280 285

Arg Leu Met Ser Glu Ser Lys Lys Pro Val Leu Tyr Val Gly Gly Gly
290 295 300

Cys Leu Asn Ser Ser Glu Glu Leu Arg Arg Phe Val Glu Leu Thr Gly
305 310 315 320

Ile Pro Val Ala Ser Thr Leu Met Gly Leu Gly Ser Tyr Pro Cys Asn
325 330 335

Asp Glu Leu Ser Leu His Met Leu Gly Met His Gly Thr Val Tyr Ala
340 345 350

Asn Tyr Ala Val Asp Lys Ala Asp Leu Leu Leu Ala Phe Gly Val Arg
355 360 365

Phe Asp Asp Arg Val Thr Gly Lys Leu Glu Ala Phe Ala Ser Arg Ala
370 375 380

Lys Ile Val His Ile Asp Ile Asp Ser Ala Glu Ile Gly Lys Asn Lys
385 390 395 400

Gln Pro His Val Ser Ile Cys Ala Asp Val Lys Leu Ala Leu Arg Gly
405 410 415

Met Asn Lys Ile Leu Glu Ser Arg Ile Gly Lys Leu Asn Leu Asp Phe
420 425 430

Ser Lys Trp Arg Glu Glu Leu Gly Glu Gln Lys Lys Glu Phe Pro Leu
435 440 445

Ser Phe Lys Thr Phe Gly Asp Ala Ile Pro Pro Gln Tyr Ala Ile Gln
450 455 460

Val Leu Asp Glu Leu Thr Asn Gly Asn Ala Ile Ile Ser Thr Gly Val
465 470 475 480

Gly Gln His Gln Met Trp Ala Ala Gln His Tyr Lys Tyr Arg Asn Pro
485 490 495

Arg Gln Trp Leu Thr Ser Gly Gly Leu Gly Ala Met Gly Phe Gly Leu
500 505 510

Pro Ala Ala Ile Gly Ala Ala Val Ala Arg Pro Asp Ala Val Val Val
515 520 525

Asp Ile Asp Gly Asp Gly Ser Phe Ile Met Asn Val Gln Glu Leu Ala

530 535 540
 Thr Ile Arg Val Glu Asn Leu Pro Val Lys Ile Met Leu Leu Asn Asn
 545 550 555 560
 Gln His Leu Gly Met Val Val Gln Leu Glu Asp Arg Phe Tyr Lys Ala
 565 570 575
 Asn Arg Ala His Thr Tyr Leu Gly Asn Pro Ser Lys Ser Ala Asp Ile
 580 585 590
 Phe Pro Asp Met Leu Lys Phe Ala Glu Ala Cys Asp Ile Pro Ser Ala
 595 600 605
 Arg Val Ser Asn Val Ala Asp Leu Arg Ala Ala Ile Gln Thr Met Leu
 610 615 620
 Asp Thr Pro Gly Pro Tyr Leu Leu Asp Val Ile Val Pro His Gln Glu
 625 630 635 640
 His Val Leu Pro Met Ile Pro Ser Gly Ala Gly Phe Lys Asp Thr Ile
 645 650 655
 Thr Glu Gly Asp Gly Arg Thr Ser
 660

<210> 5

<211> 1995

<212> DNA

<213> Beta vulgaris

<220>

<221> misc_feature

<222> (1)..(1994)

<223> Pro Mutant

<220>

<221> CDS

<222> (1)..(1995)

<400> 5

atg gcg gct acc ttc aca aac cca aca ttt tcc cct tcc tca act cca	48
Met Ala Ala Thr Phe Thr Asn Pro Thr Phe Ser Pro Ser Ser Thr Pro	
1 5 10 15	
tta acc aaa acc cta aaa tcc caa tct tcc atc tct tca acc ctc ccc	96
Leu Thr Lys Thr Leu Lys Ser Gln Ser Ser Ile Ser Ser Thr Leu Pro	
20 25 30	
ttt tcc acc cct ccc aaa acc cca act cca ctc ttt cac cgt ccc ctc	144
Phe Ser Thr Pro Pro Lys Thr Thr Pro Leu Phe His Arg Pro Leu	
35 40 45	
caa atc tca tcc tcc caa tcc cac aaa tca tcc gcc att aaa aca caa	192
Gln Ile Ser Ser Ser Gln Ser His Lys Ser Ser Ala Ile Lys Thr Gln	
50 55 60	
act gaa gaa gct tat tat gaa gct att gaa gct taa tat tta att tat	240

acc caa gca cct cct cct cca gct att gaa gat tca cct ctc gct cct Thr Gln Ala Pro Ser Ser Pro Ala Ile Glu Asp Ser Ser Phe Val Ser 65 70 75 80	270
cga ttt ggc cct gat gaa ccc aga aaa ggg tcc gat gtc ctc gtt gaa Arg Phe Gly Pro Asp Glu Pro Arg Lys Gly Ser Asp Val Leu Val Glu 85 90 95	288
gct ctt gag cgt gaa ggt gtt acc aat gtg ttt gct tac cct ggt ggt Ala Leu Glu Arg Glu Gly Val Thr Asn Val Phe Ala Tyr Pro Gly Gly 100 105 110	336
gca tct atg gaa atc cac caa gct ctc aca cgc tct aaa acc atc cgc Ala Ser Met Glu Ile His Gln Ala Leu Thr Arg Ser Lys Thr Ile Arg 115 120 125	384
aat gtc ctc cct cgc cat gaa caa ggc ggg gtt ttc gcc gcc gag gga Asn Val Leu Pro Arg His Glu Gln Gly Gly Val Phe Ala Ala Glu Gly 130 135 140	432
tat gct aga gct act gga aag gtt ggt gtc tgc att gcg act tct ggt Tyr Ala Arg Ala Thr Gly Lys Val Gly Val Cys Ile Ala Thr Ser Gly 145 150 155 160	480
cct ggt gct acc aac ctc gta tca ggt ctt gct gac get ctc ctt gat Pro Gly Ala Thr Asn Leu Val Ser Gly Leu Ala Asp Ala Leu Leu Asp 165 170 175	528
tct gtc cct ctt gtt gcc atc act ggc caa gtt tca cgc cgt atg att Ser Val Pro Leu Val Ala Ile Thr Gly Gln Val Ser Arg Arg Met Ile 180 185 190	576
ggc act gat gct ttt cag gag act cca att gtt gag gtg aca agg tct Gly Thr Asp Ala Phe Gln Glu Thr Pro Ile Val Glu Val Thr Arg Ser 195 200 205	624
att act aag cat aat tat tta gtt ttg gat gta gag gat att cct aga Ile Thr Lys His Asn Tyr Leu Val Leu Asp Val Glu Asp Ile Pro Arg 210 215 220	672
att gtt aag gaa gcc ttt ttt tta gct aat tct ggt agg cct gga cct Ile Val Lys Glu Ala Phe Phe Leu Ala Asn Ser Gly Arg Pro Gly Pro 225 230 235 240	720
gtt ttg att gat ctt cct aaa gat att cag cag caa ttg gtt gtt cct Val Leu Ile Asp Leu Pro Lys Asp Ile Gln Gln Leu Val Val Pro 245 250 255	768
gat tgg gat agg cct ttt aag ttg ggt ggg tat atg tct agg ctg cca Asp Trp Asp Arg Pro Phe Lys Leu Gly Gly Tyr Met Ser Arg Leu Pro 260 265 270	816
aag tcc aag ttt tcg acg aat gag gtt gga ctt ctt gag cag att gtg Lys Ser Lys Phe Ser Thr Asn Glu Val Gly Leu Leu Glu Gln Ile Val 275 280 285	864
agg ttg atg agt gag tcg aag aag cct gtc ttg tat gtg gga ggt ggg Arg Leu Met Ser Glu Ser Lys Lys Pro Val Leu Tyr Val Gly Gly Gly 290 295 300	912
tgt ttg aat tct agt gag gag ttg agg aga ttt gtt gag ttg aca ggg Cys Leu Asn Ser Ser Glu Glu Leu Arg Arg Phe Val Glu Leu Thr Gly 305 310 315 320	960
att ccg gtg gct agt act ttg atg ggg ttg ggg tct tac cct tgt aat Ile Pro Val Ala Ser Thr Leu Met Gly Leu Gly Ser Tyr Pro Cys Asn 325 330 335	1008
gat gaa ctg tct ctt cat atg ttg ggg atg cac ggg act gtt tat gcc Asp Glu Leu Ser Leu His Met Leu Gly Met His Gly Thr Val Tyr Ala 340 345 350	1056
aat tat gcg gtg gat aag gcg gat ttg ttg ctt gct ttc ggg gtt agg Asn Tyr Ala Val Asp Lys Ala Asp Leu Leu Leu Ala Phe Gly Val Arg 355 360 365	1104
ttt gat gat cgt gtg acc ggg aag ctc gag gcg ttt gct agc cgt gct	1152

Phe 370	Asp	Arg	Val	Thr	Gly 375	Lys	Leu	Glu	Ala	Phe 380	Ser	Arg	Ala			
aag Lys 385	att Ile	gtg Val	cat His	att Ile	gat Asp 390	att Ile	gac Asp	tct Ser	gct Ala	gag Glu 395	att Ile	ggg Gly	aag Lys	aac Asn	aag Lys 400	1200
cag Gln	ccc Pro	cat His	gtg Val	tcc Ser 405	att Ile	tgt Cys	gct Ala	gat Asp	gtt Val 410	aaa Lys	ttg Leu	gca Ala	ttg Leu	cgg Arg 415	ggt Gly	1248
atg Met	aat Asn	aag Lys	att Ile 420	ctg Leu	gag Glu	tct Ser	aga Arg	ata Ile 425	ggg Gly	aag Lys	ctg Leu	aat Asn	ttg Leu	gat Asp 430	ttc Phe	1296
tcc Ser	aag Lys	tggt Trp 435	aga Arg	gaa Glu	gaa Glu	tta Leu	ggt Gly 440	gag Glu	cag Gln	aag Lys	aag Lys	gaa Glu 445	ttc Phe	cca Pro	ctg Leu	1344
agt Ser	ttt Phe 450	aag Lys	aca Thr	ttt Phe	ggg Gly 455	gat Asp	gca Ala	att Ile	cct Pro	cca Pro	caa Gln	tat Tyr	gcc Ala	att Ile	cag Gln	1392
gtg Val 465	ctt Leu	gat Asp	gag Glu	ttg Leu	acc Thr 470	aat Asn	ggt Gly	aat Asn	gct Ala	att Ile 475	ata Ile	agt Ser	act Thr	ggt Gly	gtt Val 480	1440
ggg Gly	cag Gln	cac His	caa Gln	atg Met 485	tggt Trp	gct Ala	gcg Ala	cag Gln	cat His 490	tac Tyr	aag Lys	tac Tyr	aga Arg	aac Asn 495	cct Pro	1488
cgc Arg	caa Gln	tggt Trp	ctg Leu	acc Thr 500	tct Ser	ggt Gly	ggg Gly	ttg Leu	ggg Gly 505	gct Ala	atg Met	ggg Gly	ttt Phe 510	ggg Gly	cta Leu	1536
cca Pro	gcc Ala	gcc Ala	att Ile	gga Gly	gct Ala	gca Ala	gtt Val 520	gct Ala	cga Arg	cca Pro	gat Asp	gca Ala	gtg Val	gtt Val	gtc Val	1584
gat Asp	att Ile 530	gat Asp	ggg Gly	gat Asp	ggc Gly	agt Ser	ttt Phe 535	att Ile	atg Met	aat Asn	gtt Val 540	caa Gln	gag Glu	ttg Leu	gct Ala	1632
aca Thr	att Ile	agg Arg	gtg Val	gaa Glu	aat Asn	ctc Leu	cca Pro	gtt Val	aag Lys	ata Ile	atg Met	ctg Leu	cta Leu	aac Asn	aat Asn	1680
545					550					555					560	
caa Gln	cat His	tta Leu	ggt Gly	atg Met	gtt Val	gtc Val	caa Gln	tggt Trp	gaa Glu	gat Asp	agg Arg	ttc Phe	tat Tyr	aaa Lys	gct Ala	1728
aac Asn	cgg Arg	gca Ala	cat His	aca Thr	tac Tyr	ctt Leu	gga Gly	aac Asn	cct Pro	tcc Ser	aaa Lys	tct Ser	gct Ala	gat Asp	atc Ile	1776
ttc Phe	cct Pro	gat Asp	atg Met	ctc Leu	aaa Lys	ttc Phe	gct Ala	gag Glu	gca Ala	tgt Cys	gat Asp	att Ile	cct Pro	tct Ser	gcc Ala	1824
cgt Arg	gtt Val	agc Ser	aac Asn	gtg Val	gct Ala	gat Asp	ttg Leu	agg Arg	gcc Ala	gcc Ala	att Ile	caa Gln	aca Thr	atg Met	ttg Leu	1872
gat Asp	act Thr	cca Pro	ggg Gly	ccg Pro	tac Tyr	ctg Leu	ctc Leu	gat Asp	gtg Val	att Ile	gta Val	ccg Pro	cat His	caa Gln	gag Glu	1920
cat His	gtg Val	ttg Leu	cct Pro	atg Met	att Ile	cca Pro	agt Ser	ggt Gly	gcc Ala	ggt Gly	ttc Phe	aag Lys	gat Asp	acc Thr	att Ile	1968
aca Thr	gag Glu	ggt Gly	gat Asp	gga Gly	aga Arg	acc Thr	tct Ser	taa								1995

660

<210> 6

<211> 664

<212> PRT

<213> Beta vulgaris

<400> 6

Met Ala Ala Thr Phe Thr Asn Pro Thr Phe Ser Pro Ser Ser Thr Pro
 1 5 10 15

Leu Thr Lys Thr Leu Lys Ser Gln Ser Ser Ile Ser Ser Thr Leu Pro
 20 25 30

Phe Ser Thr Pro Pro Lys Thr Pro Thr Pro Leu Phe His Arg Pro Leu
 35 40 45

Gln Ile Ser Ser Ser Gln Ser His Lys Ser Ser Ala Ile Lys Thr Gln
 50 55 60

Thr Gln Ala Pro Ser Ser Pro Ala Ile Glu Asp Ser Ser Phe Val Ser
 65 70 75 80

Arg Phe Gly Pro Asp Glu Pro Arg Lys Gly Ser Asp Val Leu Val Glu
 85 90 95

Ala Leu Glu Arg Glu Gly Val Thr Asn Val Phe Ala Tyr Pro Gly Gly
 100 105 110

Ala Ser Met Glu Ile His Gln Ala Leu Thr Arg Ser Lys Thr Ile Arg
 115 120 125

Asn Val Leu Pro Arg His Glu Gln Gly Gly Val Phe Ala Ala Glu Gly
 130 135 140

Tyr Ala Arg Ala Thr Gly Lys Val Gly Val Cys Ile Ala Thr Ser Gly
 145 150 155 160

Pro Gly Ala Thr Asn Leu Val Ser Gly Leu Ala Asp Ala Leu Leu Asp
 165 170 175

Ser Val Pro Leu Val Ala Ile Thr Gly Gln Val Ser Arg Arg Met Ile
 180 185 190

Gly Thr Asp Ala Phe Gln Glu Thr Pro Ile Val Glu Val Thr Arg Ser
 195 200 205

Ile Thr Lys His Asn Tyr Leu Val Leu Asp Val Glu Asp Ile Pro Arg
 210 215 220

Ile Val Lys Glu Ala Phe Phe Leu Ala Asn Ser Gly Arg Pro Gly Pro
 225 230 235 240

Val Leu Ile Asp Leu Pro Lys Asp Ile Gln Gln Gln Leu Val Val Pro

	245		250		255
Asp Trp Asp Arg Pro Phe Lys Leu Gly Gly Tyr Met Ser Arg Leu Pro	260		265		270
Lys Ser Lys Phe Ser Thr Asn Glu Val Gly Leu Leu Glu Gln Ile Val	275		280		285
Arg Leu Met Ser Glu Ser Lys Lys Pro Val Leu Tyr Val Gly Gly Gly	290		295		300
Cys Leu Asn Ser Ser Glu Glu Leu Arg Arg Phe Val Glu Leu Thr Gly	305		310		315
Ile Pro Val Ala Ser Thr Leu Met Gly Leu Gly Ser Tyr Pro Cys Asn	325		330		335
Asp Glu Leu Ser Leu His Met Leu Gly Met His Gly Thr Val Tyr Ala	340		345		350
Asn Tyr Ala Val Asp Lys Ala Asp Leu Leu Leu Ala Phe Gly Val Arg	355		360		365
Phe Asp Asp Arg Val Thr Gly Lys Leu Glu Ala Phe Ala Ser Arg Ala	370		375		380
Lys Ile Val His Ile Asp Ile Asp Ser Ala Glu Ile Gly Lys Asn Lys	385		390		395
Gln Pro His Val Ser Ile Cys Ala Asp Val Lys Leu Ala Leu Arg Gly	405		410		415
Met Asn Lys Ile Leu Glu Ser Arg Ile Gly Lys Leu Asn Leu Asp Phe	420		425		430
Ser Lys Trp Arg Glu Glu Leu Gly Glu Gln Lys Lys Glu Phe Pro Leu	435		440		445
Ser Phe Lys Thr Phe Gly Asp Ala Ile Pro Pro Gln Tyr Ala Ile Gln	450		455		460
Val Leu Asp Glu Leu Thr Asn Gly Asn Ala Ile Ile Ser Thr Gly Val	465		470		475
Gly Gln His Gln Met Trp Ala Ala Gln His Tyr Lys Tyr Arg Asn Pro	485		490		495
Arg Gln Trp Leu Thr Ser Gly Gly Leu Gly Ala Met Gly Phe Gly Leu	500		505		510
Pro Ala Ala Ile Gly Ala Ala Val Ala Arg Pro Asp Ala Val Val Val	515		520		525
Asp Ile Asp Gly Asp Gly Ser Phe Ile Met Asn Val Gln Glu Leu Ala	530		535		540
Thr Ile Arg Val Glu Asn Leu Pro Val Lys Ile Met Leu Leu Asn Asn					

545 550 555 560
 Gln His Leu Gly Met Val Val Gln Trp Glu Asp Arg Phe Tyr Lys Ala
 565 570 575
 Asn Arg Ala His Thr Tyr Leu Gly Asn Pro Ser Lys Ser Ala Asp Ile
 580 585 590
 Phe Pro Asp Met Leu Lys Phe Ala Glu Ala Cys Asp Ile Pro Ser Ala
 595 600 605
 Arg Val Ser Asn Val Ala Asp Leu Arg Ala Ala Ile Gln Thr Met Leu
 610 615 620
 Asp Thr Pro Gly Pro Tyr Leu Leu Asp Val Ile Val Pro His Gln Glu
 625 630 635 640
 His Val Leu Pro Met Ile Pro Ser Gly Ala Gly Phe Lys Asp Thr Ile
 645 650 655
 Thr Glu Gly Asp Gly Arg Thr Ser
 660

<210> 7

<211> 1995

<212> DNA

<213> Beta vulgaris

<220>

<221> misc_feature

<222> (1)..(1994)

<223> 4D6834 al2 WT

<220>

<221> CDS

<222> (1)..(1995)

<400> 7

atg gcg gct acc ttc aca aac cca aca ttt tcc cct tcc tca act caa	48
Met Ala Ala Thr Phe Thr Asn Pro Thr Phe Ser Pro Ser Ser Thr Gln	
1 5 10 15	
tta acc aaa acc cta aaa tcc caa tct tcc att tct tca acc ctc ccc	96
Leu Thr Lys Thr Leu Lys Ser Gln Ser Ser Ile Ser Ser Thr Leu Pro	
20 25 30	
ttt tcc acc cct ccc aaa acc cca act cca ctc ttt cac cgt ccc ctc	144
Phe Ser Thr Pro Pro Lys Thr Pro Thr Pro Leu Phe His Arg Pro Leu	
35 40 45	
caa atc tca tcc tcc caa tcc cac aaa tca tcc gcc att aaa aca caa	192
Gln Ile Ser Ser Ser Gln Ser His Lys Ser Ser Ala Ile Lys Thr Gln	
50 55 60	
act caa gca cct tct tct cca gct att gaa gat tca tct ttc gtt tct	240
Thr Gln Ala Pro Ser Ser Pro Ala Ile Glu Asp Ser Ser Phe Val Ser	
65 70 75 80	
cga ttt ggc cct gat gaa ccc aga aaa ggg tcc gat gtc ctc gtt gaa	288

Arg Phe Gly Pro Asp Glu Pro Arg Lys Gly Ser Asp Val Leu Val Glu	
85 90 95	
gct ctt gag cgt gaa ggt gtt acc aat gtg ttt gct tac cct ggt ggt	336
Ala Leu Glu Arg Glu Gly Val Thr Asn Val Phe Ala Tyr Pro Gly Gly	
100 105 110	
gca tct atg gaa atc cac caa gct ctg acg cgc tct aaa acc atc cgc	384
Ala Ser Met Glu Ile His Gln Ala Leu Thr Arg Ser Lys Thr Ile Arg	
115 120 125	
aat gtc ctc ccc cgc cat gaa caa ggc ggg gtt ttc gcc gcc gag gga	432
Asn Val Leu Pro Arg His Glu Gln Gly Gly Val Phe Ala Ala Glu Gly	
130 135 140	
tat gct aga gct act gga aag gtt ggt gtc tgc att gcg act tct ggt	480
Tyr Ala Arg Ala Thr Gly Lys Val Gly Val Cys Ile Ala Thr Ser Gly	
145 150 155 160	
cct ggt gct acc aac ctc gta tca ggt ctt gct gac gct ctc ctt gat	528
Pro Gly Ala Thr Asn Leu Val Ser Gly Leu Ala Asp Ala Leu Asp	
165 170 175	
tct gtc cct ctt gtt gcc atc act ggc caa gtt cca cgc cgt atg att	576
Ser Val Pro Leu Val Ala Ile Thr Gly Gln Val Pro Arg Arg Met Ile	
180 185 190	
ggc act gat gct ttt cag gag act cca att gtt gag gta aca agg tct	624
Gly Thr Asp Ala Phe Gln Glu Thr Pro Ile Val Glu Val Thr Arg Ser	
195 200 205	
att act aag cat aat tat ttg gtt ttg gat gta gaa gat att cct aga	672
Ile Thr Lys His Asn Tyr Leu Val Leu Asp Val Glu Asp Ile Pro Arg	
210 215 220	
att gtt aag gaa gcc ttt ttt tta gct aat tct ggc agg cct gga cct	720
Ile Val Lys Glu Ala Phe Phe Leu Ala Asn Ser Gly Arg Pro Gly Pro	
225 230 235 240	
gtt ttg att gat ctt cct aaa gat att cag cag caa ctg gtt gtt cct	768
Val Leu Ile Asp Leu Pro Lys Asp Ile Gln Gln Leu Val Val Pro	
245 250 255	
gat tgg gat agg cct ttt aag ttg ggt ggg tat atg tct agg ctg cca	816
Asp Trp Asp Arg Pro Phe Lys Leu Gly Tyr Met Ser Arg Leu Pro	
260 265 270	
aag tcc aag ttt tcg acg aat gag gtt gga ctt ctt gag cag att gtg	864
Lys Ser Lys Phe Ser Thr Asn Glu Val Gly Leu Leu Glu Gln Ile Val	
275 280 285	
agg ttg atg agt gag tcg aag aag cct gtc ttg tat gtg gga ggt ggg	912
Arg Leu Met Ser Glu Ser Lys Lys Pro Val Leu Tyr Val Gly Gly Gly	
290 295 300	
tgt ttg aat tct agt gag gag ttg agg aga ttt gtt gag ttg aca ggg	960
Cys Leu Asn Ser Ser Glu Glu Leu Arg Arg Phe Val Glu Leu Thr Gly	
305 310 315 320	
att ccg gtg gct agt act ttg atg ggg ttg ggg tct tac cct tgt aat	1008
Ile Pro Val Ala Ser Thr Leu Met Gly Leu Gly Ser Tyr Pro Cys Asn	
325 330 335	
gat gaa ctg tct ctt cat atg ttg ggg atg cac ggc act gtt tat gcc	1056
Asp Glu Leu Ser Leu His Met Leu Gly Met His Gly Thr Val Tyr Ala	
340 345 350	
aat tat gcg gtg gat aag gcg gat ttg ttg ctt gct ttc ggg gtt agg	1104
Asn Tyr Ala Val Asp Lys Ala Asp Leu Leu Leu Ala Phe Gly Val Arg	
355 360 365	
ttt gat gat cgt gtg act ggg aag ctc gag gcg ttt gct agc cgt gct	1152
Phe Asp Asp Arg Val Thr Gly Lys Leu Glu Ala Phe Ala Ser Arg Ala	
370 375 380	

aag att gtg cat att gat att gac tct gct gag att ggg aag aac aag Lys Ile Val His Ile Asp Ile Asp Ser Ala Glu Ile Gly Lys Asn Lys 385 390 395 400	1200
cag ccc cat gtg tcc att tgt gct gat gtt aaa ttg gca ttg cgg ggt Gln Pro His Val Ser Ile Cys Ala Asp Val Lys Leu Ala Leu Arg Gly 405 410 415	1248
atg aat aag att ctg gag tct aga ata ggg aag ctg aat ttg gat ttc Met Asn Lys Ile Leu Glu Ser Arg Ile Gly Lys Leu Asn Leu Asp Phe 420 425 430	1296
tcc agg tgg aga gaa gaa tta ggt gag cag aag aag gaa ttc cca ctg Ser Arg Trp Arg Glu Glu Leu Gly Glu Gln Lys Lys Glu Phe Pro Leu 435 440 445	1344
agt ttt aag aca ttt ggg gat gca atc cct cca caa tat gcc att cag Ser Phe Lys Thr Phe Gly Asp Ala Ile Pro Pro Gln Tyr Ala Ile Gln 450 455 460	1392
gtg ctt gat gag ttg acc aat ggt aat gct att ata agt act ggt gtt Val Leu Asp Glu Leu Thr Asn Gly Asn Ala Ile Ile Ser Thr Gly Val 465 470 475 480	1440
ggg cag cac caa atg tgg gct gcg cag cat tac aag tac aga aac cct Gly Gln His Gln Met Trp Ala Ala Gln His Tyr Lys Tyr Arg Asn Pro 485 490 495	1488
cgc caa tgg ctg acc tct ggt ggg ttg ggg gct atg ggg ttt ggg cta Arg Gln Trp Leu Thr Ser Gly Gly Leu Gly Ala Met Gly Phe Gly Leu 500 505 510	1536
cca gcc gcc att gga gct gca gtt gct cga cca gat gca gtg gtt gtc Pro Ala Ala Ile Gly Ala Ala Val Ala Arg Pro Asp Ala Val Val Val 515 520 525	1584
gat att gat ggg gat ggc agt ttt att atg aat gtt caa gag ttg gct Asp Ile Asp Gly Asp Gly Ser Phe Ile Met Asn Val Gln Glu Leu Ala 530 535 540	1632
aca att agg gtg gaa aat ctc cca gtt aag ata atg ctg cta aac aat Thr Ile Arg Val Glu Asn Leu Pro Val Lys Ile Met Leu Leu Asn Asn 545 550 555 560	1680
caa cat tta ggt atg gtt gtc caa tgg gaa gat agg ttc tat aaa gct Gln His Leu Gly Met Val Val Gln Trp Glu Asp Arg Phe Tyr Lys Ala 565 570 575	1728
aat cgg gca cat aca tac ctt gga aac cct tcc aaa tct gct gat atc Asn Arg Ala His Thr Tyr Leu Gly Asn Pro Ser Lys Ser Ala Asp Ile 580 585 590	1776
ttc cct gat atg ctc aaa ttc gct gag gca tgt gat att cct tct gcc Phe Pro Asp Met Leu Lys Phe Ala Glu Ala Cys Asp Ile Pro Ser Ala 595 600 605	1824
cgt gtt agc aac gtg gct gat ttg agg gcc gcc att caa aca atg ttg Arg Val Ser Asn Val Ala Asp Leu Arg Ala Ala Ile Gln Thr Met Leu 610 615 620	1872
gat act cca ggg cgg tac ctg ctc gat gtg att gta cgg cat caa gag Asp Thr Pro Gly Pro Tyr Leu Leu Asp Val Ile Val Pro His Gln Glu 625 630 635 640	1920
cat gtg ttg cct atg att cca agt ggt gcc ggt ttc aag gat acc att His Val Leu Pro Met Ile Pro Ser Gly Ala Gly Phe Lys Asp Thr Ile 645 650 655	1968
aca gag ggt gat gga aga acc tct taa Thr Glu Gly Asp Gly Arg Thr Ser 660	1995

<210> 8

<211> 664

<212> PRT

<213> Beta vulgaris

<400> 8

Met Ala Ala Thr Phe Thr Asn Pro Thr Phe Ser Pro Ser Ser Thr Gln
1 5 10 15

Leu Thr Lys Thr Leu Lys Ser Gln Ser Ser Ile Ser Ser Thr Leu Pro
20 25 30

Phe Ser Thr Pro Pro Lys Thr Pro Thr Pro Leu Phe His Arg Pro Leu
35 40 45

Gln Ile Ser Ser Ser Gln Ser His Lys Ser Ser Ala Ile Lys Thr Gln
50 55 60

Thr Gln Ala Pro Ser Ser Pro Ala Ile Glu Asp Ser Ser Phe Val Ser
65 70 75 80

Arg Phe Gly Pro Asp Glu Pro Arg Lys Gly Ser Asp Val Leu Val Glu
85 90 95

Ala Leu Glu Arg Glu Gly Val Thr Asn Val Phe Ala Tyr Pro Gly Gly
100 105 110

Ala Ser Met Glu Ile His Gln Ala Leu Thr Arg Ser Lys Thr Ile Arg
115 120 125

Asn Val Leu Pro Arg His Glu Gln Gly Gly Val Phe Ala Ala Glu Gly
130 135 140

Tyr Ala Arg Ala Thr Gly Lys Val Gly Val Cys Ile Ala Thr Ser Gly
145 150 155 160

Pro Gly Ala Thr Asn Leu Val Ser Gly Leu Ala Asp Ala Leu Leu Asp
165 170 175

Ser Val Pro Leu Val Ala Ile Thr Gly Gln Val Pro Arg Arg Met Ile
180 185 190

Gly Thr Asp Ala Phe Gln Glu Thr Pro Ile Val Glu Val Thr Arg Ser
195 200 205

Ile Thr Lys His Asn Tyr Leu Val Leu Asp Val Glu Asp Ile Pro Arg
210 215 220

Ile Val Lys Glu Ala Phe Phe Leu Ala Asn Ser Gly Arg Pro Gly Pro
225 230 235 240

Val Leu Ile Asp Leu Pro Lys Asp Ile Gln Gln Gln Leu Val Val Pro
245 250 255

Asp Trp Asp Arg Pro Phe Lys Leu Gly Gly Tyr Met Ser Arg Leu Pro
260 265 270

Lys	Ser	Lys	Phe	Ser	Thr	Asn	Glu	Val	Gly	Leu	Leu	Glu	Gln	Ile	Val
		275					280					285			
Arg	Leu	Met	Ser	Glu	Ser	Lys	Lys	Pro	Val	Leu	Tyr	Val	Gly	Gly	Gly
	290					295					300				
Cys	Leu	Asn	Ser	Ser	Glu	Glu	Leu	Arg	Arg	Phe	Val	Glu	Leu	Thr	Gly
305					310					315					320
Ile	Pro	Val	Ala	Ser	Thr	Leu	Met	Gly	Leu	Gly	Ser	Tyr	Pro	Cys	Asn
				325					330					335	
Asp	Glu	Leu	Ser	Leu	His	Met	Leu	Gly	Met	His	Gly	Thr	Val	Tyr	Ala
			340					345					350		
Asn	Tyr	Ala	Val	Asp	Lys	Ala	Asp	Leu	Leu	Leu	Ala	Phe	Gly	Val	Arg
		355					360					365			
Phe	Asp	Asp	Arg	Val	Thr	Gly	Lys	Leu	Glu	Ala	Phe	Ala	Ser	Arg	Ala
	370					375					380				
Lys	Ile	Val	His	Ile	Asp	Ile	Asp	Ser	Ala	Glu	Ile	Gly	Lys	Asn	Lys
385					390					395					400
Gln	Pro	His	Val	Ser	Ile	Cys	Ala	Asp	Val	Lys	Leu	Ala	Leu	Arg	Gly
				405					410					415	
Met	Asn	Lys	Ile	Leu	Glu	Ser	Arg	Ile	Gly	Lys	Leu	Asn	Leu	Asp	Phe
			420					425					430		
Ser	Arg	Trp	Arg	Glu	Glu	Leu	Gly	Glu	Gln	Lys	Lys	Glu	Phe	Pro	Leu
		435					440					445			
Ser	Phe	Lys	Thr	Phe	Gly	Asp	Ala	Ile	Pro	Pro	Gln	Tyr	Ala	Ile	Gln
	450					455					460				
Val	Leu	Asp	Glu	Leu	Thr	Asn	Gly	Asn	Ala	Ile	Ile	Ser	Thr	Gly	Val
465					470					475					480
Gly	Gln	His	Gln	Met	Trp	Ala	Ala	Gln	His	Tyr	Lys	Tyr	Arg	Asn	Pro
				485					490					495	
Arg	Gln	Trp	Leu	Thr	Ser	Gly	Gly	Leu	Gly	Ala	Met	Gly	Phe	Gly	Leu
			500					505					510		
Pro	Ala	Ala	Ile	Gly	Ala	Ala	Val	Ala	Arg	Pro	Asp	Ala	Val	Val	Val
		515					520					525			
Asp	Ile	Asp	Gly	Asp	Gly	Ser	Phe	Ile	Met	Asn	Val	Gln	Glu	Leu	Ala
	530					535					540				
Thr	Ile	Arg	Val	Glu	Asn	Leu	Pro	Val	Lys	Ile	Met	Leu	Leu	Asn	Asn
545					550					555					560
Gln	His	Leu	Gly	Met	Val	Val	Gln	Trp	Glu	Asp	Arg	Phe	Tyr	Lys	Ala

```

      565      570      575
Asn Arg Ala His Thr Tyr Leu Gly Asn Pro Ser Lys Ser Ala Asp Ile
      580      585      590
Phe Pro Asp Met Leu Lys Phe Ala Glu Ala Cys Asp Ile Pro Ser Ala
      595      600      605
Arg Val Ser Asn Val Ala Asp Leu Arg Ala Ala Ile Gln Thr Met Leu
      610      615      620
Asp Thr Pro Gly Pro Tyr Leu Leu Asp Val Ile Val Pro His Gln Glu
      625      630      635      640
His Val Leu Pro Met Ile Pro Ser Gly Ala Gly Phe Lys Asp Thr Ile
      645      650      655
Thr Glu Gly Asp Gly Arg Thr Ser
      660

```

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO9510178A \[0007\] \[0049\] \[0104\] \[0105\]](#)
- [WO9802527A \[0012\] \[0103\] \[0103\] \[0130\]](#)
- [WO2012049268A \[0014\]](#)
- [EP2013076618W \[0140\]](#)

Non-patent literature cited in the description

- **KRENS et al.**Theor. appl. Genet., 1990, vol. 79, 390-396 [\[0018\]](#)
- **GUREL et al.**Critical reviews in Plant Science, 2008, vol. 27, 108-140 [\[0018\]](#)
- **HALL et al.**J. Exp. Botany, 1997, vol. 48, 255-263 [\[0018\]](#)

- Entwicklungsstadien mono- und dikotyle Pflanzen Biologische Bundesanstalt für Land und Forstwirtschaft 20010000 [0131]

Patentkrav

1. Fremgangsmåde til frembringelse af en sukkerroeplantemutant, der er resistent over for én eller flere hæmmere af enzymet acetolactatsyntase (ALS), hvilken fremgangsmåde omfatter følgende trin:

- 5 - opnåelse af protoplaster fra stomatale læbeceller isoleret fra en sukkerroeplante;
 - anvendelse, i en *in vitro*-kultur af protoplasterne, af en sammensætning, der omfatter én eller flere ALS-hæmmere ved en koncentration, der er letal for mere end 99 % af de *in vitro*-dyrkede celler; og
10 - regenerering af sukkerroeplanter fra de overlevende celler af de *in vitro*-dyrkede celler,

hvor de stomatale læbecelleprotoplaster er forudvalgte på grund af deres evne til at regenerere en sukkerroeplante, hvor de forudvalgte protoplaster har en sandsynlighed på mere end 10 % (antal voksende protoplaster:samlet antal protoplaster sat til dyrkning) for at opdeles og regenerere et levedygtigt kallus, hvor kalli opnået fra protoplasterne har mere end 10 % (antal
15 skuddannende kalli:samlet antal kalli) evne til at udvikle skud, og hvor ALS-hæmmeren/hæmmerne anvendes til mere end 20.000.000 af protoplasterne.

2. Fremgangsmåde ifølge krav 1, hvor trinnet med udvælgelse af stomatale læbecelleprotoplaster, der kan regenerere en sukkerroeplante, omfatter undertrinnene med
20 isolering af stomatale læbecelleprotoplaster fra sukkerroeplanter af forskellige genotyper og måling for hver genotype af andelen af de protoplaster, der vokser, når protoplasterne dyrkes *in vitro*.

3. Fremgangsmåde ifølge krav 1 eller 2, der endvidere omfatter trinnet med sekvensering af de
25 regenererede planters genom fra de overlevende *in vitro*-dyrkede celler og/eller med sekvensering af ALS-genet for identificering af en mutation i ALS-genet.

4. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, der endvidere omfatter trinnet med udvælgelse af regenererede sukkerroeplanter med en mutation i ALS-genet.

30 **5.** Fremgangsmåde ifølge et hvilket som helst af de foregående krav 3 til 4, hvor den regenererede sukkerroeplante har én eller flere mutationer i ALS-genet ved positioner, der koder for aminosyrerne valgt fra gruppen bestående af glycine 112, alanin 113, methionin 115, arginin 133, valin 187, arginin 190, alanin 196, phenylalanin 197, lysin 247, methionin 346,

histidin 347, arginin 368, aspartat 370, aspartat 371, arginin 372, methionin 565, valin 566, phenylalanin 573, serin 648 og glycin 649.

6. Fremgangsmåde ifølge et hvilket som helst af de foregående krav 3 til 4, hvor den regenererede sukkerroeplante har én mutation i ALS-genet ved positionen, der koder for prolin 188, og/eller én mutation i ALS-genet ved positionen, der koder for tryptophan 569.

7. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, der omfatter et forudgående trin med reduktion af den koncentration, hvorved sammensætningen, der omfatter den ene eller flere ALS-hæmmere, er letal for mindst 99 % af de *in vitro*-dyrkede celler.

8. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvor den ene eller flere ALS-hæmmere anvendes i en *in vitro*-kultur af mere end 50.000.000 stomatale læbecelleprotoplaster.

9. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvor sammensætningen, der omfatter én eller flere ALS-hæmmere, omfatter foramsulfuron, fortrinsvis hvor foramsulfuron anvendes ved en koncentration i intervallet fra 10^{-9} mol/l til 10^{-6} mol/l, og/eller ethoxysulfuron.

10. Fremgangsmåde til frembringelse af en sukkerroeplantemutant, der er resistent over for et herbicid, hvilken fremgangsmåde omfatter følgende trin:

- opnåelse af protoplaster fra stomatale læbeceller isoleret fra en sukkerroeplante;
- anvendelse, i en *in vitro*-kultur af protoplasterne, af en sammensætning, der omfatter herbicidet ved en koncentration, der er letal for mere end 99 % af de *in vitro*-dyrkede celler;
- regenerering af sukkerroeplanter fra de overlevende celler af de *in vitro*-dyrkede celler; og
- udvælgelse af regenererede sukkerroeplanter med en mutation i det gen, der koder for det peptid, der er målet for herbicidet,

hvor de stomatale læbecelleprotoplaster er forudvalgte for deres evne til at regenerere en sukkerroeplante, hvor de forudvalgte protoplaster har en sandsynlighed på mere end 10 % (antal voksende protoplaster:samlet antal protoplaster sat til dyrkning) for at deles og regenerere et levedygtigt kallus, hvor kalli opnået fra protoplasterne har mere end 10 % (antal

skuddannende kalli:samlet antal kalli) evne til at udvikle skud, og hvor herbicidet anvendes til mere end 20.000.000 af protoplasterne, hvor herbicidet er valgt fra gruppen bestående af 4-HPPD-hæmmere, hæmmere af carotenoidbiosyntesen, hæmmere af EPSP-syntase, hæmmere af fotosystem II, hæmmere af fotosystem I, hæmmere af celledeling, hæmmere af mikrotubulussamling, hæmmere af protoporphyrinogen-oxidase, hæmmere af Acetyl CoA-carboxylase, cellevægssyntesehæmmere, hæmmer af glutaminsyntetase og syntetisk auxin.


5
10 **11.** Muteret sukkerroeplante, der omfatter SEQ ID NO: 3 på én allel af ALS-genet og SEQ ID NO: 5 på den anden allel af ALS-genet, hvor den muterede sukkerroeplante ikke udelukkende er opnået ved en i alt væsentlig biologisk proces.

12. Stomatal læbecelleprotoplast isoleret fra den muterede plante ifølge krav 11.

15 **13.** Anvendelse af protoplasterne ifølge krav 12 til indføring af ét eller flere yderligere genetiske træk.

DRAWINGS

Fig. 1



Line WT

Line C

Line B

Line A