



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**


(86) Date de dépôt PCT/PCT Filing Date: 2016/11/17
(87) Date publication PCT/PCT Publication Date: 2017/05/26
(85) Entrée phase nationale/National Entry: 2018/05/16
(86) N° demande PCT/PCT Application No.: US 2016/062422
(87) N° publication PCT/PCT Publication No.: 2017/087608
(30) Priorités/Priorities: 2015/11/20 (US62/257,964);
2016/04/11 (US62/320,805)

(51) Cl.Int./Int.Cl. *C07D 401/14* (2006.01),
C07D 217/04 (2006.01), *C07D 217/16* (2006.01),
C07D 217/24 (2006.01), *C07D 231/56* (2006.01),
C07D 239/88 (2006.01), *C07D 401/04* (2006.01),
C07D 401/06 (2006.01), *C07D 401/12* (2006.01),
C07D 405/12 (2006.01), *C07D 471/04* (2006.01),
C07D 495/04 (2006.01)

(71) Demandeur/Applicant:
VITAE PHARMACEUTICALS, INC., US

(72) Inventeurs/Inventors:
CLAREMON, DAVID A., US;
DILLARD, LAWRENCE WAYNE, US;
FAN, YI, US; ...

(54) Titre : MODULATEURS DE ROR-GAMMA
(54) Title: MODULATORS OF ROR-GAMMA



(57) **Abrégé/Abstract:**

Provided are novel compounds of Formula I: pharmaceutically acceptable salts thereof, and pharmaceutical compositions thereof, which are useful in the treatment of diseases and disorders mediated by ROR γ . Also provided are pharmaceutical compositions comprising the novel compounds of Formula I and methods for their use in treating one or more inflammatory, metabolic, autoimmune and other diseases or disorders.

(72) **Inventeurs(suite)/Inventors(continued):** JIA, LANQI, US; SINGH, SURESH B., US; TICE, COLIN M., US;
XU, ZHENRONG, US; YUAN, JING, US; ZHUANG, LINGHANG, US

(74) **Agent:** GOWLING WLG (CANADA) LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property
Organization
International Bureau



WIPO | PCT



(10) International Publication Number
WO 2017/087608 A8

(43) International Publication Date
26 May 2017 (26.05.2017)

(51) International Patent Classification:

<i>C07D 401/14</i> (2006.01)	<i>C07D 401/06</i> (2006.01)
<i>C07D 217/04</i> (2006.01)	<i>C07D 401/12</i> (2006.01)
<i>C07D 217/16</i> (2006.01)	<i>C07D 405/12</i> (2006.01)
<i>C07D 217/24</i> (2006.01)	<i>C07D 239/88</i> (2006.01)
<i>C07D 231/56</i> (2006.01)	<i>C07D 471/04</i> (2006.01)
<i>C07D 401/04</i> (2006.01)	<i>C07D 495/04</i> (2006.01)

(21) International Application Number:

PCT/US2016/062422

(22) International Filing Date:

17 November 2016 (17.11.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/257,964	20 November 2015 (20.11.2015)	US
62/320,805	11 April 2016 (11.04.2016)	US

(71) Applicant: **VITAE PHARMACEUTICALS, INC.**
[US/US]; 400 Interpace Parkway, Parsippany, New Jersey
07054 (US).

(72) Inventors: **CLAREMON, David, A.**; 1508 Aidenn Lair Road, Maple Glen, PA 19002 (US). **DILLARD, Lawrence, Wayne**; 496 Kings Road, Yardley, PA 19067 (US). **FAN, Yi**; 40 Bittersweet Drive, Doylestown, PA 18901 (US). **JIA, Lanqi**; 22 Beaver Hill Road, Horsham, PA 19044 (US). **SINGH, Suresh, B.**; 4 Adams Road, Kendall Park, NJ 08824 (US). **TICE, Colin, M.**; 1325 Pinebrook Court, Maple Glen, PA 19002 (US). **XU, Zhen-rong**; 3224 Riding Court, Chalfont, PA 18914 (US). **YUAN, Jing**; 537 Candlemaker Way, Lansdale, PA 19446 (US). **ZHUANG, Linghang**; 3135 Fox Drive, Chalfont, PA 18914 (US).

(74) Agents: **DAVIS, Steven, G.** et al.; McCarter & English, LLP, 265 Franklin Street, Boston, MA 02110 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*


(48) Date of publication of this corrected version:

6 July 2017

(15) Information about Correction:

see Notice of 6 July 2017

(54) Title: MODULATORS OF ROR-GAMMA



(57) Abstract: Provided are novel compounds of Formula I: pharmaceutically acceptable salts thereof, and pharmaceutical compositions thereof, which are useful in the treatment of diseases and disorders mediated by ROR γ . Also provided are pharmaceutical compositions comprising the novel compounds of Formula I and methods for their use in treating one or more inflammatory, metabolic, autoimmune and other diseases or disorders.

WO 2017/087608 A8

MODULATORS OF ROR-GAMMA

RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application No. 62/320805, filed April 11, 2016, and U.S. Provisional Application No. 62/257964, filed November 20, 2015, both of which are incorporated herein by reference.

TECHNICAL FIELD

[0002] The present disclosure is directed to novel retinoic acid receptor-related orphan receptor gamma (“ROR γ ” or “ROR-gamma”) modulators, processes for their preparation, pharmaceutical compositions containing these modulators, and their use in the treatment of inflammatory, metabolic, autoimmune and other diseases mediated by ROR γ .

BACKGROUND

[0003] Retinoic acid receptor-related orphan receptors (RORs) are a subfamily of transcription factors in the steroid hormone nuclear receptor superfamily (Jetten & Joo (2006) Adv. Dev. Biol. 2006, 16, 313-355). The ROR family consists of ROR alpha (ROR α), ROR beta (ROR β) and ROR gamma (ROR γ), each encoded by a separate gene (in human: *RORA*, *RORB* and *RORC*, respectively; in mouse: *rora*, *rorb* and *rorc*, respectively). RORs contain four principal domains shared by the majority of nuclear receptors: an N-terminal domain, a highly conserved DNA-binding domain (DBD) consisting of two zinc finger motifs, a hinge domain, and a ligand binding domain (LBD). Each ROR gene generates several isoforms, differing only in their N-terminal domains. ROR γ has two isoforms: ROR γ 1 and ROR γ 2 (also known as ROR γ t). ROR γ refers to ROR γ 1 and/or ROR γ t. ROR γ 1 is expressed in a variety of tissues including thymus, muscle, kidney and liver, but ROR γ t is exclusively expressed in the cells of the immune system, has a critical role in thymopoiesis and the development of several secondary lymphoid tissues, and is a key regulator of Th17 cell differentiation (Jetten, 2009, Nucl. Recept. Signal., 7:e003, doi:10.1621/nrs.07003, Epub 2009 Apr 3).

[0004] Th17 cells are a subset of T helper cells which preferentially produce the pro-inflammatory cytokines IL-17A, IL-17F, IL-21 and IL-22. Th17 cells and their effector molecules, such as IL-17, IL-21, IL-22, GM-CSF and CCL20, are associated with the pathogenesis of several autoimmune and inflammatory diseases, such as rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, psoriasis, inflammatory bowel disease, allergy and asthma (Maddur *et al.*, 2012, Am. J. Pathol., 181:8-18). Recent findings support a role for IL17 and Th17 cells in the pathogenesis of acne (Thiboutot *et al.*, 2014, J. Invest. Dermatol.,

134(2):307-10, doi: 10.1038/jid.2013.400; Agak *et al.*, 2014, J. Invest. Dermatol., 134(2):366-73, doi: 10.1038/jid.2013.334, Epub 2013 Aug 7). Th17 cells are also potent inducers of inflammation associated with endometriosis, a chronic inflammatory disease (Hirata *et al.*, 2010, Endocrinol., 151:5468-5476; Hirata *et al.*, 2011, Fertil Steril., Jul;96(1):113-7, doi: 10.1016/j.fertnstert.2011.04.060, Epub 2011 May 20). Additionally, Th17 cells have a key role in the mouse autoimmune models of experimental autoimmune encephalomyelitis (EAE), collagen-induced arthritis (CIA) and adjuvant-induced arthritis (AIA) (Bedoya *et al.*, 2013, Clin. Dev. Immunol., 2013:986789. Epub 2013 Dec 26. Th17 cells are activated during inflammatory and autoimmune disease processes and are responsible for recruiting other inflammatory cell types, particularly neutrophils, to mediate pathology in target tissues (Miossec & Kolls, 2012, Nature Rev., 11:763-776; Korn *et al.*, 2009, Annu. Rev. Immunol., 27:485-517). Aberrant Th17 cell function has been implicated in a variety of autoimmune diseases, including multiple sclerosis and rheumatoid arthritis. Autoimmune disease is believed to arise from the disruption of the equilibrium between effector and regulatory T cells (Solt *et al.*, 2012, ACS Chem. Biol., 7:1515-1519, Epub 2012 July 9). The importance of ROR γ t to Th17 cell differentiation and the pathogenic role of Th17 cells is evidenced by the fact that ROR γ t-deficient mice have very few Th17 cells and have a reduction in severity of EAE (Ivanov *et al.*, 2006, Cell, 126:1121-1133).

[0005] Recently, IL-17-producing neutrophils have been identified as promoting inflammation leading to both microbial clearance and IL-17-associated tissue damage in the cornea and other tissues (Taylor *et al.*, 2014, J. Immunol, 192:3319-3327; Taylor *et al.*, 2014, Nat. Immunol., 15:143–151), supporting a role for compounds that inhibit ROR γ activity in the treatment of corneal ulcers and other diseases and disorders associated with IL-17 expressing neutrophils.

[0006] Circadian rhythms are daily cycles of behavioral and physiological changes that are regulated by endogenous circadian clocks. A number of studies have established links between nuclear receptor (including ROR γ) function and expression, the circadian regulatory circuitry, and the regulation of various physiological processes (Jetten (2009) *op. cit.*).

[0007] Obstructive sleep apnea syndrome (OSAS) is a chronic inflammatory disease regulated by T lymphocytes. OSAS patients have a significant increase in peripheral Th17 cell frequency, IL-17 and ROR γ t levels (Ye *et al.*, 2012, Mediators Inflamm., 815308, doi: 10.1155/2012/815308, Epub 2012 Dec 31).

[0008] A number of studies have provided evidence of a role of RORs in cancer. Mice deficient in the expression of ROR γ exhibit a high incidence of thymic lymphomas that

metastasize frequently to liver and spleen. High expression of Th17-associated genes (including ROR γ) and high levels of Th17 cells in the tumor microenvironment has been shown to correlate with a poor prognosis in various cancers, including lung, gastric, breast and colon cancer (Tosolini *et al.*, 2011, *Cancer Res.*, 71:1263-1271, doi: 10.1158/0008-5472.CAN-10-2907, Epub 2011 Feb 8; Su *et al.*, 2014, *Immunol. Res.*, 58:118-124, doi: 10.1007/s12026-013-8483-y, Epub 2014 Jan 9; Carmi *et al.*, 2011, *J. Immunol.*, 186:3462-3471, doi: 10.4049/jimmunol.1002901, Epub 2011 Feb 7; Chen *et al.*, 2013, *Histopathology*, 63:225-233, doi: 10.1111/his.12156, Epub 2013 Jun 6). Recent evidence also shows that ROR γ is overexpressed and amplified in metastatic castration-resistant prostate cancer tumors, and that ROR γ antagonists suppressed tumor growth in multiple androgen receptor-expressing xenograft prostate cancer models. See e.g., *Nature Medicine*, March 28, 2016, advance online publication, doi: 10.1038/nm.4070.


[0009] ROR γ has also been identified to have a regulatory role in lipid/glucose homeostasis, and has been implicated in metabolic syndrome, obesity (Meissburger *et al.*, 2011, *EMBO Mol. Med.*, 3:637–651), hepatosteatosis, insulin resistance and diabetes.

[0010] Further support for the role of ROR γ in the pathogenesis of inflammatory, metabolic, circadian effect, cancer, and autoimmune diseases and disorders can be found in the following references: Chang *et al.*, 2012, *J. Exp. Pharmacol.*, 4:141-148; Jetten *et al.*, 2013, *Frontiers Endocrinol.*, 4:1-8; Huh & Littman, 2012, *Eur. J. Immunol.*, 42:2232-2237; Martinez *et al.*, 2008, *Ann. N.Y. Acad. Sci.*, 1143:188-211; Pantelyushin *et al.*, 2012, *J. Clin. Invest.*, 122:2252-2256; Jetten & Ueda, 2002, *Cell Death Differen.*, 9:1167-1171; Solt *et al.*, 2010, *Curr. Opin. Lipidol.*, 21:204-211.

[0011] In light of the role that ROR γ plays in disease pathogenesis, inhibition of ROR γ activity and Th17 cell differentiation and activity, including IL17 production, will be of significant therapeutic benefit. It is therefore desirable to prepare compounds that inhibit ROR γ activity and hence have utility in the treatment of inflammatory, autoimmune, metabolic, circadian effect, cancer, and other diseases mediated by ROR γ , such as e.g., asthma, atopic dermatitis, acne, Crohn's disease, regional enteritis, ulcerative colitis, Sjögren's syndrome, uveitis, Behçet's disease, dermatomyositis, multiple sclerosis, ankylosing spondylitis, systemic lupus erythematosus, scleroderma, psoriasis, psoriatic arthritis, steroid resistant asthma and rheumatoid arthritis.

SUMMARY

[0012] It has now been found that compounds described herein, and pharmaceutically acceptable compositions thereof, are effective modulators of ROR γ (see *e.g.*, Tables 11-20). Such compounds include those of Formula I:



or a pharmaceutically acceptable salt thereof, wherein each of Cy², L², Hy, *, **, R¹⁰, p, X, n, L¹, and Cy¹ are as defined and described herein.


[0013] The provided compounds, and pharmaceutically acceptable compositions thereof, are modulators of ROR γ and are useful for treating a variety of diseases, disorders or conditions. Such diseases, disorders, or conditions include those described herein.

[0014] The provided compounds can be used alone (*i.e.*, as a monotherapy) or in combination with one or more other therapeutic agent effective for treating any of the indications described herein.


DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

1. General Description of Compounds

[0015] In certain embodiments, the present disclosure provides a compound of Formula I:



or a pharmaceutically acceptable salt thereof, wherein



U and V are each independently CR⁹ or N, provided that both are not CR⁹;

X is -C(O)NH- or -NHC(O)-;

n is 0, 1, 2, or 3;

p is 0, 1, 2, or 3;

* designates the attachment to L²;

** designates the attachment to X;

L² is a bond or is selected from CH₂, CH₂CH₂, CHMe, O, C(=O), CH(OH), and CH₂O, wherein the oxygen atom in CH₂O can either be attached to Cy² or Hy, provided that attachment to Hy occurs at a carbon atom on Hy;

L¹ is absent or is CR⁷R⁸;

Cy¹ is selected from aryl, heteroaryl, heterocyclyl, and cycloalkyl, wherein the aryl, heteroaryl, heterocyclyl, and cycloalkyl are each substituted with 1 to 3 groups independently selected from R⁵;

Cy² is selected from aryl, heteroaryl, monocyclic cycloalkyl, and monocyclic heterocyclyl, wherein the aryl, heteroaryl, monocyclic cycloalkyl, and monocyclic heterocyclyl are each optionally substituted with 1 to 3 groups independently selected from R⁶;

R² is (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₁-C₄)haloalkyl, or monocyclic cycloalkyl;

R^{2a} is H, (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₁-C₄)haloalkyl, or monocyclic cycloalkyl;

R⁵ and R⁶ are each independently selected from halogen, -CN, -OR^c, -NR^dR^e, -S(O)_kR^c, -NR^cS(O)₂R^c, -S(O)₂NR^dR^e, -C(=O)OR^c, -OC(=O)OR^c, -OC(=O)R^c, -OC(=S)OR^c, -C(=S)OR^c, -OC(=S)R^c, -C(=O)NR^dR^e, -NR^cC(=O)R^c, -C(=S)NR^dR^e, -NR^cC(=S)R^c, -NR^cC(=O)OR^c, -OC(=O)NR^dR^e, -NR^c(C=S)OR^c, -OC(=S)NR^dR^e, -NR^cC(=O)NR^dR^e, -NR^c(C=S)NR^dR^e, -C(=S)R^c, -C(=O)R^c, (C₁-C₆)alkyl, cycloalkyl, -(CH₂)₁₋₄-cycloalkyl, heterocyclyl, -(CH₂)₁₋₄-heterocyclyl, aryl, -NHC(=O)-heterocyclyl, -NHC(=O)-cycloalkyl, -(CH₂)₁₋₄-aryl, heteroaryl and -(CH₂)₁₋₄-heteroaryl,

wherein the alkyl, cycloalkyl, heterocyclyl, aryl or heteroaryl portion present in each of said (C₁-C₆)alkyl, cycloalkyl, -(CH₂)₁₋₄-cycloalkyl, heterocyclyl, -(CH₂)₁₋₄-heterocyclyl, aryl, -(CH₂)₁₋₄-aryl, heteroaryl and -(CH₂)₁₋₄-heteroaryl substituent for R⁶ are optionally substituted with halogen, OR^c, -NO₂, -CN, -NR^cC(=O)R^c, -NR^dR^e, -S(O)_kR^c, -C(=O)OR^c, -C(=O)NR^dR^e, -C(=O)R^c, (C₁-C₃)alkyl, halo(C₁-C₃)alkyl, (C₁-C₃)alkoxy(C₁-C₃)alkyl, (C₁-C₃)alkoxy, or halo(C₁-C₃)alkoxy;

each R^c is independently selected from hydrogen and (C₁-C₆)alkyl optionally substituted with hydroxy, (C₁-C₂)alkoxy, -C(O)NH₂, -C(O)O(C₁-C₃)alkyl, or 1 to 3 halogen;

each R^d and R^e is independently selected from hydrogen and (C_1-C_6) alkyl;

k is 0, 1 or 2;

any heterocyclyl or heteroaryl portion of Cy^1 or Cy^2 is further optionally substituted with =O;

R^7 and R^8 are each independently hydrogen, OR^c , $-C(=O)OR^c$, monocyclic heterocyclyl, halophenyl, quinolin-2(1H)one-4yl-methyl, or (C_1-C_3) alkyl, wherein the (C_1-C_3) alkyl is optionally substituted with OR^c , $-NR^dR^e$, $-C(=O)OR^c$, $-C(=O)NR^dR^e$, or halophenyl;

R^9 is hydrogen, (C_1-C_4) alkyl, (C_1-C_4) haloalkyl, (C_1-C_4) alkoxy, (C_1-C_4) haloalkoxy, halo, cyano, or monocyclic cycloalkyl; and

R^{10} is (C_1-C_4) alkyl, (C_1-C_4) haloalkyl, (C_1-C_4) alkoxy, (C_1-C_4) haloalkoxy, halo, cyano, or monocyclic cycloalkyl.

2. Compounds and Definitions

[0016] The terms “halo” and “halogen” as used herein refer to an atom selected from fluorine (fluoro, $-F$), chlorine (chloro, $-Cl$), bromine (bromo, $-Br$), and iodine (iodo, $-I$).

[0017] The term “alkyl”, used alone or as a part of a larger moiety such as *e.g.*, “haloalkyl”, means a saturated monovalent straight or branched hydrocarbon radical having, unless otherwise specified, 1-10 carbon atoms and includes, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, n-decyl and the like. “Monovalent” means attached to the rest of the molecule at one point.

[0018] The term “haloalkyl” or “halocycloalkyl” include mono, poly, and perhaloalkyl groups where the halogens are independently selected from fluorine, chlorine, and bromine.

[0019] The terms “cycloalkyl” and “cycloaliphatic”, used alone or as part of a larger moiety, refer to a saturated cyclic aliphatic monocyclic or bicyclic ring system, as described herein, having from, unless otherwise specified, 3 to 10 carbon ring atoms. Monocyclic cycloalkyl groups include, without limitation, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptyl, cycloheptenyl, and cyclooctyl. It will be understood that when specified, optional substituents on a cycloalkyl or cycloaliphatic group may be present on any substitutable position and, include, *e.g.*, the position at which the cycloalkyl or cycloaliphatic group is attached.

[0020] The term “carbocycle”, “carbocyclyl”, “carbocyclo”, or “carbocyclic” used alone or as part of a larger moiety refer to saturated, partially saturated, or aromatic ring systems comprising all carbon atoms having, unless otherwise specified, a total of 3 to 10 ring members. It will be understood that when specified, optional substituents on a carbocycle, carbocyclyl,

carbocyclo, or carbocyclic may be present on any substitutable position and, include, *e.g.*, the position at which the cycloalkyl is attached.

[0021] The term “aryl” used alone or as part of a larger moiety as in “aralkyl”, “aralkoxy”, or “aryloxyalkyl”, refers to an aromatic carbocyclic ring system having, unless otherwise specified, a total of 6 to 10 ring members. The term “aryl” may be used interchangeably with the term “aryl ring”, “aryl group”, “aryl moiety,” or “aryl radical”. In certain embodiments of the present disclosure, “aryl” refers to an aromatic ring system which includes, but is not limited to, phenyl (abbreviated as “Ph”), naphthyl and the like. It will be understood that when specified, optional substituents on an aryl group may be present on any substitutable position and, include, *e.g.*, the position at which the aryl is attached.

[0022] The term “heteroaryl” used alone or as part of a larger moiety as in “heteroarylalkyl”, “heteroarylalkoxy”, or “heteroarylaminoalkyl”, refers to a 5-10 -membered aromatic radical containing 1-4 heteroatoms selected from N, O, and S and includes, for example, thienyl, furanyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolizinyl, purinyl, naphthyridinyl, and pteridinyl. The term “heteroaryl” may be used interchangeably with the terms “heteroaryl ring”, “heteroaryl group”, or “heteroaromatic”. The terms “heteroaryl” and “heteroar-”, as used herein, also include groups in which a heteroaromatic ring is fused to one or more aryl or heteroaryl rings. Nonlimiting examples include indolyl, indazolyl, benzimidazolyl, benzthiazolyl, quinolyl, quinazolinyl, quinoxalinyl, pyrrolopyridinyl, pyrrolopyrimidinyl, thienopyridinyl, and thienopyrimidinyl. A heteroaryl group may be mono- or bicyclic. It will be understood that when specified, optional substituents on a heteroaryl group may be present on any substitutable position and, include, *e.g.*, the position at which the heteroaryl is attached.

[0023] The term “heterocyclyl” means a 4-, 5-, 6- and 7-membered saturated or partially unsaturated heterocyclic ring containing 1 to 4 heteroatoms independently selected from N, O, and S. The terms “heterocycle”, “heterocyclyl”, “heterocyclyl ring”, “heterocyclic group”, “heterocyclic moiety”, and “heterocyclic radical”, are used interchangeably herein. A heterocyclyl ring can be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure. Examples of such saturated or partially unsaturated heterocyclic radicals include, without limitation, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, pyrrolidinyl, pyrrolidonyl, piperidinyl, oxazolidinyl, piperazinyl, dioxanyl, dioxolanyl, morpholinyl, dihydrofuranyl, dihydropyranyl, dihydropyridinyl, tetrahydropyridinyl, dihydropyrimidinyl, and tetrahydropyrimidinyl. A heterocyclyl group may be mono- or bicyclic.

Unless otherwise specified, bicyclic heterocyclyl groups include, *e.g.*, unsaturated heterocyclic radicals fused to another unsaturated heterocyclic radical or aromatic or heteroaryl ring, such as for example, tetrahydronaphthyridine, indolinone, dihydropyrrolotriazole, imidazopyrimidine, quinolinone, dioxaspirodecane. It will also be understood that when specified, optional substituents on a heterocyclyl group may be present on any substitutable position and, include, *e.g.*, the position at which the heterocyclyl is attached.

[0024] As used herein the terms “subject” and “patient” may be used interchangeably, and means a mammal in need of treatment, *e.g.*, companion animals (*e.g.*, dogs, cats, and the like), farm animals (*e.g.*, cows, pigs, horses, sheep, goats and the like) and laboratory animals (*e.g.*, rats, mice, guinea pigs and the like). Typically, the subject is a human in need of treatment.

[0025] Certain of the disclosed compounds may exist in various stereoisomeric forms. Stereoisomers are compounds that differ only in their spatial arrangement. Enantiomers are pairs of stereoisomers whose mirror images are not superimposable, most commonly because they contain an asymmetrically substituted carbon atom that acts as a chiral center. “Enantiomer” means one of a pair of molecules that are mirror images of each other and are not superimposable. Diastereomers are stereoisomers that contain two or more asymmetrically substituted carbon atoms. The symbol “*” in a structural formula represents the presence of a chiral carbon center. “R” and “S” represent the configuration of substituents around one or more chiral carbon atoms. Thus, “R*” and “S*” denote the relative configurations of substituents around one or more chiral carbon atoms.

[0026] As used herein, a hyphen (“-”) at the beginning or end of a recited group designates the point at which a recited group is attached to a defined group. For example, -SO₂-(C₁-C₃)alkyl-(C₂-C₆)cycloalkyl means that the group is attached via the sulfonyl.

[0027] “Racemate” or “racemic mixture” means a compound of equimolar quantities of two enantiomers, wherein such mixtures exhibit no optical activity, *i.e.*, they do not rotate the plane of polarized light.

[0028] “Geometric isomer” means isomers that differ in the orientation of substituent atoms in relationship to a carbon-carbon double bond, to a cycloalkyl ring, or to a bridged bicyclic system. Atoms (other than H) on each side of a carbon-carbon double bond may be in an E (substituents are on opposite sides of the carbon-carbon double bond) or Z (substituents are oriented on the same side) configuration. “R,” “S,” “S*,” “R*,” “E,” “Z,” “cis,” and “trans,” indicate configurations relative to the core molecule. When a disclosed compound is named or depicted by structure without indicating a particular geometric isomer form, it is to be

understood that the name or structure encompasses one geometric isomer free of other geometric isomers, mixtures of geometric isomers, or all geometric isomers.

[0029] The compounds of the herein may be prepared as individual enantiomers by either enantio-specific synthesis or resolved from an enantiomerically enriched mixture. Conventional resolution techniques include forming the salt of a free base of each isomer of an enantiomeric pair using an optically active acid (followed by fractional crystallization and regeneration of the free base), forming the salt of the acid form of each enantiomer of an enantiomeric pair using an optically active amine (followed by fractional crystallization and regeneration of the free acid), forming an ester or amide of each of the enantiomers of an enantiomeric pair using an optically pure acid, amine or alcohol (followed by chromatographic separation and removal of the chiral auxiliary), or resolving an enantiomeric mixture of either a starting material or a final product using various well known chromatographic methods.

[0030] When the stereochemistry of a disclosed compound is named or depicted by structure, the named or depicted stereoisomer is at least 60%, 70%, 80%, 90%, 99% or 99.9% by weight pure relative to all of the other stereoisomers. Percent by weight pure relative to all of the other stereoisomers is the ratio of the weight of one stereoisomer over the weight of the other stereoisomers. When a single enantiomer is named or depicted by structure, the depicted or named enantiomer is at least 60%, 70%, 80%, 90%, 99% or 99.9% by weight optically pure. Percent optical purity by weight is the ratio of the weight of the enantiomer over the weight of the enantiomer plus the weight of its optical isomer.

[0031] When the stereochemistry of a disclosed compound is named or depicted by structure, and the named or depicted structure encompasses more than one stereoisomer (*e.g.*, as in a diastereomeric pair), it is to be understood that one of the encompassed stereoisomers or any mixture of the encompassed stereoisomers are included. It is to be further understood that the stereoisomeric purity of the named or depicted stereoisomers at least 60%, 70%, 80%, 90%, 99% or 99.9% by weight pure relative to all of the other stereoisomers. The stereoisomeric purity in this case is determined by dividing the total weight in the mixture of the stereoisomers encompassed by the name or structure by the total weight in the mixture of all of the stereoisomers.

[0032] When a disclosed compound is named or depicted by structure without indicating the stereochemistry, and the compound has one chiral center, it is to be understood that the name or structure encompasses one enantiomer of compound free from the corresponding optical isomer,

a racemic mixture of the compound and mixtures enriched in one enantiomer relative to its corresponding optical isomer.

[0033] When a disclosed compound is named or depicted by structure without indicating the stereochemistry and *e.g.*, the compound has at least two chiral centers, it is to be understood that the name or structure encompasses one stereoisomer free of other stereoisomers, mixtures of stereoisomers, and mixtures of stereoisomers in which one or more stereoisomers is enriched relative to the other stereoisomer(s). For example, the name or structure may encompass one stereoisomer free of other diastereomers, mixtures of stereoisomers, and mixtures of stereoisomers in which one or more diastereomers is enriched relative to the other diastereomer(s).


[0034] The compounds of the herein may be present in the form of pharmaceutically acceptable salts. For use in medicines, the salts of the compounds of the invention refer to non-toxic “pharmaceutically acceptable salts.” Pharmaceutically acceptable salt forms include pharmaceutically acceptable acidic/anionic or basic/cationic salts.

[0035] Pharmaceutically acceptable basic/cationic salts include, the sodium, potassium, calcium, magnesium, diethanolamine, n-methyl-D-glucamine, L-lysine, L-arginine, ammonium, ethanolamine, piperazine and triethanolamine salts.

[0036] Pharmaceutically acceptable acidic/anionic salts include, *e.g.*, the acetate, benzenesulfonate, benzoate, bicarbonate, bitartrate, carbonate, citrate, dihydrochloride, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrobromide, hydrochloride, malate, maleate, malonate, mesylate, nitrate, salicylate, stearate, succinate, sulfate, tartrate, and tosylate.


3. Description of Exemplary Compounds

[0037] In a first embodiment, the present disclosure provides a compound of Formula I:



or a pharmaceutically acceptable salt thereof, wherein the variables are as described above.

[0038] In a second embodiment, the compound of Formula I is of the Formula II:





or a pharmaceutically acceptable salt thereof, wherein the variables are as described above for Formula I.

[0039] In a third embodiment, p in Formula I or II is 0, 1, or 2; and R^{10} is (C_1-C_4) alkyl or halo, wherein the remaining variables are as described above for Formula I and the second embodiment.



[0040] In a fourth embodiment, L^2 in Formula I or II is a bond or is selected from CH_2 , O, and CH_2O , wherein the oxygen atom in CH_2O is attached to a carbon atom on Hy, wherein the remaining variables are as described above for Formula I and the second or third embodiment.

[0041] In a fifth embodiment, the compound of Formula I is of the Formula III or IV:



or a pharmaceutically acceptable salt thereof, wherein each p is independently 0 or 1, and wherein the remaining variables in Formula III and IV are as described above for Formula I and the second, third, or fourth embodiment.



[0042] In a sixth embodiment, the compound of Formula **I** is of the Formula **IIIa** or **IVa**:



or a pharmaceutically acceptable salt thereof, wherein each p is independently 0 or 1, and wherein the remaining variables in Formula **IIIa** and **IVa** are as described above for Formula **I** and the second, third, fourth, or fifth embodiment.



[0043] In a seventh embodiment, R^{10} in Formula **I**, **II**, **III**, **IIIa**, **IV**, and **IVa** is (C_1-C_4) alkyl; and L^2 is a bond or CH_2 , wherein the remaining variables in Formula **I**, **II**, **III**, **IIIa**, **IV**, and **IVa** are as described above for Formula **I** and the second, third, fourth, fifth, or sixth embodiment.

[0044] In an eighth embodiment, the compound of Formula **I** is of the Formula **V** or **VI**:





or a pharmaceutically acceptable salt thereof, wherein the variables in structural Formula **V** and **VI** are as described above for Formula **I** and the second, third, or fourth embodiment.

[0045] In a ninth embodiment, the compound of Formula **I** is of the Formula **Va** or **Vla**:



or a pharmaceutically acceptable salt thereof, wherein the variables in structural Formula **Va** and **Vla** are as described above for Formula **I** and the second, third, fourth, or eighth embodiment.

[0046] In a tenth embodiment, the compound of Formula **I** is of Formula **Va'** or **Vla'**:




or a pharmaceutically acceptable salt thereof, wherein the variables in structural Formula **Va'** and **Vla'** are as described for Formula **I** and the second, third, fourth, eighth, or ninth embodiment.

[0047] In an eleventh embodiment, R^{2a} is hydrogen or (C_1-C_4) alkyl, wherein the variables in structural Formula **Va'** and **VIa'** are as described for Formula **I** and the second, third, fourth, eighth, ninth, or tenth embodiment.



[0048] In a twelfth embodiment, L^2 in Formula **V**, **VI**, **Va**, **VIa**, **Va'**, and **VIa'** is O or CH_2O , wherein the methylene portion of CH_2O is attached to Cy^2 , and wherein the variables in Formula **V**, **VI**, **Va**, **VIa**, **Va'**, and **VIa'** are as described above for Formula **I** and the second, third, fourth, eighth, ninth, tenth, or eleventh embodiment.


[0049] In a thirteenth embodiment, the compound of Formula **I** is of the Formula **VII**:



or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1, and wherein the remaining variables in Formula **VII** are as described above for Formula **I** and the second, third, or fourth embodiment.

[0050] In a fourteenth embodiment, the compound of Formula **I** is of the Formula **VIIa**, **VIIb**, or **VIIc**:







(VIIc);

or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1, and wherein the remaining variables in Formula **VIIa**, **VIIb**, and **VIIc** are as described above for Formula **I** and the second, third, fourth, fifth, or thirteenth embodiment.


[0051] In a fifteenth embodiment, the compound of Formula **I** is of the Formula **VIIa'**, **VIIb'**, or **VIIc'**:



(VIIa');






(VIIb'); or



(VIIc');

or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1, and wherein the remaining variables in Formula **VIIa'**, **VIIb'**, and **VIIc'** are as described above for Formula **I** and the second, third, fourth, thirteenth, or fourteenth embodiment.


[0052] In a sixteenth embodiment, the compound of Formula **I** is of the Formula **VIIa''**, **VIIb''**, or **VIIc''**:



or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1, and wherein the remaining variables in Formula **VIIa''**, **VIIb''**, and **VIIc''** are as described above for Formula **I** and the second, third, fourth, thirteenth, fourteenth, or fifteenth embodiment.



[0053] In a seventeenth embodiment, L^2 in Formula **VII**, **VIIa**, **VIIb**, **VIIc**, **VIIa'**, **VIIb'**, **VIIc'**, **VIIa''**, **VIIb''**, and **VIIc''** is O or CH_2O , wherein the methylene portion of CH_2O is attached to Cy^2 , wherein the remaining variables in Formula **VII**, **VIIa**, **VIIb**, **VIIc**, **VIIa'**, **VIIb'**, **VIIc'**, **VIIa''**, **VIIb''**, and **VIIc''** are as described above for Formula **I** and the second, third, fourth, thirteenth, fourteenth, fifteenth, or sixteenth embodiment.

[0054] In an eighteenth embodiment, the compound of Formula I is of the Formula VIII:




or a pharmaceutically acceptable salt thereof, wherein the variables in Formula VIII are as described above for Formula I and the second, third, or fourth embodiment.


[0055] In a nineteenth embodiment, the compound of Formula I is of the Formula VIIIa or VIIIb:



or a pharmaceutically acceptable salt thereof, wherein the variables in Formula VIIIa and VIIIb are as described above for Formula I and the second, third, fourth, or eighteenth embodiment.



[0056] In a twentieth embodiment, the compound of Formula I is of the Formula VIIIa' or VIIIb':





or a pharmaceutically acceptable salt thereof, wherein the variables in Formula **VIIIa'** and **VIIIb'** are as described above for Formula **I** and the second, third, fourth, eighteenth, or nineteenth embodiment.


[0057] In a twenty-first embodiment, the compound of Formula **I** is of the Formula **VIIIa''** or **VIIIb''**:



or a pharmaceutically acceptable salt thereof, wherein the variables in Formula **VIIIa''** and **VIIIb''** are as described above for Formula **I** and the second, third, fourth, eighteenth, nineteenth, or twentieth embodiment.


[0058] In a twenty-second embodiment, L^2 in Formula **VIII**, **VIIIa**, **VIIIb**, **VIIIa'**, **VIIIb'**, **VIIIa''**, and **VIIIb''** is CH_2 , wherein the variables in Formula **VIII**, **VIIIa**, **VIIIb**, **VIIIa'**, **VIIIb'**, **VIIIa''**, and **VIIIb''** are as described above for Formula **I** and the second, third, fourth, eighteenth, nineteenth, twentieth, or twenty-first embodiment.

[0059] In a twenty-third embodiment, the compound of Formula I is of the Formula IX:




or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1, and wherein the remaining variables are as described above for Formula I and the second, third, or fourth embodiment.

[0060] In a twenty-fourth embodiment, the compound of Formula I is of the Formula IXa:



or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1, and wherein the variables are as described above for Formula I and the second, third, fourth, or twenty-third embodiment.

[0061] In a twenty-fifth embodiment, the compound of Formula I is of the Formula IXa':



or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1, and wherein the variables are as described above for Formula I and the second, third, fourth, twenty-third, or twenty-fourth embodiment.

[0062] In a twenty-sixth embodiment, R¹⁰ in Formula IX, IXa, and IXa' is (C₁-C₄)alkyl; and L² is CH₂O, wherein the methylene portion of CH₂O is attached to Cy², and wherein the remaining variables are as described above for Formula I and the second, third, fourth, twenty-third, twenty-fourth, or twenty-fifth embodiment.

[0063] In a twenty-seventh embodiment, R^2 in Formula **I**, **II**, **III**, **IIIa**, **IV**, **IVa**, **VIII**, **VIIIa**, **VIIIb**, **VIIIa'**, **VIIIb'**, **VIIIa''** and **VIIIb''** is (C_1-C_4) alkyl or cyclopropyl, wherein the remaining variables are as described above for Formula **I** and the second, third, fourth, fifth, sixth, seventh, eighteenth, nineteenth, twentieth, twenty-first, or twenty-second embodiment.

[0064] In a twenty-eighth embodiment, R^7 in Formula **I**, **II**, **III**, **IIIa**, **IV**, **IVa**, **V**, **VI**, **Va**, **Vla**, **Va'**, **Vla**, **VII**, **VIIa**, **VIIb**, **VIIc**, **VIIa'**, **VIIb'**, **VIIc'**, **VIIa''**, **VIIb''**, **VIIc''**, **VIII**, **VIIIa**, **VIIIb**, **VIIIa'**, **VIIIb'**, **VIIIa''**, **VIIIb''**, **IX**, **IXa**, and **IXa'** is hydrogen, OR^c , or (C_1-C_3) alkyl, wherein the (C_1-C_3) alkyl is optionally substituted with OR^c or $-NR^dR^e$; and R^8 is hydrogen, wherein the remaining variables are as described above for Formula **I** and the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, or twenty-seventh embodiment.

[0065] In a twenty-eighth embodiment, R^7 in Formula **I**, **II**, **III**, **IIIa**, **IV**, **IVa**, **V**, **VI**, **Va**, **Vla**, **Va'**, **Vla**, **VII**, **VIIa**, **VIIb**, **VIIc**, **VIIa'**, **VIIb'**, **VIIc'**, **VIIa''**, **VIIb''**, **VIIc''**, **VIII**, **VIIIa**, **VIIIb**, **VIIIa'**, **VIIIb'**, **VIIIa''**, **VIIIb''**, **IX**, **IXa**, and **IXa'** is hydrogen, $-O(C_1-C_3)$ alkyl, or (C_1-C_3) alkyl, wherein the (C_1-C_3) alkyl is optionally substituted with OH, NH_2 or $-N(C_1-C_3alkyl)_2$; and R^8 is hydrogen, wherein the remaining variables are as described above for Formula **I** and the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, or twenty-eighth embodiment.

[0066] In a thirteenth embodiment, R^7 in Formula **I**, **II**, **III**, **IIIa**, **IV**, **IVa**, **V**, **VI**, **Va**, **Vla**, **Va'**, **Vla**, **VII**, **VIIa**, **VIIb**, **VIIc**, **VIIa'**, **VIIb'**, **VIIc'**, **VIIa''**, **VIIb''**, **VIIc''**, **VIII**, **VIIIa**, **VIIIb**, **VIIIa'**, **VIIIb'**, **VIIIa''**, **VIIIb''**, **IX**, **IXa**, and **IXa'** is hydrogen or hydroxy (C_1-C_3) alkyl; and R^8 is hydrogen, wherein the remaining variables are as described above for Formula **I** and the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, twenty-eighth, or twenty-ninth.

[0067] In a thirty-first embodiment, Cy^1 in Formula **I**, **II**, **III**, **IIIa**, **IV**, **IVa**, **V**, **VI**, **Va**, **Vla**, **Va'**, **Vla**, **VII**, **VIIa**, **VIIb**, **VIIc**, **VIIa'**, **VIIb'**, **VIIc'**, **VIIa''**, **VIIb''**, **VIIc''**, **VIII**, **VIIIa**, **VIIIb**, **VIIIa'**, **VIIIb'**, **VIIIa''**, **VIIIb''**, **IX**, **IXa**, and **IXa'** is aryl or heteroaryl, each substituted with 1 to 3 groups independently selected from R^5 , wherein the remaining variables are as

described above for Formula **I** and the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, twenty-eight, twenty-ninth embodiment or thirteenth.

[0068] In a thirty-second embodiment, Cy¹ in Formula **I**, **II**, **III**, **IIIa**, **IV**, **IVa**, **V**, **VI**, **Va**, **VIa**, **Va'**, **VIa**, **VII**, **VIIa**, **VIIb**, **VIIc**, **VIIa'**, **VIIb'**, **VIIc'**, **VIIa''**, **VIIb''**, **VIIc''**, **VIII**, **VIIIa**, **VIIIb**, **VIIIa'**, **VIIIb'**, **VIIIa''**, **VIIIb''**, **IX**, **IXa**, and **IXa'** is phenyl or pyridinyl, each substituted with 1 to 3 groups independently selected from R⁵, wherein the remaining variables are as described above for Formula **I** and the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, twenty-eight, twenty-ninth, thirtieth, or thirty-first embodiment.

[0069] In a thirty-third embodiment, at least one R⁵ in Formula **I**, **II**, **III**, **IIIa**, **IV**, **IVa**, **V**, **VI**, **Va**, **VIa**, **Va'**, **VIa**, **VII**, **VIIa**, **VIIb**, **VIIc**, **VIIa'**, **VIIb'**, **VIIc'**, **VIIa''**, **VIIb''**, **VIIc''**, **VIII**, **VIIIa**, **VIIIb**, **VIIIa'**, **VIIIb'**, **VIIIa''**, **VIIIb''**, **IX**, **IXa**, and **IXa'** is -SO₂-(C₁-C₃)alkyl, wherein the remaining variables are as described above for Formula **I** and the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, twenty-eight, twenty-ninth, thirtieth, thirty-first, or thirty-second embodiment.

[0070] In a thirty-fourth embodiment, Cy² in Formula **I**, **II**, **III**, **IIIa**, **IV**, **IVa**, **V**, **VI**, **Va**, **VIa**, **Va'**, **VIa**, **VII**, **VIIa**, **VIIb**, **VIIc**, **VIIa'**, **VIIb'**, **VIIc'**, **VIIa''**, **VIIb''**, **VIIc''**, **VIII**, **VIIIa**, **VIIIb**, **VIIIa'**, **VIIIb'**, **VIIIa''**, **VIIIb''**, **IX**, **IXa**, and **IXa'** is phenyl, pyrimidinyl, cyclohexyl, pyridinyl, tetrahydropyranyl, or piperidinyl, each optionally substituted with 1 to 3 groups independently selected from R⁶, wherein the remaining variables are as described above for Formula **I** and the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, twenty-eight, twenty-ninth, thirtieth, thirty-first, thirty-second, or thirty-third embodiment.

[0071] In a thirty-fifth embodiment, Cy² in Formula **I**, **II**, **III**, **IIIa**, **IV**, **IVa**, **V**, **VI**, **Va**, **VIa**, **Va'**, **VIa**, **VII**, **VIIa**, **VIIb**, **VIIc**, **VIIa'**, **VIIb'**, **VIIc'**, **VIIa''**, **VIIb''**, **VIIc''**, **VIII**, **VIIIa**, **VIIIb**, **VIIIa'**, **VIIIb'**, **VIIIa''**, **VIIIb''**, **IX**, **IXa**, and **IXa'** is phenyl or cyclohexyl, each

optionally substituted with 1 to 3 groups independently selected from R^6 , wherein the remaining variables are as described above for Formula I and the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, twenty-eighth, twenty-ninth, thirtieth, thirty-first, or thirty-second, thirty-third, or thirty-fourth embodiment.

[0072] In a thirty-sixth embodiment, R^5 in Formula I, II, III, IIIa, IV, IVa, V, VI, Va, VIa, Va', VIa, VII, VIIa, VIIb, VIIc, VIIa', VIIb', VIIc', VIIa'', VIIb'', VIIc'', VIII, VIIIa, VIIIb, VIIIa', VIIIb', VIIIa'', VIIIb'', IX, IXa, and IXa' is selected from halogen, -CN, -OR^c, -NR^dR^e, -NR^cS(O)₂R^c, -S(O)₂NR^dR^e, -C(=O)OR^c, -C(=O)NR^dR^e, -NR^cC(=O)R^c, -NR^cC(=O)OR^c, -OC(=S)NR^dR^e, -C(=O)R^c, -SO₂-(C₁-C₃)alkyl, and (C₁-C₄)alkyl optionally substituted with halogen; and R^6 is selected from halogen, -CN, -OR^c, -NR^dR^e, -NR^cS(O)₂R^c, -S(O)₂NR^dR^e, -C(=O)OR^c, -OC(=O)OR^c, -OC(=O)R^c, -C(=O)NR^dR^e, -NR^cC(=O)R^c, -C(=S)NR^dR^e, -NR^cC(=S)R^c, -NR^cC(=O)OR^c, -OC(=O)NR^dR^e, -NR^c(C=S)OR^c, -OC(=S)NR^dR^e, -NR^cC(=O)NR^dR^e, -NR^c(C=S)NR^dR^e, -C(=S)R^c, -C(=O)R^c, -SO₂-(C₁-C₃)alkyl, and (C₁-C₄)alkyl optionally substituted with halogen, wherein the remaining variables are as described above for Formula I and the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, twenty-eighth, twenty-ninth, thirtieth, thirty-first, or thirty-second, thirty-third, thirty-fourth, or thirty-fifth embodiment.

[0073] In a thirty-seventh embodiment, R^5 in Formula I, II, III, IIIa, IV, IVa, V, VI, Va, VIa, Va', VIa, VII, VIIa, VIIb, VIIc, VIIa', VIIb', VIIc', VIIa'', VIIb'', VIIc'', VIII, VIIIa, VIIIb, VIIIa', VIIIb', VIIIa'', VIIIb'', IX, IXa, and IXa' is -SO₂-(C₁-C₃)alkyl; R^6 is selected from -CN, halo, -C(=O)OR^c, OR^c, and (C₁-C₄)alkyl optionally substituted with halogen; and R^c is (C₁-C₄)alkyl optionally substituted with halogen, wherein the remaining variables are as described above for Formula I and the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, twenty-eighth, twenty-ninth, thirtieth, thirty-first, or thirty-second, thirty-third, thirty-fourth, thirty-fifth, or thirty-sixth embodiment.

[0074] In a thirty-eighth embodiment, R^6 in Formula I, II, III, IIIa, IV, IVa, V, VI, Va, VIa, Va', VIa, VII, VIIa, VIIb, VIIc, VIIa', VIIb', VIIc', VIIa'', VIIb'', VIIc'', VIII, VIIIa,

VIIIb, VIIIa', VIIIb', VIIIa'', VIIIb'', IX, IXa, and IXa' is CF₃, wherein the remaining variables are as described above for Formula **I** and the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, eleventh, twelfth, thirteenth, fourteenth, fifteenth, sixteenth, seventeenth, eighteenth, nineteenth, twentieth, twenty-first, twenty-second, twenty-third, twenty-fourth, twenty-fifth, twenty-sixth, twenty-seventh, twenty-eighth, twenty-ninth, thirtieth, thirty-first, or thirty-second, thirty-third, thirty-fourth, thirty-fifth, thirty-sixth, or thirty-seventh embodiment..

[0075] Specific examples of compounds are provided in the EXEMPLIFICATION.

Pharmaceutically acceptable salts as well as the neutral forms of these compounds are included herein.

[0076] In certain embodiments, the present disclosure provides a method of treating a patient (*e.g.*, a human) with a disorder mediated by ROR γ comprising the step of administering to the patient an effective amount of the compound with any compound described herein, or a pharmaceutically acceptable salt or composition thereof.

4. Uses, Formulation and Administration

Pharmaceutically acceptable compositions

[0077] According to another embodiment, the present disclosure provides a method of treating a subject (*e.g.*, a human) with a disorder mediated by ROR γ using a composition comprising a compound of Formula **I** and a pharmaceutically acceptable carrier, adjuvant, or vehicle. In certain embodiments, the amount of compound of Formula **I** in a provided composition is such that it is effective as an inverse agonist or antagonist to ROR γ in a biological sample or in a subject. In certain embodiments, a provided composition is formulated for administration to a subject in need of such composition. In some embodiments, a provided composition is formulated for oral administration to a subject.

[0078] The term “pharmaceutically acceptable carrier, adjuvant, or vehicle” refers to a non-toxic carrier, adjuvant, or vehicle that does not destroy the pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or vehicles that may be used in the compositions of this disclosure include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based

substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[0079] Compositions described herein may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques.

[0080] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[0081] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[0082] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[0083] In order to prolong the effect of a provided compound, it is often desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor

water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered compound form is accomplished by dissolving or suspending the compound in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the compound in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

[0084] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar--agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[0085] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled

gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[0086] Provided compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, *e.g.*, tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[0087] Dosage forms for topical or transdermal administration of a compound herein include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, ear drops, and eye drops are also contemplated as being within the scope herein. Additionally, the present disclosure contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[0088] Pharmaceutically acceptable compositions provided herein may be formulated for oral administration. Such formulations may be administered with or without food. In some embodiments, pharmaceutically acceptable compositions of this disclosure are administered without food. In other embodiments, pharmaceutically acceptable compositions of this disclosure are administered with food.

[0089] The amount of provided compounds that may be combined with carrier materials to produce a composition in a single dosage form will vary depending upon the patient to be treated and the particular mode of administration.

[0090] It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, the judgment of the treating physician, and the severity of the particular disease being treated. The amount of a provided compound in the composition will also depend upon the particular compound in the composition.

Uses of Compounds and Pharmaceutically Acceptable Compositions

[0091] Compounds and compositions described herein are generally useful for modulating ROR γ . Thus, in some embodiments, the present disclosure provides a method of treating inflammatory, metabolic and autoimmune diseases or disorders mediated by ROR γ , comprising administering a provided compound or composition. More particularly, the compounds and compositions described herein act as inverse agonists or antagonists of ROR γ .

[0092] As used herein, the terms “treatment,” “treat,” and “treating” refer to reversing, alleviating, delaying the onset of, or inhibiting the progress of a disease or disorder, or one or more symptoms thereof, as described herein. In some embodiments, treatment may be administered after one or more symptoms have developed, *i.e.*, therapeutic treatment. In other embodiments, treatment may be administered in the absence of symptoms. For example, treatment may be administered to a susceptible individual prior to the onset of symptoms (*e.g.*, in light of a history of symptoms and/or in light of genetic or other susceptibility factors), *i.e.*, prophylactic treatment. Treatment may also be continued after symptoms have resolved, for example to prevent or delay their recurrence.

[0093] Modulation of ROR γ (or to modulate ROR γ), means that a change or alteration in the activity of ROR γ has occurred from the administration of one or more of the compounds described herein. Modulation may be an upregulation (increase) or a downregulation (decrease) in the magnitude of the activity or function of ROR γ . Exemplary activities and functions include *e.g.*, binding characteristics, enzymatic activity, cell receptor activation, transcriptional activity, and signal transduction. In one aspect, the compounds described herein inhibit ROR γ . In further aspects, the compounds described herein act as agonists, antagonists, or inverse agonists of ROR γ .

[0094] In another aspect, compounds and compositions described herein are useful for reducing the amount of IL-17 in a subject. Thus, in some embodiments, provided herein are methods of reducing the amount of IL-17 in a subject comprising administering an effective amount of a provided compound or composition. ROR γ modulators disclosed in WO

2014/179564, WO 2015/116904, WO 2016/061160, PCT/US2016/045318, and U.S. patent application Nos. 14/933,468 and 14/933,524 can also be used in such methods.

[0095] In another aspect, compounds and compositions described herein are useful for inhibiting the synthesis of IL-17 in a subject. Thus, in some embodiments, provided herein are methods of inhibiting the synthesis of IL-17 in a subject comprising administering an effective amount of a provided compound or composition. ROR γ modulators disclosed in WO 2014/179564, WO 2015/116904, WO 2016/061160, PCT/US2016/045318, and U.S. patent application Nos. 14/933,468 and 14/933,524 can also be used in such methods.

[0096] Diseases and conditions treatable according to the methods herein include, but are not limited to, inflammatory, metabolic and autoimmune diseases or disorders mediated by ROR γ . These diseases and conditions include, for example, asthma, chronic obstructive pulmonary disease (COPD), bronchitis, allergic rhinitis, atopic dermatitis, contact dermatitis, acne, urticaria, hives, angioedema, cystic fibrosis, allograft rejection, multiple sclerosis, Balo's concentric (circular) sclerosis, Balo disease, leukoencephalitis periaxialis concentrica, encephalitis periaxialis concentrica, scleroderma, limited scleroderma, CREST syndrome, arthritis, rheumatoid arthritis, juvenile rheumatoid arthritis, reactive arthritis, Reiter's syndrome, osteoarthritis, ankylosing spondylitis, systemic lupus erythematosus (SLE), psoriasis, plaque psoriasis, guttate psoriasis, inverse psoriasis, pustular psoriasis, erythrodermic psoriasis, psoriatic epidermal hyperplasia, epidermal hyperplasia, Hashimoto's disease, pancreatitis, autoimmune diabetes, type I diabetes, autoimmune ocular disease, ulcerative colitis, Crohn's disease, regional enteritis, inflammatory bowel disease (IBD), inflammatory bowel syndrome (IBS), Sjögren's syndrome, optic neuritis, obesity, hepatosteatorrhea, adipose tissue-associated inflammation, insulin resistance, type II diabetes, neuromyelitis optica, myasthenia gravis, age related macular degeneration, dry eye, uveitis, Guillain-Barré syndrome, psoriatic arthritis (PsA), steroid resistant asthma, Graves' disease, scleritis, endometriosis, obstructive sleep apnea syndrome (OSAS), Behçet's disease, dermatomyositis, polymyositis, graft versus host disease, chronic graft versus host disease, acute graft versus host disease, primary biliary cirrhosis, liver fibrosis, non-alcoholic fatty liver disease (NAFLD), sarcoidosis, primary sclerosing cholangitis, autoimmune thyroid disease, autoimmune polyendocrine syndrome type I, autoimmune polyendocrine syndrome type II, celiac disease, celiac sprue, neuromyelitis, juvenile idiopathic arthritis, systemic sclerosis, myocardial infarction, pulmonary hypertension, osteoarthritis, cutaneous leishmaniasis, sinonasal polyposis, and cancer, including but not limited to lung cancer, gastric cancer, breast cancer and colon cancer. In one aspect, an exemplified form of cancer treatable

according to the methods herein also includes prostate cancer *e.g.*, (metastatic castration-resistant prostate cancer tumors). In another aspect, an exemplified form of cancer treatable according to the methods herein includes *e.g.*, malignant tumor, angiogenesis glaucoma, infantile hemangioma, multiple myeloma, acute myeloblastic leukemia, chronic sarcoma, chronic myelogenous leukemia, metastasis melanoma, Kaposi's sarcoma, vascular proliferation, cachexia, colorectal cancer (*e.g.*, familial colorectal cancer, hereditary nonpolyposis colorectal cancer, and gastrointestinal stromal tumor), lung cancer (*e.g.*, non-small cell lung cancer, small cell lung cancer and malignant mesothelioma), mesothelioma, pancreatic cancer (*e.g.*, pancreatic duct cancer), gastric cancer (*e.g.*, papillary adenocarcinoma, mucinous adenocarcinoma and adenosquamous carcinoma), breast cancer (*e.g.*, invasive ductal carcinoma, ductal carcinoma *in situ*, inflammatory breast cancer and metastatic breast cancer), ovarian cancer (*e.g.*, ovarian epithelial carcinoma, extragonadal germ cell tumor, ovarian germ cell tumor, and ovarian low malignant potential tumor), hormone-dependent prostate cancer, non-hormone dependent prostate cancer, liver cancer (*e.g.*, primary liver cancer and extrahepatic bile duct cancer), thyroid cancer (*e.g.*, medullary thyroid carcinoma), kidney cancer (*e.g.*, renal cell carcinoma, and transitional cell carcinoma in kidney and urinary duct), uterine cancer, endometrial cancer, brain tumor (*e.g.*, pineal astrocytoma, pilocytic astrocytoma, diffuse astrocytoma and anaplastic astrocytoma), melanoma, sarcoma, urinary bladder cancer, hematologic cancer, hypophyseal adenoma, glioma, acoustic neurinoma, retinoblastoma, head and neck cancer, head and neck squamous cell carcinoma, pharyngeal cancer, laryngeal cancer, cancer of the tongue, thymoma, esophagus cancer, duodenal cancer, colorectal cancer, rectal cancer, hepatoma, pancreatic endocrine tumor, cancer of the bile duct, gallbladder cancer, penile cancer, urinary duct cancer, testis tumor, vulvar cancer, cervical cancer, endometrial cancer, uterus sarcoma, vaginal cancer, skin cancer, fungoid mycosis, basal cell tumor, soft tissue sarcoma, malignant lymphoma, Hodgkin's disease, myelodysplastic syndrome, acute lymphocytic leukemia, chronic lymphocytic leukemia, chronic myelocytic leukemia, malignant myeloma, adult T cell leukemia, chronic bone marrow proliferative disease, pancreatic endocrine tumor, fibrous histiocytoma, leiomyosarcoma, rhabdomyosarcoma, cancer of unknown primary, cancer-driven myelopoiesis, tumor growth, and metastasis.

[0097] Diseases and disorders mediated by IL-17 expression, and which are treatable using the compounds described herein also include, *e.g.*, emphysema, lung fibrosis, idiopathic pulmonary fibrosis, retroperitoneal fibrosis, giant cell arteritis, giant cell myocarditis, arteriosclerosis, hepatitis, chronic active hepatitis, alcoholic hepatitis, alcoholic liver fibrosis,

alcoholic cirrhosis, viral hepatitis, hepatitis B viral liver disorder, autoimmune hepatitis, cartilage inflammation, bone degradation, juvenile arthritis, pauciarticular juvenile rheumatoid arthritis, polyarticular juvenile rheumatoid arthritis, systemic onset juvenile rheumatoid arthritis, spondyloarthritis, juvenile ankylosing spondylitis, juvenile enteropathic arthritis, juvenile reactive arthritis, juvenile Reiter's syndrome, seronegative enthesopathy and arthropathy (SEA) syndrome, juvenile dermatomyositis, juvenile psoriatic arthritis, juvenile scleroderma, juvenile systemic lupus erythematosus, juvenile vasculitis, pauciarticular rheumatoid arthritis, polyarticular rheumatoid arthritis, systemic onset rheumatoid arthritis, enteropathic arthritis, vasculitis, leukocytoclastic vasculitis, myositis, juvenile myositis, polymyositis, autoimmune myositis, osteoarthritis, polyarteritis nodosa, arteritis, Takayasu's arteritis, temporal arteritis, giant cell arteritis, testicular autoimmunity, polymyalgia rheumatica, rheumatic fever, sclerosis, primary biliary sclerosis, primary biliary cirrhosis, sclerosing cholangitis, primary sclerosing cholangitis, enthesitis, enthesopathy, dermatitis, dermatitis herpetiformis, progesterone dermatitis, atopic eczema, contact eczema, eczema, atherosclerosis, Still's disease, Addison's disease, Raynaud's phenomenon, erythrodermic psoriasis, noninfectious uveitis, peripheral uveitis, Dressler's syndrome, eosinophilic esophagitis, eosinophilic fasciitis, erythema nodosum, experimental allergic encephalomyelitis, Evans syndrome, fibrosing alveolitis, Vogt-Koyanagi-Harada syndrome, mucosal leishmaniasis, Kawasaki disease or syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, thrombocytopenic purpura, immune thrombocytopenic purpura (also known as immune thrombocytopenia, idiopathic immune thrombocytopenia, idiopathic thrombocytopenic thrombotic purpura, primary immune thrombocytopenia, idiopathic thrombocytopenic purpura (ITP), primary immune thrombocytopenic purpura, or autoimmune thrombocytopenic purpura), agammaglobulinemia, kidney inflammation, interstitial kidney inflammation, kidney disease, chronic kidney disease, renal failure, acute renal failure, end stage kidney disease, acute kidney injury, cisplatin induced acute renal failure, sepsis induced acute renal failure, anti-glomerular basement membrane (GBM) nephritis, anti-tubular basement membrane (TBM) nephritis, antiphospholipid syndrome (APS), nephritis, nephrotoxic nephritis, glomerulonephritis, acute glomerulonephritis, antineutrophil cytoplasmic autoantibody (ANCA) associated vasculitis, microscopic polyangiitis, granulomatosis with polyangiitis (GPA), Wegener's granulomatosis, amyotrophic lateral sclerosis, lupus nephritis, allergic eczema, transplant rejection, non-radiographic spondyloarthropathy, ophthalmic disorders, organ allograft rejection, fibroid lung, renal insufficiency, diabetic complications, diabetic nephropathy, diabetic retinopathy, diabetic retinitis, diabetic microangiopathy, insulinitis, tuberculosis, invasive

staphylococchia, invasive *Staphylococcus aureus* infection, inflammation after cataract surgery, allergic conjunctivitis, alopecia, alopecia areata, chronic urticaria, allergic asthma, neutrophilic asthma, periodontal diseases, periodontitis, gingivitis, gum disease, cardiomyopathy, diastolic cardiomyopathies, cardiac infarction, myocarditis, chronic heart failure, congenital heart block, coxsackie myocarditis, postmyocardial infarction syndrome, postpericardiotomy syndrome, endocarditis, subacute bacterial endocarditis (SBE), angiostenosis, restenosis, reperfusion disorders, autoimmune pancreatitis, acute pancreatitis, chronic pancreatitis, asthma, bronchial asthma, acute respiratory distress syndrome, adult respiratory distress syndrome, inflammatory bone disease, inflammatory pulmonary disease, ischemic attack, transient ischemic attack, systemic inflammatory response syndrome, glaucoma, orbital cellulitis, sudden orbital inflammation, postoperative inflammation, posttraumatic inflammation, allergic inflammation, intestinal inflammation, mucosal inflammation, prostate inflammation, prostatitis, chronic pelvic pain syndrome, testicular inflammation, chronic testicular inflammation, orchitis, orchitis mediated infertility, liver disorder, liver injury, hepatotoxicity, pneumonia, meningitis, cystitis, interstitial cystitis, pharyngolaryngitis, gastric mucosal injury, chronic pneumonia, pulmonary infarction, silicosis, sarcoidosis, pulmonary sarcoidosis, autoimmune angioedema, autoimmune dysautonomia, autoimmune hepatitis, autoimmune hyperlipidemia, autoimmune immunodeficiency, autoimmune inner ear disease (AIED), autoimmune myocarditis, autoimmune oophoritis, autoimmune aplastic anemia, autoimmune anemia, autoimmune hemolytic anemia, autoimmune retinopathy, autoimmune thrombocytopenic purpura (ATP), autoimmune thyroid disease, autoimmune urticaria, Goodpasture's syndrome, sinusitis, chronic hypertrophic rhinitis, chronic inflammatory demyelinating polyneuropathy, mixed connective tissue disease, undifferentiated connective tissue disease (UCTD), cognitive impairment, cognitive impairment in Alzheimer's disease, Parkinson's disease, spinal muscular atrophy, spinal cerebellar atrophy, progressive supranuclear palsy, Fisher syndrome, discoid lupus, central nervous system lupus, neuromyelitis optica (NMO; also known as Devic's disease or Devic's syndrome), encephalomyelitis, acute disseminated encephalomyelitis (ADEM), transverse myelitis, acute necrotizing hemorrhagic leukoencephalitis, multiple system atrophy, Huntington's disease, cerebrovascular dementia, diffuse Lewy body disease, amyloidosis, cerebrovascular disorder, cerebral infarction, transient ischemic attack, intracerebral hemorrhage, vascular disease of the spinal cord, spinal cord infarction, Lambert-Eaton syndrome, muscular dystrophy, metabolic myopathy, inflammatory myopathy, Chagas disease, chronic inflammatory demyelinating polyneuropathy (CIDP), chronic recurrent multifocal osteomyelitis (CRMO),

Churg-Strauss syndrome, Cogan's syndrome, cold agglutinin disease, essential mixed cryoglobulinemia, demyelinating neuropathies, inclusion body myositis, encephalitis, pemphigoid, bullous pemphigoid, pemphigus, pemphigus vulgaris, pemphigus foliaceus, cicatricial pemphigoid, ocular cicatricial pemphigoid, benign mucosal pemphigoid, Castleman disease (also known as giant or angiofollicular lymph node hyperplasia, lymphoid hamartoma, and angiofollicular lymph node hyperplasia), profundus lupus erythematosus, chronic thyroiditis, autoimmune gastritis, sepsis, burn injury, axonal and neuronal neuropathies, pain, neuropathy, peripheral neuropathy, chronic pain, optic neuritis, optic neuropathy, traumatic optic neuropathy, ischemic brain injury, deep venous thrombosis, neutropenia, autoimmune neutropenia, thrombocytopenia, abnormal immunoresponse, radiodermatitis, osteoporosis, parasitic infection, clonorchiasis, *Cryptosporidium* infection, *Streptococcus pneumoniae* carriage, chronic pneumococcal carriage, an immune disorder associated with or arising from activity of pathogenic lymphocytes, Henoch-Schonlein purpura, herpes gestationis, hypogammaglobulinemia, IgA nephropathy, IgG4-related sclerosing disease, immunoregulatory lipoproteins, Lambert-Eaton syndrome, lichen planus, lichen sclerosus, ligneous conjunctivitis, linear IgA disease (LAD), chronic Lyme disease, Meniere's disease, microscopic polyangiitis, mixed connective tissue disease (MCTD), Mooren's ulcer, Mucha-Habermann disease, narcolepsy, palindromic rheumatism, pediatric autoimmune neuropsychiatric disorders associated with streptococcus (PANDAS), paraneoplastic cerebellar degeneration, paroxysmal nocturnal hemoglobinuria (PNH), Parry-Romberg syndrome, Parsonnage-Turner syndrome, pars planitis, perivenous encephalomyelitis, pernicious anemia, POEMS syndrome, type I autoimmune polyglandular syndrome, type II autoimmune polyglandular syndrome, type III autoimmune polyglandular syndrome, pyoderma gangrenosum, pure red cell aplasia, reflex sympathetic dystrophy, relapsing polychondritis, restless legs syndrome, Schmidt syndrome, sperm autoimmunity, stiff person syndrome, Susac's syndrome, sympathetic ophthalmia, Tolosa-Hunt syndrome, vesiculobullous dermatosis, and vitiligo.

[0098] Also included are diseases or disorders which are implicated by the regulation of the circadian rhythm of individuals and include, *e.g.*, major depression, seasonal affective disorder, post-traumatic stress disorder (PTSD), bipolar disorder, autism, epilepsy, Alzheimer's disease and other central nervous system (CNS) disorders associated with altered sleep and/or circadian rhythms.

[0099] Further included are diseases and disorders mediated by IL-17 expression, including STAT3-mediated IL-17 expression, in neutrophils, and include, *e.g.*, corneal fungal infection;

risk of corneal fungal infection; corneal ulcer; corneal ulcer resulting from fungal keratitis; corneal ulcer resulting from fungal infection; corneal fungal infection and related inflammation; keratitis; fungal keratitis; corneal inflammation; corneal disease; ocular disease; fungal-mediated corneal infection, corneal ulcer, corneal inflammation, or ocular ulcer, inflammation or infection; bacterial-mediated corneal infection, corneal ulcer, corneal inflammation, or ocular ulcer, inflammation or infection; microbial disease; bacterial infections; fungal infections; *Aspergillus* keratitis; *Fusarium* keratitis; cutaneous T cell lymphoma; lung inflammation; acute kidney ischemia-reperfusion injury; anthrax, including, cutaneous anthrax, inhalation anthrax, gastrointestinal anthrax and injection anthrax; aspergillosis, including, pulmonary aspergillosis, chronic pulmonary aspergillosis (CPA), chronic aspergillosis, chronic cavitary pulmonary aspergillosis (CCPA), allergic bronchopulmonary aspergillosis (ABPA), allergic *Aspergillus* sinusitis, aspergilloma, invasive aspergillosis, chronic necrotizing aspergillosis and cutaneous (skin) aspergillosis; and histoplasmosis, including, systemic histoplasmosis. In particular embodiments, the fungus or fungal infection mediating the disease or disorder described above includes one or more of *Aspergillus*, *Fusarium*, *Alternaria*, *Candida*, *Curvularia* or *Histoplasma*.

[00100] Compounds described herein can also be used to treat or reduce the risk of abnormal cortical development or psychiatric disorder, *e.g.*, autism spectrum disorder (ASD), schizophrenia, and/or depression, in a fetus. Compounds described herein can also be used to treat a pregnant female having a hyper-inflammatory condition associated with an infection, such as a viral or bacterial infection, or associated with exposure to an inflammatory or environmental toxin during pregnancy. In particular embodiments, a fetus is treated in utero in a pregnant female with a compound disclosed herein to decrease the risk of the fetus developing a psychiatric disorder, to reduce inflammation in the pregnant female, to reduce the risk of abnormal cortical development in the fetus, and/or to decrease symptoms of a psychiatric disorder in offspring of a pregnant female.

[00101] In one embodiment, a human patient is treated with a compound of Formula I and a pharmaceutically acceptable carrier, adjuvant, or vehicle, wherein said compound is present in an amount to treat or ameliorate one or more of the diseases and conditions recited above. In an alternative embodiment, the diseases and conditions treated or ameliorated by a compound of Formula I include, *e.g.*, asthma, atopic dermatitis, acne, Crohn's disease, regional enteritis, ulcerative colitis, Sjögren's syndrome, uveitis, Behçet's disease, dermatomyositis, multiple sclerosis, ankylosing spondylitis, systemic lupus erythematosus (SLE), scleroderma, psoriasis, psoriatic arthritis (PsA), steroid resistant asthma and rheumatoid arthritis in the patient.

[00102] The present disclosure further relates to a combination therapy for treating or ameliorating a disease or a disorder described herein. In some embodiments, the combination therapy comprises administering at least one compound represented by Structural Formula I in combination with one or more agents for treating or ameliorating inflammatory, metabolic and autoimmune diseases or disorders mediated by ROR γ . In some embodiments, the combination therapy comprises administering at least one compound represented by Structural Formula I in combination with one or more agents for the treatment of diseases including asthma, chronic obstructive pulmonary disease (COPD), bronchitis, allergic rhinitis, atopic dermatitis, contact dermatitis, acne, cystic fibrosis, allograft rejection, multiple sclerosis, scleroderma, arthritis, rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, systemic lupus erythematosus (SLE), psoriasis, Hashimoto's disease, pancreatitis, autoimmune diabetes, type I diabetes, autoimmune ocular disease, ulcerative colitis, Crohn's disease, regional enteritis, inflammatory bowel disease (IBD), inflammatory bowel syndrome (IBS), Sjögren's syndrome, optic neuritis, obesity, hepatosteatorrhea, adipose tissue-associated inflammation, insulin resistance, type II diabetes, neuromyelitis optica, myasthenia gravis, age related macular degeneration, dry eye, uveitis, Guillain-Barré syndrome, psoriasis, psoriatic arthritis (PsA), steroid resistant asthma, Graves' disease, scleritis, major depression, seasonal affective disorder, PTSD, bipolar disorder, autism, epilepsy, Alzheimer's, CNS disorders associated with altered sleep and/or circadian rhythms, endometriosis, obstructive sleep apnea syndrome (OSAS), Behçet's disease, dermatomyositis, polymyositis, graft versus host disease, primary biliary cirrhosis, liver fibrosis, non-alcoholic fatty liver disease (NAFLD), sarcoidosis, primary sclerosing cholangitis, autoimmune thyroid disease, autoimmune polyendocrine syndrome type I, autoimmune polyendocrine syndrome type II, celiac disease, neuromyelitis, juvenile idiopathic arthritis, systemic sclerosis, myocardial infarction, pulmonary hypertension, osteoarthritis, cutaneous leishmaniasis, sinonasal polyposis, and cancer, including but not limited to, lung cancer, gastric cancer, breast cancer and colon cancer.

[00103] The compounds herein may also be used in combination with immunotherapies for the treatment of a disease or disorder disclosed herein.

[00104] Combination therapy includes, *e.g.*, co-administration of a compound described herein and one or more other agents, sequential administration of a compound described herein and one or more other agents, administration of a composition containing a compound described herein and one or more other agents, or simultaneous administration of separate compositions containing a compound described herein and one or more other agents.

[00105] The present disclosure further provides a method of treating a subject, such as a human, suffering from one of the abovementioned disorders or diseases.

[00106] In one aspect, ROR γ modulators disclosed in WO 2014/179564, WO 2015/116904, WO 2016/061160, PCT/US2016/045318, and U.S. patent application Nos. 14/933,468 and 14/933,524 can also be used in the methods disclosed herein to treat or ameliorate, in a subject, one or more of the diseases and/or disorders and/or conditions recited herein. In one embodiment, a subject is treated with one or more ROR γ modulators disclosed in WO 2014/179564, WO 2015/116904, WO 2016/061160, PCT/US2016/045318, and U.S. patent application Nos. U.S. 14/933,468 or U.S. 14/933,524 and a pharmaceutically acceptable carrier, adjuvant, or vehicle, wherein said ROR γ modulator is present in an amount to treat or ameliorate a disease or disorder selected from corneal fungal infection; risk of corneal fungal infection; corneal ulcer; corneal ulcer resulting from fungal keratitis; corneal ulcer resulting from fungal infection; corneal fungal infection and related inflammation; keratitis; fungal keratitis; corneal inflammation; corneal disease; ocular disease; fungal-mediated corneal infection, corneal ulcer, corneal inflammation, or ocular ulcer, inflammation or infection; bacterial-mediated corneal infection, corneal ulcer, corneal inflammation, or ocular ulcer, inflammation or infection; microbial disease; bacterial infections; fungal infections; *Aspergillus* keratitis; *Fusarium* keratitis; cutaneous T cell lymphoma; lung inflammation; acute kidney ischemia-reperfusion injury; anthrax, including, cutaneous anthrax, inhalation anthrax, gastrointestinal anthrax and injection anthrax; aspergillosis, including, pulmonary aspergillosis, chronic pulmonary aspergillosis (CPA), chronic aspergillosis, chronic cavitary pulmonary aspergillosis (CCPA), allergic bronchopulmonary aspergillosis (ABPA), allergic *Aspergillus* sinusitis, aspergilloma, invasive aspergillosis, chronic necrotizing aspergillosis and cutaneous (skin) aspergillosis; histoplasmosis, including, systemic histoplasmosis; and prostate cancer. In some embodiments, the one or more ROR γ modulator disclosed in WO 2014/179564, WO 2015/116904, WO 2016/061160, PCT/US2016/045318, and U.S. patent application Nos. 14/933,468 and 14/933,524 is administered in combination with one or more additional agent for treating the disease or disorder.

[00107] The present disclosure further relates to the use of provided compounds for the production of pharmaceutical compositions which are employed for the treatment and/or prophylaxis and/or amelioration of the diseases and disorders mentioned herein.

[00108] Compounds or compositions described herein may be administered using any amount and any route of administration effective for treating or lessening the severity of one or more of

the diseases and conditions described herein. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular agent, its mode of administration, and the like. Provided compounds are preferably formulated in unit dosage form for ease of administration and uniformity of dosage. The expression "unit dosage form" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present disclosure will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts.

[00109] Pharmaceutically acceptable compositions of this disclosure can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, provided compounds may be administered orally or parenterally at dosage levels of about 0.01 mg/kg to about 50 mg/kg and preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

[00110] The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof, biopsied material obtained from a mammal or extracts thereof, and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

[00111] The amount of both, a provided compound and additional therapeutic agent (in those compositions which comprise an additional therapeutic agent as described above) that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

[00112] In those compositions which comprise an additional therapeutic agent, that additional therapeutic agent and the provided compound may act synergistically. Therefore, the amount of additional therapeutic agent in such compositions will be less than that required in a monotherapy utilizing only that therapeutic agent.

[00113] The amount of additional therapeutic agent present in the compositions of this disclosure will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent.

EXEMPLIFICATION

[00114] As depicted in the Examples below, in certain exemplary embodiments, compounds are prepared according to the following general procedures. It will be appreciated that, although the general methods depict the synthesis of certain compounds herein, the following general methods, and other methods known to one of ordinary skill in the art, can be applied to all compounds and subclasses and species of each of these compounds, as described herein.

GENERAL DESCRIPTION OF SYNTHESIS

[00115] The compounds described herein can be readily prepared according to the following reaction schemes and examples, or modifications thereof, using readily available starting materials, reagents and conventional synthesis procedures. Many of the reactions can also be carried out under microwave (MW) conditions or using conventional heating or utilizing other technologies such as solid phase reagents/scavengers or flow chemistry. In these reactions, it is also possible to make use of variants which are themselves known to those of ordinary skill in the art, but are not mentioned in greater detail. Furthermore, other methods for preparing compounds described herein will be readily apparent to a person of ordinary skill in the art in light of the following reaction schemes and examples. In cases where synthetic intermediates and final products contain potentially reactive functional groups, for example amino, hydroxy, thiol and carboxylic acid groups, that may interfere with the desired reaction, it may be advantageous to employ protected forms of the intermediate. Methods for the selection, introduction and subsequent removal of protecting groups are well known to those skilled in the art. In the discussion below variables have the meanings indicated above unless otherwise indicated. The abbreviations used in these experimental details are listed below and additional ones should be known to a person skilled in the art of synthesis. In addition, one can refer to the following references for suitable methods of synthesis as described in March, Advanced Organic Chemistry, 3rd edition, John Wiley & Sons, 1985, Greene and Wuts, Protective Groups in Organic Synthesis, 2nd edition, John Wiley & Sons, 1991, and Richard Larock, Comprehensive Organic Transformations, 4th edition, VCH publishers Inc., 1989.

[00116] Generally, reagents in the reaction schemes are used in equimolar amounts; however, in certain cases it may be desirable to use an excess of one reagent to drive a reaction to

completion. This is especially the case when the excess reagent can be readily removed by evaporation or extraction. Bases employed to neutralize HCl in reaction mixtures are generally used in slight to substantial excess (1.05 – 5 equivalents).

[00117] Where NMR data are presented, spectra were obtained on a Varian 400 (400 MHz) or 300 (300 MHz) and are reported as ppm downfield from tetramethylsilane with number of proton, multiplicities and coupling constants indicated parenthetically along with reference to deuterated solvent.

[00118] The invention is illustrated by way of the following examples, in which the following abbreviations may be employed.

Abbreviation	Meaning
ACN, MeCN, CH ₃ CN	acetonitrile
AIBN	azobisisobutyronitrile
aq	aqueous
Boc	<i>tert</i> -butoxycarbonyl or <i>t</i> -butoxycarbonyl
brine	saturated aqueous NaCl
c-Bu	cyclobutyl
Cbz	benzyloxy carbonyl
CeCl ₃	ceric chloride
Cs ₂ CO ₃	cesium carbonate
CuI	cuprous iodide
c-Pr	cyclopropyl
DCM or CH ₂ Cl ₂	methylene chloride
dcpp.2HBF ₄	1,3-Bis(dicyclohexylphosphino)propane bis(tetrafluoroborate)
DIEA	diisopropyl ethyl amine
DMF	dimethyl formamide
DMS/Me ₂ S	dimethyl sulfide
DMSO	dimethyl sulfoxide
EDCI	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
EtI	ethyl iodide
Et	ethyl
Et ₂ O	ethyl ether
Et ₃ SiH	triethylsilane

Et ₃ N	triethylamine
EtOAc , EA, AcOEt	ethyl acetate
EtOH	ethanol
FeCl ₃	ferric chloride
h, hr	hour(s)
HATU	<i>O</i> -(7-azabenzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium-hexafluorophosphate
HBTU	<i>O</i> -benzotriazole-1-yl- <i>N,N,N',N'</i> -tetramethyluronium-hexafluorophosphate
HCl	hydrochloric acid
H ₂ O	water
H ₂ O ₂	hydrogen peroxide
HPLC	high performance liquid chromatography
i-BuOCOC1	iso-butoxycarbonyl chloride
ICl	iodochloride
K ₂ CO ₃	potassium carbonate
K ₃ PO ₄	tripotassium phosphate
LC-MS	liquid chromatography–mass spectrometry
LDA	lithium diisopropylamide
LiCl	lithium chloride
LiOH	lithium hydroxide
MCPBA, <i>m</i> -CPBA	meta-chloroperoxybenzoic acid
MeOH	methanol
MeI	methyl iodide
Me	methyl
mg	milligram
MgSO ₄	magnesium sulfate (anhydrous)
min	minute(s)
mL	milliliters
mmol	millimoles
mp, m.p.	melting point
MS	mass spectrometry
MW, uwave	microwave
NaBH ₄	sodium borohydride
NaBH ₃ CN	sodium cyanoborohydride
NaH	sodium hydride
NaHCO ₃	sodium bicarbonate

NaOH	sodium hydroxide
NaOMe	sodium methoxide
Na ₂ S ₂ O ₃	sodium thiosulfate
Na ₂ S ₂ O ₅	sodium dithionate
Na ₂ SO ₄	sodium sulfate
NH ₄ OH	ammonium hydroxide
(NH ₄) ₂ CO ₃	ammonium carbonate
NH ₄ Cl	ammonium chloride
Na ₂ CO ₃	sodium carbonate
NaHCO ₃	sodium bicarbonate
NaH	sodium hydride
NBS	<i>N</i> -bromosuccinimide
<i>n</i> -BuLi	<i>n</i> -butyllithium
NMM	<i>N</i> -methyl-morpholine
NMP	<i>N</i> -methyl-pyrrolidin-2-one
OTf	trifluoromethanesulfonate
OTs	tosylate
PdCl ₂ dppf	[1,1-bis(diphenylphosphino)ferrocene] dichloropalladium(ii)
Pd ₂ (dba) ₃	tris(dibenzylideneacetone)dipalladium(0)
PE	petroleum ether
rt	room temperature
sat.	saturated
SEM	2-trimethylsilylethyoxymethyl
SFC	supercritical fluid chromatography
<i>t</i> -BuOK	potassium tert butoxide
<i>t</i> -BuLi	tert butyl lithium
<i>t</i> -BuOOH	tert butyl peroxide
TBAF	tetrabutylammonium fluoride
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TLC	thin layer chromatography
Ti(OEt) ₄	titanium tetra ethoxide
tR	retention time
Zn	zinc
Zn(CN) ₂	zinc cyanide

PREPARATION OF COMPOUNDS OF FORMULA I

[00119] Compounds of Formula I were prepared according to the general procedures outlined below.

Compounds Names^a

Cpd No	Name
Hy1A-1	(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-((tetrahydro-2H-pyran-4-yl)methoxy)isoquinoline-6-carboxamide
Hy1A-2	N-(4-(ethylsulfonyl)benzyl)-1-(4-(trifluoromethyl)phenoxy)isoquinoline-6-carboxamide
Hy1A-3.1 ^a	N-(4-(ethylsulfonyl)benzyl)-1-(((1s,4s)-4-(trifluoromethyl)cyclohexyl)oxy)isoquinoline-6-carboxamide
Hy1A-3.2 ^a	N-(4-(ethylsulfonyl)benzyl)-1-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)oxy)isoquinoline-6-carboxamide
Hy1A-4	N-(4-(ethylsulfonyl)benzyl)-1-((4-(trifluoromethyl)benzyl)oxy)isoquinoline-6-carboxamide
Hy1A-5	N-(4-(ethylsulfonyl)benzyl)-1-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide
Hy1A-6	N-(4-(ethylsulfonyl)benzyl)-3-methyl-1-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide
Hy1A-7	(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-((4-(trifluoromethyl)benzyl)oxy)isoquinoline-6-carboxamide
Hy1A-8	N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide
Hy1A-9	3-chloro-N-(4-(ethylsulfonyl)benzyl)-1-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide
Hy1A-10	N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide
Hy1A-11	3-chloro-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide
Hy1A-12	(R)-1-((3,3-difluorocyclobutyl)methoxy)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methylisoquinoline-6-carboxamide
Hy1A-13	(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-((4-fluorobenzyl)oxy)-3-methylisoquinoline-6-carboxamide

Hy1A-14	(R)-1-((4,4-difluorocyclohexyl)methoxy)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methylisoquinoline-6-carboxamide
Hy1A-15	(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-(4-(trifluoromethoxy)phenoxy)isoquinoline-6-carboxamide
Hy1A-16	N-((S)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide
Hy1A-17	(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)oxy)isoquinoline-6-carboxamide
Hy1A-18 ^b	N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-(((3S,6R)-6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methoxy)isoquinoline-6-carboxamide
Hy1A-18.2 ^b	N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-(((3R,6S)-6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methoxy)isoquinoline-6-carboxamide
Hy1A-18.3 ^b	Mixture of N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-(((3S,6S)-6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methoxy)isoquinoline-6-carboxamide and N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-(((3R,6R)-6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methoxy)isoquinoline-6-carboxamide
Hy1A-19	(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)methoxy)isoquinoline-6-carboxamide
Hy1A-20	3-ethyl-N-(4-(ethylsulfonyl)benzyl)-1-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide
Hy1A-21	3-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide
Hy1A-22.1 ^c	N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-3-methyl-1-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide
Hy1A-22.2 ^c	N-((S)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-3-methyl-1-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide
Hy1A-23	N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methoxy-1-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide

Hy1A-24	(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-((4-(trifluoromethyl)phenoxy)methyl)isoquinoline-6-carboxamide
Hy1B-1	N-(4-(ethylsulfonyl)benzyl)-2-methyl-4-((4-(trifluoromethyl)benzyl)oxy)quinoline-7-carboxamide
Hy1B-2	N-(4-(ethylsulfonyl)benzyl)-2-methyl-4-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methoxy)quinoline-7-carboxamide
Hy1B-3	(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-methyl-4-((4-(trifluoromethyl)benzyl)oxy)quinoline-7-carboxamide
Hy1B-4	N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-methyl-4-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methoxy)quinoline-7-carboxamide
Hy1C-1	N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-4-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methoxy)quinazoline-7-carboxamide
Hy1C-2	N-(4-(ethylsulfonyl)benzyl)-4-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methoxy)quinazoline-7-carboxamide
Hy2A-1.1 ^d	(R)-2-(5-cyanopyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-1.2 ^d	(S)-2-(5-cyanopyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-2	N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-3	N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-2-(4-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-4.1 ^e	ethyl (S)-2-(6-((4-(ethylsulfonyl)benzyl)carbamoyl)-1-isopropyl-3,4-dihydroisoquinolin-2(1H)-yl)pyrimidine-5-carboxylate
Hy2A-4.2 ^e	ethyl (R)-2-(6-((4-(ethylsulfonyl)benzyl)carbamoyl)-1-isopropyl-3,4-dihydroisoquinolin-2(1H)-yl)pyrimidine-5-carboxylate
Hy2A-5	ethyl 2-(6-((4-(ethylsulfonyl)benzyl)carbamoyl)-1-isopropyl-3,4-dihydroisoquinolin-2(1H)-yl)-4-(trifluoromethyl)pyrimidine-5-carboxylate
Hy2A-5.1 ^f	ethyl (S)-2-(6-((4-(ethylsulfonyl)benzyl)carbamoyl)-1-isopropyl-3,4-dihydroisoquinolin-2(1H)-yl)-4-(trifluoromethyl)pyrimidine-5-carboxylate
Hy2A-5.2 ^f	ethyl (R)-2-(6-((4-(ethylsulfonyl)benzyl)carbamoyl)-1-isopropyl-3,4-dihydroisoquinolin-2(1H)-yl)-4-(trifluoromethyl)pyrimidine-5-carboxylate

Hy2A-6	2-(4-fluorobenzyl)-1-methyl-N-(4-(methylsulfonyl)benzyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-7	2-(4-cyanobenzyl)-1-methyl-N-(4-(methylsulfonyl)benzyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-8	N-(4-(ethylsulfonyl)benzyl)-2-(4-fluorobenzyl)-1-methyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-9	2-(4-chlorobenzyl)-1-methyl-N-(4-(methylsulfonyl)benzyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-10	2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-methyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-11	2-(4-chlorobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-methyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-12.1 ^g	(S)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-2-((5-methylpyrimidin-2-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-12.2 ^g	(R)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-2-((5-methylpyrimidin-2-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-13	N-(4-(ethylsulfonyl)benzyl)-2-(4-fluorobenzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-14	2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-15.1 ^h	(S)-2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-15.2 ^h	(R)-2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-16.1 ⁱ	(S)-2-((5-cyanopyridin-2-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-16.2 ⁱ	(R)-2-((5-cyanopyridin-2-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-17.1 ^j	(S)-2-((6-cyanopyridin-3-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-17.2 ^j	(R)-2-((6-cyanopyridin-3-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-18	N-(1-(4-(ethylsulfonyl)phenyl)ethyl)-1-isopropyl-2-((5-methylpyrazin-2-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide

Hy2A-19	2-(4-chlorobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-20.1 ^k	(S)-2-(4-chlorobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-20.2 ^k	(R)-2-(4-chlorobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-21	2-((5-chloropyridin-2-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-22.1 ^l	(S)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-4,4-dimethyl-2-((5-methylpyrazin-2-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-22.2 ^l	(R)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-4,4-dimethyl-2-((5-methylpyrazin-2-yl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-23.1 ^m	(S)-2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-23.2 ^m	(R)-2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-24	methyl (S)-4-(((6-((4-(ethylsulfonyl)benzyl)carbamoyl)-1-isopropyl-3,4-dihydroisoquinolin-2(1H)-yl)methyl)piperidine-1-carboxylate
Hy2A-25.1 ⁿ	(S)-2-((6-(difluoromethoxy)pyridin-3-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-25.2 ⁿ	(R)-2-((6-(difluoromethoxy)pyridin-3-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-26.1 ^o	(S)-2-((5-bromopyrimidin-2-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-26.2 ^o	(R)-2-((5-bromopyrimidin-2-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-27	(S)-1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-28.1 ^p	(S)-1-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide

Hy2A-28.2 ^p	(S)-1-ethyl-N-((S)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-28.3 ^p	(R)-1-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-29.1 ^q	(S)-1-ethyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-29.2 ^q	(S)-1-ethyl-N-((S)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2A-29.3 ^q	(R)-1-ethyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-1,2,3,4-tetrahydroisoquinoline-6-carboxamide
Hy2B-1.1 ^r	(S)-7-(5-cyanopyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-1.2 ^r	(R)-7-(5-cyanopyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-2.1 ^s	(S)-7-(5-chloropyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-2.2 ^s	(R)-7-(5-chloropyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-3.1 ^t	(S)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-7-(5-(trifluoromethyl)pyrimidin-2-yl)-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-3.2 ^t	(R)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-7-(5-(trifluoromethyl)pyrimidin-2-yl)-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-4.1 ^u	ethyl (R)-2-(3-((4-(ethylsulfonyl)benzyl)carbamoyl)-8-isopropyl-5,8-dihydro-1,7-naphthyridin-7(6H)-yl)pyrimidine-5-carboxylate
Hy2B-4.2 ^u	ethyl (S)-2-(3-((4-(ethylsulfonyl)benzyl)carbamoyl)-8-isopropyl-5,8-dihydro-1,7-naphthyridin-7(6H)-yl)pyrimidine-5-carboxylate
Hy2B-5.1 ^v	ethyl (R)-2-(3-((4-(ethylsulfonyl)benzyl)carbamoyl)-8-isopropyl-5,8-dihydro-1,7-naphthyridin-7(6H)-yl)-4-(trifluoromethyl)pyrimidine-5-carboxylate

Hy2B-5.2 ^v	ethyl (S)-2-(3-((4-(ethylsulfonyl)benzyl)carbamoyl)-8-isopropyl-5,8-dihydro-1,7-naphthyridin-7(6H)-yl)-4-(trifluoromethyl)pyrimidine-5-carboxylate
Hy2B-6.1 ^w	rel-(R)-N-(4-(ethylsulfonyl)benzyl)-7-(4-fluorobenzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-6.2 ^w	rel-(R)-N-(4-(ethylsulfonyl)benzyl)-7-(4-fluorobenzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-7.1 ^x	rel-(R)-7-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-7.2 ^x	rel-(R)-7-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-8.1 ^y	rel-(R)-7-(4-chlorobenzyl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-8.2 ^y	rel-(R)-7-(4-chlorobenzyl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-9.1 ^z	(S)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-7-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-9.2 ^z	(R)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-7-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-10.1 ^{aa}	(S)-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-8-isopropyl-7-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy2B-10.2 ^{aa}	(R)-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-8-isopropyl-7-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide
Hy3A-1	3-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole-6-carboxamide
Hy3A-2	3-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole-6-carboxamide
Hy3A-3	3-cyclopropyl-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole-6-carboxamide

Hy3A-4	(R)-3-cyclopropyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(4-(trifluoromethyl)benzyl)-2H-indazole-6-carboxamide
Hy3A-5	3-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole-6-carboxamide
Hy3A-6	3-cyclopropyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole-6-carboxamide
Hy3A-7	(R)-3-cyclopropyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(4-(trifluoromethyl)benzyl)-2H-indazole-6-carboxamide
Hy3B-1	3-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-2H-pyrazolo[4,3-b]pyridine-6-carboxamide
Hy3B-2	3-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-2H-pyrazolo[4,3-b]pyridine-6-carboxamide
Hy3B-3	3-cyclopropyl-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-2H-pyrazolo[4,3-b]pyridine-6-carboxamide
Hy3B-4	3-cyclopropyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-2H-pyrazolo[4,3-b]pyridine-6-carboxamide
Hy4-1	N-(4-(ethylsulfonyl)benzyl)-1-methyl-7-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methoxy)-1H-pyrrolo[2,3-c]pyridine-2-carboxamide
Hy4-2	N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-methyl-7-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methoxy)-1H-pyrrolo[2,3-c]pyridine-2-carboxamide
Hy4-3	N-(4-(ethylsulfonyl)benzyl)-1-methyl-7-((4-(trifluoromethyl)benzyl)oxy)-1H-pyrrolo[2,3-c]pyridine-2-carboxamide
Hy4-4	(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-methyl-7-((4-(trifluoromethyl)benzyl)oxy)-1H-pyrrolo[2,3-c]pyridine-2-carboxamide
Hy4-5	N-(4-(ethylsulfonyl)benzyl)-7-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methoxy)-1H-pyrrolo[2,3-c]pyridine-2-carboxamide
Hy5-1	N-(4-(ethylsulfonyl)benzyl)-1-methyl-7-(4-(trifluoromethyl)phenoxy)-1H-pyrrolo[2,3-c]pyridine-3-carboxamide

Hy5-2	N-(4-(ethylsulfonyl)benzyl)-7-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methoxy)-1H-pyrrolo[2,3-c]pyridine-3-carboxamide
Hy5-3	N-(4-(ethylsulfonyl)benzyl)-1-methyl-7-((4-(trifluoromethyl)benzyl)oxy)-1H-pyrrolo[2,3-c]pyridine-3-carboxamide
Hy5-4	N-(4-(ethylsulfonyl)benzyl)-1-methyl-7-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methoxy)-1H-pyrrolo[2,3-c]pyridine-3-carboxamide
Hy5-5	N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-methyl-7-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methoxy)-1H-pyrrolo[2,3-c]pyridine-3-carboxamide
Hy6-1	N-(4-(ethylsulfonyl)benzyl)-2,5-dimethyl-4-((4-(trifluoromethyl)benzyl)oxy)thieno[2,3-d]pyrimidine-6-carboxamide
Hy6-2	N-(4-(ethylsulfonyl)benzyl)-2,5-dimethyl-4-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methoxy)thieno[2,3-d]pyrimidine-6-carboxamide
Hy6-3	(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2,5-dimethyl-4-((4-(trifluoromethyl)benzyl)oxy)thieno[2,3-d]pyrimidine-6-carboxamide
Hy6-4	N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2,5-dimethyl-4-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methoxy)thieno[2,3-d]pyrimidine-6-carboxamide
Hy7-1.1 ^{bb}	(S)-N-(4-(ethylsulfonyl)benzyl)-2-(5-fluoropyrimidin-2-yl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-1.2 ^{bb}	(R)-N-(4-(ethylsulfonyl)benzyl)-2-(5-fluoropyrimidin-2-yl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-2.1 ^{cc}	(S)-2-(5-cyanopyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-2.2 ^{cc}	(R)-2-(5-cyanopyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-3.1 ^{dd}	(S)-2-(5-chloropyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-3.2 ^{dd}	(R)-2-(5-chloropyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-4.1 ^{ee}	(S)-2-(5-cyclopropylpyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide

Hy7-4.2 ^{ee}	(R)-2-(5-cyclopropylpyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-5.1 ^{ff}	(S)-2-(5-ethoxypyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-5.2 ^{ff}	(R)-2-(5-ethoxypyrimidin-2-yl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-6.1 ^{gg}	(S)-2-(5-chloropyrimidin-2-yl)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-6.2 ^{gg}	(R)-2-(5-chloropyrimidin-2-yl)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-7.1 ^{hh}	(S)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-7.2 ^{hh}	(R)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-8.1 ⁱⁱ	(S)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(5-(trifluoromethyl)pyridin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-8.2 ⁱⁱ	(R)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(5-(trifluoromethyl)pyridin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-9.1 ^{jj}	(S)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-9.2 ^{jj}	(R)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-10.1 ^{kk}	(R)-4-ethyl-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-10.2 ^{kk}	(S)-4-ethyl-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-11.1 ^{ll}	(S)-2-(5-cyclopropylpyrimidin-2-yl)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-11.2 ^{ll}	(R)-2-(5-cyclopropylpyrimidin-2-yl)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide

Hy7- 12.1 ^{mm}	(S)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-1-oxo-2-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 12.2 ^{mm}	(R)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-1-oxo-2-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 13.1 ^{nm}	(S)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(3-methyl-5-(trifluoromethyl)pyridin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 13.2 ^{nm}	(R)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(3-methyl-5-(trifluoromethyl)pyridin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 14.1 ^{oo}	(S)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(4-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 14.2 ^{oo}	(R)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(4-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 15.1 ^{pp}	(S)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 15.2 ^{pp}	(R)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 16.1 ^{qq}	ethyl (S)-2-(7-((4-(ethylsulfonyl)benzyl)carbamoyl)-4-isopropyl-3,4-dihydroisoquinolin-2(1H)-yl)pyrimidine-5-carboxylate
Hy7- 16.2 ^{qq}	ethyl (R)-2-(7-((4-(ethylsulfonyl)benzyl)carbamoyl)-4-isopropyl-3,4-dihydroisoquinolin-2(1H)-yl)pyrimidine-5-carboxylate
Hy7- 17.1 ^{rr}	(S)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 17.2 ^{rr}	(R)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 18.1 ^{ss}	(S)-4-ethyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide

Hy7- 18.2 ^{ss}	(S)-4-ethyl-N-((S)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 18.3 ^{ss}	(R)-4-ethyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 18.4 ^{ss}	(R)-4-ethyl-N-((S)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-19.1 ^{tt}	(S)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-morpholinopyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7-19.2 ^{tt}	(R)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-morpholinopyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 20.1 ^{uu}	(S)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(5-(6-methoxypyridin-2-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 20.2 ^{uu}	(R)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(5-(6-methoxypyridin-2-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 21.1 ^{vv}	(S)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-(6-oxo-1,6-dihydropyridin-3-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 21.2 ^{vv}	(R)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-(6-oxo-1,6-dihydropyridin-3-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 22.1 ^{ww}	(S)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(3-methyl-5-(trifluoromethyl)pyridin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 22.2 ^{ww}	(R)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(3-methyl-5-(trifluoromethyl)pyridin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 23.1 ^{xx}	(S)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-(2-methoxypyridin-4-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 23.2 ^{xx}	(R)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-(2-methoxypyridin-4-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 24.1 ^{yy}	(S)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide

Hy7- 24.2 ^{yy}	(R)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 25.1 ^{zz}	(S)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-(1-methyl-2-oxo-1,2-dihydropyridin-4-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 25.2 ^{zz}	(R)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-2-(5-(1-methyl-2-oxo-1,2-dihydropyridin-4-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 26.1 ^{aaa}	ethyl (S)-2-(7-((4-(ethylsulfonyl)benzyl)carbamoyl)-4-isopropyl-3,4-dihydroisoquinolin-2(1H)-yl)-4-(trifluoromethyl)pyrimidine-5-carboxylate
Hy7- 26.2 ^{aaa}	ethyl (R)-2-(7-((4-(ethylsulfonyl)benzyl)carbamoyl)-4-isopropyl-3,4-dihydroisoquinolin-2(1H)-yl)-4-(trifluoromethyl)pyrimidine-5-carboxylate
Hy7- 27.1 ^{bbb}	(S)-N-(4-(ethylsulfonyl)benzyl)-2-(4-fluorobenzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 27.2 ^{bbb}	(R)-N-(4-(ethylsulfonyl)benzyl)-2-(4-fluorobenzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 28.1 ^{ccc}	(S)-2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 28.2 ^{ccc}	(R)-2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 29.1 ^{ddd}	(S)-2-(4-chlorobenzyl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 29.2 ^{ddd}	(R)-2-(4-chlorobenzyl)-N-(4-(ethylsulfonyl)benzyl)-4-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 30.1 ^{eee}	(S)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(tetrahydro-2H-pyran-4-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 30.2 ^{eee}	(R)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(tetrahydro-2H-pyran-4-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 31.1 ^{fff}	(S)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(5-(oxazol-2-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 31.2 ^{fff}	(R)-4-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(5-(oxazol-2-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 32.1 ^{ggg}	(S)-2-(4,4-difluorocyclohexyl)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide

Hy7- 32.2 ^{ggg}	(R)-2-(4,4-difluorocyclohexyl)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 33.1 ^{hhh}	(S)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(5-(oxazol-2-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide and
Hy7- 33.2 ^{hhh}	(R)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(5-(oxazol-2-yl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 34.1 ⁱⁱⁱ	ethyl 2-((S)-4-ethyl-7-(((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)carbamoyl)-3,4-dihydroisoquinolin-2(1H)-yl)pyrimidine-5-carboxylate
Hy7- 34.2 ⁱⁱⁱ	ethyl 2-((R)-4-ethyl-7-(((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)carbamoyl)-3,4-dihydroisoquinolin-2(1H)-yl)pyrimidine-5-carboxylate
Hy7- 35.1 ^{jjj}	(S)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((1r,4S)-4-(trifluoromethyl)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 35.2 ^{jjj}	(S)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((1s,4R)-4-(trifluoromethyl)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 35.3 ^{jjj}	(R)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((1r,4R)-4-(trifluoromethyl)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 35.4 ^{jjj}	(R)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((1s,4S)-4-(trifluoromethyl)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 36.1 ^{kkk}	(S)-N-(4-(ethylsulfonyl)benzyl)-4-(trifluoromethyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 36.2 ^{kkk}	(R)-N-(4-(ethylsulfonyl)benzyl)-4-(trifluoromethyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 37.1 ^{lll}	(S)-2-(3-cyano-5-(trifluoromethyl)pyridin-2-yl)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide
Hy7- 37.2 ^{lll}	(R)-2-(3-cyano-5-(trifluoromethyl)pyridin-2-yl)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide

Hy7B-1 N-(4-(ethylsulfonyl)benzyl)-8,8-dimethyl-6-(5-(trifluoromethyl)pyrimidin-2-yl)-5,6,7,8-tetrahydro-1,6-naphthyridine-3-carboxamide

a, b, c, d, e, f, g, h, i, j, k, l, m, n, o, p, q, r, s, t, u, v, w, x, y, z, aa, bb, cc, dd, ee, ff, gg, hh, ii, jj, kk, ll, mm, nn, oo, pp, qq, rr, ss, tt, uu, vv, ww, xx, yy, z, aaa, bbb, ccc, ddd, eee, fff, ggg, hhh, iii, jjj, kkk, lll = isomer mixtures were separated by chromatography. Unknown stereocenters were assigned arbitrarily when naming the compounds.

[00120] LC-MS Methods

[00121] Method 1

HPLC System: Waters ACQUITY
 Column: Waters ACQUITY CSH™ C18 1.7 uM
 Guard column: Waters Assy. Frit, 0.2uM, 2.1 mm.
 Column tem: 40 °C
 Mobile Phase: A: TFA : Water (1 : 1000, v:v)
 B: TFA : ACN (1: 1000, v:v)
 Gradient Program:

Time (min)	B%
0	5
1.9	95
2.20	95
2.21	5

Flow Rate: 0.65 mL/min

Mass Spectrometer Parameters

Mass Spectrometer	Waters SQD	
Ionization	Positive Electrospray Ionization (ESI)	
Mode	Scan (100-1400 m/z in every 0.2 second)	
ES Capillary Voltage:	3.5 kv	
ES Cone Voltage:	25 v	
Source Temperature	120 °C	
Disolvation Temperature:	500 °C	
Desolvation Gas Flow:	Nitrogen	Setting 650 (L/hr)
Cone Gas Flow:	Nitrogen	Setting 50 (L/hr)

[00122] Method 2

5-95AB_1.5MIN

Column	MERCK,RP-18e 25-2mm	
Mobile Phase	A:water(4L)+TFA(1.5mL)	
	B:acetonitrile(4L)+TFA(0.75mL)	
	TIME(min)	B%
	0	5
	0.7	95
	1.1	95
	1.11	5
	1.5	5
Flow Rate	1.5mL/min	
wavelength	UV 220nm	
Oven Temp	50°C	
MS ionization	ESI	

[00123] Method 3

10-80CD_3MIN

Column	Xtimate C18 2.1*30mm,3um	
Mobile Phase	A:water(1L)+NH3H2O(0.5mL)	
	B:acetonitrile	
	TIME(min)	B%
	0	10
	2	80
	2.48	80
	2.49	10
	3	10
Flow Rate	1.0mL/min	
wavelength	UV 220nm&254nm	
Oven Temp	30°C	
MS ionization	ESI	

[00124] Method 4

10-80AB_2MIN

Column	Xtimate C18 2.1*30mm,3um	
Mobile Phase	A:water(4L)+TFA(1.5mL)	
	B:acetonitrile(4L)+TFA(0.75mL)	
	TIME(min)	B%
	0	10
	0.9	80
	1.5	80
	1.51	10

	2	10
Flow Rate	1.2mL/min	
wavelength	UV 220nm	
Oven Temp	50°C	

[00125] Method 5

10-80AB 2MIN

Column	XBridge Shield RP18 2.1*50mm	
Mobile Phase	A:water(4L)+TFA(1.5mL)	
	B:acetonitrile(4L)+TFA(0.75mL)	
	TIME(min)	B%
	0	10
	0.9	80
	1.5	80
	1.51	10
	2	10
	Flow Rate	1.2mL/min
wavelength	UV 220nm	
Oven Temp	50°C	


Method 6


5-95AB 1.5MIN

Column	YMC-Pack ODS-AQ	
Mobile Phase	A:water(4L)+TFA(1.5mL)	
	B:acetonitrile(4L)+TFA(0.75mL)	
	TIME(min)	B%
	0	5
	0.7	95
	1.1	95
	1.11	5
	1.5	5
	Flow Rate	1.5mL/min
wavelength	UV 220nm	
Oven Temp	50°C	
MS ionization	ESI	

Preparation of Intermediates

Preparation 1A: (R)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethanol





Step 1: (4-bromophenyl)(ethyl)sulfane

[00126] A mixture of 4-bromobenzenethiol (50 g, 0.26 mol), bromoethane (58 g, 0.53 mol) and triethylamine (78 g, 0.78 mol) in acetonitrile (1 L) was stirred at reflux for 17 h. The mixture was cooled to rt and filtered. The filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with petroleum ether) to give (4-bromophenyl)(ethyl)sulfane (55 g, 96%) as an oil. ¹H NMR (CDCl₃, 400 MHz): δ 7.40-7.42 (dd, *J* = 6.4, 2.0 Hz, 2H), 7.18-7.20 (dd, *J* = 6.4, 2.0 Hz, 2H), 2.91-2.96 (q, *J* = 7.2 Hz, 2H), 1.30-1.33 (t, *J* = 7.2 Hz, 3H).

Step 2: 2-((tert-butyldimethylsilyl)oxy)ethanol

[00127] To a solution of ethane-1,2-diol (110 g, 1.77 mol) in anhydrous CH₂Cl₂ (1.1 L) was added triethylamine (215.2 g, 296 mL, 2.13 mol) at rt. The mixture was cooled to 0 °C, then *tert*-butylchlorodimethylsilane (267.1 g, 1.77 mol) dissolved in CH₂Cl₂ (300 mL) was added dropwise over 1 h. The mixture was stirred at rt overnight. The reaction mixture was quenched with saturated aqueous NH₄Cl solution (400 mL) and separated. The aqueous phase was extracted with MTBE (2 × 400 mL). The combined organic layers were concentrated under vacuum and the residue was redissolved in MTBE (400 mL). The MTBE layer was washed with water (2 × 500 mL) and brine (500 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give 2-((*tert*-butyldimethylsilyl)oxy)ethanol (280 g, 90%) as a slight oil, which was used for the next step directly without further purification. ¹H NMR (CDCl₃, 400 MHz): δ 3.64-3.66 (m, 2H), 3.57-3.60 (m, 2H), 0.85 (s, 9H), 0.02 (s, 6H).

Step 3: 2-((tert-butyldimethylsilyl)oxy)acetaldehyde

[00128] To a solution of CH₂Cl₂ (1.8 L) cooled to -30 °C was added oxalyl chloride (79.2 g, 52.8 mL, 624 mmol) dropwise. The mixture was cooled to -78 °C, then DMSO (62.5 g, 88.5 mL, 1.25 mmol) was added dropwise. After addition, the mixture was stirred at -78 °C for 30 min. A

solution of 2-((*tert*-butyldimethylsilyl)oxy)ethanol (100 g, 567 mmol) dissolved in CH₂Cl₂ (200 mL) was added slowly at -78 °C. The reaction mixture was stirred at -78 °C for 1 h.

Triethylamine (287 g, 395 mL, 2.84 mmol) was added dropwise at -78 °C. The mixture was stirred at -78 °C for 30 min and then rt overnight. The reaction mixture was washed with water (1 L), 1 N HCl (2 × 1 L), saturated aqueous NaHCO₃ solution (1 L) and brine (1 L). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give 2-((*tert*-butyldimethylsilyl)oxy)acetaldehyde (98.5 g, 99.8%) as a brown oil, which was used for the next step directly without further purification. ¹H NMR (CDCl₃, 400 MHz): δ 9.70 (s, 1H), 4.22 (s, 2H), 0.93 (s, 9H), 0.11 (s, 6H).

Step 4: (*R,E*)-*N*-(2-((*tert*-butyldimethylsilyl)oxy)ethylidene)-2-methylpropane-2-sulfinamide

[00129] A mixture of 2-((*tert*-butyldimethylsilyl)oxy)acetaldehyde (93.5 g, 0.54 mol), (*R*)-2-methylpropane-2-sulfinamide (78.8 g, 0.65 mol) and copper (II) sulfate (215 g, 1.35 mol) in anhydrous CH₂Cl₂ (1.5 L) was stirred at rt for 16 h. The mixture was quenched with H₂O (800 mL) and separated. The aqueous phase was extracted with CH₂Cl₂ (2 × 1 L). The combined organic layers were washed with water (1 L) and brine (1 L), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with petroleum ether: ethyl acetate = 8:1) to give (*R,E*)-*N*-(2-((*tert*-butyldimethylsilyl)oxy)ethylidene)-2-methylpropane-2-sulfinamide (38.5 g, 26%) as a yellow oil. ¹H NMR (CDCl₃, 400 MHz): δ 7.96-7.97 (t, *J* = 3.2 Hz, 1H), 4.44-4.45 (d, *J* = 2.8 Hz, 2H), 1.11 (s, 9H), 0.00 (s, 6H).

Step 5: (*R*)-*N*-((*R*)-2-((*tert*-butyldimethylsilyl)oxy)-1-(4-(ethylthio)phenyl)ethyl)-2-methylpropane-2-sulfinamide

[00130] To a solution of (4-bromophenyl)(ethyl)sulfane (28.9 g, 133.1 mmol) in anhydrous THF (500 mL) was added dropwise *n*-butyllithium (73 mL, 181.5 mmol, 2.5 M in hexanes) at -78 °C. The mixture was stirred at -78 °C for 30 min. A solution of (*R,E*)-*N*-(2-((*tert*-butyldimethylsilyl)oxy)ethylidene)-2-methylpropane-2-sulfinamide (33.5 g, 121 mmol) in anhydrous THF (100 mL) was added to the mixture at -78 °C. The mixture was stirred at -78 °C for 2 h, then allowed to warm to rt and stirred for 2 h. The mixture was quenched with saturated aqueous NH₄Cl solution (200 mL) and extracted with ethyl acetate (3 × 300 mL). The combined organic layer was washed with water (200 mL) and brine (200 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with petroleum ether: ethyl acetate = 15:1) three times to afford (*R*)-*N*-


((*R*)-2-((*tert*-butyldimethylsilyl)oxy)-1-(4-(ethylthio)phenyl)ethyl)-2-methylpropane-2-sulfinamide (22 g, 44%) as a yellow oil. ¹H NMR (CDCl₃, 400 MHz): δ 7.21-7.24 (d, *J* = 7.2 Hz, 2H), 7.18-7.21 (d, *J* = 8.4 Hz, 2H), 4.42-4.45 (dd, *J* = 8.8, 2.4 Hz, 1H), 4.21 (brs, 1H), 3.69-3.73 (dd, *J* = 10.4, 4.4 Hz, 1H), 3.51-3.56 (t, *J* = 9.6 Hz, 1H), 2.87-2.92 (q, *J* = 7.6 Hz, 2H), 1.25-1.29 (t, *J* = 7.2 Hz, 3H), 1.18 (s, 9H), 0.88 (s, 9H), 0.02 (s, 6H). LCMS *t*_R = 1.010 min in 5-95AB_1.5 min chromatography (MK RP18e 25-2mm), MS (ESI) *m/z* 437.9 [M+Na]⁺. **Isomer SFC** *t*_R = 3.607 and 4.014 min in 12 min chromatography (AD-H_5_5_40_2.3 5 ML), ee = 90.85%.

Step 6: (*R*)-2-amino-2-(4-(ethylthio)phenyl)ethanol

[00131] To a solution of (*R*)-*N*-((*R*)-2-((*tert*-butyldimethylsilyl)oxy)-1-(4-(ethylthio)phenyl)ethyl)-2-methylpropane-2-sulfinamide (22 g, 52.9 mmol) in CH₂Cl₂ (250 mL) was added HCl (26.5 mL, 4 N in dioxane) at 0 °C. The mixture was stirred at rt for 2 h. LCMS showed no starting material remaining. The mixture was concentrated under reduced pressure to afford crude (*R*)-2-amino-2-(4-(ethylthio)phenyl)ethanol HCl salt (12.3 g, 100%) as a brown solid, which was used for the next step directly without further purification. LCMS *t*_R = 1.226 min in 0-30AB_2 min chromatography (Xtimate 3um, C18, 2.1*30mm), MS (ESI) *m/z* 180.9 [M-OH]⁺.

Step 7: (*R*)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethanol

[00132] To a mixture of (*R*)-2-amino-2-(4-(ethylthio)phenyl)ethanol (15.2 g, 65.0 mmol) in methanol (200 mL) was added dropwise a solution of oxone reagent (80.0 g, 130.0 mmol) in water (200 mL) at 0 °C. The mixture was stirred at rt for 1.5 h; LCMS showed no starting material remaining. The mixture was filtered and methanol was removed under reduced pressure. The aqueous phase was extracted with EtOAc (2 × 80 mL), then the aqueous layer was basified to pH = 8-9 with solid sodium carbonate portionwise at 0 °C, then this solution was lyophilized (contained the Na₂CO₃). The solid was dissolved in CH₂Cl₂:MeOH (3:1, 600 mL) and stirred for 30 min, filtered, then concentrated under reduced pressure. The residue was purified by silica gel chromatography (eluting with CH₂Cl₂:MeOH = 1:0 to 4:1) to give (*R*)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethanol (11.5 g, 77%) as a white solid. LC-MS *t*_R = 0.738 min in 0-30CD_POS chromatography (Xtimate ODS 2.1*30mm,3um), MS (ESI) *m/z* 230.1 [M+H]⁺. **Isomer SFC** *t*_R = 6.99 min in 30 min chromatography (CD-PH_10-80_B_08 ML), ee = 97.42%. ¹H NMR (D₂O, 400 MHz): δ 7.82-7.84 (d, *J* = 8.0 Hz, 2H), 7.54-7.56 (d, *J* = 8.4 Hz, 2H), 4.33-4.35 (t, *J* = 6.4 Hz, 1H), 3.72-3.78 (m, 2H), 3.19-3.25 (q, *J* = 7.6 Hz, 2H), 1.03-1.07 (t, *J* = 7.6 Hz, 3H).

[00133] Preparation 1B: (R)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethanolStep 1: 4-(ethylsulfonyl)benzaldehyde

[00134] To a solution of 4-fluorobenzaldehyde (24.6 g, 198 mmol) in dimethylsulfoxide (60 mL) was added sodium ethanesulfinate (46 g, 396 mmol). The resulting mixture was stirred at 125°C for 20 h. After cooling to rt, the reaction mixture was triturated with 350 mL of H_2O . The product was filtered, washed with two 10-mL portions of EtOH and dried under vacuum to afford 4-(ethylsulfonyl)benzaldehyde as a light yellow solid (31.2 g, 80% yield). **LC-MS** $t_R = 1.19$ min in 2 min chromatography, MS (ESI) m/z 199.1 $[\text{M}+\text{H}]^+$. **^1H NMR** (CDCl_3) δ 10.14 (s, 1H), 8.09 (s, 4H), 3.16 (q, $J = 7.2$ Hz, 2H), 1.30 (t, $J = 7.2$ Hz, 3H).

Step 2: 2-(4-(ethylsulfonyl)phenyl)oxirane

[00135] To a solution of 4-(ethylsulfonyl)benzaldehyde (10 g, 50.5 mmol) in DMF (85 mL) at rt was added trimethylsulfonium iodide (11.9 g, 58.1 mmol) followed by potassium hydroxide powder (5.66 g, 101 mmol). The reaction mixture was stirred at rt for 20 min before quenching with H_2O (50 mL). The mixture was carefully neutralized with 1 N HCl solution (55 mL) and extracted with ethyl acetate (3 x 100 mL). The combined organic phase was washed with brine, dried over anhydrous Na_2SO_4 , and passed through a pad of silica gel (eluting with ethyl acetate). It was concentrated under reduced pressure to afford crude 2-(4-(ethylsulfonyl)phenyl)oxirane as yellow oil, which was used directly for the next step without further purification. **LC-MS** $t_R = 1.13$ min in 2 min chromatography, MS (ESI) m/z 213.2 $[\text{M}+\text{H}]^+$.


Step 3: 2-amino-2-(4-(ethylsulfonyl)phenyl)ethan-1-ol


[00136] To a solution of crude 2-(4-(ethylsulfonyl)phenyl)oxirane (50.5 mmol) in CH₃CN (200 mL) at 0 °C was slowly added concentrated sulfuric acid (5.4 mL, 101 mmol). The mixture was allowed to stir at rt for 1.5 h. LCMS showed the starting material was consumed. H₂O (15 mL) was added to the reaction mixture. Stirring continued at rt for 8 h, then at 45 °C for 10 h. After cooling to rt, the pH of the reaction mixture was adjusted to 3-4 by addition of 1 N NaOH solution (90 mL). The mixture was extracted with ethyl acetate (100 mL). The organic phase was then extracted with H₂O (2 x 30 mL). The combined aqueous layers were then basified with 1 N NaOH solution (110 mL) to pH = 9 and extracted with 1-butanol (5 x 60 mL). The combined organic layer (consisting of 1-butanol extracts) was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. It was dried under high vacuum to afford crude 2-amino-2-(4-(ethylsulfonyl)phenyl)ethan-1-ol as an off-white solid. 4 g, 35% yield over 3 steps.

Intermediate 4-(4-(ethylsulfonyl)phenyl)-2-methyl-4,5-dihydrooxazole: LC-MS t_R = 0.77, 0.81 min in 2 min chromatography, MS (ESI) m/z 254.26 [M + H]⁺. **2-amino-2-(4-(ethylsulfonyl)phenyl)ethan-1-ol:** LC-MS t_R = 0.61 min in 2 min chromatography, MS (ESI) m/z 230.21 [M + H]⁺. ¹H NMR (CD₃OD): δ 7.88 (d, J = 8.4 Hz, 2H), 7.64 (d, J = 8.4 Hz, 2H), 4.16-4.12 (m, 1H), 3.76-3.72 (m, 1H), 3.66-3.61 (m, 1H), 3.17 (q, J = 7.2 Hz, 2H), 1.19 (t, J = 7.2 Hz, 3H).

Step 4: 2-amino-2-(4-(ethylsulfonyl)phenyl)ethan-1-ol mono-mandelate salt

[00137] To a solution of 2-amino-2-(4-(ethylsulfonyl)phenyl)ethan-1-ol (238 mg, 1.0 mmol) in MeOH (3 mL) at 50 °C was added a solution of (*R*)-Mandelic acid (76 mg, 0.5 mmol) in MeOH (1 mL). The resulting solution was allowed to cool down to ambient temperature slowly. After stirring for 1 day, the resulting crystals were collected by vacuum filtration and dried under high vacuum, providing the mono-mandelate salt as a white crystal, 107 mg (28% yield), 92.5% ee. ¹H NMR (CD₃OD): δ 7.97 (d, J = 8.0 Hz, 2H), 7.71 (d, J = 8.4 Hz, 2H), 7.46 (d, J = 8.0 Hz, 2H), 7.46 (d, J = 8.0 Hz, 2H), 7.31-7.27 (m, 2H), 7.25-7.22 (m, 1H), 4.42-4.42 (m, 1H), 3.92-3.89 (m, 1H), 3.81-3.77 (m, 1H), 3.21 (q, J = 7.2 Hz, 2H), 1.21 (t, J = 7.2 Hz, 3H).

Preparation 2: (*R*)-2-amino-2-(5-(ethylsulfonyl)pyridin-2-yl)ethanol



Step 1: 2-bromo-5-(ethylthio)pyridine

[00138] To a mixture of 2-bromo-5-fluoropyridine (6.28 g, 35.66 mmol) in anhydrous DMF (60 mL) was added sodium ethanethiolate (3 g, 35.66 mmol). The mixture was stirred at 100 °C for 3 h. TLC (petroleum ether / ethyl acetate 10/1) showed that the starting material was not consumed completely. Additional sodium ethanethiolate (0.9 g, 9.56 mmol) was added to the mixture. The mixture was stirred at 100 °C for 12 h. The mixture was quenched with H₂O (150 mL) and extracted with ethyl acetate (3 x 150 mL). The combined organic layers were washed with brine (400 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (eluting with petroleum ether / ethyl acetate 80/1) to afford 2-bromo-5-(ethylthio)pyridine (7.0 g, 90%) as a colorless oil. **LC-MS** t_R = 0.717 min in 5-95AB_1.5 min chromatography (Welch Merck RP-18e 25-2mm), **MS** (ESI) m/z 217.6 [M+H]⁺.

Step 2: (R)-N-((R)-2-((tert-butyldimethylsilyl)oxy)-1-(5-(ethylthio)pyridin-2-yl)ethyl)-2-methylpropane-2-sulfinamide

[00139] To a solution of toluene (60 mL) was added *n*-BuLi (10.6 mL, 26.48 mmol, 2.5 M in hexanes) dropwise at -78 °C; the internal temperature did not exceed -50 °C. A solution of 2-bromo-5-(ethylthio)pyridine (3.85 g, 17.65 mmol) in toluene (10 mL) was then added to the reaction mixture at -78 °C; the internal temperature did not exceed -65 °C. The mixture was stirred at -78 °C for 1 h. A solution of (*R,E*)-*N*-(2-((tert-butyldimethylsilyl)oxy)ethylidene)-2-methylpropane-2-sulfinamide (4.90 g, 17.65 mmol) in toluene (10 mL) was added to the reaction mixture at -78 °C; the internal temperature did not exceed -60 °C. The mixture was stirred at -78 °C for another 2 h. The mixture was quenched with brine (150 mL) at -78 °C and extracted with ethyl acetate (3 x 150 mL). The combined organic layers were washed with brine (400 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (eluting with petroleum ether / ethyl acetate 10/1 to 3/1) to afford (*R*)-*N*-((*R*)-2-((tert-butyldimethylsilyl)oxy)-1-(5-(ethylthio)pyridin-2-yl)ethyl)-2-methylpropane-2-sulfinamide (3.0 g, 41%) as a pale yellow oil. **LC-MS** t_R = 1.014

min in 5-95AB_1.5 min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) m/z 417.2 $[M+H]^+$.


Step 3: (*R*)-2-amino-2-(5-(ethylthio)pyridin-2-yl)ethanol

[00140] Procedure same as that for (*R*)-2-amino-2-(4-(ethylthio)phenyl)ethanol with (*R*)-*N*-((*R*)-2-((tert-butyldimethylsilyl)oxy)-1-(5-(ethylthio)pyridin-2-yl)ethyl)-2-methylpropane-2-sulfonamide as the starting material.

Step 4: (*R*)-2-amino-2-(5-(ethylsulfonyl)pyridin-2-yl)ethanol


[00141] Procedure same as that for (*R*)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethanol with (*R*)-2-amino-2-(5-(ethylthio)pyridin-2-yl)ethanol as the starting material. $^1\text{H NMR}$ (CD_3OD , 400 MHz): δ 9.08 (s, 1H), 8.35 (dd, $J = 2.0, 8.4$ Hz, 1H), 7.79 (d, $J = 8.4$ Hz, 1H), 4.70 (t, $J = 5.6$ Hz, 1H), 4.03 (dd, $J = 4.8, 12.0$ Hz, 1H), 3.91 (dd, $J = 4.8, 11.6$ Hz, 1H), 3.29 (q, $J = 7.2$ Hz, 2H), 1.25 (t, $J = 7.2$ Hz, 3H).

Preparation 3: (*R*)-2-amino-2-(4-(methylsulfonyl)phenyl)ethanol



[00142] The compound was prepared analogously to (*R*)-2-amino-2-(4-(methylsulfonyl)phenyl)ethanol (B1).


Preparation 4: (*R*)-2-amino-2-(5-(methylsulfonyl)pyridin-2-yl)ethanol



[00143] The compound was prepared analogously to (*R*)-2-amino-2-(5-(ethylsulfonyl)pyridin-2-yl)ethanol (B5).

Example 1

N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide (Cpd No Hy1A-5)



[00144] Step 1

[00145] To a mixture of *trans*-4-(trifluoromethyl)cyclohexanecarboxylic acid (200 mg, 1.02 mmol) in anhydrous THF (4 mL) was added dropwise LiAlH_4 (2 mL, 2.0 mmol, 1 M in THF) at 0°C under N_2 . The mixture was stirred at 0°C for 2 h under N_2 . TLC (petroleum ether / ethyl acetate = 4/1) showed that the reaction was completed. The mixture was quenched with sat. Na_2SO_4 solution (1 mL) and diluted with CH_2Cl_2 (20 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to afford crude *trans*-4-(trifluoromethyl)cyclohexylmethanol (180 mg, 96%) as a yellow oil, which was used for the next step directly without further purification. $^1\text{H NMR}$ (CDCl_3 , 400 MHz): δ 3.41 (t, J = 5.6 Hz, 2H), 2.00-1.80 (m, 4H), 1.35-1.15 (m, 4H), 1.03-0.88 (m, 2H).

[00146] Step 2

[00147] To a mixture of crude *trans*-4-(trifluoromethyl)cyclohexyl)methanol (80 mg, 0.439 mmol) in anhydrous toluene (5 mL) was added methyl 1-chloroisoquinoline-6-carboxylate (35 mg, 0.158 mmol), Pd₂(dba)₃ (35 mg, 0.038 mmol), BINAP (35 mg, 0.0553 mmol) and Cs₂CO₃ (103 mg, 0.316 mmol). The mixture was stirred at 110 °C for 5 h under N₂. The mixture diluted with ethyl acetate (30 mL) and washed with brine (80 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC with petroleum ether / ethyl acetate = 6/1 to afford methyl 1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxylate (57 mg, 100%) as a yellow solid. **LC-MS** t_R = 0.953 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 368.0 [M+H]⁺.

[00148] Step 3

[00149] To a mixture of methyl 1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxylate (57 mg, 0.155 mmol) in MeOH (5 mL) and H₂O (1 mL) was added NaOH (124 mg, 3.106 mmol). The mixture was stirred at rt for 1.5 h. The mixture was concentrated under reduced pressure. The residue was diluted with H₂O (20 mL) and adjusted to pH = 4-5 with 1 N HCl solution. The aqueous layer was extracted with ethyl acetate (3 X 20 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to afford crude 1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxylic acid (55 mg, 100%) as a yellow solid, which was used for the next step directly without further purification. **LC-MS** t_R = 0.841 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 353.9 [M+H]⁺.

[00150] Step 4

[00151] To a mixture of crude 1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxylic acid (48 mg, 0.136 mmol) in CH₂Cl₂ (5 mL) was added (R)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethanol HCl salt (36 mg, 0.136 mmol), EDCI (52 mg, 0.272 mmol), HOBt (37 mg, 0.272 mmol) and Et₃N (41 mg, 0.408 mmol). The mixture was stirred at rt for 16 h. The mixture was quenched with H₂O (20 mL) and extracted with CH₂Cl₂ (3 X 20 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC with ethyl acetate, basic preparative HPLC separation, then dry-freezing directly to afford N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-((*trans*-4-

(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide (11.10 mg, 14%) as a white solid. **LC-MS** t_R = 0.810 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), **MS** (ESI) m/z 565.1 $[M+H]^+$. **Isomer SFC** t_R = 3.834 min in 8 min chromatography (Column: AS-H; Method Name: AS-H_S_5_5_40_3ML_8MIN_15CM..M, ee = 97%) **^1H NMR** ($CDCl_3$ 400 MHz): δ 8.26 (d, J = 8.8 Hz, 1H), 8.14 (s, 1H), 7.98 (d, J = 5.6 Hz, 1H), 7.93-7.83 (m, 3H), 7.56 (d, J = 8.0 Hz, 2H), 7.25-7.18 (m, 2H), 5.35-5.27 (m, 1H), 4.28 (d, J = 6.0 Hz, 2H), 4.15-3.95 (m, 2H), 3.04 (q, J = 7.6 Hz, 2H), 2.15-1.90 (m, 6H), 1.45-1.32 (m, 2H), 1.28-1.15 (m, 5H).

[00152] Basic preparative HPLC Method:

Mobile phase A: water with 0.05% ammonia hydroxide solution

Mobile phase B: MeCN

Flow rate: 25 mL/min.


Detection: UV 220 nm

Column: Phenomenex Gemini 150*25mm*10um

Column temperature: 40 °C

Time in min	%A	%B
0.00	49	51
10.00	19	81
10.20	0	100
13.00	0	100

[00153] The compounds listed below were prepared using analogous procedures:




Cpd No	Cy ²	L ²	R ⁷	R ⁸
Hy1A-1	4-tetrahydropyranyl	O	H	H
Hy1A-2	4-(trifluoromethyl)phenyl	O	H	H
Hy1A-3.1 ^a	4-(trifluoromethyl)cyclohexyl	O	H	H
Hy1A-3.2 ^a	4-(trifluoromethyl)cyclohexyl	O	H	H
Hy1A-4	4-(trifluoromethyl)phenyl	CH ₂ O	H	H
Hy1A-7	4-(trifluoromethyl)phenyl	CH ₂ O	CH ₂ OH	H
Hy1A-8	<i>trans</i> -4-(trifluoromethyl)cyclohexyl	CH ₂ O	CH ₂ OH	H

a cis and trans isomers separated by chromatography.

Example 2

N-((*R*)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide (Cpd No Hy1A-9)



[00154] Step 1

[00155] To a solution of 1,3-dichloroisoquinoline-6-carboxylic acid (80 mg, 0.33 mmol) in MeOH (3 mL) was added SOCl₂ (1 mL) slowly at 0 °C. The mixture was stirred at 40 °C for 1 h. LCMS showed that the reaction was completed. The reaction solution was concentrated under reduced pressure to afford crude methyl 1,3-dichloroisoquinoline-6-carboxylate (85 mg, 100%) as a white solid, which was used for the next step directly without further purification.

LC-MS: *t_R* = 0.796 min in 5-95 AB_1.5 MIN chromatography (MERCK RP-18e 25-2 mm), MS (ESI) *m/z* 255.7 [M+H]⁺.

[00156] Step 2

[00157] To a mixture of crude methyl 1,3-dichloroisoquinoline-6-carboxylate (50 mg, 0.2 mmol), Cs₂CO₃ (130 mg, 0.4 mmol), BINAP (7 mg, 0.01 mmol), and Pd₂dba₃ (10 mg, 0.01

mmol) was added a solution of (*trans*-4-(trifluoromethyl)cyclohexyl)methanol (36 mg, 0.2 mmol) in anhydrous toluene (4 mL) under N₂. The mixture was stirred at 105 °C under N₂ for 3 h. LCMS showed that the reaction was completed. The reaction mixture was filtered, the filtrate was concentrated under reduced pressure. The residue was purified by preparative TLC (petroleum ether: dichloromethane = 2:1) to afford methyl 3-chloro-1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxylate (45 mg, 56%) as a white solid. **LC-MS:** *t_R* = 1.007 min in 5-95 AB_1.5 MIN chromatography (MERCK RP-18e 25-2 mm), MS (ESI) *m/z* 402.0 [M+H]⁺. **¹H NMR** (CDCl₃ 400MHz): δ 8.38 (d, *J* = 1.2 Hz, 1H), 8.25 (d, *J* = 8.4 Hz, 1H), 8.09 (dd, *J* = 8.4, 1.2 Hz, 1H), 7.34 (s, 1H), 4.37 (d, *J* = 6.4 Hz, 2H), 4.00 (s, 3H), 2.15-1.90 (m, 5H), 1.45-1.35 (m, 2H), 1.25-1.17 (m, 3H). **NOSEY** confirmed structure.

[00158] Step 3

[00159] A mixture of methyl 3-chloro-1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxylate (140 mg, 0.35 mmol), CH₃B(OH)₂ (63 mg, 1.05 mmol), Cs₂CO₃ (228 mg, 0.70 mmol) and Pd(dppf)Cl₂ (26 mg, 0.035 mmol) in dioxane (0.5 mL) and H₂O (0.1 mL) was stirred at 100 °C in sealed tube for 8 h. TLC (petroleum ether: dichloromethane = 2:1) showed that the reaction was completed. The reaction solution was filtered. The filtrate was added with water (30 mL), extracted with ethyl acetate (3 x 15 mL). The combined organic layers were washed with brine (30 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC (petroleum ether: dichloromethane = 2:1) to afford methyl 3-methyl-1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxylate (85 mg, 64%) as a colorless oil. **LC-MS:** *t_R* = 1.017 min in 5-95 AB_1.5 MIN chromatography (MERCK RP-18e 25-2mm), MS (ESI) *m/z* 382.0 [M+H]⁺.

[00160] Step 4

[00161] To a solution of methyl 3-methyl-1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxylate (80 mg, 0.21 mmol) in MeOH (5 mL) was added 2N *aq.* NaOH (1 mL). The mixture was stirred at 40 °C for 1.5 h. TLC (petroleum ether: dichloromethane = 2:1) showed that the reaction was completed. The reaction solution was added with water (20 mL), concentrated under reduced pressure to remove MeOH. The residue was acidified by 4N HCl to pH = 3, extracted with ethyl acetate (3 x 10 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude 3-methyl-1-((*trans*-4-

(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxylic acid (77 mg, 100%) as a white solid, which was used for the next step directly without further purification.

[00162] Step 5

[00163] To a solution of crude 3-methyl-1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxylic acid (40 mg, 0.11 mmol) in CH₂Cl₂ (2 mL) was added (R)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethanol hydrochloride salt (43 mg, 0.16 mmol), HATU (60 mg, 0.16 mmol) and Et₃N (32 mg, 0.32 mmol). The mixture was stirred 25 °C for 15 h. LCMS showed that the reaction was completed. The reaction solution was concentrated under reduced pressure. The residue was purified by preparative TLC (petroleum ether: ethyl acetate = 1:1), basic preparative HPLC separation and dry-freezing to afford *N*-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-3-methyl-1-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)isoquinoline-6-carboxamide (**Hy1A-9**, 13.70 mg, 22%) as a white solid. **LC-MS**: *t_R* = 0.829 min in 5-95 AB_1.5 MIN chromatography (MERCK RP-18e 25-2 mm), MS (ESI) *m/z* 579.1 [M+H]⁺. **¹H NMR** (CDCl₃, 400MHz): δ 8.25 (d, *J* = 8.4 Hz, 1H), 8.10 (d, *J* = 1.2 Hz, 1H), 7.89 (d, *J* = 8.4 Hz, 2H), 7.82 (d, *J* = 6.8 Hz, 1H), 7.61 (d, *J* = 8.4 Hz, 2H), 7.28 (s, 1H), 7.08 (s, 1H), 5.40-5.30 (m, 1H), 4.35 (d, *J* = 6.0 Hz, 2H), 4.15-4.00 (m, 2H), 3.10 (q, *J* = 7.6 Hz, 2H), 2.54 (s, 3H), 2.12-1.88 (m, 6H), 1.45-1.35 (m, 2H), 1.29 (t, *J* = 7.6 Hz, 3H), 1.25-1.15 (m, 2H). **Isomer SFC**: *t_R* = 1.661 min in 3 min chromatography (Column: AD-H; Method Name: AD-H_3UM_4_5_40_4ML_3MIN.M, ee = 96.60%).

[00164] Basic preparative HPLC method

Instrument: DC

Mobile phase A: water (0.05% ammonia hydroxide v/v)

Mobile phase B: CH₃CN

Flow rate: 25 mL/min.


Detection: UV 220 nm / 254 nm

Column: Phenomenex Gemini 150*25mm*10um

Column temperature: 30 °C





Time in min	%A	%B
0.00	38	62
10.00	18	82
10.20	0	100
13.00	0	100

[00165] The following compound was prepared using procedures analogous to those described above:




The following compounds are prepared using procedures analogous to those described above;

Cpd No Hy1A-12	Cpd No Hy1A-13
Cpd No Hy1A-14	Cpd No Hy1A-15


	
Cpd No Hy1A-16	Cpd No Hy1A-17
	
Cpd No Hy1A-18.1, -18.2, -18.3 ^a	Cpd No Hy1A-19

^a The four isomers were separated chromatographically into three fractions. Fraction 18.1 is one trans isomer. Fraction 18.2 is the other trans isomer. Fraction 18.3 is a mixture of two cis isomers.


The following compounds are prepared following procedures analogous to those described above, using the appropriate boronic acid in Step 3 in place of methyl boronic acid:




The following compounds are prepared following procedures analogous to those described above, using 2-amino-2-(5-(ethylsulfonyl)pyridin-2-yl)ethan-1-ol in Step 5:



The following compound is prepared following procedures analogous to those described above, using $\text{Pd}_2(\text{dba})_3$, bippyphos, Cs_2CO_3 , MeOH at 60 °C in Step 3:




[00166] The following compounds were prepared using procedures analogous to those described above, omitting Step 3:



Example 3

(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-methyl-4-((4-(trifluoromethyl)benzyl)oxy)quinoline-7-carboxamide (Cpd No Hy1B-3)



[00167] Step 1

[00168] To a solution of 4-chloro-2-methylquinoline-7-carboxylic acid (150 mg, 0.679 mmol) in MeOH (3 mL) was added SOCl₂ (3 mL). The mixture was stirred at 45 °C for 3 h. The mixture was concentrated under reduced pressure to afford crude methyl 4-chloro-2-methylquinoline-7-carboxylate (158 mg, 100 %) as a white solid, which was used for next step directly without further purification. **LC-MS** *t_R* = 0.673 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), **MS** (ESI) *m/z* 235.9 [M+H]⁺.

[00169] Step 2

[00170] To a mixture of crude methyl 4-chloro-2-methylquinoline-7-carboxylate (70 mg, 0.3 mmol), (4-(trifluoromethyl)phenyl)methanol (79 mg, 0.45 mmol), BINAP (37 mg, 0.06 mmol), Cs₂CO₃ (291 mg, 0.9 mmol) in anhydrous toluene (2 mL) was added Pd₂(dba)₃ (55 mg, 0.06 mmol) under N₂. The mixture was stirred at 110 °C for 16 h under N₂. TLC (petroleum ether / ethyl acetate = 3/1) showed that the reaction was completed. The mixture was filtered through celite and the filter cake was washed with CH₂Cl₂ (20 mL). The filtrate was concentrated under reduced pressure. The residue was purified by preparative TLC (petroleum ether / ethyl acetate = 3 / 1) to afford methyl 2-methyl-4-((4-(trifluoromethyl)benzyl)oxy)quinoline-7-carboxylate (25 mg, 22.5%) as a white solid. **LC-MS t_R** = 0.685 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 375.9 [M+H]⁺.

[00171] Step 3

[00172] A solution of methyl 2-methyl-4-((4-(trifluoromethyl)benzyl)oxy)quinoline-7-carboxylate (35 mg, 0.093 mmol), NaOH (186 mg, 4.6 mmol) in MeOH (2 mL) and H₂O (1 mL) was stirred at rt for 4 h. LCMS showed that the reaction was completed. The mixture was adjusted to pH = 3 with 1 N HCl solution. The mixture was added with water (5 mL) and extracted with CH₂Cl₂ (3 X 5 mL). The combined organic layers were washed with brine (10 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude 2-methyl-4-((4-(trifluoromethyl)benzyl)oxy)quinoline-7-carboxylic acid (35 mg, 100 %) as a yellow oil, which was used for the next step directly without further purification. **LC-MS t_R** = 0.653 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 361.9 [M+H]⁺.

[00173] Step 4

[00174] To a mixture of crude 2-methyl-4-((4-(trifluoromethyl)benzyl)oxy)quinoline-7-carboxylic acid (20 mg, 0.055 mmol), (R)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethanol HCl salt (25 mg, 0.11 mmol), HATU (42 mg, 0.11 mmol) in anhydrous CH₂Cl₂ (1 mL) was added Et₃N (11 mg, 0.11 mmol). LCMS showed that the reaction was completed. The mixture was stirred at rt for 4 h. The mixture was purified by preparative TLC (petroleum ether / ethyl acetate = 1 / 4) to afford a crude product, which was separated by basic preparative HPLC separation and dry-freezing directly to afford (R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-methyl-4-((4-(trifluoromethyl)benzyl)oxy)quinoline-7-carboxamide (**Hy1B-3**, 1.80 mg, 5%) as a white solid. **LC-MS t_R** = 0.678 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS

(ESI) m/z 573.0 $[M+H]^+$. $^1\text{H NMR}$ (CDCl_3 , 400 MHz): δ 8.97-8.95 (m, 1H), 8.81 (s, 1H), 8.18 (d, $J = 9.2$ Hz, 1H), 7.99 (d, $J = 8.4$ Hz, 1H), 7.68 (d, $J = 8.4$ Hz, 1H), 7.67 (s, 1H), 7.59 (d, $J = 8.0$ Hz, 4H), 7.50 (d, $J = 8.0$ Hz, 2H), 6.85 (s, 1H), 5.44-5.40 (m, 2H), 5.30-5.20 (m, 1H), 4.09 (d, $J = 4.8$ Hz, 2H), 2.96 (q, $J = 7.2$ Hz, 2H), 2.82 (s, 3H), 1.16 (t, $J = 7.2$ Hz, 3H). **Isomer SFC** $t_R = 6.814$ min in 10 min chromatography (Column: AD-H; Method Name: AD-3_ETOH(DEA)_5_40_25ML.M, ee = 100%).

[00175] Basic preparative HPLC Method:

Mobile phase A: water with 0.05% ammonia hydroxide solution

Mobile phase B: MeCN

Flow rate: 25 mL/min.


Detection: UV 220 nm

Column: Gemini 150*25mm*10um

Column temperature: 40 °C

Time in min	%A	%B
0.00	60	40
10.00	30	70
10.20	0	100
13.00	0	100


[00176] The compounds shown below are prepared following analogous procedures:



Cpd No	Cy ²	L ²	R ⁷	R ⁸
Hy1B-1	4-(trifluoromethyl)phenyl	CH ₂ O	H	H
Hy1B-2	<i>trans</i> -4-(trifluoromethyl)cyclohexyl	CH ₂ O	H	H
Hy1B-4	<i>trans</i> -4-(trifluoromethyl)cyclohexyl	CH ₂ O	CH ₂ OH	H

Example 4

N-(4-(ethylsulfonyl)benzyl)-4-(((1*r*,4*r*)-4-(trifluoromethyl)cyclohexyl)methoxy)quinazoline-7-carboxamide (Cpd No Hy1C-2)



[00177] Step 1

[00178] To a solution of *trans*-4-(trifluoromethyl)cyclohexanecarboxylic acid (0.5 g, 2.6 mmol) in anhydrous THF (5 mL) was added LiAlH₄ (5.1 mL, 5.1 mmol, 1 M in THF) at 0 °C under N₂. The mixture was stirred at 0 - 25 °C under N₂ for 2 h. TLC (petroleum ether: ethyl acetate = 2: 1) showed the reaction was completed. The reaction solution was added with ice water (0.2 mL), 10% *aq.* NaOH (0.2 mL) and then water (0.1 mL) dropwise at 0 °C. The mixture was stirred at 0 - 20 °C for 0.5 h. Solid Na₂SO₄ (1 g) was added and then the mixture was stirred at rt for 0.5 h. The mixture was filtered and washed with ethyl acetate (3 X 10 mL). The combined organic layers were concentrated under reduced pressure to afford crude *trans*-4-(trifluoromethyl)cyclohexylmethanol (450 mg, 97%) as a colorless oil, which was used for the next step directly without further purification. ¹H NMR (CDCl₃ 400 MHz): δ 3.48 (d, *J* = 6.0 Hz, 2H), 2.01-1.90 (m, 4H), 1.35-1.30 (m, 3H), 1.02-0.94 (m, 2H).

[00179] Step 2

[00180] To a mixture of crude methyl 4-chloroquinazoline-7-carboxylate (100 mg, 0.45 mmol), *trans*-4-(trifluoromethyl)cyclohexylmethanol (123 mg, 0.68 mmol), Cs₂CO₃ (293 mg, 0.90 mmol) and BINAP (112 mg, 0.18 mmol) in toluene (5 mL) was added Pd₂(dba)₃ (165 mg, 0.18 mmol). The resulting mixture was stirred at 110 °C under N₂ for 20 h. LCMS showed the reaction was completed. The reaction solution was filtered. The filtrate was added with water (30 mL), extracted with ethyl acetate (3 x 15 mL). The mixture was washed with brine (3 X 10 mL) and then dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure.

The residue was purified by preparative TLC (petroleum ether / ethyl acetate = 5 / 1) to afford methyl 4-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)quinazoline-7-carboxylate (110 mg, 66 %) as a pale yellow solid. **LC-MS** t_R = 0.880 min in 5-95AB_1.5min chromatography (Merck RP-18e 25-2mm), MS (ESI) m/z 369.0 $[M+H]^+$.

[00181] Step 3

[00182] To a solution of methyl 4-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)quinazoline-7-carboxylate (70 mg, 0.19 mmol) in THF (3 mL) was added NaOH (76 mg, 1.9 mmol) and water (1.0 mL). The reaction mixture was stirred at rt for 5 h. LCMS showed the reaction was completed. The solvents were removed under reduced pressure. Water (10 mL) was added. The mixture was acidified with 1 N HCl solution to pH = 5. The mixture was extracted with ethyl acetate (3 X 15 mL). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to afford crude 4-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)quinazoline-7-carboxylic acid (67 mg, 100% yield) as a white solid, which was used for the next step directly without further purification. **LC-MS** t_R = 0.778 min in 5-95AB_1.5min chromatography (Merck RP-18e 25-2mm), MS (ESI) m/z 354.9 $[M+H]^+$.

[00183] Step 4

[00184] To a solution of 4-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)quinazoline-7-carboxylic acid (20 mg, 0.056 mmol), (4-(ethylsulfonyl)phenyl)methanamine hydrochloride (16 mg, 0.068 mmol) and HATU (43 mg, 0.11 mmol) in dry DMF (2 mL) was added Et₃N (17 mg, 0.17 mmol). The mixture was stirred at rt for 2 h. LCMS showed the starting material was consumed completely. Ethyl acetate (10 mL) was added. The mixture was washed with brine (2 X 10 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by basic preparative HPLC separation to afford *N*-(4-(ethylsulfonyl)benzyl)-4-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)quinazoline-7-carboxamide (**Hy1C-2**, 13.50 mg, 12%) as a white solid. **LC-MS** t_R = 0.795 min in 5-95AB_1.5min chromatography (MK RP-18e 25-2 mm), MS (ESI) m/z 536.1 $[M+H]^+$. **¹H NMR** (DMSO-*d*₆ 400 MHz): δ 9.59 (t, J = 5.6 Hz, 1H), 8.87 (s, 1H), 8.47 (s, 1H), 8.29 (d, J = 8.8 Hz, 1H), 8.13 (dd, J = 1.2, 8.4 Hz, 1H), 7.87 (d, J = 8.4 Hz, 2H), 7.64 (d, J = 8.0 Hz, 2H), 4.65 (d, J = 6.0 Hz, 2H), 4.43 (d, J = 6.0 Hz, 2H), 3.29 (q, J = 7.2 Hz, 2H), 2.34-2.28 (m, 1H), 2.04-1.90 (m, 5H), 1.40-1.24 (m, 4H), 1.10 (t, J = 7.2 Hz, 3H).

Isomer SFC t_R = 1.792 min in 3 min chromatography (Column: AD-H; Method Name: AD-H_3UM_4_5_40_4ML_3MIN.M, ee = 100%).

[00185] Basic preparative HPLC method

Mobile phase A: water with 0.05% $\text{NH}_3 \cdot \text{H}_2\text{O}$

Mobile phase B: CH_3CN

Flow rate: 25 mL/min.


Detection: UV 220 nm / 254 nm

Column: Gemini 150*25mm*5um

Column temperature: 30 °C


Time in min	%A	%B
0.00	50	50
10.00	20	80
10.20	0	100
12.00	0	100

[00186] The compound shown below is prepared following analogous procedures:



Example 5

(S)-2-((6-(difluoromethoxy)pyridin-3-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide and (R)-2-((6-(difluoromethoxy)pyridin-3-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide (Cpd Nos Hy2A-25.1 and Hy2A-25.2)



[00187] Step 1

[00188] To a solution of methyl 6-hydroxynicotinate (3.07 g, 20.0 mmol) in anhydrous acetonitrile (310 mL) was treated portionwise with NaH (60% dispersion in mineral oil, 2.16 g, 54.1 mmol) and the mixture was stirred at rt under a nitrogen atmosphere for 30 min. 2, 2-difluoro-2-(fluorosulfonyl) acetic acid (6.07 g, 34.1 mmole) was then added dropwise and the resulting heterogeneous mixture was further stirred at rt under a nitrogen atmosphere for 30 min. Water (25 mL) was added slowly, and acetonitrile was removed under reduced pressure, water (150 mL) and ethyl acetate (150 mL) were added, and the separated layer was further extracted with ethyl acetate (3 × 100 mL). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄, concentrated to dryness under reduced pressure. The residue was

purified by column chromatography on silica gel (eluting with petroleum ether : ethyl acetate = 20: 1 to 10: 1) to afford methyl 6-(difluoromethoxy)nicotinate (3.0 g, 73%) as a yellow solid. ¹H NMR: (CDCl₃ 400 MHz): δ 8.85 (s, 1H), 8.34 (d, *J* = 8.0 Hz, 1H), 7.73-7.37 (m, 1H), 6.96 (d, *J* = 7.6 Hz, 1H), 3.96 (s, 3H).

[00189] Step 2

[00190] To a solution of methyl 6-(difluoromethoxy)nicotinate (1.43 g, 7.0 mmol) in anhydrous toluene (40 mL) was treated dropwise with a solution of 1 M DIBAL-H in toluene (21.2 mL) and the resulting mixture was further stirred at -78 °C under a nitrogen atmosphere for 5 min, and then at 0 °C for 1.5 h. The mixture was treated successively with water (18.3 mL), sodium hydroxide (1 N, 4 mL) and sat. sodium bicarbonate (33 mL). The separated layer was further extracted with ethyl acetate (3 × 10 mL). The combined organic layers were dried over Na₂SO₄, filtered, concentrated to dryness under reduced pressure affording (6-(difluoromethoxy)pyridin-3-yl)methanol (1.0 g, 81%) as a colorless oil ¹H NMR: (CDCl₃ 400 MHz): δ 8.18 (s, 1H), 7.78 (d, *J* = 8.0 Hz, 1H), 7.66-7.37 (m, 1H), 6.92 (d, *J* = 8.0 Hz, 1H), 4.71 (s, 2H).

[00191] Step 3

[00192] To a solution of (6-(difluoromethoxy)pyridin-3-yl)methanol (100.0 mg, 0.56 mmol) in anhydrous dichloromethane (5 mL) was cooled down with an ice-water bath under a nitrogen atmosphere. After stirring for 5 min, PBr₃ (168.0 mg, 0.62 mmol) in anhydrous dichloromethane (0.5 mL) was added dropwise under a nitrogen atmosphere. After addition, the reaction mixture was stirred for 1 h with an ice-water bath. TLC (petroleum ether: ethyl acetate = 3:1) showed the starting material was consumed completely. The reaction mixture was quenched with sat. sodium bicarbonate (10 mL). The aqueous phase was extracted with dichloromethane (3 × 10 mL), washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated by rotary evaporation under reduced pressure to afford 5-(bromomethyl)-2-(difluoromethoxy)pyridine (94 mg, 70%) as a colorless oil.

[00193] Step 4

[00194] To an ice-cooled solution of 2-(3-bromophenyl)ethanamine (30.0 g, 0.15 mol) in pyridine (100 mL) was added isobutyryl chloride (19.2 g, 0.18 mol) dropwise. Then the mixture was stirred at rt for 17 h. The mixture was poured into ice, filtered and dried *in vacuum* to give N-(3-bromophenethyl)isobutyramide (30.0 g, 74%) as a white solid.

¹H NMR (CDCl₃ 400 MHz): δ 7.39-7.33 (m, 2H), 7.18 (t, J = 7.6 Hz, 1H), 7.12 (d, J = 7.6 Hz, 1H), 5.50 (brs, 1H), 3.52-3.45 (m, 2H), 2.80 (t, J = 7.2 Hz, 2H), 2.32-2.28 (m, 1H), 1.12 (d, J = 6.8 Hz, 6H).

[00195] Step 5

[00196] To a solution of N-(3-bromophenethyl)isobutyramide (500 mg, 1.9 mmol) in toluene (5 mL) was added P₂O₅ (657 mg, 4.6 mmol) and POCl₃ (2.7 g, 18 mmol). The mixture was refluxed for 16 h under N₂ atmosphere. TLC (petroleum ether: ethyl acetate = 5:1) showed the starting material was consumed. The mixture was added into aq. NaOH solution (20%, 50 mL) slowly. The resulting mixture was extracted with ethyl acetate (3 × 20 mL). The combined organic layers were washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (eluting with petroleum ether: ethyl acetate = 10:1-3:1) to afford 6-bromo-1-isopropyl-3,4-dihydroisoquinoline (300 mg, 62%) as an oil. **LCMS:** t_R = 1.078 min in 10-80AB_2 min chromatography (Durashell C18, XBrige Shield RP18 2.1*50mm), MS (ESI) m/z 270.0

[M+19]⁺. **¹H NMR** (CD₃OD 400 MHz): δ 7.58 (d, J = 8.4 Hz, 1H), 7.52 (d, J = 8.4 Hz, 1H), 7.48 (s, 1H), 3.60-3.56 (m, 2H), 3.36-3.32 (m, 1H), 2.71-2.67 (m, 2H), 1.20 (d, J = 6.8 Hz, 6H).

[00197] Step 6

[00198] To a solution of 6-bromo-1-isopropyl-3,4-dihydroisoquinoline (1.0 g, 4.0 mmol) in CH₃CN (15 mL) was added HCOOH-Et₃N azeotropic mixture (mole ratio 5:2, 3.5 mL) and RuCl[(R,R)-TsDPEN(P-cymene)] (30 mg, 0.05 mmol). Then the mixture was stirred at rt for 17 h under N₂ atmosphere. TLC (petroleum ether: ethyl acetate = 5:1) showed the starting material was consumed. The mixture was quenched with sat. NaHCO₃ solution (15 mL) and then extracted with ethyl acetate (3 × 20 mL). The combined organic layers were washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford (S)-6-bromo-1-isopropyl-1,2,3,4-tetrahydroisoquinoline (1.0 g, crude, 100%) as a red liquid, which was used in the next step directly. **LCMS:** t_R = 0.650 min in 5-95AB_1.5 min chromatography (MK RP-18e 25-2mm), MS (ESI) m/z 253.9 [M+H]⁺.

[00199] Step 7

[00200] To a solution of (S)-6-bromo-1-isopropyl-1,2,3,4-tetrahydroisoquinoline (4.0 g, 16.0 mmol) in CH₃OH (60 mL) was added Boc₂O (5.2 g, 23.6 mmol), K₂CO₃ (6.4 g, 46.8 mmol) and DMAP (200 mg, 1.6 mmol). Then the mixture was stirred at rt for 16 h. TLC (petroleum ether: ethyl acetate = 10:1) showed the starting material was consumed. The mixture was diluted with H₂O (60 mL) and extracted with ethyl acetate (3 × 60 mL). The combined organic layers were

washed with brine (60 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatograph on silica gel (eluting with petroleum ether: ethyl acetate = 20:1-10:1) to afford (S)-tert-butyl 6-bromo-1-isopropyl-3,4-dihydroisoquinoline-2(1H)-carboxylate (3.4 g, 60%) as an oil.

LCMS: t_R = 1.018 min in 5-95AB_1.5 min chromatography (MK RP-18e 25-2mm), MS (ESI) m/z 297.9 [M-55]⁺. **SFC:** t_R = 8.34 and 10.09 min in 20 min chromatography (AD-H_5_B_05ML), ee = 64.78%.

[00201] Step 8

[00202] A mixture of (S)-tert-butyl 6-bromo-1-isopropyl-3,4-dihydroisoquinoline-2(1H)-carboxylate (150 mg, 0.42 mmol), H₂O (23 mg, 1.26 mmol), Fu's salt (24 mg, 0.08 mmol), Mo(CO)₆ (111 mg, 0.42 mmol), DBU (152 mg, 1.26 mmol) and Pd₂(OAc)₂(P(*o*-tol)₃)₂ (20 mg, 0.02 mmol) in dioxane (2 mL) was heated in microwave at 140 °C for 20 min. The mixture was basified to pH = 10 with aq. NaOH (10%) and extracted with water (3 × 10 mL). The combined aqueous layer was acidified to pH = 2 with HCl (1N) and extracted with CH₂Cl₂ (3 × 15 mL). The combined organic layers were washed with brine (10 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford (S)-2-(tert-butoxycarbonyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxylic acid (135 mg, crude, 100%) as an oil, which was used for the next step directly. **LCMS:** t_R = 0.848 min in 5-95AB_1.5 min chromatography (MK RP-18e 25-2mm), MS (ESI) m/z 263.9 [M-55]⁺. **SFC:** t_R = 3.839 and 4.184 min in 15 min chromatography (IC-3_3_5_40_2, 35mL), ee = 65.46%.

[00203] Step 9

[00204] To a solution of (S)-2-(tert-butoxycarbonyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxylic acid (540 mg, 1.7 mmol) in CH₂Cl₂ (20 mL) was added (4-(ethylsulfonyl)phenyl)methanamine (340 mg, 1.7 mmol), HATU (988 mg, 2.6 mmol) and Et₃N (343 mg, 3.4 mmol). The mixture was stirred at rt overnight. TLC (petroleum ether: ethyl acetate = 1:1) showed the starting material was consumed. The mixture was diluted with water (20 mL) and extracted with CH₂Cl₂ (3 × 20 mL). The combined organic layers were washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC (petroleum ether: ethyl acetate = 1:1-2:1) to give (S)-tert-butyl 6-((4-(ethylsulfonyl)benzyl)carbamoyl)-1-isopropyl-3,4-dihydroisoquinoline-2(1H)-carboxylate (800 mg, 95%) as an oil. **LCMS:** t_R = 1.189 min in 5-95AB_1.5 min chromatography (MK RP-18e 25-2mm), MS (ESI) m/z 445.1 [M-55]⁺.

[00205] Step 10

[00206] To a solution of (S)-tert-butyl 6-((4-(ethylsulfonyl)benzyl)carbamoyl)-1-isopropyl-3,4-dihydroisoquinoline-2(1H)-carboxylate (0.8 g, 1.6 mmol) in dichloromethane (8.0 mL) was added TFA (1.6 mL) at 0 °C. The mixture was stirred at rt overnight. The solution was adjusted to pH = 8 with sat. NaHCO₃ solution. The mixture was extracted with dichloromethane (3 × 20 mL). The combined organic layers were washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford (S)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide (640 mg, 100%) as a yellow oil. **LCMS:** *t_R* = 0.758 min in 5-95AB_1.5 min chromatography (MK RP-18e 25-2mm), MS (ESI) *m/z* 401.1 [M+H]⁺.

[00207] Step 11

[00208] To a solution of (S)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide (50 mg, 0.125 mmol) in CH₃CN (5 mL) was added 5-(bromomethyl)-2-(difluoromethoxy)pyridine (29 mg, 0.125 mmol) and K₂CO₃ (51 mg, 0.375 mmole). The mixture was stirred at rt overnight. The mixture was diluted with ethyl acetate (10 mL) and water (10 mL), extracted with ethyl acetate (3 × 10 mL). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (eluting with petroleum ether : ethyl acetate = 10: 1 to 1: 1) to afford product which was purified by SFC and acidic preparative HPLC separation. After HPLC purification, the eluent was concentrated to remove organic solvent, the residue aqueous solution was lyophilized to give (S)-2-((6-(difluoromethoxy)pyridin-3-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide (**Hy2A-25.1**) (22.40 mg, 74%) and (R)-2-((6-(difluoromethoxy)pyridin-3-yl)methyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide (**Hy2A-25.2**) (2.60 mg, 2%) as white solids.

(**Hy2A-25.1**, 22.40 mg, 74%) as a white solid. **LC-MS:** *t_R* = 0.765 min in 5-95AB_1.50 min chromatography (MK RP-18e 25-2 mm), MS (ESI) *m/z* 558.1 [M+H]⁺. **¹H NMR** (CD₃OD 400 MHz): δ 8.52-8.51 (m, 1H), 8.05-7.91 (m, 1H), 7.90-7.84 (m, 4H), 7.66-7.61 (m, 3H), 7.35-7.33 (m, 1H), 7.14-7.12 (m, 1H), 4.72-4.70 (m, 2H), 4.45-4.14 (m, 2H), 3.93-3.90 (m, 1H), 3.50-3.33 (m, 4H), 3.23-3.19 (m, 2H), 1.30-1.15 (m, 6H), 0.82-0.80 (m, 2H), 0.66-0.64 (m, 2H).

¹⁹F NMR: (CD₃OD 400 MHz): δ -90.92. **SFC:** *t_R* = 3.799 min in 8.0 min chromatography (Column: AS-H_S_5_5_40_3ML_8 MIN_15CM) ee = 100%

[00209] Acid preparative HPLC Method:

Mobile phase A: water with 0.05% HCl solution

Mobile phase B: CH₃CN

Flow rate: 30 mL/min.

Detection: UV 220 nm

Column: Synergi Max-RP 150*30mm*4um

Column temperature: 30 °C

Time in min	%A	%B
0.00	82	18
8.00	52	48
8.20	0	100
10.00	0	100

[00210] (Hy2A-25.2, 2.60 mg, 2%) as a white solid. **LC-MS**: t_R = 0.765 min in 5-95AB_1.50 min chromatography (MK RP-18e 25-2 mm), **MS** (ESI) m/z 558.1 [M+H]⁺. **¹H NMR**: (CD₃OD 400 MHz): δ 8.29 (s, 1H), 8.05-7.91 (m, 1H), 7.89-7.84 (m, 4H), 7.66-7.61 (m, 2H), 7.34 (d, J = 8.0 Hz 1H), 7.13 (d, J = 8.0 Hz 1H), 4.72 (s, 2H), 4.51-4.37 (m, 2H), 4.24-4.17 (m, 1H), 3.90-3.89 (m, 1H), 3.52-3.50 (m, 3H), 3.23-3.19 (m, 2H), 2.24-2.23 (m, 1H), 1.28-1.15 (m, 6H), 0.82-0.80 (m, 3H). **¹⁹F NMR**: (CD₃OD 400 MHz): δ -90.80. **SFC**: t_R = 4.388 min in 8.0 min chromatography (Column: AS-H_S_5_5_40_3ML_8 MIN_15CM) ee = 100%

[00211] Acid preparative HPLC Method:

Mobile phase A: water with 0.05% HCl solution

Mobile phase B: CH₃CN

Flow rate: 30 mL/min.


Detection: UV 220 nm

Column: Synergi Max-RP 150*30mm*4um

Column temperature: 30 °C

Time in min	%A	%B
0.00	82	18
8.00	52	48
8.20	0	100
10.00	0	100


[00212] The following compounds re prepared by analogous procedures



Cpd No	Cy ²	L ²	*
Hy2A-1.1	5-cyano-2-pyrimidinyl	bond	a
Hy2A-1.2	5-cyano-2-pyrimidinyl	bond	a
Hy2A-2	5-(trifluoromethyl)-2-pyrimidinyl	bond	b
Hy2A-3	4-(trifluoromethyl)-2-pyrimidinyl	bond	b
Hy2A-4.1	5-(ethoxycarbonyl)-2-pyrimidinyl	bond	c
Hy2A-4.2	5-(ethoxycarbonyl)-2-pyrimidinyl	bond	c
Hy2A-5	5-(ethoxycarbonyl)-4-(trifluoromethyl)-2-pyrimidinyl	bond	d
Hy2A-5.1	5-(ethoxycarbonyl)-4-(trifluoromethyl)-2-pyrimidinyl	bond	e
Hy2A-5.2	5-(ethoxycarbonyl)-4-(trifluoromethyl)-2-pyrimidinyl	bond	e
Hy2A-12.1	5-methyl-2-pyrimidinyl	CH ₂	f
Hy2A-12.2	5-methyl-2-pyrimidinyl	CH ₂	f
Hy2A-13	4-fluorophenyl	CH ₂	g
Hy2A-14	4-cyanophenyl	CH ₂	h
Hy2A-15.1	4-cyanophenyl	CH ₂	i
Hy2A-15.2	4-cyanophenyl	CH ₂	i
Hy2A-16.1	5-cyano-2-pyridyl	CH ₂	j
Hy2A-16.2	5-cyano-2-pyridyl	CH ₂	j
Hy2A-17.1	6-cyano-3-pyridyl	CH ₂	k
Hy2A-17.2	6-cyano-3-pyridyl	CH ₂	k
Hy2A-18	5-methyl-2-pyrazinyl	CH ₂	l
Hy2A-19	4-chlorophenyl	CH ₂	m
Hy2A-20.1	4-chlorophenyl	CH ₂	n
Hy2A-20.2	4-chlorophenyl	CH ₂	n
Hy2A-21	5-chloro-2-pyridyl	CH ₂	o
Hy2A-24	1-methoxycarbonyl-4-piperidinyl	CH ₂	p
Hy2A-26.1	5-bromo-2-pyrimidinyl	CH ₂	q
Hy2A-26.2	5-bromo-2-pyrimidinyl	CH ₂	q

a, b, c, e, f, i, j, k, n, q = single isomers, separated by chromatography on chiral columns.


d, g, h, l, m, o, p = racemic compound.



Cpd No	Cy ²	L ²	R ^c	*
Hy2A-6	4-fluorophenyl	CH ₂	Me	a
Hy2A-7	4-cyanophenyl	CH ₂	Me	b
Hy2A-8	4-fluorophenyl	CH ₂	Et	c
Hy2A-9	4-chlorophenyl	CH ₂	Et	d
Hy2A-10	4-cyanophenyl	CH ₂	Et	e
Hy2A-11	4-chlorophenyl	CH ₂	Et	f

a, b, c, d, e, f = compounds are racemic.

The following compounds are prepared following procedures analogous to those described above except that Step 8 was replaced by reductive amination with trans-4-(trifluoromethyl)cyclohexane-1-carboxaldehyde mediated by NaCNBH₃ in the presence of HOAc, in MeOH solution at 70 °C for 1 h.




Cpd No ^a	R ⁷	R ⁸	A	*	**
Hy2A-27	H	H	H	(S)	trans
Hy2A-28.1	CH ₂ OH	H	CH	(S)	trans
Hy2A-28.2	H	CH ₂ OH	CH	(S)	trans
Hy2A-28.3	CH ₂ OH	H	CH	(R)	trans
Hy2A-29.1	CH ₂ OH	H	N	(S)	trans
Hy2A-29.2	H	CH ₂ OH	N	(S)	trans
Hy2A-29.3	CH ₂ OH	H	N	(R)	trans

a = Isomers were separated by chromatography on a chiral column; stereochemistry was assigned arbitrarily.

Example 6

(S)-2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide and (R)-2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide (Cpd Nos. Hy2A-23.1 and Hy2A-23.2)



[00213] Step 1

[00214] To a solution of 2-(3-bromophenyl)acetonitrile (5.0 g, 24.38 mmol) in anhydrous THF (60 mL) was cooled down to 0-5 °C with an ice-water bath for 10 min. NaH (60% w/w in mineral oil, 2.6 g, 63.8 mmol) was added in portions and the mixture was stirred for 10 min. MeI (16.6 g, 116.9 mmol) was added dropwise via syringe over 5 min and the reaction was stirred at 0-5 °C with an ice-water bath until the starting material was consumed by TLC (petroleum ether: ethyl acetate = 10: 1, 2 hours). The reaction was quenched with water and partitioned between ethyl acetate (3 × 30 mL). The combined organics were washed with brine (2 × 50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated by rotary evaporation under

reduced pressure to give the residue, which was purified by column chromatography on silica gel (petroleum ether: ethyl acetate = 50:1 - 15:1) to afford the 2-(3-bromophenyl)-2-methylpropanenitrile (4.8 g, 84%) as a flavescent oil. ¹H NMR: (400 MHz): δ 7.61-7.58 (m, 1H), 7.47-7.39 (m, 2H), 7.29-7.24 (m, 1H), 1.71 (s, 6H).

[00215] Step 2

[00216] To a solution of LiAlH₄ (2.7 g, 71.40 mmol) in anhydrous THF (30 mL) was cooled down to 0-5 °C with an ice-water bath under a nitrogen atmosphere for 10 min. The 2-(3-bromophenyl)-2-methylpropanenitrile (4.0 g, 17.85 mmol) in anhydrous THF (20 mL) was added dropwise via syringe over 5 min. After addition, the reaction was stirred at rt for 2 h. The reaction mixture was checked by TLC. TLC (petroleum ether: ethyl acetate = 10:1) showed the reaction was consumed completely. The reaction was cooled down to 0-5 °C with an ice-water bath for 10 min, quenched with water (30 mL). The mixture was filtered through a celite, washed with ethyl acetate (3 × 50 mL). The filtrate was concentrated by rotary evaporation under reduced pressure to give the residue, which was extracted with ethyl acetate (3 × 30 mL). The combined organics were washed with (2 × 50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated by rotary evaporation under reduced pressure to afford the 2-(3-bromophenyl)-2-methylpropan-1-amine (3.2 g, 57%, crude) as a dark oil which was used for the next step directly. **LC-MS:** t_R = 1.207 min in 10-80AB_2.0min chromatography (Welch Xtimate C18, 2.1*30 mm, 3 μ m), MS (ESI) m/z 362.1 [M+Na]⁺.

[00217] Step 3

[00218] To a solution of 2-(3-bromophenyl)-2-methylpropan-1-amine (21.0 g, 92.05 mmol) in pyridine (300 mL) was cooled down to 0-5 °C with an ice-water bath under a nitrogen atmosphere for 15 min. Isobutyryl chloride (11.6 mL, 110.46 mmol, d = 1.017 g/mL) was added dropwise via syringe over 10 min. Then the reaction was stirred at rt overnight. The reaction was cooled down to 0-5 °C with an ice-water bath, quenched with water (200 mL) and diluted with ethyl acetate (800 mL). The organic layer was washed with brine (4 × 300 mL), concentrated by rotary evaporation under reduced pressure to afford the residue which was purified by basic preparative HPLC separation to afford the N-(2-(3-bromophenyl)-2-methylpropyl)isobutyramide (13.5 g, 49%) as a yellow oil.

[00219] Basic preparative HPLC Method:

Mobile phase A: water with 0.05% ammonia hydroxide solution

Mobile phase B: MeCN

Flow rate: 80 mL/min.

Detection: UV 220 nm

Column: Phenomenex Gemini C18 250*250 mm*10um

Column temperature: 30 °C

Time in min	%A	%B
0.00	55	45
25.00	30	70
25.20	0	100
30.00	0	100

¹H NMR: (CDCl₃ 400 MHz): δ 7.51-7.48 (m, 1H), 7.42-7.37 (m, 1H), 7.33-7.28 (m, 1H), 7.27-7.21 (m, 1H), 5.19-5.08 (m, 1H), 3.46 (d, J = 6.0 Hz, 2H), 2.29-2.21 (m, 1H), 1.33 (s, 6H), 1.08 (d, J = 6.8 Hz, 6H).

[00220] Step 4

[00221] To a solution of N-(2-(3-bromophenyl)-2-methylpropyl)isobutyramide (5.0 g, 16.76 mmol) in xylene (50 mL) was added P₂O₅ (5.9 g, 41.90 mmol) with an ice-water bath under a nitrogen atmosphere. After being stirred for 5 min, POCl₃ (27.5 g, 179.35 mmol) was added dropwise via syringe over 2 min. Then the reaction mixture was heated at reflux with a 120-125 °C oil bath overnight. The reaction was cooled down to rt, quenched with 15%wt sodium hydroxide solution (3 × 100 mL), dried over anhydrous Na₂SO₄, filtered and concentrated by rotary evaporation under reduced pressure to afford the residue which was purified by purified by column chromatography on silica gel (petroleum ether: ethyl acetate = 50:1 - 20:1) to afford 6-bromo-1-isopropyl-4,4-dimethyl-3,4-dihydroisoquinoline (1.8 g, 38%) as a yellow oil. **¹H NMR:** (CDCl₃ 400 MHz): δ 7.48 (d, J = 2.0 Hz, 1H), 7.43-7.36 (m, 2H), 3.50 (d, J = 1.2 Hz, 1H), 3.29-3.19 (m, 1H), 1.22-1.16 (m, 12H).

[00222] Step 5

[00223] To a solution of 6-bromo-1-isopropyl-4,4-dimethyl-3,4-dihydroisoquinoline (500 mg, 1.78 mmol) in anhydrous MeOH (8 mL) was added NaBH₄ (136 mg, 3.56 mmol) under a nitrogen atmosphere. After addition, the reaction mixture was stirred at rt for 2 h. Then the reaction mixture was checked by TLC. TLC (petroleum ether: ethyl acetate = 10:1) showed the starting material was consumed completely. Then the reaction was diluted with sat. NH₄Cl solution (20 mL), extracted with ethyl acetate (3 × 10 mL). The combined organics were washed with brine (2 × 20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to afford the 6-bromo-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline (430 mg, 86%) as a white solid which was used for next step directly. **¹H NMR:** (CDCl₃ 400 MHz): δ 7.43 (d, J = 2.0 Hz, 1H),

7.25-7.21 (m, 1H), 7.01 (d, $J = 8.4$ Hz, 1H), 3.85 (d, $J = 4.0$ Hz, 1H), 2.79 (dd, $J = 35.6, 12.4$ Hz, 2H), 2.43-2.27 (m, 1H), 1.28 (s, 3H), 1.21 (s, 3H), 1.13 (d, $J = 7.2$ Hz, 3H), 0.72 (d, $J = 6.8$ Hz, 3H).

[00224] Step 6

[00225] To a solution of 6-bromo-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline (4.0 g, 14.17 mmol) in MeOH (50 mL) was added (Boc)₂O (4.6 g, 21.26 mmol) and K₂CO₃ (4.9 g, 35.43 mmol) under a nitrogen atmosphere. The mixture was stirred at rt overnight. After, the reaction was checked by TLC. TLC (petroleum ether: ethyl acetate = 5:1) showed the reaction was consumed completely. The reaction was diluted with water (80 mL) and extracted with ethyl acetate (3 × 30 mL). The combined organics were washed with brine (2 × 100 mL), dried over anhydrous Na₂SO₄, filtered and concentrated by rotary evaporation under reduced pressure to afford the residue which was purified by purified by column chromatography on silica gel (petroleum ether: ethyl acetate = 50:1) to afford tert-butyl 6-bromo-1-isopropyl-4,4-dimethyl-3,4-dihydroisoquinoline-2(1H)-carboxylate (4.5 g, 38%) as a colorless oil. ¹H NMR: (CDCl₃ 400 MHz): δ 7.46-7.39 (m, 1H), 7.26-7.21 (m, 1H), 7.00-6.95 (m, 1H), 4.92-4.86 (m, 0.5H), 4.79-4.72 (m, 0.5H), 4.11-4.05 (m, 0.5H), 3.90-3.83(m, 0.5H), 3.12-3.04 (m, 0.5H), 2.99-2.92 (m, 0.5H), 2.11-1.96 (m, 1H), 1.49-1.42 (m, 9H), 1.32-1.25 (m, 3H), 1.16 (s, 3H), 1.08-1.03 (m, 3H), 0.94-0.86 (m, 3H).

[00226] Step 7

[00227] To a solution of tert-butyl 6-bromo-1-isopropyl-4,4-dimethyl-3,4-dihydroisoquinoline-2(1H)-carboxylate (400 mg, 1.05 mmol) in 1,4-dioxane (2.0 mL) and H₂O(2.0 mL) was added Fu's salt (61 mg, 0.21 mmol), Mo(CO)₆ (277 mg, 1.05 mmol), DBU (480 mg, 3.15 mmol) and Pd₂(OAc)₂(P(o-tol)₃)₂ (103 mg, 0.11 mmol) under a nitrogen atmosphere. After addition, the reaction mixture was heated at 140 °C in microwave for 20 min. The reaction mixture was cooled down to rt, filtered through celite, washed with ethyl acetate (3 × 20 mL). The combined organics were washed with brine (2 × 30 mL), dried over anhydrous Na₂SO₄, concentrated by rotary evaporation under reduced pressure to afford the residue, which was purified by preparative HPLC (TFA) to afford the 2-(tert-butoxycarbonyl)-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-6-carboxylic acid (300 mg, 60% purity, 49%) as a white solid. LC-MS: $t_R = 1.217$ min in 10-80AB_2.0min chromatography (Welch Xtimate C18, 2.1*30 mm, 3 μ m), MS (ESI) m/z 292.1 [M-55]⁺.

[00228] Step 8

[00229] To a solution of 2-(tert-butoxycarbonyl)-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-6-carboxylic acid (300 mg, 0.518 mmol, 60% purity) in anhydrous DMF (8 ml) was added (4-(ethylsulfonyl)phenyl)methanamine (103 mg, 0.518 mmol), HATU (295 mg, 0.777 mmol) and DIPEA (134 mg, 1.036 mmol) under a nitrogen atmosphere. After addition, the reaction mixture was stirred at rt overnight. The reaction mixture was diluted with ethyl acetate (40 mL). The organic layer was washed with brine (3 × 20 mL), concentrated by rotary evaporation under reduced pressure to afford the residue which was purified by column chromatography on silica gel (petroleum ether: ethyl acetate = 5:1 - 1:2) to afford tert-butyl 6-((4-(ethylsulfonyl)benzyl)carbamoyl)-1-isopropyl-4,4-dimethyl-3,4-dihydroisoquinoline-2(1H)-carboxylate (220 mg, 80%) as a yellow oil. **LC-MS:** t_R = 1.242 min in 10-80AB_2.0min chromatography (Welch Xtimate C18, 2.1*30 mm, 3 μ m), MS (ESI) m/z 529.2 $[M+H]^+$.

[00230] Step 9

[00231] To a solution of tert-butyl 6-((4-(ethylsulfonyl)benzyl)carbamoyl)-1-isopropyl-4,4-dimethyl-3,4-dihydroisoquinoline-2(1H)-carboxylate (220 mg, 0.416 mmol) in CH_2Cl_2 (3 mL) was added TFA (1 mL) under a nitrogen atmosphere. After addition, the reaction was stirred at rt overnight. The reaction mixture was checked by TLC. TLC (petroleum ether: ethyl acetate = 10:1) showed the starting material was consumed completely. The reaction was basified with sat. $NaHCO_3$ solution to pH = 10-11, extracted with ethyl acetate (3 × 10 mL). The combined organic layers were dried over anhydrous Na_2SO_4 , filtered and concentrated by rotary evaporation under reduced pressure to afford the N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide (160 mg, 90%) as a brown solid. **LC-MS:** t_R = 0.816 min in 10-80AB_2.0min chromatography (Welch Xtimate C18, 2.1*30 mm, 3 μ m), MS (ESI) m/z 429.2 $[M+H]^+$.

[00232] Step 10

[00233] To a solution of N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide (60 mg, 0.14 mmol) in CH_3CN (2 mL) was added 4-(bromomethyl)benzonitrile (33 mg, 0.17 mmol) and K_2CO_3 (39 mg, 0.28 mmol) under a nitrogen atmosphere. After addition, the reaction mixture was stirred at rt overnight. The reaction mixture was diluted with ethyl acetate (20 mL) and water (10 mL). The aqueous phase was extracted with ethyl acetate (2 × 10 mL). The combined organics were washed with brine (2 × 20 mL), dried over anhydrous Na_2SO_4 , filtered and the filtrate was concentrated by rotary evaporation under reduced pressure to afford the residue which was purified by column

chromatography on silica gel (petroleum ether: ethyl acetate = 5:1 - 1:1) to afford 2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide (60 mg, 79%) as a colorless solid. **LC-MS:** t_R = 0.769 min in 5-95AB_1.5min chromatography (Welch Xtimate C18, 2.1*30 mm, 3 μ m), MS (ESI) m/z 566.1 $[M+Na]^+$.

[00234] Step 11

[00235] The 2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide (60 mg, 0.11 mmol) was separated with SFC, purified by preparative HPLC (HCl). After preparative HPLC purification, the eluent was concentrated by rotary evaporation under reduced pressure to remove organic solvents. The residual aqueous solution was lyophilized to give the two enantiomers of 2-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-1-isopropyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-6-carboxamide as white solids. (**Hy2A-23.1**, 21.50 mg, 99.79% purity, 36%) as a white solid. **LC-MS:** t_R = 0.976 min in 10-80AB_2.0min chromatography (Welch Xtimate C18, 2.1*30 mm, 3 μ m), MS (ESI) m/z 544.2 $[M+H]^+$. **^1H NMR:** (CD₃OD 400 MHz): δ 8.12 (s, 1H), 7.96-7.59 (m, 9H), 7.27 (d, J = 8.4 Hz, 1H), 4.75-4.18 (m, 4H), 4.08-3.39 (m, 2H), 3.28-2.07 (m, 4H), 1.80-1.40 (m, 6H), 1.27-1.05 (m, 6H), 0.76 (s, 3H). **SFC:** t_R = 1.490 min in 3.0 min chromatography (AD-H_3UM_5_5_40_4mL.ee = 100%).

[00236] HCl preparative HPLC Method:

Mobile phase A: water with 0.05% HCl solution (v/v)

Mobile phase B: MeCN

Flow rate: 30 mL/min.

Detection: UV 220 nm

Column: Synergi Ma-RP C18 150*30*4 μ m

Column temperature: 30 °C

Time in min	%A	%B
0.00	76	24
8.00	46	54
8.20	0	100
10.00	0	100

[00237] (**Hy2A-23.2**, 22.50 mg, 99.83% purity, 38%) as a white solid. **LC-MS:** t_R = 0.976min in 10-80AB_2.0min chromatography (Welch Xtimate C18, 2.1*30 mm, 3 μ m), MS (ESI) m/z 544.2 $[M+H]^+$. **^1H NMR:** (CD₃OD 400 MHz): δ 7.87 (m, 6.5H), 7.64 (d, J = 8.0 Hz,

3.5H), 7.26 (d, $J = 6.0$ Hz, 1H), 4.73 (s, 3H), 3.22 (dd, $J = 14.8, 7.2$ Hz, 3H), 1.50 (s, 3H), 1.39-1.28 (m, 3H), 1.25-1.18 (m, 5H), 0.77 (s, 1H). **SFC:** $t_R = 1.636$ min in 8.0 min chromatography (AD-H_3UM_5_5_40_4mL. ee = 99.38%).

[00238] HCl preparative HPLC Method:

Mobile phase A: water with 0.05% HCl solution (v/v)

Mobile phase B: MeCN

Flow rate: 30 mL/min.


Detection: UV 220 nm

Column: Synergi Ma-RP C18 150*30*4um

Column temperature: 30 °C


Time in min	%A	%B
0.00	76	24
8.00	46	54
8.20	0	100
10.00	0	100

[00239] The following compounds are prepared using analogous procedures:



Example 7

(S)-7-(4-chlorobenzyl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide and (R)-7-(4-chlorobenzyl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide (Cpd Nos Hy2B-8.1 and Hy2B-8.2)



[00240] Step 1

[00241] A mixture of (R)-2-((tert-butoxycarbonyl)amino)-3-methylbutanoic acid (19.6 g, 90.0 mmol) and CDI (15.3 g, 95.6 mmol) in THF (300 mL) was stirred at rt for 1 h. Then potassium 3-methoxy-3-oxopropanoate (15.5 g, 99.0 mmol) and MgCl_2 (9.5 g, 95.0 mmol) were added. After addition, the mixture was stirred at 50 °C for 18 h. TLC (petroleum ether: ethyl acetate = 5:1) showed the starting material was consumed. The mixture was quenched with water (500 mL), extracted with ethyl acetate (3×500 mL). The combined organic layers were washed with sat. aq. NaHCO_3 solution (200 mL), brine (200 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give crude (R)-methyl 4-((tert-butoxycarbonyl) amino)-5-methyl-3-oxohexanoate (25 g, 100%) as a yellow oil.

[00242] Step 2

[00243] To a solution of (R)-methyl 4-((tert-butoxycarbonyl)amino)-5-methyl-3-oxohexanoate (25 g, 91.5 mmol) in anhydrous THF (400 mL) was added KO^tBu (10.8 g, 96.0 mmol) at 0 °C under N₂. After stirred for 45 min, to the mixture was added DABCO (10.8 g, 96.0 mmol) and 2-chloro-1,3-bis(dimethylamino)trimethinium hexafluorophosphate (29.4 g, 96.0 mmol), and then the mixture was stirred at rt for 3 h under N₂. CH₃CO₂NH₄ (21.2 g, 275 mmol) was added to the above solution, and the resulting mixture was stirred at rt overnight. TLC (petroleum ether: ethyl acetate = 5:1) showed the starting material was consumed. The mixture was diluted with ethyl acetate (1000 mL), washed with water (3 × 300 mL), brine (300 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by column chromatograph on silica gel (eluting with petroleum ether: ethyl acetate = 5:1) to give (R)-methyl 2-(1-((tert-butoxycarbonyl)amino)-2-methylpropyl)-5-chloronicotinate (11.4 g, 36%) as a colorless oil. **LC-MS**: *t_R* = 0.913 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) *m/z* 342.9 [M+H]⁺.

[00244] Step 3

[00245] To a solution of (R)-methyl 2-(1-((tert-butoxycarbonyl)amino)-2-methylpropyl)-5-chloronicotinate (11.4 g, 33.3 mmol) in ethanol (250 mL) was added NaBH₄ (2.5 g, 66.6 mmol) and CaCl₂ (3.6 g, 33.3 mmol) at 0 °C under N₂. The mixture was stirred at the same temperature for 2 h. TLC (petroleum ether: ethyl acetate = 5:1) showed the starting material was consumed. The mixture was poured into sat. aq. NH₄Cl solution (150 mL), extracted with ethyl acetate (3 × 200 mL). The combined organic layers were washed with brine (200 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give (R)-tert-butyl (1-(5-chloro-3-(hydroxymethyl)pyridin-2-yl)-2-methylpropyl)carbamate (7.3 g, 70%) as a colorless oil, which was used for the next step directly without further purification.

[00246] Step 4

[00247] To a solution of (R)-tert-butyl (1-(5-chloro-3-(hydroxymethyl)pyridin-2-yl)-2-methylpropyl)carbamate (7.3 g, 22.2 mmol) and Et₃N (11.3 mL, 81.4 mmol, 0.726 g/mL) in CH₂Cl₂ (100 mL) was added MsCl (3.4 g, 29.6 mmol) at 0 °C under N₂. The mixture was stirred at 0 °C for 16 h. The mixture was diluted with CH₂Cl₂ (100 mL), washed with brine (100 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatograph on silica gel (eluting with petroleum ether: ethyl acetate = 5:1) to give (R)-tert-butyl (1-(5-chloro-3-(chloromethyl)pyridin-2-yl)-2-

methylpropyl)carbamate (4.2 g, 58%) as a white solid. **LC-MS**: t_R = 0.940 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 332.9 $[M+H]^+$.

[00248] Step 5

[00249] A mixture of (R)-tert-butyl (1-(5-chloro-3-(chloromethyl)pyridin-2-yl)-2-methylpropyl)carbamate (4.2 g, 12.7 mmol), NaCN (1.8 g, 36.7 mmol), and KI (0.2 g, 1.27 mmol) in DMF (50 mL) and water (5 mL) was stirred at rt overnight. TLC (petroleum ether: ethyl acetate = 5:1) showed the starting material was consumed completely. The mixture was poured into water (300 mL), extracted with ethyl acetate (3 × 300 mL). The combined organic layers were washed with brine (100 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give (R)-tert-butyl (1-(5-chloro-3-(cyanomethyl)pyridin-2-yl)-2-methylpropyl)carbamate (4.0 g, 100%) as a yellow oil. **LC-MS**: t_R = 0.873 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 267.9 $[M-55]^+$.

[00250] Step 6

[00251] To a solution of (R)-tert-butyl (1-(5-chloro-3-(cyanomethyl)pyridin-2-yl)-2-methylpropyl)carbamate (4.0 g, 12.4 mol) in ethanol (50 mL) was added dropwise a solution of NaOH (1.5 g, 10%wt in water, 37.2 mmol). The mixture was stirred at 90 °C for 16 h. The mixture was cooled to rt, poured into water (100 mL) and adjusted to pH= 3 with 1 N HCl solution. The mixture was extracted with ethyl acetate (3 × 50 mL) and the combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give (R)-2-(2-(1-((tert-butoxycarbonyl)amino)-2-methylpropyl)-5-chloropyridin-3-yl)acetic acid (2.6 g, 62%) as a colorless oil. **LC-MS**: t_R = 1.125 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 342.9 $[M+H]^+$.

[00252] Step 7

[00253] A mixture of (R)-2-(2-(1-((tert-butoxycarbonyl)amino)-2-methylpropyl)-5-chloropyridin-3-yl)acetic acid (2.6 g, 7.6 mmol) in HCl/MeOH (100 mL, 4 N) was stirred at rt overnight. The mixture was concentrated under reduced pressure. The residue was diluted with ethyl acetate (200 mL), washed with sat. aq. NaHCO₃ solution (100 mL), brine (100 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (eluting with petroleum ether: ethyl acetate = 1:1) to give (R)-3-chloro-8-isopropyl-7,8-dihydro-1,7-naphthyridin-6(5H)-one (900 mg, 53%) as a colorless oil. **LC-MS**: t_R = 0.677 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 224.9 $[M+H]^+$. **¹H NMR** (CDCl₃ 400 MHz): δ 8.41 (d, J = 2.0 Hz, 1H), 7.44 (d, J =

1.6 Hz, 1H), 4.43 (brs, 1H), 3.76-3.61 (m, 1H), 3.58-3.43 (m, 1H), 2.41-2.27 (m, 1H), 1.12-0.96 (m, 3H), 0.73 (d, $J = 6.8$ Hz, 3H).

[00254] Step 8

[00255] To a solution of (R)-3-chloro-8-isopropyl-7,8-dihydro-1,7-naphthyridin-6(5H)-one (900 mg, 4.0 mmol) in THF (40 mL) was added $\text{BH}_3\text{-Me}_2\text{S}$ (2.5 mL, 20.0 mmol) at 0 °C under N_2 . The mixture was stirred at reflux for 3 h. TLC (petroleum ether: ethyl acetate = 1:1) showed the starting material was consumed. The mixture was cooled to 0 °C, and quenched with *sat.* NH_4Cl solution (50 mL), diluted with water (100 mL), extracted with ethyl acetate (3×100 mL). The combined organic layers were washed with brine (100 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give (R)-3-chloro-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine (1.3 g, 100%) as a yellow oil, which was used for the next step directly without further purification. **LC-MS**: $t_R = 0.790$ min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 210.9 $[\text{M}+\text{H}]^+$.

[00256] Step 9

[00257] A mixture of (R)-3-chloro-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine (1.3 g, 6.2 mol), Boc_2O (3.4 g, 15.5 mmol), and Et_3N (3.1 g, 30.9 mmol) in CH_2Cl_2 (50 mL) was stirred at rt overnight. The mixture was concentrated under reduced pressure. The residue was purified by column chromatograph on silica gel (eluting with petroleum ether: ethyl acetate = 5:1) to give (R)-tert-butyl 3-chloro-8-isopropyl-5,6-dihydro-1,7-naphthyridine-7(8H)-carboxylate (276 mg, 15%) as a colorless oil. **LC-MS**: $t_R = 0.935$ min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 311.0 $[\text{M}+\text{H}]^+$.

[00258] Step 10

[00259] A mixture of (R)-tert-butyl 3-chloro-8-isopropyl-5,6-dihydro-1,7-naphthyridine-7(8H)-carboxylate (315 mg, 1.01 mmol), (4-(ethylsulfonyl)phenyl)methanamine (600 mg, 3.0 mmol), Fu's salt (29 mg, 0.1 mmol), $\text{Mo}(\text{CO})_6$ (264 mg, 1.01 mmol), DBU (456 mg, 3.0 mmol), and $\text{Pd}_2(\text{OAc})_2(\text{P}(\text{o-tol})_3)_2$ (33 mg, 0.035 mmol) in dioxane (5 mL) was stirred at 160 °C for 20 min under N_2 in microwave. The mixture was concentrated under reduced pressure. The residue was purified by preparative TLC (petroleum ether: ethyl acetate = 1:1) to give (R)-tert-butyl 3-((4-(ethylsulfonyl)benzyl)carbamoyl)-8-isopropyl-5,6-dihydro-1,7-naphthyridine-7(8H)-carboxylate (240 mg, 47%) as a colorless oil.

[00260] Step 11

[00261] A mixture of (R)-tert-butyl 3-((4-(ethylsulfonyl)benzyl)carbamoyl)-8-isopropyl-5,6-dihydro-1,7-naphthyridine-7(8H)-carboxylate (240 mg, 0.478 mmol) and TFA (1.0 mL) in

dichloromethane (5 mL) was stirred at 25 °C for 2 h. TLC (petroleum ether: ethyl acetate = 1:2) showed the starting material was consumed completely. The mixture was concentrated under reduced pressure to afford crude (R)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide (200 mg, 100%) as a red oil, which was used for the next step directly without further purification.


[00262] Step 12


[00263] A mixture of (R)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide (50 mg, 0.124 mmol), 1-(bromomethyl)-4-chlorobenzene (25 mg, 0.124 mmol), and K₂CO₃ (51 mg, 0.372 mmol) in CH₃CN (2 mL) was stirred at 25 °C for 3 h. TLC (petroleum ether: ethyl acetate = 1:2) showed the starting material was consumed completely. The mixture was filtered. The filtrate was concentrated under reduced pressure to give the crude product. The residue was purified by preparative TLC (petroleum ether: ethyl acetate = 1: 2) to give about 30 mg of the product. The product was purified by SFC separation and basic preparative HPLC to give the two enantiomers of 7-(4-chlorobenzyl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide as white solids. (**Hy2B-8.1**, 10.2 mg, 20%) as a white solid. **LCMS**: *t_R* = 0.700 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm, MS (ESI) *m/z* 526.1 [M+H]⁺. **¹H NMR** (CDCl₃ 400MHz): δ 8.82 (d, *J* = 2.0 Hz, 1H), 7.89 (m, 3H), 7.57 (d, *J* = 8.0 Hz, 2H), 7.30-7.29 (m, 4H), 6.68-6.65 (m, 1H), 4.80-4.78 (m, 2H), 3.77-3.62 (m, 2H), 3.51 (d, *J* = 6.0 Hz, 1H), 3.22-3.20 (m, 2H), 3.13 (q, *J* = 7.2 Hz, 2H), 2.79-2.78 (m, 1H), 2.77-2.70 (m, 2H), 2.20-2.15 (m, 1H), 1.30 (t, *J* = 7.6 Hz, 3H), 1.06 (d, *J* = 6.8 Hz, 3H), 0.92 (d, *J* = 6.8 Hz, 3H). **SFC**: *t_R* = 0.946 min in 3 min chromatography (AD-H_3UM_3_40_4ML_3MIN.M), ee = 100%.

[00264] (**Hy2B-8.2**, 28.5 mg, 57%) as a white solid. **LCMS**: *t_R* = 0.703 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm, MS (ESI) *m/z* 526.1 [M+H]⁺. **¹H NMR** (CDCl₃ 400 MHz): δ 8.82 (d, *J* = 2.0 Hz, 1H), 7.89 (d, *J* = 8.4 Hz, 3H), 7.57 (d, *J* = 8.4 Hz, 2H), 7.30-7.28 (m, 4H), 6.68-6.66 (m, 1H), 4.80-4.78 (m, 2H), 3.77-3.62 (m, 2H), 3.51 (d, *J* = 6.0 Hz, 1H), 3.22-3.20 (m, 2H), 3.13 (q, *J* = 7.2 Hz, 2H), 2.79-2.78 (m, 1H), 2.77-2.70 (m, 2H), 2.20-2.15 (m, 1H), 1.30 (t, *J* = 7.2 Hz, 3H), 1.06 (d, *J* = 6.8 Hz, 3H), 0.92 (d, *J* = 6.8 Hz, 3H). **SFC**: *t_R* = 1.288 min in 3 min chromatography (AD-H_3UM_3_40_4ML_3MIN.M), ee = 100%.

Example 8

(S)-7-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide and (R)-7-(4-cyanobenzyl)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide (Cpd Nos Hy2B-9.1 and Hy2B-9.2)



**[00265]** Step 1

[00266] A mixture of (R)-2-((tert-butoxycarbonyl)amino)-3-methylbutanoic acid (19.6 g, 90.0 mmol) and CDI (15.3 g, 95.6 mmol) in THF (300 mL) was stirred at rt for 1 h. Then potassium 3-methoxy-3-oxopropanoate (15.5 g, 99.0 mmol) and MgCl₂ (9.5 g, 95.0 mmol) were added. After addition, the mixture was stirred at 50 °C for 18 h. TLC (petroleum ether: ethyl acetate = 5:1) showed the starting material was consumed. The mixture was quenched with water (500 mL), extracted with ethyl acetate (3 × 500 mL). The combined organic layers were washed with sat. aq. NaHCO₃ solution (200 mL), brine (200 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give crude (R)-methyl 4-((tert-butoxycarbonyl) amino)-5-methyl-3-oxohexanoate (25 g, 100%) as a yellow oil.

[00267] Step 2

[00268] To a solution of (R)-methyl 4-((tert-butoxycarbonyl)amino)-5-methyl-3-oxohexanoate (25 g, 91.5 mmol) in THF (400 mL) was added KO^tBu (10.8 g, 96.0 mmol) at 0 °C under N₂. After stirred for 45 min, the mixture was added DABCO (10.8 g, 96.0 mmol) and 2-chloro-1,3-bis(dimethylamino)trimethinium hexafluorophosphate (29.4 g, 96.0 mmol), and then the mixture was stirred at rt for 3 h under N₂. CH₃CO₂NH₄ (21.2 g, 275 mmol) was added to the

above solution, and the resulting mixture was stirred at rt overnight. TLC (petroleum ether: ethyl acetate = 5:1) showed the starting material was consumed. The mixture was diluted with ethyl acetate (1000 mL), washed with water (3×300 mL), brine (300 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by column chromatograph on silica gel (eluting with petroleum ether: ethyl acetate = 5:1) to give (R)-methyl 2-(1-((tert-butoxycarbonyl)amino)-2-methylpropyl)-5-chloronicotinate (11.4 g, 36%) as a colorless oil. **LC-MS:** t_R = 0.913 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 342.9 $[M+H]^+$.

[00269] Step 3

[00270] To a solution of (R)-methyl 2-(1-((tert-butoxycarbonyl)amino)-2-methylpropyl)-5-chloronicotinate (11.4 g, 33.3 mmol) in ethanol (250 mL) was added NaBH_4 (2.5 g, 66.6 mmol) and CaCl_2 (3.6 g, 33.3 mmol) at 0 °C under N_2 . The mixture was stirred at the same temperature for 2 h. TLC (petroleum ether: ethyl acetate = 5:1) showed the starting material was consumed. The mixture was poured into sat. aq. NH_4Cl solution (150 mL), extracted with ethyl acetate (3×200 mL). The combined organic layers were washed with brine (200 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give (R)-tert-butyl (1-(5-chloro-3-(hydroxymethyl)pyridin-2-yl)-2-methylpropyl)carbamate (7.3 g, 70%) as a colorless oil, which was used directly for the next step without further purification.

[00271] Step 4

[00272] To a solution of (R)-tert-butyl (1-(5-chloro-3-(hydroxymethyl)pyridin-2-yl)-2-methylpropyl)carbamate (7.3 g, 22.2 mmol) and Et_3N (11.3 mL, 81.4 mmol, 0.726 g/mL) in CH_2Cl_2 (100 mL) was added MsCl (3.4 g, 29.6 mmol) at 0 °C under N_2 . The mixture was stirred at the same temperature for 16 h. The mixture was diluted with CH_2Cl_2 (100 mL), washed with brine, dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatograph on silica gel (eluting with petroleum ether: ethyl acetate = 5:1) to give (R)-tert-butyl (1-(5-chloro-3-(chloromethyl)pyridin-2-yl)-2-methylpropyl)carbamate (4.2 g, 58%) as a white solid. **LC-MS:** t_R = 0.940 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 332.9 $[M+H]^+$.

[00273] Step 5

[00274] A mixture of (R)-tert-butyl (1-(5-chloro-3-(chloromethyl)pyridin-2-yl)-2-methylpropyl)carbamate (4.2 g, 12.7 mmol), NaCN (1.8 g, 36.7 mmol), and KI (0.2 g, 1.27 mmol) in DMF (50 mL) and water (5 mL) was stirred at rt overnight. TLC (petroleum ether: ethyl acetate = 5:1) showed the starting material was consumed completely. The mixture was

poured into water (300 mL), extracted with ethyl acetate (3×300 mL). The combined organic layers were washed with brine (100 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give (R)-tert-butyl (1-(5-chloro-3-(cyanomethyl)pyridin-2-yl)-2-methylpropyl)carbamate (4.0 g, 100%) as a yellow oil. **LC-MS:** $t_R = 0.873$ min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 267.9 $[M-55]^+$.

[00275] Step 6

[00276] To a solution of (R)-tert-butyl (1-(5-chloro-3-(cyanomethyl)pyridin-2-yl)-2-methylpropyl)carbamate (4.0 g, 12.4 mol) in ethanol (50 mL) was added dropwise a solution of NaOH (1.5 g, 10%wt in water, 37.2 mmol). The mixture was stirred at 90 °C for 16 h. The mixture was cooled to rt, poured into water (100 mL) and adjusted to pH= 3 with 1N HCl solution. The mixture was extracted with ethyl acetate (3×50 mL) and the combined organic layers were washed with brine, dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure to give (R)-2-(2-(1-((tert-butoxycarbonyl)amino)-2-methylpropyl)-5-chloropyridin-3-yl)acetic acid (2.6 g, 62%) as a colorless oil. **LC-MS:** $t_R = 1.125$ min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 342.9 $[M+H]^+$.

[00277] Step 7

[00278] A mixture of (R)-2-(2-(1-((tert-butoxycarbonyl)amino)-2-methylpropyl)-5-chloropyridin-3-yl)acetic acid (2.6 g, 7.6 mmol) in HCl/MeOH (100 mL, 4N) was stirred at rt overnight. The mixture was concentrated under reduced pressure. The residue was diluted with ethyl acetate (200 mL), washed with sat. aq. $NaHCO_3$ solution (100 mL), brine (100 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (petroleum ether: ethyl acetate = 1:1) to give (R)-3-chloro-8-isopropyl-7,8-dihydro-1,7-naphthyridin-6(5H)-one (900 mg, 53%) as a colorless oil. **LC-MS:** $t_R = 0.677$ min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 224.9 $[M+H]^+$. **^1H NMR** ($CDCl_3$ 400 MHz): δ 8.41 (d, $J = 2.0$ Hz, 1H), 7.44 (d, $J = 1.5$ Hz, 1H), 4.43 (brs, 1H), 3.76-3.61 (m, 1H), 3.58-3.43 (m, 1H), 2.41-2.27 (m, 1H), 1.12-0.96 (m, 3H), 0.73 (d, $J = 6.8$ Hz, 3H).

[00279] Step 8

[00280] To a solution of (R)-3-chloro-8-isopropyl-7,8-dihydro-1,7-naphthyridin-6(5H)-one (900 mg, 4.0 mmol) in THF (40 mL) was added $BH_3 \cdot Me_2S$ (2.5 mL, 20.0 mmol) at 0 °C under N_2 . The mixture was stirred at 80 °C for 3 h. TLC (petroleum ether: ethyl acetate = 1:1) showed the starting material was consumed. The mixture was cooled to 0 °C, and quenched with sat. NH_4Cl solution (50 mL), diluted with water (100 mL), extracted with ethyl acetate (3×100

mL). The combined organic layers were washed with brine (100 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give (R)-3-chloro-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine (1.3 g, 100%) as a yellow oil, which was used directly for the next step without further purification. **LC-MS:** t_R = 0.790 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 210.9 $[M+H]^+$.

[00281] Step 9

[00282] A mixture of (R)-3-chloro-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine (1.3 g, 6.2 mol), Boc_2O (3.4 g, 15.5 mmol), and Et_3N (3.1 g, 30.9 mmol) in CH_2Cl_2 (50 mL) was stirred at rt overnight. The mixture was concentrated under reduced pressure. The residue was purified by column chromatograph on silica gel (eluting with petroleum ether: ethyl acetate = 5:1) to give (R)-tert-butyl 3-chloro-8-isopropyl-5,6-dihydro-1,7-naphthyridine-7(8H)-carboxylate (276 mg, 15%) as a colorless oil. **LC-MS:** t_R = 0.935 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 311.0 $[M+H]^+$.

[00283] Step 10

[00284] A mixture of (R)-tert-butyl 3-chloro-8-isopropyl-5,6-dihydro-1,7-naphthyridine - 7(8H)-carboxylate (200 mg, 0.65 mmol), (4-(ethylsulfonyl)phenyl)methanamine (386 mg, 1.94 mmol), Fu's salt (18.8 mg, 0.065 mmol), $Mo(CO)_6$ (170 mg, 0.65 mmol), DBU (294 mg, 1.94 mmol), and $Pd_2(OAc)_2(P(o-tol)_3)_2$ (21.2 mg, 0.0225 mmol) in dioxane (5 mL) was stirred at 160 °C for 20 min under N_2 in microwave. The mixture was concentrated under reduced pressure. The residue was purified by preparative TLC (eluting with petroleum ether: ethyl acetate = 1:1), SFC separation to give (S)-tert-butyl 3-((4-(ethylsulfonyl)benzyl)carbamoyl)-8-isopropyl-5,6-dihydro-1,7-naphthyridine-7(8H)-carboxylate (38 mg, 58%) and (R)-tert-butyl 3-((4-(ethylsulfonyl)benzyl)carbamoyl)-8-isopropyl-5,6-dihydro-1,7-naphthyridine-7(8H)-carboxylate (48 mg, 73%) as a colorless oil. **LC-MS:** t_R = 0.835 min in 5-95AB_1.5min chromatography (MK RP18e 25-2mm), MS (ESI) m/z 502.0 $[M+H]^+$.

Before SFC separation

Isomer SFC t_R = 6.334, 6.511 in 12 min chromatography (Column: OJ-H, Method Name: OJ-H_5_5_40_2, 35ML.M, ee = 0.604%).

SFC separation condition:

Instrument: Thar 80

Column: OJ 250mm*30mm, 5um

Mobile phase: A: Supercritical CO_2 , B: EtOH (0.05% NH_3H_2O), A:B =80:20 at 60 ml/min

Column Temp: 38 °C

Nozzle Pressure: 100Bar

Nozzle Temp: 60 °C

Evaporator Temp: 20 °C

Trimmer Temp: 25 °C

Wavelength: 220nm

[00285] Step 11

[00286] A mixture of (S)-tert-butyl 3-((4-(ethylsulfonyl)benzyl)carbamoyl)-8-isopropyl-5,6-dihydro-1,7-naphthyridine-7(8H)-carboxylate (38 mg, 0.076 mmol) and TFA (1.0 mL) in dichloromethane (5 mL) was stirred at rt for 2 h. TLC (petroleum ether: ethyl acetate = 1:2) showed the starting material was consumed completely. The mixture was concentrated under reduced pressure to afford crude (S)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide (crude, 100%) as a red-brown oil, which was used directly for the next step without further purification.

[00287] Step 12 (**Hy2B-9.1**)

[00288] A mixture of (S)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide (20 mg, 0.05 mmol), *trans*-4-(trifluoromethyl)cyclohexanecarbaldehyde (27 mg, 0.15 mmol), HOAc (0.23 mg) and NaBH₃CN (7.5 mg, 0.15 mmol) in MeOH (1.5 mL) was stirred at 70 °C for 2 h. TLC (petroleum ether: ethyl acetate = 1:1) showed the starting material was consumed completely. The mixture was concentrated under reduced pressure and diluted with H₂O (15 mL) and extracted with EtOAc (3 × 40 mL). The combined organic layer were washed with brine (40 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to give the crude product. The crude product was purified by preparative TLC (CH₂Cl₂: MeOH = 20:1) and acid preparative HPLC to give (S)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-7-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide (**Hy2B-9.1**, 6.8 mg, 34%) as a white solid. **LCMS:** *t_R* = 0.935 min in 10-80AB_2.0min chromatography (A: Xtimate C18, 2.1*30mm, 3um), MS (ESI) *m/z* 566.2 [M+H]⁺. **¹H NMR** (CD₃OD 400 MHz): δ 9.00 (s, 1H), 8.21 (s, 1H), 7.90 (d, *J* = 8.4 Hz, 2H), 7.66 (d, *J* = 8.4 Hz, 2H), 4.73 (s, 2H), 4.42-4.40 (m, 1H), 3.95-3.90 (m, 1H), 3.68-3.58 (m, 1H), 3.23-3.19 (m, 2H), 3.14-3.12 (m, 2H), 2.49-2.35 (m, 1H), 2.06-1.91 (m, 6H), 1.48-1.34 (m, 6H), 1.24-1.20 (m, 6H), 0.92-0.89 (m, 3H). **¹⁹F NMR** (CD₃OD 400 MHz): δ -75.40. **Isomer SFC:** *t_R* = 1.636 min in 3 min chromatography (AD-H_3UM_3_5_40_4ML_3MIN.M), ee = 100%.


[00289] Step 13

[00290] A mixture of (R)-tert-butyl 3-((4-(ethylsulfonyl)benzyl)carbamoyl)-8-isopropyl-5,6-dihydro-1,7-naphthyridine-7(8H)-carboxylate (48 mg, 0.096 mmol) and TFA (1.0 mL) in dichloromethane (5 mL) was stirred at rt for 2 h. TLC (petroleum ether: ethyl acetate = 1:2) showed the starting material was consumed completely. The mixture was concentrated under reduced pressure to afford crude (R)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide (crude, 100%) as a red-brown oil, which was used directly for the next step without further purification.

[00291] Step 14 (**Hy2B-9.2**)


[00292] A mixture of (R)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide (12 mg, 0.03 mmol), *trans*-4-(trifluoromethyl)cyclohexanecarbaldehyde (16.2 mg, 0.09 mmol), HOAc (0.18 mg) and NaBH₃CN (5.6 mg, 0.09 mmol) in MeOH (1 mL) was stirred at 70 °C for 2 h. TLC (petroleum ether: ethyl acetate = 1:1) showed the starting material was consumed completely. The mixture was concentrated under reduced pressure, diluted with H₂O (15 mL), extracted with EtOAc (3 x 40 mL). The combined organic layers were washed with brine (40 mL), dried, filtered and concentrated to give the crude product. The crude product was purified by preparative TLC (DCM: MeOH = 20:1) and acid preparative HPLC to give (R)-N-(4-(ethylsulfonyl)benzyl)-8-isopropyl-7-((*trans*-4-(trifluoromethyl)cyclohexyl) methyl)-5,6,7,8-tetrahydro-1,7-naphthyridine-3-carboxamide (**Hy2B-9.2**, 4 mg, 80%) as a white solid. **LCMS**: *t_R* = 0.929 min in 10-80AB_2.0min chromatography (A: Xtimate C18, 2.1*30mm, 3um), MS (ESI) *m/z* 566.2 [M+H]⁺. **¹H NMR** (CD₃OD 400 MHz): δ 9.41-9.38 (m, 1H), 9.00 (s, 1H), 8.20 (s, 1H), 7.91 (d, *J* = 8.4 Hz, 2H), 7.66 (d, *J* = 8.4 Hz, 2H), 4.84 (s, 2H), 4.42-4.40 (m, 1H), 3.97-3.89 (m, 1H), 3.59-3.55 (m, 1H), 3.25-3.20 (m, 2H), 3.14-3.12 (m, 2H), 2.49-2.45 (m, 1H), 2.07-1.90 (m, 6H), 1.49-1.34 (m, 6H), 1.24-1.14 (m, 6H), 0.92-0.90 (m, 3H). **¹⁹F NMR** (CD₃OD 400 MHz): δ -75.417. **Isomer SFC**: *t_R* = 1.634 min in 3 min chromatography (AD-H_3UM_3_5_40_4ML_3MIN.M), ee = 100%.

[00293] The following compounds are prepared using procedures analogous to those described in Examples 7 and 8.




Cpd No	Cy ²	L ²	*
Hy2B-1.1	5-cyano-2-pyrimidinyl	bond	a
Hy2B-1.2	5-cyano-2-pyrimidinyl	bond	a
Hy2B-2.1	5-chloro-2-pyrimidinyl	bond	b
Hy2B-2.2	5-chloro-2-pyrimidinyl	bond	b
Hy2B-3.1	5-(trifluoromethyl)-2-pyrimidinyl	bond	c
Hy2B-3.2	5-(trifluoromethyl)-2-pyrimidinyl	bond	c
Hy2B-4.1	5-(ethoxycarbonyl)-2-pyrimidinyl	bond	d
Hy2B-4.2	5-(ethoxycarbonyl)-2-pyrimidinyl	bond	d
Hy2B-5.1	5-(ethoxycarbonyl)-4-(trifluoromethyl)-2-pyrimidinyl	bond	e
Hy2B-5.2	5-(ethoxycarbonyl)-4-(trifluoromethyl)-2-pyrimidinyl	bond	e
Hy2B-6.1	4-fluorophenyl	CH ₂	f
Hy2B-6.2	4-fluorophenyl	CH ₂	f
Hy2B-7.1	4-cyanophenyl	CH ₂	g
Hy2B-7.2	4-cyanophenyl	CH ₂	g
Hy2B-10.1	4-chlorophenyl	CH ₂	h
Hy2B-10.2	4-chlorophenyl	CH ₂	h

a, b, c, d, e, f, g, h = single isomers, separated by chromatography on chiral columns.



a Isomers were separated by chromatography on a chiral column.

Example 9

3-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole-6-carboxamide (Cpd No Hy3A-1)

[00294] Step 1. (*trans*-4-(trifluoromethyl)cyclohexyl)methanol

[00295] To a solution of *trans*-4-(trifluoromethyl)cyclohexanecarboxylic acid (5.2074 g, 26.5 mmol) in THF (100 mL) was added 20 mL (40 mmol) of 2.0 M LiAlH₄ in THF at 0 °C (ice bath). The resulting reaction mixture was allowed to stir at rt for 20 h and then carefully quenched with 30 g of sodium sulfate decahydrate (Glauber's salt). After vigorously stirred at rt for 2 h, the solid was filtered and washed with DCM. The combined filtrate was evaporated under reduced pressure to afford (*trans*-4-(trifluoromethyl)cyclohexyl)methanol as a colorless liquid, which was used in the next step without further purification. ¹H NMR (CDCl₃, 400 MHz) δ 3.47 (d, *J* = 6.15 Hz, 2H), 2.02-1.89 (m, 5H), 1.57 (s, 1H), 1.52-1.44 (m, 1H), 1.38-1.27 (m, 2H), 1.04-0.94 (m, 2H); ¹⁹F NMR (CDCl₃, 376 MHz) δ -73.82 (d, *J* = 7.89 Hz).

[00296] Step 2. (*trans*-4-(trifluoromethyl)cyclohexyl)methyl 4-methylbenzenesulfonate

[00297] To a solution of (*trans*-4-(trifluoromethyl)cyclohexyl)methanol, obtained as described above, in THF (50 mL) was added 1.1350 g (28.4 mmol) of 60% NaH at 0 °C (ice bath). The resulting reaction mixture was allowed to stir at rt for 0.5 h and then 5.4140 g (28.4 mmol) of TsCl was added in portion over 10 min. The reaction mixture was stirred at rt for 17 h and then quenched with saturated NaHCO₃, extracted twice with ethyl acetate, and dried over Na₂SO₄. After the solvent was evaporated under reduced pressure, the residue was purified by chromatography on silica gel (120 g column eluted with 0→5% MeOH/DCM over 30 min) to give 5.7950 g (65% in two steps) of (*trans*-4-(trifluoromethyl)cyclohexyl)methyl 4-methylbenzenesulfonate as a solid. LC-MS t_R = 1.85 min in 2.5 min chromatography, m/z 165 (M^+ -OTs); ¹H NMR (CDCl₃, 400 MHz) δ 7.78 (d, J = 8.49 Hz, 2H), 7.35 (d, J = 8.49 Hz, 2H), 3.84 (d, J = 6.15 Hz, 2H), 2.46 (s, 3H), 1.98-1.90 (m, 3H), 1.86-1.82 (m, 2H), 1.69-1.62 (m, 1H), 1.33-1.23 (m, 2H), 1.03-0.93 (m, 2H); ¹⁹F NMR (CDCl₃, 376 MHz) δ -73.86 (d, J = 7.89 Hz).

[00298] Step 3. (4-bromo-2-fluorophenyl)(cyclopropyl)methanone and (4-bromo-2-fluorophenyl)(cyclopropyl)methanol

[00299] To a solution of 4-bromo-2-fluorobenzaldehyde (7.0400 g, 34.7 mmol) in THF (50 mL) was added 84 mL of 0.5 M cyclopropylmagnesium bromide in THF at -78 °C under nitrogen. The reaction mixture was allowed to slowly warm to 12 °C over 17 h and then quenched with saturated NH₄Cl, extracted twice with ethyl acetate, and dried over Na₂SO₄. After the solvent was evaporated under reduced pressure, the residue was purified by chromatography on silica gel (120 g column eluted with 0→60% ethyl acetate/hexanes over 40 min) to afford 3.0050 g (36%) of (4-bromo-2-fluorophenyl)(cyclopropyl)methanone, LC-MS t_R = 1.68 min in 2.5 min chromatography, m/z 243, 245 (MH^+); ¹H NMR (CDCl₃, 400 MHz) δ 7.65 (t, J = 8.20 Hz, 1H), 7.39-7.34 (m, 2H), 2.64-2.57 (m, 1H), 1.30-1.26 (m, 2H), 1.11-1.06 (m, 2H); ¹⁹F NMR (CDCl₃, 376 MHz) δ -108.96 (m) and 0.7856 g (9%) of (4-bromo-2-fluorophenyl)(cyclopropyl)methanol, LC-MS t_R = 1.51 min in 2.5 min chromatography, m/z 227, 229 (M^+ -OH); ¹H NMR (CDCl₃, 400 MHz) δ 7.45-7.41 (m, 1H), 7.30 (d, J = 8.20 Hz, 1H), 7.22 (d, J = 9.67 Hz, 1H), 4.33 (d, J = 8.20 Hz, 1H), 2.09 (m, 1H), 1.22-1.19 (m, 1H), 0.64-0.63 (m, 1H), 0.54-0.43 (m, 2H); ¹⁹F NMR (CDCl₃, 376 MHz) δ -116.28 (m).

[00300] Step 4. 6-bromo-3-cyclopropyl-2H-indazole

[00301] A mixture of (4-bromo-2-fluorophenyl)(cyclopropyl)methanone (0.5216 g, 2.15 mmol) and hydrazine hydrate (1.5360 g) in dioxane (3 mL) was heated in microwave at 160 °C for 2 h. The reaction mixture was quenched with saturated NH₄Cl, extracted twice with ethyl acetate, and dried over Na₂SO₄. After the solvent was evaporated under reduced pressure, the

crude product was used in the next step without further purification. LC-MS t_R = 1.53 min in 2.5 min chromatography, m/z 237, 239 (MH^+); ^1H NMR ($CDCl_3$, 400 MHz) δ 7.63-7.59 (m, 2H), 7.26-7.22 (m, 1H), 2.24-2.18 (m, 1H), 1.08-1.05 (m, 4H).

[00302] Step 5. 6-bromo-3-cyclopropyl-1-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indazole and 6-bromo-3-cyclopropyl-2-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole

[00303] A mixture of 6-bromo-3-cyclopropyl-2H-indazole, obtained as described above, (*trans*-4-(trifluoromethyl)cyclohexyl)methyl 4-methylbenzenesulfonate (0.8160 g, 2.42 mmol), and Cs_2CO_3 (3.1140 g) in DMF (10 mL) was heated at 80 °C for 15 h. The reaction mixture was cooled to rt and then quenched with saturated NH_4Cl , extracted twice with ethyl acetate, and dried over Na_2SO_4 . After the solvent was evaporated under reduced pressure, the residue was purified by chromatography on silica gel (40 g column eluted with 0→50% ethyl acetate/hexanes over 30 min) to afford 0.4044 g (47%) of 6-bromo-3-cyclopropyl-1-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indazole, LC-MS t_R = 2.16 min in 2.5 min chromatography, m/z 401, 403 (MH^+); ^1H NMR ($CDCl_3$, 400 MHz) δ 7.57 (d, J = 8.50 Hz, 1H), 7.46 (s, 1H), 7.20-7.17 (m, 1H), 4.06 (d, J = 7.03 Hz, 2H), 2.18-2.13 (m, 1H), 2.00-1.92 (m, 4H), 1.71 (d, J = 12.89 Hz, 2H), 1.32-1.22 (m, 2H), 1.11-1.01 (m, 6H); ^{19}F NMR ($CDCl_3$, 376 MHz) δ -73.80 (d, J = 7.89 Hz) and 0.0852 g (10%) of 6-bromo-3-cyclopropyl-2-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole, LC-MS t_R = 2.05 min in 2.5 min chromatography, m/z 401, 403 (MH^+); ^1H NMR ($CDCl_3$, 400 MHz) δ 7.79 (s, 1H), 7.53 (d, J = 8.78 Hz, 1H), 7.07 (d, J = 8.78 Hz, 1H), 4.33 (d, J = 7.32 Hz, 2H), 2.19-2.13 (m, 1H), 2.04-1.94 (m, 4H), 1.78 (d, J = 12.89 Hz, 2H), 1.37-1.26 (m, 2H), 1.19-1.08 (m, 4H), 0.97-0.93 (m, 2H); ^{19}F NMR ($CDCl_3$, 376 MHz) δ -73.80 (d, J = 9.21 Hz).

[00304] Step 6. 3-cyclopropyl-2-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole-6-carboxylic acid





[00305] A mixture of 6-bromo-3-cyclopropyl-2-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole (0.0852 g, 0.21 mmol), Molybdenumhexacarbonyl (0.1190 g), Herrmann's palladacycle (0.0579 g), Tri-*tert*-butylphosphonium tetrafluoroborate (0.0573 g), water (0.1 mL), and DBU (0.1 mL) in dioxane (2 mL) was heated in microwave at 150 °C for 20 min. The reaction mixture was purified by reverse-phase HPLC (Phenomenex® Luna 5 μ C₁₈ (2) 250 × 21.20 mm column, 10% CH_3CN/H_2O , 0.1% CF_3COOH over 1.5 min, 10% → 90% CH_3CN/H_2O , 0.1% CF_3COOH over 8 min, and then 90% CH_3CN/H_2O , 0.1% CF_3COOH over 4 min, flow rate 25 mL/min) to yield


0.0624 g (80%) of 3-cyclopropyl-2-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole-6-carboxylic acid. LC-MS t_R = 1.70 min in 2.5 min chromatography, m/z 367 (MH^+).

[00306] Step 7. 3-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole-6-carboxamide

[00307] A mixture of 3-cyclopropyl-2-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole-6-carboxylic acid (0.0208 g, 0.0567 mmol), HATU (0.0359 g), TEA (0.1 mL), and (R)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethan-1-ol (0.0176 g, 0.0767 mmol) in DMF (2 mL) was stirred at rt for 2 h. The reaction mixture was purified by reverse-phase HPLC (Phenomenex® Luna 5 μ C₁₈ (2) 250 $\times$ 21.20 mm column, 10% CH₃CN/H₂O, 0.1% CF₃COOH over 1.5 min, 10% $\rightarrow$ 90% CH₃CN/H₂O, 0.1% CF₃COOH over 8 min, and then 90% CH₃CN/H₂O, 0.1% CF₃COOH over 4 min, flow rate 25 mL/min) and the fractions were then lyophilized to afford 3-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-2H-indazole-6-carboxamide as a white fluffy solid. LC-MS t_R = 1.59 min in 2.5 min chromatography, m/z 578 (MH^+); ¹H NMR (DMSO-*d*₆, 400 MHz) δ 8.82 (d, J = 8.20 Hz, 1H), 8.21 (s, 1H), 7.84 (d, J = 8.20 Hz, 2H), 7.70-7.66 (m, 3H), 7.41 (d, J = 9.08 Hz, 1H), 5.16-5.14 (m, 1H), 4.38 (d, J = 7.33 Hz, 2H), 3.78-3.67 (m, 2H), 3.26 (q, J = 7.32 Hz, 2H), 2.20-2.09 (m, 3H), 1.85 (m, 2H), 1.66 (m, 2H), 1.24-1.14 (m, 6H), 1.09 (t, J = 7.32 Hz, 3H), 0.97-0.96 (m, 2H); ¹⁹F NMR (DMSO-*d*₆, 376 MHz) δ -72.25 (d, J = 9.21 Hz).


[00308] The following compounds are prepared using analogous procedures:

	
Cpd No Hy3A-2	Cpd No Hy3A-3
	
Cpd No Hy3A-4	Cpd No Hy3A-5






Example 9

3-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1R,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-2H-pyrazolo[4,3-b]pyridine-6-carboxamide (Cpd No Hy3B-1)




[00309] Compound **Hy3B-1** was prepared as shown in the Scheme above. (5-Bromo-3-fluoropyridin-2-yl)(cyclopropyl)methanone was prepared by addition of cyclopropylmagnesium bromide to 5-bromo-3-fluoropicolinonitrile. (5-Bromo-3-fluoropyridin-2-yl)(cyclopropyl)methanone was converted to Compound **Hy3B-1** using procedures analogous to those described in Example 8 Steps 4-7.

[00310] The following compounds are prepared using analogous procedures:

	
Cpd No Hy3B-2	Cpd No Hy3B-3
	
Cpd No Hy3B-4	

Example 10

N-(4-(ethylsulfonyl)benzyl)-1-methyl-7-(((1*r*,4*r*)-4-(trifluoromethyl)cyclohexyl)methoxy)-1*H*-pyrrolo[2,3-*c*]pyridine-2-carboxamide (Cpd No Hy4-1)



[00311] Step 1

[00312] To a mixture of 7-chloro-1*H*-pyrrolo[2,3-*c*]pyridine-2-carboxylic acid (200 mg, 1.02 mmol) in anhydrous MeOH (5 mL) was added dropwise SOCl₂ (2 mL). The mixture was stirred at 60 °C for 6 h under N₂. LCMS showed that the reaction was completed. The mixture was

concentrated under reduced pressure and dry-freezing directly to give crude methyl 7-chloro-1H-pyrrolo[2,3-c]pyridine-2-carboxylate (214 mg, 100%) as a white solid, which was used for the next step directly without further purification. **LC-MS** t_R = 0.675 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 210.9 $[M+H]^+$.

[00313] Step 2

[00314] To a mixture of crude methyl 7-chloro-1H-pyrrolo[2,3-c]pyridine-2-carboxylate (90 mg, 0.429 mmol) in anhydrous DMF (3 mL) was added NaH (34 mg, 0.857 mmol, 60% in mineral oil). The mixture was stirred at rt for 0.5 h under N_2 . MeI (183 mg, 1.29 mmol) was added to the mixture and the mixture was stirred at rt for 3 h under N_2 . TLC (petroleum ether / ethyl acetate = 4/1) showed that the starting material was consumed completely. The mixture was adjusted to pH = 4-5 with 1 M HCl solution. The mixture was concentrated under reduced pressure and dry-freezing directly to give crude 7-chloro-1-methyl-1H-pyrrolo[2,3-c]pyridine-2-carboxylic acid (150 mg, >100%, contain some salts) as a yellow solid, which was used for the next step directly without further purification. **LC-MS** t_R = 0.551 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 210.9 $[M+H]^+$.

[00315] Step 3

[00316] To a mixture of crude 7-chloro-1-methyl-1H-pyrrolo[2,3-c]pyridine-2-carboxylic acid (crude 150 mg, 0.429 mmol) in anhydrous MeOH (3 mL) was added dropwise $SOCl_2$ (0.8 mL). The mixture was stirred at 50 °C for 1 h under N_2 . LCMS showed that the reaction was completed. The mixture was concentrated under reduced pressure. The residue was diluted with H_2O (20 mL) and adjusted to pH = 7-8 with *aq.* $NaHCO_3$ solution. The aqueous layer was extracted with ethyl acetate (3 X 20 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether / ethyl acetate = 1/0-5/1 to give methyl 7-chloro-1-methyl-1H-pyrrolo[2,3-c]pyridine-2-carboxylate (50 mg, 52%) as a white solid. **LC-MS** t_R = 0.758 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 225.0 $[M+H]^+$.

[00317] Step 4

[00318] To a mixture of methyl 7-chloro-1-methyl-1H-pyrrolo[2,3-c]pyridine-2-carboxylate (20 mg, 0.0893 mmol) in dry toluene (2 mL) was added (*trans*-4-(trifluoromethyl)cyclohexyl)methanol (32 mg, 0.1786 mmol), $Pd_2(dba)_3$ (20 mg, 0.0219 mmol), BINAP (20 mg, 0.032 mmol) and Cs_2CO_3 (60 mg, 0.1786 mmol). The mixture was stirred at 110 °C for 7 h under N_2 . LCMS showed that the reaction was completed. The mixture was

quenched with water (15 mL) and extracted with ethyl acetate (3 X 15 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC with petroleum ether / ethyl acetate = 7/1 to afford methyl 1-methyl-7-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)-1H-pyrrolo[2,3-*c*]pyridine-2-carboxylate (33 mg, 100%) as a yellow solid. **LC-MS** t_R = 1.308 min in 5-95AB_1.5 min chromatography (Welch Xtimate C18, 2.1*30mm), MS (ESI) m/z 371.1 $[M+H]^+$.

[00319] Step 5

[00320] To a mixture of methyl 1-methyl-7-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)-1H-pyrrolo[2,3-*c*]pyridine-2-carboxylate (17 mg, 0.0447 mmol) in MeOH/THF/H₂O (3 mL/1 mL/0.8 mL) was added NaOH (36 mg, 0.893 mmol). The mixture was stirred at 45 °C for 2 h under N₂. LCMS showed that the reaction was completed. The mixture was concentrated under reduced pressure. The residue was diluted with H₂O (10 mL) and adjusted to pH = 5-6 with 1 M HCl solution. The aqueous layer was extracted with ethyl acetate (3 X 10 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to afford crude 1-methyl-7-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)-1H-pyrrolo[2,3-*c*]pyridine-2-carboxylic acid (16 mg, 100%) as a yellow solid, which was used for the next step directly without further purification. **LC-MS** t_R = 0.836 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 357.1 $[M+H]^+$.

[00321] Step 6

[00322] To a mixture of crude 1-methyl-7-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)-1H-pyrrolo[2,3-*c*]pyridine-2-carboxylic acid (16 mg, 0.0447 mmol) in CH₂Cl₂ (1 mL) was added (4-(ethylsulfonyl)phenyl)methanamine HCl salt (18 mg, 0.0894 mmol), HATU (34 mg, 0.0894 mmol) and Et₃N (9 mg, 0.0894 mmol). The mixture was stirred at rt for 2 h under N₂. LCMS showed that the reaction was completed. The mixture was diluted with water (10 mL) and extracted with DCM (3 X 10 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC with petroleum ether / ethyl acetate = 2/3 and basic preparative HPLC separation, then dry-freezing directly to afford N-(4-(ethylsulfonyl)benzyl)-1-methyl-7-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)-1H-pyrrolo[2,3-*c*]pyridine-2-carboxamide (**Hy4-1**, 8.10 mg, 34%) as a white solid. **LC-MS** t_R = 0.778 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 538.1 $[M+H]^+$. **¹H NMR** (CDCl₃ 400 MHz): δ 7.88 (d, J = 8.0 Hz, 2H), 7.70 (d, J = 6.0 Hz, 1H), 7.54 (d, J = 8.0 Hz, 2H), 7.07 (d, J = 5.6 Hz, 1H), 6.79

(s, 1H), 6.72-6.65 (m, 1H), 4.73 (d, $J = 6.0$ Hz, 2H), 4.34 (s, 3H), 4.32 (d, $J = 6.4$ Hz, 2H), 3.10 (q, $J = 7.6$ Hz, 2H), 2.14-1.85 (m, 6H), 1.49-1.35 (m, 2H), 1.27 (t, $J = 7.6$ Hz, 3H), 1.24-1.10 (m, 2H). **Isomer SFC** $t_R = 1.644$ min in 3 min chromatography (Column: OD-H; Method Name: OD-H_3UM_5_5_40_4ML_3MIN.M, ee = 99%).

[00323] Basic preparative HPLC Method:

Mobile phase A: water with 0.05% ammonia hydroxide solution

Mobile phase B: MeCN

Flow rate: 25 mL/min.


Detection: UV 220 nm

Column: Gemini 150*25 5u

Column temperature: 40 °C


Time in min	%A	%B
0.00	43	57
10.00	13	87
10.20	0	100
12.00	0	100

[00324] The following compounds are prepared following analogous procedures:




Cpd No	Cy ²	L ²	R ⁷	R ⁸
Hy4-2	<i>trans</i> -4-(trifluoromethyl)cyclohexyl	CH ₂ O	CH ₂ OH	H
Hy4-3	4-(trifluoromethyl)phenyl	CH ₂ O	H	H
Hy4-4	4-(trifluoromethyl)phenyl	CH ₂ O	CH ₂ OH	H

The following compound is prepared by a similar procedure using SEM protection of the indole NH.

	
Cpd No Hy4-5	

Example 11

N-(4-(ethylsulfonyl)benzyl)-1-methyl-7-((4-(trifluoromethyl)benzyl)oxy)-1H-pyrrolo[2,3-c]pyridine-3-carboxamide (Cpd No Hy5-3)



[00325] Step 1

[00326] To a solution of methyl 7-chloro-1H-pyrrolo[2,3-c]pyridine-3-carboxylate (100 mg, 0.51 mmol) in anhydrous MeOH (3 mL) was added SOCl₂ (1 mL) dropwise. The mixture was stirred at 40 °C for 16 h. LCMS showed that the reaction was completed. The mixture was concentrated under reduced pressure to afford crude methyl 7-chloro-1-methyl-1H-pyrrolo[2,3-c]pyridine-3-carboxylate (107 mg, 100%) as a yellow solid, which was used for the next step directly without further purification. **LC-MS**: t_R = 0.581 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2mm), MS (ESI) m/z 210.7 [M+H]⁺.

[00327] Step 2

[00328] To a solution of crude methyl 7-chloro-1H-pyrrolo[2,3-c]pyridine-3-carboxylate (100 mg, 0.48 mmol) in anhydrous THF (5 mL) was added KOH (67 mg, 1.20 mmol) and MeI (224 mg, 1.6 mmol). The mixture was stirred at 50 °C for 4 h. LCMS showed that the reaction was

completed. The mixture was added with water (5 mL) and extracted with CH₂Cl₂ (3 X 5 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude methyl 7-chloro-1-methyl-1H-pyrrolo[2,3-c]pyridine-3-carboxylate (110 mg, 103%) as a yellow solid, which was used for the next step directly without further purification. **LC-MS:** t_R = 0.742 min in 5-95AB_1.5 min chromatography (RP-18e, 25-2mm), MS (ESI) m/z 224.8 [M+H]⁺.

[00329] Step 3

[00330] The mixture of crude methyl 7-chloro-1-methyl-1H-pyrrolo[2,3-c]pyridine-3-carboxylate (20 mg, 0.09 mmol), (4-(trifluoromethyl)phenyl)methanol (32 mg, 0.18 mmol), Pd₂(dba)₃ (16 mg, 0.018 mmol), BINAP (11 mg, 0.018 mmol), Cs₂CO₃ (88 mg, 0.27 mmol) in anhydrous toluene (1 mL) was stirred at 110 °C for 16 h under N₂. TLC (petroleum ether / ethyl acetate = 5/1) showed that the reaction was completed. The mixture was filtered through celite and the filter cake was washed with CH₂Cl₂ (15 mL). The filtrate was concentrated under reduced pressure. The residue was purified by preparative TLC (petroleum ether / ethyl acetate = 5 / 1) to afford methyl 1-methyl-7-((4-(trifluoromethyl)benzyl)oxy)-1H-pyrrolo[2,3-c]pyridine-3-carboxylate (11 mg, 34%) as a colorless oil. **LC-MS** t_R = 0.968 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 365.0 [M+H]⁺.

[00331] Step 4

[00332] A mixture of methyl 1-methyl-7-((4-(trifluoromethyl)benzyl)oxy)-1H-pyrrolo[2,3-c]pyridine-3-carboxylate (16 mg, 0.044 mmol), NaOH (88 mg, 2.2 mmol) in THF/ MeOH/ H₂O (6 mL, v/v = 1/4/1) was stirred at 40 °C for 16 h. TLC (petroleum ether / ethyl acetate = 5/1) showed that the most of starting material was consumed. The mixture was adjusted to pH = 3 with 1 N HCl solution and extracted with CH₂Cl₂ (3 X 10 mL). The combined organic layers were washed with brine (15 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give crude 1-methyl-7-((4-(trifluoromethyl)benzyl)oxy)-1H-pyrrolo[2,3-c]pyridine-3-carboxylic acid (10 mg, 62.5%) as a white solid, which was used for next step directly without further purification.

[00333] Step 5

[00334] To a mixture of crude 1-methyl-7-((4-(trifluoromethyl)benzyl)oxy)-1H-pyrrolo[2,3-c]pyridine-3-carboxylic acid (10 mg, 0.029 mmol), (4-(ethylsulfonyl)phenyl)methanamine hydrochloride HCl salt (14 mg, 0.58 mmol) and HATU (22 mg, 0.058 mmol) in anhydrous CH₂Cl₂ (1 mL) was added with Et₃N (6 mg, 0.058 mmol). The mixture was stirred at rt for 3 h. The mixture was added water (10 mL) and extracted with CH₂Cl₂ (3 X 10 mL). The combined

organic layers were washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give crude product (15 mg, 100%), which was purified by basic preparative HPLC separation and dry-freezing directly to afford *N*-(4-(ethylsulfonyl)benzyl)-1-methyl-7-((4-(trifluoromethyl)benzyl)oxy)-1H-pyrrolo[2,3-*c*]pyridine-3-carboxamide (**Hy5-3**, 7.10 mg, 45 %) as a white solid. **LC-MS** *t_R* = 0.776 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), **MS** (ESI) *m/z* 532.0 [M+H]⁺. **¹H NMR** (CDCl₃ 400 MHz): δ 7.90-7.80 (m, 3H), 7.65-7.62 (m, 3H), 7.60-7.55 (m, 2H), 7.52 (d, *J* = 8.0 Hz, 2H), 7.46 (d, *J* = 5.6 Hz, 1H), 6.39 (t, *J* = 6.0 Hz, 1H), 5.61 (s, 2H), 4.75 (d, *J* = 6.4 Hz, 2H), 4.08 (s, 3H), 3.09 (q, *J* = 7.2 Hz, 2H), 1.25 (t, *J* = 7.2 Hz, 3H).

[00335] Basic preparative HPLC Method:

Mobile phase A: water with 0.05% ammonia hydroxide solution

Mobile phase B: MeCN

Flow rate: 25 mL/min.


Detection: UV 220 nm

Column: Phenomenex Gemini 150*25mm*10um

Column temperature: 40 °C

Time in min	%A	%B
0.00	55	45
10.00	25	75
10.20	0	100
13.00	0	100


[00336] The following compounds are prepared using analogous procedures:



Cpd No	Cy ²	L ²	R ^{2a}	R ⁷	R ⁸
Hy5-1	4-(trifluoromethyl)phenyl	O	Me	H	H
Hy5-2	<i>trans</i> -4-(trifluoromethyl)cyclohexyl	CH ₂ O	H	H	H
Hy5-4	<i>trans</i> -4-(trifluoromethyl)cyclohexyl	CH ₂ O	Me	H	H
Hy5-5	<i>trans</i> -4-(trifluoromethyl)cyclohexyl	CH ₂ O	Me	CH ₂ OH	H

Example 12

N-(4-(ethylsulfonyl)benzyl)-2,5-dimethyl-4-(((1*r*,4*r*)-4-(trifluoromethyl)cyclohexyl)methoxy)thieno[2,3-*d*]pyrimidine-6-carboxamide (Cpd No Hy6-2)



[00337] Step 1

[00338] To a solution of *trans*-4-(trifluoromethyl)cyclohexanecarboxylic acid (2.0 g, 11.4 mmol) in anhydrous THF (50 mL) was added LiAlH_4 (20 mL, 0.02 mmol, 1 M in THF) dropwise at 0 °C under N_2 . The mixture was stirred at 0 °C for 4 h. TLC (petroleum ether / ethyl acetate = 3/1) showed that the reaction was completed. The mixture was quenched with water (0.76 mL) and 10% *aq.* NaOH solution (0.76 mL) at 0 °C. The mixture was filtered. The filtrate was dried over anhydrous Na_2SO_4 , filtered, concentrated under reduced pressure to give crude (*trans*-4-(trifluoromethyl)cyclohexyl)methanol (1.73 g, 93%) as a pale yellow solid, which was used for the next step directly without further purification.

[00339] Step 2

[00340] To a solution of crude methyl 4-chloro-2,5-dimethylthieno[2,3-*d*]pyrimidine-6-carboxylate (100 mg, 0.39 mmol) in anhydrous toluene (6 mL) was added (*trans*-4-(trifluoromethyl)cyclohexyl)methanol (71 mg, 0.39 mmol), BINAP (24 mg, 0.04 mmol), Cs_2CO_3 (380 mg, 1.17 mmol) and Pd_2dba_3 (36 mg, 0.04 mmol) under N_2 . The mixture was stirred at 110 °C for 2 h. LCMS showed that the reaction was completed. The mixture was diluted with water (10 mL) and extracted with ethyl acetate (3 X 10 mL). The combined organic layers were washed with brine (20 mL), dried over anhydrous Na_2SO_4 , filtered, concentrated under reduced

pressure. The residue was purified by preparative TLC with petroleum ether / ethyl acetate = 3 / 1 to afford methyl 2,5-dimethyl-4-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)thieno[2,3-d]pyrimidine-6-carboxylate (59 mg, 38%) as a pale yellow solid. **LC-MS** t_R = 4.931 min in 10-80AB_7min chromatography (Welch Xtimate 3um, C18, 2.1*30mm), MS (ESI) m/z 403.1 $[M+H]^+$.

[00341] Step 3

[00342] To a solution of methyl 2,5-dimethyl-4-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)thieno[2,3-d]pyrimidine-6-carboxylate (50 mg, 0.12 mmol) in MeOH / H₂O (3 mL, v/v = 1/1) was added NaOH (99 mg, 2.49 mmol). The mixture was stirred at rt for 2 h. LCMS showed that the reaction was completed. The mixture was concentrated under reduced pressure. The mixture was added with water (10 mL). The aqueous layer was adjusted to pH = 3-4 with 1 N HCl solution and extracted with ethyl acetate (3 X 10 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude 2,5-dimethyl-4-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)thieno[2,3-d]pyrimidine-6-carboxylic acid (48 mg, 99%) as a pale yellow solid, which was used for the next step directly without further purification. **LC-MS** t_R = 0.844 min in 5-95 AB_1.5 min chromatography (Welch MK RP-18e 25-2mm), MS (ESI) m/z 389.0 $[M+H]^+$.

[00343] Step 4

[00344] To a solution of crude 2,5-dimethyl-4-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)thieno[2,3-d]pyrimidine-6-carboxylic acid (48 mg, 0.12 mmol) in anhydrous DMF (6 mL) was added (4-(ethylsulfonyl)phenyl)methanamine hydrochloride HCl salt (35 mg, 0.15 mmol), HATU (56 mg, 0.15 mmol) and Et₃N (37 mg, 0.37 mmol) under N₂. The mixture was stirred at rt for 2 h. LCMS showed that the reaction was completed. The mixture was added with water (10 mL) and extracted with ethyl acetate (3 X 10 mL). The combined organic layers were washed with H₂O (3 X 10 mL) and brine (10 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC with petroleum ether / ethyl acetate = 1 / 3 to give the crude product, which was purified by SFC (OJ) and basic preparative HPLC separation, then lyophilized directly to afford N-(4-(ethylsulfonyl)benzyl)-2,5-dimethyl-4-((*trans*-4-(trifluoromethyl)cyclohexyl)methoxy)thieno[2,3-d]pyrimidine-6-carboxamide (**Hy6-2**, 30.60 mg, 38%) as a white solid.

[00345] Before separation:

Isomer SFC: t_R = 1.556 and 1.636 min in 3 min chromatography (Column: OJ-H; Method Name: OJ-H_3UM_5_5_40_4ML_3MIN.M, ee = 95%).

SFC separation method:

Instrument: Berger MultiGramTM SFC, Mettler Toledo Co, Ltd

Column: OJ (250mm*30mm,5um)

Mobile phase: A: Supercritical CO₂, B: EtOH(0.05%NH₄OH), A:B =65:35 at 60 mL/min

Column Temp: 38 °C

Nozzle Pressure: 100 Bar

Nozzle Temp: 60 °C

Evaporator Temp: 20 °C

Trimmer Temp: 25 °C

Wavelength: 220 nm

LC-MS t_R = 0.985 min in 5-95 AB chromatography (Welch MK RP-18e 25-2mm), MS (ESI)

m/z 570.1 [M+H]⁺. **Isomer SFC:** t_R = 1.566 min in 3 min chromatography (Column: OJ-H;

Method Name: OJ-H_3UM_5_5_40_4ML_3MIN.M, ee = 100%). **¹H NMR** (CDCl₃ 400 MHz):

δ 7.90 (d, J = 8.4 Hz, 2H), 7.55 (d, J = 7.6 Hz, 2H), 6.33-6.26 (m, 1H), 4.73 (d, J = 6.4 Hz, 2H), 4.37 (d, J = 6.8 Hz, 2H), 3.10 (q, J = 7.6 Hz, 2H), 2.84 (s, 3H), 2.27 (s, 3H), 2.08-1.98 (m, 5H), 1.94-1.86 (m, 1H), 1.45-1.32 (m, 2H), 1.28 (t, J = 7.6 Hz, 3H), 1.21-1.11 (m, 2H).

[00346] Basic preparative HPLC Method:

Mobile phase A: water with 0.05% ammonia hydroxide v/v solution

Mobile phase B: MeCN

Flow rate: 25 mL/min.


Detection: UV 220 nm

Column: Phenomenex Synergi C18 150*25*10um

Column temperature: 30 °C

Time in min	%A	%B
0.00	40	60
10.00	10	90
10.20	0	100
13.00	0	100


[00347] The compounds shown below are prepared following analogous procedures:



Cpd No	Cy ²	L ²	R ⁷	R ⁸
Hy6-1	4-(trifluoromethyl)phenyl	CH ₂ O	H	H
Hy6-3	4-(trifluoromethyl)phenyl	CH ₂ O	CH ₂ OH	H
Hy6-4	<i>trans</i> -4-(trifluoromethyl)cyclohexyl	CH ₂ O	CH ₂ OH	H

Example 13

(S)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide and (R)-4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide (Cpd Nos Hy7-17.1 and Hy7-17.2)



[00348] Step 1

[00349] To a solution of methyl 4-(cyanomethyl)benzoate (5 g, 28.6 mmol) in anhydrous THF (50 mL) was added NaH (1.14 g, 28.6 mmol, 60% in mineral oil) at 0 °C under N₂. The mixture was stirred at 0 °C for 20 min, then iodoethane (4.01 g, 25.74 mmol) was added dropwise to the mixture at 0 °C. The mixture was stirred at rt for 2 h under N₂. TLC (petroleum ether / ethyl acetate = 3/1) showed most of methyl 4-(cyanomethyl)benzoate was consumed. The mixture was quenched with water (100 mL) and extracted with ethyl acetate (3 X 100 mL). The combined organic layers were washed with brine (200 mL), dried over anhydrous sodium

sulfate, filtered and concentrated under reduced pressure. The residue was purified by column chromatograph on silica gel eluting with petroleum ether / ethyl acetate = 20/1 - 3/1 to give methyl 4-(1-cyanopropyl)benzoate (3.1 g, 53%) as a colorless oil. ¹H NMR (CDCl₃ 400 MHz): δ 8.04 (d, *J* = 8.0 Hz, 2H), 7.40 (d, *J* = 8.4 Hz, 2H), 3.91 (s, 3H), 3.79 (t, *J* = 7.2 Hz, 1H), 2.00-1.90 (m, 2H), 1.07 (t, *J* = 7.2 Hz, 3H).

[00350] Step 2

[00351] To a solution of methyl 4-(1-cyanopropyl)benzoate (1.0 g, 4.93 mol) in methanol (10 mL) and NH₃·H₂O (5 mL) was added Raney Ni (100 mg, 10% w/w). The mixture was stirred at rt for 16 h under H₂ (30 psi). LCMS showed that the reaction was completed. The mixture was filtered through celite and the filter cake was washed with MeOH (20 mL). The filtrate was concentrated under reduced pressure to give crude methyl 4-(1-aminobutan-2-yl)benzoate (950 mg, 93%) as a green oil, which was used for the next step directly without further purification. LC-MS *t_R* = 0.530 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) *m/z* 207.9 [M+H]⁺.

[00352] Step 3

[00353] To a mixture of crude methyl 4-(1-aminobutan-2-yl)benzoate (850 mg, 4.106 mmol) in DCM (30 mL) was added Et₃N (1.244 g, 12.319 mmol) and TFAA (2.587 g, 12.319 mmol) at 0 °C. The mixture was stirred at rt for 2 h under N₂. TLC (petroleum ether / ethyl acetate = 5/1) showed that the reaction was completed. The mixture was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether / ethyl acetate = 10/1-5/1 to give methyl 4-(1-(2,2,2-trifluoroacetamido)butan-2-yl)benzoate (1.16 g, 93%) as a colorless oil. ¹H NMR (CDCl₃ 400 MHz): δ 8.01 (d, *J* = 8.4 Hz, 2H), 7.23 (d, *J* = 8.4 Hz, 2H), 6.02 (brs, 1H), 3.91 (s, 3H), 3.86-3.75 (m, 1H), 3.43-3.33 (m, 1H), 2.90-2.79 (m, 1H), 1.85-1.60 (m, 2H), 0.82 (t, *J* = 7.2 Hz, 3H).

[00354] Step 4

[00355] To a mixture of 4-(1-(2,2,2-trifluoroacetamido)butan-2-yl)benzoate (800 mg, 2.64 mmol) and (HCHO)_n (675 mg, 22.5 mmol) was added conc. H₂SO₄ (6 mL) and acetic acid (4 mL). The mixture was stirred at rt for 16 h. TLC (petroleum ether / ethyl acetate = 10/1) showed that the reaction was completed. The mixture was poured into ice-water (50 mL) and extracted with ethyl acetate (3 X 50 mL). The combined organic layers were washed with brine (100 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by column chromatograph on silica gel eluting with petroleum ether / ethyl acetate = 1/0-5/1 to give methyl 4-ethyl-2-(2,2,2-trifluoroacetyl)-1,2,3,4-

tetrahydroisoquinoline-7-carboxylate (832 mg, 100%) as a colorless oil. ¹H NMR (CDCl₃, 400 MHz): δ 7.91-7.75 (m, 2H), 7.25-7.20 (m, 1H), 4.87-4.80 (m, 1.5H), 4.66-4.56 (m, 1H), 4.32-4.31 (m, 0.5H), 3.85 (s, 3H), 3.66-3.61 (m, 0.5H), 3.40-3.33 (m, 0.5H), 2.82-2.78 (m, 1H), 1.70-1.58 (m, 2H), 1.06-0.92 (m, 3H).

[00356] Step 5

[00357] To a mixture of methyl 4-ethyl-2-(2,2,2-trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxylate (832 mg, 2.64 mmol) in methanol (2 mL) and water (0.4 mL) was added K₂CO₃ (729 mg, 5.28 mmol). The mixture was stirred at rt for 16 h. LCMS showed that the reaction was completed. The mixture was quenched with water (50 mL) and extracted with ethyl acetate (3 X 50 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give crude methyl 4-ethyl-1,2,3,4-tetrahydroisoquinoline-7-carboxylate (470 mg, 81%) as a pale yellow solid, which was used directly for the next step without further purification.

[00358] Crude methyl 4-ethyl-1,2,3,4-tetrahydroisoquinoline-7-carboxylate (180 mg, 0.82 mmol) was purified by column chromatography on silica gel eluting with petroleum ether/ ethyl acetate = 1/0-4/1 to give methyl 4-ethyl-1,2,3,4-tetrahydroisoquinoline-7-carboxylate (**1**, 133.50 mg, 74%) as a pale yellow solid. LC-MS t_R = 0.520 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 220.0 [M+H]⁺. ¹H NMR (CDCl₃, 400 MHz): δ 7.81 (d, J = 8.0 Hz, 1H), 7.72 (s, 1H), 7.26 (d, J = 8.0 Hz, 1H), 3.95-3.87 (m, 1H), 3.92 (s, 3H), 3.61-3.53 (m, 1H), 3.30-3.18 (m, 1H), 2.96-2.86 (m, 1H), 2.81-2.70 (m, 2H), 1.83-1.72 (m, 2H), 0.98 (q, J = 7.2 Hz, 3H).

[00359] Step 6

[00360] A mixture of crude methyl 4-ethyl-1,2,3,4-tetrahydroisoquinoline-7-carboxylate (100 mg, 0.46 mmol), 2-chloro-5-(trifluoromethyl)pyrimidine (99 mg, 0.55 mmol) and DIPEA (178 mg, 1.38 mmol) in CH₃CN (1 mL) was stirred at 100 °C for 2 h in a sealed tube. LCMS showed that the reaction was completed. The mixture was concentrated under reduced pressure. The residue was purified by preparative TLC with petroleum ether / ethyl acetate = 8/1 to afford methyl 4-ethyl-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxylate (140 mg, 81.8%) as a yellow oil. LC-MS t_R = 0.911 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 366.0 [M+H]⁺.

[00361] Step 7

[00362] To a mixture of methyl 4-ethyl-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxylate (140 mg, 0.38 mmol) in THF (4 mL) was added 2 N aq.

NaOH (2 mL) solution. The mixture was stirred at 40 °C for 3 h. LCMS showed that the reaction was completed. The reaction solution was added with water (10 mL) and concentrated under reduced pressure to remove THF. The residue was acidified by 1 N HCl to pH = 3, and then extracted with ethyl acetate (3 X 10 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude 4-ethyl-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid (115 mg, 85.8%) as a light yellow solid, which was used for the next step without further purification. LC-MS t_R = 0.824 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), MS (ESI) m/z 351.8 [M+H]⁺.

[00363] Step 8

[00364] To a mixture of crude 4-ethyl-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid (108 mg, 0.31 mmol), (R)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethanol hydrochloride (110 mg, 0.40 mmol) and HATU (190 mg, 0.50 mmol) in dry CH₂Cl₂ (2 mL) was added Et₃N (94 mg, 0.93 mmol). The mixture was stirred at rt for 2 h. LCMS showed that the reaction was completed. The mixture was filtered and the filtrate was concentrated under reduced pressure to give the crude product. The crude product was purified by preparative TLC with petroleum ether / ethyl acetate = 1/1, SFC separation (AD-H) and HCl preparative HPLC separation, then dry-freezing directly to give the two enantiomers of 4-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)pyrimidin-2-yl)-1,2,3,4-tetrahydroisoquinoline-7-carboxamide as white solids.

[00365] Before SFC separation :

Isomer SFC t_R = 1.029 and 1.524 min in 5 min chromatography (Column: AD-H; Method Name: AD-H_3UM_5_40_4ML_5MIN.M, ee = 0.32%).

SFC separation condition:

Instrument: SFC-80-(8)

Column: AD 250mm*30mm*10um

Mobile phase: A: Supercritical CO₂, B: ⁱPrOH (0.05% NH₄OH), A:B=50:50 at 80 mL/min

Column Temp: 40 °C

Nozzle Pressure: 100Bar

Nozzle Temp: 60 °C

Evaporator Temp: 20 °C

Trimmer Temp: 25 °C

Wavelength: 220 nm

(**Hy7-17.2**, 20.80 mg, 21.4%) as a white solid. **LC-MS** t_R = 0.909 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), **MS** (ESI) m/z 563.1 $[M+H]^+$. **^1H NMR** ($CDCl_3$ 400 MHz): δ 8.57 (s, 2H), 7.88 (d, J = 8.0 Hz, 2H), 7.73 (s, 1H), 7.67 (d, J = 8.0 Hz, 1H), 7.59 (d, J = 8.4 Hz, 2H), 7.29 (d, J = 8.0 Hz, 1H), 7.18 (d, J = 6.8 Hz, 1H), 5.33-5.27 (m, 2H), 4.79-4.67 (m, 2H), 4.11-4.00 (m, 2H), 3.56 (dd, J = 3.6, 13.2 Hz, 1H), 3.10 (q, J = 7.6 Hz, 2H), 2.91-2.88 (m, 1H), 1.65-1.56 (m, 2H), 1.28 (t, J = 7.6 Hz, 3H), 1.02 (t, J = 7.6 Hz, 3H). **Isomer SFC** t_R = 0.697 min in 3 min chromatography (Column: AD-H; Method Name: AD-H_3UM_4_40_4ML_3MIN.M, ee = 100%).

[00366] HCl preparative HPLC Method:

Mobile phase A: water(0.05%HCl)-ACN

Mobile phase B: MeCN

Flow rate: 25 mL/min.

Detection: UV 220 nm

Column: Phenomenex Synergi C18 150*25*10um

Column temperature: 40 °C

Time in min	%A	%B
0.00	45	55
10.00	15	85
10.20	0	100
13.00	0	100

[00367] (Hy7-17.1, 24.40 mg, 25.2%) as a white solid. **LC-MS** t_R = 0.909 min in 5-95AB_1.5 min chromatography (Welch MK RP-18e, 25-2 mm), **MS** (ESI) m/z 563.1 $[M+H]^+$. **^1H NMR** ($CDCl_3$ 400 MHz): δ 8.57 (s, 2H), 7.87 (d, J = 8.4 Hz, 2H), 7.73 (s, 1H), 7.67 (d, J = 8.0 Hz, 1H), 7.59 (d, J = 8.0 Hz, 2H), 7.29 (d, J = 7.6 Hz, 1H), 7.22 (d, J = 6.4 Hz, 1H), 5.33-5.25 (m, 2H), 4.77-4.68 (m, 2H), 4.09-3.98 (m, 2H), 3.57 (dd, J = 3.6, 12.8 Hz, 1H), 3.10 (q, J = 7.6 Hz, 2H), 2.90-2.88 (m, 1H), 1.68-1.56 (m, 2H), 1.29 (t, J = 7.6 Hz, 3H), 1.05 (t, J = 7.6 Hz, 3H). **Isomer SFC** t_R = 1.035 min in 3 min chromatography (Column: AD-H; Method Name: AD-H_3UM_4_40_4ML_3MIN.M, ee = 100%).

[00368] HCl preparative HPLC Method:

Mobile phase A: water(0.05%HCl)-ACN

Mobile phase B: MeCN

Flow rate: 25 mL/min.

Detection: UV 220 nm


Column: Phenomenex Synergi C18 150*25*10um

Column temperature: 40 °C





Time in min %A %B



0.00	45	55
10.00	15	85
10.20	0	100
13.00	0	100

[00369] The following compounds are prepared using procedures analogous to those described above:



Cpd No ^a	Cy ²	L ²	R ²	R ⁷	R ⁸
Hy7-1.1	5-fluoro-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-1.2	5-fluoro-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-2.1	5-cyano-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-2.2	5-cyano-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-3.1	5-chloro-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-3.2	5-chloro-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-4.1	5-cyclopropyl-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-4.2	5-cyclopropyl-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-5.1	5-ethoxy-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-5.2	5-ethoxy-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-6.1	5-chloro-2-pyrimidinyl	bond	Et	CH ₂ OH	H
Hy7-6.2	5-chloro-2-pyrimidinyl	bond	Et	CH ₂ OH	H
Hy7-7.1	4-(trifluoromethyl)phenyl	bond	Et	H	H
Hy7-7.2	4-(trifluoromethyl)phenyl	bond	Et	H	H
Hy7-8.1	5-(trifluoromethyl)-2-pyridyl	bond	Et	H	H
Hy7-8.2	5-(trifluoromethyl)-2-pyridyl	bond	Et	H	H
Hy7-9.1	5-(trifluoromethyl)-2-pyrimidinyl	bond	Et	H	H
Hy7-9.2	5-(trifluoromethyl)-2-pyrimidinyl	bond	Et	H	H
Hy7-11.1	5-cyclopropyl-2-pyrimidinyl	bond	Et	CH ₂ OH	H
Hy7-11.2	5-cyclopropyl-2-pyrimidinyl	bond	Et	CH ₂ OH	H
Hy7-13.1	5-(trifluoromethyl)-3-methyl-2-pyridyl	bond	Et	H	H
Hy7-13.2	5-(trifluoromethyl)-3-methyl-2-pyridyl	bond	Et	H	H
Hy7-14.1	4-(trifluoromethyl)-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-14.2	4-(trifluoromethyl)-2-pyrimidinyl	bond	i-Pr	H	H


Hy7-15.1	5-(trifluoromethyl)-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-15.2	5-(trifluoromethyl)-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-16.1	5-(ethoxycarbonyl)-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-16.2	5-(ethoxycarbonyl)-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-19.1	5-(4-morpholinyl)-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-19.2	5-(4-morpholinyl)-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-20.1	5-(6-methoxy-2-pyridyl)-2-pyrimidinyl	bond	Et	H	H
Hy7-20.2	5-(6-methoxy-2-pyridyl)-2-pyrimidinyl	bond	Et	H	H
Hy7-21.1		bond	i-Pr	H	H
Hy7-21.2		bond	i-Pr	H	H
Hy7-22.1	5-(trifluoromethyl)-3-methyl-2-pyridyl	bond	Et	CH ₂ OH	H
Hy7-22.2	5-(trifluoromethyl)-3-methyl-2-pyridyl	bond	Et	CH ₂ OH	H
Hy7-23.1	5-(2-methoxy-4-pyridyl)-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-23.2	5-(2-methoxy-4-pyridyl)-2-pyrimidinyl	bond	i-Pr	H	H
Hy7-24.1		bond	i-Pr	H	H
Hy7-24.2		bond	i-Pr	H	H

Hy7- 25.1		bond	i-Pr	H	H
Hy7- 25.2		bond	i-Pr	H	H
Hy7- 26.1	5-(ethoxycarbonyl)-4-(trifluoromethyl)-2-pyrimidinyl	bond	i-Pr	H	H
Hy7- 26.2	5-(ethoxycarbonyl)-4-(trifluoromethyl)-2-pyrimidinyl	bond	i-Pr	H	H
Hy7- 27.1	4-fluorophenyl	CH ₂	i-Pr	H	H
Hy7- 27.2	4-fluorophenyl	CH ₂	i-Pr	H	H
Hy7- 28.1	4-cyanophenyl	CH ₂	i-Pr	H	H
Hy7- 28.2	4-cyanophenyl	CH ₂	i-Pr	H	H
Hy7- 29.1	4-chlorophenyl	CH ₂	i-Pr	H	H
Hy7- 29.2	4-chlorophenyl	CH ₂	i-Pr	H	H
Hy7- 30.1	4-tetrahydropyranyl	bond	Et	CH ₂ OH	H
Hy7- 30.2	4-tetrahydropyranyl	bond	Et	CH ₂ OH	H
Hy7- 31.1	5-(2-oxazolyl)-2-pyrimidinyl	bond	Et	H	H
Hy7- 31.2	5-(2-oxazolyl)-2-pyrimidinyl	bond	Et	H	H
Hy7- 32.1	4,4-difluorocyclohexyl	bond	Et	CH ₂ OH	H
Hy7- 32.2	4,4-difluorocyclohexyl	bond	Et	CH ₂ OH	H
Hy7- 33.1	5-(2-oxazolyl)-2-pyrimidinyl	bond	Et	CH ₂ OH	H
Hy7- 33.2	5-(2-oxazolyl)-2-pyrimidinyl	bond	Et	CH ₂ OH	H
Hy7- 34.1	5-(ethoxycarbonyl)-2-pyrimidinyl	bond	Et	CH ₂ OH	H
Hy7- 34.2	5-(ethoxycarbonyl)-2-pyrimidinyl	bond	Et	CH ₂ OH	H
Hy7- 35.1	4-(trifluoromethyl)cyclohexyl	bond	Et	CH ₂ OH	H

Hy7-35.2	4-(trifluoromethyl)cyclohexyl	bond	Et	CH ₂ OH	H
Hy7-35.3	4-(trifluoromethyl)cyclohexyl	bond	Et	CH ₂ OH	H
Hy7-35.4	4-(trifluoromethyl)cyclohexyl	bond	Et	CH ₂ OH	H
Hy7-36.1	5-(trifluoromethyl)-2-pyrimidinyl	bond	CF ₃	H	H
Hy7-36.2	5-(trifluoromethyl)-2-pyrimidinyl	bond	CF ₃	H	H
Hy7-37.1	3-cyano-5-(trifluoromethyl)-2-pyridyl	bond	Et	CH ₂ OH	H
Hy7-37.2	3-cyano-5-(trifluoromethyl)-2-pyridyl	bond	Et	CH ₂ OH	H


^a Isomer were separated by chromatography on chiral columns.

[00370] The following compounds are prepared by procedures analogous to those described above:




^{a, b} The isomers were separated by chromatography on a chiral column.

Cpd Nos Hy7-12.1 and 12.2^a



Example 15

(R)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1-((4-(trifluoromethyl)phenoxy)methyl)isoquinoline-6-carboxamide (Hy1A-24)

[00371] LC-MS Data is presented in the following tables.

[00372] Table 1

Cpd No	LC-MS Method	tR (min)	Mass Observed
Hy1A-1	4	1.023	513.1 (M+H)
Hy1A-2	2	0.921	515.1 (M+H)
Hy1A-3.1	2	0.937	521.1 (M+H)
Hy1A-3.2	2	0.819	521.1 (M+H)
Hy1A-4	2	0.962	529.1 (M+H)
Hy1A-5	2	0.856	535.0 (M+H)
Hy1A-6	2	0.881	549.1 (M+H)
Hy1A-7	2	0.923	559.1 (M+H)
Hy1A-8	2	0.81	565.1 (M+H)
Hy1A-9	2	0.898	569.1 (M+H)
Hy1A-10	2	0.829	579.1 (M+H)
Hy1A-11	2	0.855	599.1 (M+H)
Hy1A-12	2	0.762	519.1 (M+H)
Hy1A-13	2	0.912	523.1 (M+H)

Hy1A-14	2	0.79	547.0 (M+H)
Hy1A-15	2	0.809	575.1 (M+H)
Hy1A-16	2	0.869	579.1 (M+H)
Hy1A-17	2	0.686	580.1 (M+H)
Hy1A-18.1	2	0.798	581.1 (M+H)
Hy1A-18.2	2	0.806	581.1 (M+H)
Hy1A-18.3	2	0.782	581.1 (M+H)
Hy1A-19	2	0.655	594.1 (M+H)
Hy1A-20	2	0.945	563.1 (M+H)
Hy1A-21	2	0.900	605.1 (M+H)
Hy1A-22.1	2	0.859	580.1 (M+H)
Hy1A-22.2	2	0.857	580.1 (M+H)
Hy1A-23	2	0.833	595.1 (M+H)
Hy1A-24	1	1.69	559.5 (M+H)

[00373] Table 2

Cpd No	LC-MS Method	tR (min)	Mass Observed
Hy1B-1	2	0.803	543.1 (M+H)
Hy1B-2	2	0.686	549.1 (M+H)
Hy1B-3	2	0.678	573.0 (M+H)
Hy1B-4	2	0.665	579.1 (M+H)

[00374] Table 3

Cpd No	LC-MS Method	tR (min)	Mass Observed
Hy1C-1	2	0.758	m/z 566.1 (M+H)
Hy1C-2	2	0.795	m/z 536.1 (M+H)

[00375] Table 4

Cpd No	LC-MS Method	tR (min)	Mass Observed
Hy2A-1.1	1	1.80	504
Hy2A-1.2	1	1.80	504

Hy2A-2	1	1.79	547
Hy2A-3	1	1.79	547
Hy2A-4.1	2	0.917	551.1 (M+H)
Hy2A-4.2	2	0.915	551.2 (M+H)
Hy2A-5	1	1.86	619
Hy2A-5.1	2	0.977	619.1 (M+H)
Hy2A-5.2	2	0.983	619.1 (M+H)
Hy2A-6	1	0.79	467
Hy2A-7	1	0.75	474
Hy2A-8	1	0.84	481
Hy2A-9	1	0.87	483
Hy2A-10	1	0.8	488
Hy2A-11	1	0.91	497
Hy2A-12.1	2	0.736	507.2 (M+H)
Hy2A-12.2	2	0.742	507.2 (M+H)
Hy2A-13	1	0.95	509
Hy2A-14	1	0.94	516
Hy2A-15.1	5	0.934	516.2 (M+H)
Hy2A-15.2	2	0.682	516.1 (M+H)
Hy2A-16.1	2	0.734	517.1 (M+H)
Hy2A-16.2	2	0.727	517.1 (M+H)
Hy2A-17.1	2	0.745	517.1 (M+H)
Hy2A-17.2	2	0.749	517.1 (M+H)
Hy2A-18	2	0.738	521.2 (M+H)
Hy2A-19	1	1.03	525
Hy2A-20.1	2	0.777	525.1 (M+H)
Hy2A-20.2	5	1.002	525.2 (M+H)
Hy2A-21	1	0.94	526
Hy2A-22.1	5	0.894	535.2 (M+H)
Hy2A-22.2	5	0.904	535.2 (M+H)
Hy2A-23.1	4	0.976	544.2 (M+H)
Hy2A-23.2	4	0.976	544.2 (M+H)

Hy2A-24	3	2.077	556.3 (M+H)
Hy2A-25.1	2	0.765	558.1 (M+H)
Hy2A-25.2	2	0.768	558.1 (M+H)
Hy2A-26.1	4	0.874	571.1 (M+H)
Hy2A-26.2	4	0.882	571.1 (M+H)
Hy2A-27	2	0.662	551.1 (M+H)
Hy2A-28.1	2	0.644	581.1 (M+H)
Hy2A-28.2	2	0.635	581.1 (M+H)
Hy2A-28.3	2	0.639	581.2 (M+H)
Hy2A-29.1	2	0.635	582.1 (M+H)
Hy2A-29.2	2	0.645	582.1 (M+H)
Hy2A-29.3	2	0.631	582.1 (M+H)

[00376] Table 5

Cpd No	LC-MS Method	tR (min)	Mass Observed
Hy2B-1.1	2	0.792	505.2 (M+H)
Hy2B-1.2	2	0.788	505.1 (M+H)
Hy2B-2.1	2	0.840	514.1 (M+H)
Hy2B-2.2	2	0.841	514.1 (M+H)
Hy2B-3.1	2	0.862	548.0 (M+H)
Hy2B-3.2	2	0.861	548.1 (M+H)
Hy2B-4.1	2	0.841	552.2 (M+H)
Hy2B-4.2	2	0.836	552.1 (M+H)
Hy2B-5.1	2	0.908	620.1 (M+H)
Hy2B-5.2	2	0.903	620.1 (M+H)
Hy2B-6.1	2	0.757	510.2 (M+H)
Hy2B-6.2	2	0.751	510.1 (M+H)
Hy2B-7.1	2	0.762	517.1 (M+H)
Hy2B-7.2	2	0.678	517.1 (M+H)
Hy2B-8.1	2	0.700	526.1 (M+H)
Hy2B-8.2	2	0.703	526.1 (M+H)
Hy2B-9.1	4	0.935	566.2 (M+H)

Hy2B-9.2	4	0.929	566.2 (M+H)
Hy2B-10.1	3	0.701	567.2 (M+H)
Hy2B-10.2	2	0.708	567.1 (M+H)

[00377] Table 6

Compound	LC-MS Method	tR (min)	Mass observed
Hy3A-1	1	1.59	578
Hy3A-2	1	1.67	548
Hy3A-3	1	1.63	549
Hy3A-4	1	1.52	572
Hy3A-5	1	1.61	566
Hy3A-6	1	1.49	579.6
Hy3A-7	1	1.80	573.6
Hy3B-1	1	1.54	579
Hy3B-2	1	1.65	549
Hy3B-3	1	1.57	550
Hy3B-4	1	1.77	580.6

[00378] Table 7

Compound	LC-MS Method	tR (min)	Mass observed
Hy4A-1	2	0.78	538
Hy4A-2	2	0.74	568
Hy4A-3	2	0.82	532
Hy4A-4	2	0.76	562
Hy4A-5	2	0.76	524

[00379] Table 8

Cpd No	LC-MS Method	tR (min)	Mass Observed
Hy5-1	2	0.764	518.0 (M+H)
Hy5-2	2	0.712	524.1 (M+H)
Hy5-3	2	0.776	532.0 (M+H)

Hy5-4	2	0.752	538.1 (M+H)
Hy5-5	2	0.721	568.1 (M+H)

[00380] Table 9

Cpd No	LC-MS Method	tR (min)	Mass Observed
Hy6-1	2	0.975	564.1 (M+H)
Hy6-2	2	0.985	570.1 (M+H)
Hy6-3	2	0.794	594.1 (M+H)
Hy6-4	6	0.951	600.2 (M+H)

[00381] Table 10

Cpd No	LC-MS Method	tR (min)	Mass observed
Hy7-1.1	2	0.885	497.2 (M+H)
Hy7-1.2	2	0.874	497.1 (M+H)
Hy7-2.1	2	0.894	504.1 (M+H)
Hy7-2.2	2	0.888	504.0 (M+H)
Hy7-3.1	2	0.902	513.0 (M+H)
Hy7-3.2	2	0.898	513.0 (M+H)
Hy7-4.1	2	0.843	519.2 (M+H)
Hy7-4.2	2	0.842	519.1 (M+H)
Hy7-5.1	2	0.915	523.1 (M+H)
Hy7-5.2	2	0.910	523.1 (M+H)
Hy7-6.1	2	0.879	529.1 (M+H)
Hy7-6.2	2	0.887	529.1 (M+H)
Hy7-7.1	2	0.986	531.1 (M+H)
Hy7-7.2	2	0.983	531.1 (M+H)
Hy7-8.1	2	0.806	532.1 (M+H)
Hy7-8.2	2	0.807	532.1 (M+H)
Hy7-9.1	2	0.834	533.1 (M+H)
Hy7-9.2	2	0.823	533.1 (M+H)
Hy7-10.1	2	0.935	534.1 (M+H)

Hy7-10.2	2	0.937	534.1 (M+H)
Hy7-11.1	2	0.722	535.1 (M+H)
Hy7-11.2	2	0.727	535.1 (M+H)
Hy7-12.1	2	0.656	545.1 (M+H)
Hy7-12.2	2	0.651	545.1 (M+H)
Hy7-13.1	2	0.834	546.1 (M+H)
Hy7-13.2	2	0.837	546.1 (M+H)
Hy7-14.1	2	0.856	547.1 (M+H)
Hy7-14.2	2	0.934	547.0 (M+H)
Hy7-15.1	2	0.931	547.1 (M+H)
Hy7-15.2	2	0.918	547.1 (M+H)
Hy7-16.1	2	0.932	551.1 (M+H)
Hy7-16.2	2	0.930	551.1 (M+H)
Hy7-17.1	2	0.909	563.1 (M+H)
Hy7-17.2	2	0.909	563.1 (M+H)
Hy7-18.1	2	0.797	564.1 (M+H)
Hy7-18.2	2	0.793	564.2 (M+H)
Hy7-18.3	2	0.776	564.1 (M+H)
Hy7-18.4	2	0.780	564.1 (M+H)
Hy7-19.1	4	1.081	564.3 (M+H)
Hy7-19.2	4	1.081	564.2 (M+H)
Hy7-20.1	2	0.873	572.1 (M+H)
Hy7-20.2	2	0.769	572.1 (M+H)
Hy7-21.1	2	0.687	572.1 (M+H)
Hy7-21.2	2	0.685	572.0 (M+H)
Hy7-22.1	2	0.926	576.2 (M+H)
Hy7-22.2	2	0.931	576.2 (M+H)
Hy7-23.1	2	0.811	586.0 (M+H)
Hy7-23.2	2	0.803	586.0 (M+H)
Hy7-24.1	2	0.721	586.0 (M+H)
Hy7-24.2	2	0.720	586.0 (M+H)
Hy7-25.1	2	0.709	586.0 (M+H)

Hy7-25.2	4	1.828	586.2 (M+H)
Hy7-26.1	2	0.985	619.1 (M+H)
Hy7-26.2	2	0.981	619.1 (M+H)
Hy7-27.1	2	0.704	509.2 (M+H)
Hy7-27.2	2	0.706	509.1 (M+H)
Hy7-28.1	2	0.695	516.1 (M+H)
Hy7-28.2	2	0.703	516.1 (M+H)
Hy7-29.1	2	0.724	525.1 (M+H)
Hy7-29.2	2	0.715	525.1 (M+H)
Hy7-30.1	2	0.561	501.2 (M+H)
Hy7-30.2	2	0.548	501.2 (M+H)
Hy7-31.1	2	0.771	532.0 (M+H)
Hy7-31.2	2	0.771	532.1 (M+H)
Hy7-32.1	2	0.701	535.3 (M+H)
Hy7-32.2	2	0.710	535.2 (M+H)
Hy7-33.1	2	0.747	562.1 (M+H)
Hy7-33.2	2	0.749	562.3 (M+H)
Hy7-34.1	2	0.764	567.1 (M+H)
Hy7-34.2	2	0.769	567.1 (M+H)
Hy7-35.1	2	0.639	567.1 (M+H)
Hy7-35.2	2	0.734	567.1 (M+H)
Hy7-35.3	2	0.736	567.2 (M+H)
Hy7-35.4	2	0.637	567.1 (M+H)
Hy7-36.1	2	0.919	573.1 (M+H)
Hy7-36.2	2	0.795	573.0 (M+H)
Hy7-37.1	2	0.797	587.1 (M+H)
Hy7-37.2	2	0.793	587.1 (M+H)
Hy7B-1	1	1.89	534.5 (M+H)

BIOLOGICAL ASSAYS

Radio-Ligand ROR γ Binding Assay (Assay 1)

[00382] Compounds described herein were tested for ability to bind to ROR γ in a cell-free competition assay with commercially available radio-ligand (RL), 25-hydroxy [26,27- ^3H]-cholesterol (PerkinElmer, Cat. # NET674250UC), for a ligand binding site on a recombinant ROR γ Ligand Binding Domain (LBD) protein expressed as a 6xHis-Glutathione-S-Transferase (GST) fusion. The assay was performed in 96-well SPA plates (PerkinElmer, Cat. # 1450-401) in 50 mM HEPES buffer, pH 7.4, containing 150 mM NaCl, 5 mM MgCl₂, 10% (v/v) glycerol, 2 mM CHAPS, 0.5 mM β -octylglucopyranoside and 5 mM DTT. Tested compounds were dissolved in DMSO, and semi-log (3.162x) serial dilutions of the compounds were prepared in the same solvent. Two μL of the DMSO solutions were mixed with 28 μL of 8.6 nM 25-hydroxy [26,27- ^3H]-cholesterol and 50 μL of 24 nM ROR γ LBD. The plate was shaken at 700 rpm for 20 min and incubated for 10 min at rt, after which 40 μL of poly-Lys YSi SPA beads (PerkinElmer, Cat. # RPNQ0010) were added to achieve 50 μg of the beads per well. The plate was incubated on an orbital shaker for 20 min and then for 10 min without agitation at rt. SPA signal for tritium beta radiation was registered on PerkinElmer Microbeta plate reader. Percent inhibition values were calculated based on the high signal obtained with DMSO control and the low signal observed with 10 μM standard ROR γ inverse agonist T0901317 (SigmaAldrich, Cat. # T2320). The percent inhibition vs. concentration data were fit into a four-parameter model, and IC₅₀ values were calculated from the fit as the concentrations corresponding to the inflection points on the dose-response curves. Inhibitory constants (K_i) were calculated using the following equation, where [RL] is the concentration in the assay and K_D is a dissociation constant of 25-hydroxy [26,27- ^3H]-cholesterol:

$$K_i = \frac{IC_{50}}{\left(1 + \frac{[RL]}{K_D}\right)}$$

ROR γ t 5xRORE Assay in Jurkat Cells (Assay 2)

[00383] Compounds described herein were tested for ROR γ inverse agonist activity in a cell-based, transcriptional activity assay. Secreted Nanoluc[®] luciferase was used as a reporter for transcriptional activity of the full-length ROR γ t in Jurkat cells (ATCC, Cat. # TIB-152). A reporter plasmid was constructed by inserting 5 repeats of the ROR Response Element (RORE) AAAGTAGGTCA (SEQ ID NO:1) into a commercially available promoterless plasmid pNL1.3[secNluc] (Promega, Cat. # N1021) using KpnI and HindIII restriction sites. The

expression plasmid for ROR γ t was purchased (GeneCopoeia, Cat. # EX-T6988-M02). Jurkat cells (30 million cells) were transfected with 11 μ g of EX-T6988-M02 and 26 μ g of the reporter plasmid in OptiMEM[®] media using Lipofectamine[®] LTX and Plus[™] reagents (Life Technologies, Cat. # 15338-100). After 5-6 hrs of incubation at 37 °C/5% CO₂, the cells were collected, resuspended in phenol-red free RPMI media containing 10% (v/v) delipidated FBS (Hyclone, Cat. # SH30855.03) and dispensed into 96-well clear bottom tissue culture plates (CoStar, Cat. # 3603), at 80,000 cells per well. Tested compounds were added to the cells in the same media (final concentration of DMSO was 0.1% (v/v)), and the plates were incubated at 37 °C/5% CO₂ for 16-18 hrs. Luciferase activity in the conditioned supernatants was determined with NanoGlo[®] assay reagents (Promega, Cat.# N1130). Percent inhibition values were calculated based on the fully inhibited and non-inhibited (DMSO) controls, and the values were regressed against concentrations of the tested compounds to derive IC₅₀ values using a four-parameter non-linear fitting model.

[00384] The results of assays 1 and 2 are shown in Tables 11-20.

[00385] **Table 11**

Cpd No	Avg ROR Bind Ki (nM)	Avg ROR γt/5x IC₅₀ (nM)
Hy1A-1	+++	+
Hy1A-2	++	
Hy1A-3.1	+++	+
Hy1A-3.2	+++	+
Hy1A-4	+++	++
Hy1A-5	+++	++
Hy1A-6	+++	+++
Hy1A-7	+++	++
Hy1A-8	+++	++
Hy1A-9	+++	+++
Hy1A-10	+++	+++
Hy1A-11	+++	+++
Hy1A-12	+++	+
Hy1A-13	+++	++

Hy1A-14	+++	+++
Hy1A-15	+++	++
Hy1A-16	+++	+++
Hy1A-17	+++	+
Hy1A-18.1	+++	+++
Hy1A-18.2	+++	+++
Hy1A-18.3	+++	+
Hy1A-19	+++	++
Hy1A-20	+++	+++
Hy1A-21	+++	+++
Hy1A-22.1	+++	+++
Hy1A-22.2	+++	++
Hy1A-23	+++	+++
Hy1A-24	+++	++

[00386] Table 12

Cpd No	Avg ROR Bind Ki (nM)	Avg ROR γt/5x IC50 (nM)
Hy1B-1	++	
Hy1B-2	++	
Hy1B-3	+++	++
Hy1B-4	++	

[00387] Table 13

Cpd No	Avg ROR Bind Ki (nM)	Avg ROR γt/5x IC50 (nM)
Hy1C-1	++	
Hy1C-2	++	

[00388] Table 14

Cpd No	Avg ROR Bind Ki (nM)	Avg ROR γ t/5x IC50 (nM)
Hy2A-1.1	++	
Hy2A-1.2	++	
Hy2A-2	++	+
Hy2A-3	+++	+
Hy2A-4.1	++	
Hy2A-4.2	++	
Hy2A-5	++	++
Hy2A-5.1	+++	++
Hy2A-5.2	++	
Hy2A-6	+	
Hy2A-7	+	
Hy2A-8	++	
Hy2A-9	++	
Hy2A-10	++	
Hy2A-11	+++	++
Hy2A-12.1	++	++
Hy2A-12.2	+	
Hy2A-13	+++	+++
Hy2A-14	+++	+++
Hy2A-15.1	+++	+++
Hy2A-15.2	+++	
Hy2A-16.1	+++	+++
Hy2A-16.2	++	
Hy2A-17.1	+++	+++
Hy2A-17.2	++	
Hy2A-18	+++	++
Hy2A-19	+++	+++
Hy2A-20.1	+++	+++
Hy2A-20.2	+++	++

Hy2A-21	+++	+++
Hy2A-22.1	+++	++
Hy2A-22.2	++	
Hy2A-23.1	+++	+++
Hy2A-23.2	+++	++
Hy2A-24	++	++
Hy2A-25.1	+++	+++
Hy2A-25.2	++	
Hy2A-26.1	+++	++
Hy2A-26.2	++	
Hy2A-27	+++	+++
Hy2A-28.1	+++	+++
Hy2A-28.2	+++	++
Hy2A-28.3	++	
Hy2A-29.1	++	
Hy2A-29.2	++	
Hy2A-29.3	++	

[00389] Table 15

Cpd No	Avg ROR Bind Ki	Avg ROR γ t/5x IC50
Hy2B-1.1	+	
Hy2B-1.2	+	
Hy2B-2.1	++	
Hy2B-2.2	+	
Hy2B-3.1	++	
Hy2B-3.2	++	
Hy2B-4.1	+	
Hy2B-4.2	+	
Hy2B-5.1	++	
Hy2B-5.2	++	
Hy2B-6.1	+++	+
Hy2B-6.2	++	

Hy2B-7.1	+++	++
Hy2B-7.2	+	
Hy2B-8.1	+++	++
Hy2B-8.2	++	
Hy2B-9.1	+++	++
Hy2B-9.2	++	
Hy2B-10.1	++	++
Hy2B-10.2	++	

[00390] Table 16

Compound	Avg ROR Bind Ki (nM)	Avg ROR γ t/5x IC50 (nM)
Hy3A-1	+++	+++
Hy3A-2	+++	+++
Hy3A-3	+++	+++
Hy3A-4	+++	+++
Hy3A-5	+++	+++
Hy3A-6	+++	+++
Hy3A-7	+++	+++
Hy3B-1	+++	+++
Hy3B-2	+++	+++
Hy3B-3	++	
Hy3B-4	+++	

[00391] Table 17

Cpd No	Avg ROR Bind Ki	Avg ROR γ t/5x IC50
Hy4-1	+++	++
Hy4-2	+++	+++
Hy4-3	+++	++
Hy4-4	+++	++
Hy4-5	++	

[00392] Table 18

Cpd No	Avg ROR Bind Ki (nM)	Avg ROR γt/5x IC50 (nM)
Hy5-1	++	
Hy5-2	++	
Hy5-3	+++	+++
Hy5-4	++	
Hy5-5	++	

[00393] Table 19

Cpd No	Avg ROR Bind Ki (nM)	Avg ROR γt/5x IC50 (nM)
Hy6-1	+++	++
Hy6-2	+++	++
Hy6-3	+++	+++
Hy6-4	+++	+++

[00394] Table 20

Cpd No	Avg ROR Bind Ki (nM)	Avg ROR γt/5x IC50 (nM)
Hy7-1.1	+++	+
Hy7-1.2	+++	+
Hy7-2.1	+++	++
Hy7-2.2	++	
Hy7-3.1	+++	++
Hy7-3.2	+++	+
Hy7-4.1	+++	++
Hy7-4.2	+++	+
Hy7-5.1	+++	++
Hy7-5.2	++	
Hy7-6.1	+++	+
Hy7-6.2	+++	+

Hy7-7.1	+++	+++
Hy7-7.2	+++	++
Hy7-8.1	+++	+++
Hy7-8.2	+++	++
Hy7-9.1	+++	+++
Hy7-9.2	+++	+
Hy7-10.1	+++	+++
Hy7-10.2	++	
Hy7-11.1	+++	++
Hy7-11.2	+++	+
Hy7-12.1	+++	++
Hy7-12.2	+++	++
Hy7-13.1	+++	+++
Hy7-13.2	+++	++
Hy7-14.1	+++	+
Hy7-14.2	+++	+
Hy7-15.1	+++	++
Hy7-15.2	+++	
Hy7-16.1	+++	+++
Hy7-16.2	++	
Hy7-17.1	+++	+++
Hy7-17.2	+++	+
Hy7-18.1	+++	+++
Hy7-18.2	+++	+
Hy7-18.3	++	
Hy7-18.4		
Hy7-19.1	++	
Hy7-19.2	+	
Hy7-20.1	+++	+++
Hy7-20.2	++	
Hy7-21.1	+++	++
Hy7-21.2	+	

Hy7-22.1	+++	+++
Hy7-22.2	+++	++
Hy7-23.1	+++	+++
Hy7-23.2	++	
Hy7-24.1	+++	+++
Hy7-24.2	+	
Hy7-25.1	+++	++
Hy7-25.2	+	
Hy7-26.1	+++	+++
Hy7-26.2	+++	++
Hy7-27.1	+++	+
Hy7-27.2	+++	+++
Hy7-28.1	+++	++
Hy7-28.2	+++	+++
Hy7-29.1	+++	+
Hy7-29.2	+++	++
Hy7-30.1	++	
Hy7-30.2		
Hy7-31.1	+++	+++
Hy7-31.2	+++	+
Hy7-32.1	+++	+
Hy7-32.2	++	
Hy7-33.1	+++	+++
Hy7-33.2	+++	++
Hy7-34.1	+++	+++
Hy7-34.2	+++	
Hy7-35.1	+++	+++
Hy7-35.2	+++	+
Hy7-35.3	+++	
Hy7-35.4	++	
Hy7-36.1	+++	+++
Hy7-36.2	+++	+

Hy7-37.1	+++	+++
Hy7-37.2	+++	
Hy7B-1	++	


“nt” or no value presented = not tested; + means > 1000 nM; ++ means 100 nM – 1000 nM; +++ means < 100 nM.

[00395] While we have described a number of embodiments, it is apparent that our basic examples may be altered to provide other embodiments that utilize the compounds and methods of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the appended claims rather than by the specific embodiments that have been represented by way of example.


[00396] The contents of all references (including literature references, issued patents, published patent applications, and co-pending patent applications) cited throughout this application are hereby expressly incorporated herein in their entireties by reference. Unless otherwise defined, all technical and scientific terms used herein are accorded the meaning commonly known to one with ordinary skill in the art.

CLAIMS**Listing of Claims**

1. A compound having the Formula I:



or a pharmaceutically acceptable salt thereof, wherein



U and V are each independently CR⁹ or N, provided that both are not CR⁹;

X is -C(O)NH- or -NHC(O)-;

n is 0, 1, 2, or 3;

p is 0, 1, 2, or 3;

* designates the attachment to L²;

** designates the attachment to X;

L² is a bond or is selected from CH₂, CH₂CH₂, CHMe, O, C(=O), CH(OH), and CH₂O, wherein the oxygen atom in CH₂O can either be attached to Cy² or Hy, provided that attachment to Hy occurs at a carbon atom on Hy;

L¹ is absent or is CR⁷R⁸;

Cy¹ is selected from aryl, heteroaryl, heterocyclyl, and cycloalkyl, wherein the aryl, heteroaryl, heterocyclyl, and cycloalkyl are each substituted with 1 to 3 groups independently selected from R⁵;

Cy² is selected from aryl, heteroaryl, monocyclic cycloalkyl, and monocyclic heterocyclyl, wherein the aryl, heteroaryl, monocyclic cycloalkyl, and monocyclic heterocyclyl are each optionally substituted with 1 to 3 groups independently selected from R⁶;

R² is (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₁-C₄)haloalkyl, or monocyclic cycloalkyl;

R^{2a} is H, (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₁-C₄)haloalkyl, or monocyclic cycloalkyl;

R^5 and R^6 are each independently selected from halogen, $-CN$, $-OR^c$, $-NR^dR^e$, $-S(O)_kR^c$, $-NR^cS(O)_2R^c$, $-S(O)_2NR^dR^e$, $-C(=O)OR^c$, $-OC(=O)OR^c$, $-OC(=O)R^c$, $-OC(=S)OR^c$, $-C(=S)OR^c$, $-OC(=S)R^c$, $-C(=O)NR^dR^e$, $-NR^cC(=O)R^c$, $-C(=S)NR^dR^e$, $-NR^cC(=S)R^c$, $-NR^cC(=O)OR^c$, $-OC(=O)NR^dR^e$, $-NR^c(C=S)OR^c$, $-OC(=S)NR^dR^e$, $-NR^cC(=O)NR^dR^e$, $-NR^c(C=S)NR^dR^e$, $-C(=S)R^c$, $-C(=O)R^c$, (C_1-C_6) alkyl, cycloalkyl, $-(CH_2)_{1-4}$ -cycloalkyl, heterocyclyl, $-(CH_2)_{1-4}$ -heterocyclyl, aryl, $-NHC(=O)$ -heterocyclyl, $-NHC(=O)$ -cycloalkyl, $-(CH_2)_{1-4}$ -aryl, heteroaryl and $-(CH_2)_{1-4}$ -heteroaryl,

wherein the alkyl, cycloalkyl, heterocyclyl, aryl or heteroaryl portion present in each of said (C_1-C_6) alkyl, cycloalkyl, $-(CH_2)_{1-4}$ -cycloalkyl, heterocyclyl, $-(CH_2)_{1-4}$ -heterocyclyl, aryl, $-(CH_2)_{1-4}$ -aryl, heteroaryl and $-(CH_2)_{1-4}$ -heteroaryl substituent for R^6 are optionally substituted with halogen, OR^c , $-NO_2$, $-CN$, $-NR^cC(=O)R^c$, $-NR^dR^e$, $-S(O)_kR^c$, $-C(=O)OR^c$, $-C(=O)NR^dR^e$, $-C(=O)R^c$, (C_1-C_3) alkyl, halo (C_1-C_3) alkyl, (C_1-C_3) alkoxy (C_1-C_3) alkyl, (C_1-C_3) alkoxy, or halo (C_1-C_3) alkoxy;

each R^c is independently selected from hydrogen and (C_1-C_6) alkyl optionally substituted with hydroxy, (C_1-C_2) alkoxy, $-C(O)NH_2$, $-C(O)O(C_1-C_3)$ alkyl, or 1 to 3 halogen;

each R^d and R^e is independently selected from hydrogen and (C_1-C_6) alkyl;

k is 0, 1 or 2;


any heterocyclyl or heteroaryl portion of Cy^1 or Cy^2 is further optionally substituted with $=O$;

R^7 and R^8 are each independently hydrogen, OR^c , $-C(=O)OR^c$, monocyclic heterocyclyl, halophenyl, quinolin-2(1H)one-4yl-methyl, or (C_1-C_3) alkyl, wherein the (C_1-C_3) alkyl is optionally substituted with OR^c , $-NR^dR^e$, $-C(=O)OR^c$, $-C(=O)NR^dR^e$, or halophenyl;

R^9 is hydrogen, (C_1-C_4) alkyl, (C_1-C_4) haloalkyl, (C_1-C_4) alkoxy, (C_1-C_4) haloalkoxy, halo, cyano, or monocyclic cycloalkyl; and



R^{10} is (C_1-C_4) alkyl, (C_1-C_4) haloalkyl, (C_1-C_4) alkoxy, (C_1-C_4) haloalkoxy, halo, cyano, or monocyclic cycloalkyl.

2. The compound of Claim 1, wherein the compound is of the Formula II:




or a pharmaceutically acceptable salt thereof.


3. The compound of Claim 1 or 2, wherein p is 0, 1, or 2; and R^{10} is (C_1-C_4) alkyl or halo.
4. The compound of Claim 1 or 2, wherein L^2 is a bond or is selected from CH_2 , O, and CH_2O , wherein the oxygen atom in CH_2O is attached to a carbon atom on Hy .
5. The compound of any one of Claims 1 to 4, wherein the compound is of the Formula **III** or **IV**:



or a pharmaceutically acceptable salt thereof, wherein each p is independently 0 or 1.

6. The compound of any one of Claims 1 to 5, wherein the compound is of the Formula **IIIa** or **IVa**:







or a pharmaceutically acceptable salt thereof, wherein each p is independently 0 or 1.


7. The compound of any one of Claim 1 to 6, wherein R^{10} is (C_1-C_4) alkyl; and L^2 is a bond or CH_2 .

8. The compound of any one of Claims 1 to 4, wherein the compound is of the Formula **V** or **VI**:




or a pharmaceutically acceptable salt thereof.

9. The compound of any one of Claims 1 to 4, and 8, wherein the compound is of the Formula **Va** or **Via**:




(Va); or




(Via);

or a pharmaceutically acceptable salt thereof.

10. The compound of any one of Claims 1 to 4, 8, and 9, wherein the compound is of the Formula **Va'** or **Via'**:



(Va'); or




(Via');

or a pharmaceutically acceptable salt thereof.

11. The compound of any one of Claims 1 to 4 and 8 to 10, wherein L^2 is O or CH_2O , wherein the methylene portion of CH_2O is attached to Cy^2 .




12. The compound of any one of Claims 1 to 4 and 8 to 11, wherein R^{2a} is hydrogen or (C₁-C₄)alkyl.

13. The compound of any one of Claims 1 to 4, wherein the compound is of the Formula **VII**:




or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1.

14. The compound of any one of Claims 1 to 4 and 13, wherein the compound is of the Formula **VIIa**, **VIIb**, or **VIIc**:




or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1.


15. The compound of any one of Claims 1 to 4, 13, and 14, wherein the compound is of the Formula **VIIa'**, **VIIb'**, or **VIIc'**:



(VIIa');




(VIIb'); or





(VIIc');

or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1.

16. The compound of any one of Claims 1 to 4 and 13 to 15, wherein the compound is of the Formula **VIIa''**, **VIIb''**, or **VIIc''**:




(VIIa'');



or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1.


17. The compound of any one of Claims 1 to 4 and 13 to 16, wherein L^2 is O or CH_2O , wherein the methylene portion of CH_2O is attached to Cy^2 .


18. The compound of any one of Claims 1 to 4, wherein the compound is of the Formula **VIII**:



or a pharmaceutically acceptable salt thereof.



19. The compound of any one of Claims 1 to 4 and 18, wherein the compound is of the Formula **VIIIa** or **VIIIb**:







or a pharmaceutically acceptable salt thereof.

20. The compound of any one of Claims 1 to 4, 18, and 19, wherein the compound is of the Formula **VIIIa'** or **VIIIb'**:



or a pharmaceutically acceptable salt thereof.


21. The compound of any one of Claims 1 to 4 and 18 to 20, wherein the compound is of the Formula **VIIIa''** or **VIIIb''**:



or a pharmaceutically acceptable salt thereof.


22. The compound of any one of Claims 1 to 4 and 18 to 21, wherein L² is CH₂.

23. The compound of any one of Claims 1 to 4, wherein the compound is of the Formula **IX**:




or a pharmaceutically acceptable salt thereof, wherein p is 0 or 1.

24. The compound of any one of Claims 1 to 4 and 23, wherein the compound is of the Formula **IXa**:



or a pharmaceutically acceptable salt thereof.

25. The compound of any one of Claims 1 to 4, 23, and 24, wherein the compound is of the Formula **IXa'**:



or a pharmaceutically acceptable salt thereof.

26. The compound of any one of Claims 1 to 4 and 23 to 25, wherein R¹⁰ is (C₁-C₄)alkyl; and L² is CH₂O, wherein the methylene portion of CH₂O is attached to Cy².

27. The compound of any one of Claims 1 to 7, wherein R² is (C₁-C₄)alkyl or cyclopropyl.

28. The compound of any one of Claims 1 to 27, wherein R^7 is hydrogen, OR^c , or $(C_1-C_3)alkyl$, wherein the $(C_1-C_3)alkyl$ is optionally substituted with OR^c or $-NR^dR^e$; and R^8 is hydrogen.
29. The compound of any one of Claims 1 to 28, wherein R^7 is hydrogen, $-O(C_1-C_3)alkyl$, or $(C_1-C_3)alkyl$, wherein the $(C_1-C_3)alkyl$ is optionally substituted with OH, NH_2 or $-N(C_1-C_3alkyl)_2$; and R^8 is hydrogen.
30. The compound of any one of Claims 1 to 29, wherein R^7 is hydrogen or hydroxy $(C_1-C_3)alkyl$; and R^8 is hydrogen.
31. The compound of any one of Claims 1 to 30, wherein Cy^1 is aryl or heteroaryl, each substituted with 1 to 3 groups independently selected from R^5 .
32. The compound of any one of Claims 1 to 31, wherein Cy^1 is phenyl or pyridinyl, each substituted with 1 to 3 groups independently selected from R^5 .
33. The compound of any one of Claims 1 to 32, wherein at least one R^5 is $-SO_2-(C_1-C_3)alkyl$.
34. The compound of any one of Claims 1 to 33, wherein Cy^2 is phenyl, pyrimidinyl, cyclohexyl, pyridinyl, tetrahydropyranyl, or piperidinyl, each optionally substituted with 1 to 3 groups independently selected from R^6 .
35. The compound of any one of Claims 1 to 34, wherein Cy^2 is phenyl or cyclohexyl, each optionally substituted with 1 to 3 groups independently selected from R^6 .
36. The compound of any one of Claims 1 to 35, wherein
 R^5 is selected from halogen, $-CN$, $-OR^c$, $-NR^dR^e$, $-NR^cS(O)_2R^c$, $-S(O)_2NR^dR^e$, $-C(=O)OR^c$, $-C(=O)NR^dR^e$, $-NR^cC(=O)R^c$, $-NR^cC(=O)OR^c$, $-OC(=S)NR^dR^e$, $-C(=O)R^c$, $-SO_2-(C_1-C_3)alkyl$, and $(C_1-C_4)alkyl$ optionally substituted with halogen; and
 R^6 is selected from halogen, $-CN$, $-OR^c$, $-NR^dR^e$, $-NR^cS(O)_2R^c$, $-S(O)_2NR^dR^e$, $-C(=O)OR^c$, $-OC(=O)OR^c$, $-OC(=O)R^c$, $-C(=O)NR^dR^e$, $-NR^cC(=O)R^c$, $-C(=S)NR^dR^e$, -

$\text{NR}^c\text{C}(=\text{S})\text{R}^c$, $-\text{NR}^c\text{C}(=\text{O})\text{OR}^c$, $-\text{OC}(=\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^c\text{C}(=\text{S})\text{OR}^c$, $-\text{OC}(=\text{S})\text{NR}^d\text{R}^e$, $-\text{NR}^c\text{C}(=\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^c\text{C}(=\text{S})\text{NR}^d\text{R}^e$, $-\text{C}(=\text{S})\text{R}^c$, $-\text{C}(=\text{O})\text{R}^c$, $-\text{SO}_2-(\text{C}_1-\text{C}_3)\text{alkyl}$, and $(\text{C}_1-\text{C}_4)\text{alkyl}$ optionally substituted with halogen.

37. The compound of any one of Claims 1 to 36, wherein

R^5 is $-\text{SO}_2-(\text{C}_1-\text{C}_3)\text{alkyl}$;

R^6 is selected from $-\text{CN}$, halo, $-\text{C}(=\text{O})\text{OR}^c$, OR^c , and $(\text{C}_1-\text{C}_4)\text{alkyl}$ optionally substituted with halogen; and

R^c is $(\text{C}_1-\text{C}_4)\text{alkyl}$ optionally substituted with halogen.

38. The compound of any one of Claims 1 to 37, wherein R^6 is CF_3 .

39. A pharmaceutical composition comprising a compound of any one of Claims 1 to 38, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

40. A method of treating one or more disease or disorder in a subject, comprising administering to the subject a therapeutically effective amount of a compound according to any one of Claims 1 to 38, or a pharmaceutically acceptable salt thereof.

41. The method of Claim 40, wherein the disease or disorder is selected from asthma, chronic obstructive pulmonary disease (COPD), bronchitis, allergic rhinitis, atopic dermatitis, contact dermatitis, acne, cystic fibrosis, allograft rejection, multiple sclerosis, scleroderma, arthritis, rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, systemic lupus erythematosus (SLE), psoriasis, Hashimoto's disease, pancreatitis, autoimmune diabetes, type I diabetes, autoimmune ocular disease, ulcerative colitis, Crohn's disease, regional enteritis, inflammatory bowel disease (IBD), inflammatory bowel syndrome (IBS), Sjögren's syndrome, optic neuritis, obesity, hepatosteatorrhea, adipose tissue-associated inflammation, insulin resistance, type II diabetes, neuromyelitis optica, myasthenia gravis, age related macular degeneration, dry eye, uveitis, Guillain-Barré syndrome, psoriasis, psoriatic arthritis (PsA), steroid resistant asthma, Graves' disease, scleritis, major depression, seasonal affective disorder, PTSD, bipolar disorder, autism, epilepsy, Alzheimer's, CNS disorders associated with altered sleep and/or circadian rhythms, endometriosis, obstructive sleep apnea syndrome (OSAS), Behçet's disease, dermatomyositis, polymyositis, graft versus host disease, primary biliary

cirrhosis, liver fibrosis, non-alcoholic fatty liver disease (NAFLD), sarcoidosis, primary sclerosing cholangitis, autoimmune thyroid disease, autoimmune polyendocrine syndrome type I, autoimmune polyendocrine syndrome type II, celiac disease, neuromyelitis, juvenile idiopathic arthritis, systemic sclerosis, myocardial infarction, pulmonary hypertension, osteoarthritis, cutaneous leishmaniasis, sinonasal polyposis, and cancer.

42. The method of Claim 41, wherein the disease or disorder is selected from asthma, atopic dermatitis, acne, Crohn's disease, regional enteritis, ulcerative colitis, Sjögren's syndrome, uveitis, Behçet's disease, dermatomyositis, multiple sclerosis, ankylosing spondylitis, systemic lupus erythematosus (SLE), scleroderma, psoriasis, psoriatic arthritis (PsA), steroid resistant asthma, and rheumatoid arthritis.

