

(19) **DANMARK**

(10) **DK/EP 3235821 T3**



Patent- og
Varemærkestyrelsen

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **C 07 D 491/107 (2006.01)** **A 61 K 31/407 (2006.01)** **A 61 P 29/00 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2020-04-20**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2020-04-08**
- (86) Europæisk ansøgning nr.: **17020200.6**
- (86) Europæisk indleveringsdag: **2012-07-06**
- (87) Den europæiske ansøgnings publiceringsdag: **2017-10-25**
- (30) Prioritet: **2011-07-08 EP 11005587**
- (62) Stamansøgningsnr: **12748646.2**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Grünenthal GmbH, Zieglerstrasse 6, 52078 Aachen, Tyskland**
- (72) Opfinder: **PRÜHS, Stefan, Wiesenweg 10, 69121 Heidelberg, Tyskland**
GRUSS, Michael, Hermannstraße 8, 52062 Aachen, Tyskland
Kluge, Stefan, Lörracherstraße 119, 4125 Riehen, Schweiz
- (74) Fuldmægtig i Danmark: **Orsnes Patent ApS, Forskerparken 10, 5230 Odense M, Danmark**
- (54) Benævnelse: **KRYSTALLINSK (1R,4R)-6'-FLUOR-N,N-DIMETHYL-4-PHENYL-4',9'-DIHYDRO-3'H-SPIRO[CYCLOHEXAN-1,1'-PYRAN-[3,4,B]INDOL]-4-AMIN**
- (56) Fremdragne publikationer:
WO-A1-2004/043967
WO-A1-2006/108565
WO-A2-2008/009415

DESCRIPTION

FIELD OF THE INVENTION

[0001] The invention relates to crystalline forms of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine, pharmaceutical compositions and medicaments comprising these modifications, the use of these modifications as well as to a process for the enrichment of them.

BACKGROUND OF THE INVENTION

[0002] Pharmaceutically active drugs can exist in different solid forms. For example, a drug may exist in different crystalline forms which have different physical and chemical properties.

[0003] WO 2004/043967 relates to spirocyclic cyclohexane derivatives, to a method for the production thereof, to medicaments containing said compounds and to the utilization of spirocyclic cyclohexane derivatives for the production of medicaments.


[0004] WO 2006/108565 relates to the use of spirocyclic cyclohexane derivatives for use in the production of a drug for treating substance dependency.

[0005] WO 2008/009415 relates to substituted heteroaryl derivatives, to methods for the production thereof, to medicaments containing said compounds and to the use of substituted heteroaryl derivatives for producing medicaments.

[0006] Different physical properties can cause different crystalline forms of the same drug to have largely different processing and storage performance. Such physical properties include, for example, thermodynamic stability, crystal morphology [form, shape, structure, particle size, particle size distribution, degree of crystallinity, color], ripple behavior, flowability, density, bulk density, powder density, apparent density, vibrated density, depletability, emptyability, hardness, deformability, grindability, compressability, compactability, brittleness, elasticity, caloric properties [particularly melting point], solubility [particularly equilibrium solubility, pH dependence of solubility], dissolution [particularly dissolution rate, intrinsic dissolution rate], reconstitutability, hygroscopicity, tackiness, adhesiveness, tendency to electrostatic charging, and the like.

[0007] In addition, different chemical properties can cause different crystalline forms of the same drug to have largely different performance properties. For example, a crystalline form having a low hygroscopicity (relative to other crystalline forms) can have superior chemical stability and longer shelf-life stability (cf. R. Hilfiker, Polymorphism, 2006 Wiley VCH, pp 235-242).

[0008] One particular drug that is of great interest for use in treating cancer pain (and other acute, visceral, neuropathic and chronic pain disorders) is (1*r*,4*r*)-6'-fluoro-*N,N*-dimethyl-4-phenyl-4',9'-dihydro-3'*H*-spiro[cyclohexane-1,1'-pyrano[3,4*b*]indol]-4-amine. This drug is depicted below as the compound of formula (I).



(1*r*,4*r*)-6'-fluoro-*N,N*-dimethyl-4-phenyl-4',9'-dihydro-3'*H*-spiro[cyclohexane-1,1'-pyrano[3,4-*b*]indol]-4-amine

[0009] The solid forms of (1*r*,4*r*)-6'-fluoro-*N,N*-dimethyl-4-phenyl-4',9'-dihydro-3'*H*-spiro[cyclohexane-1,1'-pyrano[3,4*b*]indol]-4-amine that are known so far are not satisfactory in every respect and there is a demand for advantageous solid forms.

[0010] In particular, there is a demand for solid forms of (1*r*,4*r*)-6'-fluoro-*N,N*-dimethyl-4-phenyl-4',9'-dihydro-3'*H*-spiro[cyclohexane-1,1'-pyrano[3,4*b*]indol]-4-amine that have properties differing from corresponding solid forms or other solid forms of the diastereomer, i.e. of (1*s*,4*s*)-6'-fluoro-*N,N*-dimethyl-4-phenyl-4',9'-dihydro-3'*H*-spiro[cyclohexane-1,1'-pyrano[3,4*b*]indol]-4-amine. Every property of one of the two diastereomers that differs from the corresponding property of the other of the two diastereomers may be useful for separating both diastereomers from one another. Isolation of pure solid forms of (1*r*,4*r*)-6'-fluoro-*N,N*-dimethyl-4-phenyl-4',9'-dihydro-3'*H*-spiro[cyclohexane-1,1'-pyrano[3,4*b*]indol]-4-amine is very desirable from a pharmaceutical point of view.

[0011] It is an object of the invention to provide forms or modifications of (1*r*,4*r*)-6'-fluoro-*N,N*-dimethyl-4-phenyl-4',9'-dihydro-3'*H*-spiro[cyclohexane-1,1'-pyrano[3,4, *b*]indol]-4-amine that have advantages compared to the forms or modifications of the prior art.

[0012] This object has been achieved by the present invention. It has surprisingly been found that different crystalline forms of (1*r*,4*r*)-6'-fluoro-*N,N*-dimethyl-4-phenyl-4',9'-dihydro-3'*H*-spiro[cyclohexane-1,1'-pyrano[3,4*b*]indol]-4-amine can be prepared which have fundamentally different properties. These inventive crystalline forms are described herein.

BRIEF DESCRIPTION OF THE DRAWINGS

[0013]

Figures 1a, 1b, 1c, 1d, 1e, 1f, 1g, 1h, 1i and 1l show the PXRD patterns of crystalline forms A, B, C, D, E, F, G, H, I and L.

Figures 2a, 2b, 2c, 2d, 2e, 2f, 2g, 2h, 2i and 2l show the Raman spectra of crystalline forms A, B, C, D, E, F, G, H, I and L.

DETAILED DESCRIPTION

[0014] The compound according to general formula (I) can systematically be referred to as "1,1-(3-dimethylamino-3-phenylpentamethylene)-6-fluoro-1,3,4,9-tetrahydropyrano[3,4-b]indole (trans)" or as "(1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine", respectively.

[0015] The compound according to general formula (I) may be present as the free base. The definition of the free base of the compound according to general formula (I) as used herein includes solvates, co-crystals and crystalline forms. For the purpose of the specification, "free base" preferably means that the compound according to general formula (I) is not present in form of a salt, particularly not in form of an acid-addition salt. The most basic functional group of the compound according to general formula (I) is its N,N-dimethylamino moiety, which thus according to the invention is preferably neither protonated nor quaternized. In other words, the free electron pair of the nitrogen atom of the N,N-dimethylamino moiety is present as a Lewis base. Methods to determine whether a chemical substance is present as the free base or as a salt are known to the skilled artisan such as ^{14}N or ^{15}N solid state NMR, x-ray diffraction, IR, Raman, XPS. ^1H -NMR recorded in solution may also be used to consider the presence of protonation.

[0016] Unless explicitly stated otherwise, all 2Θ values refer to an X-ray diffractogram measured using $\text{CuK}\alpha$ radiation having a wavelength of $1.54060 \text{ \AA}$. The terms 2Θ values and degrees 2Θ are used synonymously.

[0017] Unless explicitly stated otherwise, all values in ppm refer to ppm by weight, i.e. ppmw.

[0018] One preferred aspect of the present invention relates to a crystalline form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4b]indol]-4-amine.

[0019] In some preferred embodiments, the crystalline form according to the invention comprises an X-ray diffraction peak at about 18.9 ± 0.5 (2Θ). In some preferred embodiments, the crystalline form according to the invention comprises an X-ray diffraction peak at about 18.9 ± 0.4 (2Θ). In some preferred embodiments, the crystalline form according to the invention comprises an X-ray diffraction peak at about 18.9 ± 0.3 (2Θ). All 2Θ values refer to an X-ray

diffractogram measured using CuK α radiation having a wavelength of 1.54060 Å.

[0020] Preferably, said X-ray diffraction peak exhibits a relative intensity of at least 30%, more preferably of at least 35%, still more preferably of at least 40%, yet more preferably of at least 45%, most preferably of at least 50% and in particular, of at least 55%.

[0021] Preferably, the crystalline form according to the invention has a Raman peak at about 921 $\pm$ 5 cm⁻¹, at about 1002 $\pm$ 5 cm⁻¹ and at about 1572 $\pm$ 5 cm⁻¹.

[0022] The crystalline form according to claim 1 of the invention is an alcohol solvate.

[0023] In a reference embodiment, the crystalline form is an ansolvate.

[0024] According to some reference embodiments, the crystalline form is a solvate which is preferably selected from hydrates, solvates of dimethylsulfoxide (DMSO), or a solvate of solvent mixtures. According to the invention, the crystalline form is selected from alcohol solvates, e.g. solvates of lower alcohols such as methanol, ethanol, n-propanol, and isopropanol. Preferably, the solvate is selected from the group consisting of monosolvate, hemisolvate, disolvate, trisolvate, and mixtures thereof.

[0025] According to a reference embodiment, the crystalline form is a hydrate, preferably selected from the group consisting of monohydrate, hemi-hydrate, dihydrate, trihydrate, and mixtures thereof. In some preferred embodiments, the crystalline form is a dihydrate.

[0026] In another preferred embodiment, the crystalline form is an alcoholate, preferably selected from the group consisting of methanolate, ethanolate, propanolate (1-propanolate or 2-propanolate), and the mixtures thereof, the 2-propanolate solvate being particularly preferred.

[0027] It has been surprisingly found that some crystalline forms of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4b]indol]-4-amine disclosed herein have surprisingly higher stability than other forms, as is demonstrated in the examples. For instance, crystalline form A achieves significantly and surprisingly higher stability than other forms.

[0028] In addition, it has been surprisingly found that in some solvents the solubility of alcoholates of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano-[3,4b]indol]-4-amine may substantially differ from the solubility of the diastereomer and its solvates, respectively, namely (1s,4s)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4b]indol]-4-amine. Thus, the different solubilities may be used to separate both diastereomers from one another. For example, when the (1r,4r)-diastereomer forms an alcoholate with a comparatively low solubility and the (1s,4s)-diastereomer forms no alcoholate at all or an alcoholate with a substantially higher solubility, the (1r,4r)-diastereomer can be diastereoselectively precipitated and filtered off, thereby allowing for easy large-scale

purification of the (1r,4r)-diastereomer.

[0029] Another reference aspect of the present invention relates to a process for the production of the crystalline form according to the invention.

[0030] In a preferred embodiment, the process comprises the step of
(a-1) suspending (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in a solvent.

[0031] Conventional solvents known to persons skilled in the art may be used as solvents in a suspension of this type, such as water or organic solvents selected from the group consisting of alcohols such as methanol, ethanol, n-propanol, iso-propanol and n-butanol; esters such as ethyl acetate, n-propyl acetate, iso-propyl acetate, n-butyl acetate and iso-butyl acetate; ketones such as acetone, 2-butanone, pentan-2-one, pentan-3-one, hexan-2-one and hexan-3-one; ethers such as tert-butyl methyl ether, diethylether, tetrahydrofuran, diisopropylether and 1,4-dioxane; nitriles such as acetonitril; aromatic hydrocarbons such as toluene; saturated hydrocarbons such as n-pentane, n-hexane and n-heptane; chlorinated hydrocarbons such as dichloromethane and chloroform; and also N-methyl-2-pyrrolidone, dimethyl formamide and dimethyl sulfoxide; and mixtures thereof.

[0032] In a preferred embodiment, the solvent comprises water.

[0033] In another preferred embodiment, the solvent comprises at least one organic solvent selected from the group consisting of methanol, ethanol, n-propanol, iso-propanol, dimethyl formamide and dimethyl sulfoxid.

[0034] In still another preferred embodiment, the solvent comprises at least one organic solvent selected from the group consisting of the C4 to C6 alcohols such as n-butanol; esters such as ethyl acetate, n-propyl acetate, iso-propyl acetate, n-butyl acetate and iso-butyl acetate; ketones such as acetone, 2-butanone, pentan-2-one, pentan-3-one, hexan-2-one and hexan-3-one, ethers such as tert-butyl methyl ether, diethylether, tetrahydrofuran, diisopropylether and 1,4-dioxane; nitriles such as acetonitril; aromatic hydrocarbons such as toluene; chlorinated hydrocarbons such as dichloromethane and chloroform; and mixtures thereof. In a preferred embodiment, the solvent does neither contain water nor any solvent selected from the group consisting of methanol, ethanol, n-propanol, iso-propanol, dimethyl formamide and dimethyl sulfoxide.

[0035] Preferably, in the process, step (a-1) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80 °C, more preferably not higher than 60°C, even more preferably not higher than 40°C, and in particular in a temperature range of 15-35 °C.

[0036] Preferably, in the process, the suspension obtained in step (a-1) is stirred for a time period of at least 2 h, preferably at least 4 h, more preferably at least 8 h, still more preferably

at least 12 h, yet more preferably at least 16 h, most preferably at least 24 h, and in particular at least 2 days.

[0037] Preferably, the process further comprises the step
(b-1) separating, preferably filtering off the solid obtained in step (a-1).

[0038] Preferably, the process further comprises the step
(c-1) drying of the solid obtained in step (b-1).

[0039] Preferably, in the process, step (c-1) takes place under air. However, drying under vacuum, more preferably at a vacuum of 0 to 900 mbar, even more preferably at a vacuum of 1 to 500 mbar, and in particular at a vacuum of 10 to 200 mbar is also possible.

[0040] Preferably, in the process, step (c-1) takes place in a temperature range from 0 to 60°C, preferably from 10°C to 50°C more preferably from 20 to 40°C.

[0041] In another preferred embodiment, the process comprises the step of
(a-2) dissolving (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in a solvent.

[0042] Conventional solvents known to persons skilled in the art may be used as solvents in a suspension of this type, in particular organic solvents selected from the group consisting of alcohols such as methanol, ethanol, n-propanol, iso-propanol and n-butanol; esters such as ethyl acetate, n-propyl acetate, iso-propyl acetate, n-butyl acetate and iso-butyl acetate; ketones such as acetone, 2-butanone, pentan-2-one, pentan-3-one, hexan-2-one and hexan-3-one; ethers such as tert-butyl methyl ether, diethylether, tetrahydrofuran, diisopropylether and 1,4-dioxane; nitriles such as acetonitril; aromatic hydrocarbons such as toluene; chlorinated hydrocarbons such as dichloromethane and chloroform; and also N-methyl-2-pyrrolidone, dimethyl formamide and dimethyl sulfoxide; and mixtures thereof.

[0043] Saturated hydrocarbons, such as n-pentane, n-hexane and n-heptane, and water are less suitable, the compound (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine being only poorly soluble in these substances. However, mixtures of these substances with at least one of the solvents listed above, such as mixtures containing a saturated hydrocarbon and further at least one solvent selected from the group consisting of ketones, ethers and chlorinated hydrocarbons, may also be used. For example, n-heptane/butanone, n-heptane/dichloromethane, n-heptane/acetone, n-heptane/tetrahydrofuran, n-hexane/butanone, n-hexane/dichloromethane, n-hexane/acetone and n-hexane/tetrahydrofuran mixtures are also preferred.

[0044] Preferably, in the process, step (a-2) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80 °C, more preferably not higher than 60°C, even more preferably not higher than 40°C, and in particular in a temperature range of 20-40 °C.

[0045] In a preferred embodiment, the process further comprises the step (b-2) evaporating off the solvent of the solution obtained in step (a-2).

[0046] Suitable methods for evaporating off the solvent are known to a person skilled in the art. Preferably, in the process, the solvent is evaporated off in air, air flow, or inert gas flow, in particular argon or nitrogen flow. However, depending on the crystalline form to be obtained evaporating off the solvent under vacuum, for example by means of a rotary evaporator, is also possible.

[0047] Preferably, in the process, the solvent is evaporated off at room temperature. However, depending on the crystalline form to be obtained evaporating off the solvent at an elevated temperature, e. g. within the range of 20 °C to 60 °C, is also possible.

[0048] In another preferred embodiment, the process further comprises the step of (b-2') precipitating (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine from the solution obtained in step (a-2).

[0049] Suitable methods of precipitation are known to a person skilled in the art. In the process, step (b-2') may be carried out by reducing the volume of the solution obtained in step (a-2) and/or by cooling of the solution, preferably to a temperature of at most 15°C, more preferably at most 10°C, even more preferably at most 4-8°C and/or by cooling of the solution, preferably to a temperature of at least 10 °C, more preferably at least 30 °C, even more preferably at least 60 °C below the temperature according to step (a-2).

[0050] In a preferred embodiment, step (b-2') is carried out by the addition of a medium in which (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine is only poorly soluble ("anti-solvent") to the solution obtained in step (a-2). Said medium is preferably selected from the group consisting of saturated hydrocarbons, such as n-pentane, n-hexane and n-heptane; ethers such as tert-butyl methyl ether and diisopropyl ether; ethanol and water.

[0051] The amount of the media in which (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine is only poorly soluble, the precipitant or anti-solvent, is preferably selected in such a manner that upon its addition precipitation of the dissolved component begins. The precipitation of the dissolved component preferably begins either immediately upon the addition of the precipitant or alternatively with a delay of 2 seconds to 120 minutes. Preferably, the precipitation of the dissolved component begins within a time period of at most 60 minutes, more preferably at most 30 minutes, still more preferably at most 10 minutes, yet more preferably at most 5 minutes, most preferably at most 2 minutes, and in particular at most 30 seconds. In an especially preferred embodiment, the precipitation of the dissolved component begins immediately upon the addition of the precipitant.

[0052] Furthermore, the amount of the media in which (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-

4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine is only poorly soluble, the precipitant or anti-solvent, is preferably selected in such a manner that the dissolved component is completely or at least up to 90% of the initial amount precipitated within a time period of at most 90 minutes, more preferably at most 80 minutes, still more preferably at most 70 minutes, and most preferably at most 60 minutes after the anti-solvent has been completely added.

[0053] Preferably, in the process, after step (b-2) or respectively (b-2'), all other steps are carried out at a temperature between 40 and 0°C, preferably between 35 and 5°C, more preferably between 25 and 15°C.

[0054] Preferably, the process further comprises the step (c-2') separating, preferably filtering off the precipitate obtained in step (b-2').

[0055] Preferably, the process further comprises the step (d-2') drying of the solid obtained in step (c-2').

[0056] Preferably, in the process, step (d-2') takes place under air or inert gas flow, such as argon or nitrogen flow. However, depending on the crystalline form to be obtained evaporating off the solvent at an elevated temperature, e. g. within the range of 20 °C to 60 °C, is also possible.

[0057] In the following, any reference to a "crystalline form" refers to a crystalline form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine.

[0058] A further aspect of the present invention relates to a crystalline form A according to claim 11 of the present invention.

[0059] Preferably, the crystalline form A according to the invention has X-ray diffraction peaks at 17.6 ± 0.2 (2 Θ), 18.3 ± 0.2 (2 Θ), 18.6 ± 0.2 (2 Θ), 25.8 ± 0.2 (2 Θ) and 26.3 ± 0.2 (2 Θ). In some preferred embodiments, the crystalline form comprises an X-ray diffraction peak at about 18.3 ± 0.2 (2 Θ). In some preferred embodiments, the crystalline form comprises X-ray diffraction peaks at about 18.3 ± 0.2 (2 Θ) and about 18.6 ± 0.2 (2 Θ).

[0060] In some preferred embodiments, crystalline form A is characterized by an X-ray powder diffraction pattern comprising characteristic peaks at about 18.3 ± 0.2 degrees 2 Θ , about 18.6 ± 0.2 degrees 2 Θ and about 26.3 ± 0.2 degrees 2 Θ . In some embodiments, the X-ray powder diffraction pattern further comprises characteristic peaks at about 11.7 ± 0.2 degrees 2 Θ and about 31.6 ± 0.2 degrees 2 Θ . In some preferred embodiments, crystalline form A is characterized by an X-ray powder diffraction pattern comprising characteristic peaks at about 18.3 ± 0.2 degrees 2 Θ , about 18.6 ± 0.2 degrees 2 Θ , about 26.3 ± 0.2 degrees 2 Θ and optionally at about 17.6 ± 0.2 degrees 2 Θ and/or about 19.4 ± 0.2 degrees 2 Θ .

[0061] In some preferred embodiments, crystalline form A comprises X-ray diffraction peaks at about 17.6 ± 0.2 (2θ), about 18.3 ± 0.2 (2θ), about 18.6 ± 0.2 (2θ), about 26.3 ± 0.2 (2θ) and optionally about 25.8 ± 0.2 (2θ).

[0062] The crystalline form A according to the invention may additionally have at least one X-ray diffraction peak selected from the group consisting of about 7.8 ± 0.2 (2θ), about 13.5 ± 0.2 (2θ), about 16.8 ± 0.2 (2θ), about 19.4 ± 0.2 (2θ), about 19.7 ± 0.2 (2θ), about 27.6 ± 0.2 (2θ) and about 28.3 ± 0.2 (2θ).

[0063] Further, the crystalline form A according to the invention may be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 17.6 ± 0.2 (2θ), about 18.3 ± 0.2 (2θ), about 18.6 ± 0.2 (2θ), about 25.8 ± 0.2 (2θ) and about 26.3 ± 0.2 (2θ) and optionally one or more X-ray diffraction peaks selected from the group consisting of about 7.8 ± 0.2 (2θ), about 13.5 ± 0.2 (2θ), about 16.8 ± 0.2 (2θ), about 19.4 ± 0.2 (2θ), about 19.7 ± 0.2 (2θ), about 27.6 ± 0.2 (2θ) and about 28.3 ± 0.2 (2θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 12.2 ± 0.2 (2θ), about 20.6 ± 0.2 (2θ), about 22.3 ± 0.2 (2θ), about 22.6 ± 0.2 (2θ), about 23.5 ± 0.2 (2θ), about 23.9 ± 0.2 (2θ), about 25.0 ± 0.2 (2θ), about 29.1 ± 0.2 (2θ) and about 30.0 ± 0.2 (2θ).

[0064] The crystalline form A according to the invention may further be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 17.6 ± 0.2 (2θ), about 18.3 ± 0.2 (2θ), about 18.6 ± 0.2 (2θ), about 25.8 ± 0.2 (2θ) and about 26.3 ± 0.2 (2θ) and optionally one or more X-ray diffraction peaks selected from the group consisting of about 7.8 ± 0.2 (2θ), about 13.5 ± 0.2 (2θ), about 16.8 ± 0.2 (2θ), about 19.4 ± 0.2 (2θ), about 19.7 ± 0.2 (2θ), about 27.6 ± 0.2 (2θ) and about 28.3 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 12.2 ± 0.2 (2θ), about 20.6 ± 0.2 (2θ), about 22.3 ± 0.2 (2θ), about 22.6 ± 0.2 (2θ), about 23.5 ± 0.2 (2θ), about 23.9 ± 0.2 (2θ), about 25.0 ± 0.2 (2θ), about 29.1 ± 0.2 (2θ) and about 30.0 ± 0.2 (2θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 8.8 ± 0.2 (2θ), about 9.1 ± 0.2 (2θ), about 10.5 ± 0.2 (2θ), about 11.7 ± 0.2 (2θ), about 15.2 ± 0.2 (2θ), about 16.0 ± 0.2 (2θ), about 21.5 ± 0.2 (2θ), about 22.0 ± 0.2 (2θ), about 24.2 ± 0.2 (2θ), about 27.2 ± 0.2 (2θ), about 29.5 ± 0.2 (2θ), about 31.6 ± 0.2 (2θ), about 32.3 ± 0.2 (2θ), about 32.6 ± 0.2 (2θ), and about 33.8 ± 0.2 (2θ).

[0065] All 2θ values indicated above refer to an X-ray diffractogram measured using CuK α radiation having a wavelength of 1.54060 Å.

[0066] In DSC analyses, the crystalline form A according to the present invention preferably exhibits an endothermic event with a peak temperature at about 295-310 °C (i.e., the crystalline form has a melting endotherm at about 295-310 °C), such as for example at about 298-308 °C, or even at about 300-306 °C, or at about 302-305 °C. In some preferred embodiments, the crystalline form exhibits an endothermic event with a peak temperature at about 303-304 °C.

[0067] The crystalline form A according to the present invention may further be characterized in that it has at least a Raman peak at about $1569 \pm 2 \text{ cm}^{-1}$ and/or at least a Raman peak at about $1002 \pm 2 \text{ cm}^{-1}$. In this regard, it should be understood that all Raman peaks recited herein should be understood to additionally include values that are approximate (or about) of the recited value. For example, when it is disclosed herein that the crystalline form has a Raman peak at $1569 \pm 2 \text{ cm}^{-1}$, it should be understood that the crystalline forms has a Raman peak at about $1569 \pm 2 \text{ cm}^{-1}$.

[0068] The crystalline form A according to the present invention may further be characterized in that it has Raman peaks at $1569 \pm 2 \text{ cm}^{-1}$, $1002 \pm 2 \text{ cm}^{-1}$, $921 \pm 2 \text{ cm}^{-1}$, $1308 \pm 2 \text{ cm}^{-1}$, $1583 \pm 2 \text{ cm}^{-1}$, and $3057 \pm 2 \text{ cm}^{-1}$; and one or more Raman peaks selected from the group consisting of about $152 \pm 2 \text{ cm}^{-1}$, about $170 \pm 2 \text{ cm}^{-1}$, about $184 \pm 2 \text{ cm}^{-1}$, about $202 \pm 2 \text{ cm}^{-1}$, about $254 \pm 2 \text{ cm}^{-1}$, about $488 \pm 2 \text{ cm}^{-1}$, about $679 \pm 2 \text{ cm}^{-1}$, about $828 \pm 2 \text{ cm}^{-1}$, about $911 \pm 2 \text{ cm}^{-1}$, about $981 \pm 2 \text{ cm}^{-1}$, about $1031 \pm 2 \text{ cm}^{-1}$, about $1289 \pm 2 \text{ cm}^{-1}$, about $1453 \pm 2 \text{ cm}^{-1}$, about $1475 \pm 2 \text{ cm}^{-1}$, about $2921 \pm 2 \text{ cm}^{-1}$, about $2947 \pm 2 \text{ cm}^{-1}$, about $2960 \pm 2 \text{ cm}^{-1}$, and about $3066 \pm 2 \text{ cm}^{-1}$.

[0069] The crystalline form A according to the present invention may further be characterized in that it has one or more Raman peaks selected from the group consisting of about $365 \pm 2 \text{ cm}^{-1}$, about $420 \pm 2 \text{ cm}^{-1}$, about $519 \pm 2 \text{ cm}^{-1}$, about $544 \pm 2 \text{ cm}^{-1}$, about $609 \pm 2 \text{ cm}^{-1}$, about $620 \pm 2 \text{ cm}^{-1}$, about $636 \pm 2 \text{ cm}^{-1}$, about $694 \pm 2 \text{ cm}^{-1}$, about $714 \pm 2 \text{ cm}^{-1}$, about $785 \pm 2 \text{ cm}^{-1}$, about 8, about $872 \pm 2 \text{ cm}^{-1}$, about $943 \pm 2 \text{ cm}^{-1}$, about $1049 \pm 2 \text{ cm}^{-1}$, about $1067 \pm 2 \text{ cm}^{-1}$, about $1111 \pm 2 \text{ cm}^{-1}$, about $1128 \pm 2 \text{ cm}^{-1}$, about $1156 \pm 2 \text{ cm}^{-1}$, about $1188 \pm 2 \text{ cm}^{-1}$, about $1200 \pm 2 \text{ cm}^{-1}$, about $1235 \pm 2 \text{ cm}^{-1}$, about $1265 \pm 2 \text{ cm}^{-1}$, about $1337 \pm 2 \text{ cm}^{-1}$, about $1370 \pm 2 \text{ cm}^{-1}$, about $1405 \pm 2 \text{ cm}^{-1}$, about $1420 \pm 2 \text{ cm}^{-1}$, about $1628 \pm 2 \text{ cm}^{-1}$, about $2793 \pm 2 \text{ cm}^{-1}$, about $2851 \pm 2 \text{ cm}^{-1}$, and about $2871 \pm 2 \text{ cm}^{-1}$.

[0070] Another reference aspect of the present invention relates to a process for the production of the crystalline form A described above.

[0071] In a preferred embodiment, the process comprises the step of
(a-1) suspending (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in a solvent.

[0072] Preferably, the solvent is selected from the group consisting of the C4 to C6 alcohols such as n-butanol; esters such as ethyl acetate, n-propyl acetate, iso-propyl acetate, n-butyl acetate, and iso-butyl acetate; ketones such as acetone, 2-butanone, pentan-2-one, pentan-3-one, hexan-2-one and hexan-3-one; ethers such as tert-butyl methyl ether, diethylether, tetrahydrofuran, diisopropylether and 1,4-dioxane; nitriles such as acetonitril; aromatic hydrocarbons such as toluene; chlorinated hydrocarbons such as dichloromethane and chloroform; and mixtures thereof.

[0073] Preferably, the solvent does neither contain water nor any solvent selected from the group consisting of methanol, ethanol, n-propanol, iso-propanol, dimethyl formamide and dimethyl sulfoxid.

[0074] Preferably, in the process, step (a-1) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80 °C, more preferably not higher than 60°C, even more preferably not higher than 40°C, and in particular in a temperature range of 15-35 °C.

[0075] Preferably, the suspension obtained in step (a-1) is stirred for a time period of at least 2 h, preferably at least 4 h, more preferably at least 8 h, still more preferably at least 12 h, yet more preferably at least 16 h, most preferably at least 24 h, and in particular at least 2 days.

[0076] Preferably, the process further comprises the step (b-1) separating, preferably filtering off the solid obtained in step (a-1).

[0077] Preferably, the process further comprises the step (c-1) drying of the solid obtained in step (b-1).

[0078] Preferably, in the process, step (c-1) takes place under air. However, drying under vacuum, more preferably at a vacuum of 0 to 900 mbar, even more preferably at a vacuum of 1 to 500 mbar, and in particular at a vacuum of 10 to 200 mbar is also possible.

[0079] Preferably, in the process, step (c-1) takes place in a temperature range from 0 to 60°C, preferably from 10°C to 50°C more preferably from 20 to 40°C.

[0080] Another reference aspect of the present invention relates to a process for the production of the crystalline form A described above comprising the step of (a-2) dissolving (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in a solvent.

[0081] Preferably, the solvent is selected from the group consisting of the C4 to C6 alcohols such as n-butanol; esters such as ethyl acetate, n-propyl acetate, iso-propyl acetate, n-butyl acetate, and iso-butyl acetate; ketones such as acetone, 2-butanone, pentan-2-one, pentan-3-one, hexan-2-one and hexan-3-one; ethers such as tert-butyl methyl ether, diethylether, tetrahydrofuran, diisopropylether and 1,4-dioxane; nitriles such as acetonitril; aromatic hydrocarbons such as toluene; chlorinated hydrocarbons such as dichloromethane and chloroform; and mixtures thereof.

[0082] Mixtures of saturated hydrocarbons, such as n-pentane, n-hexane and n-heptane, further containing at least one solvent selected from the group consisting of ketones, ethers and chlorinated hydrocarbons may also be used. For instance, n-heptane/butanone, n-heptane/- dichloromethane, n-heptane/acetone, n-heptane/tetrahydrofuran, n-

hexane/butanone, n-hexane/dichloromethane n-hexane/acetone and n-hexane/tetrahydrofuran mixtures are also preferred.

[0083] Preferably, the solvent does neither contain water nor any solvent selected from the group consisting of methanol, ethanol, n-propanol, iso-propanol, dimethyl formamide and dimethyl sulfoxid.

[0084] Preferably, in the process, step (a-2) is carried out at a temperature not higher than 80 °C, more preferably not higher than 60°C, even more preferably not higher than 40°C, and in particular in a temperature range of 20-40 °C.

[0085] In a preferred embodiment, the process further comprises the step (b-2) evaporating off the solvent of the solution obtained in step (a-2).

[0086] Suitable methods for evaporating off the solvent are known to a person skilled in the art. Preferably, in the process, the solvent is evaporated off in air, air flow, or inert gas flow, in particular argon or nitrogen flow. However, evaporating off the solvent under vacuum, for example by means of a rotary evaporator, is also possible.

[0087] Preferably, in the process, the solvent is evaporated off at room temperature. However, evaporating off the solvent at an elevated temperature, e. g. within the range of 20 °C to 60 °C, is also possible.

[0088] In another preferred embodiment, the process further comprises the step (b-2') precipitating (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine from the solution obtained in step (a-2).

[0089] Suitable methods of precipitation are known to a person skilled in the art. Preferably, in the process, step (b-2') may be carried out by reducing the volume of the solution obtained according to step (a-2) and/or by cooling of the solution, preferably to a temperature of at most 15°C, more preferably at most 10°C, even more preferably at most 4-8°C and/or by cooling of the solution, preferably to a temperature of at least 10 °C, more preferably at least 30 °C, even more preferably at least 60 °C below the temperature according to step (a-2).

[0090] Preferably, in the process, after the precipitation in step (b-2'), all other steps are carried out at a temperature between 40 and 0°C, preferably between 35 and 5°C, more preferably between 25 and 15°C.

[0091] By means of the aforementioned processes any form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine, including the crystalline forms B, C, D, E, F, G, H, I and L, wherein C, D, E and F are crystalline forms according to the invention, may be converted into the crystalline form A according to the invention.

[0092] A further reference aspect of the present invention relates to a crystalline form A that is obtainable by the process as described above.

[0093] Crystal crystalline form A is thermodynamically stable up to 60 % relative humidity at room temperature. It can be obtained by suspending any other form of (1*r*,4*r*)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,*b*]indol]-4-amine at room temperature in many organic solvents, e. g. ethers such as tert-butyl methyl ether, ketones such as acetone, esters such as ethyl acetate, 1BuOH or toluene.

[0094] The thermodynamic stability is important. By using the most stable modification in a medicament it may specifically be ensured that, during storage, no crystalline conversion or polymorphic conversion of the active ingredient in the pharmaceutical formulation takes place. This is advantageous, because otherwise the properties of the medicament could change as a consequence of a conversion of a less stable modification into a more stable modification. In relation to the pharmacological properties of an administration form, this could lead for example to the solubility of the active ingredient changing, accompanied by a change in the release characteristics and thus also a change in the bioavailability. Lastly, this could result in inadequate storage stability of the medicament.

[0095] A further reference aspect of the present invention relates to a crystalline form B.

[0096] Preferably, the crystalline form B has one or more X-ray diffraction peaks selected from the group consisting of about 8.9 ± 0.2 (2θ), about 9.8 ± 0.2 (2θ), about 15.7 ± 0.2 (2θ), about 16.7 ± 0.2 (2θ), about 17.8 ± 0.2 (2θ), about 18.4 ± 0.2 (2θ), about 19.2 ± 0.2 (2θ), about 19.7 ± 0.2 (2θ), about 20.4 ± 0.2 (2θ), about 21.8 ± 0.2 (2θ), about 24.1 ± 0.2 (2θ), about 25.1 ± 0.2 (2θ), about 26.0 ± 0.2 (2θ) and about 31.1 ± 0.2 (2θ). In some preferred embodiments, the crystalline form comprises an X-ray diffraction peak at about 17.8 ± 0.2 (2θ). In some preferred embodiments, the crystalline form comprises X-ray diffraction peaks at about 9.8 ± 0.2 (2θ) and about 17.8 ± 0.2 (2θ).

[0097] In some preferred embodiments, crystalline form B comprises X-ray diffraction peaks at about 9.8 ± 0.2 (2θ), about 16.7 ± 0.2 (2θ), about 17.8 ± 0.2 (2θ), about 24.1 ± 0.2 (2θ) and optionally 19.2 ± 0.2 (2θ).

[0098] The crystalline form B may additionally have at least one X-ray diffraction peak selected from the group consisting of about 20.0 ± 0.2 (2θ), about 25.4 ± 0.2 (2θ), about 27.1 ± 0.2 (2θ), about 28.1 ± 0.2 (2θ) and about 29.2 ± 0.2 (2θ).

[0099] Further, the crystalline form B may be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 8.9 ± 0.2 (2θ), about 9.8 ± 0.2 (2θ), about 15.7 ± 0.2 (2θ), about 16.7 ± 0.2 (2θ), about 17.8 ± 0.2 (2θ), about 18.4 ± 0.2 (2θ), about 19.2 ± 0.2 (2θ), about 19.7 ± 0.2 (2θ), about 20.4 ± 0.2 (2θ), about 21.8 ± 0.2 (2θ), about 24.1 ± 0.2 (2θ), about 25.1 ± 0.2 (2θ), about 26.0 ± 0.2 (2θ) and about 31.1 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 20.0

± 0.2 (2θ), about 25.4 ± 0.2 (2θ), about 27.1 ± 0.2 (2θ), about 28.1 ± 0.2 (2θ) and about 29.2 ± 0.2 (2θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 12.0 ± 0.2 (2θ), about 16.2 ± 0.2 (2θ), about 21.4 ± 0.2 (2θ), about 22.6 ± 0.2 (2θ), about 26.7 ± 0.2 (2θ), about 27.9 ± 0.2 (2θ), about 29.7 ± 0.2 (2θ), about 30.3 ± 0.2 (2θ), about 32.7 ± 0.2 (2θ), about 32.9 ± 0.2 (2θ), about 33.5 ± 0.2 (2θ) and about 34.9 ± 0.2 (2θ).

[0100] The crystalline form B may further be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 8.9 ± 0.2 (2θ), about 9.8 ± 0.2 (2θ), about 15.7 ± 0.2 (2θ), about 16.7 ± 0.2 (2θ), about 17.8 ± 0.2 (2θ), about 18.4 ± 0.2 (2θ), about 19.2 ± 0.2 (2θ), about 19.7 ± 0.2 (2θ), about 20.4 ± 0.2 (2θ), about 21.8 ± 0.2 (2θ), about 24.1 ± 0.2 (2θ), about 25.1 ± 0.2 (2θ), about 26.0 ± 0.2 (2θ) and about 31.1 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 20.0 ± 0.2 (2θ), about 25.4 ± 0.2 (2θ), about 27.1 ± 0.2 (2θ), about 28.1 ± 0.2 (2θ) and about 29.2 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 12.0 ± 0.2 (2θ), about 21.4 ± 0.2 (2θ), about 22.6 ± 0.2 (2θ), about 26.7 ± 0.2 (2θ), about 29.7 ± 0.2 (2θ), about 30.3 ± 0.2 (2θ), about 32.7 ± 0.2 (2θ), about 32.9 ± 0.2 (2θ), about 33.5 ± 0.2 (2θ) and about 34.9 ± 0.2 (2θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 10.5 ± 0.2 (2θ), about 14.2 ± 0.2 (2θ), about 14.6 ± 0.2 (2θ), about 16.2 ± 0.2 (2θ), about 23.5 ± 0.2 (2θ), about 27.9 ± 0.2 (2θ), about 31.8 ± 0.2 (2θ), and about 33.9 ± 0.2 (2θ).

[0101] All 2θ values indicated above refer to an X-ray diffractogram measured using CuK α radiation having a wavelength of 1.54060 Å.

[0102] In DSC analyses, the crystalline form B preferably exhibits an endothermal event with a peak temperature at about 108-118 °C, more preferably at about 109-117 °C, still more preferably at about 110-116 °C, yet more preferably at about 111-115 °C and in particular at about 111-114 °C.

[0103] In DSC analyses, the crystalline form B preferably exhibits an endothermal event with a peak temperature at about 184-194 °C, more preferably at about 185-193 °C, still more preferably at about 186-192 °C, yet more preferably at about 187-191 °C and in particular at about 187-190 °C.

[0104] In DSC analyses, the crystalline form B may further exhibit an exothermal event with a peak temperature at about 202-204 °C, more preferably at about 203-213 °C, still more preferably at about 204-212 °C, yet more preferably at about 205-211 °C and in particular at about 206-210 °C.

[0105] The crystalline form B may further exhibit an endothermal event with a peak temperature at about 290-300 °C, more preferably at about 291-299 °C, still more preferably at about 292-298 °C, yet more preferably at about 293-297 °C, and in particular at about 294-297 °C.

[0106] The crystalline form B may further be characterized in that it has at least a Raman peak at about $1003 \pm 2 \text{ cm}^{-1}$ and/or at least a Raman peak at about $1571 \pm 2 \text{ cm}^{-1}$ and/or at least a Raman peak at about $1581 \pm 2 \text{ cm}^{-1}$.

[0107] The crystalline form B may further be characterized in that it has a Raman peak at about $1003 \pm 2 \text{ cm}^{-1}$ and/or at least a Raman peak at about $1571 \pm 2 \text{ cm}^{-1}$ and/or at least a Raman peak at about $1581 \pm 2 \text{ cm}^{-1}$; and/or one or more Raman peaks selected from the group consisting of about $154 \pm 2 \text{ cm}^{-1}$, about $173 \pm 2 \text{ cm}^{-1}$, about $923 \pm 2 \text{ cm}^{-1}$, about $1299 \pm 2 \text{ cm}^{-1}$, about $1476 \pm 2 \text{ cm}^{-1}$, about $3064 \pm 2 \text{ cm}^{-1}$ and about $3072 \pm 2 \text{ cm}^{-1}$; and/or one or more Raman peaks selected from the group consisting of about $217 \pm 2 \text{ cm}^{-1}$, about $259 \pm 2 \text{ cm}^{-1}$, about $370 \pm 2 \text{ cm}^{-1}$, about $492 \pm 2 \text{ cm}^{-1}$, about $683 \pm 2 \text{ cm}^{-1}$, about $825 \pm 2 \text{ cm}^{-1}$, about $1028 \pm 2 \text{ cm}^{-1}$, about $1204 \pm 2 \text{ cm}^{-1}$, about $1268 \pm 2 \text{ cm}^{-1}$, about $1374 \pm 2 \text{ cm}^{-1}$, about $1433 \pm 2 \text{ cm}^{-1}$, about $1460 \pm 2 \text{ cm}^{-1}$, about $2911 \pm 2 \text{ cm}^{-1}$, about $2950 \pm 2 \text{ cm}^{-1}$, about $2965 \pm 2 \text{ cm}^{-1}$ and about $2984 \pm 2 \text{ cm}^{-1}$.

[0108] The crystalline form B may further be characterized in that it has one or more Raman peaks selected from the group consisting of about $301 \pm 2 \text{ cm}^{-1}$, about $318 \pm 2 \text{ cm}^{-1}$, about $395 \pm 2 \text{ cm}^{-1}$, about $437 \pm 2 \text{ cm}^{-1}$, about $518 \pm 2 \text{ cm}^{-1}$, about $545 \pm 2 \text{ cm}^{-1}$, about $560 \pm 2 \text{ cm}^{-1}$, about $607 \pm 2 \text{ cm}^{-1}$, about $621 \pm 2 \text{ cm}^{-1}$, about $633 \pm 2 \text{ cm}^{-1}$, about $716 \pm 2 \text{ cm}^{-1}$, about $764 \pm 2 \text{ cm}^{-1}$, about $785 \pm 2 \text{ cm}^{-1}$, about $865 \pm 2 \text{ cm}^{-1}$, about $947 \pm 2 \text{ cm}^{-1}$, about $983 \pm 2 \text{ cm}^{-1}$, about $1039 \pm 2 \text{ cm}^{-1}$, about $1053 \pm 2 \text{ cm}^{-1}$, about $1074 \pm 2 \text{ cm}^{-1}$, about $1110 \pm 2 \text{ cm}^{-1}$, about $1119 \pm 2 \text{ cm}^{-1}$, about $1141 \pm 2 \text{ cm}^{-1}$, about $1163 \pm 2 \text{ cm}^{-1}$, about $1174 \pm 2 \text{ cm}^{-1}$, about $1191 \pm 2 \text{ cm}^{-1}$, about $1233 \pm 2 \text{ cm}^{-1}$, about $1341 \pm 2 \text{ cm}^{-1}$, about $1356 \pm 2 \text{ cm}^{-1}$, about $1630 \pm 2 \text{ cm}^{-1}$, about $2794 \pm 2 \text{ cm}^{-1}$, about $2846 \pm 2 \text{ cm}^{-1}$ and about $2879 \pm 2 \text{ cm}^{-1}$.

[0109] Another reference aspect of the present invention relates to a process for the production of the crystalline form B described above comprising the step of
(a-1) suspending (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in a solvent.

[0110] In the process for the production of the crystalline form B, the solvent preferably comprises water.

[0111] In a preferred embodiment, the solvent is water.

[0112] In another preferred embodiment, the solvent comprises water and further at least one organic solvent, preferably selected from the group consisting of the C4 to C6 alcohols such as n-butanol; esters such as ethyl acetate, n-propyl acetate, iso-propyl acetate, n-butyl acetate and iso-butyl acetate; ketones such as acetone, 2-butanone, pentan-2-one, pentan-3-one, hexan-2-one and hexan-3-one; ethers such as tert-butyl methyl ether, diethylether,

tetrahydrofuran, diisopropylether and 1,4-dioxane; THF/water mixtures are particularly preferred.

[0113] Preferably, step (a-1) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80 °C, more preferably not higher than 60°C, even more preferably not higher than 40°C, and in particular in a temperature range of 15-35 °C.

[0114] Preferably, the suspension obtained in step (a-1) is stirred for a time period of at least 2 h, preferably at least 4 h, more preferably at least 8 h, still more preferably at least 12 h, yet more preferably at least 16 h, most preferably at least 24 h, and in particular at least 2 days.

[0115] Preferably, the process further comprises the step (b-1) separating, preferably filtering off the solid obtained in step (a-1).

[0116] Preferably, the process further comprises the step (c-1) drying of the solid obtained in step (b-1).

[0117] Preferably, in the process, step (c-1) takes place under air or inert gas flow, such as argon or nitrogen flow.

[0118] Preferably, step (c-1) is performed at ambient temperature.

[0119] Another reference aspect of the present invention relates to a process for the production of the crystalline form B described above comprising the step of (a-2) dissolving (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in an organic solvent.

[0120] In some preferred embodiments, the organic solvent is selected from the group consisting of tetrahydrofuran, 1,4-dioxane and dimethylsulfoxid.

[0121] Preferably, step (a-2) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80 °C, more preferably not higher than 60°C, even more preferably not higher than 40°C, and in particular in a temperature range of 20-40 °C.

[0122] Preferably, the process further comprises the step (b-2') precipitating (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine from the solution obtained in step (a-2) by the addition of water.

[0123] The amount of the water can preferably be selected in such a manner that upon its addition precipitation of the dissolved component begins.

[0124] Preferably, the precipitation begins at most 5 minutes after the water has been added, in particular immediately upon the addition of water.

[0125] Preferably, in the process, after precipitation step (b-2') all other steps are carried out at a temperature between 40 and 0°C, preferably between 35 and 5°C, more preferably between 25 and 15°C.

[0126] Preferably, the process further comprises the step (c-2') separating, preferably filtering off the precipitate obtained in step (b-2').

[0127] Preferably, the process further comprises the step (d-2') drying of the solid obtained in step (c-2').

[0128] Preferably, in the process, step (d-2') takes place under air or inert gas flow, such as argon or nitrogen flow.

[0129] A further reference aspect of the present invention relates to a crystalline form B that is obtainable by the process as described above.

[0130] Crystal crystalline form B is thermodynamically stable at ≥ 40 % relative humidity at room temperature. It may be obtained by suspending other forms of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in water.

[0131] A further aspect of the present invention relates to a crystalline form C.

[0132] Preferably, the crystalline form C according to the invention has X-ray diffraction peaks at 9.1 ± 0.2 (2 Θ), 9.5 ± 0.2 (2 Θ), 16.8 ± 0.2 (2 Θ), 18.2 ± 0.2 (2 Θ), 18.6 ± 0.2 (2 Θ), 19.0 ± 0.2 (2 Θ), 19.3 ± 0.2 (2 Θ), 19.5 ± 0.2 (2 Θ), 22.2 ± 0.2 (2 Θ), 25.4 ± 0.2 (2 Θ), and 27.5 ± 0.2 (2 Θ). In some preferred embodiments, the crystalline form comprises an X-ray diffraction peak at about 18.2 ± 0.2 (2 Θ).

[0133] In some preferred embodiments, crystalline form C comprises X-ray diffraction peaks at about 9.1 ± 0.2 (2 Θ), about 9.5 ± 0.2 (2 Θ), about 16.8 ± 0.2 (2 Θ), about 18.2 ± 0.2 (2 Θ) and optionally 19.3 ± 0.2 (2 Θ).

[0134] The crystalline form C according to the invention may additionally have at least one X-ray diffraction peak selected from the group consisting of about 14.3 ± 0.2 (2 Θ), about 17.5 ± 0.2 (2 Θ), about 20.7 ± 0.2 (2 Θ), about 21.7 ± 0.2 (2 Θ), about 23.6 ± 0.2 (2 Θ), about 24.2 ± 0.2 (2 Θ), about 24.9 ± 0.2 (2 Θ), about 25.8 ± 0.2 (2 Θ) and about 30.3 ± 0.2 (2 Θ).

[0135] Furthermore, the crystalline form C according to the invention may be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 9.1 ± 0.2 (2 Θ), about 9.5 ± 0.2 (2 Θ), about 16.8 ± 0.2 (2 Θ), about 18.2 ± 0.2 (2 Θ), about 18.6

± 0.2 (2Θ), about 19.0 ± 0.2 (2Θ), about 19.3 ± 0.2 (2Θ), about 19.5 ± 0.2 (2Θ), about 22.2 ± 0.2 (2Θ), about 25.4 ± 0.2 (2Θ), and about 27.5 ± 0.2 (2Θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 14.3 ± 0.2 (2Θ), about 17.5 ± 0.2 (2Θ), about 20.7 ± 0.2 (2Θ), about 21.7 ± 0.2 (2Θ), about 23.6 ± 0.2 (2Θ), about 24.2 ± 0.2 (2Θ), about 24.9 ± 0.2 (2Θ), about 25.8 ± 0.2 (2Θ) and about 30.3 ± 0.2 (2Θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 14.8 ± 0.2 (2Θ), about 22.5 ± 0.2 (2Θ), about 26.2 ± 0.2 (2Θ), about 26.5 ± 0.2 (2Θ), about 28.1 ± 0.2 (2Θ), about 28.7 ± 0.2 (2Θ), about 29.4 ± 0.2 (2Θ), about 32.3 ± 0.2 (2Θ) and about 33.6 ± 0.2 (2Θ).

[0136] The crystalline form C according to the invention may further be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 9.1 ± 0.2 (2Θ), about 9.5 ± 0.2 (2Θ), about 16.8 ± 0.2 (2Θ), about 18.2 ± 0.2 (2Θ), about 18.6 ± 0.2 (2Θ), about 19.0 ± 0.2 (2Θ), about 19.3 ± 0.2 (2Θ), about 19.5 ± 0.2 (2Θ), about 22.2 ± 0.2 (2Θ), about 25.4 ± 0.2 (2Θ), and about 27.5 ± 0.2 (2Θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 14.3 ± 0.2 (2Θ), about 17.5 ± 0.2 (2Θ), about 20.7 ± 0.2 (2Θ), about 21.7 ± 0.2 (2Θ), about 23.6 ± 0.2 (2Θ), about 24.2 ± 0.2 (2Θ), about 24.9 ± 0.2 (2Θ), about 25.8 ± 0.2 (2Θ) and about 30.3 ± 0.2 (2Θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 14.8 ± 0.2 (2Θ), about 22.5 ± 0.2 (2Θ), about 26.2 ± 0.2 (2Θ), about 26.5 ± 0.2 (2Θ), about 28.1 ± 0.2 (2Θ), about 28.7 ± 0.2 (2Θ), about 29.4 ± 0.2 (2Θ), about 32.3 ± 0.2 (2Θ) and about 33.6 ± 0.2 (2Θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 7.8 ± 0.2 (2Θ), about 10.4 ± 0.2 (2Θ), about 11.1 ± 0.2 (2Θ), about 12.2 ± 0.2 (2Θ), about 13.5 ± 0.2 (2Θ), about 15.3 ± 0.2 (2Θ), about 16.1 ± 0.2 (2Θ) and about 34.5 ± 0.2 (2Θ).

[0137] All 20 values indicated above refer to an X-ray diffractogram measured using $\text{CuK}\alpha$ radiation having a wavelength of $1.54060 \text{ \AA}$.

[0138] In DSC analyses, the crystalline form C according to the present invention preferably exhibits an endothermal event with a peak temperature at about $130\text{--}140^\circ\text{C}$, more preferably at about $131\text{--}139^\circ\text{C}$, still more preferably at about $132\text{--}138^\circ\text{C}$, yet more preferably at about $133\text{--}137^\circ\text{C}$ and in particular at about $133\text{--}136^\circ\text{C}$.

[0139] In DSC analyses, the crystalline form C according to the present invention may further exhibit an endothermal event with a peak temperature at about $110\text{--}120^\circ\text{C}$, more preferably at about $111\text{--}119^\circ\text{C}$, still more preferably at about $112\text{--}118^\circ\text{C}$, yet more preferably at about $113\text{--}117^\circ\text{C}$ and in particular at about $113\text{--}116^\circ\text{C}$.

[0140] The crystalline form C according to the present invention may further exhibit an endothermal event with a peak temperature at about $243\text{--}253^\circ\text{C}$, more preferably at about $244\text{--}252^\circ\text{C}$, still more preferably at about $245\text{--}251^\circ\text{C}$, yet more preferably at about $246\text{--}250^\circ\text{C}$ and in particular at about $246\text{--}249^\circ\text{C}$.

[0141] The crystalline form C according to the present invention may further exhibit an endothermal event with a peak temperature at about $292\text{--}302^\circ\text{C}$, more preferably at about

293-301 °C, still more preferably at about 294-300 °C, and in particular at about 295-299 °C.

[0142] The crystalline form C according to the present invention may further be characterized in that it has at least a Raman peak at about $1003 \pm 2 \text{ cm}^{-1}$ and/or at least a Raman peak at about $1570 \pm 2 \text{ cm}^{-1}$ and/or at least a Raman peak at about $1587 \pm 2 \text{ cm}^{-1}$.

[0143] The crystalline form C according to the present invention may further be characterized in that it has Raman peaks at $1003 \pm 2 \text{ cm}^{-1}$, $1570 \pm 2 \text{ cm}^{-1}$, $1587 \pm 2 \text{ cm}^{-1}$, $156 \pm 2 \text{ cm}^{-1}$, $171 \pm 2 \text{ cm}^{-1}$, $183 \pm 2 \text{ cm}^{-1}$, $922 \pm 2 \text{ cm}^{-1}$, $1299 \pm 2 \text{ cm}^{-1}$, $1478 \pm 2 \text{ cm}^{-1}$, $2932 \pm 2 \text{ cm}^{-1}$, $2951 \pm 2 \text{ cm}^{-1}$ and $3070 \pm 2 \text{ cm}^{-1}$; and one or more Raman peaks selected from the group consisting of about $210 \pm 2 \text{ cm}^{-1}$, about $253 \pm 2 \text{ cm}^{-1}$, about $491 \pm 2 \text{ cm}^{-1}$, about $682 \pm 2 \text{ cm}^{-1}$, about $829 \pm 2 \text{ cm}^{-1}$, about $913 \pm 2 \text{ cm}^{-1}$, about $1028 \pm 2 \text{ cm}^{-1}$, about $1203 \pm 2 \text{ cm}^{-1}$, about $1373 \pm 2 \text{ cm}^{-1}$, about $1435 \pm 2 \text{ cm}^{-1}$, about $1462 \pm 2 \text{ cm}^{-1}$, about $2845 \pm 2 \text{ cm}^{-1}$, about $2856 \pm 2 \text{ cm}^{-1}$, about $2890 \pm 2 \text{ cm}^{-1}$, about $2977 \pm 2 \text{ cm}^{-1}$ and about $2990 \pm 2 \text{ cm}^{-1}$.

[0144] The crystalline form C according to the present invention may further be characterized in that it has one or more Raman peaks selected from the group consisting of about $371 \pm 2 \text{ cm}^{-1}$, about $394 \pm 2 \text{ cm}^{-1}$, about $432 \pm 2 \text{ cm}^{-1}$, about $520 \pm 2 \text{ cm}^{-1}$, about $542 \pm 2 \text{ cm}^{-1}$, about $560 \pm 2 \text{ cm}^{-1}$, about $608 \pm 2 \text{ cm}^{-1}$, about $621 \pm 2 \text{ cm}^{-1}$, about $633 \pm 2 \text{ cm}^{-1}$, about $712 \pm 2 \text{ cm}^{-1}$, about $786 \pm 2 \text{ cm}^{-1}$, about $885 \pm 2 \text{ cm}^{-1}$, about $948 \pm 2 \text{ cm}^{-1}$, about $983 \pm 2 \text{ cm}^{-1}$, about $1051 \pm 2 \text{ cm}^{-1}$, about $1077 \pm 2 \text{ cm}^{-1}$, about $1111 \pm 2 \text{ cm}^{-1}$, about $1119 \pm 2 \text{ cm}^{-1}$, about $1157 \pm 2 \text{ cm}^{-1}$, about $1189 \pm 2 \text{ cm}^{-1}$, $1231 \pm 2 \text{ cm}^{-1}$, about $1265 \pm 2 \text{ cm}^{-1}$, about $1339 \pm 2 \text{ cm}^{-1}$, about $1630 \pm 2 \text{ cm}^{-1}$ and about $2794 \pm 2 \text{ cm}^{-1}$.

[0145] Another reference aspect of the present invention relates to a process for the production of the crystalline form C described above comprising the step of
(a-1) suspending (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in a solvent comprising ethanol.

[0146] Preferably, the solvent is ethanol.

[0147] Preferably, step (a-1) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80 °C, more preferably not higher than 60°C, even more preferably not higher than 40°C, and in particular in a temperature range of 15-35 °C.

[0148] Preferably, the suspension obtained in step (a-1) is stirred for a time period of at least 2 h, preferably at least 4 h, more preferably at least 8 h, still more preferably at least 12 h, yet more preferably at least 16 h, most preferably at least 24 h, and in particular at least 2 days.

[0149] Preferably, the process further comprises the step (b-1) separating, preferably filtering off the solid obtained in step (a-1).

[0150] Preferably, the process further comprises the step (c-1) drying of the solid obtained in step (b-1).

Preferably, in the process, step (c-1) takes place under air or inert gas flow, such as argon or nitrogen flow. Preferably, step (c-1) is performed at ambient temperature.

[0151] Another reference aspect of the present invention relates to a process for the production of the crystalline form C described above comprising the step of

(a-2) dissolving (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in an organic solvent.

[0152] In some preferred embodiments, the organic solvent is selected from the group consisting of acetone, 2-butanone, dichloromethane, tetrahydrofuran, 1,4-dioxane and dimethyl sulfoxid.

[0153] Preferably, step (a-2) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80 °C, more preferably not higher than 60°C, even more preferably not higher than 40°C, and in particular in a temperature range of 20-40 °C.

[0154] Preferably the process further comprises the step

(b-2') precipitating (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine from the solution obtained in step (a-2) by the addition of ethanol.

[0155] The amount of the ethanol can preferably be selected in such a manner that upon its addition precipitation of the dissolved component begins.

[0156] Preferably, the precipitation starts at most 90 minutes, more preferably at most 75 minutes, most preferably at most 60 minutes, after the ethanol has been added.

[0157] Preferably, in the process, after step (b-2'), all other steps are carried out at a temperature between 40 and 0°C, preferably between 35 and 5°C, more preferably between 25 and 15°C.

[0158] Preferably, the process further comprises the step

(c-2') separating, preferably filtering off the precipitate obtained in step (b-2').

[0159] Preferably, the process further comprises the step

(d-2') drying of the solid obtained in step (c-2').

[0160] Preferably, in the process, step (d-2') takes place under air or inert gas flow, such as

argon or nitrogen flow.

[0161] A further reference aspect of the present invention relates to a crystalline form C that is obtainable by the process as described above.

[0162] A further aspect of the present invention relates to a crystalline form D.

[0163] Preferably, the crystalline form D according to the invention has X-ray diffraction peaks at 8.4 ± 0.2 (2θ), 8.8 ± 0.2 (2θ), 15.0 ± 0.2 (2θ), 15.2 ± 0.2 (2θ), 17.0 ± 0.2 (2θ), 17.6 ± 0.2 (2θ), 18.9 ± 0.2 (2θ), 21.2 ± 0.2 (2θ), 22.4 ± 0.2 (2θ), 23.2 ± 0.2 (2θ), 26.0 ± 0.2 (2θ), 29.5 ± 0.2 (2θ), and 30.7 ± 0.2 (2θ). In some preferred embodiments, the crystalline form comprises an X-ray diffraction peak at about 17.6 ± 0.2 (2θ).

[0164] In some preferred embodiments, crystalline form D comprises X-ray diffraction peaks at about 8.4 ± 0.2 (2θ), about 8.8 ± 0.2 (2θ), about 17.6 ± 0.2 (2θ), about 22.5 ± 0.2 (2θ) and optionally 15.0 ± 0.2 (2θ) and/or 15.2 ± 0.2 (2θ).

[0165] The crystalline form D according to the invention may additionally have at least one X-ray diffraction peak selected from the group consisting of about 18.1 ± 0.2 (2θ), about 20.9 ± 0.2 (2θ), about 21.6 ± 0.2 (2θ), about 22.8 ± 0.2 (2θ), about 24.9 ± 0.2 (2θ), about 25.7 ± 0.2 (2θ) and about 30.3 ± 0.2 (2θ).

[0166] Furthermore, the crystalline form D according to the invention may be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 8.4 ± 0.2 (2θ), about 8.8 ± 0.2 (2θ), about 15.0 ± 0.2 (2θ), about 15.2 ± 0.2 (2θ), about 17.0 ± 0.2 (2θ), about 17.6 ± 0.2 (2θ), about 18.9 ± 0.2 (2θ), about 21.2 ± 0.2 (2θ), about 22.4 ± 0.2 (2θ), about 23.2 ± 0.2 (2θ), about 26.0 ± 0.2 (2θ), about 29.5 ± 0.2 (2θ) and about 30.7 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 18.1 ± 0.2 (2θ), about 20.9 ± 0.2 (2θ), about 21.6 ± 0.2 (2θ), about 22.8 ± 0.2 (2θ), about 24.9 ± 0.2 (2θ), about 25.7 ± 0.2 (2θ) and about 30.3 ± 0.2 (2θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 19.4 ± 0.2 (2θ), about 19.8 ± 0.2 (2θ), about 25.2 ± 0.2 (2θ), about 26.6 ± 0.2 (2θ), about 27.5 ± 0.2 (2θ), about 28.0 ± 0.2 (2θ), about 28.5 ± 0.2 (2θ), about 31.3 ± 0.2 (2θ), about 31.9 ± 0.2 (2θ), about 32.2 ± 0.2 (2θ), about 32.8 ± 0.2 (2θ), about 34.0 ± 0.2 (2θ) and about 34.9 ± 0.2 (2θ).

[0167] The crystalline form D according to the invention may further be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 8.43 ± 0.2 (2θ), about 8.77 ± 0.2 (2θ), about 15.0 ± 0.2 (2θ), about 15.2 ± 0.2 (2θ), about 17.0 ± 0.2 (2θ), about 17.6 ± 0.2 (2θ), about 18.9 ± 0.2 (2θ), about 21.2 ± 0.2 (2θ), about 22.4 ± 0.2 (2θ), about 23.2 ± 0.2 (2θ), about 26.0 ± 0.2 (2θ), about 29.5 ± 0.2 (2θ) and about 30.7 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 18.1 ± 0.2 (2θ), about 20.9 ± 0.2 (2θ), about 21.6 ± 0.2 (2θ), about 22.8 ± 0.2 (2θ), about 24.9 ± 0.2 (2θ), about 25.7 ± 0.2 (2θ) and about 30.3 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 19.4 ± 0.2 (2θ), about 19.8 ± 0.2

(2Θ), about 25.2 ± 0.2 (2Θ), about 26.6 ± 0.2 (2Θ), about 27.5 ± 0.2 (2Θ), about 28.0 ± 0.2 (2Θ), about 28.5 ± 0.2 (2Θ), about 31.3 ± 0.2 (2Θ), about 31.9 ± 0.2 (2Θ), about 32.2 ± 0.2 (2Θ), about 32.8 ± 0.2 (2Θ), about 34.0 ± 0.2 (2Θ) and about 34.9 ± 0.2 (2Θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 11.19 ± 0.2 (2Θ), about 12.05 ± 0.2 (2Θ), about 13.65 ± 0.2 (2Θ), about 16.13 ± 0.2 (2Θ) and about 33.55 ± 0.2 (2Θ).

[0168] All 2Θ values indicated above refer to an X-ray diffractogram measured using CuK α radiation having a wavelength of 1.54060 Å.

[0169] In DSC analyses, the crystalline form D according to the present invention preferably exhibits an endothermal event with a peak temperature at about 107-117 °C, more preferably at about 108-116 °C, still more preferably at about 109-115 °C, yet more preferably at about 110-114 °C and in particular at about 110-113 °C.

[0170] Preferably, in DSC analyses, the crystalline form D according to the present invention further exhibits an endothermal event with a peak temperature at about 118-128 °C, more preferably at about 119-127 °C, still more preferably at about 120-126 °C, yet more preferably at about 121-125 °C and in particular at about 122-125 °C.

[0171] The crystalline form D according to the present invention may further be characterized in that it has one or more Raman peaks selected from the group consisting of about 169 ± 2 cm^{-1} , about 922 cm^{-1} , about $1002 \pm 2 \text{ cm}^{-1}$, about $1570 \pm 2 \text{ cm}^{-1}$, about $2957 \pm 2 \text{ cm}^{-1}$ and about $3067 \pm 2 \text{ cm}^{-1}$.

[0172] The crystalline form D according to the present invention may further be characterized in that it has Raman peaks at $169 \pm 2 \text{ cm}^{-1}$, 922 cm^{-1} , $1002 \pm 2 \text{ cm}^{-1}$, $1570 \pm 2 \text{ cm}^{-1}$, $2957 \pm 2 \text{ cm}^{-1}$, $3067 \pm 2 \text{ cm}^{-1}$, $254 \pm 2 \text{ cm}^{-1}$, $367 \pm 2 \text{ cm}^{-1}$, $491 \pm 2 \text{ cm}^{-1}$, $683 \pm 2 \text{ cm}^{-1}$, $1302 \pm 2 \text{ cm}^{-1}$, $1437 \pm 2 \text{ cm}^{-1}$, $1479 \pm 2 \text{ cm}^{-1}$ and $2935 \pm 2 \text{ cm}^{-1}$; and one or more Raman peaks selected from the group consisting of about $633 \pm 2 \text{ cm}^{-1}$, about $786 \pm 2 \text{ cm}^{-1}$, about $821 \pm 2 \text{ cm}^{-1}$, about $1028 \pm 2 \text{ cm}^{-1}$, about $1117 \pm 2 \text{ cm}^{-1}$, about $1158 \pm 2 \text{ cm}^{-1}$, about $1202 \pm 2 \text{ cm}^{-1}$, about $1264 \pm 2 \text{ cm}^{-1}$, and about $1377 \pm 2 \text{ cm}^{-1}$.

[0173] Another reference aspect of the present invention relates to a process for the production of the crystalline form D described above comprising the step of

(a-1) suspending (1*r*,4*r*)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,*b*]indol]-4-amine in a solvent comprising isopropanol.

[0174] Preferably, the solvent is isopropanol.

[0175] Preferably, step (a-1) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80 °C, more preferably not higher than 60°C, even more preferably not higher than 40°C, and in particular in a

temperature range of 15-35 °C.

[0176] Preferably, the suspension obtained in step (a-1) is stirred for a time period of at least 4 h, more preferably at least 8 h, still more preferably at least 12 h, yet more preferably at least 16 h, most preferably at least 24 h, and in particular at least 2 days.

[0177] Preferably, the process further comprises the step (b-1) separating, preferably filtering off the solid obtained in step (a-1).

[0178] Preferably, the process further comprises the step (c-1) drying of the solid obtained in step (b-1).

[0179] Preferably, in the process, step (c-1) takes place under air or inert gas flow, such as argon or nitrogen flow. Preferably, step (c-1) is performed at ambient temperature.

[0180] A further reference aspect of the present invention relates to a crystalline form D that is obtainable by the process as described above.

[0181] A further aspect of the present invention relates to a crystalline form E.

[0182] Preferably, the crystalline form E according to the invention has X-ray diffraction peaks at 8.8 ± 0.2 (2Θ), 11.9 ± 0.2 (2Θ), 17.0 ± 0.2 (2Θ), 17.7 ± 0.2 (2Θ) and 18.7 ± 0.2 (2Θ). In some preferred embodiments, the crystalline form comprises an X-ray diffraction peak at about 18.7 ± 0.2 (2Θ).

[0183] In some preferred embodiments, crystalline form E comprises X-ray diffraction peaks at about 8.8 ± 0.2 (2Θ), about 17.0 ± 0.2 (2Θ), about 17.7 ± 0.2 (2Θ), about 18.7 ± 0.2 (2Θ) and optionally 11.9 ± 0.2 (2Θ).

[0184] The crystalline form E according to the invention may additionally have at least one X-ray diffraction peak selected from the group consisting of about 22.6 ± 0.2 (2Θ), about 23.3 ± 0.2 (2Θ), about 25.7 ± 0.2 (2Θ), about 26.1 ± 0.2 (2Θ), about 26.9 ± 0.2 (2Θ), about 27.6 ± 0.2 (2Θ) and about 30.3 ± 0.2 (2Θ).

[0185] Furthermore, the crystalline form E according to the invention may be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 8.8 ± 0.2 (2Θ), about 11.9 ± 0.2 (2Θ), about 17.0 ± 0.2 (2Θ), about 17.7 ± 0.2 (2Θ) and about 18.7 ± 0.2 (2Θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 22.6 ± 0.2 (2Θ), about 23.3 ± 0.2 (2Θ), about 25.7 ± 0.2 (2Θ), about 26.1 ± 0.2 (2Θ), about 26.9 ± 0.2 (2Θ), about 27.6 ± 0.2 (2Θ) and about 30.3 ± 0.2 (2Θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 16.3 ± 0.2 (2Θ), about 18.2 ± 0.2 (2Θ), about 20.4 ± 0.2 (2Θ), about 23.9 ± 0.2 (2Θ), about 24.1 ± 0.2 (2Θ), about 26.6 ± 0.2 (2Θ), about 27.8 ± 0.2 (2Θ), about 28.2 ± 0.2 (2Θ), about 29.4 ± 0.2 (2Θ), about 30.8 ± 0.2 (2Θ), about 31.2 ± 0.2 (2Θ) and about 33.0 ± 0.2 (2Θ).

[0186] The crystalline form E according to the invention may further be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 8.8 ± 0.2 (2Θ), about 11.9 ± 0.2 (2Θ), about 17.0 ± 0.2 (2Θ), about 17.7 ± 0.2 (2Θ), and about 18.7 ± 0.2 (2Θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 22.6 ± 0.2 (2Θ), about 23.3 ± 0.2 (2Θ), about 25.7 ± 0.2 (2Θ), about 26.1 ± 0.2 (2Θ), about 26.9 ± 0.2 (2Θ), about 27.6 ± 0.2 (2Θ) and about 30.3 ± 0.2 (2Θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 16.3 ± 0.2 (2Θ), about 18.2 ± 0.2 (2Θ), about 20.4 ± 0.2 (2Θ), about 23.9 ± 0.2 (2Θ), about 24.1 ± 0.2 (2Θ), about 26.6 ± 0.2 (2Θ), about 27.8 ± 0.2 (2Θ), about 28.2 ± 0.2 (2Θ), about 29.4 ± 0.2 (2Θ), about 30.8 ± 0.2 (2Θ), about 31.2 ± 0.2 (2Θ) and about 33.0 ± 0.2 (2Θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 10.7 ± 0.2 (2Θ), about 11.3 ± 0.2 (2Θ), about 12.2 ± 0.2 (2Θ), about 13.8 ± 0.2 (2Θ), about 15.2 ± 0.2 (2Θ), about 15.8 ± 0.2 (2Θ), about 19.4 ± 0.2 (2Θ), about 19.7 ± 0.2 (2Θ), about 21.1 ± 0.2 (2Θ), about 21.9 ± 0.2 (2Θ), about 24.7 ± 0.2 (2Θ), about 25.0 ± 0.2 (2Θ), about 28.7 ± 0.2 (2Θ), about 31.5 ± 0.2 (2Θ) and about 34.4 ± 0.2 (2Θ).

[0187] All 2Θ values indicated above refer to an X-ray diffractogram measured using $\text{CuK}\alpha$ radiation having a wavelength of $1.54060 \text{ \AA}$.

[0188] The crystalline form E according to the present invention may further be characterized in that it has one or more Raman peaks selected from the group consisting of about $1003 \pm 2 \text{ cm}^{-1}$, about $1297 \pm 2 \text{ cm}^{-1}$, about $1570 \pm 2 \text{ cm}^{-1}$ and about $1585 \pm 2 \text{ cm}^{-1}$.

[0189] The crystalline form E according to the present invention may further be characterized in that it has one or more Raman peaks at $1003 \pm 2 \text{ cm}^{-1}$, $1297 \pm 2 \text{ cm}^{-1}$, $1570 \pm 2 \text{ cm}^{-1}$, $1585 \pm 2 \text{ cm}^{-1}$, $159 \pm 2 \text{ cm}^{-1}$, $188 \pm 2 \text{ cm}^{-1}$, $680 \pm 2 \text{ cm}^{-1}$, $923 \pm 2 \text{ cm}^{-1}$, $1434 \pm 2 \text{ cm}^{-1}$, $1461 \pm 2 \text{ cm}^{-1}$, $2943 \pm 2 \text{ cm}^{-1}$, $2961 \pm 2 \text{ cm}^{-1}$, $3070 \pm 2 \text{ cm}^{-1}$; and one or more Raman peaks selected from the group consisting of about $174 \pm 2 \text{ cm}^{-1}$, about $257 \pm 2 \text{ cm}^{-1}$, about $370 \pm 2 \text{ cm}^{-1}$, about $489 \pm 2 \text{ cm}^{-1}$, about $632 \pm 2 \text{ cm}^{-1}$, about $823 \pm 2 \text{ cm}^{-1}$, about $913 \pm 2 \text{ cm}^{-1}$, about $982 \pm 2 \text{ cm}^{-1}$, about $1027 \pm 2 \text{ cm}^{-1}$, about $1037 \pm 2 \text{ cm}^{-1}$, about $1169 \pm 2 \text{ cm}^{-1}$, about $1192 \pm 2 \text{ cm}^{-1}$, about $1202 \pm 2 \text{ cm}^{-1}$, about $1262 \pm 2 \text{ cm}^{-1}$, about $1476 \pm 2 \text{ cm}^{-1}$, about $2836 \pm 2 \text{ cm}^{-1}$, about $2860 \pm 2 \text{ cm}^{-1}$, about $2894 \pm 2 \text{ cm}^{-1}$, about $2994 \pm 2 \text{ cm}^{-1}$ and about $3057 \pm 2 \text{ cm}^{-1}$.

[0190] The crystalline form E according to the present invention may further be characterized in that it has one or more Raman peaks selected from the group consisting of about $271 \pm 2 \text{ cm}^{-1}$, about $315 \pm 2 \text{ cm}^{-1}$, about $394 \pm 2 \text{ cm}^{-1}$, about $423 \pm 2 \text{ cm}^{-1}$, about $434 \pm 2 \text{ cm}^{-1}$, about $518 \pm 2 \text{ cm}^{-1}$, about $541 \pm 2 \text{ cm}^{-1}$, about $557 \pm 2 \text{ cm}^{-1}$, about $604 \pm 2 \text{ cm}^{-1}$, about $621 \pm 2 \text{ cm}^{-1}$, about $710 \pm 2 \text{ cm}^{-1}$, about $760 \pm 2 \text{ cm}^{-1}$, about $784 \pm 2 \text{ cm}^{-1}$, about $870 \pm 2 \text{ cm}^{-1}$, about $945 \pm 2 \text{ cm}^{-1}$, about $1049 \pm 2 \text{ cm}^{-1}$, about $1075 \pm 2 \text{ cm}^{-1}$, about $1117 \pm 2 \text{ cm}^{-1}$, about $1135 \pm 2 \text{ cm}^{-1}$, about $1230 \pm 2 \text{ cm}^{-1}$, about $1337 \pm 2 \text{ cm}^{-1}$, about $1354 \pm 2 \text{ cm}^{-1}$, about $1376 \pm 2 \text{ cm}^{-1}$, about $1629 \pm 2 \text{ cm}^{-1}$

and about $2791 \pm 2 \text{ cm}^{-1}$.

[0191] Another reference aspect of the present invention relates to a process for the production of the crystalline form E described above comprising the step of

(a-1) suspending (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in a solvent comprising methanol.

[0192] Preferably, the solvent is methanol.

[0193] Preferably, step (a-1) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80°C , more preferably not higher than 60°C , even more preferably not higher than 40°C , and in particular in a temperature range of $15\text{--}35^\circ\text{C}$.

[0194] Preferably, the suspension obtained in step (a-1) is stirred for a time period of at least 5 minutes, preferably at least 10 minutes, more preferably at least 15 minutes, still more preferably at least 30 minutes, yet more preferably at least 1 h, and most preferably at least 2 h.

[0195] Preferably, the process further comprises the step
(b-1) separating, preferably filtering off the solid obtained in step (a-1).

[0196] Preferably, the process further comprises the step
(c-1) drying of the solid obtained in step (b-1).

[0197] Preferably, in the process, step (c-1) takes place under air or inert gas flow, such as argon or nitrogen flow. Preferably, step (c-1) is performed at ambient temperature.

[0198] Another reference aspect of the present invention relates to a process for the production of the crystalline form E described above comprising the step of

(a-3) washing (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine with methanol.

[0199] For the purpose of the specification, "washing (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine with methanol" is to be regarded as being synonymous with "bringing (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine into contact with an excess of methanol".

[0200] Preferably, the (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine is washed (brought into contact) with at least $0.5 \mu\text{L}$, more preferably $1.0 \mu\text{L}$, still more preferably at least $2 \mu\text{L}$, yet more preferably at least $3 \mu\text{L}$, most preferably at least $4 \mu\text{L}$, and in particular at least $5 \mu\text{L}$ of methanol per mg of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-

pyrano[3,4,b]indol]-4-amine.

[0201] Preferably, the (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine employed in step (a-2) is in the form of a hydrate or solvate.

[0202] A further reference aspect of the present invention relates to a crystalline form E that is obtainable by the process as described above.

[0203] A further aspect of the present invention relates to a crystalline form F.

[0204] Preferably, the crystalline form F according to the invention has X-ray diffraction peaks at 9.0 ± 0.2 (2θ), 15.4 ± 0.2 (2θ), 16.1 ± 0.2 (2θ), 17.9 ± 0.2 (2θ), 18.2 ± 0.2 (2θ), 18.7 ± 0.2 (2θ), 19.4 ± 0.2 (2θ), 20.1 ± 0.2 (2θ), 20.6 ± 0.2 (2θ), 21.8 ± 0.2 (2θ), 24.6 ± 0.2 (2θ), 25.6 ± 0.2 (2θ), 27.1 ± 0.2 (2θ), 27.4 ± 0.2 (2θ) and 29.3 ± 0.2 (2θ). In some preferred embodiments, the crystalline form comprises an X-ray diffraction peak at about 20.1 ± 0.2 (2θ).

[0205] In some preferred embodiments, crystalline form F comprises X-ray diffraction peaks at about 9.0 ± 0.2 (2θ), about 17.9 ± 0.2 (2θ), about 18.7 ± 0.2 (2θ), about 20.1 ± 0.2 (2θ) and optionally 16.1 ± 0.2 (2θ).

[0206] The crystalline form F according to the invention may additionally have at least one X-ray diffraction peak selected from the group consisting of about 21.9 ± 0.2 (2θ), about 25.0 ± 0.2 (2θ), about 27.9 ± 0.2 (2θ) and about 30.3 ± 0.2 (2θ).

[0207] Furthermore, the crystalline form F according to the invention may be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 9.0 ± 0.2 (2θ), about 15.4 ± 0.2 (2θ), about 16.1 ± 0.2 (2θ), about 17.9 ± 0.2 (2θ), about 18.2 ± 0.2 (2θ), about 18.7 ± 0.2 (2θ), about 19.4 ± 0.2 (2θ), about 20.1 ± 0.2 (2θ), about 20.6 ± 0.2 (2θ), about 21.8 ± 0.2 (2θ), about 24.6 ± 0.2 (2θ), about 25.6 ± 0.2 (2θ), about 27.1 ± 0.2 (2θ), about 27.4 ± 0.2 (2θ) and about 29.3 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 21.9 ± 0.2 (2θ), about 25.0 ± 0.2 (2θ), about 27.9 ± 0.2 (2θ) and about 30.3 ± 0.2 (2θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 14.5 ± 0.2 (2θ), about 22.9 ± 0.2 (2θ), about 23.5 ± 0.2 (2θ), about 30.0 ± 0.2 (2θ), about 30.8 ± 0.2 (2θ), about 31.4 ± 0.2 (2θ), about 31.6 ± 0.2 (2θ) and about 32.2 ± 0.2 (2θ).

[0208] The crystalline form F according to the invention may further be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 9.0 ± 0.2 (2θ), about 15.4 ± 0.2 (2θ), about 16.1 ± 0.2 (2θ), about 17.9 ± 0.2 (2θ), about 18.2 ± 0.2 (2θ), about 18.7 ± 0.2 (2θ), about 19.4 ± 0.2 (2θ), about 20.1 ± 0.2 (2θ), about 20.6 ± 0.2 (2θ), about 21.8 ± 0.2 (2θ), about 24.6 ± 0.2 (2θ), about 25.6 ± 0.2 (2θ), about 27.1 ± 0.2 (2θ), about 27.4 ± 0.2 (2θ) and about 29.3 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 21.9 ± 0.2 (2θ), about 25.0 ± 0.2 (2θ), about 27.9

± 0.2 (2Θ) and about 30.3 ± 0.2 (2Θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 14.5 ± 0.2 (2Θ), about 22.9 ± 0.2 (2Θ), about 23.5 ± 0.2 (2Θ), about 30.0 ± 0.2 (2Θ), about 30.8 ± 0.2 (2Θ), about 31.4 ± 0.2 (2Θ), about 31.6 ± 0.2 (2Θ) and about 32.2 ± 0.2 (2Θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 8.0 ± 0.2 (2Θ), about 10.6 ± 0.2 (2Θ), about 24.0 ± 0.2 (2Θ), about 32.2 ± 0.2 (2Θ), about 32.8 ± 0.2 (2Θ), about 33.3 ± 0.2 (2Θ), about 34.4 ± 0.2 (2Θ) and about 34.4 ± 0.2 (2Θ).

[0209] All 2Θ values indicated above refer to an X-ray diffractogram measured using $\text{CuK}\alpha$ radiation having a wavelength of $1.54060 \text{ \AA}$.

[0210] The crystalline form F according to the present invention may further be characterized in that it has one or more Raman peaks selected from the group consisting of about $171 \pm 2 \text{ cm}^{-1}$, about $921 \pm 2 \text{ cm}^{-1}$, about $1002 \pm 2 \text{ cm}^{-1}$, about $1299 \pm 2 \text{ cm}^{-1}$, about $1570 \pm 2 \text{ cm}^{-1}$, about $1581 \pm 2 \text{ cm}^{-1}$, about $2952 \pm 2 \text{ cm}^{-1}$ and about $3070 \pm 2 \text{ cm}^{-1}$.

[0211] The crystalline form F according to the present invention may further be characterized in that it has Raman peaks at $171 \pm 2 \text{ cm}^{-1}$, $921 \pm 2 \text{ cm}^{-1}$, $1002 \pm 2 \text{ cm}^{-1}$, $1299 \pm 2 \text{ cm}^{-1}$, $1570 \pm 2 \text{ cm}^{-1}$, $1581 \pm 2 \text{ cm}^{-1}$, $2952 \pm 2 \text{ cm}^{-1}$, $3070 \pm 2 \text{ cm}^{-1}$, $157 \pm 2 \text{ cm}^{-1}$, $183 \pm 2 \text{ cm}^{-1}$, $682 \pm 2 \text{ cm}^{-1}$, $1463 \pm 2 \text{ cm}^{-1}$, $1477 \pm 2 \text{ cm}^{-1}$, $2889 \pm 2 \text{ cm}^{-1}$, $2932 \pm 2 \text{ cm}^{-1}$, $2977 \pm 2 \text{ cm}^{-1}$ and $3058 \pm 2 \text{ cm}^{-1}$; and one or more Raman peaks selected from the group consisting of about $212 \pm 2 \text{ cm}^{-1}$, about $253 \pm 2 \text{ cm}^{-1}$, about $370 \pm 2 \text{ cm}^{-1}$, about $491 \pm 2 \text{ cm}^{-1}$, about $620 \pm 2 \text{ cm}^{-1}$, about $632 \pm 2 \text{ cm}^{-1}$, about $828 \pm 2 \text{ cm}^{-1}$, about $912 \pm 2 \text{ cm}^{-1}$, about $982 \pm 2 \text{ cm}^{-1}$, about $1027 \pm 2 \text{ cm}^{-1}$, about $1036 \pm 2 \text{ cm}^{-1}$, about $1050 \pm 2 \text{ cm}^{-1}$, about $1056 \pm 2 \text{ cm}^{-1}$, about $1110 \pm 2 \text{ cm}^{-1}$, about $1159 \pm 2 \text{ cm}^{-1}$, about $1189 \pm 2 \text{ cm}^{-1}$, about $1202 \pm 2 \text{ cm}^{-1}$, about $1373 \pm 2 \text{ cm}^{-1}$, about $1438 \pm 2 \text{ cm}^{-1}$, about $1453 \pm 2 \text{ cm}^{-1}$, about $2843 \pm 2 \text{ cm}^{-1}$, about $2860 \pm 2 \text{ cm}^{-1}$ and about $2992 \pm 2 \text{ cm}^{-1}$.

[0212] The crystalline form F according to the present invention may further be characterized in that it has one or more Raman peaks selected from the group consisting of about $291 \pm 2 \text{ cm}^{-1}$, about $394 \pm 2 \text{ cm}^{-1}$, about $424 \pm 2 \text{ cm}^{-1}$, about $471 \pm 2 \text{ cm}^{-1}$, about $519 \pm 2 \text{ cm}^{-1}$, about $542 \pm 2 \text{ cm}^{-1}$, about $560 \pm 2 \text{ cm}^{-1}$, about $602 \pm 2 \text{ cm}^{-1}$, about $607 \pm 2 \text{ cm}^{-1}$, about $712 \pm 2 \text{ cm}^{-1}$, about $762 \pm 2 \text{ cm}^{-1}$, about $786 \pm 2 \text{ cm}^{-1}$, about $848 \pm 2 \text{ cm}^{-1}$, about $870 \pm 2 \text{ cm}^{-1}$, about $894 \pm 2 \text{ cm}^{-1}$, about $946 \pm 2 \text{ cm}^{-1}$, about $970 \pm 2 \text{ cm}^{-1}$, about $1076 \pm 2 \text{ cm}^{-1}$, about $1119 \pm 2 \text{ cm}^{-1}$, about $1146 \pm 2 \text{ cm}^{-1}$, about $1172 \pm 2 \text{ cm}^{-1}$, about $1229 \pm 2 \text{ cm}^{-1}$, about $1263 \pm 2 \text{ cm}^{-1}$, about $1338 \pm 2 \text{ cm}^{-1}$, about $1353 \pm 2 \text{ cm}^{-1}$, about $1498 \pm 2 \text{ cm}^{-1}$, about $1630 \pm 2 \text{ cm}^{-1}$, about $2566 \pm 2 \text{ cm}^{-1}$, about $2748 \pm 2 \text{ cm}^{-1}$ and about $2795 \pm 2 \text{ cm}^{-1}$.

[0213] Another reference aspect of the present invention relates to a process for the production of the crystalline form F described above comprising the step of
(a-1) suspending (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-

1,1'-pyrano[3,4,b]indol]-4-amine in a solvent comprising n-propanol.

[0214] Preferably, the solvent is n-propanol.

[0215] Preferably, step (a-1) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80 °C, more preferably not higher than 60°C, even more preferably not higher than 40°C, and in particular in a temperature range of 15-35 °C.

[0216] Preferably, the suspension obtained in step (a-1) is stirred for a time period of at least 2 h, preferably at least 4 h, more preferably at least 8 h, still more preferably at least 12 h, yet more preferably at least 16 h, most preferably at least 24 h, and in particular at least 2 days.

[0217] Preferably, the process further comprises the step (b-1) separating, preferably filtering off the solid obtained in step (a-1).

[0218] Preferably, the process further comprises the step (c-1) drying of the solid obtained in step (b-1).

[0219] Preferably, in the process, step (c-1) takes place under air or inert gas flow, such as argon or nitrogen flow. Preferably, step (c-1) is performed at ambient temperature.

[0220] A further reference aspect of the present invention relates to a crystalline form F that is obtainable by the process as described above.

[0221] A further reference aspect of the present invention relates to a crystalline form G.

[0222] Preferably, the crystalline form G has one or more X-ray diffraction peaks selected from the group consisting of about 16.3 $\pm$ 0.2 (2 Θ), about 18.8 $\pm$ 0.2 (2 Θ), about 19.1 $\pm$ 0.2 (2 Θ), about 19.4 $\pm$ 0.2 (2 Θ), about 20.3 $\pm$ 0.2 (2 Θ), about 22.2 $\pm$ 0.2 (2 Θ) and about 29.1 $\pm$ 0.2 (2 Θ). In some preferred embodiments, the crystalline form comprises an X-ray diffraction peak at about 19.1 $\pm$ 0.2 (2 Θ).

[0223] In some preferred embodiments, crystalline form G comprises X-ray diffraction peaks at about 18.8 $\pm$ 0.2 (2 Θ), about 19.1 $\pm$ 0.2 (2 Θ), about 22.2 $\pm$ 0.2 (2 Θ), about 29.1 $\pm$ 0.2 (2 Θ) and optionally 20.3 $\pm$ 0.2 (2 Θ).

[0224] The crystalline form G may additionally have at least one X-ray diffraction peak selected from the group consisting of about 15.4 $\pm$ 0.2 (2 Θ), about 15.9 $\pm$ 0.2 (2 Θ), about 17.2 $\pm$ 0.2 (2 Θ), about 17.4 $\pm$ 0.2 (2 Θ), about 17.8 $\pm$ 0.2 (2 Θ), about 20.7 $\pm$ 0.2 (2 Θ), about 21.0 $\pm$ 0.2 (2 Θ), about 22.6 $\pm$ 0.2 (2 Θ), about 24.2 $\pm$ 0.2 (2 Θ), about 24.7 $\pm$ 0.2 (2 Θ), about 25.4 $\pm$ 0.2 (2 Θ), about 25.9 $\pm$ 0.2 (2 Θ), about 26.6 $\pm$ 0.2 (2 Θ), about 28.0 $\pm$ 0.2 (2 Θ), about 28.3 $\pm$ 0.2 (2 Θ), about 28.8 $\pm$ 0.2 (2 Θ), about 29.4 $\pm$ 0.2 (2 Θ) and about 33.0 $\pm$ 0.2 (2 Θ).

[0225] Furthermore, the crystalline form G may be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 16.3 ± 0.2 (2θ), about 18.8 ± 0.2 (2θ), about 19.1 ± 0.2 (2θ), about 19.4 ± 0.2 (2θ), about 20.3 ± 0.2 (2θ), about 22.2 ± 0.2 (2θ) and about 29.1 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 15.4 ± 0.2 (2θ), about 15.9 ± 0.2 (2θ), about 17.2 ± 0.2 (2θ), about 17.4 ± 0.2 (2θ), about 17.8 ± 0.2 (2θ), about 20.7 ± 0.2 (2θ), about 21.0 ± 0.2 (2θ), about 22.6 ± 0.2 (2θ), about 24.2 ± 0.2 (2θ), about 24.7 ± 0.2 (2θ), about 25.4 ± 0.2 (2θ), about 25.9 ± 0.2 (2θ), about 26.6 ± 0.2 (2θ), about 28.0 ± 0.2 (2θ), about 28.3 ± 0.2 (2θ), about 28.8 ± 0.2 (2θ), about 29.4 ± 0.2 (2θ) and about 33.0 ± 0.2 (2θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 11.5 ± 0.2 (2θ), about 15.1 ± 0.2 (2θ), about 30.2 ± 0.2 (2θ), about 31.6 ± 0.2 (2θ), about 32.3 ± 0.2 (2θ), about 33.9 ± 0.2 (2θ) and about 34.7 ± 0.2 (2θ).

[0226] The crystalline form G may further be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 16.3 ± 0.2 (2θ), about 18.8 ± 0.2 (2θ), about 19.1 ± 0.2 (2θ), about 19.4 ± 0.2 (2θ), about 20.3 ± 0.2 (2θ), about 22.2 ± 0.2 (2θ) and about 29.1 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 15.4 ± 0.2 (2θ), about 15.9 ± 0.2 (2θ), about 17.2 ± 0.2 (2θ), about 17.4 ± 0.2 (2θ), about 17.8 ± 0.2 (2θ), about 20.7 ± 0.2 (2θ), about 21.0 ± 0.2 (2θ), about 22.6 ± 0.2 (2θ), about 24.2 ± 0.2 (2θ), about 24.7 ± 0.2 (2θ), about 25.4 ± 0.2 (2θ), about 25.9 ± 0.2 (2θ), about 26.6 ± 0.2 (2θ), about 28.0 ± 0.2 (2θ), about 28.3 ± 0.2 (2θ), about 28.8 ± 0.2 (2θ), about 29.4 ± 0.2 (2θ) and about 33.0 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 11.5 ± 0.2 (2θ), about 15.1 ± 0.2 (2θ), about 30.2 ± 0.2 (2θ), about 31.6 ± 0.2 (2θ), about 32.3 ± 0.2 (2θ), about 33.9 ± 0.2 (2θ) and about 34.7 ± 0.2 (2θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 8.1 ± 0.2 (2θ), about 8.9 ± 0.2 (2θ), about 11.1 ± 0.2 (2θ), about 13.5 ± 0.2 (2θ) and about 33.5 ± 0.2 (2θ).

[0227] All 2θ values indicated above refer to an X-ray diffractogram measured using CuK α radiation having a wavelength of 1.54060 Å.

[0228] The crystalline form G may further be characterized in that it has one or more Raman peaks selected from the group consisting of about 675 ± 2 cm^{-1} , about 1569 ± 2 cm^{-1} and about 2917 ± 2 cm^{-1} .

[0229] The crystalline form G may further be characterized in that it has one or more Raman peaks selected from the group consisting of about 675 ± 2 cm^{-1} , about 1569 ± 2 cm^{-1} and about 2917 ± 2 cm^{-1} ; and/or one or more Raman peaks selected from the group consisting of about 169 ± 2 cm^{-1} , about 921 ± 2 cm^{-1} , about 1002 ± 2 cm^{-1} , about 3069 ± 2 cm^{-1} ; and/or one or more Raman peaks selected from the group consisting of about 180 ± 2 cm^{-1} , about 202 ± 2 cm^{-1} , about 254 ± 2 cm^{-1} , about 306 ± 2 cm^{-1} , about 706 ± 2 cm^{-1} , about 1029 ± 2 cm^{-1} , about 1047 ± 2 cm^{-1} , about 1292 ± 2 cm^{-1} , about 1309 ± 2 cm^{-1} , about 1418 ± 2 cm^{-1} , about 1437 ± 2 cm^{-1} ,

about $1475 \pm 2 \text{ cm}^{-1}$, about $1597 \pm 2 \text{ cm}^{-1}$, about $2945 \pm 2 \text{ cm}^{-1}$, about $2960 \pm 2 \text{ cm}^{-1}$, about $2999 \pm 2 \text{ cm}^{-1}$, about $3058 \pm 2 \text{ cm}^{-1}$.

[0230] The crystalline form G may further be characterized in that it has one or more Raman peaks selected from the group consisting of about $334 \pm 2 \text{ cm}^{-1}$, about $365 \pm 2 \text{ cm}^{-1}$, about $387 \pm 2 \text{ cm}^{-1}$, about $424 \pm 2 \text{ cm}^{-1}$, about $438 \pm 2 \text{ cm}^{-1}$, about $491 \pm 2 \text{ cm}^{-1}$, about $522 \pm 2 \text{ cm}^{-1}$, about $546 \pm 2 \text{ cm}^{-1}$, about $608 \pm 2 \text{ cm}^{-1}$, about $621 \pm 2 \text{ cm}^{-1}$, about $638 \pm 2 \text{ cm}^{-1}$, about $769 \pm 2 \text{ cm}^{-1}$, about $786 \pm 2 \text{ cm}^{-1}$, about $830 \pm 2 \text{ cm}^{-1}$, about $868 \pm 2 \text{ cm}^{-1}$, about $948 \pm 2 \text{ cm}^{-1}$, about $982 \pm 2 \text{ cm}^{-1}$, about $1038 \pm 2 \text{ cm}^{-1}$, about $1073 \pm 2 \text{ cm}^{-1}$, about $1108 \pm 2 \text{ cm}^{-1}$, about $1122 \pm 2 \text{ cm}^{-1}$, about $1136 \pm 2 \text{ cm}^{-1}$, about $1161 \pm 2 \text{ cm}^{-1}$, about $1171 \pm 2 \text{ cm}^{-1}$, about $1190 \pm 2 \text{ cm}^{-1}$, about $1201 \pm 2 \text{ cm}^{-1}$, about $1234 \pm 2 \text{ cm}^{-1}$, about $1260 \pm 2 \text{ cm}^{-1}$, about $1338 \pm 2 \text{ cm}^{-1}$, about $1373 \pm 2 \text{ cm}^{-1}$, about $1629 \pm 2 \text{ cm}^{-1}$, about $2777 \pm 2 \text{ cm}^{-1}$, about $2815 \pm 2 \text{ cm}^{-1}$, about $2841 \pm 2 \text{ cm}^{-1}$, about $2862 \pm 2 \text{ cm}^{-1}$, about $3156 \pm 2 \text{ cm}^{-1}$.

[0231] Another reference aspect of the present invention relates to a process for the production of the crystalline form G described above comprising the step of
(a-1) suspending (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in a solvent comprising dimethylsulfoxid.

[0232] Preferably, the solvent is dimethylsulfoxid.

[0233] Preferably, step (a-1) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80°C , more preferably not higher than 60°C , even more preferably not higher than 40°C , and in particular in a temperature range of $15\text{-}35^\circ\text{C}$.

[0234] Preferably, the suspension obtained in step (a-1) is stirred for a time period of at least 2 h, preferably at least 4 h, more preferably at least 8 h, still more preferably at least 12 h, yet more preferably at least 16 h, most preferably at least 24 h, and in particular at least 2 days.

[0235] Preferably, the process further comprises the step
(b-1) separating, preferably filtering off the solid obtained in step (a-1).

[0236] Preferably, the process further comprises the step
(c-1) drying of the solid obtained in step (b-1).

[0237] Preferably, in the process, step (c-1) takes place under air or inert gas flow, such as argon or nitrogen flow. Preferably, step (c-1) is performed at ambient temperature.

[0238] A further reference aspect of the present invention relates to a crystalline form G that is obtainable by the process as described above.

[0239] A further reference aspect of the present invention relates to a crystalline form H.

[0240] Preferably, the crystalline form H has one or more X-ray diffraction peaks selected from the group consisting of about 11.4 ± 0.2 (2θ), about 18.3 ± 0.2 (2θ) and about 19.2 ± 0.2 (2θ). In some preferred embodiments, the crystalline form comprises an X-ray diffraction peak at about 19.2 ± 0.2 (2θ).

[0241] In some preferred embodiments, crystalline form H comprises X-ray diffraction peaks at about 11.4 ± 0.2 (2θ), about 18.3 ± 0.2 (2θ), about 19.2 ± 0.2 (2θ), about 23.0 ± 0.2 (2θ) and optionally 18.0 ± 0.2 (2θ).

[0242] The crystalline form H may additionally have at least one X-ray diffraction peak selected from the group consisting of about 17.1 ± 0.2 (2θ), about 18.0 ± 0.2 (2θ), about 20.6 ± 0.2 (2θ), about 21.4 ± 0.2 (2θ), about 23.0 ± 0.2 (2θ), about 25.5 ± 0.2 (2θ), about 26.3 ± 0.2 (2θ) and about 27.9 ± 0.2 (2θ).

[0243] Furthermore, the crystalline form H may be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 11.4 ± 0.2 (2θ), about 18.3 ± 0.2 (2θ) and about 19.2 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 17.1 ± 0.2 (2θ), about 18.0 ± 0.2 (2θ), about 20.6 ± 0.2 (2θ), about 21.4 ± 0.2 (2θ), about 23.0 ± 0.2 (2θ), about 25.5 ± 0.2 (2θ), about 26.3 ± 0.2 (2θ) and about 27.9 ± 0.2 (2θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 7.4 ± 0.2 (2θ), about 10.8 ± 0.2 (2θ), about 15.7 ± 0.2 (2θ), about 16.2 ± 0.2 (2θ), about 19.8 ± 0.2 (2θ), about 20.3 ± 0.2 (2θ), about 22.2 ± 0.2 (2θ), about 23.8 ± 0.2 (2θ), about 24.5 ± 0.2 (2θ), about 25.8 ± 0.2 (2θ), about 27.0 ± 0.2 (2θ), about 28.3 ± 0.2 (2θ), about 28.8 ± 0.2 (2θ), about 29.4 ± 0.2 (2θ), about 30.0 ± 0.2 (2θ), about 31.0 ± 0.2 (2θ) and about 33.3 ± 0.2 (2θ).

[0244] The crystalline form H may further be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 11.4 ± 0.2 (2θ), about 18.3 ± 0.2 (2θ) and about 19.2 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 17.1 ± 0.2 (2θ), about 18.0 ± 0.2 (2θ), about 20.6 ± 0.2 (2θ), about 21.4 ± 0.2 (2θ), about 23.0 ± 0.2 (2θ), about 25.5 ± 0.2 (2θ), about 26.3 ± 0.2 (2θ) and about 27.9 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 7.4 ± 0.2 (2θ), about 10.8 ± 0.2 (2θ), about 15.7 ± 0.2 (2θ), about 16.2 ± 0.2 (2θ), about 19.8 ± 0.2 (2θ), about 20.3 ± 0.2 (2θ), about 22.2 ± 0.2 (2θ), about 23.8 ± 0.2 (2θ), about 24.5 ± 0.2 (2θ), about 25.8 ± 0.2 (2θ), about 27.0 ± 0.2 (2θ), about 28.3 ± 0.2 (2θ), about 28.8 ± 0.2 (2θ), about 29.4 ± 0.2 (2θ), about 30.0 ± 0.2 (2θ), about 31.0 ± 0.2 (2θ) and about 33.3 ± 0.2 (2θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 8.5 ± 0.2 (2θ), about 9.0 ± 0.2 (2θ), about 9.8 ± 0.2 (2θ), about 12.2 ± 0.2 (2θ), about 12.8 ± 0.2 (2θ), about 13.1 ± 0.2 (2θ), about 14.8 ± 0.2 (2θ), about 16.6 ± 0.2 (2θ), about 27.6 ± 0.2 (2θ) and about 32.7 ± 0.2 (2θ).

[0245] All 2θ values indicated above refer to a X-ray diffractogram measured using CuK α

radiation having a wavelength of 1.54060 Å.

[0246] The crystalline form H may further be characterized in that it has one or more Raman peaks selected from the group consisting of about $171 \pm 2 \text{ cm}^{-1}$, about $203 \pm 2 \text{ cm}^{-1}$, about $258 \pm 2 \text{ cm}^{-1}$, about $918 \pm 2 \text{ cm}^{-1}$, about $1002 \pm 2 \text{ cm}^{-1}$, about $1305 \pm 2 \text{ cm}^{-1}$ and about $1568 \pm 2 \text{ cm}^{-1}$.

[0247] The crystalline form H may further be characterized in that as well as one or more Raman peaks selected from the group consisting of about $171 \pm 2 \text{ cm}^{-1}$, about $203 \pm 2 \text{ cm}^{-1}$, about $258 \pm 2 \text{ cm}^{-1}$, about $918 \pm 2 \text{ cm}^{-1}$, about $1002 \pm 2 \text{ cm}^{-1}$, about $1305 \pm 2 \text{ cm}^{-1}$ and about $1568 \pm 2 \text{ cm}^{-1}$; it additionally has one or more Raman peaks selected from the group consisting of about $369 \pm 2 \text{ cm}^{-1}$, about $391 \pm 2 \text{ cm}^{-1}$, about $490 \pm 2 \text{ cm}^{-1}$, about $599 \pm 2 \text{ cm}^{-1}$, about $685 \pm 2 \text{ cm}^{-1}$, about $828 \pm 2 \text{ cm}^{-1}$, about $1030 \pm 2 \text{ cm}^{-1}$, about $1375 \pm 2 \text{ cm}^{-1}$, about $1464 \pm 2 \text{ cm}^{-1}$, about $2989 \pm 2 \text{ cm}^{-1}$.

[0248] The crystalline form H may further be characterized in that it has one or more Raman peaks selected from the group consisting of about $517 \pm 2 \text{ cm}^{-1}$, about $557 \pm 2 \text{ cm}^{-1}$, about $620 \pm 2 \text{ cm}^{-1}$, about $713 \pm 2 \text{ cm}^{-1}$, about $734 \pm 2 \text{ cm}^{-1}$, about $787 \pm 2 \text{ cm}^{-1}$, about $889 \pm 2 \text{ cm}^{-1}$, about $982 \pm 2 \text{ cm}^{-1}$, about $1048 \pm 2 \text{ cm}^{-1}$, about $1073 \pm 2 \text{ cm}^{-1}$, about $1117 \pm 2 \text{ cm}^{-1}$, about $1199 \pm 2 \text{ cm}^{-1}$, about $1219 \pm 2 \text{ cm}^{-1}$, about $1263 \pm 2 \text{ cm}^{-1}$, about $1629 \pm 2 \text{ cm}^{-1}$, about $2788 \pm 2 \text{ cm}^{-1}$, about $2921 \pm 2 \text{ cm}^{-1}$, about $2945 \pm 2 \text{ cm}^{-1}$ and about $3069 \pm 2 \text{ cm}^{-1}$.

[0249] Another reference aspect of the present invention relates to a process for the production of the crystalline form H described above comprising the step of

(a-1) dissolving (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in an organic solvent.

[0250] In a preferred embodiment, the organic solvent is 1,4-dioxane. In another preferred embodiment, the organic solvent contains at least 5 wt.-%, more preferably at least 10 wt.-%, still more preferably at least 20 wt.-%, yet more preferably at least 30 wt.-%, most preferably at least 40 wt.-%, and in particular at least 50 wt.-% 1,4-dioxane.

[0251] Preferably, in the process, step (a-1) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80°C , more preferably not higher than 60°C , even more preferably not higher than 40°C , and in particular in a temperature range of $20\text{--}40^\circ\text{C}$.

[0252] In a preferred embodiment, the further comprises the step
(b-1) evaporating off the solvent of the solution obtained in step (a-1).

[0253] Suitable methods for evaporating off the solvent are known to a person skilled in the art. Preferably, in the process, the solvent is evaporated off in air, air flow, or inert gas flow, in

particular argon or nitrogen flow. However, evaporating off the solvent under vacuum, for example by means of a rotary evaporator, is also possible.

[0254] Preferably, in the process, the solvent is evaporated off at room temperature. However, evaporating off the solvent at an elevated temperature, e. g. within the range of 20 °C to 60 °C, is also possible.

[0255] A further reference aspect of the present invention relates to a crystalline form H that is obtainable by the process as described above.

[0256] A further reference aspect of the present invention relates to a crystalline form I.

[0257] Preferably, the crystalline form I has one or more X-ray diffraction peaks selected from the group consisting of about 10.9 ± 0.2 (2 θ), about 14.6 ± 0.2 (2 θ), about 15.5 ± 0.2 (2 θ), about 17.1 ± 0.2 (2 θ), about 18.5 ± 0.2 (2 θ), about 18.8 ± 0.2 (2 θ), about 21.1 ± 0.2 (2 θ), about 21.9 ± 0.2 (2 θ), about 23.6 ± 0.2 (2 θ), about 25.9 ± 0.2 (2 θ) and about 28.0 ± 0.2 (2 θ). In some preferred embodiments, the crystalline form comprises an X-ray diffraction peak at about 17.1 ± 0.2 .

[0258] In some preferred embodiments, crystalline form I comprises X-ray diffraction peaks at about 10.9 ± 0.2 (2 θ), about 15.5 ± 0.2 (2 θ), about 17.1 ± 0.2 (2 θ), about 18.5 ± 0.2 (2 θ) and optionally 18.8 ± 0.2 (2 θ) and/or 23.6 ± 0.2 (2 θ).

[0259] The crystalline form I may additionally have at least one X-ray diffraction peak selected from the group consisting of about 16.5 ± 0.2 (2 θ), about 18.1 ± 0.2 (2 θ), about 24.0 ± 0.2 (2 θ), about 28.0 ± 0.2 (2 θ), about 28.8 ± 0.2 (2 θ), about 30.6 ± 0.2 (2 θ) and about 31.9 ± 0.2 (2 θ).

[0260] Furthermore, the crystalline form I may be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 10.9 ± 0.2 (2 θ), about 14.6 ± 0.2 (2 θ), about 15.5 ± 0.2 (2 θ), about 17.1 ± 0.2 (2 θ), about 18.5 ± 0.2 (2 θ), about 18.8 ± 0.2 (2 θ), about 21.1 ± 0.2 (2 θ), about 21.9 ± 0.2 (2 θ), about 23.6 ± 0.2 (2 θ), about 25.9 ± 0.2 (2 θ), and about 28.0 ± 0.2 (2 θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 16.5 ± 0.2 (2 θ), about 18.1 ± 0.2 (2 θ), about 24.0 ± 0.2 (2 θ), about 28.0 ± 0.2 (2 θ), about 28.8 ± 0.2 (2 θ), about 30.6 ± 0.2 (2 θ) and about 31.9 ± 0.2 (2 θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 19.4 ± 0.2 (2 θ), about 19.9 ± 0.2 (2 θ), about 20.2 ± 0.2 (2 θ), about 22.3 ± 0.2 (2 θ), about 22.8 ± 0.2 (2 θ), about 23.0 ± 0.2 (2 θ), about 25.2 ± 0.2 (2 θ), about 26.6 ± 0.2 (2 θ), about 30.0 ± 0.2 (2 θ) and about 34.8 ± 0.2 (2 θ).

[0261] The crystalline form I may further be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 10.9 ± 0.2 (2 θ), about 14.6 ± 0.2 (2 θ), about 15.5 ± 0.2 (2 θ), about 17.1 ± 0.2 (2 θ), about 18.5 ± 0.2 (2 θ), about 18.8 ± 0.2 (2 θ), about 21.1 ± 0.2 (2 θ), about 21.9 ± 0.2 (2 θ), about 23.6 ± 0.2 (2 θ), about 25.9 ± 0.2 (2 θ), and about 28.0 ± 0.2 (2 θ), and optionally one or more X-ray diffraction peaks selected from the

group consisting of about 16.5 ± 0.2 (2Θ), about 18.1 ± 0.2 (2Θ), about 24.0 ± 0.2 (2Θ), about 28.0 ± 0.2 (2Θ), about 28.8 ± 0.2 (2Θ), about 30.6 ± 0.2 (2Θ) and about 31.9 ± 0.2 (2Θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 19.4 ± 0.2 (2Θ), about 19.9 ± 0.2 (2Θ), about 20.2 ± 0.2 (2Θ), about 22.3 ± 0.2 (2Θ), about 22.8 ± 0.2 (2Θ), about 23.0 ± 0.2 (2Θ), about 25.2 ± 0.2 (2Θ), about 26.6 ± 0.2 (2Θ), about 30.0 ± 0.2 (2Θ) and about 34.8 ± 0.2 (2Θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 9.5 ± 0.2 (2Θ), about 10.1 ± 0.2 (2Θ), about 13.9 ± 0.2 (2Θ), about 27.6 ± 0.2 (2Θ), about 29.1 ± 0.2 (2Θ), about 33.1 ± 0.2 (2Θ) and about 34.2 ± 0.2 (2Θ).

[0262] All 2Θ values indicated above refer to an X-ray diffractogram measured using CuK α radiation having a wavelength of 1.54060 Å.

[0263] The crystalline form I invention may further be characterized in that it has one or more Raman peaks selected from the group consisting of about 924 ± 2 cm $^{-1}$, about 1001 ± 2 cm $^{-1}$, about 1305 ± 2 cm $^{-1}$, about 1572 ± 2 cm $^{-1}$, about 2925 ± 2 cm $^{-1}$, about 3066 ± 2 cm $^{-1}$.

[0264] The crystalline form I may further be characterized in that as well as one or more Raman peaks selected from the group consisting of about 924 ± 2 cm $^{-1}$, about 1001 ± 2 cm $^{-1}$, about 1305 ± 2 cm $^{-1}$, about 1572 ± 2 cm $^{-1}$, about 2925 ± 2 cm $^{-1}$, about 3066 ± 2 cm $^{-1}$; it additionally has one or more Raman peaks selected from the group consisting of about 155 ± 2 cm $^{-1}$, about 172 ± 2 cm $^{-1}$, about 256 ± 2 cm $^{-1}$, about 680 ± 2 cm $^{-1}$, about 1031 ± 2 cm $^{-1}$, about 1434 ± 2 cm $^{-1}$, about 1459 ± 2 cm $^{-1}$, about 1474 ± 2 cm $^{-1}$, about 1589 ± 2 cm $^{-1}$, about 1596 ± 2 cm $^{-1}$, about 2911 ± 2 cm $^{-1}$, about 2964 ± 2 cm $^{-1}$ and about 2984 ± 2 cm $^{-1}$.

[0265] The crystalline form I may further be characterized in that it has one or more Raman peaks selected from the group consisting of about 202 ± 2 cm $^{-1}$, about 280 ± 2 cm $^{-1}$, about 315 ± 2 cm $^{-1}$, about 367 ± 2 cm $^{-1}$, about 392 ± 2 cm $^{-1}$, about 421 ± 2 cm $^{-1}$, about 438 ± 2 cm $^{-1}$, about 466 ± 2 cm $^{-1}$, about 489 ± 2 cm $^{-1}$, about 519 ± 2 cm $^{-1}$, about 544 ± 2 cm $^{-1}$, about 558 ± 2 cm $^{-1}$, about 605 ± 2 cm $^{-1}$, about 621 ± 2 cm $^{-1}$, about 636 ± 2 cm $^{-1}$, about 697 ± 2 cm $^{-1}$, about 715 ± 2 cm $^{-1}$, about 767 ± 2 cm $^{-1}$, about 784 ± 2 cm $^{-1}$, about 810 ± 2 cm $^{-1}$, about 825 ± 2 cm $^{-1}$, about 895 ± 2 cm $^{-1}$, about 912 ± 2 cm $^{-1}$, about 948 ± 2 cm $^{-1}$, about 983 ± 2 cm $^{-1}$, about 1047 ± 2 cm $^{-1}$, about 1066 ± 2 cm $^{-1}$, about 1091 ± 2 cm $^{-1}$, 1113 ± 2 cm $^{-1}$, about 1123 ± 2 cm $^{-1}$, about 1141 ± 2 cm $^{-1}$, about 1159 ± 2 cm $^{-1}$, about 1188 ± 2 cm $^{-1}$, about 1199 ± 2 cm $^{-1}$, 1232 ± 2 cm $^{-1}$, about 1265 ± 2 cm $^{-1}$, about 1291 ± 2 cm $^{-1}$, about 1339 ± 2 cm $^{-1}$, about 1354 ± 2 cm $^{-1}$, about 1375 ± 2 cm $^{-1}$, 1404 ± 2 cm $^{-1}$, about 1417 ± 2 cm $^{-1}$, about 1630 ± 2 cm $^{-1}$, about 2699 ± 2 cm $^{-1}$, about 2775 ± 2 cm $^{-1}$, about 2787 ± 2 cm $^{-1}$, 2820 ± 2 cm $^{-1}$, about 2845 ± 2 cm $^{-1}$, about 2875 ± 2 cm $^{-1}$, about 2953 ± 2 cm $^{-1}$, about 2998 ± 2 cm $^{-1}$, about 3011 ± 2 cm $^{-1}$, 3051 ± 2 cm $^{-1}$ and about 3085 ± 2 cm $^{-1}$.

[0266] Another reference aspect of the present invention relates to a process for the production of the crystalline form I described above comprising the step of

(a-1) dissolving (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in an organic solvent.

[0267] Preferably, the solvent is selected from the group consisting of the C4 to C6 alcohols such as n-butanol, esters such as ethyl acetate, n-propyl acetate, iso-propyl acetate, n-butyl acetate, iso-butyl acetate, ketones such as acetone, 2-butanone, pentan-2-one, pentan-3-one, hexan-2-one, hexan-3-one, ethers such as tert-butyl methyl ether, diethylether, tetrahydrofuran, diisopropylether and 1,4-dioxane, nitriles such as acetonitril, saturated hydrocarbons such as n-pentane, n-hexane and n-heptane, aromatic hydrocarbons such as toluene, chlorinated hydrocarbons such as dichloromethane and chloroform, and mixtures thereof.

[0268] Mixtures of saturated hydrocarbons, such as n-pentane, n-hexane and n-heptane, further containing at least one solvent selected from the group consisting of ketones, ethers and chlorinated hydrocarbons are particularly preferred.

[0269] In some preferred embodiments, the organic solvent is a mixture of dichloromethane and n-heptane. Preferably, the ratio between dichloromethane and n-heptane is within the range of from 10:1 to 1:10, more preferably within the range of from 7:1 to 1:7, still more preferably within the range of from 5:1 to 1:6, yet more preferably within the range of from 3:1 to 1:5, most preferably within the range of from 1:1 to 1:3, and in particular within the range of from 1:1.5 to 1:2.5 (volume/volume).

[0270] Preferably, in the process, step (a-1) is carried out at a temperature below or at the boiling point of the respective solvent, preferably at a temperature not higher than 80 °C, more preferably not higher than 60°C, even more preferably not higher than 40°C, and in particular in a temperature range of 20-40 °C.

[0271] In a preferred embodiment, the process further comprises the step (b-1) evaporating off the solvent of the solution obtained in step (a-1).

[0272] Suitable methods for evaporating off the solvent are known to a person skilled in the art. Preferably, in the process, the solvent is evaporated off in air, air flow, or inert gas flow, in particular argon or nitrogen flow. However, evaporating off the solvent under vacuum, for example by means of a rotary evaporator, is also possible.

[0273] Preferably, in the process, the solvent is evaporated off at room temperature. However, evaporating off the solvent at an elevated temperature, e. g. within the range of 20 °C to 60 °C, is also possible.

[0274] A further reference aspect of the present invention relates to a crystalline form I that is obtainable by the process as described above.

[0275] A further reference aspect of the present invention relates to a crystalline form L.

[0276] Preferably, the crystalline form L has one or more X-ray diffraction peaks selected from the group consisting of about 8.6 ± 0.2 (2θ), about 10.3 ± 0.2 (2θ), about 16.7 ± 0.2 (2θ), about 17.2 ± 0.2 (2θ), about 18.2 ± 0.2 (2θ), about 18.8 ± 0.2 (2θ), about 21.2 ± 0.2 (2θ), about 26.0 ± 0.2 (2θ), and about 27.4 ± 0.2 (2θ). In some preferred embodiments, the crystalline form comprises an X-ray diffraction peak at about 18.8 ± 0.2 (2θ).

[0277] In some preferred embodiments, crystalline form L comprises X-ray diffraction peaks at about 16.7 ± 0.2 (2θ), about 17.2 ± 0.2 (2θ), about 18.2 ± 0.2 (2θ), about 18.8 ± 0.2 (2θ) and optionally 10.3 ± 0.2 (2θ).

[0278] The crystalline form L may additionally have at least one X-ray diffraction peak selected from the group consisting of about 14.0 ± 0.2 (2θ), about 20.7 ± 0.2 (2θ), about 23.0 ± 0.2 (2θ), about 28.9 ± 0.2 (2θ) and about 30.2 ± 0.2 (2θ).

[0279] Furthermore, the crystalline form L may be characterized in that as well as one or more X-ray diffraction peaks selected from the group consisting of about 8.6 ± 0.2 (2θ), about 10.3 ± 0.2 (2θ), about 16.7 ± 0.2 (2θ), about 17.2 ± 0.2 (2θ), about 18.2 ± 0.2 (2θ), about 18.8 ± 0.2 (2θ), about 21.2 ± 0.2 (2θ), about 26.0 ± 0.2 (2θ), and about 27.4 ± 0.2 (2θ), and optionally one or more X-ray diffraction peaks selected from the group consisting of about 14.0 ± 0.2 (2θ), about 20.7 ± 0.2 (2θ), about 23.0 ± 0.2 (2θ), about 28.9 ± 0.2 (2θ) and about 30.2 ± 0.2 (2θ), it additionally has at least one X-ray diffraction peak selected from the group consisting of about 9.1 ± 0.2 (2θ), about 9.5 ± 0.2 (2θ), about 12.2 ± 0.2 (2θ), about 22.3 ± 0.2 (2θ) and about 24.5 ± 0.2 (2θ).

[0280] All 2θ values indicated above refer to an X-ray diffractogram measured using CuK α radiation having a wavelength of 1.54060 Å.

[0281] The crystalline form L may further be characterized in that it has one or more Raman peaks selected from the group consisting of about 1001 ± 2 cm^{-1} , about 1577 ± 2 cm^{-1} , about 1590 ± 2 cm^{-1} and about 3069 ± 2 cm^{-1} .

[0282] The crystalline form L may further be characterized in that it has one or more Raman peaks selected from the group consisting of about 1001 ± 2 cm^{-1} , about 1577 ± 2 cm^{-1} , about 1590 ± 2 cm^{-1} and about 3069 ± 2 cm^{-1} ; and/or one or more Raman peaks selected from the group consisting of about 172 ± 2 cm^{-1} , about 679 ± 2 cm^{-1} , about 924 ± 2 cm^{-1} , about 1307 ± 2 cm^{-1} , about 1475 ± 2 cm^{-1} , about 2922 ± 2 cm^{-1} , about 2987 ± 2 cm^{-1} ; and/or one or more Raman peaks selected from the group consisting of about 150 ± 2 cm^{-1} , about 199 ± 2 cm^{-1} , about 249 ± 2 cm^{-1} , about 488 ± 2 cm^{-1} , about 620 ± 2 cm^{-1} , about 693 ± 2 cm^{-1} , about 828 ± 2 cm^{-1} , about 913 ± 2 cm^{-1} , about 985 ± 2 cm^{-1} , about 1029 ± 2 cm^{-1} , about 1201 ± 2 cm^{-1} , about 1293 ± 2 cm^{-1} , about 1376 ± 2 cm^{-1} , about 1438 ± 2 cm^{-1} , about 1631 ± 2 cm^{-1} , about 2843 ± 2 cm^{-1} .

cm^{-1} , about $2859 \pm 2 \text{ cm}^{-1}$, about $2879 \pm 2 \text{ cm}^{-1}$ and about $3042 \pm 2 \text{ cm}^{-1}$.

[0283] The crystalline form L may further be characterized in that it has one or more Raman peaks selected from the group consisting of about $362 \pm 2 \text{ cm}^{-1}$, about $390 \pm 2 \text{ cm}^{-1}$, about $423 \pm 2 \text{ cm}^{-1}$, about $522 \pm 2 \text{ cm}^{-1}$, about $544 \pm 2 \text{ cm}^{-1}$, about $610 \pm 2 \text{ cm}^{-1}$, about $637 \pm 2 \text{ cm}^{-1}$, about $714 \pm 2 \text{ cm}^{-1}$, about $763 \pm 2 \text{ cm}^{-1}$, about $784 \pm 2 \text{ cm}^{-1}$, about $872 \pm 2 \text{ cm}^{-1}$, about $950 \pm 2 \text{ cm}^{-1}$, about $1049 \pm 2 \text{ cm}^{-1}$, about $1077 \pm 2 \text{ cm}^{-1}$, about $1129 \pm 2 \text{ cm}^{-1}$, about $1157 \pm 2 \text{ cm}^{-1}$, about $1186 \pm 2 \text{ cm}^{-1}$, about $1235 \pm 2 \text{ cm}^{-1}$, about $1267 \pm 2 \text{ cm}^{-1}$, about $1338 \pm 2 \text{ cm}^{-1}$, about $1353 \pm 2 \text{ cm}^{-1}$, about $1418 \pm 2 \text{ cm}^{-1}$, about $2364 \pm 2 \text{ cm}^{-1}$, about $2787 \pm 2 \text{ cm}^{-1}$ and about $2828 \pm 2 \text{ cm}^{-1}$.

[0284] Another reference aspect of the present invention relates to a process for the production of the crystalline form L described above comprising the step of

1. (a) drying of the crystal crystalline form B according to the invention under vacuum,

preferably at a vacuum of at most 900 mbar, more preferably at a vacuum of at most 500 mbar, still more preferably at a vacuum of at most 300 mbar, yet more preferably at a vacuum of at most 200 mbar, and most preferably at a vacuum of at most 100 mbar.

[0285] Accordingly, the crystalline form L may be obtained according to the process for the production of the crystalline form B as described above, wherein the drying step (c-3) or respectively (d-4) takes place under vacuum, i. e. the respective drying step may be replaced by the step (a).

[0286] A further reference aspect of the present invention relates to a crystalline form L that is obtainable by the process as described above.

[0287] In some embodiments, the crystalline forms make it possible to obtain (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4b]indol]-4-amine in the form of the free base with high yields and high purity. These forms are further distinguished in that they have fundamentally different properties, which may provide advantages.

[0288] In some embodiments, the crystalline forms of the present invention are characterized by their simplicity for handling and metering of the active ingredient, i.e. of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4b]indol]-4-amine.

[0289] For the purpose of the specification, the term "active ingredient" preferably refers to the pharmacologically active ingredient (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4b]indol]-4-amine and its physiologically acceptable salts. Unless expressly stated otherwise, it preferably refers to the free base.

[0290] In some embodiments, it has been surprisingly found that (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4b]indol]-4-amine is

capable of forming two anhydrate forms (crystalline forms A and I), two different hydrates (crystalline forms B and H), a partly dehydrated form (crystalline form L) and five different solvates with organic solvents (crystalline forms C, D, E, F, G).

[0291] In some embodiments, it has been surprisingly found that crystalline form A is thermodynamically stable at a relative humidity equal or below 60%. In some embodiments, it has been found that crystalline form B is the most thermodynamically stable form at higher relative humidity (r. h. $\geq$ 40%).

[0292] In some embodiments, it has been surprisingly found that the crystalline forms according to the invention are useful for the separation of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine from its diastereomer, i.e. (1s,4s)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine. Chemical synthesis of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine typically yields mixtures of both diastereomers in various ratios and there is a demand for methods that allow easy and efficient purification of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine.

[0293] Another aspect of the invention relates to a method for the separation of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine from (1s,4s)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine comprising the process for obtaining a crystalline form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine according to the invention as described above, preferably for obtaining any of crystalline forms C, D, E or F.

[0294] Another aspect of the invention relates to a method for the preparation or for the purification of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine, the free base or a physiologically acceptable salt thereof, the method comprising the steps of

1. (i) providing a mixture of diastereomers comprising (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine and (1s,4s)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine; preferably with an excess of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine; and
2. (ii) employing said mixture of diastereomers obtained in step (i) as starting material and preparing therefrom a crystalline form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine selected from the group consisting of crystalline forms C, D, E and F as described above; and
3. (iii) separating the crystalline form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine obtained in step (ii) from the remainder of the mixture of diastereomers; preferably by precipitation and filtering;

and

4. (iv) optionally, converting the crystalline form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine into another crystalline form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine obtained in step (iii), preferably by drying, preferably into crystalline form A.

[0295] According to the invention, step (ii) involves the sub-steps of

(ii-a) dissolving said mixture of diastereomers in a solvent or solvent mixture comprising dimethylsulfoxide or another dipolar aprotic solvent;

(ii-b) adding an alcohol selected from the group consisting of methanol, ethanol, 1-propanol and 2-propanol, more preferably 2-propanol;

(ii-c) precipitating the alcoholate of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine into crystalline form C, D, E or F according to the invention; and

(ii-d) separating the precipitate obtained in step (ii-c) from the remainder of the solution containing (1s,4s)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine.

[0296] It has been surprisingly found that (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine forms crystalline alcoholates under conditions where its diastereomer, (1s,4s)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine, does not form alcoholates or forms alcoholates having a substantially different solubility, thereby allowing for easy and efficient separation of diastereomers. Further, it has been found that the precipitation can be effected by providing solutions of both diastereomers in dimethylsulfoxide (DMSO) and adding the alcohol as antisolvents. There is indication that surprisingly, the crystalline alcohol solvate of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine is formed without simultaneously forming significant amounts of crystalline dimethylsulfoxide solvate. This is advantageous for several reasons, particularly in view of the high boiling point of DMSO - there is no need to evaporate DMSO from the precipitate.

[0297] Another aspect of the invention relates to the crystalline form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine that is obtainable by the method described above.

[0298] Mixtures of the crystalline forms A, B, C, D, E, F, G, H, I and L, preferably mixtures of

two of these crystalline forms, are also included within the scope of the present invention.

[0299] For example, such mixtures of two crystalline forms may be obtained from crystalline form A, B, C, D, E, F, G, H, I or L during a crystallization process (e.g. cooling or evaporation) or respectively during a separation process (e.g. filtration), or respectively during a process where heat is applied (e.g. drying), or respectively during a process where mechanical energy is inserted (e.g. milling or grinding).

[0300] Furthermore, such mixtures of two crystalline forms may be obtained from crystalline form A, B, C, D, E, F, G, H, I or L by a partial uptake of hydrate water or respectively by a partial loss of hydrate water, or respectively by a solvent/water exchange.

[0301] The modifications A, B, C, D, E, F, G, H, I and L according to the invention may optionally also form co-crystals and/or clathrates. These are all included within the scope of the present invention.

[0302] In a preferred embodiment, the crystalline form according to the invention is subsequently converted into an amorphous form.

[0303] Suitable methods for the preparation of amorphous forms are known to a person skilled in the art. For example, amorphous forms or amorphous mixtures may be obtained by means of the following methods or combinations thereof:

1. i) precipitation from solution,
2. ii) lyophilization,
3. iii) spray drying,
4. iv) melts extrusion,
5. v) flash evaporation,
6. vi) quench cooling of the melt,
7. vii) grinding at ambient or liquid nitrogen temperatures,
8. viii) working under protection of an inert atmosphere (e.g. gaseous nitrogen or argon),
and/or
9. ix) using capillary crystallization technology.

[0304] Another aspect of the invention relates to an amorphous form, preferably to an amorphous form that is obtainable by any of the above methods or combinations thereof.

[0305] Another aspect of the invention relates to a composition comprising a mixture of at least two crystalline forms as described herein; or a mixture of at least one crystalline form as described herein with an amorphous form; or a mixture of at least one crystalline form as described herein with a salt of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro-[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine with sulfuric acid, preferably the hydrogen sulfate salt; in any mixing ratio.

[0306] In a preferred embodiment, the pharmaceutical composition according to the invention comprises a crystalline anhydrate of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine as well as crystalline solvate of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine, preferably an alcohol solvate, such as an alcohol solvate for example wherein the alcohol is selected from ethanol and isopropanol (in some especially preferred embodiments isopropanol).

[0307] Preferably, the content of the crystalline anhydrate of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine relative to the total content of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine (anhydrate(s) + solvate(s)) is at least 40 wt.-%, more preferably at least 60 wt.-%, still more preferably at least 80 wt.-%, yet more preferably at least 90 wt.-%, even more preferably at least 95 wt.-%, most preferably at least 99 wt.-%, and in particular at least 99.5 wt.-%.

[0308] Preferably, the content of the crystalline anhydrate of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine relative to the total content of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine (anhydrate(s) + solvate(s)) is at most 99.5 wt.-%, more preferably at most 99 wt.-%, still more preferably at most 95 wt.-%, yet more preferably at most 90 wt.-%, even more preferably at most 80 wt.-%, most preferably at most 60 wt.-%, and in particular at most 40 wt.-%.

[0309] In another preferred embodiment, the pharmaceutical composition according to the invention comprises crystalline form A of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine as well as crystalline form I of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine.

[0310] In a preferred embodiment, the relative weight content of crystalline form A is greater than the relative weight content of crystalline form I. In another preferred embodiment, the relative weight content of crystalline form I is greater than the relative weight content of crystalline form A.

[0311] Preferably, the content of the crystalline form A of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine relative to the total content of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine (crystalline form A + crystalline form I) is at least 40 wt.-%, more preferably at least 60 wt.-%, still more preferably at least 80 wt.-%, yet more preferably at least 90 wt.-%, even more preferably at least 95 wt.-%, most preferably at least 99 wt.-%, and in particular at least 99.5 wt.-%.

[0312] Preferably, the content of the crystalline form A of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine relative to the total content of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine (crystalline form A + crystalline form I) is at most 99.5 wt.-%, more preferably at most 99 wt.-%, still more preferably at most 95 wt.-%, yet more preferably at most 90 wt.-%, even more preferably at most 80 wt.-%, most preferably at most 60 wt.-%, and in particular at most 40 wt.-%.

[0313] In still another preferred embodiment, the pharmaceutical composition according to the invention additionally comprises a salt of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine with sulfuric acid, preferably the hydrogen sulfate salt or the sulfate salt, more preferably the hydrogen sulfate salt. Thus, according to this embodiment, the pharmaceutical composition comprises a mixture of both, the crystalline form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine (free base), preferably selected from the group consisting of crystalline forms A, B, C, D, E, F, G, H, I and L, as well as a sulfuric acid salt thereof, preferably the hydrogen sulfate salt.

[0314] Preferably, the total content of the sulfuric acid salt of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in the pharmaceutical composition, dosage form or active ingredient, such as the hydrogen sulfate salt, is at most 2000 ppm, more preferably at most 1000 ppm, still more preferably at most 750 ppm, yet more preferably at most 500 ppm, even more preferably at most 250 ppm, most preferably at most 100 ppm, and in particular at most 50 ppm, relative to the total amount (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine (free base + salts).

[0315] Preferably, the total content of the sulfuric acid salt of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine in the pharmaceutical composition, dosage form or active ingredient, preferably the hydrogen sulfate salt, is within the range of from 1 ppm to 500 ppm, more preferably 4 ppm to 440 ppm, still more preferably 7 ppm to 380 ppm, yet more preferably 10 ppm to 300 ppm, even more preferably 13 ppm to 220 ppm, most preferably 17 ppm to 140 ppm, and in particular 20 ppm to 60 ppm, relative to the total amount (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine (free base + salts).

[0316] Suitable methods for determining the content of the sulfuric acid salt of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine are known to those skilled in the art and include e.g. XRPD, elemental analysis, Raman spectroscopy, infrared spectroscopy, chromatographic methods, NMR spectroscopy, thermal analysis, electrophoresis, atom absorption spectroscopy, energy dispersive X-ray spectroscopy. Thermal methods comprise, among others, e.g. DSC, TGA, modulated temperature DSC, high-speed DSC, melting point, hot-stage XRPD, hot-stage microscopy, heat of solution, microthermal analysis, calorimetry, micro-calorimetry.

[0317] In yet another preferred embodiment, the pharmaceutical composition according to the invention comprises a crystalline form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine, preferably selected from the group consisting of crystalline forms A, B, C, D, E, F, G, H, I and L, as well as an amorphous form of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine.

[0318] Preferably, the degree of crystallinity, i.e. the content of crystalline form(s) of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine relative to the total content of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine (crystalline form(s) + amorphous form(s)) is at least 40 wt.-%, more preferably at least 60 wt.-%, still more preferably at least 80 wt.-%, yet more preferably at least 90 wt.-%, even more preferably at least 95 wt.-%, most preferably at least 99 wt.-%, and in particular at least 99.5 wt.-%.

[0319] Preferably, the degree of crystallinity, i.e. the content of crystalline form(s) of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine relative to the total content of (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine (crystalline form(s) + amorphous form(s)) is at most 99.5 wt.-%, more preferably at most 99 wt.-%, still more preferably at most 95 wt.-%, yet more preferably at most 90 wt.-%, even more preferably at most 80 wt.-%, most preferably at most 60 wt.-%, and in particular at most 40 wt.-%.

[0320] In another aspect the present invention relates to methods of treating pain, comprising administering a crystalline form as described herein to a patient in need thereof (for example, a patient who has been diagnosed with a pain disorder).

[0321] In some preferred embodiments, the present invention relates to a method of treating pain in a patient diagnosed with a pain disorder comprising orally administering to the patient an effective amount of a pharmaceutical composition comprising one or more additives or adjuvants and an active ingredient comprising the crystalline form disclosed herein. In some embodiments, the pharmaceutical composition comprises 40 ± 20 μg of the active ingredient. In some embodiments, the pharmaceutical composition comprises 400 ± 50 μg of the active ingredient. In some embodiments, the pharmaceutical composition comprises 200 ± 50 μg of the active ingredient. In some embodiments, the pharmaceutical composition comprises 600 ± 50 μg of the active ingredient. In some preferred embodiments, the crystalline form is present in the active ingredient in substantially pure form (for example, the active ingredient comprises at least about 50%, about least about 60%, about least about 70%, about least about 80%, about least about 90%, about least about 95%, or even about least about 99% of the crystalline form).

[0322] In another aspect the present invention relates to methods of treating pain, comprising administering a pharmaceutical composition that comprises a crystalline form as described

herein to a patient in need thereof (for example, a patient who has been diagnosed with a pain disorder).

[0323] The term pain as used herein preferably includes but is not limited to pain selected from the group consisting of inflammatory pain, postoperative pain, neuropathic pain, diabetic neuropathic pain, acute pain, chronic pain, visceral pain, migraine pain and cancer pain.

[0324] In some preferred embodiments, the crystalline form according to the invention is for use in the treatment of acute, visceral, neuropathic or chronic pain (cf. WO 2008/040481).

[0325] In another aspect the present invention relates to a pharmaceutical composition comprising a crystalline form as described herein and optionally one or more suitable additives and/or adjuvants such as described below.

[0326] In some preferred embodiments, the pharmaceutical composition comprises between about 0.001% by weight and about 40% by weight of one or more of the crystalline forms described herein. In some preferred embodiments, the pharmaceutical composition comprises between about 0.001% by weight and about 20% by weight of one or more of the crystalline forms described herein. In some preferred embodiments, the pharmaceutical composition comprises between about 0.001 % by weight and about 10% by weight of one or more of the crystalline forms described herein. In some preferred embodiments, the pharmaceutical composition comprises between about 0.001% by weight and about 5% by weight of one or more of the crystalline forms described herein. In some preferred embodiments, the pharmaceutical composition comprises between about 0.001% by weight and about 1% by weight of one or more of the crystalline forms described herein. In some preferred embodiments, the pharmaceutical composition comprises between about 0.01% by weight and about 1% by weight of one or more of the crystalline forms described herein. In some preferred embodiments, the pharmaceutical composition comprises between about 0.01% by weight and about 1% by weight of one or more of the crystalline forms described herein.

[0327] Preferably said pharmaceutical composition may be used for the treatment of pain.

[0328] In still another aspect the present invention relates to a medicament comprising a crystalline form as described herein, preferably a pharmaceutical composition as described herein. In a preferred embodiment, the medicament is a solid drug form. The medicament is preferably manufactured for oral administration. However, other forms of administration are also possible, e.g. for buccal, sublingual, transmucosal, rectal, intralumbal, intraperitoneal, transdermal, intravenous, intramuscular, intragluteal, intracutaneous and subcutaneous application.

[0329] Depending on the configuration, the medicament (dosage form) preferably contains suitable additives and/or adjuvants. Suitable additives and/or adjuvants in the sense of the invention are all substances known to a person skilled in the art for the formation of galenic formulations. The choice of these adjuvants and also the quantities to be used are dependent

on how the medication is to be administered, i.e. orally, intravenously, intraperitoneally, intradermally, intramuscularly, intranasally, buccally or locally.

[0330] In some preferred embodiments, the dosage form comprises $40 \pm 35 \mu\text{g}$, more preferably $40 \pm 30 \mu\text{g}$, still more preferably $40 \pm 25 \mu\text{g}$, yet more preferably $40 \pm 20 \mu\text{g}$, even more preferably $40 \pm 15 \mu\text{g}$, most preferably $40 \pm 10 \mu\text{g}$, and in particular $40 \pm 5 \mu\text{g}$ of one or more of the crystalline forms described herein. In some other preferred embodiments, the dosage form comprises $400 \pm 375 \mu\text{g}$ or $400 \pm 350 \mu\text{g}$, more preferably $400 \pm 300 \mu\text{g}$, still more preferably $400 \pm 250 \mu\text{g}$, yet more preferably $400 \pm 200 \mu\text{g}$, even more preferably $400 \pm 150 \mu\text{g}$, most preferably $40 \pm 100 \mu\text{g}$, and in particular $400 \pm 50 \mu\text{g}$ of one or more of the crystalline forms described herein. In some other preferred embodiments, the dosage form comprises $40 \pm 5 \mu\text{g}$ of the crystalline or (or of an active ingredient comprising the crystalline form). In some other preferred embodiments, the dosage form comprises $100 \pm 10 \mu\text{g}$ of the crystalline or (or of an active ingredient comprising the crystalline form). In some other preferred embodiments, the dosage form comprises $200 \pm 50 \mu\text{g}$ of the crystalline or (or of an active ingredient comprising the crystalline form). In some other preferred embodiments, the dosage form comprises $400 \pm 50 \mu\text{g}$ of the crystalline or (or of an active ingredient comprising the crystalline form). In some other preferred embodiments, the dosage form comprises $600 \pm 50 \mu\text{g}$ of the crystalline or (or of an active ingredient comprising the crystalline form).

[0331] Preparations suitable for oral administration are those in the form of tablets, chewable tablets, lozenges, capsules, granules, drops, liquids or syrups, and those suitable for parenteral, topical and inhalatory administration are solutions, suspensions, easily reconstituted dry preparations and sprays. A further possibility is suppositories for rectal administration. The application in a depot in dissolved form, a patch or a plaster, possibly with the addition of agents promoting skin penetration, are examples of suitable percutaneous forms of application.

[0332] Examples of adjuvants and additives for oral forms of application are disintegrants, lubricants, binders, fillers, mould release agents, possibly solvents, flavourings, sugar, in particular carriers, diluents, colouring agents, antioxidants etc.

[0333] Waxes or fatty acid esters, amongst others, can be used for suppositories and carrier substances, preservatives, suspension aids etc. can be used for parenteral forms of application.

[0334] Adjuvants can be, for example: water, ethanol, 2-propanol, glycerine, ethylene glycol, propylene glycol, polyethylene glycol, polypropylene glycol, glucose, fructose, lactose, saccharose, dextrose, molasses, starch, modified starch, gelatine, sorbitol, inositol, mannitol, microcrystalline cellulose, methyl cellulose, carboxymethyl-cellulose, cellulose acetate, shellac, cetyl alcohol, polyvinylpyrrolidone, paraffins, waxes, natural and synthetic rubbers, acacia gum, alginates, dextran, saturated and unsaturated fatty acids, stearic acid, magnesium stearate, zinc stearate, glyceryl stearate, sodium lauryl sulphate, edible oils, sesame oil, coconut oil, peanut oil, soybean oil, lecithin, sodium lactate, polyoxyethylene and propylene fatty acid

esters, sorbitane fatty acid esters, sorbic acid, benzoic acid, citric acid, ascorbic acid, tannic acid, sodium chloride, potassium chloride, magnesium chloride, calcium chloride, magnesium oxide, zinc oxide, silicon dioxide, titanium oxide, titanium dioxide, magnesium sulphate, zinc sulphate, calcium sulphate, potash, calcium phosphate, dicalcium phosphate, potassium bromide, potassium iodide, talc, kaolin, pectin, crosspovidon, agar and bentonite.

[0335] The production of these medicaments and pharmaceutical compositions is conducted using means, devices, methods and processes that are well known in the art of pharmaceutical technology, as described, for example, in "Remington's Pharmaceutical Sciences", A.R. Gennaro, 17th ed., Mack Publishing Company, Easton, Pa. (1985), in particular in part 8, chapters 76 to 93.

[0336] Thus, for example, for a solid formulation such as a tablet, the active substance of the drug can be granulated with a pharmaceutical carrier substance, e.g. conventional tablet constituents such as cornstarch, lactose, saccharose, sorbitol, talc, magnesium stearate, dicalcium phosphate or pharmaceutically acceptable rubbers, and pharmaceutical diluents such as water, for example, in order to form a solid composition that contains the active substance in a homogenous dispersion. Homogenous dispersion is understood here to mean that the active substances are uniformly dispersed throughout the composition, so that this can be readily divided into identically effective standard dosage forms such as tablets, capsules, lozenges. The solid composition is then divided into standard dosage forms. The tablets or pills can also be coated or otherwise compounded to prepare a slow release dosage form. Suitable coating agents include polymeric acids and mixtures of polymeric acids with materials such as shellac, cetyl alcohol and/or cellulose acetate, for example.

[0337] In one embodiment of the present invention the crystalline form as described herein is present in immediate release form.

[0338] In another embodiment of the present invention the crystalline form as described herein is at least partially present in controlled-release form. In particular, the active ingredient can be released slowly from preparations that can be applied orally, rectally or percutaneously.

[0339] The medicament can preferably be manufactured for administration once daily, twice daily (bid), or three times daily, the once daily or twice daily administration (bid) being preferred.

[0340] The term "controlled release" as used herein refers to any type of release other than immediate release such as delayed release, sustained release, slow release, extended release and the like. These terms are well known to any person skilled in the art as are the means, devices, methods and processes for obtaining such type of release.

[0341] In another embodiment of the present invention

- the medicament is manufactured for oral administration; and/or

- the medicament is a solid and/or compressed and/or film-coated drug form; and/or
- the medicament releases the crystalline form as described herein slowly from a matrix; and/or
- the medicament contains the crystalline form in a quantity of 0.001 to 99.999% by wt., more preferred 0.1 to 99.9% by wt., still more preferred 1.0 to 99.0% by wt., even more preferred 2.5 to 80% by wt., most preferred 5.0 to 50% by wt. and in particular 7.5 to 40% by wt., based on the total weight of the medicament; and/or
- the medicament contains a pharmaceutically compatible carrier and/or pharmaceutically compatible adjuvants; and/or
- the medicament has a total mass in the range of 25 to 2000 mg, more preferred 50 to 1800 mg, still more preferred 60 to 1600 mg, more preferred 70 to 1400 mg, most preferred 80 to 1200 mg and in particular 100 to 1000 mg; and/or
- the medicament is selected from the group comprising tablets, capsules, pellets and granules.

[0342] The medicament can be provided as a simple tablet and as a coated tablet (e.g. as film-coated tablet or lozenge). The tablets are usually round and biconvex, but oblong forms are also possible. Granules, spheres, pellets or microcapsules, which are contained in sachets or capsules or are compressed to form disintegrating tablets, are also possible.

[0343] In yet another one of its aspects, the present invention relates to the use of the crystalline form as described herein for the production of a medicament. Preferably said medicament is suitable for the treatment of pain.

[0344] In still another one of its aspects, the present invention relates to the use of the crystalline form as described herein for the treatment of pain.

[0345] The phrase "consisting essentially of" when used in reference to the pharmaceutical composition means that the composition contains no active pharmaceutical ingredients other than those specified, but that it may contain additional inactive components or excipients. For example, if a pharmaceutical composition is described as consisting essentially of (1*r*,4*r*)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro-[cyclohexane-1,1'-pyrano-[3,4,*b*]indol]-4-amine, it should be understood that the pharmaceutical composition contains (1*r*,4*r*)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro-[cyclohexane-1,1'-pyrano-[3,4,*b*]indol]-4-amine (*i.e.*, crystalline and/or amorphous form(s) thereof) and no other active pharmaceutical ingredients but that the dosage form may contain any number of additional inactive components or excipients. The phrase "consisting of", when used in reference to the pharmaceutical composition, means that the composition contains no other active pharmaceutical ingredient other than that specified but may contain additional components that are unrelated to the invention and/or impurities ordinarily associated with the recited components. Similarly, when the phrase "consisting of" is used to define the active ingredient, it means that the active ingredient contains substantially no other crystalline form other than that

specified but may contain additional components that are unrelated to the invention and/or impurities ordinarily associated with the recited components.

[0346] Furthermore, the present invention relates to a method for treating pain in a patient, preferably in a mammal, which comprises administering an effective amount of a crystalline form as described herein to a patient.

EXAMPLES

[0347] The following examples serve to explain the invention in more detail, but should not be interpreted as restrictive. Examples being drawn to crystalline forms A, C, D, E and F are inventive examples.

[0348] The following abbreviations are used in the examples:

iBuOAc

iso-butyl acetate

1BuOH

n-butanol (1-butanol)

DMSO

dimethylsulfoxid

EtOAc

ethyl acetate

EtOH

ethanol

Ex

example

FT-Raman

Fourier transformation Raman spectroscopy

IPE

diisopropyl ether

Δm

change in mass

MeCN

acetonitril

MEK

2-butanone

MeOH

methanol

min

minute

NMP

N-methyl-2-pyrrolidone

1PrOH	n-propanol (1-propanol)
2PrOH	iso-propanol (2-propanol)
PXRD	powder x-ray diffraction
r.h.	relative humidity
RT	room temperature, preferably 20-25°C
SCXRD	single crystal X-ray diffraction
sec	seconds
t	time (duration)
TBME	tert-butyl methyl ether
TG-FTIR	thermogravimetry coupled with Fourier transform infrared spectroscopy
THF	tetrahydrofuran
XRPD	X-ray powder diffraction

[0349] Unless otherwise specified, solvent mixtures are always volume/volume.

A) Synthesis of crystalline form A

[0350] 100 mg (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano[3,4,b]indol]-4-amine [crystalline form D according to D)] was suspended in 0.5 mL TBME. The suspension was stirred at RT for six days. The resulting solid was filtered off and dried in air. A crystalline solid of crystalline form A was obtained and characterized by FT Raman, TG-FTIR and PXRD.

B) Synthesis of crystalline form B

[0351] 100 mg crystalline form D [according to D)] was suspended in 0.5 mL THF/H₂O. The suspension was stirred at RT for six days. The resulting solid was filtered off and dried in air. A crystalline solid of crystalline form B was obtained and characterized by FT Raman, TG-FTIR

and PXRD.

C) Synthesis of crystalline form C

[0352] 107.5 g 5-Fluorotryptophol were charged into a vessel. Subsequently, 138.7 g DMAPh-cyclohexanon (4-(dimethylamino)-4-phenylcyclohexanone) and 8.40 kg dichloromethane were added. A dropping funnel was mounted on the vessel and the vessel was flushed with gaseous nitrogen. The mixture was stirred and heated to 39.4°C. 160.0 g TMS-triflate (trifluoromethane sulfonic acid trimethylsilylester) were filled under nitrogen atmosphere into the funnel and subsequently 0.2 kg dichloromethane were added to the dropping funnel and mixed with the TMS-triflate. This mixture was dosed into the vessel over 2 hrs. The reaction mixture was then stirred for 23 hrs at 40 ± 5 °C and 40 hrs at 22 ± 2 °C. A 1 N sodium hydroxide solution was prepared by diluting 224 g of a sodium hydroxide solution (30%) with 1.45 kg deionised water. This 1 N sodium hydroxide solution was added at 21 - 22 °C to the reaction mixture until the product (1r,4r)-6'-fluoro-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexane-1,1'-pyrano-[3,4,b]indol]-4-amine crystallized as a yellowish solid. The solution was cooled from 21.5 °C to 4.3°C in 1 hr and stirred for 3.5 hrs at 1 to 5°C. The resulting suspension was transferred to a nutsche, filtered off by applying pressure and washed 2 times with 0.65 L Ethanol for 15 - 20 min. Pressure was applied after filtration for further 5 min.

Purity of the product (HPLC): 99.5 %

The product was then dried in vacuum (50°C, 17.5 hrs, 2 bar) until the mass remained constant.

Yield: 206.9 g, 91%

D) Synthesis of crystalline form D

[0353] (4-(dimethylamino)-4-phenylcyclohexanone (3 g, 13.82 mmol), 2-(5-fluoro-1H-indol-3-yl)ethanol (2.47 g, 13.82 mmol) and 150 mL dichloromethane were charged to a flask at 0°C. A solution of trifluoromethane sulfonic acid trimethylsilylester (3 mL, 15.5 mmol) in 3 mL dichloromethane were added quickly. The reaction mixture changed color to violet and the temperature rose to 10°C. The reaction mixture was cooled in an ice bath and stirred for 20 min. Meanwhile a solid precipitated. The ice bath was removed and the reaction mixture was stirred for 3 to 3.5 hours at room temperature. Subsequently 50 mL of NaOH (1N) were added and the reaction mixture was stirred further 10 min. The colour changed to yellow and a solid precipitated. In order to complete the precipitation the reaction mixture (two liquid phases) was stirred for further 20 min while cooled in an ice bath. Eventually the solid was filtered off. The resulting solid (4.2 g) was subsequently recrystallized in 800 mL 2-Propanol. Yield: 3.5 g.

[0354] To enhance the yield, the liquid (Water and Dichloromethane) filtrate was separated. The aqueous solution was extracted 3 times with 20 mL Dichloromethane. The organic phases were united and dried with MgSO₄ and subsequently the solvent was stripped off until dryness.

The resulting solid (1,7 g) was subsequently recrystallized under reflux in 800 mL 2-Propanol.

Crystallization tests

Example 1

[0355] Crystalline form C [according to C)] was suspended in different solvents and the suspension was stirred at RT for eight days. The resulting solids were filtered off, dried in air and characterized by FT Raman. One sample of each obtained form was further characterized by TG-FTIR and PXRD.

[0356] The detailed experimental conditions and results are summarized in the table here below. For characterization details of the obtained forms see Section "Analysis".

Table 1

Ex.	solvent	amount crystalline form C/ amount solvent	t [days]	result	TG-FTIR	crystalline modification
1-1	acetone	120 mg / 5 mL	8	white crystalline solid	solvent free	A
1-2	CH ₂ Cl ₂	120 mg / 5 mL	8	white crystalline solid	-	A
1-3	EtOAc	120 mg / 5 mL	8	white crystalline solid	-	A
1-4	toluene	120 mg / 5 mL	8	yellowish crystalline solid	-	A
1-5	MeCN	120 mg / 5 mL	8	yellowish crystalline solid	-	A
1-6	EtOH	120 mg / 5 mL	8	yellowish crystalline solid	contains EtOH	C
1-7	MeOH	120 mg / 5 mL	8	yellowish crystalline solid	contains MeOH	E
1-8	2PrOH	120 mg / 5 mL	8	white crystalline solid	contains 2PrOH	D

Ex.	solvent	amount crystalline form C/ amount solvent	t [days]	result	TG-FTIR	crystalline modification
1-9	H ₂ O	120 mg / 5 mL	8	yellowish crystalline solid	contains H ₂ O	B
1-10	TBME	120 mg / 5 mL	8	white crystalline solid	-	A
1-11	1BuOH	150 mg / 4 mL	4	white crystalline solid	-	A
1-12	1PrOH	150 mg / 4 mL	4	yellowish crystalline solid	contains 1PrOH	F
1-13	iBuOAc	150 mg / 4 mL	4	white crystalline solid	-	A
1-14	NMP	150 mg / 4 mL	4	no precipitate was obtained		-
1-15	DMSO	150 mg / 4 mL	4	yellowish crystalline solid	contains DMSO	G
1-16	H ₂ O	400 mg / 5mL	2	yellowish crystalline solid	contains H ₂ O, EtOH (traces)	"K"
"K": a new FT Raman spectrum was obtained, which slightly differed to that of crystalline form B						

[0357] 1-17) The new "crystalline form K" (sample of Ex. 1-16) was further analyzed by the following experiments:

100 mg of "crystalline form K" were stored at RT in vacuum (100 mbar) before being characterized by FT Raman, TG-FTIR and PXRD. The analyses revealed the presence of a new crystalline form (crystalline form L), a desolvated form. After the data of crystalline form L were available, the starting material ("crystalline form K") was identified as a mixture of crystalline forms B and L by FT Raman.

[0358] TG-FTIR after four weeks of storage (RT, 100 mbar) revealed that the sample still contained H₂O. Additionally, the sample contained EtOH, indicating that the used hydrate (sample 1-16) was contaminated with traces of EtOH from the starting material. Whereas most

of the water was removed in vacuum (reduction from 9 % to 1.5 % H₂O), the remaining EtOH content changed less (reduction from 1.8 % to 0.8 % EtOH). Neither FT Raman nor PXRD showed any indication for the presence of form C (EtOH solvate).

[0359] Crystalline form L is probably metastable. At ambient conditions (RT, 40-60% r. h.) crystalline form L slowly reconverts into crystalline form B.

Example 2

[0360] Solutions of crystalline form C [according to C), 50 mg] were prepared in THF, 1,4-dioxane and DMSO. 6 mL precipitant (H₂O, EtOH, TBME, IPE) was added quickly.

[0361] A second set of experiments was performed using acetone, MEK, CH₂Cl₂ and THF as solvents and heptane, hexane and EtOH as precipitants (antisolvents). Solvents and precipitants were selected based on the approximate solubility determination in phase A. The resulting solids were filtered off, dried in air and characterized by FT Raman.

[0362] The detailed experimental conditions and results are summarized in the table here below. For characterization details of the obtained forms see Section "Analysis".

Table 2

Ex.	solvent/precipitant	amounts solvent/precipitant	result/characterization	crystalline form
2-1	THF/H ₂ O (MeOH)	3 mL/6 mL	crystalline solid	E
2-2	THF/EtOH	3 mL/6 mL	crystalline solid	C
2-3	THF/TBME	3 mL/6 mL	clear solution	-
2-4	THF/IPE	3 mL/6 mL	clear solution	-
2-5	1,4-dioxane/ H ₂ O (MeOH)	3 mL/6 mL	crystalline solid	E
2-6	1,4-dioxane/ EtOH	3 mL/6 mL	crystalline solid	C
2-7	1,4-dioxane/ TBME	3 mL/6 mL	crystalline solid	-
2-8	1,4-dioxane/ IPE	3 mL/6 mL	crystalline solid	-
2-9	DMSO/ H ₂ O	3 mL/6 mL	crystalline solid	B
2-10	DMSO/EtOH (MeOH)	3 mL/6 mL	crystalline solid	E
2-11	DMSO/TBME	3 mL/6 mL	clear solution	-
2-12	DMSO/IPE	3 mL/6 mL	clear solution	-
2-13	acetone/ n-heptane	12 mL / 20 mL	clear solution	-

Ex.	solvent/precipitant	amounts solvent/precipitant	result/characterization	crystalline form
2-14	MEKI n-heptane	10 mL / 20 mL	clear solution	-
2-15	CH ₂ Cl ₂ / n-heptane	10 mL / 20 mL	clear solution	-
2-16	THF/n-heptane + EtOH	3 mL / 15 mL / 5 mL	crystalline solid	C
2-17	acetone/n-hexane	12 mL / 20 mL	clear solution	-
2-18	MEK/n-hexane	10 mL / 20 mL	clear solution	-
2-19	CH ₂ Cl ₂ / n-hexane	10 mL / 20 mL	clear solution	-
2-20	THF/ n-hexane	3 mL/ 15 mL	crystalline solid	B
(MeOH): the samples were washed with 0.5 mL of MeOH				

[0363] Enough precipitate for characterization was obtained from experiments with H₂O or EtOH as precipitant and from THF/hexane. Whereas the addition of water resulted immediate precipitation, in case of EtOH or n-alkanes clear solutions were obtained. The precipitation started with a delay of 10 sec (THF/EtOH) to several minutes (DMSO/EtOH; THF/n-hexane; THF/n-heptane+EtOH).

[0364] Some obtained solids were very wet and hard to filter. The samples were washed with 0.5 mL of MeOH in order to improve filterability. The filtration was accelerated, but the solids were converted into the MeOH solvate (crystalline form E) indicating a very short conversion time.

[0365] From THF/n-hexane a hydrate was obtained (identified by FT Raman). The hydrate formation must be caused by residual water in the used solvents or by the high humidity on the preparation day during filtration.

[0366] When TBME or IPE was used as precipitant no or very small amounts of solid were obtained (occurring after days), not enough for characterization.

Example 3

[0367] Crystalline form C [according to C)] was dissolved in different solvents. The solvent was evaporated at RT under nitrogen flow (8-10 mL/min). Clear solutions or suspensions obtained

from precipitation experiments with not sufficient material for characterization were used for further evaporation experiment at about 50 °C.

[0368] The evaporation time was adjusted as short as possible in order to advantage metastable forms. Because of the rather low solubility in most solvent or because of low vapor pressure the evaporation time was at least one day.

[0369] Crystalline solids were obtained in all cases and characterized by FT Raman. In most cases material of crystalline form A was obtained.

[0370] Forms with new Raman spectra were obtained from dioxane (or mixtures containing dioxane, crystalline form H, hemi-hydrate) and from CH₂Cl₂/heptane (crystalline form I, ansovate 2, containing traces of water). These two samples were further characterized by TG-FTIR and PXRD.

[0371] The detailed experimental conditions and results are summarized in the table here below. For characterization details of the obtained forms see Section "Analysis".

Table 3

Ex.	solvent	T (evap.)	amount crystalline form C/ amount solvent	TG-FTIR	crystalline modification
3-1	THF	RT	40 mg / 10 mL	-	A
3-2	1,4-dioxane	RT	40 mg / 10 mL	contains H ₂ O	H
3-3	MEK	RT	40 mg / 10 mL	-	A
3-4	acetone	RT	40 mg / 10 mL	-	A
3-5	CH ₂ Cl ₂	RT	40 mg / 10 mL	-	A
3-6	THF/TBME	50 °C	50 mg / 3 mL / 6 mL	-	A
3-7	THF/IPE	50 °C	50 mg / 3 mL / 6 mL	-	A
3-8	1,4- dioxane/TBME	50 °C	50 mg / 3 mL / 6 mL	-	H
3-9	1,4- dioxane/IPE	50 °C	50 mg / 3 mL / 6 mL	-	H
3-10	CH ₂ Cl ₂ /n- heptane	50 °C	50 mg / 10 mL / 20 mL	contains H ₂ O	I
3-11	CH ₂ Cl ₂ /n- hexane	50 °C	50 mg / 10 mL / 20 mL	-	A
3-12	acetone/ n- heptane	50 °C	50 mg / 12 mL / 20 mL	-	A
3-13	acetone/ n- hexane	50 °C	50 mg / 12 mL / 20 mL	-	A
3-14	MEK/ n-	50 °C	50 mg / 10 mL / 20 mL	-	A

Ex.	solvent	T (evap.)	amount crystalline form C/ amount solvent	TG-FTIR	crystalline modification
	hexane				

Example 4

[0372] The effect of mechanical stress from grinding with an agate mortar was analyzed.

4-1) 30 mg of crystalline form A was grinded in an agate mortar for 10 min. The resulting solid was characterized by FT Raman. No effect was observed.

4-2) 30 mg of crystalline form C was grinded in an agate mortar for 10 min. The resulting solid was characterized by FT Raman. No effect was observed.

Example 5

[0373] Seven experiments with the aim to obtain single crystals of crystalline form A (ansolvate), crystalline form (di-hydrate) and crystalline form (EtOH solvate) were performed. Vapor diffusion and slow cooling of saturated solutions were used as reliable techniques for single crystal growth.

5-1) 17 mL acetone was added to 179 mg crystalline form C [according to C)] and the mixture was stirred at 45 °C for 30 min. The resulting suspension was filtered (hot) and the obtained solution was cooled from 40 °C to 5 °C with 0.5 °C/h and then stored at 5 °C for two weeks. Small crystals were obtained.

5-2) 2 ml THF was added to 179 mg crystalline form C [according to C)] and the mixture was stirred at 45 °C for 30 min. The resulting suspension was filtered (hot) and the obtained solution was cooled from 40 °C to 5 °C with 0.5 °C/h and then stored at 5 °C for two weeks. Small crystals were obtained.

5-3) 9 mL DMSO was added to 179 mg crystalline form C [according to C)] and the mixture was stirred at 45 °C for 30 min. The resulting suspension was filtered (hot) and the obtained solution was cooled from 40 °C to 5 °C with 0.5 °C/h and then stored at 5 °C for two weeks. Small crystals according to crystalline form G were obtained and characterized by FT Raman.

5-4) 12 mL DMSO was added to 179 mg crystalline form C [according to C)] and the mixture was stirred at RT for 30 min. The resulting suspension was filtered. 4 mL of the obtained solution was stored at RT in a saturated H₂O atmosphere for two weeks. Needle-like crystals according to crystalline form G were obtained and characterized by FT Raman and SCXRD.

5-5) 12 mL DMSO was added to 179 mg crystalline form C [according to C)] and the mixture was stirred at RT for 30 min. The resulting suspension was filtered. 4 mL of the obtained solution was stored at RT in a saturated EtOH atmosphere for two weeks. Long, needle-like crystals according to crystalline form C were obtained and characterized by FT Raman and SCXRD.

5-6) 12 mL DMSO was added to 179 mg crystalline form C [according to C)] and the mixture was stirred at RT for 30 min. The resulting suspension was filtered. 4 mL of the obtained solution was stored at RT in a saturated TBME atmosphere for two weeks. No precipitation was observed.

5-7) 2 mL THF was added to 77 mg crystalline form A [according to A)] and the mixture was stirred at RT for 30 min. The resulting suspension was filtered. The obtained solution was stored at RT in a saturated n-hexane atmosphere for two weeks. Small crystals according to crystalline form A were obtained and characterized by FT Raman and SCXRD.

Analysis - XRPD (X-Ray Powder Diffraction) or PXRD (Powder X-Ray Diffraction)

[0374] XRPD analyses were carried out in transmission geometry with a Philips X'pert PW 3040 X-ray powder diffractometer, monochromatised CuK α radiation being used by means of a germanium monocrystal. d-distances were calculated from the 2θ values, the wavelength of 1.54060 Å being taken as basis. The d-value analysis was performed with the software EVA version 10, 0, 0, 0. The CuK α_2 was removed by the software and only lines up to 35° 2θ were listed. In general, the 2θ values have an error rate of $\pm 0.2^\circ$ in 2θ . The experimental error in the d-distance values is therefore dependent on the location of the peak. 2θ values may be converted into d-distance values using Bragg's law.

[0375] The samples were measured without any special treatment other than the application of slight pressure to get a flat surface. An ambient air atmosphere was used.

[0376] Figure 1 shows the superposed PXRD patterns of crystalline forms A, B, C, D, E, F, G, H, I and L.

Crystalline Form A

[0377] Table 4 shows the peak list for crystalline form A. The uncertainty in the 2θ values is $\pm 0.2^\circ$ in 2θ ; rel. I is the relative intensity of the respective peaks. Maximum intensity is 100.

Table 4

2 θ	d value Å	Intensity Cps	rel. I %
7.8	11.3	324	16
8.8	10.1	156	8
9.1	9.7	120	6
10.5	8.4	40	2
11.1	8.0	300	15
11.7	7.6	64	3
12.2	7.3	280	14
13.5	6.5	343	17
15.2	5.8	140	7
16.0	5.5	134	7
16.8	5.3	347	17
17.6	5.1	735	36
18.3	4.9	2044	100
18.6	4.8	1295	63
19.4	4.6	474	23
19.7	4.5	346	17
20.6	4.3	214	11
20.8	4.3	308	15
21.5	4.1	178	9
22.0	4.0	135	7
22.3	4.0	199	10
22.6	3.9	227	11
23.5	3.8	263	13
23.9	3.7	208	10
24.2	3.7	176	9
25.0	3.6	292	14
25.8	3.5	505	25
26.3	3.4	564	28
27.2	3.3	180	9
27.6	3.2	442	22
28.3	3.2	360	18
29.1	3.1	249	12
29.5	3.0	115	6
30.0	3.0	256	13

2 θ	d value Å	Intensity Cps	rel. I %
31.6	2.8	147	7
32.3	2.8	174	9
32.6	2.8	147	7
33.8	2.7	96	5

Crystalline Form B

[0378] Table 5 shows the peak list for crystalline form B. The uncertainty in the 2 θ values is $\pm 0.2^\circ$ in 2 θ ; rel. I is the relative intensity of the respective peaks. Maximum intensity is 100.

Table 5

2 θ	d value Å	Intensity Cps	rel. I %
8.9	9.9	524	36
9.8	9.0	1241	86
10.5	8.4	59	4
12.0	7.4	189	13
14.2	6.2	91	6
14.6	6.1	85	6
15.7	5.6	485	34
16.2	5.5	142	10
16.7	5.3	798	55
17.8	5.0	1440	100
18.4	4.8	466	32
19.2	4.6	672	47
19.7	4.5	503	35
20.0	4.5	330	23
20.4	4.4	413	29
21.4	4.2	159	11
21.8	4.1	566	39
22.6	3.9	168	12
23.5	3.8	124	9
24.1	3.7	715	50
25.1	3.6	352	25
25.4	3.5	267	19
26.0	3.4	451	31

2 θ	d value Å	Intensity Cps	rel. I %
26.7	3.3	190	13
27.1	3.3	322	22
27.9	3.2	144	10
28.1	3.2	297	21
29.2	3.1	271	19
29.7	3.0	182	13
30.3	2.9	204	14
31.1 31.8	2.9	627 111	44 8
32.7	2.7	176	12
32.9	2.7	164	11
33.5	2.7	163	11
33.9	2.6	130	9
34.9	2.6	161	11

Crystalline Form C

[0379] Table 6 shows the peak list for crystalline form C. The uncertainty in the 2 θ values is $\pm 0.2^\circ$ in 2 θ ; rel. I is the relative intensity of the respective peaks. Maximum intensity is 100.

Table 6

2 θ	d value Å	Intensity Cps	rel. I %
7.8	11.4	46	3
9.1	9.7	1188	76
9.5	9.3	939	60
10.4	8.5	52	3
11.1	8.0	55	4
12.2	7.3	97	6
13.5	6.6	86	6
14.3	6.2	292	19
14.8	6.0	156	10
15.3	5.8	118	8
16.1	5.5	143	9
16.8	5.3	772	49
17.5	5.1	238	15
18.2	4.9	1572	100

2 θ	d value Å	Intensity Cps	rel. I %
18.6	4.8	508	32
19.0	4.7	518	33
19.3	4.6	731	47
19.5	4.5	387	25
20.7	4.3	259	17
21.7	4.1	240	15
22.2	4.0	605	39
22.5	3.9	210	13
23.6	3.8	256	16
24.2	3.7	254	16
24.9	3.6	328	21
25.4	3.5	586	37
25.8	3.5	329	21
26.2	3.4	198	13
26.5	3.4	186	12
27.5	3.2	634	40
28.1	3.2	222	14
28.7	3.1	206	13
29.4	3.0	158	10
30.3	3.0	322	21
32.3	2.8	151	10
33.6	2.7	209	13
34.5	2.6	143	9

Crystalline Form D

[0380] Table 7 shows the peak list for crystalline form D. The uncertainty in the 2 θ values is $\pm 0.2^\circ$ in 2 θ ; rel. I is the relative intensity of the respective peaks. Maximum intensity is 100.

Table 7

2 θ	d value Å	Intensity Cps	rel. I %
8.4	10.5	1238	88
8.8	10.1	1207	86
11.2	7.9	67	5
12.1	7.3	122	9

2 θ	d value Å	Intensity Cps	rel. I %
13.7	6.5	76	5
15.0	5.9	760	54
15.2	5.8	760	54
16.1	5.5	114	8
17.0	5.2	699	50
17.6	5.0	1408	100
18.1	4.9	297	21
18.9	4.7	690	49
19.4	4.6	157	11
19.8	4.5	239	17
20.9	4.3	281	20
21.2	4.2	399	28
21.6	4.1	286	20
22.4	4.0	1009	72
22.8	3.9	317	23
23.2	3.8	711	51
24.9	3.6	280	20
25.2	3.5	162	12
25.7	3.5	336	24
26.0	3.4	564	40
26.6	3.4	237	17
27.5	3.2	188	13
28.0	3.2	177	13
28.5	3.1	176	13
29.5	3.0	346	25
30.3	3.0	331	24
30.7	2.9	359	26
31.3	2.9	201	14
31.9	2.8	171	12
32.2	2.8	154	11
32.8	2.7	148	11
33.6	2.7	129	9
34.0	2.6	178	13
34.9	2.6	266	19

Crystalline Form E

[0381] Table 8 shows the peak list for crystalline form E. The uncertainty in the 2θ values is $\pm 0.2^\circ$ in 2θ ; rel. I is the relative intensity of the respective peaks. Maximum intensity is 100.

Table 8

2θ	d value Å	Intensity Cps	rel. I %
8.8	10.1	899	33
10.7	8.3	249	9
11.3	7.8	74	3
11.9	7.5	794	29
12.2	7.2	103	4
13.8	6.4	174	6
15.2	5.8	146	5
15.8	5.6	245	9
16.3	5.4	261	10
17.0	5.2	796	29
17.7	5.0	1594	59
18.2	4.9	316	12
18.7	4.7	2708	100
19.4	4.6	209	8
19.7	4.5	178	7
20.4	4.4	298	11
21.1	4.2	170	6
21.9	4.1	230	9
22.6	3.9	450	17
23.3	3.8	410	15
23.9	3.7	304	11
24.1	3.7	262	10
24.7	3.6	186	7
25.0	3.6	253	9
25.7	3.5	415	15
26.1	3.4	555	21
26.6	3.4	330	12
26.9	3.3	577	21

2 θ	d value Å	Intensity Cps	rel. I %
27.6	3.2	507	19
27.8	3.2	340	13
28.2	3.2	295	11
28.7	3.1	167	6
29.4	3.0	309	11
30.3	3.0	434	16
30.8	2.9	291	11
31.2	2.9	325	12
31.5	2.8	146	5
33.0	2.7	264	10
34.4	2.6	168	6

Crystalline Form F

[0382] Table 9 shows the peak list for crystalline form F. The uncertainty in the 2 θ values is $\pm 0.2^\circ$ in 2 θ ; rel. I is the relative intensity of the respective peaks. Maximum intensity is 100.

Table 9

2 θ	d value Å	Intensity Cps	rel. I %
8.0	11.1	53	4
9.0	9.8	828	69
10.6	8.4	84	7
14.5	6.1	144	12
15.4	5.7	304	25
16.1	5.5	591	49
17.9	4.9	658	55
18.2	4.9	443	37
18.7	4.8	1068	89
19.4	4.6	544	46
20.1	4.4	1196	100
20.6	4.3	333	28
21.8	4.1	294	25
21.9	4.1	244	20
22.9	3.9	166	14
23.5	3.8	157	13

2 θ	d value Å	Intensity Cps	rel. I %
24.0	3.7	111	9
24.6	3.6	411	34
25.0	3.6	188	16
25.6	3.5	328	27
27.1	3.3	315	26
27.4	3.3	357	30
27.9	3.2	190	16
29.3	3.1	328	27
30.0	3.0	163	14
30.3	3.0	249	21
30.8	2.9	128	11
31.4	2.9	140	12
31.6	2.8	122	10
32.2	2.8	122	10
32.8	2.7	106	9
33.3	2.7	100	8
34.4	2.6	82	7
35.0	2.6	99	8

Crystalline Form G

[0383] Table 10 shows the peak list for crystalline form G. The uncertainty in the 2 θ values is $\pm 0.2^\circ$ in 2 θ ; rel. I is the relative intensity of the respective peaks. Maximum intensity is 100.

Table 10

2 θ	d value Å	Intensity Cps	rel. I %
8.1	10.9	66	14
8.9	10.0	31	6
11.1	8.0	48	10
11.5	7.7	96	20
13.5	6.5	88	18
15.1	5.9	108	22
15.4	5.7	195	40
15.9	5.6	132	27
16.3	5.4	256	53

2 θ	d value Å	Intensity Cps	rel. I %
17.2	5.1	147	30
17.4	5.1	199	41
17.8	5.0	204	42
18.8	4.7	458	94
19.1	4.7	488	100
19.4	4.6	353	72
20.3	4.4	279	57
20.7	4.3	213	44
21.0	4.2	205	42
22.2	4.0	326	67
22.6	3.9	172	35
24.2	3.7	157	32
24.7	3.6	227	47
25.4	3.5	171	35
25.9	3.4	231	47
26.6	3.4	161	33
28.0	3.2	135	28
28.3	3.2	157	32
28.8	3.1	134	28
29.1	3.1	299	61
29.4	3.0	152	31
30.2	3.0	113	23
31.6	2.8	104	21
32.3	2.8	98	20
33.0	2.7	130	27
33.5	2.7	93	19
33.9	2.6	101	21
34.7	2.6	106	22

Crystalline Form H

[0384] Table 11 shows the peak list for crystalline form H. The uncertainty in the 2 θ values is $\pm 0.2^\circ$ in 2 θ ; rel. I is the relative intensity of the respective peaks. Maximum intensity is 100.

Table 11

2 θ	d value Å	Intensity Cps	rel. I %
7.4	12.0	145	10
8.5	10.4	52	3
9.0	9.9	76	5
9.8	9.1	50	3
10.8	8.2	154	10
11.4	7.7	687	45
12.2	7.2	74	5
12.8	6.9	104	7
13.1	6.8	130	9
14.8	6.0	102	7
15.7	5.6	153	10
16.2	5.5	147	10
16.6	5.3	134	9
17.1	5.2	340	22
18.0	4.9	352	23
18.3	4.8	569	38
19.2	4.6	1515	100
19.8	4.5	211	14
20.3	4.4	219	14
20.6	4.3	261	17
21.4	4.1	252	17
22.2	4.0	205	14
23.0	3.9	354	23
23.8	3.7	152	10
24.5	3.6	154	10
25.5	3.5	235	16
25.8	3.5	218	14
26.3	3.4	220	15
27.0	3.3	149	10
27.6	3.2	138	9
27.9	3.2	335	22
28.3	3.2	159	11
28.8	3.1	161	11
29.4	3.0	203	13

2 θ	d value Å	Intensity Cps	rel. I %
30.0	3.0	159	11
31.0	2.9	184	12
32.7	2.7	141	9
33.3	2.7	150	10

Crystalline Form I

[0385] Table 12 shows the peak list for crystalline form I. The uncertainty in the 2 θ values is $\pm 0.2^\circ$ in 2 θ ; rel. I is the relative intensity of the respective peaks. Maximum intensity is 100.

Table 12

2 θ	d value Å	Intensity Cps	rel. I %
9.5	9.3	48	3
10.1	8.7	73	4
10.9	8.1	1092	58
13.9	6.4	197	10
14.6	6.1	498	26
15.5	5.7	1188	63
16.5	5.4	436	23
17.1	5.2	1898	100
18.1	4.9	379	20
18.5	4.8	1207	64
18.8	4.7	744	39
19.4	4.6	229	12
19.9	4.5	273	14
20.2	4.4	268	14
21.1	4.2	503	27
21.9	4.1	501	26
22.3	4.0	349	18
22.8	3.9	267	14
23.0	3.9	240	13
23.6	3.8	735	39
24.0	3.7	391	21
25.2	3.5	227	12
25.9	3.4	486	26

2 θ	d value Å	Intensity Cps	rel. I %
26.6	3.4	229	12
27.6	3.2	144	8
28.0	3.2	474	25
28.8	3.1	442	23
29.1	3.1	154	8
30.0	3.0	223	12
30.6	2.9	443	23
31.9	2.8	408	22
33.1	2.7	192	10
34.2	2.6	150	8
34.8	2.6	247	13

Crystalline Form L

[0386] Table 13 shows the peak list for crystalline form L. The uncertainty in the 2 θ values is $\pm 0.2^\circ$ in 2 θ ; rel. I is the relative intensity of the respective peaks. Maximum intensity is 100.

Table 13

2 θ	d value Å	Intensity Cps	rel. I %
8.6	10.3	319	27
9.1	9.7	109	9
9.5	9.3	64	6
10.3	8.6	422	36
12.2	7.3	66	6
14.0	6.3	192	17
16.7	5.3	428	37
17.2	5.2	474	41
18.2	4.9	548	47
18.8	4.7	1165	100
20.7	4.3	256	22
21.2	4.2	316	27
22.3	4.0	154	13
23.0	3.9	187	16
24.5	3.6	140	12
26.0	3.4	286	25

2 θ	d value Å	Intensity Cps	rel. I %
27.4	3.3	334	29
28.9	3.1	269	23
30.2	3.0	235	20

Analysis - FT Raman Spectroscopy

[0387] FT Raman spectra were recorded on a Bruker RFS100 Raman spectrometer (Nd-YAG 100 mW laser, excitation 1064 nm, Ge detector, 64 scans, 25-3500 cm^{-1} , resolution 2 cm^{-1}).

[0388] Figure 2 shows the superposed Raman spectra of crystalline forms A, B, C, D, E, F, G, H, I and L.

[0389] Raman peak tables were generated using the software OPUS, version 3.1, build: 3, 0, 17 (20010216). The sensitivity of the peak picking function was chosen in a way that most of the peaks were found (typically between 0.5% to 3%). Features which were accidentally attributed to peaks and which were obviously noise, were removed by hand. Peaks are listed in a spectral region between 3200 cm^{-1} and 150 cm^{-1} . For the intensity classification, the absolute intensity was used and the most intense peak was scaled to 100%. The classification is as follow: very strong (vs): $I > 80\%$; strong (s): $80\% \geq I > 60\%$; medium (m): $60\% \geq I > 40\%$; weak (w): $40\% \geq I > 20\%$; and very weak (vw): $20\% \geq I$.

Crystalline form A

[0390] 3066 (w); 3057 (m); 2960 (w); 2947 (w); 2921 (w); 2871 (vw); 2851 (vw); 2793 (vw); 1628 (vw); 1583 (m); 1569 (vs); 1475 (w); 1453 (w); 1420 (vw); 1405 (vw); 1370 (vw); 1337 (vw); 1308 (m); 1289 (w); 1265 (vw); 1235 (vw); 1200 (vw); 1188 (vw); 1156 (vw); 1128 (vw); 1111 (vw); 1067 (vw); 1049 (vw); 1031 (w); 1002 (s); 981 (w); 943 (vw); 921 (m); 911 (w); 872 (vw); 828 (w); 785 (vw); 714 (vw); 694 (vw); 679 (w); 636 (vw); 620 (vw); 609 (vw); 544 (vw); 519 (vw); 488 (w); 420 (vw); 365 (vw); 254 (w); 202 (w); 184 (w); 170 (w); 152 (w).

Crystalline form B

[0391] 3072 (m); 3064 (m); 2984 (w); 2965 (w); 2950 (w); 2911 (w); 2879 (vw); 2846 (vw); 2794 (vw); 1630 (vw); 1581 (s); 1571 (vs); 1476 (m); 1460 (w); 1433 (w); 1374 (w); 1356 (vw); 1341 (vw); 1299 (m); 1268 (w); 1233 (vw); 1204 (w); 1191 (vw); 1174 (vw); 1163 (vw); 1141 (vw); 1119 (vw); 1110 (vw); 1074 (vw); 1053 (vw); 1039 (vw); 1028 (w); 1003 (s); 983 (vw); 947

(vw); 923 (m); 865 (vw); 825 (w); 785 (vw); 764 (vw); 716 (vw); 683 (w); 633 (vw); 621 (vw); 607 (vw); 560 (vw); 545 (vw); 518 (vw); 492 (w); 437 (vw); 395 (vw); 370 (w); 318 (vw); 301 (vw); 259 (w); 217 (w); 173 (m); 154 (m).

Crystalline form C

[0392] 3070 (m); 2990 (w); 2977 (w); 2951 (m); 2932 (m); 2890 (w); 2856 (w); 2845 (w); 2794 (vw); 1630 (vw); 1587 (s); 1570 (vs); 1478 (m); 1462 (w); 1435 (w); 1373 (w); 1339 (vw); 1299 (m); 1265 (vw); 1231 (vw); 1203 (w); 1189 (vw); 1157 (vw); 1119 (vw); 1111 (vw); 1077 (vw); 1051 (vw); 1028 (w); 1003 (s); 983 (vw); 948 (vw); 922 (m); 913 (w); 885 (vw); 829 (w); 786 (vw); 712 (vw); 682 (w); 633 (vw); 621 (vw); 608 (vw); 560 (vw); 542 (vw); 520 (vw); 491 (w); 432 (vw); 394 (vw); 371 (vw); 253 (w); 210 (w); 183 (m); 171 (m); 156 (m).

Crystalline form D

[0393] 3067 (s); 2957 (s); 2935 (m); 1570 (vs); 1479 (m); 1437 (m); 1377 (w); 1302 (m); 1264 (w); 1202 (w); 1158 (w); 1117 (w); 1028 (w); 1002 (s); 922 (s); 821 (w); 786 (w); 683 (m); 633 (w); 491 (m); 367 (m); 254 (m); 169 (s).

Crystalline form E

[0394] 3070 (m); 3057 (w); 2994 (w); 2961 (m); 2943 (m); 2894 (w); 2860 (w); 2836 (w); 2791 (vw); 1629 (vw); 1585 (vs); 1570 (vs); 1476 (w); 1461 (m); 1434 (m); 1376 (vw); 1354 (vw); 1337 (vw); 1297 (s); 1262 (w); 1230 (vw); 1202 (w); 1192 (w); 1169 (w); 1135 (vw); 1117 (vw); 1075 (vw); 1049 (vw); 1037 (w); 1027 (w); 1003 (s); 982 (w); 945 (vw); 923 (m); 913 (w); 870 (vw); 823 (w); 784 (vw); 760 (vw); 710 (vw); 680 (m); 632 (w); 621 (vw); 604 (vw); 557 (vw); 541 (vw); 518 (vw); 489 (w); 434 (vw); 423 (vw); 394 (vw); 370 (w); 315 (vw); 271 (vw); 257 (w); 188 (m); 174 (w); 159 (m).

Crystalline form F

[0395] 3070 (s); 3058 (m); 2992 (w); 2977 (m); 2952 (s); 2932 (m); 2889 (m); 2860 (w); 2843 (w); 2795 (vw); 2748 (vw); 2566 (vw); 1630 (vw); 1581 (vs); 1570 (vs); 1498 (vw); 1477 (m); 1463 (m); 1453 (w); 1438 (w); 1373 (w); 1353 (vw); 1338 (vw); 1299 (s); 1263 (vw); 1229 (vw); 1202 (w); 1189 (w); 1172 (vw); 1159 (w); 1146 (vw); 1119 (vw); 1110 (w); 1076 (vw); 1056 (w); 1050 (w); 1036 (w); 1027 (w); 1002 (s); 982 (w); 970 (vw); 946 (vw); 921 (s); 912 (w); 894 (vw); 870 (vw); 848 (vw); 828 (w); 786 (vw); 762 (vw); 712 (vw); 682 (m); 632 (w); 620 (w); 607 (vw); 602 (vw); 560 (vw); 542 (vw); 519 (vw); 491 (w); 471 (vw); 424 (vw); 394 (vw); 370 (w); 291 (vw); 253 (w); 212 (w); 183 (m); 171 (s); 157 (m).

Crystalline form G

[0396] 3156 (vw); 3069 (m); 3058 (w); 2999 (w); 2960 (w); 2945 (w); 2917 (vs); 2862 (vw); 2841 (vw); 2815 (vw); 2777 (vw); 1629 (vw); 1597 (w); 1569 (vs); 1475 (w); 1437 (w); 1418 (w); 1373 (vw); 1338 (vw); 1309 (w); 1292 (w); 1260 (vw); 1234 (vw); 1201 (vw); 1190 (vw); 1171 (vw); 1161 (vw); 1136 (vw); 1122 (vw); 1108 (vw); 1073 (vw); 1047 (w); 1038 (vw); 1029 (w); 1002 (m); 982 (vw); 948 (vw); 921 (m); 868 (vw); 830 (vw); 786 (vw); 769 (vw); 706 (w); 675 (s); 638 (vw); 621 (vw); 608 (vw); 546 (vw); 522 (vw); 491 (vw); 438 (vw); 424 (vw); 387 (vw); 365 (vw); 334 (vw); 306 (w); 254 (w); 202 (w); 180 (w); 169 (m).

Crystalline form H

[0397] 3069 (w); 2989 (m); 2945 (w); 2921 (w); 2788 (vw); 1629 (w); 1568 (vs); 1464 (m); 1375 (m); 1305 (s); 1263 (w); 1219 (w); 1199 (w); 1117 (w); 1073 (w); 1048 (w); 1030 (m); 1002 (s); 982 (w); 918 (s); 889 (w); 828 (m); 787 (w); 734 (w); 713 (w); 685 (m); 620 (w); 599 (m); 557 (w); 517 (w); 490 (m); 391 (m); 369 (m); 258 (s); 203 (s); 171 (vs).

Crystalline form I

[0398] 3085 (vw); 3066 (m); 3051 (vw); 3011 (vw); 2998 (vw); 2984 (w); 2964 (w); 2953 (vw); 2925 (m); 2911 (w); 2875 (vw); 2845 (vw); 2820 (vw); 2787 (vw); 2775 (vw); 2699 (vw); 1630 (vw); 1596 (w); 1589 (w); 1572 (vs); 1474 (w); 1459 (w); 1434 (w); 1417 (vw); 1404 (vw); 1375 (vw); 1354 (vw); 1339 (vw); 1305 (m); 1291 (vw); 1265 (vw); 1232 (vw); 1199 (vw); 1188 (vw); 1159 (vw); 1141 (vw); 1123 (vw); 1113 (vw); 1091 (vw); 1066 (vw); 1047 (vw); 1031 (w); 1001 (m); 983 (vw); 948 (vw); 924 (m); 912 (vw); 895 (vw); 825 (vw); 810 (vw); 784 (vw); 767 (vw); 715 (vw); 697 (vw); 680 (w); 636 (vw); 621 (vw); 605 (vw); 558 (vw); 544 (vw); 519 (vw); 489 (vw); 466 (vw); 438 (vw); 421 (vw); 392 (vw); 367 (vw); 315 (vw); 280 (vw); 256 (w); 202 (vw); 172 (w); 155 (w).

Crystalline form L

[0399] 3069 (s); 3042 (w); 2987 (m); 2922 (m); 2897 (w); 2859 (w); 2843 (w); 2828 (vw); 2787 (vw); 2364 (vw); 1631 (w); 1590 (vs); 1577 (vs); 1475 (m); 1438 (w); 1418 (vw); 1376 (w); 1353 (vw); 1338 (vw); 1307 (m); 1293 (w); 1267 (vw); 1235 (vw); 1201 (w); 1186 (vw); 1157 (vw); 1129 (vw); 1077 (vw); 1049 (vw); 1029 (w); 1001 (s); 985 (w); 950 (vw); 924 (m); 913 (w); 872 (vw); 828 (w); 784 (vw); 763 (vw); 714 (vw); 693 (w); 679 (m); 637 (vw); 620 (w); 610 (vw); 544 (vw); 522 (vw); 488 (w); 423 (vw); 390 (vw); 362 (vw); 249 (w); 199 (w); 172 (m); 150 (w).

Analysis - DSC

[0400] Differential Scanning Calorimetry (DSC): device reference Perkin Elmer DSC 7 or Perkin Elmer Pyris 1. Unless otherwise specified, the samples were weighed in a sealed gold crucible. The measurement took place in a nitrogen flow in a temperature range from -50°C up to 350°C with a heating rate of 10°C./min. The temperatures specified in relation to DSC analyses are, unless otherwise specified, the temperatures of the peak maxima.

[0401] In the following tables, "ΔH" means "specific heat", and "peak" means that a thermal event was observed at the temperature with the given peak temperature.

Table 14

	DSC
Crystalline Form A	peak, 305°C, ΔH = 137 J/g
Crystalline Form B	peak, 113°C, ΔH = 49 J/g
	peak, 189°C, ΔH = 102 J/g
	peak, 208°C, ΔH = -13 J/g
Crystalline Form C	event, 115°C, 51 mW
	peak, 135°C, ΔH = 80 J/g
Crystalline Form D	peak, 112°C, ΔH = 27 J/g
	peak, 123°C, ΔH = 46 J/g
	peak, 214°C, ΔH = 23 J/g

Analysis - TG-FTIR

[0402] Thermogravimetry analytical experiments coupled with Fourier transform infrared (TG-FTIR) spectra were recorded with a Netzsch Thermo-Microwaage TG 209 and a Bruker FT-IR spectrometer Vector 22 (aluminium crucible (open or with micro-aperture), nitrogen atmosphere, heating rate 10°C/min, 25 up to 350°C).

[0403] TG-FTIR analysis showed that crystalline form A does not contain any enclosed solvent and is thus an anhydrate form.

[0404] TG-FTIR analyses performed with samples of crystalline form B revealed that these samples contained 8-9% of water agreeing with a dihydrate.

[0405] TG-FTIR analyses performed with samples of crystalline form C revealed that these

samples contained 6-22 % of EtOH. Either the different drying times after filtration caused variation in the EtOH content or different but isomorphous solvates were obtained (EtOH content: ~ 6 % - hemi-solvate, ~ 12 % - mono-solvate).

[0406] TG-FTIR analyses performed with samples of crystalline form D revealed that these samples contained 12-13% of 2PrOH agreeing with a mono-solvate.

[0407] TG-FTIR analysis performed with a sample of crystalline form E revealed that this sample contained 7.2% of MeOH agreeing with a hemi-solvate.

[0408] TG-FTIR analysis performed with a sample of crystalline form F revealed that this sample contained 13% of 1PrOH agreeing with a hemi-solvate.

[0409] TG-FTIR analysis performed with a sample of crystalline form G revealed that this sample contained 40% of DMSO. This DMSO content is very high, probably indicating enclosed and adsorbed solvent.

[0410] TG-FTIR analysis performed with a sample of crystalline form H revealed that this sample contained 2.8% of water agreeing with the hemi-hydrate.

[0411] TG-FTIR analysis performed with a sample of crystalline form I revealed that this sample contained 0.5% of water. This could be a second anhydrate form.

Analysis - DVS

[0412] Crystalline forms A, B, C and D were each characterized by dynamic vapor sorption (DVS) using a Projekt Messtechnik SPS 11-100n multi sample water vapor sorption analyzer. For the DVS analysis, each sample was allowed to equilibrate at 50% r.h. (relative humidity) before starting a pre-defined humidity program during which the change in weight of the sample is determined. All measurements were performed according to the following program: 2 h at 50 % r.h.; 50 % r. h. → 0 % r.h. (10 %/h); 5 h at 0% r.h.; 0 → 95 % r.h. (5 %/h); 3 h at 95 % r.h.; 95 → 50 % (10 %/h), and 2 h at 50 % r.h.

[0413] Although hygroscopicity was measured in a slightly different manner, it was classified according to the European Pharmacopoeia as follows: very hygroscopic (vh): increase of the mass ≥ 15 %; hygroscopic (h): increase of the mass is less than 15 % and equal or greater than 2 %; slightly hygroscopic (sh): increase of the mass is less than 2 % and equal or greater than 0.2 %; not hygroscopic (nh): increase of the mass is less than 0.2 %; deliquescent (d): sufficient water is absorbed to form a liquid.

Crystalline form A

[0414] DVS with two cycles was performed on a sample of crystalline form A. The first cycle was not symmetric, the sample contained still water when the DVS cycle returned to 50 % r.h. (relative humidity in %). The second cycle was reversible. Below 40 % r.h. the relative mass returned to -100 % (water content = 0 %). The hysteresis between 40 % and 70 % r.h. indicates a metastable zone. The second cycle indicated the following transformations: hemi-hydrate → anhydrate (< 38% r. h.) → hemi-hydrate (> 70% r. h.). The sample was classified to be hygroscopic ($\Delta m = 3-4\%$ at 85% r. h.; Δm : change in mass).

Crystalline form B

[0415] DVS with two cycles was performed on a sample of crystalline form B. The DVS showed two reversible cycles with mass changes of 8-9% at 20 % r. h. and 80 % r. h; i. e. indicating the following transformations: dihydrate → desolvated form (< 20% r. h.; -8% mass change) → dihydrate (> 80% r. h.; +8% mass change). The sample was classified to be slightly hygroscopic ($\Delta m = 0.5\%$ at 85% r. h.).

Crystalline form C

[0416] DVS with two cycles was performed on a sample of crystalline form C. The first cycle was not symmetric and indicated a transformation of the EtOH solvate (crystalline form C) into the hemihydrate. The second cycle was reversible and indicated the following transformations: hemi-hydrate → anhydrate (< 20% r. h.) → hemi-hydrate (> 65% r. h.). The sample was classified to be hygroscopic ($\Delta m = 2\%$ at 85% r. h.).

Crystalline form D

[0417] The DVS cycle of a sample of crystalline form D was found to be not reversible and at 80-85% r. h. a mass change of 3-4% was observed. The sample was classified to be hygroscopic ($\Delta m = 3-4\%$ at 85% r. h.). The water uptake - probably combined with 2PrOH/H₂O exchange starts at approximately 65% r. h.

Analysis - "crystalline form K" (mixture of crystalline form B and L)

[0418] 60 mg of "crystalline form K" (sample 1-16; mixture of modifications B and L) was stored for two weeks at RT over saturated Mg(NO₃)₂ (55% r. h.) for two weeks. According to FT Raman, a mixture of crystalline form B and "crystalline form K" was obtained.

[0419] 60 mg of "crystalline form K" (sample 1-16; mixture of modifications B and L) was

stored for two weeks at RT over saturated NH_4Cl (79% r. h.) for two weeks. According to FT Raman, a mixture of crystalline form B and "crystalline form K" was obtained.

[0420] 60 mg of "crystalline form K" (sample 1-16; mixture of modifications B and L) was stored for two weeks at RT over saturated K_2SO_4 (97% r. h.) for two weeks. According to FT Raman, crystalline form B was obtained.

[0421] A comparison of the three FT Raman spectra revealed that "crystalline form K" is converted into crystalline form B with increasing r. h.

[0422] TG-FTIR analyses of the three samples revealed only small shifts in the water content (7.8 % at 55 % r.h. to 9.0 % at 97 % r.h.).

Analysis - hygroscopicity of crystalline form A

[0423] The hygroscopicity of crystalline form A was further studied by storing samples of crystalline form A at different relative humidity values before analyzing them by TG-FTIR.

7-1) 54 mg of crystalline form A was stored for two weeks at RT over saturated $\text{Mg}(\text{NO}_3)_2$ (55% r. h.) for two weeks.

7-2) 52 mg of crystalline form A was stored for two weeks at RT over saturated NH_4Cl (79% r. h.) for two weeks.

7-3) 51 mg of crystalline form A was stored for two weeks at RT over saturated K_2SO_4 (97% r. h.) for two weeks.

[0424] TG-FTIR confirmed that the water content of the samples at 55 % (0.7%) and 79 % r.h. (1.1%) agrees with the DVS result. At 97 % r.h. a higher water content (16.5%) was observed, probably caused by condensation of water at the powder surface. FT Raman analysis revealed that the water content did not change the crystal form or that the conversion is not detectable by Raman.

Analysis - single crystal diffraction

[0425] Measurements were realized using $\text{MoK}\alpha$ -radiation ($\lambda = 0.71073 \text{ \AA}$) and a Bruker D8-Goniometer equipped with a APEX-CCD detector.

[0426] Crystal data of crystalline forms A, C, D, G and I are summarized in the following tables 15-39.

Crystalline form A**[0427]**

Table 15: Crystal data and structure refinement for crystalline form A.

Empirical formula	C ₂₄ H ₂₇ FN ₂ O
Formula weight	378.48
Temperature	130(2) K
Wavelength	71073 Å
Crystal system	Monoclinic
Space group	P 21/c
Unit cell dimensions	a = 14.7104(14) Å alpha = 90 deg.
	b = 13.8726(14) Å beta = 101.846(2) deg.
	c = 19.4704(19) Å gamma = 90 deg.
Volume	3888.7(7) Å ³
Z	8
Density (calculated)	1.293 Mg/m ³
Absorption coefficient	0.086 mm ⁻¹
F(000)	1616
Crystal size	0.44 x 0.12 x 0.07 mm
Theta range for data collection	2.43 to 30.52 deg.
Index ranges	-20 ≤ h ≤ 20, -19 ≤ k ≤ 19, -27 ≤ l ≤ 27
Reflections collected	58460
Independent reflections	11821 [R(int) = 0.0953]
Absorption correction	None
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	11821 / 0 / 517
Goodness-of-fit on F ²	1.069
Final R indices [I > 2σ(I)]	R1 = 0.0607, wR2 = 0.0880
R indices (all data)	R1 = 0.1235, wR2 = 0.0969
Largest diff. peak and hole	.584 and -.384 e. Å ⁻³

Table 16: Atomic coordinates (x 10⁴) (i.e. (x 10⁴)) and equivalent isotropic displacement parameters (Å² × 10³) (i.e. (Å² × 10³)) for crystalline form A. U(eq) is defined as one third of the trace of the orthogonalized U_{ij} tensor.

	x	y	z	U(eq)
F(1)	2425(1)	-1986(1)	2620(1)	35(1)
O(1)	-47(1)	2184(1)	4040(1)	37(1)
N(1)	1870(1)	474(1)	4675(1)	25(1)
N(2)	904(1)	2525(1)	6396(1)	27(1)
C(1)	2315(1)	-1375(1)	3152(1)	27(1)
C(2)	1596(1)	-735(1)	3022(1)	26(1)
C(3)	1494(1)	-125(1)	3579(1)	23(1)
C(4)	2125(1)	-199(1)	4228(1)	23(1)
C(5)	1096(1)	961(1)	4316(1)	23(1)
C(6)	584(1)	1735(1)	4618(1)	26(1)
C(7)	-566(1)	1554(1)	3554(1)	42(1)
C(8)	6(1)	964(1)	3152(1)	31(1)
C(9)	845(1)	608(1)	3654(1)	24(1)
C(10)	70(1)	1336(1)	5160(1)	27(1)
C(11)	-456(1)	2118(1)	5467(1)	27(1)
C(12)	180(1)	2934(1)	5822(1)	22(1)
C(13)	550(1)	1917(1)	6899(1)	39(1)
C(14)	1554(1)	3232(1)	6772(1)	36(1)
C(15)	664(1)	3332(1)	5253(1)	27(1)
C(16)	1207(1)	2561(1)	4949(1)	27(1)
C(17)	-394(1)	3723(1)	6088(1)	24(1)
C(18)	-125(1)	4686(1)	6125(1)	30(1)
C(19)	-633(1)	5387(1)	6387(1)	35(1)
C(20)	-1423(1)	5140(1)	6622(1)	36(1)
C(21)	-1701(1)	4191(1)	6594(1)	35(1)
C(22)	-1198(1)	3497(1)	6331(1)	30(1)
C(23)	2852(1)	-863(1)	4338(1)	25(1)
C(24)	2941(1)	-1461(1)	3785(1)	27(1)
F(2)	5985(1)	9246(1)	1447(1)	50(1)
O(2)	3180(1)	4515(1)	1030(1)	28(1)
N(3)	5608(1)	5355(1)	1314(1)	23(1)
N(4)	5231(1)	1974(1)	1207(1)	23(1)
C(25)	5909(1)	8256(1)	1415(1)	34(1)
C(26)	5054(1)	7855(1)	1352(1)	32(1)

	x	y	z	U(eq)
C(27)	5004(1)	6843(1)	1322(1)	25(1)
C(28)	5836(1)	6323(1)	1358(1)	24(1)
C(29)	4661(1)	5262(1)	1243(1)	22(1)
C(30)	4174(1)	4311(1)	1167(1)	22(1)
C(31)	2921(1)	5264(1)	1462(1)	33(1)
C(32)	3258(1)	6247(1)	1272(1)	32(1)
C(33)	4270(1)	6152(1)	1259(1)	24(1)
C(34)	4452(1)	3685(1)	1819(1)	24(1)
C(35)	3981(1)	2702(1)	1725(1)	24(1)
C(36)	4208(1)	2122(1)	1108(1)	22(1)
C(37)	5672(1)	1545(1)	1878(1)	30(1)
C(38)	5509(1)	1428(1)	637(1)	33(1)
C(39)	3890(1)	2755(1)	452(1)	23(1)
C(40)	4337(1)	3754(1)	524(1)	22(1)
C(41)	3678(1)	1156(1)	1042(1)	25(1)
C(42)	3643(1)	603(1)	1637(1)	31(1)
C(43)	3172(1)	-265(1)	1586(1)	38(1)
C(44)	2721(1)	-611(1)	944(1)	41(1)
C(45)	2750(1)	-87(1)	351(1)	40(1)
C(46)	3229(1)	780(1)	399(1)	32(1)
C(47)	6694(1)	6755(1)	1427(1)	29(1)
C(48)	6727(1)	7747(1)	1455(1)	32(1)

Table 17A: Bond lengths [Å] and angles [deg] for crystalline form A.

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
F(1)- C(1)	1.3733(18)	C(15)- H(15A)	.9900	C(32)- C(33)	1.500(2)
O(1)- C(7)	1.3954(19)	C(15)- H(15B)	.9900	C(32)- H(32A)	.9900
O(1)- C(6)	1.4443(18)	C(16)- H(16A)	.9900	C(32)- H(32B)	.9900
N(1)- C(4)	1.381(2)	C(16)- H(16B)	.9900	C(34)- C(35)	1.523(2)
N(1)- C(5)	1.383(2)	C(17)- C(18)	1.392(2)	C(34)- H(34A)	.9900
N(1)-	.866(15)	C(17)-	1.395(2)	C(34)-	.9900

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
H(1)		C(22)		H(34B)	
N(2)- C(14)	1.458(2)	C(18)- C(19)	1.386(2)	C(35)- C(36)	1.539(2)
N(2)- C(13)	1.466(2)	C(18)- H(18)	.9500	C(35)- H(35A)	.9900
N(2)- C(12)	1.4886(19)	C(19)- C(20)	1.378(2)	C(35)- H(35B)	.9900
C(1)- C(2)	1.364(2)	C(19)- H(19)	.9500	C(36)- C(39)	1.541(2)
C(1)- C(24)	1.384(2)	C(20)- C(21)	1.377(2)	C(36)- C(41)	1.543(2)
C(2)- C(3)	1.408(2)	C(20)- H(20)	.9500	C(37)- H(37A)	.9800
C(2)- H(2)	.9500	C(21)- C(22)	1.376(2)	C(37)- H(37B)	.9800
C(3)- C(4)	1.409(2)	C(21)- H(21)	.9500	C(37)- H(37C)	.9800
C(3)- C(9)	1.423(2)	C(22)- H(22)	.9500	C(38)- H(38A)	.9800
C(4)- C(23)	1.394(2)	C(23)- C(24)	1.387(2)	C(38)- H(38B)	.9800
C(5)- C(9)	1.357(2)	C(23)- H(23)	.9500	C(38)- H(38C)	.9800
C(5)- C(6)	1.500(2)	C(24)- H(24)	.9500	C(39)- C(40)	1.528(2)
C(6)- C(16)	1.524(2)	F(2)- C(25)	1.3780(18)	C(39)- H(39A)	.9900
C(6)- C(10)	1.525(2)	O(2)- C(31)	1.4359(18)	C(39)- H(39B)	.9900
C(7)- C(8)	1.504(2)	O(2)- C(30)	1.4587(17)	C(40)- H(40A)	.9900
C(7)- H(7A)	.9900	N(3)- C(29)	1.3769(19)	C(40)- H(40B)	.9900
C(7)- H(7B)	.9900	N(3)- C(28)	1.3823(19)	C(41)- C(46)	1.391(2)
C(8)- C(9)	1.492(2)	N(3)- H(3)	.850(15)	C(41)- C(42)	1.399(2)
C(8)- H(8A)	.9900	N(4)- C(37)	1.4622(19)	C(42)- C(43)	1.384(2)

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(8)- H(8B)	.9900	N(4)- C(38)	1.4689(19)	C(42)- H(42)	.9500
C(10)- C(11)	1.522(2)	N(4)- C(36)	1.4911(19)	C(43)- C(44)	1.376(2)
C(10)- H(10A)	.9900	C(25)- C(26)	1.358(2)	C(43)- H(43)	.9500
C(10)- H(10B)	.9900	C(25)- C(48)	1.383(2)	C(44)- C(45)	1.373(2)
C(11)- C(12)	1.539(2)	C(26)- C(27)	1.406(2)	C(44)- H(44)	.9500
C(11)- H(11A)	.9900	C(26)- H(26)	.9500	C(45)- C(46)	1.386(2)
C(11)- H(11 B)	.9900	C(27)- C(28)	1.410(2)	C(45)- H(45)	.9500
C(12)- C(17)	1.536(2)	C(27)- C(33)	1.430(2)	C(46)- H(46)	.9500
C(12)- C(15)	1.537(2)	C(28)- C(47)	1.379(2)	C(47)- C(48)	1.379(2)
C(13)- H(13A)	.9800	C(29)- C(33)	1.365(2)	C(47)- H(47)	.9500
C(13)- H(13B)	.9800	C(29)- C(30)	1.494(2)	C(48)- H(48)	.9500
C(13)- H(13C)	.9800	C(30)- C(34)	1.525(2)		
C(14)- H(14A)	.9800	C(30)- C(40)	1.532(2)		
C(14)- H(14B)	.9800	C(31)- C(32)	1.522(2)		
C(14)- H(14C)	.9800	C(31)- H(31A)	.9900		
C(15)- C(16)	1.525(2)	C(31)- H(31B)	.9900		

Table 17B: (Table 17A continued) Bond lengths [Å] and angles [deg] for crystalline form A.

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(7)-O(1)-C(6)	115.69(13)	C(10)-C(11)- H(11B)	109.0
C(4)-N(1)-C(5)	108.25(14)	C(12)-C(11)- H(11B)	109.0

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(4)-N(1)-H(1)	123.5(11)	H(11A)-C(11)- H(11B)	107.8
C(5)-N(1)-H(1)	128.0(11)	N(2)-C(12)- C(17)	112.08(13)
C(14)-N(2)- C(13)	109.77(14)	N(2)-C(12)- C(15)	108.20(13)
C(14)-N(2)- C(12)	114.51(13)	C(17)-C(12)- C(15)	110.81(13)
C(13)-N(2)- C(12)	115.04(13)	N(2)-C(12)- C(11)	109.41(13)
C(2)-C(1)-F(1)	117.98(15)	C(17)-C(12)- C(11)	110.51(13)
C(2)-C(1)-C(24)	124.74(16)	C(15)-C(12)- C(11)	105.61(13)
F(1)-C(1)-C(24)	117.28(15)	N(2)-C(13)- H(13A)	109.5
C(1)-C(2)-C(3)	116.76(15)	N(2)-C(13)- H(13B)	109.5
C(1)-C(2)-H(2)	121.6	H(13A)-C(13)- H(13B)	109.5
C(3)-C(2)-H(2)	121.6	N(2)-C(13)- H(13C)	109.5
C(2)-C(3)-C(4)	119.42(15)	H(13A)-C(13)- H(13C)	109.5
C(2)-C(3)-C(9)	133.52(16)	H(13B)-C(13)- H(13C)	109.5
C(4)-C(3)-C(9)	107.06(14)	N(2)-C(14)- H(14A)	109.5
N(1)-C(4)-C(23)	130.50(15)	N(2)-C(14)- H(14B)	109.5
N(1)-C(4)-C(3)	107.53(14)	H(14A)-C(14)- H(14B)	109.5
C(23)-C(4)-C(3)	121.98(15)	N(2)-C(14)- H(14C)	109.5
C(9)-C(5)-N(1)	109.91(15)	H(14A)-C(14)- H(14C)	109.5
C(9)-C(5)-C(6)	124.72(15)	H(14B)-C(14)- H(14C)	109.5
N(1)-C(5)-C(6)	125.33(14)	C(16)-C(15)- C(12)	112.57(13)

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
O(1)-C(6)-C(5)	107.32(13)	C(16)-C(15)- H(15A)	109.1
O(1)-C(6)-C(16)	103.90(13)	C(12)-C(15)- H(15A)	109.1
C(5)-C(6)-C(16)	113.47(14)	C(16)-C(15)- H(15B)	109.1
O(1)-C(6)-C(10)	111.41(13)	C(12)-C(15)- H(15B)	109.1
C(5)-C(6)-C(10)	111.84(13)	H(15A)-C(15)- H(15B)	107.8
C(16)-C(6)- C(10)	108.64(13)	C(6)-C(16)- C(15)	112.03(13)
O(1)-C(7)-C(8)	114.03(15)	C(6)-C(16)- H(16A)	109.2
O(1)-C(7)- H(7A)	108.7	C(15)-C(16)- H(16A)	109.2
C(8)-C(7)- H(7A)	108.7	C(6)-C(16)- H(16B)	109.2
O(1)-C(7)- H(7B)	108.7	C(15)-C(16)- H(16B)	109.2
C(8)-C(7)- H(7B)	108.7	H(16A)-C(16)- H(16B)	107.9
H(7A)-C(7)- H(7B)	107.6	C(18)-C(17)- C(22)	116.77(15)
C(9)-C(8)-C(7)	108.37(14)	C(18)-C(17)- C(12)	122.07(15)
C(9)-C(8)- H(8A)	110.0	C(22)-C(17)- C(12)	121.12(14)
C(7)-C(8)- H(8A)	110.0	C(19)-C(18)- C(17)	121.57(17)
C(9)-C(8)- H(8B)	110.0	C(19)-C(18)- H(18)	119.2
C(7)-C(8)- H(8B)	110.0	C(17)-C(18)- H(18)	119.2
H(8A)-C(8)- H(8B)	108.4	C(20)-C(19)- C(18)	120.22(17)
C(5)-C(9)-C(3)	107.25(14)	C(20)-C(19)- H(19)	119.9
C(5)-C(9)-C(8)	122.05(15)	C(18)-C(19)- H(19)	119.9

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(3)-C(9)-C(8)	130.67(15)	C(21)-C(20)- C(19)	119.23(17)
C(11)-C(10)- C(6)	112.14(13)	C(21)-C(20)- H(20)	120.4
C(11)-C(10)- H(10A)	109.2	C(19)-C(20)- H(20)	120.4
C(6)-C(10)- H(10A)	109.2	C(22)-C(21)- C(20)	120.41(18)
C(11)-C(10)- H(10B)	109.2	C(22)-C(21)- H(21)	119.8
C(6)-C(10)- H(10B)	109.2	C(20)-C(21)- H(21)	119.8
H(10A)-C(10)- H(10B)	107.9	C(21)-C(22)- C(17)	121.79(16)
C(10)-C(11)- C(12)	112.93(13)	C(21)-C(22)- H(22)	119.1
C(10)-C(11)- H(11A)	109.0	C(17)-C(22)- H(22)	119.1
C(12)-C(11)- H(11A)	109.0	C(24)-C(23)- C(4)	117.88(15)
C(24)-C(23)- H(23)	121.1	C(34)-C(35)- C(36)	113.26(13)
C(4)-C(23)- H(23)	121.1	C(34)-C(35)- H(35A)	108.9
C(1)-C(24)- C(23)	119.22(16)	C(36)-C(35)- H(35A)	108.9
C(1)-C(24)- H(24)	120.4	C(34)-C(35)- H(35B)	108.9
C(23)-C(24)- H(24)	120.4	C(36)-C(35)- H(35B)	108.9
C(31)-O(2)- C(30)	114.21(12)	H(35A)-C(35)- H(35B)	107.7
C(29)-N(3)- C(28)	108.97(14)	N(4)-C(36)- C(35)	110.26(12)
C(29)-N(3)-H(3)	125.5(11)	N(4)-C(36)- C(39)	108.21(12)
C(28)-N(3)-H(3)	125.4(11)	C(35)-C(36)- C(39)	105.72(13)
C(37)-N(4)- C(38)	108.87(13)	N(4)-C(36)- C(41)	111.75(12)

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(37)-N(4)- C(36)	114.99(12)	C(35)-C(36)- C(41)	109.41(13)
C(38)-N(4)- C(36)	113.84(12)	C(39)-C(36)- C(41)	111.32(13)
C(26)-C(25)- F(2)	118.52(17)	N(4)-C(37)- H(37A)	109.5
C(26)-C(25)- C(48)	125.10(16)	N(4)-C(37)- H(37B)	109.5
F(2)-C(25)- C(48)	116.38(17)	H(37A)-C(37)- H(37B)	109.5
C(25)-C(26)- C(27)	116.90(17)	N(4)-C(37)- H(37C)	109.5
C(25)-C(26)- H(26)	121.6	H(37A)-C(37)- H(37C)	109.5
C(27)-C(26)- H(26)	121.6	H(37B)-C(37)- H(37C)	109.5
C(26)-C(27)- C(28)	118.12(16)	N(4)-C(38)- H(38A)	109.5
C(26)-C(27)- C(33)	134.82(16)	N(4)-C(38)- H(38B)	109.5
C(28)-C(27)- C(33)	107.05(14)	H(38A)-C(38)- H(38B)	109.5
C(47)-C(28)- N(3)	129.33(16)	N(4)-C(38)- H(38C)	109.5
C(47)-C(28)- C(27)	123.41(16)	H(38A)-C(38)- H(38C)	109.5
N(3)-C(28)- C(27)	107.26(15)	H(38B)-C(38)- H(38C)	109.5
C(33)-C(29)- N(3)	109.57(14)	C(40)-C(39)- C(36)	112.93(13)
C(33)-C(29)- C(30)	127.23(15)	C(40)-C(39)- H(39A)	109.0
N(3)-C(29)- C(30)	123.19(14)	C(36)-C(39)- H(39A)	109.0
O(2)-C(30)- C(29)	106.83(12)	C(40)-C(39)- H(39B)	109.0
O(2)-C(30)- C(34)	110.44(13)	C(36)-C(39)- H(39B)	109.0
C(29)-C(30)- C(34)	112.28(13)	H(39A)-C(39)- H(39B)	107.8

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
O(2)-C(30)- C(40)	105.56(12)	C(39)-C(40)- C(30)	113.29(13)
C(29)-C(30)- C(40)	112.04(13)	C(39)-C(40)- H(40A)	108.9
C(34)-C(30)- C(40)	109.45(13)	C(30)-C(40)- H(40A)	108.9
O(2)-C(31)- C(32)	111.56(14)	C(39)-C(40)- H(40B)	108.9
O(2)-C(31)- H(31A)	109.3	C(30)-C(40)- H(40B)	108.9
C(32)-C(31)- H(31A)	109.3	H(40A)-C(40)- H(40B)	107.7
O(2)-C(31)- H(31 B)	109.3	C(46)-C(41)- C(42)	116.62(16)
C(32)-C(31)- H(31 B)	109.3	C(46)-C(41)- C(36)	122.66(15)
H(31A)-C(31)- H(31 B)	108.0	C(42)-C(41)- C(36)	120.71(15)
C(33)-C(32)- C(31)	107.39(13)	C(43)-C(42)- C(41)	121.42(17)
C(33)-C(32)- H(32A)	110.2	C(43)-C(42)- H(42)	119.3
C(31)-C(32)- H(32A)	110.2	C(41)-C(42)- H(42)	119.3
C(33)-C(32)- H(32B)	110.2	C(44)-C(43)- C(42)	120.61(18)
C(31)-C(32)- H(32B)	110.2	C(44)-C(43)- H(43)	119.7
H(32A)-C(32)- H(32B)	108.5	C(42)-C(43)- H(43)	119.7
C(29)-C(33)- C(27)	107.11(14)	C(45)-C(44)- C(43)	119.16(18)
C(29)-C(33)- C(32)	120.28(15)	C(45)-C(44)- H(44)	120.4
C(27)-C(33)- C(32)	132.37(15)	C(43)-C(44)- H(44)	120.4
C(35)-C(34)- C(30)	111.89(13)	C(44)-C(45)- C(46)	120.31(18)
C(35)-C(34)- H(34A)	109.2	C(44)-C(45)- H(45)	119.8

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(30)-C(34)- H(34A)	109.2	C(46)-C(45)- H(45)	119.8
C(35)-C(34)- H(34B)	109.2	C(45)-C(46)- C(41)	121.86(17)
C(30)-C(34)- H(34B)	109.2	C(45)-C(46)- H(46)	119.1
H(34A)-C(34)- H(34B)	107.9	C(41)-C(46)- H(46)	119.1
C(48)-C(47)- C(28)	117.49(17)		
C(48)-C(47)- H(47)	121.3		
C(28)-C(47)- H(47)	121.3		
C(47)-C(48)- C(25)	118.99(17)		
C(47)-C(48)- H(48)	120.5		
C(25)-C(48)- H(48)	120.5		

Table 18: Hydrogen coordinates ($\times 10^4$) (i.e. ($\times 10^4$)) and isotropic displacement parameters ($\times 10^3$) (i.e. ($\times 10^3$)) for crystalline form A.

	x	y	z	U(eq)
H(1)	2191(11)	592(11)	5092(8)	31(5)
H(2)	1185	-702	2578	32
H(7A)	-1024	1933	3216	50
H(7B)	-916	1113	3803	50
H(8A)	-363	412	2922	37
H(8B)	195	1364	2783	37
H(10A)	-374	836	4938	33
H(10B)	522	1028	5544	33
H(11A)	-775	1824	5815	32
H(11B)	-936	2395	5087	32
H(13A)	1067	1714	7273	59
H(13B)	245	1347	6657	59
H(13C)	100	2284	7103	59
H(14A)	1224	3673	7029	54

	x	y	z	U(eq)
H(14B)	1829	3600	6436	54
H(14C)	2046	2900	7103	54
H(15A)	191	3614	4870	32
H(15B)	1094	3855	5457	32
H(16A)	1508	2856	4590	32
H(16B)	1701	2302	5327	32
H(18)	419	4868	5966	36
H(19)	-435	6040	6404	42
H(20)	-1773	5620	6802	43
H(21)	-2244	4014	6757	42
H(22)	-1402	2846	6314	36
H(23)	3273	-905	4778	30
H(24)	3426	-1925	3840	32
H(3)	5986(11)	4895(11)	1304(8)	24(5)
H(26)	4515	8240	1328	38
H(31A)	3190	5123	1961	39
H(31B)	2237	5276	1403	39
H(32A)	2905	6453	806	38
H(32B)	3168	6733	1624	38
H(34A)	5135	3595	1923	29
H(34B)	4282	4018	2224	29
H(35A)	3300	2794	1649	29
H(35B)	4177	2325	2163	29
H(37A)	5460	877	1898	46
H(37B)	5504	1915	2262	46
H(37C)	6348	1553	1924	46
H(38A)	6188	1379	727	49
H(38B)	5287	1761	190	49
H(38C)	5239	780	614	49
H(39A)	4045	2425	41	28
H(39B)	3206	2829	366	28
H(40A)	4084	4132	96	26
H(40B)	5014	3682	555	26
H(42)	3950	829	2085	37
H(43)	3159	-626	1999	46

	x	y	z	U(eq)
H(44)	2393	-1205	911	49
H(45)	2440	-318	-95	48
H(46)	3251	1126	-18	38
H(47)	7243	6382	1455	34
H(48)	7302	8077	1501	39

Table 19: Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) (i.e. $\text{\AA}^2 \times 10^3$) for crystalline form A. The anisotropic displacement factor exponent takes the form: $-2 \pi^2 [h^2 a^{*2} U_{11} + \dots + 2 h k a^* b^* U_{12}]$.

	U11	U22	U33	U23	U13	U12
F(1)	42(1)	31(1)	35(1)	-9(1)	13(1)	1(1)
O(1)	38(1)	32(1)	34(1)	-4(1)	-5(1)	11(1)
N(1)	24(1)	27(1)	21(1)	-1(1)	1(1)	-1(1)
N(2)	25(1)	29(1)	27(1)	4(1)	6(1)	0(1)
C(1)	30(1)	23(1)	30(1)	-6(1)	11(1)	-6(1)
C(2)	30(1)	26(1)	23(1)	0(1)	4(1)	-6(1)
C(3)	25(1)	22(1)	24(1)	0(1)	6(1)	-6(1)
C(4)	23(1)	21(1)	25(1)	-2(1)	6(1)	-6(1)
C(5)	22(1)	21(1)	26(1)	1(1)	4(1)	-1(1)
C(6)	26(1)	26(1)	24(1)	1(1)	2(1)	3(1)
C(7)	32(1)	44(1)	42(1)	-6(1)	-9(1)	7(1)
C(8)	32(1)	30(1)	27(1)	-1(1)	-2(1)	3(1)
C(9)	26(1)	22(1)	24(1)	2(1)	5(1)	0(1)
C(10)	27(1)	21(1)	35(1)	-3(1)	7(1)	-2(1)
C(11)	24(1)	22(1)	36(1)	-1(1)	9(1)	-1(1)
C(12)	23(1)	18(1)	26(1)	2(1)	6(1)	-1(1)
C(13)	48(1)	35(1)	36(1)	12(1)	12(1)	6(1)
C(14)	24(1)	48(1)	36(1)	-5(1)	3(1)	0(1)
C(15)	34(1)	21(1)	28(1)	-1(1)	11(1)	-4(1)
C(16)	31(1)	24(1)	29(1)	1(1)	12(1)	-3(1)
C(17)	26(1)	23(1)	21(1)	2(1)	4(1)	2(1)
C(18)	39(1)	24(1)	29(1)	3(1)	11(1)	0(1)
C(19)	50(1)	23(1)	29(1)	3(1)	4(1)	7(1)
C(20)	36(1)	40(1)	28(1)	-5(1)	-1(1)	17(1)
C(21)	24(1)	45(1)	36(1)	-9(1)	5(1)	0(1)
C(22)	27(1)	26(1)	37(1)	-5(1)	6(1)	-1(1)

	U11	U22	U33	U23	U13	U12
C(23)	21(1)	24(1)	28(1)	2(1)	3(1)	-4(1)
C(24)	23(1)	22(1)	36(1)	0(1)	8(1)	-3(1)
F(2)	81(1)	19(1)	51(1)	-2(1)	19(1)	-2(1)
O(2)	20(1)	32(1)	29(1)	-8(1)	2(1)	4(1)
N(3)	23(1)	17(1)	29(1)	-1(1)	5(1)	5(1)
N(4)	20(1)	27(1)	22(1)	1(1)	4(1)	2(1)
C(25)	61(1)	14(1)	26(1)	-2(1)	7(1)	-1(1)
C(26)	46(1)	24(1)	24(1)	-1(1)	5(1)	10(1)
C(27)	38(1)	22(1)	16(1)	-1(1)	4(1)	5(1)
C(28)	32(1)	18(1)	19(1)	-2(1)	4(1)	1(1)
C(29)	22(1)	24(1)	18(1)	-2(1)	1(1)	4(1)
C(30)	17(1)	26(1)	22(1)	-2(1)	1(1)	3(1)
C(31)	24(1)	40(1)	36(1)	-10(1)	10(1)	8(1)
C(32)	32(1)	32(1)	30(1)	-7(1)	4(1)	10(1)
C(33)	27(1)	24(1)	19(1)	-4(1)	1(1)	7(1)
C(34)	25(1)	26(1)	20(1)	-1(1)	3(1)	2(1)
C(35)	25(1)	25(1)	24(1)	-1(1)	6(1)	0(1)
C(36)	21(1)	22(1)	22(1)	-1(1)	3(1)	1(1)
C(37)	27(1)	34(1)	29(1)	4(1)	4(1)	1(1)
C(38)	29(1)	39(1)	31(1)	0(1)	9(1)	6(1)
C(39)	23(1)	22(1)	22(1)	-2(1)	2(1)	0(1)
C(40)	23(1)	22(1)	20(1)	1(1)	3(1)	2(1)
C(41)	20(1)	24(1)	31(1)	-2(1)	8(1)	2(1)
C(42)	30(1)	28(1)	35(1)	1(1)	9(1)	0(1)
C(43)	40(1)	28(1)	50(1)	3(1)	18(1)	1(1)
C(44)	34(1)	24(1)	67(2)	-1(1)	15(1)	-3(1)
C(45)	36(1)	31(1)	50(1)	-8(1)	2(1)	-1(1)
C(46)	31(1)	26(1)	37(1)	-2(1)	4(1)	1(1)
C(47)	35(1)	26(1)	25(1)	-2(1)	6(1)	0(1)
C(48)	44(1)	26(1)	27(1)	-3(1)	9(1)	-10(1)

Crystalline form C

[0428]

Table 20: Crystal data and structure refinement for crystalline form C.

Empirical formula	C ₂₆ H ₃₃ FN ₂ O ₂
Formula weight	424.54
Temperature	130(2) K
Wavelength	.71073 Å
Crystal system	Triclinic
Space group	P-1
Unit cell dimensions	a = 9.6987(14) Å alpha = 110.528(3) deg.
	b = 11.1173(16) Å beta = 98.919(3) deg.
	c = 12.5467(18) Å gamma = 113.367(3) deg.
Volume	1092.8(3) Å ³
Z	2
Density (calculated)	1.290 Mg/m ³
Absorption coefficient	0.087 mm ⁻¹
F(000)	456
Crystal size	0.36 x 0.13 x 0.08 mm
Theta range for data collection	2.16 to 28.49 deg.
Index ranges	-13 ≤ h ≤ 12, -14 ≤ k ≤ 14, -16 ≤ l ≤ 16
Reflections collected	14533
Independent reflections	5293 [R(int) = 0.0634]
Absorption correction	None
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	5293/0/290
Goodness-of-fit on F ²	1.005
Final R indices [I > 2σ(I)]	R1 = 0.0491, wR2 = 0.0932
R indices (all data)	R1 = 0.0681, wR2 = 0.0981
Largest diff. peak and hole	.269 and -.265 e. Å ⁻³

Table 21: Atomic coordinates (x 10⁴) (i.e. (x 10⁴)) and equivalent isotropic displacement parameters (Å² x 10³) (i.e. (Å² x 10³)) for crystalline form C. U(eq) is defined as one third of the trace of the orthogonalized U_{ij} tensor.

	x	y	z	U(eq)
F(1)	-511(1)	2563(1)	-6080(1)	28(1)
O(1)	3958(1)	7981(1)	719(1)	21(1)
N(1)	1717(1)	3970(1)	-1311(1)	18(1)

	x	y	z	U(eq)
N(2)	3395(1)	5029(1)	2670(1)	18(1)
C(1)	50(2)	2878(2)	-4885(1)	20(1)
C(2)	914(2)	4324(2)	-4035(1)	19(1)
C(3)	1442(2)	4604(2)	-2822(1)	16(1)
C(4)	1083(2)	3407(2)	-2550(1)	16(1)
C(5)	2436(2)	5478(2)	-810(1)	17(1)
C(6)	3232(2)	6469(2)	529(1)	17(1)
C(7)	3053(2)	8287(2)	-59(1)	23(1)
C(8)	2910(2)	7498(2)	-1375(1)	22(1)
C(9)	2302(2)	5914(2)	-1689(1)	17(1)
C(10)	4624(2)	6281(2)	1050(1)	18(1)
C(11)	5346(2)	7112(2)	2426(1)	18(1)
C(12)	4122(2)	6661(2)	3051(1)	16(1)
C(13)	2133(2)	4490(2)	3165(1)	26(1)
C(14)	4568(2)	4569(2)	2982(1)	25(1)
C(15)	2817(2)	6989(2)	2581(1)	18(1)
C(16)	2036(2)	6197(2)	1204(1)	18(1)
C(17)	4931(2)	7559(2)	4435(1)	18(1)
C(18)	6456(2)	7836(2)	4998(1)	21(1)
C(19)	7171(2)	8583(2)	6246(1)	24(1)
C(20)	6396(2)	9102(2)	6965(1)	26(1)
C(21)	4893(2)	8853(2)	6431(1)	27(1)
C(22)	4165(2)	8091(2)	5184(1)	23(1)
C(23)	201(2)	1954(2)	-3443(1)	19(1)
C(24)	-328(2)	1693(2)	-4629(1)	20(1)
O(2)	1709(1)	2687(1)	288(1)	26(1)
C(25)	2152(2)	1585(2)	142(1)	30(1)
C(26)	1434(2)	485(2)	-1175(1)	44(1)

Table 22A: Bond lengths [Å] and angles [deg] for crystalline form C.

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
F(1)-C(1)	1.3758(15)	C(13)-H(13A)	.9800
O(1)-C(7)	1.4291(16)	C(13)-H(13B)	.9800
O(1)-C(6)	1.4512(16)	C(13)-	.9800

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
		H(13C)	
N(1)-C(5)	1.3747(17)	C(14)- H(14A)	.9800
N(1)-C(4)	1.3787(17)	C(14)- H(14B)	.9800
N(1)-H(1)	.885(15)	C(14)- H(14C)	.9800
N(2)-C(13)	1.4700(17)	C(15)-C(16)	1.5270(18)
N(2)-C(14)	1.4737(18)	C(15)- H(15A)	.9900
N(2)-C(12)	1.5132(17)	C(15)- H(15B)	.9900
C(1)-C(2)	1.3694(19)	C(16)- H(16A)	.9900
C(1)-C(24)	1.387(2)	C(16)- H(16B)	.9900
C(2)-C(3)	1.4051(18)	C(17)-C(18)	1.394(2)
C(2)-H(2)	.9500	C(17)-C(22)	1.4012(19)
C(3)-C(4)	1.4141(19)	C(18)-C(19)	1.3848(19)
C(3)-C(9)	1.4332(19)	C(18)-H(18)	.9500
C(4)-C(23)	1.3908(19)	C(19)-C(20)	1.378(2)
C(5)-C(9)	1.3609(19)	C(19)-H(19)	.9500
C(5)-C(6)	1.5019(18)	C(20)-C(21)	1.378(2)
C(6)-C(10)	1.5299(19)	C(20)-H(20)	.9500
C(6)-C(16)	1.5367(19)	C(21)-C(22)	1.384(2)
C(7)-C(8)	1.5276(19)	C(21)-H(21)	.9500
C(7)-H(7A)	.9900	C(22)-H(22)	.9500
C(7)-H(7B)	.9900	C(23)-C(24)	1.3804(19)
C(8)-C(9)	1.4906(19)	C(23)-H(23)	.9500
C(8)-H(8A)	.9900	C(24)-H(24)	.9500
C(8)-H(8B)	.9900	O(2)-C(25)	1.4157(17)
C(10)-C(11)	1.5229(18)	O(2)-H(2A)	.934(16)
C(10)- H(10A)	.9900	C(25)-C(26)	1.503(2)
C(10)- H(10B)	.9900	C(25)- H(25A)	.9900
C(11)-C(12)	1.5392(18)	C(25)-	.9900

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
		H(25B)	
C(11)-H(11A)	.9900	C(26)-H(26A)	.9800
C(11)-H(11B)	.9900	C(26)-H(26B)	.9800
C(12)-C(17)	1.5385(19)	C(26)-H(26C)	.9800
C(12)-C(15)	1.5387(19)		

Table 22B: (Table 22A continued) Bond lengths [Å] and angles [deg] for crystalline form C.

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(7)-O(1)-C(6)	115.64(10)	N(2)-C(13)-H(13A)	109.5
C(5)-N(1)-C(4)	108.27(12)	N(2)-C(13)-H(13B)	109.5
C(5)-N(1)-H(1)	127.5(10)	H(13A)-C(13)-H(13B)	109.5
C(4)-N(1)-H(1)	124.3(10)	N(2)-C(13)-H(13C)	109.5
C(13)-N(2)-C(14)	107.81(12)	H(13A)-C(13)-H(13C)	109.5
C(13)-N(2)-C(12)	113.17(11)	H(13B)-C(13)-H(13C)	109.5
C(14)-N(2)-C(12)	113.94(11)	N(2)-C(14)-H(14A)	109.5
C(2)-C(1)-F(1)	118.28(13)	N(2)-C(14)-H(14B)	109.5
C(2)-C(1)-C(24)	124.68(13)	H(14A)-C(14)-H(14B)	109.5
F(1)-C(1)-C(24)	117.03(12)	N(2)-C(14)-H(14C)	109.5
C(1)-C(2)-C(3)	116.77(13)	H(14A)-C(14)-H(14C)	109.5
C(1)-C(2)-H(2)	121.6	H(14B)-C(14)-H(14C)	109.5
C(3)-C(2)-H(2)	121.6	C(16)-C(15)-C(12)	113.95(11)
C(2)-C(3)-C(4)	119.16(13)	C(16)-C(15)-H(15A)	108.8
C(2)-C(3)-C(9)	134.20(13)	C(12)-C(15)-	108.8

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
		H(15A)	
C(4)-C(3)-C(9)	106.63(12)	C(16)-C(15)- H(15B)	108.8
N(1)-C(4)- C(23)	130.01(13)	C(12)-C(15)- H(15B)	108.8
N(1)-C(4)-C(3)	107.84(12)	H(15A)-C(15)- H(15B)	107.7
C(23)-C(4)- C(3)	122.14(13)	C(15)-C(16)- C(6)	113.84(11)
C(9)-C(5)-N(1)	110.48(12)	C(15)-C(16)- H(16A)	108.8
C(9)-C(5)-C(6)	126.12(13)	C(6)-C(16)- H(16A)	108.8
N(1)-C(5)-C(6)	123.40(12)	C(15)-C(16)- H(16B)	108.8
O(1)-C(6)-C(5)	108.18(11)	C(6)-C(16)- H(16B)	108.8
O(1)-C(6)- C(10)	105.02(11)	H(16A)-C(16)- H(16B)	107.7
C(5)-C(6)- C(10)	111.02(11)	C(18)-C(17)- C(22)	117.14(13)
O(1)-C(6)- C(16)	110.68(10)	C(18)-C(17)- C(12)	120.72(12)
C(5)-C(6)- C(16)	111.50(11)	C(22)-C(17)- C(12)	122.12(13)
C(10)-C(6)- C(16)	110.23(11)	C(19)-C(18)- C(17)	121.38(14)
O(1)-C(7)-C(8)	111.55(12)	C(19)-C(18)- H(18)	119.3
O(1)-C(7)- H(7A)	109.3	C(17)-C(18)- H(18)	119.3
C(8)-C(7)- H(7A)	109.3	C(20)-C(19)- C(18)	120.42(14)
O(1)-C(7)- H(7B)	109.3	C(20)-C(19)- H(19)	119.8
C(8)-C(7)- H(7B)	109.3	C(18)-C(19)- H(19)	119.8
H(7A)-C(7)- H(7B)	108.0	C(21)-C(20)- C(19)	119.39(14)
C(9)-C(8)-C(7)	107.20(11)	C(21)-C(20)-	120.3

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
		H(20)	
C(9)-C(8)- H(8A)	110.3	C(19)-C(20)- H(20)	120.3
C(7)-C(8)- H(8A)	110.3	C(20)-C(21)- C(22)	120.43(14)
C(9)-C(8)- H(8B)	110.3	C(20)-C(21)- H(21)	119.8
C(7)-C(8)- H(8B)	110.3	C(22)-C(21)- H(21)	119.8
H(8A)-C(8)- H(8B)	108.5	C(21)-C(22)- C(17)	121.23(14)
C(5)-C(9)-C(3)	106.78(12)	C(21)-C(22)- H(22)	119.4
C(5)-C(9)-C(8)	120.97(12)	C(17)-C(22)- H(22)	119.4
C(3)-C(9)-C(8)	132.14(13)	C(24)-C(23)- C(4)	118.11(13)
C(11)-C(10)- C(6)	113.43(11)	C(24)-C(23)- H(23)	120.9
C(11)-C(10)- H(10A)	108.9	C(4)-C(23)- H(23)	120.9
C(6)-C(10)- H(10A)	108.9	C(23)-C(24)- C(1)	119.13(13)
C(11)-C(10)- H(10B)	108.9	C(23)-C(24)- H(24)	120.4
C(6)-C(10)- H(10B)	108.9	C(1)-C(24)- H(24)	120.4
H(10A)-C(10)- H(10B)	107.7	C(25)-O(2)- H(2A)	108.0(10)
C(10)-C(11)- C(12)	113.01(11)	O(2)-C(25)- C(26)	108.43(12)
C(10)-C(11)- H(11A)	109.0	O(2)-C(25)- H(25A)	110.0
C(12)-C(11)- H(11A)	109.0	C(26)-C(25)- H(25A)	110.0
C(10)-C(11)- H(11B)	109.0	O(2)-C(25)- H(25B)	110.0
C(12)-C(11)- H(11B)	109.0	C(26)-C(25)- H(25B)	110.0
H(11A)-C(11)-	107.8	H(25A)-C(25)-	108.4

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
H(11B)		H(25B)	
N(2)-C(12)-C(17)	110.91(11)	C(25)-C(26)-H(26A)	109.5
N(2)-C(12)-C(15)	109.29(11)	C(25)-C(26)-H(26B)	109.5
C(17)-C(12)-C(15)	111.53(11)	H(26A)-C(26)-H(26B)	109.5
N(2)-C(12)-C(11)	109.78(10)	C(25)-C(26)-H(26C)	109.5
C(17)-C(12)-C(11)	109.99(11)	H(26A)-C(26)-H(26C)	109.5
C(15)-C(12)-C(11)	105.19(11)	H(26B)-C(26)-H(26C)	109.5

[0429] Symmetry transformations used to generate equivalent atoms:

Table 23: Hydrogen coordinates ($\times 10^4$) (i.e. ($\times 10^4$)) and isotropic displacement parameters ($\times 10^3$) (i.e. ($\times 10^3$)) for crystalline form C.

	x	y	z	U(eq)
H(1)	1646(18)	3427(16)	-927(13)	26(4)
H(2)	1145	5099	-4255	23
H(7A)	3578	9361	211	28
H(7B)	1971	7969	0	28
H(8A)	2160	7612	-1913	26
H(8B)	3966	7917	-1472	26
H(10A)	4245	5221	793	21
H(10B)	5468	6629	706	21
H(11A)	5809	8180	2680	21
H(11B)	6226	6930	2698	21
H(13A)	1682	3425	2864	39
H(13B)	1289	4707	2911	39
H(13C)	2586	4979	4052	39
H(14A)	5068	5040	3864	37
H(14B)	5391	4864	2620	37
H(14C)	4026	3498	2670	37
H(15A)	1979	6700	2944	21

	x	y	z	U(eq)
H(15B)	3293	8062	2858	21
H(16A)	1274	6523	976	22
H(16B)	1418	5127	939	22
H(18)	7018	7506	4516	25
H(19)	8201	8738	6610	29
H(20)	6894	9627	7821	32
H(21)	4352	9207	6922	32
H(22)	3127	7927	4831	28
H(23)	-31	1163	-3242	23
H(24)	-942	714	-5262	24
H(2A)	2332(19)	3467(18)	1062(15)	40
H(25A)	1757	1092	638	35
H(25B)	3331	2032	412	35
H(26A)	271	74	-1442	67
H(26B)	1693	-308	-1286	67
H(26C)	1870	972	-1655	67

Table 24: Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) (i.e. $\text{\AA}^2 \times 10^3$) for crystalline form C. The anisotropic displacement factor exponent takes the form: $-2 \pi^2 [h^2 a^2 U_{11} + \dots + 2 h k a \cdot b \cdot U_{12}]$.

	U11	U22	U33	U23	U13	U12
F(1)	33(1)	32(1)	14(1)	7(1)	4(1)	14(1)
O(1)	23(1)	18(1)	17(1)	6(1)	3(1)	10(1)
N(1)	19(1)	18(1)	17(1)	8(1)	6(1)	11(1)
N(2)	17(1)	20(1)	18(1)	8(1)	7(1)	10(1)
C(1)	18(1)	28(1)	12(1)	7(1)	4(1)	13(1)
C(2)	19(1)	22(1)	19(1)	10(1)	8(1)	11(1)
C(3)	13(1)	20(1)	18(1)	7(1)	6(1)	10(1)
C(4)	12(1)	21(1)	15(1)	7(1)	6(1)	10(1)
C(5)	15(1)	17(1)	18(1)	6(1)	7(1)	9(1)
C(6)	17(1)	18(1)	15(1)	6(1)	4(1)	9(1)
C(7)	29(1)	20(1)	21(1)	9(1)	7(1)	13(1)
C(8)	25(1)	20(1)	18(1)	8(1)	5(1)	11(1)
C(9)	15(1)	20(1)	16(1)	7(1)	6(1)	10(1)
C(10)	17(1)	20(1)	16(1)	7(1)	7(1)	10(1)
C(11)	15(1)	22(1)	16(1)	8(1)	6(1)	10(1)

	U11	U22	U33	U23	U13	U12
C(12)	15(1)	19(1)	14(1)	5(1)	5(1)	10(1)
C(13)	26(1)	25(1)	28(1)	12(1)	14(1)	11(1)
C(14)	27(1)	24(1)	25(1)	9(1)	7(1)	16(1)
C(15)	17(1)	22(1)	18(1)	8(1)	9(1)	12(1)
C(16)	16(1)	21(1)	18(1)	7(1)	6(1)	11(1)
C(17)	20(1)	15(1)	17(1)	7(1)	6(1)	7(1)
C(18)	19(1)	22(1)	19(1)	9(1)	6(1)	9(1)
C(19)	24(1)	21(1)	22(1)	10(1)	2(1)	7(1)
C(20)	38(1)	20(1)	14(1)	6(1)	3(1)	12(1)
C(21)	39(1)	25(1)	20(1)	9(1)	14(1)	19(1)
C(22)	24(1)	25(1)	21(1)	10(1)	8(1)	14(1)
C(23)	18(1)	19(1)	22(1)	9(1)	8(1)	11(1)
C(24)	16(1)	20(1)	19(1)	3(1)	5(1)	9(1)
O(2)	34(1)	25(1)	19(1)	8(1)	5(1)	17(1)
C(25)	37(1)	22(1)	26(1)	9(1)	4(1)	16(1)
C(26)	66(1)	30(1)	30(1)	6(1)	-2(1)	29(1)

Crystalline form D**[0430]**

Table 25: Crystal data and structure refinement for crystalline form D.

Empirical formula	C ₂₇ H ₃₅ FN ₂ O ₂
Formula weight	438.57
Temperature	100(2) K
Wavelength	0.71073 Å
Crystal system	Triclinic
Space group	P-1
Unit cell dimensions	a = 9.909(3) Å alpha = 115.704(5) deg.
	b = 11.701(4) Å beta = 106.324(5) deg.
	c = 11.830(4) Å gamma = 94.399(6) deg.
Volume	1154.4(7) Å ³
Z	2
Density (calculated)	1.262 Mg/m ³

Absorption coefficient	0.085 mm ⁻¹
F(000)	472
Crystal size	0.29 x 0.12 x 0.08 mm
Theta range for data collection	1.98 to 25.10 deg.
Index ranges	-11 ≤ h ≤ 11, -13 ≤ k ≤ 13, -13 ≤ l ≤ 14
Reflections collected	18119
Independent reflections	4085 [R(int) = 0.0836]
Absorption correction	None
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	4085/ 1 /303
Goodness-of-fit on F ²	1.082
Final R indices [I > 2σ(I)]	R1 = 0.0651, wR2 = 0.1559
R indices (all data)	R1 = 0.0931, wR2 = 0.1710
Largest diff. peak and hole	0.510 and -0.325 eÅ ⁻³

Table 26: Atomic coordinates ($\times 10^4$) (i.e. ($\times 10^4$)) and equivalent isotropic displacement parameters ($\times 10^3$) (i.e. ($\times 10^3$)) for crystalline form D. U(eq) is defined as one third of the trace of the orthogonalized Uij tensor.

	x	y	z	U(eq)
F(1)	866(2)	-3098(2)	-2007(2)	48(1)
O(1)	3245(2)	2553(2)	5183(2)	28(1)
N(1)	2086(3)	-1043(2)	3364(3)	25(1)
N(2)	2884(3)	630(2)	7839(2)	24(1)
C(1)	1160(4)	-2626(3)	-659(3)	35(1)
C(2)	1496(3)	-1297(3)	139(3)	32(1)
C(3)	1809(3)	-831(3)	1513(3)	26(1)
C(4)	1739(3)	-1733(3)	2007(3)	24(1)
C(5)	2381(3)	267(3)	3741(3)	24(1)
C(6)	2824(3)	1330(3)	5162(3)	24(1)
C(7)	2321(4)	2703(3)	4107(3)	34(1)
C(8)	2379(3)	1766(3)	2753(3)	30(1)
C(9)	2217(3)	438(3)	2646(3)	26(1)
C(10)	4171(3)	1180(3)	6060(3)	26(1)
C(11)	4558(3)	2091(3)	7557(3)	26(1)
C(12)	3307(3)	2005(3)	8070(3)	24(1)
C(13)	1527(3)	350(3)	8042(3)	31(1)

	x	y	z	U(eq)
C(14)	4010(4)	286(3)	8665(3)	34(1)
C(15)	2055(3)	2303(3)	7200(3)	26(1)
C(16)	1596(3)	1359(3)	5713(3)	25(1)
C(17)	3729(3)	2984(3)	9560(3)	28(1)
C(18)	5164(4)	3456(3)	10423(3)	33(1)
C(19)	5532(4)	4258(3)	11782(3)	36(1)
C(20)	4475(4)	4616(3)	12336(3)	36(1)
C(21)	3047(4)	4164(3)	11498(3)	35(1)
C(22)	2673(4)	3371(3)	10135(3)	31(1)
C(23)	1402(3)	-3077(3)	1163(3)	29(1)
C(24)	1115(3)	-3518(3)	-188(3)	34(1)
O(2)	1858(3)	-1693(2)	5352(2)	38(1)
C(25A)	2047(4)	-2854(3)	5459(3)	31(1)
C(26)	3583(4)	-2995(4)	5551(5)	46(1)
C(27)	930(4)	-3962(4)	4274(4)	39(1)
C(25B)	2047(4)	-2854(3)	5459(3)	31(1)
C(28)	2190(30)	-3910(20)	4320(20)	30

Table 27A: Bond lengths [Å] and angles [deg] for crystalline form D.

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
F(1)-C(1)	1.372(4)	C(14)-H(14A)	0.9800
O(1)-C(7)	1.433(4)	C(14)-H(14B)	0.9800
O(1)-C(6)	1.447(3)	C(14)-H(14C)	0.9800
N(1)-C(4)	1.368(4)	C(15)-C(16)	1.520(4)
N(1)-C(5)	1.378(4)	C(15)-H(15A)	0.9900
N(1)-H(1 N)	0.92(3)	C(15)-H(15B)	0.9900
N(2)-C(13)	1.469(4)	C(16)-H(16A)	0.9900
N(2)-C(14)	1.469(4)	C(16)-H(16B)	0.9900
N(2)-C(12)	1.513(4)	C(17)-C(18)	1.393(4)

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(1)-C(2)	1.375(5)	C(17)- C(22)	1.404(4)
C(1)-C(24)	1.383(5)	C(18)- C(19)	1.381(5)
C(2)-C(3)	1.399(4)	C(18)- H(18)	0.9500
C(2)-H(2)	0.9500	C(19)- C(20)	1.384(5)
C(3)-C(4)	1.417(4)	C(19)- H(19)	0.9500
C(3)-C(9)	1.429(4)	C(20)- C(21)	1.381(5)
C(4)-C(23)	1.397(4)	C(20)- H(20)	0.9500
C(5)-C(9)	1.363(4)	C(21)- C(22)	1.383(4)
C(5)-C(6)	1.501(4)	C(21)- H(21)	0.9500
C(6)-C(16)	1.529(4)	C(22)- H(22)	0.9500
C(6)-C(10)	1.530(4)	C(23)- C(24)	1.381(4)
C(7)-C(8)	1.520(4)	C(23)- H(23)	0.9500
C(7)-H(7A)	0.9900	C(24)- H(24)	0.9500
C(7)-H(7B)	0.9900	O(2)- C(25A)	1.439(4)
C(8)-C(9)	1.495(4)	O(2)-H(2A)	0.8400
C(8)-H(8A)	0.9900	C(25A)- C(27)	1.484(5)
C(8)-H(8B)	0.9900	C(25A)- C(26)	1.522(5)
C(10)-C(11)	1.530(4)	C(25A)- H(25)	1.0000
C(10)- H(10A)	0.9900	C(26)- H(26A)	0.9800
C(10)- H(10B)	0.9900	C(26)- H(26B)	0.9800
C(11)-C(12)	1.539(4)	C(26)-	0.9800

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
		H(26C)	
C(11)-H(11A)	0.9900	C(27)-H(27A)	0.9800
C(11)-H(11B)	0.9900	C(27)-H(27B)	0.9800
C(12)-C(17)	1.534(4)	C(27)-H(27C)	0.9800
C(12)-C(15)	1.543(4)	C(28)-H(28A)	0.9800
C(13)-H(13A)	0.9800	C(28)-H(28B)	0.9800
C(13)-H(13B)	0.9800	C(28)-H(28C)	0.9800
C(13)-H(13C)	0.9800		

Table 27B: (Table 27A continued) Bond lengths [Å] and angles [deg] for crystalline form D.

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(7)-O(1)-C(6)	115.0(2)	C(11)-C(12)-C(15)	105.1(2)
C(4)-N(1)-C(5)	108.5(3)	N(2)-C(13)-H(13A)	109.5
C(4)-N(1)-H(1N)	126(2)	N(2)-C(13)-H(13B)	109.5
C(5)-N(1)-H(1N)	125(2)	H(13A)-C(13)-H(13B)	109.5
C(13)-N(2)-C(14)	108.2(2)	N(2)-C(13)-H(13C)	109.5
C(13)-N(2)-C(12)	114.0(2)	H(13A)-C(13)-H(13C)	109.5
C(14)-N(2)-C(12)	113.1(2)	H(13B)-C(13)-H(13C)	109.5
F(1)-C(1)-C(2)	117.9(3)	N(2)-C(14)-H(14A)	109.5
F(1)-C(1)-C(24)	117.9(3)	N(2)-C(14)-H(14B)	109.5
C(2)-C(1)-C(24)	124.2(3)	H(14A)-C(14)-H(14B)	109.5
C(1)-C(2)-C(3)	117.3(3)	N(2)-C(14)-H(14C)	109.5

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(1)-C(2)-H(2)	121.4	H(14A)-C(14)- H(14C)	109.5
C(3)-C(2)-H(2)	121.4	H(14B)-C(14)- H(14C)	109.5
C(2)-C(3)-C(4)	119.2(3)	C(16)-C(15)- C(12)	113.3(2)
C(2)-C(3)-C(9)	134.3(3)	C(16)-C(15)- H(15A)	108.9
C(4)-C(3)-C(9)	106.5(3)	C(12)-C(15)- H(15A)	108.9
N(1)-C(4)- C(23)	130.3(3)	C(16)-C(15)- H(15B)	108.9
N(1)-C(4)-C(3)	108.0(3)	C(12)-C(15)- H(15B)	108.9
C(23)-C(4)- C(3)	121.7(3)	H(15A)-C(15)- H(15B)	107.7
C(9)-C(5)-N(1)	110.1(3)	C(15)-C(16)- C(6)	113.1(2)
C(9)-C(5)-C(6)	126.0(3)	C(15)-C(16)- H(16A)	109.0
N(1)-C(5)-C(6)	123.9(3)	C(6)-C(16)- H(16A)	109.0
O(1)-C(6)-C(5)	107.7(2)	C(15)-C(16)- H(16B)	109.0
O(1)-C(6)- C(16)	110.7(2)	C(6)-C(16)- H(16B)	109.0
C(5)-C(6)- C(16)	111.4(2)	H(16A)-C(16)- H(16B)	107.8
O(1)-C(6)- C(10)	105.7(2)	C(18)-C(17)- C(22)	116.8(3)
C(5)-C(6)- C(10)	111.1(2)	C(18)-C(17)- C(12)	121.9(3)
C(16)-C(6)- C(10)	110.1(2)	C(22)-C(17)- C(12)	121.1(3)
O(1)-C(7)-C(8)	112.7(3)	C(19)-C(18)- C(17)	121.8(3)
O(1)-C(7)- H(7A)	109.0	C(19)-C(18)- H(18)	119.1
C(8)-C(7)- H(7A)	109.0	C(17)-C(18)- H(18)	119.1

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
O(1)-C(7)- H(7B)	109.0	C(18)-C(19)- C(20)	120.7(3)
C(8)-C(7)- H(7B)	109.0	C(18)-C(19)- H(19)	119.7
H(7A)-C(7)- H(7B)	107.8	C(20)-C(19)- H(19)	119.7
C(9)-C(8)-C(7)	107.4(3)	C(21)-C(20)- C(19)	118.5(3)
C(9)-C(8)- H(8A)	110.2	C(21)-C(20)- H(20)	120.7
C(7)-C(8)- H(8A)	110.2	C(19)-C(20)- H(20)	120.7
C(9)-C(8)- H(8B)	110.2	C(20)-C(21)- C(22)	121.0(3)
C(7)-C(8)- H(8B)	110.2	C(20)-C(21)- H(21)	119.5
H(8A)-C(8)- H(8B)	108.5	C(22)-C(21)- H(21)	119.5
C(5)-C(9)-C(3)	106.9(3)	C(21)-C(22)- C(17)	121.2(3)
C(5)-C(9)-C(8)	121.5(3)	C(21)-C(22)- H(22)	119.4
C(3)-C(9)-C(8)	131.5(3)	C(17)-C(22)- H(22)	119.4
C(11)-C(10)- C(6)	114.4(2)	C(24)-C(23)- C(4)	118.2(3)
C(11)-C(10)- H(10A)	108.7	C(24)-C(23)- H(23)	120.9
C(6)-C(10)- H(10A)	108.7	C(4)-C(23)- H(23)	120.9
C(11)-C(10)- H(10B)	108.7	C(23)-C(24)- C(1)	119.4(3)
C(6)-C(10)- H(10B)	108.7	C(23)-C(24)- H(24)	120.3
H(10A)-C(10)- H(10B)	107.6	C(1)-C(24)- H(24)	120.3
C(10)-C(11)- C(12)	113.7(2)	C(25A)-O(2)- H(2A)	109.5
C(10)-C(11)- H(11A)	108.8	O(2)-C(25A)- C(27)	107.4(3)

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(12)-C(11)-H(11A)	108.8	O(2)-C(25A)-C(26)	108.6(3)
C(10)-C(11)-H(11B)	108.8	C(27)-C(25A)-C(26)	113.3(3)
C(12)-C(11)-H(11B)	108.8	O(2)-C(25A)-H(25)	109.2
H(11A)-C(11)-H(11B)	107.7	C(27)-C(25A)-H(25)	109.2
N(2)-C(12)-C(17)	110.4(2)	C(26)-C(25A)-H(25)	109.2
N(2)-C(12)-C(11)	108.7(2)	H(28A)-C(28)-H(28B)	109.5
C(17)-C(12)-C(11)	111.5(2)	H(28A)-C(28)-H(28C)	109.5
N(2)-C(12)-C(15)	109.9(2)	H(28B)-C(28)-H(28C)	109.5
C(17)-C(12)-C(15)	111.2(2)		

[0431] Symmetry transformations used to generate equivalent atoms:

Table 28: Hydrogen coordinates ($\times 10^4$) (i.e. ($\times 10^4$)) and isotropic displacement parameters ($\times 10^3$) (i.e. ($\times 10^3$)) for crystalline form D.

	x	y	z	U(eq)
H(1N)	2050(30)	-1390(30)	3930(30)	31(9)
H(2)	1515	-719	-227	39
H(7A)	2608	3608	4278	40
H(7B)	1314	2557	4085	40
H(8A)	1589	1770	2024	36
H(8B)	3312	2029	2681	36
H(10A)	4019	269	5899	31
H(10B)	5001	1344	5797	31
H(11A)	4864	2999	7743	31
H(11B)	5389	1878	8062	31
H(13A)	1274	-575	7793	46
H(13B)	755	554	7483	46
H(13C)	1647	884	8986	46

	x	y	z	U(eq)
H(14A)	4172	849	9614	52
H(14B)	4910	406	8495	52
H(14C)	3700	-626	8437	52
H(15A)	1215	2271	7490	32
H(15B)	2355	3199	7350	32
H(16A)	797	1609	5217	30
H(16B)	1226	473	5554	30
H(18)	5908	3219	10067	39
H(19)	6521	4567	12343	43
H(20)	4727	5162	13272	44
H(21)	2309	4401	11864	42
H(22)	1684	3083	9578	37
H(23)	1371	-3672	1508	35
H(24)	888	-4425	-788	40
H(2A)	2477	-1047	6013	56
H(25)	1897	-2770	6290	37
H(26A)	3734	-3084	4735	70
H(26B)	3736	-3770	5647	70
H(26C)	4269	-2221	6329	70
H(27A)	-32	-3818	4272	58
H(27B)	1034	-4772	4314	58
H(27C)	1051	-4029	3455	58
H(25A)	2918	-2619	6248	37
H(25B)	1209	-3169	5631	37
H(28A)	2030	-3687	3588	45
H(28B)	1481	-4699	4026	45
H(28C)	3168	-4058	4560	45

Table 29: Anisotropic displacement parameters ($\times 10^3$) (i.e. $\text{\AA}^2 \times 10^3$) for crystalline form D. The anisotropic displacement factor exponent takes the form: $-2 \pi^2 [h^2 a^{*2} U_{11} + \dots + 2 h k a^* b^* U_{12}]$.

	U11	U22	U33	U23	U13	U12
F(1)	56(1)	51(1)	28(1)	10(1)	17(1)	9(1)
O(1)	24(1)	26(1)	35(1)	15(1)	9(1)	6(1)
N(1)	21(1)	27(1)	29(2)	14(1)	11(1)	10(1)
N(2)	17(1)	28(1)	29(1)	13(1)	9(1)	9(1)

	U11	U22	U33	U23	U13	U12
C(1)	25(2)	47(2)	29(2)	13(2)	13(1)	10(2)
C(2)	28(2)	38(2)	34(2)	19(2)	13(2)	9(2)
C(3)	15(2)	34(2)	31(2)	15(1)	10(1)	9(1)
C(4)	13(2)	29(2)	30(2)	11(1)	8(1)	7(1)
C(5)	15(2)	25(2)	32(2)	12(1)	8(1)	8(1)
C(6)	18(2)	23(2)	33(2)	14(1)	11(1)	7(1)
C(7)	30(2)	32(2)	42(2)	21(2)	11(2)	12(2)
C(8)	26(2)	30(2)	37(2)	19(2)	10(1)	7(1)
C(9)	15(2)	33(2)	33(2)	16(2)	10(1)	9(1)
C(10)	18(2)	28(2)	32(2)	14(1)	11(1)	10(1)
C(11)	18(2)	30(2)	32(2)	14(1)	10(1)	7(1)
C(12)	16(2)	24(2)	31(2)	11(1)	9(1)	7(1)
C(13)	26(2)	32(2)	38(2)	16(2)	16(2)	8(1)
C(14)	30(2)	31(2)	39(2)	16(2)	6(2)	13(2)
C(15)	16(2)	30(2)	35(2)	14(1)	12(1)	13(1)
C(16)	18(2)	26(2)	32(2)	13(1)	10(1)	10(1)
C(17)	25(2)	24(2)	32(2)	11(1)	10(1)	10(1)
C(18)	25(2)	34(2)	35(2)	14(2)	10(2)	6(1)
C(19)	27(2)	37(2)	34(2)	11(2)	6(2)	4(2)
C(20)	43(2)	27(2)	33(2)	8(2)	15(2)	11 (2)
C(21)	40(2)	31(2)	38(2)	14(2)	20(2)	18(2)
C(22)	28(2)	28(2)	35(2)	11(2)	13(1)	14(1)
C(23)	19(2)	30(2)	37(2)	13(2)	12(1)	9(1)
C(24)	22(2)	34(2)	37(2)	10(2)	12(1)	7(1)
O(2)	37(1)	33(1)	42(1)	17(1)	12(1)	11(1)
C(25A)	32(2)	28(2)	40(2)	19(2)	18(2)	15(1)
C(26)	29(2)	40(2)	70(3)	24(2)	18(2)	22(2)
C(27)	30(2)	28(2)	49(2)	10(2)	13(2)	12(2)
C(25B)	32(2)	28(2)	40(2)	19(2)	18(2)	15(1)

Crystalline form G**[0432]**

Table 30: Crystal data and structure refinement for crystalline form G.

Empirical formula	C ₂₆ H ₃₃ FN ₂ O ₂ S
Formula weight	456.60
Temperature	130(2) K
Wavelength	.71073 Å
Crystal system	Monoclinic
Space group	P 21/c
Unit cell dimensions	a = 10.8279(12) Å alpha = 90 deg.
	b = 11.5334(12) Å beta = 99.991(2) deg.
	c = 18.822(2) Å gamma = 90 deg.
Volume	2314.9(4) Å ³
Z	4
Density (calculated)	1.310 Mg/m ³
Absorption coefficient	0.174 mm ⁻¹
F(000)	976
Crystal size	0.57 x 0.35 x 0.13 mm
Theta range for data collection	2.20 to 33.19 deg.
Index ranges	-16 ≤ h ≤ 15, -17 ≤ k ≤ 16, -29 ≤ l ≤ 27
Reflections collected	37663
Independent reflections	8514 [R(int) = 0.0802]
Absorption correction	None
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	8514 / 0 / 297
Goodness-of-fit on F ²	1.070
Final R indices [I > 2σ(I)]	R1 = 0.0516, wR2 = 0.1253
R indices (all data)	R1 = 0.0639, wR2 = 0.1301
Largest diff. peak and hole	1.045 and -.349 e.Å ⁻³

Table 31: Atomic coordinates ($\times 10^4$) (i.e. ($\times 10^4$)) and equivalent isotropic displacement parameters ($\times 10^3$) (i.e. ($\times 10^3$)) for crystalline form G. U(eq) is defined as one third of the trace of the orthogonalized Uij tensor.

	x	y	z	U(eq)
F(1)	8199(1)	5868(1)	4452(1)	31(1)
O(1)	2509(1)	1818(1)	3420(1)	19(1)
N(1)	5966(1)	1682(1)	3844(1)	13(1)
N(2)	4244(1)	-1808(1)	3634(1)	14(1)
C(1)	7669(1)	4803(1)	4280(1)	20(1)

	x	y	z	U(eq)
C(2)	6385(1)	4727(1)	4144(1)	19(1)
C(3)	5870(1)	3620(1)	3988(1)	14(1)
C(4)	6686(1)	2661(1)	3990(1)	13(1)
C(5)	4726(1)	1997(1)	3749(1)	13(1)
C(6)	3658(1)	1161(1)	3598(1)	13(1)
C(7)	2417(1)	2800(1)	3872(1)	22(1)
C(8)	3370(1)	3740(1)	3784(1)	19(1)
C(9)	4621(1)	3170(1)	3833(1)	14(1)
C(10)	3743(1)	428(1)	2931(1)	16(1)
C(11)	2802(1)	-566(1)	2813(1)	16(1)
C(12)	2924(1)	-1382(1)	3469(1)	12(1)
C(13)	4584(1)	-2440(1)	4309(1)	17(1)
C(14)	4629(1)	-2483(1)	3055(1)	23(1)
C(15)	2667(1)	-604(1)	4094(1)	13(1)
C(16)	3624(1)	374(1)	4252(1)	13(1)
C(17)	2000(1)	-2397(1)	3342(1)	14(1)
C(18)	1739(1)	-3057(1)	3920(1)	15(1)
C(19)	1010(1)	-4051(1)	3811(1)	18(1)
C(20)	498(1)	-4404(1)	3120(1)	22(1)
C(21)	718(2)	-3751(1)	2541(1)	28(1)
C(22)	1463(1)	-2761(1)	2651(1)	22(1)
C(23)	7986(1)	2773(1)	4124(1)	15(1)
C(24)	8478(1)	3872(1)	4270(1)	18(1)
S(1)	2104(1)	3668(1)	948(1)	22(1)
O(2)	2254(1)	4947(1)	1046(1)	39(1)
C(25)	684(1)	3480(1)	320(1)	25(1)
C(26)	1565(2)	3135(2)	1726(1)	35(1)

Table 32A: Bond lengths [Å] and angles [deg] for crystalline form G.

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
F(1)-C(1)	1.3707(14)	C(13)-H(13A)	.9800
O(1)-C(7)	1.4303(16)	C(13)-H(13B)	.9800
O(1)-C(6)	1.4452(14)	C(13)-H(13C)	.9800

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
N(1)-C(5)	1.3718(15)	C(14)- H(14A)	.9800
N(1)-C(4)	1.3729(15)	C(14)- H(14B)	.9800
N(1)-H(1N)	.781(17)	C(14)- H(14C)	.9800
N(2)-C(13)	1.4553(15)	C(15)-C(16)	1.5258(16)
N(2)-C(14)	1.4579(15)	C(15)- H(15A)	.9900
N(2)-C(12)	1.4923(15)	C(15)- H(15B)	.9900
C(1)-C(2)	1.3716(18)	C(16)- H(16A)	.9900
C(1)-C(24)	1.3888(19)	C(16)- H(16B)	.9900
C(2)-C(3)	1.4045(16)	C(17)-C(22)	1.3940(16)
C(2)-H(2)	.9500	C(17)-C(18)	1.3969(16)
C(3)-C(4)	1.4154(16)	C(18)-C(19)	1.3870(16)
C(3)-C(9)	1.4304(16)	C(18)-H(18)	.9500
C(4)-C(23)	1.3923(16)	C(19)-C(20)	1.3842(18)
C(5)-C(9)	1.3695(16)	C(19)-H(19)	.9500
C(5)-C(6)	1.4951(16)	C(20)-C(21)	1.379(2)
C(6)-C(10)	1.5298(16)	C(20)-H(20)	.9500
C(6)-C(16)	1.5350(16)	C(21)-C(22)	1.3920(18)
C(7)-C(8)	1.5247(18)	C(21)-H(21)	.9500
C(7)-H(7A)	.9900	C(22)-H(22)	.9500
C(7)-H(7B)	.9900	C(23)-C(24)	1.3837(17)
C(8)-C(9)	1.4941(16)	C(23)-H(23)	.9500
C(8)-H(8A)	.9900	C(24)-H(24)	.9500
C(8)-H(8B)	.9900	S(1)-O(2)	1.4910(11)
C(10)-C(11)	1.5231(16)	S(1)-C(26)	1.7784(16)
C(10)- H(10A)	.9900	S(1)-C(25)	1.7825(14)
C(10)- H(10B)	.9900	C(25)- H(25A)	.9800
C(11)-C(12)	1.5401(16)	C(25)- H(25B)	.9800

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(11)-H(11A)	.9900	C(25)-H(25C)	.9800
C(11)-H(11B)	.9900	C(26)-H(26A)	.9800
C(12)-C(17)	1.5314(15)	C(26)-H(26B)	.9800
C(12)-C(15)	1.5427(15)	C(26)-H(26C)	.9800

Table 32B: (Table 32A continued) Bond lengths [Å] and angles [deg] for crystalline form G.

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(7)-O(1)-C(6)	115.09(9)	N(2)-C(13)-H(13B)	109.5
C(5)-N(1)-C(4)	108.49(10)	H(13A)-C(13)-H(13B)	109.5
C(5)-N(1)-H(1N)	127.7(12)	N(2)-C(13)-H(13C)	109.5
C(4)-N(1)-H(1N)	123.8(12)	H(13A)-C(13)-H(13C)	109.5
C(13)-N(2)-C(14)	108.86(10)	H(13B)-C(13)-H(13C)	109.5
C(13)-N(2)-C(12)	115.75(9)	N(2)-C(14)-H(14A)	109.5
C(14)-N(2)-C(12)	114.08(9)	N(2)-C(14)-H(14B)	109.5
F(1)-C(1)-C(2)	118.04(12)	H(14A)-C(14)-H(14B)	109.5
F(1)-C(1)-C(24)	117.21(11)	N(2)-C(14)-H(14C)	109.5
C(2)-C(1)-C(24)	124.74(11)	H(14A)-C(14)-H(14C)	109.5
C(1)-C(2)-C(3)	116.71(11)	H(14B)-C(14)-H(14C)	109.5
C(1)-C(2)-H(2)	121.6	C(16)-C(15)-C(12)	111.85(9)
C(3)-C(2)-H(2)	121.6	C(16)-C(15)-H(15A)	109.2
C(2)-C(3)-C(4)	119.04(11)	C(12)-C(15)-H(15A)	109.2
C(2)-C(3)-C(9)	134.39(11)	C(16)-C(15)-	109.2

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
		H(15B)	
C(4)-C(3)-C(9)	106.55(10)	C(12)-C(15)- H(15B)	109.2
N(1)-C(4)- C(23)	129.36(11)	H(15A)-C(15)- H(15B)	107.9
N(1)-C(4)-C(3)	108.03(10)	C(15)-C(16)- C(6)	113.01(9)
C(23)-C(4)- C(3)	122.60(11)	C(15)-C(16)- H(16A)	109.0
C(9)-C(5)-N(1)	110.24(10)	C(6)-C(16)- H(16A)	109.0
C(9)-C(5)-C(6)	125.60(10)	C(15)-C(16)- H(16B)	109.0
N(1)-C(5)-C(6)	124.14(10)	C(6)-C(16)- H(16B)	109.0
O(1)-C(6)-C(5)	108.22(9)	H(16A)-C(16)- H(16B)	107.8
O(1)-C(6)- C(10)	105.83(9)	C(22)-C(17)- C(18)	117.06(11)
C(5)-C(6)- C(10)	110.99(9)	C(22)-C(17)- C(12)	122.12(10)
O(1)-C(6)- C(16)	110.82(9)	C(18)-C(17)- C(12)	120.63(10)
C(5)-C(6)- C(16)	110.71(9)	C(19)-C(18)- C(17)	121.45(11)
C(10)-C(6)- C(16)	110.16(9)	C(19)-C(18)- H(18)	119.3
O(1)-C(7)-C(8)	112.34(11)	C(17)-C(18)- H(18)	119.3
O(1)-C(7)- H(7A)	109.1	C(20)-C(19)- C(18)	120.49(12)
C(8)-C(7)- H(7A)	109.1	C(20)-C(19)- H(19)	119.8
O(1)-C(7)- H(7B)	109.1	C(18)-C(19)- H(19)	119.8
C(8)-C(7)- H(7B)	109.1	C(21)-C(20)- C(19)	119.06(12)
H(7A)-C(7)- H(7B)	107.9	C(21)-C(20)- H(20)	120.5
C(9)-C(8)-C(7)	107.71(10)	C(19)-C(20)-	120.5

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
		H(20)	
C(9)-C(8)- H(8A)	110.2	C(20)-C(21)- C(22)	120.39(12)
C(7)-C(8)- H(8A)	110.2	C(20)-C(21)- H(21)	119.8
C(9)-C(8)- H(8B)	110.2	C(22)-C(21)- H(21)	119.8
C(7)-C(8)- H(8B)	110.2	C(21)-C(22)- C(17)	121.52(12)
H(8A)-C(8)- H(8B)	108.5	C(21)-C(22)- H(22)	119.2
C(5)-C(9)-C(3)	106.69(10)	C(17)-C(22)- H(22)	119.2
C(5)-C(9)-C(8)	121.33(11)	C(24)-C(23)- C(4)	117.62(11)
C(3)-C(9)-C(8)	131.94(11)	C(24)-C(23)- H(23)	121.2
C(11)-C(10)- C(6)	113.66(10)	C(4)-C(23)- H(23)	121.2
C(11)-C(10)- H(10A)	108.8	C(23)-C(24)- C(1)	119.27(11)
C(6)-C(10)- H(10A)	108.8	C(23)-C(24)- H(24)	120.4
C(11)-C(10)- H(10B)	108.8	C(1)-C(24)- H(24)	120.4
C(6)-C(10)- H(10B)	108.8	O(2)-S(1)- C(26)	106.52(8)
H(10A)-C(10)- H(10B)	107.7	O(2)-S(1)-C(25)	105.40(7)
C(10)-C(11)- C(12)	112.41(9)	C(26)-S(1)- C(25)	97.75(8)
C(10)-C(11)- H(11A)	109.1	S(1)-C(25)- H(25A)	109.5
C(12)-C(11)- H(11A)	109.1	S(1)-C(25)- H(25B)	109.5
C(10)-C(11)- H(11B)	109.1	H(25A)-C(25)- H(25B)	109.5
C(12)-C(11)- H(11B)	109.1	S(1)-C(25)- H(25C)	109.5
H(11A)-C(11)-	107.9	H(25A)-C(25)-	109.5

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
H(11B)		H(25C)	
N(2)-C(12)-C(17)	110.91(9)	H(25B)-C(25)-H(25C)	109.5
N(2)-C(12)-C(11)	108.28(9)	S(1)-C(26)-H(26A)	109.5
C(17)-C(12)-C(11)	112.27(9)	S(1)-C(26)-H(26B)	109.5
N(2)-C(12)-C(15)	109.03(9)	H(26A)-C(26)-H(26B)	109.5
C(17)-C(12)-C(15)	111.38(9)	S(1)-C(26)-H(26C)	109.5
C(11)-C(12)-C(15)	104.72(9)	H(26A)-C(26)-H(26C)	109.5
N(2)-C(13)-H(13A)	109.5	H(26B)-C(26)-H(26C)	109.5

[0433] Symmetry transformations used to generate equivalent atoms:

Table 33: Hydrogen coordinates ($\times 10^4$) (i.e. ($\times 10^4$)) and isotropic displacement parameters ($\times 10^3$) (i.e. ($\times 10^3$)) for crystalline form G.

	x	y	z	U(eq)
H(2)	5870	5392	4155	23
H(7A)	2552	2546	4382	26
H(7B)	1562	3129	3753	26
H(8A)	3134	4127	3311	23
H(8B)	3398	4330	4168	23
H(10A)	3603	935	2500	19
H(10B)	4600	104	2978	19
H(11A)	2933	-1017	2385	19
H(11B)	1943	-243	2714	19
H(13A)	4157	-3193	4270	26
H(13B)	4329	-1993	4702	26
H(13C)	5493	-2561	4410	26
H(14A)	5521	-2678	3185	35
H(14B)	4491	-2027	2609	35
H(14C)	4134	-3198	2982	35
H(15A)	1816	-268	3968	16

	x	y	z	U(eq)
H(15B)	2693	-1082	4533	16
H(16A)	4466	35	4409	16
H(16B)	3421	850	4654	16
H(18)	2068	-2819	4399	19
H(19)	861	-4493	4214	22
H(20)	2	-5087	3045	27
H(21)	360	-3977	2065	33
H(22)	1609	-2324	2246	27
H(23)	8516	2118	4116	18
H(24)	9360	3987	4362	22
H(25A)	7	3909	486	37
H(25B)	470	2654	282	37
H(25C)	795	3773	-154	37
H(26A)	2239	3196	2145	53
H(26B)	1315	2322	1650	53
H(26C)	843	3594	1811	53
H(1N)	6242(15)	1058(15)	3829(9)	19(4)

Table 34: Anisotropic displacement parameters ($^2 \times 10^3$) (i.e. ($^2 \times 10^3$)) for crystalline form G. The anisotropic displacement factor exponent takes the form: $-2 \pi^2 [h^2 a^{*2} U_{11} + \dots + 2 h k a^* b^* U_{12}]$.

	U11	U22	U33	U23	U13	U12
F(1)	32(1)	16(1)	46(1)	-5(1)	7(1)	-12(1)
O(1)	15(1)	16(1)	26(1)	1(1)	-2(1)	2(1)
N(1)	14(1)	9(1)	16(1)	0(1)	2(1)	1(1)
N(2)	14(1)	15(1)	12(1)	-2(1)	4(1)	0(1)
C(1)	26(1)	13(1)	21(1)	-1(1)	5(1)	-7(1)
C(2)	25(1)	12(1)	21(1)	0(1)	6(1)	-1(1)
C(3)	18(1)	12(1)	13(1)	1(1)	3(1)	0(1)
C(4)	17(1)	12(1)	10(1)	0(1)	3(1)	-1(1)
C(5)	14(1)	12(1)	12(1)	1(1)	2(1)	1(1)
C(6)	13(1)	11(1)	15(1)	1(1)	1(1)	0(1)
C(7)	16(1)	17(1)	32(1)	0(1)	6(1)	4(1)
C(8)	17(1)	14(1)	26(1)	0(1)	4(1)	4(1)
C(9)	16(1)	12(1)	15(1)	1(1)	3(1)	1(1)
C(10)	21(1)	16(1)	12(1)	1(1)	3(1)	-4(1)

	U11	U22	U33	U23	U13	U12
C(11)	19(1)	17(1)	10(1)	1(1)	0(1)	-4(1)
C(12)	14(1)	13(1)	10(1)	-1(1)	2(1)	-2(1)
C(13)	17(1)	16(1)	17(1)	2(1)	2(1)	2(1)
C(14)	24(1)	28(1)	20(1)	-8(1)	9(1)	4(1)
C(15)	15(1)	13(1)	12(1)	-1(1)	4(1)	-1(1)
C(16)	16(1)	13(1)	11(1)	-1(1)	2(1)	-1(1)
C(17)	14(1)	14(1)	14(1)	-2(1)	3(1)	-2(1)
C(18)	15(1)	17(1)	15(1)	-1(1)	3(1)	-3(1)
C(19)	17(1)	17(1)	21(1)	1(1)	4(1)	-4(1)
C(20)	20(1)	19(1)	26(1)	-3(1)	1(1)	-7(1)
C(21)	35(1)	28(1)	18(1)	-6(1)	0(1)	-14(1)
C(22)	29(1)	24(1)	14(1)	-2(1)	2(1)	-10(1)
C(23)	16(1)	16(1)	13(1)	1(1)	3(1)	0(1)
C(24)	18(1)	20(1)	17(1)	1(1)	4(1)	-5(1)
S(1)	18(1)	18(1)	31(1)	-1(1)	7(1)	-3(1)
O(2)	37(1)	20(1)	62(1)	-4(1)	12(1)	-13(1)
C(25)	22(1)	22(1)	30(1)	2(1)	3(1)	1(1)
C(26)	43(1)	37(1)	26(1)	1(1)	6(1)	-13(1)

Crystalline form I**[0434]**

Table 35: Crystal data and structure refinement for crystalline form I.

Empirical formula	C ₂₄ H ₂₇ FN ₂ O
Formula weight	378.48
Temperature	100(2) K
Wavelength	0.71073 Å
Crystal system	Monoclinic
Space group	P 21/c
Unit cell dimensions	a = 10.809(3) Å alpha = 90 deg.
	b = 15.946(5) Å beta = 107.632(7) deg.
	c = 11.846(4) Å gamma = 90 deg.
Volume	1945.9(10) Å ³

Z	4
Density (calculated)	1.292 Mg/m ³
Absorption coefficient	0.086 mm ⁻¹
F(000)	808
Crystal size	0.14 x 0.11 x 0.09 mm
Theta range for data collection	2.21 to 25.65 deg.
Index ranges	-13 ≤ h ≤ 12, -19 ≤ k ≤ 16, -14 ≤ l ≤ 14
Reflections collected	16947
Independent reflections	3673 [R(int) = 0.1071]
Absorption correction	None
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	3673 / 0 / 259
Goodness-of-fit on F ²	1.094
Final R indices [I > 2σ(I)]	R1 = 0.0645, wR2 = 0.1618
R indices (all data)	R1 = 0.0931, wR2 = 0.1779
Largest diff. peak and hole	0.382 and -0.320 e.Å ⁻³

Table 36: Atomic coordinates ($\times 10^4$) (i.e. ($\times 10^4$)) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) (i.e. ($\text{\AA}^2 \times 10^3$)) for crystalline form I. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U_{ij} tensor.

	x	y	z	U(eq)
F(1)	7997(1)	4716(1)	2730(1)	32(1)
O(1)	2425(2)	2871(1)	4710(2)	22(1)
N(1)	3434(2)	3016(1)	1980(2)	20(1)
N(2)	-600(2)	1697(1)	1517(2)	22(1)
C(1)	6852(2)	4284(2)	2510(2)	22(1)
C(2)	6307(2)	4202(2)	3405(2)	22(1)
C(3)	5135(2)	3753(2)	3149(2)	19(1)
C(4)	4584(2)	3413(2)	2000(2)	19(1)
C(5)	3269(2)	3107(2)	3091(2)	20(1)
C(6)	2143(2)	2746(2)	3440(2)	20(1)
C(7)	2957(2)	3679(2)	5144(2)	23(1)
C(8)	4324(2)	3767(2)	5062(2)	24(1)
C(9)	4276(2)	3545(2)	3822(2)	19(1)
C(10)	838(2)	3164(2)	2778(2)	21(1)
C(11)	-292(2)	2751(2)	3111(2)	21(1)

	x	y	z	U(eq)
C(12)	-431(2)	1809(2)	2809(2)	21(1)
C(13)	-1705(2)	2152(2)	737(2)	26(1)
C(14)	-662(3)	828(2)	1124(2)	30(1)
C(15)	875(2)	1416(2)	3535(2)	21(1)
C(16)	2035(2)	1801(2)	3242(2)	22(1)
C(17)	-1580(2)	1410(2)	3141(2)	21(1)
C(18)	-2738(2)	1849(2)	3002(2)	22(1)
C(19)	-3821(2)	1465(2)	3180(2)	23(1)
C(20)	-3778(2)	625(2)	3484(2)	25(1)
C(21)	-2633(2)	184(2)	3653(3)	31(1)
C(22)	-1553(2)	569(2)	3487(2)	27(1)
C(23)	5176(2)	3502(2)	1107(2)	22(1)
C(24)	6333(2)	3948(2)	1377(2)	24(1)

Table 37A: Bond lengths [Å] and angles [deg] for crystalline form I.

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
F(1)-C(1)	1.370(3)	C(11)-H(11A)	0.9900
O(1)-C(7)	1.441(3)	C(11)-H(11B)	0.9900
O(1)-C(6)	1.456(3)	C(12)-C(15)	1.547(3)
N(1)-C(5)	1.388(3)	C(12)-C(17)	1.548(3)
N(1)-C(4)	1.389(3)	C(13)-H(13A)	0.9800
N(1)-H(1)	0.97(4)	C(13)-H(13B)	0.9800
N(2)-C(14)	1.458(3)	C(13)-H(13C)	0.9800
N(2)-C(13)	1.462(3)	C(14)-H(14A)	0.9800
N(2)-C(12)	1.496(3)	C(14)-H(14B)	0.9800
C(1)-C(2)	1.366(4)	C(14)-H(14C)	0.9800
C(1)-C(24)	1.396(4)	C(15)-C(16)	1.528(3)
C(2)-C(3)	1.405(3)	C(15)-H(15A)	0.9900
C(2)-H(2)	0.9500	C(15)-	0.9900

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
		H(15B)	
C(3)-C(4)	1.419(4)	C(16)-H(16A)	0.9900
C(3)-C(9)	1.434(3)	C(16)-H(16B)	0.9900
C(4)-C(23)	1.400(4)	C(17)-C(18)	1.399(3)
C(5)-C(9)	1.361(3)	C(17)-C(22)	1.400(4)
C(5)-C(6)	1.512(3)	C(18)-C(19)	1.392(3)
C(6)-C(16)	1.523(4)	C(18)-H(18)	0.9500
C(6)-C(10)	1.543(3)	C(19)-C(20)	1.385(4)
C(7)-C(8)	1.517(3)	C(19)-H(19)	0.9500
C(7)-H(7A)	0.9900	C(20)-C(21)	1.384(4)
C(7)-H(7B)	0.9900	C(20)-H(20)	0.9500
C(8)-C(9)	1.496(4)	C(21)-C(22)	1.385(4)
C(8)-H(8A)	0.9900	C(21)-H(21)	0.9500
C(8)-H(8B)	0.9900	C(22)-H(22)	0.9500
C(10)-C(11)	1.541(3)	C(23)-C(24)	1.388(4)
C(10)-H(10A)	0.9900	C(23)-H(23)	0.9500
C(10)-H(10B)	0.9900	C(24)-H(24)	0.9500
C(11)-C(12)	1.540(4)		

Table 37B: (Table 37A continued) Bond lengths [Å] and angles [deg] for crystalline form I.

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(7)-O(1)-C(6)	115.28(18)	N(2)-C(12)-C(11)	109.1(2)
C(5)-N(1)-C(4)	107.9(2)	N(2)-C(12)-C(15)	109.2(2)
C(5)-N(1)-H(1)	133(2)	C(11)-C(12)-C(15)	104.96(19)
C(4)-N(1)-H(1)	119(2)	N(2)-C(12)-C(17)	110.60(19)
C(14)-N(2)-C(13)	108.7(2)	C(11)-C(12)-C(17)	111.8(2)
C(14)-N(2)-C(12)	114.8(2)	C(15)-C(12)-C(17)	111.0(2)

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(13)-N(2)- C(12)	114.3(2)	N(2)-C(13)- H(13A)	109.5
C(2)-C(1)-F(1)	118.5(2)	N(2)-C(13)- H(13B)	109.5
C(2)-C(1)-C(24)	124.5(2)	H(13A)-C(13)- H(13B)	109.5
F(1)-C(1)-C(24)	116.9(2)	N(2)-C(13)- H(13C)	109.5
C(1)-C(2)-C(3)	117.0(2)	H(13A)-C(13)- H(13C)	109.5
C(1)-C(2)-H(2)	121.5	H(13B)-C(13)- H(13C)	109.5
C(3)-C(2)-H(2)	121.5	N(2)-C(14)- H(14A)	109.5
C(2)-C(3)-C(4)	119.4(2)	N(2)-C(14)- H(14B)	109.5
C(2)-C(3)-C(9)	133.7(2)	H(14A)-C(14)- H(14B)	109.5
C(4)-C(3)-C(9)	106.9(2)	N(2)-C(14)- H(14C)	109.5
N(1)-C(4)-C(23)	130.2(2)	H(14A)-C(14)- H(14C)	109.5
N(1)-C(4)-C(3)	107.7(2)	H(14B)-C(14)- H(14C)	109.5
C(23)-C(4)-C(3)	122.1(2)	C(16)-C(15)- C(12)	112.6(2)
C(9)-C(5)-N(1)	110.4(2)	C(16)-C(15)- H(15A)	109.1
C(9)-C(5)-C(6)	125.2(2)	C(12)-C(15)- H(15A)	109.1
N(1)-C(5)-C(6)	124.4(2)	C(16)-C(15)- H(15B)	109.1
O(1)-C(6)-C(5)	107.26(18)	C(12)-C(15)- H(15B)	109.1
O(1)-C(6)-C(16)	106.06(19)	H(15A)-C(15)- H(15B)	107.8
C(5)-C(6)-C(16)	111.4(2)	C(6)-C(16)- C(15)	113.2(2)
O(1)-C(6)-C(10)	109.80(19)	C(6)-C(16)- H(16A)	108.9

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(5)-C(6)-C(10)	112.6(2)	C(15)-C(16)- H(16A)	108.9
C(16)-C(6)- C(10)	109.50(19)	C(6)-C(16)- H(16B)	108.9
O(1)-C(7)-C(8)	110.5(2)	C(15)-C(16)- H(16B)	108.9
O(1)-C(7)-H(7A)	109.5	H(16A)-C(16)- H(16B)	107.8
C(8)-C(7)-H(7A)	109.5	C(18)-C(17)- C(22)	116.9(2)
O(1)-C(7)-H(7B)	109.5	C(18)-C(17)- C(12)	121.4(2)
C(8)-C(7)-H(7B)	109.5	C(22)-C(17)- C(12)	121.4(2)
H(7A)-C(7)- H(7B)	108.1	C(19)-C(18)- C(17)	121.7(2)
C(9)-C(8)-C(7)	107.2(2)	C(19)-C(18)- H(18)	119.2
C(9)-C(8)-H(8A)	110.3	C(17)-C(18)- H(18)	119.2
C(7)-C(8)-H(8A)	110.3	C(20)-C(19)- C(18)	120.1(2)
C(9)-C(8)-H(8B)	110.3	C(20)-C(19)- H(19)	119.9
C(7)-C(8)-H(8B)	110.3	C(18)-C(19)- H(19)	119.9
H(8A)-C(8)- H(8B)	108.5	C(21)-C(20)- C(19)	119.1(2)
C(5)-C(9)-C(3)	107.0(2)	C(21)-C(20)- H(20)	120.4
C(5)-C(9)-C(8)	122.0(2)	C(19)-C(20)- H(20)	120.4
C(3)-C(9)-C(8)	130.9(2)	C(20)-C(21)- C(22)	120.7(3)
C(11)-C(10)- C(6)	111.0(2)	C(20)-C(21)- H(21)	119.7
C(11)-C(10)- H(10A)	109.4	C(22)-C(21)- H(21)	119.7
C(6)-C(10)- H(10A)	109.4	C(21)-C(22)- C(17)	121.5(2)

	bond lengths [Å] and angles [deg]		bond lengths [Å] and angles [deg]
C(11)-C(10)-H(10B)	109.4	C(21)-C(22)-H(22)	119.3
C(6)-C(10)-H(10B)	109.4	C(17)-C(22)-H(22)	119.3
H(10A)-C(10)-H(10B)	108.0	C(24)-C(23)-C(4)	117.6(2)
C(12)-C(11)-C(10)	112.8(2)	C(24)-C(23)-H(23)	121.2
C(12)-C(11)-H(11A)	109.0	C(4)-C(23)-H(23)	121.2
C(10)-C(11)-H(11A)	109.0	C(23)-C(24)-C(1)	119.4(2)
C(12)-C(11)-H(11B)	109.0	C(23)-C(24)-H(24)	120.3
C(10)-C(11)-H(11B)	109.0	C(1)-C(24)-H(24)	120.3
H(11A)-C(11)-H(11B)	107.8		

[0435] Symmetry transformations used to generate equivalent atoms:

Table 38: Hydrogen coordinates ($\times 10^4$) (i.e. ($\times 10^4$)) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) (i.e. ($\text{\AA}^2 \times 10^3$)) for crystalline form I.

	x	y	z	U(eq)
H(1)	2900(30)	2770(20)	1240(30)	51(10)
H(2)	6703	4437	4166	26
H(7A)	2970	3745	5978	28
H(7B)	2400	4127	4671	28
H(8A)	4637	4349	5247	28
H(8B)	4921	3385	5632	28
H(10A)	877	3768	2980	25
H(10B)	684	3115	1913	25
H(11A)	-150	2823	3971	25
H(11B)	-1111	3040	2685	25
H(13A)	-1764	2037	-91	39
H(13B)	-1586	2755	890	39
H(13C)	-2506	1968	888	39

	x	y	z	U(eq)
H(14A)	-1471	573	1168	44
H(14B)	79	517	1635	44
H(14C)	-637	809	305	44
H(15A)	988	1493	4390	25
H(15B)	852	805	3375	25
H(16A)	2841	1532	3740	26
H(16B)	1958	1681	2404	26
H(18)	-2786	2424	2780	26
H(19)	-4590	1780	3092	28
H(20)	-4525	354	3576	30
H(21)	-2588	-388	3885	37
H(22)	-776	255	3610	32
H(23)	4801	3265	343	26
H(24)	16766	14023	796	29

Table 39: Anisotropic displacement parameters ($^2 \times 10^3$) (i.e. ($^2 \times 10^3$)) for crystalline form I. The anisotropic displacement factor exponent takes the form: $-2 \pi^2 [h^2 a^{*2} U_{11} + \dots + 2hka^*b^*U_{12}]$.

	U11	U22	U33	U23	U13	U12
F(1)	26(1)	35(1)	35(1)	2(1)	11(1)	-8(1)
O(1)	30(1)	21(1)	15(1)	0(1)	8(1)	-4(1)
N(1)	21(1)	23(1)	16(1)	0(1)	7(1)	0(1)
N(2)	22(1)	26(1)	18(1)	-1(1)	6(1)	2(1)
C(1)	18(1)	21(1)	26(1)	4(1)	6(1)	0(1)
C(2)	23(1)	20(1)	23(1)	2(1)	6(1)	0(1)
C(3)	23(1)	14(1)	20(1)	2(1)	7(1)	3(1)
C(4)	20(1)	19(1)	19(1)	2(1)	6(1)	3(1)
C(5)	21(1)	19(1)	20(1)	0(1)	8(1)	3(1)
C(6)	24(1)	21(1)	15(1)	-1(1)	7(1)	0(1)
C(7)	33(1)	21(2)	19(1)	-3(1)	13(1)	-3(1)
C(8)	30(1)	24(2)	17(1)	-1(1)	7(1)	-4(1)
C(9)	23(1)	18(1)	18(1)	0(1)	8(1)	2(1)
C(10)	24(1)	18(1)	21(1)	3(1)	7(1)	0(1)
C(11)	22(1)	22(2)	21(1)	2(1)	8(1)	0(1)
C(12)	25(1)	21(2)	16(1)	0(1)	7(1)	0(1)
C(13)	27(1)	30(2)	22(2)	4(1)	8(1)	2(1)

	U11	U22	U33	U23	U13	U12
C(14)	31(1)	30(2)	24(2)	-6(1)	3(1)	9(1)
C(15)	24(1)	19(1)	20(1)	-1(1)	7(1)	0(1)
C(16)	22(1)	22(2)	22(1)	1(1)	8(1)	1(1)
C(17)	24(1)	21(1)	18(1)	-3(1)	7(1)	-4(1)
C(18)	29(1)	22(2)	14(1)	-2(1)	8(1)	-2(1)
C(19)	24(1)	28(2)	18(1)	-3(1)	7(1)	-2(1)
C(20)	28(1)	29(2)	19(1)	-3(1)	9(1)	-9(1)
C(21)	34(1)	24(2)	34(2)	2(1)	12(1)	-4(1)
C(22)	27(1)	22(2)	33(2)	0(1)	10(1)	0(1)
C(23)	25(1)	22(1)	18(1)	-1(1)	6(1)	3(1)
C(24)	27(1)	26(2)	23(1)	5(1)	12(1)	3(1)

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2004043967A](#) [0003]
- [WO2006108565A](#) [0004]
- [WO2008009415A](#) [0005]
- [WO2008040481A](#) [0324]

Non-patent literature cited in the description

- R. HILFIKER Polymorphism Wiley VCH20060000235-242 [0007]
- A.R. GENNARO Remington's Pharmaceutical Sciences Mack Publishing Company 19850000 [0335]

Patentkrav

1. Krystallinsk form af (1r,4r)-6'-fluor-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro-[cyclohexan-1,1'-pyran-[3,4,b]indol]-4-amin, som er et alkoholsolvat.
2. Den krystallinske form ifølge krav 1, som er form C, som har røntgendiffraktionstoppe (CuK α -stråling) ved 9,1 $\pm$ 0,2 (2 θ), 9,5 $\pm$ 0,2 (2 θ), 16,8 $\pm$ 0,2 (2 θ), 18,2 $\pm$ 0,2 (2 θ), 18,6 $\pm$ 0,2 (2 θ), 19,0 $\pm$ 0,2 (2 θ), 19,3 $\pm$ 0,2 (2 θ), 19,5 $\pm$ 0,2 (2 θ), 22,2 $\pm$ 0,2 (2 θ), 25,4 $\pm$ 0,2 (2 θ) og 27,5 $\pm$ 0,2 (2 θ).
3. Den krystallinske form C ifølge krav 2, som har Raman-toppe ved 156 $\pm$ 2 cm⁻¹, 171 $\pm$ 2 cm⁻¹, 183 $\pm$ 2 cm⁻¹, 922 $\pm$ 2 cm⁻¹, 1003 $\pm$ 2 cm⁻¹, 1299 $\pm$ 2 cm⁻¹, 1478 $\pm$ 2 cm⁻¹, 1570 $\pm$ 2 cm⁻¹, 1587 $\pm$ 2 cm⁻¹, 2932 $\pm$ 2 cm⁻¹, 2951 $\pm$ 2 cm⁻¹, 3070 $\pm$ 2 cm⁻¹.
4. Den krystallinske form ifølge krav 1, som er form D, som har røntgendiffraktionstoppe (CuK α -stråling) ved 8,4 $\pm$ 0,2 (2 θ), 8,8 $\pm$ 0,2 (2 θ), 15,0 $\pm$ 0,2 (2 θ), 15,2 $\pm$ 0,2 (2 θ), 17,0 $\pm$ 0,2 (2 θ), 17,6 $\pm$ 0,2 (2 θ), 18,9 $\pm$ 0,2 (2 θ), 21,2 $\pm$ 0,2 (2 θ), 22,4 $\pm$ 0,2 (2 θ), 23,2 $\pm$ 0,2 (2 θ), 26,0 $\pm$ 0,2 (2 θ), 29,5 $\pm$ 0,2 (2 θ) og 30,7 $\pm$ 0,2 (2 θ).
5. Den krystallinske form D ifølge krav 4, som har Raman-toppe ved 169 $\pm$ 2 cm⁻¹, 254 $\pm$ 2 cm⁻¹, 367 $\pm$ 2 cm⁻¹, 491 $\pm$ 2 cm⁻¹, 683 $\pm$ 2 cm⁻¹, 922 $\pm$ 2 cm⁻¹, 1002 $\pm$ 2 cm⁻¹, 1302 $\pm$ 2 cm⁻¹, 1437 $\pm$ 2 cm⁻¹, 1479 $\pm$ 2 cm⁻¹, 1570 $\pm$ 2 cm⁻¹, 2935 $\pm$ 2 cm⁻¹, 2957 $\pm$ 2 cm⁻¹, 3067 $\pm$ 2 cm⁻¹.
6. Den krystallinske form ifølge krav 1, som er form E, som har røntgendiffraktionstoppe (CuK α -stråling) ved 8,8 $\pm$ 0,2 (2 θ), 11,9 $\pm$ 0,2 (2 θ), 17,0 $\pm$ 0,2 (2 θ), 17,7 $\pm$ 0,2 (2 θ) og 18,7 $\pm$ 0,2 (2 θ).
7. Den krystallinske form E ifølge krav 6, som har Raman-toppe ved 159 $\pm$ 2 cm⁻¹, 188 $\pm$ 2 cm⁻¹, 680 $\pm$ 2 cm⁻¹, 923 $\pm$ 2 cm⁻¹, 1003 $\pm$ 2 cm⁻¹, 1297 $\pm$ 2 cm⁻¹, 1434 $\pm$ 2 cm⁻¹, 1461 $\pm$ 2 cm⁻¹, 1570 $\pm$ 2 cm⁻¹, 1585 $\pm$ 2 cm⁻¹, 2943 $\pm$ 2 cm⁻¹, 2961 $\pm$ 2 cm⁻¹, 3070 $\pm$ 2 cm⁻¹.
8. Den krystallinske form ifølge krav 1, som er form F, som har røntgendiffraktionstoppe (CuK α -stråling) ved 9,0 $\pm$ 0,2 (2 θ), 15,4 $\pm$ 0,2 (2 θ), 16,1 $\pm$ 0,2 (2 θ), 17,9 $\pm$ 0,2 (2 θ), 18,2 $\pm$ 0,2

(2 θ), 18,7 $\pm$ 0,2 (2 θ), 19,4 $\pm$ 0,2 (2 θ), 20,1 $\pm$ 0,2 (2 θ), 20,6 $\pm$ 0,2 (2 θ), 21,8 $\pm$ 0,2 (2 θ), 24,6 $\pm$ 0,2 (2 θ), 25,6 $\pm$ 0,2 (2 θ), 27,1 $\pm$ 0,2 (2 θ), 27,4 $\pm$ 0,2 (2 θ), og 29,3 $\pm$ 0,2 (2 θ).

9. Den krystallinske form F ifølge krav 8, som har Raman-toppe ved 157 $\pm$ 2 cm⁻¹, 171 $\pm$ 2 cm⁻¹, 183 $\pm$ 2 cm⁻¹, 682 $\pm$ 2 cm⁻¹, 921 $\pm$ 2 cm⁻¹, 1002 $\pm$ 2 cm⁻¹, 1299 $\pm$ 2 cm⁻¹, 1463 $\pm$ 2 cm⁻¹, 1477 $\pm$ 2 cm⁻¹, 1570 $\pm$ 2 cm⁻¹, 1581 $\pm$ 2 cm⁻¹, 2889 $\pm$ 2 cm⁻¹, 2932 $\pm$ 2 cm⁻¹, 2952 $\pm$ 2 cm⁻¹, 2977 $\pm$ 2 cm⁻¹, 3058 $\pm$ 2 cm⁻¹, 3070 $\pm$ 2 cm⁻¹.

10. Farmaceutisk sammensætning omfattende mindst en krystallinsk form ifølge et hvilket som helst af kravene 1 til 9.

11. Fremgangsmåde til fremstilling eller til oprensning af (1r,4r)-6'-fluor-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexan-1,1'-pyran[3,4b]indol]-4-amin, den frie base eller et fysiologisk acceptabelt salt deraf, hvilken fremgangsmåde omfatter trinnene

(i) tilvejebringelse af en blanding af diastomerer omfattende (1r,4r)-6'-fluor-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexan-1,1'-pyran[3,4b]indol]-4-amin og (1s,4s)-6'-fluor-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexan-1,1'-pyran[3,4,b]indol]-4-amin; fortrinsvis med et overskud af (1r,4r)-6'-fluor-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexan-1,1'-pyran[3,4,b]indol]-4-amin; og

(ii). anvendelse af blandingen af diastomerer opnået i trin (i) som udgangsmateriale og fremstilling derfra af en krystallinsk form af (1r,4r)-6'-fluor-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexan-1,1'-pyran[3,4,b]indol]-4-amin valgt fra gruppen bestående af krystallinske former C, D, E og F ifølge et hvilket som helst af kravene 2 til 9, hvori trin (ii) involverer undertrinnene med

(ii-a) opløsning af blandingen af diastomerer i en solvent eller solventblanding omfattende dimethylsulfoxid eller en anden dipolær aprotisk solvent;

(ii-b) tilsætning af en alkohol valgt fra gruppen bestående af methanol, ethanol, 1-propanol og 2-propanol;

(ii-c) udfældning af form C, D, E eller F; og

(ii-d) separering af præcipitatet opnået i trin (ii-c) fra den resterende opløsning indeholdende (1s,4s)-6'-fluor-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexan-1,1'-pyran[3,4,b]indol]-4-amin; og

(iii) separering af den krystallinske form af (1r,4r)-6'-fluor-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexan-1,1'-pyran[3,4,b]indol]-4-amin opnået i trin (ii) fra den resterende blanding af diastomerer; fortrinsvis ved udfældning og filtrering; og

(iv) valgfrit, konvertering af den krystallinske form af (1r,4r)-6'-fluor-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexan-1,1'-pyran[3,4,b]indol]-4-amin til en anden krystallinsk form af (1r,4r)-6'-fluor-N,N-dimethyl-4-phenyl-4',9'-dihydro-3'H-spiro[cyclohexan-1,1'-pyran[3,4,b]indol]-4-amin opnået i trin (iii), fortrinsvis ved tørring, fortrinsvis til krystallinsk form A, som har røntgendiffraktionstoppe (CuK α -stråling) ved $17,6\pm0,2$ (2θ), $18,3\pm0,2$ (2θ), $18,6\pm0,2$ (2θ), $25,8\pm0,2$ (2θ) og $26,3\pm0,2$ (2θ) og/eller Raman-toppe ved 921 ± 2 cm $^{-1}$, 1002 ± 2 cm $^{-1}$, 1308 ± 2 cm $^{-1}$, 1569 ± 2 cm $^{-1}$, 1583 ± 2 cm $^{-1}$, 3057 ± 2 cm $^{-1}$.

DRAWINGS

Figure 1a




Figure 1b




Figure 1c




Figure 1d




Figure 1e




Figure 1f




Figure 1g




Figure 1h




Figure 1i




Figure 1I




Figure 2a




Figure 2b




Figure 2c




Figure 2d




Figure 2e




Figure 2f




Figure 2g




Figure 2h




Figure 2i




Figure 2I

