

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 200120731 B2
(10) Patent No. 785148

(54) Title
Use of recombinant parainfluenza viruses (PIVs) as vectors to protect against infection and disease caused by PIV and other human pathogens

(51)⁶ International Patent Classification(s)
C12N 15/86 060408BMEP
(2006.01) A61K 39/00
A61K 39/00 20060101ALI20
(2006.01) 060408BMEP
C07K 14/115 C07K 14/115
(2006.01) 20060101ALI20
C12N 15/86 060408BMEP
20060101AFI20

(21) Application No: 200120731 (22) Application Date: 2000.12.08

(87) WIPO No: W001/42445

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/170,195	1999.12.10	US
09/458,813	1999.12.10	US
09/459,062	1999.12.10	US

(43) Publication Date : 2001.06.18

(43) Publication Journal Date : 2001.08.30

(44) Accepted Journal Date : 2006.10.05

(71) Applicant(s)
The Government of the United States of America, as represented by the Department of Health and Human Services

(72) Inventor(s)
Brian R Murphy; Peter L. Collins; Alexander C. Schmidt;
Anna P. Durbin; Mario H. Skiadopoulos; Tao Tao

(74) Agent/Attorney
PHILLIPS ORMONDE and FITZPATRICK, 367 Collins Street, MELBOURNE
VIC 3000

(56) Related Art
WO 1998053078
SKIADOPOULOS, M ETAL (1999) VACCINE 18(5-6): 503-510

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 June 2001 (14.06.2001)


PCT

(10) International Publication Number
WO 01/42445 A2

- (51) International Patent Classification: C12N 15/00
- (21) International Application Number: PCT/US00/33293
- (22) International Filing Date: 8 December 2000 (08.12.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/170,195 10 December 1999 (10.12.1999) US
09/458,813 10 December 1999 (10.12.1999) US
09/459,062 10 December 1999 (10.12.1999) US
- (71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as represented by THE DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; 6011 Executive Boulevard, Suite 325, Rockville, MD 20852 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): MURPHY, Brian,
- R, [US/US]; 5410 Tuscarawas Road, Bethesda, MD 20816 (US). COLLINS, Peter, L. [US/US]; 12304 Village Square, #401, Rockville, MD 20852 (US). SCHMIDT, Alexander, C. [US/US]; 1831 Belmont Road #203 NW, Washington, DC 20009 (US). DURBIN, Anna, P. [US/US]; 806 Hudson Avenue, Takoma Park, MD 20912 (US). SKIADOPOULOS, Mario, H. [US/US]; 8303 Aqueduct Road, Potomac, MD 20854 (US). TAO, Tao [US/US]; 10684 Weymouth Street, #104, Bethesda, MD 20814 (US).
- (74) Agents: KING, Jeffrey, J. et al.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, 8th Floor, San Francisco, CA 94111 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

[Continued on next page]

(54) Title: USE OF RECOMBINANT PARAINFLUENZA VIRUSES (PIVs) AS VECTORS TO PROTECT AGAINST INFECTION AND DISEASE CAUSED BY PIV AND OTHER HUMAN PATHOGENS



WO 01/42445 A2

(57) Abstract: Chimeric parainfluenza viruses (PIVs) are provided that incorporate a PIV vector genome or antigenome and one or more antigenic determinant(s) of a heterologous PIV or non-PIV pathogen. These chimeric viruses are infectious and attenuated in humans and other mammals and are useful in vaccine formulations for eliciting an immune responses against one or more PIVs, or against a PIV and non-PIV pathogen. Also provided are isolated polynucleotide molecules and vectors incorporating a chimeric PIV genome or antigenome which includes a partial or complete PIV vector genome or antigenome combined or integrated with one or more heterologous gene(s) or genome segment(s) encoding antigenic determinant(s) of a heterologous PIV or non-PIV pathogen. In preferred aspects of the invention, chimeric PIV incorporate a partial or complete human, bovine, or human-bovine chimeric, PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a heterologous PIV or non-PIV pathogen, wherein the chimeric virus is attenuated for use as a vaccine agent by any of a variety of mutations and nucleotide modifications introduced into the chimeric genome or antigenome.



(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

— *With (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description.*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *Without international search report and to be republished upon receipt of that report.*

**USE OF RECOMBINANT PARAINFLUENZA VIRUSES (PIVs) AS
VECTORS TO PROTECT AGAINST INFECTION AND DISEASE
CAUSED BY PIV AND OTHER HUMAN PATHOGENS**

5 BACKGROUND OF THE INVENTION

Human parainfluenza virus type 3 (HPIV3) is a common cause of serious lower respiratory tract infection in infants and children less than one year of age. It is second only to respiratory syncytial virus (RSV) as a leading cause of hospitalization for viral lower respiratory tract disease in this age group (Collins et al., p. 1205-1243. In B. N. Fields (Knipe et al., eds.), *Fields Virology*, 3rd ed., vol. 1. Lippincott-Raven Publishers, Philadelphia, 1996; Crowe et al., *Vaccine* 13:415-421, 1995; Marx et al., *J. Infect. Dis.* 176:1423-1427, 1997). Infections by this virus results in substantial morbidity in children less than 3 years of age. HPIV1 and HPIV2 are the principal etiologic agents of laryngotracheobronchitis (croup) and also can cause severe pneumonia and bronchiolitis (Collins et al., 3rd ed. In "*Fields Virology*," B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, B. Roizman, and S. E. Straus, Eds., Vol. 1, pp. 1205-1243. Lippincott-Raven Publishers, Philadelphia, 1996). In a long term study over a 20-year period, HPIV1, HPIV2, and HPIV3 were identified as etiologic agents for 6.0, 3.2, and 11.5%, respectively, of hospitalizations for respiratory tract disease accounting in total for 18% of the hospitalizations, and, for this reason, there is a need for an effective vaccine (Murphy et al., *Virus Res.* 11:1-15, 1988). The parainfluenza viruses have also been identified in a significant proportion of cases of virally-induced middle ear effusions in children with otitis media (Heikkinen et al., *N. Engl. J. Med.* 340:260-4, 1999). Thus, there is a need to produce a vaccine against these viruses that can prevent the serious lower respiratory tract disease and the otitis media that accompanies these HPIV infections. HPIV1, HPIV2, and HPIV3 are distinct serotypes which do not elicit significant cross-protective immunity.

Despite considerable efforts to develop effective vaccine therapies against HPIV, no approved vaccine agents have yet been achieved for any HPIV serotype, nor for ameliorating HPIV related illnesses. To date, only two live attenuated PIV vaccine candidates have received particular attention. One of these candidates is a bovine PIV (BPIV3) strain that is antigenically related to HPIV3 and which has been shown to protect animals against HPIV3. BPIV3 is attenuated, genetically stable and immunogenic in human

infants and children (Karron et al., J. Inf. Dis. 171:1107-14, 1995a; Karron et al., J. Inf. Dis. 172:1445-1450, 1995b). A second PIV3 vaccine candidate, JS *cp45*, is a cold-adapted mutant of the JS wildtype (*wt*) strain of HPIV3 (Karron et al., J. Inf. Dis. 172:1445-1450, 1995b; Belshe et al., J. Med. Virol. 10:235-42, 1982). This live, attenuated, cold-passaged (5 *cp*) PIV3 vaccine candidate exhibits temperature-sensitive (*ts*), cold-adaptation (*ca*), and attenuation (*att*) phenotypes which are stable after viral replication *in vivo*. The *cp45* virus is protective against human PIV3 challenge in experimental animals and is attenuated, genetically stable, and immunogenic in seronegative human infants and children (Hall et al., Virus Res. 22:173-184, 1992; Karron et al., J. Inf. Dis. 172:1445-1450, 1995b). The most promising prospects to date are live attenuated vaccine viruses since these have been shown to be efficacious in non-human primates even in the presence of passively transferred antibodies, an experimental situation that simulates that present in the very young infant who possesses maternally acquired antibodies (Crowe et al., Vaccine 13:847-855, 1995; Durbin et al., J. Infect. Dis. 179:1345-1351, 1999). Two live attenuated PIV3 vaccine candidates, a temperature-sensitive (*ts*) derivative of the wild type PIV3 JS strain (designated PIV3*cp45*) and a bovine PIV3 (BPIV3) strain, are undergoing clinical evaluation (Karron et al., Pediatr. Infect. Dis. J. 15:650-654, 1996; Karron et al., J. Infect. Dis. 171:1107-1114, 1995a; Karron et al., J. Infect. Dis. 172, 1445-1450, 1995b). The live attenuated PIV3*cp45* vaccine candidate was derived from the JS strain of HPIV3 via serial passage in cell culture at low temperature and has been found to be protective against HPIV3 challenge in experimental animals and to be satisfactorily attenuated, genetically stable, and immunogenic in seronegative human infants and children (Belshe et al., J. Med. Virol. 10:235-242, 1982; Belshe et al., Infect. Immun. 37:160-5, 1982; Clements et al., J. Clin. Microbiol. 29:1175-82, 1991; Crookshanks et al., J. Med. Virol. 13:243-9, 1984; Hall et al., Virus Res. 22:173-184, 1992; Karron et al., J. Infect. Dis. 172:1445-1450, 1995b). Because these PIV3 candidate vaccine viruses are biologically derived, there is no proven methods for adjusting the level of attenuation should this be found necessary from ongoing clinical trials.

To facilitate development of PIV vaccine candidates, recombinant DNA technology has recently made it possible to recover infectious negative-stranded RNA viruses from cDNA (for reviews, see Conzelmann, J. Gen. Virol. 77:381-89, 1996; Palese et al., Proc. Natl. Acad. Sci. U.S.A. 93:11354-58, 1996). In this context, recombinant rescue has been reported for infectious respiratory syncytial virus (RSV), rabies virus (RaV), simian virus 5 (SV5), rinderpest virus, Newcastle disease virus (NDV), vesicular stomatitis virus (VSV), measles virus (MeV), and Sendai virus (SeV) from cDNA-encoded antigenomic

RNA in the presence of essential viral proteins (see, e.g., Garcin et al., EMBO J. 14:6087-6094, 1995; Lawson et al., Proc. Natl. Acad. Sci. U.S.A. 92:4477-81, 1995; Radecke et al., EMBO J. 14:5773-5784, 1995; Schnell et al., EMBO J. 13:4195-203, 1994; Whelan et al., Proc. Natl. Acad. Sci. U.S.A. 92:8388-92, 1995; Hoffman et al., J. Virol. 71:4272-4277, 5 1997; Kato et al., Genes to Cells 1:569-579, 1996; Roberts et al., Virology 247:1-6, 1998; Baron et al., J. Virol. 71:1265-1271, 1997; International Publication No. WO 97/06270; Collins et al., Proc. Natl. Acad. Sci. U.S.A. 92:11563-11567, 1995; U.S. Patent Application No. 08/892,403, filed July 15, 1997 (corresponding to published International Application No. WO 98/02530 and priority U.S. Provisional Application Nos. 60/047,634, filed May 23, 10 1997, 60/046,141, filed May 9, 1997, and 60/021,773, filed July 15, 1996); U.S. Patent Application No. 09/291,894, filed on April 13, 1999; International Application No. PCT/US00/09695, filed April 12, 2000 (which claims priority to U.S. Provisional Patent Application Serial No. 60/129,006, filed April 13, 1999); International Application No. PCT/US00/17755, filed June 23, 2000 (which claims priority to U.S. Provisional Patent 15 Application Serial No. 60/143,132, filed by Bucholz et al. on July 9, 1999); Juhasz et al., J. Virol. 71:5814-5819, 1997; He et al. Virology 237:249-260, 1997; Peters et al. J. Virol. 73:5001-5009, 1999; Baron et al. J. Virol. 71:1265-1271, 1997; Whitehead et al., Virology 247:232-9, 1998a; Whitehead et al., J. Virol. 72:4467-4471, 1998b; Jin et al. Virology 251:206-214, 1998; Bucholz et al. J. Virol. 73:251-259, 1999; and Whitehead et al., J. Virol. 20 73:3438-3442, 1999, each incorporated herein by reference in its entirety for all purposes).

In more specific regard to the instant invention, a method for producing HPIV with a wt phenotype from cDNA was recently developed for recovery of infectious, recombinant HPIV3 JS strain (see, e.g., Durbin et al., Virology 235:323-332, 1997; U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998 (corresponding to U.S. 25 Provisional Application No. 60/059,385, filed September 19, 1997); and U.S. Provisional Application No. 60/047,575, filed May 23, 1997 (corresponding to International Publication No. WO 98/53078), each incorporated herein by reference). In addition, these disclosures allow for genetic manipulation of viral cDNA clones to determine the genetic basis of phenotypic changes in biological mutants, e.g., which mutations in the HPIV3 *cp45* virus 30 specify its *ts*, *ca* and *att* phenotypes, and which gene(s) or genome segment(s) of BPIV3 specify its attenuation phenotype. Additionally, these and related disclosures render it feasible to construct novel PIV vaccine candidates having a wide range of different mutations and to evaluate their level of attenuation, immunogenicity and phenotypic stability (see also, U.S. Application No. 09/586,479, filed June 1, 2000 (corresponding to U.S.

Provisional Patent Application Serial No. 60/143,134, filed on July 9, 1999); and U.S. Patent Application Serial No. 09/350,821, filed by Durbin et al. on July 9, 1999, each incorporated herein by reference).

Thus, infectious wild type recombinant PIV3, (r)PIV3, as well as a number of
5 *ts* derivatives, have now been recovered from cDNA, and reverse genetics systems have
been used to generate infectious virus bearing defined attenuating mutations and to study the
genetic basis of attenuation of existing vaccine viruses. For example, the three amino acid
substitutions found in the L gene of *cp45*, singularly or in combination, have been found to
specify the *ts* and attenuation phenotypes. Additional *ts* and attenuating mutations are
10 present in other regions of the PIV3 *cp45*. In addition a chimeric PIV1 vaccine candidate has
been generated using the PIV3 cDNA rescue system by replacing the PIV3 HN and F open
reading frames (ORFs) with those of PIV1 in a PIV3 full-length cDNA that contains the
three attenuating mutations in L. The recombinant chimeric virus derived from this cDNA is
designated rPIV3-1.*cp45L* (Skiadopoulos et al., J. Virol. 72:1762-8, 1998; Tao et al., J.
15 Virol. 72:2955-2961, 1998; Tao et al., Vaccine 17:1100-1108, 1999, incorporated herein by
reference). rPIV3-1.*cp45L* was attenuated in hamsters and induced a high level of resistance
to challenge with PIV1. Yet another recombinant chimeric virus, designated rPIV3-1.*cp45*,
has been produced that contains 12 of the 15 *cp45* mutations, i.e., excluding the mutations
that occur in HN and F. This recombinant vaccine candidate is highly attenuated in the
20 upper and lower respiratory tract of hamsters and induces a high level of protection against
HPiV1 infection (Skiadopoulos et al., Vaccine 18:503-510, 1999).

Recently, a number of studies have focused on the possible use of viral
vectors to express foreign antigens toward the goal of developing vaccines against a
pathogen for which other vaccine alternatives are not proved successful. In this context, a
25 number of reports suggest that foreign genes may be successfully inserted into a recombinant
negative strand RNA virus genome or antigenome with varying effects (Bukreyev et al., J.
Virol. 70:6634-41, 1996; Bukreyev et al., Proc. Natl. Acad. Sci. U.S.A. 96:2367-72, 1999;
Finke et al. J. Virol. 71:7281-8, 1997; Hasan et al., J. Gen. Virol. 78:2813-20, 1997; He et
al., Virology 237:249-60, 1997; Jin et al., Virology 251:206-14, 1998; Johnson et al., J.
30 Virol. 71:5060-8, 1997; Kahn et al., Virology 254:81-91, 1999; Kretzschmar et al., J. Virol.
71:5982-9, 1997; Mebatsion et al., Proc. Natl. Acad. Sci. U.S.A. 93:7310-4, 1996; Moriya et
al., FEBS Lett. 425:105-11, 1998; Roberts et al., J. Virol. 73:3723-32, 1999; Roberts et al., J.
Virol. 72:4704-11, 1998; Roberts et al., Virology 247:1-6, 1998; Sakai et al., FEBS Lett.

- 456:221-226, 1999; Schnell et al., Proc. Natl. Acad. Sci. U.S.A. 93:11359-65, 1996a; Schnell et al., J. Virol. 70:2318-23, 1996b; Schnell et al., Cell 90:849-57, 1997; Singh et al., J. Gen. Virol. 80:101-6, 1999; Singh et al., J. Virol. 73:4823-8, 1999; Spielhofer et al., J. Virol. 72:2150-9, 1998; Yu et al., Genes to Cells 2:457-66 et al., 1999; U.S. Patent
- 5 Application Serial No. 09/614,285, filed July 12, 2000 (corresponding to U.S. Provisional Patent Application Serial No. 60/143,425, filed on July 13, 1999, each incorporated herein by reference). When inserted into the viral genome under the control of viral transcription gene-start and gene-end signals, the foreign gene may be transcribed as a separate mRNA and yield significant protein expression. Surprisingly, in some cases foreign sequence has
- 10 been reported to be stable and capable of expressing functional protein during numerous passages *in vitro*.

However, to successfully develop vectors for vaccine use, it is insufficient to simply demonstrate a high, stable level of protein expression. For example, this has been possible since the early-to-mid 1980s with recombinant vaccinia viruses and adenoviruses,

15 and yet these vectors have proven to be disappointments in the development of vaccines for human use. Similarly, most nonsegmented negative strand viruses which have been developed as vectors do not possess properties or immunization strategies amenable for human use. Examples in this context include vesicular stomatitis virus, an ungulate pathogen with no history of administration to humans except for a few laboratory accidents;

20 Sendai virus, a mouse pathogen with no history of administration to humans; simian virus 5, a canine pathogen with no history of administration to humans; and an attenuated strain of measles virus which must be administered systemically and would be neutralized by measles-specific antibodies present in nearly all humans due to maternal antibodies and widespread use of a licensed vaccine. Furthermore, some of these prior vector candidates

25 have adverse effects, such as immunosuppression, which are directly inconsistent with their use as vectors. Thus, one must identify vectors whose growth characteristics, tropisms, and other biological properties make them appropriate as vectors for human use. It is further necessary to develop a viable vaccination strategy, including an immunogenic and efficacious route of administration.

30 The three human mononegaviruses that are currently being considered as vaccine vectors, namely measles, mumps, and rabies viruses, have additional limitations that combine to make them weak candidates for further development as vectors. For example, measles virus has been considered for use a vector for the protective antigen of hepatitis B

virus (Singh et al., *J. Virol.* 73:4823-8, 1999). However, this combined measles virus-hepatitis B virus vaccine could only be given, like the licensed measles virus vaccine, after nine months of age, whereas the current hepatitis B virus vaccine is recommended for use in early infancy. This is because the currently licensed measles virus vaccine is administered parenterally and is very sensitive to neutralization and immunosuppression by maternal antibodies, and therefore is not effective if administered before 9-15 months of age. Thus, it could not be used to vector antigens that cause disease in early infancy and therefore would not be useful for viruses such as RSV and the HPIVs. Another well known, characteristic effect of measles virus infection is virus-mediated immunosuppression, which can last several months. Immunosuppression would not be a desirable feature for a vector. The attenuated measles virus vaccine was associated with altered immune responses and excess mortality when administered at increased dose, which might be due at least in part to virus-induced immunosuppression and indicates that even an attenuated measles virus might not be appropriate as a vector. Furthermore, the use of measles virus as a vector would be inconsistent with the global effort to eradicate this pathogen. Indeed, for these reasons it would be desirable to end the use of live measles virus and replace the present measles virus vaccine with a PIV vector that expresses measles virus protective antigens, as described herein.

Rabies virus, a rare cause of infection of humans, has been considered for use as a vector (Mebatsion et al., *Proc. Natl. Acad. Sci. USA* 93:7310-4, 1996), but it is unlikely that a vector that is 100% fatal for humans would be developed for use as a live attenuated virus vector, especially since immunity to the rabies virus, which is not a ubiquitous human pathogen, is not needed for the general population. While mumps and measles viruses are less pathogenic, infection by either virus can involve undesirable features. Mumps virus infects the parotid gland and can spread to the testes, sometimes resulting in sterility. Measles virus establishes a viremia, and the widespread nature of its infection is exemplified by the associated widespread rash. Mild encephalitis during mumps and measles infection is not uncommon. Measles virus also is associated with a rare progressive fatal neurological disease called subacute sclerosing encephalitis. In contrast, PIV infection and disease in normal individuals is limited to the respiratory tract, a site that is much more advantageous for immunization than the parental route. Viremia and spread to second sites can occur in severely immunocompromised experimental animals and humans, but this is not a characteristic of the typical PIV infection. Acute respiratory tract disease is the only disease associated with PIVs. Thus, use of PIVs as vectors will, on the basis of their biological

characteristics, avoid complications such as interaction of virus with peripheral lymphocytes, leading to immunosuppression, or infection of secondary organs such as the testes or central nervous system, leading to other complications.

Among a host of human pathogens for which a vector-based vaccine approach may be desirable is the measles virus. A live attenuated vaccine has been available for more than three decades and has been largely successful in eradicating measles disease in the United States. However, the World Health Organization estimates that more than 45 million cases of measles still occur annually, particularly in developing countries, and the virus contributes to approximately one million deaths per year

Measles virus is a member of the *Morbillivirus* genus of the *Paramyxoviridae* family (Griffin et al., In "Fields Virology", B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, B. Roizman, and S. E. Straus, Eds., Vol. 1, pp. 1267-1312. Lippincott-Raven Publishers, Philadelphia, 1996). It is one of the most contagious infectious agents known to man and is transmitted from person to person via the respiratory route (Griffin et al., In "Fields Virology", B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, B. Roizman, and S. E. Straus, Eds., Vol. 1, pp. 1267-1312. Lippincott-Raven Publishers, Philadelphia, 1996). The measles virus has a complex pathogenesis, involving replication in both the respiratory tract and various systemic sites (Griffin et al., In "Fields Virology", B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, B. Roizman, and S. E. Straus, Eds., Vol. 1, pp. 1267-1312. Lippincott-Raven Publishers, Philadelphia, 1996).

Although both mucosal IgA and serum IgG measles virus-specific antibodies can participate in the control of measles virus, the absence of measles virus disease in very young infants possessing maternally-acquired measles virus-specific antibodies identifies serum antibodies as a major mediator of resistance to disease (Griffin et al., In "Fields Virology", B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, B. Roizman, and S. E. Straus, Eds., Vol. 1, pp. 1267-1312. Lippincott-Raven Publishers, Philadelphia, 1996). The two measles virus glycoproteins, the hemagglutinin (HA) and fusion (F) proteins, are the major neutralization and protective antigens (Griffin et al., In "Fields Virology", B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, B. Roizman, and S. E. Straus, Eds., Vol. 1, pp. 1267-1312. Lippincott-Raven Publishers, Philadelphia, 1996).

The currently available live attenuated measles vaccine is administered by a parenteral route (Griffin et al., In "Fields Virology", B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, B. Roizman, and S. E. Straus, Eds., Vol. 1, pp. 1267-1312. Lippincott-Raven Publishers, Philadelphia, 1996). Both the wild type
5 measles virus and the vaccine virus are very readily neutralized by antibodies, and the measles virus vaccine is rendered non-infectious by even very low levels of maternally-acquired measles virus-specific neutralizing antibodies (Halsey et al., N. Engl. J. Med. 313:544-9, 1985; Osterhaus et al., Vaccine 16:1479-81, 1998). Thus, the vaccine virus is not given until the passively-acquired maternal antibodies have decreased to undetectable levels.
10 In the United States, measles virus vaccine is not given until 12 to 15 months of age, a time when almost all children are readily infected with the measles virus vaccine. In the developing world, measles virus continues to have a high mortality rate, especially in children within the latter half of the first year of life (Gellin et al., J. Infect. Dis. 170:S3-14, 1994; Taylor et al., Am. J. Epidemiol. 127:788-94, 1988). This occurs because the measles
15 virus, which is highly prevalent in these regions, is able to infect that subset of infants in whom maternally-acquired measles virus-specific antibody levels have decreased to a non-protective level. Therefore, there is a need for a measles virus vaccine that is able to induce a protective immune response even in the presence of measles virus neutralizing antibodies with the goal of eliminating measles virus disease occurring within the first year of life as
20 well as that which occurs thereafter. Given this need, there have been numerous attempts to develop an immunization strategy to protect infants in the latter half of the first year of life against measles virus, but none of these strategies has been effective to date.

The first strategy for developing an early measles vaccine involved administration of the licensed live attenuated measles virus vaccine to infants about six
25 months of age by one of the following two methods (Cutts et al., Biologicals 25:323-38, 1997). In one general protocol, the live attenuated measles virus was administered intranasally by drops (Black et al., New Eng. J. Med. 263:165-169; 1960; Kok et al., Trans. R. Soc. Trop. Med. Hyg. 77:171-6, 1983; Simasathien et al., Vaccine 15:329-34, 1997) or into the lower respiratory tract by aerosol (Sabin et al., J. Infect. Dis. 152:1231-7, 1985), to
30 initiate an infection of the respiratory tract. In a second protocol, the measles virus was given parenterally but at a higher dose than that employed for the current vaccine. The administration of vaccines that can replicate on mucosal surfaces has been successfully achieved in early infancy for both live attenuated poliovirus and rotavirus vaccines (Melnick et al., In "Fields Virology", B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L.

Melnick, T. P. Monath, B. Roizman, and S. E. Straus, Eds., Vol. 1, pp. 655-712. 2 vols. Lippencott-Raven Publishers, Philadelphia, 1996; Perez-Schael et al., N. Engl. J. Med. 337:1181-7, 1997), presumably because passively-acquired IgG antibodies have less access to mucosal surfaces than they do to systemic sites of viral replication. In this situation, the live attenuated poliovirus vaccine viruses are able to infect the mucosal surface of the gastrointestinal tract or the respiratory tract of young infants, including those with maternal antibodies, resulting in the induction of a protective immune response.

Therefore, a plausible method is to immunize via the respiratory tract of the young infant with the live attenuated measles virus vaccine, since this is the natural route of infection with the measles virus. However, the live attenuated measles virus that is infectious by the parenteral route was inconsistently infectious by the intranasal route (Black et al., New Eng. J. Med. 263:165-169, 1960; Cutts et al., Biologicals 25:323-38, 1997; Kok et al., Trans. R. Soc. Trop. Med. Hyg. 77:171-6, 1983; Simasathien et al., Vaccine 15:329-34, 1997), and this decreased infectivity was especially apparent for the Schwartz strain of measles virus vaccine which is the current vaccine strain. Presumably, during the attenuation of this virus by passage in tissue culture cells of avian origin, the virus lost a significant amount of infectivity for the upper respiratory tract of humans. Indeed, a hallmark of measles virus biology is that the virus undergoes rapid changes in biological properties when grown *in vitro*. Since this relatively simple route of immunization was not successful, a second approach was tried involving administration of the live virus vaccine by aerosol into the lower respiratory tract (Cutts et al., Biologicals 25:323-38, 1997; Sabin et al., J. Infect. Dis. 152:1231-7, 1985).

Infection of young infants by aerosol administration of measles virus vaccine was accomplished in highly controlled experimental studies, but it has not been possible to reproducibly deliver a live attenuated measles virus vaccine in field settings by aerosol to the young uncooperative infant (Cutts et al., Biologicals 25:323-38, 1997). In another attempt to immunize six-month old infants, the measles vaccine virus was administered parenterally at a 10- to 100-fold increased dose (Markowitz et al., N. Engl. J. Med. 322:580-7, 1990). Although high-titer live measles vaccination improved seroconversion in infants 4-6 months of age, there was an associated increase in mortality in the high-titer vaccine recipients later in infancy (Gellin et al., J. Infect. Dis. 170:S3-14, 1994; Holt et al., J. Infect. Dis. 168:1087-96, 1993; Markowitz et al., N. Engl. J. Med. 322:580-7, 1990) and this approach to immunization has been abandoned.

A second strategy previously explored for a measles virus vaccine was the use of an inactivated measles virus vaccine, specifically, a formalin inactivated whole measles virus or a subunit virus vaccine prepared from measles virus (Griffin et al., In "Fields Virology", B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, B. Roizman, and S. E. Straus, Eds., Vol. 1, pp. 1267-1312. Lippincott-Raven Publishers, Philadelphia, 1996). However, the clinical use of the vaccines in the 1960's revealed a very serious complication, namely, that the inactivated virus vaccines potentiated disease rather than prevented it (Fulginiti et al., JAMA 202:1075-80, 1967). This was first observed with formalin-inactivated measles virus vaccine (Fulginiti et al., JAMA 202:1075-80, 1967). Initially, this vaccine prevented measles, but after several years vaccinees lost their resistance to infection. When subsequently infected with naturally circulating measles virus, the vaccinees developed an atypical illness with accentuated systemic symptoms and pneumonia (Fulginiti et al., JAMA 202:1075-80, 1967; Nader et al., J. Pediatr. 72:22-8, 1968; Rauh et al., Am. J. Dis. Child 109:232-7, 1965). Retrospective analysis showed that formalin inactivation destroyed the ability of the measles fusion (F) protein to induce hemolysis-inhibiting antibodies, but it did not destroy the ability of the HA (hemagglutinin or attachment) protein to induce neutralizing antibodies (Norrby et al., J. Infect. Dis. 132:262-9, 1975; Norrby et al., Infect. Immun. 11:231-9, 1975). When the immunity induced by the HA protein had waned sufficiently to permit extensive infection with wild type measles virus, an altered and sometimes more severe disease was seen at the sites of measles virus replication (Bellanti, Pediatrics 48:715-29, 1971; Buser, N. Engl. J. Med. 277:250-1, 1967). This atypical disease is believed to be mediated in part by an altered cell-mediated immune response in which Th-2 cells were preferentially induced leading to heightened disease manifestations at the sites of viral replication (Polack et al., Nat. Med. 5:629-34, 1999). Because of this experience with nonliving measles virus vaccines and also because the immunogenicity of such parenterally-administered vaccines can be decreased by passively-transferred antibodies, there has been considerable reluctance to evaluate such vaccines in human infants. It should be noted that disease potentiation appears to be associated only with killed vaccines.

Yet another strategy that has been explored for developing a vaccine against measles for use in young infants has been the use of viral vectors to express a protective antigen of the measles virus (Drillien et al., Proc. Natl. Acad. Sci. U.S.A. 85:1252-6, 1988; Fooks et al., J. Gen. Virol. 79:1027-31, 1998; Schnell et al., Proc. Natl. Acad. Sci. U.S.A. 93:11359-65, 1996a; Taylor et al., Virology 187:321-8, 1992; Wild et al., Vaccine 8:441-2,

1990; Wild et al., J. Gen. Virol. **73**:359-67, 1992). A variety of vectors have been explored including pox viruses such as the replication-competent vaccinia virus or the replication-defective modified vaccinia virus Ankara (MVA) strain. Replication-competent vaccinia recombinants expressing the F or HA glycoprotein of measles virus were efficacious in immunologically naive vaccinees. However, when they were administered parenterally in the presence of passive antibody against measles virus, their immunogenicity and protective efficacy was largely abrogated (Galletti et al., Vaccine **13**:197-201, 1995; Osterhaus et al., Vaccine **16**:1479-81, 1998; Siegrist et al., Vaccine **16**:1409-14, 1998; Siegrist et al., Dev. Biol. Stand. **95**:133-9, 1998).

10 Replication-competent vaccinia recombinants expressing the protective antigens of RSV have also been shown to be ineffective in inducing a protective immune response when they are administered parenterally in the presence of passive antibody (Murphy et al., J. Virol. **62**:3907-10, 1988a), but they readily protected such hosts when administered intranasally. Unfortunately, replication-competent vaccinia virus recombinants are not sufficiently attenuated for use in immunocompromised hosts such as persons with human immunodeficiency virus (HIV) infection (Fenner et al., World Health Organization, Geneva, 1988; Redfield et al., N. Engl. J. Med. **316**:673-676, 1987), and their administration by the intranasal route even to immunocompetent individuals would be problematic. Therefore they are not being pursued as vectors for use in human infants, some of whom could be infected with HIV.

25 The MVA vector, which was derived by more than 500 passages in chick embryo cells (Mayr et al., Infection **3**:6-14, 1975; Meyer et al., J. Gen. Virol. **72**:1031-1038, 1991), has also been evaluated as a potential vaccine vector for the protective antigens of several paramyxoviruses (Durbin et al., J. Infect. Dis. **179**:1345-51, 1999a; Wyatt et al., Vaccine **14**:1451-1458, 1996). MVA is a highly attenuated host range mutant that replicates well in avian cells but not in most mammalian cells, including those obtained from monkeys and humans (Blanchard et al., J. Gen. Virol. **79**:1159-1167, 1998; Carroll et al., Virology **238**:198-211, 1997; Drexler et al., J. Gen. Virol. **79**:347-352, 1998; Sutter et al., Proc. Natl. Acad. Sci. U.S.A. **89**:10847-10851, 1992). Avipox vaccine vectors, which have a host range restriction similar to that of MVA, also have been constructed that express measles virus protective antigens (Taylor et al., Virology **187**:321-8, 1992). MVA is non-pathogenic in immunocompromised hosts and has been administered to large numbers of humans without incident (Mayr et al., Zentralbl. Bakteriologie [B] **167**:375-90, 1978; Stickle et al.,

Dtsch. Med. Wochenschr. 99:2386-92, 1974; Werner et al., Archives of Virology 64:247-256, 1980). Unfortunately, both the immunogenicity and efficacy of MVA expressing a paramyxovirus protective antigen were abrogated in passively-immunized rhesus monkeys whether delivered by a parenteral or a topical route (Durbin et al., Virology 235:323-332, 5 1999). The immunogenicity of DNA vaccines expressing measles virus protective antigens delivered parenterally was also decreased in passively-immunized hosts (Siegrist et al., Dev. Biol. Stand. 95:133-9, 1998). Replication-defective vectors expressing measles virus protective antigens are presently being evaluated, including adenovirus-measles virus HA recombinants (Fooks et al., J. Gen. Virol. 79:1027-31, 1998). In this context, MVA 10 recombinants expressing parainfluenza virus antigens, unlike replication-competent vaccinia virus recombinants, lacked protective efficacy when given by a mucosal route to animals with passively-acquired antibodies, and it is unlikely that they, or the similar avipox vectors, can be used in infants with maternally-acquired measles virus antibodies.

Based on the reports summarized above, it appears unlikely that a replication- 15 competent or replication-defective poxvirus vector, or a DNA vaccine, expressing a measles virus protective antigen will be satisfactorily immunogenic or efficacious in infants possessing passively-acquired maternal measles virus-specific antibodies.

A recently developed replication-competent virus vector expressing measles virus HA that replicates in the respiratory tract of animal hosts has been developed, namely, 20 vesicular stomatitis virus (VSV), a rhabdovirus which naturally infects cattle but not humans (Roberts et al., J. Virol. 73:3723-32, 1999; Schnell et al., Proc. Natl. Acad. Sci. U.S.A. 93:11359-65, 1996a). Since VSV is an animal virus that can cause disease in humans, development of this recombinant for use in humans will require that a VSV backbone that is satisfactorily attenuated in human infants be first identified (Roberts et al., J. Virol. 73:3723- 25 32, 1999), but such clinical studies have not been initiated.

Although there have been numerous advances toward development of effective vaccine agents against PIV and other pathogens, including measles, there remains a clear need in the art for additional tools and methods to engineer safe and effective vaccines to alleviate the serious health problems attributable to these pathogens, particularly among 30 young infants. Among the remaining challenges in this context is the need for additional tools to generate suitably attenuated, immunogenic and genetically stable vaccine candidates for use in diverse clinical settings against one or more pathogens. To facilitate these goals,

existing methods for identifying and incorporating attenuating mutations into recombinant vaccine strains and for developing vector-based vaccines and immunization methods must be expanded. Surprisingly, the present invention fulfills these needs and provides additional advantages as described herein below.

5

Throughout the description and the claims of this specification the word "comprise" and variations of the word, such as "comprising" and "comprises" is not intended to exclude other additives, components, integers or steps.

10 The discussion of documents, acts, materials, devices, articles and the like is included in this specification solely for the purpose of providing a context for the present invention. It is not suggested or represented that any or all of these matters formed part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed in Australia before the priority date of each claim of this application.

15

SUMMARY OF THE INVENTION

20 The present invention provides chimeric parainfluenza viruses (PIVs) that are infectious in humans and other mammals and are useful in various compositions to generate desired immune responses against one or more PIVs, or against a PIV and one or more additional pathogens in a host susceptible to infection therefrom. In preferred aspects, the invention provides novel methods for designing and producing attenuated, chimeric PIVs that are useful as vaccine agents for preventing and/or treating infection and related disease symptoms attributable to PIV and one or more additional pathogens. Included within these aspects of the invention are novel, isolated polynucleotide molecules and vectors incorporating such molecules that comprise a chimeric PIV genome or antigenome including 25 a partial or complete PIV vector genome or antigenome combined or integrated with one or more heterologous genes or genome segments that encode single or multiple antigenic determinants of a heterologous pathogen or of multiple heterologous pathogens. Also provided within the invention are methods and compositions incorporating a chimeric PIV for prophylaxis and treatment of infection by both a selected PIV and one or more 30 heterologous pathogens, e.g., a heterologous PIV or a non-PIV pathogen such as a measles virus.

The invention thus involves methods and compositions for developing live vaccine candidates based on chimeras that employ a parainfluenza virus or subviral particle that is recombinantly modified to incorporate one or more antigenic determinants of a heterologous pathogen(s). Chimeric PIVs of the invention are constructed through a cDNA-based virus recovery system. Recombinant chimeric PIVs made from cDNA replicate independently and are propagated in a similar manner as biologically-derived viruses. The recombinant viruses are engineered to incorporate nucleotide sequences from both a vector (i.e., a "recipient" or "background") PIV genome or antigenome, and one or more heterologous "donor" sequences encoding one or more antigenic determinants of a different PIV or heterologous pathogen—to produce an infectious, chimeric virus or subviral particle. In this manner, candidate vaccine viruses are recombinantly engineered to elicit an immune response against one or more PIVs or a polyspecific response against a selected PIV and a non-PIV pathogen in a mammalian host susceptible to infection therefrom. Preferably the PIV and/or non-PIV pathogen(s) from which the heterologous sequences encoding the antigenic determinant(s) are human pathogens and the host is a human host. Also preferably, the vector PIV is a human PIV, although non-human PIVs, for example a bovine PIV (BPIV), can be employed as a vector to incorporate antigenic determinants of human PIVs and other human pathogens. Chimeric PIVs according to the invention may elicit an immune response against a specific PIV, e.g., HPIV1, HPIV2, HPIV3, or a polyspecific immune response against multiple PIVs, e.g., HPIV1 and HPIV2. Alternatively, chimeric PIVs of the invention may elicit a polyspecific immune response against one or more PIVs and a non-PIV pathogen such as measles virus.

The present invention provides an isolated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of one or more heterologous pathogen(s) to form a chimeric PIV genome or antigenome.

30

Exemplary chimeric PIV of the invention incorporate a chimeric PIV genome or antigenome as described above, as well as a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), and a large polymerase protein (L). Additional PIV proteins may be included in various combinations to provide a range of infectious subviral particles, up to a complete viral particle or a viral particle containing supernumerary proteins, antigenic determinants or other additional components.

The present invention also provides an isolated polynucleotide comprising a chimeric VIP genome or antigenome which includes a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of one or more heterologous pathogen(s) to form a chimeric PIV genome or antigenome.

Chimeric PIV of the invention include a partial or complete "vector" PIV genome or antigenome derived from or patterned after a human PIV or non-human PIV combined with one or more heterologous gene(s) or genome segment(s) of a different PIV or other pathogen to form the chimeric PIV genome or antigenome. In preferred aspects of the invention, chimeric PIV incorporate a partial or complete human PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a second human PIV or a non-PIV pathogen such as measles virus.

The PIV "vector" genome or antigenome typically acts as a recipient or carrier to which are added or incorporated one or more "donor" genes or genome segments of a heterologous pathogen. Typically, polynucleotides encoding one or more antigenic determinants of the heterologous pathogen are added to or substituted within the vector genome or antigenome to yield a chimeric PIV that thus acquires the ability to elicit an immune response in a selected host against the heterologous pathogen. In addition, the chimeric virus may exhibit other novel phenotypic characteristics compared to one or both of

the vector PIV and heterologous pathogens. For example, addition or substitution of heterologous genes or genome segments within a vector PIV strain may additionally, or independently, result in an increase in attenuation, growth changes, or other desired phenotypic changes as compared with a corresponding phenotype of the unmodified vector virus and/or donor. In one aspect of the invention, chimeric PIVs are attenuated for greater efficacy as a vaccine candidate by incorporation of large polynucleotide inserts which specify the level of attenuation in the resulting chimeric virus dependent upon the size of the insert.

Preferred chimeric PIV vaccine candidates of the invention bear one or more major antigenic determinants of a human PIV, e.g., of HPIV1, HPIV2 or HPIV3, and thus elicit an effective immune response against the selected PIV in human hosts. The antigenic determinant which is specific for a selected human PIV may be encoded by the vector genome or antigenome, or may be inserted within or joined to the PIV vector genome or antigenome as a heterologous polynucleotide sequence from a different PIV. The major protective antigens of human PIVs are their HN and F glycoproteins, although other proteins can also contribute to a protective or therapeutic immune response. In this context, both humoral and cell mediated immune responses are advantageously elicited by representative vaccine candidates within the invention. Thus, polynucleotides encoding antigenic determinants that may be present in the vector genome or antigenome, or integrated therewith as a heterologous gene or genome segment, may encode one or more PIV N, P, C, D, V, M, F, HN and/or L protein(s) or selected immunogenic fragment(s) or epitope(s) thereof from any human PIV.

In addition to having one or more major antigenic determinants of a selected human PIV, preferred chimeric PIV vaccine viruses of the invention bear one or more major antigenic determinants of a second human PIV or of a non-PIV pathogen. In exemplary aspects, the chimeric PIV includes a vector genome or antigenome that is a partial or complete human PIV (HPIV) genome or antigenome, for example of HPIV3, and further includes one or more heterologous gene(s) or genome segment(s) encoding antigenic determinant(s) of at least one heterologous PIV, for example HPIV1 and/or HPIV2. Preferably, the vector genome or antigenome is a partial or complete HPIV3 genome or antigenome and the heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are of one or more heterologous HPIV(s). In alternative embodiments, one or more genes or genome segments encoding one or more antigenic determinants of HPIV1

may be added to or substituted within the partial or complete HPIV3 genome or antigenome. Preferably, the antigenic determinant(s) of HPIV1 is/are selected from HPIV1 HN and F glycoproteins or comprise one or more antigenic domains, fragments or epitopes of the HN and/or F glycoproteins. In various exemplary embodiments, both of the HPIV1 genes
5 encoding the HN and F glycoproteins are substituted for counterpart HPIV3 HN and F genes in the HPIV3 vector genome or antigenome. These constructs yield chimeric PIVs that elicit a mono- or poly-specific immune response in humans to HPIV3 and/or HPIV1.

In additional exemplary embodiments, one or more genes or genome segments encoding one or more antigenic determinants of HPIV2 is/are added to, or
10 incorporated within, a partial or complete HPIV3 genome or antigenome, yielding a new or additional immunospecificity of the resultant chimera against HPIV2 alone, or against HPIV3 and HPIV2. In more detailed aspects, one or more HPIV2 genes or genome segments encoding one or more HN and/or F glycoproteins or antigenic domains, fragments or epitopes thereof is/are added to or incorporated within the partial or complete HPIV3
15 vector genome or antigenome.

In yet additional aspects of the invention, multiple heterologous genes or genome segments encoding antigenic determinants of multiple heterologous PIVs are added to or incorporated within a partial or complete PIV vector genome or antigenome, preferably an HPIV vector genome or antigenome. In one preferred embodiment, heterologous genes
20 or genome segments encoding antigenic determinants from both HPIV1 and HPIV2 are added to or incorporated within a partial or complete HPIV3 vector genome or antigenome. In more detailed aspects, one or more HPIV1 genes or genome segments encoding one or more HN and/or F glycoproteins (or antigenic domains, fragments or epitopes thereof) and one or more HPIV2 genes or genome segments encoding HN and/or F glycoproteins,
25 antigenic domains, fragments or epitopes, is/are added to or incorporated within the partial or complete HPIV3 vector genome or antigenome. In one example, both HPIV1 genes encoding HN and F glycoproteins are substituted for counterpart HPIV3 HN and F genes to form a chimeric HPIV3-1 vector genome or antigenome, which is further modified by addition or incorporation of one or more genes or gene segments encoding single or multiple
30 antigenic determinants of HPIV2. This is readily achieved within the invention, for example, by adding or substituting a transcription unit comprising an open reading frame (ORF) of an HPIV2 HN within the chimeric HPIV3-1 vector genome or antigenome. Following this method, specific constructs exemplifying the invention are provided which

yield chimeric PIVs having antigenic determinants of both HPIV1 and HPIV2, as exemplified by the vaccine candidates rPIV3-1.2HN and rPIV3-1cp45.2HN described herein below.

In alternative aspects of the invention, chimeric PIVs of the invention are based on a human PIV vector genome or antigenome which is employed as a recipient for incorporation of major antigenic determinants from a non-PIV pathogen. Pathogens from which one or more antigenic determinants may be adopted into the chimeric PIV vaccine candidate include, but are not limited to, measles virus, subgroup A and subgroup B respiratory syncytial viruses, mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses. This assemblage of pathogens that may be thus targeted for vaccine development according to the methods of the invention is exemplary only, and those skilled in the art will understand that the use of PIV vectors for carrying antigenic determinants extends broadly to a large host of additional pathogens.

This, in various alternative aspects of the invention, a human PIV genome or antigenome can be employed as a vector for incorporation of one or more major antigenic determinants from a wide range of non-PIV pathogens. Representative major antigens that can be incorporated within chimeric PIVs of the invention include, but are not limited to the measles virus HA and F proteins; the F, G, SH and M2 proteins of subgroup A and subgroup B respiratory syncytial virus, mumps virus HN and F proteins, human papilloma virus L1 protein, type 1 or type 2 human immunodeficiency virus gp160 protein, herpes simplex virus and cytomegalovirus gB, gC, gD, gE, gG, gH, gI, gJ, gK, gL, and gM proteins, rabies virus G Protein, Epstein Barr Virus gp350 protein; filovirus G protein, bunyavirus G protein, flavivirus E and NS1 proteins, and alphavirus E protein.

Various human PIV vectors can be employed to carry heterologous antigenic determinants of non-PIV pathogens to elicit one or more specific humoral or cell mediated immune responses against the antigenic determinant(s) carried by the chimeric vaccine virus and hence elicit an effective immune response against the wild-type "donor" pathogen in susceptible hosts. In preferred embodiments, one or more heterologous genes or genome segments from the donor pathogen is joined to or inserted within a partial or complete HPIV3 genome or antigenome. Alternatively, the heterologous gene or genome segment

may be incorporated within a chimeric HPIV vector genome or antigenome, for example a partial or complete HPIV3 genome or antigenome bearing one or more genes or genome segments of a heterologous PIV. For example, the gene(s) or genome segment(s) encoding the antigenic determinant(s) of a non-PIV pathogen may be combined with a partial or complete chimeric HPIV3-1 vector genome or antigenome, e.g., as described above having one or both HPIV1 genes encoding HN and F glycoproteins substituted for counterpart HPIV3 HN and F genes. Alternatively, the gene(s) or genome segment(s) encoding the antigenic determinant(s) of a non-PIV pathogen may be combined with a partial or complete chimeric genome or antigenome that incorporates single or multiple antigenic determinants of HPIV2, e.g., an HPIV2 HN gene, within an HPIV1 or HPIV3 vector genome or antigenome, or a chimeric HPIV3-1 vector genome or antigenome as described above. The heterologous gene(s) or genome segment(s) encoding one or more measles antigenic determinant(s) may be combined with any of the PIV vectors or chimeric PIV vectors disclosed herein. In the examples provided herein, the vector genome or antigenome is a partial or complete HPIV3 genome or antigenome, or a chimeric HPIV genome or antigenome comprising a partial or complete HPIV3 genome or antigenome having one or more genes or genome segments encoding antigenic determinant(s) of a heterologous HPIV added or incorporated therein. In one such chimeric construct, a transcription unit comprising an open reading frame (ORF) of a measles virus HA gene is added to a HPIV3 vector genome or antigenome at various positions, yielding exemplary chimeric PIV/measles vaccine candidates rPIV3(HA HN-L), rPIV3(HA N-P), *rcp45L*(HA N-P), rPIV3(HA F-M), or *rcp45L*(HA P-M).

In additional exemplary embodiments, the PIV vector genome or antigenome is a chimeric HPIV genome or antigenome comprising a partial or complete HPIV3 genome or antigenome having one or more gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of HPIV1 added or incorporated therein. This construct may be used as a vector, e.g., for measles virus, wherein the heterologous antigenic determinant(s) is/are selected from the measles virus HA and F proteins and antigenic domains, fragments and epitopes thereof. In one example, a transcription unit comprising an open reading frame (ORF) of a measles virus HA gene is added to or incorporated within a HPIV3-1 vector genome or antigenome having both the HPIV3 HN and F ORFs substituted by the HN and F ORFs of HPIV1. Among this category of recombinants are vaccine candidates identified herein below as rPIV3-1 HAP-M or rPIV3-1 HAP-M cp45L.

The present invention provides an isolated infectious recombinant parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a PIV genome or antigenome having a polynucleotide insertion of between 150 nucleotides (nts) and 4,000 nucleotides in length in a non-coding region (NCR) of the genome or antigenome or as a separate gene unit (GU), said polynucleotide insertion lacking a complete open reading frame (ORF) and specifying an attenuated phenotype in said recombinant PIV.

In other detailed embodiments of the invention, the partial or complete PIV vector genome or antigenome is combined with one or more "supernumerary" (i.e., additional to a full complement of genes, whether present in a wild-type vector or in a mutant, e.g., chimeric vector backbone) heterologous gene(s) or genome segment(s) to form the chimeric PIV genome or antigenome. The vector genome or antigenome is often a complete HPIV3 or HPIV3-1 chimeric genome or antigenome, and the supernumerary heterologous gene(s) or genome segment(s) are selected from HPIV1 HN, HPIV1 F, HPIV2 HN, HPIV2 F, measles HA, and/or a translationally silent synthetic gene unit. In certain exemplary embodiments, one or both of the HPIV1 HN and/or HPIV2 HN ORF(s) is/are inserted within the HPIV3 vector genome or antigenome, respectively. In more detailed embodiments, the HPIV1 HN, HPIV2 HN, and measles virus HA ORFs are inserted between the N/P, P/M, and HN/L genes, respectively. Alternatively, the HPIV1 HN and HPIV2 HN genes may be inserted between the N/P and P/M genes, respectively and a 3918-nt GU insert is added between the HN and L genes. Among this category of recombinants are vaccine candidates identified herein below as rHPIV3 1HNN-P, rHPIV3 1HNP-M, rHPIV3 2HNN-P, rHPIV3 2HNP-M, rHPIV3 1HNN-P 2HNP-M, rHPIV3 1HNN-P 2HNP-M HAHN-L, and rHPIV3 1HNN-P 2HNP-M 3918GUHN-L.

Thus designed and constructed, chimeric PIV of the invention may contain protective antigens from one, two, three, four or more different pathogens. For example, vaccine candidates are provided which contain protective antigens from one to four pathogens selected from HPIV3, HPIV1, HPIV2, and measles virus. To construct such multi-specific vaccine candidates, one or more supernumerary heterologous gene(s) or genome segment(s) can be added which may add a total length of supernumerary foreign sequence to the recombinant genome or antigenome of 30% to 50% or greater (e.g., compared to the wild-type HPIV3 genome length of 15,462 nt). The addition of one or more

supernumerary heterologous gene(s) or genome segment(s) in this context often specifies an attenuation phenotype of the chimeric PIV, which exhibits at least a 10-to 100-fold, often 100- to 1,000-fold, and up to a 1,000- to 10,000-fold or greater decrease in replication in the upper and/or lower respiratory tract.

5 To construct chimeric PIV clones of the invention, a heterologous gene or genome segment of a donor PIV or non-PIV pathogen may be added or substituted at any operable position in the vector genome or antigenome. Often, the position of a gene or gene segment substitution will correspond to a wild-type gene order position of a counterpart gene

10

15



19a

or genome segment within the partial or complete PIV vector genome or antigenome. In other embodiments, the heterologous gene or genome segment is added or substituted at a position that is more promoter-proximal or promotor-distal compared to a wild-type gene order position of a counterpart gene or genome segment within the background genome or antigenome, to enhance or reduce expression, respectively, of the heterologous gene or genome segment. In more detailed aspects of the invention, a heterologous genome segment, for example a genome segment encoding an immunogenic ectodomain of a heterologous PIV or non-PIV pathogen, can be substituted for a corresponding genome segment in a counterpart gene in the PIV vector genome or antigenome to yield constructs encoding chimeric proteins, e.g. fusion proteins having a cytoplasmic tail and/or transmembrane domain of one PIV fused to an ectodomain of another PIV or non-PIV pathogen. In alternate embodiments, a chimeric PIV genome or antigenome may be engineered to encode a polyspecific chimeric glycoprotein in the recombinant virus or subviral particle having immunogenic glycoprotein domains or epitopes from two different pathogens. In yet additional embodiments, heterologous genes or genome segments from one PIV or non-PIV pathogen can be added (i.e., without substitution) within a PIV vector genome or antigenome to create novel immunogenic properties within the resultant clone. In these cases, the heterologous gene or genome segment may be added as a supernumerary gene or genome segment, optionally for the additional purpose of attenuating the resultant chimeric virus, in combination with a complete PIV vector genome or antigenome. Alternatively, the heterologous gene or genome segment may be added in conjunction with deletion of a selected gene or genome segment in the vector genome or antigenome.

In preferred embodiments of the invention, the heterologous gene or genome segment is added at an intergenic position within the partial or complete PIV vector genome or antigenome. Alternatively, the gene or genome segment can be inserted within other noncoding regions of the genome, for example, within 5' or 3' noncoding regions or in other positions where noncoding nucleotides occur within the vector genome or antigenome. In some instances, it may be desired to insert the heterologous gene or genome segment at a non-coding site corresponding to or overlapping a cis-acting regulatory sequence within the vector genome or antigenome, e.g., within a sequence required for efficient replication, transcription, and/or translation. These regions of the vector genome or antigenome represent target sites for disruption or modification of regulatory functions associated with introduction of the heterologous gene or genome segment.

For the preferred purpose of constructing candidate vaccine viruses for clinical use, it is often desirable to adjust the attenuation phenotype of chimeric PIV of the invention by introducing additional mutations that increase or decrease the level of attenuation in the recombinant virus. Therefore, in additional aspects of the invention, attenuated, chimeric PIVs are produced in which the chimeric genome or antigenome is further modified by introducing one or more attenuating mutations that specify an attenuating phenotype in the resultant virus or subviral particle. These attenuating mutations may be generated *de novo* and tested for attenuating effects according to well known rational design mutagenesis strategies. Alternatively, the attenuating mutations may be identified in existing biologically derived mutant PIV or other viruses and thereafter incorporated into a chimeric PIV of the invention.

Preferred attenuating mutations in the latter context are readily identified and incorporated into a chimeric PIV, either by inserting the mutation within the vector genome or antigenome by cloning or mutagenizing the vector genome or antigenome to contain the attenuating mutation. Preferably, attenuating mutations are engineered within the vector genome or antigenome and are imported or copied from biologically derived, attenuated PIV mutants. These are recognized to include, for example, cold passaged (*cp*), cold adapted (*ca*), host range restricted (*hr*), small plaque (*sp*), and/or temperature sensitive (*ts*) PIV mutants. In exemplary embodiments, one or more attenuating mutations present in the well characterized JS HPIV3 *cp45* mutant strain are incorporated within chimeric PIV of the invention, preferably including one or more mutations identified in the polymerase L protein, e.g., at a position corresponding to Tyr₉₄₂, Leu₉₉₂, or Thr₁₅₅₈ of JS. Alternatively or additionally, attenuating mutations present in the JS HPIV3 *cp45* mutant strain are introduced in the N protein of chimeric PIV clones, for example which encode amino acid substitution(s) at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS. Yet additional useful attenuating mutations encode amino acid substitution(s) in the C protein, e.g., at a position corresponding to Ile₉₆ of JS and in the M protein, e.g., at a position corresponding to Pro₁₉₉ (for example a Pro₁₉₉ to Thr mutation). Other mutations identified in PIV3 JS *cp45* that can be adopted to adjust attenuation of a chimeric PIV of the invention are found in the F protein, e.g., at a position corresponding to Ile₄₂₀ or Ala₄₅₀ of JS, and in the HN protein, e.g., at a position corresponding to residue Val₃₈₄ of JS.

Attenuating mutations from biologically derived PIV mutants for incorporation into chimeric PIV of the invention also include mutations in noncoding

portions of the PIV genome or antigenome, for example in a 3' leader sequence. Exemplary mutations in this context may be engineered at a position in the 3' leader of a recombinant virus at a position corresponding to nucleotide 23, 24, 28, or 45 of JS *cp45*. Yet additional exemplary mutations may be engineered in the N gene start sequence, for example by
5 changing one or more nucleotides in the N gene start sequence, e.g., at a position corresponding to nucleotide 62 of JS *cp45*.

From PIV3 JS *cp45* and other biologically derived PIV mutants, a large "menu" of attenuating mutations is provided, each of which mutations can be combined with any other mutation(s) for finely adjusting the level of attenuation in chimeric PIV vaccine candidates of the invention. In exemplary embodiments, chimeric PIVs are constructed
10 which include one or more, and preferably two or more, mutations of HPIV3 JS *cp45*. Thus, chimeric PIVs of the invention selected for vaccine use often have two and sometimes three or more attenuating mutations from biologically derived PIV mutants or like model sources to achieve a satisfactory level of attenuation for broad clinical use. Preferably, these
15 attenuating mutations incorporated within recombinant chimeric PIVs of the invention are stabilized by multiple nucleotide substitutions in a codon specifying the mutation.

Introduction of attenuating and other desired phenotype-specifying mutations into a selected PIV vector, including chimeric bovine-human PIV vectors, may be achieved by transferring a heterologous gene or genome segment containing the mutation, e.g., a gene
20 encoding a mutant L protein, or portion thereof, into the PIV vector genome or antigenome. Alternatively, the mutation may be present in the selected vector genome or antigenome, and the introduced heterologous gene or genome segment may bear no mutations, or may bear one or more additional different mutations.

In certain examples, the vector genome or antigenome is modified at one or
25 more sites corresponding to a site of mutation in a heterologous "donor" virus (e.g., a heterologous bovine or human PIV or a non-PIV negative stranded RNA virus) to contain or encode the same, or a conservatively related, mutation (e.g., a conservative amino acid substitution) as a mutation identified in the donor virus (see, PCT/US00/09695 filed April 12, 2000 and its priority U.S. Provisional Patent Application Serial No. 60/129,006, filed
30 April 13, 1999, incorporated herein by reference). In one exemplary embodiment, a PIV vector genome or antigenome is modified at one or more sites corresponding to a site of

mutation in HPIV3 JS *cp45*, as enumerated above, to contain or encode the same or a conservatively related mutation as that identified in the *cp45* "donor."

Preferred mutant PIV strains for identifying and incorporating attenuating mutations into PIV vectors of the invention include cold passaged (*cp*), cold adapted (*ca*),
5 host range restricted (*hr*), small plaque (*sp*), and/or temperature sensitive (*ts*) mutants, for example the JS HPIV3 *cp45* mutant strain. Attenuating mutations from biologically derived PIV mutants for incorporation into human-bovine chimeric PIV of the invention also include mutations in noncoding portions of the PIV genome or antigenome, for example in a 3' leader sequence. Exemplary mutations in this context may be engineered at a position in the
10 3' leader of a recombinant virus at a position corresponding to nucleotide 23, 24, 28, or 45 of JS *cp45*. Yet additional exemplary mutations may be engineered in the N gene start sequence, for example by changing one or more nucleotides in the N gene start sequence, e.g., at a position corresponding to nucleotide 62 of JS *cp45*.

Additional mutations which can be adopted or transferred to PIV vectors of
15 the invention may be identified in non-PIV nonsegmented negative stranded RNA viruses and incorporated in PIV mutants of the invention. This is readily accomplished by mapping the mutation identified in a heterologous negative stranded RNA virus to a corresponding, homologous site in a recipient PIV genome or antigenome and mutating the existing sequence in the recipient to the mutant genotype (either by an identical or conservative
20 mutation), as described in PCT/US00/09695 filed April 12, 2000 and its priority U.S. Provisional Patent Application Serial No. 60/129,006, filed April 13, 1999, incorporated herein by reference. In accordance with this disclosure, additional attenuating mutations can be readily adopted or engineered within chimeric PIVs of the invention that are identified in other viruses, particularly other nonsegmented negative stranded RNA viruses.

25 In yet additional aspects of the invention, chimeric PIVs, with or without attenuating mutations modeled after biologically derived attenuated mutant viruses, are constructed to have additional nucleotide modification(s) to yield a desired phenotypic, structural, or functional change. Typically, the selected nucleotide modification will be made within the partial or complete PIV vector genome, but such modifications can be made
30 as well within any heterologous gene or genome segment that contributes to the chimeric clone. These modifications preferably specify a desired phenotypic change, for example a change in growth characteristics, attenuation, temperature-sensitivity, cold-adaptation,

plaque size, host range restriction, or immunogenicity. Structural changes in this context include introduction or ablation of restriction sites into PIV encoding cDNAs for ease of manipulation and identification.

In preferred embodiments, nucleotide changes within the genome or
5 antigenome of a chimeric PIV include modification of a viral gene by partial or complete deletion of the gene or reduction or ablation (knock-out) of its expression. Target genes for mutation in this context include any of the PIV genes, including the nucleocapsid protein N, phosphoprotein P, large polymerase subunit L, matrix protein M, hemagglutinin-neuraminidase protein HN, fusion protein F, and the products of the C, D and V open
10 reading frames (ORFs). To the extent that the recombinant virus remains viable and infectious, each of these proteins can be selectively deleted, substituted or rearranged, in whole or in part, alone or in combination with other desired modifications, to achieve novel deletion or knock out mutants. For example, one or more of the C, D, and/or V genes may be deleted in whole or in part, or its expression reduced or ablated (e.g., by introduction of a
15 stop codon, by a mutation in an RNA editing site, by a mutation that alters the amino acid specified by an initiation codon, or by a frame shift mutation in the targeted ORF(s)). In one embodiment, a mutation can be made in the editing site that prevents editing and ablates expression of proteins whose mRNA is generated by RNA editing (Kato et al., EMBO
16:578-587, 1997 and Schneider et al., Virology 227:314-322, 1997, incorporated herein by reference). Alternatively, one or more of the C, D, and/or V ORF(s) can be deleted in whole
20 or in part to alter the phenotype of the resultant recombinant clone to improve growth, attenuation, immunogenicity or other desired phenotypic characteristics (see, U.S. Patent Application Serial No. 09/350,821, filed by Durbin et al. on July 9, 1999, incorporated herein by reference).

Alternative nucleotide modifications in chimeric PIV of the invention include
25 a deletion, insertion, addition or rearrangement of a cis-acting regulatory sequence for a selected gene in the recombinant genome or antigenome. In one example, a cis-acting regulatory sequence of one PIV gene is changed to correspond to a heterologous regulatory sequence, which may be a counterpart cis-acting regulatory sequence of the same gene in a
30 different PIV, or a cis-acting regulatory sequence of a different PIV gene. For example, a gene end signal may be modified by conversion or substitution to a gene end signal of a different gene in the same PIV strain. In other embodiments, the nucleotide modification may comprise an insertion, deletion, substitution, or rearrangement of a translational start

site within the recombinant genome or antigenome, e.g., to ablate an alternative translational start site for a selected form of a protein.

In addition, a variety of other genetic alterations can be produced in a chimeric PIV genome or antigenome, alone or together with one or more attenuating mutations adopted from a biologically derived mutant PIV. For example, genes or genome segments from non-PIV sources may be inserted in whole or in part. In one such aspect, the invention provides methods for attenuating chimeric PIV vaccine candidates based on host range effects due to the introduction of one or more gene(s) or genome segment(s) from, e.g., a non-human PIV into a human PIV vector-based chimeric virus. For example, host range attenuation can be conferred on a HPIV-vector based chimeric construct by introduction of nucleotide sequences from a bovine PIV (BPIV) (see, e.g., as disclosed in U.S. Application Serial No. 09/586,479, filed June 1, 2000, corresponding to U.S. Provisional Application Serial No. 60/143,134 filed on July 9, 1999, incorporated herein by reference). These effects are attributed to structural and functional divergence between the vector and donor viruses and provide a stable basis for attenuation. For example, between HPIV3 and BPIV3 the percent amino acid identity for each of the N proteins is 86%, for P is 65%, M 93%, F 83%, HN 77%, and L 91%. All of these proteins are therefore candidates for introduction into a HPIV vector to yield an attenuated chimeric virus which cannot readily be altered by reversion. In exemplary embodiments, the vector genome or antigenome is an HPIV3 genome or antigenome and the heterologous gene or genome segment is a N ORF derived from a selected BPIV3 strain.

Thus, chimeric PIV are provided within the invention based on a vector genome or antigenome which is a human-bovine chimeric PIV genome or antigenome. In certain embodiments, the human-bovine chimeric vector genome or antigenome is combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of a heterologous pathogen selected from measles virus, subgroup A and subgroup B respiratory syncytial viruses, mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.

In alternate aspects of the invention, a human-bovine chimeric vector genome or antigenome comprises a partial or complete HPIV genome or antigenome combined with

one or more heterologous gene(s) or genome segment(s) from a BPIV. In one exemplary embodiment, a transcription unit comprising an open reading frame (ORF) of a BPIV3 N ORF is substituted in the vector genome or antigenome for a corresponding N ORF of a HPIV3 vector genome. Using this and similar constructs, the vector genome or antigenome is combined with a measles virus HA gene, or a selected antigenic determinant of another pathogen, as a supernumerary gene insert, as exemplified by the vaccine candidate identified below as rHPIV3-NB HAP-M.

In other alternate aspects of the invention, the human-bovine chimeric vector genome or antigenome comprises a partial or complete HPIV genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a BPIV. For example, one or more HPIV gene(s) or genome segment(s) encoding HN and/or F glycoproteins or one or more immunogenic domain(s), fragment(s) or epitope(s) thereof may be added to or incorporated within a partial or complete bovine genome or antigenome to form the vector genome or antigenome. In certain embodiments, both HPIV3 genes encoding HN and F glycoproteins are substituted for corresponding BPIV3 HN and F genes to form the vector genome or antigenome. Using this and similar constructs, the vector genome or antigenome is combined with a RSV F and/or G gene, or a selected antigenic determinant of another pathogen, as a supernumerary gene insert, as exemplified by the vaccine candidates identified below as rBHPiV3-G1 or rB/HPIV3-F1.

In yet more detailed embodiments, a chimeric human-bovine vector incorporates one or more HPIV1 HN and/or F gene(s) or genome segment(s) encoding one or more immunogenic domain(s), fragment(s) or epitope(s) thereof, and the vector is further modified by incorporation of one or more HPIV2 HN and/or F gene(s) or genome segment(s) encoding one or more immunogenic domain(s), fragment(s) or epitope(s) thereof to form the chimeric genome or antigenome which expresses protective antigen(s) from both HPIV1 and HPIV2. This category of chimeric PIV is exemplified by various vaccine candidates identified below as rB/HPIV3.1-2F; rB/HPIV3.1-2HN; or rB/HPIV3.1-2F,2HN.

In yet additional aspects of the invention, the order of genes can be changed to cause attenuation or reduce or enhance expression of a particular gene. Alternatively, a PIV genome promoter can be replaced with its antigenome counterpart to yield additional desired phenotypic changes. Different or additional modifications in the recombinant genome or antigenome can be made to facilitate manipulations, such as the insertion of

unique restriction sites in various intergenic regions or elsewhere. Nontranslated gene sequences can be removed to increase capacity for inserting foreign sequences.

5 In yet additional aspects, polynucleotide molecules or vectors encoding the chimeric PIV genome or antigenome can be modified to encode non-PIV sequences, e.g., a cytokine, a T-helper epitope, a restriction site marker, or a protein or immunogenic epitope of a microbial pathogen (e.g., virus, bacterium or fungus) capable of eliciting a protective immune response in an intended host. In one such embodiment, chimeric PIVs are constructed that incorporate a gene encoding a cytokine to yield novel phenotypic and immunogenic effects in the resulting chimera.

10 In addition to providing chimeric PIV for vaccine use, the invention provides related cDNA clones and vectors which incorporate a PIV vector genome or antigenome and heterologous polynucleotide(s) encoding one or more heterologous antigenic determinants, wherein the clones and vectors optionally incorporate mutations and related modifications specifying one or more attenuating mutations or other phenotypic changes as described
15 above. Heterologous sequences encoding antigenic determinants and/or specifying desired phenotypic changes are introduced in selected combinations, e.g., into an isolated polynucleotide which is a recombinant cDNA vector genome or antigenome, to produce a suitably attenuated, infectious virus or subviral particle in accordance with the methods described herein. These methods, coupled with routine phenotypic evaluation, provide a
20 large assemblage of chimeric PIVs having such desired characteristics as attenuation, temperature sensitivity, altered immunogenicity, cold-adaptation, small plaque size, host range restriction, genetic stability, etc. Preferred vaccine viruses among these candidates are attenuated and yet sufficiently immunogenic to elicit a protective immune response in the vaccinated mammalian host.

25 In related aspects of the invention, compositions (e.g., isolated polynucleotides and vectors incorporating a chimeric PIV-encoding cDNA) and methods are provided for producing an isolated infectious chimeric PIV. Included within these aspects of the invention are novel, isolated polynucleotide molecules and vectors incorporating such molecules that comprise a chimeric PIV genome or antigenome. Also provided is the same
30 or different expression vector comprising one or more isolated polynucleotide molecules encoding N, P, and L proteins. These proteins can alternatively be expressed directly from the genome or antigenome cDNA. The vector(s) is/are preferably expressed or coexpressed

in a cell or cell-free lysate, thereby producing an infectious chimeric parainfluenza virus particle or subviral particle.

The above methods and compositions for producing chimeric PIV yield infectious viral or subviral particles, or derivatives thereof. An infectious virus is comparable to the authentic PIV particle and is infectious as is. It can directly infect fresh cells. An infectious subviral particle typically is a subcomponent of the virus particle which can initiate an infection under appropriate conditions. For example, a nucleocapsid containing the genomic or antigenomic RNA and the N, P, and L proteins is an example of a subviral particle which can initiate an infection if introduced into the cytoplasm of cells. Subviral particles provided within the invention include viral particles which lack one or more protein(s), protein segment(s), or other viral component(s) not essential for infectivity.

The present invention provides a method for producing an infectious attenuated chimeric PIV particle from one or more isolated polynucleotide molecules encoding said PIV, comprising:

expressing in a cell or cell-free lysate an expression vector comprising an isolated polynucleotide comprising a partial or complete PIV vector genome or antigenome of a human or bovine PIV combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of one or more heterologous pathogen(s) to form a chimeric PIV genome or antigenome, and PIV N, P, and L proteins.

The present invention further provides an expression vector comprising an operably linked transcriptional promoter, a polynucleotide sequence which includes a partial or complete PIV vector genome or antigenome of a human or bovine PIV combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of one or more heterologous pathogen(s) to form a chimeric PIV genome or antigenome, and a transcriptional terminator.

In other embodiments the invention provides a cell or cell-free lysate containing an expression vector which comprises an isolated polynucleotide molecule comprising a chimeric PIV genome or antigenome as described above, and an expression vector (the same or different vector) which comprises one or more isolated polynucleotide

molecules encoding the N, P, and L proteins of PIV. One or more of these proteins also can be expressed from the genome or antigenome cDNA. Upon expression the genome or antigenome and N, P and L proteins combine to produce an infectious chimeric parainfluenza virus or subviral particle.

5 In other embodiments of the invention a cell or cell-free expression system (e.g., a cell-free lysate) is provided which incorporates an expression vector comprising an isolated polynucleotide molecule encoding a chimeric PIV, and an expression vector comprising one or more isolated polynucleotide molecules encoding N, P, and L proteins of a PIV. Upon expression, the genome or antigenome and N, P, and L proteins combine to
10 produce an infectious PIV particle, such as a viral or subviral particle.

The chimeric PIVs of the invention are useful in various compositions to generate a desired immune response against one or more PIVs, or against PIV and a non-PIV pathogen, in a host susceptible to infection therefrom. Chimeric PIV recombinants are capable of eliciting a mono- or poly-specific protective immune response in an infected mammalian host, yet are sufficiently attenuated so as to not cause unacceptable symptoms of
15 disease in the immunized host. The attenuated virus or subviral particle may be present in a cell culture supernatant, isolated from the culture, or partially or completely purified. The



20



virus may also be lyophilized, and can be combined with a variety of other components for storage or delivery to a host, as desired.

The invention further provides novel vaccines comprising a physiologically acceptable carrier and/or adjuvant and an isolated attenuated chimeric parainfluenza virus or subviral particle as described above. In preferred embodiments, the vaccine is comprised of a chimeric PIV having at least one, and preferably two or more additional mutations or other nucleotide modifications that specify a suitable balance of attenuation and immunogenicity. The vaccine can be formulated in a dose of 10^3 to 10^7 PFU of attenuated virus. The vaccine may comprise attenuated chimeric PIV that elicits an immune response against a single PIV strain or against multiple PIV strains or groups. In this regard, chimeric PIV can be combined in vaccine formulations with other PIV vaccine strains, or with other viral vaccine viruses such as RSV.

In related aspects, the invention provides a method for stimulating the immune system of an individual to elicit an immune response against one or more PIVs, or against PIV and a non-PIV pathogen, in a mammalian subject. The method comprises administering a formulation of an immunologically sufficient amount a chimeric PIV in a physiologically acceptable carrier and/or adjuvant. In one embodiment, the immunogenic composition is a vaccine comprised of a chimeric PIV having at least one, and preferably two or more attenuating mutations or other nucleotide modifications specifying a desired phenotype and/or level of attenuation as described above. The vaccine can be formulated in a dose of 10^3 to 10^7 PFU of attenuated virus. The vaccine may comprise an attenuated chimeric PIV that elicits an immune response against a single PIV, against multiple PIVs, e.g., HPIV1 and HPIV3, or against one or more PIV(s) and a non-PIV pathogen such as measles or RSV. In this context, chimeric PIVs can elicit a monospecific immune response or a polyspecific immune response against multiple PIVs, or against one or more PIV(s) and a non-PIV pathogen. Alternatively, chimeric PIV having different immunogenic characteristics can be combined in a vaccine mixture or administered separately in a coordinated treatment protocol to elicit more effective protection against one PIV, against multiple PIVs, or against one or more PIV(s) and a non-PIV pathogen such as measles or RSV. Preferably the immunogenic compositions of the invention are administered to the upper respiratory tract, e.g., by spray, droplet or aerosol. Preferably the immunogenic composition is administered to the upper respiratory tract, e.g., by spray, droplet or aerosol.

The present invention provides a method for sequential immunization to stimulate the immune system of an individual to induce protection against multiple pathogens comprising administering to a newborn to 4 month old infant an immunologically sufficient amount of a first attenuated chimeric HPIV expressing an antigenic determinant of a non-PIV pathogen and one or more antigenic determinants of HPIV3 and subsequently administering an immunologically sufficient amount of a second attenuated chimeric HPIV expressing an antigenic determinant of a non-PIV pathogen and one or more antigenic determinants of HPIV1 or HPIV2.

The invention also provides novel combinatorial vaccines and coordinate vaccination protocols for multiple pathogenic agents, including multiple PIV's and/or PIV and a non-PIV pathogen. For example, selected targets for early vaccination according to these compositions include RSV and PIV3, which each cause significant amount of illness within the first four months of life, whereas most of the illness caused by PIV1 and PIV2 occurs after six months of age (Collins et al., In *Fields Virology*, Vol. 1, pp. 1205-1243, Lippincott-Raven Publishers, Philadelphia, 1996; Reed et al., *J. Infect. Dis.* 175:807-13, 1997). A preferred immunization sequence employing live attenuated RSV and PIV vaccines is to administer RSV and PIV3 as early as one month of age (e.g., at one and two months of age) followed by a bivalent PIV1 and PIV2 vaccine at four and six months of age. It is thus desirable to employ the methods of the invention to administer multiple PIV vaccines, including one or more chimeric PIV vaccines, coordinately, e.g., simultaneously in a mixture or separately in a defined temporal sequence (e.g., in a daily or weekly sequence), wherein each vaccine virus preferably expresses a different heterologous protective antigen. Such a coordinate/sequential immunization strategy, which is able to induce secondary antibody responses to multiple viral respiratory pathogens, provides a highly powerful and extremely flexible immunization regimen that is driven by the need to immunize against each of the three PIV viruses and other pathogens in early infancy.

Importantly, the presence of multiple PIV serotypes and their unique epidemiology with PIV3 disease occurring at an earlier age than that of PIV1 and PIV2 makes it desirable to sequentially immunize an infant with different PIV vectors each expressing the same heterologous antigenic determinant such as the measles virus HA. This sequential immunization permits the induction of the high titer of antibody to the

heterologous protein that is characteristic of the secondary antibody response. In one embodiment, early infants (e. g. 2-4 month old infants) are immunized with an attenuated chimeric virus of the invention, for example a chimeric HPIV3 expressing the measles virus HA protein and also adapted to elicit an immune response against HPIV3, such as rcp45L (HA P-M). Subsequently, e. g., at four months of age the infant is again immunized but with a different, secondary vector construct, such as the rPIV3-1 cp45L virus expressing the measles virus HA gene and the HPIV1 antigenic determinants as the functional, obligate glycoproteins of the vector. Following the first vaccination, the vaccinee will elicit a primary antibody response to both the PIV3 HN and F proteins and to the measles virus HA protein, but not to the PIV 1 HN and F protein. Upon secondary immunization with the rPIV3-1 cp45L expressing the measles virus HA, the vaccinee will be readily infected with the vaccine because of the absence of antibody to the PIV 1 HN and F proteins and will develop both a primary antibody response to the PIV 1 HN and F protective antigens and a high titered secondary antibody response to the heterologous measles virus HA protein. A similar sequential immunization schedule can be developed where immunity is sequentially elicited against HPIV3 and then HPIV2 by one or more of the chimeric vaccine viruses disclosed herein, simultaneous with stimulation of an initial and then secondary, high titer protective response against measles or another non-PIV pathogen. This sequential immunization strategy, preferably employing different serotypes of PIV as primary and secondary vectors, effectively circumvents immunity that is induced to the primary vector, a factor ultimately limiting the usefulness of vectors with only one serotype. The success of sequential immunization with rPIV3 and rPIV3-1 virus vaccine candidates as described above has been demonstrated (Tao et al., Vaccine 17:1100-8, 1999).

The present invention provides an attenuated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a partial or complete PIV vector genome or antigenome, that is a HPIV3-1 vector genome or antigenome having both the HPIV3 HN and F ORFs substituted by the HN and F ORFs of HPIV 1 and further comprising an open reading frame (ORF) of a measles virus HA gene.

The present invention also provides an attenuated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid

phosphoprotein (P), a large polymerase protein (L), a partial or complete PIV vector genome or antigenome, and further comprising a translationally silent synthetic gene unit.

The present invention further provides an attenuated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of one or more heterologous pathogen(s) to form a chimeric PIV genome or antigenome,

wherein said one or more heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are added as supernumerary gene(s) or genome segment(s), and

wherein the addition of said one or more supernumerary heterologous gene(s) or genome segment(s) specifies an attenuation phenotype of the chimeric PIV which exhibits at least a 10- to 100-fold decrease in replication in the upper and/or lower respiratory tract of a host.

The present invention yet further provides an attenuated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of one or more heterologous pathogen(s) to form a chimeric PIV genome or antigenome,

wherein a transcription unit comprising an open reading frame (ORF) of a BPIV3 N ORF is substituted in the vector genome or antigenome for a corresponding N ORF of a HPIV3 vector genome.

The present invention provides an attenuated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of one or more heterologous pathogen(s) to form a chimeric PIV genome or antigenome,

wherein the vector genome or antigenome encodes a chimeric glycoprotein incorporating one or more antigenic domains, fragments, or epitopes of a glycoprotein of the vector PIV and of a glycoprotein of a heterologous PIV or non-PIV pathogen, and the chimeric PIV genome or antigenome is modified by introduction of an attenuating mutation involving an amino acid substitution of phenylalanine at position 456 of the HPIV3 L protein.

The present invention also provides an attenuated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of one or more heterologous pathogen(s) to form a chimeric PIV genome or antigenome,

wherein the chimeric PIV genome or antigenome is modified by introduction of an attenuating mutation involving an amino acid substitution of phenylalanine at position 456 of the HPIV3 L protein.

The present invention further provides an attenuated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of one or more heterologous pathogen(s) to form a chimeric PIV genome or antigenome,

wherein one or more PIV gene(s) in the vector genome or antigenome is deleted in whole or in part or expression of the gene(s) is reduced or ablated by a mutation in an RNA editing site, by a frameshift mutation, by a mutation that alters an amino acid specified by an initiation codon.

The present invention provides an isolated polynucleotide comprising a chimeric PIV genome or antigenome which includes a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) that is a transcriptionally silent synthetic gene unit to form a chimeric PIV genome or antigenome.

The present invention also provides an isolated polynucleotide comprising a chimeric PIV genome or antigenome which includes a partial or complete PIV vector

genome or antigenome combined with one or more heterologous gene(s) or genome segment(s),

wherein a transcription unit comprising an open reading frame (ORF) of a BPIV3 N ORF is substituted in the vector genome or antigenome for a corresponding N ORF of a HPIV3 vector genome.

The present invention further provides an isolated polynucleotide comprising a chimeric PIV genome or antigenome which includes a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of one or more heterologous pathogen(s) to form a chimeric PIV genome or antigenome,

wherein one or more PIV gene(s) is deleted in whole or in part or expression of the gene(s) is reduced or ablated by a mutation in an RNA editing site, by a frameshift mutation, by a mutation that alters an amino acid specified by an initiation codon.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A and 1B illustrate insertion of the HA gene of measles virus into the HPIV3 genome (Note: all of the figures presented herein and related descriptions refer to the positive-sense antigenome of HPIV3, 5' to 3').

Figure 1A provides a diagram (top; not to scale) of the 1926 nt insert containing the complete open reading frame of the hemagglutinin (HA) gene of the Edmonston wildtype strain of measles virus engineered to express the measles virus HA from an extra transcriptional unit. The insert contains, in 5' to 3' order: an *Afl*II site; nts 3699-3731 from the HPIV3 antigenome which contains the P/M gene junction, including downstream noncoding sequence for the P gene, its gene-end signal, the intergenic region, and the M gene-start signal; three additional nts (GCG); the complete measles virus HA ORF; HPIV3 nt 3594-3623 from the downstream noncoding region of the P gene; and a second *Afl*II site. Figure 1A, Panel 1 illustrates the complete antigenome of the JS wildtype strain of HPIV3 (rPIV3) with the introduced *Afl*II site in the 3'-noncoding region of the N gene before (top) and after (bottom) insertion of the measles HA ORF. Figure 1A, Panel 2 illustrates the complete antigenome of the JS wildtype strain of HPIV3 (rPIV3) with the introduced *Afl*II site in the 3'-noncoding region of the P gene before (top) and after (bottom) insertion of the measles HA ORF. SEQ ID NO. 1 and SEQ ID NO. 2 are shown in Fig. 1A.

Figure 1B provides a diagram (top; not to scale) of the 2028 nt insert containing the complete ORF of the HA gene of measles virus. The insert contains, in 5' to 3' order: a *StuI* site; nts 8602 to 8620 from the HPIV3 antigenome, which consist of downstream noncoding sequence from the HN gene and its gene-end signal; the conserved HPIV3 intergenic trinucleotide; nts 6733 to 6805 from the HPIV3 antigenome, which contains the HN gene-start and upstream noncoding region; the measles virus HA ORF; HPIV3 nts 8525-8597, which are downstream noncoding sequences from the HN gene; and a second *StuI* site. The construction is designed to, upon insertion, regenerate the HPIV3 HN gene containing the *StuI* site, and place the measles virus ORF directly after it flanked by the transcription signals and noncoding region of the HPIV3 HN gene. The complete antigenome of HPIV3 JS wildtype (rPIV3) with the introduced *StuI* site at nt position 8600 in the 3'-noncoding region of the HN gene is illustrated in the next (middle) diagram. Below is the antigenome of HPIV3 expressing the measles HA protein inserted into the *StuI* site. The HA cDNA used for this insertion came from an existing plasmid, rather than from the Edmonston wild type measles virus, which was used for the insertions in the N/P and P/M regions. This cDNA had two amino acid differences from the HA protein inserted in Fig 1A, and their location in the HA gene of measles virus is indicated by the asterisks in Figure 1B. SEQ ID NO. 3 and 4 are shown in Fig. 1B.

Figure 2 illustrates expression of the HA protein of measles virus by rHPIV3-measles virus-HA chimeric viruses in LLC-MK2 cells. The figure presents a radioimmunoprecipitation assay (RIPA) demonstrating that the measles HA protein is expressed by the recombinant chimeric viruses *rcp45L(HA P-M)* and *rcp45L(HA N-P)*, and by the Edmonston wild type strain of measles virus (Measles), but not by the rJS wild type HPIV3 (rJS). Lanes A--³⁵S-labeled infected cell lysates were immunoprecipitated by a mixture of three monoclonal antibodies specific to the HPIV3 HN protein). The 64kD band corresponding to the HN protein (open arrow) is present in each of the three HPIV3 infected cell lysates (lanes 3, 5, and 7), but not in the measles virus infected cell lysates (lane 9), confirming that the *rcp45L(HA P-M)* and *rcp45L(HA N-P)* chimeras are indeed HPIV3 and express similar levels of HN proteins. Lanes (b)--³⁵S-labeled infected cell lysates were immunoprecipitated by a mixture of monoclonal antibodies which recognizes the HA glycoprotein of measles virus (79-XV-V17, 80-III-B2, 81-1-366) (Hummel et al., *J. Virol.* 69:1913-6, 1995; Sheshberadaran et al., *Arch. Virol.* 83:251-68, 1985, each incorporated herein by reference). The 76kD band corresponding to the HA protein (closed arrow) is present in lysates from cells infected with the *rcp45L(HA)* chimeric viruses (lanes 6, 8) and

the measles virus (lane 10), but not in the lysates from rJS infected cells (lane 4), a HPIV3 wild type virus which does not encode a measles virus HA gene.

Figure 3 illustrates insertion of the HPIV2 HN gene as an extra transcription/translation unit into the antigenomic cDNA encoding rPIV3-1 or rPIV3-1cp45 chimeric virus (Note: rPIV3-1 is a rPIV3 in which the HN and F genes were replaced by those of HPIV1, and rPIV3-1cp45 is a version which contains, in addition, 12 mutations from the cp45 attenuated virus). The HPIV2 HN gene was amplified from vRNA of HPIV2 using RT-PCR with HPIV2 HN gene specific primers (Panel A). The amplified cDNA, carrying a primer-introduced *NcoI* site at its 5'-end and a *HindIII* site at its 3'-end, was digested with *NcoI-HindIII* and ligated into pLit.PIV31HNhc, that had been digested with *NcoI-HindIII*, to generate pLit.PIV32HNhc (Panel B). The pLit.PIV32HNhc plasmid was used as a template to produce a modified PIV2 HN cassette (Panel C), which has a *PpuMI* site at its 5'-end and an introduced *PpuMI* site at its 3'-end. This cassette contained, from left to right: the *PpuMI* site at the 5'-end, a partial 5'-untranslated region (UTR) of PIV3 HN, the PIV2 HN ORF, a 3'-UTR of PIV3 HN, the gene-end, intergenic, gene-start sequence that exists at the PIV3 HN and L gene junction, a portion of the 5'-untranslated region of PIV3 L, and the introduced *PpuMI* site at the 3'-end. This cDNA cassette was digested with *PpuMI* and then ligated to p38' Δ PIV31hc, that had been digested with *PpuMI*, to generate p38' Δ PIV31hc.2HN (Panel D). The 8.5 Kb *BspEI-SphI* fragment was assembled into the *BspEI-SphI* window of pFLC.2G+.hc or pFLCcp45 to generate the final full-length antigenomic cDNA, pFLC.3-1hc.2HN (Panel E) or pFLC.3-1hc.cp45.2HN (Panel F), respectively. pFLC.2G+.hc and pFLCcp45 are full-length antigenomic clones encoding wild type rPIV3-1 and rPIV3cp45, respectively, that have been described previously (Skiadopoulos et al., *J. Virol.* **73**:1374-81, 1999a; Tao et al., *J. Virol.* **72**:2955-2961, 1998, incorporated herein by reference).

Figure 4 details and verifies construction of the rPIV3-1.2HN chimeric virus carrying the PIV2 HN ORF insert between the PIV1 F and HN genes. Panel A depicts the differences in the structures of rPIV3-1 and rPIV3-1.2HN, which contains the PIV2 HN ORF insert between the PIV1 F and HN ORFs of rPIV3-1. The arrows indicate the approximate locations of the RT-PCR primers used to amplify fragments analyzed in Panels B-D. Panels B and C depict the expected sizes of the restriction enzyme digestion fragments generated from the RT-PCR products amplified from rPIV3-1 and rPIV3-1.2HN using either the *PpuMI* or *NcoI* restriction endonucleases, with the fragment sizes in base pairs (bp)

indicated, and the results presented in panel D. vRNA extracted from virus harvested from rPIV3-1.2HN or from rPIV3-1 infected LLC-MK2 cells was used as a template in the presence and absence of reverse transcriptase (RT) to amplify cDNA fragments by PCR using primers indicated in panel A. PCR fragments were absent in RT-PCR reactions

5 lacking RT indicating that the template employed for amplification of the DNA fragments was RNA and not contaminating cDNA (Lanes A and C of panel D). When the RT step was included, rPIV3-1.2HN vRNA (Lane B) yielded a fragment that was approximately 2kb larger than that of its rPIV3-1 parent (Lane D) indicating the presence of an insert of 2kb. Furthermore, digestion of this 3kb fragment with several different restriction endonucleases

10 indicated that the RT-PCR fragment from rPIV3-1.2HN (odd numbered lanes) has patterns that are different from those of the rPIV3-1 parent (even numbered lanes) for each restriction endonuclease tested. For each digestion, the number of sites and the sizes of the fragments obtained were completely consistent with the predicted sequence of the RT-PCR products of rPIV3-1 and rPIV3-1.2HN. Representative examples are presented. First, the *Pvu*MI

15 digestion of the RT-PCR product from rPIV3-1.2HN (Lane 1) produced three bands of the expected sizes indicating the presence of two *Pvu*MI sites and *Pvu*MI digestion of the RT-PCR product from rPIV3-1 produced two bands of the expected sizes for rPIV3-1 (Lane 2) indicating the presence of just one *Pvu*MI site. Second, the *Nco*I digestion of the RT-PCR product from rPIV3-1.2HN (Lane 5) produced 4 bands including the 0.5 kb fragment

20 indicative of the HPIV2 HN gene and the *Nco*I digestion of the RT-PCR product from rPIV3-1 (Lane 6) produced the expected two fragments. M identifies the lane containing the 1 kb DNA ladder used as nucleotide (nt) size markers (Life Technology). Similar results confirmed the presence of the HPIV2 HN insert in rPIV3-1cp45.2HN.

Figure 5 demonstrates that rPIV3-1.2HN expresses the HPIV2 HN protein.

25 LLC-MK2 monolayers were infected with rPIV3-1, rPIV3-1.2HN, or the PIV2/V94 wild type virus at a MOI of 5. Infected monolayers were incubated at 32°C and labeled with ³⁵S-met and ³⁵S-cys mixture from 18-36 hours post-infection. Cells were harvested and lysed, and the proteins were immunoprecipitated with anti-HPIV2 HN mAb 150S1 (Durbin et al., *Virology* 261:319-330, 1999; Tsurudome et al., *Virology* 171:38-48, 1989, incorporated

30 herein by reference) Immunoprecipitated samples were denatured, separated on a 4-12% SDS PAGE gel, and autoradiographed (Lanes: 1, rPIV3-1; 2, rPIV3-1.2HN; 3, PIV2/V9412-6). The mAb, specific to HPIV2 HN, precipitated a protein from both rPIV3-1.2HN and PIV2/V94 infected LLC-MK2 cells, but not from rPIV3-1-infected cells, with a size

expected for the 86kD Kd HN protein of HPIV2 (Rydbeck et al., *J. Gen. Virol.* 69:931-5, 1988, incorporated herein by reference).

Figure 6 depicts the location and construction of gene unit (GU) insertions or HN gene 3'-noncoding region (NCR) extensions. The nucleotide sequences and unique restriction enzyme cloning sites of the GU and NCR insertion sites are shown in panels A and B, respectively. Cis-acting transcriptional signal sequences, i.e., gene-end (GE), intergenic (IG), and gene-start (GS) signal sequences, are indicated. In Figure 6, Panel A, an oligonucleotide duplex specifying the HN GE, IG and GS signal sequences as well as the unique restriction enzyme recognition sequences are shown inserted into the introduced *StuI* restriction site (underlined nucleotides) (see Figure 1B and Example I for the location of the introduced *StuI* site). A restriction fragment from an RSV antigenome plasmid was cloned into the *HpaI* site. As necessary, a short oligonucleotide duplex was inserted into the *MluI* site of the multiple cloning site, so that the total length of the insert would conform to the rule of six. In Figure 6, Panel B, HN gene 3'-NCR insertions were cloned into the *HpaI* site of the indicated 32 nt multiple cloning site, which had been cloned into the *StuI* restriction site as described in Figure 6, Panel A. Inserted sequences were made to conform to the rule of six by insertion of short oligonucleotide duplexes into the *MluI* site in the multiple cloning site. SEQ ID NO. 5 and 6 are shown in Fig. 6.

Figure 7 illustrates open reading frames (ORFs) in the 3079 bp RSV insert. The six possible reading frames in the 3079 bp RSV fragment are shown (three in each orientation; 3, 2, 1, -1, -2, -3). Short bars represent translation start codons. Long bars represent translation stop codons. The 3079 bp fragment was inserted into the HN 3' NCR (NCR ins) or between the HN and L genes as a gene unit (GU ins) in such an orientation that the reading frames encountered by the PIV3 translation machinery correspond to -3, -2 and -1 in the figure. These reading frames contain numerous stop codons across the entire length of the sequence, and should therefore not produce any functional proteins.

Figure 8 demonstrates that rPIV3 insertion and extension mutants contain inserts of the appropriate size. RT-PCR was performed using a PIV3-specific primer pair flanking the insertion site, and RT-PCR products were separated by agarose gel electrophoresis. The expected size of the RT-PCR fragment for rPIV3wt (also referred to as rJS) is 3497 bp and that for each of the other rPIV3s GU or NCR mutants is increased in length depending on the size of the insertion. Panel A depicts GU insertion (ins) mutants: 1.

rHPIV3 wt; 2. r168 nt GU ins; 2. r678 nt GU ins; 3. r996 nt GU ins; 4. r1428 nt GU ins; 5. r1908 nt GU ins; 6. r3918 nt GU ins. M: *Hind*III restriction enzyme digestion products of lambda phage DNA. Sizes of relevant size markers are indicated. Panel B depicts NCR insertion mutants: 1. rPIV3 wt; 2. r258 nt NCR ins; 3. r972 nt NCR ins; 4. r1404 nt NCR ins; 5. r3126 nt NCR ins; 6. r3894 nt NCR ins. M: *Hind*III restriction enzyme digestion products of lambda phage DNA. Sizes of relevant size markers are indicated.

Figures 9A-9C present multi-step growth curves of GU and NCR insertion mutations compared with rHPIV3 wt and rcp45L. LLC-MK2 monolayers in 6-well plates were infected with each HPIV3 in triplicate at a multiplicity of infection (m.o.i.) of 0.01 and were washed 4 times after removal of the virus supernatant. At 0 hr and at 24 hrs intervals for 6 days post-infection, 0.5 ml virus medium from each well was harvested and 0.5 ml fresh medium was added to each well. Harvested samples were stored at -80°C. Virus present in the samples was quantified by titration on LLC-MK2 monolayers in 96-well plates incubated at 32°C. The titers of viruses are expressed as TCID₅₀/ml. The average of three independent infections from one experiment is shown. The lower limit of detection is 0.7 log₁₀TCID₅₀/ml. Figure 9A-GU insertion mutants; Figure 9B-NCR insertion mutants; Figure 9C-cp45L/GU insertion mutant.

Figure 10 illustrates the strategy for placing a supernumerary gene insert between the P and M genes of rHPIV3. The downstream (3') NCR of the rHPIV3 P gene was modified to contain an *Afl*III restriction site at antigenomic sequence positions 3693-3698 (Durbin, *J. Virol.* 74:6821-31, 2000, incorporated herein by reference). This site was then used to insert an oligonucleotide duplex (shown at the top) that contains HPIV3 cis-acting transcriptional signal sequences, i.e., gene-end (GE), intergenic (IG), and gene-start (GS) motifs. The duplex also contains a series of restriction enzyme recognition sequences available for insertion of foreign ORFs. In the case of the HPIV1 and HPIV2 HN ORFs, the cloning sites were *Nco*I and *Hind*III. Insertion of a foreign ORF into the multiple cloning sites places it under the control of a set of HPIV3 transcription signals, so that in the final recombinant virus the gene is transcribed into a separate mRNA by the HPIV3 polymerase. As necessary, a short oligonucleotide duplex was inserted into the *Mlu*I site of the multiple cloning site to adjust the final length of the genome to be an even multiple of six, which has been shown to be a requirement for efficient RNA replication (Calain et al., *J. Virol.* 67:4822-30, 1993; Durbin et al., *Virology* 234:74-83, 1997b). A similar strategy was used to

place HPIV1 and HPIV2 gene inserts between the N and P genes of rHPIV3 using an introduced *Afl*I restriction site at positions 1677-1682 9 (SEQ ID NO. 7).

Figure 11 is a diagram (not to scale) of the genomes of a series of chimeric rHPIV3s that contain one, two or three supernumerary gene inserts, each of which encodes a protective antigen of PIV1, PIV2, or measles virus. Schematic representation of rHPIV3s (not to scale) showing the relative position of the added insert(s) encoding the HN (hemagglutinin-neuraminidase) glycoprotein of HPIV1 (■) or HPIV2 (□) or the HA (hemagglutinin) glycoprotein of measles virus (▨) inserted into the rHPIV3 backbone (□). The rHPIV3 construct that is diagrammed at the bottom contains a 3918-nt insert (GU) that does not encode a protein (▨)(Skiadopoulos et al., *Virology* 272:225-34, 2000, incorporated herein by reference). Each foreign insert is under the control of a set of HPIV3 gene start and gene end transcription signals and is expressed as a separate mRNA. a. LLC-MK2 monolayers on 6 well plates (Costar) were separately infected in triplicate at an m.o.i. of 0.01 with each of the indicated viruses. Supernatants were harvested on days 5, 6 and 7 and virus was quantified as described previously (Skiadopoulos et al., *Virology* 272:225-34, 2000). The mean peak titer obtained for each virus is shown as log₁₀ TCID₅₀/ml. b. Mean of two experiments. Serially-diluted viruses were incubated at 32°C and 39°C on LLC-MK2 monolayer cultures for 7 days, and the presence of virus was determined by hemadsorption with guinea pig erythrocytes. The mean reduction in titer at 39°C compared to that of 32°C is shown.

Figure 12 provides a diagram (not to scale) illustrating insertion of a supernumerary gene insert into an rHPIV3 backbone, rHPIV3-N_B, in which the HPIV3 N ORF has been replaced by its BPIV3 counterpart, conferring an attenuation phenotype due to host range restriction (Bailly et al., *J. Virol.* 74:3188-3195, 2000a, incorporated herein by reference). Schematic representations are shown of rHPIV3 (top) and biologically derived BPIV3 (bottom). The relative position of the N ORF sequence derived from the Kansas strain of BPIV3 (■) and the measles virus hemagglutinin gene (▨) in the PIV3 backbone are shown. In each case, the foreign sequence is under the control of a set of HPIV3 transcription signals. A portion of the plasmid vector containing the NgoMIV site is shown (—/—). Designations are provided for the antigenomic cDNA clones (left) and their encoded recombinant viruses (right).

Figure 13 illustrates insertion of RSV G or F as an additional, supernumerary gene in a promoter-proximal position into the genome of rB/HPIV3. rB/HPIV3 is a recombinant version of BPIV3 in which the BPIV3 F and HN genes have been replaced by their HPIV3 counterparts (F_H and HN_H respectively). A *B*spI site was created in the B/HPIV3 backbone immediately upstream of the ATG start codon of the N ORF. The RSV G or F open reading frames (ORFs) were inserted into this *B*spI site. The downstream end of either RSV insert was designed to contain a PIV3 gene end (GE) and gene start (GS) sequences (AAGTAAGAAAAA (SEQ ID NO. 8) and AGGATTAAG, respectively, in positive sense) separated by the intergenic sequence CTT. Each insert also contained an *N*heI site that can serve as an insertion site for an additional supernumerary gene.

AGGATTAAGAAGCTTTACCGAAAGGTAAGGGGAAAGAAATCCTAAGAGCTTAGCGATG (SEQ ID NO. 9). GCTTAGCGATG (SEQ ID NO. 10). AAGCTAGCGCTTAGC (SEQ ID NO. 11). GCTTAGCAAAAAGCTAGCACAATG (SEQ ID NO. 12).

Figure 14 illustrates multicycle replication of rB/HPIV3-G1, rB/HPIV3-F1 and their recombinant and biological parent viruses in simian LLC-MK2 cells. Triplicate monolayer cultures were infected at an input MOI of 0.01 TCID₅₀ per cell with rB/HPIV3-G1, rB/HPIV3-F1, or the following control viruses: rBPIV3 Ka, which is the recombinant version of BPIV3 strain Ka; rB/HPIV3, which is the version of rBPIV3 in which the BPIV3 F and HN glycoprotein genes were replaced with their HPIV3 counterparts; HPIV3 JS, which is biologically-derived HPIV3 strain JS; and BPIV3 Ka, which is the biologically-derived version of BPIV3 strain Ka. The virus titers are shown as mean log₁₀ TCID₅₀/ml of triplicate samples. The lower limit of detection of this assay is 10^{1.45} TCID₅₀/ml.

Figure 15 is a diagram (not to scale) of the genomes of rBPIV3 (#1) and a series of chimeric rB/HPIV3s (#2-6) that contain substitutions of BPIV3 F and HN genes by those of HPIV3 (#2) or HPIV1 (#3-6), and one or two supernumerary gene inserts encoding the F and/or HN ORF of HPIV2 (#4-6). Schematic representation of the rB/HPIV3.1 chimeric viruses (not to scale) showing the relative position of the supernumerary gene encoding the F or HN glycoprotein of HPIV2 (F2 and HN2, respectively). Each foreign insert is under the control of a set of HPIV3 gene start and gene end transcription signals and is designed to be expressed as a separate mRNA.

Figure 16 provides a diagram (not to scale) illustrating the insertion of a the measles virus HA coding sequence into several different rPIV3 backbones. Three backbones

are illustrated: wild type rHPIV3 (top construct); wild type rHPIV3-1 (second construct from top) (Tao et al. *J. Virol.* 72:2955-2961, 1998, incorporated herein by reference) in which the HPIV3 F and HN glycoprotein genes have been replaced by those of HPIV1; and rHPIV3-1cp45L (third construct), a derivative of wild type rHPIV3-1 that contains three attenuating amino acid point mutations in the L gene derived from the cp45 vaccine strain (Skiadopoulos et al., *J. Virol.* 72:1762-8, 1998, incorporated herein by reference). The relative position of the HPIV1 F and HN ORF sequences () and the measles virus HA gene () in the rPIV3 backbone () are shown. In each case, each foreign ORF is under the control of a set of HPIV3 transcription signals. The relative locations of the three cp45 L amino acid point mutations in the L gene are indicated (*). A portion of the plasmid vector is containing the unique NgoMIV site is shown ().

Figure 17 illustrates construction of the PIV3-PIV2 chimeric antigenomic cDNA pFLC.PIV32hc encoding the full-length PIV2 HN and F proteins. The cDNA fragment containing the full-length PIV2 F ORF flanked by the indicated restriction sites (A1) was amplified from PIV2/V94 vRNA using RT-PCR and a PIV2 F specific primer pair (1, 2 in Table 22). This fragment was digested with NcoI plus BamHI (C1) and ligated to the NcoI-BamHI window of pLit.PIV31.fhc (B1) to generate pLit.PIV32Fhc (D1). In parallel, the cDNA fragment containing the full-length PIV2 HN ORF flanked by the indicated restriction sites (A2) was amplified from PIV2/V94 vRNA using RT-PCR and a PIV2 HN specific primer pair (3, 4 in Table 22). This fragment was digested with NcoI plus HindIII (C2) and ligated to the NcoI-HindIII window of pLit.PIV31.HNhc (B2) to generate pLit.PIV32HNhc (D2). pLit.PIV32Fhc and pLit.PIV32HNhc were digested with PpuMI and SpeI and assembled together to generate pLit.PIV32hc (E). pLit.PIV32hc was further digested with BspEI and SpeI and introduced into the BspEI-SpeI window of p38'ΔPIV31hc (F) to generate p38'ΔPIC32hc (G). The chimeric PIV3-PIV2 construct was introduced into the BspEI-SphI window of pFLC.2G+hc to generate pFLC.PIC32hc (H).

Figure 18 depicts construction of full-length PIV3-PIV2 chimeric antigenomic cDNA pFLC.PIV32TM and pFLC.PIV32TMcp45, which encode F and HN proteins containing PIV2-derived ectodomains and PIV3-derived transmembrane and cytoplasmic domains. The region of the PIV3 F ORF, in pLit.PIV3.F3a (A1), encoding the ectodomain was deleted (C1) by PCR using a PIV3 F specific primer pair (9, 10 in Table 22). The region of the PIV2 F ORF encoding the ectodomain was amplified from pLit.PIV32Fhc

(B1) using PCR and PIV2 F specific primer pair (5, 6 in Table 22). The two resulting fragments (C1 and D1) were ligated to generate pLit.PIV32FTM (E1). In parallel, the region of the PIV3 HN ORF, in pLit.PIV3.HN4 (A2), encoding the ectodomain was deleted (C2) by PCR using a PIV3 HN specific primer pair (11, 12 in Table 22). The region of the PIV2 HN ORF encoding the ectodomain was amplified from pLit.PIV32HNhc (B2) by PCR and a PIV2 HN specific primer pair (8, 9 in Table 22). Those two DNA fragments (C2 and D2) were ligated together to generate pLit.PIV32HNTM (E2). pLit.PIV32FTM and pLit.PIV32HNTM were digested with PpuMI and SpeI and assembled to generate pLit.PIV32TM (F). The BspEI-SpeI fragment from pLit.PIV32TM was ligated to the BspEI-SpeI window of p38'_PIV31hc (G) to generate p38'_PIV32TM (H). The insert containing chimeric PIV3-PIV2 F and HN was introduced as a 6.5 kb BspEI-SphI fragment into the BspEI-SphI window of pFLC.2G+.hc and pFLCcp45 to generate pFLC.PIV32TM and pFLC.PIV32TMcp45 (I), respectively.

Figure 19 shows construction of full-length PIV3-PIV2 chimeric antigenomic cDNA pFLC.PIV32CT and pFLC.PIV32Ctcp45 which encode F and HN proteins containing a PIV2-derived ectodomain, a PIV2-derived transmembrane domain, and a PIV3-derived cytoplasmic domain. The region of the PIV3 F ORF in pLit.PIV3.F3a (A1) encoding the ectodomain and the transmembrane domain was deleted (C1) by PCR using a PIV3 F specific primer pair (17, 18 in Table 22). The region of the PIV2 F ORF encoding the ectodomain plus the transmembrane domain was amplified from pLit.PIV32Fhc (B1) using PCR and a PIV2 F specific primer pair (13, 14 in Table 22). The two resulting fragments (C1 and D1) were ligated to generate pLit.PIV32FCT (E1). In parallel, the region of the PIV3 HN ORF in pLit.PIV3.HN4 (A2), encoding the ectodomain and transmembrane domain was deleted (C2) by PCR using a PIV3 HN specific primer pair (19, 20 in Table 22). The region of the PIV2 HN ORF encoding the ectodomain plus the transmembrane domain was amplified from pLit.PIV32HNhc (B2) by PCR using a PIV2 HN specific primer pair (15, 16 in Table 22). Those two DNA fragments (C2 and D2) were ligated to generate pLit.PIV32HNCT (E2). pLit.PIV32FCT and pLit.PIV32HNCT were digested with PpuMI and SpeI and assembled to generate pLit.PIV32CT (F). The BspEI-SpeI fragment from pLit.PIV32CT was ligated to the BspEI-SpeI window of p38'_PIV31hc (G) to generate p38'_PIV32CT (H). The insert containing chimeric PIV3-PIV2 F and HN was introduced as a 6.5 kb BspEI-SphI fragment into the BspEI-SphI window of pFLC.2G+.hc and pFLC.cp45 to generate pFLC.PIV32CT and pFLC.PIV32CTcp45 (I), respectively.

Figure 20 details genetic structures of the PIV3-PIV2 chimeric viruses and the gene junction sequences for rPIV3-2CT and rPIV3-2TM. Panel A illustrates the genetic structures of rPIV3-2 chimeric viruses (middle three diagrams) are compared with that of rPIV3 (top diagram) and rPIV3-1 (bottom diagram) viruses. The *cp45* derivatives are shown marked with arrows depicting the relative positions of *cp45* mutations. For the *cp45* derivatives, only the F and HN genes are different while the remaining genes remained identical, all from PIV3. From top to bottom, the three chimeric PIV3-PIV2 viruses carry decreasing amount of PIV3 glycoprotein genes. Note that rPIV3-2, carrying the complete PIV2 HN and F ORF, was not recoverable. Panel B provides the nucleotide sequences of the junctions of the chimeric F and HN glycoprotein genes for rPIV3-2TM are given along with the protein translation. The shaded portions represent sequences from PIV2. The amino acids are numbered with respect to their positions in the corresponding wild type glycoproteins. Three extra nucleotides were inserted in PIV3-PIV2 HN TM as indicated to make the construct conform to rule of six. Panel C shows the nucleotide sequences of the junctions of the chimeric F and HN glycoprotein genes for rPIV3-2CT, given along with the protein translation. The shaded portions represent sequences from PIV2. The amino acids are numbered with respect to their positions in the corresponding wild type glycoproteins. GE= gene end; I= intergenic; GS= gene start; ORF= open reading frame; TM= transmembrane domain; CT= cytoplasmic domain; *= stop codon.

Figure 21 documents multicycle replication of rPIV3-2 chimeric viruses compared with that of rPIV3/JS and PIV2/V94 wild type parent viruses. Panel A—the rPIV3-2TM and rPIV3-2TM*cp45* viruses, along with the rPIV3/JS and PIV2/V94 wt parent viruses, were used to infect LLC-MK2 cells in 6 well plates, each in triplicate, at an MOI of 0.01. All cultures were incubated at 32°C. After a 1 hour adsorption period, the inocula were removed, and the cells were washed three times with serum-free OptiMEM. The cultures were overlaid with 2 ml per well of the same medium. For rPIV3-2TM and rPIV3-2TM*cp45* infected plates, 0.5 mg/ml of p-trypsin was added to each well. Aliquots of 0.5 ml were taken from each well at 24 hour intervals for 6 days, flash frozen on dry ice, and stored at -80°C. Each aliquot was replaced with 0.5 ml of fresh medium with or without p-trypsin as indicated above. The virus present in the aliquots was titered on LLC-MK2 plates with liquid overlay at 32°C for 7 days, and the endpoints were identified with hemadsorption. Panel B—The rPIV3-2CT and rPIV3-2CT*cp45*, along with the rPIV3/JS and PIV2/V94 wt parent viruses, were used to infect LLC-MK2 in 6 well plates, each in

triplicate, as described in Panel A. Aliquots were taken and processed in the same manner as described in Panel A. Virus titers are expressed as log₁₀TCID₅₀/ml ± standard errors for both experiments presented in Panel A and B.

5 DESCRIPTION OF THE SPECIFIC EMBODIMENTS

The instant invention provides methods and compositions for the production and use of novel, chimeric parainfluenza viruses (PIVs) and associated vaccines. The chimeric viruses of the invention are infectious and immunogenic in humans and other mammals and are useful for generating immune responses against one or more PIVs, for example against one or more human PIVs (HPIVs). Alternatively, chimeric PIVs are provided that elicit an immune response against a selected PIV and one or more additional pathogens, for example against both a HPIV and measles virus. The immune response elicited can involve either or both humoral and/or cell mediated responses. Preferably, chimeric PIVs of the invention are attenuated to yield a desired balance of attenuation and immunogenicity for vaccine use.

The invention thus provides novel methods for designing and producing attenuated, chimeric PIVs that are useful as vaccine agents for preventing and/or treating infection and related disease symptoms attributable to PIV and other pathogens. In accordance with the methods of the invention, chimeric parainfluenza viruses or subviral particles are constructed using a PIV "vector" genome or antigenome that is recombinantly modified to incorporate one or more antigenic determinants of a heterologous pathogen. The vector genome or antigenome is comprised of a partial or complete PIV genome or antigenome, which may itself incorporate nucleotide modifications such as attenuating mutations. The vector genome or antigenome is modified to form a chimeric structure through incorporation of a heterologous gene or genome segment. More specifically, chimeric PIVs of the invention are constructed through a cDNA-based virus recovery system that yields recombinant viruses that incorporate a partial or complete vector or "background" PIV genome or antigenome combined with one or more "donor" nucleotide sequences encoding the heterologous antigenic determinant(s). Preferably the PIV vector comprises a HPIV genome or antigenome, although non-human PIVs, for example a bovine PIV (BPIV), can be employed as a vector to incorporate antigenic determinants of human PIVs and other human pathogens. In exemplary embodiments described herein, a human PIV3 (HPIV3)

vector genome or antigenome is modified to incorporate one or more genes or genome segments that encode antigenic determinant(s) of one or more heterologous PIVs (e.g., HPIV1 and/or HPIV2), and/or a non-PIV pathogen (e.g., measles virus). Thus constructed, chimeric PIVs of the invention may elicit an immune response against a specific PIV, e.g., HPIV1, HPIV2, and/or HPIV3, or against a non-PIV pathogen. Alternatively, compositions and methods are provided for eliciting a polyspecific immune response against multiple PIVs, e.g., HPIV1 and HPIV3, or against one or more HPIVs and a non-PIV pathogen such as measles virus.

Exemplary chimeric PIV of the invention incorporate a chimeric PIV genome or antigenome as described above, as well as a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), and a large polymerase protein (L). Additional PIV proteins may be included in various combinations to provide a range of infectious subviral particles, up to a complete viral particle or a viral particle containing supernumerary proteins, antigenic determinants or other additional components. Additional PIV proteins may be included in various combinations to provide a range of infectious subviral particles, up to a complete viral particle or a viral particle containing supernumerary proteins, antigenic determinants or other additional components.

In preferred aspects of the invention, chimeric PIV incorporate a partial or complete human PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a second human PIV or a non-PIV pathogen such as measles virus. The PIV "vector" genome or antigenome typically acts as a recipient or carrier to which are added or incorporated one or more "donor" genes or genome segments of a heterologous pathogen. Typically, polynucleotides encoding one or more antigenic determinants of the heterologous pathogen are added to or substituted within the vector genome or antigenome to yield a chimeric PIV that thus acquires the ability to elicit an immune response in a selected host against the heterologous pathogen. In addition, the chimeric virus may exhibit other novel phenotypic characteristics compared to one or both of the vector PIV and heterologous pathogens.

The partial or complete vector genome or antigenome generally acts as a backbone into which heterologous genes or genome segments of a different pathogen are incorporated. Often, the heterologous pathogen is a different PIV from which one or more gene(s) or genome segment(s) is/are of are combined with, or substituted within, the vector

genome or antigenome. In addition to providing novel immunogenic characteristics, the addition or substitution of heterologous genes or genome segments within the vector PIV strain may confer an increase or decrease in attenuation, growth changes, or other desired phenotypic changes as compared with the corresponding phenotype(s) of the unmodified vector and donor viruses. Heterologous genes and genome segments from other PIVs that may be selected as inserts or additions within chimeric PIV of the invention include genes or genome segments encoding the PIV N, P, C, D, V, M, F, HN and/or L protein(s) or one or more antigenic determinant(s) thereof.

Heterologous genes or genome segments of one PIV may be added as a supernumerary genomic element to a partial or complete genome or antigenome of a different PIV. Alternatively, one or more heterologous gene(s) or genome segment(s) of one PIV may be substituted at a position corresponding to a wild-type gene order position of a counterpart gene(s) or genome segment(s) that is deleted within the PIV vector genome or antigenome. In yet additional embodiments, the heterologous gene or genome segment is added or substituted at a position that is more promoter-proximal or promoter-distal compared to a wild-type gene order position of the counterpart gene or genome segment within the vector genome or antigenome to enhance or reduce, respectively, expression of the heterologous gene or genome segment.

The introduction of heterologous immunogenic proteins, protein domains and immunogenic epitopes to produce chimeric PIV is particularly useful to generate novel immune responses in an immunized host. Addition or substitution of an immunogenic gene or genome segment from one, donor pathogen within a recipient PIV vector genome or antigenome can generate an immune response directed against the donor pathogen, the PIV vector, or against both the donor pathogen and vector.

To achieve this purpose, chimeric PIV may be constructed that express a chimeric protein, for example an immunogenic glycoprotein having a cytoplasmic tail and/or transmembrane domain specific to a vector fused to a heterologous ectodomain of a different PIV or non-PIV pathogen to provide a fusion protein that elicits an immune response against the heterologous pathogen. For example, a heterologous genome segment encoding a glycoprotein ectodomain from a human PIV1 HN or F glycoprotein may be joined with a genome segment encoding the corresponding HPIV3 HN or F glycoprotein cytoplasmic and

transmembrane domains to form a HPIV3-1 chimeric glycoprotein that elicits an immune response against HPIV1.

Briefly, PIV of the invention expressing a chimeric glycoprotein comprise a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a HPIV vector genome or antigenome that is modified to encode a chimeric glycoprotein. The chimeric glycoprotein incorporates one or more heterologous antigenic domains, fragments, or epitopes of a second, antigenically distinct HPIV. Preferably, this is achieved by substitution within the HPIV vector genome or antigenome of one or more heterologous genome segments of the second HPIV that encode one or more antigenic domains, fragments, or epitopes, whereby the genome or antigenome encodes the chimeric glycoprotein that is antigenically distinct from the parent, vector virus.

In more detailed aspects, the heterologous genome segment or segments preferably encode a glycoprotein ectodomain or immunogenic portion or epitope thereof, and optionally include other portions of the heterologous or "donor" glycoprotein, for example both an ectodomain and transmembrane region that are substituted for counterpart glycoprotein ecto- and transmembrane domains in the vector genome or antigenome. Preferred chimeric glycoproteins in this context may be selected from HPIV HN and/or F glycoproteins, and the vector genome or antigenome may be modified to encode multiple chimeric glycoproteins. In preferred embodiments, the HPIV vector genome or antigenome is a partial HPIV3 genome or antigenome and the second, antigenically distinct HPIV is either HPIV1 or HPIV2. In one exemplary embodiment described below, both glycoprotein ectodomain(s) of HPIV2 HN and F glycoproteins are substituted for corresponding HN and F glycoprotein ectodomains in the HPIV3 vector genome or antigenome. In another exemplary embodiment, PIV2 ectodomain and transmembrane regions of one or both HN and/or F glycoproteins are fused to one or more corresponding PIV3 cytoplasmic tail region(s) to form the chimeric glycoprotein. Further details concerning these aspects of the invention are provided in United States Patent Application entitled CONSTRUCTION AND USE OF RECOMBINANT PARAINFLUENZA VIRUSES EXPRESSING A CHIMERIC GLYCOPROTEIN, filed on December 10, 1999 by Tao et al. and identified by Attorney Docket No. 17634-000340, incorporated herein by reference.

To construct chimeric PIVs of the invention carrying a heterologous antigenic determinant of a non-PIV pathogen, a heterologous gene or genome segment of the donor

pathogen may be added or substituted at any operable position in the vector genome or antigenome. In one embodiment, heterologous genes or genome segments from a non-PIV pathogen can be added (i.e., without substitution) within a PIV vector genome or antigenome to create novel immunogenic properties within the resultant clone. In these cases, the

5 heterologous gene or genome segment may be added as a supernumerary gene or genome segment, optionally for the additional purpose of attenuating the resultant chimeric virus, in combination with a complete PIV vector genome or antigenome. Alternatively, the heterologous gene or genome segment may be added in conjunction with deletion of a selected gene or genome segment in the vector genome or antigenome.

10 In preferred embodiments of the invention, the heterologous gene or genome segment is added at an intergenic position within the partial or complete PIV vector genome or antigenome. Alternatively, the gene or genome segment can be inserted within other noncoding regions of the genome, for example, within 5' or 3' noncoding regions or in other positions where noncoding nucleotides occur within the vector genome or antigenome. In

15 one aspect, the heterologous gene or genome segment is inserted at a non-coding site overlapping a cis-acting regulatory sequence within the vector genome or antigenome, e.g., within a sequence required for efficient replication, transcription, and/or translation. These regions of the vector genome or antigenome represent target sites for disruption or modification of regulatory functions associated with introduction of the heterologous gene or

20 genome segment.

As used herein, the term "gene" generally refers to a portion of a subject genome, e.g., a PIV genome, encoding an mRNA and typically begins at the upstream end with a gene-start (GS) signal and ends at the downstream end with the gene-end (GE) signal. The term gene is also interchangeable with the term "translational open reading frame", or

25 ORF, particularly in the case where a protein, such as the PIV C protein, is expressed from an additional ORF rather than from a unique mRNA. In the exemplary case of HPIV3, the genome is a single strand of negative-sense RNA 15462 nucleotides (nt) in length (Galinski et al., *Virology* 165: 499-510, 1988; Stokes et al., *Virus Res.* 25:91-103, 1992). At least eight proteins are encoded by the HPIV3 genome: the nucleocapsid protein N, the

30 phosphoprotein P, the C and D proteins of unknown functions, the matrix protein M, the fusion glycoprotein F, the hemagglutinin-neuraminidase glycoprotein HN, and the large polymerase protein L (Collins et al., 3rd ed. In "Fields Virology," B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, B. Roizman, and S. E. Straus,

Eds., Vol. 1, pp. 1205-1243. Lippincott-Raven Publishers, Philadelphia, 1996). The viral genome of all PIVs also contains extragenic leader and trailer regions, possessing all or part of the promoters required for viral replication and transcription, as well as non-coding and intergenic regions. Thus, the PIV genetic map is represented as 3' leader-N-P/C/D/V-M-F-HN-L-5' trailer. Transcription initiates at the 3' end and proceeds by a sequential stop-start mechanism that is guided by short conserved motifs found at the gene boundaries. The upstream end of each gene contains a gene-start (GS) signal, which directs initiation of its respective mRNA. The downstream terminus of each gene contains a gene-end (GE) motif which directs polyadenylation and termination. Exemplary genome sequences have been described for the human PIV3 strains JS (GenBank accession number Z11575, incorporated herein by reference) and Washington (Galinski M.S. In Kingsbury, D.W. (Ed.), The Paramyxoviruses, pp. 537-568, Plenum Press, New York, 1991, incorporated herein by reference), and for the bovine PIV3 strain 910N (GenBank accession number D80487, incorporated herein by reference).

To construct chimeric PIVs of the invention, one or more PIV gene(s) or genome segment(s) may be deleted, inserted or substituted in whole or in part. This means that partial or complete deletions, insertions and substitutions may include open reading frames and/or cis-acting regulatory sequences of any one or more of the PIV genes or genome segments. By "genome segment" is meant any length of continuous nucleotides from the PIV genome, which might be part of an ORF, a gene, or an extragenic region, or a combination thereof. When a subject genome segment encodes an antigenic determinant, the genome segment encodes at least one immunogenic epitope capable of eliciting a humoral or cell mediated immune response in a mammalian host. The genome segment may also encode an immunogenic fragment or protein domain. In other aspects, the donor genome segment may encode multiple immunogenic domains or epitopes, including recombinantly synthesized sequences that comprise multiple, repeating or different, immunogenic domains or epitopes.

Alternative chimeric PIV of the invention will contain protective antigenic determinants of HPIV1, HPIV2 and/or HPIV3. This is preferably achieved by expression of one or more HN and/or F genes or genome segments by the vector PIV, or as extra or substitute genes from the heterologous donor pathogen. In certain embodiments, a HPIV3-1 or HPIV3-2 chimeric virus may be constructed for use as a vaccine or vector strain, in which the HPIV1 or HPIV2 HN and/or F genes replace their PIV3 counterpart(s) (Skiadopoulos et

al., Vaccine 18:503-510, 1999; Tao et al., Vaccine 17:1100-1108, 1999; U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998 (and corresponding International Application published as WO 98/53078); U.S. Patent Application Serial No. 09/458,813, filed December 10, 1999; U.S. Patent Application Serial No. 09/459,062, filed December 10, 5 1999; each incorporated herein by reference). In this context, a chimeric PIV1 vaccine candidate has been generated using the PIV3 cDNA rescue system by replacing the PIV3 HN and F open reading frames (ORFs) with those of PIV1 in a PIV3 full-length cDNA that contains the three attenuating mutations in L. The recombinant chimeric virus derived from this cDNA is designated rPIV3-1.*cp45L* (Skiadopoulos et al., J. Virol. 72:1762-8, 1998; Tao 10 et al., J. Virol. 72:2955-2961, 1998; Tao et al., Vaccine 17:1100-1108, 1999, incorporated herein by reference). rPIV3-1.*cp45L* is attenuated in hamsters and induced a high level of resistance to challenge with PIV1. A recombinant chimeric virus, designated rPIV3-1.*cp45*, has also been produced that contains 12 of the 15 *cp45* mutations, i.e., excluding the mutations in HN and F, and is highly attenuated in the upper and lower respiratory tract of 15 hamsters (Skiadopoulos et al., Vaccine 18:503-510, 1999, incorporated herein by reference).

In preferred embodiments of the invention, the chimeric PIV bear one or more major antigenic determinants of a human PIV, or against multiple human PIVs, including HPIV1, HPIV2 or HPIV3. These preferred vaccine candidates elicit an effective immune response in humans against one or more selected HPIVs. As noted above, the 20 antigenic determinant(s) that elicit(s) an immune response against HPIV may be encoded by the vector genome or antigenome, or may be inserted within or joined to the PIV vector genome or antigenome as a heterologous gene or gene segment. The major protective antigens of human PIVs are their HN and F glycoproteins. However, all PIV genes are candidates for encoding antigenic determinants of interest, including internal protein genes 25 which may encode such determinants as, for example, CTL epitopes.

Preferred chimeric PIV vaccine viruses of the invention bear one or more major antigenic determinants from each of a plurality of HPIVs or from a HPIV and a non-PIV pathogen. Chimeric PIV thus constructed include a partial or complete HPIV genome or antigenome, for example of HPIV3, and one or more heterologous gene(s) or genome 30 segment(s) encoding antigenic determinant(s) of a heterologous PIV, for example HPIV1 or HPIV2. In alternative embodiments, one or more genes or genome segments encoding one or more antigenic determinants of HPIV1 or HPIV2 may be added to or substituted within a partial or complete HPIV3 genome or antigenome. In various exemplary embodiments

described below, both HPIV1 genes encoding the HN and F glycoproteins are substituted for counterpart HPIV3 HN and F genes in a chimeric PIV vaccine candidate. These and other constructs yield chimeric PIVs that elicit either a mono- or poly-specific immune response in humans to one or more HPIVs. Further detailed aspects of the invention are provided in

5 United States Patent Application entitled CONSTRUCTION AND USE OF RECOMBINANT PARAINFLUENZA VIRUSES EXPRESSING A CHIMERIC GLYCOPROTEIN, filed on December 10, 1999 by Tao et al. and identified by Attorney Docket No. 17634-000340, and U.S. Patent Application entitled USE OF RECOMBINANT PARAINFLUENZA VIRUS (PIV) AS A VECTOR TO PROTECT AGAINST DISEASE

10 CAUSED BY PIV AND RESPIRATORY SYNCYTIAL VIRUS (RSV), filed on December 10, 1999 by Murphy et al. and identified by Attorney Docket No. 17634-000330, each incorporated herein by reference.

In exemplary aspects of the invention, heterologous genes or genome segments encoding antigenic determinants from both HPIV1 and HPIV2 are added to or

15 incorporated within a partial or complete HPIV3 vector genome or antigenome. For instance, one or more HPIV1 genes or genome segments encoding HN and/or F glycoproteins, or antigenic determinant(s) thereof, and one or more HPIV2 genes or genome segments encoding HN and/or F glycoproteins or antigenic determinants can be added to or incorporated within a partial or complete HPIV3 vector genome or antigenome. In one

20 example described below, both HPIV1 genes encoding HN and F glycoproteins are substituted for counterpart HPIV3 HN and F genes to form a chimeric HPIV3-1 vector genome or antigenome. This vector construct can be further modified by addition or incorporation of one or more genes or gene segments encoding antigenic determinant(s) of HPIV2. Thus, specific constructs exemplifying the invention are provided which yield

25 chimeric PIVs having antigenic determinants of both HPIV1 and HPIV2, as exemplified by the vaccine candidates rPIV3-1.2HN and rPIV3-1cp45.2HN described herein below.

In other preferred aspects of the invention, chimeric PIV incorporate a HPIV vector genome or antigenome modified to express one or more major antigenic determinants of non-PIV pathogen, for example measles virus. The methods of the invention are

30 generally adaptable for incorporation of antigenic determinants from a wide range of additional pathogens within chimeric PIV vaccine candidates. In this regard the invention also provides for development of vaccine candidates against subgroup A and subgroup B respiratory syncytial viruses (RSV), mumps virus, human papilloma viruses, type 1 and type

2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses, among other pathogens. In this regard, pathogens that may be targeted for vaccine development according to the methods of the invention include viral and bacterial pathogens, as well as protozoans and multicellular pathogens. Useful antigenic determinants from many important human pathogens in this context are known or readily identified for incorporation within chimeric PIV of the invention. Thus, major antigens have been identified for the foregoing exemplary pathogens, including the measles virus HA and F proteins; the F, G, SH and M2 proteins of RSV, mumps virus HN and F proteins, human papilloma virus L1 protein, type 1 or type 2 human immunodeficiency virus gp160 protein, herpes simplex virus and cytomegalovirus gB, gC, gD, gE, gG, gH, gI, gJ, gK, gL, and gM proteins, rabies virus G protein, Epstein Barr Virus gp350 protein; filovirus G protein, bunyavirus G protein, flavivirus E and NS1 proteins, and alphavirus E. These major antigens, as well as other antigens known in the art for the enumerated pathogens and others, are well characterized to the extent that many of their antigenic determinants, including the full length proteins and their constituent antigenic domains, fragments and epitopes, are identified, mapped and characterized for their respective immunogenic activities.

Among the numerous, exemplary mapping studies that identify and characterize major antigens of diverse pathogens for use within the invention are epitope mapping studies directed to the hemagglutinin-neuraminidase (HN) gene of HPIV3 (van Wyke Coelingh et al., *J. Virol.* 61:1473-1477, 1987, incorporated herein by reference). This report provides detailed antigenic structural analyses for 16 antigenic variants of HPIV3 variants selected by using monoclonal antibodies (MAbs) to the HN protein which inhibit neuraminidase, hemagglutination, or both activities. Each variant possessed a single-point mutation in the HN gene, coding for a single amino acid substitution in the HN protein. Operational and topographic maps of the HN protein correlated well with the relative positions of the substitutions. Computer-assisted analysis of the HN protein predicted a secondary structure composed primarily of hydrophobic β sheets interconnected by random hydrophilic coil structures. The HN epitopes were located in predicted coil regions. Epitopes recognized by MAbs which inhibit neuraminidase activity of the virus were located in a region which appears to be structurally conserved among several paramyxovirus HN proteins and which may represent the sialic acid-binding site of the HN molecule.

This exemplary work, employing conventional antigenic mapping methods, identified single amino acids which are important for the integrity of HN epitopes. Most of these epitopes are located in the C-terminal half of the molecule, as expected for a protein anchored at its N terminus (Elango et al., *J. Virol.* 57:481-489, 1986). Previously published operational and topographic maps of the PIV3 HN indicated that the MAbs employed recognized six distinct groups of epitopes (I to VI) organized into two topographically separate sites (A and B), which are partially bridged by a third site (C). These groups of epitopes represent useful candidates for antigenic determinants that may be incorporated, alone or in various combinations, within chimeric PIVs of the invention. (See, also, Coelingh et al., *Virology* 143:569-582, 1985; Coelingh et al., *Virology* 162:137-143, 1988; Ray et al., *Virology* 148:232-236, 1986; Rydbeck et al., *J. Gen. Virol.* 67:1531-1542, 1986, each incorporated herein by reference),

Additional studies by van Wyke Coelingh et al. (*J. Virol.* 63:375-382, 1989) provide further information relating to selection of PIV antigenic determinants for use within the invention. In this study, twenty-six monoclonal antibodies (MAbs) (14 neutralizing and 12 nonneutralizing) were used to examine the antigenic structure, biological properties, and natural variation of the fusion (F) glycoprotein of HPIV3. Analysis of laboratory-selected antigenic variants and of PIV3 clinical isolates indicated that the panel of MAbs recognizes at least 20 epitopes, 14 of which participate in neutralization. Competitive binding assays confirmed that the 14 neutralization epitopes are organized into three nonoverlapping principal antigenic regions (A, B, and C) and one bridge site (AB), and that the 6 nonneutralization epitopes form four sites (D, E, F, and G). Most of the neutralizing MAbs were involved in nonreciprocal competitive binding reactions, suggesting that they induce conformational changes in other neutralization epitopes.

Other antigenic determinants for use within the invention have been identified and characterized for respiratory syncytial virus (RSV). For example, Beeler et al., *J. Virol.* 63:2941-2950, 1989, incorporated herein by reference, employed eighteen neutralizing monoclonal antibodies (MAbs) specific for the fusion glycoprotein of the A2 strain of RSV to construct a detailed topological and operational map of epitopes involved in RSV neutralization and fusion. Competitive binding assays identified three nonoverlapping antigenic regions (A, B, and C) and one bridge site (AB). Thirteen MAb-resistant mutants (MARMs) were selected, and the neutralization patterns of the MAbs with either MARMs or RSV clinical strains identified a minimum of 16 epitopes. MARMs selected with antibodies

to six of the site A and AB epitopes displayed a small-plaque phenotype, which is consistent with an alteration in a biologically active region of the F molecule. Analysis of MARMs also indicated that these neutralization epitopes occupy topographically distinct but conformationally interdependent regions with unique biological and immunological properties. Antigenic variation in F epitopes was then examined by using 23 clinical isolates (18 subgroup A and 5 subgroup B) in cross-neutralization assays with the 18 anti-F MAbs. This analysis identified constant, variable, and hypervariable regions on the molecule and indicated that antigenic variation in the neutralization epitopes of the RSV F glycoprotein is the result of a noncumulative genetic heterogeneity. Of the 16 epitopes, 8 were conserved on all or all but 1 of 23 subgroup A or subgroup B clinical isolates. These antigenic determinants, including the full length proteins and their constituent antigenic domains, fragments and epitopes, all represent useful candidates for integration within chimeric PIV of the invention to elicit novel immune responses as described above. (See also, Anderson et al., *J. Infect. Dis.* 151:626-633, 1985; Coelingh et al., *J. Virol.* 63:375-382, 1989; Fenner et al., *Scand. J. Immunol.* 24:335-340, 1986; Fernie et al., *Proc. Soc. Exp. Biol. Med.* 171:266-271, 1982; Sato et al., *J. Gen. Virol.* 66:1397-1409, 1985; Walsh et al., *J. Gen. Virol.* 67:505-513, 1986, and Olmsted et al., *J. Virol.* 63:411-420, 1989, each incorporated herein by reference).

To express antigenic determinants of heterologous PIVs and non-PIV pathogens, the invention provides numerous human and non-human PIV vectors, including bovine PIV (BPIV) vectors. These vectors are readily modified according to the recombinant methods described herein to carry heterologous antigenic determinants and elicit one or more specific humoral or cell mediated immune responses against the heterologous pathogen and vector PIV. In exemplary embodiments, one or more heterologous genes or genome segments from a donor pathogen is combined with a HPIV3 vector genome or antigenome. In other exemplary embodiments, the heterologous gene or genome segment is incorporated within a chimeric HPIV3 vector genome or antigenome, for example a chimeric HPIV3-1 vector genome or antigenome having one or both HPIV1 genes encoding the HN and F glycoproteins substituted for their counterpart HPIV3 HN and/or F gene(s). In more detailed embodiments, a transcription unit comprising an open reading frame (ORF) of the measles virus HA gene is added to a HPIV3 vector genome or antigenome at various positions, yielding exemplary chimeric PIV/measles vaccine candidates rPIV3(HA HN-L), rPIV3(HA N-P), rcp45L(HA N-P), rPIV3(HA P-M), or rcp45L(HA P-M). Alternatively, chimeric PIV

for vaccine use may incorporate one or more antigenic determinants of HPIV2, for example an HPIV2 HN gene, within a chimeric HPIV3-1 vector genome or antigenome.

In other detailed embodiments of the invention, chimeric PIVs are engineered that incorporate heterologous nucleotide sequences encoding protective antigens from respiratory syncytial virus (RSV) to produce infectious, attenuated vaccine candidates. The cloning of RSV cDNA and other disclosure is provided in U.S. Provisional Patent Application No. 60/007,083, filed September 27, 1995; U.S. Patent Application No. 08/720,132, filed September 27, 1996; U.S. Provisional Patent Application No. 60/021,773, filed July 15, 1996; U.S. Provisional Patent Application No. 60/046,141, filed May 9, 1997; U.S. Provisional Patent Application No. 60/047,634, filed May 23, 1997; U.S. Patent Application No. 08/892,403, filed July 15, 1997 (corresponding to International Publication No. WO 98/02530); U.S. Patent Application No. 09/291,894, filed on April 13, 1999; International Application No. PCT/US00/09696, filed April 12, 2000, corresponding to U.S. Provisional Patent Application Serial No. 60/129,006, filed on April 13, 1999; Collins et al., Proc Nat. Acad. Sci. U.S.A. 92:11563-11567, 1995; Bukreyev et al., J. Virol. 70:6634-41, 1996; Juhasz et al., J. Virol. 71:5814-5819, 1997; Durbin et al., Virology 235:323-332, 1997; He et al. Virology 237:249-260, 1997; Baron et al. J. Virol. 71:1265-1271, 1997; Whitehead et al., Virology 247:232-9, 1998a; Whitehead et al., J. Virol. 72:4467-4471, 1998b; Jin et al. Virology 251:206-214, 1998; and Whitehead et al., J. Virol. 73:3438-3442, 1999, and Bukreyev et al., Proc. Nat. Acad. Sci. U.S.A. 96:2367-72, 1999, each incorporated herein by reference in its entirety for all purposes). Other reports and discussion incorporated or set forth herein identify and characterize RSV antigenic determinants that are useful within the invention.

PIV chimeras incorporating one or more RSV antigenic determinants, preferably comprise a human PIV (e.g., HPIV1, HPIV2, HPIV3) vector genome or antigenome with a heterologous gene or genome segment encoding an antigenic RSV glycoprotein, protein domain (e.g., a glycoprotein ectodomain) or one or more immunogenic epitopes. In one embodiment, one or more genes or genome segments from RSV F and/or G genes is/are combined with the vector genome or antigenome to form the chimeric PIV vaccine candidate. Certain of these constructs will express chimeric proteins, for example fusion proteins having a cytoplasmic tail and/or transmembrane domain of PIV fused to an ectodomain of RSV to yield a novel attenuated virus that elicits a multivalent immune response against both PIV and RSV

As noted above, it is often desirable to adjust the phenotype of chimeric PIV for vaccine use by introducing additional mutations that increase or decrease attenuation or otherwise alter the phenotype of the chimeric virus. Detailed descriptions of the materials and methods for producing recombinant PIV from cDNA, and for making and testing various mutations and nucleotide modifications set forth herein as supplemental aspects of the present invention are provided in, e.g., Durbin et al., *Virology* 235:323-332, 1997; U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998; U.S. Patent Application Serial No. 09/458,813, filed December 10, 1999; U.S. Patent Application Serial No. 09/459,062, filed December 10, 1999; U.S. Provisional Application No. 60/047,575, filed May 23, 1997 (corresponding to International Publication No. WO 98/53078), and U.S. Provisional Application No. 60/059,385, filed September 19, 1997, each incorporated herein by reference. In particular, these documents describe methods and procedures for mutagenizing, isolating and characterizing PIV to obtain attenuated mutant strains (e.g., temperature sensitive (*ts*), cold passaged (*cp*) cold-adapted (*ca*), small plaque (*sp*) and host-range restricted (*hr*) mutant strains) and for identifying the genetic changes that specify the attenuated phenotype. In conjunction with these methods, the foregoing documents detail procedures for determining replication, immunogenicity, genetic stability and protective efficacy of biologically derived and recombinantly produced attenuated human PIV in accepted model systems, including murine and non-human primate model systems. In addition, these documents describe general methods for developing and testing immunogenic compositions, including monovalent and bivalent vaccines, for prophylaxis and treatment of PIV infection. Methods for producing infectious recombinant PIV by construction and expression of cDNA encoding a PIV genome or antigenome coexpressed with essential PIV proteins are also described in the above-incorporated documents, which include description of the following exemplary plasmids that may be employed to produce infectious PIV clones: p3/7(131) (ATCC 97990); p3/7(131)2G (ATCC 97889); and p218(131) (ATCC 97991); each deposited under the terms of the Budapest Treaty with the American Type Culture Collection (ATCC) of 10801 University Boulevard, Manassas, Virginia 20110-2209, U.S.A., and granted the above identified accession numbers.

Also disclosed in the above-incorporated references are methods for constructing and evaluating infectious recombinant PIV that are modified to incorporate phenotype-specific mutations identified in biologically-derived PIV mutants, e.g., cold passaged (*cp*), cold adapted (*ca*), host range restricted (*hr*), small plaque (*sp*), and/or temperature sensitive (*ts*) mutants, for example the JS HPIV3 *cp* 45 mutant strain. Mutations

identified in these mutants can be readily incorporated into chimeric PIV of the instant invention. In exemplary embodiments, one or more attenuating mutations occur in the polymerase L protein, e.g., at a position corresponding to Tyr₉₄₂, Leu₉₉₂, or Thr₁₅₅₈ of JS *cp45*. Preferably, these mutations are incorporated in chimeric PIV of the invention by an identical, or conservative, amino acid substitution as identified in the biological mutant. In more detailed aspects, chimeric PIV for vaccine use incorporate one or more mutation wherein Tyr₉₄₂ is replaced by His, Leu₉₉₂ is replaced by Phe, and/or Thr₁₅₅₈ is replaced by Ile. Substitutions that are conservative to these replacement amino acids are also useful to achieve desired attenuation in chimeric vaccine candidates. The HPIV3 JS *cp45* strain has been deposited under the terms of the Budapest Treaty with the American Type Culture Collection (ATCC) of 10801 University Boulevard, Manassas, Virginia 20110-2209, U.S.A. under Patent Deposit Designation PTA-2419.

Other exemplary mutations that can be adopted in chimeric PIVs from biologically derived PIV mutants include one or more mutations in the N protein, including specific mutations at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS *cp45*. In more detailed aspects, these mutations are represented as Val₉₆ to Ala or Ser₃₈₉ to Ala or substitutions that are conservative thereto. Also useful within chimeric PIV of the invention are amino acid substitution in the C protein, e.g., a mutation at a position corresponding to Ile₉₆ of JS *cp45*, preferably represented by an identical or conservative substitution of Ile₉₆ to Thr. Further exemplary mutations that can be adopted from biologically derived PIV mutants include one or more mutations in the F protein, including mutations adopted from JS *cp45* at a position corresponding to residues Ile₄₂₀ or Ala₄₅₀ of JS *cp45*, preferably represented by acid substitutions Ile₄₂₀ to Val or Ala₄₅₀ to Thr or substitutions conservative thereto. Alternatively or in addition, chimeric PIV of the invention can adopt one or more amino acid substitutions in the HN protein, as exemplified by a mutation at a position corresponding to residue Val₃₈₄ of JS *cp45*, preferably represented by the substitution Val₃₈₄ to Ala.

Yet additional embodiments of the invention include chimeric PIV which incorporate one or more mutations in noncoding portions of the PIV genome or antigenome, for example in a 3' leader sequence, that specify desired phenotypic changes such as attenuation. Exemplary mutations in this context may be engineered at a position in the 3' leader of the chimeric virus at a position corresponding to nucleotide 23, 24, 28, or 45 of JS *cp45*. Yet additional exemplary mutations may be engineered in the N gene start sequence,

for example by changing one or more nucleotides in the N gene start sequence, e.g., at a position corresponding to nucleotide 62 of JS *cp45*. In more detailed aspects, chimeric PIV incorporate a T to C change at nucleotide 23, a C to T change at nucleotide 24, a G to T change at nucleotide 28, and/or a T to A change at nucleotide 45. Additional mutations in
5 extragenic sequences are exemplified by an A to T change in the N gene start sequence at a position corresponding to nucleotide 62 of JS.

These foregoing exemplary mutations which can be engineered in a chimeric PIV of the invention have been successfully engineered and recovered in recombinant PIV—as represented by the recombinant PIV clones designated *rcp45*, *rcp45 L*, *rcp45 F*,
10 *rcp45 M*, *rcp45 HN*, *rcp45 C*, *rcp45 F*, *rcp45 3'N*, *rcp3'NL*, and *rcp45 3'NCMFHN* (Durbin et al., *Virology* 235:323-332, 1997; Skiadopoulos et al., *J. Virol.* 72:1762-1768, 1998; Skiadopoulos et al., *J. Virol.* 73:1374-1381, 1999; U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998; U.S. Patent Application Serial No. 09/458,813, filed December 10, 1999; U.S. Patent Application Serial No. 09/459,062, filed December 10,
15 1999; U.S. Provisional Application No. 60/047,575, filed May 23, 1997 (corresponding to International Publication No. WO 98/53078), and U.S. Provisional Application No. 60/059,385, filed September 19, 1997, each incorporated herein by reference). In addition, the above-incorporated references describe construction of chimeric PIV recombinants, e.g., having the HN and F genes of HPIV1 substituted into a partial HPIV3 background genome
20 or antigenome, which is further modified to bear one or more of the attenuating mutations identified in HPIV3 JS *cp45*. One such chimeric recombinant incorporates all of the attenuating mutations identified in the L gene of *cp45*. It has since been shown that all of the *cp45* mutations outside of the heterologous (HPIV1) HN and F genes can be incorporated in a HPIV3-1 recombinant to yield an attenuated, chimeric vaccine candidate.

25 From JS *cp45* and other biologically derived PIV mutants, a large "menu" of attenuating mutations is provided, each of which can be combined with any other mutation(s) for adjusting the level of attenuation, immunogenicity and genetic stability in chimeric PIV of the invention. In this context, many chimeric PIVs will include one or more, and preferably two or more, mutations from biologically derived PIV mutants, e.g.,
30 any one or combination of mutations identified in JS *cp45*. Preferred chimeric PIVs within the invention will incorporate a plurality and up to a full complement of the mutations present in JS *cp45* or other biologically derived mutant PIV strains. Preferably, these

mutations are stabilized against reversion in chimeric PIV by multiple nucleotide substitutions in a codon specifying each mutation.

Yet additional mutations that may be incorporated in chimeric PIV of the invention are mutations, e.g., attenuating mutations, identified in heterologous PIV or other nonsegmented negative stranded RNA viruses. In particular, attenuating and other desired mutations identified in one negative stranded RNA virus may be "transferred", e.g., copied, to a corresponding position within the genome or antigenome of a chimeric PIV. Briefly, desired mutations in one heterologous negative stranded RNA virus are transferred to the chimeric PIV recipient (either in the vector genome or antigenome or in the heterologous donor gene or genome segment). This involves mapping the mutation in the heterologous mutant virus, identifying by routine sequence alignment the corresponding site in the recipient PIV, and mutating the native sequence in the PIV recipient to the mutant genotype (either by an identical or conservative mutation), as described in International Application No. PCT/US00/09695, filed April 12, 2000, corresponding to U.S. Provisional Patent Application Serial No. 60/129,006, filed on April 13, 1999, incorporated herein by reference. As this disclosure teaches, it is preferable to modify the recipient chimeric PIV genome or antigenome to encode an alteration at the subject site of mutation that corresponds conservatively to the alteration identified in the heterologous mutant virus. For example, if an amino acid substitution marks a site of mutation in the mutant virus compared to the corresponding wild-type sequence, then a similar substitution can be engineered at the corresponding residue(s) in the recombinant virus. Preferably the substitution will specify an identical or conservative amino acid to the substitute residue present in the mutant viral protein. However, it is also possible to alter the native amino acid residue at the site of mutation non-conservatively with respect to the substitute residue in the mutant protein (e.g., by using any other amino acid to disrupt or impair the function of the wild-type residue). Negative stranded RNA viruses from which exemplary mutations are identified and transferred into a recombinant PIV of the invention include other PIVs (e.g., HPIV1, HPIV2, HPIV3, BPIV and MPIV), RSV, Sendai virus (SeV), Newcastle disease virus (NDV), simian virus 5 (SV5), measles virus (MeV), rinderpest virus, canine distemper virus (CDV), rabies virus (RaV) and vesicular stomatitis virus (VSV), among others. A variety of exemplary mutations are disclosed, including but not limited to an amino acid substitution of phenylalanine at position 521 of the RSV L protein corresponding to and therefore transferable to a substitution of phenylalanine (or a conservatively related amino acid) at position 456 of the HPIV3 L protein. In the case of mutations marked by deletions or

insertions, these can be introduced as corresponding deletions or insertions into the recombinant virus, however the particular size and amino acid sequence of the deleted or inserted protein fragment can vary.

Attenuating mutations in biologically derived PIV and other nonsegmented
5 negative stranded RNA viruses for incorporation within chimeric PIV of the invention may occur naturally or may be introduced into wild-type PIV strains by well known mutagenesis procedures. For example, incompletely attenuated parental PIV strains can be produced by chemical mutagenesis during virus growth in cell cultures to which a chemical mutagen has been added, by selection of virus that has been subjected to passage at suboptimal
10 temperatures in order to introduce growth restriction mutations, or by selection of a mutagenized virus that produces small plaques (*sp*) in cell culture, as described in the above incorporated references. By "biologically derived PIV" is meant any PIV not produced by recombinant means. Thus, biologically derived PIV include all naturally occurring PIV, including, e.g., naturally occurring PIV having a wild-type genomic sequence and PIV
15 having allelic or mutant genomic variations from a reference wild-type PIV sequence, e.g., PIV having a mutation specifying an attenuated phenotype. Likewise, biologically derived PIV include PIV mutants derived from a parental PIV by, *inter alia*, artificial mutagenesis and selection procedures.

As noted above, production of a sufficiently attenuated biologically derived
20 PIV mutant can be accomplished by several known methods. One such procedure involves subjecting a partially attenuated virus to passage in cell culture at progressively lower, attenuating temperatures. For example, partially attenuated mutants are produced by passage in cell cultures at suboptimal temperatures. Thus, a *cp* mutant or other partially attenuated PIV strain is adapted to efficient growth at a lower temperature by passage in culture. This
25 selection of mutant PIV during cold-passage substantially reduces any residual virulence in the derivative strains as compared to the partially attenuated parent. Alternatively, specific mutations can be introduced into biologically derived PIV by subjecting a partially attenuated parent virus to chemical mutagenesis, e.g., to introduce *ts* mutations or, in the case of viruses which are already *ts*, additional *ts* mutations sufficient to confer increased
30 attenuation and/or stability of the *ts* phenotype of the attenuated derivative. Means for the introduction of *ts* mutations into PIV include replication of the virus in the presence of a mutagen such as 5-fluorouridine according to generally known procedures. Other chemical mutagens can also be used. Attenuation can result from a *ts* mutation in almost any PIV

gene, although a particularly amenable target for this purpose has been found to be the polymerase (L) gene. The level of temperature sensitivity of replication in exemplary attenuated PIV for use within the invention is determined by comparing its replication at a permissive temperature with that at several restrictive temperatures. The lowest temperature at which the replication of the virus is reduced 100-fold or more in comparison with its replication at the permissive temperature is termed the shutoff temperature. In experimental animals and humans, both the replication and virulence of PIV correlate with the mutant's shutoff temperature.

The JS *cp45* HPIV3 mutant has been found to be relatively stable genetically, highly immunogenic, and satisfactorily attenuated. Nucleotide sequence analysis of this biologically derived virus, and of recombinant viruses incorporating various individual and combined mutations found therein, indicates that each level of increased attenuation is associated with specific nucleotide and amino acid substitutions. The above-incorporated references also disclose how to routinely distinguish between silent incidental mutations and those responsible for phenotype differences by introducing the mutations, separately and in various combinations, into the genome or antigenome of infectious PIV clones. This process coupled with evaluation of phenotype characteristics of parental and derivative viruses identifies mutations responsible for such desired characteristics as attenuation, temperature sensitivity, cold-adaptation, small plaque size, host range restriction, etc.

Mutations thus identified are compiled into a "menu" and are then introduced as desired, singly or in combination, to adjust chimeric PIV of the invention to an appropriate level of attenuation, immunogenicity, genetic resistance to reversion from an attenuated phenotype, etc., as desired. In accordance with the foregoing description, the ability to produce infectious PIV from cDNA permits introduction of specific engineered changes within chimeric PIV. In particular, infectious, recombinant PIVs are employed for identification of specific mutation(s) in biologically derived, attenuated PIV strains, for example mutations which specify *ts*, *ca*, *att* and other phenotypes. Desired mutations are thus identified and introduced into chimeric PIV vaccine strains. The capability of producing virus from cDNA allows for routine incorporation of these mutations, individually or in various selected combinations, into a full-length cDNA clone, whereafter the phenotypes of rescued recombinant viruses containing the introduced mutations to be readily determined.

By identifying and incorporating specific mutations associated with desired phenotypes, e.g., a *cp* or *ts* phenotype, into infectious chimeric PIV clones, the invention provides for other, site-specific modifications at, or within close proximity to, the identified mutation. Whereas most attenuating mutations produced in biologically derived PIVs are single nucleotide changes, other "site specific" mutations can also be incorporated by recombinant techniques into a chimeric PIV. As used herein, site-specific mutations include insertions, substitutions, deletions or rearrangements of from 1 to 3, up to about 5-15 or more altered nucleotides (e.g., altered from a wild-type PIV sequence, from a sequence of a selected mutant PIV strain, or from a parent recombinant PIV clone subjected to mutagenesis). Such site-specific mutations may be incorporated at, or within the region of, a selected, biologically derived point mutation. Alternatively, the mutations can be introduced in various other contexts within a PIV clone, for example at or near a cis-acting regulatory sequence or nucleotide sequence encoding a protein active site, binding site, immunogenic epitope, etc. Site-specific PIV mutants typically retain a desired attenuating phenotype, but may additionally exhibit altered phenotypic characteristics unrelated to attenuation, e.g., enhanced or broadened immunogenicity, and/or improved growth. Further examples of desired, site-specific mutants include recombinant PIV designed to incorporate additional, stabilizing nucleotide mutations in a codon specifying an attenuating point mutation. Where possible, two or more nucleotide substitutions are introduced at codons that specify attenuating amino acid changes in a parent mutant or recombinant PIV clone, yielding a PIV with greater genetic resistance to reversion from an attenuated phenotype. In other embodiments, site-specific nucleotide substitutions, additions, deletions or rearrangements are introduced upstream (N-terminal direction) or downstream (C-terminal direction), e.g., from 1 to 3, 5-10 and up to 15 nucleotides or more 5' or 3', relative to a targeted nucleotide position, e.g., to construct or ablate an existing cis-acting regulatory element.

In addition to single and multiple point mutations and site-specific mutations, changes to the chimeric PIV disclosed herein include deletions, insertions, substitutions or rearrangements of one or more gene(s) or genome segment(s). Particularly useful are deletions involving one or more gene(s) or genome segment(s), which deletions have been shown to yield additional desired phenotypic effects. Thus, U.S. Patent Application Serial No. 09/350,821, filed by Durbin et al. on July 9, 1999, incorporated herein by reference, describes methods and compositions whereby expression of one or more HPIV genes, for example one or more of the C, D, and/or V ORFs, is reduced or ablated by modifying the PIV genome or antigenome to incorporate a mutation that alters the coding assignment of an

initiation codon or mutation(s) that introduce one or more stop codon(s).

Alternatively, one or more of the C, D, and/or V ORFs can be deleted in whole or in part to render the corresponding protein(s) partially or entirely non-functional or to disrupt protein expression altogether. Chimeric PIV having such mutations in C, D, and/or V, or other non-essential gene(s), possess highly desirable phenotypic characteristics for vaccine development. For example, these modifications may specify one or more desired phenotypic changes including (i) altered growth properties in cell culture, (ii) attenuation in the upper and/or lower respiratory tract of mammals, (iii) a change in viral plaque size, (iv) a change in cytopathic effect, and (v) a change in immunogenicity. One exemplary "knock out" mutant PIV lacking C ORF expression, designated rC-KO, was able to induce a protective immune response against wild type HPIV3 challenge in a non-human primate model despite its beneficial attenuation phenotype.

Thus, in more detailed aspects of the instant invention, chimeric PIV incorporate deletion or knock out mutations in a C, D, and/or V ORF(s) or other non-essential gene which alters or ablates expression of the selected gene(s) or genome segment(s). This can be achieved, e.g., by introducing a frame shift mutation or termination codon within a selected coding sequence, altering translational start sites, changing the position of a gene or introducing an upstream start codon to alter its rate of expression, changing GS and/or GE transcription signals to alter phenotype, or modifying an RNA editing site (e.g., growth, temperature restrictions on transcription, etc.). In more detailed aspects of the invention, chimeric PIVs are provided in which expression of one or more gene(s), e.g., a C, D, and/or V ORF(s), is ablated at the translational or transcriptional level without deletion of the gene or of a segment thereof, by, e.g., introducing multiple translational termination codons into a translational open reading frame (ORF), altering an initiation codon, or modifying an editing site. These forms of knock-out virus will often exhibit reduced growth rates and small plaque sizes in tissue culture. Thus, these methods provide yet additional, novel types of attenuating mutations which ablate expression of a viral gene that is not one of the major viral protective antigens. In this context, knock-out virus phenotypes produced without deletion of a gene or genome segment can be alternatively produced by deletion mutagenesis, as described, to effectively preclude correcting mutations that may restore synthesis of a target protein. Several other gene knock-outs for the C, D, and/or V ORF(s) deletion and knock out mutants can be made using alternate designs and methods that are well known in the art (as described, for example, in (Kretschmer et al., *Virology* 216:309-316, 1996; Radecke et al., *Virology*

217:418-421, 1996; Kato et al., EMBO J. 16:578-587, 1987; and Schneider et al., Virology 277:314-322, 1996, each incorporated herein by reference).

Nucleotide modifications that may be introduced into chimeric PIV constructs of the invention may alter small numbers of bases (e.g., from 15-30 bases, up to 35-50 bases or more), large blocks of nucleotides (e.g., 50-100, 100-300, 300-500, 500-1,000 bases), or nearly complete or complete genes (e.g., 1,000-1,500 nucleotides, 1,500-2,500 nucleotides, 2,500-5,000, nucleotides, 5,00-6,5000 nucleotides or more) in the vector genome or antigenome or heterologous, donor gene or genome segment, depending upon the nature of the change (i.e., a small number of bases may be changed to insert or ablate an immunogenic epitope or change a small genome segment, whereas large block(s) of bases are involved when genes or large genome segments are added, substituted, deleted or rearranged.

In related aspects, the invention provides for supplementation of mutations adopted into a chimeric PIV clone from biologically derived PIV, e.g., *cp* and *ts* mutations, with additional types of mutations involving the same or different genes in a further modified PIV clone. Each of the PIV genes can be selectively altered in terms of expression levels, or can be added, deleted, substituted or rearranged, in whole or in part, alone or in combination with other desired modifications, to yield a chimeric PIV exhibiting novel vaccine characteristics. Thus, in addition to or in combination with attenuating mutations adopted from biologically derived PIV mutants, the present invention also provides a range of additional methods for attenuating or otherwise modifying the phenotype of a chimeric PIV based on recombinant engineering of infectious PIV clones. A variety of alterations can be produced in an isolated polynucleotide sequence encoding a targeted gene or genome segment, including a donor or recipient gene or genome segment in a chimeric PIV genome or antigenome for incorporation into infectious clones. More specifically, to achieve desired structural and phenotypic changes in recombinant PIV, the invention allows for introduction of modifications which delete, substitute, introduce, or rearrange a selected nucleotide or nucleotide sequence from a parent genome or antigenome, as well as mutations which delete, substitute, introduce or rearrange whole gene(s) or genome segment(s), within a chimeric PIV clone.

Thus provided are modifications in chimeric PIV of the invention which simply alter or ablate expression of a selected gene, e.g., by introducing a termination codon within a selected PIV coding sequence or altering its translational start site or RNA editing

site, changing the position of a PIV gene relative to an operably linked promoter, introducing an upstream start codon to alter rates of expression, modifying (e.g., by changing position, altering an existing sequence, or substituting an existing sequence with a heterologous sequence) GS and/or GE transcription signals to alter phenotype (e.g., growth, temperature restrictions on transcription, etc.), and various other deletions, substitutions, additions and rearrangements that specify quantitative or qualitative changes in viral replication, transcription of selected gene(s), or translation of selected protein(s). In this context, any PIV gene or genome segment which is not essential for growth can be ablated or otherwise modified in a recombinant PIV to yield desired effects on virulence, pathogenesis, immunogenicity and other phenotypic characters. As for coding sequences, noncoding, leader, trailer and intergenic regions can be similarly deleted, substituted or modified and their phenotypic effects readily analyzed, e.g., by the use of minireplicons and recombinant PIV.

In addition, a variety of other genetic alterations can be produced in a PIV genome or antigenome for incorporation into a chimeric PIV, alone or together with one or more attenuating mutations adopted from a biologically derived mutant PIV, e.g., to adjust growth, attenuation, immunogenicity, genetic stability or provide other advantageous structural and/or phenotypic effects. These additional types of mutations are also disclosed in the foregoing incorporated references and can be readily engineered into chimeric PIV of the invention. For example, restriction site markers are routinely introduced within chimeric PIVs to facilitate cDNA construction and manipulation.

In addition to these changes, the order of genes in a chimeric PIV construct can be changed, a PIV genome promoter replaced with its antigenome counterpart, portions of genes removed or substituted, and even entire genes deleted. Different or additional modifications in the sequence can be made to facilitate manipulations, such as the insertion of unique restriction sites in various intergenic regions or elsewhere. Nontranslated gene sequences can be removed to increase capacity for inserting foreign sequences.

Other mutations for incorporation into chimeric PIV constructs of the invention include mutations directed toward cis-acting signals, which can be readily identified, e.g., by mutational analysis of PIV minigenomes. For example, insertional and deletional analysis of the leader and trailer and flanking sequences identifies viral promoters and transcription signals and provides a series of mutations associated with varying degrees

of reduction of RNA replication or transcription. Saturation mutagenesis (whereby each position in turn is modified to each of the nucleotide alternatives) of these cis-acting signals also has identified many mutations which affect RNA replication or transcription. Any of these mutations can be inserted into a chimeric PIV antigenome or genome as described

5 herein. Evaluation and manipulation of trans-acting proteins and cis-acting RNA sequences using the complete antigenome cDNA is assisted by the use of PIV minigenomes as described in the above-incorporated references.

Additional mutations within chimeric PIVs of the invention may also include replacement of the 3' end of genome with its counterpart from antigenome, which is

10 associated with changes in RNA replication and transcription. In one exemplary embodiment, the level of expression of specific PIV proteins, such as the protective HN and/or F antigens, can be increased by substituting the natural sequences with ones which have been made synthetically and designed to be consistent with efficient translation. In this context, it has been shown that codon usage can be a major factor in the level of translation

15 of mammalian viral proteins (Haas et al., *Current Biol.* 6:315-324, 1996, incorporated herein by reference). Optimization by recombinant methods of the codon usage of the mRNAs encoding the HN and F proteins of PIV will provide improved expression for these genes.

In another exemplary embodiment, a sequence surrounding a translational start site (preferably including a nucleotide in the -3 position) of a selected PIV gene is

20 modified, alone or in combination with introduction of an upstream start codon, to modulate PIV gene expression by specifying up- or down-regulation of translation. Alternatively, or in combination with other recombinant modifications disclosed herein, gene expression of a chimeric PIV can be modulated by altering a transcriptional GS or GE signal of any selected gene(s) of the virus. In alternative embodiments, levels of gene expression in a chimeric

25 PIV vaccine candidate are modified at the level of transcription. In one aspect, the position of a selected gene in the PIV gene map can be changed to a more promoter-proximal or promoter-distal position, whereby the gene will be expressed more or less efficiently, respectively. According to this aspect, modulation of expression for specific genes can be achieved yielding reductions or increases of gene expression from two-fold, more typically

30 four-fold, up to ten-fold or more compared to wild-type levels often attended by a commensurate decrease in expression levels for reciprocally, positionally substituted genes. These and other transpositioning changes yield novel chimeric PIV vector virus having attenuated phenotypes, for example due to decreased expression of selected viral proteins

involved in RNA replication, or having other desirable properties such as increased antigen expression.

In other embodiments, chimeric PIVs useful in vaccine formulations can be conveniently modified to accommodate antigenic drift in circulating virus. Typically the modification will be in the HN and/or F proteins. An entire HN or F gene, or a genome segment encoding a particular immunogenic region thereof, from one PIV strain or group is incorporated into a chimeric PIV genome or antigenome cDNA by replacement of a corresponding region in a recipient clone of a different PIV strain or group, or by adding one or more copies of the gene, such that multiple antigenic forms are represented. Progeny virus produced from the modified PIV clone can then be used in vaccination protocols against emerging PIV strains.

Replacement of a human PIV coding sequence or non-coding sequence (e.g., a promoter, gene-end, gene-start, intergenic or other cis-acting element) with a heterologous counterpart yields chimeric PIV having a variety of possible attenuating and other phenotypic effects. In particular, host range and other desired effects arise from substituting a bovine PIV (BPIV) or murine PIV (MPIV) protein, protein domain, gene or genome segment imported within a human PIV background, wherein the bovine or murine gene does not function efficiently in a human cell, e.g., from incompatibility of the heterologous sequence or protein with a biologically interactive human PIV sequence or protein (i.e., a sequence or protein that ordinarily cooperates with the substituted sequence or protein for viral transcription, translation, assembly, etc.) or, more typically in a host range restriction, with a cellular protein or some other aspect of the cellular milieu which is different between the permissive and less permissive host. In exemplary embodiments, bovine PIV sequences are selected for introduction into human PIV based on known aspects of bovine and human PIV structure and function.

In more detailed aspects, the invention provides methods for attenuating chimeric PIV vaccine candidates based on the further construction of chimeras between HPIV and a non-human PIV, for example HPIV3 and BPIV3 (e.g., as disclosed in U.S. Patent Application Serial No. 09/586,479, filed June 1, 2000 by Schmidt et al.; Schmidt et al., *J. Virol.* 74:8922-9, 2000, each incorporated herein by reference). This method of attenuation is based on host range effects due to the introduction of one or more gene(s) or genome segment(s) of the non-human PIV into a human PIV vector-based chimeric virus.

For example, there are numerous nucleotide and amino acid sequence differences between BPIV and HPIVs, which are reflected in host range differences. Between HPIV3 and BPIV3 the percent amino acid identity for each of the following proteins is: N (86%), P (65%), M (93%), F (83%), HN (77%), and L (91%). The host range difference is exemplified by the highly permissive growth of HPIV3 in rhesus monkeys, compared to the restricted replication of two different strains of BPIV3 in the same animal (van Wyke Coelingh et al., *J. Infect. Dis.* 157:655-662, 1988, incorporated herein by reference). Although the basis of the host range differences between HPIV3 and BPIV3 remains to be determined, it is likely that they will involve more than one gene and multiple amino acid differences. The involvement of multiple genes and possibly cis-acting regulatory sequences, each involving multiple amino acid or nucleotide differences, gives a very broad basis for attenuation, one which cannot readily be altered by reversion. This is in contrast to the situation with other live attenuated HPIV3 viruses which are attenuated by one or several point mutations. In this case, reversion of any individual mutation may yield a significant reacquisition of virulence or, in a case where only a single residue specified attenuation, complete reacquisition of virulence.

In exemplary embodiments of the invention, the vector genome or antigenome is an HPIV3 genome or antigenome, and the heterologous gene or genome segment is a N ORF derived from, alternatively, a Ka or SF strain of BPIV3 (which are 99% related in amino acid sequence). The N ORF of the HPIV3 background antigenome is substituted by the counterpart BPIV3 N ORF—yielding a novel recombinant chimeric PIV clone. Replacement of the HPIV3 N ORF of HPIV3 with that of BPIV3 Ka or SF results in a protein with approximately 70 amino acid differences (depending on the strain involved) from that of HPIV3 N. N is one of the more conserved proteins, and substitution of other proteins such as P, singly or in combination, would result in many more amino acid differences. The involvement of multiple genes and genome segments each conferring multiple amino acid or nucleotide differences provides a broad basis for attenuation which is highly stable to reversion.

This mode of attenuation contrasts sharply to HPIV vaccine candidates that are attenuated by one or more point mutations, where reversion of an individual mutation may yield a significant or complete reacquisition of virulence. In addition, several known attenuating point mutations in HPIV typically yield a temperature sensitive phenotype. One problem with attenuation associated with temperature sensitivity is that the virus can be

overly restricted for replication in the lower respiratory tract while being under attenuated in the upper respiratory tract. This is because there is a temperature gradient within the respiratory tract, with temperature being higher (and more restrictive) in the lower respiratory tract and lower (less restrictive) in the upper respiratory tract. The ability of an attenuated virus to replicate in the upper respiratory tract can result in complications including congestion, rhinitis, fever and otitis media. Thus, attenuation achieved solely by temperature sensitive mutations may not be ideal. In contrast, host range mutations present in chimeric PIV of the invention will not in most cases confer temperature sensitivity. Therefore, the novel method of PIV attenuation provided by these kinds of modifications will be more stable genetically and phenotypically and less likely to be associated with residual virulence in the upper respiratory tract compared to other known PIV vaccine candidates.

The above-incorporated reference discloses that both Ka and SF HPIV3/BPIV3 chimeric recombinants are viable and replicate as efficiently in cell culture as either HPIV3 or BPIV3 parent—indicating that the chimeric recombinants did not exhibit gene incompatibilities that restricted replication *in vitro*. This property of efficient replication *in vitro* is important since it permits efficient manufacture of this biological. Also, the Ka and the SF HPIV3/BPIV3 chimeric recombinants (termed cKa and cSF), bearing only one bovine gene, are nearly equivalent to their BPIV3 parents in the degree of host range restriction in the respiratory tract of the rhesus monkey. In particular, the cKa and cSF viruses exhibit approximately a 60-fold or 30-fold reduction, respectively, in replication in the upper respiratory tract of rhesus monkeys compared to replication of HPIV3. Based on this finding, it is expected that other BPIV3 genes will also confer desired levels of host range restriction within chimeric PIV of the invention. Thus, according to the methods herein, a list of attenuating determinants will be readily identified in heterologous genes and genome segments of BPIV and other non-human PIVs that will confer, in appropriate combination, a desired level of host range restriction and immunogenicity on chimeric PIV selected for vaccine use.

Chimeric human-bovine PIV for use as vectors within the present invention include a partial or complete “background” PIV genome or antigenome derived from or patterned after a human or bovine PIV strain or subgroup virus combined with one or more heterologous gene(s) or genome segment(s) of a different PIV strain or subgroup virus to form the human-bovine chimeric PIV genome or antigenome. In preferred aspects of the

invention, chimeric PIV incorporate a partial or complete human PIV background genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a bovine PIV. The partial or complete background genome or antigenome typically acts as a recipient backbone or vector into which are imported heterologous genes or genome segments of the counterpart, human or bovine PIV. Heterologous genes or genome segments from the counterpart, human or bovine PIV represent "donor" genes or polynucleotides that are combined with, or substituted within, the background genome or antigenome to yield a human-bovine chimeric PIV that exhibits novel phenotypic characteristics compared to one or both of the contributing PIVs. For example, addition or substitution of heterologous genes or genome segments within a selected recipient PIV strain may result in an increase or decrease in attenuation, growth changes, altered immunogenicity, or other desired phenotypic changes as compared with a corresponding phenotype(s) of the unmodified recipient and/or donor (U.S. Patent Application Serial No. 09/586,479, filed June 1, 2000 by Schmidt et al.; Schmidt et al., *J. Virol.* 74:8922-9, 2000, each incorporated herein by reference).

Genes and genome segments that may be selected for use as heterologous substitutions or additions within human-bovine chimeric PIV vectors include genes or genome segments encoding a PIV N, P, C, D, V, M, F, SH (where appropriate), HN and/or L protein(s) or portion(s) thereof. In addition, genes and genome segments encoding non-PIV proteins, for example, an SH protein as found in mumps and SV5 viruses, may be incorporated within human-bovine PIV of the invention. Regulatory regions, such as the extragenic 3' leader or 5' trailer regions, and gene-start, gene-end, intergenic regions, or 3' or 5' non-coding regions, are also useful as heterologous substitutions or additions.

Certain human-bovine chimeric PIV vectors for use within the invention bear one or more of the major antigenic determinants of HPIV3 in a background which is attenuated by the substitution or addition of one or more BPIV3 genes or genome segments. The major protective antigens of PIVs are their HN and F glycoproteins, although other proteins can also contribute to a protective immune response. In certain embodiments, the background genome or antigenome is an HPIV genome or antigenome, e.g., an HPIV3, HPIV2, or HPIV1 background genome or antigenome, to which is added or into which is substituted one or more BPIV gene(s) or genome segment(s), preferably from BPIV3. In one exemplary embodiment described below, an ORF of the N gene of a BPIV3 is substituted for that of an HPIV. Alternatively, the background genome or antigenome may

be a BPIV genome or antigenome which is combined with one or more genes or genome segments encoding a HPIV3, HPIV2, or HPIV1 glycoprotein, glycoprotein domain or other antigenic determinant.

5 In accordance with the methods of the invention, any BPIV gene or genome segment, singly or in combination with one or more other BPIV genes, can be combined with HPIV sequences to give rise to a human-bovine chimeric PIV vaccine candidate. Any HPIV, including different strains of a particular HPIV serotype, e.g., HPIV3 will be a reasonable acceptor for attenuating BPIV gene(s). In general, the HPIV3 gene(s) or genome segment(s) selected for inclusion in a human-bovine chimeric PIV for use as a vaccine
10 against human PIV will include one or more of the HPIV protective antigens such as the HN or F glycoproteins.

In exemplary aspects of the invention, human-bovine chimeric PIV bearing one or more bovine gene(s) or genome segment(s) exhibits a high degree of host range restriction, e.g., in the respiratory tract of mammalian models of human PIV infection such
15 as non-human primates. In exemplary embodiments a human PIV is attenuated by the addition or substitution of one or more bovine gene(s) or genome segment(s) to a partial or complete human, e.g., HPIV3, PIV background genome or antigenome. In one example, the HPIV3 N gene is substituted by the BPIV3 N gene to yield a novel human-bovine chimeric PIV vector, and within this vector the measles HA gene is substituted to yield a multivalent,
20 HPIV/measles vaccine candidate, as exemplified by the recombinant rHPIV3-N_B HA_{P-M} described below.

Preferably, the degree of host range restriction exhibited by human-bovine chimeric PIV vectors for developing vaccine candidates of the invention is comparable to the degree of host range restriction exhibited by the respective BPIV parent or "donor" strain.
25 Preferably, the restriction should have a true host range phenotype, i.e., it should be specific to the host in question and should not restrict replication and vaccine preparation *in vitro* in a suitable cell line. In addition, human-bovine chimeric PIV vectors bearing one or more bovine gene(s) or genome segment(s) elicit a high level of resistance in hosts susceptible to PIV infection. Thus, the invention provides a new basis for attenuating a live virus vector
30 for developing vaccines against PIV and other pathogens, based on host range effects.

In related aspects of the invention, human-bovine chimeric PIV vectors comprise a BPIV recipient or backbone virus that incorporates one or more heterologous

gene(s) that encode an HPIV HN and/or F glycoprotein(s). Alternatively, the chimeric PIV may incorporate one or more genome segment(s) encoding an ectodomain (and alternatively a cytoplasmic domain and/or transmembrane domain), or immunogenic epitope of an HPIV HN and/or F glycoprotein(s). These immunogenic proteins, domains and epitopes are particularly useful within human-bovine chimeric PIV because they generate novel immune responses in an immunized host. In particular, the HN and F proteins, and immunogenic domains and epitopes therein, provide major protective antigens.

In certain embodiments of the invention, addition or substitution of one or more immunogenic gene(s) or genome segment(s) from a human PIV subgroup or strain to or within a bovine background, or recipient, genome or antigenome yields a recombinant, chimeric virus or subviral particle capable of generating an immune response directed against the human donor virus, including one or more specific human PIV subgroups or strains, while the bovine backbone confers an attenuated phenotype making the chimera a useful candidate for vaccine development. In one exemplary embodiment, one or more human PIV glycoprotein genes, e.g., HN and/or F, are added to or substituted within a partial or complete bovine genome or antigenome to yield an attenuated, infectious human-bovine chimera that elicits an anti-human PIV immune response in a susceptible host. Within one such exemplary vector (carrying the HPIV3 JS HN and F glycoprotein genes in BPIV3 background), the RSV A glycoprotein genes G and F were successfully inserted as additional heterologous ORF to yield multivalent, HPIV/RSV vaccine candidates exemplified by the recombinant viruses rB/HPIV3-G1 and rB/HPIV3-F1 described below.

In alternate embodiments, human-bovine chimeric PIV vectors additionally incorporate a gene or genome segment encoding an immunogenic protein, protein domain or epitope from multiple human PIV strains, for example two HN or F proteins or immunogenic portions thereof each from a different HPIV, e.g., HPIV1 or HPIV2. Alternatively, one glycoprotein or immunogenic determinant may be provided from a first HPIV, and a second glycoprotein or immunogenic determinant may be provided from a second HPIV by substitution without the addition of an extra glycoprotein- or determinant-encoding polynucleotide to the genome or antigenome. Substitution or addition of HPIV glycoproteins and antigenic determinants may also be achieved by construction of a genome or antigenome that encodes a chimeric glycoprotein in the recombinant virus or subviral particle, for example having an immunogenic epitope, antigenic region or complete ectodomain of a first HPIV fused to a cytoplasmic domain of a heterologous HPIV. For

example, a heterologous genome segment encoding a glycoprotein ectodomain from a HPIV1 or HPIV2 HN or F glycoprotein may be joined with a genome segment encoding a corresponding HPIV3 HN or F glycoprotein cytoplasmic/endodomain in the background genome or antigenome.

5 In alternate embodiments a human-bovine chimeric PIV vector genome or antigenome may encode a substitute, extra, or chimeric glycoprotein or antigenic determinant thereof in the recombinant virus or subviral particle, to yield a viral recombinant having both human and bovine glycoproteins, glycoprotein domains, or immunogenic epitopes. For example, a heterologous genome segment encoding a glycoprotein ectodomain
10 from a human PIV HN or F glycoprotein may be joined with a genome segment encoding a corresponding bovine HN or F glycoprotein cytoplasmic/endodomain in the background genome or antigenome. Alternatively, the human PIV HN or F glycoprotein or parts thereof may be joined with a genome segment encoding an HN or F glycoprotein or parts thereof from another PIV strain or serotype.

15 In combination with the host range phenotypic effects provided in the human-bovine chimeric PIV of the invention, it is often desirable to adjust the attenuation phenotype by introducing additional mutations that increase or decrease attenuation of the chimeric virus. Thus, in additional aspects of the invention, attenuated, human-bovine chimeric PIV
20 vectors are produced in which the chimeric genome or antigenome is further modified by introducing one or more attenuating mutations specifying an attenuating phenotype in the resultant virus or subviral particle. These can include mutations in RNA regulatory sequences or in encoded proteins. These attenuating mutations may be generated *de novo* and tested for attenuating effects according to a rational design mutagenesis strategy. Alternatively, the attenuating mutations may be identified in existing biologically derived
25 mutant PIV and thereafter incorporated into a human-bovine chimeric PIV of the invention.

 In preferred chimeric vaccine candidates of the invention, attenuation marked by replication in the lower and/or upper respiratory tract in an accepted animal model for PIV replication in humans, e.g., hamsters or rhesus monkeys, may be reduced by at least
30 about two-fold, more often about 5-fold, 10-fold, or 20-fold, and preferably 50-100-fold and up to 1,000-fold or greater overall (e.g., as measured between 3-8 days following infection) compared to growth of the corresponding wild-type or mutant parental PIV strain.

Infectious chimeric PIV vector clones of the invention can also be engineered according to the methods and compositions disclosed herein to enhance immunogenicity and induce a level of protection greater than that provided by infection with a wild-type, parental (i.e., vector or heterologous donor) PIV or non-PIV pathogen. For example, one or more supplemental immunogenic epitope(s), protein domains, or proteins from a heterologous PIV strain or type, or from a non-PIV pathogen such as measles or RSV, can be added to a chimeric PIV by appropriate nucleotide changes in the chimeric genome or antigenome. Alternatively, chimeric PIVs of the invention can be engineered to add or ablate (e.g., by amino acid insertion, substitution or deletion) immunogenic proteins, protein domains, or forms of specific proteins associated with desirable or undesirable immunological reactions.

Within the methods of the invention, additional genes or genome segments may be inserted into or proximate to the chimeric PIV vector genome or antigenome. These genes may be under common control with recipient genes, or may be under the control of an independent set of transcription signals. In addition to genes and genome segments encoding antigenic determinants, genes of interest in this context include genes encoding cytokines, for example, an interleukin (e.g., interleukin 2 (IL-2), interleukin 4 (IL-4), interleukin 5 (IL-5), interleukin 6 (IL6), interleukin 18 (IL-18)), tumor necrosis factor alpha (TNF α), interferon gamma (IFN γ), or granulocyte-macrophage colony stimulating factor (GM-CSF), as well as IL-2 through IL-18, especially IL-2, IL-6 and IL-12, and IL-18, gamma-interferon (see, e.g., U.S. Application No. 09/614,285, filed July 12, 2000, corresponding to U.S. Provisional Application Serial No. 60/143,425 filed July 13, 1999, incorporated herein by reference). Coexpression of these additional proteins provides the ability to modify and improve immune responses against chimeric PIV of the invention both quantitatively and qualitatively.

Deletions, insertions, substitutions and other mutations involving changes of whole viral genes or genome segments within chimeric PIV of the invention yield highly stable vaccine candidates, which are particularly important in the case of immunosuppressed individuals. Many of these changes will result in attenuation of resultant vaccine strains, whereas others will specify different types of desired phenotypic changes. For example, accessory (i.e., not essential for *in vitro* growth) genes are excellent candidates to encode proteins that specifically interfere with host immunity (see, e.g., Kato et al., EMBO J. 16:578-87, 1997, incorporated herein by reference). Ablation of such genes in vaccine viruses is expected to reduce virulence and pathogenesis and/or improve immunogenicity.

Introduction of the foregoing defined mutations into an infectious, chimeric PIV clone can be achieved by a variety of well known methods. By "infectious clone" with regard to DNA is meant cDNA or its product, synthetic or otherwise, which can be transcribed into genomic or antigenomic RNA capable of serving as template to produce the genome of an infectious virus or subviral particle. Thus, defined mutations can be introduced by conventional techniques (e.g., site-directed mutagenesis) into a cDNA copy of the genome or antigenome. The use of antigenome or genome cDNA subfragments to assemble a complete antigenome or genome cDNA as described herein has the advantage that each region can be manipulated separately (smaller cDNAs are easier to manipulate than large ones) and then readily assembled into a complete cDNA. Thus, the complete antigenome or genome cDNA, or any subfragment thereof, can be used as template for oligonucleotide-directed mutagenesis. This can be through the intermediate of a single-stranded phagemid form, such as using the Muta-gene® kit of Bio-Rad Laboratories (Richmond, CA) or a method using a double-stranded plasmid directly as template such as the Chameleon mutagenesis kit of Stratagene (La Jolla, CA), or by the polymerase chain reaction employing either an oligonucleotide primer or template which contains the mutation(s) of interest. A mutated subfragment can then be assembled into the complete antigenome or genome cDNA. A variety of other mutagenesis techniques are known and available for use in producing the mutations of interest in the PIV antigenome or genome cDNA. Mutations can vary from single nucleotide changes to replacement of large cDNA pieces containing one or more genes or genome regions.

Thus, in one illustrative embodiment mutations are introduced by using the Muta-gene phagemid *in vitro* mutagenesis kit available from Bio-Rad. In brief, cDNA encoding a portion of a PIV genome or antigenome is cloned into the plasmid pTZ18U, and used to transform CJ236 cells (Life Technologies, Gaithersburg, MD). Phagemid preparations are prepared as recommended by the manufacturer. Oligonucleotides are designed for mutagenesis by introduction of an altered nucleotide at the desired position of the genome or antigenome. The plasmid containing the genetically altered genome or antigenome fragment is then amplified and the mutated piece is then reintroduced into the full-length genome or antigenome clone.

The invention also provides methods for producing infectious chimeric PIV from one or more isolated polynucleotides, e.g., one or more cDNAs. According to the present invention cDNA encoding a PIV genome or antigenome is constructed for

intracellular or *in vitro* coexpression with the necessary viral proteins to form infectious PIV. By "PIV antigenome" is meant an isolated positive-sense polynucleotide molecule which serves as the template for the synthesis of progeny PIV genome. Preferably a cDNA is constructed which is a positive-sense version of the PIV genome, corresponding to the
5 replicative intermediate RNA, or antigenome, so as to minimize the possibility of hybridizing with positive-sense transcripts of the complementing sequences that encode proteins necessary to generate a transcribing, replicating nucleocapsid, i.e., sequences that encode N, P, and L proteins.

For purposes of the present invention the genome or antigenome of the
10 recombinant PIV of the invention need only contain those genes or portions thereof necessary to render the viral or subviral particles encoded thereby infectious. Further, the genes or portions thereof may be provided by more than one polynucleotide molecule, i.e., a gene may be provided by complementation or the like from a separate nucleotide molecule, or can be expressed directly from the genome or antigenome cDNA.

By recombinant PIV is meant a PIV or PIV-like viral or subviral particle
15 derived directly or indirectly from a recombinant expression system or propagated from virus or subviral particles produced therefrom. The recombinant expression system will employ a recombinant expression vector which comprises an operably linked transcriptional unit comprising an assembly of at least a genetic element or elements having a regulatory
20 role in PIV gene expression, for example, a promoter, a structural or coding sequence which is transcribed into PIV RNA, and appropriate transcription initiation and termination sequences.

To produce infectious PIV from cDNA-expressed genome or antigenome, the
25 genome or antigenome is coexpressed with those PIV proteins necessary to (i) produce a nucleocapsid capable of RNA replication, and (ii) render progeny nucleocapsids competent for both RNA replication and transcription. Transcription by the genome nucleocapsid provides the other PIV proteins and initiates a productive infection. Alternatively, additional PIV proteins needed for a productive infection can be supplied by coexpression.

Infectious PIV of the invention are produced by intracellular or cell-free
30 coexpression of one or more isolated polynucleotide molecules that encode a PIV genome or antigenome RNA, together with one or more polynucleotides encoding viral proteins necessary to generate a transcribing, replicating nucleocapsid. Among the viral proteins

useful for coexpression to yield infectious PIV are the major nucleocapsid protein (N) protein, nucleocapsid phosphoprotein (P), large (L) polymerase protein, fusion protein (F), hemagglutinin-neuraminidase glycoprotein (HN), and matrix (M) protein. Also useful in this context are products of the C, D and V ORFs of PIV.

5 cDNAs encoding a PIV genome or antigenome are constructed for intracellular or *in vitro* coexpression with the necessary viral proteins to form infectious PIV. By "PIV antigenome" is meant an isolated positive-sense polynucleotide molecule which serves as a template for synthesis of progeny PIV genome. Preferably a cDNA is constructed which is a positive-sense version of the PIV genome corresponding to the
10 replicative intermediate RNA, or antigenome, so as to minimize the possibility of hybridizing with positive-sense transcripts of complementing sequences encoding proteins necessary to generate a transcribing, replicating nucleocapsid.

In some embodiments of the invention the genome or antigenome of a recombinant PIV (rPIV) need only contain those genes or portions thereof necessary to
15 render the viral or subviral particles encoded thereby infectious. Further, the genes or portions thereof may be provided by more than one polynucleotide molecule, i.e., a gene may be provided by complementation or the like from a separate nucleotide molecule. In other embodiments, the PIV genome or antigenome encodes all functions necessary for viral growth, replication, and infection without the participation of a helper virus or viral function
20 provided by a plasmid or helper cell line.

By "recombinant PIV" is meant a PIV or PIV-like viral or subviral particle derived directly or indirectly from a recombinant expression system or propagated from virus or subviral particles produced therefrom. The recombinant expression system will employ a recombinant expression vector which comprises an operably linked transcriptional
25 unit comprising an assembly of at least a genetic element or elements having a regulatory role in PIV gene expression, for example, a promoter, a structural or coding sequence which is transcribed into PIV RNA, and appropriate transcription initiation and termination sequences.

To produce infectious PIV from a cDNA-expressed PIV genome or
30 antigenome, the genome or antigenome is coexpressed with those PIV N, P and L proteins necessary to (i) produce a nucleocapsid capable of RNA replication, and (ii) render progeny nucleocapsids competent for both RNA replication and transcription. Transcription by the

genome nucleocapsid provides the other PIV proteins and initiates a productive infection. Alternatively, additional PIV proteins needed for a productive infection can be supplied by coexpression.

Synthesis of PIV antigenome or genome together with the above-mentioned
5 viral proteins can also be achieved *in vitro* (cell-free), e.g., using a combined transcription-translation reaction, followed by transfection into cells. Alternatively, antigenome or genome RNA can be synthesized *in vitro* and transfected into cells expressing PIV proteins.

In certain embodiments of the invention, complementing sequences encoding
10 proteins necessary to generate a transcribing, replicating PIV nucleocapsid are provided by one or more helper viruses. Such helper viruses can be wild type or mutant. Preferably, the helper virus can be distinguished phenotypically from the virus encoded by the PIV cDNA. For example, it is desirable to provide monoclonal antibodies which react immunologically with the helper virus but not the virus encoded by the PIV cDNA. Such antibodies can be neutralizing antibodies. In some embodiments, the antibodies can be used in affinity
15 chromatography to separate the helper virus from the recombinant virus. To aid the procurement of such antibodies, mutations can be introduced into the PIV cDNA to provide antigenic diversity from the helper virus, such as in the HN or F glycoprotein genes.

In alternate embodiments of the invention, the N, P, L and other desired PIV
20 proteins are encoded by one or more non-viral expression vectors, which can be the same or separate from that which encodes the genome or antigenome. Additional proteins may be included as desired, each encoded by its own vector or by a vector encoding one or more of the N, P, L and other desired PIV proteins, or the complete genome or antigenome. Expression of the genome or antigenome and proteins from transfected plasmids can be achieved, for example, by each cDNA being under the control of a promoter for T7 RNA
25 polymerase, which in turn is supplied by infection, transfection or transduction with an expression system for the T7 RNA polymerase, e.g., a vaccinia virus MVA strain recombinant which expresses the T7 RNA polymerase (Wyatt et al., *Virology* 210: 202-205, 1995, incorporated herein by reference in its entirety). The viral proteins, and/or T7 RNA polymerase, can also be provided by transformed mammalian cells or by transfection of
30 preformed mRNA or protein.

A PIV antigenome may be constructed for use in the present invention by, e.g., assembling cloned cDNA segments, representing in aggregate the complete

antigenome, by polymerase chain reaction or the like (PCR; described in, e.g., U.S. Patent Nos. 4,683,195 and 4,683,202, and PCR Protocols: A Guide to Methods and Applications, Innis et al., eds., Academic Press, San Diego, 1990, each incorporated herein by reference in its entirety) of reverse-transcribed copies of PIV mRNA or genome RNA. For example, a
5 first construct is generated which comprises cDNAs containing the left hand end of the antigenome, spanning from an appropriate promoter (e.g., T7 RNA polymerase promoter) and assembled in an appropriate expression vector, such as a plasmid, cosmid, phage, or DNA virus vector. The vector may be modified by mutagenesis and/or insertion of synthetic polylinker containing unique restriction sites designed to facilitate assembly. For ease of
10 preparation the N, P, L and other desired PIV proteins can be assembled in one or more separate vectors. The right hand end of the antigenome plasmid may contain additional sequences as desired, such as a flanking ribozyme and tandem T7 transcriptional terminators. The ribozyme can be hammerhead type (e.g., Grosfeld et al., J. Virol. 69:5677-5686, 1995), which would yield a 3' end containing a single nonviral nucleotide, or can be any of the other
15 suitable ribozymes such as that of hepatitis delta virus (Perrotta et al., Nature 350:434-436, 1991), incorporated herein by reference in its entirety) which would yield a 3' end free of non-PIV nucleotides. The left- and right-hand ends are then joined via a common restriction site.

A variety of nucleotide insertions, deletions and rearrangements can be made
20 in the PIV genome or antigenome during or after construction of the cDNA. For example, specific desired nucleotide sequences can be synthesized and inserted at appropriate regions in the cDNA using convenient restriction enzyme sites. Alternatively, such techniques as site-specific mutagenesis, alanine scanning, PCR mutagenesis, or other such techniques well known in the art can be used to introduce mutations into the cDNA.

Alternative means to construct cDNA encoding the genome or antigenome
25 include reverse transcription-PCR using improved PCR conditions (e.g., as described in Cheng et al., Proc. Natl. Acad. Sci. U.S.A. 91:5695-5699, 1994), incorporated herein by reference) to reduce the number of subunit cDNA components to as few as one or two pieces. In other embodiments different promoters can be used (e.g., T3, SP6) or different
30 ribozymes (e.g., that of hepatitis delta virus. Different DNA vectors (e.g., cosmids) can be used for propagation to better accommodate the larger size genome or antigenome.

Isolated polynucleotides (e.g., cDNA) encoding the genome or antigenome may be inserted into appropriate host cells by transfection, electroporation, mechanical insertion, transduction or the like, into cells which are capable of supporting a productive PIV infection, e.g., HEp-2, FRhL-DBS2, LLC-MK2, MRC-5, and Vero cells. Transfection of isolated polynucleotide sequences may be introduced into cultured cells by, for example, calcium phosphate-mediated transfection (Wigler et al., Cell 14:725, 1978; Corsaro and Pearson, Somatic Cell Genetics 7:603, 1981; Graham and Van der Eb, Virology 52:456, 1973), electroporation (Neumann et al., EMBO J. 1:841-845, 1982), DEAE-dextran mediated transfection (Ausubel et al., ed., Current Protocols in Molecular Biology, John Wiley and Sons, Inc., NY, 1987), cationic lipid-mediated transfection (Hawley-Nelson et al., Focus 15:73-79, 1993) or a commercially available transfection reagent, e.g., LipofectACE® (Life Technologies) or the like (each of the foregoing references are incorporated herein by reference in its entirety).

As noted above, in some embodiments of the invention the N, P, L and other desired PIV proteins are encoded by one or more helper viruses which is phenotypically distinguishable from that which encodes the genome or antigenome. The N, P, L and other desired PIV proteins can also be encoded by one or more expression vectors which can be the same or separate from that which encodes the genome or antigenome, and various combinations thereof. Additional proteins may be included as desired, encoded by its own vector or by a vector encoding one or more of the N, P, L and other desired PIV proteins, or the complete genome or antigenome.

By providing infectious clones of PIV the invention permits a wide range of alterations to be recombinantly produced within the PIV genome (or antigenome), yielding defined mutations which specify desired phenotypic changes. By "infectious clone" is meant cDNA or its product, synthetic or otherwise, RNA capable of being directly incorporated into infectious virions which can be transcribed into genomic or antigenomic RNA capable of serving as a template to produce the genome of infectious viral or subviral particles. As noted above, defined mutations can be introduced by a variety of conventional techniques (e.g., site-directed mutagenesis) into a cDNA copy of the genome or antigenome. The use of genomic or antigenomic cDNA subfragments to assemble a complete genome or antigenome cDNA as described herein has the advantage that each region can be manipulated separately, where small cDNA subjects provide for better ease of manipulation than large cDNA subjects, and then readily assembled into a complete cDNA. Thus, the complete antigenome

or genome cDNA, or a selected subfragment thereof, can be used as a template for oligonucleotide-directed mutagenesis. This can be through the intermediate of a single-stranded phagemid form, such as using the MUTA-gen® kit of Bio-Rad Laboratories (Richmond, CA), or a method using the double-stranded plasmid directly as a template such as the Chameleon® mutagenesis kit of Stratagene (La Jolla, CA), or by the polymerase chain reaction employing either an oligonucleotide primer or a template which contains the mutation(s) of interest. A mutated subfragment can then be assembled into the complete antigenome or genome cDNA. A variety of other mutagenesis techniques are known and can be routinely adapted for use in producing the mutations of interest in a PIV antigenome or genome cDNA of the invention.

Thus, in one illustrative embodiment mutations are introduced by using the MUTA-gene® phagemid *in vitro* mutagenesis kit available from Bio-Rad Laboratories. In brief, cDNA encoding an PIV genome or antigenome is cloned into the plasmid pTZ18U, and used to transform CJ236 cells (Life Technologies). Phagemid preparations are prepared as recommended by the manufacturer. Oligonucleotides are designed for mutagenesis by introduction of an altered nucleotide at the desired position of the genome or antigenome. The plasmid containing the genetically altered genome or antigenome is then amplified.

Mutations can vary from single nucleotide changes to the introduction, deletion or replacement of large cDNA segments containing one or more genes or genome segments. Genome segments can correspond to structural and/or functional domains, e.g., cytoplasmic, transmembrane or ectodomains of proteins, active sites such as sites that mediate binding or other biochemical interactions with different proteins, epitopic sites, e.g., sites that stimulate antibody binding and/or humoral or cell mediated immune responses, etc. Useful genome segments in this regard range from about 15-35 nucleotides in the case of genome segments encoding small functional domains of proteins, e.g., epitopic sites, to about 50, 75, 100, 200-500, and 500-1,500 or more nucleotides.

The ability to introduce defined mutations into infectious PIV has many applications, including the manipulation of PIV pathogenic and immunogenic mechanisms. For example, the functions of PIV proteins, including the N, P, M, F, HN, and L proteins and C, D and V ORF products, can be manipulated by introducing mutations which ablate or reduce the level of protein expression, or which yield mutant protein. Various genome RNA structural features, such as promoters, intergenic regions, and transcription signals, can also

be routinely manipulated within the methods and compositions of the invention. The effects of trans-acting proteins and cis-acting RNA sequences can be readily determined, for example, using a complete antigenome cDNA in parallel assays employing PIV minigenomes (Dimock et al., *J. Virol.*, **67**: 2772-8, 1993, incorporated herein by reference in its entirety), whose rescue-dependent status is useful in characterizing those mutants that may be too inhibitory to be recovered in replication-independent infectious virus.

Certain substitutions, insertions, deletions or rearrangements of genes or genome segments within recombinant PIV of the invention (e.g., substitutions of a genome segment encoding a selected protein or protein region, for instance a cytoplasmic tail, transmembrane domain or ectodomain, an epitopic site or region, a binding site or region, an active site or region containing an active site, etc.) are made in structural or functional relation to an existing, "counterpart" gene or genome segment from the same or different PIV or other source. Such modifications yield novel recombinants having desired phenotypic changes compared to wild-type or parental PIV or other viral strains. For example, recombinants of this type may express a chimeric protein having a cytoplasmic tail and/or transmembrane domain of one PIV fused to an ectodomain of another PIV. Other exemplary recombinants of this type express duplicate protein regions, such as duplicate immunogenic regions.

As used herein, "counterpart" genes, genome segments, proteins or protein regions, are typically from heterologous sources (e.g., from different PIV genes, or representing the same (i.e., homologous or allelic) gene or genome segment in different PIV types or strains). Typical counterparts selected in this context share gross structural features, e.g., each counterpart may encode a comparable protein or protein structural domain, such as a cytoplasmic domain, transmembrane domain, ectodomain, binding site or region, epitopic site or region, etc. Counterpart domains and their encoding genome segments embrace an assemblage of species having a range of size and sequence variations defined by a common biological activity among the domain or genome segment variants.

Counterpart genes and genome segments, as well as other polynucleotides disclosed herein for producing recombinant PIV within the invention, often share substantial sequence identity with a selected polynucleotide "reference sequence," e.g., with another selected counterpart sequence. As used herein, a "reference sequence" is a defined sequence used as a basis for sequence comparison, for example, a segment of a full-length cDNA or

gene, or a complete cDNA or gene sequence. Generally, a reference sequence is at least 20 nucleotides in length, frequently at least 25 nucleotides in length, and often at least 50 nucleotides in length. Since two polynucleotides may each (1) comprise a sequence (i.e., a portion of the complete polynucleotide sequence) that is similar between the two

5 polynucleotides, and (2) may further comprise a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window", as used herein, refers to a conceptual segment of at least 20 contiguous nucleotide positions

10 wherein a polynucleotide sequence may be compared to a reference sequence of at least 20 contiguous nucleotides and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a

15 comparison window may be conducted by the local homology algorithm of Smith & Waterman, (Adv. Appl. Math. 2:482, 1981), by the homology alignment algorithm of Needleman & Wunsch, (J. Mol. Biol. 48:443, 1970), by the search for similarity method of Pearson & Lipman, (Proc. Natl. Acad. Sci. U.S.A. 85:2444, 1988) (each of which is incorporated by reference), by computerized implementations of these algorithms (GAP,

20 BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Dr., Madison, WI, incorporated herein by reference), or by inspection, and the best alignment (i.e., resulting in the highest percentage of sequence similarity over the comparison window) generated by the various methods is selected. The term "sequence identity" means that two polynucleotide sequences are

25 identical (i.e., on a nucleotide-by-nucleotide basis) over the window of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, U, or I) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number

30 of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The terms "substantial identity" as used herein denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 85 percent sequence identity, preferably at least 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared

to a reference sequence over a comparison window of at least 20 nucleotide positions, frequently over a window of at least 25-50 nucleotides, wherein the percentage of sequence identity is calculated by comparing the reference sequence to the polynucleotide sequence which may include deletions or additions which total 20 percent or less of the reference
5 sequence over the window of comparison. The reference sequence may be a subset of a larger sequence.

In addition to these polynucleotide sequence relationships, proteins and protein regions encoded by recombinant PIV of the invention are also typically selected to have conservative relationships, i.e. to have substantial sequence identity or sequence
10 similarity, with selected reference polypeptides. As applied to polypeptides, the term "sequence identity" means peptides share identical amino acids at corresponding positions. The term "sequence similarity" means peptides have identical or similar amino acids (i.e., conservative substitutions) at corresponding positions. The term "substantial sequence
15 identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80 percent sequence identity, preferably at least 90 percent sequence identity, more preferably at least 95 percent sequence identity or more (e.g., 99 percent sequence identity). The term "substantial
20 similarity" means that two peptide sequences share corresponding percentages of sequence similarity. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic
side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic
25 side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine. Abbreviations for the twenty naturally occurring
30 amino acids used herein follow conventional usage (*Immunology - A Synthesis*, 2nd ed., E.S. Golub & D.R. Gren, eds., Sinauer Associates, Sunderland, MA, 1991, incorporated herein by reference). Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α , α -disubstituted amino acids, N-alkyl amino acids,

lactic acid, and other unconventional amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4-hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, ω -N-methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline).
5 Moreover, amino acids may be modified by glycosylation, phosphorylation and the like.

To select candidate vaccine viruses according to the invention, the criteria of viability, attenuation and immunogenicity are determined according to well known methods. Viruses which will be most desired in vaccines of the invention must maintain viability, have
10 a stable attenuation phenotype, exhibit replication in an immunized host (albeit at lower levels), and effectively elicit production of an immune response in a vaccinee sufficient to confer protection against serious disease caused by subsequent infection from wild-type virus. The recombinant PIV of the invention are not only viable and more appropriately attenuated than previous vaccine candidates, but are more stable genetically *in vivo*--
15 retaining the ability to stimulate a protective immune response and in some instances to expand the protection afforded by multiple modifications, e.g., induce protection against different viral strains or subgroups, or protection by a different immunologic basis, e.g., secretory versus serum immunoglobulins, cellular immunity, and the like.

Recombinant PIV of the invention can be tested in various well known and
20 generally accepted *in vitro* and *in vivo* models to confirm adequate attenuation, resistance to phenotypic reversion, and immunogenicity for vaccine use. In *in vitro* assays, the modified virus (e.g., a multiply attenuated, biologically derived or recombinant PIV) is tested, e.g., for temperature sensitivity of virus replication, i.e. its phenotype, and for the small plaque or other desired phenotype. Modified viruses are further tested in animal models of PIV
25 infection. A variety of animal models have been described and are summarized in various references incorporated herein. PIV model systems, including rodents and non-human primates, for evaluating attenuation and immunogenic activity of PIV vaccine candidates are widely accepted in the art, and the data obtained therefrom correlate well with PIV infection, attenuation and immunogenicity in humans.

30 In accordance with the foregoing description, the invention also provides isolated, infectious recombinant PIV compositions for vaccine use. The attenuated virus which is a component of a vaccine is in an isolated and typically purified form. By isolated

is meant to refer to PIV which is in other than a native environment of a wild-type virus, such as the nasopharynx of an infected individual. More generally, isolated is meant to include the attenuated virus as a component of a cell culture or other artificial medium where it can be propagated and characterized in a controlled setting. For example, attenuated PIV
5 of the invention may be produced by an infected cell culture, separated from the cell culture and added to a stabilizer.

For vaccine use, recombinant PIV produced according to the present invention can be used directly in vaccine formulations, or lyophilized, as desired, using lyophilization protocols well known to the artisan. Lyophilized virus will typically be
10 maintained at about 4°C. When ready for use the lyophilized virus is reconstituted in a stabilizing solution, e.g., saline or comprising SPG, Mg⁺⁺ and HEPES, with or without adjuvant, as further described below.

PIV vaccines of the invention contain as an active ingredient an immunogenically effective amount of PIV produced as described herein. The modified virus
15 may be introduced into a host with a physiologically acceptable carrier and/or adjuvant. Useful carriers are well known in the art, and include, e.g., water, buffered water, 0.4% saline, 0.3% glycine, hyaluronic acid and the like. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile solution prior to administration, as mentioned above. The compositions may contain
20 pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, and the like. Acceptable adjuvants include incomplete Freund's adjuvant, MPL™ (3-*o*-deacylated
25 monophosphoryl lipid A; RIBI ImmunoChem Research, Inc., Hamilton, MT) and IL-12 (Genetics Institute, Cambridge MA), among many other suitable adjuvants well known in the art.

Upon immunization with a PIV composition as described herein, via aerosol, droplet, oral, topical or other route, the immune system of the host responds to the vaccine
30 by producing antibodies specific for PIV proteins, e.g., F and HN glycoproteins. As a result of the vaccination with an immunogenically effective amount of PIV produced as described herein, the host becomes at least partially or completely immune to PIV infection, or

resistant to developing moderate or severe PIV infection, particularly of the lower respiratory tract.

The host to which the vaccines are administered can be any mammal which is susceptible to infection by PIV or a closely related virus and which host is capable of generating a protective immune response to the antigens of the vaccinating strain. Accordingly, the invention provides methods for creating vaccines for a variety of human and veterinary uses.

The vaccine compositions containing the PIV of the invention are administered to a host susceptible to or otherwise at risk for PIV infection to enhance the host's own immune response capabilities. Such an amount is defined to be a "immunogenically effective dose." In this use, the precise amount of PIV to be administered within an effective dose will depend on the host's state of health and weight, the mode of administration, the nature of the formulation, etc., but will generally range from about 10^3 to about 10^7 plaque forming units (PFU) or more of virus per host, more commonly from about 10^4 to 10^6 PFU virus per host. In any event, the vaccine formulations should provide a quantity of modified PIV of the invention sufficient to effectively protect the host patient against serious or life-threatening PIV infection.

The PIV produced in accordance with the present invention can be combined with viruses of other PIV serotypes or strains to achieve protection against multiple PIV serotypes or strains. Alternatively, protection against multiple PIV serotypes or strains can be achieved by combining protective epitopes of multiple serotypes or strains engineered into one virus, as described herein. Typically when different viruses are administered they will be in admixture and administered simultaneously, but they may also be administered separately. Immunization with one strain may protect against different strains of the same or different serotype.

In some instances it may be desirable to combine the PIV vaccines of the invention with vaccines which induce protective responses to other agents, particularly other childhood viruses. In another aspect of the invention the PIV can be employed as a vector for protective antigens of other pathogens, such as respiratory syncytial virus (RSV) or measles virus, by incorporating the sequences encoding those protective antigens into the PIV genome or antigenome which is used to produce infectious PIV, as described herein.

In all subjects, the precise amount of recombinant PIV vaccine administered, and the timing and repetition of administration, will be determined based on the patient's state of health and weight, the mode of administration, the nature of the formulation, etc. Dosages will generally range from about 10^3 to about 10^7 plaque forming units (PFU) or more of virus per patient, more commonly from about 10^4 to 10^6 PFU virus per patient. In any event, the vaccine formulations should provide a quantity of attenuated PIV sufficient to effectively stimulate or induce an anti-PIV immune response, e.g., as can be determined by complement fixation, plaque neutralization, and/or enzyme-linked immunosorbent assay, among other methods. In this regard, individuals are also monitored for signs and symptoms of upper respiratory illness. As with administration to chimpanzees, the attenuated virus of the vaccine grows in the nasopharynx of vaccinees at levels approximately 10-fold or more lower than wild-type virus, or approximately 10-fold or more lower when compared to levels of incompletely attenuated PIV.

In neonates and infants, multiple administration may be required to elicit sufficient levels of immunity. Administration should begin within the first month of life, and at intervals throughout childhood, such as at two months, six months, one year and two years, as necessary to maintain sufficient levels of protection against native (wild-type) PIV infection. Similarly, adults who are particularly susceptible to repeated or serious PIV infection, such as, for example, health care workers, day care workers, family members of young children, the elderly, individuals with compromised cardiopulmonary function, may require multiple immunizations to establish and/or maintain protective immune responses. Levels of induced immunity can be monitored by measuring amounts of neutralizing secretory and serum antibodies, and dosages adjusted or vaccinations repeated as necessary to maintain desired levels of protection. Further, different vaccine viruses may be indicated for administration to different recipient groups. For example, an engineered PIV strain expressing a cytokine or an additional protein rich in T cell epitopes may be particularly advantageous for adults rather than for infants.

PIV vaccines produced in accordance with the present invention can be combined with viruses expressing antigens of another subgroup or strain of PIV to achieve protection against multiple PIV subgroups or strains. Alternatively, the vaccine virus may incorporate protective epitopes of multiple PIV strains or subgroups engineered into one PIV clone, as described herein.

The PIV vaccines of the invention elicit production of an immune response that is protective against serious lower respiratory tract disease, such as pneumonia and bronchiolitis when the individual is subsequently infected with wild-type PIV. While the naturally circulating virus is still capable of causing infection, particularly in the upper respiratory tract, there is a very greatly reduced possibility of rhinitis as a result of the vaccination and possible boosting of resistance by subsequent infection by wild-type virus. Following vaccination, there are detectable levels of host engendered serum and secretory antibodies which are capable of neutralizing homologous (of the same subgroup) wild-type virus *in vitro* and *in vivo*. In many instances the host antibodies will also neutralize wild-type virus of a different, non-vaccine subgroup.

Preferred PIV vaccine candidates of the invention exhibit a very substantial diminution of virulence when compared to wild-type virus that is circulating naturally in humans. The virus is sufficiently attenuated so that symptoms of infection will not occur in most immunized individuals. In some instances the attenuated virus may still be capable of dissemination to unvaccinated individuals. However, its virulence is sufficiently abrogated such that severe lower respiratory tract infections in the vaccinated or incidental host do not occur.

The level of attenuation of PIV vaccine candidates may be determined by, for example, quantifying the amount of virus present in the respiratory tract of an immunized host and comparing the amount to that produced by wild-type PIV or other attenuated PIV which have been evaluated as candidate vaccine strains. For example, the attenuated virus of the invention will have a greater degree of restriction of replication in the upper respiratory tract of a highly susceptible host, such as a chimpanzee, compared to the levels of replication of wild-type virus, e.g., 10- to 1000-fold less. In order to further reduce the development of rhinorrhea, which is associated with the replication of virus in the upper respiratory tract, an ideal vaccine candidate virus should exhibit a restricted level of replication in both the upper and lower respiratory tract. However, the attenuated viruses of the invention must be sufficiently infectious and immunogenic in humans to confer protection in vaccinated individuals. Methods for determining levels of PIV in the nasopharynx of an infected host are well known in the literature.

Levels of induced immunity provided by the vaccines of the invention can also be monitored by measuring amounts of neutralizing secretory and serum antibodies.

Based on these measurements, vaccine dosages can be adjusted or vaccinations repeated as necessary to maintain desired levels of protection. Further, different vaccine viruses may be advantageous for different recipient groups. For example, an engineered PIV strain expressing an additional protein rich in T cell epitopes may be particularly advantageous for adults rather than for infants.

In yet another aspect of the invention the PIV is employed as a vector for transient gene therapy of the respiratory tract. According to this embodiment the recombinant PIV genome or antigenome incorporates a sequence which is capable of encoding a gene product of interest. The gene product of interest is under control of the same or a different promoter from that which controls PIV expression. The infectious PIV produced by coexpressing the recombinant PIV genome or antigenome with the N, P, L and other desired PIV proteins, and containing a sequence encoding the gene product of interest, is administered to a patient. Administration is typically by aerosol, nebulizer, or other topical application to the respiratory tract of the patient being treated. Recombinant PIV is administered in an amount sufficient to result in the expression of therapeutic or prophylactic levels of the desired gene product. Representative gene products which may be administered within this method are preferably suitable for transient expression, including, for example, interleukin-2, interleukin-4, gamma-interferon, GM-CSF, G-CSF, erythropoietin, and other cytokines, glucocerebrosidase, phenylalanine hydroxylase, cystic fibrosis transmembrane conductance regulator (CFTR), hypoxanthine-guanine phosphoribosyl transferase, cytotoxins, tumor suppressor genes, antisense RNAs, and vaccine antigens.

The following examples are provided by way of illustration, not limitation. These examples document construction of representative chimeric PIVs bearing one or more heterologous antigenic determinant(s) according to the above described methods. In one example, the HA gene of the measles virus is inserted as an extra gene into one of three gene junctions of a JS wild type or attenuated strain of HPIV3, namely, the N/P, P/M, or HN/L junction, and recombinant chimeric viruses were recovered. Insertion of the measles HA gene at three different positions in the HPIV3 genome illustrates the range of useful constructs for transferring antigenic determinants from foreign pathogens into PIV vectors. Further, it is expected that inserted gene units that are more 3'-leader proximal will be transcribed and expressed at higher levels than the same gene units located more distally, which will allow for closer modulation of heterologous gene expression (Collins et al., 3rd ed. In "Fields Virology", B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L.

Melnick, T. P. Monath, B. Roizman, and S. E. Straus, Eds., Vol. 1, pp. 1205-1243. Lippincott-Raven Publishers, Philadelphia, 1996).

The chimeric rHPIV bearing the measles virus HA insertion in a wild type rHPIV3 background replicated efficiently *in vitro* but was restricted in replication in hamsters compared to that of the rHPIV3 virus from which it was derived. Similarly, the recombinant chimeric HPIV3 bearing the measles virus HA insertion in an attenuated rHPIV3 background replicated *in vitro* and in hamsters to a level that was also slightly less than that of the attenuated rHPIV3cp45L mutant virus from which it was derived. The amount of HA protein expressed by cells infected with the attenuated rHPIV3-measles virus HA recombinants with the HA gene in the N/P or P/M junction was very high and even exceeded that seen in cells infected with native measles virus. The level of replication of the rHPIV3cp45L with a measles virus HA insert in the N/P or P/M junction was 10-fold lower in the upper respiratory tract of the hamster than that of the rHPIV3-cp45L parent virus indicating that gene insertions can unexpectedly contribute to the attenuation of an HPIV3 vector. These results which identify a unique host range phenotype are unexpected.

Importantly, infection of hamsters with each recombinant chimeric virus tested induced high levels of antibody to both HPIV3 and to measles virus. Animals immunized with the attenuated recombinant chimeric HPIV3 carrying the HA insertion were highly resistant to replication of HPIV3 challenge virus. While the wild type measles virus does not replicate efficiently in hamsters and thus cannot be used in challenge study, the protective efficacy of the attenuated recombinant chimeric vaccine is readily apparent from the high levels of neutralizing antibody induced. These levels are associated with a high level of resistance to measles in humans (Chen et al., *J. Infect. Dis.* 162:1036-42, 1990).

It is further demonstrated in the examples that attenuated chimeric recombinant HPIV vectors, combining a backbone of HPIV3 and one or more antigenic determinants of HPIV1, can also be used as vectors to express additional foreign antigens (e.g., of HPIV2 or a non-PIV virus). This aspect of the invention takes advantage of the efficient growth and excellent attenuation properties of the HPIV3 backbone to carry antigenic determinants of multiple heterologous pathogens, as exemplified by HPIV1 and HPIV2. The cDNA encoding rPIV3-1 (a non-attenuated recombinant bearing major antigens of HPIV1) or rPIV3-1cp45 (an attenuated recombinant bearing HPIV1 major antigens) was modified by the insertion of a gene unit containing the ORF of HPIV2 HN gene between the

gene units containing the F and HN ORFs of HPIV1. The recombinant chimeric viruses, designated rPIV3-1.2HN and rPIV3-1cp45.2HN, were readily recovered and replicated efficiently in tissue culture. Each virus exhibited a level of temperature sensitivity of replication *in vitro* similar to that of its rPIV3-1 or rPIV3-1cp45 parent virus. The insertion of the PIV2 HN attenuated both the rPIV3-1 and rPIV3-1cp45 viruses in hamsters, a finding similar to that observed with the insertion of the measles viruses HA into rJS and into rPIV3cp45. Infection of hamsters with these antigenic rPIV3-1 recombinants bearing the PIV2 HN gene insert induced serum antibody responses reactive against both HPIV1 and HPIV2.

Thus, it is possible to use an attenuated rHPIV3 or rHPIV3-1 vaccine candidate as a vector to infect the respiratory tract of susceptible hosts and thereby induce a vigorous antibody response to foreign protective antigens expressed from an extra gene unit, as well as against the HPIV vector itself. The presence of three antigenic serotypes of HPIV, which do not provide significant cross-protection, allows for more effective, sequential immunization of human infants with antigenically distinct variants of HPIV each bearing the same or different heterologous antigenic determinant(s), e.g., a protective antigen, antigenic domain or epitope of measles virus or of one or more different viral or microbial pathogens. Sequential immunization permits development of a primary immune response to the foreign protein, which is boosted during subsequent infections with a secondary, antigenically-distinct HPIV bearing one or more heterologous antigenic determinants, e.g., a protective antigen, antigenic domain or epitope of measles virus or of one or more different viral or microbial pathogens. In this way, the immunity induced to one HPIV vector can be circumvented by boosting with an antigenically distinct HPIV vector. In this context, successful immunization of animals that are immune to PIV3 has been achieved with attenuated PIV3-1 vaccine candidates, confirming the feasibility of sequential immunization with serotypically distinct PIV viruses even if these PIVs share proteins other than HN and F. (Tao et al., *Vaccine* 17:1100-8, 1999). In this study, the immunogenicity and efficacy of rPIV3-1.cp45L against PIV1 challenge was examined in hamsters with and without prior immunity to PIV3. rPIV3-1.cp45L efficiently infected hamsters previously infected with wild type or attenuated PIV3, but there was approximately a five-fold reduction in replication of rPIV3-1.cp45L virus in the PIV3-immune animals. However, rPIV3-1.cp45L immunization of PIV3-immune animals induced a vigorous serum antibody response to PIV1 and reduced replication of PIV1 challenge virus 1000-fold in the lower respiratory tract and 200-fold in the upper respiratory tract. These results demonstrate that the

recombinant chimeric rPIV3-1.cp45L candidate vaccine can induce immunity to PIV1 even in animals immune to PIV3. This establishes the feasibility of employing a sequential immunization schedule in which a recombinant chimeric rPIV3-1.cp45L or other PIV vaccine virus is given following a live attenuated PIV3 vaccine. since rPIV3-1.cp45L readily induced protective immunity against itself, it would also induce an effective immune response to any vectored protective antigen that it was carrying. Also, the PIVs and RSV have the unusual property of being able to reinfect the respiratory tract, although reinfections typically are not associated with serious disease. Thus, vector based vaccine constructs of the invention are useful to boost immune responses by a second, third or fourth administration of the same HPIV vector or by sequential use of different vectors.

In preferred sequential vaccination methods of the invention, it is desirable to sequentially immunize an infant with different PIV vectors each expressing the same heterologous antigenic determinant such as the measles virus HA. This sequential immunization permits the induction of the high titer of antibody to the heterologous protein that is characteristic of the secondary antibody response. In one embodiment, early infants (e.g. 2-4 month old infants) are immunized with an attenuated chimeric HPIV3 expressing a heterologous antigenic determinant, for example the measles virus HA protein, and also adapted to elicit an immune response against HPIV3. One exemplary vaccine candidate useful in this context is the rcp45L(HA P-M) recombinant. Subsequently, e.g., at four months of age the infant is again immunized but with a different, secondary PIV vector construct antigenically distinct from the first. An exemplary vaccine candidate in this context is the rPIV3-1 cp45L virus expressing the measles virus HA gene and HPIV1 antigenic determinants as functional, obligate glycoproteins of the vector. Following the first vaccination, the vaccinee will elicit a primary antibody response to both the PIV3 HN and F proteins and to the measles virus HA protein, but not to the PIV1 HN and F protein. Upon secondary immunization with the rPIV3-1 cp45L expressing the measles virus HA, the vaccinee will be readily infected with the vaccine because of the absence of antibody to the PIV1 HN and F proteins and will develop both a primary antibody response to the PIV1 HN and F protective antigens and a high titered secondary antibody response to the heterologous measles virus HA protein. A similar sequential immunization schedule can be developed where immunity is sequentially elicited against HPIV3 and then HPIV2 by one or more of the chimeric vaccine viruses disclosed herein, simultaneous with stimulation of an initial and then secondary, high titer protective response against measles or another non-PIV pathogen. This sequential immunization strategy, preferably employing different serotypes of PIV as

primary and secondary vectors, effectively circumvents immunity that is induced to the primary vector, a factor ultimately limiting the usefulness of vectors with only one serotype.

Further in accordance with this aspect of the invention, exemplary coordinate vaccination protocols may incorporate two, three, four and up to six or more separate chimeric HPIV vaccine viruses administered simultaneously (e.g., in a polyspecific vaccine mixture) in a primary vaccination step, e.g., at one, two or four months of age. For example, two or more and up to a full panel of HPIV-based vaccine viruses can be administered that separately express one or more antigenic determinants (i.e., whole antigens, immunogenic domains, or epitopes) selected from the G protein of RSV subgroup A, the F protein of RSV subgroup A, the G protein of RSV subgroup B, the F protein of RSV subgroup B, the HA protein of measles virus, and/or the F protein of measles virus. Coordinate booster administration of these same PIV3-based vaccine constructs can be repeated at two months of age. Subsequently, e.g., at four months of age, a separate panel of 2-6 or more antigenically distinct (referring to vector antigenic specificity) live attenuated HPIV-based vaccine viruses can be administered in a secondary vaccination step. For example, secondary vaccination may involve concurrent administration of a mixture or multiple formulations that contain(s) multiple HPIV3-1 vaccine constructs that collectively express RSV G from subgroup A, RSV F from subgroup A, RSV F from subgroup B, RSV G from subgroup B, measles virus HA, and/or measles virus F, or antigenic determinants from any combination of these proteins. This secondary immunization provides a boost in immunity to each of the heterologous RSV and measles virus proteins or antigenic determinant(s) thereof. At six months of age, a tertiary vaccination step involving administration of one-six or more separate live attenuated PIV3-2 vector-based vaccine recombinants can be coordinately administered that separately or collectively express RSV G from subgroup A, RSV F from subgroup A, RSV G from subgroup B, RSV F from subgroup B, measles virus HA, and/or measles virus F, or antigenic determinant(s) thereof. Optionally at this step in the vaccination protocol, rPIV3 and rPIV3-1 vaccines may be administered in booster formulations. In this way, the strong immunity characteristic of secondary antibody to PIV1, PIV2, PIV3, RSV A, RSV B, and measles viruses are all induced within the first six months of infancy. Such a coordinate/sequential immunization strategy, which is able to induce secondary antibody responses to multiple viral respiratory pathogens, provides a highly powerful and extremely flexible immunization regimen that is driven by the need to immunize against each of the three PIV viruses and other pathogens in early infancy.

In other aspects of the invention, insertion of heterologous nucleotide sequences into HPIV vaccine candidates are employed separately to modulate the level of attenuation of candidate vaccine recombinants, e.g., for the upper respiratory tract. Thus, it is possible to insert nucleotide sequences into a rHPIV that both direct the expression of a foreign protein and that attenuate the virus in an animal host, or to use nucleotide insertions separately to attenuate candidate vaccine viruses. To define some of the rules that govern the effect of gene insertion on attenuation, gene units of varying lengths were inserted into a wild type HPIV3 backbone and the effects of gene unit length on attenuation were examined. These novel gene unit insertions were engineered to not contain a significant ORF which permitted an evaluation of the effect of gene unit length independently of an effect of the expressed protein of that gene. These heterologous sequences were inserted as an extra gene unit of sizes between 168 nt and 3918 nt between the HN and L genes. In addition, control cDNA constructions and viruses were made in which insertions of similar sizes were placed in the 3'-noncoding region of the HN gene and hence did not involve the addition of an extra gene. These viruses were made to assess the effect of an increase in the overall genome length and in gene number on attenuation. The insertion of an extra gene unit is expected to decrease the transcription of genes downstream of the insertion site which will affect both the overall abundance and ratios of the expressed proteins. As demonstrated herein, gene insertions or extensions larger than about 3000 nts in length attenuated the wild type virus for the upper and lower respiratory tract of hamsters. Gene insertions of about 2000 nts in length further attenuated the rHPIV3cp45L vaccine candidate for the upper respiratory tract. In summary, gene insertions can have the dual effect of both attenuating a candidate vaccine virus and inducing a protective effect against a second virus. Gene extensions in the 3'-noncoding region of a gene, which cannot express additional proteins, can also be attenuating in and of themselves. Within these methods of the invention, gene insertion length is a determinant of attenuation.

GU and NCR insertions within recombinant PIV of the invention produce an attenuation phenotype characterized by efficient replication *in vitro* and decreased replication *in vivo*, a phenotype not previously described for other paramyxovirus insertions. The mechanism of attenuation resulting from a GU insertion may result from one or more of the following factors acting predominantly *in vivo*. The addition of an extra gene unit may decrease the level of transcription of downstream genes since there is a transcriptional gradient in which more promoter-proximal genes are transcribed at a higher rate than the more promoter-distal genes. The decreased expression of the downstream gene products

resulting from the decreased abundance of their mRNAs could result in attenuation if their gene product is limiting or if a specific ratio of gene products that is required for efficient replication is altered. It is thought that the transcription gradient is a consequence of the transcriptase complex falling off the template during transcription as well as during the transfer across gene junctions. Alternatively, the increase in the overall length of the genome and the extra mRNAs transcribed may increase the level of viral double stranded RNA made which in turn may induce a higher level of the antiviral activity of the interferon system. Finally, the overall level of genome replication may be reduced due to the increase in length of the genome and the antigenome. This may result from a disengagement of replicase complexes from the template during replication of the genomic RNA or antigenomic RNA. The decreased amount of genome available for packaging into virions may result in a decrease in virus yield which results in attenuation.

The mechanism of attenuation resulting from a NCR insertion may result from one or more of the following factors. The extra length of the 3'-end of HN mRNA resulting from the NCR insertion may contribute to the instability of the mRNA and lead to a decrease in the expression of the HN protein. Alternatively, the increase in the overall length of the genome and the extra length of the HN mRNA may increase the level of viral double stranded RNA made that can induce a higher level of the antiviral activity of the interferon system. Alternatively or additionally, the overall level of genome replication may be reduced due to the increase in length of the genome and the antigenome. This may result from a disengagement of replicase complexes from the template during replication of the genomic RNA or antigenomic RNA. The decreased amount of genome available for packaging into virions could result in a decrease in virus yield which results in attenuation. Finally, the addition of extra nucleotides to the 3' end of the HN gene could decrease the level of transcription of downstream genes since the transcriptase complex could fall off the template during transcription of the extra nucleotides at the 3' end of the HN gene.

The *in vitro* and *in vivo* growth properties of the GU and NCR insertions into PIV3 are distinct from previous findings with other single-stranded, negative-sense RNA viruses, cited above. Previously tested insertions examined expressed proteins, whereby the independent effect of the length of insertions on viral growth *in vivo* cannot be determined. The present findings demonstrate that the GU and NCR insertions greater than 3 kb specify an attenuation phenotype that is independent of expressed protein. Shorter insertions, e.g., greater than about 2 kb, specify further attenuation in a partially attenuated recipient. Also

unexpectedly, the GU and NCR insertions specify restricted replication *in vivo* in the absence of restricted replication *in vitro*. In addition, the attenuation phenotype *in vivo* is seen when the insertion is either in the form of a GU or a NCR insertion – other documented insertions are in the form of GU only. Thus, the attenuation of replication *in vivo* specified
 5 by a GU or NCR insertion that does not encode a protein represents a unique way to attenuate members of the Mononegavirales *in vivo*.

EXAMPLE I

Construction of cDNAs Encoding a Chimeric HPIV3/Measles Virus-HA

10 Antigenome and Recovery of Infectious Virus

The full-length cDNA clones, p3/7(131)2G+, encoding the complete 15462 nucleotide antigenome of the JS PIV3 wt virus, and pFLCcp45L, which encodes the antigenome of the derivative of JS wt containing three *cp45*-specific temperature-sensitive mutations in the L ORF of PIV3, have been previously described (Durbin et al., Virology
 15 235:323-332, 1997a; Skiadopoulos et al., J. Virol. 72:1762-8, 1998, each incorporated herein by reference). These clones were used as vectors for the insertion of the HA gene of measles virus to create both wildtype and attenuated HPIV3 chimeric constructs which express a heterologous antigenic determinant, exemplified by the HA protein, of measles virus. The size of each insert containing the HA gene of measles was a multiple of six such that the
 20 chimeric virus recovered from the cDNA would conform to the rule of six (Durbin et al., Virology 234:74-83, 1997b, incorporated herein by reference).

Construction of full-length chimeric HPIV3 cDNAs encoding the HA protein of measles virus in the N/P or P/M junctions.

The *Pml* to *Bam*HI fragment of p3/7(131)2G+ (nt 1215-3903 of the PIV3
 25 antigenome) was subcloned into the plasmid pUC119 {pUC119(*Pml*-*Bam*HI)} which had been modified to include a *Pml* site in the multiple cloning region. Two independent single-stranded mutagenesis reactions were performed on pUC119(*Pml* -*Bam*HI) using Kunkel's method (Kunkel et al., Methods Enzymol. 154:367-382, 1987, incorporated herein by reference); the first reaction introduced an *Afl*II site in the 3' (downstream)-noncoding region
 30 of the N gene by mutating the CTAAAT sequence at nts 1677-1682 of the antigenome to CTTAAG (p*Afl*II N-P), and the second, separate, reaction introduced an *Afl*II site in the in

the 3'-noncoding region of the P gene by mutating the TCAATC sequence at nts 3693-3698 of the antigenome to CTTAAG (p*Afl*II P-M).

- The HA ORF of measles virus Edmonston strain was amplified from Edmonston wild type virus by reverse transcription polymerase chain reaction (RT-PCR).
- 5 The nt sequence of the Edmonston wild type HA open reading frame (ORF) is in GenBank Accession # U03669, incorporated herein by reference (note that this sequence is the ORF only without the upstream 3 nts or the stop codon). Measles virus RNA was purified from clarified medium using TRIzol-LS (Life Technologies, Gaithersburg, MD) following the manufacturer's recommended procedure. RT-PCR was performed with the Advantage RT-
- 10 for-PCR and Advantage-HF PCR kits (Clontech, Palo Alto, CA) following the recommended protocols. Primers were used to generate a PCR fragment spanning the entire ORF of the measles virus HA gene flanked by PIV3 non-coding sequence and *Afl*II restriction sites. The forward primer 5'-
- TTAATCTTAAGAATATACAAATAAGAAAACTTAGGATTAAGAGCGATGTCAC***
- 15 ***CACAACGAGACCGGATAAATGCCTTCTAC***-3' (SEQ ID NO. 13) encodes an *Afl*II site (italicized) upstream of PIV3 noncoding sequence derived from the N/P gene junction-nts 3699-3731 (underlined), containing GE, IG and GS sequences (Figure 1A) and the beginning of the measles HA ORF (bolded) preceded by three non-HPV3, non-measles virus nts designated in the primer. The reverse primer 5'-
- 20 ***ATTATTGCTTAAGGTTTGTTCGGTGTGCGTTTCCTTGTGGATCCTATCTGCGATTG***
- GTTCCATCTTC***-3' (SEQ ID NO. 14) encodes an *Afl*II site (italicized) downstream (in the positive-sense complement) of PIV3 noncoding sequence derived from the P gene, nt 3594-3623 (underlined), and the end of the measles HA ORF (bolded). The resultant PCR
- 25 fragment was then digested with *Afl*II and cloned into p(*Afl*II N-P) and p(*Afl*II P-M) to create pUC119(HA N-P) and pUC119(HA P-M) respectively. pUC119(HA N-P) and pUC119(HA P-M) were sequenced over the entire *Afl*II insert using dRhodamine Terminator Cycle Sequencing Ready Reaction (ABI prism, PE Applied Biosystems, Foster city, CA), and the sequence was confirmed to be correct.

- The *Pml*I to *Bam*HI fragments of pUC119(HA N-P) and pUC119(HA P-M)
- 30 were separately cloned into the full-length antigenome cDNA plasmid p3/7(131)2G+ as previously described (Durbin et al., *Virology* 235:323-332, 1997a, incorporated herein by reference) to create pFLC(HA N-P) and pFLC(HA P-M) (Figure 1). The *Xho*I-*Ngo*MI fragment (nt 7437-15929) of pFLCcp45L was then cloned into the *Xho*I-*Ngo*MI window of

both pFLC(HA N-P) and pFLC(HA P-M) to create pFLCcp45L(HA N-P) and pFLCcp45L(HA P-M). pFLCcp45L encodes the three amino acid changes in the L gene of PIV3 cp45 (aa position 942, 992, and 1558) which confer most of the temperature-sensitivity and attenuation of the cp45 vaccine candidate virus (Skiadopoulos et al., *J. Virol.* 72:1762-8, 1998, incorporated herein by reference), and the transfer of the *XhoI-NgoMI* fragment transferred those mutations.

Construction of full-length HPIV3 chimeric cDNAs encoding the HA protein of measles in the HN/L junction

A HPIV3 chimeric cDNA was constructed by PCR to include a heterologous polynucleotide sequence, exemplified by the measles virus HA gene, encoding a heterologous antigenic determinant of the measles virus, flanked by the transcription signals and the noncoding regions of the HPIV3 HN gene. This cDNA was designed to be combined with an rPIV3 vector as an extra gene following the HN gene. First, using Kunkel mutagenesis (Kunkel et al., *Methods Enzymol.* 154:367-382, 1987, incorporated herein by reference), a *StuI* site was introduced in the 3'-noncoding region of the HN gene by mutating the AGACAA sequence at nts 8598-8603 of the antigenome to AGGCCT yielding plasmid p3/7(131)2G-Stu (Figure 1B). A cDNA containing the measles HA ORF flanked by HPIV3 sequences (see Figure 1B) was then constructed in three pieces by PCR. The first PCR synthesized the left-hand, upstream piece of the gene. The forward primer 5'-
 20 GACAATAGGCCTAAAAGGGAAATATAAAAACTTAGGAGTAAAGTTACGCAAT
CC-3' (SEQ ID NO. 15) contains a *StuI* site (italicized) followed by HPIV3 sequence (underlined) which includes the downstream end of the HN gene (HPIV3 nts 8602-8620), an intergenic region, and the gene-start signal and sequence from the upstream end of the HN gene (HPIV3 nt 6733-6753). The reverse primer 5'-
 25 **GTAGAACGCGTTTATCCGGTCTCGTTGTGGTGACATCTCGAATTTGGATTTGT**
CTATTGGTCCCTCC-3' (SEQ ID NO. 16) contains an *MluI* site (italicized) downstream of the start of the measles HA ORF (bolded) followed by the complement to HPIV3 nts 6744-6805 (underlined), which are part of the upstream HN noncoding region. The *MluI* site present in the introduced measles virus ORF was created by changing nt 27 from T (in the wild type Edmonston HA gene) to C and nt 30 from C to G. Both of these changes are
 30 noncoding in the measles virus ORF. The PCR was performed using p3/7(131)2G-Stu as template. The resulting product, termed PCR fragment 1, is flanked by a *StuI* site at the 5'-end and an *MluI* site at the 3'-end and contains the first 36 nt of the measles HA ORF

downstream of noncoding sequence from the HPIV3 HN gene. The second PCR reaction synthesized the right-hand end of the HN gene. The forward primer **GTAGAACGCGTTATCCGGTCTCGTTGTGGTGACATCTCGAATTGGATTTGTTCTATTGGGTCCTCC-3'** (SEQ ID NO. 16) contains the *Xma*I (italics) and the end of the measles HA ORF (bold), followed by HPIV3 nts 8525-8566 (underlined) representing part of the downstream nontranslated region of the HN gene. The reverse primer 5'-CCATGTAATTGAATCCCCCAACTAGC-3', (SEQ ID NO. 17) spans HPIV3 nts 11448-11475, located in the L gene. The template for the PCR was p3/7(131)2G-Stu. PCR fragment 2 which resulted from this reaction contains the last 35 nt of the measles HA ORF and approximately 2800 nt of the L ORF of PIV3 and is flanked by an *Xma*I site and an *Sph*I site (which occurs naturally at HPIV3 position 11317). The third PCR reaction amplified the largest, central portion of the measles HA ORF from the template cDNA pTM-7, a plasmid which contains the HA ORF of the Edmonston strain of measles virus supplied by the ATCC. Sequence analysis of this plasmid showed that the measles virus HA ORF contained in PTM-7 contains 2 amino acid differences from pTM-7 of the Edmonston wild type HA sequence used for insertion into the N-P and M-P junction, and these were at amino acid positions 46 (F to S) and at position 481 (Y to N). The forward primer 5'-CGGATAAACGCGTTCTACAAAGATAACC-3' (SEQ ID NO. 18) (*Mlu*I site italicized) and reverse primer 5'-CGGATAAACGCGTTCTACAAAGATAACC-3' (SEQ ID NO. 18) (*Xma*I site italicized) amplified PCR fragment 3 which contained nts 19-1838 of the measles HA ORF. To assemble the pieces, PCR fragment 1 was digested with *Stu*I and *Mlu*I while PCR fragment 3 was digested with *Mlu*I and *Xma*I. These two digested fragments were then cloned by triple ligation into the *Stu*I-*Xma*I window of pUC118 which had been modified to include a *Stu*I site in its multiple cloning region. The resultant plasmid, pUC118(HA 1+3) was digested with *Stu*I and *Xma*I while PCR fragment 2 was digested with *Xma*I and *Sph*I. The two digested products were then cloned into the *Stu*I-*Sph*I window of p3/7(131)2G-Stu, resulting in the plasmid pFLC(HA HN-L). The *Stu*I-*Sph*I fragment, including the entire measles HA ORF, was then sequenced using dRhodamine Terminator Cycle Sequencing Ready Reaction (ABI prism, PE Applied Biosystems, Foster city, CA). The chimeric construct sequence was confirmed. In this way, the measles virus HA ORF flanked by HPIV3 transcription signals was inserted as an extra gene into the N/P, P/M, or HN/L junction of an antigenomic cDNA vector comprising a wild type HPIV3 or into the N/P or P/M junction of an antigenomic cDNA vector comprising an attenuated HPIV3.

Recovery of chimeric rPIV3 wild type and *rcp45L* expressing the HA protein of measles virus

The five full-length vector cDNAs bearing the measles HA ORF as a separate gene were transfected separately into HEP-2 cells on six-well plates (Costar, Cambridge, MA) together with the support plasmids {pTM(N), pTM(P no C), and pTM(L)}, and LipofectACE (Life Technologies), and the cells were simultaneously infected with MVA-T7, a replication-defective vaccinia virus recombinant encoding the bacteriophage T7 polymerase protein as previously described (Durbin et al., *Virology* 235:323-332, 1997; Durbin et al., *Virology* 234:74-83, 1997, each incorporated herein by reference). pTM(P no C) is a derivative of pTM(P) (Durbin et al., *Virology* 261:319-330, 1999) in which the C ORF expression has been silenced by mutation of the C start codon. After incubation at 32°C for three days, the transfection harvest was passaged onto a fresh monolayer of Vero cells in a T25 flask and incubated for 5 days at 32°C (referred to as passage 1). The presence of HPIV3 in the passage 1 harvest was determined by plaque titration on LLC-MK2 monolayer cultures with plaques visualized by immunoperoxidase staining with HPIV3 HN-specific and measles HA-specific monoclonal antibodies as previously described (Durbin et al., *Virology* 235:323-332, 1997, incorporated herein by reference).

The rPIV3(HA HN-L) virus present in the supernatant of the appropriate passage 1 harvest was biologically-cloned by plaque purification three times on LLC-MK2 cells as previously described (Hall et al., *Virus Res.* 22:173-184, 1992, incorporated herein by reference). rPIV3(HA N-P), *rcp45L*(HA N-P), rPIV3(HA P-M), and *rcp45L*(HA P-M) were biologically-cloned from their respective passage 1 harvests by terminal dilution using serial 2-fold dilutions on 96-well plates (12 wells per dilution) of Vero cell monolayers. The biologically-cloned recombinant viruses from the third round of plaque purification or from the second or third round of terminal dilution were then amplified twice in LLC-MK2 cells {rPIV3(HA HN-L) or Vero cells {rPIV3(HA N-P), *rcp45L*(HA N-P), rPIV3(HA P-M), *rcp45L*(HA P-M)} at 32°C to produce virus for further characterization. As a first step in confirming and characterizing the recombinant chimeric PIV3s expressing the HA glycoprotein of measles virus, each passage 1 harvest was analyzed by RT-PCR using three different primer pairs; one pair for each location of the HA ORF insert. The first primer pair amplified a fragment of PIV3 spanning nucleotides 1596-1968 of the full-length HPIV3 genome, which includes the N/P insertion site. This fragment size increased to 2298 nucleotides with the measles HA ORF inserted between the N and P genes. The second

primer pair amplified a fragment of PIV3 spanning nucleotides 3438-3866 of the full-length HPIV3 genome, which includes the P/M insertion site. With the measles HA ORF inserted between the P and M genes, this fragment size increased to 2352 nucleotides. The third primer pair amplified a fragment of PIV3 spanning nucleotides 8466-8649 of the full-length antigenome. With the measles HA ORF inserted between the HN and L genes, this fragment size increased to 2211 nucleotides, which includes the HN/L insertion site. All five recovered viruses contained an insert of the appropriate size at the appropriate location. The generation of each PCR product was dependent upon the inclusion of reverse transcriptase, indicating that each was derived from RNA and not from contaminating cDNA.

10 Monolayers of LLC-MK2 cells in T25 flasks were infected at a multiplicity of infection (MOI) of 5 with either *rcp45L(HA N-P)*, *rcp45L(HA P-M)*, rJS or were mock infected. Monolayers of Vero cells in T25 flasks were infected with the Edmonston wild type strain of measles virus at an MOI of 5. Vero cell monolayers were chosen for the measles Edmonston virus infection because measles virus does not grow well in LLC-MK2

15 cells. At 24 hours post-infection, the monolayer was washed with methionine-minus DMEM (Life Technologies). ³⁵S methionine was added to DMEM-minus media at a concentration of 10uCi/ml and 1 ml was added to each flask which was then incubated at 32°C for 6 hours. The cells were harvested and washed 3 times in PBS. The cell pellets were resuspended in 1 ml RIPA buffer {1% (w/v) sodium deoxycholate, 1% (v/v) Triton X-100 (Sigma), 0.2%

20 (w/v) SDS, 150mM NaCl, 50mM Tris-HCl, pH 7.4}, freeze-thawed and clarified by centrifugation at 6500 X G for 5 minutes. The cell extract was transferred to a fresh eppendorf tube and a mixture of monoclonal antibodies which recognizes the HA glycoprotein of measles virus (79-XV-V17, 80-III-B2, 81-1-366) (Hummel et al., J. Virol. 69:1913-6, 1995; Sheshberadaran et al., Arch. Virol. 83:251-68, 1985, each incorporated

25 herein by reference) or which recognizes the HN protein (101/1, 403/7, 166/11) of PIV3 (van Wyke Coelingh et al., Virology 160:465-72, 1987, incorporated herein by reference) was added to each sample and incubated with constant mixing for 2 hours at 4°C. Immune complexes were precipitated by adding 200 µl of a 10% suspension of protein A Sepharose beads (Sigma, St. Louis, MO) to each sample followed by constant mixing at 4°C overnight.

30 Each sample was suspended in 90 µl of 1X loading buffer and 10 µl of reducing agent was added. After heating at 70°C for 10 minutes, 20 µl of each sample was loaded onto a 4-12% polyacrylamide gel (NuPAGE, Novex, San Diego, CA) per the manufacturer's recommendations. The gel was dried and autoradiographed (Figure 2). *rcp45L(HA P-M)*

and *rcp45L(HA N-P)* encoded a protein precipitated by the anti-measles HA monoclonal antibodies which was the same size as the authentic measles HA protein. *rcp45L(HA P-M)* and *rcp45L(HA N-P)* expressed the measles virus HA protein to a greater extent than did the Edmonston wild type strain of measles virus indicating that these constructs efficiently expressed the measles virus HA from the N/P and P/M junctions of the attenuated strain *rcp45L*. *rcp45L(HA N-P)* and *rcp45L(HA P-M)* were confirmed to be HPIV3-based by their reactivity with the PIV3 anti-HN monoclonal antibodies.

The temperature sensitivity of replication of rPIV3 parent and rPIV3(HA) chimeric viruses *in vitro*

The level of temperature sensitivity of replication of the chimeric rPIV3s bearing the measles virus HA insertion was evaluated to assess whether acquisition of the HA insert modified the level of replication in the chimeric virus compared to the parental, vector virus at various temperatures (Table 1). Serial 10-fold dilutions of *rcp45L*, *rcp45L(N-P)*, *rcp45L(HA P-M)*, rPIV3(HA HN-L), rPIV3(HA P-M), or rJS were carried out in L-15 supplemented with 5% FBS, 4 mM glutamine, and 50 µg/ml gentamicin on LLC-MK2 cell monolayers in 96 well plates and incubated at 32, 36, 37, 38, 39, or 40°C for 6 days. Virus was detected by hemadsorption and reported as log₁₀TCID₅₀/ml. Interestingly, chimeric derivatives of both wild type vector viruses bearing the measles virus HA gene, rPIV3(HA HN-L) and rPIV3(HA P-M), were slightly restricted in replication at 40°C (Table 1). The two attenuated rPIV3s bearing the measles virus HA gene, *rcp45L(N-P)* and *rcp45L(HA P-M)*, possessed a level of temperature sensitivity similar to that of the *rcp45L* parental, vector virus with *rcp45L(HA P-M)* being slightly more ts than its parent. Thus, the viruses bearing the inserts replicated in tissue culture similarly to the parental vector rPIV3 from which they were derived, with only a slight increase in temperature sensitivity. These results indicate that rPIV3 can readily serve as a vector to accommodate the HA insert at different sites without major alteration in replication *in vitro*, and that rPIV3(HA) chimeric viruses can readily accommodate the further addition of one or more attenuating mutations.

Table 1. Replication at permissive and elevated temperatures of recombinant HPIV3s expressing the HA protein of measles virus as an extra gene in the N-P, P-M, or HN-L junctions.

Virus	Virus titer (\log_{10} TCID ₅₀ /ml) at indicated temperature					
	32°C ¹	36°C	37°C	38°C	39°C	40°C
<i>rcp45L</i> ²	8.2	8.2	7.2	<u>5.2</u> ⁶	3.4	3.0
<i>rcp45L</i> (HA P-M) ³	7.4	6.7	<u>5.2</u>	4.2	1.4	1.4
<i>rcp45L</i> (HA N-P) ³	7.4	7.2	5.7	<u>4.2</u>	2.2	≤1.2
rPIV3(HA HN-L) ⁴	7.7	8.2	7.0	7.7	6.7	<u>5.2</u>
rPIV3(HA P-M) ⁴	7.7	7.4	6.7	6.2	6.2	<u>4.7</u>
PIV3-rJS ⁵	8.7	9.0	9.0	8.4	8.2	9.0

1. Permissive temperature.
2. Recombinant *ts* derivative of the JS wild type strain of HPIV3, bearing 3 attenuating amino acid substitutions derived from *cp45*.
3. Recombinant attenuated *ts* derivative of JS wild type HPIV3 expressing the HA protein of measles virus.
4. Recombinant wild type HPIV3 expressing the HA protein of measles virus.
5. Recombinant wild type HPIV3, strain JS.
6. Underlined titer represents the lowest restrictive temperature at which a 100-fold or greater reduction in titer from that at 32°C is seen and defines the shut-off temperature of the virus.

EXAMPLE IIChimeric rPIV3s Bearing an Antigenic Determinant of Measles Virus
Replicate Efficiently in Hamsters and Induce High Titers of Antibodies Against Both HPIV3
and Measles**Determination of the level of replication and immunogenicity of the rPIV3(HA) viruses
in hamsters**

The levels of replication of chimeric rPIV3s bearing an antigenic determinant of the measles virus was compared with that of their parent rPIV3s to determine if the acquisition of the determinant, exemplified by an HA insert, significantly modified their ability to replicate and to induce an immune response *in vivo*. In two different experiments, groups of 6 or 7 4-6 week-old Golden Syrian hamsters were inoculated intranasally with 0.1 ml of EMEM (Life Technologies) containing $10^{6.0}$ PFU of rJS, *rcp45L*, *rcp45L*(HA P-M), *rcp45L*(HA N-P), rPIV3(HA HN-L), or rPIV3(HA P-M) (Tables 2 and 3). On day 4 post-inoculation the hamsters were sacrificed and the lungs and nasal turbinates were harvested. The nasal turbinates and lungs were homogenized in 10% or 20% w/v suspension of L-15 (Quality Biologicals, Gaithersburg, MD) respectively, and the samples were rapidly frozen. Virus present in the samples was titered on 96 well plates of LLC-MK2 cell monolayers and incubated at 32°C for 7 days. Virus was detected by hemadsorption, and the mean \log_{10} TCID₅₀/g was calculated for each group of hamsters. Insertion of the HA gene into wild type rJS (Table 2) restricted its replication 4 to 20-fold in the upper respiratory tract and up to five-fold in the lower respiratory tract indicating only a slight effect of the acquisition of the HA gene on replication of wild type rJS virus in hamsters. The replication of each of the two *rcp45*(HA) antigenic chimeras was 10-fold less in the upper respiratory tract of hamsters (Table 3)-than that of *rcp45L*, the recombinant parent virus bearing the three attenuating mutations in the L protein, but was the same as the *rcp45L* parent in the lower respiratory tract. Thus, for each of the two *rcp45*(HA) antigenic chimeras there was a slight, but statistically significant, reduction in replication in the upper respiratory tract of hamsters indicating that the acquisition of the HA gene by *rcp45L* increased its attenuation for the upper, but not the lower, respiratory tract. Thus, the effect of the insertion of the HA gene on the replication of wild type or attenuated PIV3 was comparable in the upper respiratory tract.

Table 2: Replication of wildtype rPIV3(HA) chimeric viruses in the upper and lower respiratory tract of hamsters

Virus ¹	# Animals	Virus Titer (\log_{10} TCID ₅₀ /gm \pm S.E. ²) [Tukey-Kramer Grouping] ³	
		Nasal Turbinates	Lungs
rcp45L	8	4.0 \pm 0.1[A]	1.5 \pm 0.1[A]
rPIV3(HA N-P)	8	5.1 \pm 0.1[B]	5.9 \pm 0.1[B]
rPIV3(HA P-M)	8	5.9 \pm 0.1[C]	6.7 \pm 0.2[C]
rPIV3(HA HN-L)	8	5.9 \pm 0.2[C]	5.8 \pm 0.1[B]
rJS	8	6.5 \pm 0.1[D]	6.6 \pm 0.2[C]

1. Animals received 10^6 TCID₅₀ of the indicated virus given intranasally in a 0.1 ml inoculum and the lungs and nasal turbinates were harvested 4 days later.
2. Standard Error.
3. Mean virus titers were assigned to statistically similar groups (A-D) by the Tukey-Kramer test. Therefore, means in each column with different letters are significantly different ($\alpha=0.05$) and those with the same letter are not significantly different.

Table 3: Replication of the rPIV3cp45L(HA) antigenic chimeric viruses in the upper and lower respiratory tract of hamsters

Virus ¹	#Animals	Virus Titer (\log_{10} TCID ₅₀ /gm \pm S.E. ²) [Tukey-Kramer Grouping] ³	
		Nasal Turbinates	Lungs
rcp45L	6	4.7 \pm 0.2[A]	2.9 \pm 0.1[A]
rcp45L(HA N-P)	6	3.7 \pm 0.2[B]	2.9 \pm 0.1[A]
rcp45L (HA P-M)	7	3.7 \pm 0.1[B]	2.9 \pm 0.2[A]
rJS	7	6.5 \pm 0.1[C]	5.6 \pm 0.2[B]

1. Animals received 10^6 pfu of the indicated virus given intranasally in a 0.1 ml inoculum and the lungs and nasal turbinates were harvested 4 days later.
2. Standard Error.
3. Mean virus titers were assigned to statistically similar groups (A-D) by the Tukey-Kramer test. Therefore, means in each column with different letters are significantly different ($\alpha=0.05$) and those with the same letter are not significantly different.

The ability of the chimeric rHPIV3(HA) viruses to induce an immune response to HPIV3 and to measles virus was studied next. Groups of 6-24 Golden Syrian hamsters (age 4-6 weeks) were infected as described above with either $10^{6.0}$ PFU rJS, rPIV3(HA P-M), *rcp45L*, *rcp45L*(HA P-M), or *rcp45L*(HA N-P) (Table 4) on day 0. Serum

5 was collected from each hamster on day -1 and on day 25 post-inoculation. The serum antibody response to HPIV3 was evaluated by hemagglutination-inhibition (HAI) assay as previously described (van Wyke Coelingh et al., *Virology* 143:569-582, 1985, incorporated herein by reference), and the serum antibody response to measles virus was evaluated by

10 60% plaque-reduction assay as previously described (Coates et al., *Am. J. Epidemiol.* 83:299-313, 1966, incorporated herein by reference). These results were compared with that from an additional control group of cotton rats that received $10^{5.0}$ of the live-attenuated measles virus (Moraten strain) administered intramuscularly on day 0. Cotton rats, rather than hamsters, were used in this group because measles virus is only weakly infectious for hamsters. As can be seen in Table 4, each of the PIV3(HA) chimeric viruses was able to

15 elicit a robust serum neutralizing antibody response against measles virus. There was no significant difference between the amount of serum neutralizing antibody elicited by the attenuated derivative *rcp45L*(HA P-M) as compared to its counterpart in the wild type background, rPIV3(HA P-M). Furthermore, the level of measles virus-neutralizing serum antibodies induced by the rPIV3(HA) recombinants were on average 5-fold greater than that

20 achieved by the intramuscular immunization with the live attenuated measles virus vaccine. In addition, the serum antibody response to HPIV3 produced by all the chimeric viruses was also robust and comparable to that produced by infection with wild type rJS.

Table 4. rPIV3(HA) antigenic chimeric viruses elicit an excellent serum antibody response to both measles virus and PIV3

Virus ¹	# Animals	Serum antibody titer to measles virus (60% plaque reduction neutralization titer, mean reciprocal log ₂ ±S.E. ²)		Serum antibody response to HPIV3 (HAI titer; mean reciprocal log ₂ ±S.E.)	
		Day 0	Day 25	Day 0	Day 25
<i>rcp45L</i> ³	18	≤3.3±0	≤3.3±0	≤2.0±0	10.7±0.2
<i>rcp45L</i> (HA P-M) ⁴	24	≤3.3±0	12.8±0.1	≤2.0±0	9.2±0.2
<i>rcp45L</i> (HA N-P) ⁵	6	≤3.3±0	13.4±0.4	≤2.0±0	10.8±0.3
rPIV3(HA P-M) ⁶	6	≤3.3±0	13.3±0.3	≤2.0±0	10.3±0.2
Measles virus (Moraten) ⁷	4	≤3.3±0	10.8±0.2	≤2.0±0	≤2.0±0
rJS ⁸	6	≤3.3±0	≤3.3±0	≤2.0±0	10.7±0.2

1. Virus was administered at a dose of 10^{6.0} PFU in a 0.1 ml inoculum intranasally on day 0 to all animals with the exception of those in the measles virus group which received virus by intramuscular injection.
2. Standard Error.
3. Recombinant attenuated HPIV3 with three temperature sensitive (*ts*) mutations in the L protein, derived from *cp45*.
4. Recombinant attenuated HPIV3 in the *cp45L* background with the HA ORF of measles virus in the P/M noncoding region of rPIV3.
5. Recombinant attenuated HPIV3 in the *cp45L* background with the HA ORF of measles virus in the N/P noncoding region of rPIV3.
6. Recombinant HPIV3 with the HA ORF of measles virus in the P/M noncoding region of wild type rPIV3.
7. The live attenuated measles vaccine virus, Moraten strain, was administered at a dose of 10⁵ pfu in a 0.1 inoculum by IM injection to 4 cotton rats in a separate study. All other animals were hamsters.
8. Recombinant wildtype HPIV3.

Six hamsters from each group and from a control group similarly infected with RSV were challenged on day 25 with $10^{6.0}$ pfu of biologically-derived HPIV3 wildtype virus given intranasally in a 0.1 ml inoculum. The lungs and nasal turbinates were harvested on day 4 and processed as described above. Virus present in the samples was titrated on 96 well plates of LLC-MK2 cell monolayers and incubated at 32°C for 7 days. Virus was detected by hemadsorption and the mean \log_{10} TCID₅₀/g was calculated for each group of hamsters. As shown in Table 5, those hamsters which had received the chimeric viruses, whether in the attenuated or wild type backbone, were highly protected against replication of challenge wild type HPIV3 in both the upper and the lower respiratory tract. Thus, despite the slight attenuating effect of the acquisition of the measles virus HA gene on replication of the *rcp45*(HA) chimeric viruses, infection with either *rcp45L*(HA P-M) or *rcp45L*(HA N-P) induced a high level of protection against HPIV3 as indicated by approximately a 1000-fold reduction of its replication in the upper and lower respiratory tract of hamsters. Since wild type measles virus does not replicate efficiently in hamsters, it cannot be used to challenge this host. However, it is expected that the attenuated chimeric *rcp45L*(HA) vaccine candidates will be highly efficacious against measles virus since high levels of neutralizing antibody, i.e., mean titer of greater than 1:5000, were induced. Comparable levels of measles virus antibodies are associated with strong resistance to measles virus disease in humans (Chen et al., *J. Infect. Dis.* 162:1036-42, 1990, incorporated herein by reference).

Table 5. Attenuated and wildtype HPIV3-measles HA chimeric viruses are highly protective against replication of challenge wildtype PIV3 in the upper and lower respiratory tracts of hamsters.

Animals Immunized with ¹	# Animals	Virus titer (\log_{10} TCID ₅₀ /g) [Tukey-Kramer Grouping ³]				Reduction in Titer (\log_{10})	
		Nasal Turbinates	Lungs	Nasal Turbinates	Lungs		
RSV	6	7.0±0.3[A]	5.7±0.4[A]	NA ²	NA		
rcp45L(HA P-M)	6	3.4±0.3[B]	2.9±0.0[B]	3.6	2.8		
rcp45L(HA N-P)	6	2.6±0.3[B]	3.4±0.2[B]	4.4	2.3		
rPIV3(HA P-M)	6	2.0±0.3[B]	3.2±0.1[B]	5.0	2.5		
rcp45L	6	1.9±0.2[B,C]	3.6±0.1[B]	5.1	2.1		
rJS	6	<1.4±0.0[C]	2.9±0.2[B]	>5.7	2.8		

1. All groups were challenged with 10^6 pfu biologically-derived JS wildtype PIV3 in a 0.1 ml inoculum given intranasally.
2. Not applicable.
3. Mean virus titers were assigned to statistically similar groups (A-C) by the Tukey-Kramer test. Therefore, means in each column with different letters are significantly different ($\alpha=0.05$) and means with the same letter are not significantly different.

EXAMPLE III**Construction of Antigenomic cDNAs Encoding a Chimeric HPIV3-1 Vector Bearing a HPIV2 HN Gene as an Extra Transcription/Translation Unit Inserted Between the F and HN Genes, and Recovery of Infectious Viruses**

5 rPIV3-1 is a recombinant chimeric HPIV3 in which the HN and F genes have been replaced by those of HPIV1 (see, e.g., Skiadopoulos et al., Vaccine 18:503-510, 1999; Tao et al., Vaccine 17:1100-1108, 1999; U.S. Patent Application Serial No. 09/083,793, filed May 22, 1998; U.S. Patent Application Serial No. 09/458,813, filed December 10, 1999; U.S. Patent Application Serial No. 09/459,062, filed December 10, 1999, each incorporated

10 herein by reference). In the present example, the HN gene of HPIV2 was inserted into the rPIV3-1 chimeric virus that served as a vector to produce a chimeric derivative virus, bearing an introduced heterologous antigenic determinant from HPIV2, able to protect

15 against both HPIV1 and HPIV2. The HPIV2 HN gene also was inserted into an attenuated derivative of rPIV3-1, designated rPIV3-1cp45, which contains 12 of the 15 cp45 mutations, i.e., those mutations on genes other than HN and F, inserted into the rPIV3 backbone

20 (Skiadopoulos et al., Vaccine 18:503-510, 1999). The source of the HPIV2 wild type virus was the wild type strain V9412-6 (designated PIV2/V94) (Tao et al., Vaccine 17:1100-1108, 1999), which was isolated in Vero cells from a nasal wash that was obtained in 1994 from a child with a natural HPIV2 infection. PIV2/V94 was plaque purified 3 times on Vero cells

25 before being amplified twice on Vero cells using OptiMEM tissue culture medium without FBS. A cDNA clone of the HN gene of PIV2/V94 was generated from virion RNA by reverse transcription (RT) using random hexamers and Superscript Preamplification System (Life Technologies) followed by PCR using Advantage cDNA Synthesis kit (Clontech, Palo Alto, CA) and synthetic primers which introduced *NcoI-HindIII* sites flanking the HN cDNA

30 (Figure 3A). The sequences of these primers were: (with HPIV specific sequences in upper case, restriction sites underlined, nts which are non-HPIV or which are altered from wt in lower case, and start and stop codons in bold), upstream HPIV2 HN 5'-
gggccATGGAAGATTACAGCAAT-3' (SEQ ID NO. 19); downstream HPIV2 HN 5'-
caataagcTTAAAGCATTAGTCCC-3' (SEQ ID NO. 20). The HN PCR fragment was

 digested with *NcoI-HindIII* and cloned into pLit.PIV31HNhc to generate pLit.32HNhc (Figure 3 B). The HPIV2 HN heterologous gene insert in pLit.32HNhc was completely sequenced using the ThermoSequenase Kit and ³³P-labeled terminators (Pharmacia

Amersham, Piscataway, NJ) and was confirmed to contain the authentic sequence of the PIV2/94 HN coding region.

The HPIV2 HN gene in pLit.32HNhc was further modified by PCR and Deep Vent thermostable DNA polymerase (New England Biolab, Beverly, MA) to introduce *PpuMI* sites for cloning into the unique *PpuMI* site in p38'ΔPIV31hc, Figure 3C (Skiadopoulos et al., *Vaccine* 18:503-510, 1999). The sequences of these primers were (with HPIV specific sequences in upper case, relevant restriction sites underlined, non-HPIV nt or nt altered from wt in lower case): upstream HPIV2 HN

5'-gcatggggcccGAGGAAGGACCCAAATAGACA-3' (SEQ ID NO. 21); downstream HPIV2 HN

5'-cccgggtcctgATTTC~~CCG~~GAGCACGCTTTG-3' (SEQ ID NO. 22). The modified cDNA bearing the HPIV2 HN ORF consists of (from left to right) a partial 5'-untranslated region (5'-UTR) of HPIV3 HN including the *PpuMI* site at the 5'-end, the HPIV2 HN ORF, the 3'-UTR of HPIV3 HN, a complete set of HPIV3 transcription signals (i.e. gene stop, intergenic region and gene start sequences) whose sequences match those at the HPIV3 HN and L gene junction, a partial 5'-UTR of HPIV3 L, and an added *PpuMI* site at its 3'-end (Figure 3C). This fragment was digested with *PpuMI* and inserted into p38'ΔPIV31hc digested with *PpuMI* to generate p38'ΔPIV31hc.2HN (Figure 3D). The inserted *PpuMI* cassette was sequenced in full and found to be as designed. The insert from p38'ΔPIV31hc.2HN was isolated as a 8.5 kb BspEI-SphI fragment and introduced into the BspEI-SphI window of pFLC.2G+.hc or pFLC*cp45* to generate pFLC.31hc.2HN or pFLC.31hc.*cp45*.2HN, respectively (Figure 3, E and F). pFLC.2G+.hc and pFLC*cp45* are full-length antigenomic clones encoding wt rPIV3-1 and rPIV3*cp45*, respectively, as described previously (Skiadopoulos et al., *J. Virol.* 73:1374-81, 1999; Tao et al., *J. Virol.* 72:2955-2961, 1998, each incorporated herein by reference).

Confluent HEp-2 cells were transfected with pFLC.31hc.2HN or pFLC.31hc.*cp45*.2HN plus the pTM(N), pTM(P no C), and pTM(L) support plasmids in the presence of MVA-T7 as previously described (Durbin et al., *Virology* 235:323-332, 1997, incorporated herein by reference). The recombinant chimeric viruses recovered from transfection were activated by addition of TPCK trypsin (Catalog No. 3741, Worthington Biochemical Corp., Freehold, NJ) as were all passages and titrations of viruses bearing the HPIV1 HN and F glycoproteins as described previously (Tao et al., *J. Virol.* 72:2955-2961,

1998, incorporated herein by reference). Recovered chimeric recombinant viruses rPIV3-1.2HN and rPIV3-1cp45.2HN were purified by plaque-to-plaque-to-plaque passage on LLC-MK2 monolayer in agarose overlay as previously described (Tao et al., *Vaccine* 17:1100-1108, 1999, incorporated herein by reference).

5 To determine if the rPIV3-1.2HN and rPIV3-1cp45.2HN recombinants contain the heterologous HPIV2 HN gene, viral RNA from each recovered recombinant chimeric virus was amplified on LLC-MK2 cells and concentrated by polyethylene glycol (PEG) precipitation (Mbiguino et al., *J. Virol. Methods* 31:161-170, 1991, incorporated herein by reference). Virion RNA (vRNA) was extracted with Trizol (Life Technologies) and used as template to synthesize first strand cDNA using Superscript Preamplification
10 system (Life Technologies, Gaithersburg, MD) and random hexamer primers as described above. The synthesized cDNA was amplified by PCR with the Advantage cDNA Synthesis kit (Clontech, Palo Alto, CA) with primers specific for HPIV1 F and HPIV1 HN coding region (for HPIV1 F 5'-AGTGGCTAATTGCATTCATCCACAT-3' (SEQ ID NO. 23) and
15 for HPIV1 HN 5'-GCCGTCTGCATGGTGAATAGCAAT-3') (SEQ ID NO. 24). The relative locations of the PIV1 F and HN primers are indicated by arrows in Figures 3 and 4. Amplified DNA fragments were digested and analyzed on agarose gels (Figure 4). Data for rPIV3-1cp45.2HN is not shown, but was comparable and confirmed in structure. rPIV3-1.2HN and rPIV3-1cp45.2HN each contained the insert of the expected size, and the
20 digestion patterns with a number of restriction enzymes confirmed the identity and authenticity of the inserts. The presence of the cp45 mutations in rPIV3-1cp45.2HN was also confirmed.

To confirm the expression of HPIV2 HN by the rPIV3-1.2HN chimeric virus, LLC-MK2 monolayers in T25 flasks were infected with PIV2/V94, rPIV3-1, or rPIV3-
25 1.2HN at a MOI of 5 in 5 ml of serum-free OptiMEM containing 0.5 µg/ml TPCK trypsin. After incubation for 18 hours at 32°C, the flasks were washed three times with 5 ml of methionine and cysteine deficient DMEM (BioWhittaker, Walkersville, MD). Cells were then fed with 1 ml of methionine and cysteine deficient DMEM supplemented with 120 µCi of ProMix 35S-methionine and 35S-cysteine mixture (Pharmacia Amersham, Piscataway,
30 NJ) and incubated for 18 hours at 32°C. Cells were scraped into medium, pelleted by brief centrifugation in a microfuge, and washed three times with cold PBS. Each cell pellet was resuspended in 1 ml RIPA buffer (1% sodium deoxycholate, 1% Triton X-100, 0.2% SDS, 150 mM NaCl, and 50 mM Tris-HCl, pH7.4) containing 250 units/ml of Benzonase (Sigma),

freeze/thawed once, and clarified by centrifugation at 12,000 X g for 5 min in a microfuge. Clarified supernatants were transferred to a clean microfuge tube, mixed with 50 μ l of anti-HPIV2 HN monoclonal antibody (mAb) 150S1 (Tsurudome et al., *Virology* 171:38-48, 1989, incorporated herein by reference), and incubated with mixing at 4°C for 3 hours. The monoclonal antibody was precipitated by the addition to each tube of 0.2 ml of 10% Protein A sepharose suspension (in RIPA buffer) and incubation with mixing at 4° for 18 hours. The beads were washed three times with RIPA buffer and pelleted by brief centrifugation in a microfuge. Each sample was suspended in 90 μ l of 1X loading buffer, and 10 μ l was resolved on a 4-12% SDS polyacrylamide gel (PAGE; NOVEX, San Diego, CA). The gel was dried and autoradiographed (Figure 5). The mAb, specific to PIV2 HN, precipitated a protein from both rPIV3-1.2HN and PIV2/V94 infected LLC-MK2 cells, but not from rPIV3-1-infected cells, with a size expected for the 86kD Kd HN protein of HPIV2 (Rydbeck et al., *J. Gen. Virol.* 69:931-5, 1988, incorporated herein by reference).

15 EXAMPLE IV

The rPIV3-1 Viruses Carrying an HPIV2 Antigenic Determinant Exhibit Temperature Sensitive Phenotypes Similar to Those of Their Parental Vector Viruses

The level of temperature sensitivity of replication of rPIV3-1.2HN and rPIV3-1.cp45.2HN in LLC-MK2 cells was evaluated to determine if the acquisition of the HN ORF of HPIV2 by rPIV3-1 wild type or attenuated viruses employed as vectors altered the level of temperature sensitivity of replication in the resultant chimeric derivatives bearing the heterologous antigenic determinant of HPIV2 compared to the parental, vector viruses (Table 6). rPIV3-1.2HN and rPIV3-1.cp45.2HN, along with control viruses, were serially diluted 1:10 in 1X L15 supplemented with 0.5 μ g/ml TPCK trypsin and used to infect LLC-MK2 monolayers in 96 well plates in quadruplicate. Infected plates were placed at various temperatures for 7 days before the virus titers were determined by hemadsorption using 0.2% guinea pig erythrocytes (in 1X PBS). The virus titers are presented as \log_{10} TCID₅₀ \pm standard error (S.E.). As shown in Table 6, rPIV3-1.2HN and rPIV3-1.cp45.2HN exhibited a level of temperature sensitivity similar to that of their parental, vector viruses, i.e. rPIV3-1 and rPIV3-1.cp45, respectively, each of which lacks the HPIV2 HN insert. This indicated that the introduction of one extra transcription/translation unit in rPIV3-1.2HN and rPIV3-

1cp45.2HN, does not significantly alter their level of temperature sensitivity of replication *in vitro*.

Table 6. The rPIV3-1 viruses carrying the PIV2 HN insertion have a temperature sensitive phenotype similar to that of their parental virus.

Virus	Titer at 32° C ^a (log ₁₀ TCID ₅₀)	Titer reduction (log ₁₀ TCID ₅₀) at various temperatures (°C) ^a					
		35° ^b	36°	37°	38°	39°	40°
PIV2/V9412	7.8	0.3	(0.1) ^c	0.0	(0.4)	(0.4)	0.0
PIV1/Wash64	8.5	1.5	1.1	1.4	0.6	0.5	0.9
rPIV3/JS	7.9	0.3	0.1	0.1	(0.3)	(0.4)	0.4
PIV3 <i>cp45</i>	7.8	0.5	0.3	1.3	<u>3.4</u> ^d	6.8	6.9
rPIV3-1	8.0	0.8	0.5	0.6	0.9	1.1	<u>2.6</u>
rPIV3-1.2HN	8.3	0.5	(0.3)	0.3	0.6	1.5	<u>2.6</u>
rPIV3-1 <i>cp45</i>	8.0	0.5	0.4	<u>3.4</u>	4.8	6.6	7.5
rPIV3-1 <i>cp45.2HN</i>	8.0	0.3	1.4	<u>2.9</u>	5.3	7.6	7.6

^a Data presented are means of two experiments.

^b Data at 35°C were from single experiment.

^c Numbers in parentheses represent titer increase.

^d Underlined value indicates shut-off temperature at which the virus titer showed a reduction of 100-fold or more in comparison to the titer at 32°C.

EXAMPLE VReplication and Immunogenicity of rHPIV3-1.2HN Chimeric Viruses in Animals

To determine the level of replication of the chimeric viruses *in vivo*, Golden Syrian hamsters in groups of six were inoculated intranasally with 0.1 ml of 1X L-15 medium containing $10^{5.3}$ TCID₅₀ (or 10^6 pfu) of virus (Table 7). Four days after infection, hamsters were sacrificed and their lungs and nasal turbinates harvested. Virus titers, expressed as mean log₁₀TCID₅₀/gram of tissue (Table 7), were determined. rPIV3-1 expressing the PIV2 HN gene, termed rPIV2-1.2HN, is more restricted in replication than its rPIV3-1 parent as indicated by a 30-fold reduction in virus titer in both the upper and lower respiratory tracts of hamsters. Thus, the insertion of a transcription/translation unit expressing the PIV2 HN protein into rPIV3-1 attenuates the virus for hamsters. The attenuating effect of insertion of a transcription/translation unit containing PIV2 HN ORF into rPIV3-1 was slightly more than that observed for the insertion of a similar unit containing the measles HA ORF into the recombinant JS strain of wild type PIV3. The rPIV3-1cp45.2HN virus was 1,000-fold more restricted in replication than the rPIV3-1cp45 parent indicating that the attenuating effect of the PIV2 HN insertion and the cp45 mutations are additive. It should be possible to adjust the level of attenuation as needed by adding fewer cp45 mutations than the 12 that are present in rPIV3-1.cp45.2HN.

Table 7. The chimeric rPIV3-1 expressing the HN glycoprotein of PIV2 (rPIV3-1.2HN) is attenuated in the respiratory tract of hamsters

Experiment No.	Virus	Virus titer in indicated tissue log ₁₀ TCID ₅₀ /g±S.E.) ^c	
		NT	Lungs
1 ^a	rPIV3-1	6.9 ± 0.1[A] ^d	6.0 ± 0.3[A]
	rPIV3-1.2HN	5.4 ± 0.2[B]	4.4 ± 0.4[C]
2 ^b	rPIV3-1	6.7 ± 0.1[A]	6.6 ± 0.2[A]
	rPIV3-1.2HN	5.1 ± 0.1[B, C]	5.2 ± 0.2[B]
	rPIV3-1cp45	4.6 ± 0.3[C]	1.8 ± 0.4[D]
	rPIV3-1cp45.2HN	1.5 ± 0.1[D]	≤1.2[D]
	rPIV3/JS	6.5 ± 0.2[A]	6.7 ± 0.1[A]
	rcp45	4.9 ± 0.2[B, C]	1.2 ± 0.04[D]

^a Groups of six animals were inoculated intranasally with 10⁶ pfu of indicated virus in 0.1 ml medium on day 0.

^b Groups of 6 hamsters were inoculated intranasally as in Experiment 1 with 10^{5.3} TCID₅₀ of indicated virus on day 0.

^c Lungs and nasal turbinates of the hamsters were harvested on day 4. Virus titers in tissue were determined and the titer expressed as log₁₀TCID₅₀/gram ± standard error (S.E.). NT = nasal turbinates.

^d Means in each column with a different letter are significantly different (α=0.05) by Duncan's Multiple Range test whereas those with the same letter are not significantly different.

Since the single rPIV3-1.2HN virus expresses protective antigens of PIV1 (the F and HN glycoprotein) and PIV2 (the HN glycoprotein only), infection with this virus will induce resistance against challenge with either PIV1 or PIV2 wild type viruses. To verify this, Golden Syrian hamsters in groups of 12 were immunized intranasally with $10^{5.3}$ TCID₅₀ of virus as described above. Half of the hamsters were challenged with PIV2 on day 29, the remaining half with PIV1 on day 32. Hamster lung and nasal turbinate tissues were harvested 4 days after challenge, and titer of challenge virus were determined as described above (Table 8). Sera were obtained before and 28 days after immunization and tested for their neutralizing antibody titer against PIV1 and PIV2.

Table 8. The chimeric rPIV3-1 virus expressing the HN glycoprotein of PIV2 (rPIV3-1.2HN) protects hamsters against challenge with both PIV1 and PIV2

Immunizing virus ^a	Serum neutralizing antibody titer against indicated virus (reciprocal mean log ₂ ±SE) ^b				Titer of challenge virus in indicated tissues (log ₁₀ TCID ₅₀ /g±SE) ^c							
	PIV1		PIV2		PIV1		PIV2		Lung		Lung	
	pre	post	pre	post	NT	Lung	NT	Lung	NT	Lung	NT	Lung
rPIV3/IS	54.0±0.0	54.0±0.0	4.5±0.1	4.6±0.2	5.4±0.2	5.1±0.1	5.4±0.2	5.1±0.1	6.8±0.2	6.0±0.3	6.8±0.2	6.0±0.3
PIV2	54.0±0.0	54.0±0.0	4.3±0.2	9.6±0.2	5.7±0.2	5.7±0.2	5.7±0.2	5.7±0.2	5.1.2	5.1.2	5.1.2	5.1.2
rPIV3-1	4.2±0.1	8.5±0.3	4.0±0.0	4.2±0.1	5.1.2	5.1.2	5.1.2	5.1.2	6.3±0.1	6.5±0.2	6.3±0.1	6.5±0.2
rPIV3-1.2HN	54.0±0.0	6.2±0.2	4.1±0.1	8.3±0.2	2.3±0.5	5.1.2	2.3±0.5	5.1.2	5.1.2	5.1.2	5.1.2	5.1.2
rPIV3-1cp45	54.0±0.0	6.2±0.4	54.0±0.0	4.0±0.0	3.6±0.3	2.7±0.5	3.6±0.3	2.7±0.5	6.0±0.1	5.7±0.4	6.0±0.1	5.7±0.4
rPIV3-1cp45.2HN	4.0±0.9	4.1±0.1	4.0±0.0	4.2±0.1	5.1±0.2	4.8±0.2	5.1±0.2	4.8±0.2	6.8±0.1	6.6±0.2	6.8±0.1	6.6±0.2

^a Hamsters in groups of 12 were immunized with 10^{5.7} TCID₅₀ of indicated virus intranasally on day 0.

^b Serum was diluted 1:10 with OptiMEM and heat-inactivated by incubation at 56° for 30 min. The serum neutralizing antibody titer was determined on LLC-MK2, and the titers are expressed as reciprocal mean log₂-standard error (SE).

^c Half of the hamsters from each immunized group were challenged with 10⁶ TCID₅₀ PIV2 on day 29, and the remaining half were challenged with 10⁶ TCID₅₀ PIV1 on day 32. Tissue samples were harvested 4 days after challenge, and challenge virus titers are expressed as log₁₀TCID₅₀/gram of tissue ± SE. NT = nasal turbinates.

As expected PIV3 provided no resistance against either PIV1 or PIV2 (Tao, Vaccine 17:1100-1108, 1999), while previous infection with PIV2 wild type virus and rPIV3-1 induced complete resistance to replication of PIV2 and PIV1 challenge viruses, respectively. In contrast to these viruses that provided protection against only one virus, rPIV3-1.2HN induced antibody to both PIV1 and PIV2 and included strong resistance to both PIV1 and PIV2 as indicated by the 1,000- to 10,000-fold reduction in replication of each virus in the upper and lower respiratory tract of rPIV3-1.2HN immunized hamsters. This indicated that a single recombinant chimeric PIV can induce resistance against two human viral pathogens. However, the derivative of rPIV3-1.2HN carrying the *cp45* mutations failed to induce significant resistance to replication of wild type PIV1 or PIV2 challenge virus indicating that this particular recombinant chimeric virus is over-attenuated in hamsters. Introduction of one or several selected *cp45* mutations, rather than the complete set of 12 mutations, into rPIV3-1.2HN can be done to adjust the level of attenuation of rPIV3-1.2HN to an appropriate level.

15

EXAMPLE VI

Construction of cDNAs Encoding rHPIV3 Viruses Containing Nucleotide Insertions

As discussed above, insertion of the measles HA ORF between either the N/P or P/M gene junction of the attenuated vector virus, rPIV3 $_{cp45L}$, as well as at the N/P, P/M, and HN/L junctions of wild type PIV3, further restricted its replication in the upper respiratory tract of hamsters, indicating that insertion of an additional gene at either location within the HPIV3 genome can augment attenuation of candidate vaccine viruses. In these exemplary aspects of the invention, the gene insert was relatively large (approximately 1900 nts). Further examples are provided herein that indicate the size of the insert specifies a selectable level of attenuation of the resulting recombinant virus. This was evaluated by introducing sequences of various lengths which were derived from a heterologous virus, exemplified by the RSV A2 strain, as single gene units (GUs) between the HPIV3 HN and L ORFs. The inserts were designed specifically to lack any significant ORF, whereby any effects observed would not be complicated by possible contribution of expressed protein. In order to distinguish between effects due to increased genome length versus expression of an additional mRNA, a second series of constructs was made in which inserts of similar sizes

20
25
30

were introduced into the downstream noncoding region (NCR) of the HN gene. Thus, two series of rPIV3s were made containing insertions of increasing length: in the GU series, the insert was added as an extra gene encoding an extra mRNA, while in the NCR series the insert was made so that the gene number was unchanged.

5 **Construction of cDNAs encoding rHPIV3 viruses containing GU and 3'-NCR insertions**

Insertion mutations were constructed in a pUC based plasmid, pUC118-Stu, containing the *XhoI* to *SphI* fragment (HPIV3 nts 7437-11317) of the full length HPIV3 clone p3/7(131)2G-Stu. Two separate plasmids were constructed as acceptor plasmids for insertion of GUs and HN gene 3'-NCR extensions (Figure 6). In each, a synthetic oligonucleotide duplex containing multiple cloning sites was inserted into the unique *Stu I* site. The inserted sequence for the GU insertion plasmid contained a HN gene-end (GE) signal sequence, the conserved intergenic (IG) trinucleotide sequence, and a L gene-start (GS) signal sequence, cis-acting sequences that direct termination of the HN gene transcription and initiation of transcription of the inserted sequence, respectively (Figure 6). Additional unique restriction endonuclease sites were included in the multiple cloning region to facilitate subsequent screening and subcloning. The 3'-NCR extension acceptor plasmid was similarly designed and constructed, but it lacked the cis-acting GE, IG, and GS sequences at its 5'-end (Figure 6B, Table 9). The RSV antigenomic plasmid d53RSV sites or subgenomic plasmid pUC118FM2 (Table 9) were digested with the appropriate restriction enzymes, and fragments of the desired sizes were isolated by electrophoresis on agarose gels and ligated individually into the unique *HpaI* site of the GU or the HN gene 3'-NCR extension acceptor plasmid (Figure 6; Table 9). Clones were screened to identify ones in which the RSV restriction fragments were inserted in the reverse orientation, an orientation in which all reading frames contained multiple stop codons (Figure 7). Short synthetic oligonucleotide duplexes ranging in size from 13 to 17 nucleotides also were inserted as necessary into the GU or 3'-NCR acceptor plasmids to modify the genome length to conform to the "rule of six" (Table 9). The specific RSV sequences and size of the short synthetic oligonucleotides added are summarized in Table 9. Plasmid clones were sequenced through all restriction enzyme sites used for subcloning, and *XhoI-SphI* fragments containing insertion mutations conforming to the rule of six, either as GUs or HN gene NCR extensions, were cloned into the full-length PIV3 cDNA plasmid p3/7(131)2G+. One insert, containing the 1908 GU insert, also was placed into an antigenomic cDNA bearing the three L mutations of *cp45*.

Table 9: Sources of nucleotides used to create the gene unit (GU) and HN gene 3' non coding region (NCR) extension insertions.

Restriction fragment size (nts)	Restriction sites and nt position in the RSV antigenome	GU multiple cloning site (58 nt) + rule of 6 oligonucleotide ^a		GU insertion (total nts inserted)	NCR multiple cloning site (32 nt) + rule 6 oligonucleotide ^a	NCR insertion (total nts inserted)
		+58 + 13	nd			
97 ^a	<i>SspI-SspI</i> ; 7272-7369	+58 + 13	nd	168	nd	nd
212 ^b	<i>HpaI-HpaI</i> ; 12243-12455	nd	nd	nd	+32 + 14	258
603 ^b	<i>SspI-SspI</i> ; 309-912	+58 + 17	nd	678	nd	nd
925 ^b	<i>HpaI-HpaI</i> ; 12455-13380	+58 + 13	nd	996	+32 + 15	972
1356 ^{b,c}	<i>HincII-HincII</i> ; 5060-6417	+58 + 14	nd	1428	+32 + 16	1404
1850 ^{b,d}	<i>HpaI-HpaI</i> ; 12455-13380	+58 + 0	nd	1908	nd	nd
3079 ^b	<i>EcoRV-EcoRV</i> ; 1403-4482	nd	nd	nd	+32 + 15	3126
3845 ^b	<i>ScaI-ScaI</i> ; 344-4189	+58 + 15	nd	3918	+32 + 17	3894

122

a. Source of RSV sequence is pUC118FPMZ, a plasmid containing a subgenomic cDNA fragment of RSV subgroup A as described previously (Juhász, K. et al, J Virol., 71:5814-5819, 1997.). b. Source of RSV sequence is D53sites, a plasmid containing the entire RSV subgroup A cDNA sequence with several introduced point mutations as described previously. The previously described D53 sites plasmid was used to derive the rASites virus described in Whitehead, S. et al. J. Virol., 72:4467-4471, 1998. c. The gel purified 1356 nt fragment contained a 1 nt deletion compared to the predicted 1357 nt restriction endonuclease cleavage product. d. The 1850 nt fragment is a product of two 3' to 3' adjoined 925 nt restriction fragments. e. The following oligonucleotides were inserted into the *MvaI* restriction site to conform all the inserted foreign sequences to the rule of six: 13mer: CGCGGCGAGGCCTG (SEQ ID NO. 25); 14mer: CGCGGCGAGGCCTG (SEQ ID NO. 26); 15mer: CGCGGCGAGGCCTCGCGG (SEQ ID NO. 27); 16mer: CGCGGCGAGGCCT (SEQ ID NO. 28); 17mer: CGCGGCGAGGCCT (SEQ ID NO. 29). nd, not done.

Recovery of recombinant PIV3s bearing insertion mutations

Full-length antigenomic cDNA derivatives bearing the insertion mutations and three support plasmids pTM(N), pTM(P no C) and pTM(L) (Durbin et al., Virology 235:323-332, 1997; Durbin et al., Virology 261:319-330, 1999, each incorporated herein by reference) were transfected into HEp-2 monolayers in 6-well plates (Costar, MA) using LipofectACE (Life Technologies, MD), and the monolayers were infected with MVA-T7 as described previously (Durbin et al., Virology 235:323-332, 1997; Skiadopoulos et al., J. Virol. 72:1762-8, 1998, each incorporated herein by reference). After incubation at 32°C for 4 days, the transfection harvest was passaged onto LLC-MK2 cells in T-25 flasks which were incubated at 32°C for four to eight days. The clarified medium supernatant was subjected to plaque purification on LLC-MK2 cells as described previously (Durbin et al., Virology 235:323-332, 1997; Hall et al., Virus Res. 22:173-184, 1992; Skiadopoulos et al., J. Virol. 72:1762-8, 1998, each incorporated herein by reference). Each biologically-cloned recombinant virus was amplified twice in LLC-MK2 cells at 32°C to produce virus for further characterization. Virus was concentrated from clarified medium by polyethylene glycol precipitation (Mbiguino et al., J. Virol. Methods 31:161-170, 1991, incorporated herein by reference), and viral RNA (vRNA) was extracted with Trizol Reagent (Life Technologies). Reverse transcription was performed on vRNA using the Superscript II Preamplification System (Life Technologies) with random hexamer primers. The Advantage cDNA PCR kit (Clontech, CA) and sense (PIV3 nt 7108-7137) and antisense primers (PIV3 nt 10605-10576) were used to amplify fragments for restriction endonuclease digestion or sequence analysis. The PCR fragments were analyzed by agarose gel electrophoresis (Figure 8) and sequencing. Each of the recovered rPIV3 insertion mutants contained insertions of the indicated sizes and they were next evaluated for their biological properties.

EXAMPLE VIIReplication of rHPIV3 Viruses Containing GU or NCR Inserts
in Animals and in Tissue Culture**Multi-step growth curves**

- 5 The growth properties of the rPIV3 GU and NCR insertion mutants were compared to rPIV3 wt and *tcp45_L* *in vitro*. As shown in Figure 9, the rate of replication and the peak virus titer of each of the rPIV3s containing either the GU or NCR insertions was indistinguishable from that of rPIV3 wt indicating that insertion of sequences of at least 3918 nts in length does not affect virus replication *in vitro*.

10 Replication in hamsters of rPIVs containing GU insertions

- Hamsters were inoculated intranasally with 10^{6.0} TCID₅₀ rPIV3wt, *tcp45_L* or with one of the indicated mutant rPIV3s bearing GU insertions (Table 10). Lungs and nasal turbinates were harvested on day four after infection and the level of replication of each virus was determined. Insertion of GUs ranging in size from 168 nt up to 1908 nt did not
- 15 significantly reduce viral replication in the respiratory tract of hamsters. However, insertion of a 3918 nt gene unit between the HN and L ORF of wild type PIV3 resulted in a 5 and 25-fold reduction in viral replication in the nasal turbinates and lungs, respectively. This indicates that gene unit insertions of this size are attenuating for a wild type virus whereas shorter sizes, e.g., below approximately 2000 nt, have little effect on replication of wild type
- 20 virus in the respiratory tract of hamsters. Thus, GU length can be altered to determine a desired level of attenuation in PIV vaccine viruses.

Table 10: Replication of rPIV3 GU insertion mutants in the respiratory tract of hamsters

Virus ^a	Mean virus titer (\log_{10} TCID ₅₀ /g \pm S.E. ^b) in:	
	Nasal Turbinates	Lungs
rPIV3 wt	5.9 \pm 0.2	6.0 \pm 0.2
r168 nt GU ins	5.9 \pm 0.1	6.4 \pm 0.1
r678 nt GU ins	6.1 \pm 0.1	6.2 \pm 0.1
r996 nt GU ins	5.5 \pm 0.2	5.4 \pm 0.2
r1428 nt GU ins	5.9 \pm 0.1	5.3 \pm 0.6
r1908 nt GU ins	5.6 \pm 0.1	5.7 \pm 0.2
r3918 nt GU ins	5.2 \pm 0.2	4.6 \pm 0.3
rcp45 _L	3.1 \pm 0.0	1.7 \pm 0.2
r1908 nt GU ins/cp45 _L	1.8 \pm 0.2	1.5 \pm 0

a. Hamsters, in groups of eight, were administered $10^{6.0}$ TCID₅₀ of virus intranasally in a 0.1 ml inoculum. Lungs and nasal turbinates were harvested four days later and virus titer was determined at 32°C. b. S.E.: Standard error.

As described above, the insertion of the HA gene of measles virus into the rJS wildtype and the attenuated *cp45L* virus further attenuated each virus for hamsters. Since the HA gene of measles virus is 1936 nt in length, we examined the effect of a similar size gene insertion (1908 nt) on replication of *rcp45L*. The 1908 gene insertion differs from the measles virus HA gene insertion in that it cannot synthesize a large polypeptide. When the 1908 nt GU insertion was combined with the *cp45 L* polymerase amino acid substitutions (r1908 nt GU ins/*cp45_L* in Table 10), attenuation was augmented approximately 20-fold in the upper respiratory tract. Considered together, these findings indicate that GU insertions of approximately 3918 nts in length can attenuate a wild type PIV3 virus for hamsters and that GU insertions of about half this size can further attenuate an attenuated PIV3 vaccine candidate. Thus, GU insertions can have dual roles in the design of recombinant vaccines. The first role is to encode a protective antigen of a pathogen, and the second role is to confer an attenuation phenotype.

Replication in hamsters of rPIV3s containing HN gene 3'-NCR insertions.

Hamsters were inoculated intranasally with rPIV3 control viruses or viruses bearing insertion mutations extending the length of the HN gene 3'-NCR (Table 11). Lungs and nasal turbinates were harvested four days after inoculation and the level of viral replication in each tissue was determined as described above. HN gene NCR insertions ranging in size from 258 nt up to 1404 nt did not significantly reduce viral replication in the respiratory tract of hamsters (Table 3). However, an insertion of 3126 nt effected a 16-fold reduction in viral titer in the upper and lower respiratory tracts of infected hamsters, and a 3894 nt HN gene NCR insertion resulted in a 12-fold reduction of viral replication in the upper and lower respiratory tracts, suggesting that increasing the genome length also confers an attenuating effect on viral replication.

Table 11: Replication of rPIV3 NCR insertion mutants in the respiratory tract of hamsters

Virus ^a	Mean virus titer (log ₁₀ TCID ₅₀ /g ± S.E. ^b) in:	
	Nasal Turbinates	Lungs
rPIV3 wt	6.2±0.1	6.4±0.1
r258 nt NCR ins	5.9±0.1	6.5±0.1
r972 nt NCR ins	5.9±0.1	6.6±0.1
r1404 nt NCR ins	5.9±0.2	6.6±0.1
r3126 nt NCR ins	5.0±0.1	5.2±0.1
r3894 nt NCR ins	5.1±0.1	5.3±0.1
rcp45 _L	3.4±0.1	1.9±0.2

a. Hamsters, in groups of eight, were administered 10^{6.0} TCID₅₀ of virus intranasally in a 0.1 ml inoculum. Lungs and nasal turbinates were harvested four days later and virus titer was determined at 32°C. b. S.E.: Standard error.

Evaluation of the level of temperature sensitivity of GU and NCR insertions

The efficiency of plaquing (EOP) at permissive and non-permissive temperatures of rPIVs was determined on LLC-MK2 monolayers as described above (Table 12). At 32°C, viruses bearing GU insertions ranging in size from 168 nt up to 3918 nt and NCR insertions ranging in size from 258 nt up to 3894 nt had a plaque morphology that was similar to that of rPIV3 wt. However, at 39°C and at higher temperatures all of the viruses bearing insertion mutations had a small plaque phenotype (Table 12). The GU insertions ranging in size from 996 nt up to 3918 nt yielded viruses that were not ts at 40°C. However, viruses bearing HN gene NCR insertions of 1404 nts or greater yielded viruses that were slightly ts at 40°C with a gradient of temperature sensitivity proportional to the size of the insertion. Addition of the 1908 nt GU insertion to the *cp45_L* backbone yielded a virus that was almost 100-fold more ts at 38°C compared to *rcp45_L*, demonstrating that the ts phenotype specified by the 1908 nt GU insertion and by the L gene *ts* mutations is additive.

Table 12: Efficiency of plaque formation of rPIV3 GU and NCR insertion mutants at permissive and non-permissive temperatures

Virus	Virus titer at indicated temperature (log ₁₀ PFU/ml)				
	32°C	37°C	38°C	39°C	40°C
rPIV3 wt	7.8	ND	ND	7.4	7.5
r168 nt GU ins	7.8	ND	ND	7.5 ^a	6.7 ^a
r678 nt GU ins	7.9	ND	ND	7.3 ^a	7.0 ^a
r996 nt GU ins	7.7	ND	ND	7.0 ^a	6.3 ^a
r1428 nt GU ins	7.8	ND	ND	7.4 ^a	6.4 ^a
r1908 nt GU ins	7.6	ND	ND	6.5 ^a	6.0 ^a
r3918 nt GU ins	6.3	ND	ND	5.7 ^a	5.0 ^a
r258 nt NCR ins	8.1	ND	ND	7.4 ^a	7.5 ^a
r972 nt NCR ins	8.2	ND	ND	7.8 ^a	7.8 ^a
r1404 nt NCR ins	6.7	ND	ND	5.2 ^a	<3.7
r3126 nt NCR ins	7.4	ND	ND	6.4 ^a	4.5 ^a
r3894 nt NCR ins	7.4	ND	ND	<u>5.3^a</u>	5.0 ^a
tcp45L	7.8	7.3	6.0	<u><0.7</u>	ND
r1908 nt GU ins/cp45L	6.7	5.0 ^a	<u>3.0^a</u>	<0.7	ND
tcp45	8.1	6.7	<u>5.7^a</u>	2.0 ^a	ND

a. Plaques were enumerated by immunoperoxidase staining after incubation for 6 days at the indicated temperature. Values which are underlined and in bold type represent the lowest restrictive temperature at which there was at least a 100-fold reduction of plaque efficiency compared to the titer at 32°C, which is defined as the shut-off temperature of plaque formation.

Since the r3918 nt GU insertion mutant as well as the r3126 nt and r3894 nt NCR insertion mutants replicated efficiently *in vitro* but were restricted in replication in the respiratory tract of hamsters, these recombinants exhibit a novel, host-range attenuation phenotype.

5 Based on the foregoing examples, it is demonstrated that recombinant HPIV3 (rHPIV3) provides an effective vector for foreign viral protective antigens expressed as additional, supernumerary genes, as exemplified by the measles virus hemagglutinin (HA) glycoprotein gene. In another embodiment, the rHPIV3-1 antigenic chimeric virus, a recombinant HPIV3 in which the PIV3 F and HN genes were replaced by their HPIV1
10 counterparts, provides an effective vector the HPIV2 hemagglutinin-neuraminidase (HN) glycoprotein. In each case, the foreign coding sequence was designed and constructed to be under the control of a set of HPIV3 gene start and gene end transcription signals, inserted into the vector genome as an additional, supernumerary gene, and expressed as a separate mRNA by the HPIV3 polymerase.

15 Expression of the measles virus HA or the HPIV2 HN glycoprotein from a supernumerary gene insert by the rHPIV3 or rHPIV3-1 vector was determined to be stable over multiple rounds of replication. Hamsters infected with the rHPIV3 vector expressing the measles virus HA or the rHPIV3-1 vector expressing the HPIV2 HN glycoprotein
20 induced a protective immune response to HPIV3 and measles virus, or to HPIV1 and HPIV2, respectively. Thus, a single rHPIV3 vector expressing the protective antigen of measles virus induced a protective immune response against two human pathogens, namely, HPIV3 via an immune response to the glycoproteins present in the vector backbone and measles virus via the HA protective antigen expressed from the extra gene inserted into rHPIV3. The measles virus glycoprotein was not incorporated into the infectious HPIV3 vector virus, and
25 hence its expression would not be expected to alter the tropism of the vector nor render it susceptible to neutralization with measles virus-specific antibodies. Similarly, a single rHPIV3-1 vector expressing the protective HN antigen of HPIV2 induced a protective immune response against two human pathogens, namely, HPIV1 via an immune response to the glycoproteins present in the vector backbone and HPIV2 via the HN protective antigen
30 expressed from the extra gene inserted into rHPIV3-1.

EXAMPLE VIIIA Single rHPIV3 Expressing Up To Three Supernumerary Foreign Viral Glycoproteins5 Induces Protective Antibodies Against Up To Three Viruses

Modification of a single recombinant vaccine virus to induce immunity against multiple pathogens has several advantages. It is much more feasible and expeditious to develop a single attenuated backbone expressing antigens against multiple pathogens than it is to develop a separate attenuated vaccine against each pathogen. Each pathogen offers
10 different challenges for manipulation, attenuation and demonstration of safety and efficacy, and it would be a daunting task to attempt to develop an attenuated version of each of a series of pathogens. It is also simpler and easier to prepare, handle, and administer a single vaccine virus than to undertake these activities with several different attenuated viruses. Reducing the number of vaccine viruses also will help simplify the crowded schedule of
15 pediatric immunizations. Several attenuated viruses can be administered as a mixture, but this complicates vaccine development, since each component must be shown to be safe separately, and then shown to be safe and efficacious as a mixture. One particular problem with the administration of mixtures of viruses is the common phenomenon of viral interference, in which one or more of the viruses in the mixture interferes with the
20 replication of one or more of the other components. This may result in reduced replication and immunogenicity for one or more components. This common problem is obviated by the use of a single vector backbone. Also, since some viruses such as measles virus have particular safety concerns, it would be safer to use a single, comparatively benign virus such as PIV as a vector bearing multiple supernumerary antigens, as opposed to a mixture of
25 separately-attenuated viruses, each of which must be developed and validated separately.

In the present example recombinant HPIVs are constructed and shown to serve as vectors for more than one supernumerary gene with satisfactory characteristics of replication and immunogenicity for development of vaccine viruses. In particular, this example describes the design, construction, recovery, and characterization of rHPIV3s
30 expressing one, two or three supernumerary genes from the following list: (i) the hemagglutinin-neuraminidase (HN) of HPIV1 (Washington/20993/1964 strain); (ii) the HN of HPIV2 (V9412 strain); (iii) the hemagglutinin (HA) of the wild type Edmonston strain of measles virus; and (iv) a 3918-nt translationally-silent synthetic gene called gene unit (GU)

(see above). The added genes were inserted into rHPIV3 between the nucleoprotein (N) and phosphoprotein (P) genes, between the P and membrane protein (M) genes, or between the HN and large polymerase (L) genes. Thus, the disclosure demonstrates the successful use of an HPIV3 vector modified into a bivalent, trivalent, or quadrivalent vaccine recombinant capable of inducing multivalent immunity, e.g., against the vector itself and one or two additional pathogens.

Insertion of the HPIV1 HN and HPIV2 HN genes between the N/P and P/M genes was performed as follows: Plasmid pUC119(*Afl*III N-P), a subclone of the HPIV3 antigenomic cDNA (Durbin, *J. Virol.* 74:6821-31, 2000, incorporated herein by reference), was modified by site directed mutagenesis to insert a unique *Afl*III site into (i) the downstream noncoding region of the HPIV3 N gene (CTAAAT to CTTAAG, HPIV3 nts 1677-1682), or (ii) the downstream noncoding region of the HPIV3 P gene (TCAATC to CTTAAG, HPIV3 nts 3693-3698). Each *Afl*III site was then modified by the insertion of an oligonucleotide duplex, creating the intermediate plasmids pUC(GE/GS-N-H)_{N,P} and pUC(GE/GS-N-H)_{P,M}, respectively. The inserted duplex contained an HPIV3 gene-end (GE) sequence, the conserved intergenic (IG) trinucleotide sequence, and an HPIV3 gene-start (GS) sequence, which are cis-acting signals that direct transcriptional termination and initiation, respectively (Figure 10). Additional unique restriction endonuclease sites were included in the multiple cloning region to facilitate subsequent subcloning and screening, including *Nco*I and *Hind*III sites for addition of the HPIV1 and HPIV2 HN ORFs. Thus, a foreign ORF inserted into the multiple cloning site would be under the control of a set of HPIV3 transcription signals and expressed as a separate mRNA by the HPIV3 polymerase. The multiple cloning site also contained an *Mlu*I site for inserting oligonucleotides of varying lengths as necessary to make the entire inserted sequence conform to the rule of six (Calain et al., *J. Virol.* 67:4822-30, 1993; Durbin et al., *Virology* 234:74-83, 1997b; 1999a Skiadopoulos et al., *Virology* 272:225-34, 2000).

The HPIV1 HN ORF, available as an *Nco*I to *Hind*III restriction fragment of p38^Δ31hc #6 (Tao et al., *J. Virol.* 72:2955-2961, 1998), was inserted into the *Nco*I to *Hind*III sites of pUC(GE/GS-N-H)_{N,P} and pUC(GE/GS-N-H)_{P,M} to generate pUC 1HN_{N,P} and pUC 1HN_{P,M}, respectively. Short oligonucleotide duplexes were inserted in the unique *Mlu*I restriction site to adjust the sequence to conform to the rule of six. These chimeric subgenomic cDNAs were then cloned into the full-length HPIV3 antigenomic cDNA p3/7(131)2G⁺, referred to here as pFLC HPIV3 wt, to yield pFLC HPIV3 1HN_{N,P} and pFLC

HPIV3 1HN_{P-M}, respectively (Figure 11, the plasmids from which the second and third recombinant viruses from the top were isolated).

The HPIV2 HN ORF, available within an *NcoI* to *HindIII* restriction fragment of p32Hnhc#3 31hc (Tao et al., *J. Virol.* 72:2955-2961, 1998, incorporated herein by reference), was inserted into the *NcoI* to *HindIII* sites of pUC(GE/GS-H-N)_{N-P} and pUC(GE/GS-H-N)_{P-M} to generate pUC 2HN_{N-P} and pUC 2HN_{P-M}, respectively. Short oligonucleotide duplexes were inserted in the unique *MluI* restriction site to adjust the sequence to conform to the rule of six. Inadvertently, the inserted oligonucleotide was one nucleotide shorter that that required to specify that the genome of the recovered virus would conform to the rule of six. Therefore, all cDNAs bearing the HIV2 HN gene insertion did not conform to the rule of six. Nonetheless, virus was recovered from each of these cDNAs. These chimeric subgenomic cDNAs were cloned into the full-length PIV3 antigenomic cDNA pFLC HPIV3 wt to yield pFLC PIV3 2HN_(N-P) and pFLC PIV3 2HN_(P-M), respectively (Figure 11, plasmids from which the fourth and fifth recombinant viruses from the top were isolated).

Additional recombinant HPIV3 antigenomic cDNAs were assembled that contained up to three supernumerary foreign genes in various combinations and locations in the HPIV3 backbone (Figure 11). These antigenomic cDNAs were assembled from the subgenomic cDNAs described above in which the HN of HPIV1 or HPIV2 was inserted between the N and P genes or the P and M genes. Other subclones used for assembly contained the measles virus HA gene between the P/M genes or HN/L genes as described above. Another subclone used in assembly contained the 3918-nt GU between the HN and L genes, as described above.

The recombinants containing two or three supernumerary inserts were as follows: rHPIV3 1HN_{N-P} 2HN_{P-M} (Figure 11, sixth recombinant from the top) contained the HPIV1 HN and HPIV2 HN genes inserted between the N/P and P/M genes, respectively; rHPIV3 1HN_{N-P} 2HN_{P-M} HA_{HN-L} (Figure 11, seventh recombinant) contained the HPIV1 HN, HPIV2 HN, and measles virus HA inserted between the N/P, P/M, and HN/L genes, respectively; and rPIV3 1HN_{N-P} 2HN_{P-M} 3918GU_{HN-L} (Figure 11, bottom), contained the HPIV1 HN and HPIV2 HN genes inserted between the N/P and P/M genes, respectively, and in addition contained the 3918-nt GU insert between the HN and L genes.

It is noteworthy that the penultimate of these constructs, rHPIV3 1HN_{N-P} 2HN_{P-M} HA_{HN-L} (Figure 11, seventh construct from the top), contained protective antigens for four pathogens: HPIV3 (HN and F), HPIV1 (HN), HPIV2 (HN), and measles virus (HA). The total length of foreign sequence inserted into this recombinant was about 5.5 kb, which is 36% of the total HPIV3 genome length of 15,462 nt. The last recombinant, rHPIV3-1HN_{N-P}2HN_{P-M}GU_{HN-L} (Figure 11, bottom), was approximately 23 kb in length. This is 50% longer than wild-type HPIV3, and longer than any previously described biologically derived or recombinant paramyxovirus.

Recovery and replication *in vitro* of recombinant rHPIV3 bearing one, two, or three supernumerary gene inserts

The full length HPIV3 antigenomic cDNAs bearing single or multiple supernumerary genes of heterologous paramyxovirus protective antigens were separately transfected into HEp-2 monolayer cultures on six-well plates (Costar, Cambridge, MA) together with the support plasmids pTM(N), pTM(P no C), and pTM(L) and LipofectACE (Life Technologies, Gaithersburg, MD) and the cells were simultaneously infected with MVA-T7, a replication-defective vaccinia virus recombinant encoding the bacteriophage T7 polymerase protein using techniques previously described (Durbin et al., *Virology* 235:323-332, 1997a; Skiadopoulos et al., *Virology* 272:225-34, 2000, each incorporated herein by reference). After incubation at 32°C for up to four days, the transfection harvest was passaged onto LLC-MK2 monolayer cultures in a 25 cm² flask and the cells were incubated for 5 days at 32°C. The virus recovered from the cell supernatant was further passaged on LLC-MK2 cells at 32°C to amplify the virus. rHPIV3s bearing single or multiple foreign gene inserts were biologically-cloned by plaque purification on LLC-MK2 cells as previously described (Skiadopoulos et al., *J. Virol.* 73:1374-81, 1999a, incorporated herein by reference). Viral suspensions derived from biologically cloned virus were amplified on LLC-MK2 cells and yielded final titers of 10⁷ and 10⁹ TCID₅₀/ml, similar to the range of titers typically obtained for wt rHPIV3. Recombinant viruses were assayed for their ability to grow at 39°C. Surprisingly several rHPIV3s bearing single or multiple foreign gene insertions (rHPIV3 1HN_{N-P}, rHPIV3 1HN_{N-P}2HN_{P-M}HA_{HN-L}, and rHPIV3 1HN_{N-P} 2HN_{P-M} 3918 GU_{HN-L}) were 100 to 1000-fold restricted for replication at 39°C compared to the replication at the permissive temperature.

Viral RNA (vRNA) was isolated from biologically cloned recombinant chimeric viruses as described above (see also, Skiadopoulos et al., *J. Virol.* 73:1374-81, 1999a, incorporated herein by reference). This was used as the template for reverse transcription and polymerase chain reaction (RT-PCR) using specific primers that border the insertion sites. The amplified products were analyzed by restriction endonuclease digestion and partial DNA sequencing of the junction regions to confirm the presence and identity of each foreign insert. In all cases, the expected, correct sequence was confirmed.

Replication in the respiratory tract of hamsters of rHPIV3s expressing one, two, or three supernumerary foreign protective antigens

It was previously shown that rHPIV3 or rHPIV3-1 viruses expressing one supernumerary viral protective antigen gene replicated efficiently *in vitro* and *in vivo* and induced protective immune responses against both the vector virus and the virus represented by the supernumerary antigen gene. However, it was unknown whether a rHPIV could accommodate two or more supernumerary genes and retain the ability to replicate efficiently *in vitro* and *in vivo* and to induce protective immune responses against both the vector and the expressed supernumerary antigens. The present example indicates that this is indeed possible.

Hamsters in groups of eight were inoculated intranasally with 10^6 TCID₅₀ of each rHPIV3 bearing single or multiple supernumerary foreign gene inserts or with control viruses (Table 13). Nasal turbinates and lungs were harvested four days post infection and the virus present in tissue homogenates was quantified by serial dilution on LLC-MK2 monolayer cultures at 32°C as described above (see also, Skiadopoulos et al., *J. Virol.* 73:1374-81, 1999a). Virus was detected by hemadsorption with guinea pig erythrocytes, and the mean virus titer for each group is expressed as log₁₀ TCID₅₀(50% tissue culture infectious dose/gram tissue ± SE).

Table 13
Replication of recombinant HPIV3s containing single or multiple supernumerary gene inserts expressing the HPIV1, HPIV2 or measles virus glycoprotein genes in the upper and lower respiratory tract of hamsters

Group ^a no.	Virus ^b	Mean virus titer ^c (\log_{10} TCID ₅₀ /g \pm S.E.) in:			
		Nasal Turbinates	titer reduction (\log_{10}) ^d	Lungs	titer reduction (\log_{10}) ^d
1	rHPIV3 1HN ₍₁₋₁₇₎	4.5 \pm 0.2	1.8	3.9 \pm 0.2	3.0
2	rHPIV3 1HN ₍₁₋₄₀₎	3.5 \pm 0.2	2.8	4.3 \pm 0.2	2.3
3	rHPIV3 2HN ₍₁₋₁₇₎	5.4 \pm 0.2	0.9	5.3 \pm 0.3	1.6
4	rHPIV3 2HN ₍₁₋₄₀₎	6.3 \pm 0.1	0.0	6.3 \pm 0.5	0.6
5	rHPIV3 HA ₍₁₋₁₇₎	5.2 \pm 0.2	1.0	5.8 \pm 0.4	1.1
6	rHPIV3 HA ₍₁₋₄₀₎	6.0 \pm 0.2	0.3	7.3 \pm 0.2	-0.4
7	rHPIV3 HA ₍₁₀₀₋₄₎	6.0 \pm 0.1	0.3	6.6 \pm 0.2	0.3
8	rHPIV3 1HN ₍₁₋₁₇₎ 2HN ₍₁₋₄₀₎	5.2 \pm 0.1	1.1	5.0 \pm 0.3	1.9
9	rHPIV3 1HN ₍₁₋₁₇₎ 2HN ₍₁₋₁₇₎ HA ₍₁₀₀₋₄₎	1.6 \pm 0.1	4.7	2.5 \pm 0.1	4.4
10	rHPIV3 1HN ₍₁₋₁₇₎ 2HN ₍₁₋₁₇₎ 3918 GU ₍₁₀₀₋₄₎	2.0 \pm 0.3	4.3	1.8 \pm 0.2	5.1
11	rHPIV3 cpA5	4.6 \pm 0.1	1.7	2.1 \pm 0.2	4.8
12	rHPIV3 wt	6.3 \pm 0.1	-	6.9 \pm 0.1	-

a. 8 hamsters per group.
 b. Each hamster was inoculated with 10^6 TCID₅₀ of virus in a 0.1 ml inoculum.
 c. Virus was titered by serial dilution on LLC-MK2 monolayer cultures at 32°C.
 d. Reduction in virus replication compared to rHPIV3 wt (group 12).

It was shown above that a rHPIV3 expressing measles virus HA from a supernumerary gene insert between the HPIV3 HN and L genes, between the N and P genes, or between the P and M genes was modestly (about 10 to 20-fold) restricted in replication in the upper and lower respiratory tract of hamsters. This was confirmed in the present
5 experiment, in which rHPIV3 containing measles virus HA as a single supernumerary gene between the N/P, P/M or HN/L genes was attenuated up to 10-fold (Table 13, groups 5, 6, and 7). Similarly, insertion of the HPIV2 HN gene between the HPIV3 N and P genes or between the P and M genes also exhibited a modest reduction (about 10 to 20-fold) in replication in the respiratory tract of hamsters (Table 13 groups 3 and 4). In contrast,
10 insertion of the HPIV1 HN gene between the P and M genes or between the N and P, resulted in over approximately 100-fold reduction of replication in the upper and lower respiratory tract of hamsters (Table 13, groups 1 and 2). Since the HPIV1 HN, HPIV2 HN, and measles virus HA gene insertions are all of approximately the same size (1794 nt, 1781 nt, and 1926 nt, respectively), this was unlikely to be due to insert length. Therefore, the
15 greater level of attenuation conferred by the introduction the HPIV1 HN gene likely is due to an additional attenuating effect that is specific to the expression of the HPIV1 HN protein on replication of the HPIV3 vector. Thus, in some cases, such as with HPIV1 HN, a supernumerary antigen can attenuate rHPIV3 for hamsters above and beyond the modest attenuation due to inserting an additional gene.

20 Inspection of the data in Table 13 indicates that the site of insertion also plays a role in the level of restriction of replication of the chimeric rHPIV3 in the respiratory tract of hamsters. Insertion of the measles virus HA gene or the HPIV2 HN gene between the rHPIV3 N and P genes resulted in a greater reduction of replication in the upper and lower respiratory tract of hamsters than did insertion between the P and M genes (Table 13,
25 compare groups 3 versus 4 and 5 versus 6). This site-specific attenuation effect on replication of the HPIV3 vector was not evident for insertions of the HPIV1 HN gene, presumably because it was masked by the more substantial attenuating effect specific to HPIV1 HN.

The rHPIV3 chimeric recombinant viruses exhibited a gradient of attenuation
30 that was a function of the number of supernumerary gene inserts. The viruses bearing three added genes exhibited the greatest effect, and were reduced approximately 10,000 - 108,000 fold in replication in the upper and lower respiratory tract of the infected hamsters (Table 13, groups 9 and 10). The rHPIV3 chimeric recombinant virus bearing two gene inserts

exhibited an intermediate level of attenuation, and was reduced approximately 12-80 fold (Table 13, group 8). rHPIV3 chimeric recombinant viruses bearing one supernumerary gene (except those bearing the HPIV1 HN gene) were reduced only approximately 10-25 fold (groups 3-7 in Table 13). Importantly, rHPIV3 chimeric recombinant viruses bearing one, two, or three supernumerary gene inserts replicated in all animals. The most attenuated of these viruses, namely those bearing three supernumerary genes, were substantially more attenuated than rcp45 (group 11) with respect to replication in the upper and lower respiratory tract.

10 Immunogenicity in hamsters of rHPIV3s expressing one, two, or three supernumerary foreign protective antigens

Hamsters were infected with HPIV1 wt, HPIV2 wt, rHPIV3 wt, or rHPIV3s bearing single, double or triple supernumerary gene inserts as described above. Serum samples were collected 3 days pre-immunization and 28 days post-immunization and were assayed for HPIV1, HPIV2, HPIV3 or measles virus-specific antibodies by virus neutralizing assay specific for either HPIV1 or measles virus, or by the hemagglutination inhibition (HAI) assay for HPIV3 or HPIV2 HN-specific antibodies (Table 14). All rHPIV3 viruses elicited a strong immune response to the HPIV3 backbone with the exception of the viruses bearing the three supernumerary gene insertions. The reduced or absent immune response in hamsters infected with either the rHPIV3 1HN_{N-P} 2HN_{N-P} HA_{HN-L} or rHPIV3 1HN_{N-P} 2HN_{N-P} 3918GU_{HN-L} was likely a result of these viruses being overly attenuated for replication in hamsters. Likewise the immune response to the vectored antigens in the viruses bearing three foreign genes was also low or undetectable. In contrast, viruses bearing single or double foreign gene insertions induced an immune response against each of the additional antigens, demonstrating that the vectored foreign genes are immunogenic in hamsters, and as in the example of rHPIV3 1HN_{N-P} 2HN_{N-P} (Table 14; group 11) can be used to induce a strong immune response to three different viruses: HPIV1, HPIV2 and HPIV3.

Table 14
Immune response in hamsters to immunization with rHPV3 vectors expressing single or multiple superimposed protective antigens of HPV1, HPV2, or measles virus^a
Serum^b antibody titer (mean log₁₀ ± SE) to the indicated virus

Group no.	Virus	HPV3 ^c	HPV1 ^d	HPV2 ^e	Measles virus ^f
1	rHPV3 wt	10.0 ± 0	-	-	-
2	HPV2 wt	<2.0 ± 0	-	8.0 ± 0.0	-
3	HPV1 wt	<2.0 ± 0	5.4 ± 0.3	-	-
4	rHPV3 HA _(P+M)	9.5 ± 0.2	-	-	12.4 ± 0.4
5	rHPV3 HA _(P+M)	8.7 ± 1.4	-	-	11.8 ± 0.2
6	rHPV3 HA _(N+L)	9.0 ± 0	-	-	8.1 ± 0.6
7	rHPV3 1HN _(P+M)	9.5 ± 0.2	3.4 ± 0.6	-	-
8	rHPV3 1HN _(P+M)	7.2 ± 0.8	2.7 ± 0.3	-	-
9	rHPV3 2HN _(P+M)	9.8 ± 0.5	-	9.3 ± 0.8	-
10	rHPV3 2HN _(P+M)	10.0 ± 0.5	-	8.3 ± 1.1	-
11	rHPV3 1HN _(P+M) 2HN _(P+M)	9.6 ± 0.7	5.5 ± 0.4	8.3 ± 0.8	-
12	rHPV3 1HN _(P+M) 2HN _(P+M) HA _(N+L)	<2.0 ± 0	1.0 ± 0.3	<2.0 ± 0.0	<3
13	rHPV3 1HN _(P+M) 2HN _(P+M) 3918 GU _(N+L)	4.3 ± 0.7	2.3 ± 0.6	<2.0 ± 0.0	-
14	rHPV3 cp45	7.7 ± 0.2	-	-	-

a. Mean antibody response in groups of hamsters (n=6) inoculated intranasally with 10⁷ TCID₅₀ rHPV3s expressing the hemagglutinin-neuraminidase protein of HPV1 (1HN), HPV2 (2HN) or measles virus hemagglutination (HA) inserted between the N and P genes (N-P), the P and M genes (P-M) or the HN and L genes (HN-L) of rHPV3.

b. Sera were collected 3 days before and 28 days after immunization.

c. Mean hemagglutination inhibiting antibody (HAI) titer to HPV3.

d. Mean neutralizing antibody titer to HPV1.

e. Mean HAI antibody titer to HPV2.

f. Mean neutralizing antibody titer to measles virus (60% plaque reduction neutralization, PRN).

EXAMPLE IXUse of rHPIV3-N_B as an Attenuated Vector for the Measles Virus HA Protein.

The use of an animal virus that is attenuated in humans because of a host range restriction as a vaccine against an antigenically-related human counterpart is the basis of the Jennerian approach to vaccine development. The Kansas (Ka) strain of bovine parainfluenza virus type 3 (BPIV3) was found to be 100- to 1000-fold restricted in replication in rhesus monkeys compared to human parainfluenza virus type 3 (HPIV3), and was also shown to be attenuated in humans (Coelingh et al., *J. Infect. Dis.* 157:655-62, 1988; Karron et al., *J. Infect. Dis.* 171:1107-14, 1995b, each incorporated herein by reference). A viable chimeric recombinant human parainfluenza virus type 3 (HPIV3) virus was previously produced containing the nucleoprotein (N) open reading frame (ORF) from BPIV3 Ka in place of the HPIV3 N ORF. This chimeric recombinant was previously designated cKa-N (Bailly et al., *J. Virol.* 74:3188-3195, 2000a, incorporated herein by reference) and is referred to here as rHPIV3-N_B. This previous study was initiated with an exchange of the N ORF because, among the PIV3 proteins, the BPIV3 and HPIV3 N proteins possess an intermediate level of amino acid sequence identity (85%) (Bailly et al., *Virus Genes* 20:173-82, 2000b, incorporated herein by reference), and it was shown that such a BPIV3/HPIV3 N recombinant is viable (Bailly et al., *J. Virol.* 74:3188-3195, 2000a, incorporated herein by reference). This represents a "modified Jennerian" approach, in which only a subset of the genes in the vaccine virus is derived from the animal counterpart. rHPIV3-N_B grew to a titer comparable to that of the rHPIV3 and BPIV3 parent viruses in LLC-MK2 monkey kidney and Madin Darby bovine kidney cells (Bailly et al., *J. Virol.* 74:3188-3195, 2000a). Thus, the heterologous nature of the N protein did not impede replication of rHPIV3-N_B *in vitro*. However, rHPIV3-N_B was restricted in replication in rhesus monkeys to a similar extent as its BPIV3 parent virus (Bailly et al., *J. Virol.* 74:3188-3195, 2000a). This identified the BPIV3 N protein as a determinant of the host range restriction of replication of BPIV3 in primates.

The rHPIV3-N_B chimeric virus thus combines the antigenic determinants of HPIV3 with the host range restriction and attenuation phenotype of BPIV3. There are 79 differences out of a total of 515 amino acids between the N proteins of HPIV3 and BPIV3 (Bailly et al., *Virus Genes* 20:173-82, 2000b). Many of these 79 amino acid differences likely contribute to the host-range attenuation phenotype of rHPIV3-N_B. Because the host range restriction is anticipated to be based on numerous amino acid differences, it is

anticipated that the attenuation phenotype of rHPIV3-N_B will be stable genetically even following prolonged replication *in vivo*. Despite its restricted replication in rhesus monkeys, rHPIV3-N_B induced a high level of resistance to challenge of the monkeys with wild type (wt) HPIV3, and this level of resistance was indistinguishable from that conferred by immunization with wt rHPIV3. The infectivity, attenuation, and immunogenicity of rHPIV3-N_B suggest that this novel chimeric virus is an excellent candidate as a HPIV3 vaccine (Bailly et al., *J. Virol.* 74:3188-3195, 2000a). Furthermore, as described below, it is shown herein that such chimeric viruses are excellent candidates to serve as an attenuated vector for the expression of supernumerary protective antigens, such as the HA of measles virus. The vector component of the resulting chimeric virus induces an immune response against HPIV3, and the added supernumerary genes induce immune responses against their respective heterologous pathogens. In this specific example, a bivalent attenuated vaccine virus is made that simultaneously induces immune response to HPIV3 and measles virus.

It is shown above that rHPIV3 can be used as a vector for expression of the measles virus hemagglutinin (HA) protein. In two examples, *rcp45_L* HA(N-P) and *rcp45* HA(HN-L), attenuated vectors expressing the measles virus HA gene possessed three attenuating amino acid point mutations in the vector backbone. The rHPIV3-N_B vector of the present invention will likely be even more stable than vectors having an attenuation phenotype based on three amino acid point mutations. Also above, it was shown that the insertion of HA as a supernumerary gene into rHPIV3 conferred some attenuation on replication of both wt HPIV3 and attenuated HPIV3*cp45_L* for hamsters. In addition, the insertion of a 1908-nt gene insert into HPIV3 did not attenuate the wild type backbone, but did increase the level of attenuation of a backbone bearing the *cp45_L* mutations for replication in hamsters. Therefore, the insertion of the measles virus HA gene into the host-range restricted rHPIV3-N_B virus is projected to further attenuate its growth *in vitro* and/or *in vivo*. Inserts that affect replication *in vitro* or *in vivo* can be problematic for development of specific vaccines such as rHPIV3-N_B. Specifically, a candidate virus that is highly restricted in replication *in vitro* would be difficult to manufacture, and one that is highly restricted in replication *in vivo* could be overattenuated and not useful as a vaccine. It was also not known whether the rHPIV3-N_B chimeric virus expressing the measles virus HA glycoprotein, designated rHPIV3-N_B HA_(P-M), would be satisfactorily immunogenic in primates against both HPIV3 and measles virus since all previous studies with HPIV3 expressing HA were conducted in a rodent model.

The present example, which details the generation of rHPIV3-N_B HA_(P-M) using reverse genetic techniques, indicates, surprisingly, that insertion of the HA gene into rHPIV3-N_B did not further restrict its replication in rhesus monkeys. Presumably the attenuating effect of insertion is masked by the genetic elements present in the N_B gene that specify the host range restriction of replication in primates. Rather, rHPIV3-N_B HA_(P-M) was satisfactorily attenuated in rhesus monkeys. Immunization of rhesus monkeys with rHPIV3-N_B HA_(P-M) induced resistance to the replication of wt HPIV3 challenge virus and stimulated high levels of neutralizing antibodies to the measles virus, levels that are known to be protective in humans (Chen et al., *J. Infect. Dis.* 162:1036-42, 1990, incorporated herein by reference).

Construction of a pFLC HPIV3-N_B HA_(P-M), a chimeric bovine/human PIV3 antigenomic cDNA encoding the BPIV3 N gene ORF in place of the rHPIV3 N gene ORF and the HA gene of measles virus as a supernumerary gene inserted between the rHPIV3 P and M genes

The full length antigenomic cDNA plasmid pFLC HPIV3-N_B HA_(P-M) (Figure 12) was constructed in two steps. First, the previously-constructed pLeft-N_B plasmid contains the 3' half of the HPIV3 antigenomic cDNA (HPIV3 nts 1-7437, the latter position being an XhoI site within the HN gene) with the HPIV3 N ORF replaced by that of BPIV3 (Baillly et al., *J. Virol.* 74:3188-3195, 2000a, incorporated herein by reference). The *PshAI*-*NgoMIV* fragment was excised from this plasmid. Note that the *PshAI* site is at position 2147 in the HPIV3 antigenome sequence (see Figure 12) and the *NgoMIV* site occurs in the vector sequence, and so this removes all of the HPIV3 sequence downstream of the *PshAI* site. This fragment was replaced by the *PshAI*-*NgoMIV* fragment from the previously-constructed plasmid pLeft HA_(P-M), which contains the measles virus HA ORF under the control of HPIV3 transcription signals and inserted between the HPIV3 N and P genes (Durbin, *J. Virol.* 74:6821-31, 2000, incorporated herein by reference). This yielded pLeft-N_B HA_(P-M). Next, the 11899 nt *NgoMIV* to *Xho* I fragment of pLeft N_B HA_(P-M), containing the 3' half of the HPIV3 antigenomic cDNA including the BPIV3 N gene ORF and the measles virus HA gene insert, was cloned into the *NgoMIV* to *Xho* I window of pRight, a plasmid encoding the 5' half of the HPIV3 antigenomic cDNA (PIV3 nts 7462-15462) (Durbin et al., *Virology* 235:323-332, 1997a). This yielded pFLC HPIV3-N_B HA_(P-M), a plasmid bearing the full length antigenomic cDNA of HPIV3 containing the BPIV3 N ORF

in place of the HPIV3 N ORF, and containing measles virus HA gene as a supernumerary gene inserted between the P and M genes of HPIV3.

5 **Recovery of chimeric rHPIV3 expressing the bovine N gene and the measles virus HA gene**

rHPIV3-N_B HA_{P-M} was recovered from HEp-2 cells transfected with pFLC HPIV3-N_B HA_{P-M}. To accomplish this, pFLC HPIV3-N_B HA_{P-M} was transfected into HEp-2 cells on six-well plates (Costar, Cambridge, MA) together with the support plasmids pTM(N), pTM(P no C), and pTM(L) and LipofectACE (Life Technologies, Gaithersburg, MD), and the cells were simultaneously infected with MVA-T7, a replication-defective vaccinia virus recombinant encoding the bacteriophage T7 polymerase protein, as described above. After incubation at 32°C for four days, the transfection harvest was passaged onto LLC-MK2 cells in a 25 cm² flask, and the cells were incubated for 5 days at 32°C. The virus recovered from the cell supernatant was amplified by a further passage on LLC-MK2 cells at 32°C. rHPIV3-N_B HA_{P-M} was biologically cloned by plaque purification on LLC-MK2 monolayer cultures as described above. Viral suspensions derived from biologically cloned virus were amplified on LLC-MK2 monolayer cultures at 32°C. Viral RNA (vRNA) was isolated from biologically cloned recombinant chimeric viruses as described above. RT-PCR was performed using specific oligonucleotide primer pairs spanning the BPIV3 N ORF or the measles virus HA gene, and the amplified cDNAs were analyzed by restriction endonuclease digestion and partial DNA sequencing as described above. This confirmed the presence of the BPIV3 N ORF substitution and the measles virus HA supernumerary gene insert.

25 Expression of the measles virus HA protein was initially confirmed by immunostaining plaques formed on LLC-MK2 monolayer cultures infected with rHPIV3-N_B HA_{P-M} using mouse monoclonal antibodies specific to the measles virus HA protein and goat anti-mouse peroxidase-conjugated antibodies, as described previously (Durbin, *J. Virol.* 74:6821-31, 2000, incorporated herein by reference).

30 **rHPIV3-N_B HA_{P-M} replicates to the same level as rHPIV3-N_B in the respiratory tract of rhesus monkeys.**

It was next determined whether the acquisition of the measles virus HA insert significantly decreased the replication of rHPIV3-N_B in the upper and lower respiratory tract, as was observed when a supernumerary gene was inserted into an attenuated HPIV3 backbone lacking a bovine chimeric component. It was also determined whether rHPIV3-N_B HA_{P-M} replicated sufficiently to induce an immune response against both HPIV3 and measles virus *in vivo*. The replication of rHPIV3-N_B HA_{P-M} in the upper and lower respiratory tract of rhesus monkeys was compared to that of its rHPIV3-N_B parent as well as wt HPIV3 and wt BPIV3 (Table 15). Rhesus monkeys that were seronegative for both HPIV3 and measles virus were inoculated simultaneously by the intranasal (IN) and intratracheal (IT) routes with one milliliter per site of L15 medium containing 10⁵ TCID₅₀ of virus suspension, as described previously (Bailly et al., *J. Virol.* 74:3188-3195, 2000a). Nasopharyngeal (NP) swab samples were collected on days 1 through 10 post-infection, and tracheal lavage (TL) samples were collected on days 2, 4, 6, 8, and 10 post-infection. Virus present in the NP and TL specimens was quantified by serial dilution on LLC-MK2 cell monolayers at 32°C, and the titer obtained was expressed as log¹⁰ TCID₅₀/ml (Table 15).

This comparison showed that the rHPIV3-N_B HA_(P-M) chimeric virus replicated to the same level in the upper and lower respiratory tract of rhesus monkeys as its rHPIV3-N_B parent virus. This level of replication was also comparable to that of the BPIV3 virus parent, demonstrating that rHPIV3-N_B HA_(P-M) retains the attenuation phenotype of rHPIV3-N_B and BPIV3 and, unexpectedly, that the insertion of the measles virus HA gene into the rHPIV3-N_B genome does not further attenuate this virus for replication in the respiratory tract of rhesus monkeys.

Table 15

A chimeric human/bovine FPIV3 containing the measles virus hemagglutinin gene is satisfactorily attenuated for replication in the upper and lower respiratory tract of rhesus monkeys, induces antibodies to both HPIV3 and measles virus, and protects against HPIV3 wild type virus challenge

Group no.	Immunizing virus ^a	No. of animals ^a	Response to immunization				Response to challenge with HPIV3 wt on day 28 or 31				Response to administration of measles virus vaccine (Moraten) on day 59	
			Virus Replication		Serum antibody response		Virus replication		Serum antibody response		Serum antibody response	
			Mean peak virus titer ^c (log ₁₀ TCID ₅₀ /ml ± SE)	Tracheal lavage	Mean HAI antibody titer (mean reciprocal logs ± S.E.) for HPIV3 on day 28/31 ^{d,e}	Mean antibody titer to measles virus (60% PRN mean reciprocal logs ± SE) (day 31 post immunization) ^f	Mean peak HPIV3 virus titer ^a (log ₁₀ TCID ₅₀ /ml ± SE)	Tracheal lavage	HAI antibody titer (mean reciprocal logs ± SE) for HPIV3 on day 50/59 ^b	Mean antibody tier to measles virus (60% PRN mean reciprocal logs ± SE) (day 87 post first immunization) ^g	Mean antibody tier to measles virus (60% PRN mean reciprocal logs ± SE) (day 87 post first immunization) ^g	Mean antibody tier to measles virus (60% PRN mean reciprocal logs ± SE) (day 87 post first immunization) ^g
1	rHPIV3 wt	6	4.9 ± 0.4	3.2 ± 0.6	9.3 ± 0.6	< 5.5 ± 0.0	0.5 ± 0.0	0.3 ± 0.0	12.0 ± 0.0	8.2 ± 0.8		
2	rHPIV3-N ₆	8	2.6 ± 0.6	2.0 ± 0.4	7.3 ± 0.3	< 5.5 ± 0.0	1.4 ± 0.9	0.5 ± 0.0	9 ± 1.0	10.1 ± 0.4		
3	rHPIV3-N ₆ HA _(r-m)	4	2.2 ± 0.6	2.8 ± 0.6	6.8 ± 0.3	9.6 ± 0.5	1.2 ± 0.7	2.3 ± 0.2	11.5 ± 0.3	10.2 ± 0.4		
4	BPIV3 Ka	8	2.3 ± 0.2	1.9 ± 0.2	5.0 ± 0.4	ND	2.9 ± 0.3	2.0 ± 0.5	11.5 ± 0.3	ND		
5	none ^b	4	ND	ND	< 2	ND	4.5 ± 0.3	4.5 ± 0.2	12.0 ± 0.6	ND		

a. The present study included 4 monkeys that received rHPIV3-N₆ HA_(r-m) and two monkeys in each of the groups that received rHPIV3 wt, rHPIV3-N₆ or BPIV3 Ka. With the exception of the group that received rHPIV3-N₆ HA_(r-m), the data presented includes historical data from studies reported in Bailey et al., *J. Virol.* 74:3188-3195, 2000, and Schmidt et al., *J. Virol.* 74:8922-8929, 2000.
 b. Historical data from Schmidt et al., *J. Virol.* 74:8922-8929, 2000.
 c. Monkeys were inoculated intranasally and intratracheally with 10⁷ TCID₅₀ of virus in a 1 ml inoculum at each site. Nasopharyngeal (NP) swab samples were collected on days 1 to 10 post-infection. Tracheal lavages (TL) samples were collected on days 2, 4, 6, 8, and 10 post-infection. Mean of the peak virus titers for each animal in its group irrespective of sampling day. S.E. = standard error. Virus titrations were performed on LLC-MK2 cells at 32°C. The limit of detection of virus titer was 10 TCID₅₀/ml.
 d. In the present study sera were collected from monkeys on day 31 post immunization and animals were then challenged with HPIV3. In the two previous studies, monkeys were sampled and challenged on day 28 post immunization.
 e. Sera collected for the present study and from the two previous studies were assayed at the same time. Serum HAI titer is expressed as the mean reciprocal logs ± standard error, SE.
 f. In the present study sera were collected from monkeys on day 31 post immunization and animals were then challenged with HPIV3. In the two previous studies, monkeys were sampled and challenged on day 28 post immunization.
 g. Sera collected for the present study and from the two previous studies were assayed at the same time. Serum HAI titer is expressed as the mean reciprocal logs ± standard error, SE.

- f. Animals were immunized on day 59 with 10^6 pfu of the measles virus Moraten vaccine strain administered parenterally (IM). Serum was collected 28 days later (i.e., 87 days after the first immunization). Data shown was obtained from samples collected only from animals in the present study. Mean neutralizing antibody titer to wt measles virus is expressed as the mean reciprocal log₂ standard error. PRN, plaque reduction neutralizing.
- g. 28 or 31 days after immunization monkeys were inoculated intranasally and intratracheally with 10^6 TCID₅₀ of wt HPIV3 in a 1 ml inoculum at each site. NP and TL samples were collected on days 0, 2, 4, 6 and 8 post challenge. The titers obtained for NP and TL samples on day 0 were $< 2.0 \log_{10}$ TCID₅₀/ml.
- h. With the exception of group 5, data shown are from the present study.

Immunization of rhesus monkeys with rHPIV3-N_B HA_{P,M} induces high titers of antibodies against both HPIV3 and measles virus and protects the monkeys from challenge with HPIV3

5 Rhesus monkeys immunized with rHPIV3-N_B HA_{P,M} developed high levels of serum antibodies against both HPIV3 and measles virus (Table 15). Serum HPIV3 antibodies were quantified by hemagglutination inhibition assay (HAI) using guinea pig erythrocytes as described previously (Durbin, *J. Virol.* 74:6821-31, 2000, incorporated herein by reference), and the titers are expressed as mean reciprocal log₂ ± SE. High levels
10 of serum HAI antibodies to HPIV3 were induced by both rHPIV3-N_B HA_{P,M} and rHPIV3-N_B, demonstrating that these attenuated recombinants can induce a strong immune response against the backbone antigens of the HPIV3 vector. It was also found that rhesus monkeys immunized with rHPIV3-N_B HA_{P,M} developed high levels of serum measles virus neutralizing antibodies 31 days after immunization, levels that are in excess of those needed
15 to protect humans against infection with measles virus (Chen et al., *J. Infect. Dis.* 162:1036-42, 1990, incorporated herein by reference). Serum neutralizing antibody titers against wild type measles virus were quantified as described previously (Durbin, *J. Virol.* 74:6821-31, 2000), and the titers are expressed as reciprocal mean log₂ ± SE (Table 15).

To compare the ability of infection with the live attenuated rHPIV3-N_B HA_{P,M}
20 and rHPIV3-N_B virus vaccine candidates to protect against wt HPIV3, the monkeys were challenged IN and IT with 10⁶ TCID₅₀ of wt HPIV3 31 days after the initial infection (Table 15). Nasopharyngeal swab and tracheal lavage samples were collected on days 2, 4, 6, and 8 post-challenge. Virus present in the specimens was quantified by serial dilution on LLC-MK2 monolayer cultures as described above. rHPIV3-N_B HA_{P,M} and rHPIV3-N_B conferred
25 a comparable, high level of protection against challenge with wt HPIV3 as indicated by a 100 to 1000-fold reduction in wt HPIV3 replication in the respiratory tract of immunized monkeys. This demonstrated that insertion of the measles virus HA gene into the chimeric bovine/human PIV3 did not diminish the level of protection induced by the HPIV3 glycoproteins present in the backbone of the attenuated virus vector.

30 Immunogenicity of rHPIV3-N_B HA_{P,M} was then compared with that of the licensed Moraten strain of live attenuated measles virus vaccine in rhesus monkeys, a species in which both PIV3 and measles virus replicate efficiently. Rhesus monkeys previously infected with a rHPIV3 virus or with rHPIV3-N_B HA_{P,M} were immunized parenterally (IM)

with 10^6 pfu of the Moraten strain of live-attenuated measles virus vaccine on day 59, and serum samples were taken on day 87 and analyzed for neutralizing antibodies against measles virus (Table 15). In animals that were naive for measles virus before receiving the Moraten vaccine (Table 15, groups 1 and 2), the titer of measles-specific antibodies induced
5 by the Moraten vaccine was approximately the same as that observed in rHPIV3-N_B HA_{P-M}-immunized animals (Table 15, group 2). Thus, rHPIV3-N_B HA_{P-M} vector expressing the HA glycoprotein measles virus was equivalent to the Moraten vaccine in the ability to induce virus-neutralizing antibodies in this primate model.

An important advantage of rHPIV3-N_B HA_{P-M} as a vaccine for measles virus
10 over the Moraten vaccine is that the PIV vector can be administered by the intranasal route, whereas live-attenuated measles virus vaccines are not consistently infectious by this route, probably as a consequence of their attenuation and adaptation to cell culture. This makes it possible to immunize with rHPIV3-N_B HA_{P-M} in early infancy, an age group that cannot be immunized with a current live attenuated measles virus vaccine such as the Moraten strain
15 because of the neutralizing and immunosuppressive effects of maternal antibodies (Durbin, *J. Virol.* 74:6821-31, 2000, incorporated herein by reference). Other advantages are also described above, including the superior growth of the PIV vector in cell culture and the lack of incorporation of measles virus HA in the virions, which should preclude changing the tropism of the PIV vector and should preclude measles virus-induced immunosuppression.

The lack of effective vaccination against measles virus infection results in the
20 deaths of over 2700 children every day worldwide. The rHPIV3-N_B HA_(P-M) candidate vaccine offers a unique opportunity to immunize against two major causes of severe pediatric disease, namely, HPIV3 and measles virus. Unlike the currently licensed measles virus vaccines, we expect that chimeric rHPIV3-N_B HA_(P-M) and other human-bovine
25 chimeric vector constructs, expressing the major antigenic determinant of measles virus or other heterologous pathogens, can be used to induce a strong immune response to, e.g., measles virus, in infants and children younger than six months of age (Durbin, *J. Virol.* 74:6821-31, 2000). An effective immunization strategy for infants and children will be required to meet the World Health Organization goal to eradicate measles by the year 2010.
30 In particular, it would be advantageous for eradication to use a measles virus vaccine that does not involve infectious measles virus.

EXAMPLE X**Use of a Recombinant Bovine-Human Parainfluenza Virus Type 3 (rB/HPIV3) as a Vector for RSV Glycoprotein Supernumerary Genes**

For use within the present invention, a recombinant chimeric human-bovine PIV was constructed in which the BPIV3 F and HN genes were replaced with those of HPIV3. This recombinant chimeric bovine-human virus rB/HPIV3 was shown to be fully competent for replication in cell culture, whereas in rhesus monkeys it displayed the host range-restricted, attenuated phenotype characteristic of BPIV3 and was highly immunogenic and protective (U.S. Patent Application Serial No. 09/586,479, filed June 1, 2000 by Schmidt et al.; Schmidt et al., *J. Virol.* 74:8922-9, 2000, each incorporated herein by reference). This is another example of a "modified Jennerian" approach that is useful within the compositions and methods of the invention, but in this case the entire set of viral "internal" genes is derived from BPIV3, with the antigenic determinants alone derived from HPIV3.

As noted above, there are numerous practical and safety considerations that favor vaccines based on a single PIV3 backbone, as opposed to a complex mixture of different viruses each of which must be separately attenuated and verified and which can interact in unpredictable ways. In addition, the host range restriction of BPIV3 confers an attenuation phenotype that should be very highly stable. In the present example, a recombinant chimeric human-bovine PIV3 (rB/HPIV3) was designed, rescued and characterized that encodes the respiratory syncytial virus (RSV) G or F glycoprotein, which are the major RSV neutralization and protective antigens. This example shows that rB/HPIV3 readily accepted the foreign RSV genes without a significant reduction of its replicative efficiency *in vitro* or *in vivo* and thus is a promising candidate vaccine and vector. This vector will be free of the problems of poor growth *in vitro* and instability that are otherwise characteristic of RSV.

Construction of antigenomic cDNAs encoding recombinant chimeric rB/HPIV3 viruses bearing an RSV subgroup A G or F ORF as an additional, supernumerary gene

A full length cDNA of the BPIV3 Kansas strain was constructed in which the F and HN glycoprotein genes of the bovine virus had been replaced with the corresponding genes of the HPIV3 JS strain (rB/HPIV3) (U.S. Patent Application Serial No. 09/586,479,

filed June 1, 2000 by Schmidt et al.; Schmidt et al., *J. Virol.* 74:8922-9, 2000, each incorporated herein by reference). For use within the present invention, this cDNA was modified to contain three additional unique restriction enzyme recognition sites. Specifically, a *BspI* site was introduced preceding the N ORF (nucleotide (nt) 103-109), an *AseI* site was introduced preceding the N gene end sequence and a *NotI* site was introduced preceding the P gene end sequence. These restriction enzyme recognition sites were introduced to facilitate the insertion of foreign, supernumerary genes into the genome of the chimeric B/HPIV3 virus genome. The sites were designed so that they did not disrupt any of the BPIV3 replication and transcription cis-acting elements. This specific example will describe insertion into the *BspI* site (Figure 13).

The previously described RSV subgroup A glycoprotein genes G and F (GenBank accession no. M74568) were modified for insertion into the promoter-proximal *BspI* site of B/HPIV3 (Figure 13). The strategy was to express each heterologous ORF as an additional, separate mRNA, and hence it was important that it be introduced into the rB/HPIV3 genome so that it was preceded by a BPIV3 gene start signal and followed by a BPIV3 gene end signal. The *BspI* insertion site followed the gene start signal of the N gene (Figure 13). Hence, for insertion at this site, the RSV ORF needed to be modified by insertion of a *BspI* site at its upstream end and addition of a BPIV3 gene end signal, intergenic region, gene start signal, and *BspI* site at its downstream end. For the RSV A G ORF, the forward PCR primer used was (5' to 3')

AAATTCGCTTAGCGATGTCCAAAAACAAGGACCAACGCACCGC (SEQ ID NO. 30),

the reverse primer was (5' to 3')

AAAAAGCTAAGCGCTAGCCTTAAATCCTAAGTTTTTCTTACTTTTTTACTACTG
GC GTGGTGTGTTGGGTGGAGATGAAGGTTGTGATGGG (SEQ ID NO. 31)(*BspI* site

underlined, ORF translational initiation and termination triplets in bold). For the RSV A F ORF, the forward PCR primer used was (5' to 3')

AAAGGCCTGCTTAGCAAAAAGCTAGCACAAATGGAGTTGCTAATCC
TCAAAGCAAAT GCAATTACC (SEQ ID NO. 32), and the reverse primer was (5' to 3')

AAAAGCTAAGCGCTAGCCTTCTTAAATCCTAAGTTTTTCTTACTTTTATTAGTTACT
AAATGCAATATTTATACCCTCAGTTGATC (SEQ ID NO. 33)(*BspI* site underlined, ORF translational initiation and termination triplets in bold).

The PCR products were digested with *BspI* and cloned into the modified full length cDNA clone using standard molecular cloning techniques. The resulting full length cDNA containing the RSV A G ORF was designated pB/HPIV3-G_A1 and the plasmid

containing the F ORF was designated pB/HPIV3-F_A1. The nucleotide sequence of each inserted gene was confirmed by restriction enzyme digestion and automated sequencing. All constructs were designed so that the final genome nucleotide length was a multiple of six, which has been shown to be a requirement for efficient RNA replication (Calain et al., *J. Virol.* 67:4822-30, 1993, incorporated herein by reference).

Recovery of rB/HPIV3-G1 and rB/HPIV3-F1 chimeric viruses from cDNA.

rB/HPIV3-G1 and rB/HPIV3-F1 viruses were recovered from the cDNAs pB/HPIV3-G_A1 and pB/HPIV3-F_A1, respectively. This was accomplished by the previously-described method in which HEp-2 cells were transfected with the respective antigenomic cDNA together with BPIV3 N, P and L support plasmids. The cells were simultaneously infected with a recombinant vaccinia virus, strain MVA, expressing the T7 RNA polymerase gene. The recovered recombinant viruses were cloned biologically by sequential terminal dilution in Vero cells. The presence of the inserted RSV G or F gene in the backbone of each recovered recombinant virus was confirmed by RT-PCR of viral RNA isolated from infected cells followed by restriction enzyme digestion and DNA sequencing. The sequence of the inserted gene and flanking regions in the recovered recombinant viruses was identical to that of the starting antigenomic cDNA.

rB/HPIV3-G1 and rB/HPIV3-F1 viruses replicate efficiently in cell culture.

The multicycle growth kinetics of rB/HPIV3-G1 and rB/HPIV3-F1 in LLC-MK2 cells were determined by infecting LLC-MK2 cell monolayers in triplicate at a multiplicity of infection (MOI) of 0.01 and harvesting samples at 24-hour intervals over a seven day period, as previously described (Bailly et al., *J. Virol.* 74:3188-3195, 2000a, incorporated herein by reference). These two viruses were compared with BPIV3 Ka, HPIV3 JS, rBIV3 Ka, and rB/HPIV3 (Figure 14). The two parental viruses bearing HPIV3 glycoproteins, namely HPIV3 and rB/HPIV3, appeared to replicate somewhat more rapidly than the others. However, the final titer achieved for each of the six viruses were similar with one exception: rB/HPIV3-F1 was approximately 8-fold reduced in its replicative capacity compared to the other viruses (Figure 14). This might be an effect of having this large gene in a promoter-proximal position, or might be an effect of the expression of a second fusogenic protein, or both. This latter possibility was suggested by the observation that rB/HPIV3-F1 induced massive syncytia, comparable to what is observed with wild type

RSV infection and greater than that observed with rB/HPIV3 or the other parental viruses. In comparison, rB/HPIV3-G1 induced less cytopathic effect and few syncytia in LLC-MK2 cells, comparable to rB/HPIV3. Nonetheless, rB/HPIV3-F1 and rB/HPIV3-G1 grew to a final titer of at least 10^7 TCID₅₀/ml in LLC-MK2 cells and in Vero cells. This indicates that each virus is fully-permissive for growth which will allow cost-efficient vaccine manufacture.

The rB/HPIV3-G1 and rB/HPIV3-F1 viruses replicate efficiently in the respiratory tract of hamsters

rB/HPIV3-G1 and rB/HPIV3-F1 were evaluated for their ability to replicate in the upper and lower respiratory tract of hamsters. The rB/HPIV3 parental virus, as well as the BPIV3 and HPIV3 biologically-derived viruses, were compared in parallel as controls (Table 16). Each virus was administered intranasally at a dose of 10^6 TCID₅₀, and one group received both rB/HPIV3-G1 and rB/HPIV3-F1. Animals from each group were sacrificed on days 4 and 5 post infection, and the virus titer in the nasal turbinates and lungs were determined by serial dilution. The level of replication of rB/HPIV3-G1 in the respiratory tract was very similar to, and statistically indistinguishable from, that of HPIV3 JS and BPIV3 Ka. Replication of rB/HPIV3-F1 appeared to be somewhat reduced on days 4 and 5 relative to the others, but this difference was not statistically significant in comparison with the biological BPIV3 virus, which in previous primate and clinical studies replicated sufficiently well to induce a protective immune response (Coelingh et al., *Virology* 162:137-143, 1988; Karron et al., *Pediatr. Infect. Dis. J.* 15:650-654, 1996, each incorporated herein by reference). Also, the titer of virus from the mixed infection of rB/HPIV3-G1 and rB/HPIV3-F1 appeared to be somewhat reduced in the lower respiratory tract on day 4, but this was not statistically significant. Replication of one of the control viruses, BPIV3 Ka, was somewhat reduced in the lower respiratory tract on day 5: this also was not statistically significant, and indicates that these small differences likely are not important. Thus, the rB/HPIV3-G1 and rB/HPIV3-F1 viruses appeared to be fully competent for replication *in vivo*, despite the presence of the 0.9 kb G or 1.8 kb F supernumerary gene next to the promoter.

Table 16
rB/HPiV3 bearing the RSV G or F ORF as a supernumerary gene in the promoter proximal position replicates efficiently in the respiratory tract of hamsters.

Immunizing virus ^a	Number of animals	Mean virus titer on day 4 ^b (log ₁₀ TCID ₅₀ /g ± S.E.) ^c		Mean virus titer on day 5 ^b (log ₁₀ TCID ₅₀ /g ± S.E.) ^c	
		Nasal turbinates	Lungs	Nasal turbinates	Lungs
rB/HPiV3-G1	6	5.9±0.1 (AB)	5.1±0.6 (A)	5.5±0.2 (A)	5.6±0.4 (AC)
rB/HPiV3-F1	6	5.1±0.3 (B)	4.6±0.2 (A)	5.7±0.2 (AB)	3.6±0.2 (BD)
rB/HPiV3-G1 & rB/HPiV3-F1	6	5.7±0.3 (BC)	4.3±0.8 (A)	5.6±0.2 (A)	5.9±0.2 (A)
rB/H PIV3	6	6.2±0.2 (AC)	5.2±0.6 (A)	6.5±0.1 (B)	5.7±0.6 (AC)
HPiV3 JS wild type	6	6.6±0.1 (A)	6.5±0.1 (A)	6.0±0.2 (AB)	6.0±0.4 (A)
BPIV3 Ka wild type	6	5.8±0.1 (AB)	6.1±0.2 (A)	5.3±0.2 (A)	4.2±0.5 (CD)

^a Hamsters were inoculated intranasally with 10⁶ TCID₅₀ of virus in a 0.1 ml inoculum.
^b Animals were sacrificed on day 4 or 5 post inoculation, as indicated, and virus titers in the nasal turbinates and lungs were determined by titration on LLC-MK2 (PIV3) or HEP-2 (RSV) cells at 32°C. The limit of detectability of virus was 10^{2.48} TCID₅₀/g tissue. S.E. = standard error.
^c Mean virus titers were assigned to similar groups (A, B, C, D) by the Tukey-Kramer test. Within each column, mean titers with different letters are statistically different (p<0.05). Titrers indicated with two letters are not significantly different from those indicated with either letter.

The rB/HPIV3-G1 and rB/HPIV3-F1 viruses induce serum antibodies to both HPIV3 and RSV

Hamsters were infected with rB/HPIV3-G1, rB/HPIV3-F1, or rB/HPIV3 as described above. An additional group received both rB/HPIV3-G1 and rB/HPIV3-F1, and another group was infected intranasally with RSV. Serum samples were collected 5 days post infection and assayed for RSV-specific antibodies by an ELISA test specific for the RSV F protein or RSV G protein (Table 17), and for HPIV3 HN-specific antibodies by the hemagglutination inhibiting (HAI) antibody assay (Table 18). The titer of F-specific or G-specific antibodies induced by the rB/HPIV3-F1 or rB/HPIV3-G1 virus, respectively, was 2- to 4-fold higher than that induced by wild type RSV. Animals inoculated with both rB/HPIV3-F1 and rB/HPIV3-G1 also had high titers of F-specific and G-specific antibodies. In addition to high ELISA titers against RSV G and F, rB/HPIV3-G1 and rB/HPIV3-F1 also induced RSV-neutralizing serum antibody titers that were higher than those induced by wt RSV (Table 18). Each of the viruses induced a titer of PIV3-specific antibody that was indistinguishable from that of their parent virus rB/HPIV3 (Table 18). Thus, the rB/HPIV3 vector bearing the F or G gene of RSV induced strong immune responses against both the RSV insert and the PIV vector.

Table 17
 Immunization of hamsters with rB/HPV3 expressing the RSV G or F ORF as a superumerary gene induces an antibody response against the RSV G or F protein.

Immunizing virus ^a	Animals per group	Serum IgG ELISA titer against RSV G protein ^b (mean recip. log ₂ S.E.)		Serum IgG ELISA titer against RSV F protein ^b (mean recip. log ₂ S.E.)		log ₂ fold increase
		Day 26		Day 26		
		Pre	Day 26	Pre	Day 26	
rB/HPV3-G1	12	6.0 ± 0.4 ^c	12.5 ± 0.5	6.7 ± 0.5 ^c	7.5 ± 0.5	0.8
rB/HPV3-F1	12	6.3 ± 0.3	7.2 ± 0.3	6.8 ± 0.3	16.2 ± 0.5	9.4
rB/HPV3-G1 & rB/HPV3-F1	12	6.5 ± 0.6	12.0 ± 0.9	7.3 ± 0.5	14.7 ± 0.4	7.4
rB/HPV3	12	6.5 ± 0.4	8.0 ± 0.4	7.3 ± 0.7	8.3 ± 0.8	1.0
RSV	12	6.8 ± 0.3	10.8 ± 0.4	7.3 ± 0.5	15.7 ± 0.4	8.2

^aHamsters were inoculated intranasally with 10⁶ TCID₅₀ of the virus in a 0.1 ml inoculum.
^bSerum samples were taken on day 26 post inoculation and analyzed by glycoprotein-specific ELISA for antibodies against RSV G or F protein, as indicated.
^cTiters in the pre serum specimen represent non-specific background levels of antibody in this sensitive ELISA.

Table 18
Immunization of hamsters with rB/HPV3s expressing the RSV G or F ORF induces neutralizing antibodies against RSV as well as hemagglutination-inhibiting (HAI) antibodies against HPV3.

Immunizing virus ^a	Animals per group	Serum neutralizing antibody response to RSV ^b (mean recip. log ₂ S.E.) ^d		Serum HAI antibody response to HPV3 ^c (mean recip. log ₂ S.E.) ^d	
		Pre	Day 26	Pre	Day 26
		rB/HPV3-G1	12	≤ 3.3	10.0±0.3 (A)
rB/HPV3-F1	12	≤ 3.3	9.3±0.3 (A)	≤ 2	8.8±0.1 (A)
rB/HPV3-G1 & rB/HPV3-F1	12	≤ 3.3	10.8±0.4 (A)	≤ 2	8.8±0.3 (A)
rB/HPV3	12	≤ 3.3	0.8±0.8 (B)	≤ 2	9.5±0.8 (A)
RSV	12	≤ 3.3	8.1±1.2 (A)	≤ 2	≤ 2 (B)

^a Hamsters were inoculated intranasally with 10⁶ TCID₅₀ of the indicated HPV3 or 10⁶ PFU of RSV in a 0.1 ml inoculum.

^b Serum samples were taken on day 26 post inoculation and antibody titers were determined by 50% plaque reduction neutralization test.

^c Serum samples were taken on day 26 post inoculation and antibody titers were determined by hemagglutination inhibition test.

^d Mean virus titers were assigned to similar groups (A, B) by the Tukey-Kramer test. Within each column, mean titers with different letters are statistically different (p<0.05).

The rB/HPIV3-G1 and rB/HPIV3-F1 viruses induce resistance to replication of HPIV3 and RSV challenge virus.

5 Hamsters immunized with rB/HPIV3, rB/HPIV3-G1, rB/HPIV3-F1, or
rB/HPIV3-G1 plus rB/HPIV3-F1 vaccine candidates were challenged 28 days later by the
intranasal inoculation of 10^6 TCID₅₀ of HPIV3 or 10^6 PFU of RSV. The animals were
sacrificed five days later and the nasal turbinates and lungs were harvested and virus titers
determined (Table 19). Animals that had received the parental rB/HPIV3 virus or the G1
and F1 derivatives exhibited a high level of resistance to the replication of the HPIV3
10 challenge virus, and there were no significant differences between experimental groups.
Animals that received rB/HPIV3-G1, or rB/HPIV3-F1, or both viruses, exhibited a high
level of resistance to replication of the RSV challenge virus. The level of protective efficacy
of the rB/HPIV3-F1 virus against the RSV challenge appeared to be marginally less than that
of the rB/HPIV3-G1 virus or of the RSV control. However, this difference was not
15 significantly different. Thus, the rB/HPIV3 vector bearing either the F or G gene of RSV
induced a level of protective efficacy that was comparable to that of complete infectious
RSV.

Table 19
 Immunization of hamsters with rB/HPiV3-G1 and/or rB/HPiV3-F1 induces resistance to challenge with HPIV3 and RSV 28 days post infection.

Immunizing virus ^a	No. of Animals	Mean HPIV3 titer ^b (log ₁₀ TCID ₅₀ /g ± S.E.) ^d		Mean RSV titer ^c (log ₁₀ PFU/g ± S.E.) ^d	
		Nasal turb.	Lungs	Nasal turb.	Lungs
rB/HPiV3-G1	6	2.3±0.1 (A)	3.1±0.2 (A)	1.9±0.2 (AB)	≤ 1.7 (A)
rB/HPiV3-F1	6	2.6±0.2 (A)	3.1±0.1 (A)	2.9±0.4 (BC)	2.1±0.2 (A)
rB/HPiV3-G1 & rB/HPiV3-F1	6	2.8±0.2 (A)	2.8±0.3 (A)	1.8±0.1 (A)	1.9±0.4 (A)
rB/HPiV3	6	2.3±0.5 (A)	3.6±0.4 (A)	4.1±0.5 (C)	3.5±0.4 (B)
RSV	6	5.6±0.2 (B)	5.2±0.2 (B)	1.9±0.3 (AB)	≤ 1.7 (A)

^a Groups of 6 hamsters were inoculated intranasally with 10⁶ TCID₅₀ of the indicated PIV3 or 10⁶ PFU of RSV in a 0.1 ml inoculum.
^b HPIV3 titrations were performed on LLC-MK2 cells. The limit of detectability of virus was 10^{1.7} TCID₅₀/g tissue.
^c Quantitation of RSV was determined by plaque enumeration on HEp-2 cells. The limit of detectability of virus was 10^{1.7} PFU/g tissue.
^d Mean virus titers were assigned to similar groups (A, B, C) by the Tukey-Kramer test. Within each column, mean titers with different letters are statistically different (p<0.05). Titrers indicated with two letters are not significantly different from those indicated with either letter.

EXAMPLE XI**Use of rB/HPIV3.1 as a Vector for the Hemagglutinin HN and F Proteins of PIV2.**

The chimeric rHPIV3-1 virus, which has a HPIV3 backbone in which the HPIV3 HN and F genes have been replaced by their HPIV1 counterparts, serves as a useful vector for the HPIV2 HN protein as a supernumerary gene. This chimeric vector, rHPIV3-1.2HN, is demonstrated herein to induce resistance to replication of both HPIV1 and HPIV2 in hamsters. These findings illustrate the surprising flexibility of the PIV expression system. For example, the rHPIV3-1.2HN recombinant virus contains elements from each of the three serotypes of HPIV that cause significant disease: the internal genes of serotype 3 combined with the HN and F glycoprotein genes of serotype 1, and the HN protective antigen of serotype 2 as a supernumerary gene.

The present example provides yet another approach to deriving a PIV-based vector vaccine to protect against both PIV1 and PIV2. In this example, the rB/HPIV3 was modified by the substitution of the human PIV3 HN and F proteins by those of HPIV1. This virus, designated rB/HPIV3.1, contains the PIV1 HN and F glycoproteins as part of the vector backbone, intended to induce neutralizing antibodies and immunity to HPIV1. This virus was used in the present example as a vector to express the HN and F proteins of HPIV2 singly or together as supernumerary gene(s). Three viruses were recovered and shown to be fully viable: rB/HPIV3.1-2F; rB/HPIV3.1-2HN; or rB/HPIV3.1-2F,2HN, and each expressed the PIV2 F and/or HN gene as a supernumerary gene or genes. rB/HPIV3.1-2F,2HN, which expresses both the PIV2 F and/or HN proteins from two supernumerary genes and the PIV1 F and HN genes from the vector backbone, thus expresses both major protective antigens, i.e., the F and HN of glycoproteins, of PIV1 and PIV2 from a single virus. This approach optimizes the vaccine's protective efficacy and minimizes manufacturing costs since it accomplishes this increased immunogenicity using only one virus. It also likely will be simpler, safer and more effective to immunize infants and children with a single multivalent virus compared to a mixture of several viruses.

30 Construction of antigenomic cDNAs encoding recombinant chimeric rB/HPIV3.1 viruses bearing the HPIV2 F and HN genes as additional, supernumerary genes

A full length cDNA of the BPIV3 Kansas strain in which the F and HN glycoprotein genes of the bovine virus had been replaced with the corresponding genes of

the HPIV3 JS strain (rB/HPIV3) was constructed as previously described (Schmidt et al., *J. Virol.* 74:8922-9, 2000, incorporated herein by reference). This cDNA was modified to contain three additional unique restriction enzyme recognition sites (Figure 15). Specifically, a *BspI* site was introduced preceding the N ORF (nucleotide (nt) 103-109), an

5 *AscI* site (nt 1676-83) was introduced preceding the N gene end sequence and a *NotI* site (nt 3674-3681) was introduced preceding the P gene end sequence. Next, the F and HN glycoprotein genes of rB/HPIV3 were substituted with the corresponding genes of HPIV1. To achieve this, the sub-clone 3.1hcR6 of the previously described rHPIV3-1 full length cDNA (Tao et al., *J. Virol.* 72:2955-2961, 1998, incorporated herein by reference), which

10 contained the ORFs of the F and HN glycoprotein genes of HPIV1 under the control of HPIV3 transcription signals was modified by PCR mutagenesis to create a *SgrAI* restriction enzyme recognition site preceding the F gene and a *BsiWI* site preceding the HN gene end sequence, analogous to the position of the *SgrAI* and *BsiWI* sites that had been introduced previously into rB/HPIV3 (Schmidt et al., *J. Virol.* 74:8922-9, 2000). The mutagenic

15 forward primer used to create the *SgrAI* site was (5' to 3')

CGGCCGTGACGCGTCTCCGCACCGGTGTATTAAGCCGAAGCAA (SEQ ID NO. 34) (*SgrAI* site underlined), and the mutagenic reverse primer was (5' to 3')

CCCGAGCACGCTTTGCTCCTAAGTTTTTATATTTCCCGTACGTCTATTGTCTGAT

20 TGC (SEQ ID NO. 35) (*BsiWI* site underlined). The *SgrAI* and *BsiWI* sites were used to replace, as a single DNA fragment, the HPIV3 F and HN genes in rB/HPIV3 with the HPIV1 F and HN genes from the modified 3.1hcR6 plasmid. This yielded the full length antigenomic cDNA pB/HPIV3.1, consisting of HPIV1 F and HN open reading frames under the control of HPIV3 transcription signals in a background that is derived from BPIV3.

In the following step, the previously described HPIV2 F and HN open reading

25 frames (GenBank accession numbers AF213351 and AF213352) were modified for insertion into the *NotI* and *AscI* sites, respectively, of pB/HPIV3.1 (Figure 15). The strategy was to express each PIV2 F and HN ORF as an additional, separate mRNA, and hence it was important that it be introduced into the rB/HPIV3 genome so that it was preceded by a PIV3 gene start signal and followed by a PIV3 gene end signal. The *NotI* insertion site precedes

30 the gene end signal of the P gene (Figure 15). Hence, for insertion at this site, the HPIV2 F ORF needed to be modified by insertion of a *NotI* site and addition of a BPIV3 gene end signal, intergenic region and gene start signal at its upstream end, and a *NotI* site at its downstream end. For the HPIV2 F ORF, the forward PCR primer used was (5' to 3')

AAAAATATAGCGGCCCAAGTAAGAAAACTTAGGATTAAGGCGGATGGATCA

CCTGCATCCAATGATAGTATGCATTTTTGTTATGTACTGG (SEQ ID NO. 36) and the

the reverse primer was (5' to 3')

5 AAAATATAGCGGCCGCTTTTACTAAGATATCCCATATATGTTCCATGATTGTTCTTGGAAAAGACGGCAGG (SEQ ID NO. 37) (*NotI* site underlined, ORF translational initiation and termination triplets in bold). For the HPIV2 HN ORF, the same cis-acting elements as described above for HPIV2 F were added, but instead of *NotI*, an *AscI* site was added on either side of the insert to facilitate cloning into the N-P gene junction. The forward PCR primer used was (5' to 3')

10 GGAAAGGCGCGCCAAAGTAAGAAAACTTAGGATTAAAGGCGGATGGAAGATTACAGCAATCTATCTCTTAAATCAATTCC (SEQ ID NO. 38), the reverse primer was (5' to 3')

GGAAAGGCGCGCCAAAATTAAGCATTAGTCCCTTAAAAATGGTATTATTTGG (SEQ ID NO. 39).

15 The PCR products were digested with *NotI* (HPIV2 F insert) or *AscI* (HPIV2 HN insert) and cloned into the modified full length cDNA clone using standard molecular cloning techniques. The resulting full length cDNA containing the HPIV2 F ORF was designated pB/HPIV3.1-2F, the full length cDNA containing the HPIV2 HN ORF was designated pB/HPIV3.1-2HN, and the plasmid containing both the F and HN inserts was designated pB/HPIV3.1-2F,2HN. The nucleotide sequence of each inserted gene was confirmed by restriction enzyme digestion and automated sequencing. All constructs were designed so that the final genome nucleotide length was a multiple of six, which has been shown to be a requirement for efficient RNA replication (Calain et al., *J. Virol.*, 67:4822-30, 1993, incorporated herein by reference). The genome nucleotide length of the recovered chimeric viruses is as follows: pB/HPIV3.1: 15492; pB/HPIV3.1-2HN: 17250; pB/HPIV3.1-2F: 17190; pB/HPIV3.1-2HN,2F: 18948.

Recovery of rB/HPIV3.1, rB/HPIV3.1-2F, rB/HPIV3.1-2HN, and rB/HPIV3.1-2F,2HN chimeric viruses from cDNA

30 rB/HPIV3.1, rB/HPIV3.1-2F, rB/HPIV3.1-2HN, and rB/HPIV3.1-2F,2HN chimeric viruses were recovered from the cDNAs pB/HPIV3.1, pB/HPIV3.1-2F, pB/HPIV3.1-2HN, and pB/HPIV3.1-2F,2HN, respectively. This was accomplished by the previously-described method in which HEp-2 cells were transfected with the respective

antigenomic cDNA together with BPIV3 N, P and L support plasmids. The cells were simultaneously infected with a recombinant vaccinia virus, strain MVA, expressing the T7 RNA polymerase gene. Porcine trypsin was added to the cell culture medium to activate the HPIV1 F protein, as previously described (Tao et al., *J. Virol.* **72**:2955-2961, 1998). The recovered recombinant viruses were cloned biologically by sequential terminal dilution in Vero cells. All of the recombinant viruses replicated efficiently, induced CPE in Vero cells within 5 days and rendered the cell monolayer positive for hemadsorption. The presence of the inserted HPIV2 F and HN gene in the backbone of each recovered recombinant virus was confirmed by RT-PCR of viral RNA isolated from infected cells followed by restriction enzyme digestion and DNA sequencing. The sequence of the inserted gene and flanking regions in the recovered recombinant viruses was identical to that of the starting antigenomic cDNA.

EXAMPLE XII

15 Use of rHPIV3-1 cp45_L as a Vector For The Measles Virus Hemagglutinin (HA) protein:
Development of a Sequential Immunization Strategy

The chimeric rHPIV3-1 virus, which has a HPIV3 backbone in which the HPIV3 HN and F genes have been replaced by their HPIV1 counterparts, was shown above to serve as a useful vector for the HPIV2 HN protein as a supernumerary gene. This chimeric vector, rHPIV3-1.2HN, was able to induce resistance to replication of both HPIV1 and HPIV2 in hamsters. This finding illustrates the surprising flexibility of the PIV expression system. For example, this particular virus, rHPIV3-1.2HN, contained elements from each of the three serotypes of HPIV: the internal genes of serotype 3 combined with the HN and F glycoprotein genes of serotype 1, and the HN protective antigen of serotype 2 as a supernumerary gene. A further derivative, rHPIV3-1.2HNcp45_L, was also made that contained attenuating mutations from the cp45 HPIV3 vaccine candidate.

Thus, a PIV vector can be represented as comprising three components: the internal vector backbone genes, which can contain attenuating mutations as desired; the vector glycoprotein genes, which can be of the same or of a heterologous serotype; and one or more supernumerary genes encoding protective antigens for additional pathogens. In most cases, these supernumerary antigens are not incorporated into the virion and hence do not change the neutralization or tropism characteristics of the virus. Thus, each PIV vector is a bivalent or multivalent vaccine in which the vector itself induces immunity against an

important human pathogen and each supernumerary antigen induces immunity against an additional pathogen.

In the present example, the flexibility of the PIV vector system is further demonstrated by using the rHPIV3-1 virus, as well as its attenuated rHPIV3-1cp45_L derivative, as vectors to express measles virus HA as a supernumerary gene. This provides a new bivalent vaccine candidate for HPIV1 and measles virus. Thus, measles virus HA can be vectored by rHPIV3 and attenuated derivatives thereof, bearing the serotype 3 antigenic determinants, or by rHPIV3-1 and attenuated derivatives thereof, bearing the serotype 1 antigenic determinants.

It is noteworthy that the three serotypes of HPIV (1, 2 and 3) do not confer significant cross-protection, and that each represents a significant human pathogen for which a vaccine is needed. This raises the possibility that the three serotypes might be used to sequentially immunize the infant against the PIVs as well as vectored protective antigens against heterologous pathogens. Specifically, immunization with a PIV vector containing the antigenic determinants of one serotype should be affected minimally or not at all by prior immunization with a vector or vectors containing the antigenic determinants of a heterologous serotype. This provides the opportunity to perform sequential immunizations and boosts (preferentially at intervals of 4-6 weeks or more) against supernumerary antigens as well as against the three HPIV serotypes, whose genes can be expressed either in the vector backbone or as supernumerary genes.

The present example details the use of the techniques of reverse genetics to develop a live-attenuated HPIV1 candidate vaccine, rPIV3-1HA_{P-M} cp45_L, expressing as a supernumerary gene the major measles virus protective antigen, the HA glycoprotein (Durbin, *J. Virol.* 74:6821-31, 2000, incorporated herein by reference), for use in infants and young children to induce an immune response against both measles virus and HPIV1. Also, a sequential immunization schedule was developed in which immunization with the attenuated rHPIV3 HA_{P-M} cp45_L candidate vaccine (bearing the serotype 3 antigenic determinants) was followed by the rHPIV3-1 HA_{P-M} cp45_L candidate vaccine (bearing the serotype 1 antigenic determinants). Hamsters immunized with these viruses developed antibodies to the HPIV3 and HPIV1 antigens present in the backbone of the vectors and also maintained high titers of antibodies to the vectored antigen, the measles virus HA expressed as a supernumerary antigen from both the HPIV3 and HPIV1 candidate vaccine viruses.

Construction of rHPIV3-1 HA_(P-M) and rHPIV3-1 HA_(P-M) cp45_L, wild type and attenuated versions of rHPIV3-1 expressing measles virus HA as a supernumerary gene.

5 Two full-length plasmids were constructed, pFLC HPIV3-1 HA_(P-M) and pFLC HPIV3-1 HA_(P-M) cp45_L (Figure 16) as described above (see also, Durbin, J. Virol. 74:6821-31, 2000; Skiadopoulos et al., J. Virol. 72:1762-8, 1998; Tao et al., J. Virol. 72:2955-2961, 1998, each incorporated herein by reference). pFLC HPIV3-1 HA_(P-M) was constructed using the above-described pFLC HPIV3 HA_(P-M) in which the wild type measles virus Edmonston strain HA gene ORF was inserted as a supernumerary gene between the P and M genes of rHPIV3. pFLC HPIV3 HA_(P-M) was digested with *BspEI* to *SphI* and the cDNA fragment lacking the 6487 bp *BspEI* to *SphI* sequence was isolated. Next, pFLC 2G+hc, a full-length antigenomic cDNA plasmid bearing the F and HN ORFs of PIV1 in place of those of HPIV3 (Tao et al., J. Virol. 72:2955-2961, 1998) was digested with *BspEI* and *SphI*, and the 6541 bp fragment (plasmid nts 4830-11371) containing the HPIV1 glycoprotein genes in the HPIV3 backbone was inserted into the *BspEI* to *SphI* window of pFLC HPIV3 HA_(P-M) to give pFLC HPIV3-1 HA_(P-M) (Figure 15). The cp45 L mutations present in the L gene ORF (point mutations encoding amino acid substitutions Ser-942 to His, Leu-992 to Phe and Thr-1558 to Ile) are the major *ts* and *att* determinants of the HPIV3 cp45 candidate vaccine (Skiadopoulos et al., J. Virol. 72:1762-8, 1998) and were previously shown to confer attenuation of replication to the rHPIV3-1 cp45_L in the respiratory tract of hamsters (Tao et al., Vaccine 17:1100-8, 1999). The pFLC HPIV3-1 HA_(P-M) was then modified to encode these three *ts* mutations to yield pFLC HPIV3-1 HA_(P-M) cp45_L (Figure 16). This was accomplished by inserting the *SphI* to *NgoMIV* restriction endonuclease fragment of pFLC HPIV3 cp45L (plasmid nts 11317 -15929) (Skiadopoulos et al., J. Virol. 72:1762-8, 1998) into the *SphI* to *NgoMIV* window of pFLC HPIV3-1 HA_(P-M).

Recovery of rHPIV3-1 HA_(P-M) and rHPIV3-1 HA_(P-M) cp45_L

30 pFLC HPIV3-1 HA_(P-M) or pFLC HPIV3-1 HA_(P-M) cp45_L was transfected separately into HEp-2 cells on six-well plates (Costar, Cambridge, MA) together with the support plasmids pTM(N), pTM(P no C), and pTM(L) and LipofectACE (Life Technologies, Gaithersburg, MD) and the cells were simultaneously infected with MVA-T7, a replication-defective vaccinia virus recombinant encoding the bacteriophage T7 polymerase protein as

previously described (Skiadopoulos et al., Vaccine 18:503-10, 1999b, incorporated herein by reference). After incubation at 32°C for four days in medium containing trypsin, the transfection harvest was passaged onto LLC-MK2 cells in a 25 cm² flask, and the cells were incubated for 5 days at 32°C. The virus recovered from the cell supernatant was further

5 passaged on LLC-MK2 monolayer cultures with trypsin at 32°C to amplify the virus. rPIV3-1 HA_{P-M} and rPIV3-1 HA_{P-M} cp45_L were biologically cloned by terminal dilution on LLC-MK2 monolayer cultures at 32°C as previously described (Skiadopoulos et al., Vaccine 18:503-10, 1999b). Viral suspensions derived from biologically cloned virus were amplified on LLC-MK2 monolayer cultures.

10 Viral RNA (vRNA) was isolated from biologically cloned recombinant chimeric viruses as described above. RT-PCR was performed using rHPIV3-1 HA_{P-M} or rHPIV3-1 HA_{P-M} cp45_L vRNA as template and specific oligonucleotide primers that spanned the HA gene insert or the cp45 mutations in the L gene. The RT-PCR products were analyzed by restriction endonuclease digestion and partial DNA sequencing of the PCR

15 products as described above. This confirmed the presence of the measles virus HA gene inserted between the P and M genes of rHPIV3-1 and the presence of the cp45 L gene mutations in its attenuated derivative.

Demonstration of the attenuation phenotype of rHPIV3-1 HA_(P-M) cp45_L in hamsters

20 Golden Syrian hamsters in groups of six were inoculated intranasally with 10⁶ TCID₅₀ of rHPIV3-1, rHPIV3-1 HA_{P-M}, rHPIV3-1 cp45_L, or rHPIV3-1 HA_{P-M} cp45_L. Four days after inoculation the lungs and nasal turbinates were harvested and titers of virus were determined as described previously (Skiadopoulos et al., Vaccine 18:503-10, 1999b). The titers are expressed as mean log₁₀ TCID₅₀/gram tissue (Table 20). The recombinant rHPIV3-

25 1 HA_{P-M} and its parent rHPIV3-1 wt replicated to comparable levels, indicating that insertion of an additional transcription unit encoding the HA gene ORF did not further attenuate this virus for hamsters. The rHPIV3-1 HA_{P-M} cp45_L and its rHPIV3-1 cp45_L parent replicated to similar levels in the upper and lower respiratory tract indicating that rHPIV3-1 HA_{P-M} cp45_L was satisfactorily attenuated for replication in hamsters and that the insertion of the measles

30 virus HA gene ORF did not further attenuate the chimeric rHPIV3-1 cp45_L parent virus.

Table 20
 Replication of wild type and attenuated versions of the rPIV3-1 and rPIV3-1 HA viruses in the respiratory tract of hamsters

Virus ^a	Mean virus titer ^b (log ₁₀ TCID ₅₀ /g ± S.E.) in:	
	Nasal Turbinates	Lungs
rPIV3-1 wt	6.3 ± 0.1	6.6 ± 0.2
rPIV3-1 HA _{PM}	6.0 ± 0.1	5.7 ± 0.7
rPIV3-1 cp45 _L	4.1 ± 0.2	1.8 ± 0.2
rPIV3-1 HA _{PM} cp45 _L	4.4 ± 0.2	1.9 ± 0.2

- a. Groups of 6 hamsters each were inoculated with 10⁶ TCID₅₀ of the indicated virus intranasally.
- b. Lungs and nasal turbinates were harvested four days later. Virus present in tissue homogenates was titered by serial dilution on LLC-MK2 monolayer cultures at 32°C. Guinea pig erythrocytes were used for hemadsorption.

A sequential immunization schedule employing immunization with the attenuated rHPIV3 HA_{P-M}cp45_L chimeric vaccine candidate followed by the attenuated rHPIV3-1 HA_{P-M}cp45_L vaccine candidate induces antibodies to the HPIV3 and HPIV1 antigens of the vector backbones and induces and maintains high titers of antibodies to the shared vectored antigen, the measles virus HA.

Immunization of a group of hamsters with rHPIV3-1 HA_{P-M}cp45_L induced a strong immune response to both the HPIV1 and to the measles virus (Table 21, group 6) indicating that rHPIV3-1, like rHPIV3, can be an efficient vector for the measles virus HA.

The feasibility of sequential immunization of hamsters with rHPIV3 HA_{P-M}cp45_L and rHPIV3-1 HA_{P-M}cp45_L was next examined. Groups of hamsters were immunized with 10⁶ TCID₅₀ of rHPIV3 HA_{P-M}cp45_L (Table 21, groups 1, 2 and 3), rHPIV3 cp45_L (group 4), or L15 medium control (group 5) (Table 21). 59 days after the first immunization, groups of hamsters were immunized with 10⁶ TCID₅₀ of rHPIV3-1 HA_{P-M}cp45_L (group 1 and 4), rHPIV3-1 cp45_L (group 2 and 5), or L15 medium control (group 3). Serum samples were collected before the first immunization, 58 days after the first immunization, and 35 days after the second immunization. Animals immunized with rHPIV3 cp45_L (Table 21, group 4) developed a strong antibody response to HPIV3, and animals immunized with rHPIV3 HA_{P-M}cp45_L (groups 1, 2 and 3) developed a strong antibody response to both HPIV3 and measles virus. Animals in Group 4, which had been previously immunized with rHPIV3 cp45_L, were subsequently immunized with rHPIV3-1 HA_{P-M}cp45_L on day 59. When assayed on day 94, these animals had high titers of antibodies against HPIV3 and measles virus and a low to moderate level of antibodies to HPIV1. This showed that the HPIV3-1 chimeric vaccine virus was able to induce an immune response to both the HPIV1 antigens of the vector and to the vectored HA protein even in the presence of immunity to HPIV3, but there was some diminution of its immunogenicity in animals immune to HPIV3. The rHPIV3-1 HA_{P-M}cp45_L vaccine was clearly immunogenic in animals previously immune to HPIV3 as indicated by the response of hamsters in Group 4. These animals, which were immunized with rHPIV3 cp45_L on day 0, developed a moderately high titer of neutralizing antibodies to measles virus on day 94, 35 days following immunization with rHPIV3-1 HA_{P-M}cp45_L on day 59. Significantly, hamsters that were first immunized with rHPIV3 HA_{P-M}cp45_L and were then immunized with rHPIV3-1 HA_{P-M}cp45_L (Group 1, Table 21) achieved a higher measles virus serum neutralizing antibody titer on day 94 than groups of hamsters that were immunized with

rHPIV3 HA_{P-M} cp45_L alone (Group 3), suggesting that rHPIV3-1 HA_{P-M} cp45_L can be used to maintain high titers of serum neutralizing antibodies to measles following immunization with rHPIV3 HA_{P-M} cp45_L. Since hamsters in Group 1 developed such a high titer of antibody to the measles virus HA following first immunization with rHPIV3 HA_{P-M} cp45_L, it
5 was not possible to detect a four-fold or greater rise of these titers following immunization with rHPIV3-1 HA_{P-M} cp45_L.

In humans, it is likely that an HPIV3 vaccine such as rHPIV3 HA_{P-M} cp45_L will be given within the first four months of life followed two months later by an HPIV1 vaccine such as rHPIV3-1 HA_{P-M} cp45_L (Skiadopoulos et al., Vaccine 18:503-10, 1999b,
10 incorporated herein by reference). In contrast to rodents, human infants characteristically develop low titers of antibodies to viral glycoprotein antigens administered within the first six months of life, due to immunologic immaturity, immunosuppression by maternal antibodies, and other factors (Karron et al., Pediatr. Infect. Dis. J. 14:10-6, 1995a; Karron et al., J. Infect. Dis. 172:1445-1450, 1995b; Murphy et al., J. Clin. Microbiol. 24:894-8, 1986,
15 each incorporated herein by reference). It therefore is very likely that a boosting effect of rPIV3-1 HA_{P-M} cp45_L on the antibody titers to measles virus HA will be needed and will be readily observed in those infants immunized with rPIV3 HA_{P-M} cp45_L within the first six months of life. The present example indicates that it is possible to sequentially immunize
20 animals with two serologically distinct live attenuated PIV vaccines, each of which expresses the measles virus HA, to develop antibodies to the HPIV3 and HPIV1 antigens of the vector backbone, and to maintain high titers of antibodies to the vectored antigen, the measles virus HA.

Table 21
 Sequential immunization of hamsters with rPIV3 HA_{cp-m} cp45_L, followed by rPIV3-1 HA_{cp-m} cp45_L, induces immunity to three viruses, namely, HPIV1, HPIV3 and measles virus, and maintains the measles virus antibody titer at high levels

Group no.	Group size	Virus given in first immunization (day 0)	Immune response to first immunization		Virus given in second immunization (day 58)	Immune response to second immunization ^d		
			Serum HAI antibody titer to HPIV3 (log ₂ ± SE) ^b (day 58)	Serum antibody titer to HPIV1 ^c (log ₂ ± SE) (day 58)		Serum HAI antibody titer to HPIV3 ^b (log ₂ ± SE) (day 94)	Serum neutralizing antibody titer to HPIV1 ^c (log ₂ ± SE) (day 94)	Serum antibody titer to measles virus ^d (60% PRN, log ₂ ± SE) (day 94)
1	8	rPIV3 HA _{cp-m} cp45 _L	10.8±0.4	≤0.5±0.0	rPIV3-1 HA _{cp-m} cp45 _L	11.5±0.5	0.9±0.2	13.1±0.3
2	8	rPIV3 HA _{cp-m} cp45 _L	10.9±0.4	≤0.5±0.0	rPIV3-1 cp45 _L	10.5±0.5	1.2±0.3	12.8±0.4
3	6	rPIV3 HA _{cp-m} cp45 _L	9.3±0.3	≤0.5±0.0	none	9.6±0.9	1.1±0.4	12.3±0.2
4	8	rPIV3 cp45 _L	9.6±0.6	≤0.5±0.0	rPIV3-1 HA _{cp-m} cp45 _L	9.0±0.7	0.9±0.3	7.3±0.3
5	6	none	< 2±0.0	≤0.5±0.0	rPIV3-1 cp45 _L	< 2±0.0	4.8±0.6	< 3.3
6	8	rPIV3-1 HA _{cp-m} cp45 _L		3.0±0.4				

a. Sera were collected 5 days before and 58 days after the first immunization. The second immunization was given 59 days after the first, and serum was collected again 35 days later (day 94).
 b. Mean serum PIV3 HAI antibody titer is expressed as the reciprocal mean log₂ ± standard error, SE.
 c. Mean serum neutralizing antibody titer to HPIV1 is expressed as the reciprocal mean log₂ ± S.E.
 d. Mean serum neutralizing antibody titer to wild type measles virus is expressed as the reciprocal mean log₂ ± standard error, PRN, plaque reduction neutralization.

EXAMPLE XIIIConstruction and Characterization of Chimeric HPIV3-2 Vaccine Recombinants Expressing Chimeric Glycoproteins

The present example details development of a live attenuated PIV2 candidate vaccine virus for use in infants and young children using reverse genetic techniques. Preliminary efforts to recover recombinant chimeric PIV3-PIV2 virus carrying full-length PIV2 glycoproteins in a wild type PIV3 backbone, as described above for HPIV3-1 chimeric constructs, did not yield infectious virus. However, viable PIV2-PIV3 chimeric viruses were recovered when chimeric HN and F ORFs rather than full-length PIV2 ORFs were used to construct the full-length cDNA. The recovered viruses, designated rPIV3-2CT in which the PIV2 ectodomain and transmembrane domain was fused to the PIV3 cytoplasmic domain and rPIV3-2TM in which the PIV2 ectodomain was fused to the PIV3 transmembrane and cytoplasmic tail domain, possessed similar, although not identical, *in vitro* and *in vivo* phenotypes. Thus, it appears that only the cytoplasmic tail of the HN or F glycoprotein of PIV3 is required for successful recovery of PIV2-PIV3 chimeric viruses.

The rPIV3-2 recombinant chimeric viruses exhibit a strong host range phenotype, i.e. they replicate efficiently *in vitro* but are strongly restricted in replication *in vivo*. This attenuation *in vivo* occurs in the absence of any added mutations from *cp45*. Although rPIV3-2CT and rPIV3-2TM replicated efficiently *in vitro*, they were highly attenuated in both the upper and the lower respiratory tract of hamsters and African green monkeys (AGMs), indicating that chimerization of the HN and F proteins of PIV2 and PIV3 itself specified an attenuation phenotype *in vivo*. A phenotype including efficient replication *in vitro* and highly restricted growth *in vivo* is greatly desired for vaccine candidates. Despite this attenuation, they were highly immunogenic and protective against challenge with PIV2 wild type virus in both species. rPIV3-2CT and rPIV3-2TM were further modified by the introduction of the 12 PIV3 *cp45* mutations located outside of the HN and F coding sequences to derive rPIV3-2CT_{*cp45*} and rPIV3-2TM_{*cp45*}. These derivatives replicated efficiently *in vitro* but were even further attenuated in hamsters and AGMs indicating that the attenuation specified by the glycoprotein chimerization and by the *cp45* mutations was additive. These findings identify the rPIV3-2CT and rPIV3-2TM recombinants as preferred candidates for use in live attenuated PIV2 vaccines.

Viruses and cells

The wild type PIV1 strain used in this study, PIV1/Washington/20993/1964 (PIV1/Wash64) (Murphy et al., Infect. Immun. 12:62-68, 1975, incorporated herein by reference), was propagated in LLC-MK2 cells (ATCC CCL 7.1) as previously described (Tao et al., J. Virol. 72:2955-2961, 1998, incorporated herein by reference). The PIV wild type virus, strain V9412-6, designated PIV2/V94, was isolated in qualified Vero cells from a nasal wash of a sick child in 1994. PIV2/V94 was plaque purified three times on Vero cells before being amplified twice on Vero cells using OptiMEM without FBS. The wild type cDNA-derived recombinant PIV3/JS strain (rPIV3/JS) was propagated as previously described (Durbin et al., Virology 235:323-332, 1997, incorporated herein by reference). The modified vaccinia Ankara virus (MVA) recombinant that expresses the bacteriophage T7 RNA polymerase was generously provided by Drs. L. Wyatt and B. Moss (Wyatt et al., Virology 210:202-205, 1995, incorporated herein by reference).

HEp-2 cells (ATCC CCL 23) were maintained in MEM (Life Technologies, Gaithersburg, MD) with 10% fetal bovine serum, 50 µg/ml gentamicin sulfate, and 2 mM glutamine. Vero cells below passage 150 were maintained in serum-free medium VP-SFM (Formula No. 96-0353SA, Life Technologies) with 50 µg/ml gentamicin sulfate and 2 mM glutamine.

Virion RNA isolation, reverse transcription and PCR amplification of viral genes, and automated sequencing

To clone viral genes or to verify genetic markers of recombinant chimeric viruses, viruses were amplified on cultured cells and concentrated by polyethylene glycol precipitation as previously described (Mbiguino et al., J. Virol. Methods 31:161-170, 1991, incorporated herein by reference). Virion RNA was extracted from the virus pellet using Trizol reagent (Life Technologies) and used as template for reverse transcription (RT) with the Superscript Preamplification system (Life Technologies). The cDNA was further PCR amplified using the Advantage cDNA kit (Clontech, Palo Alto, CA). For cloning or sequencing purposes, the RT-PCR amplified DNA was purified from agarose gels using NA45 DEAE membrane as suggested by the manufacturer (Schleicher & Schuell, Keene, NH). Sequencing was performed with the dRhodamine dye terminator cycling sequencing kit (Perkin Elmer, Forster City, CA) and an ABI 310 Gene Analyzer (Perkin Elmer, Forster City, CA).

Construction of the chimeric PIV3-PIV2 antigenomic cDNAs encoding the complete PIV2 F and HN proteins or chimeric F and HN proteins containing a PIV2-derived ectodomain and PIV3-derived cytoplasmic tail domain

A DNA encoding a full-length PIV3 antigenomic RNA was constructed in
5 which the PIV3 F and HN ORFs were replaced by their PIV2 counterparts following the
strategy described previously (Tao et al., *J. Virol.* 72:2955-2961, 1998) for PIV3-PIV1.
Details of this construction are presented in Figure 17. PIV2/V94 propagated in Vero cells
was concentrated and virion RNA (vRNA) was extracted from the virus pellet using Trizol
reagent. The F and HN ORFs of PIV2/V94 were reverse transcribed from vRNA using
10 random hexamer primers and the SuperScript Preamplification System before being
amplified by PCR using the cDNA Advantage kit and primer pairs specific to PIV2 F and
HN genes, respectively (1, 2 and 3, 4; Table 22). The amplified cDNA fragment of PIV2 F
ORF was digested with NcoI plus BamHI and ligated into the NcoI-BamHI window of
pLit.PIV31.Fhc (Tao et al., *J. Virol.* 72:2955-2961, 1998, incorporated herein by reference)
15 to generate pLit.PIV32Fhc. The BspEI site in the PIV3 full-length cDNA is unique and we
planned to use it to exchange segments between cDNAs (see Figures 17-19). Therefore, a
BspEI site that was found in the PIV2 F ORF was removed by site-directed mutagenesis
without affecting the amino acid sequence. The cDNA fragment of PIV2 HN ORF was
digested with NcoI plus HindIII and ligated into the NcoI-HindIII window of
20 pLit.PIV31.HNhc (Tao et al., *J. Virol.* 72:2955-2961, 1998) to generate pLit.PIV32HNhc.
The PIV2 ORFs in pLit.PIV32Fhc and pLit.PIV32HNhc were sequenced, and the sequence
was found to be as designed. The nucleotide sequences for the PIV2 F and HN ORFs are
submitted in the GenBank. pLit.PIV32Fhc and pLit.PIV32HNhc were each digested with
PpuMI plus SpeI and assembled to generate pLit.PIV32hc. The 4 kb BspEI-SpeI fragment
25 of pLit.PIV32hc was introduced into the BspEI-SpeI window of p38' Δ PIV31hc
(Skiadopoulos et al., *Vaccine* 18:503-510, 1999, incorporated herein by reference) to
generate p38' Δ PIV32hc. The 6.5 kb fragment, generated by BspEI and SphI digestion of
p38' Δ PIV32hc, containing the PIV2 full-length F and HN ORFs was introduced into the
BspEI-SphI window of pFLC.2G+.hc (Tao et al., *J. Virol.* 72:2955-2961, 1998) to generate
30 pFLC.PIV32hc (Figure 17; Table 23 = SEQ ID NO. 60).

Table 22. Primers used in construction of PIV3-2 full-length chimeric antigenomic cDNAs

Primer No.	Gene	Direction	Position		Used in the construction or characterization of:	Sequence ^a
			Beginning	End		
1	PIV2 F	sense	PIV2 F start codon 5070 ^b	20 bp down stream 5091	pFLC.PIV32hc	ggaccATGgATCACCCTGCATCCAAT (SEQ ID NO. 40)
2	PIV2 F	antisense	PIV2 F stop codon 6732 ^b	20 bp upstream 6705 ^b	pFLC.PIV32hc	tgfggggGTAAAGATA TCCCATATATATGTTTC (SEQ ID NO. 41)
3	PIV2 HN	sense	PIV2 HN start codon 6837 ^b	18 bp down stream 6856 ^b	pFLC.PIV32hc	ggggcATGGAAAGATTACAGCAAAT (SEQ ID NO. 19)
4	PIV2 HN	antisense	PIV2 HN stop codon 8538 ^b	17 bp upstream 8538 ^b	pFLC.PIV32hc	caatbaagcTTAAAGCATTAGTTCCCC (SEQ ID NO. 20)
5	PIV2 F	sense	5069 ^c	5088 ^c	pFLC.PIV32TM	ATGCATCACCTGCATCCAAT (SEQ ID NO. 42)
6	PIV2 F	antisense	6538 ^c	6517 ^c	pFLC.PIV32TM	TAGTGAATAAAGTGTCTTGCT (SEQ ID NO. 43)
7	PIV2 HN	sense	6962 ^c	6985 ^c	pFLC.PIV32TM	CATGAGATAAATTCATCTTGTGATGTT (SEQ ID NO. 44)
8	PIV2 HN	antisense	8560 ^c	8537 ^c	pFLC.PIV32TM	ggcTTAAAGCATTAGTTCCCTTAA (SEQ ID NO. 45)
9	PIV3 F	sense	6539 ^c	6566 ^c	pFLC.PIV32TM	ATCATAATTAITTTTGATAATGATCAITTA (SEQ ID NO. 46)
10	PIV3 F	antisense	5068 ^c	5050 ^c	pFLC.PIV32TM	GTTCAAGTGTGTTGTTGTT (SEQ ID NO. 47)
11	PIV3 HN	sense	8561 ^c	8587 ^c	pFLC.PIV32TM	TCATAATTAACCATAATATGTCATCAAT (SEQ ID NO. 48)

12	PIV3 HN	antisense	6961 ^e	6938 ^e	pFLC.PIV32FM	GATGGAATTAATTAGCACTATGAT (SEQ ID NO. 49)
13	PIV2 F	sense	5069 ^d	5088 ^d	pFLC.PIV32CT	ATGCATCACCTGCATCCAAT (SEQ ID NO. 50)
14	PIV2 F	antisense	6607 ^d	6589 ^d	pFLC.PIV32CT	GATGATGTAGGCAATCAGC (SEQ ID NO. 51)
15	PIV2 HN	sense	6887 ^d	6904 ^d	pFLC.PIV32CT	ACTGCCACAATTCCTGGC (SEQ ID NO. 52)
16	PIV2 HN	antisense	8536 ^d	8511 ^d	pFLC.PIV32CT	TTAAAGCAITTAGTCCCTTAAAAATG (SEQ ID NO. 53)
17	PIV3 F	sense	6620 ^d	6642 ^d	pFLC.PIV32CT	AAGTATTACAGAATTCAAAAGAG (SEQ ID NO. 54)
18	PIV3 F	antisense	5068 ^d	5050 ^d	pFLC.PIV32CT	GTTACGTGCTTGTGTGTT (SEQ ID NO. 47)
19	PIV3 HN	sense	8525 ^d	8551 ^d	pFLC.PIV32CT	TCATAATTAAACCATAATATGCATCAAT (SEQ ID NO. 48)
20	PIV3 HN	antisense	6898 ^d	6879 ^d	pFLC.PIV32CT	CTTATTAGTGAGCTTGTTC (SEQ ID NO. 55)
21	PIV2 F	Sense	6608 ^{e,d}	6630 ^{e,d}	Chimera confirmation	ACCGCAGCTGTAGCAATAGT (SEQ ID NO. 56)
22	PIV2 HN	antisense	7522 ^e 7501 ^d	7502 ^e 7481 ^d	Chimera confirmation	GATCCATCACTTAGGTAAT (SEQ ID NO. 57)
23	PIV3 M	sense	4759 ^{e,d}	4780 ^{e,d}	Chimera confirmation	GATACTATCCTAATAATTATTGG (SEQ ID NO. 58)
24	PIV3 L	antisense	9100 ^e 9076 ^d	9081 ^e 9057 ^d	Chimera confirmation	GCTAAATTTGATAGCACATT (SEQ ID NO. 59)

- * All the primers are annotated in that the PIV specific sequences are in uppercase, non-PIV sequences in lowercase, start and stop codons in bold, and restriction sites underlined.
- ^b The numbers are the nt positions in the full-length antigenomic cDNA construct pFLC:PIV32hc.
- ^c The numbers are the nt positions in the full-length antigenomic cDNA construct pFLC:PIV32TM and pFLC:PIV32TMcp45.
- ^d The numbers are the nt positions in the full-length antigenomic cDNA construct pFLC:PIV32CT and pFLC:PIV32CTcp45.

TABLE 23 (SEQ ID NO. 60)

Sequence of pFLC.PIV32, 15492 bp in sense orientation
(only the insert is shown)

```

1 ACCAACCAAG AGAAGAACT TGTCTGGGAA TATAAATTTA ACTTTAAATT AACTTAGGAT
61 TAAAGACATT GACTAGAAGG TCAAGAAAAG GGAACCTCTAT AATTTCAAAA ATGTTGAGCC
121 TATTTGATAC ATTTAATGCA CGTAGGCCAAG AAAACATAAC AAAATCAGCC GGTGGAGCTA
181 TCATTCTCTG ACAGAAAAAT ACTGTCTCTA TAITCGCCCT TGGACCGACA ATAACGTATG
241 ATAATGAGAA AATGACATTA GCTCTTCTAT TTCTATCTCA TTCACTAGAT AATGAGAAAC
301 AACATGCACA AAGGGCAGGG TTCTTGGTGT CTTTATTGTC AATGGCTTAT GCCAATCCAG
361 AGCTCTACCT AACAAACAAT GGAAGTAATG CAGATGTCAA GTATGTCTAT TACATGATGT
421 AGAAGATCTT AAAACGGCAA AAGTATGGAG GATTTGTGGT TAAGACGAGA GAGATGATAT
481 ATGAAAAGAC AACTGATTGG ATATTTGGAA GTGACCTGGA TTATGATCAG GAAACTATGT
541 TGCAGAACGG CAGGAACAAT TCAACAATTT AAGACCTTGT CCACACATTT GGGTATCCAT
601 CATGTTTAGG AGCTTTTATA ATACAGATCT GGATAGTTCT GGTCAAAGCT ATCACTAGTA
661 TCTCAGGGTT AAGAAAAGCC TTTTTCACCC GATTGGGAAGC TTTGAGACAA GATGGAACAG
721 TGCAGGCAGG GCTGGTATTT ACAGGTGACA CAGTGGATCA GATTGGGTCA ATCATGCCGT
781 CTCACAGCAG CTGGTAACT CTTATGGTTG AACAATTAAT AACATGAAT ACCAGCAGAA
841 ATGACCTCAC AACCATAGAA AAGAATATAC AAATTGTGGC CAACTACATA AGAGATGCAG
901 GTCCTGCTTC ATTCTTCAAT ACAATCAGAT ATGGAATTGA GACCAGAATG GCAGCTTTGA
961 CTCTATCCAC TCTCAGACCA GATATCAATA GATTAAGAGC TTTGATGGAA CTGTATTTAT
1021 CAAAGGGACC ACGCGCTCCT TTCATCTGTA TCCTCAGAGA TCCTATACAT GGTGAGTTCC
1081 CACCAGGCAA CTATCTGCC ATATGGAGCT ATGCAATGGG GGTGGCAGTT GTACAAAAATA
1141 GAGCCATGCA ACAGTATGTT ACAGGGAAGT CATATCTAGA CATTGATATG TTCAGCTAG
1201 GACAGCAGCT AGCAGCTGAT GCCGAAGCTC AATAGAGCTC AACACTGAAA GATGAACCTG
1261 GAGTGACACA CGAATCTAAA GAAAGCTTGA AGAGACATAT AAGGAACATA AACAGTTTCA
1321 AGACATCTTT CCACAAACCG ACAGGTGGAT CAGCCATAGA GATGGCAATA GATGAAGAGC
1381 CAGAACAATT CGAACATAGA GCAGATCAAG AACAAAAATG AGAACCTCAA TCATCCATAA
1441 TTCAATATGC CTGGGCAGAA GGAATAGAAA CGCATGATCA GACTGAGCAA GCTACAGAAAT
1501 CTGACAAATAT CAAGACCGAA CAACAAAAAC TCAGAGCAGC ACTAAACAAG AGACTCAACG
1561 ACAAGAGAAA ACAAGCAGT CAACCAACCA CTAACTCCAC AAACAGAAAC AACCCAGCAG
1621 AAATAGATGA TCTGTTTAA CATTGTTGAA GCAACTAATC GAATCAACAT TTTAATCTAA
1681 ATCAATAATA AATAAGAAA ACTTAGGATT AAAGAATCCT ATCATACCGG AATATAGGGT
1741 GGTAAATTTA GAGTCTGCTT GAAACTCAAT CAATAGAGAG TTGATGGAAA GCGATGCTAA
1801 AAATATCAA ATCATGGATT CTTGGGAAGA GGAATCAAGA GATAAATCAA CTAATATCTC
1861 CTCGGCCCTC AACATCATTG AATTCTACT CAGCACCGAC CCCCAGAAGG ACTTATCGGA
1921 AAACGACACA ATCAACACAA GAACCCAGCA ACTCAGTGCC ACCATCTGTC AACCGAATA
1981 CAAACCAACA GAAACAAGTG AGAAGATAG TGGATCAACT GACAAAAATA GACAGTCCGG
2041 GTCTACACAC GAATGTACAA CAGAGCAAA AGATAGAAAT ATTGATCAGG AAATCTGACA
2101 GAGAGGACCT GGGAGAAGAA GCAGCTCAGA TAGTAGAGCT GAGACTGTGG TCTCTGGAGG
2161 AATCCCCAGA AGCATCACAG ATTTAAAAA TGGAAACCAA AACACGGAGG ATATTGATCT
2221 CAATGAAATT AGAAAGATGG ATAAGGACTC TATTGAGGGG AAATGCGAC AATCTGCAAA
2281 TGTTCCAAGC GAGATATCAG GAAGTATGA CATATTTACA ACAGAACAAA GTAGAAACAG
2341 TGATCATGGA AGAAGCCTGG AATCTATCAG TACACCTGAT ACAAGATCAA TAAGTGTGTT
2401 TACTGCTGCA ACACCAAGTG ATGAGAAGA AATACTAATG AAAAATAGTA GGACAAAGAA
2461 AAGTCTTCCA ACACATCAAG AAGATGACAA AAGAATTTAA AAAGGGGAA AAGGAAAGA
2521 CTGTTTAAAG AAATCAAAAG ATACCGACAA CCAGATACCA ACATCAGACT ACAGATCCAC
2581 ATCAAAAGGG CAGAAGAAA TCTCAAGAC AACAAACACC AACACCGACA CAAAGGGGGA
2641 AACAGAAATA CAGACAGAAAT CATCAGAAAC ACAATCTTCA TCATGGAATC TCATCATCGA
2701 CAACAACACC GACCGGAACG AACRGAACAG CACAACTCCT CCAACACAA CTTCCAGATC
2761 AACTTATACA AARGAATCGA TCCGAACAAA CTCTGAATCC AAACCAAGA CACAAAGAC
2821 AATGGGAAGG GAAAGGAAGG ATCAGAAAGA GAGCAATCGA TTTACAGAGA GGGCAATTAC
2881 TCTATTGCGA AATCTTGGTG TAATTCAATC CACATCAAAA CTAGATTTAT ATCAAGACAA
2941 ACGAGTTGTA TGTGTAGCAA ATGTAATAAA CAATGTAGAT ACTGCATCAA AGATAGATTT
3001 CCTGGCAGGA TTAGTCATAG GGGTTCAAT GGACAAACGAC ACAAATTTAA CACAGATACA
3061 AAATGAAATG CTAACCTCA AAGCAGATCT AAAGAAAATG GACCAATCAC ATAGAAATTT
3121 GATAGAAAAT CAAAGAGAAC AACTGTCTAT GATCAGTCA CTAATTTCAA ATCTCAAAAT
3181 TATGACTGAG AGAGGAGGAA AGAAGATCAA AAATGAATCC AATGAGAGAG TATCATGAT
3241 CAAACAAAAA TTGAAGAGG AAAAGATCAA GAAGACCCAG TTTGACCCAC TTAGGAGGC
3301 ACAAGGCATT GACAAAGATA TACCCGATCT ATATCGACAT GCAGGAGATA CACTAGAGAA
3361 CGATGTACAA GTTAATCAG AGATATTAAG TTCAATCAAT GAGTCAAATG CAACAAGACT

```

3421 AATACCCAAA AAAGTGAGCA GTACAATGAG ATCACTAGTT GCAGTCATCA ACAACAGCAA
 3481 TCTCTCACAA AGCACAAAAC AATCATACAT AAACGAACTC AAACGTTGCA AAAATGATGA
 3541 AGAAGTATCT GAATTAATGG ACATGTTCAA TGAAGATGTC AACCAATTGCC AATGATCCAA
 3601 CAAAGAAACG ACACCGAACA AACAGACAAG AAACAACAGT AGATCAAAC CTGTCAACAC
 3661 ACACAAAATC AAGCAGAATG AAACAACAGA TATCAATCAA TATACAAATA AGAAAAACTT
 3721 AGGATTAAG AATAAATTAA TCCTTGTCCTA AAATGAGTAT AACTAACTCT GCAATATACA
 3781 CATTCCCAGA ATCATCATTG TCTGAAAATG GTCATATAGA ACCATTACCA CTCAPAGTCA
 3841 ATGAACAGAG GAAAGCAGTA CCCCACATTA GAGTTGCCAA GATCGGAAT CCACAAAAC
 3901 ACGGATCCCG GTATTAGATG GTCTTCTTAC TCGGCTTCTT CGAGATGGAA CGAATCAAAG
 3961 ACAAATACGG GAGTGTGAAT GATCTCGACA GTGACCCGAG TTACAAGTT TGTGGCTCTG
 4021 GATCATTACC AATCGGATTG GCTAAGTACA CTGGGAATGA CCGAAGATTG TTACAAGCCG
 4081 CAACCAACT GGATATAGAA GTGAGAAGAA CAGTCAAGC GAAAGAGATG GTTGTTTACA
 4141 CGGTACAAA TATAAAACCA GAACGTACC CATGGTCCAA TAGACTAAGA AAAGGAATGC
 4201 TGTCGATGC CAACAAGTT GCTTGTCTC CTCATGTCT TCCACTAGAT AGGAGCATAA
 4261 AATTTAGAGT AATCTCGTG AATTGTACGG CAATTGGATC AATAACCTTG TTCAAAATTC
 4321 CTAGTCAAT GGCATCAGTA TCTTACCCA ACACAATATC AATCAACTCG CAGGTACACA
 4381 TAAAAACAGG GGTTACAGCT GATTCTAAG GGTAGTTCA AATTTGGAT GAGAAAGCCG
 4441 AAAATCACT GAATTTCAAG GTCCATCTCG GATTGATCAA AAGAAAAGTA GGCAGAATGT
 4501 ACTCTGTTGA ATACTGTAAA CAGAAAATCG AGAAAATGAG ATTGATATT TCTTTAGGAC
 4561 TAGTTGGAGG AATCAGTCTT CATGTCAATG CAACTGGGTC CATATCAAAA ACATAGCAA
 4621 GTCAGCTGGT ATTCAAAGA GAGATTTGTT ATCCCTTAAT GGATCAAT CCGCATCTCA
 4681 ATCTAGTTAT CTGGGCTTCA TCAGTAGAGA TTACAAGAGT GGATCAAT TTCCAACCTT
 4741 CTTTACCTGG CGAGTTCAGA TACTATCTTA ATATTATTGC AAAGAGGATT GGGAAATCA
 4801 ACAATGGAA CTAGTAATCT CTATTTTAGT CCGGACGAT CTATTAAGCC GAAGCAATA
 4861 AAGGATAACT AAAAECTTAG GACAAAAGAG GTCATACCA ACAACTATTA CGACTCACAC
 4921 TCACAAGAA AAGAGAGAAG GGACCAAAA AGTCAAATAG GAGAAATCAA AACAAAAGGT
 4981 ACAGAACACC AGAACACAAA AATCAAACA TCCAACCTAC TCAAAACAAA AATTCAAAA
 5041 GAGACCCGCA ACACAACAAG CACTGAACAC CATGGATCAC CTGCATCAA TGATAGTATG
 5101 CATTTTGTGT ATGTACACTG GAATTTGAGT TTCAGATGCC ATTGCTGGAG ATCAACTCTT
 5161 CAATGTAGGG GTCATTCAAT CAAAGATAAG ATCACTCATG TACTACACTG ATGGTGGCCG
 5221 TAGCTTATT GTTGTAAAT TACTACCAA TCTTCCCCCA AGCAATGGAA CATGCAACAT
 5281 CACCACTCTA GATGCATATA ATGTTACCTT ATTTAAGTGT CTAACACCCG TGATGAGAA
 5341 CTTGAGCAAA ATTTCTGCTG TTACAGATAC CAAACCCCGC CGAGAACGAT TTGAGGAGT
 5401 CGTTATTGGG CTTGCTGCAC TAGGAGTAGC TACAGCTGCA CAAATAACCG CAGCTGTAGC
 5461 AATAGTAAAA GCCAATGCAA ATGCTGTGTC GATAAACAAT CTGTCACTT CAATTCATC
 5521 CACCAACAAG CCAGTATCCG ATGTGATAC TGCAATCAAGA ACAATTGCAA CCGCATTTCA
 5581 AGCGATTCCG GATCACATCA ATGGASCCAT TGTCAACGGG ATAACATCTG CATCATGCCG
 5641 TGCCCATGAT GCACATAATG GGTCAATATT AAATTTGTAT CTCACTGAGC TTACTACAT
 5701 ATTTATAAAT CAAATAACAA ACCCTGCGCT GACACCACCT TCCATCCAAG CTTTAAAGAT
 5761 CCTCCTCGGT AGCACCTTGC CAATTTGCTAT TGAATCCAAA CTCACACAAA AACTCAACAC
 5821 AGCAGAGCTG CTCAGTAGCG GACTGTTAAC TGGTCAAATA ATTTCCATTT CCCCAGTGA
 5881 CATGCAATG CTAATTCAAA TCAATGTTCC GACATTATA ATGCAACCCG GTGCGAAGGT
 5941 AATTGACTTA ATTGCTATCT CTGCABACCA TAAATTACAA GAGTAGTTG TACAACTTCC
 6001 TAATAGAATT CTAGAATATG CAATGAACCT ACAAACTAC CCAAGCCATG ATTGTTTCCG
 6061 GACACCAAC TCTGATTTT GTAGATACAA TGAGGGTTCC CCGATCCCTG AATCACAATA
 6121 TCAATGCTTA AGGGGGAATC TTAATTTCTG CACTTTTACC CCTATTATCG GGAACCTTCT
 6181 CAAGCGATTG GCATTTGCCA ATGGTGTGCT CTATGCCAAC TGCAAACTTT TGCTATGTAA
 6241 GTGTCCGAC CCTCCCATG TTGTGTCTCA AGATGACAAC CAAGGCATCA GCATAATTGA
 6301 TATTAGAGG TGCTCTGAGA TGATGCTTGA CACTTTTICA TTTAGGATCA CATCTACATT
 6361 CAATGCTACA TACGTGACAG ACTTCTCAAT GATTAATGCA AATATTGTAC ATCTAAGTCC
 6421 TCTAGACTTG TCAAAACAAA TCAATTCAT AAACAATCT CTTAAAAGTG CTGAGGATTG
 6481 GATTGCAGAT AGCAACTTCT TCGCTAATCA AGCCAGAACA GCCAAGACAC TTTATTCACT
 6541 AAGTGCAATC GCATTAATAC TATCAGTAT TACTTTGGTT GTTGTGGGAT TGCTGATTGC
 6601 CTACATCATC AAGCTGGTTT CTCAAATCCA TCAATTCAGA GCACAGCTG CTACAACAT
 6661 GTTCCACAGG GAGAACTCTG CCGTCTTTC CAAGAACAAT CATGGAAACA TATATGGGAT
 6721 ATCTTAGGAT CCCTACAGAT CATTAGATAT TAAATATATA AAAAECTTAG GAGTAAAGTT
 6781 ACGCAATCCA ACTTACTCA TATAATTGAG GAAGGACCCA ATAGACAAAT CCAATCCAT
 6841 GGAAGATTAC AGCAATCTAT CTCTTAAATC AATTCCTAAA AGGACATGTA GAATCATTTT
 6901 CCGAACTGCC ACAATCTTGT GCATATGCAC ATTAATTGTG CTATGTTCAA GTATTCTTCA
 6961 TGAGATAATT CATCTTGATG TTTCTCTG TCTTATGAAT TCTGATGAGT CACAGCAAGG
 7021 CATTATTCAG CCTATCATAG AATCATTAA ATCATTGAT GCTTTGGCCA ACCAGATTCT
 7081 ATATAAGTGT GCAATAGTAA TTCCCTTAA AATTGACAGT ATCGAACTG TAATACTCTC

7141 TGCTTAAAA GATATGCACA CCGGGAGTAT GTCCAATGCC AACTGCACGC CAGGAAATCT
7201 GCTTCTGCAT GATGCAGCAT ACATCAATGG AATAAACAAA TTCCTTGTAC TTGAATCATA
7261 CAATGGGACG CCTAAATATG GACCTCTCCT AAATATACCC AGCTTTATCC CCTCAGCAAC
7321 ATCTCCCAT GGGTGTACTA GAATACCATC ATTTTCACTC ATCAAGACCC ATTGGTGTTA
7381 CACTCACAAT GTAATGCTTG GAGATTGTCT TGATTTCAAG GCATCTAACC AGTATTTATC
7441 AATGGGGATA ATACAACAAT CTGCTGCAGG GTTCCCAATT TTCAGGACTA TGAAAACCAT
7501 TTACCTAAGT GATGGAATCA ATCGCAAAAG CTGTTCAGTC ACTGCTATAC CAGGAGGTTG
7561 TGCTTTGTAT TGCTATGTAG CTACAAGGTC TGAAAAAGAA GATTATGCCA CGACTGATCT
7621 AGCTGAACAG AGACTTGCTT TCTATTATTA TAATGATACC TTTATTGAAA GAGTCATATC
7681 TCTTCCAAAT ACAACAGGGC AGTGGGCCAC AATCAACCCT GCAGTCGGAA GCGGGATCTA
7741 TCATCTAGGC TTTATCTTAT TTCTGTATA TGGTGGTCTC ATAARTGGGA CTACTTCTTA
7801 CAATGAGCAG TCCTCAGCCT ATTTTATCCC AAAACATCCC AACATAACTT GTGCCGGTAA
7861 CTCCAGCAAA CAGGCTGCAA TAGCACGGAG TTCCTATGTC ATCCGTATC ACTCAACAG
7921 GTTAAATCAG AGTGCTGTTC TTATTTGTCC ATTTGCTGAC ATGCATACAG AAGAGTGTAA
7981 TCTAGTTATG TTTAAACAAT CCCAAGTCAT GATGGGTGCA GAAGGTAGGC TCTATGTAT
8041 TGGTAATAAT TTGTATTAT ATCAACGCAG TTCCTTTGG TGGTCTGCAT CGCTCTTTTA
8101 CAGGATCAAT ACAGATTTTT CTAAGGAAT TCCTCCGATC ATTGAGGCTC AATGGGTACC
8161 TCCTATCAAT GTTCTCGTC CTGGAGTCAT GCCATGCAAT GCAACAAGTT TTTGCCCTGC
8221 TAATTTGATC ACAGGGGTGT ACGCAGATGT GTGCCCGCTT AATGATCCAG AACTCATGTC
8281 ACGTAAATGCT CTGAACCCCA ACTATCGATT TGCTGGAGCC TTTCTAAAA ATGAGTCCAA
8341 CCGAAGTATC CCCACATCT ACATGTCATC GGCTAACTCC CTCTTAAAA CTACCCGGAT
8401 CAACACACCC AATCACAARAG CAGCATATAC ATCTTCAACC TGCTTAAAA ACAGTGGAAC
8461 CCAAAAAAAT TATTGTTTAA TAATAATGA AATGGCTCA TCTCTTTAG GGGAGTCCA
8521 AATAATACCA TTTTAAAGG AACTAATGCT TTAAGCTTAA TTAACCATAA TATGCATCAA
8581 TCTTCTATA ATACAAGTAT ATGATAAGTA ATCTGCAATC AGACAATAGA CAAAAGGGAA
8641 ATATAAAAAA CTTAGGACCA AAGCGTGCTC GGGAAATGGA CACTGAATCT AACAATGGCA
8701 CTGTATCTGA CATACTCTAT CCTGAGTGTG ACCTTAACTC TCCTATCGTT AAAGGTAAAA
8761 TAGCACAAAT ACACACTATT ATGAGTCTAC CTCAGCCTTA TGATATGGAT CAGGACTCAA
8821 TACTAGTTAT CACTAGACAG AAAATAAAC TTAATAAAT GGATAAAGA CACGATCTA
8881 TTAGAAGATT AAAATTAATA TTAACGTAAA AAGTGAATGA CTTAGGAAA TACACATTTA
8941 TCAGATATCC AGAAATGCA AAAGAAATGT TCAAAATATA TATACCTGGT ATTAACAGTA
9001 AAGTACTGAT ATTATTACTT AAAGCAGATA GAACATATAG TCAAAATGACT GATGATTTAA
9061 GAGATCTATG GATTAATGTG CTATCAAAAT TAGCCTCAA AAATGATGGA AGCAATATG
9121 ATCTTAATGA AGAAATTAAT AATATATCGA AAGTTCACAC AACCTATAAA TCAGATAAAT
9181 GGTATAATCC ATTCAAAACA TGGTTTACTA TCAAGTATGA TATGAGAAGA TTACAAAAG
9241 CTCGAAATGA GATCACTTTT AATGTTGGGA AGGATTTAA CTTGTTAGAA CACCAAGA
9301 ATTTCTTATT GATACATCCA GAATGGTTT TGATATTAGA TAAACAAAAC TATAATGGTT
9361 ATCTAATFAC TCCTGAATTA GTATTGATGT ATTTGACGCT AGTCGAAGGC CGATGGAATA
9421 TAAGTGCAATG TGCTAAGTTA GATCCAAAAT TACAATCTAT GTATCAGAAA GGTAATAACC
9481 TGTGGGAGT GATAGATAAA TTGTTTCCAA TTAGGGAGA AAAGACATTT GATGTGATAT
9541 CGTTATTAGA ACCACTTGCA TTATCTTAA TTCAAACTCA TGATCTGTT AAACAACATA
9601 GAGGAGCTTT TTTAAATCAT GTGTTATCCG AGATGGAAAT AATATTGAAA TCTAGAGAA
9661 CGATTAAGGA ATTTCTGAGT GTAGATTACA TTGATAAAT TTTAGATATA TTTAATAAGT
9721 CTACAATAGA TGAAATAGCA GAGATTTCT CTTTTTTAG AACATTTGGG CATCCTCCAT
9781 TAGAAGCTAG TATTGCAGCA GAAAAGTTA GAAAATATAT GTATATTGGA AAACAATTA
9841 AATTTGACAC TATTAATAAA TGTCATGCTA TCTTCTGTAC AATAATAAAT AACGGATATA
9901 GAGGAGGCCA TGGTGGACAG TGGCCTCCTG TGACATTACC TGATCATGCA CACGAATCA
9961 TCATAAATGC TTACGGTTC AACTCTGCGA TATCATATGA AAATGCTGTT GATTATTACC
10021 AGAGCTTTAT AGGAATAAAA TTCAATAAAT TCATAGAGCC TCAGTTAGAT GAGGATTTGA
10081 CAATTTATAT GAAAGATAAA GCATTTATCT CAAAAAATC AAATTTGGAC ACAGTTTATC
10141 CTGCATCTAA TTTACTGTAC CGTACTAACG CATCCAACGA ATCAGGAGA TTAGTTGAAG
10201 TATTTATAGC AGATAGTAAA TTTGATCCTC ATCAGATATT GGATATTGTA GAATCTGGGG
10261 ACTGGTTAGA TGATCCAGAA TTTAATATTT CTTATAGTCT TAAAGAAAAA GAGATCAAAC
10321 AGGAAGGTAG ACTCTTGCA AAAATGACAT ACAAAATGAG AGCTACACAA GTTTTATCAG
10381 AGACCTACT TGCAAAATAC ATAGGAARAT TCTTTCARGA AAATGGGATG GTGAGGGAG
10441 AGATGAAT ACTTAAGAGA TTAACAACCA TATCAATATC AGGAGTTCCA CGGTATAATG
10501 AAGTGTACAA TAATTTAAA AGCCATACAG ATGACCTTAA AACCTACAAT AAAATAAGTA
10561 ATCTTAATTT GTCTTCTAAT CAGAAATCAA AGAAATTTGA ATTCAGTCA ACGGATATCT
10621 ACAATGATGG ATACGAGACT GTGAGCTGTT TCCTAACCAAC AGATCTCAA AAATACTGTC
10681 TTAATTTGGG ATATGAATCA ACAGCTCTAT TTGGAGAAC TTGCAACCAA ATATTTGGAT
10741 TAAATAAAT GTTTAATGG TTACACCTC GTCTTGAAAG AAGTACAATC TATGTAGGTG
10801 ATCCTTACTG TCCTCCATCA GATAAAGAAC ATATATCATT AGAGATCAC CCTGATCTG

10861 GTTTTACGT TCATAACCCA AGAGGGGGA TAGAAGGATT TTGTCAAAA TTATGGACAC
 10921 TCATATCTAT AAGTGAATA CATCTAGCAG CTGTTAGAAT AGGCGTGAGG GTGACTGCAA
 10981 TGGTTC AAGG AGACAATCAA GCTATAGCTG TAACCACAAG AGTACCCAAC AATTATGACT
 11041 ACAGAGTTAA GAACGAGATA GTTTATAAAG ATGTAGTGAG ATTTTTGTAT TCATTAGAG
 11101 AAGTGATGGA TGATCTAGGT CATGAACCTA AATTAATGA AACGATTATA AGTAGCAAGA
 11161 TGTTTCATATA TAGCAAAAAGA ATCTATTATG ATGGGAGAAT TCCTCTCAA GCTCTAAAAG
 11221 CATTATCTAG ATGTGCTTC TGCTCAGAGA CAGTAATAGA CGAAACAAGA TCAGCATCTT
 11281 CAAATTTGGC AACATCATT GCAAAAGCAA TTGAGAATGG TTATTCACCT GTTCTAGGAT
 11341 ATGCATGCTC AATTTTTAAG AATATTCAAC AACTATATAT TGCCTTGGG ATGAATATCA
 11401 ATCCBACTAT AACACAGAAT ATCAGAGATC AGTATTTTAG GAATCCRAAT TGGATGCAAT
 11461 ATGCCTCTTT AATACCTGCT AGTGTGGGG GATTCAAITA CATGCCCATG TCAAGATGTT
 11521 TTGTAAGGAA TATTGGTAT CCATCAGTTG CCGCATTGGC TGATATTAAG AGATTTATTA
 11581 AGCGGAATCT ATTAGACCGA AGTGTCTTT ATAGGATTAT GAATCAAGAA CCAGGTGAGT
 11641 CATCTTTTT GGACTGGGCT TCAGATCCAT ATTCATGCAA TTTACCACAA TCTCAAAATA
 11701 TAACCACCAT GATAAAAAAT ATAACAGCAA GGAATGTATT ACAAGATTCA CCAATCCAT
 11761 TATTATCTGG ATTATTACA AATACAATGA TGAAGAAGA TGAAGATTA GCTGAGTCC
 11821 TGATGGACAG GAAGGTAATT CTCCCTAGAG TTGCACATGA TATTCTAGAT AATCTCTCA
 11881 CAGGAATTAG AAATGCCATA GCTGGAATGT TAGATACGAC AAAATCACTA ATTGGGGTTG
 11941 GCATAAATAG AGGAGGACTG ACATATAGTT TGTGAGGAA AATCAGTAAT TACGATCTAG
 12001 TACAATATGA AACACTAAGT AGGACTTTGC GACTAATTGT AAGTGATAAA ATCAAGTATG
 12061 AAGATATGTG TTCCGTAGAC CTTCGCATAG CATTCGCACA AAAGATGTGG ATTCATTTAT
 12121 CAGGAGGAAG GATGATAAGT GGACTTGAAA CGCCTGACCC ATTAGATTA CTATCTGGGG
 12181 TAGTAATAAC AGGATCAGAA CATTGTAATA TATGTTATC TTCAGATGGC ACAAAACCCAT
 12241 ATACTTGGAT GTATTACCCT GGTAAATATCA AAATAGGATC AGCAGAAACA GGTATATCGT
 12301 CATTAAAGAT TCCTTATTTT GGATCAGTCA CTGATGAAAG ATCTGAAGCA CAATTAGGAT
 12361 ATATCAAGAA TCTTAGTAAA CCTGCAAAAG CCGCAATAAG AATAGCAATG ATATATACAT
 12421 GGGCATTTGG TAATGATGAG ATATCTTGGG TGGAAAGCCTC ACAGATAGCA CAAACACGCTG
 12481 CAAATTTTAC ACTAGATAGT CTCAAAATTT TAACACCGGT AGCTACATCA ACAAAATTTAT
 12541 CACACGATTT AARGGATACT GCAACTCAGA TGAATTTCTC CAGTACATCA TTGATCAGAG
 12601 TCAGCAGATT CATAACAATG TCCAATGATA ACATGCTAT CARAGAAGCT AATGAAACCA
 12661 AAGATACTAA TCTTATTTAT CAACAATAA TGTTAACAGG ATTAAGTGTG TTCGAATATT
 12721 TATTTAGATT AAAAGAAACC ACAGGACACA ACCCTATAGT TATGCAATG CACATAGAAG
 12781 ATGAGTGTG TATTAAGAAA AGTTTAATG ATGAACATAT TAATCCAGAG TCTACATTAG
 12841 AATTAATTCG ATATCCTGAA AGTAATGAAT TTATTTATGA TAAAGACCCA CTCAAAGATG
 12901 TGGACTTATC AAAACTTATG GTTATTAAAG ACCATTCTTA CACAATGTAT ATGAATTTAT
 12961 GGGATGATAC TGACATCATA CATGCAATTT CAATATGTAC TGCAATTACA ATAGCAGATA
 13021 CTATGTCACA ATTAGATCGA GATAATTTAA AAGAGATAAT AGTTATTGCA AATGATGATG
 13081 ATATTAAATAG CTTAATCACT GAATTTTGA CTCTTGACAT ACTGTATTT CTCGAAGCAT
 13141 TTGGTGGGAT ATTAGTAAT CAATTTGCAT ACACTCTTTA TAGTCTAAAA ATAGAAGGTA
 13201 GGGATCTCAT TTGGGATTAT ATAATGAGAA CACTGAGAGA TACTTCCCAT TCAATATTA
 13261 AAGTATTATC TAATGCATTA TCTCATOCTA AAGTATTCAA GAGGTTCTGG GATTGTTGGG
 13321 TTTTAAACCC TATTTATGGT CCTAATACTG CTAGTCAGA CCAAGATAAA CTGGCCTAT
 13381 CTATATGTGA ATATTCACCTA GATCTATTTA TGAGAGAATG GTTGAATGGT GTATCACTTG
 13441 AAATATACAT TTGTGACAGC GATATGGAAG TTGCAATGA TAGGAAACAA GCCTTTATTT
 13501 CTAGACACCT TTCATTTGTT TGTGTTTGA CAGAAATTGC ATCTTTCGGA CCTAACCTGT
 13561 TAACTTAAAC ATACTGGAG AGACTTGATC TATGAAAACA ATATCTTGAA TAAATATTA
 13621 AAGAAGACCC TACTCTTAAA TATGTACAAA TATCTGGATT ATTAATTAAG TCGTTCCCAT
 13681 CACTGTAAC ATACGTAAGA AAGACTGCAA TCAAATATCT AAGGATTCGC GGTATTAGTC
 13741 CACCTGAGGT AATTGATGAT TGGGATCCGG TAGAAGATGA AATATGCTG GATAACATTG
 13801 TCAAAACTAT AAATGATAAC TGAATAAAG ATAATAAAGG GAATAAAAT AACAAATTTCT
 13861 GGGACTAGC ACTTAAGAAC TATCAAGTCC TTAATAACAG ATCTATAACA AGTGATTCTG
 13921 ATGATAATGA TAGACTAGAT GCTAATACAA GTGTTTGAC ACTTCTCAA GAGGGGAAT
 13981 ATCTATCGCA TCAATTGAGA TTATTCGGAA TCAACAGCAC TAGTTGTCTG AAAGCTCTTG
 14041 AGTTATCACA AATTTAATG AAGGAAGTCA ATAAAGACAA GGACAGGCTC TTCCTGGGAG
 14101 AAGGAGCAGG AGCTATGCTA GCATGTATG ATGCCACATT AGGACCTGCA GTTAAATATT
 14161 ATAATTCAGG TTTGAATATA ACAGATGTA TTGGTCAACG AGAATTGAAA ATATTTCTCT
 14221 CAGAGGTATC ATTAGTAGGT AAAAAATTAG GAATATGAC ACAGATTTCT AACAGGGTAA
 14281 AAGTACTGTT CAATGGGAAT CCTAATTCOA CATGGATAGG AAATATGGA TGTGAGAGCT
 14341 TAATATGGAG TGAATTAAT GATAAGTCCA TTGGATTAGT ACATTTGTAT ATGGAAGGAG
 14401 CTATCGGTAA ATCAGAAGAA ACTGTTCTAC ATGAACATTA TAGTGTATA AGAATTACAT
 14461 ACTTGAATGG GGATGATGAT GTTGTTTAG TTTCCAAAT TATACCTACA ATCACTCCGA
 14521 ATTGTCTAG AATACTTTAT CTATATAAAT TATATTTGAA AGATGTAAGT ATAATATCAC

14581 TCAAACTTC TAATCCTGCA TCAACAGAAT TATATCTAAT TCGAAAGAT GCATATTGTA
14641 CTATAATGGA ACCTAGTGAA ATGTGTTTAT CAAAACITAA AAGATTGTCA CTCTTGGGAG
14701 AAAATAATCT ATTAAAATGG ATCATTITAT CAAAGAGAG GAATAATGAA TGGTTACATC
14761 ATGAATCAA AGAAGGAGAA AGAGATTATG GAATCATGAG ACCATATCAT ATGCCACTAC
14821 AAATCTTGG ATTTCAAATC AATTTAATC ATCTGGCGAA AGAATTTTA TCAACCCAG
14881 ATCTGACTAA TATCAACAAT ATAATCCAAA GTTTTCAGCG AACATAAAG GATGTTTAT
14941 TTGAATGGAT TAATATAACT CATGATGATA AGAGACATAA ATTAGCCGGA AGATATAACA
15001 TATTCCCACT GAAAAATAAG GAAAAGTTAA GACTGCTATC GAGAAGACTA GTATTAAGTT
15061 GGATTTTATT ATCATTATCG ACTCGATTAC TTACAGGTGCG CTTTCCTGAT GAAAAATTG
15121 AACATAGAGC ACAGACTGGA TATGTATCAT TAGCTGATAC TGATTTAGAA TCATTAAGT
15181 TATTGTCGAR AAACATCATT AAGAATTACA GAGAGTGTAT AGGATCAATA TCATATTGGT
15241 TTCTAACCAA AGAAGTTAAA ATACTTATGA AATTGATCGG TGSTGCTAAA TTATTAGGAA
15301 TTCCAGACA ATATAAGAA CCCGAGACC AGTTATTAGA AACTACAAAT CAACATGATG
15361 AATTTGATAT CGATTA AAC ATAAATACA TGAAGATATA TCCTAACCTT TATCTTAAAG
15421 CCTAGGAATA GACAAAAGT AAGAAAACA TGTATATAT ATATACCAA CAGAGTTCIT
15481 CTCTGTTTG GT

In a second strategy (Figure 18), chimeric PIV3-PIV2 F and HN ORFs rather than the complete ORF exchange were constructed in which regions of the PIV2 F and HN ORFs encoding the ectodomains were amplified from pLit.PIV32Fhc and pLit.PIV32HNhc, respectively, using PCR, Vent DNA polymerase (NEB, Beverly, MA), and primer pairs
5 specific to PIV2 F (5, 6 in Table 22) and HN (7, 8 in Table 22). In parallel, the regions of PIV3 F and HN ORFs encoding the ectodomains were deleted from their cDNA subclones pLit.PIV3.F3a and pLit.PIV3.HN4 (Tao et al., *J. Virol.* 72:2955-2961, 1998, incorporated herein by reference), respectively, using PCR, Vent DNA polymerase, and primer pairs
10 specific to PIV3 F (9, 10 in Table 22) and HN (11, 12 in Table 22). The amplified F and HN cDNA fragments of PIV2 and PIV3 were purified from agarose gels and ligated to generate pLit.PIV32FTM and pLit.PIV32HNTM, respectively. The chimeric F and HN constructs were digested with PpuMI plus SpeI and assembled together to generate pLit.PIV32TM, which was subsequently sequenced with the dRhodamine dye terminator sequencing kit across its PIV specific region in its entirety and found to be as designed. The 4 kb BspEI-
15 SpeI fragment from pLit.PIV32TM was then introduced into the BspEI-SpeI window of p38' Δ PIV31hc to generate p38' Δ PIV32TM. The 6.5 kb BspEI-SphI fragment from p38' Δ PIV32TM, containing the PIV3-PIV2 chimeric F and HN genes, was introduced into the BspEI-SphI window of pFLC.2G+.hc and pFLCcp45 (Skiadopoulos et al., *J. Virol.* 73:1374-81, 1999, incorporated herein by reference) to generate pFLC.PIV32TM (Table 24;
20 SEQ ID NO. 61) and pFLC.PIV32TMcp45, respectively. The nucleotide sequence of the BspEI-SpeI fragment, containing the chimeric PIV3-PIV2 F and HN genes, is submitted in the GenBank.

TABLE 24 (SEQ ID NO. 61)

Sequence of pFLC.PIV32TM, 15498 bp in sense orientation
(only the antigenome is shown)

```

1 ACCAACAAG AGAAGAACT TGCTGGGAA TATAAATTA ACTTTAAAT AACTTAGGAT
61 TAAAGACATT GACTAGAAG TCAAGAAAAG GGAACCTAT AATTTCAAA ATGTTGAGCC
121 TATTTGATAC ATTTAATGCA CGTAGGCAAG AAAACATAAC AAATCAGCC GGTGGAGCTA
181 TCATTTCCTGG ACAGAAAAT ACTGTCTCTA TATTCGCCCT TGGACCGACA ATAACTGATG
241 ATAAATGAGAA AATGACATTA GCTCTTCTAT TTCTATCTCA TTCACTAGAT AATGAGAAAC
301 AACATGCACA AAGGGCAGGG TTCTTGGTGT CTTTATTGTC AATGGCTTAT GCCAATCCAG
361 AGCTCTACCT AACACAAAAT GGAAGTAATG CAGATGTCAA GTATGTCATA TACATGATTG
421 AGAAAGATCT AAAACGGCAA AAGTATGGAG GATTTGTGT TAAGCCGAGA GAGATGATAT
481 ATGAAAAGAC AACTGATTGG ATATTTGGAA GTGACCTGGA TTATGATCAG GAAACTATGT
541 TGCAGAACGG CAGGACAAT TCAACAATTG AAGACCTTGT CCACACATTI GGGTATCCAT
601 CATGTTTAGG AGCTCTTATA ATACAGATCT GGATAGTTCT GGTCAAGCTI ATCACTAGTA
661 TCTCAGGGTT AAGAAAAGGC TTTTCACCC GATTGGAAGC TTTCCAGCAA ATGGAAACAG
721 TGCAGGCAGG GCTGGTATTG AGCGGTGACA CAGTGGATCA GATTTGGTCA ATCATGCGGT
781 CTCACAGAGG CTTGGTAACT CTTATGGTGT AAACATTAAT AACATGAAT GACAGCAGAA
841 ATGACCTCAC AACCATAGAA AAGAATATAC AAATGTTGG CACTACATA AGAGATTCAG
901 GTCTCGCTTC ATTCTTCAAT ACAATCAGAT ATGGAATTGA GACCAGATG GCAGCTTTGA
961 CTCTATCCAC TCTCAGACCA GATATCAATA GATTAAAAGC TTTGATGGAA CTGTATTTAT
1021 CAAAGGGACC ACGCGTCTCT TTCATCTGTA TCCTCAGAGA TCCTATACAT GGTGAGTTCG
1081 CACCAGGCAA CTATCCTGCC ATATGGAGCT ATGCAATGGG GGTGGCAGTT GTACAAAATA
1141 GAGCCATGCA ACAGATGTGT ACGGGAGAT CATATCTAGA CATTGATATG TTCACGCTAG
1201 GACAGCAGT AGCACGTGAT GCCGAAGCTC AAATGAGCTC AACACTGGAA GATGAACCTG
1261 GAGTGACACA CGAATCTAAA GAAAGCTTGA AGAGACATAT AAGGAACATA AACAGTTCAG
1321 AGACTCTTTC CCACAAACCG ACAGGTGGAT CAGCCATAGA GATGGCAATA GATGAAGAGC
1381 CAGAACAAAT CGAACATAGA GCAGATCAAG AACAAAATGG AGAACCTCAA TCATCCATAA
1441 TTCAATATGC CTGGGCAGAA GGAATAGAAA GCGATGATCA GACTGAGCAA GCTACAGAAT
1501 CTGACAATAT CAAGACCGAA CAACAAAACA TCAGAGACAG ACTAAACAAG AGACTCAACG
1561 ACAGAAGAAA ACAAGCAGT CAACCAACCA CTAATCCCAC AAACAGAAC AACCAGGACG
1621 AAATAGATGA TCTGTTAAC GCATTTGGAA GCAACTAATC GAATCAACAT TTTAATCTAA
1681 ATCAATAATA AATAAGAAA ACTTAGGATT AAAGATCTCT ATCATACCGG AATATAGGGT
1741 GGTAAATTTA GAGTCTGCTT GAACTCAAT CAATAGAGAG TTGATGGAAA CCGATGCTAA
1801 AAATATCAA ATCATGGATT CTTGGGAAGA GGAATCAAGA GATAAATCAA CTAATATCTC
1861 CTCGGCCCTC AACATCATTG AATTCATACT CAGCACCGAC CCCCAGGAG ACTTATCGGA
1921 AAACGACACA ATCAACACAA GAACCCAGCA ACTCAGTGCC ACCATCTGTC AACAGAAAT
1981 CAAACCAACA GAAACAGTG AGAAGATAG TGGATCAACT GACAAAAATA GACAGTCCGG
2041 GTCATCACAC GAATGTACAA CAGAAGCAAA AGATAGAAAT ATTGATCAGG AAACGTGACA
2101 GAGAGGACCT GGGAGAGAAA GCAGCTCAGA TAGTAGAGCT GAGACTGTGG TCTCTGGAGT
2161 AATCCCAGA AGCATCACAG ATCTAAAAA TGGAAACCAA AACCCGAGG ATATTGATCT
2221 CAATGAAAT AGAAAGATGG ATAAGGACTC TATTGAGGGG AAAATGCGAC AATCTGCAAA
2281 TGTTCGAAGC GAGATATCAG GAAGTGATGA CATATTTACA ACAGAACAAA GTAGAAACAG
2341 TGATCATGGA AGAAGCCTGG AATCTATCAG TACACCTGAT ACAAGATCAA TAAGTGTGT
2401 TACTGCTGCA ACACCATGAT ATGAAGAAGA AATACTAATG AAAAATAGTA GGACAAAGAA
2461 AAGTCTTCA ACACATCAAG AAGATGACAA AAGAATTAAG AAAGGGGAAA AAGGGAAGAA
2521 CTGTTTAAAG AAATCAAAAG ATACCGACAA CCAGATACCA ACATCAGACT ACAGATCCAC
2581 ATCAAAAGGG CAGAAGAAA TCTCAAAGAC AACCAACACC AACCCGACA CAAAGGGGCA
2641 AACAGAAATA CAGACAGAAAT CATCAGAAAC ACAATCCTCA TCATGGAATC TCATCATCGA
2701 CAACAAACCC GACCGAACC AACAGACAAG CACAACCTCT CCAACAACAA CTCCAGATC
2761 AACTTATACA AAAGAATCGA TCCGAACAAA CTCTGAATCC AAACCCAGA CACAAAAGAC
2821 AAATGGAAGG GAAAGGAGG ATACAGAAGA GAGCAATCGA TTTACAGAGA GGGCAATTAC
2881 TCTATTGCAG AATCTGGTG TAAATCAATC CACATCAAAA CTAGATTTAT ATCAAGACAA
2941 ACGAGTTGTA TGTGTAGCAA ATGFACTAAA CAATGTAGAT ACTGCATCAA AGATAGATTT
3001 CCTGGCAGGA TTAGTCATAG GGGTTTCAAT GGACAACGAC ACAAAATTA CACAGATACA
3061 AAATGAAATG CTAACCTCA AAGCAGATCT AAAGAAAATG GACGAATCAC ATAGAAGATT
3121 GATGAAAAT CAAAGAGAAC AACTGTCAAT GATCAGCTCA CTAATTTCAA ATCTCAAAAT
3181 TATGACTGAG AGAGGAGGAA AGAAGACCA AAATGAATCC AATGAGAGAG TATCCATGAT
3241 CAAACAAAAT TTGAAGAGG AAAAGATCAA GAAGACCAGG TTTACCAC TATGGAGGC
3301 ACAAGGCATT GACAAGAAAT TACCCGATCT ATATCCACAT GCAGAGATA CACTAGAGAA

```

3361 CGATGTACAA GTTAAATCAG AGATATTAAG TTCATACAAT GAGTCAAATG CAACAAGACT
 3421 AATACCCAAA AAAGTGAGCA GTACAATGAG ATCACTAGTT GCAGTCATCA ACAACAGCAA
 3481 TCTCTCACAA AGCACAAAAC AATCATACAT AAACGAACTC AAACGTTGCA AAAATGATGA
 3541 AGAAGTATCT GAATTAATGG ACATGTTCAA TGAAGATGTC AACCAATTGCC AATGATCCAA
 3601 CAAGAGAACG ACACCGAACG AACGACACAG AACACACAGT AGATCAAAAC CTGTCAACAC
 3661 ACACAAAATC AAGCAGAATG AAACACACAG TATCAATCAA TATACAAATA AGAAAAACTT
 3721 AGGATTTAAAG AATAAATTTAA TCCTTGTCCTA AAATGAGTAT AACTAACTCT GCAATATACA
 3781 CATTCCCAGA ATCATCATTG TCTGAAAATG GTCATATAGA ACCATTACCA CTCAAAAGTCA
 3841 ATGAACAGAG GAAAGCAGTA CCCACATTA GAGTTGCCAA GATCGGAAAT CCACCAAAAC
 3901 ACGGATCCCG GTATTTAGAT GTCTTCTTAC TCGGCTTCTT CGAGATGGAA CGAATCAAAG
 3961 ACAAAATACGG GAGTGTGAAT GATCTCGACA GTGACCCGAG TTCAAAGTT TGTGGCTCTG
 4021 GATCATTACC AATCGGATTG GCTAAGTACA CTGGGAATGA CCAGGAATG TTACAAGCCG
 4081 CAACCAAACCT GGATATAGAA GTGAGAAGAA CAGTCAAAGC GAAAGAGATG GTTGTTTACA
 4141 CGGTACAAAA TATAAAACCA GAACTGTACC CATGGTCCAA TAGACTAAGA AAAGGAATGC
 4201 TGTTCGATGC CAACAAGTT GCTTTGCTC CTCAAATGCT TCCACTAGAT AGGAGCATAA
 4261 AATTTAGAGT AATCTTCGTG AATGTACCG CAATGGATC AATAACCTTG TTCAAAATTC
 4321 CTAAGTCAAT GGCATCACTA TCTTACCCTA ACACAATATC AATCAATCTG CAGGTACACA
 4381 TAAAAACAGG GGTTCAGACT GATTCTAAAG GGATAGTTC AATTTTGGAT GAGAAAGGGG
 4441 AAAAACTCACT GAATTTCTAG GTCCATCTCG GATTGATCAA AAGAAAAGTA GGCAGAATGT
 4501 ACTCTGTTGA ATACTGTAAA CAGAAAATCG AGAAAATGAG ATTGATATTT TCTTTAGGAC
 4561 TAGTTGAGG AATCAGTCTT CATGTCAATG CAACGGGTC CATATCAAAA ACACATAGCAA
 4621 GTCAGCTGGT ATTCAAAGA GAGATTTGTT ATCCTTTAAT GGATCTAAT CCGCATCTCA
 4681 ATCTAGTTAT CTGGGCTTCA TCAGTAGAGA TTCAAAGAGT GGATGCAAT TTCCAACCTT
 4741 CTTTACCTGG CGAGTTCAGA TACTATCCTA ATATTATTGC AAAAGGAGTT GGGAAAATCA
 4801 AACCAATGGAA CTAGTAATCT CTATTTTAGT CCGGACGTAT CTATTAGCC GAAGCAAATA
 4861 AAGGATAACT AAAAATCTAG GACAAAAGAG GTCAATACCA ACAACTATTA GCAGTCACAC
 4921 TCGCAAGAAT AAGAGAGAAG GGACCAAAAA AGTCAAATAG GAGAAAATCAA AACAAAAGGT
 4981 ACAGAACACC AGAACAACAA AATCAAAACA TCCAACCTAC TCAAAAACAA AATTCAAAA
 5041 GAGACCCGCA ACACAACAAG CACTGAACAT GCATCACCTG CATCCAATGA TAGTATGCAT
 5101 TTTTGTATG TACACTGGAA TTGTAGGTTT AGATGCCATT GCTGGAGATC AACTCCTCAA
 5161 TGTAGGGGTC ATTCAATCAA AGATAAGATC ACTCATGTAC TACACTGATG GTGGCGCTAG
 5221 CTTTATTTGT GTAAAATTAC TACCCAATCT TCCCCGAAGC AATGGACATG GCAACATCAC
 5281 CAGTCTAGAT GCATATAATG TTACCCTATT TAAGTTGCTA ACACCCCTGA TTGAGAACCT
 5341 GAGCAAAAT TCTGCTGTTA CAGATACCAA ACCCCGCGCA GAACGATTG CAGGAGTCTG
 5401 TATTTGGCTT GCTGCACTAG GAGTAGCTAC AGCTGCACAA ATACCCCGAG CTGTAGCAAT
 5461 AGTAAAAGCC AATGCAATG CTGCTGGCAT AAACAATCTT GCATCTTCAA TTCATCCAC
 5521 CAACAAGGCA GTATCCGATG TGATAACTGC ATCAAGAACA ATTGCAACCC CAGTTCBAGC
 5581 GATTGAGGAT CACATCAATG GAGCCATTGT CAACGGGATA ACATCTGCAT CATGCCGTGC
 5641 CCATGATGCA CTAATTTGGT CAATATTAAT TTTGTATCTC ACTGAGCTTA CTACAATATT
 5701 TCATATCAA ATAACAACCC CTGCGCTGAC ACCACTTTCC ATCCAAGCTT TAAGAAATCCT
 5761 CCTCGGTAGC ACCTTGCCAA TTGTCAATGA ATCCAACCTC AACCAAAAC TCAACACAGC
 5821 AGAGCTGCTC AGTAGCGGAC TGTAACTGG TCAAAATATT TCCATTTCCT CAATGTACAT
 5881 CCAATGCTA ATTCAAATCA ATGTCCGAC ATTTAATATG CAACCCGGTG CGAAGGTAAT
 5941 TGATCTAAT GCTATCTCTG CAAACCATAA ATTACAAGAA GTAGTTGTAC AAGTTCCTAA
 6001 TAGAATTTCTA GAATATGCAA ATGAACACTA AAACATCCCA GCCAATGATT GTTTCGTGAC
 6061 ACCAAACTCT GTATTTTGTG GATACAATGA GGGTCCCGG ATCCCTGAAT CACAATATCA
 6121 ATGCTTAAAG GGGAACTTA ATTCTTGCAC TTTTACCCCT ATATTCGGGA ACTTCTCAA
 6181 GCGATTGCA TTTGCCAATG GTGTGCTCTA TGCCAACCTG AATCTTTGC TATGTAAGTG
 6241 TGCCGACCTT CCCCATGTTG TGTCTCAGA TGACAACCAA GGCATCAGCA TAATGATAT
 6301 TAAGAGGTGC TCTGAGATGA TGCTTGACAC TTTTTCATTT AGGATCAGAT CTACATTCAA
 6361 TGCTACATAC GTGACAGACT TCTCAATGAT TAATGCAAAAT ATTGTACATC TAAGTCTCTC
 6421 AGACTTGTCA AATCAAATCA ATTCAATAAA CAAATCTCTT AAAAGTGCTG AGGATTTGGAT
 6481 TGCAAGTAGC AACTTCTTCG CTAATCAAGC CAGAACAGCC AAGACACTTT ATTCATAAT
 6541 CATAATTAAT TTGATAATGA TCATTATATT GTTTATAAAT AATATAACGA TAATTACAAT
 6601 TGCAATTAAG TATTACAGAA TTCAAAGAG AAATCGAGTG GATCAAAATG ACAAGCCATA
 6661 TGTAATAACA AACAAATAAC ATATCTACAG ATCATTAGAT ATTAATAATA TAAAAACTT
 6721 AGGAGTAAAG TTAGCAATC CAACTTACT CATATAATG AGGAAGGACC CAATAGACAA
 6781 ATCCAAATTC GAGATGGAAT ACTGGAAGCA TACCAATCAC GGAAGGATG CTGGTAATGA
 6841 GCTGAGAGCC TCTATGGCTA CTCATGGCAA CAAGCTCACT AATAAGATAA TATACATATT
 6901 ATGGACAATA ATCCTGGTGT TATTATCAAT AGTCTTCAIC ATAGTGCTAA TTAATTCAT
 6961 CCATGAGATA ATTCATCTTG ATGTTTCTC TGGTCTTATG AATCTGATG AGTCACAGCA
 7021 AGGCATTAT CAGCCTATCA TAGAATCATT AAAATCATTG ATTGCTTTGG CCAACAGAT

7081 TCTATATAAT GTTCAATAG TAATTCCTCT TAAAATTGAC AGTATCGAAA CTGTAATACT
7141 CTCTGCTTTA AAAGATATGC ACACCGGGAG TATGTCCAAT GCCAAGTACA CGCCAGGAAA
7201 TCTGTCTCTG CATGATGCAG CATAACATCA TGGAAATAAC AAATTCCTTG TACTTGAATC
7261 ATACAAATGGG ACGCTAAAT ATGGACCTCT CCTAAATATA CCCAGCTTTA TCCCTCAGC
7321 AACATCTCCC CATGGGTGTA CTAGAATACC ATCATTTTCA CTCATCAAGA CCCATTGGTG
7381 TTACACTCAC AATGTAATGC TTGGAGATTG TCTTGATTTT ACGGCATCTA ACCAGTATTT
7441 ATCAATGGGG ATAATACAAC AATCTGCTGC AGGGTTTCCA ATTTTCAGGA CTATGAAAAAC
7501 CATTTACCTA AGTGATGGAA TCAATCGCAA AAGCTGTTCA GTCAGTGTCA TACCAGGAGG
7561 TTGTCTCTTG TATTGCTATG TAGTACAAAG GTCTGAAAAA GAAGATTATG CCACGACTGA
7621 TCTAGCTGAA CTGAGACTTG CTTTCTATTA TTATAATGAT ACCTTTATTTG AAAGAGTCAT
7681 ATCTCTTCCA AATACAACAG GGCAGTGGGC CACAATCAAC CCTGCAGTCG GAAGCGGGAT
7741 CTATCATCTA GGCTTTATCT TATTTCTCTG ATATGGTGGT CTCATAAATG GGACTACTTC
7801 TTACAAATGAG CAGTCTCAC GCTATTTTAT CCCAAAACAT CCCAACATAA CTTGTGCCGG
7861 TAATCCAGC AAACAGGCTG CAATAGCAGC GAGTTCCTAT GTCATCCGGT ATCACTCAAA
7921 CAGGTTAATT CAGAGTCTG TTCTTATTTG TCCATTGCTC GACATGCATA CAGAAGAGTG
7981 TAATCTACTT ATGTTTAAACA ATTCCCAAGT CATGATGGGT GCAGAAGGTA GCCTCTATGT
8041 TATTGGTAAT AATTTGTATT ATTATCAACG CAGTTCCTCT TGGTGGTCTG CATCGCTCTT
8101 TTACAGGATC AATACAGATT TTTCTAAAGG AATTCCTCCG ATCATTGAGG CTCATGGGT
8161 ACCGCTCTAT CAAGTTCCTC GTCTGGAGT CATGCCATGC AATGCAACAA GTTTTTGCCC
8221 TGCTAATTCG ATCACAGGGG TGTACGCAGA TGTGTGGCCG CTTAATGATC CAGAATCTAT
8281 GTCACGTAAT GCTCTGAACC CCAACTATCG ATTTGCTGGA GCCTTTCTCA AAAATGAGTC
8341 CAACCGAACT AATCCACAT TCTACACTGC ATCGGCTAAC TCCCTCTTAA ATACTACCGG
8401 ATTCAACAAC ACCAATCACA AAGCAGCATA TACATCTCA ACCTGCTTAA AAAACACTGG
8461 AACCACAAAA ATTTATTGTT TAATAATAAT TGAATGGGC TCATCTCTTT TAGGGGAGTT
8521 CCAATAATAA CCATTTTAA GGAACATAAT GCTTTAAGCT TCATAATTA CCATAATATG
8581 CATCAATCTA TCTATAATAC AAGTATATGA TAAGTAAATCA GCAATCAGAC AATAGACAAA
8641 AGGGAAATAT AAAAAACTTA GGAGCAAAGC GTGCTCGGGA AATGACACT GAATCTAACA
8701 ATGGCACTGT ATCTGACATA CTCTATCTTG AGTGTCACTT TAACCTCTCT ATCTGTAAGG
8761 GTAATAATAGC ACAATTACAC ACTATTATGA GTCTACCTCA GCCTTATGAT ATGGATGAGC
8821 ACTCAATACT AGTTATCACT AGACAGAAAA TAAAACCTTA TAAATGGAT AAAAGACAC
8881 GATCTATTAG AAGATAAAA TTAATATTAA CTGAAAAAGT GAATGACTTA GGAATAATCA
8941 CATTATCAG ATATCCAGAA ATGTCAAAG AAATGTTCAA ATTATATATA CTTGGTATTA
9001 ACAGTAAAGT GACTGAATTA TTAATTAAG CAGATAGAAC ATATAGTCAA ATGACTGATG
9061 GATTAAGAGA TCTATGGATT AATGTGCTAT CAAAATTAGC CTCAAAAAAT GATGGAAGCA
9121 ATTAGTATCT TAATGAAGAA ATTAATAATA TATCGAAGT TCACACACC TATAATCAG
9181 ATAAATGGTA TAATCCATTC AAAACATGGT TTAATATCAA GTATGATATG AGAAGATTAC
9241 AAAAAAGCTG AAATGAGATC ACTTTTAAAG TTGGGAAGGA TTATAACTTG TTAGAAGACC
9301 AGAAGAAATTT CTTATTGATA CATCCAGAA TGGTTTGGAT ATTAGATAAA CAAAACATA
9361 ATGGTTATCT AATTACTCCT GAATTAGTAT TGATGTATTG TGACGTAGTC GAAAGCCGAT
9421 GGAATATAAG TGCATGTGCT AAGTTAGATC CAAAATTACA ATCTATGTAT CAGAAGGTA
9481 ATAACTGTG GSAAGTGATA GATAAATGT TTCCAATTAT GGGAGAAAAG ACATTTGATG
9541 TGATATCGTT ATTAGAACCA CTGCAATTA CCTTAATTC AACTCATGAT CCTGTTAAAC
9601 AACTAAGAGG AGCTTTTTTA AATCATGTGT TATCCGAGAT GGAATTAATA TTTGAATCTA
9661 GAGAATCGAT TAAGGAATTT CTGAGTGTAG ATTACATTGA TAAAAATTTA GATATATTTA
9721 ATAACTCTAC AATAGATGAA ATAGCAGAGA TTTTCTCTTT TTTTAGAACA TTTGGGCATC
9781 CTCCATTAGA AGCTAGTATT GCAGCAGAAA AGGTTAGAAA ATATATGTAT ATTGGAAAAC
9841 AATTAATATT TGACACTATT AATAAATGTC ATGCTATCTT CTGTACATA ATAATTAACG
9901 GATATAGAGA GAGGCATGGT GGACAGTGGC CTCCTGTGAC ATTACCTGAT CATGCACAG
9961 AATTCATCAT AAATGCTTAC GGTTCAAACT CTGCGATATC ATATGAAAAT GCTGTTGATT
10021 ATTACCAGAG CTTTATAGGA ATAAAATTC AATAAATTCAT AGAGCCTCAG TTAGATGAGG
10081 ATTTGACAAT TTATATGAAA GATAAAGCAT TATCTCCAAA AAAATCAAAT TGGGACACAG
10141 TTTATCTTGC ATCTAATTTA CTGTACCSTA CTAACGCATC CAACGAATCA CGAAGATTAG
10201 TTGAAGTATT TATAGCAGAT AGTAAATTTG ATCCTCATCA GATATTGGAT TATGTAGAAT
10261 CTGGGACTG GTTAGATGAT CCAGAATTTA ATATTCTTAA TAGTCTTAAA GAAAAAGAGA
10321 TCAACACGGA AGGTAGACTC TTTGCNAAA TGACATACAA AATGAGAGCT ACACAGTTT
10381 TATCAGAGAC CCTACTTGCA AATACATAG GAAAATCTT TCAAGAAAAT GGSATGGTGA
10441 AGGGAGAGAT TGAATTAATT AAGAGATTAA CAACCATATC AATATCAGGA GTTCCACGGT
10501 ATAAATGAAT GTACAATAAT TCTAAAAGCC ATACAGATGA CCTTAAAACC TACAATAAAA
10561 TAAGTAATCT TAATTTGCTC TCTAATCAGA AATCAAAGAA ATTTGAATTC AAGTCAACGG
10621 ATATCTACAA TGATGGATAC GAGACTGTGA GCTGTTCTCT AACACAGAT CTCAAAAAAT
10681 ACTGCTTAA TTGAGATAT GAATCAACAG CTCTATTGAG AGAACTTGC AACCAATAT
10741 TTGGATTAAT TAAATGTTT AATTGGTTAC ACCCTCGTCT TGAAGGAAGT ACAATCTATG

10801 TAGGTGATCC TTACTGTCCT CCATCAGATA AAGAACATAT ATCATTAGAG GATCACCCCTG
 10861 ATCTGGGTTT TTACGTTTAT AACCCAGAG GGGGTATAGA AGGATTTTGT CAAAAATTAT
 10921 GGACACTCAT ATCTATAAGT GCAATACATC TAGCAGCTGT TAGAATAGGC GTGAGGGTGA
 10981 CTGCATGGT TCAGGGAGAC AATCAAGCTA TAGCTGTAAC CACAAGAGTA CCCAACAT
 11041 ATGACTACAG AGTTAAGAAG GAGTAGTGT ATAAAGATGT AGTGAGATT TTTGATTCAT
 11101 TAAGAGAAGT GATGGATGAT CTAGGTCATG AACTTAAATT AAATGAAACG ATTATAAGTA
 11161 GCAAGATGTT CATATATAGC AAAAGAATCT ATTAGATGG GAGAATCTCT CCTCAAGCTC
 11221 TAAAAGCATT ATCTAGATGT GTCTCTGGT CAGAGACAGT AATAGACGAA ACAAGATCAG
 11281 CATCTTCAAA TTGGCAACA TCATTGGCAA AAGCAATGA GAATGGTTAT TCACCTGTTC
 11341 TGGGATATGC ATGCTCAATT TTTAAGATA TTCACCACT ATATATTGCC CTTGGGATGA
 11401 ATATCAATCC AACTATAACA CAGAATATCA GAGATCAGTA TTTTAGGAAT CCAATTTGGA
 11461 TGCAATATGC CTCTTTAATA CCTGCTAGTG TTGGGGGATT CAATTACATG GCCATGTCAA
 11521 GATGTTTTGT AAGGAATATT GGTGATCCAT CAGTTGCCG ATTGGCTGAT ATTTAAAGAT
 11581 TTATTAAGGC GAATCTATTA GACCGAAGTG TTCTTTATAG GATTATGAAT CAAGAACCAG
 11641 GTGATCATC TTTTTGGAC TGGGCTTCAG ATCCATATTC ATGCAATTTA CCACATCTC
 11701 AAAATATAAC CACCATGATA AAAATATAA CAGCAAGGAA TGTATTACAA GATTCACCAA
 11761 ATCCATTATT ATCTGGATTA TTCACAATA CAATGATAGA AGAAGATGAA GAATTAGCTG
 11821 AGTTCCTGAT GGACAGGAG GTAATTCTCC CTAGAGTTGC ACATGATATT CTAGATAATT
 11881 CTCTCACAGG AATTAGAAAT GCCATAGCTG GAATGTTAGA TAGCACAAA TCACATTAATC
 11941 GGGTTGGCAT AAATAGAGGA GGACTGACAT ATACTTTGTT GAGGAAAATC AGTAATTACG
 12001 ATCTAGTACA ATATGAACA CTAACTAGGA CTTTGGCACT AATTGTAAGT GATAAATCA
 12061 AGTATGAAGA TATGTGTTCC GTAGACCTTG CCATAGCATT GCGACAAAAG ATGTGGATTC
 12121 ATTTATCAGG AGGAAGGATG ATAGTGGAC TTGAACGCC TGACCCATTA GAATTACTAT
 12181 CTGGGGTAGT AATAACAGGA TCAGAACAT TAAAAATATG TTATTCTTCA GATGGCACAA
 12241 ACCCATATAC TTGGATGAT TTACCCGTA ATATCAAAAT AGGATCAGCA GAAACAGGTA
 12301 TATCGTCATT AAGAGTTCTT TATTTGGAT CAGTCACTGA TGAAGATCT GAAGCACAAT
 12361 TAGGATATAT CAAGAATCTT AGTAAACCTG CAAAAGCCGC AATAAGAATA GCAATGATAT
 12421 ATACATGGCC ATTTGGTAAAT GATGAGATA CTTGGATGGA AGCCTCACAG ATAGCACAAA
 12481 CACGTGCAAA TTTTACACTA GATAGTCTCA AAATTTAAC ACCGGTAGCT ACATCAACAA
 12541 ATTTATCACA CAGATTAAG GATACTGCAA CTCAGATGAA ATTCCTCAGT ACATCATTGA
 12601 TCAGAGTCAG CAGATTCATA ACAATGTCCA ATGATAACAT GTCTATCAAA GAAGCTAATG
 12661 AAACCAAGA TACTAATCTT ATTTATCAAC AAATAATGTT AACAGGATTA AGTGTTCG
 12721 AATATTTATT TAGATTAATA GAAACCACAG GACACAACCC TATAGTTATG CATCTGCACA
 12781 TAGAAGATGA GTGTTGATT AAAGAAAGTT TTAATGATGA ACATATTAAT CCAGAGCTCA
 12841 CATTAGAATT AATTCGATAT CCTGAAGTA ATGAATTTAT TTATGATAAA GACCCACTCA
 12901 AAGATGIGGA CTTATCAAAA CTTATGGTTA TTAAGACCA TTCTTACACA ATTGATATGA
 12961 ATATTGGGA TGATCTGAC ATCATACATG CAATTTCAAT ATGACTGCA ATTACAATAG
 13021 CAGATACTAT GTCACAATTA GATCGAGATA ATTTAAAAGA GATAATAGTT ATTGCAATG
 13081 ATGATGATAT TAATAGCTTA ATCACTGAAT TTTGACTCT TGACATACTT GTATTTCTCA
 13141 AGACATTTGG TGGATATTA GTAATCAAT TTGCATACAC TCTTTATAGT CTAAAAATAG
 13201 AAGGTAGGGA TCTCATTGG GATTATATAA TGAGAACACT GAGAGATACT TCCCATTCAA
 13261 TATTAAGAT ATATCTAAT GCATTATCTC ATCCTAAGT ATTCAGAGG TTCTGGGATT
 13321 GTGGAGTTT AAACCTATT TATGGTCTTA ATCTGCTAG TCAAGACCAG ATAAAACCTG
 13381 CCCTATCTAT ATGTGAATAT TCACTAGATC TATTTATGAG AGAATGGTTG AATGGTGTAT
 13441 CACTTGAAT ATACATTTG GACAGCGATA TGGAAAGTGC AAATGATAGG AAACAAGCCT
 13501 TTATTTCTAG ACACCTTCA TTTGTTGTT GTTTAGCAGA AATTGCATCT TTCGGACCTA
 13561 ACCTGTTAAA CTTAACATAC TTGGAGAGAC TTGATCTATT GAAACAATAT CTTGAATFAA
 13621 ATATTAAGA AGACCTACT CTTAAATATG TACAATATC TGGATTATTA ATTAATCGT
 13681 TCCCATCAAC TGTAACATAC GTAAGAAGA CTGCAATCAA ATATCTAAG ATTCCGGGTA
 13741 TTAGTCCACC TGAGGTAATT GATGATTGG ATCCGGTAGA AGATGAAAT ATGCTGGATA
 13801 ACATTGTCAA AACTATAAAT GATAACTGTA ATAAAGATAA TAAAGGGAAT AAAATTAACA
 13861 ATTTCTGGGG ACTAGCACT AAGAATATC AAGTCTTAA AATCAGATCT ATAACAGTG
 13921 ATCTGATGA TAATGATAGA CTAGATGCTA ATACAAGTGG TTTGACACTT CCTCAAGGAG
 13981 GGAATPATCT ATCGCATCAA TTGAGATTAT TCGGAATCAA CRGCACTAGT TGCTGAAAG
 14041 CTCTTGAGTT ATCACAATTT TTAATGAAG AAGTCAATAA AGACAAGGAC AGGCTCTTCC
 14101 TGGGAGAAGG AGCAGGAGCT ATGCTAGCAT GTTATGATGC CACATTAGGA CCTGCAGTTA
 14161 ATATTATAA TTCAGGTTG AATATAACAG ATGTAATTGG TCAACGAGA TTGAAATAT
 14221 TTCTTCAGA GGATCATTG GTAGGTAATA AATFAGGAAA TGTGACACAG ATTCATAACA
 14281 GGGTAAAAGT ACTGTTCAAT GGGAACTCA ATTCAACATG GATAGGAAT ATGGAATGTG
 14341 AGAGCTTAAT ATGGAGTGAA TTAATGATA AGTCCATTGG ATTAGTACAT TGTGATATGG
 14401 AAGGAGCTAT CGGTAATCA GAAGAAGCTG TTCTACATGA ACATATAGT GTTATAAGAA
 14461 TTACATACTT GATGGGGAT GATGATGTTG TTTTACTTTC CAAAATTATA CCTACATCA

14521 CTCGGAATTG GTCTAGAATA CTTTATCTAT ATAAATTATA TTGGAARGAT GTAAGTATAA
14581 TATCACTCAA AACTTCTAAT CCTGCATCAA CAGAATTATA TCTAATTTCC AAAGATGCAT
14641 ATTGTAATAT AATGGAACCT AGTGAATTG TTTTATCAAA ACTTAAAGA TTGCACTCT
14701 TGGAGAAAA TAATCTATTA AAATGGATCA TTTTATCAAA GAAGAGGAAT AATGAATGCT
14761 TACATCATGA AATCAAGAA GGAGAAAGAG ATTATGGAAT CATGAGCCA TATCATATGG
14821 CACTACAAAT CTTTGGATTT CAAATCAATT TAAATCATCT GCGAAAGAA TTTTATCAA
14881 CCCCAGATCT GACTAATATC AACAATATA TCCAAAGTTT TCAGCGAACA ATAAAGGATG
14941 TTTTATTGA ATGGATTAAT ATAACCTATG ATGATAAGAG ACATAAATTA GCGGAAGAT
15001 ATAACATATT CCCACTGAAA AATAAGGGAA AGTTAAGACT GCTATCGAGA AGACTAGTAT
15061 TAAGTTGGAT TTCATTATCA TTATCGACTC GATTACTTAC AGSTCGCTTT CCTGATGAAA
15121 AATTGAACA TAGAGCACAG ACTGGATATG TATCATTAGC TGATCTAGT TTAGAATCAT
15181 TAAAGTTATT GTCGAAAAC ATCATTAGA ATTACAGAGA GTGTATAGGA TCAATATCAT
15241 ATTGGTTTCT AACCAAGAA GTTAAATAC TTATGAAAT GATCGGTGGT GCTAATATAT
15301 TAGGAATTC CAGACAAAT AAGAACCCT AAGACCAGTT ATTAGAAAC TACAATCAAC
15361 ATGATGAATT TGATATCGAT TAAACATAA ATACAATGAA GATATATCCT AACCTTTATC
15421 TTTAAGCCTA GGAATAGACA AAAAGTAAGA AAAACATGTA ATATATATAT ACCAACAGA
15481 GTTCTTCTCT TGTTTGGT

In a third strategy (Figure 19), chimeric PIV3-PIV2 F and HN genes were constructed in which regions of the PIV2 F and HN ORFs encoding the ectodomains and the transmembrane domains were amplified from pLit.PIV32Fhc and pLit.PIV32HNhc, respectively, using PCR, Vent DNA polymerase, and primer pairs specific to PIV2 F (13, 14 in Table 22) and PIV2 HN (15, 16 in Table 22). In parallel, the partial ORFs of PIV3 F and HN genes encoding the ectodomains plus transmembrane domains were deleted from their cDNA subclones pLit.PIV3.F3a and pLit.PIV3.HN4 (Tao et al., *J. Viro.* 72:2955-2961, 1998, incorporated herein by reference), respectively, using PCR, Vent DNA polymerase, and primer pairs specific to PIV3 F (17, 18 in Table 22) and PIV3 HN (19, 20 in Table 22).

5 The F and HN cDNA fragments of PIV2 and PIV3 were gel purified and ligated to generate pLit.PIV32FCT and pLit.PIV32HNCT, respectively. The chimeric F and HN constructs were digested with PpuMI plus SpeI and assembled together to generate pLit.PIV32CT, which was sequenced across the PIV specific region in its entirety and found to be as designed. The 4 kb BspEI-SpeI fragment from pLit.PIV32CT was introduced into the

10 BspEI-SpeI window of p38' Δ PIV31hc to generate p38' Δ PIV32CT. The 6.5 kb BspEI-SphI fragment from p38' Δ PIV32CT, containing the PIV3-PIV2 F and HN chimeric genes, was introduced into the BspEI-SphI window of pFLC.2G+.hc and pFLCcp45, to generate pFLC.PIV32CT (Table 25, SEQ ID NO. 62) and pFLC.PIV32CTcp45, respectively. The nucleotide sequence of this *BspEI-SpeI* fragment is submitted in the GenBank.

15

TABLE 25 (SEQ ID NO. 62)

Sequence of pFLC.PIV32CT, 15474 bp in sense orientation
(only the insert is shown)

```

1 ACCAACAAG AGAAGAACT TGTCTGGAA TATAAATTA ACTTTAAAT AACTTAGGAT
61 TAAAGACATT GACTAGAAGG TCAAGAAAAG GGAACCTAT AATTCAAAA ATGTTGAGCC
121 TATTGATAC ATTTAATGCA CGTAGGCAAG AAAACATAAC AAAATCAGCC GGTGGAGCTA
181 TCATTCCCTGG ACAGAAAAT ACTGTCCTA TATTCGCCCT TGGACCGACA ATAACTGATG
241 ATAATGAGAA AATGACATTA GCTCTTCTAT TTCTATCTCA TTCACTAGAT AATGAGAAAC
301 AACATGCACA AAGGGCAGGG TTCTTGGTGT CTTTATTGTC AATGGCTTAT GCCAATCCAG
361 AGCTCTACCT AACAAACAAT GGAAGTAATG CAGATGTCAA GTATGTGATA TACATGATTG
421 AGAAGATCT AAAACGGCAA AAGTATGGAG GATTTGGT TAAGACGAGA GAGATGATAT
481 ATGAAAAGAC AACTGATTGG ATATTTGGAA GTGACCTGGA TTATGATCAG GAAACTATGT
541 TGCAGAACGG CAGGAACRAT TCAACAATTG AAGACCTGT CCACACATTT GGGTATCCAT
601 CATGTTTAGG AGCTCTTATA ATACAGATCT GGATAGTTCT GGTCAAAGCT ATCACTAGTA
661 TCTCAGGGTT AAGAAAAGCC TTTTTCACCC GATTGGAGGC TTTGAGCAA GATGGAAACAG
721 TGCAGGCAGG GCTGGTATTG AGCGGTGACA CAGTGGATCA GATTGGGTCA ATCATGCGGT
781 CTCAACAGAG CTTGGTAACT CTTATGTTG AAACATTAAT AACAAATGAAT ACCGAGAGAA
841 ATGACCTCAC AACCATAGAA AAGAATATAC AAATTTGGG CAACTACATA AGAGATGCAG
901 GTCTGCTTTC ATTCTCAAT ACAATCAGAT ATGGAATTGA GACCAGAAAT GCAGCTTTGA
961 CTCTATCCAC TCTCAGACCA GATATCAATA GATTAAARGC TTTGATGAAA CTGTATTTAT
1021 CAAAGGGACC ACGCGCTCCT TTCACTGTGA TCCTCAGAGA TCCTATACAT GGTGAGTTCG
1081 CACCAGGCAA CTATCCTGCC ATATGGAGCT ATGCAATGGG GGTGGCAGT GTACAAAATA
1141 GAGCCATGCA ACAGTATGTG ACGGGAAGAT CATATCTAGA CATTGATATG TTCACAGTAG
1201 GACAAGCAGT AGCACGTGAT GCCGAAGCTC AAATGAGCTC AACACTGGAA GATGAACTTG
1261 GAGTGACACA CGAATCTAAA GAAAGCTTGA AGAGACATAT AAGGAACATA AACAGTTACG
1321 AGACATCTTT CCACAAAACCG ACAGGTGGAT CAGCCATAGA GATGGCAATA GATGAAGAGC
1381 CAGAACAATT CGAACATAGA GCAGATCAAG AACAAATGG AGAACCTCAA TCATCCATAA
1441 TTCAATATGC CTGGGCAGAA GGAATAGAAA GCGATATCA GACTGAGCAA GCTACAGAAAT
1501 CTGAAATAT CAAGACCGAA CAACAAAACA TCAGAGACAG ACTAAAACAG AGACTCAACG
1561 ACAAGAGAAA ACAAGCAGT CAACCAACCA CTAATCCAC AAACAGAAAC AACCCAGGAG
1621 AAATAGATGA TCTGTTTAA CCAATTTGAA GCAACTAATC GAATCAACAT TTTAATCTAA
1681 ATCAATAATA AATAAGAAA ACTTAGGATT AAGAATCCT ATCATAACCG AATATAGGTT
1741 GGTAATAATTA GAGTCTGCTT GAACTCAAT CAATAGAGAG TTGATGAAA GCGATGCTAA
1801 AAATATCAA ATCATGATT CTGGAAGA GGAATCAGA GATAAATCAA CTAATATCTC
1861 CTCGGCCCTC AACATCATTT AATTCATACT CAGCACCGAC CCCAAGAAG ACTTATCGGA
1921 AAACGACACA ATCAACACAA GAACCCAGCA ACTCAGTGCC ACCATCTGTC AACCCAGAAAT
1981 CAACACRACA GAAACAAGT AGAAAGATAG TGGATCACT GACAAAAATA GACAGTCCGG
2041 GTCATCACAC GAATGTACAA CAGAAGCAAA AGATAGAAAT ATTGATCAGG AAACGTGACA
2101 GAGAGGACCT GGGAGAAGAA GCAGTCAGA TAGTAGAGCT GAGACTGTGG TCTCTGGAGG
2161 AATCCCCAGA AGCATCACAG ATTCTAAAAA TGGAAACCAA AACACGGAGG ATATTGATCT
2221 CAATGAAAT AGAAGATGG ATAAGACTC TATTGAGGGG AAAATGCGAC AATCTGCAAA
2281 TGTCCAAAGC GAGATATCAG GAAGTGATGA CATATTTACA ACAGAACAAA GTAGAAACAG
2341 TGATCATGGA AGAAGCCTGG AATCTATCAG TACACCTGAT ACAAGATCAA TAAGTGTGT
2401 TACTGCTGCA ACACCAAGT ATGAAGAAGA AATACTAATG AAAAATAGTA GGACAAAGAA
2461 AAGTTCTTCA ACACATCAAG AAGATGACAA AAGAATTAAA AAAGGGGAAA AAGGGAAAGA
2521 CTGTTTAAAG AATCAAAAAG ATACCGACAA CCAGATACCA ACATCAGACT ACAGATCCAC
2581 ATCAAAAAGG CAGAAGAAA TCTCAAAGAC AACACCCACC AACACCGACA CAAGGGGCA
2641 AACAGAAATA CAGACAGAAAT CATCAGAAAC ACAATCCTCA TCATGGAATC TCATCATCGA
2701 CAACAACACC GACCGGAACG AACAGACAAG CACAACCTCT CCAACACAAA CTTCAGATC
2761 AACTTATACA AARGAATCGA TCCGAACAAA CTCTGAATCC AAACCCAGA CACAAAAGAC
2821 AAATGGAAG GAAAGGAAG ATACAGAGA GAGCAATCGA TTTACAGAGA GGGCAATTAC
2881 TCTATTGACG AATCTTGGTG TAATTCATC CACATCAAAA CTAGATTTAT ATCAAGACAA
2941 ACGAGTTGTA TGTGTAGCAA ATGTACTAAA CAATGTAGAT ACTGCATCAA AGATAGATTT
3001 CCTGGCAGGA TTAGTCATAG GGGTTTCAAT GGACACGAC ACAAATTTAA CACAGATACA
3061 AAATGAAATG CTAACCTCA AAGCAGATCT AAAGAAAATG GACGAATCAC ATAGAAGATT
3121 GATAGAAAAT CAAGAGAAC AACTGTCAAT GATCAGTCA CTAATTTCAA ATCTCAAAAT
3181 TATGACTGAG AGAGGAGGAA AGAAGACCA AAATGAAATCC AATGAGAGAG TATCCATGAT
3241 CAAAACAAA TGAAGAAG AAAAGATCAA GAAGACCAGG TTTGACCCAC TTATGGAGGC
3301 ACAAGGCATT GACAAGAATA TACCCGATCT ATATCGACAT GCAGGAGATA CACTAGAGAA

```

3361 CGATGTACAA GTTAAATCAG AGATATTAAG TTCATACAAT GAGTCAAATG CAACAAGACT
 3421 AATACCCAAA AAAGTGAGCA GTACAATGAG ATCACTAGTT GCAGTCATCA ACAACAGCAA
 3481 TCTCTCACAA AGCACAAAAC AATCATACAT AAACGAACTC AAACGTTGCA AAAATGATGA
 3541 AGAAGTATCT GAATTAATGG ACATGTTCAA TGAAGATGTC AACAAATGCC AATGATCCAA
 3601 CAAAAGAACG ACACCGAACA AACAGACAAG AAACAACAGT AGATCAAAC CTGTCAACAC
 3661 ACACAAAATC AAGCAGAAATG AAACAACAGA TATCAATCAA TATACAAAATA AGAAAAACTT
 3721 AGGATTAAG AATAAATTAA TCCTTGTCCTA AAATGAGTAT AACTAACTCT GCAATATACA
 3781 CATTCACAGA ATCATCATTG TCTGAAAATG GTCATATAGA ACCATTACCA CTCAAAGTCA
 3841 ATGAACAGAG GAAAGCAGTA CCCCACATTA GAGTTGCCAA GATCGGAAAT CCACCAAAAC
 3901 ACGGATCCCG GTATTTAGAT GTCTTCTTAC TCGGCTTCTT CGAGATGGAA CGAATCAAAG
 3961 ACAAATACGG GAGTGTGAAT GATCTCGACA GTGACCCGAG TTCAAAGTT TGTGGCTCTG
 4021 GATCATTACC AATCGGATTG GCTAAGTACA CTGGGAATGA CCAGGAATTG TTACAAGCCG
 4081 CAACCAAACCT GGATATAGAA GTGAGAAGAA CAGTCAAAGC GAAAGAGATG GTTGTTTACA
 4141 CGGTACAAAA TATAAAACCA GAACGTGACC CATGGTCCAA TAGACTAAGA AAAGGAATGC
 4201 TCTTCGATCT CAACAAGATT GCTCTTGCTC CTCATGCTCT TCCACTAGAT AGGAGCATAA
 4261 AATTTAGAGT AATCTTCGCG AATTTGACGG CAATTTGATC AATAACCTTG TTCAAAATTC
 4321 CTAAGTCAAT GGCATCACTA TCTCTACCCA ACACAATATC AATCAATCTG CAGTACACA
 4381 TAAAACAGG GGTTCAGACT GATTCTAAGG GGATAGTTCA AATTTTGGAT GAGAAAGGCG
 4441 AAAAACTACT GAATTTCAAT GTCCATCTCG GATTGATCAA AAGAAAAGTA GGCAGAATGT
 4501 ACTCTGTTGA ATACTGTAAA CAGAAAATCG AGAAAATGAG ATTGATATTT TCTTTAGGAC
 4561 TAGTTGGAGG AATCAGTCTT CATGTCAATG CAACCTGGTC CATATCAAAA ACACCTAGCAA
 4621 GTCAGCTGGT ATTCAAAGA GAGATTGTTT ATCCTTAAT GGATCTAAAT CCGCACTCA
 4681 ATCTAGTTAT CTGGGCTTCA TCAGTAGAGA TTACAAGAGT GGATGCAAT TTCCACCTT
 4741 CTTTACCCTG CGAGTTCAGA TACTATCCTA ATATTATTGC AAAAGGAGTT GGGAAAATCA
 4801 AACAAATGAA CTAGTAATCT CTATTTTAGT CCGGAGCTAT CTATTAAGCC GAAGCAAATA
 4861 AAGGATAACT AAAAATCTAG GACAAAAGAG GTCAAATACCA ACAACTATTA GCAGTCACAC
 4921 TCGCAAGAAT AAGAGAGAAG GGACCAAAAA AGTCAAATAG GAGAAATCAA AACAAAAGGT
 4981 ACAGAACACC AGAACACAAA AATCAAACA TCCAACCTAC TCAAAAACAA AATTTCAAATA
 5041 GAGACCCGCA ACACAACAG CACTGAACAT GCATCACCTG CATCCAAATG TAGTATGCAT
 5101 TTTTGTATG TACACTGGA TGTAGTGTTC AGATGCCATT GCTGGAGATC AACTCTCAA
 5161 TGTAGGGGTC ATTCAATCAA AGATAAGATC ACTCAATGAT TACACTGATG GTGGCGCTAG
 5221 CTTTATTTGT GTAAAATTTAC TACCCAACTC TCCCCCAAGC AATGGAACAT GCAACATCAC
 5281 CAGTCTAGAT GCATATAATG TTACCCATTT TAAGTTGCTA ACACCCCTGA TTGAGAACCT
 5341 GAGCAAAAT TCTGCTGTTA CAGATACCAA ACCCCGCGCA GAACGATTG CAGGAGTCGT
 5401 TATTTGGGCT GCTGCACTAG GAGTAGCTAC AGCTGCACAA ATAACCCGAG CTGTAGCAAT
 5461 AGTAAAAGCC AATGCAATG CTGCTGCGAT AAACAATCTT GCATCTCAA TTCAATCCAC
 5521 CAACAAGGCA GTATCCGATG TGATAACTGC ATCAAGAACA ATTGCAACCG CAGTTCAGC
 5581 GATTGAGGAT CACATCAATG GAGCCATTTG CAACGGGATA ACATCTGCAT CATGCCGTGC
 5641 CCATGATGCA CTAATTTGGT CAATATTAAT TTTGTATCTC ACTGAGCTTA CTACAATATT
 5701 TCATATCAAA ATAACAACC CTGGGCTGAC ACCACTTTCC ATCCAAGCTT TAAGAATCCT
 5761 CCTCGGTAGC ACCTTGCCAA TTGTCAATGA ATCCAACCTC AACACAAAAC TCAACACAGC
 5821 AGAGGTGCTC AGTAGGGCAG TGTAACTGG TCAATAAAT TTCAATTTCC CAATGTACAT
 5881 GCAAAATGCTA ATTCAAATCA ATGTTCGAGC ATTTATAATG CAACCCGGTG CGAAGGTAAT
 5941 TGATCTAAT GCTATCTCTG CAAACCATAA ATTACAAGAA GTAGTTGTAC AAGTTCCTAA
 6001 TAGAATCTA GAATATGCAA ATGAATACA AAACATCCCA GCCAATGATT GTTTCGTGAC
 6061 ACCAAACTCT GTATTTTGTG ATACAATGA GGGTTCCCGG ATCCCTGAAT CACAATATCA
 6121 ATGCTTAAGG GGGAACTTAA ATTCTTGAC TTTTACCCCT ATTATCGGGA ACTTCTCAA
 6181 GCGATTGCGA TTGCGCAATG GTGTGCTCTA TGCCAACCTG AAATCTTTGC TATGTAAGTG
 6241 TGCCGACCCCT CCCCATGTTG TGTCTCAAGA TGACAACCAA GGCATCAGCA TAATGATAT
 6301 TAAGAGGTGC TCTGAGATGA TGCTTGACAC TTTTTCATTT AGGATCACAT CTACATTCAA
 6361 TGCTACATAC GTGACAGACT TCTCAATGAT TAATGCAAT ATTTGATCAT TAAGTCTCT
 6421 AGACTTGTC AATCAAATCA ATTCAATAAA CAAATCTCTT AAAAGTGCTG AGGATTGGAT
 6481 TGCAGATAGC AACTTCTCG CTAATCAAGC CAGAACAGCC AAGACACTTT ATTCACTAAG
 6541 TGCAATCGCA TTAATACTAT CAGTGATTAC TTTGGTTGTT GTGGGATTGC TGATTGCCTA
 6601 CATCATCAAG TATTACAGAA TTCAAAAGAG AATTCGAGTG GATCAAATG ACRAAGCCATA
 6661 TGTACTAACA AACAAATAC ATATCTACAG ATCATTAGAT ATTAATATTA TAAAATCTT
 6721 AGGAGTAAAG TTRCGAATC CAACTCTACT CATATAATG AGGAAGGACC CAATAGACAA
 6781 ATCCAAATTC GAGATGGAAT ACTGGAAGCA TACCAATCAC GGAAGGATG CTGGTAATGA
 6841 GCTGAGAGCC TCTATGGCTA CTATGCGAA CAAGCTCACT AATAAGACTG CCACAATCT
 6901 TGCCATATGC ACATTAATG TGCTATGTTC AAGTATCTT CATGAGATAA TTCACTTGA
 6961 GTTTCTCTCT GGTCTTATGA ATCTGATGA GTCACAGCAA GGCAATATTC AGCCTATCAT
 7021 AGAATCATTAA AATCATTTGA TTGCTTGGC CAACCGATT CTATATAATG TTGCAATAGT

7081 AATTCCTCTT AAAATTGACA GTATCGAAAC TGTAATACTC TCTGCTTTAA AAGATATGCA
 7141 CACCGGGAGT ATGTCCAATG CCAACTGCAC GCCAGGAAAT CTGCTTCTGC ATGATGCAGC
 7201 ATACATCAAT GGAATAACA AATTCCTTGT ACTTGAATCA TACAATGGGA CGCCTAATA
 7261 TGGCCTCTC CTAATAATAC CAGCTTTAT CCCCTCAGCA ACATCTCCCC ATGGGTGTAC
 7321 TAGAATACCA TCATTTTCAC TCATCAAGAC CCATTGGTGT TACACTCACA ATGTAATGCT
 7381 TGGAGATTGT CTGATTTCAC CGGCATCTAA CCAGTATTTA TCAATGGGGA TAATACAACA
 7441 ATCTGCTGCA GGGTTCCAA TTTTCAGGAC TATGAAACC ATTTACCTAA GTGATGGAAT
 7501 CAATCGCAAA AGCTGTTTCA CACTGCTAT ACCAGGAGGT TGTGCTTGT ATTGCTATGT
 7561 AGCTACAAGG TCTGAAAAG ARGATTATGC CACGACTGAT CTAGCTGAAC TGAGACTTGC
 7621 TTTCTATTAT TATAATGATA CCTTTATTGA AAGAGTCATA TCTCTCCAA ATACAACAGG
 7681 CGAGTGGGCC ACAATCAACC CTGCACTCGG AAGCGGATC TATCATCTAG GCTTTATCTT
 7741 ATTTCTGTGA TATGGTGGTC TCATAAATGG GACTACTTCT TACAATGAGC AGTCCCTCAG
 7801 CTATTTTATC CCAAAACATC CCAACAATAA TTGTGCCGGT AACTCCAGCA AACAGGCTGC
 7861 AATAGCACGG AGTTCTCTAT TCATCCGTTA TCACTCAAC AGGTTAATTC AGAGTGTCTGT
 7921 TCTTATTTGT CCATTGCTCG ACATGCATAC AGAAGAGTGT AATCTAGTTA TGTTTAACAA
 7981 TTCCCAGTCT ATGATGGGTG CAGAAGTAG GCTCTATGTT ATGGTAATA ATTTGTATTA
 8041 TTATCAACGC AGTTCCTCTT GGTGCTCTCG ATCGCTCTTT TACAGGATCA ATACAACAGG
 8101 TTCTAAAGGA ATTCTCCGA TCATTGAGGC TCAATGGGTA CCGTCTATC AAGTTCCTCG
 8161 TCCTGGAGTC ATGCCATGCA ATGCAACAAG TTTTGTCCCT GCTAATTGCA TCACAGGGGT
 8221 GTACGCAGAT GTGTGGCCGC TTAATGATCC AGAAGTCTATG TCAAGTAATG CTCTGAACCC
 8281 CAACATCGCA TTTGCTGGAG CCTTCTCAA AATAGAGTCC AACCGAATA ATCCCACATT
 8341 CTACACTGCA TCGGCTAACT CCTCTTAAA TACTACCGGA TTCAACACA CCAATCACAA
 8401 AGCAGCATAT ACATCTTCAA CCTGCTTAAA AACACTGGA ACCCAAAAAA TTTATTTGTT
 8461 AATAATAAAT GAAATGGGCT CATCTCTTTT AGGGGAGTTC CAAATAATAC CATTTTTAAG
 8521 GGAACATAAG CTTTAAATCAT AATTAACCAT AATATGCATC AATCTATCTA TAATACAAGT
 8581 ATATGATAAG TAATCAGCAA TCAGACAATA GACAAAAGGG AATATAAAAA AACTTAGGAG
 8641 CAAAGCGTGC TCGGGAATG GACACTGAAT CTAACAATGG CACTGTATCT GACATACTCT
 8701 ATCCTGAGTG TCACCTTAAC TCTCCTATCG TTAAGGTAA AATAGCACAA TTACACACTA
 8761 TTATGAGTCT ACCTCAGCCT TATGATATGG ATGACGACTC AATACTAGTT ATCACTAGAC
 8821 AGAAAATAAA ACTTAATAAA TTGATAAAA GACAACGATC TATTAGAAGA TTAATAATTA
 8881 TATTAACCTG AAAAGTGAAT GACTTAGGAA AATACACATT TATCAGATAT CCGAAATGT
 8941 CAAAAGAAAT GTTCAAAATTA TATATACCTG GTATTAACAG TAAAGTGACT GAATTAATAC
 9001 TAAAGCAGA TAGAACATAT AGTCAAAATGA CTGATGGATT AAGAGATCTA TGGATTAATG
 9061 TGCTATCAAA ATTAGCCTCA AAAAAATGAT GAAGCAATTA TGATCTTAAT GAAGAAATTA
 9121 AATAATATAT GAAAGTTCAC ACAACCTATA AATCAGATAA ATGGTATAAT CCATTCAAAA
 9181 CATGGTTTAC TATCAAGTAT GATATGAGAA GATTACAAA AGCTCGAAAT GAGATCACTT
 9241 TTAATGTTGG GAAGGATTAT AACTTGTAG AAGACCAGAA GAATTTCTTA TTGATACATC
 9301 CAGAAATGGT TTTGATATTA GATAAACAAA ACTATAATGG TTATCTAATT ACTCCTGAAT
 9361 TAGTATTGAT GTATTGTGAC GTAGTCGAAG GCCGATGGAA TATAAGTGCA TGTGCTAAGT
 9421 TAGATCCAAA ATTACAATCT ATGTATCAGA AAGGTAATAA CCTGTGGGAA GTGATAGATA
 9481 AATTGTTTCC AATTATGGGA GAAAAGACAT TTGATGTGAT ATCGTTATTA GAACCACTTG
 9541 CATTAATCCTT AATTCAAACT CATGATCCTG TTAACAACCT AAGAGGAGCT TTTTAAATC
 9601 ATGTGTTTAC CGAGATGGAA TTAATATTG AATCTAGAGA ATCGATTAG GAATTTCTGA
 9661 GTGTAGATTA CATTGATAAA ATTTTAGATA TATTTAATAA CTCTACAATA GATGAAATAC
 9721 CAGAGATTTT CTCTTTTTTT AGAACATTTG GGCATCCTCC ATTAGAAGCT AGTATTGCAG
 9781 CAGAAAAGGT TAGAAAATAT ATGTATATTG GAAAACRAAT AAAATTTGAC ACTATTAATA
 9841 AATGTCAATG TATCTTCTGT ACAATAATAA TTAACGGATA TAGAGAGAGG CATGGTGGAC
 9901 AGTGGCCTCC TGTGACATTA CCTGATCATG CACACGAATT CATCATAAAT GCTTACGGTT
 9961 CAAACTCTGC GATATCATAT GAAATGCTG TTGATTATTA CCAGAGCTTT ATAGGAATAA
 10021 AATTCARATA ATTCATAGAG CCTCAGTTAG ATGAGGATTT GACAAATTTAT ATGAAAGATA
 10081 AAGCATTTAT TCCAAAAAAA TCAARTGGG ACACAGTTTA TCCTGCATCT AATTTACTGT
 10141 ACCGTACTAA CGCATCCAAC GAATCAGCAA GATTAGTTGA AGTATTTATA GCAGATAGTA
 10201 AATTTGATCC TCATCAGATA TTGGATTATG TAGAATCTGG GGACTGGTTA GATGATCCAG
 10261 AATTTAATAT TTCTTATAGT CTTAAGAAA AAGAGATCAA ACAGGAAGGT AGACTCTTTG
 10321 CAAAATGAC ATACAAAATG AGAGCTACAC AAGTTTATC AGAGACACTA CTTGCAATAA
 10381 ACHTAGGAAA ATCTTTCAA GAAATGGGA TGGTGAAGGG AGAGATTGAA TTACTTAAGA
 10441 GATTACAAC CATATCAATA TCAGAGTTC CACGGTATAA TGAAGTGTAC AATAATCTA
 10501 AAAGCCATAC AGATGACCTT AAAACCTACA ATAAAATAAG TAATCTAAT TTGTCTCTA
 10561 ATCAGAAATC AAGAAATTT GAATTCAGT CAACGGATAT CTACAATGAT GGATACGAGA
 10621 CTGTGAGCTG TTTCCFAACA ACAGATCTCA AAAAATCTG TCTTAATTGG AGATATGAAT
 10681 CAACAGCTCT ATTTGGAGAA ACTTGCAACC AAATATTGG ATTAATAAAA TTGTTAAT
 10741 GTTTACACCC TCGTCTTGAA GGAAGTACAA TCTATGTAGG TGATCTTAC TGCTCTCAT

10801 CAGATAAAGA ACATATATCA TTAGAGGATC ACCCTGATTC TGGTTTTTAC GTTCATAACC
 10861 CAAGAGGGGG TATAGAAGGA TTTTGTCAAA AATTATGGAC ACTCATATCT ATAAGTGCAA
 10921 TACACTTAGC AGCTGTTAGA ATAGGCGTGA GGGTGACTGC AATGGTTCAA GGAGACAATC
 10981 AAGCTATAGC TGTAAACCACA AGAGTACCCA ACRATTATGA CTACAGAGTT AAGAAAGAGA
 11041 TAGTTTATAA AGATGTAGTG AGATTTTTTG ATTCATTAAAG AGAAGTGATG GATGATCTAG
 11101 GTCATGAAC TAAATTAAT GAAACGATTA TAAGTAGCAA GATGTTTATA TATAGCAAAA
 11161 GAATCTATTA TGATGGGAGA ATCTTCTCCT AAGCTCTAAA AGCATTATCT AGATGTGTCT
 11221 TCTGGTCAGA GACAGTAATA GACGAAACAA GATCAGCATC TTCAAATITG GCAACATCAT
 11281 TTGCAAAAGC AATTGAGAA TGGTATTCCAC CTGTCTTAGG ATATGCATGC TCAATTTTTA
 11341 AGAATATCCA ACAACTATAT ATTGCCCTTG GGATGAATAT CAATCCAAC TAAACACAGA
 11401 ATATCAGAGA TCAGTATTTT AGGAATCCAA ATTGGATGCA ATATGCCCTT TAAATACCTG
 11461 CTAGTGTGG GGGATTCAAT TACATGGCCA TGTCAGATG TTTTGTAAAG AATATTGGTG
 11521 ATCCATCAGT TGCCGCATTG GCTGATATTA AAAGATTAT TAAGGCCAAT CTATTAGACC
 11581 GAAGTGTCT TTATAGGATT ATGAATCAAG AACCCAGTGA GTCATCTTTT TTGGACTGGG
 11641 CTTCAGATCC ATATTATGC AATTTACCAC AATCTCAAAA TATAACCACC ATGATAAAAA
 11701 ATATAACAGC AAGGAATGTA TTACAAGATT CACCBAATCC ATTATTATCT GGATTATTCA
 11761 CAAATACAAT GATAGAAGAA GATGAAGAAT TAGCTGAGTT CCTGATGGAC AGGAAGGTAA
 11821 TTTCCCTAG AGTTGCACAT GATATTCTAG ATAATCTCT CACAGGAAT AGAAATGCCA
 11881 TAGCTGGAAT GTTAGATAC ACAAATCAC TAATTCGGGT TGGCATAAAT AGAGGAGGAC
 11941 TGACATATAG TTTGTTGAGG AAAATCAGTA ATTACGATCT AGTACAATAT GAAACACTAA
 12001 GTAGACTTT GCGACTAAT GTAAGTGATA AAATCAAGTA TGAAGATATG TGTTCGGTAG
 12061 ACCTGGCCAT AGCATTGCCG CAAAAGATGT GGATTCATTT ATCAGGAGGA AGGATGATA
 12121 GTGGACTTGA AACGCCTGAC CCATTAGAAT TACTATCTGG GGTAGTAATA ACAGGATCAG
 12181 AACATTGTAA AATATGTTAT TCTTCAGATG GCACAAACCC ATATACTTGG ATGATTTTAC
 12241 CCGGTAATAT CAAAATAGGA TCAGCAGAAA CAGGTATATC GTCATTAAGA GTTCCCTTAT
 12301 TTGGATCAGT CACTGATGAA AGATCTGAAG CACAATTAGG ATATATCAAG AATCTTAGTA
 12361 AACCTGCAAA AGCCGCAATA AGAATAGCAA TGATATATAC ATGGGCATTT GGTAAATGATG
 12421 AGATATCTTG GATGGAAGCC TCACAGATAG CACAACACG TGCAAAATTT ACACATGATA
 12481 GTCTCAAAAT TTTAACACCG GTAGCTACAT CAACAATTT ATCACACAGA TTAAGGATA
 12541 CTGCAACTCA GATGAATTC TCCAGTACAT CATTGATCAG AGTCAGCAGA TTCATAACAA
 12601 TGTCCAATGA TAACATGTCT ATCAAGAAG CTAATGAAAC CAAAGATACT AATCTTATTT
 12661 ATCAACAAAT AATGTTAACA GGATTAAGTG TTTTCGAATA TTTATTAGA TTAAGGAAAA
 12721 CCACAGGACA CAACCCATA GTTATGCATC TGCACATAGA AGATGAGTGT TGTATTAAAG
 12781 AAAGTTTTTA TGATGAACAT ATTAATCCAG AGTCTACATT AGAATTAAT CGATATCCCTG
 12841 AAAGTAATGA ATTTATTTAT GATAAAGACC CACTCAAGA TGTGGACTTA TCAAACTTA
 12901 TGGTTATTA AGACCAATCT TACACAATTG ATATGAATTA TTGGGATGAT ACTGACATCA
 12961 TACATGCAAT TTCAATATGT ACTGCAATTA CAATAGCAGA TACTATGTCA CAATTAGATC
 13021 GAGATAATTT AAAAGAGATA ATAGTTATTG CAAATGATGA TGATATTAAT AGCTTAATCA
 13081 CTGAATTTTT GACTCTTGAC ATACTTGTAT TTTCAAGAC ATTTGGTGGG TTATTAGTAA
 13141 ATCAATTTGC ATCACTCTT TATAGTCTAA AAATAGAAGG TAGGGATCTC ATTTGGGATTT
 13201 ATATAATGAG AACACTGAGA GATACTTCCC ATTCATATTT AAAAGTATTA TCTAATGCAT
 13261 TATCTCATCC TAAAGTATTC AAGAGGTCTT GGGATGTGG ACTTTTAAAC CCTATTATG
 13321 GTCCATAATC TGCTAGTCAA GACCAGATAA AACTTGCCCT ATCTATATGT GAATATTCAC
 13381 TAGATCTATT TATGAGAGAA TGGTTGAATG GTGTATCACT TGAAATATAC ATTTGTGACA
 13441 GCGATATGGA AGTTGCAAT GATAGGAAAC AAGCCTTTAT TTCTAGACAC CTTTCAATTT
 13501 TTTGTTGTTT AGCAGAAAT GCATCTTTC GACCTAACCT GTTAAACTTA ACATACTTGG
 13561 AGAGACTTGA TCTATTGAAA CAATATCTTG AATTAATAT TAAAGAAGAC CCTACTCTTA
 13621 AATATGTRCA AATATCTGGA TPATTAATTA AATCGTCCC ATCAACTGTA ACATACGTAA
 13681 GAAAGACTGC AATCAATAT CTAAGGATTC GCGGTATTAG TCCACCTGAG GTAATTGATG
 13741 ATTTGGATCC GGTAGAAGAT GAAAATATGC TGGATAACAT TTCAAAAC ATAAATGATA
 13801 ACTGTAATAA AGATAATAAA GGGAAATAAA TTAACAATTT CTGGGGACTA GCCTTAAACA
 13861 ACTATCAAGT CCTTAAATC AGATCTATAA CAAGTGATTC TGATGATAAT GATAGACTAG
 13921 ATGCTAATAC AAGTGGTTG ACATTTCTC AAGGAGGAA TTATCTATCG CATCAATTGA
 13981 GATTATTCCG AATCAACAGC ACTAGTTGTC TGAAGCTCT TGAGTTATCA CAAATTTTAA
 14041 TGAAGGAAGT CAATAAAGAC AAGSACAGC TCTTCTGGG AGAAGSAGCA GGACTATGCG
 14101 TAGCATGTTA TGATGCCACA TTAGGACCTG CAGTTAATTA TTATTAATCA GGTTTGAATA
 14161 TAACAGATGT AATTGGTCAA CGAGAATTGA AAATATTCC TTCAGAGGTA TCATTAGTAG
 14221 GTAAAAAATT AGGAAATGTG ACACAGATTC TTAACAGGGT AAAAGTACTG TTCAAATGGGA
 14281 ATCCATAATC AACATGGATA GGAATATG AATGTAGAG CTTAATATG AGTGAATTAA
 14341 ATGATAAGTC CATTTGATTA GTACATTGTG ATATGGAAGG AGCTATCGGT AAATCAGAA
 14401 AAACGTCTCT ACATGAACAT TATAGTGTTA TAAGAAATAC ATACTCTGAT GGGGATGATG
 14461 ATGTTGTTTT AGTTTCCAAA ATTATACCTA CAATCACTCC GAATTTGGTCT AGAATACTTT

14521 ATCTATATAA ATTATATTGG AAAGATGTAA GTATAATATC ACTCAAAACT TCTAATCCTG
14581 CATCAACAGA ATTATATCTA ATTCGAAAG ATGCATATTG TACTATAATG GAACCTAGTG
14641 AAATTGTTTT ATCAAACTT AAAAGATTGT CACTCTTGGG AGAAAATAAT CTATTAATAAT
14701 GGATCATTTT ATCAAAGAG AGGAATAATG AATGGTTACA TCATGAAATC AAAGAAGGAG
14761 AAAGAGATTA TCGAATCATG AGACCATATC ATATGGCACT ACAATCTTT GGATTTCAAA
14821 TCAATTTAAA TCATCTGGCG AAAGAATTTT TATCAACCCC AGATCTGACT AATATCAACA
14881 ATATAATCCA AAGTTTTAG CGAACATAA AGGATGTTTT ATTTGAATGG ATTAATATAA
14941 CTCATGATGA TAAGAGACAT AATTAGGCG GAAGATATAA CATATTCOCA CTGAAAAATA
15001 AGGGAAGTT AAGACTGCTA TCGAGAAGAC TAGTATTAG TTGATTTCA TTATCATTAT
15061 CGACTCGATT ACTTACAGGT CGCTTCTTG ATGAAAAAT TGACATAGA GCACAGACTG
15121 GATATGTATC ATTAGCTGAT ACTGATTTAG AATCATTAA GTTATTGTCG AAAACATCA
15181 TTAAGAATTA CAGAGAGTGT ATAGGATCAA TATCATATTG GTTCTAACC AAAGAAGTTA
15241 AAATACTTAT GAAATTGATC GGTGGTGCTA AATTATTAGG AATTCOCAGA CAATATAAAG
15301 AACCCGAGA CCAGTTATTA GAAAAC TACA ATCAACATGA TGAATTTGAT ATCGATTA
15361 ACATAAATAC AATGAAGATA TATCCTAACC TTTATCTTTA AGCCTAGGAA TAGACAAAAA
15421 GTAAGAAAA CATGTAATAT ATATATACCA AACAGAGTTC TTCTCTGTT TGGT

The cDNA engineering was designed so that the final PIV3-2 antigenomes conformed to the rule of six (Calain et al., *J. Virol.* 67:4822-30, 1993; Durbin et al., *Virology* 234:74-83, 1997, each incorporated herein by reference). The PIV3-2 insert in pFLC.PIV32TM is 15498 nt in length, and that in pFLC.PIV32CT is 15474 nt in length.

- 5 These total lengths do not include two 5'-terminal G residues contributed by the T7 promoter, because it is assumed that they are removed during recovery.

Transfection and recovery of recombinant chimeric PIV3-PIV2 viruses

- HEp-2 cell monolayers were grown to confluence in six-well plates, and transfections were performed essentially as described (Tao et al., 72:2955-2961, 1998, incorporated herein by reference). The HEp-2 monolayer in one well was transfected with 5 μ g PIV3-PIV2 antigenomic cDNA and three support plasmids, 0.2 μ g pTM(N), 0.2 μ g pTM(PnoC), 0.1 μ g pTM(L) in 0.2 ml of MEM containing 12 μ l LipofectACE (Life Technologies). The cells were infected simultaneously with MVA-T7 at a multiplicity of infection (MOI) of 3 in 0.8 ml of serum-free MEM containing 50 μ g/ml gentamicin and 2 mM glutamine. The chimeric antigenomic cDNA pFLC.2G+.hc (Tao et al., *J. Virol.* 72:2955-2961, 1998), was transfected in parallel as a positive control. After incubation at 15 32°C for 12 hours, the transfection medium was replaced with 1.5 ml of fresh serum-free MEM supplemented with 50 μ g/ml gentamicin and 2 mM glutamine. Transfected cells were incubated at 32°C for two additional days. Gamma-irradiated porcine trypsin (p-trypsin; 20 T1311, Sigma, St Louis, MO) was added to a final concentration of 0.5 μ g/ml on day 3 post transfection. Cell culture supernatants were harvested and passaged (referred to as passage 1) onto fresh Vero cell monolayers in T25 flasks. After overnight adsorption, the transfection harvest was replaced with fresh VP-SFM supplemented with 0.5 μ g/ml p-trypsin. Cultures from passage 1 were incubated at 32°C for 4 days, and the amplified virus 25 was harvested and further passaged on Vero cells (referred to as passage 2) for another 4 days at 32°C in the presence of 0.5 μ g/ml p-trypsin. The presence of viruses in the passage 2 cultures was determined by hemadsorption with 0.2% guinea pig red blood cells (RBCs). Viruses were further purified by three consecutive terminal dilutions performed using Vero cells maintained in VP-SFM supplemented with 2 mM glutamine, 50 μ g/ml gentamicin, and 30 0.5 μ g/ml p-trypsin. Following the third terminal dilution, virus was further amplified three times on Vero cells, and this virus suspension was used for further characterization *in vitro* and *in vivo*.

Confirmation of the chimeric nature of vRNA using sequencing and restriction analysis of PCR products

For analysis of the genetic structure of vRNAs, the recombinant PIVs were amplified on LLC-MK2 cells and concentrated. vRNA was extracted from the viral pellets and reverse transcribed using the Superscript Preamplification System. RT-PCR was performed using the Advantage cDNA synthesis kit and primer pairs specific to PIV2 or PIV3 (21, 22 or 23, 24 in Table 22). RT-PCR products were either analyzed by restriction digestion or gel purified and analyzed by sequencing.

Replication of PIVs in LLC-MK2 cells

Growth of the PIV viruses in tissue culture was evaluated by infecting confluent LLC-MK2 cell monolayers on six-well plates in triplicate at an MOI of 0.01. The inoculum was removed after adsorption for 1 hour at 32°C. Cells were washed 3 times with serum-free OptiMEM I, fed with 2 ml/well of OptiMEM I supplemented with 50 µg/ml gentamicin and 0.5 µg/ml p-trypsin, and incubated at 32°C. At each 24 hour interval, a 0.5 ml aliquot of medium was removed from each well and flash-frozen, and 0.5 ml fresh medium with p-trypsin was added to the cultures. The virus in the aliquots was titrated at 32°C on LLC-MK2 cell monolayers using fluid overlay as previously described (Tao et al., *J. Virol.* 72:2955-2961, 1998, incorporated herein by reference), and the endpoint of the titration was determined by hemadsorption, and the titers are expressed as log₁₀TCID₅₀/ml.

20 Replication of recombinant chimeric PIV3-PIV2 viruses at various temperatures

Viruses were serially diluted in 1X L15 supplemented with 2 mM glutamine and 0.5 µg/ml p-trypsin. Diluted viruses were used to infect LLC-MK2 monolayers in 96 well plates. Infected plates were incubated at various temperatures for 7 days as described (Skiadopoulos et al., *Vaccine* 18:503-510, 1999, incorporated herein by reference). Virus titers were determined as above.

Replication, immunogenicity, and protective efficacy of recombinant chimeric PIV3-PIV2 viruses in the respiratory tract of hamsters

Golden Syrian hamsters in groups of six were inoculated intranasally with 10^{5.3} TCID₅₀ of recombinant or biologically-derived viruses. Four days after inoculation, hamsters were sacrificed and their lungs and nasal turbinates were harvested and prepared

for quantitation of virus (Skiadopoulos et al., *Vaccine* 18:503-510, 1999, incorporated herein by reference). The titers are expressed as mean \log_{10} TCID₅₀/gram of tissue for each group of six hamsters.

5 Hamsters in groups of 12 were infected intranasally with $10^{5.3}$ TCID₅₀ of viruses on day 0, and six hamsters from each group were challenged four weeks later with 10^6 TCID₅₀ of PIV1 or 10^6 TCID₅₀ of PIV2. Hamsters were sacrificed 4 days after challenge and their lungs and nasal turbinates were harvested. Challenge virus titers in the harvested tissue was determined as previously described (Tao et al., *J. Virol.* 72:2955-2961, 1998, incorporated herein by reference). The virus titers are expressed as mean \log_{10} TCID₅₀/gram
10 of tissue for each group of six hamsters. Serum samples were collected three days prior to inoculation and on day 28, and hemagglutination-inhibition antibody (HAI) titers against PIV1, PIV2, and PIV3 were determined as previously described (van Wyke Coelingh et al., *Virology* 143:569-582, 1985, incorporated herein by reference). The titers are expressed as reciprocal mean \log_2 .

15 **Replication, immunogenicity, and protective efficacy of recombinant chimeric PIV3-PIV2 viruses in African green monkeys (AGMs)**

AGMs in groups of 4 were infected intranasally and intratracheally with 10^5 TCID₅₀ of virus at each site on day 0. Nasal/throat (NT) swab specimens and tracheal lavages were collected for 12 and 5 days, respectively, as previously described (van Wyke
20 Coelingh et al., *Virology* 143:569-582, 1985). On day 29, immunized AGMs were challenged intranasally and intratracheally with 10^5 TCID₅₀ of PIV2/V94 at each site. NT swab specimens and tracheal lavages were collected for 10 and 5 days, respectively. Pre-immunization, post-immunization, and post challenge serum samples were collected on days -3, 28, and 60, respectively. Virus titers in the NT swab specimens and in tracheal lavages
25 were determined as previously described (Tao et al., *J. Virol.* 72:2955-2961, 1998). Titers are expressed as \log_{10} TCID₅₀/ml. Serum neutralizing antibody titers against PIV1 and PIV2 were determined as previously described (van Wyke Coelingh et al., *Virology* 143:569-582, 1985), and the titers are expressed as reciprocal mean \log_2 .

30 **Replication and immunogenicity of recombinant chimeric PIV3-PIV2 viruses in chimpanzees**

Chimpanzees in groups of 4 were infected intranasally and intratracheally with 10^5 TCID₅₀ of PIV2/V94 or rPIV3-2TM on day 0 as previously described (Whitehead et al., *J. Virol.* 72:4467-4471, 1998, incorporated herein by reference). NT swab specimens were collected daily for 12 days and tracheal lavages were obtained on days 2, 4, 6, 8, and 10. Virus titers in the specimens were determined as previously described (Tao et al., *J. Virol.* 72:2955-2961, 1998, incorporated herein by reference). The peak virus titers are expressed as mean log₁₀TCID₅₀/ml. Pre-immunization and post-immunization serum samples were collected on days -3 and 28, respectively. Serum neutralizing antibody titers against PIV1 and PIV2 were determined as previously described (van Wyke Coelingh et al., *Virology* 143:569-582, 1985, incorporated herein by reference), and the titers are expressed as reciprocal mean log₂.

Viable recombinant chimeric virus was not recovered from PIV3-PIV2 chimeric cDNA encoding the complete PIV2 F and HN proteins

The construction of the PIV3-PIV2 chimeric cDNA, in which the F and HN ORFs of the JS wild type PIV3 were replaced by those of PIV2/V94, is described above and summarized in Figure 17. The final plasmid construct, pFLC.PIV32hc (Figure 17), encodes a PIV3-PIV2 chimeric antigenomic RNA of 15492 nt, which conforms to the rule of six.

HEp-2 cell monolayers were transfected with pFLC.PIV32hc along with the three support plasmids pTM(N), pTM(PnoC), and pTM(L) using LipofectACE, and the cells were simultaneously infected with MVA-T7 as previously described (Tao et al., *J. Virol.* 72:2955-2961, 1998, incorporated herein by reference). Virus was not recovered from several initial transfections using pFLC.PIV32hc, while chimeric viruses were recovered from all the transfections using control plasmid pFLC.2G+.hc.

Consistent with these results is the possibility that a mutation occurred outside of the 4 kb BspEI-SpeI segment of pFLC.PIV32hc that prevented the recovery of rPIV3-2 virus from cells transfected with this cDNA clone. To examine this possibility, the BspEI-SpeI fragments of p38'ΔPIV31hc and p38'ΔPIV32hc were exchanged. The regenerated p38'ΔPIV31hc and p38'ΔPIV32hc were identical to those in Figure 17 except that the SpeI-SphI fragments containing PIV3 L gene sequences were exchanged. The BspEI-SphI fragments of these two regenerated cDNAs were introduced into the BspEI-SphI window of a PIV3 full-length clone, p3/7-(131)2G+, in five separate independent ligations to give 10 pFLC.2G+.hc and pFLC.PIV32hc clones (2 clones selected from each ligation),

respectively. (Note that the PIV3 sequences outside of the BspEI-SphI window of p3/7-(131)2G+, pFLC.2G+.hc, and pFLC.PIV32hc are identical). Thus, this would have replaced any PIV3 backbone sequence which might have acquired a spurious mutation with sequence known to be functional. Furthermore, the functionality of the backbone was reevaluated in parallel. None of the 10 pFLC.PIV32hc cDNA clones yielded viable virus, but each of the 10 pFLC.2G+.hc cDNA clones yielded viable virus. Virus was not recovered from pFLC.PIV32hc despite passaging the transfection harvest in a fashion similar to that used successfully to recover the highly defective PIV3 C-knock out recombinant (Durbin et al., *Virology* 261:319-30, 1999, incorporated herein by reference). Since each of the unique components used to generate the pFLC.PIV32hc was used to successfully generate other recombinant viruses except the cytoplasmic tail domains of F and HN, it is highly unlikely that errors in the cDNA account for the failure to yield recombinant virus in this case. Rather, the favored interpretation is that the full-length PIV2 F and HN glycoproteins are not compatible with one or more of the PIV3 proteins needed for virus growth.

15 Recovery of chimeric viruses from PIV3-PIV2 chimeric cDNAs encoding the chimeric PIV3-PIV2 F and HN proteins

Using two other strategies, new chimeric PIV3-PIV2 antigenomic cDNAs were constructed, in which the ectodomain or the ectodomain and the transmembrane domain of PIV3 F and HN glycoproteins were replaced by their PIV2 counterparts. The construction of the four full-length cDNAs, namely pFLC.PIV32TM, pFLC.PIV32TMcp45, pFLC.PIV32CT, and pFLC.PIV32CTcp45, is described above and summarized in Figures 18, 19, and 20. The PIV3-2 inserts in the final plasmids pFLC.PIV32TM and pFLC.PIV32CT, in which the F and HN genes encoded chimeric glycoproteins, were 15498 nt and 15474 nt in length, respectively, and conformed to the rule of six (Calain et al., *J Virol.* 67:4822-30, 1993; Durbin et al., *Virology* 234:74-83, 1997, each incorporated herein by reference). The authenticity of those four constructs was confirmed by sequencing of the BspEI-SphI region and by restriction analysis.

Recombinant chimeric viruses were recovered from full-length clones pFLC.PIV32TM, pFLC.PIV32CT, pFLC.PIV32TMcp45, or pFLC.PIV32CTcp45 and were designated rPIV3-2TM, rPIV3-2CT, rPIV3-2TMcp45, and rPIV3-2CTcp45, respectively. These viruses were biologically cloned by 3 consecutive terminal dilutions on Vero cells and then amplified three times in Vero cells.

Genetic characterization of recombinant chimeric PIV3-PIV2 viruses

The biologically-cloned chimeric PIV3-PIV2 viruses, rPIV3-2TM, rPIV3-2CT, rPIV3-2TMcp45, and rPIV3-2CTcp45, were propagated on LLC-MK2 cells and then concentrated. Viral RNAs extracted from pelleted viruses were used in RT-PCR

5 amplification of specific gene segments using primer pairs specific to PIV2 or PIV3 (21, 22 or 23, 24 in Table 22). The restriction enzyme digestion patterns of the RT-PCR products amplified with PIV2 specific primer pairs from rPIV3-2TM, rPIV3-2CT, rPIV3-2TMcp45, and rPIV3-2CTcp45, were each distinct from that derived from PIV2/V94, and their patterns, using EcoRI, MfeI, NcoI, or PpuMI, were those expected from the designed cDNA.

10 Nucleotide sequences for the 8 different PIV3-PIV2 junctions in F and HN genes of rPIV3-2TM and rPIV3-2CT are given in Figure 20. Also, the cp45 markers present in rPIV3-2TMcp45 and rPIV3-2CTcp45, except those in the 3'-leader region and the gene start of NP, were verified with RT-PCR and restriction enzyme digestion as previously described (Skiadopoulos et al., *J. Virol.* 73:1374-81, 1999, incorporated herein by reference). These

15 results confirmed the chimeric nature of the recovered PIV3-PIV2 viruses as well as the presence of the introduced cp45 mutations.

PIV3-PIV2 recombinant chimeric viruses replicate efficiently in LLC-MK2 cells *in vitro*

The kinetics and magnitude of replication *in vitro* of the PIV3-PIV2

20 recombinant chimeric viruses were assessed by multicycle replication in LLC-MK2 cells (Figure 21). LLC-MK2 cell monolayer cultures in six-well plates were infected in triplicate with rPIV3-2TM, rPIV3-2CT, rPIV3-2TMcp45, or rPIV3-2CTcp45 at an MOI of 0.01 in the presence of p-trypsin (0.5 µg/ml). Samples were removed from culture supernate at 24 hour intervals for 6 days. Each of the recombinant chimeric viruses, except rPIV3-2CTcp45

25 (clone 2A1), replicated at the same rate and to a similar level as their PIV2/V94 parent virus indicating that PIV3-PIV2 chimerization of F and HN proteins did not alter the rates of growth of the recombinant chimeric viruses, and all reached a titer of 10⁷ TCID₅₀/ml or higher. Only the rPIV3-2CTcp45 grew slightly faster in each of two experiments and reached its peak titer earlier than PIV2/V94. This accelerated growth pattern was likely a

30 result of an unidentified mutation in this clone since a sister clone failed to exhibit this growth pattern. rPIV3-2CTcp45 clone 2A1 was used in the studies described below.

The level of temperature sensitivity of rPIV3-2 chimeric viruses and their *cp45* derivatives

The level of temperature sensitivity of replication of PIV3-PIV2 recombinant chimeric viruses was tested to determine if rPIV3-2TM and rPIV3-2CT viruses exhibit a *ts* phenotype and to determine if the acquisition of the 12 *cp45* mutations by these viruses specified a level of temperature sensitivity characteristic of *cp45* derivatives bearing these 12 PIV3 *cp45* mutations (Skiadopoulos et al., J Virol. 73:1374-81, 1999, incorporated herein by reference). The level of temperature sensitivity of the virus was determined in LLC-MK2 cell monolayers as previously described (Skiadopoulos et al., Vaccine 18:503-510, 1999, incorporated herein by reference) (Table 26). The titer of rPIV3-2TM and rPIV3-2CT was fairly constant at permissive temperature (32°C) and the various restrictive temperatures tested indicating these recombinants were *ts+*. In contrast, their *cp45* derivatives, rPIV3-2TM*cp45* and rPIV3-2CT*cp45*, were *ts* and the level of temperature sensitivity was similar to that of rPIV3-1*cp45*, the chimeric PIV3-PIV1 virus carrying the complete PIV1 F and HN glycoproteins and the same set of 12 *cp45* mutations. Thus the *in vitro* properties of rPIV3-2TM and rPIV3-2CT viruses and their *cp45* derivative are similar to those of rPIV3-1 and rPIV3-1*cp45*, respectively, suggesting that the *in vivo* properties of the rPIV3-2 and rPIV3-1 viruses would also be similar, but surprisingly this was not the case.

Table 26. The replication of rPIV3-2CT and rPIV3-2TM are not temperature sensitive in LLC-MK2 cells, whereas the inclusion of the *cp45* mutations confers the *cp45* temperature sensitive phenotype

Virus	Change in titer (log ₁₀) at various temperatures compared to that at 32° a b							
	Titer at 32° C ^c (log ₁₀ TCID ₅₀)	35° 0.3 ^e	36° 0.1	37° 0.1	38° (0.3) ^b	39° (0.4)	40° 0.4	
rPIV3/JS	7.9	0.3 ^e	0.1	0.1	(0.3) ^b	(0.4)	0.4	
PIV3 ^{cp45} ^e	7.8	0.5	0.3	1.3	<u>3.4</u> ^d	6.8	6.9	
PIV1/Wash64 ^e	8.5	1.5	1.1	1.4	0.6	0.5	0.9	
rPIV3-1	8.0	0.8	0.5	0.6	0.9	1.1	2.6	
rPIV3-1 ^{cp45}	8.0	0.5	0.4	<u>3.4</u> ^d	4.8	6.6	7.5	
PIV2/V9412 ^e	7.8	0.3	(0.1)	0.0	(0.4)	(0.4)	0.0	
rPIV3-2CT	6.9	0.3	0.3	0.6	(0.1)	0.6	0.4	
rPIV3-2TM	8.3	0.3	(0.1)	0.3	0.6	1.0	<u>2.1</u> ^d	
rPIV3-2CT ^{cp45}	8.0	0.8	(0.4)	<u>2.0</u> ^d	4.3	7.5	≥7.6	
rPIV3-2TM ^{cp45}	8.0	0.3	0.6	<u>2.4</u> ^d	5.4	7.5	≥7.6	

^a Data presented are means of two experiments.
^b Numbers not in parentheses represent titer decrease; numbers in parentheses represent titer increase.
^c Data at 35° were from one experiment only.
^d Values which are underlined represent the lowest temperature at which there was a 100-fold reduction of virus titer compared to the titer at permissive temperature (32°C). This restrictive temperature is referred to as the shut-off temperature.
^e Biologically-derived viruses.

rPIV3-2TM and rPIV3-2CT are attenuated, immunogenic, and highly protective in hamsters, and introduction of *cp45* mutations results in highly attenuated and less protective viruses

5 Hamsters in groups of six were inoculated intranasally with $10^{5.3}$ TCID₅₀ of
rPIV3-2TM, rPIV3-2CT, rPIV3-2TM_{cp45}, rPIV3-2CT_{cp45}, or control viruses. It was
previously seen that rPIV3-1 virus replicated in the upper and lower respiratory tract of
hamsters like that of its PIV3 and PIV1 parents (Skiadopoulos et al., Vaccine 18:503-510,
1999; Tao et al., J. Virol. 72:2955-2961, 1998, each incorporated herein by reference). PIV2
10 virus replicates efficiently in hamsters, but rPIV3-2TM and rPIV3-2CT viruses each
replicated to a 50- to 100-fold lower titer than their PIV2 and PIV3 parents in the upper
respiratory tract and to a 320- to 2000-fold lower titer in the lower respiratory tract (Table
27). This indicates that the chimeric PIV3-PIV2 F and HN glycoproteins specify an
unexpected attenuation phenotype in hamsters. rPIV3-2TM_{cp45} and rPIV3-2CT_{cp45}, the
15 derivatives carrying the *cp45* mutations, were 50- to 100-fold more attenuated than their
respective rPIV3-2 parents, with only barely detectable replication in the nasal turbinates,
and none in the lungs. These rPIV3-2_{cp45} viruses were clearly more attenuated than rPIV3-
1_{cp45}, exhibiting an additional 50-fold reduction of replication in the nasal turbinates. Thus,
the attenuating effects of the chimerization of F and HN glycoproteins and that specified by
cp45 mutations were additive.

20

Table 27. The rPIV3-2TM and rPIV3-2CT viruses, in contrast to rPIV3-1, are attenuated in the respiratory tract of hamsters and importation of the *cp45* mutations resulted in further attenuation.

Group ^c Virus ^a	Virus titers in the indicated tissue (log ₁₀ TCID ₅₀ /g ± S.E.) ^b [Duncan			
	NT	log ₁₀ titer reduction	Lung	log ₁₀ titer reduction
rPIV3/JS	5.9 ± 0.1[AB]	0	6.5 ± 0.1[A]	0
rPIV3 _{cp45}	4.5 ± 0.2[C]	1.4 ^c	1.8 ± 0.2[E]	4.7 ^c
PIV1/Wash64 ^d	5.7 ± 0.1[B]	-	5.5 ± 0.1[B]	-
rPIV3-1	6.4 ± 0.2[A]	0	6.6 ± 0.2[A]	0
rPIV3-1 _{cp45}	3.1 ± 0.1[D]	3.3 ^c	1.2 ± 0.0[F]	5.4 ^c
PIV2/V94 ^d	6.2 ± 0.2[A]	0	6.4 ± 0.2[A]	0
rPIV3-2CT	4.5 ± 0.4[C]	1.7 ^c	3.1 ± 0.1[D]	3.3 ^c
rPIV3-2TM	3.9 ± 0.3[C]	2.3 ^c	3.9 ± 0.4[C]	2.5 ^c
rPIV3-2CT _{cp45}	1.4 ± 0.1[E]	4.8 ^c	1.5 ± 0.2[E]	4.9 ^c
rPIV3-2TM _{cp45}	1.6 ± 0.2[E]	4.6 ^c	1.4 ± 0.1[E]	5.0 ^c

^a Hamsters in group of six were inoculated intranasally with 10^{3.5} TCID₅₀ of indicated virus on day 0.

^b Hamsters were sacrificed and their tissue samples harvested on day 4. The virus titer in hamster tissues was determined and the results are expressed as log₁₀TCID₅₀/g ± standard error (SE). NT = nasal turbinates.

^c The log₁₀ titer reduction values are derived by comparing: rPIV3_{cp45} against rPIV3/JS; rPIV3-1_{cp45} against rPIV3-1; each of the PIV3-PIV2 chimeras against PIV2/V94.

^d Biologically-derived viruses.

^e Grouping as analyzed by Duncan mult:range test.

To determine the immunogenicity and protective efficacy of the PIV3-PIV2 chimeric viruses, hamsters in groups of twelve were immunized with $10^{5.3}$ TCID₅₀ of rPIV3-2TM, rPIV3-2CT, rPIV3-2TMcp45, rPIV3-2CTcp45, or control viruses on day 0. Six of the hamsters from each group were challenged with 10^6 TCID₅₀ of PIV1 on day 29, and the remaining half were challenged with PIV2 on day 32. Hamsters were sacrificed 4 days after challenge and the lungs and nasal turbinates harvested. Serum samples were collected on day

-3 and day 28, and their HAI antibody titer against PIV1, PIV2, and PIV3 was determined. As shown in Table 28, despite their attenuated growth in hamsters, immunization with rPIV3-2TM or rPIV3-2CT each elicited a level of serum HAI antibody against PIV2 that was comparable to that induced by infection with wild type PIV2/V94. Immunization of hamsters with rPIV3-2TM and rPIV3-2CT resulted in complete restriction of the replication of PIV2 challenge virus. rPIV3-2TMcp45 and rPIV3-2CTcp45 failed to elicit a detectable serum antibody response, and immunization of hamsters with either of these two viruses resulted in only a 10- to 100-fold reduction of replication of the PIV2 challenge virus in the lower respiratory tract (Table 28).

Table 28. The rPIV3-2CT and rPIV3-2TM viruses are highly protective in hamsters against challenge with wild type PIV2, but not against PIV1

Immunizing virus ^a	HAI antibody titer ^b against indicated virus (reciprocal mean log ₂ ± SE)						Challenge virus titer ^c in indicated tissue (log ₁₀ TCID ₅₀ /g ± SE)					
	PIV1		PIV2		PIV3		PIV1		PIV2		Lung	
	NT	Lung	NT	Lung	NT	Lung	NT	Lung	NT	Lung	NT	Lung
rPIV3/S	≤1	≤1	≤1	≤1	10.2±0.1	6.2 ± 0.2	5.8 ± 0.1	5.9±0.2	5.9±0.2	5.7±0.2	5.9±0.2	5.7±0.2
rPIV3.cp45	≤1	≤1	≤1	≤1	8.6±0.2	5.9 ± 0.3	5.1 ± 0.3	5.6±0.2	5.6±0.2	4.5±0.7	5.6±0.2	4.5±0.7
PIV1	6.7±0.2	≤1	≤1	≤1	≤1.2 ± 0.0	1.3 ± 0.1	≤1.2 ± 0.0	6.1±0.2	6.1±0.2	6.2±0.3	6.1±0.2	6.2±0.3
rPIV3-1	6.4±0.2	≤1	≤1	≤1	≤1.2 ± 0.0	≤1.2 ± 0.0	≤1.2 ± 0.0	6.5±0.2	6.5±0.2	5.0±0.6	6.5±0.2	5.0±0.6
rPIV3-1.cp45	1.8±0.6	≤1	≤1	≤1	3.9 ± 0.4	5.9 ± 0.2	5.5 ± 0.1	6.2±0.2	6.2±0.2	4.5±0.6	6.2±0.2	4.5±0.6
PIV2	≤1	4.0±0.0	≤1	≤1	5.3 ± 0.1	5.2 ± 0.3	4.4 ± 0.3	≤1.2±0.0	≤1.2±0.0	≤1.2±0.0	≤1.2±0.0	≤1.2±0.0
rPIV3-2CT	≤1	3.6±0.8	≤1	≤1	5.9 ± 0.2	4.4 ± 0.3	5.7 ± 0.1	5.3±0.2	5.3±0.2	3.3±0.8	5.3±0.2	3.3±0.8
rPIV3-2TM	≤1	4.5±0.2	≤1	≤1	6.2 ± 0.2	5.8 ± 0.3	4.4 ± 0.3	5.5±0.2	5.5±0.2	3.7±0.7	5.5±0.2	3.7±0.7
rPIV3-2CT.cp45	≤1	≤1	≤1	≤1	≤1	≤1	≤1	≤1	≤1	≤1	≤1	≤1
rPIV3-2TM.cp45	≤1	≤1	≤1	≤1	≤1	≤1	≤1	≤1	≤1	≤1	≤1	≤1

^a Hamsters in groups of 12 were immunized intranasally with 10^{5.3} TCID₅₀ of the indicated virus on day 0.
^b Serum samples were collected two days before immunization and 28 days after immunization. They were tested for HAI antibody titer against the three PIVs, and the antibody titers are presented as reciprocal mean log₂ ± standard error (SE).
^c Six hamsters from each group were challenged intranasally with 10⁶ TCID₅₀ of PIV1 (on day 29) or PIV2 (on day 32). Hamster tissues were harvested 4 days after challenge, and the virus titer in indicated tissues are expressed as log₁₀TCID₅₀/g ± SE.

rPIV3-2TM and rPIV3-2CT are attenuated, immunogenic, and highly protective in AGMs, whereas introduction of *cp45* mutations results in highly attenuated and poorly protective viruses

Certain recombinant PIV3 and RSV viruses may exhibit different levels of
5 attenuation in rodents and primates (Skiadopoulos et al., *Vaccine* 18:503-510, 1999;
Skiadopoulos et al., *J. Virol.* 73:1374-81, 1999a; Skiadopoulos et al., *Virology* 272:225-34,
2000; Whitehead et al., *J. Virol.* 73:9773-9780, 1999, each incorporated herein by reference),
indicating that attenuation can be somewhat species specific. Therefore, the rPIV3-2 viruses
10 were evaluated for their level of replication and immunogenicity in AGMs. AGMs in groups
of four were intranasally and intratracheally administered 10^5 TCID₅₀ per site of rPIV3-
2TM, rPIV3-2CT, rPIV3-2TM*cp45*, rPIV3-2CT*cp45*, PIV2/V94, or rPIV3-1 on day 0. Virus
in the NT swab specimens (collected day 1 to 12) and tracheal lavages (collected on day 2, 4,
5, 8, and 10) were titered as previously described (van Wyke Coelingh et al., *Virology*
143:569-582, 1985, incorporated herein by reference). As shown in Table 29, rPIV3-2TM
15 and rPIV3-2CT were clearly attenuated in the respiratory tract of AGMs as indicated by a
peak titer of virus shedding lower in both the upper and lower respiratory tract than their
PIV2/V94 parent virus.

rPIV3-2TM*cp45* and rPIV3-2CT*cp45*, the derivatives carrying *cp45*
20 mutations, were detected at very low levels, if at all, in the NT swab and tracheal lavage
specimens suggesting that the attenuating effects of chimerization of the F and HN
glycoproteins and that specified by the *cp45* mutations were additive for AGMs as well as
for hamsters.

To determine whether immunization of AGMs with the PIV3-PIV2 chimeric
viruses is protective against PIV2 challenge, AGMs previously infected with a rPIV3-2 virus
25 were challenged with 10^5 TCID₅₀ of PIV2 on day 28 (Table 29). Virus present in the NT
swab specimens (collected day 29 to 38) and tracheal lavages fluids (collected on day 30, 32,
34, 36, and 38) was titered as previously described (Durbin et al., *Virology* 261:319-30,
1999, incorporated herein by reference). As shown in Table 29, immunization with rPIV3-
2TM and rPIV3-2CT induced a high level of restriction of the replication of PIV2/V94
30 challenge virus. In contrast, immunization of AGMs with rPIV3-2TM*cp45* and rPIV3-
2CT*cp45* failed to restrict the replication of PIV2/V94 challenge virus and these animals
developed very low levels of pre-challenge serum neutralizing antibody to PIV2. The

complete restriction of replication of PIV2/V94 challenge virus in rPIV3-2CT immunized AGMs was associated with a 2.5-fold greater level of pre-challenge serum antibody to PIV2 than that of rPIV3-2TM immunized AGMs which provided incomplete protection.

Table 29. The rPIV3-2CT or rPIV3-2TM viruses are attenuated for replication in the respiratory tract of African green monkeys, yet still induce resistance to challenge with wild type PIV2

Immunizing ^a virus	Mean peak titer ^b of immunizing virus in indicated site (log ₁₀ TCID ₅₀ /ml ± SE)		Serum neutralization antibody titer ^c against indicated virus (mean reciprocal log ₂ ± SE)		Mean peak titer ^d of PIV2/V94 challenge virus in indicated site (log ₁₀ TCID ₅₀ /ml ± SE)	
	NT	TL	PIV1	PIV2	NT	TL
rPIV3-1	2.6±0.5	3.2±0.1	6.3±0.4	3.1±0.3	3.6±0.2	3.3±0.7
PIV2/V94	2.8±0.7	5.0±0.3	3.8±0.0	7.1±0.7	≤0.2	≤0.2
rPIV3-2CT	1.5±0.4	0.5±0.2	2.9±0.1	7.2±0.1	≤0.2	≤0.2
rPIV3-2TM	1.4±0.1	1.6±0.7	4.1±0.1	5.9±0.2	1.6±0.6	1.3±0.9
rPIV3-2CT ^{op} 45	1.0±0.2	≤0.2	4.1±0.1	5.3±0.0	3.3±0.4	3.5±0.3
rPIV3-2TM ^{op} 45	0.6±0.3	≤0.2	3.4±0.2	4.6±0.6	3.0±0.5	4.1±0.2

^a African green monkeys in group of 4 were inoculated with 10⁵ TCID₅₀ of indicated virus intranasally and intratracheally on day 0.

^b Combined nasal wash and throat swab (NT) samples were collected on days 1 to 12. Tracheal lavage (TL) samples were collected on days 2, 4, 6, 8, and 10. The virus titers were determined on LLC-MK2 monolayers and expressed as log₁₀TCID₅₀/ml ± standard error (SE).

^c Serum samples collected on day 28 were assayed for their neutralizing antibody titers against PIV1 and PIV2. The titers were expressed as reciprocal mean log₂ ± SE.

^d NT specimens were collected on days 29 to 38. TL specimens were collected on days 30, 32, 34, 36, and 38.

rPIV3-2TM is attenuated in its replication in the respiratory tract of chimpanzees

Chimpanzees in groups of 4 were inoculated intranasally and intratracheally with 10^5 TCID₅₀ of rPIV3-2TM or PIV2/V94 on day 0. NT swab specimens (day 1 to 12) and tracheal lavage (days 2, 4, 6, 8, and 10) samples were collected. Virus titer was
5 determined as previously described (Durbin et al., *Virology* 261:319-30, 1999, incorporated herein by reference), and results are expressed as log₁₀TCID₅₀/ml. As shown in Table 30, rPIV3-2TM had a lower peak titer than its wild type parent PIV2/V94 and was shed for a significantly shorter duration than PIV2/V94, indicating that rPIV3-2TM is attenuated in chimpanzees. PIV2/V94 wt virus replicates to low levels in chimpanzees compared to
10 hamsters and AFGs, whereas rPIV3-2TM virus was attenuated in each of these model hosts.

Table 30. rPIV3-2TM is attenuated in the respiratory tract of chimpanzees and yet still elicits a strong serum immune response to PIV2

Inoculated virus ^a	Mean peak titer ^b of virus shed in indicated site (log ₁₀ TCID ₅₀ /ml±SE)		Mean days of virus shedding in the upper respiratory tract (days ± SE)	Serum neutralizing antibody titer ^c against indicated virus (reciprocal mean log ₂ ± SE)	
	NT	TL		PRE	POST
PIV2/V94	2.9 ± 0.6	1.2 ± 0.5	8.8 ± 1.1 ^d	≤2.8 ± 0.0	6.2 ± 0.5
rPIV3-2TM	2.0 ± 0.3	≤0.5 ± 0.0	2.5 ± 0.7 ^d	3.3 ± 0.2	4.3 ± 0.4

^a Chimpanzees in group of four were inoculated intranasally and intratracheally with 10⁷ TCID₅₀ of indicated virus.

^b Nose/throat (NT) swab specimens and tracheal lavages (TL) were collected for 12 and 10 days, respectively, and virus titer were determined. The peak titers are expressed as log₁₀TCID₅₀/ml ± standard error (SE).

^c Serum samples collected 3 days prior and 28 days after virus inoculation were assayed for their neutralizing antibody titer against indicated virus. The titers are expressed as reciprocal mean log₂ ± SE.

^d Significant difference in duration of shedding, p≤0.005, Student T test.

As noted above, the major protective antigens of PIVs are their HN and F glycoproteins. Thus, in exemplary embodiments of the invention, live attenuated PIV candidate vaccine viruses for use in infants and young children include chimeric HPIV3-1 and HPIV3-2 viruses carrying full-length PIV1 and partial PIV2 glycoproteins, respectively
5 in a PIV3 background genome or antigenome. In the latter case, chimeric HN and F ORFs rather than full-length PIV2 ORFs are used to construct the full-length cDNA. The recovered viruses, designated rPIV3-2CT in which the PIV2 ectodomain and transmembrane domain is fused to the PIV3 cytoplasmic domain and rPIV3-2TM in which the PIV2
10 ectodomain was fused to the PIV3 transmembrane and cytoplasmic tail domain, possessed similar *in vitro* and *in vivo* phenotypes. In particular, the rPIV3-2 recombinant chimeric viruses exhibit a host range phenotype, i.e. they replicate efficiently *in vitro* but are restricted in replication *in vivo*. This attenuation *in vivo* occurs in the absence of any added mutations from *cp45*. This is an unexpected host range effect which is highly desirable for vaccine
15 purposes, in part because the phenotype is not specified by point mutations which may revert to wt. At the same time, the unrestricted growth *in vitro* is highly advantageous for efficient vaccine production.

Although rPIV3-2CT and rPIV3-2TM replicate efficiently *in vitro*, they are highly attenuated in both the upper and the lower respiratory tract of hamsters and African green monkeys (AGMs), indicating that chimerization of the HN and F proteins of PIV2 and
20 PIV3 itself specified an attenuation phenotype *in vivo*. Despite this attenuation, they are highly immunogenic and protective against challenge with PIV2 wild virus in both species. rPIV3-2CT and rPIV3-2TM were further modified by the introduction of the 12 PIV3 *cp45* mutations located outside of the HN and F coding sequences to derive rPIV3-2CT $cp45$ and rPIV3-2TM $cp45$ which replicated efficiently *in vitro* but were even further attenuated in
25 hamsters and AGMs indicating that the attenuation specified by the glycoprotein chimerization and by the *cp45* mutations was additive.

The development of antigenic chimeric viruses possessing protective antigens of one virus and attenuating mutations from another virus has been reported by others for influenza viruses (Belshe et al., N. Engl. J. Med. 338:1405-1, 1998; Murphy et al., Infectious
30 Diseases in Clinical Practice 2:174-181, 1993) and for rotaviruses (Perez-Schael et al., N. Engl. J. Med. 337:1181-7, 1997). Attenuated antigenic chimeric vaccines are more readily generated for these viruses which have segmented genomes, since genome segment

reassortment occurs with high frequency during coinfection. Live attenuated influenza virus vaccine candidates are antigenically updated annually by replacement of the HA and NA genes of the attenuated donor virus with those of a new epidemic or pandemic virus. Recombinant DNA technology is also actively being used to construct live attenuated

5 antigenic chimeric virus vaccines for flaviviruses and for paramyxoviruses. For flaviviruses, a live attenuated virus vaccine candidate for Japanese encephalitis virus (JEV) has been made by the replacement of the pre-membrane (prM) and envelope (E) regions of the attenuated yellow fever virus (YFV) with those from an attenuated strain of JEV (Guirakhoo et al., Virology 257:363-72, 1999). The JEV-YFV antigenic chimeric recombinant vaccine

10 candidate was attenuated and immunogenic *in vivo* (Guirakhoo et al., Virology 257:363-72, 1999). Both the structural and the non-structural proteins of this chimeric virus came from a live attenuated vaccine virus. Antigenic chimeric vaccines have also been made between a naturally attenuated tick-borne flavivirus (Langat virus) and a wild type mosquito-borne dengue 4 virus, and the resulting recombinant was found to be significantly more attenuated

15 for mice than its tick-borne parent virus (Pletnev et al., Proc. Natl. Acad. Sci. U S A. 95:1746-51, 1998), but this chimeric virus was highly restricted in replication in Vero cells *in vitro*. This is an example of an attenuating effect that stems from partial incompatibility between the evolutionarily divergent structural proteins specified by the Langat virus and the non-structural proteins of the dengue virus. A third strategy is being pursued for the

20 production of a quadrivalent dengue virus vaccine in which a dengue 4 backbone containing an attenuating deletion mutation in the 3' non-coding region is used to construct antigenic chimeric viruses containing the protective antigens of dengue 1, 2 or 3 viruses (Bray et al., Proc. Natl. Acad. Sci. U S A 88:10342-6, 1991; J. Virol. 70:3930-7, 1996).

Antigenic chimeric viruses have also been produced for single-stranded,

25 negative-sense RNA viruses. For example, antigenic chimeric PIV1 vaccine candidates can be constructed according to the methods disclosed herein by substituting the full-length HN and F proteins of parainfluenza virus type 1 for those of PIV3 in an attenuated PIV3 vaccine candidate, and this recombinant is attenuated and protective against PIV1 challenge in experimental animals. Similarly, exemplary antigenic chimeric respiratory syncytial virus

30 (RSV) vaccine candidates can be made in which one or more of the RSV F and G protective antigens, or antigenic determinant(s) thereof, of subgroup B virus are substituted for those in an attenuated RSV subgroup A virus yielding attenuated RSV subgroup B vaccine candidates. (See also, International Publication No. WO 97/06270; Collins et al., Proc. Natl. Acad. Sci. USA 92:11563-11567 (1995); U.S. Patent Application No. 08/892,403, filed

July 15, 1997 (corresponding to published International Application No. WO 98/02530 and priority U.S. Provisional Application Nos. 60/047,634, filed May 23, 1997, 60/046,141, filed May 9, 1997, and 60/021,773, filed July 15, 1996); U.S. Patent Application Serial No. 09/291,894, filed by Collins et al. on April 13, 1999; U.S. Provisional Patent Application Serial No. 60/129,006, filed April 13, 1999; U.S. Provisional Patent Application Serial No. 60/143,132, filed by Bucholz et al. on July 9, 1999; and Whitehead et al., *J. Virol.* 73:9773-9780, 1999, each incorporated herein by reference). When the glycoprotein exchanges between the PIV1 and PIV3 viruses and between the RSV subgroup A and RSV subgroup B viruses were performed in a wild type virus background, the antigenic chimeric viruses replicated to wild type virus levels *in vitro* and *in vivo*. These findings indicate that a high level of compatibility exists between recipient and donor viruses and that only very little, if any, attenuation was achieved as a result of the process of chimerization. These findings with the PIV1 and PIV3 and the RSV A and B glycoprotein exchanges contrast strikingly in several ways with those between PIV2 and PIV3 disclosed herein.

In the present disclosure, viable recombinant virus in which the full-length PIV2 HN or F protein was used to replace those of PIV3 was not recovered in this instance, evidently attributable to incidental mutations introduced during cDNA construction, whereas this was successfully achieved for the PIV1-PIV3 glycoprotein exchange. This suggests that the PIV2 HN or F glycoprotein is poorly compatible with one or more of the PIV3 proteins encoded in the cDNA. Two viable PIV2-PIV3 chimeric viruses were obtained when chimeric HN and F ORFs rather than full-length PIV2 ORF were used to construct the full-length cDNA. One of these chimeric viruses contained chimeric HN and F glycoproteins in which the PIV2 ectodomain was fused to the PIV3 transmembrane and cytoplasmic tail region, and the other contained chimeric HN and F glycoproteins in which the PIV2 ectodomain and transmembrane region was fused to the PIV3 cytoplasmic tail region. Both rPIV3-2 recombinants possessed similar, although not identical, *in vitro* and *in vivo* phenotypes. Thus, it appeared that only the cytoplasmic tail of the HN or F glycoprotein of PIV3 was required for successful recovery of the PIV2-PIV3 chimeric viruses.

In previous studies directed to protein structure-function analysis, chimeric HN or F proteins have been constructed and expressed *in vitro* and have been used to map various functional domains of the proteins (Bousse et al., *Virology* 204:506-14, 1994; Deng et al., *Arch. Virol. Suppl.* 13:115-30, 1997; Deng, et al., *Virology* 253:43-54, 1999; Deng et

al., *Virology* 209:457-69, 1995; Mebatsion et al., *J. Virol.* 69:1444-1451, 1995; Takimoto et al., *J. Virol.* 72:9747-54, 1998; Tanabayashi et al., *J. Virol.* 70:6112-6118, 1996; Tsurudome et al., *J. Gen. Virol.* 79:279-89, 1998; Tsurudome et al., *Virology* 213:190-203, 1995; Yao et al., *J. Virol.* 69:7045-53, 1995). In one report, a chimeric glycoprotein consisting of a
5 measles virus F cytoplasmic tail fused to the transmembrane and ectodomains of the vesicular stomatitis virus G protein was inserted into a measles virus infectious clone in place of the measles virus F and HN virus glycoproteins (Spielhofer et al., *J. Virol.* 72:2150-9, 1998). A chimeric virus was obtained that was replication competent, but highly
10 restricted in replication *in vitro* as indicated by delayed growth and by low virus yields indicating a high degree of attenuation *in vitro*. This finding is in marked contrast to the phenotype exhibited by recombinant PIV of the invention expressing chimeric glycoproteins, e.g., a PIV2-PIV3 chimera, which replicate efficiently *in vitro*.

The efficient replication of rPIV3-2 and other chimeric PIV viruses of the invention *in vitro* is an important property for a live attenuated vaccine candidate that is
15 needed for large scale vaccine production. In contrast to rPIV3-2CT and rPIV3-2TM, rPIV3-1 was not attenuated *in vivo*. Thus, the chimerization of the HN and F proteins of PIV2 and PIV3 itself resulted in attenuation of replication *in vivo*, a novel finding for single-stranded, negative-sense RNA viruses. The mechanism for this host range restriction of replication *in vivo* is not known. Importantly, infection with these attenuated rPIV3-2CT
20 and rPIV3-2TM vaccine candidates induced a high level of resistance to challenge with PIV2 indicating that the antigenic structure of the chimeric glycoproteins was largely or completely intact. Thus rPIV3-2CT and rPIV3-2TM function as live attenuated PIV2 candidate vaccine viruses, exhibiting a desirable balance between attenuation and immunogenicity in both AGMs and hamsters.

25 The attenuating effects of the PIV3-PIV2 chimerization of the F and HN glycoprotein are additive with that specified by the *cp45* mutations. rPIV3-2 recombinants containing the *cp45* mutations were highly attenuated *in vivo* and provided incomplete protection in hamsters against challenge with PIV2 and little protection in AGMs. This is in contrast to the finding with rPIV3-1*cp45* which was satisfactorily attenuated *in vivo* and
30 protected animals against challenge with PIV1. The combination of the independent, additive attenuating effects of the chimerization of PIV3-PIV2 glycoproteins and the 12 *cp45* mutations appeared too attenuating *in vivo*. Clearly, if the rPIV3-2CT and rPIV3-2TM vaccine candidates are found to be insufficiently attenuated in humans, the *cp45* attenuating

mutations should be added incrementally rather than as a set of 12 to achieve a desired balance between attenuation and immunogenicity needed for a live attenuated PIV2 vaccine for use in humans. The findings presented herein thus identify a novel means to attenuate a paramyxovirus and provide the basis for evaluation of these PIV3-PIV2 chimeric live attenuated PIV2 vaccine candidates in humans. Importantly, the rPIV3-2CT or rPIV3-2TM viruses can also be used as vectors for other PIV antigens or for other viral protective antigens, e.g., the measles virus HA protein or immunogenic portions thereof.

Briefly summarizing the foregoing description and examples, recombinant chimeric PIVs constructed as vectors bearing heterologous viral genes or genome segments have been made and characterized using a cDNA-based virus recovery system. Recombinant viruses made from cDNA replicate independently and can be propagated in the same manner as if they were biologically-derived viruses. In preferred embodiments, recombinant chimeric human PIV (HPIV) vaccine candidates bear one or more major antigenic determinant(s) of a HPIV, preferably in a background that is attenuated by one or more nucleotide modifications. Preferably, chimeric PIVs of the invention also express one or more protective antigens of another pathogen, for example a microbial pathogen. In these cases, the HPIV acts as an attenuated virus vector and is used with the dual purpose of inducing a protective immune response against one or more HPIVs as well as against the pathogen(s) from which the foreign protective antigen(s) was/were derived. As mentioned above, the major protective antigens of PIVs are their HN and F glycoproteins. The major protective antigens of other enveloped viruses, for example viruses that infect the respiratory tract of humans, that can be expressed by the HPIV vector from one or more extra transcriptional units, also referred to as gene units, are their attachment proteins, e.g., the G protein of RSV, the HA protein of measles virus, the HN protein of mumps virus, or their fusion (F) proteins, e.g., the F protein of RSV, measles virus or mumps virus. It is also possible to express the protective antigens of non-enveloped viruses such as the L1 protein of human papillomaviruses which could form virus-like particles in the infected hosts (Roden et al., *J. Virol.* 70:5875-83, 1996). In accordance with these teachings, a large array of protective antigens and their constituent antigenic determinants from diverse pathogens can be integrated within chimeric PIV of the invention to generate novel, effective immune responses.

The present invention overcomes the difficulties inherent in prior approaches to vector based vaccine development and provides unique opportunities for immunization of

infants during the first year of life against a variety of human pathogens. Previous studies in developing live-attenuated PIV vaccines indicate that, unexpectedly, rPIVs and their attenuated and chimeric derivatives have properties which make them uniquely suited among the nonsegmented negative strand RNA viruses as vectors to express foreign proteins as vaccines against a variety of human pathogens. The skilled artisan would not have predicted that the human PIVs, which tend to grow substantially less well than the model nonsegmented negative strand viruses and which typically have been underrepresented with regard to molecular studies, would prove to have characteristics which are highly favorable as vectors. It is also surprising that the intranasal route of administration of these vaccines has proven a very efficient means to stimulate a robust local and systemic immune response against both the vector and the expressed heterologous antigen. Furthermore, this route provides additional advantages for immunization against heterologous pathogens which infect the respiratory tract or elsewhere.

These properties of PIV vectors are described herein above using examples of rPIV3 vectors which bear (i) a major neutralization antigen of measles virus expressed as a separate gene in wild type and attenuated backgrounds or (ii) major neutralization antigens of HPIV1 in place of the PIV3 neutralization antigens which express in addition a major neutralization antigen of HPIV2. These rPIV vectors were constructed using wild type and attenuated backgrounds. In addition, the description herein demonstrates the ability to readily modify the level of attenuation of the PIV vector backbone. According to one of these methods, varying the length of genome inserts in a chimeric PIV of the invention allows for adjustment of the attenuation phenotype, which is only apparent in wild type derivatives using very long inserts.

The present invention provides six major advantages over previous attempts to immunize the young infant against measles virus or other microbial pathogens. First, the PIV recombinant vector into which the protective antigen or antigens of measles virus or of other microbial pathogens is inserted is an attenuated rPIV bearing one or more attenuating genetic elements that are known to attenuate virus for the respiratory tract of the very young human infant (Karron et al., *Pediatr. Infect. Dis. J.* 15:650-654, 1996; Karron et al., *J. Infect. Dis.* 171:1107-1114, 1995a; Karron et al., *J. Infect. Dis.* 172:1445-1450, 1995b). This extensive history of prior clinical evaluation and practice greatly facilitates evaluation of derivatives of these recombinants bearing foreign protective antigens in the very young human infant.

The second advantage is that the rPIV backbone carrying the measles HA or other protective antigen of another human pathogen will induce a dual protective immune response against (1) the PIV, for which there is a compelling independent need for a vaccine as indicated above, and (2) the heterologous virus or other microbial pathogen whose protective antigen is expressed by the vector. This contrasts with the VSV-measles virus HA recombinant described above which will induce immunity to only one human pathogen, i.e., the measles virus, and in which the immune response to the vector itself is at best irrelevant or is potentially disadvantageous. The coding sequences of the foreign genes inserted into various members of the *Mononegavirales* Order of viruses have remained intact in the genomes of the most of the recombinant viruses following multiple cycles of replication in tissue culture cells, indicating that members of this group of viruses are excellent candidates for use as vectors (Bukreyev et al., *J. Virol.* 70:6634-41, 1996; Schnell et al., *Proc. Natl. Acad. Sci. U.S.A.* 93:11359-65, 1996a; Singh et al., *J. Gen. Virol.* 80:101-6; Yu et al., *Genes Cells* 2:457-66, 1997).

Another advantage provided by the invention is that use of a human pathogen backbone, for which there is a need for a vaccine, will favor the introduction of such a live attenuated virus vector into an already crowded early childhood immunization schedule. In addition, immunization via the mucosal surface of the respiratory tract offers various advantages. A live attenuated PIV3 was shown to replicate in the respiratory tract of rhesus monkeys and to induce a protective immune response against itself in the presence of high quantities of maternally-acquired PIV3 antibodies. The ability of two candidate PIV3 vaccines to infect and to replicate efficiently in the upper respiratory tract of the very young human infant who possess maternally-acquired antibodies has also been demonstrated (Karron et al., *Pediatr. Infect. Dis. J.* 15:650-654, 1996; Karron et al., *J. Infect. Dis.* 171:1107-1114, 1995a; Karron et al., *J. Infect. Dis.* 172:1445-1450, 1995b). This is in contrast to the currently licensed measles virus vaccine which is poorly infectious when administered to the upper respiratory tract of humans and which is highly sensitive to neutralization when administered parenterally to young children (Black et al., *New Eng. J. Med.* 263:165-169, 1960; Kok et al., *Trans. R. Soc. Trop. Med. Hyg.* 77:171-6, 1983; Simasathien et al., *Vaccine* 15:329-34, 1997). The replication of the HPIV vector in the respiratory tract will stimulate the production of both mucosal IgA and systemic immunity to the HPIV vector and to the expressed foreign antigen. Upon subsequent natural exposure to wild type virus, e.g., measles virus, the existence of vaccine-induced local and systemic

immunity should serve to restrict its replication at both its portal of entry, i.e., the respiratory tract, as well as at systemic sites of replication.

Yet another advantage of the invention is that chimeric HPIVs bearing heterologous sequences replicate efficiently *in vitro* demonstrating the feasibility for large scale production of vaccine. This is in contrast to the replication of some single-stranded, negative-sense RNA viruses which can be inhibited *in vitro* by the insertion of a foreign gene (Bukreyev et al., *J. Virol.* 70:6634-41, 1996). Also, the presence of three antigenic serotypes of HPIV, each of which causes significant disease in humans and hence can serve simultaneously as vector and vaccine, presents a unique opportunity to sequentially immunize the infant with antigenically distinct variants of HPIV each bearing the same foreign protein. In this manner the sequential immunization will permit the development of a primary immune response to the foreign protein which can be boosted during subsequent infections with the antigenically distinct HPIV also bearing the same or a different foreign protein or proteins, i.e., the protective antigen of measles virus or of another microbial pathogen. It is also likely that readministration of homologous HPIV vectors will also boost the response to both HPIV and the foreign antigen since the ability to cause multiple reinfections in humans is an unusual but characteristic attribute of the HPIVs (Collins et al., In "Fields Virology", B. N. Fields, D. M. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, B. Roizman, and S. E. Straus, Eds., Vol. 1, pp. 1205-1243. Lippincott-Raven Publishers, Philadelphia, 1996).

Yet another advantage is that the introduction of a gene unit into a PIV vector has several unexpected, but highly desirable effects, for the production of attenuated viruses. First, the insertion of gene units expressing, for example, the HA of measles virus or the HN of PIV2 can specify a host range phenotype on the PIV vector that has not been previously recognized, i.e., the resulting PIV vector replicates efficiently *in vitro* but is restricted in replication *in vivo* in both the upper and lower respiratory tracts. These findings identify the insertion of a gene unit expressing a viral protective antigen as an attenuating factor for the PIV vector, a desirable property in live attenuated virus vaccines of the invention.

The PIV vector system has unique advantages over all other members of the single-stranded, negative-sense viruses of the Order Mononegavirales. First, most other mononegaviruses that have been used as vectors are not derived from human pathogens (e.g., murine HPIV1 (Sendai virus) (Sakai et al., *FEBS Lett.* 456:221-6, 1999), vesicular stomatitis

virus (VSV) which is a bovine pathogen (Roberts et al., *J. Virol.* 72:4704-11, 1998), and canine PIV2 (SV5) He et al., *Virology* 237:249-60, 1997). For these nonhuman viruses, little or only weak immunity would be conferred against any human virus by antigens present in the vector backbone. Thus, a nonhuman virus vector expressing a supernumerary gene for a human pathogen would induce resistance only against that single human pathogen. In addition, use of viruses such as VSV, SV5, rabies, or Sendai virus as vector would expose vaccinees to viruses that they likely would not otherwise encounter during life. Infection with, and immune responses against, such nonhuman viruses would be of marginal benefit and would pose safety concerns, because there is little experience of infection with these viruses in humans.

An important and specific advantage of the PIV vector system is that its preferred, intranasal route of administration, mimicking natural infection, induces both mucosal and systemic immunity and reduces the neutralizing and immunosuppressive effects of maternally-derived serum IgG that is present in infants. While these same advantages theoretically are possible for using RSV as a vector, for example, we have found that RSV replication is strongly inhibited by inserts of greater than approximately 500 bp (Bukreyev et al., *Proc. Natl. Acad. Sci. USA* 96:2367-72, 1999). In contrast, as described herein, HPIV3 can readily accommodate several large gene inserts. The finding that recombinant RSV is unsuitable for bearing large inserts, whereas recombinant PIVs are highly suitable, represents unexpected results.

It might be proposed that some other viral vector could be given intranasally to obtain similar benefits as shown for PIV vectors, but this has not been successful to date. For example, the MVA strain of vaccinia virus expressing the protective antigens of HPIV3 was evaluated as a live attenuated intranasal vaccine against HPIV3. Although this vector appeared to be a very efficient expression system in cell culture, it was inexplicably inefficient in inducing resistance in the upper respiratory tract of primates (Durbin et al., *Vaccine* 16:1324-30, 1998) and was inexplicably inefficient in inducing a protective response in the presence of passive serum antibodies (Durbin et al., *J. Infect. Dis.* 179:1345-51, 1999). In contrast, PIV3 and RSV vaccine candidates have been found to be protective in the upper and lower respiratory tract of non-human primates, even in the presence of passive serum antibodies (Crowe et al., *Vaccine* 13:847-855, 1995; Durbin et al., *J. Infect. Dis.* 179:1345-51, 1999).

The use of PIV3 in particular as a vector offers yet additional advantages. For example, conditions have been established to obtain high titers of PIV3 in microcarrier culture that are 10 to 1000 times greater than can be achieved with viruses such as RSV and measles virus. Also, RSV infectivity is unstable, which complicates propagation, transport, storage and handling. These problems will be obviated by development of a PIV-vectored RSV vaccine.

Importantly, two versions of PIV3 have undergone extensive clinical evaluation as candidate vaccines administered intranasally, namely BPIV3 and the attenuated HPIV3 cp45 strain. Each was found to be safe, immunogenic, and phenotypically stable in children and infants. No other candidate engineered vector has been evaluated in children and infants, and in particular no other available vector has been evaluated for intranasal administration in this age group.

Another advantage of the PIV vector system is that, using HPIV3 as a model, a number attenuating mutations have been identified that can be introduced into the vector backbone singly and in combination to obtain the desired degree of attenuation. For example, the specific mutations that confer the HPIV3 cp45 attenuation phenotype have been identified directly by sequence analysis and introduction into wild type recombinant virus. Additional attenuating mutations were developed by "importing" attenuating point mutations from Sendai virus and RSV. In some cases, it was possible to introduce certain point mutations into recombinant virus using two nucleotide changes rather than one, which stabilizes the mutation against reversion to wild type. Ablation of expression of the C, D and V ORFs was shown to attenuate the virus. In addition, chimeric viruses of HPIV3 and bovine (B)PIV3 were developed to use the natural host range restriction of BPIV3 in primates as a means of attenuation. It also was found that certain sequence combinations were attenuating, such as replacement of the HPIV3 HN and F ectodomains with their counterparts from HPIV2. Thus, a large menu of PIV attenuating mutations exists that can be used to attenuate the vector backbone as desired.

Thus, one aspect of the invention disclosed herein relates to a method of using selected recombinant PIVs as vectors to express one or multiple protective antigens of a heterologous pathogen as supernumerary genes. The heterologous pathogens described herein include heterologous PIVs, measles virus, and RSV. In the examples above, rHPIV3 was engineered as a vector to express up to three separate supernumerary gene inserts each

expressing a different viral protective antigen. Furthermore, rHPIV3 readily accommodated a total aggregate insert length of at least 50% that of the wild type genome. Constructs were made with several different PIV vector backbones, namely: wild type HPIV3; an attenuated version of HPIV3 in which the N ORF was replaced by that of BPIV3; the HPIV3-1
5 chimeric virus, in which the HN and F ORFs of HPIV3 were replaced by their counterparts from HPIV1; a version of HPIV3-1 that was attenuated by the presence of three independent attenuating cp45 point mutations in the L gene; and a version of BPIV3 in which the HN and F genes were replaced by their counterparts from HPIV3. These vectors bearing one or more
10 supernumerary genes replicated efficiently *in vitro*, demonstrating feasibility for their commercial development, and they replicated and induced strong immune responses *in vivo* against both the vector and the inserts. In this way it is possible to construct a single recombinant PIV-based virus that is capable of inducing an immune response against at least four human pathogens, namely the PIV vector itself and the pathogens represented by the
15 supernumerary genes.

15 A second aspect of the invention is to use the superior characteristics of PIV as a vaccine and as a vector to make a vaccine against RSV. RSV is a pathogen that grows less well than PIV, is unstable, and tends to induce immune responses that are poorly protective for reasons that are not completely understood. The development of a live-attenuated RSV vaccine has been underway for more than 35 years, indicating the difficulty
20 of achieving an appropriate balance between immunogenicity and attenuation for this human pathogen. Thus, there are compelling reasons for developing a live attenuated RSV vaccine that is not based on infectious RSV. The RSV major protective F and G antigens were expressed as supernumerary genes from a PIV vector, in this case BPIV3, obviating the need to produce a live-attenuated vaccine based on infectious RSV.

25 A third aspect of the invention described herein has been to develop PIV-based vectors bearing the antigenic determinants of different PIV serotypes. Since there is essentially no cross protection between serotypes, this makes it possible to develop a method for sequential immunizations with a common PIV vector in which the protective antigenic determinants are changed. Thus, a single attenuated PIV vector backbone such as derived
30 from rHPIV3, bearing supernumerary genes as desired, can be used for an initial immunization. A subsequent immunization, which preferably follows the first by 4-6 or more weeks, can be achieved using a version of the same PIV vector in which the vector glycoprotein genes have been replaced with those of a heterologous PIV serotype, such as in

rHPIV3-1. This vector can contain the same supernumerary genes, which would then provide a "boost" against the supernumerary antigens, or can contain a different set. Because the second immunization is done with a version of the vector containing the glycoproteins of a heterologous PIV serotype, there is some interference by vector-specific immunity induced by the initial immunization. Alternatively, the second immunization can be performed with a PIV vector in which all of the vector genes are of a different serotype, such as HPIV1 or HPIV2. However, the advantage of using a common set of internal genes, such as in the rPIV3 and rPIV3-1 vectors that are based on HPIV3, is that a single set of attenuating mutation can be employed in each construct, and there is no need to separately develop attenuated strains for each PIV serotype. Importantly, sequential immunization follows a multivalent strategy: in each immunization, the vector itself induces immunity against an important human pathogen and each supernumerary insert induces immunity against an additional pathogen.

Although the foregoing invention has been described in detail by way of example for purposes of clarity of understanding, it will be apparent to the artisan that certain changes and modifications may be practice within the scope of the appended claims which are presented by way of illustration not limitation. In this context, various publications and other references have been cited within the foregoing disclosure for economy of description. Each of these references are incorporated herein by reference in its entirety for all purposes.

DEPOSIT OF BIOLOGICAL MATERIAL

The following materials have been deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, under the terms of the Budapest Treaty.

<u>Virus</u>	<u>Accession No.</u>	<u>Deposit Date</u>
p3/7(i131)2G	(ATCC 97989)	April 18, 1997
p3/7(131)	(ATCC 97990)	April 18, 1997
p218(131)	(ATCC 97991)	April 18, 1997
HPIV3 JS <i>cp45</i>	(ATCC PTA-2419)	August 24, 2000

EDITORIAL NOTE

APPLICATION NUMBER – 20731/01

The following sequence listing pages 1 – 35 are part of the description. The claims pages follow on pages 222 – 246

SEQUENCE LISTING

<110> THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as
Murphy, Brian R.
Collins, Peter L.
Schmidt, Alexander C.
Durbin, Anna P.
Skiadopoulos, Mario H.
Tao, Tao

<120> USE OF RECOMBINANT PARAINFLUENZA VIRUSES (PIVs) AS
VECTORS TO PROTECT AGAINST INFECTION AND DISEASE CAUSED
BY PIV AND OTHER HUMAN PATHOGENS

<130> 15280-404100PC

<140>
<141>

<150> 60/170,195
<151> 1999-12-10

<150> 09/458,813
<151> 1999-12-10

<150> 09/459,062
<151> 1999-12-10

<160> 62

<170> PatentIn Ver. 2.1

<210> 1
<211> 41
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Flanking
sequence of Measles HA gene insert for N-P and P-M
junctions

<400> 1
cttaagaata tacaaataag aaaaacttag gattaagag cg 42

<210> 2
<211> 36

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Flanking
sequence of Measles HA gene insert for N-P and P-M
junctions

<400> 2
gatccaacaa agaaacgaca ccgaacaac cttaag 36

<210> 3
<211> 101
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Flanking
sequence of Measles HA gene insert for HN-L
junction

<400> 3
aggcctaaaa gggaaatata aaaaacttag gagtaaagtt acgcaatcca actctactca 60
tataattgag gaaggaccca atagacaaat ccaaatcga g 101

<210> 4
<211> 79
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Flanking
sequence of Measles HA gene insert for HN-L
junction

<400> 4
tcataattaa ccataatatg catcaatcta tctataatac aagtatatga taagtaatca 60
gcaatcagac aataggcct 79

<210> 5
<211> 64
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Cloning site
for GU insertion

<400> 5
aggaaaaggg aatatataaa aacttaggag taaagttacg cgtgttaact tcgaagagct 60
ccct 64

<210> 6
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Cloning site
for NCR insertion

<400> 6
aggaaaaggg aacgcgtgtt aacttcgaag agotccct 38

<210> 7
<211> 63
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Cloning site
for supernumerary gene insert between the P and M
genes of rHPiV3

<400> 7
ttaacaatat acaataaga aaaacttagg attaaagagc catggcgtac gaagcttacg 60
cgt 63

<210> 8
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: FIV3 gene end
(GE) sequence

<400> 8
aagtaagaaa aa 12

<210> 9
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Cloning site
for RSV G and F gene inserts in B/H PIV3

<400> 9
aggattaaag aactttaccg aaagtaagg gaaagaaat cctaagagct tagcgatg 58

<210> 10
<211> 11
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Flanking
sequence for RSV G gene insert in B/H PIV3

<400> 10
gcttagcgat g 11

<210> 11
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Flanking
sequence of RSV G and F gene inserts in B/H PIV3

<400> 11
aagctagcgc ttagc 15

<210> 12
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Flanking
sequence for RSV F gene insert in B/H PIV3

<400> 12
gccttagcaaa aagctagcac aatg 24

<210> 13
<211> 83
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Forward primer
for PCR of measles HA gene insert for N-P and P-M
junctions

<400> 13
ttaatCttaa gaatatacaa ataagaaaa cttaggatta aagagcgatg tcaccacaac 60
gagaccgat aatgccttc tac 83

<210> 14
<211> 67
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Reverse primer
for PCR of measles HA gene insert for N-P and P-M
junctions

<400> 14
attattgctt aaggtttggt cggtgctggt tctttggttg atcctatctg cgattgggtc 60
catcttc 67

<210> 15
<211> 55
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Forward primer
for PCR of measles HA gene insert for HN-L
junction

<400> 15
gacaatagcc ctaaaaggga aatataaaaa acttaggagt aaagttacgc aatcc 55

<210> 16
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Reverse/Forward primer for PCR of measles HA gene
insert for HN-L junction

<400> 16
gtagaacgcg tttatccggt ctcggtgtgg tgacatctcg aatttgatt tgtctattgg 60
gtccttcc 68

<210> 17
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Reverse primer
for PCR of measles HA gene insert for HN-L
junction

<400> 17
ccatgtaatt gaatccccca acactagc 28

<210> 18
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Forward/Reverse primer for PCR of measles HA gene
insert for HN-L junction

<400> 18
cggataaacg cgttctacaa agataacc 28

<210> 19
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Upstream HPIV2
HN primer

<400> 19
gggccatgga agattacagc aat 23

<210> 20
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Downstream
HPIV2 HN primer

<400> 20
caataagctt aaagcattag ttccc 25

<210> 21
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Upstream HPIV2
HN primer

<400> 21
gcgatgggcc cgaggaagga cccaatagac a 31

<210> 22
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Downstream
HPIV2 HN primer

<400> 22
cccgggtcct gatttcccg gacgctttg 30

<210> 23
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: HPIV1 HN primer

<400> 23
agtggctaata tgcattgcat ccacat 26

<210> 24
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: HPIV1 HN primer

<400> 24
gccgtctgca tggatgaatag caat 24

<210> 25
<211> 13
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Oligomer insert for rule-of-six conformity

<400> 25
cgcggcaggc ctg 13

<210> 26
<211> 14
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Oligomer insert for rule-of-six conformity

<400> 26
cgcggcgagg cctg 14

<210> 27
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Oligomer
insert for rule-of-six conformity

<400> 27
cgcgaggcct ccgcg 15

<210> 28
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Oligomer
insert for rule-of-six conformity

<400> 28
cgcgccgcgg aggcct 16

<210> 29
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Oligomer
insert for rule-of-six conformity

<400> 29
cgcgccgcg gagcct 17

<210> 30
<211> 42
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Forward primer
for RSV A G gene insert

<400> 30
aattcgctta gcgatgtcca aaaacaagga ccaacgcacc gc 42

<210> 31
<211> 92
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Reverse primer
for RSV A G gene insert

<400> 31
aaaaagctaa gcgctagcct ttaatcctaa gtttttctta ctttttttac tactggcgtg 60
gtgtgttggg tggagatgaa ggttgtgatg gg 92

<210> 32
<211> 65
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Forward primer
for RSV A F gene insert

<400> 32
aaaggcctgc ttagcaaaaa gctagcacia tggagttgct aatcctcaaa gcaaatgcaa 60
ttacc 65

<210> 33
<211> 89
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Reverse primer
for RSV A G gene insert

<400> 33
aaaagctaag cgctagcttc ttaatccta agtttttctt acttttatta gttactaaat 60
gcaatattat ttataccact cagttgatc 89

<210> 34
<211> 44
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Mutagenic
forward primer for modification of rHPiV3-1 cDNA

<400> 34
cggccgtgac gcgtctccgc accggtgtat taagccgaag caaa 44

<210> 35
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Mutagenic
reverse primer for modification of rHPiV3-1 cDNA

<400> 35
cccggacag cttgtctct aagttttta tatttcccg acgtctattg tctgattgc 59

<210> 36
<211> 95
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Forward primer
for insertion of HPiV2 F ORF into rB/HPiV3 genome

<400> 36
aaaatatagc ggccgcaagt aagaaaaact taggattaaa ggcggatgga tcacctgcat 60
ccaatgatag tatgcatttt tgttatgtac actgg 95

<210> 37
<211> 72
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Reverse primer
for insertion of HPIV2 F ORF into rB/HPIV3 genome

<400> 37
aaaatagc ggccgctttt actaagatat cccatatatg tttccatgat tgttcttga 60
aaagacggca gg 72

<210> 38
<211> 81
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Forward primer
for insertion of HPIV2 HN ORF into rB/HPIV3 genome

<400> 38
ggaaaggcgc gccaaagtaa gaaaaactta ggattaaagg cggatggaag attacagcaa 60
tctatctctt aaatcaattc c 81

<210> 39
<211> 54
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Reverse primer
for insertion of HPIV2 HN ORF into rB/HPIV3 genome

<400> 39
ggaaaggcgc gccaaaatta aagcattagt tcccttaaaa atggtattat ttgg 54

<210> 40
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for
construction of PIV3-2 chimeric cDNAs, PIV2 F
(sense)

<400> 40
gtaccatgga tcacctgcat ccaat 25

<210> 41
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV2 F (antisense)

<400> 41
tgtggatcct aagatatccc atatatgttt c 31

<210> 42
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV2 F (sense)

<400> 42
atgcatcacc tgcatacaat 20

<210> 43
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV2 (antisense)

<400> 43
tagtgaataa agtgtcttgg ct 22

<210> 44
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV2 HN (sense)

<400> 44
catgagataa ttcattcttga tggt 24

<210> 45
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV2 HN (antisense)

<400> 45
agcttaaagc attagtctcc ttaa 24

<210> 46
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV3 F (sense)

<400> 46
atcataatta ttttgataat gatcatta 28

<210> 47
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV3 F (antisense)

<400> 47

WO 01/42445

PCT/US00/33293

gttcagtgct tgttggtt

19

<210> 48
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV3 HN (sense/antisense)

<400> 48
tcataattaa ccataatatg catcaat

27

<210> 49
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV3 HN (sense)

<400> 49
gatggaatta attagcacta tgat

24

<210> 50
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV2 F (antisense)

<400> 50
atgcatcacc tgcatoaat

20

<210> 51
<211> 19
<212> DNA

15

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV2 F (sense)

<400> 51

gatgatgtag gcaatcagc

19

<210> 52

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV2 HN (sense)

<400> 52

actgccacaa ttcttggc

18

<210> 53

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV2 HN (antisense)

<400> 53

ttaaagcatt agttccctta aaaatg

26

<210> 54

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV3 F (sense)

<400> 54
aagtattaca gaattcaaaa gag 23

<210> 55
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for
construction of PIV3-2 chimeric cDNAs, PIV3 HN
(antisense)

<400> 55
cttattagtg agcttggtgc 20

<210> 56
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for
construction of PIV3-2 chimeric cDNAs, PIV2 F
(sense)

<400> 56
accgcagctg tagcaatagt 20

<210> 57
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for
construction of PIV3-2 chimeric cDNAs, PIV2 HN
(antisense)

<400> 57
gattccatca cttaggtaaa t 21

<210> 58

<211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV3 M (sense)

<400> 58
 gatactatcc taatattatt gc 22

<210> 59
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer for construction of PIV3-2 chimeric cDNAs, PIV3 L (antisense)

<400> 59
 gctaattttg atagcacatt 20

<210> 60
 <211> 15492
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Sequence of pFLC.PIV32, 15492 bp in sense orientation

<400> 60
 accaacaag agaagaaact tgtctgggaa tataaattta actttaaatt aacttaggat 60
 taaagacatt gactagaagg tcaagaaaag ggaactctat aatttcaaaa atgttgagcc 120
 tatttgatac atttaatgca cgtaggcaag aaaacataac aaaatcagcc ggtggagcta 180
 tcattcctgg acagaaaaat actgtctcta tattcgcctt tggaccgaca ataactgatg 240
 ataatgagaa aatgacatta gctcttctat ttctatctca ttcactagat aatgagaaac 300
 aacatqcaca aagggcaggg ttcttgggtg ctttattgtc aatggcttat gccaatccag 360
 agctctacct aacaacaat ggaagtaatg cagatgtcaa gtatgtcata tacatgattg 420
 agaaagatct aaaacggcaa aagtatggag gatttgggt taagacgaga gagatgata 480
 atgaaaagac aactgattgg atatttggaa gtgacctgga ttatgatcag gaaactatgt 540
 tgcagaacgg caggaacaat tcaacaattg aagaccttgt ccacacattt gggtatccat 600
 catgtttagg agctcttata atacagatct ggatgttct ggtcaaagct atcactagta 660

tctcagggtt aagaaaaggc tttttcaccg gattggaagc tttcagacaa gatggaacag 720
 tgcaggcagag gctggtattg agcggtgaca cagtgatca gattgggtca atcatgagg 780
 ctcaacagag cttggtaact cttatggttg aaacattaat aacaatgaat accagcagaa 840
 atgacctcac aaccatagaa aagaatatac aaattgttgg caactacata agagatgca 900
 gtctcgcttc attcttcaat acaatcagat atggaattga gaccagaatg gcagctttga 960
 ctctatccac tctcagacca gatatacaata gattaaaagc tttgatggaa ctgtatttat 1020
 caaaggggacc acgcgctcct ttcactctgta tcctcagaga tcctatacat ggtgagttcg 1080
 caccaggcaa ctatcctgcc atatggagct atgcaatggg ggtggcagtt gtacaaaata 1140
 gagccatgca acagtatgtg acgggaagat catatctaga cattgatatg tccagctag 1200
 gacaagcagt agcacgtgat gccgaagctc aaatgagctc aacactggaa gatgaactg 1260
 gagtgcacaca cgaatctaaa gaaagcttga agagacatat aaggaaacata aacagttcag 1320
 agacatcttt ccacaaaccg acaggtggat cagccataga gatggcaata gatgaagagc 1380
 cagaacaatt cgaacataga gcagatcaag acaaaaatgg agaacctcaa tcatccataa 1440
 tccaatctgc ctgggcagaa ggaatatgaa gcgatgatca gactgagcaa gctacagaat 1500
 ctgacaatat caagaccgaa caacaaaaca tcagagacag actaaacag agactcaacg 1560
 acaagaagaa acaagcaggt caaccaccca ctaatcccac aaacagaaca aaccaggcag 1620
 aaatagatga tctgtttaac gcatttggaa gcaactaatc gaatcaacat tttaatctaa 1680
 atcaataata aataagaaaa acttaggatt aaagaatcct atcataccgg aatataggtt 1740
 ggtaaattta gagtctgctt gaaactcaat caatagagag ttgatggaaa gcgatgctaa 1800
 aaactatcaa atcatggatt ctggggaaga ggaatcaaga gataaatcaa ctaatatctc 1860
 ctccggcctc aacatcttg aattcactact cagcaccgac cccaagaag acttatcgga 1920
 aaacgcacaca atcaacacaa gaaccagca actcagtgcc acctctgtc aaccagaat 1980
 caaaccaaca gaaacaagt agaaagatag tggatcaact gacaaaaata gacagtcagg 2040
 gtcatcacac gaatgtacaa cagaagcaaa agatagaaat attgatcagg aaactgtaca 2100
 gagaggacct gggagaagaa gcagctcaga tagtagagct gagactgtg tctctggagg 2160
 aatcccaga agcatcacag attctaaaaa tggaaaccaa aacacggagg atattgatct 2220
 caatgaaatt agaaagatgg ataaggactc tattgagggg aaaaatgagc aatctgcaa 2280
 tgttccaagc gagatatcag gaagtgatga catatttaca acagaacaaa gtagaacag 2340
 tgatcatgga agaagcctgg aatctatcag tacacctgat acaagatcaa taagtgttgt 2400
 tactgtgca acaccagatg atgaagaaga aataactatg aaaaatagta ggacaaaaga 2460
 aagttcttca acacatcaag aagatgacaa aagaattaaa aaagggggaa aagggaaaaga 2520
 ctggtttaag aaatcaaaaag ataccgacaa ccagatacca acatcagact acagatccac 2580
 atcaaaaagg cagaagaaaa tctcaaaagc aacaaccacc aacaccgaca caaaggggca 2640
 aacagaata cagacagaat catcagaaac acaatcctca tcatggaatc tcatcatcga 2700
 caacaacacc gaccgggaagc aacagacaag cacaactcct ccaacaacaa ctccagatc 2760
 aacttataca aaagaatcga tccgaacaaa ctctgaatcc aaacccaaga cacaaaagac 2820
 aaatggaaag gaaagggaag atacagaaga gagcaatcga tttacagaga gggcaattac 2880
 tctattgcaag aatcttgggt taattcaatc cacatcaaaa ctagatttat atcaagacaa 2940
 acgagttgta tgtgtagcaa atgtactaaa caatgtagat actgcatcaa agatagattt 3000
 cctggcagga ttagtcatag gggtttcaat ggacaacgac acaaaaattaa cacagatata 3060
 aaatgaaatg ctaaacctca aagcagatct aaagaaaatg gacgaatcac atagaagatt 3120
 gatagaaaaat caaagagaac aactgtcatt gatcacgtca ctaatttcaa atctcaaaat 3180
 tatgactgag agaggaggaa agaaagacca aatgaaatcc aatggaggag tatccatgat 3240
 caaaaacaaa ttgaaagaag aaaaagatcaa gaagaccagg tttgaccac ttatggaggc 3300
 caaagcattt gacaagaata taccogactc atatcgacat gcaggagata cactagagaa 3360
 cgatgtacaa gttaaatcag agatattaag ttcatacaat gagtcaaatg caacaagact 3420
 aatacccaaa aaagtggca gtacaatgag atcactagtt gcagtcacata acaacagcaa 3480
 tctctcaaa agcacaaaac aatcatacat aaacgaactc aaacgttgca aaaaatgatga 3540

agaagtatct gaattaatgg acatgttcaa tgaagatgtc aacaattgcc aatgatccaa 3600
 caaagaaacg acaccgaaca aacagacaag aaacaacagt agatcaaaac ctgtcaacac 3660
 acacaaaatc aagcagaatg aaacaacaga tatcaatcaa tatcaaaata agaaaaactt 3720
 aggatataag aataaattaa tccttgcaca aaatgagat aactaactct gcaatataca 3780
 cattcccaga atcatcattc tctgaaaatg gtcataatga accattacca ctcaaagtca 3840
 atgaacagag gaaagcagta ccccacatta gagttgccaa gatcggaaat ccaccaaaac 3900
 acggatcccc gtatttagat gtcttcttac tcggcttctt cgagatggaa cgaatcaaaag 3960
 acaaatacgg gagtgtgaaat gatctcgaca gtgacccgag ttacaaaagt ttgtggctctg 4020
 gatcattacc aatcggattg gctaagtaca ctgggaatga ccaggaattg ttacaagccg 4080
 caaccaaacg ggatatagaa gtgagaagaa cagtcaaaag gaaagagatg gttgtttaca 4140
 cggtaaaaaa tataaaacca gaactgtacc catggtccaa tagactaaga aaaggaatgc 4200
 tgttcgagtc caaaaaagt gctcttgctc ctcaatgtct tccactagat aggagcataa 4260
 aatttagagt aatctctcgt aattgtacgg caattggatc aataaccttg tcaaaaattc 4320
 ctaagtcaat ggcataccta tctctacca acacaatct aatcaatctg caggtacaca 4380
 taaaaacagg ggttcagact gattctaaag ggatagtcca aattttggat gagaaaggcg 4440
 aaaaatcact gaatttcagt gtcactctcg gattgatcaa aagaaaagta ggcagaatgt 4500
 actctgttga atactgtaa cagaaaatcg agaaaatgag attgatattt tctttaggac 4560
 tagttggagg aatcagctct catgtcaatg caactgggtc catatcaaaa acactagcaa 4620
 gtcagctggt attcaaaaga gagatttgtt atcctttaat ggatctaaat ccgcatctca 4680
 atctagttat ctgggtctca tcagtagaga ttacaagagt ggatgcaatt ttccaacctt 4740
 cttcacctcg cgagttcaga tactatccta atattattgc aaaaggagtt gggaaaaatca 4800
 aacaatggaa ctagtaact ctattttagt cgggacgtat ctattaagcc gaagcaata 4860
 aaggataatc aaaaacttag gacaaaagag gtcaatacca acaactatta gcagtcacac 4920
 tcgcaagaat aagagagaag ggaccaaaaa agtcaaatag gagaaatcaa aacaaaaggt 4980
 acagaacacc agaacaaca aatcaaaaca tccaactcac tcaaaaacaa aattcaaaaa 5040
 gagaccggca acacaacaag cactgaacac catggatcac ctgcatccaa tgaatgatag 5100
 catttttgtt atgtacactg gaattgtagg ttcagatgcc attgctggag atcaactcct 5160
 caatgtaggg gtcattcaat caaagataag atcactcatg tactacactg atggtggcgc 5220
 tagctttatt gttgtaaaat tactacccaa tcttccccca agcaatggaa catgcaacat 5280
 caccagctca gatgcatata atgttaccct atttaagttg ctaacacccc tgatgagaa 5340
 cctgagcaaa atttctgctg ttacagatac caaaccccg cgagaacgat ttgcaggagt 5400
 cgttattggg cttgctgcac taggagtagc tacagctgca caataaccg cagctgtagc 5460
 aatagtaaaa gccaatgcaa atgctgctgc gataaacaat ctgcatctt caattcaatc 5520
 caccaacaag gcagtatccg atgtgataac tgcatcaaga acaattgcaa ccgagttca 5580
 agcgattcag gatcacatca atggagccat tgtcaacggg ataacatctg catcatgccg 5640
 tgcccattgat gcaactaatt ggtcaatatt aaatttggat ctcaactgagc ttactacaat 5700
 atttcataat caaataacaa accctgcgct gacaccactt tccatccaag ctttaagaat 5760
 cctcctcggt agcaccttgc caattgtcat tgaatccaaa ctcaacacaa aactcaacac 5820
 agcagagctg ctcaagtgcg gactgttaac tggtaaaata atttccattt ccccaatgta 5880
 catgcaaatg ctaattcaaa tcaatgttcc gacatttata atgcaacccg gtgcgaaggt 5940
 aattgatcta atgtctatct ctgcaaacca taaattacaa gaagtgttg tacaagtcc 6000
 taatagaatt ctagaatatg caaatgaact acaaaactac ccagccaatg attgtttcgt 6060
 gaccccaaac tctgtatatt gtatatacaa tgagggttcc ccgatccctg aatcaacaata 6120
 tcaatgctta agggggaatc ttaattcttg cacttttacc cctattatcg ggaactttct 6180
 caagcgatcc gcatttgcca atggtgtgct ctatgccaac tgcaaatctt tgctatgtaa 6240
 gtgtgcccag cctccccatg ttgtgtctca agatgacaac caaggcatca gcataattga 6300
 tattaagagg tgctctgaga tgatgcttga cactttttca tttaggatca catctacatt 6360
 caatgctaca tacgtgacag acttctcaat gattaatgca aatattgtac atctaagtcc 6420

tctagacttg tcaaatcaaa tcaattcaat aaacaaatct cttaaaagtg ctgaggattg 6480
gattgcagat agcaacttct tcgctaatac agccagaaca gccaaagcac ttatttcact 6540
aagtgaactc gcattaatac tatcagtgat tacttttggtt gttgtgggat tgctgattgc 6600
ctacatcctc aagctgggtt ctcaaatcca tcaattcaga gcactagctg ctacaacaat 6660
gttccacagg gagaatcctg ccgtctttc caagaacaat catggaaaca tatatgggat 6720
atcttaggat ccctacagat cattaagatata taaaattata aaaaacttag gagtaaaagt 6780
acgcaatcca actctactca tataattgag gaaggaccca atagacaaat ccaaatccat 6840
ggaagattac agcaatctat ctcttaaatc aattcctaaa aggacatgta gaatcatttt 6900
cogaactgcc acaattcttg gcataatcac attaatgtg ctatgttcaa gtattcttca 6960
tgagataatt catcttgatg tttcctctgg tcttatgaat tctgatgagt cacagcaagg 7020
cattatcag ccctacatag aatcattaaa atcattgatt gctttggcca accagattct 7080
atataatggt gcaatagtaa ttcctcttaa aattgacagt atcgaactg taatactctc 7140
tgcttataaa gatattgaca cggggagtat gtccaatgcc aactgcacgc caggaaactc 7200
gcttctgcat gatgcagcat acatcaatgg aataaacaaa ttctctgtac ttgaatcata 7260
caatgggagc cctaaatag gacctctcct aaatataccc agctttatcc cctcagcaac 7320
atctcccatc ggggtacta gaatacctc attttcaact atcaagacc attgggtgta 7380
cactcaaat gtaatgctg gagattgtct tgatttcacg gcactcaacc agtatttacc 7440
aatggggata atacaacaat ctgctgcagg gtttccaatt ttcaggacta tgaaaacct 7500
ttacctaat gatgaaatca atcgcacaag ctgttcagtc actgctatac caggagggtg 7560
tgctctgcat tgctatgtag ctacaaggtc tgaaaaagaa gattatgcca gagctgatct 7620
agctgaactg agacttgctt tctattatta taatgatacc tttattgaaa agtcatatc 7680
tcttccaaat acaacagggc agtggggcac aatcaacct gcagtcgga gccggatcta 7740
tcacttaggc tttatcttat ttcctgtata tgggtgctc ataaatggga ctacttctta 7800
caatgagcag tctcaccgct attttatccc aaaacatccc aacataactt gtgcccgtaa 7860
ctccagcaaa caggctgcaa tagcacggag ttcctatgct atccgttatc actcaaacag 7920
gttaattcag agtgcgtgct tttattgtcc attgtctgac atgcatacag aagagttaa 7980
tctagttatg tttacaatt cccaagtcat gatgggtgca gaagtaggc tctatgttat 8040
tggtataaat ttgtattatt atcaacgcag ttcctcttg tggctgcat cgctcttta 8100
caggatcaat acagattttt ctaaggaaat tctctccgac attgaggctc aatgggtacc 8160
gtcctatcaa gttcctcgtc ctggagtcac gccatgcaat gcaacaagt tttgccctgc 8220
taattgcatc acaggggtgt acgcagatgt gtggccgctt aatgatccag aactcatgct 8280
acgtaatgct ctgaacccca actatcgatt tgctggagcc tttctcaaaa atgagtccaa 8340
ccgaactaat cccacattct acactgcatc ggctaactcc ctcttaata ctaccggatt 8400
caacaacacc aatcacaag cagcatatac atcttcaacc tgcttataaa aactgggac 8460
ccaaaaaatt tattgttaa taataattga aatgggctca tctcttttag gggagtcca 8520
aataataca tttttaagg aactaatgct ttaagcttaa ttaaccataa tatgcatcaa 8580
tctatctata atacaagtat atgataagta atctgcaatc agacaataga caaaagggaa 8640
atataaaaaa cttaggagca aagcgtgctc gggaaatgga cactgaatct acaaatggca 8700
ctgtatctga catactctat cctgagtgct accttaactc tctatcgtt aaaggtaaaa 8760
tagcacactt acacactatt atgagctcac ctccagccta tgatatggat gacgactcaa 8820
tactagttat cactagacag aaaataaac ttaataaat ggataaaaaga caacgatcta 8880
ttagaagatt aaaattaata ttaactgaaa aagtgaatga cttaggaaaa tacacattta 8940
tcagatatcc agaaatgtca aaagaaatgt tcaaatata tatacctggt attaacgta 9000
aagtgactga attattactt aaagcagata gaacatatac tcaaatgact gatggattaa 9060
gagatctatg gattaatgct ctatcaaaat tagcctcaaa aatgatgga agcaattatg 9120
atcttaatga agaaatfaat aatatacga aagttcacac aacctataaa tcagataaat 9180
ggataaatcc attcaaaaca tggtttacta tcaagtatga tatgagaaga ttacaaaaag 9240
ctcgaatga gatcactttt aatgttggga aggattataa ctgtttagaa gaccagaaga 9300

atctcttatt gatacatcca gaattggttt tgatattaga taaacaaaac tataatggtt 9360
atctaatcacc tcctgaatta gtattgatgt attgtgacgt agtcgaaggc cgatggaata 9420
taagtgcctg tgctaagtta gatccaaaat tacaatctat gtatcagaaa ggtaataacc 9480
tgtgggaagt gatagataaa ttgtttccaa ttatgggaga aaagacattt gatgtgata 9540
cgttattaga accacttgca ttatccttaa tcaaaactca tgatcctgtt aaacaactaa 9600
gaggagcttt ttraaatcat gtgttatccg agatggaatt aatatttgaa tctagagaat 9660
cgattaagga atttctgagt gtagattaca ttgataaaat tttagatata ttttaataagt 9720
ctacaataga tgaatagca gagattttct ctttttttag aacatttggg catcctccat 9780
tagaagctag tattgcagca gaaaaggtta gaaaatata gtatattgga aaacaattaa 9840
aatttgacac tattaataaa tgcctatgta tcttctgtac aataataatt aacggatata 9900
gagagaggca tgggtggacag tggcctcctg tgacattacc tgatcagca cacgaattca 9960
tcataaatgc ttacggttca aactctgcca tatcatatga aaatgctgtt gattattacc 10020
agagctttat aggaataaaa ttcaataaat tcatagagcc tcagtttagt gaggatttga 10080
caattttatat gaaagataaa gcattatctc caaaaaaatc aaattgggac acagttttatc 10140
ctgcatctaa tttactgtac cgtactaacg catccaacga atcacgaaga ttagtgaag 10200
tatttatagc agatagtaaa ttgtatcctc atcagatatt ggattatgta gaattctgggg 10260
actggttaga tgatccgaaa ttaataattt cttatagtct taaagaaaaa gagatcaaac 10320
aggaagttag actccttgca aaaatgacat acaaaatgag agctacacaa gttttatcag 10380
agaccctact tgcaaaatac ataggaaaat tctttcaaga aaatgggatg gtgaagggag 10440
agattgaatt acttaagaga ttaacaacca tatcaatc aggaqtcca cggtataatg 10500
aagtgtacaa taattctaaa agccatacag atgaccttaa aacctacaat aaaataagta 10560
atcttaattt gtcttctaat cagaaatcaa agaaatttga attcaagtca acggatattc 10620
acaatgatgg atacgagact gtgagctgtt tcctaacaac agatctcaaa aaatactgtc 10680
ttaattggag atatgaatca acagctctat ttggagaaac ttgcaaccaa atatttggat 10740
taaataaatt gtttaattgg ttacaccctc gtcttgaagg aagtacaatc tatgtaggtg 10800
atccttactg tcctccatca gataaagaac atatatcatt agaggatcac cctgattctg 10860
gtttttacgt tcataacca agaggggta tagaaggatt ttgtcaaaaa ttatggacac 10920
tcataatctat aagtgaata catctagcag ctgttagaat aggcgtgagg gtgactgcaa 10980
tggttcaagg agacaatcaa gctatagctg taaccacaag agtaccacaac aattatgact 11040
acagagttaa gaaggagata gtttataaag atgtagttag attttttgat tcattaagag 11100
aagtgtatga tgatctaggt catgaactta aattaatga aacgattata agtagcaaga 11160
tgttcatata tagcaaaaga atctattatg atgggagaat tcttctcaa gctctaaaag 11220
cattatctag atgtgtctc tggctagaga cagtaataga cgaacaaga tcagcatctt 11280
caaatttggc aacatcattt gcaaaagcaa ttgagaatgg ttattcacct gttctaggat 11340
atgcatgctc aatttttaag aatattcaac aactatata tgccttggg atgaatatca 11400
atccaactat aacacagaat atcagagatc agtattttag gaatccaaat tggatgcaat 11460
atgctctttt aatacctgct agtgttgggg gattcaatta catggccatg tcaagatggt 11520
ttgtaaggaa tattggtgat ccatcagttg ccgcatggc tgatataaa agatttatta 11580
agggcaactc attagaccga agtgtctttt ataggattat gaatcaagaa ccagggtgag 11640
catctttttt ggactgggct tcagatccat attcatgcaa ttaccacaa tctcaaaaata 11700
taaccaccat gataaaaaat ataacagcaa ggaatgtatt acaagattca ccaaatccat 11760
tattatctg attattcaca aatacaatga tagaagaaga tgaagaatta gctgagttcc 11820
tgatggacag gaaggtaatt ctccctaqag ttgcacatga tattctagat aattctctca 11880
caggaattag aatgocata gctggaatgt tagatacagc aaaatcacta attcgggttg 11940
gcataaatag aggaggactg acatatagtt tggttgaggaa aatcagtaat tacgatctag 12000
tacaatatga aacactaagt aggactttgc gactaattgt aagtataaa atcaagtatg 12060
aagatatgtg ttcggtagac cttgccatag cattgcgaca aaagatgtgg attcatttat 12120
caggaggaag gatgataagt ggacttgaaa cgcctgacct attagaatta ctatctgggg 12180

tagtaataac aggatcagaa cattgtaaaa tatgttattc ttcagatggc acaaacccat 12240
 atacttggat gtatttacc ggtaatatca aaataggatc agcagaaaca ggtatatcgt 12300
 cattaagagt tccttatttt ggatcagtc ctagtgaag atctgaagca caattaggat 12360
 atatcaagaa tcttagtaaa cctgcataag ccgcaataag aatagcaatg atatatatc 12420
 gggcatttgg taatgatgag atatcttgg tggaaagcctc acagatagca caaacacgtg 12480
 caaattttac actagatagt ctcaaaattt taacacocgt agctacatca acaaatttat 12540
 cacacagatt aaaggatact gcaactcaga tgaattctc cagtacatca ttgatcagag 12600
 tcagcagatt cataacaatg tccaatgata acatgtctat caaagaagct aatgaaacca 12660
 aagatactaa tcttatttat caacaataa tgttaacagg attaagtgtt ttcgaattat 12720
 tatttagatt aaaagaaacc acaggacaca accctatagt tatgcactcg cacatagaag 12780
 atgagtggtg tattaagaa agtttfaatg atgaacatat taatccagag tctacattag 12840
 aattaattcg atactctgaa agtaatgaat ttatttatga taaagacca ctcaaagatg 12900
 tggacttacc aaaacttatg gttatataag accattctta cacaattgat atgaattat 12960
 gggatgatac tgacatcata catgcaattt caatagtac tgaattaca atagcagata 13020
 ctatgtcaca attagatcga gataatttaa aagagataat agttattgca aatgatgatg 13080
 atattaatag cttaactact gaatttttga ctcttgacat acttgatttt ctcaagacat 13140
 ttggtggatt attagtaaat caatttgcat acactcttta tagtctaaaa atagaaggta 13200
 gggactctcat ttgggattat ataatgagaa cactgagaga tacttcccat tcaatattaa 13260
 aagtattatc taatgcatta tctcatccta aagtattcaa gaggttctcg gattgtggag 13320
 ttttaaaccc tatttatggt cctaactctg ctagtcaaga ccagataaaa cttgcctat 13380
 ctatctgga atattcacta gatctattta tgagagaaatg gttgaatggt gttcactctg 13440
 aaatatatcat ttgtgacagc gatatggaag ttgcaaatga tgggaaacaa gcctttattt 13500
 ctagacacct ttcatattgt ttgtgtttag cagaaattgc atcttctgga cctaacctgt 13560
 taaactaac atacttggag agacttgatc tattgaaaca atacttgaa ttaaatatta 13620
 aagaagacc tactcttaa tatgtacaaa tatctggatt attaatataa tegtcccat 13680
 caactgtaac atactgtaaa aagactgcaa tcaaatatct aaggattcgc ggtattagtc 13740
 cacctgaggt aattgatgat tgggatccgg tagaagatga aaatagctg gataacattg 13800
 tcaaaactat aaatgataac tgtaataaag ataataaagg gaataaaatt aacaatttct 13860
 gggactagc acttaagaac tatcaagtc ttaaatcag atctataaca agtgattctg 13920
 atgataatga tagactagat gctaatacaa gtggtttgac acttctcaa ggagggaatt 13980
 atctatcgca tcaattgaga ttattcggaa tcaacagcac tagttgtctg aaagctctg 14040
 agttatcaca aattttaatg aaggaagtca ataaagacaa ggacaggctc ttctgggag 14100
 aaggagcagg agctatgcta gcattgtatg atgccacatt aggacctgca gttaatatt 14160
 ataattcagg tttgaatata acagatgtaa ttggtcaacg agaattgaaa atatttctt 14220
 cagaggtatc attagtaggt aaaaaattag gaaatgtgac acagattctt aacagggtaa 14280
 aagtactggt caatgggaa cctaattcaa catggatagg aaataggaa tgtgagagct 14340
 taatattgag tgaattaaat gataagtcca ttggattagt acattgtgat atggaggag 14400
 ctatcgtaa atcagaagaa actgttctac atgaacatta tagtgtata agaattacat 14460
 acttgattgg gtagatgat gttgttttag tttccaaat tatacctaca atcactccga 14520
 atgtgctag aatctttat ctatataaat tatattgaa agatgtaagt ataatatc 14580
 tcaaaacttc taatcctgca tcaacagaat tatatctaat ttcgaaagat gcatattgta 14640
 ctataatgga acctagttaa attgtttat caaaactaa aagattgtca ctctggaag 14700
 aaaataatct attaaatgg atcattttat caaagaagag gaataatgaa tggttacttc 14760
 atgaaatcaa agaaggagaa agagattatg gaatcatgag accatcat atggcactac 14820
 aaatcttgg atttcaaatc aatttaaatc atctggcga agaattttta tcaaccocag 14880
 atctgactaa tatcaaatc ataatacaaa gttttcagcg aacaataaag gatgtttat 14940
 ttgaatggat taatataact catgatgata agagacataa attagcggga agatataaca 15000
 tattccact gaaaaataag gaaaagttaa gactgctatc gagaagacta gtattaaagt 15060

```

ggatttcatt atcattatcg actcgattac ttacaggtcg ctttctgat gaaaaatttg 15120
aacatagagc acagactgga tatgtatcat tagctgatac tgatttagaa tcattaaagt 15180
tattgtcgaa aaacatcatt aagaattaca gagagtgtat aggatcaata tcattattgg 15240
ttctaaccaa agaagttaaa atacttatga aattgatcgg tgggtctaaa ttattaggaa 15300
ttcccagaca atataaagaa cccgaagacc agttattaga aaactacaat caacatgatg 15360
aatttgata cgattaaaac ataaatacaa tgaagatata tcctaaccctt tatctttaag 15420
cctaggaata gacaaaaagt aagaaaaaca tgtaatatat atataccaaa cagagttctt 15480
ctctgtttg gt 15492

```

<210> 61

<211> 15498

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sequence of
pFLC.PIV32TM, 15498 bp in sense orientation

<400> 61

```

accaaaacaag agaagaaact tgtctgggaa tataaattta actttaaatt aacttaggat 60
taaagacatt gactagaagg tcaagaaaag ggaactctat aatttcaaaa atgttgagcc 120
tatttgatac atttaatgca cgtaggcaag aaaacataac aaaatcagcc ggtggagcta 180
tcatttctcg acagaaaaat actgtctcta tattcgccct tggaccgaca ataactgatg 240
ataatgagaa aatgacatta gctcttctat ttctatctca ttcactagat aatgagaaac 300
aacatgcaca aagggcaggy ttcttggtgt ctttattgtc aatggcttat gccaatccag 360
agctctacct aacaaccaat ggaagtaatg cagatgtcaa gtatgtcata tacatgattg 420
agaaagatct aaaacggcaa aagtatggag gatttgggt taagacgaga gagatgatat 480
atgaaaagac aactgattgg atatttggaa gtgacctgga ttatgatcag gaaactatgt 540
tgcaagaacg caggaacaat tcaacaattg aagaccttgt ccacacattt gggtatccat 600
catgttttag agctcttata atacagatct ggatagttct ggtcaaaact atcactagta 660
tctcagggtt aagaaaagcc ttttccacc gattggaagc ttccagacaa gatggaacag 720
tgcaaggcag gctggtattg agcggtgaca cagtggatca gattgggtca atcatgagg 780
ctcaacagag cttgtaact cttatggttg aaaccattat aacaatgaat accagcagaa 840
atgacctcac aaccatagaa aagaatatac aaattgttgg caactacata agagatgcaq 900
gtctcgcttc attcttcaat acaatcagat atggaattga gaccagaatg gcagctttga 960
ctctatccac tctcagacca gatatcaata gattaaaagc ttgatggaa ctgtatttat 1020
caaaaggacc acgctctcct ttcattctgt. tctcagaga tctatcacat ggtgagttcg 1080
caccaggcaa ctatctgcc atatggagct atgcaatggg ggtggcagtt gtacaaaata 1140
gagccatgca acagtatgt acgggaagat catatctaga cattgatatg ttccagctag 1200
gacaagcagt agcacgtgat gccgaagctc aatgagctc aacactggaa gatgaacttg 1260
gagtgcaca cgaatctaaa gaaagcttga agagacatat aaggaacata aacagttcag 1320
agacatcttt ccacaaaacc acaggtggt cagccataga gatggcaata gatgaagagc 1380
cagaacaatt cgaacataga gcagatcaag aacaaaatgg agaacctcaa tcatccataa 1440
ttcaatatgc ctgggcagaa ggaatataga gcgatgatca gactgagcaa gctacagaat 1500
ctgcaatat caagaccgaa caacaaaaca tcagagacag actaaaacag agactcaacg 1560
acaagaagaa acaagcagt caaccacca ctaatccac aaacagaaca aaccaggagc 1620
aaatagatga tctgtttaac gcatttggaa gcaactaatc gaatcaacat tttaatctaa 1680

```

atcaataata aataagaaaa acttaggatt aaagaatcct atcataccgg aatatagggt 1740
 ggtaaaattha gagtctgctt gaaactcaat caatagagag ttgatggaaa gcgatgctaa 1800
 aaactatcaa atcatggatt cttgggaaga ggaatcaaga gataaatcaa ctaatatctc 1860
 ctcggccctc aacatcattg aattcactact cagcaccgac ccccaagaag acttatcgga 1920
 aaacgacaca atcaacacaa gaaccagca actcagtgcc accatctgtc aaccagaaat 1980
 caaaccacaa gaaacaagtg agaagatag tggatcaact gacaaaaata gacagtccgg 2040
 gtcacacac gaatgtacaa cagaagcaaa agatgaaat attgatcagg aaactgtaca 2100
 gagaggacct gggagaagaa gcagctcaga tagtagagct gagactgtg tctctggagg 2160
 aatccccaga agcatcacag attcctaaaaa tggaaaccCaa aacacggagg atattgatct 2220
 caatgaat agaaagatgg ataaggactc tattgagggg aaaaatgcgac aatctgcaaa 2280
 tgtccaagc gagatcag gaagtgatga catattaca acagaacaaa gtagaaacag 2340
 tgatcatgga agaagcctgg aatctatcag tacacctgat acaagatcaa taagtgttg 2400
 tactgtctga acaccagatg atgaagaaga aataactaat aaaaatagta ggacaaagaa 2460
 aagtctctca acacatcaag aagatgacaa aagaattaaa aaagggggaa aagggaaaga 2520
 ctgggttaag aatcaaaag ataccgacaa ccagatacca acatcagact acagatccac 2580
 atcaaaagg cagaagaaaa tctcaaaagc aacaaccacc aacaccgaca caaaggggca 2640
 aacagaaata cacagaaat catcagaaac acaatctca tcatggaatc 2700
 caacaacacc gaccggaacg aacagacaa cacaaactct ccaacaacaa cttccagatc 2760
 aacttataca aaagaatcga tccgaacaaa ctctgaatcc aaaccaca cacaagaagc 2820
 aaatggaaag gaaaggaag atacagaaga gagcaatcga ttacagaga gggcaattac 2880
 tctattgacg aatctgggtg taattcaatc cacatcaaaa ctgagttat atcaagacaa 2940
 acgagttgta tgttagcaaa atgtactaaa caatgtgat actgcatcaa agatagattt 3000
 cctggcagga ttagtcatg gggtttcaat ggacaaccgac acaaaattaa cacagataca 3060
 aaatgaaatg ctaaacctca aagcagatct aaagaaaatg gacgaatcac atagaagatt 3120
 gatagaaat caaagaaac aactgtcatt gatcaactca ctaatttcaa atctcaaaat 3180
 tatgactgag agaggaggaa agaagacca aaatgaatcc aatgagagag tatccatgat 3240
 caaaacaaaa ttgaaagaag aaaagatcaa gaagaccagg tttgaccac ttatggaggc 3300
 acaaggcatt gacaagaata taccgatct atatcgacat gcaggagata cactagagaa 3360
 cgatgtacaa gttaaatcag agatattaag ttcatacaat gagtcaaat caacaagact 3420
 aatacccaaa aaagtgaqca gtacaatgag atcaactagt gcagtcac caacaagcaa 3480
 tctctcacia agcacaaaa aatcatacat aaacgaactc aaacgttgca aaaatgatga 3540
 agaagatct gaattaatgg acatgttcaa tgaagatgac aacaattgcc aatgatccaa 3600
 caaagaaacg acaccgaaca aacagacaa aaacaacagt agatcaaaac ctgtcaaac 3660
 acacaaaatc aagcagaatg aaacaacaga tatcaatcaa tacaacaaat aaaaaactt 3720
 aggattaaa gataaattaa tcctgttcca aaatgagat aactaactct gcaatataca 3780
 cattcccaga atcatcttc tctgaaaatg gtcataatga accattacca ctcaaagtca 3840
 atgaacagag gaaaagcagta cccacatta gagttgcaa gatcggaat ccacaaaaac 3900
 acggatcccg gtatttagat gtctctctac tcggctctct cgagatggaa cgaatcaaa 3960
 acaaaatcgg gagtgtgaaat gatctcgaca gtgaccgag ttacaaagtt tgtggtctctg 4020
 gatcattacc aatcgattg gctaagtaca ctgggaatga ccaggaattg ttacaagccg 4080
 caaccaaact ggatatagaa gtgagaagaa cagtcaaacg gaaagagatg gttgtttaca 4140
 cggtacaaaa tataaaacca gaactgtacc catggtccaa tagactaaga aaaggaatgc 4200
 tggttcagtc caacaaagt gctcttqctc ctcaatgct tccactagat agggacataa 4260
 aatttagagt aatctctgtg aattgtacgg caattggatc aataaccttg ttcaaaattc 4320
 ctaaagcaat ggcacacta tctctacca acacaatc aatcaatctg caggtacaca 4380
 taaaaacagg ggttcagact gattcctaaag ggatagttca aattttgat gagaaaggcg 4440
 aaaaatcact gaatttcatg gtccatctcg gattgatcaa aagaaaagta ggcagaatgt 4500
 actctgttga atactgtaaa cagaaaatcg agaaaatgag attgatattt tctttaggac 4560

tagttggagg aatcagcttt catgtcaatg caactgggtc catatcaaaa acactagcaa 4620
 gtcagctgggt attcaaaaga gagatttgtt atcctttaat ggatctaaat cgcactctca 4680
 atctagtatt ctgggcttca tcagtagaga ttacaagagt ggatgcaatt ttccaacctt 4740
 ctttacctgg cgagttcaga tactatccta atattattgc aaaaggagt gggaaaaatca 4800
 aacaatggaa ctagtaatct ctattttagt ccggacgtat ctattaagcc gaagcaata 4860
 aaggataatc aaaaacttag gacaaaagag gtcaatacca acaactatta gcagtcacac 4920
 tcgcaagaat aagagagaag ggaccaaaaa agtcaaatag gagaaatcaa aacaaaaggt 4980
 acagaacacc agaacaacaa aatcaaaaca tccaactcac tcaaaacaaa aattccaaaa 5040
 gagaccggca acacaacaag cactgaacat gcactcactg catccaatga tagtatgcat 5100
 ttttgttatg tacactggaa ttgttagttc agatgccatt gctggagatc aactcctcaa 5160
 tgtaggggtc attcaatcaa agataagatc actcatgtac tacactgatg gtggcgctag 5220
 ctttattgtt gtaaaattac taccatctc tcccccaagc aatggaacat gcaacatcac 5280
 cagctagatg gcataataat ttacctatt taagtgtgta acacccctga ttgagaacct 5340
 gagcaaaatt tctgctgta cagatacca accccggcga gaacgattg caggagtctg 5400
 tattgggctt gctgcactag gagtagctac agctgcacaa ataaccgcag ctgtagcaat 5460
 agtaaaagcc aatgcaaatg ctgctgcgat aaacaactct gcacttcaa tcaatccac 5520
 caacaaggca gatacogag tgataactgc atcaagaaca attgcaaccg cagtcaagc 5580
 gattcaggat cacatcaatg gagccattgt caacgggata acatctgcat tagccgtgc 5640
 ccatgatgca ctaattgggt caatattaaa ttgtatctc actgagctta ctacaattt 5700
 tcataatcaa ataacaacc ctgcgtgac accactttcc atccaagctt taagaatcct 5760
 cctcggtagc accttgcaa ttgtcattga atccaaactc aacacaaaac tcaacacagc 5820
 agagctgctc agtagcggac tgttaactgg tcaataaatt tccatttccc caatgtacat 5880
 gcaaatgcta attcaatca atgttccgac atttataatg caacccgggt cgaaggtaat 5940
 tgatctaatt gctatctctg caaacctaaa attacaagaa gtagttgtac aagttcctaa 6000
 tagaattcta gaatatgcaa atgaactaca aaactacca gccaatgatt gtttctgac 6060
 accaaactct gtattttgta gatacaatga gggtecccg atccctgaat cacaatatca 6120
 atgcttaagg gggaatctta attcttgca ttttaccctc attatcgga accttctcaa 6180
 gcgattogca ttgccaatg gtgtgctcta tgccaactgc aaactttgc tatgtaagt 6240
 tgccgacctc ccccatgttg tctctcaaga tgacaaccaa ggcacagca taattgata 6300
 taagaggtgc tctgagatga tcttgacac ttttctatt aggatcacat ctacattcaa 6360
 tgetacatac gtgacagact tctcaatgat taatgcaaat attgtacatc taagtctct 6420
 agacttgta aatcaatca attcaataaa caaatctctt aaaagtctg aggattggat 6480
 tgagatagc aacttctcg ctaatcaagc cagaacagcc aagacactt attcactaat 6540
 cataattatt ttgataatga tcattatatt gtttataatt aatataacga taattacaat 6600
 tgcaattaa gattacagaa ttcaaaagag aaatcgagt gatcaaatg acaagccata 6660
 tgtactaaca acaaaataac atactacag atcattagat attaaaatta taaaaaactt 6720
 aggagtaaa ttacycaatc caactctact catataatg aggaaggacc caatagacaa 6780
 atccaaattc gagatggaat actggaagca taccaatcac ggaaggatg ctggtaatga 6840
 gctggagacy tctatggcta ctcatggcaa caagctcact aataagata tatacatatt 6900
 atggacaata atcctgggtg tattatcaat agtcttcac atagtctaa ttaattccat 6960
 ccatgagata attcatctg atgttctc tggctctatg aattctgatg agtcacagca 7020
 aggcattatt cagccatca tagaatcatt aaaaacttg attgctttg ccaaccagat 7080
 tctataaat gttgcaatg taattcctc taaaattgac agtatcgaaa ctgtaactt 7140
 ctctgcttta aaagatatgc acaccgggag tatgtccaat gccaactgca cgccaggaaa 7200
 tctgctctg catgatgag catacatcaa tggataaac aaattcctg tacttgaatc 7260
 atacaatgg acgcctaaat atggacctc ctaaatata cccagcttta toccctcagc 7320
 aacatctcc catgggtgta ctagaatacc atcatttca ctcatcaaga cccattgggt 7380
 ttacactcac aatgtaatg ttggaattg tcttgattc acggcatcta accagattt 7440

atcaatgggg ataatacaac aatctgctgc agggtttcca attttcagga ctatgaaaac 7500
catttaccta agtgatggaa tcaatcgcaa aagctgttca gtcactgcta taccaggagg 7560
ttgtgtcttg tattgctatg tagctacaag gtctgaaaaa gaagattatg ccacgactga 7620
tctagctgaa ctgagacttg ctttctatta ttataatgat acctttattg aaagagtcac 7680
atctcttcca aatacaacag ggcagtgggc cacaatcaac cctgcagtcg gaagcgggat 7740
ctatcatcta ggctttatct ttttcctgt atatggtgtg ctataaaatg ggactacttc 7800
ttacaatgag cagtccctcac gctattttat cccaaaaacat cccaacataa cttgtgcccg 7860
taactccaag aaacaggctg caatagcacg gagttcctat gtcacccgtt atcactcaaa 7920
caggtaaatc cagagtgtg ttcttatttg tccattgtct gacatgcata cagaagagtg 7980
taactagtgt atgtttaaca attcccaagt catgatgggt gcagaaggta ggctctatgt 8040
tattgtaaat aatttattt attatcaacg cagttcctct tgggtgtctg catcgctctt 8100
ttacaggatc aatacagatt tttctaaagg aattcctccg atcattgagg ctcaatgggt 8160
accgtcctat caagttcctc gtccctggagt catgccatgc aatgcaacaa gtttttgccc 8220
tgctaattgc atcacagggg tgtacgcaga tgtgtggccg cttaatgatc cagaactcat 8280
gtcacgtaat gctctgaacc ccaactatcg atttgtgga gcctttctca aaaatgagtc 8340
caaccgaaat aatcccaatc tctacactgc atcgggtaac tccctcttaa atactaccgg 8400
attcaacaac accaatcaca aagcagcata tacatcttca acctgcttca aaaaactggy 8460
aacccaaaaa atttattgtt taataataat tgaatggggc tcatctcttt taggggagt 8520
ccaaataata ccatttttaa gggaactaat gctttaaagt tcataattaa ccaataatag 8580
catcaatcta tctataatac aagtatatga taagtaatca gcaatcagac aatagacaaa 8640
agggaaatc aaaaaacta ggagcaaacg gtgctcgga aatggacact gaactcaaca 8700
atggcactgt atctgacata ctctatcctg agtgcacct taactcctct atcgtttaaag 8760
gtaaaatagc acaattcac actattatga gtctacctca gccttatgat atggatgacg 8820
actcaatact agttatcact agacagaaaa taaaacttaa taaattggat aaaagacac 8880
gatctattag aagattaaaa ttaattatga ctgaaaaagt gaatgactta ggaaaatca 8940
catttatcag ataccagaa atgtcaaaag aatgttcaa attatataa cctggatta 9000
acagtaaagt gactgaatta ttacttaaag cagatagaac atatagtcaa atgactgatg 9060
gattaagaga tctatggatt aatgtgctat caaaatagc ctcaaaaaat gatggaagca 9120
attatgatct taatgaagaa attaataata tatcgaaagt tcacacaacc tataatcag 9180
ataaatggta taatccattc aaaacatggt ttactatcaa gtatgatag agaagattac 9240
aaaaagctcg aatgagatc acttttaagt ttgggaagga ttataacttg ttagaagacc 9300
agaagaatct cttattgata catccagaat tggttttgat attagataaa caaaactata 9360
atggttatct aattactcct gaattagatc tgatgtattg tgacgtagtc gaaggccgat 9420
ggaatataag tgcatgtgct aagttagatc caaaattaca atctatgtat cagaaaagta 9480
ataacctgtg ggaagtgata gataaattgt ttccaattat gggagaaaag acatttgatg 9540
tgatctcgtt attagaacca ctgcatatc ccttaattca aactcatgat cctgtttaa 9600
aaactaaggg agctttttaa aatcatgtgt tatccgagat ggaattaata tttgaatcta 9660
gagaatcgat taaggaattt ctgagtgtag attacattga taaaattta gatatttta 9720
ataagtctac aatagatgaa atagcagaga ttttctcttt ttttagaaca tttgggcac 9780
ctccattaga agctagtatt gcagcagaaa aggttagaaa atatatgtat attgaaaaac 9840
aataaaatt tgacactatt aataaatgtc atgctatctt ctgtacaata ataattaacg 9900
gatatagaga gaggcatggt ggacagtgcc ctccctgtac attacctgat catgcacacg 9960
aattcatcat aatgcttac ggttcaaac ctgcgatatc atatgaaaat gctgttgat 10020
attaccagag ctttatagga ataaaattca ataaattcat agagcctcag ttgatgagg 10080
atttgacaat ttatatgaaa gataaagcat tatctccaaa aaaatcaaat tgggacacag 10140
tttactctgc atctaattta ctgtaccgta ctaaccgatc caacgaatca cgaagattag 10200
ttgaagtatt tatagcagat agtaaatttg atcctcatca gatattggat tatgtagaat 10260
ctgggactg gttagatgat ccagaattta atatttctta tagtcttaaa gaaaaagaga 10320

tcaaacagga aggttagactc tttgcaaaaa tgacatacaa aatgagagct acacaagttt 10380
 tatcagagac cctacttgca aataacatag gaaaattcct tcaagaaaaat gggatgggtga 10440
 agggagagat tgaattactt aagagattaa caaccatatac aatatcagga gttccacggt 10500
 ataataaggt gtacaataat tctaaaagcc atacagatga ccttaaaacc tacaataaaa 10560
 taagtaactc taatttgtct tctaatacaga aatcaaaagaa atttgaattc aagtcaacgg 10620
 atactatacaa tgaatggatc gagactgtga gctgtttcct aacaacagat ctcaaaaaat 10680
 actgtcttaa ttggagatata gaatcaacag ctctatttgg agaacttgc aaccaaata 10740
 ttggattaaa taaattgttt aattggttac accctcgtct tgaagggaagt acaatctatg 10800
 taggtgatac ttactgtcct ccatcagata aagaacatatac atcattagag gatcacccctg 10860
 atctgtgttt ttacgttcat aacccaagag ggggtataga aggatattgt caaaaattat 10920
 ggacactcat atctataagt gcaatacatc tagcagctgt tagaataaggc gtgagggtga 10980
 ctgcaatggt tcaaggagac aatcaagcta tagctgtaac cacaagagta cccaacaatt 11040
 atgactacag agttaagaag gagatagttt ataaagatgt agtgagattt tttgattcat 11100
 taaggaaggt gatggatgat ctgagtcagt aacttaaat aaatgaaacg atataagta 11160
 gcaagatggt catatatagc aaaagaatct attatgatgg gagaattcct cctcaagctc 11220
 taaaagcatt atctagatgt gtcttctggt cagagacagt aatgagacaa acaagatcag 11280
 catcttcaaa tttggcaaca tcatttgcaa aagcaattga gaatggttat tcacctgttc 11340
 taggatatac atgtcaatt ttaagaata tcaacaact atataattgcc ctggggatga 11400
 atactcaatcc aactataaca cagaataca gagatcagta ttttaggaat ccaaatgga 11460
 tgcaatatgc ctcttaata cctgctagt tgggggatt caattacatg gccatgtcaa 11520
 gatgttttgg aaggaatatt ggtgatccat cagtgtccgc attggctgat attaaaagat 11580
 ttataaagcc gaatctatta gaccgaagtg ttctttatag gattatgaat caagaaccag 11640
 gtgagtcctc ttttttggac tgggctcag atccatattc atgcaattta ccaaatctc 11700
 aaaaataaac caccatgata aaaaataaa cagcaaggaa tgtattcaaa gattcaoccaa 11760
 atccattatt atctggatta ttcacaaata caatgataga agaagatgaa gaattagctg 11820
 agttcctgat ggacaggaag gtaattctcc ctgagttgc acatgatatt ctgataaatt 11880
 ctctcacagg aattagaat gccatagctg gaatgttaga tacgcaaaaa tcaactaattc 11940
 gggttggcat aaatagagga ggactgacat atagtttgtt gaggaaaatc agtaattacg 12000
 atctagtaca atataaaca ctaagttaga ctttgcgact aatgttaagt gataaaatca 12060
 agtatgaaga tatgtgttgc gtgaccttg ccatagcatt gcgcaaaaag atgtggattc 12120
 atttatcagg aggaaggatg ataatggac ttgaaacgcc tgaccatta gaattactat 12180
 ctggggtagt aataacagga tcagaacatt gtaaaatag ttattcttca gatggcaca 12240
 acccatatac ttggatgtat ttaccggta atatacaaat aggatcagca gaaacaggta 12300
 tatcgtcatt aagagttcct tattttggat cagtcaactg tgaagatct gaagcacaat 12360
 taggatatac caagaatcct agtaaacctg caaaagccgc aataagaata gcaatgatat 12420
 atacatgggc atttggtaat gatgagatat cttggatgga agcctcacag atagcacaaa 12480
 cacgtgcaaa ttttacctca gatagtctca aaattttaac accgtagct acatcaacaa 12540
 atttatcaca cagattaaag gatactgcaa ctacagatgaa attctccagt acatcattga 12600
 tcagagtcag cagattcata acaatgtcca atgataacat gtctatcaaa gaagctaatg 12660
 aaaccaaaaga tactaatctt atttatcaac aaataatggt aacaggatta agtgttttgc 12720
 aatatttatt tagatataaa gaaaccacag gacacaaccc tatagttatg catctgcaca 12780
 tagaagatga gtgtgttatt aaagaagtt ttaatgatga acatattaat ccagagtcta 12840
 cattagaatt aattcgatat cctgaaagta atgaatttat ttatgataaa gaccactca 12900
 aagatgtgga cttatcaaaa cttatgggta ttaaagacca ttcttacaca attgatatga 12960
 attattggga tgatactgac atcatacatg caatttcaat atgtactgca attacaatag 13020
 cagatactat gtcacaatta gatcgagata atttaaaaga gataatagtt attgcaaatg 13080
 atgatgatat taatagctta atcaactgaat ttttgactct tgacatactt gtatttctca 13140
 agacatttgg tggattatta gtaaatcaat ttgcatcac ctcttatagt ctaaaaatag 13200

```

aaggtaggga tctcatttgg gattatataa tgagaacct gagagatact tcccattcaa 13260
tattaagaat attatcfaat gcattatctc atcctaaagt attcaagagg ttctgggatt 13320
gtggagtttt aaaccctatt tatggtccta atactgctag tcaagaccag ataaaaactg 13380
ccctatctat atgtgaatat tcactagatc tatttatgag agaattggtt aatggtgat 13440
caattgaaat atacatttgt gacagcgata tggaaattgc aaatgatagg aaacaagcct 13500
ttatttctag acacccttca ttgtttgtt gtttagcaga aattgcatct ttcggacct 13560
acctgttaaa cttaacatac ttggagagac ttgatctatt gaaacaatat cttgaattaa 13620
atattaaaga agaccctact cttaaatag tacaatatc tggattatta attaaatcgt 13680
tcccatcaac tgtaacatac gtaagaaaga ctgcaatcaa atactaagg attcgcggta 13740
ttagtcacc tgaggtaatt gatgattggg atccggtaga agatgaaat atgctggata 13800
acattgtcaa aactataaat gataactgta ataaagataa taaagggaaat aaaattaa 13860
atctctgggg actagcactt aagaactatc aagtccttaa aatcagatct ataacaagt 13920
atctgatga taatgataga ctgatgcta atacaatgg tttgacact cctcaaggag 13980
ggaattatct atcgcataca ttgagattat tggaaatcaa cagcactagt tgcctgaaa 14040
ctcttgagtt atcacaaatt ttaatgaagg aagtcataa agacaaggac aggcctcttc 14100
tgggagaagg agcaggagct atgctagcat gttatgatgc cacattagga cctgcaagta 14160
attattataa ttcaggttt aatataacag atgtaattg tcaacgagaa tgaataat 14220
ttccttcaga ggtatcatta gtaggtaaaa aattaggaaa tgtgacacag attccttaaca 14280
gggtaaaagt actgttcaat ggaatccta attcaacatg gataggaat atggaatgt 14340
agagcttaat atggagtga ttaaatgata agtccattgg attagatcat tgtgatatg 14400
aaggagctat cggtaaatca gaagaaactg ttctacatga acattatagt gttataagaa 14460
ttacataact gattgggat gatgatgtt ttttagttc caaaattata cctacaatca 14520
ctccgaattg gtcagaata cttatctat ataaattata ttgaaagat gtaagtataa 14580
tatcactcaa aacttctaat cctgcatcaa cagaattata tctaatttcg aaagatgat 14640
attgtactat aatggaacct agtgaattg tttatcaaa acttaaaaga ttgtactct 14700
tggagaagaa taacttatta aaatggatca tttatcaaa gaagaggaa atgaatgt 14760
tacatcatga aatcaaagaa ggagaaagag attatggaat catgagacca tatcatatg 14820
cactacaaat ctttgattt caaatcaat taaatcatct ggcgaaagaa tttttatcaa 14880
cccagatct gactaatatc aacaatataa tccaaagttt tcagcgaaca ataaaggatg 14940
ttttattga atgattaat ataactcatg atgataagag acataaatta ggcggaagt 15000
ataacatatt cccactgaaa aataaggaa agttaagact gctatcgaga agactagtat 15060
taagttggat ttcatatca ttatcgactc gattacttac aggtcgttt cctgatgaaa 15120
aattgaaac tagagcacag actggatag tatcattagc tgatactgat ttagaatcat 15180
taaagttatt gtcgaaaaac atcattaaga attacagaga gtgtatagga tcaatatcat 15240
attggtttct aaccaagaa gttaaaatc ttatgaaatt gatcgggtgt gctaaattat 15300
taggaattcc cagacaatat aaagaacctg aagaccagt attagaaac tacaatcaac 15360
atgatgaatt tgatatcgat taaacataa atacaatgaa gatatacct aacctttatc 15420
tttaagccta ggaatagaca aaaagtaaga aaaacatgta atatatatat accaaacaga 15480
gtttctctct tgtttggt 15498

```

<210> 62
<211> 15492
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sequence of

pFLC.PIV32CT, 15474 bp in sense orientation

```

<400> 62
accaaaacaag agaagaaact tgtctgggaa tataaattta actttaaatt aacttaggat 60
taaagacatt gactagaagg tcaagaaaag ggaactctat aattcaaaa atgttgagcc 120
tatttgatac atttaatgca cgtaggcaag aaaacataac aaaatcagcc ggtggagcta 180
tcattctctg acagaaaaat actgtctcta tattcgccct tggaccgaca ataactgatg 240
ataatgagaa aatgacatta gctcttctat ttctatctca ttoactagat aatgagaaac 300
aacatgcaca aagggcaggg ttcttggtgt ctttatgttc aatggcttat gccaatccag 360
agctctacct aacaacaaat ggaagtaatg cagatgtcaa gtatgtcata tacatgattg 420
agaagatct aaaacggcaa aagtatggag gatttgggt taagacgaga gagatgatat 480
atgaaaagac aactgattgg atatttggaa gtgacctgga ttatgatcag gaaactatgt 540
tgcagaacgg caggaaacat tcaacaattg aagacctgt ccacacattt gggtatccat 600
catgttttagg agctcttata atacagatct ggatgttct ggtcaaaact atcactagta 660
tctcaggggt aagaaaaggc tttttcaccg gattggaagc tttcagaca gatggaacag 720
tgcaggcagg gctggtattg agcggtgaca cagtggatca gattgggtca atcatgctg 780
ctcaacagag cttggtaact ctatggttg aaacattaat aacaatgaat accagcagaa 840
atgacctcac aacctagaa aagaatatac aaattgttg caactacata agagatgcag 900
gtctcgcctc attcttcaat acaatcagat atggaatga gaccagaatg gcagctttga 960
ctctatccac tctcagacca gatataca gattaaaagc ttgatggaa ctgtatttat 1020
caaagggacc acgctctct tcatctgta tcctcagaga tcctatacat ggtgagttcg 1080
caccaggcaa ctatcctgcc atatggagct atgcaatggg ggtggcagtt gtacaaaata 1140
gagccatgca acagtatgtg acgggaagat catatctaga cattgatatg ttccagctag 1200
gacaagcagt agcacgtgat gccgaagctc aactgagctc aacactgaa gatgaaactg 1260
gagtgcaca cgaatctaaa gaaagcttga agagacatat aaggacata aacagttcag 1320
agacatcttt ccacaaaccg acaggtggat cagccataga gatggcaata gatgaagagc 1380
cagaacaatt cgaacataga gcagatcaag acaaaaatgg agaacctcaa tcatccataa 1440
ttcaatatgc ctgggcagaa ggaatatgaa gcgatgatca gactgagcaa gctacagaat 1500
ctgacaatat caagaccgaa caacaaaaa tcagagacag actaaaacag agactcaacg 1560
acaagaagaa acaaaagcag caaccoacca ctaatccac aaacagaa caaccaggagc 1620
aaatagatga tctgtttaac gcatttggaa gcaactaatc gaatcaacat tttaatctaa 1680
atcaataata aataagaaaa acttaggatt aaagaatcct atcataccg aatatagggt 1740
ggtaaaatta gagctgctt gaaactcaat caatagagag ttgatggaaa gcgatgctaa 1800
aaactatcaa atcatggatt cttgggaaga ggaatcaaga gataaatcaa ctaatatctc 1860
ctcggccctc aacatcattg aattcatact cagcaccgac cccaagaag acttatcgga 1920
aaacgacaca atcaacaca gaaccagca actcagtgcc accatctgtc aaccagaaat 1980
caaaccaaca gaaacaagt agaaagatag tggatcaact gacaaaaata gacagtcagg 2040
gtcatcacac gaatgtacaa cagaagcaa agatagaat attgatcagg aaactgtaca 2100
gagaggacct gggagaagaa gcagctcaga tagtagagct gagactgtg tctctggagg 2160
aatcccaga agcatcacag attctaaaaa tggaaacca aacacggagg atattgatct 2220
caatgaaat agaagatgg ataaggactc tattgagggg aaatgcgac aatctgcaa 2280
tgttccaagc gagatatac gaagtatga catatttaca acagaaacaa gtgaaacag 2340
tgatcatqra aqaagcctgg aatctatcag tacacctgat acaagatcaa taagtgttg 2400
tactgtctga acaccagatg atgaagaaga aataactaatg aaaaatagta ggacaaaaga 2460
aagttcttca acacatcaag aagatgacaa aagaattaaa aaagggggaa aagggaaaga 2520
ctggtttaag aatcaaaag atacogaaa ccagatacca acatcagact acagatccac 2580
atcaaaaggg cagaagaaaa tctcaaagac acaaccacc aacaccgaca caaaggggca 2640
aacagaaata cagacagaat catcagaac acaactctca tcatggaatc tcatcatcga 2700

```

caacaacacc gaccggaacg aacagacaag cacaactcct ccaacaacaa cttccagatc 2760
aacttataca aaagaatcga tccgaacaaa ctctgaatcc aaaccaaga cacaaaagac 2820
aaatggaag gaaaggaagg atacagaaga gagcaatcga ttacagaga gggcaattac 2880
tctattgcaa aatcttgggt taattcaatc cacatcaaaa ctgattttat atcaagacaa 2940
acgagttgta tgtgtagcaa atgtactaaa caatgtagat actgcatcaa agatagattt 3000
cctggcagga ttagtcatag gggtttcaat ggacaacgac acaaaattaa cacagatata 3060
aaatgaaatg ctaaacctca aagcagatct aaagaaaatg gacgaatcac atagaagatt 3120
gatagaaaat caaagagaac aactgtcatt gatcacgtca ctaatttcaa atctcaaaat 3180
tatgactgag agagggagaa agaaagacca aaatgaatcc aatgagagag tatccatgat 3240
caaaacaaaa ttgaaagaag aaaagatcaa gaagaccagg tttgaccac ttatggaggc 3300
acaaggcatt gacaagaata taccogactc atatcgacat gcaggagata cactagagaa 3360
cgatgtacaa gttaaatcag agatattaag ttcatacaat gactcaaatg caacaagact 3420
aataccocaa aaagtggaca gtacaatgag atcactagtt gcagtcacaa acaacagcaa 3480
tctctcacia agcacaacaa aatcatacat aaacgaactc aaacgttgca aaaaatgatga 3540
agaagtatct gaattaatgg acatgttcaa tgaagatgac aacaattgcc aatgatccaa 3600
caaagaaacg acaccgaaca aacagacaag aaacaacagt agatcaaaac ctgtcaaacac 3660
acacaaaatc aagcagaatg aaacaacaga tatcaatcaa tatacaata agaaaaactt 3720
aggattaaag aataaattaa tccttgtcca aaatgagtat aactaactct gcaatataca 3780
cattcccaga atcatcattc tctgaaaatg gtcataatga accattacca ctcaaaagtc 3840
atgaacagag gaaagcagta ccccacatta gaggtgccaa gatcggaaat ccaccaaac 3900
acggatcccg gtatttagat gtcttcttac tcygcttctt cgagatggaa cgaatcaaaag 3960
acaaatcagg gagtgtgat gatctcgaca gtgaccogag ttcaaaagtt tgyggtctctg 4020
gatcattacc aatcggattg gctaagtaca ctgggaatga ccaggaattg ttcaaaagccg 4080
caaccaaaat ggatatagaa gtgagaagaa cagtcaaaag caaagagatg gttgtttaca 4140
cggtaaaaa tataaaacca gaactgtacc catggtccaa tagactaaga aaaggaatgc 4200
tgytccatgc caacaaagt gctcttctc ctcaatgtct tccactagat agggagataa 4260
aaatttagagt aatcttctg aattgtacgg caattggatc aataaccttg ttcaaaatc 4320
ctaagtcaat ggcataccta tctctacca acacaatctc aatcaatctg caggtacaca 4380
taaaaacagg ggttcagact gattctaaag ggatagttca aattttggat gagaaaggcg 4440
aaaaatcact gaatttcag gtccatctcg gattgatcaa aagaaaagta ggcagaatgt 4500
actctgttga atactgtaa cagaaaatcg agaaaatgag attgatattt tctttaggac 4560
tagttggagg aatcagttct catgtcaatg caactgggtc catatcaaaa acactagcaa 4620
gtcagctggt atcaaaaga gagatttgtt atcctttaat ggatctaat cgcactca 4680
atctagttat ctgggcttca tcagttaga ttacaagagt ggatgcaatt ttccaacctt 4740
ctttacctgg cgagttcaga tactatccta atattattgc aaaaggagtt gggaaaatca 4800
aacaatggaa ctagtaatct ctattttagt ccggacgtat ctattaagcc gaagcaata 4860
aaggataatc aaaaacttag gacaaaagag gtcaatacca acaactatta cgagtcaac 4920
tcgcaagaat aagagagaag ggacaaaaa agtcaaatag gagaatcaa aacaaaaggt 4980
acagaacacc agacaacaa aatcaaaaac tccaactcac tcaaaacaaa aattccaaa 5040
gagaccggca acacaacaag cactgaacac catggatcac ctgcatccaa tgatagtatg 5100
catttttgtt atgtacactg gaattgtagg ttcagatgcc attgctggag atcaactcct 5160
caatgtaggg gtcattcaat caaagataag atcactcatg tactacactg atggtggcgc 5220
tagctttatt gttgtaaat tactaccocaa tcttccccca agcaatggaa catgcaacat 5280
caccagtcta gatgcataa atgttaccct atttaagttg ctaaacaccc tgattgagaa 5340
cctgagcaaa atttctgctg ttacagatac caaacccgc cgagaacgat ttgcaggagt 5400
cgttattggg cttgctgac taggagtagc tacagctgca caaataaccg cagctgtagc 5460
aatagtaaaa gccaatgcaa atgctgctgc gataaacaat cttgcatctt caattcaatc 5520
caccacaag gcagatccg atgtgataac tgcatacaaga acaattgcaa ccgagttca 5580

agcgattcag gatcacatca atggagccat tgtcaacggg ataacatctg catcatgccg 5640
 tgccccatgat gcactaattg ggtcaatatt aaatttgtat ctcaactgagc ttactacaat 5700
 atttcataat caaataacaa accctgcgct gacaccaactt tccatccaag ctttaagaat 5760
 cctcctcggg agcaccttgc caattgtcat tgaatccaaa ctcaacacaa aactcaacac 5820
 agcagagctg ctocagtagc gactgttaac tggtaaaata atttccattt ccccaatgta 5880
 catgcaaatg ctaattcaaa tcaatgttcc gacatttata atgcaaccgg gtgcgaaggt 5940
 aattgatcta attgctatct ctgcaaacca taaattacaa gaagtgttg tacaagttcc 6000
 taatagaatt ctagaatatg caaatgaact acaaaactac ccagccaatg attgttctg 6060
 gacaccaaac tctgtatctt gtatatacaa tgagggttcc ccgatccctg aatcaacaata 6120
 tcaatgctta agggggaatc ttaattcttg cacttttacc cctattatcg ggaactttct 6180
 caagcgattc gcatttgcca atggtgtgct ctatgccaac tgcaaatctt tgctatgtaa 6240
 gtgtgcccag cctcccctg ttgtgtctca agatgacaac caaggcatca goataattga 6300
 tattaagagg tgctctgaga tgatgcttga cactttttca tttaggatca catctacatt 6360
 caatgtctaca tacgtgacag acttctcaat gattaatgca aatattgtac atctcaagtc 6420
 tctagacttg tcaaatcaaa tcaattcaat aaacaaatct cttaaaagtg ctgaggattg 6480
 gattgcagat agcaactctt tgcataatca agccagaaca gccaaagcac tttatctact 6540
 aagtgcactc agcattatc tatcagtgat tactttggtt gttgtgggat ctgagttgc 6600
 ctacatcctc aaqctggtt ctcaaatcca tcaattcaga gcactagctg ctacaacaat 6660
 gttcccacag gagaatcctg ccgtcttttc caagaacaat catggaaaca tatatgggat 6720
 atcttaggat cctacagat cattagat taaaattata aaaaacttag gaggtaagt 6780
 acgcaatcca actctactca tataattgag gaaggacca atagacaaat ccaaatccat 6840
 ggaagattac agcaactctat ctcttaaatc aatctctaaa aggcacatgta gaatcatttt 6900
 ccgaactgcc acaattcttg gcatatgcac attaattgtg ctatgttcaa gtattcttca 6960
 tgagataaatt catcttgatg ttctctctgg tcttatgaaat tctgatgagt cacagcaagg 7020
 cattattcag cctatcatag aatcattaaa atcattgatt gctttggcca accagattct 7080
 atataatggt gcaatagtaa ttctctttaa aattgacagt atcgaaactg taatactctc 7140
 tgctttaaaa gatatgcaca ccgggagtat gtccaatgcc aactgcacgc caggaaatct 7200
 gctctgcat gatgcagcat acatcaatgg aataaacaac ttcttctgac ttgatcata 7260
 caatggggag cctaaatag gacctctctt aaatataccc agctttatcc cctcagcaac 7320
 atctcccatc ggggttacta gaataccatc attttcactc atcaagacc cttgggtgta 7380
 cactcacaat gtaatgctg gagattgtct tgatttcacg gcatctaac agtatttacc 7440
 aatggggata atacaacaat ctgctgcagg gtttccaatt ttcaggacta tgaaaacct 7500
 ttactaaagt gatggaatca atcgcaaaag ctgttcagtc actgctatc caggagggtg 7560
 tgtcttgat tgctatgtg ctacaaggtc tgaaaaagaa gattatgcca cgactgatct 7620
 agctgaactg agacttgctt tctattatta taatgatacc ttatttgaaa gactcatatc 7680
 tcttccaaat acaacagggc agtgggccac aatcaaccct gcagtcggaa cggggatcta 7740
 tcatctaggc ttatcttat ttctgtata tgggtgtctc ataaatggga ctactctta 7800
 caatgagcag tctcagcct attttatccc aaaaatccc aacataact gtgcccgtta 7860
 ctccagcaaa caggctgcaa tagcacggag ttctatgtc atccgttatc actcaaacag 7920
 gtaattcag agtgctgttc ttattgttcc attgtctgac atgcatacag aagagtgtta 7980
 tctagttatg ttaacaatt cccaagtcag gatgggtgca gaaggtaggc tctatgttat 8040
 tggtataaat ttgtattat atcaacgcag ttctcttgg tggctgcat cgctcttata 8100
 caggatcaat acagatttt ctaaaqgaat tctctcagtc attgaggctc aatgggtacc 8160
 gtctatcaa gttctctgct ctggagtcag gccatgcaat gcaacaagt tttgcctgc 8220
 taattgcatc acaggggtgt acgagatgt gtggccgctt aatgatccag aactcatgct 8280
 acgtaatgct ctgaacccca actatcgatt tgctggagcc ttctcaaaa atgagtccaa 8340
 ccgaactaat cccacattct acactgcatc ggctaactcc ctcttaata ctaccggatt 8400
 caacaacacc aatcaaaaag cagcatatac atcttcaacc tgccttaaaa acactggaac 8460

ccaaaaaatt tattgttaa taataattga aatgggctca tctcttttag gggagttcca 8520
 aataatacca ttttaaggg aactaatgct ttaagcttaa ttaaccataa tatgcatcaa 8580
 tctatctata atacaagtat atgataagta atctgcaatc agacaataga caaaaaggaa 8640
 atataaaaaa cttaggagca aagcgtgctc gggaaatgga cactgaatct aacaatggca 8700
 ctgtatctga ctaactctat cctgagtgct accttaactc tctatctggt aaaggtaaaa 8760
 tagcacaatt acacactatt atgagctctac ctgagcctta tgatatggat gacgactcaa 8820
 tactagtatt cactagacag aaaataaaac ttaataaatt ggataaaaga caacgatcta 8880
 ttagaagatt aaaattaata ttaactgaaa aagtgaatga cttaggaaaa tacacattta 8940
 tcagatatcc agaaatgtca aaagaaatgt tcaaatata tatacctggt attaacagta 9000
 aagtgactga attattactt aaagcagata gaacatatag tcaaatgact gatggattaa 9060
 gagatctatg gattaatgct ctatcaaaat tagcctcaaa aatgatgga agcaattatg 9120
 atcttaaatga agaanaaat aatatatcga aagttcacac aacctataaa tcagataaat 9180
 ggtataatcc attcaaaa ca tggtttacta tcaagtatga tatgagaaga ttcaaaaaag 9240
 ctggaatgga gatcactttt aatgttggga aggattataa cttgttagaa gaccagaaga 9300
 atttcttatt gatacatoca gaattggttt tgatattaga taacaaaaac tataatggtt 9360
 atctaattac tctctgaatta gtattgatgt atgtgacgt agtcgaaggg cgatggaata 9420
 taagtgcatt tgctaagttg gatccaaaat tacaatctat gtatcagaaa ggtaataacc 9480
 tgtgggaatg gatagataaa ttgtttccaa ttatgggaga aaagacattt gatgtgata 9540
 cgttattaga accacttgca ttatccttaa ttcaaaactca tgatcctggt aacaactcaa 9600
 gaggagcttt tttaaatcat gtgttatccg agatggaatt aatattttaa tctagagaat 9660
 cgattaagga atttctgagt gtgattaca ttgataaaat tttagatata ttaataagt 9720
 ctacaataga tgaatagca gagattttct ctttttttag aacctttggg catcctccat 9780
 tagaagctag tattgcagca gaaaaggtta gaaaatata gtatattgga aacaattaa 9840
 aatttgacac tattaataaa tgcattgcta tctctctgac aataataatt aacggatata 9900
 gagagaggca tgggtggacag tggcctctg tgacattacc tgatcatgca cacgaattca 9960
 tcataaatgc ttacggttca aactctgca tatcatatga aatgctggt gattattacc 10020
 agagctttat aggaataaaa ttcaataaat tcatagagcc tcagttagat gaggatttga 10080
 caatttatat gaaagataaa gcattatctc caaaaaaac aatggggac acagtttctc 10140
 ctgcatctaa tttactgtac cgtactaacg catccaacga atcacaaga ttagtgaag 10200
 tatttatagc agatagtaaa ttgatcctc atcagatatt ggattatgta gaactctggg 10260
 actgggttaga tgatccagaa ttaaatattt cttatagtct taaagaaaa gagatcaaac 10320
 aggaaggtag actccttgca aaaaatgacat acaaaatgag agctacacaa gttttatcag 10380
 agaccctact tgcaataaac ataggaaaat tctttcaaga aatggggatg gtgaagggag 10440
 agattgaatt acttaagaga ttaacaacca tatcaatctc aggagtcca cggtataatg 10500
 aagtgtacaa taattctaaa agccatacag atgacctaa aacctacaat aaaataagta 10560
 atcttaattt gtcttcta at cagaatcaa agaaatttga attcaagtca acggatatct 10620
 acaatgatgg atacgagact gtgagctggt tcctaacaac agatctcaaa aataactgct 10680
 ttaattggag atatgaatca acagctctat ttggagaaac ttgcaaccaa atatttggat 10740
 taaataaatt gtttaattgg ttacacctc gtcttgaagg aagtacaatc tatgtagggt 10800
 atccttactg tctccatca gataaagaac atatatcatt agaggatcac cctgattctg 10860
 gttttactgt tcataaccca agagggggta tagaaggatt ttgtcaaaaa ttatggacac 10920
 tcatatctat aagtgaata catctagcag ctgttagaat aggcgtgaggt gtgactgcaa 10980
 tggttcaagg agacaatcaa gctatagctg taaccacaag agtaccacac aattatgact 11040
 acagagttaa gaaggagata gtttataaag atgtagttag atttttttag tcattaagag 11100
 aagtgatgga tgatctaggt catgaactta aattaaatga aacgattata agtagcaaga 11160
 tgttcatata tagcaaaaaga atctattatg atgggagaat tcttctctca gctcaaaaag 11220
 cattatctag atgtgtcttc tggtcagaga cagtaataga cgaacaaga tcagcatctt 11280
 caaatttggc aacatcattt gcaaaaagca ttgagaatgg ttatcacctt gttctaggat 11340

atgcatgctc aatttttaag aatattcaac aactatata tgccttggg atgaatatca 11400
atccaactat aacacagaat atcagagatc agtattttag gaatccaaat tggatgcaat 11460
atgcctcttt aatacctgct agtggtggg gattcaatta catggccatg tcaagatgtt 11520
ttgtaaggaa tattggtgat ccacagttg ccgcatggc tgatattaaa agatttatta 11580
aggcgaatct attagaccga agtggtcttt ataggattat gaatcaagaa ccagggtgagt 11640
catctttttt ggactgggct tcagatccat attcatgcaa tttaccacaa tctcaaaata 11700
taaccaccat gataaaaaat ataacagcaa ggaatgtatt acaagattca ccaaatccat 11760
tattatctgg attattcaca aatacaatga tagaagaaga tgaagaatta gctgagttcc 11820
tgatggacag gaaggttaatt ctccotagag ttgcacatga tattctagat aattctctca 11880
caggaattag aaatgccata gctggaatgt tagatagcac aaaatcacta attcgggttg 11940
gcataaatag aggagactg acatatagtt tggtgaggaa aatcagtaat tacgatctag 12000
tacaatatga aacactaagt aggactttgc gactaattgt aagtataaa atcaagtatg 12060
aagatatgtg ttcggtagac cttgccatag cattgcgaca aaagatgtgg attcatttat 12120
caggaggaag gatgataagt ggaactgaaa cgcctgacct attagaatta ctatctgggg 12180
tagtaataac aggatcagaa cattgtaaaa tatgttattc ttcagatggc acaaacccat 12240
atacttggat gtatttacc ccgtaatatca aataggatc agcagaaaaca ggtatatcgt 12300
cattaagagt tcttattttt ggatcagtca ctgatgaaag atctgaaagca caattaggat 12360
atatcaagaa tcttagttaa cctgcaaaaag ccgcaataag aatagcaatg atatatatc 12420
gggcattttg taatgatgag atatcttggg tggaaagcctc acagatagca caaacacgtg 12480
caaattttac actagatagt ctcaaaattt taacaccggg agctacatca acaaatttat 12540
cacacagatt aaaggatact gcaactcaga tgaattctc cagtacatca ttgatcagag 12600
tcagcagatt cataacaatg tccaatgata acatgtctat caaagaagct aatgaaacca 12660
aagatactaa tcttatttat caacaataa tgltaacagg attaagtgtt ttcgaatatt 12720
tatttagatt aaaagaacc acaggacaca accotatagt tagcatctg cacatagaag 12780
atgagtggtg tattaagaa agttttaatg atgaacatat taatccagag tctacattag 12840
aattaattcg atactctgaa agtaaatgat ttatttatga taaagaccca ctcaaatgat 12900
tggacttacc aaaacttatg gttataaag accattctta cacaattgat atgaattatt 12960
gggatgatac tgacatcata catgcaattt caaatgtac tgcaattaca atagcagata 13020
ctatgtcaca attagatcga gataatttaa aagagataat agttattgca aatgatgatg 13080
atattaatag cttaatcact gaattttgga ctcttgacat acttgatttt ctcaagacat 13140
ttggtggatt attagtaaat caatttgcac acactcttta tagtctaaaa atagaaggta 13200
gggatctcat ttgggattat ataatgagaa cactgagaga tacttcccat tcaatattaa 13260
aagtattatc taatgcatta tctcatccta aagtattcaa gagttctgg gattgtggag 13320
ttttaaaccc tatttatggt cctaatactg ctagtcaaga ccagataaaa cttgccctat 13380
ctatatgtga atattcacta gatctattta tgagagaatg gttgaatggt gtatcacttg 13440
aaatatacat ttgtgacagc gatatggaag ttgcaaatga taggaaacaa gcctttattt 13500
ctagacacct ttcatttggg ttgtgttag cagaattgc atcttccgga cctaacctgt 13560
taaacttaac atacttggag agacttgatc tattgaaaca atatctttaa ttaaatatta 13620
aagaagacc tactcttaaa tagtcaaaa tatctggatt ataataaaa tggttcccat 13680
caactgtaac atacgtaaga aagactgcaa tcaaatatct aaggattcgc ggtattagtc 13740
cacctgaggt aattgatgat tgggatccgg tagaagatga aaatagctg gataacattg 13800
tcaaaactat aaatgataac tgtaataaag ataataaagg gaataaaatt aacaatttct 13860
ggggactagc acttaagaac tatcaatgcc ttaaaatcag atctataaca agtgattctg 13920
atgataatga tagactagat gctaatacaa gtggtttgac acttctctca ggagggaatt 13980
atctatcgca tcaattgaga ttattcggaa tcaacagcac tagttgtctg aaagctcttg 14040
agttatcaca aattttaatg aaggagatca ataaagacaa ggacaggctc ttcctgggag 14100
aaggagcagg agctatgcta gcattgtatg atgccacatt aggcactgca gtttaattatt 14160
ataattcagg tttgaatata acagatgtaa ttggtcaacg agaattgaaa atatttctct 14220

cagaggtatc attagtaggt aaaaaattag gaaatgtgac acagattcctt aacagggtaa 14280
aagtactgtt caatgggaat cctaattcaa catggatagg aaatatggaa tgtgagagct 14340
taatatggag tgaattaaat gataagtcca ttggattagt acattgtgat atggaggag 14400
ctatcggtaa atcagaagaa actgttctac atgaacatta tagtgttata agaattacat 14460
acttgattgg ggatgatgat gttgttttag tttccaaaat tatactaca atcactccga 14520
attggtctag aatactttat ctatataaat tatattggaa agatgtaagt ataatacac 14580
tcaaaacttc taatcctgca tcaacagaat tatactaat ttcgaaagat gcatattgta 14640
ctataatgga acctagtgaa attgttttat caaaacttaa aagattgtca ctcttggaag 14700
aaaaataact attaaaatgg atcattttat caaagaagag gaataatgaa tggttacatc 14760
atgaaatcaa agaaggagaa agagattatg gaatcatgag accatatcat atggcactac 14820
aaatctttgg atttcaaatc aatttaaact atctggcgaa agaattttta tcaacccag 14880
atctgactaa tatcaacaat ataatacaaa gttttcagcg aacaataaag gatgttttat 14940
ttgaatggat taataaact catgatgata agagacataa attaggcga agatataaca 15000
tattcccact gaaaaataag ggaaagttaa gactgctatc gagaagacta gtattaaagt 15060
ggatttcatt atcattatcg actcgattac ttacaggtcg ctttctgat gaaaaattg 15120
aacatagagc acagactgga tatgtatcat tagctgatac tgatttagaa tcattaaagt 15180
tattgtcgaa aaacatcatt aagaattaca gagagtgtat aggatcaata tcattattgt 15240
ttctaacc aaaggtaaa atacttatga aattgatcgg tgggtctaaa ttattaggaa 15300
ttcccagaca atataaagaa cccgaagacc agttattaga aaactacaat caacatgatg 15360
aatttgat atcgataaac ataaatacaa tgaagatata tcctaacctt tatctttaag 15420
cctaggaata gacaaaaagt aagaaaaaca tgtaatat atataccaa cagagttcct 15480
ctctgtttg gt 15492

The claims defining the invention are as follows:

1. An attenuated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) encoding a complete reading frame or one or more antigenic determinant(s) of one or more heterologous pathogen(s) operably linked to regulatory sequences operable in said PIV genome or antigenome to form a chimeric PIV genome or antigenome,
- 5
- wherein said heterologous gene(s) or genome segment(s) is/are added as supernumerary gene(s) or genome segment(s) inserted into the partial or complete PIV vector genome or antigenome at one or more site(s) selected from the group consisting of a site between the P and M open reading frames, between the N and P open reading frames, a site between the HN and L open reading frames and a site between the 3' leader and the N open reading frame.
- 10
2. The chimeric PIV of claim 1, wherein said one or more heterologous pathogens is a heterologous PIV and said heterologous gene(s) or genome segments) encode(s) one or more PIV N, P, C, D, V, M, F, HN and/or L protein(s) or fragment(s) thereof.
- 15
3. The chimeric PIV of claim 1, wherein the vector genome or antigenome is a partial or complete human PIV (HPIV) genome or antigenome and the heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are of one or more heterologous PIV(s).
- 20
4. The chimeric PIV of claim 3, wherein said one or more heterologous PIV(s) is/are selected from HPIV1, HPIV2, or HPIV3.
5. The chimeric PIV of claim 3, wherein the vector genome or antigenome is a partial or complete HPIV genome or antigenome and the heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are of one or more heterologous HPIV(s).
- 25
6. The chimeric PIV of claim 5, wherein the vector genome or antigenome is a partial or complete HPIV3 genome or antigenome and the heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are of one or more heterologous HPIV(s).
- 30

7. The chimeric PIV of claim 6, wherein one or more gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of HPIV1 selected from HPIV1 HN and F glycoproteins and antigenic domains, fragments and epitopes thereof is/are added to the partial or complete HPIV3 genome or antigenome.
- 5 8. The chimeric PIV of claim 6, wherein the vector genome or antigenome is a partial or complete HPIV3 JS genome or antigenome and the heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are of one or more heterologous HPIV(s).
9. The chimeric PIV of claim 8, wherein one or more gene(s) or genome segment(s)
10 encoding one or more antigenic determinant(s) of HPIV1 selected from HPIV1 HN and F glycoproteins and antigenic domains, fragments and epitopes thereof is/are added to the partial or complete HPIV3 JS genome or antigenome.
10. The chimeric PIV of claim 6, wherein both HPIV1 genes encoding HN and F glycoproteins are added to a HPIV3 vector genome or antigenome.
- 15 11. The chimeric PIV of claim 5, wherein the chimeric genome or antigenome incorporates at least one and up to a full complement of attenuating mutations present within PIV3 JS *cp45* selected from mutations specifying an amino acid substitution in the L protein at a position corresponding to Tyr₉₄₂, Leu₉₉₂, or Thr₁₅₅₈ of JS *cp45*; in the N protein at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS *cp45*, in the C protein at a position corresponding to Ile₉₆ of JS *cp45*, a nucleotide substitution a 3' leader sequence of the chimeric virus at a position corresponding to nucleotide 23, 24, 28, or 45 of JS *cp45*, and/or a mutation in an N gene start sequence at a position corresponding to nucleotide 62 of JS *cp45*.
- 20 12. The chimeric PIV of claim 6, wherein one or more gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of HPIV2 is/are added to a partial or complete HPIV3 genome or antigenome.
- 25 13. The chimeric PIV of claim 12, wherein one or more HPIV2 gene(s) or genome segment(s) encoding one or more HN and/or F glycoprotein(s) or antigenic domain(s),

fragment(s) or epitope(s) thereof is/are added to a partial or complete HPIV3 vector genome or antigenome.

14. The chimeric PIV of claim 1, wherein a plurality of heterologous genes or genome segments encoding antigenic determinants of multiple heterologous PIVs are added to a partial or complete HPIV vector genome or antigenome.

15. The chimeric PIV of claim 14, wherein said plurality of heterologous genes or genome segments encode antigenic determinants from both HPIV1 and HPIV2.

16. The chimeric PIV of claim 15, wherein one or more HPIV1 gene(s) or genome segment(s) encoding one or more HN and/or F glycoprotein(s) or antigenic domain(s), fragment(s) or epitope(s) thereof and one or more HPIV2 gene(s) or genome segment(s) encoding one or more HN and/or F glycoprotein(s) or antigenic domain(s), fragment(s) or epitope(s) thereof is/are added to a partial or complete HPIV3 vector genome or antigenome.

17. The chimeric PIV of claim 3, wherein both HPIV1 genes encoding HN and F glycoproteins are substituted for counterpart HPIV3 HN and F genes to form a chimeric HPIV3-1 vector genome or antigenome which is further modified by addition of one or more gene(s) or gene segment(s) encoding one or more antigenic determinant(s) of HPIV2.

18. The chimeric PIV of claim 17, wherein a transcription unit comprising an open reading frame (ORF) of an HPIV2 HN gene is added to the chimeric HPIV3-1 vector genome or antigenome.

19. A chimeric PIV, which is rPIV3-1.2HN.

20. A chimeric PIV, which is rPIV3-lcp45.2HN.

21. The chimeric PIV of claim 1, wherein the vector genome or antigenome is a partial or complete human PIV (HPIV) genome or antigenome and the heterologous pathogen is selected from measles virus, subgroup A or subgroup B respiratory syncytical virus, mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.

22. The chimeric PIV of claim 21, wherein said one or more heterologous antigenic

determinant(s) is/are selected from measles HA and F proteins, subgroup A or subgroup B respiratory syncytial virus F, G, SH and M2 proteins, mumps virus HN and F proteins, human papilloma virus L1 protein, type 1 or type 2 human immunodeficiency virus gp160 protein, herpes simplex virus and cytomegalovirus gB, gC, gD, gE, gG, gH, gI, gJ, gK, gL, and gM proteins, rabies virus G protein, Epstein Barr Virus gp350 protein; filovirus G protein, bunyavirus G protein, Flavivirus pre M, E, and NS1 proteins, and alphavirus E protein, and antigenic domains, fragments and epitopes thereof.

23. The chimeric PIV of claim 22, wherein the vector genome or antigenome is a partial or complete HPIV3 genome or antigenome or a chimeric HPIV genome or antigenome comprising a partial or complete HPIV3 genome or antigenome having one or more gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of a heterologous HPIV added or incorporated therein.

24. The chimeric PIV of claim 1, wherein a transcription unit comprising an open reading frame (ORF) of a measles virus HA gene is added to a HPIV3 vector genome or antigenome.

25. The chimeric PIV of claim 1, wherein a transcription unit comprising an open reading frame encoding an antigenic determinant(s) of at least one of a measles virus HA and F protein or antigenic domains, fragments and epitopes thereof is added to a HPIV vector genome or antigenome.

26. A chimeric PIV of claim 1, which is selected from rPIV3(HA HN-L), rPIV3(HA N-P), rcp45L(HA N-P), rPF/3(HA P-M), or rcp45L(HA P-M).

27. The chimeric PIV of claim 24, wherein the vector genome or antigenome is a chimeric HPIV genome or antigenome comprising a partial or complete HPIV3 genome or antigenome having one or more gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of HPIV 1 added thereto.

28. The chimeric PIV of claim 25, wherein a transcription unit comprising an open reading frame (ORF) of a measles virus HA gene is added to a HPIV3-1 vector genome or antigenome having both the HPIV3 HN and F ORFs substituted by the HN and F ORFs of HPIV 1.

29. A chimeric PIV, which is rPIV3 - 1 HAp.M.

30. A chimeric PIV which is rPIV3-1 HAp.M cp45L.
31. The chimeric PIV of claim 1, wherein the vector genome or antigenome is a partial or complete HPIV3 genome or antigenome and said one or more supernumerary heterologous gene(s) or genome segment(s) are selected from HPIV 1 HN, HPIV1 F, HPIV2 HN, HPIV2 F, measles HA, and/or a translationally silent synthetic gene unit.
32. The chimeric PIV of claim 31, wherein one or both of the HPIV1 HN and/or HPIV2 HN ORF(s) is/are inserted within the HPIV3 vector genome or antigenome, respectively.
33. The chimeric PIV of claim 31, wherein the HPIV1 HN, HPIV2 HN, and measles virus HA ORFs are inserted between the N/P, P/M, and HN/L genes, respectively.
34. The chimeric PIV of claim 31, wherein the HPIV1 HN and HPIV2 HN genes are inserted between the N/P and P/M genes, respectively and a 3918-nt GU insert is added between the HN and L genes.
35. The chimeric PIV of claim 31, which is selected from rHPIV3-1HN_{N-P}, rHPIV3-1HN_{P-M}, rHPIV3-2HN_{N-P}, rHPIV3-2HN_{P-M}, rHPIV3-1HN_{N-P} 2HN_{P-M}, rHPIV3-1HN_{N-P} 2HN_{P-M} HA_{HN-L}, and rHPIV3-1HN_{N-P} 2HN_{P-M} 3918GU_{HN-L}.
36. The chimeric PIV of claim 1, which contains protective antigens from two, three or four pathogens.
37. The chimeric PIV of claim 2, which contains protective antigens from two to four pathogens selected from HPIV3, HPIV1, HPIV2, and measles virus.
38. The chimeric PIV of claim 31, wherein said one or more supernumerary heterologous gene(s) or genome segment(s) add a total length of foreign sequence to the recombinant genome or antigenome of 30% to 50% or greater compared to the wild-type HPIV3 genome length of 15,462 nt.
39. The chimeric PIV of claim 1, wherein the addition of said one or more supernumerary heterologous gene(s) or genome segment(s) specifies an attenuation phenotype of the chimeric PIV which exhibits at least a 10-to 100-fold decrease in replication in the upper and/or lower respiratory tract of a host of a host.

40. The chimeric PIV of claim 1, wherein the vector genome or antigenome is a human-bovine chimeric PIV genome or antigenome.
41. The chimeric PIV of claim 40, wherein the human-bovine chimeric vector genome or antigenome is combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of a heterologous pathogen selected from measles virus, subgroup A and subgroup B respiratory syncytial viruses, mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.
42. The chimeric PIV of claim 40, wherein the vector genome or antigenome comprises a partial or complete BPIV genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a BPIV.
43. The chimeric PIV of claim 42, wherein a transcription unit comprising an open reading frame (ORF) of a BPIV3 N ORF is substituted in the vector genome or antigenome for a corresponding N ORF of a HPIV3 vector genome.
44. The chimeric PIV of claim 43, wherein the supernumerary gene is a measles virus HA gene.
45. The chimeric PIV of claim 44, which is rHPIV3-N_B HA_{P-M}.
46. The chimeric PIV of claim 1, wherein the vector genome or antigenome comprises a partial or complete BPIV genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) from a HPIV.
47. The chimeric PIV of claim 46, wherein one or more HPIV gene(s) or genome segment(s) encoding HN and/or F glycoproteins or one or more immunogenic domain(s), fragments) or epitope(s) thereof is/are added to or incorporated within the partial or complete bovine genome or antigenome to form the vector genome or antigenome.
48. The chimeric PIV of claim 47, wherein both HPIV3 genes encoding HN and F glycoproteins are substituted for corresponding BPIV3 HN and F genes to form the vector genome or antigenome.

49. The chimeric PIV of claim 48, wherein the vector genome or antigenome is combined with at least one of a respiratory syncytial virus (RSV) F or G gene as a supernumerary gene insert.

50. A chimeric PIV which is selected from rBHPiV3-G1 or rB/HPIV3-F1.

5 51. The chimeric PIV of claim 47, wherein one or more HPIV 1 HN and/or F gene(s) or genome segment(s) encoding one or more immunogenic domain(s), fragment(s) or epitope(s) thereof are incorporated within the partial or complete bovine genome or antigenome to form the vector genome or antigenome, which is further modified by incorporation of one or more HPIV2 HN and/or F gene(s) or genome segment(s) encoding
10 one or more immunogenic domain(s), fragment(s) or epitope(s) thereof to form the chimeric genome or antigenome which expresses protective antigen(s) from both HPIV1 and HPIV2.

52. The chimeric PIV of claim 51, which is selected from rB/HPIV3.1-2F, rB/HPIV3.1-2HN, or rB/HPIV3.1-2F,2HN.

53. The chimeric PIV of claim 1, wherein the heterologous gene or gene segment
15 encodes a chimeric glycoprotein incorporating one or more antigenic domains, fragments, or epitopes of a glycoprotein of the vector PIV and of a glycoprotein of a heterologous PIV or non-PIV pathogen.

54. The chimeric PIV of claim 53, wherein the chimeric genome heterologous gene or
gene segment encodes a chimeric glycoprotein having antigenic domains, fragments, or
20 epitopes from two or more HPIVs.

55. The chimeric PIV of claim 53, wherein chimeric glycoprotein includes both a
glycoprotein ectodomain and transmembrane region of the same HPIV.

56. The chimeric PIV of claim 53, wherein said chimeric glycoprotein is a chimeric HPIV
glycoprotein HN or a chimeric HPIV F glycoprotein.

25 57. The chimeric PIV of claim 53, wherein the PIV vector genome or antigenome is a
partial HPIV3 genome or antigenome, and the second, antigenically distinct PIV is selected
from HPIV1 or HPIV2.

58. The chimeric PIV of claim 53, wherein the HPIV vector genome or antigenome is a

partial HPIV3 genome or antigenome and the second, antigenically distinct HPIV is HPIV2.

59. The chimeric PIV of claim 53, wherein a heterologous gene or gene segment encodes a HPIV's HN glycoprotein having an ectodomain of HPIV2 glycoprotein and a heterologous gene or gene segment encodes a HPIV3 F glycoprotein having an ectodomain of a HPIV2 F glycoprotein.

60. A chimeric PIV, which is rPF/3-2TM.

61. The chimeric PIV of claim 53, which is further modified to incorporate one or more and up to a full panel of attenuating mutations identified in HPIV3 JS cp45.

62. A chimeric PIV, which is rPIV3-2TMcp45.

10 63. The chimeric PIV of claim 53 in which the heterologous gene or gene segment encodes, PIV2 ectodomain and transmembrane regions of one or both HN and/or F glycoproteins fused to the corresponding PIV3 cytoplasmic tail region(s).

15 64. The chimeric PIV of claim 63, wherein ectodomain and transmembrane regions of both PIV2 HN and F glycoproteins are fused to corresponding PIV3 HN and F cytoplasmic tail regions.

65. A chimeric PIV which is rPIV3-2CT.

66. The chimeric PIV of claim 65, which is further modified to incorporate one or more and up to a full panel of attenuating mutations identified in HPIV3 JSep45.

67. A chimeric PIV which is rPIV3-2CTcp45.

20 68. The chimeric PIV of claim 53, which is further modified to incorporate one or more and up to a full panel of attenuating mutations identified in HPIV3 JS cp45 selected from mutations specifying an amino acid substitution in the L protein at a position corresponding to Tyr₉₄₂, Leu₉₉₂, or Thr₁₅₅₈ of JS cp45, in the N protein at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS cp45, in the C protein at a position corresponding to Ile₉₆ of JS cp45, a
25 nucleotide substitution in a 3' leader sequence of the chimeric virus at a position corresponding to nucleotide 23, 24, 28, or 45 of JS cp45, and/or a mutation in an N gene start sequence at a position corresponding to nucleotide 62 of JS cp45.

69. The chimeric PIV of claim 53, wherein a plurality of heterologous genes or genome segments encoding antigenic determinants of multiple heterologous PIVs are added to the partial or complete HPIV vector genome or antigenome.
70. The chimeric PIV of claim 69, wherein said plurality of heterologous genes or genome segments encode antigenic determinants from both HPIV1 and HPIV2.
71. The chimeric PIV of claim 53, wherein the chimeric PIV genome or antigenome is attenuated by addition or incorporation of one or more gene(s) or genome segment(s) from a bovine PIV3 (BPIV3).
72. The chimeric PIV of claim 71, wherein the gene or gene segment encodes a BPIV3 N protein.
73. The chimeric PIV of claim 53, wherein the chimeric genome or antigenome is modified by introduction of an attenuating mutation involving an amino acid substitution of phenylalanine at position 456 of the HPIV3 L protein.
74. The chimeric PIV of claim 73, wherein phenylalanine at position 456 of the HPIV3 L protein is substituted by leucine.
75. The chimeric PIV of claim 53, wherein the heterologous gene(s) or genome segment(s) encodes one or more antigenic determinants of a glycoprotein from a respiratory syncytial virus (RSV) or measles virus.
76. The chimeric PIV of claim 1, wherein the chimeric genome or antigenome is modified by addition or substitution of one or more heterologous gene(s) or genome segment(s) that confer increased genetic stability or that alter attenuation, reactogenicity in vivo, or growth in culture of the chimeric virus.
77. The chimeric PIV of claim 1, wherein the chimeric genome or antigenome is further modified by introduction of one or more attenuating mutations identified in a biologically derived mutant PIV or other mutant nonsegmented negative stranded RNA virus.
78. The chimeric PIV of claim 77, wherein the chimeric genome or antigenome incorporates at least one and up to a full complement of attenuating mutations present within PIV3 JS cp45.

79. The chimeric PIV of claim 77, wherein the chimeric genome or antigenome incorporates at least one and up to a full complement of attenuating mutations specifying an amino acid substitution in the L protein at a position corresponding to Tyr₉₄₂, Leu₉₉₂, or Thr₁₅₅₈ of JS cp45; in the N protein at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS cp45,
5 in the C protein at a position corresponding to Ile₉₆ of JS₁₀, in the F protein at a position corresponding to residues Ile₄₂₀ or Ala₄₅₀ of JS cp, in the HN protein at a position corresponding to residue Val₃₈₄ of JS cp45, a nucleotide substitution a 3' leader sequence of the chimeric virus at a position corresponding to nucleotide 23, 24, 28, or 45 of JS cp45, and/or a mutation in an N gene start sequence at a position corresponding to nucleotide 62 of
10 JS cp45.

80. The chimeric PIV of claim 77, wherein said attenuating mutation comprises an amino acid substitution of phenylalanine at position 456 of the HPIV3 L protein.

81. The chimeric PIV of claim 77, wherein the chimeric genome or antigenome includes at least one attenuating mutation stabilized by multiple nucleotide changes in a codon specifying the mutation.
15

82. The chimeric PIV of claim 1, wherein the chimeric genome or antigenome comprises an additional nucleotide modification specifying a phenotypic change selected from a change in growth characteristics, attenuation, temperature-sensitivity, cold-adaptation, plaque size, host-range restriction, or a change in immunogenicity.

83. The chimeric PIV of claim 82, wherein the additional nucleotide modification alters one or more PIV N, P, C, D, V, M, F, HN and/or L genes and/or a 3' leader, 5' trailer, and/or intergenic region within the vector genome or antigenome or within the heterologous gene(s) or gene segment(s).
20

84. The chimeric PIV of claim 83, wherein one or more PIV gene(s) is deleted in whole or in part or expression of the gene(s) is reduced or ablated by a mutation in an RNA editing site, by a frameshift mutation, by a mutation that alters an amino acid specified by an initiation codon, or by introduction of one or more stop codons in an open reading frame (ORF) of the gene.
25

85. The chimeric PIV of claim 84, wherein the additional nucleotide modification

comprises a partial or complete deletion of one or more C, D or V ORF(s) or one or more nucleotide change(s) that reduces or ablates expression of said one or more C, D or V ORF(s).

86. The chimeric PIV of claim 1, wherein the chimeric genome or antigenome comprises a heterologous gene or gene segment encoding a cytokine.
- 5 87. The chimeric PIV of claim 1, which incorporates a heterologous gene or genome segment from respiratory syncytial virus (RSV).
88. The chimeric PIV of claim 87, wherein the heterologous gene or genome segment encodes RSV F and/or G glycoprotein(s) or immunogenic domain(s), fragment(s), or epitope(s) thereof.
- 10 89. The chimeric PIV of claim 1 which is a subviral particle.
90. A method for stimulating the immune system of an individual to induce protection against PIV which comprises administering to the individual an immunologically sufficient amount of the chimeric PIV of claim 1 combined with a physiologically acceptable carrier.
- 15 91. The method of claim 90, wherein the chimeric PIV is administered in a dose of 10^3 to 10^7 PFU.
92. The method of claim 91, wherein the chimeric PIV is administered to the upper respiratory tract.
93. The method of claim 90, wherein the chimeric PIV is administered by spray, droplet or aerosol.
- 20 94. The method of claim 90, wherein the vector genome or antigenome is of human PIV3 (HPIV3) and the chimeric PIV elicits an immune response against HPIV1 and/or HPIV2.
95. The method of claim 90, wherein the chimeric PIV elicits a polyspecific immune response against multiple human PIVs and/or against a human PIV and a non-PIV pathogen.
- 25 96. The method of claim 90, wherein the vector genome or antigenome is a partial or

complete human PIV (HPIV) genome or antigenome and the heterologous pathogen is selected from measles virus, subgroup A and subgroup B respiratory syncytial viruses, mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, 5 filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.

97. The method of claim 90, wherein the chimeric PIV elicits a polyspecific immune response against a human PIV (HPIV) and measles virus.

98. The method of claim 97, wherein the chimeric PIV elicits a polyspecific immune response against HPIV3 and measles virus.

10 99. The method of claim 90, wherein a first, chimeric PIV according to claim 1 and a second PIV are administered sequentially or simultaneously to elicit a polyspecific immune response.

100. The method of claim 99, wherein the second PIV is a second, chimeric PIV according to claim 1.

15 101. The method of claim 99, wherein the first, chimeric PIV and second PIV are administered simultaneously in a mixture.

102. The method of claim 99, wherein the first, chimeric PIV and second PIV are antigenically distinct variants of HPIV.

103. The method of claim 99, wherein the first, chimeric PIV comprises a partial or 20 complete HPIV3 vector genome or antigenome.

104. The method of claim 100, wherein the first chimeric PIV and second chimeric PIV each incorporate one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of a non-PIV pathogen.

105. The method of claim 104, wherein the first and second chimeric PIV incorporate 25 one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of the same non-PIV pathogen.

106. A method for sequential immunization to stimulate the immune system of an individual to induce protection against multiple pathogens comprising administering to a newborn to 4 month old infant an immunologically sufficient amount of a first attenuated chimeric HPIV expressing an antigenic determinant of a non-PIV pathogen and one or more antigenic determinants of HPIV3 and subsequently administering an immunologically sufficient amount of a second attenuated chimeric HPIV expressing an antigenic determinant of a non-PIV pathogen and one or more antigenic determinants of HPIV1 or HPIV2.

107. The method for sequential immunization of claim 106, wherein the first attenuated chimeric HPIV is an HPIV3 expressing a measles virus antigenic determinant and wherein the second attenuated chimeric HPIV is a PIV3-1 chimeric virus expressing a measles virus antigenic determinant and incorporating one or more attenuating mutations of HPIV3 JS *cp45*.

108. The method for sequential immunization of claim 106, wherein following the first vaccination, the vaccinee elicits a primary antibody response against both PIV3 and the non-PIV pathogen, but not HPIV1 or HPIV2, and upon secondary immunization the vaccinee is readily infected with the second attenuated HPIV and develops both a primary antibody response to HPIV1 or HPIV2 and a high titered secondary antibody response against the non-PIV pathogen.

109. The method for sequential immunization of claim 106, wherein the first chimeric PIV elicits an immune response against HPIV3 and the second chimeric PIV elicits an immune response against HPIV 1 or HPIV2, and wherein both the first and second chimeric PIVs elicit an immune response against measles or RSV.

110. The method for sequential immunization of claim 106, wherein the non-PIV pathogen is selected from measles virus, subgroup A and subgroup B respiratory syncytial viruses (RSVs), mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.

111. The method for sequential immunization of claim 106, wherein the second chimeric

PIV comprises a partial or complete HPIV3 vector genome or antigenome combined with one or more gene(s) or genome segment(s) encoding one or more HPIV 1 and/or HPIV2 HN and/or F glycoprotein(s) or antigenic domain(s), fragment(s) or epitope(s) thereof.

112. The method for sequential immunization of claim 106, wherein the partial or complete vector genome or antigenome of the first, chimeric PIV incorporates at least one and up to a full complement of attenuating mutations present within HPIV3 JS cp45 selected from mutations specifying an amino acid substitution in the L protein at a position corresponding to Tyr₉₄₂, Leu₉₉₂, or Thr₁₅₃₈ of JS cp45; in the N protein at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS cp45, in the C protein at a position corresponding to Ile₉₆ of JS cp45, a nucleotide substitution a 3' leader sequence of the chimeric virus at a position corresponding to nucleotide 23, 24, 28, or 45 of JS cp45, and/or a mutation in an N gene start sequence at a position corresponding to nucleotide 62 of JS cp45.

113. An immunogenic composition to elicit an immune response against PIV comprising an immunogenically sufficient amount of the chimeric PIV of claim 1 in a physiologically acceptable carrier.

114. The immunogenic composition of claim 113, formulated in a dose of 10^3 to 10^7 PFU.

115. The immunogenic composition of claim 113, formulated for administration to the upper respiratory tract by spray, droplet or aerosol.

116. The immunogenic composition of claim 113, wherein the chimeric PIV elicits an immune response against one or more virus(es) selected from HPIV1, HPIV2 and HPIV3.

117. The immunogenic composition of claim 113, wherein the chimeric PIV elicits an immune response against HPIV3 and another virus selected from HPIV1 and HPIV2.

118. The immunogenic composition of claim 113, wherein the chimeric PIV elicits a polyspecific immune response against one or more HPIVs and a heterologous pathogen selected from measles virus, subgroup A and subgroup B respiratory syncytial viruses, mumps virus, human papilloma viruses, type I and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.

119. The immunogenic composition of claim 118, wherein the chimeric PIV elicits a polyspecific immune response against HPIV3 and measles or respiratory syncytial virus.

120. The immunogenic composition of claim 113, further comprising a second, chimeric PIV according to claim 1.

5 121. The immunogenic composition of claim 120, wherein the first and second chimeric PIVs are antigenically distinct variants of HPIV and bear the same or different heterologous antigenic determinant(s).

122. The immunogenic composition of claim 120, wherein the first chimeric PIV comprises a partial or complete HPIV3 genome or antigenome and one or more
10 heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of a non-PIV heterologous pathogen.

123. The immunogenic composition of claim 120, wherein the second chimeric PIV incorporates one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinants of the same non-PIV heterologous pathogen.

15 124. The immunogenic composition of claim 120, wherein the first chimeric PIV elicits an immune response against HPIV3 and the second chimeric PIV elicits an immune response against HPIV1 or HPIV2, and wherein both the first and second chimeric PIVs elicit an immune response against the non-PIV pathogen.

125. The immunogenic composition of claim 120, wherein the heterologous pathogen
20 is selected from measles virus, subgroup A and subgroup B respiratory syncytial viruses (RSVs), mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Ban virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.

25 126. The immunogenic composition of claim 120, wherein the heterologous pathogen is selected from measles virus or RSV.

127. The immunogenic composition of claim 120, wherein the second chimeric PIV

comprises a partial HPIV3 vector genome or antigenome and the heterologous genes(s) or genome segment(s) are one or more HPIV1 gene(s) or genome segment(s) encoding one or more antigenic determinants of HPIV1 HN and/or F glycoproteins.

128. The immunogenic composition of claim 120, wherein the second chimeric PIV
5 compresses a partial or complete HPIV3 vector genome or antigenome and the heterologous gene(s) or genome segment(s) are one or more gene(s) or genome segment(s) encoding one or more HPIV2 HN and/or F glycoprotein(s) or antigenic domain(s), fragment(s) or epitope(s) thereof.

129. An isolated polynucleotide comprising a chimeric PIV genome or antigenome which
10 includes a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) encoding a complete reading frame or one or more antigenic determinant(s) of one or more heterologous pathogen(s) operably linked to regulatory sequences operable in said PIV genome or antigenome to form a chimeric PIV genome or antigenome wherein said heterologous gene(s) or genome segment(s) encoding the
15 antigenic determinant(s) is/are added as supernumerary gene(s) or genome segment(s) inserted into said PIV genome or antigenome at one or more site(s) selected from the group consisting of a site between the P and M open reading frames, a site between the N and P open reading frames, a site between the HN and L open reading frames and a site between the 3' leader and the N open reading frame. of the partial or complete PIV vector genome or antigenome and an
20 infectious virus or subviral particle comprising said polynucleotide is attenuated.

130. The isolated polynucleotide of claim 129, wherein said one or more heterologous pathogens is a heterologous PIV and said heterologous gene(s) or genome segment(s) encode(s) one or more PIV N, P, C, D, V, M, F, HN and/or L protein(s) or immunogenic fragment(s), domain(s), or epitope(s) thereof.

131. The isolated polynucleotide of claim 129, wherein the vector genome or antigenome is
25 a partial or complete human PIV (HPIV) genome or antigenome and the heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are of one or more heterologous PIV(s).

132. The isolated polynucleotide of claim 131, wherein the vector genome or antigenome
30 is a partial or complete HPIV3 genome or antigenome and the heterologous gene(s) or

genome segment(s) encoding the antigenic determinant(s) is/are of HPIV 1 and/or HPIV2.

133. The isolated polynucleotide of claim 129, wherein the vector genome or antigenome is a partial or complete human PIV (HPIV) genome or antigenome and the heterologous pathogen is selected from measles virus subgroup A and subgroup B respiratory syncytial viruses, mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses, herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.

134. The isolated polynucleotide of claim 133, wherein said one or more heterologous antigenic determinant(s) is/are selected from measles virus HA and F proteins, subgroup A or subgroup B respiratory syncytial virus F, G, SH and M2 proteins, mumps virus HN and F proteins, human papilloma virus L1 protein, type 1 or type 2 human immunodeficiency virus gp160 protein, herpes simplex virus and cytomegalovirus gB, gC, gD, E, gG, gH, gI, gJ, gK, gL, and gM proteins, rabies virus G protein, Epstein Barr Virus gp350 protein; filovirus G protein, bunyavirus G protein, Flavivirus E and NS1 proteins, and alphavirus E protein, and antigenic domains, fragments and epitopes thereof.

135. The isolated polynucleotide of claim 129, wherein the vector genome or antigenome is a partial or complete HPIV3 genome or antigenome or a chimeric HPIV genome or antigenome comprising a partial or complete HPIV3 genome or antigenome having one or more gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of a heterologous HPIV added or incorporated therein.

136. The isolated polynucleotide of claim 135, wherein the heterologous pathogen is measles virus and the heterologous antigenic determinant(s) is/are selected from the measles virus HA and F proteins and antigenic domains, fragments and epitopes thereof.

137. The isolated polynucleotide of claim 129, wherein a transcription unit comprising an open reading frame (ORF) of a measles virus HA gene is added to a HPIV3 vector genome or antigenome.

138. The isolated polynucleotide of claim 135, wherein a transcription unit comprising an open reading frame (ORF) of a measles virus HA gene is added to or incorporated within a HPIV3-1 vector genome or antigenome having both the HPIV3 HN and F ORFs substituted

by the HN and F ORFs of HPIV1.

139. The isolated polynucleotide of claim 129, wherein the vector genome or antigenome is a partial or complete HPIV3 genome or antigenome and said one or more supernumerary heterologous gene(s) or genome segment(s) are selected from HPIV1 HN, HPIV1 F, HPIV2
5 HN, HPIV2 F, measles HA, and/or a translationally silent synthetic gene unit.

140. The isolated polynucleotide of claim 129, wherein two or all of the HPIV1 HN, HPIV2 HN, and measles virus HA ORFs are added to the vector genome or antigenome.

141. The isolated polynucleotide of claim 129, wherein two or all of the HPIV1 HN and HPIV2 HN genes and a 3918-nt GU insert is/are added to the vector genome or antigenome.

10 142. The isolated polynucleotide of claim 129, wherein said one or more supernumerary heterologous gene(s) or genome segment(s) add a total length of foreign sequence to the recombinant genome or antigenome of 30% to 50% or greater compared to the wild-type HPIV3 genome length of 15,462 nt.

15 143. The isolated polynucleotide of claim 129, wherein the vector genome or antigenome is a human-bovine chimeric PIV genome or antigenome.

144. The isolated polynucleotide of claim 143, wherein the human-bovine chimeric vector genome or antigenome is combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of a heterologous pathogen selected from measles virus, subgroup A and subgroup B respiratory syncytial viruses, mumps virus, human papilloma viruses, type 1 and type 2 human immunodeficiency viruses,
20 herpes simplex viruses, cytomegalovirus, rabies virus, Epstein Barr virus, filoviruses, bunyaviruses, flaviviruses, alphaviruses and influenza viruses.

145. The isolated polynucleotide of claim 129, wherein a transcription unit comprising an open reading frame (ORF) of a BPIV3 N ORF is substituted in the vector genome or
25 antigenome for a corresponding N ORF of a HPIV3 vector genome.

146. The isolated polynucleotide of claim 145, wherein the vector genome or antigenome is combined with a measles virus HA gene as a supernumerary gene insert.

147. The isolated polynucleotide of claim 129, wherein one or more HPIV gene(s) or genome segment(s) encoding HN and/or F glycoproteins or one or more immunogenic domain(s), fragment(s) or epitope(s) thereof is/are added to or incorporated within the partial or complete bovine genome or antigenome to form the vector genome or antigenome.

5 148. The isolated polynucleotide of claim 147, wherein both HPIV3 genes encoding HN and F glycoproteins are substituted for corresponding BPIV3 HN and F genes to form the vector genome or antigenome.

149. The isolated polynucleotide of claim 129, wherein the vector genome or antigenome is combined with a respiratory syncytial virus (RSV) F or G gene as a supernumerary gene
10 insert.

150. The isolated polynucleotide of claim 129, wherein the heterologous gene or genome segment encodes a chimeric glycoprotein having antigenic domains, fragments, or epitopes from both a human PIV (HPIV) and a heterologous pathogen.

151. The isolated polynucleotide of claim 129, wherein the heterologous gene or genome
15 segment encodes a chimeric glycoprotein having antigenic domains, fragments, or epitopes from two or more different PIVs.

152. The isolated polynucleotide of claim 129, wherein the chimeric genome or antigenome is modified by introduction of one or more attenuating mutations identified in a biologically derived mutant PIV or other mutant nonsegmented negative stranded RNA virus.

153. The isolated polynucleotide of claim 129, wherein, the chimeric genome or antigenome incorporates at least one and up to a full complement of attenuating mutations present within PIV3 JS *cp45*.
20

154. The isolated polynucleotide of claim 129, wherein the chimeric genome or antigenome incorporates an attenuating mutation from a heterologous nonsegmented negative stranded
25 RNA virus.

155. The isolated polynucleotide of claim 129, wherein the chimeric genome or antigenome comprises an additional nucleotide modification specifying a phenotypic change selected from a change in growth characteristics, attenuation, temperature-sensitivity, cold-

adaptation, plaque size, host-range restriction, or a change in immunogenicity.

156. The isolated polynucleotide of claim 129, wherein the additional nucleotide modification alters one or more PIV N, P, C, D, V, M, F, HN and/or L genes and/or a 3' leader, 5' trailer, and/or intergenic region within the vector genome or antigenome or within the
5 heterologous gene(s) or gene segment(s).

157. The isolated polynucleotide of claim 129, wherein one or more PIV gene(s) is deleted in whole or in part or expression of the gene(s) is reduced or ablated by a mutation in an RNA editing site, by a frameshift mutation, by a mutation that alters an amino acid specified by an initiation codon, or by introduction of one or more stop codons in an open reading frame
10 (ORF) of the gene.

158. A method for producing an infectious attenuated chimeric PIV particle from one or more isolated polynucleotide molecules encoding said PIV, comprising:

expressing in a cell or cell-free lysate an expression vector comprising an isolated polynucleotide comprising a partial or complete PIV vector genome or antigenome of a
15 human or bovine PIV combined with one or more heterologous gene(s) or genome segment(s) encoding a complete reading frame or one or more antigenic determinant(s) of one or more heterologous pathogen(s) operatively linked to regulatory sequences operable in said PIV to form a chimeric PIV genome or antigenome, and PIV N, P, and L proteins

wherein said one or more heterologous gene(s) or genome segment(s) encoding the
20 antigenic determinant(s) is/are added as supernumerary gene(s) or genome segment(s) inserted into the PIV genome or antigenome at a site selected from the group consisting of a site between the M and P open reading frames, a site between the N and P open reading frames, a site between the HN and L open reading frames and a site between the 3' leader and the N open reading frame of the partial or complete PIV vector genome or antigenome.

25 159. The method of claim 158, wherein the chimeric PIV genome or antigenome and the N, P, and L proteins are expressed by two or more different expression vectors.

160. An expression vector comprising an operably linked transcriptional promoter, a polynucleotide sequence which includes a partial or complete PIV vector genome or antigenome of a human or bovine PIV combined with one or more heterologous gene(s) or
30 genome segment(s) encoding a complete reading frame or one or more antigenic determinant(s) of one or more heterologous pathogen(s) operably linked to regulatory

sequences operable in said PIV to form a chimeric PIV genome or antigenome wherein said one or more heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are added as supernumerary gene(s) or genome segment(s) inserted into the PIV genome or antigenome at a site selected from the group consisting of a site between the
 5 M and P open reading frames, a site between the N and P open reading frames, a site between the HN and L open reading frames and a site between the 3' leader and the N open reading frame of the partial or complete PIV vector genome or antigenome, and a transcriptional terminator.

161. An infectious parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein,
 10 a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a PIV genome or antigenome having a polynucleotide insertion of between 150 nucleotides (nts) and 4,000 nucleotides in length in a non-coding region (NCR) of the genome or antigenome or as a separate gene unit (GU), said polynucleotide insertion lacking a complete open reading frame (ORF) and specifying an attenuated phenotype in said recombinant PIV.

15 162. The infectious PIV of claim 161, wherein said polynucleotide insert is introduced into the PIV genome or antigenome in a reverse, non-sense orientation whereby the insert does not encode protein.

163. The infectious PIV of claim 161, wherein said polynucleotide insert is approximately 2,000 nts or greater in length.

20 164. The infectious PIV of claim 161, wherein said polynucleotide insert is approximately 3,000 nts or greater in length.

165. The infectious PIV of claim 161, wherein said recombinant PIV replicates efficiently in vitro and exhibits an attenuated phenotype in vivo

25 166. A PIV according to claim 1 or 161 substantially as hereinbefore described, with reference to the Examples and Figures.

167. A method according to claim 106 or 158 substantially as hereinbefore described.

168. A composition according to claim 113 substantially as hereinbefore described.

169. A polynucleotide according to claim 129 substantially as hereinbefore described.

170. A vector according to claim 160 substantially as hereinbefore described.

171. An attenuated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a partial or complete PIV vector genome or antigenome, that is a HPIV3-1 vector genome or antigenome having both the HPIV3 HN and F ORFs substituted by the HN and F ORFs of HPIV 1 and further comprising an open reading frame (ORF) of a measles virus HA gene inserted into the PIV genome or antigenome at a site selected from the group consisting of a site between the M and P open reading frames, a site between the N and P open reading frames, a site between the HN and L open reading frames and a site between the 3' leader and the N open reading frame.
172. An attenuated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), a partial or complete PIV vector genome or antigenome, and further comprising a translationally silent synthetic gene unit.
173. An attenuated infectious chimeric parainfluenza virus (PIV) comprising a major nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large polymerase protein (L), and a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) encoding one or more antigenic determinant(s) of one or more heterologous pathogen(s) to form a chimeric PIV genome or antigenome, wherein said one or more heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) is/are added as supernumerary gene(s) or genome segment(s) inserted into the PIV genome or antigenome at a site selected from the group consisting of a site between the M and P open reading frames, a site between the N and P open reading frames, a site between the HN and L open reading frames and a site between the 3' leader and the N open reading frame, and wherein the addition of said one or more supernumerary heterologous gene(s) or genome segment(s) specifies an attenuation phenotype of the chimeric PIV which exhibits at least a 10-to 100-fold decrease in replication in the upper and/or lower respiratory tract of a host.
174. The attenuated infectious chimeric parainfluenza virus (PIV) of claim 1, wherein a transcription unit comprising an open reading frame (ORF) of a BPIV3 N ORF is substituted in the vector genome or antigenome for a corresponding N ORF of a

HPIV3 vector genome.

175. The attenuated infectious chimeric parainfluenza virus (PIV) of claim 1, wherein the vector genome or antigenome encodes a chimeric glycoprotein incorporating one or more antigenic domains, fragments, or epitopes of a glycoprotein of the vector PIV and of a glycoprotein of a heterologous PIV or non-PIV pathogen, and the chimeric PIV genome or antigenome is modified by introduction of an attenuating mutation involving an amino acid substitution of phenylalanine at position 456 of the HPIV3 L protein.

176. The attenuated infectious chimeric PIV of claim 175, wherein the phenylalanine at position 456 of the L protein is substituted by leucine.

177. The attenuated infectious chimeric parainfluenza virus (PIV) of claim 1, wherein the chimeric PIV genome or antigenome is modified by introduction of an attenuating mutation involving an amino acid substitution of phenylalanine at position 456 of the HPIV3 L protein.

178. The attenuated infectious chimeric PIV of claim 177, wherein the phenylalanine at position 456 of the L protein is substituted by leucine.

179. The attenuated infectious chimeric parainfluenza virus (PIV) of claim 1, wherein one or more PIV gene(s) in the vector genome or antigenome is deleted in whole or in part or expression of the gene(s) is reduced or ablated by a mutation in an RNA editing site, by a frameshift mutation, by a mutation that alters an amino acid specified by an initiation codon.

180. An isolated polynucleotide comprising a chimeric PIV genome or antigenome which includes a partial or complete PIV vector genome or antigenome combined with one or more heterologous gene(s) or genome segment(s) that is a transcriptionally silent synthetic gene unit to form a chimeric PIV genome or antigenome.

181. An isolated polynucleotide of claim 180, wherein a transcription unit comprising an open reading frame (ORF) of a BPIV3 N ORF is substituted in the vector genome or antigenome for a corresponding N ORF of a HPIV3 vector genome.

182. The isolated polynucleotide of claim 180, further comprising a measles virus HA gene as a supernumerary gene insert.

183. The isolated polynucleotide of claim 129, wherein one or more PIV gene(s) is deleted

in whole or in part or expression of the gene(s) is reduced or ablated by a mutation in an RNA editing site, by a frameshift mutation, by a mutation that alters an amino acid specified by an initiation codon.

184. The isolated polynucleotide of claim 180, in which the transcriptionally silent gene unit is inserted adjacent to or within a noncoding region of the partial or complete PIV vector genome or antigenome.

185. The isolated polynucleotide of claim 180, wherein the vector genome or antigenome is a partial or complete human PIV (HPIV) genome or antigenome and the chimeric genome or antigenome further comprise a heterologous gene(s) or genome segment(s) encoding the antigenic determinant(s) of one or more heterologous PIV(s).

186. The isolated polynucleotide of claim 180, wherein the chimeric PIV genome or antigenome further comprises a gene(s) or genome segment(s) encode(s) one or more of a heterologous PIV N, P, C, D, V, M, F, HN and/or L protein(s) or antigenic determinant(s) thereof.

187. The isolated polynucleotide of claim 186, wherein the chimeric PIV genome or antigenome further comprises a gene(s) or genome segment(s) encode(s) one or more of at least one heterologous PIV F and/or HN protein(s) or antigenic determinants(s) thereof.

188. An isolated polynucleotide comprising a chimeric PIV genome or antigenome which includes a partial or complete PIV vector genome or antigenome combined with one or more polynucleotide insertions of between 150 nucleotides (nts) and 4,000 nucleotides in length in a non-coding region (NCR) of the genome or antigenome or as a separate gene unit (GU), said polynucleotide insertion lacking a complete open reading frame (ORF) and specifying an attenuated phenotype in said recombinant PIV.

189. The isolated polynucleotide of claim 188, wherein said polynucleotide insert is introduced into the PIV genome or antigenome in a reverse, non-sense orientation whereby the insert does not encode protein.

190. The isolated polynucleotide of claim 188, wherein said polynucleotide insert is approximately 2,000 nts or greater in length.

191. The isolated polynucleotide of claim 188, wherein said polynucleotide insert is approximately 3,000 nts or greater in length.

192. The isolated polynucleotide of claim 188, wherein said recombinant PIV replicates efficiently in vitro and exhibits an attenuated phenotype in vivo.

2
3
4

5
6
7
8





Fig. 1A

SUBSTITUTE SHEET (RULE 26)

MEASLES HA INSERT FOR THE HN-L JUNCTION



2/24




Fig. 1B

SUBSTITUTE SHEET (RULE 26)

3/24




Fig. 2

SUBSTITUTE SHEET (RULE 26)

4/24




Fig. 3

SUBSTITUTE SHEET (RULE 26)




Fig. 4

SUBSTITUTE SHEET (RULE 26)

6/24




Fig. 5

SUBSTITUTE SHEET (RULE 26)

7/24




Fig. 6

SUBSTITUTE SHEET (RULE 26)

8/24





Fig. 7

SUBSTITUTE SHEET (RULE 26)



SUBSTITUTE SHEET (RULE 26)

Fig. 8

10/24




Fig. 9A

SUBSTITUTE SHEET (RULE 26)

11/24




Fig. 9B

SUBSTITUTE SHEET (RULE 26)

12/24




Fig. 9C

SUBSTITUTE SHEET (RULE 26)




Fig. 10

SUBSTITUTE SHEET (RULE 26)

14/24




Fig. 11

SUBSTITUTE SHEET (RULE 26)




Fig. 12

SUBSTITUTE SHEET (RULE 26)

17/24




Fig. 14

SUBSTITUTE SHEET (RULE 26)

RECOMBINANT BOVINE/HUMAN PIV3.1 EXPRESSING HPIV2 F AND HN FROM SUPERNUMERARY GENES




Fig. 15

SUBSTITUTE SHEET (RULE 26)

19/24




Fig. 16

SUBSTITUTE SHEET (RULE 26)

20/24




Fig. 17

SUBSTITUTE SHEET (RULE 26)

21/24




Fig. 18

SUBSTITUTE SHEET (RULE 26)

22/24




Fig. 19

SUBSTITUTE SHEET (RULE 26)

24/24




Fig. 21

SUBSTITUTE SHEET (RULE 26)