



(51) International Patent Classification:

C07C 17/08 (2006.01) C07C 19/10 (2006.01)
C07C 17/42 (2006.01) C07C 21/19 (2006.01)

(21) International Application Number:

PCT/US2016/033450

(22) International Filing Date:

20 May 2016 (20.05.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/164,631 21 May 2015 (21.05.2015) US

(71) Applicant: **THE CHEMOURS COMPANY FC, LLC**
[US/US]; 1007 Market Street, Wilmington, Delaware
19801 (US).

(72) Inventors: **SUN, Xuehui**; 761 Meadowbank Rd, Kennett
Square, Pennsylvania 19348 (US). **NAPPA, Mario
Joseph**; 3 Oakridge Court, Newark, Delaware 19711 (US).
KRAUSE, Karl; 572 Unionville Rd, Kennett Square,
Pennsylvania 19348 (US).

(74) Agent: **GROLZ, Edward W.**; Scully, Scott, Murphy &
Presser, 400 Garden City Plaza, Suite 300, Garden City,
New York 11530 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: HYDROFLUORINATION OF 1233xf TO 244bb BY SbF₅

(57) Abstract: The disclosure relates to a method for hydrofluorination of an olefin of the formula: RCX=CYZ to produce a hydro-fluoroalkane of formula RCXFCHYZ or RCHXCFYZ, wherein X, Y, and Z are independently the same or different and are selected from the group consisting of H, F, Cl, Br, and C₁-C₆ alkyl which is partially or fully substituted with chloro or fluoro or bromo; and R is a C₁-C₆ alkyl which is unsubstituted or substituted with chloro or fluoro or bromo, comprising reacting the olefin with HF in the liquid-phase, in the presence of SbF₅, at a temperature ranging from about -30°C to about 65°C.



WO 2016/187507 A1

HYDROFLUORINATION OF 1233xf TO 244bb BY SbF₅**BACKGROUND OF THE DISCLOSURE**

[0001] This disclosure relates to novel methods for preparing fluorinated organic compounds, and more particularly to methods of producing fluorinated hydrocarbons.

[0002] Hydrofluorocarbons (HFCs), in particular hydrofluoroalkenes or fluoroolefins, such as tetrafluoropropenes (including 2,3,3,3-tetrafluoro-1-propene (HFO-1234yf or 1234yf)) have been disclosed to be effective refrigerants, fire extinguishants, heat transfer media, propellants, foaming agents, blowing agents, gaseous dielectrics, sterilant carriers, polymerization media, particulate removal fluids, carrier fluids, buffing abrasive agents, displacement drying agents and power cycle working fluids. Unlike chlorofluorocarbons (CFCs) and hydrochlorofluorocarbons (HCFCs), both of which potentially damage the Earth's ozone layer, HFCs do not contain chlorine and, thus, pose no threat to the ozone layer.

[0003] In addition to ozone depleting concerns, global warming is another environmental concern in many of these applications. Thus, there is a need for compositions that meet both low ozone depletion standards as well as having low global warming potentials. Certain fluoroolefins are believed to meet both goals. Thus, there is a need for manufacturing processes that provide halogenated hydrocarbons and fluoroolefins that contain no chlorine that also have a low global warming potential.

[0004] One such HFO is 2,3,3,3-tetrafluoro-1-propene (HFO-1234yf or 1234yf). The preparation of HFO-1234yf starting from $CQ_2=CCl-CH_2Q$ or $CQ_3-CCl=CH_2$ or $CQ_3-CHCl-CH_2Q$ may include three reaction steps, as follows:

- (i) $(CQ_2=CCl-CH_2Q \text{ or } CQ_3-CCl=CH_2 \text{ or } CQ_3-CHCl-CH_2Q) + HF \rightarrow$ 2-chloro-3,3,3-trifluoropropene (HCFO-1233xf or 1233xf) + HCl in a vapor phase reactor charged with a solid catalyst;
- (ii) 2-chloro-3,3,3-trifluoropropene (HCFO-1233xf) + HF \rightarrow 2-chloro-1,1,1,2-tetrafluoropropane (HCFC-244bb or 244bb) in a liquid phase reactor charged with a liquid hydrofluorination catalyst; and
- (iii) 2-chloro-1,1,1,2-tetrafluoropropane (HCFC-244bb) \rightarrow 2,3,3,3-tetrafluoropropene (HFO-1234yf) in a vapor phase reactor;

wherein Q is independently selected from F, Cl, Br, and I, provided that at least one Q is not fluorine.

[0005] The hydrofluorination of 1233xf to 244bb is usually conducted in the presence of fluorinated $SbCl_5$ at temperatures above $70^\circ C$; otherwise the catalyst will freeze. Under these conditions, the 1233xf is not completely converted to 244bb because of equilibrium limitations, especially at higher temperatures. As a result, significant amounts of 1233xf are present in the product formed. Since the boiling points of 1233xf and 244bb are only about $2^\circ C$ apart, separation of these two species is difficult and expensive.

[0006] Moreover, the presence of 2-chloro-3,3,3-trifluoropropene (HCFO-1233xf) in the reaction starting materials, such as HCFC-244bb feedstock, can lead to dramatically reduced conversion of HCFC-244bb to HFO-1234yf. In addition, the 2-chloro-3,3,3-trifluoropropene copresence in the starting material, when subjected to dehydrochlorination, can lead to the formation of trifluoropropyne and oligomers, which can produce tar. This result is disadvantageous from the standpoint of a reduced yield of the desired product. Therefore, there is a need for a better catalytic reaction to achieve a higher conversion of 1233xf to 244bb to avoid and/or minimize the need for purification.

[0007] The present invention fulfills that need.

SUMMARY OF THE DISCLOSURE

[0008] The disclosure relates to a method for hydrofluorination of an olefin of the formula: $RCX=CYZ$ to produce hydrofluoroalkanes of formula $RCXFCHYZ$ and $RCHXCFYZ$, wherein X, Y and Z are independently the same or different and are selected from the group consisting of H, F, Cl, Br, and C_1-C_6 alkyl which is partly or fully substituted with chloro or fluoro or bromo, and R is a C_1-C_6 alkyl which is partially or fully substituted with chloro or fluoro or bromo, comprising reacting the fluoroolefin with HF in the liquid-phase, in the presence of SbF_5 , at a temperature ranging from about $-30^\circ C$ to about $65^\circ C$.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0009] The foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention.

[0010] As used herein, the terms “comprises,” “comprising,” “includes,” “including,” “has,” “having” or any other variation thereof, are intended to cover a non-exclusive inclusion. For example, a process, method, article, or apparatus that comprises a list of elements is not necessarily limited to only those elements but may include other elements not expressly listed or inherent to such process, method, article, or apparatus. Further, unless expressly stated to the contrary, “or” refers to an inclusive or and not to an exclusive or. For example, a condition A or B is satisfied by any one of the following: A is true (or present) and B is false (or not present), A is false (or not present) and B is true (or present), and both A and B is true (or present).

[0011] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. In case of conflict, the present specification, including definitions, will control. Although methods and materials similar or equivalent to those described

herein can be used in the practice or testing of embodiments of the present invention, suitable methods and materials are described below. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

[0012] The term “olefin”, as used herein refers to a compound containing a carbon-carbon double bond. It is defined herein relative to the formula $RCX=CYZ$.

[0013] The terms “hydrofluoroalkene” or “fluoroolefin”, as used herein, denotes a compound containing hydrogen, carbon, fluorine, and at least one carbon-carbon double bond and optionally chlorine.

[0014] “HFO”, as used herein, indicates a compound containing hydrogen, carbon, fluorine, and at least one carbon-carbon double, and no chlorine. “HCFO”, as used herein, indicates a compound containing hydrogen, carbon, chlorine, fluorine, and at least one carbon-carbon double. “HCO”, as used herein, indicates a compound containing hydrogen, carbon, chlorine, and at least one carbon-carbon double bond, and no fluorine.

[0015] The term “hydrofluorination” is understood to mean the addition reaction of hydrogen fluoride to a carbon-carbon double bond.

[0016] The term “hydrofluoroalkane”, as used herein, refers to an alkane having two or more carbon atoms containing hydrogen, fluorine, and optionally chlorine, whereby a fluorine atom and a hydrogen atom are substituted on two adjacent carbon atoms. As used herein, the hydrofluoroalkane can be the product from the hydrofluorination of the fluoroolefin.

[0017] The HF used herein is an anhydrous liquid hydrogen fluoride which is commercially available or it may be a gas that is bubbled into the reactor. Anhydrous HF is sold by, for example, Solvay S.A, The Chemours Company FC, LLC and Honeywell International, Inc.

[0018] As used herein, the term “conversion” with respect to a reactant, which typically is a limiting agent, refers to the number of moles reacted in the reaction process divided by the number of moles of that reactant initially present in the process multiplied by 100.

[0019] As used herein, percent conversion is defined as 100%, less the weight percent of starting material in the effluent from the reaction vessel.

[0020] As used herein, the term “selectivity” with respect to an organic reaction product refers to the ratio of the moles of that reaction product to the total of the moles of the organic reaction products multiplied by 100.

[0021] As used herein, “percent selectivity” is defined as the weight of a desired product formed, as a fraction of the total amount of the products formed in the reaction, and excluding the starting material.

[0022] Some fluoroolefins of this disclosure, *e.g.*, $\text{CF}_3\text{CH}=\text{CHCl}$ (HCFO-1233zd or 1233zd), exist as different configurational isomers or stereoisomers. When the specific isomer is not designated, the present disclosure is intended to include all single configurational isomers, single stereoisomers, or any combination thereof. For instance, HCFO-1233zd is meant to represent the *E*-isomer, *Z*-isomer, or any combination or mixture of both isomers in any ratio.

[0023] Described is a method for producing hydrofluoroalkanes of formula RCXFCHYZ , wherein X, Y and Z may independently be the same or different and are selected from H, F, Cl and an alkyl group having 1 to 6 carbon atoms, which alkyl group is partially or fully substituted with fluorine or chlorine; and R is an alkyl group having 1 to 6 carbon atoms, which alkyl group is partially or fully substituted with fluorine or chlorine comprising reacting a fluoroolefin of the formula $\text{RCX}=\text{CYZ}$ with HF in the liquid-phase, in the presence of a catalytic effective amount of SbF_5 .

[0024] The terms “alkyl group is partially or fully substituted with chlorine” and “chlorinated alkyl” are synonymous and it is meant that the alkyl group must be at least monosubstituted with Cl. Similarly, the terms “alkyl group is partially or fully substituted with fluorine” and “fluorinated alkyl” are synonymous and it is meant that the alkyl group must be at least monosubstituted with F. However, in both cases, the alkyl group may have one or more fluoro substituents thereon or one or more chloro substituents thereon or a combination of one or more chloro or fluoro groups thereon. Some of the carbon atoms may be substituted with one or more chloro or fluoro atoms. In an embodiment, the alkyl group is substituted with one or more fluoro atoms. In an embodiment, the alkyl group is fully substituted with chloro or fluoro or combination of both chloro and fluoro. In another embodiment, the alkyl group is perchlorinated, while in another embodiment, the alkyl group is perfluorinated.

[0025] The alkyl group may be branched or linear. In an embodiment, the alkyl group is linear. In an embodiment, the alkyl group contains 1-4 carbon atoms, and in another embodiment, it contains 1 or 2 or 3 carbon atoms and in still another embodiment 1 or 2 carbon atoms. In another embodiment, it contains only 1 carbon atom. Examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl.

[0026] As defined herein, the carbon atoms which are part of the carbon-carbon double bond are substituted with R, X, Y, and Z. R is defined, among other things, as being partially or fully substituted with chloro or fluoro and X, Y, or Z may be, among other things, partially or fully substituted with chloro or fluoro. In one embodiment, X, Y, Z are independently partially or fully substituted with chloro or fluoro, and in another embodiment, two of X, Y, and Z are partially or fully substituted with chloro or fluoro, and in another embodiment, one of X, Y, and Z is partially or fully substituted with chloro or fluoro, in still another embodiment, three of X, Y and Z are partially or fully

substituted with chloro or fluoro, and in another embodiment, none of X, Y, Z are partially or fully substituted with chloro or fluoro. With respect to X, Y, and Z, when defined as partially or fully substituted with chloro or fluoro, and with respect to R, in an embodiment, at least one carbon atom alpha or beta to the carbon atom bearing the double bond (if alkyl group contains 2 or more carbon atoms) is substituted with chloro or fluoro.

[0027] In one embodiment, X, Y, and Z are independently H or fluoro or chloro. In another embodiment, R is perchlorinated or perfluorinated. In some embodiments of this invention, R is $-\text{CF}_3$ or $-\text{CF}_2\text{CF}_3$. In another embodiment, X, Y, and Z are independently H or fluoro or chloro and R is perfluorinated or perchlorinated. In still further embodiment, X, Y, Z are independently H or fluoro or chloro and R is perfluorinated, for example, $-\text{CF}_3$ or $-\text{CF}_2\text{CF}_3$.

[0028] The process according to the invention can be carried out in any reactor made of a material that is resistant to reactants employed, especially to hydrogen fluoride. As used herein, the term "reactor" refers to any vessel in which the reaction may be performed in either a batchwise mode, or in a continuous mode. Suitable reactors include tank reactor vessels with and without agitators, or tubular reactors.

[0029] In one embodiment, the reactor is comprised of materials which are resistant to corrosion including stainless steel, Hastelloy, Inconel, Monel, gold or gold-lined or quartz. In another embodiment, the reactor is TFE or PFA-lined.

[0030] The olefin described herein has the formula $\text{RCX}=\text{CYZ}$, where R, X, Y, Z are as defined hereinabove. Examples include $\text{RCCl}=\text{CH}_2$, $\text{RCH}=\text{CHCl}$, $\text{RCCl}=\text{CHCl}$, $\text{RCH}=\text{CCl}_2$ and $\text{RCH}=\text{CH}_2$, and the like. In one embodiment, R is trifluoromethyl and in another embodiment, R is pentafluoroethyl. Representative olefins include 2-chloro-3,3,3-trifluoropropene (HCFO-1233xf), 1-chloro-3,3,3-trifluoropropene (HCFO-

1233zd), chlorotetrafluoropropenes (HCFO-1224 or 1224), 2,3,3,3-tetrafluoropropene (1234yf), dichlorotetrafluoropropenes (HCFO-1214 or 1214), 1,3,3,3-tetrafluoropropene (1234ze), 3,3,3-trifluoropropene (HFO-1243zf or 1234zf), and the like.

[0031] The hydrofluoroalkanes described herein are the addition products of HF to the fluoroolefins, as defined hereinabove. As defined herein, they have the formula $RCXFCHYZ$ or $RCXHCFYZ$, wherein R, X, Y, Z are as defined hereinabove. As described hereinabove, in one embodiment, R is trifluoromethyl and in another embodiment, R is pentafluoroethyl. Representative hydrofluoropropanes include 1,1,1,2-tetrafluoro-2-chloropropane, 1,1,1,3-tetrafluoro-3-chloropropane, 1,1,1,3,3-pentafluoro-3-chloropropane, 1,1,1,2,2-pentafluoro-3-chloropropane, 1,1,1,2,2-pentafluoropropane, 1,1,1,3,3-pentafluoropropane and the like.

[0032] The present process adds HF across the double bond of the fluoroolefin to produce a hydrofluoroalkane. The F atom may add to an internal or terminal carbon atom and the hydrogen atom may add to a terminal or internal carbon atom. Thus, for example, in accordance with the present disclosure, when the fluoroolefin is $RCCl=CH_2$, the product is $RCFCICH_3$. In another embodiment, when the fluoroolefin is $RCH=CHCl$, the product is RCH_2CHFCl . In another embodiment, when the fluoroolefin is $RCH=CCl_2$, the hydrofluoropropane is RCH_2CFCl_2 . In yet another embodiment, when the fluoroolefin is $RCH=CH_2$, the hydrofluoropropanes formed are $RCHFCH_3$ and RCH_2CH_2F . With respect to the aforementioned examples, in an embodiment, R can be CF_3 or C_2F_6 .

[0033] In one embodiment, the fluoroolefin is 2-chloro-3,3,3-trifluoropropene and the hydrofluoroalkane is 2-chloro-1,1,1,2-tetrafluoropropane. In another embodiment, the fluoroolefin is 3,3,3-trifluoropropene and the hydrofluoroalkane is 1,1,1,2-tetrafluoropropane and 1,1,1,3-tetrafluoropropane. In another embodiment, the

fluoroolefin is (Z)- or (E)- 1-chloro-3,3,3-tetrafluoropropene and the hydrofluoroalkane is 3-chloro-1,1,1,3-tetrafluoropropane. In another embodiment, the fluoroolefin is cis- or trans-1,2-dichloro-3,3,3-trifluoropropene and the hydrofluoroalkane is 1,1,1,2-tetrafluoro-2,3-dichloropropane and 1,1,1,3-tetrafluoro- 2,3-dichloropropane. In another embodiment, the fluoroolefin is 2,3,3,3-tetrafluoropropene, and the hydrofluoroalkane is 1,1,1,2,2-pentafluoropropane. In yet another embodiment, the fluoroolefin is 1,3,3,3-tetrafluoropropene and the hydrofluoroalkane is 1,1,1,3,3-pentafluoropropane.

[0034] Without wishing to be bound, it is believed that with respect to HF addition to a carbon-carbon double bond, the fluorine atom adds to the carbon atom of the double bond which has the most halogens attached thereto. Otherwise, without wishing to be bound, the HF is added to the carbon atom of the double bond in accordance with Markovnikov's rule, i.e., the hydrogen of HF will add to the carbon atom that will form the more stable carbonium ion. Thus, for example, if one of the carbon atoms of the carbon- carbon double bond has more hydrogen atoms substituted thereon than the other carbon atom of the carbon-carbon double bond, the hydrogen atom of HF will add to the carbon atom having the most hydrogen atoms substituted thereon.

[0035] The above process is conducted in the liquid phase. The fluoroolefin as well as the hydrogen fluoride are liquids at reaction conditions. Since water is used to quench the reaction, the amount of water present is minimized. For example, the hydrogen fluoride used is anhydrous. The hydrogen fluoride can be bubbled in as a gas or added as a liquid into the liquid fluoroolefin or it may be present in an anhydrous solvent, such as pyridine. Thus, for example, in an embodiment, although not necessary, the fluoroolefin is dried with a desiccant before being mixed with HF or the catalyst. By "desiccant," it is meant any material which will absorb water without dissolving in or otherwise contaminating the fluoroolefin being dried, *e.g.*, calcium sulfate or molecular sieves, and

the like. In another embodiment, the reaction can be conducted in an inert atmosphere, such as under nitrogen, helium, argon and the like. However, in an embodiment, the reaction can be conducted in air and in another embodiment, the reaction is conducted without drying the fluoroolefin.

[0036] In an embodiment, when anhydrous liquid HF is used or HF is fed as a gas, the reaction is conducted without any solvent in addition to the solvent in which the anhydrous HF is dissolved. If the HF is fed as a gas, such as, being bubbled in as a gas, the reaction may be conducted without any solvent present.

[0037] In one embodiment, the hydrofluorination reaction is conducted at a temperature ranging from about -30°C to about 65°C. In another embodiment, the hydrofluorination reaction is conducted at a temperature ranging from about -10°C to about 40°C. In another embodiment, the hydrofluorination reaction is conducted at a temperature ranging from about 0°C to about 30°C. In still another embodiment, the hydrofluorination reaction is conducted at a temperature ranging from about 0°C to about 25°C. In another embodiment, the hydrofluorination reaction is conducted at a temperature ranging from about 5°C to about 25°C. In still another embodiment, the hydrofluorination reaction is conducted at a temperature ranging from about 5°C to about 20°C. Moreover, the hydrofluorination reaction can be conducted at any temperature in-between the ranges disclosed hereinabove, and these temperatures are contemplated within the scope of the present invention. Thus, the hydrofluorination described hereinabove is conducted in a reaction vessel at about -30°C, about -29° C, about -28° C, about 27° C, about -26° C, about -25° C, about -24° C, about -23° C, about -22° C, about -21° C, about -20° C, about -19° C, about -18° C, about -17°C, about -16°C, about -15°C, about -14°C, about -13°C, about -12°C, about -11°C, about -10°C, about -9°C, about -8°C, about -7°C, about -6°C, about -5°C, about -4°C, about -3°C, about

-2°C, about -1°C, about 0°C, about 1°C, about 2°C, about 3°C, about 4°C, about 5°C, about 6°C, about 7°C, about 8°C, about 9°C, about 10°C, about 11°C, about 12°C, about 13°C, about 14°C, about 15°C, about 16°C, about 17°C, about 18°C, about 19°C, about 20°C, about 21°C, about 22°C, about 23°C, about 24°C, about 25°C, about 26°C, about 27°C, about 28°C, about 29°C, about 30°C, about 31°C, about 32°C, about 33°C, about 34°C, about 35°C, about 36°C, about 37°C, about 38°C, about 39°C, about 40°C, about 41°C, about 42°C, about 43°C, about 44°C, about 45°C, about 46°C, about 47°C, about 48°C, about 49°C, about 50°C, about 51°C, about 52°C, about 53°C, about 54°C, about 55°C, about 56°C, about 57°C, about 58°C, about 59°C, about 60°C, about 61°C, about 62°C, about 63°C, about 64°C or about 65°C.

[0038] In an embodiment, the reaction mixture is stirred using techniques known in the art. For example, the reaction mixture is spun using a stirring bar. Alternatively, the reactor in which the reaction takes place is equipped with an impeller or other stirring device which stirs the reaction mixture.

[0039] In another embodiment, mixing may be provided by alternatives to stirring devices. Such methods are known in the industry and include using the mixing provided by gas bubbles from gas added to the vessel or generated within the vessel by vaporization of liquid. Mixing can also be provided by withdrawing the liquid from the vessel to a pump and pumping the liquid back into the vessel. A static mixer or other device intended to mix the contents can be present in the circulation path of the liquid to provide additional mixing power input.

[0040] In one embodiment, the mole ratio of HF to fluoroolefin ranges from about 0.5 to about 20. In another embodiment, the mole ratio of HF to fluoroolefin is from about 1 to about 10. In another embodiment, the mole ratio of HF to fluoroolefin is from about 1 to about 5.

[0041] The SbF_5 is present in catalytic effective amounts. In one embodiment, the SbF_5 catalyst is present from about 1% to about 50% by weight of the mixture. In another embodiment, the SbF_5 catalyst is present from about 2% to about 30% by weight. In another embodiment, the SbF_5 catalyst is present from about 3% to about 15% by weight.

[0042] As described hereinabove, hydrofluoroalkanes are prepared by catalytic fluorination of the fluoroolefin. In one embodiment, the catalytic fluorination of the fluoroolefin results in a percent conversion to the hydrofluoroalkane of at least 90 mole %. In another embodiment, the catalytic fluorination of the fluoroolefin results in a percent conversion to the hydrofluoroalkane of at least 95%. In another embodiment, the catalytic fluorination of the fluoroolefin results in a percent conversion to the hydrofluoroalkane of at least 98%. In still another embodiment, the catalytic fluorination of the fluoroolefin results in a percent conversion to the hydrofluoroalkane of at least 99%.

[0043] An aspect of the invention is to replace step (ii) of the reaction for making 1234yf described in the introduction with the present process.

[0044] One of the advantages of the present disclosure is that the catalytic reaction for hydrofluorination, as described herein, takes place at lower temperatures, much lower than other catalysts for the other hydrofluorination reactions of fluoroolefin, such as SbCl_5 or fluorinated SbCl_5 . Unlike these other catalysts, SbF_5 is a liquid at these lower temperatures that are used in the present process. Therefore, less energy is required to conduct these hydrofluorination reactions. In addition, in the present process, the catalyst has substantial activity at the lower temperature. Thus, the catalytic process proceeds at a low temperature, thereby making it more efficient.

[0045] In addition, another advantage is that the ratio of the desired hydrofluoroalkane produced relative to the starting olefin is about 90:1 or greater, and in another embodiment, is about 100:1 or greater and in another embodiment is about 110:1 or greater. Thus, for another reason, this reaction is quite efficient.

[0046] Moreover, in view of the efficiency, if an olefin and the resulting hydrofluoroalkane from the hydrofluorination reaction, such as 1233xf and 244bb, were mixed together and reacted under the conditions of the present invention with SbF_5 , additional hydrofluoroalkane product would be formed. For example, in one embodiment, if the feed material ratio of olefin, such as 1233xf, to hydrofluoroalkane, such as 244bb, is greater than about 1 mole %, the present process will significantly convert the unreacted olefin to hydrofluoroalkane, thereby increasing the amount of the hydrofluoroalkane in the mixture. The present disclosure thus provides a method of maximizing the yield of the desired hydrofluoroalkane relative to the olefin. Thus, in the above example, wherein the olefin is 1233xf and the hydrofluoroalkane is 244bb, if 1233xf is present in greater than about 1 mole %, the resulting product would have significantly more 244bb present than prior to the reaction.

[0047] Thus, in one embodiment, this advantage of the present disclosure can be used to improve the yield of HFO-1234yf being produced. As described hereinabove, the preparation of HFO-1234yf may include at least three reaction steps, as follows:

- (i) $(\text{CQ}_2=\text{CCl}-\text{CH}_2\text{Q} \text{ or } \text{CQ}_3-\text{CCl}=\text{CH}_2 \text{ or } \text{CQ}_3-\text{CHCl}-\text{CH}_2\text{Q}) + \text{HF} \rightarrow$ 2-chloro-3,3,3-trifluoropropene (HCFO-1233xf) + HCl in a vapor phase reactor charged with a solid catalyst;
- (ii) 2-chloro-3,3,3-trifluoropropene (HCFO-1233xf) + HF \rightarrow 2-chloro-1,1,1,2-tetrafluoropropane (HCFC-244bb) in a liquid phase reactor charged with a liquid hydrofluorination catalyst; and

(iii) 2-chloro-1,1,1,2-tetrafluoropropane (HCFC-244bb) → 2,3,3,3-tetrafluoropropene (HFO-1234yf) in a vapor phase reactor.

[0048] The general reactions of steps (i), (ii) and (iii) are well known in the art. For example, they are described in U.S Patent No. 8,846,990, the contents of which are incorporated by reference.

[0049] In the first step, a starting composition, which comprises 1,1,2,3-tetrachloropropene (HCO-1230xa or 1230xa), reacts with anhydrous HF in a first reactor (fluorination reactor) to produce a mixture of at least HCFO-1233xf (2-chloro-3,3,3-trifluoropropene) and HCl. The reaction is carried out in a reactor in the gaseous phase at a temperature of about 200°C to about 400°C and a pressure of about 0 to about 200 psig. The effluent stream exiting in the vapor phase reactor may optionally comprise additional components, such as un-reacted HF, un-reacted starting composition, heavy intermediates, HFC-245cb, or the like.

[0050] This reaction may be conducted in any reactor suitable for a vapor phase fluorination reaction. The reactor may be constructed from materials which are resistant to the corrosive effects of hydrogen fluoride such as Hastalloy, Inconel, Monel, and the like. In the case of a vapor phase process, the reactor is filled with a vapor phase fluorination catalyst. Any fluorination catalysts known in the art may be used in this process. Suitable catalysts include, but are not limited to, metal oxides, hydroxides, halides, oxyhalides, inorganic salts thereof and their mixtures, any of which may be optionally halogenated, wherein the metal includes, but is not limited to, chromium, aluminum, cobalt, manganese, nickel, iron, and combinations of two or more thereof. Combinations of catalysts suitable for the present invention nonexclusively include Cr₂O₃, FeCl₃/C, Cr₂O₃/Al₂O₃, Cr₂O₃/AlF₃, Cr₂O₃/carbon, CoCl₂/Cr₂O₃/Al₂O₃, NiCl₂/Cr₂O₃/Al₂O₃, CoCl₂/AlF₃, NiCl₂/AlF₃ and mixtures thereof. Chromium

oxide/aluminum oxide catalysts are described in U.S. Pat. No. 5,155,082, the contents of which are incorporated herein by reference. Chromium (III) oxides such as crystalline chromium oxide or amorphous chromium oxide are preferred with amorphous chromium oxide being most preferred. Chromium oxide (Cr_2O_3) is a commercially available material which may be purchased in a variety of particle sizes. Fluorination catalysts having a purity of at least 98% are preferred. The fluorination catalyst is present in an excess but in at least an amount sufficient to drive the reaction.

[0051] This first step of the reaction is not necessarily limited to a vapor phase reaction and may also be performed using a liquid phase reaction or a combination of liquid and vapor phases, such as that disclosed in U.S. Published Patent Application No. 2007/0197842, the contents of which are incorporated herein by reference. It is also contemplated that the reaction can be carried out batch wise or in a continuous manner, or a combination of these.

[0052] For embodiments in which the reaction comprises a liquid phase reaction, the reaction can be catalytic or non-catalytic. Lewis acid catalysts, such as metal-halide catalysts, including antimony halides, tin halides, thallium halides, iron halides, and combinations of two or more of these, may be employed. In certain embodiments, metal chlorides and metal fluorides are employed, including, but not limited to, SbCl_5 , SbCl_3 , SbF_5 , SnCl_4 , TiCl_4 , FeCl_3 and combinations of two or more of these. It is noted that SbF_5 is a liquid at low temperature.

[0053] In the second step of the process for forming 2,3,3,3-tetrafluoropropene, HCFO-1233xf is converted to HCFC-244bb. In one embodiment, this step can be performed in the liquid phase in a liquid phase reactor, which may be TFE or PFA-lined. Such a process can be performed in a temperature range of about 70°C to about 120°C and at a pressure ranging from about 50 to about 120 psig. Any liquid phase fluorination catalyst

may be used that is effective at these temperatures. A non-exhaustive list includes Lewis acids, transition metal halides, transition metal oxides, Group IVb metal halides, Group Vb metal halides, or combinations thereof. Non-exclusive examples of liquid phase fluorination catalysts are antimony halide, tin halide, tantalum halide, titanium halide, niobium halide, molybdenum halide, iron halide, fluorinated chrome halide, fluorinated chrome oxide or combinations thereof. Specific non-exclusive examples of liquid phase fluorination catalysts are SbCl_5 , SbCl_3 , SbF_5 , SnCl_4 , TaCl_5 , TiCl_4 , NbCl_5 , MoCl_6 , FeCl_3 , fluorinated species of SbCl_5 , fluorinated species of SbCl_3 , fluorinated species of SnCl_4 , fluorinated species of TaCl_5 , fluorinated species of TiCl_4 , fluorinated species of NbCl_5 , fluorinated species of MoCl_6 , fluorinated species of FeCl_3 , or combinations thereof.

[0054] These catalysts can be readily regenerated by any means known in the art if they become deactivated. One suitable method of regenerating the catalyst involves flowing a stream of chlorine through the catalyst. For example, from about 0.002 to about 0.2 lb per hour of chlorine can be added to the liquid phase reaction for every pound of liquid phase fluorination catalyst. This may be done, for example, for from about 1 to about 2 hours or continuously at a temperature of from about 65°C to about 100°C.

[0055] This second step of the reaction is not necessarily limited to a liquid phase reaction and may also be performed using a vapor phase reaction or a combination of liquid and vapor phases, such as that disclosed in U.S. Published Patent Application No. 2007/0197842, the contents of which are incorporated herein by reference. To this end, the HCFO-1233xf containing feed stream is preheated to a temperature of from about 50°C to about 400°C, and is contacted with a catalyst and fluorinating agent. Catalysts may include standard vapor phase agents used for such a reaction and fluorinating agents may include those generally known in the art, such as, but not limited to, hydrogen fluoride.

[0056] In the process described in the art, such as that described in U.S. Published Patent Application No. 2007/0197842, the product from the second step is then transferred to a third reactor wherein the 244bb is dehydrohalogenated. The catalysts in the dehydrochlorination reaction may be or comprise metal halide, halogenated metal oxide, neutral (or zero oxidation state) metal or metal alloy, or activated carbon in bulk or supported form. Metal halide or metal oxide catalysts may include, but are not limited to, mono-, bi-, and tri-valent metal halides, oxides and their mixtures/combinations, and more preferably mono-, and bi-valent metal halides and their mixtures/combinations. Component metals of metal halides, oxides and their mixtures/combinations include, but are not limited to, Cr^{3+} , Fe^{3+} , Mg^{2+} , Ca^{2+} , Ni^{2+} , Zn^{2+} , Pd^{2+} , Li^+ , Na^+ , K^+ , and Cs^+ . Component halides include, but are not limited to, F, Cl, Br^- , and I. Examples of useful mono- or bi-valent metal halide include, but are not limited to, LiF, NaF, KF, CsF, MgF_2 , CaF_2 , LiCl, NaCl, KCl, and CsCl. Halogenation treatments can include any of those known in the prior art, particularly those that employ HF, F_2 , HCl, Cl_2 , HBr, Br_2 , HI, and I_2 as the halogenation source.

[0057] When the catalyst is or comprises a neutral, *i.e.*, zero valent metal, then metals and metal alloys and their mixtures are used. Useful metals include, but are not limited to, Pd, Pt, Rh, Fe, Co, Ni, Cu, Mo, Cr, Mn, and combinations of the foregoing as alloys or mixtures. The catalyst may be supported or unsupported. Useful examples of metal alloys include, but are not limited to, SS 316, Monel 400, Inconel 825, Inconel 600, and Inconel 625. Such catalysts may be provided as discrete supported or unsupported elements and/or as part of the reactor and/or the reactor walls.

[0058] Preferred, but non-limiting, catalysts include activated carbon, stainless steel (*e.g.*, SS 316), austenitic nickel-based alloys (*e.g.*, Inconel 625), nickel, fluorinated 10% CsCl/MgO, and 10% CsCl/ MgF_2 . A suitable reaction temperature is about 300°C to

about 550°C and a suitable reaction pressure may be between about 0 psig to about 150 psig. The reactor effluent may be fed to a caustic scrubber or to a distillation column to remove the byproduct of HCl to produce an acid-free organic product which, optionally, may undergo further purification using one or any combination of purification techniques that are known in the art.

[0059] The dehydrohalogenation reaction is carried out in the vapor phase. It may be carried out at a temperature range of from about 200°C to about 800°C, from about 300°C to about 600°C, or from about 400°C to about 500°C. Suitable reactor pressures range from about 0 psig to about 200 psig, from about 10 psig to about 100 psig, or from about 20 to about 70 psig.

[0060] A method of increasing the yield and conversion of 1233xf to 1234yf and to make the process more efficient is to react the product of step (ii), which contains a mixture of 1233xf and 244bb, with SbF₅ in accordance with the process of the present invention prior to the dehydrochlorination step. This increases the amount of 244bb present (decreasing the amount of 1233xf present) and the resulting product can then be subjected to step (iii) above. By conducting this additional hydrofluorination reaction, more 244bb is produced, and as a result, significantly more 1234yf is produced. The 244bb thus produced is then transferred to another reactor wherein it undergoes dehydrohalogenation, in accordance with step (iii).

[0061] Alternatively, as described above, instead of conducting step (ii) of the process, the 1233xf produced in step (i) is hydrofluorinated with HF in the presence of SbF₅, in accordance with the present invention, as described herein. The 244bb product thus formed is then dehydrochlorinated to form 1234yf, in accordance with step (iii) described hereinabove.

[0062] The following non-limiting examples further illustrate the invention.

EXAMPLESExample 1 – 1233xf hydrofluorination by HF with SbF₅ catalyst at 30°C

[0063] 13.8 g of HF and 5 g of SbF₅ were loaded into a 210 mL shaker tube reactor. The reactor was then evacuated and chilled to -15°C. 30 g of 1233xf was added into the reactor. The reactor was then head to 30°C with agitation. Once the temperature reached 30°C, water was added to the reactor to quench the catalyst. The organic layer was vapor transferred into a stainless steel cylinder and analyzed by GC-MS. Table 1 below shows the results of the GC-MS analysis.

TABLE 1

| | mol% | mol ratio 1233xf/244bb |
|---------|---------|---------------------------|
| 143a | 0.005% | |
| 245cb | 0.032% | |
| 245fa | 0.059% | |
| unknown | 0.001% | |
| 244bb | 99.420% | 0.48% |
| 1233xf | 0.474% | |
| 243ab | 0.009% | |

Example 2 – 1233xf hydrofluorination by HF with SbF₅ catalyst at 10°C

[0064] 13.8 g of HF and 5 g of SbF₅ were loaded into a 210 mL shaker tube reactor. The reactor was then evacuated and chilled to -15°C. 30 g of 1233xf was added into the reactor. The reactor was then head to 10° C with agitation. Once the temperature reached 10°C, water was added to the reactor to quench the catalyst. The organic layer was vapor transferred into a stainless steel cylinder and analyzed by GC-MS. Table 2 below shows the results of the GC-MS analysis.

TABLE 2

| | | mol ratio |
|--------------|---------|--------------|
| Compounds | mol% | 1233xf/244bb |
| 245cb | 0.017% | |
| 245fa | 0.0200% | |
| 244bb | 98.236% | |
| 1233xf | 0.794% | 0.81% |
| 1233xf dimer | 0.934% | |

Example 3 – 1233xf hydrofluorination by HF with SbF₅ catalyst at 30°C

[0065] 10.0 g of HF and 5 g of SbF₅ were loaded into a 210 mL shaker tube reactor. The reactor was then evacuated and chilled to -40°C. 30 g of 1233xf was added into the reactor. The reactor was then head to 30°C with agitation and stirred for an hour. The reactor was chilled to -30°C quickly and 75 mL of water was added to the reactor to quench the catalyst. The organic layer was vapor transferred into a stainless steel cylinder and analyzed by GC-MS. Table 3 below shows the results of the GC-MS analysis.

TABLE 3

| | | mol ratio |
|----------|---------|--------------|
| Compound | mol% | 1233xf/244bb |
| 245cb | 14.412% | |
| 245fa | 0.096% | |
| 244bb | 84.147% | |
| 1233xf | 0.768% | 0.91% |
| 243ab | 0.576% | |

Comparative Example 1 – 1233xf-244bb equilibrium by HF with fluorination SbCl_5 catalyst at 80°C

[0066] 18.0 g of HF and 14.0 g of SbCl_5 were loaded into a 210 mL shaker tube reactor and heated at 100C for 2 hours with agitation. The reactor was then evacuated and chilled to 0° C to vent off HCl. 20 g of 244bb (99.7 mol %) was added into the reactor. The reactor was then heated to 80°C for an hour and then quickly chilled to 30°C. Water was added to the reactor to quench the catalyst. The organic layer was vapor transferred into a stainless steel cylinder and analyzed by GC-MS. Table 4 below shows the results of the GC-MS analysis. The 1233xf/244bb ratio increased to 1.95 mol% from 0.3 mol%. This indicates the existence of equilibrium between 1233xf and 244bb which prevents the full conversion of 1233xf to 244bb.

TABLE 4

| | mol% | mol ratio 1233xf/244bb |
|--------|---------|---------------------------|
| 245cb | 0.089% | |
| 244bb | 92.907% | |
| 1233xf | 1.815% | 1.95% |
| 243ab | 4.905% | |
| others | 0.284% | |

[0067] Many aspects and embodiments have been described and are merely exemplary and not limiting. After reading the specification, skilled artisans appreciate that other aspects and embodiments are possible without departing from the scope of the invention.

[0068] Other features and benefits of any one or more of the embodiments will be apparent from the hereinabove detailed description and the claims.

WHAT IS CLAIMED IS:

1. A method for preparing a hydrofluoroalkane of formula RCXFCHYZ comprising reacting HF with olefin of the formula RCX=CYZ in the liquid phase in the presence of SbF₅ at a temperature ranging from about -30° C to about 65° C to produce the hydrofluoroalkane, wherein X, Y and Z are independently the same or different and are H, F, Cl, Br, or C₁-C₆ alkyl which is partially or fully substituted with chloro or fluoro or bromo, and R is a C₁-C₆ alkyl which is partially or fully substituted with chloro or fluoro, or bromo.
2. The method according to claim 1 wherein X, Y and Z are independently H, F, Cl or Br.
3. The method according to claim 1 or 2 where R is a fluorinated alkyl having 1-6 carbon atoms or chlorinated alkyl having 1-6 carbon atoms.
4. The method according to claim 3 where R is fluorinated alkyl.
5. The method according to claim 3 where R is perfluorinated alkyl.
6. The method according to claim 4 wherein R is CF₃ or C₂F₅.
7. The method of claim 1, wherein the ratio of HF to the olefin is in the range of from 1 to 10.
8. The method of claim 7, wherein the ratio of HF to the olefin is in the range of from 1 to 5.
9. The method of claim 1, wherein that catalyst is present in an amount of 1% to 50% by weight.

10. The method of claim 9, wherein that catalyst is present in an amount of 2% to 30% by weight.
11. The method of claim 10, wherein that catalyst is present in an amount of 3% to 20% by weight.
12. The method of claim 1, wherein the olefin is 2-chloro-3,3,3-trifluoropropene, 1-chloro-3,3,3-trifluoropropene, chlorotetrafluoropropenes, 2,3,3,3-tetrafluoropropene, 1,3,3,3-tetrafluoropropene, or 3,3,3-trifluoropropene.
13. The method of claim 1, wherein the hydrofluoroalkane is 2-chloro-1,1,1,2-tetrafluoropropane.
14. The method of claim 1 wherein the olefin is 2-chloro-3,3,3-trifluoropropene and the hydrofluoroalkane is 2-chloro-1,1,1,2-tetrafluoropropane.
15. The method of claim 1 wherein the olefin is (Z)- or (E)-1-chloro-3,3,3-trifluoropropene, and the hydrofluoroalkane is 3-chloro-1,1,1,3-tetrafluoropropane.
16. The method of claim 1 wherein the olefin is 1-chloro-2,3,3,3-trifluoropropene and the hydrofluoroalkane is 1,1,1,2,2-pentafluoro-3-chloropropane and 1,1,1,2,3-pentafluoro-3-chloropropane.
17. The method of claim 1 where the olefin is 2,3,3,3-tetrafluoropropene and the hydrofluoroalkane is 1,1,1,2,2-pentafluoropropane.
18. The method of claim 1 wherein the olefin is 1,3,3,3-tetrafluoropropene and the hydrofluoroalkane is 1,1,1,3,3-pentafluoropropane.
19. The method of claim 1, wherein the temperature is from about -10°C to about 40°C.

20. The method of claim 19, wherein the temperature is from about 0°C to about 30°C.
21. The method of claim 20, wherein the temperature is from about 5°C to about 25°C.
22. The method of claim 1, wherein the feed material ratio of olefin to hydrofluoroalkane is greater than 1 mole percent.
23. A method of preparing 2-chloro-1,1,1,2-tetrafluoropropane which comprises reacting 2-chloro-1,1,1-trifluoropropene with HF in the liquid phase in the presence of SbF₅ at a temperature ranging from about -30°C to about 65°C.
24. The method of claim 16, wherein the 2-chloro-1,1,1,2-tetrafluoropropane is dehydrochlorinated to form 2,3,3,3-tetrafluoro-1-propene.
25. A method for preparing 2,3,3,3-tetrafluoro-1-propene comprising reacting 2-chloro-1,1,1-trifluoropropene with HF in a liquid phase reactor charged with a hydrofluorination catalyst at a reaction temperature above 65°C to produce a mixture of 2-chloro-1,1,1-trifluoropropene and 2-chloro-1,1,1,2-tetrafluoropropane; reacting HF with said mixture in the liquid phase in the presence of SbF₅ at a temperature ranging from about -30°C to about 65°C to further react the unconverted 2-chloro-1,1,1-trifluoropropene to form additional 2-chloro-1,1,1,2-tetrafluoropropane and dehydrohalogenating 2-chloro-1,1,1,2-tetrafluoropropane in a vapor phase reactor with or without a catalyst to form 2,3,3,3-tetrafluoro-1-propene.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2016/033450**A. CLASSIFICATION OF SUBJECT MATTER****C07C 17/08(2006.01)I, C07C 17/42(2006.01)I, C07C 19/10(2006.01)I, C07C 21/19(2006.01)I**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07C 17/08; C07C 21/18; C07C 17/25; C07C 17/087; C07C 19/10; C07C 17/23; C07C 17/42; C07C 21/19

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models

Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords: 2-chloro-1,1,1,2-tetrafluoropropane, 244bb, 2-chloro-3,3,3-trifluoropropene, 1233xf, HF, hydrofluorination, SbF5, liquid phase

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| A | US 2009-0182179 A1 (MERKEL DANIEL C. et al.) 16 July 2009 See abstract, paragraphs [0014]-[0017] and example 1. | 1-25 |
| A | US 2010-0036179 A1 (MERKEL DANIEL C. et al.) 11 February 2010 See abstract, paragraphs [0020]-[0022] and example 1. | 1-25 |
| A | US 2010-0331583 A1 (JOHNSON ROBERT C. et al.) 30 December 2010 See abstract, paragraphs [0030]-[0032], [0057] and example 1. | 1-25 |
| A | US 2014-0275648 A1 (HONEYWELL INTERNATIONAL, INC.) 18 September 2014 See abstract and paragraphs [0013]-[0017]. | 1-25 |
| A | US 2009-0287026 A1 (KOPKALLI HALUK et al.) 19 November 2009 See abstract and paragraphs [0029]-[0038]. | 1-25 |

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

22 August 2016 (22.08.2016)

Date of mailing of the international search report

22 August 2016 (22.08.2016)

Name and mailing address of the ISA/KR

International Application Division

Korean Intellectual Property Office


189 Cheongsa-ro, Seo-gu, Daejeon, 35208, Republic of Korea

Facsimile No. +82-42-481-8578

Authorized officer

LEE, Sun Hwa

Telephone No. +82-42-481-5603



INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2016/033450

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|----------------------------------------|------------------|-------------------------|------------------|
| US 2009-0182179 A1 | 16/07/2009 | AU 2003-284352 A1 | 13/05/2004 |
| | | AU 2003-284352 A8 | 13/05/2004 |
| | | AU 2003-286685 A1 | 13/05/2004 |
| | | AU 2003-286685 C1 | 23/05/2013 |
| | | AU 2005-238537 A1 | 10/11/2005 |
| | | AU 2006-261816 A1 | 04/01/2007 |
| | | AU 2006-261816 B2 | 19/05/2011 |
| | | AU 2010-245671 A1 | 11/11/2010 |
| | | AU 2010-245671 B2 | 29/01/2015 |
| | | AU 2010-246160 A1 | 11/11/2010 |
| | | AU 2011-248123 A1 | 10/11/2011 |
| | | AU 2011-248123 B2 | 22/01/2015 |
| | | CA 2503421 A1 | 06/05/2004 |
| | | CA 2503421 C | 06/01/2015 |
| | | CA 2557873 A1 | 10/11/2005 |
| | | CA 2557873 C | 25/11/2014 |
| | | CA 2564897 A1 | 17/11/2005 |
| | | CA 2564897 C | 27/11/2012 |
| | | CA 2564903 A1 | 17/11/2005 |
| | | CA 2564903 C | 04/12/2012 |
| | | CA 2564991 A1 | 17/11/2005 |
| | | CA 2564991 C | 19/03/2013 |
| | | CA 2608327 A1 | 27/04/2008 |
| | | CA 2608327 C | 14/04/2015 |
| | | CA 2608675 A1 | 27/04/2008 |
| | | CA 2608675 C | 31/03/2015 |
| | | CA 2612986 A1 | 04/01/2007 |
| | | CA 2613090 A1 | 04/01/2007 |
| | | CA 2628463 A1 | 18/05/2007 |
| | | CA 2628463 C | 08/07/2014 |
| | | CA 2635917 A1 | 12/07/2007 |
| | | CA 2646990 A1 | 27/09/2007 |
| | | CA 2674256 A1 | 30/01/2010 |
| | | CA 2706774 A1 | 28/05/2009 |
| | | CA 2711861 A1 | 16/07/2009 |
| | | CA 2745762 A1 | 10/06/2010 |
| | | CA 2761418 A1 | 11/11/2010 |
| | | CA 2761478 A1 | 11/11/2010 |
| | | CA 2776375 A1 | 07/04/2011 |
| | | CA 2798620 A1 | 10/11/2011 |
| | | CA 2822739 A1 | 04/01/2007 |
| | | CA 2826532 A1 | 10/11/2005 |
| | | CA 2826545 A1 | 06/05/2004 |
| | | CN 101014680 A | 08/08/2007 |
| | | CN 101014680 B | 03/09/2014 |
| | | CN 101014680 C | 08/08/2007 |
| | | CN 101177378 A | 14/05/2008 |
| | | CN 101177378 B | 09/04/2014 |
| | | CN 101177378 C | 14/05/2008 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2016/033450

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|-------------------------------------------|---------------------|----------------------------|---------------------|
| | | CN 101182280 A | 21/05/2008 |
| | | CN 101182280 B | 27/06/2012 |
| | | CN 101182280 C | 21/05/2008 |
| | | CN 101248153 A | 20/08/2008 |
| | | CN 101248153 B | 24/07/2013 |
| | | CN 101248153 C | 20/08/2008 |
| | | CN 101260021 A | 10/09/2008 |
| | | CN 101260021 B | 06/08/2014 |
| | | CN 101260021 C | 10/09/2008 |
| | | CN 101283071 A | 08/10/2008 |
| | | CN 101283071 B | 24/07/2013 |
| | | CN 101351426 A | 21/01/2009 |
| | | CN 101351426 B | 08/05/2013 |
| | | CN 101351427 A | 21/01/2009 |
| | | CN 101351427 B | 21/08/2013 |
| | | CN 101351430 A | 21/01/2009 |
| | | CN 101395108 A | 25/03/2009 |
| | | CN 101395108 B | 13/03/2013 |
| | | CN 101448913 A | 03/06/2009 |
| | | CN 101492342 A | 29/07/2009 |
| | | CN 101495431 A | 29/07/2009 |
| | | CN 101553453 A | 07/10/2009 |
| | | CN 101553453 B | 29/05/2013 |
| | | CN 101597209 A | 09/12/2009 |
| | | CN 101665405 A | 10/03/2010 |
| | | CN 101665681 A | 10/03/2010 |
| | | CN 101665681 B | 25/06/2014 |
| | | CN 101796155 A | 04/08/2010 |
| | | CN 101925644 A | 22/12/2010 |
| | | CN 102015050 A | 13/04/2011 |
| | | CN 102015050 B | 02/07/2014 |
| | | CN 102026944 A | 20/04/2011 |
| | | CN 102112421 A | 29/06/2011 |
| | | CN 102112421 B | 26/11/2014 |
| | | CN 102140329 A | 03/08/2011 |
| | | CN 102264860 A | 30/11/2011 |
| | | CN 102264860 B | 17/06/2015 |
| | | CN 102281932 A | 14/12/2011 |
| | | CN 102281932 B | 06/01/2016 |
| | | CN 102307965 A | 04/01/2012 |
| | | CN 102307965 B | 06/01/2016 |
| | | CN 102439108 A | 02/05/2012 |
| | | CN 102596869 A | 18/07/2012 |
| | | CN 102782077 A | 14/11/2012 |
| | | CN 102803427 A | 28/11/2012 |
| | | CN 102971394 A | 13/03/2013 |
| | | CN 102971394 B | 20/04/2016 |
| | | CN 103146348 A | 12/06/2013 |
| | | CN 103146348 B | 13/01/2016 |
| | | CN 103172488 A | 26/06/2013 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2016/033450

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|----------------------------------------|------------------|-------------------------|------------------|
| | | CN 103172488 B | 21/10/2015 |
| | | CN 103215013 A | 24/07/2013 |
| | | CN 103274895 A | 04/09/2013 |
| | | CN 103467241 A | 25/12/2013 |
| | | CN 103497351 A | 08/01/2014 |
| | | CN 103497738 A | 08/01/2014 |
| | | CN 103627019 A | 12/03/2014 |
| | | CN 103642461 A | 19/03/2014 |
| | | CN 103923610 A | 16/07/2014 |
| | | CN 103923611 A | 16/07/2014 |
| | | CN 103923723 A | 16/07/2014 |
| | | CN 104045848 A | 17/09/2014 |
| | | CN 104045849 A | 17/09/2014 |
| | | CN 104045850 A | 17/09/2014 |
| | | CN 104046332 A | 17/09/2014 |
| | | CN 104059613 A | 24/09/2014 |
| | | CN 104164215 A | 26/11/2014 |
| | | CN 104164216 A | 26/11/2014 |
| | | CN 104164217 A | 26/11/2014 |
| | | CN 104178089 A | 03/12/2014 |
| | | CN 104178090 A | 03/12/2014 |
| | | CN 104194726 A | 10/12/2014 |
| | | CN 104194727 A | 10/12/2014 |
| | | CN 104194728 A | 10/12/2014 |
| | | CN 104194729 A | 10/12/2014 |
| | | CN 1732243 A | 08/02/2006 |
| | | CN 1732243 C | 08/02/2006 |
| | | CN 1852880 A | 25/10/2006 |
| | | CN 1852880 B | 15/06/2011 |
| | | CN 1852880 C | 25/10/2006 |
| | | CN 1898353 A | 17/01/2007 |
| | | CN 1898353 B | 07/05/2014 |
| | | CN 1898353 C | 17/01/2007 |
| | | CN 1902152 A | 24/01/2007 |
| | | CN 1902152 B | 13/10/2010 |
| | | CN 1902152 C | 24/01/2007 |
| | | CN 1968915 A | 23/05/2007 |
| | | CN 1968915 B | 03/08/2011 |
| | | CN 1968915 C | 23/05/2007 |
| | | CN 1972887 A | 30/05/2007 |
| | | CN 1972887 B | 13/10/2010 |
| | | CN 1972887 C | 30/05/2007 |
| | | CN 1976885 A | 06/06/2007 |
| | | CN 1976885 B | 26/05/2010 |
| | | CN 1976885 C | 06/06/2007 |
| | | EP 1563032 A2 | 17/08/2005 |
| | | EP 1563032 B1 | 02/05/2012 |
| | | EP 1578883 A2 | 28/09/2005 |
| | | EP 1658252 A2 | 24/05/2006 |
| | | EP 1658252 B1 | 04/01/2012 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2016/033450

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|-------------------------------------------|---------------------|----------------------------|---------------------|
| | | EP 1678106 A2 | 12/07/2006 |
| | | EP 1678106 B1 | 04/01/2012 |
| | | EP 1716216 A2 | 02/11/2006 |
| | | EP 1716216 B1 | 18/11/2009 |
| | | EP 1716216 B9 | 10/03/2010 |
| | | EP 1725628 A1 | 29/11/2006 |
| | | EP 1725628 B1 | 30/05/2012 |
| | | EP 1740518 A1 | 10/01/2007 |
| | | EP 1740520 A1 | 10/01/2007 |
| | | EP 1740520 B1 | 12/11/2014 |
| | | EP 1740521 A1 | 10/01/2007 |
| | | EP 1740521 B1 | 20/05/2015 |
| | | EP 1893714 A2 | 05/03/2008 |
| | | EP 1893714 B1 | 05/11/2014 |
| | | EP 1893715 A2 | 05/03/2008 |
| | | EP 1916231 A2 | 30/04/2008 |
| | | EP 1916231 A3 | 23/12/2009 |
| | | EP 1916231 B1 | 09/03/2016 |
| | | EP 1916232 A1 | 30/04/2008 |
| | | EP 1916232 B1 | 15/01/2014 |
| | | EP 1918269 A1 | 07/05/2008 |
| | | EP 1918269 B1 | 03/11/2010 |
| | | EP 1943202 A1 | 16/07/2008 |
| | | EP 1943202 B1 | 17/12/2014 |
| | | EP 1943203 A1 | 16/07/2008 |
| | | EP 1943203 B1 | 19/12/2012 |
| | | EP 1943204 A1 | 16/07/2008 |
| | | EP 1943204 B1 | 02/01/2013 |
| | | EP 1954663 A1 | 13/08/2008 |
| | | EP 1968922 A2 | 17/09/2008 |
| | | EP 1968922 B1 | 22/04/2015 |
| | | EP 1996668 A2 | 03/12/2008 |
| | | EP 2009075 A2 | 31/12/2008 |
| | | EP 2009075 A3 | 11/03/2009 |
| | | EP 2009075 B1 | 27/05/2015 |
| | | EP 2036943 A2 | 18/03/2009 |
| | | EP 2036943 A3 | 22/07/2009 |
| | | EP 2036943 B1 | 13/07/2011 |
| | | EP 2076478 A1 | 08/07/2009 |
| | | EP 2076478 B1 | 03/10/2012 |
| | | EP 2080748 A2 | 22/07/2009 |
| | | EP 2080748 A3 | 21/09/2011 |
| | | EP 2085422 A2 | 05/08/2009 |
| | | EP 2085422 A3 | 06/03/2013 |
| | | EP 2098581 A2 | 09/09/2009 |
| | | EP 2098581 A3 | 04/11/2009 |
| | | EP 2103587 A2 | 23/09/2009 |
| | | EP 2103587 A3 | 03/03/2010 |
| | | EP 2103587 B1 | 20/02/2013 |
| | | EP 2146945 A1 | 27/01/2010 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2016/033450

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|-------------------------------------------|---------------------|----------------------------|---------------------|
| | | EP 2149543 A1 | 03/02/2010 |
| | | EP 2149592 A2 | 03/02/2010 |
| | | EP 2149592 B1 | 13/04/2016 |
| | | EP 2154223 A1 | 17/02/2010 |
| | | EP 2154223 B1 | 09/01/2013 |
| | | EP 2163591 A2 | 17/03/2010 |
| | | EP 2163591 A3 | 16/11/2011 |
| US 2010-0036179 A1 | 11/02/2010 | AT 512125 T | 15/06/2011 |
| | | CN 101665403 A | 10/03/2010 |
| | | CN 103052614 A | 17/04/2013 |
| | | EP 2151425 A2 | 10/02/2010 |
| | | EP 2151425 A3 | 31/03/2010 |
| | | EP 2151425 B1 | 08/06/2011 |
| | | EP 2585423 A2 | 01/05/2013 |
| | | EP 2585423 B1 | 30/09/2015 |
| | | JP 05595689 B2 | 24/09/2014 |
| | | JP 05868969 B2 | 24/02/2016 |
| | | JP 2010-043080 A | 25/02/2010 |
| | | JP 2013-533877 A | 29/08/2013 |
| | | KR 10-1619423 B1 | 10/05/2016 |
| | | KR 10-2010-0019397 A | 18/02/2010 |
| | | MX 2009008451 A | 22/03/2010 |
| | | US 2010-0331583 A1 | 30/12/2010 |
| | | US 2014-0100393 A9 | 10/04/2014 |
| | | US 2014-012047 A1 | 09/01/2014 |
| | | US 8664455 B2 | 04/03/2014 |
| | | US 8952208 B2 | 10/02/2015 |
| | | US 9181151 B2 | 10/11/2015 |
| | | WO 2012-009114 A2 | 19/01/2012 |
| US 2010-0331583 A1 | 30/12/2010 | CN 101665403 A | 10/03/2010 |
| | | CN 103052614 A | 17/04/2013 |
| | | CN 105037080 A | 11/11/2015 |
| | | EP 2151425 A2 | 10/02/2010 |
| | | EP 2151425 A3 | 31/03/2010 |
| | | EP 2151425 B1 | 08/06/2011 |
| | | EP 2585423 A2 | 01/05/2013 |
| | | EP 2585423 A4 | 27/11/2013 |
| | | EP 2585423 B1 | 30/09/2015 |
| | | ES 2556459 T3 | 18/01/2016 |
| | | JP 05595689 B2 | 24/09/2014 |
| | | JP 05868969 B2 | 24/02/2016 |
| | | JP 2010-043080 A | 25/02/2010 |
| | | JP 2013-533877 A | 29/08/2013 |
| | | JP 2016-053075 A | 14/04/2016 |
| | | KR 10-1619423 B1 | 10/05/2016 |
| | | MX 2013000049 A | 15/02/2013 |
| | | US 2010-0036179 A1 | 11/02/2010 |
| | | US 2014-0100393 A9 | 10/04/2014 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2016/033450

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|----------------------------------------|------------------|-------------------------|------------------|
| | | US 2015-152028 A1 | 04/06/2015 |
| | | US 8664455 B2 | 04/03/2014 |
| | | US 8952208 B2 | 10/02/2015 |
| | | WO 2012-009114 A2 | 19/01/2012 |
| | | WO 2012-009114 A3 | 05/04/2012 |
| US 2014-0275648 A1 | 18/09/2014 | CN 105026350 A | 04/11/2015 |
| | | EP 2970060 A1 | 20/01/2016 |
| | | JP 2016-511284 A | 14/04/2016 |
| | | KR 10-2015-0130330 A | 23/11/2015 |
| | | MX 2015011156 A | 09/11/2015 |
| | | WO 2014-165094 A1 | 09/10/2014 |
| US 2009-0287026 A1 | 19/11/2009 | AU 2003-284352 A1 | 13/05/2004 |
| | | AU 2003-284352 A8 | 13/05/2004 |
| | | AU 2003-286685 A1 | 13/05/2004 |
| | | AU 2003-286685 C1 | 23/05/2013 |
| | | AU 2005-238537 A1 | 10/11/2005 |
| | | AU 2006-261816 A1 | 04/01/2007 |
| | | AU 2006-261816 B2 | 19/05/2011 |
| | | AU 2010-245671 A1 | 11/11/2010 |
| | | AU 2010-245671 B2 | 29/01/2015 |
| | | AU 2010-246160 A1 | 11/11/2010 |
| | | AU 2011-248123 A1 | 10/11/2011 |
| | | AU 2011-248123 B2 | 22/01/2015 |
| | | CA 2503421 A1 | 06/05/2004 |
| | | CA 2503421 C | 06/01/2015 |
| | | CA 2557873 A1 | 10/11/2005 |
| | | CA 2557873 C | 25/11/2014 |
| | | CA 2564897 A1 | 17/11/2005 |
| | | CA 2564897 C | 27/11/2012 |
| | | CA 2564903 A1 | 17/11/2005 |
| | | CA 2564903 C | 04/12/2012 |
| | | CA 2564991 A1 | 17/11/2005 |
| | | CA 2564991 C | 19/03/2013 |
| | | CA 2608327 A1 | 27/04/2008 |
| | | CA 2608327 C | 14/04/2015 |
| | | CA 2608675 A1 | 27/04/2008 |
| | | CA 2608675 C | 31/03/2015 |
| | | CA 2612986 A1 | 04/01/2007 |
| | | CA 2613090 A1 | 04/01/2007 |
| | | CA 2628463 A1 | 18/05/2007 |
| | | CA 2628463 C | 08/07/2014 |
| | | CA 2635917 A1 | 12/07/2007 |
| | | CA 2646990 A1 | 27/09/2007 |
| | | CA 2674256 A1 | 30/01/2010 |
| | | CA 2706774 A1 | 28/05/2009 |
| | | CA 2711861 A1 | 16/07/2009 |
| | | CA 2745762 A1 | 10/06/2010 |
| | | CA 2761418 A1 | 11/11/2010 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2016/033450

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|----------------------------------------|------------------|-------------------------|------------------|
| | | CA 2761478 A1 | 11/11/2010 |
| | | CA 2776375 A1 | 07/04/2011 |
| | | CA 2798620 A1 | 10/11/2011 |
| | | CA 2822739 A1 | 04/01/2007 |
| | | CA 2826532 A1 | 10/11/2005 |
| | | CA 2826545 A1 | 06/05/2004 |
| | | CN 101014680 A | 08/08/2007 |
| | | CN 101014680 B | 03/09/2014 |
| | | CN 101014680 C | 08/08/2007 |
| | | CN 101177378 A | 14/05/2008 |
| | | CN 101177378 B | 09/04/2014 |
| | | CN 101177378 C | 14/05/2008 |
| | | CN 101182280 A | 21/05/2008 |
| | | CN 101182280 B | 27/06/2012 |
| | | CN 101182280 C | 21/05/2008 |
| | | CN 101248153 A | 20/08/2008 |
| | | CN 101248153 B | 24/07/2013 |
| | | CN 101248153 C | 20/08/2008 |
| | | CN 101260021 A | 10/09/2008 |
| | | CN 101260021 B | 06/08/2014 |
| | | CN 101260021 C | 10/09/2008 |
| | | CN 101283071 A | 08/10/2008 |
| | | CN 101283071 B | 24/07/2013 |
| | | CN 101351426 A | 21/01/2009 |
| | | CN 101351426 B | 08/05/2013 |
| | | CN 101351427 A | 21/01/2009 |
| | | CN 101351427 B | 21/08/2013 |
| | | CN 101351430 A | 21/01/2009 |
| | | CN 101395108 A | 25/03/2009 |
| | | CN 101395108 B | 13/03/2013 |
| | | CN 101448913 A | 03/06/2009 |
| | | CN 101492342 A | 29/07/2009 |
| | | CN 101495431 A | 29/07/2009 |
| | | CN 101553453 A | 07/10/2009 |
| | | CN 101553453 B | 29/05/2013 |
| | | CN 101597209 A | 09/12/2009 |
| | | CN 101665405 A | 10/03/2010 |
| | | CN 101665681 A | 10/03/2010 |
| | | CN 101665681 B | 25/06/2014 |
| | | CN 101796155 A | 04/08/2010 |
| | | CN 101925644 A | 22/12/2010 |
| | | CN 102015050 A | 13/04/2011 |
| | | CN 102015050 B | 02/07/2014 |
| | | CN 102026944 A | 20/04/2011 |
| | | CN 102112421 A | 29/06/2011 |
| | | CN 102112421 B | 26/11/2014 |
| | | CN 102140329 A | 03/08/2011 |
| | | CN 102264860 A | 30/11/2011 |
| | | CN 102264860 B | 17/06/2015 |
| | | CN 102281932 A | 14/12/2011 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2016/033450

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|----------------------------------------|------------------|-------------------------|------------------|
| | | CN 102281932 B | 06/01/2016 |
| | | CN 102307965 A | 04/01/2012 |
| | | CN 102307965 B | 06/01/2016 |
| | | CN 102439108 A | 02/05/2012 |
| | | CN 102596869 A | 18/07/2012 |
| | | CN 102782077 A | 14/11/2012 |
| | | CN 102803427 A | 28/11/2012 |
| | | CN 102971394 A | 13/03/2013 |
| | | CN 102971394 B | 20/04/2016 |
| | | CN 103146348 A | 12/06/2013 |
| | | CN 103146348 B | 13/01/2016 |
| | | CN 103172488 A | 26/06/2013 |
| | | CN 103172488 B | 21/10/2015 |
| | | CN 103215013 A | 24/07/2013 |
| | | CN 103274895 A | 04/09/2013 |
| | | CN 103467241 A | 25/12/2013 |
| | | CN 103497351 A | 08/01/2014 |
| | | CN 103497738 A | 08/01/2014 |
| | | CN 103627019 A | 12/03/2014 |
| | | CN 103642461 A | 19/03/2014 |
| | | CN 103923610 A | 16/07/2014 |
| | | CN 103923611 A | 16/07/2014 |
| | | CN 103923723 A | 16/07/2014 |
| | | CN 104045848 A | 17/09/2014 |
| | | CN 104045849 A | 17/09/2014 |
| | | CN 104045850 A | 17/09/2014 |
| | | CN 104046332 A | 17/09/2014 |
| | | CN 104059613 A | 24/09/2014 |
| | | CN 104164215 A | 26/11/2014 |
| | | CN 104164216 A | 26/11/2014 |
| | | CN 104164217 A | 26/11/2014 |
| | | CN 104178089 A | 03/12/2014 |
| | | CN 104178090 A | 03/12/2014 |
| | | CN 104194726 A | 10/12/2014 |
| | | CN 104194727 A | 10/12/2014 |
| | | CN 104194728 A | 10/12/2014 |
| | | CN 104194729 A | 10/12/2014 |
| | | CN 1732243 A | 08/02/2006 |
| | | CN 1732243 C | 08/02/2006 |
| | | CN 1852880 A | 25/10/2006 |
| | | CN 1852880 B | 15/06/2011 |
| | | CN 1852880 C | 25/10/2006 |
| | | CN 1898353 A | 17/01/2007 |
| | | CN 1898353 B | 07/05/2014 |
| | | CN 1898353 C | 17/01/2007 |
| | | CN 1902152 A | 24/01/2007 |
| | | CN 1902152 B | 13/10/2010 |
| | | CN 1902152 C | 24/01/2007 |
| | | CN 1968915 A | 23/05/2007 |
| | | CN 1968915 B | 03/08/2011 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2016/033450

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|----------------------------------------|------------------|-------------------------|------------------|
| | | CN 1968915 C | 23/05/2007 |
| | | CN 1972887 A | 30/05/2007 |
| | | CN 1972887 B | 13/10/2010 |
| | | CN 1972887 C | 30/05/2007 |
| | | CN 1976885 A | 06/06/2007 |
| | | CN 1976885 B | 26/05/2010 |
| | | CN 1976885 C | 06/06/2007 |
| | | EP 1563032 A2 | 17/08/2005 |
| | | EP 1563032 B1 | 02/05/2012 |
| | | EP 1578883 A2 | 28/09/2005 |
| | | EP 1658252 A2 | 24/05/2006 |
| | | EP 1658252 B1 | 04/01/2012 |
| | | EP 1678106 A2 | 12/07/2006 |
| | | EP 1678106 B1 | 04/01/2012 |
| | | EP 1716216 A2 | 02/11/2006 |
| | | EP 1716216 B1 | 18/11/2009 |
| | | EP 1716216 B9 | 10/03/2010 |
| | | EP 1725628 A1 | 29/11/2006 |
| | | EP 1725628 B1 | 30/05/2012 |
| | | EP 1740518 A1 | 10/01/2007 |
| | | EP 1740520 A1 | 10/01/2007 |
| | | EP 1740520 B1 | 12/11/2014 |
| | | EP 1740521 A1 | 10/01/2007 |
| | | EP 1740521 B1 | 20/05/2015 |
| | | EP 1893714 A2 | 05/03/2008 |
| | | EP 1893714 B1 | 05/11/2014 |
| | | EP 1893715 A2 | 05/03/2008 |
| | | EP 1916231 A2 | 30/04/2008 |
| | | EP 1916231 A3 | 23/12/2009 |
| | | EP 1916231 B1 | 09/03/2016 |
| | | EP 1916232 A1 | 30/04/2008 |
| | | EP 1916232 B1 | 15/01/2014 |
| | | EP 1918269 A1 | 07/05/2008 |
| | | EP 1918269 B1 | 03/11/2010 |
| | | EP 1943202 A1 | 16/07/2008 |
| | | EP 1943202 B1 | 17/12/2014 |
| | | EP 1943203 A1 | 16/07/2008 |
| | | EP 1943203 B1 | 19/12/2012 |
| | | EP 1943204 A1 | 16/07/2008 |
| | | EP 1943204 B1 | 02/01/2013 |
| | | EP 1954663 A1 | 13/08/2008 |
| | | EP 1968922 A2 | 17/09/2008 |
| | | EP 1968922 B1 | 22/04/2015 |
| | | EP 1996668 A2 | 03/12/2008 |
| | | EP 2009075 A2 | 31/12/2008 |
| | | EP 2009075 A3 | 11/03/2009 |
| | | EP 2009075 B1 | 27/05/2015 |
| | | EP 2036943 A2 | 18/03/2009 |
| | | EP 2036943 A3 | 22/07/2009 |
| | | EP 2036943 B1 | 13/07/2011 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2016/033450

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|-------------------------------------------|---------------------|----------------------------|---------------------|
| | | EP 2076478 A1 | 08/07/2009 |
| | | EP 2076478 B1 | 03/10/2012 |
| | | EP 2080748 A2 | 22/07/2009 |
| | | EP 2080748 A3 | 21/09/2011 |
| | | EP 2085422 A2 | 05/08/2009 |
| | | EP 2085422 A3 | 06/03/2013 |
| | | EP 2098581 A2 | 09/09/2009 |
| | | EP 2098581 A3 | 04/11/2009 |
| | | EP 2103587 A2 | 23/09/2009 |
| | | EP 2103587 A3 | 03/03/2010 |
| | | EP 2103587 B1 | 20/02/2013 |
| | | EP 2146945 A1 | 27/01/2010 |
| | | EP 2149543 A1 | 03/02/2010 |
| | | EP 2149592 A2 | 03/02/2010 |
| | | EP 2149592 B1 | 13/04/2016 |
| | | EP 2154223 A1 | 17/02/2010 |
| | | EP 2154223 B1 | 09/01/2013 |
| | | EP 2163591 A2 | 17/03/2010 |
| | | EP 2163591 A3 | 16/11/2011 |
| | | EP 2274260 A1 | 19/01/2011 |
| | | EP 2274260 A4 | 23/04/2014 |